



- (51) **International Patent Classification:**
A61K 31/415 (2006.01) *C07D 231/56* (2006.01)
- (21) **International Application Number:**
PCT/US2014/018895
- (22) **International Filing Date:**
27 February 2014 (27.02.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
PCT/CN2013/072134 4 March 2013 (04.03.2013) CN
- (71) **Applicant: MERCK SHARP & DOHME CORP.**
[US/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).
- (72) **Inventors; and**
- (71) **Applicants (for NZ, US only): MILLER, Michael** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **BASU, Kallol** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **DE-MONG, Duane** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **SCOTT, Jack** [US/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **LIU, Hong** [CN/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **DAI, Xing** [CN/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **STAMFORD, Andrew** [AU/US]; 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US). **POIRIER, Marc** [CA/US]; 2000 Galloping Hill Road, Kenilworth, New Jersey 07033 (US). **TEMPEST, Paul** [US/CN]; 965 Halei Road, Building #10, Room 310, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN).
- (74) **Common Representative: MERCK SHARP & DOHME CORP.;** 126 East Lincoln Avenue, Rahway, New Jersey 07065-0907 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
 - *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
- Published:**
- *with international search report (Art. 21(3))*



(54) **Title:** COMPOUNDS INHIBITING LEUCINE-RICH REPEAT KINASE ENZYME ACTIVITY

(57) **Abstract:** The present invention is directed to indazole compounds which are potent inhibitors of LRRK2 kinase and useful in the treatment or prevention of diseases in which the LRRK2 kinase is involved, such as Parkinson's Disease. The invention is also directed to pharmaceutical compositions comprising these compounds and the use of these compounds and compositions in the prevention or treatment of such diseases in which LRRK2 kinase is involved.

COMPOUNDS INHIBITING LEUCINE-RICH REPEAT KINASE ENZYME ACTIVITY

BACKGROUND OF THE INVENTION

Parkinson's disease (PD) is a common neurodegenerative disease caused by progressive loss of mid-brain dopaminergic neurons leading to abnormal motor symptoms such as bradykinesia, rigidity and resting tremor. Many PD patients also experience a variety of non-motor symptoms including cognitive dysfunction, autonomic dysfunction, emotional changes and sleep disruption. The combined motor and non-motor symptoms of Parkinson's disease severely impact patient quality of life.

While the majority of PD cases are idiopathic, there are several genetic determinants such as mutations in SNCA, Parkin, PINK1, DJ-1 and LRRK2. Linkage analysis studies have demonstrated that multiple missense mutations in the Leucine-Rich Repeat Kinase 2 (LRRK2) gene lead to an autosomal late onset form of PD. LRRK2 is a 286 kDa cytoplasmic protein containing kinase and GTPase domains as well as multiple protein-protein interaction domains. See for example, Aasly *et al.*, *Annals of Neurology*, Vol. 57(5), May 2005, pp. 762-765; Adams *et al.*, *Brain*, Vol. 128, 2005, pp. 2777-85; Gilks *et al.*, *Lancet*, Vol. 365, Jan. 29, 2005, pp. 415-416, Nichols *et al.*, *Lancet*, Vol. 365, Jan. 29, 2005, pp. 410-412, and U. Kumari and E. Tan, *FEBS journal* 276 (2009) pp. 6455-6463.

In vitro biochemical studies have demonstrated that LRRK2 proteins harboring the PD associated proteins generally confer increased kinase activity and decreased GTP hydrolysis compared to the wild type protein (Guo *et al.*, *Experimental Cell Research*, Vol. 313, 2007, pp. 3658-3670) thereby suggesting that small molecule LRRK2 kinase inhibitors may be able to block aberrant LRRK2-dependent signaling in PD. In support of this notion, it has been reported that inhibitors of LRRK2 are protective in models of PD (Lee *et al.*, *Nature Medicine*, Vol 16, 2010, pp. 998-1000).

LRRK2 protein has also been demonstrated to be associated with Lewy bodies, a pathological hallmark of PD as well as other neurodegenerative diseases such as Lewy body dementia (Zhu *et al.*, *Molecular Neurodegeneration*, Vol 30, 2006, pp. 1-17) thereby suggesting that LRRK2 may be associated with the pathogenesis of these

diseases.

A growing body of evidence also suggests a role for LRRK2 in immune cell function in the brain with LRRK2 inhibition demonstrated to attenuate microglial inflammatory responses (Moehle *et al.*, The Journal of Neuroscience Vol 32, 2012, pp. 1602-1611). Neuroinflammation is a hallmark of a number of neurodegenerative diseases such as PD and Alzheimer's disease, thereby suggesting that LRRK2 inhibitors may have utility in the treatment of neuroinflammation in these disorders.

Genome-wide association studies also highlight LRRK2 in the modification of susceptibility to the chronic autoimmune Crohn's disease and leprosy (Zhang *et al.*, The New England Journal of Medicine, Vol 361, 2009, pp. 2609-2618; Umeno *et al.*, Inflammatory Bowel Disease Vol 17, 2011, pp. 2407-2415). LRRK2 is also associated with certain types of cancer, e.g. melanoma as well as renal and thyroid carcinomas (Saunders-Pullman *et al.*, Movement Disorders, Vol 25, 2010, pp. 2536-2541; Looyenga, *et al.*, Proceedings of the National Academy of Sciences, USA, Vol 108, 2011, pp. 1439-1444).


Accordingly, compounds and compositions effective at modulating LRRK2 activity may provide a treatment for neurodegenerative diseases such as Parkinson's disease, Lewy body dementia, neuroinflammation, and for disease such as Crohn's disease, leprosy and cancer.

SUMMARY OF THE INVENTION

The present invention is directed to indazole compounds which are potent inhibitors of LRRK2 kinase and may be useful in the treatment or prevention of diseases in which the LRRK2 kinase is involved, such as Parkinson's Disease. The invention is also directed to pharmaceutical compositions comprising these compounds and the use of these compounds and compositions in the prevention or treatment of such diseases in which LRRK2 kinase is involved.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compounds of Formula I:



I

or a pharmaceutically acceptable salt thereof, wherein:

X^1 , X^2 and X^3 are each independently selected from the group consisting of a bond or CR^eR^f ;

Y is O, CR^aR^b or NR^c ;

Z is O, CR^aR^b or NR^c ;

R^1 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, hydroxyl, NR^cR^d , OR^5 and $(C=O)OR^5$;

R^2 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) R^5 ,
- d) R^7 ,
- e) OR^5 , and
- f) NR^cR^d ;

R^3 is selected from the group consisting of:

- a) hydrogen,
- b) C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR^5 and NR^cR^d ,
- c) cycloalkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR^5 and NR^cR^d ,

d) heterocyclyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, oxo, R^5 , OR^5 and NR^cR^d ,

e) heteroaryl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, oxo, R^5 , OR^5 and NR^cR^d ;

f) C_{4-8} cycloalkenyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR^5 and NR^cR^d ,

g) $(C=O)R^7$,

h) $(C=O)R^5$,

i) $S(O)_mR^5$, and

j) $S(O)_mR^7$;

or R^2 and R^3 can be taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:

a) halo,

b) oxo,

c) cyano,

d) OR^5 ,

e) NR^cR^d ,

f) SO_3H ,

g) $S(O)_mR^5$,

h) $S(O)_mR^7$

i) R^5 ,

j) R^6 ,

k) R^7 ,

l) $(C=O)R^5$,

m) $(C=O)OR^5$,

n) $(C=O)R^7$, and

o) $(C=O)NR^cR^d$;

R^4 is selected from the group consisting of hydrogen, halo, cyano, OR^5 , aryl, heteroaryl, C_{3-8} cycloalkyl, C_{4-8} cycloalkenyl, C_{4-8} heterocyclyl and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OC_{1-3} alkyl, NR^cR^d and hydroxyl;

R^5 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) hydroxyl,
- c) OC_{1-6} alkyl,
- d) NR^cR^d ,
- e) $(C=O)NR^cR^d$,
- f) $S(O)_m$,
- g) $S(O)_mR^8$,
- h) $S(O)_mR^7$,
- i) R^7 and
- j) OR^7 ;

R^6 is C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo and hydroxyl;

or R^5 and R^6 can be taken together with the atoms to which they are attached to form a 4 to 6 membered heterocyclic, 3 to 8 membered carbocyclic, aryl or heteroaryl ring, wherein said heterocyclic and heteroaryl rings may contain from one to three heteroatoms selected from N, O and S, wherein said heterocyclic, carbocyclic, aryl and heteroaryl rings are optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) oxo,
- c) cyano,
- d) hydroxyl,
- e) C_{1-3} alkyl, which is optionally substituted with one to three halo,
- f) C_{3-8} cycloalkyl,

- g) OC₁₋₃ alkyl, which is optionally substituted with one to three halo, and
- h) OC₃₋₈ cycloalkyl;

R⁷ is selected from the group consisting of C₄₋₈ heterocyclyl, C₃₋₈ cycloalkyl, C₄₋₈ cycloalkenyl, aryl or heteroaryl, wherein said heterocyclyl, cycloalkyl, cycloalkenyl, aryl and heteroaryl groups are optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) hydroxyl,
- d) oxo,
- e) C₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d,
- f) OC₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl NR^cR^d and aryl,
- g) C₃₋₈ cycloalkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d,
- h) aryl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl, S(O)_mNR^cR^d, C(O)NR^cR^d and NR^cR^d,
- i) heteroaryl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl, S(O)_mNR^cR^d, C(O)NR^cR^d and NR^cR^d,
- j) heterocyclyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, OC₁₋₃ alkyl and NR^cR^d,
- k) C₄₋₈ cycloalkenyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d;

R⁸ is hydrogen or C₁₋₆ alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) hydroxyl,
- d) OC₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo and NR^cR^d,
and
- e) C₃₋₈ cycloalkyl;

R^a is selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R^b is selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R^c is selected from the group consisting of:

- a) hydrogen and
- b) C₁₋₃ alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, hydroxyl, cyano, heteroaryl, aryl, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, OC₁₋₆ alkyl, and C₃₋₈ cycloalkyl;

R^d is selected from the group consisting of:

- a) hydrogen,
- b) C₃₋₈ cycloalkyl,
- c) C₃₋₆ heterocyclyl,
- d) C₁₋₃ alkyl,
- e) (C=O)C₁₋₃ alkyl,
- f) aryl, and
- g) heteroaryl;

wherein said cycloalkyl, heterocyclyl, alkyl, aryl and heteroaryl groups are each optionally substituted with one to three substituents independently selected from the group consisting of halo, hydroxyl, cyano, R⁸, SO₂R⁸, OC₁₋₆ alkyl and C₃₋₈ cycloalkyl;

or R^c and R^d can be taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three

heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of halo, cyano, hydroxyl, C₁₋₃ alkyl and OC₁₋₃ alkyl;

R^e is selected from the group consisting of hydrogen and C₁₋₃ alkyl;

R^f is selected from the group consisting of hydrogen and C₁₋₃ alkyl; and

m is an integer from zero to two.

In a class of the invention, X¹ is a bond. In another class of the invention, X¹ is CR^eR^f.

In a class of the invention, X² is a bond. In another class of the invention, X² is CR^eR^f.

In a class of the invention, X³ is a bond. In another class of the invention, X³ is CR^eR^f.

In a class of the invention, Y is O. In another class of the invention, Y is CR^aR^b. In another class of the invention, Y is NR^c.

In a class of the invention, Z is O. In another class of the invention, Z is CR^aR^b. In another class of the invention, Z is NR^c.

In a class of the invention, R¹ is selected from the group consisting of hydrogen and C₁₋₃ alkyl. In a subclass of the invention, R¹ is hydrogen. In another subclass of the invention, R¹ is methyl.

In a class of the invention, R² and R³ are taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:

- a) halo,
- b) oxo,
- c) cyano,
- d) OR⁵,
- e) NR^cR^d,
- f) SO₃H

- g) $S(O)_mR^5$,
- h) $S(O)_mR^7$,
- i) R^5 ,
- j) R^6 ,
- k) R^7 ,
- l) $(C=O)R^5$,
- m) $(C=O)OR^5$,
- n) $(C=O)R^7$, and
- o) $(C=O)NR^cR^d$.

In a subclass of the invention, R^2 and R^3 can be taken together with the atoms to which they are attached to form a 6 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:

- a) halo,
- b) oxo,
- c) OR^5 ,
- d) NR^cR^d ,
- e) $S(O)_mR^5$,
- f) $S(O)_mR^7$,
- g) R^5 ,
- h) R^6 ,
- i) R^7 ,
- j) $(C=O)R^5$,
- k) $(C=O)OR^5$, and
- l) $(C=O)R^7$.

In a further subclass of the invention, R^2 and R^3 can be taken together with the atoms to which they are attached to form a morpholinyl group, which is optionally substituted with C_{1-6} alkyl or C_{1-6} alkylhydroxy. In a further subclass of the invention, R^2 and R^3 can be taken together with the atoms to which they are attached to form a piperazinyl group, which is optionally substituted with C_{1-6} alkyl or C_{1-6} alkyl-SO₂. In a further subclass of

the invention, R² and R³ can be taken together with the atoms to which they are attached to form a piperazinone group.

In a class of the invention, R⁴ is selected from the group consisting of hydrogen and halo. In a subclass of the invention, R⁴ is hydrogen.

In a class of the invention, R⁵ is hydrogen.

In a class of the invention, R⁶ is hydrogen.

In a class of the invention, R⁷ is heterocyclyl.

In a class of the invention, R^a is hydrogen.

In a class of the invention, R^b is hydrogen.

In a class of the invention, R^c is hydrogen.

In a class of the invention, R^d is hydrogen.

In a class of the invention, R^e is hydrogen.

In a class of the invention, R^f is hydrogen.

Reference to the preferred classes and subclasses set forth above is meant to include all combinations of particular and preferred groups unless stated otherwise.

Specific embodiments of the present invention include, but are not limited to the compounds identified herein as Examples 1 to 37, or pharmaceutically acceptable salts thereof.

The invention also encompasses a pharmaceutical composition which comprises an inert carrier and the compound of Formula I, or a pharmaceutically acceptable salt thereof.

The invention may also encompass a method of treating Parkinson's Disease in a mammalian patient in need of such treatment, which comprises administering to the patient a therapeutically effective amount of the compound of Formula I, or a pharmaceutically acceptable salt thereof.

The invention may also encompass the use of a compound of Formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, for the manufacture of a medicament for the treatment of Parkinson's Disease.

The invention is also directed to medicaments or pharmaceutical compositions which may be useful for treating diseases or disorders in which LRRK2 is involved, such

as Parkinson's Disease, which comprise a compound of Formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The invention is also directed to the use of a compound of Formula I which may be useful for treating diseases or disorders in which LRRK2 is involved, such as Parkinson's Disease.

The invention is further directed to a method for the manufacture of a medicament or a composition which may be useful for treating diseases or disorders in which LRRK2 is involved, such as Parkinson's Disease, comprising combining a compound of Formula I with one or more pharmaceutically acceptable carriers.

The compounds of the present invention may contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. Additional asymmetric centers may be present depending upon the nature of the various substituents on the molecule. Each such asymmetric center will independently produce two optical isomers and it is intended that all of the possible optical isomers and diastereomers in mixtures and as pure or partially purified compounds are included within the ambit of this invention. Unless a specific stereochemistry is indicated, the present invention is meant to comprehend all such isomeric forms of these compounds.

The independent syntheses of these diastereomers or their chromatographic separations may be achieved as known in the art by appropriate modification of the methodology disclosed herein. Their absolute stereochemistry may be determined by the x-ray crystallography of crystalline products or crystalline intermediates which are derivatized, if necessary, with a reagent containing an asymmetric center of known absolute configuration.

If desired, racemic mixtures of the compounds may be separated so that the individual enantiomers are isolated. The separation can be carried out by methods well known in the art, such as the coupling of a racemic mixture of compounds to an enantiomerically pure compound to form a diastereomeric mixture, followed by separation of the individual diastereomers by standard methods, such as fractional crystallization or chromatography. The coupling reaction is often the formation of salts using an enantiomerically pure acid or base. The diastereomeric derivatives may then be

converted to the pure enantiomers by cleavage of the added chiral residue. The racemic mixture of the compounds can also be separated directly by chromatographic methods utilizing chiral stationary phases, which methods are well known in the art.

Alternatively, any enantiomer of a compound may be obtained by stereoselective synthesis using optically pure starting materials or reagents of known configuration by methods well known in the art.

In the compounds of Formula I, the atoms may exhibit their natural isotopic abundances, or one or more of the atoms may be artificially enriched in a particular isotope having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number predominantly found in nature. The present invention is meant to include all suitable isotopic variations of the compounds of generic Formula I. For example, different isotopic forms of hydrogen (H) include protium (1H) and deuterium (2H). Protium is the predominant hydrogen isotope found in nature. Enriching for deuterium may afford certain therapeutic advantages, such as increasing *in vivo* half-life or reducing dosage requirements, or may provide a compound useful as a standard for characterization of biological samples. Isotopically-enriched compounds within generic Formula I can be prepared without undue experimentation by conventional techniques well known to those skilled in the art or by processes analogous to those described in the Schemes and Examples herein using appropriate isotopically-enriched reagents and/or intermediates.

Tautomers of compounds defined in Formula I are also included within the scope of the present invention. For example, compounds including carbonyl $-\text{CH}_2\text{C}(\text{O})-$ groups (keto forms) may undergo tautomerism to form hydroxyl $-\text{CH}=\text{C}(\text{OH})-$ groups (enol forms). Both keto and enol forms are included within the scope of the present invention.

When any variable (e.g. R^a , etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds. Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms.

If the ring system is bicyclic, it is intended that the bond be attached to any of the suitable atoms on either ring of the bicyclic moiety.

It is understood that one or more silicon (Si) atoms can be incorporated into the compounds of the instant invention in place of one or more carbon atoms by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art from readily available starting materials. Carbon and silicon differ in their covalent radius leading to differences in bond distance and the steric arrangement when comparing analogous C-element and Si-element bonds. These differences lead to subtle changes in the size and shape of silicon-containing compounds when compared to carbon. One of ordinary skill in the art would understand that size and shape differences can lead to subtle or dramatic changes in potency, solubility, lack of off-target activity, packaging properties, and so on. (Diass, J. O. *et al.* *Organometallics* (2006) 5:1188-1198; Showell, G.A. *et al.* *Bioorganic & Medicinal Chemistry Letters* (2006) 16:2555-2558).

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase “optionally substituted with one or more substituents” should be understood as meaning that the group in question is either unsubstituted or may be substituted with one or more substituents.

As used herein, “alkyl” is intended to mean linear or branched structures having no carbon-to-carbon double or triple bonds. Thus, C₁₋₄ alkyl is defined to identify the group as having 1, 2, 3 or 4 carbons in a linear or branched arrangement, such that C₁₋₄ alkyl specifically includes, but is not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl and tert-butyl.

As appreciated by those of skill in the art, “halo” or “halogen” as used herein is intended to include chloro (Cl), fluoro (F), bromo (Br) and iodo (I).

The term "cycloalkyl" shall mean cyclic rings of alkanes of three to eight total carbon atoms, unless otherwise indicated, or any number within this range (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl or cyclooctyl) and also includes bicyclic or fused spirocyclic compounds.

The term "cycloalkenyl" shall mean cyclic rings of four to eight total carbon atoms, unless otherwise indicated, or any number within this range where one or two degrees of unsaturation are present. Non-limiting examples of said cycloalkenyl groups are: cyclohexenyl, cyclopentenyl, cyclooctadienyl.

The term "carbocycle" shall mean cyclic rings of three to eight total carbon atoms, unless otherwise indicated, or any number within this range, where zero, one or two degrees of unsaturation are present and where said "carbocycle" can be bicyclic or fused spirocyclic in nature. Non-limiting examples of said carbocyclyl groups are: cyclohexenyl, cyclopentenyl, cyclooctadienyl, cyclohexyl or cyclopropyl.

If no number of carbon atoms is specified, the term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight or branched, containing from 2 to 10 carbon atoms and at least 1 carbon to carbon double bond. Preferably 1 carbon to carbon double bond is present, and up to 4 non-aromatic carbon-carbon double bonds may be present. Thus, "C2-C6 alkenyl" means an alkenyl radical having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

"Aryl" means an aromatic monocyclic or multicyclic ring system comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

The term "heteroaryl", as used herein, represents a stable monocyclic, bicyclic or tricyclic ring of up to 10 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to:

benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, indolinyl, indolyl, indolazinylyl, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydroindolyl, dihydroquinolyl, methylenedioxybenzene, benzothiazolyl, benzothienyl, quinolinyl, isoquinolinyl, oxazolyl, and tetra-hydroquinoline. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively. If the heteroaryl contains nitrogen atoms, it is understood that the corresponding N-oxides thereof are also encompassed by this definition.

The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 5- to 10-membered nonaromatic ring, unless otherwise specified, containing from 1 to 4 heteroatoms selected from the group consisting of O, N, S, SO, or SO₂ and includes bicyclic groups. The heterocyclic group also includes rings that possess one or two degrees of unsaturation. "Heterocyclyl" therefore includes, but is not limited to the following: piperazinyl, piperidinyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, dihydropiperidinyl, tetrahydrothiophenyl and the like. If the heterocycle contains a nitrogen, it is understood that the corresponding N-oxides thereof are also encompassed by this definition.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives wherein the parent compound is modified by making acid or base salts thereof. Salts in the solid form may exist in more than one crystal structure, and may also be in the form of

hydrates. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganese salts, manganous, potassium, sodium, zinc, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. In one aspect of the invention the salts are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, fumaric, and tartaric acids. It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

Exemplifying the invention is the use of the compounds disclosed in the Examples and herein. Specific compounds within the present invention include a compound which may be selected from the group consisting of the compounds disclosed in the following Examples and pharmaceutically acceptable salts thereof and individual diastereomers thereof.

The subject compounds are useful in a method of inhibition of LRRK2 receptors in a patient such as a mammal in need of such antagonism comprising the administration of an effective amount of the compound. The present invention is directed to the use of the compounds disclosed herein as inhibitors of LRRK2 receptors. In addition to

primates, especially humans, a variety of other mammals can be treated according to the method of the present invention.

Another embodiment of the present invention is directed to a method for the treatment, control, amelioration, or reduction of risk of a disease or disorder in which the LRRK2 kinase is involved in a patient that comprises administering to the patient a therapeutically effective amount of a compound that is an inhibitor of LRRK2 kinase.

The present invention is further directed to a method for the manufacture of a medicament for inhibition of LRRK2 receptors activity in humans and animals comprising combining a compound of the present invention with a pharmaceutical carrier or diluent.

The subject treated in the present methods is generally a mammal, for example a human being, male or female, in whom inhibition of LRRK2 kinase activity is desired. The term "therapeutically effective amount" means the amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

The terms "treating" or "treatment" of a disease as used herein includes: inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms; or relieving the disease, i.e., causing regression of the disease or its clinical symptoms. The term "preventing" or "prevention" of a disease as used herein includes: causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease, and the like.

The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. Such term in relation to pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical

compositions of the present invention encompass any composition made by admixing a compound of the present invention and a pharmaceutically acceptable carrier. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the terms "administration of" or "administering a" compound shall encompass the treatment of the various conditions described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier, 1985. Metabolites of these compounds include active species produced upon introduction of compounds of this invention into the biological milieu.

The ability of the compounds of the present invention to act as LRRK2 kinase inhibitors may make them useful pharmacological agents for disorders that involve LRRK2 kinase in humans and animals, but particularly in humans.

In another embodiment the invention provides a method of inhibiting LRRK2 Kinase activity (this is to say, inhibiting the kinase activity associated with Leucine-Rich Repeat Kinase 2 [LRRK2], a multidomain protein containing kinase and GTPase enzymatic activities) in a patient in need of therapy for a condition amenable to treatment by such kinase activity inhibition, for example, treatment or prevention of neurologic damage associated with Parkinson's disease, for example, improvement in dopaminergic tone and in providing symptomatic benefit, for example, in treating, alleviating, ameliorating, or managing motor and non-motor symptoms of Parkinson's disease, and other conditions that may be treated or prevented by inhibition of LRRK2 kinase. Of particular importance is the acute or prophylactic treatment of Parkinson's Disease.

The subject compounds may be further useful in a method for the prevention, treatment, control, amelioration, or reduction of risk of the diseases, disorders and conditions noted herein.

The subject compounds may be further useful in a method for the prevention, treatment, control, amelioration, or reduction of risk of the aforementioned diseases, disorders and conditions in combination with other agents.

The compounds of the present invention may be used in combination with one or more other drugs in the treatment, prevention, control, amelioration, or reduction of risk of diseases or conditions for which compounds of Formula I or the other drugs may have utility, where the combination of the drugs together are safer or more effective than either drug alone. Such other drug(s) may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with a compound of Formula I. When a compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition in unit dosage form containing such other drugs and the compound of Formula I is preferred. However, the combination therapy may also include therapies in which the compound of Formula I and one or more other drugs are administered on different overlapping schedules. It is also contemplated that when used in combination with one or more other active ingredients, the compounds of the present invention and the other active ingredients may be used in lower doses than when each is used singly. Accordingly, the pharmaceutical compositions of the present invention include those that contain one or more other active ingredients, in addition to a compound of Formula I.

For example, the present compounds may be used in conjunction with one or more additional therapeutic agents, for example: L-DOPA; dopaminergic agonists such as quinpirole, ropinirole, pramipexole, pergolide and bromocriptine; MAO-B inhibitors such as rasagiline, deprenyl and selegiline; DOPA decarboxylase inhibitors such as carbidopa and benserazide; and COMT inhibitors such as tolcapone and entacapone; or potential therapies such as an adenosine A2a antagonists, metabotropic glutamate receptor 4 modulators, or growth factors such as brain derived neurotrophic factor (BDNF), and a pharmaceutically acceptable carrier.

The above combinations include combinations of a compound of the present invention not only with one other active compound, but also with two or more other active compounds. Likewise, compounds of the present invention may be used in combination with other drugs that are used in the prevention, treatment, control, amelioration, or reduction of risk of the diseases or conditions for which compounds of the present invention are useful. Such other drugs may be administered, by a route and in an amount commonly used therefore, contemporaneously or sequentially with a compound of the present invention. When a compound of the present invention is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of the present invention is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of the present invention.

The weight ratio of the compound of the present invention to the other active ingredient(s) may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the present invention is combined with another agent, the weight ratio of the compound of the present invention to the other agent will generally range from about 1000:1 to about 1:1000, or from about 200:1 to about 1:200. Combinations of a compound of the present invention and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

In such combinations the compound of the present invention and other active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of other agent(s), and via the same or different routes of administration.

The compounds of the present invention may be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), by inhalation spray, nasal, vaginal, rectal, sublingual, buccal or topical routes of administration and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically

acceptable carriers, adjuvants and vehicles appropriate for each route of administration. In addition to the treatment of warm-blooded animals the compounds of the invention are effective for use in humans.

The pharmaceutical compositions for the administration of the compounds of this invention may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases. As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, solutions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the

gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in the U.S. Patents 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release. Oral tablets may also be formulated for immediate release, such as fast melt tablets or wafers, rapid dissolve tablets or fast dissolve films.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation.

These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, jellies, solutions or suspensions and the like, containing the compounds of the present invention are employed. Similarly, transdermal patches may also be used for topical administration.

The pharmaceutical composition and method of the present invention may further comprise other therapeutically active compounds as noted herein which are usually applied in the treatment of the above mentioned pathological conditions.

In the treatment, prevention, control, amelioration, or reduction of risk of conditions which require inhibition of LRRK2 kinase activity an appropriate dosage level will generally be about 0.01 to 500 mg per kg patient body weight per day which can be administered in single or multiple doses. A suitable dosage level may be about 0.01 to 250 mg/kg per day, about 0.05 to 100 mg/kg per day, or about 0.1 to 50 mg/kg per day. Within this range the dosage may be 0.05 to 0.5, 0.5 to 5 or 5 to 50 mg/kg per day. For oral administration, the compositions are may be provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, or may be administered once or twice per day.

When treating, preventing, controlling, ameliorating, or reducing the risk of Parkinson's Disease, or other diseases for which compounds of the present invention are indicated, generally satisfactory results are obtained when the compounds of the present invention are administered at a daily dosage of from about 0.1 milligram to about 100 milligram per kilogram of animal body weight, given as a single daily dose or in divided doses two to six times a day, or in sustained release form. For most large mammals, the total daily dosage is from about 1.0 milligrams to about 1000 milligrams, or from about 1 milligram to about 50 milligrams. In the case of a 70 kg adult human, the total daily

dose will generally be from about 7 milligrams to about 350 milligrams. This dosage regimen may be adjusted to provide the optimal therapeutic response.

It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

Methods for preparing the compounds of this invention are illustrated in the following Schemes and Examples. Starting materials are made according to procedures known in the art or as illustrated herein.

The compounds of the present invention can be prepared readily according to the following Schemes and specific examples, or modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art but are not mentioned in greater detail. The general procedures for making the compounds claimed in this invention can be readily understood and appreciated by one skilled in the art from viewing the following Schemes.


While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various adaptations, changes, modifications, substitutions, deletions, or additions of procedures and protocols may be made without departing from the spirit and scope of the invention. For example, effective dosages other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications with the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compounds selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is

intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

General Schemes

A general procedure for the preparation of cycloalkoxy substituted indazoles such as **Formula I** is shown in **Scheme 1**. Treatment of a suitable bromo-indazole **i** with iodine/KOH and the like will provide compound **ii**. The indazole can be protected with SEMCl and the like to provide the protected indazole **iii**. The iodo group in **iii** can be coupled with a boronic acid **iv** and the like to provide indazole **v**. The bromide in **v** can be converted into the pinacol boronic ester **vii** using **vi** under palladium catalyzed conditions and the like. The boronic ester **v** can be oxidized to the alcohol **viii** using acetic acid/H₂O₂ and the like. The alcohol in **viii** can be reacted with electrophiles (LG = leaving group) such as **ix** in the presence of K₂CO₃ followed by removal of the indazole protecting group with TBAF and the like to provide examples such as **Formula I**. Alternatively, the SEM protecting group can be removed by a two step method wherein the starting material is first treated with a mixture such as TFA in CH₂Cl₂ and the like followed by concentration and treatment of the residue with ammonium hydroxide and the like.


Scheme 1



Alternatively, treatment of **i** with NIS followed by a base such as NaH and the like and TrCl will provide **xi**. The iodo indazole **xi** can be converted into the pyridyl substituted indazole **xii** via palladium-catalyzed cross coupling with a boronic acid or

ester such as **iv** and the like. The aryl bromide **xii** can be converted into the alcohol **xiii** by first conversion to the pinacol boronate via pinacol diborane and a palladium catalyst and the like, followed by oxidation of the pinacol boronate with hydrogen peroxide/acetic acid and the like. The alcohol **xiii** can be converted into examples such as **Formula I** using a base such as cesium carbonate and the like and an appropriate electrophile **ix** (LG = leaving group) and the like followed by removal of the trityl group with a mixture of triethylsilane/TFA and the like. Alternatively, the trityl group can be removed with a mixture of hydrogen chloride in methanol.


Scheme 2



Treatment of **xiv** with an appropriate alcohol **xv** and base and the like will provide the ether **xvi**. The bromide **xvi** can be converted into the methylated intermediate **xvii** using palladium catalyst and $\text{Me}_3\text{B}_3\text{O}_3$ and the like. The nitro derivative **xvii** can be treated with Pd/C in the presence of HCOONH_4 to produce the


amine **xvii**. The amine **xvii** can be acylated to provide **xix**. The acylated amine **xix** can be treated with iso-amyl nitrate in the presence of Ac₂O/KOAc and the like to provide the indazole **xx**. The indazole can be treated with ammonia and the like to produce **xxi**. The indazole **xxi** can be treated with iodine and KOH and the like to provide the iodo-indazole **xxii**. The indazole can be protected with trityl chloride and base and the like to provide **xxiii**. The iodo-indazole **xxiii** can be converted into **xxiv** using the appropriate boronic acid and palladium catalyst and the like. The chloro-pyridine **xxiv** can be converted into examples such as **Formula I** using the appropriate amine (HN(R²)R³), palladium catalyst, and ligand followed by deprotection of the trityl group using standard conditions.

Scheme 3



Treatment of alcohol **xxv** with 2-chloroethyl 4-methylbenzenesulfonate and base will provide **xxvi**. The indazole **xxvi** can be converted into **xxvii** using iodine and K_2CO_3 and the like. The indazole **xxvii** can be converted into the trityl protected indazole **xxviii** using trityl chloride and a base. The chloro-ethyl group in **xxviii** can be converted into the vinyl ether **xxix** using an appropriate base and the like. The vinyl ether **xxix** can be converted into the cyclopropoxy indazole **xxx** using Et_2Zn , CH_2I_2 , and TFA and the like. The iodo-indazole **xxx** can be converted into the chloro-pyridine **xxxi** using the appropriate boronic acid (or ester) and palladium catalyst and the like. The chloro-pyridine **xxxi** can be converted into examples such as **xxxii** using a palladium catalyst, appropriate ligand, and amine followed by deprotection of the trityl group using standard conditions.


Scheme 4



Treatment of indazole **ii** with SEMCl and a base will provide the SEM protected indazole **iii**. The iodo-indazole **iii** can be converted into the fluoro-pyridine **xxxiii** using an appropriate palladium catalyst and boronic acid and the like. The fluoro-pyridine **xxxiii** can be converted into the amino-pyridine **xxxiv** using the appropriate amine ($\text{HN}(\text{R}^2)\text{R}^3$) and base under standard conditions. The bromo-indazole **xxxiv** can be converted into the boronic ester **vii** using pinacol diborane and the appropriate palladium catalyst and the like. The boronate ester **vii** can be oxidized to the alcohol **viii**

using standard conditions such as H_2O_2 in acetic acid. The alcohol **viii** can be reacted with an appropriate electrophilic reagent (where LG = leaving group) and base to provide **xxxiv** using standard conditions. Treatment of **xxxiv** with TFA will provide examples such as **Formula I**.


Scheme 5



The chloro-pyridine **xxiv** can be reacted with an amine ($\text{HN(R}^2\text{)R}^3$) and appropriate base to provide **xxxvi** using standard conditions. The trityl protected

indazole xxxvi can be converted into examples such as **Formula I** using HCl in MeOH and the like.

Scheme 6



Experimentals

Abbreviations used in the experimentals may include the following:

ACN	Acetonitrile	AcOH	Acetic acid
Aq	Aqueous	Bn	Benzyl
BOC	<i>tert</i> -Butoxycarbonyl	BOC ₂ O	BOC Anhydride
Bu	Butyl	C (or °C)	degrees Celsius
Cbz	benzyloxycarbonyl	DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DCM	Dichloromethane	DIPEA	Diisopropylethylamine
DMA	N,N-Dimethylacetamide	DMAP	4-Dimethylaminopyridine
DME	1,2-dimethoxyethane	DMF	Dimethylformamide
DMSO	Dimethyl sulfoxide	DPPF	1,1'-(bis-diphenylphosphino)ferrocene
EDCI	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride	EDC	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EI	Electron ionization	Eq	Equivalents
Et	Ethyl	EtOAc	Ethyl acetate

EtOH	Ethanol	g	grams
h, hr	hours	¹ H	proton
HATU	N,N,N',N'-Tetramethyl-O-(7-Azabenzotriazol-1-yl)uronium hexafluorophosphate	Hex	hexanes
HOBT	1-Hydroxybenzotriazole	HOBT.H ₂ O	1-Hydroxybenzotriazole hydrate
HOTS	<i>para</i> -toluene sulfonic acid (see also TsOH)	HOTS.H ₂ O	<i>para</i> -toluene sulfonic acid hydrate (see also TsOH.H ₂ O)
HMPA	hexamethylphosphoramide	HPLC	High pressure liquid chromatography
IPA	isopropanol, 2-propanol	LDA	lithium diisopropylamide
M	Molar	mmol	milimolar
mCPBA	<i>meta</i> -Chloroperoxy benzoic acid	Me	Methyl
MeCN	Acetonitrile	MeOH	Methanol
min	Minutes	mg	Milligrams
MHZ	Megahertz	mL (or ml)	Milliliter
Mol sieves	molecular sieves	N	normal
NMR	Nuclear Magnetic Resonance	MS	Mass Spectroscopy
NBS	N-Bromosuccinimide	NMM	N-Methylmorpholine
NMP	1-methyl-2-pyrrolidone	ON	Overnight
PTLC	Preparative thin layer chromatography	PyBrOP	Bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
PyBOP	(Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate	Pyr	Pyridine
Quant	quantitative	RT or rt	Room temperature
sat (or sat. or sat'd.)	Saturated	SFC	supercritical fluid chromatography
sgc	Silica gel 60 chromatography	SiO ₂	Silica gel
tBOC	<i>tert</i> -Butoxycarbonyl	<i>t</i> -Bu	<i>tert</i> -butyl
TEA	Triethylamine	Tf	Trifluoromethane sulfonyl
TFA	Trifluoroacetic acid	THF	Tetrahydrofuran
TLC	Thin layer chromatography	Ts	Toluene sulfonyl
TsOH	<i>para</i> -toluene sulfonic acid	TsOH.H ₂ O	<i>para</i> -toluene sulfonic acid hydrate

General Experimental Information:

Unless otherwise noted, all reactions are magnetically stirred.

Unless otherwise noted, when ethyl acetate, hexanes, dichloromethane, 2-propanol, and methanol are used in the experiments described below, they are Fisher Optima grade solvents.


Unless otherwise noted, when diethyl ether is used in the experiments described below, it is Fisher ACS certified material and is stabilized with BHT.


Unless otherwise noted, "concentrated to dryness" means evaporating the solvent from a solution or mixture using a rotary evaporator.

Unless otherwise noted, flash chromatography is carried out on an Isco, Analogix, or Biotage automated chromatography system using a commercially available cartridge as the column. Columns may be purchased from Isco, Analogix, Biotage, Varian, or Supelco and are usually filled with silica gel as the stationary phase.


Unless otherwise noted, all LRRK2 IC₅₀ data presented in tables refers to the LRRK2 Km ATP LanthaScreen™ Assay that is described in the Biological Assay section.

Scheme A




Step 1

5-Bromoindazole (10 grams) and KOH (5.7 g) were taken up in DMF (83 ml) at RT. Iodine (19.3 g) was added, and the resulting solution was stirred at RT for 24 h. The solution was diluted with EtOAc and 10% $NaHSO_3$ (aq.). The aqueous layer was extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over $MgSO_4$. Filtration and concentration afforded 3-iodo-5-bromoindazole.


Step 2

3-Iodo-5-bromo indazole (12.4 g), nBu_4NBr (124 mg), and KOH (19 g) were partitioned between water (19.5 g) and DCM (51 ml) at $0^\circ C$. After stirring at $0^\circ C$ for 1 h, the mixture was warmed to RT. After stirring at RT for 2 h, the mixture was extracted with DCM. The combined organic layers were washed with water, brine, and dried over $MgSO_4$. Filtration and concentration afforded the crude product. Purification via flash chromatography (hexanes/EtOAc, 0-10% EtOAc gradient over 17 minutes, 10%-50% EtOAc gradient over 5 minutes, SiO_2) provided the SEM protected indazole.

Step 3


The SEM protected indazole (2.1 g), PdCl₂(dppf) (360 mg), K₂CO₃ (1.2 g), and boronic acid (918 mg) were taken up in dioxane/water (20 ml/2 ml), and the resulting solution was stirred at 100 °C for 2 h. The mixture was extracted with EtOAc. The combined organic layers were washed with water, brine, and dried over Na₂SO₄. Filtration and concentration gave the crude product. Purification via flash chromatography (0-3% MeOH in DCM gradient, SiO₂) provided the desired arylated indazole.

Step 4




The bromo indazole (500 mg), pinacol diborane (389 mg), PdCl₂(dppf) (166 mg), and KOAc (400 mg) were taken up in dioxane (10 ml), and the resulting solution was microwaved at 120 °C for 20 minutes. The solution was concentrated, and the residue was purified via flash chromatography (0-3% MeOH in DCM gradient, SiO₂) which provided the boronate as a yellow solid.

Step 5




The boronate (200 mg) was taken up in 2 ml of THF. Acetic acid (50 μL) and H₂O₂ (180 μL of 50% in water) were added, and the resulting solution was stirred at RT for 1 h. Another portion of acetic acid/hydrogen peroxide was added. After stirring at RT

for one hour, another portion of acetic acid/hydrogen peroxide was added, and the resulting solution was stirred at RT for 24 h. The solution was concentrated and partitioned between sat. NaHCO_3 (aq.) / ether. The aqueous layer was extracted with ether. The combined ether layers were dried (Na_2SO_4), filtered, and concentrated to afford the phenol as a yellow solid.





The phenol (130 mg), K_2CO_3 (136 mg), and cyclopropyl bromide (119 mg) were taken up in DMF (1 ml), and the solution was heated at 100 °C for 3 days. The DMF was removed under reduced pressure. The residue was purified via preparative thin-layer chromatography (1/1 hexanes/EtOAc, SiO_2) provided **Example 1** as a yellow solid.




The phenol (**Step 5 of Scheme A**) was converted into **Example 2** using conditions similar to those outlined in **Step 6 of Scheme A**.


Scheme C

**Step 1**

5-Bromo-6-fluoro indazole (2 g) and NIS (2.5 g) were taken up in CH₃CN (15 ml) and the solution was heated in a microwave at 105 °C for 30 minutes. The solution was concentrated. The residue was taken up in DCM (30 ml), and TrCl (3.9 g) and TEA (1.9 g) were added. The solution was stirred at RT for 48 h and then at 40 °C for 3 hr. The solution was concentrated. Purification via flash chromatography (0-20% DCM in hexanes gradient, SiO₂) provided the iodide as a yellow solid.





The iodide was converted into the phenol using conditions outlined in **Scheme A**.




The phenol (30 mg), Cs_2CO_3 (123 mg) and cyclopropyl bromide (39 mg) were taken up in DMSO (1ml), and the resulting solution was subjected to microwave conditions (200 °C for 1 h). The mixture was filtered, and the solution was concentrated. The residue was treated with 1.5 ml of TFA and 0.5 ml of Et_3SiH . The solution was stirred at RT for 10 minutes after which DCM (1 ml) was added. The resulting solution was stirred at RT for 2 h. The solution was concentrated and 7 N NH_3 in MeOH was added (5 ml). The solution was concentrated. The residue was purified via thin layer preparative chromatography (5% MeOH in DCM, SiO_2) which provided **Example 3** as a yellow solid.

Scheme D






To a cold (0 °C), stirred mixture of 2-bromo-4-fluoronitrobenzene (10.0 g, 45.5 mmol) and 1-methylcyclopropanol (3.61 g, 50.0 mmol) in DMF (200 ml) was added NaH (2.36 g of 60% in oil, 59.1 mmol) in portions. Once the addition was complete the cold bath was removed and the mixture was stirred at rt for 5 h. The reaction was quenched with water and extracted with EtOAc (x3). The combined organic layers were washed with water (x3), brine (x2), dried, filtered and concentrated to leave an oil which was purified by column chromatography (elution with 100:0 to 20:1 hexane:EtOAc) to yield the desired product as a light yellow oil.




To a stirred mixture of the ether (2.0 g, 7.35 mmol), K_2CO_3 (2.03 g, 14.70 mmol) and Cs_2CO_3 (2.39 g, 7.35 mmol) in Dioxane (110 ml) was purged Ar for 15 min. Then trimethylboraxine (2.26 ml, 16.17 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (0.85 g, 0.74 mmol) were added and the mixture was heated at 100 °C overnight. The reaction was cooled to room temperature and concentrated under vacuum. To this residue was added 10:1 hex:EtOAc (500 mL) and filtered through a pad of silica. The solid was washed with a mixture of hexane:EtOAc (1L of 10:1 hexane:EtOAc) solution. The filtrate was concentrated under vacuum to provide the desired product, which was used in the next step without further purification.


Step 3



The nitro derivative was dissolved in absolute EtOH (110 mL). To this solution were added 10% Pd/C (0.782 g, 0.735 mmol) and ammonium formate (5.56 g, 88.0 mmol) and the mixture was stirred at room temperature for 5 h. To this solution was added a solution containing 5:1 hexane:EtOAc (500 mL) and the mixture was filtered through a pad of silica. The filtrate was concentrated and the residue was purified by column chromatography (elution with 10:1 to 5:1 hexane:EtOAc) to yield the desired amine as a dark brown oil.




To a cold (0 °C), stirred mixture of amine (1.4 g, 7.90 mmol) and triethylamine (2.20 ml, 15.80 mmol) in CH₂Cl₂ (12 ml) was added Ac₂O (1.12 ml, 11.85 mmol). The mixture was slowly warmed to room temperature and stirred overnight. Silica gel was added and the mixture was evaporated to leave a slurry which was then purified by column chromatography (elution with 1:1 hexane:EtOAc) to yield the desired acetamide as a colorless gum.



To a stirred solution of the acetamide (1.76 g, 8.03 mmol) in toluene (42 ml) were added KOAc (1.182 g, 12.04 mmol) and Ac₂O (3.48 ml, 36.9 mmol). Then the mixture was heated to 80 °C when isoamyl nitrite (4.49 ml, 32.1 mmol) was added dropwise and the resulting mixture was heated at 80 °C overnight. The insoluble material was filtered


through a pad of celite and the filtrate was concentrated to leave a residue which was purified by column chromatography (elution with 20:1 hexane:EtOAc) to yield the desired product as a yellow solid.

Step 6




To a stirred suspension of the N-acetyl indazole (6.5 g, 28.2 mmol) in MeOH (20 mL) was added NH₃ (20.16 ml of 7M solution in MeOH, 141.0 mmol) and the mixture was stirred at RT for 2 h. The reaction was concentrated and the residue was purified by column chromatography (elution with 5:1 to 1:1 hexane:EtOAc) to yield the desired indazole as a yellow solid.

Step 7




To a stirred solution of indazole (5.0 g, 26.6 mmol) in DMF (50 ml) was added KOH (5.96 g, 106 mmol) followed by I₂ (13.48 g, 53.1 mmol). The mixture was stirred at room temperature for 3 h before being quenched with a saturated aqueous solution of Na₂S₂O₃ followed by addition of water. The resulting layer was extracted with EtOAc (x4). The combined organic layers were dried over MgSO₄, filtered and concentrated under vacuum to leave a residue which was purified by column chromatography (elution with 10:1 to 1:1 hexane:EtOAc) to yield the iodide as a yellow solid.

Step 8




To a cold (0 °C), stirred solution of iodide (5.1 g, 16.24 mmol) in THF (50 ml) was added NaH (0.779 g, 19.48 mmol) in portions. After the addition was complete the mixture was stirred at 0 °C for 15 min when trityl chloride (4.98 g, 17.86 mmol) was added in one portion. The resulting mixture was then slowly warmed to room temperature and stirred at room temperature for 4 h. The reaction was quenched with water and extracted with DCM (x3). The combined organic layers were dried over MgSO₄, filtered and concentrated to leave a residue which was purified by column chromatography (elution with 100% DCM) to yield the desired product as an off-white solid.




The iodide was reacted with (2-chloropyridin-4-yl)boronic acid using conditions outlined in **Step 3** of **Scheme A** which provided the desired chloro-pyridine.

Step 10




The chloro-pyridine (150 mg), 8-oxa-3-azabicyclo[3.2.1]octane (41 mg), 1,3-bis(2,6-diisopropylphenyl)-4,5-dihydro-1H-imidazol-3-ium chloride (36 mg), $\text{Pd}_2(\text{dba})_3$ (50 mg), and NaO^tBu (53 mg) were taken up in 2 ml of dioxane in a sealed tube. The solution was heated at 100 °C for 24 h. The solution was filtered and concentrated. The residue was treated with 2 ml of TFA and 0.5 ml of Et_3SiH . After stirring at RT for 2 h, the solution was concentrated. The residue was treated with 5 ml of 7 NH_3 in MeOH, and the resulting solution was concentrated. The residue was purified via thin-layer preparative chromatography (40% EtOAc in hexanes, SiO_2) which provided **Example 4** as a yellow solid.

Scheme E




The chloro-pyridine was reacted with 3-oxa-8-azabicyclo[3.2.1]octane using conditions outlined in **Step 10** of **Scheme D** to provide **Example 5**.

Scheme E-1




Example 6

Step 1




The hydroxyl-indazole (30 g) was taken up in DMF (200 ml). Potassium carbonate (62 g) was added to the solution, and the resulting mixture was stirred at RT for 15 minutes. 2-Chloroethyl-para-toluenesulfonate (43 ml) was added, and the resulting

solution was heated at 50 °C for 20 h. The mixture was poured into water (600 ml) and quenched with acetic acid (26 ml). The mixture was filtered which provided a wet cake that was dried under reduced pressure. The brown solid was partitioned between EtOAc and water. The organic layer was dried (MgSO₄), filtered, and concentrated. After slurry of the residue in MTBE, the mixture was filtered which provided the desired product.




The chloro-indazole (36 g) was taken up in CH₃CN (500 ml). Potassium carbonate (24 g) and iodine (3 g) were added, and the resulting solution was stirred at RT for 1 h. The mixture was poured into water (1 L), and extracted with MTBE. The organic layer was washed with 10% Na₂S₂O₃ (aq.) and dried over MgSO₄. The solution was filtered and concentrated. The residue was adsorbed onto SiO₂ (90 g) and purified on a short pad of SiO₂ using a gradient elution of 10-30 % EtOAc in hexanes.. The solution was concentrated which provided the desired iodide.




The indazole (17 g) was taken up in THF (100 ml) at 0 °C. Sodium hydride (2.0 g) was added to the solution. After stirring at 0 °C for 10 minutes, trityl chloride (13.7 g) was added. The solution was warmed to RT for 2 hours. The solution was quenched with sat. NH₄Cl (aq.) and extracted with Et₂O. The aqueous layer was extracted with Et₂O. The combined organic layers were washed with brine and dried over MgSO₄. Filtration and concentration provided the crude product. The residue was slurried in hot MTBE. The solid was collected and dried. The solid was purified via gradient flash


chromatography (0-10% MTBE in hexanes, SiO₂) which provided the trityl protected indazole.




The chloro-indazole (20 g) was suspended in THF (120 mL), and potassium tert-butoxide (8.0 g) was added. After stirring at 2 hours at RT, the solution was partitioned between sat. NH₄Cl (aq.) and Et₂O. The aqueous layer was extracted with Et₂O. The combined organic layers were dried over MgSO₄. Filtration and concentration provided the desired alkene.



Diethyl zinc (57 ml of a 1.0M solution in hexane) and 20 ml of DCE were placed into a dry flask under nitrogen at RT. The solution was cooled to 0 °C, and TFA (4.2 ml) in DCE (20 ml) was added over 30 minutes (exotherm/gas evolution). The white suspension was stirred at 0 °C for 15 minutes. Diiodomethane (4.6 ml) in DCE (20 ml) was added over 15 minutes at 0 °C. The alkene (15.7 g) in DCE (100 ml) was added to the solution at 0 °C. The solution was warmed to RT. After stirring at RT for 5 h, the solution was quenched with sat. NH₄Cl (aq.) and DCM. The aqueous layer was extracted with DCM. The combined organic layers were dried over MgSO₄. Filtration and concentration provided the crude product. The residue was purified via gradient flash chromatography (0-10% MTBE in hexanes, SiO₂) which provided the cyclopropoxy indazole.




The indazole (1.4 g), boronate ester (742 mg), K_3PO_4 (3.8 ml of a 2 M aqueous solution), and $\text{PdCl}_2(\text{dppf})$ - DCM complex (211 mg) were taken up in 20 ml of DME. Argon was bubbled through the solution, and the mixture was placed into a sealed tube. The reaction was subjected to microwave irradiation (2 hours at 90 °C). The solution was cooled, filtered, and concentrated. The residue was purified by gradient flash chromatography (0-20% EtOAc in hexanes, SiO_2) which provided the chloro-pyridine.




The chloro-pyridine was reacted with 4-oxa-7-azaspiro[2.5]octane using conditions outlined in **Step 10 of Scheme D** to provide **Example 6**.

Scheme F




Step 1




The indazole (2.2 g) was taken up in THF (25 ml) at 0 °C. Sodium hydride (275 mg of a 60 wt% dispersion in oil) was added in portions. After stirring at 0 °C for 15


minutes, SEMCl (1.1 g) was added. The solution was warmed to RT. After stirring at RT for 20 hours, the reaction was partitioned between EtOAc and sat. NH₄Cl (aq.). The aqueous layer was extracted with EtOAc. The combined organic layers were dried over MgSO₄. Filtration and concentration provided the crude product. The residue was purified by gradient flash chromatography (100/1 to 15/1 hexanes/EtOAc, SiO₂) to afford the desired SEM protected indazole.




The iodo-indazole (1.6 g), boronic acid (497 mg), K₃PO₄ (2.2 g), and nBu₄NBr (228 mg) were taken up in THF (32 ml). Nitrogen was bubbled through the solution for 15 minutes, and Pd(dppf)Cl₂ DCM complex (577 mg) was added. The resulting solution was heated at 80 °C for 48 hours. The mixture was filtered through a plug of celite. The filtrate was concentrated, and then it was partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc. The combined organic layers were dried over MgSO₄. Filtration and concentration provided the crude product. The residue was purified by gradient column chromatography (100/0 to 15/1 hexanes/EtOAc, SiO₂) to afford the arylated indazole.




The indazole (810 mg), Et₃N (3.9 g), and amine (1.9 g) were taken up in DMSO (4 ml). The resulting solution was heated in a sealed tube at 100 °C for 16 hours. The solution was cooled, and it was partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc. The combined organic layers were dried (MgSO₄), filtered, and concentrated. The residue was purified by flash chromatography (30/1 DCM/MeOH, SiO₂) which provided the amino-pyridine.




The bromo-indazole was converted into the boronate ester using conditions similar to those outlined in **Step 4** of **Scheme A**.




The boronate-ester was converted into the phenol using conditions similar to those outlined in **Step 5** of **Scheme A**.



The phenol (100 mg), 3-iodooxetane (122 mg), $n\text{Bu}_4\text{NI}$ (8 mg), and Cs_2CO_3 (215 mg) were placed into DMF (2ml). The resulting mixture was stirred at 80 °C for 20 hours. The mixture was filtered and concentrated. The residue was partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc. The combined organic layers were dried (MgSO_4), filtered, and concentrated. The residue was purified by flash chromatography (30/1 DCM/MeOH, SiO_2) which provided the desired ether.












The SEM protected indazole (69 mg) was taken up in 2 ml of DCM, and TFA (1 ml) was added. The solution was stirred at RT for 5 hours. The mixture was concentrated, and the residue was taken up in DCM (2 ml), MeOH (2 ml), and 1 M NH_4OH (2 ml). The mixture was stirred at RT for 20 hours. The mixture was diluted with water and extracted with DCM. The combined organic layers were dried (MgSO_4), filtered, and concentrated. The residue was purified by reverse phase chromatography to afford **Example 7**.










Parallel preparation of **Examples 8-26**: To a set of vials each containing the requisite amine (0.152 mmol) if a solid, was added 1,3-Bis-(2,6-diisopropylphenyl)imidazolium chloride (6.47 mg, 0.015 mmol), $\text{Pd}_2(\text{dba})_3$ (3.47 mg, 3.79 μmol) and sodium *t*-butoxide (29.1 mg, 0.303 mmol). The vials were transferred to a glove bag under an atmosphere of N_2 . A solution of the chloropyridine (**Scheme E-1**) (40 mg, 0.076 mmol) in dioxane (0.500 ml) was then added to each vial followed by the requisite amine (0.152 mmol) if it was a liquid. The vials were capped and removed from the glove bag. The vials were then placed into a pre-heated aluminium block at 90°C . The mixtures were stirred at that temperature overnight. Water (2 mL) was added to each vial followed by DCM (2.0 mL). The vials were shaken at RT for 5 min. The mixture was transferred to a fritted barrel filter. The organic layer from each vial was drained into a clean vial. Additional DCM (1 mL) was added to the aqueous layer and the layers were separated. The solvent from the combined organic layers was removed *in vacuo*. To each vial was then added DCM (1 mL) followed by TFA (0.50 mL) and water (0.050 mL). The vials were shaken at RT for 1.5 hours. The solvent was removed *in vacuo*. Each crude product was redissolved in 1 mL of DMSO and filtered. The crude products were purified by mass triggered HPLC. [Waters XBridge C18 column, 5 μm , 19 x100 mm, gradient ranges from 10-30% initial to 30-70% MeCN(0.1% NH_4OH) in water (0.1% NH_4OH) 50 mL/min, 8 min run





time] to provide the examples 8-25. Example 26 was repurified using the following conditions: [Waters Sunfire C18 column, 5 μ m, 19x100 mm, gradient 10% initial to 30% final MeCN(0.1% formic acid) in water (0.1 % formic acid) 50 mL/min, 8 min run time].





Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
1		40	A	1.0	337
2		20	B	1.3	353
3		1.2	B	1.9	355
4		23	B	2.0	377

Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
5		67	B	2.1	377
6		17	B	2.1	363
7		36	B	1.8	382
8		30	C	1.07	371.2
9		16	C	0.91	307.2

Ex	Structure	LRRK2	LCMS		
		IC ₅₀ (nM)	Method	RT (min)	m/z
10		23	C	0.78	367.2
11		111	C	0.80	411.2
12		10	C	0.84	402.2


Ex	Structure	LRRK2	LCMS		
		IC ₅₀ (nM)	Method	RT (min)	m/z
13		16	C	0.97	401.2
14		19	C	0.95	357.1
15		44	C	0.77	383.2
16		71	C	0.84	399.2


Ex	Structure	LRRK2	LCMS		
		IC ₅₀ (nM)	Method	RT (min)	m/z
17		51	C	0.88	428.2
18		10	C	0.76	396.1
19		54	C	1.07	414.2
20		22	C	1.05	371.2

Ex	Structure	LRRK2	LCMS		
		IC ₅₀ (nM)	Method	RT (min)	m/z
21		19	C	1.01	357.2
22		17	C	0.91	351.2
23		12	C	0.80	402.2
24		29	C	0.77	337.2


Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
25		40	C	0.97	343.1
26		12	C	0.99	376.2

Scheme H















The chloropyridine from **Step 9** of **Scheme D** (150 mg, 0.27 mmol), morpholine, and *N,N*-diisopropylethylamine (208 mg, 1.62 mmol) were taken up into DMSO (4 mL). The reaction was sealed and heated at 140 °C for 72 h. The mixture was cooled to room temperature and diluted with water (15 mL), and then extracted with EtOAc (30 mL \times 3). The combined organic layers were washed with water (45 mL), and brine (45 mL), dried over Na_2SO_4 , filtered and concentrated to leave a residue which was purified by a silica gel chromatography column (elute: petroleum ether/EtOAc = 10/1) to yield the desired product as a yellow solid. MS (ESI) m/z = 593.1 $[\text{M}+1]^+$.



The trityl protected indazole from the previous step (110 mg, 0.18 mmol) in methanol (4 mL) was added a solution of hydrogen chloride in methanol (3.0 M, 1.0 mL, 3.0 mmol). The mixture was heated at 70 °C for 6 h. The reaction was cooled, and then concentrated to leave a residue, which was purified by a silica gel chromatography column (elute: EtOAc/MeOH = 10/1) to yield the desired product as a white solid. MS (ESI) m/z = 351.1 $[\text{M}+1]^+$; RT = 1.6 minutes (LCMS conditions D).

The following examples (**28-37**) were prepared in a similar fashion to that depicted in **Scheme H** using the appropriate amine.


Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
27		0.6	D	1.6	351.1
28		0.6	D	1.7	365.1
29		0.6	D	1.7	365.1
30		0.9	D	1.7	379.2
31		0.6	D	1.7	428.1

Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
32		0.6	D	1.6	364.1
33		1.0	D	1.7	378.1
34		0.7	D	1.6	381.1
35 ^a		1.1	D	1.6	380.2
36 ^b		2.0	D	1.6	350.0

Ex	Structure	LRRK2 IC ₅₀ (nM)	LCMS		
			Method	RT (min)	m/z
37 ^b		2.6	D	1.5	350.0

^aThe preparation of the requisite Boc protected amine for **Example 35** is shown in Scheme I. The Boc group is cleaved in Step 2 of Method D.

Scheme I



Step 1: A mixture of L-tartaric acid (250 g, 1.67 mol) and BnNH₂ (178 g, 1.67 mol) in xylene (1.5 L) was stirred under reflux for 4 h using a Dean-Stark water separator. The reaction was cooled to room temperature with stirring and filtrated. The solid was washed with EtOH to afford the desired product as a yellow solid.

Step 2: To a stirred solution of diol from Step 1 (275 g, 1.24 mol) in DMF (1.8 L) was added imidazole (254 g, 3.73 mol) followed by TBSCl (467 g, 3.11 mol). The reaction mixture was stirred at rt overnight before being diluted with EtOAc. The resulting layer was washed with water and brine, dried, filtered and concentrated to leave a residue which was purified by column chromatography on silica gel (elution with petroleum ether:EtOAc 100:1) to afford the desired product as a colorless oil.

Step 3: To a cold (-70 °C), stirred solution of compound from Step 2 (270 g, 0.6 mol) in THF (3 L) was added MeMgBr (400 ml of 1.0 M solution in THF, 1.2 mol) dropwise. The reaction mixture was slowly warmed to -10 °C and then to rt before being quenched with a saturated aqueous solution of NH₄Cl. The organic layer was separated and the aqueous layer was extracted with EtOAc. The combined organic layers were dried, filtered and concentrated under reduced pressure to leave a residue which and purified by column chromatography on silica gel (elution with 50:1 to 10:1 petroleum ether:EtOAc) to afford the desired product as a white solid.

Step 4: To a cold (-70 °C), stirred solution of compound from Step 3 (350 g, 0.75 mol) in DCM (2 L) was added Et₃SiH (870 g, 7.5 mol) followed by a solution of BF₃.Et₂O (139 ml, 1.125 mol) dropwise. The reaction mixture was warmed up to rt before being quenched with a saturated aqueous solution of NaHCO₃. The organic layer was separated and the aqueous layer was extracted with DCM. The combined organic layers were dried, filtered and concentrated under reduced pressure to leave a residue which and purified by column chromatography on silica gel (elution with petroleum ether:EtOAc 10:1) to afford the desired product as a colorless oil.

Step 5: A mixture of compound from Step 4 (325 g, 0.72 mol) and BH₃.THF (1800 mL of 1.0 M solution in THF, 1.8 mol) in THF (500 mL) was stirred at reflux for 6 h. The reaction was cooled and EtOH (300ml) was added dropwise. The mixture was stirred at reflux for additional 2 h followed by cooling to rt. The reaction mixture was concentrated

to leave a residue which was dissolved in DCM. The organic layer was washed with a saturated aqueous solution of NaHCO₃, dried, filtered and concentrated to afford the desired product as a yellow oil.

Step 6: To a stirred solution of compound from Step 5 (438 g, 1.0 mol) in THF (2.5 L) was added TBAF (657 g, 2.5 mol) and the resulting mixture was stirred at rt overnight. The reaction mixture concentrated under reduced pressure to leave a residue which was directly purified by column chromatography on silica gel (elution with 1:1 to 0:100 petroleum ether:EtOAc) to afford the diol as a white solid.

Step 7: To a solution of diol from Step 6 (95 g, 0.46 mol) in DMF (1 L) was added imidazole (47 g, 0.69 mol) followed by TBSCl (76 g, 0.5 mol). The reaction was stirred at rt overnight. The reaction was diluted with EtOAc and the resulting layer was washed with water and brine, dried, filtered and concentrated under reduced pressure to leave a residue which was purified by column chromatography on silica gel (elution with 4:1 petroleum ether:EtOAc) to afford the desired product as a yellow oil.

Step 8: To a cold (0 °C), stirred solution of alcohol from Step 7 (75 g, 0.234 mol) in DCM (750 mL) was added Et₃N (49 ml, 0.35 mol) followed by MeSO₂Cl (32 g, 0.28 mol) dropwise. The resulting mixture was stirred at 0 °C for 2 h. The mixture was washed with water and brine, dried, filtered and concentrated to afford the desired mesylate as a brown oil which was directly used in the next step without further purification.

Step 9: A mixture of mesylate from Step 8 (93 g, 0.233 mol) and LiBr·H₂O (245 g, 2.33 mol) in DMF (1 L) was stirred at 100 °C overnight. After being cooled to room temperature the reaction was diluted with EtOAc. The resulting layer was washed with water and brine, dried, filtered and concentrated under reduced pressure to leave a residue which was purified by column chromatography on silica gel (elution with 50:1 petroleum ether:EtOAc) to afford the desired product as a colorless oil.

Step 10: A mixture of bromide from Step 9 (64 g, 0.167 mol) and NaN₃ (33 g, 0.5 mol) in DMF (600 mL) and water (50 mL) was stirred at 100 °C for 2 days. After being cooled to room temperature the reaction was diluted with EtOAc, washed with water and brine, dried, filtered and concentrated to leave a brown oil which was directly used in the next step without further purification.



Step 11: A mixture of azide from Step 10 (0.344 mol) and PPh₃ (262 g, 1.0 mol) in THF (1.5 L) and water (150 mL) was stirred at 90 °C for 2 h. The reaction was concentrated under reduced pressure to leave a residue which was diluted with water (500 ml) followed by the addition of 6N HCl until pH 1-2. The aqueous layer was extracted with EtOAc. The aqueous phase was adjusted to pH 9-10 by adding an aqueous solution of NaOH (10%). The resulting layer was extracted with DCM. The organic layer was washed with brine, dried, filtered and concentrated under reduced pressure to afford the desired amine as a brown oil.

Step 12: To a stirred solution of amine from Step 11 (156 g, 0.49 mol) in DCM (2 L) was added Et₃N (103 ml, 0.73 mol) followed by Boc₂O (128 g, 0.59 mol). After being stirred at rt for 2 h the reaction was washed with water and brine, dried, filtered and concentrated under reduced pressure to leave a residue which was purified by column chromatography on silica gel (elution with 20:1 petroleum ether:EtOAc) to afford the desired product as a white solid.

Step 13: To a stirred solution of compound from Step 12 (60 g, 0.143 mol) in THF (1 L) was added TBAF (56 g, 0.214 mol) and the resulting mixture was stirred at room temperature overnight. The mixture was concentrated and the residue was purified by column chromatography on silica gel (elution with 1:1 to 1:5 Petroleum ether:EtOAc) to afford the desired product as a white solid.

Step 14: To a stirred solution of compound from Step 13 (39 g, 0.127 mol) in MeOH (800 mL) was added 10% wet Pd/C (4 g). The flask was evacuated and back-filled with H₂ (x2). The resulting mixture was then stirred at 30 °C for 24 h. The reaction was filtrated and the filtrate was concentrated to obtain the desired amine as a gray solid.

^aThe amines used to prepare **Examples 36** and **37** are shown below. The Boc group was cleaved during step 2.



LCMS Methods

A – Mobile Phase: A: water (0.1%TFA): B: ACN(0.1%TFA); Gradient: 90:10 to 2:98 (A:B) over 1.5 min, 2:98 (A:B) for 1.5 min; Flow rate: 1.5 ml/min; Column: Waters Extera (2.1 x 20 mm)

B – Mobile Phase: A: water (0.05%TFA): B: ACN(0.05%TFA); Gradient: 90:10 to 5:95 (A:B) over 1.2 min, 5:95 (A:B) for 1.2 min; Flow rate: 1.0 ml/min; Column: Agilent Zorbax SB-C18 (3.0 x 50 mm)

C – Gradient 5:95 to 100:0 MeCN (0.1 % NH₄OH): water (0.1 % NH₄OH) over 1.4 min; Waters Acquity UPLC, column: Waters BEH-C18, 1.7um, 2.1 X 50mm
1mL/min flow

D - Mobile Phase: A: water(0.01%TFA) B: ACN(0.01%TFA); Gradient: 5%-95% B in 1.5min; Flow Rate: 1.8ml/min; Column: Sunfire C18, 4.6 X 50mm

Biological Assays

The data presented for the 5mM and Km ATP LanthaScreen™ Assay represents mean IC₅₀ values based on several test results and may have reasonable deviations depending on the specific conditions and reagents used. Reagents for the LRRK2 5mM and Km ATP LanthaScreen™ Assay were purchased from Life Technologies Corporation.

LRRK2 5mM ATP LanthaScreen™ Assay

- a) 400nl of a 1:2.15 serial dilution of test compound(98μM top assay concentration) is spotted via Labcyte Echo to certain wells in a 384 well black, untreated plate. Control wells contain 400nl of either DMSO or 400nl of a known inhibitor in DMSO.
- b) 10μl of a 2.5nM LRRK2(G2019S mutation, GST-LRRK2(amino acids 970-2527)) enzyme solution in 1x assay buffer(50mM Tris pH 8.5, 10mM MgCl₂, 0.01% Brij-35, 1.0mM EGTA, 2mM DTT, 0.05mM NaVO₄) is added to all wells.
- c) A 30 minute room temperature incubation is followed by addition of 10μl of 800nM fluorescein labeled LRRKtide peptide substrate and 10mM ATP solution in 1x assay buffer to all wells.
- d) After a 35 minute room temperature incubation, 20μl of TR-FRET Dilution Buffer(Invitrogen PV3756B) containing 4nM Tb-labeled anti-phospho LRRKtide antibody and 20mM EDTA is added to all wells.
- e) Plates are incubated at room temperature for 1 hour and read on an Envision™ multi-mode plate reader with LanthaScreen™ settings. Results are analysed using Assay Data Analyzer.

LRRK2 Km ATP LanthaScreen™ Assay

- a) 400nl of a 1:2.15 serial dilution of test compound(98μM top assay concentration) is spotted via Labcyte Echo to certain wells in a 384 well black, untreated plate. Control wells contain 400nl of either DMSO or 400nl of a known inhibitor in DMSO.

- b) 10 μ l of a 2.5nM LRRK2(G2019S mutation, GST-LRRK2(amino acids 970-2527)) enzyme solution in 1x assay buffer(50mM Tris pH 8.5, 10mM MgCl₂, 0.01% Brij-35, 1mM EGTA, 2mM DTT, 0.05mM NaVO₄) is added to all wells.
- c) A 30 minute room temperature incubation is followed by addition of 10 μ l of 800nM fluorescein labeled LRRKtide peptide substrate and 186 μ M ATP solution in 1x assay buffer to all wells.
- d) After a 60 minute room temperature incubation, 20 μ l of TR-FRET Dilution Buffer(Invitrogen PV3756B) containing 4nM Tb-labeled anti-phospho LRRKtide antibody and 20mM EDTA is added to all wells.
- e) Plates are incubated at room temperature for 1 hour and read on an Envision™ multi-mode plate reader with LanthaScreen™ settings. Results are analysed using Assay Data Analyzer.

Table A: 5mM ATP LanthaScreen™ Assay Data of representative compounds

In the table below, representative examples are provided with their respective IC₅₀ in the 5 mM ATP LanthaScreen™ Assay. Preferred compounds have an IC₅₀ less than 1 μ M in the 5 mM ATP LanthaScreen Assay.

Example	IC ₅₀ (nM)
9	225
23	297
26	300

Table B: Kinase selectivity of representative compounds

Kinase selectivity was performed using Z'-LYTE™ or Adapta® assay platforms available from Life Technologies Corporation. Values in **Table B** are percent inhibition in the presence of 1 micromolar of the indicated example.


Kinase	Assay Platform	Ex. 29	Ex. 28	Ex. 30	Ex. 27
AURKB (Aurora B)	A	11	11	6	15
BRAF V599E	A	18	19	5	29
CDK1/cyclin B	A	26	25	9	33
CHEK2 (CHK2)	A	2	4	8	14
CLK2	A	53	42	49	46
DYRK1A	A	4	2	3	6

Kinase	Assay Platform	Ex. 29	Ex. 28	Ex. 30	Ex. 27
IRAK1	B	8	-5	-17	18
JAK3	A	46	49	37	56
MAPK1 (ERK2)	A	18	23	10	23
MAPK8 (JNK1)	A	54	58	39	61

A – Z-LYTE™; B – Adapta®

IN THE CLAIMS:

1. A compound of the formula:



wherein X^1 , X^2 and X^3 are each independently selected from the group consisting of a bond or CR^eR^f ;

Y is O, CR^aR^b or NR^c ;

Z is O, CR^aR^b or NR^c ;

R^1 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, hydroxyl, NR^eR^d , OR^5 and $(C=O)OR^5$;

R^2 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) R^5 ,
- d) R^7 ,
- e) OR^5 , and
- f) NR^eR^d ;

R^3 is selected from the group consisting of:

- a) hydrogen,

- b) C₁₋₆ alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR⁵ and NR^cR^d,
- c) C₃₋₈ cycloalkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR⁵ and NR^cR^d,
- d) heterocyclyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, oxo, R⁵, OR⁵ and NR^cR^d,
- e) heteroaryl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, oxo, R⁵, OR⁵ and NR^cR^d,
- f) C₄₋₈ cycloalkenyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OR⁵ and NR^cR^d,
- g) (C=O)R⁷,
- h) (C=O)R⁵,
- i) S(O)_mR⁵, and
- j) S(O)_mR⁷;

or R² and R³ can be taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:

- a) halo,
- b) oxo,
- c) cyano,
- d) OR⁵,
- e) NR^cR^d,
- f) SO₃H,
- g) S(O)_mR⁵,
- h) S(O)_mR⁷,
- i) R⁵,

- j) R^6 ,
- k) R^7 ,
- l) $(C=O)R^5$,
- m) $(C=O)OR^5$,
- n) $(C=O)R^7$, and
- o) $(C=O)NR^cR^d$;

R^4 is selected from the group consisting of hydrogen, halo, cyano, OR^5 , aryl, heteroaryl, C_{3-8} cycloalkyl, C_{4-8} cycloalkenyl, C_{4-8} heterocycloalkyl and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, cyano, OC_{1-3} alkyl, NR^cR^d and hydroxyl;

R^5 is selected from the group consisting of hydrogen and C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) hydroxyl,
- c) OC_{1-6} alkyl,
- d) NR^cR^d ,
- e) $(C=O)NR^cR^d$,
- f) $S(O)_m$,
- g) $S(O)_mR^8$,
- h) $S(O)_mR^7$,
- i) R^7 , and
- j) OR^7 ;

R^6 is C_{1-6} alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo and hydroxyl;

or R⁵ and R⁶ can be taken together with the atoms to which they are attached to form a 4 to 8 membered heterocyclic, 3 to 8 membered carbocyclic, aryl or heteroaryl ring, wherein said heterocyclic and heteroaryl rings may contain from one to three heteroatoms selected from N, O and S, wherein said heterocyclic, carbocyclic, aryl and heteroaryl rings are optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) oxo,
- c) cyano,
- d) hydroxyl,
- e) C₁₋₃ alkyl, which is optionally substituted with one to three halo,
- f) C₃₋₈ cycloalkyl,
- g) OC₁₋₃ alkyl, which is optionally substituted with one to three halo, and
- h) OC₃₋₈ cycloalkyl;

R⁷ is selected from the group consisting of C₄₋₈ heterocyclyl, C₃₋₈ cycloalkyl, C₄₋₈ cycloalkenyl, aryl or heteroaryl, wherein said heterocyclyl, cycloalkyl, cycloalkenyl, aryl and heteroaryl groups are optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) hydroxyl,
- d) oxo,
- e) C₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d,
- f) OC₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl NR^cR^d and aryl,
- g) C₃₋₈ cycloalkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d,

h) aryl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl, S(O)_mNR^cR^d, C(O)NR^cR^d and NR^cR^d,

i) heteroaryl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl, S(O)_mNR^cR^d, C(O)NR^cR^d and NR^cR^d,

j) heterocyclyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, OC₁₋₃ alkyl and NR^cR^d,

k) C₄₋₈ cycloalkenyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo, cyano, heterocyclyl, OC₁₋₃ alkyl and NR^cR^d;

R⁸ is hydrogen or C₁₋₆ alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

- a) halo,
- b) cyano,
- c) hydroxyl,
- d) OC₁₋₃ alkyl, which is optionally substituted with one to four substituents independently selected from the group consisting of hydroxyl, halo and NR^cR^d, and
- e) C₃₋₈ cycloalkyl;

R^a is selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R^b is selected from the group consisting of hydrogen and C₁₋₆ alkyl;

R^c is selected from the group consisting of:

- a) hydrogen and
- b) C₁₋₃ alkyl, which is optionally substituted with one to three substituents independently selected from the group consisting of halo, hydroxyl, cyano, heteroaryl, aryl, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, OC₁₋₆ alkyl, and C₃₋₈ cycloalkyl;

R^d is selected from the group consisting of:

- a) hydrogen,
- b) C₃₋₈ cycloalkyl,
- c) C₃₋₆ heterocyclyl,
- d) C₁₋₃ alkyl,
- e) (C=O)C₁₋₃ alkyl,
- f) aryl, and
- g) heteroaryl;

wherein said cycloalkyl, heterocyclyl, alkyl, aryl and heteroaryl groups are each optionally substituted with one to three substituents independently selected from the group consisting of halo, hydroxyl, cyano, R⁸, SO₂R⁸, OC₁₋₆ alkyl and C₃₋₈ cycloalkyl

or R^c and R^d can be taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of halo, cyano, hydroxyl, C₁₋₃ alkyl and OC₁₋₃ alkyl;

R^e is selected from the group consisting of hydrogen and C₁₋₃ alkyl;

R^f is selected from the group consisting of hydrogen and C₁₋₃ alkyl;

m is an integer from zero to two,
or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1 wherein X¹, X² and X³ are each a bond, or a pharmaceutically acceptable salt thereof.

3. The compound of Claim 2 wherein Y is O or CR^aR^b and Z is O or CR^aR^b; or a pharmaceutically acceptable salt thereof.

4. The compound of Claim 3 wherein R¹ is selected from the group consisting of: hydrogen and C₁₋₃ alkyl, or a pharmaceutically acceptable salt thereof.
5. The compound of Claim 4 wherein R⁴ is selected from the group consisting of: hydrogen and halo, or a pharmaceutically acceptable salt thereof.
6. The compound of Claim 5 wherein R² and R³ can be taken together with the atoms to which they are attached to form a 3 to 8 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:
- a) halo,
 - b) oxo,
 - c) cyano,
 - d) OR⁵,
 - e) NR^cR^d,
 - f) SO₃H,
 - g) S(O)_mR⁵,
 - h) S(O)_mR⁷,
 - i) R⁵,
 - j) R⁶,
 - k) R⁷,
 - l) (C=O)R⁵,
 - m) (C=O)OR⁵,
 - n) (C=O)R⁷, and
 - o) (C=O)NR^cR^d;

or a pharmaceutically acceptable salt thereof.


7. The compound of Claim 6 wherein R² and R³ can be taken together with the atoms to which they are attached to form a 6 membered heterocyclic ring, wherein said ring may contain from one to three heteroatoms selected from N, O and S, wherein the sulfur









is optionally oxidized to the sulfone or sulfoxide, and which ring is optionally substituted with one to four substituents each independently selected from the group consisting of:









- a) halo,
- b) oxo,
- c) OR^5 ,
- d) NR^cR^d ,
- e) $S(O)_mR^5$,
- f) $S(O)_mR^7$,
- g) R^5 ,
- h) R^6 ,
- i) R^7 ,
- j) $(C=O)R^5$,
- k) $(C=O)OR^5$, and
- l) $(C=O)R^7$.











or a pharmaceutically acceptable salt thereof.


8. A compound selected from











or a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition comprising a compound of any of claims 1 to 8, and a pharmaceutically acceptable carrier.

10. Use of a pharmaceutical composition of any of Claims 1 to 9, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of Parkinson's Disease.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 14/18895

A. CLASSIFICATION OF SUBJECT MATTER
IPC(8) - A61K 31/415; C07D 231/56 (2014.01)
USPC - 514/405; 548/361.1
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC(8): A61K 31/415; C07D 231/56 (2014.01)
USPC: 514/405; 548/361.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC: 514/337-339

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
PatBase, Google Scholar, PubWEST
indazole, pyridine, LRRK2, Leucine-rich repeat kinase, inhibit, modulate, Parkinson's

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2012/038743 A1 (CHAN et al.) 29 March 2012 (29.03.2012) pg 44, ln 26-28; pg 73, Example 190, Example 192	1-9
Y	US 2012/0329780 A1 (THORMANN et al.) 27 December 2012 (27.12.2012) para [0035], [0331], Example #36; para [1440]	1-9
Y	US 2006/0079564 A1 (JANSEN et al.) 13 April 2006 (13.04.2006) pg 177, Example 1335; pg 181, Example 1351, para [1042]	1-9

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:
 "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search
04 May 2014 (04.05.2014)

Date of mailing of the international search report
02 JUN 2014

Name and mailing address of the ISA/US
Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450
Facsimile No. 571-273-3201

Authorized officer:
Lee W. Young
PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 14/18895

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 10
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.