

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2019277560 B2**

(54) Title
Dosing regimen for the treatment of PI3K related disorders

(51) International Patent Classification(s)
A61K 31/519 (2006.01) **A61P 35/00** (2006.01)

(21) Application No: **2019277560** (22) Date of Filing: **2019.05.30**

(87) WIPO No: **WO19/232188**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/773,612	2018.11.30	US
62/714,448	2018.08.03	US
62/679,556	2018.06.01	US

(43) Publication Date: **2019.12.05**

(44) Accepted Journal Date: **2025.04.24**

(71) Applicant(s)
Incyte Corporation

(72) Inventor(s)
YELESWARAM, Krishnaswamy;ASSAD, Albert;CHEN, Xuejun

(74) Agent / Attorney
Houlihan Intellectual Property Pty Ltd, 832 High St, Kew East, VIC, 3102, AU

(56) Related Art
FORERO-TORRES ANDRES ET AL: "Results from a Phase 1/2 Study of INCB050465, a Highly Selective and Highly Potent PI3K delta Inhibitor, in Patients with Relapsed or Refractory B-Cell Malignancies", Blood (2017),130(1), pages 410 WO 2016/138363 A1



(51) International Patent Classification:

A61K 31/519 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2019/034613

(22) International Filing Date:

30 May 2019 (30.05.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/679,556	01 June 2018 (01.06.2018)	US
62/714,448	03 August 2018 (03.08.2018)	US
62/773,612	30 November 2018 (30.11.2018)	US

(71) Applicant: INCYTE CORPORATION [US/US]; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(72) Inventors: YELESWARAM, Krishnaswamy; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). ASSAD, Albert; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US). CHEN, Xuejun; 1801 Augustine Cut-Off, Wilmington, Delaware 19803 (US).

(74) Agent: MANAS, Michael et al.; Fish & Richardson, P.O. Box 1022, Minneapolis, Minnesota 55440 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: DOSING REGIMEN FOR THE TREATMENT OF PI3K RELATED DISORDERS




Figure 1A

(57) Abstract: The present application provides methods of treating PI3K δ related disorders using pyrazolopyrimidine derivatives.

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*

Dosing Regimen for the Treatment of PI3K Related Disorders

TECHNICAL FIELD

The present application provides methods of treating PI3K δ related disorders using pyrazolopyrimidine derivatives.

5

BACKGROUND

Constitutive signaling through the B-cell receptor (BCR) plays a critical role in the pathogenesis of human B-cell malignancies (Gururajan M. et al, *The Journal of Immunology*, 2006;176(10):5715-5719) and leads to downstream activation of class I phosphatidylinositol 3-kinases (PI3Ks) (Benschop R.J., Cambier J.C., *Current Opinion in Immunology*, 1999;11(2):143-151; So L., Fruman D.A. *The Biochemical Journal*, 2012;442(3):465-481). Class I PI3Ks are heterodimeric lipid kinases composed of a regulatory (p85 or p101) and a catalytic (p110) subunit (Chalhoub N. et al, *Annual review of pathology*, 2009;4:127-150). Each of the 4 tissue-specific p110 subunit isoforms (Class IA: α , β , and δ ; Class IB: γ) confers unique physiologic functions on the corresponding PI3K isoforms, including insulin signaling and angiogenesis (PI3K α) (Foukas L.C. et al, *Nature*, 2006;441:366; Knight Z.A. et al, *Cell*, 2006;125(4):733-747; and Graupera M. et al, *Nature*, 2008;453:662), platelet function and thrombosis (PI3K β) (Jackson S.P. et al, *Nature Medicine*, 2005;11:507), and white blood cell function and inflammation (PI3K γ) (Hirsch E. et al, *Thrombosis and Haemostasis*, 2006;95(1):29-35). The PI3K δ isoform functions as a critical node in signaling networks that regulate B-cell growth and survival, and its aberrant activation is a key event in malignant transformation of B-cells (Kang S. et al, *Proceedings of the National Academy of Sciences of the United States of America*, 2005;102(3):802-807; Puri K.D., Gold M.R., *Frontiers in Immunology*, 2012;3:256). Substantial interconnectivity exists between BCR and PI3K δ -mediated signaling networks and other networks important for regulating B-cell survival and proliferation, including the Janus kinase–signal transducer and activator of transcription (JAK/STAT) pathway (Lam L.T., et al, *Blood*, 2008;111(7):3701-3713; Rui L. et al, *Proceedings of the National Academy of Sciences*, 2016;113(46):E7260-E7267), suggesting potential additive or synergistic therapeutic effects in B-cell malignancies.

Prognosis for patients with relapsed or refractory B-cell non-Hodgkin lymphoma (NHL) is generally poor. For example, the 5-year overall survival (OS) rate for patients with relapsed follicular lymphoma (FL), the most common indolent NHL subtype, is only 50% (Casulo C. et al, *Journal of Clinical Oncology*, 2015;33(23):2516-2522). Prognosis is worse for patients with relapsed aggressive NHL subtypes, with a median survival of 3.6 and 4.4 months among patients with relapsed diffuse large B-cell lymphoma (DLBCL) who had failed first-line and second-line salvage regimens,

respectively (Van Den Neste E. et al, *Bone Marrow Transplant*, 2016;51(1):51-57). Current guidelines for the treatment for relapsed B-cell NHL differ according to subtype and include chemo- or radio-immunotherapy, targeted therapies with small-molecule kinase inhibitors, or immunomodulatory therapies (NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). B-cell Lymphomas. Version 7.2017; 2017; Tilly H. et al, *Annals of Oncology*, 2015;26(suppl_5):v116-v125; Dreyling M. et al, *Annals of Oncology*, 2016;27(suppl_5):v83-v90; Dreyling M. et al, *Annals of Oncology*, 2017;28(suppl_4):iv62-iv71). In addition to systemic therapy, autologous or allogeneic stem cell transplant (SCT) is often used to treat patients with relapsed B-cell NHL and allogeneic SCT is considered the only curative option for these patients (Wudhikarn K. et al

5 *Biology of Blood and Marrow Transplantation*, 2011;17(10):1497-1504; Avivi I., et al, *British Journal of Haematology*, 2009;147(5):719-728; Klyuchnikov E., et al, *Bone Marrow Transplant*, 2014;49(1):1-7; Hamadani M. et al, *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2009;15(5):547-553).

The differential tissue distribution of the PI3K isoforms factors in their distinct biological functions. Genetic ablation of either PI3K α or PI3K β results in embryonic lethality, indicating that PI3K α and PI3K β have essential and non-redundant functions, at least during development (Vanhaesebroeck, et al., 2005). In contrast, mice which lack PI3K γ and PI3K δ are viable, fertile and have a normal life span although they show an altered immune system. PI3K γ deficiency leads to impaired recruitment of macrophages and neutrophils to sites of inflammation as well as impaired T

15 cell activation (Sasaki, et al., *Science*, 2000, 287(5455):1040-6). PI3K δ -mutant mice have specific defects in B cell signaling that lead to impaired B cell development and reduced antibody responses after antigen stimulation (Clayton, et al., *J Exp Med*. 2002, 196(6):753-63; Jou, et al., *Mol Cell Biol*. 2002, 22(24):8580-91; Okkenhaug, et al., *Science*, 2002, 297(5583):1031-4).

The phenotypes of the PI3K γ and PI3K δ -mutant mice suggest that these enzymes may play a

25 role in inflammation and other immune-based diseases and this is borne out in preclinical models. PI3K γ -mutant mice are largely protected from disease in mouse models of rheumatoid arthritis (RA) and asthma (Camps, et al., *Nat Med*. 2005, 11(9):936-43; Thomas, et al., *Eur J Immunol*. 2005, 35(4):1283-91). In addition, treatment of wild-type mice with a selective inhibitor of PI3K γ was shown to reduce glomerulonephritis and prolong survival in the MRL-lpr model of systemic lupus

30 nephritis (SLE) and to suppress joint inflammation and damage in models of RA (Barber, et al., *Nat Med*. 2005, 11(9):933-5; Camps, et al., 2005). Similarly, both PI3K δ -mutant mice and wild-type mice treated with a selective inhibitor of PI3K δ have been shown to have attenuated allergic airway inflammation and hyper-responsiveness in a mouse model of asthma (Ali, et al., *Nature*. 2004, 431(7011):1007-11; Lee, et al., *FASEB J*. 2006, 20(3):455-65) and to have attenuated disease in a

35 model of RA (Randis, et al., *Eur. J. Immunol.*, 2008, 38(5):1215-24).

B cell proliferation has shown to play a major role in the development of inflammatory autoimmune diseases (Puri, *Frontiers in Immunology* (2012), 3(256), 1-16; Walsh, *Kidney International* (2007) 72, 676–682). For example, B cells support T-cell autoreactivity, an important component of inflammatory autoimmune diseases. Once activated and matured, B cells can traffic to sites of inflammation and recruit inflammatory cells or differentiate to plasmablasts. Thus, activity of B-cells can be affected by targeting B-cell stimulatory cytokines, B-cell surface receptors, or via B-cell depletion. Rituximab—an IgG1 κ mouse/human chimeric monoclonal antibody directed against the B-cell surface receptor CD20—has been shown to deplete CD20+ B cells. Use of rituximab has been shown to have efficacy in treating idiopathic thrombocytopenic purpura, autoimmune hemolytic anemia, or vasculitis. For example, treatment with rituximab resulted in remission of the disease in patients suffering from anti-neutrophil cytoplasm antibody associated (ANCA) systemic vasculitis (AASV) with demonstrated peripheral B-cell depletion (Walsh, 2007; Lovric, *Nephrol Dial Transplant* (2009) 24: 179–185). Similarly, a complete response was reported in one-third to two-thirds of patients having mixed cryoglobulinemia vasculitis after treatment with rituximab, including patients who presented with a severe form of vasculitis that was resistant or intolerant to other treatments (Cacoub, *Ann Rheum Dis* 2008;67:283–287). Similarly, rituximab has been shown to have efficacy in treating patients with idiopathic thrombocytopenic purpura (or immune thrombocytopenic purpura) (Garvey, *British Journal of Haematology*, (2008) 141, 149–169; Godeau, *Blood* (2008), 112(4), 999-1004; Medeo, *European Journal of Haematology*, (2008) 81, 165–169) and autoimmune hemolytic anemia (Garvey, *British Journal of Haematology*, (2008) 141, 149–169).

PI3K δ signaling has been tied to B cell survival, migration, and activation (Puri, *Frontiers in Immunology*, 2012, 3(256), 1-16, at pages 1-5; and Clayton, *J Exp Med*, 2002, 196(6):753-63). For example, PI3K δ is required for antigen-dependent B-cell activation driven by B cell receptor. By blocking B-cell adhesion, survival, activation, and proliferation, PI3K δ inhibition can impair the ability of B cells to activate T cells, preventing their activation and reducing secretion of autoantibodies and pro-inflammatory cytokines. Hence, by their ability to inhibit B cell activation, PI3K δ inhibitors would be expected to treat B cell mediated diseases that were treatable by similar methods such as B cell depletion by rituximab. Indeed, PI3K δ inhibitors have been shown to be useful mouse models of various autoimmune diseases that are also treatable by rituximab such as arthritis (Puri (2012)). Further, innate-like B cells, which are linked to autoimmunity are sensitive to PI3K δ activity, as MZ and B-1 cells are nearly absent in mice lacking the p110 δ gene (Puri (2012)). PI3K δ inhibitors can reduce trafficking of and activation of MZ and B-1 cells, which are implicated in autoimmune diseases.

In addition to their potential role in inflammatory diseases, all four class I PI3K isoforms may play a role in cancer. The gene encoding p110 α is mutated frequently in common cancers, including breast, prostate, colon and endometrial (Samuels, et al., *Science*, 2004, 304(5670):554; Samuels, et

al., *Curr Opin Oncol.* 2006, 18(1):77-82). Eighty percent of these mutations are represented by one of three amino acid substitutions in the helical or kinase domains of the enzyme and lead to a significant upregulation of kinase activity resulting in oncogenic transformation in cell culture and in animal models (Kang, et al., *Proc Natl Acad Sci U S A.* 2005, 102(3):802-7; Bader, et al., *Proc Natl Acad Sci U S A.* 2006, 103(5):1475-9). No such mutations have been identified in the other PI3K isoforms although there is evidence that they can contribute to the development and progression of malignancies. Consistent overexpression of PI3K δ is observed in acute myeloblastic leukemia (Sujobert, et al., *Blood*, 2005, 106(3):1063-6) and inhibitors of PI3K δ can prevent the growth of leukemic cells (Billottet, et al., *Oncogene*. 2006, 25(50):6648-59). Elevated expression of PI3K γ is seen in chronic myeloid leukemia (Hickey, et al., *J Biol Chem.* 2006, 281(5):2441-50). Alterations in expression of PI3K β , PI3K γ and PI3K δ have also been observed in cancers of the brain, colon and bladder (Benistant, et al., *Oncogene*, 2000, 19(44):5083-90; Mizoguchi, et al., *Brain Pathol.* 2004, 14(4):372-7; Knobbe, et al., *Neuropathol Appl Neurobiol.* 2005, 31(5):486-90). Further, these isoforms have all been shown to be oncogenic in cell culture (Kang, et al., 2006).

The introduction of rituximab improved treatment outcomes for patients with B-cell NHL (Chao M.P., *Cancer Management and Research.* 2013;5:251-269; Nandagopal L., Mehta A., *Expert Review of Hematology*, 2017;10(3):259-273). For patients who experience rituximab resistance and/or relapsed or refractory disease, the PI3K inhibitor class has shown promise, but clinical use has been limited by toxicities. For example, the first-in-class PI3K δ -selective inhibitor, idelalisib, has been associated with considerable toxicity, including hepatotoxicity (Coutré S.E., et al, *Leukemia & Lymphoma*, 2015;56(10):2779-2786; Gopal A., Graf S., *Expert opinion on pharmacotherapy*, 2016;17(2):265-274), as well as increased risk of opportunistic infection (Lampson B.L. et al, *Blood*, 2016;128(2):195-203), and requires frequent monitoring. Likewise, treatments targeting other PI3K isoforms have corresponding risks (e.g., hyperglycemia and hypertension) (Burris H.A., et al, *Journal of Clinical Oncology*, 2014;32(15_suppl):2513-2513; ALIQOPA™. (copanlisib) for injection, for intravenous use. U.S. Whippany, NJ: Bayer HealthCare Pharmaceuticals Inc.; 2017; Patnaik A. et al, *Annals of Oncology*, 2016;27(10):1928-1940; Dreyling M. et al, *Journal of Clinical Oncology*, 2017;35(35):3898-3905). Therefore, an unmet need remains for effective therapies with improved safety profiles in this difficult-to-treat patient population.

SUMMARY OF THE INVENTION

The present application provides, *inter alia*, a method of treating a disease in a patient, wherein said disease is associated with abnormal expression or activity of PI3K δ kinase, the method comprising:

i) administering to the patient an inhibitor of PI3K δ in a first dosage that is about 3 mg/day to about 50 mg/day for a first period of time which is about 2 weeks to about 12 weeks; and

ii) administering to the patient a second dosage of the inhibitor of PI3K δ which is less than the first dosage administered at the end of the first time period and which is:

- (a) about 2.5 mg/day or less; or
- (b) about 50 mg/week or less;

5 and wherein the second dosage is administered for a second period of time which occurs after the first period of time.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

10

DESCRIPTION OF DRAWINGS

Figures 1A-1B show PK of (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt (*i.e.*, Compound 1C) at steady state. Figure 1A: PK of Compound 1C dosing. Figure 1B: Dose-response relationship of
15 Compound 1C AUC. AUC, area under the curve; AUC_{0-T}, area under the plasma concentration-time curve; IC₅₀, half maximal inhibitory concentration; IC₉₀, 90% maximal inhibitory concentration; PK, pharmacokinetics; QD, once daily.

Figure 2 shows monotherapy best percent change from baseline in target lesion size. *Data for 3 patients with DLBCL, 2 with MZL and 1 with MCL are not shown due to non-measurable
20 disease at baseline or no valid post baseline target lesion measurements. †Best percentage change from baseline in target lesion size >100%. IB, 3 patients with MCL had received ibrutinib before the study, of whom a best overall response of CR or PR was achieved by 2 patients. DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma.

25 Figure 3 shows duration of Compound 1C monotherapy treatment. AE, adverse event; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; PD, progressive disease; QW, once weekly.

Figure 4 shows Mean percent inhibition of phosphorylated AKT at steady state. Compound 1C was dosed once daily and levels of phosphorylated AKT (pAKT; Ser473) were measured by flow
30 cytometry in SU-DHL cells added to patient whole blood after Compound 1C treatment on day 15 of cycle 1.

Figures 5A-5B shows ALT (Figure 5A) and AST (Figure 5B) over time in patients receiving Compound 1C monotherapy. ALT, alanine aminotransferase; AST, aspartate aminotransferase; QD, once daily; ULN, upper limit of normal.

Figure 6 shows Compound 1C monotherapy time to response among NHL subtypes of interest (DLBCL, FL, MCL, MZL). DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma.

Figure 7 shows representative positron emission tomography (PET) images of a patient who achieved complete response with Compound 1C monotherapy. Patient was <65 years of age, with mantle cell lymphoma with no bone marrow involvement (target lesion, 98 mm × 58 mm) and had received 1 prior treatment (R-HyperCVAD). Treated with Compound 1C, 30 mg once daily; achieved complete response at week 9.

DETAILED DESCRIPTION

The present application provides methods of treating a disease in a patient, wherein said disease is associated with abnormal expression or activity of PI3K δ kinase, the method comprising:

- i) administering to the patient an inhibitor of PI3K δ in a first dosage that is about 3 mg/day to about 50 mg/day for a first period of time which is about 2 weeks to about 12 weeks; and
- ii) administering to the patient a second dosage of the inhibitor of PI3K δ which is less than the first dosage administered at the end of the first time period and which is:

- (a) about 2.5 mg/day or less; or
- (b) about 50 mg/week or less;

and wherein the second dosage is administered for a second period of time which occurs after the first period of time.

The present application further provides methods of treating a disease in a patient, wherein said disease is associated with abnormal expression or activity of PI3K δ kinase, the method comprising:

- i) administering to the patient an inhibitor of PI3K δ in a first dosage that is about 3 mg/day to about 50 mg/day for a first period of time which is about 2 weeks to about 12 weeks; and
- ii) administering to the patient a second dosage of the inhibitor of PI3K δ which is less than the first dosage administered at the end of the first time period and which is about 7.5 mg/day or less;

and wherein the second dosage is administered for a second period of time which occurs after the first period of time.

As used herein, the term “first period of time” refer to a period of time of administration of the first dosage of the inhibitor of PI3K δ to the patient.

In some embodiments, the first period of time begins at the time of initial administration of the first dosage of the inhibitor of PI3K δ to the patient.

As used herein, the term “second period of time” refer to a period of time of administration of the second dosage of the inhibitor of PI3K δ to the patient.

In some embodiments, the second period of time begins at the time of initial administration of the second dosage of the inhibitor of PI3K δ to the patient.

In some embodiments, the time of initial administration of the second dosage of the inhibitor of PI3K δ to the patient (*i.e.*, the start of the second period of time) is within 2 weeks of the end of the first period of time, for example, within 1 week, within 6 days, within 5 days, within 4 days, within 3 days, within 2 days, or within 1 days of the end of the first period of time.

In some embodiments, the second period of time is continued for any amount of time, for example, until the treatment is terminated.

In some embodiments, each of the first dosages is administered as a single, once daily dosage.

10 In some embodiments, each of the first dosages is administered as a single, once daily oral dosage.

In some embodiments, the first dosage is about 5 mg/day to about 50 mg/day, for example, about 5 mg/day to about 40 mg/day, about 5 mg/day to about 30 mg/day, about 5 mg/day to about 20 mg/day, about 5 mg/day to about 10 mg/day, about 10 mg/day to about 40 mg/day, about 10 mg/day to about 30 mg/day, about 10 mg/day to about 20 mg/day, about 20 mg/day to about 40 mg/day, about 20 mg/day to about 30 mg/day, or about 20 mg/day to about 40 mg/day.

In some embodiments, the first dosage is about 50 mg/day, about 45 mg/day, about 40 mg/day, about 30 mg/day, about 20 mg/day, about 15 mg/day, about 10 mg/day, or about 5 mg/day.

In some embodiments, the first dosage is about 18 mg/day to about 22 mg/day.

20 In some embodiments, the first dosage is about 18 mg/day to about 22 mg/day and is administered as a single, once daily (*i.e.*, QD) dosage.

In some embodiments, the first dosage is about 18 mg/day to about 22 mg/day and is administered as a single, once daily (*i.e.*, QD) oral dosage.

In some embodiments, the first dosage is about 20 mg/day.

25 In some embodiments, the first dosage is about 20 mg/day and is administered as a single, once daily (*i.e.*, QD) dosage.

In some embodiments, the first dosage is about 20 mg/day and is administered as a single, once daily (*i.e.*, QD) oral dosage.

In some embodiments, the first dosage is about 10 mg/day.

30 In some embodiments, the first dosage is about 10 mg/day and is administered as a single, once daily (*i.e.*, QD) dosage.

In some embodiments, the first dosage is about 10 mg/day and is administered as a single, once daily (*i.e.*, QD) oral dosage.

In some embodiments, the first period of time is about 3 weeks to about 11 weeks, for example, about 8 weeks to about 12 weeks, about 4 weeks to about 10 weeks, about 5 weeks to about 9 weeks, or about 8 weeks to about 9 weeks. In some embodiments, the first period of time is about 2

weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 7 weeks, about 8 weeks, about 9 weeks, about 10 weeks, about 11 weeks, or about 12 weeks.

In some embodiments, the first period of time is about 8 weeks to about 9 weeks.

In some embodiments, the first period of time is about 8 weeks.

5 In some embodiments, the first period of time is about 9 weeks.

In some embodiments, the first dosage is reduced during the first period of time.

In some embodiments, the first dosage is reduced from the about 3 mg/day to about 50 mg/day to about 3 mg/day to about 30 mg/day, about 3 mg/day to about 20 mg/day, about 3 mg/day to about 15 mg/day, about 3 mg/day to about 10 mg/day, about 3 mg/day to about 7 mg/day, or about 3 mg/day to about 5 mg/day.

In some embodiments, the first dosage is reduced from the about 3 mg/day to about 50 mg/day to about 3 mg/day to about 7 mg/day during the first period of time.

In some embodiments, the first dosage is reduced from about 50 mg/day to about 30 mg/day, about 20 mg/day, about 15 mg/day, about 10 mg/day, about 5 mg/day, about 2 mg/day, or about 1 mg/day, during the first period of time.

In some embodiments, the first dosage is reduced from about 20 mg/day to about 15 mg/day, about 10 mg/day, about 5 mg/day, about 2 mg/day, or about 1 mg/day, during the first period of time.

In some embodiments, each of the second dosages is administered as a single, once daily (QD) dosage or as a single, once weekly (QW) dosage.

20 In some embodiments, each of the second dosages is administered as a single, once daily (QD) dosage.

In some embodiments, each of the second dosages is administered as a single, once weekly (QW) dosage.

In some embodiments, each of the second dosages is administered as a single, once daily (QD) oral dosage or as a single, once weekly (QW) oral dosage.

In some embodiments, each of the second dosages is administered as a single, once daily (QD) oral dosage.

In some embodiments, each of the second dosages is administered as a single, once weekly (QW) oral dosage.

30 In some embodiments, the second dosage is about 5.0 mg/day or less, for example, about 5.0 mg/day, about 4.0 mg/day, about 3.0 mg/day, about 2.5 mg/day, about 2.0 mg/day, about 1.75 mg/day about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day.

In some embodiments, the second dosage is about 2.5 mg/day to about 7.5 mg/day.

In some embodiments, the second dosage is about 3.0 mg/day to about 7.0 mg/day.

35 In some embodiments, the second dosage is about 4.0 mg/day to about 6.0 mg/day.

In some embodiments, the second dosage is about 5.0 mg/day.

In some embodiments, the second dosage is about 4.0 mg/day to about 6.0 mg/day and is administered as a single, once daily (QD) dosage.

In some embodiments, the second dosage is about 5.0 mg/day and is administered as a single, once daily (QD) dosage.

5 In some embodiments, the second dosage is about 4.0 mg/day to about 6.0 mg/day and is administered as a single, once daily (QD) oral dosage.

In some embodiments, the second dosage is about 5.0 mg/day and is administered as a single, once daily (QD) oral dosage.

10 In some embodiments, the second dosage is reduced from about 5.0 mg/day or less during the second period of time.

In some embodiments, the second dosage is reduced from about 5.0 mg/day or less to about 2.5 mg/day or less, 2.0 mg/day or less, about 1.75 mg/day, or less about 1.5 mg/day or less, about 1.25 mg/day or less, or about 1.0 mg/day or less during the second period of time.

15 In some embodiments, the second dosage is reduced from about 5.0 mg/day to about 2.5 mg/day, 2.0 mg/day, about 1.75 mg/day, about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day during the second period of time

In some embodiments, the second dosage is reduced from the about 5.0 mg/day to about 0.75 mg/day to about 1.25 mg/day during the second period of time.

20 In some embodiments, the second dosage is reduced from about 5.0 mg/day to about 1.25 mg/day during the second period of time.

In some embodiments, the second dosage is reduced from about 5.0 mg/day to about 2.0 mg/day, about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day.

In some embodiments, the second dosage is reduced from about 5.0 mg/day to about 1.0 mg/day.

25 In some embodiments, the second dosage is about 2.5 mg/day or less, for example, about 2.5 mg/day, about 2.0 mg/day, about 1.75 mg/day about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day.

30 In some embodiments, the second dosage is about 1.0 mg/day to about 2.5 mg/day, for example, about 1.0 mg/day to about 2.0 mg/day, about 1.0 mg/day to about 1.5 mg/day, or about 1.0 mg/day to about 1.25 mg/day.

In some embodiments, the second dosage is about 2 mg/day to about 2.5 mg/day.

In some embodiments, the second dosage is about 2.5 mg/day.

In some embodiments, the second dosage is about 2 mg/day to about 2.5 mg/day and is administered as a single, once daily (QD) dosage.

35 In some embodiments, the second dosage is about 2.5 mg/day and is administered as a single, once daily (QD) dosage.

In some embodiments, the second dosage is about 2 mg/day to about 2.5 mg/day and is administered as a single, once daily (QD) oral dosage.

In some embodiments, the second dosage is about 2.5 mg/day and is administered as a single, once daily (QD) oral dosage.

5 In some embodiments, the second dosage is reduced from about 2.5 mg/day or less during the second period of time.

In some embodiments, the second dosage is reduced from about 2.5 mg/day or less to about 2.0 mg/day or less, about 1.75 mg/day or less, about 1.5 mg/day or less, about 1.25 mg/day or less, or about 1.0 mg/day or less during the second period of time.

10 In some embodiments, the second dosage is reduced from about 2.5 mg/day to about 2.0 mg/day, about 1.75 mg/day, about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day during the second period of time.

In some embodiments, the second dosage is reduced from about 2.5 mg/day to about 1.25 mg/day during the second period of time.

15 In some embodiments, the second dosage is reduced from about 2.5 mg/day to about 2.0 mg/day, about 1.5 mg/day, about 1.25 mg/day, or about 1.0 mg/day.

In some embodiments, the second dosage is reduced from about 2.5 mg/day to about 1.0 mg/day.

20 In some embodiments, the reduced dosage of the second dosage is administered as a single, once daily (QD) dosage.

In some embodiments, the reduced dosage of the second dosage is administered as a single, once daily (QD) oral dosage.

25 In some embodiments, the second dosage is about 50 mg/week or less, for example, about 40 mg/week or less, about 30 mg/week or less, about 20 mg/week or less, about 10 mg/week or less, about 5 mg/week or less, about 2 mg/week or less, or about 1 mg/week or less.

30 In some embodiments, the second dosage is about 5 mg/week to about 50 mg/week, for example, about 5 mg/week to about 40 mg/week, about 5 mg/week to about 30 mg/week, about 5 mg/week to about 20 mg/week, about 5 mg/week to about 10 mg/week, about 10 mg/week to about 40 mg/week, about 10 mg/week to about 30 mg/week, about 10 mg/week to about 20 mg/week, about 20 mg/week to about 40 mg/week, about 20 mg/week to about 30 mg/week, or about 30 mg/week to about 40 mg/week.

In some embodiments, the second dosage is about 18 mg/week to about 20 mg/week.

In some embodiments, the second dosage is about 20 mg/week.

35 In some embodiments, the second dosage is about 18 mg/week to about 20 mg/week and is administered as a single, once weekly (QW) dosage.

In some embodiments, the second dosage is about 20 mg/week and is administered as a single, once weekly (QW) dosage.

In some embodiments, the second dosage is reduced from about 50 mg/week or less during the second period of time.

5 In some embodiments, the second dosage is reduced from about 50 mg/week or less to about 40 mg/week or less, about 30 mg/week or less, about 20 mg/week or less, about 10 mg/week or less, about 5 mg/week or less, about 2 mg/week or less, or about 1 mg/week or less during the second period of time.

10 In some embodiments, the second dosage is reduced from about 5 mg/week to about 50 mg/week, to about 5 mg/week to about 40 mg/week, about 5 mg/week to about 30 mg/week, about 5 mg/week to about 20 mg/week, about 5 mg/week to about 10 mg/week, about 10 mg/week to about 40 mg/week, about 10 mg/week to about 30 mg/week, about 10 mg/week to about 20 mg/week, about 20 mg/week to about 40 mg/week, about 20 mg/week to about 30 mg/week, or about 30 mg/week to about 40 mg/week during the second period of time.

15 In some embodiments, the second dosage is reduced from about 50 mg/week, to about 40 mg/week, about 30 mg/week, about 20 mg/week, about 10 mg/week, or about 5 mg/week during the second period of time.

In some embodiments, the second dosage is reduced from about 20 mg/week to about 10 mg/week during the second period of time.

20 In some embodiments, the second dosage is reduced from about 20 mg/week to about 10 mg/week during the second period of time.

In some embodiments, the second dosage is reduced from about 10 mg/week to about 5 mg/week during the second period of time.

25 In some embodiments, the reduced dosage of the second dosage is administered as a single, once weekly (QW) dosage.

In some embodiments, the patient has been identified as exhibiting one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs). As used herein, the severity of TEAEs described herein were assessed using Common Terminology Criteria for Adverse Events version 4.03, the disclosure of which is incorporated herein by reference in its entirety.

30 In some embodiments, the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs) during the first period of time.

In some embodiments, the first dosage is reduced from about 3 mg/day to about 50 mg/day to about 3 mg/day to about 30 mg/day, about 3 mg/day to about 20 mg/day, about 3 mg/day to about 15 mg/day, about 3 mg/day to about 10 mg/day, about 3 mg/day to about 7 mg/day, or about 3 mg/day to

35

about 5 mg/day when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the first dosage is reduced from about 50 mg/day to about 30 mg/day, about 20 mg/day, about 15 mg/day, about 10 mg/day, about 5 mg/day, about 2 mg/day, or about 1 mg/day, during the first period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the first dosage is reduced from about 20 mg/day to about 15 mg/day, about 10 mg/day, about 5 mg/day, about 2 mg/day, or about 1 mg/day, during the first period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the first dosage is reduced from about 20 mg/day to about 10 mg/day when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the first dosage is reduced from about 20 mg/day to about 5 mg/day when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the first dosage is reduced from about 10 mg/day to about 5 mg/day when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the methods provided herein further comprise stopping administration of the first dosage when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs) during the second period of time.

In some embodiments, the second dosage is reduced from about 3 mg/day to about 50 mg/day to about 3 mg/day to about 30 mg/day, about 3 mg/day to about 20 mg/day, about 3 mg/day to about 15 mg/day, about 3 mg/day to about 10 mg/day, about 3 mg/day to about 7 mg/day, or about 3 mg/day to about 5 mg/day during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the second dosage is reduced from about 50 mg/week, to about 40 mg/week, about 30 mg/week, about 20 mg/week, about 10 mg/week, about 5 mg/week, about 2 mg/week or about 1 mg/week during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the second dosage is reduced from about 50 mg/week or less to about 40 mg/week or less, about 30 mg/week or less, about 20 mg/week or less, about 10 mg/week or less, about 5 mg/week or less, about 2 mg/week or less, or about 1 mg/week or less during the second period of time during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the second dosage is reduced from about 20 mg/week to about 10 mg/week or less during the second period of time during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the second dosage is reduced from about 20 mg/week to about 5 mg/week or less during the second period of time during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the second dosage is reduced from about 10 mg/week to about 5 mg/week or less during the second period of time during the second period of time when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).

In some embodiments, the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis, nausea, fatigue, rash, neutropenia, pyrexia, hypotension, sepsis, respiratory failure, pneumonitis, pneumonia, hypertension, hyperglycemia, abdominal pain, bronchitis, dehydration, thrombocytopenia, cough, vomiting, decreased appetite, increased lacrimation, oral herpes, tachycardia, spinal cord compression, intractable pain, elevated alkaline phosphatase, elevated transaminase, hyperlipidemia, hypercalcemia, dizziness, alopecia, constipation, fluid overload, headache, hypokalemia, night sweats, encephalopathy, atrial flutter, atrial fibrillation, and dyspnea.

In some embodiments, the symptoms associated with the one or more treatment-emergent adverse events are symptoms associated with one or more of diarrhea/colitis, nausea, fatigue, rash, neutropenia, pyrexia, hypotension, sepsis, respiratory failure, pneumonitis, pneumonia, hypertension, hyperglycemia, abdominal pain, bronchitis, dehydration, thrombocytopenia, cough, vomiting, decreased appetite, increased lacrimation, oral herpes, tachycardia, spinal cord compression, intractable pain, elevated alkaline phosphatase, elevated transaminase, hyperlipidemia, hypercalcemia, dizziness, alopecia, constipation, fluid overload, headache, hypokalemia, night sweats, encephalopathy, atrial flutter, atrial fibrillation, and dyspnea.

In some embodiments, the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis, nausea, fatigue, rash, cough, vomiting, dizziness, pyrexia, hypokalemia, abdominal pain, constipation, decreased appetite, night sweats, pruritus, back pain, chills, leukopenia, neutropenia, lymphopenia, thrombocytopenia, and anemia.

In some embodiments, the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis and rash.

In some embodiments, the diarrhea/colitis comprises one or more of diarrhea, colitis, enterocolitis, gastrointestinal inflammation, colitis microscopic, and cytomegalovirus colitis.

5 In some embodiments, the rash comprises one or more of dermatitis exfoliative, rash, rash erythematous, rash macular, rash maculopapular, rash pruritic, exfoliative rash, rash generalized, rash popular, and rash pustular.

In some embodiments, the neutropenia comprises febrile neutropenia.

10 In some embodiments, the elevated transaminase comprises elevated alanine transaminase (ALT), aspartate transaminase (AST), or a combination thereof.

As used herein, the term “contacting” refers to the bringing together of indicated moieties in an *in vitro* system or an *in vivo* system. For example, “contacting” a PI3K with a PI3K δ inhibitor compound described herein includes the administration of a compound described herein to an individual or patient, such as a human, having a PI3K, as well as, for example, introducing a
15 compound described herein into a sample containing a cellular or purified preparation containing the PI3K.

In some embodiments, the inhibitor of PI3K δ is a compound of US Patent Publ. No. US 2011/0015212, filed June 28, 2010, US Patent Publ. No. 2013/0059835, filed August 31, 2013, US Patent Publ. No. 2011/0183985, filed December 17, 2010, US Patent Publ. No. 2012/0157430, filed
20 December 19, 2011, US Patent Publ. No. 2014/0249132, filed February 28, 2014, or US Patent Publ. No. 2016/0257689, filed February 26, 2016, each of which is incorporated herein by reference in its entirety.

In some embodiments, the inhibitor of PI3K δ is prepared by the methods in US Patent Publ. No. US 2011/0015212, filed June 28, 2010, US Patent Publ. No. 2013/0059835, filed August 31,
25 2013, US Patent Publ. No. 2011/0183985, filed December 17, 2010, or US Patent Publ. No. 2012/0157430, filed December 19, 2011, US Patent Publ. No. 2014/0249132, filed February 28, 2014, or US Patent Publ. No. 2016/0257689, filed February 26, 2016, each of which is incorporated herein by reference in its entirety.

In some embodiments, the inhibitor of PI3K δ is selected from:

30 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one; and

5-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one;

or a pharmaceutically acceptable salt thereof.

In some embodiments, the inhibitor of PI3K δ is 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof.

In some embodiments, the inhibitor of PI3K δ is selected from:

5 (S)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one;

(R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one;

10 (S)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one; and

(R)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one;

or a pharmaceutically acceptable salt thereof.

15 In some embodiments, the inhibitor of PI3K δ is a pharmaceutically acceptable salt of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one.

In some embodiments, the inhibitor of PI3K δ is selected from a pharmaceutically acceptable salt of:

20 (S)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one;

(R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one;

(S)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one; and

25 (R)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one.

In some embodiments, the inhibitor of PI3K δ is a pharmaceutically acceptable salt of (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one.

30 In some embodiments, the inhibitor of PI3K δ is (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt. In some embodiments, the salt is a 1:1 stoichiometric ratio of (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one to hydrochloric acid.

35 In some embodiments, the salt is crystalline (see *e.g.*, US Patent Publ. No. 2016/0257689, filed February 26, 2016 for methods and processes for preparing (R)-4-(3-((S)-1-(4-amino-3-methyl-

1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt. US Patent Publ. No. 2016/0257689 is incorporated herein by reference, in its entirety.).

In some embodiments, the hydrochloric acid salt of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one has: at least one, two, three, four or five XRPD peaks, in terms of 2-theta, selected from about 11.3°, about 16.4°, about 21.0°, about 23.0°, about 28.1°, about 31.2°, and about 32.8°; at least two XRPD peaks, in terms of 2-theta, selected from about 11.3°, about 16.4°, about 21.0°, about 23.0°, about 28.1°, about 31.2°, and about 32.8°; at least three XRPD peaks, in terms of 2-theta, selected from about 11.3°, about 16.4°, about 21.0°, about 23.0°, about 28.1°, about 31.2°, and about 32.8°. In some embodiments, the hydrochloric acid salt of the compound of Formula I has at least four XRPD peaks, in terms of 2-theta, selected from about 11.3°, about 16.4°, about 21.0°, about 23.0°, about 28.1°, about 31.2°, and about 32.8°. In some embodiments, the hydrochloric acid salt of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one has all of the listed XRPD peaks, in terms of 2-theta, at about 11.3°, about 16.4°, about 21.0°, about 23.0°, about 28.1°, about 31.2°, and about 32.8°. In some embodiments, the hydrochloric acid salt of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one has a DSC thermogram having an endothermic peak at about 207 °C.

In some embodiments, the salts and compounds described herein are substantially isolated. By “substantially isolated” is meant that the salt or compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the salts described herein. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the salts described herein, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

In some embodiments, the inhibitor of PI3K δ is 5-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one, or a pharmaceutically acceptable salt thereof.

In some embodiments, the inhibitor of PI3K δ is selected from:

(R)-5-{3-[(R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one;

(R)-5-{3-[(S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one;

(S)-5-{3-[(S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one; and

(S)-5-{3-[(R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one;

or a pharmaceutically acceptable salt thereof.

Compounds described herein, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes. In some embodiments, the compounds can be prepared as described in US Patent Nos. 9,199,982 and 9,932,341, each of which is incorporated herein by reference in its entirety.

The reactions for preparing compounds described herein can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, *e.g.*, temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

Preparation of compounds described herein can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., Wiley & Sons, Inc., New York (1999), which is incorporated herein by reference in its entirety.

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (*e.g.*, ¹H or ¹³C), infrared spectroscopy, spectrophotometry (*e.g.*, UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC), liquid chromatography-mass spectroscopy (LCMS), or thin layer chromatography (TLC). Compounds can be purified by those skilled in the art by a variety of methods, including high performance liquid chromatography (HPLC) ("*Preparative LC-MS Purification: Improved Compound Specific Method Optimization*" Karl F. Blom, Brian Glass, Richard Sparks, Andrew P. Combs *J. Combi. Chem.* **2004**, 6(6), 874-883, which is incorporated herein by reference in its entirety) and normal phase silica chromatography.

The compounds described herein can modulate activity of one or more of various kinases including, for example, phosphoinositide 3-kinases (PI3Ks). The term "modulate" is meant to refer to an ability to increase or decrease the activity of one or more members of the PI3K family. Accordingly, the compounds described herein can be used in methods of modulating a PI3K by contacting the PI3K with any one or more of the compounds or compositions described herein. In some embodiments, compounds of the present invention can act as inhibitors of one or more PI3Ks.

In further embodiments, the compounds described herein can be used to modulate activity of a PI3K in an individual in need of modulation of the receptor by administering a modulating amount of a compound described herein, or a pharmaceutically acceptable salt thereof. In some embodiments, modulating is inhibiting. Given that cancer cell growth and survival is impacted by multiple signaling pathways, the present invention is useful for treating disease states characterized by drug resistant kinase mutants. In addition, different kinase inhibitors, exhibiting different preferences in the kinases which they modulate the activities of, may be used in combination. This approach could prove highly efficient in treating disease states by targeting multiple signaling pathways, reduce the likelihood of drug-resistance arising in a cell, and reduce the toxicity of treatments for disease.

The inhibitors of PI3K δ described herein can be selective. By “selective” is meant that the compound binds to or inhibits a kinase with greater affinity or potency, respectively, compared to at least one other kinase. In some embodiments, the compounds described herein are selective inhibitors of PI3K δ over PI3K γ , PI3K α and/or PI3K β . In some embodiments, the compounds described herein are selective inhibitors of PI3K δ (*e.g.*, over PI3K α , PI3K β and PI3K γ). In some embodiments, selectivity can be at least about 2-fold, 5-fold, 10-fold, at least about 20-fold, at least about 50-fold, at least about 100-fold, at least about 200-fold, at least about 500-fold or at least about 1000-fold. Selectivity can be measured by methods routine in the art. In some embodiments, selectivity can be tested at the K_m ATP concentration of each enzyme. In some embodiments, the selectivity of compounds described herein can be determined by cellular assays associated with particular PI3K kinase activity.

Kinases to which the present compounds bind and/or modulate (*e.g.*, inhibit) include any member of the PI3K family. In some embodiments, the PI3K is PI3K α , PI3K β , PI3K γ , or PI3K δ . In some embodiments, the PI3K is PI3K γ or PI3K δ . In some embodiments, the PI3K is PI3K γ . In some embodiments, the PI3K is PI3K δ . In some embodiments, the PI3K includes a mutation. A mutation can be a replacement of one amino acid for another, or a deletion of one or more amino acids. In such embodiments, the mutation can be present in the kinase domain of the PI3K.

In some embodiments of the present methods, more than one compound described herein is used to inhibit the activity of one kinase (*e.g.*, PI3K γ or PI3K δ).

In some embodiments of the present methods, more than one compound described herein is used to inhibit more than one kinase, such as at least two kinases (*e.g.*, PI3K γ and PI3K δ).

In some embodiments of the present methods, one or more of the compounds is used in combination with another kinase inhibitor to inhibit the activity of one kinase (*e.g.*, PI3K γ or PI3K δ).

In some embodiments of the present methods, one or more of the compounds is used in combination with another kinase inhibitor to inhibit the activities of more than one kinase (*e.g.*, PI3K γ or PI3K δ), such as at least two kinases.

Another aspect of the present invention pertains to methods of treating a kinase (such as PI3K δ)-associated disease or disorder in an individual (*e.g.*, patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of one or more compounds of the present invention or a pharmaceutical composition thereof. A PI3K-associated disease can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the PI3K, including overexpression and/or abnormal activity levels. In some embodiments, the disease can be linked to Akt (protein kinase B), mammalian target of rapamycin (mTOR), or phosphoinositide-dependent kinase 1 (PDK1). In some embodiments, the mTOR-related disease can be inflammation, atherosclerosis, psoriasis, restenosis, benign prostatic hypertrophy, bone disorders, pancreatitis, angiogenesis, diabetic retinopathy, atherosclerosis, arthritis, immunological disorders, kidney disease, or cancer. A PI3K-associated disease can also include any disease, disorder or condition that can be prevented, ameliorated, or cured by modulating PI3K activity. In some embodiments, the disease is characterized by the abnormal activity of PI3K. In some embodiments, the disease is characterized by mutant PI3K. In such embodiments, the mutation can be present in the kinase domain of the PI3K.

Examples of PI3K-associated diseases include immune-based diseases involving the system including, for example, rheumatoid arthritis, allergy, asthma, glomerulonephritis, lupus, or inflammation related to any of the above.

Further examples of PI3K-associated diseases include cancers such as breast, prostate, colon, endometrial, brain, bladder, skin, uterus, ovary, lung, pancreatic, renal, gastric, or hematological cancer.

In some embodiments, the hematological cancer is acute myeloblastic leukemia (AML) or chronic myeloid leukemia (CML), or B cell lymphoma. In some embodiments, the B cell lymphoma is diffuse large B cell lymphoma.

Further examples of PI3K-associated diseases include lung diseases such as acute lung injury (ALI) and adult respiratory distress syndrome (ARDS).

Further examples of PI3K-associated diseases include osteoarthritis, restenosis, atherosclerosis, bone disorders, arthritis, diabetic retinopathy, psoriasis, benign prostatic hypertrophy, inflammation, angiogenesis, pancreatitis, kidney disease, inflammatory bowel disease, myasthenia gravis, multiple sclerosis, or Sjögren's syndrome, and the like.

Further examples of PI3K-associated diseases include autoimmune hemolytic anemia, polycythemia vera, and pemphigus.

In some embodiments, the disease is non-Hodgkin lymphoma (NHL). In some embodiments, the non-Hodgkin lymphoma (NHL) is relapsed or refractory NHL or recurrent follicular NHL.

In some embodiments, the disease is an aggressive lymphoma (*e.g.*, germinal center B cell-like (GCB) or activated B cell-like (ABC)).

In some embodiments, the disease is selected from diffuse large B-cell lymphoma, chronic lymphocytic leukemia (CLL), Non-Hodgkin lymphoma, hairy cell leukemia, Mantle cell lymphoma, small lymphocytic lymphoma, follicular lymphoma, lymphoplasmacytic lymphoma, extranodal marginal zone lymphoma, Hodgkin's lymphoma, Burkitt lymphoma, Waldenstrom's

5 macroglobulinemia, prolymphocytic leukemia, acute lymphoblastic leukemia, myelofibrosis, mucosa-associated lymphatic tissue (MALT) lymphoma, mediastinal (thymic) large B-cell lymphoma, lymphomatoid granulomatosis, splenic marginal zone lymphoma, primary effusion lymphoma, intravascular large B-cell lymphoma, plasma cell leukemia, extramedullary plasmacytoma, smouldering myeloma (aka asymptomatic myeloma), monoclonal gammopathy of undetermined

10 significance (MGUS), activated B-cell like (ABC) diffuse large B cell lymphoma (ABC-DLBCL), and germinal center B cell (GCB) diffuse large B cell lymphoma (GCB-DLBCL).

In some embodiments, the disease is myelofibrosis. In some embodiments, the myelofibrosis is selected from post-essential thrombocythemia myelofibrosis (PET-MF), primary myelofibrosis (PMF), and post-polycythemia vera myelofibrosis (PPV-MF). In some embodiments, the

15 myelofibrosis is post-essential thrombocythemia myelofibrosis (PET-MF). In some embodiments, the myelofibrosis is primary myelofibrosis (PMF). In some embodiments, the myelofibrosis is post-polycythemia vera myelofibrosis (PPV-MF).

In some embodiments, the disease is acute myeloid leukemia. In some embodiments, the disease is Burkitt lymphoma. In some embodiments, the disease is activated B-cell like (ABC) diffuse

20 large B cell lymphoma (ABC-DLBCL) or germinal center B cell (GCB) diffuse large B cell lymphoma (GCB-DLBCL).

In some embodiments, the disease is selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL) follicular lymphoma (FL) Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia

25 (WM).

In some embodiments, the marginal zone lymphoma (MZL) is selected from extranodal MZL, nodal MZL, splenic MZL, and unknown MZL subtype.

In some embodiments, the Hodgkin lymphoma (HL) is selected from classic Hodgkin lymphoma (HL) and nodular lymphocytic-predominant HL.

30 In some embodiments, the diffuse large B-cell lymphoma is selected from activated B-cell like (ABC) diffuse large B cell lymphoma (ABC-DLBCL) and germinal center B cell (GCB) diffuse large B cell lymphoma (GCB-DLBCL).

In some embodiments, the disease is selected from autoimmune hemolytic anemia, polycythemia vera, and pemphigus.

35 In some embodiments, the present application further provides a method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-

cell lymphoma (DLBCL) follicular lymphoma (FL) Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day to about 50 mg/day for a first period of time which is about 8 weeks to about 9 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 2.5 mg/day or less for a second period of time which occurs after the first period of time.

In some embodiments, the present application further provides a method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL) follicular lymphoma (FL) Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day to about 50 mg/day for a first period of time which is about 8 weeks to about 9 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 20 mg/week to about 50 mg/week for a second period of time which occurs after the first period of time.

In some embodiments, the present application further provides a method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL) follicular lymphoma (FL) Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day for a first period of time which is about 8 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 2.5 mg/day for a second period of time which occurs after the first period of time.

In some embodiments, the present application further provides a method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL) follicular lymphoma (FL) Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day for a first period of time which is about 8 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 20 mg/week for a second period of time which occurs after the first period of time.

As used herein, the term “individual” or “patient,” used interchangeably, refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

As used herein, the phrase “therapeutically effective amount” refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response that is being sought in a tissue, system, animal, individual or human by a researcher, veterinarian, medical doctor or other clinician.

As used herein, the term “treating” or “treatment” refers to one or more of (1) preventing the disease; for example, preventing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease; (2) inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology); and (3) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology) such as decreasing the severity of disease. In some embodiments, the term “treating” or “treatment” refers to inhibiting or ameliorating the disease.

As used herein, “QD” is taken to mean a dosage administered to the subject once-daily. “QW” is taken to mean a dosage administered to the subject once-weekly. “Q2W” is taken to mean a dosage administered to the subject once, every other week. “Q3W” is taken to mean a dosage administered to the subject once, every three weeks. “Q4W” is taken to mean a dosage administered to the subject once, every four weeks.

As used herein, “about” when referring to a measurable value such as an amount, a dosage, a temporal duration, and the like, is meant to encompass variations of $\pm 10\%$. In certain embodiments, “about” can include variations of $\pm 5\%$, $\pm 1\%$, or $\pm 0.1\%$ from the specified value and any variations there between, as such variations are appropriate to perform the disclosed methods.

5

Combination Therapies

I. Immune-checkpoint therapies

In some embodiments, the inhibitors of PI3K provided herein can be used in combination with one or more immune checkpoint inhibitors for the treatment of cancer as described herein. In one
 10 embodiment, the combination with one or more immune checkpoint inhibitors as described herein can be used for the treatment of a B-cell malignancy described here. Compounds of the present disclosure can be used in combination with one or more immune checkpoint inhibitors. Exemplary immune checkpoint inhibitors include inhibitors against immune checkpoint molecules such as CD20, CD28, CD40, CD122, CD96, CD73, CD47, GITR, CSF1R, JAK, PI3K δ , PI3K γ , TAM, arginase, HPK1,
 15 CD137 (also known as 4-1BB), ICOS, B7-H3, B7-H4, BTLA, CTLA-4, LAG3, TIM3, VISTA, TIGIT, PD-1, PD-L1 and PD-L2. In some embodiments, the immune checkpoint molecule is a stimulatory checkpoint molecule selected from CD27, CD28, CD40, ICOS, OX40, GITR and CD137. In some embodiments, the immune checkpoint molecule is an inhibitory checkpoint molecule selected from A2AR, B7-H3, B7-H4, BTLA, CTLA-4, IDO, KIR, LAG3, PD-1, TIM3, TIGIT, and VISTA. In
 20 some embodiments, the compounds of the disclosure provided herein can be used in combination with one or more agents selected from KIR inhibitors, TIGIT inhibitors, LAIR1 inhibitors, CD160 inhibitors, 2B4 inhibitors and TGFR beta inhibitors.

In some embodiments, the PI3K inhibitors provided herein can be used in combination with one or more agonists of immune checkpoint molecules, e.g., OX40, CD27, OX40, GITR, and CD137
 25 (also known as 4-1BB).

In some embodiments, the inhibitor of an immune checkpoint molecule is anti-PD1 antibody, anti-PD-L1 antibody, or anti-CTLA-4 antibody.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1, e.g., an anti-PD-1 monoclonal antibody. In some embodiments, the anti-PD-1 monoclonal antibody
 30 is nivolumab, pembrolizumab (also known as MK-3475), durvalumab (Imfinzi®), pidilizumab, SHR-1210, PDR001, MGA012, PDR001, AB122, or AMP-224. In some embodiments, the anti-PD-1 monoclonal antibody is nivolumab or pembrolizumab. In some embodiments, the anti-PD1 antibody is pembrolizumab. In some embodiments, the anti-PD-1 monoclonal antibody is MGA012. In some
 35 embodiments, the anti-PD1 antibody is SHR-1210. Other anti-cancer agent(s) include antibody therapeutics such as 4-1BB (e.g. urelumab, utomilumab).

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-L1, e.g., an anti-PD-L1 monoclonal antibody. In some embodiments, the anti-PD-L1 monoclonal antibody is BMS-935559, MEDI4736, MPDL3280A (also known as RG7446), or MSB0010718C. In some embodiments, the anti-PD-L1 monoclonal antibody is MPDL3280A or MEDI4736.

5 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of PD-1 and PD-L1, e.g., an anti-PD-1/PD-L1 monoclonal antibody. In some embodiments, the anti-PD-1/PD-L1 is MCLA-136.

In some embodiments, the inhibitor is MCLA-145.

10 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CTLA-4, e.g., an anti-CTLA-4 antibody. In some embodiments, the anti-CTLA-4 antibody is ipilimumab, tremelimumab, AGEN1884, or CP-675,206.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of LAG3, e.g., an anti-LAG3 antibody. In some embodiments, the anti-LAG3 antibody is BMS-986016, LAG525, or INCAGN2385.

15 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of TIM3, e.g., an anti-TIM3 antibody. In some embodiments, the anti-TIM3 antibody is INCAGN2390, MBG453, or TSR-022.

In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of GITR, e.g., an anti-GITR antibody. In some embodiments, the anti-GITR antibody is TRX518, MK-4166, INCAGN1876, MK-1248, AMG228, BMS-986156, GWN323, or MEDI1873.

20 In some embodiments, the inhibitor of an immune checkpoint molecule is an agonist of OX40, e.g., OX40 agonist antibody or OX40L fusion protein. In some embodiments, the anti-OX40 antibody is MEDI0562, MOXR-0916, PF-04518600, GSK3174998, or BMS-986178. In some embodiments, the OX40L fusion protein is MEDI6383.

25 In some embodiments, the inhibitor of an immune checkpoint molecule is an inhibitor of CD20, e.g., an anti-CD20 antibody. In some embodiments, the anti-CD20 antibody is obinutuzumab or rituximab.

The compounds of the present disclosure can be used in combination with bispecific antibodies. In some embodiments, one of the domains of the bispecific antibody targets PD-1, PD-L1, CTLA-4, GITR, OX40, TIM3, LAG3, CD137, ICOS, CD3 or TGF β receptor.

30 In some embodiments, the compounds of the disclosure can be used in combination with one or more metabolic enzyme inhibitors. In some embodiments, the metabolic enzyme inhibitor is an inhibitor of IDO1, TDO, or arginase. Examples of IDO1 inhibitors include epacadostat, NLG919, BMS-986205, PF-06840003, IOM2983, RG-70099 and LY338196.

As provided throughout, the additional compounds, inhibitors, agents, etc. can be combined with the present compound in a single or continuous dosage form, or they can be administered simultaneously or sequentially as separate dosage forms.

5 II. Cancer therapies

Cancer cell growth and survival can be impacted by multiple signaling pathways. Thus, it is useful to combine different enzyme/protein/receptor inhibitors, exhibiting different preferences in the targets which they modulate the activities of, to treat such conditions. Targeting more than one signaling pathway (or more than one biological molecule involved in a given signaling pathway) may
10 reduce the likelihood of drug-resistance arising in a cell population, and/or reduce the toxicity of treatment.

The compounds of the present disclosure can be used in combination with one or more other enzyme/protein/receptor inhibitors or one or more therapies for the treatment of diseases, such as cancer. Examples of diseases and indications treatable with combination therapies include those as
15 described herein.

One or more additional pharmaceutical or therapeutic agents such as, for example, chemotherapeutics, immune-oncology agents, metabolic enzyme inhibitors, chemokine receptor inhibitors, and phosphatase inhibitors, as well as targeted therapies such as Bcr-Abl, Flt-3, EGFR, HER2, JAK, c-MET, VEGFR, PDGFR, c-Kit, IGF-1R, RAF and FAK kinase inhibitors. The one or
20 more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

Example antibodies for use in combination therapy include but are not limited to Trastuzumab (e.g. anti-HER2), Ranibizumab (e.g. anti-VEGF-A), Bevacizumab (trade name Avastin, e.g. anti-VEGF, Panitumumab (e.g. anti-EGFR), Cetuximab (e.g. anti-EGFR), Rituxan (anti-CD20) and
25 antibodies directed to c-MET.

One or more of the following agents may be used in combination with the compounds of the present disclosure and are presented as a non-limiting list: a cytostatic agent, taxotere, taxol, camptostar, epothilones, 5-fluorouracil, SCH 66336, R115777, L778,123, BMS 214662, IRESSATM(gefitinib), TARCEVATM (erlotinib), antibodies to EGFR, GLEEVECTM (imatinib
30 mesylate), intron, ara-C, adriamycin, cytoxan, chlormethine, triethylenemelamine, triethylenethiophosphoramine, busulfan, 6-mercaptopurine, 6-thioguanine, fludarabine phosphate, ELOXATINTM (oxaliplatin), vindesine, mithramycin, deoxycoformycin, L-asparaginase, 17 alpha-ethinylestradiol, diethylstilbestrol, testosterone, prednisone, fluoxymesterone, methylprednisolone, methyltestosterone, prednisolone, triamcinolone, chlorotrianisene, hydroxyprogesterone,
35 aminoglutethimide, medroxyprogesteroneacetate, leuprolide, flutamide, goserelin, hydroxyurea, amsacrine, navelbene, reloxafine, droloxafine, hexamethylmelamine, avastin, HERCEPTINTM

(trastuzumab), BEXXAR™ (tositumomab), VELCADE™ (bortezomib), ZEVALIN™ (ibritumomab tiuxetan), TRISENOX™ (arsenic trioxide), XELODA™ (capecitabine), vinorelbine, porfimer, ERBITUX™ (cetuximab), aphidicolon, rituxan, Sml1, triapine, didox, trimidox, amidox, 3-AP, and MDL-101, 731.

5 The compounds of the present disclosure can further be used in combination with other methods of treating cancers, for example by chemotherapy, irradiation therapy, tumortargeted therapy, adjuvant therapy, immunotherapy or surgery. Examples of immunotherapy include cytokine treatment (e.g., interferons, GM-CSF, G-CSF, IL-2), CRS-207 immunotherapy, cancer vaccine, monoclonal antibody, adoptive T cell transfer, Toll receptor agonists, STING agonists, oncolytic virotherapy and
10 immunomodulating small molecules, including thalidomide or JAK1/2 inhibitor and the like. The compounds can be administered in combination with one or more anti-cancer drugs, such as a chemotherapeutics. Example chemotherapeutics include any of: abarelix, aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, anastrozole, arsenic trioxide, asparaginase, azacitidine, bevacizumab, bexarotene, baricitinib, bleomycin, bortezomib, bortezomib, busulfan intravenous,
15 busulfan oral, calusterone, capecitabine, carboplatin, carmustine, cetuximab, chlorambucil, cisplatin, cladribine, clofarabine, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, dalteparin sodium, dasatinib, daunorubicin, decitabine, denileukin, denileukin diftitox, dexrazoxane, docetaxel, doxorubicin, dromostanolone propionate, eculizumab, epirubicin, erlotinib, estramustine, etoposide phosphate, etoposide, exemestane, fentanyl citrate, filgrastim, floxuridine, fludarabine, fluorouracil,
20 fulvestrant, gefitinib, gemcitabine, gemtuzumab ozogamicin, goserelin acetate, histrelin acetate, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib mesylate, interferon alfa 2a, irinotecan, lapatinib ditosylate, lenalidomide, letrozole, leucovorin, leuprolide acetate, levamisole, lomustine, meclorethamine, megestrol acetate, melphalan, mercaptopurine, methotrexate, methoxsalen, mitomycin C, mitotane, mitoxantrone, nandrolone phenpropionate, nelarabine, nofetumomab,
25 olaparib, oxaliplatin, paclitaxel, pamidronate, panitumumab, pegaspargase, pegfilgrastim, pemetrexed disodium, pentostatin, pipobroman, plicamycin, procarbazine, quinacrine, rasburicase, rituximab, ruxolitinib, rucaparib, sorafenib, streptozocin, sunitinib, sunitinib maleate, tamoxifen, temozolomide, teniposide, testolactone, thalidomide, thioguanine, thiotepa, topotecan, toremifene, tositumomab, trastuzumab, tretinoin, uracil mustard, valrubicin, vinblastine, vincristine, vinorelbine, vorinostat,
30 niraparib, veliparib, talazoparib, and zoledronate.

Additional examples of chemotherapeutics include proteasome inhibitors (e.g., bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

Example Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts.

35 Example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts.

Example suitable RAF inhibitors include compounds, and their pharmaceutically acceptable salts.

Example suitable FAK inhibitors include compounds, and their pharmaceutically acceptable salts.

5 In some embodiments, the compounds of the disclosure can be used in combination with one or more other kinase inhibitors including imatinib, particularly for treating patients resistant to imatinib or other kinase inhibitors.

The compounds can be used in combination with tumor vaccines and CAR (Chimeric antigen receptor) T cell treatment as a booster for T cell activation. In some embodiments, tumor vaccines
10 include the proteins from viruses implicated in human cancers such as Human Papilloma Viruses (HPV), Hepatitis Viruses (HBV and HCV) and Kaposi's Herpes Sarcoma Virus (KHSV). In some embodiments, the compounds of the present disclosure can be used in combination with tumor specific antigen such as heat shock proteins isolated from tumor tissue itself. In some embodiments, the compounds can be combined with dendritic cells immunization to activate potent anti-tumor
15 responses.

Methods for the safe and effective administration of most of these chemotherapeutic agents are known to those skilled in the art. In addition, their administration is described in the standard literature. For example, the administration of many of the chemotherapeutic agents is described in the "Physicians' Desk Reference" (PDR, e.g., 1996 edition, Medical Economics Company, Montvale, NJ),
20 the disclosure of which is incorporated herein by reference as if set forth in its entirety.

In some embodiments, the additional therapeutic agent is selected from a JAK inhibitor (*e.g.*, a selective JAK1 inhibitor), a pan-PIM-selective kinase inhibitor, an inhibitor of epigenetic regulator, and an immunotherapeutic agent.

In some embodiments, the inhibitor of epigenetic regulator is selected from a bromo- and
25 extra-terminal domain inhibitor, and a lysine-specific histone demethylase 1A inhibitor.

In some embodiments, the additional therapeutic agent is an immunotherapeutic agent.

In some embodiments, the immunotherapeutic agent is R-ICE.

In some embodiments, the additional therapeutic agent is a JAK inhibitor.

In some embodiments, the JAK inhibitor is selective for JAK1 and JAK2 over JAK3 and
30 TYK2. In some embodiments, the JAK inhibitor is selective for JAK1 over JAK2, JAK3, and TYK2. For example, the JAK inhibitor may preferentially inhibit JAK1 over one or more of JAK2, JAK3, and TYK2. In some embodiments, the JAK inhibitor inhibits JAK1 preferentially over JAK2 (*e.g.*, have a JAK1/JAK2 IC₅₀ ratio >1). In some embodiments, the JAK inhibitor is 10-fold more selective for JAK1 over JAK2. In some embodiments, the JAK inhibitor is about 3-fold, about 5-fold, about
35 10-fold, about 15-fold, or about 20-fold more selective for JAK1 over JAK2 as calculated by





measuring IC_{50} at 1 mM ATP (see *e.g.*, U.S. Patent Application No.: 14/680,659, filed April 7, 2015, the disclosure of which is incorporated by reference herein in its entirety).

In some embodiments, the additional therapeutic agent is a JAK1/JAK2 inhibitor. In some embodiments, the additional therapeutic agent is ruxolitinib (*e.g.*, ruxolitinib phosphate).




- 5 In some embodiments, the additional therapeutic agent is a JAK inhibitor selected from the group provided in the following table. The compounds in provided in the table are selective JAK1 inhibitors (selective over JAK2, JAK3, and TYK2). The IC_{50} s obtained by the method of Assay A described in U.S. Patent Application No.: 14/680,659 at 1 mM ATP are shown in Table A.



10






Table A.

#	Prep.	Name	Structure	JAK1 IC_{50} (nM)	JAK2/JAK1
1	US 2016/000 079 Example 1	((2R,5S)-5-{2-[(1R)-1-hydroxyethyl]-1H-imidazo[4,5-d]thieno[3,2-b]pyridin-1-yl}tetrahydro-2H-pyran-2-yl)acetonitrile		++	>10
2	US 2016/000 079 Example 2	4-[3-(cyanomethyl)-3-(3',5'-dimethyl-1H,1'H-4,4'-bipyrazol-1-yl)azetidin-1-yl]-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide		+++	>10
3	US 2010/0298334 (Example 2) ^a	3-[1-(6-chloropyridin-2-yl)pyrrolidin-3-yl]-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile		+	>10
4	US 2010/0298334 (Example 13c)	3-(1-[1,3]oxazolo[5,4-b]pyridin-2-ylpyrrolidin-3-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
5	US 2011/ 0059951 (Example 12)	4-[(4-{3-cyano-2-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile		+	>10
6	US 2011/ 0059951 (Example 13)	4-[(4-{3-cyano-2-[3-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrrol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile		+	>10
7	US 2011/ 0224190 (Example 1)	{1-{1-[3-Fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10



#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
8	US 2011/ 0224190 (Example 154)	4-{3-(Cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-[4-fluoro-2-(trifluoromethyl)phenyl]piperidine-1-carboxamide		+	>10
9	US 2011/ 0224190 (Example 85)	[3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]-1-(1-{[2-(trifluoromethyl)pyrimidin-4-yl]carbonyl}piperidin-4-yl)azetidin-3-yl]acetonitrile		+	>10
10	US 2012/ 0149681 (Example 7b)	[<i>trans</i> -1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]-3-(4-{[2-(trifluoromethyl)pyrimidin-4-yl]carbonyl}piperazin-1-yl)cyclobutyl]acetonitrile		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
11	US 2012/ 0149681 (Example 157)	{ <i>trans</i> -3-(4-{[4-[(3-hydroxyazetidin-1-yl)methyl]-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile		+	>10
12	US 2012/ 0149681 (Example 161)	{ <i>trans</i> -3-(4-{[4-[(2S)-2-(hydroxymethyl)pyrrolidin-1-yl]methyl]-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
13	US 2012/ 0149681 (Example 162)	{ <i>trans</i> -3-(4-{[4-{[(2R)-2-(hydroxymethyl)pyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]cyclobutyl}acetonitrile		+	>10
14	US 2012/ 0149682 (Example 20) ^b	4-(4-{3-[(dimethylamino)methyl]-5-fluorophenoxy}piperidin-1-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]butanenitrile		+	>10
15	US 2013/ 0018034 (Example 18)	5-{3-(cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-isopropylpyrazine-2-carboxamide		+	>10
16	US 2013/ 0018034 (Example 28)	4-{3-(cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide		+	>10
17	US 2013/ 0018034 (Example 34)	5-{3-(cyanomethyl)-3-[4-(1H-pyrrolo[2,3-b]pyridin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-N-isopropylpyrazine-2-carboxamide		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
18	US 2013/ 0045963 (Example 45)	{1-(<i>cis</i> -4-{[6-(2-hydroxyethyl)-2-(trifluoromethyl)pyrimidin-4-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10
19	US 2013/ 0045963 (Example 65)	{1-(<i>cis</i> -4-{[4-[(ethylamino)methyl]-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10
20	US 2013/ 0045963 (Example 69)	{1-(<i>cis</i> -4-{[4-(1-hydroxy-1-methylethyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10
21	US 2013/ 0045963 (Example 95)	{1-(<i>cis</i> -4-{[4-{[(3R)-3-hydroxypyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10
22	US 2013/ 0045963 (Example 95)	{1-(<i>cis</i> -4-{[4-{[(3S)-3-hydroxypyrrolidin-1-yl]methyl}-6-(trifluoromethyl)pyridin-2-yl]oxy}cyclohexyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-3-yl}acetonitrile		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
23	US 2014/ 0005166 (Example 1)	{ <i>trans</i> -3-(4-{[4-({[(1 <i>S</i>)- 2-hydroxy-1- methylethyl]amino}met hyl)-6- (trifluoromethyl)pyridin- 2-yl]oxy}piperidin-1- yl)-1-[4-(7H- pyrrolo[2,3-d]pyrimidin- 4-yl)-1H-pyrazol-1- yl]cyclobutyl}acetonitril e		+	>10
24	US 2014/ 0005166 (Example 14)	{ <i>trans</i> -3-(4-{[4-({[(2 <i>R</i>)- 2- hydroxypropyl]amino}m ethyl)-6- (trifluoromethyl)pyridin- 2-yl]oxy}piperidin-1- yl)-1-[4-(7H- pyrrolo[2,3-d]pyrimidin- 4-yl)-1H-pyrazol-1- yl]cyclobutyl}acetonitril e		+	>10

#	Prep.	Name	Structure	JAK1 IC ₅₀ (nM)	JAK2/ JAK1
25	US 2014/ 0005166 (Example 15)	{ <i>trans</i> -3-(4-{[4-({[(2 <i>S</i>)-2-hydroxypropyl]amino}methyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl)-1 <i>H</i> -pyrazol-1-yl]cyclobutyl}acetonitrile		+	>10
26	US 2014/ 0005166 (Example 20)	{ <i>trans</i> -3-(4-{[4-(2-hydroxyethyl)-6-(trifluoromethyl)pyridin-2-yl]oxy}piperidin-1-yl)-1-[4-(7H-pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl)-1 <i>H</i> -pyrazol-1-yl]cyclobutyl}acetonitrile		+	>10

+ means <10 nM (see U.S. Patent Application No.: 14/680,659, Example A for assay conditions)

++ means ≤ 100 nM (see U.S. Patent Application No.: 14/680,659, Example A for assay conditions)

+++ means ≤ 300 nM (see U.S. Patent Application No.: 14/680,659, Example A for assay conditions)

^aData for enantiomer 1

5 ^bData for enantiomer 2

In some embodiments, the JAK inhibitor is {1-{1-[3-fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3-[4-(7H-pyrrolo[2,3-*d*]pyrimidin-4-yl)-1*H*-pyrazol-1-yl]azetidin-3-yl}acetonitrile (*i.e.*, itacitinib), or a pharmaceutically acceptable salt thereof.

10 In some embodiments, the JAK inhibitor is {1-{1-[3-fluoro-2-(trifluoromethyl)isonicotinoyl]piperidin-4-yl}-3-[4-(7H-pyrrolo[2,3-*d*]pyrimidin-4-yl)-1*H*-pyrazol-1-yl]azetidin-3-yl}acetonitrile adipic acid salt.

In some embodiments, the JAK inhibitor is 4-{3-(cyanomethyl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]azetidin-1-yl}-2,5-difluoro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]benzamide, or a pharmaceutically acceptable salt thereof.

In some embodiments, the JAK inhibitor is selected from (R)-3-[1-(6-chloropyridin-2-yl)pyrrolidin-3-yl]-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile, (R)-3-(1-[1,3]oxazolo[5,4-b]pyridin-2-yl)pyrrolidin-3-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile, (R)-4-[(4-{3-cyano-2-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile, (R)-4-[(4-{3-cyano-2-[3-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrrol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile, or (R)-4-(4-{3-[(dimethylamino)methyl]-5-fluorophenoxy}piperidin-1-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]butanenitrile, (S)-3-[1-(6-chloropyridin-2-yl)pyrrolidin-3-yl]-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile, (S)-3-(1-[1,3]oxazolo[5,4-b]pyridin-2-yl)pyrrolidin-3-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile, (S)-4-[(4-{3-cyano-2-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile, (S)-4-[(4-{3-cyano-2-[3-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrrol-1-yl]propyl}piperazin-1-yl)carbonyl]-3-fluorobenzonitrile, (S)-4-(4-{3-[(dimethylamino)methyl]-5-fluorophenoxy}piperidin-1-yl)-3-[4-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]butanenitrile; and pharmaceutically acceptable salts of any of the aforementioned.

In some embodiments, the compounds of Table A are prepared by the synthetic procedures described in US Patent Publ. No. 2010/0298334, filed May 21, 2010, US Patent Publ. No. 2011/0059951, filed August 31, 2010, US Patent Publ. No. 2011/0224190, filed March 9, 2011, US Patent Publ. No. 2012/0149681, filed November 18, 2011, US Patent Publ. No. 2012/0149682, filed November 18, 2011, US Patent Publ. 2013/0018034, filed June 19, 2012, US Patent Publ. No. 2013/0045963, filed August 17, 2012, and US Patent Publ. No. 2014/0005166, filed May 17, 2013, each of which is incorporated herein by reference in its entirety.

In some embodiments, the JAK inhibitor is selected from the compounds of US Patent Publ. No. 2010/0298334, filed May 21, 2010, US Patent Publ. No. 2011/0059951, filed August 31, 2010, US Patent Publ. No. 2011/0224190, filed March 9, 2011, US Patent Publ. No. 2012/0149681, filed November 18, 2011, US Patent Publ. No. 2012/0149682, filed November 18, 2011, US Patent Publ. 2013/0018034, filed June 19, 2012, US Patent Publ. No. 2013/0045963, filed August 17, 2012, and US Patent Publ. No. 2014/0005166, filed May 17, 2013, each of which is incorporated herein by reference in its entirety.

In some embodiments, the inhibitor of PI3K δ and additional therapeutic agent are administered simultaneously.

In some embodiments, the inhibitor of PI3K δ and additional therapeutic agent are administered sequentially.

Pharmaceutical Formulations

When employed as pharmaceuticals, the compounds used in the methods of the disclosure can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (*e.g.*, by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral, or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, *e.g.*, intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

This disclosure also includes pharmaceutical compositions which contain, as the active ingredient, the compound of the disclosure or a pharmaceutically acceptable salt thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). In some embodiments, the composition is suitable for topical administration. In making the compositions of the disclosure, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it can be milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, *e.g.* about 40 mesh.

The compounds of the disclosure may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types.

Finely divided (nanoparticulate) preparations of the compounds of the disclosure can be prepared by processes known in the art, *e.g.*, see International App. No. WO 2002/000196.

Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the disclosure can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

The compositions can be formulated in a unit dosage form. The term “unit dosage forms” refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present disclosure. When referring to these preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above.

The tablets or pills of the present disclosure can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

The liquid forms in which the compounds and compositions of the present disclosure can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions can be nebulized by use of inert gases. Nebulized solutions

may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face mask, tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

5 Topical formulations can contain one or more conventional carriers. In some embodiments, ointments can contain water and one or more hydrophobic carriers selected from, for example, liquid paraffin, polyoxyethylene alkyl ether, propylene glycol, white Vaseline, and the like. Carrier compositions of creams can be based on water in combination with glycerol and one or more other components, e.g. glycerinemonostearate, PEG-glycerinemonostearate and cetylstearyl alcohol. Gels
10 can be formulated using isopropyl alcohol and water, suitably in combination with other components such as, for example, glycerol, hydroxyethyl cellulose, and the like. In some embodiments, topical formulations contain at least about 0.1, at least about 0.25, at least about 0.5, at least about 1, at least about 2, or at least about 5 wt % of the compound of the disclosure. The topical formulations can be suitably packaged in tubes of, for example, 100 g which are optionally associated with instructions for
15 the treatment of the select indication, e.g., psoriasis or other skin condition.

The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at
20 least partially arrest the symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as by the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

The compositions administered to a patient can be in the form of pharmaceutical
25 compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients,
30 carriers, or stabilizers will result in the formation of pharmaceutical salts.

The therapeutic dosage of a compound of the present disclosure can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the disclosure in a pharmaceutical composition can
35 vary depending upon a number of factors including dosage, chemical characteristics (e.g., hydrophobicity), and the route of administration. For example, the compounds of the disclosure can

be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration.

The compositions of the disclosure can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed herein.

Labeled Compounds and Assay Methods

The methods of the present disclosure further includes the use of isotopically-labeled compounds of the disclosure. An “isotopically” or “radio-labeled” compound is a compound of the disclosure where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (i.e., naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present disclosure include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . For example, one or more hydrogen atoms in a compound of the present disclosure can be replaced by deuterium atoms (e.g., one or more hydrogen atoms of an alkyl group of a compound described herein can be optionally substituted with deuterium atoms, such as $-\text{CD}_3$ being substituted for $-\text{CH}_3$).

One or more constituent atoms of the compounds presented herein can be replaced or substituted with isotopes of the atoms in natural or non-natural abundance. In some embodiments, the compound includes at least one deuterium atom. In some embodiments, the compound includes two or more deuterium atoms. In some embodiments, the compound includes 1-2, 1-3, 1-4, 1-5, or 1-6 deuterium atoms. In some embodiments, all of the hydrogen atoms in a compound can be replaced or substituted by deuterium atoms.

In some embodiments, 1, 2, 3, 4, 5, 6, 7, or 8 hydrogen atoms, attached to carbon atoms of the compounds described herein, are optionally replaced by deuterium atoms.

Synthetic methods for including isotopes into organic compounds are known in the art (Deuterium Labeling in Organic Chemistry by Alan F. Thomas (New York, N.Y., Appleton-Century-Crofts, 1971; The Renaissance of H/D Exchange by Jens Atzrodt, Volker Derdau, Thorsten Fey and Jochen Zimmermann, Angew. Chem. Int. Ed. 2007, 7744-7765; The Organic Chemistry of Isotopic Labelling by James R. Hanson, Royal Society of Chemistry, 2011). Isotopically labeled compounds can be used in various studies such as NMR spectroscopy, metabolism experiments, and/or assays.

Substitution with heavier isotopes, such as deuterium, may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances. (see *e.g.*, A. Kerekes et. al. *J. Med. Chem.* 2011, 54, 201-210; R. Xu et. al. *J. Label Compd. Radiopharm.* 2015, 58,

308-312). In particular, substitution at one or more metabolism sites may afford one or more of the therapeutic advantages.

The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* PI3K labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I or ^{35}S can be useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br can be useful.

It is understood that a “radio-labeled” or “labeled compound” is a compound that has incorporated at least one radionuclide. In some embodiments, the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

The present disclosure can further include synthetic methods for incorporating radio-isotopes into compounds of the disclosure. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and an ordinary skill in the art will readily recognize the methods applicable for the compounds of disclosure.


Kits

The present disclosure also includes pharmaceutical kits useful, for example, in the treatment or prevention of PI3K-associated diseases or disorders described herein, which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the disclosure. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit. In certain embodiments, said instructions comprise the methods of the present disclosure including specified dosages or dosing regimens of the present disclosure.

The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.


EXAMPLES

Examples 1A-1D. Diastereoisomers of 4-{3-[1-(4-amino-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one




5

Step 1. 1-(5-Chloro-2-ethoxy-3-iodo-4-methylphenyl)ethanol



A solution of 1-(5-chloro-2-ethoxy-4-fluoro-3-iodophenyl)ethanone (20.0 g, 58.4 mmol; see Example 212, step 1 of U.S. Patent No.: 9,199,982) and 1,2-ethanediol (6.5 mL, 120 mmol) in toluene (190 mL) was treated with *p*-toluenesulfonic acid monohydrate (1.1 g, 5.8 mmol). The flask was fitted with a Dean-Stark trap that was filled with sieves, and refluxed for 3 h. The reaction mixture was cooled and added to ice cooled saturated sodium bicarbonate solution (250 mL) and extracted with ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated to a crude orange oil. The crude material was purified by flash column chromatography using ethyl acetate in hexanes (0% - 20%) to give the desired product (22 g, 99%). LCMS for $C_{12}H_{14}ClFIO_3$ (M+H)⁺: m/z = 387.0; Found: 386.9.

Step 2. Ethyl (2E)-3-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]acrylate




A mixture of 2-(5-chloro-2-ethoxy-4-fluoro-3-iodophenyl)-2-methyl-1,3-dioxolane (22 g, 58 mmol) (from Step 1), ethyl (2E)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)acrylate (16 mL, 70

mmol), and potassium carbonate (24 g, 170 mmol) in 1,4-dioxane (230 mL) and water (110 mL) was degassed with nitrogen for 10 min. The reaction mixture was treated with [1,1'-bis (diphenylphosphino)\ferrocene]dichloropalladium(II), complex with dichloromethane (1:1) (2.4 g, 2.9 mmol), degassed with nitrogen for another 10 min, and heated at 80 °C for 2 h. The reaction mixture was filtered through Celite and washed with ethyl acetate (300 mL). The filtrate was poured into water (400 mL). The aqueous layer was separated and extracted with additional ethyl acetate (300 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to a crude brown solid. The crude material was purified by flash column chromatography using ethyl acetate in hexanes (0% - 30%) to give the desired product (20 g, 96%).


¹H NMR (400 MHz, CDCl₃) δ 7.74 (d, *J* = 16.5 Hz, 1H), 7.56 (d, *J* = 8.6 Hz, 1H), 6.70 (dd, *J* = 16.5, 0.9 Hz, 1H), 4.26 (q, *J* = 7.1 Hz, 2H), 4.10 – 3.99 (m, 2H), 3.91 (q, *J* = 7.0 Hz, 2H), 3.87 – 3.76 (m, 2H), 1.73 (s, 3H), 1.44 (t, *J* = 7.0 Hz, 3H), 1.33 (t, *J* = 7.1 Hz, 3H). LCMS for C₁₇H₂₁ClFO₅ (M+H)⁺: *m/z* = 359.1; Found: 359.1.

Step 3. Ethyl 3-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]-4-nitrobutanoate




A solution ethyl (2*E*)-3-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]acrylate (10 g, 28 mmol) (from Step 2) in nitromethane (100 mL) was treated with 1,8-diazabicyclo[5.4.0]undec-7-ene (4.6 mL, 31 mmol) and stirred at 60 °C for 15 h. The reaction mixture was poured into water (400 mL) and extracted with ethyl acetate (2 x 300 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to a crude orange oil. The crude material was purified by flash column chromatography using ethyl acetate in hexanes (0% - 30%) to give the desired product as a mixture of enantiomers (10.4 g, 89%). ¹H NMR (400 MHz, CDCl₃) δ 7.52 (d, *J* = 9.1 Hz, 1H), 4.82 (ddd, *J* = 12.5, 7.6, 1.4 Hz, 1H), 4.68 (dd, *J* = 12.5, 7.2 Hz, 1H), 4.54 – 4.40 (m, 1H), 4.15 – 3.90 (m, 6H), 3.89 – 3.75 (m, 2H), 2.85 (ddd, *J* = 16.0, 8.6, 1.4 Hz, 1H), 2.73 (dd, *J* = 16.1, 6.2 Hz, 1H), 1.70 (s, 3H), 1.47 (t, *J* = 7.0 Hz, 3H), 1.21 (t, *J* = 7.1 Hz, 3H). LCMS for C₁₈H₂₄ClFNO₇ (M+H)⁺: *m/z* = 420.1; Found: 420.1.

Step 4. Enantiomers 4-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]pyrrolidin-2-one



A suspension of ethyl 3-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]-4-nitrobutanoate (1.0 g, 2.4 mmol) (from Step 3) in ethanol (16 mL) was warmed to dissolve the solid. The solution was cooled back to ambient temperature, degassed with nitrogen, and treated with a slurry of 2800 Raney Nickel in water (1.5 mL). The reaction mixture was degassed again with nitrogen and hydrogenated with a balloon of hydrogen for 3 h. The reaction mixture was filtered through Celite and concentrated to give the intermediate amino ester (0.93 g, 100%). The intermediate amino ester was dissolved in toluene (12 mL) and heated at 110 °C for 12 h. The reaction mixture was cooled to ambient temperature, at which point a solid precipitated from solution. This mixture was cooled to 0 °C, stirred for 30 min, filtered, washed with cold toluene, and dried to give the desired product as a mixture of enantiomers (0.61 g, 75%). LCMS for C₁₆H₂₀ClFNO₄ (M+H)⁺: m/z = 344.1; Found: 344.1. The mixture of enantiomers was separated by chiral HPLC to give the individual enantiomers as peak 1 and peak 2 (RT = 5.39 min and 7.01 min, respectively; Phenomenex Lux Cellulose C-1, 21.2 x 250 mm, 5 micron particle size, eluting with 20% ethanol in hexanes at 18 mL/min).

Step 5. Enantiomers of 4-(3-acetyl-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one




The separated enantiomers from step 4 were each processed individually to the final compounds. A solution of 4-[3-chloro-6-ethoxy-2-fluoro-5-(2-methyl-1,3-dioxolan-2-yl)phenyl]pyrrolidin-2-one (1.7 g, 5.0 mmol) (from Step 4) in methanol (17 mL) was treated with 6.0 M hydrogen chloride in water (11 mL, 69 mmol) dropwise and stirred 20 °C for 30 min. The reaction mixture was added dropwise to ice cooled saturated sodium bicarbonate solution (75 mL) and extracted with ethyl acetate (2 x 100 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated to give the desired products [from peak 1 (1.5 g,

99%); from peak 2 (1.5 g, 99%)] that were used without further purification. From peak 1: ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.84 (s, 1H), 7.70 (d, $J = 8.6$ Hz, 1H), 4.16 – 3.99 (m, 1H), 3.83 (q, $J = 7.0$ Hz, 2H), 3.65 – 3.54 (m, 1H), 3.30 – 3.23 (m, 1H), 2.55 (s, 3H), 2.33 (dd, $J = 16.8, 8.4$ Hz, 1H), 1.30 (t, $J = 7.0$ Hz, 3H). LCMS for $\text{C}_{14}\text{H}_{16}\text{ClFNO}_3$ ($\text{M}+\text{H}$) $^+$: $m/z = 300.1$; Found: 300.0. From peak 2:

5 ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.84 (s, 1H), 7.70 (d, $J = 8.6$ Hz, 1H), 4.13 – 4.00 (m, 1H), 3.87 – 3.77 (m, 2H), 3.65 – 3.55 (m, 1H), 3.31 – 3.23 (m, 1H), 2.55 (s, 3H), 2.32 (ddd, $J = 16.9, 8.4, 1.6$ Hz, 1H), 1.30 (t, $J = 7.0$ Hz, 3H). LCMS for $\text{C}_{14}\text{H}_{16}\text{ClFNO}_3$ ($\text{M}+\text{H}$) $^+$: $m/z = 300.1$; Found: 300.1.

Step 6. Diastereoisomers of 4-[3-chloro-6-ethoxy-2-fluoro-5-(1-hydroxyethyl)phenyl]pyrrolidin-2-one

10 one




The enantiomers from step 5 were each processed individually to the final products. A solution of 4-(3-acetyl-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one (0.402 g, 1.34 mmol) (from Step 5) in anhydrous methanol (6.7 mL) under an atmosphere of nitrogen at 0 °C was treated

15 with sodium tetrahydroborate (0.10 g, 2.7 mmol) and stirred at 0° C for 30 min. The reaction mixture was quenched with water at 0 °C and poured into water (50 mL)/ethyl acetate (100 mL) while stirring. The mixture was warmed to ambient temperature and the aqueous layer was separated and extracted with additional ethyl acetate (50 mL). The combined organic extracts were washed with


20 brine, dried over sodium sulfate, filtered, and concentrated to give white foams. The crude material were purified by flash column chromatography using acetonitrile (containing 7% methanol) in dichloromethane (0% - 100%) to give the desired products as mixtures of diastereoisomers [from peak 1 (0.40 g, 99%); from peak 2 (0.40 g, 99%)]. From peak 1: LCMS for $\text{C}_{14}\text{H}_{18}\text{ClFNO}_3$ ($\text{M}+\text{H}$) $^+$: $m/z = 302.1$; Found: 302.0. From peak 2: LCMS for $\text{C}_{14}\text{H}_{18}\text{ClFNO}_3$ ($\text{M}+\text{H}$) $^+$: $m/z = 302.1$; Found: 302.1.

Step 7. Diastereoisomers of 4-[3-chloro-5-(1-chloroethyl)-6-ethoxy-2-fluorophenyl]pyrrolidin-2-one



The mixture of diastereoisomers from step 6 were each processed individually to the final products. A solution of 4-[3-chloro-6-ethoxy-2-fluoro-5-(1-hydroxyethyl)phenyl]pyrrolidin-2-one (0.41 g, 1.4 mmol) (from Step 6) in methylene chloride (12 mL) was treated with *N,N*-dimethylformamide (0.011 mL, 0.14 mmol) followed by thionyl chloride (0.21 mL, 2.9 mmol) dropwise and stirred at 20 °C for 30 min. The reaction mixture was added dropwise to ice cooled saturated sodium bicarbonate solution and extracted with dichloromethane. The organic layer was separated and washed with brine, dried over sodium sulfate, filtered, and concentrated to give the desired products [from peak 1 (0.38 g, 87%); from peak 2 (0.39 g, 89%)] along with 17-18% of the styrene that formed from chloride elimination. These mixtures were used without further purification. From peak 1: LCMS for $C_{14}H_{17}Cl_2FNO_2$ ($M+H$)⁺: m/z = 320.1; Found: 320.0. From peak 2: LCMS for $C_{14}H_{17}Cl_2FNO_2$ ($M+H$)⁺: m/z = 320.1; Found: 320.0.

Step 8. Diastereoisomers of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one




The mixture of diastereoisomers from step 7 were each processed individually to the final products. A mixture of 4-[3-chloro-5-(1-chloroethyl)-6-ethoxy-2-fluorophenyl]pyrrolidin-2-one (0.36 g, 1.1 mmol) (from Step 7), 3-methyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine (0.19 g, 1.3 mmol), cesium carbonate (0.54 g, 1.7 mmol) and potassium iodide (18 mg, 0.11 mmol) in *N,N*-dimethylformamide (7.4 mL) was heated at 100 °C for 4.5 h. The reaction mixture was poured into water (30 mL) and extracted with ethyl acetate (3 x 50 mL) to give a mixture of diastereoisomer ((S)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one; (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one; (S)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one; and (R)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one). The mixture of diastereoisomers were purified by preparative LCMS (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.1% ammonium hydroxide, at flow rate of 60 mL/min) to give the desired products [from peak 1 were isolated peak A


(Example 1A) (0.13 g, 54%) and peak B (Example 1B) (0.11 g, 46%); from peak 2 were isolated peak A (Example 1C) (0.15 g, 63%) and peak B (Example 1D) (0.14 g, 55%).

Example 1B: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.12 (s, 1H), 7.82 (s, 1H), 7.52 (d, $J = 8.5$ Hz, 1H), 7.30 (br s, 1H), 6.23 (q, $J = 7.0$ Hz, 1H), 4.05 – 3.90 (m, 1H), 3.88 – 3.78 (m, 2H), 3.63 – 3.53 (m, 1H), 3.29 – 3.20 (m, 1H), 2.54 (s, 3H), 2.38 – 2.21 (m, 1H), 1.70 (d, $J = 7.1$ Hz, 3H), 1.39 (t, $J = 6.9$ Hz, 3H). LCMS for $\text{C}_{20}\text{H}_{23}\text{ClFN}_6\text{O}_2$ ($\text{M}+\text{H}$) $^+$: $m/z = 433.2$; Found: 433.1. Example 1C: ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.12 (s, 1H), 7.77 (s, 1H), 7.53 (d, $J = 8.5$ Hz, 1H), 7.26 (br s, 2H), 6.24 (q, $J = 7.0$ Hz, 1H), 4.04 – 3.94 (m, 1H), 3.93 – 3.85 (m, 1H), 3.84 – 3.77 (m, 1H), 3.61 – 3.53 (m, 1H), 3.27 – 3.22 (m, 1H), 2.54 (s, 3H), 2.30 (dd, $J = 18.1, 8.6$ Hz, 1H), 1.71 (d, $J = 7.1$ Hz, 3H), 1.40 (t, $J = 6.9$ Hz, 3H). LCMS for $\text{C}_{20}\text{H}_{23}\text{ClFN}_6\text{O}_2$ ($\text{M}+\text{H}$) $^+$: $m/z = 433.2$; Found: 433.1.

Examples 2A-2D. Diastereomers of 5-{3-[1-(4-amino-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one




Step 1: *tert*-Butyl [2-(3-acetyl-5-chloro-2-ethoxy-6-fluorophenyl)-2-hydroxyethyl]carbamate



0.2 M Osmium tetroxide in water (10 mL) was added to a solution of *tert*-butyl [(4-chlorobenzoyl)oxy]carbamate (Lawrence Harris, *J. Org.Chem.*, **2011**, 76, 358-372). (19 g, 70 mmol) in acetonitrile (210 mL) and stirred for 10 minutes. 1-(5-chloro-2-ethoxy-4-fluoro-3-vinylphenyl)ethanone (11.2 g, 46 mmol) (see Example 353, Step 1 of U.S. Patent No.: 9,199,982) as a solution in acetonitrile (210 mL) was added to the carbamate solution followed by the addition of water (50 mL) and the reaction was stirred for 3 hours at room temperature. The reaction was

quenched with saturated 10 M dipotassium disulfite in water (240 mL) and stirred for 5 minutes. Water was added and the reaction mixture was extracted with ethyl acetate. The extracts were washed with saturated sodium bicarbonate solution, brine and dried over sodium sulfate, filtered and evaporated. Purification on silica gel using ethyl acetate in hexanes (0-100%) gave the desired compound as a racemic mixture, 16.6 g, 95%. LCMS calculated for $C_{17}H_{23}ClFNO_5Na$ ($M+Na$)⁺: m/z = 398.1; found: 398.0.

Step 2: 5-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}-1,3-oxazolidin-2-one



The desired single enantiomer (peak 3) was prepared using the same procedure as Example 353, steps 8-12 of U.S. Patent No.: 9,199,982, except that the intermediate from step 1 in this example was racemic and thus the final separation of the four diastereomers occurred in step 12. Chiral purification on Phenomenex Lux Cellulose C-4, 21 x 250 mm (Chiral Technologies), 5 micron particle size, at flow rate of 18 mL/min using 30% ethanol in hexanes gave the peak 1: Example 2A (single enantiomer) (retention time = 12.7 minutes), peak 2: Example 2B (single enantiomer) (retention time = 14.2 minutes), peak 3: Example 2C (single enantiomer) (retention time = 20.3 minutes), and peak 4: Example 2D (single enantiomer) (retention time = 28.9 minutes); the most active enantiomer was peak 3. LCMS calculated for $C_{19}H_{21}ClFN_6O_3$ ($M+H$)⁺: m/z = 435.1; found: 435.1. ¹H NMR (500 MHz, DMSO-*d*₆): δ 8.15 (s, 1 H), 7.81 (s, 1 H), 7.71 (d, 1 H), 7.26 (bs, 1 H), 6.23 (m, 1 H), 5.84 (t, 1 H), 3.92 (m, 1 H), 3.83 (m, 1 H), 2.52 (s, 3 H), 1.75 (d, 3 H), 1.40 (m, 3 H).

Example A1: PI3K Enzyme Assay

PI3-Kinase luminescent assay kit including lipid kinase substrate, D-myo-phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂)D (+)-sn-1,2-di-O-octanoylglycerol, 3-O-phospho linked (PIP₂), biotinylated I(1,3,4,5)P₄, PI(3,4,5)P₃ Detector Protein is purchased from Echelon Biosciences (Salt Lake City, UT). AlphaScreen™ GST Detection Kit including donor and acceptor beads was purchased from PerkinElmer Life Sciences (Waltham, MA). PI3Kδ (p110δ /p85α)

is purchased from Millipore (Bedford, MA). ATP, MgCl₂, DTT, EDTA, HEPES and CHAPS are purchased from Sigma–Aldrich (St. Louis, MO).

AlphaScreen™ Assay for PI3Kδ

5 The kinase reaction are conducted in 384-well REMP plate from Thermo Fisher Scientific in a final volume of 40 µL. Inhibitors are first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay is 2%. The PI3K assays are carried out at room temperature in 50 mM HEPES, pH 7.4, 5mM MgCl₂, 50 mM NaCl, 5mM DTT and CHAPS 0.04%. Reactions are initiated by the addition of ATP, the final
10 reaction mixture consisted of 20 µM PIP2, 20 µM ATP, 1.2nM PI3Kδ are incubated for 20 minutes. 10 µL of reaction mixture are then transferred to 5 µL 50nM biotinylated I(1,3,4,5)P4 in quench buffer: 50 mM HEPES pH 7.4, 150 mM NaCl, 10 mM EDTA, 5 mM DTT, 0.1% Tween-20, followed with the addition of 10 µL AlphaScreen™ donor and acceptor beads suspended in quench buffer containing 25nM PI(3,4,5)P3 detector protein. The final concentration of both donor and acceptor
15 beads is 20 mg/ml. After plate sealing, the plate are incubated in a dark location at room temperature for 2 hours. The activity of the product is determined on Fusion-alpha microplate reader (Perkin–Elmer). IC₅₀ determination is performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software.

20 **Example A2: PI3K Enzyme Assay**

Materials

Lipid kinase substrate, phosphoinositol-4,5-bisphosphate (PIP2), are purchased from Echelon Biosciences (Salt Lake City, UT). PI3K isoforms α, β, δ and γ are purchased from Millipore (Bedford, MA). ATP, MgCl₂, DTT, EDTA, MOPS and CHAPS are purchased from Sigma–Aldrich
25 (St. Louis, MO).

The kinase reaction are conducted in clear-bottom 96-well plate from Thermo Fisher Scientific in a final volume of 24 µL. Inhibitors are first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay is 0.5%. The PI3K assays are carried out at room temperature in 20 mM MOPS, pH 6.7, 10
30 mM MgCl₂, 5 mM DTT and CHAPS 0.03%. The reaction mixture is prepared containing 50 µM PIP2, kinase and varying concentration of inhibitors. Reactions are initiated by the addition of ATP containing 2.2 µCi [γ-³³P]ATP to a final concentration of 1000 µM. The final concentration of PI3K isoforms α, β, δ and γ in the assay were 1.3, 9.4, 2.9 and 10.8 nM, respectively. Reactions are incubated for 180 minutes and terminated by the addition of 100 µL of 1 M potassium phosphate pH
35 8.0, 30 mM EDTA quench buffer. A 100 µL aliquot of the reaction solution are then transferred to 96-well Millipore MultiScreen IP 0.45 µm PVDF filter plate (The filter plate is prewetted with 200 µL

100% ethanol, distilled water, and 1 M potassium phosphate pH 8.0, respectively). The filter plate is aspirated on a Millipore Manifold under vacuum and washed with $18 \times 200 \mu\text{L}$ wash buffer containing 1 M potassium phosphate pH 8.0 and 1 mM ATP. After drying by aspiration and blotting, the plate is air dried in an incubator at 37°C overnight. Packard TopCount adapter (Millipore) is then
 5 attached to the plate followed with addition of $120 \mu\text{L}$ Microscint 20 scintillation cocktail (Perkin Elmer) in each well. After the plate sealing, the radioactivity of the product is determined by scintillation counting on Topcount (Perkin–Elmer). IC_{50} determination is performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software.

10 **Example A3: PI3K δ scintillation proximity assay**

Materials

$[\gamma\text{-}^{33}\text{P}]\text{ATP}$ (10mCi/mL) was purchased from Perkin–Elmer (Waltham, MA). Lipid kinase substrate, D-myo-Phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂)D (+)-sn-1,2-di-O-octanoylglyceryl, 3-O-phospho linked (PIP₂), CAS 204858-53-7, was purchased from Echelon
 15 Biosciences (Salt Lake City, UT). PI3K δ (p110 δ /p85 α) was purchased from Millipore (Bedford, MA). ATP, MgCl_2 , DTT, EDTA, MOPS and CHAPS were purchased from Sigma–Aldrich (St. Louis, MO). Wheat Germ Agglutinin (WGA) YSi SPA Scintillation Beads was purchased from GE healthcare life sciences (Piscataway, NJ).

20 The kinase reaction was conducted in polystyrene 384-well matrix white plate from Thermo Fisher Scientific in a final volume of $25 \mu\text{L}$. Inhibitors were first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay was 0.5%. The PI3K assays were carried out at room temperature in 20 mM MOPS, pH 6.7, 10 mM MgCl_2 , 5 mM DTT and CHAPS 0.03%. Reactions were initiated by the addition of ATP,
 25 the final reaction mixture consisted of $20 \mu\text{M}$ PIP₂, $20 \mu\text{M}$ ATP, $0.2 \mu\text{Ci}$ $[\gamma\text{-}^{33}\text{P}]\text{ATP}$, 4 nM PI3K δ . Reactions were incubated for 210 min and terminated by the addition of $40 \mu\text{L}$ SPA beads suspended in quench buffer: 150mM potassium phosphate pH 8.0, 20% glycerol. 25 mM EDTA, $400 \mu\text{M}$ ATP. The final concentration of SPA beads was 1.0mg/mL . After the plate sealing, plates were shaken overnight at room temperature and centrifuged at 1800 rpm for 10 minutes, the radioactivity of
 30 the product was determined by scintillation counting on Topcount (Perkin–Elmer). IC_{50} determination was performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software. IC_{50} data for the Examples is presented in Table B as determined by Assay A.

Table B.

Example #	PI3K δ SPA IC ₅₀ (nM)*
1A	+++
1B	+
1C	+
1D	+++
2A	+++
2B	+++
2C	+
2D	+++++

* column symbols:

+ refers to ≤ 10 nM

++ refers to >10 nM to 50 nM

5 +++ refers to >50 nM to 200 nM

++++ refers to >200 nM to 500 nM

+++++ refers to >500 nM

Example B1: B cell proliferation assay

10 To acquire B cells, human PBMC are isolated from the peripheral blood of normal, drug free donors by standard density gradient centrifugation on Ficoll-Hypaque (GE Healthcare, Piscataway, NJ) and incubated with anti-CD19 microbeads (Miltenyi Biotech, Auburn, CA). The B cells are then purified by positive immunosorting using an autoMacs (Miltenyi Biotech) according to the manufacture's instruction.

15 The purified B cells (2×10^5 /well/200 μ L) are cultured in 96-well ultra-low binding plates (Corning, Corning, NY) in RPMI1640, 10% FBS and goat F(ab')₂ anti-human IgM (10 μ g/ml) (Invitrogen, Carlsbad, CA) in the presence of different amount of test compounds for three days. [³H]-thymidine (1 μ Ci/well) (PerkinElmer, Boston, MA) in PBS is then added to the B cell cultures for an additional 12 hours before the incorporated radioactivity is separated by filtration with water
20 through GF/B filters (Packard Bioscience, Meriden, CT) and measured by liquid scintillation counting with a TopCount (Packard Bioscience).

Example B2: Pfeiffer cell proliferation assay

Pfeiffer cell line (diffuse large B cell lymphoma) are purchased from ATCC (Manassas, VA)
25 and maintained in the culture medium recommended (RPMI and 10% FBS). To measure the anti-proliferation activity of the compounds, the Pfeiffer cells are plated with the culture medium (2×10^3 cells / well/ per 200 μ l) into 96-well ultra-low binding plates (Corning, Corning, NY), in the presence or absence of a concentration range of test compounds. After 3-4 days, [³H]-thymidine (1 μ Ci/well) (PerkinElmer, Boston, MA) in PBS is then added to the cell culture for an additional 12 hours before

the incorporated radioactivity is separated by filtration with water through GF/B filters (Packard Bioscience, Meriden, CT) and measured by liquid scintillation counting with a TopCount (Packard Bioscience). IC₅₀ data for select compounds is presented in Table C.

5

Table C.

Example #	Pfeiffer IC ₅₀ (nM)*
1B	+
1C	+
2C	+

* column symbols:

+ refers to ≤ 10 nM

++ refers to >10 nM to 50 nM

10

Example C: Akt phosphorylation assay

Ramos cells (B lymphocyte from Burkitts lymphoma) are obtained from ATCC (Manassas, VA) and maintained in RPMI1640 and 10% FBS. The cells (3×10^7 cells /tube/3 mL in RPMI) are incubated with different amounts of test compounds for 2 hrs at 37 °C and then stimulated with goat F(ab')₂ anti-human IgM (5 µg/mL) (Invitrogen) for 17 minutes in a 37 °C water bath. The stimulated cells are spun down at 4 °C with centrifugation and whole cell extracts are prepared using 300 µL lysis buffer (Cell Signaling Technology, Danvers, MA). The resulting lysates are sonicated and supernatants are collected. The phosphorylation level of Akt in the supernatants are analyzed by using PathScan phospho-Akt1 (Ser473) sandwich ELISA kits (Cell Signaling Technology) according to the manufacturer's instruction.

20

Example D: Pfeiffer Model of Lymphoma

Methods

Female SCID mice, (5 to 8 weeks of age, Charles River Laboratories, Wilmington, MA) were inoculated with 1×10^7 tumor cells (Pfeiffer, ATCC #CRL-2632, Manassas, VA) and matrigel (BD Biosciences #354234) in 0.2 mL sterile saline. The inoculation was performed subcutaneously on the flank. Tumor tissue fragments (approximately 3 mm $\times$ 3 mm) were collected 3 to 6 weeks after the inoculation of cultured cells and implanted subcutaneously in lieu of cellular inoculation. Tissue fragments were implanted as solid pieces using blunt-tip forceps. The treatment of tumor bearing mice was started 15 to 25 days after tumor inoculation, depending upon the tumor size. Animals were sorted to obtain roughly equivalent mean tumor volumes in each group. Minimum mean tumor volume in all groups was 150 mm³ on the first day of treatment and groups consisted of 7 animals. Experimental therapeutic agent, Example 347, was administered to mice orally (PO). Treatment frequency was 2 times daily for a minimum of 14 days for efficacy. The size of subcutaneous tumors was measured 2 to 3 times weekly using a digital caliper. The tumor volume was calculated by

35

measuring the tumor in 2 dimensions and utilizing the equation: $\text{Volume} = [\text{Length} \times (\text{Width}^2)]/2$; where the larger number was length, and the smaller number width. If multiple tumors were formed, the final volume was the sum of the individual tumors subject to the same equation: eg, 2 tumors; $\text{Volume} = \{[L1 \times (W1)^2]/2\} + \{[L2 \times (W2)^2]/2\}$. Effects on tumor growth were reported as percent tumor growth inhibition (%TGI). Percent TGI was calculated with the equation: $(1 - (\text{Tx vol.} / \text{control vol.})) \times 100$, where control volume was the vehicle or untreated tumor volume on a given day, and Tx volume was any treatment group tumor volume on that same day. Statistical differences between treatment and vehicle controls were assessed using ANOVA: Single Factor test.

10 Results

(*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one (see Example 1) was evaluated as a single agent in the Pfeiffer human tumor xenograft model of diffuse large B-cell lymphoma, a subtype of NHL. Pfeiffer cancer cells were shown to be sensitive to the anti-proliferative effects of (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one *in vitro*. Therefore, a tumor model was established based on subcutaneous inoculation of tumor cells into immune compromised SCID mice and tumor-bearing mice received twice daily oral doses of vehicle or (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one at 0.3, 1, 3, or 10 mg/kg for 14 days. (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one treatment inhibited tumor growth by 22%, 24%, 36%, and 58% (percent tumor growth inhibition) with increasing dose (see *e.g.*, U.S. Patent No.: 9,932,341).

Example E: Dose-Escalation & Dose Expansion Studies

25 In the dose-escalation and dose-expansion studies described herein, the PK, PD, safety, and efficacy of (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt (*i.e.*, Compound 1C) (were evaluated in patients with relapsed or refractory B-cell malignancies. The PK characteristics of Compound 1C combined with the absence of dose-limiting toxicities (DLTs) permitted exposure levels that exceeded the IC_{90} throughout the once-daily (QD) dosing interval at all doses tested. In addition, the results of a food effect analysis indicated that Compound 1C may be dosed without regard to timing of meals. The near-maximal inhibition of pAKT in the *ex vivo* PD assay were consistent with exposure data. Results of the PI3K δ monotherapy and combination therapies are discussed below in Examples F-H.

Study Design and Patients

The dose-escalation and dose-expansion study was conducted in multiple parts: dose escalation of Compound 1C monotherapy (Part 1) followed by Cohort expansion (Part 3); Compound 1C + itacitinib dose escalation (Part 2) followed by Cohort expansion (Part 3); and Compound 1C + R-ICE (rituximab plus ifosfamide, carboplatin, and etoposide) dose evaluation and expansion (Part 6) (NCT02018861). Eligible patients (age ≥ 18 years) had diagnosed lymphoid malignancies of B-cell origin, including indolent or aggressive B-cell NHL malignancies, transformed NHL histologies, and Hodgkin lymphoma (HL), that had relapsed or were refractory to prior standard therapy. Patients with Burkitt lymphoma and precursor B-lymphoblastic leukemia/lymphoma were excluded. Eligible patients had Eastern Cooperative Oncology Group performance status (ECOG PS) ≤ 1 (dose escalation) or ≤ 2 (dose expansion); had adequate cardiac, liver, and kidney function; had life expectancy ≥ 12 weeks; had received ≥ 1 prior treatment regimen; and had not responded to or were not candidates for hematopoietic (H)SCT or other potentially curative therapy.

Patients were excluded if they had a history of untreated, symptomatic, or unstable brain metastases; spinal cord compression; lymphoma involving the central nervous system (permitted in the Compound 1C monotherapy expansion cohort A [B-cell malignancies] and Compound 1C + itacitinib combination expansion cohort); or received allogeneic hematopoietic stem cell transplant (HSCT) within 6 months or autologous HSCT within 3 months of enrollment. Prior treatment with PI3K δ inhibitors was excluded for Compound 1C monotherapy cohorts unless approved by the medical monitor, and prior JAK inhibitors were excluded for Compound 1C + itacitinib combination therapy cohorts.

72 patients were enrolled and treated with Compound 1C monotherapy (median age, 66 [range, 30–89] years) (Table 1). The median (range) duration of treatment was 4.0 (0.2–22.7) months. At the data cutoff, all but 11 subjects had discontinued study treatment, primarily for disease progression (n = 35 [49%]) and adverse events (AEs) (n = 14 [19%]).

Table 1. Patient demographics and baseline characteristics (ITT)

Characteristics	Compound 1C Monotherapy (N=72)	Compound 1C + Itacitinib (N=11)	Compound 1C + R-ICE (N=5)
Age, median (range), y	66 (30-89)	67 (24-82)	64 (52-72)
Age, >65 y, n (%)	37 (51)	7 (64)	1 (20)
Men, n (%)	41 (57)	6 (55)	3 (60)
Race, n (%)			
White/Caucasian	57 (79)	9 (82)	4 (80)
Black/African American	8 (11)	1 (9)	0

Characteristics	Compound 1C Monotherapy (N=72)	Compound 1C + Itacitinib (N=11)	Compound 1C + R-ICE (N=5)
Asian	0	0	0
Other*	7 (10)	1 (9)	1 (20)
Disease type, n (%)			
CLL	6 (8)	1 (9)	0
DLBCL	23 (32)	6 (55)	5 (100)
FL	14 (19)	1 (9)	0
HL	10 (14) [†]	2 (18) [‡]	0
MCL	9 (13)	1 (9)	0
MZL	9 (13) [§]	0 (0)	0
WM	1 (1)	0 (0)	0
Duration since prior diagnosis, median (range), y	4.95 (0.5-22.5)	3.2 (0.4-13.2)	0.8 (0.5-6.3)
≥3 prior systemic therapy regimens, n (%)	39 (54)	5 (45)	0
Prior HSCT, n (%)	21 (29)	1 (9)	0

*Includes American-Indian, Alaska Native, Pacific Islander, and other. [†]Includes classic HL (n=9) and nodular lymphocytic-predominant HL (n=1). [‡]Includes extranodal MZL of MALT type (n=2), nodal MZL (n=4), splenic MZL (n=2), and unknown MZL subtype (n=1). [§]Includes classic HL (n=2). CLL, chronic lymphocytic leukemia; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; HL, Hodgkin lymphoma; HSCT, hematopoietic stem cell transplant; MALT, mucosa-associated lymphatic tissue; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; R-ICE, rituximab plus ifosfamide, carboplatin, and etoposide; WM, Waldenström macroglobulinemia.

Dosing

For Compound 1C monotherapy, Compound 1C was self-administered orally, once daily (QD). Compound 1C was administered in a fasted state, except for patients participating in an optional food-effect cohort, until a protocol amendment allowed administration without regard to food. Treatment continued until unacceptable toxicity or disease progression. During dose escalation (see Table 2C), an initial single-patient cohort was treated with 5 mg QD and subsequent cohorts were enrolled using a 3 + 3 design, to identify any dose-limiting toxicities (DLTs; defined in Table 2A) within the 21-day observation period. If DLTs occurred in >1 of the first 3 patients or the total cohort of 6 patients, then the maximum tolerated dose (MTD) was deemed to be exceeded and the next lower tolerable dose level was defined as the MTD. All doses up to 45 mg QD were well tolerated over the DLT observation period of 21 days and therefore no MTD was identified. However, incidences of colitis were observed over the longer term (>3 months), which prompted a change in dosing regimen. The protocol was amended to include a modified dosing schedule wherein daily dosing was followed by weekly dosing, based on a PK simulation (see *Discussion*). Patients enrolled on or after this protocol amendment received Compound 1C 20 mg QD for the first 9 weeks followed by Compound

1C 20 mg once weekly (QW); patients who had already received Compound 1C 20 mg QD for ≥ 9 weeks were switched to the QW schedule. Per protocol amendment, ongoing and newly enrolled patients were required to take prophylactic treatment for *Pneumocystis jiroveci* pneumonia (PJP).

5

Table 2A. Definition of dose-limiting toxicity

Nonhematologic • $\geq$Grade 3 nonhematologic toxicity, excluding nausea, vomiting, diarrhea • $\geq$Grade 3 nausea, vomiting, or diarrhea uncontrolled by maximal antiemetic/antidiarrheal therapy lasting >48 hours • Any toxicity considered a DLT in the opinion of the investigator and medical monitor
Hematologic • Grade 4 neutropenia lasting ≥ 7 days* • Febrile neutropenia (ANC $<1.0 \times 10^9/L$ with a single temperature of $>38.3^\circ C$ ($101^\circ F$) or a sustained temperature of $\geq 38^\circ C$ ($100.4^\circ F$) for more than 1 hour • Grade 3 thrombocytopenia associated with clinically significant bleeding (clinically significant as determined by the investigator or resulting in the need for a transfusion of red blood cells) • Grade 4 thrombocytopenia lasting >7 days • Grade 4 anemia
General • Any specific AE that results in a dose delay or reduction in $>$ one-third of subjects

*Itacitinib can cause transient decreases in white blood cells due to margination; therefore, DLT rules required neutropenia to persist after holding itacitinib for 2 to 3 days. Where the clinical status of the subject allowed, investigators were encouraged to wait 24 hours before starting growth factors, to determine if white blood cell margination was contributing to the degree of neutropenia. AE, adverse event; ANC, absolute neutrophil count; DLT, dose-limiting toxicity.

10

For the Compound 1C + itacitinib combination, dose escalation of Compound 1C proceeded using 3 + 3 design, starting with an Compound 1C dose approximately 25% less than the recommended dose determined for Compound 1C monotherapy. Criteria for itacitinib dose interruption are summarized in Table 2A. For each cohort, itacitinib was co-administered at a dose of 300 mg QD.

15

For the Compound 1C + R-ICE combination, dose escalation of Compound 1C proceeded using 3 + 3 design with a starting dose approximately 25% below the recommended dose determined for Compound 1C. R-ICE chemotherapy components were administered either according to institutional practice or per protocol.

20

Assessments

Patients were assessed on Days 1, 8, and 15 of Cycle 1 and Day 1 (± 3) of each subsequent treatment cycle. A schedule of key study assessments is shown in Table 2B.

5

Table 2B. Study Assessments

		Treatment					Follow-up
		Cycle 1			Other cycles		
	Screening Days –30 to –1	Day 1	Day 8 (± 3 days)	Day 15 (± 3 days)	Day 1 (± 3 days)	EOT	EOT + 30 to 37 days
Prior/concomitant medications	X	X	X	X	X	X	X
Physical examination	X	X	X	X	X	X	X
Vital signs	X	X	X	X	X	X	X
12-lead ECG	X	X		X	X	X	X
Laboratory tests	X	X	X	X	X	X	X
ECOG PS	X	X	X	X	X	X	X
CT/MRI	X				X*	X	
FDG-PET	X				X**†	X	
Bone marrow examination	X				X‡		
Review AEs	X	X	X	X	X	X	X
Serum chemistry/hematology	X	X	X	X	X	X	X
Blood PK sample		X	X§	X	X§		
Blood PD sample		X		X	X¶		
Tumor tissue sample	X#				X**	X**	

*Every 9 weeks (3 cycles) or at a frequency consistent with the standard of care; performed only if measurable disease is present. †May be performed per standard of care; applicable for lymphomas only. ‡To confirm a CR.

§Not collected from patients receiving Compound 1C + R-ICE. ||Cycle 2 only; collected from patients with B-cell malignancies in the Compound 1C monotherapy expansion cohort (cohort A). ¶Through Cycle 6 for

10 Compound 1C monotherapy and Compound 1C + itacitinib combination, and in Cycles 3, 6, 9, and 12 for Compound 1C + R-ICE combination. #Archival tissue acceptable; for patients with CLL, peripheral blood acceptable. **Optional on-treatment or EOT biopsy (or peripheral blood for patients with CLL). AE, adverse

15 event; CLL, chronic lymphocytic leukemia; CR, complete response; CT, computed tomography; ECG, electrocardiogram; ECOG PS, Eastern Cooperative Oncology Group performance status; EOT, end of treatment; FDG-PET, fluorodeoxyglucose (FDG)-positron emission tomography; MRI, magnetic resonance

imaging; PD, pharmacodynamic; PK, pharmacokinetic; R-ICE, rituximab plus ifosfamide, carboplatin, and etoposide.

The primary endpoint was safety and tolerability of Compound 1C monotherapy, Compound
 20 1C + itacitinib, or Compound 1C + R-ICE as assessed by summary of treatment-emergent adverse events (TEAEs), clinical laboratory assessments, physical examination results, and 12-lead electrocardiograms. The severity of TEAEs was assessed using Common Terminology Criteria for Adverse Events version 4.03.

The secondary endpoints were efficacy and PK. Efficacy was measured by best overall
 25 response rate (ORR), defined as the proportion of patients achieving a partial response (PR) or

complete response (CR). Response was evaluated (per investigator's assessment) every 9 weeks based on the Lugano classification of lymphoma response criteria for HL and NHL (Cheson B.D. et al, *Journal of Clinical Oncology*, 2014;32(27):3059-3067), the International Working Group on Chronic Lymphocytic Leukemia criteria for chronic lymphocytic leukemia (CLL) (Hallek M. et al, *Blood*, 2008;111(12):5446-5456; Cheson B.D. et al, *J Clin Oncol*, 2012;30(23):2820-2822)), and the VIth International Workshop on Waldenström Macroglobulinemia response assessment for Waldenström Macroglobulinemia (WM) (Owen R.G. et al, *Br J Haematol*, 2013;160(2):171-176). If the patient's response to treatment was assessed by PET and CTI/MRI, then the PET result was used for response endpoints; CT/MRI result was used for an assessment if PET result was not available for the assessment; CT/MRI was used for response endpoints if it was the only form of assessment available for the patient.

Exploratory endpoints included: duration of response (DOR, duration of time from first response to death or disease progression, whichever occurred first); progression free survival (PFS, duration of time from first study dose to death or disease progression, whichever occurred first); population PK; pharmacodynamic (PD) relationship between PI3K δ inhibition (as determined by an ex vivo phosphorylated AKT [pAKT] assay).

Statistical Methods

Safety, tolerability, and efficacy analyses were performed on the safety/intent-to-treat population, which included all enrolled patients who received ≥ 1 dose of Compound 1C, itacitinib, or any of the R-ICE components. PK and PD analyses included all patients in the safety/intent to treat population with available PK/PD data. All statistical analyses were exploratory in nature and were summarized using descriptive statistics. ORRs were estimated with 95% confidence intervals (CIs) calculated based on the exact method for binomial distributions. Kaplan-Meier estimates of median DOR and PFS were presented with their respective 95% CIs.

Additional patient exclusion criteria

Additional patient exclusion criteria included radiation treatment within 4 weeks, investigational study drug within 28 days (or 5 half-lives, whichever is longer), approved anticancer drugs (except steroids at ≤ 10 mg prednisone daily) within 21 (42 for nitrosoureas) days (or 5 half-lives, whichever is longer), unresolved toxicity grade ≥ 2 , current or recent history of clinically meaningful infection, total bilirubin $\geq 1.2 \times$ upper limit of normal (ULN), alkaline phosphatase $\geq 2.5 \times$ ULN, aspartate aminotransferase (AST) or alanine aminotransferase (ALT) $\geq 2.0 \times$ ULN, or creatinine clearance < 50 mL/min based on Cockcroft-Gault formula.

Per protocol R-ICE dosing schedule

Per protocol, R-ICE was administered in 3 21-day cycles, according to the following schedule: rituximab 375 mg/m² on Days 1 and 2 of Cycle 1, and Day 1 of Cycles 2 and 3; ifosfamide 5000 mg/m² by continuous intravenous (IV) infusion over 24 h on Day 3 of each cycle; carboplatin (area under the curve = 5 mg/mL; maximum dose 800 mg) by IV infusion on Day 3; etoposide 100 mg/m² by IV on days 3 to 5.

Example F. PI3Kδ Inhibitor Monotherapy

This analysis includes data from patients enrolled into Compound 1C monotherapy dose-escalation (5-45 mg QD) and expansion (20 mg QD and 30 mg QD) cohorts.

Dose escalation and cohort expansion

The number of patients per dose level during Compound 1C monotherapy dose escalation and expansion is shown in Table 2C. No DLTs were identified, the MTD was not reached, and the 20 mg QD and 30 mg QD doses were expanded.

Table 2C. Monotherapy Dose Escalation and Cohort Expansion

				Expansion doses		
Dose	5 mg	10 mg	15 mg	20 mg	30 mg	45 mg
n	1	3	3	34	27	4
Escalation rules	In case of a grade ≥ 2 AE, the cohort was to be expanded to 3 patients		If 1 DLT was observed in the first 3 patients, 3 additional patients were to be enrolled in the cohort; if DLTs occurred in >1 of the first 3 patients or the total cohort of 6 patients, then the MTD was deemed to be exceeded			

AE = adverse event; DLT = dose limiting toxicity; MTD = maximum tolerated dose.

20 Pharmacokinetics and Pharmacodynamics of Compound 1C

All doses tested remained above the IC₉₀ for target inhibition throughout the dosing interval, as shown in Figure 1A; C_{max} and C_{trough} resulting from a 20 mg QD dose of Compound 1C were 16-fold and 2-fold above the IC₉₀ for pAKT inhibition, respectively. A PK simulation conducted to support selection of a 20 mg QW dosing schedule estimated that serum Compound 1C levels resulting from this schedule would exceed the IC₉₀ for target inhibition for ~36 hours and would have minimal to no inhibition for approximately half of the dosing interval. Following multiple-dose administration of Compound 1C alone, Compound 1C attained peak plasma concentrations rapidly, with a median

$t_{\max}$ of 0.5 to 1 hour. Subsequently, Compound 1C plasma concentrations declined in a monophasic fashion with a mean terminal-phase disposition $t_{1/2}$ of 8.6 to 11.5 hours. Dose proportionality appeared to be observed between 5 and 45 mg QD at steady state, as shown in Figure 1B.

Food effect was analyzed in 12 patients. Although co-administration of Compound 1C with high fat meal decreased $C_{\max}$ by 42% compared with the fasted state, it had a modest effect on area under the curve (AUC; 10% decrease). The fed versus fasted geometric mean ratio for AUC was 0.90 (90% CI: 0.68–1.18).

Near-maximal PI3K δ inhibition was observed at all Compound 1C doses for up to 6 hours post-dose, as measured by effects on pAKT levels in whole blood, as shown in Figure 4.

Safety and tolerability

TEAEs occurring in >10% of patients across all dose cohorts combined and in the 20 mg QD cohort are shown in Table 3.

Table 3. Nonhematologic TEAEs occurring in $\geq 15\%$ of patients receiving Compound 1C monotherapy (Safety Population)

Preferred Term (MedDRA)	20 mg* (N=34), n (%)			All doses (N=72), n (%)		
	Any Grade	Grade 3	Grade 4	Any Grade	Grade 3	Grade 4
Any TEAE	32 (94)	14 (41)	6 (18)	68 (94)	29 (40)	12 (17)
Diarrhea/Colitis [†]	12 (35)	3 (9)	1 (3)	26 (36)	6 (8)	1 (1)
Nausea	9 (26)	0	0	26 (36)	0	0
Fatigue	10 (29)	0	0	22 (31)	0	0
Rash [‡]	10 (29)	3 (9)	0	22 (31)	4 (6)	0
Cough	5 (15)	0	0	17 (24)	0	0
Vomiting	4 (12)	0	0	17 (24)	0	0
Dizziness	5 (15)	0	0	13 (18)	0	0
Pyrexia	7 (21)	0	0	13 (18)	1 (1)	0
Hypokalemia	6 (18)	1 (3)	0	12 (17)	1 (1)	0
Abdominal pain	6 (18)	1 (3)	0	11 (15)	1 (1)	0

Constipation	3 (9)	0	0	11 (15)	0	0
Decreased appetite	6 (18)	0	0	11 (15)	0	0
Night sweats	6 (18)	0	0	11 (15)	0	0
Pruritus	7 (21)	0	0	10 (14)	0	0
Back pain	6 (18)	0	0	9 (13)	0	0
Chills	5 (15)	0	0	9 (13)	0	0

*Based on starting dose. Includes patients who received QD and QW dosing. †Includes preferred terms of diarrhea, colitis, enterocolitis, gastrointestinal inflammation, colitis microscopic, and cytomegalovirus colitis. ‡Includes preferred terms of dermatitis exfoliative, rash, rash erythematous, rash macular, rash maculopapular, rash pruritic, exfoliative rash, rash generalized, rash popular, and rash pustular. MedDRA, Medical Dictionary for Regulatory Activities; TEAE, treatment-emergent adverse event.

The most common ($\geq 30\%$) any grade nonhematologic TEAEs in all patients were diarrhea/colitis (36%), nausea (36%), fatigue (31%), and rash (31%). One patient (1%) each experienced grade 3 pneumonitis and grade 3 cytomegalovirus colitis, and 7 patients (10%) experienced pneumonia (grade 3, n=3 [4%]). Treatment-emergent laboratory abnormalities occurring in $\geq 30\%$ of patients are shown in Table 4. New/worsening of any grade and grade 3/4 neutropenia occurred in 32 patients (44%) and 14 patients (19%), respectively (Table 4). SAEs experienced by >2 patients were diarrhea/colitis (n=9 [13%]), pyrexia (n=4 [6%]), hypotension (n=3 [4%]), and sepsis (n=3 [4%]). Any grade SAEs of infections and infestations (System Organ Class) occurred in 11 patients (15%), including pneumonia in 2 patients (3%).

Table 4. New or worsening hematologic laboratory abnormalities occurring in $\geq 30\%$ of patients receiving Compound 1C monotherapy (Safety Population)*

Total (N=72)	Any Grade, n (%)	Grade 3, n (%)	Grade 4, n (%)
Leukopenia	36 (50)	6 (8)	0
Neutropenia	32 (44)	10 (14)	4 (6)
Lymphopenia	25 (35)	13 (18)	2 (3)
Thrombocytopenia	25 (35)	2 (3)	5 (7)
Anemia	22 (31)	6 (8)	NA

*Based on reported laboratory values. NA, not applicable

Among TEAEs of interest, the median (range) time to onset of grade 3/4 diarrhea/colitis was 5.7 (1.6–14.9) months and median (range) time to onset of grade 3/4 rash was 2.9 (1.5–9.3) months. AST and ALT elevations occurred in 21 (29%) and 20 (28%) patients, respectively; all events were grade 1, except 4 events in 2 patients (1 patient with 1 event each of grade 3 AST and grade 2 ALT elevation, both occurring >30 days after the last dose of Compound 1C; 1 patient with 1 event each of grade 3 AST and ALT elevation considered secondary to sepsis). Bilirubin elevation occurred in 8 (11%) patients; all events were ≤grade 2 except 1 grade 3 event that occurred >30 days after date of last dose. Changes in AST and ALT levels over time are shown in Figures 5A-5B. There were no TEAEs of PJP or bowel perforation. Five patients (7%) experienced hypertension (all grade 1/2); 7 (10%) experienced hyperglycemia (all grade 1/2 except 1 grade 3).

Any grade TEAEs led to Compound 1C dose interruption in 30 patients (42%), dose reduction in 4 patients (6%), and treatment discontinuation in 14 patients (19%). The most common nonhematologic TEAEs leading to Compound 1C dose interruption were diarrhea/colitis (11/72 [15%]) and pyrexia (4/72 [6%]). TEAEs leading to death occurred in 2 patients (respiratory failure, and respiratory failure and sepsis); none was deemed related to Compound 1C.

Long-term tolerability and dosing

Prior to implementation of the QW dosing schedule, 9/31 patients (29%) with DLBCL, FL, MCL, and MZL receiving Compound 1C monotherapy had discontinued due to TEAEs, most commonly (n≥2) diarrhea/colitis (n=3 [10%]), or rash (n=2 [6%]); however, no patient had discontinued due to treatment-related AEs during the first 9 weeks of treatment. As of the data cutoff date, 26 (36%) of the 72 enrolled patients had received QW dosing of Compound 1C (after receiving QD dosing for at least 9 weeks) for a median (range) of 2.5 (1.2–9.6) months and a total of 105 patient-months; 46 patients (64%) received QD dosing only; no patients were still in the first 9 weeks of QD dosing. Four of the 26 patients initiated the QW schedule at a <20 mg dose due to previous dose reductions (1 at 5 mg and 3 at 10 mg). Among the 26 patients who received QW dosing, 8 (31%) had dose interruptions (TEAEs leading to dose interruption in >1 patient were diarrhea and neutropenia [n=2 each]), 1 (4%) had a dose reduction (due to rash pustular), and none discontinued treatment due to a TEAE. Four patients receiving QW dosing (15%) experienced a total of 6 SAEs (abdominal pain, diarrhea, pyrexia, bronchitis, sepsis, and dehydration, n=1 each) during the QW dosing period. No grade 4 nonhematologic TEAEs were reported during QW dosing. Two patients had grade 3 diarrhea and rash (n=1[4%] each); both events occurred shortly after the switch

from QD dosing. No new/worsening grade 4 neutropenia was reported, and one patient reported a new/worsening grade 4 thrombocytopenia during QW dosing.

Efficacy

5 ORR among evaluable patients varied by disease type, as shown in Table 5. Among patients having evaluable lesion size at baseline and post-baseline, deep responses (>75% decrease from baseline in target lesion size) occurred in 4/7 patients (57%) with marginal zone lymphoma (MZL), 7/14 (50%) with follicular lymphoma (FL), 3/8 (38%) with mantle cell lymphoma (MCL), and 2/20 (10%) with diffuse large B-cell lymphoma (DLBCL) (Figure 10 2). Among patients with these tumor types, the vast majority (93%) of responses occurred by the first assessment (~9 weeks), as shown in Figure 6. Durable responses (>6 months) were observed among patients receiving QW dosing and several patients remained on the QW schedule for at least 8 months, as shown in Figure 3, including 2 who were administered 10 mg QW. The median duration of response was 13.5 months (95% CI: 8.3–18.8) for DLBCL, 15 4.4 months (95% CI: 2.1–NE) for MZL, and was not reached for FL or MCL. Figure 7 shows representative positron emission tomography (PET) images of a patient who achieved complete response with Compound 1C monotherapy. The patient was <65 years of age, with mantle cell lymphoma with no bone marrow involvement (target lesion, 98 mm × 58 mm) and had received 1 prior treatment (R-HyperCVAD). Upon treatment with Compound 1C, 30 mg 20 once daily, the patient achieved complete response at week 9.

Table 5. Best overall response among patients receiving Compound 1C monotherapy*

Patients	N	ORR, n (% [95% CI]) [†]	CR/CMR, n (%)	PR/PMR, n (%)	SD, n (%)	PD/PMD, n (%)	Not evaluable/ not assessed, n (%)
DLBCL [‡]	23	7 (30 [13–53])	4 (17)	3 (13)	4 (17)	11 (48)	1 (4)
GCB	19	6 (32 [13–57])	4 (21)	2 (11)	3 (16)	9 (47)	1 (5)
ABC	2	1 (50 [1–99])	0 (0)	1 (50)	0 (0)	1 (50)	0
FL	14	10 (71 [42–92])	3 (21)	7 (50)	1 (7)	2 (14)	1 (7)
MZL [¶]	9	7 (78 [40–97])	3 (33)	4 (44)	1 (11)	1 (11)	0
MCL	9	6 (67 [30–93])	4 (44)	2 (22)	2 (22)	0 (0)	1 (11)
HL ^{**}	10	2 (20 [3–56])	0 (0)	2 (20)	3 (30)	4 (40)	1 (10)
CLL	6	2 (33 [4–78])	0 (0)	2 (33)	2 (33)	1 (17)	1 (17)
WM	1	1 (100 [3–100])	0 (0)	1 (100)	0 (0)	0 (0)	0

*Assessed by Lugano Classification, CLL IWGC Criteria, and the VIth International Workshop on WM by Disease Subtype by CT or PET. †95% CIs were calculated based on the exact method for binomial distributions. ‡Two patients had unknown DLBCL subtype. §Includes extranodal MZL of MALT type (n=2), nodal MZL (n=4), splenic MZL (n=2), and unknown MZL subtype (n=1). ¶4 patients with MCL and 3 patients with CLL had received ibrutinib before the study, of whom a best overall response of CR or PR was achieved by 2 patients with MCL and 1 patient with CLL. **Includes classic HL (n=9) and nodular lymphocytic-predominant HL (n=1). ABC, activated B cell-like; CI, confidence interval; CLL, chronic lymphocytic leukemia; CMR, complete metabolic response; CR, complete response; CT, computed tomography; DLBCL, diffuse large B-cell lymphoma; FL, follicular lymphoma; HL, Hodgkin lymphoma; GCB, germinal center B-cell-like; CLL IWGC, International Working Group for CLL; MALT, mucosa-associated lymphatic tissue; MCL, mantle cell lymphoma; MZL, marginal zone lymphoma; ORR, overall response rate; PD, progressive disease; PET, positron emission tomography; PMD, progressive metabolic disease; PMR, partial metabolic response; PR, partial response; SD, stable disease; WM, Waldenström macroglobulinemia.

Example G. PI3Kδ Inhibitor Combination Therapy

Compound 1C combined with itacitinib

Eleven patients were enrolled and treated with Compound 1C + itacitinib. The median (range) duration of both Compound 1C and itacitinib treatment was 2.1 (1.0–10.3) months. At data cutoff, treatment was ongoing in 2 patients (18%); 8 patients (73%) discontinued treatment due to disease progression, and 1 (9%) due to TEAE.

Of the 11 patients treated with Compound 1C + itacitinib, 8 received Compound 1C 20 mg QD and 3 received Compound 1C 30 mg QD; all patients received itacitinib 300 mg QD. No DLTs were observed. Nine patients (82%) experienced TEAEs, 5 (45%) had grade 3/4 TEAEs, and 2 (18%) had SAEs. The most common any grade TEAEs (n≥2) were nausea, fatigue (n=5 each), cough, hypertension, vomiting (n=3 each), decreased appetite, hyperglycemia, lacrimation increased, oral herpes, and tachycardia (n=2 each). Five patients experienced grade 3/4 TEAEs: grade 3 spinal cord compression, grade 3 intractable pain, and grade 3 elevated alkaline phosphatase (n=1); grade 3 worsening hyperlipidemia (n=1); grade 3 hypertension (n=1); grade 3 diarrhea and grade 3 dehydration (n=1); and grade 3 hypertension and grade 4 hypercalcemia (n=1). One patient had a new/worsening grade 4 hematologic laboratory abnormality (thrombocytopenia). One SAE each of grade 3 dehydration and grade 4 hypercalcemia were reported. Two patients (18%) had TEAEs leading to dose interruption of both Compound 1C and itacitinib and 1 patient (9%) had a TEAE leading to discontinuation of both Compound 1C and itacitinib; no patient had a TEAE leading to dose reduction. No TEAEs led to death.

Among evaluable patients, best overall response of CR/complete metabolic response was achieved by patients with CLL (1/1) and FL (1/1); best overall response of partial metabolic response was achieved by patients with MCL (1/1) and classic HL (1/2). All 6

patients with DLBCL and 1 of 2 patients with classic HL had best overall response of PD/progressive metabolic disease.

Compound 1C combined with R-ICE

5 Five patients with DLBCL were treated with Compound 1C + R-ICE. The median (range) duration of Compound 1C treatment was 2.3 (1.5–3.7) months; 4 patients received 3 cycles of R-ICE each, and 1 patient received 2 cycles of R-ICE. As of the data cutoff date, all patients have discontinued treatment due to physician decision (n=2), progressive disease, adverse events, and other (n=1 each). Of the 5 patients, 4 received Compound 1C 15 mg QD and 1 received 20 mg QD. No DLTs were observed. The most common any-grade nonhematologic TEAEs (≥ 2 patients) were dizziness (n=3), alopecia, constipation, fluid overload, headache, hypokalemia, and night sweats (n=2 each). Three patients reported new/worsening grade 4 thrombocytopenia and 2 patients reported new/worsening grade 4 neutropenia. Two patients (40%) experienced SAEs, 1 patient experienced grade 3 acute encephalopathy (related to ifosfamide), and 1 experienced grade 2 atrial flutter and atrial fibrillation and grade 3 dyspnea, neutropenia, and febrile neutropenia (all related to both Compound 1C and R-ICE). Three patients (60%) had TEAEs leading to dose interruption of both Compound 1C and R-ICE, 1 patient (20%) had a TEAE leading to dose interruption of only Compound 1C, and 1 patient (20%) had TEAEs leading to discontinuation of both Compound 1C and R-ICE; no patient had a TEAE leading to dose reduction. No TEAEs led to death.

Of the 5 patients treated with Compound 1C + R-ICE, 3 patients achieved a complete metabolic response (1 of the 3 patients indicated a desire to proceed to SCT) and 2 patients had stable disease as best response.

25

Example H. Results of the PI3K δ Monotherapy and Combination Therapies

Compound 1C demonstrated a differentiated safety profile relative to first-generation PI3K δ inhibitors (*e.g.*, idelalisib and duvelisib) (Coutre S.E. et al, *Leukemia & Lymphoma*, 2015;56(10):2779-2786; Flinn I.W. et al, *Blood*, 2018;131(8):877-887), most notably by the near absence of grade ≥ 2 transaminitis. Unlike copanlisib, a pan-PI3K inhibitor that is approved in the United States as monotherapy for the treatment of relapsed FL (ALIQOPA™, (copanlisib) for injection, for intravenous use. U.S. Whippany, NJ: Bayer HealthCare Pharmaceuticals Inc.; 2017), there was also no clinically meaningful hypertension or hyperglycemia, presumably due to the high selectivity of Compound 1C for the δ isoform of PI3K. Although no PJP was observed, a protocol amendment made PJP prophylaxis

35

mandatory part-way through the study in response to a PJP infection reported in a different phase 1 study that combined Compound 1C with pembrolizumab.

Diarrhea/colitis and rash, toxicities common to PI3K inhibitors, were 2 of the most frequent toxicities observed in this study; nevertheless, the majority of diarrhea events were of grade 1 or 2 and were manageable. The majority of grade 3/4 diarrhea/colitis and rash occurred after the first disease assessment; overall any grade diarrhea/colitis or rash led to treatment discontinuation in 9/72 (13%) of patients, including 3 patients with diarrhea/colitis who were positive upon rechallenge at the same dose. Grade 4 neutropenia (6%) was observed only during QD dosing (earliest occurring after 21 days of treatment). The safety profile associated with the 20 mg QD dose was generally consistent with that observed for all doses combined and with that observed for each individual dose level (data not shown), although relatively few patients were dosed at levels below 20 mg QD and above 30 mg QD.

Long-term dosing of Compound 1C identified late-onset TEAEs that led to treatment discontinuation. To improve the long-term tolerability of Compound 1C while maintaining a high response rate, a modified dosing regimen was implemented (20 mg QD for 9 weeks followed by 20 mg QW). The period of daily dosing was maintained to preserve the potential for rapid onset of response (at the time of implementation, all 11 patients with NHL who were administered 20 mg QD had achieved an objective response, 10 of which occurred at the first disease assessment). The intermittent portion of the dosing regimen, which was implemented to decrease the incidence of late-onset TEAEs, was based on comparative PK/PD simulation with copanlisib (Patnaik A., et al, *Annals of Oncology*, 2016;27(10):1928-1940). Without being bound by theory, the 20 mg QW regimen is anticipated to provide similar pAKT inhibition as expected from QW dosing of copanlisib. Encouragingly, none of the 26 patients who received QW dosing discontinued study treatment due to TEAEs, based on data accrued during a 105 patient-month period. This observation, combined with the absence of colitis, grade 4 nonhematologic TEAEs and neutropenia, and a low rate of grade 3 diarrhea and rash (n=1 each) suggest that the QW schedule is better tolerated than the continuous QD schedule using the doses evaluated herein.

Compound 1C monotherapy effected a high rate of rapid, deep, and durable objective responses in relapsed or refractory B-cell NHL at doses ranging from 10 mg QD to 45 mg QD. ORR across all subtypes ranged from 30% to 78% (DLBCL, 30%; MCL, 67%; FL, 71%, MZL, 78%) and deep responses occurred in >30% of patients. Anecdotally, objective responses were observed in 2 of the 4 patients with MCL that had been previously treated with ibrutinib; 1 of these responses was ongoing at approximately 6 months at data cutoff. Among patients with DLBCL, substantially more patients with GCB subtype (n=19) were

enrolled relative to those with ABC subtype (n=2). These small, disproportionate numbers are insufficient to determine if Compound 1C is more active in one subtype than the other.

Durable responses were observed in all 4 of these NHL subtypes including ~17 months in FL, ~10 months in MZL, ≥14 months in MCL, and ~19 months in DLBCL, all of which included QW dosing and 2 of whom (FL and MZL) had ongoing responses at data cutoff. Taken together, these data suggest that QW dosing of Compound 1C can enable durable responses in patients with B-cell NHL.

The PK and safety data from the combination studies show that Compound 1C can be safely combined with a 300 mg QD dose of itacitinib. Similarly, the combination of Compound 1C with R-ICE also appears to be tolerated. Additional studies are required to evaluate the safety of Compound 1C combined with R-ICE and the potential for increased efficacy for each of the combinations.

Compound 1C demonstrated anti-tumor activity with rapid and deep responses as well as high response rates and a differentiated toxicity profile in patients with relapsed or refractory B-cell malignancies. The modified dosing regimen described herein appeared to improve the long-term tolerability of Compound 1C and enabled durable responses in several patients. The efficacy, safety, PK, and PD data presented here support further investigation of Compound 1C monotherapy in ongoing phase 2 studies in select NHL subtypes (DLBCL [NCT02998476], FL [NCT03126019], MZL [NCT03144674], and MCL [NCT03235544]).

Phase 1 studies exploring several combination strategies in patients with NHL (NCT03039114; NCT03424122) are also underway.

Example I. Comparative Maintenance Dosing

In this study, subjects are administered 20 mg of Compound 1C once daily for 8 weeks in combination with a JAK1 inhibitor. After week 8, subjects are assigned to 1 of 2 maintenance dose groups based on block randomization. In one group, subjects are administered 5 mg of Compound 1C once daily. In the second group, subjects are administered 20 mg of Compound 1C once weekly.

The safety and tolerability of Compound 1C is assessed by summary of treatment-emergent adverse events (TEAEs), clinical laboratory assessments, physical examination results, and 12-lead electrocardiograms. The severity of TEAEs is assessed using Common Terminology Criteria for Adverse Events version 4.03. Secondary endpoints are efficacy and PK, as described in Example E.

Without being bound by theory, the 5 mg once daily maintenance dose of Compound 1C is believed to provide similar safety and tolerability as the 20 mg once weekly maintenance dose of Compound 1C.

5 **Example J. PI3K δ Combination Therapy with JAK1/2 Inhibitor in Treatment of Myelofibrosis**

Compound 1C was evaluated in combination with ruxolitinib in patients with primary or secondary myelofibrosis (MF) (*e.g.*, PMF, PPV-MF, or PET-MF) who have exhibited suboptimal response with ruxolitinib monotherapy. The study was designed in three parts, as follows:

10 Part 1, Safety Run-in: 3+3 design; ruxolitinib + Compound 1C administered in doses of 10 mg or 20 mg QD for 8 weeks, followed by 10 mg or 20 mg QW.

15 Part 2, Randomized Phase (1:1): **Group 1:** ruxolitinib + Compound 1C administered in doses of 10 mg QD for 8 weeks, followed by 10 mg QW. **Group 2:** ruxolitinib + Compound 1C administered in doses of 20 mg QD for 8 weeks followed by 20 mg QW.

Part 3: Additional dosing regimens, including continuous daily dosing.

The “suboptimal response” with ruxolitinib monotherapy was defined as follows:

- A patient treated with ruxolitinib for ≥ 6 months with stable dose for ≥ 8 weeks immediately prior to enrollment (acceptable doses of prior ruxolitinib: 5 mg – 25 mg BID); and
- Palpable spleen > 10 cm below left subcostal margin on physical examination at screening; or
- Palpable spleen 5 cm – 10 cm below left subcostal margin on physical examination and active symptoms of myelofibrosis at the screening visit, defined as 1 symptom score ≥ 5 ; or 2 symptom scores ≥ 3 each, using the following screening symptom form: 10 point scale for each of 7 symptoms. Symptoms include: night sweats, pruritus, abdominal discomfort, pain under left ribs, early satiety, bone/muscle pain, and inactivity.

25 Dosing schemes and preliminary results of the Compound 1C + ruxolitinib study are shown in Appendix A, the disclosure of which is incorporated herein by reference in its entirety. As shown Appendix A, add on therapy with Compound 1C appears to provide additional clinical benefit in patients with MF who have suboptimal responses to ruxolitinib monotherapy. A reduction in spleen volume in 56% of patients was observed at week 12, with symptoms improving as early as 4 weeks; median symptom improvement at 24 weeks was 36%. The combination of Compound 1C and ruxolitinib was well tolerated in patients, with

no dose limiting toxicities (DLTs) in Part 1 and an expected grade 3/4 adverse effect (AE) profile. Of 17 patients on therapy >6 months, only 2 (7%) discontinuations were due to AEs. AEs common to PI3K δ inhibitors, such as hepatic, rash, and colitis, were infrequent.

- 5 Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

10

WHAT IS CLAIMED IS:

1. A method of treating a disease in a patient, wherein said disease is associated with increased expression or activity of PI3K δ kinase, the method comprising:
 - i) administering to the patient an inhibitor of PI3K δ , which is 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof, in a first dosage that is about 3 mg/day to about 50 mg/day for a first period of time which is about 2 weeks to about 12 weeks; and
 - ii) administering to the patient a second dosage of the inhibitor of PI3K δ which is less than the first dosage administered at the end of the first period of time and which is:
 - (a) about 2.5 mg/day or less; or
 - (b) about 50 mg/week or less;and wherein the second dosage is administered for a second period of time which occurs after the first period of time.
2. The method of claim 1, wherein the first dosage is about 20 mg/day.
3. The method of claim 1 or 2, wherein the first period of time is about 8 weeks to about 12 weeks.
4. The method of claim 1 or 2, wherein the first period of time is about 8 weeks.
5. The method of any one of claims 1 to 4, wherein the first dosage is reduced during the first period of time.
6. The method of any one of claims 1 to 5, wherein the second dosage is about 2.5 mg/day or less.
7. The method of any one of claims 1 to 5, wherein the second dosage is about 2.5 mg/day.
8. The method of any one of claims 1 to 5, wherein the second dosage is about 50 mg/week or less.
9. The method of any one of claims 1 to 5, wherein the second dosage is about 20 mg/week.

10. The method of any one of claims 1 to 9, wherein the second dosage is reduced during the second period of time.
11. The method of any one of claims 1 to 10, wherein the patient has been identified as exhibiting one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).
12. The method of any one of claims 1 to 11, wherein the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs) during the first period of time.
13. The method of any one of claims 1 to 12, wherein the first time period ends when the patient has been identified as exhibiting the one or more symptoms associated with one or more treatment-emergent adverse events (TEAEs).
14. The method of any one of claims 1 to 13, wherein the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis, nausea, fatigue, rash, neutropenia, pyrexia, hypotension, sepsis, respiratory failure, pneumonitis, pneumonia, hypertension, hyperglycemia, abdominal pain, bronchitis, dehydration, thrombocytopenia, cough, vomiting, decreased appetite, increased lacrimation, oral herpes, tachycardia, spinal cord compression, intractable pain, elevated alkaline phosphatase, elevated transaminase, hyperlipidemia, hypercalcemia, dizziness, alopecia, constipation, fluid overload, headache, hypokalemia, night sweats, encephalopathy, atrial flutter, atrial fibrillation, and dyspnea.
15. The method of any one of claims 1 to 13, wherein the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis, nausea, fatigue, rash, cough, vomiting, dizziness, pyrexia, hypokalemia, abdominal pain, constipation, decreased appetite, night sweats, pruritus, back pain, chills, leukopenia, neutropenia, lymphopenia, thrombocytopenia, and anemia.
16. The method of any one of claims 1 to 13, wherein the one or more treatment-emergent adverse events comprise one or more of diarrhea/colitis and rash.
17. The method of any one of claims 14 to 16, wherein the diarrhea/colitis comprises one or more of diarrhea, colitis, enterocolitis, gastrointestinal inflammation, colitis microscopic, and cytomegalovirus colitis.

18. The method of any one of claims 14 to 17, wherein the rash comprises one or more of dermatitis exfoliative, rash, rash erythematous, rash macular, rash maculopapular, rash pruritic, exfoliative rash, rash generalized, rash popular, and rash pustular.
19. The method of any one of claims 14 to 18, wherein the neutropenia comprises febrile neutropenia.
20. The method of any one of claims 14 to 19, wherein the elevated transaminase comprises elevated alanine transaminase (ALT), aspartate transaminase (AST), or a combination thereof.
21. The method of any one of claims 1-20, wherein the inhibitor of PI3K δ is (S)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof.
22. The method of any one of claims 1-20, wherein the inhibitor of PI3K δ is (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof.
23. The method of any one of claims 1-20, wherein the inhibitor of PI3K δ is (S)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof.
24. The method of any one of claims 1-20, wherein the inhibitor of PI3K δ is (R)-4-(3-((R)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof.
25. The method of any one of claims 1-20, wherein the inhibitor of PI3K δ is a pharmaceutically acceptable salt of 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one.
26. The method of claim 25, wherein the inhibitor of PI3K δ is (R)-4-(3-((S)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt.

27. The method of claim 25, wherein the salt is a 1:1 stoichiometric ratio of (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one to hydrochloric acid.

28. The method of claim 25, wherein the salt is crystalline.

29. A method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day to about 50 mg/day for a first period of time which is about 8 weeks to about 9 weeks; and

ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 2.5 mg/day or less for a second period of time which occurs after the first period of time.

30. A method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day to about 50 mg/day for a first period of time which is about 8 weeks to about 9 weeks; and

ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 20 mg/week to about 50 mg/week for a second period of time which occurs after the first period of time.

31. A method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL),

Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day for a first period of time which is about 8 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 2.5 mg/day for a second period of time which occurs after the first period of time.

32. A method of treating a disease in a patient, wherein said disease selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM), the method comprising:

- i) administering to the patient (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt in a first dosage that is about 20 mg/day for a first period of time which is about 8 weeks; and
- ii) administering to the patient a second dosage of the (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt, which is about 20 mg/week for a second period of time which occurs after the first period of time.

33. A method of treating a disease in a patient, wherein said disease is associated with increased expression or activity of PI3K δ kinase, the method comprising:

- i) administering to the patient an inhibitor of PI3K δ , which is 4-{3-[1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl]-5-chloro-2-ethoxy-6-fluorophenyl}pyrrolidin-2-one, or a pharmaceutically acceptable salt thereof, in a first dosage that is about 3 mg/day to about 50 mg/day for a first period of time which is about 2 weeks to about 12 weeks; and
- ii) administering to the patient a second dosage of the inhibitor of PI3K δ which is less than the first dosage administered at the end of the first period of time and which is between about 2.5 mg/day and about 7.5 mg/day; and

wherein the second dosage is administered for a second period of time which occurs after the first period of time.

34. The method of claim 33, wherein the second dosage is about 3.0 mg/day to about 7.0 mg/day.

35. The method of claim 33, wherein the second dosage is about 4.0 mg/day to about 6.0 mg/day.

36. The method of claim 33, wherein the second dosage is about 5.0 mg/day.
37. The method of any one of claims 33-36, wherein said inhibitor of PI3K δ is (*R*)-4-(3-((*S*)-1-(4-amino-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)-5-chloro-2-ethoxy-6-fluorophenyl)pyrrolidin-2-one hydrochloric acid salt.
38. The method of any one of claims 1-28 and 33-37, wherein said disease is selected from chronic lymphocytic leukemia (CLL), diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Hodgkin lymphoma (HL), mantle cell lymphoma (MCL), marginal zone lymphoma (MZL), and Waldenström macroglobulinemia (WM).
39. The method of any one of claims 29-32 and 38, wherein the marginal zone lymphoma (MZL) is selected from extranodal MZL, nodal MZL, splenic MZL, and unknown MZL subtype.
40. The method of any one of claims 29-32 and 38, wherein the Hodgkin lymphoma (HL) is selected from classic Hodgkin lymphoma (HL) and nodular lymphocytic-predominant HL.
41. The method of any one of claims 29-32 and 38, wherein the diffuse large B-cell lymphoma is selected from activated B-cell like (ABC) diffuse large B cell lymphoma (ABC-DLBCL) and germinal center B cell (GCB) diffuse large B cell lymphoma (GCB-DLBCL).




Figure 1A




Figure 1B




Figure 2




Figure 3






Figure 4






Figure 6




Figure 7