




(86) Date de dépôt PCT/PCT Filing Date: 2011/05/27
(87) Date publication PCT/PCT Publication Date: 2011/12/08
(85) Entrée phase nationale/National Entry: 2012/11/28
(86) N° demande PCT/PCT Application No.: EP 2011/058713
(87) N° publication PCT/PCT Publication No.: 2011/151259
(30) Priorité/Priority: 2010/06/01 (EP10164587.7)

(51) Cl.Int./Int.Cl. *C07D 487/04* (2006.01),
A61K 31/4985 (2006.01)
(71) Demandeur/Applicant:
BAYER INTELLECTUAL PROPERTY GMBH, DE
(72) Inventeurs/Inventors:
KLAR, ULRICH, DE;
KOPPITZ, MARCUS, DE;
JAUTELAT, ROLF, DE;
KOSEMUND, DIRK, DE;
BOHLMANN, ROLF, DE;
BADER, BENJAMIN, DE;
LIENAU, PHILIP, DE;
...
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : IMIDAZOPYRAZINES SUBSTITUEES
(54) Title: SUBSTITUTED IMIDAZOPYRAZINES



(57) **Abrégé/Abstract:**

The present invention relates to substituted imidazopyrazine compounds of general formula (I) in which A, R¹, R³ and R⁵ are as given in the description and in the claims, to methods of preparing said compounds, to pharmaceutical compositions and combinations comprising said compounds, to the use of said compounds for manufacturing a pharmaceutical composition for the treatment or prophylaxis of a disease, as well as to intermediate compounds useful in the preparation of said compounds.



(72) Inventeurs(suite)/Inventors(continued): SIEMEISTER, GERHARD, DE; SCOTT, WILLIAM, DE

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 December 2011 (08.12.2011)

(10) International Publication Number
WO 2011/151259 A1

(51) International Patent Classification:
C07D 487/04 (2006.01) *A61K 31/4985* (2006.01)

(21) International Application Number:
PCT/EP2011/058713

(22) International Filing Date:
27 May 2011 (27.05.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10164587.7 1 June 2010 (01.06.2010) EP

(71) Applicant (for all designated States except US): **BAYER PHARMA AKTIENGESELLSCHAFT** [DE/DE]; Müller Strasse 178, 13353 Berlin (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KLAR, Ulrich** [DE/DE]; Isegrimsteig 8a, 13503 Berlin (DE). **KOPPITZ, Marcus** [DE/DE]; Scharnhorststr. 28, 10115 Berlin (DE). **JAUTELAT, Rolf** [DE/DE]; Zur Pumpstation 35, 42781 Haan (DE). **KOSEMUND, Dirk** [DE/DE]; Klutstr. 14, 12587 Berlin (DE). **BOHLMANN, Rolf** [DE/DE]; Kuehler Weg 6a, 14055 Berlin (DE). **BADER, Benjamin** [DE/DE]; Hillmannstr. 14, 13467 Berlin (DE). **LIENAU, Philip** [DE/DE]; Pücklerstr. 33, 10997 Berlin (DE). **SIEMEISTER, Gerhard** [DE/DE]; Reimerswalder Steig 26, 13503 Berlin (DE). **SCOTT, William** [US/DE]; Alsterweg 50, 14167 Berlin (DE).

(74) Common Representative: **BAYER PHARMA AKTIENGESELLSCHAFT**; Law and Patents, Patents and Licensing, Müller Strasse 178, 13353 Berlin (DE).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: SUBSTITUTED IMIDAZOPYRAZINES



(I)

(57) Abstract: The present invention relates to substituted imidazopyrazine compounds of general formula (I) in which A, R¹, R³ and R⁵ are as given in the description and in the claims, to methods of preparing said compounds, to pharmaceutical compositions and combinations comprising said compounds, to the use of said compounds for manufacturing a pharmaceutical composition for the treatment or prophylaxis of a disease, as well as to intermediate compounds useful in the preparation of said compounds.

WO 2011/151259 A1

SUBSTITUTED IMIDAZOPYRAZINES

The present invention relates to substituted imidazopyrazine compounds of general formula (I) as described and defined herein, to methods of preparing
5 said compounds, to pharmaceutical compositions and combinations comprising said compounds, to the use of said compounds for manufacturing a pharmaceutical composition for the treatment or prophylaxis of a disease, as well as to intermediate compounds useful in the preparation of said compounds.

10

BACKGROUND OF THE INVENTION

The present invention relates to chemical compounds that inhibit Mps-1 (Monopolar Spindle 1) kinase (also known as Tyrosine Threonine Kinase, TTK).

15 Mps-1 is a dual specificity Ser/Thr kinase which plays a key role in the activation of the mitotic checkpoint (also known as spindle checkpoint, spindle assembly checkpoint) thereby ensuring proper chromosome segregation during mitosis [Abrieu A *et al.*, Cell, 2001, 106, 83-93]. Every dividing cell has to ensure equal separation of the replicated chromosomes into the two daughter
20 cells. Upon entry into mitosis, chromosomes are attached at their kinetochores to the microtubules of the spindle apparatus. The mitotic checkpoint is a surveillance mechanism that is active as long as unattached kinetochores are present and prevents mitotic cells from entering anaphase and thereby completing cell division with unattached chromosomes
25 [Suijkerbuijk SJ and Kops GJ, Biochemica et Biophysica Acta, 2008, 1786, 24-31; Musacchio A and Salmon ED, Nat Rev Mol Cell Biol., 2007, 8, 379-93]. Once all kinetochores are attached in a correct amphitelic, *i.e.* bipolar, fashion with the mitotic spindle, the checkpoint is satisfied and the cell enters anaphase and proceeds through mitosis. The mitotic checkpoint consists of
30 complex network of a number of essential proteins, including members of the

MAD (mitotic arrest deficient, MAD 1-3) and Bub (Budding uninhibited by benzimidazole, Bub 1-3) families, the motor protein CENP-E, Mps-1 kinase as well as other components, many of these being over-expressed in proliferating cells (e.g. cancer cells) and tissues [Yuan B *et al.*, Clinical Cancer Research, 5 2006, 12, 405-10]. The essential role of Mps-1 kinase activity in mitotic checkpoint signalling has been shown by shRNA-silencing, chemical genetics as well as chemical inhibitors of Mps-1 kinase [Jelluma N *et al.*, PLoS ONE, 2008, 3, e2415; Jones MH *et al.*, Current Biology, 2005, 15, 160-65; Dorer RK *et al.*, Current Biology, 2005, 15, 1070-76; Schmidt M *et al.*, EMBO Reports, 2005, 6, 10 866-72].

There is ample evidence linking reduced but incomplete mitotic checkpoint function with aneuploidy and tumourigenesis [Weaver BA and Cleveland DW, Cancer Research, 2007, 67, 10103-5; King RW, Biochimica et Biophysica Acta, 2008, 1786, 4-14]. In contrast, complete inhibition of the mitotic checkpoint 15 has been recognised to result in severe chromosome missegregation and induction of apoptosis in tumour cells [Kops GJ *et al.*, Nature Reviews Cancer, 2005, 5, 773-85; Schmidt M and Medema RH, Cell Cycle, 2006, 5, 159-63; Schmidt M and Bastians H, Drug Resistance Updates, 2007, 10, 162-81]. Therefore, mitotic checkpoint abrogation through pharmacological inhibition 20 of Mps-1 kinase or other components of the mitotic checkpoint represents a new approach for the treatment of proliferative disorders including solid tumours such as carcinomas and sarcomas and leukaemias and lymphoid malignancies or other disorders associated with uncontrolled cellular proliferation.

25

Established anti-mitotic drugs such as vinca alkaloids, taxanes or epothilones activate the SAC inducing a mitotic arrest either by stabilising or destabilising microtubule dynamics. This arrest prevents separation of sister chromatids to form the two daughter cells. Prolonged arrest in mitosis forces a cell either

into mitotic exit without cytokinesis or into mitotic catastrophe leading to cell death.

In contrast, inhibitors of Mps-1 induce a SAC inactivation that accelerates progression of cells through mitosis resulting in severe chromosomal
5 missegregation and finally in cell death.

These findings suggest that Mps-1 inhibitors should be of therapeutic value for the treatment of proliferative disorders associated with enhanced uncontrolled proliferative cellular processes such as, for example, cancer, inflammation,
10 arthritis, viral diseases, neurodegenerative diseases such as Alzheimer's disease, cardiovascular diseases, or fungal diseases in a warm-blooded animal such as man.

Therefore, inhibitors of Mps-1 represent valuable compounds that should
15 complement therapeutic options either as single agents or in combination with other drugs.

Different compounds have been disclosed in prior art which show inhibitory effect on Mps-1 kinase. WO2010/124826A1 discloses substituted
20 imidazoquinoxaline compounds as inhibitors of Mps-1 kinase or TTK. WO2011/026579A1 discloses substituted aminoquinoxalines as Mps-1 inhibitors.

Protein kinase inhibitors based on substituted imidazo[1,2-a]pyrazin-8-amine compounds are well known and are claimed to specifically inhibit CDK1, CDK2,
25 MAPK, ERK, GSK3 β , JNK, CHK-1, CHK-2, VEGF-R2, EGFR, HER2, SRC, JAK, TEK, Aurora, MK1, MK2, AKT, Pim-1, Abl and Src kinases (see US 2004/0220189A1, WO 2004/026877A1, US 2005/0009832A1, US 2006/0106023A1, US 2007/0105864A1, WO 2007/056468A1, WO 2007/058942A2, WO 2007/131991A1, WO2007/145921A1, WO 2008/030795A2, WO 2008/079460A2, W O

2008/057512A2, WO 2008/082490A2, WO 2009/017701A2, WO 2009/097233A1, US 2009/0175852A1).


WO 2009/024585 A2 discloses imidazo[1,2-a]pyrazin-8-amine compounds for
5 treating infections by hepatitis C virus and other viruses.

WO 2010/069684 A1 discloses compounds for the treatment or prophylaxis of muscular dystrophy. Among the disclosed compounds there are also imidazo[1,2-a]pyrazin-8-amine compounds.

10

However, the state of the art described above does not describe the substituted imidazopyrazine compounds of general formula (I) of the present invention, or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same, as described and defined herein, and as
15 hereinafter referred to as “compounds of the present invention”, or their pharmacological activity.

WO 2011/013729 A1 discloses fused imidazole derivatives as Mps-1 inhibitors. Among the disclosed fused imidazole derivatives there are also imidazo[1,2-
20 a]pyrazin-8-amine compounds. For example, WO 2011/013729 A1 discloses compounds of general formula C1:



25

in which (X, Y, V, W) is (-N=, =CR¹-, =N-, -CR⁷=), (-CR²=, =N-, =N-, -CR⁷=), (-N=, =CR¹-, =N-, -N=) or (-N=, =CR¹-, -O-, -N=) ;

R⁸ is substituted or unsubstituted cycloalkyl ;

5

Z is a group represented by formula -NR³R⁴ or a group represented by formula -OR⁵ ;

10 A is substituted or unsubstituted aromatic hydrocarbon ring, substituted or unsubstituted aromatic heterocyclic ring, substituted or unsubstituted non-aromatic hydrocarbon ring or substituted or unsubstituted non-aromatic heterocyclic ring ;

15 R¹, R³, R⁴, R⁵, and R⁶ represent a large variety of substituents (see WO2011/013729A1, e.g. claim 1). Inter alia, R⁶ represents substituted or unsubstituted alkynyl.

However, WO 2011/013729A1 does not specifically disclose the compounds of formula (I) of the present invention.

20


It has now been found, and this constitutes the basis of the present invention, that said compounds of the present invention have surprising and advantageous properties.

25 In particular, said compounds of the present invention have surprisingly been found to effectively inhibit Mps-1 kinase and may therefore be used for the treatment or prophylaxis of diseases of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses or diseases which are accompanied with
30 uncontrolled cell growth, proliferation and/or survival, inappropriate cellular


immune responses, or inappropriate cellular inflammatory responses, particularly in which the uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses is mediated by Mps-1 kinase, such as, for example, 5 haematological tumours, solid tumours, and/or metastases thereof, e.g. leukaemias and myelodysplastic syndrome, malignant lymphomas, head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung tumours, gastrointestinal tumours, endocrine tumours, mammary and other gynaecological tumours, 10 urological tumours including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

15 SUMMARY of the INVENTION


The present invention covers compounds of general formula (I) :



in which :



Z represents a $-\text{C}(=\text{O})\text{N}(\text{H})\text{R}^2$ or $-\text{S}(=\text{O})_2\text{N}(\text{H})\text{R}^2$ group,
or a group selected from :



wherein * indicates the point of attachment of said groups with the rest
of the molecule ;

10 R¹ represents a hydrogen atom or a halogen atom, or a $-\text{CN}$, $\text{C}_1\text{-C}_6\text{-alkyl-}$,
 $-\text{N}(\text{R}^{6a})\text{R}^{6b}$, $-\text{NO}_2$, $-\text{OR}^6$ or $-\text{SR}^6$ group ;

R² represents a hydrogen atom, or a $\text{C}_1\text{-C}_6\text{-alkyl-}$, halo- $\text{C}_1\text{-C}_6\text{-alkyl-}$,
 $\text{R}^{6a}(\text{R}^{6b})\text{N-}\text{C}_1\text{-C}_6\text{-alkyl-}$, $\text{HO-}\text{C}_1\text{-C}_6\text{-alkyl-}$, $-\text{C}_1\text{-C}_6\text{-alkyl-CN}$,
15 $\text{C}_1\text{-C}_6\text{-alkoxy-}\text{C}_1\text{-C}_6\text{-alkyl-}$, halo- $\text{C}_1\text{-C}_6\text{-alkoxy-}\text{C}_1\text{-C}_6\text{-alkyl-}$, $\text{C}_2\text{-C}_6\text{-alkenyl-}$,
 $\text{C}_2\text{-C}_6\text{-alkynyl-}$ or $\text{C}_3\text{-C}_6\text{-cycloalkyl-}$ group ; wherein said $\text{C}_3\text{-C}_6\text{-cycloalkyl-}$
group is optionally substituted, one or more times, identically or
differently, with a substituent selected from: halo-, $\text{C}_1\text{-C}_3\text{-alkyl-}$, cyano-,
 $\text{C}_1\text{-C}_3\text{-alkoxy-}$;

20

R³ represents a hydrogen atom, or a $-\text{CN}$, $\text{C}_1\text{-C}_6\text{-alkyl-}$,
 $-(\text{CH}_2)_m\text{-C}_2\text{-C}_6\text{-alkenyl}$, $-(\text{CH}_2)_m\text{-C}_2\text{-C}_6\text{-alkynyl}$, $-(\text{CH}_2)_m\text{-C}_3\text{-C}_6\text{-cycloalkyl}$,
 $-(\text{CH}_2)_m\text{-(3- to 7-membered heterocycloalkyl)}$, aryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$,
heteroaryl- $\text{C}_1\text{-C}_6\text{-alkyl-}$, halo- $\text{C}_1\text{-C}_6\text{-alkyl-}$, $\text{R}^{6a}(\text{R}^{6b})\text{N-}\text{C}_1\text{-C}_6\text{-alkyl-}$,
25 halo- $\text{C}_1\text{-C}_6\text{-alkoxy-}\text{C}_1\text{-C}_6\text{-alkyl-}$, $\text{C}_3\text{-C}_6\text{-cycloalkyl-}$, 3- to 7-membered
heterocycloalkyl-, $\text{C}_2\text{-C}_6\text{-alkenyl-}$, $\text{C}_3\text{-C}_6\text{-cycloalkenyl-}$, $\text{C}_2\text{-C}_6\text{-alkynyl-}$,

aryl-, C₁-C₆-alkyl-aryl-, heteroaryl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
-C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂ or -N(H)C(=O)R⁶ group ;

5 said C₁-C₆-alkyl-, -(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
group

being optionally substituted, identically or differently, with 1, 2, 3, or 4
R⁷ groups ;

10

R^{4a}, R^{4b}, R^{4c}, R^{4d}

represent, independently from each other, a hydrogen or halogen atom,

or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,

R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-,

15 halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b},

-C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶,

-N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶,

-N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶,

-N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b},

20 -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶,

-S(=O)₂N(H)R^{6a}, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;

R⁵ represents a hydrogen atom, or a C₁-C₆-alkyl-, C₂-C₆-alkenyl-,

-(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,

25 -(CH₂)_m-(3- to 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-,

heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,

HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN, C₁-C₆-alkoxy-C₁-C₆-alkyl-,

halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered

heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-,

30 aryl- or heteroaryl- group ;

s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl-,
 -(CH₂)_n-C₂-C₆-alkynyl-, -(CH₂)_m-C₃-C₆-cycloalkyl-, -(CH₂)_m-(3- to
 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
 5 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
 C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group
 being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R⁸ groups ;

10

R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a
 C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to
 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl- or
 15 heteroaryl-C₁-C₆-alkyl- group ;

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl, 3-
 20 to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 25 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-,
 30 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,

C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 5 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;

10

m is an integer of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 ;

and

n is an integer of 1, 2, 3, 4, 5, 6 or 7 ;

15

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

The present invention further relates to methods of preparing compounds of
 20 general formula (I), to pharmaceutical compositions and combinations
 comprising said compounds, to the use of said compounds for manufacturing a
 pharmaceutical composition for the treatment or prophylaxis of a disease, as
 well as to intermediate compounds useful in the preparation of said
 compounds.

25

DETAILED DESCRIPTION of the INVENTION

The terms as mentioned in the present text have preferably the following
 30 meanings :

The term “halogen atom” or “halo-” is to be understood as meaning a fluorine, chlorine, bromine or iodine atom.

The term “C₁-C₆-alkyl” is to be understood as preferably meaning a linear or
 5 branched, saturated, monovalent hydrocarbon group having 1, 2, 3, 4, 5, or 6
 carbon atoms, *e.g.* a methyl, ethyl, propyl, butyl, pentyl, hexyl, iso-propyl,
 iso-butyl, sec-butyl, tert-butyl, iso-pentyl, 2-methylbutyl, 1-methylbutyl,
 1-ethylpropyl, 1,2-dimethylpropyl, neo-pentyl, 1,1-dimethylpropyl,
 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl,
 10 2-ethylbutyl, 1-ethylbutyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl,
 1,1-dimethylbutyl, 2,3-dimethylbutyl, 1,3-dimethylbutyl, or 1,2-dimethylbutyl
 group, or an isomer thereof. Particularly, said group has 1, 2, 3 or 4 carbon
 atoms (“C₁-C₄-alkyl”), *e.g.* a methyl, ethyl, propyl, butyl, iso-propyl, iso-butyl,
 sec-butyl, tert-butyl group, more particularly 1, 2 or 3 carbon atoms
 15 (“C₁-C₃-alkyl”), *e.g.* a methyl, ethyl, n-propyl- or iso-propyl group.

The term “halo-C₁-C₆-alkyl” is to be understood as preferably meaning a linear
 or branched, saturated, monovalent hydrocarbon group in which the term
 “C₁-C₆-alkyl” is defined *supra*, and in which one or more hydrogen atoms is
 20 replaced by a halogen atom, in identically or differently, *i.e.* one halogen
 atom being independent from another. Particularly, said halogen atom is F.
 Said halo-C₁-C₆-alkyl group is, for example, -CF₃, -CHF₂, -CH₂F, -CF₂CF₃, or
 -CH₂CF₃.

25 The term “C₁-C₆-alkoxy” is to be understood as preferably meaning a linear or
 branched, saturated, monovalent, hydrocarbon group of formula -O-alkyl, in
 which the term “alkyl” is defined *supra*, *e.g.* a methoxy, ethoxy, n-propoxy,
 iso-propoxy, n-butoxy, iso-butoxy, tert-butoxy, sec-butoxy, pentoxy,
 iso-pentoxy, or n-hexoxy group, or an isomer thereof.

30

The term “halo-C₁-C₆-alkoxy” is to be understood as preferably meaning a linear or branched, saturated, monovalent C₁-C₆-alkoxy group, as defined *supra*, in which one or more of the hydrogen atoms is replaced, in identically or differently, by a halogen atom. Particularly, said halogen atom is F. Said
 5 halo-C₁-C₆-alkoxy group is, for example, -OCF₃, -OCHF₂, -OCH₂F, -OCF₂CF₃, or -OCH₂CF₃.

The term “C₁-C₆-alkoxy-C₁-C₆-alkyl” is to be understood as preferably meaning a linear or branched, saturated, monovalent alkyl group, as defined *supra*, in
 10 which one or more of the hydrogen atoms is replaced, in identically or differently, by a C₁-C₆-alkoxy group, as defined *supra*, e.g. methoxyalkyl, ethoxyalkyl, propyloxyalkyl, iso-propoxyalkyl, butoxyalkyl, iso-butoxyalkyl, tert-butoxyalkyl, sec-butoxyalkyl, pentyloxyalkyl, iso-pentyloxyalkyl, hexyloxyalkyl group, in which the term “C₁-C₆-alkyl” is defined *supra*, or an
 15 isomer thereof.

The term “halo-C₁-C₆-alkoxy-C₁-C₆-alkyl” is to be understood as preferably meaning a linear or branched, saturated, monovalent C₁-C₆-alkoxy-C₁-C₆-alkyl group, as defined *supra*, in which one or more of the hydrogen atoms is
 20 replaced, in identically or differently, by a halogen atom. Particularly, said halogen atom is F. Said halo-C₁-C₆-alkoxy-C₁-C₆-alkyl group is, for example, -CH₂CH₂OCF₃, -CH₂CH₂OCHF₂, -CH₂CH₂OCH₂F, -CH₂CH₂OCF₂CF₃, or -CH₂CH₂OCH₂CF₃.

25 The term “C₂-C₆-alkenyl” is to be understood as preferably meaning a linear or branched, monovalent hydrocarbon group, which contains one or more double bonds, and which has 2, 3, 4, 5 or 6 carbon atoms, particularly 2 or 3 carbon atoms (“C₂-C₃-alkenyl”), it being understood that in the case in which said alkenyl group contains more than one double bond, then said double bonds
 30 may be isolated from, or conjugated with, each other. Said alkenyl group is,

for example, a vinyl, allyl, (E)-2-methylvinyl, (Z)-2-methylvinyl, homoallyl, (E)-but-2-enyl, (Z)-but-2-enyl, (E)-but-1-enyl, (Z)-but-1-enyl, pent-4-enyl, (E)-pent-3-enyl, (Z)-pent-3-enyl, (E)-pent-2-enyl, (Z)-pent-2-enyl, (E)-pent-1-enyl, (Z)-pent-1-enyl, hex-5-enyl, (E)-hex-4-enyl, (Z)-hex-4-enyl, (E)-hex-3-enyl, (Z)-hex-3-enyl, (E)-hex-2-enyl, (Z)-hex-2-enyl, (E)-hex-1-enyl, (Z)-hex-1-enyl, isopropenyl, 2-methylprop-2-enyl, 1-methylprop-2-enyl, 2-methylprop-1-enyl, (E)-1-methylprop-1-enyl, (Z)-1-methylprop-1-enyl, 3-methylbut-3-enyl, 2-methylbut-3-enyl, 1-methylbut-3-enyl, 3-methylbut-2-enyl, (E)-2-methylbut-2-enyl, (Z)-2-methylbut-2-enyl, (E)-1-methylbut-2-enyl, (Z)-1-methylbut-2-enyl, (E)-3-methylbut-1-enyl, (Z)-3-methylbut-1-enyl, (E)-2-methylbut-1-enyl, (Z)-2-methylbut-1-enyl, (E)-1-methylbut-1-enyl, (Z)-1-methylbut-1-enyl, 1,1-dimethylprop-2-enyl, 1-ethylprop-1-enyl, 1-propylvinyl, 1-isopropylvinyl, 4-methylpent-4-enyl, 3-methylpent-4-enyl, 2-methylpent-4-enyl, 1-methylpent-4-enyl, 4-methylpent-3-enyl, (E)-3-methylpent-3-enyl, (Z)-3-methylpent-3-enyl, (E)-2-methylpent-3-enyl, (Z)-2-methylpent-3-enyl, (E)-1-methylpent-3-enyl, (Z)-1-methylpent-3-enyl, (E)-4-methylpent-2-enyl, (Z)-4-methylpent-2-enyl, (E)-3-methylpent-2-enyl, (Z)-3-methylpent-2-enyl, (E)-2-methylpent-2-enyl, (Z)-2-methylpent-2-enyl, (E)-1-methylpent-2-enyl, (Z)-1-methylpent-2-enyl, (E)-4-methylpent-1-enyl, (Z)-4-methylpent-1-enyl, (E)-3-methylpent-1-enyl, (Z)-3-methylpent-1-enyl, (E)-2-methylpent-1-enyl, (Z)-2-methylpent-1-enyl, (E)-1-methylpent-1-enyl, (Z)-1-methylpent-1-enyl, 3-ethylbut-3-enyl, 2-ethylbut-3-enyl, 1-ethylbut-3-enyl, (E)-3-ethylbut-2-enyl, (Z)-3-ethylbut-2-enyl, (E)-2-ethylbut-2-enyl, (Z)-2-ethylbut-2-enyl, (E)-1-ethylbut-2-enyl, (Z)-1-ethylbut-2-enyl, (E)-3-ethylbut-1-enyl, (Z)-3-ethylbut-1-enyl, 2-ethylbut-1-enyl, (E)-1-ethylbut-1-enyl, (Z)-1-ethylbut-1-enyl, 2-propylprop-2-enyl, 1-propylprop-2-enyl, 2-isopropylprop-2-enyl, 1-isopropylprop-2-enyl, (E)-2-propylprop-1-enyl, (Z)-2-propylprop-1-enyl, (E)-1-propylprop-1-enyl, (Z)-1-propylprop-1-enyl, (E)-2-isopropylprop-1-enyl, (Z)-2-isopropylprop-1-enyl,

(E)-1-isopropylprop-1-enyl, (Z)-1-isopropylprop-1-enyl,
 (E)-3,3-dimethylprop-1-enyl, (Z)-3,3-dimethylprop-1-enyl,
 1-(1,1-dimethylethyl)ethenyl, buta-1,3-dienyl, penta-1,4-dienyl,
 hexa-1,5-dienyl, or methylhexadienyl group. Particularly, said group is vinyl or
 5 allyl.

The term “C₂-C₆-alkynyl” is to be understood as preferably meaning a linear or branched, monovalent hydrocarbon group which contains one or more triple bonds, and which contains 2, 3, 4, 5 or 6 carbon atoms, particularly 2 or 3
 10 carbon atoms (“C₂-C₃-alkynyl”). Said C₂-C₆-alkynyl group is, for example, ethynyl, prop-1-ynyl, prop-2-ynyl, but-1-ynyl, but-2-ynyl, but-3-ynyl, pent-1-ynyl, pent-2-ynyl, pent-3-ynyl, pent-4-ynyl, hex-1-ynyl, hex-2-ynyl, hex-3-ynyl, hex-4-ynyl, hex-5-ynyl, 1-methylprop-2-ynyl, 2-methylbut-3-ynyl, 1-methylbut-3-ynyl, 1-methylbut-2-ynyl, 3-methylbut-1-ynyl,
 15 1-ethylprop-2-ynyl, 3-methylpent-4-ynyl, 2-methylpent-4-ynyl, 1-methylpent-4-ynyl, 2-methylpent-3-ynyl, 1-methylpent-3-ynyl, 4-methylpent-2-ynyl, 1-methylpent-2-ynyl, 4-methylpent-1-ynyl, 3-methylpent-1-ynyl, 2-ethylbut-3-ynyl, 1-ethylbut-3-ynyl, 1-ethylbut-2-ynyl, 1-propylprop-2-ynyl, 1-isopropylprop-2-ynyl, 2,2-dimethylbut-3-ynyl, 1,1-dimethylbut-3-ynyl,
 20 1,1-dimethylbut-2-ynyl, or 3,3-dimethylbut-1-ynyl group. Particularly, said alkynyl group is ethynyl, prop-1-ynyl, or prop-2-ynyl.

The term “C₃-C₆-cycloalkyl” is to be understood as preferably meaning a saturated, monovalent, mono-, or bicyclic hydrocarbon ring which contains 3,
 25 4, 5 or 6 carbon atoms (“C₃-C₆-cycloalkyl”). Said C₃-C₆-cycloalkyl group is for example, a monocyclic hydrocarbon ring, e.g. a cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl or a bicyclic hydrocarbon ring. Said cycloalkyl ring can optionally contain one or more double bonds e.g. cycloalkenyl, such as a cyclopropenyl, cyclobutenyl, cyclopentenyl or cyclohexenyl group, wherein

the bond between said ring with the rest of the molecule may be to any carbon atom of said ring, be it saturated or unsaturated.

The term “3- to 7-membered heterocycloalkyl”, is to be understood as
 5 meaning a saturated or partially unsaturated, monovalent, mono- or bicyclic hydrocarbon ring which contains 2, 3, 4, 5, or 6 carbon atoms, and one or more heteroatom-containing groups selected from C(=O), O, S, S(=O), S(=O)₂, NR^a, in which R^a represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, H₂N-C₁-C₆-alkyl-, C₁-C₆-alkyl(H)N-C₁-C₆-alkyl-, (C₁-C₆-alkyl)₂N-C₁-C₆-alkyl-,
 10 HO-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl-, -C(=O)-C₁-C₆-alkyl, -C(=O)O-C₁-C₆-alkyl, C₃-C₇-cycloalkyl-, -C(=O)NH₂, -C(=O)N(H)C₁-C₆-alkyl, -C(=O)N(C₁-C₆-alkyl)₂, -S(=O)(=NC₁-C₆-alkyl), -S(=O)N(C₁-C₆-alkyl)₂, -S(=O)₂N(C₁-C₆-alkyl)₂, -S(=O)₂C₁-C₆-alkyl group ; it being possible for said heterocycloalkyl group to be attached to the rest of the molecule via any one
 15 of the carbon atoms or, if present, the nitrogen atom.

Particularly, said 3- to 7-membered heterocycloalkyl can contain 2, 3, 4, or 5 carbon atoms, and one or more of the above-mentioned heteroatom-containing groups (a “3- to 6-membered heterocycloalkyl”), more particularly said
 20 heterocycloalkyl can contain 4 or 5 carbon atoms, and one or more of the above-mentioned heteroatom-containing groups (a “5- to 6-membered heterocycloalkyl”).

Particularly, without being limited thereto, said heterocycloalkyl can be a
 25 4-membered ring, such as an azetidiny, oxetanyl, or a 5-membered ring, such as tetrahydrofuranyl, dioxolinyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, pyrrolinyl, or a 6-membered ring, such as tetrahydropyranyl, piperidinyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, or trithianyl, or a 7-membered ring, such as a diazepanyl ring, for example. Optionally, said
 30 heterocycloalkyl can contain one or more double bonds, e.g. 4H-pyranyl,

2H-pyranyl, 3H-diazirinyl, 2,5-dihydro-1H-pyrrolyl, [1,3]dioxolyl, 4H-[1,3,4]thiadiazinyl, 2,5-dihydrofuranyl, 2,3-dihydrofuranyl, 2,5-dihydrothiophenyl, 2,3-dihydrothiophenyl, 4,5-dihydrooxazolyl, or 4H-[1,4]thiazinyl group, or, it may be benzo fused.

5

Said heterocyclyl can be bicyclic, such as, without being limited thereto, a 5,5-membered ring, *e.g.* a hexahydrocyclopenta[c]pyrrol-2(1H)-yl ring, or a 5,6-membered bicyclic ring, *e.g.* a hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl ring, or for example.

10

As mentioned *supra*, said nitrogen atom-containing ring can be partially unsaturated, *i.e.* it can contain one or more double bonds, such as, without being limited thereto, a 2,5-dihydro-1H-pyrrolyl, 4H-[1,3,4]thiadiazinyl, 4,5-dihydrooxazolyl, or 4H-[1,4]thiazinyl ring, for example, or, it may be benzo-fused, such as, without being limited thereto, a dihydroisoquinolinyl ring, for example.

15

The term “aryl” is to be understood as preferably meaning a monovalent, aromatic or partially aromatic, mono-, or bi- or tricyclic hydrocarbon ring having 6, 7, 8, 9, 10, 11, 12, 13 or 14 carbon atoms (a “C₆-C₁₄-aryl” group), particularly a ring having 6 carbon atoms (a “C₆-aryl” group), *e.g.* a phenyl group, or a biphenyl group, or a ring having 9 carbon atoms (a “C₉-aryl” group), *e.g.* an indanyl or indenyl group, or a ring having 10 carbon atoms (a “C₁₀-aryl” group), *e.g.* a tetralinyl, dihydronaphthyl, or naphthyl group, or a ring having 13 carbon atoms, (a “C₁₃-aryl” group), *e.g.* a fluorenyl group, or a ring having 14 carbon atoms, (a “C₁₄-aryl” group), *e.g.* an anthranlyl group.

20

25

The term “heteroaryl” is understood as preferably meaning a monovalent, aromatic, mono- or bicyclic aromatic ring system having 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 ring atoms (a “5- to 14-membered heteroaryl” group), particularly

30

5 or 6 or 9 or 10 atoms, and which contains at least one heteroatom which may be identical or different, said heteroatom being such as oxygen, nitrogen or sulfur, and can be monocyclic, bicyclic, or tricyclic, and in addition in each case can be benzocondensed. Particularly, heteroaryl is selected from thienyl, furanyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, thia-4H-pyrazolyl *etc.*, and benzo derivatives thereof, such as, for example, benzofuranyl, benzothienyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, benzotriazolyl, indazolyl, indolyl, isoindolyl, *etc.*; or pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, *etc.*, and benzo derivatives thereof, such as, for example, quinolinyl, quinazolinyl, isoquinolinyl, *etc.*; or azocinyl, indolizinyll, purinyl, *etc.*, and benzo derivatives thereof; or cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthpyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, xanthenyl, or oxepinyl, *etc.* More particularly, heteroaryl is selected from pyridyl, benzofuranyl, benzisoxazolyl, indazolyl, quinazolinyl, thienyl, quinolinyl, benzothienyl, pyrazolyl, or furanyl.

In general, and unless otherwise mentioned, the heteroarylic or heteroarylenic radicals include all the possible isomeric forms thereof, e.g. the positional isomers thereof. Thus, for some illustrative non-restricting example, the term pyridinyl or pyridinylene includes pyridin-2-yl, pyridin-2-ylene, pyridin-3-yl, pyridin-3-ylene, pyridin-4-yl and pyridin-4-ylene; or the term thienyl or thienylene includes thien-2-yl, thien-2-ylene, thien-3-yl and thien-3-ylene.

The term “C₁-C₆”, as used throughout this text, e.g. in the context of the definition of “C₁-C₆-alkyl”, “C₁-C₆-haloalkyl”, “C₁-C₆-alkoxy”, or “C₁-C₆-haloalkoxy” is to be understood as meaning an alkyl group having a finite number of carbon atoms of 1 to 6, *i.e.* 1, 2, 3, 4, 5, or 6 carbon atoms. It is to be understood further that said term “C₁-C₆” is to be interpreted as any sub-range comprised therein, e.g. C₁-C₆, C₂-C₅, C₃-C₄, C₁-C₂, C₁-C₃, C₁-C₄, C₁-C₅,

C₁-C₆; particularly C₁-C₂, C₁-C₃, C₁-C₄, C₁-C₅, C₁-C₆; more particularly C₁-C₄; in the case of "C₁-C₆-haloalkyl" or "C₁-C₆-haloalkoxy" even more particularly C₁-C₂.

- 5 Similarly, as used herein, the term "C₂-C₆", as used throughout this text, *e.g.* in the context of the definitions of "C₂-C₆-alkenyl" and "C₂-C₆-alkynyl", is to be understood as meaning an alkenyl group or an alkynyl group having a finite number of carbon atoms of 2 to 6, *i.e.* 2, 3, 4, 5, or 6 carbon atoms. It is to be understood further that said term "C₂-C₆" is to be interpreted as any sub-range
10 comprised therein, *e.g.* C₂-C₆, C₃-C₅, C₃-C₄, C₂-C₃, C₂-C₄, C₂-C₅; particularly C₂-C₃.

- Further, as used herein, the term "C₃-C₆", as used throughout this text, *e.g.* in the context of the definition of "C₃-C₆-cycloalkyl", is to be understood as
15 meaning a cycloalkyl group having a finite number of carbon atoms of 3 to 6, *i.e.* 3, 4, 5 or 6 carbon atoms. It is to be understood further that said term "C₃-C₆" is to be interpreted as any sub-range comprised therein, *e.g.* C₃-C₆, C₄-C₅, C₃-C₅, C₃-C₄, C₄-C₆, C₅-C₆; particularly C₃-C₆.

- 20 The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such
25 combinations result in stable compounds.

The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties.

Ring system substituent means a substituent attached to an aromatic or nonaromatic ring system which, for example, replaces an available hydrogen on the ring system.

- 5 As used herein, the term “one or more times”, e.g. in the definition of the substituents of the compounds of the general formulae of the present invention, is understood as meaning “one, two, three, four or five times, particularly one, two, three or four times, more particularly one, two or three times, even more particularly one or two times”.

10

Where the plural form of the word compounds, salts, polymorphs, hydrates, solvates and the like, is used herein, this is taken to mean also a single compound, salt, polymorph, isomer, hydrate, solvate or the like.

- 15 By “stable compound” or “stable structure” is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The compounds of this invention may contain one or more asymmetric centre,
20 depending upon the location and nature of the various substituents desired. Asymmetric carbon atoms may be present in the (R) or (S) configuration, resulting in racemic mixtures in the case of a single asymmetric centre, and diastereomeric mixtures in the case of multiple asymmetric centres. In certain instances, asymmetry may also be present due to restricted rotation about a
25 given bond, for example, the central bond adjoining two substituted aromatic rings of the specified compounds.

Substituents on a ring may also be present in either cis or trans form. It is intended that all such configurations (including enantiomers and diastereomers), are included within the scope of the present invention.

Preferred compounds are those which produce the more desirable biological activity. Separated, pure or partially purified isomers and stereoisomers or racemic or diastereomeric mixtures of the compounds of this invention are also included within the scope of the present invention. The purification and the
5 separation of such materials can be accomplished by standard techniques known in the art.

The optical isomers can be obtained by resolution of the racemic mixtures according to conventional processes, for example, by the formation of diastereoisomeric salts using an optically active acid or base or formation of
10 covalent diastereomers. Examples of appropriate acids are tartaric, diacetyltartaric, ditoluoyltartaric and camphorsulfonic acid. Mixtures of diastereoisomers can be separated into their individual diastereomers on the basis of their physical and/or chemical differences by methods known in the art, for example, by chromatography or fractional crystallisation. The
15 optically active bases or acids are then liberated from the separated diastereomeric salts. A different process for separation of optical isomers involves the use of chiral chromatography (e.g., chiral HPLC columns), with or without conventional derivatisation, optimally chosen to maximise the separation of the enantiomers. Suitable chiral HPLC columns are
20 manufactured by Diacel, e.g., Chiracel OD and Chiracel OJ among many others, all routinely selectable. Enzymatic separations, with or without derivatisation, are also useful. The optically active compounds of this invention can likewise be obtained by chiral syntheses utilizing optically active starting materials.

25


In order to limit different types of isomers from each other reference is made to IUPAC Rules Section E (Pure Appl Chem 45, 11-30, 1976).

The present invention includes all possible stereoisomers of the compounds of
30 the present invention as single stereoisomers, or as any mixture of said


stereoisomers, in any ratio. Isolation of a single stereoisomer, e.g. a single enantiomer or a single diastereomer, of a compound of the present invention may be achieved by any suitable state of the art method, such as chromatography, especially chiral chromatography, for example.

5


Further, the compounds of the present invention may exist as tautomers. For example, any compound of the present invention which contains a pyrazole moiety as a heteroaryl group for example can exist as a 1H tautomer, or a 2H tautomer, or even a mixture in any amount of the two tautomers, or a triazole moiety for example can exist as a 1H tautomer, a 2H tautomer, or a 4H tautomer, or even a mixture in any amount of said 1H, 2H and 4H tautomers, viz. :



1H-tautomer



2H-tautomer



4H-tautomer.

15

The present invention includes all possible tautomers of the compounds of the present invention as single tautomers, or as any mixture of said tautomers, in any ratio.

20 Further, the compounds of the present invention can exist as N-oxides, which are defined in that at least one nitrogen of the compounds of the present invention is oxidised. The present invention includes all such possible N-oxides.

The present invention also relates to useful forms of the compounds as disclosed herein, such as metabolites, hydrates, solvates, prodrugs, salts, in particular pharmaceutically acceptable salts, and co-precipitates.

25

The compounds of the present invention can exist as a hydrate, or as a solvate, wherein the compounds of the present invention contain polar solvents, in particular water, methanol or ethanol for example as structural element of the crystal lattice of the compounds. The amount of polar solvents, in particular
5 water, may exist in a stoichiometric or non-stoichiometric ratio. In the case of stoichiometric solvates, *e.g.* a hydrate, hemi-, (semi-), mono-, sesqui-, di-, tri-, tetra-, penta- *etc.* solvates or hydrates, respectively, are possible. The present invention includes all such hydrates or solvates.

10 Further, the compounds of the present invention can exist in free form, *e.g.* as a free base, or as a free acid, or as a zwitterion, or can exist in the form of a salt. Said salt may be any salt, either an organic or inorganic addition salt, particularly any pharmaceutically acceptable organic or inorganic addition salt, customarily used in pharmacy.

15

The term “pharmaceutically acceptable salt” refers to a relatively non-toxic, inorganic or organic acid addition salt of a compound of the present invention. For example, see S. M. Berge, *et al.* “Pharmaceutical Salts,” J. Pharm. Sci. 1977, 66: 1-19.

20

A suitable pharmaceutically acceptable salt of the compounds of the present invention may be, for example, an acid-addition salt of a compound of the present invention bearing a nitrogen atom, in a chain or in a ring, for example, which is sufficiently basic, such as an acid-addition salt with an inorganic acid,
25 such as hydrochloric, hydrobromic, hydroiodic, sulfuric, bisulfuric, phosphoric, or nitric acid, for example, or with an organic acid, such as formic, acetic, acetoacetic, pyruvic, trifluoroacetic, propionic, butyric, hexanoic, heptanoic, undecanoic, lauric, benzoic, salicylic, 2-(4-hydroxybenzoyl)-benzoic, camphoric, cinnamic, cyclopentanepropionic, digluconic,
30 3-hydroxy-2-naphthoic, nicotinic, pamoic, pectinic, persulfuric,

3-phenylpropionic, picric, pivalic, 2-hydroxyethanesulfonate, itaconic, sulfamic, trifluoromethanesulfonic, dodecylsulfuric, ethansulfonic, benzenesulfonic, para-toluenesulfonic, methansulfonic, 2-naphthalenesulfonic, naphthalinedisulfonic, camphorsulfonic acid, citric, tartaric, stearic, lactic, 5 oxalic, malonic, succinic, malic, adipic, alginic, maleic, fumaric, D-gluconic, mandelic, ascorbic, glucoheptanoic, glycerophosphoric, aspartic, sulfosalicylic, hemisulfuric, or thiocyanic acid, for example.

Further, another suitably pharmaceutically acceptable salt of a compound of the present invention which is sufficiently acidic, is an alkali metal salt, for 10 example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically acceptable cation, for example a salt with N-methyl-glucamine, dimethyl-glucamine, ethyl-glucamine, lysine, dicyclohexylamine, 1,6-hexadamine, ethanolamine, glucosamine, sarcosine, 15 serinol, tris-hydroxy-methyl-aminomethane, aminopropandiol, sovak-base, 1-amino-2,3,4-butantriol. Additionally, basic nitrogen containing groups may be quaternised with such agents as lower alkyl halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides ; dialkyl sulfates like dimethyl, diethyl, and dibutyl sulfate ; and diamyl sulfates, long chain halides 20 such as decyl, lauryl, myristyl and strearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

Those skilled in the art will further recognise that acid addition salts of the claimed compounds may be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods. 25 Alternatively, alkali and alkaline earth metal salts of acidic compounds of the invention are prepared by reacting the compounds of the invention with the appropriate base via a variety of known methods.


The present invention includes all possible salts of the compounds of the present invention as single salts, or as any mixture of said salts, in any ratio.

As used herein, the term “*in vivo* hydrolysable ester” is understood as meaning an *in vivo* hydrolysable ester of a compound of the present invention containing a carboxy or hydroxy group, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include for example alkyl, cycloalkyl and optionally substituted phenylalkyl, in particular benzyl esters, C₁-C₆ alkoxymethyl esters, e.g. methoxymethyl, C₁-C₆ alkanoyloxymethyl esters, e.g. pivaloyloxymethyl, phthalidyl esters, C₃-C₈ cycloalkoxy-carbonyloxy-C₁-C₆ alkyl esters, e.g. 1-cyclohexylcarbonyloxyethyl ; 1,3-dioxolen-2-onylmethyl esters, e.g. 5-methyl-1,3-dioxolen-2-onylmethyl ; and C₁-C₆-alkoxycarbonyloxyethyl esters, e.g. 1-methoxycarbonyloxyethyl, and may be formed at any carboxy group in the compounds of this invention.


An *in vivo* hydrolysable ester of a compound of the present invention containing a hydroxy group includes inorganic esters such as phosphate esters and [alpha]-acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of [alpha]-acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. The present invention covers all such esters.

Furthermore, the present invention includes all possible crystalline forms, or polymorphs, of the compounds of the present invention, either as single polymorphs, or as a mixture of more than one polymorphs, in any ratio.


In accordance with a first aspect, the present invention covers compounds of general formula (I) :



in which :



Z represents a -C(=O)N(H)R² or -S(=O)₂N(H)R² group,
or a group selected from :



wherein * indicates the point of attachment of said groups with the rest of the molecule ;

- R¹ represents a hydrogen atom or a halogen atom, or a -CN, C₁-C₆-alkyl-,
-N(R^{6a})R^{6b}, -NO₂, -OR⁶ or -SR⁶ group ;
- R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
5 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
group is optionally substituted, one or more times, identically or
differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
10 C₁-C₃-alkoxy- ;
- R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
-(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,
-(CH₂)_m-(3- to 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-,
15 heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-,
aryl-, C₁-C₆-alkyl-aryl-, heteroaryl, -C(=O)R⁶, -C(=O)N(H)R^{6a},
-C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶ group ;
20 s a i d C₁-C₆-alkyl-, -(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
group
25 being optionally substituted, identically or differently, with 1, 2, 3, or 4
R⁷ groups ;

R^{4a}, R^{4b}, R^{4c}, R^{4d}

- represent, independently from each other, a hydrogen or halogen atom,
30 o r a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,

- $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$, H $O-C_1-C_6\text{-alkyl-}$, $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
 $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$, $-C(=O)N(R^{6a})R^{6b}$,
 $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$, $-N(R^{6c})C(=O)R^6$,
 $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$, $-N(H)C(=O)OR^6$,
5 $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$, $-N(H)S(=O)_2R^6$,
 $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$, $-O(C=O)N(R^{6a})R^{6b}$,
 $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$, $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$,
 $-S(=O)_2N(H)R^{6a}$, $-S(=O)_2N(R^{6a})R^{6b}$ or $-S(=O)(=NR^{6c})R^6$ group ;
- 10 R^5 represents a hydrogen atom, or a $C_1-C_6\text{-alkyl-}$, $C_2-C_6\text{-alkenyl-}$,
 $-(CH_2)_n-C_2-C_6\text{-alkenyl-}$, $-(CH_2)_n-C_2-C_6\text{-alkynyl-}$, $-(CH_2)_m-C_3-C_6\text{-cycloalkyl-}$,
 $-(CH_2)_m\text{-(3- to 7-membered heterocycloalkyl)}$, $aryl-C_1-C_6\text{-alkyl-}$,
 $heteroaryl-C_1-C_6\text{-alkyl-}$, $halo-C_1-C_6\text{-alkyl-}$, $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$,
 $HO-C_1-C_6\text{-alkyl-}$, $-C_1-C_6\text{-alkyl-CN}$, $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
15 $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $C_3-C_6\text{-cycloalkyl-}$, 3- to 7-membered
 $heterocycloalkyl-$, $C_2-C_6\text{-alkenyl-}$, $C_3-C_6\text{-cycloalkenyl-}$, $C_2-C_6\text{-alkynyl-}$,
 $aryl-$ or $heteroaryl-$ group ;
- as a i d $C_1-C_6\text{-alkyl-}$, $C_2-C_6\text{-alkenyl-}$, $-(CH_2)_n-C_2-C_6\text{-alkenyl-}$,
 $-(CH_2)_n-C_2-C_6\text{-alkynyl-}$, $-(CH_2)_m-C_3-C_6\text{-cycloalkyl-}$, $-(CH_2)_m\text{-(3- to}$
20 $7\text{-membered heterocycloalkyl)}$, $aryl-C_1-C_6\text{-alkyl-}$, $heteroaryl-C_1-C_6\text{-alkyl-}$,
 $halo-C_1-C_6\text{-alkyl-}$, $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$, $HO-C_1-C_6\text{-alkyl-}$, $-C_1-C_6\text{-alkyl-CN}$,
 $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
 $C_3-C_6\text{-cycloalkyl-}$, 3- to 7-membered $heterocycloalkyl-$, $C_2-C_6\text{-alkenyl-}$,
 $C_3-C_6\text{-cycloalkenyl-}$, $C_2-C_6\text{-alkynyl-}$, $aryl-$ or $heteroaryl-$ group
25 being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R^8 groups ;

R^6 , R^{6a} , R^{6b} , R^{6c} ,

- represent, independently from each other, a hydrogen atom, or a
30 $C_1-C_6\text{-alkyl-}$, $HO-C_1-C_6\text{-alkyl-}$, $C_3-C_6\text{-cycloalkyl-}$, $C_2-C_6\text{-alkenyl-}$, 3- to

7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or heteroaryl-C₁-C₆-alkyl- group ;

5 R^7 represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, $R^{6a}(R^{6b})N$ -C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, 10 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;

15 R^8 represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, $R^{6a}(R^{6b})N$ -C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, 20 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, 25 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;


m is an integer of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 ;

and

30 n is an integer of 1, 2, 3, 4, 5, 6 or 7.


In a preferred embodiment, the invention relates to compounds of formula (I), wherein :

5 A represents a



In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

10 A represents a




In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

15

Z represents a -C(=O)N(H)R² group.

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

20 Z represents a



In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

R¹ represents a hydrogen atom.

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

- 5 R² represents a substituted or unsubstituted C₃-C₆-cycloalkyl- group.
 Preferably, R² represents a cyclopropyl- group, wherein said cyclopropyl- group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-, C₁-C₃-alkoxy-.

10

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

- R² represents a C₁-C₄-alkyl- group ; wherein said C₁-C₄-alkyl- group is optionally substituted, one or more times, identically or differently,
 15 with a substituent selected from: halo-, cyano-, C₁-C₃-alkoxy-.
 Preferably, R² represents a substituted or unsubstituted methyl-, ethyl-, n-propyl- or iso-propyl- group. More preferably, R² represents a substituted or unsubstituted methyl- or ethyl- group.

- 20 In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

- R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
 -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl),
 aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
 25 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3-
 to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
 group ;
 said C₁-C₆-alkyl-, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered
 heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
 30 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,

C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁷ groups.

5 In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

R³ represents a hydrogen atom, or a C₁-C₆-alkyl-, -(CH₂)_m-C₃-C₆-cycloalkyl-, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
 10 halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl- group ;
 said C₁-C₆-alkyl-, -(CH₂)_m-C₃-C₆-cycloalkyl-, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 15 C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁷ groups.

20 In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

R^{4a}, R^{4b}, R^{4c}, R^{4d}
 represent, independently from each other, a hydrogen or halogen atom,
 or a -CN, C₁-C₆-alkyl, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,
 25 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl- or -OR⁶ group.

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

30 R^{4a}, R^{4b}, R^{4c}, R^{4d} :

two of which represent a hydrogen atom, while the remaining two represent, independently from each other, a hydrogen or halogen atom, or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, -OR⁶ group.

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

(R^{4a}, R^{4b}, R^{4c}, R^{4d})

represent either (H, H, H, H) or (H, C₁-C₃-alkyl-, H, H) or (H, H, C₁-C₃-alkyl-, H). Preferably the C₁-C₃-alkyl- group is a methyl- group.

In another preferred embodiment, the invention relates to compounds of formula (I), wherein :

R⁵ represents a hydrogen atom, or a C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₃-alkynyl-, -(CH₂)_m-C₃-C₆-cycloalkyl-, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group ;

s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₃-alkynyl-, -(CH₂)_m-C₃-C₆-cycloalkyl-, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-,

C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁸ groups.

In another preferred embodiment, the invention relates to compounds of
5 formula (I), wherein :

R⁵ represents a C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-,
R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, C₁-C₃-alkoxy-C₁-C₃-alkyl- or
C₃-C₆-cycloalkyl- group ;
10 said C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-,
R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, C₁-C₃-alkoxy-C₁-C₃-alkyl- or
C₃-C₆-cycloalkyl- group being optionally substituted, identically or
differently, with 1 or 2 R⁸ groups.

15

In another preferred embodiment, the invention relates to compounds of
formula (I), wherein :

m is an integer of 0, 1, 2, 3, or 4.


20 In another preferred embodiment, the invention relates to compounds of
formula (I), wherein :

n is an integer of 1, 2, 3 or 4.


It is to be understood that the present invention relates also to any
25 combination of the preferred embodiments described above.

Some examples of combinations are given hereinafter. However, the invention
shall not be limited to these combinations.

In another preferred embodiment, the invention relates to compounds of
30 formula (I), *supra*, in which :



Z represents a $-C(=O)N(H)R^2$ or $-S(=O)_2N(H)R^2$ group,
 5 or a group selected from :



wherein * indicates the point of attachment of said groups with the rest
 10 of the molecule ;

R^1 represents a hydrogen atom;

R^2 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-,
 15 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO - C_1 - C_6 -alkyl-, $-C_1$ - C_6 -alkyl-CN,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 C_2 - C_6 -alkynyl- or C_3 - C_6 -cycloalkyl- group ; wherein said C_3 - C_6 -cycloalkyl-
 group is optionally substituted, one or more times, identically or
 differently, with a substituent selected from: halo-, C_1 - C_3 -alkyl-, cyano-,
 20 C_1 - C_3 -alkoxy- ;

R^3 represents a hydrogen atom, or a $-CN$, C_1 - C_6 -alkyl-,
 $-(CH_2)_m$ - C_2 - C_6 -alkenyl, $-(CH_2)_m$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 $-(CH_2)_m$ -(3- to 7-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-,

heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-,
aryl-, C₁-C₆-alkyl-aryl-, heteroaryl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
5 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂ or -N(H)C(=O)R⁶ group ;
said C₁-C₆-alkyl-, -(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
10 group
being optionally substituted, identically or differently, with 1, 2, 3, or 4
R⁷ groups ;

R^{4a}, R^{4b}, R^{4c}, R^{4d}

15 represent, independently from each other, a hydrogen or halogen atom,
or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,
R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-,
halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b},
-C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶,
20 -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶,
-N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶,
-N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b},
-O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶,
-S(=O)₂N(H)R^{6a}, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;

25

R⁵ represents a hydrogen atom, or a C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
-(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,
-(CH₂)_m-(3- to 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-,
heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
30 HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN, C₁-C₆-alkoxy-C₁-C₆-alkyl-,

halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group ;

5 s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl-,
 -(CH₂)_n-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to
 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
 10 C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group
 being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R⁸ groups ;

15 R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a
 C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to
 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or
 heteroaryl-C₁-C₆-alkyl- group ;

20

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 25 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,

-S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
-S(=O)(=NR^{6c})R⁶ group ;

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-,
5 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
-C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
-N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
10 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
-N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶,
-O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
-S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
-S(=O)(=NR^{6c})R⁶ group ;

15

m is an integer of 0, 1, 2, 3, or 4 ;



and

n is an integer of 1, 2, 3, 4, 5, 6 or 7 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt
20 thereof, or a mixture of same.


In another preferred embodiment, the invention relates to compounds of
formula (I), *supra*, in which :

25

A represents a  or  group;

Z represents a -C(=O)N(H)R² or -S(=O)₂N(H)R² group,

or a group selected from :



5 R^1 represents a hydrogen atom ;

R^2 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO - C_1 - C_6 -alkyl-, $-C_1$ - C_6 -alkyl-CN, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
10 C_2 - C_6 -alkynyl- or C_3 - C_6 -cycloalkyl- group ; wherein said C_3 - C_6 -cycloalkyl- group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, C_1 - C_3 -alkyl-, cyano-, C_1 - C_3 -alkoxy- ;

15 R^3 represents a hydrogen atom, or a $-CN$, C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_2 - C_6 -alkenyl, $-(CH_2)_m$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 7-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 7-membered
20 heterocycloalkyl-, C_2 - C_6 -alkenyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl-, C_1 - C_6 -alkyl-aryl-, heteroaryl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$, $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$ or $-N(H)C(=O)R^6$ group ;
s a i d C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_2 - C_6 -alkenyl, $-(CH_2)_m$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-,
25 C_3 - C_6 -cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C_2 - C_6 -alkenyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl-, C_1 - C_6 -alkyl-aryl-, or heteroaryl-group

being optionally substituted, identically or differently, with 1, 2, 3, or 4 R^7 groups ;

R^{4a} , R^{4b} , R^{4c} , R^{4d}

- 5 represent, independently from each other, a hydrogen or halogen atom, or a -CN, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, -OR⁶ group ;
- 10 R^5 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-, heteroaryl- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-, HO- C_1 - C_3 -alkyl-, $-C_1$ - C_3 -alkyl-CN, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-,
- 15 halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or heteroaryl- group ;
- s a i d C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to
- 20 6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-, heteroaryl- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-, HO- C_1 - C_3 -alkyl-, $-C_1$ - C_3 -alkyl-CN, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or heteroaryl- group being
- 25 optionally substituted, identically or differently, with 1 or 2 R^8 groups ;

R^6 , R^{6a} , R^{6b} , R^{6c} ,

represent, independently from each other, a hydrogen atom, or a C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, C_2 - C_6 -alkenyl-, 3- to

7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl- or heteroaryl-C₁-C₆-alkyl- group ;

5 R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, 10 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;

15

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, 20 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, 25 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;


m is an integer of 0, 1, 2, 3 or 4 ;

and

30 n is an integer of 1, 2, 3 or 4 ;


or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

- 5 In another preferred embodiment, the invention relates to compounds of formula (I), *supra*, in which :



10

Z represents a $-\text{C}(=\text{O})\text{N}(\text{H})\text{R}^2$ or $-\text{S}(=\text{O})_2\text{N}(\text{H})\text{R}^2$ group, or a group selected from :



15

R^1 represents a hydrogen atom ;

R^2 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $\text{R}^{6a}(\text{R}^{6b})\text{N}$ - C_1 - C_6 -alkyl-, HO - C_1 - C_6 -alkyl-, $-\text{C}_1$ - C_6 -alkyl-CN, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, C_2 - C_6 -alkynyl- or C_3 - C_6 -cycloalkyl- group ; wherein said C_3 - C_6 -cycloalkyl-group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, C_1 - C_3 -alkyl-, cyano-, C_1 - C_3 -alkoxy- ;

20

R^3 represents a hydrogen atom, or a -CN, C_1 - C_6 -alkyl-,
 $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl),
 aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-,
 5 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3-
 to 6-membered heterocycloalkyl, aryl-, C_1 - C_6 -alkyl-aryl-, heteroaryl-
 group ;
 said C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 6-membered
 heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-,
 10 halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
 C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-,
 C_1 - C_6 -alkyl-aryl- or heteroaryl- group being optionally substituted,
 identically or differently, with 1 or 2 R^7 groups ;

15

R^{4a} , R^{4b} , R^{4c} , R^{4d} :

two of which represent a hydrogen atom, while the remaining two
 represent, independently from each other, a hydrogen or halogen atom,
 or a -CN, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy-, halo- C_1 - C_6 -alkyl-,
 20 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
 halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, -OR⁶ group ;

R^5 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 25 $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-,
 heteroaryl- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-,
 HO- C_1 - C_3 -alkyl-, - C_1 - C_3 -alkyl-CN, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-,
 halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered
 heterocycloalkyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or
 30 heteroaryl- group ;

s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl,
 -(CH₂)_n-C₂-C₃-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to
 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-,
 halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN,
 5 C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-,
 C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-,
 C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group being
 optionally substituted, identically or differently, with 1 or 2 R⁸ groups ;

10 R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a
 C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to
 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl- or
 heteroaryl-C₁-C₆-alkyl- group ;

15

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 20 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 25 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 30 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,

3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
5 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$,
 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$ or
 $-S(=O)(=NR^{6c})R^6$ group ;

10 m is an integer of 0, 1, 2, 3 or 4 ;


and

n is an integer of 1, 2, 3 or 4 ;


or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt
15 thereof, or a mixture of same.

In another preferred embodiment, the invention relates to compounds of
formula (I), *supra*, in which :

20

A represents a  group;

Z represents a $-C(=O)N(H)R^2$ group,
or a

25  group ;

R¹ represents a hydrogen atom ;

R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
 5 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
 group is optionally substituted, one or more times, identically or
 differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
 C₁-C₃-alkoxy- ;

10

R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
 -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl),
 aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3-
 15 to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
 group ;
 said C₁-C₆-alkyl-, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered
 heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 20 C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-,
 C₁-C₆-alkyl-aryl- or heteroaryl- group being optionally substituted,
 identically or differently, with 1 or 2 R⁷ groups ;

R^{4a}, R^{4b}, R^{4c}, R^{4d} :

25 two of which represent a hydrogen atom, while the remaining two
 represent, independently from each other, a hydrogen or halogen atom,
 or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, -OR⁶ group ;

30

R^5 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-,
heteroaryl- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-,
5 HO- C_1 - C_3 -alkyl-, $-C_1$ - C_3 -alkyl-CN, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-,
halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered
heterocycloalkyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or
heteroaryl- group ;
s a i d C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, $-(CH_2)_n$ - C_2 - C_6 -alkenyl,
10 $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to
6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-, heteroaryl- C_1 - C_3 -alkyl-,
halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-, HO- C_1 - C_3 -alkyl-, $-C_1$ - C_3 -alkyl-CN,
 C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-,
 C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-,
15 C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or heteroaryl- group being
optionally substituted, identically or differently, with 1 or 2 R^8 groups ;

R^6 , R^{6a} , R^{6b} , R^{6c} ,

represent, independently from each other, a hydrogen atom, or a
20 C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, C_2 - C_6 -alkenyl-, 3- to
7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl- C_1 - C_6 -alkyl- or
heteroaryl- C_1 - C_6 -alkyl- group ;

R^7 represents a hydrogen or halogen atom, or a HO-, -CN, C_1 - C_6 -alkoxy-,
25 halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
30 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,

-N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;

5

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 10 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 15 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b},
 -S(=O)(=NR^{6c})R⁶ group ;

m is an integer of 0, 1, 2, 3 or 4 ;

and


20 n is an integer of 1, 2, 3 or 4 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.


25

In another preferred embodiment, the invention relates to compounds of formula (I), *supra*, in which :

A represents a



Z represents a $-C(=O)N(H)R^2$ group or a



5

R^1 represents a hydrogen atom ;

R^2 represents a C_3 - C_6 -cycloalkyl- or C_1 - C_4 -alkyl- group ; wherein said C_3 - C_6 -cycloalkyl- group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, C_1 - C_3 -alkyl-, cyano-, C_1 - C_3 -alkoxy- ; and wherein said or C_1 - C_4 -alkyl- group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, cyano-, C_1 - C_3 -alkoxy- ;

15

R^3 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl-, $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-, C_1 - C_6 -alkyl-aryl- or heteroaryl- group ; said C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl-, $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-, C_1 - C_6 -alkyl-aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R^7 groups.

(R^{4a}, R^{4b}, R^{4c}, R^{4d})

represent either (H, H, H, H) or (H, methyl, H, H) or (H, H, methyl, H).

5 R⁵ represents a C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl-,
 -(CH₂)_m-C₃-C₆-cycloalkyl-, aryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, C₁-C₃-alkoxy-C₁-C₃-alkyl- or
 C₃-C₆-cycloalkyl- group ;
 s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl,
 10 -(CH₂)_m-C₃-C₆-cycloalkyl-, aryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, C₁-C₃-alkoxy-C₁-C₃-alkyl- or
 C₃-C₆-cycloalkyl- group being optionally substituted, identically or
 differently, with 1 or 2 R⁸ groups.

15 R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a
 C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to
 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or
 heteroaryl-C₁-C₆-alkyl- group ;

20

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-,
 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 25 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 30 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;

R^8 represents a hydrogen or halogen atom, or a $-CN$, C_1-C_6 -alkoxy-,
 halo- C_1-C_6 -alkyl-, $R^{6a}(R^{6b})N-C_1-C_6$ -alkyl-, $HO-C_1-C_6$ -alkyl-,
 C_1-C_6 -alkoxy- C_1-C_6 -alkyl-, halo- C_1-C_6 -alkoxy- C_1-C_6 -alkyl-, C_2-C_6 -alkenyl-,
 5 3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$,
 10 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$ or
 $-S(=O)(=NR^{6c})R^6$ group ;

m is an integer of 0, 1, 2, 3, or 4.
 15 and
 n is an integer of 1, 2, 3 or 4.

It is to be understood that the present invention relates to any
 20 sub-combination within any embodiment of the present invention of
 compounds of general formula (I), *supra*.

More particularly still, the present invention covers compounds of general
 formula (I) which are disclosed in the Example section of this text, *infra*.


25

In an embodiment of the above-mentioned aspects, the invention relates to
 compounds of formula (I), according to any of the above-mentioned
 embodiments , in the form of or a stereoisomer, a tautomer, an N-oxide, a
 hydrate, a solvate, or a salt thereof, or a mixture of same.


30

In accordance with another aspect, the present invention covers methods of preparing compounds of the present invention, said methods comprising the steps as described in the Experimental Section herein.

- 5 In an embodiment, the present invention relates to a method of preparing compounds of general formula method of preparing a compound of general formula (I) :



- 10 in which A, R¹, R³ and R⁵ are as defined for the compounds of general formula (I), *supra*,
in which method an intermediate compound of general formula (II) :




- 15 in which A, R¹ and R⁵ are as defined for the compounds of general formula (I), *supra*,
is allowed to react with an alkyne of general formula (IIa) :



in which R^3 is as defined for the compounds of general formula (I), *supra*, and
 -M represents a hydrogen or a functional group suitable to be coupled to the
 Br-bearing carbon atom of the compound of general formula (II), such as
 5 -B(OH)₂, or an ester of a boronic acid such as -B(O-C₁-C₆-alkyl)₂ or
 4,4,5,5-tetramethyl-1,3,2-dioxaborolane, for example,
 thus providing a compound of general formula (I) as defined *supra*.

10 In accordance with a further aspect, the present invention covers intermediate
 compounds which are useful in the preparation of compounds of the present
 invention of general formula (I), particularly in the method described herein.
 In particular, the present invention covers compounds of general formula (II) :




(II)

15

in which A, R¹ and R⁵ are as defined for the compounds of general formula (I),
supra.

20 In accordance with yet another aspect, the present invention covers the use of
 the intermediate compounds of general formula (II) :



in which A, R¹ and R⁵ are as defined for the compounds of general formula (I),
supra, for the preparation of a compound of general formula (I) as defined
5 *supra*.

EXPERIMENTAL SECTION

The following Table lists the abbreviations used in this paragraph, and in the Examples section.

5


Abbreviation	Meaning
DMSO	Dimethyl sulfoxide
NMR	nuclear magnetic resonance spectroscopy
Rt	Room temperature
RT	Retention time in minutes
MW	molecular weight
UPLC	ultra performance liquid chromatography

Scheme 1 and the procedures described below illustrate general synthetic routes to the compounds of general formula (I) of the invention and are not intended to be limiting. It is obvious to the person skilled in the art that the order of transformations as exemplified in Scheme 1 can be modified in various ways. The order of transformations exemplified in the Scheme 1 is therefore not intended to be limiting. In addition, interconversion of any of the substituents, R¹, R³, R⁵, or A can be achieved before and/or after the exemplified transformations. These modifications can be such as the introduction of protecting groups, cleavage of protecting groups, exchange, reduction or oxidation of functional groups, halogenation, metallation, substitution or other reactions known to the person skilled in the art. These transformations include those which introduce a functionality which allows for further interconversion of substituents. Appropriate protecting groups and their introduction and cleavage are well-known to the person skilled in the art (see for example T.W. Greene and P.G.M. Wuts in Protective Groups in Organic Synthesis, 3rd edition, Wiley 1999). Specific examples are described in the

20

subsequent paragraphs. Further, it is possible that two or more successive steps may be performed without work-up being performed between said steps, *e.g.* a “one-pot” reaction, as is well-known to the person skilled in the art.

Synthesis of compounds of general formula (I) :



Scheme 1

wherein R^1 , R^3 , R^5 and A have the meaning as given for general formula (I),
supra, and Y represents a halogen atom and X represents an oxygen atom, a
5 dialkoxy-group or an α,ω -alkylenedioxy group, and M represents a hydrogen or a
functional group suitable to be coupled to the Br-bearing carbon atom of the
compound of general formula (II), such as $-B(OH)_2$, or an ester of a boronic acid
such as $-B(O-C_1-C_6-alkyl)_2$ or 4,4,5,5-tetramethyl-1,3,2-dioxaborolane, for
example.

10

Compounds of general formula (I) can be synthesised according to the
procedure depicted in Scheme 1. Scheme 1 exemplifies the main routes that
allow variations in R^1 , R^3 , R^5 and A at different stages of the synthesis.
However, also other routes were used to synthesise the target compounds.

15

The introduction of a $NH-CH_2-R^5$ can be achieved by nucleophilic substitution
of halo or sulfonyl precursor compounds like (IV) or (V), i. e. by reaction with
suitable amines in the presence of a suitable base, such as, for example DIPEA
in a suitable solvent such as *N,N*-dimethylformamide or
20 1-methylpyrrolidin-2-one, at temperatures ranging from room temperature to
the boiling point. Introduction of substituents A is achieved by a coupling
reaction, for example, particularly a metal-catalysed coupling reaction, with a
compound of formula $A-M'$, in which A is defined as for compounds of general
formula (I) *supra*, and M' represents a functional group suitable to be coupled
25 to the iodine-bearing carbon atom of compounds like (III) or (VI) to form
compounds like (II) or (V), respectively. Examples of $A-M'$ are $A-B(OH)_2$, or
esters of boronic acids like $A-B(O-C_1-C_6-alkyl)_2$ or
4,4,5,5-tetramethyl-1,3,2-dioxaborolane of A.

Introduction of $R^3\equiv$ -groups into compounds of formula (II) is achieved in an
30 analogous manner as already described for the introduction of substituents A.

Examples of such coupling reactions may be found in the textbook entitled “Metal-Catalyzed Cross-Coupling Reactions”, Armin de Meijere (Editor), François Diederich (Editor) September 2004, Wiley Interscience ISBN: 978-3-527-30518-6.

- 5 Said coupling reactions are performed in the presence of suitable catalysts, such as, for example, palladium based catalysts like, for example, Pd(OAc)₂ tetrakis(triphenylphosphine)palladium (0), bis(triphenylphosphine)palladium (II) chloride or (1,1,-bis(diphenylphosphino)ferrocene)-dichloropalladium (II) and optionally suitable additives such as, for example, phosphines like, for
10 example, P(oTol)₃ or triphenylphosphine and, and optionally with a suitable base, such as, for example, potassium carbonate, sodium 2-methylpropan-2-olate, tetrabutylammonium fluoride or tribasic potassium phosphate in a suitable solvent, such as, for example, tetrahydrofuran.

Iodination at the five membered ring to yield compounds of general formula
15 (VI) and (VII) is achieved by using halogenation reagents like N-iodosuccinimide, for example, in an inert solvent like N,N-dimethylformamide or 1-methylpyrrolidin-2-one, for example, at temperatures ranging from room temperature to the boiling point of the solvent, for example.

- 20 The bromine atom at position 8 of the compound of formula (VIII) can be substituted by methanethiolate in an inert solvent like an alcohol to give methylsulfanyl compounds of general formula (IX).

Methylsulfones of general formula (IV) can be obtained by oxidizing methylsulfanyl compounds of general formula (VII) using oxidation reagents
25 like 3-chlorobenzenecarboxoperoxoic acid or dimethyldioxirane in inert solvents like dichloromethane or acetone, respectively.

The starting materials of general formula (VIII) may be commercially available or can be synthesized according to procedures known to persons skilled in the art or are described, for example, in the Journal of Medicinal Chemistry, 1984,

vol. 27, # 2, p. 206 - 212 by reacting 2,4-dibromopyrimidin-5-amine with compounds of the general formula (X).

The compounds and intermediates produced according to the methods of the invention may require purification. Purification of organic compounds is well known to the person skilled in the art and there may be several ways of purifying the same compound. In some cases, no purification may be necessary. In some cases, the compounds may be purified by crystallisation. In some cases, impurities may be removed by stirring using a suitable solvent. In some cases, the compounds may be purified by chromatography, particularly flash chromatography, using for example pre-packed silica gel cartridges, e.g. from Separtis such as Isolute® Flash silica gel or Isolute® Flash NH2 silica gel in combination with a suitable chromatographic system such as an Isolera system (Biotage) and eluents such as, for example, gradients of hexane/ethyl acetate or dichloromethane/methanol. In some cases, the compounds may be purified by preparative HPLC using, for example, a Waters autopurifier equipped with a diode array detector and/or on-line electrospray ionisation mass spectrometer in combination with a suitable pre-packed reverse phase column and eluants such as, for example, gradients of water and acetonitrile which may contain additives such as trifluoroacetic acid, formic acid or aqueous ammonia.

Names of compounds were generated using the Autonom 2000 add-in of ISIS/Draw [MDL Information Systems Inc. (Elsevier MDL)] or the ICS naming tool of ACD labs.

The invention disclosed herein is exemplified by the following preparations and examples which should not be construed to limit the scope of the disclosure.

General:


All reactions were run under an atmosphere of argon in degassed solvents unless stated otherwise.

Analytical UPLC-MS was performed as follows:

- 5 Method A: System: UPLC Acquity (Waters) with PDA Detector und Waters ZQ mass spectrometer; Column: Acquity BEH C18 1.7 μ m 2.1x50mm; Temperature: 60 $^{\circ}$ C; Solvent A: Water + 0.1% formic acid; Solvent B: acetonitrile; Gradient: 99 % A $\rightarrow$ 1 % A (1.6 min) $\rightarrow$ 1 % A (0.4 min) ; Flow: 0.8 mL/min; Injection Volume: 1.0 μ l (0.1mg-1mg/mL sample concentration); Detection: PDA scan
10 range 210-400 nm - Fixed and ESI (+), scan range 170-800 m/z.

Example 1

N-cyclopropyl-4-{8-[(3-hydroxypropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide




- To a solution of 150 mg (349 μ mol)
4-{6-bromo-8-[(3-hydroxypropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide which was prepared according to intermediate example 1a in 4.2 mL tetrahydrofuran were added 172 μ L piperidine, 103 μ L prop-2-yn-1-ol and
20 40 mg tetrakis(triphenylphosphine)palladium (0). The mixture was subjected to microwave radiation for 2 hours at 80 $^{\circ}$ C. Water was added and the mixture was extracted with ethyl acetate. The organic layer was washed with brine and dried over sodium sulfate. After filtration and removal of the solvent the residue was purified by chromatography to give 37.8 mg (25%) of the title
25 compound.

¹H-NMR (DMSO-d₆): δ= 0.56 (2H), 0.68 (2H), 1.74 (2H), 2.84 (1H), 3.12 (1H), 3.41-3.53 (4H), 4.25 (2H), 4.56 (1H), 5.33 (1H), 7.67-7.81 (4H), 7.94 (2H), 8.52 (1H) ppm.

5 Intermediate Example 1a

4-{6-Bromo-8-[(3-hydroxypropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide




To a solution of 500 mg (1.15 mmol)

10 N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)benzamide which was prepared according to intermediate example 1b in 15 mL

N,N-dimethylformamide were added 261 µL 3-aminopropan-1-ol and the mixture was stirred at 40°C for 6 hours. Toluene was added and the solvents removed. The residue was purified by chromatography to give 453 mg (92%) of
15 the title compound.

Intermediate Example 1b

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)benzamide




20 A mixture comprising 5.0 g (12.41 mmol)

6,8-dibromo-3-iodoimidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c, 3.31 g 4-(cyclopropylaminocarbonyl)phenylboronic

acid, 1.01 g (1,1,-bis(diphenylphosphino)ferrocene)-dichloropalladium (II), 8.7 mL aqueous 2M tribasic potassium phosphate solution and 85 mL tetrahydrofuran was stirred at 40°C for 2.5 days. Water was added and the mixture was extracted with dichloromethane. The organic layer was washed
 5 with brine and dried over sodium sulfate. After filtration and removal of the solvent the residue was purified by chromatography to give 2.49 g (46%) of the title compound.

Intermediate Example 1c

10 6,8-dibromo-3-iodo-imidazo[1,2-a]pyrazine




To a stirred solution of 8.7 g (31.4 mmol) 6,8-dibromo-imidazo[1,2-a]pyrazine which was prepared according to a procedure described in the Journal of Medicinal Chemistry, 1984, vol. 27, # 2, p. 206 - 212 in 210 mL

15 N,N-dimethylformamide was added 7.42 g N-iodopyrrolidin-2,5-dione in one portion at 23°C. After 18 hours of stirring at 60°C, the solvent was removed in vacuo and the residue was taken up in dichloromethane and washed with water and saturated sodium thiosulfate solution. The organic phase was dried over sodium sulphate, filtered and the solvent was evaporated to yield 9.46 g
 20 (74.8 of the title compound).

Example 2

N-cyclopropyl-4-[8-(isobutylamino)-6-(phenylethynyl)imidazo[1,2-a]pyrazin-3-yl]benzenesulfonamide



A mixture of 30 mg (65 μ mol)

4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzene sulfonamide which was prepared according to intermediate example 2a, 29.5


5 mg 4,4,5,5-tetramethyl-2-(phenylethynyl)-1,3,2-dioxaborolane, 1.0 mL n-propanol, 97 μ L of an aqueous 2M potassium carbonate solution, 0.85 mg triphenylphosphine, and 4.55 mg bis(triphenylphosphine)palladium (II) chloride was stirred at 120°C for 2 hours. The solution was cooled, water added and extracted with dichloromethane. The organic phase was dried over sodium sulfate. After filtration and removal of solvent the residue was purified by chromatography to give 4.6 mg (14%) of the title compound.

¹H-NMR (CDCl₃): δ = 0.63-0.73 (4H), 1.06 (6H), 2.04 (1H), 2.34 (1H), 3.53 (2H), 4.98 (1H), 6.14 (1H), 7.32-7.38 (3H), 7.54-7.60 (2H), 7.66 (1H), 7.74 (2H), 7.86 (1H), 8.08 (2H) ppm.

15

Intermediate Example 2a

4-[6-Bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzene sulfonamide




200 mg (424 μ mol)

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)benzenesulfonamide which was prepared according to intermediate example 2b were transformed in analogy to intermediate example 1a using 2-methylpropan-1-amine to give
5 after working up and purification 124 mg (63%) of the title compound.

Intermediate Example 2b


N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)benzenesulfonamide



10 3.0 g (7.45 mmol) 6,8-dibromo-3-iodoimidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c were transformed in analogy to example 1b using [4-(cyclopropylsulfamoyl)phenyl]boronic acid to give after working up and purification 1.85 g (52%) of the title compound.

15 Example 3

N-cyclopropyl-4-{6-[(2-fluorophenyl)ethynyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl}benzenesulfonamide



30 mg (65 μ mol)

20 4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzenesulfonamide which was prepared according to intermediate example 2a were


transformed in analogy to example 2 using diisopropyl

[(2-fluorophenyl)ethynyl]boronate to give after working up and purification 6.4 mg (18%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.64-0.71 (4H), 1.06 (6H), 2.04 (1H), 2.34 (1H), 3.52 (2H),
5 5.01 (1H), 6.15 (1H), 7.09 (1H), 7.15 (1H), 7.34 (1H), 7.58 (1H), 7.66 (1H), 7.74 (2H), 7.88 (1H), 8.08 (2H) ppm.

10 Example 4

N-cyclopropyl-4-[6-(3-hydroxyprop-1-yn-1-yl)-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]benzenesulfonamide




40 mg (86 μmol)

15 4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzenesulfonamide which was prepared according to intermediate example 2a were transformed in analogy to example 1 to give after working up and purification 5.7 mg (14%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.60-0.73 (4H), 1.03 (6H), 2.01 (1H), 2.38 (1H), 3.46 (2H),
20 4.50 (2H), 6.18 (1H), 7.64 (1H), 7.68 (2H), 7.76 (1H), 8.05 (2H) ppm.

Example 5

N-cyclopropyl-4-[6-(3-hydroxyprop-1-yn-1-yl)-8-(isobutylamino)imidazo[1,2-



-a]pyrazin-3-yl]benzamide

50 mg (117 μ mol)

4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzami


5 de which was prepared according to intermediate example 6.1 were transformed in analogy to example 1 to give after working up and purification 7.1 mg (14%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.68 (2H), 0.91 (2H), 4.04 (6H), 2.01 (1H), 2.58 (1H), 2.96 (1H), 3.46 (2H), 4.49 (2H), 6.13 (1H), 6.47 (1H), 7.53 (2H), 7.58 (1H), 7.68

10 (1H), 7.87 (2H) ppm.

Intermediate Example 5a

4-(6-bromo-8-isobutylamino-imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropyl-benzami
de




15

74.2 g (188 mmol) (6-bromo-3-iodo-imidazo[1,2-a]pyrazin-8-yl)-isobutyl-amine which was prepared according to intermediate example 5b were transformed in analogy to example 1b to give after working up and purification 45.2 g (56%) of the title compound.

20

Intermediate Example 5b

(6-bromo-3-iodo-imidazo[1,2-a]pyrazin-8-yl)-isobutyl-amine



To a stirred solution of 5.08 g (12.64 mmol)

6-bromo-3-iodo-8-methanesulfonyl-imidazo[1,2-a]pyrazine which was prepared

5 according to intermediate example 1c in 100 mL 1-methylpyrrolidin-2-one

were added 3.77 mL isobutylamine. After stirring at 23°C for 2 hours, 500 mL

water were added and the mixture was extracted with ethyl acetate. The


organic phase was filtered, evaporated and the residue was recrystallized from

methanol / water to yield 3.87 g (78%) of the title compound.

10

Example 6

N-cyclopropyl-4-[8-(isobutylamino)-6-(phenylethynyl)imidazo[1,2-a]pyrazin-3-yl]benzamide



15 50 mg (117 µmol)

4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzamide which was prepared according to intermediate example 5a were transformed in analogy to example 2 using


4,4,5,5-tetramethyl-2-(phenylethynyl)-1,3,2-dioxaborolane to give after

20 working up and purification 24.9 mg (45%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.66 (2H), 0.91 (2H), 1.05 (6H), 2.03 (1H), 2.94 (1H), 3.51 (2H), 6.09 (1H), 6.35 (1H), 7.30-7.37 (3H), 7.52-7.65 (5H), 7.83 (1H), 7.90 (2H) ppm.

Example 7

N-cyclopropyl-4-{6-[(2-fluorophenyl)ethynyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl}benzamide



50 mg (117 μ mol)

4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzamide which was prepared according to intermediate example 5a were transformed in analogy to example 2 using diisopropyl


10 [(2-fluorophenyl)ethynyl]boronate to give after working up and purification 24.3 mg (42%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.66 (2H), 0.91 (2H), 1.05 (6H), 2.03 (1H), 2.95 (1H), 3.51 (2H), 6.10 (1H), 6.33 (1H), 7.08 (1H), 7.14 (1H), 7.33 (1H), 7.53-7.65 (4H), 7.86 (1H), 7.91 (2H) ppm.

15

Example 8

N-cyclopropyl-4-{8-[(4-hydroxybutyl)amino]-6-(3-hydroxy-3-methylbut-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide



20 To a solution of 90 mg (203 μ mol)


4-{6-bromo-8-[(4-hydroxybutyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide which was prepared according to intermediate example 8a in 2.0

mL tetrahydrofuran were added 68.1 mg 2-methylbut-3-yn-2-ol, 1.0 mL tetrabutylammonium fluoride (1 molar in tetrahydrofuran) and 42.8 mg bis(triphenylphosphine)palladium (II) chloride. The mixture was subjected to microwave radiation for 2 hours at 130°C. Water was added and the mixture
 5 was extracted with dichloromethane. The organic layer was washed with brine and dried over sodium sulfate. After filtration and removal of the solvent the residue was purified by chromatography to give 2.1 mg (2%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.67 (2H), 0.92 (2H), 1.60 (6H), 1.70 (2H), 1.86 (2H), 2.95
 10 (1H), 3.67 (2H), 3.82 (2H), 6.31 (2H), 7.57 (2H), 7.59 (1H), 7.66 (1H), 7.89 (2H) ppm.

Intermediate example 8a

15 4-{6-Bromo-8-[(4-hydroxybutyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide




1.0 g (2.29 mmol)

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)benzamide which was
 20 prepared according to intermediate example 1b were transformed in analogy to example 1a using 4-aminobutan-1-ol to give after working up and purification 923 mg (91%) of the title compound.

Example 9

N-cyclopropyl-4-{8-[(4-hydroxybutyl)amino]-6-[(1-hydroxycyclopentyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzamide




90 mg (203 μ mol)

- 5 4-{6-bromo-8-[(4-hydroxybutyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide which was prepared according to intermediate example 8a were transformed in analogy to example 8 using 1-ethynylcyclopentanol to give after working up and purification 4.9 mg (5%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.67 (2H), 0.91 (2H), 1.62-1.94 (10H), 1.99-2.07 (4H), 2.95 (1H), 3.66 (2H), 3.80 (2H), 6.36 (1H), 6.45 (1H), 7.54 (2H), 7.56 (1H), 7.62 (1H), 7.88 (2H) ppm.

Example 10

- 15 **N-cyclopropyl-4-{8-[(2-hydroxy-2-methylpropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide**




100 mg (225 μ mol)

- 20 4-{6-bromo-8-[(2-hydroxy-2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide which was prepared according to intermediate example 10a were transformed in analogy to example 1 to give after working up and purification 21.4 mg (23%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.56 (2H), 0.68 (2H), 1.13 (6H), 2.84 (1H), 3.43 (2H), 4.25 (2H), 4.88 (1H), 5.33 (1H), 7.11 (1H), 7.72 (2H), 7.81 (1H), 8.82 (1H), 7.94 (2H), 8.51 (1H) ppm.

5 Intermediate Example 10a

4-{6-bromo-8-[(2-hydroxy-2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide




700 mg (1.61 mmol)

- 10 6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 1b were transformed in analogy to intermediate example 1a using 1-amino-2-methylpropan-2-ol to give after working up and purification 527 mg (74%) of the title compound.

15 Example 11

4-{8-[(3-Aminopropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-3-methoxybenzamide



72 mg (157 μmol)


- 20 4-{8-[(3-aminopropyl)amino]-6-bromoimidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-3-methoxybenzamide which was prepared according to intermediate example

11a were transformed in analogy to example 1 to give after working up and purification 20.3 mg (28%) of the title compound.

¹H-NMR (CD₃OD): δ= 0.66 (2H), 0.83 (2H), 2.03 (2H), 2.88 (1H), 3.03 (2H), 3.68 (2H), 3.88 (3H), 4.36 (2H), 7.43 (1H), 7.46 (1H), 7.52 (1H), 7.59 (1H), 7.60 (1H), 8.44 (1H) ppm.

Intermediate Example 11a

4-{8-[(3-aminopropyl)amino]-6-bromoimidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-3-methoxybenzamide




37.5 mg (76 μmol)

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)-3-methoxybenzamide which was prepared according to intermediate example 11b were transformed in analogy to example 1a using propane-1,3-diamine to give after working up and purification 26.6 mg (72%) of the title compound.

Intermediate Example 11b

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)-3-methoxybenzamide




72.7 mg (181 μmol) 6,8-dibromo-3-iodoimidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c were transformed in analogy to example 1b using [4-(cyclopropylcarbamoyl)-2-methoxyphenyl]boronic acid

which was prepared according to intermediate example 11c to give after working up and purification 37.5 mg (42%) of the title compound.

Intermediate Example 11c

5 [4-(cyclopropylcarbamoyl)-2-methoxyphenyl]boronic acid



A suspension of 3,0 g (15.3 mmol) 4-(dihydroxyboryl)-3-methoxybenzoic acid, 874 mg cyclopropanamine and 3.5 g


N-[3-(dimethylamino)propyl]-N'-ethylcarbodiimide in 93 mL dichloromethane

10 was stirred overnight at 23°C.








Water was added and the mixture was extracted with dichloromethane. The organic layer was washed with brine and dried over sodium sulfate. After filtration and removal of the solvent the residue was purified by chromatography to give 2.60 g (72%) of the title compound.









15








Examples 12 to 48
















4-[6-bromo-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]-N-cyclopropylbenzami
de which was prepared according to intermediate example 5a was transformed
20 in analogy to example 8 using suitable substituted ethynes to give after
working up and purification the corresponding title compounds:

Exp	Name	R ³	Yield (%)	LC/MS RT (min)	LC/MS m/z (M+1)
12	4-{6-(cyclohexylethynyl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		6	1.5	456
13	N-cyclopropyl-4-{6-(3-hydroxy-3-methylbut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		25	1.11	432
14	N-cyclopropyl-4-{6-(4-hydroxybut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		36	1.01	418
15	N-cyclopropyl-4-{6-[(1-hydroxycyclopentyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		29	1.2	458
16	N-cyclopropyl-4-{6-[3-(dimethylamino)prop-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		31	0.78	431
17	N-cyclopropyl-4-{6-(3-methoxyprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		3	1.18	418
18	N-cyclopropyl-4-{6-(3,3-dimethylbut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		14	1.4	430

19	N-cyclopropyl-4-{6-(3-hydroxybut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		11	1.06	418
20	N-cyclopropyl-4-{6-(4-hydroxypent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		17	1.06	432
21	4-{6-(5-cyanopent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		17	1.17	441
22	N-cyclopropyl-4-{6-(3-hydroxypent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		18	1.13	432
23	4-(6-{3-[benzyl(methyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropylbenzamide		23	0.91	507
24	4-{6-(3-aminoprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		6	1.04	403
25	N-cyclopropyl-4-{6-(3-hydroxyhex-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		8	1.36	446
26	N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(5-phenylpent-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide		29	1.49	492

27	6-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}hex-5-ynoic acid		9	1.18	460
28	N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(pyridin-2-ylethynyl)imidazo[1,2-a]pyrazin-3-yl}benzamide		15	1.2	451
29	N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(4-phenylbut-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide		32	1.43	478
30	4-{6-(5-cyclohexylpent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		12	1.7	498
31	N-cyclopropyl-4-{6-(4-methylpent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		28	1.42	430
32	N-cyclopropyl-4-{6-[(3-hydroxyphenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		36	1.23	466
33	4-{6-[3-(carbamoylamino)prop-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		19	0.93	446
34	N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(3-phenylprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide		4	1.5	464


35	N-cyclopropyl-4-{6-[(4-fluorophenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		34	1.41	468
36	N-cyclopropyl-4-{6-[(3S)-3-hydroxyoct-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		7	1.34	474
37	methyl 6-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}hex-5-ynoate		11	1.24	474
38	9-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}non-8-ynoic acid		9	1.24	502
39	N-cyclopropyl-4-(6-{3-[(methylcarbamoyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl) benzamide		13	0.96	460
40	N-cyclopropyl-4-(6-{4-[methyl(methylcarbamoyl)amino]but-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide		16	1.04	488
41	N-cyclopropyl-4-{6-[6-(methylamino)-6-oxohex-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		25	1.04	473

42	N-cyclopropyl-4-(6-{3-[methyl (methylcarbamoyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide		54	0.96	474
43	4-{6-[(4-carbamoylphenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		33	1.12	493
44	N-cyclopropyl-4-(6-{4-[(methylcarbamoyl)amino]but-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide		31	0.99	474
45	N-cyclopropyl-4-{6-[5-(methylamino)-5-oxopent-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide		29	1.0	459
46	N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-[(4-sulfamoylphenyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzamide		15	1.15	529
47	4-{6-(cyclopentylethynyl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide		14	1.42	441
48	N-cyclopropyl-4-{6-ethynyl-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide	H	49	1.20	374

* indicates the point of attachment of said groups with the rest of the molecule

Example 49

N-cyclopropyl-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide



32 mg (70 μ mol)


- 5 4-(6-Bromo-8-isobutylamino-imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropyl-2-methylbenzamide which was prepared according to intermediate example 49a were transformed in analogy to example 8 using prop-2-yn-1-ol to give after working up and purification 16.9 mg (56%) of the title compound.

UPLC-MS: RT = 1.02 min; m/z (ES⁺) 418.5 [MH⁺]; required MW = 417.5.

10

Intermediate Example 49a

4-(6-Bromo-8-isobutylamino-imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropyl-2-methylbenzamide



- 15 To a solution of 411 mg (1 mmol)

4-(6-Bromo-8-isobutylamino-imidazo[1,2-a]pyrazin-3-yl)-2-methylbenzoic acid which was prepared according to intermediate example 49b in 4 mL


1-methylpyrrolidin-2-one were added 114.9 mg cyclopropanamine, 760 mg bis(dimethylamino)(3-oxido-1H-[1,2,3]triazolo[4,5-b]pyridin-1-yl)methylum

- 20 hexafluorophosphate and 685 μ L N-ethyl-N-isopropylpropan-2-amine and the mixture was stirred overnight at 23°C. After filtration and removal of the

solvent the residue was purified by chromatography to give 65.5 mg (15%) of the title compound.

Intermediate Example 49b


5 4-(6-Bromo-8-isobutylamino-imidazo[1,2-a]pyrazin-3-yl)-2-methyl-benzoic acid



395 mg (1 mmol) (6-Bromo-3-iodo-imidazo[1,2-a]pyrazin-8-yl)-isobutyl-amine which was prepared according to intermediate example 5b were transformed in analogy to example 1b using 4-(dihydroxyboryl)-2-methylbenzoic acid to give
10 the title compound that was used without further purification.

Example 50

(2S)-3-({3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-8-yl}amino)propane-1,2-diol



15

69 mg (147 μ mol)


(2S)-3-({6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)propane-1,2-diol which was prepared according to intermediate example 50a were transformed in analogy to example 1 to give
20 after working up and purification 10.7 mg (16%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.82-0.92 (4H), 1.94 (1H), 3.32-3.42 (3H), 3.58 (1H), 3.71 (1H), 4.25 (2H), 4.67 (1H), 4.99 (1H), 5.33 (1H), 7.35 (1H), 7.49-7.87 (7H), 11.85+12.20 (1H) ppm.

5

Intermediate Example 50a

(2S)-3-({6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)propane-1,2-diol



10 100 mg (218 μmol)


6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to intermediate example 1a using

(2S)-3-aminopropane-1,2-diol to give after working up and purification 69 mg

15 (68%) of the title compound.

Intermediate Example 50b

6,8-dibromo-3-[4-(2-cyclopropyl-3H-imidazol-4-yl)-phenyl]-imidazo[1,2-a]pyrazine



20


1.0 g (2.48 mmol) 6,8-dibromo-3-iodo-imidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c were transformed in analogy to intermediate example 1b using

2-cyclopropyl-5-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1H-imidazole which was prepared according to intermediate example 50c to give after working up and purification 485 mg (43%) of the title compound.

Intermediate Example 50c

2-cyclopropyl-5-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1H-imidazole

dazole



A mixture comprising 42.1 g (160 mmol)

5-(4-bromophenyl)-2-cyclopropyl-1H-imidazole which was prepared according to intermediate example 50d, 60.9 g

4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane, 3.05 g

dicyclohexyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine, 1.46 g

tris-(dibenzylidenacetone)-dipalladium(0), 23.5 g potassium acetate and 267

mL 1,4-dioxane was stirred at 110°C for 4 hours. After cooling to 23°C, ethyl

acetate and 1M hydrochloric acid were added and the pH was adjusted to 8.0

by adding 5% sodium hydroxide solution. The mixture was extracted with ethyl


acetate. The organic layer was washed with brine and dried over sodium

sulfate. After filtration and removal of the solvent the residue was purified by

crystallisation to give 33.5 g (67%) of the title compound.

Intermediate Example 50d

5-(4-bromophenyl)-2-cyclopropyl-1H-imidazole




To a solution of 89.0 g (320 mmol) 2-bromo-1-(4-bromophenyl)ethanone in 2.0 L *N,N*-dimethylformamide were added cyclopropanecarboximidamide hydrochloride, 132.8 g potassium carbonate and the mixture was stirred at 23°C overnight. After removal of the solvent, water was added and the mixture was extracted with ethyl acetate. The organic layer was washed with brine and dried over sodium sulfate. After filtration and removal of the solvent the residue was purified by crystallisation to give 42.1 g (50%) of the title compound.

10

Example 51

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(4,4,4-trifluorobutyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol




15 83 mg (140 µmol)

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(4,4,4-trifluorobutyl)imidazo[1,2-a]pyrazin-8-amine which was prepared according to intermediate example 51a were transformed in analogy to example 1 to give after working up and purification 7.9 mg (11%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.97-1.06 (4H), 1.95-2.30 (6H), 3.66 (2H), 4.47 (2H), 6.48 (1H), 7.15 (1H), 7.36 (2H), 7.49 (1H), 7.62 (1H), 7.71 (2H) ppm.

Intermediate Example 51a

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(4,4,4-trifluorobutyl)imidazo[1,2-a]pyrazin-8-amine



5 100 mg (218 μ mol)


6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to example 1a using

propane-4,4,4-trifluorobutan-1-amine to give after working up and purification

10 88 mg (80%) of the title compound.

Example 52

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(2-methylprop-2-en-1-yl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



15

70 mg (156 μ mol)

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(2-methylprop-2-en-1-yl)imidazo[1,2-a]pyrazin-8-amine which was prepared according to


intermediate example 52a were transformed in analogy to example 1 to give

20 after working up and purification 12.2 mg (18%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.82-0.92 (4H), 1.71 (3H), 1.94 (1H), 3.99 (2H), 4.24 (2H), 4.74 (1H), 4.77 (1H), 5.31 (1H), 7.49-7.88 (8H), 11.85+12.20 (1H) ppm.

5 Intermediate Example 52a

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(2-methylprop-2-en-1-yl)imidazo[1,2-a]pyrazin-8-amine




100 mg (218 μmol)

- 10 6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to example 1a using 2-methylprop-2-en-1-amine to give after working up and purification 72 mg (74%) of the title compound.

15 Example 53

3-{8-(Allylamino)-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



68 mg (156 μmol)


- 20 N-allyl-6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-amine which was prepared according to intermediate example 53a were

transformed in analogy to example 1 to give after working up and purification 9.9 mg (15%) of the title compound.

¹H-NMR (DMSO-d₆): δ= 0.82-0.92 (4H), 1.94 (1H), 4.06 (2H), 4.24 (2H), 5.04 (1H), 5.14 (1H), 5.32 (1H), 5.95 (1H), 7.24+7.48-7.90 (8H), 11.85+12.20 (1H) ppm.

Intermediate Example 53a

N-allyl-6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-amine




100 mg (218 μmol)

6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to example 1a using prop-2-en-1-amine to give after working up and purification 71 mg (75%) of the title compound.

Example 54

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



140 mg (310 μ mol)

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-isobutylimidazo[1,2-a]

pyrazin-8-amine which was prepared according to intermediate example 54a


were transformed in analogy to example 1 to give after working up and

5 purification 29.5 mg (21%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.82-0.92 (10H), 1.91-2.06 (2H), 3.25 (2H), 4.23-4.27 (3H), 5.32 (1H), 7.51 (1H), 7.55 (2H), 7.68 (1H), 7.75 (1H), 7.84 (2H), 11.85+12.20 (1H) ppm.

10 Intermediate Example 54a

4-(6-bromo-8-methanesulfonyl-imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropyl-benzamide



384 mg (836 μ mol)

15 6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-


e which was prepared according to intermediate example 50b were

transformed in analogy to example 1a using 2-methylpropan-1-amine to give

after working up and purification 298 mg (79%) of the title compound.

20 Example 55

3-{3-[3-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



170 mg (377 μ mol)

6-bromo-3-[3-(2-cyclopropyl-1H-imidazol-4-yl)phenyl]-N-isobutylimidazo[1,2-a]pyrazin-8-amine which was prepared according to intermediate example 55a


5 were transformed in analogy to example 1 to give after working up and purification 15.2 mg (9%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.88-0.95 (4H), 0.98 (6H), 1.94 (2H), 3.30 (1H), 3.35 (2H), 3.54-3.82 (2H), 4.33 (2H), 7.10 (1H), 7.27 (1H), 7.43 (1H), 7.49 (1H), 7.64 (1H), 7.70 (1H), 7.77 (1H) ppm.

10

Intermediate Example 55a

6-Bromo-3-[3-(2-cyclopropyl-1H-imidazol-4-yl)phenyl]-N-isobutylimidazo[1,2-a]pyrazin-8-amine



15 316 mg (688 μ mol)


6,8-dibromo-3-[3-(2-cyclopropyl-1H-imidazol-4-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 55b were transformed in analogy to intermediate example 1a using

2-methylpropan-1-amine to give after working up and purification 293 mg (94%)

20 of the title compound.

Intermediate Example 55b

6,8-Dibromo-3-[3-(2-cyclopropyl-1H-imidazol-4-yl)phenyl]imidazo[1,2-a]pyrazine



5

689 mg (1.71 mmol) 6,8-dibromo-3-iodoimidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c were transformed in analogy to intermediate example 50b using


2-cyclopropyl-4-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1H-imidazole which was prepared according to intermediate example 55c to give after working up and purification 379 mg (48%) of the title compound.

10

Intermediate Example 55c

2-cyclopropyl-4-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1H-imidazole

15




8.41 g (32.0 mmol) 4-(3-bromophenyl)-2-cyclopropyl-1H-imidazole which was prepared according to intermediate example 55d were transformed in analogy to example 50c to give after working up and purification 690 mg (7%) of the title compound.

20

Intermediate Example 55d

4-(3-bromophenyl)-2-cyclopropyl-1H-imidazole




25.6 g (92 mmol) 2-bromo-1-(3-bromophenyl)ethanone were transformed in analogy to example 50d to give after working up and purification 8.4 g (35%) of the title compound.

5

Example 56

4-({3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-8-yl}amino)butan-1-ol



10 101 mg (216 μ mol)


4-({6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)butan-1-ol which was prepared according to intermediate example 56a were transformed in analogy to example 1 to give after working up and purification 6.5 mg (6%) of the title compound.

15 $^1\text{H-NMR}$ (CD_3OD): δ = 0.86-0.93 (4H), 1.59 (2H), 1.71 (2H), 1.92 (1H), 3.51 (2H), 3.63 (2H), 4.30 (2H), 7.08 (1H), 7.32 (2H), 7.39 (1H), 7.60 (1H), 7.64 (2H) ppm.

Intermediate Example 56a

4-({6-Bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)butan-1-ol

20




200 mg (436 μ mol)

6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were

- 5 transformed in analogy to intermediate example 1a using 4-aminobutan-1-ol to give after working up and purification 166 mg (82%) of the title compound.

Example 57

- 3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol




166 mg (338 μ mol)

- 6-Bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(3,3,3-trifluoropropyl)imidazo[1,2-a]pyrazine-8-amine which was prepared according to intermediate example 57a were transformed in analogy to example 1 to give after working up and purification 31.2 mg (19%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.81-0.93 (4H), 1.94 (1H), 2.64 (2H), 3.68 (2H), 4.26 (2H), 5.32 (1H), 7.24+7.51 (1H), 7.55+7.62 (2H), 7.70-7.92 (5H), 11.85+12.20 (1H) ppm.

Intermediate Example 57a

6-Bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(3,3,3-trifluoropropyl)imidazo[1,2-a]pyrazin-8-amine



5

240 mg (523 μ mol)


6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to intermediate example 1a using

10 3,3,3-trifluoropropan-1-amine to give after working up 340 mg of the title compound that was used without further purification.

Example 58

3-{8-[(Cyclobutylmethyl)amino]-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol

15



90 mg (194 μ mol)

6-bromo-N-(cyclobutylmethyl)-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-amine which was prepared according to intermediate


example 58a were transformed in analogy to example 1 to give after working up and purification 11.9 mg (13%) of the title compound.

¹H-NMR (CDCl₃): δ= 0.95-1.06 (4H), 1.73-2.17 (7H), 2.65 (1H), 3.59 (2H), 4.48 (2H), 6.23 (1H), 7.15 (1H), 7.32 (2H), 7.47 (1H), 7.58 (1H), 7.67 (2H) ppm.

5

Intermediate Example 58a

6-Bromo-N-(cyclobutylmethyl)-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-amine




10 229 mg (499 μmol)

6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to intermediate example 1a using 1-cyclobutylmethanamine to give after working up and purification 130 mg (56%) of the title compound.

Example 59

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(3-hydroxypropyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



135 mg (298 μ mol)


3-({6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)propan-1-ol which was prepared according to intermediate

5 example 59a were transformed in analogy to example 1 to give after working up and purification 25.9 mg (19%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.81-0.93 (4H), 1.74 (2H), 1.94 (1H), 3.43-3.53 (4H), 4.25 (2H), 4.56 (1H), 5.32 (1H), 7.48-7.87 (8H), 11.85+12.21 (1H) ppm.

10 Intermediate Example 59a

3-({6-Bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}amino)propan-1-ol




200 mg (436 μ mol)

15 6,8-dibromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazine which was prepared according to intermediate example 50b were transformed in analogy to example 1a using 3-aminopropan-1-ol to give after working up and purification 141 mg (71%) of the title compound.

Example 60

3-{8-[(3-Aminopropyl)amino]-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol



5

170 mg (376 μ mol)


N-{6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}propane-1,3-diamine which was prepared according to intermediate example 60a were transformed in analogy to example 1 to give after working up and purification 8 mg (5%) of the title compound.

10

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.81-0.91 (4H), 1.64 (2H), 1.94 (1H), 2.58 (2H), 3.47 (2H), 4.23 (2H), 7.44 (1H), 7.55 (2H), 7.65+7.68 (1H), 7.71+7.74 (1H), 7.81 (2H) ppm.

15 Intermediate Example 60a

N-{6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-8-yl}propane-1,3-diamine




720 mg (1.81 mmol) which was prepared according to intermediate example 60b were transformed in analogy to example 1b using [4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]boronic acid to give after working up and purification 420 mg (51%) of the title compound.

5

Intermediate Example 60b

N-(6-bromo-3-iodoimidazo[1,2-a]pyrazin-8-yl)propane-1,3-diamine



To a solution of 1.26 mL propane-1,3-diamine in 8 mL 2.0 g


10 N,N-dimethylformamide was added a solution of (4.98 mmol)

6-bromo-3-iodo-8-(methylsulfonyl)imidazo[1,2-a]pyrazine which was prepared according to intermediate example 60c in 20 mL N,N-dimethylformamide at 40 °C during 1.5 hours. The mixture was stirred at 40 °C for additional 1.5 hours. The solvent was removed and the residue was purified by

15 chromatography to give 1.52 g (77%) of the title compound.

Intermediate Example 60c

6-bromo-3-iodo-8-(methylsulfonyl)imidazo[1,2-a]pyrazine



20 To a stirred solution of 100 g (270.3 mmol)


6-bromo-3-iodo-8-methylsulfanylimidazo[1,2-a]pyrazine which was prepared according to intermediate example 60d in 2 L dichloromethane were added 116.6 g 3-chlorobenzenecarboxoperoxoic acid in several portions at 0 °C. After stirring for 1 hour at 23 °C, additional 46.6 g 3-chlorobenzenecarboxoperoxoic acid were added and the mixture was stirred overnight. The suspension was

25 filtered and the organic phase was washed with water, saturated sodium

hydrogen carbonate solution, brine, dried over sodium sulphate, filtered and evaporated to yield 197 g of an orange solid. The solid was refluxed in ethanol, filtered and dried to yield 104.5 g (96%) of the title compound.

5 Intermediate Example 60d


6-bromo-3-iodo-8-methylsulfanyl-imidazo[1,2-a]pyrazine



210 g (860 mmol) 6-bromo-8-methylsulfanyl-imidazo[1,2-a]pyrazine which was prepared according to intermediate example 60e were transformed in analogy to example 1c to give after working up and purification 255 g (80%) of the title compound.

Intermediate Example 60e


6-bromo-8-methylsulfanyl-imidazo[1,2-a]pyrazine



To a stirred suspension of 489 g (1766 mmol) 6,8-dibromo-imidazo[1,2-a]pyrazine which was prepared according to a procedure described in the Journal of Medicinal Chemistry, 1984, vol. 27, # 2, p. 206 - 212 in 2.9 L methanol at -20°C was dropwise added a solution of 225 g sodium methanethiolate in 800 mL water. After stirring overnight, the clear solution was poured into 30 L water, the precipitate was filtered, washed with water and dried in vacuo to yield 301 g (70%) of the title compound.

Example 61

3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-6-[3-(1,1-dioxidothiomorpholin-4-yl)prop-1-yn-1-yl]-N-(3,3,3-trifluoropropyl)imidazo[1,2-a]pyrazin-8-amine




200 mg (407 μ mol)

6-bromo-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-N-(3,3,3-trifluoropropyl)imidazo[1,2-a]pyrazin-8-amine which was prepared according to intermediate example 57a were transformed in analogy to example 1 using 4-(prop-2-yn-1-yl)thiomorpholine 1,1-dioxide to give after working up and purification 48.6 mg (19%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 0.95-1.06 (4H), 2.01 (1H), 2.55 (2H), 3.09-3.19 (8H), 3.46 (1H), 3.64 (2H), 3.88 (2H), 6.48 (1H), 7.23 (1H), 7.48 (2H), 7.55 (1H), 7.79 (1H), 7.82 (2H) ppm.

Example 62

N-cyclopropyl-4-{6-[3-(1,1-dioxidothiomorpholin-4-yl)prop-1-yn-1-yl]-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide



150 mg (311 μ mol)


4-{6-bromo-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-2-methylbenzamide which was prepared according to intermediate example 62a were transformed in analogy to example 1 using

4-(prop-2-yn-1-yl)thiomorpholine 1,1-dioxide to give after working up and purification 74.7 mg (40%) of the title compound.

¹H -NMR (CDCl₃): δ= 0.64 (2H), 0.91 (2H), 2.47-2.66 (2H), 2.53 (3H), 2.94 (1H), 3.08-3.20 (8H), 3.67 (2H), 3.92 (2H), 5.97 (1H), 6.25 (1H), 7.34 (1H), 7.35 (1H), 7.49 (1H), 7.56 (1H), 7.72 (1H) ppm.

Intermediate Example 62a


4-{6-bromo-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-2-methylbenzamide



N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)-2-methylbenzamide which was prepared according to intermediate example 62b were transformed in analogy to example 1a using 3,3,3-trifluoropropan-1-amine to give after working up and purification 464 mg (90%) of the title compound.

Intermediate Example 62b

N-cyclopropyl-4-(6,8-dibromoimidazo[1,2-a]pyrazin-3-yl)-2-methylbenzamide




1.8 g (4.47 mmol) 6,8-dibromo-3-iodoimidazo[1,2-a]pyrazine which was prepared according to intermediate example 1c were transformed in analogy to example 1b using

N-cyclopropyl-2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide to give after working up and purification 968 mg (48%) of the title compound.

5 Example 63

4-{6-Ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N,2-dimethylbenzamide




A mixture comprising 13.5 mg (35 μ mol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a, 26 μ L methanamine (2M in tetrahydrofuran), 19.8 mg N-[(dimethylamino)(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yloxy)methylene]-N-methylmethanaminium hexafluorophosphate, 6.37 mg N,N-dimethylpyridin-4-amine and 0.83 mL N,N-dimethylformamide was stirred at 23°C for 1 hour. The solvent was removed and the residue purified by chromatography to give 11.8 mg (85%) of title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 2.54 (3H), 2.57 (2H), 3.05 (3H), 3.08 (1H), 3.93 (2H), 5.81 (1H), 6.24 (1H), 7.36 (1H), 7.38 (1H), 7.52 (1H), 7.58 (1H), 7.82 (1H) ppm.

20 Example 63a

4-{6-Ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid




A mixture comprising 282 mg (700 μ mol) methyl 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate which was prepared according to intermediate example 63b, 3.1 mL aqueous lithium hydroxide (5%), 13.3 mL tetrahydrofuran and 3.6 mL methanol was stirred at 23 °C for 5.5 hours. The mixture was acidified with hydrochloric acid and extracted with ethyl acetate and methanol. The solvent of the organic layer was removed and the residue washed with ether to give 272 mg (100%) of the title compound.

10

Example 63b

Methyl 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate




A mixture comprising 410 mg (864 μ mol) methyl 2-methyl-4-{8-[(3,3,3-trifluoropropyl)amino]-6-[(trimethylsilyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzoate which was prepared according to intermediate example 63c, 842 μ L tetrabutylammoniumfluorid (1M in tetrahydrofuran) and 4.0 mL tetrahydrofuran was stirred at 23 °C for 15 minutes. Saturated aqueous ammonium chloride solution was added and the mixture extracted with ethyl acetate. The organic layer was washed with water and brine and dried over

20

sodium sulfate. After filtration and removal of the solvent the residue was purified by chromatography to give 306 mg (90%) of the title compound.

Example 63c


5 Methyl 2-methyl-4-{8-[(3,3,3-trifluoropropyl)amino]-6-[(trimethylsilyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzoate



A mixture comprising 396 mg (866 μ mol) methyl 4-{6-bromo-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate which was prepared according to intermediate example 63d, 60.8 mg dichloropalladium - triphenylphosphine (1:2), 16.5 mg copper(1+) iodide, 7.9 mL N-isopropylpropan-2-amine, 7.9 mL 1,4-dioxane and 340 mg ethynyl(trimethyl)silane was heated at 70°C under microwave irradiation for 5 minutes. The solvents were removed and the residue purified by chromatography to give 411 mg (95%) of the title compound.

Example 63d


Methyl 4-{6-bromo-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate



1.06 g (2.50 mmol) methyl 4-[6-bromo-8-(methylsulfonyl)imidazo[1,2-a]pyrazin-3-yl]-2-methylbenzoate which was prepared according to intermediate example 63e were transformed in analogy to example 60b using 3,3,3-trifluoropropan-1-amine to give after working up and purification 894 mg (74%) of the title compound.

Example 63e

Methyl 4-[6-bromo-8-(methylsulfonyl)imidazo[1,2-a]pyrazin-3-yl]-2-methylbenzoate




1.34 g (3.42 mmol) methyl 4-[6-bromo-8-(methylsulfonyl)imidazo[1,2-a]pyrazin-3-yl]-2-methylbenzoate which was prepared according to intermediate example 63f were transformed in analogy to example 60c to give after working up and purification 1.12 g (77%) of the title compound.

15

Example 63f

Methyl 4-[6-bromo-8-(methylsulfonyl)imidazo[1,2-a]pyrazin-3-yl]-2-methylbenzoate




20 4.00 g (10.8 mmol) 6-bromo-3-iodo-8-(methylsulfonyl)imidazo[1,2-a]pyrazine which was prepared according to intermediate example 60d were transformed in analogy to example 1b using [4-(methoxycarbonyl)-3-methylphenyl]boronic

acid to give after working up and purification 1.48 g (31%) of the title compound.

Example 64

5 N-ethyl-4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




13.5 mg (35 μ mol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to
 10 intermediate example 63a were transformed in analogy to example 63 using ethanamine to give after working up and purification 14 mg (100%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 1.28 (3H), 2.54 (3H), 2.57 (2H), 3.08 (1H), 3.53 (2H), 3.93 (2H), 5.77 (1H), 6.24 (1H), 7.36 (1H), 7.38 (1H), 7.52 (1H), 7.58 (1H), 7.82 (1H)
 15 ppm.

Example 65

N-(2,2-difluoroethyl)-4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




20

13.5 mg (35 μ mol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a were transformed in analogy to example 63 using 2,2-difluoroethanamine to give after working up and purification 11.4 mg (73%) of the title compound.

$^1\text{H-NMR}$ (CDCl_3): δ = 2.55 (3H), 2.55 (2H), 3.08 (1H), 3.87 (2H), 3.94 (2H), 6.02 (1H), 6.06 (1H), 6.24 (1H), 7.40 (1H), 7.41 (1H), 7.56 (1H), 7.59 (1H), 7.82 (1H) ppm.

10 Example 66

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methyl-N-(1-methylcyclopropyl)benzamide




13.5 mg (35 μ mol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a were transformed in analogy to example 63 using 1-methylcyclopropanamine to give after working up and purification 13.6 mg (89%) of the title compound.

$^1\text{H-NMR}$ (DMSO-d_6): δ = 0.57 (2H), 0.70 (2H), 1.37 (3H), 2.35 (3H), 2.64 (2H), 3.68 (2H), 4.10 (1H), 7.36 (1H), 7.42-7.50 (2H), 7.74 (1H), 7.85 (1H), 7.91 (1H), 8.45 (1H) ppm.

Example 67

N-(1-cyanocyclopropyl)-4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




13.5 mg (35μmol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a were transformed in analogy to example 63 using 1-aminocyclopropanecarbonitrile to give after working up and purification 9.6 mg (61%) of the title compound.

¹H-NMR (DMSO-d₆): δ= 1.25 (2H), 1.53 (2H), 2.39 (3H), 2.64 (2H), 3.68 (2H), 4.11 (1H), 7.45-7.56 (3H), 7.77 (1H), 7.88 (1H), 7.92 (1H), 9.21 (1H) ppm.

10 Example 68

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-(1-methoxycyclopropyl)-2-methylbenzamide




13.5 mg (35μmol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a were transformed in analogy to example 63 using 1-methoxycyclopropanamine to give after working up and purification 7.7 mg (48%) of the title compound.

¹H-NMR (DMSO-d₆): δ= 0.87 (2H), 1.00 (2H), 2.38 (3H), 2.64 (2H), 3.27 (3H), 3.68 (2H), 4.11 (1H), 7.42-7.53 (3H), 7.76 (1H), 7.87 (1H), 7.91 (1H), 9.15 (1H) ppm.

5 Example 69

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-[rel(1S,2S)-2-fluorocyclopropyl]-2-methylbenzamide




13.5 mg (35 μmol) 4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 63a were transformed in analogy to example 63 using rel(1S,2S)-2-fluorocyclopropanamine to give after working up and purification 12.3 mg (79%) of the title compound.

¹H-NMR (DMSO-d₆): δ= 0.96-1.19 (2H), 2.38 (3H), 2.64 (2H), 2.81 (1H), 3.68 (2H), 4.10 (1H), 4.72 (1H), 7.41-7.53 (3H), 7.76 (1H), 7.87 (1H), 7.91 (1H), 8.46 (1H) ppm.

Example 70

4-{6-(3-Hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N,2-dimethylbenzamide




12.8 mg (30 μ mol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 to give after working up and purification 6.4 mg (49%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 2.38 (3H), 2.64 (2H), 2.74 (3H), 3.67 (2H), 4.25 (2H), 5.33 (1H), 7.42-7.52 (3H), 7.75 (1H), 7.80 (1H), 7.91 (1H), 8.22 (1H) ppm.

10 Example 70a


4-{6-(3-Hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid



115 mg (274 μ mol) methyl 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate which was prepared according to intermediate example 70b were transformed in analogy to intermediate example 63a to give after working up and purification 107 mg (93%) of the title compound.

Example 70b

Methyl 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate




- 5 396 mg (866 μ mol) methyl 4-{6-bromo-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoate which was prepared according to intermediate example 63d were transformed in analogy to example 1 to give after working up and purification 152 mg (41%) of the title compound.

10

Example 71

N-ethyl-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




- 15 12.8 mg (30 μ mol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using ethanamine to give after working up and purification 6.7 mg (49%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 1.09 (3H), 2.38 (3H), 2.64 (2H), 3.24 (2H), 3.67 (2H), 4.26 (2H), 5.33 (1H), 7.41-7.52 (3H), 7.75 (1H), 7.79 (1H), 7.91 (1H), 8.28 (1H) ppm.

5 Example 72

N-(2,2-difluoroethyl)-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




12.8 mg (30 μmol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using 2,2-difluoroethanamine to give after working up and purification 9.3 mg (63%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 2.39 (3H), 2.63 (2H), 3.55-3.74 (4H), 4.26 (2H), 5.33 (1H), 6.11 (1H), 7.45-7.56 (3H), 7.77 (1H), 7.82 (1H), 7.92 (1H), 8.71 (1H) ppm.

Example 73

4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methyl-N-(1-methylcyclopropyl)benzamide




12.6 mg (30 μ mol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using 1-methylcyclopropanamine to give after working up and purification 12.7 mg (89%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.57 (2H), 0.70 (2H), 1.37 (3H), 2.35 (3H), 2.63 (2H), 3.68 (2H), 4.26 (2H), 5.31 (1H), 7.36 (1H), 7.42-7.51 (2H), 7.73 (1H), 7.77 (1H), 7.88 (1H), 8.45 (1H) ppm.

10

Example 74

N-(1-cyanocyclopropyl)-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide




12.6 mg (30 μ mol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using 1-aminocyclopropanecarbonitrile to give after working up and purification 5.3 mg (36%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 1.25 (2H), 1.53 (2H), 2.39 (3H), 2.64 (2H), 3.68 (2H), 4.26 (2H), 5.31 (1H), 7.45-7.56 (3H), 7.77 (1H), 7.81 (1H), 7.89 (1H), 9.20 (1H) ppm.

5 Example 75

4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-(1-methoxycyclopropyl)-2-methylbenzamide




12.6 mg (30 μmol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using 1-methoxycyclopropanamine to give after working up and purification 7.4 mg (50%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.87 (2H), 1.00 (2H), 2.38 (3H), 2.64 (2H), 3.27 (3H), 3.68 (2H), 4.25 (2H), 5.31 (1H), 7.41-7.54 (3H), 7.75 (1H), 7.79 (1H), 7.89 (1H), 9.14 (1H) ppm.

Example 76

N-[rel(1S,2S)-2-fluorocyclopropyl]-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide



12.6 mg (30 μ mol) 4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzoic acid which was prepared according to intermediate example 70a were transformed in analogy to example 63 using rel(1S,2S)-2-fluorocyclopropanamine to give after working up and purification 12.1 mg (85%) of the title compound.

$^1\text{H-NMR}$ (DMSO- d_6): δ = 0.95-1.15 (2H), 2.38 (3H), 2.63 (2H), 2.81 (1H), 3.68 (2H), 4.25 (2H), 4.73 (1H), 5.30 (1H), 7.41-7.53 (3H), 7.75 (1H), 7.79 (1H), 7.89 (1H), 8.46 (1H) ppm.

10

Further, the compounds of formula (I) of the present invention can be converted to any salt as described herein, by any method which is known to the person skilled in the art. Similarly, any salt of a compound of formula (I) of the present invention can be converted into the free compound, by any method which is known to the person skilled in the art.

15

Pharmaceutical compositions of the compounds of the invention

This invention also relates to pharmaceutical compositions containing one or more compounds of the present invention. These compositions can be utilised to achieve the desired pharmacological effect by administration to a patient in
5 need thereof. A patient, for the purpose of this invention, is a mammal, including a human, in need of treatment for the particular condition or disease. Therefore, the present invention includes pharmaceutical compositions that are comprised of a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound, or salt thereof, of the
10 present invention. A pharmaceutically acceptable carrier is preferably a carrier that is relatively non-toxic and innocuous to a patient at concentrations consistent with effective activity of the active ingredient so that any side effects ascribable to the carrier do not vitiate the beneficial effects of the active ingredient. A pharmaceutically effective amount of compound is
15 preferably that amount which produces a result or exerts an influence on the particular condition being treated. The compounds of the present invention can be administered with pharmaceutically-acceptable carriers well known in the art using any effective conventional dosage unit forms, including immediate, slow and timed release preparations, orally, parenterally,
20 topically, nasally, ophthalmically, optically, sublingually, rectally, vaginally, and the like.

For oral administration, the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, troches, lozenges, melts, powders, solutions, suspensions, or emulsions, and may be prepared according
25 to methods known to the art for the manufacture of pharmaceutical compositions. The solid unit dosage forms can be a capsule that can be of the ordinary hard- or soft-shelled gelatine type containing, for example, surfactants, lubricants, and inert fillers such as lactose, sucrose, calcium phosphate, and corn starch.

In another embodiment, the compounds of this invention may be tableted with conventional tablet bases such as lactose, sucrose and cornstarch in combination with binders such as acacia, corn starch or gelatine, disintegrating agents intended to assist the break-up and dissolution of the tablet following administration such as potato starch, alginic acid, corn starch, and guar gum, gum tragacanth, acacia, lubricants intended to improve the flow of tablet granulation and to prevent the adhesion of tablet material to the surfaces of the tablet dies and punches, for example talc, stearic acid, or magnesium, calcium or zinc stearate, dyes, colouring agents, and flavouring agents such as peppermint, oil of wintergreen, or cherry flavouring, intended to enhance the aesthetic qualities of the tablets and make them more acceptable to the patient. Suitable excipients for use in oral liquid dosage forms include dicalcium phosphate and diluents such as water and alcohols, for example, ethanol, benzyl alcohol, and polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent or emulsifying agent. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance tablets, pills or capsules may be coated with shellac, sugar or both.

Dispersible powders and granules are suitable for the preparation of an aqueous suspension. They provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example those sweetening, flavouring and colouring agents described above, may also be present.

The pharmaceutical compositions of this invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil such as liquid paraffin or a mixture of vegetable oils. Suitable emulsifying agents may be (1) naturally occurring gums such as gum acacia and gum tragacanth, (2) naturally

occurring phosphatides such as soy bean and lecithin, (3) esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan monooleate, (4) condensation products of said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate. The emulsions may
5 also contain sweetening and flavouring agents.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil such as, for example, arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent such as, for example, beeswax, hard paraffin, or cetyl
10 alcohol. The suspensions may also contain one or more preservatives, for example, ethyl or n-propyl p-hydroxybenzoate ; one or more colouring agents ; one or more flavouring agents ; and one or more sweetening agents such as sucrose or saccharin.

Syrups and elixirs may be formulated with sweetening agents such as, for
15 example, glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, and preservative, such as methyl and propyl parabens and flavouring and colouring agents.

The compounds of this invention may also be administered parenterally, that is, subcutaneously, intravenously, intraocularly, intrasynovially,
20 intramuscularly, or interperitoneally, as injectable dosages of the compound in preferably a physiologically acceptable diluent with a pharmaceutical carrier which can be a sterile liquid or mixture of liquids such as water, saline, aqueous dextrose and related sugar solutions, an alcohol such as ethanol, isopropanol, or hexadecyl alcohol, glycols such as propylene glycol or
25 polyethylene glycol, glycerol ketals such as 2,2-dimethyl-1,1-dioxolane-4-methanol, ethers such as poly(ethylene glycol) 400, an oil, a fatty acid, a fatty acid ester or, a fatty acid glyceride, or an acetylated fatty acid glyceride, with or without the addition of a pharmaceutically acceptable surfactant such as a soap or a detergent,

suspending agent such as pectin, carbomers, methycellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agent and other pharmaceutical adjuvants.

Illustrative of oils which can be used in the parenteral formulations of this invention are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, sesame oil, cottonseed oil, corn oil, olive oil, petrolatum and mineral oil. Suitable fatty acids include oleic acid, stearic acid, isostearic acid and myristic acid. Suitable fatty acid esters are, for example, ethyl oleate and isopropyl myristate. Suitable soaps include fatty acid alkali metal, ammonium, and triethanolamine salts and suitable detergents include cationic detergents, for example dimethyl dialkyl ammonium halides, alkyl pyridinium halides, and alkylamine acetates ; anionic detergents, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates ; non-ionic detergents, for example, fatty amine oxides, fatty acid alkanolamides, and poly(oxyethylene-oxypropylene)s or ethylene oxide or propylene oxide copolymers ; and amphoteric detergents, for example, alkyl-beta-aminopropionates, and 2-alkylimidazoline quarternary ammonium salts, as well as mixtures.

The parenteral compositions of this invention will typically contain from about 0.5% to about 25% by weight of the active ingredient in solution. Preservatives and buffers may also be used advantageously. In order to minimise or eliminate irritation at the site of injection, such compositions may contain a non-ionic surfactant having a hydrophile-lipophile balance (HLB) preferably of from about 12 to about 17. The quantity of surfactant in such formulation preferably ranges from about 5% to about 15% by weight. The surfactant can be a single component having the above HLB or can be a mixture of two or more components having the desired HLB.

Illustrative of surfactants used in parenteral formulations are the class of polyethylene sorbitan fatty acid esters, for example, sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

- 5 The pharmaceutical compositions may be in the form of sterile injectable aqueous suspensions. Such suspensions may be formulated according to known methods using suitable dispersing or wetting agents and suspending agents such as, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone, gum
10 tragacanth and gum acacia ; dispersing or wetting agents which may be a naturally occurring phosphatide such as lecithin, a condensation product of an alkylene oxide with a fatty acid, for example, polyoxyethylene stearate, a condensation product of ethylene oxide with a long chain aliphatic alcohol, for example, heptadeca-ethyleneoxycetanol, a condensation product of ethylene
15 oxide with a partial ester derived from a fatty acid and a hexitol such as polyoxyethylene sorbitol monooleate, or a condensation product of an ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride, for example polyoxyethylene sorbitan monooleate.

The sterile injectable preparation may also be a sterile injectable solution or
20 suspension in a non-toxic parenterally acceptable diluent or solvent. Diluents and solvents that may be employed are, for example, water, Ringer's solution, isotonic sodium chloride solutions and isotonic glucose solutions. In addition, sterile fixed oils are conventionally employed as solvents or suspending media. For this purpose, any bland, fixed oil may be employed including synthetic
25 mono- or diglycerides. In addition, fatty acids such as oleic acid can be used in the preparation of injectables.

A composition of the invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritation excipient which is

solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are, for example, cocoa butter and polyethylene glycol.

Another formulation employed in the methods of the present invention
5 employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art (see, e.g., US Patent No. 5,023,252, issued June 11, 1991, incorporated
10 herein by reference). Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Controlled release formulations for parenteral administration include liposomal, polymeric microsphere and polymeric gel formulations that are known in the art.

15 It may be desirable or necessary to introduce the pharmaceutical composition to the patient via a mechanical delivery device. The construction and use of mechanical delivery devices for the delivery of pharmaceutical agents is well known in the art. Direct techniques for, for example, administering a drug directly to the brain usually involve placement of a drug delivery catheter into
20 the patient's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of agents to specific anatomical regions of the body, is described in US Patent No. 5,011,472, issued April 30, 1991.

The compositions of the invention can also contain other conventional
25 pharmaceutically acceptable compounding ingredients, generally referred to as carriers or diluents, as necessary or desired. Conventional procedures for preparing such compositions in appropriate dosage forms can be utilized. Such ingredients and procedures include those described in the following

references, each of which is incorporated herein by reference: Powell, M.F. *et al.*, "Compendium of Excipients for Parenteral Formulations" PDA Journal of Pharmaceutical Science & Technology **1998**, 52(5), 238-311 ; Strickley, R.G "Parenteral Formulations of Small Molecule Therapeutics Marketed in the
5 United States (1999)-Part-1" PDA Journal of Pharmaceutical Science & Technology **1999**, 53(6), 324-349 ; and Nema, S. *et al.*, "Excipients and Their Use in Injectable Products" PDA Journal of Pharmaceutical Science & Technology **1997**, 51(4), 166-171.

Commonly used pharmaceutical ingredients that can be used as appropriate to
10 formulate the composition for its intended route of administration include:

acidifying agents (examples include but are not limited to acetic acid, citric acid, fumaric acid, hydrochloric acid, nitric acid) ;

alkalinizing agents (examples include but are not limited to ammonia solution, ammonium carbonate, diethanolamine, monoethanolamine,
15 potassium hydroxide, sodium borate, sodium carbonate, sodium hydroxide, triethanolamine, trolamine) ;

adsorbents (examples include but are not limited to powdered cellulose and activated charcoal) ;

aerosol propellants (examples include but are not limited to carbon dioxide,
20 CCl₂F₂, F₂ClC-CClF₂ and CClF₃)

air displacement agents (examples include but are not limited to nitrogen and argon) ;

antifungal preservatives (examples include but are not limited to benzoic acid, butylparaben, ethylparaben, methylparaben, propylparaben, sodium
25 benzoate) ;

antimicrobial preservatives (examples include but are not limited to benzalkonium chloride, benzethonium chloride, benzyl alcohol, cetylpyridinium chloride, chlorobutanol, phenol, phenylethyl alcohol, phenylmercuric nitrate and thimerosal) ;

- 5 **antioxidants** (examples include but are not limited to ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, hypophosphorus acid, monothioglycerol, propyl gallate, sodium ascorbate, sodium bisulfite, sodium formaldehyde sulfoxylate, sodium metabisulfite) ;

- binding materials** (examples include but are not limited to block polymers,
10 natural and synthetic rubber, polyacrylates, polyurethanes, silicones, polysiloxanes and styrene-butadiene copolymers) ;

buffering agents (examples include but are not limited to potassium metaphosphate, dipotassium phosphate, sodium acetate, sodium citrate anhydrous and sodium citrate dihydrate)

- 15 **carrying agents** (examples include but are not limited to acacia syrup, aromatic syrup, aromatic elixir, cherry syrup, cocoa syrup, orange syrup, syrup, corn oil, mineral oil, peanut oil, sesame oil, bacteriostatic sodium chloride injection and bacteriostatic water for injection)

- chelating agents** (examples include but are not limited to edetate disodium
20 and edetic acid)

colourants (examples include but are not limited to FD&C Red No. 3, FD&C Red No. 20, FD&C Yellow No. 6, FD&C Blue No. 2, D&C Green No. 5, D&C Orange No. 5, D&C Red No. 8, caramel and ferric oxide red) ;

clarifying agents (examples include but are not limited to bentonite) ;

emulsifying agents (examples include but are not limited to acacia, cetomacrogol, cetyl alcohol, glyceryl monostearate, lecithin, sorbitan monooleate, polyoxyethylene 50 monostearate) ;

encapsulating agents (examples include but are not limited to gelatin and
5 cellulose acetate phthalate)

flavourants (examples include but are not limited to anise oil, cinnamon oil, cocoa, menthol, orange oil, peppermint oil and vanillin) ;

humectants (examples include but are not limited to glycerol, propylene glycol and sorbitol) ;

10 **levigating agents** (examples include but are not limited to mineral oil and glycerin) ;

oils (examples include but are not limited to arachis oil, mineral oil, olive oil, peanut oil, sesame oil and vegetable oil) ;

ointment bases (examples include but are not limited to lanolin, hydrophilic
15 ointment, polyethylene glycol ointment, petrolatum, hydrophilic petrolatum, white ointment, yellow ointment, and rose water ointment) ;

penetration enhancers (transdermal delivery) (examples include but are not limited to monohydroxy or polyhydroxy alcohols, mono-or polyvalent alcohols, saturated or unsaturated fatty alcohols, saturated or unsaturated fatty esters,
20 saturated or unsaturated dicarboxylic acids, essential oils, phosphatidyl derivatives, cephalin, terpenes, amides, ethers, ketones and ureas)

plasticizers (examples include but are not limited to diethyl phthalate and glycerol) ;

solvents (examples include but are not limited to ethanol, corn oil, cottonseed
25 oil, glycerol, isopropanol, mineral oil, oleic acid, peanut oil, purified water, water for injection, sterile water for injection and sterile water for irrigation) ;

stiffening agents (examples include but are not limited to cetyl alcohol, cetyl esters wax, microcrystalline wax, paraffin, stearyl alcohol, white wax and yellow wax) ;

suppository bases (examples include but are not limited to cocoa butter and
5 polyethylene glycols (mixtures)) ;

surfactants (examples include but are not limited to benzalkonium chloride, nonoxynol 10, octoxynol 9, polysorbate 80, sodium lauryl sulfate and sorbitan mono-palmitate) ;

suspending agents (examples include but are not limited to agar, bentonite,
10 carbomers, carboxymethylcellulose sodium, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, kaolin, methylcellulose, tragacanth and veegum) ;

sweetening agents (examples include but are not limited to aspartame, dextrose, glycerol, mannitol, propylene glycol, saccharin sodium, sorbitol and
15 sucrose) ;

tablet anti-adherents (examples include but are not limited to magnesium stearate and talc) ;

tablet binders (examples include but are not limited to acacia, alginic acid, carboxymethylcellulose sodium, compressible sugar, ethylcellulose, gelatin,
20 liquid glucose, methylcellulose, non-crosslinked polyvinyl pyrrolidone, and pregelatinized starch) ;

tablet and capsule diluents (examples include but are not limited to dibasic calcium phosphate, kaolin, lactose, mannitol, microcrystalline cellulose, powdered cellulose, precipitated calcium carbonate, sodium carbonate,
25 sodium phosphate, sorbitol and starch) ;

tablet coating agents (examples include but are not limited to liquid glucose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose, ethylcellulose, cellulose acetate phthalate and shellac) ;

- 5 **tablet direct compression excipients** (examples include but are not limited to dibasic calcium phosphate) ;

tablet disintegrants (examples include but are not limited to alginic acid, carboxymethylcellulose calcium, microcrystalline cellulose, polacrillin potassium, cross-linked polyvinylpyrrolidone, sodium alginate, sodium starch
10 glycollate and starch) ;

tablet glidants (examples include but are not limited to colloidal silica, corn starch and talc) ;

tablet lubricants (examples include but are not limited to calcium stearate, magnesium stearate, mineral oil, stearic acid and zinc stearate) ;

- 15 **tablet/capsule opaquants** (examples include but are not limited to titanium dioxide) ;

tablet polishing agents (examples include but are not limited to carnuba wax and white wax) ;

thickening agents (examples include but are not limited to beeswax, cetyl
20 alcohol and paraffin) ;

tonicity agents (examples include but are not limited to dextrose and sodium chloride) ;

viscosity increasing agents (examples include but are not limited to alginic acid, bentonite, carbomers, carboxymethylcellulose sodium, methylcellulose,
25 polyvinyl pyrrolidone, sodium alginate and tragacanth) ; and

wetting agents (examples include but are not limited to heptadecaethylene oxycetanol, lecithins, sorbitol monooleate, polyoxyethylene sorbitol monooleate, and polyoxyethylene stearate).

Pharmaceutical compositions according to the present invention can be
5 illustrated as follows:

Sterile IV Solution: A 5 mg/mL solution of the desired compound of this invention can be made using sterile, injectable water, and the pH is adjusted if necessary. The solution is diluted for administration to 1 - 2 mg/mL with sterile 5% dextrose and is administered as an IV infusion over about 60 minutes.

10 Lyophilised powder for IV administration: A sterile preparation can be prepared with (i) 100 - 1000 mg of the desired compound of this invention as a lyophilised powder, (ii) 32- 327 mg/mL sodium citrate, and (iii) 300 - 3000 mg Dextran 40. The formulation is reconstituted with sterile, injectable saline or dextrose 5% to a concentration of 10 to 20 mg/mL, which is further diluted
15 with saline or dextrose 5% to 0.2 - 0.4 mg/mL, and is administered either IV bolus or by IV infusion over 15 - 60 minutes.

Intramuscular suspension: The following solution or suspension can be prepared, for intramuscular injection:

50 mg/mL of the desired, water-insoluble compound of this invention

20 5 mg/mL sodium carboxymethylcellulose

4 mg/mL TWEEN 80

9 mg/mL sodium chloride

9 mg/mL benzyl alcohol

Hard Shell Capsules: A large number of unit capsules are prepared by filling
25 standard two-piece hard galantine capsules each with 100 mg of powdered

active ingredient, 150 mg of lactose, 50 mg of cellulose and 6 mg of magnesium stearate.

Soft Gelatin Capsules: A mixture of active ingredient in a digestible oil such as soybean oil, cottonseed oil or olive oil is prepared and injected by means of a positive displacement pump into molten gelatin to form soft gelatin capsules containing 100 mg of the active ingredient. The capsules are washed and dried. The active ingredient can be dissolved in a mixture of polyethylene glycol, glycerin and sorbitol to prepare a water miscible medicine mix.

Tablets: A large number of tablets are prepared by conventional procedures so that the dosage unit is 100 mg of active ingredient, 0.2 mg. of colloidal silicon dioxide, 5 mg of magnesium stearate, 275 mg of microcrystalline cellulose, 11 mg. of starch, and 98.8 mg of lactose. Appropriate aqueous and non-aqueous coatings may be applied to increase palatability, improve elegance and stability or delay absorption.

Immediate Release Tablets/Capsules: These are solid oral dosage forms made by conventional and novel processes. These units are taken orally without water for immediate dissolution and delivery of the medication. The active ingredient is mixed in a liquid containing ingredient such as sugar, gelatin, pectin and sweeteners. These liquids are solidified into solid tablets or caplets by freeze drying and solid state extraction techniques. The drug compounds may be compressed with viscoelastic and thermoelastic sugars and polymers or effervescent components to produce porous matrices intended for immediate release, without the need of water.

Combination therapies

The compounds of this invention can be administered as the sole pharmaceutical agent or in combination with one or more other pharmaceutical agents where the combination causes no unacceptable adverse effects. The present invention relates also to such combinations. For example, the compounds of this invention can be combined with known anti-hyper-proliferative or other indication agents, and the like, as well as with admixtures and combinations thereof. Other indication agents include, but are not limited to, anti-angiogenic agents, mitotic inhibitors, alkylating agents, anti-metabolites, DNA-intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzyme inhibitors, topoisomerase inhibitors, biological response modifiers, or anti-hormones.

The additional pharmaceutical agent can be afinitor, aldesleukin, alendronic acid, alfaferone, alitretinoin, allopurinol, aloprim, aloxi, altretamine, aminoglutethimide, amifostine, amrubicin, amsacrine, anastrozole, anzmet, aranesp, arglabin, arsenic trioxide, aromasin, 5-azacytidine, azathioprine, BCG or tice BCG, bestatin, betamethasone acetate, betamethasone sodium phosphate, bexarotene, bleomycin sulfate, broxuridine, bortezomib, busulfan, calcitonin, campath, capecitabine, carboplatin, casodex, cefesone, celmoleukin, cerubidine, chlorambucil, cisplatin, cladribine, cladribine, clodronic acid, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, DaunoXome, decadron, decadron phosphate, delestrogen, denileukin diftitox, depo-medrol, deslorelin, dexrazoxane, diethylstilbestrol, diflucan, docetaxel, doxifluridine, doxorubicin, dronabinol, DW-166HC, eligard, elitek, ellence, emend, epirubicin, epoetin alfa, epogen, eptaplatin, ergamisol, estrace, estradiol, estramustine phosphate sodium, ethinyl estradiol, ethyol, etidronic acid, etopophos, etoposide, fadrozole, farston, filgrastim, finasteride, fligrastim, floxuridine, fluconazole, fludarabine, 5-fluorodeoxyuridine monophosphate, 5-fluorouracil (5-FU), fluoxymesterone, flutamide,

formestane, fosteabine, fotemustine, fulvestrant, gammagard, gemcitabine, gemtuzumab, gleevec, gliadel, goserelin, granisetron HCl, histrelin, hycamtin, hydrocortone, eyrthro-hydroxynonyladenine, hydroxyurea, ibritumomab tiuxetan, idarubicin, ifosfamide, interferon alpha, interferon-alpha 2, 5 interferon alfa-2A, interferon alfa-2B, interferon alfa-n1, interferon alfa-n3, interferon beta, interferon gamma-1a, interleukin-2, intron A, iressa, irinotecan, kytril, lentinan sulfate, letrozole, leucovorin, leuprolide, leuprolide acetate, levamisole, levofolinic acid calcium salt, levothroid, levoxyl, lomustine, lonidamine, marinol, mechlorethamine, mecobalamin, 10 medroxyprogesterone acetate, megestrol acetate, melphalan, menest, 6-mercaptopurine, Mesna, methotrexate, metvix, miltefosine, minocycline, mitomycin C, mitotane, mitoxantrone, Modrenal, Myocet, nedaplatin, neulasta, neumega, neupogen, nilutamide, nolvadex, NSC-631570, OCT-43, octreotide, ondansetron HCl, orapred, oxaliplatin, paclitaxel, pediapred, 15 pegaspargase, Pegasys, pentostatin, picibanil, pilocarpine HCl, pirarubicin, plicamycin, porfimer sodium, prednimustine, prednisolone, prednisone, premarin, procarbazine, procrit, raltitrexed, RDEA 119, rebif, rhenium-186 etidronate, rituximab, roferon-A, romurtide, salagen, sandostatin, sargramostim, semustine, sizofiran, sobuzoxane, solu-medrol, sparfosic acid, 20 stem-cell therapy, streptozocin, strontium-89 chloride, synthroid, tamoxifen, tamsulosin, tasonermin, tastolactone, taxotere, teceleukin, temozolomide, teniposide, testosterone propionate, testred, thioguanine, thiotepa, thyrotropin, tiludronic acid, topotecan, toremifene, tositumomab, trastuzumab, treosulfan, tretinoin, trexall, trimethylmelamine, trimetrexate, 25 triptorelin acetate, triptorelin pamoate, UFT, uridine, valrubicin, vesnarinone, vinblastine, vincristine, vindesine, vinorelbine, virulizin, zinecard, zinostatin stimalamer, zofran, ABI-007, acolbifene, actimmune, affinitak, aminopterin, arzoxifene, asoprisnil, atamestane, atrasentan, sorafenib (BAY 43-9006), avastin, CCI-779, CDC-501, celebrex, cetuximab, crisnatol, cyproterone 30 acetate, decitabine, DN-101, doxorubicin-MTC, dSLIM, dutasteride, edotecarin,

eflornithine, exatecan, fenretinide, histamine dihydrochloride, histrelin hydrogel implant, holmium-166 DOTMP, ibandronic acid, interferon gamma, intron-PEG, ixabepilone, keyhole limpet hemocyanin, L-651582, lanreotide, lasofoxifene, libra, lonafarnib, miproxifene, minodronate, MS-209, liposomal
5 MTP-PE, MX-6, nafarelin, nemorubicin, neovastat, nolatrexed, oblimersen, onco-TCS, osidem, paclitaxel polyglutamate, pamidronate disodium, PN-401, QS-21, quazepam, R-1549, raloxifene, ranpirnase, 13-cis -retinoic acid, satraplatin, seocalcitol, T-138067, tarceva, taxoprexin, thymosin alpha 1, tiazofurine, tipifarnib, tirapazamine, TLK-286, toremifene, TransMID-107R,
10 valspodar, vapreotide, vatalanib, verteporfin, vinflunine, Z-100, zoledronic acid or combinations thereof.

Optional anti-hyper-proliferative agents which can be added to the composition include but are not limited to compounds listed on the cancer chemotherapy drug regimens in the 11th Edition of the Merck Index, (1996),
15 which is hereby incorporated by reference, such as asparaginase, bleomycin, carboplatin, carmustine, chlorambucil, cisplatin, colaspase, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, daunorubicin, doxorubicin (adriamycine), epirubicin, epothilone, an epothilone derivative, etoposide, 5-fluorouracil, hexamethylmelamine, hydroxyurea, ifosfamide,
20 irinotecan, leucovorin, lomustine, mechlorethamine, 6-mercaptopurine, mesna, methotrexate, mitomycin C, mitoxantrone, prednisolone, prednisone, procarbazine, raloxifen, streptozocin, tamoxifen, thioguanine, topotecan, vinblastine, vincristine, and vindesine.

Other anti-hyper-proliferative agents suitable for use with the composition of
25 the invention include but are not limited to those compounds acknowledged to be used in the treatment of neoplastic diseases in Goodman and Gilman's The Pharmacological Basis of Therapeutics (Ninth Edition), editor Molinoff *et al.*, publ. by McGraw-Hill, pages 1225-1287, (1996), which is hereby incorporated by reference, such as aminoglutethimide, L-asparaginase, azathioprine,

5-azacytidine, cladribine, busulfan, diethylstilbestrol, 2',2'-difluorodeoxycytidine, docetaxel, erythrohydroxynonyl adenine, ethinyl estradiol, 5-fluorodeoxyuridine, 5-fluorodeoxyuridine monophosphate, fludarabine phosphate, fluoxymesterone, flutamide, hydroxyprogesterone caproate, idarubicin, interferon, medroxyprogesterone acetate, megestrol acetate, melphalan, mitotane, paclitaxel, pentostatin, N-phosphonoacetyl-L-aspartate (PALA), plicamycin, semustine, teniposide, testosterone propionate, thiotepa, trimethylmelamine, uridine, and vinorelbine.

- 10 Other anti-hyper-proliferative agents suitable for use with the composition of the invention include but are not limited to other anti-cancer agents such as epothilone and its derivatives, irinotecan, raloxifen and topotecan.

The compounds of the invention may also be administered in combination with protein therapeutics. Such protein therapeutics suitable for the treatment of cancer or other angiogenic disorders and for use with the compositions of the invention include, but are not limited to, an interferon (e.g., interferon .alpha., .beta., or .gamma.) supraagonistic monoclonal antibodies, Tuebingen, TRP-1 protein vaccine, Colostrinin, anti-FAP antibody, YH-16, gemtuzumab, infliximab, cetuximab, trastuzumab, denileukin diftitox, rituximab, thymosin alpha 1, bevacizumab, mecasermin, mecasermin rinfabate, oprelvekin, natalizumab, rhMBL, MFE-CP1 + ZD-2767-P, ABT-828, ErbB2-specific immunotoxin, SGN-35, MT-103, rinfabate, AS-1402, B43-genistein, L-19 based radioimmunotherapeutics, AC-9301, NY-ESO-1 vaccine, IMC-1C11, CT-322, rhCC10, r(m)CRP, MORAb-009, aviscumine, MDX-1307, Her-2 vaccine, APC-8024, NGR-hTNF, rhH1.3, IGN-311, Endostatin, volociximab, PRO-1762, lexatumumab, SGN-40, pertuzumab, EMD-273063, L19-IL-2 fusion protein, PRX-321, CNTO-328, MDX-214, tigapotide, CAT-3888, labetuzumab, alpha-particle-emitting radioisotope-linked lintuzumab, EM-1421, HyperAcute vaccine, tucotuzumab celmoleukin, galiximab, HPV-16-E7, Javelin - prostate

cancer, Javelin - melanoma, NY-ESO-1 vaccine, EGF vaccine, CYT-004-MelQbG10, WT1 peptide, oregovomab, ofatumumab, zalutumumab, cintredekin besudotox, WX-G250, Albuferon, aflibercept, denosumab, vaccine, CTP-37, efungumab, or 131I-chTNT-1/B. Monoclonal antibodies useful as the
 5 protein therapeutic include, but are not limited to, muromonab-CD3, abciximab, edrecolomab, daclizumab, gentuzumab, alemtuzumab, ibritumomab, cetuximab, bevicizumab, efalizumab, adalimumab, omalizumab, muromomab-CD3, rituximab, daclizumab, trastuzumab, palivizumab, basiliximab, and infliximab.

10 The compounds of the invention may also be combined with biological therapeutic agents, such as antibodies (e.g. avastin, rituxan, erbitux, herceptin), or recombinant proteins.

The compounds of the invention may also be in combination with antiangiogenesis agents, such as, for example, with avastin, axitinib, DAST,
 15 recentin, sorafenib or sunitinib. Combinations with inhibitors of proteasomes or mTOR inhibitors, or anti-hormones or steroidal metabolic enzyme inhibitors are also possible.

Generally, the use of cytotoxic and/or cytostatic agents in combination with a compound or composition of the present invention will serve to:

- 20 (1) yield better efficacy in reducing the growth of a tumour or even eliminate the tumour as compared to administration of either agent alone,
- (2) provide for the administration of lesser amounts of the administered chemotherapeutic agents,
- 25 (3) provide for a chemotherapeutic treatment that is well tolerated in the patient with fewer deleterious pharmacological complications than

observed with single agent chemotherapies and certain other combined therapies,

- (4) provide for treating a broader spectrum of different cancer types in mammals, especially humans,
- 5 (5) provide for a higher response rate among treated patients,
- (6) provide for a longer survival time among treated patients compared to standard chemotherapy treatments,
- (7) provide a longer time for tumour progression, and/or
- (8) yield efficacy and tolerability results at least as good as those of the
10 agents used alone, compared to known instances where other cancer agent combinations produce antagonistic effects.

Methods of Sensitizing Cells to Radiation

In a distinct embodiment of the present invention, a compound of the present
15 invention may be used to sensitize a cell to radiation. That is, treatment of a cell with a compound of the present invention prior to radiation treatment of the cell renders the cell more susceptible to DNA damage and cell death than the cell would be in the absence of any treatment with a compound of the invention. In one aspect, the cell is treated with at least one compound of the
20 invention.

Thus, the present invention also provides a method of killing a cell, wherein a cell is administered one or more compounds of the invention in combination with conventional radiation therapy.

The present invention also provides a method of rendering a cell more
25 susceptible to cell death, wherein the cell is treated one or more compounds

of the invention prior to the treatment of the cell to cause or induce cell death. In one aspect, after the cell is treated with one or more compounds of the invention, the cell is treated with at least one compound, or at least one method, or a combination thereof, in order to cause DNA damage for the purpose of inhibiting the function of the normal cell or killing the cell.

In one embodiment, a cell is killed by treating the cell with at least one DNA damaging agent. That is, after treating a cell with one or more compounds of the invention to sensitize the cell to cell death, the cell is treated with at least one DNA damaging agent to kill the cell. DNA damaging agents useful in the present invention include, but are not limited to, chemotherapeutic agents (e.g., cisplatin), ionizing radiation (X-rays, ultraviolet radiation), carcinogenic agents, and mutagenic agents.

In another embodiment, a cell is killed by treating the cell with at least one method to cause or induce DNA damage. Such methods include, but are not limited to, activation of a cell signalling pathway that results in DNA damage when the pathway is activated, inhibiting of a cell signalling pathway that results in DNA damage when the pathway is inhibited, and inducing a biochemical change in a cell, wherein the change results in DNA damage. By way of a non-limiting example, a DNA repair pathway in a cell can be inhibited, thereby preventing the repair of DNA damage and resulting in an abnormal accumulation of DNA damage in a cell.

In one aspect of the invention, a compound of the invention is administered to a cell prior to the radiation or other induction of DNA damage in the cell. In another aspect of the invention, a compound of the invention is administered to a cell concomitantly with the radiation or other induction of DNA damage in the cell. In yet another aspect of the invention, a compound of the invention is administered to a cell immediately after radiation or other induction of DNA damage in the cell has begun.

In another aspect, the cell is *in vitro*. In another embodiment, the cell is *in vivo*.

As mentioned *supra*, the compounds of the present invention have surprisingly been found to effectively inhibit Mps-1 and may therefore be used for the treatment or prophylaxis of diseases of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses, or diseases which are accompanied with uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses, particularly in which the uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses is mediated by Mps-1, such as, for example, haematological tumours, solid tumours, and/or metastases thereof, e.g. leukaemias and myelodysplastic syndrome, malignant lymphomas, head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung tumours, gastrointestinal tumours, endocrine tumours, mammary and other gynaecological tumours, urological tumours including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

20

In accordance with another aspect therefore, the present invention covers a compound of general formula (I), or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a pharmaceutically acceptable salt thereof, or a mixture of same, as described and defined herein, for use in the treatment or prophylaxis of a disease, as mentioned *supra*.

Another particular aspect of the present invention is therefore the use of a compound of general formula (I), described *supra*, or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a

30

pharmaceutically acceptable salt thereof, or a mixture of same, for the prophylaxis or treatment of a disease.

Another particular aspect of the present invention is therefore the use of a
5 compound of general formula (I) described *supra* for manufacturing a pharmaceutical composition for the treatment or prophylaxis of a disease.

The diseases referred to in the two preceding paragraphs are diseases of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular
10 immune responses, or inappropriate cellular inflammatory responses, or diseases which are accompanied with uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses, particularly in which the uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses,
15 or inappropriate cellular inflammatory responses is mediated by Mps-1, such as, for example, haematological tumours, solid tumours, and/or metastases thereof, e.g. leukaemias and myelodysplastic syndrome, malignant lymphomas, head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung
20 tumours, gastrointestinal tumours, endocrine tumours, mammary and other gynaecological tumours, urological tumours including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

The term “inappropriate” within the context of the present invention, in
25 particular in the context of “inappropriate cellular immune responses, or inappropriate cellular inflammatory responses”, as used herein, is to be understood as preferably meaning a response which is less than, or greater than normal, and which is associated with, responsible for, or results in, the pathology of said diseases.

30

Preferably, the use is in the treatment or prophylaxis of diseases, wherein the diseases are haematological tumours, solid tumours and/or metastases thereof.

5 Method of treating hyper-proliferative disorders

The present invention relates to a method for using the compounds of the present invention and compositions thereof, to treat mammalian hyper-proliferative disorders. Compounds can be utilized to inhibit, block, reduce, decrease, etc., cell proliferation and/or cell division, and/or produce
10 apoptosis. This method comprises administering to a mammal in need thereof, including a human, an amount of a compound of this invention, or a pharmaceutically acceptable salt, isomer, polymorph, metabolite, hydrate, solvate or ester thereof ; etc. which is effective to treat the disorder. Hyper-proliferative disorders include but are not limited, e.g., psoriasis,
15 keloids, and other hyperplasias affecting the skin, benign prostate hyperplasia (BPH), solid tumours, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid and their distant metastases. Those disorders also include lymphomas, sarcomas, and leukaemias.

20 Examples of breast cancer include, but are not limited to invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma in situ, and lobular carcinoma in situ.

Examples of cancers of the respiratory tract include, but are not limited to small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma
25 and pleuropulmonary blastoma.

Examples of brain cancers include, but are not limited to brain stem and hypothalamic glioma, cerebellar and cerebral astrocytoma, medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumour.

Tumours of the male reproductive organs include, but are not limited to prostate and testicular cancer. Tumours of the female reproductive organs include, but are not limited to endometrial, cervical, ovarian, vaginal, and vulvar cancer, as well as sarcoma of the uterus.

- 5 Tumours of the digestive tract include, but are not limited to anal, colon, colorectal, oesophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers.

Tumours of the urinary tract include, but are not limited to bladder, penile, kidney, renal pelvis, ureter, urethral and human papillary renal cancers.

- 10 Eye cancers include, but are not limited to intraocular melanoma and retinoblastoma.

Examples of liver cancers include, but are not limited to hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed

- 15 hepatocellular cholangiocarcinoma.

Skin cancers include, but are not limited to squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer.

- Head-and-neck cancers include, but are not limited to laryngeal, 20 hypopharyngeal, nasopharyngeal, oropharyngeal cancer, lip and oral cavity cancer and squamous cell. Lymphomas include, but are not limited to AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Burkitt lymphoma, Hodgkin's disease, and lymphoma of the central nervous system.

- 25 Sarcomas include, but are not limited to sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma.

Leukemias include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia.

These disorders have been well characterized in humans, but also exist with a similar etiology in other mammals, and can be treated by administering pharmaceutical compositions of the present invention.

The term “treating” or “treatment” as stated throughout this document is used conventionally, *e.g.*, the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of, *etc.*, of a disease or disorder, such as a carcinoma.

Methods of treating kinase disorders

The present invention also provides methods for the treatment of disorders associated with aberrant mitogen extracellular kinase activity, including, but not limited to stroke, heart failure, hepatomegaly, cardiomegaly, diabetes, Alzheimer’s disease, cystic fibrosis, symptoms of xenograft rejections, septic shock or asthma.

Effective amounts of compounds of the present invention can be used to treat such disorders, including those diseases (*e.g.*, cancer) mentioned in the Background section above. Nonetheless, such cancers and other diseases can be treated with compounds of the present invention, regardless of the mechanism of action and/or the relationship between the kinase and the disorder.

The phrase “aberrant kinase activity” or “aberrant tyrosine kinase activity,” includes any abnormal expression or activity of the gene encoding the kinase or of the polypeptide it encodes. Examples of such aberrant activity, include, but are not limited to, over-expression of the gene or polypeptide ; gene

amplification ; mutations which produce constitutively-active or hyperactive kinase activity ; gene mutations, deletions, substitutions, additions, etc.

The present invention also provides for methods of inhibiting a kinase activity, especially of mitogen extracellular kinase, comprising administering an effective amount of a compound of the present invention, including salts, polymorphs, metabolites, hydrates, solvates, prodrugs (e.g.: esters) thereof, and diastereoisomeric forms thereof. Kinase activity can be inhibited in cells (e.g., *in vitro*), or in the cells of a mammalian subject, especially a human patient in need of treatment.

10

Methods of treating angiogenic disorders

The present invention also provides methods of treating disorders and diseases associated with excessive and/or abnormal angiogenesis.

Inappropriate and ectopic expression of angiogenesis can be deleterious to an organism. A number of pathological conditions are associated with the growth of extraneous blood vessels. These include, e.g., diabetic retinopathy, ischemic retinal-vein occlusion, and retinopathy of prematurity [Aiello et al. New Engl. J. Med. **1994**, 331, 1480 ; Peer et al. Lab. Invest. **1995**, 72, 638], age-related macular degeneration [AMD ; see, Lopez et al. Invest. Ophthalmol. Vis. Sci. **1996**, 37, 855], neovascular glaucoma, psoriasis, retrolental fibroplasias, angiofibroma, inflammation, rheumatoid arthritis (RA), restenosis, in-stent restenosis, vascular graft restenosis, etc. In addition, the increased blood supply associated with cancerous and neoplastic tissue, encourages growth, leading to rapid tumour enlargement and metastasis. Moreover, the growth of new blood and lymph vessels in a tumour provides an escape route for renegade cells, encouraging metastasis and the consequence spread of the cancer. Thus, compounds of the present invention can be utilized to treat and/or prevent any of the aforementioned angiogenesis disorders,

25

e.g., by inhibiting and/or reducing blood vessel formation ; by inhibiting, blocking, reducing, decreasing, etc. endothelial cell proliferation or other types involved in angiogenesis, as well as causing cell death or apoptosis of such cell types.

5

Dose and administration

Based upon standard laboratory techniques known to evaluate compounds useful for the treatment of hyper-proliferative disorders and angiogenic disorders, by standard toxicity tests and by standard pharmacological assays for the determination of treatment of the conditions identified above in mammals, and by comparison of these results with the results of known medicaments that are used to treat these conditions, the effective dosage of the compounds of this invention can readily be determined for treatment of each desired indication. The amount of the active ingredient to be administered in the treatment of one of these conditions can vary widely according to such considerations as the particular compound and dosage unit employed, the mode of administration, the period of treatment, the age and sex of the patient treated, and the nature and extent of the condition treated.

The total amount of the active ingredient to be administered will generally range from about 0.001 mg/kg to about 200 mg/kg body weight per day, and preferably from about 0.01 mg/kg to about 20 mg/kg body weight per day. Clinically useful dosing schedules will range from one to three times a day dosing to once every four weeks dosing. In addition, "drug holidays" in which a patient is not dosed with a drug for a certain period of time, may be beneficial to the overall balance between pharmacological effect and tolerability. A unit dosage may contain from about 0.5 mg to about 1500 mg of active ingredient, and can be administered one or more times per day or less than once a day.

The average daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily rectal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily vaginal dosage regimen will preferably be from 0.01 to 200 mg/kg of total body weight. The average daily topical dosage regimen will preferably be from 0.1 to 200 mg administered between one to four times daily. The transdermal concentration will preferably be that required to maintain a daily dose of from 0.01 to 200 mg/kg. The average daily inhalation dosage regimen will preferably be from 0.01 to 100 mg/kg of total body weight.

Of course the specific initial and continuing dosage regimen for each patient will vary according to the nature and severity of the condition as determined by the attending diagnostician, the activity of the specific compound employed, the age and general condition of the patient, time of administration, route of administration, rate of excretion of the drug, drug combinations, and the like. The desired mode of treatment and number of doses of a compound of the present invention or a pharmaceutically acceptable salt or ester or composition thereof can be ascertained by those skilled in the art using conventional treatment tests.

Preferably, the diseases of said method are haematological tumours, solid tumour and/or metastases thereof.

The compounds of the present invention can be used in particular in therapy and prevention, i.e. prophylaxis, of tumour growth and metastases, especially in solid tumours of all indications and stages with or without pre-treatment of the tumour growth.

Methods of testing for a particular pharmacological or pharmaceutical property are well known to persons skilled in the art.

The example testing experiments described herein serve to illustrate the present invention and the invention is not limited to the examples given.

10 **Biological assay: Proliferation Assay**

Cultivated tumour cells (MCF7, hormone dependent human mammary carcinoma cells, ATCC HTB22; NCI-H460, human non-small cell lung carcinoma cells, ATCC HTB-177; DU 145, hormone-independent human prostate carcinoma cells, ATCC HTB-81; HeLa-MaTu, human cervical carcinoma cells, EPO-GmbH, Berlin; HeLa-MaTu-ADR, multidrug-resistant human cervical carcinoma cells, EPO-GmbH, Berlin; HeLa human cervical tumour cells, ATCC CCL-2; B16F10 mouse melanoma cells, ATCC CRL-6475) were plated at a density of 5000 cells/well (MCF7, DU145, HeLa-MaTu-ADR), 3000 cells/well (NCI-H460, HeLa-MaTu, HeLa), or 1000 cells/well (B16F10) in a 96-well multititer plate in 200 μ L of their respective growth medium supplemented 10% fetal calf serum. After 24 hours, the cells of one plate (zero-point plate) were stained with crystal violet (see below), while the medium of the other plates was replaced by fresh culture medium (200 μ L), to which the test substances were added in various concentrations (0 μ M, as well as in the range of 0.01-30 μ M; the final concentration of the solvent dimethyl sulfoxide was 0.5%). The cells were incubated for 4 days in the presence of test substances. Cell proliferation was determined by staining the cells with crystal violet: the cells were fixed by adding 20 μ L/measuring point of an 11% glutaric aldehyde solution for 15 minutes at room temperature. After three washing cycles of the fixed cells with water, the plates were dried at room temperature. The cells were stained

by adding 100 µl/measuring point of a 0.1% crystal violet solution (pH 3.0). After three washing cycles of the stained cells with water, the plates were dried at room temperature. The dye was dissolved by adding 100 µl/measuring point of a 10% acetic acid solution. The extinction was determined by photometry at a wavelength of 595 nm. The change of cell number, in percent, was calculated by normalization of the measured values to the extinction values of the zero-point plate (=0%) and the extinction of the untreated (0 µm) cells (=100%). The IC50 values were determined by means of a 4 parameter fit using the company's own software.

10

Mps-1 kinase assay

The human kinase Mps-1 phosphorylates a biotinylated substrate peptide. Detection of the phosphorylated product is achieved by time-resolved fluorescence resonance energy transfer (TR-FRET) from Europium-labelled anti-phospho-Serine/Threonine antibody as donor to streptavidin labelled with cross-linked allophycocyanin (SA-XLent) as acceptor. Compounds are tested for their inhibition of the kinase activity.

N-terminally GST-tagged human full length recombinant Mps-1 kinase (purchased from Invitrogen, Karlsruhe, Germany, cat. no PV4071) was used. As substrate for the kinase reaction a biotinylated peptide of the amino-acid sequence PWDPDADITEILG (C-terminus in amide form, purchased from Biosynthan GmbH, Berlin) was used.

For the assay 50 nL of a 100-fold concentrated solution of the test compound in DMSO was pipetted into a black low volume 384well microtiter plate (Greiner Bio-One, Frickenhausen, Germany), 2 µl of a solution of Mps-1 in assay buffer [0.1 mM sodium-ortho-vanadate, 10 mM MgCl₂, 2 mM DTT, 25 mM Hepes pH 7.7, 0.05% BSA, 0.001% Pluronic F-127] were added and the mixture was incubated for 15 min at 22°C to allow pre-binding of the test compounds to

Mps-1 before the start of the kinase reaction. Then the kinase reaction was started by the addition of 3 μ l of a solution of 16.7 μ M adenosine-tri-phosphate (ATP, 16.7 μ M => final conc. in the 5 μ l assay volume is 10 μ M) and peptide substrate (1.67 μ M => final conc. in the 5 μ l assay volume is 1 μ M) in assay
5 buffer and the resulting mixture was incubated for a reaction time of 60 min at 22°C. The concentration of Mps-1 in the assay was adjusted to the activity of the enzyme lot and was chosen appropriate to have the assay in the linear range, typical enzyme concentrations were in the range of about 1 nM (final conc. in the 5 μ l assay volume). The reaction was stopped by the addition of 3
10 μ l of a solution of HTRF detection reagents (100 mM Hepes pH 7.4, 0.1% BSA, 40 mM EDTA, 140 nM Streptavidin-XLent [# 61GSTXLB, Fa. Cis Biointernational, Marcoule, France], 1.5 nM anti-phospho(Ser/Thr)-Europium-antibody [#AD0180, PerkinElmer LAS, Rodgau-Jügesheim, Germany].

15 The resulting mixture was incubated 1 h at 22°C to allow the binding of the phosphorylated peptide to the anti-phospho(Ser/Thr)-Europium-antibody. Subsequently the amount of phosphorylated substrate was evaluated by measurement of the resonance energy transfer from the Europium-labelled anti-phospho(Ser/Thr) antibody to the Streptavidin-XLent. Therefore, the
20 fluorescence emissions at 620 nm and 665 nm after excitation at 350 nm was measured in a Viewlux TR-FRET reader (PerkinElmer LAS, Rodgau-Jügesheim, Germany). The “blank-corrected normalized ratio” (a Viewlux specific readout, similar to the traditional ratio of the emissions at 665 nm and at 622 nm, in which blank and Eu-donor crosstalk are subtracted from the 665 nm
25 signal before the ratio is calculated) was taken as the measure for the amount of phosphorylated substrate. The data were normalised (enzyme reaction without inhibitor = 0 % inhibition, all other assay components but no enzyme = 100 % inhibition). Test compounds were tested on the same microtiter plate at
30 10 different concentrations in the range of 20 μ M to 1 nM (20 μ M, 6.7 μ M, 2.2 μ M, 0.74 μ M, 0.25 μ M, 82 nM, 27 nM, 9.2 nM, 3.1 nM and 1 nM, dilution

series prepared before the assay at the level of the 100fold conc. stock solutions by serial 1:3 dilutions) in duplicate values for each concentration and IC_{50} values were calculated by a 4 parameter fit using an in-house software.

Table

Example	Mps-1 IC50 [nM]	Example	Mps-1 IC50 [nM]
1	31	39	11
2	20000	40	60
3	20000	41	4.0
4	1840	42	17
5	4.8	43	2.4
6	26	44	5.5
7	36	45	1.8
8	22	46	2.5
9	10	47	17
10	38	48	10
11	993	49	14
12	9.9	50	130
13	2.5	51	15
14	7.6	52	13
15	4.1	53	18
16	15	54	8.4
17	17	55	520
18	3.1	56	23
19	9.6	57	7.1
20	4.8	58	5.3
21	1.0	59	53
22	5.6	60	230
23	3.7	61	6.8
24	6.7	62	4.2
25	4.8	63	135
26	13	64	120
27	119	65	215
28	6.1	66	155
29	1.7	67	165
30	64	68	305
31	3.4	69	36
32	6.7	70	69
33	7.2	71	150
34	106	72	130
35	14	73	120
36	3.9	74	98
37	1.2	75	310
38	63	76	43

Spindle Assembly Checkpoint Assay

The spindle assembly checkpoint assures the proper segregation of chromosomes during mitosis. Upon entry into mitosis, chromosomes begin to condensate which is accompanied by the phosphorylation of histone H3 on serine 10. Dephosphorylation of histone H3 on serine 10 begins in anaphase and ends at early telophase. Accordingly, phosphorylation of histone H3 on serine 10 can be utilized as a marker of cells in mitosis. Nocodazole is a microtubule destabilizing substance. Thus, nocodazole interferes with microtubule dynamics and mobilises the spindle assembly checkpoint. The cells arrest in mitosis at G2/M transition and exhibit phosphorylated histone H3 on serine 10. An inhibition of the spindle assembly checkpoint by Mps-1 inhibitors overrides the mitotic blockage in the presence of nocodazole, and the cells complete mitosis prematurely. This alteration is detected by the decrease of cells with phosphorylation of histone H3 on serine 10. This decline is used as a marker to determine the capability of compounds of the present invention to induce a mitotic breakthrough.

Cultivated cells of the human cervical tumour cell line HeLa (ATCC CCL-2) were plated at a density of 2500 cells/well in a 384-well microtiter plate in 20 μ l Dulbecco's Medium (w/o phenol red, w/o sodium pyruvate, w 1000 mg/ml glucose, w pyridoxine) supplemented with 1% (v/v) glutamine, 1% (v/v) penicillin, 1% (v/v) streptomycin and 10% (v/v) fetal calf serum. After incubation overnight at 37°C, 10 μ l/well nocodazole at a final concentration of 0.1 μ g/ml were added to cells. After 24 h incubation, cells were arrested at G2/M phase of the cell cycle progression. Test compounds solubilised in dimethyl sulfoxide (DMSO) were added at various concentrations (0 μ M, as well as in the range of 0.005 μ M - 10 μ M; the final concentration of the solvent DMSO was 0.5% (v/v)). Cells were incubated for 4 h at 37°C in the presence of test compounds. Thereafter, cells were fixed in 4% (v/v) paraformaldehyde in phosphate buffered saline (PBS) at 4°C overnight then permeabilised in 0.1%

(v/v) Triton XTM 100 in PBS at room temperature for 20 min and blocked in 0.5% (v/v) bovine serum albumin (BSA) in PBS at room temperature for 15 min. After washing with PBS, 20 µl/well antibody solution (anti-phospho-histone H3 clone 3H10, FITC; Upstate, Cat# 16-222; 1:200 dilution) was added to cells, which
5 were incubated for 2 h at room temperature. Afterwards, cells were washed with PBS and 20 µl/well HOECHST 33342 dye solution (5 µg/ml) was added to cells and cells were incubated 12 min at room temperature in the dark. Cells were washed twice with PBS then covered with PBS and stored at 4°C until analysis. Images were acquired with a Perkin Elmer OPERATM High-Content
10 Analysis reader. Images were analyzed with image analysis software MetaXpressTM from Molecular devices utilizing the Cell Cycle application module. In this assay both labels HOECHST 33342 and phosphorylated Histone H3 on serine 10 were measured. HOECHST 33342 labels DNA and is used to count cell number. The staining of phosphorylated Histone H3 on serine 10
15 determines the number of mitotic cells. Inhibition of Mps-1 decreases the number of mitotic cells in the presence of nocodazole indicating an inappropriate mitotic progression. The raw assay data were further analysed by four parameter logistic regression analysis to determine the IC₅₀ value for each tested compound.

20

It will be apparent to persons skilled in the art that assays for other Mps kinases may be performed in analogy using the appropriate reagents.


Thus the compounds of the present invention effectively inhibit one or more
25 Mps-1 kinases and are therefore suitable for the treatment or prophylaxis of diseases of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses, particularly in which the uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or
30 inappropriate cellular inflammatory responses is mediated by Mps-1, more

particularly in which the diseases of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune responses, or inappropriate cellular inflammatory responses are haematological tumours, solid tumours and/or metastases thereof, *e.g.* leukaemias and myelodysplastic syndrome, 5 malignant lymphomas, head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung tumours, gastrointestinal tumours, endocrine tumours, mammary and other gynaecological tumours, urological tumours including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

10



CLAIMS

1. A compound of general formula (I) :




(I)

in which :

A represents a  or  group;

Z represents a -C(=O)N(H)R² or -S(=O)₂N(H)R² group,
or a group selected from :



wherein * indicates the point of attachment of said groups with the rest of the molecule ;

- R¹ represents a hydrogen atom or a halogen atom, or a -CN, C₁-C₆-alkyl-,
-N(R^{6a})R^{6b}, -NO₂, -OR⁶ or -SR⁶ group ;
- R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
5 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
group is optionally substituted, one or more times, identically or
differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
10 C₁-C₃-alkoxy- ;
- R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
-(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,
-(CH₂)_m-(3- to 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-,
15 heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-,
aryl-, C₁-C₆-alkyl-aryl-, heteroaryl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
-C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂ or -N(H)C(=O)R⁶ group ;
20 s a i d C₁-C₆-alkyl-, -(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl,
-(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,
C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
group
25 being optionally substituted, identically or differently, with 1, 2, 3, or 4
R⁷ groups ;

R^{4a}, R^{4b}, R^{4c}, R^{4d}

- represent, independently from each other, a hydrogen or halogen atom,
30 or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,

- $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$, H $O-C_1-C_6\text{-alkyl-}$, $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
 $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$, $-C(=O)N(R^{6a})R^{6b}$,
 $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$, $-N(R^{6c})C(=O)R^6$,
 $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$, $-N(H)C(=O)OR^6$,
5 $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$, $-N(H)S(=O)_2R^6$,
 $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$, $-O(C=O)N(R^{6a})R^{6b}$,
 $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$, $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$,
 $-S(=O)_2N(H)R^{6a}$, $-S(=O)_2N(R^{6a})R^{6b}$ or $-S(=O)(=NR^{6c})R^6$ group ;
- 10 R^5 represents a hydrogen atom, or a $C_1-C_6\text{-alkyl-}$, $C_2-C_6\text{-alkenyl-}$,
 $-(CH_2)_n-C_2-C_6\text{-alkenyl-}$, $-(CH_2)_n-C_2-C_6\text{-alkynyl-}$, $-(CH_2)_m-C_3-C_6\text{-cycloalkyl-}$,
 $-(CH_2)_m\text{-(3- to 7-membered heterocycloalkyl)}$, $aryl-C_1-C_6\text{-alkyl-}$,
 $heteroaryl-C_1-C_6\text{-alkyl-}$, $halo-C_1-C_6\text{-alkyl-}$, $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$,
 $HO-C_1-C_6\text{-alkyl-}$, $-C_1-C_6\text{-alkyl-CN}$, $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
 15 $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $C_3-C_6\text{-cycloalkyl-}$, 3- to 7-membered
 $heterocycloalkyl-$, $C_2-C_6\text{-alkenyl-}$, $C_3-C_6\text{-cycloalkenyl-}$, $C_2-C_6\text{-alkynyl-}$,
 $aryl-$, $heteroaryl-$ group ;
- s a i d $C_1-C_6\text{-alkyl-}$, $C_2-C_6\text{-alkenyl-}$, $-(CH_2)_n-C_2-C_6\text{-alkenyl-}$,
 $-(CH_2)_n-C_2-C_6\text{-alkynyl-}$, $-(CH_2)_m-C_3-C_6\text{-cycloalkyl-}$, $-(CH_2)_m\text{-(3- to}$
 20 $7\text{-membered heterocycloalkyl)}$, $aryl-C_1-C_6\text{-alkyl-}$, $heteroaryl-C_1-C_6\text{-alkyl-}$,
 $halo-C_1-C_6\text{-alkyl-}$, $R^{6a}(R^{6b})N-C_1-C_6\text{-alkyl-}$, $HO-C_1-C_6\text{-alkyl-}$, $-C_1-C_6\text{-alkyl-CN}$,
 $C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$, $halo-C_1-C_6\text{-alkoxy-}C_1-C_6\text{-alkyl-}$,
 $C_3-C_6\text{-cycloalkyl-}$, 3- to 7-membered $heterocycloalkyl-$, $C_2-C_6\text{-alkenyl-}$,
 $C_3-C_6\text{-cycloalkenyl-}$, $C_2-C_6\text{-alkynyl-}$, $aryl-$ or $heteroaryl-$ group
- 25 being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R^8 groups ;

R^6 , R^{6a} , R^{6b} , R^{6c} ,

- represent, independently from each other, a hydrogen atom, or a
 30 $C_1-C_6\text{-alkyl-}$, $HO-C_1-C_6\text{-alkyl-}$, $C_3-C_6\text{-cycloalkyl-}$, $C_2-C_6\text{-alkenyl-}$, 3- to

7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or heteroaryl-C₁-C₆-alkyl- group ;

5 R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, 10 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;

15

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, 20 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, 25 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or -S(=O)(=NR^{6c})R⁶ group ;



m is an integer of 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 ;

and


30 n is an integer of 1, 2, 3, 4, 5, 6 or 7 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

5 2. The compound according to claim 1, wherein :

A represents a  or  group;

Z represents a -C(=O)N(H)R² or -S(=O)₂N(H)R² group,
or a group selected from :



wherein * indicates the point of attachment of said groups with the rest of the molecule ;

15

R¹ represents a hydrogen atom;

R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
20 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
group is optionally substituted, one or more times, identically or
differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
C₁-C₃-alkoxy- ;

- R^3 represents a hydrogen atom, or a -CN, C_1 - C_6 -alkyl-,
 $-(CH_2)_m$ - C_2 - C_6 -alkenyl, $-(CH_2)_m$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 $-(CH_2)_m$ -(3- to 7-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-,
5 heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-,
halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C_2 - C_6 -alkenyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-,
aryl-, C_1 - C_6 -alkyl-aryl-, heteroaryl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$ or $-N(H)C(=O)R^6$ group ;
10 s a i d C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_2 - C_6 -alkenyl, $-(CH_2)_m$ - C_2 - C_6 -alkynyl,
 $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-,
 C_3 - C_6 -cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C_2 - C_6 -alkenyl-,
 C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl-, C_1 - C_6 -alkyl-aryl- or heteroaryl-
group
15 being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R^7 groups ;

R^{4a} , R^{4b} , R^{4c} , R^{4d}

- represent, independently from each other, a hydrogen or halogen atom,
20 o r a -CN, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy-, halo- C_1 - C_6 -alkyl-,
 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO - C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$, $-C(=O)N(R^{6a})R^{6b}$,
 $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$, $-N(R^{6c})C(=O)R^6$,
 $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$, $-N(H)C(=O)OR^6$,
25 $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$, $-N(H)S(=O)_2R^6$,
 $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$, $-O(C=O)N(R^{6a})R^{6b}$,
 $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$, $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$,
 $-S(=O)_2N(H)R^{6a}$, $-S(=O)_2N(R^{6a})R^{6b}$ or $-S(=O)(=NR^{6c})R^6$ group ;

R^5 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 $-(CH_2)_m$ -(3- to 7-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-,
heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-,
5 HO- C_1 - C_6 -alkyl-, $-C_1$ - C_6 -alkyl-CN, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C_2 - C_6 -alkenyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-,
aryl- or heteroaryl- group ;
s a i d C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, $-(CH_2)_n$ - C_2 - C_6 -alkenyl,
10 $-(CH_2)_n$ - C_2 - C_6 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to
7-membered heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-,
halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, $-C_1$ - C_6 -alkyl-CN,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
15 C_3 - C_6 -cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C_2 - C_6 -alkenyl-,
 C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or heteroaryl- group
being optionally substituted, identically or differently, with 1, 2, 3, or 4
 R^8 groups ;

R^6 , R^{6a} , R^{6b} , R^{6c} ,
20 represent, independently from each other, a hydrogen atom, or a
 C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, C_2 - C_6 -alkenyl-, 3- to
7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl- C_1 - C_6 -alkyl- or
heteroaryl- C_1 - C_6 -alkyl- group ;
25 R^7 represents a hydrogen or halogen atom, or a HO-, -CN, C_1 - C_6 -alkoxy-,
halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
30 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,

$-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-O(C=O)R^6$,
 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$ or
 $-S(=O)(=NR^{6c})R^6$ group ;

R^8 represents a hydrogen or halogen atom, or a $-CN$, C_1 - C_6 -alkoxy-,
 halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO - C_1 - C_6 -alkyl-,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$,
 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$,
 $-S(=O)(=NR^{6c})R^6$ group ;


m is an integer of 0, 1, 2, 3, or 4 ;

and


n is an integer of 1, 2, 3, 4, 5, 6 or 7 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

3. The compound according to claim 1 or 2, wherein :



Z represents a $-C(=O)N(H)R^2$ or $-S(=O)_2N(H)R^2$ group,
or a group selected from :



5

R¹ represents a hydrogen atom ;

R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,
10 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
group is optionally substituted, one or more times, identically or
differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
C₁-C₃-alkoxy- ;

15

R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
-(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,
-(CH₂)_m-(3- to 7-membered heterocycloalkyl), aryl-C₁-C₆-alkyl-,
heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-,
20 halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3- to 7-membered
heterocycloalkyl-, C₂-C₆-alkenyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-,
aryl-, C₁-C₆-alkyl-aryl-, heteroaryl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
-C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂ or -N(H)C(=O)R⁶ group ;
s a i d C₁-C₆-alkyl-, -(CH₂)_m-C₂-C₆-alkenyl, -(CH₂)_m-C₂-C₆-alkynyl,
25 -(CH₂)_m-C₃-C₆-cycloalkyl, aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
C₃-C₆-cycloalkyl-, 3- to 7-membered heterocycloalkyl-, C₂-C₆-alkenyl-,

C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-group

being optionally substituted, identically or differently, with 1, 2, 3, or 4 R⁷ groups ;

5

R^{4a}, R^{4b}, R^{4c}, R^{4d}

represent, independently from each other, a hydrogen or halogen atom, or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl- or -OR⁶ group ;

10

R⁵ represents a hydrogen atom, or a C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₃-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group ;

15

said C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₃-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁸ groups ;

25

R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or heteroaryl-C₁-C₆-alkyl- group ;

5

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 10 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 15 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b},
 -S(=O)(=NR^{6c})R⁶ group ;

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-,
 20 C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a},
 -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,
 -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b},
 -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶,
 25 -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -OR⁶, -O(C=O)R⁶,
 -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶,
 -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
 -S(=O)(=NR^{6c})R⁶ group ;



30 m is an integer of 0, 1, 2, 3 or 4 ;

and


n is an integer of 1, 2, 3 or 4 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

4. The compound according to any one of claims 1, 2, or 3, wherein :

A represents a  or  group;

Z represents a -C(=O)N(H)R² or -S(=O)₂N(H)R² group, or a group selected from :



R¹ represents a hydrogen atom ;

R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-group is optionally substituted, one or more times, identically or differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-, C₁-C₃-alkoxy- ;

R^3 represents a hydrogen atom, or a -CN, C_1 - C_6 -alkyl-,
 $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl),
 aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkyl-,
 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_3 - C_6 -cycloalkyl-, 3-
 5 to 6-membered heterocycloalkyl-, aryl-, C_1 - C_6 -alkyl-aryl- or heteroaryl-
 group ;
 said C_1 - C_6 -alkyl-, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to 6-membered
 heterocycloalkyl), aryl- C_1 - C_6 -alkyl-, heteroaryl- C_1 - C_6 -alkyl-,
 halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
 10 C_3 - C_6 -cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-,
 C_1 - C_6 -alkyl-aryl- or heteroaryl- group being optionally substituted,
 identically or differently, with 1 or 2 R^7 groups ;

R^{4a} , R^{4b} , R^{4c} , R^{4d} :

15 two of which represent a hydrogen atom, while the remaining two
 represent, independently from each other, a hydrogen or halogen atom,
 or a -CN, C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy-, halo- C_1 - C_6 -alkyl-,
 $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-, C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-,
 halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, -OR⁶ group ;

20

R^5 represents a hydrogen atom, or a C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 $-(CH_2)_n$ - C_2 - C_6 -alkenyl, $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl,
 $-(CH_2)_m$ -(3- to 6-membered heterocycloalkyl), aryl- C_1 - C_3 -alkyl-,
 heteroaryl- C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_3 -alkyl-,
 25 HO- C_1 - C_3 -alkyl-, - C_1 - C_3 -alkyl-CN, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-,
 halo- C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl-, C_3 - C_6 -cycloalkyl-, 3- to 6-membered
 heterocycloalkyl-, C_3 - C_6 -cycloalkenyl-, C_2 - C_6 -alkynyl-, aryl- or
 heteroaryl- group ;

said C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-, $-(CH_2)_n$ - C_2 - C_6 -alkenyl,
 30 $-(CH_2)_n$ - C_2 - C_3 -alkynyl, $-(CH_2)_m$ - C_3 - C_6 -cycloalkyl, $-(CH_2)_m$ -(3- to

6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁸ groups ;

R⁶, R^{6a}, R^{6b}, R^{6c},

represent, independently from each other, a hydrogen atom, or a C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or heteroaryl-C₁-C₆-alkyl- group ;

R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b}, -S(=O)(=NR^{6c})R⁶ group ;

25

R⁸ represents a hydrogen or halogen atom, or a -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶,

30

- 5
- $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$,
 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$,
 $-S(=O)(=NR^{6c})R^6$ group ;


m is an integer of 0, 1, 2, 3 or 4 ;

and


- 10 n is an integer of 1, 2, 3 or 4 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

- 15 5. The compound according to any one of claims 1, 2, 3, or 4 wherein :

A represents a  group;

- Z represents a $-C(=O)N(H)R^2$ group,
 20 or a group selected from :



R¹ represents a hydrogen atom ;

- 25 R² represents a hydrogen atom, or a C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
 $R^{6a}(R^{6b})N$ -C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, -C₁-C₆-alkyl-CN,

C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 C₂-C₆-alkynyl- or C₃-C₆-cycloalkyl- group ; wherein said C₃-C₆-cycloalkyl-
 group is optionally substituted, one or more times, identically or
 differently, with a substituent selected from: halo-, C₁-C₃-alkyl-, cyano-,
 5 C₁-C₃-alkoxy- ;

R³ represents a hydrogen atom, or a -CN, C₁-C₆-alkyl-,
 -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl),
 aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-, halo-C₁-C₆-alkyl-,
 10 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, 3-
 to 6-membered heterocycloalkyl-, aryl-, C₁-C₆-alkyl-aryl- or heteroaryl-
 group ;
 said C₁-C₆-alkyl-, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered
 heterocycloalkyl), aryl-C₁-C₆-alkyl-, heteroaryl-C₁-C₆-alkyl-,
 15 halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, aryl-,
 C₁-C₆-alkyl-aryl- or heteroaryl- group being optionally substituted,
 identically or differently, with 1 or 2 R⁷ groups ;

20 R^{4a}, R^{4b}, R^{4c}, R^{4d} :

two of which represent a hydrogen atom, while the remaining two
 represent, independently from each other, a hydrogen or halogen atom,
 or a -CN, C₁-C₆-alkyl-, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-,
 R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-,
 25 halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, -OR⁶ group ;

R⁵ represents a hydrogen atom, or a C₁-C₆-alkyl-, C₂-C₆-alkenyl-,
 -(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₃-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl,
 -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-,
 30 heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-,

- HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-C N , C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-, C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group ;
- 5 s a i d C₁-C₆-alkyl-, C₂-C₆-alkenyl-, -(CH₂)_n-C₂-C₆-alkenyl, -(CH₂)_n-C₂-C₃-alkynyl, -(CH₂)_m-C₃-C₆-cycloalkyl, -(CH₂)_m-(3- to 6-membered heterocycloalkyl), aryl-C₁-C₃-alkyl-, heteroaryl-C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, R^{6a}(R^{6b})N-C₁-C₃-alkyl-, HO-C₁-C₃-alkyl-, -C₁-C₃-alkyl-CN, C₁-C₃-alkoxy-C₁-C₃-alkyl-, halo-C₁-C₃-alkoxy-C₁-C₃-alkyl-,
- 10 C₃-C₆-cycloalkyl-, 3- to 6-membered heterocycloalkyl-, C₃-C₆-cycloalkenyl-, C₂-C₆-alkynyl-, aryl- or heteroaryl- group being optionally substituted, identically or differently, with 1 or 2 R⁸ groups ;

R⁶, R^{6a}, R^{6b}, R^{6c},

- 15 represent, independently from each other, a hydrogen atom, or a C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₃-C₆-cycloalkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, aryl-, heteroaryl-, aryl-C₁-C₆-alkyl-, or heteroaryl-C₁-C₆-alkyl- group ;
- 20 R⁷ represents a hydrogen or halogen atom, or a HO-, -CN, C₁-C₆-alkoxy-, halo-C₁-C₆-alkyl-, R^{6a}(R^{6b})N-C₁-C₆-alkyl-, HO-C₁-C₆-alkyl-, C₁-C₆-alkoxy-C₁-C₆-alkyl-, halo-C₁-C₆-alkoxy-C₁-C₆-alkyl-, C₂-C₆-alkenyl-, 3- to 7-membered heterocycloalkyl-, -C(=O)R⁶, -C(=O)N(H)R^{6a}, -C(=O)N(R^{6a})R^{6b}, -C(=O)O-R⁶, -N(R^{6a})R^{6b}, -NO₂, -N(H)C(=O)R⁶, -N(R^{6c})C(=O)R⁶, -N(H)C(=O)N(R^{6a})R^{6b}, -N(R^{6c})C(=O)N(R^{6a})R^{6b}, -N(H)C(=O)OR⁶, -N(R^{6c})C(=O)OR⁶, -N(H)S(=O)R⁶, -N(R^{6c})S(=O)R⁶, -N(H)S(=O)₂R⁶, -N(R^{6c})S(=O)₂R⁶, -N=S(=O)(R^{6a})R^{6b}, -O(C=O)R⁶, -O(C=O)N(R^{6a})R^{6b}, -O(C=O)OR⁶, -SR⁶, -S(=O)R⁶, -S(=O)N(H)R⁶, -S(=O)N(R^{6a})R^{6b}, -S(=O)₂R⁶, -S(=O)₂N(H)R⁶, -S(=O)₂N(R^{6a})R^{6b} or
- 25 -S(=O)(=NR^{6c})R⁶ group ;
- 30

R^8 represents a hydrogen or halogen atom, or a -CN, C_1 - C_6 -alkoxy-,
 halo- C_1 - C_6 -alkyl-, $R^{6a}(R^{6b})N$ - C_1 - C_6 -alkyl-, HO- C_1 - C_6 -alkyl-,
 C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, halo- C_1 - C_6 -alkoxy- C_1 - C_6 -alkyl-, C_2 - C_6 -alkenyl-,
 3- to 7-membered heterocycloalkyl-, $-C(=O)R^6$, $-C(=O)N(H)R^{6a}$,
 5 $-C(=O)N(R^{6a})R^{6b}$, $-C(=O)O-R^6$, $-N(R^{6a})R^{6b}$, $-NO_2$, $-N(H)C(=O)R^6$,
 $-N(R^{6c})C(=O)R^6$, $-N(H)C(=O)N(R^{6a})R^{6b}$, $-N(R^{6c})C(=O)N(R^{6a})R^{6b}$,
 $-N(H)C(=O)OR^6$, $-N(R^{6c})C(=O)OR^6$, $-N(H)S(=O)R^6$, $-N(R^{6c})S(=O)R^6$,
 $-N(H)S(=O)_2R^6$, $-N(R^{6c})S(=O)_2R^6$, $-N=S(=O)(R^{6a})R^{6b}$, $-OR^6$, $-O(C=O)R^6$,
 10 $-O(C=O)N(R^{6a})R^{6b}$, $-O(C=O)OR^6$, $-SR^6$, $-S(=O)R^6$, $-S(=O)N(H)R^6$,
 $-S(=O)N(R^{6a})R^{6b}$, $-S(=O)_2R^6$, $-S(=O)_2N(H)R^6$, $-S(=O)_2N(R^{6a})R^{6b}$ or
 $-S(=O)(=NR^{6c})R^6$ group ;

m is an integer of 0, 1, 2, 3 or 4 ;
 15 and
 n is an integer of 1, 2, 3 or 4 ;

or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

20

6. The compound according to claim 1, which is selected from the group consisting of :

N-cyclopropyl-4-{8-[(3-hydroxypropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-[8-(isobutylamino)-6-(phenylethynyl)imidazo[1,2-a]pyrazin-3-yl]benzenesulfonamide,

N-cyclopropyl-4-{6-[(2-fluorophenyl)ethynyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl}benzenesulfonamide,

N-cyclopropyl-4-[6-(3-hydroxyprop-1-yn-1-yl)-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]benzenesulfonamide,

N-cyclopropyl-4-[6-(3-hydroxyprop-1-yn-1-yl)-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl]benzamide,

N-cyclopropyl-4-[8-(isobutylamino)-6-(phenylethynyl)imidazo[1,2-a]pyrazin-3-yl]benzamide,

N-cyclopropyl-4-{6-[(2-fluorophenyl)ethynyl]-8-(isobutylamino)imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(4-hydroxybutyl)amino]-6-(3-hydroxy-3-methylbut-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(4-hydroxybutyl)amino]-6-[(1-hydroxycyclopentyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(2-hydroxy-2-methylpropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-{8-[(3-aminopropyl)amino]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropyl-3-methoxybenzamide,

4-{6-(cyclohexylethynyl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{6-(3-hydroxy-3-methylbut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(4-hydroxybut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-[(1-hydroxycyclopentyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-[3-(dimethylamino)prop-1-yn-1-yl]-8-[(2-

methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(3-methoxyprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(3,3-dimethylbut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(3-hydroxybut-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(4-hydroxypent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-{6-(5-cyanopent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{6-(3-hydroxypent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-(6-{3-[benzyl(methyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)-N-cyclopropylbenzamide,

4-{6-(3-aminoprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{6-(3-hydroxyhex-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(5-phenylpent-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

6-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}hex-5-ynoic acid,

N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(pyridin-2-ylethynyl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(4-phenylbut-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-{6-(5-cyclohexylpent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{6-(4-methylpent-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-[(3-hydroxyphenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-{6-[3-(carbamoylamino)prop-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-(3-phenylprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-[(4-fluorophenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-[(3S)-3-hydroxyoct-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

methyl 6-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}hex-5-ynoate,

9-{3-[4-(cyclopropylcarbamoyl)phenyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-6-yl}non-8-ynoic acid,

N-cyclopropyl-4-(6-{3-[(methylcarbamoyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide,

N-cyclopropyl-4-(6-{4-[methyl(methylcarbamoyl)amino]but-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide,

N-cyclopropyl-4-{6-[6-(methylamino)-6-oxohex-1-yn-1-yl]-8-[(2-

methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-(6-{3-[methyl(methylcarbamoyl)amino]prop-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide,

4-{6-[(4-carbamoylphenyl)ethynyl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-(6-{4-[(methylcarbamoyl)amino]but-1-yn-1-yl}-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl)benzamide,

N-cyclopropyl-4-{6-[5-(methylamino)-5-oxopent-1-yn-1-yl]-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{8-[(2-methylpropyl)amino]-6-[(4-sulfamoylphenyl)ethynyl]imidazo[1,2-a]pyrazin-3-yl}benzamide,

4-{6-(cyclopentylethynyl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-cyclopropylbenzamide,

N-cyclopropyl-4-{6-ethynyl-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}benzamide,

N-cyclopropyl-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(2-methylpropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

(2S)-3-({3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-6-(3-hydroxyprop-1-yn-1-yl)imidazo[1,2-a]pyrazin-8-yl}amino)propane-1,2-diol,

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(4,4,4-trifluorobutyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(2-methylprop-2-en-1-yl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{8-(Allylamino)-3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-

(isobutylamino)imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{3-[3-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-

(isobutylamino)imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

4-({3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-6-(3-hydroxyprop-1-yn-1-

yl)imidazo[1,2-a]pyrazin-8-yl}amino)butan-1-ol,

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(3,3,3-

trifluoropropyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{8-[(Cyclobutylmethyl)amino]-3-[4-(2-cyclopropyl-1H-imidazol-5-

yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{3-[4-(2-Cyclopropyl-1H-imidazol-5-yl)phenyl]-8-[(3-

hydroxypropyl)amino]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-{8-[(3-Aminopropyl)amino]-3-[4-(2-cyclopropyl-1H-imidazol-5-

yl)phenyl]imidazo[1,2-a]pyrazin-6-yl}prop-2-yn-1-ol,

3-[4-(2-cyclopropyl-1H-imidazol-5-yl)phenyl]-6-[3-(1,1-dioxidothiomorpholin-

4-yl)prop-1-yn-1-yl]-N-(3,3,3-trifluoropropyl)imidazo[1,2-a]pyrazin-8-amine,

N-cyclopropyl-4-{6-[3-(1,1-dioxidothiomorpholin-4-yl)prop-1-yn-1-yl]-8-

[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-

methylbenzamide),

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N,2-

dimethylbenzamide,

N-ethyl-4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-

yl}-2-methylbenzamide,

N-(2,2-difluoroethyl)-4-{6-ethynyl-8-[(3,3,3-

trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methyl-N-(1-methylcyclopropyl)benzamide,

N-(1-cyanocyclopropyl)-4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-(1-methoxycyclopropyl)-2-methylbenzamide,

4-{6-ethynyl-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-[rel(1S,2S)-2-fluorocyclopropyl]-2-methylbenzamide,

4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N,2-dimethylbenzamide,

N-ethyl-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

N-(2,2-difluoroethyl)-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methyl-N-(1-methylcyclopropyl)benzamide,


N-(1-cyanocyclopropyl)-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,

4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-N-(1-methoxycyclopropyl)-2-methylbenzamide,

N-[rel(1S,2S)-2-fluorocyclopropyl]-4-{6-(3-hydroxyprop-1-yn-1-yl)-8-[(3,3,3-trifluoropropyl)amino]imidazo[1,2-a]pyrazin-3-yl}-2-methylbenzamide,


or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, or a mixture of same.

- 5 7. A method of preparing a compound of general formula (I) :



in which A, R¹, R³ and R⁵ are as defined for the compounds of general formula (I) in any one of claims 1 to 6,


- 10 in which method an intermediate compound of general formula (II) :



in which A, R¹ and R⁵ are as defined for the compounds of general formula (I) in

- 15 any one of claims 1 to 6,

is allowed to react with an alkyne of general formula (IIa) :



(IIa)

in which R³ is as defined for the compounds of general formula (I) in any one of claims 1 to 5, and -M represents a hydrogen or a functional group suitable to be coupled to the Br-bearing carbon atom of the compounds of general formula (II),

5 thus providing a compound of general formula (I) as defined in any one of claims 1 to 6.

8. A compound of general formula (I), or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a pharmaceutically
10 acceptable salt thereof, or a mixture of same, according to any one of claims 1 to 6, for use in the treatment or prophylaxis of a disease.

9. A pharmaceutical composition comprising a compound of general formula (I), or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt
15 thereof, particularly a pharmaceutically acceptable salt thereof, or a mixture of same, according to any one of claims 1 to 6, and a pharmaceutically acceptable diluent or carrier.

10. A pharmaceutical combination comprising :

20 - one or more compounds of general formula (I), or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a pharmaceutically acceptable salt thereof, or a mixture of same, according to any one of claims 1 to 6 ;

and

25 - one or more agents selected from : a taxane, such as Docetaxel, Paclitaxel, or Taxol; an epothilone, such as Ixabepilone, Patupilone, or Sagopilone; Mitoxantrone; Predinisolone; Dexamethasone; Estramustin; Vinblastin; Vincristin; Doxorubicin; Adriamycin; Idarubicin; Daunorubicin; Bleomycin; Etoposide; Cyclophosphamide; Ifosfamide; Procarbazine; Melphalan;
30 5-Fluorouracil; Capecitabine; Fludarabine; Cytarabine; Ara-C;


2-Chloro-2'-deoxyadenosine; Thioguanine; an anti-androgen, such as Flutamide, Cyproterone acetate, or Bicalutamide; Bortezomib; a platinum derivative, such as Cisplatin, or Carboplatin; Chlorambucil; Methotrexate; and Rituximab.

5


11. Use of a compound of general formula (I), or a stereoisomer, a tautomer, an N-oxide, a hydrate, a solvate, or a salt thereof, particularly a pharmaceutically acceptable salt thereof, or a mixture of same, according to any one of claims 1 to 6, for the preparation of a medicament for the prophylaxis or treatment of a disease.

12. Use according to claim 11, wherein said disease is a disease of uncontrolled cell growth, proliferation and/or survival, an inappropriate cellular immune response, or an inappropriate cellular inflammatory response, particularly in which the uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune response, or inappropriate cellular inflammatory response is mediated by Mps-1, more particularly in which the disease of uncontrolled cell growth, proliferation and/or survival, inappropriate cellular immune response, or inappropriate cellular inflammatory response is a haematological tumour, a solid tumour and/or metastases thereof, *e.g.* leukaemias and myelodysplastic syndrome, malignant lymphomas, head and neck tumours including brain tumours and brain metastases, tumours of the thorax including non-small cell and small cell lung tumours, gastrointestinal tumours, endocrine tumours, mammary and other gynaecological tumours, urological tumours including renal, bladder and prostate tumours, skin tumours, and sarcomas, and/or metastases thereof.

13. Use of a compound of general formula (II) :



in which A, R¹ and R⁵ are as defined for the compounds of general formula (I) in
any one of claims 1 to 6, for the preparation of a compound of general formula
5 (I) according to any one of claims 1 to 6.



(I)