




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/00, 15/82, A01H 5/00, 5/10	A1	(11) International Publication Number: WO 99/67367 (43) International Publication Date: 29 December 1999 (29.12.99)
(21) International Application Number: PCT/US99/14022 (22) International Filing Date: 22 June 1999 (22.06.99) (30) Priority Data: 60/090,240 22 June 1998 (22.06.98) US 60/091,640 2 July 1998 (02.07.98) US (71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; Morrill Hall, 100 Church Street S.E., Minneapolis, MN 55455 (US). UNITED STATES OF AMERICA represented by THE SECRETARY, DEPARTMENT OF AGRICULTURE [US/US]; Building 005, Room 416, BARC-W, Beltsville, MD 20705-2350 (US). (71)(72) Applicants and Inventors: EGLI, Margaret, A. [US/US]; 705 Heinel Circle, Roseville, MN 55113 (US). GROH, Susanne [US/US]; 2384 Larpenner Avenue, #7, Falcon Heights, MN 55113 (US). KIANIAN, Shahryar, F. [US/US]; Apartment 230, 3201 20th Street South, Fargo, ND 58104 (US). PHILLIPS, Ronald, L. [US/US]; 819 Tanglewood Drive, Shoreview, MN 55126 (US). RINES, Howard, W. [US/US]; 4292 Nancy Place, Shoreview, MN 55126 (US). SOMERS, David, A. [US/US]; 924 West County Road B2, Roseville, MN 55113 (US).		(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lunberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: DNA ENCODING OAT ACETYL CoA CARBOXYLASE

(57) Abstract

The present invention provides the complete cDNA sequence of maize acetyl CoA carboxylase and methods for conferring herbicide tolerance and/or altering the oil content of plants by introducing and expressing a plant acetyl CoA carboxylase gene in plant cells. The method of imparting herbicide tolerance to a plant includes the steps of introducing an expression cassette encoding a plant acetyl CoA carboxylase or an antisense DNA sequence complementary to the sequence for a plant acetyl CoA carboxylase gene operably linked to a promoter functional in plant cells, into the cells of a plant tissue and expressing the plant acetyl CoA carboxylase gene in an amount effective to render the acetyl CoA carboxylase and/or plant cell tolerant to the herbicides. The method of altering the oil content in a plant includes the steps of introducing an expression cassette into plant cells and expressing the acetyl CoA carboxylase gene in an amount effective to alter the oil content of the cells. The expression cassette can also be introduced into other host cells to increase yield of a plant acetyl CoA carboxylase so that crystallized enzyme can be used to screen and identify other herbicides that bind to and inhibit the enzyme. Also provided is a partial DNA sequence encoding oat acetyl CoA carboxylase.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DNA ENCODING OAT ACETYL CoA CARBOXYLASE

Background of the Invention

Acetyl CoA carboxylase (ACCase) is an enzyme involved in many important metabolic pathways in plant, animal and bacterial cells. The enzyme is especially important in fatty acid synthesis in plants and is sensitive to inhibition by some types of herbicides. Structurally, ACCases are biotinylated and are quite large enzymes consisting of one or more subunits. For example, most ACCases of animals, higher plants, and yeast are dimers of 420 to 700 kD native MW and contain subunits of 200 to 280 kD. Diatom and algal ACCases are 700 to 740 kD tetramers of 160 to 180 kD subunits. Bacterial ACCase consists of three dissociable proteins; biotin carboxylase (51 kD), biotin carboxyl carrier protein (22.5 kD), and biotin transcarboxylase (130 kD).

Acetyl CoA Carboxylase (ACCase) catalyzes the ATP-dependent formation of malonyl-CoA from acetyl-CoA and bicarbonate in animal, plant, and bacterial cells. Malonyl-CoA is an essential substrate for (i) *de novo* fatty acid (FA) synthesis, (ii) fatty acid elongation, (iii) synthesis of secondary metabolites such as flavonoids and anthocyanins, and (iv) malonylation of some amino acids and secondary metabolites. Synthesis of malonyl-CoA is the first committed step of flavonoid and fatty acid synthesis and current evidence suggests that ACCase catalyzes the rate-limiting step of fatty acid synthesis. Formation of malonyl-CoA by ACCase occurs via two partial reactions and requires a biotin prosthetic group:



In *E. coli*, these reactions are catalyzed by three distinct components; biotin carboxylase, biotin transcarboxylase, and biotin carboxyl carrier protein, which can be separated and yet retain partial activity. Plant and animal ACCases contain all three activities on a single polypeptide.

In plants, most ACCase activity is located in plastids of green and non-green plant tissues including leaves and oil seeds. Leaf ACCase activity is primarily located in mesophyll cells, but lesser amounts have been found in C-4

bundle sheath cells and in epidermal cells. The subcellular location of ACCase activity in epidermal cells is unknown, but since synthesis of very long-chain fatty acids (VLCFA) for formation of waxes, cutin, and suberin occurs on the endoplasmic reticulum (ER), malonyl-CoA might also be derived from a
5 cytosolic ACCase. In contrast, rat ACCase is primarily cytosolic or associated with the outer mitochondrial membrane.

De novo fatty acid synthesis in chloroplasts involves successive 2-carbon additions to acetate, using malonate as the 2-C donor. All intermediates are attached to acyl carrier protein (ACP). Synthesis in plastids
10 resembles that in E. coli in that the fatty acid synthesis complex can be dissociated into separate enzymes: β -ketoacyl-ACP synthase (KAS), β -ketoacyl-ACP reductase, β -hydroxyl-ACP dehydratase, and enoyl-ACP reductase, acetyl-CoA:ACP transacylase, and malonyl-CoA:ACP transacylase. A highly active KASIII isozyme catalyzes the condensation of acetyl-CoA and
15 malonyl-ACP. Successive additions of malonyl-CoA to acyl-ACPs catalyzed by KAS I form C16 acyl-ACP, some of which is converted to C18 acyl-ACP by KAS II and then to C18:1-ACP. Fatty acid metabolism then diverges; de-esterification allows movement to the cytoplasm (eukaryotic path) where fatty acids may be further unsaturated and/or elongated by additions of malonyl-CoA
20 in the ER. Alternatively, fatty acids are linked to glycerol-3-phosphate (prokaryotic path), further unsaturated, and used for synthesis of chloroplast lipids. A portion of cytoplasmic lipids returns to the chloroplast. The relative contributions of these two paths are species-specific but appear to be relatively flexible in mutants blocked in either path. In oil-storing organs such as
25 cotyledons and monocot embryos, the triacylglycerides are stored in cytoplasmic oil bodies surrounded by a single unit membrane.

Condensation of malonyl-CoA with phenylpropionyl-CoAs or acetyl-CoA leads to synthesis of flavonoids, anthocyanins, or to polyacetates. Condensation is increased by light, elicitors, or pathogens and may be the rate-
30 limiting step in synthesis of some phytoalexins. In addition to the secondary metabolites derived by de novo synthesis, malonyl conjugates of flavonoid glycosides, formed by malonyl-CoA:flavonoid glycoside malonyltransferase, D-amino acids and 1-amino-carboxyl-cyclopropane (ethylene precursor) are

found in plants. Malonylated compounds accumulate in vacuoles, probably after synthesis in the cytoplasm.

An important property of ACCase is the central role it plays in fatty acid synthesis and accumulation in plants and seeds. Available evidence supports the idea that ACCase activity is the rate-limiting step for de novo fatty acid synthesis in plants. High rates of ACCase activity in vitro parallel or slightly precede high rates of lipid deposition or [¹⁴C]acetate incorporation into lipids in developing leaves and oil seeds. Significant changes in plant ACCase activity occur during chloroplast development and increase in ACCase activity correlates with lipid deposition in developing oil seeds. Turnham et al., Biochem. J., 212:223 (1983); and Beittenmiller et al., Plant Physiol., 100:923 (1992).

Among other properties, ACCase in most monocots is also inhibited by several herbicides. [¹⁴C]acetate incorporation into maize lipids is strongly inhibited by fluazifop and sethoxydim due to inhibition of plastid ACCase. In barley, however, fluazifop had little effect on [¹⁴C]acetate incorporation into very long-chain fatty acids. Since synthesis of very long-chain fatty acids occurs in the cytosol on the ER, and de novo fatty acid synthesis occurs in the plastids, cytosolic malonyl-CoA might be supplied by a herbicide insensitive ACCase isozyme.

There are three general mechanisms by which plants may be resistant to, or tolerant of, herbicides. These mechanisms include insensitivity at the site of action of the herbicide (usually an enzyme), rapid metabolism (conjugation or degradation) of the herbicide, or poor uptake and translocation of the herbicide. Altering the herbicide site of action from a sensitive to an insensitive form is the preferred method of conferring tolerance on a sensitive plant species. This is because tolerance of this nature is likely to be a dominant trait encoded by a single gene, and is likely to encompass whole families of compounds that share a single site of action, not just individual chemicals. Therefore, detailed information concerning the biochemical site and mechanism of herbicide action is of great importance and can be applied in two ways. First, the information can be used to develop cell selection strategies for the efficient identification and isolation of appropriate herbicide-tolerant variants. Second, it

can be used to characterize the variant cell lines and regenerated plants that result from the selections.

Tissue culture methods have been used to select for resistance (or tolerance) using a variety of herbicides and plant species (see review by Meredith and Carlson, 1982, in Herbicide Resistance in Plants, eds. Lebaron and Gressel, pp. 275-291, John Wiley and Sons, NY). For example, P. C. Anderson et al., in U.S. Patent No. 4,761,373, disclose the use of tissue culture methods to produce maize plants resistant to herbicidal imidazolidones and sulfonamides. The resistance is due to the presence of altered acetohydroxy acid synthase which is resistant to deactivation by these herbicides.

Certain 1,3-cyclohexanediones exhibit general and selective herbicidal activity against plants. One such cyclohexanedione is sethoxydim {2-[1-(ethoxyimino)-butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one}. Sethoxydim is commercially available from BASF (Parsippany, New Jersey) under the designation POAST™.

Other herbicidal cyclohexanediones include clethodim, (E,E)-(±)-2-[1-[(3-chloro-2-propenyl)oxy] imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one; available as SELECT™ from Chevron Chemical (Valent) (Fresno, California); cloproxydim, (E,E)-2-[1-[(3-chloro-2-propenyl)oxy]imino] butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one; available as SELECTONE™ from Chevron Chemical (Valent) (Fresno, California); and tralkoxydim, 2-[1-(ethoxyimino)propyl]-3-hydroxy-5-mesitylcyclohex-2-enone, available as GRASP™ from Dow Chemical USA (Midland, Michigan).

For purposes of reference in the present specification, the herbicides described in the two preceding paragraphs and other structurally related herbicidal compounds, are collectively referred to as the cyclohexanedione family of herbicides.

Certain aryloxyphenoxypropanoic acids exhibit general and selective herbicidal activity against plants. In these compounds, the aryloxy group may be phenoxy, pyridinyloxy or quinoxalinyloxy. One such herbicidal aryloxyphenoxypropanoic acid is haloxyfop, {2-[4-[(3-chloro-5-(trifluoromethyl)-2-pyridinyl)oxy]phenoxy]-propanoic acid}, which is available

as VERDICT™ from Dow Chemical USA (Midland, Michigan). Another is diclofop, {(±)-2-[4-(2,4-dichlorophenoxy)-phenoxy]propanoic acid}, available as HOELON™ from Hoechst-Roussel Agri-Vet Company (Somerville, New Jersey).

5 Other members of this family of herbicides include fenoxyprop, (±)-2-[4-[(6-chloro-2-benzoxazolyl)oxy] phenoxy]propanoic acid; available as WHIP™ from Hoechst-Roussel Agri-Vet Company (Somerville, New Jersey); fluazifop, (±)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid; available as FUSILADE™ from ICI Americas (Wilmington, Delaware);
10 fluazifop-P, (R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid; available as FUSILADE 2000™ from ICI Americas (Wilmington, Delaware); and quizalofop, (±)-2-[4[(6-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid; available as ASSURE™ from E. I. DuPont de Nemours (Wilmington, Delaware).

15 For purposes of reference in the present specification, the herbicides referred to in the two preceding paragraphs and other structurally related herbicidal compounds, are collectively referred to as herbicidal aryloxyphenoxypropanoic acids.

 Thus, there is a need for methods to develop plants that are
20 resistant or tolerant to herbicides. There is also a need to increase the oil and/or fatty acid content of the plants and seeds, as well as for methods to increase their resistance to herbicides. Moreover, there is a need to identify and clone genes important in conferring herbicide tolerance and in increasing the oil content of plants.

25

Summary of the Invention

 The present invention provides an isolated and purified DNA molecule comprising a DNA segment encoding a plant acetyl CoA carboxylase such as a monocot acetyl CoA carboxylase, e.g., a maize or oat acetyl CoA
30 carboxylase, and methods for conferring herbicide tolerance and/or altering the oil content of plants by introducing and expressing a plant acetyl CoA carboxylase gene in the plant cells. The DNA molecule encoding a plant acetyl CoA carboxylase can encode an unaltered plant acetyl CoA carboxylase or an

altered plant acetyl CoA carboxylase substantially tolerant to inhibition by cyclohexanedione or aryloxyphenoxypropanoic acid herbicides as well as encoding an antisense DNA sequence that is substantially complementary to a plant acetyl CoA carboxylase gene or to a portion thereof. A DNA molecule of the invention can also further comprise an amino terminal plant chloroplast transit peptide sequence operably linked to the plant acetyl CoA carboxylase gene. The DNA molecule of the invention preferably encodes the plastidic form of plant acetyl CoA carboxylase. The invention also provides for probes, e.g., a probe which is detectably labeled or which binds to a detectable label, or primers, e.g., a single-strand DNA molecule having at least seven nucleotide bases, that hybridize to a DNA encoding a plant acetyl CoA carboxylase, e.g., a DNA molecule comprising SEQ ID NO:23 or its complement. For example, the present invention provides an isolated and purified DNA molecule of at least seven nucleotide bases which hybridizes under high stringency conditions to a DNA molecule comprising a DNA segment encoding a plant acetyl CoA carboxylase and provides a hybridization probe comprising an isolated and purified DNA molecule of at least seven nucleotide bases, which is detectably labeled or which binds to a detectable label, which DNA molecule hybridizes under high stringency conditions to the non-coding strand of a DNA molecule comprising a DNA segment encoding a plant acetyl CoA carboxylase. High stringency conditions are defined as: hybridization at 65°C for at least 16 hours in 5xSSC, 1x Denhardt's solution, 50 mM Tris-HCl, pH 8, 0.2% SDS, 10 mM EDTA, 0.1 mg/ml salmon sperm DNA, followed by washing twice for 5 minutes in 2xSSC, 0.5% SDS at 25°C, once for 10 minutes in 0.2xSSC, 0.1% SDS at 25°C and twice for 30 minutes in 0.2xSSC, 0.1% SDS at 65°C.

The method of imparting cyclohexanedione or aryloxyphenoxypropanoic acid herbicide tolerance to a plant includes the steps of introducing a chimeric or recombinant DNA molecule comprising a gene coding for a plant acetyl CoA carboxylase or an altered or a functional mutant thereof operably linked to a promoter functional in a plant cell into cells of a susceptible plant, and regenerating the transformed plant cells to provide a differentiated plant. The promoter can be an inducible or tissue specific promoter or provide for overexpression of at least about a 2-fold amount of a

native plant acetyl CoA carboxylase. The functional linkage of a promoter to the chimeric DNA molecule results in an expression cassette. Expression of the chimeric DNA molecule is in an amount effective to render the acetyl CoA carboxylase and/or the plant tissue substantially tolerant to the herbicides relative to the native acetyl CoA carboxylase present in said plant. Herbicide tolerance can be achieved in the plants by at least two methods, including increasing the level of gene expression of a native or unaltered acetyl CoA carboxylase, or by introducing an altered gene coding for an acetyl CoA carboxylase that is less sensitive to herbicide inhibition. The level of gene expression can be increased by either combining a plant acetyl CoA carboxylase gene with a promoter that provides for a high level of gene expression, such as a 35S cauliflower mosaic virus promoter (CaMV), or by introducing multiple copies of the gene into the cell so that the multiple copies of the gene are integrated into the genome of transformed plant cells. The preferred plant cells into which to introduce the expression cassette of the invention, to achieve herbicide tolerance, are monocot plant cells. Preferably, the cells are not oat cells. Once transformed cells exhibiting herbicide tolerance are obtained, transgenic plants and seeds can then be regenerated therefrom, and evaluated for stability of the inheritance of the herbicide tolerance trait.

The invention also provides a method for altering, e.g., lowering or raising, the oil content in a plant. As described hereinbelow, a locus having a major effect on oat (*Avena sativa* L.) groat oil content was located on linkage group 11 by quantitative trait locus (QTL) dissection methods. A partial oat acetyl-CoA carboxylase (ACCase) cDNA with >84% identity to plastidic ACCase sequences from wheat or maize was linked to this locus. Similar QTL and ACCase placements were obtained with two recombinant inbred populations ($n \geq 137$) with one common parent. The locus on linkage group 11 linked to the ACCase locus accounted for up to 48% of the phenotypic variance for groat oil content. Other loci were identified which accounted for an additional 14% of the phenotypic variance. The major locus plus additional QTLs represent three potentially homoeologous genomic segments in oat that can be followed by as few as three RFLP markers to produce a large shift in groat oil content. Oat acetyl CoA carboxylase DNA is also useful in molecular marker-assisted

selection to facilitate breeding high- or low-oil oat, genetic dissection of the groat oil trait, and to identify specific genes with large effects whose activity may be manipulated via genetic transformation.

Thus, the invention provides a method which includes the steps of

5 introducing a chimeric or recombinant DNA molecule comprising a gene coding for a plant, e.g., oat, acetyl CoA carboxylase or an altered or a functional mutant thereof operably linked to a promoter functional in a plant cell into the cells of plant tissue and expressing the gene in an amount effective to alter the oil content of the plant cell. An alteration in oil content can include a change in

10 total oil content over that normally present in that type of plant cell, a change in the location of the oil in the plant, plant tissue, seed or plant part, and/or a change in the type of oil present in the cell. For example, low-oil oat includes, but is not limited to, an oat plant, plant part, plant tissue or groat having about 8 to about 40, preferably about 8 to about 30, and more preferably about 20 to

15 about 30, percent dry weight of oil relative to the corresponding parental, e.g., nontransformed or inbred, oat plant, plant part, plant tissue or groat. High-oil oat includes, but is not limited to, an oat plant, plant part, plant tissue or groat having about 2 to about 8, preferably 1 to about 4, and more preferably about 2 to about 4, percent dry weight of oil relative to the corresponding parental oat plant, plant

20 part, plant tissue or groat.

An alteration in oil content in the plant cell can be achieved by at least two methods including:

- (1) an increase or decrease in expression of an unaltered or altered plant acetyl CoA carboxylase gene; or
- 25 (2) by introducing an unaltered, altered or functional mutant plant acetyl CoA carboxylase gene.

The level of gene expression of an unaltered plant acetyl CoA carboxylase gene can be increased by either combining an unaltered plant acetyl CoA carboxylase with a promoter that provides for a high level of gene expression, or by

30 introducing multiple copies of an expression cassette into cells so that multiple copies of the gene are integrated into the genome. When an altered or a functional mutant plant acetyl CoA carboxylase gene codes for an enzyme that exhibits an increase in specific activity, it can lead to an increase in total oil

content of the plant cell. When an altered or a functional mutant acetyl CoA carboxylase gene codes for an enzyme having a decrease in specific activity, it may lead to a decrease in the total oil content of the plant cell. Preferably, the expression cassette is introduced into dicot plants such as soybeans, canola, and sunflower. In an especially preferred version, transformed cells exhibiting about a 1.2- to 5-fold increase in total oil content and/or expression or specific activity of acetyl CoA carboxylase are selected for and used to generate transgenic plants and seeds exhibiting a substantial increase in oil content. A substantial increase in oil content depends on the oil content normally present in the plant or seed and can range from about a 1.2 to a 20-fold increase.

The invention also provides for a method of producing plant acetyl CoA carboxylase in a host cell. The method includes the steps of introducing an expression cassette comprising a chimeric gene encoding a plant acetyl CoA carboxylase or an altered or a functional plant acetyl CoA carboxylase operably linked to a promoter into a host cell and expressing the gene in an amount sufficient to permit crystallization of the plant acetyl CoA carboxylase. An expression cassette can include a promoter that is functional in either a eukaryotic or a prokaryotic cell. Preferably, the expression cassette is introduced into a prokaryotic cell, such as *E. coli*, that is routinely used for production of recombinantly produced proteins. Recombinantly produced and crystallized plant acetyl CoA carboxylase can then be used to identify other herbicides and that bind to and inhibit acetyl CoA carboxylase in plants. In addition, the availability of large amounts of purified enzyme can permit the screening of the efficacy of such herbicides in terms of their ability to bind to, or otherwise inhibit, the activity of the enzyme.

The present invention also provides a method of introducing an exogenous plant acetyl CoA carboxylase gene into a host cell comprising transforming host cells *in vitro* with an expression cassette comprising a chimeric or recombinant DNA molecule encoding a plant acetyl CoA carboxylase gene operably linked to a promoter functional in the host cell, expanding the transformed host cells *in vitro*, and identifying a transformed host cell which expresses the chimeric DNA molecule.

Further provided is a method to identify a plant, e.g., an oat plant, having altered characteristics, e.g., altered groat oil content. The method comprises contacting a probe comprising at least a portion of a nucleic acid sequence encoding oat acetyl CoA carboxylase with a sample from an oat plant
5 in an amount sufficient to form complexes, wherein the sample comprises plant genomic nucleic acid. The amount of complex formation is then detected or determined. Preferably, the plant cells in the sample are monocot plant cells, e.g., wheat, rice, maize, barley, millet (e.g., foxtail or finger), sorghum, sugarcane, rye, oat or timothy cells. Also preferably, the DNA molecule encodes
10 a polypeptide comprising SEQ ID NO:24, e.g., a DNA molecule comprising SEQ ID NO:23. Thus, the invention provides a nucleic acid fragment useful in marker-assisted breeding programs such as those which identify plants having altered groat oil content or composition.

Also provided is an expression cassette comprising an
15 endosperm-specific transcriptional control region, e.g., the Dy10 or zein transcriptional control region, operably linked to a DNA encoding a plant acetyl CoA carboxylase, or subunit thereof. Other endosperm-specific promoters are also useful in the compositions and methods of the invention. These promoters include, but are not limited to, ltr1 promoter, a synthetic promoter with
20 endosperm box of HOD (Vincentcarbajosa et al., Plant J., **13**, 629 (1998)); rice glutelin gene promoter (Sindhu et al., Plant Sci., **130**, 189 (1997); Yoshihara et al., Plant Sci., **121**, 63 (1996)); barley alpha-amylase inhibitor subunit promoter (Gross et al., Plant Mol. Biol., **34**, 331 (1997)); gamma zein promoter (Towent et al., Plant Mol. Biol., **34**, 139 (1997)); maize opaque-2 locus promoter (Rossi et al., J. Biol. Chem., **272**, 13758 (1997); Gallusci et al., Mol. & Gen. Genet., **244**,
25 391 (1994)); the zmGBS, zmZ27, osAGP or osGT1 promoter (Russell et al., Transgenic Res., **6**, 157 (1997)); alpha globulin promoter (Nakase et al., Plant Mol. Biol., **33**, 513 (1997)); hordein promoter (Sorenson et al., Mol. & Gen. Genet., **250**, 750 (1996)); Viciafaba legumin and vicilin storage protein gene
30 promoter (Wobus et al., J. Plant Physiol., **145**, 592 (1995)); rice waxy gene promoter (Hirano et al., Plant & Cell Physiol., **36**, 37 (1995)); oat globulin gene osglo5 promoter (Schubert et al., Plant Mol. Biol., **26**, 203 (1994)); napA

promoter (Stalberg et al., *Plant Mol. Biol.*, 23, 671 (1993)); and rice prolamin gene promoter (Zhou et al., *Science in China*, 36, 1307 (1993)).

The term "consists essentially of" as used with respect to the present DNA molecules is defined to mean that a major portion of the nucleotide sequence encodes an ACCase, and that nucleotide sequences are not present which encode proteins other than ACCase or functional equivalents thereof.

Brief Description of the Figures

Figure 1: A schematic depiction of the fatty acid biosynthesis pathway in plants.

Figure 2: A graph depicting the effect of sethoxydim on the growth of mutant maize callus.

Figure 3: A graph depicting the shoot length growth of maize seedlings seven days after treatment with sethoxydim.

Figure 4: A graph depicting the shoot length growth of maize seedlings fourteen days after treatment with sethoxydim.

Figure 5: Total soluble and biotinylated polypeptides in ACCase purification fractions from seedling leaves of maize inbred A619. Proteins were separated by SDS-PAGE in 7.5% gels and then silver-stained (Panel A). An identical gel was Western-blotted and a longitudinal section of each lane was probed with avidin (Panel B). Lanes were 1: crude (10 μ g); 2: $(\text{NH}_4)_2\text{SO}_4$ (10 μ g); 3: S-300 (5 μ g); 4: Blue Sepharose (2 μ g); 5: Mono-Q ACCase II (5 μ g); and 6: Mono-Q ACCase I (5 μ g). Diagonal lines between lanes indicate position of molecular weight markers shown on the left.

Figure 6: Immunoprecipitation of ACCase activity from B73 leaf, embryo, endosperm, and BMS suspension cultured cells. Equal activities (0.58 nmol min⁻¹) were incubated with 16 μ L serum (immune plus preimmune), immune complexes were precipitated with Protein A-agarose, and ACCase activity remaining in the resulting supernatant fraction was measured relative to the preimmune serum control.

Figure 7: Herbicide inhibition of acetyl-(AcCoA) or propionyl-CoA (Prop-CoA)-dependent H¹⁴CO₃-incorporation into acid-stable

product by ACCase I and II Mono-Q fractions. Activities in the presence of haloxyfop (1 μ M) are expressed relative to the minus herbicide control.

Figure 8: Comparison of the peptide sequence of maize cDNA clones #15-14 and #18-5 with chicken ACCase. The approximate locations of the biotin carboxylase, biotin binding site, and biotin transcarboxylase functional domains are indicated for the chicken sequence. The percentages of amino acid identity are indicated by cross-hatched boxes for the maize coding sequence. Regions of genomic DNA Type I and Type II clone sequences that align with cDNA #18-5 are indicated by solid heavy lines. The approximate locations of subclone #28 and #16 from genomic Type I and subclone #34 from genomic Type II clones are indicated.

Figure 9: Northern blot of total RNA from leaf, immature embryo and endosperm tissue (16 days after pollination), and Black Mexican Sweet corn (BMS) cells. Lanes contain 10 μ g total RNA and were probed with the 2 kb EcoRI fragment of lambda clone #15-14.

Figure 10: DNA sequence (SEQ ID NO. 1) of a 2 kb EcoRI fragment of lambda clone #15-14 including a portion of a maize ACCase gene located at bases 2883 to 83 from the 3' stop codon.

Figure 11: Graph of ACCase activity during seed development in two high oil soybean cell lines and one low oil soybean cell line.

Figure 12: Cloning strategy to obtain the complete coding sequence of the maize ACCase gene.

Figure 13: DNA sequence (SEQ ID NO:5) of a 7470 base pair cDNA of a maize ACCase gene. (Genbank Accession No. U19183).

Figure 14: Predicted amino acid sequence of the complete ACCase gene of maize (SEQ ID NO:6).

Figure 15: Restriction map of ACCase genomic clones.

Figure 16: Partial nucleotide sequence of a Type A₁ ACCase genomic clone (SEQ ID NO:12).

Figure 17: Partial nucleotide sequence of clone 5A, a Type A ACCase genomic clone (SEQ ID NO:13).

Figure 18: Partial nucleotide sequence of six Type A₂ ACCase genomic clones (A-F) (SEQ ID NOs 14, 15, 16, 17, 18, and 19).

Figure 19: Partial nucleotide sequence of three Type B ACCase clones (SEQ ID NOs 20, 21 and 22).

Figure 20: SDS/PAGE analysis of chloroplast importation of ³⁵S-labeled ACCase polypeptides. (A) Pea and maize chloroplasts incubated with ³⁵S-labeled ACCase polypeptides for 30 minutes. (B) A time course analysis of the importation of ³⁵S-labeled ACCase polypeptides into maize chloroplasts.

Figure 21: Partial DNA sequence encoding oat ACCase (SEQ ID NO:23, Genbank Accession No. AF072737).

Figure 22: Partial amino acid sequence of an oat ACCase (SEQ ID NO:24).

Figure 23: Scan of test statistics generated by MQTL for the two recombinant inbred populations. A and B are the main effect and QTL × E interaction graphs for the Kanota X Ogle (KO) population, respectively. C and D are the main effect and QTL × E interaction graphs for the Kanota X Marion (KM) population, respectively. Thresholds for type I error rate of 5% after 1000 replications are reported above the graphs.

Figure 24: Histogram of % groat oil concentration for the mean of all environments relative to the number of recombinant inbred lines (RILs) with each parental genotype for the *XaccaseA* locus. A) The distribution for the Kanota X Ogle population, and B) distribution for the Kanota X Marion population.

Figure 25: Phylogram of ACCases.

Detailed Description of the Invention

The present invention provides a DNA molecule encoding a plant acetyl CoA carboxylase gene and methods for conferring herbicide tolerance and/or altering the oil content of plants by introducing and expressing a plant acetyl CoA carboxylase gene in the plant cells. In plants, acetyl CoA carboxylase plays a central role in regulating fatty acid synthesis and in the sensitivity of monocots to cyclohexanedione or aryloxyphenoxypropanoic acid herbicides.

In accord with the present invention, a plant acetyl CoA carboxylase gene is identified, isolated and combined with a promoter functional

in a plant cell to provide a recombinant expression cassette. A plant acetyl CoA carboxylase gene can be introduced and expressed in a plant cell. Depending on the type of plant cell, the level of gene expression, and the activity of the enzyme encoded by the gene, introduction of a plant acetyl CoA carboxylase gene into
5 the plant cell can confer herbicide tolerance and/or alteration of the oil of the plant cell.

In monocots, an exogenously introduced plant acetyl CoA carboxylase gene can be expressed at a level effective to render the cells of the plant tissue substantially tolerant to cyclohexanedione or
10 aryloxyphenoxypropanoic acid herbicide levels which normally inhibit a native or endogenous acetyl CoA carboxylase. A native acetyl CoA carboxylase is an enzyme that is normally encoded and expressed in the plant cell prior to transformation. An exogenously introduced plant acetyl CoA carboxylase gene is a gene which has been isolated and amplified from either the same or different
15 type of cell. Exogenous introduction and expression of a plant acetyl CoA carboxylase gene in both monocots and dicots can result in alteration of the oil content and quality of plant tissue and seeds. Exogenous introduction and expression in a host cell, such as a bacteria, can provide for sufficient amounts of plant acetyl CoA carboxylase to allow for crystallization and isolation of the
20 enzyme. Crystallized plant acetyl CoA carboxylase is useful to identify other herbicides that bind to and can inhibit plant acetyl CoA carboxylases. The enzyme could also be used to screen potential herbicidal compounds for efficacy.

A. Formation of an Expression Cassette

An expression cassette of the invention can comprise a chimeric
25 DNA molecule encoding a plant acetyl CoA carboxylase gene or an altered or functional mutant thereof operably linked to a promoter functional in a plant cell. The gene can code for a plant acetyl CoA carboxylase that is substantially tolerant to herbicides, preferably cyclohexanedione and/or
aryloxyphenoxypropanoic acid herbicides. An expression cassette of the
30 invention can also include an antisense DNA sequence that is substantially complementary to an acetyl CoA carboxylase gene or a portion thereof operably linked to a promoter functional in a plant cell.

1. **Isolation and Identification of a Gene Coding for a Plant Acetyl CoA Carboxylase**

A gene encoding a plant acetyl CoA carboxylase can be identified and isolated by standard methods, as described by Sambrook et al., Guide to
5 Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, NY (1989). The gene can be obtained either from monocot or dicot plant cells. When the gene encoding a plant acetyl CoA carboxylase is obtained from a dicot plant, the enzyme encoded by the gene exhibits tolerance to cyclohexanedione or aryloxyphenoxypropanoic acid herbicides. The gene can also be obtained from
10 herbicide-tolerant maize cell lines, prepared as described in U.S. Patent No. 5,162,602.

A gene encoding a plant acetyl CoA carboxylase can be identified by screening of a DNA or cDNA library generated from plant cells. Screening for DNA fragments that encode all or a portion of the gene encoding a plant
15 acetyl CoA carboxylase can be accomplished by complementation of an auxotrophic mutant of acetyl CoA carboxylase in *E. coli* (*fabE*) (Bachman, Microbiological Reviews, 47:180 (1983)) or yeast (*acc1*) (Michionada, Eur. J. Biochem., 111:79 (1980)) or by screening of plaques for binding to antibodies that specifically recognize a plant acetyl CoA carboxylase. DNA fragments that
20 can restore ACCase activity in *E. coli* or yeast and/or plaques carrying DNA fragments that are immunoreactive with antibodies to a plant ACCase can be subcloned into a vector and sequenced and/or used as probes to identify other cDNA or genomic sequences encoding all or a portion of a plant acetyl CoA carboxylase gene.

25 Specific examples of cDNA sequences encoding a portion of a plant acetyl CoA carboxylase gene include DNA fragments that include a DNA sequence that substantially corresponds to the coding sequence for the transcarboxylase active site of a plant acetyl CoA carboxylase, DNA fragments that include a DNA sequence that substantially corresponds to a coding sequence
30 for the biotin binding site of a plant acetyl CoA carboxylase, a DNA fragment encoding the 5' transcriptional start sequence of a plant acetyl CoA carboxylase gene, and a DNA fragment encoding the 3' transcriptional stop sequence for the acetyl CoA carboxylase gene. Substantially corresponding DNA sequences

share about 90% to about 100% DNA sequence homology. Especially preferred cDNA probes can be obtained from lambda clone #18-5 which include DNA sequences corresponding to the transcarboxylase active site domain and the biotin binding site domain. Lambda clone #18-5 includes EcoRI subclones of
5 3.9 kb, 1.2 kb, or 0.23 kb. Lambda subclone #18-5I is an 3.9 kb EcoRI subclone. The lambda subclone #18-5I has been deposited with the American Type Culture Collection, Rockville, MD, and given Accession No. 69236.

In a preferred version, a plant acetyl CoA carboxylase gene is identified and isolated from an herbicide tolerant maize cell line prepared as
10 described in Example II. A cDNA library can be prepared by oligo dT priming. Plaques containing DNA fragments can be screened with antibodies specific for maize acetyl CoA carboxylase. DNA fragments encoding a portion of an acetyl CoA carboxylase gene can be subcloned and sequenced and used as probes to identify a genomic acetyl CoA carboxylase gene. DNA fragments encoding a
15 portion of a maize acetyl CoA carboxylase can be verified by determining sequence homology with other known acetyl CoA carboxylases, such as chicken or yeast acetyl CoA carboxylase, or by hybridization to acetyl CoA carboxylase specific messenger RNA. Once DNA fragments encoding portions of the 5', middle and 3' ends as well as the transcarboxylase active site or biotin binding
20 site of a plant acetyl CoA carboxylase are obtained, they can be used to identify and clone a complete genomic copy of a maize acetyl CoA carboxylase gene.

To isolate a complete copy of a maize acetyl CoA carboxylase gene, a maize genomic library can then be probed with cDNA probes prepared as described above. Portions of the genomic copy or copies of a plant acetyl CoA
25 carboxylase gene can be sequenced and the 5' end of the gene are identified by standard methods including either DNA sequence homology to other acetyl CoA carboxylase genes or by RNAase protection analysis, as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, New York (1989). Once portions of the 5' end of the gene are
30 identified, complete copies of a plant acetyl CoA carboxylase gene can be obtained by standard methods, including by cloning or by polymerase chain reaction (PCR) synthesis using oligonucleotide primers complementary to the DNA sequence at the 5' end of the gene. The presence of an isolated full-length

copy of a plant acetyl CoA carboxylase gene can be verified by hybridization, partial sequence analysis, or by expression of a plant acetyl CoA carboxylase enzyme. The maize acetyl CoA carboxylase gene cloned and expressed from a maize herbicide tolerant cell line can be assessed for tolerance to
5 cyclohexanedione or aryloxyphenoxypropanoic acid herbicides by standard methods, as described in Example I.

An expression cassette of the invention can also contain an antisense DNA sequence. A antisense DNA sequence is a sequence that is substantially complementary to all or a portion of a coding sequence of a plant
10 acetyl CoA carboxylase gene. A substantially complementary sequence has about 90% to about 100% DNA sequence homology with that of the coding sequence of all or a portion of a plant acetyl CoA carboxylase. The antisense DNA sequence when expressed can act to inhibit the synthesis and expression of a native plant acetyl CoA carboxylase. Antisense sequences are preferably about
15 200 to 1000 nucleotides long in order to provide sufficient inhibition of synthesis and/or expression of a native acetyl CoA carboxylase. The inhibition of acetyl CoA carboxylase synthesis and gene expression by antisense DNA sequences can be confirmed in a transformed plant cell by standard methods for measuring the presence and/or activity of the enzyme such as described in Examples I and
20 V.

An expression cassette of the invention can also include a functional mutant of a plant acetyl CoA carboxylase gene. Mutants of a plant acetyl CoA carboxylase gene are substantially homologous to a plant acetyl CoA carboxylase gene and are functional if the acetyl CoA carboxylase expressed
25 retains significant enzyme activity. A mutant substantially homologous to a plant acetyl CoA carboxylase can share about 90% to 99.99% DNA sequence with that gene. For example, a mutant acetyl CoA carboxylase gene can code for a herbicide tolerant acetyl CoA carboxylase, or for an acetyl CoA carboxylase with altered substrate specificity so that the total amount of oil content in the
30 plants or seeds is increased, or for an enzyme with an altered substrate specificity so that synthesis of secondary metabolites such as flavonoids or anthocyanins is decreased. A preferred mutant is a gene coding for an acetyl CoA carboxylase

that is substantially tolerant to cyclohexanedione or aryloxyphenoxypropanoic acid herbicide

Altered or functional mutants of a gene coding for a plant acetyl CoA carboxylase can be obtained by several methods. The alteration or mutation of the ACCase gene can be accomplished by a variety of means including, but not limited to, the following methods.

1. spontaneous variation and direct mutant selection in cultures;
2. direct or indirect mutagenesis procedures on tissue culture of all cell types, seeds or plants; and
- 10 3. mutation of the cloned acetyl CoA carboxylase gene by methods such as site specific mutagenesis (Sambrook et al. cited supra), transposon mediated mutagenesis (Berg et al., Biotechnology, 1:417 (1983)) and deletion mutagenesis (Mitra et al., Molec. Gen. Genetic., 215:294 (1989)).

15 Mutants can be identified by a change in a functional activity of the enzyme encoded by the gene or by detecting a change in the DNA sequence using restriction enzyme mapping or partial sequence analysis.

In a preferred version, a functional mutant gene encoding for a plant acetyl CoA carboxylase tolerant to cyclohexanedione and/or aryloxyphenoxypropanoic acid herbicides is isolated from a maize herbicide tolerant cell line. The maize herbicide tolerant cell line was obtained as described in U.S. Patent No. 5,162,602, issued November 10, 1992, the disclosure of which is incorporated in Examples I-III. Briefly, partially differentiated cell cultures are grown and subcultured with continuous exposures to low herbicide levels. Herbicide concentrations are then gradually increased over several subculture intervals. Maize cells or tissues growing in the presence of normally toxic herbicide levels are repeatedly subcultured in the presence of the herbicide and characterized. Stability of the herbicide tolerance trait of the cultured cells may be evaluated by growing the selected cell lines in the absence of herbicides for various periods of time and then analyzing growth after exposing the tissue to herbicide.

Maize cell lines which are tolerant by virtue of having an altered acetyl CoA carboxylase enzyme can be selected by identifying cell lines having

enzyme activity in the presence of normally toxic levels of sethoxydim or haloxyfop. The tolerant maize cells can be further evaluated for whether acetyl CoA carboxylase is altered to a less sensitive form or increased in its level of expression.

- 5 Maize cell lines with a acetyl CoA carboxylase less sensitive to herbicide inhibition can be used to isolate a functional mutant gene of a plant acetyl CoA carboxylase. A DNA library from a maize cell line tolerant to herbicides can be generated and DNA fragments encoding all or a portion of an acetyl CoA carboxylase gene can be identified by hybridization to a cDNA probe
10 encoding a portion of the maize ACCase gene. A complete copy of the altered gene can be obtained either by cloning and ligation or by PCR synthesis using appropriate primers. The isolation of the altered gene coding for acetyl CoA carboxylase can be confirmed in transformed plant cells by determining whether the acetyl CoA carboxylase being expressed retains enzyme activity when
15 exposed to normally toxic levels of herbicides.

2. **Promoters**

Once a plant acetyl CoA carboxylase gene or functional mutant thereof or an antisense DNA sequence is obtained and amplified, it is combined with a promoter functional in a plant cell to form an expression cassette.

- 20 Most genes have regions of DNA sequence that are known as promoters and which regulate gene expression. Promoter regions are typically found in the flanking DNA sequence upstream from the coding sequence in both procaryotic and eukaryotic cells. A promoter sequence provides for regulation of transcription of the downstream gene sequence and typically includes from
25 about 50 to about 2,000 nucleotide base pairs. Promoter sequences also contain regulatory sequences such as enhancer sequences that can influence the level of gene expression. Some isolated promoter sequences can provide for gene expression of heterologous genes, that is a gene different from the native or homologous gene. Promoter sequences are also known to be strong or weak or
30 inducible. A strong promoter provides for a high level of gene expression, whereas a weak promoter provides for a very low level of gene expression. An inducible promoter is a promoter that provides for turning on and off of gene expression in response to an exogenously added agent or to an environmental or

developmental stimulus. Promoters can also provide for tissue specific or developmental regulation. An isolated promoter sequence that is a strong promoter for heterologous genes is advantageous because it provides for a sufficient level of gene expression to allow for easy detection and selection of transformed cells and provides for a high level of gene expression when desired.

The promoter in an expression cassette of the invention can provide for overexpression of acetyl CoA of a plant acetyl CoA carboxylase gene or functional mutant thereof. Overexpression of the gene is that amount of gene expression that results in an increase in tolerance of the plant cells to an herbicide or that results in an increase in the total oil content of the cells. Overexpression of an acetyl CoA carboxylase gene is preferably about a 2- to 20-fold increase in expression of an acetyl CoA carboxylase over the expression level of the native acetyl CoA carboxylase. The promoter can also be inducible so that gene expression can be turned on or off by an exogenously added agent. For example, a bacterial promoter such as the P_{tac} promoter can be induced to varying levels of gene expression depending on the level of isothiopyl galactoside added to the transformed bacterial cells. It may also be preferable to combine the gene with a promoter that provides tissue specific expression or developmentally regulated gene expression in plants.

Specific promoters functional in plant cells include the 35S cauliflower mosaic virus promoter, nopaline synthase (NOS) promoter and the like. Currently, a preferred promoter for expression in monocots is the 35S cauliflower mosaic virus promoter.

An acetyl CoA carboxylase gene can be combined with the promoter by standard methods as described in Sambrook cited *supra*. Briefly, a plasmid containing a promoter such as the 35S cauliflower mosaic virus promoter can be constructed as described in Jefferson, *Plant Molecular Biology Reporter*, 5,387 (1987) or obtained from Clontech Lab in Palo Alto, CA (e.g. pBI121 or pBI221). Typically these plasmids are constructed to provide for multiple cloning sites having specificity for different restriction enzymes downstream from the promoter. A gene for plant acetyl CoA carboxylase can be subcloned downstream from the promoter using restriction enzymes to ensure that the gene is inserted in proper orientation with respect to the promoter so that

the gene can be expressed. In a preferred version, a maize acetyl CoA carboxylase is operably linked to a 35 S Came promoter in a plasmid such as pBI121 or pBI221. Once a plant acetyl CoA carboxylase gene is operably linked to a promoter and the plasmid, the expression cassette so formed can be
5 subcloned into other plasmids or vectors.

3. Optional Sequences in the Expression Cassette

The expression cassette can also optionally contain other DNA sequences. The expression cassette can further be comprised of a chloroplast transit peptide sequence operably linked between a promoter and a plant acetyl
10 CoA carboxylase gene. If the expression cassette is to be introduced into a plant cell, the expression cassette can also contain plant transcriptional termination and polyadenylation signals and translational signals linked to the 3' terminus of a plant acetyl CoA carboxylase gene. The expression cassette can also optionally be further comprised of a plasmid.

15 Because one site of action for biosynthetic pathways involving plant acetyl CoA carboxylase is the chloroplast, an expression cassette of the invention can be combined with an exogenous DNA sequence coding for a chloroplast transit peptide, if necessary. An exogenous chloroplast transit peptide is one which is not encoded within the plant acetyl CoA carboxylase
20 gene. A chloroplast transit peptide is typically 40 to 70 amino acids in length and functions post-translationally to direct the protein to the chloroplast. The transit peptide is cleaved either during or just after import into the chloroplast to yield the mature protein. The complete copy of a gene encoding a plant acetyl CoA carboxylase may contain a chloroplast transit peptide sequence. In that
25 case, it may not be necessary to combine an exogenously obtained chloroplast transit peptide sequence into the expression cassette.

Exogenous chloroplast transit peptide encoding sequences can be obtained from a variety of plant nuclear genes, so long as the products of the genes are expressed as preproteins comprising an amino terminal transit peptide
30 and transported into chloroplast. Examples of plant gene products known to include such transit peptide sequences are the small subunit of ribulose biphosphate carboxylase, ferredoxin, chlorophyll a/b binding protein, chloroplast ribosomal proteins encoded by nuclear genes, certain heatshock proteins, amino

acid biosynthetic enzymes such as acetohydroxy acid synthase, 3-enolpyruvylphosphoshikimate synthase, dihydrodipicolinate synthase, and the like. Alternatively, the DNA fragment coding for the transit peptide may be chemically synthesized either wholly or in part from the known sequences of transit peptides such as those listed above.

Regardless of the source of the DNA fragment coding for the transit peptide, it should include a translation initiation codon and an amino acid sequence that is recognized by and will function properly in chloroplasts of the host plant. Attention should also be given to the amino acid sequence at the junction between the transit peptide and the plant acetyl CoA carboxylase enzyme where it is cleaved to yield the mature enzyme. Certain conserved amino acid sequences have been identified and may serve as a guideline. Precise fusion of the transit peptide coding sequence with the acetyl CoA carboxylase coding sequence may require manipulation of one or both DNA sequences to introduce, for example, a convenient restriction site. This may be accomplished by methods including site directed mutagenesis, insertion of chemically synthesized oligonucleotide linkers and the like.

Once obtained, the chloroplast transit peptide sequence can be appropriately linked to the promoter and a plant acetyl CoA carboxylase gene in an expression cassette using standard methods. Briefly, a plasmid containing a promoter functional in plant cells and having multiple cloning sites downstream can be constructed as described in Jefferson cited *supra*. The chloroplast transit peptide sequence can be inserted downstream from the promoter using restriction enzymes. A plant acetyl CoA carboxylase gene can then be inserted immediately downstream from and in frame with the 3' terminus of the chloroplast transit peptide sequence so that the chloroplast transit peptide is linked to the amino terminus of the plant acetyl CoA carboxylase. Once formed, the expression cassette can be subcloned into other plasmids or vectors.

When the expression cassette is to be introduced into a plant cell, the expression cassette can also optionally include 3' nontranslated plant regulatory DNA sequences. The 3' nontranslated regulatory DNA sequence preferably includes from about 300 to 1,000 nucleotide base pairs and contains plant transcriptional and translational termination sequence. Specific examples

of 3' nontranslated regulatory DNA sequences functional in plant cells include about 500 base pairs of the 3' flanking DNA sequence of the pea ribulose biphosphate carboxylase small subunit E9 gene, the 3' flanking DNA sequence of the octopine synthase gene, and the 3' flanking DNA sequence of the nopaline
5 synthase gene. These 3' nontranslated regulatory sequences can be obtained as described in An, Methods in Enzymology, 153:292 (1987) or are already present in plasmids available from commercial sources such as Clontech, Palo Alto, CA. The 3' nontranslated regulatory sequences can be operably linked to the 3' terminus of a plant acetyl CoA carboxylase gene by standard methods.

10 An expression cassette of the invention can also be further comprised of a plasmid. Plasmid vectors included additional DNA sequences that provide for easy selection, amplification and transformation of the expression cassette in procaryotic and eukaryotic cells. The additional DNA sequences include origins of replication to provide for autonomous replication of
15 the vector, selectable marker genes, preferably encoding antibiotic resistance, unique multiple cloning sites providing for multiple sites to insert DNA sequences or genes encoded in the expression cassette, and sequences that enhance transformation of prokaryotic and eukaryotic cells. The preferred vectors of the invention are plasmid vectors. The especially preferred vector is
20 the pBI121 or pBI221 vector formed as described by Jefferson cited supra.

Another vector that is useful for expression in both plant and procaryotic cells is the binary Ti vector PGA582. This binary Ti vector has been previously characterized by An, cited supra., and is available from Dr. An. This binary Ti vector can be replicated in procaryotic bacteria such as *E. coli* and
25 *Agrobacterium*. The *Agrobacterium* plasmid vectors can be used to transfer the expression cassette to plant cells. The binary Ti vectors preferably include the nopaline T DNA right and left borders to provide for efficient plant cell transformation, a selectable marker gene, unique multiple cloning sites in the T border regions, the *colE1* replication of origin and a wide host range replicon.
30 The binary Ti vectors carrying an expression cassette of the invention can be used to transform both prokaryotic and eukaryotic cells, but is preferably used to transform plant cells.

B. Method for Screening for Expression and/or Overexpression of a Plant Acetyl CoA Carboxylase Gene

A method for screening for expression or overexpression of a
5 plant acetyl CoA carboxylase gene is also provided by the invention. Once
formed, an expression cassette comprising an acetyl CoA carboxylase gene can
be subcloned into a known expression vector. The screening method in the
invention includes the steps of introducing an expression vector into a host cell
and detecting and/or quantitating expression of a plant acetyl CoA carboxylase
10 gene. This method of screening is useful to identify expression cassettes
providing for an overexpression of a plant acetyl CoA carboxylase gene,
antisense molecules that effectively inhibit acetyl CoA carboxylase synthesis,
and expression of an acetyl CoA carboxylase in the chloroplast of a transformed
plant cell.

15 Suitable known expression vectors include plasmids that
autonomously replicate in prokaryotic and eukaryotic cells. Specific examples
include plasmids such as the pBI121 or pBI221 plasmid constructed as described
by Jefferson cited *supra*, a binary Ti vector such as PG582 as described by An
cited *supra*, PUC119, or PBR322. The preferred expression system is a pBI121
20 or pBI221 plasmid.

An expression cassette of the invention can be subcloned into an
expression vector by standard methods. The expression vector can then be
introduced into prokaryotic or eukaryotic cells by standard methods including
protoplast transformation, silicon fiber-based transformation Agrobacterium
25 mediated transformation, electroporation, microprojectiles, e.g., preferably using
gold particles of about 6 to about 10 microns in diameter although diameters that
are about 100 to about 1000 times smaller may also be employed, and liposomes.
The expression vector can be introduced into plant cells such as tobacco,
Brassica, Black Mexican sweet corn, and Arabidopsis cells. The vector can also
30 be introduced into procaryotic cells such as E. coli or Agrobacterium.
Transformed cells can be selected typically using a selection marker encoded on
the expression vector.

Transient expression of a plant acetyl CoA carboxylase gene can
be detected and quantitated in the transformed cells. Gene expression can be

quantitated by a quantitative Western blot using antibodies specific for the cloned acetyl CoA carboxylase or by detecting an increase in specific activity of the enzyme. The tissue and subcellular location of the cloned acetyl CoA carboxylase can be determined by immunochemical staining methods using antibodies specific for the cloned acetyl CoA carboxylase. Sensitivity of the cloned acetyl CoA carboxylase to herbicides can also be assessed. Expression cassettes providing for overexpression of a plant acetyl CoA carboxylase or acetyl CoA carboxylase tolerant to herbicides can then be used to transform monocot and/or dicot plant tissue cells and to regenerate transformed plants and seeds.

C. Method of Imparting Cyclohexanedione or Aryloxyphenoxypropanoic Acid Herbicide Tolerance to a Plant

The invention provides a method of conferring cyclohexanedione or aryloxyphenoxypropanoic acid herbicide tolerance to a plant. The method includes the steps of introducing an expression cassette comprising a gene coding for a plant acetyl CoA carboxylase or a functional mutant thereof operably linked to a promoter into the cells of plant tissue and expressing the gene in an amount effective to render the cells of the plant tissue substantially tolerant to herbicides. An effective amount of gene expression to render the cells of the plant tissue substantially tolerant to the herbicide depends on whether the gene codes for an unaltered acetyl CoA carboxylase gene or a mutant or altered form of the gene that is less sensitive to the herbicides. Expression of an unaltered plant acetyl CoA carboxylase gene in an effective amount is that amount that provides for a 2- to 50-fold increase in herbicide tolerance and preferably increases the amount of acetyl CoA carboxylase from at least about 2- to 20-fold over that amount of the native enzyme. An altered form of the enzyme can be expressed at levels comparable to that of the native enzyme or less if the altered form of the enzyme has higher specific activity. Acetyl CoA carboxylase substantially tolerant to herbicides is an enzyme that is tolerant of levels of herbicide which normally inhibit a native acetyl CoA carboxylase and preferably can function in concentrations of herbicide of about 2- to 20-fold greater than are toxic to the native enzyme.

Herbicide tolerance can be achieved by at least two methods including: 1) by increasing the level of gene expression of a native or unaltered acetyl CoA carboxylase gene; or 2) by introducing an altered gene coding for an acetyl CoA carboxylase that is less sensitive to herbicide inhibition. The level of gene expression can be increased by either combining a plant acetyl CoA carboxylase gene with a promoter that provides for a high level of gene expression such as the 35S Came promoter or by introducing the gene into the cells so that multiple copies of the gene are integrated into the genome of the transformed plant cell. Formation of an expression cassette comprised of a plant acetyl CoA carboxylase gene operably linked to a promoter that can be expressed in an effective amount to confer herbicide tolerance has been described previously.

Most monocots, but not dicots, are sensitive to cyclohexanedione and/or aryloxyphenoxypropanoic acid herbicides. The preferred plant cells for introducing an expression cassette of the invention to achieve herbicide tolerance for the plant cells then are monocot plants. Monocot plants include corn, wheat, barley, sorghum, rice, and others. An expression cassette of the invention can be introduced by methods of transformation, especially effective for monocots including biolistic transformation of Type II embryogenic suspension cells as described by W.J. Gordon-Kamm et al., *Plant Cell*, 2, 603-618 (1990), M.E. Fromm et al., *Bio/Technology*, 8, 833-839 (1990) and D.A. Walters et al., *Plant Molecular Biology*, 18, 189-200 (1992) or by electroporation of type 1 embryogenic calluses described by D'Hafluin et al., *The Plant Cell*, 4, 1495 (1992). Transformed cells can be selected for the presence of a selectable marker gene. Transient expression of a plant acetyl CoA carboxylase gene can be detected in the transgenic embryogenic calli using antibodies specific for the cloned plant acetyl CoA carboxylase. Transformed embryogenic calli can be used to generate transgenic plants that exhibit stable inheritance of either the altered acetyl CoA carboxylase gene or overexpression of the acetyl CoA carboxylase gene. Maize cell lines exhibiting satisfactory levels of tolerance to herbicide are put through a plant regeneration protocol to obtain mature maize plants and seeds expressing the tolerance traits such as described in D'Hafluin, cited *supra.*, or An, cited *supra.* The plant regeneration protocol allows the

development of somatic embryos and the subsequent growth of roots and shoots. To determine that the herbicide-tolerance trait is expressed in differentiated organs of the plant, and not solely in undifferentiated cell culture, regenerated plants are exposed to herbicide levels which will normally inhibit shoot and root formation and growth.

5 Mature maize plants are then obtained from maize cell lines that are known to express the trait. If possible, the regenerated plants are self-pollinated. Otherwise, pollen obtained from the regenerated plants is crossed to seed-grown plants of agronomically important inbred lines. Conversely, pollen from plants of these inbred lines is used to pollinate regenerated plants. The genetics of the trait are then characterized by evaluating the segregation of the trait in the first and later generation progeny. Stable inheritance of overexpression of a plant acetyl CoA carboxylase or a functional mutant of a plant acetyl CoA carboxylase conferring herbicide tolerance to the plant is achieved if the plants maintain herbicide tolerance for at least about three to six generations.

Seed from transformed monocot plants regenerated from transformed tissue cultures is grown in the field and self-pollinated to generate true breeding plants. Progenies from these plants become true breeding lines which are evaluated for herbicide tolerance in the field under a range of environmental conditions. Herbicide tolerance must be sufficient to protect the monocot plants at the maximum labeled delivery rate under field conditions which cause herbicides to be most active. Appropriate herbicide concentrations and methods of application are those which are known and have been developed for the cyclohexanedione and/or aryloxyphenoxypropanoic acid herbicides in question.

In a preferred version, an expression cassette comprised of a maize acetyl CoA carboxylase gene isolated from a maize cell line tolerant to sethoxydim and haloxyfop and linked to the 35S Came promoter is introduced into an herbicide sensitive monocot tissue using biolistic transformation. Transformed calli are selected and used to generate transgenic plants. Transformed calli and transgenic plants can be evaluated for tolerance to sethoxydim and haloxyfop and for stable inheritance of the tolerance trait.

D. Method for Altering the Oil Content in a Plant

The invention also provides a method of altering the oil content in a plant. The method include the steps of introducing an expression cassette comprising a gene coding for plant acetyl CoA carboxylase or functional mutant thereof operably linked to a promoter functional in a plant cell into the cells of plant tissue and expressing the gene in an amount effective to alter the oil content of the plant cell. An alteration in the oil content of a plant cell can include a change in the total oil content over that normally present in that type of plant cell, or a change in the type of oil from that normally present in the plant cell. Expression of the gene in an amount effective to alter the oil content of the gene depends on whether the gene codes for an unaltered acetyl CoA carboxylase or a mutant or altered form of the gene. Expression of an unaltered plant acetyl CoA carboxylase gene in an effective amount is that amount that may provide a change in the oil content of the cell from about 1.2- to 20-fold over that normally present in that plant cell, and preferably increases the amount of acetyl CoA carboxylase about 2- to 20-fold over that amount of the enzyme normally present in that plant cell. An altered form of the enzyme can be expressed at levels comparable to that of the native enzyme or less if the altered form of the enzyme has higher specific activity.

An alteration in oil content of the plant cells according to the method of the invention can be achieved in at least two ways including:

- (1) an increase or decrease in expression of an unaltered plant acetyl CoA carboxylase gene; or
- (2) by introducing an altered or functional mutant plant acetyl CoA carboxylase gene coding for an enzyme that exhibits a change in specific activity.

The level of gene expression of an unaltered plant acetyl CoA carboxylase gene can be increased by either combining an unaltered plant acetyl CoA carboxylase gene with a promoter that provides for a high level of gene expression, such as the 35S cauliflower mosaic virus or by introducing the expression cassette and/or selecting for plant cells having multiple copies of a plant acetyl CoA carboxylase gene integrated into the genome. A decrease in expression of an unaltered acetyl CoA carboxylase can be achieved by transformation with an ACCase antisense

gene containing an expression cassette. When an altered or functional mutant plant acetyl CoA carboxylase gene codes for an enzyme that has an increase in specific activity, it may lead to an increase in total oil content of a plant cell even if the level of gene expression is comparable to that of the native enzyme. When
5 an altered or functional mutant acetyl CoA carboxylase gene codes for an enzyme having a decrease in specific activity, it may lead to a decrease in the total oil content of the plant cell compared to that normally present.

An expression cassette as described above can be introduced into either monocots or dicots. Preferably, the expression cassette is introduced into
10 dicot plants such as soybean, canola, and sunflower. An expression cassette can be introduced by standard methods including protoplast transformation, Agrobacterium-mediated transformation, microprojectiles, electroporation, and the like. Transformed cells or tissues can be selected for the presence of a selectable marker gene.

15 Transient expression of a plant acetyl CoA carboxylase gene can be detected in transformed cells or tissues by immunoreactivity with antibodies specific for the cloned acetyl CoA carboxylase. Overexpression of a plant acetyl CoA carboxylase can be detected by quantitative Western blots. A change in specific activity of the enzyme can be detected by measuring enzyme activity in
20 the transformed cells. A change in total oil content can also be examined by standard methods, as described in Clark & Snyder, *JACS*, 66:1316 (1989).

Transgenic plants and seeds can be generated from transformed cells and tissues showing a change in oil content or in the amount or specific activity of a plant acetyl CoA carboxylase using standard methods. It is
25 especially preferred that the oil content of the leaves, seeds, or fruits is increased.

In a preferred version, a maize acetyl CoA carboxylase gene is combined with a 35S cauliflower mosaic virus promoter in a vector such as pBI121 or pBI221 and introduced into soybean cells using the microprojectile method. Transformed soybean cells showing an increase in expression of acetyl
30 CoA carboxylase of at least about 2-fold or at least a 1.2-fold increase in oil content are selected. Transformed soybean cells exhibiting overexpression of acetyl CoA carboxylase or showing an increase in total oil content are used to generate transgenic plants and seeds.

E. Method of Producing Plant Acetyl CoA Carboxylase

The invention also provides a method of producing plant acetyl CoA carboxylase in a host cell. The method includes the steps of introducing an expression cassette comprised of a gene encoding a plant acetyl CoA carboxylase or functional mutant thereof into a host cell and expressing the gene in an amount sufficient to allow for crystallization of the plant acetyl CoA carboxylase. An amount sufficient to allow for crystallization of a plant acetyl CoA carboxylase is about 20- to 100-fold increase over the amount of plant acetyl CoA carboxylase that can normally be purified from plant cells, preferably about 2 to 10 mg protein. Crystallized plant acetyl CoA carboxylase can be used to identify other herbicides that can bind to and inhibit acetyl CoA carboxylase function. In addition, the availability of large amounts of purified enzyme provides for screening of the efficacy of such herbicides.

An expression cassette can include a promoter that is functional in either a eukaryotic or prokaryotic cell. The expression cassette can be introduced into a prokaryotic cell such as *E. coli*, or a eukaryotic cell such as a plant or yeast. The preferred cell is a prokaryotic cell used routinely in producing recombinant proteins such as *E. coli*. The expression cassette can be introduced and transformed cells selected by standard methods.

The plant acetyl CoA carboxylase gene can be expressed in an prokaryotic cell until sufficient amount of the enzyme is produced so that it can be crystallized. Plant acetyl CoA carboxylase can be isolated from bacterial cells using standard methods, including those described in Example V. The purified acetyl CoA carboxylase can then be crystallized and characterized by standard methods.

EXAMPLE I**Identification of Herbicide Mechanism and Site of Action**

The objective of this Example was to identify the mechanism whereby sethoxydim and/or haloxyfop inhibit fatty acid synthesis in maize. The results, reported in J. D. Burton et al., *Biochem. Biophys. Res. Comm.*, 148, 1039 (November 13, 1987), show that both sethoxydim and haloxyfop inhibit acetyl-coenzyme A carboxylase (ACCase) (EC 6.4.1.2) in maize chloroplasts.

A. Chemicals

Buffers and cofactors were purchased from Sigma Chemical Company (St. Louis, Missouri); [2-¹⁴C]acetate was purchased from Research Products International; [2-¹⁴C]pyruvate and [¹⁴C]NaHCO₃ were purchased from
5 New England Nuclear; and [2-¹⁴C]malonyl coenzyme A was purchased from Amersham. Sethoxydim was a gift from BASF (Parsippany, New Jersey), and haloxyfop was provided by Dow Chemical USA (Midland, Michigan).

B. Plant Growth Conditions

Corn (*Z. mays* L., 'B37 x Oh43') seeds were germinated in
10 darkness for 96 hours in vermiculite in an incubation chamber maintained at 30°C, 80% RH. Seedlings were then transferred to a growth chamber with a 16 hour light (25°C) and an 8 hour dark (20°C) cycle, 90% relative humidity (RH). After greening 48 hours, seedlings were returned to the dark incubation chamber for 12 hours to deplete chloroplast starch reserves. Seedlings were harvested 6
15 days after planting. Pea (*P. sativum* L., 'PI 9901-C') seedlings were grown in vermiculite in a growth chamber with a 16 hour light (21°C) and 8 hour dark (16°C) cycle, 80% RH. Peas were harvested 10 to 13 days after planting. Black Mexican Sweet (BMS) corn suspension cultures were maintained in a supplemented Murashige-Skoog (MS) medium (C. E. Green, *Hort. Sci.*, 12, 7-10
20 (1977)), and subcultured weekly by 20-fold dilution of the suspension culture into fresh medium.

C. Chloroplast Isolation

Chloroplasts from corn and pea seedlings were isolated at 4°C (K. Cline et al., *J. Biol. Chem.*, 260, 3691-3696 (1985)). Seedlings (50 g of shoots)
25 were homogenized in 200 ml buffer A (50 mM HEPES-NaOH pH 7.5, 330 mM sorbitol, 0.1% w/v BSA, 1 mM MgCl₂, 1 mM MnCl₂, 2 mM EDTA, 5 mM isoascorbate, 1.3 mM glutathione) in an omnimixer (five, 3-second bursts at full speed). The homogenate was filtered through six layers of cheesecloth and two layers of miracloth, and then centrifuged at 3000 g for 3 minutes with hand-
30 braking. The pellet was gently resuspended in buffer A and layered onto a preformed linear Percoll gradient (50 mM HEPES-NaOH pH 7.5, 330 mM sorbitol, 1.9 mM isoascorbate, 1.08 mM glutathione, 0.1% w/v BSA, 50% Percoll) which was centrifuged at 3000 g for 20 minutes in a Sorvall HB-4 rotor.

The lower band in the gradient, containing intact chloroplasts, was washed twice by gently resuspending it in 20 ml of buffer B (50 mM HEPES-NaOH, pH 7.5, and 330 mM sorbitol) followed by repelleting (3000 g, 5 minutes). The final pellet, consisting of intact chloroplasts, was resuspended in 2 to 3 ml of buffer B and stored on ice in the dark until use.

D. Fatty Acid Synthesis

[¹⁴C]acetate and [¹⁴C]pyruvate were used as precursors to measure fatty acid biosynthesis in isolated chloroplasts (B. Liedvogel et al., Planta, 169, 481-489 (1986)). [¹⁴C]acetate incorporation was assayed in a 0.5 ml-volume containing: 50 mM HEPES-NaOH (pH 7.5), 330 mM sorbitol, 5 mM KH₂PO₄, 10 mM NaHCO₃, 1 mM MgCl₂, 1 mM ATP, 0.1 mM CoA, 0.15 mM [¹⁴C]acetate (3.33 mCi/mmol), and chloroplasts (20 to 50 µg chlorophyll). [¹⁴C]pyruvate incorporation into fatty acids was assayed in the same medium except that it included 2 mM TPP, 1 mM NAD⁺, 0.15 mM [¹⁴C]-pyruvate (1.33 mCi/mmol), but no acetate. Assay suspensions were illuminated with 1400 µE/m²·second PAR at 25°C. Assays were initiated by the addition of the labeled substrate and stopped by the addition of 0.5 ml of 40% KOH. To determine the incorporation of radiolabel into a non-polar (fatty acid) fraction, each treatment was saponified at 90°C for 30 minutes in capped vials (P. B. Hoj et al., Carlsberg Res. Commun., 47, 119-141 (1982)). The vials were acidified with 0.5 ml 40% H₂SO₄, and carrier fatty acids (20 µg each of C 14:0, C 16:0, and C 18:0) were added. The assay mixture was extracted twice with 4 ml hexane. The extracts were combined, dried under N₂, and redissolved in 0.3 ml hexane. Aliquots (50 µl) were counted for radioactivity by liquid scintillation spectrometry.

Incorporation of [¹⁴C]malonyl-Coenzyme A into fatty acids (P. B. Hoj et al., *supra*; and J. B. Ohlrogge et al., Proc. Natl. Acad. Sci. USA, 76, 1194-1198 (1979)) was assayed using cell-free preparations from BMS tissue culture. Cells harvested during logarithmic growth phase were frozen in liquid nitrogen, ground with a mortar and pestle, and thawed in a medium containing: 0.1 M HEPES-KOH, pH 7.5; 0.3 M glycerol, and 5 mM DTT (buffer:tissue, 2:1, v/w). The homogenate was centrifuged at 12,000 g for 20 minutes. The supernatant was filtered through miracloth and centrifuged (125,000 g) for 60 minutes and then filtered through miracloth and assayed. Assays were conducted at 25°C in a

0.4 ml volume containing: 1.0 mM ATP, 0.32 mM NADPH, 0.38 mM NADH, 25 μ M CoA, 10 μ M acetyl-CoA, 25 μ g acyl-carrier protein, and 12 μ M malonyl-CoA (11.54 μ Ci/ μ mol). Reactions were initiated by addition of [14 C]malonyl CoA and stopped by addition of 0.4 ml 40% KOH. Label incorporation into fatty acids was determined as above. Chlorophyll (D. I. Arnon, *Plant Physiol.*, **24**, 1-15 (1949)) and protein (P. K. Smith et al., *Anal. Biochem.*, **150**, 76-85 (1985)) were determined as described therein.

E. Acetyl-Coenzyme A Carboxylase (ACCase) Activity

Maize chloroplasts, isolated as described above, were suspended in buffer C (0.1 M Tricine-KOH, pH 8.0; 0.3 M glycerol, and 1 mM DTT) and homogenized in a glass tissue homogenizer. The disrupted chloroplast fraction was centrifuged at 16,000 g for 15 minutes. The supernatant was desalted on a Sephadex G-25 column (1.5 x 5 cm equilibrated with 0.1 M Tricine-KOH, pH 8.0; and 0.3 M glycerol) and assayed directly. ACCase activity (B. J. Nikolau et al., *Arch. Biochem. Biophys.*, **211**, 605-612 (1981)) was assayed at 30°C in a 0.2 ml volume which contained 1 mM ATP, 3 mM acetyl coenzyme A, 2.5 mM MgCl_2 , 50 mM KCl, 0.5 mM DTT, and 15 mM [14 C] NaHCO_3 (0.17 mCi/mmol). Reactions were initiated by addition of acetyl coenzyme A and stopped by addition of 25 μ l of 12 N HCl. Product formation was determined by the radioactivity found in an acid stable fraction by liquid scintillation spectrometry. Enzyme activity was linear for 15 minutes.

F. Results

To probe for the site of herbicidal activity of sethoxydim and haloxyfop, labeled acetate, pyruvate, and malonyl-CoA were used individually as precursors for fatty acid synthesis. Isolated chloroplasts from corn seedlings incorporated [14 C]acetate and [14 C]pyruvate into a non-polar fraction (fatty acids). Acetate incorporation was linear for 30 min after a 5 min lag period, and dependent upon the addition of free acetyl coenzyme A. Addition of either 10 μ M sethoxydim or 1 μ M haloxyfop inhibited [14 C]acetate incorporation into fatty acids by 90% and 89%, respectively, as shown in Table I, below. Sethoxydim (10 μ M) and haloxyfop (1 μ M) also inhibited the incorporation of [14 C]pyruvate into fatty acids by 98% and 99%, respectively.

TABLE I

**Inhibition of [14 C]acetate and [14 C]pyruvate
Incorporation into Fatty Acids in
Corn Seedling Chloroplasts by
Sethoxydim (10 μ M) and Haloxyfop (1 μ M),
10 minute assay time**

	Acetate	Pyruvate
	---Activity (nmol/mg chl·min)---	
Control	4.4 ± 0.4^1	10.8 ± 2.3
	-----% Inhibition-----	
Sethoxydim	90 ± 2.5	98 ± 1.1
Haloxyfop	89 ± 3.1	99 ± 0.3

¹Results are expressed as mean of two experiments $\pm$ standard error.

The effect of 10 μ M sethoxydim and 1 μ M haloxyfop on [14 C]malonyl-CoA incorporation into fatty acids was determined using cell-free extracts from corn suspension cultures. Neither sethoxydim (10 μ M) nor haloxyfop (1 μ M) inhibited fatty acid synthetase activity. Thus, both herbicides inhibited fatty acid synthesis in intact chloroplasts from corn seedlings with either acetate or pyruvate as a precursor, but did not inhibit incorporation of malonyl-CoA into fatty acids. This suggests that ACCase which catalyzes the formation of malonyl-CoA is the site of action of these herbicides.

EXAMPLE II

Selection and Characterization of Herbicide-tolerant Cell Lines

A selection protocol to identify and isolate herbicide-tolerant maize cells was developed to minimize the adverse effects of high herbicide concentrations on somatic embryo development and plant regeneration capacity. The procedure involved exposing tissue to gradually increasing concentrations of herbicide beginning with a sethoxydim concentration representing 1/20th of lethal dose and doubling the herbicide concentration at approximately two-week intervals until the lethal dose (10 μ M sethoxydim) was reached. In this way, the

herbicide was allowed to take effect slowly with continuous selection pressure, thus permitting herbicide-tolerant cells to accumulate over time while not affecting the potential for plant regeneration.

A. Selection of a Sethoxydim-Tolerant Cell Line

5 Many selections were carried out utilizing the selection protocol described in the preceding paragraph. The selection of one such sethoxydim-tolerant cell line that was identified and characterized is described below in detail.

10 Approximately 100 grams of vigorously growing, regenerable, friable, embryogenic maize callus tissue established from an F₁ immature embryo resulting from the cross A188 x B73 were transferred to agar-solidified maintenance medium (Armstrong and Green, *Planta*, 164, 207 (1985)) in petri plates containing 0.5 μ M sethoxydim (BASF) (Parsippany, New Jersey). This callus line was designated 2167-9/2160-154. Forty plates were prepared and five
15 clumps of callus tissue weighing about 0.5 grams each were placed on each plate. The 0.5 μ M sethoxydim concentration was chosen from growth inhibition studies to provide less than 10-20% growth inhibition during the first two weeks of herbicide exposure. After 14 days, 0.25-0.5 gram pieces of tissue showing vigorous growth rate and retention of embryogenic morphology (i.e., presence of
20 somatic embryos) were subcultured on fresh medium containing 1.0 μ M sethoxydim. Eighty plates containing five pieces of tissue per plate were prepared. For each subsequent transfer, all callus tissue showing growth and somatic embryo forming ability was placed on fresh media containing a two-fold increased sethoxydim concentration. Therefore, callus was transferred at two-
25 week intervals to petri plates containing 0.5, 1.0, 2.0, 5.0 and 10.0 μ M sethoxydim. During the course of the selection process, the total number of lines decreased as the herbicide-mediated growth inhibition became more intense. Cell lines exhibiting growth on 10 μ M sethoxydim were designated as herbicide-tolerant and given an identification number. Two sethoxydim-tolerant lines were
30 recovered that exhibited uninhibited growth at 10 μ M sethoxydim. These lines were designated 2167-9/2160-154 S-1 and 2167-9/2160-154 S-2.

B. Characterization of Herbicide-Tolerant Maize Cell Line 2167-9/2160-154 S-2

Tolerant cell line 2167-9/2160-154 S-2 ("S-2") was characterized to evaluate: (1) the magnitude of sethoxydim tolerance; (2) cross-tolerance of
5 haloxyfop; and (3) the biochemical basis for the tolerance. Callus tissue from S-2 that had been maintained on 10 μ M sethoxydim was transferred to media containing up to 100 μ M sethoxydim. One-half gram of S-2 tissue was plated on a 7 cm filter paper as a lawn overlaying 50 ml agar-solidified culture medium containing 0, 0.5, 1.0, 2.0, 5.0, 10.0, 50.0 and 100 μ M sethoxydim, and cultured
10 for two weeks. Control cell line 2167-9/2160-154 was plated similarly on medium containing the same levels of sethoxydim. The results of this study are summarized in Figure 2. The control cell line growth after two weeks was inhibited 50% at 1 μ M sethoxydim. Growth of S-2 was not inhibited at 100 μ M sethoxydim, indicating that S-2 was at least 100-fold more tolerant than the
15 control callus line.

Growth of S-2 was inhibited with 0.65 μ M haloxyfop, whereas the control cell line was inhibited 50% with 0.02 μ M, indicating approximately a 30-fold increase in tolerance.

C. Acetyl-Coenzyme A Carboxylase (ACCase) Activity of Maize Cell Line S-2

Assays were conducted to determine if ACCase extracted from cell line S-2 was altered with respect to herbicide activity. ACCase activity of control tissue was 50% inhibited either by 1.5 μ M sethoxydim, or by 0.25 μ M haloxyfop. ACCase activity of S-2 tissue was inhibited 50% either by 70 μ M
25 sethoxydim, or by 1.8 μ M haloxyfop, indicating at least 40-fold and 7-fold decreases in herbicide sensitivity on concentration basis, respectively.

EXAMPLE III

Plant Regeneration and Production of Herbicide-Tolerant Seed

A. Plant Regeneration Protocol

30 Sixteen ca. 150 mg clumps of S-2 callus were transferred per 25 x 100 mm petri plate containing agar-solidified N6 basal salts and 6% sucrose and incubated 7-14 days in low light (20 μ E m⁻² s⁻¹). Several plates containing callus on plant regeneration medium were prepared. Callus was transferred to agar-

solidified Murashige-Skoog (MS) medium without hormones and incubated in high intensity light ($200 \mu\text{E m}^{-2} \text{s}^{-1}$) for shoot elongation. Developing plants (1-3 cm long) were isolated from the callus surface and transferred to magenta boxes containing agar-solidified MS salts, 2% sucrose with no hormones for two weeks of further growth. When plants reached the 2-3 leaf stage, they were transplanted to peat pots containing potting soil, and were incubated in the growth room until growing stably. Surviving plants were transferred to soil in 4" diameter plastic pots and grown in the greenhouse.

10 **B. Expression of Herbicide Tolerance in Plants Regenerated from S-2 Callus Tissue**

Groups of eight control (2167-9/2160-154 unselected) and eight S-2 plants were sprayed with either 0.0, 0.01, 0.05, 0.11, 0.22 or 0.44 kg/ha sethoxydim to determine whole plant sethoxydim-tolerance of greenhouse-grown plants. Control plants were killed by 0.05 kg/ha or more sethoxydim. 15 Plants regenerated from the S-2 cell line survived the 0.44 kg/ha sethoxydim treatment, indicating that S-2 plants exhibit at least 20-fold more tolerance of sethoxydim than control. Figure 3 shows the growth response of the regenerated plants seven days after treatment with 0.44 kg/ha sethoxydim. As shown in Figure 4, shoot height of regenerated S-2 plants was only slightly reduced 14 20 days after treatment with 0.44 kg/ha sethoxydim.

C. Seed Production from S-2 Plants

Plants surviving sethoxydim treatments of up to 0.44 kg/ha were transplanted to the genetics plot on the University of Minnesota campus, St. Paul, Minnesota. Additional S-2 plants were transplanted to the field that had 25 not been sprayed. Sixty-five 2167-9/2160-154 control plants and ninety-five S-2 plants were grown to maturity in the field. Plants were either self-pollinated or cross-pollinated to inbred maize lines A188, A619, A641, A661, A665, B37, B73, R806, and W153R. Control seed were produced by selfing 2167-9/2160-154 regenerated plants, or by crossing them with the inbreds listed above.

30 **D. Expression of Herbicide Tolerance in Progeny of Regenerated Plants**

Seeds obtained by the crossing procedure described above were viable and germinated normally. Seeds from thirty S-2 selfed plants and fifteen 2167-9/2160-154 control plants were planted in 25 x 50 cm trays of soil (28

seeds from each plant in one tray) and grown in the greenhouse. Seedlings at the 3-4 leaf stage were treated with 0.1, 0.44, and 1.1 kg/ha sethoxydim and evaluated for visual herbicide damage and shoot height. Based on visual rating of herbicide damage two weeks after treatment, selfed progeny of S-2 plants

5 segregated approximately 1:2:1 for healthy, uninjured plants: to plants showing partial injury: to dead plants, respectively, at 0.44 and 1.1 kg/ha sethoxydim treatments. All control progeny of 2167-9/2160-154 control plants were killed by 0.1 kg/ha and greater levels of sethoxydim. These results demonstrate dominant expression of sethoxydim tolerance indicating that sethoxydim

10 tolerance in S-2 plants is a heritable trait. Similar tests were conducted on progeny of S-2 plants crossed to the other inbreds. In all cases, these test cross progeny treated with 0.44 kg/ha sethoxydim segregated 1:1 for growing shoots versus dead shoots whether S-2 plants were used as male or female parents. These results confirm that sethoxydim tolerance is controlled by a single

15 dominant nuclear gene. In all cases, control plants crossed to the other inbreds were killed and therefore sethoxydim-sensitive.

E. Method for Obtaining Uniform Herbicide-Tolerant Seed

Progeny of S-2 plants surviving sethoxydim treatments of 0.44 and 1.1 kg/ha and showing no herbicide injury were transferred to the

20 greenhouse and grown to maturity. These plants may be selfed and their progeny evaluated for sethoxydim and haloxyfop tolerance to identify pure breeding herbicide-tolerant maize lines.

Progeny of S-2 plants crossed to inbred lines and exhibiting sethoxydim tolerance may be recurrently backcrossed to the same inbreds.

25 Progeny of each cross may be screened for sethoxydim-tolerance, and tolerant plants grown to maturity and again crossed to the recurrent parent. After six or seven cycles of backcrossing, sethoxydim-tolerant plants may be selfed and progeny screened for tolerance to produce homozygous sethoxydim tolerant maize inbreds.

EXAMPLE IV**Selection of Additional Herbicide-Tolerant Maize Cell Lines**

One primarily sethoxydim-tolerant maize cell line, 2167-9/2160-154 S-1, and two haloxyfop-tolerant maize cell lines, 2167-9/2160-154 H-1 and 2167-9/2160-154 H-2, were selected and characterized as follows:

A. Selection of Maize Cell Line 2167-9/2160-154 S-1

Maize cell line 2167-9/2160-154 S-1 was selected from maize cell culture using the protocol described in detail above for the selection of Line 2167-9/2160-154 S-2. Approximately 70 plants were regenerated from Line 2167-9/2160-154 S-1, and either self-pollinated or cross-pollinated to the inbred maize lines A188, A619, A641, A661, A665, B37, B73, R806, and W153R.

B. Selection of Maize Cell Line 2167-9/2160-154 H-1

Line 2167-9/2160-154 H-1 was selected from maize cell culture using a similar protocol described in detail above except maize callus tissue was selected using the herbicide haloxyfop. Maize callus tissue was initially plated on 0.01 μ M haloxyfop. At two-week intervals, surviving tissue was subcultured onto 0.05, 0.10 and 0.20 μ M haloxyfop. Approximately 50 plants were regenerated from Line 2167-9/2160-154 H-1, and were self-pollinated.

C. Selection of Maize Cell Line 2167-9/2160-154 H-2

Line 2167-9/2160-154 H-2 was selected from maize cell culture using a similar protocol described in detail for line 2167-9/2160-154 H-1. No plants have been successfully regenerated from this line.

D. Characterization of Lines 2167-9/2160-154 S-1, H-1 and H-2

The tolerant callus cultures were characterized to determine the magnitude of sethoxydim and haloxyfop tolerance. Callus tissue from these lines was evaluated in experiments as described above in the characterization of line 2167-9/2160-154 S-2. Table II summarizes the results of these studies. Line 2167-9/2160-154 S-1 and Line 2167-9/2160-154 H-2 showed a four-fold increase in haloxyfop tolerance, while Line 2167-9/2160-154 H-1 exhibited approximately a 60-fold increase in haloxyfop tolerance. Neither haloxyfop selected line showed a significant degree of sethoxydim tolerance, while the

sethoxydim selected line S-1 exhibited approximately a 100-fold increase in sethoxydim tolerance.

TABLE II

Herbicide Tolerance of Cell Lines S-1, H-1 and H-2

Cell Line	Herbicide	
	Haloxypop	Sethoxydim
2167-9/2160-154 S-1	4 ¹	100
2167-9/2160-154 H-1	61	0
2167-9/2160-154 H-2	4	0

¹The numbers represent the fold increase in herbicide concentration that results in a 50% reduction in growth of the selected cell lines compared to the unselected control cell line 2167-9/2160-154.

E. Herbicide Inhibition of Acetyl Coenzyme A Carboxylase of Maize Cell Lines S-1, H-1 and H-2

Acetyl Coenzyme A Carboxylase (ACCase) was extracted from cell lines S-1, H-1 and H-2 and assayed as described in detail for maize cell line S-2, above. Table III below summarizes the results of these studies. The ACCase from line S-1 was more tolerant of both sethoxydim and haloxypop, while the ACCase from line H-1 was more tolerant of haloxypop, but not of sethoxydim. The ACCase from line H-2 showed no difference from the unselected parent line 2167-9/2160-154 in sensitivity to either herbicide.

However, cell line H-2 exhibited approximately a five-fold higher level of ACCase activity as compared to the unselected parent line 2167-9/2160-154. Thus, selection for sethoxydim or haloxypop tolerance resulted in a less

sensitive ACCase in cell line S-1 and H-1, as well as a higher level of ACCase activity in cell line H-2.

TABLE III

**Herbicide Inhibition of ACCase of Maize
Cell Lines S-1, H-1 and H-2**

Cell Line	Herbicide	
	Haloxyfop	Sethoxydim
2167-9/2160-154 S-1	3	4
2167-9/2160-154 H-1	7	0
2167-9/2160-154 H-2	0	0

¹The numbers represent the fold increase in herbicide concentration that inhibits ACCase activity of the selected cell lines by 50% compared to the unselected parent cell line 2167-9/2160-154.

Deposit of Seeds

Seeds from representative S-2 plants (Ex. III (B)) and H-1 plants (Ex. IV(B)) have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852 USA on October 25, 1988 and assigned accession numbers ATCC 40507, and ATCC 40508, respectively.

EXAMPLE V

Formation of cDNA Clones Encoding ACCase

A. ACCase Purification

The acetyl CoA carboxylase enzyme was isolated and purified from plant tissues and characterized. The purified enzyme was used to generate antibody reagents useful in identifying cDNA clones encoding the gene or portions of the gene for ACCase.

ACCase was extracted from frozen shoots of 7-d-old maize (*Zea Mays* L. inbred A619 or B73) seedlings grown in a growth chamber (24°C, 90% RH, 16-h daylength at 210 $\mu\text{E m}^{-2} \text{s}^{-1}$). The outermost leaf and blade were removed and the remainder of the shoot was frozen in liquid N₂. Embryos and endosperm tissue from developing kernels were harvested from field-grown ears at 36 to 40 days after pollination (DAP). Black Mexican Sweet corn (BMS)

maize suspension cells were obtained from cultures as previously described (W.B. Parker et al., Plant Physiol., **92**, 1220-1225 (1990)). Tissues were stored in liquid N₂ until used.

Extraction and purification steps were performed at 0 to 4°C.

- 5 Crude extracts of leaf, bundle sheath strands, embryo, endosperm, and BMS cells were prepared from frozen tissue as described by W.B. Parker et al., Proc. Nat'l Acad. Sci. USA, **87**, 7175-7179 (1990), except that extraction buffer contained 0.1 M Tricine-KOH, pH 8.3, 0.3 M glycerol, 5 mM DTT, 2mM Na₂EDTA, and 0.5 mM phenyl methonyl sulfonyl fluoride (PMSF). Triton X-
10 100 (0.01% v/v) was added to bundle sheath strand extracts and to some whole leaf extracts. For some experiments, additional protease inhibitors (leupeptin, 2 µg mL⁻¹; pepstatin A, 100 µg mL⁻¹; benzamidine, 1 mM; ε-amino-n-caproic acid, 5 mM; and soybean trypsin inhibitor, 10 µg mL⁻¹) were included. Filtered homogenates were centrifuged 20 minutes at 30,000g. A portion of the crude
15 supernatant fraction was immediately boiled 5 minutes in 1 volume of SDS sample buffer (W.B. Parker et al., Plant Physiol., **92**, 1220-1225 (1990)) for SDS-PAGE analysis; the remainder was desalted on a 10-mL Sephadex G-25 column into extraction buffer minus PMSF.

- ACCCase was purified from the crude extract supernatant in four
20 steps. This fraction was brought to 30% saturation with solid (NH₄)₂SO₄, stirred 15 minutes, and centrifuged 20 minute at 20,000g. The supernatant was then brought to 40% saturation with (NH₄)₂SO₄ solution, stirred 30 minutes, and centrifuged. The pellet was dissolved in 5 mL extraction buffer, microfuged 5 minutes, and the resulting supernatant was applied to a Sephacryl S-300 gel
25 filtration column (Pharmacia; 2.5 x 46 cm) equilibrated with S-300 buffer (0.1 M Tricine-KOH, pH 8.3, 0.5 M glycerol, 0.5 mM DTT, 2 mM Na50 mM KCl). In later experiments a Sephacryl S-400 column was used. Fractions (2.5 mL) were eluted at 0.75 mL min⁻¹. ACCCase activity eluted shortly after the void A₂₈₀ peak (V₀ = 75 mL). Active fractions were pooled, brought to 4.25 mM MgCl₂ (from a
30 0.5 M solution), and applied at 0.2 mL min⁻¹ to a Blue Sepharose CL-6B (Pharmacia; 1.5 x 15 cm) equilibrated with Blue sepharose buffer (S-300 buffer containing 4.25 mM MgCl₂ and 10 mM NaHCO₃). The column was washed overnight with 150 mL buffer (0.45 mL min⁻¹). ACCCase activity was then eluted

with 50 mL buffer plus 10 mM ATP (0.45 mL min^{-1}). Active fractions were pooled and applied to an FPLC Mono-Q HR 5/5 anion-exchange column (Pharmacia) equilibrated with S-300 buffer minus KCl. The column was washed with 30 mL S-300 buffer minus KCl and then with a 48-mL, 0 to 500 mM KCl gradient in S-300 buffer (0.25 mL min^{-1}). Fractions (1 mL) from the two peaks of ACCase activity were pooled separately. All purification fractions were desalted into S-300 buffer and assayed for ACCase activity and protein.

ACCase was also analyzed from mesophyll chloroplasts and bundle sheath strands. Mesophyll chloroplasts from homogenates of 7- to 8-day-old seedlings that were kept in the dark 24 hours prior to harvesting were isolated on a linear Percoll gradient according to J.D. Burton et al., Pesticide Biochemistry and Physiology, **34**, 76-85 (1989), except that buffers contained 0.6 M sorbitol and centrifugation g -forces were reduced by 25%. Intact chloroplasts were taken from the discrete lower green band present after Percoll gradient centrifugation (G. Morieux et al., Plant Physiol., **67**, 470-473 (1981)). Pelleted chloroplasts were lysed by resuspending them in ACCase extraction buffer plus PMSF and 0.01% (v/v) Triton X-100. Bundle sheath strands were obtained from the original leaf homogenate material retained on a 70- μm filter after re-homogenizing the retentate five times in a total of 2 L buffer. Triton X-treated, desalted leaf, mesophyll chloroplast, and bundle sheath strand extracts were assayed for activities of Rubisco (G. Zhu et al., Plant Physiol., **97**, 1348-1353 (1991)), NADP-dependent malate dehydrogenase (M.D. Hatch et al., Biochem. Biophys. Res. Commun., **34**, 589-593 (1969)), phosphoenolpyruvate carboxylase (R.C. Leegood et al., "Isolation of Membranes and Organelles from Plant Cells," Academic Press, New York, 185-210(1983)), catalase (Worthington Biochemicals, 1972), and fumarase (R.L. Hill et al., Methods Enzymol., **13**, 91-99 (1969)), and for total chlorophyll (D.E. Arnon, Plant Physiol., **24**, 1-5 (1949)). Mesophyll chloroplast preparations were judged to be relatively free of contamination by bundle sheath chloroplasts because they contained 3-fold greater NADP-dependent malate dehydrogenase and one-tenth as much Rubisco activity (mg^{-1} chlorophyll) than bundle sheath strand extracts. Mesophyll chloroplast preparations also contained $\leq 2.6\%$ as much catalase, fumarase, and phosphoenolpyruvate carboxylase activities (mg^{-1} chlorophyll) as

did whole-leaf extracts, indicating they were relatively free of peroxisomal, mitochondrial, or cytoplasmic components.

ACCase activity as measured by acetyl-CoA-dependent $\text{H}^{14}\text{CO}_3^-$ (ICN, 2.07 GBq mmol⁻¹) incorporation into acid-stable product previously shown to be malonyl-CoA (J.D. Burton et al., Pesticide Biochemistry and Physiology, 34, 76-85 (1989)). Assays of desalted purification fractions or crude, desalted tissue extracts contained up to 50 and 25% (v/v) enzyme, respectively. In some experiments methylcrotonyl-CoA or propionyl-CoA were substituted for acetyl-CoA (E.S. Wurtele et al., Archives of Biochemistry and Biophysics, 278, 179-186 (1990)). Avidin (10 U mL⁻¹) was included in some assays. Herbicide inhibition assays contained 1% (v/v) ethanol plus or minus 1 μM haloxyfop (2-[4-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy] propanoic acid, Dow Chemical Co. analytical grade racemic mixture) or 10 μM sethoxydim (2-[1[(ethoxylmino)butyl]-5-[2-(ethylthio)-propyl]-3-hydroxy-2-cyclohexene -1-one, Li salt, BASF Corp. technical grade). Data are means plus standard error of three assays.

Protein concentrations were determined in duplicate with the Bio-Rad Coomassie blue dye-binding assay as described by the manufacturer, using BSA as the standard.

Centrifuged crude extracts and proteins in purification fractions and immunoprecipitation supernatants were separated by SDS-PAGE in 6 or 7.5% gels as previously described (W.B. Parker et al., Plant Physiol., 92, 1220-1225 (1990)). Purification fractions were precipitated in 10% (v/v) TCA, washed with 80% (v/v) acetone, and air-dried 10 minutes prior to electrophoresis. Proteins in gels were stained with silver (J. Heukeshoven et al., Electrophoresis, 6, 103-112 (1985)). High molecular weight protein standards for SDS-PAGE (Pharmacia) were used to estimate polypeptide masses.

The four-step purification procedure shown in Table IV typically yielded 30 to 190 μg of highly purified ACCase from 50 grams (fresh weight) of maize inbred A619 or B73 seedling leaves. ACCase activity in the crude supernatant fraction precipitated between 30 and 40% saturation with $(\text{NH}_4)_2\text{SO}_4$, which appeared to increase total ACCase activity approximately 38%. Crude extract components might have depressed the reaction rate shown in Table IV

because the assay mixture contained 50% enzyme (v/v). In tests of fractions from another purification, enzyme velocity was proportional to enzyme concentration in assay mixtures containing up to 25% (v/v) crude extract, but 50% (v/v) mixtures were not tested. ACCase activity eluted from the Sephacryl S-300 gel filtration column slightly after the green void peak. Approximately 56% of the S-300 fraction ACCase activity was recovered from the Blue Sepharose column, primarily in the initial ATP-containing fractions (12.5 mL). Both 10 mM NaHCO₃ and 4.25 mM MgCl₂ (1- and 0.85-fold standard assay concentrations, respectively) were included in the Blue Sepharose buffer because they improved the total and specific ACCase activity remaining after batch absorption to Blue Sepharose beads, elution with ATP, and desalting into extraction buffer minus PMSF. Neither NaHCO₃ nor MgCl₂ improved enzyme stability of crude extracts. Mono-Q anion-exchange chromatography resulted in separation of two ACCase activity peaks which eluted at approximately 210 mM (designated ACCase II) and 250 mM KCl (designated ACCase I), as previously observed for a hybrid maize variety (J.L. Howard et al., *FEBS Lett.*, **261**, 261-264 (1990)). ACCase I comprised about 85% of the total activity recovered from the column (29% of the original crude extract activity) and had high specific activity (Table IV). The specific activity of ACCase II was less than 30% that of ACCase I. Both activities were inhibited >90% by avidin, as previously reported (J.L. Howard et al., *FEBS Lett.*, **261**, 261-264 (1990)). The mass of native ACCase I was estimated to be approximately 490 kD by gel filtration on Superose 6.

TABLE IV

**Purification of ACCase I From
Maize Inbred A619 Seedling Leaves^a**

5

All fractions were desalted into S-300 buffer and assayed for protein and acetyl CoA dependent incorporation of [¹⁴C]HCO₃⁻ into acid-stable products.

10	Step	Protein	Activity	Specific Activity	Fold Purification	Activity Yield
		<i>mg</i>	<i>units^b</i>	<i>units/mg</i>		<i>%</i>
	Crude extract	215	2.45	0.0114	1	100
	30-40% (NH ₄) ₂ SO ₄	45.1	3.37	0.0748	6.56	138
	S-300	10.7	3.35	0.313	27.5	137
15	Blue Sepharose	1.50	1.86	1.24	109	76
	Mono-Q (ACCase I)	0.130	0.720	5.54	486	29

^a Data are from one purification experiment starting with 50 g fresh weight of tissue and are representative of data obtained for eight purifications.

20

^b Unit = 1 μmol acid-stable product min⁻¹.

B. Formation and Specificity of Antibodies to ACCase

25

Antibodies are sensitive reagents that allow for the identification of gene products from cDNA and other cloned genes. Antibodies to purified ACCase were prepared and used to screen for cDNA clones encoding all or a portion of a gene for ACCase.

Antiserum to maize ACCase was obtained by immunizing a female New Zealand White rabbit (Egli et al., *Plant Physiol.*, 101, 499 (1993)). An intramuscular injection of 100 μg of Mono-Q-purified, SDS-denatured ACCase I in Freund's complete adjuvant was followed by subcutaneous injections of 20 to 100 μg of gel-purified ACCase I polypeptide in acrylamide plus incomplete adjuvant every 4 to 6 weeks, for a total of six injections. Serum was stored at -20°C in 0.02% (w/v) NaN₃.

35

For Western blots, proteins in SDS gels were electrophoretically transferred to Immobilon (W.B. Parker et al., *Plant Physiol.*, 92, 1220-1225

(1990)) for 1 hour at 20 V in a Bio-Rad Transphor semi-dry blotter and then stained with Ponceau S (E. Harlow et al., "Antibodies - A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1988)).

Destained blots were blocked with Tris-buffered saline plus 0.5% (v/v) Tween-20 (Bio-Rad), and 10% (w/v) bovine serum (for antiserum blots only). ACCase and biotinylated proteins were detected with immune serum (1/10,000) plus goat anti-rabbit IgG-alkaline phosphatase conjugate or with avidin-alkaline phosphatase (W.B. Parker et al., *Plant Physiol.*, 92, 1220-1225 (1990)). Blots were repeated at least three times.

10 For immunoprecipitations, equal ACCase activities (0.58 nmol min⁻¹) in crude extracts were desalted into S-300 buffer containing 0.1 M KCl and incubated 1 hour at 25°C with 16 µL buffer or with 16 µL serum consisting of 0 to 100% ACCase antiserum in preimmune serum. Immune complexes were incubated 1 hour at 25°C with a 2-fold (IgG binding) excess of Protein
15 A-agarose and then microfuged 5 minutes to obtain immunoprecipitation supernatant fractions. ACCase activity of supernatants was expressed as a percent of the 100% preimmune serum control. Data are means plus SE of three replicate assays for each of two sets of extracts.

Western blots and silver-stained gels of purification fractions
20 separated by 7.5% SDS-PAGE showed that neither ACCase I nor ACCase II Mono-Q fractions contained biotinylated polypeptides smaller than 212 kD. A polypeptide > 212 kD was the primary protein component of the ACCase I Mono-Q fragment (Fig. 5). The ACCase II fraction contained a biotinylated polypeptide > 212 kD and a large amount of a 55 kD non-biotinylated
25 polypeptide. Fractions from earlier purification steps contained additional biotinylated proteins of approximately 74, 75, and 125 kD (Fig. 5).

To better compare the biotinylated polypeptides > 212 kD in ACCase fractions I and II, we used 6% SDS-PAGE, which showed that the mass of ACCase II was approximately 8 kD less than that of ACCase I. Molecular
30 masses were estimated to be 219 kD (ACCase II) and 227 kD (ACCase I), based on comparisons with polypeptide standards and the observation (N.R. Palosaari, *Plant Physiol.*, 99(S), 359 (1992)) that, on Phastgels (Pharmacia), ACCase I polypeptide was slightly smaller than dodecameric horse spleen ferritin (238 kD;

M. Heusterspreute et al., *FEBS Lett.*, **129**, 322-327 (1981)). All purification fractions through the Blue Sepharose step contained both ACCase I and II polypeptides. Rapid extraction of leaves in buffer containing five additional protease inhibitors, or a 4 hour incubation of extracts at 25°C, had little or no effect on the relative amounts of the two polypeptides, suggesting that ACCase II is not a breakdown product of ACCase I.

Antiserum to ACCase I strongly recognized the ACCase I polypeptide in crude extracts and showed little or no recognition of ACCase II polypeptides. No bands were recognized by preimmune serum. Assuming that avidin binds similarly to ACCase I and II polypeptides, it appears that the amount of ACCase II on the Western blot was slightly less than the amount of ACCase I. However, the relative staining with antibody compared to avidin indicated that the antibody had significantly less affinity for ACCase II than ACCase I.

To determine whether the same ACCase polypeptides were expressed in different maize cell types, proteins in mesophyll chloroplasts and crude extracts of leaves, endosperm tissue, embryos, and BMS cells were separated by SDS-PAGE. All preparations contained a predominant biotinylated polypeptide of approximately 227 kD (ACCase I) that was strongly recognized by ACCase antiserum or avidin. Similar 227 kD band densities were observed when gel lanes were probed with either avidin or ACCase antiserum. The 219 kD ACCase II polypeptide was readily detected in leaves only by avidin binding, but was in low abundance or not detected in extracts from other tissues. Only the 227 kD ACCase I polypeptide was detected in purified mesophyll chloroplasts, however, suggesting that the 219 kD ACCase II polypeptide is localized elsewhere in mesophyll cells or in other cell types of young leaves. ACCase activity and a > 212 kD biotinylated polypeptide(s) were also found in bundle sheath strand extracts, but low yields prevented us from determining the type of ACCase present. Two other major biotinylated polypeptides of 75 and 74 kD were found in all tissues. Other non-biotinylated proteins of 66 kD (faint) and 55 kD were also recognized by ACCase antiserum. The 55 kD polypeptide was only found in leaves; it was also present in both ACCase I and II Mono Q

fractions (Fig. 5) and was identified as the Rubisco large subunit based on its comigration with protein immunoprecipitated by spinach Rubisco antiserum.

ACCCase antiserum immunoprecipitated at least 75% of ACCCase activity from crude, desalted extracts of leaves, endosperm tissue, embryos, and BMS cells (Fig. 6), indicating that most of the ACCCase activity in these tissues is immunologically related to the ACCCase I polypeptide of leaves. Less activity was precipitated from leaves (75%) than from other tissues, particularly embryos (98%). Compared to immunoprecipitation, inhibition of ACCCase activity by antiserum in solution was less than 20% as effective in reducing ACCCase activity.

The substrate specificity of ACCCase from different purification fractions was examined to compare [^{14}C]HCO $_3^-$ incorporation in the presence of different acyl-CoA substrates. Both ACCCase I and II utilized propionyl-CoA 40 to 50% as rapidly as acetyl-CoA at 50 to 500 μM substrate even though they contained no biotinylated polypeptides (Figure 5) the size of known propionyl CoA carboxylases (70 to 75 kD; see E.S. Wurtele et al., Archives of Biochemistry and Biophysics, 278, 179-186 (1990)). Activities in the presence of both acetyl-CoA and propionyl-CoA (250 or 500 μM each) were approximately 90 (ACCCase I) to 130% (ACCCase II) that of 500 μM acetyl-CoA alone. Crude leaf extracts utilized propionyl-CoA and methylcrotonyl-CoA 60% as efficiently as acetyl-CoA. Methylcrotonyl CoA carboxylase activity was reduced 85% by gel filtration and was completely removed by Blue Sepharose affinity chromatography.

ACCCase I and II differed significantly in their inhibition by either haloxyfop or sethoxydim (Fig. 7). Acetyl-CoA or propionyl-CoA-dependent H $^{14}\text{CO}_3^-$ incorporation by ACCCase I was strongly inhibited (65 to 80%) by 1 μM haloxyfop or 10 μM sethoxydim, while ACCCase II activity was inhibited less than 50% for all herbicide/substrate combinations examined.

C. Cloning and Identification of Maize cDNA Clones Encoding ACCCase

Maize cDNA clones encoding a portion of the ACCCase gene were identified by screening a DNA library generated from maize. The cDNA clones were used to identify the sequence of the ACCCase gene and to identify the genomic DNA fragments encoding the gene or genes for ACCCase.

A λ gt11 cDNA library from maize inbred A188 seedlings was prepared by standard method for oligo-dT priming, as described for pea cDNA. (Gantt and Key, *Eur. J. Biochem.*, 166:119-125 (1987). Plaque lifts of the maize cDNA library were screened with maize ACCase antiserum (Egli et al., *Plant*
5 *Physiol.*, 101, 499 (1993)) to identify plaques expressing ACCase-like proteins, as described by Sambrook et al., cited *supra*. (1989). The initial screen of 800,000 plaques yielded 120 positives. Rescreening and plaque purification reduced the number of positives to 14. All 14 clones bound ACCase antibodies that, when eluted from plaque lifts (J. Hammarback et al., *J. Biol. Chem.*,
10 *265*:12763 (1990)), recognized a 227-kD biotinylated polypeptide on SDS-PAGE western blots of embryo and leaf crude extracts. The strongest western blot reaction was obtained with cDNA clone #15-14. The six best clones were digested with EcoRI to excise maize cDNA inserts. Total insert sizes ranged from 1.2 to 5.1 kb indicating the clones most likely did not contain the full
15 coding sequences for the mature 219-kD and 227-kD ACCase polypeptides (minimum estimates of 6.1 and 6.3 kb, respectively).

Clone #15-14 contained three EcoRI fragments of 2.0, 1.2 and 0.23 kb shown in Figure 8. Southern blots showed that the 1.2 and 2.0-kb fragments of clone #15-14 each hybridized to different fragments in the other
20 five clones, with the exception of clone #4-4 which only contained a 1.2-kb fragment. The six maize cDNA clones contained EcoRI fragments that hybridized to a large transcript (ca. 7.8 kb) on Northern blots of total RNA from maize leaves, embryos and endosperm (Fig.9). BMS cell culture RNA also contained a 7.8 kb transcript, but the hybridization signal is not evident on this
25 exposure (Fig. 9). The relative abundance of the 7.8-kb transcript in embryos was higher than the other sources which is consistent with their ACCase activity.

The three EcoRI fragments were subcloned from cDNA clone #15-14 into BlueScript vector and sequenced by the dideoxy chain termination method (Sequenase 2.0 USB) initially using T3 and T7 primers and then
30 oligonucleotide primers based on insert sequence. A clone #16-6 was also sequenced in a similar manner. Clone #16-6 included three EcoRI fragments of 3.1 kb, 1.2 kb, and 0.23 kb and had additional sequence located upstream from that of clone #15-14. After comparing the sequence and determining that the

sequence was the same, the additional 1.2 kb sequence at the 5' end was sequenced.

Clone #18-5 was sequenced in a similar manner. Clone #18-5 included 3.9 kb, 1.2 kb, and 0.23 kb EcoRI fragments and contains an additional
 5 1.9 kb 5' sequence upstream from clone #15-14. Subclone #18-5I (3.9 kb EcoRI fragment) has been deposited with the American Type Culture Collection and given Accession No. 69236.

GenBank, PIR-29, and Swiss-Prot 19 data banks have been searched for amino acid homology with the corresponding amino acid sequences
 10 of the three subclones of clone #18-5. Peptide sequences corresponding to the maize cDNA subclones had higher similarity to chicken, rat, and yeast ACCases than to any other peptide sequence in the data banks. Figure 8 illustrates the relative organization of the 3.9, 1.2 and 0.23-kb EcoRI fragments of clone #18-5 and their co-linearity and extent of amino acid identity with chicken ACCase
 15 cDNA sequence. This comparison shows that the maize clone #18-5 has a large region near the 3' end with high amino acid identity (40 to 61%) to chicken ACCase, a longer region with 23% identity in the middle of the 3.9-kb sequence, and a short region with 52% identity near the 5' of the 3.9 kb sequence.

Portions of the sequence of the #18-5I subclone have been
 20 identified as encoding domains of the ACCase enzyme of functional significance. Those functional regions include a fragment that spans the presumed transcarboxylase active site in the enzyme having the following presumed sequence SEQ ID NO: 2:

25 1112-856
 GTT CCT GCA AAC ATT GGT GGA CCT CTT CCT ATT ACC AAA CCT CTG GAC
 CCT CCA GAC AGA CCT GTT GCT TAC ATC CCT GAG AAC ACA TGC GAT CCA
 CGT GCA GCT ATC TGT GGT GTA GAT GAC AGC CAA GGG AAA TGG TTG GGT
 GGT ATG TTT GAC AAA GAC AGC TTT GTG GAG ACA TTT GAA GGA TGG GCA
 30 AAA ACA GTG GTT ACT GGC AGA GCA AAG CTT GGA GGA ATT CCT GTG GGC
 GTC ATA GCT GTG GAG ACA

This functional domain is contained in the sequence 1112 to 856 base pair from the 3' stop codon or carboxy terminus region of the ACCase coding sequence of maize. This transcarboxylase active sequence is also present in clone #15-14.

Another functional region that has been identified spans the
5 12 base pair sequence encoding the biotin binding site having the following peptide sequence SEQ ID NO: 3:

5' GTT ATG AAG ATG 3'
Val Met Lys Met

10

The biotin binding site is encoded approximately 30% in from the 5' (N-terminus) end of rat, chicken and yeast ACCase genes. These functional domains are useful in mapping and further identifying other cDNA and/or genomic fragments encoding ACCase genes.

15

The cDNA clones encoding portions of the acetyl CoA carboxylase genes are useful to identify the sequence of the gene or genes and are useful as probes to locate the genomic copies of the gene or genes. Because the ACCase antibodies used to screen the λ gt11 library recognize both the 219 and 227 kD ACCase polypeptides, it has not been determined which polypeptide
20 is encoded by these less than full length clones. It is likely that the majority of the clones encode the 227 kD polypeptide since that polypeptide is more abundant in the leaf tissue source of the DNA library and the antibodies have a higher affinity for the 227 kD ACCase polypeptide.

EXAMPLE VI

25

Isolation and Sequencing of Genomic Encoded ACCase Genes and a Complete cDNA Sequence of a Maize ACCase Gene

The maize genome has been analyzed to identify copy number and location of the genomic copies of ACCase gene or genes. Four distinct types of maize ACCase genomic clones have been identified, termed A1, A2, B1
30 and B2 (see below).

To obtain genomic copies of ACCase genes, a maize B73 genomic library (Clontech, Palo Alto, CA) was screened with the 2 kb subclone from #15-14 and several clones of about 15 kb were identified as having homology to the ACCase cDNA. Restriction mapping and partial sequence
35 analysis revealed two types of genomic clones (Type A and Type B) that differed

in restriction sites and in their position relative to the ACCase partial cDNA sequence as shown in Figure 8.

The 2.5 kb EcoRI-SaII fragment (#16) from the Type A genomic clone and the 3.0 kb EcoRI-EcoRI fragment (#34) from the Type B genomic clone were shown to hybridize to the 3.9 kb probe from #18-5 and were subcloned into the Bluescript vector and sequenced. Approximately 1.5 kb of DNA sequence from the genomic Type A 2.5 kb fragment were 100% identical to coding sequence from the 3.9 kb cDNA subclone #18-5I described in Example V; the remaining sequence exhibited no identity with the cDNA clone and presumably represents a noncoding intron sequence. A 350 nucleotide sequence derived from the genomic Type B 3.0 kb fragment was about 95% identical to the cDNA clone indicating that its coding sequence differs from that of genomic Type A. These results also indicate that the maize genome encodes at least two different genes encoding a polypeptide having acetyl CoA carboxylase activity.

To identify and clone the remainder of the gene representing the amino-terminus of maize ACCase, additional regions from the Type A genomic clone have been subcloned and partly sequenced. The 3.5 kb HindIII-HindIII fragment (#28) has been sequenced for about 400 nucleotides from each end. The 3' end of #28 shows significant homology to the amino acid sequence from the chicken sequence located about 0.5 kb from the start of the chicken gene.

The complete sequence for fragment #28 can be obtained and analyzed to determine whether it contains the 5' end of the ACCase coding region. The start of the transcribed region, and thus the likely start of the coding region for ACCase, can be identified by using the genomic clones in RNase protection analysis (J. Sambrook et al., "Molecular Cloning - A Laboratory Manual," Cold Spring Harbor Press, Cold Spring Harbor, New York (1989)). Based on sequence data from the genomic clone, alignment, as shown in Figure 8, with sequences of other ACCases and identification of potential open reading frames, oligonucleotide primers can be constructed to synthesize cDNA molecules representing the amino terminus of the ACCase gene. These molecules can be hybridized to genomic Type A DNA fragments such as #28 and the nonhybridizing portions digested with S1 nuclease. The end of the protected fragment are determined by analysis on a DNA sequencing gel.

To synthesize the remaining coding region between the end of the cDNA clone #18-5 and the start of transcription, two oligonucleotide primers were synthesized. Primer 1 is complementary to the DNA sequence: (SEQ ID NO:4)

5 5' GCCAGATTCC ACCAAAGCAT ATATCC 3'

near the 5' end of cDNA subclone #18-5I and can be used as a primer for synthesis of cDNA molecules from maize seedling, leaf or embryo RNA.

A primer corresponding to a DNA sequence near the transcription
10 start site can be used in combination with primer 1 for the amplification of DNA by the polymerase chain reaction (PCR). Several independent clones are then sequenced and their sequences compared to the known sequence of the Type A genomic clone to determine the exact coding sequence corresponding to that maize gene for ACCase. A similar strategy can be used to obtain the complete
15 coding sequence for genomic Type B ACCase.

The remaining cDNA sequence was obtained by three successive rounds of RT-PCR using oligonucleotide primers based on genomic apparent exon (5') and known cDNA (3') sequences. The primers used to amplify nucleotides 1-240 of the cDNA were 28sst-a5+ (SEQ ID NO. 7) and
20 28sst-6at3+ (SEQ ID NO:8), nucleotides 217-610 of the cDNA were 28sst-5+ (SEQ ID NO:9) and 28-2t3+ (SEQ ID NO:10), and nucleotides 537-2094 of the cDNA were ACCPCR5' (SEQ ID NO:11) and I55' (SEQ ID NO:30) (Table V). PCR products corresponding to nucleotides 1-240, 217-610, and 537-2094 of the final sequence were cloned into PCR-script (Stratagene).

TABLE V

cDNA Position	5' primer designation	5' primer sequence	3' primer designation	3' primer sequence
nt-1-240	28sst-a5+	GGTCTTCAATTGTGCTGTCTGG (SEQ ID NO:7)	28sst-6at3+	CCTTGACGAACAGACTGGCTGTGC (SEQ ID NO:8)
nt 217-610	28sst-5+	CACAGCCAGTCTGTTCGTCAAGG (SEQ ID NO:9)	28-2t3+	CCTCTACGTAATTGGTCAGC (SEQ ID NO:10)
nt 537-2094	ACCPCR5'	CATAGCTATGGCAACTCCGG (SEQ ID NO:11)	I55'	GGATATATGCTTTGGTGGAATCTGGC (SEQ ID NO:30)

The original 5.4-kb cDNA clone #18-5 and PCR products from at least three individual PCR per oligonucleotide pair were sequenced in both directions by the dideoxy chain-termination method, using either Sequenase II (U.S. Biochemicals) or ABI 373 (Applied Biosystems, Inc.) protocols. No
5 sequence differences were found in regions of clone overlaps. The complete sequence of the cDNA of maize ACCase (nucleotides 1-7470 SEQ ID NO:5) and its corresponding amino acid sequence (amino acids 1-2325 SEQ ID NO:6) are shown in Figures 13 and 14. The 7470 bp cDNA includes a 459 nucleotide 3' untranslated region and 36 nucleotides of 5' untranslated sequences.

10 The first Met codon in the cDNA (nucleotides 37-39) was identified as the start codon based on its similarity to consensus initiation sequences (Kozak, J. Cell. Biol., 108, 229 (1989); Lutcke et al., Embo. J., 6, 43 (1987)). An in-frame stop was found in the genomic sequence 6 nucleotides upstream of the sequenced cDNA, and RT-PCR analysis of this region suggested
15 that the in frame stop codon was also present in the cDNA. The 3' end of the coding sequence was defined by two stop codons present in the large open reading frame after nucleotide 7011. The translated coding sequence predicted a polypeptide of 2325 amino acids (257 kD; SEQ ID NO:6) which was 79 to 81% identical to the multifunctional (MF) ACCases from alfalfa (Shorrosh et al.,
20 Proc. Nat'l. Acad. Sci., 91, 4323 (1994)) and wheat (Gornicki et al., Proc. Nat'l. Acad. Sci., 91, 6860 (1994)), and to a 118-amino acid predicted polypeptide of a rice expressed sequence tag (Genbank accession # D39099, T. Sasaki), but only 53 to 55% identical to ACCase from other eukaryotes.

In a pileup alignment of plant ACCases (Genetics Computer
25 Group, Madison, Wisconsin), Met 1 of both maize and *Brassica napus* ACCases was located about 130 amino acids upstream of the conserved sequence VDEFCKALGG, compared to only 25 amino acids upstream for other plant ACCases. The predicted 2325 amino acids of maize ACCase contains a biotinylation site at position 806, within the conserved MKM motif (Ton et al.,
30 Eur. J. Biochem., 215, 687 (1993)). The arrangement and amino acid sequence of binding sites (Shorrosh et al., Proc. Nat'l. Acad. Sci., 91, 4323 (1994)) for ATP (amino acids 318-333), biotin (amino acids 799-811; biotin at 806),

acetyl-CoA (amino acids 1952-1961), and carboxybiotin (amino acids 1662-1711) were highly conserved among all MF ACCases.

EXAMPLE VII

Characterization of other Genomic Clones

5 The initial restriction fragment length polymorphism (RFLP) analysis of EcoRI-digested total DNA from three maize inbred lines showed one band when probed with the 2 kb subclone from #15-14 (internal to gene) and two bands when probed with the 1.2 kb subclone (near the 3' end of the gene). Fragments homologous to the 2 kb probe were monomorphic and the more
10 intense of the two bands hybridizing with the 1.2 kb probe was dimorphic. As discussed in Example V, these results support the view that maize contains at least two distinguishable ACCase genes and that they may be quite similar for much of the coding region. Additional genomic Southern blots of a set of recombinant inbred lines were used to map polymorphisms for the ACCase
15 probes to maize chromosomes. One polymorphism was mapped to the short arm of chromosome 2; other polymorphisms were not evident in these initial tests to identify a chromosomal location for other maize ACCase genes.

 The isolation and restriction mapping of additional genomic clones from a B73 genomic library (Clontech) resulted in the identification of
20 four different types of clones termed A1, A2, B1 and B2 (Figures 16-19) which had 96% nucleotide sequence identity. Types A and B correspond to previously published pA3 and pA4 cDNAs (Ashton et al., *Plant Mol. Biol.*, 24, 35 (1994)) and differ from pA3 and pA4 by ~ 4% in their coding sequences.

 Type A and B genomic clones have linear sequence homology
25 except for an insertion in an intron of the Type B genes about 1400 bp 3' of the A1(SEQ ID NO:5) translation start site. Analysis of the insert boundaries revealed a 3-bp target site duplication and a 6-bp direct repeat, and further sequence analysis showed the presence of two new and unique LINE elements (Long Interspersed Nuclear Elements) in B1 and B2. Mammalian LINE
30 elements are highly abundant (10^4 to 10^5 copies), 6 to 7 kb long, and have frequent 5'-end deletions and an A-rich 3' terminus. They are flanked by short direct repeats, and contain two ORFs, one encoding a reverse transcriptase. Three LINE elements (Cin4, 50-100 copies in maize; del2, 250,000 copies in

lily; BNR1, 2-5% of genome in sugarbeet) have been described in plants (Leeton et al., *Mol. Gen. Genet.*, 237, 97 (1993); Schmidt et al., *Chromo. Res.*, 3, 335 (1995); Schwarz-Sommer et al., *EMBO J.*, 6, 3873 (1987)). Maize ACCase B1 has one unique LINE element and B2 has two. The two B2 LINE elements were
 5 characterized by differences in their reverse transcriptase sequence. The B genomic clone inserts have characteristic LINE features including cysteine motifs and a possible polyA tail, and high abundance. The LINE insert also has been found in an intron of the maize *Shrunken-2* gene (Hannah et al., *Plant Physiol.*, 98, 1214 (1992)).

10 The partial nucleotide sequence (3489 nucleotide) of a Type A1 ACCase genomic clone is shown in Figure 16 (SEQ ID NO:12). The clone is a HindIII fragment which includes nucleotides 1-931 of the cDNA in Figure 13 (SEQ ID NO:5), and the first four introns within the coding region, at positions 240 (460 nucleotides), 296 (480 nucleotides), and 872 (76 nucleotides) of SEQ
 15 ID NO:5. The clone also has 1395 nucleotides 5' to the cDNA of SEQ ID NO:5 (i.e., 1431 nucleotides 5' of the translational start at nucleotide position 1432).

 The partial nucleotide sequence (1328 nucleotide) of another Type A clone is shown in Figure 17 (SEQ ID NO:13). The partial sequence is all 5' untranslated sequence and contains a 7 base insert between nucleotides
 20 279-290, but is otherwise identical to SEQ ID NO:12.

 The partial nucleotide sequence of six Type A2 clones is shown in Figure 18 (1565, 1168, 638, 558, 976 and 852 nucleotides; SEQ ID NOs 14, 15, 16, 17, 18 and 19, respectively).

 Within the A1-A2 clone pair, identified differences are in introns
 25 and 5' UTR sequences. The A2 genomic clone is weakly amplified with Type A1 PCR primers specific for the 5' UTR if the 3' primer employed is for a conserved amino acid sequence found in all ACCases (e.g., 28sst-110F, ACTGTGCGTTTGAGAAGGTC, SEQ ID NO:28, and 28sst-2T3+, CCTCTACGTAATTGGTCAGC, SEQ ID NO:29). The A2 amplified product is
 30 the same size as that from the Type A1 genomic clone, and restriction analysis indicated a difference in sequence from A1. Sequence differences in the 5' region should provide a means to distinguish between expression of A1 and A2 ACCase genes and to determine whether A2 also encodes a CTP.

The partial nucleotide sequence (231, 207 and 180 nucleotides) of three Type B clones is shown in Figure 19 (SEQ ID NOs 20, 21 and 22, respectively).

The cDNAs corresponding to genomic clones A2, B1 and B2 are
5 cloned and sequenced in a manner similar to that described above. The derived amino acid sequences are aligned with known ACCase sequences. If putative CTP sequences are identified, functionality is tested as described below. Also if the tissue specificity and developmental timing of expression differ for different ACCase genes, the sequences of the promoter regions of the corresponding
10 genomic clones are compared. Gene-specific probes for specific ACCase genes can provide more information on their roles in lipid synthesis (plastid and cytoplasmic isoforms), secondary metabolism (cytoplasmic isoforms), and herbicide resistance (likely plastid isoforms).

A 3' Type A1 ACCase cDNA probe mapped to chromosome 2S
15 (Egli et al., Maize Genetics Newsletter, 68, 92 (1994)) and to 10L (Caffrey et al., Maize Gen. Coop., 69, 3 (1995)). Two 5' Type A1 cDNA probes which span the transit peptide mapped to chromosome 2S in the same location as the 3' probe (see maize genetic map, 1996 version, Maize Genomic Database). PCR primers 28sst-97F (CCTTTTATGGCACTGTGCG, SEQ ID NO:25) and 28sst-6t3+;
20 (CATCGTAGCCTATATGAGGACG, SEQ ID NO:26) located in non-coding regions of A1 that span the chloroplast transit peptide were used to amplify a B73 chromosome-specific product which segregated with the resistance trait. A nearby 5' primer (28sst-a5+, SEQ ID NO:7) amplified all genotypes and functioned as a positive control. Herbicide resistance due to the *Acc1-S3*
25 mutation segregates (29/29 individuals to date) with production of a Type A 5' end-specific PCR product derived from the mutant parent while herbicide sensitive plants lack the transit sequence (15/17 progeny). Two individual plants which contained B73-specific DNA at this location died of unknown causes while grown in the presence of herbicide.

30 Mutations in maize that confer resistance to cyclohexanedione and aryloxyphenoxypropionate herbicides by means of an altered ACCase target are found at two non-allelic loci, *Acc1* and *Acc2*. A1 and A2 appear to encode plastidic ACCases and correspond to the *Acc1* and *Acc2* herbicide resistance

loci. *Acc2* has been mapped to 10 L (VanDee, M.S. Thesis, University of Minnesota (1994)). *Acc1* is the site of five allelic mutations including *Acc1-S2* and *-S3* (Marshall et al., *Theor. Appl. Genet.*, 83, 435 (1992)), and has been mapped to chromosome 2.

5 Only one plastidic ACCase polypeptide was identified by SDS-PAGE of maize leaf extracts, although 2-D gel analyses might provide evidence for a second, highly similar isoform. Of the two ACCase isoforms, only ACCase I shows altered herbicide inhibition in *Acc1-S2* mutants, and most of the ACCase activity in leaves and developing embryos is herbicide-resistant and
10 thus attributed to the *Acc1-S2* gene product.

 Although a 3' ACCase probe has been mapped both to 10L near *Acc2-S5* and to 2S, the conserved sequence of ACCase genes and lack of polymorphism in multiple bands complicates identification of genes encoded at these loci. The Type A1 ACCase gene is probably located on chromosome 2,
15 since (i) 5' untranslated and chloroplast transit peptide probes from Type A1 hybridize to two bands (dark and light) in maize inbreds, and (ii) analysis of maize-oat addition lines carrying maize chromosomes 2 through 9 indicates the dark band is on chromosome 2 and the light band is on chromosome 1 or 10.

 Type B ACCase genes are likely to encode cytosolic isoforms.
20 Given that cytosolic malonyl-CoA is a precursor in the synthesis of many secondary metabolites including flavonoids (e.g. maysin, a corn silk component associated with corn earworm resistance), these cytosolic ACCases can have agronomic utility.

 Northern blot analysis of total maize RNA with an ACCase probe
25 (nucleotides 3400-5932) showed a single 8.3 kilobase band. To determine whether the expression of ACCase RNAs was developmentally regulated, blots of total RNA from 16 to 42 DAP (days after pollination) embryos were probed with an ACCase cDNA fragment. Transcript abundance peaked about 23 DAP and the steady state pattern was similar to *in vitro* ACCase enzyme activities and
30 protein measured from developing embryos. Type A- and B- specific ³²P-CTP-labeled antisense transcripts were 780 nt long (662 nt of ACCase sequence + 118 nt of vector/promoter sequence) and were identical except for 15 base mismatches scattered along their length. Each antisense transcript was

hybridized to total RNA from embryos at 16, 20, 23, and 42 DAP and digested with RNase A/TI mixture to yield a 662-base fragment specific to the probe used. The results showed that the Type A transcript was more abundant than Type B at all tested stages, and that only Type A remained high in older embryos. Types A and B had similar expression patterns and peaked around 20-23 DAP. The ratio of Type A:B mRNA in leaves was about 2:1, similar to its relative abundance in cDNA expression libraries.

EXAMPLE VIII

Expression of the Maize ACCase Chloroplast Transit Peptide

The N-terminus of the predicted maize ACCase polypeptide is longer than that of predicted cytosolic ACCase isoforms and has several characteristics typical of chloroplast transit peptides within the first approximately 73 amino acids of the predicted N-terminal sequence. The CTP cleavage site motif is not found in the putative maize ACCase CTP, although only about 30% of known CTPs contain this consensus sequence (Gavel and von Hejne, FEBS Lett., 261, 455 (1990)). However, the maize ACCase N-terminus appears to have several other properties typical of known CTPs: (1) a lack of acidic residues in amino acids 1-49, (2) high Ser + Thr content (69% within amino acid residues 23-35), (3) an R-rich region between S- and D-rich regions in amino acid residues 36-49, and (4) a predicted turn $\rightarrow\beta$ sheet within amino acid residues 58-73 (von Hejne and Nishikawa, FEBS Lett., 278, 1 (1991)).

The ability of the amino acid sequence contained within the N-terminal 100 amino acids of the translated maize acetyl-CoA carboxylase (ACCase) cDNA to direct the N-terminal portion of the maize ACCase biotin carboxylase domain into chloroplasts was tested *in vitro* by methods used extensively in the literature (see Cline et al., J. Biol. Chem., 260, 3691 (1985); Lubben and Keegstra, Proc. Nat'l. Acad. Sci., 83, 5502 (1986)). The criteria for import was that (1) *in vitro*-synthesized, ^{35}S -labeled protein was imported into chloroplasts, and (2) the transported protein was smaller than the original translation product, by an amount which corresponds to the removal of the expected CTP. Import studies utilized either maize or pea chloroplasts. Pea chloroplasts have been reported to correctly import proteins from many different species, including maize (Nieto-Sotelo et al., Plant Physiol., 93, 1321 (1990)).

Alternatively, the function of the putative maize ACCase CTP is tested by inserting the first 258 coding nucleotides of maize ACCase in frame with and 5' of a GUS reporter gene in pBI221 (Clontech). This construct and the pBAR plasmid are used to co-transform maize "Black Mexican Sweet" suspension cells by particle bombardment. Basta-resistant transformants are selected, and GUS activity and/or protein is assayed in surviving cultures or in plasmids isolated from transformants.

A partial ACCase construct consisting of nucleotides 1-833 of SEQ ID NO: 5 including the putative CTP (nucleotides 37 to 256) and the first domain within the biotin carboxylase region (identified by amino acid sequence comparison with *E. coli* biotin carboxylase; see Waldrop et al., *Biochem.*, **33**, 6249 (1994)) was amplified by PCR and cloned into the EcoRV site of PCR-script (Stratagene) to create the plasmid pBCN1. A corresponding plasmid lacking CTP sequences (nucleotides 278-833) was also made (-pBCN1). The protein encoded by -pBCN1 begins at amino acid residue 83 (Val → Met).

Constructs were transformed into *E. coli* SURE cells (Stratagene). Restriction analysis of pBCN1 with BamH1 and HindIII indicated that the 5' end of the ACCase was located adjacent to the T7 RNA polymerase binding site in PCR-Script. A partial sequence of pBCN1 obtained by using the T7 sequencing primer and the ABI373 automated sequencing protocol confirmed this orientation and showed that the pBCN1 insert sequence was identical to maize ACCase cDNA for at least the first 300 nucleotides and that it included the maize ACCase Met 1 ATG. An acyl carrier protein clone containing a CTP (spinach ACPII, a gift of Dr. John Ohlrogge, Michigan State University) can be used as a positive control. These constructs are used for *in vitro* transcription, translation, chloroplast import, and SDS-PAGE analysis of products in the same manner as pBCN1.

Purified pBCN1 was digested with EcoRI to linearize the plasmid at the 3' end of the BCN1 insert, electrophoresed in 1.5% agarose, and the plasmid band at approximately 3.8 kb was excised and Gene-Cleaned (BioLab 101). The purified band was digested with 20 µg proteinase K to remove any residual RNase, extracted with phenol and then chloroform under RNase-free conditions. DNA content was estimated by ethidium bromide fluorescence in

droplets, relative to λ DNA standards (Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed. (1989). One μ g of pBCN1 DNA was transcribed into capped RNA with the T7/mMessage mMachine kit (Ambion). Uncapped transcripts (Sp6 RNA polymerase; Promega) of pea RUBISCO small subunit (SSU; Anderson et al., Biochem. J., **240**, 709 (1986)) were also transcribed. RNA yield was estimated by determining the % incorporation of a ^{32}P -ATP (Amersham) into a precipitable product, according to the Ambion kit instructions. Electrophoresis and autoradiography of ^{32}P -labeled product showed that it contained a single RNA band of approximately 895 nucleotides, as expected.

The RNA transcripts were translated into ^{35}S -labeled polypeptides with Ambion's wheat germ IVT kit and approximately 45 μCi ^{35}S -methionine (Amersham; 37 TBq/mmol) in a 50- μl reaction. Labeled proteins were held on ice 6 hours prior to their use in chloroplast import experiments.

Pea (cv. "Little Marvel") and maize (inbred A188) plants were grown in a growth chamber at 25°C, 16 hour day length. Chloroplasts were isolated from pea and maize leaves 7 days after planting, respectively, as previously described (Burton et al., Pestic Biochem. and Physiol., **34**, 76 (1989); Egli et al., Plant Physiol., **101**, 499 (1993)). Intact mesophyll chloroplasts were washed in resuspension buffer [50 mM HEPES-KOH, pH 7.8 plus 0.33 M (pea) or 0.66M (maize) sorbitol] in preparation for import assays. Suspensions were diluted to obtain 75 μg chlorophyll/0.3 ml (Arnon, Plant Physiol., **24**, 1 (1949)).

Import experiments were carried out essentially as described by Cline et al. (J. Biol. Chem., **260**, 3691 (1985)). Import reactions containing 0.3 ml pea or maize chloroplast suspension, 40 μl ^{35}S -translation mixture, 3 mM Mg-ATP and 10 mM Met were incubated under light for 1-30 minutes at 25°C. Unimported proteins were digested for 30 minutes with 40 μg of thermolysin, and proteolysis was stopped with 10 mM EDTA.

Chloroplasts were re-isolated by centrifuging them through 1-ml 40% v/v Percoll gradients in the presence of resuspension buffer plus 3 mM Mg-ATP, 10 mM Met, and 20 mM EDTA, washed twice in the same buffer, and resuspended in 65 μl of 1 mM MgCl_2 /10 mM Tris buffer, pH 8.0. Chloroplasts were lysed by three cycles of freeze-thawing in liquid N_2 , microfuged, and

aliquots of the supernatants and of the original *in vitro*-translated proteins were analyzed by SDS-PAGE in 8-25% gradient Phast gels (Pharmacia), followed by direct detection of radiolabeled proteins in the wet gels (AMBIS) (Figure 20).

As estimated by SDS-PAGE, a 30 minute import converted the
5 original 32-kD BCN1 polypeptide to a doublet of 27.2 and 27.5 kD in maize and produced an additional 30-kD band in pea (Figure 20A). Neither maize or pea chloroplasts imported -pBCN1-derived polypeptides. Formation of the 27.2-kD polypeptide likely resulted from cleavage after amino acid #47-49, a likely cleavage site because it lies between S- and D-rich regions, and R residues are
10 located at -2, -7, and -8 (S. Gavel and G. Von Heijne, FEBS Lett., 261, 455 (1990)).

Time-dependence of BCN1 import was further examined (1-30 minutes) (Figure 20B) to determine if any imported polypeptides were a result of incomplete processing or proteolysis. Import was maximal after 15
15 minutes, but import time had no effect on the relative amounts of different-sized import products. Higher amounts of ATP (5 mM) stimulated import relative to lower amounts of ATP (< 0.2 mM). The data suggest that, in maize, efficient cleavage of BCN1 occurs at two closely adjacent sites and that partially processed products are also formed during BCN1 import by pea chloroplasts.
20 Therefore, nucleotides 1-833 of the maize ACCase gene encode a CTP.

EXAMPLE IX

Expression of a cDNA Clone or Genomic Clones Encoding the ACCase Gene

The cDNA and genomic clones encoding all or a portion of the
25 ACCase gene can be subcloned into a known expression system and the gene products reactive with the antibodies specific for maize ACCase can be identified using a Western blot. For example, the ACCase cDNA clones are inserted into two transformation plasmids: (i) Glb1exp which contains the embryo-specific maize Globulin1 (*Glb1*) promoter and 3' regions (Belanger et al., Genetics, 129, 863 (1991)); and (ii) pAHC17 which contains the maize ubiquitin (*Ubi-1*) constitutive promoter and first exon and intron, and the NOS 3' terminator (Christensen et al., Plant Mol. Biol., 18, 675 (1992); Toki et al., Plant Physiol., 100, 1503 (1992)). The 3' end of the A₁ cDNA has a unique *SaII* site

just 3' of the stop codon which is used to ligate into a *SaI*I site in both plasmids ahead of the construct terminator. Other cloning sites will be added as needed to the plasmids or cDNA to complete the ligation of the 5' end. The gene products can also be further characterized structurally and/or enzymatically. This will
5 ensure that the genomic and cDNA clones that encode acetyl CoA carboxylase can be screened for promoters that provide for overproduction of the native or herbicide tolerant ACCase enzyme in plants.

For example, the 2 kb EcoRI fragment from clone #15-14 can be subcloned into a plant transformation plasmid pBI121 or pBI221 downstream
10 from the 35S CaMV promoter and upstream from the nopaline 3' polyadenylation signal sequence, as described in Jefferson, Plant Molec. Biol. Repr., 5, 387-405 (1987). This plasmid can then be used to transform plant cells such as tobacco, Brassica and Arabidopsis cells using protoplast or biolistic transformation, as described by W.J. Gordon-Kamm et al., Plant Cell, 2, 603-618
15 (1990); M.E. Fromm et al., Bio/Technology, 8, 833-839 (1990); An, Methods in Enzymology, 153, 292 (1987); and D'Haflin, The Plant Cell, 4, 1495 (1992). An increase in transient expression can be detected using quantitative Western blotting with antibodies specific for the ACCase enzymes. Polyclonal antibodies to maize ACCase most likely do not substantially crossreact with ACCase from
20 dicots like tobacco or Arabidopsis.

Alternatively, the ACCase gene can be subcloned along with the 35S Came promoter into a binary Ti vector pGA482, as described in An, cited supra., which is a binary Ti vector system and can be used to transform plant cells by Agrobacterium-mediated transformation. Stably transformed plants can
25 be generated by standard methods as described in Example III, and levels of expression of ACCase genes can be determined by quantitative Western blots, as described in Harlow and Lane, Antibodies, Cold Spring Harbor Laboratories (1988). The ability to monitor expression of cloned ACCase genes will permit the identification of promoters that provide for enhanced expression of the
30 ACCase gene. The expression system can be used to screen for those promoters that enhance gene expression of the ACCase gene at least about 5 to 10-fold over the endogenous levels of ACCase produced normally in the plant cells. Because the 35S Came promoter is known as a strong promoter, it is likely this promoter

will provide for at least a 5-fold increase in the expression of ACCase over that normally produced in the plant cell.

In addition, this expression system can be used to screen antisense DNA sequences. For example, an antisense sequence can be obtained that is
5 complementary to an about 0.5 kb region of the maize ACCase cDNA that has high homology with a portion of the chicken ACCase gene and contains the sequence for the presumed transcarboxylase active site domain, as shown in FIG. 8. The antisense sequence could be subcloned into a pBI121 or pBI221 expression under the control of an inducible plant promoter, such as nitrite
10 reductase promoter (Back et al., Plant Molec. Biol., 17:9-18 (1991)). The ability of the antisense sequence to inhibit expression of the native ACCase gene can be evaluated in transformed cells, for example as described in Hamilton et al., Nature, 346:284-287 (1990).

EXAMPLE X

15 **Identification and Cloning of the Gene From Herbicide Resistant Maize Cell Lines**

Herbicide resistant maize cell lines were generated as described in Examples I, II, and IV. These herbicide resistant cell lines have been shown to produce an ACCase enzyme that is less sensitive to inhibition by sethoxydim or
20 haloxyfop. The genes encoding the herbicide resistant forms of the ACCase will be identified and cloned using standard methods as described in Sambrook et al., Guide to Molecular Cloning: A Laboratory Manual (1989). The genes encoding the herbicide resistant forms of ACCase can then be introduced into herbicide sensitive plant species by standard methods to confer herbicide resistance. For
25 example, the ACCase enzyme in the maize cell line 2167-9/2160-154 S-1 is at least 100-fold less sensitive to sethoxydim than the wild-type.

DNA from the cell line or plants will be obtained and digested with EcoRI and/or other appropriate restriction enzymes, according to standard methods. The restriction enzyme digest will be separated out by agarose gel
30 electrophoresis and probed with either the 2 kb or the 3.9 kb cDNA ACCase probe described in Example V. Fragments hybridizing to the 2 kb or 3.9 kb probe will be subcloned into a Bluescript vector and portions of the gene will be

sequenced, as described in Example V, to verify that the entire ACCase gene has been isolated.

To confirm that the clone encodes the ACCase gene, it will be subcloned into the pBI121 or pBI221 expression vector, as described in Example

- 5 VIII. The ACCase gene product expressed by the clone in either Black Mexican sweet corn cells or tobacco cells will be evaluated for reactivity with ACCase specific antibodies, by enzyme activity, and/or resistance of the enzyme activity to inhibition with sethoxydim and/or haloxyfop. It is likely that the cloned gene will encode an ACCase which is resistant to inhibition by sethoxydim and
- 10 haloxyfop. This gene can then be introduced into an herbicide-sensitive embryogenic plant cell or an embryo, including maize cells or immature embryos, to confer herbicide resistance to that plant species upon regeneration.

- The complete coding sequence encoding the herbicide resistant form of the ACCase enzyme will be cloned into a plant transformation vector
- 15 such as pBI121 or pBI221 as described in Jefferson, Plant Molec. Biol. Reporter, 5:387-405 (1987). This vector contains the 35S Came constitutive promoter, the β -glucuronidase structural gene, and the nopaline synthase 3' polyadenylation signals. The β -glucuronidase gene is replaced with a cloned ACCase gene. Optionally, the cloned ACCase gene can be combined with natural or
- 20 synthetically produced chloroplast transit peptide sequence from pea, as described in Keegstra & Olsen, Ann. Rev. Plant. Physiol./Mol. Biol., 40:471-501 (1989) and/or unique restriction sites introduced so the cloned gene can be distinguished from the endogenous maize ACCase gene. Standard methods of subcloning will be utilized as described in Sambrook et al., cited *supra*.

- 25 For transformation of maize cells, type II calli can be transformed using biolistic transformation, as described by W.J. Gordon-Kamm et al., Plant Cell, 2, 603-618 (1980); M.E. Fromm et al., Bio/Technology, 8, 833-839 (1990); and D.A. Walters et al., Plant Molecular Biology, 18, 189-200 (1992).

- Alternatively, type I embryogenic calli can be transformed using electroporation
- 30 after mechanically or enzymatically wounding calli, as described by D'Hafluin et al., The Plant Cell, 4:1495 (1992). Once the cloned gene is introduced into these cells and transformants are selected, typically by antibiotic resistance, fertile transgenic maize plants can be regenerated, as described by D'Hafluin et al. cited

supra. Fertile transgenic plants can be evaluated for herbicide tolerance, as described in Example III. It is likely that the fertile transgenic plants having and expressing a cloned ACCase gene as an ACCase resistant to sethoxydim and/or haloxyfop will exhibit herbicide tolerance as compared to the corresponding
5 untransformed plant.

EXAMPLE XI

Generation of Transgenic Plants Having an Increase in Oil Content

Once identified and cloned, the gene or genes from maize acetyl CoA carboxylase can be introduced into monocot or dicot plant species,
10 including maize, under the control of a promoter that provides for overexpression of the ACCase enzyme. The overexpression of the ACCase enzyme is likely to lead to an increase in the oil content of the plants and seeds.

Naturally occurring soybeans that have a high oil content and soybeans that have a low oil content have been identified. The acetyl CoA
15 carboxylase from both types of soybeans was isolated, as described in Example V. The activity of the enzyme was measured as a function of the time of seed development and the results are shown in Figure 11.

The results in the Figure 11 indicate that higher oil content soybean is associated with a 2-fold increase in the ACCase activity during early
20 to mid stages of development when compared with a low oil content soybean. Thus, increased expression of the ACCase gene correlates with an increase in the oil content of the seed. Total oil content of the seed was also measured at maturity (60 days). The high oil producing cell lines, Anoka and PI28C.134, have a total oil content of 21.8% and 19.9%, respectively. In contrast, the low
25 oil soybean line of M76-395, has an oil content of 13.6% oil. Thus, the increase of ACCase expression early in seed development correlates with a higher total oil content in the seed at maturity.

A gene encoding a genomic maize acetyl CoA carboxylase can be isolated, as described in Example V, and used to transform plant species by
30 protoplast or biolistic transformation. If the gene is combined with a strong promoter, such as the 35S cauliflower mosaic virus promoter, overexpression of the ACCase gene is likely. Alternatively, selecting transformed cells with multiple copies of the gene can also result in transformed cells overexpressing

the ACCase gene. The gene can be cloned into a vector such as pBI121 or pBI221, as described by Jefferson, cited *supra*. This vector contains the 35S cauliflower mosaic virus promoter, the β -glucuronidase structural gene, and the nopaline synthase 3' polyadenylation signals. The cloned ACCase gene can
5 replace the β -glucuronidase gene and then be used to transform plant cells, including maize, as described in Example VIII.

Transformed cells can be screened for overproduction of ACCase. The presence of the cloned gene can be verified by identifying the unique restriction enzyme sites incorporated into the cloned gene. ACCase levels can
10 be assessed by standard enzyme assay methods and quantitative Western blots using antibodies specific for maize ACCase. Fatty acid and lipid content in cell lines overproducing ACCase are likely to be elevated and can be assessed using standard methodologies, as described in Clark & Snyder, *JACS*, 66:1316 (1989). Transformed cell lines overproducing ACCase and having increased total oil
15 content will be used to regenerate fertile transgenic plants and seeds, as described in D'Haflin, cited *supra*.

EXAMPLE XII

Generation of Transgenic Plants Having an Increase in Plastidic ACCase Activity

20 Maize embryos are transformed with sense and antisense cDNA constructs encoding the plastidic A1 ACCase. Selected transgenic cultures and regenerated transgenic plants and progeny are subjected to detailed analyses of: ACCase transcript levels; activity of ACCase I (plastidic) and ACCase II
25 (presumably cytosolic) in various plant tissues; fatty acid synthesis; lipid accumulation in kernels (primarily embryos); and other plant traits. The culmination of these tests indicates whether plastidic ACCase activity can be modified via transformation and whether fatty acid synthesis is affected.

1. Transformation of maize embryos and plant regeneration

30 Immature maize embryos of the Hi-II genotype are transformed by particle bombardment according to slight modifications from previously described procedures (Fromm et al., *Biotech.*, 8, 833 (1990); Koziel et al., *Biotech.*, 11, 194 (1993)). This procedure has been employed in transformation studies for bombardment of approximately 15,000 embryos which were then

selected for Basta-resistant callus (*bar* selectable marker gene expression), and regenerated into plants. Transformed (Basta-resistant) plants are obtained from 1-2% of the initial embryos and, when separate plasmids are used for co-transformation, the nonselected transgene is recovered in about 50% of the

5 Basta-resistant plants.

Basta-resistant, hemizygous transformed ($T_0 = F1$) plants will be tested by PCR or Southern blots for the presence of A1 ACCase sequences unique to the transformation vector, grown to maturity in growth chambers, greenhouse or field, and self-pollinated when possible, or backcrossed to a
10 nontransformed parent. F2 or backcross progeny are grown in the greenhouse and field and tested for Basta resistance and presence of the A1 ACCase transgene to identify homozygous transgenic plants. Homozygous inbred transgenic lines are then developed.

2. Analysis of ACCase A1 transformants

15 Plants recovered from at least 100 independent transformation events (i.e., from different bombarded embryos) for both the UBI1 and GLB1 vectors are recovered. Regenerated plants are tested for the presence of the intact ACCase A1 transgene and its cosegregation with the Basta-resistance marker. Homozygous and heterozygous transgenic lines are assayed for total
20 ACCase activity in leaves for UBI1 transformants and in developing embryos (22-26 DAP) for both UBI1 and GLB1 transformants. Sethoxydim and haloxyfop inhibition are used to quickly determine the levels of herbicide-sensitive ACCase I (plastidic) and the herbicide-insensitive ACCase II activity in both leaves and embryos. Increased expression of the ACCase A1 transgene
25 contributes to plastidic ACCase activity and not to ACCase II activity. Kernel fatty acid and oil content are analyzed at maturity and their relationship with ACCase I determined by methods well known to the art.

Transformants that differ in kernel ACCase activity and/or fatty acid and oil content are then selected for more detailed analysis of embryos
30 throughout development (4-day intervals from 16 DAP to maturity). These analyses include RNase protection assays to determine total A1 + A2 transcript levels using a non-specific probe and to determine relative levels of endogenous A1 versus transgene A1 transcripts by use of antisense riboprobes spanning the

5' UTR region of the A1 transgene constructs. Western blots of total proteins separated by SDS-PAGE gels are probed with the ACCase I-specific antibody described hereinabove or with avidin and analyzed by densitometry to distinguish changes in the 227-kD ACCase I and 219-kD ACCase II isoforms.

- 5 ACCase I activity, fatty acid and lipid content are determined in embryos at each stage of development. These analyses determine whether expression of an additional gene(s) for plastidic ACCase increases ACCase activity and consequently fatty acid and oil content in maize tissues, especially in embryos.

3. Transformation with maize ACCase A1 antisense transformation vectors

- 10 Antisense transformation vectors were constructed by blunt-end ligation of nucleotides 1-833 of SEQ ID NO:5 in reverse orientation into multicloning sites of both the GLB1 and UBI1 plasmids. A sense construct with the same 833-bp cDNA sequence also was made with the GLB1 plasmid to serve as a transformation control. Insert orientations were verified by restriction
15 mapping. UBI1 antisense, GLB1 antisense, and GLB1 sense constructs were introduced into > 2100, > 2900 and > 2000 embryos, respectively, and Basta-resistant callus were selected.

- If antisense expression results in significant reduction in ACCase activity, it may not be possible to obtain viable callus or plants from the
20 constitutive UBI1 antisense transformants. Similarly, plants transformed with the embryo-specific GLB1 antisense construct may exhibit deleterious effects on embryo development. Thus, failure to obtain transgenic progeny containing the antisense ACCase gene from these transformations may indicate that ACCase activity cannot be downregulated without loss of viability.

25 4. Analysis of ACCase A1 antisense transformants

- All Basta-resistant cultures will be regenerated. The presence of the UBI1 and GLB1 antisense constructs will be determined by PCR analysis for unique transgene sequences such as the *Ubi-1* intron/ACCase A1 junction or *Glb1* 5' UTR/ACCase A1 junction, or by Southern blotting to detect unique
30 fragments. Plants and lines homozygous or heterozygous for the antisense transgene are analyzed for steady state level of the ACCase A1 antisense transcripts in appropriate tissues/organs (such as leaves, tassels, ears, embryos and endosperm for UBI1; leaves and embryos for GLB1) by using ACCase A1

sense riboprobes for hybridization on RNA blots or for RNase protection assays. Total ACCase activity (both ACCase I and ACCase II isoforms) and fatty acid and lipid content are determined for the antisense transgenic lines and for corresponding tissues from nontransformed control plants. These analyses show
5 whether ACCase A1 antisense transgenes are expressed in plants and, if so, whether expression is associated with reduced ACCase activity and altered fatty acid and lipid content in maize.

EXAMPLE XIII

Expression of Plastidic and Cytosolic ACCases during Plant Development

10 Intact embryos are isolated from developing kernels of field-grown inbred B73 at 2 to 4 day intervals between 16-42 DAP and frozen immediately in liquid nitrogen. Samples also are saved for fresh and dry weight determinations. Subsamples from each stage are analyzed for total lipid and fatty acid content. Seedling leaves are sampled along the leaf blade ranging
15 from the etiolated, meristematic basal region to the fully expanded, green tip. Leaves and other tissues (e.g., epidermis) of maize genotypes that accumulate anthocyanin pigments are also analyzed to assess whether a specific ACCase (such as a cytosolic ACCase) is more highly expressed in tissues in which malonyl-CoA also is required as a substrate for chalcone synthase in the
20 flavonoid pathway leading to anthocyanin synthesis.

Gene-specific antisense riboprobes in RNase protection assays are employed to determine A1, A2, B1 and B2 transcript levels. The corresponding sense transcripts are produced *in vitro* and used as standards to verify specificity and quantitate the sample transcript levels. Quantitation is
25 done on an AMBIS radioanalytic image system. Herbicide inhibition of total ACCase activity provides an assessment of levels of herbicide-sensitive ACCase I (plastidic) and the herbicide-insensitive ACCase II activity in these tissues. ACCase I and II isoforms are separated by ion-exchange chromatography. Total proteins are separated by SDS-PAGE and Western blots probed with avidin to
30 detect the biotinylated 227-kD ACCase I and 219-kD ACCase II isoforms or probed with ACCase I-specific antibodies.

EXAMPLE XIV

Identification and Isolation of an Oat Gene Associated with Groat Oil**Content**

Cultivated oat, *Avena sativa* L. ($2n=6x=42$), is grown mostly in temperate regions. In many parts of the world, oat grain is produced for many purposes such as for animal feed, human food, and cosmetic and pharmaceutical additives (Webster, 1986). Seventy-eight percent of the world oat production is used for livestock feed (Schricket, 1986). Among the cereals, oat groats have the highest lipid concentration (Morrison, 1978), but due to their high fiber content oat is lower in energy value than many other cereals when used for livestock feed. Oat is also unique among the cereals because more than 50% of total seed lipid is deposited in the starchy endosperm rather than the embryo of developing caryopses (Peterson and Wood, 1997; Youngs, 1986). An increase in oil content is desirable for making oat a higher-energy feed grain. On the other hand, a decrease in oil content would make oat a more attractive commodity for use in many current human diets where oat is desired for the type and concentration of its fiber (Ripsin et al., 1992).

A polygenic pattern of inheritance has been indicated for groat oil content with heritability estimates ranging from 63 to 93% (Schipper and Frey, 1991a) with primarily additive gene action (Thro and Frey, 1985). The average groat oil concentration in contemporary cultures is 6 to 7% but variation from 2 to 16% oil occurs in diverse germplasm (e.g., Kanota has about 11%, Ogle about 9% and Marion about 8% of oil (tryglycerides plus polar lipids) content, per dry weight). Genotype by environment interaction for this trait is small (Gullard, 1986; Branson and Frey, 1989). After five cycles of a recurrent selection program, Schipper and Frey (1991b) were able to produce oat lines with groat oil content greater than 16%. This response to selection is similar to the classical Illinois long-term selection experiments for oil content in maize (*Zea mays* L.) that resulted in dramatic progress toward altering the oil content in the original variety (Dudley et al., 1974). These studies demonstrate that, except when physiological limits are encountered, long-term selections for a quantitative, highly heritable and environmentally stable trait can prove successful. Analyses of 5987 oat samples representing a wide range of *Avena* germplasm showed a range of 2 to 12% for free non-polar lipids, with polar lipids contributing an

additional 2 to 3% (Youngs et al., 1982). Combinations of alleles for changing the groat oil content beyond this range may exist because this is mostly an unselected trait that exhibits large phenotypic diversity. Therefore, genetic dissection of this trait could expedite a program aimed at modifying the oil level
5 of oat.

The recent explosion in development of restriction fragment length polymorphism (RFLP) linkage maps and their use in identifying genomic regions which affect quantitatively inherited traits is well documented (Phillips and Vasil, 1994). Molecular genetic studies relating to seed oil content have
10 been reported for major oil-producing crop plants such as rape seed (*Brassica rapa* spp. *oleifera*; Tanhuanpaa et al., 1995a and 1995b), soybean (*Glycine max* L., Diers et al., 1992; Mansur et al., 1993; Brummer et al., 1997) and maize (Goldman et al., 1994; Alrefai et al., 1995). These studies indicate that genomic regions can be identified which significantly influence seed oil content. In the
15 maize long-term Illinois selection material, two populations, the high oil by low oil population (Alrefai et al., 1995) and the high protein by low protein population (Goldman et al., 1994) were analyzed to identify genomic regions influencing kernel oil content. In both populations a region on chromosome 6 was identified near the linoleic acid1 locus. In the one case, this region explains
20 63% of the phenotypic variation in the ratio of oleic to linoleic (18:1 to 18:2) fatty acids and also influenced the total kernel oil concentration (Alrefai et al., 1995). To date, oat plants with an altered fatty acid composition as a result of a qualitative genetic mutation have not been identified, possibly due to the hexaploid nature of the genome.

25 In monocots, graminicides cause plant cell death by inhibiting ACCase and cessation of fatty acid biosynthesis (Burton et al., 1989). Maize mutants have been isolated that grow in the presence of normally lethal concentrations of ACCase-inhibiting herbicides, exhibit up to 20-fold increase in wild-type ACCase activity, and synthesize fatty acids in the presence of
30 herbicide (Parker et al., 1990). Other results point to the role of ACCase in regulating seed oil deposition. A herbicide-insensitive ACCase mutation, Acc1, exhibits increased ACCase activity in corn embryos paralleling the reporting time course of oil deposition in this tissue (Somers et al., 1993). In developing

seeds of castor bean and rape, measured activity of ACCase correlates well with lipid accumulation providing additional evidence as to the role of this enzyme in fatty acid biosynthesis (Simcox et al., 1979; Turnham and Northcote, 1983).

Moreover, expression of *Arabidopsis* cytosolic ACCase in *B. napus* plastids

5 alters fatty acid composition, with the largest effect being an increase in oleic acid, and increasing total seed oil content by about 5% (Roesler et al., 1997).

This increase in oleic acid is consistent with studies in maize indicating a positive correlation between percentage of oleic acid and total seed oil content (Alrefai et al., 1995).

10 Materials and Methods

Genetic Material

Two oat F_{2:6}-derived recombinant inbred populations developed by single seed descent (SSD) were used in this study: a population of 137 recombinant inbred lines (RILs) developed from the cross of a facultative winter-

15 type cultivar, Kanota, by a spring cultivar, Ogle, (KO) and a second population
of 139 RILs from the cross of Kanota by another spring cultivar, Marion, (KM).

A portion (71 RILs) of the first population had been employed in previously reported oat mapping and QTL studies (O'Donoughue et al., 1995;

Siripoonwiwat et al., 1996; Holland et al., 1997). The second population was

20 chosen on the basis of having a common parent, Kanota, to provide a degree of
“biological replication”. These materials were grown in 1992-1995 in a
randomized complete block design at the following locations;

Kanota X Ogle (KO): Aberdeen, Idaho, 1992 and 1993; Ithaca, New York, 1993; and St. Paul and Rosemount, Minnesota, 1994.

Kanota X Marion (KM): St. Paul and Rosemount, Minnesota, 1994; and St. Paul, Minnesota, 1995.

At Aberdeen, Idaho, the materials were grown in 3 replicates of four-row plots of 2.5 m in length, 30 cm between rows with the center 2 rows harvested. At

30 Ithaca, New York, the materials were grown in 3 replicates of six-row plots of 4
m length and 18 cm between rows. At St. Paul and Rosemount, Minnesota, the
materials were grown in 4 replicates of hill plots, with 30 seeds planted per hill

on 30 cm grid spacing. The exact parents used to generate the populations were included as single or multiple entries in each replicate.

Oil Content Analysis

To reduce the number of samples needed for analysis, equal
5 amounts of sample from each replicate at a given location/year were bulked to represent a single balanced sampling per line. This resulted in five measurements for each RIL in the Kanota X Ogle population and three for the Kanota X Marion population.

Groat oil content was measured by a Bran+Luebbe InfraAlyzer
10 500 on dehulled ground samples. The Near Infrared (NIR) calibration was based on total fat as determined by acid hydrolysis. The calibration set contained 348 oat samples (representing commercial and elite breeding lines from North American oat breeding programs) with fat contents that ranged from 4.95 to 14.01%. The calibration equation had a correlation coefficient of 0.985, a
15 Standard Error of Estimate of 0.299, and a F-value of 11500. Acid hydrolysis extracts the total lipid content of the groat. Thus, the reported values represent the sum of the neutral lipids (triglycerides) plus the polar lipid fractions (Youngs et al., 1982). Thus, these values are higher than those obtained from ether extracts which only contain the neutral lipid fraction.

20 Oat ACCase Clones or PCR Products

An oat groat cDNA clone was isolated based on its hybridization to a maize cDNA fragment which included portions of the less conserved mid-peptide sequence and the transcarboxylase domain (Shorrosh et al., 1994). Sequence identity indicated that the oat cDNA (3031 bp) corresponded to maize
25 ACCase nt 3124-6169 (Egli et al., 1995). The 1010-aa predicted polypeptide was nearly identical to portions of wheat (89%) and maize (84%) plastidic multifunctional ACCases; it had lower identity to cytosolic ACCases, including wheat (67.5%) and alfalfa (64.6%) (Gornicki et al., 1994; Shorrosh et al., 1994).

An amplified 'Prairie' oat developing groat cDNA library (R.
30 Skadsen, USDA, Madison, WI) cloned into the EcoRI site of λ ZAPII (Stratagene) was screened with a [32 P-dCTP]-labelled maize ACCase cDNA probe (Ausabel et al., 1997) derived from an EcoRI digest of clone 18-5 (nt 2020-5927; Egli et al., 1995). Plaques from eleven partially purified positives

were eluted in SM medium (Sambrook et al., 1989) and excised *in vivo* to obtain pBlueScript plasmids carrying oat cDNA inserts, according to the manufacturer's instructions (Stratagene Corp.). Five colonies per transformation reaction were screened (as colony lifts) with an AflIII-BbrPI fragment of maize ACCase cDNA (4281-5797; Egli et al., 1995). One clone, oat-91, hybridized to the probe. Oat-91 was sequenced using the ABI Prizm system (Perkin-Elmer Corp.) at the Advanced Genetic Analysis Center (Univ. of Minnesota, St. Paul, MN). Sequence identity indicated the oat-91 EcoRI insert corresponded to maize ACCase nt 3124-6169 (Egli et al., 1995). Oat ACCase cDNA sequence was compared to other plant ACCases with the BESTFIT program (BESTFIT, Wisconsin Package V. 9.1, Genetics Computer Group (GCG), Madison, WI).

For PCR, Pileup alignments of maize, wheat, barley, and rice ACCase coding sequence (GCG, Madison, WI; 12, 18, F, R) were used to identify conserved regions (i) for design of new primers or (ii) regions of identity to available maize ACCase oligos, to be used for PCR amplification of oat ACCase. Two maize primers, PCR2 (5'-ggcagagcaa agcttgagg; SEQ ID NO:25) and PCR1 (5'-tccaccagag aagcctctcc; SEQ ID NO:26) appeared like to amplify oat cDNA based on their identity to other monocots and previous use to amplify soybean ACCase. These primers are located within the transcarboxylase domain, surrounding the acetyl-CoA binding site of ACCase. An amplified 'Prairie' oat seedling leaf cDNA library cloned into the EcoRI site of IZAPII (Skadsen, University of Wisconsin) was used as a source of cDNA. One I of the oat cDNA library was amplified with maize primers (PCR2/PCR1), using the Expand High Fidelity PCR system (BMB) in a GeneAmp PCR System 9600 thermal cycler (Perkin-Elmer) after denaturing phage heads at 94°C for 10 minutes. The single, 222-bp product was purified by Wizard PCR Plus (Promega Corp.) and sequenced in an ABI377 (Perkin-Elmer) using PCR1 as primer. The resulting sequence appeared to be from oat since it had high identity to maize plastidic ACCase (89%). Additional 5' cDNA was obtained by (i) PCR of the oat cDNA library with T3/PCR1 primers to generate a complex band mixture and (ii) reamplification of 0.1 I products with maize ACCase primers 1324+ (5'-caatcacgtcagactg; SEQ ID NO:27)/PCR1 to obtain a 1.9-kb product with the same 3' sequence as the PCR2/PCR1 product.

RFLP and Statistical Analysis

Briefly, two statistical procedures were used to determine the location, effect and genotype by environment interaction (GxE) for significant QTLs. The first procedure was single factor ANOVA (ANOVA and GLM procedures; SAS Institute, 1990) with the added criteria for consistency in association of genomic regions with a trait over environments and linkage distances (Kianian et al., 1998). If the association were significant when considering the average of all the environments, the association was also significant over the majority of the environments (3 of 5 in KO and 2 of 3 in KM), and closely linked markers within that region, if available, also showed significant associations. A “backward elimination” process as described by Kennard and Havey (1995) was used to construct a multi-locus model containing the most important markers identified by the ANOVA procedure. Marker loci were excluded one at a time from this model, based on the criterion of least significant ($P < 0.05$) Type-III sum of squares (Kennard et al., 1994). In addition, MQTL software (Tinker and Mather, 1995a & b) was used to strengthen and validate the ANOVA results. This software performs Simple Interval Mapping (SIM) and Simplified Composite Interval Mapping (sCIM). It also has the advantages of searching for one QTL while simultaneously accounting for the effect of others, providing separate analyses for each environment, and calculation of a threshold to control the type-I error rate. One thousand permutations were performed to estimate the threshold for a type-I error rate below 5%. Significant QTL regions were determined as described by Tinker et al., 1996. These regions were grouped in a multi-locus linear model to estimate the overall phenotypic variance explained by the model. However, additional criteria were used to include markers in the final MQTL model, which gave 10 cM spacing between QTLs, and minimized the number of regions excluding those markers explaining less than 5% of the phenotypic variance, while maintaining most of the overall R^2 .

Results

Trait Data

Groat oil concentration for both populations showed a relatively continuous distribution with many of the individuals having values within one standard deviation of the parents (Table VI). Among the three parents, Kanota showed the highest concentration of groat oil content followed by Marion and then Ogle. The groat oil content value for the Kanota line derived from the parent used in generating the KM population was higher than that used for the KO population (Table VI). This is not unexpected: Kanota is a heterogeneous population that shows variation for other traits, but individual parent lines were used in population development and were maintained as controls for each population. Large numbers of individuals having values outside the range of parents were observed for both populations (Table VI). Transgressive segregants tended more toward the lower groat oil concentration than the high. The KO population contained individuals with the highest and lowest groat oil concentrations, possibly due to a larger difference in parental values (3% between Kanota and Ogle, 2% between Kanota and Marion). Overall, the KM population contained more transgressive segregants than the KO population (Table VI).

Table VI. Groat oil concentration and percent transgressive segregants in two recombinant inbred oat populations

Population (P1 × P2)	Location	Parents			Population			% Transgressive segregants			
		P1	P2	Mean	SD	Min.	Max.	X<P2	X>P1	X<P2-SD	X>P1+SD
Kanota X Ogle	Aberdeen 92	11.5	8.3	9.7	1.3	7.5	14.0	13	8	0	2
	Aberdeen 93	10.5	8.4	10.1	1.6	7.2	15.4	18	35	0	8
	Ithaca 93	10.9	7.1	9.0	1.4	6.5	13.6	8	8	0	2
	St. Paul 94	10.9	7.7	9.2	1.4	6.6	13.4	13	10	0	2
	Rosemount 94	11.1	8.4	9.0	1.3	6.7	12.6	30	5	5	1
	Average	11.0	8.0	9.4	1.4	6.9	13.8	16	13	1	3
Kanota X Marion	St. Paul 94	12.9	10.2	10.2	1.0	8.0	13.3	47	1	12	0
	Rosemount 94	12.2 ^a	9.2	10.4	1.2	7.5	14.2	10	7	1	3
	St. Paul 95	10.0	8.4	8.8	0.7	7.1	10.8	26	2	4	1
	Average	11.5	9.3	9.8	1.0	7.5	12.8	28	3	6	1

^a Adjusted average value for the parents calculated based on the mean of population at that location divided by the overall population mean times the average parental values was used in this calculation (i.e., (10.4/9.8)11.5).

SD = Standard Deviation

Table VII. Phenotypic correlations for groat oil concentration between environments*

Population	Location	Aberdeen 92	Aberdeen 93	Ithaca 93	St. Paul 94
Kanota X Ogle	Aberdeen 93	0.94			
	Ithaca 93	0.94	0.97		
	St. Paul 94	0.90	0.91	0.92	
	Rosemount 94	0.91	0.95	0.96	0.94
Kanota X Marion		St. Paul 94	Rosemount 94		
	Rosemount 94	0.87			
	St. Paul 95	0.80	0.71		

* All values are considered significant at $P < 0.0001$.

The correlation among the different environments for oil content was high (Table VII). Measurements for the KO population had an average R of 0.93 with the lowest value of 0.90 between material grown at Aberdeen, ID 1992 and St. Paul, MN 1994 ($P < 0.0001$). Values for the KM population were less well correlated (average R of 0.79) with the lowest correlation between the samples grown in Rosemount, MN 1994 and St. Paul, MN 1995 ($P < 0.0001$). The difference in the size of the population correlations could be due to a smaller range of values for the KM population (5%) as opposed to the KO population (7%), or fewer measurements (3 vs. 5 environments), or both.

10 Kanota X Ogle Population
RFLP Data

A hexaploid oat RFLP map that contains 561 loci has been generated from 71 RILs in the KO population (O'Donoghue et al., 1995). Initial quantitative analysis was performed using this subset of the KO population, the available trait data, and 360 of the 561 loci that are co-dominantly segregating and evenly distributed across the oat genome. To take advantage of the entire population and to add power to the analysis, the remaining 66 RILs in this population were also mapped. To date, 153 loci have been mapped with these individuals. Detection of QTL with either the subset (360 markers and 71 individuals) or the entire data set (153 markers and 137 individuals) of the KO population did not make a major difference, except in two situations. In both cases, QTLs were detected in the subset, but were not detected in the entire population due to the absence of RFLP data for the remaining 66 RILs. Results of the entire population are presented herein except for the aforementioned differences.

An RFLP map containing all 137 individuals and the 153 mapped loci was generated and was in accord with the published map (O'Donoghue et al., 1995) except for linkage group 3. An inter-varietal translocation is believed to involve this linkage group. In the final analysis, the published map was used for this region. Additional information from aneuploid analysis (Kianian et al., 1997) was also taken into account when two linkage groups were assigned to the same chromosome. Thus, linkage groups 5 and 7 assigned to chromosome 5C, linkage groups 6 and 20 assigned to chromosome 16, and linkage groups 4 and

12 assigned to chromosome 21, were analyzed as three distinct units. In order to have a complete coverage of the genome for SIM and sCIM, 344 mapped loci (191 loci scored on only 71 RILs) were used in the MQTL analysis.

Quantitative Analysis

- 5 Analysis of variance using the criteria mentioned revealed eight linkage groups and one unlinked marker significantly associated with groat oil content ($P < 0.05$ for individual environment and $P < 0.0001$ using means across environments). These regions are on linkage groups 6 and 7 (Table VIII). Two additional genomic segments on linkage group 37 and an unlinked locus
- 10 *Coleoptal* were significant in tests of the first 71 individuals of the mapping population but could not be verified on the whole population due to a lack of RFLP data for the entire set. The positive alleles (those that cause an increase in groat oil content) for all of the genomic regions influencing this trait were from the Kanota parent (Table 3). Overall, the alleles from Kanota contributed 2.8%
- 15 (4.1% if linkage group 37 and *Coleoptal* loci were included) to the groat oil content. Regions on linkage groups 6 and 11 are considered highly significant and remained in the full model after the backward elimination process (Table 4). However, the two additional loci from linkage group 37 and *Coleoptal* were also
- 20 being analyzed. However, the two additional loci from linkage groups 37 and *Coleoptal* were also included in the model if only the first 71 individuals in this population were being analyzed. Locus Xcdo665B on linkage group 11 accounted for 36% to 48% (depending on the environment) of the phenotypic variance for this trait ($P < 0.0001$). For these linked loci, a substitution of
- 25 Kanota alleles for their Ogle counterparts caused an average increase of 2% in groat oil content. Substitution of Kanota alleles for Ogle alleles at the locus *Xcdol414B* (or *Xumn51B*) on linkage group 37 caused an average increase of 1% in groat oil content and was the second most significant locus considering only the first 71 individuals in this population.

Table VIII. Genomic regions significantly influencing groat oil content in oil identified by single factor analysis of variance

Kanota X Ogle					Kanota X Marion				
Linkage Group	Locus ^a	Source	R ²	% Oil ^b	Linkage Group	Locus ^a	Source	R ²	% Oil ^b
6	<i>Xcdo1357C</i>	Kanota	0.10	0.8	3	<i>Xbcd1562A</i>	Kanota	0.05	0.4
11	<i>Xcdo665B</i>	Kanota	0.38	2.0	11	<i>Xcdo665B</i>	Kanota	0.18	0.7
37 ^c	<i>Xcdo1414B</i>	Kanota	0.10	1.0	22	<i>Xumn5004</i>	Kanota	0.05	0.3
Unlinked ^c	<i>Coleoptal</i>	Kanota	0.13	0.3	5X ^d	<i>Xcdo1199</i> C	Kanota	0.11	0.7
Total	2.8 (4.1 ^b)				2.1				

^a Locus with the highest R².^b Substitution of the source alleles will cause on average (based on average of values from each of five environments in KO and three in KM) this change in groat oil concentration. P < 0.0001 for the mean across all environments and P < 0.05 for each individual environment.^c Significant region identified only in the first 71 individuals of this population.^d More markers are needed to accurately place this linkage group on the hexaploid oat RFLP map.

Table IX. Multi-locus models for groat oil concentration in oat

Population	ANOVA Model ^a	MQTL Model ^b
Kanota X Ogle	<i>Xcdo1357C</i> (6, K)	<i>Xcdo665B</i> (11, K) ^c
	<i>Kcdo665B</i> (11, K) ^c	<i>Xcdo1414B</i> (37, K)
	<i>Xcdo1414B</i> (37, K) ^d	<i>Coleoptal</i> (Unlinked, K)
	<i>Coleoptal</i> (Unlinked, K) ^d	
Total R ²	0.45 (0.67) ^e	0.47
Kanota X Marion	<i>Xcdo665B</i> (11, K) ^c	<i>Xcdo665B</i> (11, K) ^c
	<i>Xumn5004</i> (22, K)	<i>Xumn5004</i> (22, K)
	<i>Xcdo1199C</i> (5X, K)	<i>Xcdo1199C</i> (5X, K)
Total R ²	0.28	0.20

^a The linkage group and parent contributing the positive allele are depicted in parentheses (i.e., linkage group 11 with Kanota allele being positive (11, K)). The models are based on five environments in KO and three in KM.

^b The VM/VP (variance explained by the QTL(s) over the phenotypic variance) is presented as the R² for MQTL.

^c *Xaccase4* is closely linked to *Xcdo665B* and gives the same relative effects as *Xcdo665B*.

^d These additional loci fit the model when the analysis included only the first 71 individuals in this population.

^e Total R² with markers marked d is presented in parentheses.

Quantitative analysis using the MQTL software gave a slightly different result. Like the ANOVA results, a highly significant locus ($P < 0.0001$), Xcdo665B on linkage group 11, was identified at a test statistic (TS) value of 290.2 that is roughly equal to a LOD score of 64 ($0.22 \times TS$). This locus overshadowed any other significant region and accounted for more than 37% of the phenotypic variance [Variance QTL Main effect (VM)/Variance Phenotypic (VP) = 38%, Variance QTL $\times$ environment Interaction and VM (VI)/VP = 38% and Variance Genetic from background markers (VG)/VP = 7%] and a change of 2% in groat oil content (Figure 23). However, secondary loci that showed significance below the threshold value (threshold TS = 58.2 after 1000 permutations) were inferred when either SIM or sCIM gave evidence for QTLs. Xumn51B on linkage group 37 and the Coleoptal locus, which is unlinked, were significant at test statistic values of 49.9 and 44.8 respectively. The three loci together accounted for 47% of the phenotypic variance and a change of 3.3% in groat oil content (VM/VP = 47%, VI/VP = 48% and VG/VP = 7%). For all three loci the Kanota parent contributed the positive allele. These loci showed a remarkable consistency over the five environments. Thus, there were no significant peaks inferred from the QTL $\times$ E interaction scan (Figure 23).

20 Kanota X Marion Population RFLP Data

The results from an initial quantitative analysis of the KO population (first 71 RILs) for groat physical and chemical traits were used to determine markers for mapping the KM population. The markers mapped were not chosen for their association with groat oil content in the KO subset population, but for their association with other groat chemical and physical characteristics (Kianian et al., 1998). Since the level of polymorphism in hexaploid oat is relatively high, most probes detected sequences mapping to more than one location. A total of 60 loci were mapped with the 139 RILs in the KM population. These markers mapped to 20 of the 38 linkage groups that had been identified in the KO population (linkage groups 1, 2, 4, 5, 8, 9, 18, 19, 21, 24, 25, 26, 30, 33, 34, 35, 36, 37, 38 were not covered). The linkage groups not covered were, in general, the smaller ones consisting of only two to a few

markers in the KO map. The data from the KM population generated an additional linkage group of two loci, and 13 markers remained unlinked. Except for linkage group 37 and *Coleoptal* locus, this KM map covered all regions with significant associations with groat oil concentration in the KO population.

5 Quantitative Analysis

Single factor ANOVA revealed four linkage groups significantly associated with groat oil content in the KM population. These were groups 3, 11, 22, and 5X ($P < 0.05$ for individual environment and $P < 0.0001$ using means across environments). As in KO, all of the genomic regions influencing this trait were from the Kanota parent. Overall, the Kanota genomic regions contributed 2.1% to the groat oil content. Regions on linkage groups 11, 22, and 5X (group 5X has been tentatively assigned to linkage group 5 but more markers are needed to accurately place this linkage group on the hexaploid oat RFLP map) are considered highly significant and remained in the full model after the backward elimination process. As in the KO population, locus *Xcdo665B* on linkage group 11 was highly significant for its effect on groat oil content ($P < 0.0001$), accounting for 16% to 20% (depending on the environment) of the phenotypic variance for this trait. Overall, the three significant loci explained 15% of the phenotypic variance and together accounted for a total difference of 1.7% in groat oil content.

Analysis by SIM or sCIM gave the same result as ANOVA. Linkage groups 11, 22, and 5X were significant in the KM population. Locus *Xcdo665B* on linkage group 11 was significant at a test statistic value of 70.5 (LOD = 15.5). This locus was not as dominant in the KM population as it was in the KO population. This locus alone explained 18% of the phenotypic variance (VM/VP = 18%, VI/VP = 19% and VG/VP = 3%) and accounted for 0.9% change in groat oil content. The other significant locus above the threshold value was *Xcdo1199C* on linkage group 5X (TS = 33.1) accounting for a change of 0.2% in groat oil concentration. A secondary locus inferred was *Xcdo506B* on linkage group 22, which accounted for 0.4% change in oil content. These three loci together accounted for 20% of the phenotypic variance and 1.5% in groat oil content (VM/VP = 20%, VI/VP = 22% and VG/VP = 3%). As expected from the trait data, these loci showed a varied effect over the three environments and one

interaction on linkage group 11 is notable (Figure 23). The QTL $\times$ E interaction peak on linkage group 11 was at TS = 6.6, slightly below the significant threshold, and was due to a range of 0.5% in groat oil content explained [the values for the groat oil content were 0.6% (St. Paul, Minnesota, 1995) and 1.1% (Rosemount, Minnesota, 1994)].

ACCCase and its Relationship to Groat Oil Content QTLs

An oat groat cDNA was isolated based on its hybridization to a maize cDNA fragment which included portions of the less conserved mid-peptide sequence and the transcarboxylase domain (Shorrosh et al., 1994). The 3031-bp oat cDNA was nearly identical to wheat (90%) and maize (84%) genes encoding plastidic ACCCase (Egli et al., 1995; Gornick et al., 1997). A 424 nucleotide sequence from the 3' end of the insert was translated and compared to other plant ACCCase peptides. The oat peptide sequence was 93% identical to plastidic isozymes from maize and wheat (Egli et al., 1995; Gornicki et al., 1997), but only 78 to 83% identical to cytosolic ACCCase from maize, wheat (Gornicki et al., 1994), or alfalfa (Shorrosh et al., 1994).

The 3031-bp oat ACCCase cDNA clone identified a single polymorphic locus in the KO population (*AccaseA*) and two polymorphic loci in the KM population (*AccaseA* and *AccaseB*). The *AccaseA* locus maps 2.3 cM from *Xcdo665B* and 3.7 cM from *Xisu2287* on linkage group 11. The *AccaseB* locus maps 11 cM from *Xcdo665D* and 7.3 cM from *Xbcd1729B*. In the KO population *AccaseA* explains on average 37% of the phenotypic variance and in KM 18% of the phenotypic variance for groat oil content (for both cases $P < 0.0001$ using means across all environments). The individuals in both populations can be divided into two groups based on their genotype for this locus and groat oil content using the mean across all environments (Figure 3). These two classes differ in their oil content by an average of 1.67% in KO and 0.78% in KM. Results of QTL analysis for the *AccaseA* locus are the same as that reported for *Xcdo665B* (Table 4).

30

Table X. Peptide sequence identity of oat and other multifunctional ACCCases. Partial, internal sequences are 1010 amino acids unless noted otherwise. Presence of an N-terminal plastid transit peptide is noted, if known.

	ACCCase Gene source	GenBank #	Transit Peptide?	% identity
	<i>Avena sativa</i>	AF072737	?	100
	<i>Triticum aestivum</i>	AF029895	yes	89.2
	<i>Zea mays</i>	U19183	yes	84.1
5	<i>Triticum aestivum</i>	U10187	no	67.5
	<i>Zea mays</i>	Egli, unpubl.	?	78.6 #
	<i>Hordeum vulgare</i>	X99102	?	71.9 §
	<i>Arabidopsis thaliana</i>	L27074	no	63.7
	<i>Brassica napus</i>	X77576	yes	63.9

10 # based on 145 aa at the C-terminus of oat-91
 § based on 498 aa at the C-terminus of oat-91
 ? partial cDNA, unknown N-terminus

15

Discussion

The ACCase cDNA clone from developing oat seeds hybridizes to a genomic region that is linked to a QTL locus on linkage group 11 having a major influence on the total lipid content of oat groats. The influence of this locus was detected in various environments and two genetic backgrounds, thus confirming the value of biological as well as environmental replication. These results provide strong support for the hypothesis that ACCase has a major role in determining groat oil content in crop plants. Either allelic variation in ACCase activity controls oil content in oat groats, or a gene(s) closely linked to ACCase is responsible. In other studies of oat directed at transferring a major crown rust resistance locus on linkage group 11 (Bush et al., 1994), lines with the introduced rust resistance gene often had altered groat oil content. These results further indicate the likely presence on linkage group 11 of a genetic locus with a critical role in the fatty acid biosynthetic pathway in oat. Mutations, which cause a gradual but demonstrable major effect on a biochemical pathway, are believed to be the most common type in polyploids (Sears, 1972).

Identification of a major quantitative trait locus influencing the biochemical composition of grain is not unusual. Studies of fatty acid, starch, and protein concentrations and composition provide good examples (Goldman et al., 1993 and 1994; Alrefai et al., 1995; Hu et al., 1995). In many of these cases,

the significant region is in close proximity to a known genetic locus with a demonstrated effect on the trait of interest. For instance, in maize, a major genomic region for fatty acid concentration and composition has been identified near the *linoleic acid-1* locus, and one influencing protein and starch

- 5 concentration was identified near the *Shrunken-2* locus (Goldman et al., 1993 and 1994; Alrefai et al., 1995). These and numerous other reports bolster the hypothesis that allelic variants at known genetic loci may be responsible for quantitative effects (Robertson, 1985).

- Reliable identification of genomic regions with an influence on a
10 particular quantitative trait has always been difficult. In a number of studies conducted on a given quantitatively inherited trait in various populations and environments, the most significant regions tended to be in common (Goldman et al., 1994; Alrefai et al., 1995). However, the accuracy of identifying a quantitative trait locus will depend, in part on the degree to which it influences
15 the trait in a given population. This is demonstrated in our study by the locus on linkage group 11 which has similar but varied effects in different populations. Marion has a higher concentration of groat oil than Ogle and more loci with positive influence on this trait. The locus on linkage group 11 increased the groat oil content to a lesser degree in the KM population than in the KO
20 population. Correspondingly, the relative proportion of variance due to the environmental influence on this locus was much higher in the KM population than in the KO population. This example illustrates the power of biological replication (i.e., having populations with a common parent) in identifying quantitative trait loci.

- 25 Identification of the most significant locus does not preclude the identification of other less important loci due to the influence of genetic background and environment on the overall effect. The consistency in identification of significant regions by various quantitative analysis methods in the KM population is a good example. The regions on linkage groups 5, 11 and
30 22 and Xcdo395C locus are considered to make important contributions to the groat oil content in this population. The Xcdo1199C locus located near the Xacor242 locus on linkage group 5 of the published hexaploid oat RFLP map (O'Donoghue et al., 1995) has as great an effect as Xcdo665B-XaccaseA region

on linkage group 11 in this population. This locus was not detected in the KO population presumably because the Ogle allele had a similar influence as Kanota and in the KM population the Marion parent was carrying the allele with positive influence on the groat oil content. The other two significant loci in the KM
5 population have a relatively minor effect on this trait and would be more difficult to detect in a population segregating for regions with a larger influence on oil such as in the KO population.

Results from other QTL studies can help identify possible pleiotropic effects of other traits such as heading date and vernalization response
10 on groat oil concentration. A region on linkage group 3 with alleles being contributed by Kanota is implicated as having a major influence on heading date and possibly yield, thus reducing its importance as having a direct effect on groat oil concentration (Siripoonwiwat et al., 1996; Holland et al., 1997). The cultivar Ogle was found to have loci on linkage groups 22 and 37 that influence plant
15 height in the KO population (Siripoonwiwat et al., 1996). Loci on linkage group 11 is believed to have an influence on test weight and other groat physical characteristics, thus providing additional evidence of their role in oat groat morphophysiology (Siripoonwiwat et al., 1996; Cakir et al., 1996). These and other studies of quantitative traits in oat are helping identify genomic regions
20 that influence the growth, development and physiology of grain, the primary product. Thus, genetic and molecular manipulation of the oat genome to improve grain quality and quantity can be directed at critical segments.

Cultivated oat is a hexaploid composed of three ancestral genomes designated A, C, and D (Rajhathy and Thomas, 1974). Recent studies
25 have indicated that the organization of hexaploid oat genomes into discrete homoeologous groupings such as that of hexaploid wheat is difficult (Rooney et al., 1994; O'Donoghue et al., 1995; Kianian et al., 1997). However, C-genome chromosomes, due to their distinctiveness, have been used as a basis to identify homoeologous segments among the other chromosomes (Kianian et al., 1997).
30 Thus, it is believed that a type of segmental homology instead of whole chromosome homology best describes the genome organization in hexaploid oat (Kianian et al., 1997). Association of homoeologous genomic segments with a quantitative trait further substantiates that relationship.

The region marked by the probe CDO1414 can identify both the significant region on linkage group 11 and the one on linkage group 37 indicating possible homoeology of these regions. The same is true for probes CDO1199 for linkage groups 6 and 5X and CDO708 for linkage groups 6 and 22. Thus, the association of groat oil content with RFLP markers in oat is explained by at least three sets of homoeologous genomic segments.

In hexaploid wheat plastidic ACCase maps to the short arm of homoeologous group 2 near the telomere (Gornicki et al., 1997). Plastidic ACCase genes in wheat have been shown to be transcriptionally active in seedling leaves but their effect on seed lipid content are unknown (Gornicki et al., 1997). ACCase in maize has been mapped to short arm of chromosome 2 near the centromere and long arm of chromosome 10 (Egli et al., 1995; Van Dee, 1994). A QTL with moderate effect on oil concentration in progeny of Illinois High and Low oil maize lines was located near the ACCase locus on maize chromosome 2 (Berke and Rocheford, 1995). Additionally, there are numerous *defective kernel (dek)*, *reduced endosperm (ren)*, and *Endosperm factor (Ef)* loci influencing grain morphophysiology located on chromosomes 2 and 10 of maize near these synthetic segments (MaizeDB). In maize, oil is stored in embryos and embryo size is associated with oil content of whole seeds (Dudley, 1974). Unlike maize, oat grains contain significant amounts of lipid in the endosperm (Peterson and Wood, 1997), and thus QTLs affecting pathway flux could be more important than those altering seed anatomy. The regions in wheat and maize agree well in terms of conserved syntony and correspond to chromosome 4 of rice, linkage group C of diploid oat and linkage group 11 of hexaploid oat map (Ahn and Tanksley, 1993; Van Deynze et al., 1995). This comparative evidence from other grass species and homoeologous evidence from oat further help reinforce the validity of the association of these genomic regions with groat oil content and composition in oat.

30 Summary

Oat groats are unique among cereals for the high level and the embryo-plus-endosperm localization of lipids. Genetic manipulation of groat quality traits such as oil is desired for optimizing the value of oat in human and

livestock diets. A locus having a major effect on oil content in oat (*Avena sativa* L.) groats was located on linkage group 11 by single factor analysis of variance, simple interval mapping and simplified composite interval mapping. A partial oat acetyl-CoA carboxylase (ACCase) cDNA clone with >84% identity to genes
5 for plastidic ACCase of maize and wheat, which catalyzes the first committed step in *de novo* fatty acid synthesis, identified a polymorphism linked to this major locus. Similar QTL and ACCase locus placements were obtained with two recombinant inbred populations having one parent in common (Kanota X Ogle and Kanota X Marion) and containing 137 and 139 individual lines,
10 respectively. By having a common parent, these populations provide biological replication of the results in that significant genomic regions should be evident in analysis of multiple cross combinations.

Similar QTL and ACCase placements were obtained with two recombinant inbred populations having one parent in common (Kanota X Ogle
15 and Kanota X Marion) and containing 137 and 139 individual lines, respectively. By having a common parent, these populations provide biological replication of the results in that significant genomic regions should be evident in analysis of multiple cross combinations. These populations were mapped with RFLP loci (150 and 60) distributed over the genome and were grown over diverse locations
20 (4 and 2) and years (3 and 2) to provide replication for measurements of groat oil content. The locus on linkage group 11 linked to *AccaseA* accounted for up to 48% of the phenotypic variance for groat oil content. Additional QTLs in both populations were identified which accounted for an additional 10% of the phenotypic variance. Thus, the major locus plus additional QTLs represent three
25 homologous genomic segments in oat that can be followed by as few as three RFLP markers to produce a large shift in the groat oil content.

In contrast to oat, a maize ACCase gene was linked only to a moderate oil QTL. Unlike maize, where embryo size affects seed lipid concentration, oat stores significant amounts of lipid in the endosperm, thus
30 QTLs affecting pathway flux may be more important than those altering seed anatomy. If an ACCase gene is indeed the major groat oil QTL, differential ACCase gene expression and/or enzyme activity may control groat oil content.

Moreover, if an ACCase DNA is a QTL in oat, it may be useful as a marker for marker-assisted breeding in oat and other cereals, e.g., maize and wheat.

Table XI. Genomic regions significantly influencing groat oil content in oat.

5 Groat oil values were averages of five (KXO) or three (KxM) environments.

Kanota x Ogle

Locus	Linkage Group	High-Oil Parent	Coefficient of variation	Groat oil, % dry wt
<i>Xcdo665B</i>	11	K	0.38	2.0 #
<i>XaccaseA</i>	11	K	0.38	2.0
<i>Coleoptal</i>	unlinked	K	0.13	0.3
<i>Xcdo1414B</i>	37	K	0.10	1.0
<i>Xcdo1357C</i>	6	K	0.10	0.8
				4.1 Total

Kanota x Marion

Locus	Linkage Group	High-Oil Parent	Coefficient of variation	Groat oil, % dry wt
<i>Xcdo665B</i>	11	K	0.18	0.70 #
<i>XaccaseA</i>	11	K	0.18	0.70
<i>Xcdo1199C</i>	5X	K	0.11	0.70
<i>Xbcd1562A</i>	3	K	0.05	0.40
				1.80 Total

Xcdo665B and *XaccaseA* were associated with the same QTL.

Groat oil values were averages of five (KxO) or three (KxM) environments.

QTLs were identified by single factor analysis of variance.

EXAMPLE XV

Method to Decrease Oil Content in Oat Seed

Youngs (1986) reviewed 11 studies reporting lipid content of oat groats and found an overall range of 2.0 to 11.8% oil concentration in oat groats. Contemporary oat cultivars exhibit an average of 6 to 7% oil (1994-96 Midwest Regional Uniform Nursery data). Recently, Peterson and Wood (1997) reported on the composition and structure of high-oil oat selections ranging from 6.9 to 18.1% oil. While this level of lipid is beneficial for animal feeding because high

oil is associated with higher energy value in the feed, high oil content in oats causes rancidity problems during processing. Rapid liberation of fatty acids from triglycerides by endogenous lipases results in fatty acid oxidation.

Furthermore, use of current oat cultivars results in relatively high-fat food

- 5 products, e.g., oatmeal. Oat processors indicate that a 50% reduction of groat lipid content, while maintaining the superior agronomic characteristics of the currently existing cultivars, would be highly desirable.

- Oat is unique among the cereals in regard to seed lipid deposition (Welch, 1995; Youngs, 1986). Most other cereals accumulate storage lipid
10 primarily in the embryo of developing caryopses. Mature oat groats contain more than 50% of total seed lipid content in the starchy endosperm (Peterson and Wood, 1997). Additionally, significant amounts of oil are found in the bran fraction of processed oats. A substantial portion of the bran is in starchy
15 as in wheat. Together the oat endosperm and bran fractions contain >90% of the total lipid in the groat. Thus, the endosperm is the major source of the oil in the groat.

- Two distinct types of ACCase are known. The first type is typical of eukaryotic organisms where a single, nuclear gene encodes the biotin
20 carboxylase, biotin carboxyl carrier protein, and carboxyl transferase domains within a single, multifunctional polypeptide of 210-280 kDa that, in plants, forms a dimeric active enzyme. Plant lipid synthesis begins in plastids, where ACCase synthesizes the malonyl-CoA needed to initiate and elongate fatty acids of up to 18 carbons (Guenther, 1996). To date, multifunctional ACCase genes
25 from maize, wheat and *Brassica* have been shown to encode functional chloroplast transit peptides. Both monocots and dicots also appear to contain a cytoplasmic form of the multifunctional ACCase. The cytoplasmic ACCase may be involved in the synthesis of malonyl-CoA needed for such reactions as the synthesis of very long chain fatty acids and flavonoids in plants.

- 30 The second, multisubunit type of ACCase consists of three functional domains formed from four separable polypeptides and is found in dicot plastids and in prokaryotic organisms, but appears not to be present in the *Gramineae*. The biotinylated polypeptide component of this multisubunit form

of ACCase is absent from the *Gramineae* and the plastid-localized multifunctional ACCase is believed to be responsible for fatty acid biosynthesis.

Measurements of the levels of acetyl-ACP and malonyl-ACP in spinach leaves (Post-Beittenmiller et al., 1991) and in isolated spinach and pea chloroplasts (Post-Beittenmiller et al., 1991) in the light and dark have provided evidence that ACCase plays a regulatory role in chloroplast fatty acid biosynthesis in vegetative tissues. In monocots, there also is evidence that ACCase activity regulates fatty acid biosynthesis. Several related cyclohexanedione and aryloxyphenoxypropionate graminicides specifically inhibit monocot multifunctional, plastid-localized ACCase. Complete herbicide inhibition of ACCase results in plant cell death due to cessation of fatty acid biosynthesis (Burton et al., 1989). Based on the apparent flux control coefficient for ACCase in barley and maize leaves exposed to low levels of graminicides, ACCase was shown to be the major flux-controlling enzyme for light-stimulated lipid synthesis in these leaves (Page et al., 1994). Parker et al. (1990b) isolated maize tissue cultures that were selected for growth in the presence of normally lethal concentrations of ACCase inhibiting herbicides. Incubation of wild-type cell cultures in the presence of herbicide inhibited incorporation of acetate into the lipid fraction, while variant cell lines exhibiting up to 20-fold increases in wild-type ACCase activity synthesized fatty acid in the presence of herbicide (Parker et al., 1990b). These results provide further evidence that ACCase is rate-limiting in monocot vegetative tissues.

Monocot plastid-localized, multifunctional ACCase is likely involved in regulating seed oil deposition. Herbicide-resistant maize mutant plants were isolated using tissue culture selection (Parker et al., 1990a). The herbicide-resistant phenotype of these plants is conferred by mutations converting plastid-localized, multifunctional, herbicide-sensitive ACCase to herbicide-resistant ACCase (Marshall et al., 1992; Parker et al., 1990a). Two mutant genes for herbicide-insensitive ACCase, *Acc1* and *Acc2*, have been characterized and both are partially dominant (VanDee et al., 1992). Mutant alleles of *Acc1* were used to study the role of ACCase in regulating kernel development and fatty acid synthesis. ACCase activity in corn embryos increased during development paralleling the reported time course of oil

deposition in this tissue (Somers et al., 1993). Furthermore, the maximum value of corn embryo ACCase was two-fold greater than the average ACCase activity determined in seedling leaf extracts, suggesting that the enzyme was involved in elevated synthesis of fatty acids required for oil deposition in the embryo. Using the herbicide-insensitivity of the mutant ACCase as a marker for the *Acc1* gene, we determined the tissue-specific expression pattern of that gene. ACCase activity assayed in extracts of developing kernels harvested from herbicide-resistant plants was tested for herbicide inhibition. The embryo and endosperm ACCase activities exhibited similar reductions of herbicide inhibition as ACCase extracted from the leaves of these herbicide-resistant mutants, indicating that the *Acc1* gene was the major ACCase expressed in the embryo and endosperm of the developing corn kernel. To further investigate this gene-enzyme relationship, homozygous herbicide-resistant and wild-type susceptible kernels were isolated four days after pollination and cultured *in vitro* to maturity in the presence and absence of a range of herbicide concentrations (Somers et al., 1993). Wild-type kernels cultured on all levels of herbicide tested were shrunk and ceased development presumably because of herbicide inhibition of kernel ACCase activity and thus fatty acid synthesis. Mutant kernels developed normally in the presence of herbicide, indicating that the *Acc1*-encoded ACCase is the major isoform expressed in the developing kernel. The results from the corn studies indicate that multifunctional, plastid-localized ACCase is a rate-limiting enzyme in fatty acid biosynthesis in vegetative and seed tissues.

Complete coding sequences of several multifunctional ACCases from higher plants have been described. They show high amino acid homology within the ATP, carboxybiotin, and acetyl-CoA binding domains and an overall amino acid similarity of 70% or greater. Maize multifunctional ACCase isoforms are encoded by at least four genes, A1, A2, B1, and B2 (Egli et al., 1997). Type A cDNAs are > 98% identical, and are 95% identical to a type B cDNA. Types A and B encode highly similar N-termini and are both likely to encode plastid-targeted isoforms (Egli and Gengenbach, 1996; Egli et al., 1997; Egli et al., 1995). Herbicide resistance in corn (Marshall et al., 1992; VanDee et al., 1992) has been mapped to the same chromosomal locations as an ACCase cDNA probe with high identity to both A and B types (Caffrey et al., 1995; Egli

et al., 1994; VanDee et al., 1992). Herbicide-tolerant ACCase is expressed in developing corn kernels (Somers et al., 1993) and both A- and B-type cDNAs are found in immature embryos at the time of peak ACCase activity (Egli and Gengenbach, 1996; Somers et al., 1993). In hexaploid wheat, six cDNAs
5 encoding cytosolic ACCase were attributed either to three homoeologous sets of duplicate genes or to allelic differences in each genome (Podkowinski et al., 1996). Hexaploid wheat also contains three transcriptionally active genes encoding plastidic ACCase with $\geq 98\%$ exon sequence identity. These genes are located on homoeologous chromosomes (Gornicki et al., 1997).

10 Oat is used for both human food and animal feed. Crop improvement goals for oat involve two divergent end uses in that high oil oat is desired as an animal feed whereas a lower oil oat is desired to make low-fat oat products for human consumption. Although progress in decreasing oil content in oat can be achieved by plant breeding, a genetic engineering approach may also
15 be used to create a single, dominant low-oil gene that can be introduced into any oat variety either directly by genetic engineering or via plant breeding for oil reduction in contemporary and future cultivars. Low oil in oat may be a value-added trait for the crop that may allow processors to produce low-fat oat products that contain previously identified cholesterol-lowering constituents
20 including, but not limited to, β -glucans, thus making oats a more healthful human food.

Targeted antisense down-regulation of ACCase in the endosperm using an endosperm-specific promoter may avoid problems that would be caused by inhibition of ACCase in the embryo. Determination of the antisense
25 phenotype by detection of reduced ACCase activity and by quantitation of groat oil content in transgenic lines provides direct evidence for the involvement of ACCase in oat seed oil deposition.

Current oat varieties contain about 7% oil which causes rancidity problems during processing and results in high-fat food products. To reduce oil
30 (triglyceride) content in groats, ACCase cDNA, e.g., oat ACCase DNA, is employed in an antisense orientation to specifically down-regulate oat endosperm ACCase activity and thus reduce groat oil content. Preferably, a plant having a dominantly inherited low-oil gene useful for production of low-oil

varieties is prepared. As oat oil is primarily localized in the starchy endosperm, down-regulation of endosperm ACCase activity is less likely to cause deleterious effects on the embryo. Moreover, converting oil reduction to a single, dominantly inherited antisense trait is useful in rapid production of low oil varieties in this allohexaploid. Seed from transgenic plants are analyzed for expression of the transgene during development and for oil content at maturity. In addition, low oil oat varieties are produced.

Reduction of groat oil content may decrease processing problems associated with oil rancidity and can decrease caloric value resulting in a lower fat food. Reduced-fat oatmeal and other oat products may increase the use of oat as a human food and create a value-enhanced crop for producers of the low-fat oat. In addition, the preparation of high-energy (high oil) oat feed through use of a high-oil gene is also contemplated.

Isolation and Characterization of Oat ACCase cDNAs.

An oat groat cDNA clone was isolated based on its hybridization to a maize cDNA fragment which included portions of the less conserved mid-peptide sequence and the transcarboxylase domain (Shorrosh et al., 1994). Sequence identity indicated that the oat cDNA (3031 bp) corresponded to maize ACCase nt 3124-6169 (Egli et al., 1995). The 1010-aa predicted polypeptide was nearly identical to portions of wheat (89%) and maize (84%) plastidic multifunctional ACCases; it had lower identity to cytosolic ACCases, including wheat (67.5%) and alfalfa (64.6%) (Gornicki et al., 1994; Shorrosh et al., 1994). Therefore, the oat-91 cDNA is likely to encode a portion of a plastid-targeted, multifunctional ACCase polypeptide.

An amplified 'Prairie' oat developing groat cDNA library (R. Skadsen, USDA, Madison, Wisconsin) cloned into the *EcoRI* site of IZAPII (Stratagene) was screened with a [³²P-dCTP]-labeled maize ACCase cDNA probe (Ausabel et al., 1997) derived from an *EcoRI* digest of clone 18-5 (nt 2020-5927; Egli et al., 1995). One to three plaques/plate of eleven partially purified positives were eluted in SM medium plus CHC13. *In vivo* excision was carried out to transform SOLR cells with pBlueScript plasmids carrying oat cDNA inserts, according to the manufacturer's instructions (Stratagene Corp.). Five colonies per transformation reaction were rescreened (as colony lifts) with

an AflIII-BbrPI fragment of maize ACCase cDNA (4281-5797; Egli et al., 1995). One clone, oat-91, hybridized to the probe. Oat-91 was sequenced using the ABI Prizm system (Perkin-Elmer Corp.) at the Advanced Genetic Analysis Center (University of Minnesota, St. Paul, Minnesota). Sequence identity indicated the
5 3.05-kb *EcoRI* insert of oat-91 corresponded to maize ACCase nt 3124-6169 (Egli et al., 1995). Oat ACCase cDNA sequence was compared to other plant ACCases with the BESTFIT program (BESTFIT, Wisconsin Package Version 9.1, Genetics Computer Group (GCG), Madison, Wisconsin).

For PCR, Pileup alignments of maize, wheat, barley, and rice
10 ACCase coding sequences (GCG, Madison, Wisconsin; 12, 18, F, R) were used to identify conserved regions (i) for design of new primers or (ii) regions of identity to available maize ACCase oligos, to be used for PCR amplification of oat ACCase. Two maize primers, PCR2 (5'-ggcagagcaa agcttgagg) and PCR1 (5'-tccaccagag aagcctctcc) appeared likely to amplify oat cDNA based on their
15 identity to other monocots and previous use to amplify soybean ACCase. These primers are located within the transcarboxylase domain, surrounding the acetyl-CoA binding site of ACCase (50). An amplified 'Prairie' oat seeding leaf cDNA library cloned into the *EcoRI* site of IZAPII (Skadsen, University of Wisconsin) was used as a source of cDNA. One I of the oat cDNA library was amplified
20 with maize primers (PCR2/PCR1), using the Expand High Fidelity PCR system (BMB) in a GeneAmp PCR System 9600 thermal cycler (Perkin-Elmer) after denaturing phage heads at 94°C for 10 minutes. The single, 222-bp product was purified by Wizard PCR Plus (Promega Corp.) and sequenced in an AB1377 (Perkin-Elmer) using PCR1 as primer. The resulting sequence appeared to be
25 from oat since it had high identity to maize plastidic ACCase (89%; 4, 12, Egli, unpublished). Additional 5' cDNA was obtained by (i) PCR of the oat cDNA library with T3/PCR1 primers to generate a complex band mixture and (ii) reamplification of 0.1 I products with maize ACCase primers 1324+ (5'-caatcacgtcagactg)/PCR1 to obtain a 1.9-kb product with the same 3' sequence as
30 the PCR2/PCR1 product.

To test whether the oat ACCase cDNA clone oat-91 encodes a plastidic isoform, 422 nt of 3' insert sequence was translated and compared to predicted amino acid sequences of maize plastidic ACCases, a putative maize

cytosolic ACCase, and to multifunctional ACCases from wheat, barley, alfalfa, soybean, *Arabidopsis*, *Brassica*, chicken, human, *Ustilago*, yeast and *Cyclotella*. Aligned, degapped peptides were used to construct a phylogram by the UPGMA method based on Kimura protein distances (GCG, Madison, Wisconsin). The
5 oat ACCase sequence was $\geq 93\%$ identical to plastidic maize and wheat ACCase sequences, but only 78-83% identical to cytosolic ACCases from maize, wheat, or dicots. The corresponding nucleotide identities were 88-92% and 73-78%, respectively. In the phylogram, monocot ACCases were separated from dicots and grouped according to their known or expected cellular locations (Figure 25).

- 10 Both sequence identity and phylogenetic grouping indicate that oat cDNA encodes a plastidic ACCase.

Additional 5' sequence is obtained by screening the cDNA library with the most 5' oat probes or by PCR of the library with near 5' (upstream) oat plus vector-specific oligonucleotide primers. Alternatively, the 5' end of
15 ACCase message may be obtained by 5' RACE PCR (Frohman, 1993). Total RNA is isolated from developing groats, and digested with DNase.

Complementary DNA is synthesized with Superscript II reverse transcriptase (Gibco BRL) by priming with oat ACCase primers. PCR amplification of cDNA utilizes oat primers plus 5' maize primers. Because chloroplast transit peptide
20 sequences are not well conserved between ACCase genes of maize, wheat, and *Brassica*, or between different gene families of a single species, and because 5' non-coding sequences also are unlikely to be conserved, completion of the 5' sequence of oat ACCase cDNA is accomplished by 5' RACE-PCR (GIBCO-BRL; Gornicki et al., 1994). 5' RACE products and downstream sequences are

- 25 sequenced to verify that a chloroplast transit peptide coding sequence is present and to identify primers that may be used to generate a complete or partial oat ACCase cDNA product by PCR. Oat cDNA is reamplified using either the Expand High Fidelity or Long Template PCR kits (Boehringer Mannheim Biochemicals), depending on the length of desired product. Products are blunt-
30 cloned into pBluescript (Stratagene) or a similar vector, or extended with dATP and cloned into pGEM-T (Promega). Adjacent sites in vector polycloning regions or unique sites introduced by PCR are utilized to move oat ACCase cDNA into expression vectors.

Co-segregation of Oat ACCase cDNA and a Major Groat Oil QTL

A major quantitative trait locus (QTL) controlling groat oil content in oat was recently identified by the oat genome mapping program at the University of Minnesota (See Example XIV). Single factor ANOVA showed
5 that RFLP marker *Xcdo665B* on linkage group 11 was associated with a QTL accounting for approximately 37% of the variance in groat oil in a Kanota by Ogle mapping population (O'Donoghue et al., 1995). The 1.9 kb oat ACCase cDNA was placed on this map to within 2.6 cM of *Xcdo665B* in the same population (n = 136). This ACCase polymorphism also accounted for about
10 37% of the variation in groat oil content. These results provide strong support for the hypothesis that ACCase has a major role in determining groat oil content. Either allelic variation in ACCase activity controls oil content in oat groats or a gene(s) closely linked to ACCase is responsible. Suppression of ACCase expression by antisense transformation can provide a direct test of whether
15 ACCase activity indeed has a major effect on groat oil.

Antisense ACCase Transformation Construct Production

Antisense ACCase cDNAs are constructed using standard procedures in the plasmids that are routinely used for oat transformation (Torbert et al., 1995). The cDNA is ligated in antisense orientation relative to an
20 endosperm-specific promoter and is terminated using the nopaline synthase (NOS) 3' untranslated sequence of the *Agrobacterium tumefaciens* NOS gene, although other termination regions may be employed. This antisense cassette is ligated into a plasmid backbone carrying a neomycin phosphotransferase II (nptII) gene under control of the cauliflower mosaic virus 35S promoter as a
25 selectable marker (Torbert et al., 1995).

An investigation of down-regulated phenotypes conferred by expression of various granule-bound starch synthase antisense constructs in potato indicated that the size of the antisense transcript did not affect efficacy of inhibition (Kuipers et al., 1995). Choice of promoter sequences controlling
30 expression of the antisense construct appeared to have important effects in the degree of antisense inhibition of potato granule-bound starch synthase. Effective antisense RNAs have been reported which (relative to the mRNA) are either full length or fragments designed to block 5' end capping, intron splicing, or the poly

A tail (Guerineau et al., 1994; Inoye, 1988). Preferably, the longest oat ACCase clones are employed in an anti-sense construct.

Complete inhibition of vegetative- or seed-expressed ACCase is likely to be lethal because the enzyme is required for general fatty acid biosynthesis, as shown by the lethality of herbicide inhibition of ACCase in susceptible monocots. In oat, it is not known whether tissue-specific isozymes or homologs of ACCase exist, or if they do, whether they vary significantly in nucleotide sequence. Therefore, it is important to target antisense down-regulation of ACCase to the developing oat endosperm and that the antisense gene exert a partial down-regulation of endosperm ACCase homologs to avoid deleterious effects of possible endosperm ablation due to complete inhibition of ACCase. Because antisense suppression of heterologous genes or members of a gene family can occur (deLang et al., 1993), and because the partial inhibition of plastidic ACCases but not cytosolic ACCases is desired, partial antisense cDNAs designed from functional domain sequences may be used, or conversely, less-conserved regions between functional domains. Comparison of wheat genes for cytosolic and plastidic ACCases (Gornicki et al., 1997; Gornicki et al., 1994) may aid in the design of antisense cDNAs to specifically target plastidic ACCase.

The wheat Dy10 high molecular weight glutenin promoter (Belchi and Anderson, 1994) confers strong expression of GUS in oat and expression is specifically limited to the endosperm. Because of its tissue specificity and the likelihood that its expression pattern does not directly coincide with the expression of ACCase during oat endosperm development, the Dy10 promoter may be useful in restricting the ACCase antisense transcripts to the oat endosperm resulting in a partial inhibition of ACCase expression and the normal development of endosperm cells.

Oat Transformation

Elite cultivars of oat can be transformed and on average one transgenic tissue culture is produced that regenerates fertile, transgenic plants per two microprojectile bombardments. Generally, from 15 to 30 transgenic plants are produced from the microprojectile bombardments conducted during one experiment. The oat ACCase antisense plasmid is introduced into regenerable

oat tissue cultures of oat genotypes that exhibit high oil content using microprojectile bombardment as described by Torbert et al. (1995). Tissue cultures are selected for resistance to paromomycin conferred by the nptII selectable marker gene. Paromomycin-resistant tissue cultures are regenerated
5 and plants analyzed for the nptII protein using an NPTII ELISA kit (3' > 5' Inc., Boulder, CO). Plants that express the nptII protein are grown to maturity.

The progeny from at least 100 independent transgenic events are then evaluated.

Characterization of the ACCase Antisense Phenotype

10 Homozygous T₂ seed are divided for maintenance, advancement to produce greater quantities of seed for more detailed analyses, and for preliminary analysis. Single oat plants may produce up to 100 seed. Therefore, sufficient seed should be available to determine if a transgene phenotype is detectable. Dried mature seed of antisense homozygotes and nontransgenic
15 sister lines are analyzed for oil content initially using nondestructive Near Infrared Reflectance (NIR) spectroscopy. This method requires only 1 g of dehulled groats and is inexpensive. NIR is employed to rapidly identify transgenic lines that exhibit lowered oil content. Selected lines identified by NIR are reanalyzed using either NMR or the gravimetric procedure (Peterson
20 and Wood, 1997). If differences in oil content are observed, oil composition is analyzed by gas chromatography. If the results of these analyses warrant further investigation of the materials, a replicated greenhouse trial is conducted to confirm the reduced oil phenotype in the transgenic plants. This test is conducted on T₃ seed produced from multiple T₂ plants of each transgenic line.
25 If a low-oil phenotype is observed, these materials are entered into a breeding program to initiate variety development.

Selected low-oil lines are advanced an additional generation. Seed is isolated at different stages of development and stored under conditions to preserve ACCase activity and mRNA for Northern blot analysis. ACCase
30 activity levels and sense and antisense transcript levels are determined to further characterize the antisense phenotype of these plants.

While oat is considered a beneficial alternative crop with multiple uses, production acreage continues to shrink due to poor profitability and

international competition. Yet oat is used as a high quality animal feed and for production of nutritious human food. Numerous positive human health qualities are attributed to consumption of oatmeal. Foremost among these is the cholesterol-lowering property of oats. A major problem confronting oat
5 processors is that current oat varieties have elevated oil content that precludes the production of a low-fat oat product. Thus, in the production of oat products such as oatmeal, the high oil content offsets the benefit of the cholesterol-lowering constituents of the product. Oat oil content is relatively easy to manipulate using conventional plant breeding. However, the trend has been to
10 increase oil content of oats for use as animal feed and only recently has low-oil oat been recognized as important for food products.

Thus, oat varieties with reduced oil content may be prepared using genetic engineering. Moreover, a single dominant gene system can vastly simplify conversion of existing agronomically adapted oat varieties to low oil as
15 well as being useful in the production of new low-oil varieties.

Based on the demand by processors for low-oil oat varieties, it is warranted to consider low oil as a value-added trait for oat. Increasing the value of oat may increase the acreage devoted to growing oats in the U.S. because of possible increased profitability of the crop.

20

References

- Abu-Elheiga, L., Jayakumar, A., Baldini, A., Chirala S. S., Wakil, S. J. Human acetyl-CoA carboxylase: characterization, molecular cloning and evidence for two isoforms. Proc. Nat. Acad. Sci. U.S.A., 92:4011-4015 (1995).
- 25 Ahn, S., Tanksley, S. D. Comparative linkage maps of the rice and maize genomes. Proc. Natl. Acad. Sci. U.S.A., 90:7980-7984 (1993).
- Alban, C., Baldet, P., Douce, R. Localization and characterization of two structurally different forms of acetyl-CoA carboxylase in young pea leaves, of which one is sensitive to aryloxyphenoxypropionate herbicides. Biochem. J.,
30 300:557-565 (1994).

- Al-Feel, W., Chirala, S. S., Wakil, S. J. Cloning of the yeast FAS3 gene and primary structure of yeast acetyl-CoA carboxylase. Proc. Natl. Acad. Sci. U.S.A., **89**:4534-4538 (1992).
- Alrefai, R., Berke, T. G., Rocheford, T. R. Quantitative trait locus
5 analysis of fatty acid concentrations in maize. Genome, **38**:894-901 (1995).
- Anderson, J. V., Lutz, S. M., Gengenbach, B. G., Gronwald, J. W. Genomic sequence for a nuclear gene encoding acetyl-coenzyme A carboxylase (GenBank L42814) in soybean. Plant Physiol., **109**:338 (1995).
- Ashton, A., Jenkins, C. L. D., Whitfeld, P. R. Molecular cloning of two
10 different cDNAs for maize acetyl-CoA carboxylase. Plant Mol. Biol., **24**:35-49 (1994).
- Ausubel F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A., Struhl, K., eds. Current Protocols in Molecular Biology, John Wiley & Sons, Inc. (1997).
- 15 Bailey, A., Keon, J., Owen, J., Hargreaves, J. The ACC1 gene, encoding acetyl-CoA carboxylase, is essential for growth in *Ustilago maydis*. Mol. Gen. Genet., **249**:191-201 (1995).
- Belchl, A. E., Anderson, O. D. Expression of a novel high-molecular-weight glutenin subunit gene in transgenic wheat. Nature Biotechnology, **14**
20 (1996).
- Berke, T.G., Rocheford, T.R. Quantitative trait loci for flowering, plant and ear height, and kernel traits in maize, Crop Sci., **35**, 1542-1549 (1995).
- Betty, M., Ireland, R. J., Smith, A. M. Purification and characterization of acetyl-CoA carboxylase from developing pea embryos. Plant Physiol. J.,
25 **140**:513-520 (1992).
- Branson, V., Frey, K. J. Recurrent selection for groat oil content in oat. Crop Sci., **28**:1382-1387 (1989).
- Brummer, E. C., Graef, G. L., Orf, J., Wilcox, J. R., Shoemaker, R. C. Mapping QTL for seed protein and oil content in eight soybean populations.
30 Crop Sci., **37**:370-378 (1997).
- Burton, J. D., Gronwald, J. W., Somers, D. A., Gengenbach, B. G., Wyse, D. L. Inhibition of corn acetyl-CoA carboxylase by cyclohexanedione and aryloxyphenoxypropionate herbicides. Pest Biochem. Physiol., **34**:76-85 (1989).

- Bush, A. L., Wise, R. P., Rayapati, P. J., Lee, M. Restriction fragment length polymorphisms linked to genes for resistance to crown rust (*Puccinia coronata*) in near-isogenic lines of hexaploid oat (*Avena sativa*). Genome, **37**:823-831 (1994).
- 5 Caffrey, J. J., Wurtele, E. S., Nikolau, B. J. Genetic mapping of two acetyl-CoA carboxylase genes. Maize Genetics Cooperation Newsletter, **69**:3-4 (1995).
- Cakir, M., Reeves, D. L., Kahler, A. L. Linkage analysis of RFLP markers and quantitative traits in hexaploid oats. Agronomy Abstracts.
- 10 American Society of Agronomy, Madison, Wisconsin, pp. 163 (1996).
- Charles, D. J., Cherry, J. H. Purification and characterization of acetyl-CoA carboxylase from developing soybean seeds. Phytochem., **25**:1067 (1986).
- de Lange, P., de Boer, G-J, Mol, J. M. N., Kooter, J. M. Conditional inhibition of β -glucuronidase expression by antisense gene fragments in petunia
- 15 protoplasts. Plant Molec. Biol., **23**:45-55 (1993).
- Diers, B. W., Keim, P., Fehr, W. R., Shoemaker, R. C. RFLP analysis of soybean seed protein and oil content. Theor. Appl. Genet., **83**:608-612 (1992).
- Dudley, J. W., Lambert, R. J., Alexander, D. E. Seventy generations of selection for oil and protein concentration in the maize kernel. In: Dudley, J. W.
- 20 (ed.) Seventy generations of selection for oil and protein concentration in the maize kernel. Crop Sci. Soc. Am., Madison, Wisconsin, pp. 181-212 (1974).
- Egli, M., Gengenbach, B. Plastid localization of a multifunctional acetyl-CoA carboxylase in maize. Maize Genetics Cooperation Newsletter, **70**:58-59 (1996).
- 25 Egli, M. A., Gengenbach, B. G., Gronwald, J. W., Somers, D. A., Wyse, D. L. Characterization of maize acetyl-coenzyme A carboxylase. Plant Physiol., **101**:499-506 (1993).
- Egli, M., Lutz, S., Gengenbach, B. Genetic analysis of maize plastidic acetyl-CoA carboxylase. 1997 Biochemistry and Molecular Biology of Plant
- 30 Fatty Acid and Glycerolipids Symposium, South Lake Tahoe, CA, P-4, (1997).
- Egli, M., Lutz, S., Somers, D., Gengenbach, B. Identification and mapping of maize acetyl-CoA carboxylase genes. Maize Genetics Cooperation Newsletter, **68**:92-93 (1994).

- Egli, M. A., Lutz, S. M., Somers, D. A., Gengenbach, B. G. A maize acetyl-coenzyme A carboxylase cDNA sequence. Plant Physiol., **108**:1299-1300 (1995).
- Elborough, K. M., Simon, J. W., Swinhoe, R., Ashton, A. R., Slabas, A.
5 R. Studies on wheat acetyl CoA carboxylase and the cloning of a partial cDNA. Plant Mol. Biol., **24**:21-34 (1994).
- Evenson, K. J., Gronwald, J. W., Wyse, D. L. Purification and characterization of acetyl-coenzyme A carboxylase from diclofop-resistant and -susceptible *Lolium multiflorum*. Plant Physiol., **105**:671-680 (1994).
- 10 Frohman, M. A. Rapid amplification of complementary DNA ends for generation of full-length complementary DNAs: Thermal RACE. Methods in Enzymology, **218**:340-356 (1993).
- Gale, M. D., Devos, K. M., Proc. Natl. Acad. Sci. USA, **95**, 1971 (1998).
- Goldman, I. L., Rocheford, T. R., Dudley, J. W. Quantitative trait loci
15 influencing protein and starch concentration in the Illinois Long Term Selection maize strains. Theor. Appl. Genet., **87**:217-224 (1993).
- Goldman, I. L., Rocheford, T. R., Dudley, J. W. Molecular markers associated with maize kernel oil concentration in an Illinois high protein X Illinois low protein cross. Crop. Sci., **34**:908-915 (1994).
- 20 Gornicki, P., Faris, J., King, I., Podkowinski, J., Gill, B., Haselkorn, R. Plastid-localized acetyl-CoA carboxylase of bread wheat is encoded by a single gene on each of the three ancestral chromosome sets. Proc. Natl. Acad. Sci. U.S.A., **94**:14179-14184 (1997).
- Gornicki, P., Haselkorn, R. Wheat acetyl-CoA carboxylase. Plant Mol.
25 Biol., **22**:547-552 (1993).
- Gornicki, P., Podkowinski, J., Scappino, L. A., DiMaio, J., Ward, E., Haselkorn, R. Wheat acetyl-coenzyme A carboxylase: cDNA and protein structure. Proc. Natl. Acad. Sci. U.S.A., **91**:6860-6864 (1994).
- Gornicki, P., Faris, J., King, I., Podkowinski, J., Gill, B., Haselkorn, R.
30 Plastid-localized acetyl-CoA carboxylase of bread wheat is encoded by a single gene on each of the three ancestral chromosome sets, Proc. Nat. Acad. Sci. USA, **94**, 14179-14184 (1997).

- Gornicki, P., Podkowinski, J., Scappino, L.A., DiMaio, J., Ward, E., Haselkorn, R. Wheat acetyl-coenzyme A carboxylase: cDNA and protein structure, Proc. Natl. Acad. Sci. USA, **91**, 6860-6864 (1994).
- Guenther, S. Genbank accession #X99102 (barley) (1996).
- 5 Guerineau, F., Waugh, R. Brown, J. W. S. Parameters affecting the activity of antisense RNA sequences in tobacco protoplasts. Plant Cell Rep., **13**:703-708 (1994).
- Guchhait, R. B., Polakis, E., Dimroth, P., Stoll, E., Moss, J., Lane, M. D. Acetyl coenzyme A carboxylase system of *Escherichia coli*. J. Biol. Chem.,
10 **249**:6633-6645 (1974).
- Gullord, M. Oil and protein content in oats (*Avena sativa* L.). In: Lawes, D. A., and Thomas, H. (eds.) Proc. 2nd Int. Oats Conf. Aberystwyth, Wales. 15-18 July 1985. Martinus Nijhoff Publishers, Dordrecht, The Netherlands, pp. 210-213 (1986).
- 15 Harwood, J. L. Fatty acid metabolism. Ann. Rev. Plant Physiol. Plant Mol. Biol., **39**:101-138 (1988).
- Holland, J. B., Moser, H.S., O'Donoghue, L. S., Lee, M. QTLs and epistasis associated with vernalization responses in oat. Crop Sci., **37**:1306-1316 (1997).
- 20 Hosaka, K., Kianian, S. F., McGrath, J. M., Quiros, C. F. Development and chromosomal localization of genome-specific DNA markers of *Brassica* and the evolution of amphidiploids and n = 9 diploid species. Genome, **33**:131-142 (1990).
- Hu, J., Quiros, C. F., Arus, P., Struss, D., Robbelen, G. Mapping of a
25 gene determining linolenic acid concentration in rapeseed with DNA-based markers. Theor. Appl. Genet., **90**:258-262 (1995).
- Inouye, M. Antisense RNA: its functions and applications in gene regulation – a review. Gene, **72**:25-34 (1988).
- Keegstra, K., Olsen, L. J., Theg, S. M. Chloroplastic precursors and their
30 transport across the envelope membranes. Annu. Rev. Plant Physiol. Plant Mol. Biol., **40**:471-501 (1989).

- Kennard, W. C., Havey, M. J. Quantitative trait analysis of fruit quality in cucumber: QTL detection, confirmation, and comparison with mating-design variation. *Theor. Appl. Genet.*, **91**:53-61 (1995).
- Kennard, W.C., Slocum, M.K., Figdore, S.S., Osborn, T.C. Genetic
5 analysis of morphological variation in *Brassica oleracea* using molecular markers, *Theor. Appl. Genet.*, **87**, 721-732 (1994).
- Kianian, S. F., Wu, B-C, Fox, S. L., Rines, H. W., Phillips, R. L. Aneuploid marker assignment in hexaploid oat with the C genome as a reference for determining remnant homoeology. *Genome*, **40**:386-396 (1997).
- 10 Kolattukudy, P. E. Chemistry and Biochemistry of Natural Waxes. P. E. Kolattukudy, ed., Elsevier, New York, pp. 289-347 (1976).
- Kondo, H., Shiratsuchi, K., Yoshimoto, T., Mauda, T., Kitazono, A., Tsuru, D., Anai, M., Sekiguchi, M., Tanabe, T. Acetyl-CoA carboxylase from *E. coli*: gene organization and nucleotide sequence of the biotin carboxylase
15 subunit. *Proc. Natl. Acad. Sci. U.S.A.*, **88**:9730-9733 (1991).
- Konishi, T., Shinohara, K., Yamada, K., Sasaki, Y. Acetyl-CoA carboxylase in higher plants: most plants other than Gramineae have both the prokaryotic and the eukaryotic forms of this enzyme. *Plant Cell Physiol.*, **37**:117-122 (1996).
- 20 Kuipers, A. G. S., Soppe, W. J. J., Jacobsen, E., Visser, R. G. F. Factors affecting inhibition by antisense RNA of granule-bound starch synthase gene expression in potato. *Mol. Gen. Genet.*, **246**:745-755 (1995).
- Lee, E-S J., Gronwald, J. W., Orf, J. H. Acetyl-CoA carboxylase activity in high-oil and low-oil soybeans (*Glycine max* L.). 11th International Meeting
25 on Plant Lipids. Paris, France. Abstract A14 (1994).
- Li, S-J, Cronan, J. E. The gene encoding the biotin carboxylase subunit of *E. coli* acetyl-CoA carboxylase. *J. Biol. Chem.*, **267**:855-863 (1992).
- Li, S-J, Cronan, J. E. The genes encoding the two carboxyltransferase subunits of *E. coli* acetyl-CoA carboxylase. *J. Biol. Chem.*, **267**:16841-16847
30 (1992).
- Lopez-Casillas, F., Bai, D-H, Luo, X., Kong, I-S, Hermodson, M. A., Kim, K-H. Structure of the coding sequence and primary amino acid sequence

of acetyl-coenzyme A carboxylase. Proc. Natl. Acad. Sci. U.S.A., **85**:5784-5788 (1988).

- 5 Mansur, L. M., Lark, K. G., Kross, H., Oliveira, A. Interval mapping of quantitative trait loci for reproductive, morphological, and seed traits of soybean (*Glycine max* L.). Theor. Appl. Genet., **86**:907-913 (1993).

Marshall, L. C., Somers, D. A., Dotray, P. A., Gengenbach, B. G., Wyse, D. L., Gronwald, J. W. Allelic mutations in acetyl-coenzyme A carboxylase confer herbicide tolerance in maize. Theor. Appl. Genet., **83**:435-442 (1992).

- 10 Marten, G. C., Shenk, J. S., Barton II, F. E. (ed.) Near infrared reflectance spectroscopy (NIRS): Analysis of forage quality. Agric. Handb. 643. USDA-ARS, Washinton, D. C. (1989).

Morrison, W. R. Cereal lipids. In: Pomeranz, Y. (ed.) Advances in cereal science and technology. Vol. 2. Am. Assoc. Cereal Chem., St. Paul, Minnesota, pp. 221-348 (1978).

- 15 Nikolau, B.J., Hawke, J.C. Purification and characterization of maize leaf acetyl-CoA carboxylase, Arch. Biochem. Biophys., **228**, 86-96 (1984).

- O'Donoghue, L. S., Kianian, S. F., Rayapati, P. J., Penner, G. A., Sorrells, M. E., Tanksley, S. D., Phillips, R. L., Rines, H. W., Lee, M., Fedak, G., Molnar, S. J., Hoffman, D., Salas, C. A., Wu, B., Autrique, E., Van Deynze, A. A molecular linkage map of cultivated oat. Genome, **38**:368-380 (1995).
- 20

Ohlrogge, H. B., Jaworski, J. B., Regulation of fatty acid synthesis, Annu. Rev. Plant Physiol. Plant Mol. Biol., **48**, 109-136 (1997).

- Page, R. A., Okada, S., Harwood, J. L. Acetyl-CoA carboxylase exerts strong flux control over lipid synthesis in plants. Biochem. Biophys. Acta, **1210**:369-372 (1994).
- 25

Page, R.A., Okada, S., Harwood, J.L. Acetyl-CoA carboxylase exerts strong flux control over lipid synthesis in plants, Biochem. Biophys. Acta, **1210**, 369-372 (1994).

- Parker, W. B., Marshall, L. C., Burton, J. B., Somers, D. A., Wyse, D. L., Gronwald, J. W., Gengenbach, B. G. Dominant mutations causing alterations in acetyl-coenzyme A carboxylase confer tolerance to cyclohexanedione and aryloxyphenoxypropionate herbicides in maize. Proc. Natl. Acad. Sci. U.S.A., **87**:7175-7179 (1990a).
- 30

- Parker, W. B., Somers, D. A., Wyse, D. L., Keith, R. A., Burton, J. D., Gronwald, J. W., Gengenbach, B. G. Selection and characterization of sethoxydim-tolerant maize tissue cultures. Plant Physiol., **92**:1220-1225 (1990b).
- 5 Peterson, D. M., Wood, D. F. Composition and structure of high oil oat. J. Cereal Sci., **26**:121-128 (1997).
- Phillips, R. L., Vasil, I. K. (eds.) DNA-based markers in plants. Kluwer Academic Publishers. Dordrecht, The Netherlands (1994).
- 10 Podkowinski, J., Sroga, G. E., Haselkorn, R., Gornicki, P. Structure of a gene encoding a cytosolic acetyl-CoA carboxylase of hexaploid wheat. Proc. Natl. Acad. Sci. U.S.A., **93**:1870-1874 (1996).
- Post-Beittenmiller, D. J., Jaworski, J. G., Ohlrogge, J. B. *In vivo* pools of free and acylated acyl carrier proteins in spinach. Evidence for sites of regulation of fatty acid biosynthesis. J. Biol. Chem., **226**:1858-1865 (1991).
- 15 Post-Beittenmiller, D., Roughan, P. G., Ohlrogge, J. B. Regulation of plant fatty acid biosynthesis: Analysis of acyl-coenzyme A and acyl-acyl carrier substate pools in spinach and pea chloroplast. Plant Physiol., **100**:923-930 (1992).
- 20 Rajhathy, T., Thomas, H. Cytogenetics of oats (*Avena* L.). Wilkes, A. (ed.). Genetics Society of Canada Miscellaneous Publication No. 2. Genetics Society of Canada, Ottawa, Ontario (1974).
- Ripsin, C. M., Keenan, J. M., Jacobs, D. R., Elmer, P. J., Welch, R. R., Van Horn, L., Liu, K., Turnbull, W. H., Thye, F. W., Kestin, M., Hegsted, M., Davidson, D. M., Davidson, M. H., Dugan, L. D., Demark-Wahnefried, W.,
- 25 Beling, S. Oat products and lipid lowering. J. Am. Med. Assoc., **267**:3317-3325 (1992).
- Robertson, D. S. A possible technique for isolating genic DNA for quantitative traits in plants. J. Theor. Biol., **117**:1-10 (1985).
- 30 Rocheford, T. R. Chromosome Regions Associated with Control of Maize Kernel Composition. Proceedings of the 49th Annual Sorghum Research Conference, pp. 239-249 (1994).

- Roesler, K., Shintani, D., Savage, L., Boddupalli, S., Ohlrogge, J. Targeting of the *Arabidopsis* homomeric acetyl-coenzyme A carboxylase to plastids of rapeseeds. *Plant Physiol.*, 113:75-81 (1997).
- Roesler, K. R., Shorrosh, B. S., Ohlrogge, J. B. Structure and expression
5 of an *Arabidopsis* acetyl-coenzyme A carboxylase gene. *Plant Physiol.*,
105:611-617 (1994).
- Roessler, P. G., Ohlrogge, J. B. Cloning and characterization of the gene that encodes acetyl-coenzyme A carboxylase in the alga *Cyclotella cryptica*. *J. Biol. Chem.*, 268:19254-19259 (1993).
- 10 Rooney, W. L., Jellen, E. N., Phillips, R. L., Rines, H. W., Kianian, S. F. Identification of homoeologous chromosomes in hexaploid oat (*A. byzantina* cv. Kanota) using monosomics and RFLP analysis. *Theor. Appl. Genet.*, 89:329-335 (1994).
- Sambrook, J., Fritsch, E.F., Maniatis, T. Molecular cloning: A laboratory
15 manual, 2nd ed., Cold Spring Harbor Laboratory Press (1989).
- Siripoonwiwat, W., O'Donoghue, L.S., Wesenberg, D., Hoffman, D.L., Barbosa-Neto, J.F., Sorrells, M.E. Chromosomal regions associated with quantitative traits in oats. JQTL (<http://probe.nalusda.gov:8000/otherdocs/jqtl/jqtl1996-03>) (1996).
- 20 SAS Institute, Inc. SAS procedures guide, Version 6, 3rd ed. SAS Institute Inc., Cary, N.C. (1990).
- Schipper, H., Frey, K. J. Selection for groat-oil content in oats grown in field and greenhouse. *Crop Sci.*, 31:661-665 (1991a).
- Schipper, H., Frey, K. J. Observed gains for three recurrent selection
25 regimes for increased groat-oil content. *Crop Sci.*, 31:1505-1510 (1991b).
- Schrackel, D. J. Oat production, value and use. In: Webster F. H. (ed.) Oat chemistry and technology. Assoc. Cereal Chem., St. Paul, Minnesota, pp. 1-11 (1986).
- Schulte, W., Schell, J., Toepfer, R. A gene encoding acetyl-coenzyme A
30 carboxylase from *Brassica napus*. *Plant Physiol.*, 106:793-794 (1994).
- Schulte, W., Topfer, R., Stracke, R., Schell, J., Martini, N. Multi-functional acetyl-CoA carboxylase from *Brassica napus* is encoded by a multi-

- gene family: Indication for plastidic localization of at least one isoform. Proc. Natl. Acad. Sci. U.S.A., **94**:3465-3470 (1997).
- Sears, E. R. The nature of mutation in hexaploid wheat. Symp. Biol. Hung., **12**:73-82 (1972).
- 5 Shintani, D., Roesler, K., Shorrosh, B., Savage, L., Ohlrogge, J. Antisense expression and overexpression of biotin carboxylase in tobacco leaves. Plant Physiol., **114**:881-886 (1997).
- Shorrosh, B. S., Dixon, R. A., Ohlrogge, J. B. Molecular cloning, characterization, and elicitation of acetyl-CoA carboxylase from alfalfa. Proc. Natl. Acad. Sci. U.S.A., **91**:4323-4327 (1994).
- 10 Simcox, P. D., Gerland, W., De Luca, V., Canvin, D. T., Dennis, D. T. Respiratory pathways and fat synthesis in the developing castor oil seed. Can. J. Bot., **57**:1008-1014 (1979).
- Siripoonwiwat, W., O'Donoghue, L. S., Wesenberg, D., Hoffman, D.
- 15 L., Barbosa-Neto, J. F., Sorrells, M. E. Chromosomal regions associated with quantitative traits in oat. JQTL (<http://probe.nalusda.gov:8000/otherdocs/jqtl/jqtl1996-03>) (1996).
- Slabas, A. R., Fawcett, T. The biochemistry and molecular biology of plant lipid biosynthesis. Plant Mol. Biol., **19**:169-191 (1992).
- 20 Slabas, A. R., Hellyer, A., Bambridge, H. E. The basic polypeptide subunit of rape leaf acetyl-CoA carboxylase is a 220 kDa protein. Biochem. Soc. Trans., **14**:714-715 (1986).
- Somers, D. A., Keith, R. A., Egli, M. A., Marshall, L. C., Gengenbach, B. G., Gronwald, J. W., Wyse, D. L. Expression of the Acc1 gene-encoded
- 25 acetyl-coenzyme-A carboxylase in developing maize (*Zea mays* L.) kernels. Plant Physiol., **101**:1097-1101 (1993).
- Somers, D. A., Rines, H. W., Gu, W., Kaeppler, H. F., Bushnell, W. R. Fertile, transgenic oat plant. Bio/Technology, **10**:1589-1594 (1992).
- Somers, D. A., Torbert, K. A., Pawlowski, W. P., Rines, H. W., Progress
- 30 in genetic engineering of oat. Proceedings of the 5th International Oat Conference and the 7th International Barley Genetics Symposium, Scoles, G., Rossnagel, B., eds. University Extension Press, University of Saskatchewan, pp. 230-234 (1996).

- Takai, T., Yokayama, C., Wada, K., Tanabe, T. Primary structure of chicken liver acetyl-CoA carboxylase deduced from cDNA sequence. *J. Biol. Chem.*, **263**:2651-2657 (1988).
- 5 Tanhuanpaa, P. K., Vikki, J. P., Vikki, H. J. Identification of a RAPD marker for palmitic-acid concentration in the seed of spring turnip rape (*Brassica rapa* spp. *oleifera*). *Theor. Appl. Genet.*, **91**:477-480 (1995a).
- Tanhuanpaa, P. K., Vikki, J. P., Vikki, H. J. Association of a RAPD marker with linolenic acid concentration in the seed oil of rapeseed (*Brassica napus* L.). *Genome*, **38**:414-416 (1995b).
- 10 Thro, A. M., Frey, K. J. Inheritance of groat-oil content and high-oil selection in oats (*Avena sativa* L.). *Euphytica*, **34**:251-263 (1985).
- Tinker, N. A., Mather, D. E. Methods for QTL analysis with progeny replicated in multiple environments. *IQTL* (<http://probe.nalusda.gov:8000/otherdocs/jqtl/jqtl1995-01>) (1995a).
- 15 Tinker, N. A., Mather, D. E. MQTL: software for simplified composite interval mapping of QTL in multiple environments. *IQTL* (<http://probe.nalusda.gov:8000/otherdocs/jqtl/jqtl1995-02>) (1995b).
- Tinker, N. A., Mather, D. E., Rossnagel, B. G., Kasha, K. J., Kleinhofs, A., Hayes, P. M., Falk, D. E., Ferguson, T., Shugar, L. P., Legge, W. G., Irvine, R. B., Choo, T. M., Briggs, K. G., Ullrich, S. E., Franckowiak, J. D., Blake, T. K., Graf, R. J., Dofing, S. M., Saghai Maroof, M. A., Scoles, G. J., Hoffman, D., Dahleen, L. S., Kilian, A., Chen, F., Biyashev, R. M., Kudrna, D. A., Steffenson, B. J. Regions of the genome that affect agronomic performance in two-row barley. *Crop Sci.*, **36**:1053-1062 (1996).
- 20 Torbert, K. A., Rines, H. W., Somers, D. A. Use of paramomycin as a selective agent for oat transformation. *Plant Cell Reports*, **14**:635-640 (1995).
- Torbert, K. A., Rines, H. W., Somers, D. A. Transformation of oat using mature embryo-derived tissue cultures. *Crop Sci.*, **38**:226-231 (1998).
- Turnham, E., Northcote, D. N. Changes in the activity of acetyl-CoA carboxylase during rapeseed formation. *Biochem. J.*, **212**:223-229 (1983).
- 30 Van Dee, K. L., Somers, D. A., Wyse, D. L. RFLP mapping of *Acc1* in *Zea mays* L. *Agronomy Abstracts*, p.198 (1992).

Van Dee, K.L. Evidence for two independently segregating loci encoding acetyl-CoA carboxylase in *Zea mays*, M.S. Thesis, Univ. of Minnesota, 62 p. (1994).

- VanDeynze, A. E., Dubcovsky, J., Gill, K. S., Nelson, J. C., Sorrells, M. E., Dvorak, J., Gill, B. S., Lagudah, E. S., McCouch, S. R., Appels, R. Molecular-genetic maps for group 1 chromosomes of Triticeae species and their relation to chromosomes in rice and oat. *Genome*, 38:45-59 (1995).

- Webster, F. H. Oat utilization: past, present and future. In: Webster F. H. (ed.) Oat chemistry and technology. Assoc. Cereal Chem., St. Paul, Minnesota, pp. 413-427 (1986).

Welch, R. W. The chemical composition of oats. In: The Oat Crop, Welch, R.W., ed. Chapman and Hall, pp. 279-320 (1995).

- Youngs, V. L. Oat lipids and lipid in related enzymes. In: Oats: Chemistry and Technology, Webster, F. H., ed. American Association of Cereal Chemists, Inc. pp. 205-226 (1986).

Youngs, V. L. Oat lipids and lipid-related enzymes. In: Webster, F. H. (ed.) Oat chemistry and technology. Assoc. Cereal Chem., St. Paul, Minnesota, pp. 205-226 (1986).

- Youngs, V. L., Peterson, D. M., Brown, C. M. Oats. In: Pomeranz, Y. (ed.) Advances in cereal science and technology. Vol. 5. Am. Assoc. Cereal Chem., St. Paul, Minnesota, pp. 49-105 (1982).

Yuzo, M., Takuji, S. Genbank accession #D24786 (rice) (1993).

- The complete disclosure of all patents, patent documents and publications cited herein are incorporated herein by reference as if individually incorporated. The foregoing detailed description and examples have been given for clarity of understanding only. No unnecessary limitations are to be understood therefrom. The invention is not limited to the exact details shown and described for variations obvious to one skilled in the art will be included within the invention defined by the claims.

WHAT IS CLAIMED IS:

1. An isolated and purified DNA molecule comprising a DNA segment encoding an oat acetyl CoA carboxylase, or a fragment thereof.
5
2. The DNA molecule of claim 1 which comprises a DNA segment comprising SEQ ID NO:23.
3. The DNA molecule of claim 1 or 2 further comprising a DNA molecule encoding an amino terminal chloroplast transit peptide, wherein the chloroplast transit peptide is operably linked to the DNA encoding the oat acetyl CoA carboxylase.
10
4. The DNA molecule of claim 1 which encodes a polypeptide comprising SEQ ID NO:24.
15
5. A method of imparting cyclohexanedione or aryloxyphenoxypropanoic acid herbicide tolerance to a plant tissue comprising:
 - (a) introducing an expression cassette comprising a chimeric DNA molecule encoding an oat acetyl CoA carboxylase comprising SEQ ID NO:24, operably linked to a promoter functional in a plant cell into cells of a susceptible plant so as to yield transformed plant cells;
20
 - (b) regenerating the transformed plant cells to provide a differentiated transformed plant; and
25
 - (c) expressing the chimeric DNA molecule encoding the oat acetyl CoA carboxylase in the cells of the differentiated plant in an amount effective to render the plant tolerant to levels of a cyclohexanedione or an aryloxyphenoxypropanoic acid herbicide which are toxic to a susceptible plant which does not comprise the expression cassette.
30

6. The method of claim 5 wherein the chimeric DNA molecule encodes an acetyl CoA carboxylase, the activity of which is not inhibited by an amount of cyclohexanedione or aryloxyphenoxypropanoic acid herbicide that inhibits the activity of an acetyl CoA carboxylase which is sensitive to inhibition by a cyclohexanedione or aryloxyphenoxypropanoic acid herbicide.
7. The method of claim 5 wherein the chimeric DNA molecule is expressed in the cells at about a 2' to about a 20-fold amount over that of a cyclohexanedione or aryloxyphenoxypropanoic acid herbicide-sensitive acetyl CoA carboxylase.
8. The method of claim 5 wherein the chimeric DNA molecule comprises a DNA sequence comprising SEQ ID NO:23.
9. A transformed plant prepared by the method of claim 5.
10. A transformed seed of the transformed plant of claim 9.
11. A transformed monocot plant, which plant is tolerant to herbicides comprising:
 - (a) an acetyl CoA carboxylase gene that encodes an acetyl CoA carboxylase the activity of which is inhibited by a cyclohexanedione or aryloxyphenoxypropanoic acid herbicide, and
 - (b) a recombinant DNA molecule encoding an oat acetyl CoA carboxylase operably linked to a promoter functional in a plant, wherein the recombinant DNA molecule is expressed in the plant in an amount effective to confer tolerance to the levels of a cyclohexanedione herbicide or an aryloxyphenoxypropanoic acid herbicide which inhibit the activity of the acetyl CoA carboxylase of step (a).

12. A method for altering the oil content in a plant comprising:
- (a) introducing an expression cassette comprising a recombinant DNA molecule encoding an oat acetyl CoA carboxylase operably linked to a promoter functional in a plant cell into the cells of a plant so as to yield transformed plant cells; and
- (b) regenerating the transformed plant cells to provide a differentiated transformed plant, wherein the plant expresses the DNA molecule encoding the oat acetyl CoA carboxylase in cells of said plant so as to alter the oil content of the differentiated transformed plant cells relative to the oil content of cells of a corresponding untransformed plant.
13. The method according to claim 12, wherein the recombinant DNA molecule is expressed at about a 2- to about a 20-fold increase over the level of the native acetyl CoA carboxylase.
14. The method according to claim 13 wherein the oil content of the plant cells is increased about 1.2- to about 20-fold.
15. A transformed plant having an altered oil content in the plant cells comprising: a recombinant DNA molecule encoding an oat acetyl CoA carboxylase, wherein the recombinant DNA molecule is expressed in the cells of the plant in an amount of acetyl CoA carboxylase effective to alter the oil content of the plant cells relative to the oil content in the cells of a corresponding untransformed plant.
16. The transformed plant of claim 15 wherein the transformed plant has an increase in oil content of about 1.2- to about 20-fold in its leaves, seeds, or fruit above that present in a corresponding untransformed plant.
17. A transformed plant prepared by the method of claim 12.
18. A seed of the transformed plant of claim 17.

19. The seed of claim 18 wherein the oil content is increased in the endosperm.
20. A method of introducing an exogenous oat acetyl CoA carboxylase gene
5 into a host cell comprising:
(a) introducing to host cells *in vitro* an expression cassette comprising a DNA molecule encoding an oat acetyl CoA carboxylase operably linked to a promoter functional in the host cell so as to yield transformed cells; and
10 (b) identifying transformed host cells which express the DNA molecule.
21. The method of claim 20 wherein the host cells are plant cells.
- 15 22. The method of claim 21 wherein the plant cells can be regenerated into a differentiated plant.
23. The method of claim 21 wherein the plant cells are monocot cells.
- 20 24. The method of claim 21 wherein the plant cells are dicot cells.
25. The method of claim 12, wherein the DNA molecule encoding the oat acetyl CoA carboxylase is introduced into plant cells by a method selected from the group consisting of electroporation, microinjection,
25 protoplast transformation, microprojectile bombardment, and liposomal encapsulation, *Agrobacterium*-mediated transformation.
26. A method for altering the oil content in a plant comprising:
30 (a) introducing an expression cassette comprising a recombinant DNA molecule encoding an antisense oat acetyl CoA carboxylase transcript operably linked to a promoter functional in a plant cell into the cells of a plant so as to yield transformed plant cells; and

- (b) regenerating the transformed plant cells to provide a differentiated transformed plant, wherein the transcript is expressed in the cells of the differentiated transformed plant so as to alter the oil content of the plant cells relative to the cells of a corresponding untransformed plant.
- 5
27. A transformed plant having an altered oil content in the plant cells comprising: a recombinant DNA molecule encoding an antisense oat acetyl CoA carboxylase transcript wherein the transcript is expressed in the cells of the plant so as to alter the oil content of the plant cells relative to the cells of a corresponding untransformed plant.
- 10
28. The method of claim 12 or 26 wherein the recombinant DNA molecule further comprises a DNA segment encoding a chloroplast transit peptide which is operably linked to the DNA molecule encoding the acetyl CoA carboxylase.
- 15
29. The transformed plant of claim 15 or 27 wherein the recombinant DNA molecule further comprises a DNA segment encoding a chloroplast transit peptide which is operably linked to the DNA molecule encoding the acetyl CoA carboxylase.
- 20
30. The method of claim 12 or 26 wherein the recombinant DNA molecule further comprises a DNA segment comprising a endosperm-specific promoter.
- 25
31. The plant of claim 15 or 27 wherein the recombinant DNA molecule further comprises a DNA segment comprising a endosperm-specific promoter.
- 30
32. The method of claim 12 or 26 wherein the cells are monocot plant cells.

33. The method of claim 32 wherein the cells are wheat, rice, maize, barley, millet, sorghum, sugarcane, rye, oat or timothy cells.
34. The method of claim 12 or 26 wherein the cells are dicot plant cells.
- 5 35. The method of claim 34 wherein the cells are soybean, rape, sunflower, tobacco, Arabidopsis, petunia, pea, Canola, bean, tomato, bean, potato, lettuce, spinach, alfalfa, cotton or carrot.
- 10 36. The method of claim 12, 20, or 25 wherein the DNA molecule encodes a polypeptide comprising SEQ ID NO:24.
37. The plant of claim 11, 15, or 27 wherein the DNA molecule encodes a polypeptide comprising SEQ ID NO:24.
- 15 38. An expression cassette comprising an endosperm-specific transcriptional control region operably linked to a DNA encoding a plant ACCase, or a biologically active subunit thereof.
- 20 39. The expression cassette of claim 38 wherein the endosperm-specific transcriptional control region comprises Dy10.
40. The plant of claim 15 which has increased lipid content, lipid biosynthesis or lipid deposition in groat.
- 25 41. The plant of claim 27 which has decreased lipid content, lipid biosynthesis or lipid deposition in groat.
42. The plant of claim 15 which has increased storage of seed lipid in endosperm.
- 30 43. The plant of claim 27 which has decreased storage of seed lipid in endosperm.

44. A method to identify an oat plant having altered groat oil content, comprising:
- 5 (a) contacting a probe comprising at least a portion of a nucleic acid sequence encoding oat acetyl CoA carboxylase with a nucleic acid sample from a plant in an amount sufficient to form complexes, wherein the sample comprises genomic nucleic acid; and
- (b) detecting or determining the amount of complex formation.
- 10 45. A method to identify a plant having altered oil content, β -glucan content, seed weight, starch content or protein content, comprising:
- (a) contacting a probe comprising at least a portion of a nucleic acid sequence encoding oat acetyl CoA carboxylase with a sample from a plant in an amount sufficient to form complexes, wherein
- 15 the sample comprises genomic nucleic acid; and
- (b) detecting or determining the amount of complex formation.
46. The method of claim 45 wherein the plant is an oat, maize or wheat plant.
- 20 47. A method for marker-assisted selection of plants having desired properties, comprising:
- (a) contacting a probe comprising at least a portion of a nucleic acid sequence encoding oat acetyl CoA carboxylase with a sample from a plant in an amount sufficient to form complexes, wherein
- 25 the sample comprises genomic nucleic acid; and
- (b) detecting or determining the amount of complex formation.
48. An isolated and purified nucleic acid segment associated with altered oil content or composition in monocots.
- 30 49. A method to map a restriction fragment length polymorphism in plants, comprising:

- (a) contacting a probe comprising at least a portion of a nucleic acid sequence encoding oat acetyl CoA carboxylase with a sample from a plant in an amount sufficient to form complexes, wherein the sample comprises genomic nucleic acid; and
 - 5 (b) detecting or determining the amount of complex formation.
- 50. The method of claim 44, 45, 47 or 49 wherein the genomic nucleic acid is from oat, maize, rice or wheat.

FATTY ACID BIOSYNTHESIS




FIG. 1




FIG. 2




FIG. 3

4/37




FIG. 4




FIG. 5

6/37




FIG. 6




FIG. 7




FIG. 8




FIG. 9

AGA GAT GAA GCT CGC ATG CCA ATG CGC CAC ACA TTC CTC TGG TTG GAT
GAC AAG AGT TGT TAT GAA GAA GAG CAG ATT CTC CGG CAT GTG GAG CCT
CCC CTC TCT ACA CTT CTT GAA TTG GAT AAG TTG AAG GTG AAA GGA TAC
AAT GAA ATG AAG TAT ACT CCT TCG CGT GAC CGC CAA TGG CAT ATC TAC
ACA CTA AGA AAT ACT GAA AAC CCC AAA ATG TTG CAT AGG GTG TTT TTC
CGA ACT ATT GTC AGG CAA CCC AAT GCA GGC AAC AAG TTT AGA TCG GCT
CAG ATC AGC GAC GCN AAG GTA GGA TGT CCC GAA GAA TCT CTT TCA TTT
ACA TCA AAT AGC ATC TTA AGA TCA TTG ATG ACT GCT ATT GAA GAA TTA
GAG CTT CAT GCA ATT AGG ACA GGT CAT TCT CAC ATG TAT TTG TGC ATA
CTG AAA GAG CAA AAG CTT CTT GAC CTC ATT CCA TTT TCA GGG AGT ACA
ATT GTT GAT GTT GGC CAA GAT GAA GCT ACC GCT TGT TCA CTT TTA AAA
TCA ATG GCT TTG AAG ATA CAT GAG CTT GTT GGT GCA AGG ATG CAT CAT
CTG TCT GTA TGC CAG TGG GAG GTG AAA CTC AAG TTG GAC TGT GAT GGC
CCT GCA AGT GGT ACC TGG AGA GTT GTA ACT ACA AAT GTT ACT GGT CAC
ACC TGC ACC ATT GAT ATA TAC CGA GAA GTG GAG GAA ATA GAA TCA CAG
AAG TTA GTG TAC CAT TCA GCC AGT TCG TCA GCT GGA CCA TTG CAT GGT
GTT GCA CTG AAT AAT CCA TAT CAA CCT TTG AGT GTG ATT GAT CTA AAG
CGC TGC TCT GCT AGG AAC AAC AGA ACA ACA TAT TGC TAT GAT TTT CCG
CTG GCC TTT GAA ACT GCA CTG CAG AAG TCA TGG CAG TCC AAT GGC TCT
ACT GTT TCT GAA GGC AAT GAA AAT AGT AAA TCC TAC GTG AAG GCA ACT
GAG CTA GTG TTT GCT GAA AAA CAT GGG TCC TGG GGC ACT CCT ATA ATT
CCG ATG GAA CGC CCT GCT GGG CTC AAC GAC ATT GGT ATG GTC GCT TGG
ATC ATG GAG ATG TCA ACA CCT GAA TTT CCC AAT GGC AGG CAG ATT ATT
GTT GTA GCA AAT GAT ATC ACT TTC AGA GCT GGA TCA TTT GGC CCA AGG
GAA GAT GCA TTT TTT GAA ACT GTC ACT AAC CTG GCT TGC GAA AGG AAA
CTT CCT CTT ATA TAC TTG GCA GCA AAC TCT GGT GCT AGG ATT GGC ATA
GCT GAT GAA GTA AAA TCT TGC TTC CGT GTT GGA TGG TCT GAC GAA GGC
AGT CCT GAA CGA GGG TTT CAG TAC ATC TAT CTG ACT GAA GAA GAC TAT
GCT CGC ATT AGC TCT TCT GTT ATA GCA CAT AAG CTG GAG CTA GAT AGT
GGT GAA ATT AGG TGG ATT ATT GAC TCT GTT GTG GGC AAG GAG GAT GGG
CTT GGT GTC GAG AAC ATA CAT GGA AGT GCT GCT ATT GCC AGT GCT TAT
TCT AGG GCA TAT GAG GAG ACA TTT ACA CTT ACA TTT GTG ACT GGG CGG
ACT GTA GGA ATA GGA GCT TAT CTT GCT CGA CTT GGT ATA CGG TGC ATA
CAG CGT CTT GAC CAG CCT ATT ATT TTA ACA GGG TTT TCT GCC CTG AAC
AAG CTC CTT GGG CGG GAA GTG TAC AGC TCC CAC ATG CAG CTT GGT GGT
CCT AAG ATC ATG GCG ACC AAT GGT GTT GTC CAC CTC ACT GTT CCA GAT

FIG. 10

GTC CTT GAA GGT GTT TCC AAT ATA TTG AGG TGG CTC AGC TAT GTT CCT
GCA AAC ATT GGT GGA CCT CTT CCT ATT ACC AAA CCT CTG GAC CCT CCA
GAC AGA CCT GTT GCT TAC ATC CCT GAG AAC ACA TGC GAT CCA CGT GCA
GCT ATC TGT GGT GTA GAT GAC AGC CAA GGG AAA TGG TTG GGT GGT ATG
TTT GAC AAA GAC AGC TTT GTG GAG ACA TTT GAA GGA TGG GCA AAA ACA
GTG GTT ACT GGC AGA GCA AAG CTT GGA GGA ATT




FIG. 11

ACCase A cloning strategy and position of start and stop codons




FIG. 12

1 GGTCTTCAAT TGTGCTGTCT GGGCCACGGA ACGACAATGT CACAGCTTGG
51 ATTAGCCGCA GCTGCCTCAA AGGCCTTGCC ACTACTCCCT AATCGCCAGA
101 GAAGTTCAGC TGGGACTACA TTCTCATCAT CTTCATTATC GAGGCCCTTA
151 AACAGAAGGA AAAGCCATAC TCGTTCACTC CGTGATGGCG GAGATGGGGT
201 ATCAGATGCC AAAAAGCACA GCCAGTCTGT TCGTCAAGGT CTTGCTGGCA
251 TTATCGACCT CCCAAGTGAG GCACCTTCCG AAGTGGATAT TTCACATGGA
301 TCTGAGGATC CTAGGGGGCC AACAGATTCT TATCAAATGA ATGGGATTAT
351 CAATGAAACA CATAATGGAA GACATGCCTC AGTGTCCAAG GTTGTGTAAT
401 TTTGTGCGGC ACTAGGTGGC AAAACACCAA TTCACAGTAT ATTAGTGGCC
451 AACAATGGAA TGGCAGCAGC AAAATTTATG AGGAGTGTCC GGACATGGGC
501 TAATGATACT TTTGGATCTG AGAAGGCAAT TCAACTCATA GCTATGGCAA
551 CTCCGGAAGA CATGAGGATA AATGCAGAAC ACATTAGAAT TGCTGACCAA
601 TTCGTAGAGG TGCCTGGTGG AACAAACAAT AATAACTACG CCAATGTTCA
651 ACTCATAGTG GGGATGGCAC AAAAACTAGG TGTTTCTGCT GTTTGGCCTG
701 GTTGGGGTCA TGCTTCTGAG AATCCTGAAC TGCCAGATGC ATTGACCGCA
751 AAAGGGATCG TTTTCTTGG CCCACCTGCA TCATCAATGA ATGCTTTGGG
801 AGATAAGGTC GGCTCAGCTC TCATTGCTCA AGCAGCCGGG GTCCCAACTC
851 TTGCTTGGAG TGGATCACAT GTTGAAGTTC CATTAGAGTG CTGCTTAGAC
901 GCGATACCTG AGGAGATGTA TAGAAAAGCT TCGGTTACTA CCACAGAGGA
951 AGCAGTTGCA AGTTGTCAAG TGGTTGGTTA TCCTGCCATG ATTAAGGCAT
1001 CCTGGGGAGG TGGTGGTAAA GGAATAAGAA AGGTTCATAA TGATGATGAG
1051 GTTAGAGCGC TGTTTAAGCA AGTACAAGGT GAAGTCCCTG GCTCCCAAT
1101 ATTTGTCATG AGGCTTGCAT CCCAGAGTCG GCATCTTGAA GTTCAGTTGC
1151 TTTGTGATCA ATATGGTAAT GTAGCAGCAC TTCACAGTCG TGATTGCAGT
1201 GTGCAACGGC GACACCAGAA GATTATTGAA GAAGGTCCAG TTAGTGTGTC
1251 TCCTCGTGAG ACAGTTAAAG CACTTGAGCA GGCAGCAAGG AGGCTTGCTA
1301 AGGCTGTGGG TTATGTTGGT GCTGCTACTG TTGAGTATCT TTACAGCATG
1351 GAAACTGGAG ACTACTATTT TCTGGAACCT AATCCCCGAC TACAGGTTGA
1401 GCATCCAGTC ACTGAGTGGA TAGCTGAAGT GAATCTGCCT GCAGCTCAAG
1451 TTGCTGTTGG AATGGGCATA CCTCTTTGGC AGATTCCAGA AATCAGACGT
1501 TTCTATGGAA TGGACTATGG AGGAGGGTAT GACATTTGGA GGAAAACAGC
1551 AGCTCTTGCT ACACCATTTA ATTTTGATGA AGTAGATTCT CAATGGCCAA
1601 AGGGCCATTG TGTAGCAGTT AGAATTACTA GTGAGGACCC AGATGATGGT
1651 TTCAAACCTA CTGGTGGGAA AGTGAAGGAG ATAAGTTTTA AAAGCAAGCC
1701 TAATGTTTGG GCCTACTTCT CAGTAAAGTC TGGTGGAGGC ATTCATGAAT
1751 TTGCTGATTC TCAGTTTGGG CATGCTTTTG CATATGGACT CTCTAGACCA

FIG. 13

1801 GCAGCTATAA CAAACATGTC TCTTGCATTA AAAGAGATTC AGATTCGTGG
1851 AGAAATTCAT TCAAATGTTG ATTACACAGT TGACCTCTTA AACGCTTCAG
1901 ACTTCAGAGA AAACAAGATC CACACTGGTT GGCTGGATAC AAGAATAGCT
1951 ATGCGTGTTT AAGCTGAGAG GCCCCCATGG TATATCTCAG TGGTTGGAGG
2001 TGCTTTATAT AAAACAGTAA CCACCAATGC AGCCACTGTT TCTGAATATG
2051 TTAGTTATCT CACCAAGGGC CATATTCCAC CAAAGCATAT ATCCCTTGTC
2101 AATTCTACAG TTAATTTGAA TATAGAAGGG AGCAAATACA CAATTGAAAC
2151 TGTAAGGACT GGACATGGTA GCTACAGGTT GAGAATGAAT GATTCAACAG
2201 TTGAAGCGAA TGTACAATCT TTATGTGATG GTGGCCTCTT AATGCAGTTG
2251 GATGGAAACA GCCATGTAAT TTATGCAGAA GAAGAAGCTG GTGGTACACG
2301 GCTTCAGATT GATGGAAAGA CATGTTTATT GCAGAATGAC CATGATCCAT
2351 CGAAGTTATT AGCTGAGACA CCCTGCAAAC TTCTTCGTTT CTGTTGCT
2401 GATGGTGCTC ATGTTGATGC GGATGTACCA TACGCGGAAG TTGAGGTTAT
2451 GAAGATGTGC ATGCCTCTCT TGTCACCTGC TTCTGGTGTC ATTCATTGTA
2501 TGATGTCTGA GGGCCAGGCA TTGCAGGCTG GTGATCTTAT AGCAAGGTTG
2551 GATCTTGATG ACCCTTCTGC TGTGAAAAGA GCTGAGCCAT TTGATGGAAT
2601 ATTTCCACAA ATGGAGCTCC CTGTTGCTGT CTCTAGTCAA GTACACAAAA
2651 GATATGCTGC AAGTTTGAAT GCTGCTCGAA TGGTCCTTGC AGGATATGAG
2701 CACAATATTA ATGAAGTCGT TCAAGATTTG GTATGCTGCC TGGACAACCC
2751 TGAGCTTCCT TTCCTACAGT GGGATGAACT TATGTCTGTT CTAGCAACGA
2801 GGCTTCCAAG AAATCTCAAG AGTGAGTTAG AGGATAAATA CAAGGAATAC
2851 AAGTTGAATT TTTACCATGG AAAAAACGAG GACTTTCCAT CCAAGTTGCT
2901 AAGAGACATC ATTGAGGAAA ATCTTTCTTA TGGTTCAGAG AAGGAAAAGG
2951 CTACAAATGA GAGGCTTGTT GAGCCTCTTA TGAACCTACT GAAGTCATAT
3001 GAGGGTGGGA GAGAGAGCCA TGCACATTTT GTTGTCAAGT CTCTTTTCGA
3051 GGAGTATCTT ACAGTGGAAG AACTTTTTAG TGATGGCATT CAGTCTGACG
3101 TGATTGAAAC ATTGCGGCAT CAGCACAGTA AAGACCTGCA GAAGGTTGTA
3151 GACATTGTGT TGTCTACCA GGGTGTGAGG AACAAAGCTA AGCTTGTAAC
3201 GGCACCTATG GAAAAGCTGG TTTATCCAAA TCCTGGTGGT TACAGGGATC
3251 TGTTAGTTCG CTTTTCTTCC CTCAATCATA AAAGATATTA TAAGTTGGCC
3301 CTTAAAGCAA GTGAACTTCT TGAACAAACC AAATAAGTG AACTCCGTGC
3351 AAGCGTTGCA AGAAGCCTTT CGGATCTGGG GATGEATAAG GGAGAAATGA
3401 GTATTAAGGA TAACATGGAA GATTTAGTCT CTGCCCCATT ACCTGTTGAA
3451 GATGCTCTGA TTTCTTGTG TGATTACAGT GATCGAACTG TTCAGCAGAA
3501 AGTGATTGAG ACATACATAT CACGATTGTA CCAGCCTCAT CTTGTAAAGG
3551 ATAGCATCCA AATGAAATTC AAGGAATCTG GTGCTATTAC TTTTGGGAA

FIG. 13 (continued)

3601 TTTTATGAAG GGCATGTTGA TACTAGAAAT GGACATGGGG CTATTATTGG
3651 TGGGAAGCGA TGGGGTGCCA TGGTCGTTCT CAAATCACTT GAATCTGCGT
3701 CAACAGCCAT TGTGGCTGCA TTAAAGGATT CGGCACAGTT CAACAGCTCT
3751 GAGGGCAACA TGATGCACAT TGCATTATTG AGTGCTGAAA ATGAAAGTAA
3801 TATAAGTGGG ATAAGCAGTG ATGATCAAGC TCAACATAAG ATGGAAAAGC
3851 TTAGCAAGAT ACTGAAGGAT ACTAGCGTTG CAAGTGATCT CCAAGCTGCT
3901 GGT TTGAAGG TTATAAGTTG CATTGTTCAA AGAGATGAAG CTCGCATGCC
3951 AATGCGCCAC ACATTCCTCT GGT TGGATGA CAAGAGTTGT TATGAAGAAG
4001 AGCAGATTCT CCGGCATGTG GAGCCTCCCC TCTCTACACT TCTTGAATTG
4051 GATAAGTTGA AGGTGAAAGG ATACAATGAA ATGAAGTATA CTCCTTCGCG
4101 TGACCGCCAA TGGCATATCT ACACACTAAG AAATACTGAA AACCCCAAAA
4151 TGTTGCATAG GGTGTTTTTC CGAACTATTG TCAGGCAACC CAATGCAGGC
4201 AACAAGTTTA GATCGGCTCA GATCAGCGAC GCTGAGGTAG GATGTCCCGA
4251 AGAATCTCTT TCATTTACAT CAAATAGCAT CTTAAGATCA TTGATGACTG
4301 CTATTGAAGA ATTAGAGCTT CATGCAATTA GGACAGGTCA TTCTCACATG
4351 TATTTGTGCA TACTGAAAGA GCAAAAGCTT CTTGACCTCA TTCCATTTTC
4401 AGGGAGTACA ATTGTTGATG TTGGCCAAGA TGAAGCTACC GCTTGTTTAC
4451 TTTTAAAATC AATGGCTTTG AAGATACATG AGCTTGTTGG TGCAAGGATG
4501 CATCATCTGT CTGTATGCCA GTGGGAGGTG AAACCTCAAGT TGGACTGTGA
4551 TGGCCCTGCA AGTGGTACCT GGAGAGTTGT AACTACAAAT GTTACTGGTC
4601 ACACCTGCAC CATTGATATA TACCGAGAAG TGGAGGAAAT AGAATCACAG
4651 AAGTTAGTGT ACCATTCAGC CAGTTCGTCA GCTGGACCAT TGCATGGTGT
4701 TGCATGAAT AATCCATATC AACCTTTGAG TGTGATTGAT CTAAAGCGCT
4751 GCTCTGCTAG GAACAACAGA ACAACATATT GCTATGATTT TCCGCTGGCC
4801 TTTGAACTG CACTGCAGAA GTCATGGCAG TCCAATGGCT CTACTGTTTC
4851 TGAAGGCAAT GAAAATAGTA AATCCTACGT GAAGGCAACT GAGCTAGTGT
4901 TTGCTGAAAA ACATGGGTCC TGGGGCACTC CTATAATTCC GATGGAACGC
4951 CCTGCTGGGC TCAACGACAT TGGTATGGTC GCTTGGATCA TGGAGATGTC
5001 AACACCTGAA TTTCCCAATG GCAGGCAGAT TATTGTTGTA GCAAATGATA
5051 TCACTTTCAG AGCTGGATCA TTTGGCCCAA GGAAGATGC ATTTTTTGAA
5101 ACTGTCAC TAACCTGGCTTG CGAAAGGAAA CTTCTCTTA TATACTTGGC
5151 AGCAAACCTCT GGTGCTAGGA TTGGCATAGC TGATGAAGTA AAATCTTGCT
5201 TCCGTGTTGG ATGGTCTGAC GAAGGCAGTC CTGAACGAGG GTTTCAGTAC
5251 ATCTATCTGA CTGAAGAAGA CTATGCTCGC ATTAGCTCTT CTGTTATAGC
5301 ACATAAGCTG GAGCTAGATA GTGGTGAAAT TAGGTGGATT ATTGACTCTG
5351 TTGTGGGCAA GGAGGATGGG CTTGGTGTGCG AGAACATACA TGGAAGTGCT

FIG. 13 (continued)

5401 GCTATTGCCA GTGCTTATTC TAGGGCATAT GAGGAGACAT TTACACTTAC
5451 ATTTGTGACT GGGCGGACTG TAGGAATAGG AGCTTATCTT GCTCGACTTG
5501 GTATACGGTG CATAACGCGT CTTGACCAGC CTATTATTTT AACAGGGTTT
5551 TCTGCCCTGA ACAAGCTCCT TGGGCGGGAA GTGTACAGCT CCCACATGCA
5601 GCTTGGTGGT CCTAAGATCA TGGCGACCAA TGGTGTGTGC CACCTCACTG
5651 TTCCAGATGT CCTTGAAGGT GTTTCCAATA TATTGAGGTG GCTCAGCTAT
5701 GTTCCTGCAA ACATTGGTGG ACCTCTTCCT ATTACCAAAC CTCTGGACCC
5751 TCCAGACAGA CCTGTTGCTT ACATCCCTGA GAACACATGC GATCCACGTG
5801 CAGCTATCTG TGGTGTAGAT GACAGCCAAG GGAAATGGTT GGGTGGTATG
5851 TTTGACAAAG ACAGCTTTGT GGAGACATTT GAAGGATGGG CAAAAACAGT
5901 GGTTACTGGC AGAGCAAAGC TTGGAGGAAT TCCTGTGGGC GTCATAGCTG
5951 TGGAGACACA GACCATGATG CAGATCATCC CTGCTGATCC AGGTCAGCTT
6001 GATTCCCATG AGCGATCTGT CCCTCGTGCT GGACAAGTGT GGTTCCCAAG
6051 TTCTGCAACC AAGACCGCTC AGGCATTATT AGACTTCAAC CGTGAAGGAT
6101 TGCCTCTGTT CATCCTGGCT AATTGGAGAG GCTTCTCTGG TGGACAAAGA
6151 GATCTCTTTG AAGGAATTCT TCAGGCTGGG TCAACAATTG TCGAGAACCT
6201 TAGGACATAT AATCAGCCTG CTTTTGTGTA CATTCCCTATG GCTGGAGAGC
6251 TTCGTGGAGG AGCTTGGGTT GTGGTCGATA GCAAAATAAA TCCAGACCGC
6301 ATTGAGTGTT ATGCTGAAAG GACTGCCAAA GGTAATGTTT TCGAACCTCA
6351 AGGGTTAATT GAAATCAAGT TCAGGTCAGA GGAACCTCAA GACTGTATGG
6401 GTAGGCTTGA CCCAGAGTTG ATAAATCTGA AAGCAAACT CCAAGATGTA
6451 AATCATGGAA ATGGAAGTCT ACCAGACATA GAAGGGATTC GGAAGAGTAT
6501 AGAAGCACGT ACGAAACAGT TGCTGCCTTT ATATACCCAG ATTGCAATAC
6551 GGTTTGCTGA ATTGCATGAT ACTTCCCTAA GAATGGCAGC TAAAGGTGTG
6601 ATTAAGAAAG TTGTAGACTG GGAAGAATCA CGCTCGTTCT TCTATAAAAG
6651 GCTACGGAGG AGGATCGCAG AAGATGTTCT TGCAAAAGAA ATAAGGCAGA
6701 TAGTCGGTGA TAAATTTACG CACCAATTAG CAATGGAGCT CATCAAGGAA
6751 TGGTACCTTG CTTCTCAGGC CACAACAGGA AGCACTGGAT GGGATGACGA
6801 TGATGCTTTT GTTGCCCTGGA AGGACAGTCC TGAAAACCTAC AAGGGGCATA
6851 TCCAAAAGCT TAGGGCTCAA AAAGTGTCTC ATTCGCTCTC TGATCTTGCT
6901 GACTCCAGTT CAGATCTGCA AGCATTCTCG CAGGGTCTTT CTACGCTATT
6951 AGATAAGATG GATCCCTCTC AGAGAGCGAA GTTTGTTCAG GAAGTCAAGA
7001 AGGTCCTTGA TTGATGATAC CAACACATCC AACACAATGT GTGCATGTCA
7051 CATCTTTTTG TTCTAGTACA TACATAGAAG GATATTGCTT GGTCTTGATT
7101 GATCATGTCT GATTTAAGTC GACTATTATT TCTTGGAAAT TTCTTTTGGG
7151 CCTGGTGCTA TGGTTGATGG ATGTATATTG GATATGTGCG TTCTGCCAGG

FIG. 13 (continued)

17/37

7201 TGTAAGCACA AAGGTTTAGA CARAMMRARA RCAAGAGCGA GTGAACCTGT
7251 TCTGGTTTTG CAGTGGTTCA GTAAGGCAGA AAGTTGTTAA ACCGTAGTTC
7301 TGAGATGTAT TACCAGTGNC GCCATGCTGT ACTTTTAGGG TGTATAATGC
7351 GGATACAAAT AAACAATTTA GCGGTTTCATT AAAGTTTGAA CTCAAATAAC
7401 ATGTTCTTTG TAAGCATATG TACCGTACCT CTACGTGAAA TAAAGTTGTT
7451 GAATTAGCAT TCGAAAAAAA

FIG. 13 (continued)

1 MSQGLGAAAA SKALPLLPNR QRSSAGTTFS SSSLSRPLNR RKSHTRSLRD
51 GGDGVSDAKK HSQSVRQGLA GIIDL PSEAP SEVDISHGSE DPRGPTDSYQ
101 MNGIINETHN GRHASVSKVV EFCAALGGKT PIHSILVANN GMAAAKFMRS
151 VRTWANDTFG SEKAIQLIAM ATPEDMRINA EHIRIADQFV EVPGGTNNNN
201 YANVQLIVGM AQKLGVS AVW PGWGHASENP ELPDALTAKG IVFLGPPASS
251 MNALGDKVGS ALIAQAAGVP TLAWSGSHVE VPLECCLD AI PEEMYRKACV
301 TTTEEAVASC QVVGYPAMIK ASWGGGGKGI RKVHNDDEVR ALFKQVQGEV
351 PGSPIFVMRL ASQSRHLEVQ LLC DQYGNVA ALHSRDCSVQ RRHQKIIEEG
401 PVTVAPRETV KALEQAARRL AKAVGYVGAA TVEYLYSMET GDYYFLELNP
451 RLQVEHPVTE WIAEVNLPAA QVAVGMGIPL WQIPEIRRFY GMDYGGGYDI
501 WRKTAALATP FNFDEVDSQW PKGHC VAVRI TSEDPDDGFK PTGGKVKEIS
551 FKSKPNVWAY FSVKSGGGIH EFADSQFGHA FAYGLSRPAA ITNMSLALKE
601 IQIRGEIHSN VDYTVDLLNA SDFRENKIHT GWLDTRIAMR VQAERPPWYI
651 SVVGGALYKT VTTNAATVSE YVSYLTKGHI PPKHISLVNS TVNLNIEGSK
701 YTIETVRTGH GSYRLRMNDS TVEANVQSLC DGGLLMQLDG NSHVIYAEED
751 AGGTRLQIDG KTCLLQNDHD PSKLLAETPC KLLRFLVADG AHVDADVPIA
801 EVEVMKMCMP LLSPASGVII CMMSEGQALQ AGDLIARLDL DDPSAVKRAE
851 PFDGIFPQME LPVAVSSQVH KRYAASLNAA RMVLAGYEHN INEVVQDLVC
901 CLDNPELPFL QWDELMSVLA TRLPRNLKSE LEDKYKEYKL N FYH GKNEDE
951 PSKLLRDIIE ENLSYGSEKE KATNERLVEP LMNLLKSYEG GRESHAHFVV
1001 KSLFEEYLTV EELFSDGIQS DVIETLRHQH SKDLQKVVDI VLSHQGVRNK
1051 AKLVTALMEK LVYPNPGGYR DLLVRFSSLN HKRYYKLALK ASELLEQTKL
1101 SELRASVARS LSDLGMHKGE MSIKDNMEDL VSAPLPVEDA LISLFDYS DR
1151 TVQQKVIETY ISRLYQPHLV KDSIQMKFKE SGAITFWEFY EGHVDTRNGH
1201 GAIIGGKRWG AMVVLKSLES ASTAIVAALK DSAQFNSSEG NMMHIALLSA
1251 ENESNISGIS SDDQAQHKME KLSKILKDS VASDLQAAGL KVISCIVQRD
1301 EARMPPMRHTF LWLDDKSCYE EEQILRHVEP PLSTLLELDK LKVKGYNEMK
1351 YTPSRDRQWH IYTLRNTENP KMLHRVFFRT IVRQPNAGNK FRSAQISDAE
1401 VGCPEESLSF TSNSILRSLM TAIEEELHA IRTGHSHMYL CILKEQKLLD
1451 LIPFSGSTIV DVGQDEATAC SLLKSMALKI HELVGARMHH LSVCQWEVKL
1501 KLDCDGPASG TWRVVTTNVT GHTCTIDIYR EVEEIESQKL VYHSASSSAG
1551 PLHGVALNNP YQPLSVIDLK RCSARNRRT YCYDFPLAFE TALQKSWQSN
1601 GSTVSEGNEN SKSYVKATEL VFAEKHGSWG TPIIPMERPA GLNDIGMVAW
1651 IMEMSTPEFP NGRQIIVVAN DITFRAGSFG PREDAFFETV TNLACERKLP
1701 LIYLAANS GA RIGIADEVKS CFRVGWSDEG SPERGFQYIY LTEEDYARIS
1751 SSVIAHKLEL DSGEIRWIID SVVGKEDGLG VENIHGSAAI ASAYSRAYEE

FIG. 14

19/37

1801 TFTLTFTVGR TVGIGAYLAR LGIRCIQRLD QPIILTGFS A LNKLLGREVY
1851 SSHMQLGGPK IMATNGVVHL TVPDVLEGVS NILRWLSYVP ANIGGPLPIT
1901 KPLDPPDRPV AYIPENTCDP RAAICGVDDS QGKWLGGMFD KDSFVETFEG
1951 WAKTVVTGRA KLG GIPVGVI AVETQTMMQI IPADPGQLDS HERSVPRAGQ
2001 VWFPDSATKT AQALLDFNRE GLPLFILANW RGFSGGQDL FEGILQAGST
2051 IVENLRTYNQ PAFVYIPMAG ELRGGAWVVV DSKINPDRIE CYAERTAKGN
2101 VLEPQGLIEI KFRSEELQDC MGRLDPELIN LKAKLQDVNH GNGSLPDIEG
2151 IRKSIEARTK QLLPLYTQIA IRFAELHDT S LRMAAKGVK KVV DWEE SRS
2201 FFYKRLRRRI AEDVLAKEIR QIVGDKFTHQ LAMELIK EY LASQATTGST
2251 GWDDDDAFVA WKDSPENYKG HIQKLRAQKV SHSLSDLADS SSDLQAFSQG
2301 LSTLLDKMDP S QRAK FVQEV KKVLD

FIG. 14 (continued)




FIG. 15

21/37

1 AAGCTTGGTA TGGATTcGTC AGCGCCAAgC CGGGGTTTTG CATGCGCCCCg
 51 ACTGGaArCs GAATTCCgTg AgCCcGTaC rrCaATGGCA ACCCCAsGGT
 101 TACTggGGTG GCTGAATGGT CTCsGCTTAC GCAATTGTTT GTGGCAGCwG
 151 CGTGGGCTAA ATGTArGTTG TCTCTTGTTG CACTGCArGA TGGATGGGTA
 201 gCCTCTGGGC CGCCTCTGCT ArTGTCTArC GtTTGCTGAC TGTGGTTTTat
 251 TCAgGGATGC CCATGCCCCAT GCTAgATTGA tAGGTGCCAT TCTAATGGTA
 301 GGTGGCGGTA AGGTTTATTA AGCTGyAGyA TCAGTAGGTA ACCTCATGAA
 351 TCAGGGTTTA aGCACACCTT TTCCTTTGTG TGGGTGCATA AGGAATGCAC
 401 TTGGCTTCGT TCCCTGATAG TCTTTGStCA TGTGTCATTC TACCAAGTGG
 451 GTTACTGTAA CATTGCACTC TATGATGGTT GGTGGTtGTG CATCTTTyTG
 501 CTyCCCCTGG yTGTCTAATA CCTGCATGTa ACTGATGACC yyCyTTTATG
 551 TATCATATAG ATTACATCCT TTTGTTGTAC ATCTCAATTC TGAAAAACA
 601 ATGTTTTGCA TTCTTAGCGc TCTGTGCaCA AgGaAAaGGa gGTTTTACCT
 651 gCAAcTtTTT TTTTCGAGAA AAAACAAACC TTTCTGAaAg gCAGTGATCA
 701 TTTAGtATAA AGAAAATTTG ATTTACTTTC TTCAGAGAGA AtATkCCAAr
 751 CAAACAATTT TCTTACTGTC TGAGCCACGA AATTTGATCT TGATCtTACT
 801 TTCACAAGCC ACATGAAGCC tTATCATCGC TcTGATAAAA AArCCAAaTA
 851 GGTGATTCAT AGAATGAGar AAAGAACCTG TTGCCATTTG GGGaCCTTGT
 901 TGTGTACTCA TTATCCCCC TGCTCAGGTT GAGGTTTCCT TGCCACTGCC
 951 ACCCCTTGGC CCCTTCTTAT ACAACCATCT CCATTGAAAa AGATTTTGCA
 1001 CtACATTTGG GCTTcGTATG aCAAAAaAGG aAAATaAAaC TaAaCAGCAG
 1051 AAACATAGTA TAaTTATAGg TAAAAGGTTc TGGCAAGTTT GAGTGGaAGA
 1101 GACCTTTGTA TATTTGGACA TATTTCACTA GTAAATAGTT TTCTAAAATc
 1151 TTCATGAATG GTGGCCAATA AACTTGATAA GATCTCAACA TGGCAGGTTC
 1201 CTCmAAATG AGAGGAAAAC TGGAAACATC ACAAATATTT TTTAGCGAGT
 1251 GGCCTATAAA TTATAATGTT GCTTTCATTT CTTTGATATT CaAAACTTCC
 1301 TAAGAGTATT CTGCTAGAGC TCTGATGGTG TCTTTTGCCT CTGTCAGATT
 1351 TTCCAGGaGT TTTCTTCCCT TTTtATGGCA CTGTGCGTTT GAGAAGGTCT
 1401 TCAATTGTGC TGTCTGGGCC ACGGAACGAC AATGTCACAG CTTGGATTAG
 1451 CCGCAGCTGC CTCAAAGGCC TTGCCACTAC TCCCTAATCG CCAGAGAAGT
 1501 TCAGCTGGGA CTACATTCTC ATCATCTTCA TTATCGAGGC CCTTAAACAG
 1551 AAGGAAAAGC CATACTCGtT CACTCCGTGA TGGCGGAGAT GGgGTATCAG
 1601 ATGCCAAAAA GCACAGCCAG TCTGTTCGTC AAGGTACTGt GAATATCTTT
 1651 tGaTACAAGc tAAAATTTtG cTACAGAATA TaTAtTtaAA GAGTtCTtTC
 1701 TTGGcTGGtG ttGtTTATTT GttT.aACat sCGAAAGGGC CtctAgttGg
 1751 attGgttaAG tggSCTGAAT ACCACTCCTT AAGGTCTTGA GTTTGCTTTT

FIG. 16

1801 CCCChCGGAG CGAATTTTAg GCTAGGGTTA CCCCCCACC CCCACCCGAA
1851 TCTGCACAGy CCGGyCGyGG yCGyCCTCAT ATAGGCTACG ATGTCATTGT
1901 GTATCGGCGG GCCAGGGGTT TAAGAGTTTT CTTGACCTTT GTTAGAAGAT
1951 CTTAATAATA CAATGTCCAA GGGCTGTCTT ACCCTGTAGG TCGAGTTTTT
2001 AGTTGTTTTA ACATGGTAAT GTTTGAAGCC TCATTCTAGG TrCCAATATA
2051 GATATGCTCA CTGCTCAGTT TCAAATGTTT GTCTGCATGT AGGTCTTGCT
2101 GGCATTATCG ACCTCCCAAG TGAGGCACCT TCCGAAGTGG ATATTTTACA
2151 GTAAGGACTA CAATATTTTG CGTACGTTTG TTTTGGAAAA AGAAAATATT
2201 CTCAGCTTAT TTATACTAGC TTCGCTAATA CTGAAATGCT GTCTTAATGT
2251 CCTGGTGCTG TATGCTCAAT CTTTCATAGT AAATGCTGCA AAATATGTGA
2301 TGTAACTGTT GCAACACAGC CAGGGACCTG TTATTTAGAG CATGGTGAAT
2351 GCTCTGGTTC AGTTATATGA TGTAAGTTATA GCTCATGTTG AAGAATTAGT
2401 TGCAGTGTTT GCTGGACAAT GGTCACTTAT TATAAATCAT ATCTGCATAC
2451 ACATTTGTGA CTTCTGTTGC TGTAATGCC CGCATTTTTT GAGAAAAATT
2501 TAAATGCTTG GCCTAAATTG GACATATATG ATAGACAAAG CTGATTTGAA
2551 CTTTGTTTAT TTTTGACATC CATGCATATT GTCAGTGTTG TGAAAACAAT
2601 ACTAATCCTT TTTTTTTGTC TTTTCCAGT GGATCTGAGG ATCCTAGGGG
2651 GCCAACAGAT TCTTATCAA TGAATGGGAT TATCAATGAA ACACATAATG
2701 GAAGACATGC CTCAGTGTC AAGGTTGTTG AATTTTGTC GGCCTAGGT
2751 GGCAAAACAC CAATTCACAG TATATTAGTG GCCAACAATG GAATGGCAGC
2801 ACCCAAATTT ATGAGGAGTG TCCGGACATG GGCTAATGAT ACTTTTGGAT
2851 CTGAGAAGGC AATTCAACTC ATAGCTATGG CAACTCCGGA AGACATGAGG
2901 ATAAATGCAG AACACATTAG AATTGCTGAC CAATTACGTA GAGGTGCCTG
2951 GTGGAACAAA CAATAATAAC TACGCCAATG TTCAACTCAT AGTGGAGGTT
3001 AGCCTTGCTA ATCTGTTAGT TTACTACTGG TCTGCTGTTT CCTTTATTTG
3051 TTGTATAATG ATTGACATAT TTAAGTAGAG AAATTTATAT TTCTCCTCTG
3101 CTGTTGTGGA AGTCCAATTG TCATCATTAA CTGTGAAATA TTGCAGATGG
3151 CACAAAAACT AGGTGTTTCT GCTGTTTGGC CTGGTTGGGG TCATGCTTCT
3201 GAGAATCCTG AACTGCCAGA TGCATTGACC GCAAAAGGGA TCGTTTTTCT
3251 TGGCCCACCT GsATCATCAA TGAATGCTTT GGGAGATAAG GTCGGCTCAG
3301 CTCTCATTGC TCAAGCAGCC GGGGnCCCAA CTCTTGCTTG GAGTGGATCA
3351 CATGTGAGTC TCACTCTTTG ATTACTATCC GCCTGFCTCA TTGCTCTCTC
3401 TTTTCATATC TAATGACACT AAATTTAGGT TGAAGTCCA TTAGAGTGCT
3451 GCTTAGACGC GATACCTGAG GAGATGTATA GAAAAGCTT

FIG. 16 (continued)

23/37

1 GAATTCCTGTG AGCCCTGTAC GGCAATGGCA ACCCCAGGGT TACTGGGGTG
51 GCTGAATGGT CTCGGCTTAC GCAATTGTTT GTGGCAGCTG CGTGGGCTAA
101 ATGTAGGTTG TCTCTTGTTG CACTGCAGGA TGGATGGGTA GCCTCTGGGC
151 CGCCTCTGCT AGTGTCTAGC GTTGCTGACT GTGGTTTATT CAGGGATGCC
201 CATGCCCATG CTAGATTGAT AGGTCATAGG TGCCATTCTA ATGGTAGGTG
251 GCGGTAAGGT TTATTAAGCT GTCGTATCAG TAGGTAACCT CATGAATCAG
301 GGTTTAAGCC CACCTTCTCC TTTGTGTGGG TGCATAAGGA ATGCACTTGG
351 CTTGTTCCC TGCTAGTCTT TGCTCATGTG TCATTCTACC AAGTGGGTTA
401 CTGTAACATT GCACTCTATG ATGGTTGGTG GTTGTGCATC TTTTGTCTC
451 CCCTGGTTGT CTAATACCTG CATGTAACCTG ATGACCTTCT TTTATGTATC
501 ATATAGATTA CATCTTTTGT TGTACATCTC AATTCTGAAA AACAAATGTTT
551 TGCATTCTTA GCGCTCTGTG CACAAGGAAA AGGAGGTTTT ACCTGCAACT
601 TTTTTTTTCG AGAAAAACA AACCTTCTG AAAGGCAGTG ATCATTTAGT
651 ATAAAGAAAA TTTGATTTAC TTTCTTCAGA GAGAATATTC CAAACAAACA
701 ATTTTCTTAC AGTCTGAGCC ACGAAATTTG ATCTTGATCT TACTTTCACA
751 AGCCACATGA AGCCTTATCA TCGCTCTGAT AAAAAACCA AATAGGTGAT
801 TCATAGAATG AGAAAAAGAA CCTGTTGCCA TTTGGGGACC TTGTTGTGTA
851 CTCATTATCC CCCCTGCTCA GGTTGAGGTT TCCTTGCCAC TGCCACCCCT
901 TGGCCCCCTC TTATACAACC ATCTCCATTG AAAAAGATTT TGCACACAT
951 TTGGGCTTCG TATAACAAAA AAGGAAAATA AACTAAACA GCAGAAACAT
1001 AGTATAATTA TAGGTAAAAG GTTNTGGCAA GTTTGAGTGG TAGAGACCTT
1051 TGTATATTTG GACATATTTT ACTAGTAAAT AGTTTTCTAA AATGTTCATG
1101 AATGGTGGCC AATAAACTTG ATAAGATCTC AACATGGCAG GTTCCTTCAA
1151 AATGAGAGGA AACTGGAAA CATCACAAAT ATTTTTTAGC GAGTGGCCTA
1201 TAAATTATAA TGTTGCTTTC ATTTCTTTGA TATTCAAAC TTCCTAAGAG
1251 TATTCTGCTA GAGCTCTGAT GGTGTCTTTT GCCTCTGTCA GATTTTCCAG
1301 GAGTTTTCTT CCCTTTTTAT GGCACGTG

FIG. 17

24/37

1 AGCATCCCTT GGGATTGTGA TnACTCACAT AAATTCTTGC GAAnTGTTGA
 51 CATTCTAGTG ATTTGAGTTC CGTTCTAGTG TGCTAGTCAn TTGAGCTCAA
 101 GTCTTGTTTT TATGTGTGCG TATTTACTGT GATCTTTGTG TCGTGTGTGA
 151 GTTGTTGATC CTTCCCTTGC TCCGTGATTC TTTGTGAAAT CTTTTGAAAG
 201 GGCGAGAGGC TCCAAGCTGT GGAGATTCTT CGCAAGTGGG ATTAAGAAAA
 251 GCAAAGCAAC ACCGTGGTAT TCAAGTTGGT CTTTGGACCG CTTGAGAGGG
 301 GTTGATTGCA ACCCTCGTCC GTTGGGACGC CACAACGTGG AGTAGGCAAG
 351 CGTTGGTCTT GGCCGAACCA CGGGATAACC ACCGTGCCAT CTCTGTGATT
 401 GATATCTCTT GGTTATTGTG TTGTGTTGAG ATCCTTCTCT AGCCACTTGG
 451 CAAATTACTG TGCTAACAAT TAATCAAGTT TTGTGGCTTA AGATTTTGAA
 501 GTATTACAGG ATCTGCATCA TGGTCTGTGT CTCCACAGCT ATGACACCCA
 551 CAGGAATTCA TGTGTTCCCT GGAGCCACTC TTGGATGACC TAAAGGAATT
 601 ATTTCTAACC GGCTTGTACA CATATGATGC ATCAAGAGAT GAGTGTTTTA
 651 CTATGCGAGG GGCCATGCTT ATGACCATAA GTAATCTTCC TGGTTTAGAA
 701 ATGCTTGCTT CTCATATGGT TCATGGGAAA TTCGCATGCC TCCTTGTGGT
 751 GAAAATGTCT GGACAAAACA GCTGAAGAAT GGTCGTAAAT CTTGTTTTAT
 801 GGGAAATCGC CAATATATTG ATCTTGATCA TTCTTATTGC TTGGATGCAG
 851 ACTCCGTTTG ATGGAACGAT AGACTTCGAA CAAAACCTAA AACCTATTAT
 901 GATCGTCCAA TTTTGGATGA AATCATCACA CTTGGTGATT TCAAGAACTC
 951 AAAAaYTTAC AGTTAATTGG ATATAGGAGG GnGCAAAAAC ACAGTAAGTT
 1001 GGACATTCCA TAAGGGGATT TATTTTAGTT GACAATAAAG TAGATGGGCA
 1051 TCATCCTGAG TTTnGTTTGG CATCGTGTCT TAGATTGAAA CTGTAAGGAT
 1101 GGACATGGTA GnTAACAGGT TGAGATGAAT GATTCAACAG TTGAAGCGAA
 1151 TGTACAATCT TTATGTGATG GTGGCTnTTA ATGCAGGTAA CTAGTTTTTT
 1201 TTTATGCTTT ATTATTAATT AGTTGGATAA ATGGTTTnGA TTTnTGATTG
 1251 TTAAAnTGCA ATGGCTCCAG TTGGATGGnA ACAGCCAnGT AATTTATGCA
 1301 GAAGnAGnAG nTGGTGGTAC ACGGnTTCAG ATTGATGGAA AGAnATGTTT
 1351 ATTGCAGGTA AATAnTCCCT TnTTCCTTTA TATTTTTGTT GTnTGATTGT
 1401 ATAAntTTGn TAGATTATTT GTATAATTTA TTATTGCATT TCACCCCACT
 1451 AAnTTATTTT TAAAAGATGG GTTTTGTGTG TTGnTTCAGC nGGCGACATC
 1501 ACATAAGnAA ATTGTGATTA ATTTTTGTTT TTTTGCAGnA TGACCATGAT
 1551 CCATCAAAGT TATTA

FIG. 18A

1 ACATAAGCTG GGTTAGTAGT GGTGAATTAG TGGATTATTG ATCTGTTGTG
51 GGCAAGGAGn TGGCTTGGTG TCGAGAAAAA ACATGGAAgT GTGTATTGCC
101 AGTGcTTaTT yTrGGGATAT GAGGGAATTw AmATTAcATT TGTGAcTGGG
151 CGGACTGTAG GATAGGAGTT ATCTTGyTcG ATTGGTATAC GGTGCATACA
201 GskyTTGACC AGCTATTATT TTAACAGGgT TTTCTGCCCT GAACAAGTCC
251 TTGGGCGGGA AGTGTACAGc TCCCACATGC AGCTTGGTGG TCCTAAGATC
301 ATGGCGACCA ATGGTGTTGT CCACCTCACT GTTCCAGATG ACCTTGAAGG
351 TGTTTCCAAT ATATTGAGGT GGCTCAGCTA TGTCCTGCA AACATTGGTG
401 GACCTcTTCC TATTACCAA CCTCTGGACC CTCCAGACAG ACCTGTTGCT
451 TACATCCCTG AGAACACATG CGATCCACGT GCAGCTATCT GTGGTGTAGA
501 TGACAGCCAA GGGAAATGGT TGGGTGGTAT GTTTGACAAA GACAGCTTTG
551 TGGAGACATT TGAAGGATGG GCAAAACAG TGGTTACTGG CAGAGCAAAG
601 CTTGGAGGAA TTCCATGCAT CTTAATAAAC ACAGTTGGCC CTTAAAGCAA
651 GTGAACCTTCT TGAACAAACC AAATAAGTG AACTCCGTGC AAGCGTTGCA
701 AGAAGCCTTT CGGATCTGGG GATGCATAAG GGAGAAATGA GTATTAAGGA
751 TAACATGGAA GATTTAGTCT CTGCCCCATT ACCTGTTGAA GATGCTCTGA
801 TTTCTTTGTT TGATTACAGT GATCGAACTG TTCAGCAGAA AGTGATTGAG
851 ACATACATAT CACGATTGTA CCAGGTATTA TATCAACTAA CTTAATGTCT
901 TCCATAGTCT CACTAAGCAT ATCTGATATG tTTAGATACC CTACATGGAA
951 TGCTCATCTT TTCATTTGAC ACAAAGAAAC ATTGAGAAAT GAGATGCTGA
1001 CGATTGGCTG AAATTAAGTG GGTnTGAGAA ATTGTGATCT CCCAACTTGT
1051 TAATGCACAA TGTTCTGGCT AACTTGCCAA TATTTTTTCA GCCTCATCTT
1101 GTTnAGGATA GCAnCCAAAT GAAATCCAAG GATCTGGTGC TATTACTTTT
1151 TGGGAATTTA TGAAGGGC

FIG. 18B

26/37

1 CTCCCAATAT TGT CATGAGG CTTGCATCCC AGGTTAGTTT TTTTTCCTTT
51 CTGAAATTTA TATTCATAC CTTTTCACCT TTAGTTATCC TTGTATTTTC
101 TGGAAGCTTC ATCTGATGCA TTATTGACAA ATGCACTAAT GGTCATCATA
151 TTTGGAGATT AACATATTTA TCTTAATTGA TGGGAACCTCT TGAAAATGAC
201 AATGGTTGAG CAGATAATTA ACAGTTTTTTT AATAAAAAAA CATGCATTTTC
251 TAGGAGTTGG ACTAAGCTTT TCTTAGTATG AAGTGCCATG TTTTACATGG
301 TCCATTTGTG TCAATTTACA GTCGGTATCA TGGAAAGGTT GTCATAATGG
351 CTGGAGANAA ACAACACATC TTGTTTCTCA ACACTTGTGG GAGAAGANGT
401 TTTACCTTTT TTCCTAAAAT TACTTTTTGT ACTAAATTGT ATAATTTTTC
451 CAATATTCTC CATGATTATT GAACTCTGCT GTGTTCAAAC AGCCAAAACA
501 TGTTTCCATA CTTTACACCT TTATTTTTTA GATGGAACCT GGAATTGTGC
551 TCTGTTATCT GTATCATGCA TATATTGATC TTAAACCTAT CTCTATTGTA
601 GAATCCGCAC TTGAATTCAG TTGCTTGTGA TCAATATG

FIG. 18C

1 GGTAACCACC ACACCCGCGG CGCTTAATGG CCGTACAGGG NGGTCCCATT
51 CGCCATTTCAG GTGCGCAACT GTTGGGAAGG GCGATCGGTG CGGGCTTCTT
101 CGNTATTACG CCAGCTGGCG AAAGGGGGAT GTGCTGCAAG GCGATTAAGT
151 TGGGTAACGC CAGGGTTTTTC CCAGTCACGA CGTTGTAAAA CGACGGCCAG
201 TGAGCGCGCG TAATACGACT CACTATAGGG CGAATTGGGT ACCGGGCCCC
251 CCCTCGAGGT CGACCTGCAG GTCAACGGAT CCTAGGGGGC CAACAGATTC
301 TTATCAAATG AATGGGATTA TCAATGAAAC ACATAATGGA AGACATGCCT
351 CAGTGTCCAA GGTTGTTGAA TTTTGTGCGG CACTAGGTGG CAAAACACCA
401 ATTCACAGTA TATTAGTGGC CAACAATGGA ATGGCAGCAG CAAAATTTAT
451 GAGGAGTGTC CGGACATGGG CTAATGATAC TTTTGGATCT GAGAAGGCAA
501 TTCAACTCAT AGCTATGGCA ACTCCGGAAG ACATGAGGTA AATGCAGAAC
551 ACATTAGA

FIG. 18D.

1 GAATAATCTG CCTGCAGCTC AAGTTGCTGT TGGAATGGGC ATACCTCTTT
51 GGCAGATTCC AGGTAATTAC CAATTTACCA ACTTATTTAG TTCCTTATTG
101 TTTTATTCTC TAATTTTCTA CTTATGTAgA AATCAGACGT TTCTATGGAA
151 TGGACTATGG AGGAGGGTAT GACATTTGGA GGAAAACAGC AGCTCTTGCT
201 ACACCATTTA ATTTTGATGA AGTAgtATTcT CAATGGCCAA AGGGCCATTG
251 TGTAgtCAGTT AGAATTACTA GTGAGGACCC AGATGATGGT TTCAAACcTA
301 CTGGTGGGAA AGTGAAGGTA AGTTTTCTAG ATGACATGTA TTATATATCG
351 TTCAAgtAgA TTAAGTTTGG TTAAATGAcT AGGTCTTGAT TTTTtATCTT
401 TCAGGAGATA AGTTTTTAAA GCAAGCCTAA TGTTTGGGCC TaCTTCTCAG
451 TAAAGGTaAC TTGTTAActT TAGTACGCTG TCACATTATt ctTCsTTGTG
501 .AAAATAAtT TGAACGGTtC TCTTTGTATT TTaACCAAtCC AtCgtTCTCAT
551 TTAsCagAgC ACACAAATAT TtGCACTGAC CCCCcTcCCC tTATCtGctT
601 TCAGtCTGGT GGAgGCATtC AtGAATTTtGC TGATTCTCAG TTCGGTATGT
651 GTAAACCAAG AGTATTCTTT GTAATTTATA TTGGTCCTCA ATTTTGAAAT
701 ATTGcTCTTT CCGTTACAGG ACaWGTTTTT GCATATGGGC TCTCTAGATC
751 AGCAGCAATA ACAAACATGA CTCTTGcATT AAAAnAGATT CAAATTCGTG
801 GAGAAATTCA TTCAAATGTT TGATTACACA GTTGATCTCT TAAATGTTAA
851 GAAATATTAA CCACCTTTTA AATCACATTT TCCATTATGT TTGATTCCAT
901 ATCATTAATT TTGATTTTCT ATTATGGCTA AACCTGTGGT GCTATTTTCC
951 TATTATCCCA GGCTTCCGAC TTTAGA

FIG. 18E

GGATCC TAgGGGGCCA

1401 ACAGATTCTT ATCAAATGrA aTGGGATTAT CAATGAAACA CATAATGGAA
1451 GACATGCCTC AGTGTCCAAG GTTGTGAAT TTTGTGCGGC ACTAGGTGGC
1501 AAAACACCAA TTCACAGTAT ATTAGTGGCC AACAATGGAA TGGCAGCAgC
1551 aAAATTTATG AggAGTGTCC GGACATGGGC TAATGATACT TTTGGATCTG
1601 AGAAgGCAAT TCAACTCATA GCTATgGCAA CTCCGGAAGA CATGAGGATA
1651 AATGCAGAAC ACATTAGAAT TGCTGACCAA TTaCGTAgrAr gTGcctgGTG
1701 gaacaaACAA tA.ATAActA cgCCaATGTT CAACtCaTAg TGGAAgTTAG
1751 CCTTGcTAAT CTGTTAgtTTT ACTACTGgTC tGctGTTtCC TTtATTtGtt
1801 GTaTAATGAT tGACaTATtt AagTAgtAgAA atTTATAtTT CtCctCtgCt
1851 GTTGTGgAag TCCAatTGTC acCATtAAct GTgAAaTATt gCAgATgGca
1901 CaAAAActAG gTgtTTCTgC TGTTTGGCCT gGTTGGGGTC ATGCTTCTGA
1951 GAATCcTGAA CTGCCAGATG CATTGACCGC AAAAGGGATC GTTTTTCTTG
2001 GCCCACCTGc ATCATCAATG AATGCTTTGG GAGATAAGGT CGGCTCAGCT
2051 CTCATTGCTC AAGCAGCCGG GgtCCCAACT CTTGCTTGGA GTGGATCACA
2101 TGTGAGTCTC ACTCTTTGAT TACTATCCGC CTGTCTCATT GCTCTCTCTT
2151 TCATATTCTA ATGACACTAA ATTTAGGTTG AAGTTCCATT AGAGTGCTGC
2201 TTAGACGCGA TACCTGAGGA GATGTATAGA AAAGCTT

FIG. 18F

1 AATTCCTGTG GGTGTTATAG CTGTGGAGAC ACAGACCATG ATGCAGCTCA
51 TCCCTGCTGA TCCAGGTCAA CTTGATTCCC ATGAGCGATG TGTTCCTCGG
101 GCTGGACAAG TGTGGTTCCC AGATNCTGCA ACCAAGACAG CTCAGGCATT
151 ATTAGACTTC AACCGTGAAG GATTGCCTCT GTTCATCCTG GCTAACTGGA
201 GAGGCTtCTC TGGGGgACAG AGAGATCTCT T

FIG. 19A

1 AATTCATGCA TCTTAATAAA CACAGTTGGC CCTTAAAGCA AGTGAAGTTC
51 TTGAACAAAC CAAACTAAGT GAACTCTGTT CCAGCATTGC AAGAAGCCTT
101 TCAGATCTGG GGATGCATAA GGGAGAAATG ACTATTAAGG ATAGCATGGA
151 AGATTTAGTC TCTGNCCCAT TGCCTGTTGA AGATGCTCTT ATTTCTTTGT
201 TTGATTA

FIG. 19B

1 ATAGACCTGT CGCATACATC CCTGAGAACA CATGCGATCC GCGTGCAGCC
51 ATCCGTGGnG TAGATGACAG CCAAGGGAAA TGGTTGGGTG GTATGTTTGA
101 CAAAGACAGC TTTGTGGAGA CATTTGAAGG ATGGGCAAAA ACAGTGGTTA
151 CTGGTAGAGC AAAGCTTGGA GGAAGGAATT

FIG. 19C



A**B**

FIG.20

32/37

1 ACAAATAGT AAAGACCTCC AGAAGGTTGT AGACATTGTT TTGTCTCACC
51 AGGGTGTGAG AAGCAAAACC AAGCTgAtAC TCaCGCiCAT gGAgAAACTG
101 GTcAtCCaA ACCCTgcTGC CtATAGGGaC CAGcTgATTc GCtTTTcTTC
151 CCTCAACCAT aAAAGaTaTT aCAAGcTGGC CCTTaAAgcT AgTgAACTTc
201 TcGaACaAAC CaAGcTCAgT gaACtCCGca CAAGCATTGT AaGGAaCcTT
251 tcAGCgCTGg AgATGTTTAC TGAAGAAAGG GCAGGTTTCT CGTTGCAAGA
301 GAAGAAATTG GCCATTAATG AGAGCATGGG AGATTTAGTC AGTGCCCCAC
351 TGCCAGTTGA AGATGCACTT ATATCTTTGT TTGATTGTAT TGATcAAACT
401 CTTcAGCAGA GAGTGATCGA GACGTACATA TCTCGATTAT ACCAGCCTcA
451 ACTTGTGAAG GATAGTATCC AGCTGAAATA TCAGGATTCT GCTGtATCG
501 CTTTATGGGA ATTCAACGAA GGGCATCCTG ATAATAGATT GGGTGCTATG
551 GTTATCCTGA AGTCTCTAGA ATCTGTGTCA ACAGCAATTG GAGCTGCTCT
601 AAAGGATAGA TCACATTATG CAAGCTCTGC GGGTAACACA GTGCATATTG
651 TTTTGTTGGA TGCTGATACT CAGACGAATA CAACTGAAGA TAGTGGTGAT
701 AACGACCAAG CTCAAGACAG GATGGACAAA CTCTCTGTAA TACTGAAACA
751 AGATATTGTC ACGGCTGATC TATGTGCTGC TGGTGTCAAG GTTATTAGTT
801 GCATTGtCA AAGAGATGGA GCACTCATGC CTATgCGCCG TACCTTCCTC
851 TTGTCTGAAG AAAAGCTTTG TTATGAGGAA GAGCCGATTc TTCGGCATGT
901 GGAGCCTCCA CTTTCTGCAC TTCTTGAGTT GGATAAATTG AAAGTGAAAG
951 GATaCAATGA GATGAAGTAT ACACCGTCAC GTGATCGTCA GTGGCATATA
1001 TACACACTTA GAAATACTGA AAACCCCAAA ATGTTGCACA GGGTATTTTT
1051 CCGAACTCTT GTCAGACAGC CGAGTGCTGG CAACAGGTTT ATGTCAGGCC
1101 ATATCAATGA TGTGAAGTA GGACATGCTG AGGAATCTCT TTCATTTACA
1151 TCAAGCAGCA TATTAAGATC TTTGATGACT GCTATAGAAG AATTGGAGCT
1201 TCACGCGATT AGGACTGGTC ATTCTCATAT GTACTTGTGC ATATTGAAAG
1251 AGCAAAAGCT TCTTGACCTT GtCCTGTTT CAgGGAACAC AGTTGTGGAT
1301 GTTGGCCAAG ATGAAGCTAC TGCGTGCTCT CTTATGAAAG AAATGGCTCT
1351 AAAGATACAT GAACTTGTTG GTGCAAGAAT GCATCATCTT TCTGtATGCC
1401 AGTGGGAAGT gAAACTTAAG TTGGACAGTG ATGGACCTGC CAGTGGTAGC
1451 TGGAGAGTTG TAACAACCAA TGtCACTAGT CACACCTGCA CTGTGGATAT
1501 CTACCGGGAG GTCGAAGATa CAGATTCACA GAAGCTAGTA TACCACTATG


Figure 21

33/37

1551 CTGCATCGTC ACCTGGTCCT TTGCATGGTG TCACATTGAG TAATTCATAC
1601 CAGCCTTTGA GCGTTATTGA TCTAAAACGA TGCTCTGCCA GGAACAACAG
1651 AACTACATAC TGCTACGATT TTCCATTGGC ATTTGAAACT GCAGTGC GGA
1701 AGTCATGGGC TAACGTTTCt AGTgAAAaCA aCCAATGTtA TGTTAAgCA
1751 aCAgAgcTGG TGTTtGCTgA CAAGAATGGG TCGTGGGGCA CTCCTATAGT
1801 TCCTATGCAG CGTGCTGCTG GGCTCAATGA CATCGGTATG GtTGCgTGGA
1851 TCtTGGaCAT GTCCaCTCCT GAATTTCCCA GTGGCAGACA GATTATTGTT
1901 ATCGCAAATG ATATTACTTT CAGAGCTGGA TCATTTGGCC CAAGGGAAGA
1951 TGCATTTTTT GAAGCTGTTA CCAACCTGGC TTGTGAGAGA AAGCTTCCTC
2001 TTATCTACAT GGCAGCAAAC TCTGGTGCTC GGATTGGCAT TGCTGATGAA
2051 GTAAATCTT GCTTCCGTGT TGAATGGGT GATCCTGCCa ACCCTGAACG
2101 TGGATTTAAG tACATTTATC TGAATgAAGA AGaCTATGGT CGTAtTAgtCT
2151 cTTCTGtTA tAGCGCACAAg aCaCAgcTag ATagtGGCgA AAtAaGgtgG
2201 GtTAICGATt CTGTTGIGGg CAAGGAGGAT GGaCTAGGTG TGGAGAACAT
2251 ACATGGAAgT GCTGcTAItG CCAGTGCTTA TTCTAGGgCA tACgAgGAGA
2301 CaTTtACaCT TACATTtGTG AGTGGtCgAA CGGTTgGAAT AGgAgCcTAT
2351 cTTGCTcGGc TTGGcAtaCG GtGCAtACAG cGTATTGaCC AGcCCATTAT
2401 CTtAAcTgGG TTTTCTGCCc TGaACAAGCT TCTCGGGCGG gAAGTGTACA
2451 GCTCCACAT GCAGTTGGGT GGTCCCAAAA TCATGGCTAC AAAtGGTGtT
2501 GTCCATCTGA CTGTTCCAGA TGACCTTgAA GGTGtTtCtA ATATAItGAG
2551 GTGGCTCAgT TATGTTCTG CTAACATTGG TGGACCTCTT cCTATTaCaA
2601 AATCTTTAGA CCCAATaGAC AGACCTGtTG cATACATcCC TGaGAATaCa
2651 TGtGAICCTc GTGCAGCCAT CAGTGGCATT GATGACAGCC AAGGGAAATG
2701 GTTGGGTGGT ATGTTTGACA AAGACAGTTT TGTGGAGACA TTTGAAGGAT
2751 GGGCAAAGAC AGTAGTTACT GGCAGAGCAA AACTTGGAGG GATTCCTGTG
2801 GGTGTTATAG CTGTGGAGAC ACAGACCATG ATGCAGCTCG TCCCTGCAGA
2851 TCCAGGCCAG CCTGATTCCC ATGAGCGGTC TGTTCTCGT GCGGGGCAAG
2901 TCTGGTTTCC AGATTCAGCT ACCAAGACAG CGCAGCAAT GTTGGACTTC
2951 AACCGTgAAG GATTACCTCT GTTCATCCTt GCTAAcTGA GAGGCTTCTC
3001 TGGTGGGCAA AGAGACCTTT TTGAAGGAAT T

1 QNSKDLQKV DIVLSHQGVR SKTKLILTL EKLVPNPAA YRDQLIRFSS
51 LNHKRYKLA LKASELLEQT KLSELRTSIV RNLSALEMFT EERAGFSLQE
101 KKLAINESMG DLVSAPLPVE DALISLFDCI DQTLQQRVIE TYISRLYQPQ
151 LVKDSIQLKY QDSAVIALWE FNEGHPDNRL GAMVILKSLE SVSTAIGAAL
201 KDRSHYASSA GNTVHIVLLD ADTQTNTTED SGDNDQAQDR MDKLSVILKQ
251 DIVTADLCAA GVKVISCIVQ RDGALMPMRR TFLSEEKLC YEEEPILRHV
301 EPPLSALLEL DKLKVKGyne MKYTPSRDRQ WHIYTLRNTE NPKMLHRVFF
351 RTLVRQPSAG NRFMSGHIND VEVGHAEESL SFTSSSILRS LMTAIEEEL
401 HAIRTGHSHM YLCILKEQKL LDLPVSGNT VVDVGQDEAT ACSLMKEMAL
451 KIHVLGARM HHL SVCQWEV KLKLDSDGPA SGSWRVVTTN VTSHTCTVDI
501 YREVEDTDSQ KLVYHYAASS PGPLHGVTL NSYQPLSVID LKRCSARNNR
551 TTYCYDFPLA FETAVRKSWA NVSSENNQCY VKATELVFAD KNGSWGTPIV
601 PMQRAAGLND IGMVAWILDM STPEFPSGRQ IIVIANDITF RAGSFGPRED
651 AFFEAVTNLA CERKLPLIYM AANS GARIGI ADEVKSCFRV EWVDPANPER
701 GFKYIYLNNE DYGRISSEVI AHKTQLDSGE IRWVIDSVVG KEDGLGVENI
751 HGSAAIASAY SRAYEETFTL TFSGRTVGI GAYLARLGIR CIQRIDQPII
801 LTGFSALNKL LGREYSSHM QLGGPKIMAT NGVHLLTVPD DLEGVSNILR
851 WLSYVPANIG GPLPITKSLD PIDRPVAYIP ENTCDPRAAI SGIDDSQGKW
901 LGGMFDKDSF VETFE GWAKT VVTGRAKLGG IPVGVI AVET QTMMQLVPAD
951 PGQPDSHERS VPRAGQVWFP DSATKTAHAM LDFNREGLPL FILANWRGFS
1001 GGQRDLFEGI

Figure 22



Example 3








Figure 2A




A



B

25
Figure 7.
Phylogram of ACCases based on
UPGMA analysis of Kimura
protein distances for 121
ungapped amino acids.

-Chloroplast transit peptides
on completely sequenced plant
ACCases are noted as present
(+) or absent (-).



SEQUENCE LISTING

<110> Regents of the University of Minnesota, et al.

5 <120> DNA encoding oat acetyl CoA carboxylase

<130> 600.439WO1

10 <150> US 60/091,640

<151> 1998-07-02

<150> US 60/090,240

<151> 1998-06-22

15

<160> 30

<170> FastSEQ for Windows Version 3.0

20 <210> 1

<211> 2001

<212> DNA

<213> Zea mays

25 <220>

<221> unsure

<222> (303)...(303)

<223> "n" represents "a" or "g" or "c" or "t"

30 <400> 1

agagatgaag	ctcgcatgcc	aatgcgccac	acattcctct	ggttgatga	caagagttgt	60
tatgaagaag	agcagattct	ccggcatgtg	gagcctcccc	tctctacact	tcttgaattg	120
gataagttga	aggtgaaagg	atacaatgaa	atgaagtata	ctccttcgcg	tgaccgccaa	180
tggcatatct	acacactaag	aaatactgaa	aacccccaaa	tggtgcatag	ggtgtttttc	240
35 cgaactattg	tcaggcaacc	caatgcaggc	aacaagttta	gatcgggtca	gatcagcgac	300
gчнаaggtag	gatgtcccga	agaatctctt	tcatttacat	caaatagcac	cttaagatca	360
ttgatgactg	ctattgaaga	attagagctt	catgcaatta	ggacagggtca	ttctcacatg	420
tatttggtga	tactgaaaga	gcaaaagctt	cttgacctca	ttccattttc	agggagtaca	480
attggtgatg	ttggccaaga	tgaagctacc	gcttggtcac	ttttaaaatc	aatggctttg	540

aagatacatg agcttggttg tgcaaggatg catcatctgt ctgtatgcca gtgggaggtg 600
 aaactcaagt tggactgtga tggccctgca agtgggtacct ggagagttgt aactacaaat 660
 gttactggtc acacctgcac cattgatata taccgagaag tggaggaaat agaatacacag 720
 aagttagtgt accattcagc cagttcgtca gctgggacct tgcattggtgt tgcactgaat 780
 5 aatccatatt aacctttgag tgtgattgat ctaaagcgct gctctgctag gaacaacaga 840
 acaacatatt gctatgattt tccgctggcc tttgaaactg cactgcagaa gtcattggcag 900
 tccaatggct ctactgtttc tgaaggcaat gaaaatagta aatcctacgt gaaggcaact 960
 gagctagtgt ttgctgaaaa acatgggtcc tggggcactc ctataattcc gatggaacgc 1020
 cctgctgggc tcaacgacat tgggtatggtc gcttggatca tggagatgtc aacacctgaa 1080
 10 tttcccaatg gcaggcagat tattgttgta gcaaatgata tcactttcag agctggatca 1140
 tttggcccaa gggaagatgc attttttgaa actgtcacta acctggcttg cgaaaggaaa 1200
 cttcctctta tatacttggc agcaaactct ggtgctagga ttggcatagc tgatgaagta 1260
 aaatcttgct tccgtgttg atggtctgac gaaggcagtc ctgaacgagg gtttcagtac 1320
 atctatctga ctgaagaaga ctatgctcgc attagctctt ctgttatagc acataagctg 1380
 15 gagctagata gtggtgaaat taggtggatt attgactctg ttgtgggcaa ggaggatggg 1440
 cttggtgtcg agaacatata tgggaagtgc gctattgcca gtgcttattc tagggcatat 1500
 gaggagacat ttacacttac atttgtgact gggcggactg taggaatagg agcttatctt 1560
 gctcgacttg gtatacggtg cataeagcgt cttgaccagc ctattatctt aacagggttt 1620
 tctgccctga acaagctcct tgggcgggaa gtgtacagct cccacatgca gcttggtggt 1680
 20 cctaagatca tggcgaccaa tgggtgtgtc cacctcactg ttccagatgt ccttgaagggt 1740
 gtttccaata tattgaggtg gctcagctat gttcctgcaa acattggtgg acctcttctt 1800
 attaccaaac ctctggaccc tccagacaga cctgttgctt acatccctga gaacacatgc 1860
 gatccacgtg cagctatctg tgggtgtgat gacagccaag ggaaatggtt ggggtggtatg 1920
 tttgacaaag acagctttgt ggagacattt gaaggatggg caaaaacagt ggttactggc 1980
 25 agagcaaagc ttggaggaat t 2001

<210> 2

<211> 258

<212> DNA

30 <213> Zea mays

<400> 2

gttcctgcaa acattggtgg acctcttctt attaccaaac ctctggaccc tccagacaga 60
 cctgttgctt acatccctga gaacacatgc gatccacgtg cagctatctg tgggtgtgat 120
 35 gacagccaag ggaaatggtt ggggtggtatg tttgacaaag acagctttgt ggagacattt 180
 gaaggatggg caaaaacagt ggttactggc agagcaaagc ttggaggaat tcctgtgggc 240
 gtcatactg tggagaca 258

<210> 3

40 <211> 4

<212> PRT

<213> Zea mays

<400> 3

5 Val Met Lys Met

1

<210> 4

<211> 26

10

<212> DNA

<213> Zea mays

<400> 4

gccagattcc accaaagcat atatcc

26

15

<210> 5

<211> 7470

<212> DNA

<213> Zea mays

20

<220>

<221> unsure

<222> (7319) ... (7319)

<223> "n" represents "a" or "g" or "c" or "t"

25

<400> 5

ggctctcaat tgtgctgtct gggccacgga acgacaatgt cacagcttgg attagccgca	60
gctgcctcaa aggccttgcc actactccct aatcgccaga gaagttcagc tgggactaca	120
ttctcatcat cttcattatc gaggccctta aacagaagga aaagccatac tcgttcactc	180
30cgtgatggcg gagatggggt atcagatgcc aaaaagcaca gccagtctgt tcgtcaaggt	240
cttgctggca ttatcgacct cccaagttag gcaccttccg aagtggatat ttcacatgga	300
tctgaggatc ctagggggcc aacagattct tatcaaata atgggattat caatgaaaca	360
cataatggaa gacatgcctc agtgtccaag gttgttgaat tttgtgcggc actaggtggc	420
aaaacaccaa ttcacagtat attagtggcc aacaatggaa tggcagcagc aaaatztatg	480
35aggagtgtcc ggacatgggc taatgatact tttggatctg agaaggcaat tcaactcata	540
gctatggcaa ctccgaaga catgaggata aatgcagaac acattagaat tegtaccaa	600
ttcgtagagg tgcctggtgg aacaaacaat aataactacg ccaatgttca actcatagtg	660
gggatggcac aaaaactagg tgtttctgct gtttggcctg gttgggtca tgcttctgag	720
aatcctgaac tgccagatgc attgaccgca aaagggatcg tttttcttgg cccacctgca	780
40tcatcaatga atgctttggg agataaggtc ggctcagctc tcattgctca agcagccggg	840

	gtcccaactc	ttgcttggag	tggatcacat	gttgaagttc	cattagagtg	ctgcttagac	900
	gcgatacctg	aggagatgta	tagaaaagct	tgcgttacta	ccacagagga	agcagttgca	960
	agttgtcaag	tggttgggta	tcctgccatg	attaaggcat	cctggggagg	tggtggtaaa	1020
	ggaataagaa	aggttcataa	tgatgatgag	gttagagcgc	tgtttaagca	agtacaaggt	1080
5	gaagtccctg	gctccccaat	atttgtcatg	aggcttgcac	cccagagtcg	gcatcttgaa	1140
	gttcagttgc	tttgtgatca	atatggtaat	gtagcagcac	ttcacagtcg	tgattgcagt	1200
	gtgcaacggc	gacaccagaa	gattattgaa	gaagggtccag	ttactgttgc	tcctcgtgag	1260
	acagttaaag	cacttgagca	ggcagcaagg	aggcttgcta	aggctgtggg	ttatgttggt	1320
	gctgctactg	ttgagtatct	ttacagcatg	gaaactggag	actactatct	tctggaactt	1380
10	aatccccgac	tacaggttga	gcatccagtc	actgagtggg	tagctgaagt	gaatctgcct	1440
	gcagctcaag	ttgctgttgg	aatgggcata	cctctttggc	agattccaga	aatcagacgt	1500
	ttctatggaa	tggtactatg	aggagggtat	gacatttgga	ggaaaacagc	agctcttgct	1560
	acaccattta	attttgatga	agtagattct	caatggccaa	agggccattg	tgtagcagtt	1620
	agaattacta	gtgaggaccc	agatgatggg	ttcaaacctt	ctgggtgggaa	agtgaaggag	1680
15	ataagtttta	aaagcaagcc	taatgtttgg	gcctacttct	cagtaaagtc	tggtggaggc	1740
	attcatgaat	ttgctgattc	tcagtttgga	catgcttttg	catatggact	ctctagacca	1800
	gcagctataa	caaacatgtc	tcttgcatca	aaagagattc	agattcgtgg	agaaattcat	1860
	tcaaagtgtg	attacacagt	tgacctctta	aacgcttcag	acttcagaga	aaacaagatc	1920
	cacactgggt	ggctggatac	aagaatagct	atgcgtgttc	aagctgagag	gccccatgg	1980
20	tatatctcag	tggttggagg	tgctttatat	aaaacagtaa	ccaccaatgc	agccactgtt	2040
	tctgaatatg	ttagttatct	caccaagggc	catattccac	caaagcatat	atcccttgct	2100
	aattctacag	ttaatttgaa	tatagaaggg	agcaaataca	caattgaaac	tgttaaggact	2160
	ggacatggta	gctacagggt	gagaatgaat	gattcaacag	ttgaagcgaa	tgtacaatct	2220
	ttatgtgatg	gtggcctctt	aatgcagttg	gatggaaaca	gccatgtaat	ttatgcagaa	2280
25	gaagaagctg	gtggtacacg	gcttcagatt	gatggaaaga	catgtttatt	gcagaatgac	2340
	catgatccat	cgaagttatt	agctgagaca	ccctgcaaac	ttcttcgttt	cttggttgct	2400
	gatggtgctc	atgttgatgc	ggatgtacca	tacgcggaag	ttgaggttat	gaagatgtgc	2460
	atgcctctct	tgtcacctgc	ttctggtgtc	attcattgta	tgatgtctga	gggccaggca	2520
	ttgcaggctg	gtgatcttat	agcaagggtg	gatcttgatg	acccttctgc	tgtgaaaaga	2580
30	gctgagccat	ttgatggaat	atttccacaa	atggagctcc	ctgttgctgt	ctctagtcaa	2640
	gtacacaaaa	gatatgctgc	aagtttgaat	gctgctcgaa	tggtccttgc	aggatatgag	2700
	cacaatatta	atgaagtcgt	tcaagatttg	gtatgctgcc	tggaacaacc	tgagcttcct	2760
	ttcctacagt	gggatgaact	tatgtctgtt	ctagcaacga	ggcttccaag	aaatctcaag	2820
	agtgaagttag	aggataaata	caaggaatac	aagttgaatt	tttaccatgg	aaaaaacgag	2880
35	gactttccat	ccaagttgct	aagagacatc	attgaggaaa	atctttctta	tggttcagag	2940
	aaggaaaagg	ctacaaatga	gaggcttggt	gagcctctta	tgaacctact	gaagtcatat	3000
	gagggtggga	gagagagcca	tgacattttt	gttgtcaagt	ctcttttcga	ggagtatctt	3060
	acagtggaa	aacttttttag	tgatggcatt	cagtctgacg	tgattgaaac	attgcggcat	3120
	cagcacagta	aagacctgca	gaaggttgta	gacattgtgt	tgtctcacca	gggtgtgagg	3180
40	aacaaagcta	agcttgtaac	ggcacttatg	gaaaagctgg	tttatccaaa	tcctggtggg	3240

tacagggatc	tgtagttcg	cttttcttcc	ctcaatcata	aaagatatta	taagttggcc	3300
cttaaagcaa	gtgaacttct	tgaacaaacc	aaactaagt	aactccgtgc	aagcgttgca	3360
agaagccttt	cggatctggg	gatgcataag	ggagaaatga	gtattaagga	taacatggaa	3420
gatttagtct	ctgccccatt	acctgttgaa	gatgctctga	tttctttgtt	tgattacagt	3480
5gatcgaactg	ttcagcagaa	agtgattgag	acatacatat	cacgattgta	ccagcctcat	3540
cttgtaaagg	atagcatcca	aatgaaattc	aaggaatctg	gtgctattac	tttttgggaa	3600
ttttatgaag	ggcatgttga	tactagaaat	ggacatgggg	ctattattgg	tggaagcga	3660
tggggtgcca	tggtcgttct	caaactcactt	gaatctgcgt	caacagccat	tgtggctgca	3720
ttaaaggatt	cggcacagtt	caacagctct	gagggcaaca	tgatgcacat	tgcatatttg	3780
10agtgctgaaa	atgaaagtaa	tataagtggg	ataagcagt	atgatcaagc	tcaacataag	3840
atggaaaagc	ttagcaagat	actgaaggat	actagcgttg	caagtgatct	ccaagctgct	3900
ggtttgaagg	ttataagttg	cattgttcaa	agagatgaag	ctcgcatgcc	aatgcgccac	3960
acattcctct	ggttgatga	caagagttgt	tatgaagaag	agcagattct	ccggcatgtg	4020
gagcctcccc	tctctacact	tcttgaattg	gataagttga	aggtgaaagg	atacaatgaa	4080
15atgaagtata	ctccttcgcg	tgaccgccaa	tggcatatct	acacactaag	aaatactgaa	4140
aacccccaaa	tggtgcatag	ggtgtttttc	cgaactattg	tcaggcaacc	caatgcaggc	4200
aacaagttta	gatcggctca	gatcagcgac	gctgaggtag	gatgtcccga	agaatctctt	4260
tcattttacat	caaatagcat	cttaagatca	ttgatgactg	ctattgaaga	attagagctt	4320
catgcaatta	ggacaggtga	ttctcacatg	tatttgtgca	tactgaaaga	gcaaaagctt	4380
20cttgacctca	ttccattttc	agggagtaca	attgttgatg	ttggccaaga	tgaagctacc	4440
gcttggtcac	ttttaaaatc	aatggctttg	aagatacatg	agcttggttg	tgcaaggatg	4500
catcatctgt	ctgtatgcc	gtgggaggtg	aaactcaagt	tgactgtga	tgccctgca	4560
agtggtagct	ggagagttgt	aactacaaat	gttactggtc	acacctgcac	cattgatata	4620
taccgagaag	tggaggaaat	agaatcacag	aagttagtgt	accattcagc	cagttcgtca	4680
25gctggaccat	tgcatggtgt	tgcactgaat	aatccatata	aacctttgag	tgtgattgat	4740
ctaaagcgct	gctctgctag	gaacaacaga	acaacataat	gctatgattt	tccgctggcc	4800
tttgaaactg	cactgcagaa	gtcatggcag	tccaatggct	ctactgtttc	tgaaggcaat	4860
gaaaatagta	aatcctacgt	gaaggcaact	gagctagtgt	ttgctgaaaa	acatgggtcc	4920
tggggcactc	ctataattcc	gatggaacgc	cctgctgggc	tcaacgacat	tggtatggtc	4980
30gcttgatca	tggagatgtc	aacacctgaa	tttcccaatg	gcaggcagat	tattgttgta	5040
gcaaatgata	tcactttcag	agctggatca	tttggcccaa	gggaagatgc	attttttgaa	5100
actgtcacta	acctggcttg	cgaaaggaaa	cttctcttta	tatacttggc	agcaaaactct	5160
ggtgctagga	ttggcatagc	tgatgaagta	aaatcttgct	tccgtgttgg	atggtctgac	5220
gaaggcagtc	ctgaacgagg	gtttcagtac	atctatctga	ctgaagaaga	ctatgctcgc	5280
35attagctctt	ctgttatagc	acataagctg	gagctagata	gtggtgaaat	taggtggatt	5340
attgactctg	ttgtgggcaa	ggaggatggg	cttgggtgctg	agaacataca	tggaagtgtc	5400
gctattgcca	gtgcttattc	tagggcatat	gaggagacat	ttacacttac	atttgtgact	5460
gggaggactg	taggaatagg	agcttatctt	gctcgacttg	gtatacggtg	catacagcgt	5520
cttgaccagc	ctattatttt	aacagggttt	tctgccctga	acaagctcct	tgggcgggaa	5580
40gtgtacagct	cccacatgca	gcttggtggt	cctaagatca	tggcgaccaa	tggtgttgct	5640

```

cacctcactg ttccagatgt cettgaaggt gtttccaata tattgaggtg gctcagctat 5700
gttcctgcaa acattggtgg acctcttcct attaccaaac ctctggaccc tccagacaga 5760
cctggttgctt acatccctga gaacacatgc gatccacgtg cagctatctg tgggtgtagat 5820
gacagccaag ggaaatgggt ggggtggtatg tttgacaaag acagctttgt ggagacattt 5880
5gaaggatggg caaaaacagt ggttactggc agagcaaagc ttggaggaat tcctgtgggc 5940
gtcatagctg tggagacaca gaccatgatg cagatcatcc ctgctgatcc aggtcagctt 6000
gattcccatg agcgatctgt ccctcgtgct ggacaagtgt ggttcccaga ttctgcaacc 6060
aagaccgctc aggcattatt agacttcaac cgtgaaggat tgccctctgtt catcctggct 6120
aattggagag gcttctctgg tggacaaaga gatctctttg aagggaattct tcaggctggg 6180
10tcaacaattg tcgagaacct taggacatat aatcagcctg cttttgtgta cattcctatg 6240
gctggagagc ttcgtggagg agcttgggtt gtggtcgata gcaaaataaa tccagaccgc 6300
attgagtgtt atgctgaaag gactgccaaa ggtaatgttc tcgaacctca agggttaatt 6360
gaaatcaagt tcaggtcaga ggaactccaa gactgtatgg gtaggcttga cccagagttg 6420
ataaatctga aagcaaaact ccaagatgta aatcatggaa atggaagtct accagacata 6480
15gaagggattc ggaagagtat agaagcacgt acgaaacagt tgctgccttt atatacccgag 6540
attgcaatac ggtttgctga attgcatgat acttccctaa gaatggcagc taaaggtgtg 6600
attaagaaag ttgtagactg ggaagaatca cgctcgttct totataaaag gctacggagg 6660
aggatcgag aagatgttct tgcaaaagaa ataaggcaga tagtcggtga taaattttacg 6720
caccaattag caatggagct catcaaggaa tggtagcttg cttctcaggc cacaacagga 6780
20agcactggat gggatgacga tgatgctttt gttgcctgga aggacagtcc tgaaaactac 6840
aaggggcata tccaaaagct tagggctcaa aaagtgtctc attcgctctc tgatcttgct 6900
gactccagtt cagatctgca agcattctcg cagggtcttt ctacgctatt agataagatg 6960
gatccctctc agagagcgaa gtttgctcag gaagtcaaga aggtccttga ttgatgatac 7020
caacacatcc aacacaatgt gtgcatgtca catctttttg ttctagtaca tacatagaag 7080
25gatattgctt ggtcttgatt gatcatgtct gatttaagtc gactattatt tcttggaatt 7140
ttctttttgga cctggtgcta tggttgatgg atgtatatg gatatgtgcg ttctgccagg 7200
tgtaagcaca aaggtttaga carammrara rcaagagcga gtgaacctgt tctggttttg 7260
cagtggttca gtaaggcaga aagttgttaa accgtagttc tgagatgtat taccagtgnc 7320
gccatgctgt acttttaggg tgtataatgc ggatacaaat aaacaattta gcggttcatt 7380
30aaagtttgaa ctcaaataac atgttctttg taagcatatg taccgtacct ctacgtgaaa 7440
taaagttggt gaattagcat tcgaaaaaaa 7470

```

35 <210> 6
 <211> 2325
 <212> PRT
 <213> Zea mays

40 <400> 6

Met Ser Gln Leu Gly Leu Ala Ala Ala Ala Ser Lys Ala Leu Pro Leu
 1 5 10 15
 Leu Pro Asn Arg Gln Arg Ser Ser Ala Gly Thr Thr Phe Ser Ser Ser
 20 25 30
 5 Ser Leu Ser Arg Pro Leu Asn Arg Arg Lys Ser His Thr Arg Ser Leu
 35 40 45
 Arg Asp Gly Gly Asp Gly Val Ser Asp Ala Lys Lys His Ser Gln Ser
 50 55 60
 Val Arg Gln Gly Leu Ala Gly Ile Ile Asp Leu Pro Ser Glu Ala Pro
 10 65 70 75 80
 Ser Glu Val Asp Ile Ser His Gly Ser Glu Asp Pro Arg Gly Pro Thr
 85 90 95
 Asp Ser Tyr Gln Met Asn Gly Ile Ile Asn Glu Thr His Asn Gly Arg
 100 105 110
 15 His Ala Ser Val Ser Lys Val Val Glu Phe Cys Ala Ala Leu Gly Gly
 115 120 125
 Lys Thr Pro Ile His Ser Ile Leu Val Ala Asn Asn Gly Met Ala Ala
 130 135 140
 Ala Lys Phe Met Arg Ser Val Arg Thr Trp Ala Asn Asp Thr Phe Gly
 20 145 150 155 160
 Ser Glu Lys Ala Ile Gln Leu Ile Ala Met Ala Thr Pro Glu Asp Met
 165 170 175
 Arg Ile Asn Ala Glu His Ile Arg Ile Ala Asp Gln Phe Val Glu Val
 180 185 190
 25 Pro Gly Gly Thr Asn Asn Asn Asn Tyr Ala Asn Val Gln Leu Ile Val
 195 200 205
 Gly Met Ala Gln Lys Leu Gly Val Ser Ala Val Trp Pro Gly Trp Gly
 210 215 220
 His Ala Ser Glu Asn Pro Glu Leu Pro Asp Ala Leu Thr Ala Lys Gly
 30 225 230 235 240
 Ile Val Phe Leu Gly Pro Pro Ala Ser Ser Met Asn Ala Leu Gly Asp
 245 250 255
 Lys Val Gly Ser Ala Leu Ile Ala Gln Ala Ala Gly Val Pro Thr Leu
 260 265 270
 35 Ala Trp Ser Gly Ser His Val Glu Val Pro Leu Glu Cys Cys Leu Asp
 275 280 285
 Ala Ile Pro Glu Glu Met Tyr Arg Lys Ala Cys Val Thr Thr Thr Glu
 290 295 300
 Glu Ala Val Ala Ser Cys Gln Val Val Gly Tyr Pro Ala Met Ile Lys
 40 305 310 315 320

Val Gln Ala Glu Arg Pro Pro Trp Tyr Ile Ser Val Val Gly Gly Ala
 645 650 655
 Leu Tyr Lys Thr Val Thr Thr Asn Ala Ala Thr Val Ser Glu Tyr Val
 660 665 670
 5 Ser Tyr Leu Thr Lys Gly His Ile Pro Pro Lys His Ile Ser Leu Val
 675 680 685
 Asn Ser Thr Val Asn Leu Asn Ile Glu Gly Ser Lys Tyr Thr Ile Glu
 690 695 700
 Thr Val Arg Thr Gly His Gly Ser Tyr Arg Leu Arg Met Asn Asp Ser
 10 705 710 715 720
 Thr Val Glu Ala Asn Val Gln Ser Leu Cys Asp Gly Gly Leu Leu Met
 725 730 735
 Gln Leu Asp Gly Asn Ser His Val Ile Tyr Ala Glu Glu Glu Ala Gly
 740 745 750
 15 Gly Thr Arg Leu Gln Ile Asp Gly Lys Thr Cys Leu Leu Gln Asn Asp
 755 760 765
 His Asp Pro Ser Lys Leu Leu Ala Glu Thr Pro Cys Lys Leu Leu Arg
 770 775 780
 Phe Leu Val Ala Asp Gly Ala His Val Asp Ala Asp Val Pro Tyr Ala
 20 785 790 795 800
 Glu Val Glu Val Met Lys Met Cys Met Pro Leu Leu Ser Pro Ala Ser
 805 810 815
 Gly Val Ile His Cys Met Met Ser Glu Gly Gln Ala Leu Gln Ala Gly
 820 825 830
 25 Asp Leu Ile Ala Arg Leu Asp Leu Asp Asp Pro Ser Ala Val Lys Arg
 835 840 845
 Ala Glu Pro Phe Asp Gly Ile Phe Pro Gln Met Glu Leu Pro Val Ala
 850 855 860
 Val Ser Ser Gln Val His Lys Arg Tyr Ala Ala Ser Leu Asn Ala Ala
 30 865 870 875 880
 Arg Met Val Leu Ala Gly Tyr Glu His Asn Ile Asn Glu Val Val Gln
 885 890 895
 Asp Leu Val Cys Cys Leu Asp Asn Pro Glu Leu Pro Phe Leu Gln Trp
 900 905 910
 35 Asp Glu Leu Met Ser Val Leu Ala Thr Arg Leu Pro Arg Asn Leu Lys
 915 920 925
 Ser Glu Leu Glu Asp Lys Tyr Lys Glu Tyr Lys Leu Asn Phe Tyr His
 930 935 940
 Gly Lys Asn Glu Asp Phe Pro Ser Lys Leu Leu Arg Asp Ile Ile Glu
 40 945 950 955 960

Val Ala Ser Asp Leu Gln Ala Ala Gly Leu Lys Val Ile Ser Cys Ile
 1285 1290 1295
 Val Gln Arg Asp Glu Ala Arg Met Pro Met Arg His Thr Phe Leu Trp
 1300 1305 1310
 5 Leu Asp Asp Lys Ser Cys Tyr Glu Glu Glu Gln Ile Leu Arg His Val
 1315 1320 1325
 Glu Pro Pro Leu Ser Thr Leu Leu Glu Leu Asp Lys Leu Lys Val Lys
 1330 1335 1340
 Gly Tyr Asn Glu Met Lys Tyr Thr Pro Ser Arg Asp Arg Gln Trp His
 10 1345 1350 1355 1360
 Ile Tyr Thr Leu Arg Asn Thr Glu Asn Pro Lys Met Leu His Arg Val
 1365 1370 1375
 Phe Phe Arg Thr Ile Val Arg Gln Pro Asn Ala Gly Asn Lys Phe Arg
 1380 1385 1390
 15 Ser Ala Gln Ile Ser Asp Ala Glu Val Gly Cys Pro Glu Glu Ser Leu
 1395 1400 1405
 Ser Phe Thr Ser Asn Ser Ile Leu Arg Ser Leu Met Thr Ala Ile Glu
 1410 1415 1420
 Glu Leu Glu Leu His Ala Ile Arg Thr Gly His Ser His Met Tyr Leu
 20 1425 1430 1435 1440
 Cys Ile Leu Lys Glu Gln Lys Leu Leu Asp Leu Ile Pro Phe Ser Gly
 1445 1450 1455
 Ser Thr Ile Val Asp Val Gly Gln Asp Glu Ala Thr Ala Cys Ser Leu
 1460 1465 1470
 25 Leu Lys Ser Met Ala Leu Lys Ile His Glu Leu Val Gly Ala Arg Met
 1475 1480 1485
 His His Leu Ser Val Cys Gln Trp Glu Val Lys Leu Lys Leu Asp Cys
 1490 1495 1500
 Asp Gly Pro Ala Ser Gly Thr Trp Arg Val Val Thr Thr Asn Val Thr
 30 1505 1510 1515 1520
 Gly His Thr Cys Thr Ile Asp Ile Tyr Arg Glu Val Glu Glu Ile Glu
 1525 1530 1535
 Ser Gln Lys Leu Val Tyr His Ser Ala Ser Ser Ser Ala Gly Pro Leu
 1540 1545 1550
 35 His Gly Val Ala Leu Asn Asn Pro Tyr Gln Pro Leu Ser Val Ile Asp
 1555 1560 1565
 Leu Lys Arg Cys Ser Ala Arg Asn Asn Arg Thr Thr Tyr Cys Tyr Asp
 1570 1575 1580
 Phe Pro Leu Ala Phe Glu Thr Ala Leu Gln Lys Ser Trp Gln Ser Asn
 40 1585 1590 1595 1600

	Gly	Ser	Thr	Val	Ser	Glu	Gly	Asn	Glu	Asn	Ser	Lys	Ser	Tyr	Val	Lys	
					1605					1610					1615		
	Ala	Thr	Glu	Leu	Val	Phe	Ala	Glu	Lys	His	Gly	Ser	Trp	Gly	Thr	Pro	
					1620					1625					1630		
5	Ile	Ile	Pro	Met	Glu	Arg	Pro	Ala	Gly	Leu	Asn	Asp	Ile	Gly	Met	Val	
					1635					1640					1645		
	Ala	Trp	Ile	Met	Glu	Met	Ser	Thr	Pro	Glu	Phe	Pro	Asn	Gly	Arg	Gln	
					1650					1655					1660		
	Ile	Ile	Val	Val	Ala	Asn	Asp	Ile	Thr	Phe	Arg	Ala	Gly	Ser	Phe	Gly	
10	1665					1670					1675					1680	
	Pro	Arg	Glu	Asp	Ala	Phe	Phe	Glu	Thr	Val	Thr	Asn	Leu	Ala	Cys	Glu	
						1685					1690					1695	
	Arg	Lys	Leu	Pro	Leu	Ile	Tyr	Leu	Ala	Ala	Asn	Ser	Gly	Ala	Arg	Ile	
					1700					1705					1710		
15	Gly	Ile	Ala	Asp	Glu	Val	Lys	Ser	Cys	Phe	Arg	Val	Gly	Trp	Ser	Asp	
					1715					1720					1725		
	Glu	Gly	Ser	Pro	Glu	Arg	Gly	Phe	Gln	Tyr	Ile	Tyr	Leu	Thr	Glu	Glu	
					1730					1735					1740		
	Asp	Tyr	Ala	Arg	Ile	Ser	Ser	Ser	Val	Ile	Ala	His	Lys	Leu	Glu	Leu	
20	1745					1750					1755					1760	
	Asp	Ser	Gly	Glu	Ile	Arg	Trp	Ile	Ile	Asp	Ser	Val	Val	Gly	Lys	Glu	
						1765					1770					1775	
	Asp	Gly	Leu	Gly	Val	Glu	Asn	Ile	His	Gly	Ser	Ala	Ala	Ile	Ala	Ser	
					1780					1785					1790		
25	Ala	Tyr	Ser	Arg	Ala	Tyr	Glu	Glu	Thr	Phe	Thr	Leu	Thr	Phe	Val	Thr	
					1795					1800					1805		
	Gly	Arg	Thr	Val	Gly	Ile	Gly	Ala	Tyr	Leu	Ala	Arg	Leu	Gly	Ile	Arg	
					1810					1815					1820		
	Cys	Ile	Gln	Arg	Leu	Asp	Gln	Pro	Ile	Ile	Leu	Thr	Gly	Phe	Ser	Ala	
30	1825					1830					1835					1840	
	Leu	Asn	Lys	Leu	Leu	Gly	Arg	Glu	Val	Tyr	Ser	Ser	His	Met	Gln	Leu	
						1845					1850					1855	
	Gly	Gly	Pro	Lys	Ile	Met	Ala	Thr	Asn	Gly	Val	Val	His	Leu	Thr	Val	
					1860					1865					1870		
35	Pro	Asp	Val	Leu	Glu	Gly	Val	Ser	Asn	Ile	Leu	Arg	Trp	Leu	Ser	Tyr	
					1875					1880					1885		
	Val	Pro	Ala	Asn	Ile	Gly	Gly	Pro	Leu	Pro	Ile	Thr	Lys	Pro	Leu	Asp	
					1890					1895					1900		
	Pro	Pro	Asp	Arg	Pro	Val	Ala	Tyr	Ile	Pro	Glu	Asn	Thr	Cys	Asp	Pro	
40	1905					1910					1915					1920	

Arg Ala Ala Ile Cys Gly Val Asp Asp Ser Gln Gly Lys Trp Leu Gly
 1925 1930 1935
 Gly Met Phe Asp Lys Asp Ser Phe Val Glu Thr Phe Glu Gly Trp Ala
 1940 1945 1950
 5 Lys Thr Val Val Thr Gly Arg Ala Lys Leu Gly Gly Ile Pro Val Gly
 1955 1960 1965
 Val Ile Ala Val Glu Thr Gln Thr Met Met Gln Ile Ile Pro Ala Asp
 1970 1975 1980
 Pro Gly Gln Leu Asp Ser His Glu Arg Ser Val Pro Arg Ala Gly Gln
 10 1985 1990 1995 2000
 Val Trp Phe Pro Asp Ser Ala Thr Lys Thr Ala Gln Ala Leu Leu Asp
 2005 2010 2015
 Phe Asn Arg Glu Gly Leu Pro Leu Phe Ile Leu Ala Asn Trp Arg Gly
 2020 2025 2030
 15 Phe Ser Gly Gly Gln Arg Asp Leu Phe Glu Gly Ile Leu Gln Ala Gly
 2035 2040 2045
 Ser Thr Ile Val Glu Asn Leu Arg Thr Tyr Asn Gln Pro Ala Phe Val
 2050 2055 2060
 Tyr Ile Pro Met Ala Gly Glu Leu Arg Gly Gly Ala Trp Val Val Val
 20 2065 2070 2075 2080
 Asp Ser Lys Ile Asn Pro Asp Arg Ile Glu Cys Tyr Ala Glu Arg Thr
 2085 2090 2095
 Ala Lys Gly Asn Val Leu Glu Pro Gln Gly Leu Ile Glu Ile Lys Phe
 2100 2105 2110
 25 Arg Ser Glu Glu Leu Gln Asp Cys Met Gly Arg Leu Asp Pro Glu Leu
 2115 2120 2125
 Ile Asn Leu Lys Ala Lys Leu Gln Asp Val Asn His Gly Asn Gly Ser
 2130 2135 2140
 Leu Pro Asp Ile Glu Gly Ile Arg Lys Ser Ile Glu Ala Arg Thr Lys
 30 2145 2150 2155 2160
 Gln Leu Leu Pro Leu Tyr Thr Gln Ile Ala Ile Arg Phe Ala Glu Leu
 2165 2170 2175
 His Asp Thr Ser Leu Arg Met Ala Ala Lys Gly Val Ile Lys Lys Val
 2180 2185 2190
 35 Val Asp Trp Glu Glu Ser Arg Ser Phe Phe Tyr Lys Arg Leu Arg Arg
 2195 2200 2205
 Arg Ile Ala Glu Asp Val Leu Ala Lys Glu Ile Arg Gln Ile Val Gly
 2210 2215 2220
 Asp Lys Phe Thr His Gln Leu Ala Met Glu Leu Ile Lys Glu Trp Tyr
 40 2225 2230 2235 2240

```

Leu Ala Ser Gln Ala Thr Thr Gly Ser Thr Gly Trp Asp Asp Asp Asp
                2245                                2250                    2255

Ala Phe Val Ala Trp Lys Asp Ser Pro Glu Asn Tyr Lys Gly His Ile
                2260                                2265                    2270

5  Gln Lys Leu Arg Ala Gln Lys Val Ser His Ser Leu Ser Asp Leu Ala
                2275                                2280                    2285

Asp Ser Ser Ser Asp Leu Gln Ala Phe Ser Gln Gly Leu Ser Thr Leu
                2290                                2295                    2300

Leu Asp Lys Met Asp Pro Ser Gln Arg Ala Lys Phe Val Gln Glu Val
10 2305                                2310                    2315                    2320

Lys Lys Val Leu Asp
                2325

```

15 <210> 7
 <211> 22
 <212> DNA
 <213> Zea mays

<400> 7
20 ggtcttcaat tgtgctgtct gg 22

25 <210> 8
 <211> 24
 <212> DNA
 <213> Zea mays

<400> 8
ccttgacgaa cagactgqct qtgc 24

30 <210> 9
 <211> 23
 <212> DNA
 <213> Zea mays

35 <400> 9
cacagccagt ctgttcgtca agg

23

40 <210> 10
 <211> 20
 <212> DNA

<213> Zea mays

<400> 10
cctctacgta attggtcagc 20

5

<210> 11
<211> 20
<212> DNA
<213> Zea mays

10

<400> 11
catagctatg gcaactccgg 20

<210> 12
15 <211> 3488
<212> DNA
<213> Zea mays

<220>
20 <221> unsure
<222> (1803) ... (1803)
<223> "n" represents "a" or "g" or "c" or "t"

<221> unsure
25 <222> (3324) ... (3324)
<223> "n" represents "a" or "g" or "c" or "t"

<400> 12
aagcttggtg tggattcgtc agcgccaagc cgggggttttg catgcgcccg actggaarcs 60
30 gaattccgtg agccctgtac rrcaatggca accccasggg tactgggggtg gctgaatggt 120
ctcsgcttac gcaattgttt gtggcagcwg cgtgggctaa atgtargttg tctcttgttg 180
cactgcarga tggatgggta gcctctgggc cgcctctgct artgtctarc gtttgctgac 240
tgtggtttat tcagggatgc ccatgcccac gctagattga taggtgccat tctaattggtg 300
gggtggcggta aggtttatta agctgyagya tcagtaggta acctcatgaa tcaggggttta 360
35 agcacacctt ttcctttgtg tgggtgcata aggaatgcac ttggcttcgt tccctgatag 420
tctttgstca tgtgtcattc taccaagtgg gttactgtaa cattgcactc tatgatgggt 480
gggtggttggt catcttlytg ctycccctgg ytgtctaata cctgcatgta actgatgacc 540
yycytttatg tatcatatag attacatcct tttgtgtgac atctcaattc tgaaaaaaca 600
atgttttgca ttcttagcgc tctgtgcaca aggaaaagga gggttttacct gcaacttttt 660
40 ttttcgagaa aaaacaaacc tttctgaaag gcagtgatca tttagtataa agaaaatttg 720

	atttacttttc	ttcagagaga	atatkccaar	caaacaattt	tcttactgtc	tgagccacga	780
	aatttgatct	tgatcttact	ttcacaagcc	acatgaagcc	ttatcatcgc	tctgataaaa	840
	aarccaaata	ggtgattcat	agaatgagar	aaagaacctg	ttgccatttg	gggaccttgt	900
	tgtgtactca	ttatccccc	tgctcagggt	gagggttccct	tgccactgcc	acccttggc	960
5	cccttcttat	acaacctatct	ccattgaaaa	agatttttgca	ctacatttgg	gcttcgtatg	1020
	acaaaaaagg	aaaataaaaac	taaacagcag	aaacatagta	taattatagg	taaaagggttc	1080
	tggcaagttt	gagtgggaaga	gacctttgta	tatttggaca	tatttcacta	gtaaatagtt	1140
	ttctaaaatc	ttcatgaatg	gtggccaata	aacttgataa	gatctcaaca	tggcagggttc	1200
	cttcmaaata	agaggaaaac	tggaaacatc	acaaatattt	tttagcgagt	ggcctataaa	1260
10	ttataatggt	gctttcattt	ctttgatatt	caaaacttcc	taagagtatt	ctgctagagc	1320
	tctgatgggt	tcttttgcct	ctgtcagatt	ttccaggagt	tttcttccct	ttttatggca	1380
	ctgtgcgttt	gagaaggctc	tcaattgtgc	tgtctggggc	acggaacgac	aatgtcacag	1440
	cttgatttag	ccgcagctgc	ctcaaaggcc	ttgccactac	tccctaatac	ccagagaagt	1500
	tcagctggga	ctacattctc	atcatcttca	ttatcgaggc	ccttaaacag	aaggaaaagc	1560
15	catactcggt	cactccgtga	tggcggagat	gggggtatcag	atgccaaaaa	gcacagccag	1620
	tctgttcgtc	aaggactgtg	gaatatcttt	tgatacaagc	taaaattttg	ctacagaata	1680
	tatatattaaa	gagttctttc	ttggctggtg	ttgtttattt	gtttaacats	cgaaagggcc	1740
	tctagttaga	ttggttaggt	ggsctgaata	ccactcctta	aggtcttgag	tttgcttttc	1800
	ccnccggagc	gaatttttag	ctagggttac	ccccccaccc	ccaccgaat	ctgcacagyc	1860
20	cggycgyggy	cgycctcata	taggctacga	tgtcattgtg	tatcggcggg	ccaggggttt	1920
	aagagttttc	ttgacctttg	ttagaagatc	ttaataatac	aatgtccaag	ggctgtctta	1980
	ccctgtaggt	cgagttttta	gttgttttta	catggtaatg	tttgaagcct	cattctaggt	2040
	rccaatatag	atatgctcac	tgctcagttt	caaagtgttg	tctgcatgta	ggtcttgctg	2100
	gcattatcga	cctcccaagt	gaggcacctt	ccgaagtggg	tatttcacag	taaggactac	2160
25	aatattttgc	gtacgtttgt	tttggaaaaa	gaaaatattc	tcagcttatt	tatactagct	2220
	tcgctaatac	tgaaatgctg	tcttaatgtc	ctgggtgctg	atgctcaatc	tttcatagta	2280
	aatgctgcaa	aatatgtgat	gtaactgttg	caacacagcc	agggacctgt	tatttagagc	2340
	atgggtgaat	ctctggttca	gttatatgat	gtagttatag	ctcatgttga	agaattagtt	2400
	gcagtgtttg	ctggacaatg	gtcacttatt	ataaatcata	tctgcataca	catttgtgac	2460
30	ttctgttgct	gtaaatgccc	gcattttttg	agaaaaattt	aaatgcttgg	cctaaattgg	2520
	acatatatga	tagacaaagc	tgatttgaac	tttgtttatt	tttgacatcc	atgcatattg	2580
	tcagtgttgt	gaaaacaata	ctaatacctt	ttttttgtct	ttttccagtg	gatctgagga	2640
	tcctaggggg	ccaacagatt	cttatcaaata	gaatgggatt	atcaatgaaa	cacataatgg	2700
	aagacatgcc	tcagtgtcca	agggtgttga	atgttgtgct	gcactagggt	gcaaaacacc	2760
35	aattcacagt	atattagtg	ccaacaatgg	aatggcagca	cccaaattta	tgaggagtgt	2820
	ccggacatgg	gctaatagata	cttttggatc	tgagaaggca	attcaactca	tagctatggc	2880
	aactccggaa	gacatgagga	taaatgcaga	acacattaga	attgctgacc	aattacgtag	2940
	agggtgctgg	tggaaacaaac	aataataact	acgccaatgt	tcaactcata	gtggagggtta	3000
	gccttgctaa	tctgttagtt	tactactggg	ctgctgtttc	ctttatttgt	tgtataatga	3060
40	ttgacatatt	taagtagaga	aatttatatt	tctcctctgc	tgttgtggaa	gtccaattgt	3120

17

catcattaac tgtgaaatat tgcagatggc acaaaaaacta ggtgtttctg ctgtttggcc 3180
 tggttggggg catgcttctg agaatcctga actgccagat gcattgaccg caaaagggat 3240
 cgtttttctt ggcccacctg satcatcaat gaatgctttg ggagataagg tcggctcagc 3300
 tctcattgct caagcagccg gggncccaac tcttgcttgg agtggatcac atgtgagtct 3360
 5 cactctttga ttactatccg cctgtctcat tgctctctct ttcatttct aatgacacta 3420
 aatttaggtt gaagttccat tagagtgtg cttagacgag atacctgagg agatgtatag 3480
 aaaagctt 3488

<210> 13

10 <211> 1328

<212> DNA

<213> Zea mays

<220>

15 <221> unsure

<222> (1024) ... (1024)

<223> "n" represents "a" or "g" or "c" or "t"

<400> 13

20 gaattccgtg agccctgtac ggcaatggca accccagggg tactgggggtg gctgaatggg 60
 ctcggttac gcaattgttt gtggcagctg cgtgggctaa atgtagggtg tctcttgttg 120
 cactgcagga tggatgggta gcctctgggc cgcctctgct agtgtctagc gttgctgact 180
 gtggtttatt cagggatgcc catgcccattg ctagattgat aggtcatagg tgccattcta 240
 atggtaggtg gcggttaagg ttattaagct gtcgtatcag taggtaacct catgaatcag 300
 25 ggtttaagcc caccctctcc tttgtgtggg tgcataagga atgcacttgg cttcgttccc 360
 tgctagtctt tgctcatgtg tcattctacc aagtgggtta ctgtaacatt gcactctatg 420
 atggttgggtg gttgtgcac tttttgcttc ccttggttgt ctaataacctg catgtaactg 480
 atgaccttct tttatgtatc atatagatta catcttttgt tgtacatctc aattctgaaa 540
 aacaatgttt tgcattctta gcgtctgtg cacaaggaaa aggaggtttt acctgcaact 600
 30 ttttttttcg agaaaaaaca aacctttctg aaaggcagt atcatttagt ataaagaaaa 660
 tttgatttac tttcttcaga gagaatatc caaacaaca attttcttac agtctgagcc 720
 acgaaatttg atcttgatct tactttcaca agccacatga agccttatca tcgctctgat 780
 aaaaaacca aataggtgat tcatagaatg agaaaaagaa cctgttgcca tttggggacc 840
 ttgttgtgta ctcatatcc cccctgctca ggttgaggtt tccttgccac tgccaccctt 900
 35 tggccccttc ttatacaacc atctccattg aaaaagattt tgcactacat ttgggcttcg 960
 tataacaaaa aaggaaaata aaactaaaca gcagaaacat agtataatta taggtaaaag 1020
 gttntggcaa gtttgagtgg tagagacctt tgtatatattg gacatatctc actagtaaat 1080
 agttttctaa aatgttcatg aatgggtggc aataaacttg ataagatctc aacatggcag 1140
 gttccttcaa aatgagagga aaactggaaa catcacaat attttttagc gagtggccta 1200
 40 taaattataa tgttgctttc atttctttga tattcaaac ttcctaagag tattctgcta 1260

gagctctgat ggtgtctttt gcctctgtca gattttccag gagttttctt ccctttttat 1320
ggcactgt 1328

- 5 <210> 14
 <211> 1565
 <212> DNA
 <213> Zea mays
- 10 <220>
 <221> unsure
 <222> (22) ... (22)
 <223> "n" represents "a" or "g" or "c" or "t"
- 15 <221> unsure
 <222> (44) ... (44)
 <223> "n" represents "a" or "g" or "c" or "t"
- 20 <221> unsure
 <222> (90) ... (90)
 <223> "n" represents "a" or "g" or "c" or "t"
- 25 <221> unsure
 <222> (982) ... (982)
 <223> "n" represents "a" or "g" or "c" or "t"
- 30 <221> unsure
 <222> (1064) ... (1064)
 <223> "n" represents "a" or "g" or "c" or "t"
- 35 <221> unsure
 <222> (1112) ... (1112)
 <223> "n" represents "a" or "g" or "c" or "t"
- 40 <221> unsure
 <222> (1177) ... (1177)
 <223> "n" represents "a" or "g" or "c" or "t"
- <221> unsure
 <222> (1238) ... (1238)
 <223> "n" represents "a" or "g" or "c" or "t"

- <221> unsure
<222> (1244)...(1244)
<223> "n" represents "a" or "g" or "c" or "t"
- 5 <221> unsure
 <222> (1256)...(1256)
 <223> "n" represents "a" or "g" or "c" or "t"
- 10 <221> unsure
 <222> (1279)...(1279)
 <223> "n" represents "a" or "g" or "c" or "t"
- 15 <221> unsure
 <222> (1288)...(1288)
 <223> "n" represents "a" or "g" or "c" or "t"
- 20 <221> unsure
 <222> (1305)...(1305)
 <223> "n" represents "a" or "g" or "c" or "t"
- 25 <221> unsure
 <222> (1308)...(1308)
 <223> "n" represents "a" or "g" or "c" or "t"
- 30 <221> unsure
 <222> (1311)...(1311)
 <223> "n" represents "a" or "g" or "c" or "t"
- 35 <221> unsure
 <222> (1325)...(1325)
 <223> "n" represents "a" or "g" or "c" or "t"
- 40 <221> unsure
 <222> (1344)...(1344)
 <223> "n" represents "a" or "g" or "c" or "t"
- <221> unsure
 <222> (1365)...(1365)
 <223> "n" represents "a" or "g" or "c" or "t"

<221> unsure
<222> (1372)...(1372)
<223> "n" represents "a" or "g" or "c" or "t"

5 <221> unsure
<222> (1393)...(1393)
<223> "n" represents "a" or "g" or "c" or "t"

10 <221> unsure
<222> (1405)...(1405)
<223> "n" represents "a" or "g" or "c" or "t"

15 <221> unsure
<222> (1410)...(1410)
<223> "n" represents "a" or "g" or "c" or "t"

20 <221> unsure
<222> (1453)...(1453)
<223> "n" represents "a" or "g" or "c" or "t"

25 <221> unsure
<222> (1484)...(1484)
<223> "n" represents "a" or "g" or "c" or "t"

30 <221> unsure
<222> (1491)...(1491)
<223> "n" represents "a" or "g" or "c" or "t"

35 <221> unsure
<222> (1508)...(1508)
<223> "n" represents "a" or "g" or "c" or "t"

<221> unsure
<222> (1539)...(1539)
<223> "n" represents "a" or "g" or "c" or "t"

<400> 14
agcatccctt gggattgtga tnactcacat aaattcttgc gaantgttga cattctagtg 60
atttgagttc cggttctagtg tgctagtcan ttgagctcaa gtcttggttt tatgtgtgcg 120
40 tattcactgt gatctttgtg tcgtgtgtga gttgttgatc cttcccttgc tccgtgattc 180

```

tttgtgaaat cttttgaaag ggcgagaggg tccaagctgt ggagattcct cgcaagtggg      240
attaagaaaa gcaaagcaac accgtgggtat tcaagttggg ctttggaccg cttgagaggg      300
gttgattgca accctcgtcc gttgggacgc cacaacgtgg agtaggcaag cgttgggtctt      360
ggccgaacca cgggataacc accgtgccat ctctgtgatt gatatctctt ggttattgtg      420
5 ttgtgttgag atccttctct agccacttgg caaattactg tgctaacaat taatcaagtt      480
ttgtggctta agattttgaa gtattacagg atctgcatca tggctctgtgt ctccacagct      540
atgacaccca caggaattca tgtgttcctt ggagccactc ttggatgacc taaaggaatt      600
atctctaacc ggcttgtaga catatgatgc atcaagagat gagtgtttta ctatgcgagg      660
ggccatgctt atgaccataa gtaatcttcc tggtttagaa atgcttgctt ctcatatggt      720
10 tcatgggaaa ttcgcatgcc tccttgtggg gaaaatgtct ggacaaaaca gctgaagaat      780
ggtcgtaaat cttgttttat gggaaatcgc caatatattg atcttgatca ttcttattgc      840
ttggatgcag actccgtttg atggaacgat agacttcgaa caaacctaa aacctattat      900
gatcgtccaa ttttggatga aatcatcaca cttggtgatt tcaagaactc aaaaayttac      960
agttaattgg atataggagg gngcaaaaac acagtaagtt ggacattcca taaggggatt     1020
15 tatttttagtt gacaataaag tagatgggca tcatcctgag tttngtttgg catcgtgtcg     1080
tagattgaaa ctgtaaggat ggacatggta gntaacagggt tgagatgaat gattcaacag     1140
ttgaagcgaa tgtacaatct ttatgtgatg gtggctntta atgcaggtaa ctagtttttt     1200
tttatgcttt attattaatt agttggataa atggtttnga tttntgattg ttaaantgca     1260
atggctccag ttggatggna acagccangt aatttatgca gaagnagnag ntgggtgtac     1320
20 acggnnttcag attgatggaa aganatgttt attgcaggta aatantccct tnttccttta     1380
tatttttgtt gtntgattgt ataantttgn tagattatnt gtataattta ttattgcatt     1440
tcacccact aanttatntt taaaagatgg gttttgttgt ttgnttcagc nggcgacatc     1500
acataagnaa attgtgatta atttttgttt ttttgcagna tgaccatgat ccatcaaagt     1560
tatta

```

25

```

<210> 15
<211> 1168
<212> DNA
<213> Zea mays

```

30

```

<220>
<221> unsure
<222> (60) ... (60)
<223> "n" represents "a" or "g" or "c" or "t"

```

35

```

<221> unsure
<222> (1024) ... (1024)
<223> "n" represents "a" or "g" or "c" or "t"

```

40

```

<221> unsure

```

<222> (1104)...(1104)

<223> "n" represents "a" or "g" or "c" or "t"

<221> unsure

5 <222> (1114)...(1114)

<223> "n" represents "a" or "g" or "c" or "t"

<400> 15

	acataagctg ggttagtagt ggtgaattag tggattattg atctgttgtg ggcaaggagn	60
10	tggcttggtg tgcagaaaaa acatggaagt gtgtattgcc agtgcttatt ytrgggatat	120
	gagggaattw amattacatt tgtgactggg cggactgtag gataggagtt atcttgytcg	180
	attggtatac ggtgcataca gskyttgacc agctattatt ttaacagggt tttctgcctt	240
	gaacaagtcc ttgggcgga agtgtagc tcccacatgc agcttggtgg tcctaagatc	300
	atggcgacca atgggtgtgt ccacctcact gttccagatg acctgaagg tgtttccaat	360
15	atattgaggt ggctcagcta tgttcctgca aacattggtg gacctcttcc tattacaaa	420
	cctctggacc ctccagacag acctgttgct tacatccctg agaacacatg cgatccacgt	480
	gcagctatct gtggtgtaga tgacagccaa gggaaatggt tgggtggtat gtttgacaaa	540
	gacagctttg tggagacatt tgaaggatgg gcaaaaacag tggttactgg cagagcaaag	600
	cttgaggaa ttccatgcat cttaataaac acagttggcc cttaaagcaa gtgaacttct	660
20	tgaacaaacc aaactaagt aactccgtgc aagcgttgca agaagccttt cggatctggg	720
	gatgcataag ggagaaatga gtattaagga taacatggaa gatttagtct ctgccccatt	780
	acctgttgaa gatgctctga tttctttgtt tgattacagt gatcgaactg ttcagcagaa	840
	agtgattgag acatacatat cacgattgta ccaggatatta tatcaactaa cttaatgtct	900
	tccatagtct cactaagcat atctgatatg ttagatatac ctacatggaa tgctcatctt	960
25	ttcatttgac acaaagaaac attgagaaat gagatgctga cgattggctg aaattaactg	1020
	ggtntgagaa attgtgatct cccaacttgt taatgcacaa tgttctggct aacttgccaa	1080
	tattttttca gcctcatctt gtnaggata gcanccaaat gaaatccaag gatctggtgc	1140
	tattactttt tgggaattta tgaagggc	1168

30 <210> 16

<211> 638

<212> DNA

<213> Zea mays

35 <220>

<221> unsure

<222> (358)...(358)

<223> "n" represents "a" or "g" or "c" or "t"

40 <221> unsure

<222> (398) ... (398)

<223> "n" represents "a" or "g" or "c" or "t"

<400> 16

5	ctcccaatat	tgtcatgagg	cttgcacccc	aggtagttt	tttttccttt	ctgaaattta	60
	tattccatac	cttttcacct	ttagttatcc	ttgtattttc	tggaagcttc	atctgatgca	120
	ttattgacaa	atgcactaat	ggatcatcata	tttgagatt	aacatattta	tcttaattga	180
	tgggaactct	tgaaaatgac	aatgggttgag	cagataatta	acagtttttt	aataaaaaaa	240
	catgcatttc	taggagttgg	actaagcttt	tcttagtatg	aagtgccatg	ttttacatgg	300
10	tccattttgtg	tcaattttaca	gtcgggtatca	tggaagggtt	gtcataatgg	ctggaganaa	360
	acaacacatc	ttgtttctca	acacttgtgg	gagaagangt	tttacctttt	ttcctaaaaat	420
	tactttttgt	actaaattgt	ataatttttc	caatattctc	catgattatt	gaactctgct	480
	gtgttcaaac	agccaaaaca	tggtttccata	ctttcacact	ttatttttta	gatggaacct	540
	ggaattgtgc	tctgttatct	gtatcatgca	tatattgatc	ttaaacctat	ctctattgta	600
15	gaatccgcac	ttgaattcag	ttgcttgtga	tcaatatg			638

<210> 17

<211> 558

<212> DNA

20 <213> Zea mays

<220>

<221> unsure

<222> (41) ... (41)

25 <223> "n" represents "a" or "g" or "c" or "t"

<221> unsure

<222> (103) ... (103)

<223> "n" represents "a" or "g" or "c" or "t"

30

<400> 17

	ggtaaccacc	acaccgcg	cgcttaatgg	ccgtacaggg	nggtcccatt	cgccattcag	60
	gtgcgcaact	gttgggaagg	gcgatcgggtg	cgggcttctt	cgntattacg	ccagctggcg	120
	aaaggggat	gtgctgcaag	gcgattaagt	tgggtaacgc	cagggttttc	ccagtcacga	180
35	cgttgtaaaa	cgacggccag	tgagcgcgcg	taatacgact	cactataggg	cgaattgggt	240
	accgggcccc	ccctcgaggt	cgacctgcag	gtcaacggat	cctagggggc	caacagattc	300
	ttatcaaatg	aatgggatta	tcaatgaaac	acataatgga	agacatgcct	cagtgtccaa	360
	ggttgttgaa	ttttgtgcgg	cactaggtgg	caaaacacca	attcacagta	tattagtggc	420
	caacaatgga	atggcagcag	caaaatttat	gaggagtgtc	cggacatggg	ctaattgatac	480
40	ttttgatct	gagaaggcaa	ttcaactcat	agctatggca	actccggaag	acatgaggta	540

aatgcagaac acattaga

558

<210> 18

<211> 975

5

<212> DNA

<213> Zea mays

<220>

<221> unsure

10

<222> (784) ... (784)

<223> "n" represents "a" or "g" or "c" or "t"

<400> 18

	gaataatctg cctgcagctc aagttgctgt tggaaatgggc atacctcttt ggcagattcc	60
15	aggtaattac caatttacca acttatttag ttcttatttg ttttattctc taattttcta	120
	cttatgtaga aatcagacgt ttctatggaa tggactatgg aggagggtat gacatttgga	180
	ggaaaacagc agctcttgct acaccattta attttgatga agtagattct caatggccaa	240
	agggccattg tgtagcagtt agaattacta gtgaggaccc agatgatggg ttcaaacctt	300
	ctggtgggaa agtgaaggta agttttctag atgacatgta ttatatatcg ttcaaagaga	360
20	ttaagtttgg ttaaatagact aggtcttgat tttttatctt tcaggagata agttttaaaa	420
	gcaagcctaa tgtttggggt tactttctcag taaaggtaac ttgttaactt tagtacgtg	480
	tcacattatt cttcttcttg aaaataattt gaacggttct ctttgtattt taaccatcca	540
	tcgtctcatt tascagagca cacaaatatt tgcactgacc cccctcccct tatctgcttt	600
	cagtctggtg gaggcattca tgaatttgct gattctcagt tcggtatgtg taaaccaaga	660
25	gtattctttg taatttatat tggctctcaa ttttgaaata ttgctctttc cgttacagga	720
	cawgtttttg catatggggt ctctagatca gcagcaataa caaacatgac tcttgcatta	780
	aaanagattc aaattcgtgg agaaattcat tcaaagtgtt gattacacag ttgatctctt	840
	aatgtttaag aaatattaac caccttttaa atcacatttt ccattatggt tgattccata	900
	tcattaattt tgattttcta ttatggctaa acctgtgggt ctattttcct attatcccag	960
30	gcttccgact ttaga	975

<210> 19

<211> 852

<212> DNA

35

<213> Zea mays

<400> 19

	ggatcctagg gggccaacag attcttatca aatgraatgg gattatcaat gaaacacata	60
	atggaagaca tgcctcagtg tccaagggtt ttgaattttg tgcggcacta ggtggcaaaa	120
40	caccaattca cagtatatta gtggccaaca atggaatggc agcagcaaaa tttatgagga	180

25

gtgtccggac atgggctaata gatacttttg gatctgagaa ggcaattcaa ctcatagcta 240
 tggcaactcc ggaagacatg aggataaatg cagaacacat tagaattgct gaccaattac 300
 gtagargtgc ctggtggaac aaacaataat aactacgcca atgttcaact catagtggaa 360
 gttagccttg ctaatctgtt agtttactac tggctctgctg tttcctttat ttgttgata 420
 5 atgattgaca tatttaagta gagaaattta ttttctcct ctgctgttgtt ggaagtccaa 480
 ttgtcaccat taactgtgaa atattgcaga tggcacaaaa actaggtgtt tctgctgttt 540
 ggctctggttg gggatcatgct tctgagaatc ctgaactgcc agatgcattg accgcaaaag 600
 ggatcgtttt tcttgcccca cctgcatcat caatgaatgc tttgggagat aaggctggct 660
 cagctctcat tgctcaagca gccgggggtcc caactcttgc ttggagtgga tcacatgtga 720
 10 gtctcactct ttgattacta tccgcctgtc tcattgctct ctctttcata ttctaataagac 780
 actaaattta ggttgaagtt ccattagagt gctgcttaga cgcgatacct gaggagatgt 840
 atagaaaagc tt 852

<210> 20
 15 <211> 231
 <212> DNA
 <213> Zea mays
 <220>
 20 <221> unsure
 <222> (125)... (125)
 <223> "n" represents "a" or "g" or "c" or "t"

<400> 20
 25 aattcctgtg ggtgttatag ctgtggagac acagaccatg atgcagctca tccctgctga 60
 tccaggtaaa cttgattccc atgagcgatg tgttctctcg gctggacaag tgtggttccc 120
 agatnctgca accaagacag ctcaggcatt attagacttc aaccgtgaag gattgcctct 180
 gttcatcctg gctaactgga gaggtctctc tgggggacag agagatctct t 231

30 <210> 21
 <211> 207
 <212> DNA
 <213> Zea mays
 35 <220>
 <221> unsure
 <222> (165)... (165)
 <223> "n" represents "a" or "g" or "c" or "t"

40 <400> 21

26

	aattcatgca tcttaataaa cacagttggc ccttaaagca agtgaacttc ttgaacaaac	60
	caaactaagt gaactctgtt ccagcattgc aagaagcctt tcagatctgg ggatgcataa	120
	gggagaaatg actattaagg atagcatgga agatttagtc tctgncccat tgctgttga	180
	agatgctctt atttctttgt ttgatta	207
5		
	<210> 22	
	<211> 180	
	<212> DNA	
	<213> Zea mays	
10		
	<220>	
	<221> unsure	
	<222> (59) ... (59)	
	<223> "n" represents "a" or "g" or "c" or "t"	
15		
	<400> 22	
	atagacctgt cgcatacatc cctgagaaca catgcatcc gcgtgcagcc atccgtggng	60
	tagatgacag ccaagggaaa tggttgggtg gtatgtttga caaagacagc tttgtggaga	120
	catttgaagg atgggcaaaa acagtgggta ctggtagagc aaagcttga ggaaggaatt	180
20		
	<210> 23	
	<211> 3031	
	<212> DNA	
	<213> Avena sativa	
25		
	<400> 23	
	acaaaatagt aaagacctcc agaaggttgt agacattgtt ttgtctcacc aggggtgtgag	60
	aagcaaaacc aagctgatac tcacgctcat ggagaaactg gtctatccaa accctgctgc	120
	ctatagggac cagctgattc gcttttcttc cctcaaccat aaaagatatt acaagctggc	180
30	ccttaaagct agtgaacttc tcgaacaaac caagctcagt gaactccgca caagcattgt	240
	aaggaacctt tcagcgctgg agatgtttac tgaagaaagg gcaggtttct cgttgcaaga	300
	gaagaaattg gccattaatg agagcatggg agatttagtc agtgccccac tgccagttga	360
	agatgcactt atatctttgt ttgattgtat tgatcaaaact cttcagcaga gagtgatcga	420
	gacgtacata tctcgattat accagcctca acttgtgaag gatagtatcc agctgaaata	480
35	tcaggattct gctgttatcg ctttatggga attcaacgaa gggcatcctg ataatagatt	540
	gggtgctatg gttatcctga agtctctaga atctgtgtca acagcaattg gagctgctct	600
	aaaggataga tcacattatg caagctctgc gggtaacaca gtgcatattg ttttgttga	660
	tgctgatact cagacgaata caactgaaga tagtggtgat aacgaccaag ctcaagacag	720
	gatggacaaa ctctctgtaa tactgaaaca agatattgtc acggctgatc tatgtgctgc	780
40	tggtgtcaag gttattagtt gcattgttca aagagatgga gcactcatgc ctatgcgccg	840

	taccttcctc	ttgtctgaag	aaaagctttg	ttatgaggaa	gagccgattc	ttcggcatgt	900
	ggagcctcca	ctttctgcac	ttcttgagtt	ggataaattg	aaagtgaag	gatacaatga	960
	gatgaagtat	acaccgtcac	gtgatcgta	gtggcatata	tacacactta	gaaatactga	1020
	aaaccccaaa	atgttgacac	gggtattttt	ccgaactctt	gtcagacagc	cgagtgtctg	1080
5	caacagggtt	atgtcaggcc	atatcaatga	tgttgaagta	ggacatgctg	aggaatctct	1140
	ttcattttaca	tcaagcagca	tattaagatc	tttgatgact	gctatagaag	aattggagct	1200
	tcacgcgatt	aggactggtc	attctcatat	gtacttgtgc	atattgaaag	agcaaaagct	1260
	tcttgacctt	gttctgtttt	cagggaacac	agttgtggat	gttgccaag	atgaagctac	1320
	tgcgtgctct	cttatgaaag	aaatggctct	aaagatacat	gaacttgttg	gtgcaagaat	1380
10	gcatcatctt	tctgtatgcc	agtgggaagt	gaaacttaag	ttggacagt	atggacctgc	1440
	cagtggtagc	tggagagttg	taacaaccaa	tgtcactagt	cacacctgca	ctgtggatat	1500
	ctaccgggag	gtcgaagata	cagattcaca	gaagctagta	taccactatg	ctgcatcgtc	1560
	acctggctct	ttgcatgggt	tcacattgag	taattcatac	cagcctttga	gcgttattga	1620
	tctaaaacga	tgctctgcc	ggaacaacag	aactacatac	tgctacgatt	ttccattggc	1680
15	atgtgaaact	gcagtgcgga	agtcattggc	taacgtttct	agtgaataca	accaatgtta	1740
	tggttaaagca	acagagctgg	tggttgctga	caagaatggg	tgtgggggca	ctcctatagt	1800
	tcctatgcag	cgtgctgctg	ggctcaatga	catcggtatg	gttgctgga	tcttgacat	1860
	gtccactcct	gaatttccca	gtggcagaca	gattattgtt	atcgcaaatg	atattacttt	1920
	cagagctgga	tcatttggtc	caagggaaga	tgcatTTTTT	gaagctgtta	ccaacctggc	1980
20	ttgtgagaga	aagcttcctc	ttatctacat	ggcagcaaac	tctgggtgctc	ggattggcat	2040
	tgctgatgaa	gtaaaatctt	gcttccgtgt	tgaatgggtt	gatcctgcca	acctgaacg	2100
	tggttttaag	tacatttatc	tgaatgaaga	agactatggt	cgtattagct	cttctgttat	2160
	agcgacacag	acacagctag	atagtggcga	aataagggtg	gttatcgatt	ctgttgtggg	2220
	caaggaggat	ggactagggt	tggagaacat	acatggaagt	gctgctattg	ccagtgttta	2280
25	ttctagggca	tacgaggaga	catttacact	tacatttgtg	agtggctgaa	cggttggaat	2340
	aggagcctat	cttgctcggc	ttggcatacg	gtgcatacag	cgtattgacc	agccatttat	2400
	cttaactggg	ttttctgccc	tgaacaagct	tctcgggcgg	gaagtgtaca	gctcccacat	2460
	gcagttgggt	gggtcccaaaa	tcatggctac	aaatgggtgt	gtccatctga	ctgttccaga	2520
	tgaccttgaa	gggttttcta	atatattgag	gtggctcagt	tatgttctctg	ctaacttggg	2580
30	tggacctctt	cctattacaa	aatctttaga	cccaatagac	agacctgttg	catacatccc	2640
	tgagaataca	tgtgatcctc	gtgcagccat	cagtggcatt	gatgacagcc	aagggaaatg	2700
	gttgggtggg	atgtttgaca	aagacagttt	tgtggagaca	tttgaaggat	gggcaaagac	2760
	agtagttact	ggcagagcaa	aacttgaggg	gattcctgtg	gggtttatag	ctgtggagac	2820
	acagaccatg	atgcagctcg	tccctgcaga	tccaggccag	cctgattccc	atgagcggtc	2880
35	tgttcctcgt	gcggggcaag	tctggtttcc	agattcagct	accaagacag	cgcacgcaat	2940
	gttggacttc	aaccgtgaag	gattacctct	gttcatcctt	gctaactgga	gaggcttctc	3000
	tggtgggcaa	agagaccttt	ttgaaggaat	t			3031

<210> 24

40

<211> 1010

<212> PRT

<213> Avena sativa

<400> 24

5 Gln Asn Ser Lys Asp Leu Gln Lys Val Val Asp Ile Val Leu Ser His
 1 5 10 15
 Gln Gly Val Arg Ser Lys Thr Lys Leu Ile Leu Thr Leu Met Glu Lys
 20 25 30
 Leu Val Tyr Pro Asn Pro Ala Ala Tyr Arg Asp Gln Leu Ile Arg Phe
 10 35 40 45
 Ser Ser Leu Asn His Lys Arg Tyr Tyr Lys Leu Ala Leu Lys Ala Ser
 50 55 60
 Glu Leu Leu Glu Gln Thr Lys Leu Ser Glu Leu Arg Thr Ser Ile Val
 65 70 75 80
 15 Arg Asn Leu Ser Ala Leu Glu Met Phe Thr Glu Glu Arg Ala Gly Phe
 85 90 95
 Ser Leu Gln Glu Lys Lys Leu Ala Ile Asn Glu Ser Met Gly Asp Leu
 100 105 110
 Val Ser Ala Pro Leu Pro Val Glu Asp Ala Leu Ile Ser Leu Phe Asp
 20 115 120 125
 Cys Ile Asp Gln Thr Leu Gln Gln Arg Val Ile Glu Thr Tyr Ile Ser
 130 135 140
 Arg Leu Tyr Gln Pro Gln Leu Val Lys Asp Ser Ile Gln Leu Lys Tyr
 145 150 155 160
 25 Gln Asp Ser Ala Val Ile Ala Leu Trp Glu Phe Asn Glu Gly His Pro
 165 170 175
 Asp Asn Arg Leu Gly Ala Met Val Ile Leu Lys Ser Leu Glu Ser Val
 180 185 190
 Ser Thr Ala Ile Gly Ala Ala Leu Lys Asp Arg Ser His Tyr Ala Ser
 30 195 200 205
 Ser Ala Gly Asn Thr Val His Ile Val Leu Leu Asp Ala Asp Thr Gln
 210 215 220
 Thr Asn Thr Thr Glu Asp Ser Gly Asp Asn Asp Gln Ala Gln Asp Arg
 225 230 235 240
 35 Met Asp Lys Leu Ser Val Ile Leu Lys Gln Asp Ile Val Thr Ala Asp
 245 250 255
 Leu Cys Ala Ala Gly Val Lys Val Ile Ser Cys Ile Val Gln Arg Asp
 260 265 270
 Gly Ala Leu Met Pro Met Arg Arg Thr Phe Leu Leu Ser Glu Glu Lys
 40 275 280 285

Leu Cys Tyr Glu Glu Glu Pro Ile Leu Arg His Val Glu Pro Pro Leu
 290 295 300
 Ser Ala Leu Leu Glu Leu Asp Lys Leu Lys Val Lys Gly Tyr Asn Glu
 305 310 315 320
 5 Met Lys Tyr Thr Pro Ser Arg Asp Arg Gln Trp His Ile Tyr Thr Leu
 325 330 335
 Arg Asn Thr Glu Asn Pro Lys Met Leu His Arg Val Phe Phe Arg Thr
 340 345 350
 Leu Val Arg Gln Pro Ser Ala Gly Asn Arg Phe Met Ser Gly His Ile
 10 355 360 365
 Asn Asp Val Glu Val Gly His Ala Glu Glu Ser Leu Ser Phe Thr Ser
 370 375 380
 Ser Ser Ile Leu Arg Ser Leu Met Thr Ala Ile Glu Glu Leu Glu Leu
 385 390 395 400
 15 His Ala Ile Arg Thr Gly His Ser His Met Tyr Leu Cys Ile Leu Lys
 405 410 415
 Glu Gln Lys Leu Leu Asp Leu Val Pro Val Ser Gly Asn Thr Val Val
 420 425 430
 Asp Val Gly Gln Asp Glu Ala Thr Ala Cys Ser Leu Met Lys Glu Met
 20 435 440 445
 Ala Leu Lys Ile His Glu Leu Val Gly Ala Arg Met His His Leu Ser
 450 455 460
 Val Cys Gln Trp Glu Val Lys Leu Lys Leu Asp Ser Asp Gly Pro Ala
 465 470 475 480
 25 Ser Gly Ser Trp Arg Val Val Thr Thr Asn Val Thr Ser His Thr Cys
 485 490 495
 Thr Val Asp Ile Tyr Arg Glu Val Glu Asp Thr Asp Ser Gln Lys Leu
 500 505 510
 Val Tyr His Tyr Ala Ala Ser Ser Pro Gly Pro Leu His Gly Val Thr
 30 515 520 525
 Leu Ser Asn Ser Tyr Gln Pro Leu Ser Val Ile Asp Leu Lys Arg Cys
 530 535 540
 Ser Ala Arg Asn Asn Arg Thr Thr Tyr Cys Tyr Asp Phe Pro Leu Ala
 545 550 555 560
 35 Phe Glu Thr Ala Val Arg Lys Ser Trp Ala Asn Val Ser Ser Glu Asn
 565 570 575
 Asn Gln Cys Tyr Val Lys Ala Thr Glu Leu Val Phe Ala Asp Lys Asn
 580 585 590
 Gly Ser Trp Gly Thr Pro Ile Val Pro Met Gln Arg Ala Ala Gly Leu
 40 595 600 605

Asn Asp Ile Gly Met Val Ala Trp Ile Leu Asp Met Ser Thr Pro Glu
 610 615 620
 Phe Pro Ser Gly Arg Gln Ile Ile Val Ile Ala Asn Asp Ile Thr Phe
 625 630 635 640
 5 Arg Ala Gly Ser Phe Gly Pro Arg Glu Asp Ala Phe Phe Glu Ala Val
 645 650 655
 Thr Asn Leu Ala Cys Glu Arg Lys Leu Pro Leu Ile Tyr Met Ala Ala
 660 665 670
 Asn Ser Gly Ala Arg Ile Gly Ile Ala Asp Glu Val Lys Ser Cys Phe
 10 675 680 685
 Arg Val Glu Trp Val Asp Pro Ala Asn Pro Glu Arg Gly Phe Lys Tyr
 690 695 700
 Ile Tyr Leu Asn Glu Glu Asp Tyr Gly Arg Ile Ser Ser Ser Val Ile
 705 710 715 720
 15 Ala His Lys Thr Gln Leu Asp Ser Gly Glu Ile Arg Trp Val Ile Asp
 725 730 735
 Ser Val Val Gly Lys Glu Asp Gly Leu Gly Val Glu Asn Ile His Gly
 740 745 750
 Ser Ala Ala Ile Ala Ser Ala Tyr Ser Arg Ala Tyr Glu Glu Thr Phe
 20 755 760 765
 Thr Leu Thr Phe Val Ser Gly Arg Thr Val Gly Ile Gly Ala Tyr Leu
 770 775 780
 Ala Arg Leu Gly Ile Arg Cys Ile Gln Arg Ile Asp Gln Pro Ile Ile
 785 790 795 800
 25 Leu Thr Gly Phe Ser Ala Leu Asn Lys Leu Leu Gly Arg Glu Val Tyr
 805 810 815
 Ser Ser His Met Gln Leu Gly Gly Pro Lys Ile Met Ala Thr Asn Gly
 820 825 830
 Val Val His Leu Thr Val Pro Asp Asp Leu Glu Gly Val Ser Asn Ile
 30 835 840 845
 Leu Arg Trp Leu Ser Tyr Val Pro Ala Asn Ile Gly Gly Pro Leu Pro
 850 855 860
 Ile Thr Lys Ser Leu Asp Pro Ile Asp Arg Pro Val Ala Tyr Ile Pro
 865 870 875 880
 35 Glu Asn Thr Cys Asp Pro Arg Ala Ala Ile Ser Gly Ile Asp Asp Ser
 885 890 895
 Gln Gly Lys Trp Leu Gly Gly Met Phe Asp Lys Asp Ser Phe Val Glu
 900 905 910
 Thr Phe Glu Gly Trp Ala Lys Thr Val Val Thr Gly Arg Ala Lys Leu
 40 915 920 925

31

Gly Gly Ile Pro Val Gly Val Ile Ala Val Glu Thr Gln Thr Met Met
 930 935 940
 Gln Leu Val Pro Ala Asp Pro Gly Gln Pro Asp Ser His Glu Arg Ser
 945 950 955 960
 5 Val Pro Arg Ala Gly Gln Val Trp Phe Pro Asp Ser Ala Thr Lys Thr
 965 970 975
 Ala His Ala Met Leu Asp Phe Asn Arg Glu Gly Leu Pro Leu Phe Ile
 980 985 990
 Leu Ala Asn Trp Arg Gly Phe Ser Gly Gly Gln Arg Asp Leu Phe Glu
 10 995 1000 1005
 Gly Ile
 1010

<210> 25
 15 <211> 20
 <212> DNA
 <213> Zea mays

 <400> 25
 20 ccttttttatg gcactgtgcg 20

 <210> 26
 <211> 22
 <212> DNA
 25 <213> Zea mays

 <400> 26
 catcgtagcc tatatgagga cg 22

30 <210> 27
 <211> 16
 <212> DNA
 <213> Zea mays

35 <400> 27
 caatcacgtc agactg 16

 <210> 28
 <211> 20
 40 <212> DNA

<213> Zea mays

<400> 28
actgtgcggtt tgagaaggctc 20

5

<210> 29
<211> 20
<212> DNA
<213> Zea mays

10

<400> 29
cctctacgta attggtcagc 20

<210> 30
15 <211> 26
<212> DNA
<213> Zea mays

<400> 30
20 ggatatatgc tttggtggaa tctggc 26

INTERNATIONAL SEARCH REPORT

International Application No

PL /US 99/14022

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/00 C12N15/82 A01H5/00 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL EBI Accession No: 048959, 1 June 1998 (1998-06-01) GORNICKI P. ET AL.: "ACETYL-COENZYME A CARBOXYLASE; from Triticum Aestivum" XP002122229 see sequence abstract --- -/--	1-50

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

10 November 1999

Date of mailing of the international search report

26/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Chakravarty, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/14022

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL EBI Accession No. Q08367, 1 November 1996 (1996-11-01) ASHTON A.R. ET AL.: "ACETYL-COA CARBOXYLASE (ACC); from Zea Mays" XP002122230 see sequence abstract ----	1-50
X	DATABASE EMBL EBI Accession no. AF029895, 3 February 1998 (1998-02-03) GORNICKI P. ET AL.: "Triticum aestivum acetyl-coenzyme A carboxylase (Acc-1) mRNA, nuclear gene encoding plastid protein, complete cds." XP002122231 see sequence abstract ----	1-50
X	DATABASE EMBL EBI Accession no. U58598, 31 May 1996 (1996-05-31) ASHTON A.R. ET AL.: "Zea mays acetyl CoA carboxylase mRNA, partial cds." XP002122232 see sequence abstract ----	1-50
X	WO 93 11343 A (GAHAN JOHN PETER) 10 June 1993 (1993-06-10) figure 3 ----	1-50
X	WO 96 31609 A (UNIV MINNESOTA ;GENGENBACH BURLE G (US); SOMERS DAVID A (US); WYSE) 10 October 1996 (1996-10-10) the whole document -----	1-50

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PL./US 99/14022

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9311343	A	10-06-1993	AU	2937192 A	28-06-1993
WO 9631609	A	10-10-1996	BR	9608013 A	02-03-1999
			CA	2217367 A	10-10-1996
			EP	0820513 A	28-01-1998