



(12) UK Patent (19) GB (11) 2103614 B

(54) Title of invention

Benzazepin-2-ones

(51) INT CL³; C07D 223/16 A61K 31/55 C07D 401/12 405/12 (C07D 401/12 213/30)
(C07D 405/12 223/16 307/819)

(21) Application No
8222564

(22) Date of filing
5 Aug 1982

(30) Priority data

(31) **291907**
319863

(32) **11 Aug 1981**
9 Nov 1981

(33) **United States of America**
(US)

(43) Application published
23 Feb 1983

(45) Patent published
21 Nov 1984

(52) Domestic classification
C2C 1200 1474 1530 1732 213
215 21X 220 222 225 226 227
22Y 246 247 250 251 253 25Y
271 282 28X 292 29X 29Y 302
305 30Y 313 314 31Y 321 322
323 324 32Y 332 337 338 340 342
346 34Y 351 352 360 362 364 366
367 368 36Y 386 387 388 389
40Y 43X 440 594 620 621 624
625 627 628 62X 634 635 638
645 648 650 658 65X 661 670
672 699 711 719 722 72Y 760 761
762 768 802 80Y AA KA KN LK
LW LZ MA MV TT TW
U1S 1328 2415 C2C

(continued on next page)

(73) Proprietor
Ciba-Geigy AG (Switzerland),
Klybeckstrasse 141
4002 Basle
Switzerland

(72) Inventor
Jeffrey William Herbert
Watthey

(74) Agent and/or
Address for Service
Kenneth D. Sparrow,
Ciba-Geigy plc
Patent Dept.
Tenax Road
Trafford Park
Manchester M17 1WT

GB 2103614 B

2 103 614 B — continuation

(56) Documents cited
None

(58) Field of search
C2C


Benzazepin-2-ones

The present invention is based upon the discovery that certain substituted 3-amino-[1]benzazepin-2-one-1-alkanoic acids and derivatives represent a new class of potent angiotensin-converting enzyme (ACE) inhibitors.


The foregoing attributes render the 3-amino-[1]benzazepin-2-ones of this invention particularly useful when administered, alone or in combination, to mammals, e.g. for the treatment or prevention of diseases responsive to inhibition of angiotensin converting enzyme e.g., cardiovascular disorders such as hypertension and cardiac conditions such as congestive heart failure.

This invention relates to novel 3-amino-[1]benzazepin-2-one-1-alkanoic acids, and derivatives useful as angiotensin-converting enzyme inhibitors, processes for preparing same, pharmaceutical compositions comprising said compounds, and to said compounds and compositions for use in a therapeutic method of treating diseases responsive to inhibition of angiotensin-converting enzyme in humans and mammals.

The compounds of the invention are characterized by the general formula I



wherein R_A and R_B are radicals of the formula



in which R_0 is carboxy or a functionally modified carboxy; R_1 is hydrogen, lower alkyl, amino(lower) alkyl, aryl, aryl (lower) alkyl, cycloalkyl or cycloalkyl (lower) alkyl; R_2 is hydrogen or lower alkyl; R_3 and R_4 , each independently, represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxy, halogen, trifluoromethyl, or R_3 and R_4 taken together represent lower alkylene-dioxy; R_5 is hydrogen or lower alkyl, and X represents oxo, two hydrogens, or one hydroxy together with one hydrogen; and wherein the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro; and salts and complexes thereof.

The functionally modified carboxyl group in the meaning of the symbol R_0 is e.g. an esterified carboxyl group or a carbamoyl group optionally substituted on the nitrogen atom.

More specifically one or both of R_0 represented by COR_6 in radical R_A and represented by COR_7 in radical R_B independently represent carboxy, esterified carboxy, carbamoyl or substituted carbamoyl.

The salts and complexes of the compounds of formula I are derived from those compounds which have salt forming properties and are preferably pharmaceutically acceptable salts and complexes.

A carboxyl group R_0 is represented by COR_6 (in radical R_A) wherein R_6 is hydroxy or COR_7 (in radical R_B) wherein R_7 is hydroxy.

An esterified carboxyl group R_0 is especially one in which the esterifying radical represents optionally substituted lower alkyl or optionally substituted phthalidyl and is represented by the partial formula $-\text{COR}_6$ (in radical R_A) or the partial formula $-\text{COR}_7$

(in radical R_B), wherein one or both of R_6 and R_7 represents lower alkoxy; (amino, mono- or di-lower alkylamino)-substituted lower alkoxy; carboxy-substituted lower alkoxy, e.g. α -carboxy substituted lower alkoxy; lower alkoxy carbonyl-substituted lower alkoxy, e.g. α -lower alkoxy carbonyl-substituted lower alkoxy; aryl-substituted lower alkoxy, e.g. optionally substituted benzyloxy or pyridylmethoxy; (hydroxy, lower alkanoyloxy or lower alkoxy) substituted lower alkoxy, e.g. pivaloyloxy methoxy; (hydroxy, lower alkanoyloxy or lower alkoxy)-substituted lower alkoxy methoxy; bicycloalkoxy carbonyl-substituted lower alkoxy, e.g. bicyclo [2,2,1]heptyloxy carbonyl-substituted lower alkoxy, especially bicyclo[2,2,1]heptyloxy carbonyl-substituted methoxy; 3-phthalidoxy; (lower alkyl, lower alkoxy, halo)-substituted 3-phthalidoxy.


An optionally N-substituted carbamoyl group R_8 is especially one which is represented by the partial formula $-COR_6$ (in radical R_A) or the partial formula $-COR_7$ (in radical R_B), wherein one or both of R_6 and R_7 represent amino; lower alkylamino; di-lower alkylamino; di-lower alkylamino in which both alkyl groups are linked by a carbon to carbon bond and together with the amino nitrogen form a 5-, 6- or 7-membered heterocyclic ring, e.g. pyrrolidino, piperidino, or perhydroazepino; (amino or acylamino)-substituted lower alkylamino; α -(carboxy or lower alkoxy carbonyl)-substituted lower alkylamino; aryl substituted lower alkylamino in which aryl is preferably phenyl or indolyl and which can be substituted on the α -carbon atom by carboxy or lower alkoxy carbonyl.

Any prodrug derivatives of compounds of this invention e.g. any pharmaceutically acceptable esters and amides of the mono- or di-carboxylic acids of this invention that may be convertible by solvolysis or under physiological conditions to the said carboxylic acids, e.g. esters and amides cited above, represent a particular object of the invention.

Said esters are preferably, e.g., the straight chain or branched lower alkyl esters unsubstituted or suitably substituted such as the pivaloyloxymethyl, bornyloxycarbonylmethyl, benzyl, pyridylmethyl, α -carboxyethyl or suitably esterified α -carboxyethyl esters, and the like.

Said amides are preferably e.g. simple primary and secondary amides and amides derived from the amino acids or derivatives thereof, such as the amides derived from alanine, phenylalanine and the like.

More particularly, the invention relates to compounds of formula IA



wherein R_1 is hydrogen, lower alkyl, amino(lower)alkyl, aryl, aryl(lower)alkyl, cycloalkyl(lower)alkyl, R_2 and R_5 represent hydrogen or lower alkyl, R_3 and R_4 represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxy, halogen, trifluormethyl; or R_3 and R_4 taken together represent lower alkylendioxy, X represents oxo, two hydrogens or one hydroxy group and one hydrogen, R_6 and R_7 independently represent hydroxy, amino, mono- or di-(lower)alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy, (amino, mono- or di-lower alkylamino, carboxy, or lower alkoxy carbonyl)-lower alkoxy; or the pharmaceutically acceptable salts or complexes thereof.

Preferred embodiments of this invention relate to compounds of formula IA, wherein R_1 is hydrogen, lower alkyl, amino(lower)-alkyl, aryl(lower)alkyl where aryl represents phenyl unsubstituted or mono- or disubstituted by lower alkyl, hydroxy, lower alkoxy, lower alkylendioxy, lower alkanoyloxy, halogen or trifluoromethyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 and R_4 are hydrogen, lower alkoxy, lower alkyl, halogen or trifluoromethyl;

or R_3 and R_4 taken together represent alkylendioxy, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower) alkoxy, lower alkoxycarbonyl(lower)alkoxy, or pharmaceutically acceptable salts thereof.


Very useful are compounds of formula IA, wherein R_1 is hydrogen, lower alkyl, ω -amino(lower)alkyl, aryl(lower)alkyl where aryl represents phenyl unsubstituted or mono-substituted by lower alkyl, hydroxy; lower alkoxy, lower alkanoyloxy, halogen or trifluoromethyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 and R_4 are hydrogen, lower alkoxy, lower alkyl, halogen, or trifluoromethyl; or R_3 and R_4 taken together represent lower alkylendioxy, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower)alkoxy, lower alkoxycarbonyl(lower)alkoxy, or pharmaceutically acceptable salts thereof.

Particularly useful are compounds of formula IA wherein R_1 is hydrogen, lower alkyl, ω -amino(lower)alkyl, aryl(lower) alkyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 is hydrogen, R_4 is hydrogen, lower alkoxy, lower alkyl, halogen, or trifluoromethyl, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower) alkoxy, lower alkoxycarbonyl(lower)alkoxy, or pharmaceutically acceptable salts thereof.

Especially useful are compounds of formula IA wherein R_1 is hydrogen, methyl, ethyl, isopropyl, ω -aminopropyl, ω -aminobutyl, aryl(methyl, ethyl, propyl) where aryl represents phenyl unsubstituted or substituted by one methyl, hydroxy, methoxy, methylenedioxy, acetoxy, chloro or trifluoromethyl group, R_2 and R_5 are hydrogen or methyl, R_3 and R_4 represents hydrogen, methoxy, methyl, chloro or

trifluoromethyl, X represents oxo, one hydroxy and one hydrogen or 2 hydrogens, R₆ and R₇ independently represent hydroxy, amino, ethoxy, methoxy, benzyloxy, ethoxycarbonylmethoxy or pivaloyloxy-methoxy; or pharmaceutically acceptable salts thereof.

Exceedingly useful are compounds of formula IB



wherein n represents an integer from 1 to 4, R₈ is hydrogen, phenyl unsubstituted or monosubstituted by lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, hydroxy, or trifluoromethyl, R₆ and R₇ independently represent hydroxy, lower alkoxy of up to 4 carbon atoms, benzyloxy, or amino, or pharmaceutically acceptable salts thereof.


Especially valuable are compounds of formula IB, wherein C_nH_{2n} represents ethylene, R₈ represents phenyl or phenyl mono-substituted by lower alkoxy with up to 4 carbon atoms, lower alkyl with up to 4 carbon atoms, halogen or trifluoromethyl, R₆ and R₇ independently represent hydroxy or lower alkoxy with up to 4 carbon atoms, or pharmaceutically acceptable salts thereof.

The present invention also relates to the stereoisomers of compounds of formula I. A number of racemates are obtainable when, e.g. in

formula IA at least one of R_1 and R_2 is not hydrogen and/or X represents H(OH).

The individual enantiomers of said racemates may in turn be obtained. Certain specific said isomers are preferred as angiotensin-converting enzyme inhibitors.

Outstanding are compounds of formula IC



wherein S represents the chirality, n represents an integer from 1 to 4, R_8 is hydrogen, phenyl unsubstituted or monosubstituted by lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, hydroxy, or trifluoromethyl, R_6 and R_7 independently represent hydroxy, lower alkoxy of up to 4 carbon atoms, benzyloxy or amino, or pharmaceutically acceptable salts thereof.

The general definitions used herein have the following meanings within the scope of the present invention.

Aryl represents a carbocyclic or heterocyclic aromatic radical preferably being phenyl, unsubstituted or mono or di-substituted by lower alkyl, lower alkoxy, lower alkylenedioxy, lower alkanoyloxy, hydroxy, halogen or trifluoromethyl.

The term cycloalkyl represents a cyclic hydrocarbon radical which preferably contains 3 to 8 carbons and is, for example, cyclopentyl or cyclohexyl.

The term aryl(lower)alkyl represents preferably benzyl, 1- or 2-phenylethyl, 1-,2- or 3-phenylpropyl 1-,2-,3- or 4-phenylbutyl, wherein the phenyl ring is unsubstituted or mono- or disubstituted by lower alkyl, hydroxy, lower alkoxy, lower alkylenedioxy, lower alkanoyloxy, halogen or trifluoromethyl.

The term cycloalkyl(lower)alkyl represents preferably 1- or 2-(cyclopentyl or cyclohexyl)ethyl, 1-,2- or 3-(cyclopentyl or cyclohexyl)propyl, or 1-,2-,3- or 4-(cyclopentyl or cyclohexyl)-butyl.

The term "lower" referred to above and hereinafter in connection with organic radicals or compounds respectively defines such with up to and including 7, preferably up and including 4 and advantageously one or two carbon atoms.

A lower alkyl group preferably contains 1-4 carbon atoms and represents for example ethyl, propyl, butyl or advantageously methyl.

A lower alkoxy group preferably contains 1-4 carbon atoms and represents for example methoxy, propoxy, isopropoxy or advantageously ethoxy. A mono-(lower)alkylamino group preferably contains 1-4 carbon atoms in the alkyl portion and is for example N-methylamino, N-propylamino or advantageously N-ethylamino. A di-(lower)alkylamino group preferably contains 1-4 carbon atoms in each lower alkyl portion and represents, for example, N,N-dimethylamino, N-methyl-N-ethylamino and advantageously N,N-diethylamino.

Lower alkanoyloxy represents preferably acetoxy, propionyloxy or pivaloyloxy.

Alkylenedioxy represents preferably ethylenedioxy, and advantageously methylenedioxy.

Aryl lower alkoxy represents advantageously e.g. benzyloxy, benzyloxy substituted by methyl, methoxy or chloro, and pyridylmethoxy.

Carboxy lower alkoxy represents advantageously e.g. 1-carboxyethoxy.

Lower alkoxycarbonyl lower alkoxy represents advantageously e.g. 1-(ethoxycarbonyl)ethoxy.

Amino(lower)alkoxy, mono-(lower)alkylamino lower alkoxy, di-(lower) alkylamino lower alkoxy advantageously represent respectively e.g. aminoethoxy, ethylaminoethoxy, diethylaminoethoxy.

Lower alkanoyloxymethoxy represents advantageously e.g. pivaloyloxymethoxy.

Bicycloalkyloxycarbonyl-(lower)alkoxy preferably represents bicyclo-[2,2,1]heptyloxycarbonyl-(lower)alkoxy unsubstituted or substituted by lower alkyl advantageously bornyloxycarbonylmethoxy.

Amino(lower)alkyl and ω -amino(lower)alkyl represent preferably amino(ethyl, propyl or butyl) and ω -amino(ethyl, propyl or butyl) respectively.

Halogen preferably represents chlorine, but may also be bromine, fluorine or iodine.

According to the present invention one or both of the carboxyl groups of the dicarboxylic acids, i.e. compounds of formula IA or IB wherein R₆ and R₇ are hydroxy, may be functionalized as esters or amides. These functional derivatives are preferably the mono or bis lower alkyl esters e.g. methyl, ethyl, n- or i-propyl, butyl or benzyl esters; the mono- or bis-amides, the mono- or di N-alkylated amides, e.g. mono- or diethylamides; the mono or bis substituted lower alkyl esters, e.g. the ω -(amino, mono- or dimethylamino, carboxy or carbethoxy) -(ethyl, propyl or butyl) esters. Highly preferred functional derivatives are the mono esters of formula IA, e.g. wherein one of R₆ and R₇ represents hydroxy and the other represents lower alkoxy.

Pharmaceutically acceptable salts are preferably metal or ammonium salts of said compounds of formula I wherein R₀ represents carboxy or of formula IA wherein COR₆ and/or COR₇ represent carboxy, more particularly alkali or alkaline earth metal salts, e.g., the sodium, potassium, magnesium or calcium salt; or advantageously easily crystallizing ammonium salts derived from ammonia or organic amines, such as mono-, di- or tri-lower (alkyl, cycloalkyl or hydroxyalkyl)-amines, lower alkylendiamines or lower hydroxyalkyl or aralkyl)-alkylammonium bases, e.g., methylamine, diethylamine, triethylamine, dicyclohexylamine, triethanolamine, ethylenediamine, tris-(hydroxymethyl)aminomethane or benzyltrimethylammonium hydroxide. Said compounds of Formula I form acid addition salts, which are preferably such of therapeutically acceptable inorganic or organic acids, such as strong mineral acids, for example hydrohalic, e.g. hydrochloric or hydrobromic acid; sulfuric, phosphoric, nitric or perchloric acid; aliphatic or aromatic carboxylic or sulfonic acids, e.g. formic, acetic, propionic, succinic, glycolic, lactic, malic, tartaric, gluconic, citric, ascorbic, maleic, fumaric, hydroxymaleic, pyruvic, phenylacetic, benzoic, 4-aminobenzoic, anthranilic, 4-hydroxybenzoic, salicylic, 4-aminosalicylic, pamoic, nicotinic; methanesulfonic,

ethanesulfonic, hydroxyethanesulfonic, benzenesulfonic, p-toluene-sulfonic, naphthalenesulfonic, sulfanilic or cyclohexylsulfamic acid.

The compounds of formula I exhibit valuable pharmacological properties, e.g. cardiovascular effects, by inter alia inhibiting the release of Angiotensin II through selective inhibition of angiotensin-converting enzyme in mammals. The compounds are thus useful for treating diseases responsive to angiotensin-converting enzyme inhibition in mammals including man.

The compounds of this invention exhibit primarily hypotensive/antihypertensive and cardiac effects. These properties are demonstrable by in vivo or in vitro tests, using advantageously mammals, e.g., rats, cats, dogs or isolated organs thereof, as test objects. The animals may either be normotensive or hypertensive e.g., genetically spontaneous hypertensive rats, or renal hypertensive rats and dogs, and sodium-depleted dogs. The compounds can be applied to the test animals enterally or parenterally, advantageously orally or intravenously, for example within gelatin capsules or in the form of starchy suspensions or aqueous solutions. The applied dosage may range between about 0.01 and 100 mg/kg/day, preferably between about 0.05 and 50 mg/kg/day, advantageously between about 0.1 and 25 mg/kg/day.

The in vivo lowering effect on the blood pressure is recorded, either directly by means of a catheter, placed in the test animal's femoral artery, or indirectly by sphygmomanometry at the rat's tail and a transducer. The blood pressure is recorded prior to and after dosing in mm Hg.

Thus the antihypertensive effects are demonstrable in spontaneously hypertensive rats by indirect measurement of systolic pressure. Conscious rats are placed individually in restraint cages within a

gently warmed chamber. A pulse sensor is placed distal to an inflatable occlusive cuff on each rat's tail. The cuff is periodically inflated to occlude the tail artery. The pressure in the cuff is continuously reduced and the systolic pressure corresponds to the pressure in the cuff, at which the pulse waves reappear. After obtaining control values of blood pressure and heart rate, test compounds are administered orally once daily for 4 consecutive days. Additional blood pressure measurements are usually made at 2.0, 4.0 and 23.5 hours after each daily dosing, and responses are compared to those of rats dosed with the treatment vehicle.

As an illustration of the invention, the antihypertensive effect of the "higher melting" 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenyl-propylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one of example 1 is reported: at a dose of 3 mg/kg p.o. it lowers blood pressure by 40 mm Hg as the average effect measured at 2 and 4 hours after the last two daily dosings. The corresponding S,S enantiomer of example 12 at a dose of 1 mg/kg p.o. lowers blood pressure by 30 mm Hg.

The compounds of this invention when administered intravenously or orally also exhibit an inhibitory effect against the Angiotensin I induced pressor response of normotensive rats. Angiotensin I is hydrolyzed by the reaction of said converting enzyme to the potent pressor substance Angiotensin II. The inhibition of said enzyme prevents the generation of Angiotensin II from Angiotensin I. In this manner the increase of blood pressure provoked by Angiotensin I is attenuated.

The corresponding in vivo test for intravenously administered compounds is performed with male, normotensive rats, which are anesthetized with sodium 5-ethyl-5-(1-methylpropyl)-2-thiobarbiturate. A femoral artery and saphenous vein are cannulated respectively for direct blood pressure measurement and the i.v. administration of Angiotensin I and a compound of this invention. After the basal

blood pressure is stabilized, pressor responses to 3 challenges of 333 ng/kg Angiotensin I i.v., at 5 minute intervals, are obtained. Such pressure responses are usually again obtained at 5, 10, 15, 30 and 60 minutes after i.v. administration of the compound to be tested, and compared with the initial responses. Any observed decrease of said pressor response is an indication of Angiotensin I converting enzyme inhibition. Illustrative of this invention, the "higher melting" 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H[1]benzazepin-2-one of example 1 and the corresponding S,S enantiomer of example 12 completely inhibit the pressor response following Angiotensin I challenge through 30 minutes after administration of either of the said compounds at a dose of 1 mg/kg i.v..

The in vitro inhibition of the angiotensin-converting enzyme by the compounds of this invention can be demonstrated by a method analogous to *Biochim. Biophys. Acta* 293, 451 (1973). According to this method, said compounds are dissolved at about 1 mM concentration in phosphate buffer. To 100 microliters of solutions of the test compound in phosphate buffer, diluted to the desired concentration, are added 100 microliters of 5 mM hippuryl-histidyl-leucine in phosphate buffer, followed by 50 microliters of the angiotensin- converting enzyme preparation (from lungs of adult male rabbits) in Tris buffer, containing potassium and magnesium chloride, as well as sucrose. Said solutions are incubated at 37°C for 30 minutes and combined with 0.75 ml of 0.6 N aqueous sodium hydroxide to stop further reaction. Then 100 microliters of a 0.2% solution of o-phthalaldehyde in methanol are added at room temperature, and 10 minutes later 100 microliters of 6N hydrochloric acid. These samples are read against water in a spectrophotometer set at 360 nm, and the optical densities thereof estimated. They are corrected for the standard curve via conversion factor expressing nanomoles of histidyl-leucine formed during said 30 minute incubation period. The results are plotted

against drug concentration to determine the IC_{50} , i.e., the drug concentration which gives half the activity of the control sample containing no drug. Illustrative of the invention, the "higher melting" 1-carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one of example 9 and the corresponding S,S enantiomer of example 19 show an IC_{50} of 5.2×10^{-9} M and 1.7×10^{-9} M respectively. The corresponding "lower melting" 1-carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one of example 8 shows an IC_{50} of 5.8×10^{-8} M.


Angiotensin-converting enzyme not only participates in the conversion of Angiotensin I to Angiotensin II, but also plays a role in the control of bradykinin and aldosterone levels. The effect of the compounds of this invention on these factors may also contribute to the antihypertensive and cardiac effects of these new compounds.

The aforementioned advantageous properties render the compounds of this invention of great value as specific therapeutic agents for mammals including man.

Accordingly, the compounds of this invention are valuable antihypertensive agents, especially useful for ameliorating hypertension (regardless of etiology) and/or cardiac conditions, such as congestive heart failure, and/or other edemic or ascitic diseases. They are also useful intermediates in the preparation of other valuable products, especially of corresponding pharmaceutical compositions.

The compounds of formula I according to the invention can be prepared in a manner which is known per se, in that, e.g.

a) in a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein X , R_B , R_3 , R_4 and R_5 have the meanings given hereinbefore, R_A is introduced by alkylation with a compound of the formula




wherein Z is a reactive esterified hydroxyl group and R_A has the meanings given hereinbefore, or with a compound of the formula



wherein R_1 and R_O have the meanings given hereinabove, in the presence of a reducing agent, with a temporary protection of any primary and secondary amino groups and/or, optionally, hydroxyl and/or oxo groups, which may be present in any one of the substituents X , R_A , R_B , R_1 , R_3 , R_4 and R_5
or

b) a compound of the formula




in which the carbocyclic ring may also be hexahydro or 6,7,8,9-

tetrahydro, and wherein X , R_3 , R_4 and R_5 have the meanings given hereinabove and R'_A is hydrogen or R_A as defined hereinabove, is alkylated with a compound of the formula



wherein Z is a reactive esterified hydroxyl group and R_B has the meanings given hereinabove, while protecting temporarily any primary and secondary amino groups and/or, optionally, hydroxyl and/or oxo groups which may be present in any one of the residues X , R_A , R_B , R_3 , R_4 and R_5 , or

c) a compound of the formula




in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro and wherein Y is oxo or a reactive esterified hydroxyl group Z together with hydrogen, and X , R_B , R_3 and R_4 have the meanings given hereinabove, is condensed with an amine of the formula




wherein R_A and R_5 have the meanings given hereinabove, with the proviso that in the case Y is oxo, the condensation is carried out in the presence of a reducing agent and with a temporary protection of the oxo group which may be present as the substituent X , or

d) in a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein X and R₁ to R₅ have the meanings given hereinabove, one of the symbols R_o' and R_o'' is cyano and the other one is cyano or R_o as defined hereinabove, the cyano group(s) is (are) subjected to solvolysis, or


e) a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro and wherein X, R_A, R_B, R₃, R₄ and R₅ have the meanings given hereinabove, or an ester thereof, is cyclised, or

f) a compound which is structurally identical with a compound of formula I specified above, except for having an additional double bond located at C-3, or between the nitrogen atom and the adjacent carbon atom within the group R_A, is treated with a reducing agent in order to saturate this double bond, or

g) in order to produce a compound of formula I as specified hereinabove, in which X is oxo, condensing a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein R_5 , R_3 and R_4 have the meanings given hereinabove, with an amine of the formula



wherein R_A and R_5 have the meaning given hereinabove, and

h) if desired, a resulting compound of formula I as specified above is converted into another compound of formula I within its above-specified scope, and/or


i) if desired, a resulting compound of formula I as specified above and having salt-forming properties is converted into a salt thereof or a free compound is liberated from such a salt, and/or

j) if desired, a resulting compound of formula I as specified above and having complex-forming properties is converted into a complex thereof, and/or

k) if so required, an optical isomer which has a specific configuration with respect to at least one center of chirality is enriched from a mixture of stereoisomeric forms of a resulting compound of formula I.

The alkylation according to processes a) and b), which serves for introduction of residues R_A and R_B , respectively, is carried out in a conventional manner, advantageously by treating a corresponding starting material of formulae II and V, respectively, with an alkylating agent of the formula R_A-Z (IIIA) or R_B-Z (IIIB), respectively, wherein R_A or R_B have the meanings given hereinabove and Z is a reactive esterified hydroxyl group, such as a hydroxyl group esterified with a strong organic acid, e.g. an aliphatic or aromatic sulfonic acid (such as a lower alkane sulfonic acid, especially methane sulfonic, trifluoromethanesulfonic acid, especially benzenesulfonic, p-toluenesulfonic, p-bromobenzenesulfonic and p-nitrobenzenesulfonic acid) or with a strong inorganic acid, such as, especially, sulfuric acid, or a hydrohalic acid, such as hydrochloric or, most preferably, hydriodic or hydrobromic acid. The alkylation is carried out under conventional general conditions at temperatures ranging between about 0°C up to the boiling temperature of the reaction mixture, preferably at temperatures between room temperature to about 100°C. The reaction takes place advantageously in the presence of a solvent which is inert with respect to the reactants, such as chlorinated lower alkane (e.g. chloroform or methylene chloride), an acyclic or cyclic ether (e.g. diethyl ether, 1,2-dimethoxyethane, dioxane or tetrahydrofuran) and, in particular, a low-molecular weight tertiary amide (e.g. N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, N-ethylpiperidone and hexamethylphosphoric acid triamide). Advantageously, the strong acid HZ liberated during the reaction is bound by the addition of an acid-binding agent, such as, preferably, an inorganic acid-scavenger such as an alkali metal bicarbonate, carbonate or hydroxide, an organic quaternary ammonium salt (e.g. a tetrabutylammonium salt) or an organic tertiary base, such as triethylamine, N-ethylpiperidine, pyridine or quinoline.

In process a), the alkylation can also be carried out under the conditions of reductive alkylation in the manner generally known and used in the art. In carrying out the alkylation, a compound of the general formula



in which R_1 and R_2 have the meanings given hereinabove, is reacted with the starting bicyclic compound II and, simultaneously or in a subsequent step, with a reducing agent. Among reducing agents which are used simultaneously with the alkylating agent, mention should be made of formic acid and complex metal hydrides such as sodium cyano-borohydride; among reducing agents used predominantly in a separate subsequent operation, i.e. reduction of a preformed imine (Schiff's base), mention should be made of diborane and complex metal hydrides, such as, sodium borohydride, sodium cyanoborohydride which are added advantageously to the primary reaction mixture without isolating an intermediate, e.g. the imine. In this case, the alkylation is carried out advantageously in an organic solvent inert to the reducing agent, such as in an aliphatic or cyclic ether (such as diethyl ether, diisopropyl ether, 1,2-dimethoxyethane, dioxane or tetrahydrofuran) or an aliphatic alcohol (such as methanol, ethanol, isopropyl alcohol, glycol, glycol monomethyl ether or diethyleneglycol), preferably at about 0°-80°C. A principal reducing agent, however, which can be used both simultaneously and subsequently, is hydrogen, especially catalytically activated hydrogen. The catalysts are those conventionally used as hydrogenation catalysts, i.e. preferably those of the class of precious metals (such as palladium, platinum and rhodium) on a carrier (such as calcium carbonate, aluminium oxide or barium sulfate), in a finely dispersed suspension without carrier or, in form of complexes, in a homogeneous phase. Also, finely dispersed transition metals, such as Raney metals, especially Raney nickel, are very suitable catalysts for the reductive alkylation. The specific reaction conditions depend, to a

large extent, on the particular hydrogenation catalyst and its precise activity, and do not differ from those generally known for hydrogenation. Temperatures ranging from room temperature to about 150°C, and pressures of hydrogen ranging from atmospheric pressure to about 300 atmospheres are applicable according to the standard procedures of the art. In addition to the inert solvents which were mentioned above in connection with the hydride reduction, also low-molecular weight amides, especially tertiary amides (such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, N-ethylpiperidone, hexamethylphosphoric acid triamide) but also formamide and acetamide can be used as suitable solvents. Special measures have to be taken with starting materials of formula II which have an easily reducible functional group, such as the 5-oxo group; in order to preserve these groups, selective reduction conditions, as known in the prior art, have to be applied, or, if a simultaneous reduction of these groups is desired or required, vigorous reagents and/or conditions are employed accordingly.

The preformed imines referred to above are preferably prepared by condensing an amine of formula II with a compound of formula IV in an inert solvent, e.g. toluene or methylene chloride, advantageously in the presence of a dehydrating catalyst, e.g. boron trifluoride etherate, p-toluenesulfonic acid or molecular sieves.

Process b) is preferably carried out in the presence of very strong bases, such as alkali metal hydrides (e.g. sodium or potassium hydride), alkoxides (e.g. sodium methoxide or ethoxide, potassium tert-butoxide) or amides (e.g. lithium diisopropylamide), whereby ethers and amides mentioned above are preferred as solvents. In a special modification of process b), starting materials are used in which R'_A is hydrogen, and at least two equivalents of the reactant IIIB is employed. In the resulting product, both R_A and R_B are identical and within the scope of the meanings of R_B .

In any of the alkylation processes, primary and secondary amino groups in starting materials, except for the secondary amino group to be alkylated, must be in a temporarily protected form during the alkylation. Suitable protecting groups, as well as procedures for their introduction and removal are well known in the art being elaborated in great detail in particular as general methods for the synthesis of peptides, cf. Houben-Weyl: Methoden der organischen Chemie; 4th edition, vol. 15/I and II, E.Wünsch (editor): Synthese von Peptiden (Georg Thieme Verlag, Stuttgart; 1974). The narrower selection of the protecting groups depends on the specific purpose, it being necessary to take into account in particular the specific properties of the particular starting materials and the reaction conditions of the specific process. In the case of several functional groups to be protected, advantageous combinations can be selected. Preferably, for example, similar or, even better, identical amino protecting groups, are used both in the radicals R_o and in the radical R_1 and are simultaneously removed following alkylation.

Suitable as amino-protecting groups are especially amino-protecting groups that can be removed by reduction, for example especially those of the benzyloxycarbonyl type in which the benzyloxycarbonyl group may be substituted in the aromatic moiety by halogen atoms, lower alkoxy groups and/or lower alkyl radicals and, especially, by nitro groups, such as the p-chloro- and p-bromobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-methylbenzyloxycarbonyl and, especially, p-nitrobenzyloxycarbonyl group, or alternatively the isonicotinyloxy-carbonyl group. An advantageous amino-protecting group is an ethoxy-carbonyl group which carries in the β -position a silyl group substituted by three hydrocarbon radicals, such as triphenylsilyl, dimethyl-tert.butylsilyl or, especially, trimethylsilyl. A β -(trihydrocarbonyl-silyl)-ethoxycarbonyl group of this type, such as a β -(tri-lower alkylsilyl)-ethoxycarbonyl group, for example, especially β -(tri-methylsilyl)-ethoxycarbonyl, forms with the amino group to be protected a corresponding β -trihydrocarbysilylethoxycarbonylamino group

(for example the β -trimethylsilylethoxycarbonylamino group), which may be removed under very specific, very mild conditions by the action of fluoride ions.

It is also possible to use groups that can be removed by acidolysis, such as the tert-butoxycarbonyl groups and analogous groups, as well as those of the aralkyl type, such as benzhydryl, di-(4-methoxy)-benzhydryl and triphenylmethyl (trityl), or certain aralkoxycarbonyl groups of the 2-(*p*-biphenylyl)-2-propoxycarbonyl type, which are described in Swiss Patent Specification No. 509 266. It should be noted that protecting groups derived from esters of carbonic acids are in most cases also removable by basic hydrolysis.

For the optional temporary protection of hydroxy groups, protecting groups may be used advantageously that can be removed by reduction, cf. the above-cited text (Houben-Weyl), and also groups that can be removed by acidolysis, such as 2-tetrahydropyranyl, tert-butoxycarbonyl and tert-butyl. Preferred hydroxy-protecting groups that can be removed by reduction are, for example, benzyl groups that may be substituted in the aromatic moiety by halogen, lower alkyl, lower alkoxy and/or, especially, nitro, especially the 4-nitrobenzyl group. It is also possible to use acyl groups that can be removed under weakly basic conditions, such as formyl or trifluoroacetyl.

For the optional protection of oxo groups, these are preferably protected as ketals, especially as ketals derived from lower alkanols, such as methanol or ethanol, or advantageously of ethylene glycol, or as corresponding thioketals preferably those of 1,2-ethanedithiol. All these groups can liberate oxo groups under the conditions indicated further below.

The subsequent removal of protecting groups in accordance with the invention depends on their nature and is carried out in each case in a conventional manner known per se taking into consideration the

general properties of the derived product. If the protecting groups for amino, hydroxy and oxo have been so selected that they can be removed under similar conditions (especially preferred here are the groups removable by acidolysis or, for amino and hydroxy, by reduction, that have already been given special mention), then all of these protecting groups are advantageously removed in a single operation; in special cases, however, it is possible to use different types of groups and remove each of them individually.

The groups that can be removed by reduction, especially those that contain halogenated lower alkyl radicals (for example 2,2,2-trichlorethyl radicals), isonicotinyl radicals (for example isonicotinyloxy-carbonyl) and, especially, substituted benzyl radicals, especially 4-nitrobenzyl radicals of any kind, are preferably removed by zinc reduction, usually in the presence of an acid, preferably acetic acid, and with or without the addition of an inert organic solvent, usually at room temperature. The removal of a protecting group by acid hydrolysis (acidolysis) is carried out in the case of groups of the tert-butyl type by means of hydrogen chloride, hydrogen fluoride or trifluoroacetic acid, and in the case of acid-sensitive protecting groups chiefly by means of a lower aliphatic carboxylic acid, such as formic acid and/or acetic acid, in the presence of water and, optionally, a polyhalogenated lower alkanol or lower alkanone, such as 1,1,1,3,3-hexafluoropropan-2-ol or hexafluoroacetone. In this manner it is possible, for example, for an N-trityl group to be removed by an organic acid, such as formic acid, acetic acid, chloroacetic acid or trifluoroacetic acid, in aqueous or absolute trifluoroethanol as solvent (cf. German Offenlegungsschrift DT 2 346 147) or by aqueous acetic acid; for the tert-butoxycarbonyl group to be removed by trifluoroacetic acid or hydrochloric acid; and for the 2-(p-bi-phenyl)-isopropoxycarbonyl group to be removed by aqueous acetic acid or, for example, by a mixture of glacial acetic acid, formic acid (82.8% strength) and water (7:1:2) or in accordance with the process in DT 2 346 147. The β -silylethyl ester groups are pre-

ferably removed by fluoride ion-yielding reagents, for example fluorides of quaternary organic bases, such as tetraethylammonium fluoride.

Ketalized and thioketalized oxo groups are converted into free oxo groups by acidolysis with usual strong inorganic acids, or with oxalic acid, in the presence of water, the latter ones advantageously by treatment with a sulfur-binding agent, e.g. a mercury II - salt and/or cadmium carbonate. Protecting groups that are unstable to basic conditions, for example formyl, trifluoroacetyl and carbonic acid ester groups, can be carefully removed by the action of an aqueous sodium or potassium bicarbonate or carbonate solution or, also, aqueous ammonia, in an organic solvent, usually at room temperature. The protecting groups are preferably removed under the reaction conditions of the examples, or under analogous conditions.

Those of the end products according to the invention that contain basic groups are obtained, depending on the manner of isolation, in the form of bases or acid addition salts; analogously, end products having acidic groups may also be obtained in the form of salts. Each form can be converted into the other in known manner. The bases can be obtained from the acid addition salts in a manner known per se. From the bases it is in turn possible to obtain acid addition salts, especially therapeutically useful acid addition salts, by reaction with acids, for example with acids of the type that form the above-mentioned salts. Acids and their salts also stand in a similar relationship to one another. Compounds that have both a free carboxy group and a basic group may be in the form of inner salts and these are obtained, for example, by establishing the isoelectric point.

The starting materials of formula IIIA, IIIB and IV, that is to say the alkylating agents, are known or, if they are unknown, can be simply obtained by conventional synthetic processes.

The starting materials of formula II and V can be obtained by conventional synthetic processes, and advantageously in the manner which is described in more detail and exemplified for specific intermediates hereinafter.

Process c), also being an alkylation reaction is performed according to the same general considerations and under the same experimental conditions as the above processes a) and b) as described in detail above for the treatment with an alkylating agent of formula IIIA, IIIB or IV (i.e. substitutive alkylation or reductive alkylation). Starting materials of formula VI can be obtained by conventional processes known per se, e.g. in the manner described more specifically hereinafter. The amines of formula VII are known, or if unknown, they are easily accessible by conventional synthetic methods.

Process d), is also carried out in a conventional manner under the general conditions of solvolysis, which are known to convert cyanides (nitriles) into free carboxylic acids or their salts, esters or imides. - For the conversion into a free acid, hydrolysis with water is carried out advantageously in an inert organic solvent which is at least partially miscible with water, such as ethers (e.g. diethyl and diisopropyl ether, 1,2-dimethoxyethane or, especially dioxane or tetrahydrofuran) or lower alkanols (e.g. methanol, ethanol, isopropyl alcohol, butyl alcohols, especially tert-butyl alcohol), a larger amount of water being required in the latter cases in order to prevent alcoholysis. The hydrolysis can be catalysed both by strong acids, especially inorganic acids such as sulfuric acid or, preferably hydrohalic acids (e.g. hydrobromic or, as a first choice, hydrochloric acid), or by bases, especially inorganic bases such as hydroxides and carbonates of alkali metals, e.g. sodium and potassium hydroxide. The bases are usually employed in at least stoichiometric quantities giving rise to carboxylic acid salts as primary products.

The acidic catalysts are advantageously applied as dilute aqueous solution for the best result. Final products of formula I, in which R_o represents an esterified carboxyl group, can be obtained by carrying out the solvolysis of the nitrile with the corresponding alcohol (alcoholysis) in the presence of a catalytic amount of an anhydrous strong acid, advantageously gaseous hydrogen chloride. Usually, excess alcohol is used as solvent; however, inert organic solvents can be added, such as acyclic and cyclic ethers (especially these mentioned above), and/or halogenated lower alkanes (especially chloroform and dichloromethane). If the alcoholysis is carried out under strictly anhydrous conditions, the primary product (imino ester) is to be hydrolyzed, advantageously by adding water to the reaction mixture; otherwise, by carrying out the alcoholysis in the presence of an approximately stoichiometric equivalent of water, the desired ester is obtained directly. In order to obtain a corresponding amide (i.e. a compound of formula I, wherein R_o is carbamoyl), a corresponding nitrile of formula VIII can preferably be subjected to alkaline hydrolysis in the presence of hydrogen peroxide.

The starting materials of formula VIII can be obtained by conventional methods known per se, e.g. by a condensation analogous to that of process c), in which a starting material of the above-defined formula VI is treated with an amine of the formula



wherein R_1 and R_5 have the meanings given hereinabove, and which corresponds to the above-defined amine of formula VII. Also, processes a) and b) can analogously be used for the preparation of the nitriles of formula VIII.

The cyclization according to process variant e) can also be carried out in the manner known per se, e.g. by dehydration. Especially useful general methods for this purpose are those developed in connection with the formation of the amide bond in peptides, as reviewed in compilative works, e.g. Houben-Weyl, Volumes 15/1 and 15/2 as cited hereinabove. According to one preferred modification, the amino group to be cyclized is rendered inactive by protonation (i.e. in the form of an acid addition salt), and the carboxyl group is converted into an activated ester, such as that with 2,4,5-trichlorophenol, pentachlorophenol, pentafluorophenol, 2-nitrophenol or, especially, 4-nitrophenol, or with an N-hydroxy compound, such as N-hydroxysuccinimide, 1-hydroxybenztriazole or N-hydroxypiperidine, or alternatively with an N,N'-di-substituted isourea, such as, especially, N,N'-dicyclohexylisourea, or a similar generally known activating agent. The cyclization is effected by basification preferably by the addition of an organic base, for example a quaternary ammonium salt, or especially a tertiary amine, such as triethylamine, N-ethylmorpholine or N-methylpiperidine, in order to re-activate the amino group to be cyclized by converting it into the unprotonated form. The reaction temperature is usually from -20° to +50°C, preferably approximately at room temperature, and customary solvents are used, for example, dioxan, tetrahydrofuran, acetonitrile, pyridine, dimethylformamide, dimethylacetamide, dimethyl sulfoxide, N-methylpyrrolidone, hexamethylphosphoric acid triamide, as well as chloroform and methylene chloride, and expedient mixture thereof. In a special variant of the process, the carboxy group can be directly activated in situ by the action of the free acid with a carbodiimide, such as N,N'-dicyclohexylcarbodiimide (optionally with the addition of N-hydroxysuccinimide, an unsubstituted or, for example, halogen-, methyl- or methoxy-substituted 1-hydroxybenztriazole or 4-hydroxybenzo-1,2,3-triazine-3-oxide or N-hydroxy-5-norbornene-2,3-dicarboximide), or with N,N'-carbonyldiimidazole.

Starting materials of formula IX can be obtained according to general methods known per se, e.g. as discussed in more specific examples hereinafter.

Also, reduction according to process f) can be carried out in a manner generally known per se for saturation of such double bonds. More specifically, the double bond in the unsaturated starting materials corresponding to formula I can be located between C-3 and C-4 or between C-3 and the adjacent nitrogen atom, or between the nitrogen atom and the adjacent carbon atom within a group R_A. The saturation of the double bond is advantageously carried out by catalytic hydrogenation, e.g. under the preferred conditions discussed in detail hereinbefore, and also by metal reduction, such as zinc reduction in neutral or acidic medium, or, especially in the case of the C-N double bond, by diborane or complex hydrides such as sodium borohydride, as mentioned hereinbefore. The unsaturated starting materials for this process variant are obtained according to known general methods, e.g. those discussed in processes a) and c) and/or, in a more specific form hereinafter.

The condensation according to process g) is carried out under conventional general conditions at temperatures ranging between about 0°C and 100°C in a solvent which is inert to the reactants, e.g. methylene chloride, 1,2-dimethoxyethane, N,N-dimethylformamide optionally in the presence of a base, e.g. a tertiary amine such as triethylamine or an alkali metal hydride such as sodium hydride.

In performing the optional interconversions of a resulting final product of formula I, into another compound within the above-specified scope of formula I, transformations such as the following are carried out: an amino group is alkylated, and/or an oxo group, especially that of the symbol X, is converted into hydroxyl (plus hydrogen) or into two hydrogens by reduction and/or hydroxyl is converted into oxo by oxidation or into hydrogen by reduction,

and/or a free hydroxyl or carboxyl group is liberated from its esterified form by hydrolysis or hydrogenolysis and/or a hydroxyl or amino group is acylated and/or a free carboxyl is esterified, and/or the aromatic carbocyclic ring in formula I is hydrogenated to hexahydro or 6,7,8,9-tetrahydro, and/or the hexahydro carbocyclic ring is dehydrogenated to the 6,7,8,9-tetrahydro or aromatic carbocyclic ring.

All these optional interconversions are carried out by well-known conventional methods. By the alkylation reaction, e.g. the lower alkyl as represented by R_5 can be introduced into the final product of formula I, wherein R_5 is hydrogen, using any of the modifications discussed in detail in connection with process variant a). Both substitutive and reductive alkylation can be employed, the former with alkyl halides, the latter with lower aliphatic aldehydes and ketones and catalytically activated hydrogen or, in the case of formaldehyde, advantageously with formic acid as the reducing agent. By the substitutive alkylation, lower alkyls can also be introduced into an amino group which is a component of the carbamoyl group represented by symbol R_6 . Also the reduction of the 5-oxo group to hydroxy is carried out in the usual manner, e.g. using a complex metal hydride, especially a mild one, such as an alkali metal borohydride (e.g. sodium borohydride), or according to the method of Meerwein-Ponndorf, or a modification thereof using an alkanol, especially isopropyl alcohol, as both solvent and reducing agent and a metal alkoxide, preferably one corresponding to the reducing alcohol, such as aluminium isopropoxide, as a catalyst. The reduction of the oxo group to two hydrogens can advantageously be accomplished e.g. by treatment with amalgamated zinc and hydrochloric acid, or by Raney-nickel desulfurization of a corresponding dithioketal. The oxidation of hydroxyl to form oxo can be preferably carried out with a derivative of hexavalent chromium such as chromic acid and its salts, with a permanganate salt (especially potassium permanganate) or under the conditions of

the Oppenauer oxidation, with acetone or cyclohexanone as oxidant and aluminium isopropoxide as catalyst. Esterified hydroxyl groups are liberated in particular by methods discussed in detail hereinabove in connection with removing hydroxyl-protecting groups; the acylation of both hydroxyl and amino groups is carried out in the usual way, preferably using a corresponding acid anhydride or halide. For esterification, a carboxyl group can be reacted directly with a diazoalkane, especially diazomethane, or with a corresponding alcohol in the presence of a strong acid catalyst (e.g. sulfuric acid or an organic sulfonic acid) and/or a dehydrating agent (e.g. dicyclohexyl-carbodiimide). Alternatively, the carboxyl group can be converted into a reactive derivative thereof, such as an active ester mentioned in connection with process e), or into a mixed anhydride, e.g. with an acid halide (i.e., especially acid chloride) or with trifluoroacetic acid, and this activated intermediate reacted with the desired alcohol.

The free carboxyl group can be liberated from an esterified carboxyl in a manner generally known, especially by base-catalyzed hydrolysis. Of special interest, however, are methods capable of selectively liberating one particular carboxy group represented by the symbols $-\text{COR}_6$ and $-\text{COR}_7$. In such a case, use can be made of a proper combination of ester groups known in the art especially as carboxyl-protecting groups and developed in a great variety in particular for the synthesis of peptides, cf. Houben-Weyl, Volumes 15/1 and 15/2 as cited hereinabove. Radicals suitable for selective removal with liberation of the carboxyl are esters derived, for example, from alcohols that yield radicals that can be removed by acidolysis, such as cyanomethyl alcohol, benzoylmethyl alcohol or tert-butyl alcohol, but especially alcohols that yield radicals which can be removed by reduction, such as 2,2,2-trichloroethanol, benzyl alcohol, and especially 4-nitrobenzyl alcohol, or alternatively isonicotinyl alcohol. An especially advantageous class of substituted

alkanols are ethyl alcohols which carry in the β -position a tri-substituted silyl group, such as triphenylsilyl, dimethylbutylsilyl or, especially, trimethylsilyl. As is described, for example, in Belgian Patent No. 851.576, these alcohols are particularly suitable for selective removal because the corresponding β -silylethyl esters, for example β -(trimethylsilyl)-ethyl ester, have the stability of customary alkyl esters but can selectively be removed under mild conditions by the action of fluoride ions to retain other esterified carboxyl groups, for example alkoxy carbonyl groups.


The removal of esterifying groups depends on their nature and is carried out in each case in a conventional manner known per se taking into consideration the properties of the other radicals involved. The groups that can be removed by reduction, especially those that contain halogenated lower alkyl radicals (for example 2,2,2-trichloroethyl radicals), isonicotinyl radicals (for example isonicotinyloxycarbonyl) and, optionally substituted benzyl radicals, especially 4-nitrobenzyl radicals of any kind, are preferably removed by zinc reduction, usually in the presence of an acid, preferably acetic acid, and with or without the addition of an inert organic solvent, usually at room temperature, those of the benzyl type, especially unsubstituted benzyl esters, also by hydrogenolysis techniques conventionally used for benzyl groups.

The removal of an ester group by acid hydrolysis (acidolysis) can be carried out especially in the case of groups of the tert-butyl type, by means of hydrogen chloride, hydrogen fluoride or trifluoroacetic acid. The β -silylethyl ester groups are preferably removed by fluoride-ion-yielding reagents, for example fluorides of quaternary organic bases, such as tetraethylammonium fluoride. Ester groups that are base-unstable can be carefully removed by the rapid action of an aqueous sodium or potassium bicarbonate solution or, preferably, aqueous ammonia in an organic solvent, usually at room temperature. The ester groups are preferably removed under the


reaction conditions of the examples, or under analogous conditions.

A proper combination of the ester groups can be chosen in the earlier stages of the synthesis, or by a proper choice of starting materials and reactants, e.g. in process a), a selectively removable ester group being introduced with a carboxyl which is to be liberated in the last stage.

The compounds of formula I in general, and IA in particular, are prepared advantageously according to reaction sequence 1, which involves an advantageous selection of starting materials and intermediates, and comprises the following steps: a) condensing under conditions of basic catalysis, a compound of the formula




wherein R₃' and R₄' represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, trifluoromethyl or R₃' and R₄' taken together represent lower alkylendioxy, X' presents 2 hydrogens, one hydrogen and one etherified or esterified hydroxy, oxo or oxo protected in form of a ketal or thioketal and R₉ is amino, lower alkyl-amino, azido or acylamino, e.g. lower alkanoylamino or alkyloxy-carbonylamino, with a compound of the formula




wherein R₂' represents hydrogen or lower alkyl, Z represents


reactively esterified hydroxy, and R'_7 represents hydroxy, di(lower) alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy or lower alkoxycarbonyl(lower)alkoxy, b) optionally reducing, hydrogenolyzing, hydrolyzing or alkylating the resulting intermediate to obtain a compound of the formula II'



wherein R'_3 , R'_4 , X' are as defined for formula XI; R'_2 and R'_5 represent hydrogen or lower alkyl, R'_7 represents hydroxy, amino, mono- or di(lower)alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy, di(lower alkylamino)lower alkoxy or lower alkoxycarbonyl(lower)alkoxy, c) condensing a compound of formula II' above under conditions of reductive alkylation with a compound of the formula IV'



wherein R'_1 is hydrogen, lower alkyl, acylated amino (lower) alkyl, aryl, aryl(lower)alkyl, cycloalkyl(lower)alkyl and R'_6 represents hydroxy, di(lower)alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy or lower alkoxycarbonyl(lower)alkoxy, or condensing under alkylation conditions a compound of formula II' above with a compound of the formula III'A



wherein R'_1 and R'_6 have meanings given above in formula IV' and Z represents reactively esterified hydroxy, d) optionally hydrolyzing or derivatizing the resulting product; e) converting any resulting compound of formula IA into another compound of the invention.

Compounds of formula XI are obtained from the corresponding optionally substituted and/or derivatized 2,3,4,5-tetrahydro-1H[1]benzazepin-2-ones (J. Chem. Soc. 1937, 456; British patent 1,359,285; Liebigs's Annalen Chemie 574, 171 (1951). Novel appropriately derivatized starting [1]benzazepin-2-ones are advantageously prepared by Beckmann rearrangement of the correspondingly derivatized naphthalen-1-ones using procedures known to the art and exemplified herein.

Said tetrahydro-[1]benzazepin-2-ones are converted to the 3-halo-, e.g. 3-chloro-2,3,4,5-tetrahydro-1H[1]benzazepin-2-one under conditions exemplified herein, e.g. by treatment with phosphorus pentachloride followed by hydrogenation. Substitution of said halo derivative with a metal azide, e.g. sodium azide and optional reduction, or substitution with ammonia or a lower alkylamine and optional acylation, yields compounds of formula XI.

Alternatively, compounds of formula XI wherein R_9 represents amino, alkylamino or acylamino are obtained by reduction and cyclization of the appropriately substituted and/or derivatized 4-(o-nitro-phenyl)-2-aminobutyric acid and optional subsequent N-alkylation or N-acylation.

An alternate synthesis for the optically active compounds of this invention starts with the natural amino acid tryptophane. Specifically L-4-(o-aminophenyl)-4-oxo-2-amino-butyric acid (L-kynurenine, J.Am.Chem. Soc. 76, 1708 (1954), derived from L-tryptophane) is converted to an optically active starting material of formula XI wherein R_9 is acylamino, e.g. 3-(S)-t-butyloxycarbonylamino-2,3,4,5-


tetrahydro-1H[1]benzazepin-2,5-dione as described in the Australian Journal of Chemistry 33, 633-40 (1980). The lactam alkylation of a compound of formula XI with a reactant of formula III'B, well known in the art, is preferably carried out in the presence of bases such as alkali metal hydrides, e.g. sodium or potassium hydride, alkali metal alkoxides, e.g. potassium t-butoxide or sodium methoxide, organometallic reagents, e.g. lithium diisopropylamide or under conditions of phase transfer catalysis e.g. in the presence of a tetrabutylammonium salt, preferably in a solvent e.g. tetrahydrofuran, dimethylformamide, at a temperature preferably between about 0° and 75°.

Condensation of intermediates of formula II' with the known α -keto-acid derivatives of formula IV' (e.g. Chem. Ber. 31, 551, 3133) by reductive N-alkylation is carried out under conditions known to the art, e.g. by catalytic hydrogenation with hydrogen in the presence of platinum, palladium or nickel catalysts or with chemical reducing agents such as simple or complex light metal hydrides, advantageously an alkali metal cyanoborohydride such as sodium cyanoborohydride. The reductive amination with an alkali metal cyanoborohydride is preferably carried out in an inert solvent, e.g. methanol or acetonitrile, advantageously in the presence of an acid, e.g. hydrochloric acid or acetic acid at a temperature between about 0 and 50°, preferably room temperature.


Alkylation of intermediate amines of formula II' with a reactant of formula III'A, well known to the art, is carried out with or without basic catalysts such as triethylamine or potassium carbonate in an inert solvent.

The compounds of formula I in general, and IA in particular, can also be prepared by sequences 2 and 3.

Sequence 2 comprises the following steps: a) condensing under conditions of reductive alkylation a compound of the formula




wherein R'_3 , R'_4 and X' have meanings as defined for formula XI, and R'_5 is hydrogen or lower alkyl, with a compound of the formula IV'




wherein R'_1 and R'_6 have meanings as previously defined, or under alkylation conditions with a compound of formula III'A



wherein R'_1 , R'_6 and Z have meanings as previously defined, to obtain a compound of the formula V'




wherein R'_1 , R'_3 , R'_4 , R'_5 , R'_6 and X' have meanings as previously defined, b) condensing under conditions of basic catalysis a resulting compound of the formula V' with a compound of the formula III'B




wherein R'_2 and R'_7 and Z have meanings as previously defined,
c) optionally hydrolyzing or derivatizing the resulting product,
d) optionally converting any resulting compound of formula I into
another compound of the invention.

Sequence 3 comprises the following steps: a) condensing a compound of the formula VII'



wherein R''_1 is hydrogen, lower alkyl, acylated amino(lower)alkyl, aryl, aryl(lower)alkyl, cycloalkyl(lower)alkyl; R''_5 represents hydrogen or lower alkyl; and R''_6 represents hydroxy, di(lower)alkyl-amino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy or lower alkoxy carbonyl(lower)alkoxy, with a compound of the formula VI'



wherein R''_2 represents hydrogen or lower alkyl, R''_3 and R''_4 represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, trifluoromethyl or R''_3 and R''_4 taken together represent lower

alkylenedioxy; X" represents 2 hydrogens, one hydrogen and one etherified or esterified hydroxy, oxo or oxo protected in the form of a ketal or thioketal; R₇" represents hydroxy, di(lower)alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy or lower alkoxy-carbonyl(lower)alkoxy; and Y represents oxo or dichloro- under conditions of reductive N-alkylation, or condensing a compound of formula VII' with a compound of the above formula VI' wherein X" represents oxo, Y represents hydrogen and one reactively esterified or etherified hydroxy, or with a 3,4-dehydro elimination product of said compound or with a 3,4-dehydro derivative of said compound; b) optionally reducing, hydrolyzing or derivatizing the resulting product; c) optionally converting any resulting compound into another compound of the invention.

In the preceding sequences 2 and 3 the steps of lactam alkylation, reductive N-alkylation and alkylation of amines are advantageously carried out under the conditions described for process 1.

In sequences 1, 2 and 3 described herein, reactants of e.g. formulae III'A, III'B and VII' may be replaced with the corresponding nitriles, e.g. R₂'CH(Z)CN, R₁'CH(Z)CN and R₅NHCH(R₁')CN respectively. The nitriles thus obtained may be converted to the carboxylic acids, esters and amides of formula I using methods well known to the art.

The starting materials of formula VII' represent amino acids and derivatives well known to the art. It is noteworthy that the optically active compounds of this invention may be synthesized starting with an optically active compound of formula VII', e.g. L- α -aminophenylbutyric acid, L-phenylalanine and derivatives thereof.

In the case of reactants of formula III'A, III'B, IV' and VII' wherein R₇', R₆' or R₆" represents hydroxy, an appropriate carboxylate salt is prepared, preferably in situ, before condensation with the described intermediates cited above.

Certain terms used in the foregoing processes have the meanings as defined below.

A reactively esterified hydroxy represents such esterified by a strong inorganic or organic acid, above all a hydrohalic acid, e.g. hydrochloric, hydrobromic or hydriodic acid, an aliphatic or aromatic sulfonic acid, e.g. methanesulfonic acid or p-toluenesulfonic acid.

Etherified hydroxy represents preferably lower alkoxy, e.g. methoxy, ethoxy or t-butoxy.

The optional steps of reducing, hydrogenolyzing hydrolyzing or derivatizing the initial products of the aforesaid processes and the conversion of a resulting product into another compound of this invention are performed by chemical methodology known to the art and exemplified herein.

Compounds of formula I or IA wherein R₆ and/or R₇ is lower alkoxy may be amidized with ammonia, mono- or di-(lower)alkylamines to yield compounds of formula I or IA wherein R₆ and/or R₇ represents unsubstituted, mono- or di-(lower)alkylamino.

Conversion of compounds of formula I or IA wherein R₆ and/or R₇ is lower alkoxy, aryl(lower)alkoxy, amino, mono- or di-(lower)amino to compounds of formula I or IA wherein R₆ and/or R₇ represents hydroxy is advantageously carried out by hydrolysis with inorganic acids such as hydrohalic or sulfuric acid or with aqueous alkalies preferably alkali metal hydroxides such as lithium or sodium hydroxide.

The selective conversion of compounds of formula I or IA wherein R₆ and/or R₇ represents α -aryl(lower)alkoxy, e.g. benzyloxy to compounds of formula I or IA wherein R₆ and/or R₇ represents hydroxy

is advantageously carried out by hydrogenolysis using hydrogen in the presence of a catalyst, e.g. palladium.

Compounds of formula I or IA wherein neither R_6 nor R_7 represents hydroxy may be converted to monocarboxylic acids of formula I or IA wherein one of R_6 and R_7 is hydroxy. Such conversion is carried out by selective hydrolytic or hydrogenolytic procedures well known to the art and based on the chemical character of the R_6 and R_7 substituents.

Free carboxylic acids of formula I or IA wherein R_6 and/or R_7 represent hydroxy or salts thereof may be esterified with the appropriate alcohols or reactive derivatives thereof well known to the art to give the corresponding mono- or bis-ester, namely compounds of formula I or IA wherein R_6 and/or R_7 is lower alkoxy, aryl(lower) alkoxy, lower alkanoyloxymethoxy, or lower alkoxycarbonyl(lower) alkoxy. Furthermore the free carboxylic acids may be converted via reactive intermediates to mono- or bis-amides of formula I wherein R_6 and/or R_7 represents amino, mono- or di-(lower)alkylamino.

Compounds of formula I or IA, and intermediates therefor, e.g. of formulae X and V', wherein X or X' represents oxo may be converted to the corresponding compounds wherein X or X' represents one hydrogen and one hydroxy by reduction, e.g. by catalytic hydrogenation, e.g. with hydrogen in the presence of a platinum catalyst, or with a metal hydride reducing agent such as sodium borohydride. Resulting compounds wherein X or X' represents one hydrogen and one hydroxy may be converted to compounds wherein X or X' represents two hydrogens, e.g. by catalytic hydrogenation of the adduct of a carbodiimide, e.g. the adduct formed by condensation of a compound wherein X or X' represents one hydrogen and one hydroxy with dicyclohexylcarbodiimide in the presence of cuprous chloride according to the general method described in *Chem. Ber.*, 107, 1353 (1974).

Alternatively, the compounds wherein X or X' represents one hydrogen and one hydroxy may be first converted to the corresponding compounds wherein X or X' represents one hydrogen and one acyloxy (e.g. acetoxy) and subsequently reduced, e.g. by catalytic hydrogenation in the presence of a palladium catalyst, to compounds wherein X or X' represents two hydrogens.

The above-mentioned reactions are carried out according to standard methods, in the presence or absence of diluents, preferably such as are inert to the reagents and are solvents thereof, of catalysts, condensing or said other agents respectively and/or inert atmospheres, at low temperatures, room temperature or elevated temperatures, preferably at the boiling point of the solvents used, at atmospheric or superatmospheric pressure.

The invention further includes any variant of the present processes, in which an intermediate product obtainable at any stage thereof is used as starting material and the remaining steps are carried out, or the process is discontinued at any stage thereof, or in which the starting materials are formed under the reaction conditions, or in which the reaction components are used in the form of their salts or optically pure antipodes. Mainly those starting materials should be used in said reactions, that lead to the formation of those compounds indicated above as being especially useful.

Depending on the choice of starting materials and methods, the new compounds may be in the form of one of the possible isomers or mixtures thereof, for example, depending on the number of asymmetric carbon atoms, as pure optical isomers, such as antipodes, or as

mixtures of optical isomers such as racemates or mixtures of diastereoisomers.

Resulting mixtures of diastereoisomers and mixtures of racemates can be separated on the basis of the physicochemical differences of the constituents, in known manner, into the pure isomers, diastereoisomers or racemates, for example by chromatography and/or fractional crystallisation.

Resulting racemates can furthermore be resolved into the optical antipodes by known methods, for example by recrystallisation from an optically active solvent, by means of microorganisms or by reacting an acidic end product with an optically active base that forms salts with the racemic acid, and separating the salts obtained in this manner, for example on the basis of their different solubilities, into the diastereoisomers, from which the antipodes can be liberated by the action of suitable agents. Basic racemic products can likewise be resolved into the antipodes, for example, by separation of diastereomeric salts thereof, e.g. by the fractional crystallization of d- or l-tartrates. Any racemic intermediates or starting materials can likewise be resolved.

Advantageously, the more active of the two antipodes is isolated.

Finally, the compounds of the invention are either obtained in the free form, or as a salt thereof. Any resulting base can be converted into a corresponding acid addition salt, preferably with the use of a pharmaceutically acceptable acid or anion exchange preparation, or resulting salts can be converted into the corresponding free bases, for example, with the use of a stronger base, such as a metal or ammonium hydroxide or a basic salt, e.g. an alkali metal hydroxide or carbonate, or a cation exchange preparation. A compound of formula I wherein R_o represents carboxy or of formula IA wherein

COR_6 and/or COR_7 represent carboxy can thus also be converted into the corresponding metal or ammonium salts. These or other salts, for example, the picrates, can also be used for purification of the bases obtained; the bases are converted into salts, the salts are separated and the bases are liberated from the salts. In view of the close relationship between the free compounds and the compounds in the form of their salts, whenever a compound is referred to in this context, a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for the crystallization.

The pharmaceutical compositions according to the invention are those suitable for enteral, such as oral or rectal, and parenteral administration to mammals, including man, for the treatment or prevention of diseases responsive to inhibition of angiotensin-converting enzyme, e.g. cardiovascular diseases such as hypertension and congestive heart failure comprising an effective amount of a pharmacologically active compound of formula I, or pharmaceutically acceptable salts thereof, alone or in combination with one or more pharmaceutically acceptable carriers.

The pharmacologically active compounds of the invention are useful in the manufacture of pharmaceutical compositions comprising an effective amount thereof in conjunction or admixture with excipients or carriers suitable for either enteral or parenteral application. Preferred are tablets and gelatin capsules comprising the active ingredient together with a) diluents, e.g. lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine, b) lubricants, e.g. silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol, for tablets also c) binders, e.g.

magnesium aluminium silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and/or polyvinyl-pyrrolidone, if desired, d) disintegrants, e.g. starches, agar, alginic acid or its sodium salt, or effervescent mixtures and/or e) absorbents, colorants, flavors and sweeteners. Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. Said compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1 to 75%, preferably about 1 to 50%, of the active ingredient. A unit dosage for a mammal of about 50 to 70 kg may contain between about 10 to 200 mg of the active ingredient.

The following Examples are intended to illustrate the invention and are not to be construed as being limitations thereon.

Temperatures are given in degrees Centigrade, and all parts wherever given are parts by weight. If not mentioned otherwise, all evaporation are performed under reduced pressure, preferably between about 15 and 100 mmHg.

In the case of compounds of formula I or IA wherein more than one asymmetric center exists the resulting diastereoisomeric compounds are denoted as A, B, etc., in the said examples. The respective diastereoisomeric compounds are characterized by physical properties, e.g. melting point, relative migration on chromatography, infrared, or nuclear magnetic resonance spectral properties.

In the case of compounds of formula I or IA wherein X is H₂ and an asymmetric center exists in the side chain at the carbon atom bearing the nitrogen atom, the symbols A and B have been assigned as follows to the respective isomers on the basis of their relative migration on chromatography. On the basis of migration on thin-layer chromatography and normal phase high pressure liquid chromatography employing silica gel as the stationary phase, the fast moving isomer is called isomer A and the slow moving isomer is called isomer B. On the basis of migration on reverse phase high pressure liquid chromatography the slow moving isomer is called isomer A and the fast moving isomer is called isomer B.

Example 1:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (Higher melting isomer)

A solution of 3-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (10.0 g) and ethyl benzylpyruvate (26.4 g) in acetic acid (75 ml) and methanol (75 ml) is stirred at room temperature under nitrogen for 1 hour. Sodium cyanoborohydride (3.4 g) in methanol (25 ml) is added dropwise over 4 hours. The reaction mixture is stirred at room temperature for 24 hours. Concentrated hydrochloric acid (4 ml) is added dropwise, and the mixture stirred at room temperature for 1 hour. The reaction mixture is evaporated to dryness. The residue is partitioned between 150 ml of water and 50 ml of ether and adjusted to pH 9 with 40 % aqueous sodium hydroxide solution. The layers are separated and the ether layer is discarded. The aqueous layer is adjusted to pH 4.3 with concentrated hydrochloric acid and extracted with 3 x 75 ml of ethyl acetate. The organic portions are dried (magnesium sulfate) and concentrated to dryness. Hydrogen chloride gas is bubbled into a solution of the crude product in 310 ml of methylene chloride for 5 minutes. The solution is evaporated and the residue is stirred in 225 ml of ether. The product is collected by filtration to give a 70:30 diastereomeric mixture as determined by high pressure liquid chromatography. The product is recrystallized from ethanol/ethyl acetate (1:3) to give 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride melting at 246-248° (decomposition) and corresponding to the racemic isomer B.

A solution of the above hydrochloride salt (0.9 g) and propylene oxide (10 ml) in ethanol (150 ml) is stirred under nitrogen for 18 hours. The solution is evaporated to dryness, and the residue is dissolved in 3 ml of ethanol. Ether (75 ml) is added, precipitating a small

quantity of the starting hydrochloride. The filtrate is evaporated to dryness and stirred with ether/petroleum ether (1:9). The solid is filtered off to give 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenyl-propylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one melting at 139-141°, and being the higher melting racemic isomer B of the compound of formula IB wherein C_nH_{2n} is ethylene, R_6 is ethoxy, R_7 is hydroxy and R_8 is phenyl.

Resolution under standard conditions with an optically active amine and separation of the diastereoisomeric salts yields pure enantiomer, e.g. 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one of Example 12.

Using high pressure liquid chromatography on a reverse phase column (solvent system: methanol, water (3:1) containing 0.025 % acetic acid) isomer B is faster moving than lower melting racemic isomer A of Example 5.

The starting material, 3-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is prepared as follows:

A mixture of 2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (48.3 g, see Briggs et al., J. Chem. Soc. 1937, 456), phosphorus pentachloride (188 g), and xylene (1300 ml) is heated with stirring under an atmosphere of nitrogen to 90° (oil bath temperature) during 30 min with pauses at 30° (to allow the phosphorus pentachloride to dissolve) and at 50°. There is a copious evolution of hydrogen chloride. The temperature is maintained at 90° for 30 minutes. The reaction mixture is filtered while hot to remove a small amount of suspended solid, and the filtrate is evaporated under reduced pressure until all the solvent is removed. The residue is added with stirring to saturated aqueous sodium carbonate (100 ml). The product is filtered after the

solidification process is complete, then slurried in ethanol (150 ml), filtered, washed with ethanol (50 ml) and ether (50 ml) and dried to give 3,3-dichloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 185-187°.

A solution of 3,3-dichloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (20 g, 0.174 mol) and anhydrous sodium acetate (15.4 g, 0.188 mol) in glacial acetic acid (920 ml) is hydrogenated at atmospheric pressure using 5 % Pd-C (1.72 g) as catalyst until the uptake of hydrogen ceases. The catalyst is filtered off and the acetic acid evaporated under reduced pressure. The residue is equilibrated between 10 % NaHCO_3 (900 ml) and dichloromethane (300 ml). The aqueous layer (pH 8) is further extracted with dichloromethane (3 x 300 ml) and the combined organic solutions are dried over anhydrous sodium sulfate and evaporated to give 3-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 163-167°.

A solution of 3-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (15.9 g, 0.08 mol) and sodium azide (6.36 g, 0.10 mol) in dimethylsulfoxide (320 ml) is maintained at 80° under an atmosphere of nitrogen for 3 hours. At this time, the IR spectrum of an aliquot shows a strong peak at 2150 cm^{-1} characteristic of the azide group. The reaction mixture is poured into 1000 ml of ice/water and the suspension is stirred for 30 min. The solid is filtered off, washed with water (250 ml) and dried to give 3-azido-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 142-145°.

A solution of 3-azido-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (8.7 g, 0.043 mol), in dry dimethylformamide (75 ml) is added during 30 min to a stirred suspension of sodium hydride [from 60 % mineral oil dispersion (1.9 g) washed with petroleum ether (3 x 150 ml)] in dry dimethylformamide (250 ml) maintained at 0° under a nitrogen at-

mosphere. Stirring is continued for an additional 1.5 hours, then benzyl bromoacetate (10.8 g; 0.047 mol) in dry dimethylformamide (75 ml) is added during 45 minutes, the temperature being maintained at 0°. The reaction mixture is then allowed to warm to room temperature while stirring for an additional 18 hours. The dimethylformamide is removed under reduced pressure and the residue partitioned between water (500 ml) and dichloromethane (500 ml). The aqueous phase is extracted with additional dichloromethane (3 x 500 ml). The combined extracts are dried over sodium sulfate and the solvent is removed under reduced pressure to give the crude ester-azide as an oil. This material is dissolved in toluene (500 ml) and silica gel (48 g) is added. Filtration and removal of the solvent under reduced pressure gives 3-azido-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one, as an oil, used without further purification in the next synthetic step.

A suspension of Raney nickel active catalyst in water (15 ml) is washed with ethanol (5 x 100 ml) and added to a mechanically stirred solution of 3-azido-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5.0 g) in ethanol (300 ml), and the suspension is stirred for 18 hours at room temperature under nitrogen. The catalyst is filtered off and the solvent removed under reduced pressure. The residue is dissolved in 2H hydrochloric acid (200 ml) and the solution extracted with ether (2 x 250 ml). The aqueous solution is made basic (pH 9) with concentrated aqueous ammonia, and the solution extracted with ether (3 x 200 ml). The combined ether solutions are dried over sodium sulfate and evaporated under reduced pressure to give 3-amino-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as an oil, used without further purification for the next synthetic step.

3-Amino-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benz-

azepin-2-one is also prepared as follows: A solution of 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5.0 g, 0.028 mol), in dimethylformamide (100 ml) is added under a nitrogen atmosphere to a stirred suspension of sodium hydride [prepared from the 60 % mineral oil dispersion (1.2 g) by washing with petroleum ether (3 x 150 ml)] in dimethylformamide (400 ml) to which tetrabutylammonium bromide (10.0 g, 0.031 mol) has been added. The reaction mixture is maintained at 50° for 15 minutes, then a solution of benzyl bromoacetate (7.2 g, 0.031 mol) in dimethylformamide (25 ml) is added. The reaction mixture is stirred for an additional 18 hours at 50°, then cooled to room temperature, and the dimethylformamide removed under high vacuum. The residue is stirred with toluene/dichloromethane (1:1, 500 ml) to precipitate inorganic salts. After filtration, the solution is evaporated under reduced pressure, and the residue chromatographed on silica gel (200 g). Elution with 0-15 % ethyl acetate in toluene gives 3-amino-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as a major product.

A solution of 3-amino-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one (1.3 g) in ethanol (250 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % Pd-C (0.20 g) as catalyst, until uptake of hydrogen ceases. The catalyst is filtered off and the solvent removed under reduced pressure to give a white foam (0.90 g). This material is triturated with ether to give 3-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 147-150°.

A solution of 3-azido-1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (14.0 g, 0.04 mol) in ethanol (300 ml) is hydrogenated for 25 hours at 3.1 at room temperature using 5 % Pd-C (2.0 g) as catalyst. The catalyst is filtered off and the solvent removed under reduced pressure. The residue is dissolved in water

(500 ml) and the solution extracted with dichloromethane (2 x 400 ml). The aqueous solution is filtered, and evaporated under reduced pressure. Ethanol (50 ml) is added and the solution evaporated under reduced pressure. More ethanol (50 ml) is added, and the evaporation repeated. The residue is recrystallized from ethanol/ethyl acetate to give 3-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 147-150°.

Example 2:

1-Benzylloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

Sodium cyanoborohydride (0.152 g, 0.0014 mol) is added to a solution of 1-benzylloxycarbonylmethyl-3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.45 g, 0.0014 mol) and benzylpyruvic acid (0.48 g, 0.0028 mol) in methanol (35 ml). The reaction mixture is stirred at room temperature under nitrogen for 2 hours. Additional benzylpyruvic acid (0.48 g, 0.0028 mol) is added, and the reaction mixture stirred for an additional 18 hours. Concentrated hydrochloric acid (0.5 ml) is added and the resulting solution stirred for 1 hour. The solvents are removed under reduced pressure and the residue is treated with dichloromethane (100 ml) to precipitate sodium chloride. After filtration, the solvent is removed under reduced pressure and the residue chromatographed on silica gel (30 g). Elution with ethyl acetate/methanol/acetic acid (90:10:0.2) gives 1-benzylloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as an oil; NMR(CDC₃) δ 7.35(m,14H), 5.10(s,2H), 4.60(m,2H), 3.00(m,12H).

Example 3:

1-Benzylloxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 1-benzylloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.364 g, 0.00075 mol), sodium bicarbonate (0.190 g, 0.0022 mol), and ethyl iodide (0.315 g, 0.002 mol) in dimethylacetamide (15 ml) is stirred at room temperature under nitrogen for 72 hours. The reaction mixture is filtered and evaporated under reduced pressure. Water (100 ml) is added, and the resulting solution extracted with dichloromethane (4 x 50 ml). The combined extracts are dried over sodium sulfate and the solvent removed under reduced pressure to give the diester as an oil. This material is separated by high pressure liquid chromatography into three fractions, using ethyl acetate/toluene (30:70) as solvent. The first fraction yields isomer A of the title compound as an oil; the second fraction contains a mixture of isomers A and B and the third fraction yields isomer B of the title compound. Using high pressure liquid chromatography on a reverse phase column (solvent system: methanol, water (3:1) containing 0.025 % acetic acid) isomer A moves more slowly than isomer B.

Example 4:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (Higher melting isomer)

A solution of 1-benzylloxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer B of example 3, 0.9 g) in ethanol (150 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % palladium on charcoal (0.5 g) as catalyst. After uptake of hydrogen has ceased, the catalyst is filtered off, and the solvent removed under reduced pressure to give a solid. This material is triturated with ether (8 ml) to give the title compound melting at 138-140° and identical

to the compound obtained in Example 1.

Example 5:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (Lower melting isomer)

A solution of 1-benzyloxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer A of example 3; 1.2 g) in ethanol (125 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % palladium on charcoal (0.5 g) as catalyst. After uptake of hydrogen ceases, the catalyst is filtered off, and the solvent removed under reduced pressure to give a solid. This material is triturated with ether (8 ml) to give 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one melting at 126-129°, and being the lower melting racemic isomer A.

Using high pressure liquid chromatography on a reverse phase column [solvent system: methanol, water (3:1) containing 0.025 % acetic acid] isomer A moves more slowly than higher melting racemic isomer B of Example 1.

Example 6:

1-Benzyloxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5.0 g), in dry dimethylformamide (20 ml) is added under a nitrogen atmosphere to a stirred suspension of sodium hydride [prepared from the 60 % mineral oil dispersion (0.6 g) by washing with petroleum ether (3 x 75 ml)] in dry dimethylformamide (85 ml) to which tetrabutylammonium bromide (4.4 g) has been added. The reaction mixture is stirred at room temperature for

30 minutes, then a solution of benzyl bromoacetate (3.2 g) in dry dimethylformamide (10 ml) is added. The reaction mixture is stirred for an additional 30 minutes at room temperature, heated to 60°, and maintained at that temperature for 18 hours. The reaction mixture is cooled to room temperature, and the solvent removed under high vacuum. Water (150 ml) is added, and the resulting solution extracted with ethyl acetate (2 x 250 ml). The combined ethyl acetate extracts are washed with water (100 ml), dried over magnesium sulfate, and the solvent removed under reduced pressure to give a brown oil. This material is chromatographed on silica gel (150 g). Elution with toluene/ethyl acetate (3:1) first gives isomer A of 1-benzyloxy-carbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one followed by isomer B. Isomer A and B are identical to compounds of Example 3 as determined by high pressure liquid chromatography on a reverse phase column (solvent system: methanol, water (3:1) containing 0.025 % acetic acid).

The starting material is prepared as follows:

A solution of diethyl acetamidomalonate (33.2 g) in ethanol (150 ml) is added to a solution of sodium ethoxide in ethanol [prepared from sodium (3.8 g) and ethanol (200 ml)]. The reaction mixture is stirred at room temperature for 30 minutes and a solution of 2-nitrophenethyl bromide (J. Med. Chem. 20, 1020 (1977), 40.0 g) in ethanol (100 ml) is added dropwise during 20 minutes. After addition is complete, the reaction mixture is refluxed for 18 hours, then cooled to room temperature and evaporated under reduced pressure. The residue is dissolved in water (350 ml) and the solution extracted with ethyl acetate (2 x 350 ml). The combined ethyl acetate extracts are washed with water (200 ml) and dried over magnesium sulfate. Removal of the solvent under reduced pressure gives diethyl 2-acetamido-2-(o-nitrophenethyl)-malonate as a low melting solid, used without further

purification for the next synthetic step.

A solution of diethyl 2-acetamido-2-(o-nitrophenethyl)-malonate (80 g) in 3N hydrochloric acid (900 ml) is refluxed for 12 hours. The solution is cooled and extracted with ethyl acetate (200 ml). The aqueous solution is filtered, and evaporated to dryness under reduced pressure. The residue is recrystallized from ethanol/ether to give 2-amino-4-(2-nitrophenyl)butyric acid hydrochloride, m.p. 219-221° (decomposition).

A solution of 2-amino-4-(2-nitrophenyl)butyric acid hydrochloride (38.0 g) in 10 % ethanolic hydrogen chloride (1200 ml) is refluxed with stirring for 18 hours. The reaction mixture is evaporated to dryness under reduced pressure, water (250 ml) is added, and the aqueous solution made basic by the addition of 2N sodium hydroxide. The solution is extracted with dichloromethane (2 x 500 ml), and the combined dichloromethane solutions washed with water (2 x 150 ml), and dried over anhydrous magnesium sulfate. Evaporation gives ethyl 2-amino-4-(2-nitrophenyl)butyrate, used without further purification for the next synthetic step.

A solution of ethyl 2-amino-4-(2-nitrophenyl)butyrate (27 g) in ethanol (600 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % palladium on charcoal (2.5 g) as catalyst, until hydrogen uptake ceases. The catalyst is filtered off and evaporation to dryness gives ethyl 2-amino-4-(2-aminophenyl)butyrate used without purification for the next synthetic step.

A solution of ethyl 2-amino-4-(2-aminophenyl)butyrate (35.0 g) in methanol (100 ml) is added to a solution of sodium methoxide in methanol [prepared from sodium (1.0 g) and methanol (400 ml)] with stirring, under a nitrogen atmosphere. The reaction mixture is

refluxed for 65 hours and evaporated under reduced pressure. The residue is distributed between water (100 ml) and dichloromethane (400 ml). The aqueous solution is extracted with dichloromethane (400 ml), and the combined organic solutions washed with water (100 ml) and dried over magnesium sulfate. Evaporation to dryness and titration with ether (250 ml) gives 3-amino-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one, m.p. 161-162°.

Alternatively, a solution of 2-amino-4-(2-nitrophenyl)-butyric acid hydrochloride (2.5 g) in water (200 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % Pd-C (0.5 g) as catalyst. After uptake of hydrogen ceases, the catalyst is filtered off, and the filtrate evaporated to dryness. The residue is dissolved in water (50 ml) and the pH adjusted to 7 by the addition of 10 % sodium hydroxide. The solid is filtered off, washed with water, and dried to give 2-amino-4-(2-aminophenyl)butyric acid. A solution of the 2-amino-4-(2-aminophenyl)butyric acid (1.0 g), hexamethyldisilazane (5.4 g), and chlorotrimethylsilane (0.1 g) in xylene (125 ml) is refluxed for 65 hours. The reaction mixture is cooled, poured into ethanol (200 ml) and evaporated under reduced pressure. Water (100 ml) is added, and the solution extracted with dichloromethane (2 x 125 ml). The combined dichloromethane solutions are washed with water (50 ml), dried over magnesium sulfate, and evaporated under reduced pressure to give 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as above.

3-Amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is also prepared as follows:

To a solution of 3-azido-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (see Example 1) (27 g) in ethanol (3500 ml) while stirring at room temperature under an atmosphere of nitrogen, a suspension of Raney nickel in water (50 ml, washed with 10 volumes of ethanol) is added.

The mixture is stirred at room temperature for 2 hours when an additional 30 ml of Raney nickel suspension is added. After stirring for an additional 30 minutes, the catalyst is filtered off and the solvent removed under reduced pressure to give an oil which solidifies on addition of ether to give 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, melting at 161-162°.

A solution of 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (8.0 g) and benzylpyruvic acid (18.0 g) in methanol (450 ml) is stirred at room temperature under nitrogen for 30 minutes. Sodium cyanoborohydride (4.5 g) is added, and the resulting solution stirred at room temperature for 48 hours. Concentrated hydrochloric acid (7 ml) was added dropwise during 10 minutes and stirring is maintained for an additional 1 hour. The reaction mixture is evaporated to dryness, dichloromethane (150 ml) is added, and the mixture stirred for 30 minutes. The solid is filtered off, stirred with water (100 ml) for 15 minutes, then filtered, washed with water (50 ml), and dried to give 3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 173-175° as a mixture of isomers.

A solution of 3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (6.0 g), sodium bicarbonate (4.0 g), and ethyl iodide (11.6 g) in dimethylacetamide (200 ml) is stirred at room temperature under nitrogen for 72 hours. The reaction mixture is filtered and evaporated under high vacuum. Water (250 ml) is added, and the resulting solution extracted with dichloromethane (2 x 400 ml). The combined extracts are dried over magnesium sulfate and the solvent removed under reduced pressure to give 3-(1-ethoxycarbonyl-3-phenyl-propylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as a mixture of isomers. NMR(CDCl₃) δ 9.22(s, 1H), 4.10(2 superimposed quartets, 2H), 1.13(2 superimposed triplets, 3H).

Example 7:

1-Benzylloxycarbonylmethyl-3-(1-benzylloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 3-(1-benzylloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one (4.0 g) in dry dimethylformamide is added under a nitrogen atmosphere to a stirred suspension of sodium hydride [from the 60 % mineral oil dispersion (0.42 g) washed with petroleum ether (3 x 80 ml)] in dry dimethylformamide (100 ml) at room temperature to which tetrabutylammonium bromide (3.1 g) has been added. Stirring is continued for an additional 30 minutes at room temperature, when a solution of benzyl bromoacetate (2.2 g) in dry dimethylformamide (10 ml) is added. After an additional 30 minutes at room temperature, the reaction mixture is heated to 50°, and maintained at that temperature for 18 hours. The reaction mixture is cooled to room temperature, and the solvent removed under high vacuum. Water (150 ml) is added and the solution extracted with ethyl acetate (2 x 300 ml). The combined ethyl acetate solutions are washed with water (100 ml), dried over magnesium sulfate, and the solvent removed under reduced pressure to give a brown oil which is chromatographed on silica gel (250 g). Elution with toluene/ethyl acetate (1:1, 600 ml) gives an oil, characterized as isomer A of the title compound; NMR (CDCl_3) δ 5.12(s, 4H), 4.50(q, 2H). Elution with an additional 2000 ml of the solvent mixture gives an oil characterized as isomer B of the title compound; NMR(CDCl_3) δ 5.17(s, 2H), 5.03(d, 2H), 4.60(q, 2H).

The starting material is prepared as follows:

A solution of 3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (as described in Example 6, 13.0 g), sodium bicarbonate (10.0 g), and benzyl bromide (19.0 g) in dimethylacetamide (750 ml) is stirred at room temperature under a nitrogen

atmosphere for 72 hours. The reaction mixture is filtered and evaporated under high vacuum. Water (150 ml) is added, and the resulting solution extracted with dichloromethane (2 x 400 ml). The combined extracts are washed with water (100 ml), dried over magnesium sulfate and evaporated under reduced pressure to give the crude benzyl ester. Recrystallization from ethyl acetate gives 3-(1-benzyloxy-carbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 139-141°.

Example 8:

1-Carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (Lower melting isomer)

A solution of 1-benzyloxycarbonylmethyl-3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer A of example 7, 2.7 g) in ethanol (800 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % palladium on charcoal (0.5 g) as catalyst. After uptake of hydrogen has terminated, the catalyst is filtered off, and the solvent removed under reduced pressure to give the title diacid, characterized as isomer A, m.p. 256-259°.

The identical compound is obtained on hydrolysis of the compound of Example 5.

Example 9:

1-Carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (Higher melting isomer)

A solution of 1-benzyloxycarbonylmethyl-3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer B of Example 7, 5.0 g) in ethanol (950 ml) is hydrogenated at room temperature and atmospheric pressure, using palladium on charcoal

(0.5 g) as catalyst. After uptake of hydrogen has terminated, the catalyst is filtered off, and the solvent removed under reduced pressure to give the title diacid, characterized as isomer B, m.p. 280-282°.

The identical compound is obtained on hydrolysis of the compound of Example 1 (isomer B) or compound of Example 10 (isomer B).

Example 10:

1-Ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (see Example 6, 3.0 g) in dry dimethylformamide (10 ml) is added dropwise during 10 minutes to a stirred suspension of sodium hydride [from the 60 % mineral oil dispersion (0.36 g) washed with petroleum ether (3 x 75 ml)] in dry dimethylformamide (100 ml) at room temperature under nitrogen. Stirring is maintained for an additional 30 minutes, a solution of ethyl bromoacetate (1.4 g) in dimethylformamide (15 ml) is added and the reaction mixture is maintained at 60° for 48 hours. After the reaction mixture is cooled to room temperature, the solvent is removed under high vacuum. Water (100 ml) is added, and the solution extracted with ethyl acetate (2 x 200 ml). The combined ethyl acetate solutions are washed with water (50 ml), dried over magnesium sulfate, and the solvent removed under reduced pressure to give a yellow oil (3.8 g). This material is chromatographed on silica gel (120 g). Elution with toluene/ethyl acetate (1:1; 250 ml) gives isomer A of the desired product. Elution with an additional 250 ml of solvent mixture gives an oil which contains mostly isomer B and some isomer A of the desired product as determined by analytical high pressure liquid chromatography (see Example 6). Elution with a further 250 ml of solvent mixture gives an oil which is essentially pure isomer B.

(slower moving). This material is dissolved in methanol (25 ml) and converted to the maleate salt by addition of an equimolar quantity of maleic acid in methanol. Evaporation of the solvent and recrystallization of the residue from methanol/ether yields pure isomer B of 1-ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as the maleate salt melting at 114-116°.

Example 11:

1-Carboxymethyl-3-carboxymethylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 1-benzyloxycarbonylmethyl-3-benzyloxycarbonylmethylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (4.8 g; 0.01 mol) in ethanol (550 ml) is hydrogenated at room temperature and atmospheric pressure using 5 % Pd-C (0.85 g) as catalyst until uptake of hydrogen ceases. Water (300 ml) is added, the catalyst filtered off, and the solvent removed under reduced pressure. The residue is triturated with ether to give the title diacid, m.p. 232-236°.

The starting material is prepared as follows:

A solution of 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5.0 g, 0.028 mol) in dimethylformamide (100 ml) is added under a nitrogen atmosphere to a stirred suspension of sodium hydride [prepared from the 60 % mineral oil dispersion (1.2 g) by washing with petroleum ether (3 x 150 ml)] in dimethylformamide (400 ml) to which tetrabutylammonium bromide (10.9 g, 0.031 mol) has been added. The reaction mixture is maintained at 50° for 15 minutes, then a solution of benzyl bromoacetate (7.2 g, 0.031 mol) in dimethylformamide (25 ml) is added. The reaction mixture is stirred for an additional 18 hours at 50°, then cooled to room temperature, and the dimethylformamide removed under high vacuum. The residue is stirred with toluene/dichloro-

methane (1:1, 500 ml) to precipitate inorganic salts. After filtration, the solution is evaporated under reduced pressure, and the residue chromatographed on silica gel (200 g). Elution with 0-15 % ethyl acetate in toluene gives 1-benzyloxycarbonylmethyl-3-benzyloxycarbonylmethylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as the first fraction. Further elution gives 3-benzyloxycarbonylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 124-127° and 3-amino-1-benzyl-oxy carbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (see Example 1).

Example 12:

1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

3(S)-Amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one when treated with ethyl benzylpyruvate in the presence of sodium cyanoborohydride by the procedure described in Example 1 for the racemic compound gives after purification 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, as described below.

A solution of sodium hydroxide (2.1 g) in water (5 ml) is added to a solution of 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (14.0 g) in methanol (150 ml) at room temperature, and the solution is stirred for two hours. The solvents are evaporated and the residue is thoroughly dried, then slurried with ether, to give 3(S)-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one sodium salt. This is used without further purification.

A solution of the above sodium salt (12.9 g) and ethyl benzylpyruvate (31 g) in acetic acid (100 ml) and methanol (75 ml) is stirred at room temperature under a dry nitrogen atmosphere for one hour. A solution

of sodium cyanoborohydride (3.8 g) in methanol (30 ml) is then added dropwise over a 4 hour period. The combined solutions are stirred overnight at room temperature. Concentrated hydrochloric acid (10 ml) is added dropwise and the mixture stirred at room temperature for 1 hour followed by the evaporation of solvents. The residue is partitioned between water (400 ml) and ether (100 ml) and the pH adjusted to 9.3 with 40 % sodium hydroxide. The layers are separated and the ether layer is discarded. The aqueous layer is adjusted to pH 4.3 with concentrated hydrochloric acid and extracted with ethyl acetate (3 x 100 ml). The organic phases are combined, dried (magnesium sulfate), and evaporated. Hydrogen chloride gas is bubbled through a solution of the crude product in methylene chloride (150 ml) for 5 minutes. The solvent is evaporated and the resulting foam is dissolved in hot methyl ethyl ketone (100 ml). The solid which precipitated is collected by filtration to give a 95:5 diastereomeric mixture as determined by high pressure liquid chromatography. The product is recrystallized from 3-pentanone/methanol (10:1) to give 1-carboxy-methyl-3(S)-(1(S)-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride, m.p. 188-190°, $[\alpha]_D = -141.0^\circ$ ($c=0.9$ in ethanol), of formula IIa wherein $C_{nH_{2n}}$ is ethylene, R_6 is ethoxy, R_7 is hydroxy and R_8 is phenyl.

A solution of the above hydrochloride salt (0.035 g) and propylene oxide (0.5 ml) in ethanol (4 ml) is stirred under nitrogen overnight at room temperature. The solution is evaporated to dryness. Ether (2 ml) is added, and the solid is filtered off to give 1-carboxy-methyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 148-149°, $[\alpha]_D = -159^\circ$ ($c = 1.2$ in ethanol).

The optically active starting material is prepared as indicated below.

a) A solution of 0.4 g of 3(S)-t-butyloxycarbonylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione [prepared from L-kynurenine as described in Australian J. Chemistry Vol. 33, 633-40 (1980)], and ethyl bromoacetate (0.23 g) in dry tetrahydrofuran (30 ml) is stirred at 0° under a dry nitrogen atmosphere. Potassium t-butoxide (0.254 g) is added in one portion. After 1 hour at 0°, an additional quantity of ethyl bromoacetate (0.23 g) is added and the reaction mixture is stirred at 0° for a further 1 hour. Water (100 ml) is added and the mixture is extracted with ethyl acetate (2 x 50 ml). The combined ethyl acetate solutions are washed with water (100 ml) and dried over magnesium sulfate. Removal of the solvent under reduced pressure gives a yellow gum which on trituration with ether/petroleum ether (bp 30-60°) gives 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione, m.p. 86-88°, $[\alpha]_D = -203^\circ$ (c = 1 in dimethylformamide).

A solution of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione (0.14 g) and sodium borohydride (7 mg) in ethanol (10 ml) is stirred at room temperature for 18 hours. The ethanol is removed under reduced pressure, and the residue dissolved in dichloromethane (25 ml). The solution is extracted with 2N hydrochloric acid (2 x 20 ml) and saturated aqueous sodium chloride solution (20 ml), and dried over sodium sulfate. The solvent is removed under reduced pressure, and the residue triturated with ether to give 3(S)-t-butyloxycarbonylamino-1-ethoxy-carbonylmethyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 167-169.5°, $[\alpha]_D = -193^\circ$ (c = 0.52 in dimethylformamide). The substance is also obtained by hydrogenation of the benzazepin-2,5-dione derivative with H₂/Pt in ethanol.

A mixture of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.076 g), dicyclo-

hexylcarbodiimide (0.064 g) and cuprous chloride (7 mg) is heated at 60° under nitrogen for 32 hours. The reaction mixture is allowed to cool to room temperature. The residue is dissolved in methylene chloride (50 ml) and washed with dilute ammonium hydroxide (2 x 15 ml) followed by water (20 ml). The organic phase is dried over sodium sulfate and evaporated to give a mixture of the desired adduct and excess dicyclohexylcarbodiimide.

This mixture (0.100 g) is dissolved in ethyl acetate (40 ml) and placed in a pressure bottle. 10 % Pd/C (0.010 g) is added and the mixture is hydrogenated at 3 atmospheres pressure and at 40° for 16 hours. The catalyst is filtered off and the filtrate evaporated. The residue is triturated with ether, and the ether solution evaporated to give 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 115-116.5°, $[\alpha]_D = -182^\circ$ (c = 2.6 in dimethylformamide).

b) Tartaric acid (12.6 g) and racemic 3-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (22 g) are dissolved in hot ethanol (200 ml). This solution is cooled and allowed to stand overnight at room temperature. The solid which precipitates is collected by filtration and recrystallized twice from ethanol (200 ml) to give 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one tartrate salt. This is dissolved in water (100 ml) and the pH adjusted to 9 with dilute ammonium hydroxide and extracted with methylene chloride (2 x 50 ml). The combined extracts are washed with water (75 ml), dried (magnesium sulfate) and evaporated to give 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 104-106°, $[\alpha]_D = -285.5^\circ$ (c = 0.99 in ethanol).

c) Hydrogen chloride gas is bubbled through a solution of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one (under a) above, 0.225 g) in ethyl acetate (25 ml)

for 45 minutes. Nitrogen is then bubbled through this solution for 30 minutes. The ethyl acetate is washed with water (30 ml) and 1N hydrochloric acid (30 ml). The ethyl acetate layer is discarded and the aqueous phases are combined. The aqueous solution is adjusted to pH 9 with dilute ammonium hydroxide, extracted with ethyl acetate (3 x 50 ml); the organic phases are combined, dried (sodium sulfate) and evaporated to give 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 101-102°, $[\alpha]_D = -298^\circ$ (c = 0.46 in ethanol).

Treatment with ethanedithiol/boron trifluoride etherate or trifluoroacetic acid/anisole to remove the protecting group yields 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

Alternately 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is also prepared as follows:

d) A solution of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (previously described, 1.0 g) in acetic anhydride (20 ml) is maintained at 80° for 3 hours. The reaction mixture is cooled to room temperature and the solvents are removed under reduced pressure. Ether (100 ml) is added, and the resulting solution washed with water (50 ml) and dried over magnesium sulfate. The solvent is removed under reduced pressure to give 5-acetoxy-3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as a pale yellow oil which is used without further purification.

A solution of 5-acetoxy-3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.7 g) in ethanol (50 ml) is hydrogenated at 2.9 atmospheres for 24 hours at 70° using

10 % palladium on charcoal (0.5 g) as catalyst. The catalyst is filtered off and the solvent removed under reduced pressure to give 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one which, without further purification is converted to 3(S)-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one by the procedure described above; m.p. 99-101°, $[\alpha]_D = -297^\circ$ (c = 1 in ethanol).

e) A solution of 3(S)-t-butyloxycarbonylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione (12.5 g) prepared from L-kynurenine as described in Australian J. Chemistry Vol. 33, 633-40 (1980), and t-butyl bromoacetate (10.1 g) in acetone (700 ml) is stirred at room temperature under a dry nitrogen atmosphere. Potassium carbonate (12.5 g) is added in one portion and the resulting suspension is stirred at room temperature for 16 hours. The potassium salts are filtered off and the filtrate evaporated to dryness. The residue is partitioned between ethyl acetate (250 ml) and water (250 ml). The layers are separated and the organic phase is dried (sodium sulfate). The residue is triturated with petroleum ether (350 ml; bp 30-60°) to give 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione, m.p. 75-77°, $[\alpha]_D = -172^\circ$ (c = 0.96 in dimethylformamide).

A solution of 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,5-dione (8.0 g) in ethanol (500 ml) containing platinum oxide (800 mg) is hydrogenated at atmospheric pressure and at room temperature for two hours. The catalyst is filtered off and the filtrate evaporated to give 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, $[\alpha]_D = -173^\circ$ (c = 1.8 in dimethylformamide).

A suspension of 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonyl-methyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (3.0 g), dicyclohexylcarbodiimide (5.0 g), and cuprous chloride (500 mg) is mechanically stirred and heated at 80° for 16 hours under a dry nitrogen atmosphere. The mixture is cooled, diluted with methylene chloride (100 ml), and filtered. The solids are discarded. The filtrate is washed with 7 % ammonium hydroxide (4 x 75 ml), followed by 1 x 100 ml with water and saturated aqueous sodium chloride solution (100 ml). The organic phase is dried (sodium sulfate) and evaporated to give a mixture of the desired adduct and excess dicyclohexylcarbodiimide.

This mixture (5.5 g) is dissolved in ethyl acetate (200 ml) and placed in a pressure bottle. 10 % Pd/C (3.0 g) is added and the mixture is hydrogenated at 3 atmospheres pressure and at 40° for 16 hours. The catalyst is filtered off and the filtrate evaporated. The residue is triturated with ether (75 ml) to give a white solid, 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one, m.p. 145-147°, $[\alpha]_D = -194^\circ$ (c = 0.46 in dimethylformamide).

A solution of 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (described above, 3.0 g) in acetic anhydride (50 ml) is heated at 80° under a dry nitrogen atmosphere for 2 hours. The acetic anhydride is evaporated. The residue is dissolved in ethyl acetate (75 ml) and washed with saturated aqueous sodium bicarbonate solution (50 ml), water (50 ml), and saturated aqueous sodium chloride solution (50 ml). The organic phase is dried (sodium sulfate), evaporated, and the residue triturated with ether (50 ml) to give 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-5-acetoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 164-166.5°, $[\alpha]_D = -169^\circ$ (c = 0.36 in dimethylformamide).

A solution of 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-5-acetoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (2.2 g) in ethanol (300 ml) containing 10 % Pd/C (2.0 g) is placed in a pressure bottle and hydrogenated at 3 atmospheres pressure and 70° for 3 days. The catalyst is filtered off and the filtrate evaporated to give 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 164-165°, $[\alpha]_D = -200.6^\circ$ (c = 0.64 in dimethylformamide).

Hydrogen chloride gas is bubbled through a solution of 3(S)-t-butyloxycarbonylamino-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one (0.85 g) in ethyl acetate (40 ml) for 2 hours. Nitrogen is then bubbled through the solution for 0.5 hour. The ethyl acetate is evaporated and the white solid residue immediately dissolved in ethanol (40 ml). Propylene oxide (5 ml) is added and the mixture is stirred at room temperature for 16 hours. The white solid which precipitates is collected by filtration to give 3(S)-amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 275-276°, $[\alpha]_D = -287^\circ$ (c = 0.71, in 1N hydrochloric acid) which is condensed with ethyl benzylpyruvate in the presence of sodium cyanoborohydride as described above.

Example 13:

1-Carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

To a solution of 1-benzyloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,5-dihydro-1H-[1]benzazepin-2,5-dione (1.00 g) in glacial acetic acid (50 ml) is added platinum oxide (0.10 g). The resulting mixture contained in a pressure bottle is hydrogenated at 2.9 atmospheres for 5 hours. The catalyst is removed by filtration, the filtrate concentrated, and the resulting oil triturated with anhydrous ethanol. The resulting solid is collected, dried, and suspended in

water (10 ml). The suspension is stirred for 1.5 hours. The solid is collected and dried to give impure 1-carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one melting with decomposition at 179°.

The starting material is prepared as follows:

Benzyl bromoacetate (9.16 g, 0.04 mole) is added dropwise to a mixture of 3-methoxy-2,5-dihydro-1H-[1]benzazepine-2,5-dione [8.13 g, 0.04 mol, prepared as described in the Canadian J. Chem., 52, 610 (1974)] powdered potassium hydroxide (2.24 g, 0.04 mole) and tetrabutyl-ammonium bromide (1.29 g, 0.004 mol) in 1000 ml of acetonitrile with stirring at room temperature. Upon complete addition, the suspension is stirred at room temperature for 64 hours, filtered, and the filtrate concentrated under reduced pressure to give a partially crystalline oil. This oil is triturated with ether to give a solid which is suspended and stirred in ethyl acetate (100 ml) for 1.5 hours. The insoluble material is filtered off and the filtrate concentrated to give the crude 1-benzyloxycarbonylmethyl-3-methoxy-2,5-dihydro-1H-[1]-benzazepin-2,5-dione which is used directly in the next step.

To a 1.0M solution of potassium t-butoxide (0.64 g, 0.0057 mol) in t-butanol (5.7 ml), while stirring under nitrogen at room temperature is added (+)-homophenylalanine (1.02 g; 0.0057 mol) in one portion. The resulting suspension and t-butanol (4.3 ml) is heated until most of the suspended solid is dissolved. Upon cooling, a suspension is obtained. This suspension is added, in portions, via pipette to a refluxing solution of 1-benzyloxycarbonylmethyl-3-methoxy-2,5-dihydro-1H-[1]benzazepin-2,5-dione (2.00 g) in t-butanol (40 ml) stirring under nitrogen over a period of 10 minutes. During the addition, a yellow precipitate forms. Upon complete addition, the resulting suspension is refluxed for 3 hours. The suspension is filtered, the

resulting gummy solid is washed with petroleum ether and dissolved in water (20 ml). The solution is filtered, acidified to pH 5 with 3N hydrochloric acid, the resulting crude 1-benzyloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,5-dihydro-1H-[1]benzazepin-2,5-dione is collected and used directly for the preparation of the title compound.

Example 14:

Analogous to the methods disclosed herein, the following compounds of formula IA wherein X = H₂, R₂ and R₅ = H, R₆ = OC₂H₅ and R₇ = OH are prepared:

No.	<u>R₁</u>	<u>R₃</u>	<u>R₄</u>
1	C ₆ H ₅ CH ₂	H	H
2	C ₆ H ₅ CH ₂ CH ₂	7-OCH ₃	8-OCH ₃
3	C ₆ H ₅ CH ₂ CH ₂	7-Cl	H
4	C ₆ H ₅ CH ₂ CH ₂	8-CH ₃	H
5	C ₆ H ₅ CH ₂ CH ₂	8-OCH ₃	H
6	p-ClC ₆ H ₄ CH ₂ CH ₂	H	H
7	CH ₃	H	H

The starting substituted 2,3,4,5-tetrahydro-1H-[1]benzazepin-2-ones for compounds 2-5 are prepared as follows:

The 7-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 164-165°, is prepared as described in British Patent 1,359,285.

The 8-methyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is prepared by the method of Huisgen, Liebigs Ann. Chem. 574, 171 (1951), m.p. 153-154°.

The 7,8 dimethoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is prepared as follows:

A solution of 24 g of 6,7-dimethoxy- α -tetralone [Snider, T. et al, Org. Prep. Proced. Int., 5, 291 (1973)] in ethanol (300 ml) and water (60 ml) is treated at reflux for two hours with hydroxylamine hydrochloride (16 g) and sodium hydroxide (25 g) to form the oxime. The reaction mixture is poured into 500 ml of an ice/water mixture and extracted with 3 x 300 ml portions of dichloromethane. The combined extracts are washed with 200 ml water, dried over anhydrous magnesium sulfate and evaporated to yield 25 g of the oxime, m.p. 154-156°.

The oxime is redissolved in 170 ml of dichloromethane and 170 ml of polyphosphate ester (Fieser and Fieser: Reagents for Organic Synthesis, Wiley N.Y. 1967, P. 892) was added. The reaction mixture is refluxed for 18 hours. The dichloromethane layer is separated, treated with charcoal and dried over magnesium sulfate to yield the 7,8-dimethoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 153-156°.

The 8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 132-134° is similarly prepared from 7 methoxy- α -tetralone.

3-Amino-7-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is synthesized as follows:

A solution of 3-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (4.0 g), 2-t-butylloxycarbonyloxyimino-2-phenylacetonitrile (6.1 g) and triethylamine (5 ml) in water (20 ml) and dioxane (25 ml) is stirred at room temperature for 18 hours. The resulting solid is filtered off and washed with water. Recrystallization from ethyl acetate gives 3-t-butylloxycarbonylamino-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one, m.p. 199-201°.

Chlorine gas is bubbled through a solution of 3-t-butylloxycarbonyl-amino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (1.5 g) in acetic acid

(20 ml) for 10 minutes. The reaction mixture is stirred for an additional 10 minutes. The solid which precipitates is collected, suspended in water (30 ml) and aqueous ammonia is added until basic. Filtration gives 3-amino-7-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 170-171°.

Example 15: Preparation of 10,000 tablets each containing 10 mg of the active ingredient of Example 1:

Formula:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropyl-amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one	100 g
Lactose	1,157 g
Corn starch	75 g
Polyethylene glycol 6,000	75 g
Talcum powder	75 g
Magnesium stearate	18 g
Purified water	q.s.

Procedure:

All the powders are passed through a screen with openings of 0.6 mm. Then the drug substance, lactose, talcum, magnesium stearate and half of the starch are mixed in a suitable mixer. The other half of the starch is suspended in 40 ml of water and the suspension added to the boiling solution of the polyethylene glycol in 150 ml of water. The paste formed is added to the powders which are granulated, if necessary, with an additional amount of water. The granulate is dried overnight at 35°, broken on a screen with 1.2 mm openings and compressed into tablets using concave punches with 6.4 mm diameter, uppers bisected.

Example 16: Preparation of an injectable formulation containing 25 mg of the active ingredient of Example 1 per 5 ml of solution:

Formula:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropyl-amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride	25.0 g
Propylparaben	1.0 g
Water for injection q.s.	5000.0 ml

Procedure:

The active ingredient and preservative are dissolved in 3500 ml of water for injection and the solution is diluted to 5000 ml. The solution is filtered through a sterile filter and filled into injection vials under sterile conditions each vial containing 5 ml of the solution.

Example 17: Preparation of 10,000 capsules each containing 20 mg of the active ingredient of Example 9.

Formula:

1-Carboxymethyl-3-(1-carboxy-3-phenylpropyl-amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one	200 g
Lactose	1,700 g
Talcum powder	100 g

Procedure: All the powders are passed through a screen with openings of 0.6 mm. Then the drug substance is placed in a suitable mixer and mixed first with the talcum, then with the lactose until homogenous. No. 3 capsules are filled with 200 mg; using a capsule filling machine.

Analogously, tablets, injectable formulations or capsules are prepared from the remaining compounds of the invention, e.g., those illustrated by the Examples herein.

Example 18:

1-Carboxymethyl-3S-(1R-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

The methyl ethyl ketone filtrate from the crystallization of 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride in Example 12 is evaporated, and the residue triturated with ethyl acetate (50 ml). The resulting solid is distributed between ethyl acetate (100 ml) and water (100 ml), and adjusted to pH 4.3 with concentrated hydrochloric acid. The layers are separated and the aqueous phase is extracted with ethyl acetate (2 x 100 ml). The combined ethyl acetate solutions are dried over sodium sulfate and the solvent removed under reduced pressure. The residue is separated into its components by high pressure liquid chromatography with a C₁₈ reverse phase preparative column and using water/methanol (3:7) containing 0.05 % acetic acid as the solvent. An additional quantity of the S,S isomer of Example 12 is thus obtained, as well as the S,R isomer. The material corresponding to the S,R isomer is dissolved in dichloromethane (75 ml), and hydrogen chloride gas bubbled in for five minutes. The solvent is evaporated under reduced pressure and the residue recrystallized from methyl ethyl ketone to give 1-carboxymethyl-3S-(1R-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride, m.p. 181-183°, [α]_D = -188° (c = 0.8 in ethanol).

Example 19:

1-Carboxymethyl-3S-(1S-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of sodium hydroxide (0.27 g) in water (2 ml) is added to a solution of 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropyl-

amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride (1 g) in methanol (10 ml). The reaction mixture is stirred for 18 hours at room temperature and the solvents removed under reduced pressure. The residue is dissolved in water (25 ml), and the pH adjusted to 3 by the addition of 4N hydrochloric acid. The resulting solid is filtered off, washed with water, and dried to give 1-carboxymethyl-3S-(1S-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 270-272°, $[\alpha]_D = -200.5^\circ$ (c = 1, in 3 % aqueous ammonia).

Example 20:

1-Ethoxycarbonylmethyl-3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5.0 g) in dry dimethylformamide is added under a nitrogen atmosphere to a stirred suspension of sodium hydride [from the 60 % mineral oil dispersion (0.5 g) washed with petroleum ether (3 x 80 ml)] in dry dimethylformamide (100 ml) at room temperature. Stirring is continued for an additional 30 minutes at room temperature, when a solution of ethyl bromoacetate (2.0 g) in dry dimethylformamide (10 ml) is added. After an additional 30 minutes at room temperature the reaction mixture is heated to 50°, and maintained at that temperature for 18 hours. The reaction mixture is cooled to room temperature and the solvent removed under high vacuum. Water (150 ml) is added and the solution extracted with ethyl acetate (2 x 300 ml). The combined ethyl acetate solutions are washed with water (100 ml), dried over magnesium sulfate, and the solvent removed under reduced pressure to give a brown oil which is chromatographed on silica gel (250 g). Elution with toluene/ethyl acetate (9:1; 600 ml) gives an oil, characterized as isomer A of the title compound. Elution with an additional 1000 ml of the solvent mixture gives an oil characterized as isomer B of the title compound.

Example 21:

1-Ethoxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

1-Ethoxycarbonylmethyl-3-(1-benzyloxycarbonyl-3-phenylpropylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer B of Example 20, 1.1 g) in ethanol (150 ml) is hydrogenated at room temperature and atmospheric pressure using palladium on charcoal (0.5 g) as catalyst. After uptake of hydrogen terminates, the catalyst is filtered off, and the solvent removed under reduced pressure to give a semi-solid. Trituration with ether (30 ml) yielded isomer B of the title compound, m.p. 175-177°.

Example 22:

1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (isomer B)

A solution of 3-amino-1-carboxymethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (4.0 g) and ethyl benzylpyruvate (9.4 g) in a mixture of acetic acid (35 ml) and methanol (35 ml) is stirred for 1 hour. A solution of sodium cyanoborohydride (1.1 g) in methanol (50 ml) is then added slowly over the course of 5 hours. After stirring an additional 16 hours, concentrated hydrochloric acid (4 ml) is added and stirring is continued for 1 hour. The solvents are removed at reduced pressure and the residue is partitioned between water (75 ml) and ether (35 ml). The pH is adjusted to 9.4 and the ether layer is separated and discarded. The aqueous layer is acidified to pH 4.3 and extracted with ethyl acetate (3 x 50 ml). The combined ethyl acetate solutions are dried over magnesium sulfate and the solvent is removed at reduced pressure. Hydrogen chloride gas is bubbled into a solution of the crude product in methylene chloride (100 ml) for 5 minutes. The solution is evaporated and the residue is stirred in ether (75 ml). The product is collected by filtration to

give an approximately 70:30 diastereomeric mixture as determined by high pressure liquid chromatography.

The product is recrystallized from 3-pentanone to give 1-carboxymethyl-8-methoxy-3-(1-ethoxycarbonyl-3-phenylpropyl-amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride (isomer B) melting at 240-245° (decomposition).

The starting material is prepared as follows: A solution of 8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (7.0 g, described in Example 14) and phosphorus pentachloride (30.0 g) in xylene (200 ml) is heated with stirring under an atmosphere of nitrogen to 90° (oil bath temperature) during 30 minutes with pauses at 30° and at 50°. There is a copious evolution of hydrogen chloride gas. The temperature is maintained at 90° for 30 minutes. The reaction mixture is filtered while hot to remove a small amount of suspended solid, and the filtrate is evaporated under reduced pressure until all the solvent is removed. The residue is added with stirring to saturated aqueous sodium carbonate (20 ml). The product is filtered after the solidification process is complete, slurried in ethanol (30 ml), washed with ethanol (10 ml) and ether (10 ml) and dried to give 3,3-dichloro-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 148-150°.

A solution of 3,3-dichloro-8-methoxy-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one (20g) and anhydrous sodium acetate (13.2 g) in glacial acetic acid (250 ml) is hydrogenated at atmospheric pressure using 10 % Pd/C (1g) as catalyst, until the uptake of hydrogen ceases. The catalyst is filtered off and the acetic acid is evaporated under reduced pressure. Water (100 ml) is added to the residue and the suspension stirred for 1 hour. The solid is filtered, washed with

water (50 ml), and dried to give 3-chloro-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 162-163°.

A solution of 3-chloro-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (12.5 g) and sodium azide (4.3 g) in dimethylsulfoxide (150 ml) is maintained at 80° under an atmosphere of nitrogen for 3 hours. The reaction mixture is poured into ice/water (300 ml) and the suspension is stirred for 30 minutes. The solid is filtered off, washed with water (50 ml) and dried to give 3-azido-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 136-138°.

3-Azido-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5g) is added in one portion to a stirred suspension of potassium hydroxide (1.3 g) and tetrabutylammonium bromide (0.7 g) in tetrahydrofuran (50 ml) maintained at 0° under a nitrogen atmosphere. Stirring is continued for 5 minutes, then a solution of ethyl bromoacetate (3.6 g) in tetrahydrofuran (15 ml) is added during 5 minutes. The reaction mixture is allowed to warm to room temperature while stirring for an additional 2 hours. The reaction mixture is filtered and the tetrahydrofuran is removed at reduced pressure. The residue is partitioned between water (50 ml) and ether (100 ml). The organic phase is washed with 2N hydrochloric acid (10 ml), dried over magnesium sulfate and the solvent removed under reduced pressure to give 3-azido-1-ethoxycarbonylmethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 90-91°.

A suspension of 3-azido-1-ethoxycarbonylmethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (13.8 g) in methanol (75 ml) is treated with a solution of sodium hydroxide (1.9 g) in water (75 ml). The reaction mixture is stirred at 40-45° for 2 hours. Water (100 ml) is added and the mixture is acidified with concentrated hydrochloric acid (10 ml) and extracted with methylene chloride (3 x 75 ml). The

combined methylene chloride solutions are dried over magnesium sulfate and evaporated at reduced pressure to give 3-azido-1-carboxymethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 145-147°.

A solution of 3-azido-1-carboxymethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (11g) in a mixture of ethanol (250 ml) and water (50 ml) is hydrogenated for 3 hours at 3 atmospheres pressure and room temperature using 10 % Pd-C (0.5 g) as catalyst. 2N Hydrochloric acid (50 ml) is added, and the catalyst is filtered off. The solvent is removed at reduced pressure, and the residue dissolved in a mixture of water (50 ml) and ethanol (50 ml). Propylene oxide (25 ml) is added and the mixture is stirred for 1 hour. The solvents are removed under reduced pressure to give 3-amino-1-carboxymethyl-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. > 300°.

Example 23:

1-(1-Carboxyethyl)-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride

A solution of 3-amino-1-(1-carboxyethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride (3 g) and ethyl benzylpyruvate (6.5 g) in acetic acid (30 ml) and methanol (30 ml) is stirred at room temperature for 1 hour. Sodium cyanoborohydride (0.8 g) in methanol (10 ml) is added over 4 hours. The reaction mixture is stirred at room temperature for 24 hours. Concentrated hydrochloric acid (2 ml) is added and the mixture is stirred for 1 hour. The solvents are removed at reduced pressure and the residue is partitioned between water (50 ml) and ether (30 ml). The pH is adjusted to 9.4, the ether layer is separated and discarded. The aqueous solution is adjusted to pH 4.3 and extracted with ethyl acetate (3 x 50 ml). The combined ethyl acetate solutions are dried over magnesium sulfate and the solvent removed under reduced pressure. Hydrogen chloride is bubbled

into a solution of the crude product in methylene chloride (10 ml) for 2 minutes. The solution is evaporated to give 1-(1-carboxyethyl)-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one hydrochloride, as a mixture of diastereomers, m.p. 87-94°.

The starting material is prepared as follows: 3-Azido-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (as prepared in Example 1, 5 g) is added in one portion to a stirred suspension of potassium hydroxide (1.8 g) and tetrabutylammonium bromide (0.8 g) in tetrahydrofuran (50 ml) maintained at 0° under a nitrogen atmosphere. Stirring is continued for 5 minutes, then (R)-t-butyl 2-bromopropionate [J.P. Greenstein et al., J. Am. Chem. Soc. 76, 6054 (1954), H. Niedrich and G. Koller, J. Prakt. Chem. 316, 729 (1974)] (5.2 g) in tetrahydrofuran (15 ml) is added during 5 minutes. The reaction mixture is allowed to warm to room temperature while stirring for an additional 2 hours. The reaction mixture is filtered and the tetrahydrofuran removed at reduced pressure. The residue is partitioned between water (50 ml) and ether (100 ml). The organic phase is washed with 2N hydrochloric acid (10 ml), dried over manganese sulfate, and the solvent evaporated under reduced pressure to give 3-azido-1-(1-t-butyloxycarbonylethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as an oil that is used without further purification.

A solution of 3-azido-1-(1-t-butyloxycarbonylethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (7g) in ethanol (70 ml) is hydrogenated at 3 atmospheres pressure for 3 hours using 10 % Pd-C (0.5 g) as catalyst. The catalyst is removed by filtration and the ethanol removed under reduced pressure to give 3-amino-1-(1-t-butyloxycarbonylethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one as an oil. High pressure liquid chromatography (HPLC) indicates that the product is an approximately 1:1 mixture of diastereomers. This material is

used without further purification.

A solution of the above 3-amino-1-(1-t-butyloxycarbonylethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (4.7 g) in trifluoroacetic acid (25 ml) is stirred at room temperature for 1 hour. The trifluoroacetic acid is removed under reduced pressure and the residue dissolved in ether (100 ml). Hydrogen chloride gas is bubbled into the solution until precipitation ceases. The solid is collected by filtration to give 3-amino-1-(1-carboxyethyl)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride, m.p. 165-176°. HPLC indicated that the product is an approximately 1:1 mixture of diastereomers.

Example 24:

1-Ethoxycarbonylmethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of 3S-amino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (1.5 g), ethyl 2-bromo-4-phenylbutyrate (1.6 g), and triethylamine (0.8 ml) in dimethylformamide (37 ml) is stirred under nitrogen for 18 hours at 70°. The dimethylformamide is then removed under reduced pressure. The residue is taken up in ethyl acetate (70 ml), washed with water (5 x 25 ml), dried over magnesium sulfate, and evaoprated. The product mixture is then separated on a silica gel chromatography system with ethyl acetate/hexane (40:60) as solvent to yield about equal quantities of 1-ethoxycarbonylmethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one [NMR (CDCl_3) δ 4.52 (q, 2H)] the S,S enantiomer of the compound of Example 10 and its diastereomer, 1-ethoxycarbonylmethyl-3S-(1R-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one; NMR (CDCl_3): δ 4.50 (q, 2H). TLC: (silica gel, ethyl acetate/hexane 40:60): the (S,S) isomer has R_f = 0.24 and the (S,R) isomer R_f = 0.33.

Example 25:

1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

2N Potassium hydroxide (0.26 ml) is added dropwise to a solution of 1-ethoxycarbonylmethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.25 g) in ethanol (5 ml), while stirring at room temperature under a nitrogen atmosphere. After stirring for one hour the ethanol is evaporated and the residue is dissolved in water (5 ml), acidified with 2N hydrochloric acid to pH 2 and extracted with ethyl acetate (2 x 30 ml). The combined ethyl acetate solutions are washed with saturated sodium chloride solution (5 ml), dried over magnesium sulfate and evaporated to dryness to yield 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, the compound of Example 12.

Example 26:

1-Carboxymethyl-7-chloro-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, isomer B

Chlorine is bubbled through a solution of 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepine-2-one (isomer B, 1.5 g) in acetic acid (25 ml), with stirring at room temperature. A white solid precipitates out; chlorine is bubbled through the reaction mixture until the reaction is complete. The solid is filtered off and separated by reverse phase HPLC using a C₁₈ column and methanol/0.1 % aqueous ammonium carbonate (1:1) as solvent. The appropriate fraction is dissolved in methanol/ethyl acetate (1:1, 50 ml) and hydrogen chloride bubbled through the solution. The solution is evaporated, the residue is suspended in ether (100 ml) and the suspension is filtered to give 1-carboxymethyl-7-chloro-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one hydrochloride, m.p. 149-151° (isomer B).

Example 27:

1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride

3(S)-Amino-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one sodium salt (619 g) having $[\alpha]_D^{25} = -304.4^\circ$ (c=1.08 in water), ethyl benzylpyruvate (1960 g), anhydrous ethyl alcohol (5880 ml) and glacial acetic acid (5880 ml) are combined and stirred at 20-25° for 1.5 hours. A solution of sodium cyanoborohydride (179 g) in anhydrous ethyl alcohol (2200 ml) is added at a constant slow rate over 24 hours. After addition is complete, the reaction mixture is stirred for 24 hours. 12N Hydrochloric acid (500 ml) is added to the reaction mixture and the solvent is evaporated at 35-40°/3 mm Hg. The oil which remains is combined with ice (3000 g) water (3000 ml) and diethyl ether (3000 ml), and the pH of the mixture is adjusted to 9-9.5 with 10N sodium hydroxide solution (1735 ml). The aqueous portion is removed and an additional 8000 ml of diethyl ether is added to the ether portion to oil out additional product. The ether immiscible portion is removed and combined with the aqueous portion. The ether extract is then washed with water (2 x 1000 ml), the washes are incorporated with the aqueous/oil portions from above and the mixture is adjusted to pH 4.25-4.35 with 12N hydrochloric acid (550-650 ml). The mixture is extracted with ethyl acetate (3x2000 ml), the combined ethyl acetate portions are washed with water (2000 ml) and dried with anhydrous magnesium sulfate (500 g). The drying agent is removed by filtration and the solvent is thoroughly removed by evaporation at 40°/3 mm Hg. The resulting oil is dissolved in ethyl acetate (4500 ml) and 28 % ethereal hydrogen chloride (309 g) was added with vigorous stirring. Diethyl ether (1500 ml) is added and the mixture is stirred for 1 hour. The solid is collected and is washed with ethyl acetate (2x500 ml) and diethyl ether (3x1000 ml). Drying at 50°/3 mm Hg affords crude product consisting of approximately 65 % of the desired 1-carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-

tetrahydro-1H-[1]benzazepin-2-one, identical to the material of Example 12, as determined by reverse phase HPLC on a C₁₈ column with a mixture of methanol, water, and acetic acid (75:25:0.02) as eluent.

Hydrogen chloride gas is added in a steady stream to a suspension of the above crude product in dichloromethane (26900 ml). A solution is obtained after 40 minutes when the addition of the gas is stopped. The solution is filtered to remove trace insolubles and diethyl ether (10750 ml) was added.

The suspension is stirred overnight at ambient temperature and the solid is collected by filtration and washed with dichloromethane (4 x 500 ml) and diethyl ether (3 x 1000 ml). Drying affords purer product as the hydrochloride salt, m.p. 175-178°.

1880 g of above hydrochloride salt is combined with dichloromethane (18000 ml). The suspension is again treated with hydrogen chloride gas to complete solution. Diethyl ether (7200 ml) is added. The suspension is stirred for 3 hours and filtered. The collected solid is washed with dichloromethane (2 x 1000 ml) and diethyl ether (2 x 1000 ml) and is dried to give product m.p. 183-185° (HPLC indicated that the product was approximately 96 % pure).

1280 g of the above salt is combined with chloroform (4000 ml) and the mixture is heated at reflux temperature for 10 minutes. Heating is discontinued and the mixture is stirred for 4 hours and filtered. The solid is washed with chloroform (2 x 200 ml) and diethyl ether (3 x 500 ml), dried and sieved to give 1-carboxymethyl-3S-(1S-ethoxy-carbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride, m.p. 184-186°, $[\alpha]_D^{25} = -139.26^\circ$ (c = 0.92, absolute ethanol), and identical to the hydrochloride salt of Example 12.

Example 28:

3-(1-benzyloxycarbonyl-3-phenylpropylamino)-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride (Isomer B)

Dry hydrogen chloride gas is bubbled through a solution of 3-(1-benzyloxycarbonyl-3-phenylpropylamino)-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (4.0 g, see Example 7) in ethyl acetate (100 ml) for 20 minutes while stirring at 0°. The reaction mixture is evaporated under reduced pressure and the resulting solid triturated with ether (50 ml). The solid is filtered off, washed with ether (15 ml) and ethyl acetate (15 ml), and then boiled with ethyl acetate (50 ml). The product is recrystallized from methanol/ethyl acetate to give the title compound. m.p. 197-199° (isomer B).

The starting material is prepared as follows: Potassium t-butoxide (1.2 g) is added to a solution of 3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (3.0 g) and t-butyl bromoacetate (2.2 g) in tetrahydrofuran (100 ml) stirring at room temperature under an atmosphere of dry nitrogen. The reaction mixture is stirred for 20 hours at room temperature, then poured into water (250 ml) and extracted with dichloromethane (2 x 150 ml). The combined dichloromethane solutions are washed with water (100 ml) and dried over magnesium sulfate. Evaporation of the solvent gives 3-(1-benzyloxycarbonyl-3-phenylpropylamino)-1-t-butyloxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

Example 29:

1-Ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

A solution of ethyl 2-(1-ethoxycarbonyl-3-phenylpropylamino)-4-[o-(ethoxycarbonylmethylamino)-phenyl]-butyrate (5.6 g) in methanol (100 ml) is added to a solution of sodium methoxide in methanol

[prepared from sodium (0.25 g) and methanol (50 ml)] with stirring under a nitrogen atmosphere. The reaction mixture is refluxed for 65 hours, then evaporated under reduced pressure. The residue is distributed between water (50 ml) and dichloromethane (200 ml). The aqueous solution is extracted with dichloromethane (200 ml) and the combined organic solutions washed with water (50 ml) and dried over potassium carbonate. Evaporation of the solvent gives as a mixture of isomers A and B of 1-ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, which is separated by chromatography on silica gel and converted to the individual maleate salts as described in Example 10.

The starting material is obtained as follows: To a solution of ethyl 2-amino-4-(o-nitrophenyl)-butyrate (17.4 g) in 50 % aqueous dioxane (130 ml) is added triethylamine (10.5 g) and 2-(tert-butyloxycarbonyloxyimino)-2-phenylacetonitrile (18.7 g). The reaction mixture is stirred at room temperature for 4 hours and then diluted with water (300 ml). The mixture is extracted with ether (2 x 150 ml), the aqueous phase acidified with ice-cold 2N hydrochloric acid and extracted with ethyl acetate (2 x 250 ml). The ethyl acetate layers are combined, washed with water (150 ml) and dried over sodium sulfate. The solvent is removed under reduced pressure to give ethyl 2-t-butyloxycarbonylamino-4-(o-nitrophenyl)-butyrate, used without further purification.

A solution of ethyl 2-t-butyloxycarbonylamino-4-(o-nitrophenyl)-butyrate (13.0 g) in ethanol (300 ml) is hydrogenated at room temperature and atmospheric pressure, using 10 % palladium on charcoal (1 g) as catalyst, until uptake ceases. The catalyst is filtered off. Evaporation of the solvent gives ethyl 2-t-butyloxycarbonylamino-4-(o-aminophenyl)-butyrate which is used without further purification for the next step.

A solution of ethyl 2-t-butyloxycarbonylamino-4-(o-aminophenyl)-butyrate (10.0 g) and ethyl glyoxylate (4.2 g) in ethanol (120 ml) is hydrogenated at 80° and 3 atmospheres pressure for 72 hours using 10 % palladium on charcoal (3 g) as catalyst. The reaction mixture is cooled to room temperature and the catalyst filtered off. The solvent is removed under reduced pressure and the residue distributed between ethyl acetate (150 ml) and water (75 ml). The organic phase is dried over sodium sulfate and the solvent removed under reduced pressure to give ethyl 2-t-butyloxycarbonylamino-4-[o-(ethoxycarbonylmethylamino)-phenyl]-butyrate which is used without further purification for the next step.

Hydrogen chloride gas is bubbled through a solution of ethyl 2-t-butyloxycarbonylamino-4-[o-(ethoxycarbonylmethylamino)-phenyl]-butyrate (8.5 g) in ethyl acetate (150 ml) for 30 minutes at room temperature. The solution is evaporated under reduced pressure and the residue dissolved in ethyl acetate (100 ml). The solution is washed with water (3 x 100 ml) and dried over sodium sulfate. The solvent is removed under reduced pressure to give ethyl 2-amino-4-[o-(ethoxycarbonylmethylamino)-phenyl]-butyrate used without further purification for the next step.

A solution of ethyl 2-amino-4-[o-(ethoxycarbonylmethylamino)-phenyl]-butyrate (4.7 g) and ethyl benzylpyruvate (12.4 g) in acetic acid (35 ml) and methanol (35 ml) is stirred at room temperature under nitrogen for 1 hour. Sodium cyanoborohydride (1.6 g) in methanol (15 ml) is added dropwise over 4 hours. The reaction mixture is stirred at room temperature for 24 hours. Concentrated hydrochloric acid (2 ml) is added dropwise, and the mixture stirred at room temperature for 1 hour. The reaction mixture is evaporated to dryness, and the residue partitioned between water (75 ml) and ether (75 ml) and adjusted to pH 2 with 6N hydrochloric acid. The layers are separated,

and the aqueous phase extracted with ether (2 x 75 ml). The ether extracts are discarded and the aqueous layer adjusted to pH 9 with 40 % sodium hydroxide, and extracted with ethyl acetate (3 x 50 ml). The ethyl acetate extracts are dried over sodium sulfate and the solvent removed under reduced pressure to give ethyl 2-(1-ethoxy-carbonyl-3-phenylpropylamino)-4-(o-ethoxycarbonylmethylamino)-phenyl]butyrate which is used directly for preparing the final product above.

Example 30:

Ethyl 2-amino-4-phenylbutyrate is treated under conditions of reductive alkylation as described in the previous Examples with 1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,3-dione to give 1-ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropyl-amino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one of Example 10.

The starting material is prepared as follows: A solution of 3,3-dichloro-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (1.0 g, 4.32 mmol) and ethyl bromoacetate (0.51 ml) in tetrahydrofuran (30 ml) is added dropwise with stirring during 15 minutes to a solution of sodium hydride (4.76 mmol) in tetrahydrofuran (20 ml) at room temperature under a nitrogen atmosphere. Stirring is continued for an additional 2 hours. The solution is quenched by addition of saturated aqueous ammonium chloride and the solvents are removed under reduced pressure. The residue is extracted with ether (3 x 20 ml), the combined ether solutions washed with saturated aqueous sodium chloride solution (20 ml) and dried over magnesium sulfate. Removal of the solvent under reduced pressure gives 3,3-dichloro-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one. NMR(CDC₃): δ 1.27 (t, 3H); 3.22 (m, 4H); 4.25 (q, 2H); 4.65 (s, 2H) and 7.3 (m, 4H).

A mixture of morpholine (0.315 ml, 3.6 mmol) and 3,3-dichloro-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (0.5 g) is stirred under nitrogen at 110° for 18 hours. The solution is diluted to 10 ml with chloroform and cooled to 0°. 20 % sulfuric acid (1 ml) is added and the solution stirred for 2 hours at 0°. The solution is extracted with chloroform (2 x 20 ml) and the extracts are washed with 2 N hydrochloric acid (2 x 10 ml) and saturated aqueous sodium chloride solution (5 ml). The solution is dried over magnesium sulfate and evaporated under reduced pressure to yield 1-ethoxy- carbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2,3-dione. NMR (CDCl₃): δ 1.25 (t, 3H); 2.6 (m, 2H); 3.6 (m, 2H); 4.2 (q, 2H) and 7.3 (m, 4H).

Example 31:

Ethyl 2-amino-4-phenylbutyrate is treated in the presence of potassium carbonate in methylene chloride with 3-bromo-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one to give 1-ethoxy-carbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H[1]benzazepin-2-one of Example 10.

The starting material is prepared as follows: To a solution of 2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (2.5 g) in chloroform (30 ml), phosphorous pentachloride (3.2 g) is added in portions, while maintaining the temperature at 0-5°. When the addition is complete, iodine (30 mg) is added followed by bromine (2.5 g), which is added dropwise over 5 minutes. The mixture is then refluxed for 4 hours. The chloroform solution is evaporated and the residue is partitioned between ice-water (30 ml) and dichloromethane (75 ml). The organic phase is dried over magnesium sulfate and evaporated under reduced pressure. The crude residue is purified by chromatography over silica gel, eluting with ether and hexane (7:3). Concentration of the appropriate fractions yields 3-bromo-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one, m.p. 146-148°.

3-Bromo-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (300 mg) is added in one portion to a stirred suspension of potassium hydroxide (90 mg) and tetrabutylammonium bromide (40 mg) in tetrahydrofuran (10 ml) maintained at 0° under a nitrogen atmosphere. Stirring is continued for 5 minutes, then ethyl bromoacetate (200 mg) is added in one portion. The reaction mixture is allowed to warm to room temperature while stirring for an additional 3 hours. The tetrahydrofuran is removed under reduced pressure and the residue partitioned between water (5 ml) and ether (25 ml). The organic phase is washed with 2N hydrochloric acid (5 ml), dried over magnesium sulfate, and the solvent removed under reduced pressure to give 3-bromo-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, m.p. 114-116°.

3-Chloro-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is similarly prepared.

A solution of 3-chloro-2,3,4,5-tetrahydro-1H-[1]benzazepine-2-one (1.95 g) in dimethylformamide (10 ml) is added dropwise with stirring to a solution of potassium t-butoxide (1.12 g) in dimethylformamide (10 ml) at 5°. The solution is stirred for an additional 15 minutes at 5°, then ethyl bromoacetate (1.78 g) in dimethylformamide (5 ml) is added dropwise. Stirring is continued for an additional 30 minutes at 5° and then for 3 hours at room temperature. The reaction mixture is cooled to 10° and water (100 ml) is added. The solution is extracted with chloroform (100 ml) and the chloroform solution washed with water (2 x 10 ml) and dried over sodium sulfate. The solvent is removed under reduced pressure to yield 3-chloro-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one; NMR (DMSO-d₆): δ1.2 (t, 3H); 2.65 (m, 4H); 4.15 (q, 2H); 2.6 (d, 2H) and 7.3 (m).

Example 32:

1-carboxymethyl-3S-(1S-pivaloyloxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

1-Benzylloxycarbonylmethyl-3S-(1S-pivaloyloxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (3 g) is dissolved in ethanol (50 ml) and 10 % Pd-C (0.3 g) is added and the solution hydrogenated at 1 atmosphere pressure and room temperature for 2 hours. The reaction mixture is filtered and evaporated to yield 1-carboxymethyl-3S-(1S-pivaloyloxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

The starting material is prepared as follows: 1-benzylloxycarbonylmethyl-3S-(1S-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one (5 g, Example 2) is dissolved in 2N potassium hydroxide solution (5.15 ml) and the solution evaporated to dryness. Iodomethyl pivalate (2.3 g) and dimethylformamide (50 ml) are added, and the reaction mixture is stirred at room temperature for 18 hours under a nitrogen atmosphere. The dimethylformamide is evaporated, the residue is taken up in ethyl acetate (100 ml) and washed with saturated sodium bicarbonate (3 x 25 ml), water (3 x 25 ml), and saturated sodium chloride (25 ml), and dried over magnesium sulfate. Evaporation gives 1-benzylloxycarbonylmethyl-3S-(1S-pivaloyloxy-methoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]-benzazepin-2-one.

Similarly prepared are:

- a) 1-carboxymethyl-3S-(1S-*l*-bornyloxycarbonylmethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one using *l*-bornyl iodoacetate as starting material.

b) 1-carboxymethyl-3S-(1S- β -methoxyethoxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one using β -methoxyethoxymethyl chloride as starting material.

c) 1-carboxymethyl-3S-[1S-(3-phthalidoxycarbonyl)-3-phenylpropylamino]-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one using 3-bromophthalide as starting material.

d) 1-carboxymethyl-3S-[1S-(3-pyridylmethoxycarbonyl)-3-phenylpropylamino]-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one using 3-pyridylmethyl chloride as starting material.

Example 33:

1-Carboxymethyl-3S-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5,5a,-6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one

A solution of 3(S)-amino-1-carboxymethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one sodium salt (0.6 g) and ethyl benzylpyruvate (1.5 g) in acetic acid (5 ml) and methanol (3 ml) is stirred at room temperature under an atmosphere of dry nitrogen for 1 hour. A solution of sodium cyanoborohydride (0.2 g) in methanol (2 ml) is then added over a 4 hour period. The reaction mixture is stirred at room temperature for 18 hours. Concentrated hydrochloric acid (0.5 ml) is added and the mixture stirred at room temperature for 1 hour. The solvents are removed under reduced pressure and the residue partitioned between water (20 ml) and ether (20 ml). The pH is adjusted to 9.3 with 40 % sodium hydroxide. The layers are separated and the ether layer discarded. The aqueous phase is adjusted to pH 4.3 with concentrated hydrochloric acid and extracted with ethyl acetate (3 x 25 ml). The extracts are dried over magnesium sulfate and the solvent removed under reduced pressure. Hydrogen chloride is bubbled into a solution of the residue in dichloromethane (70 ml) for

5 minutes. The solution is evaporated and the residue recrystallized from ethanol/ether to give 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one-hydrochloride as a mixture of isomers.

The starting material is prepared as follows: A solution of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5-tetrahydro-1H-[1]benzazepine-2,5-dione (3.6 g) in acetic acid (50 ml) is hydrogenated for 120 hours at 3 atmospheres pressure using platinum oxide (1.2 g) as catalyst. The catalyst is filtered off and the filtrate evaporated under reduced pressure. The residue is distributed between dichloromethane (200 ml) and saturated aqueous sodium bicarbonate (100 ml). The dichloromethane solution is washed with water (50 ml), dried over sodium sulfate and the solvent removed under reduced pressure. The residue is chromatographed on silica gel eluting with 0-50 % ethyl acetate in toluene. The fraction eluting with 50 % ethyl acetate in toluene is collected to give 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2,5-dione used without further purification for the next synthetic step.

A solution of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2,5-dione (2.7 g) and sodium borohydride (0.2 g) in ethanol (100 ml) is stirred at room temperature for 18 hours. The solvent is removed under reduced pressure, and the residue dissolved in dichloromethane (100 ml). The solution is extracted with ice-cold 2N hydrochloric acid (2 x 50 ml) and saturated aqueous sodium chloride solution (50 ml) and dried over sodium sulfate. The solvent is removed under reduced pressure and the residue triturated with ether to give 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-5-hydroxy-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one.

A mixture of 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-5-hydroxy-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one (2.1 g), dicyclohexylcarbodiimide (1.8 g) and cuprous chloride (0.2 g) is heated at 80° under nitrogen for 32 hours. The reaction mixture is cooled to room temperature, the residue is dissolved in methylene chloride (200 ml), washed with dilute ammonium hydroxide (2 x 50 ml) and water (50 ml). The organic phase is dried over sodium sulfate and evaporated to give a mixture of the desired adduct and excess dicyclohexylcarbodiimide. This mixture is dissolved in ethyl acetate (100 ml) and placed in a pressure bottle. 10 % Pd/C (0.4 g) is added and the mixture hydrogenated at 3 atmospheres pressure and 40° for 16 hours. The catalyst is filtered off and the filtrate is evaporated to give 3(S)-t-butyloxycarbonylamino-1-ethoxycarbonylmethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one, used without further purification for the next synthetic step.

Hydrogen chloride gas is bubbled through a solution of the above compound (1.1 g) in ethyl acetate (50 ml) for 45 minutes. The reaction mixture is evaporated under reduced pressure, the residue dissolved in ethyl acetate (50 ml) and washed with water (3 x 30 ml). The ethyl acetate solution is dried over sodium sulfate and the solvent removed under reduced pressure to give 3(S)-amino-1-ethoxy-carbonylmethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one, which is used without further purification for the next step.

A solution of sodium hydroxide (0.1g) in water (0.25 ml) is added to a solution of the above amine (0.6 g) in methanol (7.5 ml) at room temperature, and the solution is stirred for 2 hours. The solvents are evaporated and the residue thoroughly dried, then slurried with ether, to give the sodium salt of 3(S)-amino-1-carboxymethyl-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one.

Example 34:

N-[1-(1-carboxymethyl)-2,3,4,5-tetrahydro-2-oxo-1H-[1]benzazepin-3S-ylamino)-3-phenylpropyl-1-carbonyl]-L-phenylalanine.

L-Phenylalanine methyl ester hydrochloride is condensed with 1-benzyl-oxycarbonylmethyl-3S-(1S-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one in methylene chloride in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride at room temperature to yield after workup the N-[1-(1-benzyloxycarbonylmethyl-2,3,4,5-tetrahydro-2-oxo-1H-[1]benzazepin-3S-ylamino)-3-phenylpropyl-1-carbonyl]-L-phenylalanine methyl ester.

Hydrogenation using 10 % Pd/C catalyst in ethanol gives N-[1-(1-carboxymethyl-2,3,4,5-tetrahydro-2-oxo-1H-[1]benzazepin-3S-ylamino)-3-phenylpropyl-1-carbonyl]-L-phenylalanine methyl ester.

Hydrolysis with dilute aqueous sodium hydroxide at room temperature for 18 hours yields the N-[1-(1-carboxymethyl-2,3,4,5-tetrahydro-2-oxo-1H-[1]benzazepin-3S-ylamino)-3-phenylpropyl-1-carbonyl]-L-phenylalanine.

Example 35:

1-Ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one

Treatment of 3-(1-carboxy-3-phenylpropylamino)-1-cyanomethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one with ethanol-ether (1:1) saturated with hydrogen chloride at room temperature for 48 hours gives after workup 1-ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one identical to the compound of Example 10.

The starting material is prepared as follows: 3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one is alkylated with bromoacetonitrile in dimethylformamide solution in the presence of sodium hydride to yield after work-up, 3-(1-carboxy-3-phenylpropylamino)-1-cyanomethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one, used directly in the next step.

Example 36:

Preparation of 10,000 tablets each containing 10 mg of the active ingredient of Example 12:

Formula:

1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one	100 g
Lactose	1,157 g
Corn starch	75 g
Polyethylene glycol 6,000	75 g
Talcum powder	75 g
Magnesium stearate	18 g
Purified water	q.s

Procedure: As described in Example 15.

Example 37:

Preparation of 10,000 capsules each containing 20 mg of the hydrochloride salt of the active ingredient of Example 12.

Formula:

1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one hydrochloride	200 g
Lactose	1,700 g
Talcum powder	100 g

Procedure: As described in Example 17.

Cardiovascular pharmacology of compounds of the invention

Testing of compounds is carried out by methods for evaluation of the inhibition of the angiotensin converting enzyme (ACE). Biochemical assessment of in vitro ACE inhibition (ACEI) gauges the inhibition of peptidolytic activity of a compound in rabbit lung tissue. In in vivo studies angiotensin I (AI) pressor response inhibition of the compounds are conducted in rats.

In the in vivo test method an increase in the blood pressure is first caused by administration of angiotensin I (AI) to the test animal. The inhibitory action of the individual compounds on this increase in blood pressure is then determined.

Biochemical Testing Methodology

A rabbit lung tissue preparation [Das and Saffer, J. Biol. Chem. 250: 6762, (1975)] was used for assessment of ACE by the method of Cheung and Cushman [Cheung and Cushman, Biochim. Biophys. Acta 293: 451, (1973)]. This test system incorporates spectrophotometric evaluation of the amount of histidyl-leucine liberated from a synthetic substrate after 30 min. of 37°C incubation. IC₅₀ values for ACE inhibition were determined graphically as the concentration of test drug required to reduce the amount of histidyl-leucine formed to 50% of that generated in the absence of the test compound.

Methodology of angiotensin I (AI) pressor response inhibition following intravenous administration of test compounds (% AI)


In these studies catheters were placed in a femoral artery and a saphenous vein of anesthetized rats as described above. Arterial pressure was continuously recorded from the arterial catheter, while AI and the test compounds were injected through the venous catheter. AI pressor response inhibition was expressed as percent decrease of the response from pretreatment control values and tabulated as the average inhibition recorded within 30 minutes after test drug administration.

Results:


Compound of Example	in vitro ACEI IC ₅₀ (M)	Angiotensin I pressor response inhibition in rats	
		i.v. Dose (mg/kg)	% AI inhibition
1	6×10^{-7}	10	100
		1.0	100
5	2×10^{-7}	0.1	50
		0.1	37
9	5×10^{-9}	0.3	93
		0.1	80
10	1×10^{-5}	0.03	40
		1.0	80
Maleate Salt			
12	4×10^{-7}	1.0	100
		0.3	95
HCl Salt		0.1	82
		0.06	74
19	2×10^{-9}	0.03	29
		0.1	93
28	1×10^{-7}	0.06	84
		0.03	70
Isomer B, HCl Salt		0.02	69
		0.01	28
		0.007	14
		0.1	92

Claims:

1. A compound of the general formula I



wherein R_A and R_B are radicals of the formula




in which R_o is carboxy or a functionally modified carboxy; R_1 is hydrogen, lower alkyl, amino(lower) alkyl, aryl, aryl (lower) alkyl, cycloalkyl or cycloalkyl (lower) alkyl; R_2 is hydrogen or lower alkyl; R_3 and R_4 , each independently, represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxy, halogen, trifluoromethyl, or R_3 and R_4 taken together represent lower alkylene-dioxy; R_5 is hydrogen or lower alkyl, and X represents oxo, two hydrogens, or one hydroxy together with one hydrogen; and wherein the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro; complexes thereof; and stereoisomers of all these compounds.

2. A compound of claim 1 wherein one or both of R_o represented by COR_6 in radical R_A and represented by COR_7 in radical R_B independently represent carboxy, esterified carboxy, carbamoyl or substituted carbamoyl.

3. A compound of claim 2 wherein one or both of R_6 and R_7 represent hydroxy; lower alkoxy; (amino, mono- or di-lower alkylamino)-substituted lower alkoxy; carboxy-substituted lower alkoxy; lower alkoxycarbonyl-substituted lower alkoxy; aryl-substituted lower alkoxy; (hydroxy, lower alkanoyloxy or lower alkoxy)-substituted lower alkoxy; (hydroxy, lower alkanoyloxy or lower alkoxy)-substituted lower alkoxymethoxy; bicycloalkoxycarbonyl-substituted lower alkoxy; 3-phthalidoxy; (lower alkyl, lower alkoxy, halo)-substituted 3-phthalidoxy; amino; lower alkylamino; di-lower alkylamino; di-lower alkylamino in which both alkyl groups are linked by a carbon to carbon bond and together with the amino nitrogen form a 5, 6- or 7-membered heterocyclic ring; (amino or acylamino)-substituted lower alkylamino; α -(carboxy or lower alkoxycarbonyl)-substituted lower alkylamino; aryl-substituted lower alkylamino which can be substituted on the α -carbon by carboxy or lower alkoxycarbonyl.

4. A compound of the formula IA



wherein R_1 is hydrogen, lower alkyl, amino(lower)alkyl, aryl, aryl-(lower)alkyl, cycloalkyl(lower)alkyl, R_2 and R_5 represent hydrogen or lower alkyl, R_3 and R_4 represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxy, halogen, trifluormethyl; or R_3 and R_4 taken together represent lower alkylendioxy, X represents oxo, two hydrogens or one hydroxy group and one hydrogen, R_6 and R_7 independently represent hydroxy, amino, mono- or di-(lower)alkylamino, lower alkoxy, aryl(lower)alkoxy, lower alkanoyloxymethoxy, (amino, mono- or di-lower alkylamino, carboxy, or lower alkoxycarbonyl)-lower alkoxy; or complexes thereof.

5. A compound of the formula IA shown in claim 4, wherein R_1 is hydrogen, lower alkyl, amino(lower)alkyl, aryl(lower)alkyl where aryl represents phenyl unsubstituted or mono- or disubstituted by lower alkyl, hydroxy, lower alkoxy, lower alkylenedioxy, lower alkanoyloxy, halogen or trifluoromethyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 and R_4 are hydrogen, lower alkoxy, lower alkyl, halogen or trifluoromethyl; or R_3 and R_4 taken together represent alkylenedioxy, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower)alkoxy, lower alkoxy carbonyl(lower)alkoxy.


6. A compound of the formula IA shown in claim 4, wherein R_1 is hydrogen, lower alkyl, ω -amino(lower)alkyl, aryl(lower)alkyl where aryl represents phenyl unsubstituted or mono-substituted by lower alkyl, hydroxy; lower alkoxy, lower alkanoyloxy, halogen or trifluoromethyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 and R_4 are hydrogen, lower alkoxy, lower alkyl, halogen, or trifluoromethyl; or R_3 and R_4 taken together represent lower alkylendioxy, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower)alkoxy, lower alkoxy carbonyl(lower)alkoxy.

7. A compound of the formula IA shown in claim 4, wherein R_1 is hydrogen, lower alkyl, ω -amino(lower)alkyl, aryl(lower)alkyl, R_2 and R_5 are hydrogen or lower alkyl, R_3 is hydrogen, R_4 is hydrogen, lower alkoxy, lower alkyl, halogen, or trifluoromethyl, X represents oxo, one hydroxy and one hydrogen, or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, lower alkoxy, phenyl(lower)alkoxy, lower alkoxy carbonyl(lower)alkoxy.

8. A compound of the formula IA shown in Claim 4, wherein R_1 is hydrogen, methyl, ethyl, isopropyl, ω -aminopropyl, ω -aminobutyl, aryl(methyl, ethyl, propyl) where aryl represents phenyl unsubstituted or substituted by one methyl, hydroxy, methoxy, methylenedioxy,

acetyloxy, chloro or trifluoromethyl group, R_2 and R_5 are hydrogen or methyl, R_3 and R_4 represents hydrogen, methoxy, methyl, chloro or trifluoromethyl, X represents oxo, one hydroxy and one hydrogen or 2 hydrogens, R_6 and R_7 independently represent hydroxy, amino, ethoxy, methoxy, benzyloxy, ethoxycarbonylmethoxy or pivaloyloxy-methoxy.

9. A compound of the formula IB




wherein n represents an integer from 1 to 4, R_8 is hydrogen, phenyl unsubstituted or monosubstituted by lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, hydroxy, or trifluoromethyl, R_6 and R_7 independently represent hydroxy, lower alkoxy of up to 4 carbon atoms, benzyloxy, or amino.

10. A compound of the formula IB shown in claim 9, wherein C_nH_{2n} represents ethylene, R_8 represents phenyl or phenyl mono-substituted by lower alkoxy with up to 4 carbon atoms, lower alkyl with up to 4 carbon atoms, halogen or trifluoromethyl, R_6 and R_7 independently represent hydroxy or lower alkoxy with up to 4 carbon atoms.

11. A compound of the formula IA shown in claim 4, wherein $X = H_2$, R_2 and $R_5 = H$, $R_6 = OC_2H_5$ and $R_7 = OH$ and the other symbols have the following meanings:

No.	<u>R₁</u>	<u>R₃</u>	<u>R₄</u>
1	C ₆ H ₅ CH ₂	H	H
2	C ₆ H ₅ CH ₂ CH ₂	7-OCH ₃	8-OCH ₃
3	C ₆ H ₅ CH ₂ CH ₂	7-Cl	H
4	C ₆ H ₅ CH ₂ CH ₂	8-CH ₃	H
5	C ₆ H ₅ CH ₂ CH ₂	8-OCH ₃	H
6	P-CIC ₆ H ₄ CH ₂ CH ₂	H	H
7	CH ₃	H	H

12. A compound of the formula IC



wherein S represents the chirality, n represents an integer from 1 to 4, R₈ is hydrogen, phenyl unsubstituted or monosubstituted by lower alkyl, lower alkoxy, lower alkanoyloxy, halogen, hydroxy, or trifluoromethyl, R₆ and R₇ independently represent hydroxy, lower alkoxy of up to 4 carbon atoms, benzyloxy or amino, or pharmaceutically acceptable salts thereof.

13. 1-Carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one or stereoisomers thereof.

14. The higher melting racemic compound of the compound of claim 13 and an enantiomer thereof.

15. The lower melting racemic compound of the compound claimed in claim 13 and an enantiomer thereof.

16. 1-Carboxymethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

17. 1-Benzylloxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

18. 1-Benzylloxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

19. 1-Benzylloxycarbonylmethyl-3-(1-benzylloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

20. 1-Carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one or stereoisomers thereof.

21. The lower melting racemic compound of the compound claimed in claim 20 or an enantiomer thereof.

22. The higher melting racemic compound of the compound claimed in claim 20 or an enantiomer thereof.

23. The racemic isomer B of 1-ethoxycarbonylmethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

24. 1-Carboxymethyl-3-carboxymethylamino-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

25. 1-Carboxymethyl-3-(1-carboxy-3-phenylpropylamino)-5-hydroxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

26. 1-Carboxymethyl-3S-(1R-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
27. 1-Carboxymethyl-3S-(1S-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
28. 1-Ethoxycarbonylmethyl-3-(1-benzyloxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
29. The racemic isomer A of the compound claimed in claim 28.
30. The racemic isomer B of the compound claimed in claim 28.
31. 1-Ethoxycarbonylmethyl-3-(1-carboxy-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one or stereoisomers thereof.
32. The racemic isomer compound B of the compound claimed in claim 31.
33. The racemic isomer compound of 1-carboxymethyl-3-(1-ethoxycarbonyl-3-phenylpropylamino)-8-methoxy-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
34. 1-(1-Carboxyethyl)-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
35. 1-Ethoxycarbonylmethyl-3S-(1S-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
36. 1-Ethoxycarbonylmethyl-3S-(1R-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.
37. The racemic isomer compound B of 1-carboxymethyl-7-chloro-3-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

38. 3-(1-Benzylloxycarbonyl-3-phenylpropylamino)-1-carboxymethyl-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one or stereoisomers thereof.

39. 1-Carboxymethyl-3S-(1S-pivaloyloxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

40. 1-Carboxymethyl-3S-(1S-*β*-bornyloxycarbonylmethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

41. 1-Carboxymethyl-3S-(1S- β -methoxyethoxymethoxycarbonyl-3-phenylpropylamino)-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

42. 1-Carboxymethyl-3S-[1S-(3-phthalidoxycarbonyl)-3-phenylpropylamino]-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

43. 1-Carboxymethyl-3S-[1S-(3-pyridylmethoxycarbonyl)-3-phenylpropylamino]-2,3,4,5-tetrahydro-1H-[1]benzazepin-2-one.

44. 1-Carboxymethyl-3S-(1-ethoxycarbonyl-3-phenylpropylamino)-2,3,4,5,5a,6,7,8,9,9a-decahydro-1H-[1]benzazepin-2-one.

45. N-[1-(1-carboxymethyl)-2,3,4,5-tetrahydro-2-oxo-1H-[1]benzazepin-3S-ylamino)-3-phenylpropyl-1-carbonyl]-L-phenylalanine.

46. A salt of a compound having a salt forming group, as claimed in anyone of claims 4, 6-9 and 11-25.

47. A salt of a compound having a salt forming group, as claimed in anyone of claims 5, 10 and 26-32.

48. A salt of a compound having a salt forming group, as claimed in anyone of claims 1-3 and 33-45.

49. A pharmaceutically acceptable salt of a compound having a salt forming group, as claimed in any one of claims 4, 6-9 and 11-25.

50. A pharmaceutically acceptable salt of a compound having a salt forming group, as claimed in any one of claims 5, 10 and 26-32.

51. A pharmaceutically acceptable salt of a compound having a salt forming group, as claimed in any one of claims 1-3 and 33-45.

52. A compound of formula I substantially as described with reference to any of Examples 1 to 35.


53. A pharmaceutical preparation comprising a compound claimed in any one of claims 4, 6-9, 11-25 and 49, in admixture or conjunction with a pharmaceutically suitable carrier.

54. A pharmaceutical preparation comprising a compound claimed in any one of claims 5, 10, 26-32 and 50 in admixture or conjunction with a pharmaceutically suitable carrier.


55. A pharmaceutical preparation comprising a compound claimed in any one of claims 1-3, 33-45 and 51 in admixture or conjunction with a pharmaceutically suitable carrier.

56. A pharmaceutical preparation according to claim 53, 54 or 55 substantially as described with reference to Examples 36 or 37.

57. Process for the manufacture of 3-amino-[1]benzazepin-2-one-1-alkanoic acids of the general formula I




wherein R_A and R_B are radicals of the formula



, respectively,

in which R_0 is carboxy or a functionally modified carboxy; R_1 is hydrogen, lower alkyl, amino(lower) alkyl, aryl, aryl (lower) alkyl, cycloalkyl or cycloalkyl (lower) alkyl; R_2 is hydrogen or lower alkyl; R_3 and R_4 , each independently, represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxy, halogen, trifluoromethyl, or R_3 and R_4 taken together represent lower alkylene-dioxy; R_5 is hydrogen or lower alkyl, and X represents oxo, two hydrogens, or one hydroxy together with one hydrogen; and wherein the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro; salts, complexes and stereoisomers of all these compounds, which consists in that

a) in a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein X , R_B , R_3 , R_4 and R_5 have the meanings given hereinbefore, R_A is introduced by alkylation with a compound of the formula




wherein Z is a reactive esterified hydroxyl group and R_A has the meanings given hereinbefore, or with a compound of the formula



wherein R_1 and R_0 have the meanings given hereinabove, in the presence of a reducing agent, with a temporary protection of any primary and secondary amino groups and/or, optionally, hydroxyl and/or oxo groups, which may be present in any one of the substituents X , R_A , R_B , R_1 , R_3 , R_4 and R_5 or

b) a compound of the formula




in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein X , R_3 , R_4 and R_5 have the meanings given hereinabove and R' is hydrogen or R_A as defined hereinabove, is alkylated with a compound of the formula



wherein Z is a reactive esterified hydroxyl group and R_B has the meanings given hereinabove, while protecting temporarily any primary and secondary amino groups and/or, optionally, hydroxyl and/or oxo groups which may be present in any one of the residues X , R_A , R_B , R_3 , R_4 and R_5 , or

c) a compound of the formula




in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro and wherein Y is oxo or a reactive esterified hydroxyl group Z together with hydrogen, and X, R₃, R₄ and R₅ have the meanings given hereinabove, is condensed with an amine of the formula




wherein R_A and R₅ have the meanings given hereinabove, with the proviso that in the case Y is oxo, the condensation is carried out in the presence of a reducing agent and with a temporary protection of the oxo group which may be present as the substituent X, or

d) in a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein X and R₁ to R₅ have the meanings given hereinabove, one of the symbols R'_o and R''_o is cyano and the other one is cyano or R_o as defined hereinabove, the cyano group(s) is (are) subjected to solvolysis, or


e) a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro and wherein X, R_A, R_B, R₃, R₄ and R₅ have the meanings given hereinabove, or an ester thereof, is cyclised, or

f) a compound which is structurally identical with a compound of formula I specified above, except for having an additional double bond located at C-3, or between the nitrogen atom and the adjacent carbon atom within the group R_A , is treated with a reducing agent in order to saturate this double bond, or

g) in order to produce a compound of formula I as specified hereinabove, in which X is oxo, condensing a compound of the formula



in which the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro, and wherein R_B , R_3 and R_4 have the meanings given hereinabove, with an amine of the formula



wherein R_A and R_5 have the meaning given hereinabove, and

- h) if desired, a resulting compound of formula I as specified above is converted into another compound of formula I within its above-specified scope, and/or
- i) if desired, a resulting compound of formula I as specified above and having salt-forming properties is converted into a salt thereof or a free compound is liberated from such a salt, and/or
- j) if desired, a resulting compound of formula I as specified above and having complex-forming properties is converted into a complex thereof, and/or
- k) if so required, an optical isomer which has a specific configuration with respect to at least one center of chirality is enriched from a mixture of stereoisomeric forms of a resulting compound of formula I.

58. A process for the preparation of compounds of formula I substantially as described in any one of Examples 1 to 14.

59. A process for the preparation of compounds of formula I substantially as described in any one of Examples 18 to 21.

60. A process for the preparation of compounds of formula I substantially as described in any one of Examples 22 to 35.

61. A compound of formula I when prepared according to a process of claim 57 or 58.

62. A compound of formula I when prepared according to a process of claim 57 or 59.

63. A compound of formula I when prepared according to a process of claim 57 or 60.

64. A compound of formula I for use in a therapeutic method of treating diseases responsive to inhibition of angiotensin-converting enzyme in humans and mammals.

2103614

Dated:	Application No.:	Published:
2103614 5 August 1982	8222564	23 February 1983
Priority:	United States of America	291,907
	United States of America	319,863
11 August 1981		
9 November 1981		

CIBA-GEIGY AG, (Basle-City), Klybeckstrasse 141, 4002 Basle, Switzerland,

JEFFREY WILLIAM HERBERT WATTHEY, 61 Deepwood Drive, Chappaqua, New York 10514, United States of America,

Benzazepin-2-ones:

Address for Service:

Kenneth D Sparrow, Ciba-Geigy PLC, Patent Department, Tenax Road, Trafford Park, Manchester M17 1WT.

Request for examination:	5/11/1982
Application refused	
or withdrawn:	
Patent granted:	21 NOV 1984
Renewal Fee paid in respect of	
5th Year	
6th Year	
7th Year	
8th Year	
9th Year	
10th Year	
11th Year	
12th Year	
13th Year	
14th Year	
15th Year	
16th Year	
17th Year	
18th Year	
19th Year	
20th Year	
Patent ceased or	
expired:	