



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**


(86) Date de dépôt PCT/PCT Filing Date: 2019/04/18
(87) Date publication PCT/PCT Publication Date: 2019/10/24
(85) Entrée phase nationale/National Entry: 2020/10/15
(86) N° demande PCT/PCT Application No.: US 2019/028207
(87) N° publication PCT/PCT Publication No.: 2019/204666
(30) Priorités/Priorities: 2018/04/18 (US62/659,676);
2019/03/20 (US62/821,084)

(51) Cl.Int./Int.Cl. *A61K 47/06* (2006.01),
A61K 47/14 (2017.01), *A61K 47/18* (2017.01),
A61K 47/24 (2006.01), *A61K 47/28* (2006.01),
A61K 9/14 (2006.01)
(71) Demandeur/Applicant:
OISIN BIOTECHNOLOGIES, INC., US
(72) Inventeur/Inventor:
SCHOLZ, MATTHEW REIN, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : NANOPARTICULES LIPIDIQUES FUSOGENES ET LEURS PROCEDES DE FABRICATION ET LEURS METHODES D'UTILISATION POUR LA PRODUCTION SPECIFIQUE A UNE CELLULE CIBLE D'UNE PROTEINE THERAPEUTIQUE ET POUR LE TRAITEMENT D'UNE MALADIE, D'UNE AFFECTION OU D'UN TROUBLE ASSOCIE A UNE CELLULE CIBLE

(54) Title: FUSOGENIC LIPID NANOPARTICLES AND METHODS FOR THE MANUFACTURE AND USE THEREOF FOR THE TARGET CELL-SPECIFIC PRODUCTION OF A THERAPEUTIC PROTEIN AND FOR THE TREATMENT OF A DISEASE, CONDITION, OR DISORDER ASSOCIATED WITH A TARGET CELL

FIG. 1



(57) **Abrégé/Abstract:**

Provided nucleic acid- based expression construct for the target cell-specific production of a therapeutic protein, such as a pro-apoptotic protein, within a target cell, including a target cell that is associated with aging, disease, or other condition, in particular a target cell that is a senescent cell or a cancer cell. Also provided are formulations and systems, including fusogenic lipid nanoparticle (LNP) formulations and systems, for the delivery of nucleic acid-based expression constructs as well as methods for making and using such nucleic acid-based expression constructs, formulations, and systems for reducing, preventing, and/or eliminating the growth and/or survival of a cell, such as a senescent cell and/or a cancer cell, which is associated with aging, disease, or other condition as well as methods for the treatment of aging, disease, or other conditions by the in vivo administration of a formulation, such as a fusogenic LPN formulation, comprising an expression construct for the target cell- specific, production of a therapeutic protein, such as a pro-apoptotic protein, in a target cell that is associated with aging, disease, or other condition, in particular a target cell that is a senescent cell or a cancer cell.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
24 October 2019 (24.10.2019)



(10) International Publication Number
WO 2019/204666 A1

(51) International Patent Classification:

A61K 31/7088 (2006.01) *A61K 47/18* (2017.01)
A61K 31/7105 (2006.01) *A61K 47/24* (2006.01)
A61K 31/713 (2006.01)

(21) International Application Number:

PCT/US2019/028207

(22) International Filing Date:

18 April 2019 (18.04.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/659,676	18 April 2018 (18.04.2018)	US
62/821,084	20 March 2019 (20.03.2019)	US

(71) Applicant: **OISIN BIOTECHNOLOGIES, INC.**
[US/US]; 454 N. 34th Street, Seattle, WA 98103 (US).


(72) Inventor: **SCHOLZ, Matthew, Rein**; 454 N. 34th Street, Seattle, WA 98103 (US).

(74) Agent: **MYLES, Gary, M.**; Myles Intellectual Property Law, P.O. Box 2049, Issaquah, WA 98027 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

(54) Title: FUSOGENIC LIPID NANOPARTICLES AND METHODS FOR THE MANUFACTURE AND USE THEREOF FOR THE TARGET CELL-SPECIFIC PRODUCTION OF A THERAPEUTIC PROTEIN AND FOR THE TREATMENT OF A DISEASE, CONDITION, OR DISORDER ASSOCIATED WITH A TARGET CELL

FIG. 1



(57) Abstract: Provided nucleic acid-based expression construct for the target cell-specific production of a therapeutic protein, such as a pro-apoptotic protein, within a target cell, including a target cell that is associated with aging, disease, or other condition, in particular a target cell that is a senescent cell or a cancer cell. Also provided are formulations and systems, including fusogenic lipid nanoparticle (LNP) formulations and systems, for the delivery of nucleic acid-based expression constructs as well as methods for making and using such nucleic acid-based expression constructs, formulations, and systems for reducing, preventing, and/or eliminating the growth and/or survival of a cell, such as a senescent cell and/or a cancer cell, which is associated with aging, disease, or other condition as well



WO 2019/204666 A1

WO 2019/204666 A1 

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

as methods for the treatment of aging, disease, or other conditions by the in vivo administration of a formulation, such as a fusogenic LPN formulation, comprising an expression construct for the target cell-specific, production of a therapeutic protein, such as a pro-apoptotic protein, in a target cell that is associated with aging, disease, or other condition, in particular a target cell that is a senescent cell or a cancer cell.

FUSOGENIC LIPID NANOPARTICLES AND METHODS FOR THE MANUFACTURE AND USE THEREOF FOR THE TARGET CELL-SPECIFIC PRODUCTION OF A THERAPEUTIC PROTEIN AND FOR THE TREATMENT OF A DISEASE, CONDITION, OR DISORDER ASSOCIATED WITH A TARGET CELL

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This PCT patent application was filed on April 18, 2019 and claims the benefit of U.S. Provisional Patent Application 62/659,676, filed April 18, 2018, and U.S. Provisional Patent Application 62/821,084, filed March 20, 2019. The entirety of the 62/659,676 and 62/821,084 Provisional Patent Applications are incorporated by reference herein.

SEQUENCE LISTING

[0002] The present application includes a Sequence Listing in electronic format as a txt file entitled "OSIN-01-0304WO01_2019-04-18_SEQLIST_ST25," which was created on April 18, 2019 and which has a size of 68,831 bytes. The present application also includes a Sequence Listing as a PDF file entitled "OSIN-01-0304WO01_2019-04-18_SEQLIST_ST25" and created on April 18, 2019. The contents of the txt and PDF files entitled "OSIN-01-0304WO01_2019-04-18_SEQLIST_ST25" are identical and are incorporated by reference herein.

BACKGROUND OF THE DISCLOSURE

Technical Field

[0001] The present disclosure relates, generally, to the field of medicine, including the treatment of disease, promotion of longevity, anti-aging, and health extension. More specifically, this disclosure concerns compositions and methods for reducing the growth and/or survival of cells that are associated with aging, disease, and other conditions. Provided are expression constructs for target cell specific expression of therapeutic proteins, which constructs exploit unique intracellular functionality, including transcription regulatory functionality, that is present within a target cell but is either absent from or substantially reduced in a normal, non-target cell. Such expression constructs are used in systems that include a vector for the delivery of a nucleic acid to a target cell, which vectors may comprise, but do not necessarily require, a fusogenic lipid nanoparticle and, optionally, a targeting moiety for enhancing the delivery of an expression construct to a target cell.

Description of the Related Art

[0002] Cancer cells, senescent cells, and other cells having an undesirable phenotype can accumulate over the course of a person's life and, without appropriate treatment, such cells can contribute to or even cause a person's morbidity and, ultimately, mortality.

[0003] The role of senescent cells in disease and the potential benefits of eliminating senescent cells has been discussed in scientific publications such as Baker *et al. Nature* 479:232–6 (2011). Systems and methods have been described that purport to address the problem of accumulating senescent cells. For example, Grigg, PCT Patent Publication No. WO 1992/009298, describes a system for preventing or reversing cell senescence with chemical compounds similar to carnosine and Gruber, U.S. Patent Publication No. 2012/0183534, describes systems for killing senescent cells with radiation, ultrasound, toxins, antibodies, and antibody-toxin conjugates, which systems include senescent cell-surface proteins for use in targeting of therapeutic molecules.

[0004] The selective killing of senescent cells has proven impractical in mammals other than genetically-modified laboratory research animals. Currently-available systems and methods exhibit substantial systemic toxicity, inadequate targeting of cells of interest, and a lack of adequate safety features. These shortcomings in the art have hampered the development of safe and effective therapies for the treatment of certain cancers and for slowing the effects of aging.

SUMMARY OF THE DISCLOSURE

[0005] The present disclosure is based upon the discovery that a cell, such as a cell that is associated with aging, a disease, and/or another condition (collectively, "a target cell"), can be selectively killed, in a target cell-specific manner, without the need for the targeted delivery of a therapeutic agent to the target cell. The expression constructs, systems, and methods described herein overcome safety and efficacy concerns that are associated with existing technologies that employ targeted delivery of therapeutic agents, which technologies have yielded limited therapeutic benefit to patients in need thereof.

[0006] As described herein, the present disclosure provides expression cassettes, systems, and methods for inducing, in a target cell-specific manner, the expression of a nucleic acid that encodes a protein that, when produced in a cell, reduces or eliminates the growth and/or survival of a cell, such as a cell that is associated with aging, disease, and/or other condition.

[0007] The expression cassettes, systems, and methods described herein exploit the unique transcription regulatory machinery that is intrinsic to certain cells that are associated with age

(such as senescent cells), disease (such as cancers, infectious diseases, and bacterial diseases), as well as other conditions, which transcription regulatory machinery is not operative, or exhibits substantially reduced activity, in a normal cell (*i.e.*, “a non-target cell”) that is not associated with aging, disease, or other condition.

[0008] The presently-disclosed expression cassettes, systems, and methods achieve a high degree of target cell specificity as a consequence of intracellular functionality that is provided by, and unique to, the target cell, which intracellular functionality is not provided by, or is substantially reduced in, a normal, non-target cell. Thus, the presently disclosed systems and methods employ nucleic acid delivery vectors that are non-specific with respect to the cell type to which the nucleic acid is delivered and, indeed, the vectors described herein need not be configured for target cell-specific delivery of a nucleic acid (*e.g.*, an expression cassette) to achieve target cell specificity and, consequently, the therapeutically effective reduction, prevention, and/or elimination in the growth and/or survival of a target cell.

[0009] Within certain embodiments, the present disclosure provides expression constructs for the targeted production of therapeutic proteins within a target cell, such as a cell that is associated with aging, disease, and/or another condition. The expression constructs disclosed herein comprise: (1) a transcriptional promoter that is activated in response to one or more factors each of which is produced within a target cell and (2) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of a cell, including the target cell.

[0010] Within certain aspects of these embodiments, the transcriptional promoter is activated in a target cell that is associated with a disease, condition, or age but is not activated in a normal mammalian cell that is not associated with the disease, condition, or aging. Target cell-specific transcriptional activation is achieved by the action of one or more factors that are produced in the target cell but not produced in a normal mammalian cell, including a normal human cell, such as normal skeletal myoblasts, normal adipose cells, normal cells of the eye, normal brain cells, normal liver cells, normal colon cells, normal lung cells, normal pancreas cells, and/or normal heart cells, which normal cells are not associated with the disease, condition, or aging.

[0011] Within other aspects of these embodiments, the target cell can be a mammalian cell or a bacterial cell. Target mammalian cells can include human cells such as senescent cells, cancer cells, precancerous cells, dysplastic cells, and cells that are infected with an infectious agent.

[0012] In certain aspects of these embodiments wherein the human target cell is a senescent cell, the transcriptional promoter can include a transcriptional promoter, such as the p16INK4a/CDKN2A transcriptional promoter, which is responsive to activation by transcription factors such as SP1, ETS1, and/or ETS2. In other aspects of these embodiments wherein the human target cell is a senescent cell, the transcriptional promoter can include a transcriptional promoter, such as the p21/CDKN1A transcriptional promoter, which is responsive to p53/TP53.

[0013] In a target cell, such as a senescent cell, transcriptional promoters induce the expression of a nucleic acid that encodes a therapeutic protein such as, for example, Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase as well as inducible and self-activating variants of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase which therapeutic protein reduces, prevents, and/or eliminates the growth and/or survival of the senescent cell, such as, for example, by inducing cell death in the senescent cell via a cellular process including apoptosis. Other therapeutic proteins may be employed that reduce, prevent, and/or eliminate the growth and/or survival of a senescent cell by, for example, inducing cell death via a cellular process including necrosis/necroptosis, autophagic cell death, endoplasmic reticulum-stress associated cytotoxicity, mitotic catastrophe, paraptosis, pyroptosis, pyronecrosis, and entosis.

[0014] In other aspects of these embodiments wherein the human target cell is a cancer cell, such as a brain cancer cell, a prostate cancer cell, a lung cancer cell, a colorectal cancer cell, a breast cancer cell, a liver cancer cell, a hematologic cancer cell, and a bone cancer cell, the transcriptional promoter can include the p21^{cip1/waf1} promoter, the p27^{kip1} promoter, the p53^{kip2} promoter, the TdT promoter, the Rag-1 promoter, the B29 promoter, the Blk promoter, the CD19 promoter, the BLNK promoter, and/or the λ 5 promoter, which transcriptional promoter is responsive to activation by one or more transcription factors such as an EBF3, O/E-1, Pax-5, E2A, p53, VP16, MLL, HSF1, NF-IL6, NFAT1, AP-1, AP-2, HOX, E2F3, and/or NF- κ B transcription factor, and which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein such as, for example, Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase as well as inducible and self-activating variants of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase which therapeutic protein reduces, prevents, and/or eliminates the growth and/or survival of the senescent cell, such as, for

example, by inducing cell death in the senescent cell via a cellular process including apoptosis. Other therapeutic proteins may be employed that reduce, prevent, and/or eliminate the growth and/or survival of a senescent cell by, for example, inducing cell death via a cellular process including necrosis/necroptosis, autophagic cell death, endoplasmic reticulum-stress associated cytotoxicity, mitotic catastrophe, paraptosis, pyroptosis, pyronecrosis, and entosis.

[0015] In still further aspects of these embodiments wherein the target cell is a human cell that is infected with an infectious agent, such as a virus, including, for example, a herpes virus, a polio virus, a hepatitis virus, a retrovirus virus, an influenza virus, and a rhino virus, or the target cell is a bacterial cell, the transcriptional promoter can be activated by a factor that is expressed by the infectious agent or bacterial cell, which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein such as, for example, Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase as well as inducible and self-activating variants of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase which therapeutic protein reduces, prevents, and/or eliminates the growth and/or survival of the senescent cell, such as, for example, by inducing cell death in the senescent cell via a cellular process including apoptosis. Other therapeutic proteins may be employed that reduce, prevent, and/or eliminate the growth and/or survival of a senescent cell by, for example, inducing cell death via a cellular process including necrosis/necroptosis, autophagic cell death, endoplasmic reticulum-stress associated cytotoxicity, mitotic catastrophe, paraptosis, pyroptosis, pyronecrosis, and entosis.

[0016] Within other embodiments, the present disclosure provides systems for the targeted production of a therapeutic protein within a target cell. These systems comprise a vector that is capable of delivering a nucleic acid to a cell, including a target cell as well as a non-target cell, wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with age, disease, or other condition) but not within a non-target cell, wherein the expression construct comprises a transcriptional promoter that is activated in response to one or more factors each of which is produced within said target cell; and a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of a cell in which it is produced, including a target cell.

[0017] Within certain aspects of these embodiments, formulations and systems include lipid nanoparticle (LNP) formulations and systems wherein an LPN encapsulates a polynucleotide

construct (e.g., a plasmid DNA) comprising a coding region for a pro-apoptotic protein, such as a caspase protein, and wherein the coding region is under the regulatory control of a target cell-specific transcriptional promoter, such as a senescent cell-specific transcriptional promoter or a cancer cell-specific transcriptional promoter. Exemplary cell-specific transcriptional promoters include p16, p22, p53. Exemplary coding regions for pro-apoptotic proteins include coding regions for Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase proteins. Pro-apoptotic proteins include inducible Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase proteins and self-activating Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase proteins, which are exemplified herein by an inducible Caspase 9 (iCasp9) or a self-activating Caspase 9 (saCasp9).

[0018] Inducible pro-apoptotic proteins, including iCasp9 proteins, can include a dimerization domain, such as an FKBP or FK506 binding protein domain, that binds to a chemical inducer of dimerization (CID), such as AP1903 or AP20187. Clackson, *Proc Natl Acad Sci U S A*, 95:10437-10442 (1998). Inducible Caspase 9 (iCasp9; Ariad, Erie, PA) may be activated in the presence of AP1903. U.S. Patent No. 5,869,337 and Straathof, *Blood* 105:4247-4254 (2005). Exemplary human genes encoding FKBP domains include AIP, AIPL1, FKBP1A, FKBP1B, FKBP2, FKBP3, FHBP5, FKBP6, FKBP7, FKBP8, FKBP8, FKBP9L, FKBP10, FKBP11, FKBP14, FKBP15, FKBP52, and LOC541473.

[0019] Within other aspects of these embodiments, lipid nanoparticles (LNP) are fusogenic lipid nanoparticles, such as fusogenic lipid nanoparticles comprising a fusogenic protein, such as a fusogenic p14 FAST fusion protein from reptilian reovirus to catalyze lipid mixing between the LNP and target cell plasma membrane. Suitable fusogenic proteins are described in PCT Patent Publication Nos. WO2012/040825A1 and WO2002/044206A2, Lau, *Biophys. J.* 86:272 (2004), Nesbitt, Master of Science Thesis (2012), Zijlstra, *AACR* (2017), Mrlouah, *PAACRAM* 77(13Suppl):Abst 5143 (2017), Krabbe, *Cancers* 10:216 (2018), Sanchez-Garcia, *ChemComm* 53:4565 (2017), Clancy, *J Virology* 83(7):2941 (2009), Sudo, *J Control Release* 255:1 (2017), Wong, *Cancer Gene Therapy* 23:355 (2016), and Corcoran, *JBC* 281(42):31778 (2006) and are exemplified by the P14 and P14e15 proteins having the amino acid sequences presented in Table 1.

Table 1
Fusogenic Protein Sequences

P14	MMSGPSNFMVNHAPGEAIVTGLEKGVAGTISHTIWEVIAG LVALLTFLAFGFWLFKYLQKRRERRRQLTEFQKRYLRNSYR LSEIQRPISEQHEYEDPYEPPSRRKPPPPYSTYVNIIDNVSAI*	SEQ ID NO: 16
P14e15	MMSGPSNFMVNHAPGEAIVTGLEKGVAGTISHTIWEVIAG LVALLTFLAFGFWLFKYLQWYNRKSKNKRRKEQIREQIELG LLSYGAGVASLPLLNVIHNP GSVISATPIYKGPCTGVPNSRL LQITSGTAEENTRILNHDGRNPDGSINV*	SEQ ID NO: 17

[0020] Contacting a cell expressing an iCasp9 protein with a CID facilitates the dimerization of the iCasp9 protein, which triggers apoptosis in a target cell. AP1903 has been used in humans multiple times, its intravenous safety has been confirmed, and its pharmacokinetics determined. Iuliucci, *J Clin Pharmacol* **41**(8):870-9 (2001) and Di Stasi, *N Engl J Med* **365**:1673-83 (2011). iCasp9 + AP1903 were used successfully in humans to treat GvHD after allogeneic T cell transplant. Di Stasi, *N Engl J Med* **365**:1673-83 (2011).

[0021] Within certain embodiments, a polynucleotide encoding a self-activating caspase, such as a self-activating Caspase 9 (saCasp9), may be employed wherein expression of the caspase polynucleotide is under the regulatory control of a factor that is active in a target cell population, such as a senescent cell population or a cancer cell population. Self-activating caspases activate in the absence of a chemical inducer of dimerization (CID). Cells expressing self-activating caspases, such as saCasp9, apoptose almost immediately. It will be appreciated by those of skill in the art that such self-activating caspases may be advantageously employed for the induction of apoptosis in a rapidly dividing cell, such as a rapidly dividing tumor cell, where an inducible caspase protein would be diluted out before administration of a CID. Moreover, because cell death with a self-activating caspase occurs over a longer period of time as compared to an inducible caspase, the risk of tumor lysis syndrome is reduced with a self-activating caspase.

[0022] Formulations comprising a plasmid DNA encapsulated with a LNP formulation are non-toxic and non-immunogenic in animals at doses of >15 mg/kg and exhibit an efficiency in excess of 80x greater than that achievable with neutral lipid formulations and 2-5x greater than that achievable with cationic lipid formulations. LNP cargo is deposited directly into the cytoplasm thereby bypassing the endocytic pathway.

[0023] Within further aspect of these embodiments, the system further comprises one or more safety features that permit additional control over the expression of the nucleic acid within the

expression construct or the functionality of a therapeutic protein encoded by the nucleic acid such as, for example, by requiring the contacting of a target cell with a chemical or biological compound that, in addition to the intracellular factor that promotes transcriptional activation of the promoter within the expression construct or promotes the functionality of the therapeutic protein, such as by promoting the dimerization of as well as inducible variants of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, and cytosine deaminase.

[0024] A further safety element that may be employed in the expression constructs and systems of the present disclosure includes a tamoxifen-inducible Cre construct using Life Technologies Gateway Cloning Vector System employing a pDEST26 plasmid for mammalian expression. For example, a fusion protein of Cre and estrogen receptor can be constitutively expressed and induced upon the addition of tamoxifen, which permits activated Cre to re-orient the transcriptional promoter, thereby expressing the therapeutic protein.

[0025] Within yet other aspects of these embodiments, the system may further comprise a nucleic acid that encodes a detectable marker, such as a bioluminescent marker, thereby allowing the identification of cells that express the therapeutic protein and, in the case of an inducible therapeutic protein such as an inducible Casp3, Casp8, Casp9, will be killed by the administration of a compound that promotes activity of the therapeutic protein, such as by inducing the dimerization of an inducible Casp3, Casp8, Casp9.

[0026] Within further embodiments, the present disclosure provides methods for reducing, preventing, and/or eliminating the growth of a target cell, which methods comprise contacting a target cell with a system for the targeted production of a therapeutic protein within a target cell, wherein the system comprises a vector that is capable of delivering a nucleic acid to a cell, wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with age, disease, or other condition) but not within a non-target cell, wherein the expression construct comprises: (a) a transcriptional promoter that is activated in response to one or more factors each of which factors is produced within a target cell and (b) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that is produced upon expression of the nucleic acid and wherein production of the therapeutic protein in the target cell (*i.e.*, the cell that is associated with age, disease, or other condition) reduces, prevents, and/or eliminates growth and/or survival of the target cell.

[0027] Within still further embodiments, the present disclosure provides methods for the treatment of an aging human or a human that is afflicted with a disease or another condition, wherein the aging, disease, or other condition is associated with a target cell within the human, the methods comprising administering to the human a system for the targeted production of a therapeutic protein within a target cell, wherein the system comprises a vector that is capable of delivering a nucleic acid to a cell, wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with age, disease, or other condition) but not within a non-target cell, wherein the expression construct comprises: (a) a transcriptional promoter that is activated in response to one or more factors each of which factors is produced within a target cell and (b) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that is produced upon expression of the nucleic acid and wherein production of the therapeutic protein in the target cell (*i.e.*, the cell that is associated with age, disease, or other condition) reduces, prevents, and/or eliminates growth and/or survival of the target cell thereby slowing aging in the human and/or slowing, reversing, and/or eliminating the disease or condition in the human.

[0028] Within further embodiments, the present disclosure provides lipid nanoparticle (LNP) formulation for the targeted production of a therapeutic protein within a target cell, which LNP formulation comprise: (a) a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, which mammalian cells include both target cells or non-target cells, wherein said lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s), and (b) an expression construct for the preferential production of a therapeutic protein within a target cell.

[0029] LNP formulations according to certain aspects of these embodiments include one or more lipid(s) at a concentration ranging from 1 mM to 100 mM, or from 5 mM to 50 mM, or from 10 mM to 30 mM, or from 15 mM to 25 mM. LNP formulations exemplified herein include one or more lipid(s) at a concentration of about 20 mM.

[0030] Within certain illustrative LNP formulations, one or more lipid(s) is selected from 1,2-dioleoyl-3-dimethylammonium-propane (DODAP), 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), Cholesterol, and 1,2-dimyristoyl-rac-glycero-3-methoxypolyethylene glycol (DMG-PEG). LNP formulations

may two or more lipids selected from the group consisting of DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG.

[0031] Exemplified herein are LNP formulations including DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG at a molar ratio of 35-55 mole % DODAP:10-20 mole % DOTAP:22.5-37.5 mole % DOPE:4-8 mole % Cholesterol:3-5 mole % DMG-PEG; or at a molar ratio of about 45 mole % DODAP:about 15 mole % DOTAP:about 30 mole % DOPE:about 6 mole % Cholesterol:about 4 mole % DMG-PEG. Within certain aspects, the LNP formulations include DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG at a molar ratio of 45 mole % DODAP:15 mole % DOTAP:30 mole % DOPE:6 mole % Cholesterol:4 mole % DMG-PEG.

[0032] LNP formulations according to other aspects of these embodiments include one or more fusogenic protein(s) at a concentration ranging from 0.5 μM to 20 μM , or from 1 μM to 10 μM , or from 3 μM to 4 μM . Exemplified herein are LNP formulations wherein fusogenic protein(s) are present at a concentration of 3.5 μM . Exemplary, suitable fusogenic protein(s) include the p14 fusogenic protein (SEQ ID NO: 16) and a the p14e15 fusogenic protein (SEQ ID NO: 17).

[0033] Within additional aspects of these embodiments, LNP formulations include expression constructs comprising (i) a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within said target cells as compared to said non-target cells and (ii) a nucleic acid that is operably linked to and under regulatory control of said transcriptional promoter, wherein said nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced within said target cells but is not produced in said non-target cells.

[0034] Exemplified herein are LNP formulations including expression constructs at a concentration ranging from 20 $\mu\text{g}/\text{mL}$ to 1.5 mg/mL , of from 100 $\mu\text{g}/\text{mL}$ to 500 $\mu\text{g}/\text{mL}$, or at a concentration of 250 $\mu\text{g}/\text{mL}$.

[0035] A suitable exemplary LNP formulation includes the following: for each 1mL of LNP, the lipid concentration is 20mM, the DNA content is 250 μg , and the fusogenic protein (e.g., p14 or p14e15) is at 3.5 μM wherein the lipid formulation comprises DODAP:DOTAP:DOPE:Cholesterol:DMG-PEG at a mole % ratio of 45:15:30:6:4, respectively.

[0036] Within still further aspects of these embodiments, LNP formulations include expression constructs having a transcriptional promoter selected from a p16 transcriptional promoter, a p21 transcriptional promoter, and a p53 transcriptional promoter, and include transcriptional promoters that are responsive to a factor selected from SP1, ETS1, ETS2, and p53/TP53. Exemplified herein are LNP formulations wherein said transcriptional promoter is a p16INK4a/CDKN2A transcriptional promoter or a p21/CDKN1A transcriptional promoter.

[0037] Within related aspects of these embodiments, LNP formulations include expression constructs having a transcriptional promoter that is responsive to a factor selected from EBF3, O/E-1, Pax-5, E2A,p53, VP16, MLL, HSF1, NF-IL6, NFAT1, AP-1, AP-2, HOX, E2F3, and/or NF- κ B. Exemplified herein are LNP formulations wherein said transcriptional promoter is a p21^{cip1/waf1} promoter, the p27^{kip1} promoter, the p57^{kip2} promoter, the TdT promoter, the Rag-1 promoter, the B29 promoter, the Blk promoter, the CD19 promoter, the BLNK promoter, and the λ 5 promoter.

[0038] Within other related aspects of these embodiments, LNP formulations include expression constructs that include a nucleic acid that encodes a therapeutic protein, such as a therapeutic protein selected from a caspase (Casp), an inducible caspase (iCasp), a self-activating caspase (saCasp), BAX, DFF40, HSV-TK, and cytosine deaminase. Exemplified herein are LNP formulations that include expression constructs having a nucleic acid that encodes a Casp9, including, for example, an inducible Casp9 (iCasp9) or a self-activating Casp9 (saCasp9).

[0039] Other embodiments of the present disclosure provide methods for reducing, preventing, and/or eliminating the growth of a target cell, which comprise contacting a target cell with an LNP formulation having (a) a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, which mammalian cells include both target cells or non-target cells, wherein said lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s), and (b) an expression construct for the preferential production of a therapeutic protein within a target cell.

[0040] Within certain aspects of these embodiments the methods employ LNP formulations comprising (i) a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within target cells as compared to non-target cells and (ii) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate

the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced within the target cells but is not produced in the non-target cells.

[0041] Other embodiments of the present disclosure provide methods for the treatment of a disease or condition in a patient, including a human patient, having a target cell, wherein the method comprises administering to the patient an LNP formulation having (a) a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, wherein the mammalian cells include both target cells or non-target cells, and wherein the lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s) and (b) an expression construct for the preferential production of a therapeutic protein within a target cell.

[0042] Within certain aspects of these embodiments the methods employ LNP formulations comprising (i) a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within target cells as compared to non-target cells and (ii) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced within the target cells but is not produced in the non-target cells.

[0043] These and other related aspects of the present disclosure will be better understood in light of the following drawings and detailed description, which exemplify certain aspects of the various embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a diagrammatic representation of conventional and fusogenic liposomes, including stealth fusogenic liposomes, including lipid nanoparticles employing Innovascreen's Fusogenix™ Platform according to certain aspects of the present disclosure. Shown are Fusogenix™ lipid nanoparticles utilizing a p14 FAST fusion protein from reptilian reovirus and including a plasmid vector encoding an inducible Caspase 9 (iCasp9) under a promoter that is active in a target cell population, such as a senescent target cell population or a cancer target cell population. Exemplified in this diagram are Casp9 fusion peptides that are activated via a small molecule dimerizer such as AP1903.

FIG. 2 is a diagrammatic representation of the liposomal delivery to the cytoplasm of a

target cell, according to certain aspects of the present disclosure. Shown are Fusogenix™ lipid nanoparticles (LNPs) that are configured for the delivery of nucleic acids, such as those encoding a pro-apoptotic protein, such as Caspase 9, under the regulatory control of a target cell-specific transcriptional promoter, such as a target senescent cell encoding p16 or a target cancer cell encoding p53. Exemplified are Fusogenix™ lipid nanoparticles comprising a p14 FAST protein to catalyze the rapid lipid mixing between the lipid nanoparticle (LNP) and the target cell plasma membrane. Such Fusogenix™ lipid nanoparticles (i) deliver the cargo nucleic acids directly into the cytoplasm thereby bypassing the endocytic pathway, (ii) are non-toxic (i.e., non-immunogenic) in animals at doses of ≥ 15 mg/kg, (iii) are 80x more efficient than neutral lipid formulations, (iv) are 2-5x more efficient than cationic lipid formulations, and (iv) are manufacturable at scale.

FIG. 3 is a table comparing the reported maximum tolerated dose (MTD) for clinical stage lipid-based *in vivo* delivery technologies. The MTD of >15 mg/kg for fusogenic lipid nanoparticles of the present disclosure was estimated from rat toxicity data.

FIG. 4A is a diagrammatic representation of the induction of an inducible Caspase 9 homodimer (iCasp9), which iCasp9 is a fusion protein comprising a drug-binding domain for binding to a chemical inducer of dimerization (CID) and an active portion of Caspase 9. A CID, as exemplified by CIDs designated AP1903 and AP20187, binds to the drug-binding domain of the iCasp9 fusion protein to dimerize and, thereby, activate iCasp9, which results in the intracellular activation of pro-apoptotic molecules and the induction of apoptosis within a target cell.

FIG. 4B is a diagrammatic representation of an exemplary apoptosome according to certain aspects of the present disclosure.

FIG. 5 depicts the chemical structure of an exemplary chemical inducer of dimerization (CID), which is a homodimerizer designated herein as AP1903 (APExBIO, Houston, TX) that may be employed in various embodiments of the present disclosure for inducing the activity of an inducible pro-apoptotic protein, such as an inducible caspase protein (e.g., iCasp9).

FIG. 6 depicts the chemical structure of an exemplary chemical inducer of dimerization (CID), which is a homodimerizer designated herein as AP20187 (APExBIO, Houston, TX) that may be employed in various embodiments of the present disclosure for inducing the activity of an inducible pro-apoptotic protein, such as an inducible caspase protein (e.g., iCasp9).

FIG. 7 presents data obtained in mice that were administered intravenously Fusogenix lipid nanoparticles labeled with ^{64}Cu -NOTA [1,4,7-triazacyclononane-1,4,7-triacetic acid]. See, Fournier, *EJNMMI Research* 2:8 (2012). ^{64}Cu was detected via positron emission tomography (PET). **FIG. 7A** presents PET data obtained from a mouse to which ^{64}Cu -NOTA-liposomes without protein were administered. **FIG. 7B** presents PET data obtained from a mouse to which ^{64}Cu -NOTA-liposome-p14 were administered.

FIG. 8 is a bar graph of data obtained with Fusogenix lipid nanoparticles comparing $\text{SUV}_{\text{mean}, 24\text{h}}$ for ^{64}Cu -NOTA-liposomes without protein and ^{64}Cu -NOTA-liposome-p14. The data presented in **FIGs. 7** and **8** demonstrate a 50% increase in gene/siRNA delivery to prostate tumors as compared to a competing formulation.

FIG. 9 is a bar graph of the biodistribution of labelled pegylated liposomes in nude mice expressed after 24 hours as discussed in Example 1.

FIGs. 10 and **11** are graphs of optical density at 405nm as a function of concentration ($\mu\text{g}/\text{ml}$; **FIG. 10**) and of anti-p14 and anti-LNP antibody responses (**FIG. 11**), which demonstrate the safety and tolerability of exemplary fusogenic lipid nanoparticles utilizing a reptilian reovirus p14 FAST fusion protein (FusogenixTM). As shown, virtually no antibody response was observed in immune competent mice (with and without adjuvant).

FIGs. 12 and **13** are bar graphs of data from *in vitro* anti-p14 and anti-LNP antibody neutralization assays showing that lipid nanoparticle formulations according to the present disclosure are non-reactive with C4d (**FIG. 12**) and less reactive with iC3b (**FIG. 13**) as compared to Doxil in 8 out of 10 human samples tested for Complement activation-related pseudoallergy (CARPA) using C4d and iC3b complement ELISA assays as described in Szebani, *Mol Immunol* 61(2):163-73 (2014).

FIG. 14 is a restriction map of the plasmid vector pVAX1TM which is employed in certain aspects of the expression constructs, systems, formulations, and methods of the present disclosure for the target cell-specific production of a therapeutic protein, such as a pro-apoptotic protein, including a caspase protein, such as Caspase 9, as well as inducible and self-activating variants of a pro-apoptotic protein, including inducible and self-activating variants of caspase proteins, such as inducible Caspase 9 (iCasp9) and self-activating Caspase 9 (saCasp9). In certain embodiments, expression constructs and formulations may additionally include a safety element, such as a tamoxifen-inducible Cre construct (e.g., Life Technologies Gateway Cloning Vector

System). A fusion protein of Cre and estrogen receptor is constitutively expressed and induced upon the addition of tamoxifen, which permits activated Cre to re-orient the p16-promoter, thereby expressing caspase 9 or inducible/self-activating variant thereof. pVAX1 is commercially available from ThermoFisher Scientific (Waltham, MA).

FIG. 15 is a diagrammatic representation of an exemplary p16-targeting construct for the target cell-specific expression of an inducible Caspase 9 (iCasp9) or a self-activating Caspase 9 (saCasp9) protein in target cells expressing p16, such as target cells that are associated with aging and/or senescence, which p16-targeting construct comprises a p16s transcriptional promoter in operable connection to iCasp9 or saCasp9. An exemplary p16 transcriptional promoter is described in Baker *et al.*, *Nature* 479(7372):232-67 (2011)).

FIG. 16 is a restriction map of the plasmid vector pVAX1-16s-iCasp9-MX (SEQ ID NO: 6), which comprises an exemplary p16-targeting construct for the target cell-specific expression of an inducible Caspase 9 (iCasp9) protein in target cells expressing p16, such as target cells that are associated with aging and/or senescence, which p16-targeting construct comprises a p16s transcriptional promoter in operable connection to iCasp9.

FIG. 17 is a plasmid map of the vector p10-p16-iCasp9 (SEQ ID NO: 12), which comprises an exemplary p16-targeting construct for the target cell-specific expression of an inducible Caspase 9 (iCasp9) protein in target cells expressing p16, such as target cells that are associated with aging and/or senescence, which p16-targeting construct comprises a p16e transcriptional promoter in operable connection to iCasp9.

FIG. 18 is a plasmid map of the vector p10-p16-saCasp9 (SEQ ID NO: 13), which comprises an exemplary p16-targeting construct for the target cell-specific expression of a self-activating Caspase 9 (saCasp9) protein in target cells expressing p16, such as target cells that are associated with aging and/or senescence, which p16-targeting construct comprises a p16e transcriptional promoter in operable connection to saCasp9.

FIG. 19 is a diagrammatic representation of the *in vivo* administration of an exemplary p16-targeting construct in an mouse model system for aging, wherein the aging mouse model exhibits a senescent cell burden (as defined by the presence of p16⁺ cells) and secretion of factors associated with a senescence-associated secretory phenotype (SASP; van Deursen, *Nature* 509(7501):439-446 (2014)). A formulation comprising a vector and an expression construct, such as a lipid nanoparticle (LNP) vector, *e.g.*, a fusogenic LNP comprising a fusogenic protein such

as p14 FAST, encompassing a p16-Casp9 expression construct, *e.g.*, pVAX1-16s-iCasp9, p10-p16e-iCasp9, p10-p16e-saCasp9, or variant thereof expressing luciferase (for visualization), is administered *in vivo* to an aged mouse via injection into a tail vein and the LNP+ expression construct transfects target and non-target cells without specificity. Upon subsequent *in vivo* administration of a chemical inducer of dimerization (CID), such as AP20187, p16+ target cells (*e.g.*, senescent cells) expressing an iCasp9 protein undergo apoptosis, resulting in a reduction in SASP levels, while p16- cells remain viable.

FIGs. 20A-D is a photomicrograph of the histological staining of senescent-associated β -gal in kidney cells from an *in vivo* aged mouse model either untreated (upper panel) or treated (low dose – middle panel and high dose – lower panel) following the *in vivo* administration (16 animals at 80 weeks of age) of a formulation comprising a vector and an expression construct, such as a lipid nanoparticle (LNP) vector, *e.g.*, a fusogenic LNP comprising a fusogenic protein such as p14 FAST, encompassing a p16-Casp9 expression construct, *e.g.*, pVAX1-16s-iCasp9, p10-p16e-iCasp9, p10-p16e-saCasp9, or variant thereof, is administered *in vivo* to an aged mouse and kidney cells stained for β -gal. These data demonstrated a dose-dependent reduction of p16+ senescent kidney cells.

FIGs. 20E-H is a photomicrograph of the histological staining of senescent-associated β -gal in seminal vesicle cells from an *in vivo* aged mouse model either untreated (upper panel) or treated (low dose – middle panel and high dose – lower panel) following the *in vivo* administration (16 animals at 80 weeks of age) of a formulation comprising a vector and an expression construct, such as a lipid nanoparticle (LNP) vector, *e.g.*, a fusogenic LNP comprising a fusogenic protein such as p14 FAST, encompassing a p16-iCasp9 expression construct, *e.g.*, pVAX1-16s-iCasp9, p10-p16e-iCasp9, p10-p16e-saCasp9, or variant thereof, is administered *in vivo* to an aged mouse and seminal vesicle cells stained for β -gal. These data demonstrated a dose-dependent reduction of p16+ senescent seminal vesicle cells.

FIG. 21 is a bar graph demonstrating the dose-dependent targeting of p16+ kidney cells in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Kidney cells were subjected to a qRT-PCR reaction to detect p16^{lnk4a} transcripts. Relative expression was calculated using $2^{-\Delta\Delta Ct}$ (Livak, *Methods* 25:402-408 (2001)).

FIG. 22 is a bar graph demonstrating the dose-dependent targeting of p16+ spleen cells

in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Spleen cells were subjected to a qRT-PCR reaction to detect p16^{lnk4a} transcripts. Relative expression was calculated using $2\Delta\Delta Ct$ (Livak, *Methods* 25:402-408 (2001)).

FIG. 23 is a bar graph demonstrating the dose-dependent targeting of p16+ seminal vesicle cells in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Seminal vesicle cells were subjected to a qRT-PCR reaction to detect p16^{lnk4a} transcripts. Relative expression was calculated using $2\Delta\Delta Ct$ (Livak, *Methods* 25:402-408 (2001)).

FIG. 24 is a bar graph demonstrating the dose-dependent targeting of p16+ inguinal fat cells in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Inguinal fat cells were subjected to a qRT-PCR reaction to detect p16^{lnk4a} transcripts. Relative expression was calculated using $2\Delta\Delta Ct$ (Livak, *Methods* 25:402-408 (2001)).

FIG. 25 is a bar graph demonstrating the dose-dependent targeting of p16+ lung cells in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Lung cells were subjected to a qRT-PCR reaction to detect p16^{lnk4a} transcripts. Relative expression was calculated using $2\Delta\Delta Ct$ (Livak, *Methods* 25:402-408 (2001)).

FIG. 26 is a bar graph of data demonstrating the remediation of chemotherapy-induced damage (as determined by the clearance of damaged cells (*i.e.*, senescent cells) after treatment with doxorubicin). Senescence was induced in B6 mice with doxorubicin. Animals were treated with murine p53-iCasp9 and dimerizer or controls (dimerizer only and LNP only) and sacrificed. Tissues were assayed for p53 expression via rt-PCR.

FIG. 27 is a diagrammatic representation of an exemplary p53-targeting cassette for use in treatment of cancers (oncology) by the selective killing of tumor cells according certain embodiments of the present disclosure. The p53-targeting cassette comprises a p53 transcriptional promoter, which drives the expression an inducible caspase 9 protein (iCasp9) or a self-activating caspase 9 protein (saCasp9).

FIG. 28 is a restriction map of a plasmid (pVAX1-p53-iCasp9-MX; SEQ ID NO: 7)

comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of an iCasp9 nucleic acid encoding an inducible Casp9 protein is regulated by the p53 transcriptional promoter.

FIG. 29 is a restriction map of a plasmid (pVAX1-p53-saCasp9; SEQ ID NO: 8) comprising a p53-targeting cassette. Expression of a nucleic acid encoding a self-activating Caspase 9 (saCasp9) protein is regulated by the p53 transcriptional promoter.

FIG. 30 is a restriction map of a plasmid (pVAX1-p53-iCasp9-OVA; SEQ ID NO: 11) comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of a nucleic acid encoding an inducible Casp9 protein is regulated by the p53 transcriptional promoter.

FIG. 31 is a restriction map of a plasmid (pVAX1-p53-iCasp9-G-O; SEQ ID NO: 9) comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of an iCasp9 nucleic acid encoding an inducible Casp9 protein is regulated by the p53 transcriptional promoter.

FIG. 32 is a restriction map of a plasmid (pVAX1-p53-iCasp9-huCD40L; SEQ ID NO: 10) comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of an iCasp9 nucleic acid encoding an inducible Casp9 protein is regulated by the p53 transcriptional promoter. Additional targeting cassettes and plasmid constructs have been developed for advanced oncology applications, as disclosed herein, which constructs employ nucleic acids encoding, for example, one or more immunostimulatory cytokines (such as huCD40L, as shown in **FIG. 32**, as well as GMCSF and IL12) and/or one or more antigens (such as chicken ovalbumin (OVA), as shown in **FIG. 30**, as well as Nt1, tetanus antigens, and influenza antigens).

FIG. 33 is a map of a plasmid (p10-p53e-iCasp9; SEQ ID NO: 14) comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of an iCasp9 nucleic acid encoding an inducible Casp9 protein is regulated by the p53 transcriptional promoter. Additional targeting cassettes and plasmid constructs have been developed for advanced oncology applications, as disclosed herein, which constructs employ nucleic acids encoding, for example, one or more immunostimulatory cytokines (such as huCD40L, as shown in **FIG. 32**, as well as GMCSF and IL12) and/or one or more antigens (such as chicken ovalbumin (OVA), as shown in **FIG. 30**, as well as Nt1, tetanus antigens, and influenza antigens).

FIG. 34 is a map of a plasmid (p10-p53e-saCasp9; SEQ ID NO: 15) comprising a p53-targeting cassette as depicted in **FIG. 27**. Expression of an saCasp9 nucleic acid encoding a self-activating Casp9 protein is regulated by the p53 transcriptional promoter. Additional targeting

cassettes and plasmid constructs have been developed for advanced oncology applications, as disclosed herein, which constructs employ nucleic acids encoding, for example, one or more immunostimulatory cytokines (such as huCD40L, as shown in **FIG. 32**, as well as GM-CSF and IL12) and/or one or more antigens (such as chicken ovalbumin (OVA), as shown in **FIG. 30**, as well as Nt1, tetanus antigens, and influenza antigens).

FIG. 35 is a diagram showing the rationale for targeting p53+ tumors with expression constructs comprising a p53 promoter in operable combination with a pro-apoptotic protein, such as a caspase protein, *e.g.*, a Caspase 9 protein. Cancer cells often mutate or delete it so they can grow uncontrollably. However, even when the p53 gene is mutated, the transcription factors that bind to it are almost always still active.

FIG. 36 is a Western blot of iCasp 9 and Casp 9 protein levels obtained with p53-expressing cells (pVax-p53) and control cells (pcDNA3-GFP). Human prostate cancer PC-3 cells were treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc (luciferin) plasmid (in the presence and absence of the homodimerizer AP201870) and assessed for iCasp9 expression. These data demonstrate that addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolishes the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

FIGs. 37A-D are microscopic images of human prostate cancer (LNCaP, DU145, PC-3) or normal epithelial (RWPE) cells treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc plasmid and assessed for iCasp9 expression by Western blot and luminescence assays.

FIGs. 38-41 are bar graphs of data obtained with the p53-expressing cells presented in **FIG. 37**. Human prostate cancer (LNCaP (**FIG. 38**), DU145 (**FIG. 39**), PC-3 (**FIG. 40**) or normal epithelial (RWPE (**FIG. 41**)) cells were treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc plasmid and assessed for iCasp9 expression by Western blot and luminescence assays. These data demonstrate that addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolishes the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

FIG. 42 is a bar graph of data from a luminescence assay of iCasp 9 and Casp 9 protein levels obtained with the p53-expressing cells presented in **FIG. 36** (pVax-p53) and control cells (pcDNA3-GFP). Human prostate cancer PC-3 cells were treated with Fusogenix lipid

nanoparticles carrying the pVax-p53-iCasp9-luc (luciferin) plasmid (in the presence and absence of the homodimerizer AP20187) and assessed for iCasp9 expression. These data demonstrate that addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolishes the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

FIGs. 43A, 43B, 44A, and 44B are flow cytometry apoptosis data (Annexin V) from human prostate cancer PC-3 cells treated with pVax-p53 Fusogenix lipid nanoparticles (in the absence and presence of AP20187, **FIG. 43A and 44A** and **43B and 44B**, respectively). The data presented in these figures demonstrates that suicide gene therapy selectively kills p53-expressing human prostate cancer cells in culture by inducing apoptosis (Luciferase-Annexin V flow cytometry).

FIG. 45 is a flow diagram depicting a pre-clinical oncology study according to the present disclosure with 30x NSG mice implanted with human prostate tumor cells.

FIG. 46 is a graph of tumor volume (mm^3) from the pre-clinical oncology study depicted in **FIG. 33** in which NSG mice bearing a subcutaneous human prostate PC-3 tumor was injected intratumorally (IT) with 100 μg Fusogenix pVax-p53 formulation, followed 96 hours later by intravenous (IV) administration of 2 mg/kg of the homodimerizer AP20187.

FIGs. 47A-47C are photographs of tumors from the IT injection oncology study of **FIG. 46** in which NSG mouse bearing a subcutaneous human prostate PC-3 tumor was injected intratumorally with 100 μg Fusogenix pVax-p53 formulation, followed 96 hours by 2 mg/kg AP20187 IV.

FIG. 48 is a graph from the first of four NSG mice bearing subcutaneous human prostate cancer PC-3 tumors that were injected intravenously (IV) with 4x 100 μg doses of Fusogenix pVax-p53 formulation, followed 24 hours later by 2 mg/kg AP20187 IV.

FIG. 49 is a graph from the second of four NSG mice bearing subcutaneous human prostate cancer PC-3 tumors that were injected intravenously (IV) with 4x 100 μg doses of Fusogenix pVax-p53 formulation, followed 24 hours later by 2 mg/kg AP20187 IV.

FIG. 50 is a graph from the third of four NSG mice bearing subcutaneous human prostate cancer PC-3 tumors that were injected intravenously (IV) with 4x 100 μg doses of Fusogenix pVax-p53 formulation, followed 24 hours later by 2 mg/kg AP20187 IV.

FIG. 51 is a graph from the fourth of four NSG mice bearing subcutaneous human prostate cancer PC-3 tumors that were injected intravenously (IV) with 4x 100 µg doses of Fusogenix pVax-p53 formulation, followed 24 hours later by 2 mg/kg AP20187IV.

FIG. 52 is a graph showing the percentage change in tumor volume as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in NSG mice (N=6 for all groups) bearing a prostate tumor that were treated with intravenous p14 LNP pVAX.

FIG. 53 is a survival curve showing the percent survival as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in NSG mice (N=6 for all groups) bearing a prostate tumor that were treated with intravenous p14 LNP pVAX.

FIG. 54 is a graph of dose escalation data showing the percentage change in tumor volume as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in NOD-SCID mice (N=6 for all groups) bearing a prostate tumor that were treated with 100 µg, 400 µg, and 1000 µg of intravenous p14 LNP pVAX. NOD-SCID mice were implanted subcutaneously with 500,000 PC-3 cells and randomized into treatment groups when their tumors reached 200 mm³, (N = 2 for all groups). Animals were injected with their assigned dose of p53-iCasp9 LNP IV twice followed by 2 mg/kg dimerizer. Tumors were measured directly every 24 hours.

FIG. 55 is a graph showing the suppression of metastatic tumor growth with repeat treatment of a p53-iCasp9 LNP with or without a chemical inducer of dimerization (CID). NOD-SCID mice were injected with 500,000 PC-3M-luciferase cells on Day 0, LNP dosing was started on Day 22 with 150µg p53-iCasp9 LNP. Dimerizer doses started Day 24 at 2 mg/kg. Mice were imaged every 24-48 hours to detect whole animal luminescence.

FIGs. 56 and 57 are graphs showing the percentage change in tumor volume (**FIG. 56**) and percent survival (**FIG. 57**) as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in isogenic C57B6 mice implanted with B16 murine melanoma cells treated with LNPs containing a construct encoding iCasp9 and murine CD40L under control of the murine p53 promoter. Even though the rapid (10 hour) doubling time of the B16 cells made them largely refractory to the iCasp9-induced apoptosis, they still secreted enough CD40L to effectively halt the tumor's growth. A construct encoding GMCSF + OVA antigen was also tested and determined to be more effective than iCasp9 alone, but less effective than the CD40L version. N=3 for both groups.

FIGs. 58A-58D and 59 are photographs and a bar graph, respectively, of a B15F10 lung metastasis model data in which 100 µg of a control LNP or a p53-iCasp9 LNP was administered intravenously at days 3, 6, 9, and 12 following the intravenous injection of 75,000 B16F10 cells. At days 5, 8, 11, and 13, a chemical inducer of dimerization (CID) was administered intraperitoneally. Animals were sacrificed at day 14 and lung metastases were quantified.

FIGs. 60 and 61 are DEXA scans, which were performed monthly, after *in vivo* administration of LNP formulations targeting p16, p53, or the combination (p16 + p53) (N=10 for all groups). Mice were treated monthly starting at 728 days (104 weeks) of age.

FIG. 62 and 63 are graphs showing the change in bone density in male (**FIG. 62**) and female (**FIG. 63**) naturally aged mice after *in vivo* administration of LNP formulations targeting p16, p53, or the combination (p16 + p53) (N=10 for all groups). Mice were treated monthly starting at 728 days (104 weeks) of age (arrows). At 896 days (128 weeks), the increase in bone density benefit for treated mice is apparent in the male mice.

FIG. 64 is a survival curve showing the percent survival as a function of time after *in vivo* administration of LNP formulations targeting p16, p53, or the combination (p16 + p53) (N=10 for all groups). Mice were treated monthly starting at 728 days (104 weeks) of age (arrows). At 931 days (133 weeks), the survival benefit for treated mice is apparent (>50% survival difference between combination treatment and control).

DETAILED DESCRIPTION

[0044] The present disclosure provides expression cassettes, systems, and methods for the selective reduction, prevention, and/or elimination in the growth and/or survival of a cell that is associated with aging, disease, or another condition (collectively “a target cell”), which expression cassettes, systems, and methods overcome the safety and efficacy concerns that are associated with existing technologies that rely on targeted delivery of a therapeutic compound and, as a result of, for example, inefficient target cell delivery and/or off-target effects, have limited therapeutic benefit.

[0045] More specifically, the expression cassettes, systems, and methods disclosed herein exploit the cell-specific transcription regulatory machinery that is intrinsic to a target cell and, thereby, achieve a target cell-specific therapeutic benefit without the need for targeted- delivery of a therapeutic compound. These expression cassettes, systems, and methods permit the target cell-specific induction of expression of a nucleic acid that encodes a therapeutic protein, which

protein can reduce, prevent, and/or eliminate the growth and/or survival of a cell in which it is produced.

[0046] Thus, the various embodiments that are provided by the present disclosure include:

1. Expression constructs for the targeted production of therapeutic proteins within a target cell, such as a cell that is associated with aging, disease, and/or another condition, the expression construct comprising:

a. transcriptional promoter that is activated in response to one or more factors each of which is produced within a target cell and

b. a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of a cell, including the target cell.

2. Systems for the targeted production of a therapeutic protein within a target cell, the systems comprising a vector for delivering a nucleic acid to a cell, including a target cell as well as a non-target cell,

wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with aging, cancer, and/or other disease and/or condition) but not within a non-target cell,

wherein the expression construct comprises (i) a transcriptional promoter that is activated in response to one or more factors each of which is produced within a target cell and (ii) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter,

wherein the nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of a cell in which it is produced, including a target cell.

3. Methods for reducing, preventing, and/or eliminating the growth of a target cell, the methods comprising contacting a target cell with a system for the targeted production of a therapeutic protein within a target cell,

wherein the system comprises a vector for delivery of a nucleic acid to a cell,

wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with age, disease, or other condition) but not within a non-target cell,

wherein the expression construct comprises (i) a transcriptional promoter that is activated in response to one or more factors each of which factors is produced within a target cell

and (ii) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter,

wherein the nucleic acid encodes a therapeutic protein that is produced upon expression of the nucleic acid and

wherein production of the therapeutic protein in the target cell (*i.e.*, the cell that is associated with age, disease, or other condition) reduces, prevents, and/or eliminates growth and/or survival of the target cell.

4. Methods for the treatment of aging, disease, or other condition in a human, wherein aging, disease, or other condition is associated with a target cell, the methods comprising administering to the human a system for the targeted production of a therapeutic protein within a target cell,

wherein the system comprises a vector that is capable of delivering a nucleic acid to a cell,

wherein the vector comprises an expression construct for the targeted production of a therapeutic protein within a target cell (*e.g.*, a cell that is associated with age, disease, or other condition) but not within a non-target cell,

wherein the expression construct comprises (i) a transcriptional promoter that is activated in response to one or more factors each of which factors is produced within a target cell and (ii) a nucleic acid that is operably linked to and under regulatory control of the transcriptional promoter,

wherein the nucleic acid encodes a therapeutic protein that is produced upon expression of the nucleic acid and

wherein production of the therapeutic protein in the target cell (*i.e.*, the cell that is associated with age, disease, or other condition) reduces, prevents, and/or eliminates growth and/or survival of the target cell thereby slowing aging in the human and/or slowing, reversing, and/or eliminating the disease or condition in the human.

Definitions

[0047] These and other aspects of the present disclosure can be better understood by reference to the following non-limiting definitions.

[0048] As used herein, the term “transcriptional promoter” refers to a region of DNA that initiates transcription of a particular gene. Promoters are located near transcription start sites of genes, on the same strand and upstream on the DNA (towards the 3' region of the anti-sense

strand, also called template strand and non-coding strand). Promoters can be about 100–1000 base pairs long. For the transcription to take place, the enzyme that synthesizes RNA, known as RNA polymerase, must attach to the DNA near a gene. Promoters contain specific DNA sequences and response elements that provide a secure initial binding site for RNA polymerase and for proteins called transcription factors that recruit RNA polymerase. These transcription factors have specific activator or repressor sequences of corresponding nucleotides that attach to specific promoters and regulate gene expressions. The process is more complicated, and at least seven different factors are necessary for the binding of an RNA polymerase II to the promoter. Promoters represent critical elements that can work in concert with other regulatory regions (enhancers, silencers, boundary elements/insulators) to direct the level of transcription of a given gene.

[0049] Eucaryotic transcriptional promoters comprise a number of essential elements, which collectively constitute a core promoter (*i.e.*, the minimal portion of a promoter that is required to initiate transcription). Those elements include (1) a transcription start site (TSS), (2) an RNA polymerase binding site (in particular an RNA polymerase II binding site in a promoter for a gene encoding a messenger RNA), (3) a general transcription factor binding site (*e.g.*, a TATA box having a consensus sequence TATAAA, which is a binding site for a TATA-binding protein (TBP)), (4) a B recognition element (BRE), (5) a proximal promoter of approximately 250 bp that contains regulatory elements, (6) transcription factor binding sites (*e.g.*, an E-box having the sequence CACGTF, which is a binding site for basic helix- loop-helix (bHLH) transcription factors including BMAL1-Clock and cMyc), and (7) a distal promoter containing additional regulatory elements. As used herein, the term “transcriptional promoter” is distinct from the term “enhancer,” which refers to a regulatory element that is distant from the transcriptional start site.

[0050] Eucaryotic promoters are often categorized according to the following classes: (1) AT-based class, (2) CG-based class, (3) ATCG-compact class, (4) ATCG-balanced class, (5) ATCG-middle class, (6) ATCG-less class, (7) AT-less class, (8) CG-spike class, (9) CG-less class, and (10) ATspike class. See, Gagniuc and Ionescu-Tirgoviste, *BMC Genomics* 13:512 (2012). Eucaryotic promoters can be “unidirectional” or “bidirectional.” Unidirectional promoters regulate the transcription of a single gene and are characterized by the presence of a TATA box. Bidirectional promoters are short (<1 kbp), intergenic regions of DNA between the 5' ends of genes in a bidirectional gene pair (*i.e.*, two adjacent genes coded on opposite strands having 5' ends oriented toward one another). Bidirectional genes are often functionally related and because they share a single promoter, can be co-regulated and co-expressed. Unlike unidirectional

promoters, bidirectional promoters do not contain a TATA box but do contain GpC islands and exhibit symmetry around a midpoint of dominant Cs and As on one side and Gs and Ts on the other. CCAAT boxes are common in bidirectional promoters as are NRF-1, GABPA, YY1, and ACTACAnnTCCC motifs.

[0051] Transcriptional promoters often contain two or more transcription factor binding sites. Thus, the efficient expression of a nucleic acid that is downstream of a promoter having multiple transcription factor binding sites typically requires the cooperative action of multiple transcription factors. Accordingly, the specificity of transcriptional regulation, and hence expression of an associated nucleic acid, can be increased by employing transcriptional promoters having two or more transcription factor binding sites.

[0052] As used herein, the term “transcription factor” refers to sequence-specific DNA-binding factors that bind to specific sequences within a transcriptional promoter thereby regulating the transcription of a nucleic acid that is in operable proximity to and downstream of the promoter. Transcription factors include activators, which promote transcription, and repressors, which block transcription by preventing the recruitment or binding of an RNA polymerase. Transcription factors typically contain (1) one or more DNA-binding domains (DBDs), which facilitate sequence specific binding to a cognate transcription factor binding site (a/k/a response element) within a transcriptional promoter; (2) one or more signal-sensing domains (SSDs), which includes ligand binding domains that are responsive to external signals; and (3) one or more transactivation domains (TADs), which contain binding sites for other proteins, including transcription coregulators.

[0053] As used herein, the term “transcription factor” refers exclusively to those factors having one or more DBDs and is not intended to include other regulatory proteins such as coactivators, chromatin remodelers, histone acetylases, deacetylases, kinases, and methylases, which do not contain DBDs.

[0054] Of the approximately 2,600 human proteins that contain DNA-binding domains, the majority are believed to be transcription factors. Transcription factors are categorized according to structural features of the DNA-binding domain, which include basic helix-loop-helix domains, basic-leucine zipper (bZIP domains), C-terminal effector domains of bipartite response regulators, GCC box domains, helix-turn-helix domains, homeodomains, lambda repressor-like domains, serum response factor-like (srf-like) domains, paired box domains, winged helix

domains, zinc finger domains, multi-Cys₂His₂ zinc finger domains, Zn₂Cys₆ domains, and Zn₂Cys₈ nuclear receptor zinc finger domains.

[0055] Many transcription factors are either tumor suppressors or oncogenes, and, thus, mutations within and the aberrant expression of such transcription factors is associated with some cancers and other diseases and conditions. For example, transcription factors within (1) the NF-kappaB family, (2) the AP-1 family, (3) the STAT family, and (4) the steroid receptor family have been implicated in the neurodevelopmental disorder Rett syndrome (the MECP2 transcription factor), diabetes (hepatocyte nuclear factors (HNFs) and insulin promoter factor-1 (IPF1/Pdx1)), developmental verbal dyspraxia (the FOXP2 transcription factor), autoimmune diseases (the FOXP3 transcription factor), Li-Raumeni syndrome (the p53 tumor suppressor), and multiple cancers (the STAT and HOX family of transcription factors). Clevenger, *Am. J. Pathol.* 165(5):1449-60 (2004); Carrithers *et al.*, *Am J Pathol* 166(1):185- 196 (2005); Herreros-Villanueva *et al.*, *World J Gastroenterology* 20(9):2247-2254 (2014); and Campbell *et al.*, *Am J Pathol* 158(1):25-32 (2001). Olsson *et al.*, *Oncogene* 26(7):1028-37 (2007) describe the upregulation of the transcription factor E2F3, which is a key regulator of the cell cycle, in human bladder and prostate cancers. Cantile *et al.*, *Curr Med Chem* 18(32):4872-84 (2011) describe the upregulation of HOX genes in urogenital cancers; Cillo *et al.*, *Int J. Cancer* 129(11):2577-87 (2011) describe the upregulation of HOX genes in hepatocellular carcinoma; Cantile *et al.*, *Int J. Cancer* 125(7):1532-41 (2009) describe HOX D13 expression across 79 tumor tissue types; Cantile *et al.*, *J Cell Physiol* 205(2):202-10 (2005) describe upregulation of HOX D expression in prostate cancers; Cantile *et al.*, *Oncogene* 22(41):6462-8 (2003) describe the hyperexpression of locus C genes in the HOX network in human bladder transitional cell carcinomas; Morgan *et al.*, *BioMed Central* 14:15 (2014), describe HOX transcription factors as targets for prostate cancer; and Alharbi *et al.*, *Leukemia* 27(5):1000-8 (2013) describe the role of HOXC genes in hematopoiesis and acute leukemia.

[0056] The AP-2 family includes five transcription factors that can act as both repressors and activators. AP-2 γ regulates cancer cell survival by blocking p53 activation of the p21CIP gene. High levels of AP-2 γ are associated with poor prognosis in breast cancer. Gee *et al.*, *J Pathol* 217(1):32-41 (2009) and Williams *et al.*, *EMBO J* 28(22):3591-601 (2009). A further transcription factor that promotes cell survival are the forkhead transcription factors (FOX), which can promote the expression of proteins involved in drug resistance and also block programmed cell death and may therefore protect cancer cells from chemotherapeutic drugs.

Gomes *et al.*, *Chin J. Cancer* 32(7):365-70 (2013) describe the role of FOXO3a and FOXM1 in carcinogenesis and drug resistance.

[0057] Transcription factors can bind to promoters as well as to enhancers. As used in the present disclosure, the term transcription factor refers to the subset of transcription factors that bind to transcription factor binding sites within a promoter and excludes those factors that bind to enhancer sequences. Transcription factors can also upregulate or downregulate the expression of an associated nucleic acid. The present disclosure employs transcriptional promoters having transcription factor binding sites for transcription factors that promote rather than inhibit expression and therefore cause the upregulation in the expression of an associated nucleic acid. Such transcription factors that upregulate nucleic acid expression include, for example and not limitation, transcription factors that (1) stabilize RNA polymerase binding to its cognate binding site, (2) recruit coactivator or corepressor proteins to a transcription factor DNA complex, and/or (3) catalyze the acetylation of histone proteins (or recruit one or more other proteins that catalyze the acetylation of histone proteins). Such histone acetyltransferase (HAT) activity reduces the affinity of histone binding to DNA thereby making the DNA more accessible for transcription.

[0058] As used herein, the term “necrosis” refers to a process leading to cell death that occurs when a cell is damaged by an external force, such as poison, a bodily injury, an infection, or loss of blood supply. Cell death from necrosis causes inflammation that can result in further distress or injury within the body. As used herein, the term “apoptosis” refers to a process leading to cell death in which a programmed sequence of events leads to the elimination of cells without releasing harmful substances. Apoptosis plays a crucial role in developing and maintaining the health of the body by eliminating old cells, unnecessary cells, and unhealthy cells. Apoptosis is mediated by proteins produced by suicide genes, including the caspase proteins, which break down cellular components needed for survival and induce the production of DNAses, which destroy nuclear DNA.

[0059] As used herein, the term “suicide gene” refers to a class of genes that produce proteins that induce p53-mediated apoptotic cell killing. Suicide genes that can be employed in the expression constructs and systems of the present disclosure include the caspases, Casp3, Casp8, Casp9, BAX, DFF40, Herpes Simplex Virus Thymidine Kinase (HSV-TK), and cytosine deaminase and inducible variants of Casp3, Casp8, Casp9, BAX, DFF40, Herpes HSV-TK, and cytosine deaminase.

[0060] The presently disclosed expression constructs and systems are used in methods for the treatment of aging, cancer infectious disease, bacterial infections, and/or other conditions as well as in methods for the killing of cells that are associated with aging, cancer, infectious disease, bacterial infections, and/or other conditions and employ a therapeutic protein that reduces the growth and/or proliferation of a target cell. In certain embodiments, the therapeutic protein can be expressed by a suicide gene, which encodes Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase as well as a inducible variants of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase. The expression cassettes and systems can also be used in conjunction with conventional chemotherapeutics to enhance the effectiveness of therapeutic regimen for the treatment of aging, cancers, infectious diseases, bacterial infections, and other diseases and conditions.

[0061] Within certain aspects of the present disclosure, expression constructs are pVAX1 (**FIG. 14**) plasmid expression constructs comprising a polynucleotide encoding a pro-apoptotic protein under the regulatory control of a target cell-specific promoter, such as a senescent cell-specific promoter or a cancer cell-specific promoter.

[0062] Exemplary pVAX1TM plasmid expression constructs include pVAX-16s-iCasp9-MX (**FIG. 16**; SEQ ID NO: 6) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) under the regulatory control of a p16s promoter, pVAX1-53-iCasp9-MX (**FIG. 26**; SEQ ID NO: 7) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) under the regulatory control of a p53 promoter, pVax1-p53-saCasp9-5 (**FIG. 27**; SEQ ID NO: 8) for the target cell-specific expression of a self-activating Caspase 9 protein (saCASP9) under under the regulatory control of a p53 promoter, pVax1-p53-iCasp9-OVA (**FIG. 28**; SEQ ID NO: 11) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) and an ovalbumin protein under the regulatory control of a p53 promoter, pVax1-p53-iCasp9-G-O (**FIG. 29**; SEQ ID NO: 9) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) and an ovalbumin protein under the regulatory control of a p53 promoter, pVax1-p53-iCasp9-huCD40L (**FIG. 30**; SEQ ID NO: 10) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) and a CD40 ligand protein (CD40L) under the regulatory control of a p53 promoter.

[0063] Exemplary p10 plasmid expression constructs include p10-p16e-iCasp9 (**FIG. 17**; SEQ ID NO: 12) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) under the regulatory control of a p16e promoter, p10-p16e-saCasp9 (**FIG. 18**; SEQ ID NO: 13)

for the target cell-specific expression of a self-activating Caspase 9 protein (saCasp9) under the regulatory control of a p16e promoter, p10-p53-iCasp9 (FIG. 33; SEQ ID NO: 14) for the target cell-specific expression of an inducible Caspase 9 protein (iCasp9) under the regulatory control of a p53 promoter, and p10-p53-saCasp9 (FIG. 34; SEQ ID NO: 15) for the target cell-specific expression of a self-activating Caspase 9 protein (saCasp9) under the regulatory control of a p53 promoter.

[0064] Within other aspects of the present disclosure, expression constructs are NTC-based plasmid expression constructs, including NTC8385, NTC8685, and NTC9385 plasmid expression constructs, comprising a polynucleotide encoding a pro-apoptotic protein under the regulatory control of a target cell-specific promoter, such as a senescent cell-specific promoter or a cancer cell-specific promoter.

[0065] Within further aspects of the present disclosure, expression constructs are gWiz-based plasmid expression constructs comprising a polynucleotide encoding a pro-apoptotic protein under the regulatory control of a target cell-specific promoter, such as a senescent cell-specific promoter or a cancer cell-specific promoter.

[0066] The practice of the present disclosure will employ, unless indicated specifically to the contrary, conventional methodology and techniques that are in common use in the fields of virology, oncology, immunology, microbiology, molecular biology, and recombinant DNA, which methodology and techniques are well known by and readily available to those having skill of the art. Such methodology and techniques are explained fully in laboratory manuals as well as the scientific and patent literature. See, *e.g.*, Sambrook, *et al.*, "Molecular Cloning: A Laboratory Manual" (2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989); Maniatis *et al.*, "Molecular Cloning: A Laboratory Manual" (1982); "DNA Cloning: A Practical Approach, vol. I & II" (Glover, ed.); "Oligonucleotide Synthesis" (Gait, ed., 1984); Ausubel *et al.* (eds.), "Current Protocols in Molecular Biology" (John Wiley & Sons, 1994); "Nucleic Acid Hybridization" (Hames & Higgins, eds., 1985); "Transcription and Translation" (Hames & Higgins, eds., 1984); "Animal Cell Culture" (Freshney, ed., 1986); and Perbal, "A Practical Guide to Molecular Cloning" (1984). All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

*Systems and Expression Constructs for Reducing, Preventing, and/or
Eliminating the Growth and/or Survival of a Target cell*

[0067] Within certain embodiments, the present disclosure provides expression constructs and systems comprising a delivery vector and an expression construct for achieving a target cell specific reduction, prevention, and/or elimination in the growth and/or survival of the target cell.

Systems

[0068] Systems of the present disclosure comprise (1) a vector that is capable of non-specific delivery of a nucleic acid to a cell, whether that cell is a target cell or a non-target cell, and (b) an expression construct comprising a target cell specific transcriptional promoter and a nucleic acid that encodes a therapeutic protein, which expression constructs achieve the target cell specific production of a therapeutic protein. The systems disclosed herein will find utility in a broad range of therapeutic applications in which it is desirable to effectuate the growth or survival characteristics of a target cell, such as a cell that is associated with aging, disease, or another condition, but, at the same time, to not effectuate the growth or survival characteristics of a normal, a non-target cell that is not associated with aging, disease, or another condition.

[0069] The present disclosure provides systems for effectuating the growth and/or survival of a broad range of cells that are associated with aging, disease, or other conditions that similarly comprises (1) a non-specific nucleic acid delivery vector and (2) an expression construct comprising (a) a target cell specific transcriptional promoter and (b) a nucleic acid that encodes a therapeutic protein. Each of these aspects of the presently disclosed systems are described in further detail herein.

[0070] Within certain embodiments, provided herein are systems for effectuating the growth and/or survival of target cells, which systems comprise: (1) a non-specific nucleic acid delivery vector and (2) an expression construct comprising: (a) a transcriptional promoter, which transcriptional promoter is activated in target cells but not in normal, non-target cells, and (b) a nucleic acid that is under the control of the transcriptional promoter, which nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of a target cell, for example by inducing a mechanism of programmed cell death in a cell in which it is produced. Thus, these systems achieve the selective killing of target cells by exploiting transcriptional machinery that is produced in, and intrinsic to, target cells; without the use of toxins and in the absence of target cell specific delivery of the expression construct.

[0071] In certain aspects of these embodiments wherein the human target cell is a senescent cell, the transcriptional promoter can include at least a transcription factor binding site (*i.e.*, a

response element) of p16INK4a/CDKN2A as described in Wang *et al.*, *J. Biol. Chem.* 276(52):48655-61 (2001), which transcriptional promoter is responsive to activation by a factor such as SP1, ETS1, and ETS2. The transcriptional promoter can also include at least a transcription factor binding site (*i.e.*, a response element) of p21/CDKN1A, which transcriptional promoter is responsive to activation by a factor such as p53/TP53. Transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein such as Casp3, Casp8, Casp9, DFF40, BAX, HSV-TK, or carbonic anhydrase or an inducible variant of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase.

[0072] In other aspects of these embodiments wherein the human target cell is a cancer cell, such as a brain cancer cell, a prostate cancer cell, a lung cancer cell, a colorectal cancer cell, a breast cancer cell, a liver cancer cell, a hematologic cancer cell, and a bone cancer cell, the transcriptional promoter can include at least a transcription factor binding site (*i.e.*, a response element) of the p21^{cip1/waf1} promoter, the p27^{kip1} promoter, the p53^{kip2} promoter, the TdT promoter, the Rag-1 promoter, the B29 promoter, the Blk promoter, the CD19 promoter, the BLNK promoter, and/or the $\lambda 5$ promoter, which transcriptional promoter is responsive to activation by one or more transcription factors such as an EBF3, O/E-1, Pax-5, E2A, p53, VP16, MLL, HSF1, NF-IL6, NFAT1, AP-1, AP-2, HOX, E2F3, and/or NF- κ B transcription factor, and which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein such as, for example, Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase or an inducible variant of Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase which therapeutic protein reduces, prevents, and/or eliminates the growth and/or survival of the cancer cell, such as, for example, by inducing cell death in the senescent cell via a cellular process including apoptosis. Other therapeutic proteins may be employed that reduce, prevent, and/or eliminate the growth and/or survival of a cancer cell by, for example, inducing cell death via a cellular process including necrosis/necroptosis, autophagic cell death, endoplasmic reticulum-stress associated cytotoxicity, mitotic catastrophe, paraptosis, pyroptosis, pyronecrosis, and entosis. In still further aspects of these embodiments wherein the target cell is a human cell that is infected with an infectious agent, such as a virus, including, for example, a herpes virus, a polio virus, a hepatitis virus, a retrovirus, an influenza virus, and a rhino virus, or the target cell is a bacterial cell, the transcriptional promoter can be activated by a factor that is expressed by the infectious agent or bacterial cell, which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein such as, for example, Casp3, Casp8, Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase or an inducible variant of Casp3, Casp8,

Casp9, BAX, DFF40, HSV-TK, or cytosine deaminase which therapeutic protein reduces, prevents, and/or eliminates the growth and/or survival of the senescent cell, such as, for example, by inducing cell death in the senescent cell via a cellular process including apoptosis. Other therapeutic proteins may be employed that reduce, prevent, and/or eliminate the growth and/or survival of a senescent cell by, for example, inducing cell death via a cellular process including necrosis/necroptosis, autophagic cell death, endoplasmic reticulum-stress associated cytotoxicity, mitotic catastrophe, paraptosis, pyroptosis, pyronecrosis, and entosis.

[0073] Each of these aspects of the presently disclosed systems are described in further detail herein.

1. Non-specific Nucleic Acid Delivery Vectors

[0074] The systems of the present disclosure achieve target cell specificity by exploiting transcriptional machinery that is unique to a target cell. Thus, the systems described herein employ nucleic acid delivery vectors that can be readily adapted for the non-specific delivery of expression constructs to a cell, including but not limited to a target cell.

[0075] A wide variety of both non-viral and viral nucleic acid delivery vectors are well known and readily available in the art and may be adapted for use for the non-specific cellular delivery of the expression constructs disclosed herein. See, for example, Elsabahy *et al.*, *Current Drug Delivery* **8(3)**:235-244 (2011) for a general description of viral and non-viral nucleic acid delivery methodologies. The successful delivery of a nucleic acid into mammalian cells relies on the use of efficient delivery vectors. Viral vectors exhibit desirable levels of delivery efficiency, but often also exhibit undesirable immunogenicity, inflammatory reactions, and problems associated with scale-up, all of which can limit their clinical use. The ideal vectors for the delivery of a nucleic acid are safe, yet ensure nucleic acid stability and the efficient transfer of the nucleic acid to the appropriate cellular compartments.

[0076] Non-limiting examples of non-viral and viral nucleic acid delivery vectors are described herein and disclosed in scientific and patent literature. More specifically, the presently disclosed systems may employ one or more liposomal vectors, viral vectors, nanoparticles, polyplexes, dendrimers, each of which has been developed for the non-specific delivery of nucleic acids, can be adapted for the non-specific delivery of the expression constructs described herein, and can be modified to incorporate one or more agents for promoting the targeted delivery of a

system to a target cell of interest thereby enhancing the target cell specificity of the presently disclosed systems.

2. Liposomal Vectors and Nanoparticles

[0077] An expression cassette may be incorporated within and/or associated with a lipid membrane, a lipid bi-layer, and/or a lipid complex such as, for example, a liposome, a vesicle, a micelle and/or a microsphere. Suitable methodology for preparing lipid-based delivery systems that may be employed with the expression constructs of the present disclosure are described in Metselaar *et al.*, *Mini Rev. Med. Chem.* 2(4):319-29 (2002); O'Hagen *et al.*, *Expert Rev. Vaccines* 2(2):269-83 (2003); O'Hagan, *Curr. Drug Targets Infect. Disord.* 1(3):273-86 (2001); Zho *et al.*, *Biosci Rep.* 22(2):355-69 (2002); Chikh *et al.*, *Biosci Rep.* 22(2):339-53 (2002); Bungener *et al.*, *Biosci. Rep.* 22(2):323-38 (2002); Park, *Biosci Rep.* 22(2):267-81 (2002); Ulrich, *Biosci. Rep.* 22(2):129-50; Lofthouse, *Adv. Drug Deliv. Rev.* 54(6):863-70 (2002); Zhou *et al.*, *J. Immunother.* 25(4):289-303 (2002); Singh *et al.*, *Pharm Res.* 19(6):715-28 (2002); Wong *et al.*, *Curr. Med. Chem.* 8(9):1123-36 (2001); and Zhou *et al.*, *Immunomethods* 4(3):229-35 (1994). Midoux *et al.*, *British J. Pharmacol* 157:166-178 (2009) describe chemical vectors for the delivery of nucleic acids including polymers, peptides and lipids. Sioud and Sorensen, *Biochem Biophys Res Commun* 312(4):1220-5 (2003) describe cationic liposomes for the delivery of nucleic acids.

[0078] Due to their positive charge, cationic lipids have been employed for condensing negatively charged DNA molecules and to facilitate the encapsulation of DNA into liposomes. Cationic lipids also provide a high degree of stability to liposomes. Cationic liposomes interact with a cell membrane and are taken up by a cell through the process of endocytosis. Endosomes formed as the results of endocytosis, are broken down in the cytoplasm thereby releasing the cargo nucleic acid. Because of the inherent stability of cationic liposomes, however, transfection efficiencies can be low as a result of lysosomal degradation of the cargo nucleic acid.

[0079] Helper lipids (such as the electroneutral lipid DOPE and L- α -dioleoyl phosphatidyl choline (DOPC)) can be employed in combination with cationic lipids to form liposomes having decreased stability and, therefore, that exhibit improved transfection efficiencies. These electroneutral lipids are referred to as Fusogenic lipids. See, Gruner *et al.*, *Biochemistry* 27(8):2853-66 (1988) and Farhood *et al.*, *Biochim Biophys Acta* 1235(2):289-95 (1995). DOPE forms an HII phase structure that induces supramolecular arrangements leading to the fusion of

a lipid bilayer at a temperature greater than 5°C to 10°C. The incorporation of DOPE into liposomes also helps the formation of HII phases that destabilize endosomal membranes.

[0080] Cholesterol can be employed in combination with DOPE liposomes for applications in which a liposomal vector is administered intravenously. Sakurai *et al.*, *Eur J Pharm Biopharm* 52(2):165-72 (2001). The presence of one unsaturation in the acyl chain of DOPE is a crucial factor for membrane fusion activity. Talbot *et al.*, *Biochemistry* 36(19):5827-36 (1997).

[0081] Fluorinated helper lipids having saturated chains, such as DF4C11PE (rac-2,3- Di[11-(F-butyl)undecanoyl] glycerol-1-phosphoethanolamine) also enhance the transfection efficiency of lipopolyamine liposomes. Boussif *et al.*, *J Gene Med* 3(2):109-14 (2001); Gaucheron *et al.*, *Bioconj Chem* 12(6):949-63 (2001); and Gaucheron *et al.*, *J Gene Med* 3(4):338-44 (2001).

[0082] The helper lipid 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP) enhances efficient of in vitro cell transfection as compared to DOPE lipoplexes. Prata *et al.*, *Chem Commun* 13:1566-8 (2008). Replacement of the double bond of the oleic chains of DOPE with a triple bond as in Distear-4-ynoyl L- α -phosphatidylethanolamine [DS(9-yne)PE] has also been shown to produce more stable lipoplexes. Fletcher *et al.*, *Org Biomol Chem* 4(2):196-9 (2006).

[0083] Amphiphilic anionic peptides that are derived from the N-terminal segment of the HA-2 subunit of influenza virus haemagglutinin, such as the IFN7 (GLFEAIEGFIE NGWEGMIDGW YG) and E5CA (GLFEAIAEFI EGGWEGLIEG CA) peptides, can be used to increase the transfection efficiency of liposomes by several orders of magnitude. Wagner *et al.*, *Proc Natl Acad Sci U.S.A.* 89(17):7934-8 (1992); Midoux *et al.*, *Nucl Acids Res.* 21(4):871-8 (1993); Kichler *et al.*, *Bioconj Chem* 8(2):213-21 (1997); Wagner, *Adv Drug Deliv Rev* 38(3):279-289 (1999); Zhang *et al.*, *J Gene Med* 3(6):560-8 (2001). Some artificial peptides such as GALA have been also used as fusogenic peptides. See, for example, Li *et al.*, *Adv Drug Deliv Rev* 56(7):967-85 (2004) and Sasaki *et al.*, *Anal Bioanal Chem* 391(8):2717-27 (2008). The fusogenic peptide of the glycoprotein H from herpes simplex virus improves the endosomal release of DNA/Lipofectamine lipoplexes and transgene expression in human cell (Tu and Kim, *J Gene Med* 10(6):646-54 (2008).

[0084] PCT Patent Publication Nos. WO 1999024582A1 and WO 2002/044206 describe a class of proteins derived from the family Reoviridae that promote membrane fusion. These proteins are exemplified by the p14 protein from reptilian reovirus and the p16 protein from aquareovirus. PCT Patent Publication No. WO 2012/040825 describes recombinant

polypeptides for facilitating membrane fusion, which polypeptides have at least 80% sequence identity with the ectodomain of p14 fusion-associated small transmembrane (FAST) protein and having a functional myristoylation motif, a transmembrane domain from a FAST protein and a sequence with at least 80% sequence identity with the endodomain of p15 FAST protein. The '825 PCT further describes the addition of a targeting ligand to the recombinant polypeptide for selective fusion. The recombinant polypeptides presented in the '825 PCT can be incorporated within the membrane of a liposome to facilitate the delivery of nucleic acids. Fusogenic liposomes for delivering therapeutic compounds, including nucleic acids, to the cytoplasm of a mammalian cell, which reduce liposome disruption and consequent systemic dispersion of the cargo nucleic acid and/or uptake into endosomes and resulting nucleic acid destruction are available commercially from Innovascreen Inc. (Halifax, Nova Scotia, CA).

[0085] A wide variety of inorganic nanoparticles, including gold, silica, iron oxide, titanium, hydrogels, and calcium phosphates have been described for the delivery of nucleic acids and can be adapted for the delivery of the expression constructs described herein. See, for example Wagner and Bhaduri, *Tissue Engineering* 18(1):1-14 (2012) (describing inorganic nanoparticles for delivery of nucleic acid sequences); Ding *et al.*, *Mol Ther* e-pub (2014) (describing gold nanoparticles for nucleic acid delivery); Zhang *et al.*, *Langmuir* 30(3):839-45 (2014) (describing titanium dioxide nanoparticles for delivery of DNA oligonucleotides); Xie *et al.*, *Curr Pharm Biotechnol* 14(10):918-25 (2014) (describing biodegradable calcium phosphate nanoparticles for gene delivery); Sizovs *et al.*, *J Am Chem Soc* 136(1):234-40 (2014) (describing sub-30 nm monodisperse oligonucleotide nanoparticles).

[0086] Among the advantages of inorganic vectors are their storage stability, low immunogenicity, and resistance to microbial attack. Nanoparticles of less than 100 nm can efficiently trap nucleic acids and allow its escape from endosomes without degradation. Inorganic nanoparticles exhibit improved *in vitro* transfection for attached cell lines due to their high density and preferential location on the base of the culture dish. Quantum dots have been described that permit the coupling of nucleic acid delivery with stable fluorescence markers.

[0087] Hydrogel nanoparticles of defined dimensions and compositions, can be prepared via a particle molding process referred to as PRINT (Particle Replication in Non-wetting Templates), and can be used as delivery vectors for the expression constructs disclosed herein. Nucleic acids can be encapsulated in particles through electrostatic association and physical entrapment. To prevent the disassociation of cargo nucleic acids from nanoparticles following

systemic administration, a polymerizable conjugate with a degradable, disulfide linkage can be employed.

[0088] The PRINT technique permits the generation of engineered nanoparticles having precisely controlled properties including size, shape, modulus, chemical composition and surface functionality for enhancing the targeting of the expression cassette to a target cell. See, *e.g.*, Wang *et al.*, *J Am Chem Soc* 132:11306-11313 (2010); Enlow *et al.*, *Nano Lett* 11:808-813 (2011); Gratton *et al.*, *Proc Natl Acad Sci USA* 105:11613-11618 (2008); Kelly, *J Am Chem Soc* 130:5438-5439 (2008); Merkel *et al.* *Proc Natl Acad Sci USA* 108:586-591 (2011). PRINT is also amenable to continuous roll-to-roll fabrication techniques that permit the scale-up of particle fabrication under good manufacturing practice (GMP) conditions.

[0089] Nanoparticles can be encapsulated with a lipid coating to improve oral bioavailability, minimize enzymatic degradation and cross blood brain barrier. The nanoparticle surface can also be PEGylated to improve water solubility, circulation *in vivo*, and stealth properties.

3. Viral Vectors

[0090] A wide variety of viral vectors are well known by and readily available to those of skill in the art, including, for example, herpes simplex viral vectors, lentiviral vectors, adenoviral vectors, and adeno-associated viral vectors, which viral vectors can be adapted for use in the systems disclosed herein for the delivery of nucleic acids, in particular nucleic acids comprising an expression cassette for the target cell specific expression of a therapeutic protein.

[0091] The tropisms of natural or engineered viruses towards specific receptors are the foundations for constructing viral vectors for delivery of nucleic acids. The attachment of these vectors to a target cell is contingent upon the recognition of specific receptors on a cell surface by a ligand on the viral vector. Viruses presenting very specific ligands on their surfaces anchor onto the specific receptors on a cell. Viruses can be engineered to display ligands for receptors presented on the surface of a target cell of interest. The interactions between cell receptors and viral ligands are modulated *in vivo* by toll like receptors.

[0092] The entry of a viral vector into a cell, whether via receptor mediated endocytosis or membrane fusion, requires a specific set of domains that permit the escape of the viral vector from endosomal and/or lysosomal pathways. Other domains facilitate entry into nuclei. Replication, assembly, and latency determine the dynamics of interactions between the vector

and the cell and are important considerations in the choice of a viral vector, as well as in engineering therapeutic cargo carrying cells, in designing cancer suicide gene therapies.

[0093] Herpes simplex virus (HSV) belongs to a family of herpesviridae, which are enveloped DNA viruses. HSV binds to cell receptors through orthologs of their three main ligand glycoproteins: gB, gH, and gL, and sometimes employ accessory proteins. These ligands play decisive roles in the primary routes of virus entry into oral, ocular, and genital forms of the disease. HSV possesses high tropism towards cell receptors of the nervous system, which can be utilized for engineering recombinant viruses for the delivery of expression cassettes to target cells, including senescent cells, cancer cells, and cells infected with an infectious agent. Therapeutic bystander effects are enhanced by inclusion of connexin coding sequences into the constructs. Herpes Simplex Virus vectors for the delivery of nucleic acids to target cells have been reviewed in Anesti and Coffin, *Expert Opin Biol Ther* 10(1):89-103 (2010); Marconi *et al.*, *Adv Exp Med Biol* 655:118-44 (2009); and Kasai and Saeki, *Curr Gene Ther* 6(3):303-14 (2006).

[0094] Lentivirus belongs to a family of retroviridae, which are enveloped, single stranded RNA retroviruses and include the Human immunodeficiency virus (HIV). HIV envelope protein binds CD4, which is present on the cells of the human immune system such as CD4⁺ T cells, macrophages, and dendritic cells. Upon entry into a cell, the viral RNA genome is reverse transcribed into double-stranded DNA, which is imported into the cell nucleus and integrated into the cellular DNA. HIV vectors have been used to deliver the therapeutic genes to leukemia cells. Recombinant lentiviruses have been described for mucin-mediated delivery of nucleic acids into pancreatic cancer cells, to epithelial ovarian carcinoma cells, and to glioma cells, without substantial non-specific delivery to normal cells. Lentiviral vectors for the delivery of nucleic acids to target cells have been reviewed in Primo *et al.*, *Exp Dermatol* 21(3):162-70 (2012); Staunstrup and Mikkelsen, *Curr Gene Ther* 11(5):350-62 (2011); and Dreyer, *Mol Biotechnol* 47(2):169-87 (2011).

[0095] Adenovirus is a non-enveloped virus consisting of a double-stranded, linear DNA genome and a capsid. Naturally, adenovirus resides in adenoids and may be a cause of upper respiratory tract infections. Adenovirus utilizes a cell's coxsackie virus and adenovirus receptor (CAR) for the adenoviral fiber protein for entry into nasal, tracheal, and pulmonary epithelia. CARs are expressed at low levels on senescent and cancer cells. Recombinant adenovirus can be generated that are capable of nucleic acid deliver to target cells. Replication-competent

adenovirus-mediated suicide gene therapy (ReCAP) is in the clinical trials for newly-diagnosed prostate cancer. Adenoviral vectors for the delivery of nucleic acids to target cells have been reviewed in Huang and Kamihira, *Biotechnol Adv.* 31(2):208-23 (2013); Alemany, *Adv Cancer Res* 115:93-114 (2012); Kaufmann and Nettelbeck, *Trends Mol Med* 18(7):365-76 (2012); and Mowa *et al.*, *Expert Opin Drug Deliv* 7(12):1373-85 (2010).

[0096] Adeno-associated virus (AAV) is a small virus that infects humans and some other primate species. AAV is not currently known to cause disease and consequently the virus causes a very mild immune response. Vectors using AAV can infect both dividing and quiescent cells and persist in an extrachromosomal state without integrating into the genome of the host cell. These features make AAV a very attractive candidate for creating viral vectors for use in the systems of the present disclosure. Adeno-associated virus (AAV) vectors for the delivery of nucleic acids to target cells have been reviewed in Li *et al.*, *J. Control Release* 172(2):589-600 (2013); Hajitou, *Adv Genet* 69:65-82 (2010); McCarty, *Mol Ther* 16(10):1648-56 (2008); and Grimm *et al.*, *Methods Enzymol* 392:381-405 (2005).

4. Polyplexes

[0097] Polyplexes are complexes of polymers with DNA. Polyplexes consist of cationic polymers and their fabrication is based on self-assembly by ionic interactions. One important difference between the methods of action of polyplexes and liposomes and lipoplexes is that polyplexes cannot directly release their nucleic acid cargo into the cytoplasm of a target cell. As a result co-transfection with endosome-lytic agents such as inactivated adenovirus is required to facilitate escape from the endocytic vesicle made during particle uptake. better understanding of the mechanisms by which DNA can escape from endolysosomal pathway (*i.e.*, the proton sponge effect) has triggered new polymer synthesis strategies such as the incorporation of protonable residues in polymer backbone and has revitalized research on polycation-based systems. See, *e.g.*, Parhamifar *et al.*, *Methods* e-pub (2014); Rychgak and Kilbanov, *Adv Drug Deliv Rev* e-pub (2014); Jafari *et al.*, *Curr Med Chem* 19(2):197-208 (2012).

[0098] Due to their low toxicity, high loading capacity, and ease of fabrication, polycationic nanocarriers exhibit substantial advantages over viral vectors, which show high immunogenicity and potential carcinogenicity and lipid-based vectors which cause dose dependent toxicity. Polyethyleneimine, chitosan, poly(beta-amino esters), and polyphosphoramidate have been described for the delivery of nucleic acids. See, *e.g.*, Buschmann *et al.*, *Adv Drug Deliv Rev*

65(9):1234-70 (2013). The size, shape, and surface chemistry of these polymeric nano-carriers can be easily manipulated.

5. Dendrimers

[0099] Dendrimers are highly branched macromolecules having a spherical shape. The surface of dendrimer particles may be functionalized such as, for example, with positive surface charges (cationic dendrimers), which may be employed for the delivery of nucleic acids. Dendrimer-nucleic acid complexes are taken into a cell via endocytosis. Dendrimers offer robust covalent construction and extreme control over molecule structure and size. Dendrimers are available commercially from Dendritic Nanotechnologies Inc. (Priostar; Mt Pleasant, MI), who produce dendrimers using kinetically driven chemistry, which can be adapted for the delivery of nucleic acids and can transfect cells at a high efficiency with low toxicity.

[00100] It will be understood that, while targeted delivery of an expression construct is not required by the systems of the present disclosure and that the targeted reduction, prevention, and/or elimination in the growth and/or survival of a target cell may be achieved by exploiting the intracellular transcriptional machinery of a target cell that is unique to the target cell, it may be desirable, depending upon the precise application contemplated, to incorporate into an otherwise non-specific delivery vector one or more components that facilitate the targeted delivery to a subset of cells at least some of which include a target cell that is susceptible to the growth and/or survival inhibition by the expression constructs of the present disclosure.

[00101] The targeted delivery of nucleic acids by liposome, nanoparticle, viral and other vectors described herein has been described in the scientific and patent literature and is well known by and readily available to those of skill in the art. Such targeted delivery technologies may, therefore, be suitably adapted for targeting the delivery of expression constructs of the present disclosure to enhance the specificity of the growth and/or survival reduction, prevention, and/or elimination that is achieved within a target cell. The following examples of targeted delivery technologies are provided herein to exemplify, not to limit, the targeted delivery vectors that may be adapted to achieve the systems of the present disclosure.

Expression Constructs

[00102] Expression constructs of the present disclosure comprise: (a) a transcriptional promoter that is responsive to a factor or factors that are produced in a target cell, one or more of which factors is not produced, is produced at a substantially reduced level, is inactive, and/or exhibits

a substantially reduced activity in a non-target cell; and (b) a nucleic acid that is operably linked to and under the regulatory control of the transcriptional promoter, wherein the nucleic acid encodes a protein that is capable of reducing, preventing, and/or eliminating the growth and/or survival of a cell in which it is produced, including a target cell.

1. Target Cell Specific Transcriptional Promoters

[00103] The present disclosure provides systems comprising a vector for delivering a nucleic acid to a cell wherein the nucleic acid is under the transcriptional control of a promoter that is derepressed or activated in a target cell, but is repressed or inactivated in a normal cell, non-target cell.

[00104] It will be understood the specificity of the presently disclosed systems toward a target cell is achieved, therefore, through the target cell-specific transcriptional activation of a nucleic acid that encodes a protein that reduces, prevents, and/or eliminates the growth and/or survival of a cell without regard to whether that cell is a target cell. Thus, the target cell specificity of the presently-disclosed systems derives from the transcriptional promoter that regulates the expression of the nucleic acid within the expression cassette in conjunction with transcription-regulatory machinery that is provided by, and unique to, the target cell.

[00105] Thus, transcriptional promoters that may be suitably employed in the expression constructs, systems, and methods of the present disclosure include those transcriptional promoters that are capable of promoting the expression of a nucleic acid in a target cell (*i.e.*, a cell that is associated with aging, disease, or other condition), but incapable of, or exhibit a substantially reduced capability of, promoting expression of that nucleic acid in a non-target cell.

[00106] Exemplified herein are expression constructs and systems comprising expression constructs wherein the transcriptional promoter is activated in a target cell that is associated with aging, disease, or another condition.

[00107] In some embodiments, the present disclosure provides expression constructs and systems that may be employed in methods for the treatment of aging reducing, preventing, and/or eliminating the growth and/or survival of a cell, such as a senescent cell, which is associated with aging. In certain aspects of those embodiments, expression constructs employ a transcriptional promoter that is responsive to one or more factors that are produced within a target cell, such as a senescent cell, but are not produced in a non-target cell wherein those one or more

factors derepress and/or activate the transcriptional promoter and, as a consequence, promote the expression of a nucleic acid encoding a therapeutic protein that reduces, prevents, and/or eliminates the growth and/or survival of a cell that is associated with aging, including a senescent cell.

[00108] The transcriptional promoter itself is the primary mechanism by which senescent cells are preferentially targeted by the systems described in this disclosure. A prototypic example of a target specific transcriptional promoter for use with the systems in this disclosure is a promoter that is only active or mostly active in senescent cells. A number of promoters known by artisans to be active in senescent cells may be used with this system.

[00109] In certain aspects of these embodiments wherein the human target cell is a senescent cell, the transcriptional promoter can include the promoter region of p16INK4a/CDKN2A as described in Wang *et al.*, *J. Biol. Chem.* 276(52):48655-61 (2001), which transcriptional promoter is responsive to activation by a factor such as SP1, ETS1, and ETS2. The transcriptional promoter can also include the promoter region of p21/CDKN1A, which transcriptional promoter is responsive to activation by a factor such as p53/TP53.

[00110] In other aspects of these embodiments wherein the human target cell is a cancer cell, such as a brain cancer cell, a prostate cancer cell, a lung cancer cell, a colorectal cancer cell, a breast cancer cell, a liver cancer cell, a hematologic cancer cell, and a bone cancer cell, the transcriptional promoter can include the p21^{cip1/waf1} promoter, the p27^{kip1} promoter, the p57^{kip2} promoter, the TdT promoter, the Rag-1 promoter, the B29 promoter, the Blk promoter, the CD19 promoter, the BLNK promoter, and/or the $\lambda 5$ promoter, which transcriptional promoter is responsive to activation by one or more transcription factors such as an EBF3, O/E-1, Pax-5, E2A, p53, VP16, MLL, HSF1, NF-IL6, NFAT1, AP-1, AP-2, HOX, E2F3, and/or NF- κ B transcription factor, and which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein.

[00111] In still further aspects of these embodiments wherein the target cell is a human cell that is infected with an infectious agent, such as a virus, including, for example, a herpes virus, a polio virus, a hepatitis virus, a retrovirus virus, an influenza virus, and a rhino virus, or the target cell is a bacterial cell, the transcriptional promoter can be activated by a factor that is expressed by the infectious agent or bacterial cell, which transcriptional activation induces the expression of a nucleic acid that encodes a therapeutic protein.

2. The p16 Transcriptional Promoter

[00112] In one embodiment, the suicide gene could be placed under control of a p16 promoter, such as a p16Ink4a gene promoter, which is transcriptionally active in senescent, but not in non-senescent cells.

[00113] In humans, p16 is encoded by the *CDKN2A* gene, which gene is frequently mutated or deleted in a wide variety of tumors. p16 is an inhibitor of cyclin dependent kinases such as CDK4 and CDK6, which phosphorylate retinoblastoma protein (pRB) thereby causing the progression from G1 phase to S phase. p16 plays an important role in cell cycle regulation by decelerating cell progression from G1 phase to S phase, and therefore acts as a tumor suppressor that is implicated in the prevention of cancers, including, for example, melanomas, oropharyngeal squamous cell carcinomas, and esophageal cancers. The designation p16Ink4A refers to the molecular weight (15,845) of the protein encoded by one of the splice variants of the *CDKN2A* gene and to its role in inhibiting CDK4.

[00114] In humans, p16 is encoded by *CDKN2A* gene, located on chromosome 9 (9p21.3). This gene generates several transcript variants that differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4. The remaining transcript includes an alternate exon 1 located 20 kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein that is structurally unrelated to the products of the other variants. The ARF product functions as a stabilizer of the tumor suppressor protein p53, as it can interact with and sequester MDM2, a protein responsible for the degradation of p53. In spite of their structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by this gene, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in control of the G1 phase of the cell cycle. This gene is frequently mutated or deleted in a wide variety of tumors and is known to be an important tumor suppressor gene.

[00115] Concentrations of p16INK4a increase dramatically as tissue ages. Liu et al., *Aging Cell* 8(4):439–48 (2009) and Krishnamurthy et al., *Nature* 443(7110):453–7 (2006). The increased expression of the p16 gene with age reduces the proliferation of stem cells thereby increasing the cellular senescence-associated health risks in a human.

[00116] p16 is a cyclin-dependent kinase (CDK) inhibitor that slows down the cell cycle by prohibiting progression from G1 phase to S phase. Normally, CDK4/6 binds cyclin D and forms an active protein complex that phosphorylates retinoblastoma protein (pRB). Once phosphorylated, pRB disassociates from the transcription factor E2F1, liberating E2F1 from its cytoplasm bound state allowing it to enter the nucleus. Once in the nucleus, E2F1 promotes the transcription of target genes that are essential for transition from G1 to S phase.

[00117] p16 acts as a tumor suppressor by binding to CDK4/6 and preventing its interaction with cyclin D. This interaction ultimately inhibits the downstream activities of transcription factors, such as E2F1, and arrests cell proliferation. This pathway connects the processes of tumor oncogenesis and senescence, fixing them on opposite ends of a spectrum.

[00118] On one end, the hypermethylation, mutation, or deletion of p16 leads to downregulation of the gene and can lead to cancer through the dysregulation of cell cycle progression. Conversely, activation of p16 through the ROS pathway, DNA damage, or senescence leads to the build up of p16 in tissues and is implicated in aging of cells.

[00119] Regulation of p16 is complex and involves the interaction of several transcription factors, as well as several proteins involved in epigenetic modification through methylation and repression of the promoter region. PRC1 and PRC2 are two protein complexes that modify the expression of p16 through the interaction of various transcription factors that execute methylation patterns that can repress transcription of p16. These pathways are activated in cellular response to reduce senescence.

3. The p21 Transcriptional Promoter

[00120] A nucleic acid encoding a therapeutic protein could be placed under the control of the p21/CDKN1A transcriptional promoter that is often transcriptionally active in senescent, and cancerous or pre-cancerous cells. p53/TP53 plays a central role in the regulation of p21 and, therefore, in the growth arrest of cells when damaged. p21 protein binds directly to cyclin-CDK complexes that drive the cell cycle and inhibits their kinase activity thereby causing cell cycle arrest to allow repair to take place. p21 also mediates growth arrest associated with differentiation and a more permanent growth arrest associated with cellular senescence. The p21 gene contains several p53 response elements that mediate direct binding of the p53 protein, resulting in transcriptional activation of the gene encoding the p21 protein. The role of p53 gene

regulation in cellular senescence is described in Kelley *et al.*, *Cancer Research* 70(9):3566–75. (2010).

Nucleic Acids and Therapeutic Proteins Encoded Thereby

[00121] Nucleic acids that may be suitably employed in the expression constructs, systems, and methods of the present disclosure encode a protein that is capable of reducing, preventing, and/or eliminating the growth and/or survival of a cell in which it is produced, including a target cell. Thus, the target cell specificity of the presently disclosed expression constructs and systems is achieved by the expression within a target cell, but not within a non-target cell, of a nucleic acid that encodes a therapeutic protein.

[00122] Nucleic acids encoding therapeutic proteins that may be employed in the expression constructs and systems of the present disclosure include nucleic acids encoding one or more protein that induces apoptosis in a cell in which it is produced. Exemplified herein are expression constructs and systems comprising one or more “suicide genes,” such as a nucleic acid encoding Herpes Simplex Virus Thymidine Kinase (HSV-TK), cytosine deaminase, Casp3, Casp8, Casp9, BAX, DFF40, cytosine deaminase, or other nucleic acid that encodes a protein that is capable of inducing apoptosis in a cell.

[00123] Apoptosis, or programmed cell death (PCD), is a common and evolutionarily conserved property of all metazoans. In many biological processes, apoptosis is required to eliminate supernumerary or dangerous (such as pre-cancerous) cells and to promote normal development. Dysregulation of apoptosis can, therefore, contribute to the development of many major diseases including cancer, autoimmunity and neurodegenerative disorders. In most cases, proteins of the caspase family execute the genetic programme that leads to cell death.

[00124] Apoptosis is triggered in a mammalian cell, in particular in a human cell, through the activation of caspase proteins, in particular the caspase proteins CASP3, CASP8, and CASP9. See, for example, Xie *et al.*, *Cancer Res* 61(18):186-91 (2001); Carlotti *et al.*, *Cancer Gene Ther* 12(7):627-39 (2005); Lowe *et al.*, *Gene Ther* 8(18):1363-71 (2001); and Shariat *et al.*, *Cancer Res* 61(6):2562-71 (2001).

[00125] DNA fragmentation factor (DFF) is a complex of the DNase DFF40 (CAD) and its chaperone/inhibitor DFF45 (ICAD-L). In its inactive form, DFF is a heterodimer composed of a 45kDa chaperone inhibitor subunit (DFF45 or ICAD), and a 40kDa latent endonuclease subunit (DFF40 or CAD). Upon caspase-3 cleavage of DFF45, DFF40 forms active

endonuclease homo-oligomers. It is activated during apoptosis to induce DNA fragmentation. DNA binding by DFF is mediated by the nuclease subunit, which can also form stable DNA complexes after release from DFF. The nuclease subunit is inhibited in DNA cleavage but not in DNA binding. DFF45 can also be cleaved and inactivated by caspase-7 but not by caspase-6 and caspase-8. The cleaved DFF45 fragments dissociate from DFF40, allowing DFF40 to oligomerise, forming a large complex that cleaves DNA by introducing double strand breaks. Histone H1 confers DNA binding ability to DFF and stimulates the nuclease activity of DFF40. Activation of the apoptotic endonuclease DFF-40 is described in Liu *et al.*, *J Biol Chem* 274(20):13836-40 (1999).

[00126] Thymidine kinase (TK) is an ATP-thymidine 5'-phosphotransferase that is present in all living cells as well as in certain viruses including herpes simplex virus (HSV), varicella zoster virus (VZV), and Epstein-Barr virus (EBV). Thymidine kinase converts deoxythymidine into deoxythymidine 5'-monophosphate (TMP), which is phosphorylated to deoxythymidine diphosphate and to deoxythymidine triphosphate by thymidylate kinase and nucleoside diphosphate kinase, respectively. Deoxythymidine triphosphate is incorporated into cellular DNA by DNA polymerases and viral reverse transcriptases.

[00127] When incorporated into DNA, certain dNTP analogs, such as synthetic analogues of 2'-deoxy-guanosine (*e.g.*, Ganciclovir), cause the premature termination of DNA synthesis, which triggers cellular apoptosis.

[00128] Within certain embodiments, the expression cassettes and systems of the present disclosure employ a nucleic acid that encodes HSV-TK. Following the administration to a human of a system employing a nucleic acid encoding HSV-TK, an analogue of a 2'-deoxy-nucleotide, such as 2'-deoxy-guanosine, is administered to the human. The HSV-TK efficiently converts the 2'-deoxy-nucleotide analogue into a dNTP analogue, which when incorporated into the DNA induces apoptosis in the target cell.

[00129] Cytosine deaminase (CD) catalyzes the hydrolytic conversion in DNA of cytosine to uracil and ammonia. If a CD-modified site is recognized by an endonuclease, the phosphodiester bond is cleaved and, in a normal cell, is repaired by incorporating a new cytosine. In the presence of 5-fluorocytosine (5-FC), cytosine deaminase converts 5-FC into 5-fluorouracil (5-FU), which can inhibit target cell growth. Transgenic expression of CD in a target cell, therefore, reduces the growth and/or survival of the target cell.

[00130] The present disclosure provides expression constructs and systems that further comprise one or more safety features to ensure that the expression of a nucleic acid encoding a therapeutic protein is upregulated in appropriate cells, over a desired time period, and/or to a specified level.

[00131] Within one such embodiments, expression constructs and systems of the present disclosure employ nucleic acids that encode inducible variants of therapeutic proteins, including, for example, inducible variants of Casp3, Casp8, Casp9, which require the further contacting of a cell with or administration to a human of a chemical or biological compound that activates the therapeutic protein.

[00132] Inducible suicide gene systems are well known and readily available in the art and have been described, for example, in Miller *et al.*, PCT Patent Publication No. WO 2008/154644 and Brenner, US Patent Publication No. 2011/0286980. In addition, Shah *et al.*, *Genesis* 45(4):104-199 (2007) describe a double-inducible system for Caspase 3 and 9 that employs RU486 and chemical inducers of dimerization (CID). Straathof *et al.*, *Blood* 105(11):4247-4254 (2005) describe an inducible caspase 9 system in which caspase 9 is fused to a human FK506 binding protein (FKBP) to allow the conditional dimerization using the small molecule AP20187 (ARIAD Pharmaceuticals, Cambridge, MA), which is a non-toxic synthetic analog of FK506. Carlotti *et al.*, *Cancer Gene Ther* 12(7):627-39 (2005) describe an inducible caspase 8 system by employing the ARIADTM homodimerization system (FKC8; ARIAD Pharmaceuticals).

[00133] Full-length inducible caspase 9 (F¹F-C-Casp9.I.GFP) comprises a full-length caspase 9, including its caspase recruitment domain (CARD; GenBank NM001 229) linked to two 12 kDa human FK506 binding proteins (FKBP12; GenBank AH002 818) that contain an F36V mutation as described in Clackson *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 95:10437-10442 (1998) and are connected by a Ser-Gly-Gly-Gly-Ser linker that connects the FKBP12s and caspase 9 to enhance flexibility.

[00134] In a further embodiment, the inducible suicide gene could be linked to the nucleic acid sequence for a detectable biomarker such as luciferase or green fluorescent protein to permit the detection of the targeted cells prior to administering a compound to activate an inducible therapeutic protein.

Compositions and Formulations of Systems Comprising
Vectors and Expression Cassettes

[00135] The present disclosure provides systems comprising a vector and an expression cassette wherein the expression cassette comprises a transcriptional promoter that is responsive to one or more transcription factors that are expressed in a target cell and a nucleic acid encoding a therapeutic protein. Systems can be administered to a human patient by themselves or in pharmaceutical compositions where they are mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a disease or condition as described herein. Mixtures of these systems can also be administered to the patient as a simple mixture or in pharmaceutical compositions.

[00136] Compositions within the scope of this disclosure include compositions wherein the therapeutic agent is a system comprising a vector and an expression cassette in an amount effective to reduce or eliminate the growth and/or survival of a target cell such as a senescent cell, a cancer cell, a cell infected with an infectious agent, a bacterial cell, or a cell that is associated with another disease or condition. Determination of optimal ranges of effective amounts of each component is within the skill of the art. The effective dose is a function of a number of factors, including the specific system, the presence of one or more additional therapeutic agent within the composition or given concurrently with the system, the frequency of treatment, and the patient's clinical status, age, health, and weight.

[00137] Compositions comprising a system may be administered parenterally. As used herein, the term "parenteral administration" refers to modes of administration other than enteral and topical administration, usually by injection, and include, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid, intraspinal, and intrasternal injection and infusion. Alternatively, or concurrently, administration may be orally.

[00138] Compositions comprising a system may, for example, be administered intravenously via an intravenous push or bolus. Alternatively, compositions comprising a system may be administered via an intravenous infusion.

[00139] Compositions include a therapeutically effective amount of a system, and a pharmaceutically acceptable carrier. As used herein, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more

particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skimmed milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

[00140] These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. Such compositions will contain a therapeutically effective amount of the inhibitor, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[00141] Compositions can be formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to a human. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00142] The systems disclosed herein can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from

hydrochloric, phosphoric, acetic, oxalic, tartaric acids, and the like, and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

Methods for Treatment of a Disease or Condition Associated with, and for Reducing, Inhibiting, and/or Preventing the Growth and/or Survival of, a Cell that is Associated with Aging, Cancer, Infectious Disease, Bacterial Infection, and/or other Disease or Condition

[00143] The present disclosure provides methods for reducing, inhibiting, and/or preventing the growth and or survival of a cell that is associated with aging, cancer, infectious disease, bacterial infection, and/or other disease or condition, which methods comprise contacting a target cell or a population of cell comprising a target cell with a system as described herein, which system comprises a vector and an expression construct, which expression construct comprises a transcriptional promoter and a nucleic acid.

[00144] The present disclosure also provides methods for the treatment of aging, cancer, infectious disease, bacterial infection, and/or other disease or condition in a patient, which methods comprise the administration of a system as described herein, which system comprises a vector and an expression construct, which expression construct comprises a transcriptional promoter and a nucleic acid.

[00145] The present therapeutic methods involve contacting a target cell with, or administering to a human patient, a composition comprising one or more system comprising a vector and an expression cassette to a human patient for reducing and/or eliminating the growth and/or survival of a cell that is associated with senescence, cancer, an infectious disease, a bacterial infection or another disease or condition.

[00146] The amount of the system that will be effective in the treatment, inhibition, and/or prevention of aging, cancer, infectious disease, bacterial infection, or other disease or condition that is associated with the elevated expression of one or more transcription factors can be determined by standard clinical techniques. *In vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00147] The systems or pharmaceutical compositions of the present disclosure can be tested *in vitro*, and then *in vivo* for the desired therapeutic or prophylactic activity, prior to use in

humans. For example, *in vitro* assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include the effect of a system on a cell line or a patient tissue sample. The effect of the system or pharmaceutical composition thereof on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to proliferation and apoptosis assays. In accordance with the present disclosure, *in vitro* assays that can be used to determine whether administration of a specific compound is indicated, include *in vitro* cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

[00148] The present disclosure provides methods for the treatment and growth and/or survival inhibition by administration to a subject of an effective amount of a system or pharmaceutical composition thereof as described herein. In one aspect, the system is substantially purified such that it is substantially free from substances that limit its effect or produce undesired side-effects.

[00149] Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The systems or compositions thereof may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the inhibitors or compositions into the central nervous system by any suitable route, including intraventricular and intrathecal injection. Intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, for example, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[00150] It may be desirable to administer the systems or compositions thereof locally to the area in need of treatment; this may be achieved by, for example, local infusion during surgery, topical application, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

[00151] The system can be delivered in a controlled release system placed in proximity of the therapeutic target, thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, in *Medical Applications of Controlled Release* 2:115-138 (1984)).

[00152] Intravenous infusion of a compositions comprising a system may be continuous for a duration of at least about one day, or at least about three days, or at least about seven days, or at least about 14 days, or at least about 21 days, or at least about 28 days, or at least about 42 days, or at least about 56 days, or at least about 84 days, or at least about 112 days.

[00153] Continuous intravenous infusion of a composition comprising a system may be for a specified duration, followed by a rest period of another duration. For example, a continuous infusion duration may be from about 1 day, to about 7 days, to about 14 days, to about 21 days, to about 28 days, to about 42 days, to about 56 days, to about 84 days, or to about 112 days. The continuous infusion may then be followed by a rest period of from about 1 day, to about 2 days to about 3 days, to about 7 days, to about 14 days, or to about 28 days. Continuous infusion may then be repeated, as above, and followed by another rest period.

[00154] Regardless of the precise infusion protocol adopted, it will be understood that continuous infusion of a composition comprising a system will continue until either desired efficacy is achieved or an unacceptable level of toxicity becomes evident.

* * * * *

[00155] It will be understood that, unless indicated to the contrary, terms intended to be "open" (*e.g.*, the term "including" should be interpreted as "including but not limited to," the term "having" should be interpreted as "having at least," the term "includes" should be interpreted as "includes but is not limited to," etc.). Phrases such as "at least one," and "one or more," and terms such as "a" or "an" include both the singular and the plural.

[00156] It will be further understood that where features or aspects of the disclosure are described in terms of Markush groups, the disclosure is also intended to be described in terms of any individual member or subgroup of members of the Markush group. Similarly, all ranges disclosed herein also encompass all possible sub-ranges and combinations of sub-ranges and that language such as "between," "up to," "at least," "greater than," "less than," and the like include the number recited in the range and includes each individual member.

[00157] All references cited herein, whether *supra* or *infra*, including, but not limited to, patents, patent applications, and patent publications, whether U.S., PCT, or non-U.S. foreign, and all technical and/or scientific publications are hereby incorporated by reference in their entirety.

EXAMPLES

[00158] While various embodiments have been disclosed herein, other embodiments will be apparent to those skilled in the art. The various embodiments disclosed herein are for purposes of illustration and are not intended to be limiting, with the true scope and spirit being indicated by the claims. The present disclosure is further described with reference to the following examples, which are provided to illustrate certain embodiments and are not intended to limit the scope of the present disclosure or the subject matter claimed.

Example 1

p14 FAST Fusogenic Lipid Nanoparticles (LNP) Enhance Gene Delivery to Tumors

[00159] This Example demonstrates that Fusogenix™ (Innovascreen, Halifax, Nova Scotia, Canada) lipid nanoparticles utilizing a p14 FAST fusion from reptilian reovirus are effective at delivering a plasmid DNA construct to a target tumor.

[00160] Fusogenix lipid nanoparticles labeled with ⁶⁴Cu (⁶⁴Cu NOTA-liposomes) either with or without a p14 FAST fusion protein (described in PCT Patent Publication Nos. WO2002044206A2 and WO2012040825A1) were administered intravenously to a M16 mouse model system for prostate cancer (PC3 cells). Seo, *Bioconjug Chem* 19(12):2577-2585 (2009) and Reeves, *Cancer Therapy* 136(7):1731-1740 (2014). 24 hours post-immunization, PC3 tumors were visualized using positron emission tomography (PET). **FIGs. 7A and 7B.**

[00161] The data presented in **FIG. 8** demonstrate a 50% increase in PC3 prostate tumor uptake of ⁶⁴Cu NOTA-liposomes with p14 FAST fusion protein as compared to ⁶⁴Cu NOTA-liposomes without p14 FAST fusion protein. The biodistribution of labelled pegylated liposomes in nude mice expressed after 24 hours is presented in **FIG. 9.**

Example 2

In vivo Administered p14 FAST Fusogenic Lipid Nanoparticles are Non-toxic and Well Tolerated

[00162] This Example demonstrates that Fusogenix™ (Innovascreen, Halifax, Nova Scotia, Canada) lipid nanoparticles utilizing a p14 FAST fusion from reptilian reovirus do not exhibit

adverse side-effects in any of the major mammalian organ systems examined when administered *in vivo* to Sprague-Dawley rats. are effective at delivering a plasmid DNA construct to a target tumor.

[00163] Presented herein are comparative studies that were performed with N=20 male rats treated with either (i) no LNPs (PBS), (ii) LNPs without p14, or (iii) p14 containing Fusogenix lipid nanoparticles (LNPs). Each animal received a total of three injections of 15 mg/kg in their tail, over a 4 day period. Treatment of the animals with p14 containing LNPs did not result in any acute changes in animal behavior and animal growth was not affected by treatment with p14 containing LNPs. Animals treated with p14 containing LNPs had similar organ weights as compared to all other animal groups studied.

[00164] Treatment with p14 containing LNPs did not affect the microscopic appearance of tissues from major organ systems. Tissues from the lungs, brain, heart, kidney, liver, reproductive organs, gut, endocrine system, lymph nodes, spleen, pancreas, bladder and tail were all independently examined and p14 did not elicit any visible signs of toxicity. Importantly, the liver appeared to be unaffected by exposure to p14. Moreover, no differences were identified between the tissues of p14 treated animals versus control groups.

[00165] A number of blood chemistry values were measured to determine the impact of p14 on physiological function and inflammation. Parameters such as ALT and AST that denote acute liver function were all within normal ranges. Fusogenix LNPs containing p14 do not show any adverse side-effects in any of the major mammalian organ systems examined. Histological appearance of tissues was also normal.

[00166] Mice were injected three (3) times at 10 day intervals with purified p14 mixed with Freund's adjuvant. A first dose contained CFA (complete Freund's adjuvant) while second and third doses contained IFA (incomplete Freund's adjuvant). Each injection was with 50 µg of p14. Mice were sacrificed after 30 days and sera was analyzed for anti-p14 antibodies. p14 lipid nanoparticles were also tested in two (2) mice via intravenous injection of 400 µg of p53-iCasp9 Fusogenix lipid nanoparticles containing 240 µg of p14. Mice were sacrificed after 30 days of injection and serum was analyzed for anti-p14 antibodies. A positive control included purified antibodies spiked in serum at a high dose of 250 ng/ml and a low dose of 50 ng/ml. The data presented in **FIGs. 10 and 11** demonstrate the safety and tolerability of Fusogenix™ lipid nanoparticles utilizing a reptilian reovirus p14 FAST fusion protein. Anti-p14 and anti-LNP

antibody assays demonstrated that virtually no antibody response was observed in immune competent mice (with and without adjuvant).

[00167] Ten (10) human serum samples were tested for Complement activation-related psuedoallergy (CARPA) using C4d and iC3b complement ELISA assays as described in Szebeni, *Mol Immunol* 61(2):163-73 (2014). The data presented in **FIGs. 12** and **13** demonstrate that LNP formulations according to the present disclosure were non-reactive with C4d (**FIG. 12**) and less reactive with iC3b (**FIG. 13**) as compared to Doxil in 8 out of 10 human samples (approximately 5-10% of humans exhibit a CARPA reaction to nanomedicines such as Doxil).

[00168] *In vitro* anti-p14 and anti-LNP antibody neutralization assays revealed that vector neutralization required very high antibody concentrations. Moreover, vaccination or pretreatment with p14-LNPs did not result in a decrease in therapeutic efficacy and repeated *in vivo* dosing was effective and well tolerated. CARPA assays with Fusogenix™ p14 FAST lipid nanoparticles elicit less complement activity as compared to a control pegylated liposomal doxorubicin (Doxil).

Example 3

In vivo Suppression of p16-positive Senescent Cell Burden in Aged Mice

[00169] This Example demonstrates the target-cell specific suppression in p16-positive senescent cell burden following the *in vivo* administration of an exemplary p16-targeting construct in an mouse model system for aging.

[00170] The aging mouse model exhibits a senescent cell burden (as defined by the presence of p16⁺ cells) and secretion of factors associated with a senescence-associated secretory phenotype (SASP; van Deursen, *Nature* 509(7501):439-446 (2014)).

[00171] A formulation comprising a vector and an expression construct, such as a lipid nanoparticle (LNP) vector, *e.g.*, a fusogenic LNP comprising a fusogenic protein such as p14 FAST, encompassing a p16-iCasp9 expression construct (pVAX1-16s-iCasp9; SEQ ID NO: 06; **FIG. 16**) which comprises an exemplary p16-targeting construct for the target cell-specific expression of an inducible Caspase 9 (iCasp9) protein in target cells expressing p16, such as target cells that are associated with aging and/or senescence, which p16-targeting construct comprises a p16s transcriptional promoter in operable connection to iCasp9 or variant thereof expressing luciferase (for visualization), was administered *in vivo* to an aged mouse via injection into a tail vein and the LNP+expression construct transfects target and non-target cells without specificity. **FIG. 19**. Upon subsequent *in vivo* administration of the chemical inducer of dimerization (CID),

AP20187, p16⁺ target cells (*e.g.*, senescent cells) underwent apoptosis, resulting in a reduction in SASP levels, while p16⁻ cells remained viable.

[00172] Histological staining of senescent-associated β -gal in kidney cells from an *in vivo* aged mouse model either untreated (upper left panel) or treated (upper right panel) following the *in vivo* administration (16 animals at 80 weeks of age) of a formulation comprising a vector and an expression construct, such as a lipid nanoparticle (LNP) vector, *e.g.*, a fusogenic LNP comprising a fusogenic protein such as p14 FAST, encompassing a p16-iCasp9 expression construct, *e.g.*, pVAX1-16s-iCasp9 or variant thereof, was administered *in vivo* to an aged mouse and kidney cells stained for β -gal. FIGs. 20A-D. The lower panel is a photomicrograph of the histological staining of senescent-associated β -gal in 4-month old kidney cells from a normal mouse. These data demonstrated a dose-dependent reduction of p16⁺ senescent kidney cells.

[00173] The dose-dependent targeting of p16⁺ kidney cells (FIG. 21), spleen cells (FIG. 22), seminal vesicle cells (FIG. 23), inguinal fat cells (FIG. 24), and lung cells (FIG. 25) was demonstrated in naturally aged mice following the *in vivo* administration of a fusogenic lipid nanoparticle (LNP) formulation comprising a pVAX1-p16 expression construct. Kidney cells were subjected to a qRT-PCR reaction to detect p16^{ink4a} transcripts. Relative expression was calculated using $2^{-\Delta\Delta Ct}$ (Livak, *Methods* 25:402-408 (2001)).

Example 4

In vivo Oncology Study with NSG Mice Implanted with a Human Prostate Tumor

[00174] This Example demonstrates the target-cell specific suppression of p53-expressing prostate cancer cells in NSG mice implanted with a human prostate tumor (*i.e.*, a PC-3 xenograft).

[00175] Human prostate cancer PC-3 cells were treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc (luciferin) plasmid (in the presence and absence of the homodimerizer AP201870) and assessed for iCasp9 expression and subjected to Western blot analysis of iCasp 9 and Casp 9 protein levels obtained with p53- expressing cells (pVax-p53) and control cells (pcDNA3-GFP). FIG. 36. These data demonstrated that the addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolishes the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

[00176] Human prostate cancer cells (LNCaP (FIG. 38), DU145 (FIG. 39), and PC-3 (FIG. 40)) and normal epithelial cells (RWPE (FIG. 41)) were treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc plasmid and assessed for iCasp9 expression by

Western blot and luminescence assays. These data demonstrated that addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolished the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

[00177] Human prostate cancer PC-3 cells were treated with Fusogenix lipid nanoparticles carrying the pVax-p53-iCasp9-luc (luciferin) plasmid (in the presence and absence of the homodimerizer AP20187) and assessed for iCasp9 expression. The data presented in **FIG. 42** demonstrated that the addition of the chemical inducer of dimerization (CID; *e.g.*, AP20187 and AP1903) abolished the expression of iCasp9 and luciferase in p53-expressing cells engineered to express iCasp9 or luciferase.

[00178] Flow cytometry apoptosis data (Annexin V) from human prostate cancer PC-3 cells treated with pVax-p53 Fusogenix lipid nanoparticles (in the absence and presence of AP20187, **FIG. 43A** and **44A** and **43B** and **44B**, respectively) demonstrated that suicide gene therapy selectively killed p53-expressing human prostate cancer cells in culture by inducing apoptosis (Luciferase-Annexin V flow cytometry).

[00179] A pre-clinical oncology study according to the present disclosure was conducted with 30x NSG mice implanted with human prostate tumor cells. **FIG. 45**. NSG mice bearing a subcutaneous human prostate PC-3 tumor were injected intratumorally (IT) with 100 µg Fusogenix pVax-p53 formulation, followed 96 hours later by intravenous (IV) administration of 2 mg/kg of the homodimerizer AP20187. **FIG. 46**. Tumors from the NSG mice bearing subcutaneous human prostate PC-3 tumors injected intratumorally with 100 µg Fusogenix pVax-p53 formulation, followed 96 hours by 2 mg/kg AP20187 IV, were photographed (**FIGs 47A-47C**).

[00180] Four NSG mice bearing subcutaneous human prostate cancer PC-3 tumors that were injected intravenously (IV) with 4x 100 µg doses of Fusogenix pVax-p53 formulation, followed 24 hours later by 2 mg/kg AP20187 IV. Tumor volume was measured and plotted as a function of time following IV injection. **FIGs. 48-51**.

[00181] The percentage change in tumor volume was determined and plotted as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in NSG mice (N=6 for all groups) bearing a prostate tumor that were treated with intravenous p14 LNP pVAX. **FIG. 52**. The percent survival was determined and plotted as a function of time after *in vivo*

administration of a chemical inducer of dimerization (CID) in NSG mice (N=6 for all groups) bearing a prostate tumor that were treated with intravenous p14 LNP pVAX. **FIG. 53.**

[00182] A dose escalation study was carried out in which the percentage change in tumor volume as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in NOD-SCID mice (N=6 for all groups) bearing a prostate tumor that were treated with 100 μ g, 400 μ g, and 1000 μ g of intravenous p14 LNP pVAX. NOD-SCID mice were implanted subcutaneously with 500,000 PC-3 cells and randomized into treatment groups when their tumors reached 200 mm³, (N = 2 for all groups). Animals were injected with their assigned dose of p53-iCasp9 LNP IV twice followed by 2 mg/kg dimerizer. Tumors were measured directly every 24 hours. **FIG. 54.**

[00183] In total, the data presented herein demonstrate that apoptosis can be reliably induced in a p53+ prostate cancer cell-specific manner by the intravenous administration of fusogenic lipid nanoparticle formulations comprising a p53-iCasp9 expression construct.

Example 5

In vivo Suppression of Metastases in NOD-SCID Mice Implanted with a Metastatic Tumor

[00184] The suppression of metastatic tumor growth with repeat treatment of a p53-iCasp9 LNP with or without a chemical inducer of dimerization (CID) was demonstrated in a NOD-SCID mouse model system.

[00185] NOD-SCID mice were injected with 500,000 PC-3M-luciferase cells on Day 0, LNP dosing was started on Day 22 with 150 μ g p53-iCasp9 LNP. Dimerizer doses started Day 24 at 2 mg/kg. Mice were imaged every 24-48 hours to detect whole animal luminescence. **FIG. 55.**

Example 6

In vivo Suppression of Melanoma in Isogenic C57B6 Mice Implanted with B16 Murine Melanoma Cells

[00186] Isogenic C57B6 mice implanted with B16 murine melanoma cells were treated with LNPs containing a construct encoding iCasp9 and murine CD40L under control of the murine p53 promoter followed by the AP20187 dimerizer.

[00187] The percentage change in tumor volume (**FIG. 56**) and percent survival (**FIG. 57**) was measured as a function of time after *in vivo* administration of a chemical inducer of dimerization (CID) in isogenic C57B6 mice implanted by subcutaneous injection with 250,000 B16 murine

melanoma cells treated (grown to 400 mm³) with LNPs containing a construct encoding iCasp9 and murine CD40L under control of the murine p53 promoter.

[00188] These data demonstrated that, even though the rapid (10 hour) doubling time of the B16 cells made them largely refractory to the iCasp9-induced apoptosis, they still secreted enough CD40L to effectively halt the tumor's growth. A construct encoding GMCSF + OVA antigen was also tested and determined to be more effective than iCasp9 alone, but less effective than the CD40L version. N=3 for both groups.

Example 7

In vivo Suppression of Lung Cancer Metastasis in Mice Implanted with B16F10 Murine Melanoma Cells

[00189] This Example demonstrates the *in vivo* p53+ target cell suppression murine p53+ B16F10 melanoma target cells implanted in a lung metastasis mouse model system.

[00190] A B16F10 lung metastasis model system was employed in which 100 µg of a control LNP or a p53-iCasp9 LNP was administered intravenously at days 3, 6, 9, and 12 following the intravenous injection of 75,000 B16F10 cells. At days 5, 8, 11, and 13, a chemical inducer of dimerization (CID) was administered intraperitoneally. Animals were sacrificed at day 14 and lung metastases were quantified. FIGs. 58A-58D and 59.

CLAIMS

What is claimed is:

1. A lipid nanoparticle (LNP) formulation for the targeted production of a therapeutic protein within a target cell, said LNP formulation comprising:
 - a. a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, which mammalian cells include both target cells or non-target cells, wherein said lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s), and
 - b. an expression construct for the preferential production of a therapeutic protein within a target cell, said expression construct comprising:
 - i. a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within said target cells as compared to said non-target cells and
 - ii. a nucleic acid that is operably linked to and under regulatory control of said transcriptional promoter, wherein said nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced within said target cells but is not produced in said non-target cells.
2. The LNP formulation of claim 1 wherein said one or more lipid(s) are present in said LNP formulation at a concentration ranging from 1 mM to 100 mM.
3. The LNP formulation of claim 2 wherein said one or more lipid(s) are present in said LNP formulation at a concentration ranging from 5 mM to 50 mM.
4. The LNP formulation of claim 3 wherein said one or more lipid(s) are present in said LNP formulation at a concentration ranging from 10 mM to 30 mM.
5. The LNP formulation of claim 4 wherein said one or more lipid(s) are present in said LNP formulation at a concentration ranging from 15 mM to 25 mM.
6. The LNP formulation of claim 5 wherein said one or more lipid(s) are present in said LNP formulation at a concentration of about 20 mM.
7. The LNP formulation of claim 1 wherein one or more of said lipid(s) is

selected from the group consisting of 1,2-dioleoyl-3-dimethylammonium-propane (DODAP), 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), Cholesterol, and 1,2-dimyristoyl-rac-glycero-3-methoxypolyethylene glycol (DMG-PEG).

8. The LNP formulation of claim 7 wherein said lipid(s) include two or more lipids selected from the group consisting of DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG.

9. The LNP formulation of claim 8 wherein said lipid(s) include DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG.

10. The LNP formulation of claim 9 wherein said DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG are present in said LNP formulation at a molar ratio of 35-55 mole % DODAP:10-20 mole % DOTAP:22.5-37.5 mole % DOPE:4-8 mole % Cholesterol:3-5 mole % DMG-PEG.

11. The LNP formulation of claim 10 wherein said DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG are present in said LNP formulation at a molar ratio of about 45 mole % DODAP:about 15 mole % DOTAP:about 30 mole % DOPE:about 6 mole % Cholesterol:about 4 mole % DMG-PEG.

12. The LNP formulation of claim 11 wherein said DODAP, DOTAP, DOPE, Cholesterol, and DMG-PEG are present in said LNP formulation at a molar ratio of 45 mole % DODAP:15 mole % DOTAP:30 mole % DOPE:6 mole % Cholesterol:4 mole % DMG-PEG.

13. The LNP formulation of claim 1 wherein said one or more fusogenic protein(s) are present in said LNP formulation at a concentration ranging from 0.5 μM to 20 μM .

14. The LNP formulation of claim 13 wherein said one or more fusogenic protein(s) are present in said LNP formulation at a concentration ranging from 1 μM to 10 μM .

15. The LNP formulation of claim 14 wherein said one or more fusogenic protein(s) are present in said LNP formulation at a concentration ranging from 3 μM to 4 μM .

16. The LNP formulation of claim 15 wherein said one or more fusogenic protein(s) are present in said LNP formulation at a concentration of 3.5 μM .

17. The LNP formulation of claim 1 wherein one or more of said fusogenic protein(s) is selected from the group consisting of a p14 fusogenic protein (SEQ ID NO: 16) and a p14e15 fusogenic protein (SEQ ID NO: 17).

18. The LNP formulation of claim 1 wherein said fusogenic protein is a p14 fusogenic protein (SEQ ID NO: 16).

19. The LNP formulation of claim 1 wherein said fusogenic protein is a p14e15 fusogenic protein (SEQ ID NO: 17).

20. The LNP formulation of claim 1 wherein said expression construct is present in said LNP formulation at a concentration ranging from 20 µg/mL to 1.5 mg/mL.

21. The LNP formulation of claim 20 wherein said expression construct is present in said LNP formulation at a concentration ranging from 100 µg/mL to 500 µg/mL.

22. The LNP formulation of claim 21 wherein said expression construct is present in said LNP formulation at a concentration of 250 µg/mL.

23. The LNP formulation of claim 21 wherein said expression construct is present in said LNP formulation at a concentration of 250 µg/mL.

24. The LNP formulation of claim 1 wherein said transcriptional promoter is selected from the group consisting of a p16 transcriptional promoter, a p21 transcriptional promoter, and a p53 transcriptional promoter.

25. The LNP formulation of claim 24 wherein said transcriptional promoter is responsive to a factor selected from the group consisting of SP1, ETS1, ETS2, and p53/TP53.

26. The LNP formulation of claim 24 wherein said transcriptional promoter is selected from a p16INK4a/CDKN2A transcriptional promoter and a p21/CDKN1A transcriptional promoter.

27. The LNP formulation of claim 24 wherein said transcriptional promoter is responsive to a factor selected from the group consisting of an EBF3, O/E-1, Pax-5, E2A, p53, VP16, MLL, HSF1, NF-IL6, NFAT1, AP-1, AP-2, HOX, E2F3, and/or NF-κB transcription factor.

28. The LNP formulation of claim 27 wherein said transcriptional promoter is selected from the group consisting of the p21^{cip1/waf1} promoter, the p27^{kip1} promoter, the p53^{kip2} promoter, the TdT promoter, the Rag-1 promoter, the B29 promoter, the Blk promoter, the CD19 promoter, the BLNK promoter, and the λ 5 promoter.

29. The LNP formulation of claim 1 wherein said nucleic acid encodes a therapeutic protein selected from the group consisting of a caspase (Casp), an inducible caspase (iCasp), a self-activating caspase (saCasp), BAX, DFF40, HSV-TK, and cytosine deaminase.

30. The LNP formulation of claim 29 wherein said nucleic acid encodes a caspase selected from the group consisting of a Casp3, a Casp8, and a Casp9.

31. The LNP formulation of claim 30 wherein said nucleic acid encodes a Casp9.

32. The LNP formulation of claim 31 wherein said nucleic acid encodes an inducible Casp9 (iCasp9).

33. The LNP formulation of claim 31 wherein said nucleic acid encodes a self-activating Casp9 (saCasp9).

34. A method for reducing, preventing, and/or eliminating the growth of a target cell, said method comprising contacting a target cell with an LNP formulation comprising:

a. a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, which mammalian cells include both target cells or non-target cells, wherein said lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s), and

b. an expression construct for the preferential production of a therapeutic protein within a target cell, said expression construct comprising:

i. a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within said target cells as compared to said non-target cells and

ii. a nucleic acid that is operably linked to and under regulatory control of said transcriptional promoter, wherein said nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced

within said target cells but is not produced in said non-target cells.

35. A method for the treatment of a disease or condition in a patient having a target cell, said method comprising: administering to said patient an LNP formulation comprising:

a. a lipid nanoparticle vector for the non-specific delivery of a nucleic acid to mammalian cells, which mammalian cells include both target cells or non-target cells, wherein said lipid nanoparticle includes one or more lipid(s) and one or more fusogenic protein(s), and

b. an expression construct for the preferential production of a therapeutic protein within a target cell, said expression construct comprising:

i. a transcriptional promoter that is activated in response to one or more factors that are preferentially produced within said target cells as compared to said non-target cells and

ii. a nucleic acid that is operably linked to and under regulatory control of said transcriptional promoter, wherein said nucleic acid encodes a therapeutic protein that can reduce, prevent, and/or eliminate the growth and/or survival of mammalian cells, including both target cells and non-target cells and wherein said therapeutic protein is produced within said target cells but is not produced in said non-target cells.

FIG. 1




FIG. 2




FIG. 3


Company	Technology	Type	MTD	Clinical Development
Various	Lipid-uz, DOTAP, others	CL	0.045 mg/kg	Phase I (various - all terminated due to toxicity)
Colson/Amgen/Novartis	RONDEL lipopolymer	CL	0.81 mg/kg	Phase I (CALAA-01)
ECEN	PEG-PEI-Cholesterol Lipopolymer	CL	0.65 mg/kg	Phase II (ESEN-001, failed due to toxicity)
Marina Biotech, ProNAI	Smarticles (amphoteric liposomes)	CCL	3.2 mg/kg	Phase I (PNT2258)
Marina Biotech	DILA*	CCL	1 mg/kg	Phase I
Arbutus (Iskemia) /Angion	SNALP	CCL	0.9 mg/kg	7 programs in Phase I and II
MD Anderson (Aral Sood)	DOPC liposomes	NL	>10 mg/kg	Phase I (siRNA-DOPC-EphA2)
 innovoscreen	Fusogenix LNPs	NL	>15 mg/kg	Estimated from rat toxicity study

FIG. 4A




FIG. 4B






FIG. 5

FIG. 6

FIGs. 7A-7B

⁶⁴Cu-NOTA-liposome
w/o protein

⁶⁴Cu-NOTA-liposome-
p14





FIG. 7A

FIG. 7B


FIG. 8**PC3 tumor uptake (24 h p.i.)**

FIG. 9





FIG. 10


FIG. 11

FIG. 12

Detection of C4d fragment in human serum




FIG. 13

Detection of iC3b fragment in human serum




FIG. 14




FIG. 15

p16-targeting construct (aging/senescence)



p16 promoter

iCasp9 or *SA-Casp9*

FIG. 17




FIG. 18









FIG. 19



FIGs. 20A-D

Kidney



FIGs. 20E-H

Seminal Vesicle





FIG. 21

KIDNEY




FIG. 22

SPLEEN




FIG. 23




FIG. 24

INGUINAL FAT




FIG. 25




FIG. 26

Level of p53 RNA normalized to group control




FIG. 27

p53-targeting construct (oncology)



FIG. 28





FIG. 29



pVax1-p53-iCasp9-5 (saCasp9 - M&M's Version) (GenArt 16ADG21C_1977690)
3740 bp

FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37A-37D




FIG. 38




FIG. 39




FIG. 40




FIG. 41




FIG. 42




FIG. 43





FIG. 43A


FIG. 43B

FIG. 44

Annexin V flow cytometry

**FIG. 44A**

pVax-p53 alone

**FIG. 44B**

pVax-p53 + AP20187

FIG. 45




FIG. 46




FIG. 47




FIG. 47A




FIG. 47B




FIG. 47C

FIG. 48




FIG. 49




FIG. 50




FIG. 51





FIG. 52**Change in Tumor Volume After Dimerizer Treatment (IV Group)**

*=p<0.05, **=p<0.01, ***=p<0.001


FIG. 53

FIG. 54




FIG. 55




FIG. 56

CD40L/iCasp9 Treatment









FIG. 57



FIGs. 58A-58D



Control LNP

P53-iCasp9 LNP


FIG. 59

FIG. 60

DEXA scans performed monthly




FIG. 62




FIG. 63




FIG. 64




FIG. 1

