



(86) Date de dépôt PCT/PCT Filing Date: 1997/10/01
(87) Date publication PCT/PCT Publication Date: 1998/04/23
(45) Date de délivrance/Issue Date: 2006/11/28
(85) Entrée phase nationale/National Entry: 1999/04/09
(86) N° demande PCT/PCT Application No.: SE 1997/001651
(87) N° publication PCT/PCT Publication No.: 1998/016228
(30) Priorité/Priority: 1996/10/11 (SE9603725-4)

(51) Cl.Int./Int.Cl. *A61K 31/44* (2006.01),
A61K 31/415 (2006.01), *A61K 31/00* (2006.01),
A61K 31/4184 (2006.01), *A61K 31/435* (2006.01),
A61K 31/4439 (2006.01), *A61K 31/444* (2006.01),
A61K 31/56 (2006.01), *A61K 31/57* (2006.01),
A61P 29/00 (2006.01)

(72) Inventeurs/Inventors:
LINDBERG, PER, SE;
PINAS-MASSO, JOAN, ES;
SERRA-CARRERAS, JORDI, ES;
TROFAST, JAN, SE

(73) Propriétaire/Owner:
ASTRAZENECA AB, SE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : UTILISATION D'UN INHIBITEUR DE H⁺, K⁺-ATPASE DANS LE TRAITEMENT DE POLYPES DU NEZ
(54) Title: USE OF AN H⁺, K⁺-ATPASE INHIBITOR IN THE TREATMENT OF NASAL POLYPS

(57) **Abrégé/Abstract:**

The invention provides a method for the treatment of polyposis which comprises treating a subject suffering from polyposis with an H⁺, K⁺-ATPase inhibitor and, optionally, a glucocorticoid. The invention also relates to a pharmaceutical formulation for simultaneous, separate or sequential administration in the treatment of Widal's Syndrome and in the treatment of asthma.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 31/415	A1	(11) International Publication Number: WO 98/16228 (43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/SE97/01651 (22) International Filing Date: 1 October 1997 (01.10.97) (30) Priority Data: 9603725-4 11 October 1996 (11.10.96) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG (publ) [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): LINDBERG, Per [SE/SE]; Gundas Gata 40, S-431 51 Mölndal (SE). PINAS-MASSO, Joan [ES/ES]; 2 ^o -3 ^o piso, Placa Francesc Masia, 8, E-08029 Barcelona (ES). SERRA-CARRERAS, Jordi [ES/ES]; 1 ^o -3 ^o piso, Paseo Pere III, 71ü, E-08240 Manresa (ES). TRO- FAST, Jan [SE/SE]; Vapenkroken 34, S-226 47 Lund (SE). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: USE OF AN H ⁺ , K ⁺ -ATPase INHIBITOR IN THE TREATMENT OF NASAL POLYPS (57) Abstract The invention provides a method for the treatment of polyposis which comprises treating a subject suffering from polyposis with an H ⁺ , K ⁺ -ATPase inhibitor and, optionally, a glucocorticoid. The invention also relates to a pharmaceutical formulation for simultaneous, separate or sequential administration in the treatment of Widal's Syndrome and in the treatment of asthma.		

23940-1065

1

USE OF AN H^+,K^+ -ATPASE INHIBITOR IN THE TREATMENT OF NASAL POLYPS

Field of the Invention

The present invention provides a new treatment for polyposis
5 using proton pump inhibitors (PPIs), i.e., H^+,K^+ -ATPase inhibitors.

Background of the Invention

Polyposis can generally arise in the nose and the
gastrointestinal tract. In the nose, polyps are pale bags
10 of tissue that arise in the nasal cavity. Their paleness is
generally due to poor blood supply. It is not known what
causes the polyps to be formed but their presence is often
associated with certain medical conditions, for example
asthma and aspirin intolerance. Within the general
15 population the incidence of nasal polyps is low at around
only 1% but 13% of asthma sufferers and 36% of aspirin
intolerant asthmatics suffer from nasal polyposis. The
triple condition of nasal polyposis, aspirin intolerance and
asthma is known as Widal's Syndrome.

20 Nasal polyposis is generally treated in two stages.
Initially a reduction in size of the polyps is achieved
either by surgery or by the application of a topical
intranasal steroid preparation, for example betamethasone
sodium phosphate. Once a reduction in size has been
25 obtained then long term maintenance of the reduction is
necessary by regular use of an intranasal steroid spray such
as beclomethasone dipropionate, budesonide, or fluticasone
propionate. When rapid amelioration is required, oral
steroids such as prednisolone or dexamethasone or synthetic
30 adrenocorticotrophic hormones are used (see V J Lund
Diagnosis and treatment of nasal polyps Brit Med J 1995,

23940-1065

1a

311, 1411-4). There are also proposals that non-steroidal antiinflammatory drug can be used in the treatment of nasal polyposis (see WO 9703659-A).

Summary of the Invention

- 5 In one aspect, the invention provides use of an H^+, K^+ -ATPase inhibitor and a glucocorticoid in the manufacture of a pharmaceutical formulation intended for simultaneous, separate or sequential administration in the treatment of Widal's Syndrome.
- 10 In a further aspect, the invention provides use of an H^+, K^+ -ATPase inhibitor and a glucocorticoid in the manufacture of a pharmaceutical formulation intended for simultaneous, separate or sequential administration in the treatment of nasal polyps.
- 15 In a still further aspect, the invention provides use of an H^+, K^+ -ATPase inhibitor and a glucocorticoid for the treatment of Widal's Syndrome.

In a yet further aspect, the invention provides use of an H^+, K^+ -ATPase inhibitor and a glucocorticoid for the treatment
20 of nasal polyps.

In another aspect, the invention provides a pharmaceutical formulation for simultaneous, separate or sequential use in the treatment of Widal's Syndrome which formulation comprises an H^+, K^+ -ATPase inhibitor and a glucocorticoid.

- 25 The invention also provides a pharmaceutical formulation for simultaneous, separate or sequential administration in the treatment of nasal polyps which pharmaceutical formulation comprises an H^+, K^+ -ATPase inhibitor and a glucocorticoid.

23940-1065

1b

Detailed Description of the Invention

According to the invention there is provided a method for the treatment of nasal polyps which method comprises treating a subject suffering from the said condition with an

5 H⁺,

K^+ -ATPase inhibitor. The invention further provides the use of an H^+ , K^+ -ATPase inhibitor in the manufacture of a medicament for the treatment of nasal polyps.

H^+ , K^+ -ATPase inhibitors are a known class of pharmaceutical agents generally used in therapy for the treatment of gastric acid related diseases. Examples of H^+ , K^+ -ATPase inhibitors are for instance compounds known under the generic names omeprazole, lansoprazole, pantoprazole, rabeprazole and leminoprazole. Some of these compounds are for instance disclosed in EP-A1-0005129, EP-A1-174726, EP-A1-166287 and GB 2163747.

10


These pharmaceutical substances are generally known to be useful for inhibiting gastric acid secretion in mammals and man by controlling gastric acid secretion at the final step of the acid secretory pathway. Thus, in a more general sense, they may be used for prevention and treatment of gastric-acid related diseases in mammals and man, including e.g. reflux oesophagitis, gastritis, duodenitis, gastric ulcers and duodenal ulcers.

15

It has now surprisingly been found that H^+ , K^+ -ATPase inhibitors are useful in the treatment of nasal polyps, particularly where known treatments have failed.

The H^+ , K^+ -ATPase inhibitors preferably used in the invention are compounds of the general formula

20




WO 98/16228


PCT/SE97/01651

3


wherein Het₁ is



or




Het₂ is




5

and X is



or



wherein N in the benzimidazole moiety of Het₂ means that one of the ring carbon atoms substituted by R₆-R₉ optionally may be exchanged for a nitrogen atom without any substituents;

R₁ and R₃ each independently represent hydrogen, alkyl, or alkoxy on the condition that R₁ and R₃ do not simultaneously represent alkoxy; and R₂ represents alkyl, alkoxy optionally substituted by fluorine, alkylthio or alkoxyalkoxy; or one of R₁ and R₃ is halogen and the other is hydrogen and R₂ is 1-morpholino, 1-piperidino or dialkylamino;

R₄ and R₅ are the same or different and selected from hydrogen and alkyl;

R_6 - R_9 are the same or different and selected from hydrogen, halogen, alkyl, alkoxy, haloalkoxy, alkylcarbonyl, and alkoxycarbonyl;

5 R_{10} is hydrogen or R_{10} and R_3 together complete a ring containing 6 to 8 carbon atoms; and


R_{11} represents hydrogen, halogen or alkyl;






10 wherein the compound of formula (I) is optionally in the form of an pharmaceutically acceptable alkaline salt or in its neutral form or is a single enantiomer or a racemic mixture thereof;




wherein each alkyl or alkylenyl moiety has a branched or straight chain and has 1 to 6,
15 preferably 1 to 4, carbon atoms;

wherein a halogen atom is preferably a fluorine, chlorine, or bromine atom, preferably a fluorine or chlorine atom.


20 Examples of particularly preferred compounds according to formula I for use in the invention are







5 and



The H^+ , K^+ -ATPase inhibitor used in the invention is preferably of formula (Ia): in other words it is preferably omeprazole, or an alkaline salt of omeprazole, the (-)-enantiomer of omeprazole or an alkaline salt thereof.

The compound of formula (I) when optionally in the form of a pharmaceutically acceptable alkaline salt is preferably the Mg^{2+} , Ca^{2+} , Na^{+} or K^{+} salt, more preferably the Mg^{2+} salt.

The H^{+} , K^{+} -ATPase inhibitor used in the invention can be administered orally, rectally or parenterally. While the effect of the inhibitors on the nasal polyps has been established in patients who have taken omeprazole by the oral route, it is believed that the effect of the inhibitor on the polyps is a systemic effect that is not dependent on what mode of administration is used. Accordingly a reduction in size of the polyps should be obtainable with other routes of administration.

Commercially available pharmaceutical preparations of H^{+} , K^{+} -ATPase inhibitors are suitably used in the invention. Examples of such preparations for omeprazole include enteric coated pellets of omeprazole filled in capsules, or formulated into a multiple unit tableted dosage form; enteric coated tablets of omeprazole or an alkaline salt thereof; and solutions for parenteral administration comprising an alkaline salt of omeprazole.

The dose of the H^{+} , K^{+} -ATPase inhibitor to be administered will vary according to the type of nasal polyps to be treated and the condition of the patient. However the dosage for oral, rectal or i.v. administration is generally in the range of from 1 to 100 mg of H^{+} , K^{+} -

ATPase inhibitor per day. Normally an amount of from 10 to 40 mg per day is used for oral administration.

The invention may be applied in combination with other treatments known to ameliorate the other symptoms generally associated with nasal polyps, for example asthma. In other words, the invention can be applied in the treatment of Widal's Syndrome which consists of the conditions of nasal polyps, asthma and aspirin intolerance. The invention may also be applied in the treatment of other inflammatory diseases in the upper respiratory tract such as acute and chronic rhinosinusitis, allergic and non-allergic rhinitis, as well as in the lower respiratory tract such as asthma. Therefore according to the invention there is further

provided a method for treating Widal's Syndrome and other respiratory tract inflammatory diseases which method comprises simultaneously, separately or sequentially administration to a subject suffering from the syndrome or the diseases a pharmaceutical formulation comprising an H^+ , K^+ -ATPase inhibitor and a glucocorticoid. According to the invention
 5 there is also provided a pharmaceutical formulation for simultaneous, separate or sequential administration to be used in the treatment of Widal's Syndrome or in the treatment of asthma which formulation comprises an H^+ , K^+ -ATPase inhibitor and a glucocorticoid. The invention further provides the use of an H^+ , K^+ -ATPase inhibitor and a glucocorticoid in the manufacture of such a pharmaceutical formulation.

10

Preferred glucocorticoids are topically active anti-inflammatory steroids. Examples of suitable steroids include budesonide; rofleponide; rofleponide palmitate; ciclesonide; mometasone furoate; fluticasone propionate; 16α , 17α -butylidenedioxy- 6α , 9α -difluoro- 11β , 21 -dihydroxypregna- $1,4$ -diene- $3,20$ -dione; 6α , 9α -difluoro- 11β -hydroxy- 16α , 17α -
 15 dibutylidenedioxy- 17α -methylthio-androsta- 4 -ene- 3 -one; S-methyl- 16α , 17α -butylidenedioxy- 6α , 9α -difluoro- 11β -hydroxy- 3 -oxo-androsta- $1,4$ -diene 17β -carbothioate; methyl 9α -chloro- 6α -fluoro- 11α -hydroxy- 16α -methyl- 3 -oxo- 17α -propionyloxy-androsta- $1,4$ -diene- 17α -carboxylate; 6α , 9α -difluoro- 11β -hydroxy- 16α -methyl- 3 -oxo- 17α -propionyloxy-androsta- $1,4$ -diene- 17β -carbothioic acid S-(2 -oxo-
 20 tetrahydro-furan- $3S$ -yl) ester; tipredane; fluocinolone acetonide; flunisolide; flumethasone; dexamethasone; betamethasone; beclomethasone dipropionate; deflazacort; cortivazol; or cortisol and/or hydrocortisol, optionally in their pure isomeric forms (where such forms exist) and in the forms of their pharmaceutically acceptable salts.

25 The steroids for use in the invention may be applied using conventional dosing rates, e.g. 40 to $3000 \mu\text{g}$ per day. Administration may be by inhalation orally or intranasally. The steroids can optionally be adapted to be administered from a dry powder inhaler, a pressurised metered dose inhaler, or a nebuliser.

When the steroids are administered from a pressurised inhaler, they are preferably in micronised form. They are suspended or dissolved in a liquid propellant mixture. The propellants which can be used include chlorofluorocarbons, hydrocarbons or hydrofluoroalkanes. Especially preferred propellants are P134a (tetrafluoroethane) and
5 P227 (heptafluoropropane) each of which may be used alone or in combination. They are optionally used in combination with other propellants and/or surfactants and/or other excipients, for example ethanol, surfactants, lubricants, anti-oxidants and stabilising agents.

When the steroids are administered via a nebuliser they may be in the form of a nebulised
10 aqueous suspension or solution, with or without a suitable pH or tonicity adjustment, either as a unit dose or multidose device.

The invention is described more in detail with reference to the following examples.

15 *Example 1*

A 53 year old woman who had had Widal's Syndrome for several years but who had refused surgical treatment of her polyps suffered mild upper abdominal pain and no improvement in the polyps after treatment by topical and systemic corticosteroids.
20 However within two weeks of being prescribed 20 mg of omeprazole per day in addition to 100 µg of budesonide (Aqua preparation) per nostril/b.i.d. and 6 mg of deflazacort per day, she experienced a progressive improvement in her nasal respiratory problem. Eventually she recovered completely from the polyps.

25 *Examples 2 to 10*

Nine patients, each with the conditions shown in the following Table 1, were treated during a two week period. The treatment consisted of 20 mg of omeprazole, 100 µg of intranasal budesonide and 3 to 15 mg of oral deflazacort (deflazacort was used in such a small
30 quantity to ensure the patients' compliance). The results are also shown in Table 1.

Table 1

Example No.	Condition	Result
2	Widal's Syndrome	Temporary benefit
3	Widal's Syndrome	Positive effect
4	Widal's Syndrome	Positive effect
5	Nasal Polyposis	Long term benefit
6	Nasal Polyposis	No benefit
7	Nasal Polyposis	Positive effect
8	Nasal Polyposis	Positive effect
9	Nasal Polyposis and aspirin intolerance	Positive effect
10	Nasal Polyposis and asthma	No benefit

Where a positive effect is indicated, this means that the patient experienced a decrease in rhinorrhoea, a marked improvement in nasal respiratory ventilation and a reduction in the size of the polyps. The patient who experienced a temporary benefit by the treatment suffered a recurrence of the polyposis following the withdrawal of omeprazole and deflazacort (topical anti-inflammatory steroids, i.e. budesonide, were taken as required). However after treatment with the same regimen was resumed, a marked reduction in the size of the polyps was achieved.

10


The patient of Example 5 who experienced a long term benefit initially experienced no benefit at the end of the initial 2 week treatment period but continued with omeprazole and was rewarded with a positive effect at the end of 2 months.

23940-1065


11

CLAIMS:


1. Use of an H^+,K^+ -ATPase inhibitor and a glucocorticoid in the manufacture of a pharmaceutical formulation intended for simultaneous, separate or
5 sequential administration in the treatment of Widal's Syndrome.
2. Use of an H^+,K^+ -ATPase inhibitor and a glucocorticoid in the manufacture of a pharmaceutical formulation intended for simultaneous, separate or
10 sequential administration in the treatment of nasal polyps.
3. Use according to claim 1 or 2, wherein the glucocorticoid is a topically active anti-inflammatory steroid.
4. Use according to claim 3, wherein the
15 glucocorticoid is budesonide, beclomethasone dipropionate or fluticasone propionate.
5. Use according to any one of claims 1 to 4, wherein the H^+,K^+ -ATPase inhibitor is a compound of formula:



20


wherein Het_1 is

or




23940-1065


12

Het₂ is

5 and X is



or



wherein N in the benzimidazole moiety of Het₂ means
that one of the ring carbon atoms substituted by R₆-R₉
10 optionally may be exchanged for a nitrogen atom without any
substituents;

R₁ and R₃ each independently represent hydrogen,
alkyl, or alkoxy on the condition that R₁ and R₃ do not
simultaneously represent alkoxy; and R₂ represents alkyl,
15 alkoxy optionally substituted by fluorine, alkylthio or
alkoxyalkoxy; or one of R₁ and R₃ is halogen and the other is
hydrogen and R₂ is 1-morpholino, 1-piperidino or
dialkylamino;

R₄ and R₅ are the same or different and selected
20 from hydrogen and alkyl;

R₆-R₉ are the same or different and selected from
hydrogen, halogen, alkyl, alkoxy, haloalkoxy, alkylcarbonyl,
and alkoxycarbonyl;

R₁₀ is hydrogen or R₁₀ and R₃ together complete a
25 ring containing 6 to 8 carbon atoms; and

R₁₁ represents hydrogen, halogen or alkyl;


23940-1065

13

wherein the compound of formula (I) is optionally in the form of a pharmaceutically acceptable alkaline salt or in its neutral form or is a single enantiomer or a racemic mixture thereof;

5 wherein each alkyl or alkylenyl moiety has a branched or straight chain and has 1 to 6 carbon atoms.

6. Use according to claim 5, wherein the compound of formula (I) is a compound of formula



or an alkaline salt thereof, or the (-)-enantiomer or an alkaline salt of the (-)-enantiomer.

7. Use of an H^+, K^+ -ATPase inhibitor and a
15 glucocorticoid for the treatment of Widal's Syndrome.

8. Use of an H^+, K^+ -ATPase inhibitor and a glucocorticoid for the treatment of nasal polyps.

9. Use according to claim 7 or 8, wherein the H^+, K^+ -ATPase inhibitor and the glucocorticoid are used
20 simultaneously.

10. Use according to claim 7 or 8, wherein the H^+, K^+ -ATPase inhibitor and the glucocorticoid are used separately.

11. Use according to claim 7 or 8, wherein the H^+, K^+ -ATPase inhibitor and the glucocorticoid are used
25 sequentially.

23940-1065

14

12. Use according to any one of claims 7 to 11, wherein the glucocorticoid is a glucocorticoid defined in claim 3 or 4.

13. Use according to any one of claims 7 to 12,
5 wherein the H^+,K^+ -ATPase inhibitor is a compound of the formula (I) defined in claim 5 or 6.

14. A pharmaceutical formulation for simultaneous, separate or sequential use in the treatment of Widal's Syndrome which formulation comprises an H^+,K^+ -ATPase
10 inhibitor and a glucocorticoid.

15. A pharmaceutical formulation for simultaneous, separate or sequential administration in the treatment of nasal polyps which pharmaceutical formulation comprises an H^+,K^+ -ATPase inhibitor and a glucocorticoid.

15 16. A pharmaceutical formulation according to claim 14 or 15, wherein the glucocorticoid is a glucocorticoid defined in claim 3 or 4.

17. A pharmaceutical formulation according to any one of claims 14 to 16, wherein the H^+,K^+ -ATPase inhibitor is a
20 compound of the formula (I) defined in claim 5 or 6.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA

PATENT AGENTS