

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 March 2009 (12.03.2009)

PCT

(10) International Publication Number
WO 2009/031896 A2

- (51) **International Patent Classification:**
A61K 9/127 (2006.01)
- (21) **International Application Number:**
PCT/NL2008/050587
- (22) **International Filing Date:**
5 September 2008 (05.09.2008)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
07115897.6 7 September 2007 (07.09.2007) EP
- (71) **Applicant (for all designated States except US):** SYN-VOLUX IP B.V. [NL/NL]; Zielstraweg 1, NL-9713 GX Groningen (NL).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** KAMPS, Johannes Adrianus Antonius Maria [NL/NL]; Galkemaheerd 37, NL-9736 BE Groningen (NL). MOLEMA, Grietje [NL/NL]; H.W. Mesdagstraat 53, NL-9718 HD Groningen (NL). RUITERS, Marcel Herman Josef [NL/NL]; Cunera van Wahrenlaan 17, NL-9766 PJ Eelderwolde (NL). ADRIAN, Joanna Ewa [NL/NL]; Grote Lelystraat 97, NL-9712 SR Groningen (NL).
- (74) **Agent:** HATZMANN, M.J.; Vereenigde, Johan de Wittlaan 7, NL-2517 JR Den Haag (NL).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report



WO 2009/031896 A2

(54) **Title:** IMPROVED LIPOSOMES AND USES THEREOF

(57) **Abstract:** The invention relates to the field of molecular medicine and pharmacology. More specifically, it relates to liposomes and their use as delivery vehicle for therapeutic compounds. Provided is a liposome comprising at least one lipid bilayer enclosing an interior compartment, wherein said lipid bilayer comprises at least one synthetic pyridinium-derived amphiphile, for instance a Saint-molecule.

Title: Improved liposomes and uses thereof.

The invention relates to the field of molecular medicine and pharmacology. More specifically, it relates to liposomes and their use as delivery vehicle for therapeutic compounds. Also provided is a method for preparing liposomes.

5 The therapeutic benefit of many compounds is limited by low uptake of the compound by the target cells. Generally, for maximum therapeutic benefit, delivery of the compound to the cytoplasmic or nuclear compartment of the cell is desired, where gene transcription, and translation of mRNA into protein take place. Lipophilic compounds, including many small, uncharged
10 compounds, can permeate across the cell membrane and allow relatively efficient uptake by the cell. However, for a variety of larger and/or charged compounds, such as proteins, nucleic acids, and highly water soluble charged organic compounds, passive uptake by permeation across the cell membrane is limited.

15 Several methods for improving uptake of such compounds into cells have been proposed. For example, a drug can be administered in a modified or prodrug form for transport into cells and then undergo enzymatic conversion to an active form within the cells. However, it will be understood that the preparation of a suitable prodrug form is not possible for each and every
20 therapeutic compound.

 Alternatively, the cellular processes of phagocytosis or endocytosis can be used, where drug-containing particles are taken up by the target cells. With most of the currently available drug delivery particles, this approach is however limited to certain cell types, for example, phagocytosis is limited to
25 cells of monocyte lineage and to certain other myeloid cells, such as neutrophils, while endocytosis is limited in a number of other cells, among other cells from mesenchymal lineage, such as vascular endothelial cells,

epithelial cells and fibroblasts. Thus, unlike macrophages and other cells that are specialized in scavenging and processing particles from the blood, there are many cells that are not or much less equipped for handling drug-containing particles.

5 Still another approach to enhancing drug uptake by cells involves the use of fusogenic particles designed to fuse with the surface membrane of a target cell, releasing the particle's contents into the cytoplasmic compartment of the cell. Inactivated and reconstituted virus particles have been proposed for this purpose, particularly in gene therapy where large nucleic acid strands are
10 introduced into cells. Virus-like particles composed of fusion-promoting viral proteins embedded in an artificial lipid bilayer represent another example. However, safety concerns and the expenses associated with growing, isolating, and deactivating viral components limit a broad application of these approaches.

15 Endothelial cells, covering the vascular wall of all blood vessels, are an example of cells that are hampered with respect to the handling of drug-containing particles, e.g., drug-loaded liposomes. Endothelial cells play a pivotal role in whole body homeostasis and the patho(physio)logy of important diseases, including (chronic) inflammation and cancer. They are also attractive
20 target cells /target tissue for drug delivery as they are fully accessible for any compound transported by the blood. In addition, the heterogeneity of the endothelium with respect to appearance and function allows for tissue and disease-specific drug delivery approaches. Endothelial cells are a promising target for cancer therapy because angiogenesis is vital for the supply of oxygen
25 and nutrients to solid tumours. Moreover, in inflammatory diseases they guide the movement of white blood cells from the blood into the tissue, and as such are indispensable for disease development. The present inventors observed that liposomes targeted to E-selectin or VCAM-1 expressed on TNF α -activated endothelial cells are readily endocytosed, in amounts that are comparable to
30 the endocytosis capacity of macrophages. However, unlike macrophages,

endothelial cells do not extensively process or degrade the liposome and hence accumulate the drug-loaded vehicle inside vesicular bodies. The absence of liposome degradation and/or destabilization results in retention of the encapsulated drug in the liposome and, consequently, in inferior
5 pharmacological efficacy.

Therefore, a goal of the present invention is to provide a lipid-based drug delivery vehicle that allows for an improved intracellular drug release in target cells as compared to known delivery vehicles. A specific aim is to provide a lipid-based drug delivery vehicle for an efficient intracellular drug release in
10 target cells that are not specialized in scavenging and/or processing lipid based-particles, such as endothelial cells and epithelial cells, muscle cells, brain cells, nerve cells, skin cells, hair cells, subsets of progenitor cells, and pericytes and all other cells not specialized in processing delivery vehicles.


Using a method which distinguishes between a) liposome uptake by a
15 target cell and b) drug release from a liposome in a target cell, the present inventors surprisingly found that the incorporation of a synthetic pyridinium-derived amphiphile in the bilayer of a "classical" or "traditional" liposome (i.e. an aqueous compartment surrounded by a lipid bilayer) yields a lipid-based delivery vehicle that ensures efficacious intracellular drug delivery in cells
20 that are not specialized in processing lipid-based particles. Only liposomes comprising a bilayer comprising one or more synthetic pyridinium-derived amphiphile(s) in an amount of 2 to 25 mol% based on the total lipid content were found to be sufficiently stable. Accordingly, the invention provides a liposome comprising at least one lipid bilayer enclosing an aqueous interior
25 compartment, wherein said lipid bilayer comprises at least one synthetic pyridinium-derived amphiphile in an amount of 2 to 25 mol% based on the total lipid content of the liposome.

The term "liposome" is known in the art to refer to an aqueous compartments enclosed by phospholipid bilayer membrane. Phospholipid
30 molecules consist of an elongated nonpolar (hydrophobic) structure with a

polar (hydrophilic) structure at one end. When dispersed in water, they spontaneously form bilayer membranes, also called lamellae, which are composed of two monolayer layer sheets of lipid molecules with their nonpolar (hydrophobic) surfaces facing each other and their polar (hydrophilic) surfaces facing the aqueous medium.


An amphiphile is a compound consisting of molecules having a polar water-soluble group attached to a water-insoluble hydrocarbon chain. Pyridinium refers to the cationic form of pyridine. This can either be due to protonation of the ring nitrogen or because of addition of a substituent to the ring nitrogen, typically via alkylation. The lone pair of electrons on the nitrogen atom of pyridine is not delocalized, and thus pyridine can be protonated easily. The expression "pyridinium-derived" as used herein refers to any amphiphile having a pyridinium-moiety in its polar group.

Synthetic pyridinium-derived amphiphiles are known in the art. Of particular interest for use in the present invention is the group of synthetic pyridinium-derived amphiphiles covered by Synvolux' proprietary technology, SAINT (Synthetic, Amphiphilic, INTeractive). See for instance European Patent EP0755924-B1. In one embodiment of the present invention, at least one synthetic pyridinium-derived amphiphile is selected from the group of SAINT compounds with the general formula (I):



in which:


R₁ is a (C₁-C₅)alkyl, ar(alkyl) or an alkyl group with a cationic functional group, like



5 or R₁ is (C₁-C₅ alkylene) R₅ in which R₅ is a structure with the general formula I except R₁;


X is a halide counter ion, chosen from Cl⁻, I⁻, Br⁻;

R₃ is hydrogen and R₂ and R₄ are identical or different and are chosen from the group, comprising branched or linear (C₁₀-C₂₀)alkyl, a mono- or polyunsaturated (C₁₀-C₂₀)alkenyl, O=C-O-alkyl,



or ar(alkyl),

or R₂ and R₄ are hydrogen and R₃ is -CH(R₅)₂ with R₅, comprising (C₁₀-C₂₀)alkyl, mono- or polyunsaturated (C₁₀-C₂₀)alkenyl, O=C-O-alkyl,



or aralkyl.

In a further embodiment, a SAINT molecule as defined above is incorporated, wherein disclaimed are the compounds with the general formula I in
20 which R₁ is CH₃, R₂ and R₄ are hydrogen, R₃ is (C₁₆H₃₃)₂CH and X is all mentioned counter ions and disclaimed are the compounds in which R₁ is CH₃, R₂ and R₄ are C₁₆H₃₃-O-C(O), R₃ is hydrogen and X is all mentioned counter ions.

Liposomes according to the present invention comprising a SAINT molecule are also referred to as "SAINT-o-somes". Specific embodiments of the present invention use SAINT-18 (1-methyl-4-(cis-9-dioleyl)methyl-pyridinium-chloride) and/or SAINT-12 (1-methyl-4-(pentacosan-13-yl)-pyridinium-chloride).

The incorporation of synthetic pyridinium-derived amphiphiles, such as SAINT, in traditional liposomes has not been described or suggested in the art. In contrast, SAINT molecules have thus far been used in non-liposomal delivery vehicles (called "lipoplexes") for the delivery of macromolecules such as nucleic acids or proteins to mammalian cells. EP-A-0755924 for example discloses complexing of the macromolecule to SAINT, optionally in a 1:1 mixture with a helper lipid, to form a lipoplex particle of up to several μm in size. In the lipoplex formed only non-covalent interactions are present between SAINT and the macromolecule. The cationic amphiphiles on the surface of the particle have high affinity for the negatively charged cell surface. Whereas the introductory section of EP-A-0755924 refers briefly to "liposomes which consist of a bilayer of phospholipids", it must be emphasized that the invention disclosed in EP-A-0755924 clearly does not relate to such liposomes. Rather, it relates to the lipoplexes whose structural organization was studied in detail by Oberle *et al.* using atomic force microscopy (2000, Biophysical Journal 79(3), 1447-1454). Figure 7 of Oberle *et al.* illustrates that the SAINT-containing lipoplexes, as disclosed for example in EP-A-0755924, consist of DNA wrapped into several layers of amphiphile, which subsequently merge into a large complex. It is of note that some DNA "sticks out" of the lipoplex and is this not shielded from the environment as it would be when incorporated in the interior compartment of a SAINT-containing liposome as disclosed in the present invention. Rejman *et al.* (Biochimica et Biophysica Acte 1660 (2004) 41-52), Zuhorn *et al.* (Biochimica et Biophysica Acta 1560 (2002) 25-36), Van de Woude *et al.* (Proc. Natl. Acad. Sci. USA, vol. 94, pp. 1160-1165, 1997) and

WO2006/043809 likewise disclose lipoplexes rather than liposomes comprising an aqueous interior compartment.

In contrast, the present invention relates to liposomes comprising an aqueous interior space surrounded by a lipid bilayer, wherein one or more synthetic pyridinium-derived amphiphiles are incorporated. Without wishing to be bound by any theory, the properties of the cationic pyridinium headgroup of the amphiphile makes the liposomal lipid bilayer less rigid and more prone to lipid mixing (e.g. fusing) with intracellular, endosomal membrane lipids, thereby promoting intracellular release of the encapsulated compound. As is demonstrated herein below, this effect is not observed for just any cationic amphiphile, since liposomes with similar amounts of DOTAP (N-[1-(2,3-Dioleoyloxy)]-N,N,N-trimethylammonium propane) a well known cationic liposomal transfection reagent, exhibited inferior performance.

Liposomes according to the invention have one or more pyridinium-derived amphiphiles. The exact composition of the liposomes will depend on the particular circumstances for which they are to be used. Those of ordinary skill in the art will find it a routine matter to determine a suitable lipid composition. General information on liposomes can for example be found Lasic, D.D. "Liposomes: from physics to application", Elsevier Science B.V. Amsterdam, 1993 (ISBN 0444895485).

The relative amount of pyridinium-derived amphiphile in a liposome of the invention can vary, but is typically up to about 5-25 mole% based on the total amount of other lipid constituents of the liposome. This ensures that the liposome is stable outside the target cell (e.g., in the blood stream) while it allows for efficient intracellular release of its encapsulated contents upon mixing or fusion with a target cell membrane. Very good results were obtained when liposomes comprising 5-20 mole%, preferably 10-20 mole%, such as 11, 12, 13, 14, 15, 16, 17, 18 19 or 20 mole%, pyridinium-derived amphiphile relative to the other lipids was used. Specific embodiments include liposomes

comprising 10-20 mole% SAINT molecules, such as 10, 11, 12, 14, 15, 17, 19 or 20 mole% SAINT-18 or SAINT-12.

In one embodiment, the liposomes of the present invention consist essentially of a single type of lipid in addition to the at least one pyridinium-
5 derived amphiphile. In a preferred embodiment, the liposomes comprise a mixture of two or more lipids, preferably selected from the group consisting of glycerophospholipids, sphingolipids and cholesterol. Naturally occurring and/or synthetic lipids may be used. In fact, any type of liposome-forming lipid or molecule may be used in combination with at least one pyridinium-derived
10 amphiphile to form a liposome according to the invention. The lipid constituting the liposomes of the present invention includes phosphatidylcholines, phosphatidylethanolamines, phosphatidic acids, gangliosides, glycolipids, phosphatidylglycerols, and cholesterol. The phosphatidylcholines preferably include dimyristoylphosphatidylcholine,
15 dipalmitoylphosphatidylcholine, palmitoyloleoylphosphatidylcholine (POPC) and distearoylphosphatidylcholine. The phosphatidylethanolamines preferably include dimyristoylphosphatidylethanolamine, dipalmitoylphosphatidylethanolamine, and distearoylphosphatidylethanolamine. The phosphatidic acids preferably include dimyristoylphosphatidic acid, dipalmitoylphosphatidic
20 acid, distearoylphosphatidic acid, and dicetylphosphoric acid. The gangliosides preferably include ganglioside GM1, ganglioside GD1a, and ganglioside GT1b. The glycolipids preferably include galactosylceramide, glucosylceramide, lactosylceramide, phosphatide, and globoside. The phosphatidylglycerols preferably include dimyristoylphosphatidylglycerol,
25 dipalmitoylphosphatidylglycerol, and distearoylphosphatidylglycerol.

In vivo studies have shown that conventional liposomes are rapidly removed from the circulation by cells of the reticuloendothelial system, i.e., tissue-resident phagocytes present in a number of organs, particularly the spleen and the liver. This interaction is reduced when using sterically
30 stabilized liposomes, which can be prepared by insertion of

monosialoganglioside (GM1) or poly(ethylene glycol) (PEG) derivatized lipids within the lipid bilayer of liposomes. Such liposomes coated with inert polymers show a substantial improvement in their blood circulation half life (in human in the order of days as opposed to minutes for conventional liposomes).

Sterically stabilized immunoliposomes with cell specific recognition properties are often prepared by coupling of antibodies to the distal ends of PEG chains. Using the PEG chains as linker between the liposome and antibody leads to an enhanced antibody-antigen recognition and binding since the antibody is not shielded by the steric barrier activity of PEG. Several covalent coupling methods have been developed for attaching (derivatized) antibodies at the PEG terminus. They make use of functionalized PEG-lipids with a chemically reactive endgroup such as hydrazide, N-(3'-(pyridyldithio)propionate, maleimide, succinyl, p-nitrophenylcarbonyl or cyanuric chloride. Thus, in one embodiment of the invention a derivatized (e.g., PEGylated) phospholipid may be used to improve the *in vivo* circulation time of a liposome formulation, and/or to allow for coupling of proteins to the liposomal surface. In a specific aspect, a liposome of the present invention, comprises a mixture of phosphatidylcholine and cholesterol in relative amounts from 1:1 to 2:1 (mole%), optionally in a mixture with one or more natural or synthetic lipids, for example a PEG-ylated phospholipid.

The liposome itself can be produced through any conventional method including a thin film method, a reverse phase evaporation method, an ethanol injection method, and a dehydration-rehydration method. Accordingly, the invention provides a method for preparing a liposome according to the invention, comprising mixing conventional lipids with a suitable amount of at least one synthetic pyridinium-derived amphiphile, and preparing a liposome according to conventional procedures. Also provided are liposomes obtainable by a method of the invention.

For example, a mixture of the above-mentioned lipids, from which the solvents have been removed, can be emulsified by the use of a homogenizer, lyophilized, and melted to obtain multilamellar liposomes. Alternatively, unilamellar liposomes can be produced by the reverse phase
5 evaporation method (Szoka and Papahadjopoulos, 1978, Proc. Natl. Acad. Sci. USA 75:4194-4198). Unilamellar vesicles can also be prepared by sonication or extrusion. Sonication is generally performed with a bath-type sonifier, or a Branson tip sonifier at a controlled temperature as determined by the melting point of the lipid.

10 Following liposome preparation, the liposomes that have not been sized during formation may be sized by extrusion to achieve a desired size range and relatively narrow distribution of liposome sizes. The particle size of the liposome can be controlled through an ultrasonic radiation method, an extrusion method, a French press method, a homogenization method or any
15 other suitable conventional method.

Extrusion may be carried out by biomembrane extruders, such as the Lipex Biomembrane Extruder (Northern Lipids Inc, Vancouver, British Columbia, Canada). Defined pore size in the extrusion filters may generate unilamellar liposomal vesicles of specific sizes. A size range of about 200-400
20 nm will allow the liposome suspension to be sterilized by filtration through a conventional filter (e.g., a 0.22 micron filter). The filter sterilization method can be carried out on a high throughput basis. The liposomes can also be formed by extrusion through an asymmetric ceramic filter, such as a Ceraflow Microfilter (commercially available from the Norton Company, Worcester,
25 Mass.).

For therapeutic applications, it is preferred that a liposome according to the invention has a size (i.e. diameter) of up to about 300 nm. In one embodiment, the liposome size ranges from between about 20 to 250 nm, preferably 30-200 nm, such as 50, 60, 75, 100, 120, 135, 150 or 180 nm. A wide
30 variety of liposomes may be used in the invention (including oligo- or

multilamellar vesicles), but preferably the liposomes are small unilamellar vesicles (SUVs) having an external diameter of from about 30 to about 300 nanometers (nm), most preferably 50 to 150 nm.

A liposome of the invention may comprise one or more other
5 beneficial constituents, for example means for targeting the liposome to a specific cell. Site-specific delivery of drugs to diseased cells can lead to increased therapeutic effects and to a significant reduction of toxicity. Drug targeting by antibody-conjugated liposomes or immunoliposomes represents a technology which has been applied to the targeting of specific sites of drug
10 action such as the brain, lung, cancer cells, HIV-infected cells or cells of the immune system. Liposomal targeting means, or targeting ligands, are known in the art, and include proteins, peptides, such as antibodies or fragments thereof, having specificity for surface antigen of a particular target cell or for a specific intracellular compartment, such as mitochondria. Liposomes provided
15 at their surface with a targeting antibody are referred to in the art as immunoliposomes. Site specific targeting is often mediated by the high affinity binding of monoclonal antibodies to their specific antigens. US patent no. 5,258,499 discloses delivery vehicle formulations comprising active agents encapsulated within liposomal vesicles to which are attached protein hormones
20 (ligands) such as interleukin-2. The ligands are capable of showing affinity for specific cell receptors resulting in delivery of the encapsulated active agent to target cells, enabling delivery of active agents to particular cell populations in the treatment of conditions such as immune system disorders. In a preferred embodiment, the liposome carrying a targeting moiety, e.g., a ligand, is
25 internalized by a target cell. In yet another embodiment, a targeting moiety is a ligand that specifically interacts with a tyrosine kinase receptor such as, for example, EGFR, HER2, HER3, HER4, PD-GFR, VEGFR, bFGFR or IGFR receptors. In still another embodiment, the targeting moiety specifically interacts with a growth factor receptor, an angiogenic factor receptor, a
30 transferrin receptor, a scavenger receptor, a cell adhesion molecule, or a

vitamin receptor. Exemplary cell-specific ligands include the RGD-peptide, NGR-peptide, ATWLPPR-peptide, APRPG-peptide, SMSIARL-peptide, TAASGVRSMH-peptide, LTLRWVGLMS-peptide, CDSDSITWDQLWDLMK-peptide, GPLPLR-peptide, HWGF-peptide, recombinant VEGF, antibodies and
5 monoclonal antibodies, bispecific antibodies and single chain fragments against e.g., E-selectin, VCAM-1, endoglin, MHCII, VEGF:VEGFR complex, $\alpha\beta_3$, moc-31, cd-90, and other cellular target epitopes. Furthermore receptor specific ligand such as transferrin, apolipoprotein E, lactoferrin, modified albumins etc.

10 In one embodiment, a liposome comprises at least one targeting ligand that is bound via a linker-molecule to a SAINT-molecule incorporated in the liposomal lipid bilayer. See for exemplary SAINT-linker-ligand molecules WO2006/043811. Preferably, a liposome according to the invention comprises an unmodified (i.e. not comprising a a targeting means) SAINT molecule as
15 well as a SAINT molecule to which at least one cell or organlle-specific ligand is bound via a linker.

In view of sterical hindrance, the relative amount of modified versus unmodified SAINT molecule is preferably less than 5%, more preferably less than 1%, such as approximately 0,1%. For example, liposomes are prepared
20 comprising as bilayer constituents 35.8 mol% POPC, 40 mol% cholesterol, 4 mol% PEG-DSPE, 20 mol% SAINT C18 and 0.2 mol% SAINT C18-linker.

As mentioned above, a liposome of the invention comprises an interior space that can be used to deliver one or more molecules of interest, for instance a biologically active compound, to a target cell. For the purposes of
25 this invention, the term "biologically-active compound" is intended to encompass all naturally-occurring or synthetic compounds capable of eliciting a biological response or having an effect, either beneficial or deleterious, including cytotoxic, on biological systems, particularly tissues, cells and cellular organelles. These compounds are intended to include all varieties of
30 drugs, including antibiotic, antibacterial, antiviral, antimycotic, anti-

inflammatory, antiproliferative and antineoplastic drugs, and any inhibitor of intracellular signal transduction developed as therapeutics, such as a MAPK or SAPK inhibitor; hormones, including peptide hormones and steroid hormones; genes, recombinant nucleic acids, oligonucleotides or other nucleic acids encoding all or a portion of a mammalian gene, including custom small
5 interfering RNA (also known as short interfering RNA or siRNA) and specially engineered short RNA molecules which can effectively silence the action of microRNA's in regulating gene expression, such as antagomirs, a viral gene or a gene from a microorganism; antigens; enzymes; nutrients; and most
10 particularly any biologically active compound, that is inefficiently taken up by passive permeation across the cell membrane of a target cell of interest. In one aspect, the liposome comprises in its interior compartment a biologically active compound having a neutral or positive net charge, for instance a proteinaceous substance.

15 A further aspect of the present invention relates to compositions comprising a liposome according to the invention. In general, the liposome composition of the present invention is quite stable during storage, e.g., as measured by the percentage of entrapped entity released outside of the liposomes or still maintained inside of the liposomes after a certain time period
20 from the initial loading of the entity inside the liposomes of the present invention. For example, the liposome composition of the present invention is stable at 4° C for at least 6 months, e.g., less than 10% of entrapped entity is released 6 months after the initial loading of the entity. In one embodiment, the liposome composition of the present invention is stable at 4°C for at least 2
25 years, e.g., less than 20% of entrapped entity is released 2 years after the initial loading of the entity.

According to another embodiment of the present invention, the liposome composition of the present invention can be provided as a pharmaceutical composition containing the liposome composition of the
30 present invention and a carrier, e.g., pharmaceutically acceptable carrier.

Examples of pharmaceutically acceptable carriers are normal saline, isotonic dextrose, isotonic sucrose, Ringer's solution, and Hanks' solution. A buffer substance can be added to provide pH optimal for storage stability. For example, pH between about 6.0 and about 7.5, more preferably pH about 6.5, is
5 optimal for the stability of liposome membrane lipids, and provides for excellent retention of the entrapped entities. Histidine, hydroxyethylpiperazine-ethylsulfonate (HEPES), morpholino-ethylsulfonate (MES), succinate, tartrate, and citrate, typically at 2-20 mM concentration, are exemplary buffer substances. Other suitable carriers include, e.g., water,
10 buffered aqueous solution, 0.4% NaCl, 0.3% glycine, and the like. Protein, carbohydrate, or polymeric stabilizers and tonicity adjusters can be added, e.g., gelatin, albumin, dextran, or polyvinylpyrrolidone. The tonicity of the composition can be adjusted to the physiological level of 0.25-0.35 mol/kg with glucose or a more inert compound such as lactose, sucrose, mannitol, or
15 dextrin. These compositions may be sterilized by conventional, well known sterilization techniques, e.g., by filtration. The resulting aqueous solutions may be packaged for use or filtered under aseptic conditions and lyophilized, the lyophilized preparation being combined with a sterile aqueous medium prior to administration.

20 The pharmaceutical liposome compositions can also contain other pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, etc. Additionally, the
25 liposome suspension may include lipid-protective agents which protect lipids against free-radical and lipid-peroxidative damages on storage. Lipophilic free-radical quenchers, such as alpha-tocopherol and water-soluble iron-specific chelators, such as ferrioxamine, are suitable.

The concentration of the liposomes of the present invention in the
30 fluid pharmaceutical formulations can vary widely, i.e., from less than about

0.05% usually or at least about 2-10% to as much as 30 to 50% by weight and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected. For example, the concentration may be increased to lower the fluid load associated with treatment. This may
5 be particularly desirable in patients having atherosclerosis-associated congestive heart failure or severe hypertension. Alternatively, liposome pharmaceutical compositions composed of irritating lipids may be diluted to low concentrations to lessen inflammation at the site of administration.

The amount of liposome pharmaceutical composition administered
10 will depend upon several factors, for example the particular therapeutic entity entrapped inside the liposomes, the disease state being treated, the type of liposomes being used, and / or the judgment of the clinician. Generally, the amount of pharmaceutical liposome composition administered will be sufficient to deliver a therapeutically effective dose of the particular therapeutic entity.

15 The quantity of pharmaceutical liposome composition necessary to deliver a therapeutically effective dose can be determined by routine *in vitro* and *in vivo* methods, common in the art of drug testing. See, for example, D. B. Budman, A. H. Calvert, E. K. Rowinsky (editors). Handbook of Anticancer Drug Development, LWW, 2003. Therapeutically effective dosages for various
20 therapeutic entities are well known to those of skill in the art; and according to the present invention a therapeutic entity delivered via the pharmaceutical liposome composition of the present invention provides at least the same, or 2-fold, 4-fold, or 10-fold higher activity than the activity obtained by administering the same amount of the therapeutic entity in a conventional
25 liposome formulation not comprising at least one synthetic pyridinium-derived amphiphile. Typically the dosages for the liposome pharmaceutical composition of the present invention range between about 0.005 and about 500 mg of the therapeutic entity per kilogram of body weight, most often, between about 0.1 and about 100 mg therapeutic entity/kg of body weight or targeted
30 organ weight.

In one embodiment, the pharmaceutical liposome composition of the present invention is prepared as a topical or an injectable, either as a liquid solution or suspension. However, solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared. The
5 composition can also be formulated into an enteric-coated tablet or gel capsule according to known methods in the art.

The liposome composition of the present invention can be administered in any way which is medically acceptable which may depend on the condition or disease being treated. Possible administration routes include
10 injections, by parenteral routes such as intramuscular, subcutaneous, intravenous, intra-arterial, intraperitoneal, intra-articular, intra-epidural, intrathecal, or others, as well as oral, nasal, ophthalmic, rectal, vaginal, topical, or pulmonary, e.g., by inhalation. For the delivery of liposomally drugs formulated according to the invention, to tumors of the central nervous system,
15 a slow, sustained intracranial infusion of the liposomes directly into the tumor (a convection-enhanced delivery, or CED) is of particular advantage. See Saito, et al., *Cancer Research*, vol. 64, p. 2572-2579, 2004; Mamot, et al., *J. Neuro-Oncology*, vol. 68, p. 1-9, 2004. The compositions may also be directly applied to tissue surfaces. Sustained release, pH dependent release, or other specific
20 chemical or environmental condition mediated release administration is also specifically included in the invention, e.g., by such means as depot injections, or erodible implants.

It is desirable that the average particle diameter is 400 nm or less in the case of its intravenous administration. This is because a liposome having a
25 particle diameter of exceeding 400 nm is cleared by the liver, spleen and the like reticuloendothelial system and the lungs.

Also provided is a method for facilitating the delivery of a biologically active compound to a target cell that is not specialized in scavenging and processing particles from the blood, comprising the
30 intravenous administration of a liposome according to the invention which

comprises said biologically active compound. A further aspects relates to the use of a liposome according to the invention as delivery vehicle to *in vitro* or *in vivo* introduce a substance of interest in a target cell that is not specialized in scavenging and/or processing lipid-based particles. Preferably, said target cell
5 is selected from the group consisting of endothelial cells, epithelial cells, muscle cells, brain cells, nerve cells, skin cells, hair cells, subsets of progenitor cells, and pericytes. More preferably, the target cell is an endothelial cell. A liposome and liposome-based delivery method of the invention for instance facilitates the (targeted) delivery of a biologically active compound to
10 endothelial cells, for instance TNF α -activated endothelial cells. Thus, a liposome as disclosed herein is advantageously used as drug delivery vehicle.

Also provided is the use of a liposome according to the invention comprising at least one biologically active compound, either in its interior
15 space and/or inserted in the lipid bilayer and/or attached covalently or otherwise to one of the liposome constituents, for the manufacture of a medicament for the treatment of cancer, chronic inflammation, tissue regeneration/repair, diabetes or metabolic disease associated cell dysfunction.

In one embodiment, a liposome provided with an EPCAM targeting
20 means and encapsulating a cytostatic compound is used for the manufacture of a medicament for the treatment of cancer. In another embodiment, a liposome provided with an E-selectin targeting means and comprising a p38MAPK inhibitor is used for the manufacture of a medicament for the treatment of glomerulonephritis or reuma.

25

LEGENDS TO THE FIGURES

Figure 1A: The intracellular processing of anti-E-selectin immunoliposomes in HUVEC targeted to E-selectin (HES) or VCAM-1 (VCAM) was investigated by using double radiolabeled conventional liposomes, i.e. without synthetic pyridinium-derived amphiphile, containing the metabolically degradable ester cholesteryl [¹⁴C]oleate in addition to the non-degradable [³H]cholesteryloleyl ether. In the case of proper metabolism, following endocytosis the cholesterylester is hydrolyzed and the liberated [¹⁴C]oleic acid will be released from the cells into the culture medium. The [³H]cholesteryloleyl ether will remain associated and thus the ³H/¹⁴C ratio is a convenient measure of liposome degradation [5]. As shown in panel A, the ³H/¹⁴C ratio does not change upon prolonged incubation, indicating the absence of hydrolysis of cholesteryl-[¹⁴C]oleate. Even a 24 h incubation did not result in any significant hydrolysis the double labeled anti-E-selectin immunoliposomes. In panel B, the degradation of immunoliposomes by HUVEC is compared to the degradation of similar liposomes by IC21 cells (peritoneal macrophages). It is shown that the same liposomes can efficiently be degraded by specialized cells. Incubations in the presence of inhibitors (chloroquine, NH₄Cl) of lysosomal degradation indicated that degradation was lysosomal. (data not shown).

Figure 2: TNF α activated HUVEC were incubated for 24 h with anti-E-selectin immunoliposomes (without synthetic pyridinium-derived amphiphile) that were fluorescently labeled with the lipid bilayer marker DiI. The liposomes have accumulated in distinct intracellular vesicles, while there is no indication of redistribution of the fluorescent label over cellular membranes, i.e., the liposomes are not degraded after being endocytosed by the endothelial cells.

Figure 3: Liposomes according to the invention were prepared as described in the general method and comprising the indicated amount of synthetic pyridinium-derived amphiphile (in this case SAINT-18) or the reference cationic amphiphile DOTAP. Liposomes were stored at 4°C under argon during

the time periods indicated. Liposome particle size was analyzed at the indicated times by dynamic light scattering using a Nicomp model 380 submicron particle analyzer in the volume weighing mode. Data are presented as single value for 1, or mean \pm sd of 2 to five independent liposome
5 preparations.

Figure 4: Liposomes were prepared as described in the general method. Liposomes were kept in thermostated water bath at 37°C or on the laboratory bench at room temperature (20°C) in the absence (panel A) or presence (panel
10 B) of 10% (v/v) serum . Particle size was analyzed at the indicated times by dynamic light scattering using a Nicomp model 380 submicron particle analyzer in the volume weighing mode. Data are presented as mean \pm sd of 3 or 4 independent liposome preparations.

Figure 5: Liposomes were prepared as described with increasing amounts of pyridinium-derived amphiphile. Liposomes are fully size stable when the bilayer is formulated with 0 to 20 mol% SAINT C18. Above around 20 to 25 mol% pyridinium-derived amphiphile, the liposomes start to loose their size
15 stability.

20

Figure 6: Calceine release from liposomes as a function of pH. Fluorescence of the liposomes was monitored while incubated in buffers with the indicated pH in 96 well plates in a standard fluorimeter. Calceine release was related to total fluorescence of the liposomes measured after treatment with Triton-X100.
25 Data are presented as relative release compared to liposomes without SAINT or DOTAP (mean \pm sd of 3 to 4 independent experiments).

Figure 7: Calceine release from liposomes formulated with 20 mol% SAINT or DOTAP at pH 7.4 and pH 5. Fluorescence of the liposomes was monitored in
30 buffers with the indicated pH in 96 well plates in a standard fluorimeter.

Calceine release was related to total fluorescence of the liposomes measured after treatment with Triton-X100. Data are presented as relative release compared to liposomes without SAINT or DOTAP (mean \pm sd of 3 to 4 independent experiments). * $p < 0.005$ compared to pH 7.4; # $p < 0.05$ compared to liposomes containing 20 mol% SAINT.

Figure 8: Effect of 10% serum (v/v) on calceine release from liposomes. Fluorescence of the liposomes was monitored in 96 well plates in a standard fluorimeter. Calceine release was calculated after determination of total fluorescence of the liposomes measured after treatment with Triton-X100. Data are presented as relative release compared to liposomes without SAINT or DOTAP (mean \pm sd of 3 to 4 independent experiments)

Figure 9: TNF α activated HUVEC were incubated for 24 h with liposomes with encapsulated calceine. Panel A: anti-E-selectin immunoliposomes formulated with 20 mol% SAINT. Panel B: anti-E-selectin immunoliposomes without SAINT. Panel C: untargeted liposomes with 20 mol% SAINT. Calceine release is most prominent with targeted liposomes that contain SAINT.

Figure 10: Uptake of DiI (panel A) and calcein (panel B) labeled liposomes by HUVEC. Non- and TNF- α stimulated cells were incubated with the indicated liposomes for 3h. Uptake of liposomes was analyzed by FACS. Data are representative for three independent experiments.

Figure 11: Effect of E-selectin targeted liposomes formulated with 20 mol% SAINT containing Interleukine-8 (IL-8) siRNA on expression of IL-8 in conditionally immortalized human glomerular endothelial cells (ciGEnC). TNF- α activated ciGEnC were incubated with E-selectin SAINT immunoliposomes containing siRNA (either specific for IL-8 or scrambled siRNA) for 48h.

Suppression of IL-8 expression was analyzed by real-time PCR. Presented data come from one experiment.

Figure 12: Down-regulation of VE-cadherin expression **Figure 11:** Effect of E-selectin targeted liposomes formulated with 20 mol% SAINT and containing siRNA on the H5V polyoma middle-T oncogene-transformed endothelioma cell line. E-sel Saint-o-somes containing specific siRNA for VE-cadherin and scrambled siRNA were incubated with H5V cells at the concentration of siRNA 1000 pmol/ml for 48h. The effects of siRNA were analysed by real-time PCR. The expression of VE-cadherin in cells treated with liposomes was compared to TNF- α treated cells (arbitrary set to 1). Data are presented as \pm SD.

15 EXAMPLES

Abbreviations used:

TNF α , tumor necrosis factor α

VCAM-1, vascular cell adhesion molecule 1

HUVEC, human umbilical vein endothelial cells

20 DiI, 1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate
SAINT, N-methyl-4-alkylpyridium chloride; SATA, (N-succinimidyl-S-acetylthioacetate)

DOTAP, N-[1-(2,3-Dioleoyloxy)]-N,N,N-trimethylammonium propane

25 **Liposome preparation.**

Liposomes were prepared as follows. Lipids from stock solutions of 1-Palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), cholesterol (Chol), 2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000] (DSPE-PEG) and 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-
30 [methoxy(polyethylene glycol)-2000]-maleimide (DSPE-PEG-Mal) in

chloroform : methanol (9 : 1, by volume), were mixed in a molar ratio of 55 : 40 : 4 : 1, dried under reduced nitrogen pressure, dissolved in cyclohexane and lyophilized. Where indicated, 1-methyl-4-(cis-9-dioleyl)methyl-pyridinium-chloride (SAINT-18) was added to the lipid mixture in the indicated molar%,
5 always at the cost of the amount of POPC. When appropriate, trace amounts of [3H]cholesteryloleyl-ether and cholesteryl-[¹⁴C]-oleate were added to the preparation as a non-degradable marker and degradable marker, respectively. For fluorescence microscopy and confocal laser scanning microscopy, 0.5 mol% DiI, was added to the lipid mixture as indicated. The lipids were then hydrated
10 in a Hepes buffer (10 mM N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (Hepes), 135 mM NaCl, pH 6.7) or in an aqueous solution containing 100 mM Calcein. The liposomes formed were sized by repeated extrusion (13 times) through polycarbonate filters (Costar, Cambridge MA, USA), pore size 50 nm, using a high pressure extruder (Lipex, Vancouver, Canada).

15 The monoclonal mouse anti-human E-selectin antibody (H18/7, kindly provided by dr. M. Gimbrone, jr., Boston, MA, USA) was thiolated by means of N-succinimidyl-S-acetylthioacetate and coupled to a maleimide group at the distal end of the polyethylene glycol chain by a sulfhydryl-maleimide coupling technique [2], exactly as described before for albumin [3]. Control
20 immunoliposomes were prepared as described above using irrelevant rat IgG (Sigma-Aldrich Chemie, Zwijndrecht, The Netherlands). Control liposomes without antibody were prepared from the same lipid mixture but, instead of being incubated with antibody, they were incubated with cystein in a molar amount twice that of DSPE-PEG-Mal to block reactive maleimido groups. The
25 immuno-liposomes were characterized by determining protein using mouse immunoglobulin G as a standard 4 and phospholipid phosphorus content 5. Total liposomal lipid concentrations were adjusted for the amount of cholesterol and SAINT present in the liposome preparations.

Particle size was analyzed by dynamic light scattering using a Nicomp model 380 ZLS submicron particle analyzer in the volume weighing mode (NICOMP particle sizing systems, Santa Barbara, CA, USA).

5 **Cell targeting**

HUVEC were activated with TNF α for the indicated time period. Liposomes, either radiolabeled or fluorescently labelled, were added to the cells for the indicated time. For analysis of cell association of radiolabeled liposomes, cells were placed on ice at the end of the incubation, the medium was removed and
10 cells were washed for 5 times with PBS. Cells were then lysed and radioactivity was determined by liquid scintillation counting. For fluorescence microscopy and confocal laser scanning microscopy incubations were performed as described above. Microscopy was performed within 1 hour after finishing the incubation. During this time period there was no change in cell
15 morphology and images of similar incubations were reproducible during this period.

RESULTS

Immunoliposomes targeted to E-selectin or VCAM-1 on TNF α activated
20 endothelial cells are readily endocytosed in amounts that are comparable to the endocytosis capacity of macrophages. However, unlike macrophages, endothelial cells do not process (degrade) the liposomes (Figure 1) and accumulate them inside vesicular bodies (Figure 2). The absence of liposome degradation and/or destabilization results in retention of the incorporated
25 drugs inside the liposome and consequently in inferior pharmacological efficacy.

Liposomes formulated with increasing mol% SAINT are size stable for a period of over 2 months when stored under argon at 4°C. Formulation of liposomes
30 with SAINT results in a slight increase in particle diameter from 90 nm for

liposomes without SAINT to 130 nm for liposomes formulated with 20 mol% SAINT (Figure 3). The results were compared to liposomes containing 20 mol% DOTAP, a non-pyridinium cationic lipid often used in lipid based nucleotide delivery.

5

Liposomes formulated with SAINT display also size stability at 37°C for at least 24 h (Figure 4A). When incubated in the presence of 10% (v/v) serum, an increase in liposome size after 7 h of incubation, especially at room temperature (Figure 4B). However, at the physiologically relevant temperature of 37°C the increase in diameter after 24 h of incubation was very limited for liposomes formulated with a synthetic pyridinium-derived amphiphile. For liposomes formulated with 20 mol% SAINT the mean diameter increased to 130 nm as compared to time 0.

15 As can be derived from figure 5, liposomes formulated with up to 20 mol% of pyridinium-derived amphiphile remained size stable. Increasing amounts lead to reduced stability, while at 30 mol% resulted in giant, fused particles having a diameter of around 500 nm.

20 The release of liposomal content was determined using calceine as a model for a water soluble encapsulated compound. Fluorescence of calceine is quenched at the concentration at which it is encapsulated inside the liposomes (100 mM). Upon dilution calceine emits fluorescence. This allows to distinguish encapsulated calceine (no fluorescence) from released calcein (fluorescence).

25 The endocytotic pathway by which (immuno)liposomes are taken up involves intracellular transport through endosomes, where a drop in pH occurs. Therefore, the release of calceine from liposomes formulated with increasing amounts of synthetic pyridinium-derived amphiphile was determined as a function of pH (see Figure 6). Retention of calceine in liposomes formulated
30 with SAINT is stable at neutral pH. Below pH 6 calceine release increases,

where liposomes formulated with 20 mol% show most release in a pH range from 4.5-6. Liposomes formulated with 20 mol% DOTAP display a less favourable release pattern (Figure 6). In Figure 7 the release patterns for liposomes formulated with 20 mol% SAINT are compared at pH 7.4 and pH 5 (relevant pH with regard to pH within endosomes). At pH 5 there is a significant increase in calcein release from the liposomes. Additionally, release from SAINT liposomes at this pH is significantly better than from liposomes formulated with DOTAP.

10 The addition of 10% serum (v/v) to liposomes formulated with SAINT led to an increased calcein release in time for liposomes containing > 10mol% SAINT. (Figure 8).

Next, *in vitro* calcein release from liposomes was investigated. Endothelial cells, HUVEC, were activated with TNF α and incubated with (a) anti-E-selectin immunoliposomes formulated with 20 mol% SAINT, (b) control anti-E-selectin immunoliposomes without SAINT or (c) liposomes (without anti-E-selectin) formulated with 20 mol% SAINT. Figure 9 clearly shows that cells incubated with immunoliposomes formulated with SAINT showed the most calcein fluorescence, indicating preferential intracellular release of the fluorophore from the SAINT liposomes that were taken up by the cells. Uptake of anti-E-selectin immunoliposomes is not influenced by incorporation of SAINT (determined by FACS using DiI as a label, see figure 10), thus the amount of liposomes taken up in Figure 9A and Figure 9B was similar. Liposomes formulated with SAINT but without cell targeting means are taken up to a low extent (Figure 9C).

The result from figure 9 was confirmed in short term incubations with liposomes that were double labeled with DiI as a lipid label and calcein as encapsulated label. HUVEC that were activated for 4 h with TNF α were

incubated for 5, 15, 30 or 60 minutes with anti-E-selectin immunoliposomes without SAINT or formulated with 20 mol% SAINT. In cells incubated with liposomes without SAINT primarily red fluorescence is observed at all time points. In contrast, in cells incubated with liposomes formulated with SAINT,
5 apart from red fluorescence also yellow fluorescence was observed at the early time points. The yellow fluorescence indicates release of calceine which fluorescence is co-localized with the liposomes. At 30 and 60 minutes there was also green fluorescence visible, indicating a different intracellular routing of the liposome content as compared to the liposome itself (data not shown).

10

Uptake of liposomes and release of encapsulated calceine from liposomes was (semi)quantified using FACS. Uptake of E-selectin targeted liposomes by TNF α activated HUVEC was comparable for liposomes that were formulated with 20 mol% SAINT and for liposomes without SAINT, as determined by

15

measurement of the mean fluorescence intensity (arbitrary units) of liposomal DiI (figure 10A). Calceine release from E-selectin targeted liposomes formulated with 20 mol% SAINT was, however, 8-10 times higher as compared to liposomes without SAINT (figure 10B).

20

E-selectin targeted liposomes formulated with 20 mol% SAINT, containing specific siRNA were very effective in downregulation of target genes (figures 11 and 12). In TNF α activated glomerular endothelial cells E-selectin targeted liposomes formulated with 20 mol% SAINT and containing siRNA against interleukin-8 (IL-8), inhibited IL-8 expression for almost 60%, while there was no effect of liposomes formulated without SAINT or liposomes containing a
25 control (scrambled) siRNA (figure 11). The same holds true for the expression of VE-cadherin, which could be efficiently inhibited for more than 60% using E-selectin targeted liposomes formulated with 20 mol% SAINT, in a mouse derived endothelial cell line.

30 **References**

- (1) Bevilacqua MP, Pober JS, Mendrick DL, Cotran RS, Gimbrone MA, Jr. Identification of an inducible endothelial-leukocyte adhesion molecule. *Proc Natl Acad Sci U S A*. 1987;84:9238-9242.
- 5 (2) Derksen J, Scherphof G. An improved method for the covalent coupling of proteins to liposomes. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 1985;814:151-155.
- (3) Kamps JA, Swart PJ, Morselt HW et al. Preparation and characterization of conjugates of (modified) human serum albumin and
10 liposomes: drug carriers with an intrinsic anti-HIV activity. *Biochim Biophys Acta*. 1996;1278:183-190.
- (4) Peterson GL. A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal Biochem*. 1977;83:346-356.
- 15 (5) Böttcher CJF, Van Gent CM, Pries C. A rapid and sensitive sub-micro phosphorus determination. *Anal Chim Acta*. 1961;24:203-204.

Claims

1. A liposome comprising at least one lipid bilayer enclosing an interior aqueous compartment, wherein said lipid bilayer comprises at least one synthetic pyridinium-derived amphiphile in an amount of up to 25 mol% based on the total lipid content of the liposome.
2. Liposome according to claim 1, wherein the total amount of pyridinium-derived amphiphile is 2 to 25 mol%, preferably 5 to 20 mol%.
3. Liposome according to claim 2, wherein the total amount of pyridinium-derived amphiphile is 10-20 mole%.
4. Liposome according to claim 1 or 2, having an external diameter of from about 30 to about 300 nanometers (nm), most preferably 50 to 150 nm.
5. Liposome according to any one of the preceding claims, wherein said at least one synthetic pyridinium-derived amphiphile is a Saint-molecule, preferably a Saint-molecule selected from those claimed in European Patent EP0755924.
6. Liposome according to claim 4, comprising Saint-18 (1-methyl-4-(cis-9-dioleyl)methyl-pyridinium-chloride) and/or Saint-12 (1-methyl-4-(pentacosan-13-yl)-pyridinium-chloride).
7. Liposome according to any one of claims 1 to 6, being sterically stabilized, preferably by insertion of a monosialoganglioside (GM1) and/or or poly(ethylene glycol) (PEG) derivatized lipid within the lipid bilayer.

8. Liposome according to any one of the preceding claims, comprising at its surface at least one targeting means capable of recognizing and binding to a surface molecule of a particular target cell or an intracellular compartment of a target cell.
9. Liposome according to claim 8, wherein said targeting means has specificity for the surface of a target cell that is not specialized in scavenging and processing particles, preferably an endothelial cell.
10. Liposome according to any one of claims 7 to 9, wherein said targeting means is the RGD-peptide motif, anti-E-selectin antibody or anti-VCAM-1 antibody.
11. Liposome according to any one of claims 7 to 10, wherein said at least one targeting means is bound via a linker molecule to a Saint-molecule inserted in the lipid bilayer.
12. Liposome according to any one of the preceding claims, comprising in its interior compartment and/or associated with its lipid bilayer at least one biologically active compound.
13. Liposome according to claim 12, wherein said biologically active compound is selected from the group consisting of antibiotic, antibacterial, antiviral, antimycotic, anti-inflammatory, antiproliferative and antineoplastic drugs; hormones; nucleic acids, including genes, recombinant nucleic acids, oligonucleotides, siRNA, a viral gene or a gene from a microorganism; and antigens.
14. A method for preparing a liposome according to any one of claims 1-13, comprising mixing conventional lipids with a suitable amount of at least one

synthetic pyridinium-derived amphiphile, and preparing a liposome according to conventional procedures.

15. A pharmaceutical composition comprising a liposome according to claim 12 or 13 and a pharmaceutically acceptable carrier.
16. Use of a liposome according to any one of claims 1-13 as delivery vehicle to *in vitro* or *in vivo* introduce a substance of interest in a target cell that is not specialized in scavenging and/or processing lipid-based particles, in particular wherein said target cell is selected from the group consisting of endothelial cells, epithelial cells, muscle cells, brain cells, nerve cells, skin cells, hair cells, subsets of progenitor cells, and pericytes.
17. Use of a liposome according to any one of claims 1-13 as a drug delivery vehicle.
18. Use of a liposome according to claims 12 or 13 in the treatment of cancer, chronic inflammation, atherosclerosis, tissue regeneration/repair, diabetes or metabolic disease associated cell dysfunction.

Figure 1A




Figure 1B




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 5

liposome diameter as function of SAINT
content




Figure 6




Figure 7




Figure 8







Figure 9



A



B



C

Figure 10A




Figure 10B




Figure 11




Figure 12

