




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 229/28, A61K 31/195, C07C 233/47, A61K 31/16</b>		<b>A1</b>	(11) International Publication Number: <b>WO 97/33858</b>
			(43) International Publication Date: 18 September 1997 (18.09.97)
(21) International Application Number: <b>PCT/US97/02295</b>		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 18 February 1997 (18.02.97)		<b>Published</b> <i>With international search report.</i>	
(30) Priority Data: 60/013,432 14 March 1996 (14.03.96) US			
(71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HORWELL, David, Christopher [GB/GB]; 8 West Hill Road, Foxton, Cambridge CB2 6SZ (GB). BRYANS, Justin, S. [GB/GB]; Dean Cottage, 3 W. Wickham Road, Balsham CB1 6DZ (GB). KNEEN, Clare, O. [GB/GB]; Slade Cottage, Petts Lane, Little Walden, Essex CB10 1XH (GB). MORRELL, Andrew, I. [GB/GB]; 4 Monet Close, St. Ives, Huntingdon, Cambridgeshire PE17 6EH (GB). RATCLIFFE, Giles, S. [GB/GB]; 60 High Street, Melbourn, Near Royston, Hertfordshire SG8 6AB (GB). HARTENSTEIN, Johannes [DE/DE]; Fohrenbuhl 23, D-79252 Stegen-Wittental (DE).			
(74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.			

(54) Title: NOVEL SUBSTITUTED CYCLIC AMINO ACIDS AS PHARMACEUTICAL AGENTS

## (57) Abstract

Novel substituted cyclic amino acids of formula (I) are disclosed and are useful as agents in the treatment of epilepsy, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, depression, anxiety, panic, pain, and neuropathological disorders. Processes for the preparation and intermediates useful in the preparation are also disclosed.



(I)

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

NOVEL SUBSTITUTED CYCLIC AMINO ACIDS AS  
PHARMACEUTICAL AGENTS

## 5 BACKGROUND OF THE INVENTION

Compounds of formula




wherein  $R_1$  is hydrogen or a lower alkyl radical and  $n$  is 4, 5, or 6 are known in United States Patent Number 4,024,175 and its divisional United States  
15 Patent Number 4,087,544. The uses disclosed are: protective effect against cramp induced by thiosemicarbazide; protective action against cardiazole cramp; the cerebral diseases, epilepsy, faintness attacks, hypokinesia, and cranial traumas; and  
20 improvement in cerebral functions. The compounds are useful in geriatric patients. The patents are hereby incorporated by reference.

## 25 SUMMARY OF THE INVENTION

The novel substituted cyclic amino acids, their derivatives, prodrugs, and pharmaceutically acceptable salts are useful in a variety of disorders. The  
30 disorders include: epilepsy, faintness attacks, hypokinesia, cranial disorders, neurodegenerative disorders, depression, anxiety, panic, pain, and neuropathological disorders.

- 2 -

The compounds are those of formula



5 a pharmaceutically acceptable salt thereof or a prodrug thereof wherein

$R_1$  to  $R_{10}$  are each independently selected from straight or branched alkyl of from 1 to 6 carbon atoms, unsubstituted or substituted benzyl or phenyl which substituents are selected from halogen, alkoxy, alkyl, hydroxy, carboxy, carboalkoxy, trifluoromethyl, and nitro, and

15 any  $R_1$  to  $R_{10}$ , which is not one of the above, is hydrogen.

20 Especially preferred compounds of the invention are:

(1-aminomethyl-4-tert-butyl-cyclohexyl)-acetic acid;

(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid;

25 (1-aminomethyl-3-methyl-cyclohexyl)-acetic acid [1R-(1 $\alpha$ ,3 $\beta$ )];

(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid [1S-(1 $\alpha$ ,3 $\beta$ )];

cis (1-aminomethyl-4-methyl-cyclohexyl)-acetic acid;

30 cis (1-aminomethyl-4-isopropyl-cyclohexyl)-acetic acid;

(1-aminomethyl-2-methyl-cyclohexyl)-acetic acid;

( $\pm$ )-(1-aminomethyl-3,3-dimethyl-cyclohexyl)-acetic acid;

35 acid;

- 3 -

(1-aminomethyl-3,3,5,5-tetramethyl-cyclohexyl)-  
acetic acid;  
(1-aminomethyl-4-methyl-cyclohexyl)-acetic acid;  
(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid  
5 methyl ester monohydrochloride;  
[1-(acetylamino-methyl)-3-methyl-cyclohexyl]-  
acetic acid; and  
[2-(1-Aminomethyl-3-methyl-cyclohexyl)-  
acetylamino]-acetic acid monohydrochloride.

10 Novel intermediates useful in the preparation of  
the final products are disclosed as well as a novel  
process for the preparation of the compounds.

15 DETAILED DESCRIPTION

The compounds of the instant invention and their  
pharmaceutically acceptable salts are as defined by  
Formula I.

20 The term "alkyl" is a straight or branched group  
of from 1 to 6 carbon atoms including but not limited  
to methyl, ethyl, propyl, n-propyl, isopropyl, butyl,  
2-butyl, tert-butyl, pentyl, hexyl, and n-hexyl.

Preferred groups are methyl and tert-butyl.

25 The benzyl and phenyl groups may be unsubstituted  
or substituted by from 1 to 3 substituents selected  
from halogen, alkyl, alkoxy, hydroxy, carboxy,  
carboalkoxy, trifluoromethyl, and nitro.

Halogen includes fluorine, bromine, chlorine, and  
30 iodine.

Since amino acids are amphoteric, pharmaco-  
logically compatible salts when R is hydrogen can be  
salts of appropriate inorganic or organic acids, for  
example, hydrochloric, sulphuric, phosphoric, acetic,  
35 oxalic, lactic, citric, malic, salicylic, malonic,  
maleic, succinic, and ascorbic. Starting from

-4-

corresponding hydroxides or carbonates, salts with alkali metals or alkaline earth metals, for example, sodium, potassium, magnesium, or calcium are formed. Salts with quaternary ammonium ions can also be prepared with, for example, the tetramethyl-ammonium ion. The carboxyl group of the amino acids can be esterified by known means.

Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

Certain of the compounds of the present invention possess one or more chiral centers and each center may exist in the R(D) or S(L) configuration. The present invention includes all enantiomeric and epimeric forms as well as the appropriate mixtures thereof. For example, the compound of Example 1 is a mixture of all four possible stereoisomers. The compound of Example 6 is one of the isomers. The configuration of the cyclohexane ring carbon centers may be R or S in these compounds where a configuration can be defined.


The compounds of the invention may be synthesized, for example, by utilizing the general strategy (Scheme 1 below) outlined by Griffiths G., et al., Helv. Chim. Acta, 74:309 (1991). Alternatively, they may also be made as shown (in Scheme 2 below), analogously to the published procedure for the synthesis of 3-oxo-2,8-diazaspiro[4,5]decane-8-carboxylic acid tert-butyl ester (1) (Smith P.W., et al., J. Med. Chem., 38:3772 (1995)). The compounds may also be synthesized by the methods outlined by Satzinger G., et al., (US 4,024,175 and US 4,152,326) (Schemes 3 and 4 below). The compounds may also be synthesized by the route outlined by Griffiths G.,

-5-

et al., Helv. Chim. Acta, 74:309 (1991) as in Scheme 5  
below.

- 6 -

Scheme 1




- (i) Ethyl cyanoacetate, piperidine (Cope, et al., J. Am. Chem. Soc., 63:3452 (1941))
- (ii) NaCN, EtOH/H<sub>2</sub>O,
- (iii) EtOH, HCl,
- (iv) H<sub>2</sub>O/H<sup>+</sup>,
- (v) H<sub>2</sub>, Rh/C, MeOH,
- (vi) HCl




- 7 -

Scheme 2




- (i)  $\text{Ph}_3\text{P} = \text{CHCO}_2\text{Me}$ ,  
(ii)  $\text{MeNO}_2$ , 1,1,3,3-tetramethylguanidine,  
(iii) Raney nickel,  $\text{EtOH}/\text{H}_2\text{O}$ ,  
(iv)  $\text{HCl}$



- 8 -


Scheme 3



- (i) Ethylcyanoacetate, ammonia then  $H_3O^+$ ;  
(ii)  $H_2SO_4$ ;  
(iii)  $Ac_2O$ ;  
(iv)  $MeOH$ ;  
(v) Curtius Reaction;  
(vi)  $HCl$ ,  $H_2O$ , then anion exchange

- 9 -


Scheme 4



- (i) Ethylcyanoacetate, ammonia then  $\text{H}_3\text{O}^+$ ;  
(ii)  $\text{H}_2\text{SO}_4$ ;  
(iii)  $\text{Ac}_2\text{O}$ ;  
(iv)  $\text{H}_2\text{NOH}$ ;  
(v)  $\text{PhSO}_2\text{Cl}$ ;  
(vi)  $\text{Et}_3\text{N}$ ,  $\text{MeOH}$ ;  
(vii)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ , then anion exchange

- 10 -


Scheme 5



- (i) Ethyl cyanoacetate, piperidine (Cope, et al., J. Am. Chem. Soc., 63:3452 (1941));
- (ii) NaCN, EtOH/H<sub>2</sub>O;
- (iii) BnOH, HCl;
- (iv) H<sub>2</sub>O/H<sup>+</sup>;
- (v) H<sub>2</sub>, Rh/C, MeOH

-11-


Examples of prodrugs are:



These can be synthesized, for example, via the routes outlined in Schemes 6 through 8 below.

- 12 -


Scheme 6



(i) MeOH, HCl reflux

-13-


Scheme 7



(i)  $\text{MeCOCl}$ ,  $\text{NaOH}$ ,  $\text{H}_2\text{O}$

-14-

Scheme 8



- (i)  $\text{BnOCOCl}$ ,  $\text{H}_2\text{O}$ , 1,4-dioxan,  $\text{NaOH}$ ;  
 (ii) a) Dicyclohexylcarbodiimide,  
 pentafluorophenol, ethyl acetate,  
 b) Glycine benzyl ester, triethylamine;  
 (iii)  $\text{Pd}(\text{OH})_2/\text{C}$ ,  $\text{HCl}$ ,  $\text{EtOH}$ ,  $\text{H}_2$



-15-






The radioligand binding assay using [ $^3\text{H}$ ]gabapentin and the  $\alpha_2\delta$  subunit derived from porcine brain tissue was used ("The Novel Anti-convulsant Drug, Gabapentin, Binds to the  $\alpha_2\delta$  Subunit of a Calcium Channel", Gee N., et al., J. Biological Chemistry, in press).

TABLE 1

Compound	Structure	$\text{IC}_{50}$ ( $\mu\text{M}$ )
10 (1-aminomethyl-4-tert-butyl-cyclohexyl)-acetic acid		200
15 (1-aminomethyl-3-methyl-cyclohexyl)-acetic acid		0.13
20 (1-aminomethyl-3-methyl-cyclohexyl)-acetic acid [1R-(1 $\alpha$ ,3 $\beta$ )]		13
25 (1-aminomethyl-3-methyl-cyclohexyl)-acetic acid [1S-(1 $\alpha$ ,3 $\beta$ )]		.030
30 cis (1-aminomethyl-4-methyl-cyclohexyl)-acetic acid		10

-16-


TABLE 1. (con't)

Compound	Structure	IC <sub>50</sub> (μM)
5 cis (1-aminomethyl-4-isopropyl-cyclohexyl)-acetic acid		10
10 (1-aminomethyl-2-methyl-cyclohexyl)-acetic acid		7
15 (±)-(1-aminomethyl-3,3-dimethyl-cyclohexyl)-acetic acid		0.5
20 (1-aminomethyl-3,3,5,5-tetramethyl-cyclohexyl)-acetic acid		10
25 (1-aminomethyl-4-methyl-cyclohexyl)-acetic acid		0.33

-17-

Table 1 above shows the binding affinity of the compounds of the invention to the  $\alpha_2\delta$  subunit. Gabapentin (Neurontin®) is about 0.10 to 0.12  $\mu\text{M}$  in this assay. The compounds of the instant invention are expected, therefore, to exhibit pharmacologic properties comparable to gabapentin. For example, as agents for convulsions, anxiety, and pain.

The compounds of the invention are related to Neurontin®, a marketed drug effective in the treatment of epilepsy. Neurontin® is 1-(aminomethyl)-cyclohexanecarboxylic acid of structural formula



The compounds of the invention are also expected to be useful in the treatment of epilepsy. See Table 1 above for  $\text{IC}_{50}$  data as compared to Neurontin®.

The present invention also relates to therapeutic use of the compounds of the mimetic as agents for neurodegenerative disorders.

Such neurodegenerative disorders are, for example, Alzheimer's disease, Huntington's disease, Parkinson's disease, and Amyotrophic Lateral Sclerosis.

The present invention also covers treating neurodegenerative disorders termed acute brain injury. These include but are not limited to: stroke, head trauma, and asphyxia.

Stroke refers to a cerebral vascular disease and may also be referred to as a cerebral vascular incident (CVA) and includes acute thromboembolic stroke. Stroke includes both focal and global ischemia. Also, included are transient cerebral ischemic attacks and

-18-

other cerebral vascular problems accompanied by cerebral ischemia. A patient undergoing carotid endarterectomy specifically or other cerebrovascular or vascular surgical procedures in general, or diagnostic  
5 vascular procedures including cerebral angiography and the like.

Other incidents are head trauma, spinal cord trauma, or injury from general anoxia, hypoxia, hypoglycemia, hypotension as well as similar injuries  
10 seen during procedures from embolus, hyperfusion, and hypoxia.

The instant invention would be useful in a range of incidents, for example, during cardiac bypass surgery, in incidents of intracranial hemorrhage, in  
15 perinatal asphyxia, in cardiac arrest, and status epilepticus.

A skilled physician will be able to determine the appropriate situation in which subjects are susceptible to or at risk of, for example, stroke as well as  
20 suffering from stroke for administration by methods of the present invention.

The compounds of the invention are also expected to be useful in the treatment of depression. Depression can be the result of organic disease,  
25 secondary to stress associated with personal loss, or idiopathic in origin. There is a strong tendency for familial occurrence of some forms of depression suggesting a mechanistic cause for at least some forms of depression. The diagnosis of depression is made  
30 primarily by quantification of alterations in patients' mood. These evaluations of mood are generally performed by a physician or quantified by a neuropsychologist using validated rating scales, such as the Hamilton Depression Rating Scale or the Brief Psychiatric Rating Scale. Numerous other scales have  
35 been developed to quantify and measure the degree of

-19-

mood alterations in patients with depression, such as insomnia, difficulty with concentration, lack of energy, feelings of worthlessness, and guilt. The standards for diagnosis of depression as well as all psychiatric diagnoses are collected in the Diagnostic and Statistical Manual of Mental Disorders (Fourth Edition) referred to as the DSM-IV-R manual published by the American Psychiatric Association, 1994.

GABA is an inhibitory neurotransmitter with the central nervous system. Within the general context of inhibition, it seems likely that GABA-mimetics might decrease or inhibit cerebral function and might therefore slow function and decrease mood leading to depression.

The compounds of the instant invention may produce an anticonvulsant effect through the increase of newly created GABA at the synaptic junction. If gabapentin does indeed increase GABA levels or the effectiveness of GABA at the synaptic junction, then it could be classified as a GABA-mimetic and might decrease or inhibit cerebral function and might, therefore, slow function and decrease mood leading to depression.

The fact that a GABA agonist or GABA-mimetic might work just the opposite way by increasing mood and thus, be an antidepressant, is a new concept, different from the prevailing opinion of GABA activity heretofore.

The compounds of the instant invention are also expected to be useful in the treatment of anxiety and of panic as demonstrated by means of standard pharmacological procedures.

-20-

## MATERIAL AND METHODS

Carrageenin-Induced Hyperalgesia

Nociceptive pressure thresholds were measured in the rat paw pressure test using an analgesymeter (Randall-Selitto method: Randall L.O. and Sellitto J.J., A method for measurement of analgesic activity on inflamed tissue. Arch. Int. Pharmacodyn., 4:409-419 (1957)). Male Sprague Dawley rats (70-90 g) were trained on this apparatus before the test day. Pressure was gradually applied to the hind paw of each rat and nociceptive thresholds were determined as the pressure (g) required to elicit paw withdrawal. A cutoff point of 250 g was used to prevent any tissue damage to the paw. On the test day, two to three baseline measurements were taken before animals were administered 100 µL of 2% carrageenin by intraplantar injection into the right hind paw. Nociceptive thresholds were taken again 3 hours after carrageenin to establish that animals were exhibiting hyperalgesia. Animals were dosed with either gabapentin (3-300 mg, s.c.), morphine (3 mg/kg, s.c.) or saline at 3.5 hours after carrageenin and nociceptive thresholds were examined at 4, 4.5, and 5 hours postcarrageenin.

Semicarbazide-Induced Tonic Seizures

Tonic seizures in mice are induced by subcutaneous administration of semicarbazide (750 mg/kg). The latency to the tonic extension of forepaws is noted. Any mice not convulsing within 2 hours after semicarbazide are considered protected and given a maximum latency score of 120 minutes.

Animals

Male Hooded Lister rats (200-250 g) are obtained from Interfauna (Huntingdon, UK) and male TO mice

-21-

(20-25 g) are obtained from Bantin and Kingman (Hull, UK). Both rodent species are housed in groups of six. Ten Common Marmosets (*Callithrix Jacchus*) weighing between 280 and 360 g, bred at Manchester University Medical School (Manchester, UK) are housed in pairs. All animals are housed under a 12-hour light/dark cycle (lights on at 07.00 hour) and with food and water ad libitum.

#### 10 Drug Administration

Drugs are administered either intraperitoneally (IP) or subcutaneously (SC) 40 minutes before the test in a volume of 1 mL/kg for rats and marmosets and 10 mL/kg for mice.

15

#### Mouse Light/Dark Box

The apparatus is an open-topped box, 45 cm long, 27 cm wide, and 27 cm high, divided into a small (2/5) and a large (3/5) area by a partition that extended 20 cm above the walls (Costall B., et al., Exploration of mice in a black and white box: validation as a model of anxiety. Pharmacol. Biochem. Behav., 32:777-785 (1989)).

There is a 7.5 × 7.5 cm opening in the center of the partition at floor level. The small compartment is painted black and the large compartment white. The white compartment is illuminated by a 60-W tungsten bulb. The laboratory is illuminated by red light. Each mouse is tested by placing it in the center of the white area and allowing it to explore the novel environment for 5 minutes. The time spent in the illuminated side is measured (Kilfoil T., et al., Effects of anxiolytic and anxiogenic drugs on exploratory activity in a simple model of anxiety in mice. Neuropharmacol., 28:901-905 (1989)).

-22-

Rat Elevated X-Maze

A standard elevated X-maze (Handley S.L., et al., Effects of alpha-adrenoceptor agonists and antagonists in a maze-exploration model of 'fear'-motivated behavior. Naunyn-Schiedeberg's Arch. Pharmacol., 327:1-5 (1984)), was automated as previously described (Field, et al., Automation of the rat elevated X-maze test of anxiety. Br. J. Pharmacol., 102(Suppl):304P (1991)). The animals are placed on the center of the X-maze facing one of the open arms. For determining anxiolytic effects the entries and time spent on the end half sections of the open arms is measured during the 5-minute test period (Costall, et al., Use of the elevated plus maze to assess anxiolytic potential in the rat. Br. J. Pharmacol., 96(Suppl):312p (1989)).

Marmoset Human Threat Test

The total number of body postures exhibited by the animal towards the threat stimulus (a human standing approximately 0.5 m away from the marmoset cage and staring into the eyes of the marmoset) is recorded during the 2-minute test period. The body postures scored are slit stares, tail postures, scent marking of the cage/perches, piloerection, retreats, and arching of the back. Each animal is exposed to the threat stimulus twice on the test day before and after drug treatment. The difference between the two scores is analyzed using one-way analysis of variance followed by Dunnett's t-test. All drug treatments are carried out SC at least 2 hours after the first (control) threat. The pretreatment time for each compound is 40 minutes.

Rat Conflict Test

Rats are trained to press levers for food reward in operant chambers. The schedule consists of alternations of four 4-minute unpunished periods on



-23-

variable interval of 30 seconds signalled by chamber lights on and three 3-minute punished periods on fixed ratio 5 (by footshock concomitant to food delivery) signalled by chamber lights off. The degree of footshock is adjusted for each rat to obtain approximately 80% to 90% suppression of responding in comparison with unpunished responding. Rats receive saline vehicle on training days.

The compounds of the instant invention are also expected to be useful in the treatment of pain and phobic disorders (Am. J. Pain Manag., 5:7-9 (1995)).

The compounds of the instant invention are also expected to be useful in treating the symptoms of manic, acute or chronic, single episode, or recurring. They are also expected to be useful in treating and/or preventing bipolar disorder (United States Patent Application Number 08/440,570 filed May 15, 1995).

The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of Formula I or a corresponding pharmaceutically acceptable salt of a compound of Formula I.

For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid.

-24-

Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

-25-

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsules, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

-26-

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 1 g according to the particular application and the potency of the active component. In medical use the drug may be administered three times daily as, for example, capsules of 100 or 300 mg. The composition can, if desired, also contain other compatible therapeutic agents.


In therapeutic use, the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 0.01 mg to about 100 mg/kg daily. A daily dose range of about 0.01 mg to about 100 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

The following examples are illustrative of the instant invention; they are not intended to limit the scope.

30

- 27 -

## EXAMPLE 1

(i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ ,  $\text{NH}_4\text{Ac}$ ,  $\text{AcOH}$ , Toluene,  $120^\circ\text{C}$ (ii) a.  $\text{NaCN}$ ,  $\text{EtOH}$  (95%),  $\text{H}_2\text{O}$ ,  $115^\circ\text{C}$ ; b.  $\text{HCl}$  (g)25 (iii)  $\text{EtOH}$ ,  $\text{HCl}$  (g), Toluene(iv)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ (v)  $\text{H}_2$ ,  $\text{EtOH}/\text{NH}_3$ , Raney Nickel,  $30-50^\circ\text{C}$ (vii)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ,  $140^\circ\text{C}$ 

30

General method, exemplified by synthesis of trans  
(R)-3-ethyl gabapentin

Step (i) Cyanoacetate

35

A mixture of 3-(R)-Methylcyclohexanone (125 mmol),  
 ethyl cyanoacetate (124 mmol), ammonium acetate

-28-

(12.5 mmol) and glacial acetic acid (24 mmol) were refluxed with a Dean Stark trap for 24 hours. The mixture was cooled and washed with H<sub>2</sub>O. The H<sub>2</sub>O washes were extracted with toluene. The toluene extracts were combined with the original organic layer, dried over MgSO<sub>4</sub>, and the solvent evaporated. The crude oil was purified by Kugelrohr distillation to give an oil. Bpt oven temperature 150-160°C. Yield 86%.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 1.01-1.05 (3H, m), 1.17-1.32 (1H, m), 1.35 (3H, t, J = 7 Hz), 1.42-2.30 (6H, m), 2.98 (1H, d, J = 13 Hz), 3.74 (1H, d, J = 13 Hz), 4.27 (2H, q, J = 7 Hz).

MS (CI) m/z: 85, 91, 95, 135, 162, 178, 180, 200, 208 (100% MH<sup>+</sup>), 209.

IR (Film) ν<sub>max</sub>, cm<sup>-1</sup>: 3437, 2956, 2930, 2870, 2223, 1729, 1603, 1448, 1367, 1347, 1313, 1290, 1262, 1246, 1218, 1101, 1084, 1046, 1023, 974, 957, 914, 859, 822, 780.

Microanalysis: C<sub>12</sub>H<sub>17</sub>NO<sub>2</sub>:

Calc: C, 69.54; H, 8.27; N, 6.76.

Found: C, 69.44; H, 8.22; N, 6.76.

#### Step (ii) Bisnitrile

To a solution of NaCN (40 mmol) in 6 mL H<sub>2</sub>O and 160 mL Ethanol (95%) was added the cyanoacetate (40 mmol). After 22 hours at reflux the cooled solution was filtered, the filtrate acidified with gaseous HCl, and filtered again. The solvent was removed, and the crude oil was purified by column chromatography to give a pale yellow crystalline solid. Yield 88% Mpt.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 0.90 (1H, m), 0.98 (3H, d, J = 6 Hz), 1.11 (1H, t, J = 12 Hz), 1.38 (1H, dt, J = 4.9 Hz), 1.60-1.90 (4H, m), 2.07 (2H, m), 2.68 (2H, s).

MS (CI) m/z: 91 (100%), 92, 108, 130, 136, 163, (50% MH<sup>+</sup>), 180.

-29-

IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu_{\max}$  cm<sup>-1</sup>: 2956, 2932, 2862, 2234, 1714, 1457, 1447, 1427, 1386, 1358.

Microanalysis: C<sub>10</sub>H<sub>14</sub>N<sub>2</sub>:

Calc: C, 74.04; H, 8.70; N, 17.27.

5 Found: C, 74.05; H, 8.71; N, 17.25.

Step (iii) Imidate

To a solution of the binitrile (6.2 mmol) in 30 mL ethanol (absolute) was added 30 mL dried toluene. The  
10 solution was chilled in ice while saturating with gaseous HCl. The stoppered solution was then left to stand at room temperature for 24 hours. The solvent was removed, and the solid residue was triturated with diethyl ether to obtain a ppt which was dried to give a  
15 white crystalline solid. Yield 50%. Mpt 118-120°C.

<sup>1</sup>H NMR (DMSO) 400 MHz:  $\delta$  0.8-0.89 (1H, m), 0.91 (3H, d, J = 6.3 Hz), 1.06-1.12 (1H, m), 1.24-1.35 (1H, m), 1.37 (3H, t, J = 7 Hz), 1.41-1.95 (6H, m), 3.02 (2H, s), 4.49 (2H, q, J = 7 Hz).

20 MS (CI) m/z: 91, 133, 154, 164, 179, 181, (100% MH<sup>+</sup> - CN), 195 (MH<sup>+</sup>), 209.

IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu_{\max}$  cm<sup>-1</sup>: 2957, 2938, 2858, 2233, 1651, 1573, 1446, 1388, 1361, 1137, 1103, 1022, 1005, 952, 933, 874, 834.

25 Microanalysis: C<sub>12</sub>H<sub>20</sub>N<sub>2</sub>O·1.08 HCl:

Calc: C, 58.19; H, 8.58; N, 11.31.

Found: C, 58.25; H, 8.59; N, 11.59.

Step (iv) Ester

30 The imidate (1.1 mmol) was dissolved in ice cold H<sub>2</sub>O (40 mL) and the pH adjusted with 1N HCl to pH 1.5. The solution was stirred at room temperature for 20 hours. Ethylacetate was added (30 mL), and the organic layer was washed with H<sub>2</sub>O, dried, and the  
35 solvent removed to leave a clear oil. Yield 82%.

-30-

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.78-0.90 (1H, m), 0.93 (3H, d,  $J = 6$  Hz), 0.97-1.00 (1H, m), 1.23-1.25 (1H, m), 1.29 (3H, t,  $J = 7.2$  Hz), 1.59-1.80 (4H, m), 2.05-2.08 (2H, Br t), 2.54 (2H, s), 4.20 (2H, q,  $J = 7.2$  Hz).

5 MS (CI)  $m/z$ : 88, 95, 109, 122, 137, 160, 164 (100%  $\text{M}^+$ -EtOH), 182, 183, 199, 210 (60%  $\text{MH}^+$ ), 230.

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2930, 2870, 2235, 1737, 1458, 1414, 1375, 1345, 1264, 1196, 1171, 1096, 1041, 1026, 959, 847.

10 Microanalysis:  $\text{C}_{12}\text{H}_{19}\text{NO}_2$ :

Calc: C, 68.87; H, 9.15; N, 6.69.

Found: C, 68.87; H, 9.11; N, 6.90.

#### Step (v) Lactam

15 The ester (8.9 mmol) was dissolved in  $\text{NH}_3/\text{EtOH}$  (7%, 40 mL) along with prewashed Raney Nickel ( $\text{H}_2\text{O}$  followed by EtOH) in a 250 mL Parr flask. The solution was hydrogenated at  $30^\circ\text{C}$ , 46 psi for 24 hours. The cooled solution was filtered through a pad of celite, washing with ethylacetate. The solvent was removed from the filtrate to leave a white solid. Yield 30%. Mpt  $92-98^\circ\text{C}$ .

20  $^1\text{H}$  NMR (DMSO) 400 MHz:  $\delta$  0.75-0.82 (1H, m), 0.84 (3H, d,  $J = 6.4$  Hz), 0.88-0.94 (1H, m), 1.14-1.19 (1H, m), 1.20-1.50 (2H, m), 1.50-1.63 (4H, m), 1.91 (2H, s), 3.03 (2H, s), 7.42 (1H, s).

25 MS (CI)  $m/z$ : 166, 167, 168 (100%  $\text{MH}^+$ ), 182, 196.

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3260, 2907, 1695, 1652, 1446, 1318, 1255, 1210, 1068.

30 Microanalysis:  $\text{C}_{10}\text{H}_{17}\text{NO}$ :

Calc: C, 71.81; H, 10.25; N, 8.37.

Found: C, 71.80; H, 10.29; N, 8.31.

#### Step (vi) 3-Methyl gabapentin

35 The lactam (2.17 mmol) was dissolved in a solution of 10 M HCl (5 mL) and  $\text{H}_2\text{O}$  (5 mL), and the mixture was



-31-

refluxed at approximately 140°C for 5 hours. The cooled solution was diluted with 10 mL H<sub>2</sub>O and 10 mL DCM and the aqueous layer was further washed with 2 × 15 mL DCM. The aqueous layer was then reduced to dryness to leave a white solid. Yield 76%. Mpt 148-155°C.  $[\alpha]_D = -2.5$  (T-20°C, c = 1, MeOH). One isomer (RR).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 06.9-0.79 (1H, m), 0.82 (3H, d, J = 6 Hz), 0.87-0.90 (1H, m), 1.12-1.20 (1H, dt, J = 4.5, 13.3 Hz), 1.34-1.50 (3H, m), 1.60-1.63 (3H, m), 2.30 (2H, s), 3.01 (2H, s), 7.93 (3H, Br s).

MS (CI) m/z: 95, 109, 121, 151, 167, 168 (100% MH<sup>+</sup> - H<sub>2</sub>O), 186 (MH<sup>+</sup>).

IR (MeOH)  $\nu_{\max}$  cm<sup>-1</sup>: 2924, 2353, 1708, 1599, 1523, 1454, 1216.


Microanalysis: C<sub>10</sub>H<sub>19</sub>NO<sub>2</sub>·1.1 HCl:

Calc: C, 53.29; H, 8.99; N, 6.21.

Found: C, 53.23; H, 8.99; N, 6.45.

- 32 -

## EXAMPLE 2



(i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ ,  $\text{NH}_4\text{Ac}$ ,  $\text{AcOH}$ , toluene,  $120^\circ\text{C}$ ;

(ii) a.  $\text{NaCN}$ ,  $\text{EtOH}$  (95%),  $\text{H}_2\text{O}$ ,  $115^\circ\text{C}$ ; b.  $\text{HCl}$  (g);

25 (iii)  $\text{EtOH}$ ,  $\text{HCl}$  (g), toluene;

(iv)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ;

(v)  $\text{H}_2$ ,  $\text{EtOH}/\text{NH}_3$ , Raney Nickel,  $30-50^\circ\text{C}$ ;

(vi)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ,  $140^\circ\text{C}$

30 Cis/trans (RS)-2-methyl gabapentin

Step (i) Cyanoacetate

(±)-2-Methylcyclohexanone (80 mmol), ethyl cyanoacetate (80 mmol), ammonium acetate (8 mmol), and glacial acetic acid (16 mmol) were reacted as in the general method Step (i), to give a clear oil. Yield 76%. Bpt oven temperature  $120-140^\circ\text{C}$ , 3 mbar.

35

-33-

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  1.23 (3H, dd,  $J = 7, 10$  Hz),  
1.35 (3H, t,  $J = 7$  Hz), 1.55-1.82 (5H, m), 1.93-2.05  
(1H, m), 2.17 (1H, dt,  $J = 5, 14$  Hz), 2.47 (1H,  
dt,  $J = 5, 9$  Hz), 2.92-2.97 (1H, Br d,  $J = 15$  Hz),  
5 3.30-3.35 (1H, m), 3.81-3.86 (1H, Br d,  $J = 15$  Hz),  
4.06-4.14 (1H, m), 4.23-4.30 (3H, dq,  $J = 1, 6$  Hz).  
MS (CI)  $m/z$ : 91, 105, 120, 162, 180, 184, 189, 208  
( $\text{MH}^+$ ), 216, 233, 234, 242, 261, 262 (100%), 263.  
IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3438, 2978, 2938, 2864, 2223,  
10 1732, 1596, 1463, 1447, 1391, 1368, 1334, 1311, 1289,  
1247, 1224, 1164, 1144, 1103, 1077, 1058, 1032, 993,  
982, 957, 907, 892, 858, 781.  
Microanalysis:  $\text{C}_{12}\text{H}_{17}\text{NO}_2$ :  
Calc: C, 69.54; H, 8.27; N, 6.76.  
15 Found: C, 69.26; H, 8.26; N, 6.66.

#### Step (ii) Bisnitrile

The cyanoacetate (37 mmol) and NaCN (37 mmol) were  
reacted as in the general method Step (ii). The crude  
20 solid was purified by column chromatography (3:1,  
heptane:ethylacetate) to give a clear oil. Yield 76%.  
 $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  1.06 (3H, d,  $J = 6.8$  Hz),  
1.11 (3H, d,  $J = 6.8$  Hz), 1.20-2.20, (18 H, m), 2.77  
(2H, dd,  $J = 16.8$  Hz), 2.63 (2H, dd,  $J = 16.8$  Hz).  
25 MS (CI)  $m/z$ : 91, 95, 108, 109, 136, 16.3 (100%  $\text{MH}^+$ ).  
IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2939, 2865, 2255, 2237, 1750,  
1720, 1450, 1425, 1387, 1356, 1337, 1316, 1269, 1160,  
1097, 992, 929, 879.  
Microanalysis:  $\text{C}_{10}\text{H}_{14}\text{N}_2 \cdot 0.1 \text{H}_2\text{O}$ :  
30 Calc: C, 73.49; H, 8.69; N, 16.86.  
Found: C, 73.24; H, 8.73; N, 17.08.

#### Step (iii) Imidate

The binitrile (7.3 mmol) was reacted as in the  
35 general method Step (iii) to give a white solid.  
Yield 70%. Mpt 107-114°C.

-34-

$^1\text{H}$  NMR (DMSO) 400 MHz:  $\delta$  1.00-1.06 (3H, 2  $\times$  t,  $J$  = 6.4 Hz), 1.10-1.38 (2H, m), 1.38 (3H, t,  $J$  = 6.8 Hz), 1.40-2.10 (7H, m), 2.86, 2.92, 3.10, 3.28 (2H, 4  $\times$  d,  $J$  = 14, 14.4, 14.8, 14 Hz, respectively), 4.48 (2H, q,  $J$  = 6.8 Hz).

MS (CI)  $m/z$ : 87, 95, 154, 163, 181, 195, 209 (100%  $\text{MH}^+$ ), 210.

IR ( $\text{CH}_2\text{Cl}_2$ )  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2938, 2864, 2664, 2235, 1656, 1575, 1446, 1389, 1367, 1139, 1100, 1007, 948, 881, 837, 809.

Microanalysis:  $\text{C}_{12}\text{H}_{20}\text{N}_2\text{O} \cdot 1.06 \text{ HCl}$ :

Calc: C, 58.37; H, 8.60; N, 11.34.

Found: C, 58.15; H, 8.63; N, 11.60.

#### 15 Step (iv) Ester

The imidate (4.1 mmol) was reacted as in the general method Step (iv) to give a clear oil.

Yield 82%.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  1.03, 1.09 (3H, 2  $\times$  d,  $J$  = 7 Hz), 1.27-1.30 (3H, m), 1.32-2.00 (8H, m), 2.10-2.20 (1H, m), 2.44, 2.82 (3H, 2  $\times$  d,  $J$  = 14.8 Hz), 2.54 (1H, m), 4.16-4.22 (2H, m).

MS (CI)  $m/z$ : 88, 95, 109, 122, 164, 182, 210 ( $\text{MH}^+$  100%).

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2936, 2864, 2234, 1737, 1449, 1418, 1385, 1372, 1345, 1270, 1225, 1186, 1128, 1098, 1029, 1001, 932, 883, 864, 808, 732.

Microanalysis:  $\text{C}_{12}\text{H}_{19}\text{NO}_2$ :

Calc: C, 68.87; H, 9.15; N, 6.69.

Found: C, 68.84; H, 9.13; N, 6.75.

#### 30 Step (v) Lactam

The ester (8.4 mmol) was reacted as in the general method Step (v) for 24 hours at 10°C, 50 psi. The crude oil was purified by column chromatography

-35-

(ethylacetate), to give a white solid. Yield 34%.

Mpt 85-90°C.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.88-0.91 (3H, dd,  $J = 4$ , 6.8 Hz), 1.41-1.78 (9H, m), 2.00-2.30 (2H, m),  
5 3.06-3.23 (2H, m), 7.27 (1H, Br s).

MS (CI)  $m/z$ : 81, 95, 108, 137, 166, 167, 168  
(100  $\text{MH}^+$ ), 169, 182, 196.

IR ( $\text{CH}_2\text{Cl}_2$ )  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3210, 2920, 2846, 1690, 1491,  
1446, 1379, 1298, 1242, 1070.

10 Microanalysis:  $\text{C}_{10}\text{H}_{17}\text{NO}$ :

Calc: C, 71.81; H, 10.24; N, 8.37.

Found: C, 71.83; H, 10.19; N, 8.27.

#### Step (vi) 2-Methyl gabapentin

15 The lactam (2.5 mmol) was reacted as in the  
general method Step (vi) to give a white solid.  
Yield 42%. Mpt 108-110°C.  $[\alpha]_{\text{D}} = 0$  ( $T = 20.5^\circ\text{C}$ ,  $C = 1$ ,  
MeOH). Two diastereomers 3:1.

$^1\text{H}$  NMR ( $\text{DMSO} + \text{D}_2\text{O}$ ) 400 MHz:  $\delta$  0.79, 0.85 (3H, 2  $\times$  d,  
20  $J = 6.8$  Hz), 1.21-1.65 (9H, m), 2.22, 2.43 (1H, 2  $\times$  d,  
 $J = 15$  Hz), 2.46, 2.49 (1H, 2  $\times$  d,  $J = 15$  Hz),  
2.83-2.92 (1H, 2  $\times$  d,  $J = 13.6$  Hz), 3.05, 3.15  
(1H, 2  $\times$  d,  $J = 13.6$  Hz).

MS (CI)  $m/z$ : 95, 109, 137, 166, 168 (100% lactam), 169  
25 ( $\text{MH}^+ - \text{H}_2\text{O}$ ), 186 ( $\text{MH}^+$ ), 196.

IR (MeOH)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3384, 2931, 2861, 1703, 1608,  
1506, 1456, 1406, 1232, 1206, 1068, 999.


Microanalysis:  $\text{C}_{10}\text{H}_{19}\text{NO}_2 \cdot 1.3 \text{HCl}$ .

Calc: C, 51.64; H, 8.79; N, 6.02.

30 Found: C, 51.66; H, 8.91; N, 6.16.

- 36 -

## EXAMPLE 3



- (i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ ,  $\text{NH}_4\text{Ac}$ ,  $\text{AcOH}$ , toluene,  $120^\circ\text{C}$ ;  
 (ii) a.  $\text{NaCN}$ ,  $\text{EtOH}$  (95%),  $\text{H}_2\text{O}$ ,  $115^\circ\text{C}$ ; b.  $\text{HCl}$  (g);  
 (iii)  $\text{EtOH}$ ,  $\text{HCl}$  (g), toluene;  
 (iv)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ;  
 (v)  $\text{H}_2$ ,  $\text{EtOH}/\text{NH}_3$ , Raney Nickel,  $30\text{-}50^\circ\text{C}$ ;  
 (vi)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ,  $140^\circ\text{C}$ ;

Step (i) Cyanoacetate

The 4-Methylcyclohexanone (125 mmol), ethyl cyanoacetate (124 mmol), ammonium acetate (12.4 mmol), and glacial acetic acid (24.4 mmol) were reacted as in the general method Step (i) for 8 hours to give a clear

-37-

oil. Yield 82%. Bpt oven temperature 160-190°C,  
4 mbar.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.95 (3H, d,  $J$  = 6.8 Hz),  
1.20-1.31 (2H, m), 1.35 (3H, t,  $J$  = 7.2 Hz), 1.80-1.90  
5 (1H, m), 1.90-2.10 (2H, m), 2.15 (1H, dt,  $J$  = 4.8,  
13.6 Hz), 2.34 (1H, dt,  $J$  = 4.8, 13.6 Hz), 3.02  
(1H, dd,  $J$  = 2.4, 14 Hz), 3.84 (1H, dd,  $J$  = 2.4,  
14 Hz), 4.27 (2H, q,  $J$  = 7.2 Hz).

MS (CI)  $m/z$ : 114, 134, 151, 162, 179, 180, 207, 208  
10 (100%  $\text{MH}^+$ ), 209, 236.

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2927, 2225, 1728, 1601, 1456,  
1367, 1288, 1242, 1192, 1095, 1028, 959, 857, 779.

Microanalysis:  $\text{C}_{12}\text{H}_{17}\text{NO}_2$ :

Calc: C, 69.54; H, 8.27; N, 6.76.

15 Found: C, 69.39; H, 8.27; N, 6.77.

#### Step (ii) Binitrile

The cyanoacetate (30 mmol) and NaCN (30 mmol) were  
reacted as in the general method Step (ii) to give a  
20 crude oil. The oil was purified by column  
chromatography (3:1, heptane:ethylacetate) to give a  
clear oil. Yield 66%.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.98 (3H, d,  $J$  = 5.6 Hz),  
1.30-1.40 (3H, m), 1.50 (2H, m), 1.73-1.92 (2H, m),  
25 2.10 (2H, d,  $J$  = 12.4 Hz), 2.68 (2H, s).

MS (CI)  $m/z$ : 95, 136, 163 (100%  $\text{MH}^+$ ), 164, 182.

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3628, 3288, 2932, 2859, 2252,  
2238, 1779, 1748, 1721, 1626, 1455, 1423, 1381, 1371,  
1332, 1287, 1263, 1194, 1170, 1143, 1109, 1004, 953,  
30 893, 852.

Microanalysis:  $\text{C}_{10}\text{H}_{14}\text{N}_2 \cdot 0.6 \text{H}_2\text{O}$ :

Calc: C, 72.74; H, 8.74; N, 16.97.

Found: C, 72.98; H, 8.61; N, 16.65.

-38-

Step (iii) Imidate

The binitrile (12.4 mmol) was reacted as in the general method Step (i) to give a slightly impure white solid. No purification was attempted, and solid was used in next step.

Step (iv) Ester

The imidate (4.7 mmol) was reacted as in the general method Step (iv) to give a low melting solid.

Yield 75%, based on binitrile.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.92-1.01 (3H, m), 1.27-1.31 (3H, m), 1.37 (5H, m), 1.70-1.73 (2H, m), 2.10-2.13 (2H, m), 2.54 (2H, s), 4.21 (2H, q,  $J = 7.2$  Hz).

MS (CI)  $m/z$ : 95, 112, 122, 164, 182 (100%  $\text{MH}^+ - \text{C}_2\text{H}_5$ ), 210 ( $\text{MH}^+$ ).

IR ( $\text{CH}_2\text{Cl}_2$ )  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2926, 2856, 2235, 1735, 1733, 1452, 1373, 1345, 1253, 1191, 1033, 953.

Microanalysis:  $\text{C}_{12}\text{H}_{19}\text{N}_2\text{O}_2 \cdot 0.12 \text{ H}_2\text{O}$ :

Calc: C, 68.16; H, 9.17; N, 6.62.

Found: C, 68.14; H, 8.91; N, 6.77.

Step (v) Lactam

The ester (2.9 mmol) was reacted as in the general method Step (v) to give a white fibrous solid.

Yield 95%. Mpt 150-152°C.

$^1\text{H NMR}$  (DMSO) 400 MHz:  $\delta$  0.86 (3H, d,  $J = 6$  Hz), 0.93-1.06 (2H, m), 1.27-1.30 (3H, m), 1.51 (2H, d,  $J = 11.6$  Hz), 1.62 (2H, d,  $J = 13.2$  Hz), 1.92 (2H, s), 3.02 (2H, s), 7.43 (1H, Br s).

MS (CI)  $m/z$ : 81, 95, 110, 166, 167, 168 (100%  $\text{MH}^+$ ), 169, 182, 196.

IR ( $\text{CH}_2\text{Cl}_2$ )  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3189, 3093, 2945, 2921, 2864, 1679, 1486, 1447, 1417, 1260.

Microanalysis:  $\text{C}_{10}\text{H}_{17}\text{NO} \cdot 0.15 \text{ H}_2\text{O}$ :

Calc: C, 70.67; H, 10.17; N, 8.24.

Found: C, 70.69; H, 10.05; N, 7.87.



-39-

Step (vi) 4-Methyl gabapentin

The lactam (2.5 mmol) was reacted as in the general method Step (vi) to give an off-white hygroscopic solid. Yield 92%. Mpt 146-151°C

5  $[\alpha]_D = 0$  (T = 21°C, C = 1, MeOH). One diastereomer (cis).

$^1\text{H}$  NMR (DMSO) 400 MHz:  $\delta$  0.88 (3H, d, J = 6 Hz), 1.02-1.12 (2H, m), 1.25-1.32 (3H, m), 1.43-1.47 (2H, m), 2.33 (2H, s), 2.99 (2H, s), 8.03 (3H, Br s), 12.33 (1H, Br s).

10 MS (CI) m/z: 81, 95, 109, 166, 167, 168 (100%  $\text{MH}^+ - \text{H}_2\text{O}$ ), 169, 182, 186 ( $\text{MH}^+$ ), 196.

IR (MeOH)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 3393, 2925, 2862, 1714, 1613, 1514, 1451, 1387, 1251, 1232, 1192, 1151, 1119, 864.


15 Microanalysis:  $\text{C}_{10}\text{H}_{19}\text{NO}_2 \cdot 1 \text{ HCl} \cdot 1 \text{ H}_2\text{O}$ :

Calc: C, 50.04; H, 9.26; N, 5.84.

Found: C, 50.04; H, 9.18; N, 5.82.

- 40 -

## EXAMPLE 4



- (i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ ,  $\text{NH}_4\text{Ac}$ ,  $\text{AcOH}$ , toluene,  $120^\circ\text{C}$ ;
- (ii) a.  $\text{NaCN}$ ,  $\text{EtOH}$  (95%),  $\text{H}_2\text{O}$ ,  $115^\circ\text{C}$ ; b.  $\text{HCl}$  (g);
- 30 (iii)  $\text{EtOH}$ ,  $\text{HCl}$  (g), toluene;
- (iv)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ;
- (v)  $\text{H}_2$ ,  $\text{EtOH}/\text{NH}_3$ , Raney Nickel,  $30-50^\circ\text{C}$ ;
- (vi)  $\text{HCl}$ ,  $\text{H}_2\text{O}$ ,  $140^\circ\text{C}$ ;

-41-

Cis 4-isopropyl gabapentinStep (i) Cyanoacetate

5 The 4-Isopropyl-cyclohexanone (57 mmol), ethylacetate (57 mmol), ammonium acetate (58 mmol), and glacial acetic acid (11.3 mmol) were reacted as in the general method Step (i). Kugelrohr distillation gave a clear oil. Yield 83%. Bpt oven temperature 170-19°C, 4 mbar.

10  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.89 (6H, d,  $J = 6.8$  Hz), 1.20-1.33 (2H, m), 1.35 (3 H, t,  $J = 7.2$  Hz), 1.37-1.50 (2H, m), 2.00-2.11 (3H, m), 2.30 (1H, dt,  $J = 5$ , 14 Hz), 3.10 (1H, m), 3.92 (1H, m), 4.27 (2H, q,  $J = 7.2$  Hz).

15 MS (CI)  $m/z$ : 163, 179, 190, 207, 208, 235, 236 (100%  $\text{MH}^+$ ), 237, 264.

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2959, 2871, 2225, 1730, 1603, 1448, 1387, 1368, 1291, 1264, 1239, 1214, 1190, 1140, 1101, 1029, 918, 852, 777.

Microanalysis:  $\text{C}_{14}\text{H}_{21}\text{NO}_2$ :

20 Calc: C, 71.46; H, 8.99; N, 5.95.

Found: C, 71.28; H, 8.95; N, 5.90.

Step (ii) Dinitrile

25 The cyanoacetate (37 mmol) and NaCN (37 mmol) were reacted as in the general method Step (ii) to give a yellow solid. Yield 100%. Mpt 79-81°C.

30  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.91 (6H, d,  $J = 6.8$  Hz), 1.00-1.20 (1H, m), 1.3-1.6 (5H, m), 1.85 (2H, d,  $J = 12.8$  Hz), 2.14 (2H, d,  $J = 12$  Hz), 2.70 (2H, m). MS (CI)  $m/z$ : 95, 121, 148, 164, 191 (100%  $\text{MH}^+$ ), 192, 209, 210, 219, 231.

IR ( $\text{CH}_2\text{Cl}_2$ )  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2961, 2933, 2868, 2250, 2237, 1468, 1451, 1388, 1370, 1344, 1318, 1266, 1238, 1216, 1146, 1093, 1065, 1035, 998, 966, 934, 909, 738.

-42-

Microanalysis:  $C_{12}H_{18}N_2$ :

Calc: C, 75.74; H, 9.53; N, 14.72.

Found: C, 75.45; H, 9.51; N, 14.64.

5 Step (iii) Imidate

The binitrile (12.3 mmol) was reacted as in the general method Step (iii) to give a slightly impure white solid. No purification was attempted and solid was used in next step.

10

Step (iv) Ester

The imidate (4.4 mmol) was reacted as in the general method Step (iv) to leave a low melting solid. Yield 76% based on binitrile.

15  $^1H$  NMR ( $CDCl_3$ ) 400 MHz:  $\delta$  0.89 (6H, d,  $J = 6.8$  Hz), 0.91-1.04 (1H, m), 1.29 (3H, t,  $J = 7$  Hz), 1.33-1.51 (5H, m), 1.74-1.78 (2H, m), 2.14-2.17 (2H, m), 2.54 (2H, s), 4.17-4.22 (2H, q,  $J = 7$  Hz).

MS (CI)  $m/z$ : 88, 123, 150, 192 ( $MH^+$  - EtOH), 210 ( $MH^+$  - CO), 238 (100%  $MH^+$ ).

20 IR (Film)  $\nu_{max}$   $cm^{-1}$ : 2955, 2927, 2863, 2235, 1733, 1450, 1369, 1244, 1187, 1030, 933.

Microanalysis:  $C_{14}H_{23}NO_2 \cdot 0.12 H_2O$ :

Calc: C, 70.21; H, 9.78; N, 5.85.

25 Found: C, 70.18; H, 9.82; N, 6.03.

Step (v) Lactam

The ester (2.9 mmol) was hydrogenated as in the general method Step (v) at 50°C, 50 psi, to give a crude solid. The solid was purified by column chromatography to give a white solid. Yield 38%. Mpt 130-134°C.

30  $^1H$  NMR ( $CDCl_3$ ) 400 MHz:  $\delta$  0.85-0.90 (6H, dd,  $J = 0.8$ , 6.8 Hz), 1.00-1.05 (3H, m), 1.34-1.45 (3H, m), 1.63-1.65 (2H, m), 1.73-1.81 (2H, m), 2.13 (2H, d,  $J = 0.8$  Hz), 3.19 (2H, s), 5.91 (1H, Br s).

35

-43-

MS (CI) m/z: 95, 152, 194, 195, 196 (100% MH<sup>+</sup>), 197, 210, 224.

IR (CH<sub>2</sub>Cl<sub>2</sub>)  $\nu_{\max}$  cm<sup>-1</sup>: 3210, 3094, 2931, 2857, 1699, 1493, 1449, 1382, 1322, 1301, 1265, 919, 788.

5 Microanalysis: C<sub>12</sub>H<sub>21</sub>NO:

Calc: C, 73.80; H, 10.84; N, 7.77.

Found: C, 73.83; H, 10.90; N, 7.11.

Step (vi) 4-isopropyl gabapentin

10 The lactam (1 mmol) was reacted as in the general method Step (vi) to give a white powder. Yield 60%. Mpt 167-170°C.  $[\alpha]_D = 0$  (T = 20°C, C = 1, MeOH). One diastereomer (cis).

<sup>1</sup>H NMR (DMSO) 400 MHz:  $\delta$  0.84 (6H, d, J = 6.8 Hz),  
15 0.90-1.00 (1H, m), 1.00-1.56 (2H, m), 1.23-1.30 (2H, m), 1.38-1.48 (3H, m), 1.66-1.70 (2H, m), 2.32 (2H, s), 2.97 (2H, s), 8.00 (3H, Br s), 12.00 (1H, Br s).

MS (CI) m/z: 190, 196 (100% lactam H<sup>+</sup>), 214 (MH<sup>+</sup>).

20 IR (MeOH)  $\nu_{\max}$  cm<sup>-1</sup>: 3557, 3144, 3027, 2949, 2865, 2354, 1712, 1591, 1507, 1455, 1468, 1409, 1322, 1286, 1246, 1199, 1077, 852.


Microanalysis: C<sub>12</sub>H<sub>23</sub>NO<sub>2</sub>·1.12 HCl:

Calc: C, 56.71; H, 9.57; N, 5.51.

25 Found: C, 56.77; H, 9.56; N, 5.51.

- 44 -

## EXAMPLE 5



- (i)  $\text{CuI}$ ,  $\text{MeLi}$ ,  $\text{NH}_4\text{Cl}$ ,  $\text{NH}_3$  (92%);  
 (ii)  $\text{NCCH}_2\text{CO}_2\text{Et}$ ,  $\text{NH}_4\text{OAc}$ ,  $\text{AcOH}$ , toluene (83%);  
 (iii)  $\text{NaCN}$ ,  $\text{EtOH}$ ,  $\text{H}_2\text{O}$  (57%);  
 (iv)  $\text{HCl}$ ,  $\text{EtOH}$ , toluene (93%);  
 (v)  $\text{H}_2$ , Raney Nickel,  $\text{EtOH}$ ,  $\text{NH}_3$  (84%);  
 (vi)  $\text{HCl}$ ,  $\text{H}_2\text{O}$  (64%)

Step (i) 3,3-Dimethyl-cyclohexanone

Synthesised via the method outlined by  
 Pelletier S.W. and Mody N.V., J. Org. Chem., 41:1069  
 (1969).

-45-

A solution of lithium dimethyl cuprate was prepared by the addition of methyl lithium (1.4 M in ether, 77.25 mL, 2.45 mol) to copper (I) iodide (8.8 g, 0.046 mol) under argon. The solution was cooled to 0°C, and 3-methyl-cyclohexen-1-one (5 mL, 0.044 mol) was added dropwise, with stirring, and a deep yellow precipitate was formed.

The suspension was stirred at room temperature for 1 hour before being poured into a solution of aqueous ammonia (100 mL) and ammonium acetate (ca. 5 g). The layers were separated and the aqueous layer was washed with diethyl ether (3 × 50 mL). The combined organics were washed with saturated brine (3 × 100 mL), dried (MgSO<sub>4</sub>), and the solvent removed in vacuo to leave a dark yellow liquid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: 0.98 (6H, s, 2 × Me), 1.59 (2H, m), 1.88 (2H, m), 2.14 (2H, m), 2.26 (2H, m).

IR (Film)  $\nu_{\max}$  cm<sup>-1</sup>: 2956, 1711 (C = O), 1457, 1368, 1292, 1226, 1076.

#### Step (ii) Cyanoacetate

To a solution of 3,3-dimethyl-cyclohexanone (4 g, 0.032 mol) in toluene (25 mL) was added ethyl cyanoacetate (3.37 mL, 0.032 mol, 1 eq.), ammonium acetate (0.24 g, 0.003 mol, 0.1 eq.), and acetic acid (0.36 mL, 0.006 mol, 0.2 eq.). The yellow solution was heated to reflux while attached to a Dean-Stark trap, and heating was continued until no more water condensed in the trap. After cooling, the now orange solution was washed with water (3 × 2.5 mL) and the organic layer dried (MgSO<sub>4</sub>). Filtration and removal of the solvent in vacuo gave the crude product as a deep orange liquid. Purification was achieved by Kugelrohr distillation to leave the mixture of cis and trans products as a pale yellow liquid, bp 160-170°C, 4 mbar (5.83 g, 83%).

-46-

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 0.96 (6H, s, 2  $\times$  Me), 0.99 (6H, s, 2  $\times$  Me), 1.34 (6H, m, 2  $\times$  Me of ester), 1.49 (4H, m), 1.75 (2H, quin,  $J = 6.4$ ), 1.82 (2H, quin,  $J = 6.4$ ), 2.46 (2H, s), 2.60 (2H, t,  $J = 6.4$ ), 2.80 (2H, s) 2.93 (2H, t,  $J = 6.4$ ), 4.27 (4H, m, 2  $\times$   $\text{CH}_2$  ester).

MS (CI)  $z/e$ : 222 ( $\text{M}^+ = 1$ , 100%), 221 (5), 206 (4), 194 (6), 176 (5).

IR (Film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 2958, 2870, 2224 (CN), 1731 (C = O), 1606 (C = C), 1277, 1223.

Microanalysis:  $\text{C}_{13}\text{H}_{19}\text{O}_2\text{N}$ :

Calc: C, 70.56; H, 8.65; N, 6.32.

Found: C, 70.35; H, 8.79; N, 6.25.

#### Step (iii) Bisnitrile

To a solution of the unsaturated cyanoester (1.26 g, 0.006 mol) in ethanol (100 mL) and water (4 mL) was added sodium cyanide (0.28 g, 0.006 mol, 1 eq.). The yellowish solution was heated to reflux for 8 hours and then cooled, during which time an off-white precipitate was formed. The suspension was filtered under vacuum and the filtrate acidified with HCl gas until the pH was approximately 2. The mixture was then filtered a second time and then the solvent removed in vacuo to leave the crude product as a pale green solid. Flash column chromatography, after absorption of the crude product on to silica, eluting with 0% to 50% EtOAc in heptane gave the binitrile as a colorless solid (0.57 g, 57%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 0.99 (3H, s, Me), 1.13 (1H, td,  $J = 13.2$ , 4.2 Hz), 1.21 (3H, s, Me), 1.32 (2H, m), 1.54 (1H, m), 1.82 (3H, m), 2.15 (1H, m), 2.65 (2H, s,  $\text{CH}_2\text{CN}$ ).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 19.61, 25.17, 30.79, 31.18, 33.77, 34.79, 35.37, 37.92, 46.26, 115.06, 122.19.



-47-

MS (CI)  $z/e$ : 177 ( $M^+ + 1$ , 100%), 161 (10), 150 (20), 136 (5), 120 (4), 109 (5).

IR (Film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 2988, 2937, 2869, 2237 ( $2 \times \text{CN}$ ), 1749, 1456, 1423, 1369, 1202, 1180, 1031, 972.

5 Microanalysis:  $\text{C}_{11}\text{H}_{16}\text{N}_2$ :

Calc: C, 74.96; H, 9.15; N, 15.89.

Found: C, 75.08; H, 9.32; N, 15.80.

Step (iv) Cyanoester

10 The binitrile (0.50 g, 2.84 mmol) was dissolved in absolute ethanol (20 mL) at room temperature and then cooled to  $0^\circ\text{C}$ . Toluene (20 mL) was added to the solution and then the reaction mixture was acidified by passing HCl gas through it at a gentle rate for  
15 ca. 45 minutes. The flask was then stoppered and left to stand at room temperature for 24 hours.

The yellow solution was partitioned between ethyl acetate and water and the layers separated. The aqueous layer was extracted with ethyl acetate ( $3 \times$   
20 30 mL), and the combined organics washed with aqueous saturated sodium hydrogen carbonate solution ( $3 \times$  50 mL), brine ( $3 \times$  50 mL), dried ( $\text{MgSO}_4$ ), and the solvent removed under reduced pressure to leave a pale yellow liquid (0.59 g, 93%).

25  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 0.94 (3H, s, Me), 1.16 (3H, m), 1.21 (3H, s, Me), 1.29 (3H, t,  $J = 7.2$ ,  $\text{CH}_2\text{CH}_3$ ), 1.50 (1H, m), 1.65 (1H, dt,  $J = 14.4$ , 7.6), 1.84 (1H, qt,  $J = 13.3$ , 3.2), 1.96 (1H, dt,  $J = 13.7$ , 2.2), 2.16 (1H, m), 2.48 (1H, d,  $J = 15.6$ , C-2H), 2.54  
30 (1H, d,  $J = 15.6$ , C-2H), 4.20 (2H, q,  $J = 7.2$ ,  $\text{CH}_2\text{CH}_3$ ).  
 $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 14.21, 19.65, 25.42, 31.03, 34.04, 34.14, 36.08, 38.44, 46.14, 46.80, 61.02, 123.67, 169.00.

MS (CI)  $z/e$ : 224 ( $M^+ + 1$ , 100%), 196 (12), 178 (35),  
35 136 (13), 109 (12).

-48-

IR (Film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 2998, 2937, 2868, 2234 (CN),  
1738 (C = O), 1457, 1372, 1217, 1181, 1154, 1026.

Microanalysis:  $\text{C}_{13}\text{H}_{21}\text{NO}_2$ :

Calc: C, 69.92; H, 9.48; N, 6.27.

5 Found: C, 69.63; H, 9.45; N, 6.15.

Step (v) Lactam

The cyanoester (0.5 g, 2.23 mmol) was hydrogenated  
in ethanolic ammonia (600 mL) with Raney nickel as  
10 catalyst (ca. 0.25 g) at 50°C and 50 psi for 48 hours.

The catalyst was then removed by filtration  
through Celite, and the solvent removed in vacuo to  
leave a greenish crystalline solid.

Flash column chromatography, eluting with 0% to  
15 100% ethyl acetate in heptane, gave the pure lactam as  
a colorless solid (340 mg, 84%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 0.89 (3H, s, Me), 0.92 (3H, s,  
Me), 1.25 (2H, m), 1.36 (2H, m), 1.51 (3H, m),  
1.68 (1H, s), 2.18 (1H, d,  $J = 16.4$ ,  $\text{CH}_2\text{NH}$ ), 2.24 (1H,  
20 d,  $J = 16.8$ ,  $\text{CH}_2\text{NH}$ ), 3.15 (2H, s,  $\text{CH}_2\text{CO}$ ).

$^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) 400 MHz: 19.16, 29.88, 30.36, 31.28,  
36.57, 39.05, 39.61, 44.58, 49.54, 54.79, 177.72.

MS (CI)  $z/e$ : 182 ( $\text{M}^+ + 1$ , 100%), 181 (15), 180 (5),  
166 (3).

25 IR (Film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 3203, 3100 (NH), 2914, 2860,  
1698 (C = O), 1486, 1374, 1317, 1289, 1257, 1076.

Microanalysis:  $\text{C}_{11}\text{H}_{19}\text{NO}$ :

Calc: C, 72.88; H, 10.56; N, 7.73.

Found: C, 72.38; H, 10.47; N, 7.56.

30

Step (vi) 3,3-Dimethyl Gabapentin Hydrochloride

The lactam (0.3 g, 1.66 mmol) was dissolved in a  
mixture of HCl (concentrated, 5 mL) and water (5 mL),  
and the resultant colorless solution heated to reflux  
35 for 20 hours. The solution was cooled and then  
partitioned between water and dichloromethane, and the

-49-

layers separated. The aqueous layer was washed with dichloromethane (3 × 20 mL) and the water/HCl removed by rotary evaporation to leave the crude product as an off-white solid. Trituration of this solid with ethyl acetate and filtration of the product gave  
5 3,3-dimethylgabapentin, hydrochloride salt as a colorless solid (140 mg, 42%, 64% based on recovered starting material).

<sup>1</sup>H NMR (DMSO) 400 MHz: 0.90 (3H, s, Me), 0.92 (3H, s, Me), 1.15-1.49 (8H, m), 2.45 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.90 (2H, br q, J = 13.5, CH<sub>2</sub>NH<sub>3</sub>), 7.96 (3H, br s, NH<sub>3</sub>), 12.36 (1H, br s, OH).

IR (Film)  $\nu_{\max}$  cm<sup>-1</sup>: 2930, 1728 (C = O), 1272, 1123.

Microanalysis: C<sub>11</sub>H<sub>22</sub>NO<sub>2</sub>Cl:

15 Calc: C, 56.04; H, 9.41; N, 5.94.

Found: C, 55.79; H, 9.61; N, 6.23.

20

25


30

35


- 50 -

## EXAMPLE 6

5



10



15

(i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ , MeOH,  $\text{NH}_3$  (67%);(ii)  $\text{H}_2\text{SO}_4$ ,  $180^\circ\text{C}$  (79%);(iii)  $\text{Ac}_2\text{O}$ ,  $120^\circ\text{C}$  (82%);

20

(iv) MeOH, NaOMe (83%);

(v)  $\text{N}_3\text{P}(\text{O})(\text{OPh})_2$ ,  $\text{Et}_3\text{N}$ , MeCN,  $60^\circ\text{C}$  then HCl reflux, then crystallization (13%)

Steps (i) and (ii)

25

(R)-3-Methylcyclohexanone (10.92 mL, 89.2 mmol)

was dissolved in methanol (25 mL) with ethylcyanoacetate (18.96 mL, 178 mmol) and cooled to  $0^\circ\text{C}$ .

Ammonia gas was bubbled through the solution for

30

25 minutes, after which the solution was stoppered and

stored at  $-20^\circ\text{C}$ . After 66 hours, diethyl ether

(100 mL) was added to the mixture, and the white solid

which formed was filtered off, washed with diethyl

ester ( $2 \times 50$  mL), and dried to give 15.71 g (67%) of a white solid.

35

Without further purification, a sample of the solid (4.0 g, 15.3 mmol) was dissolved in concentrated

-51-

H<sub>2</sub>SO<sub>4</sub> (40 mL) with gentle warming and allowed to stand overnight. Water (40 mL) was then cautiously added and the resulting mixture heated to 170°C. After 5 hours, all the solid had dissolved. The mixture was cooled to room temperature, diluted with water (200 mL), and extracted with diethyl ether (3 × 150 mL). The ether extracts were combined, dried over magnesium sulphate, and the solvent removed in vacuo. The oily residue was triturated with heptane to obtain a precipitate which was filtered off and dried to give 2.57 g (79%) of a buff colored solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 0.85-0.94 (2H, m), 0.87 (3H, d, J = 6 Hz), 1.15 (1H, m), 1.39-1.61 (3H, m), 1.71 (1H, br d, J = 12.8 Hz), 1.87 (2H, m), 2.48 (2H, ABq, J = 4 Hz), 2.67 (2H, s).

MS (ES) z/e: 214 ([M]<sup>+</sup>, 13%), 213 (100%).

IR (thin film) ν<sub>max</sub> cm<sup>-1</sup>: 1204, 1290, 1413, 1458, 1702, 2924.

Microanalysis: C<sub>11</sub>H<sub>18</sub>O<sub>4</sub>:

Calc: C, 61.66; H, 8.47.

Found: C, 61.67; H, 8.51.

#### Step (iii) Anhydride

The diacid (2.5 g, 11.68 mmol) was heated to reflux in acetic anhydride (30 mL). After 3 hours, the solvent was removed in vacuo. The residue was dissolved in dichloromethane (50 mL) and washed with saturated aqueous sodium bicarbonate, dried (MgSO<sub>4</sub>), and the solvent removed in vacuo to obtain 1.83 g (82%) of a brown oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 0.84, 0.89 (3H, d, J = 6 Hz), 0.98 (1H, m), 1.38-1.60 (4H, m), 1.64-1.80 (2H, m), 2.53 (2H, s), 2.74 (2H, s).

MS (APCI+) z/e: 197 ([MH]<sup>+</sup>, 100%), 126 (32%).

IR (thin film) ν<sub>max</sub> cm<sup>-1</sup>: 947, 1073, 1181, 1761, 1810, 2925.

-52-

Microanalysis:  $C_{11}H_{16}O_3$ :

Calc: C, 67.32; H, 8.22.

Found: C, 66.98; H, 8.07.

5 Step (iv) Half Ester, cis/trans Mixture

The anhydride (1.865 g, 9.5 mmol) was dissolved in dry methanol (10 mL) with sodium methoxide (0.5 M in MeOH, 20 mL, 10 mmol) and stirred at room temperature. After 3 hours the solvent was removed in vacuo and the residue partitioned between ethyl acetate (150 mL) and 1N HCl (50 mL). The organic phase was separated and the aqueous phase re-extracted with ethyl acetate (2 × 100 mL). The organic extracts were combined, dried ( $MgSO_4$ ), and the solvent removed in vacuo to give 1.8 g (83%) of a pale brown oil which contained a -1:1 mixture of the cis and trans isomers.

$^1H$  NMR ( $CDCl_3$ ) 400 MHz:  $\delta$  0.85-0.93 (2H, m); 0.86 (3H, d,  $J = 6$  Hz); 1.17 (1H, m); 1.39-1.62 (3H, m); 1.64-1.80 (3H, m); 2.48 (2H, m); 2.64-2.65 (2H, 2 × s one from each isomer).

MS (ES-)  $z/e$ : 227 ( $[M-H]^+$ , 100%).

IR (thin film)  $\nu_{max}$   $cm^{-1}$ : 1163, 1194, 1440, 1705, 1738, 2926, 3200.

Microanalysis:  $C_{12}H_{20}O_4$ :

25 Calc: C, 63.13; H, 8.83.

Found: C, 63.29; H, 8.83.

Step (v) (1-Aminomethyl-3-methyl-cyclohexyl)-acetic acid [(1S-(1 $\alpha$ ,3 $\beta$ )]

30 The mixture of half ester isomers (515 mg, 2.26 mmol) was dissolved in acetone (6 mL) and cooled to  $-10^\circ C$ . Triethylamine (377  $\mu L$ , 2.7 mmol) was added followed by ethyl chloroformate (259  $\mu L$ , 2.7 mmol). The mixture was stirred at  $-10^\circ C$  for 40 minutes, after which a solution of sodium azide (220 mg, 3.39 mmol) in water (1 mL) was added and the mixture allowed to warm

35

-53-

to 0°C. After 40 minutes, the mixture was poured into ice cold water (20 mL) and extracted with ice cold toluene (3 × 20 mL). The toluene extracts were combined and dried over magnesium sulfate at 0°C. The toluene solution was then added dropwise into a flask preheated to 180°C in an oil bath at 180°C. The solvent was removed via distillation. Once the addition was complete, the mixture was stirred at 180°C for a further 20 minutes, until all the solvent had been removed. Dioxane (5 mL) and concentrated HCl (5 mL) were then added and the mixture refluxed for 3 hours. The mixture was then cooled to room temperature, diluted with water (30 mL), and washed with dichloromethane (2 × 30 mL). The aqueous phase was collected and the solvent removed in vacuo to give a brown gum, which was triturated with ethyl acetate to give a buff colored solid. The solid was recrystallized from a mixture of methanol, ethyl acetate, and heptane to yield 35 mg (7%) of a white solid.

<sup>1</sup>H NMR (d<sub>6</sub> DMSO) 400 MHz: δ 0.70-0.88 (2H, m), 0.83 (3H, d, J = 6 Hz), 1.06-1.17 (1H, m), 1.36-1.69 (6H, m), 2.44 (2H, s), 2.84 (2H, s), 7.92 (4H, br s). MS (ES+) z/e: 186 ([MH-HCl]<sup>+</sup>, 100%). IR (thin film) ν<sub>max</sub> cm<sup>-1</sup>: 1211, 1408, 1709, 2925, 3200.

Microanalysis: C<sub>10</sub>H<sub>20</sub>NO<sub>2</sub>Cl·0.25 H<sub>2</sub>O:


Calc: C, 53.09; H, 9.13; N, 6.19; Cl, 15.67.

Found: C, 53.24; H, 9.26; N, 6.23; Cl, 15.43.

30

- 54 -

## EXAMPLE 7



- 20 (i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ , MeOH,  $\text{NH}_3$ ;  
 (ii)  $\text{H}_2\text{SO}_4$ ,  $180^\circ\text{C}$ ;  
 (iii)  $\text{Ac}_2\text{O}$ ,  $120^\circ\text{C}$ ;  
 (iv) MeOH, NaOMe;  
 (v)  $\text{EtOCOCl}$ ,  $\text{NEt}_3$ , then  $\text{NaN}_3$ , then heat, then HCl  
 25 reflux

Cis/trans 3,5-dimethyl gabapentinSteps (i) and (ii) Diacid

30 Cis-3,5-dimethyl-cyclohexanone (11.24 g, 89.2 mmol) was dissolved in methanol (25 mL) ethyl cyanoacetate (18.96 mL, 178.2 mmol) and cooled to  $0^\circ\text{C}$ . Ammonia gas was then bubbled through the solution for 30 minutes. The solution was then stored at  $-20^\circ\text{C}$ . After 66 hours, the solid was filtered off, washed with  
 35 ether, and dried to yield 18.46 g (75%) of a white solid.



-55-

Without further purification, a portion of the solid prepared above (6.0 g, 21.7 mmol) was dissolved in concentrated sulphuric acid (40 mL) with warming and left to stand overnight. Water (40 mL) was then  
5 cautiously added and the resulting solution heated to 180°C. After 5 hours, the mixture was cooled to room temperature, diluted with water (200 mL), and extracted with diethyl ether (3 × 150 mL). The organic extracts were combined, washed with brine, dried (MgSO<sub>4</sub>), and  
10 the solvent removed in vacuo. The residue was triturated with heptane to obtain a solid which was recrystallized from a dichloromethane/heptane mixture to obtain 3.122 g (63%) of a buff colored solid.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 0.49 (1H, m), 0.80 (2H, m),  
15 0.87 (6H, d, J = 6 Hz), 1.55-1.76 (3H, m), 1.85 (2H, br, d, J = 13.2 Hz), 2.50 (2H, s), 2.67 (2H, s).  
MS (ES) z/e: 228 ([M]<sup>+</sup>, 14%), 227 ([M-H]<sup>+</sup>, 100%).  
IR (thin film) ν<sub>max</sub> cm<sup>-1</sup>: 893, 1147, 1208, 1284, 1312, 1337, 1407, 1450, 1699, 2846, 2914, 2947, 3100.  
20 Microanalysis: C<sub>12</sub>H<sub>20</sub>O<sub>4</sub>:  
Calc: C, 63.13; H, 8.83.  
Found: C, 63.22; H, 8.95.

Step (iii) Anhydride

25 The diacid (3.0 g, 13.16 mmol) was dissolved in acetic anhydride (40 mL) and heated to reflux. After 3 hours, the mixture was cooled to room temperature and the solvent removed in vacuo. The residue was dissolved in dichloromethane (150 mL) and washed once  
30 with saturated aqueous sodium bicarbonate. The organic phase was separated, dried (MgSO<sub>4</sub>), and the solvent removed in vacuo to obtain 2.60 g (94%) of a brown oil which solidified on standing.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>) 400 MHz: δ 0.53 (1H, m), 0.81-0.96  
35 (2H, m, and 6H, d, J = 6 Hz), 1.43-1.71 (4H, m), 1.76 (1H, m), 2.54 (2H, s), 2.73 (2H, s).

-56-

MS (APCI+)  $z/e$ : 211 ( $[MH]^+$ , 100%).

IR (thin film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 950, 1073, 1183, 1459, 1756, 1767, 1812, 2910, 2952.

Microanalysis:  $\text{C}_{12}\text{H}_{18}\text{O}_3$ :

5 Calc: C, 68.55; H, 8.75.

Found: C, 68.32; H, 8.75.

Step (iv) Cis/trans Half Ester

10 The anhydride (2.556 g, 12.17 mmol) was dissolved in dry methanol (15 mL) and stirred with in vacuo and the residue partitioned between 1N HCl (150 mL) and ethyl acetate (150 mL). The organic phase was separated, washed with brine, dried ( $\text{MgSO}_4$ ), and the solvent removed in vacuo to give a yellow oil. This  
15 was purified by flash chromatography (silica, ethyl acetate:heptane, 1:1) to give 2.68 g (91%) of a colorless oil.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.47 (2H, m), 0.82 (4H, m), 0.87 (12H, d,  $J = 6$  Hz), 1.57-1.80 (10H, m), 2.46 (2H, s, isomer A), 2.48 (2H, s, isomer B), 2.63 (2H, s, isomer B), 2.64 (2H, s, isomer A), 3.67 (6H, s).  
20

MS (ES-)  $z/e$ : 241 ( $[M-H]^+$ , 100%).

IR (thin film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 1163, 1197, 1437, 1459, 1706, 1736, 2913, 2951, 3100.

25 Microanalysis:  $\text{C}_{13}\text{H}_{22}\text{O}_4$ :

Calc: C, 64.44; H, 9.15.

Found: C, 64.17; H, 9.17.

Step (v) Cis/trans-3,5-Dimethyl gabapentin

30 The cis/trans mixture of half esters (1.09 g, 4.5 mmol) was dissolved in acetone (15 mL) and cooled to  $-10^\circ\text{C}$ . Triethylamine (predried over lithium aluminum hydride) (660  $\mu\text{L}$ , 4.74 mmol) was then added followed by ethyl chloroformate (453  $\mu\text{L}$ , 4.764 mmol).  
35 After 40 minutes at  $10^\circ\text{C}$ , a solution of sodium azide (337 mg, 5.19 mmol) in water (2.5 mL) was added and the

-57-

mixture allowed to warm to 0°C. After 40 minutes, the mixture was poured into ice cold water (30 mL) and extracted with ice cold toluene (3 × 20 mL). The organic extracts were combined, dried (MgSO<sub>4</sub>), and stored at 0°C. The toluene solution was then added dropwise to a flask set up for distillation in an oil bath set at 180°C. The solvent was removed by distillation during the additions. After the addition was complete, the mixture was stirred at 180°C for 1 hour, after which a gentle stream of nitrogen was passed through the apparatus to remove the last traces of solvent. Hydrochloric acid (75% v/v, 20 mL) was then added cautiously, and the resulting solution refluxed for 3 hours. The mixture was cooled to room temperature and stored at room temperature overnight. The mixture was diluted with water (20 mL) and extracted with dichloromethane (2 × 15 mL). The aqueous phase was collected and the solvent removed in vacuo. The residue was triturated with ethyl acetate to obtain 255 mg (24%) of a white solid.

<sup>1</sup>H NMR (d<sub>6</sub> DMSO) 400 MHz: δ 0.46 (2H, m), 0.76-0.90 (16H, m), 1.50-1.70 (10H, m), 2.30 (2H, s, isomer A), 2.44 (2H, s, isomer B), 2.84 (2H, s, isomer B), 3.00 (2H, s, isomer A), 7.91 (6H, br s), 12.40 (2H, br s).

MS (ES+) z/e: 200 ([MH-HCl]<sup>+</sup>, 100%).

IR (thin film) ν<sub>max</sub> cm<sup>-1</sup>: 1201, 1458, 1715, 2949, 3200.


Microanalysis: C<sub>11</sub>H<sub>22</sub>NO<sub>2</sub>Cl:

Calc: C, 56.04; H, 9.41; N, 5.94.

Found: C, 55.75; H, 9.46; N, 5.87.

- 58 -

## EXAMPLE 8



- 20 (i)  $\text{EtO}_2\text{CCH}_2\text{CN}$ , MeOH,  $\text{NH}_3$ ;  
 (ii)  $\text{H}_2\text{SO}_4$ ,  $180^\circ\text{C}$ ;  
 (iii)  $\text{Ac}_2\text{O}$ ,  $120^\circ\text{C}$ ;  
 (iv) MeOH, NaOMe;  
 (v)  $\text{EtOCOCl}$ ,  $\text{NEt}_3$ , then  $\text{NaN}_3$ , then heat, then HCl  
 25 reflux

Cis/trans 4-methyl gabapentinSteps (i) and (ii) Diacid

30 4-Methylcyclohexanone (5 mL, 40.74 mmol) was dissolved in methanol (15 mL) with ethyl cyanoacetate (8.67 mL, 81.48 mmol) and cooled to  $0^\circ\text{C}$ . Ammonia gas was bubbled through the solution for 25 minutes, after which the solution was stoppered and stored at  $-20^\circ\text{C}$ . After 20 hours, diethyl ether (100 mL) was added to the mixture, and the white solid which formed was filtered

35

-59-

off, washed with diethyl ether ( $2 \times 50$  mL), and dried to give 7.51 g (70%) of a white solid.

Without further purification, a sample of the solid (4.0 g, 15.3 mmol) was dissolved in concentrated  $\text{H}_2\text{SO}_4$  (40 mL) with gentle warming and allowed to stand overnight. Water (40 mL) was then cautiously added and the resulting mixture heated to  $170^\circ\text{C}$ . After 3 hours, all the solid had dissolved. The mixture was cooled to room temperature, diluted with water (150 mL), and extracted with diethyl ether ( $3 \times 100$  mL). The ether extracts were combined, dried over magnesium sulphate, and the solvent removed in vacuo. The oily residue was triturated with heptane to obtain a precipitate which was filtered off and dried to give 2.3 g (73%) of a buff colored solid.

$^1\text{H}$  NMR ( $\text{d}_6$  DMSO) 400 MHz:  $\delta$  0.87 (3H, d,  $J = 6$  Hz); 1.1 (2H, m); 1.27 (3H, m); 1.44 (2H, m); 1.70 (2H, br d,  $J = 13$  Hz); 2.34 (2H, s); 2.45 (2H, s).

MS (ES-)  $z/e$ : 214 ( $[\text{M}]^+$ , 13%), 213 ( $[\text{M}-\text{H}]^+$ , 100%).

IR (thin film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 917, 1183, 1215, 1289, 1349, 1399, 1455, 1704, 2858, 2925, 3100.

Microanalysis:  $\text{C}_{11}\text{H}_{18}\text{O}_4$ :

Calc: C, 61.66; H, 8.47.

Found: C, 61.54; H, 8.47.

#### Step (iii) Anhydride

The diacid (2.30 g, 10.75 mmol) was heated to reflux in acetic anhydride (30 mL). After 3.5 hours, the solvent was removed in vacuo. The residue was dissolved in dichloromethane (50 mL) and washed with saturated aqueous sodium bicarbonate, dried ( $\text{MgSO}_4$ ), and the solvent removed in vacuo to obtain 2.07 g (98%) of a brown oil which solidified on standing.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.93 (3H, d,  $J = 6$  Hz), 1.07 (2H, m), 1.37 (3H, m), 1.49-1.71 (4H, m), 2.56 (2H, s), 2.72 (2H, s).

-60-

MS (APCI+)  $z/e$ : 197 ( $[MH]^+$ , 100%).

IR (thin film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 953, 1064, 1183, 1241, 1455, 1761, 1810, 2924.

Microanalysis:  $\text{C}_{11}\text{H}_{16}\text{N}_3$ :

5 Calc: C, 67.32; H, 8.22.

Found: C, 67.41; H, 8.29.

Step (iv) Cis/trans Half Ester

The anhydride (2.06 g, 10.5 mmol) was dissolved in  
10 dry methanol (40 mL) and stirred with sodium methoxide  
(624 mg, 11.55 mmol). After 4 hours, the solvent was  
removed in vacuo and the residue partitioned between  
1N HCl (150 mL) and dichloromethane (150 mL). The  
organic phase was separated, washed with brine, dried  
15 ( $\text{MgSO}_4$ ), and the solvent removed in vacuo to give a  
yellow oil. This was purified by flash chromatography  
(silica, ethyl acetate:heptane, 1:1) to give  
1.98 g (83%) of a colorless oil.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ ) 400 MHz:  $\delta$  0.83-0.92 (2H, m), 0.91 (6H,  
20 d,  $J = 6$  Hz), 1.14 (4H, m), 1.21-1.42 (4H, m), 1.54  
(4H, m), 1.77 (4H, m), 2.49 (2H, s, isomer A), 2.50  
(2H, s, isomer B), 2.62 (2H, s, isomer B), 2.64 (2H, s,  
isomer A), 3.66 (3H, s, isomer A), 3.67 (3H, s,  
isomer B).

25 MS (ES-)  $z/e$ : 227 ( $[M-H]^+$ , 100%).

IR (thin film)  $\nu_{\max}$   $\text{cm}^{-1}$ : 1162, 1193, 1434, 1699,  
1731, 2922, 3200.

Microanalysis:  $\text{C}_{12}\text{H}_{20}\text{O}_4$ :

Calc: C, 63.13; H, 8.83.

30 Found: C, 63.12; H, 8.71.

Step (v) Cis/trans 4-methyl gabapentin

The cis/trans mixture of half esters (1.90 g,  
8.3 mmol) was dissolved in acetone (20 mL) and cooled  
35 to  $-10^\circ\text{C}$ . Triethylamine (predried over lithium  
aluminium hydride) (1.21 mL, 8.7 mmol) was then added

-61-

followed by ethyl chloroformate (832  $\mu$ L, 8.7 mmol). After 50 minutes at  $-10^{\circ}\text{C}$ , a solution of sodium azide (630 mg, 9.69 mmol) in water (5 mL) was added and the mixture allowed to warm to  $0^{\circ}\text{C}$ . After 40 minutes, the mixture was poured into ice cold water (50 mL) and extracted with ice cold toluene ( $3 \times 50$  mL). The organic extracts were combined, dried ( $\text{MgSO}_4$ ), and kept at  $0^{\circ}\text{C}$ . The toluene solution was then added dropwise to a flask set up for distillation in an oil bath set at  $180^{\circ}\text{C}$ . The solvent was removed by distillation during the addition. After the addition was complete, the mixture was stirred at  $180^{\circ}\text{C}$  for 1 hour, after which a gentle stream of nitrogen was passed through the apparatus to remove the last traces of solvent. Hydrochloric acid (75% v/v, 40 mL) was then added cautiously, and the resulting solution refluxed for 3 hours. The mixture was cooled to room temperature and sorted at room temperature overnight. The mixture was diluted with water (30 mL) and extracted with dichloromethane ( $3 \times 30$  mL). The aqueous phase was collected and the solvent removed in vacuo. The residue was triturated with ethyl acetate to obtain 590 mg (32%) of a white solid.

$^1\text{H}$  NMR ( $d_6$  DMSO) 400 MHz:  $\delta$  0.87 (6H, d,  $J = 6$  Hz), 1.07 (4H, m), 1.19-1.40 (6H, m), 1.41-1.58 (6H, m), 1.61 (2H, m), 2.32 (2H, s, isomer A), 2.44 (2H, s, isomer B), 2.85 (2H s, isomer B), 2.99 (2H, s, isomer A), 7.96 (6H, br s), 12.36 (2H br s).

MS (ES+)  $z/e$ : 186 ( $[\text{MH} - \text{HCl}]^+$ , 100%).


IR (thin film)  $\nu_{\text{max}}$   $\text{cm}^{-1}$ : 1195, 1404, 1457, 1506, 1607, 1712, 2924, 3200.

Microanalysis:  $\text{C}_{10}\text{H}_{20}\text{NO}_2\text{Cl}$ :  
Calc: C, 54.17; H, 9.09; N, 6.32.  
Found: C, 54.13; H, 9.18; N, 6.45.

- 62 -

## CLAIMS

1. A compound of formula



a pharmaceutically acceptable salt thereof or a prodrug thereof wherein

$R_1$  to  $R_{10}$  are each independently selected from straight or branched alkyl of from 1 to 6 carbon atoms, unsubstituted or substituted benzyl or phenyl which substituents are selected from halogen, alkyl, alkoxy, hydroxy, carboxy, carboalkoxy, trifluoromethyl, and nitro, and any  $R_1$  to  $R_{10}$ , which is not one of the above, is hydrogen.

2. A compound according to Claim 1 wherein  $R_1$  to  $R_{10}$  is selected from methyl, ethyl, propyl, butyl straight or branched.
3. A compound according to Claim 1 wherein  $R_5$  is tert-butyl.
4. A compound according to Claim 1 wherein  $R_3$  is methyl.
5. A compound according to Claim 1 selected from a compound of Formula I, wherein  $R_1$  is methyl or  $R_5$  is methyl.



-63-

6. A compound according to Claim 1 wherein R<sub>5</sub> is isopropyl.
7. A compound according to Claim 1 selected from  
(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid and  
(1-aminomethyl-3-cyclohexyl)-acetic acid  
5 [1R-(1 $\alpha$ ,3 $\beta$ )].
8. A compound according to Claim 1 selected from  
(1-aminomethyl-4-tert-butyl-cyclohexyl)-acetic acid and  
cis (1-aminomethyl-4-methyl-cyclohexyl)-  
5 acetic acid.
9. A compound according to Claim 1 and named  
(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid [1S-(1 $\alpha$ ,3 $\beta$ )].
10. A compound according to Claim 1 selected from  
(1-aminomethyl-3-methyl-cyclohexyl)-acetic acid methyl ester monohydrochloride;  
[1-(acetyl-amino-methyl)-3-methyl-cyclohexyl]-  
5 acetic acid; and  
[2-(1-Aminomethyl-3-methyl-cyclohexyl)-acetyl-amino]-acetic acid monohydrochloride.
11. A compound according to Claim 1 selected from  
( $\pm$ )-(1-aminomethyl-3,3-dimethyl-cyclohexyl)-acetic acid;  
(1-aminomethyl-3,3,5,5-tetramethyl-cyclohexyl)-acetic acid; and  
5 (1-aminomethyl-4-methyl-cyclohexyl)-acetic acid.

-64-

12. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to Claim 1 and a pharmaceutically acceptable carrier.
13. A method for treating epilepsy comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in need of said treatment.
14. A method for treating faintness attacks, hypokinesia, and cranial disorders comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in  
5 need of said treatment.
15. A method for treating neurodegenerative disorders comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in need of said treatment.
16. A method for treating depression comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in  
need of said treatment.
17. A method for treating anxiety comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in  
need of said treatment.
18. A method for treating panic comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in  
need of said treatment.

-65-

19. A method for treating pain comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in need of said treatment.
20. A method for treating neuropathological disorders comprising administering a therapeutically effective amount of a compound according to Claim 1 to a mammal in need of said treatment.

# INTERNATIONAL SEARCH REPORT

Int'l Application No.  
PCT/US 97/02295

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07C229/28 A61K31/195 C07C233/47 A61K31/16

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 087 544 A (SATZINGER GERHARD ET AL) 2 May 1978 cited in the application see claims ---	1-20
A	US 5 025 035 A (WALLACE JAN D) 18 June 1991 see claims ---	1-20
A	EP 0 340 677 A (WARNER LAMBERT CO) 8 November 1989 see claims -----	1-20

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

28 May 1997

Date of mailing of the international search report

05.06.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Sánchez García, J.M.

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 97/02295

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4087544 A	02-05-78	DE 2460891 A	01-07-76
		AT 340892 B	10-01-78
		AU 8774175 A	23-06-77
		BE 836835 A	18-06-76
		CA 1052811 A	17-04-79
		CH 612665 A	15-08-79
		CH 612666 A	15-08-79
		CH 612664 A	15-08-79
		DE 2543821 A	14-04-77
		FR 2294697 A	16-07-76
		GB 1465229 A	23-02-77
		JP 941538 C	20-02-79
		JP 51088940 A	04-08-76
		JP 53024064 B	18-07-78
		LU 74058 A	20-07-76
		NL 7514900 A,B,	23-06-76
		SE 423385 B	03-05-82
		SE 7514442 A	22-06-76
		US 4024175 A	17-05-77
US 5025035 A	18-06-91	CA 2092416 A	13-04-92
		EP 0552240 A	28-07-93
		JP 6502148 T	10-03-94
		WO 9206686 A	30-04-92
EP 0340677 A	08-11-89	US 4894476 A	16-01-90
		AU 624130 B	04-06-92
		AU 3267889 A	02-11-89
		CA 1306755 A	25-08-92
		DE 68912819 D	17-03-94
		DE 68912819 T	19-05-94
		ES 2061774 T	16-12-94
		IE 62958 B	08-03-95
		JP 2011546 A	16-01-90
		US 4960931 A	02-10-90