



(86) Date de dépôt PCT/PCT Filing Date: 2015/04/29
(87) Date publication PCT/PCT Publication Date: 2015/11/05
(45) Date de délivrance/Issue Date: 2023/01/17
(85) Entrée phase nationale/National Entry: 2016/10/26
(86) N° demande PCT/PCT Application No.: IB 2015/053110
(87) N° publication PCT/PCT Publication No.: 2015/166427
(30) Priorités/Priorities: 2014/04/29 (US61/985,589);
2014/05/16 (US61/994,310)


(51) Cl.Int./Int.Cl. *C12N 5/10* (2006.01),
C12N 15/09 (2006.01), *C12N 15/57* (2006.01),
C12N 15/67 (2006.01), *C12N 15/85* (2006.01),
C12N 15/87 (2006.01), *C12N 5/071* (2010.01),
C12N 9/64 (2006.01), *C12P 21/02* (2006.01),
C12Q 1/00 (2006.01), *C12Q 1/37* (2006.01)

(72) Inventeurs/Inventors:
LAUX, HOLGER, CH;
ROMAND, SANDRINE, CH;
BODENDORF, URSULA, CH

(73) Propriétaire/Owner:
NOVARTIS AG, CH

(74) Agent: SMART & BIGGAR LP

(54) Titre : NOUVELLES CELLULES DE VERTEBRE ET PROCEDES D'EXPRESSION PAR RECOMBINAISON D'UN
POLYPEPTIDE D'INTERET
(54) Title: NOVEL VERTEBRATE CELLS AND METHODS FOR RECOMBINANTLY EXPRESSING A POLYPEPTIDE OF
INTEREST



(57) Abrégé/Abstract:

The present application pertains inter alia to an isolated vertebrate cell suitable for recombinant expression of a polypeptide of interest, wherein the vertebrate cell is altered to impair the function of the endogenous protease matriptase and wherein said cell comprises at least one heterologous polynucleotide encoding a polypeptide of interest and wherein the polypeptide of interest is secreted by the cell. It was found that using respective vertebrate cells for producing a recombinant polypeptide of interest significantly reduces clipping of the polypeptide of interest that is secreted into the cell culture medium. Also provided are improved production and screening methods.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
5 November 2015 (05.11.2015)

WIPO | PCT

(10) International Publication Number
WO 2015/166427 A3

(51) International Patent Classification:

C12N 5/00 (2006.01) *C12N 15/87* (2006.01)
C12N 5/071 (2010.01) *C12N 15/85* (2006.01)
C12N 15/67 (2006.01)

(21) International Application Number:

PCT/IB2015/053110

(22) International Filing Date:

29 April 2015 (29.04.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/985,589 29 April 2014 (29.04.2014) US
61/994,310 16 May 2014 (16.05.2014) US

(71) Applicant (for all designated States except US): **NO-VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(71) Applicants (for US only): **LAUX, Holger** [DE/CH]; c/o Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **RO-MAND, Sandrine** [FR/CH]; c/o Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **BODENDORF, Ursula** [DE/CH]; c/o Novartis Pharma AG, Postfach, CH-4002 Basel (CH).

(74) Agent: **LEHMEIER, Thomas**; Novartis Pharma AG, Postfach, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
— with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:

14 January 2016

(54) Title: NOVEL VERTEBRATE CELLS AND METHODS FOR RECOMBINANTLY EXPRESSING A POLYPEPTIDE OF INTEREST

(57) Abstract: The present application pertains *inter alia* to an isolated vertebrate cell suitable for recombinant expression of a polypeptide of interest, wherein the vertebrate cell is altered to impair the function of the endogenous protease matriptase and wherein said cell comprises at least one heterologous polynucleotide encoding a polypeptide of interest and wherein the polypeptide of interest is secreted by the cell. It was found that using respective vertebrate cells for producing a recombinant polypeptide of interest significantly reduces clipping of the polypeptide of interest that is secreted into the cell culture medium. Also provided are improved production and screening methods.



WO 2015/166427 A3

“Novel vertebrate cells and methods for recombinantly expressing a polypeptide of interest “

FIELD OF THE DISCLOSURE

5

[1] The present disclosure concerns the field of recombinant expression technologies. It *inter alia* pertains to altered vertebrate cells and their use in recombinant expression methods. When recombinantly expressing a polypeptide of interest in said altered cells which is then secreted into the cell culture medium, clipping of the recombinantly expressed polypeptide of interest in the cell culture medium is significantly reduced or even completely prevented.

BACKGROUND OF THE DISCLOSURE

15 [2] The market for biopharmaceuticals continues to grow at a high rate as biopharmaceuticals become more and more important for today's medicine. Currently, an increasing number of biopharmaceuticals is produced in vertebrate cells such as in particular mammalian cells. Biopharmaceuticals include but are not limited to antibodies, antibody fragments, ADCs (antibody drug conjugate), nanobodies, Fc-fusion proteins, growth factors, hormones, cytokines, enzymes and other therapeutic polypeptides. Especially expression of recombinant non-antibody therapeutic proteins is of increasing importance. Furthermore, the recombinant expression of polypeptides in mammalian cells is also for other fields of use of high interest. Considering the production costs for recombinantly expressed polypeptides, it is important to have high expressing mammalian cell lines. The polypeptide of interest is expressed and secreted by the mammalian cells into the cell culture medium from which it is then harvested.

[3] One major problem that is encountered when using vertebrate cells such as mammalian cells as host cells for recombinant expression is the proteolytic degradation of the expressed and secreted polypeptide of interest in the cell culture medium, also referred to as “clipping”. Proteases originating from the vertebrate cells used for production are active in the cell culture medium whose proteolytic activity may degrade the recombinantly expressed and secreted polypeptide of interest, thereby rendering an altered, e.g. non- or less functional polypeptide of interest. This host cell related proteolytic degradation is one of the major hurdles in the recombinant expression of polypeptides. Even though IgG antibodies may also be affected, e.g. because IgG antibodies may be clipped in the variable regions, proteolytic degradation, i.e. clipping, occurs at a much higher frequency with non-IgG polypeptides. Such non-IgG polypeptides, in particular glycopolypeptides, due to their relatively exposed three-dimensional structure, can be very sensitive to proteolytic degradation. Clipping of recombinantly expressed non-IgG polypeptides can reach values up to 100%. Clipping can lead to inactive and/or immunogenic polypeptides which are not useful for the intended purpose. Furthermore, even if proteolytic degradation only occurs to a certain percentage, the clipped polypeptide cannot be used for the intended purpose and therefore reduces the

yield of useful recombinant polypeptide of interest. Additionally, for many applications the clipped polypeptide has to be removed during purification. This may require the specific development of purifications methods that allow separating the clipped from the intact polypeptide, which can be labor and cost intensive and sometimes are even not successful so that clipped protein remains as impurity. Therefore, proteolytic degradation of the expressed polypeptide of interest in the cell culture medium is a major issue when recombinantly producing a polypeptide of interest in vertebrate cells.

[4] Several studies have investigated the phenomenon of clipping in order to provide solutions to this problem. Clipping was found to be time and temperature dependent and could be inactivated by heat treatment. However, so far little is known about the proteases expressed by the vertebrate host cells used for recombinant expression, and about the proteases responsible for clipping. One of the major challenges is the number of proteases expressed by vertebrate host cells. E.g. more than 700 proteases are known to be present in rodent cell genomes, many of which could be involved in clipping.

[5] Several approaches were suggested in the prior art to overcome the problems of clipping. One approach to avoid clipping is the reengineering of the affected protein in order to eliminate amino acid motifs that are prone to clipping. However, this approach is time consuming, needs extensive testing whether the chosen approach was successful and does not guarantee that the reengineered protein is still functional, does not show clipping at other sites and is not immunogenic. A different way to control protease activity in the cell culture is to add inhibitors or substitute substrates into the culture medium. However, in large-scale manufacturing processes such additives are expensive and have to be removed during purification which may require specific development of purification methods and sensitive assays to monitor that no additives remain. Other ways to reduce proteolytic degradation of the expressed polypeptide of interest in the cell culture medium include timing of the harvest (early harvest) to limit the exposure of the polypeptides of interest to proteases in the cell culture medium, reducing the temperature of the culture, optimizing the pH conditions or improving the cell viability to reduce the release of intracellular proteases into the cell culture medium. Furthermore, it was found that the choice of the vertebrate cell line used as production host cell may have an influence on clipping depending on the proteases that are expressed by the respective cell line and the individual polypeptide of interest to be expressed. Therefore, one option is to screen different cell lines to identify a cell line wherein clipping of the polypeptide of interest does not occur or occurs to a lesser extent, which, however, is time-consuming and may require the adaptation of the production process to the chosen cell line. Even though these different options can reduce clipping, they will rarely completely eliminate the proteolytic degradation. Furthermore, in many cases adaptations of the expression and production process to the individual polypeptide of interest to be expressed is required in order to develop a production method that avoids or at least reduces the problems of clipping and provides the intact polypeptide of interest with sufficient yield and purity. Such adaptations are, however, time consuming and costly. Therefore, improved ways are needed to address the problem of clipping.

[6] Therefore, is an object of the invention to provide an improved method for the recombinant production of a polypeptide of interest in vertebrate cells wherein clipping of the expressed and secreted polypeptide of interest in the cell culture medium is reduced or eliminated. In particular, it is an object of the present invention to provide novel vertebrate cells, wherein clipping of a polypeptide of interest expressed and secreted by said cells is reduced or eliminated.

SUMMARY OF THE DISCLOSURE

[7] The present disclosure is *inter alia* based on the unexpected finding that altering a vertebrate cell to impair the effect of the endogenous protease matriptase, e.g. by reducing or eliminating the functional expression of the matriptase gene, significantly decreases the proteolytic degradation ("clipping") of a recombinant polypeptide of interest that is expressed and secreted by said cell into the cell culture medium. Thus, impairing the effect of matriptase in said cell reduces clipping of the secreted recombinant polypeptide of interest compared to a corresponding vertebrate cell in which the effect of matriptase is not impaired. With matriptase, a key protease responsible for clipping of recombinantly expressed and secreted polypeptides was identified. Altering the vertebrate cell to impair the effect of matriptase allows to significantly improve the recombinant production of a polypeptide of interest by reducing or eliminating clipping of the recombinantly expressed and secreted polypeptide of interest in the cell culture medium. Thereby, the yield of intact polypeptide of interest is increased. As it is demonstrated by the examples, these advantageous effects are not seen when impairing the function of other proteases which confirms that matriptase is the major protease responsible for clipping. The novel vertebrate cells provided by the present disclosure obviate the need to reengineer the polypeptide of interest to be expressed in order to eliminate proteolytic sites or to laboriously adapt the production process to reduce or prevent clipping or to design specific purification processes to remove clipped protein. Advantageously, these altered vertebrate cells described herein can be used as universal host cells for expressing different polypeptides of interest, including polypeptides of interest that are particularly prone to clipping. Specific adaptations of the polypeptide to be expressed or the expression system become obsolete, which saves time and costs. Furthermore, as is demonstrated by the examples, the altered vertebrate cell provided by the present disclosure show good growth and expression characteristics. Therefore, impairing the effect of matriptase still provides host cells with favorable expression characteristics that are important for the recombinant expression of a polypeptide of interest. Hence, the present invention makes an important contribution to the prior art.

[8] According to a first aspect, the present disclosure provides an isolated vertebrate cell suitable for recombinant expression of a polypeptide of interest, wherein the vertebrate cell is altered to impair the effect of matriptase and comprises at least one heterologous polynucleotide encoding a polypeptide of interest, wherein the vertebrate cell secretes the polypeptide of interest. The vertebrate cell is altered to impair the effect of matriptase e.g. by reducing or eliminating functional expression of the matriptase gene in said vertebrate cell, e.g. by gene silencing, gene deletion or by mutating the matriptase gene so that a non- or

less functional protein is expressed. Impairing the effect of matriptase in the vertebrate cell reduces clipping of the secreted recombinant polypeptide of interest compared to a corresponding vertebrate cell in which the effect of matriptase is not impaired. As is shown by the examples *inter alia* based on spike-in experiments, in the supernatant of the cell culture medium that is obtained when culturing respectively altered vertebrate cells in which the effect of matriptase is impaired, clipping of recombinant polypeptides of interest is surprisingly highly reduced even though other proteases are still active in the supernatant. These results were confirmed by expressing several polypeptides of interest in respectively altered cells wherein the effect of matriptase is impaired. These results confirm that matriptase is a key protease responsible for clipping of a recombinantly expressed and secreted polypeptide of interest. Thus, the vertebrate cells according to the first aspect are particularly suitable as host cells for recombinant production technologies and can be used for recombinant production of a polypeptide of interest that is secreted by the vertebrate cell into the cell culture medium from which it can then be harvested.

[9] According to a second aspect, the present disclosure provides a method for producing a vertebrate cell according to the first aspect, comprising altering a vertebrate cell to impair the effect of matriptase and introducing a polynucleotide encoding a polypeptide of interest to be expressed, wherein said polypeptide of interest is secreted by the vertebrate cell. Impairment of the effect can be achieved e.g. by reducing or eliminating functional expression of the matriptase gene in said cell, e.g. by gene silencing, gene deletion or by mutating the matriptase gene so that a non- or less functional protein is expressed.

[10] According to a third aspect, a method is provided for recombinantly producing a polypeptide of interest, comprising utilizing a vertebrate cell according to the first aspect as host cell for recombinant expression of the polypeptide of interest. As described above, due to the achieved reduced level of polypeptide clipping in the cell culture medium, these altered vertebrate cells are particularly suitable as host cells for recombinant production of a polypeptide of interest, in particular of polypeptides that are prone to clipping such as glycopolypeptides. As preferred embodiment, a method for recombinantly producing a polypeptide of interest is provided, comprising

- (a) culturing vertebrate host cells according to the first aspect under conditions that allow for the expression and secretion of the polypeptide of interest into the cell culture medium;
- (b) isolating the polypeptide of interest from the cell culture medium; and
- (c) optionally processing the isolated polypeptide of interest.

[11] According to a fourth aspect, a method for recombinantly producing a polypeptide of interest is provided, comprising

- (a) culturing vertebrate host cells which comprise at least one heterologous polynucleotide encoding a polypeptide of interest under conditions that allow for the expression and secretion of the polypeptide of interest into the cell culture medium, wherein the cell culture medium comprises a protease inhibitor that is selective for matriptase;

- (b) isolating the polypeptide of interest from the cell culture medium; and
- (c) optionally processing the isolated polypeptide of interest.

5 [12] According to a fifth aspect, a method is provided for selecting a host cell which recombinantly expresses a polypeptide of interest, comprising

- (a) providing vertebrate cells according to the first aspect as host cells; and
- (b) selecting one or more host cells expressing the polypeptide of interest.

10 [13] According to a sixth aspect, the present disclosure pertains to the use of a vertebrate cell for recombinant production of a polypeptide of interest that is secreted from the vertebrate cell, wherein the used cell is altered to impair the effect of the endogenous protease matriptase. Respectively altered cells can be e.g. transfected with a polynucleotide encoding a polypeptide of interest that is supposed to be expressed and secreted by said
15 cell. When using respective vertebrate cells wherein the effect of the endogenous protease matriptase is impaired, no or reduced amounts of functional matriptase are active in the cell culture medium. This significantly reduces or even eliminates proteolytic degradation of the recombinant polypeptide of interest that is secreted by the cells into the cell culture medium. Therefore, it is advantageous to use such altered vertebrate cell for recombinant protein
20 expression.

[14] According to a seventh aspect, the present disclosure pertains to a method for selecting a vertebrate cell for recombinant production of a polypeptide of interest, comprising analyzing if the endogenous protease matriptase is functionally expressed in the vertebrate cell and
25 selecting a vertebrate cell in which the effect of such endogenous matriptase is impaired for recombinant production of the polypeptide of interest. This selection process allows identifying vertebrate cells that are capable of producing a recombinant polypeptide of interest, wherein clipping of the polypeptide of interest in the cell culture medium is reduced. Respective vertebrate cells are particularly suitable for recombinant protein production.

30 [15] Other objects, features, advantages and aspects of the present application will become apparent to those skilled in the art from the following description and appended claims. It should be understood, however, that the following description, appended claims, and specific examples, while indicating preferred embodiments of the application, are given by way of illustration only. Various changes and modifications within the spirit and scope of the
35 disclosed invention will become readily apparent to those skilled in the art from reading the following.

- [15a] In an embodiment, there is provided a method for selecting a vertebrate cell to be used for recombinant production of a secreted polypeptide of interest, comprising: analyzing if the endogenous protease matriptase is functionally expressed in the vertebrate cell; and selecting a vertebrate cell in which the effect
- 5 of such endogenous matriptase is impaired by reduction or elimination of functional expression of the matriptase gene by gene knock-out, gene mutation, gene deletion, gene silencing or a combination of any of the foregoing; wherein impairment of such endogenous matriptase is indicative that the vertebrate cell is suitable for use in recombinant production of the secreted polypeptide of interest.
- 10 [15b] In an embodiment, there is provided the method as described herein, wherein the selected cell is a vertebrate cell as described herein.

BRIEF DESCRIPTION OF THE FIGURES

[16] **Fig. 1** shows a Western Blot for two therapeutic proteins (upper panel: IgG (mAb), lower panel: Fc-fusion protein) incubated for several days in conditioned medium obtained by cultivating CHO cells transfected with siRNAs directed against different protease target genes (7 days incubation for mAb, 3 days incubation for the Fc-fusion protein). The first control "(+)" represents a sample of the polypeptide of interest incubated for the same time in a chemically defined culture medium which was not in contact with cells. The chemically defined culture medium used as control was the same culture medium as the culture medium in which the cells were cultured to obtain the conditioned medium. The second control "(-)" represents a sample of the polypeptide of interest incubated in the conditioned medium obtained from CHO cells transfected with a non-effective siRNA (125pmol) which served as siRNA negative control. The names of the target trypsin-like serine proteases and homologs whose expression were silenced by siRNA transfection are given above the upper panel (MT-SP1 (matriptase, also referred to herein as St14), C1r (also referred to as C1ra), C1s (also referred to as Gm5077), Plat, and Prss35). For each of the combinations "polypeptide of interest/conditioned medium of cells with silenced protease expression" results of two experimental set ups with different siRNA concentrations are shown. Also determined and indicated in Fig. 1 below in parenthesis is the percentage of residual protease gene-expression in relation to the protease gene-expression in the siRNA negative control cells which was set as 100%. This means e.g. with respect to MT-SP1 (18.6%) that there was 18.6% residual gene-expression upon silencing. MT-SP1: 125pmol (18.6%) and 150pmol (18.2%), C1r (C1ra): 125pmol (6.3%) and 150pmol (6.7%), C1s (Gm5077): 100pmol (11.9%) and 125pmol (14.4%), Plat: 125pmol (8.3%) and 150pmol (5.1%) and Prss35: 100pmol (18.4%) and 125pmol (14.8%). The results demonstrate that when the MT-SP1 mRNA level is down-regulated by RNA interference, significantly less clipping occurred in the conditioned medium (detected for mAb and Fc-fusion protein). When other expressed trypsin-like serine proteases were down-regulated, clipping remains the same. The results thus demonstrate that matriptase is the key protease responsible for clipping.

[17] **Fig. 2** shows the sequence of exon 2 of the matriptase gene plus flanking intron regions in CHO-K1 derived cells. Sequencing primers (cursive, bold and underlined) were designed to target flanking introns to confirm CHO-K1-derived MT-SP1 exon 2 sequence (shaded in grey). Binding sites of DNA binding domains of TALENs used to obtain matriptase knock-out cells are depicted in bold and underlined and the target region for knock-out mutation is shown in bold and double-underlined.

[18] **Fig. 3** shows a column diagram representing the relative matriptase mRNA expression measured in the CHO-K1 derived matriptase knock-out cell clones KO-1 to KO-9 and clone $\Delta 7/\Delta 15$ and a CHO-K1 derived wildtype cell line (WT). The matriptase mRNA expression in the wildtype CHO-K1 derived cell line was defined as 100%. As can be seen, matriptase mRNA expression was reduced in all knock-out clones and clone $\Delta 7/\Delta 15$.

[19] **Fig. 4** shows a Western Blot of three different polypeptides of interest that are prone to clipping incubated in the cell culture medium supernatant of wild-type CHO-K1 cells (WT) and a variety of different CHO-K1 derived cell clones in which matriptase was knocked out by different mutations (see Example 2, in particular Table 5). The polypeptides of interest (concentration 0.7 μ M each) were incubated in conditioned medium obtained from wildtype CHO-K1 cells (WT), the knock-out clones KO-1 to KO-9 shown in Table 5 (indicated in Fig. 4 by the numbers 1 to 9, respectively) or a chemically defined medium that was not in contact with cells (+). The polypeptide of interest in Fig. 4A is an IgG (mAb) and the incubation time was 48h. The polypeptide of interest in Fig. 4B is a Fc-fusion protein, the incubation time was 24h. The polypeptide of interest in Fig. 4C is a further recombinant protein with two glycovariants, the incubation time was 1h. Both glycovariants were clipped in conditioned medium obtained from wildtype CHO-K1 cells (WT). The intact and the clipped proteins are indicated by the arrows. Fig. 4D shows the result of a repetition of the experiment with the mAb using conditioned medium obtained from respective cells that had been cultivated for three month. The incubation conditions were identical as in Fig 4A. The results demonstrate that clipping is efficiently prevented in the conditioned medium obtained from the matriptase knock-out cell clones KO-1 to KO-9 and that these advantageous results are maintained during prolonged culturing.

[20] **Fig. 5A and B** show Western Blots of two different viral glycoproteins (polypeptides of interest) that are prone to clipping. The polypeptides of interest were incubated in conditioned medium obtained from a CHO-K1 derived cell line which expresses matriptase (WT) and different cell clones in which matriptase was knocked out (KO-1 to KO-3). The polypeptides of interest were incubated in a concentration of 0.7 μ M in conditioned medium obtained from wildtype cells (WT), conditioned medium from the knock-out clones KO-1 to KO-3 shown in Table 5 (indicated by the numbers 1 to 3, respectively) or a chemically defined culture medium that was not in contact with cells (+). D1 indicates an incubation time of 24h and D2 an incubation time of 48h. Intact and clipped proteins are indicated by arrows (white and black arrows). The numbers on the left side of the Western Blot in **Fig. 5B** represent the molecular weight in kDa as determined. The results again demonstrate that clipping is significantly reduced in the conditioned medium obtained from matriptase KO clones.

[21] **Fig. 6** shows a Western Blot of a Fc-fusion protein (**Fig. 6A**) and a IgG (mAb) (**Fig. 6B**) incubated in the cell culture medium supernatant obtained from CHO cells expressing wildtype matriptase ("WT1" and "WT2") and a CHO clone in which the matriptase was mutated (" Δ 7/ Δ 15") so that functional expression of matriptase was reduced. A corresponding chemically defined culture medium ("(+)" that was not in contact with the cells served as positive control. The Fc-fusion protein and the mAb (concentration of 0.7 μ M each) were incubated for 2h and 24h, respectively. The numbers at the side of the Western Blot represent the approx. molecular weight in kDa as determined on the gel. The results demonstrate that clipping is reduced in the conditioned medium obtained from the mutant cell clone Δ 7/ Δ 15 in which functional expression of active matriptase is reduced.

[22] **Fig. 7** shows the results of a coincubation of different polypeptides of interest with two different trypsin-like proteases, namely matriptase from mouse (MT-SP1) and human Htra1. **Fig. 7A** shows a Western Blot analysis of mAb (incubation time 24h), **Fig. 7B** a capillary gel electrophoresis analysis (Caliper LabChip®) of an Fc-fusion protein (incubation time 2h) and **Fig. 7C** a Western Blot analysis of a further recombinant protein (incubation time 1h). The concentration of the polypeptide of interest was 0.7µM in all experimental set ups. Each polypeptide of interest was tested with decreasing amounts of the MT-SP1 and Htra1: Molar ratios of protease to polypeptide of interest from left to right are 1/10, 1/100 and 1/1000 for MT-SP1 and 1/3, 1/10 and 1/100 for Htra1. The polypeptides of interest were also incubated for the same time in a chemically defined culture medium "(+)" and in conditioned medium (supernatant) obtained from CHO-K1 derived wildtype cells "(-)". The intact proteins (larger) and the clipped proteins (smaller) are indicated by the arrows in Fig. 7A and 7C. In Fig. 7B, there are two types of clipped polypeptides beside the intact polypeptide (all marked with an arrow and illustrated at the side). Fig. 7A to Fig. 7C demonstrates that clipping occurs in the presence of matriptase, thereby confirming that matriptase is a key protease responsible for clipping.

[23] **Fig. 8** shows a Western Blot analysis of two polypeptides of interest, a monoclonal IgG-antibody (upper panel, incubation time 24h) and a recombinant protein (lower panel, incubation time 1h). The polypeptides of interest (concentration 0.7µM each) were incubated in either a chemically defined culture medium "(+)", in conditioned medium obtained from CHO-K1 derived wildtype cells "(-)", or in a chemically defined culture medium with recombinant matriptase from Mus Musculus added ((+) + MT-SP1). To several samples an anti-MT-SP1 inhibitory Fab fragment was added as indicated (" + Fab"). The concentration of the Fab in the sample is shown above the lane (1µM, 10µM or 50µM). Arrows indicate protein bands representing the intact or the clipped protein. The protein band at approx. 25kD in the lanes wherein the Fab fragment was added represent the Fab fragment. The results demonstrate that addition of the anti-MT-SP1 Fab fragment (which is a selective matriptase inhibitor) to a cell culture medium in which matriptase is active reduces or even completely prevents clipping. The results further confirm that matriptase is the key protease responsible for clipping.

[24] **Fig. 9** shows the result of a microchip capillary electrophoresis (Caliper LabChip®). A recombinant mAb was expressed in CHO-K1 derived wild-type cells (WT) or in matriptase knock-out cells (KO-4, see also Table 5) and was purified using affinity chromatography (protein A). The numbers above the lanes present different mAb production cell pools obtained from parallel transfection and selection processes. In each lane three protein bands are visible as indicated by arrows. The upper protein band represents the intact mAb heavy chain, the second band (directly below the first) represents the clipped antibody heavy chain and the band at the bottom represents the light chain of the mAb. The results show that clipping is significantly reduced when expressing a polypeptide of interest in matriptase knock-out cells.

DETAILED DESCRIPTION

[25] The present disclosure is *inter alia* based on the unexpected finding that an altered vertebrate cell, in which the effect of the endogenous protease matriptase is impaired, e.g. by reducing or eliminating functional expression of the matriptase gene, is capable of expressing and secreting a recombinant polypeptide of interest into the cell culture medium, wherein, however, clipping of the recombinant polypeptide of interest in the cell culture medium is significantly reduced. Thus, the major protease responsible for clipping of recombinantly expressed and secreted polypeptides was identified among hundreds of different proteases. It was also highly surprising that targeting and impairing the activity of a single protease is sufficient to significantly reduce or even eliminate clipping in the cell culture medium. Using e.g. respectively altered vertebrate cells according to the invention allows to significantly improve the recombinant production of a polypeptide of interest by increasing the yield of intact polypeptide of interest that can be obtained after expression and secretion from the cell culture medium. Therefore, when using the altered vertebrate cell according to the invention for recombinantly producing a polypeptide of interest, it is not necessary to perform additional measures to reduce proteolytic degradation and hence prevent clipping of the polypeptide of interest. Therefore, the present invention makes an important contribution to the prior art.

[26] The individual aspects and suitable and preferred embodiments of the present disclosure will now be described in detail.

A. Altered vertebrate cells

[27] According to a first aspect, the present disclosure provides an isolated vertebrate cell suitable for recombinant expression of a polypeptide of interest, wherein the vertebrate cell is altered to impair the effect of matriptase and comprises at least one heterologous polynucleotide encoding a polypeptide of interest, wherein the vertebrate cell secretes the polypeptide of interest. Impairing the effect of matriptase in the vertebrate cell reduces clipping of the secreted recombinant polypeptide of interest compared to a corresponding vertebrate cell in which the effect of matriptase is not impaired.

[28] Matriptase, was first described in 1993 as a new gelatinolytic activity in cultured breast cancer cells. Matriptase belongs to the family of type II transmembrane serine proteases (TTSPs). Orthologs of matriptase are present in different vertebrate species, including mammalian species, and were identified for example in human, chimpanzee, dog, mouse, rat, chicken, zebrafish, spotted green pufferfish and tiger pufferfish which suggests a conserved evolutionary function. Matriptase is listed in the IUBMB Enzyme nomenclature as EC 3.4.21.109. Matriptase is also known as membrane-type serine protease 1 (MT-SP1) and suppressor of tumorigenicity-14 (ST14) (see Chen et al, The Transmembrane Serine Protease Matriptase: Implications for Cellular Biology and Human Diseases J Med Sci 2012; 32 (3): 097-108). It is an integral membrane protein with a single-span transmembrane domain close to the cytoplasmatic N-Terminus. The extracellular part consists of a stem region (including a single SEA, 2 CUB and 4 LDLRA domains) and the C-terminal serine

protease domain that is structurally highly similar to other TTSPs and includes a conserved histidine/aspartic acid/serine (HDS) catalytic triad essential for catalytic activity (see e.g. List et al, Matriptase: Potent Proteolysis on the cell Surface; MOL MED 12 (1-3) 1-7, JANUARY-MARCH 2006 and Chen et al, The Transmembrane Serine Protease Matriptase: Implications for Cellular Biology and Human Diseases J Med Sci 2012; 32 (3): 097-108). Matriptase is described as being expressed in the epithelia in many organ systems such as skin, breast, lung, epidermis, cornea, salivary gland, oral and nasal cavities, thyroid, thymus, esophagus, trachea, bronchioles, alveoli, stomach, pancreas, gallbladder, duodenum, small intestine, colon, rectum, kidney, adrenals, urinary bladder, ureter, seminal vesicles, epididymis, prostate, ovaries, uterus and vagina (see List et al, 2006 and Chen et al, 2012). Matriptase is synthesized as an inactive zymogen and is converted to its active form via a complicated process. Details regarding the activation process which involves endoproteolytic cleavages are described for the human matriptase in List et al 2006 and Chen et al 2012. Matriptase is bound to the membrane as type II transmembrane protein with the catalytic domain orientated into the extracellular space. Furthermore, it is described in the literature that a significant shedding of matriptase, respectively its extracellular part, occurs *in vivo* (see List et al, 2006 and Chen et al 2012). It is described in the literature that matriptase is shed in form of a complex, e.g. complexed to the Kunitz-type serine protease inhibitor HAI-1. Different studies suggest that in human cells the specific inhibitor HAI-1 facilitates the transport of the matriptase to the cell membrane as it was shown that removal or even single point mutations in HAI-1 lead to an accumulation of the matriptase in the Golgi compartment. In the literature, several different endogenous inhibitors of matriptase besides HAI-1 have been described such as HAI-2, antithrombin, alpha-1 antitrypsin and alpha -2-antiplasmin. Furthermore, also other inhibitors of matriptase have been described (see e.g. Chen et al, 2012). It is described in the literature that matriptase may play numerous roles in normal physiology such as skin barrier function, epithelial integrity, hair follicle development, and thymus homeostasis, and in human pathologies, such as osteoarthritis, atherosclerosis, and tumor progression, invasion, and metastasis.

[29] Against this scientific background which is unrelated to the recombinant production of a polypeptide of interest, the present finding that matriptase is a key protease responsible for clipping of recombinantly produced polypeptides of interest that are secreted by the host cells into the cell culture medium was highly surprising. Considering the large number and variety of proteases expressed in vertebrate cells, such as in particular mammalian cells, it was even more surprising that impairing the function of this single protease – matriptase – is sufficient to significantly reduce or even eliminate clipping of the secreted polypeptide of interest in the cell culture medium. These advantageous effects are not seen with other, even closely related proteases what supports the importance of the finding that matriptase is the key enzyme responsible for clipping of secreted recombinant polypeptides in the cell culture medium. As is shown by the examples, when using altered vertebrate cells, wherein the effect of the endogenous protease matriptase is impaired, clipping of a recombinant polypeptide of interest that is secreted by the cell into the cell culture medium is significantly reduced. E.g., impairing the function of the matriptase either by gene silencing (e.g. RNAi as shown in Example 1) or mutation of the endogenous matriptase gene (e.g. knock-out of one

or both of the two matriptase alleles as shown in Example 2 and 5) leads to a significant reduction or even complete elimination of clipping of the recombinantly expressed and secreted polypeptide of interest in the cell culture medium. A significant reduction of clipping was found for all tested polypeptides that are prone to clipping, such as IgG antibodies, Fc-fusion proteins, glycosylated viral proteins and other therapeutically active proteins. Therefore, using a matriptase deficient cell line for producing a polypeptide of interest by recombinant expression is advantageous. It is further apparent from the examples that matriptase directly cleaves the polypeptide of interest that is secreted into the cell culture medium. Thus, any impairment of the effect of the matriptase reduces clipping of the recombinantly expressed polypeptide that is secreted into the cell culture medium. Furthermore, as is demonstrated by the examples, adding a selective matriptase inhibitor to the cell culture medium may also reduce clipping.

[30] Because the effect of the endogenous protease matriptase is impaired in the vertebrate cell according to the first aspect of the present disclosure, no or less functionally active matriptase is present in the cell culture medium in which said cells are cultured, e.g. because such cells present no or reduced amounts of functional matriptase on the cell surface and/or release (e.g. due to shedding) no or reduced amounts of functional matriptase into the cell culture medium. Thereby, the proteolytic degradation of the recombinant polypeptide of interest that is secreted into the cell culture medium is significantly reduced or even eliminated what is an important advantage when recombinantly producing a polypeptide of interest. Because clipping is significantly reduced, the yield of intact polypeptide of interest is increased. Less or even no non-functional and potentially immunogenic clipped by-product is produced. In addition, it was found that these novel vertebrate host cells generally show good expression yields and have good growth characteristics, which makes them particularly suitable as production cell lines. Further advantages are described in the following and are also apparent from the examples. Thus, these advantageous vertebrate cells allow to recombinantly produce a polypeptide of interest with improved product quality and yield. Furthermore, using the vertebrate cells according to the invention, which preferably are mammalian cells, reduces the time required for developing a production cell line for recombinant production of a polypeptide of interest. Less or even no optimization of the polypeptide of interest is necessary in order to avoid or reduce clipping e.g. by changing the amino acid sequence. In particular, removal of clipping sites becomes obsolete. Furthermore, time consuming purification processes for removing clipped protein can be avoided when using these cells as production cell line. Thus, these vertebrate cells have important advantages when being used as host cell lines for recombinant production technologies.

[31] The sequence listing shows exemplary amino acid sequences of matriptase of different vertebrate species such as hamster (SEQ ID NO: 1 – NCBI reference sequence: XP_003495890), human (SEQ ID NO: 2 – NCBI reference sequence: NP_068813), mouse (SEQ ID NO: 3 – NCBI Reference sequence: NP_035306), rat (SEQ ID NO: 4 – NCBI reference sequence: NP_446087) and chimpanzee (SEQ ID NO: 5 – NCBI reference sequence: NP_001189434). As is evident from Table 1, matriptase is currently also referred to as “suppressor of tumorigenicity 14 protein” (e.g. for human) and “suppressor of

tumorigenicity 14 protein homolog" (e.g. in mouse and Chinese hamster). The term "matriptase" as used herein in particular encompasses any protein that shares at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to one or more of the proteins shown in SEQ ID NO: 1 to 5 as matriptase reference protein and which has the same proteolytic activity as said matriptase reference protein. Identity as used herein is calculated over the entire length of the reference protein. Matriptase is distinguished from other proteases. In the literature, matriptase has been assigned a variety of names, a selection of which is provided in Table 1. The corresponding genes encoding the enzyme matriptase have also been assigned a variety of names, a selection of which is provided in Table 1. In the context of the invention, the protease is referred to as "matriptase", "MT-SP1", "suppressor of tumorigenicity 14 protein" or "suppressor of tumorigenicity 14 protein homolog" for the ease of simplicity. However, the term "matriptase" also refers to and encompasses any alternative names of said protein or the corresponding gene e.g. used to characterize the corresponding protein or gene in different species. Homologs and orthologs of matriptase which have the same function are included in the term "matriptase". In this context it is mentioned that matriptase-2 and matriptase-3 relate to proteases that are distinct from matriptase (although structurally related) and thus, are not covered by the term "matriptase" as used herein.

Table 1: Exemplary alternative names of matriptase gene and/or the encoded protein product matriptase used in the literature (alphabetical order)

Breast cancer 80 kDa protease
CAP3
Channel-activating protease 3
EC 3.4.21.109
Epithin
HAI
Matriptase
Matriptase-1
Membrane-type serine protease 1
MT-SP1; MTSP1
Prostamin
PRSS14 g.p. (Homo sapiens), PRSS14
Serine endopeptidase SNC19
Serine protease 14
Serine protease TADG-15; TADG-15; TADG15
SNC19
ST14 (official gene name in human according to HGNC)
St14 (official gene name in mouse according to MGI)
Suppression of tumorigenicity-14 Protein; suppression of tumorigenicity 14 (colon carcinoma, matriptase, epithin)
Suppressor of tumorigenicity 14 protein

Suppressor of tumorigenicity 14 protein homolog (mouse)
TMPRSS14
Tumor associated differentially expressed gene 15 protein

[32] A gene encoding a matriptase protein is also referred to as "matriptase gene" herein. The genomic gene sequence of different mammalian species is known, and is e.g. described in Chinese hamster (NCBI Gene-ID: 100755225); Homo sapiens (NCBI Gene-ID: 6768); Mus musculus (NCBI Gene-ID: 19143); Rattus norvegicus (NCBI Gene-ID: 114093); Pan Troglodytes (NCBI Gene-ID: 100188950) and others. Synonyms for the matriptase gene are listed in Table 1, commonly used is "ST14" or "St14". The term "matriptase gene" as used herein in particular encompasses any endogenous gene of a vertebrate cell which encodes a matriptase protein as shown in SEQ ID NO: 1 to 5 or which encodes a matriptase protein that shares at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to one or more of the matriptase proteins shown in SEQ ID NO: 1 to 5 as matriptase reference protein and which has the same proteolytic activity as said matriptase reference protein. Identity as used herein is calculated over the entire length of the reference protein.

[33] The present disclosure *inter alia* pertains to a modified vertebrate cell, such as preferably a mammalian cell, wherein the effect of the endogenous matriptase, which usually is endogenously expressed by a corresponding unmodified vertebrate cell, is impaired in said cell. Because of this alteration which impairs the effect of the endogenous protease matriptase, the altered vertebrate cell described herein presents or releases no or reduced amounts of functional matriptase into the cell culture medium, whereby clipping of the secreted polypeptide of interest in the cell culture medium is reduced or even completely prevented. As is demonstrated by the examples, this alteration, which can be transient or permanent, significantly reduces proteolytic degradation of a recombinant polypeptide of interest that is secreted by said altered cell into the cell culture medium.

[34] There are several possibilities to alter and hence modify a vertebrate cell to impair the effect of the endogenous matriptase in said cell. The effect of the matriptase may be impaired e.g. on the gene or on protein level. The effect of matriptase can be impaired, for example, by modification of the structure/sequence of matriptase, the transcription, translation and/or cellular trafficking of matriptase. Non-limiting options are described in the following.

[35] According to one embodiment, the effect of matriptase is impaired because functional expression of the matriptase gene is reduced or eliminated in said cell. As is shown by the examples, altering the expression of the matriptase gene by reducing or eliminating the functional expression of the matriptase gene, e.g. by gene silencing or by gene knock-out, is a very efficient measure to provide a vertebrate cell that is suitable for expressing a recombinant polypeptide of interest wherein, however, clipping of the secreted polypeptide of interest in the cell culture medium is reduced or even completely avoided. It was found that when the functional expression of the matriptase gene is reduced or eliminated in a

vertebrate cell, clipping of a secreted recombinant polypeptide of interest in the cell culture medium is likewise decreased. This correlation is an unexpected finding.

5 [36] Reduction or elimination of functional expression of the matriptase may be achieved by various means. Functional expression can be reduced for example by reducing the expression level of the matriptase or by reducing the catalytic activity of the matriptase or by a combination of both. According to one embodiment, the cell is altered so that the functional expression of the matriptase gene is reduced or eliminated by gene knock-out, gene mutation, gene deletion, gene silencing or a combination of any of the foregoing. According to one embodiment, the genome of the vertebrate cell is altered to impair the effect of matriptase.

15 [37] According to one embodiment, functional expression of the matriptase gene is reduced or eliminated in the cell by gene knockout. A gene knockout is a genetic technique by which a gene is made inoperative by disrupting its function. E.g. a nucleic acid can be inserted into the coding sequence, thereby disrupting the gene function. Furthermore, the complete matriptase gene or a portion thereof can be deleted, whereby no or no functional protein is expressed by the respectively altered cell. Another option is to introduce one or more knock-out mutations into the coding sequence, which renders a non- or a less functional expression product. E.g. one or more frameshift mutations can be introduced that result in a non- or less-functional protein. Alternatively or additionally, one or more stop codons can be introduced into the coding sequence so that a truncated, non- or less functional protein is obtained. Hence, according to one embodiment, the matriptase gene comprises one or more mutations which provide a non- or less functional expression product. According to one embodiment, said one or more mutations are frame-shift or stop codon mutations. According to one embodiment, all or part of the protease domain located at the C-terminus of matriptase is not present due to the introduced one or more mutations. Other options include but are not limited to introducing one or more mutations in the promoter, in the 5'UTR, the 3' UTR and/or other regulatory elements. According to one embodiment, the promoter function of the matriptase gene is disrupted, e.g. by introducing a promoter deletion or by introducing a construct between the promoter and the transcription start. Methods for achieving a gene knockout to suppress or eliminate functional expression of a target gene are also well-known to the skilled person and thus, do not need any detailed description herein. Some non-limiting examples are nevertheless described below.

35 [38] According to one embodiment, the matriptase gene is functionally knocked out by genetic engineering. Examples include but are not limited to genome editing, such as genome editing with engineered nucleases (GEEN). This is a type of genetic engineering in which DNA is inserted, replaced or removed from a genome using artificially engineered nucleases, or "molecular scissors." The nucleases create specific double-stranded breaks (DSBs) at desired locations in the genome, and harness the cell's endogenous mechanisms to repair the induced break by natural processes of homologous recombination (HR) and nonhomologous end-joining (NHEJ). There are at least four families of engineered nucleases that can be used for this purpose: Zinc finger nucleases (ZFNs), Transcription Activator-Like

40

Effector Nucleases (TALENs), a nuclease recognizing Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR), and engineered meganuclease re-engineered homing endonucleases. TALEN technology was also used in the examples to provide altered mammalian cells wherein the matriptase gene was fully or partially knocked out, thereby impairing the effect of matriptase in said cells.

[39] According to one embodiment, one or more copies of the matriptase gene present in the genome of the vertebrate cell are altered, e.g. knocked-out or deleted, to reduce or eliminate and hence impair the effect of matriptase in the vertebrate cell. Thus, according to one embodiment, at least one copy of the matriptase gene is deleted or functionally inactivated in the genome of the vertebrate cell. According to one embodiment, the vertebrate cell comprises one or more mutations in at least one copy of the matriptase gene to provide a non- or less-functional expression product. In general, a vertebrate cell comprises two allelic copies of the matriptase gene. One or more mutations may be inserted into one or both copies of the matriptase gene. Preferably, one or more mutations are inserted into both, respectively all copies of the matriptase gene to provide a non- or less-functional expression product and, hence to impair the effect of matriptase in the vertebrate cell. Thereby, all copies of matriptase gene are basically impaired or inactivated in the genome. As is shown in the examples, different mutations can be introduced into the different alleles of the matriptase gene to achieve impairment.

[40] According to one embodiment, said one or more mutations are comprised in a coding region of the matriptase gene and result in a non- or less functional expression product. E.g. the function of matriptase may be impaired because the altered vertebrate cell comprises one or more mutations in a region of the matriptase gene that encodes an amino acid sequence present in more than one catalytically active and hence functional splice variant of matriptase. Several splice variants of matriptase have been identified in different species. Several of the matriptase exons are found in the majority or even all identified functional splicing variants. It is advantageous to introduce one or more mutations into a region of the matriptase gene which encodes an exon that is present in more than one, preferably in the majority of, or even all functional matriptase splicing variants of the respective vertebrate cell.

[41] According to one embodiment, one or more mutations are comprised in a polynucleotide sequence that encodes exon 2 of matriptase. Choosing exon 2 as target for alteration to impair the function of the endogenous protease matriptase in the altered vertebrate cell has the advantage that this approach covers several different functional splicing variants. Exons close to the N-terminus of matriptase such as e.g. exon 1, exon 2 and exon 3 are advantageous targets for introducing one or more mutations, in particular one or more frameshift mutations. A frameshift mutation in one of these exons most likely leads to a stop codon early in the sequence. The truncated protein encoded by the mutated gene is most likely short and assumed to be located intracellular and/or be non-active. An inactive truncated protein is advantageous as it can be assumed that its expression is not toxic for the cells. However, one or more mutations may also be introduced in one of the subsequent exons e.g. selected from exons 4 to 19.

[42] In the examples, CHO cells comprising different frameshift mutations in exon 2 were generated. The examples show that respectively altered cells sustained cell growth. Therefore, the encoded truncated version(s) of matriptase tested in the examples were apparently not toxic for the cells. Furthermore, it was found that the overall mRNA expression of the respectively mutated matriptase gene was lower, so that lower levels of the truncated version were obtained (see Fig. 3). Hence, the examples demonstrate that a respective alteration is efficient in order to impair the function of matriptase in a vertebrate cell such as preferably a mammalian cell while maintaining other characteristics that are important for the recombinant production of a polypeptide of interest. Thus, according to one embodiment, the isolated vertebrate cell comprises at least one frameshift mutation in exon 2 of the matriptase gene which results in that a non- or less functional truncated polypeptide is expressed. According to a preferred embodiment, the vertebrate cell is a CHO cell which comprises one or more frameshift mutations in exon 2 of one or preferably both alleles of the matriptase gene, whereby the effect of matriptase is impaired in said CHO cell.

[43] According to one embodiment, the vertebrate cell comprises one or more mutations in a polynucleotide sequence of the matriptase gene that encodes at least part of the catalytic domain of matriptase, whereby a non- or less functional expression product is obtained. The catalytic domain is the region of an enzyme that interacts with its substrate to cause the enzymatic reaction. One or more mutations can be introduced into this domain so that the catalytic activity of the protein is reduced or eliminated. The catalytic domain is coded by amino acids in exons 16, 17, 18 and 19. Thus, according to one embodiment, the vertebrate cell comprises one or more mutations in one or more exons selected from exon 16, exon 17, exon 18 and exon 19. According to one embodiment, the one or more mutations in the catalytic domain lead to a reduction or elimination of the catalytic activity of the matriptase. This may be achieved e.g. by a frameshift mutation, by a specific point mutation, a stop codon mutation and/or a deletion or insertion in the catalytic domain. According to one embodiment, one or more mutations are introduced so that the catalytic triad of matriptase is altered, thereby providing a non- or less functional protein. Catalytic inactive mutants of matriptase such as e.g. G827R-matriptase or S805A-matriptase have also been described in the literature (see Désilets et al, The Journal of Biological Chemistry Vol. 283, No. 16, pp. 10535-10542, 2008). Furthermore, the crystal structure of the catalytic domain of a recombinant matriptase is known. From this structure and sequence data the skilled person can derive further specific targets for mutations to impair the catalytic function of the matriptase.

[44] Altered cells wherein functional expression of matriptase is impaired may also be obtained using random mutagenesis or screening approaches. Respective methods are known in the prior art and therefore, do not need to be described in detail. Altered cells wherein functional expression of matriptase is impaired may then be identified using e.g. the method according to the seventh aspect.

[45] Functional expression of matriptase can also be influenced by altering the promoter and/or enhancer of the matriptase gene so that less or no transcript is produced, or by gene silencing technologies such as transcriptional or post-transcriptional gene silencing. According to one embodiment the isolated vertebrate cell comprises one or more mutations in the promoter region of the matriptase gene. For example, the promoter region may be altered to provide a less functional or non-functional promoter, the promoter may also be completely eliminated. Alternatively or in addition thereto, it is possible to add a polynucleotide sequence encoding a polypeptide including a stop codon between the promoter and the start codon of the matriptase gene which leads to the expression of the other polypeptide instead of the matriptase. The polypeptide encoded by the polynucleotide sequence inserted between the promoter and the start codon may for example be a reporter polypeptide such as green fluorescence protein (GFP). The signal of the reporter will indicate that the heterologous polynucleotide encoding the reporter is expressed instead of the matriptase thereby allowing to easily identify cells wherein the functional expression of matriptase is impaired.

[46] Reduction of functional gene expression may achieve a level wherein functional expression is even eliminated.

[47] Post-transcriptional gene silencing can be achieved e.g. by using antisense molecules or molecules that mediate RNA interference. Non-limiting examples will be briefly described in the following.

[48] Antisense polynucleotides may be designed to specifically bind to RNA, resulting in the formation of RNA-DNA or RNA-RNA hybrids, with an arrest of reverse transcription or messenger RNA translation. Many forms of antisense have been developed and can be broadly categorized into enzyme-dependent antisense or steric blocking antisense. Enzyme-dependent antisense includes forms dependent on RNase H activity to degrade target mRNA, including single-stranded DNA, RNA, and phosphorothioate antisense. Antisense polynucleotides are typically generated within the cell by expression from antisense constructs that contain the antisense strand as the transcribed strand. Trans-cleaving catalytic RNAs (ribozymes) are RNA molecules possessing endoribonuclease activity. Ribozymes may be specifically designed for a particular target and may be engineered to cleave any RNA species site-specifically in the background of cellular RNA. The cleavage event renders the mRNA unstable and prevents protein expression. The genome of the vertebrate cell can be altered so that a respective antisense molecule is e.g. permanently expressed.

[49] Another suitable option for reducing functional expression of the matriptase gene on a post-transcriptional level is based on RNA interference (RNAi). As is shown by the examples based on spike-in experiments, reducing the expression of matriptase by RNAi is effective in order to decrease the degree of clipping of recombinant polypeptides of interest that are expressed and secreted into the cell culture medium by the respectively altered vertebrate cells. Significantly more recombinant polypeptide stays intact in the supernatant/cell culture

medium upon silencing of the matriptase gene by RNAi and hence, can be harvested therefrom. In contrast, reduction of the expression levels of other proteases, even closely related serine proteases, had no or little effect on clipping. This emphasizes the importance of having identified with matriptase the major protease responsible for clipping when expressing a recombinant polypeptide of interest in a vertebrate cell such as preferably in a mammalian cell. Methods for silencing a target gene by RNAi are well known to the skilled person and thus, do not need any detailed description here. Examples of RNAi inducing compounds that can be used to silence the expression of the matriptase gene thereby providing an altered cell in which the effect of matriptase is impaired, include but are not limited to short interfering nucleic acids (siNA), short interfering RNA (siRNA), microRNA (miRNA), short hairpin RNAs (shRNA) as well as precursors thereof which are processed in the cell to the actual RNAi inducing compound. According to one embodiment, a siRNA is used for silencing. The siRNA may be provided as double-stranded molecule having 3' overhangs on each strand. Blunt ended molecules may also be used. Said siRNA may comprise desoxy- as well as ribonucleotides and furthermore, may comprise modified nucleotides. Several embodiments and variations of siRNA compounds are known in the prior art and can be used to reduce expression of the matriptase. Suitable siRNAs targeting the chosen/identified target sequences of the target genes on the RNA level can be identified by using proper computational methods, applying certain design-algorithms. In order to obtain a siRNA against the target transcript, the double-stranded molecule can be transfected directly into the cell. As is shown by the examples, even such transient methods that reduce expression of the matriptase are effective to prevent clipping of a polypeptide of interest in the conditioned medium obtained from such vertebrate cells. Alternatively, the siRNA may result from processing by dicer, an enzyme that converts either long dsRNAs or small hairpin RNAs (shRNAs) into siRNAs. These precursors or the final siRNA molecules can be produced exogenously (artificially) and can then be introduced into the vertebrate cells by various transfection methods. According to a further embodiment, the RNAi inducing compound is expressed by a vector that is transfected into the vertebrate cell. For siRNA, this can be done e.g. by the introduction of a loop between the two strands, thus producing a single transcript, which can be then processed into a functional siRNA in the vertebrate cell. Such transcription cassettes typically use an RNA polymerase III promoter (for example U6 or H1) which usually direct the transcription of small nuclear RNAs (e.g. for expressing shRNAs). It is assumed that the resulting shRNA transcript from the vector is then processed by dicer, thereby producing the double-stranded siRNA molecules, preferably having the characteristic 3' overhangs. According to one embodiment, such shRNA providing vector is stably integrated into the genome of the vertebrate cell. This embodiment is advantageous, as the downregulation of the matriptase gene is due to the constantly produced siRNA rather stable and not transient. E.g. cells comprising a respective shRNA providing vector can then be transfected with an expression vector comprising a polynucleotide encoding the polypeptide of interest that is to be expressed and secreted by the vertebrate cell. Furthermore, co-transfection strategies can be used, wherein the vector generating the shRNA is co-transfected with the expression vector comprising the polynucleotide encoding the polypeptide of interest.

[50] Transcriptional gene silencing may e.g. include epigenetic modifications. Furthermore, the sequence of the matriptase gene can be changed to reduce the half-life of the mRNA. This can also achieve a reduction in the functional expression of matriptase.

5 [51] According to one embodiment, matriptase expression is reduced or eliminated by targeting a regulatory element involved in the regulation of expression of the matriptase gene. E.g. a transcription factor, promoter (see also above), enhancer, UTRs or other regulatory element can be targeted e.g. by knock-out, deletion, mutation, down-regulation or
10 preventing or reducing functional expression of the matriptase gene and thereby impairing the effect of matriptase in said cells.

[52] According to one embodiment the vertebrate cell is altered to impair the function of the matriptase by heterologous expression of a mutant matriptase which is non- or less
15 functional than the endogenous matriptase. In this embodiment, the isolated vertebrate cell comprises in addition to the heterologous polynucleotide encoding the polypeptide of interest a further heterologous polynucleotide encoding a mutant matriptase. The mutant matriptase has a decreased or even no catalytic activity compared to the endogenous matriptase. By overexpressing a respective mutant matriptase the likelihood is increased that the mutant,
20 inactive form is inserted into the plasma membrane instead of the endogenous matriptase, in order to create a dominant negative phenotype. A further option to impair and hence reduce the effect of matriptase that is normally expressed by the cell is the heterologous expression of a protein such as an antibody or matriptase inhibitor which neutralizes and/or inhibits matriptase and hence impairs the effect of matriptase.

25 [53] According to one embodiment, the effect of matriptase is impaired in the cell by altering the functional expression of molecules that functionally interact with matriptase in the cell.

[54] According to one embodiment, the function of the matriptase is impaired in the cell by
30 impairing intracellular trafficking of the matriptase.

[55] According to another embodiment the function of the matriptase is impaired by impairing activation of the matriptase zymogen. According to one embodiment the function of the matriptase is impaired by upregulation of one or more endogenous cellular matriptase
35 inhibitors and/or by co-expression of a matriptase inhibitor. Several endogenous inhibitors of the matriptase have been described to date in different vertebrate cells such as HAI-1, HAI-2, alpha1-anti-trypsin, alpha2-antiplasmin, antithrombin (see review by Chen et al, 2012). If expressed by the vertebrate cell to be altered, an upregulation of the expression of these inhibitors and/or a co-expression of these inhibitors may also impair the function of the
40 endogenous protease matriptase, thereby reducing clipping of the secreted polypeptide of interest in the cell culture medium.

[56] According to one embodiment, functional expression of matriptase is impaired by recombinant expression of an antagonist such as an antibody or a binding domain thereof. E.g. a respective antagonist can be overexpressed by the vertebrate cell.

5 [57] According to one embodiment, the coding sequence of matriptase is altered so that the protein remains in the ER. E.g. the matriptase protein can be altered to include a KDEL motif which has the effect that matriptase remains in the ER. Similar approaches may also be used.

10 [58] Impairing the effect of the endogenous protease matriptase results in an altered cell wherein clipping of the recombinant polypeptide of interest that is expressed and secreted by said altered cell is reduced or even absent compared to a corresponding vertebrate cell in which the function of the endogenous protease matriptase is not impaired. As described herein, when using respectively altered cells wherein the effect of matriptase is impaired as
15 host cells for recombinant expression, no or less functional matriptase is active in the cell culture medium containing said host cells and into which the polypeptide of interest is secreted.

[59] According to one embodiment, expression of the matriptase gene is 40% or less, 35% or less, 30% or less, 25% or less, 20% or less, 15% or less, 12.5% or less, 10% or less, 5% or less, 2.5% or less, 1.5% or less, 1% or less or 0.05% or less compared to the expression of the matriptase gene in an unaltered corresponding reference vertebrate cell in which the effect of matriptase is not impaired (set as 100%). According to one embodiment, in the altered cell according to the first aspect expression of the matriptase gene is 0.1% or less, or less, 0.075% or less, 0.05% or less, 0.045% or less, 0.04% or less, 0.035% or less, 0.03% or less, 0.025% or less, 0.02% or less, 0.015% or less, 0.01% or less, 0.0075% or less, 0.005% or less, 0.0045% or less, 0.004% or less, 0.0035% or less, 0.003% or less, 0.0025% or less, 0.002% or less, 0.0015% or less or 0.001% or less compared to the expression of the 18S RNA (set as 100%) in said cell. According to one embodiment, expression of the matriptase
20 gene is 0.00075% or less, 0.0005% or less or 0.0004% or less compared to the expression of the 18S RNA (set as 100%) in said cell. The functional expression of the matriptase gene is reduced or eliminated such that clipping of a recombinant polypeptide of interest that is secreted by said cell into the cell culture medium is reduced compared to a corresponding reference vertebrate cell (from which the altered cell derived) wherein the functional
25 expression of the matriptase gene is not reduced or eliminated. E.g. an altered CHO cell in which functional expression of matriptase is reduced is compared to a CHO wildtype cell as reference vertebrate cell in order to analyze said feature. As test polypeptide, a polypeptide that is prone to clipping when expressed by the reference vertebrate cell is used. According to one embodiment, clipping is reduced by at least 2fold, at least 3fold, at least 5fold, at least
30 7.5fold, at least 8fold or at least 10fold compared to a corresponding reference vertebrate cell in which functional expression of the matriptase gene is not reduced or eliminated. This can be tested using e.g. the assays described in the examples.

[60] The altered vertebrate cell is derived from a species and cell type which endogenously expresses matriptase. The vertebrate cell preferably is a mammalian cell. Thus, all embodiments described herein for a vertebrate cell in general apply and specifically refer to the preferred embodiment wherein a mammalian cell is used. The term "isolated" is merely used to render clear that the vertebrate cell is not contained in a living organism such as an animal or human. As described herein, the vertebrate cell can be provided in form of a cell culture, cell line, cell clone and the like. Examples are also described below. As is described above, said vertebrate cell is altered to impair the effect of the matriptase compared to a corresponding unaltered vertebrate cell which endogenously expresses matriptase. Impairment is preferably achieved by reducing or eliminating functional expression of the matriptase gene. Embodiments are described above. In order to provide a production cell line with stable, uniform and thus predictable characteristics, it is preferred to alter the genome of the vertebrate cell to achieve matriptase impairment. Suitable embodiments are described above. The respectively altered vertebrate cell can then be transfected with an expression vector comprising a polynucleotide encoding a polypeptide of interest to provide the host cells according to the present disclosure which comprise a heterologous polynucleotide encoding the polypeptide of interest and which secrete the polypeptide of interest into the cell culture medium. The vertebrate cell preferably is a mammalian cell and may be e.g. selected from the group consisting of rodent cells, human cells and monkey cells. Preferred vertebrate cells are rodent cells such as e.g. cells derived from hamster or mouse. The rodent cell can be a cell line selected from the group consisting of a Chinese hamster cell line (such as e.g. a Chinese Hamster Ovary (CHO) cell line), a BHK cell line, a NS0 cell line, a C127 cell line, a mouse 3T3 fibroblast cell line, and a SP2/0 cell line. Particularly preferred is a CHO cell such as a CHO-K1 derived CHO cell. As is shown in the examples, reducing or eliminating the functional expression of matriptase in a CHO cell provides a conditioned cell culture medium in which clipping of the secreted polypeptide of interest is significantly decreased or even eliminated. Matriptase is also expressed in human cells. Thus, according to one embodiment, the vertebrate cell is derived from a human cell, which may be e.g. selected from the group consisting of a HEK293 cell, a MCF-7 cell, a PerC6 cell, a CAP cell, hematopoietic cells and a HeLa cell. Another alternative are monkey cells, which, e.g., may be selected from the group consisting of a COS cells, COS-1, a COS-7 cell and a Vero cell. According to one embodiment the vertebrate cell is provided as cell clone, cell line or cell culture.

[61] The "polypeptide of interest" is the recombinant polypeptide that is supposed to be expressed and secreted by the vertebrate cell in large quantity. The polypeptide of interest is encoded by a heterologous polynucleotide comprised in said cell. Said host cell may comprise more than one heterologous polynucleotide encoding a polypeptide of interest. The polypeptide of interest is secreted by the vertebrate cell into the cell culture medium from which it can be harvested, e.g. isolated and purified.

[62] A "heterologous polynucleotide" or "heterologous nucleic acid" and likewise expressions used herein in particular refer to a polynucleotide sequence that has been introduced into the vertebrate cell e.g. by the use of recombinant techniques such as transfection. A

“polynucleotide” in particular refers to a polymer of nucleotides which are usually linked from one deoxyribose or ribose to another and refers to DNA as well as RNA, depending on the context. The term “polynucleotide” does not comprise any size restrictions.

5 [63] The vertebrate cell may comprise a heterologous polynucleotide encoding a selectable marker and/or a heterologous polynucleotide encoding a reporter in addition to the at least one heterologous polynucleotide encoding the polypeptide of interest. This simplifies the selection of host cells which are successfully transfected and thus express the polypeptide of interest. Furthermore, the vertebrate cell may comprise several polynucleotides encoding
10 different selectable markers and/or reporter polypeptides.

[64] A “selectable marker” allows under appropriate selective culture conditions the selection of host cells expressing said selectable marker. A selectable marker provides the carrier of said marker under selective conditions with a survival and/or growth advantage. Typically, a
15 selectable marker gene will confer resistance to a selection agent such as a drug, e.g. an antibiotic or other toxic agent, or compensate for a metabolic or catabolic defect in the host cell. It may be a positive or negative selection marker. According to one embodiment, the selectable marker is a drug resistance marker encoding a protein that confers resistance to selection conditions involving said drug. A variety of selectable marker genes have been
20 described (see, e.g., WO 92/08796, WO 94/28143, WO2004/081167, WO2009/080759, WO2010/097240). E.g. at least one selectable marker may be used which confers resistance against one or more antibiotic agents. The selectable marker may according to one embodiment be an amplifiable selectable marker. An amplifiable selectable marker allows the selection of vector containing host cells and may promote gene amplification of said
25 vector in the host cells. Selectable marker genes commonly used with vertebrate cells include the genes for aminoglycoside phosphotransferase (APH), hygromycin phosphotransferase (hyg), dihydrofolate reductase (DHFR), thymidine kinase (tk), glutamine synthetase, asparagine synthetase, and genes encoding resistance to neomycin (G418), puromycin, hygromycin, zeocin, ouabain, blasticidin, histidinol D, bleomycin, phleomycin and
30 mycophenolic acid. According to one embodiment, a folate receptor is used as selectable marker in conjunction with the novel vertebrate cells described herein (see e.g. WO2009/080759), which preferably are mammalian cells. The folate receptor can also be used in combination with DHFR as selectable marker as is described in WO10/097240. A “reporter polypeptide” allows the identification of a cell expressing said reporter polypeptide
35 based on the reporting characteristics (e.g. fluorescence). Reporter genes usually do not provide the host cells with a survival advantage. However, the expression of the reporter polypeptide can be used to differentiate between cells expressing the reporter polypeptide and those cells which do not. Therefore, also a reporter gene enables the selection of successfully transfected host cells. Suitable reporter polypeptides include but are not limited
40 to as e.g. green fluorescence protein (GFP), YFP, CFP and luciferase. According to one embodiment, the reporter polypeptide has characteristics that enable the selection by flow cytometry.

[65] According to one embodiment, the at least one heterologous polynucleotide encoding the polypeptide of interest is integrated into the genome of said cell and wherein optionally, at least one heterologous polynucleotide encoding a selectable marker or reporter polypeptide is additionally integrated into the genome of said cell.

5

[66] An expression vector can be used to introduce a heterologous polynucleotide into the host cell. The polynucleotides can be comprised in an expression cassette. The polynucleotide(s) encoding the polypeptide of interest and the polynucleotide(s) encoding a selectable marker or reporter polypeptide may be located on the same or on different expression vectors. If they are located on different expression vectors, the expression vectors are co-transfected into the host cell. Such co-transfection strategies likewise enable selection as is well-known in the prior art. Introduction into the vertebrate cell may be achieved e.g. by transfecting a suitable expression vector comprising the polynucleotide encoding the polypeptide of interest into the host cells. The expression vector preferably integrates into the genome of the host cell (stable transfection). In case the heterologous nucleic acid is not inserted into the genome, the heterologous nucleic acid can be lost at the later stage e.g. when the cells undergo mitosis (transient transfection). Stable transfection is preferred for generating high expressing cell clones for producing a polypeptide of interest on industrial scale. This is particularly important for therapeutically active or diagnostic polypeptides of interest. Several appropriate methods are known in the prior art for introducing a heterologous nucleic acid such as an expression vector into vertebrate host cells, which preferably are mammalian host cells, and thus, do not need any detailed description herein. Respective methods include but are not limited to calcium phosphate transfection, electroporation, lipofection, biolistic- and polymer-mediated genes transfer and the like. Besides traditional random integration based methods also recombination mediated approaches can be used to transfer the heterologous polynucleotide into the host cell genome. As respective methods are well known in the prior art, they do not need any detailed description here. Non-limiting embodiments of suitable vector designs are also described subsequently and it is referred to the respective disclosure.

30

[67] Expression vectors used to achieve expression of a polypeptide of interest in the host cell usually contain transcriptional control elements suitable to drive transcription such as e.g. promoters, enhancers, polyadenylation signals, transcription pausing or termination signals usually as element of an expression cassette. Suitable translational control elements are preferably included in the vector, such as e.g. 5' untranslated regions leading to 5' cap structures suitable for recruiting ribosomes and stop codons to terminate the translation process. The resultant transcripts harbour functional translation elements that facilitate protein expression (i.e. translation) and proper translation termination. A functional expression unit, capable of properly driving the expression of an incorporated polynucleotide is also referred to as an "expression cassette". It is well-known to the skilled person how an expression cassette shall be designed in order to allow the expression and secretion of a polypeptide of interest in a vertebrate cell.

40

[68] To achieve secretion of the recombinant polypeptide of interest into the cell culture medium, an appropriate leader peptide is provided in the polypeptide of interest. Leader sequences and expression cassette designs to achieve secretion of the polypeptide of interest are well known in the prior art and therefore, do not need to be described herein.

5

[69] Any polypeptide of interest can be expressed in the vertebrate cell according to the invention. The term "polypeptide" refers to a molecule comprising a polymer of amino acids linked together by (a) peptide bond(s). Polypeptides include polypeptides of any length, including proteins (e.g. having more than 50 amino acids) and peptides (e.g. 2 – 49 amino acids). Polypeptides include proteins and/or peptides of any activity, function or size, and may include e.g. enzymes (e.g. kinases, phosphatases), receptors, transporters, bactericidal and/or endotoxin-binding proteins, structural polypeptides, membrane-bound polypeptides, glycopolypeptides, globular proteins, immune polypeptides, toxins, antibiotics, hormones, growth factors, blood factors, vaccines, viral glycopolypeptides and the like. The polypeptide of interest that is expressed according to the teachings described herein may also be a subunit or domain of a polypeptide, such as e.g. a heavy chain or a light chain of an antibody or a functional fragment or derivative thereof. The term "polypeptide of interest" may refer to such individual subunit or domain or the final protein that is composed of the respective subunits or domains, depending on the context.

10

15

20

25

30

35

40

[70] According to one embodiment, the polypeptide of interest is selected from a therapeutic or diagnostic polypeptide. Therapeutic and hence therapeutically active polypeptides are particularly important. The term therapeutic polypeptides also encompasses prophylactic polypeptides, e.g. used for vaccination. The polypeptide may be selected from the group consisting of peptide hormones, interleukins, tissue plasminogen activators, cytokines, growth factors, immunoglobulins, in particular antibodies or functional antibody fragments or variants or derivatives thereof and Fc-fusion proteins. In one embodiment the polypeptide of interest is an immunoglobulin molecule such as an antibody. The term "antibody" as used herein particularly refers to a protein comprising at least two heavy chains and two light chains connected by disulfide bonds. The term "antibody" includes naturally occurring antibodies as well as all recombinant forms of antibodies, e.g., humanized antibodies, fully human antibodies and chimeric antibodies. Each heavy chain is usually comprised of a heavy chain variable region (VH) and a heavy chain constant region (CH). Each light chain is usually comprised of a light chain variable region (VL) and a light chain constant region (CL). The term "antibody", however, also includes other types of antibodies such as single domain antibodies, heavy chain antibodies, i.e. antibodies only composed of one or more, in particular two heavy chains, and nanobodies, i.e. antibodies only composed of a single monomeric variable domain. Nanobodies may also be linked to form multivalent structures. As discussed above, the polynucleotide encoding the polypeptide of interest may also encode one or more subunits or domains of an antibody, e.g. a heavy or a light chain or a functional fragment or derivative thereof, as polypeptide of interest. Said subunits or domains can be expressed either from the same or different expression cassettes.

[71] A "functional fragment or derivative" of an antibody in particular refers to a polypeptide which is derived from an antibody and is capable of binding to the same antigen, in particular to the same epitope as the antibody. It has been shown that the antigen-binding function of an antibody can be executed by fragments of a full-length antibody or derivatives thereof.

5 Examples of fragments or derivatives of an antibody include (i) Fab fragments, monovalent fragments consisting of the variable region and the first constant domain of each the heavy and the light chain; (ii) F(ab)₂ fragments, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) Fd fragments consisting of the variable region and the first constant domain CH1 of the heavy chain; (iv) Fv fragments consisting of
10 the heavy chain and light chain variable region of a single arm of an antibody; (v) scFv fragments, Fv fragments consisting of a single polypeptide chain; (vi) (Fv)₂ fragments consisting of two Fv fragments covalently linked together; (vii) a heavy chain variable domain; and (viii) multibodies consisting of a heavy chain variable region and a light chain variable region covalently linked together in such a manner that association of the heavy
15 chain and light chain variable regions can only occur intermolecular but not intramolecular.

[72] According to one embodiment the polypeptide of interest that is expressed by the altered vertebrate cell is susceptible to clipping by proteases. According to one embodiment, the polypeptide of interest comprises at least one clipping site that is recognized by matriptase.

20 Several polypeptides selected from the group consisting of glycoproteins, antibodies, non-IgG proteins, Fab fragments, protein complexes, peptidases, signal peptides, Fc-fusion proteins, nanobodies, growth factors, hormones, cytokines, viral glycopolypeptides, blood factors and enzymes have been identified that are prone to clipping. Many of these polypeptides contain more than one clipping site and are particularly sensitive to clipping.
25 Expressing a polypeptide that is prone to clipping as polypeptide of interest in an altered vertebrate cell according to the present disclosure is advantageous because clipping of the secreted polypeptide of interest in the cell culture medium is significantly reduced or even eliminated as is demonstrated by the examples.

30 [73] The vertebrate cell may or may not comprise an endogenous polynucleotide corresponding to, respectively being identical to the polynucleotide encoding the polypeptide of interest. According to one embodiment, the vertebrate cell does not comprise an endogenous gene corresponding to the polypeptide of interest.

35 [74] According to one embodiment, the polypeptide of interest is not or does not comprise matriptase. According to one embodiment, the polypeptide of interest is not or does not comprise HAI-1.

40 [75] The expression vector or the combination of expression vectors that are comprised in the vertebrate cell may additionally comprise further vector elements. E.g. at least one additional polynucleotide encoding a further product of interest can be comprised. As explained above and as becomes apparent from the above described examples of polypeptides that can be expressed according to the present teachings, the final polypeptide that is to be produced and secreted by the host cell can also be a protein that is composed of

several individual subunits or domains. An example of a respective protein is an immunoglobulin molecule, in particular an antibody that comprises e.g. heavy and light chains. There are several options for producing a respective protein that is composed of different individual subunits or domains and appropriate vector designs are known in the art.

5 According to one embodiment, two or more subunits or domains of said protein are expressed from one expression cassette. In this embodiment, one long transcript is obtained from the respective expression cassette that comprises the coding regions of the individual subunits or domains of the protein. According to one embodiment, at least one IRES element (internal ribosomal entry site) is functionally located between the coding regions of the individual subunits or domains and each coding region is preceded by a secretory leader sequence. Thereby, it is ensured that separate translation products are obtained from said transcript and that the final protein can be correctly assembled and secreted. Respective technologies are known in the prior art and thus, do not need any detailed description herein. For some embodiments such as the expression of antibodies it is even preferred to express the individual subunits or domains from different expression cassettes. According to one embodiment, the expression cassette used for expressing the product of interest is a monocistronic expression cassette. All expression cassettes comprised in the expression vector or combination of expression vectors may be monocistronic. According to one embodiment, accordingly, each expression cassette designed for expressing a product of interest comprises a polynucleotide encoding one subunit or domain of the protein to be expressed as polypeptide of interest. E.g. in case of antibodies, one expression cassette may encode the light chain of an antibody and another expression cassette may encode the heavy chain of the antibody. After expression of the individual subunits or domains from the individual expression cassettes, the final protein such as an antibody is assembled from said subunits or domains and secreted by the host cell. This embodiment is particularly suitable for expressing immunoglobulin molecules such as antibodies.

[76] As described above, the polynucleotide(s) encoding the polypeptide of interest and the polynucleotides encoding the selectable marker(s) and/or reporter polypeptide(s) are preferably comprised in expression cassettes. Several embodiments are suitable. For example, each of said polynucleotide(s) can be comprised in a separate expression cassette. This is also referred to as monocistronic setting. However, it is also within the scope of the present invention that at least two of the respective polynucleotides are comprised in one expression cassette. According to one embodiment, at least one internal ribosomal entry site (IRES) element is functionally located between the polynucleotides that are expressed from the same expression cassette. Respective IRES based expression technologies and other bi- and polycistronic systems are well known and thus need no further description here.

B. Method for producing a vertebrate cell according to the first aspect

40 [77] According to a second aspect, a method is provided for producing a vertebrate cell according to the first aspect, said method comprising altering a vertebrate cell to impair the effect of matriptase and introducing a polynucleotide encoding a polypeptide of interest into said cell, wherein said polypeptide of interest is then secreted by the vertebrate cell.

[78] Suitable and preferred embodiments to impair the function of the endogenous protease matriptase are described above in conjunction with the vertebrate cells according to the first aspect and it is referred to the above disclosure, which also applies here. Therefore, the method may comprise impairing the effect of matriptase as described above in conjunction with the altered vertebrate cells according to the first aspect. Non-limiting embodiments are again briefly described in the following.

[79] According to one embodiment, the method comprises reducing or eliminating the functional expression of the matriptase gene thereby impairing the effect of the matriptase. As described above, this impairment has the effect that no or reduced amounts of functional matriptase is present in active form in the cell culture medium containing the altered cells, e.g. because no or reduced amounts of functional matriptase is presented on the cell surface and/or is released, e.g. shedded, by the altered cell. Thereby, no or less matriptase is present in proteolytically active form in the cell culture medium into which the recombinant polypeptide of interest is secreted. As the present disclosure shows that matriptase is a key protease responsible for clipping of the secreted polypeptide of interest, this alteration of the cell advantageously reduces or even eliminates clipping of the polypeptide of interest. Suitable ways for reducing or eliminating functional expression of matriptase are described above in conjunction with the vertebrate cells according to the first aspect and it is referred thereto. According to one embodiment, the genome of the vertebrate cell is altered to reduce or eliminate the functional expression of matriptase gene. For example, a gene knock-out may be introduced into the matriptase gene. According to one embodiment, such gene knock-out is introduced into all copies of the matriptase gene. According to one embodiment, the matriptase gene is deleted or knocked-out by introducing one or more mutations such as e.g. frameshift and/or stop codon mutations. All copies of the matriptase gene may be respectively altered in the genome.

[80] According to one embodiment, said one or more mutations are comprised in a coding region of the matriptase gene and result in a non- or less functional expression product. Details are described above in conjunction with the altered vertebrate cells according to the first aspect and it is referred to the respective disclosure which also applies here. According to one embodiment, one or more mutations are introduced into a region of the matriptase gene that encodes an amino acid sequence present in one or more than one catalytically active splice variant of matriptase. Several of the matriptase exons are found in the majority or even all identified catalytically active splicing variants. As described above and demonstrated in the examples, the polynucleotide sequence of the matriptase gene that encodes exon 2 of matriptase is a suitable target for introducing one or more mutations such as e.g. frame-shift mutations, because exon 2 is expressed in the majority of splice variants of matriptase and a respective mutation renders a non- or less functional expression product. Moreover, exons close to the N-terminus of matriptase such as exon 2 are advantageous targets for introducing one or more frameshift mutations leading to the expression of a non- or less functional truncated protein. Furthermore, one or more mutations can be introduced into the catalytical domain in order to disrupt the function of matriptase.

[81] According to one embodiment, the vertebrate cell is a mammalian cell. According to one embodiment, the vertebrate cell is a rodent cell. Preferably, the rodent cell is a hamster cell such as a CHO cell. According to one embodiment, a CHO cell, preferably derived from the cell line K1, is used in order to provide an altered vertebrate cell line wherein the effect of the endogenous protease matriptase is impaired, preferably by reducing or eliminating the functional expression of the matriptase gene.

[82] If the genome of the host cell is altered to achieve impairment, it is preferred to first alter the vertebrate cell to basically permanently impair the effect and hence function of the endogenous protease matriptase and then introduce the heterologous polynucleotide encoding the polypeptide of interest. In case the alteration is not permanent as is e.g. the case when transiently silencing the matriptase gene, e.g. via RNAi, one may first introduce the heterologous polynucleotide encoding the polypeptide of interest and then alter the vertebrate cell to impair the function of the endogenous protease matriptase, e.g. by post-transcriptional gene silencing. Suitable methods were described above and are known to the skilled person. As described above, according to a preferred embodiment, the genome of the vertebrate cell is altered to achieve impairment. The method according to the seventh aspect can be used in order to identify and select a host cell wherein the effect of matriptase is impaired.

[83] According to one embodiment, the method according to the second aspect further comprises introducing into a vertebrate cell which is altered so that the functional expression of matriptase gene is reduced or eliminated, at least one polynucleotide encoding a polypeptide of interest and preferably at least one polynucleotide encoding a selectable marker. According to one embodiment, the polynucleotide encoding a polypeptide of interest and the polynucleotide encoding a selectable marker are located on the same or on different expression vectors. Suitable and preferred embodiments are described above and it is referred to the respective disclosure, which also applies here. Introduction can be achieved by transfection, wherein stable transfection is preferred. A vertebrate host cell that successfully expresses the polypeptide of interest can be selected using e.g. the method according to the fifth aspect. It is referred to the subsequent disclosure.

C. Methods for recombinantly producing a polypeptide of interest

[84] According to a third aspect, a method is provided for recombinantly producing a polypeptide of interest, comprising utilizing a vertebrate cell according to the first aspect as host cell for recombinant expression of the polypeptide of interest. As described above, due to the decreased level of proteolytic degradation of the polypeptide of interest in the cell culture medium that is achieved when using these novel vertebrate cells for recombinant expression, these novel vertebrate cells, which preferably are mammalian cells, are particularly suitable as host cells for recombinantly producing a polypeptide of interest. Suitable and preferred examples of the vertebrate host cell, which is altered to impair the effect of the matriptase, preferably by reducing or eliminating functional expression of the matriptase gene, as well as suitable and preferred examples of the polypeptide of interest

are described in detail above and it is referred to the above disclosure which also applies here.

5 [85] According to one embodiment, the method comprises introducing into a vertebrate host cell which is altered so that the effect of the endogenous protease matriptase is impaired at least one polynucleotide encoding a polypeptide of interest and selecting a vertebrate host cell which recombinantly expresses the polypeptide of interest. Introduction can be achieved by transfection as is well-known and also described herein. Selection may occur using the method according to the fifth aspect of the present disclosure. Preferably, host cells are
10 selected wherein the heterologous polynucleotide encoding the polypeptide of interest is stably integrated into the genome of the host cell.

[86] According to one embodiment, the method comprises
15 (a) culturing vertebrate host cells according to first aspect under conditions that allow for the expression and secretion of the polypeptide of interest;
(b) isolating the polypeptide of interest from the cell culture medium; and
(c) optionally processing the isolated polypeptide of interest.

Hence, as preferred embodiment of the method according to the third aspect, a method for
20 recombinantly producing a polypeptide of interest is provided, comprising
(a) culturing vertebrate host cells according to the first aspect under conditions that allow for the expression and secretion of the polypeptide of interest into the cell culture medium;
(b) isolating the polypeptide of interest from the cell culture medium; and
(c) optionally processing the isolated polypeptide of interest.
25

[87] Said host cells may be cultured under serum-free conditions. The polypeptide is expressed and secreted into the culture medium and can be obtained therefrom. For the purpose of secretion, an appropriate leader peptide is provided so that the polypeptide of interest is secreted. Leader sequences and expression cassette designs to achieve secretion
30 are well known in the prior art and therefore do not need to be described herein.

[88] As described above, when using the altered vertebrate cells according to the first aspect for production host cells, no or reduced amounts of functional matriptase are effective in the cell culture medium containing said cells, whereby proteolytic degradation of the recombinant
35 polypeptide of interest that is secreted into the cell culture medium is significantly reduced or is even completely avoided. This, because it was surprisingly found that among the hundreds of different proteases expressed by vertebrate cells, matriptase is the key protease that causes clipping of recombinantly expressed polypeptides in the cell culture medium. Therefore, when using the novel vertebrate host cells according to the present disclosure as
40 production host cells, it is not necessary to perform additional measures to reduce or avoid proteolytic degradation and hence clipping of the polypeptide of interest in the cell culture medium. Therefore, according to one embodiment, no protease inhibitor is added to the cell culture medium. According to one embodiment, the cultivation temperature is not lowered to reduce the proteolytic activity in the cell culture medium. The present disclosure

encompasses methods wherein the polypeptide of interest is reengineered to remove one or more motives prone to clipping by proteases. However, according to one embodiment, the polypeptide of interest is not reengineered in order to remove one or more amino acid motives that are prone to clipping by matriptase in order to reduce or prevent clipping.

5 According to this embodiment, a respective reengineering may be performed if desired in order to remove one or more amino acid motives that are prone to clipping by a protease different than matriptase. However, according to one embodiment, the polypeptide of interest is not reengineered at all in order to remove one or more amino acid motives that are prone to clipping by a protease in order to reduce or prevent clipping. As is shown by the examples,
10 clipping is effectively reduced or even completely eliminated when using the altered cells described herein so that respective reengineering approached become basically obsolete.

[89] Examples of polypeptides of interest that may be produced with the altered vertebrate cell are described above in conjunction with the first aspect of the invention and it is referred
15 to the respective disclosure, which also applies here. According to one embodiment, the production method is used for producing a polypeptide of interest which has one or more of the following characteristics:

- a) it is therapeutic or diagnostic polypeptide;
- 20 b) it is susceptible to clipping by proteases;
- c) it comprises at least one clipping site for matriptase;
- d) it is a glycopolypeptide;
- e) it is selected from the group consisting of glycoproteins, antibodies, non-IgG proteins, Fab fragments, protein complexes, peptidases, signal peptides, Fc-
25 fusion proteins, nanobodies, growth factors, hormones, cytokines, blood factors and enzymes;
- f) it is not or does not comprise matriptase; and/or
- g) it is not or does not comprise HAI-1.

30 [90] The polypeptide of interest that is produced is isolated from the cell culture medium and optionally further processed by methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, ultra-filtration, extraction or precipitation. Further processing steps such as purification steps may be performed by a variety of procedures known in the art
35 including, but not limited to, chromatography (e.g. ion exchange, affinity, hydrophobic, chromatofocusing, protein A or protein G chromatography and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g. ammonium sulphate precipitation) or extraction. Furthermore, the isolated and purified polypeptide of interest may be further processed, such as e.g. modified and/or formulated
40 into a composition, e.g. a pharmaceutical composition.

[91] According to a fourth aspect, a method for recombinantly producing a polypeptide of interest is provided, comprising

(a) culturing vertebrate host cells which comprise at least one heterologous polynucleotide encoding a polypeptide of interest under conditions that allow for the expression and secretion of the polypeptide of interest into the cell culture medium, wherein the cell culture medium comprises a protease inhibitor that is selective for matriptase;

- 5 (b) isolating the polypeptide of interest from the cell culture medium; and
(c) optionally processing the isolated polypeptide of interest.

[92] The method according to the fourth aspect is also based on the important finding described herein that matriptase is the key protease responsible for the proteolytic degradation of the secreted recombinant polypeptide of interest in the cell culture medium. In the method according to the fourth aspect, the proteolytic degradation of the polypeptide of interest in the cell culture medium, also referred to as "clipping" (see above) is avoided by adding a protease inhibitor that is selective for matriptase to the cell culture medium. A protease inhibitor that is selective for matriptase is also referred to herein as selective matriptase inhibitor. Of course, also more than one selective matriptase inhibitors can be added. As is demonstrated by Example 4, adding a selective matriptase inhibitor to the cell culture medium allows to significantly reduce or even abolish clipping of the polypeptide of interest even when using vertebrate cells that normally express matriptase. Thus, this embodiment allows to use unaltered vertebrate cells wherein accordingly, the effect of matriptase is not impaired because impairment is achieved in the cell culture medium.

[93] The selective matriptase inhibitor present in the cell culture medium leads to a decrease or abolishment of the proteolytic activity of matriptase in the cell culture medium. The matriptase inhibitor may e.g. be a competitive inhibitor that competes with the substrate for the active site of the matriptase or an allosteric inhibitor that modifies the structure of the matriptase to reduce its activity. A matriptase inhibitor is considered selective, if its inhibitory activity against matriptase is higher than its inhibitory activity against other serine proteases, in particular other type II transmembrane serine proteases.

[94] The selective matriptase inhibitor may be of any kind as long as it does not show harmful effects on the host cell that is used for producing the recombinant polypeptide of interest. The selective matriptase inhibitor may be selected from i) biological matriptase inhibitors which include but are not limited to antibody based or antibody derived selective matriptase inhibitors, peptides or proteins and ii) chemical matriptase inhibitors such as small molecules.

[95] According to one embodiment, the selective matriptase inhibitor is at least 5 fold, at least 10 fold, at least 25 fold, at least 50 fold, at least 100 fold or at least 250 fold more selective and thus specific for matriptase than to other proteases. However, the selectivity may also be substantially higher, such as e.g. at least 500 fold, at least 5000 fold, at least 10000 fold, at least 50000 fold, at least 500000 fold and at least 5000000 fold. Ranges include but are not limited to 5 fold to 10 000 000 fold, 10 fold to 5000000 fold, 25 fold to 1000000 fold, 50 fold to 5000000 fold. A higher selectivity of e.g. at least 5000 fold or at least 10 000 fold and higher may be achieved e.g. with selective biological matriptase inhibitors, such as e.g. matriptase specific antibodies or Fab-fragments.

[96] Selectivity may be determined e.g. based on the k_i -value, which is for a selective matriptase inhibitor lower than for other tested proteases. According to one embodiment, the matriptase inhibitor inhibits matriptase selective with a k_i value of 200nM or less, 150nM or less, 100nM or less, 50nM or less, 25nM or less, 20nM or less, 15nM or less, 10nM or less, 7.5nM or less, 5nM or less, 2.5nM or less, 1nM or less or 0.75nM or less.

[97] As mentioned above, several selective matriptase inhibitors are described in the prior art. E.g. Faraday et al describe different antibody based selective matriptase inhibitors, namely a Fab fragment (Farady et al., 2008 J.Mol.Biol. (2008) 380, 351-360) and two scFv antibody inhibitors (J Mol Biol. 2007 June 15; 369 (4): 1041-1051). The single chain variable fragment (scFv) antibody inhibitors E2 and S4 with K_i values of 12 pM and 160 pM, were described as particularly selective. An antibody-based selective matriptase inhibitor may be active against matriptase derived from different species. As is demonstrated by Example 4, e.g. the Fab-fragment against human matriptase was also effective against matriptase derived from mouse and hamster. However, respective antibody-based selective matriptase inhibitors may also be generated against the specific matriptase expressed by the concerned host cell in order to ensure a maximal inhibitory effect and allow to use lower concentrations of the selective matriptase inhibitor in the cell culture medium. Moreover, antibodies specific for the catalytic site of the matriptase have been described (US 2006/171884 A1). Peptidomimetic molecule selective matriptase inhibitors such as the 3-amidinophenyl-alanine type molecules CJ-730 and CJ-697 have k_i -values for matriptase of 46 nM and 26 nM, respectively (described in Förbs et al. International Journal of Oncology 27: 1061-1070 2005). As is described in Förbs, addition of these compounds to cell culture of tumour cell lines reduces the matriptase activity significantly without influencing the proliferation of the cells. Selective inhibitors of matriptase are also described in Steinmetzer et al, J. Med. Chem. 2006, 49, 4116-4126 and Goswami et al, ACS Med. Chem. Lett., 2013 4 (12) pp 1152-1157. Furthermore, structural variants of the eglin c, a small monomeric protein which is a serine protease inhibitor, have been identified as respective matriptase inhibitors (Desilets et al. FEBS Letters 580 (2006) 2227-2232). Here, in a screening assay different structural variants of eglin c were created and tested for their specificity for matriptase. A single amino acid exchange (L45R) in the reactive site loop makes eglin c matriptase specific (k_i = 18 nM). Thus, many selective matriptase inhibitors are known that can be used for the purpose of the present disclosure in order to selectively inhibit matriptase in the cell culture medium thereby reducing or preventing clipping of the polypeptide of interest that is recombinantly expressed and secreted into the cell culture medium by the vertebrate host cell.

[98] For impairing the effect of the matriptase in the cell culture medium, the selective matriptase inhibitor may be added to the cell culture medium in which the expression host cells are cultivated. The concentration of an individual selective matriptase inhibitor required to achieve sufficient matriptase inhibition in the cell culture medium so that the proteolytic degradation of the polypeptide of interest is reduced can be determined based on routine experiments. The selective matriptase inhibitor may be added to the cell culture medium

before or after addition of the host cells. According to one embodiment, the selective matriptase inhibitor is added at the start of cell cultivation. According to one embodiment, the selective matriptase inhibitor is added at a point in time, wherein the host cells begin secretion of the polypeptide of interest. The selective matriptase inhibitor may also be added continuously. For example, in fed batch cultivation the selective matriptase inhibitor may be included in the feed.

[99] Details with respect to the polypeptide of interest, the isolation of the polypeptide of interest from the cell culture medium as well as the subsequent processing are described above e.g. in conjunction with the method according to the third aspect and it is referred to the above disclosure which also applies here.

[100] Considering the cell culture volume required for large scale production, large quantities of the (one or more) selective matriptase inhibitor must be added to the cell culture medium even if a matriptase inhibitor with a high selectivity is used in order to efficiently reduce or prevent clipping of the polypeptide of interest in the cell culture medium. This can be costly. Additionally, for many polypeptides of interest such as biopharmaceuticals, the added selective matriptase inhibitor must be eliminated during the subsequent isolation and purification of the polypeptide of interest to avoid contaminations of the final polypeptide of interest with the matriptase inhibitor. Such removal step may comprise any known purification step such as ion exchange, affinity, size exclusion, or reversed phase chromatography. The eligible method of removal is dependent on the type of selective matriptase inhibitor used. As such steps can be laborious and may increase costs, the method according to the third aspect which involves the use of altered vertebrate cells according to the first aspect is preferred.

D. Selection method

[101] According to a fifth aspect, a method is provided for selecting a host cell which recombinantly expresses a polypeptide of interest, comprising

- (a) providing vertebrate cells according the first aspect of the invention as host cells; and
- (b) selecting one or more host cells expressing the polypeptide of interest.

[102] The vertebrate host cells according to the first aspect, including suitable and preferred embodiments, as well as their advantages are described in detail above and it is referred to the respective disclosure which also applies here. Preferably, the vertebrate cell is a mammalian cell. According to one embodiment, stage (a) of the selection method according to the fifth aspect comprises transfecting vertebrate cells, which are altered to impair the effect of the endogenous protease matriptase with at least one polynucleotide encoding the polypeptide of interest, to provide vertebrate host cells according to the first aspect. According to one embodiment, the cells do not comprise a heterologous polynucleotide encoding a polypeptide of interest prior to transfection of the polynucleotide encoding the polypeptide of interest to be expressed and secreted by the host cell. As

described, mammalian cells are preferably used as vertebrate host cells. The polynucleotide encoding the product of interest may be comprised in an expression vector that is then transfected into the vertebrate cell.

5 [103] According to one embodiment said host cells provided in stage (a) additionally comprise at least one heterologous polynucleotide encoding a selectable marker and stage (b) comprises culturing said plurality of host cells under conditions selective for the selectable marker. Selection stage (b) may be a multi-step selection process comprising several selection steps in order to select and thus identify host cells that express the polypeptide of interest. Thus, according to one embodiment selection stage (b) may comprise several selection steps. For example, stage (b) may include one or more selection steps to select cells that were successfully transfected as well as one or more subsequent selection steps to select high expressing cells from the pool of successfully transfected cells. The appropriate selection strategy depends on the design of the expression vector that is used for introducing the polynucleotide encoding the polypeptide of interest and in particular depends on the used selection marker(s) and/or reporter(s). Non-limiting embodiments will be described in the following.

20 [104] As described above, the vertebrate host cells may comprise at least one heterologous polynucleotide encoding a selectable marker. The polynucleotide encoding the selectable marker can be introduced into the host cell together with the polynucleotide encoding the polypeptide of interest using either the same or a different, co-transfected expression vector. Stage (b) then comprises culturing said plurality of host cells under conditions providing a selection pressure to the host cells to select successfully transfected host cells, e.g. using an appropriate selection medium. As used herein, a "selection medium" in particular refers to a cell culture medium useful for the selection of host cells that express the selectable marker. It may include e.g. a selection agent such as a toxic agent which allows selecting successfully transfected host cells. Alternatively, an essential compound can be absent or its concentration can be reduced in the selection medium. According to one embodiment, host cells which were not successfully transfected and hence, do not express the selection marker(s) or wherein expression is low cannot proliferate or die under the selective cultivation conditions. In contrast, host cells which were successfully transfected with the expression vector(s) and which express the selection marker(s) (and accordingly the co-introduced polypeptide of interest) with sufficient yield are resistant to or are less affected by the selection pressure and therefore, can proliferate, thereby outgrowing the host cells which were not successfully transfected or wherein the integration site into the genome of the cell is not favourable. The selectable marker may be selected from the group consisting of antibiotic resistance markers, drug resistance markers and metabolic markers. Suitable examples for selectable markers and selection principles are described above in conjunction with the first aspect and appropriate selection conditions for the individual selectable markers are also well-known to the skilled person. As described above, alternatively or additionally, a reporter polypeptide based selection can be performed.

[105] According to one embodiment, the vertebrate cells provided in stage (a) are mammalian cells. According to one embodiment, the mammalian cells are rodent cells, preferably hamster cells such as CHO cells. Suitable and preferred embodiments are described above in conjunction with the first aspect and it is referred to the above disclosure.

5 Further preferred embodiments in particular with respect to the vertebrate host cells according to the first aspect, the expression vector, or combination of expression vectors are likewise described in detail above. It is referred to the above disclosure.

10 [106] Cells obtained as a result of the selection method according to the fifth aspect can be isolated and cultured as individual cells. It is, however, also possible to use an enriched population of genetically different host cells, i.e. a cell pool, in the downstream process. The obtained host cells can also be subjected to additional qualitative or quantitative analysis, or can be used e.g. in the development of a clonal cell line for protein production. A clonal cell line may be established from a selected host cell which stably expresses the polypeptide of
15 interest with high yield.

[107] According to one embodiment, selected cells are cultivated to provide cell clones, in particular in the form of clonal cell cultures. A clonal cell culture is a cell culture derived from one single ancestral cell. In a clonal cell culture, all cells are clones of each other. Preferably,
20 all the cells in a cell culture contain the same or substantially the same genetic information. In certain embodiments, the amount or concentration of the polypeptide of interest in the cell culture is determined to evaluate the productivity. E.g. the titer can be measured by analysing the culture supernatant. Furthermore, a stability study can be performed with the obtained cell clones.

25

E. Use of altered vertebrate cells for recombinant production

[108] According to a sixth aspect, the use of an isolated vertebrate cell is provided for recombinant production of a polypeptide of interest that is secreted from the vertebrate cell, wherein the used cell is altered to impair the effect of the endogenous protease matriptase.
30 According to one embodiment of the sixth aspect the effect of the matriptase is impaired as is described in detail above in conjunction with the first aspect, preferably by reducing or eliminating functional expression of the matriptase gene as is described above in further detail. The vertebrate cell is preferably a mammalian cell. Thus, according to one embodiment, the effect of the endogenous protease matriptase is impaired as described
35 above in conjunction with the altered vertebrate cells according to the first aspect. According to one embodiment, a polynucleotide encoding a polypeptide of interest is introduced into said cell. After introduction, a vertebrate cell is provided which comprises a heterologous polynucleotide encoding a polypeptide of interest that is then secreted from the vertebrate cell into the cell culture medium. Thus, the use may comprise introducing a heterologous
40 polynucleotide encoding a polypeptide of interest into said vertebrate cell, wherein said polypeptide of interest is secreted by the vertebrate cell. Details with respect to the polypeptide of interest are likewise described in detail above and it is referred to the respective disclosure which also applies here. Methods for introducing a polynucleotide into a vertebrate cell are known to the skilled person and are also briefly described above.

[109] According to one embodiment, the vertebrate cell does not comprise a heterologous polypeptide encoding a polypeptide of interest prior to introducing the polynucleotide encoding the polypeptide of interest to be expressed. According to one embodiment, said cell further does not comprise a heterologous polynucleotide encoding a selectable marker and/or a heterologous polynucleotide encoding a reporter polypeptide. According to one embodiment, the vertebrate cell does not comprise any heterologous polynucleotide prior to introduction of the polynucleotide encoding the polypeptide of interest. A respective "empty" vertebrate cell which is altered to impair the effect of the endogenous protease matriptase can be used e.g. as cloning cell line for recombinant production technologies. A respective cell line can be transfected with a heterologous polynucleotide encoding a polypeptide of interest, e.g. using an appropriate expression vector. Such "empty" vertebrate cells which do not yet express and secrete a recombinant product can thus be transfected with different expression vectors, depending on the desired polypeptide of interest that is supposed to be recombinantly produced. Thus, such vertebrate cell line can be used for different projects, i.e. for the production of different polypeptides of interest. According to one embodiment, the vertebrate cell is a vertebrate cell according to the first aspect. Details are described above and it is referred to the above disclosure.

F. Method for selecting vertebrate host cells with impaired matriptase function

[110] According to a seventh aspect, the present disclosure pertains to a method for selecting a vertebrate cell for recombinant production of a polypeptide of interest, comprising analyzing if the endogenous protease matriptase is functionally expressed in the vertebrate cell and selecting a vertebrate cell in which the effect of such endogenous matriptase is impaired for recombinant production of the polypeptide of interest. This selection process allows identifying a host cell that is capable of expressing and secreting a recombinant polypeptide of interest, wherein clipping of the secreted polypeptide of interest in the cell culture medium is reduced. Respective host cells are particularly suitable for recombinant production.

[111] This analytical method can be advantageously used e.g. in combination with the method according to the second aspect of the present disclosure in order to identify whether a vertebrate cell was produced wherein the effect of the matriptase gene was impaired. According to a preferred embodiment, the method comprises analyzing whether the functional expression of the matriptase gene is reduced or eliminated in said cells. Non-limiting embodiments are described in the following. Which analytical method is suitable also depends on how the cells are altered to achieve a reduction or elimination of the functional expression of the matriptase.

[112] For example, when introducing a gene knock-out into the matriptase gene in order to reduce or eliminate functional expression of matriptase, one can amplify the corresponding DNA section and sequence the amplified DNA in order to confirm that the gene knock-out was introduced into the matriptase gene. The introduction of one or more mutations can be detected e.g. by sequencing. If functional expression of the matriptase is reduced or

eliminated by completely or partly deleting said gene one can detect the deletion on the DNA level, e.g. using suitable amplification based detection methods to detect the deletion (such methods are known to the skilled person).

5 [113] Furthermore, the method can be used in order to select a suitable cell line wherein naturally functional expression of matriptase is reduced or eliminated, e.g. because the overall expression is reduced or eliminated and/or because the activity of the expressed matriptase is reduced or eliminated e.g. due to at least one mutation. Respective cells can be identified using the method according to the seventh aspect as is described herein. E.g.
10 expression of matriptase can be analysed using PCR based techniques such as RT-PCR and/or sequencing. Alternatively or additionally, the activity of matriptase can be analysed using conditioned medium obtained from the cells to be analysed.

[114] According to one embodiment, cells with no or reduced amount of matriptase on the
15 cell surface are selected by flow cytometry. E.g. an antibody or other detection agent that binds matriptase can be used to mark the cells that express matriptase. Respective cells can then be labelled by adding a labelled agent that binds the antibody or other detection agent, or the antibody or other detection agent may be labelled directly. According to one embodiment, the label is a fluorescent label. Thereby, the cells are marked according to their
20 expression level of matriptase which allows to then identify and sort the cells which show reduced or no expression of matriptase, e.g. by fluorescence activated cell sorting (FACS).

[115] According to one embodiment, the expression profile of the vertebrate cells is analyzed to determine whether functional expression of the matriptase gene is reduced or
25 eliminated. For example, the analysis may comprise performing a qualitative or quantitative RT (reverse transcription) PCR in order to detect the presence, absence, amount or length of matriptase mRNA. In addition or alternatively to the analysis of functional matriptase expression in the cell, the selection process may comprise a step of testing whether there is matriptase activity in conditioned medium obtained from said cell. For example, spike-in
30 experiments as described in the examples may be performed. Accordingly, the cell culture medium may be collected from a variety of potential cell lines in order to analyse whether there is matriptase activity in the respectively obtained conditioned medium. E.g. the conditioned medium may be tested for its clipping activity on a polypeptide that is prone to clipping. By using one or more polypeptides that are known to be clipped as standard for
35 testing the clipping activity, it is possible to qualify or quantify the clipping activity of the potential cell lines relative to each other or to a standard cell line.

[116] According to one embodiment the method according to the seventh aspect is for selecting mammalian cells, in particular rodent cells such as hamster cells, preferably CHO
40 cells.

[117] According to one embodiment, the method according to the seventh aspect comprises selecting at least one cell wherein the function of the matriptase is impaired, preferably by reduction or elimination of functional expression of the matriptase gene for

30483-378

38

recombinant expression of a polypeptide of interest. Cells having the respective characteristics are particularly suitable for recombinant expression as shown by the examples. Further embodiments of respective cells are also described in detail above and it is referred to the respective disclosure.

5

[118] Numeric ranges described herein are inclusive of the numbers defining the range. The headings provided herein are not limitations of the various aspects or embodiments of this disclosure which can be read by reference to the specification as a whole. According to one embodiment, subject-matter described herein as comprising certain elements also refers to subject-matter consisting of the respective elements. In particular, the polynucleotides described herein as comprising certain sequences may also consist of the respective sequences. It is preferred to select and combine preferred embodiments described herein and the specific subject-matter arising from a respective combination of preferred embodiments also belongs to the present disclosure.

15

EXAMPLES

[119] The following examples serve to illustrate the present invention without in any way limiting the scope thereof. In particular, the examples relate to preferred embodiments of the present invention.

20

I. Materials and Methods

25 1. Cell culture

[120] Unless described otherwise herein, as cell culture, suspension growing CHO cells derived from CHO-K1 were cultivated in shake flasks in a standard medium and cell culture process as disclosed e.g. in WO2011/134920. Cells were passaged 2 times per week into fresh media and were maintained in logarithmic growth phase throughout the study.

30

The same culture medium that was used for culturing the cells was used as positive control in the examples.

2. Spike-in experiments using conditioned medium

[121] Cells which do not comprise a heterologous polynucleotide encoding a polypeptide of interest were passaged at a density of 2×10^5 viable cells/ml into standard medium (30 ml culture) and grown at 37°C. At the peak of viable cell density (usually 7 or 8 days after passaging; viability still above 95%) cells were removed from the medium. Under these conditions and at this stage of cell growth, the maximum amounts of secreted proteases are expected to be active in the cell culture medium without release of intracellular proteases due to cell death. Also the maximum amount of secreted polypeptide of interest is expected under these conditions if the cells were transfected to express a recombinant polypeptide of interest. For obtaining conditioned medium, the cell cultures were centrifuged for 15 min at

40

90 g (gentle enough that cells are not releasing intracellular proteases due to centrifugation stress). After centrifugation, the supernatant was transferred and passed through a 0.22 µm filter to remove remaining cell particles from the conditioned medium.

5 [122] The polypeptide of interest was added to the so obtained conditioned medium with a final concentration of 0.7 µM and incubated at 37°C with continuous shaking at 500 rpm. The incubation time for each polypeptide of interest was previously determined by removing aliquots periodically in the wild-type sample and analyzing them using SDS-PAGE and Western Blot. A time point at which at least 50% degradation was observed in the wild-type
10 conditioned medium was chosen for each polypeptide of interest. After incubation, all samples of the polypeptide of interest were analyzed by SDS-PAGE usually followed by Western Blot analysis to determine the amount of clipping. In the examples, different therapeutic polypeptides were used as polypeptide of interest in order to analyze clipping.

15 3. SDS-PAGE and Western Blot

[123] Protein samples were diluted in Pierce Lane Marker Reducing Sample Buffer (10 mM DTT, cat. 39000) and boiled for 5 min at 95°C. About 0.1-0.6 µg protein/lane was loaded into a pre-cast 4-12 % Bis-Tris Gel (Invitrogen, cat. NP0322BOX). After electrophoresis, gels were transferred on nitrocellulose membrane (Invitrogen, LC2001) using a Bio-Rad wet
20 electro-blotting system. The polypeptide of interest was detected using an HRP-coupled specific antibody and ECL (Pierce, cat. 32109).

II. Example 1: Clipping of different polypeptides of interest when silencing expression of different proteases by RNA interference (RNAi)

25

[124] First, the proteolytic cleavage site(s) of different recombinant proteins that are prone to clipping were analyzed using different techniques including CE-SDS analysis, mass spectrometry and in silico analysis (data not shown). This analysis combined with further analyses using different protease inhibitors revealed that the majority of tested proteins that
30 are prone to clipping are cleaved by trypsin-like proteases, which is a subfamily of the serine proteases. The RNAi experiment of example 1 was set-up to demonstrate that matriptase is the key protease responsible for clipping and that silencing of the matriptase gene significantly reduces clipping while silencing of other genes encoding different proteases does not reduce clipping of the polypeptide of interest. The siRNAs were designed against
35 the following target mRNAs expressed in CHO cells: 1. Matriptase (MT-SP1; also referred to herein as St14), 2. C1r (complement component activated C1r; also referred to as Ctra), 3. C1s (complement component C1sB; also referred to as Gm5077), 4. Plat (t-plasminogen activator) and 5. Prss35 (protease, serine, 35). These target genes encode trypsin-like proteases that are either secreted or transmembrane proteins and thus could be involved in
40 clipping of recombinantly expressed proteins that are secreted into the cell culture medium. It was confirmed in advance that these proteins are expressed by the CHO cells. Table 2 provides an overview over the respective target genes:

Table 2

Gene symbol	NCBI gene ID (CriGri_1.0 (BioProject: PRJNA72741))	NCBI mRNA Reference Sequence ID (CHO cell line)	Ensembl gene ID (Mus musculus)
St14	100755225	XM_007644507.1; XM_007644511.1 (XM_003495842.1 was replaced by NCBI as a result of standard genome annotation processing)	ENSMUSG00000031995
C1ra/C1r	100768476	XM_003496099.1	ENSMUSG00000055172
Gm5077 (C1s)	100759777	XM_007613477.1 XM_007646821	ENSMUSG00000079343
Plat	100750775	XM_003503315.1	ENSMUSG00000031538
Prss35	100761989	XM_003495638	ENSMUSG00000033491

[125] The sense and antisense sequence of the siRNA against matriptase and the other target proteases are listed in Table 3.

5

Table 3

Target Gene	Sense sequence of siRNA	Antisense sequence of siRNA
Matriptase (MT-SP1, also referred to as St14)	GCAAGAUCACUGUUCGCUUTT (SEQ ID NO: 6)	AAGCGAACAGUGAUCUUGCTG (SEQ ID NO: 7)
C1ra (also referred to as C1r)	GAUGUCUUUUCUAAAAUATT (SEQ ID NO: 8)	UAUUUUGAGAAAAGACAUCAT (SEQ ID NO: 9)
Gm5077 (also referred to as C1s)	CGCUGAACGUGUGAUUAUUTT (SEQ ID NO: 10)	AAUAAUCACACGUUCAGCGGT (SEQ ID NO: 11)
Plat (siRNA sequences were designed based on internal information on the sequence of the transcript for Chinese hamster)	GAAACAAGAUGAAGACAGATT (SEQ ID NO: 12)	UCUGUCUUCAUCUUGUUUCCC (SEQ ID NO: 13)
Prss35	GGCCUUGACUACGACUAUTT (SEQ ID NO: 14)	AUAGUCGUAGUCUAAGGCCGG (SEQ ID NO: 15)

The detailed set-up of the experiment is described in the following.

10 1.1. siRNA transfection

[126] RNAi transfection of CHO-K1 derived cells was performed using Lipofectamine® RNAiMAX Reagent (Life Technology, cat. 13778), following the 'Reverse Transfection Protocol' of the manual MAN0001182, scaled up for 6-well plates. For each mRNA target, the two most efficient concentrations of RNAi duplex molecules were used among 100, 125 and 150 pmol: 125 and 150 pmol for MT-SP1 (St14), 125 and 150 pmol for C1r (C1ra), 100 and 125 pmol for C1s (Gm5077), 125 and 150 pmol for Plat and 100 and 125 pmol for Prss35. RNAi duplex molecules of the two selected concentrations were diluted in 1ml Opti-MEM® (Gibco, cat. 31985-062). As RNAi negative control (siRNA negative control), the Silencer® Negative Control No.1 siRNA (AM4611) was used (at 125 pmol). 5µl Lipofectamine® RNAiMAX Reagent was added in each well. 2ml of a cell suspension with a concentration of 0.5×10^6 cells/ml were added and plates were incubated at 37°C and 10% CO₂. Three days after transfection, cell density was measured, total RNA of 3×10^6 viable

20

cells was extracted using RNeasy Plus Mini Kit (Qiagen, cat. 74134) and conditioned medium was collected from each well. To control the silencing effect on the target genes, mRNA expression was determined by real-time RT-PCR.

5 **Table 4:** Primers and probe sequences to determine protease mRNAs expression

Matriptase/MT-SP1 Forward Primer Sequence	CGCTGAGTACCTGTCCTACGA (SEQ ID NO: 16)
Matriptase/MT-SP1 Reverse Primer Sequence	ACCGTCCAGTGTTACACATGAAC (SEQ ID NO: 17)
Matriptase/MT-SP1 Reporter 1 Sequence	CCAATGACCCATGCCC (SEQ ID NO: 18)
C1ra (C1r) Forward Primer Sequence	ACCTGCAAACAAGGCTACCA (SEQ ID NO: 19)
C1ra (C1r) Reverse Primer Sequence	TGGCAAACAGCTGTGAAGGA (SEQ ID NO: 20)
C1ra (C1r) Reporter 1 Sequence	CAGCACCTGGTTTCC (SEQ ID NO: 21)
Gm5077 (C1s) Forward Primer Sequence	TGCGAGGAGCCATATTACTACATG (SEQ ID NO: 22)
Gm5077 (C1s) Reverse Primer Sequence	GCAGCGCAGCGATACTC (SEQ ID NO: 23)
Gm5077 (C1s) Reporter 1 Sequence	CCGCCGTGTTCTTCAT (SEQ ID NO: 24)
Plat Forward Primer Sequence	AGCTGACATGGAATACTGTGATG (SEQ ID NO: 25)
Plat Reverse Primer Sequence	CCTTTAATTCGAACTGTGGCTGTT (SEQ ID NO: 26)
Plat Reporter 1 Sequence	CCGTGCTCCACCTGC (SEQ ID NO: 27)
Prss35 Forward Primer Sequence	AGGAGAGCACACACAAAGAC (SEQ ID NO: 28)
Prss35 Reverse Primer Sequence	ACACGAGTCCACTGGAAGGA (SEQ ID NO: 29)
Prss35 Reporter 1 Sequence	CCCCGGACCCCTCCTG (SEQ ID NO: 30)

[127] The residual protease gene-expression in relation to the protease gene-expression in the siRNA negative control cells which was set as 100% was as follows: MT-SP1: 125pmol (18.6%) and 150pmol (18.2%), C1r (C1ra): 125pmol (6.3%) and 150pmol (6.7%), C1s (Gm5077): 100pmol (11.9%) and 125pmol (14.4%), Plat: 125pmol (8.3%) and 150pmol (5.1%) and Prss35: 100pmol (18.4%) and 125pmol (14.8%).

1.2. Spike-in experiments in conditioned medium

[128] To collect conditioned medium, 200µl cell culture medium from the 6-well plates was centrifuged (300g for 3min) and the obtained supernatant was used as conditioned medium for spike-in experiments as described in Materials and Methods above. The polypeptide of interest was either a monoclonal IgG antibody (mAb) or a Fc-fusion protein. As described above, a time point at which at least 50% degradation was observed in the wild-type conditioned medium was chosen for each polypeptide of interest. The results led to an incubation time of 3 days for the Fc-fusion protein (wherein, however, degradation was

already far above 50%) and to an incubation time of 7 days for the mAb. After incubation, samples of the polypeptide of interest in conditioned medium were analyzed by SDS-PAGE and Western Blot analysis to determine the amount of clipping as described in Material and Methods.

1.3. Protein clipping: Results of Spike-In experiments

[129] **Fig. 1** shows the Western Blot of the different analyzed polypeptides of interest after incubation as defined above, in the upper panel the mAb and in the lower panel the Fc-fusion protein.

[130] The first lane of the Western Blot in the upper panel shows the mAb after incubation for 7 days in untreated (chemically defined) medium (the same medium was also used for culturing the cells), wherein accordingly, no clipping should occur as the cell culture medium was not in contact with cells and hence, no cellular proteases are present in said medium (positive control (+)). The mAb is displayed as a single strong protein band (marked with an arrow) and no clipping occurred.

[131] The second lane shows the mAb sample after 7 days incubation in the conditioned medium of negative control cells ((-)), which were transfected with a siRNA negative control (no effect on gene expression). As the siRNA negative control has no effect on gene expression, the conditioned medium corresponds in essence to the conditioned medium that is obtained after incubation of the unaltered cells and hence cells which show normal target gene expression and wherein accordingly, the encoded proteases are normally expressed. As can be seen, in addition to the protein band of the intact mAb a second strong protein band appears below that represents the clipped mAb (marked with an arrow). Both protein bands have a similar intensity, thus it can be assumed that about 50% of the mAb was clipped. The same result - significant clipping - is found for the mAb incubated in the cell supernatant of cells in which C1r (C1ra), C1s (Gm5077), Plat and Prss35 were silenced by RNAi (see lanes 5 to 12). That the target genes were effectively silenced was confirmed by real-time RT-PCR. For C1r (C1ra) the strongest mRNA reduction was found with 93.7% and 93.3%. The reduction found for Plat was 91.7% and 94.9%, for C1s (Gm5077) 88.1% and 85.6% and for Prss35 81.6% and 85.2%. Accordingly, all these target genes were successfully silenced by more than 80%. Nevertheless, in all these cases the protein band of the clipped mAb had at least about the same signal intensity as the protein band of the intact mAb. Thus, as in the negative control (-), at least about 50% of the mAb was clipped. Accordingly, downregulation of these protease genes did not reduce clipping of the polypeptide of interest in the conditioned medium.

[132] In contrast, reducing matriptase (MT-SP1) expression in the cells by RNAi (see lanes 3 and 4) considerably reduced clipping of the mAb. In both set-ups in which MT-SP1 was suppressed by 81.4% and 81.8% in comparison to the negative control (remaining MT-SP1 expression 18.6% and 18.2%), the protein band of the clipped mAb is much weaker than the protein band of the intact mAb. This demonstrates that significantly less mAb was clipped and it shows that altering a vertebrate cell to reduce functional expression of the matriptase

gene, here by RNAi, significantly reduces the amount of clipping of a polypeptide of interest in the conditioned cell culture medium obtained from said cells.

[133] Results for the Fc-fusion protein as a further example of a polypeptide of interest are shown in the lower panel of **Fig. 1**. In contrast to the mAb, the comparative sample “(+)” of the Fc-fusion protein contains in addition to the protein band of the intact protein a second strong band corresponding to the Fc portion alone, which is also the main clipping product (both bands are marked with an arrow). The clipped Fc portion is also found in the positive control, because the starting material (Fc-fusion protein produced in CHO cells), contains even after purification a high percentage of the clipped Fc-fusion protein. During the purification process of the Fc-protein, not all of the clipped material can be removed. Therefore, the Fc-fusion protein used in the positive control as starting material already contains significant amounts of the clipped Fc portion as impurity. In the conditioned medium of cells wherein expression of C1r (C1ra), C1s (Gm5077), Plat or Prss35 was silenced, the signal for the intact Fc-fusion protein ranges from weak (Plat) to very weak (Prss35). Accordingly, clipping of the Fc-fusion protein was very high in these cases and reached values up to almost 100 % even though expression of these proteases was suppressed. Therefore, silencing of these protease genes was ineffective in order to reduce clipping. In contrast, in the conditioned medium obtained from cells in which expression of matriptase (MT-SP1) was silenced (only one siRNA concentration was used), about the same intensity of the intact protein as in the positive control (+) is found. Therefore, clipping was significantly reduced. The lane representing the siRNA negative control (-) also comprises intact protein. However, this result is a contamination of the siRNA negative control lane by the positive control which spilled over upon loading.

[134] In summary, this RNAi experiment demonstrates that silencing expression of matriptase in the host cells results in a significant decrease in clipping of different exemplary polypeptides of interest in the conditioned medium obtained from said cells. In contrast, silencing expression of other trypsin-like proteases expressed by the host cells does not have any positive effect on clipping. Example 1 therefore supports the importance of the finding that matriptase is the major protease responsible for clipping of recombinantly expressed and secreted polypeptides of interest. Furthermore, it was found that the cells wherein matriptase was down-regulated by RNAi had the same growth characteristics as the siRNA negative control cells. Therefore, down-regulation of matriptase expression did not affect cell growth.

III. Example 2: Matriptase gene knock-out (KO) in CHO cells leads to a decrease in clipping

A. KO performed with TALEN technology

[135] Nine matriptase (MT-SP1) knock-out cell clones on the basis of CHO-K1 derived cells were generated using TALEN (Transcription Activator-Like Effector Nucleases) technology. For the knockout, matriptase exon 2 was targeted on a region located before the coding region of the transmembrane domain. Exon 2 was chosen because it covers different

alternative splicing variants. Adding frameshift mutations in exon 2 has the advantage that the truncated protein will be short and will be located intracellularly and furthermore, unstable and not toxic for the cells.

5 **2.A.1. Design/production and use of TALENs which are specific for exon 2 of matriptase**

[136] Exon 2 of matriptase and the flanking introns were sequenced in the CHO-K1 derived parental cell line (see Fig. 2, SEQ ID NO: 31).

10 [137] Two truncated TAL FokI targeting matriptase exon 2 were designed. Each TALEN is targeting and binding to 19 nucleotides on either the 5' (on the forward) or the 3' (on the reverse) DNA strand, respectively. The two binding sites are separated by the sixteen nucleotides of the cutting site. Each designed TAL was synthesized and cloned in Gateway® entry vector and subcloned in a pEXP3-DEST_A302 destination vector. Product description and methods are available from Life Technology/GeneArt. Plasmids coding for the 5' (left) or
15 the 3' (right) strand-recognizing TALEN (subcloned in the backbone vector pEXP3-DEST) were produced. pEXP3-DEST contains a T7 promoter upstream of the TALEN coding sequence, allowing in vitro transcription (IVT) of mRNA coding for TAL-FokI.

20 **2.A.2. In-vitro transcription of the TALEN vectors**

[138] TALEN mRNA was produced by in-vitro transcription (IVT) of the TALEN vectors. TALEN vectors were previously linearized with HindIII restriction enzyme (Roche, cat. 10656321001) and purified using isopropanol followed by 70 % ethanol precipitations. mRNA was generated using procedures known by the person skilled in the art. Capped IVT product
25 was purified with NH₄-Ac precipitation and generated mRNA (containing polyA tail and five-prime cap) was purified using the Qiagen RNeasy Micro kit (cat. 74004).

2.A.3. Transfection of TALEN mRNA

[139] Parental CHO-K1 cells in exponential growth phase with viability over 95% were used
30 for transfection. Electroporation (nucleofection) was performed using the Amaxa™, Nucleofector™ Technology according to the instructions of the manufacturer (Lonza). The transfected cells were expanded at day 4 after transfection and single cells separated at day 7 in 30 x 96-well plates. Monoclonality and confluence were controlled with the CloneSelect™ Imager (Genetix).

35

2.A.4. Cel-I-assay and screening strategy

[140] The Cel-I-assay was performed according to the manual of SAFC Biosciences. The Cel-I-assay is a standard assay in order to determine the cutting efficiency. In brief, 3 days after transfection genomic DNA was isolated from the cells and a PCR was performed using
40 the following primers:

Fwd: tttttgcccagtcctggtt (SEQ ID NO: 32)

Rev: ccctttggtctgtcctctga (SEQ ID NO: 33)

[141] The amplification product was denatured and allowed to renature. Then, nuclease S and nuclease S enhancer were added and incubated. The digested product was analyzed. Two smaller bands were present indicating TALEN activity within that region of the genome and therefore, supporting that cells wherein the matriptase gene is altered by mutation were present in the analyzed cell pools. From the most positive cell pools (stronger intensity of the two smaller bands), single cells were sorted in 96 well plates.

[142] Genomic DNA (gDNA) was extracted from each clone in 96 well plates using the Extract-N-Amp™ Blood PCR Kits (cat. XNAB2R, Sigma). The gDNA extracts were used to screen for mutated clones, in a 'CutSite PCR' assay using the Reverse Primer (Rev) shown above (SEQ ID NO: 33) in combination with a primer binding on the cutting site (referred to as Cut. Primer) which had the following sequence:

Cut. Primer: GTGGAGTTTCTGCCTGTGAA (SEQ ID NO: 34)

If in the cutting site region a mutation occurred due to the activities of the TALENs, the Cut. Primer will not bind resulting in that no PCR product is obtained. Clones with no PCR product were analysed via sequencing to determine the introduced mutation. Alternatively or additionally, the Surveyor Mutation Detection assay (Transgenomics, cat. 706025) using the Forward (Fwd) and Reverse (Rev) primers shown above was performed. Clones with mutations were transferred in 125ml shake-flasks and sequenced. In said screening assays clone $\Delta 7/\Delta 15$ was identified. Furthermore, after the first round of transfection and screening using the Cel-I assay, one clone with the genotype wt/ $\Delta 4$ was generated, which had a frameshift mutation in one matriptase allele.

[143] To obtain knock-out clones which comprise frameshift mutations in both alleles, two TALEN mRNA transfection and cloning rounds were performed. To obtain knock-out clones with frameshift mutations in both alleles of MT-SP1, the clone with the genotype wt/ $\Delta 4$ was used in a second TALEN mRNA transfection – cloning round as described above to generate frameshift mutations also in the second allele. In total, nine matriptase knock-out clones (KO-1 to KO-9) containing frameshift mutations in both alleles were generated. The genotypes of the nine clones are disclosed in Table 5. The MT-SP1 exon 2 sequences referring to the wildtype and these mutations are shown in Table 6. Partial sequences of the MT-SP1 product resulting from the mutations are shown in Table 7. The amino acids encoded by exon 2 are highlighted in bold.

[144] According to Table 5 there are four clones, KO-1, KO-4 and KO-7 and KO-9, with the genotype $\Delta 4/\Delta 4$. The matriptase genotype of these clones is identical. KO-1, KO-4 and KO-7 are derived from the same TALEN re-transfected pool, KO-9 is derived from a different one. FISH analysis of clones showed that there are some differences in the karyogram of KO-1, KO-4 and KO-7. Thus, it is assumed that they are not derived from the same mother cell.

[145] A frameshift mutation is inserted into both alleles of the matriptase gene in all clones KO-1 to KO-9. Furthermore, mutated clone $\Delta 7/\Delta 15$ was obtained which has the genotype $\Delta 7/\Delta 15$ and has a frameshift mutation in only one allele ($\Delta 7$). In the second allele, the mutation $\Delta 15$, a deletion of 15 base pairs, results in the deletion of five amino acids in the short intracellular domain of matriptase. However, no frameshift occurs due to the $\Delta 15$ deletion. The affected domain does not seem to be necessary for the catalytic function, as it remains attached to the membrane when the extracellular part comprising the catalytic protease domain is shed. Therefore, it is assumed that matriptase expressed from the $\Delta 15$ allele is still catalytically active. Therefore, in the cell clone $\Delta 7/\Delta 15$, functional expression of matriptase is reduced, but not abolished. It is further assumed that approx. half of the remaining mRNA expressed by said clone contains the frameshift mutation.

Table 5: Mutations in exon 2 of both alleles of the matriptase gene

clone #	Genotype
KO-1	$\Delta 4 / \Delta 4$
KO-2	$\Delta 4 / \Delta 4'$
KO-3	$\Delta 4 / \Delta 17$
KO-4	$\Delta 4 / \Delta 4$
KO-5	$\Delta 4 / \Delta 14$
KO-6	$\Delta 4 / \Delta 4'$
KO-7	$\Delta 4 / \Delta 4$
KO-8	$\Delta 4 / \Delta 11$
KO-9	$\Delta 4 / \Delta 4$
$\Delta 7 / \Delta 15$	$\Delta 7 / \Delta 15$

Table 6: Matriptase gene sequences encoding exon 2 from CHO-K1 derived WT and the different KO clones. Δ indicates a deletion. The number behind the Δ indicates how many nucleotides were deleted. In parentheses and marked in bold, the deleted nucleotides are shown.

WT	aacatgaatggcttggaggagggtgaggagttctgcctgtgaataatccaagaaagtggaagcg aggcccccggcgctgtgtggtgctgtggtcctgctggtcagttcctcttctcactcgtggcttcc tggtgtggcacttctct (SEQ ID NO: 35)
$\Delta 4$	aacatgaatggcttggaggagggtgaggagtt(Δtctg)cctgtgaataatccaagaaagtggaagaa gagaggcccccggcgctgtgtggtgctgtggtcctgctggtcagttcctcttctcactcgtggctggc ttcctggtgtggcacttctct (SEQ ID NO: 36)
$\Delta 4'$ KO-2 nucleotide exchange (c/t)	aacatgaatggcttggaggagggtgaggagtt(Δtctg)tctgtgaataatccaagaaagtggaagaa gagaggcccccggcgctgtgtggtgctgtggtcctgctggtcagttcctcttctcactcgtggctggc ttcctggtgtggcacttctct (SEQ ID NO: 37)
$\Delta 17$	aacatgaatggcttggaggagggtg(Δtggagtttctgcctgtg[#])aataatccaagaaagtggaag

mRNA expression measured in the individual clones by qRT-PCR was compared to the mRNA expression in a wild-type CHO-K1 derived cell. The result of this experiment is shown in Fig. 3. According to Fig. 3, in the cell clones with mutated matriptase, mRNA is only expressed in the range of 5 to 40% compared to the wild-type. This result suggests that expression of the mutated matriptase is reduced. Thus less (non-functional) matriptase protein is expressed by the cells. Also cell clone $\Delta 7/\Delta 15$ shows only low matriptase mRNA expression in the cells.

2.A.6. Spike-in experiments with cell supernatant from the nine matriptase (MT-SP1) knockout clones

[147] To evaluate the effect of the matriptase knock-out, spike-in experiments were performed with five different polypeptides of interest, which are prone to clipping. Conditioned medium of cell cultures from the CHO knockout clones was obtained according to the protocol described under Material and Methods. For the spike-in experiments, the polypeptide of interest was added to the conditioned medium with a final concentration of 0.7 μ M and incubated at 37°C with continuous shaking at 500 rpm. The incubation time was dependent on the type of polypeptide of interest tested. After incubation, samples of the polypeptide of interest in conditioned medium were analyzed by SDS page and Western Blot analysis as described under Material and Methods.

[148] Fig. 4A shows the Western Blot analysis of a spike-in experiment with a monoclonal IgG antibody (mAb) as polypeptide of interest. As reference control, the first lane from the left of the Western Blot (indicated by "(+)") shows the mAb after incubation in chemically defined medium (the same culture medium was also used for obtaining the conditioned medium). No clipping of the product is detected; the intact protein is marked with an arrow. The second lane (WT) shows the monoclonal antibody after 48h of incubation in conditioned medium obtained from CHO-K1 derived cells which normally express matriptase. Here, a second strong band is detected, representing the clipped antibody (marked by an arrow). As the protein band of the clipped antibody is stronger than the protein band of the intact protein, more than 50 % of the protein is clipped. In contrast, after incubation of the monoclonal antibody in the conditioned medium obtained from the nine knockout cell lines KO-1 to KO-9 (1 to 9 in Fig. 4), respectively, no clipped product is detected. Therefore, a knock-out of the matriptase gene in the vertebrate host cell is a significant improvement, because clipping of the mAb in the cell culture medium could be efficiently prevented.

[149] The results of a corresponding spike-in experiment with an Fc-fusion protein are shown in Fig. 4B. The incubation time was 24h. In contrast to the monoclonal antibody, already the positive control in chemical defined medium (+) contained a large portion of clipped protein (marked with an arrow). This is, because the Fc-fusion protein that was added as polypeptide of interest was produced in CHO cells and is heavily clipped in the culture medium from which it is harvested. Not all clipped protein could be removed during the purification process so that the starting material already contained some clipped protein contamination. While in the experiments with the matriptase knockout clones (1 to 9), the amounts of intact protein and clipped protein are comparable to the starting material

incubated in the positive control (+), incubation in the conditioned medium of the unaltered CHO-K1 derived wildtype cells (WT) completely abolished and hence degraded the intact protein. Thus, basically 100% clipping was observed in the conditioned medium from cells which were not altered to impair the function of matriptase.

5

[150] A comparable result is found for a further recombinant therapeutic protein that was incubated in the conditioned media for 1h (**Fig. 4C**). Two glycovariants of the protein are present. Again, in the conditioned medium of the wildtype CHO-K1 cells (WT) more than 50% of the both protein glycovariants are clipped (marked by arrows), while incubation in the conditioned media of the KO-1 to KO-9 cells (1 to 9), wherein the function of the endogenous protease matriptase is impaired due to the gene knock-out, preserved the state of the intact protein as observed after incubation in a chemical defined medium (positive control (+)).

10

[151] The KO clones were also analysed over months in order to analyse whether prevention, respectively reduction of clipping is a stable characteristic of the altered vertebrate cells in which the matriptase gene was knocked out. Accordingly, the spike-in experiment with the mAb was repeated with conditioned medium obtained from the matriptase knock-out cell clones which had been cultivated for 3 month. The result of the experiment, again compared with conditioned medium from CHO wildtype cell as negative control "(WT)" and chemically defined medium as positive control "(+)", is shown in **Fig. 4D**. The Western Blot reveals the exact same protein band pattern as the seen in **Fig. 4A**. The same result was also seen after 3 month for the Fc-fusion protein (data not shown). Hence, it was confirmed that with the KO clones, no clipping appeared after several months. Furthermore, it was found that the KO cell lines grew well and cell growth even improved over 3 month culture.

20

25

[152] The results of the spike-in experiments with two glycosylated viral proteins using conditioned medium of the matriptase knock-out clones KO-1 to KO-3 (1-3) are shown in **Fig. 5** after 24h (D1) and 48h of incubation (D2). In the first lane of the Western Blot (**Fig. 5A**) representing the protein after incubation in chemical defined medium (+), two protein bands are visible. The upper band represents the intact protein, the lower band a clipped version of the protein which makes up about 5 % of the total protein in the sample. In the conditioned medium of CHO-K1 wildtype (WT lanes) a strongly reduced amount of intact protein is present after 24h (D1) of incubation. After 48h (D2) the protein band is almost invisible, i.e. almost all protein was clipped. In addition, the protein band of the clipped protein completely disappeared suggesting that the protein was further degraded. In contrast, no difference in the protein band pattern is found for proteins incubated for 24h or 48h in the conditioned media of the matriptase knockout clones KO-1 to KO-3 (see 1 to 3). Accordingly, with conditioned medium obtained from the matriptase knock-out cell lines, no or significantly reduced proteolytic degradation of the therapeutic proteins was detected. The second viral protein shows a comparable result (**Fig 5B**). Here, the intact protein is represented by the single protein band shown in lane "(+)". Already after 24h of incubation in conditioned medium of CHO-K1 derived wildtype cells (WT, D1) almost no trace of the intact protein can be detected. Instead, two lower protein bands appear, representing clipped versions of the

30

35

40

protein. These protein bands are also seen in the experiment with the medium obtained from matriptase knockout clones KO-1 to KO-3. However, the majority of the viral protein is preserved with all tested knock-out clones even after 48h of incubation.

- 5 [153] Confirming the experiments above, spike-in experiment with five additional Fc-fusion proteins, which are prone to clipping, were performed as described above. The cell free harvest supernatant was captured by Protein A affinity liquid chromatography using 1 mL HiTrap MabSelect Sure (GE Healthcare). A residence time of 2 minutes was applied for equilibration, load and wash. For the elution a residence time of 4 minutes was applied. The
- 10 eluate was titrated to pH 5 with 1 M Tris prior to sterile filtration with a Millex-GV Syringe Filter Unit (0.22 μ m, PVDF, 13 mm; Millipore). The protein concentration was determined using a NanoDrop 2000 spectrophotometer (Thermo Scientific) at 280 nm wavelength. All steps were performed at room temperature. Table 8 shows the mass spectrometry analysis data. As can be seen, all tested Fc-fusion proteins were completely or almost completely
- 15 clipped when expressed in the parental cells. In contrast thereto, clipping was considerably reduced when being expressed in the knock-out cell line.

Table 8: Clipping analysis of different Fc-fusion proteins

Candidates	WT	KO-4
	Clipped Fc-fusion protein [%]	Clipped Fc-fusion protein [%]
Fc-fusion protein 1	100 %	81 %
Fc-fusion protein 2	97 %	48 %
Fc-fusion protein 3	98 %	63 %
Fc-fusion protein 4	100 %	68 %
Fc-fusion protein 5	100 %	67 %

- 20 [154] These examples demonstrate that the altered vertebrate cells according to the present disclosure wherein the function of the endogenous protease matriptase is impaired, present or release no or reduced amounts of functional matriptase into the cell culture medium. Therefore, proteolytic degradation of the recombinant polypeptide of interest that is present in (normally secreted into) the cell culture medium is significantly reduced when
- 25 using these altered cells for producing a polypeptide of interest.

2.A.7. Spike-in experiments with cell supernatant from the matriptase mutant clone $\Delta 7/\Delta 15$

- 30 [155] To evaluate the effect of reduced functional matriptase expression, the spike-in experiments with the Fc-fusion protein and the monoclonal IgG antibody (mAb) were repeated with conditioned medium obtained from the $\Delta 7/\Delta 15$ clone. Conditioned medium of the cell culture was obtained according to the protocol described under Material and Methods. For the spike-in experiment, the polypeptide of interest was added to the conditioned medium with a final concentration of 0.7 μ M and incubated at 37°C with
- 35 continuous shaking at 500 rpm. The Fc-fusion protein was incubated for 2h, the mAb for 24h. After incubation, samples of the polypeptides of interest in conditioned medium were analysed by SDS-PAGE and Western Blot analysis as described under Materials and Methods.

[156] The results of the spike-in experiment with the Fc-fusion protein are shown in **Fig. 6A**. As reference, the first lane of the Western Blot shows the Fc-fusion protein after incubation in chemically defined medium (indicated by "(+)"). The second and third lane ("WT1" and "WT2") shows the Fc-fusion protein after incubation in conditioned medium from two different CHO-K1 derived wild-type cell lines which normally express the unmodified, functional matriptase. As discussed above, the starting material contained the intact Fc-fusion protein as well as clipped Fc-fusion protein as impurity. In the protein samples derived from incubation with conditioned medium from CHO cells with wild-type matriptase ("WT1" and "WT2"), basically no intact Fc-fusion protein is observed. In contrast, incubation in conditioned medium obtained from the $\Delta 7/\Delta 15$ cell clone (" $\Delta 7/\Delta 15$ ") in which functional expression of matriptase is significantly reduced (for details regarding the clone see Example 2.4), clipping is significantly reduced as can be concluded from the strong protein band representing the intact Fc-fusion protein.

[157] **Fig. 6B** shows the Western Blot analysis of a spike-in experiment with the IgG mAb as polypeptide of interest. As reference, the first lane of the Western Blot (indicated by "(+)") shows the mAb after incubation in chemically defined medium. One strong protein band representing the intact antibody is visible. The second and third lane ("WT1" and "WT2") shows the monoclonal antibody after incubation in conditioned medium from two different CHO-K1 derived wild-type cell lines which express the unmodified matriptase. In this second and third lane, an additional strong protein band having a lower molecular weight appears representing the clipped antibody. According to the signal strength, more than 50% of the antibody appears to be clipped. In the fourth lane (" $\Delta 7/\Delta 15$ ") showing the mAb after incubation in conditioned medium from the cell clone $\Delta 7/\Delta 15$, the intensities of the signals are reversed. Thus, although clipping is not completely abolished, it is strongly reduced compared to the matriptase wild-type cell clones. This result confirms that reducing functional expression of matriptase, significantly reduces clipping of the polypeptide of interest in the cell culture medium.

[158] These experiments show that when using a cell clone wherein only one matriptase allele is knocked-out and wherein functional matriptase expression is reduced also leads to a reduced clipping of the polypeptide of interest in the cell culture medium. Thus, it can be concluded that the clipping activity is proportional to the degree of functional matriptase expression.

B. KO performed with ZFN technology

[159] One ZFN (MT-SP1) knock-out cell clone on the basis of CHO parental cells was generated using ZFN (Zinc Finger Nucleases) technology. For the knockout, matriptase exon 2 was targeted on a region located before the coding region of the transmembrane domain (as described in Example 2 A).

2.B.1. Design/production and use of TALENs which are specific for exon 2 of matriptase

[160] A pair ZFNs targeting matriptase exon 2 were designed. One ZFN is targeting and binding to 12 nucleotides, the other one is targeting 18 nucleotides. The two binding sites are separated by the five nucleotides of the cutting site.

2.B.2. In-vitro transcription of the TALEN vectors

[161] ZFN mRNA was produced by in-vitro transcription (IVT) of the ZFN vectors. mRNA was generated using procedures known by the person skilled in the art. Capped IVT product was purified with NH4-Ac precipitation and generated mRNA (containing polyA tail and five-prime cap) was purified using the Qiagen RNeasy Micro kit (cat. 74004).

2.B.3. Transfection of ZFN mRNA

[162] Parental CHO cells in exponential growth phase with viability over 95% were used for transfection. Electroporation (nucleofection) was performed using the Amaxa™, Nucleofector™ Technology according to the instructions of the manufacturer (Lonza).

2.B.4. Cel-I-assay and screening strategy

[163] The Cel-I-assay was performed according to the manual of SAFC Biosciences and a similar screen as described in example 2A.4 was performed.

[164] To obtain a knock-out clone which comprise frameshift mutations in both alleles, two ZFN mRNA transfection and cloning rounds were performed. One matriptase knock-out clones (KO-10) containing frameshift mutations in both alleles was generated ($\Delta 13/\Delta 13$).

2.B.5. Spike-in experiments with cell supernatant from matriptase (MT-SP1) knockout clone

[165] To evaluate the effect of the matriptase knock-out, spike-in experiments were performed with two different polypeptides of interest, which are both prone to clipping. Conditioned medium of cell cultures from the CHO knockout clones was obtained according to the protocol described under Material and Methods. For the spike-in experiments, the polypeptides of interest (one Fc-fusion protein and one recombinant therapeutic protein) was added to the conditioned medium with a final concentration of 0.7 μ M and incubated at 37°C with continuous shaking at 500 rpm. The incubation time was dependent on the type of polypeptide of interest tested. After incubation, samples of the polypeptide of interest in conditioned medium were analyzed by SDS page as described under Material and Methods.

[166] The results are corresponding to the spike-in experiments performed with KO clones generated with TALEN technology. An Fc-fusion protein and a recombinant therapeutic protein as described in example section 2.A.6 were used. While in the experiments with the matriptase knockout clone, the amounts of intact proteins and clipped proteins are comparable to the starting material incubated in the positive control (+), incubation in the conditioned medium of the unaltered CHO-K1 derived wildtype cells (WT) completely abolished and hence degraded the intact proteins.

[167] The KO clones were also analysed over several months in order to analyse whether prevention, respectively reduction of clipping is a stable characteristic of the altered vertebrate cells in which the matriptase gene was knocked out. Accordingly, the spike-in experiment with the Fc-fusion protein was repeated with conditioned medium obtained from the matriptase knock-out cell clones which had been cultivated for 6 months. The result of the experiment confirmed that with the KO clone, no clipping appeared even after several months. Furthermore, it was found that the KO cell line grew well and cell growth even improved over culture time.

IV. Example 3: Recombinant proteins are directly cleaved by matriptase

[168] In order to further demonstrate that matriptase directly cleaves recombinantly expressed and secreted proteins (clipping targets), commercially available proteases were used, namely mouse MT-SP1 and human Htra1. A monoclonal IgG antibody mAb (**Fig. 7A**), an Fc-fusion protein (**Fig. 7B**) and a further recombinant protein (**Fig. 7C**) were incubated for 24h, 2h and 1h, respectively, with the two trypsin-like proteases mouse MT-SP1 and human Htra1, which were added to the chemically defined culture medium. Coincubation was performed at 37°C with continuous shaking at 500 rpm. The polypeptides of interest were used at a concentration of 0.7µM each. Each polypeptide of interest was tested with decreasing amounts of the proteases MT-SP1 and Htra1: Molar ratios of protease/polypeptide of interest from left to right are 1/10, 1/100, 1/1000 for MT-SP1 and 1/3, 1/10 and 1/100 for Htra1. As controls, additional samples of the polypeptide of interest are incubated with conditioned medium from CHO-K1 wildtype derived cells as negative control (lane “(-)”) and a chemically defined medium which was not in contact with cells as positive control (lane “(+)”).

[169] From the Western Blots in **Fig. 7A to C** it is evident that the recombinant Htra1 that was added to the cell culture medium cleaves none of the tested polypeptides of interest. Even at higher Htra1 concentrations, the protein bands look exactly like in the positive control (“(+)”). In contrast, all recombinant polypeptides of interest were significantly clipped even at the lowest concentration of MT-SP1 added. The protein band either resembled that of the negative control (“(-)”) or was even worse. Therefore, significant degrees of clipping occurred under all conditions tested when MT-SP1 was added.

[170] In the experiment with the Fc-fusion protein a concentration dependent effect of the protease MT-SP1 was observed (**Fig. 7B**). While at the lowest concentration of matriptase/MT-SP1 the protein band of the intact Fc-fusion protein at about 85 kDa is only slightly reduced in comparison to the positive control, at higher concentrations of MT-SP1 the protein band at approx. 85 kDa completely disappears. Consequently, the degree of clipping of the Fc-fusion protein correlates with the concentration of matriptase/MT-SP1 in the cell culture medium.

[171] To determine the clipping site induced by the commercial mouse MT-SP1 used in this example, the antibody sample (mAb) with MT-SP1 concentration 1/100 was subjected to mass spectrometry. The identified clipping site was the same as when the mAb was produced by the wildtype CHO cell line. This shows that the clipping site of mouse matriptase on the mAb is the same as the clipping site of the protease responsible for clipping when producing the mAb in CHO cells. These results further confirm that matriptase is the clipping protease.

V. Example 4: Inhibition of the matriptase clipping action by a selective matriptase inhibitor

[172] To further confirm that clipping of the recombinant polypeptide of interest is caused by matriptase as key protease responsible for clipping, spike-in experiments were performed with a specific anti-MT-SP1 Fab fragment which specifically inhibits human matriptase/MT-SP1. The inhibitory Fab structure and binding details on human MT-SP1 are published in Farady et al., 2008 J. Mol. Biol. (2008) 380, 351-360).

[173] The therapeutic polypeptides of interest as clipping targets were in this case a monoclonal IgG antibody (mAb) and a recombinant non-antibody glycoprotein with two glycovariants. The polypeptide of interest was added in a concentration of 0.7 μ M in each case. Table 9 describes the further details of experimental set-up corresponding to the Western Blot results shown in Fig. 8 for the mAb (Fig. 8, upper panel) and the recombinant protein (Fig. 8, lower panel):

Table 9

Lane 1	Marker
Lane 2	Chemically defined medium (+)
Lane 3	Conditioned medium obtained from CHO-K1 derived wildtype cells (-)
Lane 4	Chemically defined medium with 3.45nM recombinant mouse MT-SP1 added
Lane 5	Chemically defined medium with 3.45nM recombinant mouse MT-SP1 and 1 μ M anti-MT-SP1 Fab fragment added
Lane 6	Chemically defined medium with 3.45nM recombinant mouse MT-SP1 and 10 μ M anti-MT-SP1 Fab fragment added
Lane 7	Chemically defined medium with 3.45nM recombinant mouse MT-SP1 and 50 μ M anti-MT-SP1 Fab fragment added
Lane 8	Conditioned medium obtained from CHO-K1 derived wildtype cells
Lane 9	Conditioned medium obtained from CHO-K1 derived wildtype cells and 1 μ M anti-MT-SP1 Fab fragment added
Lane 10	Conditioned medium obtained from CHO-K1 derived wildtype cells and 10 μ M anti-MT-SP1 Fab fragment added

Lane 11	Conditioned medium obtained from CHO-K1 derived wildtype cells and 50µM anti-MT-SP1 Fab fragment added
---------	--

[174] Conditioned medium was prepared as described under Materials and Methods. The incubation time was 24h in case of the monoclonal IgG antibody and 1h for the recombinant non-antibody protein.

5

[175] **Fig. 8** (see upper panel) shows that a strong protein band of the intact mAb is present in the positive control (+) (see lane 2). No clipped mAb is observed. In the negative control (-) (see lanes 3 and 8), a second protein band of higher intensity appears slightly below the intact protein indicating that the monoclonal antibody is significantly clipped in the conditioned medium of CHO-K1 derived wildtype cells. The intact and the clipped mAb are marked with arrows. The same pattern as in the negative control (-) is seen in the chemically defined medium wherein mouse MT-SP1 was added (see lane 4). With respect to the recombinant non-antibody protein (see **Fig. 8**, lower panel), in the positive control (+) (see lane 2), a strong band is seen for the two intact protein glycovariants at about 37 kDa. Those protein bands for the intact protein glycovariants seen in the positive control disappeared in the negative control (-) and new protein bands for the clipped protein glycovariants appeared (see lanes 3 and 8).

[176] With respect to the samples that were incubated in the presence of anti-MT-SP1 Fab, already 1µM of the Fab was sufficient to completely abolish degradation of the mAb by the mouse MT-SP1 (**Fig. 8**, upper panel, lane 5). 1µM of the anti-MT-SP1 Fab also reduced but could not completely prevent clipping of the mAb in conditioned medium (see upper panel, lane 9). Clipping of the mAb was abolished in conditioned medium at a Fab concentration of 10µM and 50µM (**Fig. 8**, upper panel, lanes 10 and 11). For the recombinant protein incubated with recombinant mouse MT-SP1 in chemically defined medium, clipping was reduced already at a concentration of 1µM Fab (see **Fig. 8**, lower panel, lane 5). A further reduction of clipping was seen at 10µM Fab and a complete stop of clipping was seen with 50µM Fab in the chemically defined medium with recombinant mouse MT-SP1 added as well as in the conditioned medium (see **Fig. 8**, lower panel, lanes 6, 7 and 10 and 11). Therefore, the Fab raised against human matriptase also effectively inhibited mouse as well as hamster matriptase. However, higher concentrations were necessary to see full inhibition of the mouse and hamster matriptase. It is assumed that this is attributable to amino acid differences between the human and the mouse and hamster matriptase within the epitope binding site so that the inhibitory Fab against the human matriptase is less potent on mouse and hamster matriptase so that higher concentrations of the Fab are needed to inhibit them.

VI. Example 5: Protein production in matriptase knock-out cell lines

[177] In order to confirm the results from the spike-in experiments, several polypeptide of interest were expressed in a matriptase knock-out cell clone in fed batch cultivation and the clipping of the expressed and secreted polypeptide of interest was analyzed and compared

40

to the results obtained with a corresponding CHO wildtype cell line which endogenously expresses intact matriptase.

5.1. Transfection of the mAb coding vector

5 [178] A CHO wild-type cell line which expresses matriptase (derived from CHO-K1) and the matriptase knock-out CHO cell clone 4 (KO-4, see Table 5) were transfected with an expression vector comprising a polynucleotide encoding a monoclonal antibody (mAb) and two selectable marker genes, namely neo and DHFR. For transfection, cells were grown to exponential phase and 5×10^6 cells were transfected with $3\mu\text{g}$ vector DNA. Five
10 transfection replicates were performed using the wild-type cell line and four transfection replicates were performed using the KO-4 cell line. Selection of cells stably transfected with the vector encoding the protein of interest was performed with G418 (G418 concentration 0.8 mg/ml) followed by two consecutive steps of MTX selection (500nM and $1\mu\text{M}$ MTX). The selection conditions were identical for all pools and both cell lines. At the end of the
15 selection (cell viability > 95 %) cells were frozen. During the selection process, titers of expressed mAb in the medium were determined. The titer in the supernatant of the matriptase knock-out cells was compared to the titer of the CHO wild-type cells. The results demonstrated that the matriptase knockout has no negative effect on the production capacity of the cells but even had a tendency to increase the titer.

5.2. Fed batch production procedure

20 [179] For fed batch production, all four pools of the matriptase knock-out clone KO-4 and three CHO wild-type cell line pools with the highest antibody titer were selected. The frozen cells of the seven pools were thawed at the same time and passaged once before
25 inoculation into the individual fed batch reactors.

[180] The cells were cultivated in fed batch shakers containing a chemical defined medium enriched in amino acids, vitamins and trace elements (Fed Batch Medium). The fed batch cultivation was performed at temperature of 37°C and shaking. During the fed batch cultivation, a feed containing glucose, and amino acids (Fed Batch Feed) was
30 regularly added along the process. During the fed batch cultivation process, samples of the fed batch culture material were regularly collected to determine the viable cell density (vcd) using a Vi-Cell cell viability analyzer (Beckman Coulter) and to determine the protein titers in the cell culture medium. At the end of the fedbatch (day 13 or 14), the cultivation process was stopped. The conditioned medium from the shake-flask (100ml culture) was harvested
35 and filtered using a $0.22\mu\text{m}$ Steriflip filter. The monoclonal antibody was purified from the filtered conditioned medium using protein A affinity liquid chromatography (Protein A Mini Columns, Proteus, cat. PUR008). The purification was performed using the vendor's protocol. The protein concentration of the samples after protein A purification was determined using NanoDropTM system (Thermo Scientific) according to the vendors
40 protocol. Analysis of the fed batch cultures revealed that over the 14 days cultivation period, the titer of expressed mAb in the culture medium of the KO-4 pools was similar or even higher than in the culture medium of CHO wild-type cell pools. Moreover, the cell viability was similar or even higher for the KO-4 cell pools. The KO cells also grew at similar or even

better cell densities. These results indicate that the matriptase knock-out does not negatively influence the protein expression rate and might even enhance it. In addition cell growth was not negatively affected but may even be improved.

5.3. Clipping analysis

- 5 [181] For clipping analysis two parallel approaches were used: Microchip electrophoresis (ME) - SDS and mass spectrometry.

Microchip electrophoresis

- 10 [182] 5µL of the purified protein is used for sample preparation. Depending on the concentration of the samples, the amount of protein is different. For the sample preparation of reducing microchip assay, 5 µL is mixed with 35 µL of reducing sample buffer. Then the mixed solution is incubated in the heat block for 15 min at 70°C. After cooling it down, 70 µL of MilliQ water was added to a final volume of 110 µL. The LabChip® GX II instrument from PerkinElmer was used as system. The used protocol is a revised version from the vendor's protocol. The main difference is the ratio of sample/reducing sample buffer, which is 1:7 in the present protocol for better denaturation instead of 1:3.5 in the vendor's protocol.

- 15 [183] **Fig. 9** shows the result of the microchip electrophoresis. The first three lanes from the left (identified as WT 1, 2 and 3) represent the protein sample composition of the protein
- 20 A purified cell culture from the three productions with CHO wild-type cell pools. In each sample the same protein band pattern is observed. A broad protein band representing the intact heavy chain (HC) of the monoclonal antibody was observed (the size from the Labchip assay is not accurate and usually overestimated). Directly below this protein band a second thinner protein band representing the clipped heavy chain is observed. The
- 25 clipped HC species co-migrates with non-glycosylated HC. In addition, a third protein band representing the light chain is observed at about 29kDa. In the protein samples of the monoclonal antibody produced in the matriptase knock-out cell pools (identified as KO-4 1, 2, 3, and 4), both, the protein band of the intact heavy chain and of the light chain, are at least comparable if not stronger compared to the protein bands found in the samples
- 30 produced by the CHO wild-type cell lines. However, the band of the clipped heavy chain identified by the second arrow from the top in **Fig. 9** is hardly visible in all four antibody samples KO4-1, KO4-2, KO4-3 and KO4-4. This demonstrates that clipping of the mAb is almost abolished when being produced in matriptase knock-out cell lines.

- 35 [184] In order to quantify the result the intensity of the protein bands of clipped antibody heavy chain and intact antibody heavy chain was determined and from this the percentage of clipping was calculated. The result of this analysis is summarized in Table 10. It shows that the amount of clipping of the monoclonal antibody is drastically reduced when using the matriptase knock-out clone as production cell line. According to the analysis, on average
- 40 16.1% of the monoclonal antibody heavy chain produced by the three CHO wild-type cell pools was clipped. In contrast, only 1% or less of clipping occurred on average for the mAb HC produced in the four matriptase knock-out fed batch cultivations. Due to the comigration

of the non-glycosylated HC, exact values cannot be determined based on LabChip® analysis.

Table 10: Clipping Analysis based on protein band intensity

Sample	Clipped mAb Heavy Chain [%]
WT 1	20.2
WT 2	15.6
WT 3	12.4
KO-4 1	0.9
KO-4 2	0.9
KO-4 3	1.0
KO-4 4	0.7

5

Thus, using microchip electrophoresis, merely 1% or less clipping is detected in mAb expressed by matriptase KO cells while 12% to 20% clipping is detected in WT cells expressing the same mAb (clipping detected in HC).

10 **Mass spectrometry analysis**

[185] Samples were deglycosylated and reduced prior to mass spectrometry analysis. From each of the seven protein A purified protein samples an aliquot with a total protein of 50µg was incubated overnight at 37°C with 1.25µl PNGase F at a concentration of 0.3 mg/ml in 50 mM Tris/HCl pH 7.5. The final concentration of the sample was about 0.5 mg/ml. After deglycosylation 88µl of the PNGase F digested samples were reduced by adding 112µl reducing buffer (8M GuHCl, 10µl 1M Tris/HCl pH 7.5 and 2µl 1M DTT) and incubation for one hour at 37°C. The reduction was quenched by addition of 2µl of 10 % trifluoroacetic acid.

15

[186] The so obtained samples with a total volume of 202µl and a concentration of about 0.22mg/ml protein was then used for analysis by combined liquid chromatography and mass spectroscopy. The LC/MS instrument was Waters Synapt G2 system coupled with UPLC (software: MassLynx 4.1). LC method used a flow rate of 0.2 mL/min and buffers MPA (0.1% TFA in water) and MPB (0.09% TFA in acetonitrile). UV detection was performed at 214 nm and 280 nm. Sample temperature was ~5°C. 14µl of the deglycosylated and reduced sample (protein amount 3 µg) were loaded onto a BEH C4 1.7 µm, 2.1 X 100 mm column (Waters). The column temperature was 80°C. Prior to sample loading the column was equilibrated with the buffer MPA. Elution was performed with buffer MPB.

25

30

[187] The eluted protein was then further analyzed by mass spectrometry. Using LC/MS data, the clipping sites were identified and the percentage of clipped species quantified. Table 11 shows the percentage of clipped antibody heavy chains based on LC/MS assay in each protein sample. While in the wild-type samples 20 to 29% clipping was observed, only 2% or less of the antibody heavy chain was clipped when using the matriptase knock-out

35

cells for production. The results calculated based on UV214nm data are slightly higher compared to the ones obtained from reducing LabChip assay (see Table 10). However, the strong reduction of clipping that is achieved when applying the teachings of the present disclosure is again confirmed. The LC/MS results demonstrate a 15-fold decrease in clipping for the mAbs produced in matriptase knock-out cell lines. Analysis of the clipping site revealed that the mAb heavy chain produced in the matriptase knock-out cells is clipped at the same position as the mAb HC produced in the CHO wild-type cell lines. Protein modelling further revealed that the clipping site is located in a very flexible area of the mAb which is exposed to the cell culture medium. Thus, even proteases with a low affinity for this clipping site can easily access and cleave it. These findings may explain the residual clipping events in matriptase knock-out strains.

Table 11: Clipping analysis based on LC/MS assay

Sample	Clipped mAb Heavy Chain [%]
WT 1	28.5
WT 2	23.5
WT 3	19.6
KO-4 1	2.0
KO-4 2	1.5
KO-4 3	1.8
KO-4 4	1.7

5.4. Transfection of WT and KO-4 with further glycoproteins

[188] Further examples were performed to again demonstrate that the introduced matriptase knock-out does not negatively affect the production level and product quality of recombinant glycoproteins. Two glycoproteins different from an antibody (here a Fc-fusion protein as well as one recombinant therapeutic protein) were expressed. The matriptase knock-out cell line KO-4 and the CHO wildtype cell line from which clone KO-4 was derived were transfected with suitable expression vectors. After transfection and selection, titer and protein analytics (e.g. as described above in chapter 5.3) were performed.

[189] The protein titers achieved by the matriptase KO cell line KO-4 were in the same range or higher as in the corresponding WT cell line from which clone KO-4 was derived (see also Table 12). Clipping was again significantly reduced in the proteins such as the Fc-fusion protein obtained from the KO cells vs. the proteins such as the Fc-fusion protein obtained from the WT cells (see also Table 13). This clearly shows that a KO of the matriptase gene reduced proteolytic degradation for a variety of further polypeptides.

Table 12: Expression analysis of a Fc-fusion protein and a recombinant therapeutic protein

Candidates	WT	KO-4
Fc-fusion protein	1.32 g/L	1.24 g/L
Recombinant therapeutic protein	0.59 g/L	1.11 g/L

Table 13: Clipping analysis of a Fc-fusion protein and a recombinant therapeutic protein

Candidates	WT Clipped protein [%]	KO-4 Clipped protein [%]
Fc-fusion protein	12.4 %	4.5 %
Recombinant therapeutic protein	70.2 %	53.9 %

[190] In general, the matriptase KO cell line showed similar or even improved features for the production of glycoproteins. Therefore, the respective cell lines are suitable for the production of proteolytic sensitive polypeptides, as well as for the production of proteolytic insensitive polypeptides. Thus, a universal cell line is provided which simplifies the production of different polypeptides of interest.

VII. Example 6: Upstream process suitability and bioreactor scale-up

[191] In order to show suitability of the matriptase knock-out cell line to be scaled-up for large scale therapeutic production, 12 parental clones derived from two different matriptase KO approaches (ZFN and TALEN technology) were evaluated in a three stage screening approach. CHO wildtype cell line was used for comparison.

6.1. Assessment of parental clone performance

6.1.1. Comparison of parental clone performance during cell expansion

[192] In order to assess the performance of 12 not transfected KO clones (seven ZFN subclones (generated through single cell sorting of the ZFN KO clone described in Example 2b) and five TALEN-derived) and the comparator wild-type cell line during expansion for large-scale production, the seed train was evaluated in shake-flasks using two cell culture expansion media, which differed in the concentration of an essential vitamin. Therefore the clones were inoculated to both expansion media at defined viable cell density and cultured for four days at 36.5 °C. The clones were analysed with respect to the final viable cell densities, average growth rates, as described in Table 14. Along with the results from a suitability assessment of the parental clones using production stage medium (Chapter 6.1.2), the seven best performing clones were selected for further comparison with respect to production capabilities (see Chapter 6.2).

Table 14: Results of parental clone performances during cell expansion

Clone ID	Mue [d ⁻¹] in expansion. medium No. 1	Mue [d ⁻¹] in expansion. medium No. 2	VCD [cells/ml] in expansion. medium No. 1	VCD [cells/ml] in expansion. medium No. 2
ZFN KO1	0.87	0.83	6.75E+06	4.57E+06
ZFN KO1	0.83	0.80	5.50E+06	6.88E+06
ZFN KO3	0.79	0.84	4.06E+06	8.90E+06
ZFN KO4	0.73	0.91	6.00E+06	7.70E+06
ZFN KO5	0.86	0.86	8.99E+06	5.03E+06

ZFN KO6	0.85	0.84	7.23E+06	5.57E+06
ZFN KO7	0.78	0.90	4.08E+06	6.48E+06
TALEN KO-1	0.87	0.85	6.24E+06	6.35E+06
TALEN KO-2	0.88	0.80	4.97E+06	6.01E+06
TALEN KO-3	0.85	0.81	5.55E+06	4.11E+06
TALEN KO-4	0.80	0.86	5.91E+06	7.08E+06
TALEN KO-7	0.84	N.A.	6.19E+06	N.A.
wildtype	0.81	N.A.	6.18E+06	N.A.

6.1.2. Comparison of parental clone performance under production conditions

[193] The 12 parental clones and the reference wild-type cell line were further compared in a shake flask fed-batch production process. The cells from the expansion study described in chapter 6.1.1. were cultivated at 36.5 °C for 14 days using two different cultivation conditions (two different chemically defined production media, inoculation cell densities and feeding regimens). Feeding with two independent feed solutions was carried out throughout the process in pre-defined profile and a temperature shift was applied. Cell densities, viabilities and key metabolites were monitored on a daily basis and used to compare the KO clones to the WT clones with respect to suitability for production. Seven clones were identified which achieved a growth performance similar or superior to the wildtype cell line, while the metabolite profiles were comparable.

6.2. Shake flask screening of transfected pools

[194] The best 7 parental clones as identified in Chapter 6.1 and the CHO wildtype were transfected in triplicate with a monoclonal IgG antibody as polypeptide of interest that was known to be sensitive to proteolytic degradation in the background of the wildtype cell line. The transfected cells were assessed using shake flask cultivation and two different conditions as described in Chapter 6.1.2. Cell growth (viable cell densities, cell viabilities) and product formation was monitored during the cultivation. Cells derived from the TALEN-knock-out approach showed similar or increased cell growth compared to the CHO wildtype cells, whereby ZFN-derived pools showed slightly lower cell growth during exponential phase compared to the TALEN-derived pools. However, cell viability at end of cultivation was comparable for all clones. Volumetric productivity at end of cultivation was on average 19 % higher for TALEN and 9 % higher for zinc-finger-nuclease pools compared to the productivity of the CHO wildtype pools.

[195] At the end of the cultivation (day 14), the monoclonal IgG antibody was analysed for integrity. Therefore, the supernatants were harvested by centrifugation and sterile filtered. The monoclonal IgG antibody was captured from the supernatants and analyzed by CE-SDS (Capillary Electrophoresis-Sodium Dodecyl Sulfate) for proteolytic degradation and by CEC (cation exchange chromatography) to assess charge variant distribution. All pools derived from knock-out clones showed a similarly low level of ca. 1% of polypeptide clipping, whereas the product derived from the CHO wild-type was cleaved at around 21%.

6.3. Bioreactor screening of parental subclones and transfected pools

[196] The four best performing parental clones and transfected pools as identified in chapters 6.1. and 6.2. were selected for in-depth characterization using the controlled cultivation conditions of 7L glass bioreactors. The cultivation process was essentially as described in Chapter 6.1.2., though only the preferred cultivation conditions as identified in Chapter 6.2 was used. A slightly lower growth was seen for the transfected pools compared to the non-transfected parental clones, which is a known phenomenon ascribed to the metabolic burden caused by the polypeptide expression. However, no significant differences in cell growth behavior were found compared to the WT cell line. Metabolites such as lactate and ammonium were within the common ranges and similar to the wildtype cell line.

[197] The product quality was analytically assessed for purity by SEC, CEC and CE-SDS and N-glycosylation by a proprietary method. The results showed that all KO clones achieved comparable product quality in terms of aggregation and degradation products, charge variant distribution and glycosylation pattern. CE-SDS reduced results showed that clipped species are presents for all KO subclones at a low level of 0.7 %. All KO clones not only showed similar or preferred growth characteristics compared the reference wildtype cell line and but also produced polypeptide of consistently better quality.

20

CLAIMS:

1. An isolated vertebrate cell suitable for recombinant expression of a polypeptide of interest, wherein the vertebrate cell is altered to impair the effect of matriptase, wherein the effect of matriptase is impaired because functional
5 expression of the matriptase gene is reduced or eliminated in said cell by gene knock-out, gene mutation, gene deletion, gene silencing or a combination of any of the foregoing, and wherein the vertebrate cell comprises at least one heterologous polynucleotide encoding a polypeptide of interest, wherein the vertebrate cell secretes the polypeptide of interest.
- 10 2. The isolated vertebrate cell according to claim 1, wherein the genome of the vertebrate cell is altered to impair the function of the endogenous protease matriptase, and/or wherein at least one or all copies of the matriptase gene are deleted or functionally inactivated.
3. The isolated vertebrate cell according to claim 2, having one or more
15 of the following features
 - a) the vertebrate cell comprises one or more mutations in at least one copy or all copies of the matriptase gene to provide a non- or less-functional expression product; and/or
 - b) the vertebrate cell comprises one or more mutations in the
20 promoter, in the 5'UTR, the 3'UTR and/or other regulatory elements of the matriptase gene.
4. The isolated vertebrate cell according to claim 3, wherein said one or more mutations are comprised in a coding region of the matriptase gene and result in a non- or less functional expression product, wherein optionally the one or more
25 mutations are comprised in a polynucleotide sequence of the matriptase gene that encodes exon 2 of matriptase or wherein the one or more mutations are comprised in a polynucleotide sequence of the matriptase gene that encodes at least part of the catalytic domain of matriptase whereby a non- or less functional expression product is obtained.

5. The isolated vertebrate cell according to any one of claims 1 to 4, wherein the unaltered endogenous matriptase shares at least 70 %, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to one or more of the amino acid sequences shown in SEQ ID NO: 1 to 5 as matriptase reference protein and has the same proteolytic activity as said matriptase reference protein.
6. The isolated vertebrate cell according to any one of claims 1 to 5, wherein the vertebrate cell has one or more of the following characteristics:
- a) the vertebrate cell is a mammalian cell;
 - 10 b) the vertebrate cell is a human cell;
 - c) the vertebrate cell is a rodent cell;
 - d) the vertebrate cell is a hamster cell;
 - e) the vertebrate cell is:
 - (i) a rodent cell,
 - 15 (ii) a human cell selected from the group consisting of a HEK293 cell, a MCF-7 cell, a PerC6 cell, a CAP cell, human hematopoietic cells and a HeLa cell, or
 - (iii) a monkey cell selected from the group consisting of a COS cell, a COS-1 cell, a COS- 7 cell and a Vero cell,
 - 20 f) the vertebrate cell is a CHO cell; and/or
 - g) the vertebrate cell is a mammalian cell provided as cell clone or cell line.
7. The isolated vertebrate cell according to claim 1, wherein the vertebrate cell is a CHO cell which comprises one or more frame-shift mutations in exon 2 of one or both alleles of the matriptase gene.
- 25 8. The isolated vertebrate cell according to any one of claims 1 to 7, wherein the polypeptide of interest has one or more of the following characteristics:

- a) it is a therapeutically active or diagnostic polypeptide;
- b) it is susceptible to clipping by proteases;
- c) it comprises at least one clipping site for matriptase;
- d) it is a glycopolypeptide; and/or

5 e) it is selected from the group consisting of glycoproteins, antibodies, non-IgG proteins, Fc-fusion proteins, Fab fragments, protein complexes, peptidases, signal peptides, nanobodies, growth factors, hormones, cytokines, blood factors and enzymes.

9. The isolated vertebrate cell according to any one of claims 1 to 8,
10 wherein the at least one heterologous polynucleotide encoding the polypeptide of interest is integrated into the genome of said cell and wherein optionally, at least one heterologous polynucleotide encoding a selectable marker or reporter polypeptide is additionally integrated into the genome of said cell.

10. The isolated vertebrate cell according to any one of claims 1 to 9,
15 wherein the at least one heterologous polynucleotide encoding the polypeptide of interest is comprised in an expression cassette.

11. The isolated vertebrate cell according to any one of claims 1 to 10, wherein impairing the effect of matriptase in said cell reduces or eliminates clipping of the secreted polypeptide of interest.

20 12. A method for producing a vertebrate cell according to any one of claims 1 to 11, comprising altering a vertebrate cell to impair the effect of matriptase by reducing or eliminating functional expression of the matriptase gene in said cell by gene knock-out, gene mutation, gene deletion, gene silencing or a combination of any of the foregoing, and introducing a polynucleotide encoding a polypeptide of
25 interest, wherein said polypeptide of interest is secreted by the vertebrate cell.

13. A method for recombinantly producing a polypeptide of interest, comprising

(a) culturing vertebrate host cells according to any one of claims 1 to 11 under conditions that allow for the expression and secretion of the polypeptide of interest into the cell culture medium;

5 and
(b) isolating the polypeptide of interest from the cell culture medium;

(c) optionally processing the isolated polypeptide of interest.

14. A method for selecting a host cell which recombinantly expresses a polypeptide of interest, comprising

10 (a) providing vertebrate cells according to any one of claims 1 to 11 as host cells; and

(b) selecting one or more host cells expressing the polypeptide of interest;

wherein optionally stage (a) comprises transfecting vertebrate cells in which the function of the endogenous protease matriptase is impaired with at least one
15 polynucleotide encoding a polypeptide of interest to provide vertebrate host cells according to any one of claims 1 to 11.

15. The method according to claim 14, having one or more of the following characteristics

20 a) the vertebrate cells are mammalian cells;
b) said host cells provided in stage (a) additionally comprise at least one heterologous polynucleotide encoding a selectable marker and stage (b) comprises culturing said plurality of host cells under conditions selective for the selectable marker;

25 c) polynucleotides are introduced into the vertebrate cells by transfecting one or more expression vectors;

d) stage (b) comprises one or multiple selection steps; and/or

e) stage (b) comprises performing a flow cytometry based selection.

16. A method for selecting a vertebrate cell to be used for recombinant production of a secreted polypeptide of interest, comprising:

analyzing if the endogenous protease matriptase is functionally
5 expressed in the vertebrate cell; and

selecting a vertebrate cell in which the effect of such endogenous matriptase is impaired by reduction or elimination of functional expression of the matriptase gene by gene knock-out, gene mutation, gene deletion, gene silencing or a combination of any of the foregoing;

10 wherein impairment of such endogenous matriptase is indicative that the vertebrate cell is suitable for use in recombinant production of the secreted polypeptide of interest.

17. The method of claim 16, wherein the vertebrate cell has one or more of the following characteristics:

15 a) the vertebrate cell is a mammalian cell;

b) the vertebrate cell is a human cell;

c) the vertebrate cell is a rodent cell;

d) the vertebrate cell is a hamster cell;

e) the vertebrate cell is:

20 (i) a rodent cell,

(ii) a human cell selected from the group consisting of a HEK293 cell, a MCF-7 cell, a PerC6 cell, a CAP cell, human hematopoietic cells and a HeLa cell, or

(iii) a monkey cell selected from the group consisting of a COS cell, a
25 COS-1 cell, a COS- 7 cell and a Vero cell,

f) the vertebrate cell is a CHO cell; and/or

g) the vertebrate cell is a mammalian cell provided as cell clone or cell line.

18. The method of claim 16, wherein the selected cell is a vertebrate cell according to any one of claims 1 to 9.

5 19. Use of a vertebrate cell for recombinant production of a polypeptide of interest that is secreted from the vertebrate cell, wherein the used cell is altered to impair the effect of the endogenous protease matriptase, wherein the effect of matriptase is impaired because functional expression of the matriptase gene is reduced or eliminated in said cell by gene knock-out, gene mutation, gene deletion,
10 gene silencing or a combination of any of the foregoing.

20. The use of claim 19, wherein the vertebrate cell is a cell according to any one of claims 1 to 11.

Fig. 1





Fig. 2

ttttttgcccagtccttggttttcttcctccaaactcacaggtgtcactgaacctccgggctgggaagctattgtcggggagcgtctcttgagtcctgaaa
tatctttctgttttagaacatgaatggcttttgaggagggtgtggagttttcttgccctgtgaataaatgccaaagaaagtgggagaagagcggagggccccggcg
ctgtgtggtgcttgtggtcctgtggtcagtttccctctttctctcactctgtggctgggttccctctgtggtggcacttccctctctgtgagtacagtgggg
gctgtgggaggcgacagaggggtagtgtctctctctctctctcagaggacagaccaaagg

Fig. 3






Fig. 4

Fig. 5




Fig. 6

A





Fig. 7






Fig. 8

Fig. 9

