




(86) Date de dépôt PCT/PCT Filing Date: 2003/07/14
(87) Date publication PCT/PCT Publication Date: 2004/01/22
(45) Date de délivrance/Issue Date: 2014/02/04
(85) Entrée phase nationale/National Entry: 2004/12/29
(86) N° demande PCT/PCT Application No.: US 2003/021923
(87) N° publication PCT/PCT Publication No.: 2004/006846
(30) Priorités/Priorities: 2002/07/15 (US60/396,269);
2003/02/13 (US60/447,212)

(51) Cl.Int./Int.Cl. *C07D 239/93* (2006.01),
A61K 31/517 (2006.01), *A61P 35/00* (2006.01),
C07D 239/94 (2006.01), *C07D 401/06* (2006.01),
C07D 401/12 (2006.01), *C07D 403/06* (2006.01),
C07D 403/12 (2006.01), *C07D 405/12* (2006.01),
C07D 413/12 (2006.01), *C07D 417/12* (2006.01),
C07D 451/06 (2006.01), ...

(72) Inventeurs/Inventors:
RICE, KENNETH D., US;
ANAND, NEEL KUMAR, US;
BUSSENIUS, JOERG, US;
COSTANZO, SIMONA, US;
KENNEDY, ABIGAIL R., US; ...

(54) Titre : MODULATEURS DE KINASE DE TYPE RECEPTEUR ET PROCEDES D'UTILISATION
(54) Title: RECEPTOR-TYPE KINASE MODULATORS AND METHODS OF USE



(57) Abrégé/Abstract:

The present invention provides compounds for modulating receptor kinase activity, particularly ephrin and EGFR, and methods of treating diseases mediated by receptor kinase activity utilizing the compounds and pharmaceutical compositions thereof. Diseases mediated by receptor kinase activity include, but are not limited to, diseases characterized in part by abnormal levels of cell proliferation (i.e. tumor growth), programmed cell death (apoptosis), cell migration and invasion and angiogenesis associated with tumor growth. Compounds of the invention include "spectrum selective" kinase modulators, compounds that inhibit, regulate and/or modulate signal transduction across subfamilies of receptor-type tyrosine kinases, including ephrin and EGFR.

(51) Cl.Int./Int.Cl. (suite/continued) *C07D 471/04* (2006.01), *C07D 487/04* (2006.01), *C07D 493/04* (2006.01),
C07D 498/04 (2006.01)

(72) Inventeurs(suite)/Inventors(continued): KIM, ANGIE I., US; PETO, CSABA J., US; TSANG, TSZE H., US;
BLAZEY, CHARLES M., US

(73) Propriétaires(suite)/Owners(continued):SYMPHONY EVOLUTION, INC., US

(74) Agent: BENNETT JONES LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 January 2004 (22.01.2004)

PCT

(10) International Publication Number
WO 2004/006846 A3

(51) International Patent Classification⁷: **C07D 239/93**,
239/94, 401/06, 401/12, 403/06, 403/12, A61K 31/517,
A61P 35/00

Charles, M. [US/US]; 201 Cornwall Street, #3, San
Francisco, CA 94118 (US).

(21) International Application Number:
PCT/US2003/021923

(74) Agents: **GRIEDEL, Brian** et al.; Exelixis, Inc., P.O. Box
511, 170 Harbor Way, South San Francisco, CA 94083-
0511 (US).

(22) International Filing Date: 14 July 2003 (14.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/396,269 15 July 2002 (15.07.2002) US
60/447,212 13 February 2003 (13.02.2003) US

(71) Applicant (*for all designated States except US*): **EX-
ELIXIS, INC.** [US/US]; P.O. Box 511, 170 Harbor Way,
South San Francisco, CA 94083-0511 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RICE, Kenneth,
D.** [US/US]; 802 Autumn Lane, Mill Valley, CA 94941
(US). **ANAND, Neel, Kumar** [GB/US]; 2713 South
Norfolk Street, Apt.#206, San Mateo, CA 94403 (US).
BUSSENIUS, Joerg [DE/US]; 1042 Helm Lane, Foster
City, CA 94404 (US). **COSTANZO, Simona** [IT/US]; 51
Jennings Lane, Atherton, CA 94027 (US). **KENNEDY,
Abigail, R.** [US/US]; 325 West Lowell Avenue, Tracy,
CA 95376 (US). **KIM, Angie, I.** [US/US]; 1000 Foster
City Boulevard #4305, Foster City, CA 94404 (US).
PETO, Csaba, J. [HU/US]; 965 Shorepoint Court, #305,
Alameda, CA 94501 (US). **TSANG, Tsze, H.** [US/US];
8367 Kent Drive, El Cerrito, CA 94530 (US). **BLAZEY,**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments*

(88) Date of publication of the international search report:

15 July 2004

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: RECEPTOR-TYPE KINASE MODULATORS AND METHODS OF USE

(57) Abstract: The present invention provides compounds for modulating receptor kinase activity, particularly ephrin and EGFR, and methods of treating diseases mediated by receptor kinase activity utilizing the compounds and pharmaceutical compositions thereof. Diseases mediated by receptor kinase activity include, but are not limited to, diseases characterized in part by abnormal levels of cell proliferation (i.e. tumor growth), programmed cell death (apoptosis), cell migration and invasion and angiogenesis associated with tumor growth. Compounds of the invention include "spectrum selective" kinase modulators, compounds that inhibit, regulate and/or modulate signal transduction across subfamilies of receptor-type tyrosine kinases, including ephrin and EGFR.



WO 2004/006846 A3

RECEPTOR-TYPE KINASE MODULATORS AND METHODS OF USE

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention relates to compounds for modulating protein kinase enzymatic activity for modulating cellular activities such as proliferation, differentiation, programmed cell death, migration and chemoinvasion. Even more specifically, the invention relates to quinazolines which inhibit, regulate and/or modulate kinase receptor signal transduction pathways related to the changes in cellular activities as mentioned above, compositions which contain these compounds, and methods of using them to treat kinase-dependent diseases and conditions.

Summary of Related Art

[0002] Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms.

[0003] Protein kinases are enzymes that catalyze the phosphorylation of proteins, in particular, hydroxy groups on tyrosine, serine and threonine residues of proteins. The consequences of this seemingly simple activity are staggering; cell differentiation and proliferation; i.e., virtually all aspects of cell life in one-way or another depend on protein kinase activity. Furthermore, abnormal protein kinase activity has been related to a host of disorders, ranging from relatively non-life threatening diseases such as psoriasis to extremely virulent diseases such as glioblastoma (brain cancer).

[0004] Protein kinases can be categorized as receptor type or non-receptor type. Receptor-type tyrosine kinases have an extracellular, a transmembrane, and an intracellular domain, while non-receptor type tyrosine kinases are wholly intracellular.

[0005] Receptor-type tyrosine kinases are comprised of a large number of transmembrane receptors with diverse biological activity. In fact, about twenty different subfamilies of receptor-type tyrosine kinases have been identified. One tyrosine kinase subfamily, designated the HER subfamily, is comprised of EGFR (HER1), HER2, HER3, and HER4.

Ligands of this subfamily of receptors identified so far include epithelial growth factor, TGF-alpha, amphiregulin, HB-EGF, betacellulin and heregulin. Another subfamily of these receptor-type tyrosine kinases is the insulin subfamily, which includes INS-R, IGF-IR, and IR-R. The PDGF subfamily includes the PDGF-alpha and beta-receptors, CSFIR, c-kit and FLK-II. Additionally there is the FLK family, which is comprised of the kinase insert domain receptor (KDR), fetal liver kinase-1 (FLK-1), fetal liver kinase-4 (FLK-4) and the fms-like tyrosine kinase-1 (flt-1). The PDGF and FLK families are usually considered together due to the similarities of the two groups. For a detailed discussion of the receptor-type tyrosine kinases, see Plowman et al., DN&P 7(6): 334-339, 1994.

[0006] The non-receptor type of tyrosine kinases is also comprised of numerous subfamilies, including Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack, and LIMK. Each of these subfamilies is further sub-divided into varying receptors. For example, the Src subfamily is one of the largest and includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr, and Yrk. The Src subfamily of enzymes has been linked to oncogenesis. For a more detailed discussion of the non-receptor type of tyrosine kinases, see Bolen, Oncogene, 8:2025-2031 (1993).

[0007] Since protein kinases and their ligands play critical roles in various cellular activities, deregulation of protein kinase enzymatic activity can lead to altered cellular properties, such as uncontrolled cell growth associated with cancer. In addition to oncological indications, altered kinase signaling is implicated in numerous other pathological diseases. These include, but are not limited to: immunological disorders, cardiovascular diseases, inflammatory diseases, and degenerative diseases. Therefore, both receptor and non-receptor protein kinases are attractive targets for small molecule drug discovery.

[0008] One particularly attractive goal for therapeutic use of kinase modulation relates to oncological indications. For example, modulation of protein kinase activity for the treatment of cancer has been demonstrated successfully with the FDA approval of Gleevec® (imatinib mesylate, produced by Novartis Pharmaceutical Corporation of East Hanover, NJ) for the treatment of Chronic Myeloid Leukemia (CML) and gastrointestinal stroma cancers (GIST). Gleevec is a c-Kit and Abl kinase inhibitor.

[0009] Modulation (particularly inhibition) of cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A. Drug Disc Technol

2001 6, 1005-1024), is an attractive goal for development of small-molecule drugs. Anti-angiogenic therapy represents a potentially important approach for the treatment of solid tumors and other diseases associated with dysregulated vascularization, including ischemic coronary artery disease, diabetic retinopathy, psoriasis and rheumatoid arthritis. As well, cell antiproliferative agents are desirable to slow or stop the growth of tumors.

[0010] Inhibition of EGF, VEGF and ephrin signal transduction will prevent cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A. Drug Disc Technol 2001 6, 1005-1024). VEGF receptors are previously described targets for small molecule inhibition.

[0011] The Eph receptors comprise the largest family of receptor tyrosine kinases and are divided into two groups, EphA and EphB, based on their sequence homology. The ligands for the Eph receptors are ephrin, which are membrane anchored. Ephrin A ligands bind preferentially to EphA receptors whilst ephrin B ligands bind to EphB receptors. Binding of ephrin to Eph receptors causes receptor autophosphorylation and typically requires a cell-cell interaction since both receptor and ligand are membrane bound.

[0012] Overexpression of Eph receptors has been linked to increased cell proliferation in a variety of tumors (Zhou R 1998 Pharmacol Ther. 77, 151-181; Kiyokawa E, Takai S, Tanaka M et al 1994 Cancer Res 54, 3645-3650; Takai N Miyazaki T, Fujisawa K, Nasu K and Miyakawa. 2001 Oncology reports 8, 567-573). The family of Eph receptor tyrosine kinases and their ephrin ligands play important roles in a variety of processes during embryonic development and also in pathological angiogenesis and potentially metastasis. Therefore modulation of Eph receptor kinase activity should provide means to treat or prevent disease states associated with abnormal cell proliferation such as those described above.

[0013] The epidermal growth factor receptor (EGFR, HER1, erbB1) is part of a family of plasmam membrane receptor tyrosine kinases that control cellular growth, proliferation and apoptosis. The ligand for EGFR is the epidermal growth factor and dysregulation of the EGFR signal transduction pathway has been implicated in tumorigenesis and cancer progression thus making it a clinically relevant target for novel anticancer treatments Drevs J et al 2003 Curr Drug Targets 4, 113-121; Ciardiello F and Tortora G. 2001 Clin. Cancer Res. 7, 2958-2970; Thomas M. 2002 Semin Onc. Nurs. 18, 20-27).

[0014] EGFR is overexpressed in different human cancers especially non-small cell lung cancer and glioblastomas. In these cancers, EGFR overexpression is commonly

associated with advanced disease and poor prognosis (Baselga J et al 1999 Semin. Oncol. 26, 78-83).

[0015] Accordingly, the identification of small-molecule compounds that specifically inhibit, regulate and/or modulate the signal transduction of tyrosine kinases, particularly ephrin and EGFR receptor kinases, is desirable as a means to treat or prevent disease states associated with abnormal cell proliferation and is an object of this invention.

SUMMARY OF THE INVENTION

[0016] In one aspect, the present invention provides compounds for modulating kinase activity, particularly ephrin receptor kinase (from herein denoted “ephrin”) activity and/or EGFR activity; methods of treating diseases mediated by such activity utilizing the compounds and pharmaceutical compositions thereof. Diseases mediated by ephrin include, but are not limited to, diseases characterized in part by abnormal cell migration and invasion and angiogenesis associated with tumor growth. Diseases mediated by EGF driven signal transduction include, but are not limited to, diseases characterized by abnormal levels of cell proliferation (i.e. tumor growth) and programmed cell death (apoptosis). Diseases mediated by both ephrin and EGF include diseases characterized by both abnormal cell proliferation and angiogenesis which are associated with tumor growth.

[0017] Inhibitors that are selective for a particular kinase, for example ephrin and EGFR, are included in this invention. However, another aspect of the invention are compounds that inhibit, regulate and/or modulate the signal transduction receptor tyrosine kinase families, family members, or otherwise related sets of kinases. Such related kinase families may include receptor-type tyrosine kinases of the HER, FLK and insulin subfamilies, which demonstrate similarity in both structure and broad biochemical function. Thus quinazolines of the invention include “spectrum selective” kinase modulators. “Spectrum selective” kinase modulators are defined as quinazolines of the invention that inhibit, regulate and/or modulate signal transduction across various subfamilies of receptor-type tyrosine kinases including those of ephrin and EGFR receptor tyrosine kinase subfamilies.

[0018] In another aspect, the invention provides methods of screening for modulators of receptor tyrosine kinase activity, for example activity of ephrin and EGFR. The methods comprise combining a composition of the invention, a receptor tyrosine kinase, and at

least one candidate agent and determining the effect of the candidate agent on the kinase activity.

[0019] In yet another aspect, the invention also provides pharmaceutical kits comprising one or more containers filled with one or more of the ingredients of pharmaceutical compounds and/or compositions of the present invention, including, one or more tyrosine receptor kinase activity modulators as described herein. Such kits can also include, for example, other compounds and/or compositions (for example, diluents, permeation enhancers, lubricants, and the like), a device(s) for administering the compounds and/or compositions, and written instructions in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflect approval by the agency of manufacture, use or sale for human administration.

[0020] In still yet another aspect, the invention also provides a diagnostic agent comprising a compound of the invention and, optionally, pharmaceutically acceptable adjuvants and excipients.


[0021] These and other features and advantages of the present invention will be described in more detail below with reference to the associated drawings.

DETAILED DESCRIPTION OF THE INVENTION

[0022] The compositions of the invention are used to treat diseases associated with abnormal and or unregulated cellular activities. Disease states which can be treated by the methods and compositions provided herein include, but are not limited to, cancer (further discussed below), immunological disorders such as rheumatoid arthritis, graft-host diseases, multiple sclerosis, psoriasis; cardiovascular diseases such as arteriosclerosis, myocardioinfarction, ischemia, stroke and restenosis; other inflammatory and degenerative diseases such as interbowel diseases, osteoarthritis, macular degeneration, diabetic retinopathy.

[0023] It is appreciated that in some cases the cells may not be in a hyper- or hypo-proliferative and/or migratory state (abnormal state) and still require treatment. For example, during wound healing, the cells may be proliferating "normally", but proliferation and migration enhancement may be desired. Alternatively, reduction in "normal" cell proliferation and/or migration rate may be desired.

[0024] The present invention comprises a compound for modulating tyrosine kinase activity of Formula I,



or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein,

R^1 is C_1 - C_3 alkyl optionally substituted with between one and three R^{50} substituents;

R^2 is selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -N(R³)R⁴, -S(O)₀₋₂R⁴, -SO₂N(R³)R⁴, -CO₂R³, -C(=O)N(R³)R⁴, -N(R³)SO₂R⁴, -N(R³)C(=O)R³, -N(R³)CO₂R⁴, -C(=O)R³, optionally substituted lower alkyl, optionally substituted lower alkenyl, and optionally substituted lower alkynyl;

R^3 is -H or R^4 ;

R^4 is selected from optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, and optionally substituted lower heterocyclylalkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form an optionally substituted five- to seven-membered heterocyclyl, said optionally substituted five- to seven-membered heterocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P;

q is zero to five;

Z is selected from -OCH₂-, -O-, -S(O)₀₋₂-, -N(R⁵)CH₂-, and -NR⁵-;

R^5 is -H or optionally substituted lower alkyl;

M^1 is -H, C_1 - C_8 alkyl- L^2 - L^1 - optionally substituted by R^{50} , $G(CH_2)_{0-3}$, or $R^{53}(R^{54})N(CH_2)_{0-3}$; wherein G is a saturated five- to seven-membered heterocyclyl containing one or two annular heteroatoms and optionally substituted with between one and three R^{50} substituents; L^1 is -C=O- or -SO₂-; L^2 is a direct bond, -O-, or -

NH-; and R^{53} and R^{54} are independently C_1 - C_3 alkyl optionally substituted with between one and three R^{50} substituents;

M^2 is a saturated or mono- or poly-unsaturated C_3 - C_{14} mono- or fused-polycyclic hydrocarbyl optionally containing one, two, or three annular heteroatoms per ring and optionally substituted with between zero and four R^{50} substituents; and

M^3 is $-NR^9$ -, $-O$ -, or absent;

M^4 is $-CH_2$ -, $-CH_2CH_2$ -, $-CH_2CH_2CH_2$ -, or absent;

R^9 is $-H$ or optionally substituted lower alkyl;

R^{50} is $-H$, halo, trihalomethyl, $-OR^3$, $-N(R^3)R^4$, $-S(O)_{0-2}R^4$, $-SO_2N(R^3)R^4$, $-CO_2R^3$, $-C(=O)N(R^3)R^4$, $-C(=NR^{25})N(R^3)R^4$, $-C(=NR^{25})R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(=O)R^3$, optionally substituted alkoxy, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, and optionally substituted lower heterocyclylalkyl; or

two of R^{50} , when taken together on the same carbon are oxo; or

two of R^{50} , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P; and


R^{25} is selected from $-H$, $-CN$, $-NO_2$, $-OR^3$, $-S(O)_{0-2}R^4$, $-CO_2R^3$, optionally substituted lower alkyl, optionally substituted lower alkenyl, and optionally substituted lower alkynyl.

[0025] Embodiment [0025]: In one example, the compound is according to Formula I, wherein M^4 is $-CH_2$ -.

[0026] Embodiment [0026]: In another example, the compound is according to Embodiment [0025], wherein Z is $-NR^5$ -.

[0027] Embodiment [0027]: In another example, the compound is according to Embodiment [0026], wherein R^1 is CH_3 -.

[0028] Embodiment [0028]: In another example, the compound is according to Embodiment [0027], wherein




[0029] Embodiment [0029]: In one example, the compound is according to Embodiment [0028], wherein M^2 is a monocyclic five- to seven-membered heterocyclyl or a five- to six-membered heteroaryl, each optionally substituted with between one and three of R^{50} .

[0030] Embodiment [0030]: In another example, the compound is according to Embodiment [0029], wherein M^2 is selected from the group consisting of morpholinyl, thiazolyl, oxadiazolyl, tetrahydropyranyl, and oxazepanyl, each optionally substituted with between one and three of R^{50} .


[0031] Embodiment [0031]: In another example, the compound is according to Embodiment [0030], wherein M^1 is selected from the group consisting of -H, dimethylaminomethyl, (4-methylpiperazin-1-yl)methyl, piperidinyl, 1-methylpiperidin-4-yl, morpholin-4-ylmethyl, and phenylmethyl.

[0032] Embodiment [0032]: In another example, the compound is according to Embodiment [0031], wherein




[0033] Embodiment [0033]: In another example, the compound is according to Embodiment [0028], wherein M^1 is either a three- to seven-membered saturated carbocyclyl or a heterocyclyl with one or two annular heteroatoms, wherein the either of the aforementioned are optionally substituted with at least one of C_1 - C_3 alkyl, C_1 - C_3 alkoxy, C_1 - C_3 hydroxyalkyl, $R^{10}(R^{11})N$ -, and hydroxy, provided there are no geminal heteroatom substitutions; and wherein R^{10} and R^{11} are each independently C_1 - C_3 alkyl.

[0034] Embodiment [0034]: In another example, the compound is according to Embodiment [0033], wherein



[0035] Embodiment [0035]: In another example, the compound is according to Embodiment [0024], wherein M^1 - M^2 - M^3 - M^4 - together are according to formula II;



II

wherein X^1 , X^2 , and optionally X^3 , represent the atoms of a saturated bridged ring system, said saturated bridged ring system containing up to three annular heteroatoms represented by any of X^1 , X^2 , and X^3 ; wherein,

each X^1 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

each X^2 is independently a bridgehead methine optionally substituted with R^6 , or a bridgehead nitrogen;

each X^3 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

provided, for X^1 , X^2 , and X^3 , there are no nitrogen-nitrogen annular bonds nor geminal di-nitrogen substitutions;

E is selected from $-NR^9-$, $-O-$, and absent;

Y is either:

- a C_{1-3} alkylene linker, between the oxygen at the 7-position of the quinazoline ring system of I and either E, or when E is absent, any ring atom of the saturated bridged ring system except X^2 , when X^2 is a bridgehead nitrogen; provided there are at least two carbon atoms between the oxygen at the 7-position of the quinazoline ring system of I and either E, or when E is absent, any heteroatom represented by X^1 , X^2 or X^3 ; or

Y is absent, when Y is absent, E is also absent; said saturated bridged ring system is directly attached to the oxygen at the 7-position of the quinazoline ring system of I via a carbon atom of said saturated bridged ring system;

m and p are each independently between one and four;

n is between zero and two, when n is zero, then there is a direct single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -N(R³)R⁴, -S(O)₀₋₂R⁴, -SO₂N(R³)R⁴, -CO₂R³, -C(O)N(R³)R⁴, -N(R³)SO₂R⁴, -N(R³)C(O)R³, -NCO₂R³, -C(O)R³, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, optionally substituted lower heterocyclylalkyl; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P; and

R^8 is selected from R³, -SO₂N(R³)R⁴, -CO₂R³, -C(O)N(R³)R⁴, -SO₂R⁴, and -C(O)R³.

with the proviso that when Y is a C₁₋₃ alkylene linker; E is absent, Z is -NH- or -N(CH₃)-; R¹ is a C₁₋₃ alkyl; R² is -H or halogen; n = 0; and, the atoms, X¹, of one bridge of the saturated bridged ring system, when combined with both bridgehead atoms, X², of the saturated bridged ring system, represent:

either a pyrrolidine ring or a piperidine ring, and any atom, X¹ or X², of either of said pyrrolidine ring or said piperidine ring is attached to Y, then the other bridge of said saturated bridged ring system cannot be any one of -OC(O)CH₂-, -CH₂OC(O)-, -OC(O)CH₂CH₂-, -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, -OC(O)CH₂NH-, -OC(O)CH₂N(C₁₋₄alkyl)-, and -OC(O)CH₂O-; or

either a piperazine ring or a 4-(C₁₋₄ alkyl)-piperazine ring, and any atom, X¹ or X², of either of said piperazine ring or said 4-(C₁₋₄ alkyl)-piperazine ring is attached to Y, then the other bridge of said saturated bridged ring system, only when attached via the 2- and the 3-position of either of said piperazine ring or said 4-(C₁₋₄ alkyl)-piperazine ring, cannot be one of -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂alkyl groups; or

a piperazine ring, and any atom, X¹ or X², of said piperazine ring is attached to Y, then the other bridge of said saturated bridged ring system, only when

attached via the 3- and the 4-position of said piperazine ring, cannot be one of $-\text{C}(\text{O})\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{OC}(\text{O})\text{CH}_2-$, and either of the two aforementioned bridges optionally substituted by one or two C_{1-2} alkyl groups, and only when either of the two aforementioned bridges are attached to the 3-position of said piperazine ring via their left-hand end as depicted above; or

a 2-oxomorpholine ring, said 2-oxomorpholine ring attached to Y via its 4-position, then the other bridge of said saturated bridged ring system, only when attached via the 5- and the 6-position of said 2-oxomorpholine ring, cannot be one of $-(\text{CH}_2)_g-$, $-\text{CH}_2\text{WCH}_2-$, $-\text{CH}_2\text{WCH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{WCH}_2-$, wherein W is $-\text{O}-$, $-\text{S}(\text{O})_{0-2}-$, $-\text{NH}-$, or $-\text{N}(\text{C}_{1-4} \text{ alkyl})-$ wherein g is 2, 3, or 4.

[0036] Embodiment [0036]: In one example, the compound is according to Embodiment [0035], wherein Z is $-\text{NR}^5-$.

[0037] Embodiment [0037]: In another example, the compound is according to Embodiment [0036], wherein R^2 is selected from $-\text{H}$, halogen, trihalomethyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{OR}^3$, and optionally substituted lower alkyl.

[0038] Embodiment [0038]: In another example, the compound is according to Embodiment [0037], wherein R^1 is an unsubstituted C_{1-3} alkyl.

[0039] Embodiment [0039]: In another example, the compound is according to Embodiment [0038], wherein the saturated bridged ring system has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1].


[0040] Embodiment [0040]: In another example, the compound is according to Embodiment [0039], wherein Y is selected from $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-$, and absent.

[0041] Embodiment [0041]: In another example, the compound is according to Embodiment [0040], wherein q is one, two, or three.

[0042] Embodiment [0042]: In another example, the compound is according to Embodiment [0041], wherein R^5 is $-\text{H}$.

[0043] Embodiment [0043]: In another example, the compound is according to Embodiment [0042], wherein R^1 is methyl.

[0052] Embodiment [0052]: In another example, the compound is according to Formula Ia:



or a pharmaceutically acceptable salt or hydrate thereof, wherein

q is 1, 2, or 3;

R^2 is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -OR³, and optionally substituted lower alkyl;

E is selected from -NR⁹-, -O-, and absent;

Y is selected from -CH₂CH₂-, -CH₂-, and absent;

R^{10} is selected from -H, optionally substituted alkyl, and -OR¹³; and R^{11} and R^{12} are each independently selected from -H, -CF₃, -F, -N(R³)R⁴, -N(C=O)R³, -N(R³)SO₂R³, -S(O)₀₋₂R¹³, and -OR¹³; or

R^{10} is selected from -H, and -OR¹³; and R^{11} and R^{12} , when taken together, are oxo, exo-alkenyl, or when taken together with the carbon to which they are attached, form a three- to seven-membered spirocyclyl;

R^{13} is selected from -H, -C(=O)R⁴, optionally substituted lower alkynyl, optionally substituted lower arylalkynyl, optionally substituted lower heterocyclalkynyl, optionally substituted lower alkenyl, optionally substituted lower arylalkynyl, optionally substituted lower heterocyclalkynyl, optionally substituted lower alkyl, optionally substituted lower arylalkyl, optionally substituted aryl, optionally substituted lower heterocyclalkyl, and optionally substituted heterocycl;

two R^{13} 's, when taken together, form 1) a corresponding spirocyclic ketal from R^{11} , R^{12} and the carbon to which they are attached, when R^{11} and R^{12} are both -OR¹³, or 2) a corresponding cyclic ketal from R^{10} and one of R^{11} and R^{12} , and the corresponding carbons to which they are attached, when R^{10} is -OR¹³, and at least one of R^{11} and R^{12} is also -OR¹³.

[0053] Embodiment [0053]: In another example, the compound is according to Embodiment [0052], wherein Y is either -CH₂- or absent.

[0054] Embodiment [0054]: In another example, the compound is according to Embodiment [0053], wherein one of R^{11} and R^{12} is $-OR^{13}$, wherein R^{13} is selected from $-H$, $-C(O)R^4$, and optionally substituted lower alkyl; and R^{10} and the other of R^{11} and R^{12} are both $-H$.


[0055] Embodiment [0055]: In another example, the compound is according to Embodiment [0053], wherein one of R^{11} and R^{12} is $-F$; and R^{10} and the other of R^{11} and R^{12} are both $-H$.

[0056] Embodiment [0056]: In another example, the compound is according to Embodiment [0053], wherein R^{13} is an alkyl group containing at least one fluorine substitution thereon.

[0057] Embodiment [0057]: In another example, the compound is according to Embodiment [0053], wherein q is two or three.

[0058] Embodiment [0058]: In another example, the compound is according to Embodiment [0057], wherein each R^2 is independently selected from $-F$, $-Cl$, $-Br$, $-CF_3$, $-CH_3$, and $-OR^{25}$; wherein R^{25} is either methyl or aryl, each optionally substituted with one to three halogens.


[0059] Embodiment [0059]: In another example, the compound is according to Embodiment [0045], wherein said saturated bridged ring system is according to either formula V or formula VI;



wherein R^8 is selected from $-H$, optionally substituted lower alkyl, $-CO_2R^3$, $-C(O)N(R^3)R^4$, $-SO_2R^4$, and $-C(O)R^3$.

[0060] Embodiment [0060]: In another example, the compound is according to Embodiment [0059], wherein Y is either $-CH_2-$ or absent.


[0061] Embodiment [0061]: In another example, the compound is according to Embodiment [0060], wherein



[0062] Embodiment [0062]: In another example, the compound is according to Embodiment [0061], wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br.

[0063] Embodiment [0063]: In another example, the compound is according to Embodiment [0062], wherein R^8 is methyl or ethyl.

[0064] Embodiment [0064]: In another example, the compound is according to Embodiment [0046], wherein said saturated bridged ring system is according to formula **VII**;



VII

wherein A is selected from -O-, -S(O)₀₋₂-, -NR⁸-, -CR⁶R⁷-, and absent.

[0065] Embodiment [0065]: In another example, the compound is according to Embodiment [0064], wherein R^3 is selected from -H and optionally substituted alkyl.


[0066] Embodiment [0066]: In another example, the compound is according to Embodiment [0065], wherein A is either -C(R⁶)R⁷- or absent.

[0067] Embodiment [0067]: In another example, the compound is according to Embodiment [0066], wherein A is either -CH₂- or absent.

[0068] Embodiment [0068]: In another example, the compound is according to Embodiment [0067], wherein Y is -CH₂-.

[0069] Embodiment [0069]: In another example, the compound is according to Embodiment [0068], wherein q is 3.

[0070] Embodiment [0070]: In another example, the compound is according to Embodiment [0069], wherein




[0071] Embodiment [0071]: In another example, the compound is according to Embodiment [0070], wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br.


[0072] Embodiment [0072]: In another example, the compound is according to Embodiment [0043], wherein the saturated bridged ring system has a geometry selected from the group consisting of [3.3.1], [3.2.1], and [2.2.1].

[0073] Embodiment [0073]: In another example, the compound is according to Embodiment [0072], wherein said saturated bridged ring system contains one or two annular nitrogens, said one or two annular nitrogens are selected from $-NR^8-$, when X^1 , and a bridgehead nitrogen, when X^2 .

[0074] Embodiment [0074]: In another example, the compound is according to Embodiment [0073], wherein said saturated bridged ring system is according to formula VIII or formula IX;



VIII




IX

wherein R^8 is selected from -H, optionally substituted lower alkyl, $-CO_2R^3$, $-C(O)N(R^3)R^4$, $-SO_2R^4$, and $-C(O)R^3$; and R^{26} is C_{1-3} alkyl.

[0075] Embodiment [0075]: In another example, the compound is according to Embodiment [0074], wherein Y is $-CH_2CH_2-$; and E is either absent or $-N(R^9)-$.

[0076] Embodiment [0076]: In another example, the compound is according to Embodiment [0075], wherein q is 3.

[0077] Embodiment [0077]: In another example, the compound is according to Embodiment [0076], wherein



[0078] Embodiment [0078]: In another example, the compound is according to Embodiment [0077], wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br.

[0079] Embodiment [0079]: In another example, the compound is according to Embodiment [0078], wherein R^8 is methyl or ethyl.

[0080] Embodiment [0080]: In another example, the compound is according to Formula I or Ia, selected from Table 1:

Table 1




Entry	Name	Structure
1	N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,5r,6aS)-2-(1-methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-(1-methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
2	N-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3aR,5r,6aS)-2-(1-methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-(1-methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
3	7-({[(3aR,5r,6aS)-2-acetyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine; 7-({[(3aR,6aS)-2-acetyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine	

Table 1






4	N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-{{(3aR,5r,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl}oxy}quinazolin-4-amine; N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-{{(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl}oxy}quinazolin-4-amine	
5	ethyl (3aR,6aS)-5-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate	
6	N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,5r,6aS)-2-(methylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine; N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-2-(methylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine	
7	N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,5r,6aS)-2-ethyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-ethyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
8	N-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,5r,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine; N-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine	

Table 1





9	N-(3,4-dichloro-2-fluorophenyl)-7- ([(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(3,4-dichloro-2-fluorophenyl)-7- ([(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
10	N-(4-bromo-3-chloro-2-fluorophenyl)-7- ([(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(4-bromo-3-chloro-2-fluorophenyl)-7- ([(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
11	N-(3-chloro-2,4-difluorophenyl)-7- ([(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(3-chloro-2,4-difluorophenyl)-7- ([(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
12	N-(4,5-dichloro-2-fluorophenyl)-7- ([(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; N-(4,5-dichloro-2-fluorophenyl)-7- ([(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine	

Table 1





13	N-(4-bromo-5-chloro-2-fluorophenyl)-7- ({(3aR,5s,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine; N-(4-bromo-5-chloro-2-fluorophenyl)-7- ({(3aR,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine	
14	N-(4-bromo-2,3-dichlorophenyl)-7- ({(3aR,5s,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine; N-(4-bromo-2,3-dichlorophenyl)-7- ({(3aR,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine	
15	N-(3,4-dichlorophenyl)-7-((3aR,5s,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine; N-(3,4-dichlorophenyl)-7-((3aR,6aS)-2- methyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine	
16	N-(4-bromo-3-chloro-2-fluorophenyl)-7- ({(3aR,5r,6aS)-2- ethyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine; N-(4-bromo-3-chloro-2-fluorophenyl)-7- ({(3aR,6aS)-2- ethyloctahydrocyclopenta[c]pyrrol-5- yl)methyl}oxy)-6-(methyloxy)quinazolin-4- amine	

Table 1

17	N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methoxy)-7-({[(3aR,5r,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine; N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methoxy)-7-({[(3aR,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine	
18	N-(4-bromo-2,3-dichlorophenyl)-7-{{[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
19	N-(4,5-dichloro-2-fluorophenyl)-7-{{[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
20	N-(4-bromo-5-chloro-2-fluorophenyl)-7-{{[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
21	N-(3-chloro-2,4-difluorophenyl)-7-{{[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
22	N-(3,4-dichloro-2-fluorophenyl)-7-{{[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
23	N-(4-bromo-3-chloro-2-fluorophenyl)-7-{{[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	
24	N-(3-chloro-2,4-difluorophenyl)-7-{{[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methoxy)quinazolin-4-amine	

Table 1

25	N-(3,4-dichlorophenyl)-7-[(hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl)oxy]-6-(methyloxy)quinazolin-4-amine	
26	N-(4,5-dichloro-2-fluorophenyl)-7- {[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
27	N-(4-bromo-2,3-dichlorophenyl)-7- {[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
28	N-(4-bromo-5-chloro-2-fluorophenyl)-7- {[(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
29	N-(3,4-dichloro-2-fluorophenyl)-7- {[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
30	N-(4-bromo-3-chloro-2-fluorophenyl)-7- {[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
31	N-(3,4-dichlorophenyl)-7- {[(3R,8aR)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
32	N-(4-bromo-5-chloro-2-fluorophenyl)-7- {[(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine	

Table 1

33	N-(3,4-dichlorophenyl)-7-([(3S,8aR)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
34	N-(3,4-dichlorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
35	N-(3,4-dichlorophenyl)-7-([(3R,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
36	N-(3,4-dichloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
37	N-(4-bromo-3-chloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
38	N-(3-chloro-2,4-difluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
39	N-(4-bromo-2,3-dichlorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
40	N-(4,5-dichloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	
41	1,4:3,6-dianhydro-5-([4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol	

Table 1

42	1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol	
43	1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-xylo-hexitol	
44	1,4:3,6-dianhydro-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol	
45	1,4:3,6-dianhydro-5-([4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol	
46	1,4:3,6-dianhydro-5-([4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-glucitol	
47	1,4:3,6-dianhydro-2-deoxy-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-O-methyl-D-threo-hexitol	
48	1,4:3,6-dianhydro-5-deoxy-5-([4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol	
49	(3S,9aS)-3-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydro-2H-pyrido[1,2-a]pyrazin-1(6H)-one	
50	(3S,9aR)-3-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydro-2H-pyrido[1,2-a]pyrazin-1(6H)-one	

Table 1

51	(3S,8aS)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	
52	(3S,8aR)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	
53	(3S,8aS)-3-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	
54	(3S,8aS)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	
55	N-(3,4-dichlorophenyl)-7-({[2-[(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)ethyl]oxy]-6-(methyloxy)quinazolin-4-amine	
56	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{(8aR)-tetrahydro-1H-[1,3]thiazolo[4,3-c][1,4]oxazin-6-ylmethyl}oxy}quinazolin-4-amine; N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydro-1H-[1,3]thiazolo[4,3-c][1,4]oxazin-3-ylmethyl)oxy]quinazolin-4-amine	
57	N-(3,4-dichlorophenyl)-7-{{[2-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	

Table 1

58	N-(3,4-dichlorophenyl)-7-[[8-methyl-8-azabicyclo[3.2.1]oct-3-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
59	N-(3,4-dichlorophenyl)-7-[[3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]oxy}-6-(methyloxy)quinazolin-4-amine	
60	N-(3,4-dichlorophenyl)-7-[(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)oxy]-6-(methyloxy)quinazolin-4-amine	
61	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
62	1,4:3,6-dianhydro-2-O-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
63	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
64	1,4:3,6-dianhydro-2-O-methyl-5-O-[6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl]-L-iditol	
65	1,4:3,6-dianhydro-5-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-D-xylo-hexitol	
66	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	

Table 1

67	dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)-quinazolin-7-yl]-L-sorbose ethylene glycol acetal	
68	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
69	1,4:3,6-dianhydro-2-O-[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
70	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-(difluoromethyl)-L-iditol	
71	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
72	1,4:3,6-dianhydro-2-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
73	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
74	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-ethyl-L-iditol	
75	1,4:3,6-dianhydro-2-O-[4-[(3-bromo-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	

Table 1

76	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	
77	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-D-xylohexitol	
78	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-D-glucitol	
79	methyl 3,6-anhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-alpha-L-idofuranoside	
80	3,6-anhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2-O-(1-methylethylidene)-beta-L-xylo-hexofuranose	
81	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-5-methylidene-D-xylo-hexitol	
82	methyl 3,6-anhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-beta-L-idofuranoside	
83	N-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[(octahydro-2H-quinolizin-3-ylmethyl)oxy]quinazolin-4-amine	
84	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[6-(methyloxy)-4-[(2,3,4-trifluorophenyl)amino]quinazolin-7-yl]-D-iditol	

Table 1

85	1,4:3,6-dianhydro-5-O-[4-[(2-chloro-4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
86	1,4:3,6-dianhydro-5-O-[4-[(2-bromo-4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
87	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,6-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
88	1,4:3,6-dianhydro-5-O-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
89	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-[[4-fluoro-3-(trifluoromethyl)phenyl]amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol	
90	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
91	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,5-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
92	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,3-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
93	1,4:3,6-dianhydro-5-O-[4-[(5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	

Table 1

94	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(3,5-difluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-fluoro-D-iditol	
95	1,4:3,6-dianhydro-5-O-[4-[(3-chloro-4-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
96	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-2-chlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
97	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-fluoro-D-iditol	
98	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
99	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[6-(methoxy)-4-[(2,4,5-trifluorophenyl)amino]quinazolin-7-yl]-D-iditol	
100	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[6-(methoxy)-4-[(2,4,6-trifluorophenyl)amino]quinazolin-7-yl]-D-iditol	
101	1,4:3,6-dianhydro-5-O-[4-[(4-[(4-chlorophenyl)oxy]-3,5-difluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
102	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	

Table 1

103	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
104	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chloro-5-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
105	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
106	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl]-D-iditol	
107	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[6-(methyloxy)-4-[(3,4,5-trichlorophenyl)amino]quinazolin-7-yl]-D-iditol	
108	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
109	1,4:3,6-dianhydro-5-O-[4-[(4-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
110	1,4:3,6-dianhydro-5-O-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
111	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(3,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	

Table 1

112	1,4:3,6-dianhydro-5-O-[4-[(2-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
113	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-[(2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol	
114	1,4:3,6-dianhydro-5-O-[4-[(3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
115	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-[(4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol	
116	1,4:3,6-dianhydro-5-O-[4-[(4-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
117	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
118	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(2,5-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
119	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
120	1,4:3,6-dianhydro-5-O-[4-[(2-bromo-4,6-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	

Table 1

121	1,4:3,6-dianhydro-5-O-[4-{[4-chloro-3-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
122	1,4:3,6-dianhydro-5-O-[4-{[2-chloro-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
123	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-{[2-fluoro-3-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-D-iditol	
124	1,4:3,6-dianhydro-5-O-[4-{[2-bromo-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
125	1,4:3,6-dianhydro-5-O-[4-{[2-bromo-4-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
126	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-{[4-fluoro-2-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-D-iditol	
127	1,4:3,6-dianhydro-5-O-[4-{[3-bromo-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
128	1,4:3,6-dianhydro-5-O-[4-{[2-bromophenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
129	1,4:3,6-dianhydro-5-O-[4-{[3-bromophenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	

Table 1

130	1,4:3,6-dianhydro-5-O-[4-[(4-bromophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
131	1,4:3,6-dianhydro-5-O-[4-[(3-bromo-4-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
132	1,4:3,6-dianhydro-5-O-[4-[(5-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
133	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[(3,5-dimethylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
134	1,4:3,6-dianhydro-5-O-[4-[[2,5-bis(methyloxy)phenyl]amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
135	1,4:3,6-dianhydro-5-O-[4-[[5-chloro-2,4-bis(methyloxy)phenyl]amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
136	1,4:3,6-dianhydro-5-O-[4-[[4-chloro-2,5-bis(methyloxy)phenyl]amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
137	1,4:3,6-dianhydro-5-O-[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol	
138	N-(3,4-dichlorophenyl)-7-[[5-[(dimethylamino)methyl]-1,2,4-oxadiazol-3-yl]methyl]oxy]-6-(methyloxy)quinazolin-4-amine	

Table 1

139	N-(3,4-dichlorophenyl)-7-[(3-[(dimethylamino)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
140	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-[(4-methylpiperazin-1-yl)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	
141	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-piperidin-4-yl-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
142	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-(1-methylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
143	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-(morpholin-4-ylmethyl)-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	
144	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	
145	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-piperidin-2-yl-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
146	N-(3,4-dichlorophenyl)-7-[(2-[(dimethylamino)methyl]-1,3-thiazol-4-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
147	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4-(phenylmethyl)morpholin-2-yl)methyl]oxy]quinazolin-4-amine	

Table 1

148	1,1-dimethylethyl 2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholine-4-carboxylate	
149	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(morpholin-4-ylmethyl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
150	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-[(4-methylpiperazin-1-yl)methyl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
151	N-(3,4-dichlorophenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
152	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(1,4-oxazepan-2-ylmethyl)oxy]quinazolin-4-amine	
153	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-piperidin-3-yl-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	
154	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-(1-methylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	
155	N-(3,4-dichlorophenyl)-7-({[4-methyl-1,4-oxazepan-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
156	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-(1-methylpiperidin-3-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	

Table 1

157	N-(3,4-dichlorophenyl)-7-({[5-(1,1-dimethylethyl)-1,2,4-oxadiazol-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
158	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-phenyl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
159	7-[(2,1,3-benzothiadiazol-4-ylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
160	N-(3,4-dichlorophenyl)-7-({[5-methylisoxazol-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
161	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-methyl-4-phenylisoxazol-3-yl]methyl}oxy)quinazolin-4-amine	
162	7-[(1,3-benzothiazol-2-ylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
163	7-[(2,1,3-benzoxadiazol-5-ylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
164	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(2-thienyl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
165	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[1-phenyl-1H-pyrazol-4-yl]methyl}oxy)quinazolin-4-amine	

Table 1









166	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
167	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
168	7-([3-(4-chlorophenyl)-1,2,4-oxadiazol-5-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
169	7-([6-bromo-2-(methyloxy)naphthalen-1-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
170	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	
171	7-[(6-chloropyridin-3-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
172	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(pyridin-4-yl)methyl]oxy]quinazolin-4-amine	
173	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-methyl-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	

Table 1

174	7-[[[6-chloro-4H-1,3-benzodioxin-8-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
175	7-[[[5-chloro-1-methyl-3-phenyl-1H-pyrazol-4-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
176	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[1-methyl-3-(trifluoromethyl)-1H-thieno[2,3-c]pyrazol-5-yl)methyl]oxy]quinazolin-4-amine	
177	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[3-phenylisoxazol-5-yl)methyl]oxy]quinazolin-4-amine	
178	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[2,4,6-trimethylphenyl)methyl]oxy]quinazolin-4-amine	
179	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[pyridin-3-yl)methyl]oxy]quinazolin-4-amine	
180	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[3-[4-(methyloxy)phenyl]isoxazol-5-yl)methyl]oxy]quinazolin-4-amine	
181	N-(3,4-dichlorophenyl)-7-([5-[(2,4-dichlorophenyl)oxy]-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	

Table 1

182	7-[(cyclopropylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
183	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydrofuran-2-ylmethyl)oxy]quinazolin-4-amine	
184	7-(cyclopentyloxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
185	7-[(2-cyclohexylethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
186	7-[(cyclohexylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
187	7-[(cyclobutylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
188	N-(3,4-dichlorophenyl)-7-{[2-(1,3-dioxolan-2-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
189	N-(3,4-dichlorophenyl)-7-{[2-(1,3-dioxan-2-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
190	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-morpholin-4-ylethyl)oxy]quinazolin-4-amine	

Table 1

191	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-pyrrolidin-1-ylethyl)oxy]quinazolin-4-amine	
192	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-piperidin-1-ylethyl)oxy]quinazolin-4-amine	
193	2-(2-[[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy]ethyl)-1H-isoindole-1,3(2H)-dione	
194	methyl 6-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-alpha-D-glucopyranoside	
195	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-morpholin-4-yl-2-oxoethyl)oxy]quinazolin-4-amine	
196	1,1-dimethylethyl 2-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate	
197	1,1-dimethylethyl 4-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate	
198	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-pyrrolidin-1-ylphenyl)-1,3-thiazol-2-yl]methyl}oxy)quinazolin-4-amine	

Table 1

199	N-(3,4-dichlorophenyl)-7-({[4-[4-(diethylamino)phenyl]-1,3-thiazol-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
200	5-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-4-yl]-2-hydroxybenzamide	
201	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-pyridin-3-yl-1,3-thiazol-2-yl)methyl]oxy}quinazolin-4-amine	
202	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-pyridin-2-yl-1,3-thiazol-2-yl)methyl]oxy}quinazolin-4-amine	
203	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-pyridin-4-yl-1,3-thiazol-2-yl)methyl]oxy}quinazolin-4-amine	
204	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(2-morpholin-4-yl-1,3-thiazol-4-yl)methyl]oxy}quinazolin-4-amine	
205	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-(3-morpholin-4-yl-1,2,4-oxadiazol-5-yl)methyl]oxy}quinazolin-4-amine	
206	N-(3,4-dichlorophenyl)-7-({[3-(dimethylamino)-1,2,4-oxadiazol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
207	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[(4-methylpiperazin-1-yl)methyl]-1,3-thiazol-2-yl]methyl}oxy)quinazolin-4-amine	

Table 1

208	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-ylmethyl)oxy]quinazolin-4-amine	
209	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(morpholin-4-ylmethyl)-1,3-thiazol-2-yl]methyl}oxy)quinazolin-4-amine	
210	N-(3,4-dichlorophenyl)-7-({[4-[(4-methyl-1,4-diazepan-1-yl)methyl]-1,3-thiazol-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
211	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-{{[phenylmethyl]oxy}methyl}-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	
212	N-(3,4-dichlorophenyl)-7-({[4-ethylmorpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
213	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-piperidin-4-yl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
214	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(1-methylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
215	1,1-dimethylethyl 4-[5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate	
217	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-piperazin-1-yl-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine	

Table 1

218	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([3-(4-methylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl)oxy)quinazolin-4-amine	
219	N-(3,4-dichlorophenyl)-7-([5-(1-ethylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	
220	N-(3,4-dichlorophenyl)-7-([3-(4-ethylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	
221	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-[4-(methyloxy)phenyl]-1,2,4-oxadiazol-3-yl]methyl)oxy)quinazolin-4-amine	
222	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl]methyl)oxy)quinazolin-4-amine	
223	7-([2-(4-chlorophenyl)-1,3-thiazol-4-yl]methyl)oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
224	N-(3,4-dichlorophenyl)-7-([5-(3,5-dimethylisoxazol-4-yl)-1,2,4-oxadiazol-3-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	
225	7-([(5-chloro-1-benzothien-3-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	

Table 1

226	N-(3,4-dichlorophenyl)-7-[(3-[4-(1,1-dimethylethyl)phenyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	
227	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(5-[2-(methoxy)phenyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	
228	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([5-(4-methylphenyl)-1,3,4-oxadiazol-2-yl]methyl)oxy]quinazolin-4-amine	
229	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([1-(phenylmethyl)-1H-imidazol-2-yl]methyl)oxy]quinazolin-4-amine	
230	N-(3,4-dichlorophenyl)-7-([3-(2,6-dichlorophenyl)-5-methylisoxazol-4-yl]methyl)oxy]-6-(methoxy)quinazolin-4-amine	
231	N-(3,4-dichlorophenyl)-7-([(6-fluoro-4H-1,3-benzodioxin-8-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
232	7-([(3,5-dibromophenyl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
233	N-(3,4-dichlorophenyl)-7-([(2,6-difluorophenyl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	

Table 1

234	N-(3,4-dichlorophenyl)-6-(methoxy)-7-(((3-((pyridin-2-ylsulfonyl)methyl)-1,2,4-oxadiazol-5-yl)methyl)oxy)quinazolin-4-amine	
235	N-(3,4-dichlorophenyl)-6-(methoxy)-7-(((5-phenyl-1,2,4-oxadiazol-3-yl)methyl)oxy)quinazolin-4-amine	
236	7-((4-chloro-2-(trifluoromethyl)quinolin-6-yl)methyl)oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
237	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([2-(1-methylpyrrolidin-2-yl)ethyl]oxy)quinazolin-4-amine	
238	N-(3,4-dichlorophenyl)-7-([5-(1-ethylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
239	N-(3,4-dichlorophenyl)-7-([5-(1-ethylpiperidin-3-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
240	N-(3,4-dichlorophenyl)-7-([2-(dimethylamino)-1,3-thiazol-4-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
241	N-(3,4-dichlorophenyl)-7-([4-ethyl-1,4-oxazepan-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
242	N-(3,4-dichlorophenyl)-7-([2-(1-ethylpiperidin-4-yl)-1,3-thiazol-4-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	

Table 1

243	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-[(2S)-pyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	
244	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-[(2S)-pyrrolidin-2-yl]-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	
245	[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]methyl benzoate	
246	[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]methanol	
247	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)methyl]oxy]quinazolin-4-amine	
248	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-[(4S)-1,3-thiazolidin-4-yl]-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	
249	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-(1-piperidin-2-yl)-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	
250	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-(1-methylpiperidin-2-yl)-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	

Table 1

251	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-piperidin-3-yl-1,3-thiazol-4-yl)methyl]oxy}quinazolin-4-amine	
252	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-(1-methylpiperidin-3-yl)-1,3-thiazol-4-yl)methyl]oxy}quinazolin-4-amine	
253	N-(3,4-dichlorophenyl)-7-([2-(1-ethylpiperidin-2-yl)-1,3-thiazol-4-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
254	N-(3,4-dichlorophenyl)-7-([2-(1-ethylpiperidin-3-yl)-1,3-thiazol-4-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
255	N-(3,4-dichlorophenyl)-7-([3-[(2S)-1-ethylpyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
256	N-(3,4-dichlorophenyl)-7-([2-[(2S)-1-ethylpyrrolidin-2-yl]-1,3-thiazol-4-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
257	N-(3,4-dichlorophenyl)-7-([5-ethyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
258	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-propyl-1,4-oxazepan-2-yl)methyl]oxy}quinazolin-4-amine	

Table 1

259	7-({[4-(cyclopropylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
260	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-{2-(methyloxy)ethyl}-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	
261	N-(3,4-dichlorophenyl)-7-({[4-(1-methylethyl)-1,4-oxazepan-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
262	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-piperazin-1-yl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
263	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-pyrrolidin-2-yl-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	
264	N-(3,4-dichlorophenyl)-7-({[5-(1-ethylpyrrolidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
265	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-[(2S)-1-methylpyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine	
266	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-[(2S)-1-methylpyrrolidin-2-yl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
267	N-(3,4-dichlorophenyl)-7-({[2-(4-ethylpiperazin-1-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	

Table 1

268	N-(3,4-dichlorophenyl)-7-({[(1,4-dimethylpiperazin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
269	7-({[(4-cyclopentylmorpholin-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
270	N-(3,4-dichlorophenyl)-7-({[4-(1-methylethyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
271	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(3-phenylpropyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
272	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[2-(methyloxy)ethyl]morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
273	ethyl 2-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]propanoate	
274	N-(3,4-dichlorophenyl)-7-({[(4-hex-5-en-1-yl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
275	2-({2-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]ethyl}oxy)ethanol	
276	methyl 3-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]propanoate	

Table 1

277	6-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]hexanenitrile	
278	N-(3,4-dichlorophenyl)-6-(methoxy)-7-({[4-(tetrahydro-2H-pyran-2-ylmethyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
279	4-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]butanenitrile	
280	N-(3,4-dichlorophenyl)-7-({[4-[(4-fluorophenyl)methyl]morpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	
281	methyl 5-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]pentanoate	
282	N-(3,4-dichlorophenyl)-6-(methoxy)-7-({[4-(oct-7-en-1-yl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
283	N-(3,4-dichlorophenyl)-6-(methoxy)-7-({[4-(propyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
284	6-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]hexan-1-ol	
285	7-({[4-(acetylmorpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	

Table 1

286	7-({[4-(cyclopropylmethyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
287	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-prop-2-yn-1-ylmorpholin-2-yl]methyl}oxy)quinazolin-4-amine	
288	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-pyridin-4-ylmorpholin-2-yl]methyl}oxy)quinazolin-4-amine	
289	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(pyridin-2-ylmethyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
290	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-pent-2-yn-1-ylmorpholin-2-yl]methyl}oxy)quinazolin-4-amine	
291	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(4-methylpiperazin-1-yl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
292	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-(1-methylpyrrolidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	
293	N-(3-chloro-4-fluorophenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
294	7-({[4-butyl-1,4-oxazepan-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	

Table 1

295	(3,4-dichlorophenyl)[7-(methoxy)-6-({[4-(2-methylpropyl)-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	
296	7-({[4-acetyl-1-ethylpiperazin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
297	(3,4-dichlorophenyl)(6-(methoxy)-7-({[4-pentyl-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	
298	(3,4-dichlorophenyl)[6-(methoxy)-7-({[4-(tetrahydro-2H-pyran-2-ylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	
299	(3,4-dichlorophenyl)[6-(methoxy)-7-({[4-(3-thienylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	
300	N-[4-chloro-2,5-bis(methoxy)phenyl]-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	
301	N-(3-bromo-2-methylphenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	
302	7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methoxy)-N-(3,4,5-trichlorophenyl)quinazolin-4-amine	
303	N-(3-chloro-2-methylphenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	

Table 1










304	N-(3,4-dichlorophenyl)-7-[[[4-ethanimidoyl-1,4-oxazepan-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
305	N-(4-bromo-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
306	N-(5-chloro-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
307	N-(4-chloro-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
308	N-(2,4-dichlorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
309	N-(2,4-dibromophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
310	7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)-N-(2,3,4-trichlorophenyl)quinazolin-4-amine	
311	N-(3,4-dichlorophenyl)-7-[[[1-ethyl-4-methylpiperazin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
312	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholine-4-carboximidamide	

Table 1

313	N-(3,4-dichlorophenyl)-6-(methoxy)-7-({[2-(pyrrolidin-1-ylmethyl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	
314	N-(3,4-dichlorophenyl)-6-(methoxy)-7-({[4-(tetrahydro-2H-pyran-4-yl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
315	N-(3,4-dichlorophenyl)-7-({[4-(2-ethylbutyl)morpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	
316	7-({[4-(cyclohexylmethyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
317	2-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]ethanol	
318	7-({[4-but-2-yn-1-ylmorpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
319	7-({[4-(cyclobutylmorpholin-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	
320	N-(3,4-dichlorophenyl)-7-({[4-[2-(1,3-dioxolan-2-yl)ethyl]morpholin-2-yl]methyl}oxy)-6-(methoxy)quinazolin-4-amine	
321	7-({[4-(2-cyclohexylethyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	

Table 1

322	N-(3,4-dichlorophenyl)-7-[(4-[2-(1,3-dioxan-2-yl)ethyl]morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
323	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-pent-4-en-1-yl)morpholin-2-yl)methyl]oxy}quinazolin-4-amine	
324	N-(3,4-dichlorophenyl)-7-[(4-[(2R)-2-methylbutyl]morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
325	N-(3,4-dichlorophenyl)-7-[(4-(4-fluorobutyl)morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
326	3-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]butan-2-one	
327	1-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]butan-2-one	
328	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-pentylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	
329	N-(3,4-dichlorophenyl)-7-[(4-hexylmorpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
330	N-(3,4-dichlorophenyl)-7-[(4-heptylmorpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	

Table 1










331	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([(4-octylmorpholin-2-yl)methyl]oxy)quinazolin-4-amine	
332	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([(4-(2-phenylethyl)morpholin-2-yl)methyl]oxy)quinazolin-4-amine	
333	7-([(4-butylmorpholin-2-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
334	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([(4-prop-2-en-1-yl)morpholin-2-yl)methyl]oxy)quinazolin-4-amine	
335	2-[2-([(4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]-1-phenylethanone	
336	N-(3,4-dichlorophenyl)-7-([(4-(2-fluoroethyl)morpholin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
337	N-(3,4-dichlorophenyl)-7-([(4-(3-methylbut-2-en-1-yl)morpholin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	
338	7-([(4-[(2E)-3-bromoprop-2-en-1-yl]morpholin-2-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
339	2-[2-([(4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]acetamide	

Table 1

340	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([4-[3-(tetrahydro-2H-pyran-2-yloxy)propyl]-1,4-oxazepan-2-yl]methyl)oxy)quinazolin-4-amine	
341	N-(3,4-dichlorophenyl)-7-([4-(3-methylbutyl)-1,4-oxazepan-2-yl]methyl)oxy)-6-(methoxy)quinazolin-4-amine	
342	7-([4-(cyclohexylmethyl)-1,4-oxazepan-2-yl]methyl)oxy)-4-[(3,4-dichlorophenyl)methyl]-6-(methoxy)quinazoline	
343	7-([4-(2-cyclohexylethyl)-1,4-oxazepan-2-yl]methyl)oxy)-4-[(3,4-dichlorophenyl)methyl]-6-(methoxy)quinazoline	
345	N-(3,4-dichlorophenyl)-7-([4-(2-ethylbutyl)-1,4-oxazepan-2-yl]methyl)oxy)-6-(methoxy)quinazolin-4-amine	
346	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([4-(methylsulfonyl)-1,4-oxazepan-2-yl]methyl)oxy)quinazolin-4-amine	
347	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([4-(1-methylpiperidin-4-yl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	
348	N-(3-chloro-2-fluorophenyl)-7-([4-(4-methylmorpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	
349	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)-1,4-oxazepan-4-carboximidamide	

Table 1

350	N-(3-bromo-4-methylphenyl)-7-({[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
351	N-(3,4-dichlorophenyl)-7-({[1,4-diethylpiperazin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
352	4-({[4-[(4-bromo-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-N'-cyanopiperidine-1-carboximidamide	
353	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(methylsulfonyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
354	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[(phenylmethyl)sulfonyl]morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
355	N-(3,4-dichlorophenyl)-7-({[4-[(4-fluorophenyl)sulfonyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
356	N-(3,4-dichlorophenyl)-7-({[4-(ethylsulfonyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
357	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(phenylsulfonyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	

Table 1

358	7-[(4-[(3-chloropropyl)sulfonyl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
359	7-[(4-(butylsulfonyl)morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
360	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4-[(4-methylphenyl)sulfonyl]morpholin-2-yl)methyl]oxy]quinazolin-4-amine	
361	N-(3,4-dichlorophenyl)-7-[(4-[(3,5-dimethylisoxazol-4-yl)carbonyl]morpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	
362	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4-[(3-(methyloxy)phenyl)acetyl]morpholin-2-yl)methyl]oxy]quinazolin-4-amine	
363	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4-(2-methylpentanoyl)morpholin-2-yl)methyl]oxy]quinazolin-4-amine	
364	7-[(4-[(4-butylphenyl)carbonyl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
365	7-[(4-[(4-chlorophenyl)acetyl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
366	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4-(2-propylpentanoyl)morpholin-2-yl)methyl]oxy]quinazolin-4-amine	

Table 1

367	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-methylpentanoyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
368	N-(3,4-dichlorophenyl)-7-({[4-[(2,5-difluorophenyl)carbonyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
369	7-({[4-(cyclopentylcarbonyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
370	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(2-phenylbutanoyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
371	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[(2,3,6-trifluorophenyl)carbonyl]morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
372	N-(3,4-dichlorophenyl)-7-({[4-(furan-3-ylcarbonyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
373	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(propanoyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	
374	N-(3,4-dichlorophenyl)-7-({[4-(hexanoyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	
375	N-(3,4-dichlorophenyl)-7-({[4-(2-ethylhexanoyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	

Table 1










376	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(3-phenylpropanoyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	
377	N-(3,4-dichlorophenyl)-7-([4-(2,2-dimethylpropanoyl)morpholin-2-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	
378	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(naphthalen-1-ylcarbonyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	
379	7-([4-[(2-chloropyridin-3-yl)carbonyl]morpholin-2-yl]methyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
380	7-([4-[(6-chloropyridin-3-yl)carbonyl]morpholin-2-yl]methyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
381	7-([4-[(1,3-benzodioxol-5-yl)carbonyl]morpholin-2-yl]methyl)oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	
382	N-(3,4-dichlorophenyl)-6-[(1-methylethyl)oxy]-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	
383	N-(3,4-dichlorophenyl)-6-[[2-(methyloxy)ethyl]oxy]-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	
384	N-(3,4-dichlorophenyl)-6-(ethyloxy)-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	

Table 1

385	N-(3,4-dichlorophenyl)-6-(ethoxy)-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	
386	N-(4-bromo-2-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
387	N-(4-chloro-3-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
388	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)-N-methylmorpholine-4-carboximidamide	
389	N-(4-bromo-3-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
390	N-(3,4-dichlorophenyl)-6-[(1-methylethyl)oxy]-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	
391	N-(3,4-dichlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-[[2-(methoxy)ethyl]oxy}quinazolin-4-amine	
392	N-(4-bromo-2-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
393	7-[[4-(4-acetyl-1,4-oxazepan-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	

Table 1

394	4-[(3,4-dichlorophenyl)amino]-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-6-ol	
395	N-(3-bromo-4-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
396	3-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]-3-oxopropanoic acid	
397	methyl 4-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]-4-oxobutanoate	
398	N-(3,4-dichlorophenyl)-7-[[4-methylmorpholin-3-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
399	N-(3-bromo-2-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
400	N'-cyano-2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-N-[2-(methyloxy)ethyl]morpholine-4-carboximidamide	
401	N'-cyano-2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-N-ethylmorpholine-4-carboximidamide	
402	[(1E)-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl](piperidin-1-yl)methylidene]cyanamide	

Table 1

403	[(1E)-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl](pyrrolidin-1-yl)methylidene]cyanamide	
404	[(1E)-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl](4-methylpiperazin-1-yl)methylidene]cyanamide	
405	N-(3,4-dichlorophenyl)-7-{{[(6-ethyl-4,6-dimethylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
406	N-(4-bromo-3-methylphenyl)-7-{{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
407	N-(3,4-dichlorophenyl)-7-{{[(6,6-dimethylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	
408	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(4,6,6-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	
409	N-(3,4-dichlorophenyl)-7-{{[2-(5,5-dimethylmorpholin-2-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	
410	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[2-(4,5,5-trimethylmorpholin-2-yl)ethyl]oxy}quinazolin-4-amine	
411	1,1-dimethylethyl 2-(2-{{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl}-5,5-dimethylmorpholine-4-carboxylate	

Table 1

412	N-(3,4-dichlorophenyl)-6-(methoxy)-7- {[(4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	
413	N-(4-bromo-2,3-dichlorophenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
414	N-(4,5-dichloro-2-fluorophenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
415	N-(3,4-dichlorophenyl)-6-(methoxy)-7- {[2-(4,6,6-trimethylmorpholin-2-yl)ethyl]oxy}quinazolin-4-amine	
416	N-(4-bromo-2,3-difluorophenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
417	N-(4-bromo-2,5-difluorophenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
418	N-(4-bromo-3,5-difluorophenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
419	N-(3,4-dichloro-2-methylphenyl)-7- {[(4-methylmorpholin-2-yl)methyl]oxy}-6- (methoxy)quinazolin-4-amine	
420	N-(3,4-dichlorophenyl)-7- {[(2R,5S,6S)-5,6- dimethylmorpholin-2-yl]methyl]oxy}-6- (methoxy)quinazolin-4-amine	

Table 1

421	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([[(2R,5S,6S)-4,5,6-trimethylmorpholin-2-yl]methyl]oxy)quinazolin-4-amine	
422	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([[(2S,5S,6S)-4,5,6-trimethylmorpholin-2-yl]methyl]oxy)quinazolin-4-amine	
423	N-(4-bromo-3-chloro-2-methylphenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
424	N-(4-bromo-5-chloro-2-fluorophenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
425	N-(4-bromo-3-chloro-2-fluorophenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
426	N-(3,4-dichloro-2-fluorophenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
427	N-(3-chloro-2,4-difluorophenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
428	N-(2,3-dichloro-4-methylphenyl)-7-{[(4-methylmorpholin-2-yl)methyl]oxy}-6-(methoxy)quinazolin-4-amine	
429	6-([4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl-3,3,4-trimethylmorpholin-2-one	

Table 1

430	N-(4-bromo-2,3-dichlorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
431	N-(4-bromo-5-chloro-2-fluorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
432	N-(4,5-dichloro-2-fluorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
433	N-(3,4-dichloro-2-fluorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
434	N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
435	N-(3-chloro-2,4-difluorophenyl)-6-(methoxy)-7-[[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	
436	(6S)-6-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)-4-methylpiperazin-2-one	
437	(6S)-6-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)-4-methylpiperazin-2-one	
438	(6S)-6-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)-1,4-dimethylpiperazin-2-one	

Table 1

439	(6S)-6-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-1,4-dimethylpiperazin-2-one	
440	N-(4-bromo-3-chlorophenyl)-7-[[[(3a'S,4R,6'S,6a'R)-2,2-dimethyltetrahydrospiro[1,3-dioxolane-4,3'-furo[3,2-b]furan]-6'-yl]oxy]-6-(methyloxy)quinazolin-4-amine	
441	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-5-C-[(methyloxy)methyl]-L-glucitol	
442	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-(methylsulfonyl)-L-glucitol	
443	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-glucitol	
444	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-S-methyl-5-thio-D-iditol	
445	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-morpholin-4-yl-D-iditol	
446	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(4-methylpiperazin-1-yl)-D-iditol	

Table 1

447	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-pyrrolidin-1-yl-D-iditol	
448	2-O-acetyl-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol	
449	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol	
450	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(methylsulfonyl)-D-iditol	
451	2-amino-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-D-iditol	
452	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(dimethylamino)-D-iditol	
453	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(diethylamino)-D-iditol	
454	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-piperidin-1-yl-D-iditol	
455	2-(acetamino)-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-D-iditol	

Table 1

456	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-5-O-methyl-5-C-(trifluoromethyl)-L-glucitol	
457	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-2-deoxy-2-[(methylsulfonyl)amino]-D-iditol	
458	N-(4-bromo-3-chlorophenyl)-6-(methoxy)-7-[(1-methylpyrrolidin-3-yl)oxy]quinazolin-4-amine	
459	N-(4-bromo-3-chlorophenyl)-6-(methoxy)-7-[(3R)-tetrahydrofuran-3-yloxy]quinazolin-4-amine	
460	N-(4-bromo-3-chlorophenyl)-6-(methoxy)-7-[[[(3S,4R)-4-(methoxy)tetrahydrofuran-3-yl]oxy]quinazolin-4-amine	
461	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-(6-(methoxy)-4-[[4-(4-methylpiperazin-1-yl)phenyl]amino]quinazolin-7-yl)-D-iditol	
462	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-[[3-fluoro-4-(4-methylpiperazin-1-yl)phenyl]amino]-6-(methoxy)quinazolin-7-yl]-D-iditol	
463	1,4:3,6-dianhydro-2-deoxy-5-O-[4-[[2,3-dichloro-4-(4-methylpiperazin-1-yl)phenyl]amino]-6-(methoxy)quinazolin-7-yl]-2-fluoro-D-iditol	

Table 1

464	1,4:3,6-dianhydro-2-deoxy-5-O-[4-{[3,4-dichloro-2-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol	
465	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-C-(trifluoromethyl)-D-glucitol	
466	(3,4-dichlorophenyl)[6-(methyloxy)-7-({[4-(tetrahydrofuran-2-ylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	


- [0081] Embodiment 81: Another aspect of the invention is a pharmaceutical composition comprising a compound according to any one of embodiments [0024]-[0080] and a pharmaceutically acceptable carrier.
- [0082] Embodiment 82: Another aspect of the invention is a metabolite of the compound, and optionally together with a pharmaceutically acceptable carrier, according to any one of embodiments [0024]-[0081].
- [0083] Embodiment 83: Another aspect of the invention is a method of modulating the *in vivo* activity of a kinase, the method comprising administering to a subject an effective amount of the compound, and optionally together with a pharmaceutically acceptable carrier, according to any of embodiments [0024]-[0081].
- [0084] Embodiment 84: Another aspect of the invention is a method according to embodiment [0083], wherein the kinase is selected from ephrin and EGFR.
- [0085] Embodiment 85: Another aspect of the invention is a method of modulating the *in vivo* activity of a plurality of kinases, the method comprising administering to a subject an effective amount of the compound, and optionally together with a pharmaceutically acceptable carrier, according to any of embodiments [0024]-[0081].
- [0086] Embodiment 86: Another aspect of the invention is a method according to embodiment [0085], wherein the plurality of kinases comprises at least one of ephrin and EGFR.
- [0087] Embodiment 87: Another aspect of the invention is a method according to embodiment [0084], wherein modulating the *in vivo* activity of the kinase comprises inhibition of said kinase.
- [0088] Embodiment 88: Another aspect of the invention is a method according to embodiment [0086], wherein modulating the *in vivo* activity of the plurality of kinases comprises inhibition of at least one of ephrin and EGFR.
- [0089] Embodiment 89: Another aspect of the invention is a method of treating diseases or disorders associated with uncontrolled, abnormal, and/or unwanted cellular activities, the method comprising administering, to a mammal in need thereof, a therapeutically effective amount of the compound, and optionally together with a pharmaceutically acceptable carrier, as described in any one of embodiments [0024]-[0081].

Definitions

- [0090] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in


which they are used indicates otherwise or they are expressly defined to mean something different.

[0091] The atom numbering convention for the quinazoline structure is as follows:




[0092] The symbol “-” means a single bond, “=” means a double bond, “≡” means a triple bond. The symbol “” refers to a group on a double-bond as occupying either position on the terminus of a double bond to which the symbol is attached; that is, the geometry, *E*- or *Z*-, of the double bond is ambiguous. When a group is depicted removed from its parent formula, the “” symbol will be used at the end of the bond which was theoretically cleaved in order to separate the group from its parent structural formula.

[0093] If a group “R” is depicted as “floating” on a ring system, as for example in the formula:



then, unless otherwise defined, a substituent “R” may reside on any atom of the ring system, assuming replacement of a depicted, implied, or expressly defined hydrogen from one of the ring atoms, so long as a stable structure is formed.


[0094] If a group “R” is depicted as floating on a fused ring system, as for example in the formulae:



then, unless otherwise defined, a substituent “R” may reside on any atom of the fused ring system, assuming replacement of a depicted (for example the -NH- in the formula above), implied (for example as in the formula above, where the hydrogens are not shown but understood to be present), or expressly defined hydrogen (for example where in the formula above, “X” equals -CH-) from one of the ring atoms, so long as a stable structure is formed. In the example depicted, the “R” group may reside on either the 5-membered or the 6-membered ring of the fused ring system. In the formula depicted above, when *y* is 2 for example, then the two “R’s” may reside on any two atoms of the ring system,


again assuming each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0095] When there are more than one such depicted “floating” groups, as for example in the formulae:




where there are two groups, namely, the “R” and the bond indicating attachment to a parent structure; then, unless otherwise defined, the “floating” groups may reside on any atoms of the ring system, again assuming each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0096] When a group “R” is depicted as existing on a ring system containing saturated carbons, as for example in the formula:



where, in this example, “y” can be more than one, assuming each replaces a currently depicted, implied, or expressly defined hydrogen on the ring; then, unless otherwise defined, where the resulting structure is stable, two “R’s” may reside on the same carbon. A simple example is when R is a methyl group; there can exist a geminal dimethyl on a carbon of the depicted ring (an “annular” carbon). In another example, two R’s on the same carbon, including that carbon, may form a ring, thus creating a spirocyclic ring (a “spirocyclyl” group) structure with the depicted ring as for example in the formula:



[0097] “Alkyl” is intended to include linear, branched, or cyclic hydrocarbon structures and combinations thereof, inclusively. For example, “C₈ alkyl” may refer to an *n*-octyl, *iso*-octyl, cyclohexylethyl, and the like. Lower alkyl refers to alkyl groups of from one to six carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *s*-butyl, *t*-butyl, isobutyl, pentyl, hexyl and the like. Higher alkyl refers to alkyl groups containing more than eight carbon atoms. Exemplary alkyl groups are those of C₂₀ or below. Cycloalkyl is a subset of alkyl and includes cyclic hydrocarbon groups of from three to thirteen carbon atoms. Examples of cycloalkyl groups include *c*-propyl, *c*-butyl, *c*-pentyl, norbornyl, adamantyl and the like. In this application, alkyl refers to alkanyl, alkenyl, and alkynyl residues (and combinations thereof); it is intended

to include cyclohexylmethyl, vinyl, allyl, isoprenyl, and the like. Thus when an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, either “butyl” or “C₄ alkyl” is meant to include *n*-butyl, *sec*-butyl, isobutyl, *t*-butyl, isobutenyl and but-2-yne radicals; and “propyl” or “C₃ alkyl” each include *n*-propyl, propenyl, and isopropyl.

[0098] “Alkylene” refers to straight or branched chain divalent radical consisting solely of carbon and hydrogen atoms, containing no unsaturation and having from one to ten carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene and the like. Alkylene is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, fully saturated. Examples of alkylene include ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), dimethylpropylene (-CH₂C(CH₃)₂CH₂-), and cyclohexylpropylene (-CH₂CH₂CH(C₆H₁₃)).

[0099] “Alkylidene” refers to a straight or branched chain unsaturated divalent radical consisting solely of carbon and hydrogen atoms, having from two to ten carbon atoms, for example, ethylidene, propylidene, *n*-butylidene, and the like. Alkylidene is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, double bond unsaturation. The unsaturation present includes at least one double bond and a double bond can exist between the first carbon of the chain and a carbon atom of the rest of the molecule to which it is attached.

[00100] “Alkylidyne” refers to a straight or branched chain unsaturated divalent radical consisting solely of carbon and hydrogen atoms having from two to ten carbon atoms, for example, propylid-2-ynyl, *n*-butylid-1-ynyl, and the like. Alkylidyne is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, triple bond unsaturation. The unsaturation present includes at least one triple bond and a triple bond can exist between the first carbon of the chain and a carbon atom of the rest of the molecule to which it is attached.

[00101] Any of the above radicals, “alkylene,” “alkylidene” and “alkylidyne,” when optionally substituted, may contain alkyl substitution which itself contains unsaturation. For example, 2-(2-phenylethynyl-but-3-enyl)-naphthalene (IUPAC name) contains an *n*-butylid-3-ynyl radical with a vinyl substituent at the 2-position of said radical.

[00102] “Alkoxy” or “alkoxyl” refers to the group -O-alkyl, for example including from one to eight carbon atoms of a straight, branched, cyclic configuration, unsaturated chains, and combinations thereof attached to the parent structure through an oxygen.

Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy, cyclohexyloxy and the like. Lower-alkoxy refers to groups containing one to six carbons.

[00103] “Substituted alkoxy” refers to the group -O-(substituted alkyl), the substitution on the alkyl group generally containing more than only carbon (as defined by alkoxy). One exemplary substituted alkoxy group is “polyalkoxy” or -O-optionally substituted alkylene-optionally substituted alkoxy, and includes groups such as -OCH₂CH₂OCH₃, and glycol ethers such as polyethyleneglycol and -O(CH₂CH₂O)_xCH₃, where x is an integer of between about two and about twenty, in another example, between about two and about ten, and in a further example between about two and about five. Another exemplary substituted alkoxy group is hydroxyalkoxy or -OCH₂(CH₂)_yOH, where y is for example an integer of between about one and about ten, in another example y is an integer of between about one and about four.

[00104] “Acyl” refers to groups of from one to ten carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples include acetyl, benzoyl, propionyl, isobutyryl, t-butoxycarbonyl, benzyloxycarbonyl and the like. Lower-acyl refers to groups containing one to six carbons.


[0100] “α-Amino Acids” refer to naturally occurring and commercially available amino acids and optical isomers thereof. Typical natural and commercially available α-amino acids are glycine, alanine, serine, homoserine, threonine, valine, norvaline, leucine, isoleucine, norleucine, aspartic acid, glutamic acid, lysine, ornithine, histidine, arginine, cysteine, homocysteine, methionine, phenylalanine, homophenylalanine, phenylglycine, ortho-tyrosine, meta-tyrosine, para-tyrosine, tryptophan, glutamine, asparagine, proline and hydroxyproline. A “side chain of an α-amino acid” refers to the radical found on the α-carbon of an α-amino acid as defined above, for example, hydrogen (for glycine), methyl (for alanine), benzyl (for phenylalanine), and the like.

[0101] “Amino” refers to the group -NH₂. “Substituted amino,” refers to the group -N(H)R or -N(R)R where each R is independently selected from the group: optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted heterocyclyl, acyl, carboxy, alkoxycarbonyl, sulfanyl, sulfinyl and sulfonyl, for example, diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino.

[0102] “Aryl” refers to aromatic six- to fourteen-membered carbocyclic rings include, for example, benzene, naphthalene, indane, tetralin, fluorene and the like.

[0103] “Arylalkyl” refers to a residue in which an aryl moiety is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne radical. Examples include benzyl, phenethyl, phenylvinyl, phenylallyl and the like. The aryl, alkylene, alkylidene, or alkylidyne radical portion of an arylalkyl group may be optionally substituted. “Lower arylalkyl” refers to an arylalkyl where the “alkyl” portion of the group has one to six carbons.

[0104] “Exo-alkenyl” refers to a double bond that emanates from an annular carbon, and is not within the ring system, for example the double bond depicted in the formula below.



[0105] “Fused-polycyclic” or “fused ring system” refers to a polycyclic ring system that contains bridged or fused rings; that is, where two rings have more than one shared atom in their ring structures. Typically, but not necessarily, fused-polycyclics share a vicinal set of atoms. Typically, a spiro ring system is not a fused-polycyclic by this definition, but fused polycyclic ring systems of the invention may themselves have spiro rings attached thereto via a single ring atom of the fused-polycyclic.

[0106] “Halogen” or “halo” refers to fluorine, chlorine, bromine or iodine. Dihaloaryl, dihaloalkyl, trihaloaryl etc. refer to aryl and alkyl substituted with a plurality of halogens, but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl.

[0107] “Heteroatom” refers to O, S, N, or P.

[0108] “Heterocyclyl” refers to a stable three- to fifteen-membered ring radical that consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, phosphorus, oxygen and sulfur. For purposes of this invention, the heterocyclyl radical may be a monocyclic, bicyclic or tricyclic ring system, which may include fused or bridged ring systems, and the nitrogen, phosphorus, carbon or sulfur atoms in the heterocyclyl radical may be optionally oxidized to various oxidation states. In addition, the nitrogen atom may be optionally quaternized; and the ring radical may be partially or fully saturated or aromatic. Examples of such heterocyclyl radicals include, but are not limited to, azetidiny, acridiny, benzodioxolyl, benzodioxanyl, benzofuranyl, carbazoyl, cinnolinyl, dioxolanyl, indoliziny, naphthyridiny, perhydroazepiny,


phenaziny, phenothiaziny, phenoxaziny, phthalaziny, pteridiny, puriny, quinazolinyl, quinoxaliny, quinolinyl, isoquinolinyl, tetrazoyl, tetrahydroisoquinolyl, piperidinyl, piperaziny, 2-oxopiperaziny, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepiny, azepiny, pyrroly, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, dihydropyridinyl, tetrahydropyridinyl, pyridinyl, pyraziny, pyrimidinyl, pyridaziny, oxazolyl, oxazolinyl, oxazolidinyl, triazolyl, indanyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolinyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, isoindolyl, indolinyl, isoindolinyl, octahydroindolyl, octahydroisoindolyl, quinolyl, isoquinolyl, decahydroisoquinolyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothieliyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, dioxaphospholanyl, and oxadiazolyl.

- [0109] “Heteroalicyclic” refers specifically to a non-aromatic heterocyclyl system radical.
- [0110] “Heteroaryl” refers specifically to an aromatic heterocyclyl system radical.
- [0111] “Heterocyclylalkyl” refers to a residue in which a heterocyclyl is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne radical. Examples include (4-methylpiperazin-1-yl) methyl, (morpholin-4-yl) methyl, 2-(oxazolin-2-yl) ethyl, 4-(4-methylpiperazin-1-yl)-2-butenyl, and the like. The heterocyclyl, alkylene, alkylidene, or alkylidyne radical portion of an arylalkyl group may be optionally substituted. “Lower heterocyclylalkyl” refers to an arylalkyl where the “alkyl” portion of the group has one to six carbons.
- [0112] “Hydrocarbyl” refers to a hydrocarbon residue, generally. The term “hydrocarbyl” can be modified to mean more specific structures, for example “a saturated or mono- or poly-unsaturated C₃-C₁₄ mono- or fused-polycyclic hydrocarbyl optionally containing one, two, or three annular heteroatoms per ring” means a mono- or polycyclic (for example a bridged bicyclic) ring system, having between three and fourteen-ring atoms, that contains only carbon ring atoms, but optionally can contain up to three heteratoms per ring and/or unsaturation.
- [0113] “Optional” or “optionally” means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. It will be understood by those skilled in the art with respect to any group containing one or more substituents that such groups are not intended to introduce any substitution or substitution patterns (for example, substituted alkyl includes optionally substituted cycloalkyl groups,

which in turn are defined as including optionally substituted alkyl groups) that are sterically impractical and/or synthetically non-feasible. “Optionally substituted” refers to all subsequent modifiers in a term, for example in the term “optionally substituted arylC₁₋₈ alkyl,” optional substitution may occur on both the “C₁₋₈ alkyl” portion and the “aryl” portion of the molecule; and for example, optionally substituted alkyl includes optionally substituted cycloalkyl groups, which in turn are defined as including optionally substituted alkyl groups. Examples of optional substitution include, but are not limited to alkyl, halogen, alkoxy, hydroxy, oxo, carbamyl, acylamino, sulfonamido, carboxy, alkoxycarbonyl, acyl, alkylthio, alkylsulfonyl, nitro, cyano, amino, alkylamino, cycloalkyl and the like.

[0114] “Saturated bridged ring system” refers to a bicyclic or polycyclic ring system that is not aromatic. Such a system may contain isolated or conjugated unsaturation, but not aromatic or heteroaromatic rings in its core structure (but may have aromatic substitution thereon). For example, 2,3,3a,4,7,7a-hexahydro-1H-indene, 7-aza-bicyclo[2.2.1]heptane and 1,2,3,4,4a,5,8,8a-octahydro-naphthalene are all included in the class “saturated bridged ring system.”

[0115] “Spirocyclyl” or “spirocyclic ring” refers to a ring originating from a particular annular carbon of another ring. For example, a ring atom of the aforementioned saturated bridged ring system, but not a bridgehead atom, can be a shared atom between the saturated bridged ring system (rings B and B’) and a spirocyclyl (ring A) attached thereto, as depicted below. A spirocyclyl can be either carbocyclic or heterocyclic



[0116] “Substituted” alkyl, aryl, and heterocyclyl, refer respectively to alkyl, aryl, and heterocyclyl, wherein one or more (for example up to about five, in another example, up to about three) hydrogen atoms are replaced by a substituent independently selected from the group: optionally substituted alkyl (for example, fluoroalkyl), optionally substituted alkoxy, alkylendioxy (for example methylenedioxy), optionally substituted amino (for example, alkylamino and dialkylamino), optionally substituted amidino, optionally substituted aryl (for example, phenyl), optionally substituted arylalkyl (for example, benzyl), optionally substituted aryloxy (for example, phenoxy), optionally substituted arylalkyloxy (for example, benzyloxy), carboxy (-COOH), carboalkoxy (i.e., acyloxy or

-OOCR), carboxyalkyl (i.e., esters or -COOR), carboxamido, aminocarbonyl, benzyloxycarbonylamino (CBZ-amino), cyano, carbonyl, halogen, hydroxy, optionally substituted heterocyclalkyl, optionally substituted heterocycl, nitro, sulfanyl, sulfinyl, sulfonyl, and thio.

[0117] "Sulfanyl" refers to the groups: -S-(optionally substituted alkyl), -S-(optionally substituted aryl), and -S-(optionally substituted heterocycl).

[0118] "Sulfinyl" refers to the groups: -S(O)-H, -S(O)-(optionally substituted alkyl), -S(O)-(optionally substituted aryl), and -S(O)-(optionally substituted heterocycl).

[0119] "Sulfonyl" refers to the groups: -S(O₂)-H, -S(O₂)-(optionally substituted alkyl), -S(O₂)-(optionally substituted aryl), -S(O₂)-(optionally substituted heterocycl), -S(O₂)-(optionally substituted alkoxy), -S(O₂)-(optionally substituted aryloxy), and -S(O₂)-(optionally substituted heterocycloxy).

[0120] "Yield" for each of the reactions described herein is expressed as a percentage of the theoretical yield.

[0121] In some embodiments, as will be appreciated by those skilled in the art, two adjacent groups on an aromatic system may be fused together to form a ring structure. The fused ring structure may contain heteroatoms and may be optionally substituted with one or more groups. It should additionally be noted that saturated carbons of such fused groups (i.e. saturated ring structures) can contain two substitution groups.

[0122] Some of the compounds of the invention may have imino, amino, oxo or hydroxy substituents off aromatic heterocycl systems. For purposes of this disclosure, it is understood that such imino, amino, oxo or hydroxy substituents may exist in their corresponding tautomeric form, i.e., amino, imino, hydroxy or oxo, respectively.

[0123] Compounds of the invention are generally named using ACD/Name (available from Advanced Chemistry Development, Inc. of Toronto, Canada). This software derives names from chemical structures according to systematic application of the nomenclature rules agreed upon by the International Union of Pure and Applied Chemistry (IUPAC), International Union of Biochemistry and Molecular Biology (IUBMB), and the Chemical Abstracts Service (CAS).

[0124] The compounds of the invention, or their pharmaceutically acceptable salts, may have asymmetric carbon atoms, oxidized sulfur atoms or quaternized nitrogen atoms in their structure.

[0125] The compounds of the invention and their pharmaceutically acceptable salts may exist as single stereoisomers, racemates, and as mixtures of enantiomers and

diastereomers. The compounds may also exist as geometric isomers. All such single stereoisomers, racemates and mixtures thereof, and geometric isomers are intended to be within the scope of this invention.

[0126] It is assumed that when considering generic descriptions of compounds of the invention for the purpose of constructing a compound, such construction results in the creation of a stable structure. That is, one skilled in the art would recognize that there can theoretically be some constructs which would not normally be considered as stable compounds. Stable constructs for the saturated bridged ring system as represented by X^1 , X^2 , and optionally X^3 of formula **II** include but are not limited to motifs such as: 1) where heteroatoms of a ring or bridge thereon are bonded directly to each other, for example a bridge containing a sulfonamide, 2) where heteroatoms of a ring or bridge thereon are separated by only one carbon, for example a urea, carbamate, acetal, aminal, thioacetal, thioaminal, amidine, guanidine, and the like, 3) where heteroatoms of a ring or bridge thereon are separated by two or more carbons, for example an $-NHCH_2CH_2O-$ bridge, and the like, and 4) where heteroatoms in the bridged ring system are separated by more than two carbon atoms, for example wherein the bridged ring system is a decahydro-isoquinoline.

[0127] When a particular group with its bonding structure is denoted as being bonded to two partners, for example a linking group such as $-OCH_2-$, then it is understood that either of the two partners may be bound to the particular group at one end, and the other partner is necessarily bound to the other end of the particular group, unless stated explicitly otherwise.

[0128] Methods for the preparation and/or separation and isolation of single stereoisomers from racemic mixtures or non-racemic mixtures of stereoisomers are well known in the art. For example, optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When desired, the R- and S-isomers may be resolved by methods known to those skilled in the art, for example by: formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallization; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral

solvent. It will be appreciated that where a desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation. For a mixture of enantiomers, enriched in a particular enantiomer, the major component enantiomer may be further enriched (with concomitant loss in yield) by recrystallization.

[0129] "Patient" for the purposes of the present invention includes humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In a preferred embodiment the patient is a mammal, and in a most preferred embodiment the patient is human.

[0130] "Kinase-dependent diseases or conditions" refer to pathologic conditions that depend on the activity of one or more protein kinases. Kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion, migration, differentiation and invasion. Diseases associated with kinase activities include tumor growth, the pathologic neovascularization that supports solid tumor growth, and associated with other diseases where excessive local vascularization is involved such as ocular diseases (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

[0131] While not wishing to be bound to theory, phosphatases can also play a role in "kinase-dependent diseases or conditions" as cognates of kinases; that is, kinases phosphorylate and phosphatases dephosphorylate, for example protein substrates. Therefore compounds of the invention, while modulating kinase activity as described herein, may also modulate, either directly or indirectly, phosphatase activity. This additional modulation, if present, may be synergistic (or not) to activity of compounds of the invention toward a related or otherwise interdependent kinase or kinase family. In any case, as stated previously, the compounds of the invention are useful for treating diseases characterized in part by abnormal levels of cell proliferation (i.e. tumor growth), programmed cell death (apoptosis), cell migration and invasion and angiogenesis associated with tumor growth.

[0132] "Therapeutically effective amount" is an amount of a compound of the invention, that when administered to a patient, ameliorates a symptom of the disease. The amount

of a compound of the invention which constitutes a “therapeutically effective amount” will vary depending on the compound, the disease state and its severity, the age of the patient to be treated, and the like. The therapeutically effective amount can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

[0133] “Cancer” refers to cellular-proliferative disease states, including but not limited to:

Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiforme, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian

carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, SertoliLeydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Karposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal lands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

[0134] "Pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, as well as organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

[0135] "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Exemplary salts are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins, and the like. Exemplary organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine,

choline, and caffeine. (See, for example, S. M. Berge, et al., "Pharmaceutical Salts," J. Pharm. Sci., 1977;66:1-19)

- [0136] "Prodrug" refers to compounds that are transformed (typically rapidly) *in vivo* to yield the parent compound of the above formulae, for example, by hydrolysis in blood. Common examples include, but are not limited to, ester and amide forms of a compound having an active form bearing a carboxylic acid moiety. Examples of pharmaceutically acceptable esters of the compounds of this invention include, but are not limited to, alkyl esters (for example with between about one and about six carbons) wherein the alkyl group is a straight or branched chain. Acceptable esters also include cycloalkyl esters and arylalkyl esters such as, but not limited to benzyl. Examples of pharmaceutically acceptable amides of the compounds of this invention include, but are not limited to, primary amides, and secondary and tertiary alkyl amides (for example with between about one and about six carbons). Amides and esters of the compounds of the present invention may be prepared according to conventional methods. A thorough discussion of prodrugs is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.
- [0137] "Metabolite" refers to the break-down or end product of a compound or its salt produced by metabolism or biotransformation in the animal or human body; for example, biotransformation to a more polar molecule such as by oxidation, reduction, or hydrolysis, or to a conjugate (see Goodman and Gilman, "The Pharmacological Basis of Therapeutics" 8.sup.th Ed., Pergamon Press, Gilman et al. (eds), 1990 for a discussion of biotransformation). As used herein, the metabolite of a compound of the invention or its salt may be the biologically active form of the compound in the body. In one example, a prodrug may be used such that the biologically active form, a metabolite, is released *in vivo*. In another example, a biologically active metabolite is discovered serendipitously, that is, no prodrug design *per se* was undertaken. An assay for activity of a metabolite of a compound of the present invention is known to one of skill in the art in light of the present disclosure.
- [0138] In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

[0139] In addition, it is intended that the present invention cover compounds made either using standard organic synthetic techniques, including combinatorial chemistry or by biological methods, such as bacterial digestion, metabolism, enzymatic conversion, and the like.

[0140] "Treating" or "treatment" as used herein covers the treatment of a disease-state in a human, which disease-state is characterized by abnormal cellular proliferation, and invasion and includes at least one of: (i) preventing the disease-state from occurring in a human, in particular, when such human is predisposed to the disease-state but has not yet been diagnosed as having it; (ii) inhibiting the disease-state, i.e., arresting its development; and (iii) relieving the disease-state, i.e., causing regression of the disease-state. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

General Administration

[0141] Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally (intravenous, intramuscular, or subcutaneous), topically, transdermally, intravaginally, intravesically, intracistemally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages.

[0142] The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, etc. Compositions of the invention may be used in combination with anticancer or other agents that are generally administered to a patient being treated for cancer. Adjuvants include preserving, wetting, suspending, sweetening, flavoring, perfuming, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars,

sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0143] If desired, a pharmaceutical composition of the invention may also contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, etc.

[0144] Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

[0145] One preferable route of administration is oral, using a convenient daily dosage regimen that can be adjusted according to the degree of severity of the disease-state to be treated.

[0146] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, cellulose derivatives, starch, alginates, gelatin, polyvinylpyrrolidone, sucrose, and gum acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, croscarmellose sodium, complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, magnesium stearate and the like (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

- [0147] Solid dosage forms as described above can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They may contain pacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedded compositions that can be used are polymeric substances and waxes. The active compounds can also be in microencapsulated form, if appropriate, with one or more of the above-mentioned excipients.
- [0148] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. Such dosage forms are prepared, for example, by dissolving, dispersing, etc., a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like; solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide; oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols and fatty acid esters of sorbitan; or mixtures of these substances, and the like, to thereby form a solution or suspension.
- [0149] Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances, and the like.
- [0150] Compositions for rectal administrations are, for example, suppositories that can be prepared by mixing the compounds of the present invention with for example suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol or a suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore, melt while in a suitable body cavity and release the active component therein.
- [0151] Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is admixed under sterile conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

[0152] Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. In one example, the composition will be between about 5% and about 75% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

[0153] Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pa., 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state in accordance with the teachings of this invention.

[0154] The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount which will vary depending upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of the compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular disease-states, and the host undergoing therapy. The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per day. For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is an example. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well known to those skilled in the art.

Utility of compounds of the invention as screening agents

[0155] To employ the compounds of the invention in a method of screening for candidate agents that bind to, for example ephrin or EGFR receptor kinase, the protein is bound to a support, and a compound of the invention is added to the assay. Alternatively, the compound of the invention is bound to the support and the protein is added. Classes of candidate agents among which novel binding agents may be sought include specific

antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[0156] The determination of the binding of the candidate agent to, for example ephrin or EGFR protein, may be done in a number of ways. In one example, the candidate agent (the compound of the invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, thus may be done by attaching all or a portion of ephrin or EGFR protein to a solid support, adding a labeled agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

[0157] By “labeled” herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, for example, radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[0158] In some embodiments, only one of the components is labeled. For example, ephrin or EGFR protein may be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the candidate agents.

[0159] The compounds of the invention may also be used as competitors to screen for additional drug candidates. “Candidate bioactive agent” or “drug candidate” or grammatical equivalents as used herein describe any molecule, for example, protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular

proliferation protein binding and/or activity is screened. In the case where protein binding or activity is screened, some embodiments exclude molecules already known to bind to that particular protein. Exemplary embodiments of assays described herein include candidate agents, which do not bind the target protein in its endogenous native state, termed herein as “exogenous” agents. In one example, exogenous agents further exclude antibodies to ephrin or EGFR.

[0160] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules having a molecular weight of more than about 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, for example at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclyl structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs, or combinations thereof.

[0161] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

[0162] In one example, the binding of the candidate agent is determined through the use of competitive binding assays. In this example, the competitor is a binding moiety known to bind to ephrin or EGFR, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[0163] In some embodiments, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to ephrin or EGFR for a time sufficient to allow

binding, if present. Incubations may be performed at any temperature that facilitates optimal activity, typically between 4°C and 40°C.

[0164] Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[0165] In one example, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to ephrin or EGFR and thus is capable of binding to, and potentially modulating, the activity of ephrin or EGFR. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[0166] In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to ephrin or EGFR with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to ephrin or EGFR.

[0167] It may be of value to identify the binding site of, for example, ephrin or EGFR. This can be done in a variety of ways. In one embodiment, once ephrin or EGFR has been identified as binding to the candidate agent, ephrin or EGFR is fragmented or modified and the assays repeated to identify the necessary components for binding.

[0168] Modulation is tested by screening for candidate agents capable of modulating the activity of ephrin or EGFR comprising the steps of combining a candidate agent with ephrin or EGFR, as above, and determining an alteration in the biological activity of ephrin or EGFR. Thus, in this embodiment, the candidate agent should both bind to (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both *in vitro* screening methods and *in vivo* screening of cells for alterations in cell viability, morphology, and the like.

[0169] Alternatively, differential screening may be used to identify drug candidates that bind to native ephrin or EGFR, but cannot bind to modified ephrin or EGFR.

[0170] Positive controls and negative controls may be used in the assays. For example, all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of samples is for a time sufficient for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[0171] A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, for example, albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

[0172] Abbreviations and their Definitions

The following abbreviations and terms have the indicated meanings throughout:

Ac	=	acetyl
ATP	=	adenosine triphosphate
BNB	=	4-bromomethyl-3-nitrobenzoic acid
Boc	=	t-butyloxycarbonyl
br	=	broad
Bu	=	butyl
C	=	degrees Celsius
c-	=	cyclo
CBZ	=	carbobenzoxyl = benzyloxycarbonyl
d	=	doublet
dd	=	doublet of doublet
dt	=	doublet of triplet
DBU	=	diazabicyclo[5.4.0]undec-7-ene
DCM	=	dichloromethane = methylene chloride = CH ₂ Cl ₂


DCE	=	1,2-dichloroethane
DEAD	=	diethyl azodicarboxylate
DIC	=	diisopropylcarbodiimide
DIEA	=	N,N-diisopropylethylamine
DMAP	=	4-N,N-dimethylaminopyridine
DMF	=	N,N-dimethylformamide
DMSO	=	dimethyl sulfoxide
DVB	=	1,4-divinylbenzene
EEDQ	=	2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline
EI	=	Electron Impact ionization
Et	=	ethyl
Fmoc	=	9-fluorenylmethoxycarbonyl
g	=	gram(s)
GC	=	gas chromatography
h or hr	=	hour(s)
HATU	=	0-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HMDS	=	hexamethyldisilazane
HOAc	=	acetic acid
HOBt	=	hydroxybenzotriazole
HPLC	=	high pressure liquid chromatography
L	=	liter(s)
M	=	molar or molarity
m	=	multiplet
Me	=	methyl
mesyl	=	methanesulfonyl
mg	=	milligram(s)
MHz	=	megahertz (frequency)

Min	=	minute(s)
mL	=	milliliter(s)
mM	=	millimolar
mmol	=	millimole(s)
mol	=	mole(s)
MS	=	mass spectral analysis
MTBE	=	methyl t-butyl ether
N	=	normal or normality
NBS	=	N-bromosuccinimide
NCS	=	N-chlorosuccinimide
nM	=	nanomolar
NMO	=	N-methylmorpholine oxide
NMR	=	nuclear magnetic resonance spectroscopy
PEG	=	polyethylene glycol
pEY	=	poly-glutamine, tyrosine
Ph	=	phenyl
PhOH	=	phenol
PfP	=	pentafluorophenol
PfPy	=	pentafluoropyridine
PPTS	=	pyridinium p-toluenesulfonate
Py	=	pyridine
PyBroP	=	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
q	=	quartet
RT	=	room temperature
Sat'd	=	saturated
s	=	singlet
s-	=	secondary
t-	=	tertiary

t or tr	=	triplet
TBDMS	=	t-butyldimethylsilyl
TES	=	triethylsilane
TFA	=	trifluoroacetic acid
THF	=	tetrahydrofuran
TMOF	=	trimethyl orthoformate
TMS	=	trimethylsilyl
tosyl	=	p-toluenesulfonyl
Trt	=	triphenylmethyl
uL	=	microliter(s)
uM	=	micromole(s) or micromolar

Synthesis of Compounds

Scheme 1



[0173] Scheme 1 depicts a general synthetic route for compounds of the invention and is not intended to be limiting. Specific examples are described subsequently to this general synthetic description. A benzoic ester **1**, where R is typically but not necessarily a methyl

radical and P is typically but not necessarily an alkyl group, is O-alkylated at the oxygen *para* to the carboxylate group with an electrophile to afford a substituted derivative 2. P is typically a lower alkyl group, but can be a protecting group that is removed later in a synthesis. When P is a lower alkyl group it can possess functionality initially, or be derivitized to contain such functionality at various stages of the synthesis. The group, E¹, represents either a protecting group, for example benzyl, or a group that either has moieties present in compounds of the invention or possesses functionality that serve as precursors to such groups. Aromatic ring nitration and reduction of the corresponding nitro group are carried out in a regio- and chemoselective manner by methods well known in the art to give anthranilate derivative 3. Formation of quinazolin-4-one 4 is carried out by methods well known in the art, for example by heating 3 in formamide solution in the presence of ammonium formate, or for another example by heating directly with formamidine hydrochloride. Introduction of 4-position functionality is carried out by methods known in the art. For example, quinazolin-4-one 4 is converted to an intermediate quinazoline 5, where "L" represents a leaving group, for example chlorine. Quinazoline 5 is then converted to 6 by reaction with a range of nucleophiles, for example amines, alcohols, and thiols. After formation of 6, group "Z" is either left "as is" or converted at some subsequent stage to a derivative thereof. For example when Z is -NH-, then the hydrogen on the nitrogen may optionally be replaced with an alkyl group, or when Z is sulfur, then that sulfur atom may be oxidized to, for example, a sulfone. Compound 6 may represent a compound of the invention or, for example when E¹ serves as a protecting group, E¹ may be removed to provide phenol 7. Introduction of a group E² is carried out by methods well established in the art, for example alkylation with an appropriately derivitized alkyl halide.

Examples

[0174] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. Generally, each example set out below describes a multi-step synthesis as outlined above.

Example 1

1,4:3,6-Dianhydro-2-*O*-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol

[0175] 1,4:3,6-dianhydro-2-*O*-methyl-5-*O*-(methanesulfonyl)-D-glucitol: To a solution of 1,4:3,6-dianhydro-2-*O*-methyl-D-glucitol (1.19g, 7.4 mmol) in dichloromethane was added pyridine (1mL, 12.36 mmol) followed by methanesulfonyl chloride (0.69mL, 8.92 mmol) and the mixture was allowed to stir at room temperature over 12 hours. The solvent was removed and the amorphous residue was partitioned between ethyl acetate and aqueous hydrochloric acid (for example 0.1M HCl). The aqueous phase was extracted once with additional ethyl acetate and the combined organic layers were washed with saturated aqueous sodium chloride then dried over anhydrous magnesium sulfate. Filtration and concentration followed by drying *in vacuo* afforded 1,4:3,6-dianhydro-2-*O*-methyl-5-*O*-(methanesulfonyl)-D-glucitol (1.67g, 94% yield) as a colorless oil. GCMS calculated for C₈H₁₄SO₆: 238 (M⁺).

[0176] 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol: 4-Chloro-6-(methyloxy)-7-[(phenylmethyl)oxy]quinazoline hydrochloride (22.91g, 67.9 mmol) was suspended in isopropanol followed by addition of 3,4-dichloroaniline (13.2g, 81.5 mmol) and concentrated aqueous hydrochloric acid (1mL). The mixture was brought to reflux over 12 hours and diluted with ethyl ether (150mL). The solid was collected by filtration, washed with additional ethyl ether and dried. The material was then taken into trifluoroacetic acid (150mL) and brought to reflux over 1 hour. The solution was cooled to room temperature then concentrated *in vacuo* to give a crystalline residue. The residue was suspended in acetonitrile (100mL) followed by addition of ethyl ether (100mL). The solid was collected by filtration and washed with additional ethyl ether then dried *in vacuo* to give 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol (21.49g, 64% yield) as a tan solid. ¹H NMR (400 MHz, d₆-DMSO): 11.09 (br s, 1H), 8.87 (s, 1H), 8.07 (d, 1H), 8.00 (s, 1H), 7.23 (s, 1H), 3.98 (s, 3H); MS (EI) for C₁₅H₁₁N₃O₂Cl₂: 337 (MH⁺).

[0177] 1,4:3,6-dianhydro-2-*O*-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol hydrochloride: A suspension of 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol (1.70g, 3.78 mmol), 1,4:3,6-dianhydro-2-*O*-methyl-5-*O*-(methanesulfonyl)-D-glucitol (1.00g, 4.20 mmol), and potassium carbonate (2.64g, 19.10 mmol) in DMF (20mL) was stirred at 80°C under nitrogen for 15 h. The reaction mixture was poured into water (100mL), and extracted

with ethyl acetate (3 x 50mL). The organic layers were washed with 5% LiCl (2 x 50mL), and brine (50mL) then dried over anhydrous sodium sulfate. Filtration, concentration and column chromatography on silica (97:3 dichloromethane/methanol) gave a solid, which was dissolved in methanol (50mL), and treated with 4M HCl in 1,4-dioxane (5mL). The resulting precipitation was filtered, washed with methanol (2 x 20mL), and dried to afford 0.99g (51%) of 1,4:3,6-dianhydro-2-*O*-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol hydrochloride as a yellow solid. ¹H NMR (400 MHz, d₆-DMSO): 11.51 (br s, 1H), 8.91 (s, 1H), 8.38 (s, 1H), 8.16 (d, 1H), 7.82 (dd, 1H), 7.76 (d, 1H), 7.42 (s, 1H), 5.03 (m, 1H), 4.66 (m, 2H), 4.11 (m, 1H), 4.04 (s, 3H), 4.02 (m, 1H), 3.90 (m, 3H), 3.31 (s, 3H); MS (EI) for C₂₂H₂₁N₃O₅Cl₂: 478 (MH⁺).

[0178] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0179] 1,4:3,6-dianhydro-2-*O*-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ¹H NMR (400 MHz, d₆-DMSO): 8.76 (s, 1H), 8.09 (s, 1H), 7.79 (dd, 1H), 7.55 (t, 1H), 7.39 (s, 1H), 5.10-5.06 (m, 1H), 4.65 (s, 2H), 4.11 (dd, 1H), 4.04-4.02 (m, 1H), 4.00 (s, 3H), 3.94-3.91 (m, 1H), 3.90-3.87 (m, 2H), 3.31 (s, 3H); MS (EI) for C₂₂H₂₀BrClFN₃O₅: 540 (MH⁺).

[0180] 1,4:3,6-dianhydro-2-*O*-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ¹H NMR (400 MHz, d₆-DMSO): 8.77 (s, 1H), 8.13 (s, 1H), 7.96 (d, 1H), 7.56 (d, 1H), 7.42 (s, 1H), 5.10-5.06 (m, 1H), 4.66 (s, 2H), 4.12 (dd, 1H), 4.05-4.02 (m, 1H), 4.00 (s, 3H), 3.94-3.91 (m, 1H), 3.90-3.87 (m, 2H), 3.31 (s, 3H); MS (EI) for C₂₂H₂₀BrCl₂N₃O₅: 556 (MH⁺).

[0181] 1,4:3,6-Dianhydro-2-*O*-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ¹H NMR (400 MHz, d₆-DMSO): 9.63 (s, 1H), 8.30 (s, 1H), 7.86 (s, 1H), 7.41 (m, 1H), 7.30 (m, 2H), 7.26 (s, 1H), 5.05 (br s, 1H), 4.63 (dd, 2H), 4.03 (ddd AB, 2H), 3.95 (s, 3H), 3.91 (s, 1H), 3.86 (d, 2H), 3.31 (s, 3H), 2.19 (s, 3H); MS (EI) for C₂₃H₂₄N₃O₅Cl: 458 (MH⁺).

[0182] 1,4:3,6-Dianhydro-2-*O*-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ¹H NMR (400 MHz, d₆-DMSO): 9.69 (s, 1H), 8.42 (s, 1H), 7.94-7.91 (m, 2H), 7.82 (s, 1H), 7.29 (s, 1H), 5.06 (br s, 1H), 4.63 (dd, 2H), 4.03 (ddd AB, 2H), 3.95 (s, 3H), 3.91-3.86 (m, 3H), 3.31 (s, 3H); MS (EI) for C₂₂H₂₀N₃O₅BrClF: 542 (MH⁺).

- [0183] 1,4:3,6-Dianhydro-2-*O*-[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 11.20 (br s, 1H), 8.81 (s, 1H), 8.06 (s, 1H), 7.68-7.62 (m, 1H), 7.51 (d tr, 1H), 7.43 (s, 1H), 5.09 (br s, 1H), 4.66 (s, 2H), 4.07 (ddd AB 2H), 3.94 (s, 3H), 4.00-3.88 (m, 3H) 3.31 (s, 3H); MS (EI) for $\text{C}_{22}\text{H}_{20}\text{N}_3\text{O}_5\text{ClF}_2$: 480 (MH^+).
- [0184] 1,4:3,6-Dianhydro-2-*O*-[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.69 (s, 1H), 8.42 (s, 1H), 7.94 (d, 1H), 7.84 (d, 1H), 7.82 (s, 1H), 7.29 (s, 1H), 4.63 (m, 2H), 4.03 (ddd AB, 2H), 3.95 (s, 3H), 3.91-3.86 (m, 3H), 3.31 (s, 3H); MS (EI) for $\text{C}_{22}\text{H}_{20}\text{N}_3\text{O}_5\text{Cl}_2\text{F}$: 496 (MH^+).
- [0185] 1,4:3,6-Dianhydro-2-*O*-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.68 (s, 1H), 8.37 (s, 1H), 7.82 (s, 1H), 7.52-7.45 (m, 2H), 7.27 (d tr, 1H), 5.04 (br s, 1H), 4.63 (dd, 2H), 4.02 (ddd AB, 2H), 3.94 (s, 3H), 3.89 (br s, 1H), 3.87 (d 2H), 3.30 (s, 3H); MS (EI) for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_5\text{ClF}$: 462 (MH^+).
- [0186] 1,4:3,6-Dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.58 (s, 1H), 8.52 (s, 1H), 8.23 (d, 1H), 7.82-7.79 (m, 2H) 7.74 (d, 1H), 7.26 (s, 1H) 5.03 (br s, 1H), 4.62 (dd AB, 2H), 4.02 (ddd AB, 2H), 3.96 (s, 3H), 3.89 (br s, 1H), 3.85 (d, 2H) 3.30 (s, 3H); MS (EI) for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_5\text{BrCl}$: 524 (MH^+).
- [0187] 1,4:3,6-Dianhydro-2-*O*-[4-[(3-bromo-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.65 (s, 1H), 8.30 (s, 1H), 7.86 (s, 1H), 7.57 (dd, 1H), 7.33 (d, 1H), 7.26-7.21 (m, 2H), 5.04 (br s, 1H), 4.63 (dd AB, 2H), 4.03 (ddd AB, 2H), 3.94 (s, 3H), 3.91 (br s, 1H), 3.87 (d, 2H), 3.31 (s, 3H); MS (EI) for $\text{C}_{23}\text{H}_{24}\text{N}_3\text{O}_5\text{Br}$: 502 (MH^+).
- [0188] 1,4:3,6-Dianhydro-5-*O*-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-*O*-methyl-D-glucitol: MS (EI) for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_5\text{Cl}_2$: 478 (MH^+).
- [0189] 1,4:3,6-Dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-(difluoromethyl)-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.64 (s, 1H), 8.56 (s, 1H), 8.26 (d, 1H), 7.86-7.82 (m, 2H), 7.77 (d, 1H), 7.31 (s, 1H), 6.84 (tr, 1H), 5.12 (br s, 1H), 4.74 (m, 2H), 4.06 (ddd AB, 2H), 3.98-3.90 (m, 6H); MS (EI) for $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_5\text{BrClF}_2$: 558 (MH^+).
- [0190] 1,4:3,6-Dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-ethyl-L-iditol: ^1H NMR (400 MHz, $\text{d}_6\text{-DMSO}$): 9.62

(s, 1H), 8.55 (s, 1H), 8.26 (d, 1H), 7.85-7.82 (m, 2H), 7.76 (d, 1H), 5.04 (br s, 1H), 4.62 (dd AB, 2H), 4.15-4.09 (m, 1H), 4.00-3.95 (m, 5H), 3.95-3.82 (m, 2H), 3.57-3.48 (m, 2H), 1.13 (tr, 3H); MS (EI) for $C_{23}H_{23}N_3O_5BrCl$: 536 (MH^+).

[0191] 1,4:3,6-Dianhydro-5-deoxy-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-*D*-xylo-hexitol: ^1H NMR (400 MHz, d_6 -DMSO): 9.62 (s, 1H), 8.55 (s, 1H), 8.26 (s, 1H), 7.85-7.82 (m, 2H), 7.76 (d, 1H), 7.30 (s, 1H), 4.98 (m, 1H), 4.80 (m, 1H), 4.51 (m, 1H), 4.18-4.14 (m, 1H), 3.97 (s, 3H), 3.92-3.81 (m, 3H), 2.01-1.97 (m, 2H); MS (EI) for $C_{21}H_{19}N_3O_4BrCl$: 492 (MH^+).

[0192] 1,4:3,6-Dianhydro-2-*O*-methyl-5-*O*-{6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl}-*L*-iditol: ^1H NMR (400MHz; DMSO- d_6): 9.82 (br s, 1H), 8.36 (s, 1H), 8.86 (s, 1H), 7.75-7.73 (d, 1H), 7.60-7.58 (d, 1H), 7.29 (s, 1H), 5.06 (br s, 1H), 5.64-5.62 (m, 2H), 4.10-4.07 (dd, 1H), 4.02-4.01 (d, 1H), 3.97-3.94 (m, 1H), 3.95 (s, 3H), 3.93-3.90 (m, 1H), 3.88 (br m, 1H), 3.31 (s, 3H); MS (EI) for $C_{22}H_{20}Cl_3N_3O_5$: 511.91 (MH^+).

[0193] 1,4:3,6-Dianhydro-2-*O*-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-*O*-methyl-*L*-iditol hydrochloride: ^1H NMR (400MHz; DMSO- d_6): 8.75 (s, 1H), 8.03 (s, 1H), 7.70-7.53 (m, 2H), 7.38 (s, 1H), 5.09-5.07 (m, 1H), 4.64-4.63 (br, 1H), 4.13-4.10 (dd, 1H), 4.02-4.01 (d, 1H), 3.99 (s, 3H), 3.93-3.92 (m, 1H), 3.89-3.88 (m, 2H), 3.31 (s, 3H); MS (EI) for $C_{22}H_{20}Cl_2FN_3O_5$: 495.96 (MH^+).

[0194]

Example 2

1,4:3,6-Dianhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)-quinazolin-7-yl]-*L*-sorbose ethylene glycol acetal

[0195] 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-*D*-fructose ethylene glycol acetal: A solution of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-*D*-fructose (2.00g, 8.06 mmol), ethylene glycol (5.00g, 80.6 mmol), and *p*-toluenesulfonic acid (1.53g, 8.06 mmol) in benzene (100mL) was refluxed for 90 min using a Dean-Stark Trap apparatus. The reaction mixture was diluted with ethyl acetate (100mL), washed with saturated aqueous

sodium bicarbonate (2 x 50mL) then brine (50mL), and dried over anhydrous sodium sulfate. Filtration, concentration and column chromatography on silica (1:1 hexane/ethyl acetate) provided 1.44g (61% yield) of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose ethylene glycol acetal as a colorless solid. ¹H NMR (400 MHz; CDCl₃): 8.08 (m, 2H), 7.58 (m, 1H), 7.54 (m, 2H), 5.38 (dd, 1H), 4.97 (t, 1H), 4.21-4.02 (m, 7H), 3.86 (d, 1H), 3.75 (d, 1H).

[0196] 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal: To a solution of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose ethylene glycol acetal (1.44g, 4.93 mmol) in methanol (40mL) was added 50% aqueous sodium hydroxide (0.38 g, 4.75 mmol) and the mixture was stirred at room temperature for 30 minutes. Neutralization with 1M HCl, followed by concentration and column chromatography on silica (1:2 hexane/ethyl acetate) provided 0.74g (80% yield) of 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal as a colorless solid. ¹H NMR (400 MHz; CDCl₃): 4.60 (t, 1H), 4.32 (m, 1H), 4.14 (d, 1H), 4.05-3.98 (m, 5H), 3.82 (s, 2H), 3.62 (dd, 1H), 2.65 (d, 1H).

[0197] 1,4:3,6-dianhydro-5-*O*-(methanesulfonyl)-D-fructose ethylene glycol acetal: To a solution of 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal (0.74g, 3.93 mmol) and triethylamine (1.20g, 11.86 mmol) in dichloromethane (40mL) was added methanesulfonyl chloride (0.90g, 7.88 mmol) at 0°C under nitrogen. The solution was warmed to room temperature and stirred for 13 h. Dichloromethane (50mL) was added, and the organic layer was washed with saturated aqueous sodium bicarbonate (30mL), water (30mL), and brine (30mL) then dried over anhydrous sodium sulfate. Filtration and concentration provided 1.02g (97%) of 1,4:3,6-dianhydro-5-*O*-(methanesulfonyl)-D-fructose ethylene glycol acetal as a yellow oil. ¹H NMR (400 MHz; CDCl₃): 5.08 (m, 1H), 4.82 (t, 1H), 4.13 (dd, 1H), 4.04 (m, 4H), 3.93 (dd, 1H), 3.87 (d, 1H), 3.81 (d, 1H), 3.13 (s, 3H).

[0198] 1,4:3,6-dianhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)-quinazolin-7-yl]-L-sorbose ethylene glycol acetal: A suspension of 4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol (235mg, 0.48 mmol), 1,4:3,6-dianhydro-5-*O*-(methanesulfonyl)-D-fructose ethylene glycol acetal (190mg, 0.71 mmol), and potassium carbonate (329mg, 2.38 mmol) in DMF (10 mL) was stirred at 130°C under nitrogen for 14 h. The reaction mixture was poured into water (50mL), and extracted with ethyl acetate (3 x 30mL). The organic layers were washed with 5% LiCl (2 x 25mL), and brine (25mL) then dried over anhydrous sodium sulfate and concentrated. Filtration and column chromatography on silica (9:1

dichloromethane/acetone to 7:3 dichloromethane/acetone) gave 77mg (29%) of 1,4:3,6-dianhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]-L-sorbose ethylene glycol acetal as a off-white solid. ¹H NMR (400 MHz, CDCl₃): 8.70 (s, 1H), 8.00 (d, 1H), 7.61 (d, 1H), 7.52 (dd, 1H), 7.31 (s, 1H), 7.14 (s, 1H), 7.00 (s, 1H), 4.98 (m, 1H), 4.86 (d, 1H), 4.42 (d, 1H), 4.32-4.23 (m, 2H), 4.10-4.05 (m, 4H), 4.00 (s, 3H), 3.86 (d, 1H), 3.78 (d, 1H); MS (EI) for C₂₃H₂₁N₃O₆BrCl: 550 (MH⁺).

Example 3

1,4:3,6-Dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)-quinazolin-7-yl]-5-deoxy-5-methylidene-D-xylo-hexitol:

[0199] 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-*arabino*-hexitol: To a solution of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(phenylcarbonyl)-D-*arabino*-hexitol (329mg, 1.34 mmol) in methanol (10mL) was added 50% aqueous sodium hydroxide (95mg, 1.19 mmol) and the mixture was stirred at room temperature for 30 minutes. Neutralization with 4M hydrogen chloride in 1,4-dioxane, followed by concentration and column chromatography on silica (1:1 hexane/ethyl acetate) provided 141mg (74%) of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-*arabino*-hexitol as a colorless solid. ¹H NMR (400 MHz; CDCl₃): 5.37 (m, 1H), 5.20 (m, 1H), 4.80 (m, 1H), 4.54 (m, 2H), 4.43 (m, 1H), 4.26 (m, 1H), 3.95 (dd, 1H), 3.54 (dd, 1H), 2.70 (d, 1H).

[0200] 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(methylsulfonyl)-D-*arabino*-hexitol: To a solution of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-*arabino*-hexitol (135mg, 0.95 mmol) and triethylamine (288mg, 2.85 mmol) in dichloromethane (10mL) was added methanesulfonyl chloride (222mg, 1.94 mmol) at 0°C under nitrogen. The solution was warmed to room temperature and stirred for 18 h. Dichloromethane (50mL) was added and the organic layer was washed with saturated aqueous sodium bicarbonate (2 x 25mL), water (25mL) and brine (25mL) then dried over anhydrous sodium sulfate. Filtration and concentration provided 213mg (72%) of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(methylsulfonyl)-D-*arabino*-hexitol as a yellow oil. ¹H NMR (400 MHz; CDCl₃): 5.40 (m, 1H), 5.23 (m, 1H), 5.04 (m, 1H), 4.85 (m, 1H), 4.73 (t, 1H), 4.58 (m, 1H), 4.41 (m, 1H), 4.08 (dd, 1H), 3.86 (dd, 1H), 3.14 (s, 3H).

[0201] 1,4:3,6-dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)-quinazolin-7-yl]-5-deoxy-5-methylidene-D-xylo-hexitol: A suspension of 4-[(4-bromo-3-chlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (425mg, 0.86 mmol), 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(methylsulfonyl)-D-*arabino*-hexitol (208mg,

0.94 mmol), and potassium carbonate (594mg, 4.30 mmol) in DMF (10 mL) was stirred at 130°C under nitrogen for 15 h. The reaction mixture was poured into water (50mL), and extracted with ethyl acetate (3 x 30mL). The organic layers were washed with 5% LiCl (2 x 25mL), and brine (25mL), dried over anhydrous sodium sulfate then filtered and concentrated. Column chromatography on silica (97:3 dichloromethane/ methanol) gave 234mg (54%) of 1,4:3,6-dianhydro-2-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-5-methylidene-D-xylo-hexitol as a yellow solid. ¹H NMR (400 MHz, CDCl₃): 8.70 (s, 1H), 8.01 (d, 1H), 7.61 (d, 1H), 7.51 (dd, 1H), 7.36 (s, 1H), 7.14 (s, 1H), 7.01 (s, 1H), 5.42 (m, 1H), 5.23 (m, 1H), 5.04 (d, 1H), 4.97 (t, 1H), 4.74 (d, 1H), 4.55 (m, 1H), 4.35 (m, 1H), 4.22 (m, 2H), 4.01 (s, 3H); MS (EI) for C₂₂H₁₉N₃O₄BrCl: 504 (MH⁺).

Example 4

Methyl 3,6-anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-*O*-methyl- α -L-idofuranoside

[0202] To a mixture of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-(D)-glucitol (4.32g, 17.3 mmol), triethylamine (4.91 mL, 35.3 mmol) and 4-dimethylaminopyridine (0.63g, 5.2 mmol) in dichloromethane (50 mL) at -10 ° to -15° was added trifluoromethanesulfonic anhydride (3.48mL, 20.7 mmol) dropwise over ten minutes and the resulting mixture was stirred at this temperature for 3 hours. The mixture was poured into 100 mL of ice-water and extracted with dichloromethane (3 x 50 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered then concentrated. The crude triflate was suspended in toluene (50 mL) followed by addition of 1,8-diazabicyclo[4,5,0]undec-7-ene (5.25 mL, 34.6 mmol) and the mixture was stirred at room temperature for 18 hours. The reaction mixture was poured into ice-water and partitioned then the aqueous portion was extracted with dichloromethane (3 x 50 mL). The combined organic portion was washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by flashed chromatography (silica gel, 5-20% ethyl acetate-hexane) to give 1,4:3,6-dianhydro-2-deoxy-5-*O*-(phenylcarbonyl)-D-*arabino*-hex-1-enitol, as a white solid, 3.10g, 77% yield. ¹H NMR (400MHz; CDCl₃): 8.08-8.06 (m, 2H), 7.61-7.57 (m, 1H), 7.56-7.43 (m, 2H), 6.62-6.61 (d, 1H), 5.48-5.46 (m,1H), 5.32-5.26 (m,1H), 5.13-5.10 (m, 2H), 4.18-4.14 (tr,1H), 3.61-3.56 (tr, 1H).

- [0203] Methyl 3,6-anhydro-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside: To a solution of 1,4:3,6-dianhydro-2-deoxy-5-*O*-(phenylcarbonyl)-D-arabino-hex-1-enitol (1.00g, 4.3 mmol) in methanol (17 mL) at -4°C was added 3-chloroperoxybenzoic acid (85%, 1.35g, 8.6 mmol), and the resulting mixture was slowly warmed to room temperature and stirred for 18 hours. The reaction mixture was concentrated, diluted with dichloromethane (50 mL), washed with saturated aqueous sodium bicarbonate solution, dried over sodium sulfate, filtered and concentrated. The residue was purified by flash chromatography (silica gel, 25-60% ethyl acetate-hexane) to give methyl 3,6-anhydro-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside as a white solid, 1.03g, 83% yield. ¹H NMR (400MHz; CDCl₃): 8.11-8.08 (d, 2H), 7.61-7.56 (tr, 1H), 7.48-7.44 (m, 2H), 5.24-5.17 (m, 2H), 4.96 (s, 1H), 4.57-4.56 (d, 1H), 4.27 (s, 1H), 4.22-4.18 (dd, 1H), 4.08-4.04 (dd, 1H) 3.36 (s, 3H).
- [0204] Methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside: A mixture of methyl 3,6-anhydro-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside (1.03g, 3.7 mmol), silver (I) oxide (0.85g, 3.7 mmol) and methyl iodide (0.34 mL, 5.5 mmol) in DMF (2 mL) was heated at 60°C for 1 hour. After cooling to room temperature the reaction mixture was diluted with ethyl acetate (50 mL), filtered over celite™, adsorbed on silica gel (10g) and purified by flash chromatography (silica gel, 5-30% ethyl acetate-hexane) to give methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside as a colorless oil, 0.82g, 76% yield. ¹H NMR (400MHz; CDCl₃): 8.11-8.09 (d, 2H), 7.60-7.56 (m, 1H), 7.46-7.44 (m, 2H), 5.24-5.20 (m, 1H), 5.18-5.09 (tr, 1H), 4.99 (s, 1H), 4.61-4.60 (d, 1H), 4.21-4.17 (tr, 1H), 4.08-4.03 (tr, 1H), 3.81 (s, 1H), 3.40 (s, 3H), 3.57 (s, 3H).
- [0205] Methyl 3,6-anhydro-2-*O*-methyl- β -D-glucofuranoside: A solution of methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(phenylcarbonyl)- β -D-glucofuranoside (820mg, 3.1mmol) and 50% sodium hydroxide (248 mg, 3.1 mmol) in methanol (10mL) was stirred at room temperature for 30 minutes. The material was adsorbed on silica gel (5g) and passed through a short column (15% ethyl acetate in hexanes to 5% methanol in ethyl acetate) to give methyl 3,6-anhydro-2-*O*-methyl- β -D-glucofuranoside as a colorless oil, 420 mg, 85% yield. ¹H NMR (400MHz; CDCl₃): 5.04 (s, 1H), 5.84-5.81 (tr, 1H), 4.44-4.42 (tr, 1H), 4.25-4.19 (m, 1H), 3.85-3.75 (m, 1H), 3.49 (s, 3H), 3.43 (s, 3H), 2.75-2.72 (d, 1H).
- [0206] Methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(methylsulfonyl)- β -L-glucofuranoside: Methyl 3,6-anhydro-2-*O*-methyl- β -D-glucofuranoside (420 mg, 2.6 mmol) was dissolved

in dichloromethane (10 mL) and pyridine (0.36 mL, 3.7 mmol) at 0°C. Methanesulfonyl chloride (0.14 mL, 3.1 mmol) was added and the resulting mixture was stirred at 0°C for 1 hour then at room temperature for 2 hours. The reaction mixture was washed with water and saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate, filtered and concentrated to give methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(methylsulfonyl)-β-D-glucofuranoside as a colorless oil, 669mg, 95% yield, which was used without further purification.

[0207] Methyl 3,6-anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-*O*-methyl-α-L-idofuranoside: Methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(methylsulfonyl)-β-D-glucofuranoside (314 mg, 1.1 mmol) was dissolved in DMF (3mL). To this solution was added potassium carbonate (404 mg, 2.9 mmol) and 4-[(4-bromo-3-chlorophenyl)amino]-6-methyloxy-7-hydroxyquinazoline trifluoroacetate (280 mg, 0.59 mmol). The resulting mixture was heated at 135°C for 18h. After cooling to room temperature the reaction mixture was diluted with ethyl acetate (15 mL), washed with water, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by flash chromatography (silica gel, 2-7% methanol in 1:1 ethyl acetate:hexanes) to give methyl 3,6-anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-*O*-methyl-α-L-idofuranoside as a white solid, 181 mg, 28% yield. ¹H NMR (400MHz; Methanol-d₄): 8.75 (s, 1H), 8.04–8.06 (d, 1H), 7.99 (s, 1H), 7.78-7.75 (d, 1H), 7.64-7.61 (d, 1H), 7.35 (s, 1H), 5.16-5.14 (d, 1H), 5.02 (s, 1H), 4.89 (br, 1H), 4.69-4.68 (d, 1H) 4.46-4.42 (dd, 1H), 4.09 (br, 1H), 4.06 (s, 3H), 3.69(s, 1H), 3.48(s, 3H), 3.42 (s, 3H); MS (EI) for C₂₃H₂₃BrClN₃O₆: 551.88 (MH⁺).

[0208]

[0209]

[0210]

[0211]

Example 6

3,6-Anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2-*O*-(1-methylethylidene)-β-D-idofuranose

[0212] 3,6-Anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)-α-L-glucofuranose A mixture of 3,6-anhydro-5-*O*-(phenylcarbonyl)-α-L-glucofuranose

(1.00g), 2,2-dimethoxy propane (0.63 mL), p-toluenesulfonic acid (20 mg) and benzene (10 mL) was heated at reflux for 3 hours. The reaction mixture was cooled then adsorbed on silica gel (10g) and purified by flash chromatography (silica gel, 5-35% ethyl acetate in hexanes) to give 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)- α -L-glucofuranose as colorless oil, 0.85g, 74% yield. ^1H NMR (400MHz; CDCl_3): 8.08-8.06 (d, 2H), 7.59-7.56 (tr, 1H), 7.46-7.42 (m, 2H), 5.99-5.98 (d, 1H), 5.35-5.31 (tr, 1H), 5.10-5.08 (d, 1H), 4.66-4.65 (d, 1H), 4.61-4.60 (d, 1H), 4.20-4.16 (dd, 1H), 3.91-3.74 (tr, 1H), 1.50 (s, 3H), 1.34 (s, 3H).

[0213] 3,6-Anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(methylsulfonyl)- α -L-glucofuranose: A solution of 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)- α -L-glucofuranose (850mg) and 50% sodium hydroxide (111 mg) in methanol (10mL) was stirred at room temperature for 30 minutes. The mixture was then adsorbed on silica gel (5g) and passed through a short column (15% ethyl acetate in hexanes to 5% methanol in ethyl acetate) and the alcohol intermediate, 390 mg, 70% yield, was used immediately in the next step. The alcohol was dissolved in dichloromethane (10 mL) and pyridine (0.32 mL) at 0°C. Methanesulfonyl chloride (0.12 mL) was added and the resulting mixture was stirred at 0°C for 1 hour then at room temperature for 2 hours. The reaction mixture was washed with water and saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate, filtered and concentrated to give 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(methylsulfonyl)- α -L-glucofuranose as a colorless oil, 485 mg, 90% yield, which was immediately employed in the next step.

[0214] 3,6-Anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2-*O*-(1-methylethylidene)- β -D-idofuranose: 3,6-Anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(methylsulfonyl)- α -L-glucofuranose, (85 mg, 0.30 mmol) was dissolved in DMF (3mL). To this solution was added potassium carbonate (168 mg, 1.21 mmol) and 4-[(4-bromo-3-chlorophenyl)amino]-6-methyloxy-7-hydroxyquinazoline trifluoroacetate (145 mg, 0.30 mmol). The resulting mixture was heated at 135°C for 18h. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate (15 mL), washed with water, dried over anhydrous sodium sulfate, filtered, concentrated. The residue was purified by flash chromatography (silica gel, 2-7% Methanol in 1:1 ethyl acetate:hexanes) to give 3,6-anhydro-5-*O*-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2-*O*-(1-methylethylidene)- β -D-idofuranose, 121 mg, 77%

yield, as a white solid. ^1H NMR (400MHz; Methanol- d_4): 8.48 (s, 1H), 8.17-8.16 (d, 1H), 7.76 (s, 1H), 7.70-7.61 (m, 2H), 7.19 (s, 1H), 5.95-5.94 (d, 1H), 5.18-5.17 (d, 1H), 4.93-4.91 (m, 1H), 4.70-4.62 (m, 2H), 4.28-4.22 (dd, 1H), 4.08-4.06 (d, 1H), 4.03 (s, 3H), 1.44 (s, 3H), 1.32 (s, 3H); MS (EI) for $\text{C}_{24}\text{H}_{23}\text{BrClN}_3\text{O}_6$: 563.83 (MH^+).

Example 7

(3*S*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine and (3*R*,8*aS*)-3-(chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine

[0215] (S)-(+)-Prolinol (6.00 g, 59.3 mmol) was added to epichlorohydrin (47 mL, 600 mmol) at 0°C. The solution was stirred at 40°C for 0.5 h and then concentrated *in vacuo*. The residual oil was cooled in an ice bath and concentrated sulfuric acid (18 mL) was added dropwise with stirring. The mixture was heated at 170-180°C for 1.5 h, poured into ice (300 mL) and then basified with sodium carbonate to pH~8. The mixture was partitioned with ethyl acetate/hexanes and filtered. The filtrate was separated and the aqueous portion was extracted twice with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford an oil which was purified by column chromatography (ethyl acetate for less polar product and then 30% methanol in ethyl acetate). (3*S*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine (less polar product) (1.87 g, 10.7 mmol, 18% yield): ^1H NMR (400 MHz, CDCl_3): 4.06 (dd, 1H), 3.79-3.71 (m, 1H), 3.60-3.48 (m, 2H), 3.36 (dd, 1H), 3.15 (dd, 1H), 3.13-3.06 (m, 1H), 2.21-2.01 (m, 3H), 1.90-1.68 (m, 3H), 1.39-1.24 (m, 1H); MS (EI) for $\text{C}_8\text{H}_{14}\text{NOCl}$: 176 (MH^+). (3*R*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine (1.54 g, 8.77 mmol, 15% yield): ^1H NMR (400 MHz, CDCl_3): 3.94-3.77 (m, 4H), 3.55 (dd, 1H), 3.02-2.93 (m, 2H), 2.45 (dd, 1H), 2.29-2.15 (m, 2H), 1.88-1.64 (m, 3H), 1.49-1.38 (m, 1H); MS (EI) for $\text{C}_8\text{H}_{14}\text{NOCl}$: 176 (MH^+).

[0216] Using the same or analogous synthetic techniques and/or substituting with alternative starting materials, the following reagents were prepared:

[0217] (3*R*,8*aR*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine: ^1H NMR (400 MHz, CDCl_3): 4.05 (dd, 1H), 3.79-3.70 (m, 1H), 3.61-3.48 (m, 2H), 3.35 (dd, 1H), 3.15 (dd, 1H), 3.13-3.07 (m, 1H), 2.21-2.01 (m, 3H), 1.89-1.67 (m, 3H), 1.39-1.25 (m, 1H); MS (EI) for $\text{C}_8\text{H}_{14}\text{NOCl}$: 176 (MH^+).

[0218] (3*S*,8*aR*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine: ^1H NMR (400 MHz, CDCl_3): 3.93-3.77 (m, 4H), 3.55 (dd, 1H), 3.02-2.93 (m, 2H), 2.45 (dd, 1H),

2.30-2.15 (m, 2H), 1.88-1.64 (m, 3H), 1.49-1.37 (m, 1H); MS (EI) for C₈H₁₄NOCl: 176 (MH⁺).

Example 8

***N*-(4-Bromo-2,3-dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0219] *N*-(4-bromo-2,3-dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: (3*S*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine (115 mg, 0.655 mmol) and 4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetic acid salt (347 mg, 0.655 mmol) were dissolved in dimethylacetamide (0.8 mL) and potassium carbonate (452 mg, 3.28 mmol) was added. The mixture was stirred at 130°C for 36 h. The mixture was cooled to room temperature and the mixture was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate and the combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a brown oil which was purified by column chromatography (ethyl acetate-ethanol 1:1). The purified material was dissolved in ethanol and treated with 4M solution of HCl in 1,4-dioxane (0.25 mL) and the mixture was concentrated *in vacuo*. The residue was dissolved in water and lyophilized to afford *N*-(4-bromo-2,3-dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride as a brown solid (131 mg, 0.222 mmol, 34% yield). ¹H NMR (400 MHz, d₆-DMSO): 11.9-11.5 (m, 2H), 8.79 (s, 1H), 8.34 (s, 1H), 7.96 (d, 1H), 7.54 (d, 1H), 7.41 (s, 1H), 4.47-4.25 (m, 4H), 4.03 (s, 3H) 3.96-3.00 (m, 6H), 2.18-1.88 (m, 3H), 1.73-1.57 (m, 1H); MS (EI) for C₂₃H₂₃N₄O₃Cl₂Br: 553 (MH⁺).

[0220] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0221] *N*-(3,4-Dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 11.5-11.2 (m, 2H), 8.86 (s, 1H), 8.32 (s, 1H), 8.13 (s, 1H), 7.79 (dd, 1H), 7.73 (d, 1H), 7.37 (s, 1H), 4.45-4.24 (m, 4H), 4.03 (s, 3H) 3.93-3.00 (m, 6H), 2.20-1.90 (m, 3H), 1.74-1.56 (m, 1H); MS (EI) for C₂₃H₂₄N₄O₃Cl₂: 475 (MH⁺).

- [0222] *N*-(3,4-Dichlorophenyl)-7-{[(3*R*,8*aR*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 11.6-11.3 (m, 2H), 8.86 (s, 1H), 8.32 (s, 1H), 8.14 (s, 1H), 7.80 (dd, 1H), 7.73 (d, 1H), 7.35 (s, 1H), 4.45-4.25 (m, 4H), 4.03 (s, 3H), 3.96-2.98 (m, 6H), 2.19-1.89 (m, 3H), 1.72-1.57 (m, 1H); MS (EI) for C₂₃H₂₄N₄O₃Cl₂: 475 (MH⁺).
- [0223] *N*-(3,4-Dichlorophenyl)-7-{[(3*R*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 11.6-11.4 (m, 2H), 8.89 (s, 1H), 8.42 (s, 1H), 8.18 (d, 1H), 7.84 (dd, 1H), 7.75 (d, 1H), 7.40 (s, 1H), 4.32 (d, 2H), 4.23-4.15 (m, 1H), 4.09-3.95 (m, 2H), 4.05 (s, 3H), 3.70-3.03 (m, 5H), 2.14-2.04 (m, 4H); MS (EI) for C₂₃H₂₄N₄O₃Cl₂: 475 (MH⁺).
- [0224] *N*-(3,4-Dichlorophenyl)-7-{[(3*S*,8*aR*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 11.7-11.5 (m, 2H), 8.90 (s, 1H), 8.49 (s, 1H), 8.18 (d, 1H), 7.86 (dd, 1H), 7.75 (d, 1H), 7.42 (s, 1H), 4.31 (d, 2H), 4.24-4.16 (m, 1H), 4.09-3.95 (m, 2H), 4.06 (s, 3H), 3.69-3.04 (m, 5H), 2.14-2.03 (m, 4H); MS (EI) for C₂₃H₂₄N₄O₃Cl₂: 475 (MH⁺).

Example 9

N-(3,4-Dichloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride

- [0225] (3*S*,8*aS*)-Hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl acetate: (3*S*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine (2.30 g, 13.1 mmol) and potassium acetate (12.8 g, 131 mmol) were stirred in dimethylformamide (25 mL) at 140°C for 20 h. The mixture was partitioned between ethyl acetate and water. The organic portion was washed twice with water, then with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl acetate as a brown oil (2.53 g, 12.7 mmol, 97% yield). ¹H NMR (400 MHz, CDCl₃): 4.14-4.02 (m, 3H), 3.81-3.72 (m, 1H), 3.37-3.31 (m, 1H), 3.09 (dt, 1H), 3.00 (dd, 1H), 2.21-2.00 (m, 3H), 2.10 (s, 3H), 1.90-1.67 (m, 3H), 1.39-1.24 (m, 1H); MS (EI) for C₁₀H₁₇NO₃: 200 (MH⁺).
- [0226] (3*S*,8*aS*)-Hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethanol: (3*S*,8*aS*)-Hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl acetate (2.36 g, 11.9 mmol) was

treated with sodium methoxide (25 wt% solution in methanol; 2.7 mL) for 0.5 h. The mixture was cooled in an ice bath and a solution of 4M HCl in 1,4-dioxane (3 mL, 12.0 mmol) was added slowly. The mixture was stirred at room temperature for 5 minutes and then was concentrated *in vacuo* to afford a suspension which was diluted with dichloromethane, filtered and the filtrate was concentrated *in vacuo* to afford (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethanol as a brown oil (1.93 g, >100% yield). ¹H NMR (400 MHz, CDCl₃): 4.05 (dd, 1H), 3.73-3.65 (m, 2H), 3.62-3.56 (m, 1H), 3.39-3.34 (m, 1H), 3.10 (dt, 1H), 3.00-2.95 (m, 1H), 2.24-1.98 (m, 4H), 1.97-1.70 (m, 3H), 1.44-1.28 (m, 1H); MS (EI) for C₈H₁₅NO₂: 158 (MH⁺).

[0227] (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl methanesulfonate: (3*S*,8*aS*)-Hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethanol (1.00 g, 6.37 mmol) was dissolved in dichloromethane (10 mL) and triethylamine (2.4 mL, 17.3 mmol) was added at 0°C followed by dropwise addition of methanesulfonyl chloride (0.93 mL, 12.0 mmol). The solution was warmed to room temperature and stirred for 1.25 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The organic portion was washed with saturated sodium bicarbonate solution. The combined aqueous portion was extracted with ethyl acetate. The combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl methanesulfonate as an orange-brown oil (1.20 g, 5.1 mmol, 80% yield). MS (EI) for C₉H₁₇NO₄S: 236 (MH⁺).

[0228] *N*-(3,4-dichloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: 4-[(3,4-Dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetic acid salt (307 mg, 0.655 mmol) was dissolved in dimethylformamide (1 mL) and potassium carbonate (452 mg, 3.28 mmol) was added followed by (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl methanesulfonate (250 mg, 1.06 mmol). The mixture was stirred at 70°C for 41 h and then was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford an orange oil which was purified by column chromatography (ethyl acetate-ethanol 1:1). The purified material was dissolved in methanol and treated with 4M solution of HCl in 1,4-dioxane (0.1 mL) and the mixture was concentrated *in vacuo* to afford *N*-(3,4-

dichloro-2-fluorophenyl)-7- $\{[(3S,8aS)$ -hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy $\}$ -6-(methyloxy)quinazolin-4-amine hydrochloride as a pale yellow solid (66 mg, 0.125 mmol, 19% yield). ^1H NMR (400 MHz, d_6 -DMSO): 11.9-11.5 (m, 2H), 8.83 (s, 1H), 8.39-8.35 (m, 1H), 7.69 (dd, 1H), 7.62 (dd, 1H), 7.43 (s, 1H), 4.48-4.24 (m, 4H), 4.04 (s, 3H), 3.97-3.85 (m, 1H), 3.78-2.96 (m, 5H), 2.17-1.90 (m, 3H), 1.72-1.58 (m, 1H); MS (EI) for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_3\text{FCl}_2$: 493 (MH^+).

[0229] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0230] *N*-(3-Chloro-2,4-difluorophenyl)-7- $\{[(3S,8aS)$ -hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy $\}$ -6-(methyloxy)quinazolin-4-amine hydrochloride: ^1H NMR (400 MHz, d_6 -DMSO): 11.9-11.6 (m, 2H), 8.84 (s, 1H), 8.41-8.37 (m, 1H), 7.67-7.60 (m, 1H), 7.51 (dt, 1H), 7.41 (s, 1H), 4.48-4.24 (m, 4H), 4.04 (s, 3H), 3.97-3.86 (m, 1H), 3.80-2.96 (m, 5H), 2.18-1.90 (m, 3H), 1.72-1.59 (m, 1H); MS (EI) for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_3\text{F}_2\text{Cl}$: 477 (MH^+).

[0231] *N*-(4-Bromo-3-chloro-2-fluorophenyl)-7- $\{[(3S,8aS)$ -hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy $\}$ -6-(methyloxy)quinazolin-4-amine: ^1H NMR (400 MHz, CDCl_3): 8.70 (s, 1H), 8.49 (dd, 1H), 7.49 (dd, 1H), 7.29 (s, 1H), 6.98 (s, 1H), 4.25 (dd, 1H), 4.17 (dd, 1H), 4.13-4.06 (m, 2H), 4.03 (s, 3H), 3.46-3.38 (m, 1H), 3.20 (dd, 1H), 3.14 (dt, 1H), 2.28-2.17 (m, 2H), 2.17-2.07 (m, 1H), 1.90-1.71 (m, 3H), 1.42-1.30 (m, 1H); MS (EI) for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_3\text{FClBr}$: 537 (MH^+).

[0232] *N*-(4,5-Dichloro-2-fluorophenyl)-7- $\{[(3S,8aS)$ -hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy $\}$ -6-(methyloxy)quinazolin-4-amine: ^1H NMR (400 MHz, CDCl_3): 8.91 (d, 1H), 8.71 (s, 1H), 7.29-7.26 (m, 2H), 6.94 (s, 1H), 4.24 (dd, 1H), 4.16 (dd, 1H), 4.11-4.04 (m, 2H), 4.02 (s, 3H), 3.44-3.38 (m, 1H), 3.19 (dd, 1H), 3.13 (dt, 1H), 2.28-2.17 (m, 2H), 2.16-2.07 (m, 1H), 1.91-1.69 (m, 3H), 1.42-1.30 (m, 1H); MS (EI) for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_3\text{FCl}_2$: 493 (MH^+).

[0233] *N*-(4-Bromo-5-chloro-2-fluorophenyl)-7- $\{[(3S,8aS)$ -hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy $\}$ -6-(methyloxy)quinazolin-4-amine: ^1H NMR (400 MHz, CDCl_3): 8.95 (d, 1H), 8.72 (s, 1H), 7.43 (d, 1H), 7.28 (s, 1H), 6.93 (s, 1H), 4.24 (dd, 1H), 4.15 (dd, 1H), 4.12-4.04 (m, 2H), 4.02 (s, 3H), 3.44-3.37 (m, 1H), 3.19 (dd, 1H), 3.13 (dt, 1H), 2.27-2.16 (m, 2H), 2.16-2.06 (m, 1H), 1.90-1.69 (m, 3H), 1.42-1.28 (m, 1H); MS (EI) for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_3\text{FClBr}$: 537 (MH^+).

Example 10

***N*-(3,4-Dichloro-2-fluorophenyl)-6-(methyloxy)-7-[(octahydro-2*H*-quinolizin-3-ylmethyl)oxy]quinazolin-4-amine**

- [0234] Octahydro-2*H*-quinolizin-3-ylmethanol: Ethyl octahydro-2*H*-quinolizine-3-carboxylate (2.35 g, 11.1 mmol) was added dropwise to a stirred suspension of lithium aluminum hydride (1 M solution in tetrahydrofuran, 33 mL, 33 mmol) in tetrahydrofuran (50 mL) at 0°C. The reaction was stirred at room temperature for 3 h. The mixture was cooled in an ice bath and ethyl acetate (6 mL) was added slowly, followed by water (1.25 mL), 15% aqueous sodium hydroxide solution (5 mL) and water (1.25 mL). The mixture was filtered through a pad of celite and washed with ether. The filtrate was concentrated *in vacuo* and dried rigorously to afford octahydro-2*H*-quinolizin-3-ylmethanol as a yellow oil (1.66 g, 9.82 mmol, 88% yield). MS (EI) for C₁₀H₁₉NO: 170 (MH⁺).
- [0235] Octahydro-2*H*-quinolizin-3-ylmethyl methanesulfonate: Octahydro-2*H*-quinolizin-3-ylmethanol (600 mg, 3.55 mmol) was dissolved in dichloromethane (8 mL) and triethylamine (1.5 mL, 10.8 mmol) was added at 0°C followed by dropwise addition of methanesulfonyl chloride (0.56 mL, 7.16 mmol). The solution was warmed to room temperature and stirred for 1.25 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The aqueous portion was extracted with ethyl acetate. The combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford octahydro-2*H*-quinolizin-3-ylmethyl methanesulfonate as an orange oil (796 mg, 3.22 mmol, 91% yield). MS (EI) for C₁₁H₂₁NO₃S: 248 (MH⁺).
- [0236] *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[(octahydro-2*H*-quinolizin-3-ylmethyl)oxy]quinazolin-4-amine: 4-[(3,4-Dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol hydrochloride (469 mg, 1.20 mmol) was dissolved in dimethylformamide (1 mL) and potassium carbonate (828 mg, 6.00 mmol) was added followed by octahydro-2*H*-quinolizin-3-ylmethyl methanesulfonate (466 mg, 1.89 mmol) in dimethylformamide (1 mL). The mixture was stirred at 70°C for 38 h and then was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a brown oil which was purified by column chromatography (15-20% methanol in dichloromethane). The purified material was crystallized from methanol to afford *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[(octahydro-2*H*-quinolizin-3-ylmethyl)oxy]quinazolin-4-amine as a cream colored solid

(83.4 mg, 0.165 mmol, 14% yield). ^1H NMR (400 MHz, CDCl_3): 8.69 (s, 1H), 8.53 (t, 1H), 7.34 (dd, 1H), 7.28-7.22 (m, 1H), 7.23 (s, 1H), 6.98 (s, 1H), 4.06-3.95 (m, 2H), 4.02 (s, 3H), 3.09 (d, 1H), 2.87 (d, 1H), 2.43-2.27 (m, 1H), 2.10-1.97 (m, 1H), 1.95-1.84 (m, 2H), 1.80-1.52 (m, 5H), 1.46-0.95 (m, 5H); MS (EI) for $\text{C}_{25}\text{H}_{27}\text{N}_4\text{O}_2\text{FCl}_2$: 505 (MH^+).

Example 11

(3*S*,8*aS*)-3-({[4-[(3,4-Dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one

[0237] (3*S*,8*aS*)-3-(Hydroxymethyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: A solution of methyl 1-[(2*S*)-3-hydroxy-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (3.50 g, 10.4 mmol) in methanol was added to 5% palladium on carbon (50 wt.% in water) in methanol and treated with hydrogen at 40 psi for 1 h. The mixture was filtered and the filtrate was brought to reflux briefly and then cooled and concentrated *in vacuo* to afford (3*S*,8*aS*)-3-(hydroxymethyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a colorless solid (1.50 g, 8.83 mmol, 85% yield). ^1H NMR (400 MHz, CDCl_3): 7.28-7.22 (m, 1H), 3.83-3.75 (m, 1H), 3.69 (dd, 1H), 3.56 (dd, 1H), 3.31 (t, 1H), 3.08 (dd, 1H), 2.92 (dt, 1H), 2.76-2.70 (m, 1H), 2.66 (dd, 1H), 2.28-2.16 (m, 1H), 2.02-1.73 (m, 3H); MS (EI) for $\text{C}_8\text{H}_{14}\text{N}_2\text{O}_2$: 171 (MH^+).

[0238] (3*S*,8*aS*)-3-({[(1,1-Dimethylethyl)(dimethyl)silyl]oxy}methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: To a solution of (3*S*,8*aS*)-3-(hydroxymethyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one (1.49 g, 8.82 mmol) in dimethylformamide (20 mL) was added triethylamine (2.45 mL, 17.6 mmol) and 4-dimethylaminopyridine (90 mg, 0.882 mmol). The solution was cooled in an ice bath and *tert*-butyldimethylsilyl chloride (2.66 g, 17.6 mmol) was added. The mixture was warmed to room temperature and stirred for 14 h. The mixture was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The aqueous portion was extracted twice with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a pale brown solid which was triturated with ethyl acetate to afford (3*S*,8*aS*)-3-({[(1,1-dimethylethyl)(dimethyl)silyl]oxy}methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as an off-white solid (1.74 g, 5.84 mmol, 66% yield). ^1H NMR (400 MHz, CDCl_3): 6.09-5.90 (m, 1H), 3.86-3.76 (m, 1H), 3.63 (dd, 1H), 3.44 (dd, 1H), 3.25 (t, 1H), 3.10 (ddd, 1H), 2.98-2.90 (m, 1H), 2.68-2.60 (m, 1H), 2.52 (dd, 1H), 2.28-2.18 (m, 1H), 2.06-1.95 (m, 1H), 1.93-1.74 (m, 2H), 0.90 (s, 9H), 0.07 (s, 6H); MS (EI) for $\text{C}_{14}\text{H}_{28}\text{N}_2\text{O}_2\text{Si}$: 285 (MH^+).

[0239] (3*S*,8*aS*)-3-({[(1,1-Dimethylethyl)(dimethyl)silyl]oxy}methyl)-2-methylhexahydro pyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: (3*S*,8*aS*)-3-({[(1,1-Dimethylethyl)(dimethyl)silyl]oxy} methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one (1.51g, 5.32mmol) in dimethylformamide (8 mL) was added to an ice-cooled suspension of sodium hydride (60 wt.% dispersion in oil; 213 mg, 5.32 mmol) in dimethylformamide (8 mL). The mixture was stirred at 0°C for 0.25 h and then iodomethane (0.332 mL, 5.32 mmol) was added dropwise. The mixture was stirred at room temperature for 0.5 h and then was stirred at 70°C for 2 h. The mixture was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3*S*,8*aS*)-3-({[(1,1-dimethylethyl)(dimethyl)silyl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a yellow oil (1.552 g, 5.21 mmol) which was dissolved in tetrahydrofuran (20 mL) and treated with tetrabutylammonium fluoride (1.0M solution in tetrahydrofuran; 10.4 mL, 10.4 mmol) for 2 h at room temperature. The mixture was concentrated *in vacuo* and purified by column chromatography (10% methanol in dichloromethane) to afford (3*S*,8*aS*)-3-(hydroxymethyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a yellow oil (496mg, 2.70mmol, 51% yield from (3*S*,8*aS*)-3-({[(1,1-dimethylethyl)(dimethyl)silyl]oxy} methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one). ¹H NMR (400 MHz, CDCl₃): 3.98-3.93 (m, 1H), 3.86 (dd, 1H), 3.61-3.55 (m, 1H), 3.29-3.25 (m, 1H), 3.09-3.03 (m, 1H), 3.03-2.97 (m, 1H), 3.02 (s, 3H), 2.93 (dd, 1H), 2.87-2.79 (m, 1H), 2.32-2.21 (m, 1H), 2.00-1.86 (m, 2H), 1.83-1.64 (m, 1H); MS (EI) for C₉H₁₆N₂O₂: 185 (MH⁺).

[0240] (3*S*,8*aS*)-3-({[4-[(3,4-Dichloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: (3*S*,8*aS*)-3-(Hydroxymethyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one (505 mg, 2.74 mmol) was dissolved in dichloromethane (10 mL) and triethylamine (0.8 mL, 5.75 mmol) was added at 0°C followed by dropwise addition of methanesulfonyl chloride (0.45 mL, 5.81 mmol). The solution was warmed to room temperature and stirred for 1.25 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The aqueous portion was extracted with ethyl acetate. The combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford [(3*S*,8*aS*)-2-methyl-1-

oxooctahydropyrrolo[1,2-*a*]pyrazin-3-yl)methyl methanesulfonate as an orange oil (538 mg, 2.05 mmol, 75% yield). 4-[(3,4-Dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol hydrochloride (469 mg, 1.20 mmol) was dissolved in dimethylformamide (1 mL) and potassium carbonate (828 mg, 6.00 mmol) was added followed by [(3*S*,8*aS*)-2-methyl-1-oxooctahydropyrrolo[1,2-*a*]pyrazin-3-yl)methyl methanesulfonate (538 mg, 2.05 mmol) in dimethylformamide (1 mL). The mixture was stirred at 70°C for 34 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a brown oil which was purified by column chromatography (6-8% methanol in dichloromethane) to afford a yellow foam (300 mg, 0.577 mmol, 48% yield). The yellow foam (100 mg) was purified further by column chromatography (ethyl acetate-ethanol 1:1) to afford (3*S*,8*aS*)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a yellow solid (60 mg). ¹H NMR (400 MHz, CDCl₃): 8.71 (s, 1H), 8.52 (dd, 1H), 7.36-7.32 (m, 2H), 7.01 (s, 1H), 4.49 (dd, 1H), 4.34 (dd, 1H), 4.03 (s, 3H), 3.90-3.84 (m, 1H), 3.47 (t, 1H), 3.13 (s, 3H), 3.05 (dd, 1H), 2.95 (dd, 1H), 2.93-2.83 (m, 2H), 2.29-2.19 (m, 1H), 2.03-1.84 (m, 2H), 1.83-1.70 (m, 1H); MS (EI) for C₂₄H₂₄N₅O₃FCl₂: 520 (MH⁺).

Example 12

(3*S*,8*aS*)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one

[0241] 1,2-Dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidiny]-2-[[[(phenylmethoxy)carbonyl]amino]-D-*glycero*-hexitol: To a solution of 2-deoxy-2-[[[(phenylmethyloxy)carbonyl]amino]-D-*glycero*-hexopyranose (5.0 g, 0.016 mol) in methanol (500 mL) was added L-proline methyl ester hydrochloride (2.8 g, 0.022 mol) and sodium cyanoborohydride (3.4 g, 0.054 mol). The solution was heated to 64 °C for 14 h. After cooling to room temperature, the reaction mixture was concentrated *in vacuo* to afford 1,2-dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidiny]-2-[[[(phenylmethoxy)carbonyl]amino]-D-*glycero*-hexitol (6.81 g, 100%) as a clear and colorless oil. MS (EI) for C₂₀H₃₁N₂O₈: 427 (MH⁺).

[0242] Methyl 1-[(2*S*)-3-hydroxy-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate: 1,2-dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidinyl]-2-[[[(phenylmethoxy)carbonyl]amino]-D-*glycero*-hexitol (6.81 g, 0.016 mol) was taken into water (100 mL) and the resulting solution was cooled to 0°C. Sodium periodate (14.8 g, 0.069 mol) dissolved in water was added dropwise and the resulting mixture was stirred at 0°C for 2 h. The reaction mixture was partitioned with dichloromethane (3x100 mL), dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The residue was taken up in methanol (200 mL) and the resulting solution was cooled to 0°C. Sodium borohydride (1.98 g, 0.052 mol) was added and the reaction mixture was stirred for 1 h at 0°C. The reaction mixture was concentrated *in vacuo* and partitioned with dichloromethane and saturated aqueous ammonium chloride. The organic layer was dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The resulting crude product was purified by column chromatography (5% methanol in dichloromethane) to yield methyl 1-[(2*S*)-3-hydroxy-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (4.9 g, 92%) as a white solid. MS (EI) for C₁₇H₂₅N₂O₅: 337 (MH⁺).

[0243] Methyl 1-[(2*S*)-3-[(methylsulfonyl)oxy]-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate: Methyl 1-[(2*S*)-3-hydroxy-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (200 mg, 0.594 mmol) was dissolved in dichloromethane (5 mL) followed by the addition of 4-(dimethylamino)pyridine (3.6 mg, 0.039 mmol) and triethylamine (0.125 mL, 0.891 mmol) and the resulting mixture was cooled to 0 °C. Methanesulfonyl chloride (0.060 mL, 0.773 mmol) was added dropwise and the reaction mixture was stirred for 1 h at 0°C. The mixture was partitioned between dichloromethane and saturated aqueous sodium bicarbonate. The organic layer was dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to afford methyl 1-[(2*S*)-3-[(methylsulfonyl)oxy]-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (246 mg, 100%) as a clear and colorless oil. MS (EI) for C₁₈H₂₇N₂O₇S: 415 (MH⁺).

[0244] methyl (1*R*)-2-[(2*S*)-3-{[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]cyclopentane carboxylate: 4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol hydrochloride (400 mg, 1.02 mmol) and methyl 1-[(2*S*)-3-[(methylsulfonyl)oxy]-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (603 mg, 1.45 mmol) were suspended in DMF (5 mL) and powdered potassium carbonate (705 mg, 5.10 mmol) was

added. The mixture was stirred at 70°C for 12 h. The reaction mixture was filtered and concentrated *in vacuo*. The crude residue was purified by column chromatography (10% methanol in dichloromethane) to yield methyl (1*R*)-2-[(2*S*)-3-{[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]cyclopentanecarboxylate (686 mg, 100%) as a yellow oil. MS (EI) for C₃₂H₃₂Cl₂FN₅O₆: 672 (M⁺).

[0245] (3*S*,8*aS*)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: Methyl (1*R*)-2-[(2*S*)-3-{[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]cyclopentanecarboxylate (686mg, 1.02 mmol) was diluted with glacial acetic acid (3 mL) and 30wt% hydrogen bromide in acetic acid (2 mL) was added. The resulting mixture was stirred for 13 h and then concentrated *in vacuo*. The crude residue was taken up in methanol (5 mL) followed the addition of powdered potassium carbonate (700 mg, 5.07 mmol) at room temperature. The resulting mixture was stirred for 7 h, filtered, and concentrated *in vacuo*. The crude residue was purified by column chromatography (10% methanol in dichloromethane) to yield the title compound (181 mg, 35%) as a yellow solid. ¹H NMR (400 MHz, d₄-MeOH): 8.34 (s, 1H), 7.75 (s, 1H), 7.76-7.55 (m, 1H), 7.45-7.40 (dd, 1H), 7.17 (s, 1H), 4.30-4.25 (m, 1H), 4.20-4.15 (m, 2H), 4.03 (s, 3H), 3.78-3.70 (m, 1H), 3.65-3.60 (m, 1H), 3.55-3.50 (m, 1H), 3.32-3.29 (m, 2H), 3.00-2.95 (m, 2H), 2.83-2.78 (m, 2H), 2.25-2.15 (m, 2H); MS (EI) for C₂₃H₂₃Cl₂FN₅O₃: 507 (MH⁺).

[0246] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0247] (3*S*,8*aR*)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: C₂₃H₂₃Cl₂FN₅O₃: 507 (MH⁺).

[0248] (3*S*,8*aS*)-3-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}-methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: ¹H NMR (400 MHz, d₄-MeOH): 8.36 (s, 1H), 7.71 (s, 1H), 7.60-7.55 (m, 2H), 7.18 (s, 1H), 4.29-4.22 (m, 1H), 4.19-4.14 (m, 1H), 4.02 (s, 3H), 3.99-3.92 (m, 1H), 3.36-3.30 (m, 1H), 3.32-3.90 (m, 2H), 2.82-2.74 (m, 1H), 2.26-2.10 (m, 1H), 2.19-2.18 (m, 3H), 1.30-1.20 (m, 2H), 0.90-0.80 (m, 1H); MS (EI) for C₂₃H₂₃BrClN₅O₃: 551 (MH⁺).

- [0249] (3*S*, 9*aS*)-3-({[4-[(3, 4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2*H*-pyrido[1,2-*a*]pyrazin-1(6*H*)-one: C₂₄H₂₅Cl₂FN₅O₃: 521 (MH⁺).
- [0250] (3*S*, 9*aR*)-3-({[4-[(3, 4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2*H*-pyrido[1,2-*a*]pyrazin-1(6*H*)-one: C₂₄H₂₅Cl₂FN₅O₃: 521 (MH⁺).

Example 13

***N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

- [0251] 1,1-Dimethylethyl (3*aR*,6*aS*)-5-(hydroxymethyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: Under a nitrogen atmosphere, borane tetrahydrofuran complex (1*M* in THF, 42 mL, 41.9 mmol) was diluted with tetrahydrofuran (42 mL) and cooled with an ice bath. Neat 2,3-dimethylbut-2-ene (5.0 mL, 41.9 mmol) was added in portions over 0.25 h and the solution was stirred at 0°C for 3 h. A solution of 1,1-dimethylethyl (3*aR*,6*aS*)-5-methylidenehexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (1.98 g, 8.88 mmol) in tetrahydrofuran (10 mL) was added slowly, and the solution was warmed to room temperature and stirred 12 h. After cooling to 0°C, 10% aqueous sodium hydroxide (17 mL, 41.7 mmol) was added slowly, followed by 30% aqueous hydrogen peroxide (13 mL, 128 mmol) and the solution was warmed to room temperature. The solvent was removed *in vacuo* and the solution was partitioned between water and diethyl ether. The layers were separated and the aqueous layer was further extracted (3 x 50 mL diethyl ether). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide 2.04 (95%) of 1,1-dimethylethyl (3*aR*,6*aS*)-5-(hydroxymethyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate, which was used without purification. ¹H NMR (400 MHz, CDCl₃): 8.50 (broad s, 1H), 3.66-3.46 (m, 3H), 3.20-3.00 (m, 2H), 2.70-2.59 (m, 2H), 2.37-2.18 (m, 1H), 2.04 (m, 1H), 1.84 (broad s, 1H), 1.70-1.55 (m, 1H), 1.46 (s, 9H), 1.17 (m, 1H), 0.93 (m, 1H).
- [0252] 1,1-Dimethylethyl (3*aR*,6*aS*)-5-{{[(methanesulfonyl)oxy]methyl}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate Methanesulfonyl chloride (0.2mL, 2.48mmol), was added dropwise to a solution of 1,1-dimethylethyl (3*aR*,6*aS*)-5-(hydroxymethyl)hexahydro cyclopenta[*c*] pyrrole-2(1*H*)-carboxylate (0.40 g, 1.65 mmol) and triethylamine (0.69 mL, 4.95 mmol) in 20 mL dichloromethane at 0°C and the reaction mixture was stirred for 1 h at room temperature. The solvent was evaporated, the

resulting crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), 1M aqueous sodium hydroxide, brine, 1M aqueous hydrochloric acid and brine again. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The resulting 1,1-dimethylethyl (3a*R*,6a*S*)-5-[[[(methylsulfonyl)oxy]methyl]hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate was used without further purification. MS (EI) for C₁₄H₂₅NO₅S: 320 (MH⁺), 264 (M-tBu).

[0253] 1,1-Dimethylethyl (3a*R*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: A solution of 4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetate (salt) (0.217g, 0.425mmol), 1,1-dimethylethyl (3a*R*,6a*S*)-5-[[[(methylsulfonyl)oxy]methyl]hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (0.163g, 0.510 mmol), potassium carbonate (0.290 g, 2.12 mmol) in *N,N*-dimethylacetamide (1.6 mL) was heated in a sealed reaction tube at 90°C for 12 h. The crude reaction mixture was diluted with 100 mL of 10% methanol in ethyl acetate and washed with water (5 x 30 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 3:1 hexanes:acetone) provided 1,1-dimethylethyl (3a*R*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate which was used directly in the next step. MS (EI) for C₂₈H₃₁N₄O₄FCIBr: 623 (MH⁺).

[0254] *N*-(4-Bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-[[[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-ylmethyl]oxy]quinazolin-4-amine hydrochloride: 1,1-Dimethylethyl (3a*R*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate derivative was taken up in methanol (50 mL) and treated with 4.0M hydrogen chloride in dioxane (excess) and heated briefly to reflux. Concentration *in vacuo* gave *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-[[[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-ylmethyl]oxy]quinazolin-4-amine hydrochloride which was used directly in the next step. ¹H NMR (400 MHz, d₄-MeOH): 8.70 (s, 1H), 7.97 (s, 1H), 7.68 (d, 1H), 7.49 (t, 1H), 7.28 (s, 1H), 4.25 (m, 2H), 4.08 (s, 3H), 3.57 (m, 1H), 3.02 (m, 4H), 2.80-2.60 (m, 2H), 2.35 (m, 1H), 1.89 (m, 4H), 1.40 (m, 1). MS (EI) for C₂₃H₂₃N₄O₂FCIBr: 522 (MH⁺).

[0255] *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-[[[(3a*R*,5*r*,6a*S*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: *N*-(4-Bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-[[[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta-

[*c*]pyrrol-5-ylmethyl}oxy}quinazolin-4-amine hydrochloride was solubilized in formic acid (3.0 mL) and 37% aqueous formaldehyde (0.5 mL, 2.55 mmol) was added. The solution was heated to 95°C for 12 h and additional formaldehyde (1.0 mL, 5.10 mmol) was added. After heating an additional 12 h, the reaction mixture was concentrated *in vacuo*. The residue was taken up in methanol and treated with Bio-Rad AG 1-X8 resin hydroxide form until pH 8. The product was filtered, concentrated *in vacuo*, and purified by HPLC (reverse-phase, water/acetonitrile/0.1% TFA). Upon removal of solvent, the product was taken up in methanol and treated with Bio-Rad AG 1-X8 resin hydroxide form until pH 8. The product was filtered and concentrated *in vacuo* then taken up in fresh methanol and treated with 4.0 M hydrogen chloride in dioxane (0.05 mL). Removal of solvent *in vacuo* provided 54.1 mg (24%) of *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydro-cyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride. ¹H NMR (400 MHz, d₆-DMSO): 8.83 (d, 1H), 8.33 (s, 1H), 7.80 (d, 1H), 7.56 (t, 1H), 7.40 (s, 1H), 4.16 (m, 2), 4.01 (s, 3H), 3.80-3.68 (m, 1H), 3.05 (m, 2H), 2.90-2.70 (m, 5H), 2.34 (m, 1H), 2.15 (m, 2H), 1.75 (m, 1H), 1.57 (m, 2H), 1.35 (m, 1H). MS (EI) for C₂₄H₂₅N₄O₂FClBr: 537 (MH⁺).

[0256] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0257] *N*-(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.83 (d, 1H), 8.33 (s, 1H), 8.03 (d, 1H), 7.93 (d, 1H), 7.41 (m, 1H), 4.16 (m, 2), 4.02 (s, 3H), 3.70 (m, 1H), 3.05 (m, 2H), 2.91-2.75 (m, 5H), 2.34 (m, 1H), 2.16 (m, 2H), 1.75 (m, 1H), 1.57 (m, 2H), 1.35 (m, 1H). MS (EI) for C₂₄H₂₅N₄O₂FClBr: 537 (MH⁺).

[0258] *N*-(3-chloro-2,4-difluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.81 (d, 1H), 8.39 (d, 1H), 7.61 (m, 1H), 7.47 (m, 1H), 7.42 (s, 1H), 4.17 (m, 2), 4.02 (s, 3H), 3.67 (m, 1H), 3.05 (m, 2H), 2.91-2.75 (m, 5H), 2.34 (m, 1H), 2.16 (m, 2H), 1.75 (m, 1H), 1.57 (m, 2H), 1.35 (m, 1H). MS (EI) for C₂₄H₂₅N₄O₂F₂Cl: 475 (M⁺).

Example 14

***N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride:**

[0259] 1,1-Dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: A solution of 4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl trifluoroacetate (salt) (0.22 g, 0.47 mmol), 1,1-dimethylethyl (3a*R*,6a*S*)-5-{{[(methysulfonyl)oxy]methyl}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (0.16 g, 0.51 mmol), K₂CO₃ (0.33 g, 2.36 mmol) in N,N-dimethylacetamide (5 mL) was heated in a sealed reaction tube at 90°C for 12 h. The crude reaction mixture was diluted with 100 mL 10% methanol in ethyl acetate and washed with saturated aqueous sodium bicarbonate (1x 30 mL), water (1 x 30 mL) and brine (1 x 30 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 3:2 hexanes:acetone) provided 1,1-dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta-*c*]pyrrole-2(1*H*)-carboxylate which was used directly in the next step. MS (EI) for C₂₈H₃₁Cl₂FN₄O₄: 577, 579 (MH⁺).

[0260] *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]methyl]oxy}quinazolin-4-amine hydrochloride: 1,1-Dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate was taken up in methanol (10 mL) and treated with 4.0M hydrogen chloride in dioxane (excess) and heated briefly to reflux. Concentration *in vacuo* provided *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]methyl]oxy}quinazolin-4-amine hydro-chloride. MS (EI) for C₂₃H₂₃Cl₂FN₄O₂: 477, 479 (MH⁺).

[0261] *N*-(3,4-Dichloro-2-fluorophenyl)-7-({[(3a*R*,5*r*,6a*S*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: *N*-(3,4-Dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]methyl]oxy}quinazolin-4-amine hydrochloride was solubilized in formic acid (5.0 mL) and 37% aqueous formaldehyde (1 mL) was added. The solution was heated to 95°C for 12 h. The reaction mixture was concentrated *in vacuo*. The residue was taken up in a mixture of 10% methanol in ethyl acetate (100 mL) and washed with saturated aqueous sodium bicarbonate (2x 30 mL) and brine. The organic layer was dried over anhydrous sodium sulfate, filtered then concentrated and purified by HPLC (reverse-phase, water/acetonitrile/0.1% TFA). Upon removal of solvent the product was taken up in 10% methanol in ethyl acetate (100 mL) and washed

with saturated aqueous sodium bicarbonate (2x 30 mL) and brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated then taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane (1eq.). Removal of solvent *in vacuo* provided 78.3 mg (25%) of *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride. ¹H NMR (400 MHz, d₆-DMSO): 11.00 (bs, 1H), 8.36 (d, 1H), 8.10 (s, 1H), 7.58 (s, 2H), 7.20 (d, 1H), 4.16 (m, 2H), 4.00 (s, 3H), 3.35 (bs, 3H), 2.50 (m, 2H), 2.21 (m, 3H), 2.03 (m, 2H), 1.60 (m, 2H), 1.12 (m, 2H). MS (EI) for C₂₄H₂₅Cl₂FN₄O₂: 491, 493 (MH⁺).

[0262] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0263] *N*-(4,5-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 10.9 (bs, 1H), 8.46 (d, 1H), 8.10 (s, 1H), 8.04 (s, 1H), 7.93 (s, 1H), 7.54 (s, 1H), 4.18 (m, 2H), 4.01 (s, 3H), 3.33 (bs, 3H), 2.46 (m, 2H), 2.23 (s, 3H), 2.04 (m, 2H), 1.58 (m, 2H), 1.14 (m, 2H). MS (EI) for C₂₄H₂₅Cl₂FN₄O₂: 491 (MH⁺).

[0264] *N*-(4-bromo-2,3-dichlorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 11.0 (bs, 1H), 8.60 (s, 1H), 8.14 (s, 1H), 7.76 (d, 1H), 7.44 (t, 1H), 7.24 (s, 1H), 4.16 (m, 2H), 4.00 (s, 3H), 3.35 (bs, 3H), 2.50 (m, 2H), 2.18 (m, 3H), 2.03 (m, 2H), 1.60 (m, 2H), 1.12 (m, 2H). MS (EI) for C₂₄H₂₅BrCl₂N₄O₂: 550, 552 (MH⁺).

[0265] *N*-(3,4-dichlorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 10.98 (bs, 1H), 8.88 (s, 1H), 8.41 (s, 1H), 8.20 (d, 1H), 7.86 (d, 1H), 7.75 (s, 1H), 7.46 (s, 1H), 4.08 (m, 2H), 3.98 (s, 3H), 3.28 (m, 2H), 2.54 (m, 2H), 2.20 (s,m, 4H), 2.18 (m, 2H), 1.62 (m, 2H), 1.24 (m, 2H). MS (EI) for C₂₄H₂₆Cl₂N₄O₂: 473 (MH⁺).

Example 15

***N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-(1-methylethyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0266] *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-(1-methylethyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine

hydrochloride: A solution of *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-
 {[(3*aR*,5*r*,6*aS*)-octahydrocyclopenta[*c*]pyrrol-5-ylmethyl]oxy} quinazolin-4-amine
 hydrobromide (0.1 g, 0.166 mmol), acetone (0.024 mL, 0.332 mmol), and glacial acetic
 acid (5 drops) in acetonitrile:water (3:1) was cooled to 0°C and sodium
 triacetoxyborohydride (53.0 mg, 0.249 mmol) was added. The solution was warmed to
 room temperature and stirred 12 h. Additional acetic acid (5 drops), acetone (0.30 mL,
 6.54 mmol), sodium triacetoxyborohydride (0.300 g, 1.42 mmol) was added in portions
 over 12 h. The acetonitrile was removed *in vacuo* and the aqueous layer was diluted with
 saturated aqueous sodium bicarbonate and 10% methanol in ethyl acetate then the layers
 were separated. The aqueous layer was extracted with 10% methanol in ethyl acetate (2 x
 75 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered
 and concentrated. Column chromatography (SiO₂, gradient of 30-50% methanol in
 chloroform) followed by concentration and treatment in methanol with 4.0 M hydrogen
 chloride in dioxane (0.05 mL) and concentration provided *N*-(4-bromo-3-chloro-2-
 fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-(1-methylethyl)octahydrocyclopenta[*c*]pyrrol-5-
 yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride (75.7 mg, 76% yield).
¹H NMR (400 MHz, d₄-MeOH): 8.69 (s, 1H), 7.97 (s, 1H), 7.67 (d, 1H), 7.50 (t, 1H),
 7.28 (s, 1H), 4.27 (d, 2H), 4.07 (s, 3H), 3.84-3.20 (m, 4H), 3.01 (m, 3H), 2.80 (m, 1H)
 2.34 (m, 3H), 1.52 (m, 2H), 1.42 (dd, 6H); MS (EI) for C₂₆H₂₉N₄O₂FClBr: 565 (MH⁺).

[0267] Using the same synthetic techniques and/or substituting with alternative reagents,
 the following compound of the invention was also prepared.

[0268] *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-(1-
 methylethyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-
 4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.68 (broad s, 1H), 7.98 (broad
 s, 1H), 7.54 (m, 3H), 7.32 (broad s, 1H), 4.27 (d, 2H), 4.07 (s, 3H), 3.84-3.20 (m, 4H),
 3.01 (m, 3H), 2.80 (m, 1H) 2.34 (m, 3H), 1.93-1.75 (m, 2H), 1.42 (dd, 6H); MS (EI) for
 C₂₆H₂₉N₄O₂FCl₂: 519 (MH⁺).

Example 16

***N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-ethyloctahydrocyclo-
 penta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0269] *N*-(4-Bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-ethyloctahydrocyclo-
 penta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: A
 solution of *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3*aR*,5*r*,6*aS*)-

octahydrocyclopenta[*c*]pyrrol-5-ylmethyl}oxy}quinazolin-4-amine hydrobromide (0.1 g, 0.166 mmol) and acetaldehyde (0.010 mL, 0.249 mmol) in 50% methanol in tetrahydrofuran was cooled to 0°C and sodium cyanoborohydride (1 M in THF, 0.10 mL, 0.200 mmol) was added. The solution was warmed to room temperature and stirred for 1.5 h. The solvents were removed and the residue was partitioned between water and 10% methanol in ethyl acetate. The layers were separated and the aqueous layer was extracted with 10% methanol in ethyl acetate (3 x 50 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography (SiO₂, gradient of 5-10% methanol in chloroform), followed by treatment in methanol with 4.0 M hydrogen chloride in dioxane (0.05 mL) and concentration provided *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-ethyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy) quinazolin-4-amine hydrochloride (37.5 mg, 36% yield). ¹H NMR (400 MHz, d₆-DMSO): 8.75 (d, 1H), 8.21 (broad s, 1H), 7.75 (d, 3H), 7.54 (t, 1H), 7.34 (m, 1H), 7.12 (d, 1H), 4.16 (d, 2H), 4.00 (s, 3H), 3.75 (m, 1H), 3.11-2.65 (m, 3H), 2.40 (m, 1H), 2.15 (m, 2H), 1.61 (m, 2H), 1.26 (m, 5); MS (EI) for C₂₅H₂₇N₄O₂FClBr: 551 (MH⁺).

[0270] Using the same synthetic techniques and/or substituting with alternative reagents, the following compound of the invention were prepared:

[0271] *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3*aR*,5*r*,6*aS*)-2-(2-methylpropyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.73 (d, 1H), 8.32 (broad s, 1H), 7.76 (d, 1H), 7.54 (t, 1H), 7.41 (m, 1H), 7.22 (d, 1H), 4.18 (d, 2H), 4.01 (s, 3H), 3.74 (m, 1H), 3.11 (m, 1H), 2.94 (m, 5H), 2.65 (m, 1H), 2.40 (m, 1H), 2.13 (m, 2H), 2.00 (m, 1H), 1.69 (m, 1H), 1.36 (m, 1H), 0.98 (t, 6H); MS (EI) for C₂₇H₃₁N₄O₂FClBr: 579 (MH⁺).

[0272] *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-ethyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.77 (d, 1H), 8.36 (broad s, 1H), 7.63 (m, 2H), 7.42 (m, 1H), 7.20 (d, 1H), 4.17 (d, 2H), 4.02 (s, 3H), 3.74 (m, 1H), 3.11-2.75 (m, 4H), 2.66 (m, 1H), 2.36 (m, 1H), 2.14 (m, 2H), 1.80 (m, 1H), 1.65 (m, 1H), 1.28 (m, 5H); MS (EI) for C₂₅H₂₇N₄O₂FCl₂: 505 (MH⁺).

[0273] *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3*aR*,5*r*,6*aS*)-2-(2-methylpropyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.83 (d, 1H), 8.36 (d, 1H), 7.68 (d, 1H), 7.62 (t, 1H), 7.43 (d, 1H), 7.30 (d, 1H), 4.18 (d, 2H), 4.01 (s, 3H), 3.75 (m, 1H), 3.11 (m,

1H), 2.95 (m, 5H), 2.67 (m, 1H), 2.40 (m, 1H), 2.14 (m, 2H), 2.00 (m, 1H), 1.69 (m, 1H), 1.36 (m, 1H), 0.98 (t, 6H); MS (EI) for C₂₇H₃₁N₄O₂FCl₂: 533 (MH⁺).

Example 17

Ethyl (3a*R*,5*r*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate hydrochloride

[0274] **[0274]** Ethyl (3a*R*,5*r*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate hydrochloride: A solution of *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-[(3a*R*,5*r*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-ylmethyl]oxy}quinazolin-4-amine hydrobromide (0.050 g, 0.0830 mmol), triethylamine (0.046 mL, 0.0332 mmol) in 2.0 mL dichloromethane was cooled to 0°C and ethyl chloridocarbonate (0.010 mL, 0.0913 mmol) was added. The solution was stirred for 0.5 h at low temperature and quenched with saturated aqueous sodium bicarbonate. The reaction mixture was then partitioned between dichloromethane and saturated aqueous sodium bicarbonate. The layers were separated and the aqueous layer was extracted with dichloromethane (2 x 75 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 5% methanol in dichloromethane), followed by treatment in methanol with 4.0 M hydrogen chloride in dioxane (0.05 mL) and concentration provided ethyl (3a*R*,5*r*,6a*S*)-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate hydrochloride (27.7 mg, 53% yield). ¹H NMR (400 MHz, d₄-MeOH): 8.59 (s, 1H), 7.83 (s, 1H), 7.60 (d, 1H), 7.41 (t, 1H), 7.12 (s, 1H), 4.14 (d, 2H), 4.11 (m, 2H), 4.09 (s, 3H), 3.45 (dd, 2H), 3.30 (dd, 2H), 2.67 (m, 2H), 2.58 (m, 1H), 2.12 (m, 2H), 1.74 (m, 1H), 1.36 (m, 2H), 1.18 (t, 3H); MS (EI) for C₂₆H₂₇N₄O₄FClBr: 595 (MH⁺).

[0275] Using the same synthetic techniques and/or substituting with alternative reagents, the following compound of the invention were prepared:

[0276] *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-([[(3a*R*,5*r*,6a*S*)-2-(methylsulfonyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl]oxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.70 (s, 1H), 7.94 (s, 1H), 7.69 (d, 1H), 7.49 (t, 1H), 7.23 (s, 1H), 4.24 (d, 2H), 4.18 (m, 2H), 4.09 (s, 3H), 3.45 (dd, 2H), 2.90 (s,

3H), 2.87 (m, 3H), 2.59 (m, 1H), 2.28 (m, 2H), 1.43 (m, 2H); MS (EI) for $C_{24}H_{25}N_4O_4FSClBr$: 601 (MH^+).

[0277] 7-({[(3a*R*,5*r*,6a*S*)-2-acetyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-*N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride: ^1H NMR (400 MHz, d_4 -MeOH): 8.70 (s, 1H), 7.97 (s, 1H), 7.69 (d, 1H), 7.49 (t, 1H), 7.28 (s, 1H), 4.14 (d, 2H), 4.25 (m, 2H), 4.08 (s, 3H), 3.31 (m, 1H), 3.02 (m, 4H), 2.78 (m, 2H), 2.36 (m, 1H), 1.93 (m, 3H), 1.43 (m, 2H); MS (EI) for $C_{25}H_{25}N_4O_3FClBr$: 565 (MH^+).

Example 18

***N*-(3,4-dichlorophenyl)-7-({[(3a*R*,6a*S*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride:**

[0278] 1,1-Dimethylethyl (3a*R*,6a*S*)-5-(hydroxy)-hexahydrocyclopenta[*c*] pyrrole-2(1*H*)-carboxylate: Sodium borohydride (0.15 g, 4.00 mmol), was added to a solution of 1,1-dimethylethyl (3a*R*,6a*S*)-5-oxo-hexahydrocyclopenta[*c*] pyrrole-2(1*H*)-carboxylate (0.45 g, 2.00 mmol) in 10 mL methanol at 0°C and the reaction mixture was stirred for 1 h at this temperature. The solvent was evaporated, the crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), 1M aqueous hydrochloric acid and brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated to give 1,1-dimethylethyl (3a*R*,6a*S*)-5-(hydroxy)-hexahydrocyclopenta[*c*] pyrrole-2(1*H*)-carboxylate (0.44g, 98%). ^1H NMR (400 MHz, d_6 -DMSO): 4.08 (m, 1H), 3.40 (m, 2H), 3.30 (m, 2H), 2.50 (m, 2H), 1.98 (m, 2H), 1.40 (s, 9H), 1.30 (m, 2H). MS (EI) for $C_{12}H_{21}NO_3$: 228 (MH^+).

[0279] 1,1-Dimethylethyl (3a*R*,6a*S*)-5-{{[(methylsulfonyl)oxy]}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: Methanesulfonyl chloride (0.18 mL, 2.33 mmol), was added dropwise to a solution of 1,1-dimethylethyl (3a*R*,6a*S*)-5-(hydroxy)-hexahydrocyclopenta[*c*] pyrrole-2(1*H*)-carboxylate (0.44 g, 1.94 mmol) and triethylamine (0.81 mL, 5.81 mmol) in 10 mL dichloromethane at 0°C and the reaction mixture was stirred for 1 h at room temperature. The solvent was evaporated, the resulting crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), brine, 1M aqueous hydrochloric acid and brine again. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The resulting crude 1,1-dimethylethyl (3a*R*,6a*S*)-5-{{[(methylsulfonyl)oxy]}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate was used without further purification. MS (EI) for $C_{13}H_{23}NO_5S$: 306 (MH^+).

[0280] 1,1-dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy})hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: A solution of 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetate (salt) (0.22 g, 0.49 mmol), 1,1-dimethylethyl (3a*R*,6a*S*)-5-{{[(methysulfonyl)oxy]}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (0.15 g, 0.45 mmol), potassium carbonate (0.34 g, 2.50 mmol) in *N,N*-dimethylacetamide (5 mL) was heated in a sealed reaction tube at 90°C for 12 h. The crude reaction mixture was diluted with 100 mL 10% methanol in ethyl acetate and washed with saturated aqueous sodium bicarbonate (1x 30 mL), water (1 x 30 mL) and brine (1 x 30 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. Column chromatography (SiO₂, 3:2 hexanes:acetone) provided 1,1-dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy})hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (0.23 g, 98%). ¹H NMR (400 MHz, d₆-DMSO): 9.57 (s, 1H), 8.52 (s, 1H), 8.24 (d, 1H), 7.88 (dd, 1H), 7.78 (s, 1H), 7.62 (d, 1H), 7.13 (s, 1H), 5.15 (m, 1H), 3.96 (s, 3H), 3.42 (m, 2H), 3.36 (m, 2H), 2.80 (bs, 2H), 2.06 (m, 2H), 1.94 (m, 2H), 1.40 (s, 9H). MS (EI) for C₂₇H₃₀Cl₂N₄O₄: 547 (MH⁺).

[0281] *N*-(3,4-dichloro-phenyl)-6-(methyloxy)-7-{{[(3a*R*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]oxy} quinazolin-4-amine hydrochloride: 1,1-Dimethylethyl (3a*R*,6a*S*)-5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy})hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (0.23 g, 0.42 mmol) was taken up in methanol (10 mL) and treated with 4.0M hydrogen chloride in dioxane (excess) and heated briefly to reflux. Concentration *in vacuo* provided *N*-(3,4-dichloro-phenyl)-6-(methyloxy)-7-{{[(3a*R*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]oxy} quinazolin-4-amine hydrochloride (0.20 g, 100%). MS (EI) for C₂₂H₂₂Cl₂N₄O₂: 445 (MH⁺).

[0282] *N*-(3,4-Dichlorophenyl)-7-({[(3a*R*,6a*S*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: *N*-(3,4-Dichloro-phenyl)-6-(methyloxy)-7-{{[(3a*R*,6a*S*)-octahydrocyclopenta[*c*]pyrrol-5-yl]oxy} quinazolin-4-amine hydrochloride (0.20 g, 0.42 mmol) was solubilized in formic acid (5.0 mL) and 37% aqueous formaldehyde (1 mL) was added. The solution was heated to 95°C for 12 h. The reaction mixture was concentrated *in vacuo*. The residue was taken up in a mixture of 10% methanol in ethyl acetate (100 mL) and washed with saturated aqueous sodium bicarbonate (2x 30 mL) and brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by HPLC (reverse-phase,

acetonitrile/water/0.1% TFA). Upon removal of solvent the product was taken up in a mixture of 10% methanol in ethyl acetate (100 mL) and washed with saturated aqueous sodium bicarbonate (2x 30 mL) and brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated then the product was taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane (1eq.). Removal of solvent *in vacuo* provided 116 mg (56%) of *N*-(3,4-dichlorophenyl)-7-({[(3*aR*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride. ¹H NMR (400 MHz, d₆-DMSO): 11.05 (bs, 1H), 8.90 (s, 1H), 8.44 (s, 1H), 8.18 (d, 1H), 7.84 (dd, 1H), 7.76 (s, 1H), 7.48 (s, 1H), 5.30 (m, 1H), 4.00 (s, 3H), 3.35 (m, 2H), 2.90 (m, 2H), 2.24 (m, 5H), 2.10 (m, 2H), 1.24 (m, 2H). MS (EI) for C₂₃H₂₄Cl₂N₄O₂: 459, 461 (MH⁺).

Example 19

***N*-(4-bromo-3-chloro-2-fluorophenyl)-7-{{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0283] 3-(Chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine: A solution of (3*R*)-morpholin-3-ylmethanol (4.21 g, 36.0 mmol) in 2-(chloromethyl)oxirane (28.2 mL, 0.360 mol) was heated to 40°C for 3 h and then the solution was concentrated *in vacuo*. The intermediate was cooled in an ice bath and treated with 30.0 mL of concentrated sulfuric acid. The mixture was heated to 170°C for 2 h and then allowed to cool to room temperature. The mixture was poured into ice-water and solid sodium bicarbonate was carefully added until the solution was basic. 10% methanol in ethyl acetate was added and the biphasic mixture was filtered. The layers were separated and the aqueous layer was extracted (3 x 100 mL 10% methanol in ethyl acetate). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 2:5 hexanes:ethyl acetate) provided 3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine 2.44g (35%) as two separated diastereomers. (3*R*,9*aS*)-3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine: (0.886 g, 13% yield): ¹H NMR (400 MHz, CDCl₃): 3.91 (m, 3H), 3.82 (m, 1H), 3.68 (dt, 1H), 3.61 (dd, 1H), 3.47 (dd, 1H), 3.35 (t, 1H), 3.19 (t, 1H), 2.80 (d, 1H), 2.54 (m, 2H), 2.40 (m, 2H); MS (EI) for C₈H₁₄NO₂Cl: 192 (MH⁺). (3*S*,9*aS*)-3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine: (1.55 g, 22% yield): ¹H NMR (400 MHz, CDCl₃): 3.85 (m, 2H), 3.73 (m, 3H), 3.50 (m, 2H), 3.29 (t, 1H), 3.18 (t, 1H), 2.85 (dd, 1H), 2.64 (dd, 1H), 2.40 (m, 2H), 2.17 (t, 1H); MS (EI) for C₈H₁₄NO₂Cl: 192 (MH⁺).

[0284] Hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate: A suspension of (3*R*,9*aS*)-3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine (1.97 g, 10.3 mmol) and potassium acetate (10.1 g, 102 mmol) in DMF (20.0 mL) was stirred at 140°C for 16 h, and then at 150°C for another 12 h. The reaction mixture was partitioned between water (250 mL) and ethyl acetate (250 mL), the organic layer was washed with 5% lithium chloride (2 x 100 mL) and brine (100 mL) then dried over anhydrous sodium sulfate and concentrated *in vacuo*. Column chromatography (SiO₂, 1:1 hexane:ethyl acetate, then 100% ethyl acetate) afforded 0.92 g (42%) of hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate as a yellow oil. Distinct diastereomers as described above were converted in this step to give: (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate: ¹H NMR (400 MHz, CDCl₃): 4.18 (dd, 1H), 4.00 (m, 1H), 3.80 (dd, 1H), 3.68 (dt, 1H), 3.60 (dd, 1H), 3.46 (m, 2H), 3.22 (t, 1H), 2.64 (dd, 1H), 2.53 (m, 2H), 2.43-2.35 (m, 2H), 2.10 (s, 3H), and (3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate: ¹H NMR (400 MHz, CDCl₃): 4.09 (d, 2H), 3.90-3.82 (m, 2H), 3.75-3.64 (m, 3H), 3.27 (t, 1H), 3.18 (t, 1H), 2.69 (dd, 1H), 2.63 (m, 1H), 2.46-2.33 (m, 2H), 2.16 (t, 1H), 2.10 (s, 3H).

[0285] (3*R*,9*aS*)-Hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl methanesulfonate: To a solution of (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate (0.922 g, 4.28 mmol) in methanol (14.0 mL) was added 1.03 mL (4.50 mmol) of sodium methoxide (25% wt. in methanol) dropwise at room temperature. After 5 min., 1.6 mL (6.43 mmol) of 4.0M hydrogen chloride in dioxane was added and a pink precipitate formed. The solution was concentrated *in vacuo* and the pink solid was taken up in 30.0 mL dichloromethane. This slurry was cooled in an ice bath and triethylamine (3.0 mL, 21.5 mmol) was added, followed by methanesulfonyl chloride (0.37 mL, 4.71 mmol). The resultant yellow solution was stirred for 30 minutes at room temperature. The mixture was then partitioned between dichloromethane and saturated aqueous sodium bicarbonate then the aqueous layer was extracted (3 x 50 mL dichloromethane). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide crude (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl methanesulfonate which was taken on to the following reaction without purification.

[0286] *N*-(4-Bromo-3-chloro-2-fluorophenyl)-7-{-[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: A solution of (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-

ylmethyl methanesulfonate (0.215 g, 0.856 mmol), 4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-ol hydrochloride (0.247 g, 0.570 mmol), and potassium carbonate (0.400 g, 2.90 mmol) in DMF (1.9 mL) was heated in a sealed reaction tube at 75°C for 12 h, then 90°C for 12 h. The reaction mixture was concentrated *in vacuo* and the residue was partitioned between 10% methanol in ethyl acetate and saturated aqueous sodium bicarbonate. The layers were separated and the aqueous layer was extracted (3 x 50 mL 10% methanol in ethyl acetate). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The crude residue was purified by HPLC (reverse-phase, acetonitrile/water/0.1% TFA). Upon removal of solvent the product was taken up in methanol and treated with Bio-Rad AG 1-X8 resin (hydroxide form) until pH 8. The product was filtered and concentrated *in vacuo* then taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane (0.10 mL). Removal of solvent *in vacuo* provided 32.1 mg (10%) of *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride. MS (EI) for C₂₃H₂₃N₄O₄FClBr: 554 (M⁺).

[0287] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0288] *N*-(3,4-dichlorophenyl)-7-[(hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl)oxy]-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₆-DMSO): 8.90 (s, 1H), 8.50 (s, 1H), 8.19 (d, 1H), 7.85 (d, 1H), 7.75 (d, 1H), 7.42 (s, 1H), 4.51 (m, 1H), 4.32 (m, 2H), 4.04 (s, 3H), 4.00-3.62 (m, 4H); MS (EI) for C₂₃H₂₄N₄O₄Cl₂B: 491 (MH⁺).

[0289] *N*-(3,4-dichloro-2-fluorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.71 (s, 1H), 7.99 (s, 1H), 7.58-7.52 (m, 2H), 7.33 (s, 1H), 4.50 (m, 1H), 4.44 (d, 2H), 4.17-3.94 (m, 4H), 4.09 (s, 3H), 3.82-3.59 (m, 5H), 3.54-3.37 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄Cl₂F: 509 (MH⁺).

[0290] *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.71 (s, 1H), 7.99 (s, 1H), 7.69 (d, 1H), 7.49 (t, 1H), 7.32 (s, 1H), 4.49 (m, 1H), 4.44 (m, 2H), 4.16-3.95 (m, 4H), 4.10 (s, 3H), 3.82-3.58 (m, 5H), 3.54-3.35 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄BrClF: 553 (MH⁺).

- [0291] *N*-(3-chloro-2,4-difluorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: MS (EI) for C₂₃H₂₃N₄O₄ClF₂: 493 (MH⁺).
- [0292] *N*-(4,5-dichloro-2-fluorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.73 (s, 1H), 7.99 (s, 1H), 7.86 (d, 1H), 7.64 (d, 1H), 7.33 (s, 1H), 4.51 (m, 1H), 4.44 (d, 2H), 4.16-3.94 (m, 4H), 4.10 (s, 3H), 3.84-3.60 (m, 5H), 3.54-3.36 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄ClF₂: 509 (MH⁺).
- [0293] *N*-(4-bromo-5-chloro-2-fluorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.73 (s, 1H), 7.98 (s, 1H), 7.85 (d, 1H), 7.76 (d, 1H), 7.33 (s, 1H), 4.49 (m, 1H), 4.44 (d, 2H), 4.16-3.94 (m, 4H), 4.09 (s, 3H), 3.82-3.60 (m, 5H), 3.53-3.35 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄BrClF: 553 (MH⁺).
- [0294] *N*-(4-bromo-2,3-dichlorophenyl)-7-{[(3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.67 (s, 1H), 8.00 (s, 1H), 7.84 (d, 1H), 7.48 (d, 1H), 7.34 (s, 1H), 4.51 (m, 1H), 4.44 (d, 2H), 4.09 (s, 3H), 4.15-4.00 (m, 4H), 3.82-3.63 (m, 5H), 3.63-3.38 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄Cl₂Br: 570 (MH⁺).
- [0295] *N*-(4-bromo-2,3-dichlorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: MS (EI) for C₂₃H₂₃N₄O₄Cl₂Br: 570 (MH⁺).
- [0296] *N*-(3,4-dichloro-2-fluorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: MS (EI) for C₂₃H₂₃N₄O₄FCl₂: 509 (MH⁺).
- [0297] *N*-(4-bromo-5-chloro-2-fluorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: MS (EI) for C₂₃H₂₃N₄O₄FClBr: 554 (MH⁺).
- [0298] *N*-(4,5-dichloro-2-fluorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: ¹H NMR (400 MHz, d₄-MeOH): 8.74 (s, 1H), 8.00 (s, 1H), 7.86 (d, 1H), 7.64 (d, 1H), 7.35 (broad s, 1H), 4.51 (m, 2H), 4.44 (m, 1H), 4.25-3.95 (m, 4H), 4.09 (s, 3H), 3.82-3.63 (m, 5H), 3.63-3.38 (m, 2H); MS (EI) for C₂₃H₂₃N₄O₄FCl₂: 509 (MH⁺).

[0299] *N*-(3-chloro-2,4-difluorophenyl)-7-[[[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy]-6-(methyloxy)quinazolin-4-amine hydrochloride: MS (EI) for C₂₃H₂₃N₄O₄F₂Cl: 493 (MH⁺).

Example 20

N-(3,4-dichlorophenyl)-7-[(2-[[[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]amino}ethyl]oxy]-6-(methyloxy)quinazolin-4-amine

[0300] 7-[(2-Aminoethyl)oxy]-*N*-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride: A solution of 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetate (salt) (1.00 g, 2.15 mmol), 1,1-dimethylethyl (2-bromoethyl)carbamate (0.480 g, 2.15 mmol), and potassium carbonate (1.78 g, 12.9 mmol) in *N,N*-dimethylacetamide (2.2 mL) was heated to 100°C for 2.5 h. An additional 0.23 g (1.03 mmol) of 1,1-dimethylethyl (2-bromoethyl)carbamate was added and the reaction mixture was further heated to 100°C for a total of 7 h. The crude reaction mixture was partitioned between water and ethyl acetate and the layers were separated. The aqueous layer was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo*. Column chromatography (SiO₂, 3:2 hexanes:acetone) gave *N*-Boc product, which was then taken up in methanol and treated with 4M hydrogen chloride in dioxane while heating. Dilution with ethyl ether precipitated a pale yellow solid, which was collected by filtration and dried to give 0.761 g (94%) of 7-[(2-aminoethyl)oxy]-*N*-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride. ¹H NMR (400 MHz, d₆-DMSO): 12.01 (s, 1H), 8.94 (s, 1H), 8.67 (s, 1H), 8.35 (broad s, 2H), 8.21 (s, 1H), 7.90 (dd, 1H), 7.75 (d, 1H), 7.53 (s, 1H), 4.43 (t, 2H), 4.08 (s, 3H), 3.36 (m, 2H).

[0301] *N*-(3,4-Dichlorophenyl)-7-[(2-[[[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]amino}ethyl]oxy]-6-(methyloxy)quinazolin-4-amine: To a DMF solution (3.0 mL) of 7-[(2-aminoethyl)oxy]-*N*-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride (56.8 mg, 0.137 mmol) at room temperature, was added glacial acetic acid (3 drops), (1*R*,5*S*)-8-methyl-8-azabicyclo[3.2.1]octan-3-one (49.0 mg, 0.137 mmol), and sodium triacetoxyborohydride (43.0 mg, 0.205 mmol). After stirring for 12 h, additional (1*R*,5*S*)-8-methyl-8-azabicyclo[3.2.1]octan-3-one (50.0 mg, 0.140 mmol), acetic acid (3 drops) and sodium triacetoxyborohydride were added. The solution was quenched with water, filtered and purified by HPLC (reverse-phase, acetonitrile/water/0.1% TFA). Upon removal of solvent, the product was taken up in methanol and treated with Bio-Rad

AG 1-X8 resin (hydroxide form) until pH 8. The product was filtered and concentrated *in vacuo*, to provide 45.1 mg (66%) of *N*-(3,4-dichlorophenyl)-7-[(2-[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]amino}ethyl)oxy]-6-(methyloxy)quinazolin-4-amine. ¹H NMR (400 MHz, d₆-DMSO): 9.63 (s, 1H), 8.55 (s, 1H), 8.26 (d, 1H), 7.90 (dd, 1H), 7.83 (s, 1H), 7.65 (s, 1H), 7.23 (d, 1H), 4.20 (t, 1H), 4.15 (t, 3H), 3.97 (s, 1H), 3.03 (m, 1H), 2.91 (t, 2H), 2.81 (t, 1H), 2.17 (s, 3H), 2.00-1.84 (m, 6H), 1.67-1.32 (m, 4H); MS (EI) for C₂₅H₂₉N₅O₂Cl₂: 502 (MH⁺).

Example 21

***N*-(3,4-dichlorophenyl)-7-[(3-*exo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0302] *N*-(3,4-Dichlorophenyl)-7-[(3-*exo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride: A solution of 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetate (salt) (0.150 g, 0.322 mmol), (3-*endo*)-3-[(methylsulfonyl)oxy]-8-azabicyclo[3.2.1]octane (0.106 g, 0.483 mmol), and potassium carbonate (0.220 g, 1.60 mmol) in *N,N*-dimethylacetamide (1.1 mL) was heated in a sealed tube at 100°C for 12 h, followed by 48 h at room temperature. The crude reaction mixture was filtered through celite using methanol eluent, and the solvents were removed *in vacuo*. The residue was purified by HPLC (reverse-phase, acetonitrile/water/0.1% TFA). Upon removal of solvent, the product was taken up in methanol and treated with Bio-Rad AG 1-X8 resin (hydroxide form) until pH 8. The product was filtered and concentrated *in vacuo*, then taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane (0.050 mL). Removal of solvent *in vacuo* provided 48.7 mg (31%) of *N*-(3,4-dichlorophenyl)-7-[(3-*exo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride. ¹H NMR (400 MHz, d₆-DMSO): 10.69 (s, 1H), 8.92 (s, 1H), 8.32 (s, 1H), 8.17 (d, 1H), 7.81 (m, 2H), 7.75 (d, 1H), 5.05 (m, 1H), 4.02 (s, 3H), 2.69 (d, 2H), 2.39 (m, 1H), 2.29-2.18 (m, 6H); MS (EI) for C₂₃H₂₄N₄O₂Cl₂: 459 (MH⁺).

Example 22

***N*-(3,4-dichlorophenyl)-7-[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0303] 7-[(3-*endo*)-8-Azabicyclo[3.2.1]oct-3-ylmethyl]oxy}-*N*-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride: A solution of 4-[(3,4-

dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl trifluoroacetate (salt) (0.200 g, 0.429 mmol), 1,1-dimethylethyl (3-*endo*)-3-[[[(methanesulfonyl)oxy]methyl]-8-azabicyclo[3.2.1]octane-8-carboxylate (0.200 g, 0.626 mmol), and potassium carbonate (0.300 g, 2.17 mmol) in N,N-dimethylacetamide (1.4 mL) was heated in a sealed tube at 110°C for 12 h. Additional mesylate (0.430 g, 1.35 mmol) was added and the mixture was heated for 2 h at 110°C. The crude reaction mixture was partitioned between 10% methanol in ethyl acetate (50 mL) and water (50 mL). The layers were separated and the organic layer was washed with water (2 x 50 mL) and 1M aqueous sodium hydroxide (1 x 50 mL). The organic layer was dried over anhydrous sodium sulfate and the solvent was removed *in vacuo*. The residue was purified by column chromatography (SiO₂, 2:1 hexanes:ethyl acetate), followed by HPLC (reverse-phase, acetonitrile/water/0.1% TFA). Solvent was removed *in vacuo* and the residue partitioned with 10% methanol in ethyl acetate and water. The aqueous layer was made basic with saturated aqueous sodium bicarbonate. The layers were separated and the aqueous layer was further extracted with 10% methanol in ethyl acetate (2x). The combined organic extracts were dried over anhydrous sodium sulfate, filtered and the concentrated *in vacuo*. The residue was taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane then concentrated to provide 7-[[[(3-*endo*)-8-azabicyclo[3.2.1]oct-3-ylmethyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride (0.104 g, 53%). MS (EI) for C₂₃H₂₄N₄O₂Cl₂: 459 (MH⁺).

[0304] N-(3,4-Dichlorophenyl)-7-([[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: 7-[[[(3-*Endo*)-8-azabicyclo[3.2.1]oct-3-ylmethyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride, (0.104 g, 0.210 mmol) was combined with 37% aqueous formaldehyde (0.10 mL, 1.26 mmol) in formic acid (1.0 mL) and the solution was heated to 110°C for 12 h. The solvent was removed *in vacuo* and the residue was taken up in methanol and treated with Bio-Rad AG 1-X8 resin (hydroxide form) until pH 8. The product was filtered and concentrated *in vacuo*. The residue was purified by HPLC (reverse-phase, acetonitrile/water/0.1% TFA). Upon removal of solvent, the product was taken up in methanol and treated with Bio-Rad AG 1-X8 resin (hydroxide form) until pH 8. The product was filtered and concentrated *in vacuo*, then taken up in methanol and treated with 4.0 M hydrogen chloride in dioxane (0.10 mL). Removal of solvent *in vacuo* provided 41.4 mg (39%) of N-(3,4-dichlorophenyl)-7-([[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride.

¹H NMR (400 MHz, d₆-DMSO): 8.89 (s, 2H), 8.45 (s, 1H), 8.20 (s, 1H), 7.86 (d, 1H), 7.75 (d, 1H), 7.46 (s, 1H), 4.24 (m, 2H), 4.04 (s, 3H), 3.97 (broad s, 1H), 3.85 (broad s, 1H), 2.65 (d, 1H), 2.25-2.51 (m, 6H), 2.03-1.80 (m, 5H); MS (EI) for C₂₄H₂₆N₄O₂Cl₂: 473 (MH⁺).

Example 23

***N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-{[(8*aR*)-tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazin-6-ylmethyl]oxy}quinazolin-4-amine trifluoroacetate**

[0305] (8*aR*)-6-(Chloromethyl)tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazine: A solution of (4*R*)-1,3-thiazolidin-4-ylmethanol (0.300 g, 2.52 mmol) in 2-(chloromethyl)oxirane (2.0 mL, 25.5 mmol) was heated under nitrogen to 40°C for 12 h. The solution was then cooled to room temperature and 2-(chloromethyl)oxirane was removed *in vacuo*. The crude intermediate was cooled in ice, and was taken up in 2.0 mL of concentrated sulfuric acid. The resulting mixture was heated to 200°C for 0.5 h then poured carefully onto wet ice, which was allowed to melt. The aqueous solution was carefully made basic using solid sodium bicarbonate and the resulting mixture was filtered using water and 10% methanol in ethyl acetate as eluent. The layers were separated and the aqueous layer was extracted with 10% methanol in ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give 11.6 mg (2.4% yield) of crude (8*aR*)-6-(chloromethyl)tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazine as a mixture of diastereomers which was directly taken on to the next step.

[0306] *N*-(3,4-Dichlorophenyl)-6-(methyloxy)-7-{[(8*aR*)-tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazin-6-ylmethyl]oxy}quinazolin-4-amine trifluoroacetate: A solution of (8*aR*)-6-(chloromethyl)tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazine (11.6 mg, 0.0599 mmol), 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol trifluoroacetate (salt) (30.0 mg, 0.0644 mmol), and potassium carbonate (45.0 mg, 0.326 mmol) in *N,N*-dimethylacetamide (1.0 mL) was heated in a sealed tube to 150°C for 12 h. The crude reaction mixture was directly purified via reverse-phase preparative HPLC (acetonitrile/water/0.1% TFA). Lyophilization of the pure fractions yielded 3.5 mg (8.9%) of *N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-{[(8*aR*)-tetrahydro-1*H*-[1,3]thiazolo[4,3-*c*][1,4]oxazin-6-ylmethyl]oxy}quinazolin-4-amine trifluoroacetate. ¹H NMR (400 MHz, d₆-DMSO): 8.80 (s, 1H), 8.11 (s, 1H), 7.99 (s, 1H), 7.74 (s, 2H), 7.29

(s, 1H), 4.29 (d, 2H), 4.11 (m, 2H), 4.00 (s, 3H), 3.96 (m, 1H), 2.99 (m, 2H), 2.56 (t, 1H), 2.367 (m, 1H); MS (EI) for C₂₂H₂₂N₄O₃SCl₂: 492 (MH⁺).

Example 24

***N*-(3,4-dichlorophenyl)-7-({2-[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0307] 1,1-Dimethylethyl (3-*endo*)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate: To a solution of 1,1-dimethylethyl (3-*endo*)-3-(2-hydroxyethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate (30.3 mg, 1.19 mmol) in dichloromethane (4.0 mL), was added triethylamine (0.5 mL, 3.56 mmol) and the solution was cooled to 0°C under nitrogen. Methanesulfonyl chloride (0.11 mL, 1.42 mmol) was added slowly and mixture was allowed to warm to room temperature and stirred for 1h. The reaction mixture was partitioned between dichloromethane and water. The aqueous phase was extracted with dichloromethane (2 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide 35.1 mg (89%) of 1,1-dimethylethyl (3-*endo*)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate, which was carried forward without purification.

[0308] 1,1-Dimethylethyl (3-*endo*)-3-(2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate: To a solution of 1,1-dimethylethyl (3-*endo*)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate (0.175 g, 0.526 mmol) in N,N-dimethylacetamide (3.5 mL) was added 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl trifluoroacetate (salt) (0.490 g, 1.05 mmol) and potassium carbonate (0.728 g, 5.26 mmol), and the reaction was stirred at 110°C for 18h. An additional portion of 1,1-dimethylethyl (3-*endo*)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate (0.175 g, 0.526 mmol) was added and the mixture was stirred at 140°C for 2h. Another portion of the mesylate (0.300 g, 1.05 mmol) in N,N-dimethylacetamide (4.0 mL) was added and the mixture continued to stir at 140°C for a further 18h. The reaction mixture was concentrated *in vacuo*, and the residue was partitioned between 10% methanol in ethyl acetate and water. The organic layer was washed (3 x 50 mL water) and the combined aqueous portions were extracted (2 x 100 mL 10% methanol in ethyl acetate). All organic layers were combined, dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by HPLC (reverse-phase,

acetonitrile/water/0.1% TFA). Upon concentration the remaining aqueous layer was neutralized with solid sodium bicarbonate, extracted (100 mL 10% methanol in ethyl acetate), dried over anhydrous sodium sulfate then filtered and concentrated *in vacuo* to afford 1,1-dimethylethyl (3-*endo*)-3-(2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate (39.9 mg, 66% yield). ¹H NMR (400 MHz; d₆-DMSO): 9.43 (broad s, 1H), 8.48 (s, 1H), 7.87 (s, 1H), 7.61 (d, 1H), 7.53 (s, 1H), 7.22 (d, 1H), 7.02 (s, 1H), 4.23-3.82 (m, 4H), 3.80 (s, 3H), 2.19 (m, 1H), 1.93 (s, 6H), 1.69-1.42 (m, 3H), 1.36 (s, 9H), 1.22 (m, 1H).

[0309] 7-({2-[(3-*endo*)-8-Azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-*N*-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine hydrochloride: 1,1-Dimethylethyl (3-*endo*)-3-(2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate was solubilized in methanol (2.3 mL) and treated with 4.0 M hydrogen chloride in dioxane (2.3 mL). The solution was heated to reflux then immediately allowed to cool to room temperature. The solution was then concentrated *in vacuo* to give 7-({2-[(3-*endo*)-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-*N*-(3,4-dichlorophenyl)-6-(methyloxy) quinazolin-4-amine hydrochloride (34.6 mg, 98% yield). MS (EI) for C₂₄H₂₆Cl₂N₄O₂: 473 (MH⁺).

[0310] *N*-(3,4-Dichlorophenyl)-7-({2-[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: To a solution of 7-({2-[(3-*endo*)-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-*N*-(3,4-dichlorophenyl)-6-(methyloxy) quinazolin-4-amine hydrochloride (0.346 g, 0.678 mmol) in formic acid (2.7 mL), was added aqueous formaldehyde (37%, 0.27 mL, 4.07 mmol) and the mixture was heated to 110°C for 5h, then allowed to cool to room temperature. The solution was concentrated *in vacuo* and the residue was taken up in methanol and treated with AG 1-X8 resin (hydroxide form) to pH 8. The mixture was filtered and concentrated then the residue purified by HPLC (reverse-phase, acetonitrile/water/0.1% TFA) and the pure fractions lyophilized. The residue was taken up in methanol and neutralized with AG 1-X8 resin (hydroxide form) to pH 8 then filtered and concentrated. The residue was taken into methanol (3 mL) and treated with 4.0 M hydrogen chloride in dioxane to pH 2. Concentration *in vacuo* afforded the title compound *N*-(3,4-dichlorophenyl)-7-({2-[(3-*endo*)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride (12.5 mg, 36% yield). ¹H NMR (400 MHz; d₆-DMSO): 8.86 (s, 2H), 8.40 (s, 1H), 8.16 (t, 1H), 7.82 (d, 1H), 7.73 (d, 1H), 7.38 (s, 1H), 4.22 (m, 2H), 4.03 (s, 3H),

3.92 (broad s, 2H), 2.28 (m, 2H), 2.12-1.91 (m, 6H), 1.88-1.58 (m, 6H); MS (EI) for $C_{25}H_{28}Cl_2N_4O_2$: 485 (MH^+).

Example 25

1,4:3,6-Dianhydro-5-O-{4-[(3-chloro-2-fluorophenyl)amino]-6-(methoxy)quinazolin-7-yl}-2-deoxy-2-fluoro-L-iditol

[0311] 1,4:3,6-dianhydro-2-O-[4-(methoxy)carbonyl-2-(methoxy)phenyl]-5-O-(methylsulfonyl)-D-glucitol: 1,4:3,6-dianhydro-2,5-bis-O-(methylsulfonyl)-D-mannitol (15.6 g, 51.7 mmol) and methyl vanillate (9.40 g, 51.7 mmol) were dissolved in DMF (60 mL) and cesium carbonate (33.7 g, 103 mmol) was added. The mixture was stirred at 100°C for 2.5 h then cooled to room temperature and filtered. The filtrate was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. Hexanes were added to the biphasic mixture and the precipitate was removed by filtration and discarded. From the filtrate a second crop of precipitate was obtained (8.7g) and the material was purified by silica gel column chromatography using 30% ethyl acetate in chloroform eluent to afford 1,4:3,6-dianhydro-2-O-[4-(methoxy)carbonyl-2-(methoxy)phenyl]-5-O-(methylsulfonyl)-D-glucitol as a colorless solid (3.08 g, 15% yield). ^1H -NMR (400 MHz, $CDCl_3$): 7.65 (dd, 1H), 7.57 (d, 1H), 6.95 (d, 1H), 5.12 (q, 1H), 4.95 (t, 1H), 4.93-4.90 (m, 1H), 4.67 (d, 1H), 4.28-4.23 (m, 1H), 4.19 (dd, 1H), 4.03 (dd, 1H), 3.95 (dd, 1H), 3.90 (2 x s, 6H), 3.15 (s, 3H).

[0312] 1,4:3,6-dianhydro-2-O-[4-(methoxy)carbonyl-2-(methoxy)phenyl]-5-O-(methylsulfonyl)-D-glucitol (4.38 g, 11.3 mmol) was taken into methyl sulfoxide (30 mL) followed by addition of potassium fluoride (7.5 g, 128 mmol) and the mixture was heated to 180°C over 12 hours. The mixture was cooled to room temperature and partitioned with ethyl acetate and water. The organic layer was washed with water (3x) then brine and dried over anhydrous magnesium sulfate. Filtration and concentration of the organic solution followed by further purification by silica gel column chromatography using hexanes:ethyl acetate (2:1) eluent afforded 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-{2-(methoxy)-4-[(methoxy)carbonyl]phenyl}-L-iditol (1.5g, 43% yield) as a colorless oil which was employed in the next step without further purification. GCMS: Calculated for $C_{15}H_{17}O_6F$: 312 (M^+).

[0313] 1,4:3,6-Dianhydro-2-deoxy-2-fluoro-5-O-{2-(methoxy)-4-[(methoxy)carbonyl]phenyl}-L-iditol (4.8 g, 15.4 mmol) was taken into dichloromethane (45 mL) and the solution cooled to 0°C. Fuming nitric acid (90% reagent, 1.3 mL) was added dropwise to

the solution followed by addition of concentrated sulfuric acid (0.3 mL). Two additional aliquots of both acids were added at ten minute intervals and the mixture was allowed to warm to room temperature with stirring an additional 20 minutes. An excess of ethyl acetate (100 mL) was added to the mixture followed by water (50 mL) and the organic layer subsequently washed with water (1x), saturated aqueous sodium bicarbonate (2x) and brine then dried over anhydrous magnesium sulfate, filtered and concentrated to give 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-{2-(methyloxy)-4-[(methyloxy)carbonyl]-5-nitrophenyl}-L-iditol (5.0g, 91% yield) as a yellow amorphous residue. The material was hydrogenated at 50 psi hydrogen gas pressure in a Parr apparatus in methanol solution (50 mL) using 10% Pd/C catalyst (1.0 g) over 12 hours. Filtration of the catalyst and concentration of the organic solution afforded 5-*O*-{5-amino-2-(methyloxy)-4-[(methyloxy)carbonyl]phenyl}-1,4:3,6-dianhydro-2-deoxy-2-fluoro-L-iditol (4.5 g, 90% overall yield) as a solid. ¹H-NMR (400 MHz, CDCl₃): 7.32 (s, 1H), 6.24 (s, 1H), 5.55-5.62 (br s, 2H), 5.10 (dd, 1H), 4.82 (d, 2H), 4.79 (m, 1H), 4.81-4.03 (m, 4H), 3.93 (dd, 1H), 3.85 (s, 3H), 3.78 (s, 3H). MS (EI) for C₁₅H₁₈NO₆F: 328 (MH⁺).

[0314] 5-*O*-{5-Amino-2-(methyloxy)-4-[(methyloxy)carbonyl]phenyl}-1,4:3,6-dianhydro-2-deoxy-2-fluoro-L-iditol (4.5 g 13.7 mmol.) was taken into formamide (40 mL) followed by addition of ammonium formate (1.7g, 27.5 mmol.) and the mixture was heated to 165°C for 2.5 hours. The mixture was then cooled to room temperature and partitioned with ethyl acetate:hexanes (1:1) and water to give a biphasic suspension. The residue was collected by filtration, washed with water then ethyl ether and dried *in vacuo* to give 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-[6-(methyloxy)-4-oxo-3,4-dihydroquinazolin-7-yl]-L-iditol (3.61 g, 82% yield) as a tan solid. MS (EI) for C₁₅H₁₅N₂O₅F: 323 (MH⁺).

[0315] 1,4:3,6-Dianhydro-2-deoxy-2-fluoro-5-*O*-[6-(methyloxy)-4-oxo-3,4-dihydroquinazolin-7-yl]-L-iditol (3.61 g, 11.2 mmol.) was suspended in chloroform (50 mL) followed by addition of DMF (1.0 mL) and oxalyl chloride (2.0 mL) then the mixture was brought to reflux for 10 minutes then cooled followed by addition of DMF (0.5 mL) and oxalyl chloride (1.0 mL) and the mixture was brought to reflux an additional hour. The mixture was again allowed to cool to room temperature and neutralized by slow addition of saturated aqueous sodium bicarbonate. The mixture was extracted with chloroform (2x) then ethyl acetate (1x) and the combined organic layers were dried over anhydrous magnesium sulfate, filtered and concentrated to give 1,4:3,6-dianhydro-5-*O*-[4-chloro-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-L-iditol (3.34 g, 88% yield). ¹H-NMR (400 MHz, d₆-DMSO): 8.91 (s, 1H), 7.57 (s, 1H), 7.45 (s, 1H), 5.24 (dd, 1H), 5.24 (br s,

1H), 4.84-4.79 (m, 2H), 4.17-3.87 (m, 4H), 4.01 (s, 3H). MS (EI) for $C_{15}H_{14}N_2O_4FCl$: 341 (MH^+).

[0316] 1,4:3,6-Dianhydro-5-*O*-[4-chloro-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-L-iditol (3.34 g, 9.8 mmol.) and 3-chloro-2-fluoroaniline hydrochloride (2.0 g, 10.8 mmol.) were taken into acetonitrile (50 mL) and the mixture was brought to reflux for 2.5 hours. The resulting suspension was cooled to room temperature and diluted with an excess of ethyl ether. The solid product was collected by filtration and recrystallized from a minimum of warm methanol with addition of ethyl ether. The solid was collected by filtration and dried *in vacuo* to give 1,4:3,6-dianhydro-5-*O*-{4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl}-2-deoxy-2-fluoro-L-iditol (4.2 g, 95% yield) as an off white solid. ^1H -NMR (400 MHz, d_6 -DMSO): 12.11 (s, 1H), 8.84 (s, 1H), 8.52 (s, 1H), 7.64 (tr, 1H), 7.53 (tr, 1H), 7.50 (s, 1H), 7.35 (tr, 1H), 5.24 (dd, 1H), 5.11 (tr, 1H), 4.16-3.89 (m, 4H), 4.02 (s, 3H). MS (EI) for $C_{21}H_{18}N_3O_4F_2Cl$: 450 (MH^+).

[0317]

Example 26

N-(3,4-dichlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine

[0372] A solution of 2-(chloromethyl)-4-(phenylmethyl)morpholine (498 mg, 2.21 mmol) in methanol (20 mL) was hydrogenated over 10% Pd-C (120 mg) for 17 h. The catalyst was filtered off, and the filtrate was concentrated to give 2-(chloromethyl)morpholine as a colorless oil. To a solution of this oil in methanol (20 mL) was added di-*tert*-butyl dicarbonate (425 mg, 1.95 mmol), and the reaction mixture was stirred at room temperature for 5 h. Concentration and purification by column chromatography on silica (9:1 hexanes/ethyl acetate) provided 1,1-dimethylethyl 2-(chloromethyl)morpholine-2-carboxylate 283 mg (54%) as a colorless solid. ^1H NMR (400 MHz, d_4 -MeOH): 4.02 (d, 1H), 3.88 (d, 1H), 3.81 (m, 1H), 3.61-3.47 (m, 4H), 2.95 (br. s, 1H), 2.75 (br. s, 1H), 1.46 (s, 9H).

[0373] To a solution of 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol (12.95 g, 38.52 mmol) in dimethylacetamide (80 mL) was added 1,1-dimethylethyl 2-(chloromethyl)morpholine-2-carboxylate (11.94 g, 50.66 mmol) and potassium carbonate (15.97 g, 115.55 mmol), and the reaction mixture was stirred at 130°C under nitrogen for 13 h. After cooling to room temperature, the mixture was partitioned between water (500 mL) and ethyl acetate (250 mL). The layers were separated, the organic layer diluted

with hexanes (250 mL) and washed with water (200 mL). The combined aqueous layers were further extracted with ethyl acetate (2 x 200 mL). Some product precipitated from the combined organic layers and was filtered. The solid was washed with methanol (2 x 50 mL) and dried to give 4.77 g (25%) of 1,1-dimethylethyl 2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}morpholine-4-carboxylate as a tan solid. The methanol washes were concentrated, combined with the filtrate, washed with 5% lithium chloride (2 x 200 mL) and brine (200 mL), dried over sodium sulfate and concentrated. Crystallization from methanol gave another 6.86 g (36%). ¹H NMR (400 MHz, d₄-MeOH): 8.47 (s, 1H), 8.15 (d, 1H), 7.74 (dd, 1H), 7.70 (s, 1H), 7.49 (d, 1H), 7.15 (s, 1H), 4.24-4.06 (m, 3H), 4.02 (s, 3H), 3.98-3.82 (m, 3H), 3.59 (m, 1H), 3.00 (br.s, 2H), 1.48 (s, 9H); MS (EI) for C₂₅H₂₈N₄O₅Cl₂: 535 (MH⁺).

[0374] To a suspension of 1,1-dimethylethyl 2-({[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}morpholine-4-carboxylate (4.77 g, 8.91 mmol) in methanol (50 mL) was added a 4M solution of HCl in 1,4-dioxane (50 mL), and the reaction mixture was refluxed for 5 min. After cooling to room temperature, diethyl ether (100 mL) was added and the precipitate was filtered and dried. The solid was dissolved in methanol (200 mL) and treated with Bio-Rad 1-X8 resin hydroxide form until pH 8. Filtration and concentration *in vacuo* provided 3.32 g (86%) of N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine as a brown solid. ¹H NMR (400 MHz, d₄-MeOH): 8.47 (s, 1H), 8.14 (d, 1H), 7.73 (m, 2H), 7.49 (d, 1H), 7.15 (s, 1H), 4.24-4.13 (m, 2H), 4.09-3.97 (m, 5H), 3.76 (m, 1H), 3.22 (dd, 1H), 3.02-2.90 (m, 3H); MS (EI) for C₂₀H₂₀N₄O₃Cl₂: 435 (MH⁺).

[0375] To a solution of N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine (3.32 g, 7.7 mmol) in formic acid (33.2 mL) was added a 37wt% solution of formaldehyde in water (3.32 mL), and the reaction mixture was heated at 95°C for 2.5 h. After cooling to room temperature, the reaction mixture was concentrated. The residue was dissolved in methanol and treated with Bio-Rad AG[®]1-X8 hydroxide form resin until pH 8. Filtration and concentration *in vacuo* afforded 3.11 g (91%) of the title compound as a yellow-brown solid. ¹H NMR (400 MHz, d₄-MeOH): 8.46 (s, 1H), 8.14 (d, 1H), 7.73 (dd, 1H), 7.68 (s, 1H), 7.48 (d, 1H), 7.13 (s, 1H), 4.21-4.09 (m, 2H), 4.02 (s, 3H), 4.01-3.91 (m, 3H), 2.96 (m, 1H), 2.75 (m, 1H), 2.35 (s, 3H), 2.22 (m, 1H), 2.11 (t, 1H); MS (EI) for C₂₁H₂₂N₄O₃Cl₂: 449 (MH⁺).

[0376] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0377] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(phenylmethyl)morpholin-2-yl]methyl)oxy]quinazolin-4-amine: ^1H NMR (400 MHz, $\text{d}_4\text{-MeOH}$): 8.48 (s, 1H), 8.15 (d, 1H), 7.74 (m, 2H), 7.50 (d, 1H), 7.38-7.24 (m, 5H), 7.16 (s, 1H), 4.21-4.09 (m, 2H), 4.06-3.99 (m, 4H), 3.91 (m, 1H), 3.72 (m, 1H), 3.58 (s, 2H), 2.96 (m, 1H), 2.73 (m, 1H), 2.25 (m, 1H), 2.16 (t, 1H); MS (EI) for $\text{C}_{27}\text{H}_{26}\text{N}_4\text{O}_3\text{Cl}_2$: 525 (MH^+).

[0378] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(1,4-oxazepan-2-ylmethyl)oxy]quinazolin-4-amine: ^1H NMR (400 MHz, $\text{d}_4\text{-MeOH}$): 8.47 (s, 1H), 8.14 (d, 1H), 7.73 (m, 2H), 7.49 (d, 1H), 7.15 (s, 1H), 4.20-4.00 (m, 7H), 3.80 (m, 1H), 3.21 (dd, 1H), 2.95 (m, 3H), 1.91 (m, 2H); MS (EI) for $\text{C}_{21}\text{H}_{22}\text{N}_4\text{O}_3\text{Cl}_2$: 449 (MH^+).

[0379] N-(3,4-dichlorophenyl)-7-([(4-methyl-1,4-oxazepan-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine: ^1H NMR (400 MHz, $\text{d}_4\text{-MeOH}$): 8.46 (s, 1H), 8.14 (d, 1H), 7.73 (dd, 1H), 7.70 (s, 1H), 7.49 (d, 1H), 7.14 (s, 1H), 4.23-4.13 (m, 2H), 4.09-3.93 (m, 5H), 3.87 (m, 1H), 3.08 (d, 1H), 2.88 (m, 1H), 2.65 (m, 2H), 2.46 (s, 3H), 2.05 (m, 1H), 1.92 (m, 1H); MS (EI) for $\text{C}_{22}\text{H}_{24}\text{N}_4\text{O}_3\text{Cl}_2$: 463 (MH^+).

[0380] N-(3,4-dichlorophenyl)-7-([(4-ethyl-1,4-oxazepan-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride: ^1H NMR (400 MHz, $\text{d}_4\text{-MeOH}$): 8.77 (s, 1H), 8.06-8.05 (m, 2H), 7.72 (dd, 1H), 7.60 (d, 1H), 7.31 (s, 1H), 4.62-4.32 (m, 3H), 4.30 (s, 3H), 4.04-3.36 (m, 8H), 2.40-2.08 (m, 2H), 1.45 (tr, 3H); MS (EI) for $\text{C}_{23}\text{H}_{26}\text{N}_4\text{O}_3\text{Cl}_2$: 477 (MH^+).

[0381] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]quinazolin-4-amine: ^1H NMR (400MHz, $\text{d}_6\text{-DMSO}$): 9.55 (s, 1H), 8.51 (s, 1H), 8.24 (d, 1H), 7.88 (dd, 1H), 7.77 (s, 1H), 7.61 (d, 1H), 7.17 (s, 1H), 4.05 (m, 2H), 3.97 (s, 3H), 3.92-3.88 (m, 1H), 3.73-3.65 (m, 1H), 3.42-3.38 (m, 2H), 1.88-1.81 (m, 1H), 1.70-1.64 (m, 1H), 1.53-1.35 (m, 3H); MS (EI) for $\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_3\text{Cl}_2$: 434 (MH^+).

Example 27

N-(3,4-dichlorophenyl)-7-([(2-[(dimethylamino)methyl]-1,3-thiazol-4-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine

[0382] (Dimethylamino)acetonitrile (1.16 mL, 11.9 mmol) was added to a mixture of DMF (5 mL) and triethylamine (2 mL). Hydrogen sulfide gas was bubbled into the solution until the solution was dark green. The solution was then heated at 70°C for 0.5

h. Excess hydrogen sulfide was removed from the system by bubbling nitrogen into the solution and then the volatiles were removed *in vacuo*. to afford 2-(dimethylamino)ethanethioamide as burgundy crystals (0.660 g, 5.59 mmol, 47% yield). ^1H NMR (400 MHz; CDCl_3): 2.30 (s, 6H); 3.37 (s, 2H); 7.80 (br s, 1H); 8.80 (br s, 1H); GCMS for $\text{C}_4\text{H}_{10}\text{N}_2\text{S}$: 118 (M^+).

[0383] 2-(Dimethylamino)ethanethioamide (0.382 g, 3.24 mmol), 1,3-dichloroacetone (0.453 g, 3.56 mmol) and sodium bicarbonate (0.301 g, 3.56 mmol) was stirred in 1,2-dichloroethane (4 mL) at room temperature for 24 h. The reaction mixture was filtered and the filter cake was washed with 1,2-dichloroethane. The filtrate was added dropwise to a cooled (ice bath) solution of thionyl chloride (0.260 mL, 3.56 mmol) in 1,2-dichloroethane (2 mL). The solution was stirred at 70°C for 0.5 h and then cooled to room temperature. The brown mixture was filtered and washed with 1,2-dichloroethane to afford N-{[4-(chloromethyl)-1,3-thiazol-2-yl]methyl}-N,N-dimethylamine hydrochloride as a brown solid (0.550 g, 2.42 mmol, 75% yield). ^1H NMR (400 MHz; D_2O): 2.79 (s, 6H); 4.52 (s, 2H); 4.61 (s, 2H); 7.60 (s, 1H).

[0384] N-{[4-(Chloromethyl)-1,3-thiazol-2-yl]methyl}-N,N-dimethylamine hydrochloride (0.050 g, 0.220 mmol) and 4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (0.074 g, 0.220 mmol) were suspended in DMF (4 mL) and potassium carbonate (0.152 g, 1.10 mmol) was added. The mixture was stirred at room temperature for 50 h and then at 70°C for 3.5 h. An additional portion of N-{[4-(chloromethyl)-1,3-thiazol-2-yl]methyl}-N,N-dimethylamine hydrochloride (0.019 g, 0.084 mmol) was added and the mixture was stirred at 70°C for a further 25 h. The reaction mixture was concentrated *in vacuo*. and the residue was partitioned between ethyl acetate and brine. The organic portion was washed with 1 N aqueous sodium hydroxide, brine, dried over sodium sulfate, filtered and concentrated *in vacuo*. to afford a brown solid. Recrystallization from ethyl acetate / diethyl ether afforded the title compound as a pale brown solid (0.016 g, 0.033 mmol, 15% yield). ^1H NMR (400 MHz; CDCl_3): 2.37 (s, 6H); 3.79 (s, 2H); 4.00 (s, 3H); 5.34 (s, 2H); 7.00 (s, 1H); 7.15 (br s, 1H); 7.34-7.37 (m, 2H); 7.55 (d, 1H); 7.57 (d, 1H); 7.96 (d, 1H); 8.66 (s, 1H); MS (EI) for $\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{N}_5\text{O}_2\text{S}$: 490 (MH^+).

Example 28

N-(3,4-dichlorophenyl)-6-(methoxy)-7-([2-(morpholin-4-ylmethyl)-1,3-thiazol-4-yl]methyl)oxy)quinazolin-4-amine

- [0385] Morpholin-4-ylacetonitrile (1.03 g, 8.17 mmol) was added to a mixture of DMF (3.5 mL) and triethylamine (1.4 mL). Hydrogen sulfide gas was bubbled into the solution until the solution was light green. The solution was then heated at 70°C for 0.5 h. Excess hydrogen sulfide was removed from the system by bubbling nitrogen into the solution and then the volatiles were removed *in vacuo* to afford brown crystals. These crystals were triturated with ethanol to afford 2-morpholin-4-ylethanethioamide as very pale brown crystals (0.525 g, 3.28 mmol, 40% yield). ¹H NMR (400 MHz; CDCl₃): 2.53-2.58 (m, 4H); 3.44 (s, 2H); 3.70-3.74 (m, 4H); 7.74 (br s, 1H); 8.70 (br s, 1H); GCMS for C₆H₁₂N₂OS: 160 (M⁺).
- [0386] 2-Morpholin-4-ylethanethioamide (0.403 g, 2.52 mmol), 1,3-dichloroacetone (0.353 g, 2.77 mmol) and sodium bicarbonate (0.234 g, 2.77 mmol) was stirred in 1,2-dichloroethane (4 mL) at room temperature for 74 h at which point 1,4-dioxane was added in an attempt to improve solubility and the mixture was stirred for a further 30 h. The reaction mixture was filtered and the filter cake was washed with 1,2-dichloroethane. The filtrate was added dropwise to a cooled (ice bath) solution of thionyl chloride (0.202 mL, 2.77 mmol) in 1,2-dichloroethane (3 mL). The solution was stirred at 70°C for 0.5 h and then cooled to room temperature. The brown mixture was filtered and washed with 1,2-dichloroethane to afford 4-{[4-(chloromethyl)-1,3-thiazol-2-yl]methyl}morpholine hydrochloride as a very pale brown solid (0.024 g, 0.091 mmol, 4% yield). ¹H NMR (400 MHz; D₂O): 3.29-3.39 (m, 4H); 3.78-3.92 (m, 4H); 4.63 (s, 2H); 4.67 (s, 2H); 7.67 (s, 1H); MS (EI) for C₉H₁₃ClN₂OS: 233 (MH⁺).
- [0387] 4-{[4-(Chloromethyl)-1,3-thiazol-2-yl]methyl}morpholine hydrochloride (0.024 g, 0.089 mmol) and 4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (0.030 g, 0.089 mmol) were suspended in DMF (2 mL) and potassium carbonate (0.062 g, 0.449 mmol) was added. The mixture was stirred at 70°C for 21 h. The reaction mixture was concentrated *in vacuo* and the residue was purified by reverse phase HPLC to afford the title compound as a colorless solid (0.026 g, 0.049 mmol, 54% yield). ¹H NMR (400 MHz; CDCl₃): 2.50 (br s, 1H); 2.57-2.64 (m, 4H); 3.75 (t, 4H); 3.84 (s, 2H); 3.96 (s, 3H); 5.28 (s, 2H); 7.09 (s, 1H); 7.31 (s, 1H); 7.37 (s, 1H); 7.43 (d, 1H); 7.60 (dd, 1H); 7.97 (d, 1H); 8.61 (s, 1H); MS (EI) for C₂₄H₂₃Cl₂N₅O₃S: 532 (MH⁺).

Example 29

N-(3,4-dichlorophenyl)-6-(methoxy)-7-([2-[(4-methylpiperazin-1-yl)methyl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine

- [0388]** (4-Methylpiperazin-1-yl)acetonitrile (1.03 g, 7.42 mmol) was added to a mixture of DMF (5 mL) and triethylamine (2 mL). Hydrogen sulfide gas was bubbled into the solution until the solution was dark green. The solution was then heated at 70°C for 1 h. Excess hydrogen sulfide was removed from the system by bubbling nitrogen into the solution and then the volatiles were removed *in vacuo* to afford brown crystals. These crystals were triturated with ethanol to afford 2-(4-methylpiperazin-1-yl)ethanethioamide as very pale brown crystals (0.455 g, 2.63 mmol, 35% yield). ¹H NMR (400 MHz; CDCl₃): 2.29 (s, 3H); 2.38-2.66 (m, 8H); 3.44 (s, 2H); 7.72 (br s, 1H); 8.76 (br s, 1H); GCMS for C₇H₁₅N₃S: 173 (M⁺).
- [0389]** 2-(4-Methylpiperazin-1-yl)ethanethioamide (0.438 g, 2.53 mmol), 1,3-dichloroacetone (0.355 g, 2.78 mmol) and sodium bicarbonate (0.236 g, 2.78 mmol) was stirred in chloroform (4 mL) at room temperature for 48 h. The reaction mixture was filtered and the filter cake was washed with chloroform. The filtrate was added dropwise to a cooled (ice bath) solution of thionyl chloride (0.205 mL, 2.78 mmol) in chloroform (2 mL). The solution was stirred at 60°C for 0.5 h and then cooled to room temperature. The brown mixture was filtered and washed with chloroform to afford 1-{[4-(chloromethyl)-1,3-thiazol-2-yl]methyl}-4-methylpiperazine hydrochloride as a brown solid (0.421 g, 1.49 mmol, 59% yield). ¹H NMR (400 MHz; D₂O): 2.89 (s, 3H); 3.18-3.43 (m, 4H); 3.56-3.78 (m, 4H); 4.56 (s, 2H); 4.67 (s, 2H); 7.66 (s, 1H); MS (EI) for C₁₀H₁₆ClN₃S: 246 (MH⁺).
- [0390]** 1-{[4-(Chloromethyl)-1,3-thiazol-2-yl]methyl}-4-methylpiperazine hydrochloride (0.067 g, 0.238 mmol) and 4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (0.080 g, 0.238 mmol) were suspended in DMF (4 mL) and potassium carbonate (0.164 g, 1.19 mmol) was added. The mixture was stirred at 70°C for 21 h. The reaction mixture was concentrated *in vacuo* and the residue was purified by reverse phase HPLC to afford the title compound as a colorless solid (0.025 g, 0.046 mmol, 19% yield). ¹H NMR (400 MHz; CDCl₃): 2.08 (s, 5H); 2.39 (s, 3H); 2.59-2.74 (m, 6H); 3.44-3.78 (m, 5H); 3.86 (s, 2H); 3.98 (s, 3H); 5.30 (s, 2H); 7.10 (s, 1H); 7.33 (s, 1H); 7.37 (s, 1H); 7.43 (d, 1H); 7.61 (dd, 1H); 7.99 (d, 1H); 8.60 (s, 1H); MS (EI) for C₂₅H₂₆Cl₂N₆O₂S: 545 (MH⁺).

Example 30**N-(3,4-dichlorophenyl)-6-(methoxy)-7-([5-(1-methylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine (Method 1)**

- [0391]** Bromoacetonitrile (2.1 ml, 30.1 mmol) was added to a mixture of 4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (10.12 g; 30.1 mmol) and potassium carbonate (16.6g; 120.1 mmol) in DMF (200 ml). The mixture was stirred at room temperature overnight. The solvent was removed under high vacuum and the residue was triturated with water, filtered, washed with a mixture of hexane and ether (1/1) and dried under vacuum to give {[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}acetonitrile (9.6 g; 91% yield). ¹H NMR (400MHz; DMSO-d₆): 3.99 (s, 3H), 5.40 (s, 2H), 7.41 (s, 1H), 7.64 (d, 1H), 7.87 (dd, 1H), 7.90 (s, 1H), 8.23 (d, 1H), 8.57 (s, 1H), 9.69 (s, 1H); MS (EI) for C₁₇H₁₂Cl₂N₄O₂: 375.06 (MH⁺).
- [0392]** To a suspension of {[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}acetonitrile (14.7g; 39.3 mmol) in EtOH (600 ml) was added a 50% aqueous solution of NH₂OH (24.1 ml; 393 mmol) and the reaction mixture was refluxed for 2 hours. The solvent was evaporated off and the residue was triturated with ether, collected by filtration and dried under vacuum to give 2-{[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}-N-hydroxyethanimidamide (14.5 g; 90% yield). ¹H NMR (400MHz; DMSO-d₆): 3.98 (s, 3H), 4.58 (s, 2H), 5.70 (s, 2H), 7.38 (s, 1H), 7.65 (d, 1H), 7.84 (s, 1H), 7.89 (dd, 1H), 8.26 (d, 1H), 8.55 (s, 1H), 9.41 (s, 1H), 9.64 (s, 1H); MS (EI) for C₁₇H₁₅Cl₂N₅O₃: 408.05 (MH⁺).
- [0393]** To a solution of Boc-isonipecotic acid (4.72g; 20.6 mmol) and 4-methylmorpholine (5.7 ml; 51.5 mmol) in DMF (200ml) were added HOBt (3.06g; 22.7 mmol) and EDCI (4.35g; 22.7 mmol) and the solution was stirred at room temperature for 30 minutes. To the reaction mixture 2-{[4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy}-N-hydroxyethanimidamide (7g; 17.1 mmol) was added and the solution was stirred at room temperature overnight. The solvent was removed under vacuum and the residue was dissolved in ethyl acetate. The organic layer was washed with saturated aqueous NaHCO₃, brine, dried over NaSO₄ and concentrated.
- [0394]** Procedure a) To the residue was added p-xylene (200ml) and the suspension was refluxed for two hours. The solvent was removed under vacuum and the residue was purified by column chromatography (SiO₂, hexanes/acetone from 4/1 to 1/1) to give 1,1-

dimethylethyl 4-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate (8.3 g; 81% yield over two steps)

[0395] Procedure b) The residue was dissolved in THF (400ml) and *n*-Bu₄NF (1M in THF; 17.1ml; 17.1mmol) was added. The reaction mixture was stirred at room temperature for 2h. The solvent was evaporated off and the residue was dissolved in ethyl acetate; the organic layer was washed with H₂O, brine, dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by column chromatography (SiO₂, hexanes/acetone) to give 1,1-dimethylethyl 4-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate (6.9 g; 66% overall yield). ¹H NMR (400MHz; DMSO-*d*₆): 1.40 (s, 9H), 1.55-1.67 (m, 2H), 2.01-2.08 (m, 2H), 2.83-3.05 (br s, 2H), 3.33 (m, 1H), 3.88-3.96 (m, 2H), 3.99 (s, 3H), 5.53 (s, 2H), 7.43 (s, 1H), 7.73 (s, 2H), 8.02 (s, 1H), 8.1 (s, 1H), 8.81 (s, 1H), 10.74 (br s, 1H); MS (EI) for C₂₈H₃₀Cl₂N₆O₅: 601.09 (MH⁺)

[0396] 1,1-Dimethylethyl 4-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate (12.3g, 20.4 mmol) was dissolved in MeOH (100ml) and HCl (4M in dioxane; 65ml) was added. The solution was stirred at room temperature for 10 min and concentrated to half the initial volume. Ethyl ether was added to the suspension obtained and the precipitate was collected by filtration, washed with ether and concentrated to dryness to give 8g of N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(5-piperidin-4-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy}quinazolin-4-amine (78% yield). ¹H NMR (400MHz; DMSO-*d*₆): 1.61 (qd, 2H), 1.94 (m, 2H), 2.07-2.19 (br s, 1H), 2.57 (td, 2H), 2.95 (dt, 2H), 3.17 (m, 1H), 3.97 (s, 3H), 5.44 (s, 2H), 7.40 (s, 1H), 7.63 (d, 1H), 7.85 (s, 1H), 7.87 (dd, 1H), 8.23 (d, 1H), 8.54 (s, 1H), 9.64 (s, 1H); MS (EI) for C₂₃H₂₂Cl₂N₆O₃: 501.07 (MH⁺)

[0397] Aqueous formaldehyde (37%, 894 ml, 12.0 mmol) was added to a solution of N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(5-piperidin-4-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy}quinazolin-4-amine (1g, 2.0mmol) in formic acid (2ml). The mixture was heated at 95°C for two hours. The volatiles were removed under vacuum, the residue was dissolved in CH₃OH and the pH of the solution was adjusted to 9 by addition of Biorad AG[®] 1-X8 hydroxide form resin. The solution was filtered, concentrated *in vacuo* and the residue was crystallized from MeOH/H₂O to give N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(5-(1-methylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy}quinazolin-4-amine

(714 mg; 69% yield). ^1H NMR (400MHz; DMSO- d_6): 1.69-1.82 (m, 2H), 1.97-2.05 (m, 4H), 2.16 (s, 3H), 2.71-2.78 (m, 2H), 2.98-3.10 (m, 1H), 3.96 (s, 3H), 5.44 (s, 2H), 7.39 (s, 1H), 7.62 (d, 1H), 7.83 (s, 1H), 7.87 (dd, 1H), 8.22 (d, 1H), 8.53 (s, 1H), 9.63 (s, 1H); MS (EI) for $\text{C}_{24}\text{H}_{24}\text{Cl}_2\text{N}_6\text{O}_3$: 515.07 (MH^+)

[0398] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0399] N-(3,4-dichlorophenyl)-7-[(5-[(dimethylamino)methyl]-1,2,4-oxadiazol-3-yl)methyl]oxy-6-(methyloxy)quinazolin-4-amine: ^1H NMR (400MHz; DMSO- d_6): 2.26 (s, 6H), 3.87 (s, 2H), 3.97 (s, 3H), 5.49 (s, 2H), 7.39 (s, 1H), 7.85-7.90 (m, 2H), 8.23 (d, 1H), 8.53 (s, 1H), 9.67 (s, 1H); MS (EI) for $\text{C}_{21}\text{H}_{20}\text{Cl}_2\text{N}_6\text{O}_3$: 475.06 (MH^+).

[0400] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-piperidin-3-yl-1,2,4-oxadiazol-3-yl)methyl]oxy}quinazolin-4-amine: ^1H NMR (400MHz; DMSO- d_6): 1.37-1.82 (m, 4H), 2.04-2.14 (m, 1H), 2.20-2.34 (m, 1H), 2.70-2.87 (m, 2H), 3.07-3.22 (m, 2H), 3.97 (s, 3H), 5.44 (s, 2H), 7.40 (s, 1H), 7.63 (d, 1H), 7.85 (s, 1H), 7.87 (dd, 1H), 8.23 (d, 1H), 8.54 (s, 1H), 9.64 (s, 1H); MS (EI) for $\text{C}_{23}\text{H}_{22}\text{Cl}_2\text{N}_6\text{O}_3$: 501.03 (MH^+).

[0401] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(1-methylpiperidin-3-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine: ^1H NMR (400MHz; DMSO- d_6): 1.51-1.76 (m, 4H), 1.92-2.12 (m, 2H), 2.18 (s, 3H), 2.30-2.40 (m, 1H), 2.51-2.58 (m, 1H), 2.87-2.94 (m, 1H) 3.97 (s, 3H), 5.44 (s, 2H), 7.40 (s, 1H), 7.63 (d, 1H), 7.83-7.90 (m, 2H), 8.23 (d, 1H), 8.54 (s, 1H), 9.64 (s, 1H); MS (EI) for $\text{C}_{24}\text{H}_{24}\text{Cl}_2\text{N}_6\text{O}_3$: 515.06 (MH^+).

[0402] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(piperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine: ^1H NMR (400MHz; DMSO- d_6): 1.26-1.50 (m, 4H), 1.52-1.72 (m, 2H), 1.82-1.90 (m, 1H), 2.50-2.64 (m, 1H), 2.80-2.90 (m, 1H), 3.91 (s, 3H), 5.41 (s, 2H), 7.35 (s, 1H), 7.58 (d, 1H), 7.80-7.86 (m, 2H), 8.19 (d, 1H), 8.49 (s, 1H), 9.60 (s, 1H); MS (EI) for $\text{C}_{23}\text{H}_{22}\text{Cl}_2\text{N}_6\text{O}_3$: 501.11 (MH^+).

[0403] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(1-methylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine hydrochloride: ^1H NMR (400MHz; DMSO- d_6): 1.61 (br m, 1H), 1.75-1.88 (br m, 3H), 2.08 (br m, 1H), 2.28 (br d, 1H), 2.75 (s, 3H), 3.24 (br m, 1H), 3.49 (br m, 2H), 4.05 (s, 3H), 5.08 (br s, 1H), 5.64 (s, 2H), 7.62 (s, 1H), 7.71 (d, 1H), 7.87 (dd, 1H), 8.18 (d, 1H), 8.65 (s, 1H), 8.88 (s, 1H), 11.95 (s, 1H); MS (EI) for $\text{C}_{24}\text{H}_{24}\text{Cl}_2\text{N}_6\text{O}_3$: 515.11 (MH^+).

[0404] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(1-ethylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine hydrochloride: ^1H NMR (400MHz;

DMSO- d_6): 11.80 (br s, 1H), 8.87 (s, 1H), 8.56 (s, 1H), 8.18 (d, 1H), 7.85 (dd, 1H), 7.73 (d, 1H), 7.60 (s, 1H), 5.65 (s, 2H), 5.10 (br s, 1H), 4.05 (s, 3H), 3.80-3.50 (br m, 4H), 3.20-3.00 (br m, 2H), 2.30-2.00 (br m, 1H), 1.95 (br m, 2H), 1.65 (br m, 1H), 1.20 (br m, 3H); MS (EI) for $C_{25}H_{26}Cl_2N_6O_3$: 529 (MH^+).

Example 31

N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(5-piperidin-2-yl-1,2,4-oxadiazol-3-yl)methyl]oxy}quinazolin-4-amine hydrochloride (Method 2)

[0405] To a suspension of 50.0 g (0.41 mol) benzoic acid in 150 mL N,N-dimethylformamide 70.0g (0.50 mol) potassium carbonate was added and the mixture was stirred until the evolution of gas has ceased. The reaction mixture was cooled to 0°C and a solution of 29.0 mL (0.41 mol) bromoacetonitrile in 50 mL N,N-dimethylformamide was added dropwise. The reaction mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was dissolved in 500 mL ethyl acetate. It was washed with water (3x150 mL), saturated aqueous sodium hydrogencarbonate (150 mL) and brine. The organic layer was separated and dried over anhydrous sodium sulfate. The drying agent was filtered off and the solvent was evaporated. The resulting crude product was purified by column chromatography (hexane-ethyl acetate 9:1 to 4:1) to yield cyanomethyl benzoate 62.6 g (95%). ^1H -NMR (400MHz; DMSO- d_6): 7.98 (m, 2H), 7.69 (m, 1H), 7.54 (m, 2H), 5.20 (s, 2H); GCMS for $C_9H_7NO_2$: 161 (M^+).

[0406] Cyanomethyl benzoate (22.2g, 138 mmol) was dissolved in ethanol (250mL) followed by addition of 50% aqueous hydroxylamine (16.9mL, 276mmol) and the mixture was stirred for 15 minutes at room temperature. Concentration followed by addition of water (250mL) afforded a colorless crystalline solid that was collected by filtration and washed with additional water then dried *in vacuo* to give 2-(hydroxyamino)-2-iminoethyl benzoate (23.69g, 88% yield). ^1H -NMR (400MHz; DMSO- d_6): 9.37 (s, 1H), 7.98 (m, 2H), 7.65 (m, 1H), 7.54 (tr, 2H), 5.69 (br s, 2H), 4.64 (s, 2H).

Example 32

N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(3-[(4-methylpiperazin-1-yl)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy}quinazolin-4-amine

[0407] A mixture of 4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-ol (3.67g), methyl bromoacetate (0.81 mL), and potassium carbonate (4.66g) in DMF (21mL) was

stirred at room temperature for 18 hours. The reaction mixture was diluted with ethyl acetate and filtered. The solid was combined with the filtrate and washed with water, dried over sodium sulfate and evaporated to give a brown solid which was recrystallized in methanol to give methyl {[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}acetate (2.80g). ¹H-NMR (400MHz; DMSO-d₆): 9.62 (s, 1H), 8.46 (s, 1H), 8.24 (s, 1H), 7.94-7.85 (m, 2H), 7.63 (d, 1H), 7.17 (s, 1H), 5.04 (s, 2H), 4.00 (s, 3H), 3.72 (s, 3H); MS (EI) for C₁₈H₁₅N₃O₄Cl₂: 408 (MH⁺).

[0408] A mixture (4-methylpiperazin-1-yl)acetonitrile (400 mg), hydroxylamine (50% aqueous solution, 2.0mL) and ethanol (10.0 mL), was stirred at room temperature for 18 hours. The reaction mixture was evaporated to dryness to give a crystalline solid that was washed with hexane and dried under vacuum to give N-hydroxy-2-(4-methylpiperazin-1-yl)ethanimidamide (405 mg). ¹H-NMR (400MHz; CDCl₃): 5.19 (br s, 3H), 3.00 (s, 2H), 2.75-2.28 (br m, 8H), 2.70 (s, 3H).

[0409] Sodium hydride (60% oil dispersion, 30 mg) was added to a mixture of N-hydroxy-2-(4-methylpiperazin-1-yl)ethanimidamide (50mg) and 3A molecular sieves (110 mg) in 1 mL of DMF. After the initial hydrogen evolution had stopped, the mixture was heated at 60°C for 30 minutes. {[4-[(3,4-Dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}acetate (50mg) was added to the reaction mixture and the heating was continued at 90°C for one hour. The reaction was cooled to room temperature, diluted with methanol, and filtered. The filtrate was purified by reverse phase HPLC, and the eluent containing the expected product was concentrated to dryness and isolated as the TFA salt. The salt was taken into methanol and treated with Biorad AG[®] 1-X8 hydroxide form resin then filtered and the filtrate concentrated and dried *in vacuo* to give the title compound (25.0 mg) ¹H-NMR (400MHz, DMSO-d₆): 10.60 (br s, 1H), 8.78 (s, 1H), 8.17 (s, 1H), 8.02 (s, 1H), 7.79 (d, 1H), 7.70 (d, 1H), 7.22 (s, 1H), 5.78 (s, 2H), 4.00 (s, 3H), 3.82 (s, 2H), 3.15-2.90 (br m, 4H), 2.80 (s, 3H), 2.60-2.45 (br m, 4H); MS (EI) for C₂₄H₂₅N₇O₃Cl₂: 530 (MH⁺).

[0410] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0411] N-(3,4-dichlorophenyl)-7-[(3-[(dimethylamino)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine: ¹H-NMR (400MHz; DMSO-d₆): 10.48 (br s, 1H), 8.75 (s, 1H), 8.20 (s, 1H), 8.05 (s, 1H), 7.81 (d, 1H), 7.72 (d, 1H), 7.25

(s, 1H), 5.85 (s, 2H), 4.85 (s, 2H), 4.00 (s, 3H), 2.88 (s, 6H); MS (EI) for $C_{21}H_{20}N_6O_3Cl_2$: 475 (MH^+).

[0412] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([3-(morpholin-4-ylmethyl)-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine: ^1H -NMR (400MHz, DMSO- d_6): 10.50 (br s, 1H), 8.78 (s, 1H), 8.18 (s, 1H), 8.04 (s, 1H), 7.79 (d, 1H), 7.70 (d, 1H), 7.22 (s, 1H), 5.82 (s, 2H), 4.37 (s, 2H), 4.00 (s, 3H), 3.78-3.70 (br m, 4H), 2.15-2.95 (br m, 4H); MS (EI) for $C_{24}H_{25}N_7O_3Cl_2$: 517 (MH^+).

Example 33

N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([3-(4-methylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine

[0413] 1,1-Dimethylethyl 1-piperazinecarboxylate (25g, 134.2 mmol) was taken into THF (100mL) followed by addition of triethylamine (25mL, 178 mmol). The solution was cooled to 0°C followed by addition dropwise addition of cyanogen bromide (15.6g, 147.6 mmol) in THF (100mL) and the resulting mixture was allowed to warm to room temperature then stirred an additional 12 hours. The reaction mixture was concentrated *in vacuo* and partitioned with ethyl ether and water. The organic layer was washed once with saturated aqueous NaCl then dried over anhydrous magnesium sulfate followed by concentration to afford 1,1-dimethylethyl 4-cyanopiperazine-1-carboxylate (24.5g, 86% yield) as a colorless crystalline solid. ^1H -NMR (400MHz, $CDCl_3$): 3.51 (tr, 4H), 3.19 (tr, 4H), 1.46 (s, 9H).

[0414] 1,1-Dimethylethyl 4-cyanopiperazine-1-carboxylate (10g, 47.1 mmol) was taken into ethanol (100mL) and the resulting solution was cooled to 0°C. Hydroxylamine (50% aqueous solution, 3.5mL, 56.5 mmol) was added and the mixture was allowed to warm to room temperature then stirred an additional hour. The mixture was then concentrated *in vacuo* to a paste and suspended in 1:1 ethyl ether / hexane (100mL) and the solid product collected by filtration. The solid was washed with an additional portion of ethyl ether / hexane then dried to give 1,1-dimethylethyl 4-[(hydroxyamino)(imino)methyl]piperazine-1-carboxylate (9.64g, 83% yield) as a white solid. ^1H -NMR (400MHz, DMSO- d_6): 8.34 (s, 1H), 5.19 (s, 2H), 3.30 (tr, 4H), 2.91 (tr, 4H), 1.39 (s, 9H).

[0415] 1,1-Dimethylethyl 4-[(hydroxyamino)(imino)methyl]piperazine-1-carboxylate (3.58g, 14.6 mmol) was suspended in THF (50mL) followed by the addition of diisopropylethylamine (3.0mL, 17.5 mmol) and the mixture was cooled to 0°C.

Acetoxyacetyl chloride (1.6mL, 14.6 mmol) was added by syringe and the mixture was allowed to stir an additional 30 minutes at 0°C. The resulting homogeneous solution was allowed to warm to room temperature then concentrated *in vacuo* to afford a white solid residue. The material was suspended in water (50ml) and the solid product was collected by filtration, washed with additional water then hexanes and dried *in vacuo*. The intermediate O-acyl derivative was then suspended in THF (50mL) followed by addition of TBAF (1.0M in THF, 3.5mL) and the mixture was stirred for 30 minutes at room temperature. The homogeneous solution obtained was concentrated *in vacuo* and the residue partitioned with ethyl ether and water. The organic layer was washed twice with additional water then with saturated aqueous sodium chloride and dried over anhydrous magnesium sulfate. Filtration and concentration gave 1,1-dimethylethyl 4-{5-[(acetoxymethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (3.6g, 76% yield) as a colorless crystalline solid. ¹H-NMR (400MHz, CDCl₃): 5.14 (s, 2H), 3.51 (tr, 4H), 3.42 (tr, 4H), 2.18 (s, 3H), 1.48 (s, 9H).

[0416] 1,1-Dimethylethyl 4-{5-[(acetoxymethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (13.57g, 41.6 mmol) was dissolved in methanol (100mL) followed by addition of aqueous sodium hydroxide (4M, 10.4mL) and the mixture was stirred for 30 minutes at room temperature. The reaction mixture was then concentrated *in vacuo* and the residue partitioned with ethyl acetate and 0.1M aqueous hydrochloric acid. The organic layer was then washed with saturated aqueous sodium chloride then dried over anhydrous magnesium sulfate. Filtration and concentration afforded 1,1-dimethylethyl 4-[5-(hydroxymethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (10.51g, 89% yield) as a colorless crystalline solid. ¹H-NMR (400MHz, CDCl₃): 4.75 (s, 2H), 3.53-3.50 (m, 4H), 3.44-3.42 (m, 4H), 3.20 (br s, 1H), 1.48 (s, 9H).

[0417] 1,1-Dimethylethyl 4-[5-(hydroxymethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (7.4g, 26 mmol) was taken into dichloromethane (100mL) followed by addition of pyridine (5.3mL, 65 mmol) and the solution was cooled to 0°C. Thionyl chloride (2.3mL, 31.2 mmol) was added by syringe and the mixture was allowed to warm to room temperature and stirred an additional 12 hours. The mixture was then concentrated *in vacuo* and the residue partitioned with ethyl acetate and water. The aqueous phase was extracted twice with ethyl acetate and the combined organic layers were washed once with 0.1M aqueous hydrochloric acid then saturated aqueous sodium chloride and dried over anhydrous magnesium sulfate. Filtration, concentration and purification of the residue by silica gel flash chromatography using 3:1 hexanes:ethyl

acetate provided 1,1-dimethylethyl 4-[5-(chloromethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (4.82g, 61% yield) as a slightly yellow crystalline solid. ¹H-NMR (400MHz, CDCl₃): 4.53 (s, 2H), 3.51 (tr, 4H), 3.43 (tr, 4H), 1.48 (s, 9H).

[0418] 1,1-dimethylethyl 4-[5-(chloromethyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (173.6mg, 0.57 mmol), 4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-ol (154mg, 0.46 mmol) and potassium carbonate (315mg, 2.85 mmol) were taken into DMF (2mL) and the mixture was heated to 50°C for one hour. The mixture was then cooled to room temperature and partitioned with ethyl acetate and water. The organic layer was washed once more with water and the organic layer was partitioned again with water. A precipitate forms in the organic layer at this point and hexanes was added to the biphasic mixture in portions until formation of the precipitate is complete. The solid product was then collected by filtration. The solid residue was suspended in hot methanol and diluted with water followed by collection of the solid by filtration. The solid was washed with ethyl ether and dried *in vacuo* to give 1,1-dimethylethyl 4-[5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (170.7mg, 62% yield) as a tan solid. ¹H-NMR (400MHz, DMSO-d₆): 9.65 (s, 1H), 8.54 (s, 1H), 8.23 (d, 1H), 7.88-7.86 (m, 2H), 7.63 (d, 1H), 7.32 (s, 1H), 5.58 (s, 2H), 3.98 (s, 3H), 3.43-3.40 (m, 4H), 3.34-3.32 (m, 4H), 1.41 (s, 9H); MS (EI) for C₂₇H₂₉Cl₂N₇O₅: 602 (MH⁺).

[0419] 1,1-dimethylethyl 4-[5-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate (170mg, 0.28 mmol) was taken into methanol (2.5mL) followed by addition of anhydrous hydrogen chloride in dioxane (4M, 2.5mL) and the mixture was brought to reflux then immediately allowed to cool to room temperature over 5 minutes. The mixture was then concentrated *in vacuo* to a slurry followed by addition of excess ethyl ether. The solid residue was collected by filtration and dried *in vacuo* to give N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(3-piperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl}oxy}quinazolin-4-amine hydrochloride (145mg, 100% yield) as a tan solid. ¹H-NMR (400MHz, DMSO-d₆): (hydrochloride) 11.98 (s, 1H), 9.54 (br s, 2 H), 8.87 (s, 1H), 8.66 (s, 1H), 8.17 (d, 1H), 7.86 (dd, 1H), 7.70 (d, 1H), 7.47 (s, 1H), 5.66 (s, 2H), 4.06 (s, 3H), 3.64-3.61 (m, 4H), 3.19-3.16 (m, 4H); (free base) 9.69 (s, 1H), 8.53 (s, 1H), 8.24 (d, 1H), 7.90-7.87 (m, 2H), 7.62 (d, 1H), 7.31 (s, 1H), 5.56 (s, 2H), 3.99 (s, 3H), 3.31-3.29 (m, 4H), 2.81-2.79 (m, 4H); MS (EI) for C₂₂H₂₁Cl₂N₇O₃: 502 (MH⁺).

[0420] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{[(3-piperazin-1-yl)-1,2,4-oxadiazol-5-yl)methyl}oxyquinazolin-4-amine hydrochloride (66mg, 0.12 mmol) was suspended in methanol (5mL) followed by addition of Bio-Rad AG[®]1-X8 hydroxide form resin until pH 8. The resin was then removed by filtration and the methanol solution was concentrated *in vacuo*. The residue was taken into formic acid (2mL) and 37W% aqueous formaldehyde (50uL, 0.6 mmol) was added. The mixture was heated to 90°C for 2.5 hours then concentrated *in vacuo*. The residue was taken into 1mL of methanol and the solution acidified by addition of 4M anhydrous hydrogen chloride in dioxane to pH 2. Portionwise addition of ethyl ether afforded a crystalline solid that was collected by filtration and dried to give N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-(4-methylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine hydrochloride (53.9mg, 80% yield) as a tan solid. ¹H-NMR (400MHz, DMSO-d₆): 11.98 (s, 1H), 11.39 (s, 1H), 8.87 (s, 1H), 8.65 (s, 1H), 8.17 (d, 1H), 7.86 (dd, 1H), 7.69 (d, 1H), 7.48 (s, 1H), 5.67 (s, 2H), 4.06 (s, 3H), 3.95 (br d, 2H), 3.49-3.46 (m, 4H), 3.15 (br m, 2H), 2.78 (s, 3H); MS (EI) for C₂₃H₂₃Cl₂N₇O₃: 516 (MH⁺).

[0421] Using the same or analogous synthetic techniques and/or substituting with alternative reagents, the following compounds of the invention were prepared:

[0422] N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-(4-ethylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine hydrochloride: ¹H-NMR (400MHz, DMSO-d₆):

Example 34

N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{[(5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)methyl}oxy}quinazolin-4-amine hydrochloride

[0423] To a solution of {[4-[3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}acetonitrile (0.450 g, 1.20 mmol) in DMF (2 mL) was added NEt₃ (0.30 mL, 2.2 mmol). Hydrogen sulfide (g) was bubbled through the solution until saturation (5 min.). The solution was then heated to 70°C and it turned dark green. After 0.5 h, the solution was cooled to room temperature, 0.20 mL NEt₃ (1.4 mmol) was added, and H₂S (g) was bubbled through for 5 min. The solution was again heated to 70°C for 1.5 h. The solvent was removed *in vacuo* and the product was crystallized from MeOH to yield 0.459 g (94%) of 2-{{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethanethioamide as a yellow solid. ¹H NMR (400 MHz, d₆-DMSO): 10.07 (s,

1H), 9.68 (s, 1H), 9.29 (s, 1H), 8.55 (s, 1H), 8.26 (d, 1H), 7.91 (m, 2H), 7.66 (d, 1H), 7.10 (s, 1H), 4.96 (s, 1H), 4.01 (s, 3H), 2.73 (d, 2H); MS (EI) for $C_{17}H_{14}N_4O_2SCl_2$: 409 (M^+).

[0424] To a solution of 2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethanethioamide (0.300 g, 0.733 mmol) in DMF (5 mL) was added 1,1-dimethylethyl 3-bromo-4-oxopiperidine-1-carboxylate (0.310 g, 1.11 mmol). The solution was stirred at 60°C for 72 h. An additional 0.100 g (0.360 mmol) of 1,1-dimethylethyl 3-bromo-4-oxopiperidine-1-carboxylate was added to the reaction mixture, which was then heated at 60°C for an additional 24 h. The solvent was removed *in vacuo* and the crude reaction mixture was purified via column chromatography (SiO_2 , 50% hexanes/ethyl acetate). This intermediate was then subjected to 10% TFA in CH_2Cl_2 at room temperature. After removal of the solvent, the crude amine salt was taken up in fresh MeOH and treated with Bio-Rad AG 1-X8 resin hydroxide form until pH 8. Filtration and concentration *in vacuo* provided 0.247 g (69%) of *N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-ylmethyl)oxy]quinazolin-4-amine as a brown oil.

[0425] ^1H NMR (400 MHz, d_6 -DMSO): 9.68 (s, 1H), 8.43 (s, 1H), 8.24 (d, 1H), 7.89 (d, 1H), 7.87 (d, 1H), 7.62 (d, 1H), 7.37 (s, 1H), 5.54 (s, 2H), 3.98 (s, 3H), 3.87 (s, 2H), 2.97 (t, 2H), 2.66 (t, 2H); MS (EI) for $C_{22}H_{19}N_5O_2SCl_2$: 488 (M^+)

[0426] *N*-(3,4-Dichlorophenyl)-6-(methyloxy)-7-[(4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-ylmethyl)oxy]quinazolin-4-amine (0.247 g, 0.506 mmol) was combined with 37% aqueous formaldehyde (0.23 mL, 3.1 mmol) in formic acid (2 mL) and the solution was heated at 95°C for 1 h. The solvent was removed *in vacuo* and the residue was purified via HPLC (reverse-phase, CH_3CN/H_2O with 0.1% TFA). Upon removal of CH_3CN/H_2O , the product was taken up in MeOH and treated with Bio-Rad AG 1-X8 resin hydroxide form until pH 8. The product was filtered and concentrated *in vacuo*, then taken up in fresh MeOH and treated with 4.0 M HCl/dioxane (0.050 mL). Removal of solvent *in vacuo* provided 0.085 g (31%) of *N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-yl)methyl]oxy]quinazolin-4-amine hydrochloride as a pale yellow solid. ^1H NMR (400 MHz, d_6 -DMSO): 8.69 (s, 1H), 8.24 (d, 1H), 8.11 (s, 1H), 7.89 (dd, 1H), 7.70 (d, 1H), 7.42 (s, 1H), 5.65 (s, 3H), 4.65 (m, 1H), 4.40 (m, 1H), 4.02 (s, 3H), 3.70 (m, 1H), 3.12 (m, 2H), 2.94 (s, 4H); MS (EI) for $C_{23}H_{21}N_5O_2SCl_2$: 502 (M^+).

Example 35

***N*-(3,4-dichlorophenyl)-7-[(5-ethyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0427] *N*-(3,4-Dichlorophenyl)-6-(methyloxy)-7-[(4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-ylmethyl)oxy]quinazolin-4-amine (0.249 g, 0.510 mmol) was taken up in 50% THF/MeOH (10 mL) and the solution was cooled with an ice bath. Acetaldehyde (0.057 mL, 1.0 mmol), followed by NaCNBH₃ (0.038 g, 0.61 mmol) were added. The solution was allowed to warm to room temperature. After 4 h, additional acetaldehyde (0.050 mL, 0.89 mmol) and NaCNBH₃ (0.040 g, 0.64 mmol) were added and the solution was stirred for 12 h. An additional 0.050 mL (0.89 mmol) of acetaldehyde and NaCNBH₃ (0.020 g, 0.32 mmol) were added. The solution was poured into 10% MeOH/ethyl acetate (100 mL) and washed once with H₂O (100 mL). The organic layer was dried (Na₂SO₄); filtered and the solvent was removed *in vacuo*. The crude product was purified via preparative HPLC (reverse phase, CH₃CN/H₂O/NH₄OAc/AcOH). The solvents were removed, the product was taken up in MeOH and treated with 4.0M HCl/dioxane (0.025 mL), and lyophilized to provide 0.036 g (12%) of *N*-(3,4-dichlorophenyl)-7-[(5-ethyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-*c*]pyridin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine hydrochloride as a yellow solid.

[0428] ¹H NMR (400 MHz, d₆-DMSO): 9.56 (broad s, 1H), 8.85 (s, 1H), 8.37 (broad s, 1H), 8.15 (d, 1H), 7.81 (dd, 1H), 7.73 (d, 1H), 7.48 (s, 1H), 5.66 (s, 2H), 4.71 (d, 1H), 4.36 (m, 1H), 4.04 (s, 3H), 3.74 (m, 1H), 3.13 (m, 3H), 1.32 (t, 3H); MS (EI) for C₂₄H₂₃N₅O₂SCl₂: 516 (M⁺).

Example 36

***N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-(1-methylpiperidin-4-yl)-1,3-thiazol-4-yl)methyl]oxy)quinazolin-4-amine hydrochloride**

[0429] 1,1-Dimethylethyl 4-(aminocarbonothioyl)piperidine-1-carboxylate (1.50 g, 6.14 mmol), NaHCO₃ (0.570 g, 6.78 mmol) and 1,3-dichloroacetone (0.860 g, 6.77 mmol) were combined in 1,2-dichloroethane (4 mL) and the reaction mixture was stirred at room temperature for 12 h. The crude reaction mixture was filtered using CH₂Cl₂ and the filtrate was concentrated *in vacuo* until approximately 30 mL of solvent remained. To this solution was added pyridine (0.75 mL, 9.2 mmol), and the solution was cooled with an ice bath. Thionyl chloride (0.49 mL, 6.8 mmol) was added and the solution was allowed to warm slowly to room temperature. The solvent was removed *in vacuo* and the residue was taken up in 10% MeOH/ethyl acetate (100 mL). The organic layer was

washed with H₂O (100 mL) and brine (100 mL), dried (Na₂SO₄), filtered, and concentrated *in vacuo* to yield 2.24 g (>100%) of crude 1,1-dimethylethyl 4-[4-(chloromethyl)-1,3-thiazol-2-yl]piperidine-1-carboxylate as a colorless oil. ¹H NMR (400 MHz, d₆-DMSO): 7.62 (s, 1H), 4.60 (s, 2H), 3.98 (m, 1H), 3.63 (dd, 2H), 3.16 (m, 1H), 2.90 (broad s, 2H), 2.01 (dd, 1H), 1.51 (m, 2H), 1.40 (s, 9H). MS (EI) for C₁₄H₂₁N₂O₂SCl: 261 (M - *t*Bu).

[0430] To a solution of 4-[3,4-dichlorophenyl]amino]-6-(methyloxy)quinazolin-7-ol (1.00 g, 2.97 mmol) in DMF (10 mL) was added 1,1-dimethylethyl 4-[4-(chloromethyl)-1,3-thiazol-2-yl]piperidine-1-carboxylate (1.04 g, 3.28 mmol) and K₂CO₃ (2.06 g, 14.9 mmol). The solution was heated to 70°C for 12 h, and then the solvent was removed *in vacuo*. The residue was taken up in 10% MeOH/ethyl acetate (100 mL) and washed with H₂O (100 mL). The organic layer was dried (Na₂SO₄), filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 50% hexanes/acetone) yielded 0.642 g (35%) of crude 1,1-dimethylethyl 4-[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]piperidine-1-carboxylate. Half of this intermediate (0.300 g) was then subjected to 10% TFA in CH₂Cl₂ at room temperature for 1 h, then concentrated *in vacuo*. The residue was dissolved in MeOH (50 mL) and treated with Bio-Rad AG 1-X8 resin hydroxide form to pH 8. Filtration and concentration *in vacuo* provided 0.230 g (92% from 1,1-dimethylethyl 4-[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]piperidine-1-carboxylate) of *N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(2-piperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy}quinazolin-4-amine as a brown oil. MS (EI) for C₂₄H₂₃N₅O₂SCl₂: 516 (M⁺).

[0431] *N*-(3,4-Dichlorophenyl)-6-(methyloxy)-7-{{[(2-piperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy}quinazolin-4-amine (0.078 g, 0.15 mmol) was combined with 37% aqueous formaldehyde (0.025 mL, 0.34 mmol) in formic acid (2 mL) and the solution was heated to 95°C for 1 h. The solvent was removed *in vacuo* and the residue was taken up in 10% MeOH/ethyl acetate (100 mL) and washed with saturated NaHCO₃ (aq) (100 mL). The organic layer was dried (Na₂SO₄), filtered, and concentrated *in vacuo*. Trituration from MeOH and treatment of the resultant yellow solid with 4.0 M HCl/dioxane (0.050 mL) in MeOH, followed by lyophilization, provided 0.032 g (37%) of *N*-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(1-methylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine hydrochloride as a pale yellow solid. ¹H NMR (400 MHz, d₆-DMSO): 8.55 (s, 1H), 8.25 (d, 1H), 7.89 (dd, 2H), 7.76 (s, 1H), 7.63 (d, 1H),

7.41 (s, 1H), 5.30 (s, 2H), 3.96 (s, 3H), 3.52 (d, 2H), 3.29 (m, 2H), 3.08 (m, 1H), 2.79 (s, 3H), 2.29 (d, 2H), 1.95 (m, 2H); MS (EI) for $C_{26}H_{25}N_5O_2SCl_2$: 530 (M^+).

Example 37

***N*-(3,4-dichlorophenyl)-7-([2-(1-ethylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride**

[0432] *N*-(3,4-Dichlorophenyl)-6-(methyloxy)-7-[(2-piperidin-4-yl-1,3-thiazol-4-yl)methyl]oxy}quinazolin-4-amine (0.230 g, 0.445 mmol) was taken up in 50% MeOH/THF (10 mL) and cooled with an ice bath. Acetaldehyde (0.050 mL, 0.89 mmol) was added, followed by NaCNBH₃ (0.034 g, 0.54 mmol) and the solution was allowed to warm to room temperature. After 1.5 h, 0.016 g (0.25 mmol) of additional NaCNBH₃ was added and the solution was stirred for 12 h. The reaction mixture was poured into ethyl acetate (100 mL), washed with saturated NaHCO₃ (aq) (50 mL) and dried (Na₂SO₄). Filtration and concentration *in vacuo* was followed by preparative HPLC (reverse-phase, CH₃CN/H₂O with 0.01% TFA). This product was neutralized (NaHCO₃) to give 0.167 g (0.306 mmol) of free amine, which was taken up in MeOH (50 mL) and treated with 4.0 M HCl/dioxane (0.077 mL, 0.31 mmol). Lyophilization provided 0.105 g (41%) of *N*-(3,4-dichlorophenyl)-7-([2-(1-ethylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine hydrochloride as a pale yellow solid. ¹H NMR (400 MHz, d₆-DMSO): 8.65 (broad s, 1H), 8.23 (d, 1H), 8.08 (broad s, 1H), 7.88 (d, 1H), 7.78 (s, 1H), 7.66 (d, 1H), 7.47 (s, 1H), 5.32 (s, 2H), 3.98 (s, 3H), 3.55 (d, 2H), 3.01-3.11 (m, 5H), 2.29 (m, 2H), 2.04 (m, 2H), 1.26 (t, 3H); MS (EI) for $C_{26}H_{27}N_5O_2SCl_2$: 544 (M^+).

Assays

[0433] Generally for assay of activity, either ephrin, EGFR, or a compound according to the invention is non-diffusably bound to an insoluble support having isolated sample-receiving areas (for example, a microtiter plate, an array, etc.). The insoluble support may be made of any composition to which the compositions can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (for example, polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out

simultaneously, using small amounts of reagents and samples. The particular manner of binding of the composition is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the composition and is nondiffusible. Exemplary methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to “sticky” or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the protein or agent, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine serum albumin (BSA), casein or other innocuous protein or other moiety.

[0434] One measure of inhibition is K_i . For compounds with IC_{50} 's less than 1 μ M, the K_i or K_d is defined as the dissociation rate constant for the interaction of the agent with ephrin or EGFR. Exemplary compositions have K_i 's of, for example, less than about 100 μ M, less than about 10 μ M, less than about 1 μ M, and further for example having K_i 's of less than about 100 nM, and still further, for example, less than about 10 nM. The K_i for a compound is determined from the IC_{50} based on three assumptions. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[I - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4E_0 I_0}}{2E_0} \right]$$

Where V is the observed rate, $V_{\max}$, is the rate of the free enzyme, I_0 is the inhibitor concentration, E_0 is the enzyme concentration, and K_d is the dissociation constant of the enzyme-inhibitor complex.

[0435] Another measure of inhibition is GI_{50} , defined as the concentration of the compound that results in a decrease in the rate of cell growth by fifty percent. Exemplary compounds have GI_{50} 's of, for example, less than about 1 mM, less than about 10 μ M, less than about 1 μ M, and further, for example, having GI_{50} 's of less than about 100 nM, still further having GI_{50} 's of less than about 10 nM. Measurement of GI_{50} is done using a cell proliferation assay.

[0436] Tyrosine kinase activity is determined by 1) measurement of kinase-dependent ATP consumption by in the presence of a generic substrate such as polyglutamine, tyrosine (pEY), by luciferase/luciferin-mediated chemiluminescence or; 2) incorporation of radioactive phosphate derived from ^{33}P -ATP into a generic substrate which has been adsorbed onto the well surface of polystyrene microtiter plates. Phosphorylated substrate products are quantified by scintillation spectrometry.

Materials and Methods

[0437] Kinase activity and compound inhibition are investigated using one or more of the three assay formats described below. A brief summary of assay conditions is listed in Table 2. The ATP concentrations are selected near its Michaelis-Menten constant (K_M) for each individual kinase. Dose-response experiments are performed at ten different inhibitor concentrations in a 384-well plate format. The data are fitted to a standard four-parameter equation listed below:

$$Y = \text{Min} + (\text{Max} - \text{Min}) / (1 + (X/\text{IC}_{50})^H)$$

where Y is the observed signal, X is the inhibitor concentration, Min is the background signal in the absence of enzyme (0% enzyme activity), Max is the signal in the absence of inhibitor (100% enzyme activity), IC_{50} is the inhibitor concentration at 50% enzyme inhibition and H represents the empirical Hill's slope to measure the cooperativity. Typically H is close to unity. These parameters are obtained by nonlinear regression algorithm built into ActivityBase software (available from ID Business Solutions Ltd., of Guildford, Surrey, UK).

^{33}P phosphoryl transfer assay (radiometric)

[0438] Greiner 384-well white clear bottom high binding plates (available from Greiner Bio-One, Inc., of Longwood, Florida) are coated with 2 μg /well of protein or peptide substrate in a 50 μL volume overnight at ambient temperature. The coating buffer contains 40 $\mu\text{g}/\text{mL}$ substrate, 22.5 mM Na_2CO_3 , 27.5 mM NaHCO_3 , 150 mM NaCl and 3 mM NaN_3 . The coating solution is aspirated and the plates are washed once with 50 μL of assay buffer and padded dry. Subsequently compounds and enzymes are mixed with γ ^{33}P -ATP (3.3 $\mu\text{Ci}/\text{nmol}$) in a total volume of 20 μL in suitable assay buffers (see Table 2). For example the final EphB4 reaction solution contains 20 mM TrisHCl , pH 7.5, 10 mM MgCl_2 , 0.01% Triton X-100, 0.1 mM NaVO_3 , 5 nM EphB4 enzyme and 5 μM ATP.

The mixture is incubated at ambient temperature for 1.5-2.5 hrs as indicated in Table 2 and stopped by aspirating using an EMBLA 96-head washer. The plates are subsequently washed 6-12 times with PBST or TBS buffer. Scintillation fluid (50µl/well) is then added, the plates are sealed and activity assessed by liquid scintillation spectrometry on a Perkin Elmer MicroBeta TriLux (available from PerkinElmer Life and Analytical Sciences, Inc., of Boston Massachusetts).

Luciferase-coupled chemiluminescent assay (LCCA)

[0439] In the LCCA assays, kinase activity is measured by the ATP consumption that is accurately measured by luciferase-coupled chemiluminescence. Greiner 384-well white clear bottom medium binding plates are used for LCCA. Briefly the kinase reaction is initiated by mixing compounds, ATP and kinases in a 20 µL volume. The mixture is incubated at ambient temperature for 2-4 hrs as indicated in Table 2. At the end of the kinase reaction, a 20 µL luciferase-luciferin mix is added and the chemiluminescent signal is read on a Wallac Victor² reader. The luciferase-luciferin mix consists of 50 mM HEPES, pH 7.8, 8.5 ug/mL oxalic acid (pH 7.8), 5 (or 50) mM DTT, 0.4% Triton X-100, 0.25 mg/mL coenzyme A, 63 µM AMP, 28 ug/mL luciferin and 40,000 units of light/mL luciferase. For the LCCA assays, the ATP consumption has been kept at 25-45%, where the decrease in substrate concentration has less than 35% effect on IC₅₀ values compared to the "theoretical" values with no substrate turnover. The IC₅₀ values correlates well with those of radiometric assays.

AlphaScreen

[0440] In AlphaScreen, when the donor and acceptor beads are close in proximity, a series of photochemical events will give rise to a fluorescent signal upon light activation. Here we use biotinylated poly-(Glu, Tyr) 4:1 as the kinase substrate, streptavidin-coated donor beads and anti-phosphotyrosine antibody PY100-coated acceptor beads. Upon phosphorylation, the peptide substrate can bind to both donor and acceptor beads, thus gives rise to fluorescence. Compounds, ATP, biotinylated poly-(Glu, Tyr) and kinases are mixed in a volume of 20 µL for 1 hr at ambient temperature using Greiner 384-well white clear bottom medium binding plates. Then 10 uL solution containing 15-30 mg/mL AlphaScreen beads, 75 mM Hepes, pH 7.4, 300 mM NaCl, 120 mM EDTA, 0.3% BSA and 0.03% Tween-20 is added to each well. After 2-16 hr incubation of the beads, plates

are read in a Perkin Elmer AlphaQuest reader (available from PerkinElmer Life and Analytical Sciences, Inc., of Boston Massachusetts). The IC₅₀ values correlate well with those of radiometric assays.

[0441] Enzymes were purchased from ProQinase (of Freiburg, Germany) and Panvera (of Madison, Wisconsin).

Table 2

Enzyme	Assay format	[E]	[ATP]	Substrate	[Substrate]	Incubation Time (min)	Assay Buffer	Enzyme Construct/Preparation	Enzyme Source
EphB4	Radiometric	5nM	5 μ M	poly-AEKY	2 μ g/well	150	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 0.1 mM NaVO ₃ , 0.01% Triton	Kinase domain E605-E890 with N-terminal His6 tag is expressed in baculovirus and purified by IMAC chromatography.	PanVera
EphA2	LCCA	20nM	3 μ M	poly-EY	1.6 μ M	180	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 3 mM MnCl ₂ , 0.01% Triton	N598-R890 with N-terminal His6 tag is expressed in baculovirus and purified by IMAC chromatography.	PanVera
EGFR	LCCA	7nM 11nM	3 μ M	poly-EY	1.6 μ M	210	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 3 mM MnCl ₂ , 1 mM DTT, 0.01% Triton	amino acids H672-A1210, N-terminally fused to GST-HIS6-Thrombin cleavage site, expressed in baculovirus, one-step affinity purification using GSH-agarose	ProQinase
ErbB2	Alphascreen	1nM	3 μ M	poly-EY	5nM	60	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 3 mM MnCl ₂ , 1 mM DTT, 0.01% Triton	Kinase domain Q679-V1255 with N-terminal GST-HIS6 tag is expressed in baculovirus infected Sf9 cells and purified by GSH-agarose	ProQinase
KDR	LCCA	5nM 6nM	3 μ M	poly-EY	1.6 μ M	240	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 3 mM MnCl ₂ , 1 mM DTT, 0.01% Triton	Human KDR c-DNA, Amino Acids D807-V1356, N-terminally fused to GST. One-step affinity purification using GSH-agarose	ProQinase
Flt-1	Radiometric	6nM	5 μ M	poly-EY	2 μ g/well	120	20 mM TrisHCl, pH 7.5, 10 mM MgCl ₂ , 3 mM MnCl ₂ , 1 mM DTT, 0.01% Triton	Human VEGF-R1 c-DNA, cytoplasmic domain, N-terminally fused to GST-Factor X cleavage site. One-step affinity purification using GSH-agarose	ProQinase

Structure Activity Relationships

[0442] Tables 3 - 6 show structure activity relationship data for selected compounds of the invention. As the data indicates, compounds of the invention fall into different classes; some are “spectrum selective” (*supra*), some are selective for Ephrin, some selective for EGFR, and some inhibit Ephrin and EGFR selectively, for example. Inhibition is indicated as IC₅₀ with the following key: A = IC₅₀ less than 50 nM, B = IC₅₀ greater than 50 nM, but less than 1000 nM, C = IC₅₀ greater than 1000 nM, but less than 20,000 nM, and D = IC₅₀ greater than 20,000 nM. Abbreviations for enzymes listed in Tables 3 - 6 are defined as follows: EphB4 and EphA2 refer to ephrin receptor tyrosine kinase family members ephrin B4 and A2; KDR, kinase insert domain receptor tyrosine kinase, and Flt-1, fms-like tyrosine kinase-1, are representative of the FLK family or receptor tyrosine kinases; EGFR, epidermal growth factor receptor tyrosine kinase, and ErbB2 are representative of the HER family of receptor tyrosine kinases.

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
1	N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	A
8	N-(4-bromo-2,3-dichlorophenyl)-7-({[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	C	A	C
9	N-(4,5-dichloro-2-fluorophenyl)-7-({[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	B	A	B
10	N-(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	B	A	B
11	N-(3-chloro-2,4-difluorophenyl)-7-({[(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	B	A	B

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
12	N-(3,4-dichloro-2-fluorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	A	A	A
13	N-(4-bromo-3-chloro-2-fluorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	A	A	A
14	N-(3-chloro-2,4-difluorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	B	A	B
15	N-(3,4-dichlorophenyl)-7-[(hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	B	A	A
16	N-(4,5-dichloro-2-fluorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	B	A	B
17	N-(4-bromo-2,3-dichlorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	B	A	C
18	N-(4-bromo-5-chloro-2-fluorophenyl)-7-([(3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	B	A	B
19	N-(3,4-dichloro-2-fluorophenyl)-7-([(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	B	A	B
20	N-(4-bromo-3-chloro-2-fluorophenyl)-7-([(3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	A	B	A	B
21	N-(3,4-dichlorophenyl)-7-([(3R,8aR)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	B	A	B
22	N-(4-bromo-5-chloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	B

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
23	N-(3,4-dichlorophenyl)-7-([(3S,8aR)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	A
24	N-(3,4-dichlorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	B
25	N-(3,4-dichlorophenyl)-7-([(3R,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	A
26	N-(3,4-dichloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	B
27	N-(4-bromo-3-chloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	A	A	B
28	N-(3-chloro-2,4-difluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	B	B	A	B
29	N-(4-bromo-2,3-dichlorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	B
30	N-(4,5-dichloro-2-fluorophenyl)-7-([(3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	B	A	B
32	1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol	B	C	B	C	A	B
36	1,4:3,6-dianhydro-5-([4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-glucitol	A	C	B	C	A	C
38	1,4:3,6-dianhydro-5-deoxy-5-([4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol	B	B	B	C	A	B

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
39	(3S,9aS)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2H-pyrido[1,2-a]pyrazin-1(6H)-one	A	A	A	B	A	C
40	(3S,9aR)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2H-pyrido[1,2-a]pyrazin-1(6H)-one	A	B	B	B	A	C
41	(3S,8aS)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	A	B	A	B	A	B
42	(3S,8aR)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	A	A	A	B	A	C
43	(3S,8aS)-3-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	A	B	A	B	A	C
44	(3S,8aS)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-a]pyrazin-1(2H)-one	A	B	B	B	A	C
46	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[[(8aR)-tetrahydro-1H-[1,3]thiazolo[4,3-c][1,4]oxazin-6-ylmethyl]oxy}quinazolin-4-amine	A	B	B	B	A	B
49	N-(3,4-dichlorophenyl)-7-[[[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]oxy]-6-(methyloxy)quinazolin-4-amine	B	B	B	C	A	B
51	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	C
52	1,4:3,6-dianhydro-2-O-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	A	C	C	C	A	B
53	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	B	C	A	B

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
54	1,4:3,6-dianhydro-2-O-methyl-5-O-{6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl}-L-iditol	B	C	C	C	A	C
55	1,4:3,6-dianhydro-5-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-D-xylo-hexitol	B		B	C	A	B
56	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	B
57	dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-sorbose ethylene glycol acetal	B	C	C	C	A	B
58	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	C
59	1,4:3,6-dianhydro-2-O-[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	C
60	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-(difluoromethyl)-L-iditol	B	C	C	B	A	B
61	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	B
62	1,4:3,6-dianhydro-2-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	C	A	B
63	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	B	C	C	D	A	B
64	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-ethyl-L-iditol	B	C	C	B	A	B

Table 3

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
65	1,4:3,6-dianhydro-2-O-[4-[(3-bromo-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	C	C	D	D	A	C
66	1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol	C	C	D	D	A	C
67	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-D-xylo-hexitol	C	C	C	C	A	B
68	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-D-glucitol	C	C	C	C	A	C

Table 5

Entry	Name	EphB4 IC ₅₀
1	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]quinazolin-4-amine	A
2	N-(3,4-dichlorophenyl)-7-[(5-[(dimethylamino)methyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	A
3	N-(3,4-dichlorophenyl)-7-[(3-[(dimethylamino)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	B
4	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-[(4-methylpiperazin-1-yl)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	B
5	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-piperidin-4-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	A
6	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(5-(1-methylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	A
7	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-(morpholin-4-yl)methyl)-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	B
8	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(morpholin-2-yl)methyl]oxy]quinazolin-4-amine	A

Table 5

Entry	Name	EphB4 IC ₅₀
9	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-piperidin-2-yl-1,2,4-oxadiazol-3-yl)methyl]oxy}quinazolin-4-amine	A
10	N-(3,4-dichlorophenyl)-7-[[2-[(dimethylamino)methyl]-1,3-thiazol-4-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	B
11	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(phenylmethyl)morpholin-2-yl)methyl]oxy)quinazolin-4-amine	B
12	1,1-dimethylethyl 2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholine-4-carboxylate	B
13	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-(morpholin-4-ylmethyl)-1,3-thiazol-4-yl)methyl]oxy)quinazolin-4-amine	B
14	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-[(4-methylpiperazin-1-yl)methyl]-1,3-thiazol-4-yl)methyl]oxy)quinazolin-4-amine	A
15	N-(3,4-dichlorophenyl)-7-([4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A
16	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(1,4-oxazepan-2-ylmethyl)oxy]quinazolin-4-amine	A
17	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-piperidin-3-yl-1,2,4-oxadiazol-3-yl)methyl]oxy}quinazolin-4-amine	A
18	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(1-methylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy)quinazolin-4-amine	A
19	N-(3,4-dichlorophenyl)-7-([4-methyl-1,4-oxazepan-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A
20	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-(1-methylpiperidin-3-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy)quinazolin-4-amine	A

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
1	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(1,3-thiazol-4-ylmethyl)oxy]quinazolin-4-amine	B	B	B	B	A	-
2	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(pyridin-3-ylmethyl)oxy]quinazolin-4-amine	-	B	B	B	-	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
3	7-[(cyclopropylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	B
4	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydrofuran-2-ylmethyl)oxy]quinazolin-4-amine	A	B	B	-	A	-
5	7-(cyclopentyloxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	C	-	A	-
6	methyl 6-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-alpha-D-glucopyranoside	A	B	B	-	A	B
7	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-morpholin-4-yl-2-oxoethyl)oxy]quinazolin-4-amine	C	B	C	-	A	-
8	1,1-dimethylethyl 2-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate	B	C	C	-	A	-
9	1,1-dimethylethyl 4-[3-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,2,4-oxadiazol-5-yl]piperidine-1-carboxylate	B	B	B	-	A	-
10	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(4-pyrrolidin-1-ylphenyl)-1,3-thiazol-2-yl]methyl}oxy)quinazolin-4-amine	C	C	C	-	A	-
11	N-(3,4-dichlorophenyl)-7-({[4-[4-(diethylamino)phenyl]-1,3-thiazol-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-
12	5-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-4-yl]-2-hydroxybenzamide	C	C	C	-	A	-
13	7-[(2-cyclohexylethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-
14	7-[(cyclohexylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	C	C	-	A	-
15	7-[(cyclobutylmethyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	C	C	-	A	-
16	N-(3,4-dichlorophenyl)-7-{[2-(1,3-dioxolan-2-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
17	N-(3,4-dichlorophenyl)-7-([2-(1,3-dioxan-2-yl)ethyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
18	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-morpholin-4-ylethyl)oxy]quinazolin-4-amine	B	B	B	-	A	-
19	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-pyrrolidin-1-ylethyl)oxy]quinazolin-4-amine	B	B	B	-	A	-
20	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-piperidin-1-ylethyl)oxy]quinazolin-4-amine	B	B	B	-	A	-
21	2-(2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl)-1H-isoindole-1,3(2H)-dione	A	B	B	-	A	B
22	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-pyridin-3-yl-1,3-thiazol-2-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
23	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-pyridin-2-yl-1,3-thiazol-2-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
24	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-pyridin-4-yl-1,3-thiazol-2-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
25	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[2-morpholin-4-yl-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	-	C		-	A	C
26	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[3-morpholin-4-yl-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	-	C	C	-	A	-
27	N-(3,4-dichlorophenyl)-7-([3-(dimethylamino)-1,2,4-oxadiazol-5-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	A	C	C	-	A	-
28	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-[(4-methylpiperazin-1-yl)methyl]-1,3-thiazol-2-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	-
29	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	B
30	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(morpholin-4-yl)methyl]-1,3-thiazol-2-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
31	N-(3,4-dichlorophenyl)-7-[(4-[(4-methyl-1,4-diazepan-1-yl)methyl]-1,3-thiazol-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	-
32	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-[(phenylmethyl)oxy]methyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	B	C	B	-	A	-
33	N-(3,4-dichlorophenyl)-7-[(4-ethylmorpholin-2-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	B
34	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(2-piperidin-4-yl-1,3-thiazol-4-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	A
35	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[2-(1-methylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl]oxy]quinazolin-4-amine	B	B	A	-	A	B
36	1,1-dimethylethyl 4-[5-({4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]-1,2,4-oxadiazol-3-yl]piperazine-1-carboxylate	C	C	C	-	A	-
37	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[(3-piperazin-1-yl-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	B	A	A	-	A	A
38	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([3-(4-methylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl]oxy)quinazolin-4-amine	B	A	A	-	A	A
39	N-(3,4-dichlorophenyl)-7-([5-(1-ethylpiperidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	A
40	N-(3,4-dichlorophenyl)-7-([3-(4-ethylpiperazin-1-yl)-1,2,4-oxadiazol-5-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	-
41	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([5-[4-(methyloxy)phenyl]-1,2,4-oxadiazol-3-yl]methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
42	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([2-[4-(trifluoromethyl)phenyl]-1,3-thiazol-4-yl]methyl]oxy]quinazolin-4-amine	C	C	C	--	B	-
43	7-([2-(4-chlorophenyl)-1,3-thiazol-4-yl]methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
44	N-(3,4-dichlorophenyl)-7-([5-(3,5-dimethylisoxazol-4-yl)-1,2,4-oxadiazol-3-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
45	7-[[5-chloro-1-benzothien-3-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
46	N-(3,4-dichlorophenyl)-7-[[3-[4-(1,1-dimethylethyl)phenyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
47	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-[2-(methyloxy)phenyl]-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
48	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-(4-methylphenyl)-1,3,4-oxadiazol-2-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
49	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[1-(phenylmethyl)-1H-imidazol-2-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
50	N-(3,4-dichlorophenyl)-7-[[3-(2,6-dichlorophenyl)-5-methylisoxazol-4-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
51	N-(3,4-dichlorophenyl)-7-[[6-fluoro-4H-1,3-benzodioxin-8-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-
52	7-[[3,5-dibromophenyl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
53	N-(3,4-dichlorophenyl)-7-[[2,6-difluorophenyl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	B	C	C	-	A	-
54	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[3-[(pyridin-2-yl)sulfonyl)methyl]-1,2,4-oxadiazol-5-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	-
55	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[5-phenyl-1,2,4-oxadiazol-3-yl)methyl]oxy]quinazolin-4-amine	C	C	C	-	A	-
56	7-[[4-chloro-2-(trifluoromethyl)quinolin-6-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	B	-
57	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[2-(1-methylpyrrolidin-2-yl)ethyl]oxy]quinazolin-4-amine	B	B	B	-	A	-
58	N-(3,4-dichlorophenyl)-7-[[5-(1-ethylpiperidin-4-yl)-1,2,4-oxadiazol-3-yl)methyl]oxy]-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	A

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
59	N-(3,4-dichlorophenyl)-7-({[5-(1-ethylpiperidin-3-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	A
60	N-(3,4-dichlorophenyl)-7-({[2-(dimethylamino)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-
61	N-(3,4-dichlorophenyl)-7-({[4-ethyl-1,4-oxazepan-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	B
62	N-(3,4-dichlorophenyl)-7-({[2-(1-ethylpiperidin-4-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	-
63	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-[(2S)-pyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine	A	B	B	-	A	-
64	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-[(2S)-pyrrolidin-2-yl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
65	[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]methyl benzoate	C	C	C	-	A	-
66	[4-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,3-thiazol-2-yl]methanol	B	C	B	-	A	-
67	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-methyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl]methyl}oxy)quinazolin-4-amine	A	B	B	-	A	A
68	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-[(4S)-1,3-thiazolidin-4-yl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
69	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-piperidin-2-yl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	A	A	A	-	A	A
70	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(1-methylpiperidin-2-yl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	A	-	A	-
71	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-piperidin-3-yl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	
72	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-(1-methylpiperidin-3-yl)-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	B

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
73	N-(3,4-dichlorophenyl)-7-({[2-(1-ethylpiperidin-2-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	-
74	N-(3,4-dichlorophenyl)-7-({[2-(1-ethylpiperidin-3-yl)-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	C	B	-	A	-
75	N-(3,4-dichlorophenyl)-7-({[3-[(2S)-1-ethylpyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
76	N-(3,4-dichlorophenyl)-7-({[2-[(2S)-1-ethylpyrrolidin-2-yl]-1,3-thiazol-4-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
77	N-(3,4-dichlorophenyl)-7-({[5-ethyl-4,5,6,7-tetrahydro[1,3]thiazolo[5,4-c]pyridin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	A
78	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-propyl-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	A	A	A	-	B	B
79	7-({[4-(cyclopropylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	-
80	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[2-(methyloxy)ethyl]-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	A	B	B	-	A	-
81	N-(3,4-dichlorophenyl)-7-({[4-(1-methylethyl)-1,4-oxazepan-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
82	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-piperazin-1-yl-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
83	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[5-pyrrolidin-2-yl-1,2,4-oxadiazol-3-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
84	N-(3,4-dichlorophenyl)-7-({[5-(1-ethylpyrrolidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
85	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[3-[(2S)-1-methylpyrrolidin-2-yl]-1,2,4-oxadiazol-5-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
86	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[2-[(2S)-1-methylpyrrolidin-2-yl]-1,3-thiazol-4-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
87	N-(3,4-dichlorophenyl)-7-([2-(4-ethylpiperazin-1-yl)-1,3-thiazol-4-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	-
88	N-(3,4-dichlorophenyl)-7-([(1,4-dimethylpiperazin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	C	-	A	-
89	7-[(4-cyclopentylmorpholin-2-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	A	A	-	A	-
90	N-(3,4-dichlorophenyl)-7-([4-(1-methylethyl)morpholin-2-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	-
91	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(3-phenylpropyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	B	B	-	A	-
92	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-[2-(methyloxy)ethyl]morpholin-2-yl]methyl)oxy)quinazolin-4-amine	A	B	B	-	A	-
93	ethyl 2-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]propanoate	B	B	B	-	A	-
94	N-(3,4-dichlorophenyl)-7-([(4-hex-5-en-1-yl)morpholin-2-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	-
95	2-([2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]ethyl)oxy)ethanol	B	B	B	-	A	-
96	methyl 3-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]propanoate	A	A	A	-	A	-
97	6-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]hexanenitrile	A	B	A	-	A	B
98	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(tetrahydro-2H-pyran-2-yl)methyl]morpholin-2-yl]methyl)oxy)quinazolin-4-amine	A	A	A	-	A	-
99	4-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]butanenitrile	A	A	A	-	A	-
100	N-(3,4-dichlorophenyl)-7-([4-[(4-fluorophenyl)methyl]morpholin-2-yl]methyl)oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
101	methyl 5-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]pentanoate	B	B	B	-	A	-
102	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-oct-7-en-1-ylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	B	B	-	A	-
103	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-propylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	B	B	-	A	-
104	6-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]hexan-1-ol	A	B	A	-	A	-
105	7-{{[4-acetylmorpholin-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
106	7-{{[4-(cyclopropylmethyl)morpholin-2-yl]methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	-
107	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-prop-2-yn-1-ylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	-
108	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-pyridin-4-ylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	-
109	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-(pyridin-2-yl)methyl)morpholin-2-yl]methyl]oxy}quinazolin-4-amine	B	B	A	-	A	-
110	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[4-pent-2-yn-1-ylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	B	B	-	A	-
111	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[2-(4-methylpiperazin-1-yl)-1,3-thiazol-4-yl]methyl]oxy}quinazolin-4-amine	B	B	B	-	A	B
112	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[5-(1-methylpyrrolidin-2-yl)-1,2,4-oxadiazol-3-yl]methyl]oxy}quinazolin-4-amine	B	B	B	-	A	-
113	N-(3-chloro-4-fluorophenyl)-7-{{[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
114	7-{{[4-acetyl-1-ethylpiperazin-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	C	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
115	N-[4-chloro-2,5-bis(methoxy)phenyl]-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	C	-	B	-
116	N-(3-bromo-2-methylphenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
117	7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)-N-(3,4,5-trichlorophenyl)quinazolin-4-amine	A	B	B	-	A	B
118	N-(3-chloro-2-methylphenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	A	A	B	-	A	-
119	N-(3,4-dichlorophenyl)-7-[[[4-ethanimidoyl-1,4-oxazepan-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	A	A	A	-	A	A
120	N-(4-bromo-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	A	-	A	-
121	N-(5-chloro-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
122	N-(4-chloro-2-fluorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	A	-	A	-
123	N-(2,4-dichlorophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
124	N-(2,4-dibromophenyl)-7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
125	7-[[[4-methylmorpholin-2-yl)methyl]oxy]-6-(methoxy)-N-(2,3,4-trichlorophenyl)quinazolin-4-amine	A	A	A	-	A	B
126	N-(3,4-dichlorophenyl)-7-[[[1-ethyl-4-methylpiperazin-2-yl)methyl]oxy]-6-(methoxy)quinazolin-4-amine	B	B	C	-	A	-
127	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methoxy)quinazolin-7-yl]oxy)methyl)morpholine-4-carboximidamide	A	A	A	-	A	A
128	N-(3,4-dichlorophenyl)-6-(methoxy)-7-([2-(pyrrolidin-1-yl)methyl]-1,3-thiazol-4-yl)methyl]oxy)quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
129	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(tetrahydro-2H-pyran-4-yl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	A	B	B	-	A	-
130	N-(3,4-dichlorophenyl)-7-({[4-(2-ethylbutyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
131	7-({[4-(cyclohexylmethyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
132	2-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]ethanol	B	B	B	-	A	-
133	7-({[4-but-2-yn-1-ylmorpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
134	7-({[4-(cyclobutylmorpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	-
135	N-(3,4-dichlorophenyl)-7-({[4-[2-(1,3-dioxolan-2-yl)ethyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
136	7-({[4-(2-cyclohexylethyl)morpholin-2-yl]methyl}oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
137	N-(3,4-dichlorophenyl)-7-({[4-[2-(1,3-dioxan-2-yl)ethyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
138	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-pent-4-en-1-ylmorpholin-2-yl]methyl}oxy)quinazolin-4-amine	A	B	A	-	A	-
139	N-(3,4-dichlorophenyl)-7-({[4-[(2R)-2-methylbutyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
140	N-(3,4-dichlorophenyl)-7-({[4-(4-fluorobutyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	-
141	3-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]butan-2-one	B	B	B	-	A	-
142	1-[2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)morpholin-4-yl]butan-2-one	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
143	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-pentylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	B	B	-	A	-
144	N-(3,4-dichlorophenyl)-7-[[4-hexylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
145	N-(3,4-dichlorophenyl)-7-[[4-heptylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	C	B	-	A	-
146	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4-octylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	C	C	-	A	-
147	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(2-phenylethyl)morpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	C	B	-	A	-
148	7-[[4-butylmorpholin-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	B
149	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-prop-2-en-1-ylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B	B	B	-	A	-
150	2-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]-1-phenylethanone	B	B	B	-	A	-
151	N-(3,4-dichlorophenyl)-7-([4-(2-fluoroethyl)morpholin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
152	N-(3,4-dichlorophenyl)-7-([4-(3-methylbut-2-en-1-yl)morpholin-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
153	7-([4-[(2E)-3-bromoprop-2-en-1-yl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
154	2-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]morpholin-4-yl]acetamide	A	B	B	-	A	A
155	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-[3-(tetrahydro-2H-pyran-2-yloxy)propyl]-1,4-oxazepan-2-yl)methyl]oxy}quinazolin-4-amine	A	B	A	-	A	-
156	N-(3,4-dichlorophenyl)-7-([4-(3-methylbutyl)-1,4-oxazepan-2-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
157	7-({[4-(cyclohexylmethyl)-1,4-oxazepan-2-yl]methyl}oxy)-4-[(3,4-dichlorophenyl)methyl]-6-(methyloxy)quinazoline	B	B	A	-	A	-
158	7-({[4-(2-cyclohexylethyl)-1,4-oxazepan-2-yl]methyl}oxy)-4-[(3,4-dichlorophenyl)methyl]-6-(methyloxy)quinazoline	B	B	B	-	A	-
159	N-(3,4-dichlorophenyl)-7-({[4-(2-ethylbutyl)-1,4-oxazepan-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
160	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(methylsulfonyl)-1,4-oxazepan-2-yl]methyl}oxy)quinazolin-4-amine	A	B	B	-	A	A
161	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(1-methylpiperidin-4-yl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	A	B	A	-	A	-
162	N-(3-chloro-2-fluorophenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
163	N'-cyano-2-({[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-1,4-oxazepane-4-carboximidamide	A	B	B	-	A	A
164	N-(3-bromo-4-methylphenyl)-7-({[4-methylmorpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	B
165	N-(3,4-dichlorophenyl)-7-({[1,4-diethylpiperazin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	B	C	-	A	-
166	4-({[4-[(4-bromo-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-N'-cyanopiperidine-1-carboximidamide	B	B	A	-	A	-
167	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-(methylsulfonyl)morpholin-2-yl]methyl}oxy)quinazolin-4-amine	A	A	B	-	A	A
168	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-({[4-[(phenylmethyl)sulfonyl]morpholin-2-yl]methyl}oxy)quinazolin-4-amine	B	B	B	-	A	-
169	N-(3,4-dichlorophenyl)-7-({[4-[(4-fluorophenyl)sulfonyl]morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	B	C	B	-	A	-
170	N-(3,4-dichlorophenyl)-7-({[4-(ethylsulfonyl)morpholin-2-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine	A	A	B	-	A	A

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
171	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(phenylsulfonyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	C	B	-	A	-
172	7-([4-([3-chloropropyl)sulfonyl]morpholin-2-yl]methyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	B
173	7-([4-(butylsulfonyl)morpholin-2-yl]methyl)oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
174	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-([4-methylphenyl)sulfonyl]morpholin-2-yl]methyl)oxy]quinazolin-4-amine	C	C	B	-	A	-
175	N-(3,4-dichlorophenyl)-7-([4-([3,5-dimethylisoxazol-4-yl)carbonyl]morpholin-2-yl]methyl)oxy]-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	A
176	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-([3-(methyloxy)phenyl]acetyl]morpholin-2-yl]methyl)oxy]quinazolin-4-amine	B	B	B	-	A	-
177	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(2-methylpentanoyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	B	B	-	A	-
178	7-([4-([4-butylphenyl]carbonyl]morpholin-2-yl]methyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	C	C	C	-	A	-
179	7-([4-([4-chlorophenyl]acetyl]morpholin-2-yl]methyl)oxy]-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	B	C	B	-	A	-
180	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(2-propylpentanoyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	B	B	-	A	-
181	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(4-methylpentanoyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	B	B	-	A	-
182	N-(3,4-dichlorophenyl)-7-([4-([2,5-difluorophenyl]carbonyl]morpholin-2-yl]methyl)oxy]-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
183	7-([4-(cyclopentylcarbonyl)morpholin-2-yl]methyl)oxy)-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	A
184	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-([4-(2-phenylbutanoyl)morpholin-2-yl]methyl)oxy)quinazolin-4-amine	B	C	B	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
185	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-[(2,3,6-trifluorophenyl)carbonyl]morpholin-2-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	-
186	N-(3,4-dichlorophenyl)-7-[(4-(furan-3-ylcarbonyl)morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	A	B	B	-	A	-
187	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-propanoylmorpholin-2-yl)methyl]oxy]quinazolin-4-amine	A	B	B	-	A	A
188	N-(3,4-dichlorophenyl)-7-[(4-hexanoylmorpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
189	N-(3,4-dichlorophenyl)-7-[(4-(2-ethylhexanoyl)morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
190	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-(3-phenylpropanoyl)morpholin-2-yl)methyl]oxy]quinazolin-4-amine	B	B	B	-	A	-
191	N-(3,4-dichlorophenyl)-7-[(4-(2,2-dimethylpropanoyl)morpholin-2-yl)methyl]oxy)-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
192	N-(3,4-dichlorophenyl)-6-(methoxy)-7-[(4-(naphthalen-1-ylcarbonyl)morpholin-2-yl)methyl]oxy]quinazolin-4-amine	B	C	B	-	A	-
193	7-[(4-[(2-chloropyridin-3-yl)carbonyl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	A	B	A	-	A	A
194	7-[(4-[(6-chloropyridin-3-yl)carbonyl]morpholin-2-yl)methyl]oxy]-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	A	B	B	-	A	-
195	7-[(4-(1,3-benzodioxol-5-ylcarbonyl)morpholin-2-yl)methyl]oxy)-N-(3,4-dichlorophenyl)-6-(methoxy)quinazolin-4-amine	B	B	B	-	A	-
196	N-(3,4-dichlorophenyl)-6-[(1-methylethyl)oxy]-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	A	A	A	-	A	A
197	N-(3,4-dichlorophenyl)-6-[[2-(methoxy)ethyl]oxy]-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	A	B	A	-	A	-
198	N-(3,4-dichlorophenyl)-6-(ethoxy)-7-[(morpholin-2-ylmethyl)oxy]quinazolin-4-amine	A	A	A	-	A	A

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
199	N-(3,4-dichlorophenyl)-6-(ethyloxy)-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	A
200	N-(4-bromo-2-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
201	N-(4-chloro-3-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
202	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-N-methylmorpholine-4-carboximidamide	A	B	B	-	A	A
203	N-(4-bromo-3-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	B
204	N-(3,4-dichlorophenyl)-6-[(1-methylethyl)oxy]-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	A
205	N-(3,4-dichlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-[[2-(methyloxy)ethyl]oxy}quinazolin-4-amine	A	B	B	-	A	-
206	N-(4-bromo-2-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
207	7-[[4-acetyl-1,4-oxazepan-2-yl)methyl]oxy}-N-(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	A
208	4-[(3,4-dichlorophenyl)amino]-7-[[4-methylmorpholin-2-yl)methyl]oxy}quinazolin-6-ol	B	B	B	-	A	-
209	N-(3-bromo-4-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
210	3-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl]-3-oxopropanoic acid	A	A	A	-	A	-
211	methyl 4-[2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl]-4-oxobutanoate	B	B	B	-	A	-
212	N-(3,4-dichlorophenyl)-7-[[4-methylmorpholin-3-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	C	-	A	-

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
213	N-(3-bromo-2-chlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	-
214	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-N-[2-(methyloxy)ethyl]morpholine-4-carboximidamide	A	B	B	-	A	-
215	N'-cyano-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-N-ethylmorpholine-4-carboximidamide	A	B	B	-	A	A
216	[(1E)-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl](piperidin-1-yl)methylidene]cyanamide	A	B	B	-	A	A
217	[(1E)-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl](pyrrolidin-1-yl)methylidene]cyanamide	A	A	B	-	A	A
218	[(1E)-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)morpholin-4-yl](4-methylpiperazin-1-yl)methylidene]cyanamide	A	B	A	-	A	-
219	N-(3,4-dichlorophenyl)-7-[[6-ethyl-4,6-dimethylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	-
220	N-(4-bromo-3-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	B
221	N-(3,4-dichlorophenyl)-7-[[6,6-dimethylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B		A	-	A	B
222	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4,6,6-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	B		B	-	A	B
223	N-(3,4-dichlorophenyl)-7-[[2-(5,5-dimethylmorpholin-2-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	B
224	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[2-(4,5,5-trimethylmorpholin-2-yl)ethyl]oxy}quinazolin-4-amine	B	B	B	-	A	B
225	1,1-dimethylethyl 2-(2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)ethyl)-5,5-dimethylmorpholine-4-carboxylate	C	C	C	-	A	C
226	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	B	A	-	A	A

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
227	N-(4-bromo-2,3-dichlorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	B
228	N-(4,5-dichloro-2-fluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	B
229	N-(3,4-dichlorophenyl)-6-(methyloxy)-7-[[2-(4,6,6-trimethylmorpholin-2-yl)ethyl]oxy}quinazolin-4-amine	B	C	B	-	A	B
230	N-(4-bromo-2,3-difluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	B
231	N-(4-bromo-2,5-difluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	B
232	N-(4-bromo-3,5-difluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	B
233	N-(3,4-dichloro-2-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	B	-	A	C
234	N-(4-bromo-3-chloro-2-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	B	-	A	B
2235	N-(4-bromo-5-chloro-2-fluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	B	A	-	A	B
236	N-(4-bromo-3-chloro-2-fluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	A
237	N-(3,4-dichloro-2-fluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	A
238	N-(3-chloro-2,4-difluorophenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	A	A	A	-	A	A
239	N-(2,3-dichloro-4-methylphenyl)-7-[[4-methylmorpholin-2-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine	B	B	A	-	A	C
240	6-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-3,3,4-trimethylmorpholin-2-one	A	B	B	-	A	B

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
241	N-(4-bromo-2,3-dichlorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	B	-	A	B
242	N-(4-bromo-5-chloro-2-fluorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	B
243	N-(4,5-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	A
244	N-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	A
245	N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	A	-	A	A
246	N-(3-chloro-2,4-difluorophenyl)-6-(methyloxy)-7-[[4,5,5-trimethylmorpholin-2-yl)methyl]oxy}quinazolin-4-amine	A	A	B	-	A	B
247	(6S)-6-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-4-methylpiperazin-2-one	B	B	B	-	A	B
248	(6S)-6-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-4-methylpiperazin-2-one	B	B	B	-	A	B
249	(6S)-6-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-1,4-dimethylpiperazin-2-one	B	B	B	-	A	C
250	(6S)-6-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-1,4-dimethylpiperazin-2-one	B	B	B	-	A	C
251	N-(4-bromo-3-chlorophenyl)-7-[[[(3a'S,4R,6'S,6a'R)-2,2-dimethyltetrahydrospiro[1,3-dioxolane-4,3'-furo[3,2-b]furan]-6'-yl]oxy]-6-(methyloxy)quinazolin-4-amine	B	C	C	-	A	C
253	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-(methylsulfonyl)-D-glucitol	C	C	C	-	A	C
254	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-glucitol	B	C	C	-	A	B
255	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-S-methyl-5-thio-L-iditol	B	C	C	-	A	C

Table 6


Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
256	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-morpholin-4-yl-L-iditol	B	C	C	-	A	C
257	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(4-methylpiperazin-1-yl)-L-iditol	B	C	C	-	A	C
258	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-pyrrolidin-1-yl-L-iditol	B	C	C	-	A	C
259	2-O-acetyl-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol	B	C	C	-	A	C
260	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol	B	C	C	-	A	B
261	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(methylsulfonyl)-L-iditol	B	C	C	-	A	B
262	2-amino-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-L-iditol	B	C	C	-	A	C
263	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(dimethylamino)-L-iditol	B	C	C	-	A	C
264	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(diethylamino)-L-iditol	B	C	B	-	A	C
265	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-piperidin-1-yl-L-iditol	B	C	C	-	A	C
266	2-(acetilamino)-1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-L-iditol	B	C	C	-	A	C
267	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-5-C-(trifluoromethyl)-D-glucitol	C	C	C	-	A	B
268	1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-[(methylsulfonyl)amino]-L-iditol	B	C	C	-	A	B
269	N-(4-bromo-3-chlorophenyl)-6-(methyloxy)-7-[(1-methylpyrrolidin-3-yl)oxy]quinazolin-4-amine	B	C	B	-	A	C

Table 6

Entry	Name	EphB4 IC ₅₀	EphA2 IC ₅₀	KDR IC ₅₀	Flt-1 IC ₅₀	EGFR IC ₅₀	ErbB2 IC ₅₀
270	N-(4-bromo-3-chlorophenyl)-6-(methyloxy)-7-[(3R)-tetrahydrofuran-3-yloxy]quinazolin-4-amine	B	C	B	-	A	B
271	1,4:3,6-dianhydro-5-O-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-L-iditol	B	C	C	-	A	B
272	N-(4-bromo-3-chlorophenyl)-6-(methyloxy)-7-[(3S,4R)-4-(methyloxy)tetrahydrofuran-3-yl]oxy]quinazolin-4-amine	B	C	C	-	A	B
273	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-(6-(methyloxy)-4-{[4-(4-methylpiperazin-1-yl)phenyl]amino}quinazolin-7-yl)-L-iditol	C	C	C	-	A	C
274	1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-O-[4-{[3-fluoro-4-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-L-iditol	C	C	C	-	A	C
275	1,4:3,6-dianhydro-2-deoxy-5-O-[4-{[2,3-dichloro-4-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-fluoro-L-iditol	C	C	C	-	A	C
276	1,4:3,6-dianhydro-2-deoxy-5-O-[4-{[3,4-dichloro-2-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-fluoro-L-iditol	C	C	C	-	A	C
277	1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-C-(trifluoromethyl)-D-glucitol	B	C	C	-	A	B

What is claimed is:

1. A compound of Formula I,



or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof, wherein,

R^1 is C_1 - C_3 alkyl optionally substituted with between one and three R^{50} substituents;

R^2 is selected from halogen, trihalomethyl, $-CN$, $-NH_2$, $-NO_2$, $-OR^3$, $-N(R^3)R^4$, $-S(O)_{0-2}R^4$, $-SO_2N(R^3)R^4$, $-CO_2R^3$, $-C(=O)N(R^3)R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(=O)R^3$, $-N(R^3)CO_2R^4$, $-C(=O)R^3$, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R^3 is $-H$ or R^4 ;

R^4 is selected from C_1 - C_6 alkyl; C_1 - C_6 alkyl substituted with one, two, or three halogen; aryl; aryl substituted with one, two, or three halogen; unsubstituted aryl C_1 - C_6 alkyl, heterocyclyl, and heterocyclyl C_1 - C_6 alkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form a five- to seven-membered heterocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P where the five- to seven-membered heterocyclyl is optionally substituted by one, two, or three alkyl;

q is 0, 1, 2, 3, 4, or 5;

Z is selected from $-OCH_2-$, $-O-$, $-S(O)_{0-2}-$, $-N(R^5)CH_2-$, and $-NR^5-$;

R^5 is $-H$ or C_1 - C_6 alkyl;


R^{50} is $-H$, halo, trihalomethyl, $-OR^3$, $-N(R^3)R^4$, $-S(O)_{0-2}R^4$, $-SO_2N(R^3)R^4$, $-CO_2R^3$, $-C(=O)N(R^3)R^4$, $-C(=NR^{25})N(R^3)R^4$, $-C(=NR^{25})R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(=O)R^3$, alkoxy, C_1 - C_6 alkyl, aryl, unsubstituted aryl C_1 - C_6 alkyl, heterocyclyl, and heterocyclyl C_1 - C_6 alkyl optionally substituted with one alkyl; or

two of R^{50} , when taken together on the same carbon are oxo; or

two of R^{50} , when taken together with a common carbon to which they are attached, form a three- to seven-membered spirocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P;

R^{25} is selected from -H, -CN, -NO₂, -OR³, -S(O)₀₋₂R⁴, -CO₂R³, C₁-C₆alkyl, C₂-C₆alkenyl, and C₂-C₆alkynyl;

M¹-M²-M³-M⁴- together are according to formula II:



II

wherein X^1 , X^2 , and optionally X^3 , represent the atoms of a saturated bridged ring system, said saturated bridged ring system containing up to three annular heteroatoms represented by any of X^1 , X^2 , and X^3 ; wherein,

each X^1 is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

each X^2 is independently a bridgehead methine optionally substituted with R⁶, or a bridgehead nitrogen;

each X^3 is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

provided, for X^1 , X^2 , and X^3 , there are no nitrogen-nitrogen annular bonds nor geminal di-nitrogen substitutions;

E is selected from -NR⁹-, -O-, and absent;

Y is either:

- a C₁₋₃ alkylene linker, between the oxygen at the 7-position of the quinazoline ring system of I and either E, or when E is absent, any ring atom of the saturated bridged ring system except X^2 , when X^2 is a bridgehead nitrogen; provided there are at least two carbon atoms between the oxygen at the 7-position of the quinazoline ring system of I and either E, or when E is absent, any heteroatom represented by X^1 , X^2 or X^3 ; or

Y is absent, when Y is absent, E is also absent; said saturated bridged ring system is directly attached to the oxygen at the 7-position of the

quinazoline ring system of I via a carbon atom of said saturated bridged ring system;

m and p are each independently 1, 2, 3, or 4;

n is 0, 1, or 2, when n is zero, then there is a direct single bond between the two bridgehead X²'s;

R⁶ and R⁷ are each independently selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -N(R³)R⁴, -S(O)₀₋₂R⁴, -SO₂N(R³)R⁴, -CO₂R³, -C(O)N(R³)R⁴, -N(R³)SO₂R⁴, -N(R³)C(O)R³, -NCO₂R³, -C(O)R³, C₁-C₆alkyl, aryl, unsubstituted aryl C₁-C₆alkyl, heterocyclyl optionally substituted with one alkyl, heterocyclyl C₁-C₆alkyl; or

R⁶ and R⁷, when taken together are oxo; or

R⁶ and R⁷, when taken together with a common carbon to which they are attached, form a three- to seven-membered spirocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P and which spirocyclic ring is optionally substituted with one or two alkyl; and

R⁸ is selected from R³, -SO₂N(R³)R⁴, -CO₂R³, -C(O)N(R³)R⁴, -SO₂R⁴, and -C(O)R³; and

R⁹ is -H or C₁-C₆alkyl;

with the proviso that when Y is a C₁₋₃ alkylene linker, E is absent, Z is -NH- or -N(CH₃)-, R¹ is a C₁₋₃ alkyl, R² is -H or halogen, n = 0, and the atoms X¹ of one bridge of the saturated bridged ring system, when combined with both bridgehead atoms, X², of the saturated bridged ring system, represent:

either a pyrrolidine ring or a piperidine ring, and any atom, X¹ or X², of either of said pyrrolidine ring or said piperidine ring is attached to Y; then the other bridge of said saturated bridged ring system cannot be any one of -OC(O)CH₂-, -CH₂OC(O)-, -OC(O)CH₂CH₂-, -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, -OC(O)CH₂NH-, -OC(O)CH₂N(C₁₋₄alkyl)-, and -OC(O)CH₂O-; and

either a piperazine ring or a 4-(C₁₋₄ alkyl)-piperazine ring, and any atom, X¹ or X², of either of said piperazine ring or said 4-(C₁₋₄ alkyl)-piperazine ring is attached to Y; then the other bridge of said saturated bridged ring system,

only when attached via the 2- and the 3-position of either of said piperazine ring or said 4-(C₁₋₄ alkyl)-piperazine ring, cannot be one of -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, and either of the two aforementioned bridges cannot be optionally substituted by one or two C₁₋₂alkyl groups; and


a piperazine ring, and any atom, X¹ or X², of said piperazine ring is attached to Y; then the other bridge of said saturated bridged ring system, only when attached via the 3- and the 4-position of said piperazine ring, cannot be one of -C(O)OCH₂CH₂-, -CH₂OC(O)CH₂-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂ alkyl groups, and only when either of the two aforementioned bridges are attached to the 3-position of said piperazine ring via their left-hand end as depicted above; and

a 2-oxomorpholine ring, said 2-oxomorpholine ring attached to Y via its 4-position; then the other bridge of said saturated bridged ring system, only when attached via the 5- and the 6-position of said 2-oxomorpholine ring, cannot be one of -(CH₂)_g-, -CH₂WCH₂-, -CH₂WCH₂CH₂-, and -CH₂CH₂WCH₂-, wherein W is -O-, -S(O)₀₋₂-, -NH-, or -N(C₁₋₄ alkyl)- and wherein g is 2, 3, or 4.

2. The compound according to claim 1, wherein Z is -NR⁵-; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
3. The compound according to claim 2, wherein R² is selected from halogen, trihalomethyl, -CN, -NO₂, -OR³, and C₁-C₆alkyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
4. The compound according to claim 3, wherein R¹ is an unsubstituted C₁₋₃ alkyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.


5. The compound according to claim 4, wherein the saturated bridged ring system has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1]; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
6. The compound according to claim 5, wherein Y is selected from $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-$, and absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
7. The compound according to claim 6, wherein q is 1, 2, or 3; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
8. The compound according to claim 7, wherein R^5 is $-\text{H}$; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
9. The compound according to claim 8, wherein R^1 is methyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
10. The compound according to claim 9, wherein the saturated bridged ring system has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], and [3.1.0]; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
11. The compound according to claim 10, wherein said saturated bridged ring system contains one or two annular nitrogens, said one or two annular nitrogens are selected from $-\text{NR}^8-$, when X^1 , and a bridgehead nitrogen, when X^2 ; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

12. The compound according to claim 11, wherein E is absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
13. The compound according to claim 12, wherein said saturated bridged ring system is according to formula III;





III

- wherein A is selected from -O-, -S(O)₀₋₂-, -NR⁸-, and absent; and e is 0 or 1; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
14. The compound according to claim 13, wherein Y is -CH₂-; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
15. The compound according to claim 14, wherein A is selected from -NR⁸-, wherein R⁸ is selected from -H, C₁-C₆alkyl, -CO₂R³, -C(O)N(R³)R⁴, -SO₂R⁴, and -C(O)R³; -O-; and absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
16. The compound according to claim 15, wherein




geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

17. The compound according to claim 16, wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
18. The compound according to claim 11, wherein said saturated bridged ring system is according to either formula V or formula VI;



wherein R^8 is selected from -H, C_1 - C_6 alkyl, $-CO_2R^3$, $-C(O)N(R^3)R^4$, $-SO_2R^4$, and $-C(O)R^3$; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.


19. The compound according to claim 18, wherein Y is either $-CH_2-$ or absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
20. The compound according to claim 19, wherein



R^{2b} , and R^{2c} are each independently selected from F, Cl, and Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

21. The compound according to claim 20, wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

22. The compound according to claim 21, wherein R^8 is methyl or ethyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
23. The compound according to claim 12, wherein said saturated bridged ring system is according to formula VII;




VII

- wherein A is selected from -O-, -S(O)₀₋₂-, -NR⁸-, -CR⁶R⁷-, and absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
24. The compound according to claim 23, wherein R^3 is selected from -H and alkyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
25. The compound according to claim 24 wherein A is either -C(R⁶)R⁷- or absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
26. The compound according to claim 25, wherein A is either -CH₂- or absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
27. The compound according to claim 26, wherein Y is -CH₂-; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

28. The compound according to claim 27, wherein q is 3; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

29. The compound according to claim 28, wherein





30. The compound according to claim 29, wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

31. The compound according to claim 9, wherein the saturated bridged ring system has a geometry selected from the group consisting of [3.3.1], [3.2.1], and [2.2.1]; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

32. The compound according to claim 31, wherein said saturated bridged ring system contains one or two annular nitrogens, said one or two annular nitrogens are selected from $-NR^8-$, when X^1 , and a bridgehead nitrogen, when X^2 ; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

33. The compound according to claim 32, wherein said saturated bridged ring system is according to formula VIII or formula IX;




VIII

IX

wherein R^8 is selected from -H, C_1 - C_6 alkyl, $-CO_2R^3$, $-C(O)N(R^3)R^4$, $-SO_2R^4$, and $-C(O)R^3$; and R^{26} is C_{1-3} alkyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.


34. The compound according to claim 33, wherein Y is $-CH_2CH_2-$; and E is either absent or $-N(R^9)-$; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
35. The compound according to claim 34, wherein q is 3; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
36. The compound according to claim 35, wherein



R^{2b} , and R^{2c} are each independently selected from F, Cl, and Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

37. The compound according to claim 36, wherein R^{2a} is F, R^{2b} is Cl, and R^{2c} is either Cl or Br; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
38. The compound according to claim 37, wherein R^8 is methyl or ethyl; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

39. A compound of Formula Ia,



or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof, wherein, q is 1, 2, or 3;

R^2 is selected from halogen, trihalomethyl, -CN, -NO₂, -OR³, and C₁-C₆alkyl;

Y is either:

-CH₂- or -CH₂CH₂- provided there are at least two carbon atoms between the oxygen at the 7-position of the quinazoline ring system of Ia and E when E is -NR⁹- or -O-; or

Y is absent; and when Y is absent, E is also absent;

E is selected from -NR⁹-, -O-, and absent;

R^3 is -H or R^4 ;

R^4 is selected from C₁-C₆alkyl; C₁-C₆alkyl substituted with one, two, or three halogen; aryl; aryl substituted with one, two, or three halogen; unsubstituted aryl C₁-C₆alkyl; heterocyclyl; and heterocyclyl C₁-C₆alkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form a five- to seven-membered heterocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P where the five- to seven-membered heterocyclyl is optionally substituted by one, two, or three alkyl;

R^9 is -H or C₁-C₆alkyl;

R^{10} is selected from -H, alkyl, and -OR¹³; and R^{11} and R^{12} are each independently selected from -H, -CF₃, -F, -N(R^3) R^4 , -N(C=O) R^3 , -N(R^3)SO₂ R^3 , -OR¹³ and -S(O)₀₋₂ R^{13} ; or

R^{10} is selected from -H, and -OR¹³; and R^{11} and R^{12} , when taken together, are oxo, exo-alkenyl, or when taken together with the carbon to which they are attached, form a three- to seven-membered spirocyclyl; and

R^{13} is selected from -H, $-C(=O)R^4$, C_2-C_6 alkynyl, unsubstituted aryl C_2-C_6 alkynyl, heterocyclyl C_2-C_6 alkynyl, C_2-C_6 alkenyl, unsubstituted aryl C_2-C_6 alkenyl, heterocyclyl C_2-C_6 alkenyl, C_1-C_6 alkyl, C_1-C_6 alkyl substituted with one, two, or three halogen, unsubstituted aryl C_1-C_6 alkyl, aryl, heterocyclyl C_1-C_6 alkyl optionally substituted with one alkyl, and heterocyclyl; or
 two R^{13} 's, when taken together, form 1) a corresponding spirocyclic ketal from R^{11} , R^{12} and the carbon to which they are attached, when R^{11} and R^{12} are both $-OR^{13}$, or 2) a corresponding cyclic ketal from R^{10} and one of R^{11} and R^{12} , and the corresponding carbons to which they are attached, when R^{10} is $-OR^{13}$, and at least one of R^{11} and R^{12} is also $-OR^{13}$.

40. The compound of claim 39 wherein

q is 1, 2, or 3;

R^2 is selected from halogen, trihalomethyl, -CN, $-NO_2$, $-OR^3$, and C_1-C_6 alkyl;

Y is either:

-CH₂- or -CH₂CH₂- provided there are at least two carbon atoms between the oxygen at the 7-position of the quinazoline ring system of Ia and E when E is $-NR^9$ - or -O-; or

Y is absent; and when Y is absent, E is also absent;

E is selected from $-NR^9$ -, -O-, and absent;

R^3 is -H or R^4 ;

R^4 is selected from C_1-C_6 alkyl; C_1-C_6 alkyl substituted with one, two, or three halogen; aryl; aryl substituted with one, two, or three halogen; unsubstituted aryl C_1-C_6 alkyl, heterocyclyl, and heterocyclyl C_1-C_6 alkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form a five- to seven-membered heterocyclyl optionally containing at least one additional heteroatom selected from N, O, S, and P where the five- to seven-membered heterocyclyl is optionally substituted by one, two, or three alkyl;

R^9 is -H or C_1-C_6 alkyl;

R^{10} is selected from -H, alkyl, and $-OR^{13}$; and R^{11} and R^{12} are each independently selected from -H, $-CF_3$, -F, $-N(R^3)R^4$, $-N(C=O)R^3$, $-N(R^3)SO_2R^3$, $-S(O)_{0-2}R^{13}$, and $-OR^{13}$; or

R^{10} is selected from -H, and $-OR^{13}$; and R^{11} and R^{12} , when taken together, are oxo, exo-alkenyl, or when taken together with the carbon to which they are attached, form a three- to seven-membered spirocyclyl; and

R^{13} is selected from -H, $-C(=O)R^4$, C_2-C_6 alkynyl, unsubstituted aryl C_2-C_6 alkynyl, heterocyclyl C_2-C_6 alkynyl, C_2-C_6 alkenyl, unsubstituted aryl C_2-C_6 alkenyl, heterocyclyl C_2-C_6 alkenyl, C_1-C_6 alkyl, C_1-C_6 alkyl substituted with one, two, or three halogen, unsubstituted aryl C_1-C_6 alkyl, aryl, heterocyclyl C_1-C_6 alkyl optionally substituted with one alkyl, and heterocyclyl; or

two R^{13} 's, when taken together, form 1) a corresponding spirocyclic ketal from R^{11} , R^{12} and the carbon to which they are attached, when R^{11} and R^{12} are both $-OR^{13}$, or 2) a corresponding cyclic ketal from R^{10} and one of R^{11} and R^{12} , and the corresponding carbons to which they are attached, when R^{10} is $-OR^{13}$, and at least one of R^{11} and R^{12} is also $-OR^{13}$; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

41. The compound according to claim 40, wherein Y is either $-CH_2-$ or absent; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
42. The compound according to claim 41, wherein one of R^{11} and R^{12} is $-OR^{13}$, wherein R^{13} is selected from -H, $-C(O)R^4$, C_1-C_6 alkyl, and C_1-C_6 alkyl substituted with one, two, or three halogen; and R^{10} and the other of R^{11} and R^{12} are both -H; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
43. The compound according to claim 41, wherein one of R^{11} and R^{12} is -F; and R^{10} and the other of R^{11} and R^{12} are both -H; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
44. The compound according to claim 41, wherein R^{13} is an alkyl group containing at least one fluorine substitution thereon; or a single geometric isomer, stereoisomer, racemate,

enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

45. The compound according to claim 41, wherein q is 2 or 3; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
46. The compound according to claim 45, wherein each R^2 is independently selected from -F, -Cl, -Br, -CF₃, -CH₃, and -OR²⁵; wherein R²⁵ is either methyl or aryl, each optionally substituted with one to three halogens; or a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
47. A compound selected from 1,4:3,6-dianhydro-2-*O*-[4-(methyloxy)carbonyl-2-(methyloxy)phenyl]-5-*O*-(methanesulfonyl)-D-glucitol; 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-{2-(methyloxy)-4-[(methyloxy)carbonyl]phenyl}-L-iditol; 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-{2-(methyloxy)-4-[(methyloxy)carbonyl]-5-nitrophenyl}-L-iditol; 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-{5-amino-2-(methyloxy)-4-[(methyloxy)carbonyl]-phenyl}-L-iditol; 1,4:3,6-dianhydro-2-deoxy-2-fluoro-5-*O*-[6-(methyloxy)-4-oxo-3,4-dihydroquinazolin-7-yl]-L-iditol; and 1,4:3,6-dianhydro-5-*O*-[4-chloro-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-L-iditol; and a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.
48. A compound selected from 1,1-dimethylethyl (3*R*,6*aS*)-5-(hydroxymethyl)hexahydrocyclopenta[*c*] pyrrole-2(1*H*)-carboxylate; 1,1-dimethylethyl (3*R*,6*aS*)-5-[[[(methanesulfonyl)oxy]methyl]hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate; (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl methanesulfonate; (3*S*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate; (3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate; hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl acetate; (3*S*,9*aS*)-3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine; (3*R*,9*aS*)-3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine; 3-(chloromethyl)hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazine; [(3*S*,8*aS*)-2-methyl-1-oxooctahydropyrrolo[1,2-*a*]pyrazin-3-yl]methyl

methanesulfonate; (3*S*,8*aS*)-3-(hydroxymethyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one; (3*S*,8*aS*)-3-(hydroxymethyl)-hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one; (3*S*,8*aS*)-3-({[(1,1-dimethylethyl)(dimethyl)silyl]oxy}methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one; (3*S*,8*aS*)-3-({[(1,1-dimethylethyl)(dimethyl)silyl]oxy}methyl) hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one; (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl methanesulfonate; (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethanol; (3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl acetate; (3*S*,8*aR*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine; (3*R*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine; (3*R*,8*aR*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine; and (3*S*,8*aS*)-3-(Chloromethyl)hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazine; and a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

49. The compound of claim 2 selected from

<i>N</i> -(3,4-dichlorophenyl)-6-(methyloxy)-7-{{[(8 <i>aR</i>)-tetrahydro-1 <i>H</i> -[1,3]thiazolo[4,3- <i>c</i>][1,4]oxazin-6-ylmethyl]oxy}quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-6-(methyloxy)-7-[(tetrahydro-1 <i>H</i> -[1,3]thiazolo[4,3- <i>c</i>][1,4]oxazin-3-ylmethyl)oxy]quinazolin-4-amine
3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2- <i>O</i> -(1-methylethylidene)-beta-L-xylo-hexofuranose;
3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-1,2- <i>O</i> -(1-methylethylidene)-β-D-idofuranose;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-[(octahydro-2 <i>H</i> -quinolizin-3-ylmethyl)oxy]quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chlorophenyl)-7-{{[(3 <i>a'S</i> ,4 <i>R</i> ,6' <i>S</i> ,6 <i>a'R</i>)-2,2-dimethyltetrahydrospiro[1,3-dioxolane-4,3'-furo[3,2- <i>b</i>]furan]-6'-yl]oxy}-6-(methyloxy)quinazolin-4-amine;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -(methylsulfonyl)-L-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -(methylsulfonyl)-D-glucitol;
2-amino-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-D-iditol;

2-amino-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-L-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-{[3,4-dichloro-2-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-{[3,4-dichloro-2-(4-methylpiperazin-1-yl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-fluoro-L-iditol; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

50. The compound of claim 15 selected from

<i>N</i> -(4-bromo-2,3-dichlorophenyl)-7-{[(3 <i>R</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4,5-dichloro-2-fluorophenyl)-7-{[(3 <i>R</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-5-chloro-2-fluorophenyl)-7-{[(3 <i>R</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3-chloro-2,4-difluorophenyl)-7-{[(3 <i>R</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3-chloro-2,4-difluorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-[(hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy]-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4,5-dichloro-2-fluorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-2,3-dichlorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-5-chloro-2-fluorophenyl)-7-{[(3 <i>S</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-{[(3 <i>R</i> ,9 <i>aS</i>)-hexahydro-1 <i>H</i> -[1,4]oxazino[3,4- <i>c</i>][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(4-bromo-3-chloro-2-fluorophenyl)-7-{[(3*R*,9*aS*)-hexahydro-1*H*-[1,4]oxazino[3,4-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine; and

a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

51. The compound of claim 15 selected from

N-(3,4-dichlorophenyl)-7-{[(3*R*,8*aR*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(4-bromo-5-chloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(3,4-dichlorophenyl)-7-{[(3*S*,8*aR*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(3,4-dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(3,4-dichlorophenyl)-7-{[(3*R*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(3,4-dichloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(4-bromo-3-chloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(3-chloro-2,4-difluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(4-bromo-2,3-dichlorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine;

N-(4,5-dichloro-2-fluorophenyl)-7-{[(3*S*,8*aS*)-hexahydro-1*H*-pyrrolo[2,1-*c*][1,4]oxazin-3-ylmethyl]oxy}-6-(methyloxy)quinazolin-4-amine; and

a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

52. The compound of claim 19 selected from

N-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,6*aS*)-2-(1-methylethyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;

N-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,6*aS*)-2-(1-methylethyl)octahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;

7-({[(3aR,6aS)-2-acetyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy}quinazolin-4-amine;
N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-2-(methylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-ethyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
N-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(3-chloro-2,4-difluorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4,5-dichloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-2,3-dichlorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(3,4-dichlorophenyl)-7-({[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3aR,6aS)-2-ethyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
N-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3aR,6aS)-2-(2-methylpropyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

53. The compound of claim 19 selected from

N-(3-chloro-2,4-difluorophenyl)-7-({[(3aR,5s,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;

<i>N</i> -(3-chloro-2,4-difluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta- [c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-2,3-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2- methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4- amine;
<i>N</i> -(4-bromo-2,3-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclo- penta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[c]pyrrol-5- yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[c]pyrrol-5- yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-{{[(3 <i>aR</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]oxy}- 6-(methyloxy)quinazolin-4-amine; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

54. The compound of claim 21 selected from

<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(1- methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4- amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(1- methylethyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4- amine;
7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-acetyloctahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)- <i>N</i> -(4-bromo- 3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)- octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy} quinazolin-4-amine;
ethyl (3 <i>aR</i> ,6 <i>aS</i>)-5-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6- (methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[c]pyrrole-2(1 <i>H</i>)- carboxylate;
ethyl (3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-5-[[{4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6- (methyloxy)quinazolin-7-yl}oxy)methyl]hexahydrocyclopenta[c]pyrrole-2(1 <i>H</i>)- carboxylate;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2- (methylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-ethyloctahydrocyclopenta[c]pyrrol- 5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;

<i>N</i> -(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(2-methylpropyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4,5-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4,5-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(2-methylpropyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
1,1-dimethylethyl (3 <i>aR</i> ,6 <i>aS</i>)-5-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyl-oxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[<i>c</i>]pyrrole-2(1 <i>H</i>)-carboxylate;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
1,1-dimethylethyl (3 <i>aR</i> ,6 <i>aS</i>)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl) hexahydrocyclopenta-[<i>c</i>]pyrrole-2(1 <i>H</i>)-carboxylate; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

55. The compound of claim 19 selected from

<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(1-methylethyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(1-methylethyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-acetyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)- <i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
ethyl (3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-5-[(4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl]hexahydrocyclopenta[<i>c</i>]pyrrole-2(1 <i>H</i>)-carboxylate;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(methylsulfonyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-ethyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(2-methylpropyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>s</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3-chloro-2,4-difluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4,5-dichloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-5-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-2,3-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-methyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;

<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-ethyloctahydrocyclopenta[<i>c</i>]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-({[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-2-(2-methylpropyl)octahydrocyclopenta[<i>c</i>]pyrrol-5-yl)methyl}oxy)quinazolin-4-amine; and
1,1-dimethylethyl (3 <i>aR</i> ,6 <i>aS</i>)-5-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyl-oxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[<i>c</i>]pyrrole-2(1 <i>H</i>)-carboxylate;
<i>N</i> -(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3 <i>aR</i> ,5 <i>r</i> ,6 <i>aS</i>)-octahydrocyclopenta[<i>c</i>]pyrrol-5-yl)methyl]oxy}quinazolin-4-amine
1,1-dimethylethyl (3 <i>aR</i> ,6 <i>aS</i>)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl) hexahydrocyclopenta-[<i>c</i>]pyrrole-2(1 <i>H</i>)-carboxylate; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

56. The compound of claim 55 selected from *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; *N*-(4-bromo-3-chloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; and *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*s*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine; and optionally as a pharmaceutically acceptable salt or hydrate thereof.
57. A pharmaceutical composition comprising a compound of claim 56 and a pharmaceutically acceptable carrier.
58. The compound of claim 55 selected from 1,1-dimethylethyl (3*aR*,6*aS*)-5-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyl-oxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate; *N*-(4-bromo-3-chloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3*aR*,5*r*,6*aS*)-octahydrocyclopenta[*c*]pyrrol-5-yl)methyl]oxy}quinazolin-4-amine; *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3*aR*,5*r*,6*aS*)-octahydrocyclopenta[*c*]pyrrol-5-yl)methyl]oxy}quinazolin-4-amine; 1,1-dimethylethyl (3*aR*,6*aS*)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl) hexahydrocyclopenta-[*c*]pyrrole-2(1*H*)-carboxylate; and a single geometric isomer, stereoisomer, racemate, enantiomer, or

diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

59. The compound of claim 56 named *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta[*c*]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine optionally as a pharmaceutically acceptable salt or hydrate thereof.
60. A pharmaceutical composition comprising a compound of claim 59 and a pharmaceutically acceptable carrier.
61. The compound of claim 30 selected from

(3 <i>S</i> ,9 <i>aS</i>)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2 <i>H</i> -pyrido[1,2- <i>a</i>]pyrazin-1(6 <i>H</i>)-one;
(3 <i>S</i> ,9 <i>aR</i>)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydro-2 <i>H</i> -pyrido[1,2- <i>a</i>]pyrazin-1(6 <i>H</i>)-one;
(3 <i>S</i> ,8 <i>aS</i>)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2- <i>a</i>]pyrazin-1(2 <i>H</i>)-one;
(3 <i>S</i> ,8 <i>aR</i>)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2- <i>a</i>]pyrazin-1(2 <i>H</i>)-one;
(3 <i>S</i> ,8 <i>aS</i>)-3-({[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydropyrrolo[1,2- <i>a</i>]pyrazin-1(2 <i>H</i>)-one;
(3 <i>S</i> ,8 <i>aS</i>)-3-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)-2-methylhexahydropyrrolo[1,2- <i>a</i>]pyrazin-1(2 <i>H</i>)-one; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

62. The compound of claim 33 selected from

<i>N</i> -(3,4-dichlorophenyl)-7-({2-[(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)amino]ethyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-[(2-{[(3- <i>endo</i>)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]amino}ethyl)oxy]-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-{[2-(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)ethyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({2-[(3- <i>endo</i>)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)-6-(methyloxy)quinazolin-4-amine;

<i>N</i> -(3,4-dichlorophenyl)-7-{[(8-methyl-8-azabicyclo[3.2.1]oct-3-yl)methyl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({[(3- <i>endo</i>)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-{[8-methyl-8-azabicyclo[3.2.1]oct-3-yl]oxy}-6-(methyloxy)quinazolin-4-amine;
<i>N</i> -(3,4-dichlorophenyl)-7-({[(3- <i>exo</i>)-8-methyl-8-azabicyclo[3.2.1]oct-3-yl]oxy}-6-(methyloxy)quinazolin-4-amine;
7-{[(3- <i>endo</i>)-8-azabicyclo[3.2.1]oct-3-ylmethyl]oxy}- <i>N</i> -(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine;
1,1-dimethylethyl (3- <i>endo</i>)-3-(2-{[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}ethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate; and
7-({2-[(3- <i>endo</i>)-8-azabicyclo[3.2.1]oct-3-yl]ethyl}oxy)- <i>N</i> -(3,4-dichlorophenyl)-6-(methyloxy)quinazolin-4-amine; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

63. The compound of claim 43 selected from

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(2-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -[4-[(2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -[4-[(4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(2-bromophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-bromophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol; and

a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

64. The compound of claim 46 selected from

1,4:3,6-dianhydro-5-([4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol;
1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2-O-methyl-D-glucitol;
1,4:3,6-dianhydro-2-deoxy-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-O-methyl-D-threo-hexitol;
1,4:3,6-dianhydro-5-deoxy-5-([4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-methyl-5-O-{6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl}-L-iditol;
1,4:3,6-dianhydro-5-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-O-methyl-D-glucitol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]- <i>L</i> -sorbose ethylene glycol acetal;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -(difluoromethyl)- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -ethyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(3-bromo-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>L</i> -iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy- <i>D</i> -xylo-hexitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl- <i>D</i> -glucitol;
methyl 3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2- <i>O</i> -methyl- α - <i>L</i> -idofuranoside;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-5-methylidene- <i>D</i> -xylo-hexitol;
methyl 3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2- <i>O</i> -methyl- β - <i>L</i> -idofuranoside;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -{6-(methyloxy)-4-[(2,3,4-trifluorophenyl)amino]quinazolin-7-yl}- <i>D</i> -iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(2-chloro-4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro- <i>D</i> -iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(2-bromo-4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,6-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -[4-{[4-fluoro-3-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,5-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,3-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(3,5-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-chloro-4-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-2-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -{6-(methyloxy)-4-[(2,4,5-trifluorophenyl)amino]quinazolin-7-yl}-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -{6-(methyloxy)-4-[(2,4,6-trifluorophenyl)amino]quinazolin-7-yl}-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-({4-[(4-chlorophenyl)oxy]-3,5-difluorophenyl}amino)-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chloro-5-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -{6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl}-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -{6-(methyloxy)-4-[(3,4,5-trichlorophenyl)amino]quinazolin-7-yl}-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(3,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(2,5-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(2-bromo-4,6-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[4-chloro-3-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[2-chloro-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -[4-{[2-fluoro-3-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[2-bromo-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-{[2-bromo-4-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-2-fluoro-5- <i>O</i> -[4-{[4-fluoro-2-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[3-bromo-5-(trifluoromethyl)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-bromo-4-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(5-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-2-deoxy-5- <i>O</i> -[4-[(3,5-dimethylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[2,5-bis(methyloxy)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[5-chloro-2,4-bis(methyloxy)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-{[4-chloro-2,5-bis(methyloxy)phenyl]amino}-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-fluoro-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -{4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl}-2-deoxy-2-fluoro-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-5- <i>C</i> -[(methyloxy)methyl]-L-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>S</i> -methyl-5-thio-D-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>S</i> -methyl-5-thio-L-iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-morpholin-4-yl-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-morpholin-4-yl-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(4-methylpiperazin-1-yl)-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(4-methylpiperazin-1-yl)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-pyrrolidin-1-yl-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-pyrrolidin-1-yl-L-iditol;
2- <i>O</i> -acetyl-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol;
2- <i>O</i> -acetyl-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(methylsulfonyl)-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(methylsulfonyl)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(dimethylamino)-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(dimethylamino)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(diethylamino)-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(diethylamino)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-piperidin-1-yl-D-iditol;

1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-piperidin-1-yl-L-iditol;
2-(acetylamino)-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-D-iditol;
2-(acetylamino)-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-5- <i>C</i> -(trifluoromethyl)-L-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-5- <i>C</i> -(trifluoromethyl)-D-glucitol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-[(methanesulfonyl)amino]-D-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-[(methanesulfonyl)amino]-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>C</i> -(trifluoromethyl)-D-glucitol;
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

65. The compound of claim 64 selected from

1,4:3,6-dianhydro-5-([4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2- <i>O</i> -methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2- <i>O</i> -methyl-D-glucitol;
1,4:3,6-dianhydro-5-deoxy-5-([4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2- <i>O</i> -methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2- <i>O</i> -methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2- <i>O</i> -methyl-D-xylo-hexitol;
1,4:3,6-dianhydro-5-([4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5-deoxy-2- <i>O</i> -methyl-D-glucitol;
1,4:3,6-dianhydro-2-deoxy-2-([4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-5- <i>O</i> -methyl-D-threo-hexitol;

1,4:3,6-dianhydro-5-deoxy-5-([4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy)methyl)-2-O-methyl-D-glucitol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-5-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-methyl-5-O-{6-(methyloxy)-4-[(2,3,4-trichlorophenyl)amino]quinazolin-7-yl}-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-2,3-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-5-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-sorbose ethylene glycol acetal;
1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2,4-difluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4,5-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-(difluoromethyl)-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-ethyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3-bromo-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(3-chloro-2-methylphenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-O-methyl-L-iditol;
1,4:3,6-dianhydro-2-O-[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-D-xylo-hexitol;

1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-D-glucitol;
methyl 3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2- <i>O</i> -methyl- α -L-idofuranoside;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-deoxy-5-methylidene-D-xylo-hexitol;
methyl 3,6-anhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2- <i>O</i> -methyl- β -L-idofuranoside;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-C-(trifluoromethyl)-D-glucitol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(3,4-dichlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2- <i>O</i> -methyl-D-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-D-glucitol;
1,4:3,6-dianhydro-5- <i>O</i> -{4-[(3-chloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl}-2-deoxy-2-fluoro-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-D-glucitol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5-S-methyl-5-thio-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-morpholin-4-yl-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(4-methylpiperazin-1-yl)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-pyrrolidin-1-yl-L-iditol;
2- <i>O</i> -acetyl-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(methylsulfonyl)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(dimethylamino)-L-iditol;


1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-(diethylamino)-L-iditol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-piperidin-1-yl-L-iditol;
2-(acetylamino)-1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-L-iditol;
1,4:3,6-dianhydro-2- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-5- <i>O</i> -methyl-5- <i>C</i> -(trifluoromethyl)-D-glucitol;
1,4:3,6-dianhydro-5- <i>O</i> -[4-[(4-bromo-3-chlorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]-2-deoxy-2-[(methylsulfonyl)amino]-L-iditol; and
a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof and optionally as a pharmaceutically acceptable salt or hydrate thereof.

66. The compound of claim 48 selected from 1,1-dimethylethyl (3*aR*,6*aS*)-5-(hydroxymethyl)hexahydro cyclopenta[*c*] pyrrole-2(1*H*)-carboxylate; 1,1-dimethylethyl (3*aR*,6*aS*)-5- {[(methylsulfonyl)oxy]methyl}hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate; and a single geometric isomer, stereoisomer, racemate, enantiomer, or diastereomer, thereof; and optionally as a pharmaceutically acceptable salt or hydrate thereof.
67. A pharmaceutical composition comprising a compound of any one of claims 1-56, 58 and 61-65 and a pharmaceutically acceptable carrier.
68. Use of a compound according to any one of claims 1-56, 58 and 61-65 optionally together with a pharmaceutically acceptable carrier for modulating the *in vivo* activity of a kinase selected from ephrin and EGFR.
69. Use of a compound according to any one of claims 1-56, 58 and 61-65 optionally together with a pharmaceutically acceptable carrier for treating a disease or a disorder associated with abnormal cellular activities.
70. The use of claim 69 where the disease is cancer.
71. The use of claim 70 where the cancer is modulated by one or more kinases selected from ephrin, KDR, Flt-1, EGFR, and ErbB2.

72. The use of claim 70 where the cancer is selected from non-small cell lung cancer, glioblastoma, pancreatic cancer, cancer of the nervous system, cancer of the large bowel, multiple myeloma, undifferentiated small cell bronchogenic carcinoma, gastrointestinal cancer, esophageal cancer, malignant melanoma, neuroblastoma, osteosarcoma, ovarian cancer, endometrial cancer, cervical cancer, bladder cancer, urethral cancer, and prostate cancer.
73. The use of claim 70 where the cancer is selected from non-small cell lung cancer, glioblastoma, pancreatic cancer, cancer of the nervous system, cancer of the large bowel, neuroblastoma, and gastrointestinal cancer.
74. The use of claim 70 where the cancer is selected from ovarian cancer, cervical cancer, bladder cancer, esophageal cancer, malignant melanoma, and prostate cancer.
75. The use of claim 70 where the cancer is non-small cell lung cancer.
76. The use of claim 70 where the cancer is glioblastoma.
77. The use of claim 72 where the gastrointestinal cancer is stomach cancer.
78. The use of claim 69 where the disease is selected from diabetic retinopathy, psoriasis and rheumatoid arthritis.
79. Use of a compound according to claim 59 optionally together with a pharmaceutically acceptable carrier for modulating the *in vivo* activity of a kinase selected from ephrin and EGFR.
80. Use of a compound according to claim 59 optionally together with a pharmaceutically acceptable carrier for treating a disease or a disorder associated with abnormal cellular activities.
81. The use of claim 80 where the disease is cancer.
82. The use of claim 81 where the cancer is modulated by one or more kinases selected from ephrin, KDR, Flt-1, EGFR, and ErbB2.
83. The use of claim 81 where the cancer is selected from non-small cell lung cancer, glioblastoma, pancreatic cancer, cancer of the nervous system, cancer of the large bowel, multiple myeloma, undifferentiated small cell bronchogenic carcinoma, gastrointestinal

cancer, esophageal cancer, malignant melanoma, neuroblastoma, osteosarcoma, ovarian cancer, endometrial cancer, cervical cancer, bladder cancer, urethral cancer, and prostate cancer.

84. The use of claim 81 where the cancer is selected from non-small cell lung cancer, glioblastoma, pancreatic cancer, cancer of the nervous system, cancer of the large bowel, neuroblastoma, and gastrointestinal cancer.
85. The use of claim 81 where the cancer is selected from ovarian cancer, cervical cancer, bladder cancer, esophageal cancer, malignant melanoma, and prostate cancer.
86. The use of claim 81 where the cancer is non-small cell lung cancer.
87. The use of claim 81 where the cancer is glioblastoma.
88. The use of claim 83 where the gastrointestinal cancer is stomach cancer.
89. The use of claim 80 where the disease is selected from diabetic retinopathy, psoriasis and rheumatoid arthritis.
90. A process for preparing a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein


P is methyl;

Ar is 3,4-dichloro-2-fluorophenyl;

Z is -NH-;


E² is selected from [(3a*R*,5*r*,6a*S*)-2-methyloctahydrocyclopenta-[*c*]pyrrol-5-yl]methyl
and [(3a*R*,5*s*,6a*S*)-2-methyloctahydrocyclopenta-[*c*]pyrrol-5-yl]methyl;

where the process comprises introducing a group E^2 as defined above for Formula 8, into a compound of formula 7



wherein P, Ar and Z are as defined above for Formula 8.

91. A process according to claim 90, where the compound of formula 8 is



or a pharmaceutically acceptable salt thereof.

92. A process according to claim 90, where the compound of formula 8 is *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*s*,6*aS*)-2-methyloctahydrocyclopenta-*c*]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine or a pharmaceutically acceptable salt thereof.
93. A process according to claim 90, where the compound of formula 8 is *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3*aR*,5*r*,6*aS*)-2-methyloctahydrocyclopenta-*c*]pyrrol-5-yl)methyl}oxy)-6-(methyloxy)quinazolin-4-amine or a pharmaceutically acceptable salt thereof.
94. A process according to claim 93, wherein the introducing is carried out in *N,N*-dimethylacetamide.
95. A process according to claim 93 or claim 94, wherein the introducing is carried out in the presence of potassium carbonate.

96. A process according to any of claims 93 to 95, wherein E² is introduced by reacting 1,1-dimethylethyl (3aR,6aS)-5-{{[(methylsulfonyl)oxy]methyl}hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate with the compound of formula 7 to provide 1,1-dimethylethyl (3aR,6aS)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta-[c]pyrrol-2(1H)-carboxylate.
97. A process according to claim 96, further comprising the step of deprotecting the 1,1-dimethylethyl (3aR,6aS)-5-({[4-[(3,4-dichloro-2-fluorophenyl)amino]-6-(methyloxy)quinazolin-7-yl]oxy}methyl)hexahydrocyclopenta-[c]pyrrol-2(1H)-carboxylate to provide *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3aR,5r,6aS)-octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy}-quinazolin-4-amine.
98. A process according to claim 97, further comprising the step of methylating the *N*-(3,4-dichloro-2-fluorophenyl)-6-(methyloxy)-7-{{[(3aR,5r,6aS)-octahydrocyclopenta[c]pyrrol-5-yl]methyl}oxy}-quinazolin-4-amine to provide *N*-(3,4-dichloro-2-fluorophenyl)-7-({[(3aR,5r,6aS)-2-methyloctahydrocyclopenta-[c]pyrrol-5-yl]methyl}oxy)-6-(methyloxy)quinazolin-4-amine.

