

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number  
**WO 2004/015071 A2**

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/US2003/024551
- (22) International Filing Date: 6 August 2003 (06.08.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/401,739 7 August 2002 (07.08.2002) US
- (71) Applicant (for all designated States except US): **EX-ELIXIS, INC.** [US/US]; P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **FRANCIS-LANG, Helen** [GB/US]; 1782 Pacific Avenue, Apt. 2, San Francisco, CA 94109 (US). **FRIEDMAN, Lori** [US/US]; 113 Arundel Road, San Carlos, CA 94070 (US). **KIDD, Thomas** [GB/US]; 643 Lake Street, San Francisco, CA 94118 (US). **ROCHE, Siobhan** [IE/IE]; 30 Moatfield Park, Coolock, Dublin 5 (IE). **ZHANG, HaiGuang** [US/US]; 4833 El Grande Place, El Sobrante, CA 94803 (US).
- (74) Agents: **SHAYESTEH, Laleh** et al.; Exelixis, Inc., P. O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 2004/015071 A2

(54) Title: CSNK1GS AS MODIFIERS OF THE p21 PATHWAY AND METHODS OF USE

(57) Abstract: Human CSNK1G genes are identified as modulators of the p21 pathway, and thus are therapeutic targets for disorders associated with defective p21 function. Methods for identifying modulators of p21, comprising screening for agents that modulate the activity of CSNK1G are provided.

**CSNK1GS AS MODIFIERS OF THE p21 PATHWAY AND METHODS OF USE****REFERENCE TO RELATED APPLICATIONS**

This application claims priority to U.S. provisional patent application US  
5 60/401,739 filed 8/7/2002. The content of the prior application is hereby incorporated in  
its entirety.

**BACKGROUND OF THE INVENTION**

The p21/CDKN1/WAF1/CIP1 protein(El-Deiry, W. S.; et al. Cell 75: 817-825,  
10 1993; Harper, J. W.; et al. Cell 75: 805-816, 1993; Huppi, Ket al. Oncogene 9: 3017-3020,  
1994) is a cell cycle control protein that inhibits cyclin-kinase activity, is tightly regulated  
at the transcriptional level by p53, and mediates p53 suppression of tumor cell growth.  
Along with p53, p21 appears to be essential for maintaining the G2 checkpoint in human  
cells (Bunz, F.; Dutriau, A.; et al. Science 282:1497-1501, 1998). Sequences of P21 are  
15 well-conserved throughout evolution, and have been identified in species as diverse as  
human ( Genbank Identifier 13643057), *Drosophila melanogaster* (GI# 1684911),  
*Caenorhabditis elegans* (GI#4966283), and yeast (GI#2656016).

Casein kinase I is the most abundant serine/threonine kinase in eukaryotic cell  
extracts, and preferentially phosphorylate acidic substrates using ATP as a phosphate  
20 donor. Multiple isoforms of the enzyme exist. The gamma-1 isoform (CSNK1G1) is  
involved in growth and morphogenesis of eukaryotic cells (Kusuda, J., et al (2000)  
Cytogenet Cell Genet 90:298-302), the gamma-2 (CSNK1G2) and gamma-3 (CSNK1G3)  
isoforms may play a role in signal transduction (Kitabayashi, A. N., et al. (1997)  
Genomics 46, 133-7; Kusuda, J., et al (1998) Cytogenet Cell Genet 83:101-3).

25 The ability to manipulate the genomes of model organisms such as *Drosophila*  
provides a powerful means to analyze biochemical processes that, due to significant  
evolutionary conservation, have direct relevance to more complex vertebrate organisms.  
Due to a high level of gene and pathway conservation, the strong similarity of cellular  
processes, and the functional conservation of genes between these model organisms and  
30 mammals, identification of the involvement of novel genes in particular pathways and  
their functions in such model organisms can directly contribute to the understanding of the  
correlative pathways and methods of modulating them in mammals (see, for example,  
Mechler BM et al., 1985 EMBO J 4:1551-1557; Gateff E. 1982 Adv. Cancer Res. 37: 33-  
74; Watson KL., et al., 1994 J Cell Sci. 18: 19-33; Miklos GL, and Rubin GM. 1996 Cell

86:521-529; Wassarman DA, et al., 1995 *Curr Opin Gen Dev* 5: 44-50; and Booth DR. 1999 *Cancer Metastasis Rev.* 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a “genetic entry point”) that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a “modifier” involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as p21, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

## SUMMARY OF THE INVENTION

We have discovered genes that modify the p21 pathway in *Drosophila*, and identified their human orthologs, hereinafter referred to as casein kinase I gamma isoforms (CSNK1G). The invention provides methods for utilizing these p21 modifier genes and polypeptides to identify CSNK1G-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired p21 function and/or CSNK1G function. Preferred CSNK1G-modulating agents specifically bind to CSNK1G polypeptides and restore p21 function. Other preferred CSNK1G-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress CSNK1G gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

CSNK1G modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with a CSNK1G polypeptide or nucleic acid. In one embodiment, candidate CSNK1G modulating agents are tested with an assay system comprising a CSNK1G polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate p21 modulating agents. The assay system may be cell-based or cell-free. CSNK1G-modulating agents include CSNK1G related proteins (e.g. dominant negative mutants, and biotherapeutics); CSNK1G -specific antibodies; CSNK1G -specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with CSNK1G or

compete with CSNK1G binding partner (e.g. by binding to a CSNK1G binding partner). In one specific embodiment, a small molecule modulator is identified using a kinase assay. In specific embodiments, the screening assay system is selected from a binding assay, an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction  
5 assay.

In another embodiment, candidate p21 pathway modulating agents are further tested using a second assay system that detects changes in the p21 pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may  
10 use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the p21 pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the CSNK1G function and/or the p21 pathway in a mammalian cell by contacting the mammalian cell with an  
15 agent that specifically binds a CSNK1G polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated the p21 pathway.

20

### DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the p21 pathway in *Drosophila*, where a dominant loss of function screen was carried out to identify genes that interact with the cyclin dependent kinase inhibitor, p21 (Bourne HR, et al., Nature  
25 (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95). Expression of the p21 gene in the eye causes deterioration of normal eye morphology. Modifiers of the eye phenotype were identified as members of the p21 pathway. The GISH (CG6963) gene was identified as a modifier of the p21 pathway. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, CSNK1G genes (i.e., nucleic acids  
30 and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective p21 signaling pathway, such as cancer.

In vitro and in vivo methods of assessing CSNK1G function are provided herein. Modulation of the CSNK1G or their respective binding partners is useful for understanding the association of the p21 pathway and its members in normal and disease

conditions and for developing diagnostics and therapeutic modalities for p21 related pathologies. CSNK1G-modulating agents that act by inhibiting or enhancing CSNK1G expression, directly or indirectly, for example, by affecting a CSNK1G function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided  
5 herein. CSNK1G modulating agents are useful in diagnosis, therapy and pharmaceutical development.

### **Nucleic acids and polypeptides of the invention**

Sequences related to CSNK1G nucleic acids and polypeptides that can be used in  
10 the invention are disclosed in Genbank (referenced by Genbank identifier (GI) number) as GI#s 11545750 (SEQ ID NO:1), 16878051 (SEQ ID NO:2), 10439939 (SEQ ID NO:3), 21733126 (SEQ ID NO:4), 27375099 (SEQ ID NO:5), 21314777 (SEQ ID NO:6), 18088089 (SEQ ID NO:7), 2199528 (SEQ ID NO:8), 9956036 (SEQ ID NO:9), 4758079 (SEQ ID NO:10), 4590041 (SEQ ID NO:11), and 28837292 (SEQ ID NO:12) for nucleic  
15 acid, and GI#s 11545751 (SEQ ID NO:13), 5579454 (SEQ ID NO:14), 21314778 (SEQ ID NO:15), and 4758080 (SEQ ID NO:16) for polypeptides.

The term "CSNK1G polypeptide" refers to a full-length CSNK1G protein or a functionally active fragment or derivative thereof. A "functionally active" CSNK1G fragment or derivative exhibits one or more functional activities associated with a full-  
20 length, wild-type CSNK1G protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of CSNK1G proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one  
25 embodiment, a functionally active CSNK1G polypeptide is a CSNK1G derivative capable of rescuing defective endogenous CSNK1G activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of a CSNK1G, such as a kinase domain or a binding domain.  
30 Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). For example, the kinase domain (PFAM 00069) of CSNK1G from GI#s 11545751, 21314778, and 4758080 (SEQ ID NOs:13, 15, and 16, respectively) is located respectively at approximately amino acid residues 44-312, 46-312, and 43-308. Methods for obtaining CSNK1G polypeptides are also further described below. In some

embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of any one of SEQ ID NOs:13-16 (a CSNK1G). In further preferred embodiments, the fragment comprises the entire kinase  
5 (functionally active) domain.

The term "CSNK1G nucleic acid" refers to a DNA or RNA molecule that encodes a CSNK1G polypeptide. Preferably, the CSNK1G polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more  
10 preferably 90%, and most preferably at least 95% sequence identity with human CSNK1G. Methods of identifying orthologs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are  
15 assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight  
20 conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by  
25 ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Drosophila*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion  
30 of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all

the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, *Advances in Applied Mathematics* 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, *J. of Molec.Biol.*, 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" ([www.psc.edu](http://www.psc.edu)) and references cited therein.; W.R. Pearson, 1991, *Genomics* 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 *Nucl. Acids Res.* 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of any of SEQ ID NOs:1-12. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing.

Conditions routinely used are set out in readily available procedure texts (*e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of any one of SEQ ID NOs:1-12 under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

### **Isolation, Production, Expression, and Mis-expression of CSNK1G Nucleic Acids and Polypeptides**

CSNK1G nucleic acids and polypeptides are useful for identifying and testing agents that modulate CSNK1G function and for other applications related to the involvement of CSNK1G in the p21 pathway. CSNK1G nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the

particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of a CSNK1G protein for assays used to assess CSNK1G function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (*e.g.*, Higgins SJ and Hames BD (eds.) *Protein Expression: A Practical Approach*, Oxford University Press Inc., New York 1999; Stanbury PF et al., *Principles of Fermentation Technology*, 2<sup>nd</sup> edition, Elsevier Science, New York, 1995; Doonan S (ed.) *Protein Purification Protocols*, Humana Press, New Jersey, 1996; Coligan JE et al, *Current Protocols in Protein Science* (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant CSNK1G is expressed in a cell line known to have defective p21 function, such as HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding a CSNK1G polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native CSNK1G gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the CSNK1G gene product, the expression vector can comprise a promoter operably linked to a CSNK1G gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the CSNK1G gene product based on the

physical or functional properties of the CSNK1G protein in *in vitro* assay systems (*e.g.* immunoassays).

The CSNK1G protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the CSNK1G gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native CSNK1G proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of CSNK1G or other genes associated with the p21 pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

### 25 **Genetically modified animals**

Animal models that have been genetically modified to alter CSNK1G expression may be used in *in vivo* assays to test for activity of a candidate p21 modulating agent, or to further assess the role of CSNK1G in a p21 pathway process such as apoptosis or cell proliferation. Preferably, the altered CSNK1G expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal CSNK1G expression. The genetically modified animal may additionally have altered p21 expression (*e.g.* p21 knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include

zebrafish, *C. elegans*, and *Drosophila*. Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, i.e. mosaic animals (see, for example, techniques described by Jakobovits, 1994, *Curr. Biol.* 4:761-763.) or stably integrated into its germ line DNA (i.e., in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster et al., *Proc. Nat. Acad. Sci. USA* 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Pat. No. 4,873,191 by Wagner et al., and Hogan, B., *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, *Science* (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) *Nature* 402:370-371; for transgenic Zebrafish see Lin S., *Transgenic Zebrafish*, *Methods Mol Biol.* (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, *Experientia* (1991) 47:897-905; for transgenic rats see Hammer *et al.*, *Cell* (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., *Teratocarcinomas and Embryonic Stem Cells, A Practical Approach*, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) *Nature* 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous CSNK1G gene that results in a decrease of CSNK1G function, preferably such that CSNK1G expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse CSNK1G gene is

used to construct a homologous recombination vector suitable for altering an endogenous CSNK1G gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH *et al.*, (1994) Scan J Immunol 40:257-264; Declerck PJ *et al.*, (1995) J Biol Chem. 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the CSNK1G gene, e.g., by introduction of additional copies of CSNK1G, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the CSNK1G gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the *cre/loxP* recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the *FLP* recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X *et al.* (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the p21 pathway, as animal models of disease and disorders implicating defective p21 function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in

5 screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered CSNK1G function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered CSNK1G expression that receive candidate therapeutic agent.

10 In addition to the above-described genetically modified animals having altered CSNK1G function, animal models having defective p21 function (and otherwise normal CSNK1G function), can be used in the methods of the present invention. For example, a p21 knockout mouse can be used to assess, *in vivo*, the activity of a candidate p21 modulating agent identified in one of the *in vitro* assays described below. p21 knockout mice are described in the literature (Umanoff H, et al., Proc Natl Acad Sci U S A 1995 Feb 28;92(5):1709-13). Preferably, the candidate p21 modulating agent when administered to a model system with cells defective in p21 function, produces a detectable phenotypic change in the model system indicating that the p21 function is restored, i.e., the cells exhibit normal cell cycle progression.

### Modulating Agents

20 The invention provides methods to identify agents that interact with and/or modulate the function of CSNK1G and/or the p21 pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the p21 pathway, as well as in further analysis of the CSNK1G protein and its contribution to the p21 pathway. Accordingly, the invention also provides methods for modulating the p21 pathway comprising the step of specifically modulating CSNK1G activity by administering a CSNK1G-interacting or -modulating agent.

25 As used herein, an "CSNK1G-modulating agent" is any agent that modulates CSNK1G function, for example, an agent that interacts with CSNK1G to inhibit or enhance CSNK1G activity or otherwise affect normal CSNK1G function. CSNK1G function can be affected at any level, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the CSNK1G - modulating agent specifically modulates the function of the CSNK1G. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the CSNK1G polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the CSNK1G. These

phrases also encompass modulating agents that alter the interaction of the CSNK1G with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of a CSNK1G, or to a protein/binding partner complex, and altering CSNK1G function). In a further preferred embodiment, the CSNK1G- modulating agent is a modulator of the p21 pathway (e.g. it restores and/or upregulates p21 function) and thus is also a p21-modulating agent.

Preferred CSNK1G-modulating agents include small molecule compounds; CSNK1G-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19<sup>th</sup> edition.

#### 15            **Small molecule modulators**

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the CSNK1G protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for CSNK1G-modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the p21 pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary

functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

### Protein Modulators

Specific CSNK1G-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the p21 pathway and related disorders, as well as in validation assays for other CSNK1G-modulating agents. In a preferred embodiment, CSNK1G-interacting proteins affect normal CSNK1G function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, CSNK1G-interacting proteins are useful in detecting and providing information about the function of CSNK1G proteins, as is relevant to p21 related disorders, such as cancer (e.g., for diagnostic means).

A CSNK1G-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with a CSNK1G, such as a member of the CSNK1G pathway that modulates CSNK1G expression, localization, and/or activity. CSNK1G-modulators include dominant negative forms of CSNK1G-interacting proteins and of CSNK1G proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous CSNK1G-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3<sup>rd</sup>, Trends Genet (2000) 16:5-8).

An CSNK1G-interacting protein may be an exogenous protein, such as a CSNK1G-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). CSNK1G antibodies are further discussed below.

In preferred embodiments, a CSNK1G-interacting protein specifically binds a CSNK1G protein. In alternative preferred embodiments, a CSNK1G-modulating agent binds a CSNK1G substrate, binding partner, or cofactor.

## 5            *Antibodies*

In another embodiment, the protein modulator is a CSNK1G specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify CSNK1G modulators. The antibodies can also be used in dissecting the portions of the CSNK1G pathway responsible for various cellular responses and in the general processing and maturation of the CSNK1G.

Antibodies that specifically bind CSNK1G polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of CSNK1G polypeptide, and more preferably, to human CSNK1G. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab').sub.2 fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of CSNK1G which are particularly antigenic can be selected, for example, by routine screening of CSNK1G polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Natl. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89; Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence shown in any of SEQ ID NOs:13-16. Monoclonal antibodies with affinities of  $10^8 \text{ M}^{-1}$  preferably  $10^9 \text{ M}^{-1}$  to  $10^{10} \text{ M}^{-1}$ , or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of CSNK1G or substantially purified fragments thereof. If CSNK1G fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of a CSNK1G protein. In a particular embodiment, CSNK1G-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of CSNK1G-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding CSNK1G polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

5 Chimeric antibodies specific to CSNK1G polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing  
10 together genes that encode the appropriate regions from each species (Morrison et al., Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies  
15 into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun.  
20 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

CSNK1G-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No.  
25 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As  
30 used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells

that express the targeted protein (Menard S, et al., *Int J. Biol Markers* (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about 10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

### **Nucleic Acid Modulators**

Other preferred CSNK1G-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit CSNK1G activity. Preferred nucleic acid modulators interfere with the function of the CSNK1G nucleic acid such as DNA replication, transcription, translocation of the CSNK1G RNA to the site of protein translation, translation of protein from the CSNK1G RNA, splicing of the CSNK1G RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the CSNK1G RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to a CSNK1G mRNA to bind to and prevent translation, preferably by binding to the 5' untranslated region. CSNK1G-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other 5 embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide 10 may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of 15 which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton 20 J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred CSNK1G nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded 25 RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239- 30 245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al.*, Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, a CSNK1G-specific nucleic acid modulator is used in an assay to further elucidate the role of the CSNK1G in the p21 pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, a CSNK1G-specific antisense oligomer is used as a therapeutic agent for treatment of p21-related disease states.

### Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of CSNK1G activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the CSNK1G nucleic acid or protein. In general, secondary assays further assess the activity of a CSNK1G modulating agent identified by a primary assay and may confirm that the modulating agent affects CSNK1G in a manner relevant to the p21 pathway. In some cases, CSNK1G modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising a CSNK1G polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates CSNK1G activity, and hence the p21 pathway. The CSNK1G polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

## Primary Assays

The type of modulator tested generally determines the type of primary assay.

### *Primary assays for small molecule modulators*

5 For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, *Curr Opin Chem Biol* (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead  
10 cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding),  
15 transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicity and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular  
20 molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of CSNK1G and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that  
25 retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when CSNK1G-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the CSNK1G protein may be assayed by various known  
30 methods such as substrate processing (*e.g.* ability of the candidate CSNK1G-specific binding agents to function as negative effectors in CSNK1G-expressing cells), binding equilibrium constants (usually at least about  $10^7 \text{ M}^{-1}$ , preferably at least about  $10^8 \text{ M}^{-1}$ , more preferably at least about  $10^9 \text{ M}^{-1}$ ), and immunogenicity (*e.g.* ability to elicit CSNK1G specific antibody in a heterologous host such as a mouse, rat, goat or rabbit).

For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a CSNK1G polypeptide, a fusion protein thereof, or to cells or  
5 membranes bearing the polypeptide or fusion protein. The CSNK1G polypeptide can be full length or a fragment thereof that retains functional CSNK1G activity. The CSNK1G polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The CSNK1G polypeptide is preferably human CSNK1G, or  
10 is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of CSNK1G interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has CSNK1G –specific binding activity, and can be used to assess normal CSNK1G gene function.

Suitable assay formats that may be adapted to screen for CSNK1G modulators are  
15 known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, *Curr Opin Chem Biol* (1998) 2:597-603; Sundberg SA, *Curr Opin Biotechnol* 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved  
20 fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.*, Selvin PR, *Nat Struct Biol* (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, *Curr Opin Chem Biol* (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate CSNK1G  
25 and p21 pathway modulators (*e.g.* U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); and U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

30

**Kinase assays.** In some preferred embodiments the screening assay detects the ability of the test agent to modulate the kinase activity of a CSNK1G polypeptide. In further embodiments, a cell-free kinase assay system is used to identify a candidate p21 modulating agent, and a secondary, cell-based assay, such as an apoptosis or hypoxic

induction assay (described below), may be used to further characterize the candidate p21 modulating agent. Many different assays for kinases have been reported in the literature and are well known to those skilled in the art (e.g. U.S. Pat. No. 6,165,992; Zhu et al., Nature Genetics (2000) 26:283-289; and WO0073469). Radioassays, which monitor the transfer of a gamma phosphate are frequently used. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from gamma -<sup>33</sup>P ATP to a biotinylated peptide substrate; the substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand.

Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133).

Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

**Apoptosis assays.** Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the caspase cleavage activity as part of a cascade of events that occur during programmed cell

death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-ONE™ Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death assay known to those skilled in the art, and available commercially (Roche, Cat# 1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses monoclonal antibodies directed against DNA and histones respectively, thus specifically determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due to the fact that DNA fragmentation occurs several hours before the plasma membrane breaks down, allowing for accumulation in the cytoplasm. Nucleosomes are not present in the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether CSNK1G function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the CSNK1G plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

25

**Cell proliferation and cell cycle assays.** Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, *Int. J. Cancer* 38, 369; Campana *et al.*, 1988, *J. Immunol. Meth.* 107, 79), or by other means.

30

Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specific to the phosphorylated form

of the serine 10 residue of histone H3. (Chadlee, D.N. 1995, *J. Biol. Chem* 270:20098-105). Cell Proliferation may also be examined using [<sup>3</sup>H]-thymidine incorporation (Chen, J., 1996, *Oncogene* 13:1395-403; Jeoung, J., 1995, *J. Biol. Chem.* 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, 5 cells synthesizing DNA will incorporate [<sup>3</sup>H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL et al., 1998, *In Vitro Cell Dev Biol Anim* 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt, MTS. MTS assays are commercially available, for example, the Promega CellTiter 96<sup>®</sup> Aqueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

15 Cell proliferation may also be assayed by colony formation in soft agar (Sambrook et al., *Molecular Cloning*, Cold Spring Harbor (1989)). For example, cells transformed with CSNK1G are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Cell proliferation may also be assayed by measuring ATP levels as indicator of 20 metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo<sup>™</sup>, which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) *Int J Radiat Biol Relat Stud Phys Chem Med* 49:237-55). Cells transfected with a CSNK1G may be stained with propidium iodide and evaluated in a flow 25 cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no 30 test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test

whether CSNK1G function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the CSNK1G plays a direct role in cell proliferation or cell cycle.

**Angiogenesis.** Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether CSNK1G function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the CSNK1G plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

**Hypoxic induction.** The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glycolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with CSNK1G in hypoxic conditions (such as with 0.1% O<sub>2</sub>, 5% CO<sub>2</sub>, and balance N<sub>2</sub>, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic

conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic  
5 induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether CSNK1G function plays a direct  
10 role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the CSNK1G plays a direct role in hypoxic induction.

15 **Cell adhesion.** Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The  
20 wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in  
25 a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells  
30 expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF, and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators  
5 against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., *Bioconjug Chem.* 2001 May-Jun;12(3):346-53).

**Tubulogenesis.** Tubulogenesis assays monitor the ability of cultured cells,  
10 generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include Matrigel<sup>TM</sup> (Becton Dickinson), an extract of basement membrane proteins containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4° C and forms a solid gel at 37° C. Other suitable matrices comprise extracellular components  
15 such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, *Nature Medicine* 5:1418-1423). These assays have traditionally involved  
20 stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond  
25 differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF- $\alpha$ . Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing a CSNK1G's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF- $\alpha$ , ephrin, etc.

**Cell Migration.** An invasion/migration assay (also called a migration assay) tests  
30 the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, *J Biol Chem* 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as

described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing a CSNK1G's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

15

**Sprouting assay.** A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900 $\mu$ l of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100  $\mu$ l of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

25  
30

### ***Primary assays for antibody modulators***

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the CSNK1G protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 5 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting CSNK1G-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

10

### ***Primary assays for nucleic acid modulators***

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance CSNK1G gene expression, preferably mRNA expression. In general, expression analysis comprises comparing CSNK1G expression in like 15 populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express CSNK1G) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that 20 CSNK1G mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most 25 commonly detected with specific antibodies or antisera directed against either the CSNK1G protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve CSNK1G mRNA expression, may also be used 30 to test nucleic acid modulators.

### **Secondary Assays**

Secondary assays may be used to further assess the activity of CSNK1G-modulating agent identified by any of the above methods to confirm that the modulating

agent affects CSNK1G in a manner relevant to the p21 pathway. As used herein, CSNK1G-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or  
5 to test the specificity of the modulating agent's interaction with CSNK1G.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express CSNK1G) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate CSNK1G-modulating agent results in  
10 changes in the p21 pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use “sensitized genetic backgrounds”, which, as used herein, describe cells or animals engineered for altered expression of genes in the p21 or interacting pathways.

#### 15 *Cell-based assays*

Cell based assays may use a variety of mammalian cell lines known to have defective p21 function such as HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). Cell based assays may detect endogenous p21 pathway activity or may rely on recombinant expression of p21  
20 pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

#### *Animal Assays*

25 A variety of non-human animal models of normal or defective p21 pathway may be used to test candidate CSNK1G modulators. Models for defective p21 pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the p21 pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration,  
30 injection, etc.

In a preferred embodiment, p21 pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal p21 are used to test the candidate modulator's affect on CSNK1G in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin,

collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the CSNK1G. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on CSNK1G is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, e.g., Ogawa K et al., 2000, *Oncogene* 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the CSNK1G endogenously are injected in the flank,  $1 \times 10^5$  to  $1 \times 10^7$  cells per mouse in a volume of 100  $\mu$ L using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorigenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for

evaluation of the candidate modulator. Tumorogenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorogenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorogenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

#### **Diagnostic and therapeutic uses**

Specific CSNK1G-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the p21 pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the p21 pathway in a cell, preferably a cell pre-determined to have defective or impaired p21 function (e.g. due to overexpression, underexpression, or misexpression of p21, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates CSNK1G activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the p21 function is restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought

closer to normal compared to untreated cells. For example, with restored p21 function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired p21 function by administering a therapeutically effective amount of a CSNK1G -modulating agent that modulates the p21 pathway. The invention further provides methods for modulating CSNK1G function in a cell, preferably a cell pre-determined to have defective or impaired CSNK1G function, by administering a CSNK1G -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired CSNK1G function by administering a therapeutically effective amount of a CSNK1G -modulating agent.

The discovery that CSNK1G is implicated in p21 pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the p21 pathway and for the identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether CSNK1G expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, eds., John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective p21 signaling that express a CSNK1G, are identified as amenable to treatment with a CSNK1G modulating agent. In a preferred application, the p21 defective tissue overexpresses a CSNK1G relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial CSNK1G cDNA sequences as probes, can determine whether particular tumors express or overexpress CSNK1G. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of CSNK1G expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the CSNK1G oligonucleotides, and antibodies directed against a CSNK1G, as described above for: (1) the detection of the presence of CSNK1G gene mutations, or the detection of either over- or under-expression of CSNK1G mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-

abundance of CSNK1G gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by CSNK1G.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in CSNK1G expression; the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for CSNK1G expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer, most preferably a cancer as shown in TABLE 1. The probe may be either DNA or protein, including an antibody.

### EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

#### I. Drosophila p21 screen

A dominant loss of function screen was carried out in *Drosophila* to identify genes that interact with the cyclin dependent kinase inhibitor, p21 (Bourne HR, et al., Nature (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95). Expression of the p21 gene from GMR-p21 transgene (Hay, B. A., et al. (1994) Development 120:2121-2129) in the eye causes deterioration of normal eye morphology, resulting in reduced, rough eyes. Flies carrying this transgene were maintained as a stock (P 1025 F, genotype: y w; P{p21-pExp-gl-w[+]Hsp70(3'UTR)-5}). Females of this stock were crossed to a collection of males carrying piggyBac insertions (Fraser M *et al.*, Virology (1985) 145:356-361). Resulting progeny carrying both the transgene and transposons were scored for the effect of the transposon on the eye phenotype, i.e. whether the transposon enhanced or suppressed (or had no effect) the eye phenotype. All data was recorded and all modifiers were retested with a repeat of the original cross, and the retests were scored at least twice. Modifiers of the eye phenotype were identified as members of the p21 pathway. GISH (CG6963) was an enhancer of the eye phenotype. Orthologs of the modifiers are referred to herein as CSNK1G.

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of *Drosophila* modifiers.

Various domains, signals, and functional subunits in proteins were analyzed using the PSORT (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6; Kenta Nakai, Protein sorting signals and prediction of subcellular localization, Adv. Protein Chem. 54, 277-344 (2000)), PFAM (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2),  
5 SMART (Ponting CP, et al., SMART: identification and annotation of domains from signaling and extracellular protein sequences. Nucleic Acids Res. 1999 Jan 1;27(1):229-32), TM-HMM (Erik L.L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow,  
10 T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998), and clust (Remm M, and Sonnhammer E. Classification of transmembrane protein families in the Caenorhabditis elegans genome and identification of human orthologs. Genome Res. 2000 Nov;10(11):1679-89) programs. For example, the kinase domain (PFAM 00069) of CSNK1G from GI#s 11545751, 21314778, and 4758080 (SEQ  
15 ID NOs:13, 15, and 16, respectively) is located respectively at approximately amino acid residues 44-312, 46-312, and 43-308.

## II. High-Throughput In Vitro Fluorescence Polarization Assay

Fluorescently-labeled CSNK1G peptide/substrate are added to each well of a 96-  
20 well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of CSNK1G activity.

25

## III. High-Throughput In Vitro Binding Assay.

<sup>33</sup>P-labeled CSNK1G peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM MgCl<sub>2</sub>, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a  
30 Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate p21 modulating agents.

#### IV. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins,  $3 \times 10^6$  appropriate recombinant cells containing the CSNK1G proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM HEPES, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at  $15,000 \times g$  for 15 min. The cell lysate is incubated with 25  $\mu$ l of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

#### V. Kinase assay

A purified or partially purified CSNK1G is diluted in a suitable reaction buffer, e.g., 50 mM HEPES, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10  $\mu$ g/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100  $\mu$ l. The reaction is initiated by the addition of  $^{33}\text{P}$ -gamma-ATP (0.5  $\mu$ Ci/ml) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation ( $\text{Mg}^{2+}$  or  $\text{Mn}^{2+}$ ) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is

defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

#### VI. Expression analysis

5 All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, Ardais, Genome Collaborative, and Ambion.

10 TaqMan analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/ $\mu$ l. Single stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of 15 Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan protocols, and the following criteria: a) primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis was performed using a 20 7900HT instrument.

Taqman reactions were carried out following manufacturer's protocols, in 25  $\mu$ l total volume for 96-well plates and 10  $\mu$ l total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis was prepared using a universal pool of human cDNA samples, which is a 25 mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched normal tissues from the same patient. A gene was considered overexpressed in a tumor 30 when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue was not available, a universal pool of cDNA samples was used instead. In these cases, a gene was considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type was greater

than 2 times the standard deviation of all normal samples (i.e., Tumor – average(all normal samples) > 2 x STDEV(all normal samples) ).

Results are shown in Table 1. Number of pairs of tumor samples and matched normal tissue from the same patient are shown for each tumor type. Percentage of the samples with at least two-fold overexpression for each tumor type is provided. A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

15

Table 1

<b>SEQ ID NO</b>	1	8	11
<b>Breast</b>	7%	17%	7%
<b># of Pairs</b>	30	30	30
<b>Colon</b>	3%	12%	8%
<b># of Pairs</b>	36	33	36
<b>Kidney</b>	0%	5%	0%
<b># of Pairs</b>	20	20	20
<b>Lung</b>	3%	9%	0%
<b># of Pairs</b>	35	32	35
<b>Ovary</b>	11%	26%	0%
<b># of Pairs</b>	19	19	19
<b>Prostate</b>	7%	7%	7%
<b># of Pairs</b>	15	14	15
<b>Uterus</b>	5%	5%	5%
<b># of Pairs</b>	19	19	19

## VII. CSNK1G functional assays

RNAi experiments were carried out to knock down expression of CSNK1Gs in various cell lines using small interfering RNAs (siRNA, Elbashir et al, *supra*).

20

Effect of CSNK1G RNAi on cell proliferation. BrdU and Cell Titer-Glo™ assays, as described above, were employed to study the effects of decreased CSNK1G expression on cell proliferation. The results of these experiments indicated that RNAi of CSNK1G of SEQ ID NOs:1, 8, and 11 decreases proliferation in LX1 small cell lung cancer cells, and  
5 in 231T breast cancer cells. MTS cell proliferation assay, as described above, was also employed to study the effects of decreased CSNK1G expression on cell proliferation. The results of this experiment indicated that RNAi of CSNK1G of SEQ ID NOs:1, 8, and 11 decreased proliferation in the above cell lines in addition to A549lung cancer cells.

Effect of CSNK1G RNAi on apoptosis. Nucleosome ELISA apoptosis assay, as  
10 described above, was employed to study the effects of decreased CSNK1G expression on apoptosis. Results indicated that RNAi of SEQ ID NO:1 caused increased apoptosis in A549 cells; RNAi of SEQ ID NO:8 did not have an affect on apoptosis; and RNAi of SEQ ID NO:11 caused increased apoptosis in LX1 cells.

## WHAT IS CLAIMED IS:

1. A method of identifying a candidate p21 pathway modulating agent, said method comprising the steps of:
  - 5 (a) providing an assay system comprising a CSNK1G polypeptide or nucleic acid;
  - (b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and
  - (c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as
- 10 a candidate p21 pathway modulating agent.
2. The method of Claim 1 wherein the assay system comprises cultured cells that express the CSNK1G polypeptide.
- 15 3. The method of Claim 2 wherein the cultured cells additionally have defective p21 function.
4. The method of Claim 1 wherein the assay system includes a screening assay comprising a CSNK1G polypeptide, and the candidate test agent is a small molecule
- 20 modulator.
5. The method of Claim 4 wherein the assay is a kinase assay.
6. The method of Claim 1 wherein the assay system is selected from the group consisting
- 25 of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.
7. The method of Claim 1 wherein the assay system includes a binding assay comprising a CSNK1G polypeptide and the candidate test agent is an antibody.
- 30 8. The method of Claim 1 wherein the assay system includes an expression assay comprising a CSNK1G nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.
10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.
- 5 11. The method of Claim 1 additionally comprising:  
(d) administering the candidate p21 pathway modulating agent identified in (c) to a model system comprising cells defective in p21 function and, detecting a phenotypic change in the model system that indicates that the p21 function is restored.
- 10 12. The method of Claim 11 wherein the model system is a mouse model with defective p21 function.
13. A method for modulating a p21 pathway of a cell comprising contacting a cell defective in p21 function with a candidate modulator that specifically binds to a CSNK1G  
15 polypeptide, whereby p21 function is restored.
14. The method of Claim 13 wherein the candidate modulator is administered to a vertebrate animal predetermined to have a disease or disorder resulting from a defect in  
20 p21 function.
15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.
16. The method of Claim 1, comprising the additional steps of:  
25 (e) providing a secondary assay system comprising cultured cells or a non-human animal expressing CSNK1G ,  
(f) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and  
30 (g) detecting an agent-biased activity of the second assay system,  
wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate p21 pathway modulating agent,  
and wherein the second assay detects an agent-biased change in the p21 pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.
18. The method of Claim 16 wherein the secondary assay system comprises a non-human  
5 animal.
19. The method of Claim 18 wherein the non-human animal mis-expresses a p21 pathway gene.
- 10 20. A method of modulating p21 pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a CSNK1G polypeptide or nucleic acid.
21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the p21 pathway.  
15
22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.
23. A method for diagnosing a disease in a patient comprising:  
20 (a) obtaining a biological sample from the patient;  
(b) contacting the sample with a probe for CSNK1G expression;  
(c) comparing results from step (b) with a control;  
(d) determining whether step (c) indicates a likelihood of disease.
- 25 24. The method of Claim 23 wherein said disease is cancer.
25. The method according to Claim 24, wherein said cancer is a cancer as shown in Table 1 as having >25% expression level.
- 30

## SEQUENCE LISTING

<110> EXELIXIS, INC.  
 <120> CSNK1GS AS MODIFIERS OF THE P21 PATHWAY AND METHODS OF USE  
 <130> EX03-057C-PC  
 <150> US 60/401,739  
 <151> 2002-08-07  
 <160> 16  
 <170> PatentIn version 3.2  
 <210> 1  
 <211> 1950  
 <212> DNA  
 <213> Homo sapiens  
 <400> 1  
 aatatggtgc taaaagaact ccttaccttt ctctgactac aatttatttg gacatacttt 60  
 tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag actctgtcct 120  
 gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca actaaacca 180  
 tggcacaaaag gagtgcacac tgctctcgac catctggctc ctcatcgtcc tctggggttc 240  
 ttatggtggg acccaacttc agggttggca agaagatagg atgtgggaac ttcggagagc 300  
 tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg gaaccaataa 360  
 aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt ggcagtgcag 420  
 gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat gccatggtgc 480  
 tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga acatttactt 540  
 tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac gtgcactcaa 600  
 agaacctcat ttaccgagat gtcaagccag agaacttctt gattgggtcga caaggcaata 660  
 agaaagagca tgttatacac attatagact ttggactggc caaggaatac attgaccccg 720  
 aaaccaaaaa acacatacct tatagggaac acaaaagttt aactggaact gcaagatata 780  
 tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg gaagccctag 840  
 gccatagtgt catgtatttc cttcgaggca gcctcccctg gcaaggactc aaggctgaca 900  
 cattaaaaga gagatatcaa aaaattggtg acaccaaaag gaatactccc attgaagctc 960  
 tctgtgagaa ctttcagag gagatggcaa cctaccttcg atatgtcagg cgactggact 1020  
 tctttgaaaa acctgattat gagtatttac ggaccctctt cacagacctc tttgaaaaga 1080  
 aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct actccagtag 1140  
 ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac acacataggg 1200  
 atcggccatc acaacagcag cctcttcgaa atcagagcct taggactggt acagctgagc 1260

attatgatgt taacaactca gccatctggc acaggggaag aggcacctaa acagattcct 1320  
 ctggacaaaa gtcacatatg agcaagatca gaatagtagc agatgtaaaa agcagaatgt 1380  
 atcatcagag cgccgaggag agtgggaaat tcagcccagc cggcagacca atacctcata 1440  
 cctaacgtct cacttggctg cagaccgcca tgggggatca gtgcaggtgg ttagctcaac 1500  
 caatggagag ctgaatgttg atgatcccac gggagcccac tccaatgcac caatcacagc 1560  
 tcatgccgag gtggaggtag tggaggaagc taagtgtctg tgtttcttta agaggaaaag 1620  
 gaagaagact gctcagcgcc acaagtgacc agtgcctccc aggagtcctc aggccttggg 1680  
 gactctgact caattgtacc tgcagctcct gccatttctc attggaaggg actcctcttt 1740  
 gggggagggt ggatatccaa actaaaaaga agaaaacaga tgccccaga aggggcccagt 1800  
 gcgggcagcc agggcctagt gggtcattgg ccatctccgc ctgcctaagg ctctgagcag 1860  
 gtcccagagc tgctgttctt ccaactgctt cccatagggc tgcttgggtg actctccttc 1920  
 ccattgttta cagtgaagggt gtcattcaca 1950

<210> 2  
 <211> 2066  
 <212> DNA  
 <213> Homo sapiens

<400> 2  
 ggcacgaggc tgctctcgac catctggctc ctcatcgtcc tctgggggttc ttatgggtggg 60  
 acccaacttc agggttggca agaagatagg atgtgggaac ttcggagagc tcagattagg 120  
 taaaaatctc tacaccaatg aatatgtagc aatcaactg gaaccaataa aatcacgtgc 180  
 tccacagctt catttagagt acagatttta taaacagctt ggcagtgagc gtgaaggctc 240  
 cccacaggtg tattactttg gaccatgtgg gaaatataat gccatgggtg tggagctcct 300  
 tggccctagc ttggaggact tgtttgacct ctgtgaccga acatttactt tgaagacggt 360  
 gttaatgata gccatccagc tgctttctcg aatggaatac gtgcactcaa agaacctcat 420  
 ttaccgagat gtcaagccag agaacttctt gattggctga caaggcaata agaaagagca 480  
 tgttatacac attatagact ttggactggc caaggaatac attgaccccg aaaccaaaaa 540  
 acacatacct tataggaac acaaaagttt aactggaact gcaagatata tgtctatcaa 600  
 cacgcatctt ggcaaagagc aaagccggag agatgatttg gaagccctag gccatatggt 660  
 catgtatttc cttcgaggca gcctcccctg gcaaggactc aaggctgaca cattaagaaga 720  
 gagatatcaa aaaattggtg acacaaaag gaatactccc attgaagctc tctgtgagaa 780  
 ctttcagag gagatggcaa cctaccttcg atatgtcagg cgactggact tctttgaaaa 840  
 acctgattat gagtatttac ggacctctt cacagacctc tttgaaaaga aaggctacac 900

ctttgactat gcctatgatt gggttgggag acctattcct actccagtag ggtcagttca 960  
 cgtagattct ggtgcatctg caataactcg agaaagccac acacataggg atcggccatc 1020  
 acaacagcag cctcttcgaa atcaggtggg tagctcaacc aatggagagc tgaatggtga 1080  
 tgatcccacg ggagcccact ccaatgcacc aatcacagct catgccgagg tggaggtagt 1140  
 ggaggaagct aagtgctgct gtttctttaa gaggaaaagg aagaagactg ctcagcgcca 1200  
 caagtgacca gtgcctccca ggagtcctca ggccctgggg actctgactc aattgtacct 1260  
 gcagctoctg ccattttctca ttggaagggg ctctctcttg ggggagggtg gatatccaaa 1320  
 ccaaaaagaa gaaaacagat gccccagaa ggggccagtg cgggcagcca gggcctagtg 1380  
 ggtcattggc catctcggcc tgcctaaggc tctgagcagg tcccagagct gctgttcctc 1440  
 cactgcttgc ccatagggct gcctgggtga ctctccttcc cattgtttac agtgaaggtg 1500  
 tcattcacia aaactcaagg actgctatc tccttcttcc ccttagttta ctctgggttt 1560  
 ttaccccacc ctcaaccctc tccagcataa aacctagtga gctaaaggct ttgtctgcag 1620  
 aaggagatca agaggctggg ggtaaggcca agaaggtagg aggaaaatgg cagacctggg 1680  
 ctggagaaga accttctccg tatcccaggt gtgcctggca gtatggtttc ctcttctct 1740  
 gtgcctgtgc agcattcatc ccagctggcc ttggggttca ggttccttct tccctccctc 1800  
 ctgtgaagtt aactgttagg acacaagctg tgagcaatct gcagtctact gtccctgtgt 1860  
 gttggcgttc ttagcttttt tgacaaaactc ttttctccag gtagtaggac aatgaaaatt 1920  
 gttctaagca aaggaaaagaa aactgacttt gttgcacttt tagttttttt aaaaaaaca 1980  
 aaaaacaaaa catggcagat gcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040  
 aaaaaaaaaa aaaaaaaaaa aaaaaa 2066

<210> 3  
 <211> 3552  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 ggagacggtg ttaatgatag ccatccagct gctttctcga atggaatacg tgcactcaaa 60  
 gaacctcatt taccgagatg tcaagccaga gaacttctctg attggctcgac gaggcaataa 120  
 gaaagagcat gttatacaca ttatagactt tggactggcc aaggaataca ttgacccccga 180  
 aaccaaaaaa cacatacctt atagggaaaca caaaagttta actggaactg caagatatat 240  
 gtctatcaac acgcatcttg gcaaagagca aagccggaga gatgatttgg aagccctagg 300  
 ccatatgttc atgtatttcc ttcgaggcag cctcccctgg caaggactca aggctgacac 360  
 attaaaagag agatatcaaa aaattggtga caccaaaagg aatactccca ttgaagctct 420  
 ctgtgagaac tttccagagg agatggcaac ctaccttcga tatgtcaggg gactggactt 480

ctttgaaaaa cctgattatg agtattttacg gaccctcttc acagacctct ttgaaaagaa	540
aggctacacc tttgactatg cctatgattg ggttgggaga cctattccta ctccagtagg	600
gtcagttcac gtagattctg gtgcatctgc aataactcga gaaagccaca cacatagggga	660
tcggccatca caacagcagc ctcttcgaaa tcaggtgggtt agctcaacca atggagagct	720
gaatgttgat gatcccacgg gagcccactc caatgcacca atcacagctc atgccgaggt	780
ggaggttagtg gaggaagcta agtgctgctg tttctttaag aggaaaagga agaagactgc	840
tcagcgccac aagtgaccag tgccctcccag gagtcctcag gccctgggga ctctgactca	900
attgtacctg cagctcctgc cattttctcat tggaaggac tcctctttgg gggaggggtg	960
atatccaaac caaaaagaag aaaacagatg cccccagaag gggccagtgc gggcagccag	1020
ggcctagtgg gtcattggcc atctccgct gcctaaggct ctgagcaggt cccagagctg	1080
ctgttcctcc actgcttgcc catagggctg cctggttgac tctcctccc attgtttaca	1140
gtgaaggtgt cattcacaaa aactcaagga ctgctattct cttcttccc cttagtttac	1200
tcctggtttt taccaccacc tcaaccctct ccagcataaa acctagttag ctaaaggctt	1260
tgtctgcaga aggagatcaa gaggctgggg gtaaggccaa gaaggtagga ggaaaatggc	1320
agacctgggc tggagaagaa ctttctccgt atcccaggtg tgcctggcag tatggtttcc	1380
tcttctctg tgccctgtgca gcattcatcc cggctggcct tggggttcag gttccttctt	1440
ccctccctcc tgtgaagtta cactgtagga cacaagctgt gagcaatctg cagtctactg	1500
tccccgtgtg taggogttct tagctttttt gacaaactct tttctccagg tagtaggaca	1560
atgaaaattg ttctaagcaa aggaaagaaa actgactttg ttgcactttt agttttttta	1620
aaaaaaacaa aaacaaaaac atggcagatg catattgtgt ctggttatat tgggggtttt	1680
acttttacct gttttgaggg ggatggggcc ggccaagcca ttcagagaga acatgggtcc	1740
agaggacatt ctgagtgaa agagtgtgat ctgcagcacc cagaagagaa gccaaactcg	1800
gtgtcattct gagtgaacac tcaggttggc aagaaaacat acttgaattt tcattcatct	1860
tctcagcagc tgaagaatgt ccctgccaga gcatcttgac ctaatcagct tacagtttga	1920
aaacctagct ctccagaaca tgagatgagc cagccgagcc agactgtgac caggaaacag	1980
ctcatcccag agaaggagat gcttaacaaa aaaaaattga aattgtttcc catgctgcca	2040
gggacttcca actagatagc catgtgacgt cctgggtgact tgagggaaaa attagtgatg	2100
aaacagccac caccatattg ccattagtga aaaaaagag gacagtgaac ctgccttcca	2160
cctgccagag ggacctcagg gtgtggcatt atagggccag gaaaagaaaa tcgggtgtatc	2220
ctatctgccc caatagctga gctgtagcat ttgggctggc ctgccttate agaaaccaag	2280
cttatgaaga tcttctccca gcaggtccat agcagtaggc ttaggatgca gtatatgggg	2340

ccgcatttaa aaggagggaa agattgtttg gtgctggaac attccaggggaa aaaggagact 2400  
 ggaatgaaag gtctgaaatt atcttctcaa ttggactcct tccagaaagg tggccgtgcc 2460  
 tctaagcatg tttttcccag tatgccctag gcctcccccc atgggtgtttt catatgaggt 2520  
 acaactgtga aggatctggt tcctcattca ctggttgaca agtctttcat gtgtggagtt 2580  
 actcttctca tgcccaattt tcatttgagt ttagtggtt aaccaaacia tgactcctca 2640  
 ttccagcggg gacagaagag aaagggtcat ttacatcagg aaagaggtct tgtatctggg 2700  
 agtagagagc taaccatgga gcacagtggc tgggtgggtga cttagtctga tggtttgtgg 2760  
 accatagaag tcttcacctc tggtttgagg tgcagggctg tcttttgtag tggaggggtgt 2820  
 ggggatattt tctgatagtt gccatttctt gaaaaattcc cttgatgtac cttacacaga 2880  
 gcagaaataa cattaacatg gatcagaggt actgggcttc atctgttcca ttggaccttg 2940  
 gctaggggaat atcatttcac tggcatcaaa cctgcttagc ttatgaaaag atggtaatat 3000  
 gtcatttcta taaatgtttc tataatgaa acataaagtgc gcaggagat acaatatcac 3060  
 accccttccc cacaaggact gtgaatattg ggatttatgt ccttgccatt acctagtggg 3120  
 tacagcccta tcactaaaat ttacatcgtt tctcagttgg gatttgggca ttgctaactt 3180  
 actgtataga aagtttaact tttcctcacc cctgtataga aaatgccttg cctctcaaga 3240  
 gagggcagag ggggggcccag gtgcagtggc tcacgcctgt aatcccagca gtttgggagg 3300  
 ccaaggcaag tggatcatgt gaggtcaaga gttcgagacc agcctggcca acatggtgaa 3360  
 acccctgtct tacaaaaaat acaaaaatta gctgggcatg gtggcatgct cccgtagtcc 3420  
 cagctactcg ggaggctgag gcaggagaat cacttgagcc tgggaggcag aagttgcagt 3480  
 gagccgagat cgcaccactg cactccagcc tgggcaacag agtgagactc tgtctaaaaa 3540  
 aaaaaaaaaa aa 3552

<210> 4  
 <211> 4222  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 gcctcatttg cgccttgcag cactgctgga ccagggtaca agatgttcac ctaagattga 60  
 gacctagtga ctacatttcc tacgggaaca aataaatggt ttttcatctc ccggagatac 120  
 attacaaaca aatatggtgc taaaagaact ccttaccttt ctctgactac aatttatttg 180  
 gacatacttt tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag 240  
 actctgtcct gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca 300  
 actaaacca tggcacaag gagtgcacac tgctctcgac catctggctc ctcatcgtcc 360

tctggggttc ttatggtggg acccaacttc agggttggca agaagatagg atgtgggaac	420
ttcgggagagc tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg	480
gaaccaataa aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt	540
ggcagtgcag gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat	600
gccatggtgc tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga	660
acatttactt tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac	720
gtgcactcaa agaacctcat ttaccgagat gtcaagccag agaacttctt gattggtcga	780
caaggcaata agaaagagca tgttatacac attatagact ttggactggc caaggaatac	840
attgaccccg aaaccaaaaa acacatacct tatagggaac acaaaagttt aactggaact	900
gcaagatata tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg	960
gaagccctag gccatatgtt catgtatttc ctctgaggca gcctcccctg gcaaggactc	1020
aaggctgaca cattaaaaga gagatatcaa aaaattgggtg acaccaaag gaatactccc	1080
attgaagctc tctgtgagaa ctttccagag gagatggcaa cctaccttcg atatgtcagg	1140
cgactggact tctttgaaaa acctgattat gagtatttac ggaccctctt cacagacctc	1200
tttgaaaaga aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct	1260
actccagtag ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac	1320
acacataggg atcggccatc acaacagcag cctcttcgaa atcaggtggt tagctcaacc	1380
aatggagagc tgaatggtga tgatcccacg ggagcccact ccaatgcacc aatcacagct	1440
catgccgagg tggaggtagt ggaggaagct aagtgctgct gtttctttaa gaggaaaagg	1500
aagaagactg ctcagcgcca caagtgacca gtgcctccca ggagtcctca ggccctgggg	1560
actctgactc aattgtacct gcagctcctg ccatttctca ttggaagga ctctctttg	1620
ggggagggtg gatatccaaa ccaaaaagaa gaaaacagat gccccagaa ggggccagtg	1680
cgggcagcca gggcctagtg ggtcattggc catctccgcc tgcctaaggc tctgagcagg	1740
tcccagagct gctgttcctc cactgcttgc ccatagggct gcctggttga ctctccttc	1800
cattgtttac agtgaaggtg tcattcacia aaactcaagg actgctattc tccttcttc	1860
ccttagttta ctcttggtt ttaccccacc ctcaaccctc tccagcataa aacctagtga	1920
gctaaaggct ttgtctgcag aaggagatca agaggctggg ggtaaggcca agaaggtagg	1980
aggaaaatgg cagacctggg ctggagaaga accttctccg tatcccaggt gtgcctggca	2040
gtatggtttc ctcttctct gtgcctgtgc agcattcatc ccagctggcc ttggggttca	2100
ggttcttct tccctcctc ctgtgaagtt aactgtagg acacaagctg tgagcaatct	2160
gcagtctact gtcctgtgt gttggcgttc ttagcttttt tgacaaactc ttttctccag	2220

gtagtaggac aatgaaaatt gttctaagca aaggaaagaa aactgacttt gttgcacttt	2280
tagttttttt aaaaaaaca aaaacaaaaa catggcagat gcatattgtg tctgggtata	2340
ttggggggtt tacttttacc tgttttgagg gggatggggc cggccaagcc attcagagag	2400
aacatgggtc cagaggacat tctcagtga aagagtttga tctgcagcac ccagaagaga	2460
agccaaactc ggtgtcattc tgagtgaaca ctcaggttgg caagaaaaca tacttgaatt	2520
ttcattcatc ttctcagcag ctgaagaatg tccctaccag agcatcttga cctaatacagc	2580
ttacagtttg aaaacctagc tctccagaac atgagatgag ccagccgagc cagactgtga	2640
ccaggaaaca gctcatccca gagaaggaga tgcttaaca aaaaaattg aaattgtttc	2700
ccatgctgcc agggacttcc aactagatag ccatgtgacg tcctggtgac ttgggggaaa	2760
aattagtgat gaaacagcca ccacatatt gccattagtg gaaaaaaga ggacagtga	2820
cctgccttcc acctgccaga gggacctcag ggtgtggcat tatagggcca ggaaaagaaa	2880
atcgggtgat cctatctgcc ccaatagctg agctgtagca tttgggctgg cctgccttat	2940
cagaaaccaa gcttatgaag atcttctccc agcaggtcca tagcagtagg cttaggatgc	3000
agtatatggg gccgcattta aaaggaggga aagattgttt ggtgctggaa cattccaggg	3060
aaaaggagac tggaatgaaa ggtctgaaat tatcttctca attggactcc ttccagaaag	3120
gtggccgtgc ctctaagcat gtttttcca gtatgccta ggcctcccc catggtgttt	3180
tcatatgagg tactactgtg aaggatctgg ttctcattc actgtttgac aagtctttca	3240
tgtgtggagt tactcttctc atgcccatt ttcatctgag tttagtggtc taaccaaca	3300
atgactctc attccagcgg tgacagaaga gaaagggtca tttacatcag gaaagagtc	3360
ttgtatctgg gagtagagag ctaacctgag agcacagtgg ctggtgggtg acttagtctg	3420
atggtttgtg gaccatagaa gtcttcacct ctggtttgag gtgcagggct gtcttttgta	3480
ctggaggggtg tggggatatt ttctgatagt tgccatttct tgaaaaattc cttgatgta	3540
ccttacacag agcagaaata acattaacat ggatcagagg tactgggctt catctgttcc	3600
attggacctt ggctagggaa tatcatttca ctggcatcaa acctgcttag cttatgaaaa	3660
gatggaata tgtcatttct ataaatgttt ctatatatga aacataaagt ggcagggaga	3720
tacaatatca cacccttcc ccacaaggac tgtgaatatt gggatttatg tccttgccat	3780
tacctagtgg ttacagccct atcactaaaa tttacatcgt ttctcagttg ggatttgggc	3840
attgctaact tactgtatag aaagtttaac ttttctcac ccctgtatag aaaatgcctt	3900
gcctctcaag agagggcaga gggggggcca ggtgcagtgg ctcacgcctg taatcccagc	3960
agtttgggag gccaaggcaa gtggatcatg tgagggtcaag agttcgagac cagcctggcc	4020
aacatggtga aacccctct ctacaaaaa tacaaaaatt agctgggcat ggtggcatgc	4080

tcccgtagtc ccagctactc gggaggctga ggcaggagaa tcacttgagc ctgggaggca 4140  
 gaagttgcag tgagccgaga tcgcaccact gcactccagc ctgggcaaca gagtgagact 4200  
 ctgtctaaaa aaaaaaaaaa aa 4222

<210> 5  
 <211> 1754  
 <212> DNA  
 <213> Homo sapiens

<400> 5  
 gacctagtga ctacatttcc tacgggaaca aataaatggt ttttcatctc ccggagatac 60  
 attacaaaca aatatggtgc taaaagaact ccgtaccttt ctctgactac aatttatttg 120  
 gacatacttt tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag 180  
 actctgtcct gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca 240  
 actaaacca tggcaciaag gagtgcacac tgctctcgac catctggctc ctcacgtcc 300  
 tctgggggttc ttatgggtggg acccaacttc agggttggca agaagatagg atgtgggaac 360  
 ttcggagagc tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg 420  
 gaaccaataa aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt 480  
 ggcagtgcag gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat 540  
 gccatggtgc tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga 600  
 acatttactt tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac 660  
 gtgcactcaa agaacctcat ttaccgagat gtcaagccag agaacttcct gattggtcga 720  
 caaggcaata agaaagagca tgttatacac attatagact ttggactggc caaggaatac 780  
 attgaccccg aaaccaaaaa acacatacct tatagggaaac acaaaaagttt aactggaact 840  
 gcaagatata tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg 900  
 gaagccctag gccatatggt catgtatttc ctctgaggca gcctcccctg gcaaggactc 960  
 aaggctgaca cattaagaaga gagatatcaa aaaattggtg acaccaaaag gaatactccc 1020  
 attgaagctc tctgtgagaa ctttccagag gagatggcaa cctaccttcg atatgtcagg 1080  
 cgactggact tctttgaaaa acctgattat gagtatttac ggaccctctt cacagacctc 1140  
 tttgaaaaga aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct 1200  
 actccagtag ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac 1260  
 acacataggg atcggccatc acaacagcag cctcttcgaa atcaggtggt tagctcaacc 1320  
 aatggagagc tgaatggtga tgatcccacg ggagcccact ccaatgcacc aatcacagct 1380  
 catgccgagg tgggggtagt ggaggaagct aaatgtataa tgtttcataa acctggcac 1440  
 tggtcacaaa gctcagctgt gaaaatgaaa tttgtagtat ttttaaacat gaatgtcaat 1500

ttcaagtgta tttgaaatgg ttcctccagg agagatttgt gcaccattag gaaaatcttc 1560  
 tctgcagagg aagtagcctt ctttggagaa aatggaaaat gggttctgat atgtgatctc 1620  
 agagtagccc atttcctagg gcaccatgga aaacacaaat gtgatcttta agtatacctc 1680  
 ttccccagtt tggggaggaa aggactcagt ttgcaccctt tttgtatgta aaataaaatg 1740  
 tcttaccttt cttg 1754

<210> 6  
 <211> 2446  
 <212> DNA  
 <213> Homo sapiens

<400> 6  
 gggatttgca cggcagcaga gtcaccgtgg agaggccagg gtatcacaaa cttatggatt 60  
 ttgacaagaa aggagggaaa ggggagacgg aggagggccg gagaatgtcc aaggccggcg 120  
 ggggccggag cagccacggc atccggagct cggggaccag ctccgggggtc ctgatgggtgg 180  
 gccccaaactt ccgcgtcggc aagaagatcg gctgcggcaa ctccggggag ctccgcctag 240  
 gaaagaatct ctatacaaat gaatacgtgg ctatcaaatt ggagccgatc aagtcccggg 300  
 ccccgagct gcacctggag taccggttct acaagcagct cagcgccaca gagggcgtcc 360  
 ctcaaggctta ctacttcggg ccgtgcggga attacaacgc catggtgctg gagctgctgg 420  
 ggcccagcct ggaggacctg ttcgacctgt gcgaccggac cttcacgctc aagacgggtgc 480  
 tgatgatcgc catccagctg atcacgcgca tggagtatgt gcacaccaag agcctaactc 540  
 accgggacgt gaagcccagag aacttcctgg tgggccgccc ggggaccaag cggcagcatg 600  
 ccatccacat catcgacttc gggctggcca aggagtacat cgaccccag accaagaagc 660  
 acatcccgtta ccgcgagcac aagagcctga cgggcacggc gcgctacatg agcatcaaca 720  
 cgcacctggg caaggagcag agccgcccgg acgacctgga ggcgctgggc cacatgttca 780  
 tgtacttcct gcgcggcagc ctcccctggc aggggctcaa ggccgacacg ctcaaggagc 840  
 ggtaccagaa gatcggggac accaaacgcg ccacgcccac cgaggtgctc tgcgagaact 900  
 tcccagagga gatggccacg tacctgcgct atgtgcggcg cctggacttc ttcgagaagc 960  
 ccgactatga ctacctgagg aagctcttca ccgacctctt cgaccgcagt ggcttcgtgt 1020  
 tcgactatga gtacgactgg gccgggaagc ccctgccgac ccccatcggc accgtccaca 1080  
 ccgacctgcc ctcccagcct cagctccggg acaaaaacca gccgcacagc aaaaaccagg 1140  
 cgttgaactc caccaacggg gagctgaatg cggacgacct cacggccggc cactccaacg 1200  
 ccccgatcac agcgcctgca gaggtggagg tggccgatga aaccaaagtc tgctgtttct 1260  
 tcaagaggag aaagagaaaa tcgctgcagc gacacaagtg accctggggc cgtgcagccc 1320

cctgaatctt ctccgtgcag ccccttgggg cgcgaccttg tgcgaggccc tcggggccca 1380  
 cccacagcgg cccagggcca gaccctgggt ggaagccaga acgcagactg cagggggccgc 1440  
 gcctggctca ggcggcccca cccccgggac gtgggggtcac ttccttcacg taagactttg 1500  
 gccgaaattt ctacacctgt gtctagtctt cccctccaag agcattaact atttaaaaca 1560  
 aggaaaagag gaaaaaaaaa acagaggccc gccctacccc actcctgccc ctccgtttct 1620  
 ttgctgaagt gagtagtggt atcctggagg cccccgggcc tggccccgcc ccgccagccg 1680  
 cccccgttag cgtcataaag tccagcttgt ctccctcgat ccaaaggccg ttttctcgag 1740  
 gggagggcag gcccggcctg gaggggtgct gtggagctgt cttgcccagg ccctcctggg 1800  
 agggggacag gcattgttgc caggggtgag gccgtgcccc aggcctcccc gaaaccaaag 1860  
 ggggaaggcag ggggtggggc gtggctgaag ccggctcccc aaccaaagtg ctgcacaaa 1920  
 gctcggggcg cgcgggcacg gctgctgcag tctcttccca gcctggccct ggcaaggggc 1980  
 ggggtggggc tgccaggcgg gtgcttctcg acgcacttgc tcccggaggc tgcgccccgg 2040  
 cgcctggaac ccgaggtggg aggaccggtt ggtgtcacc tgctcggccc tcagccctgc 2100  
 cgcgtggggc gcgtggggcac ggagcttctt gcctctgctc cgacaccggg caagcagccg 2160  
 gagacaaaac gccttaaagc cccccggcca gccctgcagg tatattgcag gggcctgggg 2220  
 ggggccctgg actggcgggc ggttccccag tggggtgccc tggaggctgc cgggcagagt 2280  
 ggagcagctt ggggccgtgc ccagggcggg ggctgtgagt ctagtttttg ctttaccag 2340  
 tgtacagaaa tggcatttac gtttctctga tgctcccttg aagccataga atttaggggc 2400  
 ttttttaaaa aaataaaaga aaaatgaaac caaaaaaaaa aaaaaa 2446

<210> 7  
 <211> 2446  
 <212> DNA  
 <213> Homo sapiens

<400> 7  
 gggatttgca cggcagcaga gtcaccgtgg agaggccagg gtatcacaaa cttatggatt 60  
 ttgacaagaa aggagggaaa ggggagacgg aggagggccg gagaatgtcc aaggccggcg 120  
 ggggccggag cagccacggc atccggagct cggggaccag ctccgggggtc ctgatgggtg 180  
 gcccactt ccgcgtcggc aagaagatcg gctgcggcaa ctccggggag ctccgcctag 240  
 gaaagaatct ctatacaaat gaatacgtgg ctatcaaatt ggagccgatc aagtcccggg 300  
 ccccgcagct gcacctggag taccggttct acaagcagct cagcgccaca gagggcgtcc 360  
 ctcaggtcta ctacttcggg ccgtgcggga attacaacgc catgggtgctg gagctgctgg 420  
 ggcccagcct ggaggacctg ttcgacctgt gcgaccggac cttcacgctc aagacgggtg 480  
 tgatgatcgc catccagctg atcacgcgca tggagtatgt gcacaccaag agcctaactc 540

accgggacgt gaagccccgag aacttcctgg tgggcccgcc ggggaccaag cggcagcatg	600
ccatccacat catcgacttc gggctggcca aggagtacat cgacccccgag accaagaagc	660
acatccccgta ccgcgagcac aagagcctga cgggcacggc gcgctacatg agcatcaaca	720
cgcacctggg caaggagcag agccgccgag acgacctgga ggcgctgggc cacatgttca	780
tgtacttcct gcgcggcagc ctcccctggc aggggctcaa ggccgacacg ctcaaggagc	840
ggtaccagaa gatcggggac accaaacgag ccacgccccat cgagggtgctc tgcgagaact	900
tcccagagga gatggccacg tacctgcgct atgtgcggcg cctggacttc ttcgagaagc	960
ccgactatga ctacctgagg aagctcttca ccgacctctt cgaccgcagt ggcttcgtgt	1020
tcgactatga gtacgactgg gccgggaagc ccctgccgac ccccatcggc accgtccaca	1080
ccgacctgcc ctcccagcct cagctccggg acaaaacca gccgcacagc aaaaaccagg	1140
cgttgaactc caccaacggg gagctgaatg cggacgacct cacggccggc cactccaacg	1200
ccccgatcac agcgcctgca gaggtggagg tggccgatga aaccaaagtc tgctgtttct	1260
tcaagaggag aaagagaaaa tcgctgcagc gacacaagtg accctgggag cgtgcagccc	1320
cctgaatctt ctccgtgcag ccccttgggg cgcgacctg tgcgaggccc tcggggccca	1380
cccacagcgg cccagggcca gaccctggct ggaagccaga acgcagactg caggggccgc	1440
gcctggctca ggcggcccca cccccgggac gtgggggtcac ttccttcag taagactttg	1500
gccgaaattt ctacacctgt gtctagtcct cccctccaag agcattaact atttaaaaca	1560
aggaaaagag gaaaaaaaaa acagaggccc gccctacccc actcctgccc ctccgtttct	1620
ttgctgaagt gagtagtgtg atcctggagg cccccggcc tggccccgcc ccgccagccg	1680
ccccgtag cgtcataaag tccagcttgt ctccctcgat ccaaaggccg ttttctcgag	1740
gggagggcag gcccgccctg gaggggtgct gtggagctgt cttgcccagg ccctcctggg	1800
agggggacag gcattgttgc caggggtgag gccgtgcccc aggcctcccc gaaaccaaag	1860
gggaaggcag ggggtggggc gtggctgaag ccggctcccc aaccaaagtc ctgcaccaa	1920
gctcgggcgc gcgcggcacg gctgctgcag tctcttccca gcctggccct ggcaaggggc	1980
gggtgggcgc tgccaggcgg gtgcttctcg acgcacttgc tcccggaggc tgcgccccgg	2040
cgctggaac ccgaggtggg aggaccggtt ggtgtcacc cgtctggccc tcagccctgc	2100
cgctggggc gcgtgggcac ggagcttctt gcctctgctc cgacaccgg caagcagccg	2160
gagacaaaac gccttaaagc ccccggcca gccctgcagg tatattgcag gggcctgggg	2220
gcggccctgg actggcgggc ggttccccag tggggtgccc tggaggctgc cgggcagagt	2280
ggagcagctt ggggccgtgc ccagggcggg ggctgtgagt ctagtttttg ctttaccag	2340
tgtacagaaa tggcatttac gtttctctga tgctcccttg aagccataga atttaggggc	2400

ttttttaaaa aaataaaaga aaaatgaaac caaaaaaaaa aaaaaa 2446

<210> 8

<211> 2773

<212> DNA

<213> Homo sapiens

<400> 8

cggcaccgagc agcagaatgt ctccctgcccc cgagagcgcac cccgaggcca ctgagaagag 60  
cagcgcgggcc tggccggccc gaacgcctgc gtctcagtag ctgggagcca cgggcccacg 120  
cccgcccacc ggccgcagtg atgttctagc cacagaggag ccaagacctc aggtttccag 180  
agacttggga tttgcacggc agcagagtca ccgtggagag gccagggtat cacaaactta 240  
tggatthttga caagaaagga gggaaagggg agacggagga gggccggaga atgtccaagg 300  
ccggcggggg ccggagcagc cacggcatcc ggagctcggg gaccagctcg ggggtcctga 360  
tgggtgggccc caacttccgc gtcggcaaga agatcggctg cggcaacttc ggggagctcc 420  
gcctagtgca gagtagaaga gccgggaagg accccatcac catcagatgc atcagagctt 480  
caaaggcccc cccatcaaga cgctgccaag agcatgaggc cagccaggct ggggaggacg 540  
ttggcgagca tccgcctgcc cgtcaggag ccacagctgg atgatacatc gacagacagc 600  
agagggacgg tgtagcacac aggcgagggc ttgtcctcct cagcctgggg gagcgcagct 660  
cagtgccttg tggtcagcac gtgactgaca ctttgccgga gaaggggacg gctgcccgca 720  
ggaagagcta cacagccgtg ttcgccgga ctccgctctg gagagtcaga actggagacg 780  
ggagggcgctg cctcaagggc agatgggcaa accaaacccc atcctccctg tgagccgcgg 840  
cacagcacca ggcgggcaca cgcagcacga cacggcgagg agcccaccgc aggggcagcc 900  
ttttgggccc ggaagggggg atgaatccac tgtgaatccc acctccgaag ggcagcgcag 960  
gaaggacacg caaatgctgg actgaacctg cagctgattc agggggcagt gtcctgagct 1020  
ctaaaacttc aaacctgggg cagtgtcctg agctctaaaa cttcgaaatc tgaccaggca 1080  
cagtggctca cgctgtaat cccaacactt tgggaggctg atgtgggagg atcacctgac 1140  
gtcaggagtt cgagaccagc ctggacaaca tggagaaagc ccatctctag taaaaataca 1200  
aaaattagcc aggttgatgg tgccctgctg taattccagc tactcaggag gctgaggcag 1260  
gagaatcgct tgaaccagc aggcagaggt tccagtgagc cgtgatcgcg ccagtgcact 1320  
ccagcctggg caacagaacg agactctgct caaaaagaa aaaaactttg aaatccctct 1380  
aaaaataaaa tggaccggtg gatggtagct gtgaggatgg accgtgtggc aggaaagaat 1440  
ctctatacaa atgaatacgt ggctatcaaa ttggagccga tcaagtcccc ggccccgag 1500  
ctgcacctgg agtaccggtt ctacaagcag ctcagcgcca cagagggcgt ccctcaggtc 1560

tactacttcg gtccgtgagg gaagtacaac gccatggtgc tggagctgct ggggccagc 1620  
ctggaggacc tgttcgacct gtgagaccgg accttcacgc tcaagacggt gctgatgatc 1680  
gccatccagc tgatcacgag catggagtat gtgcacacca agagcctaata ctaccgggac 1740  
gtgaagcccc agaacttcct ggtggggccg ccggggacca agcggcagca tgccatccac 1800  
atcatcgact tcgggctggc caaggagtac atcgaccccc agaccaagaa gcacatcccc 1860  
taccgagagc acaagagcct gacggggcag gcgagctaca tgagcatcaa cacgcacctg 1920  
ggcaaggagc agagccgccc cgacgacctg gagggcgtgg gccacatggt catgtacttc 1980  
ctgagcggca gcctccccct gcaggggctc aaggccgaca cgctcaagga gcggtaccag 2040  
aagatcgggg acaccaaacc cgccacgccc atcgaggtgc tctgagagaa cttcccagag 2100  
gagatggcca cgtacctgag ctatgtgagg cgcctggact tcttcgagaa gcccgactat 2160  
gactacctgc ggaagctctt caccgacctc ttgagccgca gtggcttcgt gttcgactat 2220  
gagtacgact gggccgggaa gccctgcccg acccccacgc gcaccgtcca caccgacctg 2280  
ccctcccagc ctcagctccg ggacaaaacc cagccgcaca gcaaaaacca ggcgttgaac 2340  
tccaccaacc gggagctgaa tgcggacgac cccagggccc gccactcaa cggcccagtc 2400  
acagcgcctg cagaggtgga ggtggccgat gaaaccaaata gctgctgttt cttcaagagg 2460  
agaaagagaa aatcgctgca gcgacacaag tgaccctggg cgcgtgcagc cccctgaatc 2520  
ttctccgtgc agccccttgg ggagcagcct tgtgagaggc cctcggggcc caccacagc 2580  
ggcccagggc cagaccctgg ctggaagcca gaacgcagac tgcagggggc gcgctggct 2640  
caggcggccc cccccggg acgtggggtc acttccttca tgtaagactt tggccgaaat 2700  
ttctacacct gtgtctagtc ctcccctcca agagcattaa ctatttaaaa caaggaaaaa 2760  
aaaaaaaaa aaa 2773

<210> 9  
<211> 1615  
<212> DNA  
<213> Homo sapiens

<400> 9  
agtacatcga ccccgagacc aagaagcaca tcccgtaccg cgagcacaag agcctgacgg 60  
gcacggcgcg ctacatgagc atcaacacgc acctgggcaa ggagcagagc cgccgagacg 120  
acctggaggc gctgggcccac atgttcatgt acttcctgag cggcagcctc ccctggcagg 180  
ggctcaaggc cgacacgctc aaggagcggg accagaagat cggggacacc aaacgcgcca 240  
cgcccacatcga ggtgctctgc gagaacttcc cagaggagat ggccacgtac ctgagctatg 300  
tgaggcgcct ggacttcttc gagaagcccc actatgacta cctgaggaag ctcttcaccg 360  
acctcttcga ccgagctggc ttctgtgttc actatgagta cgactggggc gggaagcccc 420

tgccgacccc catcggcacc gtccacaccg acctgccctc ccagcctcag ctccgggaca 480  
 aaaccagcc cctgaatct tctccgtgca gcccttggg gcgacacctt gtgcgaggcc 540  
 ctccggggccc acccagcgcg gccagggccc agaccctggc tggaagccag aacgcagact 600  
 gcagggggccc cgcttggtc aggcggcccc accccggga cgtgggggtca cttccttcat 660  
 gtaagacttt ggccgaaatt tctacacctg tgtctagtcc tcccctcaa gagcattaac 720  
 tatttaaac aaggaaaaga ggaaaaaaaa aacagaggcc cgccctacc cactcctgcc 780  
 cctccgtttc tttgctgaag tgagtagtgt gatcctggag gcccccggc ctggccccgc 840  
 cccgccagcc gccccgtta gcgtcataaa gtccagcttg tctccctcga tccaaaggcc 900  
 gttttctcga ggggaggga ggcccggcct ggaggggtgc tgtggagctg tcttgcccag 960  
 gccctcctgg gagggggaca ggcatgttg ccaggggtga ggccgtgccc caggcctccc 1020  
 cgaaaccaa ggggaaggca ggggtggggc cgtggctgaa gccggctccc caacaaaat 1080  
 gctgcaccaa agctcgggcg ccgcgggcac ggctgctgca gtctcttccc agcctggccc 1140  
 tggcaagggg cgggtgggcg ctgccaggcg ggtgcttctc gacgcacttg ctcccggagg 1200  
 ctgcgccccg gcgctggaa cccgaggtgg gaggaccggt tgggtgcacc ctgctcggcc 1260  
 ctcagccctg ccgctgggg cgctgggca cggagcttcc tgctctctg ctccgacacc 1320  
 cggcaagcag ccggagaaa aacgcctaa agccccggc ccagccctgc aggtatattg 1380  
 caggggcctg gggcgggccc tggactggcg ggcggttccc cagtggggtg ccctggaggc 1440  
 tgccgggcag agtggagcag cttggggccc tgcccagggc ggtggctgtg agtctagttt 1500  
 ttgctttacc aagtgtacag aaatggcatt tacgtttctc tgatgctccc ttgaagccat 1560  
 agaatttagg ggctttttta aaaaaataa aagaaaaatg aaacaaaaaa aaaaa 1615

<210> 10

<211> 2421

<212> DNA

<213> Homo sapiens

<400> 10

gaattcaaag tggagtaccg caaacttgat atggaaaata aaaagaaaga caaggacaaa 60  
 tcagatgata gaatggcacg acctagtggc cgatcgggac acaacactcg aggaactggg 120  
 tcttcatcgt ctggagtttt aatgggtgga cctaacttta gagttgaaa aaaaattgga 180  
 tgtggcaatt ttggagaatt acgattaggg aaaaatttat acacaaatga atatgtggca 240  
 attaagttgg agcccatgaa atcaagagca ccacagctac atttggaata cagattctat 300  
 aagcagttag gatctggaga tggataacct caagtttact atttcggccc ttgtggtaaa 360  
 tacaatgcta tgggtgctgga actgctggga cctagtttgg aagacttggt tgacttggtg 420

gacagaacat tttctcttaa aacagttctc atgatagcta tacaactgat ttctcgcgatg 480  
gaatatgtcc attcaaagaa cttgatatac agagatgtaa aacctgagaa cttcttaata 540  
ggacgaccaa gaaacaaaac ccagcaagtt attcacatta tagatthttgg tttggcaaag 600  
gaatatattg atccggagac aaagaaacac ataccataca gagaacacaa gagccttaca 660  
ggaacagcta gatatatgag cataaacaca catttaggaa aagaacaaag tagaagagac 720  
gatttagaag ctttaggtca tatgttcatg tathttctga gaggcagtct tccttggcaa 780  
ggcttaaagg ctgacacatt aaaggagagg tatcagaaaa ttggagatac aaaacgggct 840  
acaccaatag aagtgttatg tgaaaattht ccagaaatgg caacatatct tcgttatgta 900  
agaaggctag atthttttga aaaaccagac tatgaatact taagaaagct thttactgac 960  
ttgtttgatc gaaaaggata tatgtttgat tatgaatatg actggattgg taaacagttg 1020  
cctactccag tgggtgcagt tcagcaagat cctgctctgt catcaaacag agaagcacat 1080  
caacacagag ataagatgca acaatccaaa aaccagtcgg cagaccacag ggcagcttgg 1140  
gactcccagc aggcaaatcc ccaccatttg agagctcacc ttgcagcaga cagacatgg 1200  
ggctcggtac aggttgtaag ttctacaaat ggagagthaa acacagatga ccccaccgca 1260  
ggacgttcaa atgcacccat cacagcccct actgaagtag aagtgatgga tgaaaccaag 1320  
tgctgctgct thttcaaacg aaggaaaagg aaaaccatac agcgccacaa atgactctgg 1380  
acacagacag atcctgggga gttacttaca tgttcatctg ctgtcttgtg atthaaatca 1440  
tctctgtagt gaccacgtat atthtcaagg actcactctt agaaacaaaa atgtcatact 1500  
atcatacttc atthtgtgg igtcttacet tctthttctt thttthtttc tctaathtaa 1560  
cctthtatgga agctthaaag thttgtcaaa acatgagtg ttgcccac agtgaatgga 1620  
atggaccaat gaggtggat caatgaatat agttccatag aacathttcc agaagthctt 1680  
ctgttgtaga aagcagtaca gtatcttaag tgtcaaccag ttatatacct aatctggtht 1740  
thttataactt ctgtaagagc ataatcaaac aggaathttc thttctcagt ggataataca 1800  
acagagaaaa cagagthgcc caaatatth aaagaagtta thccttgaga agthcatatt 1860  
thgtgacatc tgcattgatt tcagtatthc tgatggthct gthattcata agthcatatta 1920  
acathctctc cgtgaaatca tggthacagtc actgcccaga gthactgagg aaaagcaata 1980  
thggthtcggc agatggthgg gthaaaatga atctthagga gthgtggthaa thgtgthctc 2040  
gctthtthgtg catcactatg tgaagthctg thttgcagaa thggcaaaaag cgtthathtt 2100  
thaaaatgca aathaththgt acaatgthac ththtgcttc caaataataa thgtatgthtag 2160  
acagcaagaa atgaathctt thaaaagthg thththgtgg agththaaag aathacacta 2220  
aggagagthg thaaatgthg acctthgttc agthgthaaag thggaaagct aagaaathct 2280

caccgaaact tactgctgaa tgattacatt ctcccttaag cagaaaactt tggatgtgcc 2340  
 atgcaatggg gtctgtgtaa ttatcttgct ctttgattaa aaaaaagacc cccagcaata 2400  
 aaaagtgggt cactctatgc c 2421

<210> 11  
 <211> 2453  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 gaattcaaag tggagtaccg caaacttgat atggaaaata aaaagaaaga caaggacaaa 60  
 tcagatgata gaatggcacg acctagtggg cgatcgggac acaacactcg aggaactggg 120  
 tcttcatcgt ctggagtttt aatggttgga cctaacttta gagttgaaa aaaaattgga 180  
 tgtggcaatt ttggagaatt acgattaggg aaaaatttat acacaaatga atatgtggca 240  
 attaagttgg agcccatgaa atcaagagca ccacagctac atttggata cagattctat 300  
 aagcagttag gatctggaga tgggtatacct caagtttact atttcggccc ttgtggtaaa 360  
 tacaatgcta tgggtgctgga actgctggga cctagtttgg aagacttgtt tgacttgtgt 420  
 gacagaacat tttctcttaa aacagttctc atgatagcta tacaactgat ttctcgcag 480  
 gaatatgtcc attcaaagaa cttgatatac agagatgtaa aacctgagaa cttcttaata 540  
 ggacgaccaa gaaacaaaac ccagcaagtt attcacatta tagatcttgg tttggcaaag 600  
 gaatatattg atccggagac aaagaaacac ataccataca gagaacacaa gagccttaca 660  
 ggaacagcta gatatatgag cataaacaca catttaggaa aagaacaaag tagaagagac 720  
 gatttagaag ctttaggtca tatgttcatg tattttctga gaggcagtct tccttggcaa 780  
 ggcttaaagg ctgacacatt aaaggagagg tatcagaaaa ttggagatac aaaacgggct 840  
 acaccaatag aagtgttatg tgaaaatfff ccagaaatgg caacatatct tcgttatgta 900  
 agaaggctag atttttttga aaaaccagac tatgaatact taagaaagct ttttactgac 960  
 ttgtttgatc gaaaaggata tatgtttgat tatgaatatg actggattgg taaacagttg 1020  
 cctactccag tgggtgcagt tcagcaagat cctgctctgt catcaaacag agaagcacat 1080  
 caacacagag ataagatgca acaatccaaa aaccagtcgg cagaccacag ggcagcttgg 1140  
 gactcccagc aggcaaacc ccaccattg agagctcacc ttgcagcaga cagacatggg 1200  
 ggctcggtag aggttgtaag ttctacaaat ggagagttaa acacagatga ccccaccgca 1260  
 ggacgttcaa atgcacccat cacagcccct actgaagtag aagtgatgga tgaaccaaac 1320  
 tgccagaaag tgttgaacat gtgggtctgc tgctttttca aacgaaggaa aaggaaaacc 1380  
 atacagcgcc acaaatgact ctggacacag acagatcctg gggagtact tacatgttca 1440  
 tctgctgtct tgtgattaaa atcatctctg tagtgaccac gtatattttc aaggactcac 1500

tcttagaaac aaaaatgtca tactatcata cttcattttg tggttgtcctt acattcctttt 1560  
 tctttttttt tttctcctaat ttaaccttta tggaagcttt aaagttttgt caaaacatga 1620  
 gtgctttgcc catcagtgaa tggaatggac caatgagggtg gtatcaatga atatagttcc 1680  
 atagaacatt ttccagaagt tcttctggtg tagaaagcag tacagtatct taagtgtcaa 1740  
 ccagttatat acctaactctg gttttttata acttctgtaa gagcataatc aacaggaat 1800  
 tttcttttct cagtggataa tacaacagag aaaacagagt tgcccaaata tttaaaagaa 1860  
 gttattcctt gagaagttca tattttgtga catctgcatt gatttcagta ttactgatgg 1920  
 tactgttatt cataagtcat attaacattc tctccgtgaa atcatggtag agtcaactgcc 1980  
 cagaggtact gaggaaaagc aatatgggtt cggcagatgg tgggtgtaaa atgaatctta 2040  
 aggagtgtgg taaatatgtg ctccgctttt gttgcatcac tatgtgaagt actgtgttgc 2100  
 agaagtggca aaagcgctta tttttaaaaa tgcaaaatat ttgtacaatg taactttatg 2160  
 cttccaaaata ataatgtatg ttagacagca agaaatgaat actttaaaaa gtgatatatg 2220  
 ttggagttat aaagaaatac actaaggaga ggtagtaa atgtgaaccttg ttgcagtgtg 2280  
 taaggtggaa gcctaaagaa atctcaccga aacttactgc tgaatgatta cattctcctt 2340  
 taagcagaaa actttggatg tgccatgcaa tgggtgtctgt gtaattattt tgctctttga 2400  
 ttaaaaaaaaa gacccccagc aataaaaagt gggtcactct atgccctctg tgc 2453

<210> 12

<211> 2717

<212> DNA

<213> Homo sapiens

<400> 12

gctttcctcc tccggccgcc ggcgggtgtg atgtgccgcc gccgctgccc ccgcccggcg 60  
 tgacggactg gggcgccgc ccgccccgg gaccgacccc tctgctcggg gccgcgcctt 120  
 tgagctctct atcaatatca gtcacatca ttgaaaagat aattttgaag acatgttttg 180  
 ctgaaaagac actaagaaaa attttacgaa tgggatgaac atgctccagt taattgacta 240  
 cctactgcaa tttgaatggt aacattacc atctggtaca gttacctagt gatgtaccta 300  
 ttttcacaat accctgtttc agtgtgcttg tcttgattaa agaattcaaa gtggagtacc 360  
 gcaaacttga tatggaaaat aaaaagaaag acaaggacaa atcagatgat agaatggcac 420  
 gacctagtgg tcgatcggga cacaacactc gaggaactgg gtcttcatcg tctggagttt 480  
 taatggttgg acctaacttt agagttggaa aaaaaattgg atgtggcaat tttggagaat 540  
 tacgattagg gaaaaattta tacacaaatg aatatgtggc aattaagttg gagcccatga 600  
 aatcaagagc accacagcta catttggaa acagattcta tgagcagtta ggatctggag 660

atggtatacc tcaagtttac tatttcggcc cttgtggtaa atacaatgct atggtgctgg	720
aactgctggg acctagtttg gaagacttgt ttgacttgtg tgacagaaca ttttctctta	780
aaacagttct catgatagct atacaactga tttctcgcat ggaatatgtc cattcaaaga	840
acttgatata cagagatgta aaacctgaga acttcttaat aggacgacca ggaaacaaaa	900
cccagcaagt tattcacatt atagattttg gtttggcaaa ggaatatatt gatccggaga	960
caaagaaaca cataccatac agagaacaca agagccttac aggaacagct agatataatga	1020
gcataaacac acatttagga aaagaacaaa gtagaagaga cgatttagaa gctttaggtc	1080
atatgttcat gtattttctg agaggcagtc ttccttggca aggcttaaag gctgacacat	1140
taaaggagag gtatcagaaa attggagata caaacgggc tacaccaata gaagtgttat	1200
gtgaaaattt tccagaaatg gcaacatatac ttcgttatgt aagaaggcta gatTTTTTTTg	1260
aaaaaccaga ctatgactac ttaagaaagc tttttactga cttgtttgat cgaaaaggat	1320
atatgtttga ttatgaatat gactggattg gtaaacagtt gcctactcca gtgggtgcag	1380
ttcagcaaga tctgctctg tcatcaaaca gagaagcaca tcaacacaga gataagatgc	1440
aacaatccaa aaaccaggtt gtaagttcta caaatggaga gttaaacaca gatgacccca	1500
ccgcaggacg ttcaaatgca cccatcacag cccctactga agtagaagtg atggatgaaa	1560
ccaactgcc aagagtgtg aacatgtggt gctgctgttt tttcaaacga aggaaaagga	1620
aaaccataca gcgccacaaa tgactctgga cacagacaga tcctgggggag ttacttacat	1680
gttcatctgc tgtcttgtga ttaaaatcat ctctgtagtg accacgtata ttttcaagga	1740
ctcactctta gaaacaaaaa tgcataactt tcatacttca ttttgtgggt gtcttacatt	1800
ctttttcttt tttttttttt ctctaattta acatttatgg aagctttaaa gttttgtcaa	1860
aacatgagtg ctttgcccat cagtgaatgg aatggaccaa tgagggtgga tcaatgaata	1920
tagttccata gaacattttc cagaagttct tctgtttag aaagcagtac agtatcttaa	1980
gtgtcaacca gttatatacc taatctggtt ttttataact tctgtaagag cataatcaaa	2040
caggaatttt cttttctcag tggataatac aacagagaaa acagagttgc ccaaatattt	2100
aaaagaagtt attccttgag aagttcatat tttgtgacat ctgcattgat ttcagtatta	2160
ctgatggtac tgttattcat aagtcataat aacattctct ccgtgaaatc atggtacagt	2220
cactgcccag aggtactgag gaaaaagcaa tatgggttcg gcagatggtg gtggtaaaat	2280
gaatcttaag gagtgtggta aatatgtgct ccgcttttgt tgcataccta tgtgaagtac	2340
tgtgttcag aagtggcaaa agcgcttatt tttaaaaatg caaaatattt gtacaatgta	2400
actttatgct tccaaataat aatgtatggt agacagcaag aaatgaatac tttaaaaagt	2460
gatgtatggt ggagttataa agaaatacac taaggagagg tagtaaagt gaaccttgtt	2520

gcagtgtata aggtggaagc ctaaagaaat ctcaccaaaa cttactgctg aatgattaca 2580  
 ttctccctta agcagaaaac tttggatgtg ccatgcaatg gtgtctgtgt aattatattg 2640  
 ctctttgatt aaaaaaaga cccccagcaa taaaaagtgg gtcactctat gccctctgtg 2700  
 caaaaaaaaa aaaaaaa 2717

<210> 13  
 <211> 393  
 <212> PRT  
 <213> Homo sapiens

<400> 13

Met Asp His Pro Ser Arg Glu Lys Asp Glu Arg Gln Arg Thr Thr Lys  
 1 5 10 15

Pro Met Ala Gln Arg Ser Ala His Cys Ser Arg Pro Ser Gly Ser Ser  
 20 25 30

Ser Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val Gly Lys  
 35 40 45

Lys Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys Asn Leu  
 50 55 60

Tyr Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Ile Lys Ser Arg  
 65 70 75 80

Ala Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu Gly Ser  
 85 90 95

Ala Gly Glu Gly Leu Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly Lys  
 100 105 110

Tyr Asn Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp Leu  
 115 120 125

Phe Asp Leu Cys Asp Arg Thr Phe Thr Leu Lys Thr Val Leu Met Ile  
 130 135 140

Ala Ile Gln Leu Leu Ser Arg Met Glu Tyr Val His Ser Lys Asn Leu  
 145 150 155 160

Ile Tyr Arg Asp Val Lys Pro Glu Asn Phe Leu Ile Gly Arg Gln Gly  
 165 170 175

Asn Lys Lys Glu His Val Ile His Ile Ile Asp Phe Gly Leu Ala Lys  
 180 185 190

Glu Tyr Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu His  
 195 200 205

Lys Ser Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His Leu  
 210 215 220

Gly Lys Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His Met  
 225 230 235 240

Phe Met Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys Ala  
 245 250 255

Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg Asn  
 260 265 270

Thr Pro Ile Glu Ala Leu Cys Glu Asn Phe Pro Glu Glu Met Ala Thr  
 275 280 285

Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp Tyr  
 290 295 300

Glu Tyr Leu Arg Thr Leu Phe Thr Asp Leu Phe Glu Lys Lys Gly Tyr  
 305 310 315 320

Thr Phe Asp Tyr Ala Tyr Asp Trp Val Gly Arg Pro Ile Pro Thr Pro  
 325 330 335

Val Gly Ser Val His Val Asp Ser Gly Ala Ser Ala Ile Thr Arg Glu  
 340 345 350

Ser His Thr His Arg Asp Arg Pro Ser Gln Gln Gln Pro Leu Arg Asn  
 355 360 365

Gln Ser Leu Arg Thr Val Thr Ala Glu His Tyr Asp Val Asn Asn Ser  
 370 375 380

Ala Ile Trp His Arg Gly Arg Gly Thr  
 385 390

<210> 14  
 <211> 415  
 <212> PRT  
 <213> Homo sapiens

<400> 14

Met Asp Phe Asp Lys Lys Gly Gly Lys Gly Glu Thr Glu Glu Gly Arg



Ala Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg  
 260 265 270

Ala Thr Pro Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Glu Met Ala  
 275 280 285

Thr Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp  
 290 295 300

Tyr Asp Tyr Leu Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Ser Gly  
 305 310 315 320

Phe Val Phe Asp Tyr Glu Tyr Asp Trp Ala Gly Lys Pro Leu Pro Thr  
 325 330 335

Pro Ile Gly Thr Val His Thr Asp Leu Pro Ser Gln Pro Gln Leu Arg  
 340 345 350

Asp Lys Thr Gln Pro His Ser Lys Asn Gln Ala Leu Asn Ser Thr Asn  
 355 360 365

Gly Glu Leu Asn Ala Asp Asp Pro Thr Ala Gly His Ser Asn Ala Pro  
 370 375 380

Ile Thr Ala Pro Ala Glu Val Glu Val Ala Asp Glu Thr Lys Cys Cys  
 385 390 395 400

Cys Phe Phe Lys Arg Arg Lys Arg Lys Ser Leu Gln Arg His Lys  
 405 410 415

<210> 15  
 <211> 415  
 <212> PRT  
 <213> Homo sapiens

<400> 15

Met Asp Phe Asp Lys Lys Gly Gly Lys Gly Glu Thr Glu Glu Gly Arg  
 1 5 10 15

Arg Met Ser Lys Ala Gly Gly Gly Arg Ser Ser His Gly Ile Arg Ser  
 20 25 30

Ser Gly Thr Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val  
 35 40 45

Gly Lys Lys Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys  
 50 55 60

Asn Leu Tyr Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Ile Lys  
 65 70 75 80

Ser Arg Ala Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu  
 85 90 95

Ser Ala Thr Glu Gly Val Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly  
 100 105 110

Asn Tyr Asn Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp  
 115 120 125

Leu Phe Asp Leu Cys Asp Arg Thr Phe Thr Leu Lys Thr Val Leu Met  
 130 135 140

Ile Ala Ile Gln Leu Ile Thr Arg Met Glu Tyr Val His Thr Lys Ser  
 145 150 155 160

Leu Ile Tyr Arg Asp Val Lys Pro Glu Asn Phe Leu Val Gly Arg Pro  
 165 170 175

Gly Thr Lys Arg Gln His Ala Ile His Ile Ile Asp Phe Gly Leu Ala  
 180 185 190

Lys Glu Tyr Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu  
 195 200 205

His Lys Ser Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His  
 210 215 220

Leu Gly Lys Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His  
 225 230 235 240

Met Phe Met Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys  
 245 250 255

Ala Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg  
 260 265 270

Ala Thr Pro Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Glu Met Ala  
 275 280 285

Thr Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp  
 290 295 300

Tyr Asp Tyr Leu Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Ser Gly  
 305 310 315 320

Phe Val Phe Asp Tyr Glu Tyr Asp Trp Ala Gly Lys Pro Leu Pro Thr  
 325 330 335

Pro Ile Gly Thr Val His Thr Asp Leu Pro Ser Gln Pro Gln Leu Arg  
 340 345 350

Asp Lys Thr Gln Pro His Ser Lys Asn Gln Ala Leu Asn Ser Thr Asn  
 355 360 365

Gly Glu Leu Asn Ala Asp Asp Pro Thr Ala Gly His Ser Asn Ala Pro  
 370 375 380

Ile Thr Ala Pro Ala Glu Val Glu Val Ala Asp Glu Thr Lys Cys Cys  
 385 390 395 400

Cys Phe Phe Lys Arg Arg Lys Arg Lys Ser Leu Gln Arg His Lys  
 405 410 415

<210> 16  
 <211> 447  
 <212> PRT  
 <213> Homo sapiens

<400> 16

Met Glu Asn Lys Lys Lys Asp Lys Asp Lys Ser Asp Asp Arg Met Ala  
 1 5 10 15

Arg Pro Ser Gly Arg Ser Gly His Asn Thr Arg Gly Thr Gly Ser Ser  
 20 25 30

Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val Gly Lys Lys  
 35 40 45

Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys Asn Leu Tyr  
 50 55 60

Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Met Lys Ser Arg Ala  
 65 70 75 80

Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu Gly Ser Gly  
 85 90 95

Asp Gly Ile Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly Lys Tyr Asn  
 100 105 110

Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp Leu Phe Asp  
 115 120 125

Leu Cys Asp Arg Thr Phe Ser Leu Lys Thr Val Leu Met Ile Ala Ile  
 130 135 140

Gln Leu Ile Ser Arg Met Glu Tyr Val His Ser Lys Asn Leu Ile Tyr  
 145 150 155 160

Arg Asp Val Lys Pro Glu Asn Phe Leu Ile Gly Arg Pro Arg Asn Lys  
 165 170 175

Thr Gln Gln Val Ile His Ile Ile Asp Phe Gly Leu Ala Lys Glu Tyr  
 180 185 190

Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu His Lys Ser  
 195 200 205

Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His Leu Gly Lys  
 210 215 220

Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His Met Phe Met  
 225 230 235 240

Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys Ala Asp Thr  
 245 250 255

Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg Ala Thr Pro  
 260 265 270

Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Met Ala Thr Tyr Leu Arg  
 275 280 285

Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp Tyr Glu Tyr Leu  
 290 295 300

Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Lys Gly Tyr Met Phe Asp  
 305 310 315 320

Tyr Glu Tyr Asp Trp Ile Gly Lys Gln Leu Pro Thr Pro Val Gly Ala  
 325 330 335

Val Gln Gln Asp Pro Ala Leu Ser Ser Asn Arg Glu Ala His Gln His  
 340 345 350

Arg Asp Lys Met Gln Gln Ser Lys Asn Gln Ser Ala Asp His Arg Ala

355

360

365

Ala Trp Asp Ser Gln Gln Ala Asn Pro His His Leu Arg Ala His Leu  
370 375 380

Ala Ala Asp Arg His Gly Gly Ser Val Gln Val Val Ser Ser Thr Asn  
385 390 395 400

Gly Glu Leu Asn Thr Asp Asp Pro Thr Ala Gly Arg Ser Asn Ala Pro  
405 410 415

Ile Thr Ala Pro Thr Glu Val Glu Val Met Asp Glu Thr Lys Cys Cys  
420 425 430

Cys Phe Phe Lys Arg Arg Lys Arg Lys Thr Ile Gln Arg His Lys  
435 440 445