

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 2408802 T3**

-
- (51) Int.Cl.: *C 07 K 1/00 (2006.01)* *C 07 D 487/04 (2006.01)* *C 07 D 487/14 (2006.01)*
C 07 F 7/18 (2006.01) *C 07 K 5/02 (2006.01)* *C 07 K 5/12 (2006.01)*
C 07 K 7/64 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2017-08-28**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-05-03**
- (86) Europæisk ansøgning nr.: **10753047.9**
- (86) Europæisk indleveringsdag: **2010-03-16**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-01-25**
- (86) International ansøgning nr.: **CA2010000408**
- (87) Internationalt publikationsnr.: **WO2010105363**
- (30) Prioritet: **2009-03-16 US 160571 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **The Governing Council of the University of Toronto, 27 King's College Circle, Toronto, ON M5S 3G4, Canada**
- (72) Opfinder: **YUDIN, Andrei, 561 Wingrove Crescent, Oakville, Ontario L6L 4R7, Canada**
HILI, Ryan, 10 Smith Avenue, Apt 3, Summerville, Massachusetts 02143, USA
- (74) Fuldmægtig i Danmark: **Orsnes Patent ApS, Forskerparken 10, 5230 Odense M, Danmark**
- (54) Benævnelse: **CYKLISKE AMINOSYREMOLEKYLER OG FREMGANGSMÅDER TIL FREMSTILLING HERAF**
- (56) Fremdragne publikationer:
WO-A1-2008/046232
OTILIE E. VERCILLO ET AL.: "Design and Synthesis of Cyclic RGD Pentapeptoids by Consecutive Ugi Reactions", ORGANIC LETTERS, vol. 10, no. 2, 1 January 2008 (2008-01-01), pages 205-208, XP055055694, ISSN: 1523-7060, DOI: 10.1021/ol702521g
HILI, R. ET AL.: 'Macrocyclization of linear peptides enabled by amphoteric molecules' J. AM. CHEM. SOC. vol. 132, 15 February 2010, pages 2889 - 2891, XP055055687
VERCILLO, O.E. ET AL.: 'Design and synthesis of cyclic RGD pentapeptoids by consecutive Ugi reactions' ORG. LETT. vol. 10, 19 December 2007, pages 205 - 208, XP055055694
BAKTHARAMAN, S. ET AL.: 'Amino carbonyl compounds in organic synthesis' ALDRZCHIMICA ACTA vol. 41, 2008, pages 109 - 117, XP008166926
YUDIN, A.K. ET AL.: 'Overcoming the demons of protecting groups with amphoteric molecules' CHEMISTRY: EUR. J. vol. 13, 10 July 2007, pages 6539 - 6542, XP055098088
HILI, R. ET AL.: 'Readily available unprotected amino aldehydes' J. AM. CHEM. SOC. vol. 128, 27 October 2006, pages 14772 - 14773, XP055098301
Tomoyasu Hirose ET AL.: "Total Synthesis and Determination of the Absolute Configuration of Guadinomines B

Fortsættes ...

and C 2", Chemistry - A European Journal, vol. 14, no. 27, 19 September 2008 (2008-09-19), pages 8220-8238, XP55113663, ISSN: 0947-6539, DOI: 10.1002/chem.200801024

LAURA QUARTARA ET AL: "Influence of lipophilicity on the biological activity of cyclic pseudopeptide NK-2 receptor antagonists", JOURNAL OF MEDICINAL AND PHARMACEUTICAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, US, vol. 37, 1 January 1994 (1994-01-01), pages 3630-3638, XP002198370, DOI: 10.1021/JM00047A020

ACHMATOWICZ M S ET AL: "The synthesis of l-proline derived hexaazamacrocyclic ligands of C3 symmetry via intramolecular methyl ester aminolysis", TETRAHEDRON ASYMMETRY, PERGAMON PRESS LTD, OXFORD, GB, vol. 12, no. 3, 5 March 2001 (2001-03-05), pages 487-495, XP004233491, ISSN: 0957-4166, DOI: 10.1016/S0957-4166(01)00068-4

MURRAY P J ET AL: "The synthesis of cyclic tetrapeptoid analogues of the antiprotozoal natural product apicidin", BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, PERGAMON, AMSTERDAM, NL, vol. 11, no. 6, 26 March 2001 (2001-03-26), pages 773-776, XP004230929, ISSN: 0960-894X, DOI: 10.1016/S0960-894X(01)00049-X

John E Burden ET AL: "Synthesis and biological activities of ykfa analogues: effects of position 4 substitutions and altered ring size on In vitro opioid activity", Bioorganic & Medicinal Chemistry Letters, vol. 12, no. 2, 1 January 2002 (2002-01-01), pages 213-216, XP55113667, ISSN: 0960-894X, DOI: 10.1016/S0960-894X(01)00706-5

Joost Pijl ET AL: "Synthesis and electrophysiological characterization of cyclic morphiceptin analogues", Biochemical Pharmacology, vol. 67, no. 10, 1 May 2004 (2004-05-01), pages 1887-1895, XP55113668, ISSN: 0006-2952, DOI: 10.1016/j.bcp.2004.02.011

DESCRIPTION

FIELD

[0001] The present invention relates to cyclic amino acid molecules and methods of preparing the same, and in particular the macrocyclization of amino acids or linear peptides upon reaction with amphoteric amino aldehydes and isocyanides.


BACKGROUND

[0002] Peptides play vital roles by mediating a wide range of biological processes, acting as hormones, antibiotics, and signaling molecules. Due to the highly specific interaction with their biological targets, peptides have been widely used in medicine. However, the enormous therapeutic potential of peptides is not always easy to realize due to their low bioavailability. This shortcoming is a consequence of the degradation of peptides by endo- and exopeptidases, which results in poor in vivo stability of peptides. Compared to their linear counterparts, cyclic peptides are more resistant to degradation. There are two main reasons for this stability. Firstly, exopeptidases cannot cleave the cyclic peptide at its (non-existent) ends. Secondly, cyclic peptides, especially those with a small-to-medium ring size, are protected against endopeptidases because the constrained cyclic peptide backbone prevents the adaptation of the required extended conformation during proteolysis. In addition, the reduced charge and intramolecular hydrogen bonding within cyclic peptides facilitate passive membrane permeability, which contributes to their enhanced bioavailability. Most significantly, conformational constraints imposed on the amino acid sequence by the cyclic topology maximize enthalpic interactions between cyclic peptides and their biochemical targets while ensuring favourable entropy of binding.

[0003] There has been enormous interest in both naturally occurring and synthetic cyclic peptides as scaffolds that pre-organize an amino acid sequence into a rigid conformation.¹ Amongst the vast number of known cyclic peptides, rigid small-to-medium sized rings have been of particular interest. Various cyclolactamization and non-peptidic cyclization methods² have been developed.

[0004] The macrocyclization of linear precursors is afflicted by several thermodynamic and kinetic challenges that arise from the conformational preferences of linear peptides. The chain/ring conformational equilibrium is the central obstacle facing synthesis of cyclic molecules from acyclic precursors. Short linear peptides can easily adopt a circular conformation, which is driven by ion pairing between the N- and C-termini (Scheme 1, **A**).³ Despite the unfavorable entropy, these circular conformations are thermodynamically favoured due to the enthalpy garnered through electrostatic and other polar interactions. As shown in

Scheme 1, conventional activation reagents tend to remove the zwitterionic character of the peptide, rendering it incapable of forming ion pairs. Consequently, without enthalpic contribution from electrostatics and other polar interactions, the activated peptide adopts a random linear conformation (Scheme 1, **B**). In order for macrocyclization to occur, the activated peptide must adopt a pre-cyclization conformation (**C**) prior to forming the desired cyclic molecule (**D**). High dilution, on the order of 10^{-4} or greater, is essential to limiting the formation of by-products arising from cyclodimerization,⁴ cyclotrimerization, and polymerization.⁵ Unfortunately, dilution brings about long reaction times, which in turn provoke background processes such as epimerization. Amongst the most challenging cyclizations are those attempted on linear peptides containing less than seven residues.^{6,7}




Scheme 1. A common peptide macrocyclization strategy.

[0005] Another common challenge in exploring macrocyclic chemistry space has to do with late-stage modification. This is a historic challenge for macrocyclic compound libraries built in the biotechnology and pharmaceutical companies. Their typical cyclization techniques (ring-closing metathesis, Huisgen cycloaddition) do not naturally lend themselves to further elaboration. Functional group handles must be built in prior to cyclization to achieve this goal.

SUMMARY OF THE INVENTION

[0006] Described herein is a method of forming a cyclic peptide or amino acid comprising the steps of:


1. (i) reacting an amine end group of a peptide with an aziridine aldehyde having the following structure:



and

2. (ii) reacting the mixture of step (i) with an isocyanide.

[0007] In further examples described herein there is provided a process to produce a cyclic amino acid molecule comprising reacting an amino acid molecule, having an amino terminus and a carboxyl terminus, with an isocyanide and a compound having formula (Ia) and/or (Ib):





wherein:

$n = 0$ or 1 , and R_1, R_2, R_3, R_4 and R_5 are independently selected from H; lower alkyl; alkenyl; heterocycle; cycloalkyl; esters of the formula $-C(O)OR^*$ wherein R is selected from alkyl and aryl; amides of the formula $-C(O)NR$, wherein R and R are independently selected from alkyl and aryl; $-CH_2C(O)R$, wherein R is selected from $-OH$, lower alkyl, aryl, $-lower\ alkyl-aryl$, or $-NR_3R_b$, where R_3 and R_b are independently selected from H, lower alkyl, aryl or $-lower\ alkyl-aryl$; $-C(O)R_c$, wherein R_o is selected from lower alkyl, aryl or $-lower\ alkyl-aryl$; or $-lower\ alkyl-OR_a$, wherein R_d is a suitable protecting group or OH group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents; and the aldehyde component thereof may optionally be in its bisulfite adduct form; and the amino acid molecule is an amino acid, a linear peptide or a salt of the foregoing, provided that if the amino acid molecule is a linear peptide, the compound comprises an aziridine chiral center proximal to the aldehyde with matching stereochemistry to the carbon atom proximal to the amino terminus of the peptide.

[0008] In further examples described herein, there is provided a cyclic amino acid molecule prepared using further examples of processes described herein.

[0009] In further examples described herein, there is provided a cyclic amino acid molecule of formula (II):





wherein,

$n = 0$ or 1 , and R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from H; lower alkyl; aryl; heteroaryl; alkenyl; heterocycle; esters of the formula $-C(O)OR^*$ wherein R^* is selected from alkyl and aryl; amides of the formula $-C(O)NR^{**}R^{***}$, wherein R^{**} and R^{***} are independently selected from alkyl and aryl; $-CH_2C(O)R$, wherein R is selected from $-OH$, lower alkyl, aryl, -loweralkyl- aryl, or $-NR_3R_b$, where R_3 and R_b are independently selected from H, lower alkyl, aryl or -loweralkyl-aryl; $-C(O)R_0$, wherein R_0 is selected from lower alkyl, aryl or -lower alkyl-aryl; or -lower alkyl- OR_d , wherein R_j is a suitable protecting group or OH group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents,


bonds [a] and [b] are syn to each other;

R' is an amino acid side chain of the amino terminus amino acid;

R'' is an optionally substituted amide;

and the amino acid molecule is an amino acid or a linear peptide, wherein N' is the nitrogen at the amino terminus end of the amino acid molecule and C is the carbon at the carboxy terminus end of the amino acid molecule, and provided that if the amino acid molecule is a linear peptide, bonds [a] and [c] are anti to each other.

[0010] In further examples described herein, there is provided use of a compound for cyclizing an amino acid molecule, wherein the compound has the formula (Ia)/(Ib):



comprises an α -stereocenter at the carbon atom proximal to the aldehyde group with matching stereochemistry to the carbon atom proximal to the amino terminus of the amino acid molecule; and

$n = 0$ or 1 , and R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from H; lower alkyl; aryl; heteroaryl; alkenyl; heterocycle; esters of the formula $-C(O)OR^*$ wherein R^* is selected from alkyl and aryl; amides of the formula $-C(O)NR^*R^*$, wherein R and R are independently selected from alkyl and aryl; $-CH_2C(O)R$, wherein R is selected from $-OH$, lower alkyl, aryl, -loweralkyl-aryl, or $-NR_aR_b$, where R_a and R_b are independently selected from H, lower alkyl, aryl or -loweralkyl-aryl; $-C(O)R_c$, wherein R

is

selected from lower alkyl, aryl or -loweralkyl-aryl; or -lower alkyl- OR_d , wherein R_d is a suitable protecting group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents; and the amino acid molecule is an amino acid,

a linear peptide or a salt of the foregoing, provided that if the amino acid molecule is a linear peptide, the compound comprises an aziridine chiral center proximal to the aldehyde with matching stereochemistry to the carbon atom proximal to the amino terminus of the peptide.

BRIEF DESCRIPTION OF THE FIGURES:

[0011] Embodiments of the invention may best be understood by referring to the following description and accompanying drawings. In the description and drawings, like numerals refer to like structures or processes. In the drawings:

FIG. 1 is a ^1H NMR comparison of crude reaction mixtures using isomers of leucine.

FIG. 2 is a scheme depicting a mismatched reaction leading to amination products.

FIG. 3 is a matched reaction leading to a cyclic product.

FIG. 4 shows the chemical structures of the cyclic products synthesized by Methods 1 - 3.

FIG. 5 is the ^1H and ^{13}C NMR spectra for cyclic product 1.

FIG. 6 is the ^1H and ^{13}C NMR spectra for cyclic product 2.

FIG. 7 is the ^1H and ^{13}C NMR spectra for cyclic product 3.

FIG. 8 is the ^1H and ^{13}C NMR spectra for cyclic product 4.

FIG. 9 is the ^1H and ^{13}C NMR spectra for cyclic product 5.

FIG. 10 is the ^1H and ^{13}C NMR spectra for cyclic product 6.

FIG. 11 is the ^1H and ^{13}C NMR spectra for cyclic product 7.

FIG. 12 is the ^1H and ^{13}C NMR spectra for cyclic product 8.

FIG. 13 is the ^1H NMR spectrum for cyclic product 9.

FIG. 14. is the ^1H and ^{13}C NMR spectra for cyclic product **10**.

FIG. 15 is the ^1H and ^{13}C NMR spectra for cyclic product **11**.

FIG. 16 is the ^1H and ^{13}C NMR spectra for cyclic product **12**.

FIG. 17 is the ^1H and ^{13}C NMR spectra for cyclic product **13**.

FIG. 18 is the ^1H and ^{13}C NMR spectra for cyclic product **14**.

FIG. 19 is the ^1H and ^{13}C NMR spectra for cyclic product **15**.

FIG. 20 is the ^1H , COSY, and ^{13}C NMR spectra for cyclic product **16**.

FIG. 21 is the ^1H and ^{13}C NMR spectra for cyclic product **17**.

FIG. 22 is the ^1H and ^{13}C NMR spectra for cyclic product **18**.

FIG. 23 is the ^1H and ^{13}C NMR spectra for cyclic product **19**.

FIG. 24 is the ^1H and ^{13}C NMR spectra for cyclic product **20**.

FIG. 25 is the ^1H and ^{13}C NMR spectra for cyclic product **21**.

FIG. 26 is the ^1H and COSY NMR spectra for cyclic product **22**.

FIG. 27 is the ^1H NMR spectrum for cyclic product **23**.

FIG. 28 depicts a crude LC-MS(ESI) analysis of HATU-mediated cyclization of Pro-Gly-Gly-Gly at 0.2M.

FIG. 29 depicts a crude LC-MS(ESI) analysis of isocyanide/amino aldehyde-mediated cyclization of Pro-Gly-Gly-Gly at 0.2M.

FIG. 30 shows the nucleophilic ring-opening of aziridine-containing cyclic peptides. FIG. 31 depicts the pK_a 's of thiols and ammonium ions and ΔpK_a of ion pairs.

FIG. 32 depicts a crude LC-MS(ESI) analysis for a reaction mixture of thiol opened product.


DETAILED DESCRIPTION

[0012] There is described herein a one-step process that provides cyclic peptides in high yields and selectivities while evading the problems typically encountered during traditional cyclization reactions. The late stage diversification of macrocyclic molecules can now be achieved in a seamless fashion. The products of the macrocyclization are equipped with specific modification sites, which enable late-stage structural modification of cyclic peptides. The post cyclization diversification has been a historic challenge for macrocycle libraries in both biotechnology and pharmaceutical industries. The present macrocyclization approach solves this problem, among others.

[0013] There is demonstrated herein a use for a class of stable amphoteric amino aldehydes that contain synthetically useful unprotected aziridine and aldehyde groups.⁸ Amphoteric amino

aldehydes are coaxed into undergoing a reversible reaction with the amine and carboxyl ends of the peptide. The resulting electrophilic intermediate rapidly reacts with the nucleophilic aziridine portion due to the high effective molarity. As a result, the unfavorable chain/ring equilibrium is shifted using the Le Chatelier principle, securing efficient conversion to cyclic products. In another aspect, the cyclization tolerates aliphatic, acidic, and basic amino acid side chain residues. In particular, peptides, including those containing both acidic and basic residues, are found to be within the scope of this process, delivering a single diastereomeric product in each case. Trifluoroethanol ("TFE"), known for its capacity to stabilize peptide secondary structure and promote polar interactions,⁹ is the preferred reaction medium. The Ugi four-component condensation, a well-known reaction involving carboxylic acids, amines, aldehydes, and isocyanides¹⁰, is the preferred means of generating zwitterionic macrocyclization precursors. Mechanistically, the Ugi reaction is known to proceed through a series of reversible transformations that tend towards the thermodynamic driving force of amide bond formation. When an α -amino acid, aldehyde, and isocyanide are used as starting materials, the reaction traverses a zwitterionic iminium ion intermediate, which upon attack by the isocyanide produces an electrophilic mixed anhydride. Subsequent reaction with methanol gives a linear peptide ester as a mixture of diastereoisomers.¹¹ This reaction has been attempted in the synthesis of cyclic peptides from linear peptides and conventional (monofunctional) aldehydes. However, low yields are accompanied by lack of diastereoselectivity and dominant cyclodimerization products.^{12,13}

[0014] In further examples described herein, there is provided a process to produce a cyclic amino acid molecule comprising reacting an amino acid molecule, having an amino terminus and a carboxyl terminus, with an isocyanide and a compound having formula (Ia) and/or (Ib):



wherein:

$n = 0$ or 1 , and R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from H; lower alkyl; alkenyl; heterocycle; cycloalkyl; esters of the formula $-C(O)OR^*$ wherein R^* is selected from alkyl and aryl; amides of the formula $-C(O)NR^{**}R^{***}$, wherein R^{**} and R^{***} are independently selected from alkyl and aryl; $-CH_2C(O)R$, wherein R is selected from $-OH$, lower alkyl, aryl, $-lower\ alkyl-aryl$, or $-NR_3R_b$, where R_a and R_b are independently selected from H, lower alkyl, aryl or $-lower\ alkyl-aryl$; $-C(O)R_c$, wherein R_o is selected from lower alkyl, aryl or $-lower\ alkyl-aryl$; or $-lower\ alkyl-OR_d$, wherein R_a is a suitable protecting group or OH group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents; and the aldehyde component thereof may optionally be in its bisulfite adduct form; and the amino acid molecule is an amino acid, a linear peptide or a salt of the foregoing, provided that if the amino acid molecule is a linear peptide, the compound comprises an

aziridine chiral center proximal to the aldehyde with matching stereochemistry to the carbon atom proximal to the amino terminus of the peptide.

[0015] Examples herein describe that if the amino acid molecule is an amino acid then the amino terminus is a primary amino group or a secondary amino group but wherein the amino acid molecule is a peptide, then the amino terminus is a secondary amino group.

[0016] Also described are examples wherein any one of $R_1 - R_5$ is H. Preferably, $n=0$ and R_2 and R_3 is H or $R_1 - R_3$ is H.

[0017] Also described are examples wherein R_1 is CH_2OTBDMS or CH_2^1Pr .

[0018] Also described are examples wherein the amino acid molecule is a linear peptide. Preferably, the amino terminus amino acid of the linear peptide is selected from the group consisting of proline and an amino acid with an amino group substituted with NHBn , $\text{NHCH}_2\text{CH}_2\text{SO}_2\text{Ph}$ or $\text{NHCH}_2\text{CH}_2\text{CN}$.

[0019] Also described are examples wherein the amino acid molecule is a D or L amino acid selected from the group consisting of: alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, selenocysteine, serine, tyrosine, threonine, tryptophan and valine.

[0020] The amino acid molecule may be an alpha-amino acid, beta-amino acid or gamma-amino acid.

[0021] Also described are examples wherein the isocyanide is selected from the group consisting of: (S)-(-)- α - Methylbenzyl isocyanide; 1,1,3,3,-Tetramethylbutyl isocyanide; 1-Pentyl isocyanide; 2,6-Dimethylphenyl isocyanide; 2-Morpholinoethyl isocyanide; 2-Naphthyl isocyanide; 2-Pentyl isocyanide; 4-Methoxyphenyl isocyanide; Benzyl isocyanide; Cutyl isocyanide; Cyclohexyl isocyanide; Isopropyl isocyanide; p-Toluenesulfonylmethyl isocyanide; Phenyl isocyanide dichloride; tert-Butyl isocyanide; (Trimethylsilyl)methyl isocyanide; 1H-Benzotriazol-1-ylmethyl isocyanide; 2-Chloro-6-methylphenyl isocyanide; Di-tert-butyl 2-isocyanosuccinate; tert-Butyl 2-isocyano-3-methylbutyrate; tert-Butyl 2-isocyano-3-phenylpropionate; tert-Butyl 2-isocyanopropionate; and tert-Butyl 3-isocyanopropionate, preferably, tert-Butyl isocyanide.

[0022] Also described are examples wherein the process is conducted in a non-nucleophilic reaction medium, preferably trifluoroethanol or HFIP mixed with water.

[0023] Also described are examples wherein if the amino acid molecule is an amino acid, the process is conducted in water.


[0024] In some examples described herein, the process further comprises conjugating a fluorescent tag to the cyclic amino acid molecule by nucleophilic ring-opening of the aziridine moiety.

[0025] In some examples described herein, the peptide is between 2 and 30 amino acids in length.

[0026] In further examples described herein, there is provided a cyclic amino acid molecule prepared using examples of the process described herein.

[0027] One would understand that cyclization, in some cases, may require and would include protecting certain peptide or amino acid side chains in manner known to a person skilled in the art.

[0028] In further examples described herein, there is provided a cyclic amino acid molecule of formula (II): amino acid molecule



wherein,

$n = 0$ or 1 , and R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from H; lower alkyl; aryl; heteroaryl; alkenyl; heterocycle; esters of the formula $-C(O)OR^*$ wherein R is selected from alkyl and aryl; amides of the formula $-C(O)NR$, wherein R and R are independently selected from alkyl and aryl; $-CH_2C(O)R$, wherein R is selected from - OH, lower alkyl, aryl, -loweralkyl-aryl, or $-NR_aR_b$, where R_a and R_b are independently selected from H, lower alkyl, aryl or - loweralkyl-aryl; $-C(O)R_c$, wherein R_c is selected from lower alkyl, aryl

or -lower alkyl-aryl; or -lower alkyl- OR_d , wherein R_d is a suitable protecting group or OH group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents,

bonds [a] and [b] are syn to each other;

R' is an amino acid side chain of the amino terminus amino acid;

R'' is an optionally substituted amide;

and the amino acid molecule is an amino acid or a linear peptide, wherein N' is the nitrogen at the amino terminus end of the amino acid molecule and C is the carbon at the carboxy

terminus end of the amino acid molecule, and provided that if the amino acid molecule is a linear peptide, bonds [a] and [c] are anti to each other.

[0029] In examples described herein, if the amino acid molecule is an amino acid then the amino terminus is a primary amino group or a secondary amino group but wherein the amino acid molecule is a linear peptide, then the amino terminus is a secondary amino group.

[0030] In further examples described herein, any one of $R_1 - R_5$ is H. Preferably, $n=0$ and R_2 and R_3 is H or $R_1 - R_3$ is H.

[0031] In examples described herein, R_1 is CH_2OTBDMS or CH_2^1Pr .


[0032] In further examples described herein, the amino acid molecule is a linear peptide. Preferably, the amino terminus amino acid of the linear peptide is selected from the group consisting of proline and an amino acid with an amino group substituted with NHBn , $\text{NHCH}_2\text{CH}_2\text{SO}_2\text{Ph}$ or $\text{NHCH}_2\text{CH}_2\text{CN}$.

[0033] In further examples described herein, the amino acid molecule is a D or L amino acid selected from the group consisting of: alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, selenocysteine, serine, tyrosine, threonine, tryptophan and valine.

[0034] The amino acid molecule may be an alpha-amino acid, beta-amino acid or gamma-amino acid.

[0035] In further examples described herein, R'' is tert-Butyl amide.

[0036] In further examples described herein, there is provided use of a compound for cyclizing an amino acid molecule, wherein the compound has the formula (Ia)/(Ib):



(Ia)

(Ib)

comprises an α -stereocenter at the carbon atom proximal to the aldehyde group with matching stereochemistry to the carbon atom proximal to the amino terminus of the amino acid molecule; and

$n = 0$ or 1 , and R_1 , R_2 , R_3 , R_4 and R_5 are independently selected from H; lower alkyl; aryl; heteroaryl; alkenyl; heterocycle; esters of the formula $-\text{C}(\text{O})\text{OR}^*$ wherein R^* is selected from alkyl and aryl; amides of the formula $-\text{C}(\text{O})\text{NR}$, wherein R and R are independently selected from alkyl and aryl; $-\text{CH}_2\text{C}(\text{O})\text{R}$, wherein R is selected from $-\text{OH}$, lower alkyl, aryl, -loweralkyl-aryl, or $-\text{NR}_3\text{R}_b$, where R_a and R_b are independently selected from H, lower alkyl, aryl or -

loweralkyl-aryl; $-C(O)R_c$, wherein R_c is selected from lower alkyl, aryl or -loweralkyl-aryl; or - lower alkyl-OR_d, wherein R_a is a suitable protecting group; all of which are optionally substituted at one or more substitutable positions with one or more suitable substituents; and the amino acid molecule is an amino acid, a linear peptide or a salt of the foregoing, provided that if the amino acid molecule is a linear peptide, the compound comprises an aziridine chiral center proximal to the aldehyde with matching stereochemistry to the carbon atom proximal to the amino terminus of the peptide.

[0037] As used herein, the term "amino acid molecule" is meant to include single amino acids and also peptides.

[0038] As used herein, the term "amino acid" refers to molecules containing an amine group, a carboxylic acid group and a side chain that varies. Amino acid is meant to include not only the twenty amino acids commonly found in proteins but also non-standard amino acids and unnatural amino acid derivatives known to those of skill in the art, and therefore includes, but is not limited to, alpha, beta and gamma amino acids. Peptides are polymers of at least two amino acids and may include standard, non-standard, and unnatural amino acids.

[0039] The term "suitable substituent" as used in the context of the present invention is meant to include independently H; hydroxyl; cyano; alkyl, such as lower alkyl, such as methyl, ethyl, propyl, n-butyl, t-butyl, hexyl and the like; alkoxy, such as lower alkoxy such as methoxy, ethoxy, and the like; aryloxy, such as phenoxy and the like; vinyl; alkenyl, such as hexenyl and the like; alkynyl; formyl; haloalkyl, such as lower haloalkyl which includes CF_3 , CCl_3 and the like; halide; aryl, such as phenyl and naphthyl; heteroaryl, such as thienyl and furanyl and the like; amide such as $C(O)NR_aR_b$, where R_a and R_b are independently selected from lower alkyl, aryl or benzyl, and the like; acyl, such as $C(O)-C_6H_5$, and the like; ester such as $-C(O)OCH_3$ the like; ethers and thioethers, such as O-Bn and the like; thioalkoxy; phosphino; and $-NR_aR_b$, where R_a and R_b are independently selected from lower alkyl, aryl or benzyl, and the like. It is to be understood that a suitable substituent as used in the context of the present invention is meant to denote a substituent that does not interfere with the formation of the desired product by the processes of the present invention.

[0040] As used in the context of the present invention, the term "lower alkyl" as used herein either alone or in combination with another substituent means acyclic, straight or branched chain alkyl substituent containing from one to six carbons and includes for example, methyl, ethyl, 1-methylethyl, 1-methylpropyl, 2-methylpropyl, and the like. A similar use of the term is to be understood for "lower alkoxy", "lower thioalkyl", "lower alkenyl" and the like in respect of the number of carbon atoms. For example, "lower alkoxy" as used herein includes methoxy, ethoxy, t-butoxy.

[0041] The term "alkyl" encompasses lower alkyl, and also includes alkyl groups having more than six carbon atoms, such as, for example, acyclic, straight or branched chain alkyl substituents having seven to ten carbon atoms.

[0042] The term "aryl" as used herein, either alone or in combination with another substituent, means an aromatic monocyclic system or an aromatic polycyclic system. For example, the term "aryl" includes a phenyl or a naphthyl ring, and may also include larger aromatic polycyclic systems, such as fluorescent (eg. anthracene) or radioactive labels and their derivatives.

[0043] The term "heteroaryl" as used herein, either alone or in combination with another substituent means a 5, 6, or 7-membered unsaturated heterocycle containing from one to 4 heteroatoms selected from nitrogen, oxygen, and sulphur and which form an aromatic system. The term "heteroaryl" also includes a polycyclic aromatic system comprising a 5, 6, or 7-membered unsaturated heterocycle containing from one to 4 heteroatoms selected from nitrogen, oxygen, and sulphur.

[0044] The term "cycloalkyl" as used herein, either alone or in combination with another substituent, means a cycloalkyl substituent that includes for example, but is not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

[0045] The term "cycloalkyl-alkyl-" as used herein means an alkyl radical to which a cycloalkyl radical is directly linked; and includes, but is not limited to, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 1-cyclopentylethyl, 2-cyclopentylethyl, cyclohexylmethyl, 1-cyclohexylethyl and 2-cyclohexylethyl. A similar use of the "alkyl" or "lower alkyl" terms is to be understood for aryl-alkyl-, aryl-loweralkyl- (eg. benzyl), -lower alkyl-alkenyl (eg. allyl), heteroaryl-alkyl-, and the like as used herein. For example, the term "aryl-alkyl-" means an alkyl radical, to which an aryl is bonded. Examples of aryl-alkyl- include, but are not limited to, benzyl (phenylmethyl), 1-phenylethyl, 2-phenylethyl and phenylpropyl.

[0046] As used herein, the term "heterocycle", either alone or in combination with another radical, means a monovalent radical derived by removal of a hydrogen from a three- to seven-membered saturated or unsaturated (including aromatic) heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur. Examples of such heterocycles include, but are not limited to, aziridine, epoxide, azetidine, pyrrolidine, tetrahydrofuran, thiazolidine, pyrrole, thiophene, hydantoin, diazepine, imidazole, isoxazole, thiazole, tetrazole, piperidine, piperazine, homopiperidine, homo-piperazine, 1,4-dioxane, 4-morpholine, 4-thiomorpholine, pyridine, pyridine-N-oxide or pyrimidine, and the like.

The term "alkenyl", as used herein, either alone or in combination with another radical, is intended to mean an unsaturated, acyclic straight chain radical containing two or more carbon atoms, at least two of which are bonded to each other by a double bond. Examples of such radicals include, but are not limited to, ethenyl (vinyl), 1-propenyl, 2-propenyl, and 1-butenyl.

[0047] The term "alkynyl", as used herein is intended to mean an unsaturated, acyclic straight chain radical containing two or more carbon atoms, at least two of which are bonded to each other by a triple bond. Examples of such radicals include, but are not limited to, ethynyl, 1-propynyl, 2-propynyl, and 1-butyne. The term "alkoxy" as used herein, either alone or in combination with another radical, means the radical -O-(C_{1-n})alkyl wherein alkyl is as defined

above containing 1 or more carbon atoms, and includes for example methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy and 1,1-dimethylethoxy. Where n is 1 to 6, the term "lower alkoxy" applies, as noted above, whereas the term "alkoxy" encompasses "lower alkoxy" as well as alkoxy groups where n is greater than 6 (for example, n = 7 to 10). The term "aryloxy" as used herein alone or in combination with another radical means -O-aryl, wherein aryl is defined as noted above.

[0048] A peptide is a polymer of two or more amino acids.

[0049] The following examples are illustrative of various examples of the invention.

EXAMPLES

Example 1

Materials and Methods

[0050] Anhydrous toluene and dimethylformamide (DMF) were purchased and used as received. Tetrahydrofuran (THF) was distilled from sodium benzophenone ketyl under argon. All other solvents including TFE (2,2,2,-trifluoroethanol) and HFIP (1,1,1,3,3,3-hexafluoro-2-isopropanol) were of reagent grade quality. Melting points were obtained on a MelTemp melting-point apparatus and are uncorrected.

Method 1: amino acid/peptide cyclization reaction was conducted using the following method. In a screw-cap vial equipped with a magnetic stirring bar was added peptide (0.2 mmol) and 1 ml of TFE and stirred until a homogeneous solution has been obtained. Aziridine aldehyde dimer (0.1 mmol) and isocyanide (0.2 mmol) were then added sequentially and the resulting mixture was stirred for the time specified in Table 1. Reactions were monitored by electrospray ionization mass spectrometry ("ESI-MS") at 60 eV and/or thin layer chromatography ("TLC") analysis. After completion of the reaction, 1 ml of water and 1 ml of Et₂O were added and the mixture was shaken vigorously and then cooled on ice. The resulting precipitate was filtered and washed with hexanes and cold Et₂O (1 ml) to afford the cyclic peptide. For products that are water soluble or do not precipitate, the reaction mixture was concentrated under reduced pressure and then triturated with Et₂O and hexanes (0.2 ml) to afford the cyclic peptide product.

Method 2: amino acid/peptide cyclization reaction was conducted using HFIP in place of TFE (Method 2). In a screw-cap vial equipped with a magnetic stirring bar was added peptide (0.2 mmol) and 1 ml of HFIP and stirred until homogeneous solution has been obtained. Aziridine aldehyde dimer (0.1 mmol) and isocyanide (0.2 mmol) were then added sequentially and the

resulting mixture was stirred for the time specified in Table 1. Reactions were monitored by ESI-MS at 60 eV and/or TLC analysis. After completion of the reaction, the mixture was concentrated under reduced pressure and then triturated with Et₂O and hexanes (0.2 ml) to afford the cyclic peptide product.

Method 3: amino acid/peptide cyclization reaction was carried out using HFIP and water. In a screw-cap vial equipped with a magnetic stirring bar was added peptide (0.2 mmol) and 1 ml of HFIP and 47 microliters of H₂O and stirred until homogeneous. Aziridine aldehyde dimer (0.1 mmol) and isocyanide (0.2 mmol) were then added sequentially and the resulting mixture was stirred for the time specified in Table 1. Reactions were monitored by ESI-MS at 60 eV and/or TLC analysis. After completion of the reaction, the mixture was concentrated under reduced pressure and then triturated with Et₂O and hexanes to afford the cyclic peptide product.

[0051] Note: In reactions involving either cysteine or peptides containing thiol residues, the solvent was degassed with argon for two hours prior to reaction. The reaction was thereafter carried out under an atmosphere of argon.

[0052] Some of the cyclic products that may be synthesized from each of Methods 1 - 3 are summarized in Table 3 and depicted in Figure 4

Table 3. Methods used for amino acid/peptide cyclization reactions.

Method	Cyclic Products
1	1, 2, 3, 4, 7, 9, 11, 12, 15, 16, 17, 18, 19, 20, 21, 22
2	5, 8, 10, 13, 14
3	6

[0053] The relative stereochemistry of each cyclic product listed in Table 3 was established by correlating the methine region of each ¹H NMR spectrum with that of piperazinone 1.

[0054] Chromatography: Flash column chromatography was carried out using Silicycle 230-400 mesh silica gel. Thin-layer chromatography (TLC) was performed on Macherey Nagel pre-coated glass backed TLC plates (SIL G/UV254, 0.25 mm) and visualized using a UV lamp (254 nm) and iodine stain.

[0055] Nuclear magnetic resonance spectra: ¹H and ¹³C NMR spectra were recorded on Varian Mercury 400 or 500 MHz spectrometers. ¹H NMR spectra were referenced to TMS (0 ppm) and ¹³C NMR spectra were referenced to CDC13 (77.23 ppm). Peak multiplicities are designated by the following abbreviations: s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quartet; m, multiplet; ds, doublet of singlets; dd, doublet of doublets; ddd, doublet of doublet of

doublets; bt, broad triplet; td, triplet of doublets; tdd, triplet of doublets of doublets.

[0056] Mass Spectroscopy: High-resolution mass spectra were obtained on a VG 70-250S (double focusing) mass spectrometer at 70 eV or on an ABI/Sciex Qstar mass spectrometer with ESI source, MS/MS and accurate mass capabilities. Low resolution mass spectra (ESI) were obtained at 60 eV, 70 eV and 100 eV.

[0057] Synthetic routes toward aziridine aldehyde dimers: All dimeric amino aldehydes used in this study were prepared according to the synthetic route outlined in Supplementary Figures 1 and 2.

Results

[0058] Diverse piperazinones were produced in a one-step process (Table 1).

Table 1. The scope of piperazinone synthesis (L-amino acid-derived product is shown).

Entry ^a	R ¹	Amino Acid	Time	Product ^b
1	CH ₂ OTBDMS	L-Phe	1	1 (92)
2	CH ₂ OTBDMS	D-Phe	1.5	2 (90)
3	CH ₂ OTBDMS	L-Ala	1.5	3 (82)
4	CH ₂ OTBDMS	L-Pro	1	4 (98)
5	CH ₂ ⁱ Pr	D-Arg	1	5 (83)
6 ^c	CH ₂ ⁱ Pr	L-Lys HCl ^g	1	6 (76)
7	CH ₂ OTBDMS	L-Gly	1	7 (80)
8 ^e	CH ₂ ¹ Pr	L-Asp	3	8 (76)
9 ^f	CH ₂ OTBDMS	L-Cys	2	9 (82)
10 ^d	CH ₂ OTBDMS	L-His	1	10 (88)
11	CH ₂ OTBDMS	L-Ser	3	11 (77)

^a Unless specified otherwise, reactions were performed at room temperature using 0.2 mmol of isocyanide and amino acid, and 0.1 mmol of amino aldehyde dimer in TFE (0.2 M).

^b % isolated yield.

^c HFIP/H₂O (20:1) was used as solvent (0.2 M).


^d HFIP was used as solvent (0.2 M).

^e Cyclization occurs at the more acidic α -carboxylic acid.

^f TFE was degassed prior to use.

^g Lysine side chain amine is protonated.


[0059] The step that shifts the ring/chain equilibrium is transannular collapse of the nucleophilic NH aziridine onto the electrophilic mixed anhydride generated when an amphoteric amino aldehyde is subjected to the Ugi reaction condition (Scheme 2). The reaction takes place under stoichiometric conditions and delivers cyclic peptide product. High dilution, a critical condition normally required in order to achieve high yields in cyclic peptide synthesis, is not necessary.




Scheme 2. A. Macrocyclization with a monofunctional aldehyde; B. macrocyclization mediated by an amphoteric amino aldehyde.

[0060] The normally challenging medium-sized rings can be made with great facility and in good yields within several hours (Table 2).

Table 2. Representative scope of linear peptide macrocyclization.





^a Unless specified otherwise, reactions were performed at room temperature using 0.2 mmol of isocyanide and amino acid, and 0.1 mmol of amino aldehyde dimer in TFE (0.2 M). ^b Isolated yield. ^c Diastereoselectivity of >20:1 was confirmed by ¹H NMR analysis of each crude reaction mixture. The relative stereochemistry was established by correlating the methine region of each ¹H NMR spectrum with that of piperazinone **1** (see X-ray crystal structure of **1**). ^d HFIP was used as solvent (0.2 M).

[0061] ¹H and ¹³C NMR spectra for the cyclic products of Table 3 are shown in Figures 5 - 27.

[0062] A one-step process for making versatile cyclic peptides with high stereo- and chemoselectivity for linear peptides, isocyanides, and amphoteric amino aldehydes is provided. The Ugi reaction is run with an amphoteric amino aldehyde in place of a monofunctional aldehyde. When L-phenyl alanine was reacted with tert-butyl isocyanide, and an aziridine aldehyde, a cyclic piperazinone was obtained in 92% yield in approximately one hour (Table 1, Entry 1). The relative stereochemistry of **1** was established using X-ray analysis. A range of amino acids were subjected to this reaction and in all cases the corresponding piperazinone products were obtained as single diastereoisomers without formation of linear peptides (Figure 1).

[0063] The existence of epimeric products were investigated through the use of ¹H NMR (DMSO-d₆ or CDCl₃) analysis of crude reaction mixtures. In all of the reactions studied, there were no detectable signals corresponding to the epimer. As such, the resulting crude cyclic peptides were obtained in >20:1 selectivity.

[0064] The effect of medium-sized peptide chain length on the reaction outcome was studied (Table 2). The challenging medium-sized rings are readily prepared; the reaction times are less than 10 hours, proceeding with high yields and diastereoselectivities. The preferred work-up procedure involves product precipitation from diethyl ether and hexanes. The cyclic peptides require no further purification by HPLC. In addition, racemization was not detected throughout the course of the reaction or during product isolation. The lack of epimerization is further evidenced by high stereoselectivity; aziridine aldehydes with S stereocenters next to the carbonyl group undergo macrocyclization with the peptides that contain an L-amino acid residue at the N-terminus. The "mismatched" reaction with the D-amino acid-terminated peptide is unproductive, leading to the formation of stable amins.

[0065] It has been determined that the α-stereocenter of the N-terminal amino acid, in this case proline, should match the α-stereocenter of the aziridine aldehyde in order for cyclization to occur, with respect to peptides. In a mismatched reaction, the only product formed is the corresponding aminal (Figure 2). Even after an extended reaction time and additional equivalents of isocyanide, cyclic peptide formation was not observed. This restriction is not present for cyclization of single amino acids.

[0066] When the opposite enantiomer of amino aldehyde was used in the same reaction, the proline and the amino aldehyde both had S-configuration at their α -stereocenters. This matched case readily underwent peptide cyclization to yield the corresponding cyclic product in high yield (Figure 3).

[0067] This stereoselectivity process is significant in the context of possible epimerization at the N-terminus. Since matched stereochemistry is required for the cyclization to occur, the resulting cyclic peptides will necessarily contain matching stereocenters. Furthermore, the fact that no cyclic peptide formation was observed in mismatched reactions suggests that no epimerization, which would result in matched substrates, occurs under our macro conditions.

[0068] Consistent yields in this chemistry are likely obtained due to the mechanism that governs cyclization. It is at this point that a departure from the Ugi reaction with monofunctional aldehydes becomes apparent. When monofunctional aldehydes are used in the reaction with isocyanides and peptides, low diastereoselectivities are observed. More importantly, the undesired cyclodimerization occurs during the cyclization of linear peptides containing less than six residues; the cyclization of tripeptides yields only the cyclodimers.¹³ This low selectivity is due to a slow *transannular* attack of the amine onto the mixed anhydride (Scheme 2, **A**), thus allowing the intermolecular process to be kinetically competitive. By adding an amphoteric amino aldehyde to the cyclization reaction mixture containing a secondary amine-terminated peptide, the slow transannular attack is being replaced with a fast attack by the nucleophilic aziridine, which is positioned exocyclic to the mixed anhydride (Scheme 2, **B**). This provides an unencumbered trajectory of attack. Since TFE is a non-nucleophilic solvent, premature solvolysis of the mixed anhydride is not observed. The reaction mechanism ensures that the C-terminus is activated only upon formation of the intermediate cyclic mixed anhydride, which is then attacked by the exocyclic aziridine. This not only secures selectivity for the intramolecular macrocyclization, but also avoids prolonged C-terminus activation and potential epimerization. Accordingly, high dilution, a critical condition normally required in order to achieve high yields in conventional cyclic peptide synthesis, is not necessary.⁶ Furthermore, no oligomeric or polymeric by-products were detected in the subject experiments. A side-by side comparison of the subject reaction with the traditional lactamization of a linear tetrapeptide, widely used in the synthesis of cyclic peptides, was conducted. The amino aldehyde mediated macrocyclization delivered rapid, selective, and efficient formation of the cyclic peptide. In contrast, only trace amount of the cyclic peptide was detected among myriad structures formed in the course of the lactamization. The undesired cyclodimerization⁵ has dominated this process, which is a testament to sluggish kinetics of the conventional intramolecular process. Peptide macrocyclization reactions are normally performed under extremely dilution. Typical protocols necessitate molar concentrations of 10^{-4} M. This large dilution increases the selectivity for intramolecular cyclizations, thereby limiting undesired cyclodimerization and trimerization.

[0069] Characterization of the relative stereochemistry of each cyclic product listed in Table 3 is set out below.

(3*S*,5*R*,6*R*,7*S*)-3-benzyl-*N*-tert-butyl-7-((tert-butyldimethylsilyloxy)methyl)-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (1)

[0070] ^1H NMR (CDCl_3 , 400MHz) δ : 7.36 - 7.18 (m, 5H), 6.75 (bs, 1H), 3.86 (dd, J = 11.6 Hz, 4.8 Hz, 1H), 3.74 (dd, J = 11.2 Hz, 4.8 Hz, 1H), 3.54 (bs, 1H), 3.47 (dd, J = 3.6 Hz, 1.2 Hz, 1H), 3.25 (dd, J = 14.4 Hz, 3.2 Hz, 1H), 3.10 (m, 1H), 2.52 (dd, J = 14.4 Hz, 10.4 Hz, 1H), 2.40 (m, 1H), 1.80 (bs, 1H), 1.02 (s, 9H), 0.90 (s, 9H), 0.10 (s, 3H), 0.08 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 100 MHz) δ : 184.8, 169.0, 138.5, 129.7, 128.9, 127.0, 63.9, 57.5, 53.6, 50.7, 44.6, 39.0, 35.5, 28.5, 26.1, 18.6, -5.0 ppm. HRMS (ESI) $[\text{MH}]^+$ calcd. 446.2833, found 446.2831

(3*R*,5*R*,6*R*,7*S*)-3-benzyl-*N*-tert-butyl-7-((tert-butyldimethylsilyloxy)methyl)-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (2)

[0071] ^1H NMR (CDCl_3 , 400MHz) δ : 7.35 - 7.20 (m, 5H), 6.32 (bs, 1H), 3.86 (m, 1H), 3.73 (dd, J = 11.5 Hz, 4.2 Hz, 1H), 3.86 (bs, 1H), 3.73 (dd, J = 11.5 Hz, 4.2 Hz, 1H), 3.25 (dd, J = 14.4 Hz, 3.2 Hz, 1H), 3.10 (m, 1H), 2.52 (dd, J = 14.4 Hz, 10.4 Hz, 1H), 2.40 (m, 1H), 1.80 (bs, 1H), 1.31 (s, 9H), 0.90 (s, 9H), 0.10 (s, 3H), 0.08 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 75 MHz) δ : 184.0, 169.1, 136.2, 129.2, 129.1, 127.3, 63.0, 59.5, 51.4, 49.6, 42.2, 38.7, 36.0, 28.8, 26.1, 18.7, -5.1 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 446.3, found 446.3

(3*S*,5*R*,6*R*,7*S*)-*N*-tert-butyl-7-((tert-butyldimethylsilyloxy)methyl)-3-methyl-2-oxo-1,4-diazabicyclo [4.1.0] heptane-5-carboxamide (3)

[0072] ^1H NMR (CDCl_3 , 400MHz) δ : 7.42 (s, NH), 3.88 (dd, J = 11.2 Hz, 4.8 Hz, 1H), 3.70 (dd, J = 11.2 Hz, 5.2 Hz, 1H), 3.55 (d, J = 0.8 Hz, 1H), 3.46 (dd, J = 3.6 Hz, 1.6 Hz, 1H), 3.07 (m, 1H), 2.36 (td, J = 4.8 Hz, 3.6 Hz, 1H), 1.8 (bs, NH), 1.39 (s, 9H), 1.23 (d, J = 6.8 Hz, 3H), 0.90 (s, 9H), 0.09 (s, 3H), 0.08 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 100 MHz) δ : 185.5, 169.3, 63.9, 53.8, 51.2, 51.0, 44.5, 39.3, 28.9, 26.1, 16.6, 15.0, -5.0 ppm. HRMS (ESI) $[\text{MH}]^+$ calcd. 370.2520, found 370.2520

(1*S*,3*aS*,8*R*,8*aS*)-*N*-tert-butyl-1-((tert-butyldimethylsilyloxy)methyl)-3-oxooctahydroazirino [1,2-*a*] pyrrolo[1,2-*d*]pyrazine-8-carboxamide (4)

[0073] ^1H NMR (CDCl_3 , 400MHz) δ : 6.17 (s, 1H), 3.76 (dd, J = 11.6 Hz, 4.4 Hz, 1H), 3.67 (dd, J = 11.6 Hz, 4.8 Hz, 1H), 3.70 (d, J = 6.2 Hz, 1H), 3.12 (dt, J = 4.8 Hz, 9.2 Hz, 1H), 2.97 (dd, J

= 6.4 Hz, 3.6 Hz, 1H), 2.93 - 2.89 (m, 1H), 2.60 (q, J = 4.4 Hz, 1H), 2.25 - 2.05 (m, 2H), 1.90 - 1.75 (m, 3H), 1.70 - 1.51 (m, 1H), 1.48 (s, 9H), 0.85 (s, 9H), 0.09 (s, 3H), 0.08 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 100 MHz) δ : 183.0, 168.6, 65.0, 63.4, 63.2, 54.6, 51.3, 43.4, 41.0, 29.0, 26.1, 22.2, 22.0, 18.9, -5.1 ppm. HRMS (ESI) $[\text{MH}]^+$ calcd. 396.2676, found 396.2656

(3*R*,5*S*,6*S*,7*S*)-*N*-tert-butyl-3-(3-guanidinopropyl)-7-isobutyl-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (5)

[0074] ^1H NMR (D_2O , 400MHz) δ : 3.59 (t, J = 5.6 Hz, 1H), 3.46 (m, 1H), 3.13 (m, 1H), 2.94 (m, 1H), 3.37 (m, 1H), 1.78 - 1.44 (m, 8H), 1.22 (s, 9H), 0.83 (d, J = 2.8 Hz, 3H), 0.81 (d, J = 2.8 Hz, 3H) ppm. ^{13}H NMR (D_2O , 400MHz) δ : 186.7, 172.0, 156.9, 54.9, 54.5, 54.0, 44.5, 42.8, 28.7, 27.9, 26.5, 25.7, 25.0, 24.1, 22.2, 21.5 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 367.3, found 367.2 and 184.1 for $[\text{M}+2\text{H}]^{2+}/2$.

(3*S*,5*S*,6*S*,7*S*)-3-(4-aminobutyl)-*N*-tert-butyl-7-isobutyl-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (6)

[0075] NMR (D_2O with 10% v/v CD_3COOD , 400MHz) δ : 5.14 (ddd, J = 9.6 Hz, 4.4 Hz, 2.4 Hz, 1H), 4.12 (d, J = 2.4 Hz, 1H), 3.07 (dd, J = 7.6 Hz, 4.4 Hz, 1H), 2.85 (t, J = 7.6 Hz, 2H), 1.82 - 1.46 (m, 10H), 1.32 (s, 9H), 0.83 (d, J = 6.8 Hz, 3H), 0.81 (d, J = 6.8 Hz, 3H) ppm. ^{13}C NMR (D_2O with 10% v/v CD_3COOD , 100 MHz) δ : 178.0, 173.8, 97.2, 58.6, 54.5, 53.0, 40.6, 39.3, 29.3, 27.8, 27.6, 27.2, 26.8, 24.6, 22.2, 21.9, 20.8 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 339.3, found 339.2 and 170.1 for $[\text{M}+2\text{H}]^{2+}/2$

(5*R*,6*R*,7*S*)-*N*-tert-butyl-7-((tert-butyl)dimethylsilyloxy)methyl-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (7)

[0076] ^1H NMR (400 MHz, CDCl_3) δ : 6.98 (s, 1H), 3.87 (dd, J = 11.5, 4.5 Hz, 1H), 3.78 (dd, J = 11.5, 4.5 Hz, 1H), 3.34 - 3.24 (m, 2H), 3.13 (d, J = 4.6 Hz, 1H), 2.40 (td, J = 4.5, 3.4 Hz, 1H), 2.17 - 1.96 (m, 1H), 1.43 - 1.36 (m, 9H), 0.94 - 0.86 (m, 9H), 0.13 - 0.05 (m, 6H). ^{13}C NMR (101 MHz, CDCl_3) δ : 183.93, 168.37, 77.48, 77.16, 76.84, 63.02, 55.81, 51.25, 48.72, 45.27, 39.64, 28.87, 26.06, 18.55, 0.15, -5.09, -5.12.

2-((3*S*,5*R*,6*R*,7*R*)-5-(tert-butylcarbamoyl)-7-isobutyl-2-oxo-1,4-diazabicyclo[4.1.0]heptan-3-yl)acetic acid (8)

[0077] ^1H NMR (400 MHz, CD_3OD) δ : 4.69 - 4.58 (m, 1H), 3.92 (d, J = 2.0 Hz, 1H), 3.76 (dd, J = 9.8, 3.6 Hz, 1H), 3.66 (dd, J = 7.8, 4.0 Hz, 1H), 2.90 (dd, J = 17.5, 3.6 Hz, 1H), 2.81 (dd, J = 16.7, 4.0 Hz, 1H), 2.69 - 2.53 (m, 2H), 1.36 - 1.32 (m, 9H), 0.98 (dt, J = 8.5, 4.2 Hz, 6H). ^{13}C NMR (101 MHz, CD_3OD) δ 188.21, 172.40, 158.66, 158.62, 56.05, 55.25, 55.23, 51.95, 44.17, 43.55, 43.35, 42.55, 42.39, 42.16, 41.98, 29.85, 28.90, 28.85, 28.10, 27.37, 27.16, 26.85, 26.51, 23.19, 22.63, 21.05.

(3R,5R,6R,7S)-N-(tert-butyl)-7-(((tert-butyldimethylsilyl)oxy)methyl)-3-(mercaptopomethyl)-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (9)

[0078] *Note: proper spectroscopic characterization of this compound is difficult as spectra are characterized by extremely broad peaks. This is attributed to inherent instability of 9 due to the presence of nucleophilic thiol and its susceptibility to oxidation and aziridine ring-opening.*

[0079] ^1H NMR (CDCl_3 , 400MHz) δ : 3.47 (s, 1H), 3.41 (d, J = 6.5 Hz, 1H), 4.35-2.42 (m, 7H), 1.70-1.42 (bs, 10H), 0.90 (s, 9H), 0.10 (s, 3H), 0.08 (s, 3H) ppm. MS (ESI) $[\text{MH}]^+$ calcd. 402.6, found 402.2

(3S,5R,6R,7S)-3-((1H-imidazol-4-yl)methyl)-N-(tert-butyl)-7-(((tert-butyldimethylsilyl)oxy)methyl)-2-oxo-1,4-diazabicyclo [4.1.0] heptane-5-carboxamide (10)

[0080] ^1H NMR (CDCl_3 , 400MHz) δ : 7.56 (s, 1H), 7.25 (s, 1H), 6.87 (s, 1H), 6.22 (s, 1H), 4.46 (m, 1H), 3.72 (dd, J = 11.2 Hz, 4.8 Hz, 1H), 3.60 (m, 1H), 3.44 (dd, J = 3.6 Hz, 1.5 Hz, 1H), 3.27 (dd, J = 14.1 Hz, 3.1 Hz, 1H), 2.99 (dd, J = 14.7 Hz, 4.2 Hz, 1H), 2.80 (dd, J = 15.2 Hz, 8.1 Hz, 1H), 2.40 (m, 1H), 2.05 (bs, 1H), 1.36 (s, 9H), 0.87 (s, 9H), 0.07 (s, 3H), 0.05 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 100 MHz) δ : 185.4, 171.4, 169.5, 168.6, 135.1, 64.3, 62.7, 60.6, 54.7, 51.2, 44.5, 39.3, 28.9, 26.0, 18.4, -5.3 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 436.6, found 436.2

(3S,5R,6R,7S)-N-(tert-butyl)-7-(((tert-butyldimethylsilyl)oxy)methyl)-3-(hydroxymethyl)-2-oxo-1,4-diazabicyclo[4.1.0]heptane-5-carboxamide (11)

[0081] ^1H NMR (CDCl_3 , 400MHz) δ : 6.80 (bs, 1H), 6.15 (bs, 1H), 4.02 (d, J = 11.6 Hz, 1H), 3.85 (dd, J = 13.3 Hz, 6.7 Hz, 1H), 3.76 (dd, J = 14.2 Hz, 3.0 Hz, 1H), 3.68 (bs, 1H), 3.53 (d, J = 5.0 Hz, 1H), 3.45 (dd, J = 3.5 Hz, 2.4 Hz, 1H), 3.13 (m, 1H), 2.55 (m, 1H), 2.42 (m, 1H), 1.39

(s, 9H), 0.90 (s, 9H), 0.10 (s, 3H), 0.08 (s, 3H) ppm. ^{13}C NMR (CDCl_3 , 100 MHz) δ : 185.3, 168.9, 63.5, 61.4, 55.8, 53.9, 51.2, 44.9, 39.6, 28.9, 26.1, 18.6, -5.1 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 386.6, found 386.2

(1S,4S,6aS,11R,11aS)-N-tert-butyl-1-((tert-butyldimethylsilyloxy)methyl)-4-isobutyl-3,6-dioxodecahydro-1H-azirino[1,2-a]pyrrolo[1,2-d][1,4,7]triazonine-11-carboxamide (12)

[0082] ^1H NMR (CD_3OD , 400MHz) δ : 4.47 (m, 1H), 4.14 (dd, J = 8.4 Hz, 6 Hz, 1H), 4.07 - 3.99 (m, 2H), 3.84 (dd, J = 12 Hz, 3.2 Hz, 1H), 3.32 (d, J = 5.6 Hz, 1H), 3.26 - 3.15 (m, 2H), 2.42 - 2.30 (m, 2H), 1.92 - 1.78 (m, 3H), 1.74 (q, J = 6.8 Hz, 2 H), 1.59 (sept, J = 6.8 Hz, 1H), 1.22 (s, 9H), 0.90 (d, J = 6.8 Hz, 3H), 0.87 (d, J = 6.4 Hz, 3H), 0.84 (s, 9H), 0.06 (s, 3H), 0.03 (s, 3H) ppm. ^{13}C NMR (CD_3OD , 100 MHz) δ : 174.1, 164.1, 157.4, 86.5, 62.4, 60.7, 60.6, 57.7, 54.2, 52.1, 50.2, 39.6, 29.0, 25.8, 25.2, 25.1, 22.1, 21.0, 20.5, 17.9, -6.5 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 509.3, found 509.3

(1S,4S,6aR,11S,11aR)-N-tert-butyl-4-(3-guanidinopropyl)-1-isobutyl-3,6-dioxodecahydro-1H-azirino[1,2-a]pyrrolo[1,2-d][1,4,7]triazonine-11-carboxamide (13)

[0083] ^1H NMR (CD_3OD , 400MHz) δ : 4.26 (t, J = 5.6 Hz, 1H), 4.18 (dd, J = 8.8 Hz, 2.4 Hz, 1H), 3.58 (dd, J = 8.8 Hz, 5.2 Hz, 1H), 3.28 - 3.24 (m, 2H), 3.00 (d, J = 5.6 Hz, 1H), 2.70 (dd, J = 10.4 Hz, 6.4 Hz, 1H), 2.32 - 2.18 (m, 2H), 1.98 - 1.52 (m, 11H), 1.28 (s, 9H), 1.01 (d, J = 6.4 Hz, 3H), 0.96 (d, J = 6.4 Hz, 3H) ppm. ^{13}C NMR (CD_3OD , 100 MHz) δ : 178.0, 160.8, 159.1, 157.4, 86.9, 64.3, 63.5, 60.2, 54.5, 53.7, 51.0, 42.6, 41.4, 29.4, 28.8, 26.1, 24.9, 24.5, 22.7, 21.3, 20.4 ppm. MS(ESI) $[\text{MH}]^+$ calcd. 464.3, found 464.3 and 232.6 for $[\text{M}+2\text{H}]^{2+}/2$

2-((1S,4S,6aR,11S,11aR)-11-(tert-butylcarbamoyl)-1-isobutyl-3,6-dioxodecahydro-1H-azirino[1,2-a]pyrrolo[1,2-d][1,4,7]triazonin-4-yl)acetic acid (14)

[0084] ^1H NMR (CD_3OD , 400MHz) δ : 4.46 (t, J = 9.2 Hz, 1H), 4.34 (dd, J = 8.4 Hz, 2.0 Hz, 1H), 3.81 (dd, J = 8.4 Hz, 5.6 Hz, 1H), 3.38 - 3.30 (m, 1H), 3.02 (dd, J = 17.2 Hz, 2.4 Hz, 1H), 2.85 - 2.73 (m, 1H), 2.57 (dd, J = 17.2 Hz, 8.4 Hz, 1H), 2.45 - 2.32 (m, 2H), 2.05 - 1.62 (m, 7H), 1.58 - 1.48 (m, 2H), 1.27 (s, 9H), 1.01 (d, J = 6.4 Hz, 3H), 0.98 (d, J = 6.4 Hz, 3H) ppm. MS (ESI) $[\text{MH}]^+$ calcd. 423.3, found 423.2

(1S,4S,9aS,14R,14aS)-N-tert-butyl-1-((tert-butyldimethylsilyloxy)methyl)-4-isobutyl-3,6,9-trioxotetradecahydroazirino [1,2-a] pyrrolo [1,2-d][1,4,7,10]tetraazacyclododecine-14-carboxamide (15)

[0085] ^1H NMR (CD_3OD , 400MHz) δ : 4.55 (ddd, $J = 8$ Hz, 3.2 Hz, 1.6 Hz, 1H), 4.32 (dd, $J = 10.4$ Hz, 4 Hz, 1H), 4.21 (dd, $J = 14$ Hz, 6Hz, 1H), 4.15 - 4.05 (m, 3H), 3.91 (dd, $J = 12$ Hz, 3.2 Hz, 1H), 3.41 Hz (dd, $J = 6$ Hz, 1 H), 3.36 - 3.28 (m, 1H), 2.50 - 2.30 (m, 2H), 2.00 - 1.50 (m, 7H), 1.31 (s, 9 H), 0.98 - 0.92 (m, 6H), 0.92 (s, 9 H), 0.13 (s, 3H), 0.11 (s, 3H) ppm. ^{13}C NMR (CD_3OD , 100 MHz) δ : 178.3, 166.4, 165.1, 157.6, 86.7, 62.7, 61.0, 60.1, 54.2, 53.7, 52.6, 50.5, 44.0, 41.6, 28.9, 25.6, 25.2, 25.1, 25.1, 22.6, 20.7, 20.5, 17.9, -6.5, -6.5 ppm. MS (ESI) $[\text{MH}]^+$ calcd. 566.4, found 566.4

(1R,1aS,2R,6aS,12S,18S)-18-benzyl-N-tert-butyl-1,12-diisobutyl-7,10,13,16,19-pentaoxoicosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16]hexaazacyclooctadecine-2-carboxamide (16)

[0086] ^1H NMR (CDCl_3 , 500MHz) δ : 7.82 (bs, NH), 7.40 - 7.20 (m, 5H), 7.18 (bs, NH), 7.05 (bs, NH), 6.81 (bs, 2 NH), 4.64 (q, $J = 6.8$ Hz, 1H), 4.13 (dd, $J = 16$ Hz, 7.6 Hz, 1H), 4.10 - 3.90 (m, 2H), 3.57 (dd, $J = 16$ Hz, 4.8 Hz, 1H), 3.50 - 3.39 (m, 2H), 3.26 (dd, $J = 14$ Hz, 5.6 Hz, 1H), 3.12 (dd, $J = 14$ Hz, 7.2 Hz, 1H), 3.12 - 3.08 (m, 1H), 3.00 - 2.92 (m, 2H), 2.63 (t, $J = 3.6$ Hz, 1H), 2.28 - 2.18 (m, 1H), 1.90 - 1.70 (m, 2H), 1.37 - 1.25 (m, 9H), 1.10 - 0.88 (m, 12H) ppm. ^{13}C NMR (CDCl_3 , 125 MHz) δ : 181.4, 176.1, 175.7, 172.1, 170.1, 169.2, 136.4, 130.1, 128.7, 126.8, 56.2, 63.0, 43.2, 42.5, 40.2, 38.6, 37.9, 31.2, 30.0, 29.8, 29.5, 28.8, 28.7, 27.2, 25.1, 24.9, 24.3, 23.3, 23.1, 22.8, 22.7, 22.6, 22.4, 22.3, 21.9, MS (ESI) $[\text{MH}]^+$ calcd. 682.4, found 682.4

(1S,12aS,17R,17aR)-N-tert-butyl-1-isobutyl-3,6,9,12-tetraoxohexadecahydro-1H-azirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13]pentaazacyclopentadecine-17-carboxamide (17)

[0087] ^1H NMR (400 MHz, CDCl_3) δ : 4.57 - 4.44 (m, 1H), 4.32 - 3.70 (m, 10H), 3.38 (d, $J = 5.4$ Hz, 1H), 3.28 (s, 1H), 3.14 (dd, $J = 11.2$, 5.6 Hz, 1H), 2.52 - 2.41 (m, 1H), 2.37 (d, $J = 6.9$ Hz, 1H), 2.15 - 1.65 (m, 8H), 1.64 - 1.48 (m, 2H), 1.42 - 1.23 (m, 8H), 1.00 (dt, $J = 21.4$, 7.4 Hz, 6H).

[0088] ^{13}C NMR (100 MHz, CD_3OD) δ : 176.18, 171.23, 168.97, 165.90, 159.24, 86.37, 65.77, 63.49, 61.48, 59.76, 55.41, 51.43, 47.38, 45.65, 45.23, 44.59, 44.43, 44.19, 43.41, 42.46, 29.85, 29.43, 28.97, 28.58, 28.46, 27.72, 27.13, 26.74, 25.68, 23.84, 23.68, 23.14, 23.06, 22.16, 21.62.

tert-butyl 2-((1S,4S,7S,13S,15aS,20R,20aS)-20-(tert-butylcarbamoyl)-1-(((tert-butylidimethylsilyl)oxy)methyl)-4-methyl-3,6,9,12,15-pentaoxo-13-(3-(3-((2,2,4,6,7-

pentamethyl-2,3-dihydrobenzofuran-5-yl)sulfonyl)guanidino)propyl)icosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16]hexaazacyclooctadecin-7-yl)acetate (18)

[0089] ^1H NMR (400 MHz, CDCl_3) δ : 7.84 (d, J = 9.7 Hz, 2H), 7.58 (dd, J = 18.1, 9.1 Hz, 2H), 7.07 (s, 1H), 6.41 (s, 2H), 5.68 (s, 1H), 4.85 (dq, J = 13.7, 6.8 Hz, 1H), 4.70 - 4.58 (m, 1H), 4.3 (q, J = 8.1 Hz, 1H), 4.17 - 4.07 (m, 1H), 3.97 (dd, J = 14.2, 7.5 Hz, 2H), 3.74 (s, 1H), 3.69 - 3.57 (m, 1H), 3.56 - 3.42 (m, 2H), 3.42 - 3.13 (m, 4H), 2.96 (s, 2H), 2.93 (d, J = 2.3 Hz, 1H), 2.89 - 2.76 (m, 4H), 2.66 - 2.53 (m, 7H), 2.42 - 2.15 (m, 2H), 2.10 (s, 3H), 2.07 - 1.98 (m, 1H), 1.87 - 1.55 (m, 8H), 1.46 (s, 6H), 1.44 (s, 9H), 1.34 (s, 9H), 0.87 (s, 9H), 0.09 (s, 3H), 0.05 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 187.07, 176.14, 174.46, 171.46, 171.04, 169.11, 166.93, 158.77, 156.39, 138.63, 133.66, 132.55, 124.65, 117.55, 105.24, 86.49, 82.64, 63.92, 63.84, 60.81, 57.07, 51.69, 51.31, 50.37, 49.56, 45.47, 43.51, 40.43, 37.55, 35.99, 30.28, 29.17, 28.83, 28.22, 26.11, 24.61, 19.49, 18.71, 18.19, 17.31, 12.69, -5.11, -5.44. MS (ESI) $[\text{MH}]^+$ calcd. 1104.6, found 1104.6

tert-butyl (4-((1S,4S,7S,10S,13S,15aS,20R,20aS)-13-((R)-1-(tert-butoxy)ethyl)-20-(tert-butylcarbamoyl)-1-(((tert-butyl)dimethylsilyl)oxy)methyl)-4-methyl-10-(2-(methylthio)ethyl)-3,6,9,12,15-pentaoxoicosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16] hexaazacyclooctadecin-7-yl)butyl)carbamate (19)

[0090] ^1H NMR (400 MHz, CDCl_3) δ : 7.83 (d, J = 8.1 Hz, 1H), 7.77 (s, 1H), 7.50 (d, J = 21.3 Hz, 1H), 7.35 (m, 2H), 7.08 (dd, J = 15.7, 7.5 Hz, 1H), 6.90 (s, 1H), 6.82 (s, 1H), 4.98 (d, J = 27.6 Hz, 1H), 4.84 - 4.67 (m, 1H), 4.60 (s, 1H), 4.40 - 4.30 (m, 1H), 4.16 - 3.93 (m, 4H), 3.82 (s, 2H), 3.77 - 3.69 (m, 3H), 3.67 (d, J = 6.3 Hz, 1H), 3.45 - 3.28 (m, 2H), 3.27 - 2.99 (m, 6H), 3.00 - 2.86 (m, 2H), 2.84 (s, 1H), 2.78 - 2.55 (m, 3H), 2.45 - 2.19 (m, 5H), 2.12 (s, 4H), 2.06 - 1.97 (m, 3H), 1.89 (d, J = 4.7 Hz, 7H), 1.71 - 1.55 (m, 3H), 1.54 - 1.39 (m, 21H), 1.36 (s, 13H), 1.31 - 1.26 (m, 12H), 1.06 (t, J = 10.5 Hz, 4H), 0.93 - 0.82 (m, 14H), 0.06 (s, 3H), 0.05 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 182.99, 175.20, 171.94, 170.75, 170.37, 168.93, 156.44, 77.43, 75.69, 66.32, 64.35, 63.93, 61.77, 57.52, 56.58, 54.28, 42.68, 40.37, 31.21, 30.95, 30.08, 29.88, 29.36, 28.85, 28.66, 26.01, 24.92, 24.16, 18.91, 18.48, 17.89, 15.48, -5.25, -5.29. MS (ESI) $[\text{MH}]^+$ calcd. 983.6, found 983.6

(1S,7S,10S,15aS,20R,20aS)-10-((1H-imidazol-5-yl)methyl)-7-(4-(tert-butoxy)benzyl)-N-(tert-butyl)-1-(((tert-butyl)dimethylsilyl)oxy)methyl)-3,6,9,12,15-pentaoxoicosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16]hexaazacyclooctadecine-20-carboxamide (20)

[0091] ^1H NMR (400 MHz, CDCl_3) δ : 9.92 (s, 1H), 7.49 (d, J = 4.8 Hz, 2H), 7.39 - 7.27 (m, 11H), 7.17 - 7.02 (m, 11H), 6.78 (d, J = 8.4 Hz, 2H), 6.69 (s, 1H), 4.48 (p, J = 6.9 Hz, 1H), 4.42 - 4.18 (m, 2H), 4.06 - 3.99 (m, 1H), 3.91 (dd, J = 11.4, 3.8 Hz, 1H), 3.78 (dd, J = 11.3, 4.4 Hz, 1H), 3.61 (dd, J = 9.8, 4.8 Hz, 1H), 3.48 - 3.37 (m, 1H), 3.32 - 2.99 (m, 8H), 2.83 (dd, J = 6.8, 3.4 Hz, 1H), 2.41 - 1.95 (m, 2H), 1.91 - 1.72 (m, 2H), 1.61 (s, 12H), 1.41 (s, 10H), 1.35 - 1.24 (m, 20H), 0.72 (s, 1H), 0.07 (s, 3H), 0.05 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 184.31, 175.20, 172.61, 170.67, 169.47, 168.12, 154.53, 142.31, 138.19, 136.81, 131.91, 129.97, 129.85, 128.43, 128.37, 120.09, 77.43, 63.49, 62.17, 61.55, 57.84, 54.97, 52.19, 51.15, 48.67, 43.36, 42.53, 41.75, 36.36, 31.84, 30.99, 29.92, 29.88, 29.44, 29.05, 28.67, 26.07, 24.49, 19.80, 18.56, -5.19. MS (ESI) $[\text{MH}]^+$ calcd. 866.5, found 866.5

(1S,4S,7S,13S,15aS,20R,20aS)-13-benzyl-N-(tert-butyl)-1-(((tert-butyl)dimethylsilyl)oxy)methyl)-4-methyl-3,6,9,12,15-pentaoxo-7-(2-oxo-2-(tritylamino)ethyl)icosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16]hexaazacyclooctadecine-20-carboxamide (21)

[0092] ^1H NMR (400 MHz, CDCl_3) δ : 7.94 (t, J = 9.8 Hz, 2H), 7.73 (d, J = 8.7 Hz, 1H), 7.51 (t, J = 6.7 Hz, 1H), 7.34 - 7.11 (m, 23H), 6.79 (s, 1H), 4.92 - 4.72 (m, 2H), 4.59 (dd, J = 16.3, 8.6 Hz, 1H), 4.25 (d, J = 11.5 Hz, 1H), 4.01 (d, J = 10.7 Hz, 1H), 3.82 (dd, J = 14.2, 7.6 Hz, 1H), 3.73 (s, 1H), 3.59 - 3.51 (m, 2H), 3.48 - 3.24 (m, 3H), 3.09 (dd, J = 13.8, 8.9 Hz, 1H), 2.96 - 2.68 (m, 6H), 2.59 (dd, J = 15.8, 3.6 Hz, 1H), 2.45 - 2.08 (m, 2H), 2.07 - 1.96 (m, 1H), 1.62 (s, 5H), 1.52 - 1.17 (m, 18H), 0.94 (s, 9H), 0.15 (s, 3H), 0.13 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ : 184.06, 175.64, 173.93, 171.01, 170.49, 169.15, 167.20, 144.28, 136.98, 129.61, 128.81, 128.49, 128.30, 127.48, 126.75, 77.55, 77.23, 76.91, 71.17, 63.63, 60.72, 57.12, 52.96, 51.54, 50.58, 49.71, 49.27, 45.35, 44.53, 37.27, 36.80, 33.93, 30.22, 29.24, 26.24, 24.32, 18.78, 17.89, 0.22, -5.11, -5.34. MS (ESI) $[\text{MH}]^+$ calcd. 1027.5, found 1027.6

(4S,10S,15aS,20R,20aS)-4-benzyl-N-(tert-butyl)-10-isobutyl-3,6,9,12,15-pentaoxoicosahydroazirino[1,2-a]pyrrolo[1,2-d][1,4,7,10,13,16]hexaazacyclooctadecine-20-carboxamide (22)

[0093] ^1H NMR (CDCl_3 , 400MHz) δ : 8.33 (bs, 1H), 8.17 (d, J = 9.1 Hz, 1H), 8.03 (d, J = 7.4 Hz, 1H), 7.54 (s, 1H), 7.21 (m, 5H), 6.04 (bs, 1H), 5.08 (bs, 1H), 4.58 (m, 1H), 3.87 (dd, J = 8.1 Hz, 5.0 Hz, 1H), 3.80 (bs, 1H), 2.48 (dd, J = 3.6 Hz, 1.8 Hz, 2H), 2.16 (m, 2H), 1.62-1.57 (m, 7H), 1.48 (m, 2H), 1.42 (m, 2H), 1.35 (m, 2H), 1.20 (m, 9H), 1.07 (m, 2H), 0.81 (m, 6H) ppm. MS (ESI) $[\text{MH}]^+$ calcd. 626.8, found 626.3 and 313.7 for $[\text{M}+2\text{H}]^{2+}/2$

7-mm conjugated 22 (23)

[0094] Cyclic peptide **22** was reacted as a crude material to yield conjugated peptide **23**. (see page 39) ^1H NMR (300 MHz, DMSO) δ 8.95 (s, 1H), 8.68 (d, J = 8.2 Hz, 1H), 8.32 (t, J = 6.3 Hz, 1H), 7.68 (d, J = 8.4 Hz, 1H), 7.51 (d, J = 9.4 Hz, 1H), 7.42 - 7.08 (m, 5H), 6.46 (s, 1H), 6.31 (s, 1H), 4.47 - 4.15 (m, 3H), 3.73 (dd, J = 20.5, 14.1 Hz, 3H), 3.45 - 3.33 (m, 2H), 3.17 (dd, J = 20.9, 13.0 Hz, 2H), 2.98 - 2.64 (m, 4H), 2.40 (s, 2H), 2.22 - 2.10 (m, 1H), 1.91 (ddd, J = 22.9, 13.4, 6.6 Hz, 2H), 1.79 - 1.40 (m, 5H), 1.30 - 1.15 (m, 9H), 0.93 - 0.74 (m, 6H). MS (ESI) $[\text{MH}]^+$ calcd. 818.4, found 818.3

HATU-mediated cyclization

[0095] Advantageously, the cyclization reaction can be run at least at 0.002M concentration of the starting amino acid/peptide. Preferably the concentration is run between 0.002M to 0.2M.

[0096] In another embodiment the process is run at a concentration of at least 0.1M of the starting amino acid/peptide. In another embodiment the process is run at a concentration of around 0.2M of the starting amino acid/peptide.

[0097] Traditionally, prior macrocyclizations cannot be performed at such high concentrations. This was demonstrated using a direct comparison with HATU process which fails in that the major products observed are derived from oligomerization and polymerization.

[0098] O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate ("HATU")-mediated cyclization of Pro-Gly-Gly-Gly was compared with the amino aldehyde/isocyanide cyclization method at 0.2 M. Upon completion, crude reaction mixtures were analyzed using liquid chromatography electrospray ionization mass spectrometry ("LC-MS(ESI)") to assess the level of selectivity for the desired cyclic peptide.

[0099] Mass spectrometry ("MS")-analysis of the HATU reaction revealed that cyclodimerization predominated at 0.2M (Figure 28). The major peaks at m/z = 537.2 and 559.2 correspond to the cyclodimer product, and its sodium chelate, respectively, while only trace amounts of the desired cyclic peptide was detected at m/z = 269. The cyclotrimer (m/z = 805) was also detected. The crude reaction mixture also contained several other unidentified byproducts.

[0100] Results from the MS peak analysis found 6 components.

Component 1: Peak at Scan 27.4. Top ions are 655 566 923

Component 2: Peak at Scan 28.4. Top ions are 982 744 714

Component 3: Peak at Scan 29.9. Top ions are 235 848 538

Component 4: Peak at Scan 30.9. Top ions are 559 827 560

Component 5: Peak at Scan 31.9. Top ions are 752 931

Component 6: Peak at Scan 32.7. Top ions are 537 295 446

[0101] The crude LC-MS(ESI) analysis for the isocyanide/amino aldehyde mediated cyclization revealed a high level of selectivity for the desired cyclic peptide ($m/z = 479$) at 0.2M (Figure 29). No trace peaks were detected for either dimer or trimer byproduct. Only characteristic fragmentation peaks and residual Pro-Gly-Gly-Gly remained in the reaction mixture.

[0102] Results from the MS peak analysis found 2 components.

Component 1: Peak at Scan 29.7. Top ions are 287 212 383 (Pro-Gly-Gly-Gly)

Component 2: Peak at Scan 30.9. Top ions are 479 396 240 (cyclic peptide)

Example 2

Methods and Materials

[0103]


Method A: ring-opening of aziridine-containing cyclic peptide is conducted using an aromatic thiol as the nucleophile (Method A). In a screw-cap vial equipped with a magnetic stirring bar was added aziridine-containing cyclic peptide (0.06 mmol) and aromatic thiol (0.066 mmol) were added to 0.2 ml of degassed CHCl_3 . NEt_3 (0.06 mmol) was then added to the solution and the reaction was stirred at room temperature for 3-4 hours. The reaction was then diluted with CH_2Cl_2 , and washed twice with saturated aqueous NH_4Cl (3 ml) followed by one wash with brine (3 ml). The organic layer was subsequently dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The crude ring-opened products were of high purity, but analytically pure samples were obtained following purification by flash column chromatography (5 % MeOH in EtOAc).

Method B: ring-opening of aziridine-containing cyclic peptide is carried out using an aliphatic thiol or imide as the nucleophile (Method B). In a screw-cap vial equipped with a magnetic stirring bar was added aziridine-containing cyclic peptide (0.06 mmol) and aliphatic thiol or imide (0.066 mmol) to 0.2 ml of degassed CHCl_3 . DBU (0.06 mmol) was then added to the

solution and the reaction was stirred at room temperature for 3 -4 hours. The reaction was then diluted with CH_2Cl_2 , and washed twice with saturated aqueous NH_4Cl (3 ml) followed by one wash with brine (3 ml). The organic layer was subsequently dried over Na_2SO_4 , filtered, and concentrated under reduced pressure. The crude ring-opened products were of high purity, but analytically pure samples were obtained following purification by flash column chromatography (5 % MeOH in EtOAc).

Results

[0104] Side chains including, but not limited to, fluorescent substituents have been appended to the products of macrocyclization (Scheme 3).



Scheme 3. Late-stage, site-specific attachment of fluorescent tags to a cyclic peptide.

[0105] The incorporation of an activated aziridine ring into the framework of a cyclic peptide provides a useful point for conjugation to various side-chains via nucleophilic ring-opening, a well developed methodology that has been demonstrated using nucleophilic biomolecules ranging from carbohydrates to biotin and farnesyl derivatives.¹⁴ As an example, cyclic peptide conjugation strategy can be demonstrated through the nucleophilic ring-opening of an aziridine moiety with the widely used fluorescent tag 7-mercapto-4-methyl-coumarin (7-mmc) (Scheme 3). In another aspect, a late-stage synthesis of a thioester residue by ring-opening of the cyclic peptide with commercially available thiobenzoic acid can also be used. The reaction proceeded smoothly, delivering the ring opened product in 98% yield. The late-stage incorporation of the thioester functionality avoids steps typically required for its protection from both oxidation and electrophilic reagents. Many other amino acid side chains, both natural and unnatural, can be installed by similar ring-opening protocols, opening doors for conformational optimization of cyclic peptides at a late stage of synthesis. Other nucleophiles such as aliphatic thiols, acids, secondary amines such as morpholine, and imides worked equally well during ring-opening.

[0106] In another aspect, cyclic peptides equipped with aziridine rings are subjected to ring-opening (Figure 30). In the course of this process, a side-chain of interest can be appended to the cyclic peptide scaffold at a late stage of synthesis. The ring-opening reactions were carried out using an appropriate base/nucleophile system that was selected on the basis of pK_a . The preferred reactions involved the employment of a base/nucleophile pair satisfying $\Delta pK_a = pK_a[NR_3H^+] - pK_a[Nucleophile] \approx -1$ (Figure 31).

[0107] These data suggest that strong ion pairing ($\Delta pK_a > 1$) between the base and nucleophile may result in decreased reactivity. The optimized conditions for ring-opening were found to proceed cleanly to afford the modified cyclic peptides in high purity following extractive work-up (Figure 32).

Example 3

[0108] X-ray analysis of cyclic product 1 was conducted.

Results

[0109]




Table 4. Crystal data and structure refinement for k07227.

Identification code	k07227
Empirical formula	C ₂₄ H ₃₈ N ₃ O ₃ Si
Formula weight	445.67
Temperature	150(2) K
Wavelength	0.71073 Å
Crystal system	Monoclinic
Space group	P 21
Unit cell dimensions	a = 14.590(2) Å α = 90°.
	b = 12.326(2) Å β = 94.718(7)°.
	c = 15.088(2) Å γ = 90°.
Volume	2704.2(7) Å ³
Z	4
Density (calculated)	1.097 Mg/m ³

Absorption coefficient	0.115 mm ⁻¹
F(000)	968
Crystal size	0.32 x 0.10 x 0.08 mm ³
Theta range for data collection	2.62 to 23.05°.
Index ranges	-15 ≤ h ≤ 15, -13 ≤ k ≤ 12, -15 ≤ l ≤ 16
Reflections collected	10115
Independent reflections	3828 [R(int) = 0.1359]
Completeness to theta = 23.05°	95.9 %
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	1.000 and 0.770
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	3828/1/559
Goodness-of-fit on F ²	1.044
Final R indices [I > 2σ(I)]	R1 = 0.0982, wR2 = 0.2560
R indices (all data)	R1 = 0.1715, wR2 = 0.3055
Largest diff. peak and hole	0.492 and -0.316 e.Å ⁻³

Table 5. Atomic coordinates (x 10⁴) and equivalent isotropic displacement parameters (Å² × 10³) for k07227. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
Si(1A)	7245(3)	4768(4)	8396(3)	68(1)
O(1A)	3310(7)	4696(12)	6317(8)	94(4)
O(2A)	4778(6)	4429(8)	4729(7)	62(3)
O(3A)	4070(7)	3807(11)	8551(7)	82(3)
O(4A)	6881(6)	4731(9)	7332(6)	64(3)
N(1A)	5222(7)	4298(9)	6198(7)	53(3)
N(2A)	2735(8)	3545(12)	7673(8)	74(4)
C(1A)	4577(10)	4187(12)	5473(9)	56(4)
C(2A)	3624(9)	3854(14)	5700(9)	61(4)
C(3A)	3884(8)	4758(14)	7161(9)	61(4)
C(4A)	4916(9)	4610(11)	7073(9)	57(4)
C(5A)	5474(9)	5392(12)	6570(9)	55(4)
C(6A)	6464(10)	5591(15)	6823(10)	72(5)
C(7A)	7044(16)	3423(18)	8820(12)	118(8)
C(8A)	6568(12)	5820(20)	9009(13)	105(6)
C(9A)	8501(12)	5097(14)	8489(14)	86(6)

	x	y	z	U(eq)
C(10A)	8607(13)	6160(30)	8022(19)	150(11)
C(11A)	9045(13)	4350(30)	8043(17)	169(14)
C(12A)	8846(14)	5210(20)	9454(15)	133(9)
C(13A)	3590(10)	3973(17)	7871(11)	76(5)
C(14A)	2268(10)	2780(16)	8263(10)	74(5)
C(15A)	2819(13)	1767(18)	8419(13)	103(6)
C(16A)	2093(11)	3415(16)	9107(10)	86(5)
C(17A)	1325(12)	2540(20)	7748(12)	110(7)
C(18A)	2957(9)	3707(15)	4897(9)	66(4)
C(19A)	2151(10)	3006(18)	5091(10)	77(5)
C(20A)	2294(13)	1877(19)	5232(12)	90(5)
C(21A)	1548(17)	1230(20)	5361(14)	112(7)
C(22A)	725(19)	1640(30)	5502(16)	128(9)
C(23A)	525(13)	2770(30)	5385(13)	116(9)
C(24A)	1270(12)	3480(20)	5217(11)	101(7)
Si(1B)	8204(3)	2344(4)	2871(3)	68(1)
O(1B)	6478(7)	2195(11)	6755(7)	86(3)
O(2B)	4724(6)	1912(7)	5174(6)	53(2)
O(3B)	8659(6)	1106(10)	6205(7)	75(3)
O(4B)	7241(6)	2259(8)	3360(6)	66(3)
N(1B)	6249(7)	1815(8)	4890(7)	48(3)
N(2B)	7884(8)	963(11)	7449(9)	69(3)
C(1B)	5494(10)	1740(12)	5431(9)	55(4)
C(2B)	5816(9)	1389(12)	6398(9)	53(3)
C(3B)	7344(9)	2232(14)	6288(8)	59(4)
C(4B)	7165(9)	2098(12)	5256(9)	60(4)
C(5B)	6589(10)	2897(14)	4693(11)	70(4)
C(6B)	6790(11)	3147(14)	3790(11)	74(5)
C(7B)	8766(12)	968(16)	3042(13)	97(6)
C(8B)	8926(11)	3466(19)	3390(11)	98(6)
C(9B)	7893(10)	2620(14)	1666(10)	71(5)
C(10B)	7510(13)	3777(14)	1548(11)	85(5)
C(11B)	7165(14)	1807(19)	1279(12)	103(6)
C(12B)	8748(12)	2536(19)	1143(11)	100(6)

	x	y	z	U(eq)
C(13B)	8015(10)	1332(15)	6630(11)	69(4)
C(14B)	8453(10)	163(13)	7952(11)	66(4)
C(15B)	8439(13)	-842(17)	7445(15)	111(7)
C(16B)	9438(11)	564(17)	8135(13)	97(6)
C(17B)	8040(11)	0(20)	8825(11)	103(7)
C(18B)	5021(9)	1225(13)	6976(9)	61(4)
C(19B)	5242(8)	526(15)	7787(10)	65(5)
C(20B)	5337(10)	-610(14)	7633(11)	65(4)
C(21B)	5477(11)	-1260(20)	8343(13)	101(7)
C(22B)	5524(12)	-940(20)	9202(18)	105(7)
C(23B)	5439(11)	130(30)	9374(13)	100(7)
C(24B)	5292(10)	878(19)	8643(11)	86(6)

Table 6. Bond lengths [Å] and angles [°] for k07227.

Si(1A)-O(4A)	1.649(10)
Si(1A)-C(7A)	1.81(2)
Si(1A)-C(9A)	1.871(18)
Si(1A)-C(8A)	1.91(2)
O(1A)-C(3A)	1.468(17)
O(1A)-C(2A)	1.491(18)
O(2A)-C(1A)	1.221(15)
O(3A)-C(13A)	1.211(17)
O(4A)-C(6A)	1.418(18)
N(1A)-C(1A)	1.390(17)
N(1A)-C(4A)	1.479(16)
N(1A)-C(5A)	1.494(18)
N(2A)-C(13A)	1.36(2)
N(2A)-C(14A)	1.50(2)
C(1A)-C(2A)	1.52(2)
C(2A)-C(18A)	1.502(19)
C(3A)-C(13A)	1.53(2)
C(3A)-C(4A)	1.533(18)
C(4A)-C(5A)	1.507(19)
C(5A)-C(6A)	1.48(2)
C(9A)-C(11A)	1.42(3)
C(9A)-C(10A)	1.50(3)

C(9A)-C(12A)	1.51(3)
C(14A)-C(15A)	1.49(3)
C(14A)-C(16A)	1.53(2)
C(14A)-C(17A)	1.55(2)
C(18A)-C(19A)	1.51(2)
C(19A)-C(20A)	1.42(3)
C(19A)-C(24A)	1.44(2)
C(20A)-C(21A)	1.38(3)
C(21A)-C(22A)	1.34(3)
C(22A)-C(23A)	1.44(4)
C(23A)-C(24A)	1.44(3)
Si(1B)-O(4B)	1.643(9)
Si(1B)-C(9B)	1.868(16)
Si(1B)-C(8B)	1.871(19)
Si(1B)-C(7B)	1.893(18)
O(1B)-C(2B)	1.457(18)
O(1B)-C(3B)	1.496(16)
O(2B)-C(1B)	1.177(15)
O(3B)-C(13B)	1.214(16)
O(4B)-C(6B)	1.457(18)
N(1B)-C(1B)	1.428(17)
N(1B)-C(4B)	1.446(17)
N(1B)-C(5B)	1.461(19)
N(2B)-C(13B)	1.344(19)
N(2B)-C(14B)	1.461(19)
C(1B)-C(2B)	1.56(2)
C(2B)-C(18B)	1.522(18)
C(3B)-C(13B)	1.54(2)
C(3B)-C(4B)	1.567(18)
C(4B)-C(5B)	1.51(2)
C(5B)-C(6B)	1.45(2)
C(9B)-C(12B)	1.53(2)
C(9B)-C(10B)	1.54(2)
C(9B)-C(11B)	1.54(2)
C(14B)-C(15B)	1.45(3)

C(14B)-C(17B)	1.51(2)
C(14B)-C(16B)	1.52(2)
C(18B)-C(19B)	1.51(2)
C(19B)-C(24B)	1.36(2)
C(19B)-C(20B)	1.43(2)
C(20B)-C(21B)	1.34(2)
C(21B)-C(22B)	1.35(3)
C(22B)-C(23B)	1.35(3)
C(23B)-C(24B)	1.44(3)
O(4A)-Si(1A)-C(7A)	105.7(8)
O(4A)-Si(1A)-C(9A)	108.3(7)
C(7A)-Si(1A)-C(9A)	111.0(9)
O(4A)-Si(1A)-C(8A)	110.6(8)
C(7A)-Si(1A)-C(8A)	110.1(10)
C(9A)-Si(1A)-C(8A)	111.0(9)
C(3A)-O(1A)-C(2A)	113.3(11)
C(6A)-O(4A)-Si(1A)	126.6(10)
C(1A)-N(1A)-C(4A)	119.6(10)
C(1A)-N(1A)-C(5A)	121.0(11)
C(4A)-N(1A)-C(5A)	60.9(8)
C(13A)-N(2A)-C(14A)	124.4(13)
O(2A)-C(1A)-N(1A)	120.1(13)
O(2A)-C(1A)-C(2A)	124.6(12)
N(1A)-C(1A)-C(2A)	115.0(11)
O(1A)-C(2A)-C(18A)	112.0(12)
O(1A)-C(2A)-C(1A)	106.7(12)
C(18A)-C(2A)-C(1A)	113.3(11)
O(1A)-C(3A)-C(13A)	113.7(13)
O(1A)-C(3A)-C(4A)	114.4(11)
C(13A)-C(3A)-C(4A)	108.6(12)
N(1A)-C(4A)-C(5A)	60.0(8)
N(1A)-C(4A)-C(3A)	118.7(11)
C(5A)-C(4A)-C(3A)	122.9(12)
C(6A)-C(5A)-N(1A)	116.8(12)
C(6A)-C(5A)-C(4A)	122.1(12)
N(1A)-C(5A)-C(4A)	59.1(8)

O(4A)-C(6A)-C(5A)	112.4(12)
C(11A)-C(9A)-C(10A)	105(2)
C(11A)-C(9A)-C(12A)	111.3(18)
C(10A)-C(9A)-C(12A)	109(2)
C(11A)-C(9A)-Si(1A)	113.9(16)
C(10A)-C(9A)-Si(1A)	107.0(12)
C(12A)-C(9A)-Si(1A)	109.9(13)
O(3A)-C(13A)-N(2A)	125.1(16)
O(3A)-C(13A)-C(3A)	121.6(14)
N(2A)-C(13A)-C(3A)	113.2(14)
C(15A)-C(14A)-N(2A)	110.8(13)
C(15A)-C(14A)-C(16A)	114.8(16)
N(2A)-C(14A)-C(16A)	106.7(15)
C(15A)-C(14A)-C(17A)	111.4(17)
N(2A)-C(14A)-C(17A)	104.4(13)
C(16A)-C(14A)-C(17A)	108.1(14)
C(2A)-C(18A)-C(19A)	112.2(12)
C(20A)-C(19A)-C(24A)	120.2(19)
C(20A)-C(19A)-C(18A)	118.9(16)
C(24A)-C(19A)-C(18A)	121(2)
C(21A)-C(20A)-C(19A)	119(2)
C(22A)-C(21A)-C(20A)	122(3)
C(21A)-C(22A)-C(23A)	122(3)
C(22A)-C(23A)-C(24A)	118(2)
C(23A)-C(24A)-C(19A)	118(2)
O(4B)-Si(1B)-C(9B)	107.5(6)
O(4B)-Si(1B)-C(8B)	109.4(7)
C(9B)-Si(1B)-C(8B)	110.9(8)
O(4B)-Si(1B)-C(7B)	105.0(7)
C(9B)-Si(1B)-C(7B)	111.3(9)
C(8B)-Si(1B)-C(7B)	112.5(9)
C(2B)-O(1B)-C(3B)	114.0(11)
C(6B)-O(4B)-Si(1B)	125.9(10)
C(1B)-N(1B)-C(4B)	122.0(11)
C(1B)-N(1B)-C(5B)	117.7(11)

C(4B)-N(1B)-C(5B)	62.6(9)
C(13B)-N(2B)-C(14B)	126.0(13)
O(2B)-C(1B)-N(1B)	124.1(13)
O(2B)-C(1B)-C(2B)	124.4(12)
N(1B)-C(1B)-C(2B)	111.4(11)
O(1B)-C(2B)-C(18B)	113.1(11)
O(1B)-C(2B)-C(1B)	107.3(11)
C(18B)-C(2B)-C(1B)	112.9(11)
O(1B)-C(3B)-C(13B)	111.0(12)
O(1B)-C(3B)-C(4B)	112.7(10)
C(13B)-C(3B)-C(4B)	107.8(12)
N(1B)-C(4B)-C(5B)	59.2(8)
N(1B)-C(4B)-C(3B)	118.6(10)
C(5B)-C(4B)-C(3B)	122.0(13)
C(6B)-C(5B)-N(1B)	118.9(14)
C(6B)-C(5B)-C(4B)	121.4(13)
N(1B)-C(5B)-C(4B)	58.2(9)
C(5B)-C(6B)-O(4B)	113.1(13)
C(12B)-C(9B)-C(10B)	107.8(14)
C(12B)-C(9B)-C(11B)	109.0(15)
C(10B)-C(9B)-C(11B)	109.1(14)
C(12B)-C(9B)-Si(1B)	110.2(11)
C(10B)-C(9B)-Si(1B)	109.8(12)
C(11B)-C(9B)-Si(1B)	110.8(12)
O(3B)-C(13B)-N(2B)	125.6(15)
O(3B)-C(13B)-C(3B)	119.1(14)
N(2B)-C(13B)-C(3B)	114.7(13)
C(15B)-C(14B)-N(2B)	108.6(13)
C(15B)-C(14B)-C(17B)	110.9(16)
N(2B)-C(14B)-C(17B)	107.0(13)
C(15B)-C(14B)-C(16B)	110.2(15)
N(2B)-C(14B)-C(16B)	111.3(13)
C(17B)-C(14B)-C(16B)	108.9(15)
C(19B)-C(18B)-C(2B)	114.7(11)
C(24B)-C(19B)-C(20B)	118.0(16)
C(24B)-C(19B)-C(18B)	125.4(18)

C(20B)-C(19B)-C(18B)	116.5(14)
C(21B)-C(20B)-C(19B)	117.9(18)
C(20B)-C(21B)-C(22B)	125(3)
C(23B)-C(22B)-C(21B)	119(2)
C(22B)-C(23B)-C(24B)	119(2)
C(19B)-C(24B)-C(23B)	121(2)

Table 7. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for k07227. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Si(1A)	59(2)	75(3)	67(3)	-5(2)	-2(2)	-7(2)
O(1A)	67(6)	116(10)	101(8)	-29(8)	23(6)	-1(7)
O(2A)	66(6)	55(6)	64(7)	-6(5)	11(5)	0(5)
O(3A)	61(6)	118(10)	66(7)	14(7)	8(5)	-7(6)
O(4A)	55(5)	64(6)	76(6)	5(6)	11(4)	-3(5)
N(1A)	54(7)	54(7)	54(7)	-6(6)	19(6)	1(6)
N(2A)	51(7)	99(11)	74(8)	17(8)	12(6)	6(7)
C(1A)	77(10)	49(9)	40(8)	1(7)	-2(7)	3(7)
C(2A)	60(9)	70(10)	52(8)	-9(8)	3(7)	6(8)
C(3A)	44(7)	70(10)	70(9)	-7(9)	5(6)	7(7)
C(4A)	61(8)	46(9)	65(9)	9(7)	7(7)	0(7)
C(5A)	51(8)	48(9)	64(9)	4(7)	4(7)	4(7)
C(6A)	62(10)	86(13)	69(10)	20(10)	18(8)	-7(9)
C(7A)	170(20)	110(17)	71(12)	20(12)	-25(12)	-17(16)
C(8A)	83(12)	124(18)	109(14)	-12(13)	23(10)	3(12)
C(9A)	69(10)	59(11)	130(16)	44(11)	17(11)	11(9)
C(10A)	47(11)	180(30)	220(30)	30(20)	-10(13)	-16(14)
C(11A)	61(13)	300(50)	140(20)	-60(20)	-2(13)	13(19)
C(12A)	101(14)	170(20)	122(17)	57(17)	-40(13)	-18(15)
C(13A)	49(9)	122(15)	59(9)	-5(10)	12(8)	-2(10)
C(14A)	58(9)	101(14)	64(10)	15(9)	17(7)	-2(9)
C(15A)	95(14)	103(16)	113(15)	37(13)	27(11)	15(12)
C(16A)	85(11)	105(15)	68(11)	-27(10)	17(8)	-17(10)
C(17A)	82(12)	150(20)	95(13)	10(14)	15(10)	-60(13)
C(18A)	63(9)	86(12)	51(8)	3(8)	8(7)	-16(9)
C(19A)	59(10)	122(18)	49(9)	5(10)	-2(7)	2(10)

	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
C(20A)	97(13)	92(15)	85(12)	4(11)	31(10)	-19(12)
C(21A)	118(18)	121(19)	100(15)	8(13)	21(13)	-48(16)
C(22A)	95(19)	190(30)	102(17)	11(19)	0(13)	-26(19)
C(23A)	50(11)	210(30)	86(14)	9(17)	1(9)	1(16)
C(24A)	63(12)	170(20)	68(11)	-7(13)	-7(8)	0(13)
Si(1B)	60(3)	75(3)	70(3)	8(3)	13(2)	2(2)
O(1B)	71(7)	106(9)	83(7)	14(7)	14(5)	3(7)
O(2B)	51(6)	44(5)	65(6)	-3(5)	3(4)	3(4)
O(3B)	52(6)	99(9)	75(7)	4(6)	9(5)	9(6)
O(4B)	69(6)	60(6)	71(6)	11(6)	24(5)	5(5)
N(1B)	60(7)	36(6)	49(6)	-1(5)	6(5)	13(5)
N(2B)	50(7)	75(9)	84(9)	10(8)	13(6)	3(7)
C(1B)	57(10)	48(9)	61(9)	-17(7)	6(7)	-4(7)
C(2B)	45(8)	50(9)	65(9)	-11(7)	4(6)	0(7)
C(3B)	51(8)	72(10)	54(8)	-1(8)	3(6)	5(8)
C(4B)	47(8)	62(10)	73(9)	25(8)	17(7)	8(7)
C(5B)	59(9)	65(11)	86(12)	6(9)	8(8)	-9(8)
C(6B)	82(11)	66(11)	77(11)	15(9)	21(8)	17(9)
C(7B)	84(12)	87(14)	123(15)	34(12)	28(10)	36(11)
C(8B)	64(10)	147(19)	84(12)	-19(13)	10(8)	-22(11)
C(9B)	56(9)	78(13)	82(11)	10(9)	14(8)	-12(8)
C(10B)	113(13)	64(11)	77(11)	23(10)	11(9)	-7(10)
C(11B)	121(16)	104(16)	84(12)	-14(12)	11(11)	-22(13)
C(12B)	97(13)	117(17)	89(12)	32(13)	36(10)	13(12)
C(13B)	48(9)	85(12)	75(11)	-3(10)	18(8)	-9(8)
C(14B)	66(10)	50(10)	81(11)	11(9)	3(8)	0(8)
C(15B)	88(13)	88(17)	152(19)	37(15)	-11(12)	29(11)
C(16B)	62(10)	112(17)	116(15)	14(13)	-1(10)	-1(10)
C(17B)	80(11)	150(20)	74(11)	55(13)	12(9)	23(12)
C(18B)	48(8)	73(11)	61(9)	-5(8)	7(6)	6(7)
C(19B)	34(7)	105(15)	56(10)	14(10)	8(6)	-16(8)
C(20B)	67(10)	64(11)	64(10)	11(9)	3(7)	1(8)
C(21B)	76(12)	150(20)	74(13)	14(16)	6(10)	-11(12)
C(22B)	67(12)	120(20)	120(20)	35(17)	13(11)	-20(13)

	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
C(23B)	60(11)	180(30)	64(12)	9(16)	10(8)	-26(14)
C(24B)	59(10)	138(17)	63(11)	-3(12)	14(8)	-8(10)

Table 8. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for k07227.

	x	y	z	U(eq)
H(2A)	2439	3731	7164	89
H(2AB)	3677	3148	6028	73
H(3AA)	3805	5507	7399	73
H(4AA)	5261	4254	7594	69
H(5AA)	5126	6016	6284	66
H(6AA)	6788	5688	6277	86
H(6AB)	6530	6272	7171	86
H(7AA)	7391	2890	8499	177
H(7AB)	6386	3254	8734	177
H(7AC)	7246	3395	9455	177
H(8AA)	5915	5621	8958	157
H(8AB)	6644	6535	8745	157
H(8AC)	6797	5834	9638	157
H(10A)	9261	6348	8041	225
H(10B)	8274	6725	8319	225
H(10C)	8356	6093	7402	225
H(11A)	9691	4578	8109	254
H(11B)	8832	4327	7410	254
H(11C)	8989	3625	8301	254
H(12A)	9506	5375	9499	199
H(12B)	8742	4534	9768	199
H(12C)	8516	5804	9723	199
H(15A)	3407	1942	8750	154
H(15B)	2933	1440	7847	154
H(15C)	2478	1254	8764	154
H(16A)	2680	3561	9449	128
H(16B)	1699	2986	9469	128
H(16C)	1788	4103	8942	128
H(17A)	1421	2134	7206	166
H(17B)	1014	3229	7588	166

	x	y	z	U(eq)
H(17C)	944	2115	8123	166
H(18A)	3279	3371	4415	80
H(18B)	2729	4426	4688	80
H(20A)	2894	1576	5238	108
H(21A)	1622	461	5349	135
H(22A)	261	1165	5684	154
H(23A)	-81	3040	5417	140
H(24A)	1185	4248	5190	122
H(2B)	7412	1227	7705	83
H(2BB)	6146	681	6365	64
H(3BA)	7647	2950	6413	71
H(4BA)	7677	1752	4955	72
H(5BA)	6321	3505	5026	84
H(6BA)	7190	3796	3799	89
H(6BB)	6209	3326	3435	89
H(7BA)	8930	851	3678	146
H(7BB)	9322	938	2721	146
H(7BC)	8336	402	2817	146
H(8BA)	8613	4161	3276	147
H(8BB)	9522	3478	3134	147
H(8BC)	9023	3346	4033	147
H(10D)	7339	3913	916	127
H(10E)	7981	4300	1769	127
H(10F)	6967	3855	1883	127
H(11D)	7007	1971	649	154
H(11E)	6613	1863	1604	154
H(11F)	7412	1069	1336	154
H(12D)	8576	2679	512	149
H(12E)	9009	1805	1211	149
H(12F)	9206	3070	1372	149
H(15D)	8712	-718	6882	166
H(15E)	7802	-1087	7323	166
H(15F)	8792	-1398	7788	166
H(16D)	9719	649	7571	146

	x	y	z	U(eq)
H(16E)	9792	36	8509	146
H(16F)	9437	1264	8442	146
H(17D)	7413	-280	8715	154
H(17E)	8026	694	9140	154
H(17F)	8414	-521	9189	154
H(18C)	4814	1945	7173	73
H(18D)	4502	892	6609	73
H(20B)	5304	-894	7046	78
H(21B)	5548	-2016	8234	121
H(22B)	5615	-1446	9672	126
H(23B)	5475	386	9970	120
H(24B)	5229	1630	8761	104

REFERENCES

[0110]

1. (1) Drahl, S. Chem. Eng. News 2009, 87, 31.
2. (2) a) Shao, Y.; Lu, W.; Kent, S. B. H. Tetrahedron. Lett. 1998, 39, 3911. b) Tam, J. P.; Lu, Y.-A.; Yu, Q. J. Am. Chem. Soc. 1999, 121, 4316. c) Bernal, F.; Tyler, A. F.; Korsmeyer, S. J.; Walensky, L.D.; Verdine, G. L. J. Am. Chem. Soc. 2007, 129, 2456. d) Home, W.S.; Stout, C. D.; Ghadiri, M. R. J. Am. Chem. Soc. 2003, 125, 9372. e) Blackwell, H. E.; Sadowsky, J. D.; Howard, R. J.; Sampson, J. N.; Chao, J. A.; Steinmetz, W. E.; O'Leary, D. J.; Grubbs, R. H. J. Org. Chem. 2001, 66, 5291. f) Kirshenbaum, K.; Arora, P. S. Nat. Chem. Bio. 2008, 4, 527. g) Hebach, C.; Kazmaier, U. Chem. Comm. 2003, 9, 596. h) Reid, R. C.; Kelso, M. J.; Scanlon, M. J.; Fairlie, D. P. J. Am. Chem. Soc. 2002, 124, 5673. i) Al-Obeidi, F.; de Castrucci, A. M. L.; Hadley, M. E.; Hruby, V. J. J. Med. Chem. 1989, 32, 2555. j) Geyer, A.; Mueller, G.; Kessler, H. J. Am. Chem. Soc. 1994, 116, 7735.
3. (3) Schmuck, C.; Wienand, W. J. Am. Chem. Soc. 2003, 125, 452.
4. (4) A. F. Spatola and P. Romanovskis, in Combinatorial Peptide and Nonpeptide Libraries; Jung, G., Ed.; VCH: Weinheim, 1996; p 328.
5. (5) H. C. Gilon, C. Mang, E. Lohof, A. Friedler, H. Kessler, in Synthesis of Peptides and Peptidomimetics; Goodman, M., Ed.; Thieme: Stuttgart, 2004; pp 461.
6. (6) Zimmer, S.; Hoffmann, E.; Jung, G.; Kessler, H. Liebigs. Ann. Chem. 1993, 497.
7. (7) Schmidt, U.; Langner, J. J. Pept. Res. 1997, 49, 67.
8. (8) a) Hili, R.; Yudin, A. K. J. Am. Chem. Soc. 2009, 131, 16404. b) Baktharaman, S.; Hili, R.; Yudin, A. K. Aldrichimica Acta 2008, 41, 109. c) Hili, R.; Baktharam, S.; Yudin, A.

- K. Eur. J. Org. Chem. 2008, 5201. d) Hili, R.; Yudin, A. K. Angew. Chem. Int. Ed. 2008, 47, 4188. e) Yudin, A. K.; Hili, R. Chem. Eur. J. 2007, 13, 6538. f) Hili, R.; Yudin, A. K. J. Am. Chem. Soc. 2006, 128, 14772.
9. (9) Roccatano, D., Colombo, G., Fioroni, M., Mark, A.E. Proc. Nat. Acad. Sci. 2002, 99, 12179.
10. (10) a) Domling, A.; Ugi, I. Angew. Chem. Int. Ed. 1997, 39, 3168. b) Dömling, A.; Ugi, I. Angew. Chem. Int. Ed. 2000, 39, 3169. c) Thompson, M. J.; Chen, B. J. Org. Chem. 2009, 74, 7084.
11. (11) Demharter, A.; Hörl, W.; Herdtweck, E.; Ugi, I. Angew. Chem. Int. Ed. 1996, 36, 173.
12. (12) For a recent example describing consecutive Ugi reactions to access glycine-based peptoid structures from formaldehyde, see: Vercillo, O. E., Andrade, C. K. Z., Wessjohann, L. A. Org. Lett. 2008, 10, 205.
13. (13) A. Failli, H. Immer, M. Götz, Can. J. Chem. 1979, 57, 3257.
14. (14) Galonić, D. P.; Ide, N. D.; van der Donk, W. A.; Gin, D. Y. J. Am. Chem. Soc. 2005, 127, 7359.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description


- **DRAHL, S.** Chem. Eng. News, 2009, vol. 87, 31- [0110]
- **SHAO, Y. LU, W. KENT, S. B. H.** Tetrahedron. Lett., 1998, vol. 39, 3911- [0110]
- **TAM, J. P. LU, Y.-A. YU, Q. J.** Am. Chem. Soc., 1999, vol. 121, 4316- [0110]
- **KORSMEYER, S. J. WALENSKY, L. D. VERDINE, G. L. J.** Am. Chem. Soc., 2007, vol. 129, 2456- [0110]
- **HOME, W. S. STOUT, C. D. GHADIRI, M. R. J.** Am. Chem. Soc., 2003, vol. 125, 9372- [0110]
- **BLACKWELL, H. E. SADOWSKY, J. D. HOWARD, R. J. SAMPSON; J. N. CHAO; J. A. STEINMETZ, W. E. O'LEARY, D. J. GRUBBS, R. H. J.** Org. Chem., 2001, vol. 66, 5291- [0110]
- **KIRSHENBAUM, K. ARORA, P. S.** Nat. Chem. Bio., 2008, vol. 4, 527- [0110]
- **HEBACH, C. KAZMAIER, U.** Chem. Comm., 2003, vol. 9, 596- [0110]
- **REID, R. C. KELSO, M. J. SCANLON, M. J. FAIRLIE, D. P. J.** Am. Chem. Soc., 2002, vol. 124, 5673- [0110]

- **AL-OBEIDI, F.DE CASTRUCCI, A. M. L.HADLEY, M. E.HRUBY, V. J.J.** Med. Chem., 1989, vol. 32, 2555- [\[0110\]](#)
- **GEYER, A.MUELLER, G.KESSLER, H.J.** Am. Chem. Soc., 1994, vol. 116, 7735- [\[0110\]](#)
- **SCHMUCK, C.WIENAND, W.J.** Am. Chem. Soc., 2003, vol. 125, 452- [\[0110\]](#)
- **A. F. SPATOLAP. ROMANOVSKIS**Combinatorial Peptide and Nonpeptide LibrariesVCH19960000328- [\[0110\]](#)
- **H. C. GILONC. MANGE. LOHOFA. FRIEDLERH. KESSLER**Synthesis of Peptides and PeptidomimeticsThieme20040000461- [\[0110\]](#)
- **ZIMMER, S.HOFFMANN, E.JUNG, G.KESSLER, H.**Liebigs. Ann. Chem., 1993, 497- [\[0110\]](#)
- **SCHMIDT, U.LANGNER, J.J.** Pept. Res., 1997, vol. 49, 67- [\[0110\]](#)
- **HILI, R.YUDIN, A. K.**J. Am. Chem. Soc., 2009, vol. 131, 16404- [\[0110\]](#)
- **BAKTHARAMAN, S.HILI, R.YUDIN, A. K.**Aldrichimica Acta, 2008, vol. 41, 109- [\[0110\]](#)
- **HILI, R.BAKTHARAM, S.YUDIN, A. K.**Eur. J. Org. Chem., 2008, 5201- [\[0110\]](#)
- **HILI, R.YUDIN, A. K.**Angew. Chem. Int. Ed., 2008, vol. 47, 4188- [\[0110\]](#)
- **YUDIN, A. K.HILI, R.**Chem. Eur. J., 2007, vol. 13, 6538- [\[0110\]](#)
- **HILI, R.YUDIN, A. K.**J. Am. Chem. Soc., 2006, vol. 128, 14772- [\[0110\]](#)
- **ROCCATANO, D.COLOMBO, G.FIORONI, M.MARK, A.E.**Proc. Nat. Acad. Sci., 2002, vol. 99, 12179- [\[0110\]](#)
- **DOMLING, A.UGI, I.**Angew. Chem. Int. Ed., 1997, vol. 39, 3168- [\[0110\]](#)
- **DÖMLING, A.UGI, I.**Angew. Chem. Int. Ed., 2000, vol. 39, 3169- [\[0110\]](#)
- **THOMPSON, M. J.**Chen, B. J. Org. Chem., 2009, vol. 74, 7084- [\[0110\]](#)
- **DEMARTER, A.HÖRL, W.HERDTWECK, E.UGI, I.**Angew. Chem. Int. Ed., 1996, vol. 36, 173- [\[0110\]](#)
- **VERCILLO, O. E.ANDRADE, C. K. Z.WESSJOHANN, L. A.**Org. Lett., 2008, vol. 10, 205- [\[0110\]](#)
- **A. FAILLIH. IMMERM. GÖTZ**Can. J. Chem., 1979, vol. 57, 3257- [\[0110\]](#)
- **GALONIĆ, D. P.IDE, N. D.VAN DER DONK, W. A.GIN, D. Y.**J. Am. Chem. Soc., 2005, vol. 127, 7359- [\[0110\]](#)

Patentkrav

1. Fremgangsmåde til dannelse af et cyklisk peptid, hvilken fremgangsmåde omfatter følgende trin:

(i) omsætning af en aminendegruppe af et peptid eller en aminosyre med at aziridinaldehyd med følgende struktur:



og

(ii) omsætning af blandingen i trin (i) med et isocyanid.

2. Fremgangsmåde ifølge krav 1, hvor aziridinaldehydet er aziridinaldehydbisulfit.

3. Fremgangsmåde ifølge et hvilket som helst af kravene 1 til 2, hvor isocyanidet er tertær-butylisocyanid.

4. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor fremgangsmåden er en ettrinsfremgangsmåde.

5. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor omsætningerne finder sted under støkiometriske betingelser.

6. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor omsætningen foretages med en aminosyre i vand.


7. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvilken fremgangsmåde yderligere omfatter følgende trin: (iii) omsætning af blandingen i trin (ii) med henblik på at åbne aziridinringen.

8. Fremgangsmåde ifølge krav 7, hvor blandingen i trin (ii) omsættes til en thioester.

9. Fremgangsmåde ifølge krav 7, hvor thiolsyren er thiobenzoesyre.

10. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvilken fremgangsmåde yderligere omfatter det trin, at der inkorporeres fluorescerende mærker i det cykliske peptid ved nukleofil ringåbning af aziridingruppen.

11. Forbindelse ifølge nedenstående almene formel:




hvor,

R¹ og Rⁿ hver især uafhængigt af hinanden er valgt fra en


aminosyresidekædegruppe, og R^2 er en alkylgruppe.

12. Forbindelse ifølge krav 11, hvor R^2 er en tertær-butylgruppe.


13. Forbindelse med en formel ifølge formel (I) til (VII):




Formel (I)




Formel (II)




Formel (III)




Formel (IV)



Formel (V)




Formel (VI)




Formel (VII)


DRAWINGS




Formula (I)




Formula (II)




Formula (III)




Formula (IV)



Formula (V)




Formula (VI)




Formula (VII)

Crude reaction from D-Leu



Crude reaction from L-Leu



Mixture of isomers




Figure 1. ^1H NMR comparison of crude reactions mixtures using isomers of leucine.




Figure 2. Mismatched reaction leads only to aminor products.




Figure 3. Matched reaction is fast and high yielding




Figure 4. Chemical structures of cyclic products synthesized by Methods 1 – 3.




Figure 5. ^1H and ^{13}C NMR spectra for cyclic product **1**.




Figure 6. ^1H and ^{13}C NMR spectra for cyclic product 2.




Figure 7. ^1H and ^{13}C NMR spectra for cyclic product 3.




Figure 8. ^1H and ^{13}C NMR spectra for cyclic product 4.




Figure 9. ^1H and ^{13}C NMR spectra for cyclic product **5**.




Figure 10. ^1H and ^{13}C NMR spectra for cyclic product 6.




Figure 11. ^1H and ^{13}C NMR spectra for cyclic product 7.




Figure 12. ^1H and ^{13}C NMR spectra for cyclic product **8**.




Figure 13. ^1H NMR spectrum for cyclic product 9.




Figure 14. ^1H and ^{13}C NMR spectra for cyclic product 10.




Figure 15. ^1H and ^{13}C NMR spectra for cyclic product 11.




Figure 16. ^1H and ^{13}C NMR spectra for cyclic product **12**.




Figure 17. ^1H and ^{13}C NMR spectra for cyclic product 13.




Figure 18. ^1H and ^{13}C NMR spectra for cyclic product 14.




Figure 19. ^1H and ^{13}C NMR spectra for cyclic product **15**.




Figure 20. ^1H , COSY and ^{13}C NMR spectra for cyclic product 16.




Figure 20. Continued.




Figure 21. ^1H and ^{13}C NMR spectra for cyclic product 17.




Figure 22. ^1H and ^{13}C NMR spectra for cyclic product 18.




Figure 23. ^1H and ^{13}C NMR spectra for cyclic product **19**.




Figure 24. ^1H and ^{13}C NMR spectra for cyclic product 20.




Figure 25. ^1H and ^{13}C NMR spectra for cyclic product **21**.




Figure 26. ^1H and COSY NMR spectra for cyclic product **22**.




Figure 27. ^1H NMR spectrum for cyclic product 23.





Figure 28. Crude LC-MS(ESI) analysis of HATU-mediated cyclization of Pro-Gly-Gly-Gly at 0.2M





Figure 29. Crude LC-MS(ESI) analysis of isocyanide/amino aldehyde-mediated cyclization of Pro-Gly-Gly at 0.2M




Figure 30. Nucleophilic ring-opening of aziridine-containing cyclic peptides.




Figure 31. pK_a 's of thiols and ammonium ions and ΔpK_a of ion pairs.




Figure 32. Crude reaction mixture of thiol opened product.