



(43) International Publication Date
26 July 2012 (26.07.2012)

(10) International Publication Number
WO 2012/098049 A1

- (51) **International Patent Classification:**
C07D 413/12 (2006.01) *C07D 419/12* (2006.01)
- (21) **International Application Number:**
PCT/EP2012/050470
- (22) **International Filing Date:**
13 January 2012 (13.01.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
11151280.2 18 January 2011 (18.01.2011) EP
11184685.3 11 October 2011 (11.10.2011) EP
- (71) **Applicant (for all designated States except US):** **DSM SINOCHEM PHARMACEUTICALS NETHERLANDS B.V.** [NL/NL]; P.O. Box 245, Alexander Fleminglaan 1, NL-2613 AX Delft (NL).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **BESSEMBINDER, Karin, Henderika, Maria** [NL/NL]; P.O. Box 4, NL-6100 AA Echt (NL). **HEEMSKERK, Dennis** [NL/NL]; P.O. Box 4, NL-6100 AA Echt (NL). **LANGE, DE, Ben** [NL/NL]; P.O. Box 4, NL-6100 AA Echt (NL).
- (74) **Agents:** **VROOM, DE, Erik** et al.; DSM Intellectual Property, P.O. Box 130, NL-6100 AC Echt (NL).

- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** PROCESS FOR THE PREPARATION OF STATINS IN THE PRESENCE OF BASE

(57) **Abstract:** The present invention relates to a process for the preparation of statins by means of a Julia-Kocienski reaction between an aldehyde and a sulfone derivative in the presence of an alkaline metal alkoxo base. The resulting derivatives are suitable as building blocks for statin type compounds such as cerivastatin, fluvastatin, pitavastatin and rosuvastatin.



WO 2012/098049 A1

PROCESS FOR THE PREPARATION OF STATINS IN THE PRESENCE OF BASE

Field of the invention

5

The present invention relates to a process for the preparation of statins by means of a Julia-Kocienski reaction between an aldehyde and a sulfone derivative in the presence of an alkaline metal alkoxy base. The resulting derivatives are suitable as building blocks for statin type compounds such as cerivastatin, fluvastatin, pitavastatin and rosuvastatin.

10

Background of the invention

15

Chiral diol sulfones are advanced intermediates used in preparing statins, a class of compounds useful as HMG CoA reductase inhibitors. In particular, chiral diol sulfones are employed in preparing statins in which an unsaturated carbon-carbon bond ($R_1\text{-CH=CH-R}_2$) is to be formed such as is the case in the antilipemic drugs cerivastatin, fluvastatin, pitavastatin and rosuvastatin.

20

A method for preparing chiral diol sulfones is described in WO 2002/098854 and WO 2001/096311. In these citations, a sulfone is prepared from an alcohol, more in particular *tert*-butyl 2-((4*R*,6*S*)-6-(hydroxymethyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate known as "Kaneka alcohol". The preparation of such an alcohol is described in EP 1024139.

25

Formation of unsaturated carbon-carbon bonds can be realized by the so-called Julia-Kocienski olefination (for a review see Aïssa (Eur. J. Org. Chem. 2009, 1831-1844)) between an aldehyde and the sulfones mentioned above in the presence of a base. Well-advocated bases in this respect are lithium hexamethyldisilazane (LiHMDS), potassium hexamethyldisilazane (KHMDS) and sodium hexamethyldisilazane (NaHMDS) as these bases are known for their versatility in controlling the E/Z-ratio of the products. A major drawback of these bases is that they can easily form unwanted side products which results in lower yields and difficult recovery and purification procedures.

30


It is an object of the present invention to provide a process wherein improved yields are obtained in comparison to the Julia-Kocienski olefination in the presence of KHMDS, LiHMDS or NaHMDS.

5

Detailed description of the invention


In a first aspect of the present invention, a compound of formula (1) or the corresponding lactone form (1') can be used as starting material

10



(1)

15




(1')

wherein X stands for halogen, like bromine or chlorine, preferably chlorine.


Prior to use in the process of the invention the hydroxyl groups and the carboxyl group of the above compounds may be protected as ketal and ester as outlined in general formula (1a) or as lactone and ether as outlined in general formula (1b)

20



(1a)

25





(1b)

wherein R_3 and R_4 each independently stand for an alkyl with for instance 1 to 12 C-atoms, preferably 1 to 6 C-atoms, an alkenyl with for instance 1 to 12 C-atoms, preferably 1 to 6 C-atoms, a cycloalkyl with for instance 3 to 7 C-atoms, a cycloalkenyl with for instance 3 to 7 C-atoms, an aryl with for instance 6 to 10 C-atoms or an aralkyl with for instance 7 to 12 C-atoms, each of R_3 and R_4 may be substituted and wherein R_3 and R_4 may form a ring together with the C-atom to which they are bound, use being

30



made of a suitable acetal forming agent, in the presence of an acid catalyst, for example as described in WO 2002/06266. The groups R_3 and R_4 are for example halogens or hydrocarbon groups with for instance 1 to 10 C-atoms, optionally containing one or more heteroatoms, for instance Si, N, P, O, S, F, Cl, Br or I. In practice, $R_3 = R_4$ is methyl is most preferred. In the compound of general formula (1a) R_5 is an alkyl or alkenyl group with one, two, three or four carbon atoms. Such relatively small substituents are generally known as being sterically unhindered or at least not very bulky. Suitable examples are allyl, *iso*-butenyl, *n*-butyl, *sec*-butyl, *tert*-butyl, ethyl, methyl, *n*-propyl, *iso*-propyl and vinyl. Preferably R_5 is a group that is easily introduced, small and easily removed under acidic conditions such as ethyl, methyl or *iso*-propyl. In the compound of general formula (1b) R_6 is hydrogen or an alcohol protecting group. Such a group can be any alcohol protecting group known to the skilled person such as described in, for example "Protective Groups in Organic Synthesis" (T.W. Greene, 1981, Wiley-Interscience Publication, ISBN 0-471-05764-9). These protecting groups are for example esters or ethers. These protection groups are preferred because in the final stage of conversion of these building blocks to statins, these generally acid labile or basic labile (in the case of esters) protection groups have the advantage to be removed either simultaneously with the opening of the lactone ring or by a pH shift. Hence, suitable groups R_6 are allyl, benzyloxymethyl, *tert*-butoxymethyl, *tert*-butyl, methoxymethyl, 1-ethoxyethyl, methoxyethoxymethyl, 4-methoxytetrahydropyranyl, methylthiomethyl, 1-(*iso*-propoxy)ethyl, tetrahydrofuranyl, tetrahydropyranyl, 2-methoxypropanyl, 1-propenyl, acetate, chloroacetate or benzoate. The compounds of formula (1) and (1'), and hence the compounds of formula (1a) and (1b) can be either enantiomerically pure or enriched in one of the enantiomers or racemic.

The compounds of formula (1a) or (1b) are reacted with a thiol compound of general formula R_7-S-X' (2) to give a compound of general formula (3a) or (3b), respectively with R_3 , R_4 , R_5 and R_6 as defined above.



In the compound of general formula (2), X' represents a proton or another cation like, for example, an alkali metal ion, like sodium or potassium or lithium cation, or an ammonium ion, like tetraalkylammonium, or a phosphonium ion, like tetraalkylphosphonium. R₇ is an aryl group that for instance is suitable for a one-pot or modified Julia-Kocienski olefination. The Julia-Kocienski olefination is a reaction in which an aldehyde of formula R₁-CH=O is reacted with a sulfone of formula R₂-CH₂-SO₂-R₇ to form an olefinic bond, *i.e.* R₁-CH=CH-R₂. The original Julia olefination requires two steps. In the modified reaction (Julia-Kocienski olefination), the intermediate obtained in the reaction with an aldehyde undergoes spontaneous elimination to give the olefin.

Preferably, the aryl group R₇ is a residue sufficiently π -electron deficient to be suitable for the modified (or one-pot) Julia-Kocienski olefination. In particular, it is preferred, that the aryl group is capable to promote a so-called Smiles rearrangement. Preferably, the thiol-aryl compound contains as an aryl group an aromatic moiety having a hetero atom, more preferably nitrogen. More in particular, the aromatic residue contains an electrophilic imine-like moiety within the heterocycle. Suitable aryl groups are *e.g.* described in P.R. Blakemore, J. Chem. Soc., Perkin Trans. 1, 2002, 2563. Particularly suitable in this respect are substituents R₇ of general formula R_{7'} and R_{7''}.

(R_{7'})(R_{7''})

wherein A-B is C=C or N=N or wherein A-B is part of a aromatic five- or six-membered ring, wherein Y is sulfur or alkyl-substituted nitrogen, preferably *tert*-butyl-substituted nitrogen, ethyl-substituted nitrogen, methyl-substituted nitrogen or phenyl-substituted nitrogen and wherein Z is nitrogen or CH. Preferred aryl groups include tetrazole, substituted phenyl and benzimidazole type compounds. Specific examples of preferred aryl groups include 3,5-bis(trifluoromethyl)phenyl-1-yl, pyridine-2-yl, pyrimidin-2-yl, benzothiazol-2-yl, 1-methyl-1*H*-tetrazol-5-yl, 1-phenyl-1*H*-tetrazol-5-yl, 1-*tert*-butyl-1-*H*-tetrazol-5-yl, 1-methylimidazol-2-yl, benzimidazol-2-yl,

4-methyl-1,2,4-triazol-3-yl and *iso*-quinolin-1-yl. Most preferred aryl groups are 3,5-bis(trifluoromethyl)phenyl-1-yl, 1-methyl-1*H*-tetrazol-5-yl, 1-phenyl-1*H*-tetrazol-5-yl, 1-*tert*-butyl-1-*H*-tetrazol-5-yl and benzothiazol-2-yl.

The reaction from (1a) or (1b) to (3a) and (3b), respectively will take place under
5 suitable conditions wherein it is generally important to preclude harsh conditions (like temperatures above 130°C or exceedingly long reaction times or application of strongly basic or acidic conditions) in order to preclude degradation of the starting compound or the thiol obtained. Suitable reaction conditions are temperatures about 50°C or higher, preferably about 80°C or higher, and more in particular about 100°C or higher.
10 Generally, the temperature will be about 150°C or lower, preferably about 140°C or lower, and more in particular about 130°C or lower. In case a temperature at the higher end-range is chosen, care should be taken to choose the time period such, that limited degradation occurs. Limited degradation is less than 10% of the starting halomethyl compound of general formula (1a) or (1b), preferably less than 5%, more in particular
15 less than about 3%. Generally, a reaction time of less than about 20 h, preferably less than about 10 h should be possible in case the reaction conditions are chosen properly. However, the time period is not critical, and may be up to 30 h or longer. Generally, the reaction takes longer than about 1 h, but this is strongly dependant on the reaction conditions, reaction engineering aspects (like reactor design or application of rate-
20 accelerating means like application of ultrasound or microwave irradiation) and amounts of reagents used, and this is not critical.

The reaction to obtain the thio-ether can be performed in a solvent or without the presence of a solvent. In case a solvent is used, the concentration of halomethyl starting compound of general formula (1a) or (1b) is generally about 10 wt% or higher,
25 preferably about 30 wt% or higher, more in particular about 40 wt% or higher. Preferably, the reaction is carried out with a relatively high concentration of halomethyl starting compound of general formula (1a) or (1b) of 70 to 99 wt%.

Suitable solvents are dimethylsulfoxide (DMSO), N-methyl pyrrolidone (NMP), dimethylformamide (DMF), sulfolane, acetonitrile, glymes (alkyl-capped or uncapped
30 mono-, oligo-, or poly-ethylene glycol ethers of varying chain length) or other polar non-protic solvents or alcohols like methanol, ethanol, 2-propanol, or halogenated hydrocarbons like dichloromethane, chloroform, 1,2-dichloroethane, optionally in combination with non-polar solvents like toluene or methyl *tert*-butyl ether (MTBE). It is

also possible to use biphasic solvent systems consisting of an aqueous phase and an organic phase in the presence of a phase-transfer catalyst, like quaternary ammonium salt or quarterly phosphonium salt (like tetraalkylammonium halide, e.g. tetrabutylammonium bromide) or crown ether (like 18-crown-6). These phase transfer-catalysts are also very suitable for use in mono-phasic solvent systems. Another class of suitable solvents comprises ionic liquids like, for example, 1,3-dialkyl imidazolium salts or *N*-alkyl pyridinium salts of acids like hexafluorophosphoric acid or tetrafluoroboric acid or trifluoromethane sulfonic acid, or with $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ as anionic counterpart.



The amounts of reagents can be chosen from a wide range. It is preferred to use a rate-enhancing excess of thiol compound, as the excess thiol that remains after reaction with the halomethyl derivative can be easily removed by washing with water at high pH. The molar amount of thiol to halogen compound generally is about 0.5 to 1 or higher, preferably 1 to 1 or higher, more preferably 1.1 to 1 or higher. Generally, the amount of thiol to halogen compound will be 3 to 1 or lower, preferably 2 to 1 or lower, most preferably 1.5 to 1 or lower. Preferably excess thiol is recovered for re-use which is easily achieved with the thiols of the present invention.

The thio-ether compound of general formula (3a) or (3b) can be isolated from the reaction mixture, or the mixture can be used as such in a subsequent oxidation reaction. Preferably, the reaction mixture is treated so as to remove excess thiol compound or excess halogen compound as the case may be. Any excess thiol compound can be easily removed by extraction with water at pH higher than 7, preferably higher than 8, more preferably of about 9 or higher. Suitable extraction agents are for example saturated caustic soda solution, saturated bicarbonate solution, or diluted sodium hydroxide solution. After extraction, the thio-ether compound of general formula (3a) or (3b) can be isolated by removal of the solvent by distillation, or by crystallization or precipitation, e.g. by addition of (or solvent switch to) an anti-solvent like hexane, heptane, *iso*-octane, or water. It is however not necessary to remove the solvent, as the oxidation can be performed in the same solvent. It is however preferred, to remove water from the reaction mixture, in case water interferes with the oxidation reaction. Hence, in a preferred embodiment of the invention, the oxidation is carried out without purification of the thio-ether compound of general

formula (3a) or (3b), more preferably in the same solvent as was used in the etherification reaction.

For the purpose of the present invention, a halogen derivative can be used as starting compound. This is advantageous because the Kaneka alcohol generally is prepared from such a halogen derivative. Therefore, the present invention provides a process, in which additional steps in the prior art are made obsolete if the chiral diol sulfone is to be used in a Julia-Kocienski olefination.

The thio-ether compound of general formula (3a) or (3b) is oxidized in manners known in the art, for example by oxidation with hydrogen peroxide or peracids (e.g. 3-chloroperoxybenzoic acid, peroxyacetic acid, monoperoxyphthalic acid), bleach, *tert*-BuOCl, perborates, *N*-oxides, permanganate, chromate, chlorate, bromate, perchlorate, periodate, *tert*-butyl hydroperoxide, oxone, peroxodisulfates and air/oxygen. The oxidation can be carried out in the presence of an appropriate catalyst, such as salts or oxides of the metals V, Ce, Mn, Ni, Fe, Cu, Os, Mo, W, Re, or Ru or organic catalysts like *iso*-butyraldehyde in the case of air/oxygen or tetramethylpiperidine *N*-oxide (TEMPO) in the case of bleach. The resulting sulfones are of general formula (4a) and (4b), respectively, with R₃, R₄, R₅, R₆ and R₇ as defined above.




The oxidation is performed in a solvent such as dichloromethane, chloroform, 1,2-dichloroethane, methanol, ethanol, 2-propanol, acetonitrile, acetic acid, toluene, water, NMP, DMSO, DMF, tetrahydrofuran (THF) or MTBE. Alternatively, biphasic solvent systems may be used. These may be an aqueous phase and an organic phase in the presence of a phase-transfer catalyst, like quaternary ammonium salt or quarterly phosphonium salt (like tetraalkylammonium halide, e.g. tetrabutylammonium bromide) or crown ether (like 18-crown-6). Other suitable solvents are ionic liquids like, 1,3-dialkyl imidazolium salts or *N*-alkyl pyridinium salts of acids like hexafluorophosphoric acid or

tetrafluoroboric acid or trifluoromethane sulfonic acid or with $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ as anionic counterpart. A reaction temperature of about -20°C or higher is effective.

For the oxidation, a temperature of about 0°C or higher is applied, more preferably a temperature close to ambient temperature ($18-25^\circ\text{C}$ *i.e.* around 20°C). A temperature of about 150°C or lower (preferably 100°C or lower, more preferably 60°C or lower, most preferably 40°C or lower) usually is effective. The molar amount of oxidant to thio-ether generally is about 1 to 1 or higher, preferably about 2 to 1 or higher, more preferably about 3 to 1 or higher. Generally, the amount of oxidant to thio-ether will be about 20 to 1 or lower, preferably about 10 to 1 or lower, most preferably about 5 to 1 or lower. The sulfone of general formula (4a) or (4b) can be isolated by aqueous extraction of excess oxidant/catalyst and subsequent removal of the solvent by evaporation. If water-miscible solvents like alcohols or aprotic polar solvents are applied as reaction medium, the reaction mixture can be partitioned between an aqueous and an organic phase prior to this operation, in order to extract the solvent to the aqueous phase. If ionic liquids are applied as reaction medium, the sulfone can be isolated by extraction with an organic solvent immiscible with the ionic liquid, followed by evaporation of the solvent. Alternatively, the sulfone can be isolated from the reaction mixture by precipitation or crystallization, *e.g.* by addition of (or solvent switch to) an anti-solvent like hexane, heptane, *iso*-octane, or water. If desired, purification of the sulfone can be performed by chromatography or, preferably, by re-crystallization from (or trituration with) a suitable solvent, like 2-propanol or another solvent, depending on the aryl group used with the thiol compound of formula (2) and the residues R_3 , R_4 , R_5 and R_6 used with the initial halomethyl compounds of formula (1a) or (1b).

In the present invention, the sulfone of general formula (4a) or (4b) is treated with an aldehyde $\text{R}_1\text{-CH=O}$, in which R_1 is chosen so as to obtain suitable precursors to useful statin-type compounds including pitavastatin, rosuvastatin, fluvastatin, and cerivastatin, or in which R_1 is a suitable precursor to these moieties (*cf.* WO 2002/098854 and WO 2001/096311). Preferred examples of aldehyde $\text{R}_1\text{-CH=O}$ are 4-(4-fluorophenyl)-2,6-diisopropyl-5-(methoxymethyl)nicotinaldehyde, 3-(4-fluorophenyl)-1-isopropyl-1H-indole-2-carbaldehyde, 2-cyclopropyl-4-(4-fluorophenyl)-quinoline-3-carbaldehyde and N-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-N-methylmethanesulfonamide as these aldehydes are the precursors for cerivastatin,

fluvastatin, pitavastatin and rosuvastatin, respectively. Thus, R_1 preferably is a radical chosen from the list consisting of the radicals of formula (A), (B), (C) and (D):




The above Julia-Kocienski olefination between compounds (4a) or (4b) and aldehyde $R_1\text{-CH=O}$ is carried out in the presence of a base, examples of which are lithium hydride, potassium hydride, sodium hydride, KHMDS, LiHMDS, NaHMDS, solid potassium hydroxide, solid sodium hydroxide, alkaline metal alkoxy bases, lithium bis-trimethylsilylamide ($\text{LiN}(\text{TMS})_2$), sodium bis-trimethylsilylamide ($\text{NaN}(\text{TMS})_2$), potassium bis-trimethylsilylamide ($\text{KN}(\text{TMS})_2$), sodium amide, P4-tBu, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) and the like. Whereas the E/Z-ratio in the final product depends on various parameters, such as type of base, thio-substituent (R_7) and solvents, as for instance outlined in P.R. Blakemore *et al.* (Synlett 1998, 26-28), this ratio normally varies between 40:60 and 80:20 in customary solvents such as diethyl ether, dimethoxyethane, tetrahydrofuran and toluene.

Surprisingly it was found that in the condensation reaction of the present invention, *i.e.* starting from compounds of general formula (4a) and (4b) and aldehydes $R_1\text{-CH=O}$, a marked improvement is achieved when using bases having a pKa value between 15 and 22, preferably between 16 and 21 and more preferably between 17 and 20. When using bases having a higher pKa value, such as described in the prior art (KHMDS, LiHMDS or NaHMDS) in the synthesis of rosuvastatin, wherein R_1 is a radical of formula (D), significant amounts of side products are found. For example, de-protonation of the methyl group α to the sulfone moiety in (D) leads to 15-20% of unwanted impurities. Surprisingly, when the bases of the present invention are used, these amounts are only 1-2%, under otherwise same conditions. Particularly suitable examples of bases in this respect are alkaline metal alkoxy bases such as lithium *tert*-


butoxide, lithium ethoxide, lithium methoxide, potassium *tert*-butoxide, potassium ethoxide, potassium methoxide, sodium *tert*-butoxide, sodium ethoxide and sodium methoxide. These alkaline metal alkoxy bases have significant additional advantages as, compared to the prior art bases, they are less sensitive towards traces of water, can be used in solvents that are difficult to dry (like *tert*-butanol), are easier to handle, are less expensive and lead to smaller and less toxic waste streams as is the case with the silicon-comprising bases such as KHMDS, LiHMDS and NaHMDS.

In addition a marked difference is observed between lithium-comprising bases and sodium-comprising bases where the latter strongly favors extremely high E/Z-ratio's. This is advantageous as the E-configuration is the required configuration in cerivastatin, fluvastatin, pitavastatin and rosuvastatin. Hence, the use of a sodium-comprising base precludes laborious removal and/or recycling of undesired Z-isomer.

Following the Julia-Kocienski olefination between compounds (4a) or (4b) and aldehyde $R_1-CH=O$ the resulting products (5a) and (5b), respectively may be isolated and purified after which they are deprotected to give product (6). Alternatively deprotection may be carried out without isolation and/or purification of intermediate products (5a) and (5b). Deprotection is carried out according to procedures known to the skilled person, for instance by using acid such as hydrochloric acid as described in US 6,844,437 or WO 2007/000121.



In a second aspect of the invention there is disclosed a compound of general formula (5a) wherein R₁ is a radical chosen from the list consisting of the radicals of formula (A), (B), (C) and (D)



and wherein R_3 and R_4 each independently stand for an alkyl with for instance 1
 10 to 12 C-atoms, preferably 1 to 6 C-atoms, an alkenyl with for instance 1 to 12 C-atoms,
 preferably 1 to 6 C-atoms, a cycloalkyl with for instance 3 to 7 C-atoms, a cycloalkenyl
 with for instance 3 to 7 C-atoms, an aryl with for instance 6 to 10 C-atoms or an aralkyl
 with for instance 7 to 12 C-atoms, each of R_3 and R_4 may be substituted and wherein
 R_3 and R_4 may form a ring together with the C-atom to which they are bound, use being
 15 made of a suitable acetal forming agent, in the presence of an acid catalyst, for
 example as described in WO 2002/06266. The groups R_3 and R_4 are for example
 halogens or hydrocarbon groups with for instance 1 to 10 C-atoms, optionally
 containing one or more heteroatoms, for instance Si, N, P, O, S, F, Cl, Br or I.
 Preferably $R_3 = R_4$ is methyl. In the compound of general formula (1a) R_5 is an alkyl or
 20 alkenyl group with one, two, three, four, five or six carbon atoms. Novel and preferred
 examples are those wherein R_5 is an alkyl with four, five or six carbon atoms. It was
 found that, apart from compounds wherein R_5 is an alkyl with one, two or three carbon
 atoms, also the compounds with four, five or six carbon atoms are suitable from a
 preparative and economic point of view. Particular suitable examples in this respect are
 25 *iso*-butenyl, *n*-butyl, *sec*-butyl, *tert*-butyl, 3-methylbutan-2-yl, 4-methylpentan-2-yl and
 2-pentyl.


EXAMPLES

30

Example 1

Preparation of 2-(4*R*,6*S*)-6-((benzo[d]thiazole-2-ylthio)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid *sec*-butyl ester (B) from (4*R*,6*S*)-6-(chloromethyl)-2,2-

dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (A) and 2-mercapto-1H-benzothiazole (2-MBT)



5


A reactor was charged with 297.5 g (1.07 mol) of compound A and 1160 mL of N-methylpyrrolidone. To this solution was added 214 g 2-MBT (1.2 eq. 1.28 mol), 117 g NaHCO₃ (1.3 eq., 1.39 mol) and 0.69 g tetra-n-butyl ammonium bromide (TBAB). The reaction mixture was stirred for 23 h at 90°C and then cooled to RT, diluted with 1.5 L of methyl-*tert*-butyl ether and washed with 1 L of saturated aqueous NaHCO₃. The organic layer was washed again with 2 x 700 mL of saturated aqueous NaHCO₃ and 2 x 700 mL of water. The resulting brown organic layer was treated with 20 g active carbon. After removal of the carbon by filtration, the organic layer was concentrated to give 2-(4*R*,6*S*)-6-((benzo[d]thiazole-2-ylthio)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (compound B), as a yellow oil. Yield 301.8 g (69%). ¹H NMR (300 MHz, CDCl₃) δ 7.91 (d, 1H), 7.76 (d, 1H), 7.49-7.39 (m, 1H), 7.36-7.28 (m, 1H), 4.96-4.77 (m, 1H), 4.40-4.21 (m, 2H), 3.53 (add, 2H), 2.46 (ddd, 2H), 1.84 (dt, 1H), 1.65-1.49 (m, 2H), 1.46 (s, 3H), 1.39 (s, 3H), 1.36-1.22 (m, 1H), 1.20 (dd, 3H), 0.89 (m, 3H).

20

Example 2

Preparation of 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (C) from 2-(4*R*,6*S*)-6-((benzo[d]thiazole-2-ylthio)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (B)

25



A) Procedure using *m*-CPBA

2-((4*R*,6*S*)-6-((benzo[d]thiazole-2-ylthio)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (B, 2.48 g, 6.0 mmol) and 1.9 g NaHCO₃ (22.8 mmol; 3.8 equiv.) were added to 50 mL of dichloromethane at 0°C. Then 4.1 g *m*-CPBA (16.8 mmol; 2.8 equiv) was added slowly keeping the temperature below 5°C (slightly exothermic). When addition was completed, the temperature was allowed to raise to 20-25°C and stirred for 18 h. The mixture was diluted with 50 mL of dichloromethane and the precipitated salts were removed by filtration. The organic layers was washed with 2 x 100 mL of 10 w/w% aqueous NaHSO₃ and 4 x 100 mL of 5 w/w% aqueous Na₂CO₃. After drying on Na₂SO₄, the dichloromethane was evaporated. The resulting thick oil was stirred in isopropanol to give compound 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester as a white solid (C, 0.54 g, yield 20%). ¹H NMR (300 MHz, CDCl₃) δ 8.22 (d, 1H), 8.01 (d, 1H), 7.70-7.51 (m, 2H), 4.85 (m, 1H), 4.59 (ddt, 1H), 4.40-4.24 (m, 1H), 3.87 (dd, 1H), 3.42 (dd, 1H), 2.45-2.35 (m, 2H), 1.71 (d, 1H), 1.43 (d, 2H), 1.36-1.33 (m, 4H), 1.18 (dd, 3H), 0.87 (dt, 3H), 0.78 (s, 3H).

B) Procedure using Na₂WO₄•2H₂O and 30% H₂O₂

2-((4*R*,6*S*)-6-((Benzo[d]thiazole-2-ylthio)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester (B, 4 g, 9.8 mmol) was dissolved in 20 mL of methanol and Na₂WO₄•2H₂O (0.3 g, 10 mol%) was added. Next, 3 mL of 30% H₂O₂ solution was added in 1 h at 25°C while keeping the pH at about 8.5 using aqueous 4M NaOH. After the addition was completed, the mixture was stirred for 18 h. The precipitated solid was isolated by filtration. The solid was dissolved in ethylacetate and washed 2 times with 10 w/w% aqueous Na₂SO₃. The organic layer was evaporated and the resulting solid was re-crystallized from isopropanol to give 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid sec-butyl ester as a white solid (2.46 g, yield 57%). NMR identical as given under procedure A using *m*-CPBA.


C) Procedure using Na₂WO₄•2H₂O and 30% H₂O₂ (270 g scale)

A reactor was filled with 270 g (0.66 mol) of compound B, 21.7 g Na₂WO₄•2H₂O (10 mol%) and 1940 mL of methanol. To this mixture was added in 2 h, 222 mL of a 30% H₂O₂ solution (3.3 eq.), keeping the temperature at 20°C and the pH at 8 using

aqueous 4M NaOH. When the addition was completed, the reaction mixture was stirred for 23 h. The precipitated product was filtered and washed with methanol (100 mL). To the filtrate was added water (500 mL) to precipitate more solid. The product was filtered. The combined solids were dried to give 164 g of crude product. The crude material was re-crystallized from isopropanol, to give 125 g of 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetic acid *sec*-butyl ester (C). The filtrate was concentrated to about 150 mL to give another 19 g of compound C. In total 144 g of compound C (yield = 41%) was obtained.

Example 3

Preparation of 2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methylmethylsulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate, *sec*-butyl ester (E) from *N*-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-*N*-methylmethanesulfonamide (D) and 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate *sec*-butyl ester (C)



A) Procedure using KO-*t*Bu

2-((4*R*,6*S*)-6-((Benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate *sec*-butyl ester (C, 8.0 g 18.1 mmol) and *N*-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-*N*-methylmethanesulfonamide (D, 6.4 g, 18.1 mmol) were added to 110 mL of tetrahydrofuran. The mixture was heated until complete dissolution was obtained. The solution was cooled to -50°C, followed by addition of 19.9 mL of 1 M

KO-*t*Bu in tetrahydrofuran (19.9 mmol) in 1 h. The temperature was allowed to increase to -10°C and quenched with 100 mL of 10 w/w% of aqueous NH₄Cl. HPLC analysis indicated 77% product formation. The layers were separated and the organic phase washed again with 10 w/w% aqueous NH₄Cl. Then the reaction mixture was extracted three times with brine at a pH of 12-13 (adjusted with 4 M aqueous NaOH). After drying over Na₂SO₄, the organic layer was evaporated to give an oil, which slowly solidified. The solid was re-crystallized from isopropanol to give 2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methanesulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate, *sec*-butyl ester (compound E) as a white solid (5.0 g, yield 48%). ¹H NMR (300 MHz, CDCl₃) δ 7.58 (dd, 2H), 7.01 (t, 2H), 6.45 (dd, 1H), 5.40 (dd, 1H), 4.81, (m, 1H), 4.44-4.18 (m, 2H), 3.54-3.47 (m, 3H), 3.47-3.38 (m, 3H), 3.35-3.25 (m, 1H), 2.37 (ddd, 2H), 1.59-1.43 (m, 4H), 1.41 (s, 3H), 1.33 (s, 3H), 1.20, (dd, 6 H), 1.14 (d, 3H), 0.83 (t, 3H).

B) Procedure using NaO-*t*Bu

2-((4*R*,6*S*)-6-((Benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate *sec*-butyl ester (C, 2.0 g 4.53 mmol) and *N*-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-*N*-methanesulfonamide (D, 1.43 g, 4.08 mmol) were added to 26 mL of tetrahydrofuran. The mixture was heated until complete dissolution was obtained. The solution was cooled to -50°C, followed by addition of 6.8 mL of 1 M NaO-*t*Bu in tetrahydrofuran (6.8 mmol, 1.5 eq) in 1 h. The temperature was allowed to increase to -10°C and quenched with 100 mL of 10 w/w% of aqueous NH₄Cl. HPLC analysis indicated 90% product formation. The product can be isolated as described in the procedure using KO-*t*Bu.

C) Comparative procedure using NaHMDS

N-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-*N*-methanesulfonamide (5.7 g, 16.3 mmol) and 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate *sec*-butyl ester (8.0 g, 18.1 mmol) were added to 104 mL of tetrahydrofuran. The reaction mixture was heated until all the reagents were dissolved and then cooled to -70°C. At this temperature 27.2 mL of a NaHMDS solution (20% in tetrahydrofuran, total 27.2 mmol, 1.5 eq.) was added in 1 h at -70°C. When dosing was completed, the reaction mixture was stirred for 1 h at -70°C. HPLC analysis showed

68% of product formed. The reaction mixture was quenched with 100 mL of 10% aqueous NH₄Cl, the aqueous phase separated and the organic phase washed 2 times with 100 mL of 10% aqueous NH₄Cl. Next, the organic phase was washed 3 times with water at pH 12 (using 1M aqueous NaOH). The organic phase was evaporated and the residue was crystallized from isopropanol to give 2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methylmethylsulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate, *sec*-butyl ester as a solid. ¹H NMR (300 MHz, CDCl₃) δ 7.58 (dd, 2H), 7.01 (t, 2H), 6.45 (dd, 1H), 5.40 (dd, 1H), 4.81, (m, 1H), 4.44-4.18 (m, 2H), 3.54-3.47 (m, 3H), 3.47-3.38 (m, 3H), 3.35-3.25 (m, 1H), 2.37 (ddd, 2H), 1.59-1.43 (m, 4H), 1.41 (s, 3H), 1.33 (s, 3H), 1.20, (dd, 6 H), 1.14 (d, 3H), 0.83 (t, 3H).

Tentative Example 3

Preparation of 2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methylmethylsulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate 4-methylpentan-2-yl ester from *N*-(4-(4-fluorophenyl)-5-formyl-6-isopropylpyrimidin-2-yl)-*N*-methylmethanesulfonamide (D) and 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate 4-methylpentan-2-yl ester

The 4-methylpentan-2-yl ester of compound (5a) with R₁ is a radical of formula (D), R₃ and R₄ are both methyl and R₅ is 4-methylpentan-2-yl can be prepared according to the same procedures as outlined in Example 3 starting from 2-((4*R*,6*S*)-6-((benzo[d]thiazol-2-ylsulfonyl)methyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate 4-methylpentan-2-yl ester instead of the corresponding *sec*-butyl ester using the same molar amount.


For reference purposes 2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methylmethylsulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate 4-methylpentan-2-yl ester was prepared from rosuvastatin methyl ester (EP 521471). Thus, rosuvastatin methyl ester (30 g, 61 mmol) was added to 200 mL of acetonitrile and 2N aqueous NaOH was added until the pH was stable at 12.5. The reaction mixture was stirred for 2 h at 20°C. Then the pH was lowered to 5.0 with 2N aqueous HCl. To the reaction mixture was added 200 mL of ethyl acetate and the organic phase was separated and washed 2 times with 100 mL of water. The ethyl acetate phase was dried over Na₂SO₄, filtered and evaporated to a sirup (≈ 30 g of rosuvastatin acid). Part of this sirup (20 g) was dissolved in toluene and heated to reflux under azeotropic water

removal for 4 h. The reaction mixture was cooled to 20°C and stirred for 18h. The precipitated solid was filtered, washed with toluene (2 x 10 mL) and dried to give 16.1 g (35 mmol) of N-(4-(4-fluorophenyl)-5-((E)-2-((2*S*,4*R*)-4-hydroxy-6-oxotetrahydro-2H-pyran-2-yl)vinyl)-6-isopropylpyrimidin-2-yl)-N-methylmethanesulfonamide as a white solid. ¹H NMR (300 MHz, CDCl₃) δ 7.62 (dd, 2H), 7.11 (dd, 2H), 6.72 (dd, 1H), 5.48 (dd, 1H), 5.28-5.20 (m, 1H), 4.38-4.30 (m, 1H), 3.58 (s, 3H), 3.52 (s, 3H), 3.38-3.30 (m, 1H), 2.80-2.60 (m, 2H), 2.10-2.00 (m, 1H), 1.95-1.85 (m, 1H), 1.73-1.68 (m, 1H), 1.28 (d, 3H), 1.26 (d, 3H).

Of this compound, 2.3 g (5.0 mmol) was added to 25 mL of 2-(4-methyl)-pentanol. Then 2 drops of methanesulphonic acid were added and the reaction mixture was heated to 60°C and stirred for 1h. The reaction mixture was cooled to 20°C and stirred for 18h. Next 2,2-dimethoxypropane (0.78 g, 7.5 mol) was added and the mixture was stirred for 2h. The reaction mixture was quenched with 20 mL of saturated aqueous NaHCO₃ followed by addition of 25 mL of ethyl acetate. The organic phase was separated and washed 2 times with 20 mL of saturated aqueous NaHCO₃. The organic phase was evaporated and the residue slowly solidified to give the title compound as a solid (2.9 g, 96% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.58 (dd, 2H), 7.01 (t, 2H), 6.46 (dd, 1H), 5.42 (dd, 1H), 4.87 (m, 1H), 4.43 - 4.20 (m, 2H), 3.50 (s, 3H), 3.45 (s, 3H), 3.35 - 3.25 (m, 1H), 2.38 (ddd, 2H), 1.59 - 1.43 (m, 4H), 1.40 (s, 3H), 1.35 (s, 3H), 1.20 (dd, 4H), 1.14 (d, 6H), 0.83 (t, 6H).

Example 4


Preparation of rosuvastatin-Ca (F) from-((4*R*,6*S*)-6-((E)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methylmethanesulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate, sec-butyl ester (E)




2-((4*R*,6*S*)-6-((*E*)-2-(4-(4-fluorophenyl)-6-isopropyl-2-(*N*-methylethylsulfonamido)pyrimidin-5-yl)vinyl)-2,2-dimethyl-1,3-dioxan-4-yl)acetate, *sec*-butyl ester (compound E, 2.0 g, 3.46 mmol) was added to 40 mL of methanol. The mixture was heated to 35°C until complete dissolution was obtained. The solution was cooled to 20°C and 9 mL of 0.2 N HCl was added over a period of 2 h. The mixture was stirred for 18 h, followed by addition of 1 N NaOH in 15 min. After stirring for 1 h, 0.2 g of dicalite was added and the mixture filtered. The solution was concentrated to about 15 g, 10 mL of water was added and the mixture again concentrated to 15 g. Then 10 mL of water was added. To the obtained clear solution was added in portions over a period of 1 h, 7 mL of a solution of 4.5 w/w% Ca(OAc)₂•H₂O (1.2 equiv.) in water. Upon addition white precipitate was formed. After 1 h the precipitate was filtered and dried to give 1.34 g (2.68 mmol) of the calcium salt of rosuvastatin as a slightly yellow solid (yield 77%). ¹H NMR (300 MHz, DMSO) δ 7.72 (dd, 2H), 7.29 (t, 2H), 6.51 (d, 1H), 5.54 (dd, 1H), 4.21 (dd, 1H), 3.71 (m, 1H), 3.55 (s, 3H), 3.51-3.41 (m, 4H), 2.09 (dd, 1H), 1.92 (dd, 1H), 1.57-1.42 (m, 1H), 1.36-1.25 (m, 1H), 1.22 (dd, 6H).

CLAIMS

1. Process for the preparation of a compound of general formula $R_1-C=C-R_2$ wherein R_1 is a radical chosen from the list consisting of the radicals of formula (A), (B), (C) and (D)





and wherein R_2 is a radical of general formula (E) or (F)



wherein R_3 and R_4 each independently stand for an alkyl with 1 to 12 carbon atoms or an alkenyl with 1 to 12 carbon atoms or a cycloalkyl with 3 to 7 carbon atoms or a cycloalkenyl with 3 to 7 carbon atoms or an aryl with 6 to 10 carbon atoms or an aralkyl with 7 to 12 carbon atoms or wherein R_3 and R_4 form a ring together with the carbon atom to which they are bound, wherein R_5 is an alkyl or alkenyl group with 1 to 4 carbon atoms and wherein R_6 is an alcohol protecting group,

comprising reacting a compound of formula $R_1-CH=O$ with a compound of formula $R_2-CH_2-SO_2-R_7$ wherein R_1 and R_2 are as outlined above and R_7 is a radical of general formula R_7' or R_7''



wherein A-B is C=C or N=N or wherein A-B is part of a aromatic five- or six-membered ring, optionally substituted, wherein Y is sulfur or alkyl-substituted nitrogen and wherein Z is nitrogen or CH,
in the presence of a base characterized in that said base has a pKa value of
5 from 16 to 21.


2. Process according to claim 1 wherein said base is an alkaline metal alkoxy base.

10 3. Process according to claim 2 wherein said base is chosen from the list consisting of lithium *tert*-butoxide, lithium ethoxide, lithium methoxide, potassium *tert*-butoxide, potassium ethoxide, potassium methoxide, sodium *tert*-butoxide, sodium ethoxide and sodium methoxide.

15 4. Process according to claim 3 wherein R₃ is ethyl or methyl and R₄ is ethyl or methyl or R₃ and R₄ form a cyclopentyl ring or a cyclohexyl ring together with the carbon atom to which they are bound, R₅ is *sec*-butyl, *tert*-butyl, ethyl, methyl, or *iso*-propyl, R₆ is methoxyethoxymethyl, tetrahydrofuranyl or tetrahydropyranyl and R₇ is 1-methyl-1*H*-tetrazol-5-yl, 1-phenyl-1*H*-tetrazol-5-yl, 1-*tert*-butyl-1-*H*-
20 tetrazol-5-yl, benzothiazol-2-yl or 3,5-bis(trifluoromethyl)phenyl-1-yl.


5. Process according to claim 4 wherein R₁ is a radical of formula (D).

25 6. Process according to any one of claims 1 to 5 followed by deprotection and isolation to give a compound of general formula R₁-C=C-R₂ or a salt thereof wherein R₁ is as outlined above and R₂ is a radical of general formula (G)




30

7. Process according to claim 6 wherein said compound obtained after deprotection and isolation is rosuvastatin or the calcium salt of rosuvastatin.



8. A compound of general formula (5a)
 wherein R_1 is a radical chosen from the list consisting of the radicals of formula
 (A), (B), (C) and (D)



wherein R_3 and R_4 each independently stand for an alkyl with 1 to 12 carbon atoms or an alkenyl with 1 to 12 carbon atoms or a cycloalkyl with 3 to 7 carbon atoms or a cycloalkenyl with 3 to 7 carbon atoms or an aryl with 6 to 10 carbon atoms or an aralkyl with 7 to 12 carbon atoms or wherein R_3 and R_4 form a ring together with the carbon atom to which they are bound,
 characterized in that R_5 is an alkyl or alkenyl group with 4 to 6 carbon atoms.

9. Compound according to claim 8 wherein R_1 is a radical of formula (D), R_3 and R_4 are methyl and R_5 is chosen from the list consisting of *iso*-butenyl, *n*-butyl, *sec*-butyl, *tert*-butyl, 3-methylbutan-2-yl, 4-methylpentan-2-yl and 2-pentyl.
10. Compound according to claim 9 wherein R_5 is *sec*-butyl.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/050470

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D413/12 C07D419/12
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EP0-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02/098854 A2 (SQUIBB BRISTOL MYERS CO [US]; BRODFUEHRER PAUL R [US]; SATTELBERG THOM) 12 December 2002 (2002-12-12) cited in the application scheme 1, reaction 7 to 8; scheme 2, reaction 7a to 6a; claim 16; example 3 -----	1-7
A	WO 01/96311 A2 (SQUIBB BRISTOL MYERS CO [US]; ROBL JEFFREY A [US]; CHEN BANG CHI [US];) 20 December 2001 (2001-12-20) cited in the application scheme 6, reaction 17 to 18; -----	1-7
X	WO 2010/140765 A2 (CHONG KUN DANG PHARM CORP [KR]; LEE HONGWOO [KR]; PARK DAEJONG [KR]; Y) 9 December 2010 (2010-12-09) page 4, compound [3] ----- -/--	8-10



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 March 2012

Date of mailing of the international search report

03/04/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Schmid, Arnold

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/050470

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/077062 A2 (HANMI PHARM IND CO LTD [KR]; KIM HEE SEOCK [KR]; KIM WON JEOUNG [KR];) 8 July 2010 (2010-07-08) claims; examples 5,8 -----	8-10
X	WO 2005/042522 A1 (ASTRAZENECA UK LTD [GB]; SHIONOGI & CO [JP]; OKADA TETSUO [JP]; HORBUR) 12 May 2005 (2005-05-12) compound 4; compounds 7a; page 19, ines 15-17; intermediate of preparations table bridging pages 19, 20 -----	8-10
X	BAN-CHI CHEN ET AL: "Synthesis of d3-cerivastatin for use as internal standard in a LC/MS/MS method developed for quantitation of the drug in human serum", JOURNAL OF LABELLED COMPOUNDS AND RADIOPHARMACEUTICALS, vol. 49, 2006, pages 311-319, XP002671557, page 313, compound 13 -----	8-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/050470

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 02098854	A2	12-12-2002	BR 0209942 A 30-03-2004
		CA 2449813 A1 12-12-2002	
		CN 1656077 A 17-08-2005	
		CZ 20033329 A3 10-11-2004	
		DE 02737324 T1 15-07-2004	
		EP 1392656 A2 03-03-2004	
		ES 2214169 T1 16-09-2004	
		HU 0401724 A2 28-12-2004	
		JP 2004536813 A 09-12-2004	
		MX PA03011195 A 18-03-2004	
		PL 367133 A1 21-02-2005	
		TR 200400600 T3 21-06-2004	
		TW 1256391 B 11-06-2006	
		WO 02098854 A2 12-12-2002	
-----	-----	-----	-----
WO 0196311	A2	20-12-2001	AR 028732 A1 21-05-2003
		AU 6686001 A 24-12-2001	
		BR 0111571 A 01-07-2003	
		CA 2412979 A1 20-12-2001	
		CN 1436174 A 13-08-2003	
		CZ 20023931 A3 12-03-2003	
		EC SP024386 A 10-03-2003	
		EP 1294696 A2 26-03-2003	
		HU 0302955 A2 29-12-2003	
		JP 2004503541 A 05-02-2004	
		MX PA02012415 A 06-06-2003	
		NO 20026011 A 12-02-2003	
		PE 13642001 A1 27-02-2002	
		PL 362477 A1 02-11-2004	
		UY 26776 A1 31-01-2002	
		WO 0196311 A2 20-12-2001	
		ZA 200210102 A 12-03-2004	
-----	-----	-----	-----
WO 2010140765	A2	09-12-2010	EP 2439198 A2 11-04-2012
		KR 20100131245 A 15-12-2010	
		WO 2010140765 A2 09-12-2010	
-----	-----	-----	-----
WO 2010077062	A2	08-07-2010	KR 20100080432 A 08-07-2010
		WO 2010077062 A2 08-07-2010	
-----	-----	-----	-----
WO 2005042522	A1	12-05-2005	AR 046191 A1 30-11-2005
		AU 2004285750 A1 12-05-2005	
		BR PI0415681 A 19-12-2006	
		CA 2543358 A1 12-05-2005	
		CN 1898233 A 17-01-2007	
		CN 101654453 A 24-02-2010	
		EP 1682536 A1 26-07-2006	
		EP 2272842 A1 12-01-2011	
		IS 8478 A 22-05-2006	
		JP 2007509119 A 12-04-2007	
		JP 2008024712 A 07-02-2008	
		JP 2008044948 A 28-02-2008	
		KR 20060132829 A 22-12-2006	
		NZ 547094 A 31-05-2009	
		RU 2372349 C2 10-11-2009	
		TW 1344957 B 11-07-2011	
		US 2007255060 A1 01-11-2007	
		US 2011301348 A1 08-12-2011	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/050470

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		UY 28577 A1	30-06-2005
		WO 2005042522 A1	12-05-2005
		ZA 200603173 A	25-07-2007
