



US00PP19620P2

(12) **United States Plant Patent**
Kerley et al.

(10) **Patent No.:** **US PP19,620 P2**

(45) **Date of Patent:** **Jan. 6, 2009**

(54) **PETUNIA×HYBRIDA PLANT NAMED**
‘KERWHIFAN’

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: *Petunia×hybrida*
Varietal Denomination: **Kerwhifan**

(52) **U.S. Cl.** **Plt./356**

(58) **Field of Classification Search** **Plt./356**
See application file for complete search history.

(76) Inventors: **David W. Kerley**, Bethany, 49 Station
Road, Over, Cambridge CB24 5NJ (GB);
Priscilla G. Kerley, Bethany, 49 Station
Road, Over, Cambridge CB24 5NJ (GB)

Primary Examiner—Howard J Locker
(74) *Attorney, Agent, or Firm*—Audrey Charles

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 192 days.

(57) **ABSTRACT**

A new and distinct cultivar of *Petunia* plant named
‘Kerwhifan’, characterized by its white-colored flowers,
medium green-colored foliage, moderately vigorous,
spreading, and semi-trailing growth habit.

(21) Appl. No.: **11/656,621**

(22) Filed: **Jan. 23, 2007**

1 Drawing Sheet

1

2

Latin name of genus and species of plant claimed: *Petunia×hybrida*.
Variety denomination: ‘Kerwhifan’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Petunia* plant botanically known as *Petunia×hybrida* and
hereinafter referred to by the cultivar name ‘Kerwhifan’.

The new cultivar originated in a controlled breeding pro-
gram in Cambridge, England during August 2002. The
objective of the breeding program was the development of
early and large flowered *Petunia* varieties.

The new *Petunia* cultivar is the result of cross-pollination.
The female (seed) parent of the new cultivar is the prop-
rietary *Petunia×hybrida* breeding selection designated
02-114-6, not patented, characterized by its white-colored
flowers, darkgreen-colored foliage, upright growth habit.
The male (pollen) parent of the new cultivar is the prop-
rietary *Petunia×hybrida* breeding selection designated
02-109-5, not patented, characterized by its white-colored
flowers, medium green-colored foliage, and upright growth
habit. The new cultivar was discovered and selected as a
single flowering plant within the progeny of the above stated
cross-pollination during May 2003 in a controlled environ-
ment at Cambridge, England.

Asexual reproduction of the new cultivar by terminal stem
cuttings since September 2003 at Cambridge, England has
demonstrated that the new cultivar reproduces true to type
with all of the characteristics, as herein described, firmly
fixed and retained through successive generations of such
asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have
been repeatedly observed and can be used to distinguish
‘Kerwhifan’ as a new and distinct cultivar of *Petunia* plant:

1. White-colored flowers;
2. Medium green-colored foliage; and
3. Moderately vigorous, spreading, and semi-trailing
growth habit.

Plants of the new cultivar differ from plants of the female
parent primarily in growth habit and from plants of the male
parent primarily in growth habit.

Of the many commercially available *Petunia* cultivars
known to the inventor, the most similar in comparison to the
new cultivar is ‘Kesupite’, not patented. However, in side by
side comparisons, plants of the new cultivar differ from
plants of ‘Kesupite’ in the following characteristics:

1. Plants of the new cultivar have larger flowers than
plants of ‘Kesupite’; and
2. Plants of the new cultivar have flower throats with
green-colored venation, whereas plants of the cultivar
‘Kesupite’ have flower throats with purple-colored
venation.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. Colors in the photographs differ slightly from
the color values cited in the detailed description, which
accurately describes the colors of ‘Kerwhifan’. The plants
were grown in 8-inch hanging pots with three plants per pot
for about 12 weeks in a greenhouse at Cambridge, England.

FIG. 1 illustrates a side view of the overall growth and
flowering habit of ‘Kerwhifan’.

FIG. 2 illustrates a close-up view of an individual flower
of ‘Kerwhifan’.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible
environmental conditions to date. Accordingly, it is possible
that the phenotype may vary somewhat with variations in the
environment, such as temperature, light intensity, and day
length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 1995 edition, except where general color terms of ordinary significance are used. The color values were determined on Sep. 21, 2006 between 10.00 a.m. and 12.00 p.m. under natural light conditions in Cambridge, England.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at Cambridge, England in 8-inch hanging pots for 12 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 64° F. to 95° F. (18° C. to 35° C.) during the day and approximately 57° F. to 68° F. (14° C. to 20° C.) during the night. Greenhouse light levels of 3,200 footcandles to 6,000 footcandles were maintained during the day.

Botanical classification: *Petunia×hybrida* cultivar Kerwhifan.

Parentage:

Female parent.—Proprietary *Petunia×hybrida* breeding selection designated 02-114-6.

Male parent.—Proprietary *Petunia×hybrida* breeding selection designated 02-109-5.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 10 to 12 days.

Time to produce a rooted cutting.—Approximately 28 to 35 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 5 to 7 weeks from a rooted cutting.

Growth habit and general appearance.—Spreading, semi-trailing.

Size.—Height from soil level to top of plant plane: Approximately 20.0 cm. Width: Approximately 55.0 cm.

Branching habit.—Freely branching; however, one pinch is required for commercial product. Quantity of main branches per plant: Approximately 4 to 5.

Branch.—Length: Approximately 47.0 cm. Diameter: Approximately 2.9 mm. Length of central internode: Approximately 2.7 cm. Texture: Pubescent. Color of mature stem: 144A.

Foliage description:

General description.—Fragrance: Not noticeable. Form: Simple. Arrangement on non-flowering stem: Alternate. Arrangement on flowering stem: Opposite.

Leaves.—Aspect: Perpendicular to obtuse angle to stem. Shape: Elliptic. Margin: Slightly undulating. Apex: Acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 6.5 cm. Width of mature leaf: Approximately 2.9 cm. Texture of upper surface: Sparsely pubescent. Texture of lower surface: Sparsely pubescent. Color of upper surface of young foliage: 137C with venation of 137C. Color of lower surface of young foliage: 146B with venation of 144A. Color of upper surface of mature foliage: 137A with venation of 137A. Color of lower surface of mature foliage: 147B with venation of 144A.

Petiole.—Length: Approximately 9.0 mm. Width: Approximately 2.5 mm. Texture: Pubescent. Color upper surface: 144A. Color lower surface: 144B.

Flowering description:

Flowering habit.—‘Kerwhifan’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 10 to 12 days, flowers not persistent.

Flower description:

General description.—Type: Salverform. Arrangement: Flowers face outward; single, axillary. Fragrance: Not noticeable.

Bud.—Rate of opening: Generally takes 3 to 4 days for bud to progress from first color to fully open flower.

Bud just before opening.—Shape: Oblong with ruffled apices. Diameter: Approximately 6.6 mm. Color: 150C to 145A.

Corolla.—Diameter: Approximately 7.2 cm. Flower depth: Approximately 4.9 cm.

Petals.—Quantity: 5, fused to form a tube. Shape: Roughly spatulate; the midrib rises away from the throat, then recurves near the tip; the lamina rises away from the midrib. Appearance: Dull, smooth texture. Margin: Slightly undulating. Apex: Acute. Length from tube: Approximately 3.9 cm. Width: Approximately 3.7 cm. Texture of upper surface: Glabrous, slightly rugose. Texture of lower surface: Sparsely pubescent, slightly rugose. Color of upper surface when first open: 155C to 157D with midveins of 144B. Color of lower surface when first open: 155A with midveins of 144C. Color of upper surface when fully open: Whiter than 155C with midveins of 144B. Color of lower surface when fully open: Whiter than 155C with midveins of 144C.

Corolla tube.—Length: Approximately 3.2 cm. Diameter at distal end: Approximately 1.6 cm. Diameter at proximal end: Approximately 3.0 mm. Texture of inner surface: Fairly smooth. Texture of outer surface: Pubescent, somewhat coarse, and strongly ribbed at veins. Color of inner surface: 157D with venation of 144A to 144B. Color of outer surface: 157A to 157C with venation of 144C.

Peduncle.—Strength: Moderate. Aspect: Approximately a 25 to 35 degree angle to stem, becoming a 60 to 80 degree angle as flower ages. Length: Approximately 2.7 cm. Diameter: Approximately 1.6 mm. Texture: Pubescent. Color: 144A to 144B.

Sepals.—Quantity per flower: 5, fused at base. Shape: Narrow, irregular oblong, undulating margin. Apex: Blunt. Length: Approximately 1.9 cm. Width: Approximately 5.6 mm. Texture of upper surface: Pubescent. Texture of lower surface: Coarse, pubescent. Color of upper surface: 146A. Color of lower surface: 146B.

Reproductive organs.—Androecium: Stamen quantity: 5, partially fused to inside of corolla tube. Anther shape: Bilobed. Anther length: Approximately 2 to 2.5 mm. Anther color: 4D. Pollen amount: Abundant. Pollen color: 158B to 158C. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 2.3 cm. Stigma shape: Funnel. Stigma color: 144A to 144B. Style length: Approximately 1.9 cm. Style color: 145B to 145C. Ovary color: 144A.

Seed and fruit production: Neither seed nor fruit production has been observed.

US PP19,620 P2

5

Disease and pest resistance: Resistance to pathogens and pests common to *Petunia* has not been observed.
Weather tolerance: Exceptional resistance to wet weather has been observed.

6

What is claimed is:
1. A new and distinct cultivar of *Petunia* plant named 'Kerwhifan', substantially as herein shown and described.
* * * * *




FIG. 1




FIG. 2