

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
16 August 2007 (16.08.2007)

PCT

(10) International Publication Number  
WO 2007/090401 A2

## (51) International Patent Classification:

C12Q 1/68 (2006.01)

## (21) International Application Number:

PCT/DK2007/000060

## (22) International Filing Date: 5 February 2007 (05.02.2007)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

PA 2006 00181 8 February 2006 (08.02.2006) DK

PA 2007 00165 31 January 2007 (31.01.2007) DK

(71) Applicants (for all designated States except US):  
**AARHUS UNIVERSITET** [DK/DK]; Nordre Ringgade 1, DK-8000 Århus C (DK). **KVÆGAVLSFORENINGEN DANSIRE** [DK/DK]; Ebeltoftvej 16, DK-8900 Randers (DK).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): **LUND, Mogens**, Sandø [DK/DK]; Vestergade 7, Ørum, DK-8830 Tjele (DK). **BENDIXEN, Christian** [DK/DK]; Amstrupvej 7, DK-8860 Ulstrup (DK). **JENSEN, Helle** [DK/DK]; Middagsbøjvej 38, 1., DK-8800 Viborg (DK). **THOMSEN, Bo** [DK/DK]; Ryhaven 49, DK-8210 Århus V (DK). **SØRENSEN, Peter** [DK/DK]; Baldersvej 13, DK-8800

Viborg (DK). **SVENDSEN, Søren** [DK/DK]; Rindsvej 13, st. tv., DK-8900 Randers (DK). **THOMASEN, Jørn**, Rind [DK/DK]; Nørre Hvamvej 2, Hvam, DK-7500 Holstebro (DK). **NIELSEN, Vivi**, Hunnicke [DK/DK]; Veldsparken 72, Ørum, DK-8830 Tjele (DK). **MAJGREN, Bente**, Flügel [DK/DK]; Slænvej 10, Hørby, DK-9500 Hobro (DK). **GULDBRANDTSEN, Bernt** [DK/DK]; Fjældevænget 124, 1. th., DK-8210 Århus V (DK).


(74) Agent: **HØIBERG A/S**; Store Kongensgade 59 A, DK-1264 Copenhagen K (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

## (54) Title: CALVING CHARACTERISTICS



(57) Abstract: The invention relates to a method for determining calving characteristics in bovine subjects, wherein calving characteristics comprise stillbirth, calving difficulty and calf size at birth, which are all economically important factors. In particular, the method of the invention involves identification of genetic markers and/or Quantitative Trait Locus (QTL) for the determination of calving characteristics in a bovine subject. The determination of calving characteristics involves resolution of the specific microsatellite status. Furthermore, the invention relates to a diagnostic kit for detection of genetic marker(s) associated with calving characteristics. The method and kit of the present invention can be applied for selection of bovine subjects for breeding purposes. Thus, the invention provides a method of genetically selecting bovine subjects with calving characteristics that will yield cows less prone to stillbirth, calving difficulties and undesired calf size at birth.

WO 2007/090401 A2



European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

- *without international search report and to be republished upon receipt of that report*

## Calving characteristics

### Field of invention

The present invention relates to calving characteristics in bovine subjects. In particular, 5 the invention relates to genetic markers for the determination of calving characteristics in a bovine subject and a diagnostic kit for detection of genetic marker(s) associated with calving characteristics.

### Background of invention

10 Stillbirth, calving difficulty and calf size at birth are economic important calving traits, which are included in the Danish dairy cattle breeding program (Pedersen et al., 2003). The incidence of stillbirths for Holstein cattle has increased in several Holstein populations during the last two decades (Hansen et al., 2004). The increased incidence of stillbirths reduces the potential number of replacement heifers in dairy cattle herds 15 and is associated with ethical problems.

Both direct and maternal genetic components are associated with the calving traits. In Danish Holstein, the heritability ( $h^2$ ) estimates of the calving traits, measured as a direct sire effect ( $h^2=0.05-0.19$ ) are higher than the heritability estimates of the calving traits 20 measured as a maternal grand sire effect ( $h^2=0.04-0.06$ ). The genetic correlation between calving traits measured as a direct sire effect (0.69-0.93) are markedly higher than the genetic correlation between calving traits measured as a maternal grand sire effect (0.01-0.62). The genetic component associated with the calving traits may be due to the segregation of one or more quantitative trait loci (QTL).

25 Quantitative trait locus (QTL) is a region of DNA that is associated with a particular trait (e.g. a disease or calving characteristics). A QTL is not necessarily a gene itself, but rather a DNA region that is closely linked to the genes that underlie the trait in question. Most likely, a QTL is a set of genes that collectively encode a quantitative trait that 30 varies continuously across a population. Thus, the allelic variation of the QTL is associated with variation in a quantitative trait. The presence of QTL is inferred from genetic mapping, in which the genetic location of the QTL is determined relative to known genetic markers.

The identification of genetic markers that are linked to a particular phenotype, such as calving traits or to a heritable disease, has been facilitated by the discovery of 5 microsatellite markers as a source of polymorphic markers and single nucleotide polymorphisms linked to a mutation causing a specific phenotype. Markers linked to the mutation or the mutation itself causing a specific phenotype of interest are localised by use of genetic analysis in pedigrees and also by exploiting linkage disequilibrium (LD) when looking at populations

Linkage disequilibrium reflects recombination events dating back in history and the use 10 of LD mapping within families increases the resolution of mapping. LD exists when observed haplotypes in a population do not agree with the haplotype frequencies predicted by multiplying together the frequency of individual genetic markers in each haplotype. In this respect the term haplotype means a set of closely linked genetic markers present on one chromosome which tend to be inherited together.

15 In order for LD mapping to be efficient the density of genetic markers needs to be compatible with the distance across which LD extends in the given population. In a study of LD in dairy cattle population using a high number of genetic markers (284 autosomal microsatellite markers) it was demonstrated that LD extends over several tens of centimorgans for intrachromosomal markers (Farnir et al. 2000). Similarly,

20 Georges, M (2000) reported that the location of a genetic marker that is linked to a particular phenotype in livestock typically has a confidence interval of 20-30 cM (corresponding to maybe 500-1000 genes) (Georges, M., 2000). The existence of linkage disequilibrium is taken into account in order to use maps of particular regions of interest with high confidence.

25 Several QTL have been detected for calving traits in other Holstein populations (e.g. Kühn et al., 2003; Schrooten et al., 2000; Elo et al., 1999). Some QTL may affect more than one trait, and some QTL may even be located at the same chromosomal position for different traits. If the QTL affects multiple traits then it is important, for e.g. selection purposes, to test if it is a pleiotropic or linked QTL affecting the traits.

30 Calving traits such as stillbirth, calving difficulty and calf size are not easily predicted. The use of genetic analysis and genetic selection appears to be a possible method for prognostication of these calving traits. Once mapped, a QTL can be usefully applied in 35 marker assisted selection.

**Summary of invention**

It is an object of the present invention to provide an application method for marker assisted selection of polymorphisms in the bovine genome, wherein polymorphisms are 5 associated with calving trait characteristics, such as still birth, calving difficulties, and calf size; and/or provide genetic markers for use in such a method, and/or to provide animals selected using the method of the invention.

One aspect of the present invention relates to a method of determining calving 10 characteristics in a bovine subject, comprising detecting in a sample from said bovine subject the presence or absence of at least one genetic marker that is linked to at least one trait indicative of increased risk of stillbirth and/or increased risk of calving difficulties and/or increased risk of non-desired calf size, wherein said at least one genetic marker is located on the bovine chromosome BTA3 in a region flanked by and 15 including polymorphic microsatellite markers INRA006 and BM7225 and/or

BTA4 in the region flanked by and including polymorphic microsatellite markers BMS1788 and MGTG4B and/or,

BTA5 in the region flanked by and including polymorphic microsatellite markers BMS1095 and BM2830 and/or,

BTA7 in a region flanked by and including polymorphic microsatellite markers BM7160 and BL1043 and/or,

BTA8 in a region flanked by and including polymorphic microsatellite markers IDVGA-11 and BMS836 and/or,

BTA9 in a region flanked by and including polymorphic microsatellite markers BMS2151 and BMS1967 and/or,

BTA10 in a region flanked by and including polymorphic microsatellite markers DIK2658 and BMS2614 and/or,

BTA11 in the region flanked by and including polymorphic microsatellite markers BM716 and HEL13 and/or,

BTA12 in a region flanked by and including polymorphic microsatellite markers BMS410 and BMS2724 and/or,

BTA15 in a region flanked by and including polymorphic microsatellite markers BR3510 and BMS429 and/or,

BTA18 in a region flanked by and including polymorphic microsatellite markers IDVGA-31 and DIK4013 and/or,

25

30

35

BTA19 in a region flanked by and including polymorphic microsatellite markers BM9202 and BMS601 and/or,

BTA20 in a region flanked by and including polymorphic microsatellite markers BM3517 and UWCA26 and/or,

5 BTA21 in a region flanked by and including polymorphic microsatellite markers DIK5182 and IDVGA-30 and/or,

BTA22 in a region flanked by and including polymorphic microsatellite markers CSSM26 and BM4102 and/or,

BTA24 in a region flanked by and including polymorphic microsatellite markers 10 BMS917 and BMS3024 and/or,

BTA25 in a region flanked by and including polymorphic microsatellite markers ILSTS102 and AF5 and/or,

BTA26 in a region flanked by and including polymorphic microsatellite markers BMS651 and BM7237 and/or,

15 BTA28 in a region flanked by and including polymorphic microsatellite markers, BMC6020 and BMC2208, , wherein the presence of said at least one genetic marker is indicative of calving characteristics of said bovine subject and/or off-spring therefrom.

20 A second aspect of the present invention relates to diagnostic kit for use in detecting the presence in a bovine subject of at least one genetic marker associated with bovine calving characteristics, comprising at least one oligonucleotide sequence, wherein the nucleotide sequences are selected from any of SEQ ID NO.: 1 to SEQ ID NO.: 558 and/or any combination thereof.

25

**Description of drawings**

Fig. 1: Genome scan of BTA3 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL 30 correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect 35 the investigated calving traits.

Fig. 2: Genome scan of BTA4 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 3: Genome scan of BTA7 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 4: Genome scan of BTA7 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 5: Genome scan of BTA8 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL

analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 6: Genome scan of BTA8 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 7: Genome scan of BTA9 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 8: Genome scan of BTA10 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 9: Genome scan of BTA12 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the

positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 10: Genome scan of BTA12 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the

positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 11: Genome scan of BTA15 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are

designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 12: Genome scan of BTA18 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL

correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 13: Genome scan of BTA18 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL

correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis

represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

5 Fig. 14: Genome scan of BTA18 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis  
10 represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

15 Fig. 15: Genome scan of BTA18 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis  
20 represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

25 Fig. 16: Genome scan of BTA19 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis  
30 represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

35 Fig. 17: Genome scan of BTA20 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in

calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 18: Genome scan of BTA21 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in

calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 19: Genome scan of BTA22 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in

calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 20: Genome scan of BTA22 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in

calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 21: Genome scan of BTA24 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL

5

10

15

20

25

30

35

correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

5 Fig. 22: Genome scan of BTA25 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

10 Fig. 23: Genome scan of BTA25 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

15 Fig. 24: Genome scan of BTA26 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

20 Fig. 25: Genome scan of BTA26 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are

designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 26: Genome scan of BTA26 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are

designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 27: Genome scan of BTA28 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. Calving parameters are designated by D: Direct effect, M Maternal effect, while LK corresponds to stillbirth, FL

correspond to calving difficulty, and ST correspond to calf size. The number 1 in calving parameter designates that data is derived from first calving. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 28: Genome scan of BTA5 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

Fig. 29: Genome scan of BTA11 in relation to calving characteristics. Numbers refer to 'herdbook number' and calving parameter, respectively. The X-axis represents the distance of the chromosome expressed in Morgan according to the positions employed in this analysis. The Y-axis represents the test-statistics of the QTL analysis expressed

in the F-value. High F-values are indicative of genes, which affect the investigated calving traits.

#### **Detailed description of the invention**

5 The present invention relates to genetic determinants of calving characteristics in dairy cattle. Calving traits, such as calving difficulties, stillbirths and calf size are economically important factors in the dairy industry. Therefore, it is of economic interest to identify those bovine subjects that have a genetic predisposition for specific calving characteristics. Bovine subjects with genetic predisposition for calving characteristics  
10 are carriers of non-desired traits, which both complicate calving, and can be passed on to their offspring.

The term "bovine subject" refers to cattle of any breed and is meant to include both cows and bulls, whether adult or newborn animals. No particular age of the animals are

15 denoted by this term. One example of a bovine subject is a member of the Holstein breed. In one embodiment, the bovine subject is a member of the Holstein-Friesian cattle population. In another embodiment, the bovine subject is a member of the Holstein Swartbont cattle population. In another embodiment, the bovine subject is a member of the  
20 Deutsche Holstein Schwarzbunt cattle population. In another embodiment, the bovine subject is a member of the US Holstein cattle population. In one embodiment, the bovine subject is a member of the Red and White Holstein breed.

In another embodiment, the bovine subject is a member of the Deutsche Holstein Schwarzbunt cattle population. In one embodiment, the bovine subject is a member of any family, which include members of the Holstein breed. In one embodiment the  
25 bovine subject is a member of the Danish Red population. In another embodiment the bovine subject is a member of the Finnish Ayrshire population. In yet another embodiment the bovine subject is a member of the Swedish Red population. In a further embodiment the bovine subject is a member of the Danish Holstein population.

In another embodiment, the bovine subject is a member of the Swedish Red and White population. In yet another embodiment, the bovine subject is a member of the Nordic Red population.  
30

In one embodiment of the present invention, the bovine subject is selected from the group consisting of Swedish Red and White, Danish Red, Finnish Ayrshire, Holstein-Friesian, Danish Holstein and Nordic Red. In another embodiment of the present  
35

invention, the bovine subject is selected from the group consisting of Finnish Ayrshire and Swedish Red cattle. In another embodiment of the present invention, the bovine subject is selected from the group consisting of Finnish Ayrshire and Swedish Red cattle.

5

In one embodiment, the bovine subject is selected from the group of breeds shown in table 1a

10

15

20

25

30 Table 1a Breed names and breed codes assigned by ICAR (International Committee for Animal Recording)

Breed	Breed Code	National Breed Names Annex
Abondance	AB	-
Tyrol Grey	AL	2.2
Angus	AN	2.1
Aubrac	AU	
Ayrshire	AY	2.1
Belgian Blue	BB	
Blonde d'Aquitaine	BD	
Beefmaster	BM	
Braford	BO	
Brahman	BR	
Brangus	BN	
Brown Swiss	BS	2.1
Chianina	CA	
Charolais	CH	
Dexter	DR	
Galloway	GA	2.2
Guernsey	GU	
Gelbvieh	GV	
Hereford, horned	HH	
Hereford, polled	HP	
Highland Cattle	HI	
Holstein	HO	2.2
Jersey	JE	
Limousin	LM	
Maine-Anjou	MA	
Murray-Grey	MG	
Montbéliard	MO	
Marchigiana	MR	
Normandy	NO**	
Piedmont	PI	2.2
Pinzgau	PZ	
European Red Dairy Breed	[RE]*	2.1, 2.2
Romagnola	RN	
Holstein, Red and White	RW***	2.2
Salers	SL**	
Santa Gertrudis	SG	
South Devon	SD	
Shorthorn	[SH]*	2.2
Simental	SM	2.2
Sahiwal	SW	
Tarentaise	TA	
Welsh Black	WB	
Buffalo (Bubalis bubalis)	BF	
* new breed code		
** change from earlier code because of existing code in France		
*** US proposal WW		

In one embodiment, the bovine subject is a member of a breed selected from the group of breeds shown in table 1b

Table 1b Breed names

<u>National Breed Names</u>		
<u>English Name</u>		<u>National names</u>
Angus	Including	Aberdeen Angus Canadian Angus American Angus German Angus
Ayrshire	Including	Ayrshire in Australia Canada Colombia Czech Republic Finland Kenya New Zealand Norway (NRF) Russia South Africa Sweden (SRB) and SAB UK US Zimbabwe
Belgian Blue	French: Flemish:	Blanc-bleu Belge Witblauw Ras van Belgie
Brown Swiss	German: Italian: French: Spanish: Serbo-Croatian: Czech: Romanian: Russian: Bulgarian:	Braunvieh Razza Bruna Briue Bruna, Parda Alpina Slovenscko belo Husky Karpatsky Shivitskaja Bruna Bijarska kafyava
European Red Dairy Breed	Including	Danish Red Angeln Swedish Red and White Norwegian Red and White Estonian Red Latvian Brown Lithuanian Red Byelorussian Red Polish Red Lowland

In one embodiment, the bovine subject is a member of a breed selected from the group of breeds shown in table 1c

Table 1c Breed names

English Name	National Breed Names	
	English Name	National names
<i>European Red Dairy Breed (continued)</i>		Ukrainian Polish Red (French Rouge Flamande?) (Belgian Flamande Rouge?)
Galloway:	Including	Black and Dun Galloway Belted Galloway Red Galloway White Galloway
Holstein, Black and White:	Dutch: German: Danish: British: Swedish: French: Italian: Spanish:	Holstein Schwarzbunt Deutsche Holstein, schwarzbuunt Sortbroget Dansk Melkekvæg Holstein Friesian Svensk Läglands Boskap Prüm Holstein Holstein Frisona Holstein Frisona
Holstein, Red and White	Dutch: German: Danish:	Holstein, roodbunt Holstein, rotbunt Roedbroget Dansk Melkekvæg
Piedmont	Italian:	Piemontese
Shorthorn	Including	Dairy Shorthorn Beef Shorthorn Polled Shorthorn
Simmental	Including dual purpose and beef use	
	German: French: Italian: Czech: Slovakian: Romanian: Russian:	Fleckvieh Simmental Française Razza Pezzata Rossa Český strakář Slovenský strakář Baltata românească Simmentalskjære
Tyrol Grey	German: Italian:	Tiroler Grauvieh Oberinntaler Grauvieh Rätisches Grauvieh Razza Grigia Alpina

The term "genetic marker" refers to a variable nucleotide sequence (polymorphism) of

5 the DNA on the bovine chromosome. The variable nucleotide sequence can be identified by methods known to a person skilled in the art, for example by using specific oligonucleotides in for example amplification methods and/or hybridization techniques and/or observation of a size difference. However, the variable nucleotide sequence may also be detected by sequencing or for example restriction fragment length polymorphism analysis. The variable nucleotide sequence may be represented by a deletion, an insertion, repeats, and/or a point mutation. Thus, a genetic marker 10 comprises a variable number of polymorphic alleles.

One type of genetic marker is a microsatellite marker that is linked to a quantitative trait locus. Microsatellite markers refer to short sequences repeated after each other. In short sequences are for example one nucleotide, such as two nucleotides, for example three nucleotides, such as four nucleotides, for example five nucleotides, such as six nucleotides, for example seven nucleotides, such as eight nucleotides, for example nine nucleotides, such as ten nucleotides. However, changes sometimes occur and the number of repeats may increase or decrease. The specific definition and locus of the polymorphic microsatellite markers can be found in the USDA genetic map (Kappes et al. 1997; or by following the link to U.S. Meat Animal Research Center

10 <http://www.marc.usda.gov/>).

In one embodiment of the present invention, specific marker alleles are linked to quantitative trait loci affecting calving characteristics.

15 It is furthermore appreciated that the nucleotide sequences of the genetic markers of the present invention are genetically linked to traits for calving in a bovine subject. Consequently, it is also understood that a number of genetic markers may be generated from the nucleotide sequence of the DNA region(s) flanked by and including the genetic markers according to the method of the present invention.

20

### **Calving trait characteristics**

Calving in a bovine subject is affected by a number of characteristics. Traits that affect calving according to the present invention are for example the occurrence of stillbirth (SB), calving difficulty (CD) and the size of the calf at birth (CS). The traits are

25 assessed by a direct effect (D) of the sire in the calf. However, the traits are also assessed as a maternal effect (M) of the sire in the mother of the calf.

By the term calving characteristics is meant traits which affect calving in the bovine subject or its off-spring. Thus, calving characteristics of a bull are physically manifested by its off-spring – both female and male.

30 In the present invention calving characteristics comprise the traits SB, CD, and CS, which refer to the following characteristics:

SB: Designates stillbirths.

CS: Size of calves.

CD: Calving difficulties, which are based on registrations from the farmers where it is subjectively registered how difficult the calving is. The calving difficulties consist of four categories:

- 5 1: easy with no help
- 2: easy with assistance
- 3: difficult but without veterinary assistance
- 4: difficult with veterinary assistance

In one embodiment of the present invention, the method and kit described herein relates to still births, calving difficulties as categorized herein and/or calf size. In one embodiment of the present invention, the method and kit described herein relates to still births. In another embodiment, the method and kit of the present invention pertains to calving difficulties, such as detected by the calving difficulty categories described above. In yet another embodiment, the method and kit of the present invention relates to calf size. In another embodiment of the present invention, the method and kit described herein relates to any combination of still birth, calving difficulties and/or calf size.

#### **Granddaughter design**

20 The granddaughter design includes analysing data from DNA-based markers for grandsires that have been used extensively in breeding and for sons of grandsires where the sons have produced offspring. The phenotypic data that are to be used together with the DNA-marker data are derived from the daughters of the sons. Such phenotypic data could be for example milk production features, features relating to 25 calving, meat quality, or disease. One group of daughters has inherited one allele from their father whereas a second group of daughters has inherited the other allele from their father. By comparing data from the two groups information can be gained whether a fragment of a particular chromosome is harbouring one or more genes that affect the trait in question. It may be concluded whether a QTL is present within this fragment of 30 the chromosome.

A prerequisite for performing a granddaughter design is the availability of detailed phenotypic data. In the present invention such data have been available (<http://www.lr.dk/kvaeg/diverse/principles.pdf> ).

In contrast, DNA markers can be used directly to provide information of the traits passed on from parents to one or more of their offspring when a number of DNA markers on a chromosome have been determined for one or both parents and their offspring. The markers may be used to calculate the genetic history of the chromosome 5 linked to the DNA markers.

### **Frequency of recombination**

The frequency of recombination is the likelihood that a recombination event will occur between two genes or two markers. The frequency of recombination may be calculated 10 as the genetic distance between the two genes or the two markers. Genetic distance is measured in units of centiMorgan (cM). One centiMorgan is equal to a 1% chance that a marker at one genetic locus will be separated from a marker at a second locus due to crossing over in a single generation. One centiMorgan is equivalent, on average, to one million base pairs.

15

### **Chromosomal regions and markers**

BTA is short for *Bos taurus* autosome.

One aspect of the present invention relates to a method of determining calving characteristics in a bovine subject, comprising detecting in a sample from said bovine 20 subject the presence or absence of at least one genetic marker that is linked to at least one trait indicative of increased risk of stillbirth and/or increased risk of calving difficulties and/or increased risk of non-desired calf size, wherein said at least one genetic marker is located on the bovine chromosome BTA3 in a region flanked by and including polymorphic microsatellite markers INRA006 and BM7225 and/or BTA4 in the 25 region flanked by and including polymorphic microsatellite markers BMS1788 and MGTG4B and/or, BTA5 in the region flanked by and including polymorphic microsatellite markers BMS1095 and BM2830 and/or, BTA7 in a region flanked by and including polymorphic microsatellite markers BM7160 and BL1043 and/or, BTA8 in a region flanked by and including polymorphic microsatellite markers IDVGA-11 and 30 BMS836 and/or, BTA9 in a region flanked by and including polymorphic microsatellite markers BMS2151 and BMS1967 and/or, BTA10 in a region flanked by and including polymorphic microsatellite markers DIK2658 and BMS2614 and/or, BTA11 in the region flanked by and including polymorphic microsatellite markers BM716 and HEL13 and/or, BTA12 in a region flanked by and including polymorphic microsatellite markers 35 BMS410 and BMS2724 and/or, BTA15 in a region flanked by and including

polymorphic microsatellite markers BR3510 and BMS429 and/or, BTA18 in a region flanked by and including polymorphic microsatellite markers IDVGA-31 and DIK4013 and/or, BTA19 in a region flanked by and including polymorphic microsatellite markers BM9202 and BMS601 and/or, BTA20 in a region flanked by and including polymorphic 5 microsatellite markers BM3517 and UWCA26 and/or, BTA21 in a region flanked by and including polymorphic microsatellite markers DIK5182 and IDVGA-30 and/or, BTA22 in a region flanked by and including polymorphic microsatellite markers CSSM26 and BM4102 and/or, BTA24 in a region flanked by and including polymorphic microsatellite markers BMS917 and BMS3024 and/or, BTA25 in a region flanked by and including polymorphic microsatellite markers ILSTS102 and AF5 and/or, BTA26 in a region 10 flanked by and including polymorphic microsatellite markers BMS651 and BM7237 and/or, BTA28 in a region flanked by and including polymorphic microsatellite markers, BMC6020 and BMC2208, , wherein the presence of said at least one genetic marker is indicative of calving characteristics of said bovine subject and/or off-spring therefrom.

15

In order to determine calving characteristics in a bovine subject, wherein the at least one genetic marker is located on a bovine chromosome in the region flanked by and including the polymorphic microsatellite marker, it is appreciated that more than one genetic marker may be employed in the present invention. For example the at least one 20 genetic marker may be a combination of at least two or more genetic markers such that the accuracy may be increased, such as at least three genetic markers, for example four genetic markers, such as at least five genetic markers, for example six genetic markers, such as at least seven genetic markers, for example eight genetic markers, such as at least nine genetic markers, for example ten genetic markers.

25

The at least one genetic marker may be located on at least one bovine chromosome, such as two chromosomes, for example three chromosomes, such as four chromosomes, for example five chromosomes, and/or such as six chromosomes.

30

In a preferred embodiment the at least one marker is selected from any of the individual markers of the tables shown herein.

BTA3

In one embodiment of the invention the at least one genetic marker is located on the 35 bovine chromosome BTA3. In one specific embodiment of the present invention, the at

least one genetic marker is located in the region from about 17.1 cM to about 101.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers INRA006 and BM7225. The at least one genetic 5 marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 2a:

10 Table 2a

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA006	17.1
UWCA7	17.4
ILSTS096	27.4
DIK4403	32.5
RME23	32.5
BMS963	32.9
BMS819	33.5
FCGR1	34.6
BL41	43.3
DIK4353	52.5
INRA003	59.4
BMS2790	62.4
ILSTS029	64.9
BM220	66.3
INRA123	66.3
BMS862	67.4
HUJ246	68.0
BMS937	68.0
DIK4664	68.3
DIK2702	77.6
HUJII77	87.3
DIK2686	95.5

BM7225	101.8
--------	-------

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 34.6 cM to about 87.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers FCGR1 and HUJII77. The at least one genetic marker is selected from the group of markers shown in Table 2b:

5

Table 2b

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
FCGR1	34.6
BL41	43.3
DIK4353	52.5
INRA003	59.4
BMS2790	62.4
ILSTS029	64.9
BM220	66.3
INRA123	66.3
BMS862	67.4
HUJ246	68.0
BMS937	68.0
DIK4664	68.3
DIK2702	77.6
HUJII77	87.3

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 32.5 cM to about 59.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers DIK4403 and INRA003. The at least one genetic marker is selected from the group of markers shown in Table 2c:

15

Table 2c

Marker on BTA3	Relative position (cM)
----------------	------------------------

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK4403	32.5
RME23	32.5
BMS963	32.9
BMS819	33.5
FCGR1	34.6
BL41	43.3
DIK4353	52.5
INRA003	59.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.6 cM to about 101.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers DIK2702 and BM7225. The at least one genetic marker is selected from the group of markers shown in Table 2d:

Table 2d

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2702	77.6
HUJII77	87.3
DIK2686	95.5
BM7225	101.8

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 52.5 cM to about 68.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers DIK4353 and DIK4664. The at least one genetic marker is selected from the group of markers shown in Table 2e:

Table 2e

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK4353	52.5

INRA003	59.4
BMS2790	62.4
ILSTS029	64.9
BM220	66.3
INRA123	66.3
BMS862	67.4
HUJ246	68.0
BMS937	68.0
DIK4664	68.3

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 59.4 cM to about 66.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers INRA003 and INRA123. The at least one genetic marker is selected from the group of markers shown in Table 2f:

Table 2f

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA003	59.4
BMS2790	62.4
ILSTS029	64.9
BM220	66.3
INRA123	66.3

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 32.5 cM to about 52.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA3 in the region flanked by and including the markers DIK4403 and DIK4353. The at least one genetic marker is selected from the group of markers shown in Table 2g:

Table 2g

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>

DIK4403	32.5
RME23	32.5
BMS963	32.9
BMS819	33.5
FCGR1	34.6
BL41	43.3
DIK4353	52.5

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.6 cM to 101.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA3. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA3 in the region flanked by and including the marker FCGR1 and HUJII77. The at least one genetic marker is selected from the group of markers shown in Table 2h:

Table 2h

Marker on BTA3	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2702	77.6
HUJII77	87.3
DIK2686	95.5
BM7225	101.8

10 BTA4

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA4. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 12.5 cM to about 112.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the

15 at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers BMS1788 and MGTG4B. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected 20 from the group of markers shown in Table 3a:

Table 3a

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1788	12.5
BMS2646	43.2
TGLA116	52.5
INRA072	63.0
BM8233	73.4
BMS648	91.2
BR6303	104.9
MGTG4B	112.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 12.5 cM to about 91.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers BMS1788 and BMS648. The at least one genetic marker is selected from the group of markers shown in Table 3b:

Table 3b

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1788	12.5
BMS2646	43.2
TGLA116	52.5
INRA072	63.0
BM8233	73.4
BMS648	91.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 43.2 cM to about 91.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers BMS2646 and BMS648. The at least one genetic marker is selected from the group of markers shown in Table 3c:

Table 3c

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2646	43.2
TGLA116	52.5
INRA072	63.0
BM8233	73.4
BMS648	91.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 43.2 cM to about 63.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA4 in the region flanked by and including the markers BMS2646 and INRA072. The at least one genetic marker is selected from the group of markers shown in Table 3d:

Table 3d

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2646	43.2
TGLA116	52.5
INRA072	63.0

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 52.2 cM to about 73.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers TGLA116 and BM8233. The at least one genetic marker is selected from the 15 group of markers shown in Table 3e:

Table 3e

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
TGLA116	52.5
INRA072	63.0
BM8233	73.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 63.0 cM to about 91.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers INRA072 and BMS648. The at least one genetic marker is selected from the group of markers shown in Table 3f:

Table 3f

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA072	63.0
BM8233	73.4
BMS648	91.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 63.0 cM to about 73.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA4. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA4 in the region flanked by and including the markers INRA072 and BM8233. The at least one genetic marker is selected from the group of markers shown in Table 3g:

Table 3g

Marker on BTA4	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA072	63.0
BM8233	73.4

## BTA5

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA5. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 116.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers BMS1095 and BM2830. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is

significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 4a:

Table 4a

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1095	0.0
BM6026	6.0
MNB-33	7.4
BMS610	12.0
BP1	17.3
DIK4747	18.3
DIK2718	30.1
AGLA293	32.3
DIK5002	33.7
DIK4759	40.3
BMC1009	41.7
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9
BMS1248	90.8
BM315	103.2
BMS1658	105.7
BM2830	116.9

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 103.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers BMS1095 and BM315. The at least one genetic marker is selected from the group of markers shown in Table 4b:

10

Table 4b

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1095	0.0
BM6026	6.0
MNB-33	7.4
BMS610	12.0
BP1	17.3
DIK4747	18.3
DIK2718	30.1
AGLA293	32.3
DIK5002	33.7
DIK4759	40.3
BMC1009	41.7
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9
BMS1248	90.8
BM315	103.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.1 cM to about 103.2 cM (<http://www.marc.usda.gov/>) on the

5 bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers DIK2718 and BM315. The at least one genetic marker is selected from the group of markers shown in Table 4c:

Table 4c

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>

DIK2718	30.1
AGLA293	32.3
DIK5002	33.7
DIK4759	40.3
BMC1009	41.7
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9
BMS1248	90.8
BM315	103.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.1 cM to about 78.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers DIK2718 and BMS1216. The at least one genetic marker is selected from the group of markers shown in Table 4d:

Table 4d

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2718	30.1
AGLA293	32.3
DIK5002	33.7
DIK4759	40.3
BMC1009	41.7
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2

ETH10	71.8
CSSM022	74.2
BMS1216	78.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 18.3 cM to about 56.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA5 in the region flanked by and including the markers DIK4747 and RM500. The at least one genetic marker is selected from the group of markers shown in Table 4e:

Table 4e

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK4747	18.3
DIK2718	30.1
AGLA293	32.3
DIK5002	33.7
DIK4759	40.3
BMC1009	41.7
CSSM034	45.5
RM500	56.3

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 17.3 cM to about 33.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers BP1 and DIK5002. The at least one genetic marker is selected from the group of markers shown in Table 4f:

15 Table 4f

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BP1	17.3
DIK4747	18.3
DIK2718	30.1

AGLA293	32.3
DIK5002	33.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 45.5 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers CSSM034 and DIK2943. The at least one genetic marker is selected from the group of markers shown in Table 4g:

Table 4g

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 45.5 cM to about 66.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers CSSM034 and DIK5046. The at least one genetic marker is selected from the group of markers shown in Table 4h:

Table 4h

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
CSSM034	45.5
RM500	56.3
BMS1617	56.3
DIK5046	66.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 66.2 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers DIK5046 and DIK2943. The at least one genetic marker is selected from the group of markers shown in Table 4i:

Table 4i

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5046	66.2
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 71.8 cM to about 90.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA5. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA5 in the region flanked by and including the markers ETH10 and BMS1248. The at least one genetic marker is selected from the group of markers shown in Table 4j:

Table 4j

Marker on BTA5	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ETH10	71.8
CSSM022	74.2
BMS1216	78.2
DIK2943	82.9
BMS1248	90.8

## BTA7

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA7. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 135.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the

at least one genetic marker is located on the bovine chromosome BTA7 in the region flanked by and including the markers BM7160 and BL1043. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, 5 for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 5a:

Table 5a

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM7160	0.0
BL1067	14.7
BMS713	16.8
DIK5321	22.3
DIK4421	22.7
DIK2207	26.7
DIK5412	30.2
IL4	32.0
BM6105	37.9
TGLA303	39.3
DIK2819	47.9
DIK4606	55.3
BM7247	57.3
UWCA20	58.6
BM6117	62.2
BMS2840	65.3
DIK2915	76.2
BMS2258	77.2
OARAE129	95.9
DIK2895	103.1
ILSTS006	116.6
BL1043	135.6

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.2 cM to about 95.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA7 in the region flanked by and including the markers DIK5412 and OARAE129. The at least one genetic marker is selected from the group of markers shown in Table 5b:

Table 5b

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5412	30.2
IL4	32.0
BM6105	37.9
TGLA303	39.3
DIK2819	47.9
DIK4606	55.3
BM7247	57.3
UWCA20	58.6
BM6117	62.2
BMS2840	65.3
DIK2915	76.2
BMS2258	77.2
OARAE129	95.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.2 cM to about 55.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA7 in the region flanked by and including the markers DIK5412 and DIK4606. The at least one genetic marker is selected from the group of markers shown in Table 5c:

Table 5c

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5412	30.2
IL4	32.0

BM6105	37.9
TGLA303	39.3
DIK2819	47.9
DIK4606	55.3

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 58.6 cM to about 95.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA7 in the region flanked by and including the markers UWCA20 and OARAE129. The at least one genetic marker is selected from the group of markers shown in Table 5d:

Table 5d

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
UWCA20	58.6
BM6117	62.2
BMS2840	65.3
DIK2915	76.2
BMS2258	77.2
OARAE129	95.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.2 cM to about 135.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA7 in the region flanked by and including the markers BMS2258 and BL1043. The at least one genetic marker is selected from the group of markers shown in Table 5e:

Table 5e

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2258	77.2
OARAE129	95.9
DIK2895	103.1
ILSTS006	116.6

BL1043	135.6
--------	-------

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.2 cM to about 116.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA7 in the region flanked by and including the markers BMS2258 and ILSTS006. The at least one genetic marker is selected from the group of markers shown in Table 5f:

Table 5f

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2258	77.2
OARAE129	95.9
DIK2895	103.1
ILSTS006	116.6

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.2 cM to about 95.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA7. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA7 in the region flanked by and including the markers BMS2258 and OARAE129. The at least one genetic marker is selected from

15 the group of markers shown in Table 5g:

Table 5g

Marker on BTA7	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2258	77.2
OARAE129	95.9

#### BTAA8

In one embodiment of the invention the at least one genetic marker is located on the

20 bovine chromosome BTA8. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 11.3 cM to about 122.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA8. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA8 in the region flanked by and including the markers IDVGA-11 and BMS836. The at least one genetic

marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 6a:

Table 6a

Marker on BTA8	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-11	11.3
BMS1591	31.4
BMS678	41.6
INRA129	54.6
BMS2072	66.0
BMS887	68.5
URB037	69.0
MCM64	71.1
CSSM047	118.7
BMS836	122.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 11.3 cM to about 71.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA8. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA8 in the region flanked by and including the markers IDVGA-11 and MCM64. The at least one genetic marker is selected from the group of markers shown in Table 6b:

Table 6b

Marker on BTA8	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-11	11.3
BMS1591	31.4
BMS678	41.6
INRA129	54.6
BMS2072	66.0
BMS887	68.5

URB037	69.0
MCM64	71.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 41.6 cM to about 66.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA8. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA8 in the region flanked by and including the markers BMS678 and BMS2072. The at least one genetic marker is selected from the group of markers shown in Table 6c:

Table 6c

Marker on BTA8	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS678	41.6
INRA129	54.6
BMS2072	66.0

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 71.1 cM to about 122.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA8. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA8 in the region flanked by and including the markers MCM64 and BMS836. The at least one genetic marker is selected from the 15 group of markers shown in Table 6d:

Table 6d

Marker on BTA8	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
MCM64	71.1
CSSM047	118.7
BMS836	122.9

20 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 11.3 cM to about 41.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA8. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA8 in the region flanked by and including the markers IDVGA-11 and BMS678. The at least one genetic marker is selected from the group of markers shown in Table 6e:

Table 6e

Marker on BTA8	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-11	11.3
BMS1591	31.4
BMS678	41.6

## BTA9

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA9. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 8.49 cM to about 109.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers BMS2151 and BMS1967. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 7a:

15 Table 7a

Marker on BTA9	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2151	8.49
ETH225	12.8
ILSTS037	26.3
BM2504	30.9
DIK2892	30.9
DIK3003	36.5
DIK3002	36.5
BMS1267	38.7
DIK5142	43.8
BMS555	43.8
DIK5364	45.7
UWCA9	50.0

DIK4720	54.0
BMS1290	64.9
DIK2816	68.1
BM6436	77.6
BMS2753	79.2
BM4208	90.7
BMS2819	91.0
BMS2295	98.6
BMS1967	109.3

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 12.8 cM to about 90.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers ETH225 and BM4208. The at least one genetic marker is selected from the group of markers shown in Table 7b:

Table 7b

Marker on BTA9	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ETH225	12.8
ILSTS037	26.3
BM2504	30.9
DIK2892	30.9
DIK3003	36.5
DIK3002	36.5
BMS1267	38.7
DIK5142	43.8
BMS555	43.8
DIK5364	45.7
UWCA9	50.0
DIK4720	54.0
BMS1290	64.9
DIK2816	68.1
BM6436	77.6

BMS2753	79.2
BM4208	90.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 12.8 cM to about 64.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers ETH225 and BMS1290. The at least one genetic marker is selected from the group of markers shown in Table 7c:

Table 7c

Marker on BTA9	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ETH225	12.8
ILSTS037	26.3
BM2504	30.9
DIK2892	30.9
DIK3003	36.5
DIK3002	36.5
BMS1267	38.7
DIK5142	43.8
BMS555	43.8
DIK5364	45.7
UWCA9	50.0
DIK4720	54.0
BMS1290	64.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 50.0 cM to about 91.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers UWCA9 and BMS2819. The at least one genetic marker is selected from the group of markers shown in Table 7d:

Table 7d

Marker on BTA9	Relative position (cM)

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
UWCA9	50.0
DIK4720	54.0
BMS1290	64.9
DIK2816	68.1
BM6436	77.6
BMS2753	79.2
BM4208	90.7
BMS2819	91.0

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 50.0 cM to about 79.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA9 in the region flanked by and including the markers UWCA9 and BMS2753. The at least one genetic marker is selected from the group of markers shown in Table 7e:

Table 7e

Marker on BTA9	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
UWCA9	50.0
DIK4720	54.0
BMS1290	64.9
DIK2816	68.1
BM6436	77.6
BMS2753	79.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 45.7 cM to about 68.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers DIK5364 and DIK2816. The at least one genetic marker is selected from the group of markers shown in Table 7f:

Table 7f

Marker on BTA9	Relative position (cM)

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5364	45.7
UWCA9	50.0
DIK4720	54.0
BMS1290	64.9
DIK2816	68.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 12.8 cM to about 43.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA9. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA9 in the region flanked by and including the markers ETH225 and DIK5142. The at least one genetic marker is selected from the group of markers shown in Table 7g:

Table 7g

Marker on BTA9	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ETH225	12.8
ILSTS037	26.3
BM2504	30.9
DIK2892	30.9
DIK3003	36.5
DIK3002	36.5
BMS1267	38.7
DIK5142	43.8

10 BTA10

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA10. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 2.7 cM to about 104.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers DIK2658 and BMS2614. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least

one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 8a:

Table 8a

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2658	2.7
DIK2503	9.0
CSSM38	11.0
BMS528	24.0
BM1237	24.7
MB077	35.1
DIK2000	37.5
BMS2742	44.3
BMS529	55.6
DIK2361	56.5
BM888	60.0
TGLA433	74.0
INRA037	79.0
BMS1620	80.4
ILSTS070	81.4
BMS2641	87.5
BMS614	100.0
BMS2614	109.4

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 9.0 cM to about 35.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers DIK2503 and MB077. The at least one genetic marker is selected from the

10 group of markers shown in Table 8b:

Table 8b

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2503	9.0

CSSM38	11.0
BMS528	24.0
BM1237	24.7
MB077	35.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 11.0 cM to about 37.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers CSSM38 and DIK2000. The at least one genetic marker is selected from the group of markers shown in Table 8c:

Table 8c

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
CSSM38	11.0
BMS528	24.0
BM1237	24.7
MB077	35.1
DIK2000	37.5

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 24.0 cM to about 35.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers BMS528 and MB077. The at least one genetic marker is selected from the group of markers shown in Table 8d:

Table 8d

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS528	24.0
BM1237	24.7
MB077	35.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 37.5 cM to about 80.4 cM (<http://www.marc.usda.gov/>) on the

bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers DIK2000 and BMS1620. The at least one genetic marker is selected from the group of markers shown in Table 8e:

5 Table 8e

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2000	37.5
BMS2742	44.3
BMS529	55.6
DIK2361	56.5
BM888	60.0
TGLA433	74.0
INRA037	79.0
BMS1620	80.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 44.3 cM to about 74.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is

10 located on the bovine chromosome BTA10 in the region flanked by and including the markers BMS2742 and TGLA433. The at least one genetic marker is selected from the group of markers shown in Table 8f:

Table 8f

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2742	44.3
BMS529	55.6
DIK2361	56.5
BM888	60.0
TGLA433	74.0

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 56.5 cM to about 74.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the

markers DIK2361 and TGLA433. The at least one genetic marker is selected from the group of markers shown in Table 8g:

Table 8g

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2361	56.5
BM888	60.0
TGLA433	74.0

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 74.0 cM to about 87.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers TGLA433 and BMS2641. The at least one genetic marker is selected from the group of markers shown in Table 8h:

Table 8h

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
TGLA433	74.0
INRA037	79.0
BMS1620	80.4
ILSTS070	81.4
BMS2641	87.5

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 87.5 cM to about 109.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA10. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA10 in the region flanked by and including the markers BMS2641 and BMS2614. The at least one genetic marker is selected from the group of markers shown in Table 8i:

Table 8i

Marker on BTA10	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2641	87.5

BMS614	100.0
BMS2614	109.4

## BTA11

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA11. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 19.4 cM to about 122.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers BM716 and HEL13. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table9a:

Table 9a

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM716	19.4
BMS2569	21.1
BM2818	30.0
INRA177-2	34.8
INRA177	34.8
RM096	40.5
INRA131	47.3
BM7169	50.3
BM6445	61.6
ILSTS036	61.6
BMS1822	65.9
TGLA58	73.1
BMS2047	78.5
HUJV174	92.2
BMS989	92.2
TGLA436	105.2

BMS460	109.4
ILSTS045	114.2
DIK4819	115.0
HEL13	122.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 19.4 cM to about 92.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA11 in the region flanked by and including the markers BM716 and BMS989. The at least one genetic marker is selected from the group of markers shown in Table 9b:

Table 9b

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM716	19.4
BMS2569	21.1
BM2818	30.0
INRA177-2	34.8
INRA177	34.8
RM096	40.5
INRA131	47.3
BM7169	50.3
BM6445	61.6
ILSTS036	61.6
BMS1822	65.9
TGLA58	73.1
BMS2047	78.5
HUJV174	92.2
BMS989	92.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 19.4 cM to about 50.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the

markers BM716 and BM7169. The at least one genetic marker is selected from the group of markers shown in Table 9c:

Table 9c

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM716	19.4
BMS2569	21.1
BM2818	30.0
INRA177-2	34.8
INRA177	34.8
RM096	40.5
INRA131	47.3
BM7169	50.3

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.0 cM to about 50.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers BM2818 and BM7169. The at least one genetic marker is selected from the group of markers shown in Table 9d:

10

Table 9d

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM2818	30.0
INRA177-2	34.8
INRA177	34.8
RM096	40.5
INRA131	47.3
BM7169	50.3

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 34.8 cM to about 47.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the

markers INRA177-2 and INRA131. The at least one genetic marker is selected from the group of markers shown in Table 9e:

Table 9e

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA177-2	34.8
INRA177	34.8
RM096	40.5
INRA131	47.3

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 50.3 cM to about 92.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers BM7169 and BMS989. The at least one genetic marker is selected from the group of markers shown in Table 9f:

10

Table 9f

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM7169	50.3
BM6445	61.6
ILSTS036	61.6
BMS1822	65.9
TGLA58	73.1
BMS2047	78.5
HUJV174	92.2
BMS989	92.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 61.6 cM to about 92.2 cM (<http://www.marc.usda.gov/>) on the

15 bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers BM6445 and BMS989. The at least one genetic marker is selected from the group of markers shown in Table 9g:

Table 9g

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM6445	61.6
ILSTS036	61.6
BMS1822	65.9
TGLA58	73.1
BMS2047	78.5
HUJV174	92.2
BMS989	92.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 73.3 cM to about 92.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers TGLA58 and BMS989. The at least one genetic marker is selected from the group of markers shown in Table 9h:

Table 9h

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
TGLA58	73.1
BMS2047	78.5
HUJV174	92.2
BMS989	92.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 92.2 cM to about 109.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA11. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA11 in the region flanked by and including the markers HUJV174 and BMS460. The at least one genetic marker is selected from the group of markers shown in Table 9i:

Table 9i

Marker on BTA11	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
HUJV174	92.2

BMS989	92.2
TGLA436	105.2
BMS460	109.4

## BTA12

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA12. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 109.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA12. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA12 in the region flanked by and including the markers BMS410 and BMS2724. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 10a:

Table 10a

Marker on BTA12	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS410	0.0
BM6108	15.1
BM860	50.4
BMS975	63.8
BMS1316	102.0
BMS2724	109.0

15

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 50.4 cM to about 109.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA12. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA12 in the region flanked by and including the markers BM860 and BMS2724. The at least one genetic marker is selected from the group of markers shown in Table 10b:

Table 10b

Marker on BTA12	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>

BM860	50.4
BMS975	63.8
BMS1316	102.0
BMS2724	109.0

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 50.4 cM to about 102.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA12. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA12 in the region flanked by and including the markers BM860 and BMS1316. The at least one genetic marker is selected from the group of markers shown in Table 10c:

Table 10c

Marker on BTA12	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM860	50.4
BMS975	63.8
BMS1316	102.0

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 63.8 cM to about 102.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA12. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA12 in the region flanked by and including the markers BMS975 and BMS1316. The at least one genetic marker is selected from the group of markers shown in Table 10d:

Table 10d

Marker on BTA12	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS975	63.8
BMS1316	102.0

#### BT15

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BT15. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 9.4 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BT15. In one

20

embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BR3510 and BMS429. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table11a:

Table 11a

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BR3510	9.4
BMS2533	13.9
INRA050	31.1
JAB8	31.2
BMS2684	48.2
DIK1106	51.9
INRA145	67.8
IDVGA-10	67.8
DIK4850	74.1
DIK2768	77.9
ILSTS027	83.4
BMS812	84.9
BMS2076	91.8
BL1095	94.8
BMS820	98.2
BMS927	105.0
BMS429	109.8

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 48.2 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS2684 and BMS429. The at least one genetic marker is selected from the group of markers shown in Table 11b:

15

Table 11b

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2684	48.2
DIK1106	51.9
INRA145	67.8
IDVGA-10	67.8
DIK4850	74.1
DIK2768	77.9
ILSTS027	83.4
BMS812	84.9
BMS2076	91.8
BL1095	94.8
BMS820	98.2
BMS927	105.0
BMS429	109.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 48.2 cM to about 91.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS2684 and BMS2076. The at least one genetic marker is selected from the group of markers shown in Table 11c:

Table 11c

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2684	48.2
DIK1106	51.9
INRA145	67.8
IDVGA-10	67.8
DIK4850	74.1
DIK2768	77.9
ILSTS027	83.4
BMS812	84.9
BMS2076	91.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 77.9 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers 77.9 and 109.8. The at least one genetic marker is selected from the group of markers shown in Table 11d:

Table 11d

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2768	77.9
ILSTS027	83.4
BMS812	84.9
BMS2076	91.8
BL1095	94.8
BMS820	98.2
BMS927	105.0
BMS429	109.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 84.9 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS812 and BMS429. The at least one genetic marker is selected from the group of markers shown in Table 11e:

Table 11e

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS812	84.9
BMS2076	91.8
BL1095	94.8
BMS820	98.2
BMS927	105.0
BMS429	109.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 84.9 cM to about 94.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS812 and BL1095. The at least one genetic marker is selected from the group of markers shown in Table 11f:

Table 11f

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS812	84.9
BMS2076	91.8
BL1095	94.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 91.8 cM to about 105.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS2076 and BMS927. The at least one genetic marker is selected from the group of markers shown in Table 11g:

Table 11g

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2076	91.8
BL1095	94.8
BMS820	98.2
BMS927	105.0

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 98.2 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS820 and BMS429. The at least one genetic marker is selected from the group of markers shown in Table 11h:

Table 11h

Marker on BTA15	Relative position (cM)

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS820	98.2
BMS927	105.0
BMS429	109.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 105.0 cM to about 109.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA15. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA15 in the region flanked by and including the markers BMS927 and BMS429. The at least one genetic marker is selected from the group of markers shown in Table 11i:

Table 11i

Marker on BTA15	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS927	105.0
BMS429	109.8

## 10 BTA18

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA18. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 84.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment

15 the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers IDVGA-31 and DIK4013. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one

20 genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table12a:

Table 12a

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-31	0.0
BMS1355	2.9
BMS1322	13.5

TEXAN-10	20.7
BMS2213	24.5
INRA121	30.2
BR4406	33.4
BMS2554	40.2
MNB-27	44.0
BM7109	47.0
INRA063	48.0
ILSTS002	54.7
BMS2639	55.5
DIK4960	56.3
DIK4849	57.0
BMON117	57.6
DIK4232	61.2
BMS2785	72.0
DIK4569	73.8
BM2078	76.8
BM6507	78.8
TGLA227	84.1
DIK4013	84.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 13.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA18 in the region flanked by and including the markers IDVGA-31 and BMS1322. The at least one genetic marker is selected from the group of markers shown in Table 12b:

Table 12b

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-31	0.0
BMS1355	2.9
BMS1322	13.5

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 2.9 cM to about 13.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BMS1355 and BMS1322. The at least one genetic marker is selected from the group of markers shown in Table 12c:

Table 12c

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1355	2.9
BMS1322	13.5

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.2 cM to about 61.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers INRA121 and DIK4232. The at least one genetic marker is selected from the group of markers shown in Table 12d:

Table 12d

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA121	30.2
BR4406	33.4
BMS2554	40.2
MNB-27	44.0
BM7109	47.0
INRA063	48.0
ILSTS002	54.7
BMS2639	55.5
DIK4960	56.3
DIK4849	57.0
BMON117	57.6
DIK4232	61.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 33.4 cM to about 54.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BR4406 and ILSTS002. The at least one genetic marker is selected from the group of markers shown in Table 12e:

Table 12e

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BR4406	33.4
BMS2554	40.2
MNB-27	44.0
BM7109	47.0
INRA063	48.0
ILSTS002	54.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 57.6 cM to about 84.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BMON117 and DIK4013. The at least one genetic marker is selected from the group of markers shown in Table 12f:

Table 12f

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMON117	57.6
DIK4232	61.2
BMS2785	72.0
DIK4569	73.8
BM2078	76.8
BM6507	78.8
TGLA227	84.1
DIK4013	84.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 61.2 cM to about 84.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers DIK4232 and DIK4013. The at least one genetic marker is selected from the group of markers shown in Table 12g:

Table 12g

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK4232	61.2
BMS2785	72.0
DIK4569	73.8
BM2078	76.8
BM6507	78.8
TGLA227	84.1
DIK4013	84.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 72.0 cM to about 76.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BMS2785 and BM2078. The at least one genetic marker is selected from the group of markers shown in Table 12h:

Table 12h

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2785	72.0
DIK4569	73.8
BM2078	76.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 76.8 cM to about 84.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the

20

markers BM2078 and DIK4013. The at least one genetic marker is selected from the group of markers shown in Table 12i:

Table 12i

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM2078	76.8
BM6507	78.8
TGLA227	84.1
DIK4013	84.4

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 76.8 cM to about 78.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BM2078 and BM6507. The at least one genetic marker is selected from the group of markers shown in Table 12j:

10

Table 12j

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM2078	76.8
BM6507	78.8

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 78.8 cM to about 84.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA18. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA18 in the region flanked by and including the markers BM6507 and DIK4013. The at least one genetic marker is selected from the group of markers shown in Table 12k:

Table 12k

Marker on BTA18	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM6507	78.8
TGLA227	84.1
DIK4013	84.4

## BTA19

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA19. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 108.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers BM9202 and BMS601. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table13a:

Table 13a

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM9202	0.0
BMS745	16.0
BP20	45.9
IDVGA-46	47.0
BMS2389	52.2
CSSM065	69.8
ETH3	90.0
BMS601	108.0

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 90.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers BM9202 and ETH3. The at least one genetic marker is selected from the group of markers shown in Table 13b:

Table 13b

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM9202	0.0
BMS745	16.0

BP20	45.9
IDVGA-46	47.0
BMS2389	52.2
CSSM065	69.8
ETH3	90.0

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 45.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers BM9202 and BP20. The at least one genetic marker is selected from the group of markers shown in Table 13c:

Table 13c

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM9202	0.0
BMS745	16.0
BP20	45.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 16.0 cM to about 45.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers BMS745 and BP20. The at least one genetic marker is selected from the group of markers shown in Table 13d:

Table 13d

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS745	16.0
BP20	45.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 47.0 cM to about 90.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the

markers IDVGA-46 and ETH3. The at least one genetic marker is selected from the group of markers shown in Table 13e:

Table 13e

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-46	47.0
BMS2389	52.2
CSSM065	69.8
ETH3	90.0

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 52.2 cM to about 108.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers BMS2389 and BMS601. The at least one genetic marker is selected from the group of markers shown in Table 13f:

Table 13f

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2389	52.2
CSSM065	69.8
ETH3	90.0
BMS601	108.0

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 69.8 cM to about 90.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA19. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA19 in the region flanked by and including the markers CSSM065 and ETH3. The at least one genetic marker is selected from the group of markers shown in Table 13g:

Table 13g

Marker on BTA19	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
CSSM065	69.8

ETH3	90.0
------	------

## BTA20

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA20. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 77.1 cM (http://www.marc.usda.gov/) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers BM3517 and UWCA26. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table14a:

Table 14a

Marker on BTA20	Relative position (cM) http://www.marc.usda.gov/
BM3517	0.0
HEL12	0.6
BMS1282	19.1
BMS1754	26.3
TGLA126	31.9
BMS2361	49.7
AGLA29	55.1
BMS703	60.1
BM5004	71.8
UWCA26	77.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 71.8 cM (http://www.marc.usda.gov/) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers BM3517 and BM5004. The at least one genetic marker is selected from the group of markers shown in Table 14b:

Table 14b

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM3517	0.0
HEL12	0.6
BMS1282	19.1
BMS1754	26.3
TGLA126	31.9
BMS2361	49.7
AGLA29	55.1
BMS703	60.1
BM5004	71.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.0 cM to about 26.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA20 in the region flanked by and including the markers BM3517 and BMS1754. The at least one genetic marker is selected from the group of markers shown in Table 14c:

Table 14c

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM3517	0.0
HEL12	0.6
BMS1282	19.1
BMS1754	26.3

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 0.6 cM to about 19.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers HEL12 and BMS1282. The at least one genetic marker is selected from the group of markers shown in Table 14d:

Table 14d

Marker on BTA20	Relative position (cM)

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
HEL12	0.6
BMS1282	19.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 19.1 cM to about 55.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA20 in the region flanked by and including the markers BMS1282 and AGLA29. The at least one genetic marker is selected from the group of markers shown in Table 14e:

Table 14e

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1282	19.1
BMS1754	26.3
TGLA126	31.9
BMS2361	49.7
AGLA29	55.1

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 31.9 cM to about 49.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers TGLA126 and BMS2361. The at least one genetic marker is selected from the group of markers shown in Table 14f:

Table 14f

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
TGLA126	31.9
BMS2361	49.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 49.7 cM to about 55.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is

20 located on the bovine chromosome BTA20 in the region flanked by and including the

markers BMS2361 and AGLA29. The at least one genetic marker is selected from the group of markers shown in Table 14g:

Table 14g

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2361	49.7
AGLA29	55.1

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 55.1 cM to about 77.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers AGLA29 and UWCA26. The at least one genetic marker is selected from the group of markers shown in Table 14h:

Table 14h

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
AGLA29	55.1
BMS703	60.1
BM5004	71.8
UWCA26	77.1

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 60.1 cM to about 71.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA20. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA20 in the region flanked by and including the markers BMS703 and BM5004. The at least one genetic marker is selected from the group of markers shown in Table 14i:

Table 14i

Marker on BTA20	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS703	60.1
BM5004	71.8

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA21. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 5.6 cM to about 76.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment 5 the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the markers DIK5182 and IDVGA-30. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one 10 genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table15a:

Table 15a

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5182	5.5
BMS1117	11.0
RM151	12.6
DIK2492	18.3
AGLA233	21.2
ILSTS095	23.7
DIK4602	24.3
BM103	29.8
DIK4001	30.0
IDVGA-45	30.9
DIK2481	33.7
INRA103	35.9
BMS2815	41.7
DIK2842	41.7
DIK3036	47.8
DIK4391	52.1
DIK2913	57.1
BM846	61.2
TGLA122	62.7
ILSTS054	65.8

BMS743	75.3
IDVGA-30	76.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 11.0 cM to about 61.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the markers BMS1117 and BM846. The at least one genetic marker is selected from the group of markers shown in Table 15b:

Table 15b

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1117	11.0
RM151	12.6
DIK2492	18.3
AGLA233	21.2
ILSTS095	23.7
DIK4602	24.3
BM103	29.8
DIK4001	30.0
IDVGA-45	30.9
DIK2481	33.7
INRA103	35.9
BMS2815	41.7
DIK2842	41.7
DIK3036	47.8
DIK4391	52.1
DIK2913	57.1
BM846	61.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 18.3 cM to about 57.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the

markers DIK2492 and DIK2913. The at least one genetic marker is selected from the group of markers shown in Table 15c:

Table 15c

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2492	18.3
AGLA233	21.2
ILSTS095	23.7
DIK4602	24.3
BM103	29.8
DIK4001	30.0
IDVGA-45	30.9
DIK2481	33.7
INRA103	35.9
BMS2815	41.7
DIK2842	41.7
DIK3036	47.8
DIK4391	52.1
DIK2913	57.1

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 18.3 cM to about 30.0 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the markers DIK2492 and DIK4001. The at least one genetic marker is selected from the group of markers shown in Table 15d:

Table 15d

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2492	18.3
AGLA233	21.2
ILSTS095	23.7
DIK4602	24.3
BM103	29.8

DIK4001	30.0
---------	------

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 30.9 cM to about 47.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the markers IDVGA-45 and DIK3036. The at least one genetic marker is selected from the group of markers shown in Table 15e:

Table 15e

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-45	30.9
DIK2481	33.7
INRA103	35.9
BMS2815	41.7
DIK2842	41.7
DIK3036	47.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 33.7 cM to about 41.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the markers DIK2481 and BMS2815. The at least one genetic marker is selected from the group of markers shown in Table 15f:

Table 15f

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2481	33.7
INRA103	35.9
BMS2815	41.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 5.5 cM to about 61.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA21. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA21 in the region flanked by and including the

markers DIK5182 and BM846. The at least one genetic marker is selected from the group of markers shown in Table 15g:

Table 15g

Marker on BTA21	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK5182	5.5
DIK3036	47.8
DIK4391	52.1
DIK2913	57.1
BM846	61.2

## 5 BTA22

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA22. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 0.0 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment

10 the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the markers CSSM26 and BM4102. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one

15 genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table16a:

Table 16a

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
CSSM26	0.0
INRA026	2.9
BM1558	19.1
BM3628	47.1
BMS875	64.1
BM4102	82.9

In a preferred embodiment of the invention, the at least one genetic marker is located

20 in the region from about 2.9 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the

bovine chromosome BTA22. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the markers INRA026 and BM4102. The at least one genetic marker is selected from the group of markers shown in Table 16b:

5

Table 16b

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA026	2.9
BM1558	19.1
BM3628	47.1
BMS875	64.1
BM4102	82.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 2.9 cM to about 47.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment the at least one genetic marker is

10

located on the bovine chromosome BTA22 in the region flanked by and including the markers INRA026 and BM3628. The at least one genetic marker is selected from the group of markers shown in Table 16c:

Table 16c

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA026	2.9
BM1558	19.1
BM3628	47.1

15

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 19.1 cM to about 47.1 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the markers BM1558 and BM3628. The at least one genetic marker is selected from the group of markers shown in Table 16d:

20

Table 16d

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>

BM1558	19.1
BM3628	47.1

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 19.1 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the markers BM1558 and BM4102. The at least one genetic marker is selected from the group of markers shown in Table 16e:

Table 16e

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM1558	19.1
BM3628	47.1
BMS875	64.1
BM4102	82.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 47.1 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the markers BM3628 and BM4102. The at least one genetic marker is selected from the group of markers shown in Table 16f:

Table 16f

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM3628	47.1
BMS875	64.1
BM4102	82.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 64.1 cM to about 82.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA22. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA22 in the region flanked by and including the

markers BMS875 and BM4102. The at least one genetic marker is selected from the group of markers shown in Table 16g:

Table 16g

Marker on BTA22	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS875	64.1
BM4102	82.9

## 5 BTA24

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA24. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 6.2 cM to about 65.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment 10 the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BMS917 and BMS3024. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one 15 genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table17a:

Table 17a

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS917	6.2
BM7151	8.2
BM226	8.2
BMS2526	8.2
TGLA351	11.1
BM7228	19.3
CSSM23	20.6
BMS2270	23.7
ILSTS065	27.4
BMS1862	35.5
BMS466	48.8

INRA090	56.3
BMS1926	61.2
BMS3024	65.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 8.2 cM to about 65.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BM7151 and BMS3024. The at least one genetic marker is selected from the group of markers shown in Table 17b:

Table 17b

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM7151	8.2
BM226	8.2
BMS2526	8.2
TGLA351	11.1
BM7228	19.3
CSSM23	20.6
BMS2270	23.7
ILSTS065	27.4
BMS1862	35.5
BMS466	48.8
INRA090	56.3
BMS1926	61.2
BMS3024	65.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 8.2 cM to about 35.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BM7151 and BMS1862. The at least one genetic marker is selected from the group of markers shown in Table 17c:

Table 17c

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM7151	8.2
BM226	8.2
BMS2526	8.2
TGLA351	11.1
BM7228	19.3
CSSM23	20.6
BMS2270	23.7
ILSTS065	27.4
BMS1862	35.5

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 11.1 cM to about 23.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA24 in the region flanked by and including the markers TGLA351 and BMS2270. The at least one genetic marker is selected from the group of markers shown in Table 17d:

Table 17d

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
TGLA351	11.1
BM7228	19.3
CSSM23	20.6
BMS2270	23.7

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 35.5 cM to about 65.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BMS1862 and BMS3024. The at least one genetic marker is selected from the group of markers shown in Table 17e:

15

Table 17e

Marker on BTA24	Relative position (cM)

	<a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1862	35.5
BMS466	48.8
INRA090	56.3
BMS1926	61.2
BMS3024	65.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 48.8 cM to about 61.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BMS466 and BMS1926. The at least one genetic marker is selected from the group of markers shown in Table 17f:

Table 17f

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS466	48.8
INRA090	56.3
BMS1926	61.2

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 48.8 cM to about 56.3 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA24 in the region flanked by and including the markers BMS466 and INRA090. The at least one genetic marker is selected from the group of markers shown in Table 17g:

Table 17g

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS466	48.8
INRA090	56.3

20 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 56.3 cM to about 61.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA24. In one embodiment the at least one genetic marker is

located on the bovine chromosome BTA24 in the region flanked by and including the markers INRA090 and BMS1926. The at least one genetic marker is selected from the group of markers shown in Table 17h:

Table 17h

Marker on BTA24	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
INRA090	56.3
BMS1926	61.2

5

## BTA25

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA25. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 7.2 cM to about 61.7 cM

10 (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the markers ILSTS102 and AF5. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table18a:

Table 18a

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ILSTS102	7.2
BMS2843	22.6
BM737	31.6
ILSTS046	33.3
BMS1353	46.4
AF5	61.7

20 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 7.2 cM to about 31.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the

markers ILSTS102 and BM737. The at least one genetic marker is selected from the group of markers shown in Table 18b:

Table 18b

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ILSTS102	7.2
BMS2843	22.6
BM737	31.6

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 7.2 cM to about 22.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the markers ILSTS102 and BMS2843. The at least one genetic marker is selected from the 10 group of markers shown in Table 18c:

Table 18c

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ILSTS102	7.2
BMS2843	22.6

15 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 31.6 cM to about 61.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the markers BM737 and AF5. The at least one genetic marker is selected from the group of markers shown in Table 18d:

Table 18d

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BM737	31.6
ILSTS046	33.3
BMS1353	46.4
AF5	61.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 33.3 cM to about 46.4 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the markers ILSTS046 and BMS1353. The at least one genetic marker is selected from the group of markers shown in Table 18e:

Table 18e

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ILSTS046	33.3
BMS1353	46.4

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 46.4 cM to about 61.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA25. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA25 in the region flanked by and including the markers BMS1353 and AF5. The at least one genetic marker is selected from the group of markers shown in Table 18f:

Table 18f

Marker on BTA25	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1353	46.4
AF5	61.7

## BTA26

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA26. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 2.8 cM to about 66.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers BMS651 and BM7237. The at least one genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one

genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table 19a:

Table 19a

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS651	2.8
HEL11	22.9
BMS332	31.7
RM026	37.6
BM9284	41.6
RME40	43.2
IDVGA-59	53.1
BMS882	53.5
BM804	60.5
BM7237	66.8

5 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 2.8 cM to about 60.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers BMS651 and BM804. The at least one genetic marker is selected from the 10 group of markers shown in Table 19b:

Table 19b

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS651	2.8
HEL11	22.9
BMS332	31.7
RM026	37.6
BM9284	41.6
RME40	43.2
IDVGA-59	53.1
BMS882	53.5
BM804	60.5

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 2.8 cM to about 37.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers BMS651 and RM026. The at least one genetic marker is selected from the group of markers shown in Table 19c:

Table 19c

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS651	2.8
HEL11	22.9
BMS332	31.7
RM026	37.6

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 22.9 cM to about 31.7 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers HEL11 and BMS332. The at least one genetic marker is selected from the group of markers shown in Table 19d:

Table 19d

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
HEL11	22.9
BMS332	31.7

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 31.7 cM to about 41.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers BMS332 and BM9284. The at least one genetic marker is selected from the group of markers shown in Table 19e:

Table 19e

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>

BMS332	31.7
RM026	37.6
BM9284	41.6

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 37.6 cM to about 66.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is

5 located on the bovine chromosome BTA26 in the region flanked by and including the markers RM026 and BM7237. The at least one genetic marker is selected from the group of markers shown in Table 19f:

Table 19f

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
RM026	37.6
BM9284	41.6
RME40	43.2
IDVGA-59	53.1
BMS882	53.5
BM804	60.5
BM7237	66.8

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 37.6 cM to about 43.2 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers RM026 and RME40. The at least one genetic marker is selected from the

15 group of markers shown in Table 19g:

Table 19g

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
RM026	37.6
BM9284	41.6
RME40	43.2

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 43.2 cM to about 66.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers RME40 and BM7237. The at least one genetic marker is selected from the group of markers shown in Table 19h:

Table 19h

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
RME40	43.2
IDVGA-59	53.1
BMS882	53.5
BM804	60.5
BM7237	66.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 53.1 cM to about 60.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA26. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA26 in the region flanked by and including the markers IDVGA-59 and BM804. The at least one genetic marker is selected from the group of markers shown in Table 19i:

Table 19i

Marker on BTA26	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
IDVGA-59	53.1
BMS882	53.5
BM804	60.5

## BTA28

In one embodiment of the invention the at least one genetic marker is located on the bovine chromosome BTA28. In one specific embodiment of the present invention, the at least one genetic marker is located in the region from about 8.0 cM to about 59.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers BMC6020 and BMC2208. The at least one

genetic marker is significant for the calving traits SB, CD and/or CS. In a particular embodiment the at least one genetic marker is significant for for example the trait SB, such as CD, for example CS. However, in a further embodiment the at least one genetic marker is significant for the traits in any combination. The at least one genetic marker is selected from the group of markers shown in Table20a:

Table 20a

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMC6020	8.0
ETH1112	16.9
BL25	24.8
DIK2955	38.0
BMS2608	38.5
BMS2658	43.0
DIK713	45.9
BMS1714	49.4
DIK5056	50.5
DIK5323	55.9
DIK4862	59.6
BMC2208	59.6

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 8.0 cM to about 24.8 cM (<http://www.marc.usda.gov/>) on the

10 bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers BMC6020 and BL25. The at least one genetic marker is selected from the group of markers shown in Table 20b:

Table 20b

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMC6020	8.0
ETH1112	16.9
BL25	24.8

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 16.9 cM to about 24.8 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers ETH1112 and BL25. The at least one genetic marker is selected from the group of markers shown in Table 20c:

5 Table 20c

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
ETH1112	16.9
BL25	24.8

In a preferred embodiment of the invention, the at least one genetic marker is located 10 in the region from about 24.8 cM to about 50.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers BL25 and DIK5056. The at least one genetic marker is selected from the group of markers shown in Table 20d:

15 Table 20d

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BL25	24.8
DIK2955	38.0
BMS2608	38.5
BMS2658	43.0
DIK713	45.9
BMS1714	49.4
DIK5056	50.5

In a preferred embodiment of the invention, the at least one genetic marker is located 20 in the region from about 38.0 cM to about 45.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers DIK2955 and DIK713. The at least one genetic marker is selected from the group of markers shown in Table 20e:

Table 20e

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2955	38.0
BMS2608	38.5
BMS2658	43.0
DIK713	45.9

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 38.0 cM to about 43.0 cM (<http://www.marc.usda.gov/>) on the 5 bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers DIK2955 and BMS2658. The at least one genetic marker is selected from the group of markers shown in Table 20f:

Table 20f

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK2955	38.0
BMS2608	38.5
BMS2658	43.0

10

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 43.0 cM to about 59.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the 15 markers BMS2658 and BMC2208. The at least one genetic marker is selected from the group of markers shown in Table 20g:

Table 20g

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS2658	43.0
DIK713	45.9
BMS1714	49.4
DIK5056	50.5

DIK5323	55.9
DIK4862	59.6
BMC2208	59.6

In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 45.9 cM to about 55.9 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers DIK713 and DIK5323. The at least one genetic marker is selected from the group of markers shown in Table 20h:

Table 20h

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
DIK713	45.9
BMS1714	49.4
DIK5056	50.5
DIK5323	55.9

10 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 49.4 cM to about 50.5 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the markers BMS1714 and DIK5056. The at least one genetic marker is selected from the group of markers shown in Table 20i:

Table 20i

Marker on BTA28	Relative position (cM) <a href="http://www.marc.usda.gov/">http://www.marc.usda.gov/</a>
BMS1714	49.4
DIK5056	50.5

20 In a preferred embodiment of the invention, the at least one genetic marker is located in the region from about 55.9 cM to about 59.6 cM (<http://www.marc.usda.gov/>) on the bovine chromosome BTA28. In one embodiment the at least one genetic marker is located on the bovine chromosome BTA28 in the region flanked by and including the



21	X	X	X										
22	X	X											X
23	X			X									X
24	X									X			X

Table 20k2.

Table 20k3.

Embedment	BTA																		
	3	4	5	7	8	9	10	11	12	15	18	19	20	21	22	24	25	26	28
1			X	X															
2			X		X														
3			X			X													
4			X				X												
5			X					X											
6			X						X										
7			X							X									
8			X								X								
9			X									X							
10			X										X						
11			X											X					
12			X												X				
13			X													X			
14			X														X		

Table 20k4.

Embedment	BTA																		
	3	4	5	7	8	9	1	1	1	1	1	2	2	2	2	2	2	2	2
1			X	X															
2			X		X														
3			X			X													
4			X				X												
5			X					X											
6			X						X										
7			X							X									
8			X								X								
9			X									X							
10			X										X						
11			X											X					
12			X												X				
13			X													X			
14			X																X
15		X	X	X		X		X		X						X		X	X
16		X	X		X	X	X		X		X		X		X		X	X	
17		X	X		X		X				X		X		X	X			
18			X		X		X		X		X		X				X		X
19			X	X	X							X							X
20			X	X	X							X							
21			X	X	X														
22			X	X								X							
23				X	X							X							X
24				X								X							X

Table 20k5.

Embodiment	BTA																	
	3	4	5	7	8	9	1	1	1	1	1	2	2	2	2	2	2	2
1					X	X												
2					X		X											
3					X			X										
4					X				X									
5					X					X								
6					X						X							
7					X							X						
8					X								X					

Table 20k6.

Table 20k7.

3						X			X																
4						X			X																
5						X				X															
6						X					X														
7						X						X													
8						X							X												
9						X								X											
10						X									X										
11						X										X					X				
12						X																X			
13	X	X	X	X	X		X	X							X	X	X	X	X	X	X	X	X		
14		X	X		X	X	X		X		X		X		X		X	X	X						
15	X				X	X	X				X		X		X		X	X	X						
16		X	X	X	X	X		X		X		X				X		X		X	X	X			
17	X				X	X		X		X		X				X									
18						X																			
19	X	X				X				X														X	
20	X	X				X				X															
21	X	X				X																			
22	X						X			X															
23					X		X			X														X	
24						X			X															X	

Table 20k8.

Embody ment	BTA																							
	3	4	5	7	8	9	1 0	1 1	1 2	1 5	1 8	1 9	2 0	2 1	2 1	2 2	2 4	2 5	2 6	2 8				
1							X	X																
2							X		X															
3							X				X													
4							X					X												
5							X						X											
6							X							X										
7							X									X								
8							X										X							
9							X											X						
10							X												X					
11								X												X				
12	X	X	X	X	X	X	X		X							X	X	X	X	X	X	X		
13		X	X		X	X	X		X		X		X		X		X	X	X					
14	X				X	X	X				X		X		X		X	X	X					
15			X	X	X	X	X		X		X		X					X	X	X				
16	X				X	X	X	X	X		X		X		X									
17		X	X			X							X								X			
18			X	X		X	X		X		X		X		X		X	X	X					
19	X	X				X					X												X	
20	X	X				X					X													
21	X	X				X																		
22	X						X				X													
23					X		X				X												X	
24							X			X		X											X	

Table 20k9.

Table 20k10.



16	X	X								X	X								
17		X	X							X	X								X
18		X									X								
19		X	X							X	X								
20		X	X							X									X
21		X								X	X								X
22		X								X									X
23				X						X	X								
24	X									X	X								X

Table 20k13.

Embodiment	BTA																		
	3	4	5	7	8	9	1 0	1 1	1 2	1 5	1 8	1 9	2 0	2 1	2 2	2 4	2 5	2 6	2 8
1														X	X				
2													X		X				
3													X		X				
4													X			X			
5													X				X		
6													X						X
7		X		X			X		X				X		X	X	X		X
8	X		X			X		X	X			X	X	X	X		X		
9	X			X		X		X	X	X		X							X
10	X	X			X		X		X				X	X		X		X	
11	X	X	X										X	X					X
12	X	X	X										X	X					
13	X	X	X										X	X					
14	X	X											X	X					
15		X	X										X	X					X
16		X											X	X					
17		X	X										X						
18		X	X										X						X
19		X											X						X
20		X											X	X					X
21				X									X						
22	X												X	X					X
23				X									X						X
24													X						X

Table 20k14.

Embodiment	BTA																		
	3	4	5	7	8	9	1 0	1 1	1 2	1 5	1 8	1 9	2 0	2 1	2 2	2 4	2 5	2 6	2 8
1															X	X			
2														X		X			
3														X			X		
4														X				X	
5														X					X
6		X		X			X		X	X			X		X	X	X		X
7	X		X			X		X	X	X			X	X	X	X		X	
8			X			X		X	X	X			X						X
9				X		X							X		X	X	X	X	X

10		X	X		X	X		X		X	X	X									
11	X		X		X				X		X									X	
12	X		X		X				X		X										
13	X		X		X						X										
14	X		X							X		X									
15			X		X				X		X									X	
16		X										X									
17		X		X					X		X										
18		X		X							X		X							X	
19		X								X		X								X	
20		X									X			X						X	
21				X					X		X										
22	X									X		X								X	
23	X									X		X								X	
24									X		X									X	

Table 20k15.

Embodiment	BTA																											
	3	4	5	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
1																					X	X						
2																				X		X						
3																				X			X					
4																				X				X				
5		X			X			X		X						X		X	X	X	X					X		
6	X		X			X		X	X	X				X	X	X	X	X	X							X		
7		X			X		X	X	X	X				X			X									X		
8			X	X							X					X		X	X	X						X		
9		X	X			X		X				X						X	X	X						X		
10	X		X	X								X						X									X	
11	X		X	X								X						X										
12	X		X	X															X									
13	X		X									X						X										
14		X		X								X					X										X	
15		X																	X									
16		X		X								X					X											
17		X	X																X			X						
18		X										X					X			X							X	
19		X																	X			X					X	
20				X								X					X			X								
21	X											X					X			X							X	
22	X											X					X			X							X	
23												X					X			X							X	
24												X					X			X							X	

Table 20k16.

Embodiment	BTA																										
	3	4	5	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
1																				X	X						
2																			X		X						
3																			X			X					

4		X		X		X		X		X		X
5		X		X		X		X	X	X		X
6			X		X		X	X	X	X		X
7				X	X				X		X	X
8		X	X		X	X		X			X	X
9		X	X	X				X			X	X
10		X	X	X				X			X	
11		X	X	X							X	
12		X	X					X			X	
13			X	X				X			X	X
14			X								X	
15			X	X				X			X	
16			X	X							X	X
17			X					X			X	X
18			X								X	X
19				X				X			X	
20		X						X			X	X
21		X						X			X	X
22								X			X	X
23								X			X	X
24								X			X	X

Table 20k17.

Table 20k18.

Embod BTA

Embodyment	3	4	5	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
							0	1	2	5	8	9	0	1	2	2	2	2	2	2	
1																				X	X
2	X		X		X						X									X	
3	X		X		X						X									X	
4	X		X		X															X	
5	X		X								X									X	
6			X		X						X									X	
7			X		X															X	
8			X								X									X	
9			X		X															X	
10			X								X									X	
11			X																	X	
12					X						X									X	
13	X										X									X	
14	X				X						X									X	
15	X		X		X		X		X	X	X	X	X	X	X	X	X	X	X	X	X
16	X																			X	
17					X			X		X	X				X		X	X	X	X	
18		X	X			X			X		X	X				X	X	X	X	X	
19		X	X								X									X	
20		X	X								X									X	
21		X	X		X	X					X									X	X
22	X	X									X									X	
23						X					X		X		X					X	
24											X		X		X					X	

Table 20k19.

Embodyment	BTA																				
	3	4	5	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
							0	1	2	5	8	9	0	1	2	2	2	2	2	2	
1		X			X			X		X										X	X
2	X			X			X		X	X	X			X	X	X					X
3			X			X		X	X	X	X			X							X
4				X	X									X				X			X
5		X	X			X		X												X	X
6	X	X	X											X						X	X
7	X	X	X											X						X	
8	X	X	X		X	X								X	X	X	X				X
9	X	X												X							X
10		X	X											X		X				X	X
11		X														X					X
12		X	X											X							X
13		X	X													X	X			X	X
14		X												X						X	X
15		X														X	X			X	X
16				X										X							X
17	X													X				X		X	X
18	X													X						X	X
19		X	X	X										X						X	X
20					X									X		X				X	X
21		X	X	X	X									X						X	X
22	X	X												X							X

23	X			X					X				X	
24	X								X				X	

### Detection

The detection of the presence or absence of a genetic marker allele according to the

5 present invention may be conducted on the DNA sequence of the bovine chromosomes BTA3, BTA4, BTA5, BTA7, BTA8, BTA9, BTA10, BTA11, BTA12, BTA15, BTA18, BTA19, BTA20, BTA21, BTA22, BTA24, BTA25, BTA26, and/or BTA28 specified elsewhere herein according to the present invention or a complementary sequence as well as on transcriptional (mRNA) and translational

10 products (polypeptides, proteins) therefrom.

It will be apparent to the person skilled in the art that there are a large number of analytical procedures which may be used to detect the presence or absence of variant nucleotides at one or more of positions mentioned herein in the specified region.

15 Mutations or polymorphisms within or flanking the specified region can be detected by utilizing a number of techniques. Nucleic acid from any nucleated cell can be used as the starting point for such assay techniques, and may be isolated according to standard nucleic acid preparation procedures that are well known to those of skill in the art. In general, the detection of allelic variation requires a mutation discrimination technique,

20 optionally an amplification reaction and a signal generation system.

A number of mutation detection techniques are listed in Table 21. Some of the methods listed in Table 21 are based on the polymerase chain reaction (PCR), wherein the

25 method according to the present invention includes a step for amplification of the nucleotide sequence of interest in the presence of primers based on the nucleotide sequence of the variable nucleotide sequence. The methods may be used in combination with a number of signal generation systems, a selection of which is also listed in Table 22.

Table 21

General techniques	DNA sequencing, Sequencing by hybridisation, SNAPshot
Scanning techniques	Single-strand conformation polymorphism analysis, Denaturing gradient gel electrophoresis, Temperature gradient gel electrophoresis, Chemical mismatch

	cleavage, cleavage, heteroduplex analysis, enzymatic mismatch cleavage
Hybridisation based techniques	<p>Solid phase hybridisation: Dot blots, Multiple allele specific diagnostic assay (MASDA), Reverse dot blots, Oligonucleotide arrays (DNA Chips)</p> <p>Solution phase hybridisation: Taqman -U.S. Pat. No. 5,210,015 &amp; 5,487,972 (Hoffmann-La Roche), Molecular Beacons -- Tyagi et al (1996), Nature Biotechnology, 14, 303; WO 95/13399 (Public Health Inst., New York), Lightcycler, optionally in combination with Fluorescence resonance energy transfer (FRET).</p>
Extension based techniques	Amplification refractory mutation system (ARMS), Amplification refractory mutation system linear extension (ALEX) - European Patent No. EP 332435 B1 (Zeneca Limited), Competitive oligonucleotide priming system (COPS) - Gibbs et al (1989), Nucleic Acids Research, 17, 2347.
Incorporation based techniques	Mini-sequencing, Arrayed primer extension (APEX)
Restriction Enzyme based techniques	Restriction fragment length polymorphism (RFLP), Restriction site generating PCR
Ligation based techniques	Oligonucleotide ligation assay (OLA)
Other	Invader assay
Various Signal Generation or Detection Systems	<p>Fluorescence:</p> <p>Fluorescence resonance energy transfer (FRET),</p> <p>Fluorescence quenching, Fluorescence polarisation-- United Kingdom Patent No. 2228998 (Zeneca Limited)</p>
Other	Chemiluminescence, Electrochemiluminescence, Raman, Radioactivity, Colorimetric, Hybridisation protection assay, Mass spectrometry

Further amplification techniques are listed in Table 2. Many current methods for the detection of allelic variation are reviewed by Nollau et al., Clin. Chem. 43, 1114-1120, 1997; and in standard textbooks, for example "Laboratory Protocols for Mutation 5 Detection", Ed. by U. Landegren, Oxford University Press, 1996 and "PCR", 2nd Edition by Newton & Graham, BIOS Scientific Publishers Limited, 1997.

The detection of genetic markers can according to one embodiment of the present invention be achieved by a number of techniques known to the skilled person, including 10 typing of microsatellites or short tandem repeats (STR), restriction fragment length polymorphisms (RFLP), detection of deletions or insertions, random amplified polymorphic DNA (RAPIDs) or the typing of single nucleotide polymorphisms by methods such as restriction fragment length polymerase chain reaction, allele-specific 15 oligomer hybridisation, oligomer-specific ligation assays, hybridisation with PNA or locked nucleic acids (LNA) probes.

Table 22

Further amplification techniques	Self sustained replication (SSR), Nucleic acid sequence based amplification (NASBA), Ligase chain reaction (LCR), Strand displacement amplification (SDA)
----------------------------------	---

A primer of the present invention is a nucleic acid molecule sufficiently complementary 20 to the sequence on which it is based and of sufficiently length to selectively hybridise to the corresponding region of a nucleic acid molecule intended to be amplified. The primer is able to prime the synthesis of the corresponding region of the intended nucleic acid molecule in the methods described above. Similarly, a probe of the present 25 invention is a molecule for example a nucleic acid molecule of sufficient length and sufficiently complementary to the nucleic acid sequence of interest which selectively binds to the nucleic acid sequence of interest under high or low stringency conditions.

### Sample

The method according to the present invention includes analyzing a sample of a bovine 30 subject, wherein said sample may be any suitable sample capable of providing the

bovine genetic material for use in the method. The bovine genetic material may for example be extracted, isolated and purified if necessary from a blood sample, a tissue samples (for example spleen, buccal smears), clipping of a body surface (hairs or nails), milk and/or semen. The samples may be fresh or frozen.

5

The sequence polymorphisms of the invention comprise at least one nucleotide difference, such as at least two nucleotide differences, for example at least three nucleotide differences, such as at least four nucleotide differences, for example at least five nucleotide differences, such as at least six nucleotide differences, for example at least seven nucleotide differences, such as at least eight nucleotide differences, for example at least nine nucleotide differences, such as 10 nucleotide differences. The nucleotide differences comprise nucleotide differences, deletion and/or insertion or any combination thereof.

10

### 15 **Primers**

The primers that may be used according to the present invention are shown in Table 22. The in Table 22 specified primer pairs may be used individually or in combination with one or more primer pairs of Table 22.

20

The design of such primers or probes will be apparent to the molecular biologist of ordinary skill. Such primers are of any convenient length such as up to 50 bases, up to 40 bases, more conveniently up to 30 bases in length, such as for example 8-25 or 8-15 bases in length. In general such primers will comprise base sequences entirely complementary to the corresponding wild type or variant locus in the region. However, if required one or more mismatches may be introduced, provided that the discriminatory power of the oligonucleotide probe is not unduly affected. The primers/probes of the invention may carry one or more labels to facilitate detection.

25

In one embodiment, the primers and/or probes are capable of hybridizing to and/or amplifying a subsequence hybridizing to a single nucleotide polymorphism containing the sequence delineated by the markers as shown herein.

30

The primer nucleotide sequences of the invention further include: (a) any nucleotide sequence that hybridizes to a nucleic acid molecule of the delineated region(s) or its complementary sequence or RNA products under stringent conditions, e.g., hybridization to filter-bound DNA in 6x sodium chloride/sodium citrate (SSC) at about

35

45°C followed by one or more washes in 0.2x SSC/0.1% Sodium Dodecyl Sulfate (SDS) at about 50-65°C, or (b) under highly stringent conditions, e.g., hybridization to filter-bound nucleic acid in 6x SSC at about 45°C followed by one or more washes in 0.1x SSC/0.2% SDS at about 68°C, or under other hybridization conditions which are 5 apparent to those of skill in the art (see, for example, Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at pp. 6.3.1-6.3.6 and 2.10.3). Preferably the 10 nucleic acid molecule that hybridizes to the nucleotide sequence of (a) and (b), above, is one that comprises the complement of a nucleic acid molecule of the region s or r or a complementary sequence or RNA product thereof. In a preferred embodiment, nucleic acid molecules comprising the nucleotide sequences of (a) and (b), comprises nucleic acid molecule of RAI or a complementary sequence or RNA product thereof.

15 Among the nucleic acid molecules of the invention are deoxyoligonucleotides ("oligos") which hybridize under highly stringent or stringent conditions to the nucleic acid molecules described above. In general, for probes between 14 and 70 nucleotides in length the melting temperature (TM) is calculated using the formula:

$$T_m(\text{°C}) = 81.5 + 16.6(\log [\text{monovalent cations (molar)}]) + 0.41(\% \text{ G+C}) - (500/N)$$

20 where N is the length of the probe. If the hybridization is carried out in a solution containing formamide, the melting temperature is calculated using the equation  
$$T_m(\text{°C}) = 81.5 + 16.6(\log[\text{monovalent cations (molar)}]) + 0.41(\% \text{ G+C}) - (0.61\% \text{ formamide}) - (500/N)$$
 where N is the length of the probe. In general, hybridization is carried out at 25 about 20-25 degrees below Tm (for DNA-DNA hybrids) or 10-15 degrees below Tm (for RNA-DNA hybrids).

30 Exemplary highly stringent conditions may refer for example to washing in 6x SSC/0.05% sodium pyrophosphate at 37°C (for about 14-base oligos), 48°C (for about 17-base oligos), 55°C (for about 20-base oligos), and 60°C (for about 23-base oligos). Accordingly, the invention further provides nucleotide primers or probes which detect 35 the r region polymorphisms of the invention. The assessment may be conducted by means of at least one nucleic acid primer or probe, such as a primer or probe of DNA, RNA or a nucleic acid analogue such as peptide nucleic acid (PNA) or locked nucleic acid (LNA).

According to one aspect of the present invention there is provided an allele-specific oligonucleotide probe capable of detecting a polymorphism at one or more of positions in the delineated regions 1.

5 The allele-specific oligonucleotide probe is preferably 5-50 nucleotides, more preferably about 5-35 nucleotides, more preferably about 5-30 nucleotides, more preferably at least 9 nucleotides.

### **Determination of linkage**

In order to detect whether the genetic marker is present in the genetic material, 10 standard methods well known to persons skilled in the art may be applied, for example by the use of nucleic acid amplification. In order to determine whether the genetic marker is genetically linked to the calving traits, a permutation test can be applied when the regression method is used (Doerge and Churchill, 1996), or the Piepho-method can be applied (Piepho, 2001) when the variance components method is used. The 15 principle of the permutation test is well described by Doerge and Churchill (1996), whereas the Piepho-method is well described by Piepho (2001). Significant linkage in the within family analysis using the regression method, a 1000 permutations were made using the permutation test (Doerge and Churchill, 1996). A threshold at the 5% chromosome wide level was considered to be significant evidence for linkage between 20 the genetic marker and the calving traits. In addition, the QTL was confirmed in different sire families. For the across family analysis and multi-trait analysis with the variance component method the piepho method was used to determine the significance level (Piepho, 2001). A threshold at the 5% chromosome wide level was considered to be significant evidence for linkage between the genetic marker and the 25 calving traits.

### **Kit**

Another aspect of the present invention relates to a diagnostic kit for use in detecting 30 the presence or absence in a bovine subject of at least one genetic marker associated with bovine calving characteristics, comprising at least one oligonucleotide sequence, wherein the nucleotide sequences are selected from any of SEQ ID NO.: 1 to SEQ ID NO.: 558 and/or any combination thereof.

Genotyping of a bovine subject in order to establish the genetic determinants of calving traits for that subject according to the present invention can be based on the analysis of genomic DNA which can be provided using standard DNA extraction methods as described herein. The genomic DNA may be isolated and amplified using standard techniques such as the polymerase chain reaction using oligonucleotide primers corresponding (complementary) to the polymorphic marker regions. Additional steps of purifying the DNA prior to amplification reaction may be included. Thus, a diagnostic kit for establishing calving characteristics comprises, in a separate packing, at least one oligonucleotide sequence selected from the group of sequences shown in table 23 and any combinations thereof.

### Examples

#### Experimental design

A total genome scan for QTL affecting calving traits, was carried out in the Danish Holstein population. Marker and phenotypic data were collected according to the 15 granddaughter design (Weller et al., 1990), which included 34 sires with 2042 progeny-tested sons. Numbers of sons per sire ranged from 20 to 106. Sires and their sons were genotyped for marker information whereas phenotypic records were taken from granddaughter performances. Numbers of daughters of each son ranged between 70 20 and 100. The marker data set included a total of 384 microsatellites covering all 29 Bos Taurus chromosomes.

#### Purification of genomic DNA

Genomic DNA was purified from semen according to the following protocol:

25 After thawing the semen-straw, both ends of the straw were cut away with a pair of scissors and the content of semen transferred to a 1.5 ml eppendorf tube. 1 ml of 0.9% NaCl was used to flush the straw into the tube. The tube was then centrifuged for 5 minutes at 2000 rpm, followed by removal of the supernatant. This washing step was repeated twice.

30 Then 300 µl buffer S (10 mM Tris HCl pH 8, 100 mM NaCl, 10 mM EDTA pH 8; 0,5 % SDS), 20 µl 1 M DTT and 20 µl pronase (20 mg/ml) (Boehringer )are added to the tube. After mixing the tubes are incubated over night with slow rotation where after 180 µl saturated NaCl is added followed by vigorous agitation for 15 seconds. The tube is then centrifuged for 15 minutes at 11000 rpm. 0.4 ml of the supernatant is transferred to 35 a 2 ml tube and 1 ml of 96% ethanol is added, mixing is achieved by slow rotation of

the tube. The tube is then centrifuged for 10 minutes at 11000 rpm. Remove the supernatant by pouring away the liquid, wash the pellet with 70% ethanol (0.2 ml) and centrifuge again for 10 minutes at 11000 rpm. Pour away the ethanol, dry the pellet and resuspend in 0.5 ml of TE-buffer) for 30 minutes at 55°C.

5

#### Amplification procedures

PCR reactions were run in a volume of 8 µl using TEMPase (GeneChoice) polymerase and reaction buffer I as provided by the supplier (GeneChoice). Usually 5 different markers are included in each multiplex PCR. 1 µl DNA, 0.1 µl TEMPase enzyme, 0.2 mM dNTPs, 1.2 mM MgCl<sub>2</sub>, 0.3 µM each primer.

The PCR mixtures were subjected to initial denaturation at 94°C for 15 min (for TEMPase). Subsequently, the samples were cycled for 10 cycles with touchdown, i.e. the temperature is lowered 1°C at each cycle (denaturation at 94°C 30", annealing at 67°C 45", elongation 72°C 30"), after which the samples were cycled for 20 cycles with normal PCR conditions (denaturation at 94°C 30", annealing at 58°C 45", elongation 72°C 30") PCR cycling was terminated by 1 cycle at 72°C 30' and the PCR machine was programmed to cooling down the samples at 4°C for 'ever'.

20 The nucleotide sequence of the primers used for detecting the markers is shown in Table 23. The sequence is listed from the 5' end.

Table 23	Forward Primer F	SEQ ID NO.:
Marker name	Reverse Primer R	
<b>BTA3:</b>		
INRA006	<b>F</b> AGGAATATCTGTATCAACCTCAGTC	SEQ ID NO.: 1
	<b>R</b> CTGAGCTGGGTGGAGCTATAAATA	SEQ ID NO.: 2
UWCA7	<b>F</b> TGTAGCTCCCTGGAGGGAGAA	SEQ ID NO.: 3
	<b>R</b> GCAAATACAACCCAGTCTGGTG	SEQ ID NO.: 4
ILSTS096	<b>F</b> GTGACCTGGAGAAGTTTCC	SEQ ID NO.: 5
	<b>R</b> ACCACGCTCTGACTTGTAGC	SEQ ID NO.: 6
DIK4403	<b>F</b> CGTGCTGCAACTGAGAGTTC	SEQ ID NO.: 7
	<b>R</b> GCTGTATAGCAAAGTGACCCAGT	SEQ ID NO.: 8
RME23	<b>F</b> AGAACAAATGTGACACTCACA	SEQ ID NO.: 9
	<b>R</b> GTGAGTACAGGCGCTTCTG	SEQ ID NO.: 10
BMS963	<b>F</b> GGAGGATGAAGGAGTCTTG	SEQ ID NO.: 11
	<b>R</b> AATTTACCACAGTCCACCGC	SEQ ID NO.: 12

BMS819	<b>F</b> AAAGAATTGGACCTGACTGAGC <b>R</b> GCTTCACTCTGCTGGCTT	SEQ ID NO.: 13 SEQ ID NO.: 14
FCGR1	<b>F</b> GGTCTTCATTGGTGTTCCTCC <b>R</b> GAGCTGCCCTAGATGAGGTG	SEQ ID NO.: 15 SEQ ID NO.: 16
BL41	<b>F</b> CCTCTGCCATCTTATTCCG <b>R</b> AAGATCAACTTATTCCACAGTGG	SEQ ID NO.: 17 SEQ ID NO.: 18
DIK4353	<b>F</b> TGAACTTAGGGCAGCATGA <b>R</b> AAGACTGAGATGTGGGGAAAA	SEQ ID NO.: 19 SEQ ID NO.: 20
INRA003	<b>F</b> CTGGAGGTGTGTGAGCCCCATTAA <b>R</b> CTAAGAGTCGAAGGTGTGACTAGG	SEQ ID NO.: 21 SEQ ID NO.: 22
BMS2790	<b>F</b> AAGACAAGGACTTCAGCCC <b>R</b> AAAGAGTCGGACATTACTGAGC	SEQ ID NO.: 23 SEQ ID NO.: 24
ILSTS029	<b>F</b> TGTGTTGATGGAACACAGCC <b>R</b> TGGATTTAGACCAGGGTTGG	SEQ ID NO.: 25 SEQ ID NO.: 26
BM220	<b>F</b> TTTTCTACTGCCAACAAAGTG <b>R</b> TAGGTACCATAGCCTAGCCAAG	SEQ ID NO.: 27 SEQ ID NO.: 28
INRA123	<b>F</b> TCTAGAGGATCCCCGCTGAC <b>R</b> AGAGAGCAACTCCACTGTGC	SEQ ID NO.: 29 SEQ ID NO.: 30
BMS862	<b>F</b> TATAATGCCCTAGATCCACTCA <b>R</b> ATGGAAAAATAAGATGTGGTATGTG	SEQ ID NO.: 31 SEQ ID NO.: 32
HUJ246	<b>F</b> ACTCCAGTTTCTTCCTGG <b>R</b> TGCCATGTAGTAGCTGTGTGC	SEQ ID NO.: 33 SEQ ID NO.: 34
BMS937	<b>F</b> GTAGCCATGGAGACTGGACTG <b>R</b> CATTATCCCCGTACACACACC	SEQ ID NO.: 35 SEQ ID NO.: 36
DIK4664	<b>F</b> AACTGGCTCCAAGGTCAATG <b>R</b> TCCCCTGTACACACACCTGTA	SEQ ID NO.: 37 SEQ ID NO.: 38
DIK2702	<b>F</b> TGCGATATTAAATGGATGTCT <b>R</b> TTCCCTTCTCCGAACTGCTC	SEQ ID NO.: 39 SEQ ID NO.: 40
HUJI77	<b>F</b> TCCATCAAGTATTGAGTGCAA <b>R</b> ATAGCCCTACCCACTGTTCTG	SEQ ID NO.: 41 SEQ ID NO.: 42
DIK2686	<b>F</b> ATGTTTTCAGGCCAATCCA <b>R</b> TGCCCTGATTCTCATACCC	SEQ ID NO.: 43 SEQ ID NO.: 44
BM7225	<b>F</b> GGTGTTATGCATTCTCTAGGTGC <b>R</b> AAGAGTTAGACATGACTGAGCACG	SEQ ID NO.: 45 SEQ ID NO.: 46
<b>BTA4:</b>		
BMS1788	<b>F</b> ACGTCCAGATTCAAGATTCTTG <b>R</b> GGAGAGGAATCTTGCAAAGG	SEQ ID NO.: 47 SEQ ID NO.: 48
BMS2646	<b>F</b> CAAAGCCATAAGAAGCAATTATG	SEQ ID NO.: 49

BMS2646	R CCTTCTATAGTGTGGTACTACCC F GCACAGTAATAAGAGTGATGGCAGA	SEQ ID NO.: 50 SEQ ID NO.: 51
TGLA116	R TGGAGAAGATTGGCTGTACCCA F CTTAACTCATTCACCTCAACTG	SEQ ID NO.: 52 SEQ ID NO.: 53
INRA072	R AGTGATTGAGCACATTGCGCAT F GCATTGGCAAGTGGATTCTT	SEQ ID NO.: 54 SEQ ID NO.: 55
BM8233	R AAGGCAATTAACACACATACATCACC F ACTTCCCATTCCATCCATCAG	SEQ ID NO.: 56 SEQ ID NO.: 57
BMS648	R CTTCCATTCTCAGCCATCTAGC F TGAGCCATAGAATTAAAGATTCAAGC	SEQ ID NO.: 58 SEQ ID NO.: 59
BR6303	R TTTGTTCCCTCTTATTTCATCTGC F GAGCAGCTTCTTCATCTTCTCATCTT	SEQ ID NO.: 60 SEQ ID NO.: 61
MGTG4B	R GCTCTTGGAAAGCTTATTGTATAAAG	SEQ ID NO.: 62
<b>BTA5:</b>		
BMS1095	F AGGGATTGGTTATGCTCTCTC R GTTGCAGAGTCGGACATGAC	SEQ ID NO.: 63 SEQ ID NO.: 64
BM6026	F GCAACTAACGACCCAACCAAC R ACTGATGTGCTCAGGTATGACG	SEQ ID NO.: 65 SEQ ID NO.: 66
MNB-33	F GCTTGGTACACCCCTTAAGC R GAACAAATTACAAGGGAAAAC	SEQ ID NO.: 67 SEQ ID NO.: 68
BMS610	F TTTCACTGTCATCTCCCTAGCA R ATGTATTCATGCACACCAACACA	SEQ ID NO.: 69 SEQ ID NO.: 70
BP1	F AAAATCCCTTCATAACAGTGCC R CATCGTGAATTCCAGGGTTC	SEQ ID NO.: 71 SEQ ID NO.: 72
DIK4747	F CCAAAAATTCTGGCACCAAT R CCTGGGCTTGTGACTAGCAT	SEQ ID NO.: 73 SEQ ID NO.: 74
DIK2718	F AGGAAGGACAAGGACATTGC R AGAGGGTCAAAGGCTTAATGG	SEQ ID NO.: 75 SEQ ID NO.: 76
AGLA293	F GAAACTCAACCCAAGACAACTCAAG R ATGACTTTATTCTCCACCTAGCAGA	SEQ ID NO.: 77 SEQ ID NO.: 78
DIK5002	F TGTGCTGGAGGTGATAGCTG R TGCAGGAATATGAGAGCTGAGA	SEQ ID NO.: 79 SEQ ID NO.: 80
DIK4759	F AGTTGGACCTGCCATTGTT R ACTTATGTGCGTGCCTGCT	SEQ ID NO.: 81 SEQ ID NO.: 82
BMC1009	F GCACCAGCAGAGAGGACATT R ACCGGCTATTGTCCATCTTG	SEQ ID NO.: 83 SEQ ID NO.: 84
CSSM034	F CCATAACTCTGGACTTTCTCA R ATGTTCAGCCATCTCCTGGTCC	SEQ ID NO.: 557 SEQ ID NO.: 558

RM500	<b>F</b> CAGACACGACTAAGCGACCA <b>R</b> CCTACAATAAAGCACGGGA	SEQ ID NO.: 85 SEQ ID NO.: 86
BMS1617	<b>F</b> GCCTGCATGTGTCTGTGG <b>R</b> TCTGTGTCGGAATACCCCTCC	SEQ ID NO.: 87 SEQ ID NO.: 88
DIK5046	<b>F</b> TGAATTGTTCTGCTTCTTGGAA <b>R</b> TGCATGACTCCCCTCTCTCT	SEQ ID NO.: 89 SEQ ID NO.: 90
ETH10	<b>F</b> GTTCAGGACTGGCCCTGCTAACAA <b>R</b> CCTCCAGCCCACTTCTCTTCTC	SEQ ID NO.: 91 SEQ ID NO.: 92
CSSM022	<b>F</b> TCTCTCTAATGGAGTTGGTTTTG <b>R</b> ATATCCCAC TGAGGATAAGAATTCA	SEQ ID NO.: 93 SEQ ID NO.: 94
BMS1216	<b>F</b> GAGTAGAACACAACTGAGGACACA <b>R</b> CAATGCTGTGGGTACTGAGG	SEQ ID NO.: 95 SEQ ID NO.: 96
DIK2943	<b>F</b> GGTTTCCTCAGGACATGGTG <b>R</b> CAGTCCATGAGGTTGCAGAA	SEQ ID NO.: 97 SEQ ID NO.: 98
BMS1248	<b>F</b> GTAATGTAGCCTTTGTGCCG <b>R</b> TCACCAACATGAGATAGTGTGC	SEQ ID NO.: 99 SEQ ID NO.: 100
BM315	<b>F</b> TGGTTTAGCAGAGAGCACATG <b>R</b> GCTCCTAGCCCTGCACAC	SEQ ID NO.: 101 SEQ ID NO.: 102
BMS1658	<b>F</b> ATTGATGCTTATGATCCTCATG <b>R</b> CCCACTAAGAGAGGGAGGAGG	SEQ ID NO.: 103 SEQ ID NO.: 104
BM2830	<b>F</b> AATGGCGTATAAACACAGATG <b>R</b> TGAGTCCTGTCACCACATCAGC	SEQ ID NO.: 105 SEQ ID NO.: 106
<b>BT47:</b>		
BM7160	<b>F</b> TGGATTTAAACACAGAACATGTGG <b>R</b> TCAGCTTCTCTTAAATTCTCTGG	SEQ ID NO.: 107 SEQ ID NO.: 108
BL1067	<b>F</b> AGCCAGTTCTCAAATCAACC <b>R</b> ATGGTTCCGCAGAGAAACAG	SEQ ID NO.: 109 SEQ ID NO.: 110
BMS713	<b>F</b> CCAAGGGAGGAAAAATAAGTTAA <b>R</b> ACCAGCAGTAGGTTGAGGTTAA	SEQ ID NO.: 111 SEQ ID NO.: 112
DIK5321	<b>F</b> AACCTTCACAGGCTCCTCC <b>R</b> CCCATCTCTTGTCGCAAATC	SEQ ID NO.: 113 SEQ ID NO.: 114
DIK4421	<b>F</b> CATCTGAATGGCCAGAACATGA <b>R</b> GTCCCTGCATGTGTCTCTC	SEQ ID NO.: 115 SEQ ID NO.: 116
DIK2207	<b>F</b> ACATTGGCTTACGCTCACACT <b>R</b> CCTGTCTGGTTTGTGCT	SEQ ID NO.: 117 SEQ ID NO.: 118
DIK5412	<b>F</b> ATGGACAGAACAGCCTGACA <b>R</b> TGGTGAACTCAGCCTCACTG	SEQ ID NO.: 119 SEQ ID NO.: 120
IL4	<b>F</b> GTGCTGGACATCTGCAAGTG	SEQ ID NO.: 121

	R ACATTCAGGTCTGTGATCCATG	SEQ ID NO.:122
	F ACTAATAAGAAATTCTGCATGTGTG	SEQ ID NO.:123
BM6105	R CCACCATGACTCAGAAGTAGTTC	SEQ ID NO.:124
	F TAATCATAAGTCAAAGTAACAGTT	SEQ ID NO.:125
TGLA303	R GATCTGGACATACAAAAGTATTAC	SEQ ID NO.:126
	F TTACTTTCTGGGCCAGAG	SEQ ID NO.:127
DIK2819	R GGAACTGTGCCACATAGCAA	SEQ ID NO.:128
	F TCTGGAAAGGGGAAAAAGC	SEQ ID NO.:129
DIK4606	R TGCTTCATAGCACTTATCTCTTCA	SEQ ID NO.:130
	F AGTAAGGCCTGCAGTATTATATCC	SEQ ID NO.:131
BM7247	R AATCTTCCCTAGAACTTACAAAGG	SEQ ID NO.:132
	F CTGAAACACTCTAAAGGGTATGC	SEQ ID NO.:133
UWCA20	R ATCCAACATCCACCCATTCC	SEQ ID NO.:134
	F GTTCTGAGGTTGTAAAGCCC	SEQ ID NO.:135
BM6117	R GGTGAGCTACAATCCATAGGG	SEQ ID NO.:136
	F AGGAACCCATAGGCAGACAC	SEQ ID NO.:137
BMS2840	R GCCTGGCAAAGAGAAAATTG	SEQ ID NO.:138
	F TCTCACCTCACATGGTTCA	SEQ ID NO.:139
DIK2915	R GTGGAGCCAAGGTGAAAGAA	SEQ ID NO.:140
	F CCAGCAGAAGAGAAAGATACTGA	SEQ ID NO.:141
BMS2258	R AGTGGTAGAACCTCCATCTCACA	SEQ ID NO.:142
	F AATCCAGTGTGTGAAAGACTAATCCAG	SEQ ID NO.:143
OARAE129	R GTAGATCAAGATATAGAATATTTTCAACACC	SEQ ID NO.:144
	F CTCAATGACGTTGGCTTCA	SEQ ID NO.:145
DIK2895	R GGTGCCTGACTCCAATTGAT	SEQ ID NO.:146
	F TGTCTGTATTCTGCTGTGG	SEQ ID NO.:147
ILSTS006	R ACACGGAAGCGATCTAACCG	SEQ ID NO.:148
	F AGTGCCAAAAGGAAGCGC	SEQ ID NO.:149
BL1043	R GACTTGACCGTTCCACCTG	SEQ ID NO.:150
<b>BTA8:</b>		
	F CCTCTGGGTCTATCCATGTTG	SEQ ID NO.:151
IDVGA-11	R TGGATGAATGAAGAAAGATGCC	SEQ ID NO.:152
	F GACAAGATAGGCTTGCATGA	SEQ ID NO.:153
BMS1591	R GATAGAAATATACCAGGAGCTCACA	SEQ ID NO.:154
	F ACCATCTACTGTGCTATGGCTT	SEQ ID NO.:155
BMS678	R GCAGAAACACAATACTCAGTGC	SEQ ID NO.:156
	F GGGTAGCCTGTTAAAATGCAG	SEQ ID NO.:157
INRA129	R CAGTGCTGACCTCTGAAGTAAG	SEQ ID NO.:158

BMS2072	F TGTTCAGTGCTTGCTTAGCTG R TCTTCAAAGCCATCAATCATC	SEQ ID NO.:159 SEQ ID NO.:160
BMS887	F AAGCTAACTGATATTCTGCCACA R TTCCCTCTCTCCCTCTCC	SEQ ID NO.:161 SEQ ID NO.:162
URB037	F ACTGGAGACGACTGAAGCAACC R GAGTGGCTGTTGCTAAATTGG	SEQ ID NO.:163 SEQ ID NO.:164
MCM64	F TACAGTCCATGGGTACAAAGAG R TCTGAATCTACTCCCTCCTCAGAGC	SEQ ID NO.:165 SEQ ID NO.:166
CSSM047	F TCTCTGTCTCTATCACTATATGGC R CTGGCACCTGAAACTATCATCAT	SEQ ID NO.:167 SEQ ID NO.:168
BMS836	F GAAACTCTTTCACTCTGCGC R GCTCTTAGGGATTGCTTCACC	SEQ ID NO.:169 SEQ ID NO.:170
<b>BTAG9:</b>		
BMS2151	F CCATTAAGAGGAAATTGTGTTCA R ATGGAGTCACTGAAAGGTACTGA	SEQ ID NO.:171 SEQ ID NO.:172
ETH225	F GATCACCTGCCACTATTCCT R ACATGACAGCCAGCTGCTACT	SEQ ID NO.:173 SEQ ID NO.:174
ILSTS037	F TAGGCTATGTACTGACCATGC R CTGAACTGAGATGACTTGGC	SEQ ID NO.:175 SEQ ID NO.:176
BM2504	F CAGCTTCCATCCCCTTTC R CTCCCATCCAAACACAGAC	SEQ ID NO.:177 SEQ ID NO.:178
DIK2892	F TTGACCCCTGAAAGATGTCCA R CACGGTTTATCAGCTTGGTA	SEQ ID NO.:179 SEQ ID NO.:180
DIK3003	F ACTTCAGTTTGGCTGAC R TGTCACTAGGTAAATTGGTG	SEQ ID NO.:181 SEQ ID NO.:182
DIK3002	F AAATGGAGGTAATGAAATAAAATA R CAAACCCATGGACTGTAACCT	SEQ ID NO.:183 SEQ ID NO.:184
BMS1267	F TTCTGAATTGATTCCCAACA R ACTGTTCTTAAAGCTTCCC	SEQ ID NO.:185 SEQ ID NO.:186
DIK5142	F TGGGTAAGTGGAAAGGATG R CTCAGCCAGGTTGTCCTCTC	SEQ ID NO.:187 SEQ ID NO.:188
BMS555	F GGAAAGAGTAGGTGATTCCCTG R ATTTAATTGTCATCCCAGGTGA	SEQ ID NO.:189 SEQ ID NO.:190
DIK5364	F CCTCTGAAACCCCAGACTTG R AAAAACCCAAAACAACACACAA	SEQ ID NO.:191 SEQ ID NO.:192
UWCA9	F CCTTCTCTGAATTGGTTGAAAGC R GGACAGAAGTGAGTGACTGAGA	SEQ ID NO.:193 SEQ ID NO.:194
DIK4720	F CATGATATTACCCGTGTGTC R	SEQ ID NO.:195

	R GAGGAGCTGGAGGGCTAAAG	SEQ ID NO.:196
	F TTGGCACTTACTACCTCATATGTT	SEQ ID NO.:197
BMS1290	R TTTCTGGATGTTGAGCCTATT	SEQ ID NO.:198
	F ACCTTGGGAATCAAGGTCAT	SEQ ID NO.:199
DIK2816	R CCCAGTAGTCCAGTGGCTCA	SEQ ID NO.:200
	F AAAGACTGCTGCCTGAAGC	SEQ ID NO.:201
BM6436	R CAACCACTGATGCTGTACTCTG	SEQ ID NO.:202
	F TCAAAAAGTGGACATGACTGA	SEQ ID NO.:203
BMS2753	R AGGTTTCAAATGAGAGACTTTTC	SEQ ID NO.:204
	F TCAGTACACTGGCCACCATG	SEQ ID NO.:205
BM4208	R CACTGCATGCTTTCCAAAC	SEQ ID NO.:206
	F GCTCACAGGTTCTGAGGACTC	SEQ ID NO.:207
BMS2819	R AACTTGAAGAAGGAATGCTGAG	SEQ ID NO.:208
	F GCTCTGGTACCCAGGTG	SEQ ID NO.:209
BMS2295	R CTGGCAGGAGATGAGAGGGAG	SEQ ID NO.:210
	F GGGCAGATGTGAGTAATTTCC	SEQ ID NO.:211
BMS1967	R AACTGAGCTGTATGGTGGACG	SEQ ID NO.:212
<b>BTA10</b>		
	F GCACATTGGGATCTCTCCTG	SEQ ID NO.:213
DIK2658	R AAAGTCCCACCCACAATCA	SEQ ID NO.:214
	F TCCTTACAACACACCATGCAA	SEQ ID NO.:215
DIK2503	R CACACCCAGGCATCCATAC	SEQ ID NO.:216
	F TTCATATAAGCAGTTATAAACGC	SEQ ID NO.:217
CSSM38	R ATAGGATCTGGTAACTTACAGATG	SEQ ID NO.:218
	F CTCACTCCACTGGGCTTCTC	SEQ ID NO.:219
BMS528	R TGTGTTCTCACCTCGACCAC	SEQ ID NO.:220
	F TCATCTGGGCATAAGACAGG	SEQ ID NO.:221
BM1237	R ATTGTTCCCAGCATCTTAGAGG	SEQ ID NO.:222
	F CACCCGTACCCCTCACTGC	SEQ ID NO.:223
MB077	R TCACAACCCCTTCTCACCC	SEQ ID NO.:224
	F TGGCTTGCAACACTGCAC	SEQ ID NO.:225
DIK2000	R CCCACCTACGACTGGACTTA	SEQ ID NO.:226
	F GCTTCAGTTCTGCTTTCAACC	SEQ ID NO.:227
BMS2742	R CTTCAGCATCTTGATTGTTGC	SEQ ID NO.:228
	F CTCCAGGTAAGACAGGCCAC	SEQ ID NO.:229
BMS529	R CCCGATCTGTGTGGGT	SEQ ID NO.:230
	F TGTGGGTTTGATCTCTGAGT	SEQ ID NO.:231
DIK2361	R TGTGTCCTCCTTGTGGTAGAA	SEQ ID NO.:232

	<b>F</b> ACTAGGAGGCCATATAGGAGGC	SEQ ID NO.:233
BM888	<b>R</b> GAGCTAAAACGAGGGACAG	SEQ ID NO.:234
	<b>F</b> ATTCTATGAAGTAGTCTTCTGACT	SEQ ID NO.:235
TGLA433	<b>R</b> ATTTAAAACTAGTCACGAGTGCCT	SEQ ID NO.:236
	<b>F</b> GATCCTGCTTATATTAACCAC	SEQ ID NO.:237
INRA037	<b>R</b> AAAATTCCATGGAGAGAGAAC	SEQ ID NO.:238
	<b>F</b> TATGAACTCACATGGTTACCACA	SEQ ID NO.:239
BMS1620	<b>R</b> TTGCCCAAAAATAGACCTAAA	SEQ ID NO.:240
	<b>F</b> GGTATTTGAGAATGTGGGC	SEQ ID NO.:241
ILSTS070	<b>R</b> TCTTGACCACTACCTATCC	SEQ ID NO.:242
	<b>F</b> GTGCGGAAAGGAACAGAGTC	SEQ ID NO.:243
BMS2641	<b>R</b> AAAGCCGGACTGGAGTGTC	SEQ ID NO.:244
	<b>F</b> AATGCGTGGACTTGT	SEQ ID NO.:245
BMS614	<b>R</b> CAATTGCTGAAGCAGTCACA	SEQ ID NO.:246
	<b>F</b> ACTTTCTTTCCTGTGGCTCG	SEQ ID NO.:247
BMS2614	<b>R</b> CAGAGCTGGCACCAAGAGG	SEQ ID NO.:248
<b>BTA11:</b>		
	<b>F</b> AGTACTTGGCTTGCTTGCTC	SEQ ID NO.:249
BM716	<b>R</b> TTAAATTCCATCTCACCCCTGG	SEQ ID NO.:250
	<b>F</b> AGAGAGGCCAAAGCTGGG	SEQ ID NO.:251
BMS2569	<b>R</b> TTCCCTGGGCTTCAGGAG	SEQ ID NO.:252
	<b>F</b> TTCTGTGGTTGAAGAGTGT	SEQ ID NO.:253
BM2818	<b>R</b> CAATGGCTAACAGAGGCCAGTG	SEQ ID NO.:254
	<b>F</b> TCCAAAAGTTCGTACATATTG	SEQ ID NO.:255
INRA177-2	<b>R</b> CACCAGGCTTCTCTGTTGAA	SEQ ID NO.:256
	<b>F</b> TCCAAAAGTTCGTACATATTG	SEQ ID NO.:257
INRA177	<b>R</b> CACCAGGCTTCTCTGTTGAA	SEQ ID NO.:258
	<b>F</b> TCGAAAAAGTTGGACAAGAC	SEQ ID NO.:259
RM096	<b>R</b> TTAGCAGGGTGCCTGACACTT	SEQ ID NO.:260
	<b>F</b> GGTAAAATCCTGCAAAACACAG	SEQ ID NO.:261
INRA131	<b>R</b> TGACTGTATAGACTGAAGCAAC	SEQ ID NO.:262
	<b>F</b> TGGTATGTAGTTACAGCAGCCC	SEQ ID NO.:263
BM7169	<b>R</b> CCATTGAAACAGACATGAATGC	SEQ ID NO.:264
	<b>F</b> GTGTCTGCAAAAGATGAATGG	SEQ ID NO.:265
BM6445	<b>R</b> GACAACGTCTCGTTGGG	SEQ ID NO.:266
	<b>F</b> GAGTATTATGCTGGGAGGC	SEQ ID NO.:267
ILSTS036	<b>R</b> AGACAGGATGGGAAGTCACC	SEQ ID NO.:268
	<b>F</b> AAAGGCTTCTATTTGTGGTGG	SEQ ID NO.:269

	R TTGATGCTTTATTGTTTCCTCT	SEQ ID NO.:270
TGLA58	F TTCTACTCTCCAGCCTCCTCC	SEQ ID NO.:271
	R GTTGGCTCCAAGAGCAAGTC	SEQ ID NO.:272
	F ACTATGGACATTGGGGCAG	SEQ ID NO.:273
BMS2047	R AGTAGGTGGAGATCAAGGATGC	SEQ ID NO.:274
	F CAGACCAGTTCTCAGACAAGC	SEQ ID NO.:275
HUJV174	R TCATTCTGTGTCAATACAGCC	SEQ ID NO.:276
	F TTTGAGAACCTTTGTTCTGAGC	SEQ ID NO.:277
BMS989	R TTATTTGCTTTCTGATTTGTG	SEQ ID NO.:278
	F TGTATGGCTGAATGATATTCCATT	SEQ ID NO.:279
TGLA436	R CTACTGACAGATGATTAGATAAAGA	SEQ ID NO.:280
	F TGCCCCATAGTGTAGTGCTC	SEQ ID NO.:281
BMS460	R GCCAGCAGAGAATTGTAGCA	SEQ ID NO.:282
	F TTCTGGCAAACATATTCCACC	SEQ ID NO.:283
ILSTS045	R CATGAAAGACACAGATGACC	SEQ ID NO.:284
	F ATTTTCCCAGCGCCTCTC	SEQ ID NO.:285
DIK4819	R AAACAGAAGACTCAGGAAGACGA	SEQ ID NO.:286
	F TAAGGACTTGAGATAAGGAG	SEQ ID NO.:287
HEL13	R CCATCTACCTCCATCTAAC	SEQ ID NO.:288
<b>BTA12:</b>		
BMS410	F GGCTGAAAAGCTGTGGTGT	SEQ ID NO.:289
	R TTGCCACATTACCTTCTTCA	SEQ ID NO.:290
	F TTCTAATGTAGAGCAAAGTGATTGA	SEQ ID NO.:291
BM6108	R TGTAGGAGGGACAGATTGGG	SEQ ID NO.:292
	F ACCAGATTGGTGGTAGTGGT	SEQ ID NO.:293
BM860	R CATGCCGTGGCTAACGACC	SEQ ID NO.:294
	F TGGAGCTAAATCAATGCGTG	SEQ ID NO.:295
BMS975	R CCAATGGCCAATTAAAGTACC	SEQ ID NO.:296
	F CCTTCATGGAAGAAATTGTG	SEQ ID NO.:297
BMS1316	R GGAGTTACAGTCCATGGGTT	SEQ ID NO.:298
	F GGCTGATACACAGAGACATGC	SEQ ID NO.:299
BMS2724	R CCTCTCTGCCTTCTATCAGGT	SEQ ID NO.:300
<b>BTA15:</b>		
BR3510	F GCTGGTGGGTTGTTACAC	SEQ ID NO.:301
	R ACCCCGTGGACTGTAGTCTG	SEQ ID NO.:302
	F TGAAGTAAGTAAGCACACAAGCA	SEQ ID NO.:303
BMS2533	R TTGATCATCTTAGGTCCATCC	SEQ ID NO.:304
	F ACAGGCTACAGTCCATGGGTT	SEQ ID NO.:305

	R TATAGAACAGAAAAATGACTACACG	SEQ ID NO.:306
	F CACGTCACCCGCTTCTCTTG	SEQ ID NO.:307
JAB8	R GGTGAGTGTAAACACCTGTGTGCG	SEQ ID NO.:308
	F CCAAGGTCATTGTTGCAGC	SEQ ID NO.:309
BMS2684	R TGGGGATTGCTTCTCAGTC	SEQ ID NO.:310
	F CAAGAGTCAGACATGACTTAGTGAC	SEQ ID NO.:311
DIK1106	R TCTACCTTTGATAGCGTGAGC	SEQ ID NO.:312
	F TAATAAAACTGGTCCCTCTGGC	SEQ ID NO.:313
INRA145	R TGCTGGCTCTCCAGTATGC	SEQ ID NO.:314
	F TCTCCTGGCTACAGGGCTAA	SEQ ID NO.:315
IDVGA-10	R CCCACTGGCCTAGAACCC	SEQ ID NO.:316
	F AGGGGCGAAGTGAGGATTA	SEQ ID NO.:317
DIK4850	R TTGCATGGTTCTGCAGATGT	SEQ ID NO.:318
	F AGCCTTCCCAGTACCTGTCA	SEQ ID NO.:319
DIK2768	R TAAGGGAGCTCAAAACCACA	SEQ ID NO.:320
	F GGTGTGTTGGTTAACAGTGG	SEQ ID NO.:321
ILSTS027	R GAATCATAGACCTGACTTCC	SEQ ID NO.:322
	F TGGACAGGACTGAGTATGCA	SEQ ID NO.:323
BMS812	R AGGTATCCAACTAACACAGCCA	SEQ ID NO.:324
	F AGCACCTGTACCATCTGTTCC	SEQ ID NO.:325
BMS2076	R TCCATAGGCTCACAAAGAGTTG	SEQ ID NO.:326
	F TCCCTCTACCATATATTCCCC	SEQ ID NO.:327
BL1095	R CATTAGCATGGAAAAACCTCTG	SEQ ID NO.:328
	F CCACTACTGCCTCAGGGAG	SEQ ID NO.:329
BMS820	R ACAGGACTCTCAAGCATCAGC	SEQ ID NO.:330
	F GATGATCCACCATAACTACCAGA	SEQ ID NO.:331
BMS927	R TGGCTCTCAAAGGTATTGT	SEQ ID NO.:332
	F TACATTAACCCCAAAATTAAATGC	SEQ ID NO.:333
BMS429	R CCCTTGATTCTCTCATGAGTATT	SEQ ID NO.:334
<b>BTA18:</b>		
	F CCTTGAGATGAATGTTGAGGATG	SEQ ID NO.:335
IDVGA-31	R AACGCAGCCAGCAGGGTCAGG	SEQ ID NO.:336
	F TAAAACCCCAAAAGAACCC	SEQ ID NO.:337
BMS1355	R ATATTGCGACATTGGATGAA	SEQ ID NO.:338
	F TGATGCTGATTGATTGTGTG	SEQ ID NO.:339
BMS1322	R TATCTTGCTCACTCTTCCCC	SEQ ID NO.:340
	F TGTGGCTAGGTTCAAGCTCC	SEQ ID NO.:341
TEXAN-10	R TCTCTCTGGTGCATCCATTG	SEQ ID NO.:342

BMS2213	F ATGGGCAGCTTAGGGATTG R CTTCAAGAGCCTTCAGTGGG	SEQ ID NO.:343 SEQ ID NO.:344
INRA121	F GGAAACCCATTGGAGGATTG R CTTCACTATTCCCCACAAAGC	SEQ ID NO.:345 SEQ ID NO.:346
BR4406	F TACCTACCAGTTTCCAGCACC R AGAAGAGCCTGGAGGGCTAC	SEQ ID NO.:347 SEQ ID NO.:348
BMS2554	F GGGCTGTAAAGAGTAGGACACA R ATCATCTGCTTCCAGTCACAG	SEQ ID NO.:349 SEQ ID NO.:350
MNB-27	F GAGTAAATAAAGCTGCATGATGTC R GGATCAGGAGATTCAACACAG	SEQ ID NO.:351 SEQ ID NO.:352
BM7109	F CAGGTAAAAGAGCGGCTTG R CAGCTTCATGCCCTAGAAGG	SEQ ID NO.:353 SEQ ID NO.:354
INRA063	F ATTTGCACAAGCTAAATCTAAC R AAACCACAGAAATGCTTGGAAAG	SEQ ID NO.:355 SEQ ID NO.:356
ILSTS002	F TCTATACACATGTGCTGTGC R CTTAGGGGTGAAGTGACACG	SEQ ID NO.:357 SEQ ID NO.:358
BMS2639	F ATATCGTTTCAGATTCTTTGC R GAGAGATAAATTGGGAGTTGAGA	SEQ ID NO.:359 SEQ ID NO.:360
DIK4960	F CGCAACTTCCAAGTCCATCT R GGACACCTTCCCTGTCTCAA	SEQ ID NO.:361 SEQ ID NO.:362
DIK4849	F CCATCTTCCCCATTGTGTA R CCCCTCTTCATCTCAAAACA	SEQ ID NO.:363 SEQ ID NO.:364
BMON117	F TAGGGCCGTGATACTGTGT R CTCTACCATCCAGCACCCCTAAT	SEQ ID NO.:365 SEQ ID NO.:366
DIK4232	F TTGTGAGGTAAAGGGACATGA R GCCAGATTGCCAACTGTTT	SEQ ID NO.:367 SEQ ID NO.:368
BMS2785	F ACAAACCTGTGCGCCTTG R GGCAATCAGTCGGACACAC	SEQ ID NO.:369 SEQ ID NO.:370
DIK4569	F TCCCCCTAAGGCTCAGAGTT R CTAACCTCCCCCTCGGAACC	SEQ ID NO.:371 SEQ ID NO.:372
BM2078	F CCCAAAAGAAGCCAGGAAG R TCAGAGTTGGGGTCCTCAG	SEQ ID NO.:373 SEQ ID NO.:374
BM6507	F ACTTAGCACAATGCCCTCTAGG R ATGTTATTCCATCAGGAGGAGC	SEQ ID NO.:375 SEQ ID NO.:376
TGLA227	F CGAATTCCAAATCTGTTAATTGCT R ACAGACAGAAACTCAATGAAAGCA	SEQ ID NO.:377 SEQ ID NO.:378
DIK4013	F GAAATTGTGACCCCTGCAT R CTAAAGCTGCCTCCCAAG	SEQ ID NO.:379 SEQ ID NO.:380

**BTA19:**

BM9202	<b>F</b> TCTATGAAGACTTCAGGACCTC	SEQ ID NO.:381
BMS745	<b>R</b> GCATCCCGGTCTCCTATG	SEQ ID NO.:382
	<b>F</b> TAGGGACTTGTACCCGTGG	SEQ ID NO.:383
	<b>R</b> TGCAAGCTGTGAGGAGGAG	SEQ ID NO.:384
BP20	<b>F</b> TCTGTGGGTGAACAAGCAAG	SEQ ID NO.:385
	<b>R</b> GGCTCCCTAAAGACCCACTC	SEQ ID NO.:386
	<b>F</b> AAATCCTTCAAGTATGTTTCA	SEQ ID NO.:387
IDVGA-46	<b>R</b> ACTCACTCCAGTATTCTGTCTG	SEQ ID NO.:388
	<b>F</b> AATGTTAGGTTACATGCAGCC	SEQ ID NO.:389
BMS2389	<b>R</b> AGGCAATAGGATCTCCACTAGC	SEQ ID NO.:390
	<b>F</b> TTCCTGCTTGGTAAACTTGAAC	SEQ ID NO.:391
CSSM065	<b>R</b> CAACTCAAAGCTCAACAGCAGCC	SEQ ID NO.:392
	<b>F</b> GAACCTGCCTCTCCTGCATTGG	SEQ ID NO.:393
ETH3	<b>R</b> ACTCTGCCTGTGGCCAAGTAGG	SEQ ID NO.:394
	<b>F</b> CACTAGGACGATGCTCTCAGG	SEQ ID NO.:395
BMS601	<b>R</b> TCACAAGAGCAATGACGAGG	SEQ ID NO.:396

**BTA20:**

BM3517	<b>F</b> GTGTGTTGGCATCTGGACTG	SEQ ID NO.:397
	<b>R</b> TGTCAAATTCTATGCAGGATGG	SEQ ID NO.:398
	<b>F</b> GCATTAGGTTCTCCAGAGAA	SEQ ID NO.:399
HEL12	<b>R</b> CAGACTTGTCAAGACTCCATA	SEQ ID NO.:400
	<b>F</b> ACTCTTCCACAGTTGGCCTG	SEQ ID NO.:401
BMS1282	<b>R</b> CCTCCTTCCTCCAGAGCC	SEQ ID NO.:402
	<b>F</b> GCATTATTCTTGTCTTTGGG	SEQ ID NO.:403
BMS1754	<b>R</b> GTTCTGCTCCTGATCTCCTG	SEQ ID NO.:404
	<b>F</b> CTAATTAGAACATGAGAGAGGCTCT	SEQ ID NO.:405
TGLA126	<b>R</b> TTGGTCTCTATTCTCTGAATATTCC	SEQ ID NO.:406
	<b>F</b> ACACAACCCAAATGTTACCAA	SEQ ID NO.:407
BMS2361	<b>R</b> ATTGTGCAGAGACCAAGTGC	SEQ ID NO.:408
	<b>F</b> AGGAAGCCGAGTGAGATATGTAAGC	SEQ ID NO.:409
AGLA29	<b>R</b> TTACAGCCTGTGTGAATGTCCTCTA	SEQ ID NO.:410
	<b>F</b> CAATGAGCTCAGATTGTTGCA	SEQ ID NO.:411
BMS703	<b>R</b> ATACATGTAGTCAAAAGGCTCATCC	SEQ ID NO.:412
	<b>F</b> TCTGGAGTGAATGTTCTGAGG	SEQ ID NO.:413
BM5004	<b>R</b> TTGTGATGAGCACCTGAAGG	SEQ ID NO.:414
	<b>F</b> TGGGGTCTAAAGAGTCAGAG	SEQ ID NO.:415
UWCA26	<b>R</b> TTCAAGTCTGCCCTTGGTTCGT	SEQ ID NO.:416

**BTA21:**

	<b>F</b> CGATGTAAAGGGCAGGTTCT	SEQ ID NO.:417
DIK5182	<b>R</b> CTCTAGAATCCTGTTTAGGG	SEQ ID NO.:418
	<b>F</b> TGTGTGCTCTCACACATGC	SEQ ID NO.:419
BMS1117	<b>R</b> AACCAAAGCAGGGATCAGG	SEQ ID NO.:420
	<b>F</b> CCCAGAGGTGACAACATTCCAG	SEQ ID NO.:421
RM151	<b>R</b> GATCCACCAAAAACCAGCTGGA	SEQ ID NO.:422
	<b>F</b> AATCCATCCATTCAGCCTTC	SEQ ID NO.:423
DIK2492	<b>R</b> GAAATGACAGCCCACCTCCAG	SEQ ID NO.:424
	<b>F</b> TGCAAACATCCACGTAGCATAAATA	SEQ ID NO.:425
AGLA233	<b>R</b> GCATGAACAGCCAATAGTGTCATC	SEQ ID NO.:426
	<b>F</b> GAAAGATGTTGCTAGTGGGG	SEQ ID NO.:427
ILSTS095	<b>R</b> ATTCTCCTGTGAACCTCTCC	SEQ ID NO.:428
	<b>F</b> GACTGCGACCAGGTCTTTTC	SEQ ID NO.:429
DIK4602	<b>R</b> AGGCCCATACGCATTGTTA	SEQ ID NO.:430
	<b>F</b> CTAGCTGCTGGCTACTTGGG	SEQ ID NO.:431
BM103	<b>R</b> GGCTGCTCTGGCTATTG	SEQ ID NO.:432
	<b>F</b> TTCTCCAACCCGGTTATGC	SEQ ID NO.:433
DIK4001	<b>R</b> CTGATTGGTCACTCCATCCA	SEQ ID NO.:434
	<b>F</b> GTGGTGGCAAAGAGTCAGA	SEQ ID NO.:435
IDVGA-45	<b>R</b> AACAGCCCTGATTCCATA	SEQ ID NO.:436
	<b>F</b> CCGTGTGCTTCCTCTGA	SEQ ID NO.:437
DIK2481	<b>R</b> TGACAGCAGCCAAGATATGG	SEQ ID NO.:438
	<b>F</b> TTGTCCAGCCCAGCATTAGC	SEQ ID NO.:439
INRA103	<b>R</b> GGAGAAGACTTATGGGAGC	SEQ ID NO.:440
	<b>F</b> TGATATTCAAACCTCAATGAACCC	SEQ ID NO.:441
BMS2815	<b>R</b> CTTGCATATGCTCATCATTATCA	SEQ ID NO.:442
	<b>F</b> GGATTTAGCTGCCATTGCT	SEQ ID NO.:443
DIK2842	<b>R</b> AATCCCATGGACAGAAAAGC	SEQ ID NO.:444
	<b>F</b> TGTGTGGCTTAGCACAT	SEQ ID NO.:445
DIK3036	<b>R</b> CAGAAAGGGAAATCACATCC	SEQ ID NO.:446
	<b>F</b> CCCTTCCAATAGGCAAATCTC	SEQ ID NO.:447
DIK4391	<b>R</b> TCCAACAAGCTTCTTCC	SEQ ID NO.:448
	<b>F</b> AACGTCCAGTCGCTCAAAT	SEQ ID NO.:449
DIK2913	<b>R</b> TCACACACCTGAACCTCAAAGC	SEQ ID NO.:450
	<b>F</b> GACCACTGGACCACCAGG	SEQ ID NO.:451
BM846	<b>R</b> CTGGTAAAAGCAATGATGCC	SEQ ID NO.:452
TGLA122	<b>F</b> CCCTCCTCCAGGTAAATCAGC	SEQ ID NO.:453

	R AATCACATGGCAAATAAGTACATAC	SEQ ID NO.:454
ILSTS054	F GAGGATCTGATTTGATGTCC	SEQ ID NO.:455
	R AGGCCACTATGGTACTTCC	SEQ ID NO.:456
BMS743	F AGCTACCCCTGGTATACAACACG	SEQ ID NO.:457
	R GCTCTGAAATTCTGGCAGTG	SEQ ID NO.:458
IDVGA-30	F GCATCTGGGAGCCTCGTATCTC	SEQ ID NO.:459
	R TTGTAAAACTCGGGGCATAAGCA	SEQ ID NO.:460
<b>BTA22:</b>		
CSSM26	F GACTTCTGCTTGTGGTTCCAAGT	SEQ ID NO.:461
	R TTTCCCATTATGGTTATCCCAG	SEQ ID NO.:462
INRA026	F TAGTTCCAATGAGACACGAACA	SEQ ID NO.:463
	R TAGGAGCACGGAGGTAAAACA	SEQ ID NO.:464
BM1558	F TGAGGAAAGCCTTGGCAG	SEQ ID NO.:465
	R ACTGGGCCTAGCTCCTTCTC	SEQ ID NO.:466
BM3628	F CTGAGATGGACTCAGGGAGG	SEQ ID NO.:467
	R GTTGGATTGGAAAGGTTAGGC	SEQ ID NO.:468
BMS875	F TCCAGCTTGAATCCCTTCC	SEQ ID NO.:469
	R AAGCAAAGGCTGGAACAC	SEQ ID NO.:470
BM4102	F CCAAATTCCACTGTGCTGC	SEQ ID NO.:471
	R GAGCGGCCTATCAACCCCTAC	SEQ ID NO.:472
<b>BTA24:</b>		
BMS917	F TAATGCCTCTGGAAGGTTGA	SEQ ID NO.:473
	R CAAGCTGGTTGTTCTTTGC	SEQ ID NO.:474
BM7151	F AAATGTCCACTGCTCAAAGATG	SEQ ID NO.:475
	R ACTTGGAGATAGAACTGGCAGG	SEQ ID NO.:476
BM226	F ATTGCCTGTCCGTGTATCC	SEQ ID NO.:477
	R CCGGCTGAATTGCTATAAGC	SEQ ID NO.:478
BMS2526	F CAGGCTCCATGTTGGACAC	SEQ ID NO.:479
	R CATCAGGTTGGCAGAGTCG	SEQ ID NO.:480
TGLA351	F GCACATCTGGTGGCCACATCAG	SEQ ID NO.:481
	R CTCTAGGGGATTCAGTCTCAGT	SEQ ID NO.:482
BM7228	F TTAAATCCTCAAGTAAAGGAAGGC	SEQ ID NO.:483
	R GCAAACCTAAGAACCTCATTTC	SEQ ID NO.:484
CSSM23	F CACTGGAGTGGGTTGCCATTGTCT	SEQ ID NO.:485
	R GTTCGCAATATGATCTGATTG	SEQ ID NO.:486
BMS2270	F CTGCGTTAACACCCCCACC	SEQ ID NO.:487
	R GCAGGAAGGCTGATGCAC	SEQ ID NO.:488
ILSTS065	F GCTGCAAAGAGTTGAACACC	SEQ ID NO.:489

	R AACTATTACAGGAGGCTCCC	SEQ ID NO.:490
	F GCACATGCAATCTTGAAAGG	SEQ ID NO.:491
BMS1862	R ACCAGAGATGATGAAGAACCCC	SEQ ID NO.:492
	F AGCAGAGGGCAAATGTTATG	SEQ ID NO.:493
BMS466	R GGATGTAAGAGGATGCAGACC	SEQ ID NO.:494
	F GGTCACTTCCATTATGACAGCAG	SEQ ID NO.:495
INRA090	R GGTGTTACCTTTTAGTCTCC	SEQ ID NO.:496
	F CAACTAGCTCTCAATGCCTTT	SEQ ID NO.:497
BMS1926	R TTCTCCCAATCTGTAAGTGCA	SEQ ID NO.:498
	F CCAAACCAGTGTGACTGACA	SEQ ID NO.:499
BMS3024	R TTGCTCATTAACTTCATTACAACA	SEQ ID NO.:500
<b>BTA25:</b>		
	F CAGGACTGAGTAACTAAGGC	SEQ ID NO.:501
ILSTS102	R AGGAGACAGCTACAAACCCCC	SEQ ID NO.:502
	F ATCCAAGGAGGTCCCAGG	SEQ ID NO.:503
BMS2843	R TCCTCCAGTGGAAATATGG	SEQ ID NO.:504
	F TGGGATAGACCACATTGGAA	SEQ ID NO.:505
BM737	R GAATGCTTTGGGAGGGTA	SEQ ID NO.:506
	F TAAAGTCCTGCAAGAGAAGG	SEQ ID NO.:507
ILSTS046	R TTTCTGTCTTGAGTCTCTCC	SEQ ID NO.:508
	F TTTCAGGACTAATAGGGCATGG	SEQ ID NO.:509
BMS1353	R ATTCAAGACCTGCCTGGTGAC	SEQ ID NO.:510
	F GCAGAAGGAAAAAGCAATGG	SEQ ID NO.:511
AF5	R GATCCTGCGAGCCACAAG	SEQ ID NO.:512
<b>BTA26:</b>		
	F AATATGTAAAACAAGTCAAAGCA	SEQ ID NO.:513
BMS651	R CCTGGCAAGCAACAGTTAAT	SEQ ID NO.:514
	F CTTTGTGGAAGGCTAAGATG	SEQ ID NO.:515
HEL11	R TCCCACATGATCTATGGTGC	SEQ ID NO.:516
	F GACAAAACCTTTAGCACAGG	SEQ ID NO.:517
BMS332	R AATTGCATGGAAAGTTCTCAGC	SEQ ID NO.:518
	F TTGTACATTCTGTCAATGCCTT	SEQ ID NO.:519
RM026	R ACAATGTCATTGGTCAATTCTT	SEQ ID NO.:520
	F AGGTGCTGGAATGGCAAC	SEQ ID NO.:521
BM9284	R TGTGATTTGGTCTCCTTGC	SEQ ID NO.:522
	F TCTGTGAGCATGTGCAGAAT	SEQ ID NO.:523
RME40	R CTCACAGGTAAATGGGTGAT	SEQ ID NO.:524
IDVGA-59	F AACCCAAATATCCATCAATAG	SEQ ID NO.:525

	R CAGTCCCTCAACCCTCTTTC	SEQ ID NO.:526
BMS882	F TAGTGTCCACCAGAGACCCC	SEQ ID NO.:527
	R CCAAAGACACAGTTAAAGGGC	SEQ ID NO.:528
BM804	F CCAGCATCAACTGTCAGAGC	SEQ ID NO.:529
	R GGCAGATTCTTGCCTTCTG	SEQ ID NO.:530
BM7237	F CCAGCATCAACTGTCAGAGC	SEQ ID NO.:531
	R GGCAGATTCTTGCCTTCTG	SEQ ID NO.:532
<b>BTA28:</b>		
BMC6020	F ATTGCATGTAGCTCTGGGG	SEQ ID NO.:533
	R AAGTGGGTGGCTTCAACACT	SEQ ID NO.:534
ETH1112	F AGTGGATCCTGCATGTTATGCCG	SEQ ID NO.:535
	R CCAGACGGACCTTGTGGGCAA	SEQ ID NO.:536
BL25	F AACAGTGGCAATGGAAGTGG	SEQ ID NO.:537
	R AGTCAGGATCTAGTGGGTGAGTG	SEQ ID NO.:538
DIK2955	F CATTGAACACTGAAAGGAAAGC	SEQ ID NO.:539
	R TCACAAGGGCTTGAAGTGA	SEQ ID NO.:540
BMS2608	F GACTAAGCATATGAACCTGGC	SEQ ID NO.:541
	R CTGCCCCTTGTCATCTCATC	SEQ ID NO.:542
BMS2658	F TCCCTGGACTTCTGCAGAG	SEQ ID NO.:543
	R CTGGCCCCAGACACAATC	SEQ ID NO.:544
DIK713	F CACTTGCTGTGGACCTGAA	SEQ ID NO.:545
	R ACCCAGGAACTGAACCCAT	SEQ ID NO.:546
BMS1714	F TTTATCCCAAGAGGTTCCACC	SEQ ID NO.:547
	R AGGTGCTTGCAGTGAATCTG	SEQ ID NO.:548
DIK5056	F CCACCAAGGCTAATGGTAAA	SEQ ID NO.:549
	R TGGTGTGTCATCTGCATTCT	SEQ ID NO.:550
DIK5323	F CTGGGAAGCCTTTGATCTG	SEQ ID NO.:551
	R ATGGACCAGATGGTGGAAAT	SEQ ID NO.:552
DIK4862	F CTTTCCCATCCTTCACCAA	SEQ ID NO.:553
	R AAGTAGGGTGTGTGGGGTA	SEQ ID NO.:554
BMC2208	F GTTGAGCAGGGGGTAACAAAG	SEQ ID NO.:555
	R ACGAGTCCCTGCTGCTCTAC	SEQ ID NO.:556

0.5 µl PCR-product is added to 9.5 µl formamide and analysed on an ABI-3730XL sequencing Instrument (Applied Biosystems Inc.).

The calving traits considered were stillbirth (SB), calving difficulty (CD) and the size of calf at birth (CS) after first calving. The traits were assessed both as a "direct" effect (D) of the sire in the calf and as a "maternal" effect (M) of the sire in the mother of the calf, giving a total of 6 traits for the QTL analysis. Breeding values for each trait were obtained from the Danish Agricultural Advisory Service database. The breeding values were obtained from the routine breeding value estimation procedure by the exception that information from correlated traits and pedigree information were ignored.

#### Statistical Analysis

The calving traits were analyzed using the linear regression mapping procedure of Haley & Knott (1992). Significant QTL were found by using permutation tests developed by Churchill & Doerge (1994). In this procedure traits and chromosomes were analyzed separately and tested for the presence of a single QTL affecting a particular trait. If the test: (1) exceeds the 5% chromosome-wise significance threshold and (2) the QTL-region affecting two or more traits, then the QTL is retained for further characterization. The variance component QTL mapping approach was used to test if it is a single pleiotropic QTL affecting two traits or two linked QTL affecting different traits. The QTL is modeled as a random effect in a bivariate linear mixed model that adjusts for polygenic and overall trait means. The IBD matrices were computed using a recursive algorithm (Sørensen et al., 2003, Wang et al., 1995), conditional on the most likely marker linkage phase in the sire. The IBD matrices were computed for every 2 cM along the chromosomes and used in the subsequent variance component estimation procedure.

Baysian information criterion (BIC) and correlation between the QTL ( $r_q$ ) were used to compare the pleiotropic and linked model.

#### Example 1

The chromosome-wise regression test (table 24) showed a total of 27 significant QTL for calving traits in first lactation on 17 different chromosomes. 15 of the QTL were related to direct calving ease and 12 QTL was related to the maternal effects.

Average number of informative markers per grandsire family varied from 3.0 (BTA25) to 8.5 (BTA3) informative markers per chromosome.

Chromosome wise regressions analysis across families for calving traits after first calving. QTL are shown for traits that exceed 5 % chromosome wise threshold level. Numbers of segregating families are shown in brackets for each trait and chromosome.

BTA	Inform. Level	D_CD	D_SB	D_CS	M_CD	M_SB	M_CS
BTA3	8.5 <sup>a</sup> (34) <sup>b</sup>		0.010 <sup>c</sup> (5) <sup>d</sup>				
BTA4	5.0 (19)	0.023 (3)					
<b>BTA7</b>	6.4 (34)		0,003 (6)	0,042 (5)			
<b>BTA8</b>	3.6 (34)	0.042 (2)				0.030 (3)	
BTA9	6.0 (19)						0.027 (3)
BTA10	6.1 (34)				0,035 (3)		
<b>BTA12</b>	5.1 (19)		0.031 (1)			0.028 (2)	
BTA15	6.5 (34)			0,02 (3)			
<b>BTA18</b>	7.0 (34)	0.010 (5)	0.026 (4)	0 (4)		0.015 (7)	
BTA19	5.2 (19)			0.007 (3)			
BTA20	3.5 (19)			0.005 (5)			
BTA21	5.3 (34)				0.044 (2)		
<b>BTA22</b>	4.1 (19)			0.010 (2)			0.029 (3)
BTA24	4.6 (19)					0.041 (2)	
<b>BTA25</b>	3.0 (19)	0.006 (2)		0.002 (4)			
<b>BTA26</b>	4.7 (34)		0.021 (3)			0.00 (7)	
<b>BTA28</b>	3.5 (33)				0.025 (3)	0.045 (0)	

5 D\_CD: direct calving difficulty, D\_SB: direct stillbirth, D\_CS: direct calf size, M\_CD: maternal calving difficulty, M\_SB: maternal stillbirth, M\_CS: maternal calf size.

<sup>a</sup>: numbers of informative markers, <sup>b</sup> : number of analyzed grandsires, <sup>c</sup> : p-values

Each QTL was detected significant in 0 to 7 Holstein families when the test was performed within family analysis. BTA 28 showed no significant families for M\_SB, but four families were candidates to significance ( $p<0.10$ ). Seven chromosomes showed more than one significant QTL in the same region and were further examined for the presence of pleiotropic or linked QTL. Only BTA 18 showed more than two significant QTL.

15

## Example 2

Table 25 shows results of tests to distinguish between pleiotropic and linked QTL. Two regions (BTA 12, BTA25) indicate QTL with pleiotropic effects with strong correlations

between the traits (close to 1 or -1). For BTA7 and BTA26 the linkage model is in favor with correlations closer to 0 and high BIC-values. The analysis on BTA22 and BTA28 could not clarify whether it is linked or pleiotropic QTL. BTA8 did not give useful results because the likelihood did not converge to a maximum. On BTA 18 there may be a pleiotropic QTL affecting all the direct calving traits and probably one QTL affecting maternal stillbirth (M\_SB).

Table 25

Multi-trait analysis with pleiotropic and linked QTL models for calving traits on BTA 7, 8, 10 12, 18, 22, 25, 26, and 28 where QTL were identified for more than one calving trait in first lactation.

BTA		$r_q$	Dist (cM)	No. markers	Bayes factor <sup>b</sup>
BT	AT7	D_SB, D_CS	0.35	26	1.15
BT	AT8	D_CD, M_SB	NC	38	1.12
BT	AT12	D_SB, M_SB	0.99	4	0
BT	AT18	D_CD, D_SB	0.87	0	27
		D_CD, D_CS	0.93	0	109848
		D_CD, M_SB	0.71	14	0.7
		D_SB, D_CS	0.95	0	1806411
		D_SB, M_SB	NC	14	1.15
		D_CS, M_SB	0.49	14	1.15
BT	AT22	D_CS, M_SB	0.72	14	0.68
BT	AT25	D_CD, D_CS	1.00	0	548
BT	AT26	D_SB, M_SB	0.1	10	0.32
BT	AT28	M_CD, M_SB	0.78	10	0.39
					3.7

D\_CD: direct calving difficulty, D\_SB: direct stillbirth, D\_CS: direct calf size, M\_CD: maternal calving difficulty, M\_SB: maternal stillbirth, M\_CS: maternal calf size.

<sup>a</sup> : average number of informative markers between QTL, <sup>b</sup> : probability of a pleiotropic model over the linked model

Several QTL affecting both direct and maternal calving traits were identified. The QTL for D\_CD on BTA8 confirmed the result in Ashwell et al (2003) and the QTL for direct and maternal stillbirth on BTA7 and BTA18 confirmed the results in Kühn et al (2003).

20 The multi-trait and multiple QTL variance component approach detected two pleiotropic QTL affecting both direct calving size and calving difficulties, and two pleiotropic QTL

affecting both direct and maternal stillbirth. The identified QTL could have important implications for the Danish Holstein breeding program because of relative high economic weight in the combined selection index. In particular, QTL affecting survival and stillbirth without affecting calf size will be an efficient way to improve genetic 5 progress for calving traits. More marker information is needed to get a more precise characterization of the QTL, before it can be used for effective selection purposes.

**Claims**

1. A method of determining calving characteristics in a bovine subject, comprising  
detecting in a sample from said bovine subject the presence or absence of at  
least one genetic marker that is linked to at least one trait indicative of  
increased risk of stillbirth and/or increased risk of calving difficulties and/or  
increased risk of non-desired calf size, wherein said at least one genetic marker  
is located on the bovine chromosome BTA3 in a region flanked by and including  
polymorphic microsatellite markers INRA006 and BM7225 and/or  
BTA4 in the region flanked by and including polymorphic microsatellite markers  
BMS1788 and MGTG4B and/or,  
10 BTA5 in the region flanked by and including polymorphic microsatellite markers  
BMS1095 and BM2830 and/or,  
BTA7 in a region flanked by and including polymorphic microsatellite markers  
BM7160 and BL1043 and/or,  
15 BTA8 in a region flanked by and including polymorphic microsatellite markers  
IDVGA-11 and BMS836 and/or,  
BTA9 in a region flanked by and including polymorphic microsatellite markers  
BMS2151 and BMS1967 and/or,  
BTA10 in a region flanked by and including polymorphic microsatellite markers  
20 DIK2658 and BMS2614 and/or,  
BTA11 in the region flanked by and including polymorphic microsatellite  
markers BM716 and HEL13 and/or,  
BTA12 in a region flanked by and including polymorphic microsatellite markers  
BMS410 and BMS2724 and/or,  
25 BTA15 in a region flanked by and including polymorphic microsatellite markers  
BR3510 and BMS429 and/or,  
BTA18 in a region flanked by and including polymorphic microsatellite markers  
IDVGA-31 and DIK4013 and/or,  
BTA19 in a region flanked by and including polymorphic microsatellite markers  
30 BM9202 and BMS601 and/or,  
BTA20 in a region flanked by and including polymorphic microsatellite markers  
BM3517 and UWCA26 and/or,  
BTA21 in a region flanked by and including polymorphic microsatellite markers  
DIK5182 and IDVGA-30 and/or,

BTA22 in a region flanked by and including polymorphic microsatellite markers CSSM26 and BM4102 and/or,

BTA24 in a region flanked by and including polymorphic microsatellite markers BMS917 and BMS3024 and/or,

5 BTA25 in a region flanked by and including polymorphic microsatellite markers ILSTS102 and AF5 and/or,

BTA26 in a region flanked by and including polymorphic microsatellite markers BMS651 and BM7237 and/or,

BTA28 in a region flanked by and including polymorphic microsatellite markers, BMC6020 and BMC2208, , wherein the presence of said at least one genetic marker is indicative of calving characteristics of said bovine subject and/or offspring therefrom.

10 1. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA3 in the region from about 17.1 to 101.8 cM, or between genetic markers INRA006 and BM7225.

15 2. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA4 in the region from about 12.5 to 112.8 cM, or between genetic markers BMS1788 and MGTG4B.

20 3. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA5 in the region from about 0.0 to 116.9 cM, or between genetic markers BMS1095 and BM2830

25 4. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA7 in the region from about 0.0 to 135.6 cM, or between genetic markers BM7160 and BL1043.

5. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA8 in the region from about 11.3 to 122.9 cM, or between genetic markers IDVGA-11 and BMS836.

30 6. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA9 in the region from about 8.49 to 109.3 cM, or between genetic markers BMS2151 and BMS1967.

7. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA10 in the region from about 2.7 to 109.4 cM, or between genetic markers DIK2658 and BMS2614.

8. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA11 in the region from about 19.4 to 122.4 cM, or between genetic markers BM716 and HEL13.
9. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA12 in the region from about 0.0 to 109.0 cM, or between genetic markers BMS410 and BMS2724.
- 5 10. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA15 in the region from about 9.4 to 109.8 cM, or between genetic markers BR3510 and BMS429.
- 10 11. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA18 in the region from about 0.0 to 84.4 cM, or between genetic markers IDVGA-31 and DIK4013.
12. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA19 in the region from about 15 0.0 to 108.0 cM, or between genetic markers BM9202 and BMS601.
13. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA20 in the region from about 0.0 to 77.1 cM, or between genetic markers BM3517 and UWCA26.
14. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA21 in the region from about 20 5.5 to 76.8 cM, or between genetic markers DIK5182 and IDVGA-30.
15. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA22 in the region from about 0.0 to 82.9 cM, or between genetic markers CSSM26 and BM4102.
- 25 16. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA24 in the region from about 6.2 to 65.9 cM, or between genetic markers BMS917 and BMS3024.
17. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA25 in the region from about 30 7.2 to 61.7 cM, or between genetic markers ILSTS102 and AF5.
18. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA26 in the region from about 2.8 to 66.8 cM, or between genetic markers BMS651 and BM7237.

19. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA28 in the region from about 8.0 to 59.6 cM, or between genetic markers BMC6020 and BMC2208.
20. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA3 in the region from about 32.6 to 59.4 cM, or between genetic markers DIK4403 and INRA003.
- 5 21. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA3 in the region from about 77.6 to 101.8 cM, or between genetic markers DIK2702 and BM7225.
- 10 22. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA4 in the region from about 43.2 to 91.2 cM, or between genetic markers BMS2646 and BMS648.
23. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA4 in the region from about 15 52.5 to 73.4 cM, or between genetic markers TGLA116 and BM8233.
24. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA5 in the region from about 18.3 to 56.3 cM, or between genetic markers DIK4747 and BMS1617.
25. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA5 in the region from about 20 45.5 to 82.9 cM, or between genetic markers CSSM034 and BMS1248.
26. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA7 in the region from about 30.2 to 55.3 cM, or between genetic markers DIK5412 and DIK4606.
27. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA7 in the region from about 25 77.2 to 116.6 cM, or between genetic markers BMS2258 and ILSTS006.
28. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA8 in the region from about 30 41.6 to 66.0 cM, or between genetic markers BMS678 and BMS2072.
29. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA8 in the region from about 71.1 to 122.9 cM, or between genetic markers MCM64 and BMS836.

30. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA9 in the region from about 12.8 to 64.9 cM, or between genetic markers ETH225 and BMS1290.
31. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA9 in the region from about 5 50.0 to 79.2 cM, or between genetic markers UWCA9 and bms2753.
32. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA10 in the region from about 11.0 to 37.5 cM, or between genetic markers CSSM38 and DIK2000.
- 10 33. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA10 in the region from about 44.3 to 74.0 cM, or between genetic markers BMS2742 and TGLA433.
34. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA10 in the region from about 15 87.5 to 109.4 cM, or between genetic markers BMS2641 and BMS2614.
35. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA11 in the region from about 19.4 to 50.3 cM, or between genetic markers BM716 and BM7169.
- 20 36. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA11 in the region from about 61.6 to 92.2 cM, or between genetic markers BM6445 and BMS989.
37. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA12 in the region from about 50.4 to 109.0 cM, or between genetic markers BM860 and BMS2724.
- 25 38. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA12 in the region from about 63.8 to 102.0 cM, or between genetic markers BMS975 and BMS1316.
39. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA15 in the region from about 30 91.8 to 105.0 cM, or between genetic markers BMS2076 and BMS927.
40. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA15 in the region from about 98.2 to 109.8 cM, or between genetic markers BMS820 and BMS429.

41. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA18 in the region from about 30.2 to 61.2 cM, or between genetic markers INRA121 and DIK4232.
- 5 42. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA18 in the region from about 61.2 to 84.4 cM, or between genetic markers DIK4232 and DIK4013.
43. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA19 in the region from about 10 16.0 to 45.9 cM, or between genetic markers BMS745 and BP20.
44. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA19 in the region from about 47.0 to 90.0 cM, or between genetic markers IDVGA-46 and ETH3.
- 15 45. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA20 in the region from about 19.1 to 55.1 cM, or between genetic markers BMS1282 and AGLA29.
46. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA20 in the region from about 55.1 to 77.1 cM, or between genetic markers AGLA29 and UWCA26.
- 20 47. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA21 in the region from about 18.3 to 30.0 cM, or between genetic markers DIK2492 and DIK4001.
48. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA21 in the region from about 25 30.0 to 47.8 cM, or between genetic markers DIK4001 and DIK3036.
49. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA22 in the region from about 2.9 to 47.1 cM, or between genetic markers INRA026 and BM3628.
50. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA22 in the region from about 30 47.1 to 82.9 cM, or between genetic markers BM3628 and BM4102.
51. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA24 in the region from about 8.2 to 35.5 cM, or between genetic markers BM7151 and BMS1862.

52. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA24 in the region from about 35.5 to 65.9 cM, or between genetic markers BMS1862 and BMS3024.
53. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA25 in the region from about 7.2 to 31.6 cM, or between genetic markers ILSTS102 and BM737.
54. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA25 in the region from about 46.4 to 61.7 cM, or between genetic markers BMS1353 and AF5.
- 10 55. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA26 in the region from about 2.8 to 37.6 cM, or between genetic markers BMS651 and RM026.
56. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA26 in the region from about 15 43.2 to 66.8 cM, or between genetic markers RME40 and BM7237.
57. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA28 in the region from about 24.8 to 50.5 cM, or between genetic markers BL25 and DOK5056.
58. The method according to claim 1, wherein the at least one genetic marker is located in the region of the bovine chromosome BTA28 in the region from about 20 43.0 to 59.6 cM, or between genetic markers BMS2658 and BMS2208.
59. The method according to claim 1, wherein the at least one marker is a combination of genetic markers.
- 25 60. The method according to claim 1, wherein a significance level chromosome wise is at least 5%.
61. A diagnostic kit for use in detecting the presence or absence in a bovine subject of at least one genetic marker associated with bovine calving characteristics, comprising at least one oligonucleotide sequence, wherein the nucleotide sequences are selected from any of SEQ ID NO.: 1 to SEQ ID NO.: 558 and/or 30 any combination thereof.

1/29




Fig. 1

2/29




Fig. 2

3/29




Fig. 3

4/29




Fig. 4

5/29




Fig.5

6/29




Fig.6

7/29




Fig. 7

8/29




Fig.8

9/29




Fig. 9

10/29




Fig. 10

11/29




Fig. 11

12/29




Fig. 12

13/29




Fig. 13

14/29




Fig. 14

15/29




Fig. 15

16/29




Fig. 16

17/29




Fig. 17

18/29




Fig. 18

19/29




Fig. 19

20/29




Fig. 20

21/29




Fig. 21

22/29




Fig. 22

23/29




Fig. 23

24/29





Fig. 24

25/29



SUBSTITUTE SHEET (RULE 26)

Fig. 25

26/29




Fig. 26

27/29




Fig. 27

28/29




Fig. 28

29/29




Fig. 29