

Example 139

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-(2-thienyl)benzamide

Example 140

5 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-(3-thienyl)benzamide

Example 141

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl]-2-(2-thienyl)benzamide

10 Example 142

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl]-2-(2-thienyl)benzamide

Example 143

15 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl][1,1'-biphenyl]-2-carboxamide

Example 144

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl][1,1'-biphenyl]-2-carboxamide

20 Example 145

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dichlorophenyl][1,1'-biphenyl]-2-carboxamide

Example 146

25 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methyl-6-chlorophenyl][1,1'-biphenyl]-2-carboxamide

Example 147

30 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chloro-6-fluorophenyl][1,1'-biphenyl]-2-carboxamide

Example 148

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-methylphenyl][1,1'-biphenyl]-2-carboxamide

Example 149

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-chlorophenyl][1,1'-biphenyl]-2-carboxamide

Example 150

5 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-(2-pyridinyl)benzamide

Example 151

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-(3-pyridinyl)benzamide

10

Example 152

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-(4-pyridinyl)benzamide

Example 153

15 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chlorophenyl]-2-(2-pyridinyl)benzamide

Example 154

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chlorophenyl]-2-(3-pyridinyl)benzamide

Example 155

20 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chlorophenyl]-2-(4-pyridinyl)benzamide

Example 156

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl]-2-(2-pyridinyl)benzamide

25

Example 157

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl]-2-(3-pyridinyl)benzamide

Example 158

30 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl]-2-(4-pyridinyl)benzamide

Example 159

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methyl-6-fluorophenyl]-2-(2-thienyl)benzamide

Example 160

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl]-2-(2-pyridinyl)benzamide

Example 161

5 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl]-2-(3-pyridinyl)benzamide

Example 162

10 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl]-2-(4-pyridinyl)benzamide

Example 163

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-methoxypyridine-3-carboxamide

Example 164

15 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chlorophenyl]-2-methylthiopyridine-3-carboxamide

Example 165

20 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-methylphenyl]-2-methylpyridine-3-carboxamide

Example 166

25 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3,6-dimethylphenyl]-2-methylpyridine-3-carboxamide

Example 167

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl]-2-methylpyridine-3-carboxamide

Example 168

30 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chlorophenyl]-2-methylpyridine-3-carboxamide

Example 169

35 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-3-chloro-5-methylphenyl]-2-fluoropyridine-3-carboxamide

Example 170

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-3-chlorophenyl]-2-fluoropyridine-3-
carboxamide

5

Example 171

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-3-methylphenyl]-2-chloropyridine-3-
carboxamide

Example 172

10 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-3,6-dimethylphenyl]-2-chloropyridine-3-
carboxamide

Example 173

15 N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-phenyl]-3-methylpyridine-2-carboxamide

Example 174

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-3-chlorophenyl]-3-methylpyridine-2-
carboxamide

20

Example 175

N-[4-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-3-chlorophenyl]-2-chloropyridine-3-
carboxamide

Example 176

25 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-2-pyridinyl][1,1'-biphenyl]-2-carboxamide,

m.p. 278°C-281°C

Example 177

30 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-2-pyridinyl]-2-(2-thienyl)benzamide

Example 178

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-2-pyridinyl]-2-(3-thienyl)benzamide

Example 179

35 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-
ylcarbonyl)-2-pyridinyl]-2-fluorobenzamide

Example 180

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(2-pyridinyl)benzamide

Example 181

5 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(3-pyridinyl)benzamide

Example 182

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(4-pyridinyl)benzamide

10 Example 183

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(2-furanyl)benzamide

Example 184

15 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(3-furanyl)benzamide

Example 185

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-3-chloropyridine-2-carboxamide

Example 186

20 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylpyridine-3-carboxamide

Example 187

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-5-fluoro-2-methylbenzamide

25 Example 188

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chlorobenzamide

Example 189

30 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chloro-5-fluorobenzamide

Example 190

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylbenzamide

Example 191

35 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2,5-dimethylbenzamide

Example 192

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chloro-4-fluorobenzamide

Example 193

5 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chloro-6-fluorobenzamide

Example 194

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methyl-3-fluorobenzamide

10

Example 195

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-hydroxybenzamide

Example 196

15 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-acetyloxybenzamide

Example 197

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-aminobenzamide

Example 198

20 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(methylamino) benzamide

Example 199

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(dimethylamino)benzamide

25

Example 200

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-aminomethylbenzamide

Example 201

30 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(diethylamino)benzamide

Example 202

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(dimethylaminomethyl)benzamide

Example 203

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(methylthio)benzamide

Example 204

5 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chloropyridine-3-carboxamide

Example 205

10 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-fluoropyridine-3-carboxamide

Example 206

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methoxypyridine-3-carboxamide

Example 207

15 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylthiopyridine-3-carboxamide

Example 208

20 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-aminopyridine-3-carboxamide

Example 209

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylamino-pyridine-3-carboxamide

Example 210

25 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-(dimethylamino)pyridine-3-carboxamide

Example 211

30 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]thiophene-2-carboxamide

Example 212

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]thiophene-3-carboxamide

Example 213

35 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]furane-2-carboxamide

Example 214

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylthiophene-3-carboxamide

Example 215

5 N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-3-methylthiophene-2-carboxamide

Example 216

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-chlorothiophene-3-carboxamide

10 Example 217

N-[5-(6,11-Dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-2-pyridinyl]-2-methylthiophene-3-carboxamide

Example 218

15 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide

A mixture of 0.196 g of 5,11-dihydro-10H-dibenzo[b,e][1,4]diazepine, 0.155 g of N,N-diisopropylethylamine and 0.444 g of 4-[[[1,1'-biphenyl]-2-carbonyl)amino]-2-chlorobenzoyl chloride in 20 12 ml of dichloromethane is stirred at room temperature overnight. The mixture is poured into water and extracted with dichloromethane. The extract is washed with 2N K₂CO₃, H₂O, brine and dried (Na₂SO₄). The 25 solution is filtered through a thin pad of hydrous magnesium silicate. The filtrate is concentrated to dryness and the residue triturated with ether and the solvent removed. The residue is triturated with dichloromethane to give 0.31 g of solid, m.p. 158°C- 30 184°C. Anal. Found for C₃₃H₂₄ClN₃O₂ 1/2 H₂O; C, 73.7; H, 4.6; N, 7.5; Cl, 6.9.

As described for Example 218, the following compounds can be prepared by the reaction of 5,11-dihydro-10H-dibenz[b,e][1,4]diazepine with the 35 appropriate substituted or unsubstituted [(aryl-

carbonyl)amino]benzoyl chloride or the appropriate substituted or unsubstituted [(aryl-carbonyl)amino]pyridinylcarbonyl chloride.

Example 219

5 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-phenyl][1,1'-biphenyl]-2-carboxamide

Example 220

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-
10 carboxamide

Example 221

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl][1,1'-biphenyl]-2-
15 carboxamide

Example 222

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(2-thienyl)benzamide

Example 223

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(3-thienyl)benzamide
20

Example 224

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(3-furanyl)benzamide

Example 225

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(2-furanyl)benzamide
25

Example 226

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-(2-thienyl)benzamide

30 Example 227

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-(3-thienyl)benzamide

Example 228

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-carboxamide
35

Example 229

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl][1,1'-biphenyl-2-carboxamide

5

Example 230

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methyl-6-chlorophenyl][1,1'-biphenyl-2-carboxamide

Example 231

10 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chloro-6-fluorophenyl][1,1'-biphenyl-2-carboxamide

Example 232

15 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-methylphenyl][1,1'-biphenyl-2-carboxamide

Example 233

20 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chloro-6-methylphenyl][1,1'-biphenyl-2-carboxamide

Example 234

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-chlorophenyl][1,1'-biphenyl-2-carboxamide

25

Example 235

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-phenyl]-2-(2-pyridinyl)benzamide

Example 236

30 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-phenyl]-2-(3-pyridinyl)benzamide

Example 237

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-phenyl]-2-(4-pyridinyl)benzamide

Example 238

35 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(2-pyridinyl)benzamide

Example 239

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(3-pyridinyl)benzamide

Example 240

5 N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(4-pyridinyl)benzamide

Example 241

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-(2-pyridinyl)benzamide

10 Example 242

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-(3-pyridinyl)benzamide

Example 243

15 N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-(4-pyridinyl)benzamide

Example 244

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl]-2-(2-pyridinyl)benzamide

20 Example 245

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl]-2-(3-pyridinyl)benzamide

Example 246

25 N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl]-2-(4-pyridinyl)benzamide

Example 247

30 N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]phenyl]-2-phenylmethyl)benzamide

Example 248

N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]phenyl]-2-(3-chlorophenylmethyl)benzamide

Example 249

35 N-[4-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(phenylmethyl)benzamide

Example 250

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-methoxypyridine-3-carboxamide

5

Example 251

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-chlorophenyl]-2-(methylthio)pyridine-3-carboxamide

Example 252

10 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-methylpyridine-3-carboxamide

Example 253

15 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-3-methylpyridine-2-carboxamide

Example 254

20 N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methylphenyl]-2-chloropyridine-3-carboxamide

Example 255

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3,6-dimethylphenyl]-2-fluoropyridine-3-carboxamide

25

Example 256

N-[4-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-3-methyl-6-chlorophenyl]-2-chloropyridine-3-carboxamide

Example 257

30 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide,

m.p. 280°C-285°C

Example 258

35 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(2-thienyl)benzamide

Example 259

N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(3-thienyl)benzamide

Example 260

5 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(2-furanyl)benzamide

Example 261

10 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(2-pyridinyl)benzamide

Example 262

N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(3-pyridinyl)benzamide

Example 263

15 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(4-pyridinyl)benzamide

Example 264

N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(2-furanyl)benzamide

Example 265

20 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(3-furanyl)benzamide

Example 266

25 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methoxypyridine-6-carboxamide

Example 267

N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]pyridine-3-carboxamide

Example 268

30 N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-pyridinecarboxamide

Example 269

N-[5-(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methyl-5-fluorobenzamide

Example 270

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-chlorobenzamide

Example 271

5 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-chloro-5-fluorobenzamide

Example 272

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methylbenzamide

10 Example 273

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2,5-dimethylbenzamide

Example 274

15 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-chloro-4-fluorobenzamide

Example 275

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-chloro-6-fluorobenzamide

Example 276

20 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methyl-3-fluorobenzamide

Example 277

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-hydroxybenzamide

25 Example 278

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-acetyloxybenzamide

Example 279

30 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-aminobenzamide

Example 280

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(methylamino)benzamide

Example 281

35 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(aminomethyl)benzamide

Example 282

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(dimethylamino)benzamide

Example 283

5 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-chloropyridine-3-carboxamide

Example 284

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-fluoropyridine-3-carboxamide

10 Example 285

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methoxypyridine-3-carboxamide

Example 286

15 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(methylthio)pyridine-3-carboxamide

Example 287

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-aminopyridine-3-carboxamide

20 Example 288

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(methylamino)pyridine-3-carboxamide

Example 289

25 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-(dimethylamino)pyridine-3-carboxamide

Example 290

30 N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-2-methylthiophene-3-carboxamide

Example 300

N-[5-[(5,11-Dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)carbonyl]-2-pyridinyl]-3-methylthiophene-2-carboxamide

Example 301

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-phenyl][1,1'-biphenyl]-2-carboxamide

A mixture of 6,11-dihydro-5H-dibenz[b,e]
5 azepine (0.195g), 4-[(1,1'-biphenyl)-2-carbonyl)amino]
benzoyl chloride (0.41g) and 0.155 g of N,N-
diisopropylethylamine in 12 ml of dichloromethane is
stirred at room temperature for 3 hours. The mixture is
poured into water and extracted with dichloromethane.
10 The extract is washed with H₂O, saturated NaHCO₃, H₂O,
brine and dried (Na₂SO₄). The solution is filtered
through a thin pad of hydrous magnesium silicate and the
filter pad washed with dichloromethane. The filtrate is
concentrated to dryness to give 0.66 g of a yellow
15 solid. Chromatography on thick layer silica gel plates
with hexane-ethyl acetate (1.5:1) gives crystals
(0.165g) (from dichloromethane-ethyl acetate), m.p.
224°C-225°C

Example 302

20 N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide

A mixture of 0.195 g of 6,11-dihydro-5H-
dibenz[b,e]azepine, 0.444 g of 4-[(1,1'-biphenyl)-2-
carbonyl)amino]-2-chlorobenzoyl chloride and 0.155 g of
25 N,N-diisopropylethylamine is stirred at room temperature
for 3 hours. The mixture is poured into water and
extracted with dichloromethane. The extract is washed
with H₂O, saturated NaHCO₃, H₂O, brine and dried
(Na₂SO₄). The solvent is removed and the residue
30 chromatographed on thick layer silica gel plates with
solvent hexane-ethyl acetate (1.5:1) to give 0.32 g of
crystals, m.p. 120°C-125°C.

As described for Example 302, the following
compounds can be prepared by reaction of 6,11-dihydro-
35 5H-dibenz[b,e]azepine with the appropriate substituted
or unsubstituted 4-[(arycarbonyl)amino]benzoyl chloride

or the appropriate substituted or unsubstituted 6-
[(arycarbonyl)amino]pyridine-3-carbonyl chloride

Example 303

5 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]phenyl]-2-(2-thienyl)benzamide

Example 304

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]phenyl]-2-(3-thienyl)benzamide

Example 305

10 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]phenyl]-3,6-dichlorophenyl]-2-(2-
 thienyl)benzamide

Example 306

15 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3-chlorophenyl]-2-(2-thienyl)benzamide

Example 307

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3-chlorophenyl]-2-(3-thienyl)benzamide

Example 308

20 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3-chloro-6-methylphenyl]-2-(2-
 thienyl)benzamide

Example 309

25 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3-methylphenyl]-2-(2-thienyl)benzamide

Example 310

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3,6-dimethylphenyl]-2-(2-thienyl)benzamide

Example 311

30 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-carboxamide

Example 312

35 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
 carbonyl]-3,6-dimethylphenyl][1,1'-biphenyl]-2-
 carboxamide

Example 313

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-methyl-6-chlorophenyl][1,1'-biphenyl]-2-
carboxamide

5

Example 314

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chloro-6-methylphenyl][1,1'-biphenyl]-2-
carboxamide

Example 315

10 N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-methylphenyl][1,1'-biphenyl]-2-carboxamide

Example 316

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-chlorophenyl][1,1'-biphenyl]-2-carboxamide

15

Example 317

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-2-(2-pyridinyl)benzamide

Example 318

20 N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-2-(3-pyridinyl)benzamide

Example 319

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-2-(4-pyridinyl)benzamide

Example 320

25 N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl]-2-(2-pyridinyl)benzamide

Example 321

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl]-2-(3-pyridinyl)benzamide

30

Example 322

N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-methylphenyl]-2-(2-pyridinyl)benzamide

Example 323

35 N-[4-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-methylphenyl]-2-(3-pyridinyl)benzamide

Example 324

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3,6-dimethylphenyl]-2-(2-pyridinyl)benzamide

Example 325

5 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3,6-dimethylphenyl]-2-(3-pyridinyl)benzamide

Example 326

10 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3,6-dimethylphenyl]-2-(4-pyridinyl)benzamide

Example 327

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl]-[3'-methylthio-1,1'-biphenyl]-
2-carboxamide

Example 328

15 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl]-[3'-methoxy-1,1'-biphenyl]-2-
carboxamide

Example 329

20 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-[4'-dimethylamino-1,1'-biphenyl]-2-
carboxamide

Example 330

25 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-methylphenyl]-[3'-chloro-1,1'-biphenyl]-2-
carboxamide

Example 331

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl]-2-(3-furanyl)benzamide

Example 332

30 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-2-(2-furanyl)benzamide

Example 333

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]phenyl]-[3'-chloro-1,1'-biphenyl]-2-carboxamide

Example 334

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl][3'-chloro-1,1'-biphenyl]-2-
carboxamide

5

Example 335

N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-methylphenyl][3'-chloro-1,1'-biphenyl]-2-
carboxamide

Example 336

10 N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl][3'-chloro-1,1'-biphenyl]-2-
carboxamide

Example 337

15 N-[4-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-3-chlorophenyl][4'-fluoro-1,1'-biphenyl]-2-
carboxamide

Example 338

N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-2-chloropyridine-3-carboxamide

20

Example 339

N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-2-fluoropyridine-3-carboxamide

Example 340

25 N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-2-aminopyridine-3-carboxamide

Example 341

N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-2-(methylamino)pyridine-3-
carboxamide

30

Example 342

N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-2-(dimethylamino)pyridine-3-
carboxamide

Example 343

35 N-[5-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)
carbonyl]-2-pyridinyl]-3-methylthiophene-2-carboxamide

Example 344

N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-methylthiophene-3-carboxamide

Example 345

5 N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-chlorobenzamide

Example 346

N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-chloro-5-fluorobenzamide

10 Example 347

N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-chloro-6-fluorobenzamide

Example 348

15 N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-methyl-3-fluorobenzamide

Example 349

N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-(methylamino)benzamide

Example 350

20 N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-hydroxybenzamide

Example 351

N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl]-2-(aminomethyl)benzamide

25 Example 352

N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide

30 As described for Example 1, 6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepine (2 mmol) is reacted with 6-[(5-fluoro-2-methylbenzoyl)amino]pyridine-3-carbonyl chloride (2.1 mmol) in the presence of triethylamine (4 mmol) in dichloromethane to give the product as a solid, m.p. 102°C-104°C.

Example 353

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide

5 As described for Example 134, 6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepine (0.197 g) is reacted with 4-[(1,1'-biphenyl)-2-carbonyl]amino)-2-chlorobenzoyl chloride (0.444 g) in the presence of N,N-diisopropylethylamine (0.155 g) in 12 ml of
10 dichloromethane to give the product as a solid.

As described for Example 352, the following compounds can be prepared by reaction of 6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepine with the appropriate
15 substituted or unsubstituted 4-[(arylcarbonyl)amino]benzoyl chloride or the appropriate substituted or unsubstituted 6-[(arylcarbonyl)amino]pyridine-3-carbonyl chloride

Example 354

20 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl][1,1'-biphenyl]-2-carboxamide

Example 355

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-(2-thienyl)benzamide

25 Example 356

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-

carboxamide

Example 357

30 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methyl-6-chlorophenyl][1,1'-biphenyl]-2-carboxamide

Example 358

35 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3,6-dimethylphenyl][1,1'-biphenyl]-2-carboxamide

Example 359

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methylphenyl]-2-(2-thienyl)benzamide

Example 360

5 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-chlorophenyl]-2-(2-thienyl)benzamide

Example 361

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-(2-pyridinyl)benzamide

10 Example 362

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-(3-pyridinyl)benzamide

Example 363

15 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-chlorophenyl]-2-(2-pyridinyl)benzamide

Example 364

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-chlorophenyl]-2-(3-pyridinyl)benzamide

Example 365

20 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methylphenyl]-2-(2-pyridinyl)benzamide

Example 366

25 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3,6-dimethylphenyl]-2-(3-pyridinyl)benzamide

Example 367

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-(4-pyridinyl)benzamide

Example 368

30 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-chloropyridine-3-carboxamide

Example 369

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-fluoropyridine-3-carboxamide

Example 370

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-chlorophenyl]-2-chloropyridine-3-carboxamide

5

Example 371

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-methoxypyridine-3-carboxamide

Example 372

10 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methylphenyl]-2-(methylthio)pyridine-3-carboxamide

Example 373

15 N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-3-methylphenyl]-2-chloropyridine-3-carboxamide

Example 374

N-[4-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]phenyl]-2-aminopyridine-3-carboxamide

Example 375

20 N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide

Example 376

N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-2-(2-thienyl)benzamide

25

Example 377

N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-2-(3-thienyl)benzamide

Example 378

30 N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-2-(2-pyridinyl)benzamide

Example 379

N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-2-(3-pyridinyl)benzamide

Example 380

35 N-[5-[(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-2-(4-pyridinyl)benzamide

Example 381

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-(2-furanyl)benzamide

Example 382

5 N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-(3-furanyl)benzamide

Example 383

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-chlorobenzamide

10

Example 384

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-chloro-5-fluorobenzamide

Example 385

15 N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2,5-dimethylbenzamide

Example 386

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-chloro-6-fluorobenzamide

Example 387

20 N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-methyl-3-fluorobenzamide

Example 388

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-hydroxybenzamide

25

Example 389

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-aminobenzamide

Example 390

30 N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-(methylamino)benzamide

Example 391

N-[5-(6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl-2-(dimethylamino)benzamide

Example 392

N-[5-[6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl]carbonyl]-2-pyridinyl]-2-chloropyridine-3-carboxamide

5

Example 393

N-[5-[6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl]carbonyl]-2-pyridinyl]-2-(dimethylamino)pyridine-3-carboxamide

Example 394

10 N-[5-[6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl]carbonyl]-2-pyridinyl]-2-methylthiophene-3-carboxamide

Example 395

15 N-[5-[6,11-Dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl]carbonyl]-2-pyridinyl]-3-methylthiophene-2-carboxamide

Example 396

N-[4-[4,5-Dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl]carbonyl]phenyl]-[1,1'-biphenyl]-2-carboxamide

20

A mixture of 0.278 g of 4,5-dihydropyrazolo[4,3-d][1]benzazepine, 1.11 g of 4-[[[1,1'-biphenyl]-2-carbonyl]amino]benzoyl chloride and 0.426 g of N,N-diisopropylethylamine in 12 ml of dichloromethane-tetrahydrofuran (1:1) is stirred at room temperature overnight. The mixture is poured into water and extracted with dichloromethane. The extract is washed with 2N Na₂CO₃, H₂O, brine and dried (Na₂SO₄).

25

The solvent is removed under vacuum to give 1.45 g of solid. To the preceding solid in 25 ml methanol-tetrahydrofuran (1:1) is added 2.78 ml of 2N NaOH and the solution stirred at room temperature for 3.5 hours. The solvent is removed under vacuum, water added to the residue and the mixture extracted with dichloromethane. The extract is washed with H₂O, 0.5 N citric acid; H₂O, brine and dried (Na₂SO₄). The solution is passed through a thin pad of hydrous magnesiim silicate and the

30

35

filtrate concentrated to dryness. The residue (0.95 g) is triturated with ether-dichloromethane to give 0.17 g of crystals, m.p. 255°C-260°C; Anal. found for $C_{31}H_{24}N_4O_2 \cdot 1/2H_2O$: C, 75.9; H, 5.1; N, 10.8.

5

Example 397

N-[4-[(4,5-Dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide

A mixture of 0.185 g of 4,5-dihydropyrazolo[4,3-d][1]benzazepine, 0.814 g of 4-[[[1,1'-biphenyl]-2-carbonyl)amino]-2-chlorobenzoyl chloride and 0.284 g of N,N-diisopropylethylamine in 9 ml of dichloromethane-tetrahydrofuran (1:1) is stirred at room temperature overnight. The mixture is poured into water and extracted with dichloromethane. The extract is washed with 2N NaOH, H₂O, brine and dried (Na₂SO₄). The solvent is removed under vacuum to give 0.99 g of a solid. To the preceding solid in 15 ml of methanol-tetrahydrofuran (1:1) is added 1.75 ml of 2N NaOH and the solution stirred at room temperature for 2 hours. The volatiles are removed under vacuum and the mixture extracted with chloroform. The extract is washed with 1N citric acid, H₂O, brine and dried (Na₂SO₄). The solution is filtered through a thin pad of hydrous magnesium silicate and the filtrate concentrated to dryness. The residue (0.76 g) is chromatographed on thick layer silica gel plates with hexane-ethyl acetate (1:2) as solvent to give 0.37 g of white solid, m.p. 172-202°C, Anal. found for 1/2 hydrate: C, 70.1; H, 5.0; N, 9.8; Cl, 6.4.

Example 398

N-[5-[(4,5-Dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide

As described for Example 396, (0.1 mmol) of 4,5-dihydropyrazolo[4,3-d][1]benzazepine is reacted with (0.21 mmol) of 6-[[[1,1'-biphenyl]-2-carbonyl)amino]

pyridine-3-carbonyl chloride to give the product as a light tan solid. Recrystallized from ethyl acetate/hexane, m.p. 228°C-234°C as white crystals.

Example 399

5 N-[5-[(4,5-Dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide

As described for Example 396, 0.10 mmol of 4,5-dehydropyrazolo[4,3-d][1]benzazepine is reacted with 0.21 mmol) of 6-[(5-fluoro-2-methylbenzoyl)amino]-2-pyridine-3-carbonyl chloride to give the product as a tan solid, (0.15 g) m.p. 126°C-176°C: Mass Spec (FAB)-found 442(M⁺+H).

Example 400

15 N-[5-(4H-Thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-2-pyridinyl]-5-fluoro-2-methylbenzamide

As described by J.B. Press et al in J. Med. Chem., 22, 725 (1979), 9,10-dihydro-4H-thieno[3,4-b][1,5]benzodiazepin-10(9H)-one is prepared. This intermediate (4.8 g) is dissolved in tetrahydrofuran under nitrogen and 4.2 g of lithium aluminum hydride (LAH) is added portionwise with stirring. The mixture is refluxed for 18 hours and quenched by the dropwise addition of water. The mixture is extracted with chloroform and the extract is filtered through diatomaceous earth. The organic layer is washed with water (200 ml), dried (Na₂SO₄) and the solvent removed. The residual oil is chromatographed on thick layer silica gel plates with chloroform as eluent to give 1.2 g of 9,10-dihydro-4H-thieno[3,4-b][1,5]benzodiazepine as a solid. The proceeding compound (0.5 g) is reacted with 1.06 g of 6-[(5-fluoro-2-methylbenzoyl)amino]pyridine-3-carbonyl chloride, hydrochloride in dichloromethane which contains 7 ml of N,N-diisopropylethylamine. The mixture is stirred at room temperature for 16 hours and then is washed with water, 1N HCl, saturated sodium bicarbonate solution, water and

dried (Na₂SO₄). The solvent is removed and the residue purified by chromatography on silica gel with ethyl acetate-hexane (1:3) to give 1.1 g of a solid, m.p. 89°C-92°C.

5

Example 401

N-[4-(4H-Thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-phenyl][1,1'-biphenyl]-2-carboxamide

As described for Example 400, 9,10-dihydro-4H-thieno[3,4-b][1,5]benzodiazepine is reacted with 4-
10 [([1,1'-biphenyl]-2-carbonyl)amino]benzoyl chloride to give the product as a solid.

Example 402

N-[4-(4H-Thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide

15

As described for Example 400, 9,10-dihydro-4H-thieno[3,4-b][1,5]benzodiazepine is reacted with 4-
[([1,1'-biphenyl]-2-carbonyl)amino]-2-chlorobenzoyl chloride to give the product as a solid.

Example 403

20 N-[5-(4H-Thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-2-pyridinyl][1,1'-biphenyl]-2-carboxamide

As described for Example 400, 9,10-dihydro-4H-thieno[3,4-b][1,5]benzodiazepine is reacted with 6-
[([1,1'-biphenyl]-2-carbonyl)amino]-pyridine-3-carbonyl
25 chloride to give the product as a solid.

Example 404

5,11-Dihydro-10-[4-(2-thienyl)benzoyl]-10H-dibenz[b,e][1,4]diazepine

A mixture of 3 g of 4-(2-thienyl)benzoic acid
30 and 30 ml of sulfonyl chloride is refluxed for 45 minutes and the solvent removed. The residue is dissolved in carbon tetrachloride and the solvent removed under vacuum (2-times) to give 4-(2-thienyl)benzoyl chloride. To a cooled (0°C) solution of 2.0 g
35 of 5,11-dihydro-10H-dibenz[b,e][1,4]diazepine and 7 ml of N,N-diisopropylethylamine in 30 ml of dichloromethane

is added dropwise a solution of 3.15 g of 4-(2-thienyl)benzoyl chloride in 30 ml of dichloromethane. The mixture is stirred at room temperature for 16 hours and diluted with 50 ml of chloroform. The solution is washed with 30 ml each of water, 2N HCl, saturated NaHCO₃, water and dried (Na₂SO₄). The solvent is removed under vacuum and the residue (3.1 g) chromatographed on silica gel (column) with hexane-ethyl acetate (2:1) as eluent to give 1.8 g of solid, m.p. 114°C-116°C.

Example 405

5,11-Dihydro-10-[4-(3-thienyl)benzoyl]-10H-dibenz[b,e][1,4]diazepine

To a mixture of 2.0 g of 5,11-dihydro-10H-dibenz[b,e][1,4]diazepine and 7 ml of N,N-diisopropylethylamine in 30 ml of dichloromethane is added dropwise a solution of 3.15 g of 4-(3-thienyl)benzoyl chloride in 30 ml of dichloromethane. The mixture is stirred 16 hours at room temperature and diluted with dichloromethane (50 ml). The solution is washed with 30 ml each of H₂O, 2N HCl, saturated NaHCO₃, water and dried (Na₂SO₄). The solvent is removed under vacuum and the residue purified by chromatography on silica gel with hexane-ethyl acetate as eluent to give a solid.

Binding Assay to Rat Hepatic V₁ Receptors

Rat liver plasma membranes expressing the vasopressin V₁ receptor subtypes are isolated by sucrose density gradient according to the method described by Lesko et al., (1973). These membranes are quickly suspended in 50.0 mM Tris.HCl buffer, pH 7.4, containing 0.2% bovine serum albumin (BSA) and 0.1 mM phenylmethylsulfonylfluoride (PMSF) and kept frozen at -70°C until used in subsequent binding experiments. For binding experiments, the following is added to the wells of a ninety-six well format microtiter plate: 100 ml of

100.0 mM Tris.HCl buffer containing 10.0 mM MgCl₂, 0.2% heat inactivated BSA and a mixture of protease inhibitors: leupeptin, 1.0 mg%; aprotinin, 1.0 mg %; 1,10-phenanthroline, 2.0 mg %; trypsin inhibitor, 10.0 mg % and 0.1 mM PMSF, 20.0 ml of [phenylalanyl-3,4,5,-³H] vasopressin (S.A. 45.1 Ci/mmol) 0.8 mM, and the reaction initiated by the addition of 80 ml of tissue membranes containing 20 mg of tissue protein. The plates are kept undisturbed on the bench top at room temperature for 120 min. to reach equilibrium. Non-specific samples are assayed in the presence of 0.1 mM of the unlabeled antagonist phenylalanylvasopressin, added in 20.0 ml volume.

For test compounds, these are solubilized in 50% dimethylsulfoxide (DMSO) and added in 20.0 ml volume to a final incubation volume of 200 ml. Upon completion of binding, the content of each well is filtered off, using a Brandel® cell Harvester (Gaithersburg, MD). The radioactivity trapped on the filter disk by the ligand-receptor complex is assessed by liquid scintillation counting in a Packard LS Counter, with an efficiency of 65% for tritium. The data are analyzed for IC₅₀ values by the LUNDON-2 program for competition (LUNDON SOFTWARE, OH) and displayed in Table I.

Binding Assay to Rat Kidney Medullary V₂ Receptors

Medullary tissues from rat kidneys are dissected out, cut into small pieces and soaked in a 0.154 mM sodium chloride solution containing 1.0 mM EDTA with many changes of the liquid phase, until the solution is clear of blood. The tissue is homogenized in a 0.25 M sucrose solution containing 1.0 mM EDTA with many changes of the liquid phase, until the solution is clear of blood. The tissue is homogenized in a 0.25 M sucrose solution containing 1.0 mM EDTA and 0.1 mM PMSF using a Potter-Elvehjem homogenizer with a teflon pestle. The homogenate is filtered through several

layers (4 layers) of cheese cloth. The filtrate is rehomogenized using a dounce homogenizer, with a tight fitting pestle. The final homogenate is centrifuged at 1500 X g for 15 min. The nuclear pellet is discarded and the supernatant fluid recentrifuged at 40,000 x g for 30 min. The resulting pellet formed contains a dark inner part with the exterior, slightly pink. The pink outer part is suspended in a small amount of 50.0 mM Tris.HCl buffer, pH 7.4. The protein content is determined by the Lowry's method (Lowry et al., J. Biol. Chem., 1953). The membrane suspension is stored at -70°C, in 50.0 mM Tris.HCl, containing 0.2% inactivated BSA and 0.1 mM PMSF in aliquots of 1.0 ml containing 10.0 mg protein per ml of suspension until use in subsequent binding experiments.

For binding experiments, the following is added in ml volume to wells of a 96 well format of a microtiter plate: 100.0 ml of 100.0 mM Tris.HCl buffer containing 0.2 % heat inactivated BSA, 10.0 mM MgCl₂ and a mixture of protease inhibitors: leupeptin, 1.0 mg %; aprotinin, 1.0 mg % 1,10-phenanthroline, 2.0 mg %; trypsin inhibitor, and 0.1 mM PMSF, 20.0 ml of [³H] Arginine⁸, vasopressin (S.A. 75.0 Ci/mmol) at 0.8 nM and the reaction initiated by the addition of 80.0 ml of tissue membranes (200.0 mg tissue protein). The plates are left undisturbed on the bench top for 120 minutes to reach equilibrium. Non-specific binding is assessed in the presence of 1.0 mM of unlabeled ligand, added in 20 ml volume. For test compounds, these are solubilized in 50% dimethylsulfoxide (DMSO) and added in 20.0 ml volume to a final incubation volume of 200 ml. Upon completion of binding, the content of each well is filtered off, using a Brandel® cell Harvester (Gaithersburg, MD). The radioactivity trapped on the filter disk by the ligand-receptor complex is assessed by liquid scintillation counting in a Packard LS Counter, with an efficiency of

65% for tritium. The data are analyzed for IC₅₀ values by the LUNDON-2 program for competition (LUNDON SOFTWARE, OH) and displayed in Table I.

5 Radioligand Binding Experiments with Human Platelet Membranes

(a) Platelet Membrane Preparation:

Frozen platelet rich plasma (PRP), received from the Hudson Valley Blood Services, are thawed to room temperature. (Platelet Source: Hudson Valley Blood Services, Westchester Medical Center, Valhalla, NY). The tubes containing the PRP are centrifuged at 16,000 x g for 10 minutes at 4°C and the supernatant fluid discarded. The platelets resuspended in an equal volume of 50.0 mM Tris. HCL, pH 7.5 containing 120 mM NaCl and 20.0 mM EDTA. The suspension is recentrifuged at 16,000 x g for 10 minutes. This washing step is repeated one more time. The wash discarded and the lysed pellets homogenized in low ionic strength buffer of Tris. HCl, 5.0 m, pH 7.5 containing 5.0 mM EDTA. The homogenate is centrifuged at 39,000 x g for 10 minutes. The resulting pellet is resuspended in Tris.HCl buffer, 70.0 mM, pH 7.5 and recentrifuged at 39,000 x g for 10 minutes. The final pellet is resuspended in 50.0 mM Tris.HCl buffer pH 7.4 containing 120 mM NaCl and 5.0 mM KCl to give 1.0-2.0 mg protein per ml of suspension.

(b) Binding to Vasopressin V₁ receptor subtype in Human Platelet Membranes:

In wells of a 96 well format microtiter plate, add 100 µl of 50.0 mM Tris. HCl buffer containing 0.2% BSA and a mixture of protease inhibitors (aprotinin, leupeptin etc.) Then add 20 µl of [³H]Ligand (Manning or Arg⁸Vasopressin), to give final concentrations ranging from 0.01 to 10.0 nM. Initiate the binding by adding 80.0 µl of platelet suspension (approx. 100 mg protein). Mix all reagents by pipetting the mixture up

and down a few times. Non specific binding is measured in the presence of 1.0 mM of unlabeled ligand (Manning or Arg⁸Vasopressin). Let the mixture stand undisturbed at room temperature for ninety (90) minutes. Upon this
5 time, rapidly filter off the incubate under vacuum suction over GF/B filters, using a Brandel Harvester. The radioactivity caught on the filter disks is determined by the addition of liquid scintillant and counting in a liquid scintillator.

Binding to Membranes of Mouse Fibroblast Cell Line (LV-2)
Transfected with the cDNA Expressing the Human V₂ Vasopressin Receptor

(a) Membrane Preparation

5 Flasks of 175 ml capacity, continuing attached cells grown to confluence, are cleared of culture medium by aspiration. The flasks containing the attached cells are rinsed with 2 x 5 ml of phosphate buffered saline (PBS) and the liquid aspirated off each time. Finally,
10 5 ml of an enzyme free dissociation Hank's based solution (Specialty Media, Inc., Lafayette, NJ) is added and the flasks are left undisturbed for 2 minutes. The content of all flasks is poured into a centrifuge tube and the cells pelleted at 300 x g for 15 minutes. The
15 Hank's based solution is aspirated off and the cells homogenized with a polytron at setting #6 for 10 sec in 10.0 mM Tris.HCl buffer, pH 7.4 containing 0.25 M sucrose and 1.0 mM EDTA. The homogenate is centrifuged at 1500 x g for 10 minutes to remove ghost membranes.
20 The supernatant fluid is centrifuged at 100,000 x g for 60 minutes to pellet the receptor protein. Upon completion, the pellet is resuspended in a small volume of 50.0 mM Tris.HCl buffer, pH 7.4. The protein content is determined by the Lowry method and the receptor
25 membranes are suspended in 50.0 mM Tris.HCl buffer containing 0.1 mM phenylmethylsulfonylfluoride (PMSF) and 0.2% bovine serum albumin (BSA) to give 2.5 mg receptor protein per ml of suspension.

(b) Receptor Binding

30 For binding experiments, the following is added in ml volume to wells of a 96 well format of a microtiter plate: 100.0 ml of 100.0 mM Tris.HCl buffer containing 0.2% heat inactivated BSA, 10.0 mM MgCl₂ and a mixture of protease inhibitors: leupeptin, 1.0 mg%;
35 aprotinin, 1.0 mg%; 1,10-phenanthroline, 2.0 mg %; trypsin inhibitor, 10.0 mg % and 0.1 mM PMSF., 20.0 ml

of [³H] Arginine⁸, vasopressin (S.A. 75.0 Ci/mmol) at 0.8 nM and the reaction initiated by the addition of 80.0 ml of tissue membranes (200.0 mg tissue protein). The plates are left undisturbed on the bench top for 120 minutes to reach equilibrium. Non specific binding is assessed in the presence of 1.0 mM of unlabeled ligand, added in 20 ml volume. For test compounds, these are solubilized in 50% dimethylsulfoxide (DMSO) and added in 20.0 ml volume to a final incubation volume of 200 ml. Upon completion of binding, the content of each well is filtered off, using a Brandel® cell Harvester (Gaithersburg, MD). The radioactivity trapped on the filter disk by the ligand-receptor complex is assessed by liquid scintillation counting in a Packard LS Counter, with an efficiency of 65% for tritium. The data are analyzed for IC₅₀ values by the LUNDON-2 program for competition (LUNDON SOFTWARE, OH) and the data is displayed in Table I.

Vasopressin V₂ Antagonist Activity in Conscious Hydrated

20 Rats:

Conscious hydrated rats are treated with compounds under study from 0.1 to 100 mg/kg orally or vehicle. Two to four rats are used for each compound. One hour later, arginine vasopressin (AVP, antidiuretic hormone, ADH) dissolved in peanut oil is administered at 0.4 mg/kg intraperitoneally. Two rats in each test would not receive arginine vasopressin but only the vehicle (peanut oil) to serve as water-loading control. Twenty minutes later each rat is given 30 mL/kg of deionized water orally by gavage and is placed individually in a metabolic cage equipped with a funnel and a graduated glass cylinder to collect urine for four hours. Urine volume is measure and osmolality analyzed by use of a Fiske One-Ten osmometer (Fiske Assoc., Norwood, MA, USA). Urinary sodium, potassium, and chloride are analyzed by use of ion-specific electrodes

in a Beckman E3 (Electrolyte 3) Analyzer. In the following results, decreased urine volume and decreased osmolality relative to AVP-control indicates activity.

Vasopressin V₁ Antagonist Activity in Conscious Rats:

5 Conscious rats are restrained in a supine position with elastic tape. The area at the base of the tail is locally anesthetized by subcutaneous infiltration with 2% procaine (0.2 ml). Using aseptic technique to the ventral caudal tail artery is isolated and a cannula made of PE 10 and 20 (heat-fused) tubing is passed into the lower abdominal aorta. The cannula is secured, heparinized (1000 iu/cc), sealed and the wound closed with one or two stitches of Dexon 4-0. The caudal vein is also cannulated in the same manner for intravenous drug administration. The duration of the surgery is approximately 5 minutes. Additional local anesthesia (2% procaine or lidocaine) is provided as needed.

The animals are placed in plastic restraining cages in an upright position. The cannula is attached to a Statham P23Db pressure transducer and pulsatile blood pressure is recorded. Increase of systolic blood pressure responses to arginine vasopressin 0.01 and 0.2 international unit (I.U.) (350 I.U. = 1 mg) injections are recorded prior to any drug (compound) administration, after which each rat is dosed orally with compounds under study 0.1-100 mg/kg (10 cc/kg) or intravenously 0.1-30 mg/kg (1 cc/kg). The vasopressin injections are repeated 30, 60, 90, 120, 180, 240 and 300 minutes later. Percentage of antagonism by the compound is calculated using the pre-drug vasopressin vasopressor response as 100%.

Oxytocin Receptor Binding

(a) Membrane Preparation

35 Female Sprague-Dawley rats weighing approximately 200-250 g are injected intramuscularly

(i.m.) with 0.3 mg/kg of body weight of diethylstilbestrol (DES). The rats are sacrificed 18 hours later under pentobarbital anesthesia. The uteri are dissected out, cleaned of fat and connective tissues and rinsed in 50 ml of normal saline. The tissue pooled from six rats is homogenized in 50 ml of 0.01 mM Tris.HCl, containing 0.5 mM dithiothreitol and 1.0 mM EDTA, adjusted to pH 7.4, using a polytron at setting 6 with three passes of 10 sec each. The homogenate is passed through two (2) layers of cheesecloth and the filtrate centrifuged at 1000 x g for 10 minutes. The clear supernatant is removed and recentrifuged at 165,000 x g for 30 minutes. The resulting pellet containing the oxytocin receptors is resuspended in 50.0 mM Tris.HCl containing 5.0 mM MgCl₂ at pH 7.4, to give a protein concentration of 2.5 mg/ml of tissue suspension. This preparation is used in subsequent binding assays with [³H]oxytocin.




(b) Radioligand Binding

Binding of 3,5-[³H]Oxytocin ([³H]OT) to its receptors is done in microtiter plates using [³H]OT, at various concentrations, in an assay buffer of 50.0 mM Tris.HCl, pH 7.4 and containing 5.0 mM MgCl₂, and a mixture of protease inhibitors: BSA, 0.1 mg; aprotinin, 1.0 mg; 1,10-phenanthroline, 2.0 mg; trypsin, 10.0 mg; and PMSF, 0.3 mg per 100 ml of buffer solution. Non-specific binding is determined in the presence of 1.0 uM unlabeled OT. The binding reaction is terminated after 60 minutes, at 22°C, by rapid filtration through glass fiber filters using a Brandel® cell harvester (Biomedical Research and Development Laboratories, Inc., Gaithersburg, MD). Competition experiments are conducted at equilibrium using 1.0 nM [³H]OT and varying the concentration of the displacing agents. The concentrations of agent displacing 50% of [³H]OT at its

sites (IC₅₀) are calculated by a computer assisted LUNDON-2 program (LUNDON SOFTWARE INC., Ohio, USA).



Binding Assay to Rat Hepatic V₁ Receptors and Rat Kidney Medullary V₂ Receptors or *Binding to V₁ Receptor Subtype in Human Platelet and **Binding to membranes of Mouse Fibroblast Cell Line (LV-2) Transfected with the cDNA Expressing the Human V₂ Receptor

Table I

Ex. No.	Structure	V ₁ IC ₅₀ (μM)	V ₂ IC ₅₀ (μM)
1		0.24	0.054
2		0.059	0.029
4		44% at 10μM	20% at 10μM

Cont'd.

Ex. No.	Structure	V_1 IC ₅₀ (μM)	V_2 IC ₅₀ (μM)
5			
12		100% at 1μM	90% at μM
10			
15	24	*72% (μM)	**26% (10μM)
20			
25	25	100% (1μM)	39% (1μM)
25			
30			

Cont'd.			
Ex. No.	Structure	V ₁ IC ₅₀ (μM)	V ₂ IC ₅₀ (μM)
5			
26		46% (1μM)	29% (10μM)
10			
15			
27		*0.014	**1.8
20			

Cont'd.




Ex. No.	Structure	V ₁ IC ₅₀ (μM)	V ₂ IC ₅₀ (μM)
32		53% (10μM)	33% (10μM)
33		10% (10μM)	16% (10μM)
45		34% (10μM)	62% (10μM)

Table II
Vasopressin V₂ Antagonist Activity
In Conscious Hydrated Rats

	Ex. No.	Dose (mg/kg)	N	Urine Volume (ml/4 hrs)	Osmolality (mOsm/kg)
5	*		78	13.3 ± 0.3	229 ± 6
	**		6	12.1 ± 1	497 ± 53
			4	12.4 ± 0.8	361 ± 30
	***		76	2 ± 0.2	1226 ± 58
10	26	10	2	4.5	1058
	45	10	2	6.6	979
	4	10	2	6.8	878
	2	10	2	16.5	591
15	32	10	2	9.3	726
	2	10	2	16.5	591
	24	10	2	4.3	1492
	27	10	2	3.3	1317

- 20 * Water-load control
 ** Water-load
 Control +DMSO (10%)
 (20%)
- 25 *** AVP-control

Vasopressin Antidiuretic (V₂) Response in Conscious Rats
with Free Access to Water Drinking Before But Not During
the Experiment:

5

Male or female normotensive Sprague-Dawley rats (Charles River Laboratories, Inc., Kingston, NY) of 400-450 g of body weight were supplied with Laboratory Rodent Feed #5001 (PMI Feeds, Inc., Richmond, IN) and water ad libitum. On the day of test, rats were placed individually into metabolic cages equipped with stainless steel screens (to separate the feces from the urine) and funnels for collection of urine. Compounds, vehicle, or reference agent was given at various oral doses. During the test, rats were provided with no water or food. After dosing, urine was collected in graduated cylinders for four hours. Urine volume was measured. Urinary osmolality was determined using a Fiske One-Ten Osmometer (Fiske Associates, Norwood, MA 02062). An aliquot of each urine collection was analyzed for Na⁺, K⁺ and Cl⁻ using ion specific electrodes in Beckman E3 (Electrolyte 3) analyzer. The vehicle used for testing compounds was 20% dimethylsulfoxide (DMSO) in 2.5% preboiled starch.

25

Table IV list results with compounds tested by this procedure.

Table IV
Vasopressin V₂ Antagonist Activity in Conscious Rats

	Ex. No.	Dose (mg/Kg)	N	Urine Volume (ml/4 hrs)	Osmolality (MOSm/kg)
5	*		16	7-10±2	981±34
	47	10	2	22.0	394
	66	10	2	17.0	442
	67	10	2	21.5	402
	134	10	2	40.5	333
10		3	2	28	396
		1	2	18.2	596
	133	10	2	27.5	234
	135	10	2	39.5	284
		3	2	26.8	391
		1	2	19.5	526
	176	10	2	12.8	567
15	218	10	2	34	222
	257	10	2	22.5	317
	301	10	2	41.5	363
	302	10	2	40	356
	352	10	2	9.3	779
	396	10	2	21.8	238
	397	10	2	29.8	288
20	398	10	2	20.5	316
	399	10	2	17.0	404
	400	10	2	24.8	270
	404	10	2	6	909

*Control DMSO (20%)-2.5% corn starch

25 Compounds were dissolved in DMSO and then diluted in 2.5% corn starch (final concentration of DMSO was 20%). All rats were orally dosed with this mixture at 10mV kg, by gavage.

Table III
Oxytocin Binding Assay

	Ex. No.	Dose (μ M)	% Inhibition	IC ₅₀ (μ M)
5	1	1	12	-
	2	10	86	1.1
	4	10	20	-
	12	10	76	0.61
	24	10	97	1.8
10	25	10	94	0.113
	26	10	73	2.5
	27	1	83	-
	32	10	88	1.8
	33	1	37	-
15	45	1	54	-

The compounds of the present invention can be used in the form of salts derived from pharmaceutically or physiologically acceptable acids or bases. These salts include, but are not limited to, the following: salts with inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid and, as the case may be, such organic acids as acetic acid, oxalic acid, succinic acid, and maleic acid. Other salts include salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium or magnesium or with organic bases. The compounds can also be used in the form of esters, carbamates and other conventional "pro-drug" forms, which, when administered in such form, convert to the active moiety in vivo.

When the compounds are employed for the above utilities, they may be combined with one or more pharmaceutically acceptable carriers, for example, solvents, diluents and the like, and may be administered orally in such forms as tablets, capsules, dispersible

powders, granules, or suspensions containing, for example, from about 0.05 to 5% of suspending agent, syrups containing, for example, from about 10 to 50% of sugar, and elixirs containing, for example, from about 20 to 50% ethanol, and the like, or parenterally in the form of sterile injectable solutions or suspensions containing from about 0.05 to 5% suspending agent in an isotonic medium. Such pharmaceutical preparations may contain, for example, from about 25 to about 90% of the active ingredient in combination with the carrier, more usually between about 5% and 60% by weight.

The effective dosage of active ingredient employed may vary depending on the particular compound employed, the mode of administration and the severity of the condition being treated. However, in general, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.5 to about 500 mg/kg of animal body weight, preferably given in divided doses two to four times a day, or in a sustained release form. For most large mammals the total daily dosage is from about 1 to 100 mg, preferably from about 2 to 80 mg. Dosage forms suitable for internal use comprise from about 0.5 to 500 mg of the active compound in intimate admixture with a solid or liquid pharmaceutically acceptable carrier. This dosage regimen may be adjusted to provide the optimal therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

These active compounds may be administered orally as well as by intravenous, intramuscular, or subcutaneous routes. Solid carriers include starch, lactose, dicalcium phosphate, microcrystalline cellulose, sucrose and kaolin, while liquid carriers include sterile water, polyethylene glycols, non-ionic surfac-

tants and edible oils such as corn, peanut and sesame oils, as are appropriate to the nature of the active ingredient and the particular form of administration desired. Adjuvants customarily employed in the preparation of pharmaceutical compositions may be advantageously included, such as flavoring agents, coloring agents, preserving agents, and antioxidants, for example, vitamin E, ascorbic acid, BHT and BHA.

The preferred pharmaceutical compositions from the standpoint of ease of preparation and administration are solid compositions, particularly tablets and hard-filled or liquid-filled capsules. Oral administration of the compounds is preferred.

These active compounds may also be administered parenterally or intraperitoneally. Solutions or suspensions of these active compounds as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid, polyethylene glycols and mixtures thereof in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacterial and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol (e.g., glycerol, propylene glycol and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oil.

The new tricyclic non-peptide vasopressin antagonists of this invention are useful in treating conditions where decreased vasopressin levels are desired, such as in congestive heart failure, in disease
5 conditions with excess renal water reabsorption and in conditions with increased vascular resistance and coronary vasoconstriction.


In particular, the vasopressin antagonists of this invention are therapeutically useful in the
10 treatment and/or prevention of hypertension, cardiac insufficiency, coronary vasospasm, cardiac ischemia, renal vasospasm, liver cirrhosis, congestive heart failure, nephritic syndrome, brain edema, cerebral ischemia, cerebral hemorrhage-stroke, thrombosis-
15 bleeding and abnormal states of water retention.

In particular, the oxytocin antagonists of this invention are useful in the prevention of preterm labor and premature birth which is a significant cause
20 of infant health problems and infant mortality.

The claims defining the invention are as follows:

1. A compound selected from those of Formula

I:




Formula I

wherein Y is selected from (CH₂)_n, O, S, NH, NCOCH₃, N-lower alkyl (C₁-C₃), CH-lower alkyl (C₁-C₃), CHNH-lower alkyl (C₁-C₃), CHNH₂, CHN[lower alkyl (C₁-C₃)]₂, CHO-lower alkyl (C₁-C₃), CHS-lower alkyl (C₁-C₃),

wherein n is an integer from 0-2;

A-B is




wherein m is an integer from 1-2, provided that when Y is $-(CH_2)_n-$ and n=2, m may also be zero and when n is zero, m may also be three, provided also that when Y is $-(CH_2)_n-$ and n is 2, m may not also be two;

R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, -S-lower alkyl (C₁-C₃), -SH, -SO lower alkyl (C₁-C₃), -SO₂ lower alkyl (C₁-C₃), -CO-lower alkyl (C₁-C₃), -CF₃, lower alkyl (C₁-C₃), O-lower alkyl (C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl (C₁-C₃), -N-[lower alkyl (C₁-C₃)]₂, -SO₂NH₂, -SO₂NH lower alkyl (C₁-C₃), or -SO₂N[lower alkyl (C₁-C₃)]₂;


R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl (C₁-C₃), O-lower alkyl (C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;

R₃ is the moiety




wherein Ar is a moiety selected from the group


5




10



15



20








and X is O, S, -NCH₃ or -NH

R₄ is selected from hydrogen, lower alkyl (C₁-C₃), -CO-lower alkyl (C₁-C₃);


25

R₅ and R₇ are selected from hydrogen, (C₁-C₃) lower alkyl, (C₁-C₃) lower alkoxy and halogen


R₆ is selected from (a) moieties of the formula:



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



5




10 $-(CH_2)_2-O$ -lower alkyl(C₁-C₃) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$; and


(b) a moiety of the formula:

$-X-R_{10}$, wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), $-(CH_2)_p$ -cycloalkyl(C₃-C₆),


15




20



25




30




and p is zero to three;

X is C, S, NH, NCH₃,




and R₅ and R₇ are as previously defined.

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl (C₃-C₈) branched or unbranched, lower alkenyl (C₃-C₈) branched or unbranched, -O-lower alkyl (C₃-C₈) branched or unbranched, -O-lower alkenyl (C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties



or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:





15




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally

substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃) lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;


- 5 (d) a moiety selected from those of the formulae:



- wherein R_c is selected from halogen, (C₁-C₃) lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;
- 10 wherein Ar' is selected from the group:




- R₂ and R₃ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 5 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 W' is selected from O, S, NH, N-lower alkyl (C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃)
 10 R₂₅ is selected from the moieties



and the moiety $Z \bigcirc$

represents: (1) phenyl or substituted phenyl optionally substituted by one or two substituents selected from (C₁-C₃)lower alkyl, halogen, amino, (C₁-C₃)lower alkoxy, and (C₁-C₃)lower alkylamino; (2) a 5-membered aromatic (unsaturated) heterocyclic ring having one heteroatom selected from O, N and S; (3) a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom; (4) a 5 or 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen atoms; (5) a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom together with either one oxygen or one sulfur atom; wherein the 5 or 6-membered heterocyclic rings are optionally substituted by (C₁-C₃)lower alkyl, formyl, a moiety of the formula:



halogen or (C₁-C₃)lower alkoxy; and the pharmaceutically acceptable salts, esters and pro-drug forms thereof.

2. A compound according to Claim 1 wherein

the moiety $Z \bigcirc$


represents a phenyl ring optionally substituted by one or two substituents selected from (C₁-C₃)lower alkyl, halogen, amino, (C₁-C₃)lower alkoxy and (C₁-C₃)lower alkyl amino and n, m, W', X, Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

3. A compound according to Claim 1 wherein



5 represents a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom and n, m, W', X, Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

10 4. A compound according to Claim 1 wherein




represents a 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen heteroatoms and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

5. A compound according to Claim 1 wherein



20 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one sulfur heteroatom and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.


6. A compound according to Claim 1 wherein



25 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one oxygen heteroatom and Y,

A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

7. A compound according to Claim 1 wherein




5

represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

10


8. A compound according to Claim 1 wherein



15

represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

9. A compound according to Claim 1 wherein




20

represents a fused 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen and one sulfur heteroatom and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

25


10. A compound according to Claim 1 wherein




represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen and one oxygen
heteroatom and Y, A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆,
R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim


5 1.

11. A compound according to Claim 1 wherein



10 represents a phenyl ring, optionally substituted by one
or two substituents selected from (C₁-C₃) lower alkyl,
halogen, amino, (C₁-C₃) lower alkoxy and (C₁-C₃) lower
alkyl amino, Y is selected from -CH₂-,




15 

and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉,
R₁₀, R₂₅ are as previously defined in Claim 1.


12. A compound according to Claim 1 wherein

20




represents a 6-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen heteroatom, Y is
selected from -CH₂-,

25




and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

5 13. A compound according to Claim 1 wherein



10 represents a 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen heteroatoms, Y is selected from -CH₂-,




15 and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

14. A compound according to Claim 1 wherein

the moiety



represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one sulfur heteroatom wherein Y
5 is selected from -CH₂-,




and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀,
10 R₂₅ are as previously defined in Claim 1.

15. A compound according to Claim 1 wherein

the moiety




represents a 5-membered aromatic (unsaturated)
15 heterocyclic ring having one oxygen heteroatom wherein Y
is selected from -CH₂-,




and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

5 16. A compound according to Claim 1 wherein



10 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom, Y is selected from -CH₂-,






15 and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

17. A compound according to Claim 1 wherein

the moiety



represents a 5-membered aromatic (unsaturated)
heterocyclic ring having two nitrogen heteroatoms, Y is
5 selected from -CH₂-,




and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀,
R₂₅ are as previously defined in Claim 1.

10 18. A compound according to Claim wherein

the moiety




represents a 5-membered aromatic (unsaturated)
15 heterocyclic ring having one nitrogen and one sulfur
heteroatom, Y is selected from -CH₂-,




and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

5 19. A compound according to Claim 1 wherein



represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen and one oxygen heteroatom, Y is selected from -CH₂-,

10



and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

15

20. A compound according to Claim 1 wherein

the moiety



represents a phenyl ring, optionally substituted by one or two substituents selected from (C₁-C₃)lower alkyl,

- 5 halogen, amino, (C₁-C₃)lower alkoxy and (C₁-C₃)lower alkyl amino, Y is -(CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

21. A compound according to Claim 1 wherein

10

the moiety



represents a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom, Y is -

- 15 (CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

22. A compound according to Claim 1 wherein

the moiety



- 20 represents a 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen heteroatoms, Y is -(CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

25

23. A compound according to Claim 1 wherein

the moiety



represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one sulfur heteroatom, Y is -
5 (CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅,
R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in
Claim 1.

24. A compound according to Claim 1 wherein

the moiety



10

represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one oxygen heteroatom, Y is -
(CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅,
R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in
15 Claim 1.

25. A compound according to Claim 1 wherein

the moiety



20

represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen heteroatom, Y is -
(CH₂)_n, n is zero and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅,
R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in
Claim 1.

26. A compound according to claim 1 wherein

the moiety



represents a 5-membered aromatic (unsaturated)
heterocyclic ring having two nitrogen heteroatoms, Y is
5 $-(CH_2)_n$, n is zero and A-B, R_a , R_b , R_1 , R_2 , R_3 , R_4 , R_5 ,
 R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as previously defined in
Claim 1.

27. A compound according to Claim 1 wherein

the moiety



10

represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen and one sulfur
heteroatom, Y is $-(CH_2)_n$, n is zero and A-B, R_a , R_b , R_1 ,
 R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as
15 previously defined in Claim 1.

28. A compound according to Claim 1 wherein

the moiety



20


represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen and one oxygen
heteroatom, Y is $-(CH_2)_n$, n is zero and A-B, R_a , R_b , R_1 ,
 R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as
previously defined in Claim 1.

29. A compound according to Claim 1 wherein

the moiety 


represents a phenyl or substituted phenyl ring, Y is selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

30. A compound according to Claim 1 wherein

the moiety 


represents a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom, Y is selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

31. A compound according to Claim 1 wherein

the moiety 

represents a 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen heteroatoms, Y is selected from O, S, NH, NCOCH₃, and N-lower alkyl (C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

32. A compound according to Claim 1 wherein

the moiety 

represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one sulfur heteroatom, Y is
selected from O, S, NH, NCOCH₃, and N-lower alkyl
(C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈,
5 R₉, R₁₀, R₂₅ are as previously defined in Claim 1.


33. A compound according to Claim 1 wherein



represents a 5-membered aromatic (unsaturated)
10 heterocyclic ring having one oxygen heteroatom, Y is
selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃)
and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀,
R₂₅ are as previously defined in Claim 1.


34. A compound according to Claim 1 wherein

15



represents a 5-membered aromatic (unsaturated)
heterocyclic ring having one nitrogen heteroatom, Y is
selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃)
20 and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀,
R₂₅ are as previously defined in Claim 1.


35. A compound according to Claim 1 wherein



25 represents a 5-membered aromatic (unsaturated)
heterocyclic ring having two nitrogen heteroatoms, Y is
selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃)


and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

36. A compound according to Claim 1 wherein




5 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen and one sulfur heteroatom, Y is selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅,
10 R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.

37. A compound according to Claim 1 wherein




15 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen and one oxygen heteroatom, Y is selected from O, S, NH, NCOCH₃ and N-lower alkyl (C₁-C₃) and A-B, R_a, R_b, R₁, R₂, R₃, R₄, R₅,
20 R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 1.


38. A compound according to Claim wherein Y is selected from -(CH₂)_n-,





10 wherein n is an integer zero or one;
 R₃ is the moiety




15 wherein Ar is a moiety selected from the group



and R₆ is selected from the group



wherein Ar' is selected from the group



5

W' is O or S; A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X, cycloalkyl and

the moiety




10

are as previously defined in Claim 1.


39. A compound according to Claim 1 wherein Y is selected from -CH₂-,

15




10


R₃ is the moiety




wherein Ar is a moiety selected from the group



5 and R_6 is selected from the group



wherein Ar' is selected from the group



W' is O or S; A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl and


5

the moiety




are as previously defined in Claim 1.

40. a compound according to Claim 1 wherein Y
10 is selected from -CH₂-,




R₃ is the moiety




15


wherein Ar is a moiety selected from the group




and R₆ is selected from the group



wherein Ar' is selected from the group




5 and W' is O or S; and




10 represents a phenyl ring optionally substituted by one or two substituents selected from (C₁-C₃)lower alkyl, halogen, amino, (C₁-C₃)lower alkoxy and (C₁-C₃)lower alkyl amino and A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl are as previously defined in Claim 1.

15 41. A compound according to claim 1 wherein Y is -(CH₂)_n-, wherein n is an integer zero; R₃ is the moiety





wherein Ar is a moiety selected from the group

20




and R₆ is selected from the group



5

wherein Ar' is selected from the group



10 and W' is O or S; and


the moiety




represents a phenyl ring optionally substituted by one
5 or two substituents selected from (C₁-C₃)lower alkyl,
halogen, amino, (C₁-C₃)lower alkyl amino and A-B, R_a, R_b,
R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl
are as previously defined in Claim 1.

42. A compound according to Claim 1 wherein Y
10 is selected from O, S, NH, NCOCH₃ and N-lower alkyl
(C₁-C₃);


R₃ is the moiety



15 wherein Ar is a moiety selected from the group




and R_5 is selected from the group



5

wherein Ar' is selected from the group



and W' is O or S; and


the moiety



represents a phenyl ring optionally substituted by one
or two substituents selected from (C₁-C₃)lower alkyl,
5 halogen, amino, (C₁-C₃)lower alkoxy and (C₁-C₃)lower
alkyl amino and A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉,
R₁₀, R₂₅, X and cycloalkyl are as previously defined in
Claim 1.


43. A compound according to Claim 1 wherein Y
10 is -(CH₂)_n-; n is an integer zero;

R₃ is the moiety





5



wherein Ar is a moiety selected from the group




10



15





20





25


and R₆ is selected from the group




30



wherein Ar' is selected from the group



and W' is O or S; and




5

represents a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl are as previously defined in Claim 1.

10


44. A compound according to claim 1 wherein Y is -CH₂-; R₃ is the moiety




15

wherein Ar is a moiety selected from the group


5



10




15




and R_6 is selected from the group

20




25



wherein Ar' is selected from the group

30



35


and W' is O or S; and

the moiety




5 represents a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen heteroatom A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl are as previously defined in Claim 1.



10 45. A compound according to Claim 1 wherein Y is -(CH₂)_n-; n is an integer zero; R₃ is the moiety





15 wherein Ar is a moiety selected from the group




20




25




30



and R₆ is selected from the group




wherein Ar' is selected from the group




5

and W' is O or S; and



10 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one sulfur heteroatom A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl are as previously defined in Claim 1.


46. A compound according to Claim 1 wherein Y
15 is -CH₂-;
R₃ is the moiety





wherein Ar is a moiety selected from the group

20




5



10





15






and R₆ is selected from the group

20






25



wherein Ar' is selected from the group

30



35

and W' is O or S; and


the moiety



5 represents a 5-membered aromatic (unsaturated) heterocyclic ring having one sulfur heteroatom A-B, R_a, R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and cycloalkyl are as previously defined in Claim 1.


10 47. A compound according to Claim 1 wherein Y is selected from O, S, NH, NCOCH₃, N-lower alkyl (C₁-C₃);

R₃ is the moiety





15


wherein Ar is a moiety selected from the group



and R₆ is selected from the group




wherein Ar' is selected from the group



5


and W' is O or S; and



represents a 5-membered aromatic (unsaturated)
 10 heterocyclic ring having one sulfur heteroatom A-B, R_a,
 R_b, R₁, R₂, R₄, R₅, R₇, R₈, R₉, R₁₀, R₂₅, X and
 cycloalkyl are as previously defined in Claim 1.


48. A compound selected from those of the
 formulae:

15




wherein m is an integer one or two;

- R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)];
- R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;
- R₃ is the moiety:










- wherein Ar is selected from moieties of the formula:




R₅ and R₇ are selected from hydrogen, (C₁-C₃)lower alkyl, (C₁-C₃)lower alkoxy and halogen.

R₆ is selected from (a) moieties of the formula:




wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:




-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅; and

5 (b) a moiety of the formula:

-X-R₁₀, wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




10



and p is zero to three;


X is O, S, NH, NCH₃,

15



R5 and R7 are as previously defined.


(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl (C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties

10







15 or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂- lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH, R_b is as hereinbefore defined;

Ar' is a moiety selected from the group

15




R_8 and R_9 are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 5 NH-lower alkyl(C₁-C₃); N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties



10 W' is O, S, NH, N-lower alkyl(C₁-C₃), NCO-lower
 alkyl(C₁-C₃) or NSO₂-lower alkyl(C₁-C₃) or NSO₂ lower

alkyl(C₁-C₃) and the pharmaceutically acceptable salts thereof.

49. A compound selected from those of the formula:



5

wherein Y is selected from O, S, NH, and N-lower alkyl(C₁-C₃);


R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower
10 alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂, -NH lower alkyl(C₁-C₃) -SO₂NH₂; -SO₂NH lower alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
15 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;

R₃ is the moiety:




20

wherein Ar is selected from moieties of the formula:




R₅ and R₇ are selected from hydrogen, (C₁-C₃)lower alkyl, (C₁-C₃)lower alkoxy and halogen;


5 R₆ is selected from (a) moieties of the formula:




5




10





15




20



25




30



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃,





35 C₂H₅, moieties of the formulae:



- (CH₂)₂-O-lower alkyl (C₁-C₃) or -CH₂CH₂OH; q is one, two or three R_b is hydrogen, -CH₃ or -C₂H₅; and


(b) a moiety of the formula:

-X-R₁₀, wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




and p is zero to three;

X is O, S, NH, NCH₃




and R₅ and R₇ are as previously defined.

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl (C₃-C₈) branched or unbranched, lower alkenyl (C₃-C₈) branched or unbranched, -O-lower alkyl (C₃-C₈) branched or unbranched, -O-lower alkenyl (C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







15 or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH, R_b is as hereinbefore defined;

Ar' is a moiety selected from the group

15



R_8 and R_9 are independently hydrogen, lower alkyl
 (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 5 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 NH-lower alkyl(C₁-C₃); N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties



10

W' is O, S, NH, N-lower alkyl(C₁-C₃), NCO-lower alkyl(C₁-C₃) or NSO₂-lower alkyl(C₁-C₃) or NSO₂ lower alkyl(C₁-C₃) and the pharmaceutically acceptable salts thereof.

5 50. The compound according to claim 1, N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.

 51. The compound according to claim 1, N-[5-[(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-
10 pyridinyl]-2-dimethylamino pyridine-3-carboxamide.

 52. The compound according to claim 1, N-[5-[(9,10-Dihydro-4H-thieno[2,3-c][1]benzazepin-9-yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.

 53. The compound according to claim 1, N-[5-
15 [(9,10-Dihydro-4H-thieno[2,3-c][1]benzazepin-9-yl)carbonyl]-2-pyridinyl]-2-dimethylamino pyridine-3-carboxamide.

 54. The compound according to claim 1, N-[5-[(4,10-Dihydro-5H-thieno[3,2-c][1]benzazepin-5-
20 yl)carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.

 55. The compound according to claim 1, N-[5-[(4,10-Dihydro-5H-thieno[3,2-c][1]benzazepin-5-yl)carbonyl]-2-pyridinyl]-2-dimethylamino pyridine-3-carboxamide.

25 56. The compound according to claim 1, N-[5-[9,10-Dihydro-4H-thieno[2,3-c][1]benzazepin-9-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide.

 57. The compound according to claim 1, N-[5-
30 [(4,10-Dihydro-5H-thieno[3,2,-c][1]benzazepin-5-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide.

 58. The compound according to claim 1, N-[5-[(4,5-Dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)carbonyl]-2-pyridinyl]-2-dimethylamino pyridine-3-carboxamide.


59. The compound according to claim 1, *N*-[5-[(4,5-Dihydropyrazolo[4,3-*d*][1]benzazepin-6(1*H*-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide.

60. The compound according to claim 1, *N*-[5(peri[2,3-*b*][1,4]benzoxazepin-5-(6*H*)-ylcarbonyl)-2-pyridinyl]-5-fluoro-2-methyl benzamide.

5 61. A pharmaceutical composition comprising a suitable pharmaceutical carrier and an effective amount of a compound of any one of claims 1 to 60.

62. A method of treating diseases characterised by excess renal reabsorption of water as well as congestive heart failure, liver cirrhosis, nephrotic syndrome, central nervous system injuries, lung disease and hyponatremia in a mammal comprising
10 administering a compound of any one of claims 1 to 60 or of a composition of claim 61 to said mammal in an amount effective to alleviate the disease.


63. A process for preparing a compound of the formula:



Formula I

15 wherein Y is (CH₂)_n, O, S, NH, NCOCH₃, N-lower alkyl (C₁-C₃), CH-lower alkyl (C₁-C₃), CHNH-lower alkyl (C₁-C₃), CHNH₂, CHN[lower alkyl (C₁-C₃)]₂, CHO-lower alkyl (C₁-C₃), CHS-lower (C₁-C₃),
wherein n is an integer from 0-2;

A-B is




wherein m is an integer from 1-2, provided that when Y is $-(\text{CH}_2)_n\text{---}$ and $n=2$, m may also be zero and when n is


- 5 zero, m may also be three, provided also that when Y is $-(\text{CH}_2)_n\text{---}$ and n is 2, m may not also be two;

R₁ is selected from the group of hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, -S-lower alkyl(C₁-C₃), -SH, -SO lower alkyl(C₁-C₃), -SO₂ lower
10 alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂, -SO₂NH lower alkyl(C₁-C₃), -SO₂N[lower alkyl(C₁-C₃)]₂;

- 15 R₂ is selected from the group of hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy; R₃ is the moiety



- 20 wherein Ar is a moiety selected from the group









R_4 is hydrogen, lower alkyl (C1-C3); -CO-lower alkyl (C1-C3);


R_5 and R_7 are selected from the group, hydrogen,

5 (C1-C3) lower alkyl, (C1-C3) lower alkoxy and halogen

R_6 is selected from (a) moieties of the formulae:



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



5




10

-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅;


and (b) a moiety of the formula:

-X-R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈),

15 -(CH₂)_p -cycloalkyl(C₃-C₆),



20




25

30


and p is zero to three;

35 X is O, S, NH, NCH₃,




and R₅ and R₇ are as previously defined.

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







15 or -CH₂-K wherein K is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

wherein Ar' is selected from the group

15




- R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 5 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q -N(R_b)₂;
 R₂₅ is selected from the moieties




- 10 W' is selected from O, S, NH, N-lower alkyl (C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃);

the moiety $\text{Z} \bigcirc$


represents: (1) phenyl or substituted phenyl optionally substituted by one or two substituents selected from (C₁-C₃)lower alkyl, halogen, amino, (C₁-C₃)lower alkoxy, and (C₁-C₃)lower alkylamino; (2) a 5-membered aromatic (unsaturated) heterocyclic ring having one heteroatom selected from O, N and S; (3) a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom; (4) a 5 or 6-membered aromatic (unsaturated) heterocyclic ring having two nitrogen atoms; (5) a 5-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom together with either one oxygen or one sulfur atom; wherein the 5 or 6-membered heterocyclic rings are optionally substituted by (C₁-C₃)lower alkyl, formyl, a moiety of the formula:



halogen or (C₁-C₃)lower alkoxy; which comprises reacting a compound of the formulae:




with a compound of the formula:




wherein the moiety represented by the formula is an
5 aracyl chloride or an aryl carboxylic acid which has been
activated by conversion to a mixed anhydride or
activated with a peptide coupling reagent to give
compounds of the Formula I.

64. A compound selected from those of the
10 formula:




wherein R₁ is hydrogen, halogen (chlorine, bromine,
fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-
15 lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃),
-N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower
alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
20 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
methylenedioxy or ethylenedioxy;
R₃ is the moiety:




25

wherein Ar is selected from moieties of the formulae:



and X is O, S, $-NCH_3$, or $-NH$:

R is independently selected from hydrogen, lower
5 alkyl (C_1 - C_3),











$-(CH_2)_q-OH$, $-(CH_2)_q-O$ -alkyl (C_1 - C_3); q is one, two or
three;

10 R_4 is selected from hydrogen, lower alkyl (C_1 - C_3), $-CO$ -
lower alkyl (C_1 - C_3),


R_5 and R_7 are selected from hydrogen, (C_1 - C_3) lower
alkyl, (C_1 - C_3) lower alkoxy and halogen;

R₆ is selected from (a) moieties of the formulae:



5


wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:




$-(\text{CH}_2)_2\text{-O-lower alkyl}(\text{C}_1\text{-C}_3)$ or $-\text{CH}_2\text{CH}_2\text{OH}$; q is one, two or three; R_b is hydrogen, $-\text{CH}_3$ or $-\text{C}_2\text{H}_5$;

5 and (b) a moiety of the formula:

$-\text{X-R}_{10}$; wherein R_{10} is lower alkyl($\text{C}_3\text{-C}_8$), lower alkenyl($\text{C}_3\text{-C}_8$), $-(\text{CH}_2)_p\text{-cycloalkyl}(\text{C}_3\text{-C}_6)$,




10



and p is zero to three:


X is O, S, NH, NCH_3 ,




and R₅ and R₇ are as previously defined.

(c) a moiety of the formula:

5




wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched,
10 -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







15

or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

wherein Ar' is selected from the group:

15




- R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
- 5 -NH-lower alkyl(C₁-C₃); N-[lower alkyl(C₁-C₃)]₂, -N(R_b)(CH₂)_q -N(R_b)₂;
- R₂₅ is selected from the moieties.




- 10 W' is selected from O, S, NH, N-lower alkyl(C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and the pharmaceutically acceptable salts thereof.


65. A compound selected from those of the formula:




- 5 wherein R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃),
 10 -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;
 15 R₃ is the moiety:



wherein Ar is selected from moieties of the formula:



R is independently selected from hydrogen, lower alkyl (C₁-C₃),












5


-(CH₂)_q-OH, -(CH₂)_q-O-alkyl (C₁-C₃); q is one or two;
R₄ is selected from hydrogen, lower alkyl (C₁-C₃), -CO-
lower alkyl (C₁-C₃);

10 R₅ and R₇ are selected from hydrogen, (C₁-C₃) lower
alkyl, (C₁-C₃) lower alkoxy and halogen

R₆ is selected from (a) moieties of the formula:




wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl,
 5 cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃,
 C₂H₅, moieties of the formulae:




$-(CH_2)_2-O$ -lower alkyl(C₁-C₃) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$;

5 and (b) a moiety of the formula:

$-X-R_{10}$; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), $-(CH_2)_p$ -cycloalkyl(C₃-C₆),




10



and p is zero to three:


X is O, S, NH, NCH₃,



and R₅ and R₇ are as previously defined


(c) a moiety of the formula:

5




wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched,

10 -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







15

or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;


(d) a moiety selected from those of the formulae:




10

wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

wherein Ar' is selected from the group:




R_8 and R_9 are independently hydrogen, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃,
 5 -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties




10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
-NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
pharmaceutically acceptable salts thereof.


66. A compound selected from those of the
5 formula:



wherein R₁ is hydrogen, halogen (chlorine, bromine,
fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-
lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
10 alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-
[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower
alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
15 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
methylenedioxy or ethylenedioxy;
R₃ is the moiety:




20 wherein Ar is selected from moieties of the formula:



R is independently selected from hydrogen, halogen lower alkyl(C₁-C₃),


5







-(CH₂)_q-OH, -(CH₂)_q-O-alkyl(C₁-C₃); q is one or two; R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower
 10 alkyl(C₁-C₃);
 R₅ is hydrogen, -CH₃, -C₂H₅, Cl, Br, F, -O-CH₃, or -O-C₂H₅;

R₅ and R₇ are selected from hydrogen, (C₁-C₃) lower alkyl, (C₁-C₃) lower alkoxy and halogen:


R₆ is selected from (a) moieties of the formula:



5



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:





5

-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅;

and (b) a moiety of the formula:

-X-R₁₀; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl


10 (C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),



15


and p is zero to three:

X is O, S, NH, NCH₃,




and R₅ and R₇ are as previously defined

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







15 or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally

substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;


- 5 (d) a moiety selected from those of the formulae:



- wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;
- 10 wherein Ar_i is selected from the group:




R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 5 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R₂₅ is selected from the moieties




10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
 pharmaceutically acceptable salts thereof.

67. A compound selected from those of the
 5 formula:




wherein R₁ is hydrogen, halogen (chlorine, bromine,
 fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-
 10 lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
 alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
 alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-
 [lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-
 C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
 15 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
 methylenedioxy or ethylenedioxy;
 R₃ is the moiety:




20

wherein Ar is selected from moieties of the formula:



R is independently selected from hydrogen, halogen, lower alkyl(C₁-C₃),

5








-(CH₂)_q-OH, -(CH₂)_q-O-alkyl(C₁-C₃); q is one, two or three;

10 R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃);


R₅ and R₇ are selected from hydrogen, (C₁-C₃)lower alkyl, (C₁-C₃)lower alkoxy and halogen.

R₆ is selected from (a) moieties of the formula:



5


wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:






-(CH₂)₂-O-lower alkyl (C₁-C₃) or -CH₂CH₂OH; q is one or two; R_b is hydrogen, -CH₃ or -C₂H₅;

5 and (b) a moiety of the formula:

-X-R₁₀; wherein R₁₀ is lower alkyl (C₃-C₈), lower alkenyl (C₃-C₈), -(CH₂)_p-cycloalkyl (C₃-C₆),




10



and p is zero to three:


X is C, S, NH, NCH₃;

15




and R₅ and R₇ are as previously defined

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moities







15

or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃) lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

Ar' is selected from the group:

15




R_8 and R_9 are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 5 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties




10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
 pharmaceutically acceptable salts thereof.

68. A compound selected from those of the
 5 formula:



wherein A-B is




10

wherein R₁ is hydrogen, halogen (chlorine, bromine,
 fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-
 lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
 alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
 15 alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃),
 -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower
 alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;


R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are

20 methylenedioxy or ethylenedioxy;

R₃ is the moiety:




wherein Ar is selected from moieties of the formula:



R is independently selected from hydrogen, halogen, lower alkyl(C_1-C_3),

5








$-(CH_2)_q-OH$, $-(CH_2)_q-O-alkyl(C_1-C_3)$; q is one, two or three;


10 R_4 is selected from hydrogen, lower alkyl(C_1-C_3), $-CO-$ lower alkyl(C_1-C_3),

R_5 and R_7 are selected from hydrogen(C_1-C_3) lower alkyl, (C_1-C_3) lower alkoxy and halogen;

R₆ is selected from (a) moieties of the formula:




wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:




$-(CH_2)_2-O$ -lower alkyl (C_1-C_3) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$;

5 and (b) a moiety of the formula:

$-X-R_{10}$; wherein R_{10} is lower alkyl (C_3-C_8), lower alkenyl (C_3-C_8), $-(CH_2)_p$ -cycloalkyl (C_3-C_6),




10




and p is zero to three:

X is O, S, NH, NCH_3 ,




and R₅ and R₇ are as previously defined
(c) a moiety of the formula:

5




wherein J is R_a, lower alkyl(C₃-C₈) branched or
unbranched, lower alkenyl(C₃-C₈) branched or unbranched,
10 -O-lower alkyl(C₁-C₈) branched or unbranched, -O-lower
alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran,
tetrahydrothiophene,
the moieties







15

or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran,
tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:




10


wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

Ar' is selected from the group:

15




- R_8 and R_9 are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 5 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties




- 10 W' is selected from O, S, NH, N-lower alkyl(C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and pharmaceutically acceptable salts thereof.


69. A compound selected from those of the formulae:




- 5 wherein Y is $-(CH_2)_n-$ and n is an integer zero or one;
A-B is



- 10 wherein m is an integer one when n is one and m is an integer one or two when n is zero;
R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
15 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;
20 R₃ is the moiety:



- 25 wherein Ar is selected from moieties of the formula:












R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃);


R₅ and R₇ are selected from hydrogen(C₁-C₃)lower

5 alkyl(C₁-C₃) lower alkoxy and halogen;

R₆ is selected from (a) moieties of the formula:




- 5 wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:






-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅;

5 and (b) a moiety of the formula:

-X-R₁₀; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




10



and p is zero to three:

X is C, S, NH, NCH₃,


15




and R₅ and R₇ are as previously defined

(c) a moiety of the formula:

5




wherein J is R_a, lower alkyl(C₃-C₈) branched or
unbranched, lower alkenyl(C₃-C₈) branched or unbranched,
10 -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower
alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran,
tetrahydrothiophene,
the moieties







15

or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran,
tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-
 5 lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;


(d) a moiety selected from those of the formulae:




10

wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

wherein Ar' is selected from the group:




R₈ and R₉ are independently hydrogen, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃,
 5 -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R₂₅ is selected from the moieties



10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
 pharmaceutically acceptable salts thereof.

70. A compound selected from those of the
 5 formula:




wherein Y is selected from O, S, NH, and N-lower
 alkyl(C₁-C₃);

R₁ is hydrogen, halogen (chlorine, bromine, fluorine,
 10 iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower
 alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
 alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
 alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃),
 -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-
 15 C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;


R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
 methylenedioxy or ethylenedioxy;

R₃ is the moiety:



20






wherein Ar is selected from moieties of the formula:



R₄ is selected from hydrogen, lower alkyl(C₁-3), -CO-lower alkyl(C₁-C₃);

R₅ and R₇ are selected from hydrogen(C₁-C₃) lower alkyl(C₁-C₃) lower alkoxy and halogen

R₆ is selected from (a) moieties of the formula:



5


wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:




-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅;

5 and (b) a moiety of the formula:

-X-R₁₀; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




10



and p is zero to three:


X is O, S, NH, NCH₃,



15

and R₅ and R₇ are as previously defined


(c) a moiety of the formula:



5


wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties

10



or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:

15







wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃) lower alkoxy,

20

-CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;


(d) a moiety selected from those of the formulae:




5

wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

10 wherein Ar' is selected from the group:




R_8 and R_9 are independently hydrogen, lower alkyl
 (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 5 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;
 R_{25} is selected from the moieties




10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
-NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
pharmaceutically acceptable salts thereof.


71. A compound selected from those of the
5 formula:




wherein Y is O, S, NH and N-lower alkyl; and



R₁ is hydrogen, halogen (chlorine, bromine, fluorine,
iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower
alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
15 alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃),
-N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower
alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
20 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
methylenedioxy or ethylenedioxy;
R₃ is the moiety:












25
wherein Ar is selected from moieties of the formula:




R_4 is selected from hydrogen, lower alkyl(C₁-C₃),
-CO-lower alkyl(C₁-C₃)

- 5 R_5 and R_7 are selected from hydrogen(C₁-C₃)lower
alkyl(C₁-C₃) lower alkoxy and halogen;

R_6 is selected from (a) moieties of the formula:




- wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl,
 5 cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃,
 C₂H₅, moieties of the formulae:






-(CH₂)₂O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, CH₃ or -C₂H₅;

5 (b) a moiety of the formula:

-X-R₁₀; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




10



and p is zero to three:


X is O, S, NH, NCH₃,

15




and R₅ and R₇ are as previously defined

(c) a moiety of the formula:




5

wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, O-lower alkyl(C₃-C₈) branched or unbranched, O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydroguran, tetrahydrothiophene, the moieties







15 or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally

substituted with halogen, (C₁-C₃) lower alkyl, hydroxy,
 -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy,
 -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as
 hereinbefore defined;

- 5 (d) a moiety selected from those of the formulae:




- wherein R_c is selected from halogen, (C₁-C₃)lower alkyl,
 10 -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore
 defined;
 wherein Ar' is selected from the group:



R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃),
 5 -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or
 -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂,
 -N(R_b)(CH₂)_q-N(R_b)₂;


R₂₅ is selected from the moieties




10

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃),
 -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and
 pharmaceutically acceptable salts thereof.

72. A compound selected from those of the formula:




wherein Y is selected from O, S, NH, and N-lower
5 alkyl(C₁-C₃);




R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower
10 alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower
alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower
alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -
N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-
C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;

R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃),
15 O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are
methylenedioxy or ethylenedioxy;

R₃ is the moiety:




wherein Ar is selected from moieties of the formula:




R₄ is selected from hydrogen, lower alkyl(C₁-3), -CO-lower alkyl(C₁-C₃);

R₅ and R₇ are selected from hydrogen(C₁-C₃) lower alkyl(C₁-C₃) lower alkoxy and halogen

R₆ is selected from (a) moieties of the formula:



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



5




10 $-(CH_2)_2-O$ -lower alkyl(C₁-C₃) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$;


and (b) a moiety of the formula:

$-X-R_{10}$; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl (C₃-C₈), $-(CH_2)_i$, cycloalkyl(C₃-C₆),


15




20



25




30




and p is zero to three:

X is O, S, NH, NCH₃,




and R5 and R7 are as previously defined


5 (c) a moiety of the formula:



10 wherein J is R_a, lower alkyl(C3-C8) branched or unbranched, lower alkenyl(C3-C8) branched or unbranched, -O-lower alkyl(C3-C8) branched or unbranched, -O-lower alkenyl(C3-C8) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties







or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C1-C3)


lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;
 (d) a moiety selected from those of the formulae:



wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;
 wherein Ar' is selected from the group:



5




10

R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃);-N-[lower alkyl(C₁-C₃)]₂,


15 -N(R_b)(CH₂)_q-N(R_b)₂;

R₂₅ is selected from the moieties



20

25




30

W' is selected from O, S, NH, N-lower alkyl(C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and pharmaceutically acceptable salts thereof.


73. A compound according to claim 72 wherein

5




R₃ is the moiety:

10



wherein Ar is selected from moieties of the formula:

15




20

and Y, R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 72.

74. A compound according to claim 72 wherein

25




Y is O; and


R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 72.

30


75. A compound according to claim 72 wherein



5 R_3 is the moiety:




wherein Ar is selected from moieties of the formula:



15 Y is O; and

$\text{R}_a, \text{R}_b, \text{R}_c, \text{R}_1, \text{R}_2, \text{R}_3, \text{R}_4, \text{R}_5, \text{R}_6, \text{R}_7, \text{R}_8, \text{R}_9, \text{R}_{10}, \text{R}_{25}$ are as previously defined in Claim 72.


76. A compound according to claim 72 wherein




Y is NH; and

25 $\text{R}_a, \text{R}_b, \text{R}_c, \text{R}_1, \text{R}_2, \text{R}_3, \text{R}_4, \text{R}_5, \text{R}_6, \text{R}_7, \text{R}_8, \text{R}_9, \text{R}_{10}, \text{R}_{25}$ are as previously defined in Claim 72.


77. A compound according to claim 72 wherein



R_3 is the moiety:




wherein Ar is selected from moieties of the formula:



10 and Y is NH; and, R_a , R_b , R_c , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as previously defined in Claim 72.

78. A compound selected from those of Formula I:




Formula I

20 wherein Y is selected from $(\text{CH}_2)_n$, O, S, NH, NCOCH_3 , N-lower alkyl (C_1 - C_3), CH-lower alkyl(C_1 - C_3), CHNH -lower alkyl(C_1 - C_3), CHNH_2 , $\text{CHN}[\text{lower alkyl}(\text{C}_1$ - $\text{C}_3)]_2$, CHO -lower alkyl(C_1 - C_3), CHS -lower alkyl(C_1 - C_3),


wherein n is an integer from 0-2;

25 A-B is









30 wherein m is an integer from 1-2, provided that when Y is $-(\text{CH}_2)_n$ - and $n=2$, m may also be zero and when n is zero, m may also be three, provided also that when Y is $-(\text{CH}_2)_n$ - and n is 2, m may not also be two;

R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, -S-
 lower alkyl(C₁-C₃), -SH, -SO lower alkyl(C₁-C₃), -SO₂ lower alkyl(C₁-C₃),
 -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃),
 -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂,
 5 -SO₂NH₂, -SO₂NH lower alkyl (C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃),
 or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;
 R₃ is the moiety




wherein Ar is a moiety selected from the group





and X is O, S, -NCH₃ or -NH

R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃);
 R₅ and R₇ are selected from hydrogen, (C₁-C₃) lower alkyl, (C₁-C₃)lower
 alkoxy and halogen

R₆ is selected from (a) moieties of the formula:





wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅; and


(b) a moiety of the formula:

15 -X-R₁₀, wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl(C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),



and p is zero to three;


X is O, S, NH, NCH₃,



and R₅ and R₇ are as previously defined.

5


(c) a moiety of the formula:




10

wherein J is R_a, lower alkyl (C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties

15




20



25





or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:

30




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

5 (d) a moiety selected from those of the formulae:



wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

25 wherein Ar' is selected from the group:





R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂, -N(R_b)(CH₂)_q-N(R_b)₂;



W' is selected from O, S, NH, N-lower alkyl (C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃);

R₂₅ is selected from the moieties



5



10



15




and the moiety $Z \bigcirc$

represents: a 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom, wherein the 6-membered aromatic (unsaturated) heterocyclic ring having one nitrogen atom is optionally substituted by (C₁-C₃)lower alkyl, formyl, a moiety of the formula:




halogen or (C₁-C₃)lower alkoxy; and the pharmaceutically acceptable salts, esters and pro-drug forms thereof.

79. A compound selected from those of the formulae:




wherein Y is selected from -(CH₂)-, O, S, NH, NCOCH₃, N-lower alkyl (C₁-C₃), CH-lower alkyl(C₁-C₃), CHNH-lower alkyl(C₁-C₃), CHNH₂, CHN[lower alkyl(C₁-C₃)]₂, CHO-lower alkyl(C₁-C₃), CHS-lower alkyl(C₁-C₃);

25 A-B is




30 R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃),

-NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂,
 -SO₂NH₂; -SO₂NH lower alkyl (C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂;
 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃),
 or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;
 5 R₃ is the moiety:




wherein Ar is selected from moieties of the formula:




R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃);
 R₅ and R₇ are selected from hydrogen(C₁-C₃)lower alkyl(C₁-C₃) lower
 alkoxy and halogen;
 R₆ is selected from (a) moieties of the formula:

30



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



5




10 $-(CH_2)_2-O$ -lower alkyl(C₁-C₃) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$;


and (b) a moiety of the formula:

$-X-R_{10}$; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl (C₃-C₈), $-(CH_2)_p$ -cycloalkyl(C₃-C₆),


15




20



25




30




and p is zero to three:

X is O, S, NH, NCH₃,




and R₅ and R₇ are as previously defined

5 (c) a moiety of the formula:




10 wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties






or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:

30





wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:






wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined; wherein Ar' is selected from the group:




R₈ and R₉ are independently hydrogen, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂, -N(R_b)(CH₂)_q-N(R_b)₂;

R₂₅ is selected from the moieties




W' is selected from O, S, NH, N-lower alkyl(C₁-C₃), -NCO-lower alkyl(C₁-C₃), or N(SO₂-lower alkyl(C₁-C₃)); and pharmaceutically acceptable salts thereof.


5 80. A compound according to claim 79 wherein



10 R₃ is the moiety:




wherein Ar is selected from moieties of the formula:



20 and Y, R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.

81. A compound according to claim 79 wherein




Y is -(CH₂)_n-; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.

30 82. A compound according to claim 79 wherein

5




Y is O; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.

10

83. A compound according to claim 79 wherein




15


Y is NH; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.


84. A compound according to claim 79 wherein



R₃ is the moiety:




wherein Ar is selected from moieties of the formula:




Y is $-(\text{CH}_2)-$; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.

85. A compound according to claim 79 wherein




R₃ is the moiety:



wherein Ar is selected from moieties of the formula:

5




10

Y is O; and


R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previous defined in Claim 79.

86. A compound according to claim 79 wherein

15




R₃ is the moiety:



20

wherein Ar is selected from moieties of the formula:


25



Y is NH; and


R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previous defined in Claim 79.

87. A compound selected from those of Formula I:



Formula I


wherein Y is selected from $(CH_2)_n$, O, S, NH, NCOCH₃, N-
 5 lower alkyl (C₁-C₃), CH-lower alkyl (C₁-C₃), CHNH-lower
 alkyl (C₁-C₃), CHNH₂, CHN[lower alkyl (C₁-C₃)]₂, CHO-lower
 alkyl (C₁-C₃), CHS-lower alkyl (C₁-C₃),
 wherein n is an integer from 0-2;
 A-B is




10

wherein m is an integer from 1-2, provided that when Y
 is $-(CH_2)_n$ and n=2, m may also be zero and when n is
 zero, m may also be three, provided also that when Y is
 $-(CH_2)_n$ and n is 2, m may not also be two;


15 R₁ is hydrogen, halogen (chlorine, bromine, fluorine,
 iodine), OH, -S-lower alkyl (C₁-C₃), -SH, -SO lower
 alkyl (C₁-C₃), -SO₂ lower alkyl (C₁-C₃), -CO-lower
 alkyl (C₁-C₃), -CF₃, lower alkyl (C₁-C₃), O-lower
 alkyl (C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl (C₁-C₃), -N-
 20 [lower alkyl (C₁-C₃)]₂, -SO₂NH₂, -SO₂NH lower alkyl
 (C₁-C₃), or -SO₂N[lower alkyl (C₁-C₃)]₂;
 R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl (C₁-C₃),
 O-lower alkyl (C₁-C₃), or R₁ and R₂ taken together are
 methylenedioxy or ethylenedioxy;
 25 R₃ is the moiety




wherein Ar is a moiety selected from the group



- 5 and X is O, S, -NCH₃ or -NH
- R₄ is selected from hydrogen, lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃);
- R₅ and R₇ are selected from hydrogen, (C₁-C₃) lower alkyl, (C₁-C₃) lower alkoxy and halogen
- 10 R₆ is selected from (a) moieties of the formula:



wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



5




10 $-(CH_2)_2-O$ -lower alkyl(C1-C3) or $-CH_2CH_2OH$; q is one, two or three; R_b is hydrogen, $-CH_3$ or $-C_2H_5$; and


(b) a moiety of the formula:

$-X-R_{10}$, wherein R_{10} is lower alkyl(C3-C8), lower alkenyl(C3-C8), $-(CH_2)_f$ cycloalkyl(C3-C6),


15




20




25



30




and p is zero to three;
X is O, S, NH, NCH₃,




and R5 and R7 are as previously defined.

5 (c) a moiety of the formula:



10 wherein J is R_a, lower alkyl (C₃-C₈) branched or unbranched, lower alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched, tetrahydrofuran, tetrahydrothiophene, the moieties



or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran, tetrahydrothiophene or the heterocyclic ring moiety:




wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃)


lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;

(d) a moiety selected from those of the formulae:


5




10



15





20



wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;

25


wherein Ar' is selected from the group:



15 R₈ and R₉ are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂, -N(R_b)(CH₂)_q-N(R_b)₂;

W' is selected from O, S, NH, N-lower alkyl (C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃)

20 R₂₅ is selected from the moieties









25

20

15

10

5




and the moiety $Z \bigcirc$

5

represents: a 5-membered aromatic (unsaturated) heterocyclic ring having one S heteroatom wherein the 5-membered aromatic (unsaturated) heterocyclic ring is optionally substituted by (C₁-C₃)lower alkyl, formyl, a moiety of the formula:

10




15

halogen or (C₁-C₃)lower alkoxy; and the pharmaceutically acceptable salts, esters and pro-drug forms thereof.


88. A compound selected from those of the formula:

20



wherein Y is selected from $-(CH_2)-$, O, S, NH, and N-lower alkyl(C₁-C₃);


25




30

R₁ is hydrogen, halogen (chlorine, bromine, fluorine, iodine), OH, S-lower alkyl(C₁-C₃), -SH, -SO-lower alkyl(C₁-C₃), -SO₂-lower alkyl(C₁-C₃), -CO-lower alkyl(C₁-C₃), -CF₃, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), -NO₂, -NH₂, -NHCO lower alkyl(C₁-C₃), -N-[lower alkyl(C₁-C₃)]₂, -SO₂NH₂; -SO₂NH lower alkyl(C₁-C₃), or -SO₂N[lower alkyl(C₁-C₃)]₂; R₂ is hydrogen, Cl, Br, F, I, -OH, lower alkyl(C₁-C₃), O-lower alkyl(C₁-C₃), or R₁ and R₂ taken together are methylenedioxy or ethylenedioxy;


R₃ is the moiety:




5 wherein Ar is selected from moieties of the formula:




10



15












20





R₄ is selected from hydrogen, lower alkyl(C₁-3), -CO-lower alkyl(C₁-C₃);
R₅ and R₇ are selected from hydrogen(C₁-C₃) lower alkyl(C₁-C₃) lower
alkoxy and halogen

25

R₆ is selected from (a) moieties of the formula:




wherein cycloalkyl is defined as C₃ to C₆ cycloalkyl, cyclohexenyl or cyclopentenyl; R_a is hydrogen, CH₃, C₂H₅, moieties of the formulae:



-(CH₂)₂-O-lower alkyl(C₁-C₃) or -CH₂CH₂OH; q is one, two or three; R_b is hydrogen, -CH₃ or -C₂H₅;


and (b) a moiety of the formula:

15 -X-R₁₀; wherein R₁₀ is lower alkyl(C₃-C₈), lower alkenyl (C₃-C₈), -(CH₂)_p-cycloalkyl(C₃-C₆),




and p is zero to three:


X is O, S, NH, NCH₃,




5 and R₅ and R₇ are as previously defined
(c) a moiety of the formula:






10 wherein J is R_a, lower alkyl(C₃-C₈) branched or unbranched, lower
alkenyl(C₃-C₈) branched or unbranched, -O-lower alkyl(C₃-C₈) branched
or unbranched, -O-lower alkenyl(C₃-C₈) branched or unbranched,
15 tetrahydrofuran, tetrahydrothiophene,
the moieties





25 or -CH₂-K' wherein K' is halogen, -OH, tetrahydrofuran,
30 tetrahydrothiophene or the heterocyclic ring moiety:



wherein D, E, F and G are selected from carbon or nitrogen and wherein the carbon atoms may be optionally substituted with halogen, (C₁-C₃) lower alkyl, hydroxy, -CO-lower alkyl(C₁-C₃), CHO, (C₁-C₃)lower alkoxy, -CO₂-lower alkyl(C₁-C₃), and R_a and R_b are as hereinbefore defined;
 5 (d) a moiety selected from those of the formulae:




wherein R_c is selected from halogen, (C₁-C₃)lower alkyl, -O-lower alkyl(C₁-C₃) and OH; R_b is as hereinbefore defined;
 wherein Ar' is selected from the group:




15 R8 and R9 are independently hydrogen, lower alkyl (C₁-C₃), O-lower alkyl(C₁-C₃), S-lower alkyl(C₁-C₃), -CF₃, -CN, -OH, -SCF₃, -OCF₃, halogen, NO₂, amino, or -NH-lower alkyl(C₁-C₃); -N-[lower alkyl(C₁-C₃)]₂, -N(R_b)(CH₂)_q-N(R_b)₂;

R₂₅ is selected from the moieties




W' is selected from O, S, NH, N-lower alkyl(C₁-C₃), -NCO-lower alkyl(C₁-C₃), or NSO₂-lower alkyl(C₁-C₃); and pharmaceutically acceptable salts thereof.


5 89. A compound according to claim 88 wherein



10 R₃ is the moiety:




wherein Ar is selected from moieties of the formula:



20 and Y, R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.

90. A compound according to claim 88 wherein




Y is -(CH₂)-; and

30 R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.

91. A compound according to claim 88 wherein

A-B is




Y is O; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.

92. A compound according to claim 88 wherein

A-B is




Y is NH; and


R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.

93. A compound according to claim 88 wherein


A-B is



R₃ is the moiety:



wherein Ar is selected from moieties of the formula:




Y is -(CH₂)-; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.


94. A compound according to claim 88 wherein

5




R₃ is the moiety:

10



wherein Ar is selected from moieties of the formula:

15




and Y is NH;

20

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.


95. A compound according to claim 88 wherein

25




R₃ is the moiety:

30



wherein Ar is selected from moieties of the formula:



and Y is O;

R_a , R_b , R_c , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25}
 5 are as previously defined in Claim 88.

96. The compound according to claim 1, N-[4-(dibenz[b,f][1,4]oxazepin-10(11H)-ylcarbonyl)-phenyl]-[1,1'-biphenyl]-2-carboxamide.

97. The compound according to claim 1, N-[4-(dibenz[b,f][1,4]oxazepin-10(11H)-ylcarbonyl)-3-chloro-
 10 phenyl][1,1'-biphenyl]-2-carboxamide.

98. The compound according to claim 1, N-[5-(dibenz[b,f][1,4]oxazepin-10(11H)-ylcarbonyl)-2-pyridinyl]-5-fluoro-2-methylbenzamide.

99. The compound according to claim 1, N-[5-(dibenz[b,f][1,4]oxazepin-10(11H)-ylcarbonyl)-2-pyridinyl]-2-(4-pyridinyl)benzamide.
 15

100. The compound according to claim 1, N-[5-(pyrido[2,3-b][1,5]benzoxazepin-6(5H)-ylcarbonyl)-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.
 20

101. The compound according to claim 1, N-[5-(pyrido[2,3-b][1,4]benzoxazepin-5(6H)-ylcarbonyl)-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.

102. The compound according to claim 1, N-[4-(pyrido[2,3-b][1,4]benzoxazepin-5(6H)-ylcarbonyl)-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.
 25

103. The compound according to claim 1, N-[4-(6,11-dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-ylcarbonyl)-phenyl][1,1'-biphenyl]-2-carboxamide.

104. The compound according to claim 1, N-[4-(6,11-dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-yl-carbonyl)-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

105. The compound according to claim 1, N-[4-(6,11-dihydropyrido[2,3-b][1,5]benzodiazepin-6(5H)-yl-carbonyl)phenyl][1,1'-biphenyl]-2-carboxamide, hydrochloride.

106. The compound according to claim 1, N-[4-[(5,11-dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)-carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

107. The compound according to claim 1, N-[4-[(5,11-dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)-carbonyl]-phenyl][1,1'-biphenyl]-2-carboxamide.

108. The compound according to claim 1, N-[4-[(5,11-dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)-carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-carboxamide.

109. The compound according to claim 1, N-[4-[(5,11-dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)-carbonyl]-2-methylphenyl][1,1'-biphenyl]-2-carboxamide.

110. The compound according to claim 1, N-[4-[(5,11-dihydro-10H-dibenz[b,e][1,4]diazepin-10-yl)-carbonyl]-2-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

111. The compound according to claim 1, N-[4-[(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-phenyl][1,1'-biphenyl]-2-carboxamide.

112. The compound according to claim 1, N-[4-[(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

113. The compound according to claim 1, N-[4-[(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-carboxamide.

114. The compound according to claim 1, N-[4-[(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)carbonyl]-2-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

115. The compound according to claim 1, N-[5-
[(6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide.

116. The compound according to claim 1, N-[4-
5 [(6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)-
carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

117. The compound according to claim 1, N-[4-
[(6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)-
carbonyl]phenyl][1,1'-biphenyl]-2-carboxamide.

10 118. The compound according to claim 1, N-[4-
[(6,11-dihydro-5H-pyrido[2,3-b][1,4]benzodiazepin-5-yl)-
carbonyl]-3-methylphenyl][1,1'-biphenyl]-2-carboxamide.

119. The compound according to claim 1, N-[4-
[(4,5-dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)-
15 carbonyl]phenyl][1,1'-biphenyl]-2-carboxamide.

120. The compound according to claim 1, N-[4-
[(4,5-dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)-
carbonyl]-3-chlorophenyl][1,1'-biphenyl]-2-carboxamide.

121. The compound according to claim 1, N-[5-
20 [(4,5-dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)-
carbonyl]-2-pyridinyl][1,1'-biphenyl]-2-carboxamide.

122. The compound according to claim 1, N-[5-
[(4,5-dihydropyrazolo[4,3-d][1]benzazepin-6(1H)-yl)-
carbonyl]-2-pyridinyl]-5-fluoro-2-methylbenzamide.

25 123. The compound according to claim 1, N-[5-
(4H-thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-2-
pyridinyl]-5-fluoro-2-methylbenzamide.

124. The compound according to claim 1, N-[4-
(4H-thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-phenyl]-
30 [1,1'-biphenyl]-2-carboxamide.


125. The compound according to claim 1, N-[4-
(4H-thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-3-chloro-
phenyl][1,1'-biphenyl]-2-carboxamide.

35 126. The compound according to claim 1, N-[5-
(4H-thieno[3,4-b][1,5]benzodiazepin-9(10H)-yl)-2-
pyridinyl][1,1'-biphenyl]-2-carboxamide.


127. The compound according to claim 1, 5,11-dihydro-10-[4-(2-thienyl)benzoyl]-10H-dibenz[b,e][1,4]-diazepine.

128. The compound according to claim 1, 5,11-dihydro-10-[4-(3-thienyl)benzoyl]-10H-dibenz[b,e][1,4]-diazepine.

129. A compound according to claim 79 wherein




R₃ is the moiety:




10

wherein Ar is selected from moieties of the formula:



15 and Y, R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.


130. A compound according to claim 79 wherein



20 Y is -(CH₂)-; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.


131. A compound according to claim 79 wherein



Y is O; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 79.


5 132. A compound according to claim 79 wherein




Y is NH; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 79.


10 133. A compound according to claim 79 wherein



R₃ is the moiety:



wherein Ar is selected from moieties of the formula:




15


Y is $-(\text{CH}_2)-$; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 79.


134. A compound according to claim 79 wherein



5 R₃ is the moiety:




wherein Ar is selected from moieties of the formula:




Y is O; and

10 R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀,
R₂₅ are as previously defined in Claim 79.


135. A compound according to claim 79 wherein



R₃ is the moiety:




15 wherein Ar is selected from moieties of the formula:



Y is NH; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 79.


5 136. A compound according to claim 88 wherein



R₃ is the moiety:




wherein Ar is selected from moieties of the formula:



10

and Y, R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅ are as previously defined in Claim 88.


137. A compound according to claim 88 wherein



15 Y is $\text{---(CH}_2\text{)---}$; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 88.


138. A compound according to claim 88 wherein



5 Y is O; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 88.


139. A compound according to claim 88 wherein




10 Y is NH; and

R_a, R_b, R_c, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, R₉, R₁₀, R₂₅
are as previously defined in Claim 88.


140. A compound according to claim 88 wherein



15 R₃ is the moiety:




wherein Ar is selected from moieties of the formula:



Y is $-(\text{CH}_2)-$; and

R_a , R_b , R_c , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as previously defined in Claim 88.


5 141. A compound according to claim 88 wherein



R_3 is the moiety:



wherein Ar is selected from moieties of the formula:




10

and Y is NH;


R_a , R_b , R_c , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as previously defined in Claim 88.

142. A compound according to claim 88 wherein




15

R_3 is the moiety:



wherein Ar is selected from moieties of the formula:



and Y is O;

5 R_a , R_b , R_c , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{25} are as previously defined in Claim 88.

143. A tricyclic benzazepine vasopressin antagonist, substantially as hereinbefore described with reference to any one of the Examples.

144. A pharmaceutical composition comprising a suitable pharmaceutical carrier
10 and an effective amount of a compound of any one of claims 64 to 143.

145. A method of treating diseases characterised by excess renal reabsorption of water as well as congestive heart failure, liver cirrhosis, nephrotic syndrome, central nervous system injuries, lung disease and hyponatremia in a mammal comprising administering a compound of any one of claims 64 to 143, or of a composition of claim
15 144 to said mammal in an amount effective to alleviate the disease.

146. A process for the preparation of a tricyclic benzazepine vasopressin antagonist, substantially as hereinbefore described with reference to any one of the Examples.

20 147. Use of a compound of any one of claims 1 to 60 or 64 to 143 for the manufacture of a medicament for treating diseases characterised by excess renal reabsorption of water as well as congestive heart failure, liver cirrhosis, nephrotic syndrome, central nervous system injuries, lung disease and hyponatremia.

25 148. A compound of any one of claims 1 to 60 or 64 to 143 when used for treating diseases characterised by excess renal reabsorption of water as well as congestive heart failure, liver cirrhosis, nephrotic syndrome, central nervous system injuries, lung disease and hyponatremia in a mammal.

Dated 10 December, 1999
American Cyanamid Company

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON