



- (51) International Patent Classification:
C07D 487/00 (2006.01)
- (21) International Application Number:
PCT/US2013/059108
- (22) International Filing Date:
11 September 2013 (11.09.2013)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/701,242 14 September 2012 (14.09.2012) US
- (71) Applicant: **DOW AGROSCIENCES LLC** [US/US];
9330 Zionsville Road, Indianapolis, IN 46268 (US).
- (72) Inventors: **BLAND, Douglas C.**; 331 Broadhead Drive,
Midland, MI 48642 (US). **HAMILTON, Christopher T.**;
3708 East Acorn Lane, Midland, MI 48642 (US).
- (74) Agent: **ASAM, Michael**; Dow AgroSciences LLC, 9330
Zionsville Rd, Indianapolis, Indiana 46268 (US).
- (81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,

KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

- (84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

(54) Title: AN IMPROVED PROCESS FOR THE PREPARATION OF 2-AMINO-5,8-DIMETHOXY[1,2,4]TRIAZOLO[1,5-C]PYRIMIDINE FROM 4-AMINO-2,5-DIMETHOXPYRIMIDINE

(57) Abstract: 5-Substituted-8-alkoxy[1,2,4]triazolo[1,5-c]pyrimidin-2-amines are manufactured from 4-amino-2,5-dialkoxypyrimidines in an improved process in which the generation of gaseous by-products is controlled by the continuous addition of hydroxylamine as a free base.



AN IMPROVED PROCESS FOR THE PREPARATION OF 2-AMINO-5,8-DIMETHOXY[1,2,4]TRIAZOLO[1,5-*c*]PYRIMIDINE FROM 4-AMINO-2,5-DIMETHOXYPYRIMIDINE


Background

5 Provided herein is an improved process for the preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine from 4-amino-2,5-dimethoxypyrimidine.

U.S. Patent 8,143,395 B2 describes a method to manufacture certain 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidines that avoids hydrazine and cyanogen halide. In this process a 4-amino-2,5-dialkoxypyrimidine is reacted with, for example, ethoxy carbonyl-
10 isothiocyanate (S=C=N-CO₂Et) in a polar aprotic solvent to provide an ethyl [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate, which in turn is reacted with an hydroxylamine salt in the presence of a base to provide an ethyl [(2,5-dialkoxypyrimidin-4-yl)amino](hydroxyimino)methylcarbamate. Upon heating, this intermediate cyclizes via a series of reactions to provide the desired 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]-
15 pyrimidines. While the process described in U.S. Patent 8,143,395 B2 eliminates the need for hydrazine and cyanogen halide, it introduces new challenges with respect to the generation of several equivalents of gaseous by-products. It would be desirable to have a more streamlined and safer process by which to produce 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidines.

20 Summary

Provided herein is an improved process for the preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine from 4-amino-2,5-dimethoxypyrimidine. In some embodiments, 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidines of the formula (I),




I

in which

R represents C₁-C₄ alkyl

are prepared in a process which comprises:


i) contacting a 4-amino-2,5-dialkoxypyrimidine of the formula



5


in which R is as previously defined

with a C₁-C₄ alkoxy carbonylthiocyanate of the formula



in which R is as previously defined,

10 in a polar or nonpolar aprotic solvent at a temperature from about 60 °C to about 110 °C to provide a [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate of the formula



in which R is as previously defined; and


ii) continuously adding to the [(2,5-dialkoxypyrimidin-4-yl)amino]carbono-thioylcarbamate

15 in the polar or nonpolar aprotic solvent an aqueous solution of hydroxylamine at a temperature from about 60 °C to about 110 °C to provide the 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidine.

Detailed Description

The term alkyl and derivative terms such as alkoxy, as used herein refer to straight chain or branched chain groups. Typical alkyl groups are methyl, ethyl, propyl, 1-methylethyl, butyl, 1,1-dimethylethyl and 1-methylpropyl. Methyl and ethyl are often preferred.

The present invention concerns an improved process for the preparation of a 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidine from 4-amino-2,5-dialkoxypyrimidine. It is particularly useful for the preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine from 4-amino-2,5-dimethoxypyrimidine.



U.S. Patent 8,143,395 B2 teaches that, in the first step of the reaction, the alkoxy carbonylthiocyanate should be added to the 4-amino-2,5-dialkoxypyrimidine in a polar aprotic solvent, preferably acetonitrile or ethyl acetate, at a temperature from 0 °C to room temperature. It has now been found that the reaction of the alkoxy carbonylthiocyanate need not be conducted at or below room temperature and that nonpolar aprotic solvents such as toluene perform just as well as polar aprotic solvents and offer advantages with respect to recovery and recycle.

Similarly, U.S. Patent 8,143,395 B2 recommends using at least an equivalent of hydroxylamine, preferably as a salt, with a base (not uncommonly up to 4 equivalents each of hydroxylamine salt and base) at a temperature between 0 °C and 35 °C, preferably at room temperature. The use of hydroxylamine as a salt with a base suggests the approach of frontloading the entire amount of hydroxylamine reagent. Low temperatures would be needed to control the release of gaseous by-products generated during the cascade of cyclization reactions. By continuously adding an aqueous solution of hydroxylamine as a free base, higher temperatures from about 60 °C to about 110 °C can be safely used and the release of gaseous by-products mitigated by the controlled addition of hydroxylamine.

Polar or nonpolar aprotic solvents include both aromatic and aliphatic hydrocarbons and halogenated hydrocarbons, esters, nitriles and amides.

The first step of the present invention (*i*) concerns the conversion of a 4-amino-2,5-dialkoxypyrimidine to a [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate. This is accomplished using at least one equivalent and preferably a small excess of C₁-C₄ alkoxy carbonylthiocyanate in a polar or a nonpolar aprotic solvent. In certain embodiments, the polar or nonpolar aprotic solvent is acetonitrile, ethyl acetate or toluene. The alkoxy carbonylthiocyanate is added at a temperature from about ambient to about 110 °C; the mixture is then heated at a temperature from about 60 °C to about 110 °C. In certain embodiments, the reaction mixture is heated to the reflux temperature of the solvent(s). In certain embodiments, the temperature is at least 60 °C, at least 70 °C, at least 80 °C, at least 90 °C, or at least 100 °C. The product can be isolated by conventional techniques, such as by filtration of a precipitated or crystallized material, but is generally used as is in the next step.

In certain embodiments, the 4-amino-2,5-dialkoxypyrimidine is dissolved or suspended in the polar or nonpolar aprotic solvent and then treated with the appropriate amount of the C₁-C₄ alkoxy carbonylthiocyanate. After heating to reflux, the reaction mixture can be cooled for storage or used immediately for step (*ii*).

The second step of the present invention (*ii*) concerns the conversion of the [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate to the 2-amino-5,8-dimethoxy[1,2,4]-triazolo[1,5-*c*]pyrimidine. This is accomplished using at least an equivalent, and in certain embodiments, a small excess, of hydroxylamine as the free base. The aqueous solution of hydroxylamine is continuously added to the mixture of [(2,5-dialkoxypyrimidin-4-


yl)amino]carbonothioyl-carbamate in a polar or a nonpolar aprotic solvent from step (i) at a temperature from about 60 °C to about 110 °C at a rate slow enough to safely handle the release of gaseous by-products generated during the cascade of cyclization reactions leading to the 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidine. In certain embodiments, the temperature is at least 60 °C, at least 70 °C, at least 80 °C, at least 90 °C, or at least 100 °C. The reaction mixture is cooled and can be optionally treated with sodium sulfite as a deodorizer. In some embodiments, the product is isolated by conventional techniques, such as collection by filtration and drying.

The described embodiments and following examples are for illustrative purposes and are not intended to limit the scope of the claims. Other modifications, uses, or combinations with respect to the compositions described herein will be apparent to a person of ordinary skill in the art without departing from the spirit and scope of the claimed subject matter.

EXAMPLES

All reagents described were obtained commercially and used without additional purification.

Example 1. Preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine (Ia).



To a 700 milliliter (mL) jacketed vessel equipped with a mechanical stirrer, a dual pH/temperature probe, a nitrogen inlet, and a reflux condenser was added sequentially 19.4 grams (g; 0.125 moles (mol)) of 4-amino-2,5-dimethoxypyrimidine followed by 151 g (1.717 mol) of ethyl acetate. The reaction mixture was heated to reflux (~80 °C) and then 18.7 g (0.143 mol) of ethoxy carbonylthiocyanate was continuously added via addition funnel over a 22 minute (min) period. The addition funnel was rinsed with 1.8 g of ethyl acetate and then the reaction mixture was heated at reflux for 9 hours (h). The reaction mixture was cooled to 25 °C and allowed to stand overnight. The mixture was heated to 60 °C and then 100 g (5.500 mol) of deionized water was added to the mixture. After heating to reflux (~71

°C), 9.4 g (0.142 mol) of a 50 weight percent (wt%) aqueous hydroxylamine solution was continuously added over a 1 h period. During the course of the amine addition, the reaction pH rose from 4.00 to 6.60. The reaction mixture was then heated an additional 3 h during which time the reaction pH naturally lowered to 6.30. To this reaction mixture was added a solution of 4.5 g (0.036 mol) of sodium sulfite in 20 g (1.110 mol) of deionized water over an 8 min period. The reaction pH rose from 6.30 to 7.44 during the sodium sulfite addition. The reaction mixture was then cooled to ambient temperature and allowed to stand overnight. The reaction mixture was suction filtered through a medium coarse fritted glass funnel (filtration time less than 2.0 min), and then the reaction vessel was washed with 150 g of deionized water and this rinse was used to wash the isolated product cake. A final displacement wash with 30 g of fresh deionized water was performed and the product was partially dried by suction for 3.5 h to afford 18.01 g of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine as a light cream colored wet cake. NMR analysis (using benzyl acetate as an internal standard) indicated a 91.6% purity which corresponds to a 67.5% yield. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 3.89 (s, 3H), 4.06 (s, 3H), 6.27 (br s, 2H), 7.47 (s, 1H); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 55.37, 57.03, 123.06, 138.59, 143.72, 148.49, 166.01.

Example 2. Preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine (Ia)

To a 700 mL jacketed vessel equipped with a mechanical stirrer, a dual pH/temperature probe, a nitrogen inlet, and a reflux condenser was added sequentially 19.4 g (0.125 mol) of 4-amino-2,5-dimethoxypyrimidine followed by 151 g (1.717 mol) of ethyl acetate. The reaction mixture was heated to reflux (~80 °C) and then 18.7 g (0.143 mol) of ethoxy carbonylisothiocyanate was continuously added via addition funnel over an 8 min period. The addition funnel was rinsed with 1.8 g of ethyl acetate and then the reaction mixture was heated at reflux for 9 h. The reaction mixture was cooled to 25 °C and allowed to stand overnight. The mixture was heated to 60 °C and then 100 g (5.50 mol) of deionized water was added to the mixture. After heating to reflux (~71 °C), 9.4 g (0.142 mol) of a 50 wt% aqueous hydroxylamine solution was continuously added over a 46 min period. During the course of the amine addition, the reaction pH rose from 4.20 to 6.35. The reaction mixture was then heated an additional 3 h during which time the reaction pH naturally lowered to 6.30. To this reaction mixture was added a solution of 4.5 g (0.036 mol) of sodium sulfite in 20 g (1.110 mol) of deionized water over a 6 min period. The reaction pH

rose from 6.35 to 7.51 during the sodium sulfite addition. The reaction mixture was then cooled to 18 °C and allowed to stir an additional 30 min at this temperature. The reaction mixture was suction filtered through a medium coarse fritted glass funnel (filtration time less than 2.0 min), and then the reaction vessel was washed with 30 g of deionized water and this
5 rinse was used to wash the isolated product cake. A final displacement wash with 30 g of fresh deionized water was performed and the product was partially dried by suction for 30 min and then allowed to dry overnight under a gentle stream of nitrogen to afford 19.1 g of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine as a light cream colored wet cake. NMR analysis (using benzyl acetate as an internal standard) indicated a 88.25% purity of 2-
10 amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine active which corresponds to a 69.1% yield.

Example 3. Preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine (Ia)

To a 700 mL jacketed vessel equipped with a mechanical stirrer, a dual pH/temperature probe, a nitrogen inlet, and a reflux condenser was added sequentially 19.4 g
15 (0.125 mol) of 4-amino-2,5-dimethoxypyrimidine followed by 151 g (1.639 mol) of toluene. The reaction was heated to gentle reflux (~80 °C) and then 19.24 g (0.144 mol) of 98% ethoxy carbonylisothiocyanate was added, and the reaction mixture was heated at gentle reflux (89 °C) for 7 h. The reaction mixture was cooled to 26 °C and allowed to stand overnight. The mixture was heated to 60 °C and then 100 g (5.500 mol) of deionized water
20 was added to the mixture. After heating to reflux (~69 °C), 9.6 g (0.145 mol) of a 50 wt% aqueous hydroxylamine solution was continuously added over a 1 h period. During the course of the amine addition, the reaction pH rose from 4.00 to 6.67. After completing addition of hydroxylamine, the reaction mixture was heated to 77 °C and then stirred an additional 3 h during which time the reaction pH naturally raised to 7.42. To this reaction
25 mixture was added a solution of 4.5 g (0.036 mol) of sodium sulfite in 20 g (1.110 mol) of deionized water over a one min period. The reaction pH rose from 7.34 to 7.81 during the sodium sulfite addition. The reaction mixture was stirred an additional 1 h at 77 °C and then cooled to ambient temperature and allowed to stand overnight. The reaction mixture was suction filtered through a medium coarse fritted glass funnel (filtration time was 43 min), and
30 then the reaction vessel was washed with 30 g of deionized water and this rinse was used to wash the isolated product cake. A final displacement wash with 30 g of fresh deionized water was performed and the product was dried to afford 19.62 g of 2-amino-5,8-


dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine as a light cream colored solid. NMR analysis (using benzyl acetate as an internal standard) indicated an 83.3% purity of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine active which corresponds to a 67.0% yield.

Example 4. Preparation of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine (Ia)

5 To a 700 mL jacketed vessel equipped with a mechanical stirrer, a dual pH/temperature probe, a nitrogen inlet, and a reflux condenser was added sequentially 27.9 g (0.180 mol) of 4-amino-2,5-dimethoxypyrimidine followed by 165.4 g (0.207 mol) of 16.4 wt% ethoxy carbonylisothiocyanate solution in toluene. The reaction mixture was heated to gentle reflux (87 °C) for 7 h at which time liquid chromatographic (LC) analysis indicated
10 ~95% conversion of starting 4-amino-2,5-dimethoxypyrimidine. The reaction mixture was cooled to 27 °C and allowed to stand overnight. The mixture was heated to 40 °C and then 114.2 g (6.34 mol) of deionized water was added to the mixture. After heating to reflux (~68 °C), 14.3 g (0.217 mol) of a 50 wt% aqueous hydroxylamine solution was continuously added over a 2 h 15 min period via a peristaltic pump. During the course of the amine
15 addition, the reaction pH rose from 4.44 to 6.95. After complete addition of hydroxylamine, the pump line was flushed with 4.8 g (0.266 mol) of deionized water, the reaction mixture was heated to 81 °C, and then stirred an additional 3 h during which time the reaction pH naturally raised to 7.40. The reaction mixture was cooled to ambient temperature (26 °C). The reaction mixture was then suction transferred into a temporary holding vessel. The
20 reactor was washed with two 30 g portions of water. These water washes were combined with the reaction mixture. The combined mixture was suctioned filtered through a coarse Buchner funnel (filtration time about 30 seconds), and the filtrate was collected and filtered a second time through the cake. A final displacement cake wash with ~40 g of methanol was performed and the product was dried at 60 °C under vacuum (~<10 mm Hg; 1333Pa) to
25 afford 25.37 g of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine as a light cream colored solid. NMR analysis (using benzyl acetate as an internal standard) indicated an 97.3% purity of 2-amino-5,8-dimethoxy[1,2,4]triazolo[1,5-*c*]pyrimidine active which corresponds to a 70.4% yield.


WHAT IS CLAIMED IS:

1. A process for the preparation of 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]-pyrimidines of the formula (I),



5 in which R is C₁-C₄ alkyl

comprising continuously adding to a [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate of the following formula:



in which R is as previously defined

10

in a polar or nonpolar aprotic solvent an aqueous solution of hydroxylamine at a temperature from about 60 °C to about 110 °C.

2. The process of claim 1, wherein R is CH₃.

3. The process of any of claims 1-2, wherein the aprotic solvent is nonpolar.

15


4. The process of any of claims 1-3, wherein the aprotic solvent is toluene.

5. The process of any of claims 1-4, wherein the temperature of steps (i) and (ii) is at least 60 °C.

6. The process of any of claims 1-5, wherein the temperature of steps (i) and (ii) is at least 80 °C.


20

7. The process of any of claims 1-6, wherein the [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate is formed by contacting a 4-amino-2,5-dialkoxypyrimidine of the formula



in which R is as previously defined


with a C₁-C₄ alkoxy carbonylisothiocyanate of the formula



5 wherein R is as previously defined,

in a polar or nonpolar aprotic solvent at a temperature from about 60 °C to about 110 °C.


8. A process for the preparation of 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]-pyrimidines of the formula (I),



10 in which R represents C₁-C₄ alkyl


comprising:

i) contacting a 4-amino-2,5-dialkoxypyrimidine of the formula




wherein R is as previously defined

15 with a C₁-C₄ alkoxy carbonylisothiocyanate of the formula



in which R is as previously defined

in a polar or nonpolar aprotic solvent at a temperature from about 60 °C to about 110 °C to provide a [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate of the formula



5 in which R is as previously defined; and

ii) continuously adding to the [(2,5-dialkoxypyrimidin-4-yl)amino]carbonothioylcarbamate in the polar or nonpolar aprotic solvent an aqueous solution of hydroxylamine at a temperature from about 60 °C to about 110 °C to provide the 2-amino-5,8-dialkoxy[1,2,4]triazolo[1,5-*c*]pyrimidine.

- 10 9. The process of claim 8, wherein R represents CH₃.
10. The process of any of claims 8-9, wherein the aprotic solvent is nonpolar.
11. The process of any of claims 8-10, wherein the aprotic solvent is toluene.
12. The process of any of claims 8-11, wherein the temperature of steps (i) and (ii) is at least 60 °C.
- 15 13. The process of any of claims 8-12, wherein the temperature of steps (i) and (ii) is at least 80 °C.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/59108

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C07D 487/00 (2013.01)

USPC - 544/280,281

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - C07D 487/00 (2013.01)

USPC - 544/280,281

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
Patents, non-patent literature, search term limitedElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Patbase (pgpb, uspt, usoc, epab, jpab, dwpi, tdbd), Dialog Proquest (npl), Google Patents (pl, npl), Google scholar (pl, npl); Search
Terms: triazolo, pyrimidine, hydroxylamine, benzene, toluene, dioxane, DCM, dichloromethane, hexane, chloroform, aprotic, solvent,
nucleophile, cyclization, nonpolar, non-polar, douglas, dow, agrosience, constant, sl

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2012/0088913 A1 (BOTT, et al.) 12 April 2012 (12.04.2012) entire document, especially para [0005]-[0016], [0033], [0037]-[0038]	1-3, 8-10
Y	US 2003/0134873 A1 (NETTEKOVEN, et al.) 17 July 2003 (17.07.2003) entire document, especially para [0049]	1-3, 8-10
Y	US 2010/0093698 A1 (BAHMANYAR, et al.) 15 April 2010 (15.04.2010) entire document, especially para [0670]	3, 10

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 January 2014 (10.01.2014)

Date of mailing of the international search report

28 JAN 2014

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/59108

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-7 and 11-13
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

摘要

用改进的方法由 4-氨基 2,5-二甲氧基嘧啶制备 5-取代的-8-烷氧基[1,2,4]三唑[1,5-c]嘧啶-2 胺，其中通过连续加入羟胺作为游离碱来控制气态副产物的生成。