

(51) International Patent Classification:
C07D 231/14 (2006.01)(21) International Application Number:
PCT/EP2011/062313(22) International Filing Date:
19 July 2011 (19.07.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10356021.5 20 July 2010 (20.07.2010) EP
61/388,880 1 October 2010 (01.10.2010) US(71) Applicant (for all designated States except US): **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Strasse 50, 40789 Monheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BENTING, Jürgen** [DE/DE]; Amselstrasse 7, 42799 Leichlingen (DE). **DAHMEN, Peter** [DE/DE]; Altebrückerstraße 63, 41470 Neuss (DE). **DESBORDES, Philippe** [FR/FR]; 14 rue du Docteur Mouisset, F-69006 Lyon (FR). **GARY, Stéphanie** [FR/FR]; 9 rue Simon Buisson, F-69410 Champagne Au Mont D'or (FR). **WACHENDORFF-NEUMANN, Ulrike** [DE/DE]; Oberer Markweg 85, 56566 Neuwied (DE).(74) Agent: **GUITTON, Carole**; Bayer Sas, Patents & Licensing Department, 14 impasse Pierre Baizet, CS 99163, F-69263 LYON cedex 09 (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: BENZOCYCLOALKENES AS ANTIFUNGAL AGENTS




(57) Abstract: The present invention relates to fungicidal benzocycloalkene carboxamides or their thiocarboxamide derivatives of formula (I), their process of preparation and intermediate compounds for their preparation, their use as fungicides, particularly in the form of fungicidal compositions and methods for the control of phytopathogenic fungi of plants using these compounds or their compositions.

BENZOCYCLOALKENES AS ANTIFUNGAL AGENTS

DESCRIPTION


The present invention relates to fungicidal benzocycloalkene carboxamides or their thiocarboxamide derivatives, their process of preparation and intermediate compounds for their preparation, their use as fungicides, particularly in the form of fungicidal compositions and methods for the control of phytopathogenic fungi of plants using these compounds or their compositions.

In international patent application DE3410925 [US4628058] certain fungicidal azolylcarboxamides derivatives are generically embraced in a broad disclosure of numerous compounds of the following formula:




wherein R¹ can represent various substituents among which a cycloalkyl group, Y is CH or N, and R² and R³ independently can represent a hydrogen atom or an alkyl group. However, this document does not disclose compounds wherein the urea moiety can be replaced by a carboxamide or thiocarboxamide moiety. Furthermore, there is no specific disclosure in this document of any compound including a cyclopropyl group linked to the nitrogen atom of the azolylcarboxamide residue.

In international patent application WO-2001/064644 certain fungicidal thiocarboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:




wherein Z can represent the 3,4-dichloro-1,2-thiazol-5-yl group and R can represent various substituents among which a indan-1-yl residue or a 1-(2,3-dihydro-1-benzofuran-2-yl)ethyl residue. However, this document does not disclose compounds wherein the nitrogen atom of the thiocarboxamide residue can be substituted by a cycloalkyl group.

In international patent applications WO-2000/015622 and WO-2001/055124 certain fungicidal carboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:




wherein R^1 can represent a hydrogen or a cycloalkyl group, Q can represent a substituted methylene group, k can represent 0 or 1 and Z can represent a 5-7-membered heterocyclic ring comprising 1 to 4 nitrogen atoms or a phenyl group. However, this document does not disclose compounds wherein Z can represent a partially saturated benzo-fused carbocycle or a partially saturated nitrogen-free benzo-fused heterocycle. Moreover, this document does not specifically disclose compounds wherein the nitrogen atom of the thiocarboxamide residue can be substituted by a cycloalkyl group.

In international patent application WO-2009/016218 certain fungicidal (thio)carboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:



wherein A can represent a 5-membered heterocyclic ring, T can represent an oxygen atom or a sulfur atom, Z^1 can represent a C_3 - C_7 -cycloalkyl group, Z^2 can represent various substituents among which a hydrogen atom or a C_1 - C_8 -alkyl group, m can be 1 to 3, and L^1 and L^2 independently can represent various groups or atoms among which an oxygen atom or a methylene group. However, this document does not disclose compounds wherein the nitrogen atom is always linked to the aromatic ring by at least a 2-atoms spacer.

In international patent application WO-2009/016222 certain fungicidal 5-membered fused heteroaryl-methylene carboxamide derivatives are generically embraced in a broad disclosure of numerous compounds of the following formula:




wherein A can represent a 5-membered heterocyclic ring, T can represent an oxygen atom or a sulfur atom, Z^1 can represent a C_3 - C_7 -cycloalkyl group, Z^2 and Z^3 independently can represent various substituents among which a hydrogen atom or a C_1 - C_8 -alkyl group, and the bicyclic ring residue can represent various bicycles among which a benzofurane or a benzothiophene. However, this document does not disclose compounds wherein the the bicyclic ring residue can represent a 2,3-dihydrobenzofurane or a 2,3-dihydrobenzothiophene. Furthermore, there is no disclosure in this document of any compounds wherein the nitrogen atom is always linked to the bicyclic ring by at least a 2-atoms spacer.

It is always of high-interest in the field of agrochemicals to use pesticidal compounds more active than the compounds already known by the man ordinary skilled in the art whereby reduced amounts of compound can be used whilst retaining equivalent efficacy.

Furthermore, the provision of new pesticidal compounds with a higher efficacy strongly reduces the risk of appearance of resistant strains in the fungi to be treated.

We have now found a new family of compounds which show enhanced fungicidal activity over the general known family of such compounds.

Accordingly, the present invention provides a benzocycloalkene carboxamide or its thiocarboxamide derivative of formula (I)



5 wherein

- A represents a carbo-linked, unsaturated or partially saturated, 5-membered heterocyclyl group that can be substituted by up to four groups R that can be the same or different providing that A do not represent the 3,4-dichloro-1,2-thiazol-5-yl group ;

- T represents O or S ;

10 • n represents 0 or 1 ;

- L¹ represents CZ¹Z², NZ⁷, O, S, S(O) or S(O)₂ ;

- L², L³ and L⁴ independently represent a direct bond, CZ¹Z², NZ⁷, O, S, S(O) or S(O)₂

providing that when L² represents NZ⁷, O, S, S(O) or S(O)₂ then L³, L⁴ independently represent a direct bond or CZ¹Z²; or

15 providing that when L³ represents NZ⁷, O, S, S(O) or S(O)₂ then L² represent CZ¹Z² and L⁴ represent direct bond or CZ¹Z²; or

providing that when L⁴ represents NZ⁷, O, S, S(O) or S(O)₂ then L², L³ independently represent CZ¹Z²;

- m represents 0, 1, 2 or 3 ;

- 20 • X represents a halogen atom ; nitro ; cyano ; isonitrile ; hydroxy ; amino ; sulfanyl ; pentafluoro-λ⁶-sulfanyl ; formyl ; formyloxy ; formylamino ; substituted or non-substituted (hydroxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₁-C₈-alkoxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₂-C₈-alkenyloxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₂-C₈-alkynyloxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (benzyloxyimino)-C₁-C₈-alkyl ;
- 25 carboxy ; carbamoyl ; N-hydroxycarbamoyl ; carbamate ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C₂-C₈-alkenyl ; C₂-C₈-halogenoalkenyl having 1 to 5 halogen atoms ; substituted or non-substituted C₂-C₈-alkynyl ; C₂-C₈-halogenoalkynyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxy ; C₁-C₈-halogenoalkoxy having 1 to 5 halogen atoms ; substituted or non-substituted
- 30 C₁-C₈-alkylsulfanyl ; C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylsulfanyl ; C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylsulfonyl ; C₁-C₈-halogenoalkylsulfonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylamino ; substituted or non-substituted di-C₁-C₈-alkylamino ; substituted or non-substituted C₂-C₈-alkenyloxy ; C₂-C₈-halogenoalkenyloxy

having 1 to 5 halogen atoms ; substituted or non-substituted C₃-C₈-alkynyloxy ; C₂-C₈-halogenoalkynyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₃-C₇-cycloalkyl ; C₃-C₇-halogenocycloalkyl having 1 to 5 halogen atoms ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₁-C₈-alkyl ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₂-C₈-alkenyl ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₂-C₈-alkynyl ; substituted or non-substituted tri(C₁-C₈)alkylsilyl ; substituted or non-substituted tri(C₁-C₈)alkylsilyl-C₁-C₈-alkyl ; substituted or non-substituted C₁-C₈-alkylcarbonyl ; C₁-C₈-halogenoalkylcarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbonyloxy ; C₁-C₈-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbonylamino ; C₁-C₈-halogenoalkyl-carbonylamino having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxycarbonyl ; C₁-C₈-halogenoalkoxycarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkyloxycarbonyloxy ; C₁-C₈-halogenoalkyloxycarbonyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbamoyl ; substituted or non-substituted di-C₁-C₈-alkylcarbamoyl ; substituted or non-substituted C₁-C₈-alkylaminocarbonyloxy ; substituted or non-substituted di-C₁-C₈-alkylaminocarbonyloxy ; substituted or non-substituted N-(C₁-C₈-alkyl)hydroxy carbamoyl ; substituted or non-substituted C₁-C₈-alkoxycarbamoyl ; substituted or non-substituted N-(C₁-C₈-alkyl)-C₁-C₈-alkoxycarbamoyl ; aryl that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkyl that can be substituted by up to 6 groups Q which can be the same or different ; C₂-C₈-arylalkenyl that can be substituted by up to 6 groups Q which can be the same or different ; C₂-C₈-arylalkynyl that can be substituted by up to 6 groups Q which can be the same or different ; aryloxy that can be substituted by up to 6 groups Q which can be the same or different ; arylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; arylamino that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkyloxy that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; or C₁-C₈-arylalkylamino that can be substituted by up to 6 groups Q which can be the same or different ; or

- two substituent X together with the consecutive carbon atoms to which they are linked can form a 5- or 6-membered, saturated carbocycle or saturated heterocycle, which can be substituted by up to four groups Q which can be the same or different ;
- Z¹, Z², Z³, Z⁴ and Z⁵ independently represents a hydrogen atom ; halogen atom ; cyano ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxy; substituted or non-substituted C₁-C₈-alkylsulfanyl; or substituted or non-substituted C₁-C₈-alkoxycarbonyl ; or Z⁴ and Z⁵ are a C₂-C₅-alkylene group that can be substituted by up to four C₁-C₈-alkyl groups ;
- Z⁶ represents a non-substituted C₃-C₇-cycloalkyl or a C₃-C₇-cycloalkyl substituted by up to 10 atoms or groups that can be the same or different and that can be selected in the list consisting of halogen atoms, cyano, C₁-C₈-alkyl, C₁-C₈-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkoxy, C₁-C₈-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkoxycarbonyl, C₁-C₈-halogenoalkoxycarbonyl comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkylaminocarbonyl and di-C₁-C₈-alkylaminocarbonyl ;

- Z^7 represents a hydrogen atom ; a substituted or non-substituted C_1 - C_8 -alkyl ; a C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_2 - C_8 -alkenyl ; a C_2 - C_8 -halogenoalkenyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_8 -alkynyl ; a C_3 - C_8 -halogenoalkynyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_7 -cycloalkyl ; a C_3 - C_7 -halogeno-cycloalkyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_7 -cycloalkyl- C_1 - C_8 -alkyl ; or formyl ;
- Q independently represents a halogen atom ; cyano ; nitro ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; C_1 - C_8 -halogenoalkylsulfanyl having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl- C_1 - C_8 -alkyl ; substituted or non-substituted (C_1 - C_8 -alkoxyimino)- C_1 - C_8 -alkyl ; or substituted or non-substituted (benzyloxyimino)- C_1 - C_8 -alkyl ;
- R independently represents hydrogen atom ; halogen atom ; nitro ; cyano ; hydroxy ; amino ; sulfanyl ; pentafluoro- λ^6 -sulfanyl ; substituted or non-substituted (C_1 - C_8 -alkoxyimino)- C_1 - C_8 -alkyl ; substituted or non-substituted (benzyloxyimino)- C_1 - C_8 -alkyl ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C_2 - C_8 -alkenyl ; C_2 - C_8 -halogenoalkenyl having 1 to 5 halogen atoms ; substituted or non-substituted C_2 - C_8 -alkynyl ; C_2 - C_8 -halogenoalkynyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; C_1 - C_8 -halogenoalkylsulfanyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfinyl ; C_1 - C_8 -halogenoalkylsulfinyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfonyl ; C_1 - C_8 -halogenoalkylsulfonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylamino ; substituted or non-substituted di- C_1 - C_8 -alkylamino ; substituted or non-substituted C_2 - C_8 -alkenyloxy ; substituted or non-substituted C_3 - C_8 -alkynyloxy ; substituted or non-substituted C_3 - C_7 -cycloalkyl ; C_3 - C_7 -halogenocycloalkyl having 1 to 5 halogen atoms ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl ; substituted or non-substituted C_1 - C_8 -alkylcarbonyl ; C_1 - C_8 -halogenoalkylcarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkoxycarbonyl ; C_1 - C_8 -halogenoalkoxycarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylcarbamoyl ; substituted or non-substituted di- C_1 - C_8 -alkylcarbamoyl ; phenoxy ; phenylsulfanyl ; phenylamino ; benzyloxy ; benzylsulfanyl ; or benzylamino ;

as well as its salts, N-oxydes, metallic complexes, metalloidic complexes and optically active isomers.

For the compounds according to the invention, the following generic terms are generally used with the following meanings:

- halogen means fluorine, bromine, chlorine or iodine.
- carboxy means $-C(=O)OH$;

carbonyl means $-C(=O)-$;

carbamoyl means $-C(=O)NH_2$;

N-hydroxycarbamoyl means $-C(=O)NHOH$;

S(O) represents a sulfoxide group ;

S(O)₂ represents a sulfone group ;

- an alkyl group, an alkenyl group and an alkynyl group as well as moieties containing these terms, can be linear or branched;
- the aryl moiety contained in an aryl group, an arylalkyl group, an arylalkenyl group and an arylalkynyl group as well as moieties containing these terms, can be a phenyl group that can be substituted by up to 5 groups Q which can be the same or different, a naphthyl group that can be substituted by up to 7 groups Q which can be the same or different or a pyridyl group that can be substituted by up to 4 groups Q which can be the same or different ;
- and, heteroatom means sulfur, nitrogen or oxygen.
- in the case of an amino group or the amino moiety of any other amino-comprising group, substituted by two substituent that can be the same or different, the two substituent together with the nitrogen atom to which they are linked can form a heterocyclyl group, preferably a 5- to 7-membered heterocyclyl group, that can be substituted or that can include other hetero atoms, for example a morpholino group or piperidinyl group.
- unless indicated otherwise, a group or a substituent that is substituted according to the invention can be substituted by one or more of the following groups or atoms: a halogen atom, a nitro group, a hydroxy group, a cyano group, an amino group, a sulfanyl group, a pentafluoro- λ^6 -sulfanyl group, a formyl group, a formyloxy group, a formylamino group, a carbamoyl group, a N-hydroxycarbamoyl group, a carbamate group, a (hydroxyimino)-C₁-C₆-alkyl group, a C₁-C₈-alkyl, a tri(C₁-C₈-alkyl)silyl-C₁-C₈-alkyl, C₁-C₈-cycloalkyl, tri(C₁-C₈-alkyl)silyl-C₁-C₈-cycloalkyl, a C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms, a C₁-C₈-halogenocycloalkyl having 1 to 5 halogen atoms, a C₂-C₈-alkenyl, a C₂-C₈-alkynyl, a C₂-C₈-alkenyloxy, a C₂-C₈-alkynyloxy, a C₁-C₈-alkylamino, a di-C₁-C₈-alkylamino, a C₁-C₈-alkoxy, a C₁-C₈-halogenoalkoxy having 1 to 5 halogen atoms, a C₁-C₈-alkylsulfanyl, a C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms, a C₂-C₈-alkenyloxy, a C₂-C₈-halogenoalkenyloxy having 1 to 5 halogen atoms, a C₃-C₈-alkynyloxy, a C₃-C₈-halogenoalkynyloxy having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbonyl, a C₁-C₈-halogenoalkylcarbonyl having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbamoyl, a di-C₁-C₈-alkylcarbamoyl, a N-C₁-C₈-alkyloxycarbamoyl, a C₁-C₈-alkoxycarbamoyl, a N-C₁-C₈-alkyl-C₁-C₈-alkoxycarbamoyl, a C₁-C₈-alkoxycarbonyl, a C₁-C₈-halogenoalkoxycarbonyl having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbonyloxy, a C₁-C₈-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms, a C₁-C₈-alkylcarbonylamino, a C₁-C₈-halogenoalkylcarbonylamino having 1 to 5 halogen atoms, a C₁-C₈-alkylaminocarbonyloxy, a di-C₁-C₈-alkylaminocarbonyloxy, a C₁-C₈-alkyloxycarbonyloxy, a C₁-C₈-alkylsulfinyl, a C₁-C₈-halogenoalkylsulfinyl having 1 to 5 halogen atoms, a C₁-C₈-alkylsulfonyl, a C₁-C₈-halogenoalkylsulfonyl having 1 to 5 halogen atoms, a C₁-C₈-alkylaminosulfamoyl, a di-C₁-C₈-alkylaminosulfamoyl, a (C₁-C₆-alkoxyimino)-C₁-C₆-alkyl, a (C₁-C₆-alkenyloxyimino)-C₁-C₆-alkyl, a (C₁-C₆-alkynyloxyimino)-C₁-C₆-alkyl, a 2-oxopyrrolidin-1-yl,

(benzyloxyimino)-C₁-C₈-alkyl, C₁-C₈-alkoxyalkyl, C₁-C₈-halogenoalkoxyalkyl having 1 to 5 halogen atoms, benzyloxy, benzylsulfanyl, benzylamino, phenoxy, phenylsulfanyl, or phenylamino.

Any of the compounds of the present invention can exist in one or more optical or chiral isomer forms depending on the number of asymmetric centres in the compound. The invention thus relates equally to all the optical isomers and to their racemic or scalemic mixtures (the term "scalemic" denotes a mixture of enantiomers in different proportions) and to the mixtures of all the possible stereoisomers, in all proportions. The diastereoisomers and/or the optical isomers can be separated according to the methods which are known *per se* by the man ordinary skilled in the art.


Any of the compounds of the present invention can also exist in one or more geometric isomer forms depending on the number of double bonds in the compound. The invention thus relates equally to all geometric isomers and to all possible mixtures, in all proportions. The geometric isomers can be separated according to general methods, which are known *per se* by the man ordinary skilled in the art.

Any of the compounds of the present invention can also exist in one or more geometric isomer forms depending on the relative position (syn/anti or cis/trans) of the substituents of ring B. The invention thus relates equally to all syn/anti (or cis/trans) isomers and to all possible syn/anti (or cis/trans) mixtures, in all proportions. The syn/anti (or cis/trans) isomers can be separated according to general methods, which are known *per se* by the man ordinary skilled in the art.

Any of the compounds of formula (I) wherein X represents a hydroxy, a sulfanyl group or an amino group may be found in its tautomeric form resulting from the shift of the proton of said hydroxy, sulfanyl or amino group. Such tautomeric forms of such compounds are also part of the present invention. More generally speaking, all tautomeric forms of compounds of formula (I) wherein X represents a hydroxy, a sulfanyl group or an amino group, as well as the tautomeric forms of the compounds which can optionally be used as intermediates in the preparation processes and which will be defined in the description of these processes, are also part of the present invention.

Preferred compounds according to the invention are compounds of formula (I) wherein A is selected in the list consisting of:


- a heterocycle of formula (A¹)



wherein :

R¹ to R³ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;


- a heterocycle of formula (A²)



wherein :

R⁴ to R⁶ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A³)




wherein :

R⁷ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R⁸ represents a hydrogen atom or a substituted or non-substituted C₁-C₅-alkyl ;


- a heterocycle of formula (A⁴)



wherein :

R⁹ to R¹¹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; amino ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₁-C₅-alkylsulfanyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A⁵)




wherein :

R¹² and R¹³ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; amino ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R¹⁴ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; amino ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A⁶)




wherein :

R¹⁵ represents a hydrogen atom ; a halogen atom ; a cyano ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R¹⁶ and R¹⁸ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkoxycarbonyl ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R¹⁷ represent a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

- a heterocycle of formula (A⁷)




wherein :

R^{19} represents a hydrogen atom or a C_1 - C_5 -alkyl

R^{20} to R^{22} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

5

- a heterocycle of formula (A^8)




wherein :

10 R^{23} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{24} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different;

15 - a heterocycle of formula (A^9)




wherein :

20 R^{25} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{26} represents a hydrogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A^{10})




25

wherein :

R^{27} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

30 R^{28} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C_1 - C_5 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C_1 - C_5 -alkylamino or substituted or non-substituted di(C_1 - C_5 -alkyl)amino ;

- a heterocycle of formula (A¹¹)




5 wherein :

R²⁹ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

10 R³⁰ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C₁-C₅-alkylamino or substituted or non-substituted di(C₁-C₅-alkyl)amino ;

15 - a heterocycle of formula (A¹²)



wherein :


R³¹ represents a hydrogen atom or a substituted or non-substituted C₁-C₅-alkyl

20 R³² represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R³³ represents a hydrogen atom ; a halogen atom ; a nitro ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

25

- a heterocycle of formula (A¹³)




30 wherein :

R^{34} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; substituted or non-substituted C_3 - C_5 -cycloalkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_5 -alkoxy ; substituted or non-substituted C_2 - C_5 -alkynyloxy or C_1 - C_5 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R^{35} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; a cyano ; substituted or non-substituted C_1 - C_5 -alkoxy ; substituted or non-substituted C_1 - C_5 -alkylsulfanyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C_1 - C_5 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C_1 - C_5 -alkylamino or substituted or non-substituted di(C_1 - C_5 -alkyl)amino;

R^{36} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl ;

-a heterocycle of formula (A¹⁴)




wherein :

R^{37} and R^{38} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_5 -alkoxy or a substituted or non-substituted C_1 - C_5 -alkylsulfanyl ;

R^{39} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl ;


- a heterocycle of formula (A¹⁵)



wherein :

R^{40} and R^{41} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A¹⁶)




wherein :

R^{42} and R^{43} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or amino ;

5


-a heterocycle of formula (A^{17})



wherein :

10 R^{44} and R^{45} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A^{18})




15

wherein :

R^{47} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

20 R^{46} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or substituted or non-substituted C_1 - C_5 -alkylsulfanyl ;

- a heterocycle of formula (A^{19})




25

wherein :

R^{49} and R^{48} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; substituted or non-substituted C_1 - C_5 -alkoxy ; C_1 - C_5 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

30


- a heterocycle of formula (A^{20})



wherein :

- 5 R⁵⁰ and R⁵¹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;


- 10 -a heterocycle of formula (A²¹)



wherein :

- 15 R⁵² represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.


- a heterocycle of formula (A²²)



- 20 wherein :

R⁵³ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.

- a heterocycle of formula (A²³)



- 25


wherein :

R^{54} and R^{56} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{55} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl ;

5

-a heterocycle of formula (A^{24})




wherein :

10 R^{57} and R^{59} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{58} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl ;

15 -a heterocycle of formula (A^{25})




wherein :

20 R^{60} and R^{61} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{62} represents a hydrogen atom or substituted or non-substituted C_1 - C_5 -alkyl ;

-a heterocycle of formula (A^{26})



25

wherein :

R^{65} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; substituted or non-substituted C_3 - C_5 -cycloalkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be

the same or different ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₂-C₅-alkynyloxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; R⁶³ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; a cyano ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₁-C₅-alkylsulfanyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C₁-C₅-alkylamino or di(C₁-C₅-alkyl)amino ; R⁶⁴ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl.

More preferred compounds according to the invention are compounds of formula (I) wherein A is selected in the list consisting of A² ; A⁶ ; A¹⁰ and A¹³ as herein-defined.

Even more preferred compounds according to the invention are compounds of formula (I) wherein A represents A¹³ wherein R³⁴ represents a substituted or non-substituted C₁-C₅-alkyl, C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; or a substituted or non-substituted C₁-C₅-alkoxy ; R³⁵ represents a hydrogen atom or a halogen atom and R³⁶ represents a substituted or non-substituted C₁-C₅-alkyl.

Other preferred compounds according to the invention are compounds of formula (I) wherein T represents O.

Other preferred compounds according to the invention are compounds of formula (I) wherein L¹ represents O or CZ¹Z².

Other preferred compounds according to the invention are compounds of formula (I) wherein L² represents O or CZ¹Z².

Other preferred compounds according to the invention are compounds of formula (I) wherein L³ represents a direct bond, O or CZ¹Z².

Other preferred compounds according to the invention are compounds of formula (I) wherein L⁴ represents a direct bond, O or CZ¹Z². More preferably, L⁴ represents a direct bond.

Other preferred compounds according to the invention are compounds of formula (I) wherein X independently represents a halogen atom ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted tri(C₁-C₈-alkyl)silyl ; substituted or non-substituted C₁-C₈-alkoxy or C₁-C₈-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; or wherein two consecutive substituents X together with the phenyl ring form a substituted or non substituted 1,3-benzodioxolyl ; 1,2,3,4-tetrahydro-quinoxalinylyl ; 3,4-dihydro-2H-1,4-benzoxazinyl ; 1,4-benzodioxanylyl ; indanylyl ; 2,3-dihydrobenzofuranylyl ; or indolinyl.

Other preferred compounds according to the invention are compounds of formula (I) wherein Z^1 , Z^2 , Z^3 , Z^4 and Z^5 independently represents a hydrogen atom, halogen, substituted or non-substituted C_1 - C_8 -alkyl or substituted or non-substituted C_1 - C_8 -alkoxy.

5

Other preferred compounds according to the invention are compounds of formula (I) wherein Z^6 represents a non-substituted C_3 - C_7 cycloalkyl or a C_3 - C_7 cycloalkyl substituted by up to 10 groups or atoms that can be the same or different and that can be selected in the list consisting of halogen atoms, C_1 - C_8 -alkyl, C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, C_1 - C_8 -alkoxy and C_1 - C_8 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different. More preferably Z^6 represents a non-substituted C_3 - C_7 -cycloalkyl ; even more preferably Z^6 represents cyclopropyl.

15

Other preferred compounds according to the invention are compounds of formula (I) wherein Z^7 independently represents a substituted or non-substituted C_1 - C_8 -alkyl.

20

Other preferred compounds according to the invention are compounds of formula (I) wherein R independently represents a hydrogen atom ; halogen atom ; cyano ; substituted or non-substituted C_1 - C_8 -alkylamino ; substituted or non-substituted di- C_1 - C_8 -alkylamino ; substituted or non-substituted tri(C_1 - C_8 -alkyl)silyl ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; amino ; hydroxyl ; nitro ; substituted or non-substituted C_1 - C_8 -alkoxycarbonyl ; substituted or non-substituted C_2 - C_8 -alkynyloxy.

25

The above mentioned preferences with regard to the substituents of the compounds according to the invention can be combined in various manners. These combinations of preferred features thus provide sub-classes of compounds according to the invention. Examples of such sub-classes of preferred compounds according to the invention can be combined:

30

- preferred features of A with preferred features of L^1 to L^4 , Z^1 to Z^7 , X, T and R;

- preferred features of L^1 with preferred features of A, L^2 to L^4 , Z^1 to Z^7 , X, T and R;

- preferred features of L^2 with preferred features of A, L^1 , L^3 to L^4 , Z^1 to Z^7 , X, T and R;

- preferred features of L^3 with preferred features of A, L^1 to L^2 , L^4 , Z^1 to Z^7 , X, T and R;

- preferred features of L^4 with preferred features of A, L^1 to L^3 , Z^1 to Z^7 , X, T and R;

35

- preferred features of Z^1 with preferred features of A, L^1 to L^4 , Z^2 to Z^7 , X, T and R;

- preferred features of Z^2 with preferred features of A, L^1 to L^4 , Z^1 , Z^3 to Z^7 , X, T and R;

- preferred features of Z^3 with preferred features of A, L^1 to L^4 , Z^1 to Z^2 , Z^4 to Z^7 , X, T and R;

- preferred features of Z^4 with preferred features of A, L^1 to L^4 , Z^1 to Z^3 , Z^5 to Z^7 , X, T and R;

- preferred features of Z^5 with preferred features of A, L^1 to L^4 , Z^1 to Z^4 , Z^6 to Z^7 , X, T and R;

40

- preferred features of Z^6 with preferred features of A, L^1 to L^4 , Z^1 to Z^5 to Z^7 , X, T and R;

- preferred features of Z^7 with preferred features of A, L^1 to L^4 , Z^1 to Z^6 , X, T and R;
- preferred features of X with preferred features of A, L^1 to L^4 , Z^1 to Z^6 , T and R;
- preferred features of T with preferred features of A, L^1 to L^4 , Z^1 to Z^6 , X and R;
- preferred features of R with preferred features of A, L^1 to L^4 , Z^1 to Z^6 , X and T;

5


In these combinations of preferred features of the substituents of the compounds according to the invention, the said preferred features can also be selected among the more preferred features of each of A, L^1 to L^4 , Z^1 to Z^7 , X, T and R, so as to form most preferred subclasses of compounds according to the invention.

10


The present invention also relates to a process for the preparation of the compound of formula (I).

Thus, according to a further aspect of the present invention there is provided a process P1 for the preparation of a compound of formula (I) as herein-defined and wherein T represents O and that comprises reacting a benzocycloalkenylamine or benzocycloalkenylalkylamine derivative of formula (II) or one of its salts:

15



wherein X, m, n, L^1 , L^2 , L^3 , L^4 , Z^3 , Z^4 , Z^5 and Z^6 are as herein-defined; with a carboxylic acid derivative of formula (III):



20

wherein A is as herein-defined and Y represents a leaving group selected in the list consisting of a halogen atom, a hydroxyl group, $-OR^a$, $-OC(=O)R^a$, R^a being a substituted or non-substituted C_1 - C_6 -alkyl, a substituted or non-substituted C_1 - C_6 -haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl group, or a group of formula $O-C(=O)A$; in the presence of a catalyst and in the presence of a condensing agent in case Y represents a hydroxyl group, and in the presence of an acid binder in case Y represents a halogen atom.

25

Benzocycloalkenylamine or benzocycloalkenylalkylamine derivatives of formula (II) are known or can be prepared by known processes such as reductive amination of an aldehyde or ketone (Bioorganics and Medicinal Chemistry Letters (2006), 2014 and WO-2005/012291), or reduction of imines (Tetrahedron (2005), 11689), or nucleophilic substitution of a halogen, mesylate or tosylate (Journal of Medicinal Chemistry (2002), 3887).

30

More specifically, N-[(7-methylbicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl]cyclopropanamine derivatives of formula (II) wherein L^1 , L^2 and L^3 represent a direct bond, can be prepared by a palladium mediated cyclisation of an alkyl 2-(2-halogenophenyl)-2-methylpropanoate derivative into an alkyl 7-methylbicyclo[4.2.0]octa-1,3,5-triene-7-carboxylate derivative followed by an amidification by cyclopropylamine and further reduction of the obtained amide to N-[(7-methylbicyclo[4.2.0]octa-1,3,5-trien-7-yl)methyl]-

35

cyclopropanamine derivatives of formula (II) (WO-2010/007253 and Journal of the American Chemical Society (2010), 132, 10706)

Carboxylic acid derivatives of formula (III) can be prepared by known processes.

5

In case Y represents a hydroxy group, the process according to the present invention is conducted in the presence of condensing agent. Suitable condensing agent may be selected in the non limited list consisting of acid halide former, such as phosgene, phosphorous tribromide, phosphorous trichloride, phosphorous pentachloride, phosphorous trichloride oxide or thionyl chloride; anhydride former, such as ethyl chloroformate, methyl chloroformate, isopropyl chloroformate, isobutyl chloroformate or methanesulfonyl chloride; carbodiimides, such as N,N'-dicyclohexylcarbodiimide (DCC) or other customary condensing agents, such as phosphorous pentoxide, polyphosphoric acid, N,N'-carbonyl-diimidazole, 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine/tetrachloromethane, 4-(4,6-dimethoxy[1.3.5]-triazin-2-yl)-4-methylmorpholinium chloride hydrate or bromotripyrrolidino-phosphonium-hexafluorophosphate.

15

The process according to the present invention is conducted in the presence of a catalyst. Suitable catalyst may be selected in the list consisting of 4-dimethyl-aminopyridine, 1-hydroxy-benzotriazole or dimethylformamide.

In case Y represents a halogen atom, the process according to the present invention is conducted in the presence of an acid binder. Suitable acid binders for carrying out process P1 according to the invention are in each case all inorganic and organic bases that are customary for such reactions. Preference is given to using alkaline earth metal, alkali metal hydride, alkali metal hydroxides or alkali metal alkoxides, such as sodium hydroxide, sodium hydride, calcium hydroxide, potassium hydroxide, potassium tert-butoxide or other ammonium hydroxide, alkali metal carbonates, such as cesium carbonate, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate, alkali metal or alkaline earth metal acetates, such as sodium acetate, potassium acetate, calcium acetate and also tertiary amines, such as trimethylamine, triethylamine, diisopropylethylamine, tributylamine, N,N-dimethylaniline, pyridine, N-methylpiperidine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).


25

30

It is also possible to work in the absence of an additional condensing agent or to employ an excess of the amine component, so that it simultaneously acts as acid binder agent.

35

According to a further aspect according to the invention, there is provided a process P2 for the preparation of a compound of formula (I) wherein T represents S, starting from a compound of formula (I) wherein T represents O and illustrated according to the following reaction scheme :



Scheme P2

wherein A, X, m, n, L¹, L², L³, L⁴, Z³, Z⁴, Z⁵ and Z⁶ are as herein-defined, in the optional presence of a catalytic or stoichiometric or more, quantity of a base such as an inorganic and organic base. Preference is given to using alkali metal carbonates, such as sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate; heterocyclic aromatic bases, such as pyridine, picoline, lutidine, collidine; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylaminopyridine or N-methyl-piperidine.

Process P2 according to the invention is performed in the presence of a thionating agent.


Starting amide derivatives of formula (I) can be prepared according to processes P1.

Suitable thionating agents for carrying out process P2 according to the invention can be sulfur (S), sulfhydic acid (H₂S), sodium sulfide (Na₂S), sodium hydrosulfide (NaHS), boron trisulfide (B₂S₃), bis(diethylaluminium) sulfide ((AlEt₂)₂S), ammonium sulfide ((NH₄)₂S), phosphorous pentasulfide (P₂S₅), Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,2,3,4-dithiadiphosphetane 2,4-disulfide) or a polymer-supported thionating reagent such as described in Journal of the Chemical Society, Perkin 1 (2001), 358.

The compound according to the present invention can be prepared according to the general processes of preparation described above. It will nevertheless be understood that, on the basis of his general knowledge and of available publications, the skilled worker will be able to adapt this method according to the specifics of each of the compounds, which it is desired to synthesize.

Still in a further aspect, the present invention relates to compounds of formula (II) useful as intermediate compounds or materials for the process of preparation according to the invention.

The present invention thus provides compounds of formula (II) :



wherein X, m, n, L¹, L², L³, L⁴, Z³, Z⁴ and Z⁵ are as herein-defined, and Z⁶ represents a cyclopropyl group with the exclusion of

- substituted N-cyclopropyl-1,2,3,4-tetrahydronaphthalen-2-amines,
- substituted N-cyclopropylchroman-3-amines,
- N-cyclopropyl-2,3-dihydro-1H-inden-2-amine,

- N-(3,4-dihydro-2H-chromen-3-ylmethyl)cyclopropanamine,
- N-(2,3-dihydro-1-benzofuran-2-ylmethyl)cyclopropanamine.

On the basis of the present description and his general knowledge and of available publications as for example the international patent applications WO-2005/012291 and WO-2006/122955, the skilled person can prepare intermediate compound of formula (II) according to the present invention.

In a further aspect, the present invention also relates to a fungicide composition comprising an effective and non-phytotoxic amount of an active compound of formula (I).

The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention that is sufficient to control or destroy the fungi present or liable to appear on the crops and that does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be controlled, the type of crop, the climatic conditions and the compounds included in the fungicide composition according to the invention. This amount can be determined by systematic field trials that are within the capabilities of a person skilled in the art.

Thus, according to the invention, there is provided a fungicide composition comprising, as an active ingredient, an effective amount of a compound of formula (I) as herein defined and an agriculturally acceptable support, carrier or filler.

According to the invention, the term "support" denotes a natural or synthetic, organic or inorganic compound with that the active compound of formula (I) is combined or associated to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support can be a solid or a liquid. Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports can also be used.

The composition according to the invention can also comprise additional components. In particular, the composition can further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention can be made, for example, of polyacrylic acid salts, lignosulfonic acid salts, phenolsulfonic or naphthalenesulfonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulfosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols and derivatives of the above compounds containing sulfate, sulfonate and phosphate functions. The presence of at least one surfactant is generally essential when the active compound and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content can be comprised from 5% to 40% by weight of the composition.

Optionally, additional components can also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active

compounds can be combined with any solid or liquid additive, that complies with the usual formulation techniques.

In general, the composition according to the invention can contain from 0.05 to 99% by weight of active compound, preferably 10 to 70% by weight.

Compositions according to the invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), ultra low volume (ULV) liquid, ultra low volume (ULV) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder. These compositions include not only compositions that are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions that must be diluted before application to the crop.

The compounds according to the invention can also be mixed with one or more insecticide, fungicide, bactericide, attractant, acaricide or pheromone active substance or other compounds with biological activity. The mixtures thus obtained have normally a broadened spectrum of activity. The mixtures with other fungicide compounds are particularly advantageous.

Examples of suitable fungicide mixing partners can be selected in the following lists:

(1) Inhibitors of the ergosterol biosynthesis, for example (1.1) aldimorph (1704-28-5), (1.2) azaconazole (60207-31-0), (1.3) bitertanol (55179-31-2), (1.4) bromuconazole (116255-48-2), (1.5) cyproconazole (113096-99-4), (1.6) diclobutrazole (75736-33-3), (1.7) difenoconazole (119446-68-3), (1.8) diniconazole (83657-24-3), (1.9) diniconazole-M (83657-18-5), (1.10) dodemorph (1593-77-7), (1.11) dodemorph acetate (31717-87-0), (1.12) epoxiconazole (106325-08-0), (1.13) etaconazole (60207-93-4), (1.14) fenarimol (60168-88-9), (1.15) fenbuconazole (114369-43-6), (1.16) fenhexamid (126833-17-8), (1.17) fenpropidin (67306-00-7), (1.18) fenpropimorph (67306-03-0), (1.19) fluquinconazole (136426-54-5), (1.20) flurprimidol (56425-91-3), (1.21) flusilazole (85509-19-9), (1.22) flutriafol (76674-21-0), (1.23) furconazole (112839-33-5), (1.24) furconazole-cis (112839-32-4), (1.25) hexaconazole (79983-71-4), (1.26) imazalil (60534-80-7), (1.27) imazalil sulfate (58594-72-2), (1.28) imibenconazole (86598-92-7), (1.29) ipconazole (125225-28-7), (1.30) metconazole (125116-23-6), (1.31) myclobutanil (88671-89-0), (1.32) naftifine (65472-88-0), (1.33) nuarimol (63284-71-9), (1.34) oxpoconazole (174212-12-5), (1.35) paclobutrazol (76738-62-0), (1.36) pefurazoate (101903-30-4), (1.37) penconazole (66246-88-6), (1.38) piperalin (3478-94-2), (1.39) prochloraz (67747-09-5), (1.40) propiconazole (60207-90-1), (1.41) prothioconazole (178928-70-6), (1.42) pyributicarb (88678-67-5), (1.43) pyrifenoxy (88283-41-4), (1.44) quinconazole (103970-75-8), (1.45) simeconazole (149508-90-7), (1.46) spiroxamine (118134-30-8), (1.47) tebuconazole (107534-96-3), (1.48) terbinafine (91161-71-6), (1.49) tetraconazole (112281-77-3),

(1.50) triadimefon (43121-43-3), (1.51) triadimenol (89482-17-7), (1.52) tridemorph (81412-43-3), (1.53) triflumizole (68694-11-1), (1.54) triforine (26644-46-2), (1.55) triticonazole (131983-72-7), (1.56) uniconazole (83657-22-1), (1.57) uniconazole-p (83657-17-4), (1.58) viniconazole (77174-66-4), (1.59) voriconazole (137234-62-9), (1.60) 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol (129586-32-9), (1.61) methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate (110323-95-0),
5 (1.62) N'-{5-(difluoromethyl)-2-methyl-4-[3-(trimethylsilyl)propoxy]phenyl}-N-ethyl-N-methylimidiformamide, (1.63) N-ethyl-N-methyl-N'-{2-methyl-5-(trifluoromethyl)-4-[3-(trimethylsilyl)propoxy]phenyl}imidiformamide and (1.64) O-[1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl] 1H-imidazole-1-carbothioate (111226-71-2).

10

(2) inhibitors of the respiratory chain at complex I or II, for example (2.1) bixafen (581809-46-3), (2.2) boscalid (188425-85-6), (2.3) carboxin (5234-68-4), (2.4) diflumetorim (130339-07-0), (2.5) fenfuram (24691-80-3), (2.6) fluopyram (658066-35-4), (2.7) flutolanil (66332-96-5), (2.8) fluxapyroxad (907204-31-3), (2.9) furametpyr (123572-88-3), (2.10) furmecyclox (60568-05-0), (2.11) isopyrazam (mixture of syn-
15 epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR) (881685-58-1), (2.12) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.13) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.14) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.15) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (2.16) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.17) isopyrazam (syn-epimeric enantiomer 1S,4R,9S), (2.18) mepronil (55814-41-0), (2.19) oxycarboxin (5259-88-1),
20 (2.20) penflufen (494793-67-8), (2.21) penthiopyrad (183675-82-3), (2.22) sedaxane (874967-67-6), (2.23) thifluzamide (130000-40-7), (2.24) 1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (2.25) 3-(difluoromethyl)-1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-1H-pyrazole-4-carboxamide, (2.26) 3-(difluoromethyl)-N-[4-fluoro-2-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-1-methyl-1H-pyrazole-4-carboxamide, (2.27) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (1092400-95-7) (WO
25 2008148570), (2.28) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (2.29) N-(4'-chlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.30) N-(2',4'-dichlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.31) 3-(difluoromethyl)-1-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (2.32) N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (2.33) 3-(difluoromethyl)-1-methyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide (known from WO 2004/058723),
30 (2.34) 5-fluoro-1,3-dimethyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.35) 2-chloro-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide (known from WO 2004/058723), (2.36) 3-(difluoromethyl)-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.37) N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.38) 3-(difluoromethyl)-N-(4'-ethynylbiphenyl-2-yl)-1-methyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.39) N-(4'-ethynylbiphenyl-2-yl)-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.40) 2-chloro-N-(4'-ethynylbiphenyl-2-yl)pyridine-3-carboxamide
40 (known from WO 2004/058723), (2.41) 2-chloro-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide (known from WO 2004/058723), (2.42) 4-(difluoromethyl)-2-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1,3-thiazole-5-carboxamide (known from WO 2004/058723), (2.43) 5-fluoro-

N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.44) 2-chloro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide (known from WO 2004/058723), (2.45) 3-(difluoromethyl)-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.46) 5-fluoro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide (known from WO 2004/058723), (2.47) 2-chloro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]pyridine-3-carboxamide (known from WO 2004/058723) and salts thereof

(3) inhibitors of the respiratory chain at complex III, for example (3.1) ametocetradin (865318-97-4), (3.2) amisulbrom (348635-87-0), (3.3) azoxystrobin (131860-33-8), (3.4) cyazofamid (120116-88-3), (3.5) dimoxystrobin (141600-52-4), (3.6) enestroburin (238410-11-2) (WO 2004/058723), (3.7) famoxadone (131807-57-3) (WO 2004/058723), (3.8) fenamidone (161326-34-7) (WO 2004/058723), (3.9) fluoxastrobin (361377-29-9) (WO 2004/058723), (3.10) kresoxim-methyl (143390-89-0) (WO 2004/058723), (3.11) metominostrobin (133408-50-1) (WO 2004/058723), (3.12) orysastrobin (189892-69-1) (WO 2004/058723), (3.13) picoxystrobin (117428-22-5) (WO 2004/058723), (3.14) pyraclostrobin (175013-18-0) (WO 2004/058723), (3.15) pyrametostrobin (915410-70-7) (WO 2004/058723), (3.16) pyraoxystrobin (862588-11-2) (WO 2004/058723), (3.17) pyribencarb (799247-52-2) (WO 2004/058723), (3.18) trifloxystrobin (141517-21-7) (WO 2004/058723), (3.19) (2E)-2-(2-([6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl]oxy)phenyl)-2-(methoxyimino)-N-methylethanamide (WO 2004/058723), (3.20) (2E)-2-(methoxyimino)-N-methyl-2-(2-([[(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene]amino)oxy]methyl]phenyl)ethanamide (WO 2004/058723) and salts thereof., (3.21) (2E)-2-(methoxyimino)-N-methyl-2-{2-[(E)-{1-[3-(trifluoromethyl)phenyl]ethoxy}imino)methyl]phenyl}ethanamide (158169-73-4), (3.22) (2E)-2-{2-([[(1E)-1-(3-[[[(E)-1-fluoro-2-phenylethenyl]oxy]phenyl]ethylidene]amino)oxy]methyl]phenyl)-2-(methoxyimino)-N-methylethanamide (326896-28-0), (3.23) (2E)-2-{2-([[(2E,3E)-4-(2,6-dichlorophenyl)but-3-en-2-ylidene]amino)oxy]methyl]phenyl)-2-(methoxyimino)-N-methylethanamide, (3.24) 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)pyridine-3-carboxamide (119899-14-8), (3.25) 5-methoxy-2-methyl-4-(2-([[(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene]amino)oxy]methyl]phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, (3.26) methyl (2E)-2-{2-([cyclopropyl[(4-methoxyphenyl)imino]methyl]sulfanyl)methyl]phenyl}-3-methoxyprop-2-enoate (149601-03-6), (3.27) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-(formylamino)-2-hydroxybenzamide (226551-21-9), (3.28) 2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide (173662-97-0), (3.29) (2R)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide (394657-24-0) and salts thereof.

(4) Inhibitors of the mitosis and cell division, for example (4.1) benomyl (17804-35-2), (4.2) carbendazim (10605-21-7), (4.3) chlorfenazole (3574-96-7), (4.4) diethofencarb (87130-20-9), (4.5) ethaboxam (162650-77-3), (4.6) fluopicolide (239110-15-7), (4.7) fuberidazole (3878-19-1), (4.8) pencycuron (66063-05-6), (4.9) thiabendazole (148-79-8), (4.10) thiophanate-methyl (23564-05-8), (4.11) thiophanate (23564-06-9), (4.12) zoxamide (156052-68-5), (4.13) 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine (214706-53-3), (4.14) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine (1002756-87-7) and salts thereof.

(5) Compounds capable to have a multisite action, like for example (5.1) bordeaux mixture (8011-63-0), (5.2) captafol (2425-06-1), (5.3) captan (133-06-2) (WO 02/12172), (5.4) chlorothalonil (1897-45-6), (5.5) copper hydroxide (20427-59-2), (5.6) copper naphthenate (1338-02-9), (5.7) copper oxide (1317-39-1), (5.8) copper oxychloride (1332-40-7), (5.9) copper(2+) sulfate (7758-98-7), (5.10) dichlofluanid (1085-98-9), (5.11) dithianon (3347-22-6), (5.12) dodine (2439-10-3), (5.13) dodine free base, (5.14) ferbam (14484-64-1), (5.15) fluorofolpet (719-96-0), (5.16) folpet (133-07-3), (5.17) guazatine (108173-90-6), (5.18) guazatine acetate, (5.19) iminoctadine (13516-27-3), (5.20) iminoctadine albesilate (169202-06-6), (5.21) iminoctadine triacetate (57520-17-9), (5.22) mancopper (53988-93-5), (5.23) mancozeb (8018-01-7), (5.24) maneb (12427-38-2), (5.25) metiram (9006-42-2), (5.26) metiram zinc (9006-42-2), (5.27) oxine-copper (10380-28-6), (5.28) propamidine (104-32-5), (5.29) propineb (12071-83-9), (5.30) sulfur and sulfur preparations including calcium polysulfide (7704-34-9), (5.31) thiram (137-26-8), (5.32) tolylfluanid (731-27-1), (5.33) zineb (12122-67-7), (5.34) ziram (137-30-4) and salts thereof.

(6) Compounds capable to induce a host defence, like for example (6.1) acibenzolar-S-methyl (135158-54-2), (6.2) isotianil (224049-04-1), (6.3) probenazole (27605-76-1), (6.4) tiadinil (223580-51-6) and salts thereof.

(7) Inhibitors of the amino acid and/or protein biosynthesis, for example , (7.1) andoprim (23951-85-1), (7.2) blasticidin-S (2079-00-7), (7.3) cyprodinil (121552-61-2), (7.4) kasugamycin (6980-18-3), (7.5) kasugamycin hydrochloride hydrate (19408-46-9), (7.6) mepanipyrim (110235-47-7), (7.7) pyrimethanil (53112-28-0) and salts thereof.

(8) Inhibitors of the ATP production, for example (8.1) fentin acetate (900-95-8), (8.2) fentin chloride (639-58-7), (8.3) fentin hydroxide (76-87-9) and (8.4) silthiofam (175217-20-6).

(9) Inhibitors of the cell wall synthesis, for example (9.1) benthiavalicarb (177406-68-7), (9.2) dimethomorph (110488-70-5), (9.3) flumorph (211867-47-9), (9.4) iprovalicarb (140923-17-7), (9.5) mandipropamid (374726-62-2), (9.6) polyoxins (11113-80-7), (9.7) polyoxorim (22976-86-9), (9.8) validamycin A (37248-47-8) and (9.9) valifenalate (283159-94-4; 283159-90-0).

(10) Inhibitors of the lipid and membrane synthesis, for example (10.1) biphenyl (92-52-4), (10.2) chloroneb (2675-77-6), (10.3) dicloran (99-30-9), (10.4) edifenphos (17109-49-8), (10.5) etridiazole (2593-15-9), (10.6) iodocarb (55406-53-6), (10.7) iprobenfos (26087-47-8), (10.8) isoprothiolane (50512-35-1), (10.9) propamocarb (25606-41-1), (10.10) propamocarb hydrochloride (25606-41-1), (10.11) prothiocarb (19622-08-3), (10.12) pyrazophos (13457-18-6), (10.13) quintozone (82-68-8), (10.14) tecnazene (117-18-0) and (10.15) tolclofos-methyl (57018-04-9).

(11) Inhibitors of the melanine biosynthesis, for example (11.1) carpropamid (104030-54-8), (11.2) diclocymet (139920-32-4), (11.3) fenoxanil (115852-48-7), (11.4) phthalide (27355-22-2), (11.5) pyroquilon (57369-32-1) and (11.6) tricyclazole (41814-78-2).

(12) Inhibitors of the nucleic acid synthesis, for example (12.1) benalaxyl (71626-11-4), (12.2) benalaxyl-M (kiralaxyl) (98243-83-5), (12.3) bupirimate (41483-43-6), (12.4) clozylacon (67932-85-8), (12.5) dimethirimol (5221-53-4), (12.6) ethirimol (23947-60-6), (12.7) furalaxyl (57646-30-7), (12.8) hymexazol (10004-44-1), (12.9) metalaxyl (57837-19-1), (12.10) metalaxyl-M (mefenoxam) (70630-17-0), (12.11) ofurace (58810-48-3), (12.12) oxadixyl (77732-09-3) and (12.13) oxolinic acid (14698-29-4).

(13) Inhibitors of the signal transduction, for example (13.1) chlozolate (84332-86-5), (13.2) fenpiclonil (74738-17-3), (13.3) fludioxonil (131341-86-1), (13.4) iprodione (36734-19-7), (13.5) procymidone (32809-16-8), (13.6) quinoxifen (124495-18-7) and (13.7) vinclozolin (50471-44-8).

(14) Compounds capable to act as an uncoupler, like for example (14.1) binapacryl (485-31-4), (14.2) dinocap (131-72-6), (14.3) ferimzone (89269-64-7), (14.4) fluazinam (79622-59-6) and (14.5) meptyldinocap (131-72-6).

(15) Further compounds, like for example (15.1) benthiazole (21564-17-0), (15.2) bethoxazin (163269-30-5), (15.3) capsimycin (70694-08-5), (15.4) carvone (99-49-0), (15.5) chinomethionat (2439-01-2), (15.6) chlazafenone (688046-61-9), (15.7) cufraneb (11096-18-7), (15.8) cyflufenamid (180409-60-3), (15.9) cymoxanil (57966-95-7), (15.10) cyprosulfamide (221667-31-8), (15.11) dazomet (533-74-4), (15.12) debacarb (62732-91-6), (15.13) dichlorophen (97-23-4), (15.14) diclomezine (62865-36-5), (15.15) difenzoquat (49866-87-7), (15.16) difenzoquat methylsulfate (43222-48-6), (15.17) diphenylamine (122-39-4), (15.18) ecomate, (15.19) fenpyrazamine (473798-59-3), (15.20) flumetover (154025-04-4), (15.21) fluoroimide (41205-21-4), (15.22) flusulfamide (106917-52-6), (15.23) flutianil (304900-25-2), (15.24) fosetyl-aluminium (39148-24-8), (15.25) fosetyl-calcium, (15.26) fosetyl-sodium (39148-16-8), (15.27) hexachlorobenzene (118-74-1), (15.28) irumamycin (81604-73-1), (15.29) methasulfocarb (66952-49-6), (15.30) methyl isothiocyanate (556-61-6), (15.31) metrafenone (220899-03-6), (15.32) mildiomyacin (67527-71-3), (15.33) natamycin (7681-93-8), (15.34) nickel dimethyldithiocarbamate (15521-65-0), (15.35) nitrothal-isopropyl (10552-74-6), (15.36) octhilinone (26530-20-1), (15.37) oxamocarb (917242-12-7), (15.38) oxyfenthiin (34407-87-9), (15.39) pentachlorophenol and salts (87-86-5), (15.40) phenothrin, (15.41) phosphorous acid and its salts (13598-36-2), (15.42) propamocarb-fosetilate, (15.43) propanosine-sodium (88498-02-6), (15.44) proquinazid (189278-12-4), (15.45) pyrrolnitrine (1018-71-9) (EP-A 1 559 320), (15.46) tebufloquin (376645-78-2), (15.47) tecloftalam (76280-91-6), (15.48) tolnifanide (304911-98-6), (15.49) triazoxide (72459-58-6), (15.50) trichlamide (70193-21-4), (15.51) zarilamid (84527-51-5), (15.52) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-79-6) (WO 2008013622), (15.53) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003319-80-9) (WO 2008013622), (15.54) 1-(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone (1003318-67-9) (WO 2008013622), (15.55) 1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate (111227-17-9), (15.56) 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine (13108-52-6), (15.57) 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)-one (221451-58-7), (15.58) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5R)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-

53-7) (WO 2008013622), (15.59) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5S)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone (1003316-54-8) (WO 2008013622), (15.60) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-{4-[4-(5-phenyl-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}ethanone (1003316-51-5) (WO 2008013622), (15.61) 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, (15.62) 2-chloro-5-[2-chloro-1-(2,6-difluoro-4-methoxyphenyl)-4-methyl-1H-imidazol-5-yl]pyridine, (15.63) 2-phenylphenol and salts (90-43-7), (15.64) 3,4,5-trichloropyridine-2,6-dicarbonitrile (17824-85-0), (15.65) 3-[5-(4-chlorophenyl)-2,3-dimethyl-1,2-oxazolidin-3-yl]pyridine, (15.66) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (15.67) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (15.68) 5-amino-1,3,4-thiadiazole-2-thiol, (15.69) 5-chloro-N'-phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide (134-31-6), (15.70) 5-methyl-6-octyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, (15.71) ethyl (2Z)-3-amino-2-cyano-3-phenylprop-2-enoate, (15.72) N-(4-chlorobenzyl)-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.73) N-[(4-chlorophenyl)(cyano)methyl]-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.74) N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloropyridine-3-carboxamide, (15.75) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloropyridine-3-carboxamide, (15.76) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodopyridine-3-carboxamide, (15.77) N-{(E)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (15.78) N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide (221201-92-9), (15.79) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-1,3-thiazole-4-carboxamide (922514-49-6) (WO 2007014290), (15.80) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-07-6) (WO 2007014290), (15.81) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1S)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide (922514-48-5) (WO 2007014290), (15.82) pentyl {6-[[[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylidene]amino]oxy)methyl}pyridin-2-yl}carbamate, (15.83) phenazine-1-carboxylic acid, (15.84) quinolin-8-ol (134-31-6) and (15.85) quinolin-8-ol sulfate (2:1) (134-31-6), (15.86) (5-bromo-2-methoxy-4-methylpyridin-3-yl)(2,3,4-trimethoxy-6-methylphenyl)methanone (known from EP-A 1 559 320) and (9.10) N-[2-(4-{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl)ethyl]-N2-(methylsulfonyl)valinamide (220706-93-4).

The composition according to the invention comprising a mixture of a compound of formula (I) with a bactericide compound can also be particularly advantageous. Examples of suitable bactericide mixing partners can be selected in the following list: bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin, othillinone, furancarboxylic acid, oxytetracycline, probenazole, streptomycin, tecloftalam, copper sulfate and other copper preparations.

The compounds of formula (I) and the fungicide composition according to the invention can be used to curatively or preventively control the phytopathogenic fungi of plants or crops.

Thus, according to a further aspect of the invention, there is provided a method for curatively or preventively controlling the phytopathogenic fungi of plants or crops characterised in that a compound of formula (I) or a fungicide composition according to the invention is applied to the seed, the plant or to the fruit of the plant or to the soil wherein the plant is growing or wherein it is desired to grow.

The method of treatment according to the invention can also be useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruit of the concerned plant.

According to the invention all plants and plant parts can be treated. By plants is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. By plant parts is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

Among the plants that can be protected by the method according to the invention, mention may be made of major field crops like corn, soybean, cotton, *Brassica* oilseeds such as *Brassica napus* (e.g. canola), *Brassica rapa*, *B. juncea* (e.g. mustard) and *Brassica carinata*, rice, wheat, sugarbeet, sugarcane, oats, rye, barley, millet, triticale, flax, vine and various fruits and vegetables of various botanical taxa such as *Rosaceae* sp. (for instance pip fruit such as apples and pears, but also stone fruit such as apricots, cherries, almonds and peaches, berry fruits such as strawberries), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantings), *Rubiaceae* sp. (for instance coffee), *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruit); *Solanaceae* sp. (for instance tomatoes, potatoes, peppers, eggplant), *Liliaceae* sp., *Compositae* sp. (for instance lettuce, artichoke and chicory - including root chicory, endive or common chicory), *Umbelliferae* sp. (for instance carrot, parsley, celery and celeriac), *Cucurbitaceae* sp. (for instance cucumber - including pickling cucumber, squash, watermelon, gourds and melons), *Alliaceae* sp. (for instance onions and leek), *Cruciferae* sp. (for instance white cabbage, red cabbage, broccoli, cauliflower, brussel sprouts, pak choi, kohlrabi, radish, horseradish, cress, Chinese cabbage), *Leguminosae* sp. (for instance peanuts, peas and beans - such as climbing beans and broad beans), *Chenopodiaceae* sp. (for instance mangold, spinach beet, spinach, beetroots), *Malvaceae* (for instance okra), *Asparagaceae* (for instance asparagus); horticultural and forest crops; ornamental plants; as well as genetically modified homologues of these crops.

The method of treatment according to the invention can be used in the treatment of genetically modified organisms (GMOs), e.g. plants or seeds. Genetically modified plants (or transgenic plants) are plants of which a heterologous gene has been stably integrated into genome. The expression "heterologous gene" essentially means a gene which is provided or assembled outside the plant and when introduced in the nuclear, chloroplastic or mitochondrial genome gives the transformed plant new or improved agronomic or other

properties by expressing a protein or polypeptide of interest or by downregulating or silencing other gene(s) which are present in the plant (using for example, antisense technology, cosuppression technology or RNA interference – RNAi - technology). A heterologous gene that is located in the genome is also called a transgene. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive (“synergistic”) effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the active compounds and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, bigger fruits, larger plant height, greener leaf color, earlier flowering, higher quality and/or a higher nutritional value of the harvested products, higher sugar concentration within the fruits, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

At certain application rates, the active compound combinations according to the invention may also have a strengthening effect in plants. Accordingly, they are also suitable for mobilizing the defense system of the plant against attack by unwanted microorganisms. This may, if appropriate, be one of the reasons of the enhanced activity of the combinations according to the invention, for example against fungi. Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances or combinations of substances which are capable of stimulating the defense system of plants in such a way that, when subsequently inoculated with unwanted microorganisms, the treated plants display a substantial degree of resistance to these microorganisms. In the present case, unwanted microorganisms are to be understood as meaning phytopathogenic fungi, bacteria and viruses. Thus, the substances according to the invention can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period of time within which protection is effected generally extends from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active compounds.

Plants and plant cultivars which are preferably to be treated according to the invention include all plants which have genetic material which impart particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

Plants and plant cultivars which are also preferably to be treated according to the invention are resistant against one or more biotic stresses, i.e. said plants show a better defense against animal and microbial pests, such as against nematodes, insects, mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

Examples of nematode resistant plants are described in e.g. US Patent Application Nos 11/765,491, 11/765,494, 10/926,819, 10/782,020, 12/032,479, 10/783,417, 10/782,096, 11/657,964, 12/192,904, 11/396,808, 12/166,253, 12/166,239, 12/166,124, 12/166,209, 11/762,886, 12/364,335, 11/763,947, 12/252,453, 12/209,354, 12/491,396 or 12/497,221.

Plants and plant cultivars which may also be treated according to the invention are those plants which are resistant to one or more abiotic stresses. Abiotic stress conditions may include, for example, drought, cold temperature exposure, heat exposure, osmotic stress, flooding, increased soil salinity, increased mineral exposure, ozone exposure, high light exposure, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients, shade avoidance.

Plants and plant cultivars which may also be treated according to the invention, are those plants characterized by enhanced yield characteristics. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can furthermore be affected by improved plant architecture (under stress and non-stress conditions), including but not limited to, early flowering, flowering control for hybrid seed production, seedling vigor, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content, protein content, oil content and composition, nutritional value, reduction in anti-nutritional compounds, improved processability and better storage stability.

Examples of plants with the above-mentioned traits are non-exhaustively listed in Table A.

Plants that may be treated according to the invention are hybrid plants that already express the characteristic of heterosis or hybrid vigor which results in generally higher yield, vigor, health and resistance towards biotic and abiotic stresses). Such plants are typically made by crossing an inbred male-sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). Hybrid seed is typically harvested from the male sterile plants and sold to growers. Male sterile plants can sometimes (e.g. in corn) be produced by detasseling, i.e. the mechanical removal of the male reproductive organs (or males flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants it is typically useful to ensure that male fertility in the hybrid plants is fully restored. This can be accomplished by ensuring that the male parents have appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic determinants responsible for male-sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described in Brassica species (WO 92/05251, WO 95/09910, WO 98/27806, WO 05/002324, WO 06/021972 and US 6,229,072). However, genetic determinants for male sterility can also be located in the nuclear genome. Male sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 89/10396 in which, for example, a ribonuclease such as barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar (e.g. WO 91/02069).

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated according to the invention are herbicide-tolerant plants, i.e. plants made tolerant to one or

more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

Herbicide-resistant plants are for example glyphosate-tolerant plants, i.e. plants made tolerant to the herbicide glyphosate or salts thereof. Plants can be made tolerant to glyphosate through different means.

5 For example, glyphosate-tolerant plants can be obtained by transforming the plant with a gene encoding the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are the AroA gene (mutant CT7) of the bacterium *Salmonella typhimurium* (Comai et al., 1983, Science 221, 370-371), the CP4 gene of the bacterium *Agrobacterium sp.* (Barry et al., 1992, Curr. Topics Plant Physiol. 7, 139-145), the genes encoding a Petunia EPSPS (Shah et al., 1986, Science 233, 478-481), a
10 Tomato EPSPS (Gasser et al., 1988, J. Biol. Chem. 263, 4280-4289), or an Eleusine EPSPS (WO 01/66704). It can also be a mutated EPSPS as described in for example EP 0837944, WO 00/66746, WO 00/66747 or WO02/26995. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate oxido-reductase enzyme as described in U.S. Patent Nos. 5,776,760 and 5,463,175. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a
15 glyphosate acetyl transferase enzyme as described in for example WO 02/36782, WO 03/092360, WO 05/012515 and WO 07/024782. Glyphosate-tolerant plants can also be obtained by selecting plants containing naturally-occurring mutations of the above-mentioned genes, as described in for example WO 01/024615 or WO 03/013226. Plants expressing EPSPS genes that confer glyphosate tolerance are described in e.g. US Patent Application Nos 11/517,991, 10/739,610, 12/139,408, 12/352,532,
20 11/312,866, 11/315,678, 12/421,292, 11/400,598, 11/651,752, 11/681,285, 11/605,824, 12/468,205, 11/760,570, 11/762,526, 11/769,327, 11/769,255, 11/943801 or 12/362,774. Plants comprising other genes that confer glyphosate tolerance, such as decarboxylase genes, are described in e.g. US patent applications 11/588,811, 11/185,342, 12/364,724, 11/185,560 or 12/423,926.

25 Other herbicide resistant plants are for example plants that are made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition, e.g. described in US Patent Application No 11/760,602. One such efficient detoxifying enzyme is an enzyme encoding a phosphinothricin acetyltransferase (such as the bar or pat
30 protein from *Streptomyces* species). Plants expressing an exogenous phosphinothricin acetyltransferase are for example described in U.S. Patent Nos. 5,561,236; 5,648,477; 5,646,024; 5,273,894; 5,637,489; 5,276,268; 5,739,082; 5,908,810 and 7,112,665.

Further herbicide-tolerant plants are also plants that are made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvatedioxygenase (HPPD). Hydroxyphenylpyruvatedioxygenases are enzymes
35 that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is transformed into homogentisate. Plants tolerant to HPPD-inhibitors can be transformed with a gene encoding a naturally-occurring resistant HPPD enzyme, or a gene encoding a mutated or chimeric HPPD enzyme as described in WO 96/38567, WO 99/24585, WO 99/24586, WO 2009/144079, WO 2002/046387, or US 6,768,044..

Tolerance to HPPD-inhibitors can also be obtained by transforming plants with genes encoding certain
40 enzymes enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD-inhibitor. Such plants and genes are described in WO 99/34008 and WO 02/36787. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an

enzyme having prephenate deshydrogenase (PDH) activity in addition to a gene encoding an HPPD-tolerant enzyme, as described in WO 2004/024928. Further, plants can be made more tolerant to HPPD-inhibitor herbicides by adding into their genome a gene encoding an enzyme capable of metabolizing or degrading HPPD inhibitors, such as the CYP450 enzymes shown in WO 2007/103567 and WO

2008/150473.

Still further herbicide resistant plants are plants that are made tolerant to acetolactate synthase (ALS) inhibitors. Known ALS-inhibitors include, for example, sulfonylurea, imidazolinone, triazolopyrimidines, pyrimidinyoxy(thio)benzoates, and/or sulfonylaminocarbonyltriazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxyacid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides, as described for example in Tranel and Wright (2002, Weed Science 50:700-712), but also, in U.S. Patent No. 5,605,011, 5,378,824, 5,141,870, and 5,013,659. The production of sulfonylurea-tolerant plants and imidazolinone-tolerant plants is described in U.S. Patent Nos. 5,605,011; 5,013,659; 5,141,870; 5,767,361; 5,731,180; 5,304,732; 4,761,373; 5,331,107; 5,928,937; and 5,378,824; and international publication WO 96/33270. Other imidazolinone-tolerant plants are also described in for example WO 2004/040012, WO 2004/106529, WO 2005/020673, WO 2005/093093, WO 2006/007373, WO 2006/015376, WO 2006/024351, and WO 2006/060634. Further sulfonylurea- and imidazolinone-tolerant plants are also described in for example WO 07/024782 and US Patent Application No 61/288958.

Other plants tolerant to imidazolinone and/or sulfonylurea can be obtained by induced mutagenesis, selection in cell cultures in the presence of the herbicide or mutation breeding as described for example for soybeans in U.S. Patent 5,084,082, for rice in WO 97/41218, for sugar beet in U.S. Patent 5,773,702 and WO 99/057965, for lettuce in U.S. Patent 5,198,599, or for sunflower in WO 01/065922.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are insect-resistant transgenic plants, i.e. plants made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such insect resistance.

An "insect-resistant transgenic plant", as used herein, includes any plant containing at least one transgene comprising a coding sequence encoding:

1) an insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins listed by Crickmore et al. (1998, Microbiology and Molecular Biology Reviews, 62: 807-813), updated by Crickmore et al. (2005) at the *Bacillus thuringiensis* toxin nomenclature, online at:

http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/, or insecticidal portions thereof, e.g., proteins of the Cry protein classes Cry1Ab, Cry1Ac, Cry1B, Cry1C, Cry1D, Cry1F, Cry2Ab, Cry3Aa, or Cry3Bb or insecticidal portions thereof (e.g. EP 1999141 and WO 2007/107302), or such proteins encoded by synthetic genes as e.g. described in and US Patent Application No 12/249,016 ; or

2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein from *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cry34 and Cry35 crystal proteins (Moellenbeck et al. 2001, Nat. Biotechnol. 19: 668-72; Schnepf et al. 2006, Applied Environm. Microbiol. 71, 1765-1774) or

the binary toxin made up of the Cry1A or Cry1F proteins and the Cry2Aa or Cry2Ab or Cry2Ae proteins (US Patent Appl. No. 12/214,022 and EP 08010791.5); or

3) a hybrid insecticidal protein comprising parts of different insecticidal crystal proteins from *Bacillus thuringiensis*, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2)

above, e.g., the Cry1A.105 protein produced by corn event MON89034 (WO 2007/027777); or

4) a protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in corn events MON863 or MON88017, or the Cry3A protein in corn event MIR604; or

5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal (VIP) proteins listed at:

http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html, e.g., proteins from the VIP3Aa protein class; or

6) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B. cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins (WO 94/21795); or

7) a hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the proteins in 2) above; or

8) a protein of any one of 5) to 7) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding an insecticidal protein), such as the VIP3Aa protein in cotton event COT102; or

9) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a crystal protein from *Bacillus thuringiensis*, such as the binary toxin made up of VIP3 and Cry1A or Cry1F (US Patent Appl. No. 61/126083 and 61/195019), or the binary toxin made up of the VIP3 protein and the Cry2Aa or Cry2Ab or Cry2Ae proteins (US Patent Appl. No. 12/214,022 and EP 08010791.5).

10) a protein of 9) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding an insecticidal protein)

Of course, an insect-resistant transgenic plant, as used herein, also includes any plant comprising a combination of genes encoding the proteins of any one of the above classes 1 to 10. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the above classes 1 to 10, to expand the range of target insect species affected when using different proteins directed at different target insect species, or to delay insect resistance development to the plants by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

An "insect-resistant transgenic plant", as used herein, further includes any plant containing at least one transgene comprising a sequence producing upon expression a double-stranded RNA which upon ingestion by a plant insect pest inhibits the growth of this insect pest, as described e.g. in WO 2007/080126, WO 2006/129204, WO 2007/074405, WO 2007/080127 and WO 2007/035650.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are tolerant to abiotic stresses. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress tolerance plants include:

1) plants which contain a transgene capable of reducing the expression and/or the activity of poly(ADP-ribose) polymerase (PARP) gene in the plant cells or plants as described in WO 00/04173, WO/2006/045633, EP 04077984.5, or EP 06009836.5.

2) plants which contain a stress tolerance enhancing transgene capable of reducing the expression and/or the activity of the PARG encoding genes of the plants or plants cells, as described e.g. in WO 2004/090140.

3) plants which contain a stress tolerance enhancing transgene coding for a plant-functional enzyme of the nicotinate adenine dinucleotide salvage synthesis pathway including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenylyl transferase, nicotinamide adenine dinucleotide synthetase or nicotine amide phosphorybosyltransferase as described e.g. in EP 04077624.7, WO 2006/133827, PCT/EP07/002433, EP 1999263, or WO 2007/107326.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention show altered quantity, quality and/or storage-stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as :

1) transgenic plants which synthesize a modified starch, which in its physical-chemical characteristics, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the side chain distribution, the viscosity behaviour, the gelling strength, the starch grain size and/or the starch grain morphology, is changed in comparison with the synthesised starch in wild type plant cells or plants, so that this is better suited for special applications. Said transgenic plants synthesizing a modified starch are disclosed, for example, in EP 0571427, WO 95/04826, EP 0719338, WO 96/15248, WO 96/19581, WO 96/27674, WO 97/11188, WO 97/26362, WO 97/32985, WO 97/42328, WO 97/44472, WO 97/45545, WO 98/27212, WO 98/40503, WO99/58688, WO 99/58690, WO 99/58654, WO 00/08184, WO 00/08185, WO 00/08175, WO 00/28052, WO 00/77229, WO 01/12782, WO 01/12826, WO 02/101059, WO 03/071860, WO 2004/056999, WO 2005/030942, WO 2005/030941, WO 2005/095632, WO 2005/095617, WO 2005/095619, WO 2005/095618, WO 2005/123927, WO 2006/018319, WO 2006/103107, WO 2006/108702, WO 2007/009823, WO 00/22140, WO 2006/063862, WO 2006/072603, WO 02/034923, EP 06090134.5, EP 06090228.5, EP 06090227.7, EP 07090007.1, EP 07090009.7, WO 01/14569, WO 02/79410, WO 03/33540, WO 2004/078983, WO 01/19975, WO 95/26407, WO 96/34968, WO 98/20145,

WO 99/12950, WO 99/66050, WO 99/53072, US 6,734,341, WO 00/11192, WO 98/22604, WO 98/32326, WO 01/98509, WO 01/98509, WO 2005/002359, US 5,824,790, US 6,013,861, WO 94/04693, WO 94/09144, WO 94/11520, WO 95/35026, WO 97/20936

2) transgenic plants which synthesize non starch carbohydrate polymers or which synthesize non starch carbohydrate polymers with altered properties in comparison to wild type plants without genetic modification. Examples are plants producing polyfructose, especially of the inulin and levan-type, as disclosed in EP 0663956, WO 96/01904, WO 96/21023, WO 98/39460, and WO 99/24593, plants producing alpha-1,4-glucans as disclosed in WO 95/31553, US 2002031826, US 6,284,479, US 5,712,107, WO 97/47806, WO 97/47807, WO 97/47808 and WO 00/14249, plants producing alpha-1,6 branched alpha-1,4-glucans, as disclosed in WO 00/73422, plants producing alternan, as disclosed in e.g. WO 00/47727, WO 00/73422, EP 06077301.7, US 5,908,975 and EP 0728213,

3) transgenic plants which produce hyaluronan, as for example disclosed in WO 2006/032538, WO 2007/039314, WO 2007/039315, WO 2007/039316, JP 2006304779, and WO 2005/012529.

4) transgenic plants or hybrid plants, such as onions with characteristics such as 'high soluble solids content', 'low pungency' (LP) and/or 'long storage' (LS), as described in US Patent Appl. No. 12/020,360 and 61/054,026.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as cotton plants, with altered fiber characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered fiber characteristics and include:

- a) Plants, such as cotton plants, containing an altered form of cellulose synthase genes as described in WO 98/00549
- b) Plants, such as cotton plants, containing an altered form of rsw2 or rsw3 homologous nucleic acids as described in WO 2004/053219
- c) Plants, such as cotton plants, with increased expression of sucrose phosphate synthase as described in WO 01/17333
- d) Plants, such as cotton plants, with increased expression of sucrose synthase as described in WO 02/45485
- e) Plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fiber cell is altered, e.g. through downregulation of fiber-selective β -1,3-glucanase as described in WO 2005/017157, or as described in EP 08075514.3 or US Patent Appl. No. 61/128,938
- f) Plants, such as cotton plants, having fibers with altered reactivity, e.g. through the expression of N-acetylglucosaminetransferase gene including nodC and chitin synthase genes as described in WO 2006/136351

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be obtained by genetic

transformation, or by selection of plants contain a mutation imparting such altered oil profile characteristics and include:

- a) Plants, such as oilseed rape plants, producing oil having a high oleic acid content as described e.g. in US 5,969,169, US 5,840,946 or US 6,323,392 or US 6,063,947
- b) Plants such as oilseed rape plants, producing oil having a low linolenic acid content as described in US 6,270,828, US 6,169,190, or US 5,965,755
- c) Plant such as oilseed rape plants, producing oil having a low level of saturated fatty acids as described e.g. in US Patent No. 5,434,283 or US Patent Application No 12/668303

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered seed shattering characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered seed shattering characteristics and include plants such as oilseed rape plants with delayed or reduced seed shattering as described in US Patent Appl. No. 61/135,230 WO09/068313 and WO10/006732.

Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or combination of transformation events, that are the subject of petitions for non-regulated status, in the United States of America, to the Animal and Plant Health Inspection Service (APHIS) of the United States Department of Agriculture (USDA) whether such petitions are granted or are still pending. At any time this information is readily available from APHIS (4700 River Road Riverdale, MD 20737, USA), for instance on its internet site (URL http://www.aphis.usda.gov/brs/not_reg.html). On the filing date of this application the petitions for nonregulated status that were pending with APHIS or granted by APHIS were those listed in table B which contains the following information:

- Petition : the identification number of the petition. Technical descriptions of the transformation events can be found in the individual petition documents which are obtainable from APHIS, for example on the APHIS website, by reference to this petition number. These descriptions are herein incorporated by reference.
- Extension of Petition : reference to a previous petition for which an extension is requested.
- Institution : the name of the entity submitting the petition.
- Regulated article : the plant species concerned.
- Transgenic phenotype : the trait conferred to the plants by the transformation event.
- Transformation event or line : the name of the event or events (sometimes also designated as lines or lines) for which nonregulated status is requested.
- APHIS documents : various documents published by APHIS in relation to the Petition and which can be requested with APHIS.

Additional particularly useful plants containing single transformation events or combinations of transformation events are listed for example in the databases from various national or regional regulatory agencies (see for example http://gmoinfo.jrc.it/gmp_browse.aspx and <http://www.agbios.com/dbase.php>).

Further particularly transgenic plants include plants containing a transgene in an agronomically neutral or beneficial position as described in any of the patent publications listed in Table C.

Table A

Trait	Reference	
Water use efficiency	WO 2000/073475	
Nitrogen use efficiency	WO 1995/009911	WO 2007/076115
	WO 1997/030163	WO 2005/103270
	WO 2007/092704	WO 2002/002776
Improved photosynthesis	WO 2008/056915	WO 2004/101751
Nematode resistance	WO 1995/020669	WO 2003/033651
	WO 2001/051627	WO 1999/060141
	WO 2008/139334	WO 1998/012335
	WO 2008/095972	WO 1996/030517
	WO 2006/085966	WO 1993/018170
Reduced pod dehiscence	WO 2006/009649	WO 1997/013865
	WO 2004/113542	WO 1996/030529
	WO 1999/015680	WO 1994/023043
	WO 1999/000502	
Aphid resistance	WO 2006/125065	WO 2008/067043
	WO 1997/046080	WO 2004/072109
Sclerotinia resistance	WO 2006/135717	WO 2005/000007
	WO 2006/055851	WO 2002/099385
	WO 2005/090578	WO 2002/061043
Botrytis resistance	WO 2006/046861	WO 2002/085105
Bremia resistance	US 20070022496	WO 2004/049786
	WO 2000/063432	
Erwinia resistance	WO 2004/049786	
Closterovirus resistance	WO 2007/073167	WO 2002/022836
	WO 2007/053015	
Stress tolerance (including drought tolerance)	WO 2010/019838	WO2008/002480
	WO 2009/049110	WO2005/033318
Tobamovirus resistance	WO 2006/038794	

Table B

**Petitions of Nonregulated Status Granted or Pending by APHIS
as of March 31, 2010**

NOTE: To obtain the most up-to-date list of Crops No Longer Regulated, please look at the Current Status of Petitions. This list is automatically updated and reflects all petitions received to date by APHIS, including petitions pending, withdrawn, or approved.

Abbreviations:

CMV-cucumber mosaic virus; CPB-colorado potato beetle; PLRV- potato leafroll virus;
PRSV-papaya ringspot virus; PVY-potato virus Y; WMV2- watermelon mosaic virus 2
ZYMV-zucchini yellow mosaic virus

Petitions for Nonregulated Status Pending					
Applicant Documents		Institution	Regulated Article	Transgenic Phenotype	Transformation Event or Line
Petition	Extension of Petition Number ***				
10-070-01p		Virginia Tech	Peanut	Sclerotinia blight resistant	N70, P39, and W171
09-349-01p		Dow AgroSciences	Soybean	Herbicide Tolerant	DAS-68416-4
09-328-01p		Bayer Crop Science	Soybean	Herbicide Tolerant	FG72
09-233-01p		Dow	Corn	Herbicide Tolerant	DAS-40278-9
09-201-01p		Monsanto	Soybean		MON-877Ø5-6
09-183-01p		Monsanto	Soybean		MON-87769
09-082-01p		Monsanto	Soybean	Lepidopteran resistant	MON 87701
09-063-01p		Stine Seed	Corn	Glyphosate tolerant	HCEM485
09-055-01p		Monsanto	Corn	Drought Tolerant	MON 87460
09-015-01p		BASF Plant Science, LLC	Soybean	Herbicide Tolerant	BPS-CV127-9 Soybean
08-366-01p		ArborGen	Eucalyptus	Freeze Tolerant, Fertility Altered	ARB-FTE1-08
08-340-01p		Bayer	Cotton	Glufosinate Tolerant, Insect Resistant	T304-40XGHB119
08-338-01p		Pioneer	Corn	Male Sterile, Fertility Restored, Visual Marker	DP-32138-1
08-315-01p		Florigene	Rose	Altered Flower Color	IFD-524Ø1-4 and IFD-529Ø1-9
07-253-01p		Syngenta	Corn	Lepidopteran resistant	MIR-162 Maize
07-108-01p		Syngenta	Cotton	Lepidopteran Resistant	COT67B
06-354-01p		Pioneer	Soybean	High Oleic Acid	DP-3Ø5423-1
05-280-		Syngenta	Corn	Thermostable alpha-	3272

01p				amylase	
04-110-01p		Monsanto & Forage Genetics	Alfalfa	Glyphosate Tolerant	J101, J163
03-104-01p		Monsanto & Scotts	Creeping bentgrass	Glyphosate Tolerant	ASR368
Petitions for Nonregulated Status Granted					
Applicant Documents					
Petition	Extension of Petition Number ***	Institution	Regulated Article	Transgenic Phenotype	Transformation Event or Line
07-152-01p		Pioneer	Corn	glyphosate & Imidazolinone tolerant	DP-098140-6
04-337-01p		University of Florida	Papaya	Papaya Ringspot Virus Resistant	X17-2
06-332-01p		Bayer CropScience	Cotton	Glyphosate tolerant	GHB614
06-298-01p		Monsanto	Corn	European Corn Borer resistant	MON 89034
06-271-01p		Pioneer	Soybean	Glyphosate & acetolactate synthase tolerant	356043 (DP-356Ø43-5)
06-234-01p	98-329-01p	Bayer CropScience	Rice	Phosphinothricin tolerant	LLRICE601
06-178-01p		Monsanto	Soybean	Glyphosate tolerant	MON 89788
04-362-01p		Syngenta	Corn	Corn Rootworm Protected	MIR604
04-264-01p		ARS	Plum	Plum Pox Virus Resistant	C5
04-229-01p		Monsanto	Corn	High Lysine	LY038
04-125-01p		Monsanto	Corn	Corn Rootworm Resistant	88017
04-086-01p		Monsanto	Cotton	Glyphosate Tolerant	MON 88913
03-353-01p		Dow	Corn	Corn Rootworm Resistant	59122
03-323-01p		Monsanto	Sugar Beet	Glyphosate Tolerant	H7-1
03-181-01p	00-136-01p	Dow	Corn	Lepidopteran Resistant & Phosphinothricin tolerant	TC-6275
03-155-01p		Syngenta	Cotton	Lepidopteran Resistant	COT 102
03-036-01p		Mycogen/Dow	Cotton	Lepidopteran Resistant	281-24-236
03-036-02p		Mycogen/Dow	Cotton	Lepidopteran Resistant	3006-210-23

02-042-01p		Aventis	Cotton	Phosphinothricin tolerant	LLCotton25
01-324-01p	98-216-01p	Monsanto	Rapeseed	Glyphosate tolerant	RT200
01-206-01p	98-278-01p	Aventis	Rapeseed	Phosphinothricin tolerant & pollination control	MS1 & RF1/RF2
01-206-02p	97-205-01p	Aventis	Rapeseed	Phosphinothricin tolerant	Topas 19/2
01-137-01p		Monsanto	Corn	Corn Rootworm Resistant	MON 863
01-121-01p		Vector	Tobacco	Reduced nicotine	Vector 21-41
00-342-01p		Monsanto	Cotton	Lepidopteran resistant	Cotton Event 15985
00-136-01p		Mycogen c/o Dow & Pioneer	Corn	Lepidopteran resistant phosphinothricin tolerant	Line 1507
00-011-01p	97-099-01p	Monsanto	Corn	Glyphosate tolerant	NK603
99-173-01p	97-204-01p	Monsanto	Potato	PLRV & CPB resistant	RBMT22-82
98-349-01p	95-228-01p	AgrEvo	Corn	Phosphinothricin tolerant and Male sterile	MS6
98-335-01p		U. of Saskatchewan	Flax	Tolerant to soil residues of sulfonyl urea herbicide	CDC Triffid
98-329-01p		AgrEvo	Rice	Phosphinothricin tolerant	LLRICE06, LLRICE62
98-278-01p		AgrEvo	Rapeseed	Phosphinothricin tolerant & Pollination control	MS8 & RF3
98-238-01p		AgrEvo	Soybean	Phosphinothricin tolerant	GU262
98-216-01p		Monsanto	Rapeseed	Glyphosate tolerant	RT73
98-173-01p		Novartis Seeds & Monsanto	Beet	Glyphosate tolerant	GTSB77
98-014-01p	96-068-01p	AgrEvo	Soybean	Phosphinothricin tolerant	A5547-127
97-342-01p		Pioneer	Corn	Male sterile & Phosphinothricin tolerant	676, 678, 680
97-339-01p		Monsanto	Potato	CPB & PVY resistant	RBMT15-101, SEMT15-02, SEMT15-15
97-336-01p		AgrEvo	Beet	Phosphinothricin tolerant	T-120-7
97-287-01p		Monsanto	Tomato	Lepidopteran resistant	5345
97-265-01p		AgrEvo	Corn	Phosphinothricin tolerant & Lep. resistant	CBH-351
97-205-		AgrEvo	Rapeseed	Phosphinothricin	T45

01p			d	tolerant	
97-204-01p		Monsanto	Potato	CPB & PLRV resistant	RBMT21-129 & RBMT21-350
97-148-01p		Bejo	Cichorium intybus	Male sterile	RM3-3, RM3-4, RM3-6
97-099-01p		Monsanto	Corn	Glyphosate tolerant	GA21
97-013-01p		Calgene	Cotton	Bromoxynil tolerant & Lepidopteran resistant	Events 31807 & 31808
97-008-01p		Du Pont	Soybean	Oil profile altered	G94-1, G94-19, G-168
96-317-01p		Monsanto	Corn	Glyphosate tolerant & ECB resistant	MON802
96-291-01p		DeKalb	Corn	European Corn Borer resistant	DBT418
96-248-01p	92-196-01p	Calgene	Tomato	Fruit ripening altered	1 additional FLAVRSAVR line
96-068-01p		AgrEvo	Soybean	Phosphinothricin tolerant	W62, W98, A2704-12, A2704-21, A5547-35
96-051-01p		Cornell U	Papaya	PRSV resistant	55-1, 63-1
96-017-01p	95-093-01p	Monsanto	Corn	European Corn Borer resistant	MON809 & MON810
95-352-01p		Asgrow	Squash	CMV, ZYMV, WMV2 resistant	CZW-3
95-338-01p		Monsanto	Potato	CPB resistant	SBT02-5 & -7, ATBT04-6 & -27, -30, -31, -36
95-324-01p		Agritope	Tomato	Fruit ripening altered	35 1 N
95-256-01p		Du Pont	Cotton	Sulfonylurea tolerant	19-51a
95-228-01p		Plant Genetic Systems	Corn	Male sterile	MS3
95-195-01p		Northrup King	Corn	European Corn Borer resistant	Bt11
95-179-01p	92-196-01p	Calgene	Tomato	Fruit ripening altered	2 additional FLAVRSAVR lines
95-145-01p		DeKalb	Corn	Phosphinothricin tolerant	B16
95-093-01p		Monsanto	Corn	Lepidopteran resistant	MON 80100
95-053-01p		Monsanto	Tomato	Fruit ripening altered	8338
95-045-01p		Monsanto	Cotton	Glyphosate tolerant	1445, 1698
95-030-01p	92-196-01p	Calgene	Tomato	Fruit ripening altered	20 additional FLAVRSAVR lines

94-357-01p		AgrEvo	Corn	Phosphinothricin tolerant	T14, T25
94-319-01p		Ciba Seeds	Corn	Lepidopteran resistant	Event 176
94-308-01p		Monsanto	Cotton	Lepidopteran resistant	531, 757, 1076
94-290-01p		Zeneca & Petoseed	Tomato	Fruit polygalacturonase level decreased	B, Da, F
94-257-01p		Monsanto	Potato	Coleopteran resistant	BT6, BT10, BT12, BT16, BT17, BT18, BT23
94-230-01p	92-196-01p	Calgene	Tomato	Fruit ripening altered	9 additional FLAVRSAVR lines
94-228-01p		DNA Plant Tech	Tomato	Fruit ripening altered	1345-4
94-227-01p	92-196-01p	Calgene	Tomato	Fruit ripening altered	Line N73 1436-111
94-090-01p		Calgene	Rapeseed	Oil profile altered	pCGN3828-212/86- 18 & 23
93-258-01p		Monsanto	Soybean	Glyphosate tolerant	40-3-2
93-196-01p		Calgene	Cotton	Bromoxynil tolerant	BXN
92-204-01p		Upjohn	Squash	WMV2 & ZYMV resistant	ZW-20
92-196-01p		Calgene	Tomato	Fruit ripening altered	FLAVR SAVR

*** Extension of Petition Number: Under 7CFR 340.6(e) a person may request that APHIS extend a determination of non-regulated status to other organisms based on their similarity of the previously deregulated article. This column lists the previously granted petition of that deregulated article.

**** Preliminary EA: The Environmental Assessment initially available for Public comment prior to finalization.

Table C

5

Plant species	Event	Trait	Patent reference
Corn	PV-ZMGT32 (NK603)	Glyphosate tolerance	US 2007-056056
Corn	MIR604	Insect resistance (Cry3a055)	EP 1 737 290
Corn	LY038	High lysine content	US 7,157,281
Corn	3272	Self processing corn (alpha-amylase)	US 2006-230473
Corn	PV-ZMIR13 (MON863)	Insect resistance (Cry3Bb)	US 2006-095986
Corn	DAS-59122-7	Insect resistance (Cry34Ab1/Cry35Ab1)	US 2006-070139
Corn	TC1507	Insect resistance (Cry1F)	US 7,435,807
Corn	MON810	Insect resistance (Cry1Ab)	US 2004-180373
Corn	VIP1034	Insect resistance	WO 03/052073

Corn	B16	Glufosinate resistance	US 2003-126634
Corn	GA21	Glyphosate resistance	US 6,040,497
Corn	GG25	Glyphosate resistance	US 6,040,497
Corn	GJ11	Glyphosate resistance	US 6,040,497
Corn	FI117	Glyphosate resistance	US 6,040,497
Corn	GAT-ZM1	Glufosinate tolerance	WO 01/51654
Corn	MON87460	Drought tolerance	WO 2009/111263
Corn	DP-098140-6	Glyphosate tolerance / ALS inhibitor tolerance	WO 2008/112019
Wheat	Event 1	Fusarium resistance (trichothecene 3-O-acetyltransferase)	CA 2561992
Sugar beet	T227-1	Glyphosate tolerance	US 2004-117870
Sugar beet	H7-1	Glyphosate tolerance	WO 2004-074492
Soybean	MON89788	Glyphosate tolerance	US 2006-282915
Soybean	A2704-12	Glufosinate tolerance	WO 2006/108674
Soybean	A5547-35	Glufosinate tolerance	WO 2006/108675
Soybean	DP-305423-1	High oleic acid / ALS inhibitor tolerance	WO 2008/054747
Rice	GAT-OS2	Glufosinate tolerance	WO 01/83818
Rice	GAT-OS3	Glufosinate tolerance	US 2008-289060
Rice	PE-7	Insect resistance (Cry1Ac)	WO 2008/114282
Oilseed rape	MS-B2	Male sterility	WO 01/31042
Oilseed rape	MS-BN1/RF-BN1	Male sterility/restoration	WO 01/41558
Oilseed rape	RT73	Glyphosate resistance	WO 02/36831
Cotton	CE43-67B	Insect resistance (Cry1Ab)	WO 2006/128573
Cotton	CE46-02A	Insect resistance (Cry1Ab)	WO 2006/128572
Cotton	CE44-69D	Insect resistance (Cry1Ab)	WO 2006/128571
Cotton	1143-14A	Insect resistance (Cry1Ab)	WO 2006/128569
Cotton	1143-51B	Insect resistance (Cry1Ab)	WO 2006/128570
Cotton	T342-142	Insect resistance (Cry1Ab)	WO 2006/128568
Cotton	event3006-210-23	Insect resistance (Cry1Ac)	WO 2005/103266
Cotton	PV-GHGT07 (1445)	Glyphosate tolerance	US 2004-148666
Cotton	MON88913	Glyphosate tolerance	WO 2004/072235
Cotton	EE-GH3	Glyphosate tolerance	WO 2007/017186
Cotton	T304-40	Insect-resistance (Cry1Ab)	WO2008/122406
Cotton	Cot202	Insect resistance (VIP3)	US 2007-067868
Cotton	LLcotton25	Glufosinate resistance	WO 2007/017186
Cotton	EE-GH5	Insect resistance (Cry1Ab)	WO 2008/122406
Cotton	event 281-24-236	Insect resistance (Cry1F)	WO 2005/103266
Cotton	Cot102	Insect resistance (Vip3A)	US 2006-130175
Cotton	MON 15985	Insect resistance (Cry1A/Cry2Ab)	US 2004-250317
Bent Grass	Asr-368	Glyphosate tolerance	US 2006-162007

Brinjal

EE-1

Insect resistance (Cry1Ac)

WO 2007/091277

Among the diseases of plants or crops that can be controlled by the method according to the invention, mention can be made of :

Powdery mildew diseases such as :

- 5 Blumeria diseases, caused for example by *Blumeria graminis* ;
 Podosphaera diseases, caused for example by *Podosphaera leucotricha* ;
 Sphaerotheca diseases, caused for example by *Sphaerotheca fuliginea* ;
 Uncinula diseases, caused for example by *Uncinula necator* ;

Rust diseases such as :

- 10 Gymnosporangium diseases, caused for example by *Gymnosporangium sabinae* ;
 Hemileia diseases, caused for example by *Hemileia vastatrix* ;
 Phakopsora diseases, caused for example by *Phakopsora pachyrhizi* or *Phakopsora*
 meibomia ;
 Puccinia diseases, caused for example by *Puccinia recondite*, *Puccinia graminis* or
 15 *Puccinia striiformis* ;
 Uromyces diseases, caused for example by *Uromyces appendiculatus* ;

Oomycete diseases such as :

- Albugo diseases caused for example by *Albugo candida* ;
 Bremia diseases, caused for example by *Bremia lactucae* ;
 20 Peronospora diseases, caused for example by *Peronospora pisi* or *P. brassicae* ;
 Phytophthora diseases, caused for example by *Phytophthora infestans* ;
 Plasmopara diseases, caused for example by *Plasmopara viticola* ;
 Pseudoperonospora diseases, caused for example by *Pseudoperonospora humuli* or
 Pseudoperonospora cubensis ;
 25 Pythium diseases, caused for example by *Pythium ultimum* ;

Leafspot, leaf blotch and leaf blight diseases such as :

- Alternaria diseases, caused for example by *Alternaria solani* ;
 Cercospora diseases, caused for example by *Cercospora beticola* ;
 Cladosporium diseases, caused for example by *Cladosporium cucumerinum* ;
 30 Cochliobolus diseases, caused for example by *Cochliobolus sativus* (Conidiaform:
 Drechslera, Syn: Helminthosporium) or *Cochliobolus miyabeanus* ;
 Colletotrichum diseases, caused for example by *Colletotrichum lindemuthianum* ;
 Cycloconium diseases, caused for example by *Cycloconium oleaginum* ;
 Diaporthe diseases, caused for example by *Diaporthe citri* ;
 35 Elsinoe diseases, caused for example by *Elsinoe fawcettii* ;
 Gloeosporium diseases, caused for example by *Gloeosporium laeticolor* ;
 Glomerella diseases, caused for example by *Glomerella cingulata* ;
 Guignardia diseases, caused for example by *Guignardia bidwelli* ;
 Leptosphaeria diseases, caused for example by *Leptosphaeria maculans* ; *Leptosphaeria nodorum* ;
 40 Magnaporthe diseases, caused for example by *Magnaporthe grisea* ;

Mycosphaerella diseases, caused for example by *Mycosphaerella graminicola* ; *Mycosphaerella arachidicola* ; *Mycosphaerella fijiensis* ;

Phaeosphaeria diseases, caused for example by *Phaeosphaeria nodorum* ;

Pyrenophora diseases, caused for example by *Pyrenophora teres*, or *Pyrenophora tritici repentis*;

Ramularia diseases, caused for example by *Ramularia collo-cygni* , or *Ramularia areola*;

Rhynchosporium diseases, caused for example by *Rhynchosporium secalis* ;

Septoria diseases, caused for example by *Septoria apii* or *Septoria lycopersici* ;

Typhula diseases, caused for example by *Typhula incarnata* ;

Venturia diseases, caused for example by *Venturia inaequalis* ;

Root, Sheath and stem diseases such as :

Corticium diseases, caused for example by *Corticium graminearum* ;

Fusarium diseases, caused for example by *Fusarium oxysporum* ;

Gaeumannomyces diseases, caused for example by *Gaeumannomyces graminis* ;

Rhizoctonia diseases, caused for example by *Rhizoctonia solani* ;

Sarocladium diseases caused for example by *Sarocladium oryzae*;

Sclerotium diseases caused for example by *Sclerotium oryzae*;

Tapesia diseases, caused for example by *Tapesia acuformis* ;

Thielaviopsis diseases, caused for example by *Thielaviopsis basicola* ;

Ear and panicle diseases such as :

Alternaria diseases, caused for example by *Alternaria spp.* ;

Aspergillus diseases, caused for example by *Aspergillus flavus* ;

Cladosporium diseases, caused for example by *Cladosporium spp.* ;

Claviceps diseases, caused for example by *Claviceps purpurea* ;

Fusarium diseases, caused for example by *Fusarium culmorum* ;

Gibberella diseases, caused for example by *Gibberella zeae* ;

Monographella diseases, caused for example by *Monographella nivalis* ;

Smut and bunt diseases such as :

Sphacelotheca diseases, caused for example by *Sphacelotheca reiliana* ;

Tilletia diseases, caused for example by *Tilletia caries* ;

Urocystis diseases, caused for example by *Urocystis occulta* ;

Ustilago diseases, caused for example by *Ustilago nuda* ;

Fruit rot and mould diseases such as :

Aspergillus diseases, caused for example by *Aspergillus flavus* ;

Botrytis diseases, caused for example by *Botrytis cinerea* ;

Penicillium diseases, caused for example by *Penicillium expansum* ;

Rhizopus diseases caused by example by *Rhizopus stolonifer*

Sclerotinia diseases, caused for example by *Sclerotinia sclerotiorum* ;

Verticillium diseases, caused for example by *Verticillium alboatrum* ;

Seed and soilborne decay, mould, wilt, rot and damping-off diseases :

Alternaria diseases, caused for example by *Alternaria brassicicola*

Aphanomyces diseases, caused for example by *Aphanomyces euteiches*

Ascochyta diseases, caused for example by *Ascochyta lentis*
 Aspergillus diseases, caused for example by *Aspergillus flavus*
 Cladosporium diseases, caused for example by *Cladosporium herbarum*
 Cochliobolus diseases, caused for example by *Cochliobolus sativus*
 5 (Conidiaform: *Drechslera*, *Bipolaris* Syn: *Helminthosporium*);
 Colletotrichum diseases, caused for example by *Colletotrichum coccodes*;
 Fusarium diseases, caused for example by *Fusarium culmorum*;
 Gibberella diseases, caused for example by *Gibberella zeae*;
 Macrophomina diseases, caused for example by *Macrophomina phaseolina*
 10 Monographella diseases, caused for example by *Monographella nivalis*;
 Penicillium diseases, caused for example by *Penicillium expansum*
 Phoma diseases, caused for example by *Phoma lingam*
 Phomopsis diseases, caused for example by *Phomopsis sojae*;
 Phytophthora diseases, caused for example by *Phytophthora cactorum*;
 15 Pyrenophora diseases, caused for example by *Pyrenophora graminea*
 Pyricularia diseases, caused for example by *Pyricularia oryzae*;
 Pythium diseases, caused for example by *Pythium ultimum*;
 Rhizoctonia diseases, caused for example by *Rhizoctonia solani*;
 Rhizopus diseases, caused for example by *Rhizopus oryzae*
 20 Sclerotium diseases, caused for example by *Sclerotium rolfsii*;
 Septoria diseases, caused for example by *Septoria nodorum*;
 Typhula diseases, caused for example by *Typhula incarnata*;
 Verticillium diseases, caused for example by *Verticillium dahliae* ;

Canker, broom and dieback diseases such as :

25 Nectria diseases, caused for example by *Nectria galligena* ;

Blight diseases such as :

Monilinia diseases, caused for example by *Monilinia laxa* ;

Leaf blister or leaf curl diseases such as :

Exobasidium diseases caused for example by *Exobasidium vexans*

30 Taphrina diseases, caused for example by *Taphrina deformans* ;

Decline diseases of wooden plants such as :

Esca diseases, caused for example by *Phaemoniella clamydospora* ;

Eutypa dyebark, caused for example by *Eutypa lata* ;

Ganoderma diseases caused for example by *Ganoderma boninense*;

35 Rigidoporus diseases caused for example by *Rigidoporus lignosus*

Diseases of Flowers and Seeds such as

Botrytis diseases caused for example by *Botrytis cinerea*;

Diseases of Tubers such as

Rhizoctonia diseases caused for example by *Rhizoctonia solani*;

40 Helminthosporium diseases caused for example by *Helminthosporium solani*;

Club root diseases such as

Plasmodiophora diseases, cause for example by *Plasmodiophora brassicae*.

Diseases caused by Bacterial Organisms such as

Xanthomonas species for example *Xanthomonas campestris* pv. *oryzae*;

Pseudomonas species for example *Pseudomonas syringae* pv. *lachrymans*;

Erwinia species for example *Erwinia amylovora*.

5

The composition according to the invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds according to the invention or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

10

The dose of active compound usually applied in the method of treatment according to the invention is generally and advantageously from 10 to 800 g/ha, preferably from 50 to 300 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously from 2 to 200 g per 100 kg of seed, preferably from 3 to 150 g per 100 kg of seed in the case of seed treatment.

15

It is clearly understood that the doses indicated herein are given as illustrative examples of the method according to the invention. A person skilled in the art will know how to adapt the application doses, notably according to the nature of the plant or crop to be treated.

20


The compounds or mixtures according to the invention can also be used for the preparation of composition useful to curatively or preventively treat human or animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton diseases and candidiases or diseases caused by *Aspergillus spp.*, for example *Aspergillus fumigatus*.

25

The various aspects of the invention will now be illustrated with reference to the following table of compound examples and the following preparation or efficacy examples.

Table 1 illustrates in a non-limiting manner examples of compounds of formula (I) according to the invention :

30



In table 1, unless otherwise specified, M+H (Apcl+) means the molecular ion peak plus 1 a.m.u. (atomic mass unit) as observed in mass spectroscopy via positive atmospheric pressure chemical ionisation.









35











In table 1, the logP values were determined in accordance with EEC Directive 79/831 Annex V.A8 by HPLC (High Performance Liquid Chromatography) on a reversed-phase column (C 18), using the method described below :







Temperature: 40°C ; Mobile phases : 0.1% aqueous formic acid and acetonitrile ; linear gradient from 10% acetonitrile to 90% acetonitrile.

Calibration was carried out using unbranched alkan-2-ones (comprising 3 to 16 carbon atoms) with known logP values (determination of the logP values by the retention times using linear interpolation between two successive alkanones). lambda-max-values were determined using UV-spectra from 200 nm to 400 nm and the peak values of the chromatographic signals.

Table 1:

Example	A	T	n	Z ⁴	Z ⁵	Z ⁶		logP	Mass (M+H)
1		O	0			cyclopropyl	indan-2-yl	3.35	366
2		O	0			cyclopropyl	indan-2-yl	3.19	350
3		O	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	4.01	310
4		O	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.00	328
5		O	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.35	342
6		O	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.08	346
7		O	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.48	364

Example	A	T	n	Z ⁴	Z ⁵	Z ⁶		logP	Mass (M+H)
8		S	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.73	362
9		S	0			cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	4.21	380
10		O	1	Me	H	cyclopropyl	indan-2-yl	4.34	324
11		O	1	Me	H	cyclopropyl	indan-2-yl	3.25	342
12		O	1	Me	H	cyclopropyl	indan-2-yl	3.67	356
13		O	1	Me	H	cyclopropyl	indan-2-yl	3.33	360
14		O	1	Me	H	cyclopropyl	indan-2-yl	3.76	378
15		O	1	Me	H	cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	4.82	452
16		O	1	Me	H	cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	3.99	392

Example	A	T	n	Z ⁴	Z ⁵	Z ⁶		logP	Mass (M+H)
17		O	1	Et	H	cyclopropyl	1,2,3,4-tetrahydronaphthalen-2-yl	4.41	406
18		O	1	H	H	cyclopropyl	2,3-dihydro-1-benzofuran-2-yl	2.54	348
19		O	1	H	H	cyclopropyl	2,3-dihydro-1-benzofuran-2-yl	2.86	366
20		O	1	H	H	cyclopropyl	3,4-dihydro-2H-chromen-3-yl	3.00	380
21		O	1	H	H	cyclopropyl	2,3-dihydro-1-benzofuran-2-yl	3.04	382

The following examples illustrate in a non-limiting manner the preparation and efficacy of the compounds of formula (I) according to the invention.

5

Preparation example 1 : preparation of N-cyclopropyl-3-(difluoromethyl)-N-[1-(indan-2-yl)ethyl]-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide (compound 14)

Step 1 : preparation of N-[1-(indan-2-yl)ethyl]cyclopropanamine

10 To a cooled solution of 3.5 g (21.8 mmol) of 1-(indan-2-yl)ethanone in 100 ml of methanol, are added 10 g of 3Å molecular sieves and 2.3 g (40.2 mmol) of cyclopropylamine followed by a slow addition of 2.87 ml (50.2 mmol) of acetic acid. The reaction mixture is stirred for 2.5 hrs at reflux. The reaction mixture is then cooled to 0°C and 1.9 g (30 mmol) of sodium cyanoborohydride are slowly added and the reaction mixture is further stirred for 2 hrs at reflux. The cooled reaction mixture is then filtered over a cake of

15 diatomaceous earth. The cake is washed twice by 80 ml of methanol and the combined methanolic extracts are concentrated under vacuum. 100 ml of water are then added to the residue and the pH is

adjusted to 12 with a 0.5 N solution of sodium hydroxyde. The watery layer is extracted with 300 ml of ethyl acetate. The organic layer is washed twice by brine and filtered over a phase separator paper to yield after concentration 5 g of a yellow oil. Column chromatography on silica gel (gradient heptane/ethyl acetate) yields 1.2 g (28% yield) of N-[1-(indan-2-yl)ethyl]cyclopropanamine as a colourless oil (M+H = 202).

Step 2 : preparation of N-cyclopropyl-3-(difluoromethyl)-N-[1-(indan-2-yl)ethyl]-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide

At ambient temperature, a solution of 217 mg (1.02 mmol) of 3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carbonyl chloride in 1 ml of tetrahydrofurane is added dropwise to a solution of 202 mg (0.93 mmol) of N-[1-(indan-2-yl)ethyl]cyclopropanamine and 103 mg (1.02 mmol) of triethylamine in 5 ml of tetrahydrofurane. The reaction mixture is stirred for 1 hr at 70°C. The solvent is removed under vacuum and 100 ml of water are then added to the residue. The watery layer is extracted twice with ethyl acetate (2 x 150 ml) and the combined organic layers are successively washed by a 1 N solution of HCl, a saturated solution of potassium carbonate and filtered over a chemelut™ cardridge to yield after concentration 190 mg of a beige oil. Column chromatography on silica gel (gradient heptane/ethyl acetate) yields 120 mg (30% yield) of N-cyclopropyl-3-(difluoromethyl)-N-[1-(indan-2-yl)ethyl]-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide as a colourless oil (M+H = 378).

General preparation example: thionation of amide of formula (I) on Chemspeed apparatus

In a 13 ml Chemspeed vial is weighted 0.27mmole of phosphorous pentasulfide (P₂S₅). 3 ml of a 0.18 molar solution of the amide (I) (0.54mmole) in dioxane is added and the mixture is heated at reflux for two hours. The temperature is then cooled to 80°C and 2.5 ml of water are added. The mixture is heated at 80°C for one more hour. 2 ml of water are then added and the reaction mixture is extracted twice by 4 ml of dichloromethane. The organic phase is deposited on a basic alumina cartridge (2g) and eluted twice by 8 ml of dichloromethane. The solvents are removed and the crude thioamide derivative is analyzed by LCMS and NMR. Insufficiently pure compounds are further purified by preparative LCMS.

Example A : *in vivo* preventive test on *Alternaria solani* (tomato)

Solvent: 49 parts by weight of N,N-dimethylformamide

Emulsifier: 1 part by weight of Alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound at the stated rate of application. One day after this treatment, the plants are inoculated with an aqueous spore suspension of *Alternaria solani*. The plants remain for one day in an incubation cabinet at approximately 22°C and a relative atmospheric humidity of 100%. Then the plants are placed in an incubation cabinet at approximately 20°C and a relative atmospheric humidity of 96%.

The test is evaluated 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control while an efficacy of 100% means that no disease is observed.

Under these conditions, good (at least 80 %) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table A :

Table A :

Example	Efficacy	Example	Efficacy	Example	Efficacy
1	89	9	93	14	95
2	94	10	95	16	95
5	95	11	100	17	80
6	95	12	100	20	80
7	100	13	100		

Example B : *in vivo* preventive test on *Blumeria graminis* (barley)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has been dried, the plants are dusted with spores of *Blumeria graminis f.sp. hordei*.

The plants are placed in the greenhouse at a temperature of approximately 18°C and a relative atmospheric humidity of approximately 80% to promote the development of mildew pustules.

The test is evaluated 7 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed..

Under these conditions, total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table B :

Table B :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	13	100	16	100
12	100	14	100		

Example C : *in vivo* curative test on *Fusarium nivale* (var. *majus*) (wheat)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for curative activity, young plants are slightly injured by using a sandblast and afterwards they are sprayed with a conidia suspension of *Fusarium nivale* (var. *majus*) and placed for 24 hours in a greenhouse under a translucent incubation cabinet at a temperature of approximately 10°C and a relative atmospheric humidity of approximately 100% and are subsequently sprayed with the preparation of active compound at the stated rate of application.

After the spray coating has been dried, the plants remain in the greenhouse under translucent incubation cloches at a temperature of approximately 10°C and a relative atmospheric humidity of approximately 100%.

The test is evaluated 5 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table C :

Table C :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	13	100	16	100
12	100	14	100	17	100

Example D: *in vivo* preventive test on *Leptosphaeria nodorum* (wheat)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or the compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for protective activity, young plants are sprayed with a preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has dried on, the plants are sprayed with a spore suspension of *Leptosphaeria nodorum*. The plants remain for 48 hours in an incubation cabinet at 20°C and a relative atmospheric humidity of 100%.

The plants are placed in a greenhouse at a temperature of approximately 22°C and a relative atmospheric humidity of approximately 80%.

The test is evaluated 8 days after the inoculation. 0% means an efficacy which corresponds to that of the control, while an efficacy of 100% means that no disease is observed.

Under these conditions, good (at least 75 %) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table D :

Table D :

Example	Efficacy	Example	Efficacy	Example	Efficacy
2	94	11	100	14	100
5	78	12	100	16	86
7	94	13	100	17	80

Under the same conditions, excellent protection (greater than 94%) is observed at a dose of 500 ppm of active ingredient with compound 2, whereas poor protection (less than 30%) is observed with the isosteric analogue compound CMP1 claimed in WO-2009/016218 as in table D2.

Table D2 :

Example	dose (ppm)	Efficacy
2 from this invention	500	94
CMP1	500	30

The isosteric analogue compound CMP1 claimed in WO-2009/016218 corresponds to N-cyclopropyl-3-(difluoromethyl)-N-(indan-1-yl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide.

These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds.

Example E: in vivo preventive test on *Puccinia triticina* (wheat)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has been dried, the plants are sprayed with a spore suspension of *Puccinia triticina*. The plants remain for 48 hours in an incubation cabinet at approximately 20°C and a relative atmospheric humidity of approximately 100%.

The plants are placed in the greenhouse at a temperature of approximately 20°C and a relative atmospheric humidity of approximately 80%.

The test is evaluated 8 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, high (at least 85 %) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table E :

Table E :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	13	100	16	100
12	100	14	100	17	88

Example F: in vivo preventive test on *Pyrenophora teres* (barley)

Solvent: 49 parts by weight of N,N-dimethylformamide

Emulsifier: 1 part by weight of Alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound at the stated rate of application. One day after this treatment, the plants are inoculated with an aqueous spore suspension of *Pyrenophora teres*.

The plants remain for 48 hours in an incubation cabinet at 22°C and a relative atmospheric humidity of 100%. Then the plants are placed in a greenhouse at a temperature of approximately 20°C and a relative atmospheric humidity of approximately 80%.

The test is evaluated 7-9 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control while an efficacy of 100% means that no disease is observed.

Under these conditions, high (at least 90 %) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table F :

Table F :

Example	Efficacy	Example	Efficacy	Example	Efficacy
4	90	10	90	17	100
5	95	11	100	18	95
6	100	12	100	19	100
7	100	13	100	20	100
8	95	14	100		
9	100	16	100		

Example G: *in vivo* preventive test on *Septoria tritici* (wheat)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has dried on, the plants are sprayed with a spore suspension of *Septoria tritici*. The plants remain for 48 hours in an incubation cabinet at approximately 20°C and a relative atmospheric humidity of approximately 100% and then 60 hours at approximately 15°C in a translucent incubation cabinet at a relative atmospheric humidity of approximately 100%.

The plants are placed in a greenhouse at a temperature of approximately 15°C and a relative atmospheric humidity of approximately 80%.

The test is evaluated 21 days after the inoculation. 0% means an efficacy which corresponds to that of the control, while an efficacy of 100% means that no disease is observed.

Under these conditions, total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table G :

Table G :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	13	100	16	100
12	100	14	100	17	100

Example H: *in vivo* preventive test on *Sphaerotheca fuliginea* (cucumbers)

Solvent: 24.5 parts by weight of acetone

24.5 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for protect activity, young plants are sprayed with the preparation of active compound at the stated rate of application. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of *Sphaerotheca fuliginea*. The plants are then placed in a greenhouse at approximately 23°C and a relative atmospheric humidity of approximately 70 %.

The test is evaluated 7 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, high (at least 90 %) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table H :

Table H :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	92	13	95	16	100
12	95	14	100		

Under the same conditions, high protection (greater than 80%) is observed at a dose of 500 ppm of active ingredient with compound 7, whereas poor protection (less than 20%) is observed with the isosteric analogue compound CMP2 claimed in WO-2009/016218 as in table H2.

Table H2 :

Example	dose (ppm)	Efficacy
7 from this invention	500	80
CMP2	500	18

The isosteric analogue compound CMP2 claimed in WO-2009/016218 corresponds to N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-1H-pyrazole-4-carboxamide.

These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds.

Example I: *in vivo* preventive test on *Uromyces appendiculatus* (beans)

Solvent: 24.5 parts by weight of acetone

24.5 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound at the stated rate of application. After the spray coating has dried on, the plants are inoculated with an aqueous

spore suspension of the causal agent of bean rust (*Uromyces appendiculatus*) and then remain for 1 day in an incubation cabinet at approximately 20°C and a relative atmospheric humidity of 100 %.

The plants are then placed in a greenhouse at approximately 21°C and a relative atmospheric humidity of approximately 90 %.

- 5 The test is evaluated 10 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, excellent (at least 95 %) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table I :

10 Table I :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	12	100	13	98

Example J: *in vivo* preventive test on *Venturia inaequalis* (apples)

Solvent: 24.5 parts by weight of acetone

- 15 24.5 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

- 20 To test for preventive activity, young plants are sprayed with the preparation of active compound at the stated rate of application. After the spray coating has dried on, the plants are inoculated with an aqueous conidia suspension of the causal agent of apple scab (*Venturia inaequalis*) and then remain for 1 day in an incubation cabinet at approximately 20°C and a relative atmospheric humidity of 100 %.

- 25 The plants are then placed in a greenhouse at approximately 21°C and a relative atmospheric humidity of approximately 90 %.

The test is evaluated 10 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table J :


30

Table J :

Example	Efficacy	Example	Efficacy	Example	Efficacy
11	100	13	100	16	100
12	100	14	100		

CLAIMS

1. A compound of formula (I)



5 wherein

- A represents a carbo-linked, unsaturated or partially saturated, 5-membered heterocyclyl group that can be substituted by up to four groups R that can be the same or different providing that A do not represent the 3,4-dichloro-1,2-thiazol-5-yl group ;

- T represents O or S ;

10

- n represents 0 or 1 ;

- L¹ represents CZ¹Z², NZ⁷, O, S, S(O) or S(O)₂ ;

- L², L³ and L⁴ independently represents a direct bond, CZ¹Z², NZ⁷, O, S, S(O) or S(O)₂ ;

providing that when L² represents NZ⁷, O, S, S(O) or S(O)₂ then L³, L⁴ independently represent a direct bond or CZ¹Z²; or

15

providing that when L³ represents NZ⁷, O, S, S(O) or S(O)₂ then L² represent CZ¹Z² and L⁴ represent direct bond or CZ¹Z²; or

providing that when L⁴ represents NZ⁷, O, S, S(O) or S(O)₂ then L², L³ independently represent CZ¹Z²;

- m represents 0, 1, 2 or 3 ;

20

- X represents a halogen atom ; nitro ; cyano ; isonitrile ; hydroxy ; amino ; sulfanyl ; pentafluoro-λ⁶-sulfanyl ; formyl ; formyloxy ; formylamino ; substituted or non-substituted (hydroxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₁-C₈-alkoxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₂-C₈-alkenyloxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (C₂-C₈-alkynyloxyimino)-C₁-C₈-alkyl ; substituted or non-substituted (benzyloxyimino)-C₁-C₈-alkyl ;
- 25 carboxy ; carbamoyl ; N-hydroxycarbamoyl ; carbamate ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C₂-C₈-alkenyl ; C₂-C₈-halogenoalkenyl having 1 to 5 halogen atoms ; substituted or non-substituted C₂-C₈-alkynyl ; C₂-C₈-halogenoalkynyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxy ; C₁-C₈-halogenoalkoxy having 1 to 5 halogen atoms ; substituted or non-substituted
- 30 C₁-C₈-alkylsulfanyl ; C₁-C₈-halogenoalkylsulfanyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylsulfinyl ; C₁-C₈-halogenoalkylsulfinyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylsulfonyl ; C₁-C₈-halogenoalkylsulfonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylamino ; substituted or non-substituted di-C₁-C₈-alkylamino ; substituted or non-substituted C₂-C₈-alkenyloxy ; C₂-C₈-halogenoalkenyloxy

having 1 to 5 halogen atoms ; substituted or non-substituted C₃-C₈-alkynyloxy ; C₂-C₈-halogenoalkynyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₃-C₇-cycloalkyl ; C₃-C₇-halogenocycloalkyl having 1 to 5 halogen atoms ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₁-C₈-alkyl ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₂-C₈-alkenyl ; substituted or non-substituted (C₃-C₇-cycloalkyl)-C₂-C₈-alkynyl ; substituted or non-substituted tri(C₁-C₈)alkylsilyl ; substituted or non-substituted tri(C₁-C₈)alkylsilyl-C₁-C₈-alkyl ; substituted or non-substituted C₁-C₈-alkylcarbonyl ; C₁-C₈-halogenoalkylcarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbonyloxy ; C₁-C₈-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbonylamino ; C₁-C₈-halogenoalkyl-carbonylamino having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxycarbonyl ; C₁-C₈-halogenoalkoxycarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkyloxycarbonyloxy ; C₁-C₈-halogenoalkyloxycarbonyloxy having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkylcarbamoyl ; substituted or non-substituted di-C₁-C₈-alkylcarbamoyl ; substituted or non-substituted C₁-C₈-alkylaminocarbonyloxy ; substituted or non-substituted di-C₁-C₈-alkylaminocarbonyloxy ; substituted or non-substituted N-(C₁-C₈-alkyl)hydroxy carbamoyl ; substituted or non-substituted C₁-C₈-alkoxycarbamoyl ; substituted or non-substituted N-(C₁-C₈-alkyl)-C₁-C₈-alkoxycarbamoyl ; aryl that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkyl that can be substituted by up to 6 groups Q which can be the same or different ; C₂-C₈-arylalkenyl that can be substituted by up to 6 groups Q which can be the same or different ; C₂-C₈-arylalkynyl that can be substituted by up to 6 groups Q which can be the same or different ; aryloxy that can be substituted by up to 6 groups Q which can be the same or different ; arylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; arylamino that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkyloxy that can be substituted by up to 6 groups Q which can be the same or different ; C₁-C₈-arylalkylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; or C₁-C₈-arylalkylamino that can be substituted by up to 6 groups Q which can be the same or different ;

- two substituent X together with the consecutive carbon atoms to which they are linked can form a 5- or 6-membered, saturated carbocycle or saturated heterocycle, which can be substituted by up to four groups Q which can be the same or different ;
- Z¹, Z², Z³, Z⁴ and Z⁵ independently represents a hydrogen atom ; halogen atom ; cyano ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C₁-C₈-alkoxy; substituted or non-substituted C₁-C₈-alkylsulfanyl; or substituted or non-substituted C₁-C₈-alkoxycarbonyl ; or Z⁴ and Z⁵ are a C₂-C₅-alkylene group that can be substituted by up to four C₁-C₈-alkyl groups ;
- Z⁶ represents a non-substituted C₃-C₇-cycloalkyl or a C₃-C₇-cycloalkyl substituted by up to 10 atoms or groups that can be the same or different and that can be selected in the list consisting of halogen atoms, cyano, C₁-C₈-alkyl, C₁-C₈-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkoxy, C₁-C₈-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkoxycarbonyl, C₁-C₈-halogenoalkoxycarbonyl comprising up to 9 halogen atoms that can be the same or different, C₁-C₈-alkylaminocarbonyl and di-C₁-C₈-alkylaminocarbonyl ;


- Z^7 represents a hydrogen atom ; a substituted or non-substituted C_1 - C_8 -alkyl ; a C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_2 - C_8 -alkenyl ; a C_2 - C_8 -halogenoalkenyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_8 -alkynyl ; a C_3 - C_8 -halogenoalkynyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_7 -cycloalkyl ; a C_3 - C_7 -halogeno-cycloalkyl comprising up to 9 halogen atoms that can be the same or different ; a substituted or non-substituted C_3 - C_7 -cycloalkyl- C_1 - C_8 -alkyl ; or formyl ;
- Q independently represents a halogen atom ; cyano ; nitro ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; C_1 - C_8 -halogenoalkylsulfanyl having 1 to 9 halogen atoms that can be the same or different ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl- C_1 - C_8 -alkyl ; substituted or non-substituted (C_1 - C_8 -alkoxyimino)- C_1 - C_8 -alkyl ; substituted or non-substituted (benzyloxyimino)- C_1 - C_8 -alkyl ;
- R independently represents hydrogen atom ; halogen atom ; nitro ; cyano ; hydroxy ; amino ; sulfanyl ; pentafluoro- λ^6 -sulfanyl ; substituted or non-substituted (C_1 - C_8 -alkoxyimino)- C_1 - C_8 -alkyl ; substituted or non-substituted (benzyloxyimino)- C_1 - C_8 -alkyl ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C_2 - C_8 -alkenyl ; C_2 - C_8 -halogenoalkenyl having 1 to 5 halogen atoms ; substituted or non-substituted C_2 - C_8 -alkynyl ; C_2 - C_8 -halogenoalkynyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; C_1 - C_8 -halogenoalkylsulfanyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfinyl ; C_1 - C_8 -halogenoalkylsulfinyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylsulfonyl ; C_1 - C_8 -halogenoalkylsulfonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylamino ; substituted or non-substituted di- C_1 - C_8 -alkylamino ; substituted or non-substituted C_2 - C_8 -alkenyloxy ; substituted or non-substituted C_3 - C_8 -alkynyloxy ; substituted or non-substituted C_3 - C_7 -cycloalkyl ; C_3 - C_7 -halogenocycloalkyl having 1 to 5 halogen atoms ; substituted or non-substituted tri(C_1 - C_8)alkylsilyl ; substituted or non-substituted C_1 - C_8 -alkylcarbonyl ; C_1 - C_8 -halogenoalkylcarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkoxycarbonyl ; C_1 - C_8 -halogenoalkoxycarbonyl having 1 to 5 halogen atoms ; substituted or non-substituted C_1 - C_8 -alkylcarbamoyl ; substituted or non-substituted di- C_1 - C_8 -alkylcarbamoyl ; phenoxy ; phenylsulfanyl ; phenylamino ; benzyloxy ; benzylsulfanyl ; or benzylamino ;

as well as its salts, N-oxydes, metallic complexes, metalloidic complexes and optically active isomers.

2. A compound of formula (I) according to claim 1 wherein A is selected in the list consisting of:

- a heterocycle of formula (A^1)


61



wherein :

R¹ to R³ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;


- a heterocycle of formula (A²)



wherein :

R⁴ to R⁶ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A³)




wherein :

R⁷ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R⁸ represents a hydrogen atom or a substituted or non-substituted C₁-C₅-alkyl ;

- a heterocycle of formula (A⁴)




(A⁴)

wherein :

R⁹ to R¹¹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; amino ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₁-C₅-alkylsulfanyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A⁵)




wherein :

R¹² and R¹³ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; amino ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R¹⁴ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; amino ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A⁶)




wherein :

R¹⁵ represents a hydrogen atom ; a halogen atom ; a cyano ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R¹⁶ and R¹⁸ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkoxycarbonyl ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R¹⁷ represent a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

- a heterocycle of formula (A⁷)




5 wherein :

R¹⁹ represents a hydrogen atom or a C₁-C₅-alkyl

R²⁰ to R²² that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

10

- a heterocycle of formula (A⁸)




wherein :

15 R²³ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R²⁴ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different;

20 - a heterocycle of formula (A⁹)




wherein :

25 R²⁵ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R²⁶ represents a hydrogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A¹⁰)

64




wherein :

R²⁷ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R²⁸ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C₁-C₅-alkylamino or substituted or non-substituted di(C₁-C₅-alkyl)amino ;

- a heterocycle of formula (A¹¹)




wherein :

R²⁹ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R³⁰ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C₁-C₅-alkylamino or substituted or non-substituted di(C₁-C₅-alkyl)amino ;

- a heterocycle of formula (A¹²)



wherein :


R³¹ represents a hydrogen atom or a substituted or non-substituted C₁-C₅-alkyl

R³² represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R³³ represents a hydrogen atom ; a halogen atom ; a nitro ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms

that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A¹³)




wherein :

R³⁴ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₃-C₅-cycloalkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₂-C₅-alkynyloxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; R³⁵ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; a cyano ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₁-C₅-alkylsulfanyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C₁-C₅-alkylamino or substituted or non-substituted di(C₁-C₅-alkyl)amino;

R³⁶ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

-a heterocycle of formula (A¹⁴)




wherein :

R³⁷ and R³⁸ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C₁-C₅-alkoxy or a substituted or non-substituted C₁-C₅-alkylsulfanyl ;

R³⁹ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

- a heterocycle of formula (A¹⁵)




wherein :

R^{40} and R^{41} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

5


- a heterocycle of formula (A^{16})



wherein :

10 R^{42} and R^{43} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or amino ;

-a heterocycle of formula (A^{17})




15

wherein :

R^{44} and R^{45} that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

20

- a heterocycle of formula (A^{18})



25

wherein :


R^{47} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl or C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R^{46} represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C_1 - C_5 -alkyl ; C_1 - C_5 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or substituted or non-substituted C_1 - C_5 -alkylsulfanyl ;

30

- a heterocycle of formula (A^{19})


67



wherein :

R⁴⁹ and R⁴⁸ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;


- a heterocycle of formula (A²⁰)



wherein :

R⁵⁰ and R⁵¹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted or non-substituted C₁-C₅-alkoxy ; C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;


-a heterocycle of formula (A²¹)



wherein :

R⁵² represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.


-a heterocycle of formula (A²²)



wherein :

R⁵³ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.

-a heterocycle of formula (A²³)




5 wherein :

R⁵⁴ and R⁵⁶ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R⁵⁵ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

10

-a heterocycle of formula (A²⁴)




wherein :

15 R⁵⁷ and R⁵⁹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R⁵⁸ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

20 -a heterocycle of formula (A²⁵)




wherein :

25 R⁶⁰ and R⁶¹ that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl or C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R⁶² represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl ;

-a heterocycle of formula (A²⁶)



wherein :

R⁶⁵ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; substituted
 5 or non-substituted C₃-C₅-cycloalkyl ; C₁-C₅-halogenoalkyl comprising up to 9 halogen atoms that can be
 the same or different ; substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₂-C₅-
 alkynyloxy or C₁-C₅-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;
 R⁶³ represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C₁-C₅-alkyl ; a cyano ;
 substituted or non-substituted C₁-C₅-alkoxy ; substituted or non-substituted C₁-C₅-alkylsulfanyl ; C₁-C₅-
 10 halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C₁-C₅-halogenoalkoxy
 comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-
 substituted C₁-C₅-alkylamino or di(C₁-C₅-alkyl)amino ;
 R⁶⁴ represents a hydrogen atom or substituted or non-substituted C₁-C₅-alkyl.

3. A compound of formula (I) according to any one of claims 1 to 2 wherein A is selected in the list
 consisting of A² ; A⁶ ; A¹⁰ and A¹³.

4. A compound of formula (I) according to any one of claims 1 to 3 wherein A represents A¹³ wherein R³⁴
 represents a substituted or non-substituted C₁-C₅-alkyl, C₁-C₅-halogenoalkyl comprising up to 9 halogen
 20 atoms that can be the same or different ; or a substituted or non-substituted C₁-C₅-alkoxy ; R³⁵ represents
 a hydrogen atom or a halogen atom and R³⁶ represents a substituted or non-substituted C₁-C₅-alkyl.

5. A compound of formula (I) according to any one of claims 1 to 4 wherein T represents O.

6. A compound of formula (I) according to any one of claims 1 to 5 wherein L¹ or L² independently
 represents O or CZ¹Z².

7. A compound of formula (I) according to any one of claims 1 to 6 wherein L³ or L⁴ independently
 represents a direct bond, O or CZ¹Z².

8. A compound of formula (I) according to anyone of claims 1 to 7 wherein X independently represents a
 halogen atom ; substituted or non-substituted C₁-C₈-alkyl ; C₁-C₈-halogenoalkyl comprising up to 9
 halogen atoms that can be the same or different ; substituted or non-substituted tri(C₁-C₈-alkyl)silyl ;
 substituted or non-substituted C₁-C₈-alkoxy or C₁-C₈-halogenoalkoxy comprising up to 9 halogen atoms
 35 that can be the same or different; or wherein two consecutive substituents X together with the phenyl ring
 form a substituted or non substituted 1,3-benzodioxolyl ; 1,2,3,4-tetrahydro-quinoxaliny ; 3,4-dihydro-2H-
 1,4-benzoxazinyl ; 1,4-benzodioxanyl ; indanyl ; 2,3-dihydrobenzofuranyl ; or indoliny.

9. A compound of formula (I) according to any one of claims 1 to 8 wherein Z^1 , Z^2 , Z^3 , Z^4 and Z^5 independently represents a hydrogen atom, halogen, substituted or non-substituted C_1 - C_8 -alkyl or substituted or non-substituted C_1 - C_8 -alkoxy.

10. A compound of formula (I) according to any one of claims 1 to 9 wherein Z^6 represents a non-substituted C_3 - C_7 cycloalkyl or a C_3 - C_7 cycloalkyl substituted by up to 10 groups or atoms that can be the same or different and that can be selected in the list consisting of halogen atoms, C_1 - C_8 -alkyl, C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, C_1 - C_8 -alkoxy and C_1 - C_8 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different.


11. A compound of formula (I) according to any one of claims 1 to 10 wherein Z^6 represents a non-substituted C_3 - C_7 -cycloalkyl.

12. A compound of formula (I) according to any one of claims 1 to 11 wherein Z^6 represents cyclopropyl.

13. A compound of formula (I) according to any one of claims 1 to 12 wherein Z^7 represents a substituted or non-substituted C_1 - C_8 -alkyl.

14. A compound of formula (I) according to any one of claims 1 to 13 wherein R independently represents a hydrogen atom ; halogen atom ; cyano ; substituted or non-substituted C_1 - C_8 -alkylamino ; substituted or non-substituted di- C_1 - C_8 -alkylamino ; substituted or non-substituted tri(C_1 - C_8 -alkyl)silyl ; substituted or non-substituted C_1 - C_8 -alkyl ; C_1 - C_8 -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkoxy ; C_1 - C_8 -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C_1 - C_8 -alkylsulfanyl ; amino ; hydroxyl ; nitro ; substituted or non-substituted C_1 - C_8 -alkoxycarbonyl ; or substituted or non-substituted C_2 - C_8 -alkynyloxy.

15. A compound of formula (II)



(II)

wherein X, m, n, L^1 , L^2 , L^3 , L^4 , Z^3 , Z^4 and Z^5 are as defined in any one of claims 1 to 12, and Z^6 represents a cyclopropyl group with the exclusion of

- substituted N-cyclopropyl-1,2,3,4-tetrahydronaphthalen-2-amines,
- substituted N-cyclopropylchroman-3-amines,
- N-cyclopropyl-2,3-dihydro-1H-inden-2-amine,
- N-(3,4-dihydro-2H-chromen-3-ylmethyl)cyclopropanamine,
- N-(2,3-dihydro-1-benzofuran-2-ylmethyl)cyclopropanamine.

16. A fungicide composition comprising, as an active ingredient, an effective amount of a compound of formula (I) according to claims 1 to 14 and an agriculturally acceptable support, carrier or filler.

- 5 17. A method for controlling phytopathogenic fungi of crops, characterized in that an agronomically effective and substantially non-phytotoxic quantity of a compound according to claims 1 to 14 or a composition according to claim 16 is applied to the soil where plants grow or are capable of growing, to the leaves and/or the fruit of plants or to the seeds of plants.