



(51) International Patent Classification:

A61K 33/38 (2006.01) A61P 17/02 (2006.01)
A61K 36/00 (2006.01)

(21) International Application Number:

PCT/GB2018/050869

(22) International Filing Date:

29 March 2018 (29.03.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1705140.0 30 March 2017 (30.03.2017) GB

(71) Applicant: **PHYTO PHARM LIMITED** [GB/GB]; 5 Jupiter House, Calleva Park, Aldermaston, Reading Berkshire RG7 8NN (GB).

(72) Inventor: **DEWAR, Laura**; 1B Camden Wood, 48 Yester Road, Chislehurst Kent BR7 5HR (GB).

(74) Agent: **EIP**; Fairfax House, 15 Fulwood Place, London Greater London WC1V 6HU (GB).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COLLOIDAL SILVER COMBINED WITH PLANT EXTRACTS FOR USE IN TREATING WOUNDS AND OTHER SKIN CONDITIONS




FIG 1

(57) Abstract: In one aspect, a composition for topical application is provided. The composition comprises a base. Dispersed within the base is elemental silver as a colloidal suspension, and plant extract from at least 6 plants. The plants are selected from the group consisting of Echinacea purpurea, Stellaria media, Aloe vera, Matricaria recutita, Hypericum perforatum, Calendula officinalis, Equisetum arvense, Symphytum officinale, Panax ginseng, Rumex crispus, Arctium lappa, Trifolium pratense, Chelidonium majus, Thuja occidentalis, Urtica dioica, Mahonia aquifolium, and Galium aparine.



COLLOIDAL SILVER COMBINED WITH PLANT EXTRACTS FOR USE IN
TREATING WOUNDS AND OTHER SKIN CONDITIONS

Technical Field

The present invention relates to compositions for topical application. In particular, the
5 present invention relates to compositions for topical application comprising colloidal
silver and a combination of plant extracts, for use as cosmetics and/or medicaments.

Background

There are a number of dermatological conditions and cosmetic skin conditions which
10 may cause distress to humans or animals. Said distress may be due to sensations such
as pain, itching, or burning. Said distress may alternatively or additionally be due to an
unsightly appearance associated with the condition, such as reddening of the skin, dry
skin, or pigmentation of the skin.

15 Sufferers presenting with these conditions often wish to alleviate their distress by using
some form of remedy for the condition. One option for addressing these conditions may
be the use of prescription or over-the-counter pharmaceutical preparations. However,
there is a consumer demand for natural remedies, or non-pharmaceutical remedies.

20 Given that these conditions may affect only a portion of the surface of the body of a
sufferer, there is a consumer demand for a local approach of addressing the
dermatological condition or cosmetic skin condition, rather than a systemic approach.
One such suitable approach may be a topical application of a remedy which may reduce
or diminish the dermatological condition or cosmetic skin condition.

25 Accordingly, there is a need for compositions suitable for topical application to
dermatological conditions and/or cosmetic skin conditions which may at least partially
alleviate pain, itching, burning, reddening of the skin, dry skin, or pigmentation of the
skin, as appropriate.

30

Summary

According to a first aspect of the present invention, there is provided a composition for topical application comprising a base, and dispersed within the base:

elemental silver as a colloidal suspension; and

- 5 plant extract from at least 6 plants selected from the group consisting of *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, *Equisetum arvense*, *Symphytum officinale*, *Panax ginseng*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Mahonia aquifolium*, and *Galium aparine*.

10

According to further aspects, there is provided:

- 2) A composition according to 1), wherein the composition comprises:

base in an amount of from 80% to 99.9% by volume/volume of the composition;

elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by

- 15 weight/volume of the composition; and

plant extract in an amount of from 0.1% to 10% by weight/volume of the composition.

- 3) A composition according to 1) or 2), wherein the composition is an aqueous
20 suspension, a cream, an ointment, a spray, a gel, or an aerosol.

- 4) A composition according to 3), wherein the composition is an aqueous suspension, and the base is water.

- 25 5) A composition according to 4), wherein the composition consists essentially of:
elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by
weight/volume of the composition;

plant extract in an amount of from 0.1% to 10% by weight/volume of the composition; and

- 30 water as the remainder of the composition up to 100%.

- 6) A composition according to any of 1) to 5), wherein the plant extract is an extract in an extractant selected from C₁-C₄ alkanol, acetic acid, glycerine, propylene glycol, honey, water, and mixtures thereof.
- 5 7) A composition according to 6), wherein the extractant consists of aqueous ethanol.
- 8) A composition according to 6) or 7), wherein the plant extract is an extract of plant material in a ratio of from 1:1 to 1:12 weight/volume of plant material to
10 extractant.
- 9) A composition according to any of 1) to 5), wherein the extract is an extractant-free extract.
- 15 10) A composition according to any of 1) to 9), wherein the plant extracts are from *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 20 11) A composition according to any of 1) to 9), wherein the plant extracts are from *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 25 12) A composition according to any of 1) to 9), wherein the plant extracts are from *Symphytum officinale*, *Echinacea purpurea*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

- 13) A composition according to any of 1) to 9), wherein the plant extracts are from *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 5 14) A composition according to any of 1) to 9), wherein the plant extracts are from *Chelidonium majus*, *Thuja occidentalis*, *Echinacea purpurea*, *Galium aparine*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 10 15) A composition according to any of 1) to 9), wherein the plant extracts are from *Equisetum arvense*, *Symphytum officinale*, *Mahonia aquifolium*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 15 16) A composition according to any of 1) to 9), wherein the plant extracts are from *Arctium lappa*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 20 17) A composition according to any of 1) to 9), wherein the plant extracts are from *Chelidonium majus*, *Thuja occidentalis*, *Mahonia aquifolium*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*.
- 25 18) A composition according to any of 1) to 9), wherein the plant extracts are from *Arctium lappa*, *Urtica dioica*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 30 19) A composition according to any of 1) to 9), wherein the plant extracts are from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*.
- 20) A composition according to any of 1) to 9), wherein the plant extracts are from *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*,

Stellaria media, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

21) A composition according to any of 1) to 9), wherein the plant extracts are from
5 *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

22) A composition according to any of 1) to 9), wherein the plant extracts are from
10 *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*.

23) A composition according to any of 1) to 9), wherein the plant extracts are from
15 *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, and *Calendula officinalis*.

24) A composition according to any of 1) to 23) which is a cosmetic composition.

25) A composition according to any of 1) to 23) for use as a medicament.
20

26) A composition according to any of 1) to 10) for use in the treatment of dermatitis.

27) A composition according to any of 1) to 9) or 11) for use in the treatment of
25 severed skin.

28) A composition according to any of 1) to 9) or 12) for use in the treatment of burns.

30 29) A composition according to any of 1) to 9) or 13) for use in the treatment of sunburn.

- 30) A composition according to any of 1) to 9) or 14) for use in the treatment of tinea pedis.
- 5 31) A composition according to any of 1) to 9) or 15) for use in the treatment of blisters.
- 32) A composition according to any of 1) to 9) or 18) for use in the treatment of insect bites.
- 10 33) A composition according to any of 1) to 9) or 19) for use in the treatment of irritant diaper dermatitis.
- 34) A composition according to any of 1) to 9) or 20) for use in the treatment of contact dermatitis related to incontinence.
- 15 35) A method of manufacturing a composition for topical application, the method comprising:
- providing plant extract from at least 6 plants selected from the group consisting of *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*;
- 20 providing a colloidal suspension of elemental silver in a base, the elemental silver content of the colloidal suspension being of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by weight/volume of the base; and
- 25 combining the plant extract and the colloidal suspension of elemental silver to provide a composition for topical application.
- 30 36) A method according to 35), wherein the base is water.

- 37) A method according to 35) or 36), wherein the plant extract and colloidal suspension of elemental silver are combined in a ratio of from 1:20 to 1:4 weight/volume plant extract to colloidal suspension of elemental silver.
- 5 38) A dressing comprising a composition according to any of 1) to 34).
- 39) A porous sheet impregnated with a composition according to any of 1) to 34).
- 40) A method of treating an area affected by a dermatological condition comprising:
10 providing a composition according to any of 1) to 34); and
topically applying the composition to the area affected by the dermatological condition.
- 41) A non-therapeutic cosmetic method comprising:
15 providing a composition according to any of 1) to 34);
selecting an area of the body to which the composition should be applied; and
topically applying the composition to the selected area.

Further features and advantages of the invention will become apparent from the
20 following description of preferred embodiments of the invention, given by way of
example only, which is made with reference to the accompanying drawings.

Brief Description of the Drawings

Figure 1 shows a subject presenting with an open wound.
25

Figures 2, 3 and 4 show the variation in appearance of the wound of Figure 1 over time after application of a composition of the present invention to the affected area.

Figures 5 and 6 show the variation in appearance of a nevi over a period of time after
30 application of a composition of the present invention to the affected area.

Figure 7 shows a subject presenting with two insect bites on the skin.

Figures 8 and 9 show the variation in appearance of the insect bites of Figure 7 over time after application of a composition of the present invention to the affected area.

5

Figure 10 shows a subject presenting with an open wound.

Figures 11 and 12 show the variation in appearance of the wound of Figure 10 over time after application of a composition of the present invention to the affected area.

10

Detailed Description

In the specification and claims, a volume to volume ratio or percentage may be indicated by “(v/v)”. A weight to volume ratio or percentage may be indicated by “(w/v)”. A weight to weight ratio or percentage may be indicated by “(w/w)”. Unless otherwise specified, ratios or percentages are given in relation to the weight or volume of the total composition, as appropriate.

15

The unit “ppm” refers to “parts per million”, and may be used herein to indicate a concentration of a material in a fluid phase. Unless otherwise specified, all values given as ppm are given as weight to volume (w/v) percentages. Accordingly, 1 ppm is equivalent to $1 \times 10^{-4} \%$ (w/v).

20

The compositions of the present invention are suitable for topical application. That is, the compositions may be applied to body surfaces such as the skin or mucous membranes.

25

Base

The composition comprises a base, or carrier. The base is used as a vehicle for the other components of the composition. For the avoidance of doubt, the term ‘base’ as used herein does not mean a Brønsted-Lowry base or a Lewis base.

30

The base is preferably a pharmaceutically acceptable base. The nature of the base may influence the formulation of the composition. For example, a suitable base may comprise water (in particular water purified by distillation, ion exchange, reverse osmosis or any other suitable method), or a mixture of water and C₁-C₄ alkanol. A composition comprising such a base may suitably provide a topically applicable aqueous suspension formulation. Depending on the ratio of water to C₁-C₄ alkanol, a topically applicable aqueous suspension may be more suitable for spray application (that is, the suspension is applied to the skin as droplets in an aerosol), or may be more suitable for manual application (that is, the user applies the suspension to the skin or mucous membrane with a cloth, pad, or part of their body, for example their hand).

In another example, a suitable base may be an oil, or emulsion of oil and water. In some embodiments, the base may be an ointment base.

Any suitable ointment base may be used. A suitable ointment base may comprise an oleaginous base. Oleaginous bases typically comprise hard paraffin, soft paraffin (also known as petroleum jelly, petrolatum, white petrolatum or multi-hydrocarbon), microcrystalline wax or ceresine. Such bases are typically anhydrous, or have a low water content. These bases may further comprise a film forming agent to provide a protective film after application to the skin. Suitable film forming agents include polyvinylpyrrolidone, polyvinyl alcohol and cellulose derivatives.

Another suitable ointment base may comprise an absorption base (also known as an emulsion base). Absorption ointment bases typically comprise either an oleaginous base as described above and a water-in-oil surfactant, or a hydrophilic oleaginous base, to allow for the formation of a water-in-oil emulsion upon addition of water. Such bases may comprise any of the oleaginous bases described above and/or components selected from: lanolin, beeswax, cetyl alcohol, stearyl alcohol, cetostearyl alcohol, and sodium lauryl ether sulfate. Examples of appropriate absorption bases include hydrophilic petrolatum, anhydrous lanolin, Aquabase®, Aquaphor® and Polysorb®.

Another suitable ointment base may comprise a water-in-oil emulsion base. Water-in-oil emulsion bases typically comprise water and any of the ointment base components described above. In particular, water-in-oil emulsion bases typically comprise a water-in-oil surfactant. Examples of appropriate water-in-oil emulsion bases include cold
5 creams, hydrous lanolin, rose water ointment, aqueous cream, Hydrocream®, Eucerin® and Nivea®.

Another suitable ointment base may comprise an oil-in-water emulsion base. Oil-in-water emulsion bases typically comprise water in an amount greater than that in a water-
10 in-oil emulsion base, and any of the ointment base components described above. In particular, oil-in-water emulsion bases typically comprise an oil-in-water surfactant. Examples of appropriate oil-in-water emulsion bases include hydrophilic ointment, Dermabase®, Velvachol® and Unibase®.

15 Another suitable ointment base may comprise a water-miscible ointment base. Water-miscible ointment bases typically comprise a combination of low and high molecular weight polyethylene glycols. Examples of appropriate water-miscible ointment bases include PEG ointment and Polybase®.

20 A composition comprising such ointment bases as described above may suitably provide a topically applicable ointment formulation, or topically applicable cream formulation.

In another example, a suitable base may be a gel. 'Gel' as used herein refers to any
25 dispersion of inorganic or organic molecules in a liquid system wherein there is at least some cross-linking or association between the inorganic or organic molecules. Gels may be prepared by combining a liquid such as water or aqueous C₁-C₄ alkanol with suitable gelling agent(s), such as carbomers, carboxymethyl cellulose, ethylcellulose, gelatin, hydroxyethyl cellulose, hydroxypropyl cellulose, magnesium aluminium
30 silicate, methylcellulose, poloxamers, polyvinyl alcohol, sodium alginate, tragacanth,

and xanthan gum. A composition comprising such a base may suitably provide a topically applicable gel formulation.

5 In another example, a suitable base may comprise water and/or C₁-C₄ alkanol and a propellant. Suitable propellants may include hydrocarbons, hydrofluorocarbons, and compressed gases. A composition comprising such a base may suitably provide a topically applicable aerosol formulation.

10 The base may be present in the composition in any suitable amount. Preferably, the base is the majority component of the composition. For example, the base may be present in the composition in an amount of from 80 to 99.9% (v/v), or 85 to 99.5% (v/v), or from 90 to 99% (v/v). The base may be present in the composition in an amount less than or equal to 99.9% (v/v), or 99.5% (v/v), or 99% (v/v), or 98% (v/v), or 97% (v/v), or 96% (v/v), or 95% (v/v), or 90% (v/v). The base may be present in the composition in an
15 amount greater than or equal to 80% (v/v), 85% (v/v), 90%, 95% (v/v), 96% (v/v), 97% (v/v), 98% (v/v), 99% (v/v), 99.5% (v/v), or 99.9% (v/v).

In a preferred embodiment, the base comprises water. In a particular embodiment, the base essentially consists of water.

20

Silver

The composition also comprises elemental silver as colloidal silver, namely particulate elemental silver dispersed within the base as a colloidal suspension. The silver may be present in the composition in any suitable amount, for example in an amount of from
25 0.1 ppm to 100 ppm, or from 0.5 ppm to 50 ppm, or from 1 ppm to 10 ppm, or from 5 ppm to 10 ppm; that is, from 1×10^{-5} to 1×10^{-2} % (w/v), or from 5×10^{-5} to 5×10^{-3} % (w/v), or from 1×10^{-4} to 1×10^{-3} % (w/v), or from 5×10^{-4} to 1×10^{-3} % (w/v). The silver may be present in the composition in an amount less than or equal to 100 ppm, or 50 ppm, or 10 ppm; that is, less than or equal to 1×10^{-2} % (w/v), or
30 5×10^{-3} % (w/v), or 1×10^{-3} % (w/v). The silver may be present in the composition in an amount greater than or equal to 0.1 ppm, or 0.5 ppm, or 1 ppm; that is, greater than

or equal to 1×10^{-5} % (w/v), or 5×10^{-5} % (w/v), or 1×10^{-4} % (w/v). Suitably, the silver may be present in the composition in an amount of approximately 5 ppm; that is, 5×10^{-4} % (w/v).

5 **Plant extract**

Compositions according to the present invention comprise plant extract. A plant extract refers to a substance that has been removed from plant material. Plant extracts may be prepared by treating any plant material with an extractant to extract at least some components of the plant material from the plant material into the extractant. Extractant
10 may be referred to as solvent. The resulting extractant containing at least some components extracted from the plant material is referred to herein as “plant extract”. Put another way, “plant extract” may refer to components removed from plant material by treating the plant material with an extractant.

15 The plant material which is treated with extractant to provide a plant extract may be or derive from any part of a plant, for example, the leaves, bark, roots, seeds or fruits of the plant. In the preparation of plant extract, the plant components extracted from the plant material will be at least partially soluble or suspendable in the extractant. After the plant material has been treated with the extractant, the insoluble plant material may
20 be left in the extractant, or the insoluble plant material may be separated from the extractant. The insoluble plant material may be separated from the extractant according to any appropriate method, for example, decantation, filtration or centrifugation. In preferred embodiments of the present invention, the insoluble plant material has been separated from the extractant; the plant extract does not comprise insoluble
25 components. In a particularly preferred embodiment, the plant extract is prepared by cold percolation extraction. Cold percolation extraction includes placing plant material in a perforated flask or vessel, and treating the plant material with cold extractant. The resulting plant extract will pass through the perforation, leaving the insoluble and/or insuspendable plant material in the flask or vessel.

Suitable plant extract extractants include any which may be used to extract components from the plant material; that is, any extractant in which at least some of the components of the plant material are soluble or suspendable. In one embodiment, the extractant may comprise a C₁-C₄ alkanol. Suitable C₁-C₄ alkanols include methanol, ethanol, *n*-propanol, isopropanol, *n*-butanol, isobutanol, *sec*-butanol, and *tert*-butanol. In one
5 embodiment, the extractant may comprise methanol and/or ethanol. In a particular embodiment, the extractant may comprise ethanol.

In another embodiment, the extractant may comprise acetic acid, glycerine (in particular
10 vegetable glycerine), propylene glycol, honey, or water. Plant extracts may be prepared with hot water or cold water.

The extractant may comprise a combination of extractants. For example, the extractant may comprise a combination of any of the extractants given above. In a particular
15 embodiment, the extractant may comprise water and a C₁-C₄ alkanol. Such a combination may be referred to as an aqueous C₁-C₄ alkanol. In a preferred embodiment, the extractant comprises water and ethanol. Such a combination may be referred to as aqueous ethanol.

In a further embodiment, the extractant may essentially consist of any one of the
20 extractants given above, or a combination thereof. In a particular embodiment, the extractant may consist of water and a C₁-C₄ alkanol. In a preferred embodiment, the extractant consists of water and ethanol.

Aqueous ethanol may have any appropriate ethanol content. For example, the aqueous
25 ethanol may have an ethanol content of from 5 to 95% (v/v), 10 to 90% (v/v), 20 to 80% (v/v), 30 to 70% (v/v), or 40 to 60% (v/v). The aqueous ethanol may have an ethanol content of greater than or equal to 5% (v/v), 10% (v/v), 20% (v/v), 30% (v/v), 40% (v/v), 50% (v/v), 60% (v/v), 70% (v/v), 80% (v/v), 90% (v/v), or 95% (v/v). The
30 aqueous ethanol may have an ethanol content of less than or equal to 95% (v/v), 90%

(v/v), 80% (v/v), 70% (v/v), 60% (v/v), 50% (v/v), 40% (v/v), 30% (v/v), 20% (v/v), 10% (v/v), or 5% (v/v).

5 The plant extracts suitable for use in the present invention may be prepared by treating plant material with extractant in any appropriate ratio. For example, plant extracts for use in the present invention may be prepared by treating plant material with extractant in a ratio of from 2:1 to 1:20 (w/v) plant material to extractant, or from 1:1 to 1:10 (w/v) plant material to extractant, or from 1:2 to 1:9 (w/v) plant material to extractant.

10 Plant extracts suitable for use in the present invention may be referred to by a variety of names. Plant extracts suitable for use in the present invention particularly include products which may commercially be referred to as extracts (typically an aqueous ethanol extraction of plant material carried out in a ratio of approximately 1:1 (w/v) plant material to extractant), tinctures (typically an aqueous ethanol extraction of plant
15 material carried out in a ratio of approximately 1:3 to 1:5 (w/v) plant material to extractant), mother tinctures (typically an aqueous ethanol extraction of plant material carried out in a ratio of approximately 1:9 (w/v) plant material to extractant), and tisanes such as infusions (typically a hot water extraction of plant material), decoctions (typically an aqueous extraction of plant material carried out in boiling water over an
20 extended period of time) and macerations (typically an extraction of chopped plant material in cold water).

Plant extracts suitable for use with the present invention may be purchased from the vendor G Baldwin & Co.

25

In some embodiments, the plant extract is an extractant-free extract. That is, a plant extract has been prepared according to the above methods, and then the extractant has been removed to leave a dry substance. The dry substance may correspond to the plant components extracted from the plant material or derivatives thereof only.

30

The compositions of the present invention comprise plant extract from plants selected from *Equisetum arvense* (also known as common horsetail, or field horsetail), *Rumex crispus* (also known as curly dock, or yellow dock), *Arctium lappa* (also known as burdock, or greater burdock), *Trifolium pratense* (also known as red clover),
5 *Chelidonium majus* (also known as greater celandine, tetterwort, nipplewort, or swallowwort), *Thuja occidentalis* (also known as northern white-cedar or eastern arborvitae), *Urtica dioica* (also known as nettle), *Symphytum officinale* (also known as comfrey, in particular, common comfrey), *Mahonia aquifolium* (also known as Oregon grape), *Echinacea purpurea*, *Stellaria media* (also known as chickweed), *Galium*
10 *aparine* (also known as cleavers), *Aloe vera* (also known as *Aloe barbadensis*), *Matricaria recutita* (also known as chamomile), *Hypericum perforatum* (also known as St John's wort), *Calendula officinalis* (also known as common marigold), and *Panax ginseng*.

15 In some embodiments, the composition may contain other plant extracts in addition to those recited hereinabove. In other embodiments, the composition may contain only the plant extracts recited hereinabove and no other plant extracts.

The composition comprises a combination of extracts from at least 6 plants selected
20 from the group given above. In some embodiments, the composition comprises a combination of extracts from at least 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17 plants selected from the group given above.

Without being bound by theory, it is believed that the combinations of plant extracts
25 with colloidal silver as described herein may provide an alternative or better effect than the corresponding cumulative effect of the plant extracts or colloidal silver used individually.

Given that the plant extract in the composition described herein is invariably an extract
30 from a combination of plants, it may be referred to as 'the plant extract combination'.

Plant extract from a combination of plants can be provided as a single extraction from a combination of plants, or a combination of extracts from individual plants. Preferably, plant extract from a combination of plants is provided as a combination of extracts from individual plants. In embodiments comprising a combination of extracts from
5 individual plants, the extracts from individual plants may be referred to as components of the plant extract combination.

The composition may comprise plant extract in any suitable amount. For example, the composition may comprise plant extract in an amount of from 0.1% to 10% (w/v), or
10 0.25% to 5% (w/v), or 0.5% to 2.5% (w/v), or 0.75% to 2% (w/v). The plant extract may be present in the composition in an amount less than or equal to 10% (w/v), 5% (w/v), 2.5% (w/v), 2% (w/v), or 1.5% (w/v). The plant extract may be present in the composition in an amount greater than or equal to 0.1% (w/v), or 0.25% (w/v), or 0.5% (w/v), or 0.75% (w/v), or 1% (w/v).

15

In embodiments wherein the plant extract is provided as a combination of extracts from individual plants, each component of the plant extract may be present in the composition any suitable amount. For example, each component of the plant extract may be present in the composition in an amount of from 0.05% to 10% (w/v), or 0.1%
20 to 5% (w/v), or 0.25% to 2.5% (w/v), or 0.5% to 2% (w/v), or 0.6% (w/v) to 1% (w/v). Each component of the plant extract may be present in the composition in an amount less than or equal to 10% (w/v), 5% (w/v), 2.5% (w/v), 2% (w/v), or 1.5% (w/v), or 1% (w/v). Each component of the plant extract may be present in the composition in an amount greater than or equal to 0.05% (w/v), or 0.1% (w/v), or 0.25% (w/v), or 0.5%
25 (w/v), or 0.6% (w/v), or 0.75% (w/v).

In embodiments wherein the plant extract is provided as an extractant-free plant extract, each component of the extractant-free plant extract may be present in the composition in an amount of from 0.01% to 1% (w/v), or 0.025% to 0.5% (w/v), or 0.05% to 0.25%
30 (w/v), or 0.075% to 0.2% (w/v). The plant extract may be present in the composition in an amount less than or equal to 1% (w/v), 0.5% (w/v), 0.25% (w/v), 0.2% (w/v), or

0.15% (w/v). The plant extract may be present in the composition in an amount greater than or equal to 0.01% (w/v), or 0.025% (w/v), or 0.05% (w/v), or 0.075% (w/v), or 0.1% (w/v).

- 5 Plant extracts from individual plants may be combined in any ratio relative to each other to provide the plant extract combination. In some embodiments, the ratio between the component of the plant extract combination present in the largest amount and the component of the plant extract combination present in the smallest amount may be less than 10:1 (w/w), less than 8:1 (w/w), less than 6:1 (w/w), or less than 4:1 (w/w).

10

Use of composition

- The compositions of the present invention may be suitable for use with humans. For example, the compositions may be suitable for application to the skin and/or mucous membrane of a human body. In other embodiments, the compositions may be suitable for use with non-human animals. For example, the composition may be suitable for application to the skin, mucous membrane, hair and/or coat of animals. The composition may be suitable for application to the skin, mucous membrane, hair and/or coat of non-human animals such as fish, amphibians, reptiles, birds, and non-human mammals. In particular, the composition may be suitable for application to the skin, mucous membrane, hair and/or coat of household pets such as cats, dogs, rabbits, lizards, snakes, parrots, budgerigars and the like. Alternatively or additionally, the composition may be particularly suitable for application to the skin, mucous membrane, hair and/or coat of farmyard animals such as cows, pigs, sheep, horses, goats, donkeys and the like.

25

- The compositions of the present invention may be used as medicaments. The compositions may be used as medicaments for a range of indications, for example dermatitis, psoriasis, wounds, tinea pedis (also known as Athlete's foot), blisters, insect bites, and inflammation. It may be that one combination of plant extracts is not optimal for treating all of the given indications. Rather, it may be that particular plant extract combinations are particularly suitable for use as medicaments for particular indications.

30

Particular combinations and the indications they may be particularly suitable for are described herein.

5 Dermatititis may also be referred to as eczema. Dermatititis includes atopic dermatitis, contact dermatitis, xerotic dermatitis, and seborrhoeic dermatitis. Compositions of the present invention may particularly be used as a medicament for contact dermatitis including irritant diaper dermatitis (also known as nappy rash), and contact dermatitis deriving related to incontinence.

10 Wounds include open wounds such as incisions, lacerations, abrasions, avulsions, and puncture wounds. Open wounds may also be referred to as severed skin, or cuts. Wounds also include closed wounds such as hematomas. Hematomas include contusions (also referred to as bruises), petechiae, purpura and ecchymoses. Compositions of the present invention may particularly be used as a medicament for
15 open wounds, and/or contusions. Some embodiments may increase the rate of healing of such wounds.

Burns include thermal burns, chemical burns and radiation burns. Compositions of the present invention may particularly be used as a medicament for radiation burns such as
20 sunburn. Alternatively or additionally, compositions of the present invention may particularly be used as a medicament for thermal burns such as oil burns, iron burns, oven burns, and steam burns. Some embodiments may increase the rate of healing of such wounds.

25 The present invention also relates to a method of treating a dermatological condition. Said method comprises providing a composition as described herein, and applying the composition to the affected area. For example, the method may comprise providing a composition for use as a medicament for dermatitis, and applying said composition to a portion of skin presenting with dermatitis. Alternatively, the method may comprise
30 providing a composition for use as a medicament for psoriasis, and applying said composition to a portion of skin presenting with psoriasis. Alternatively, the method

may comprise providing a composition for use as a medicament for wounds, and applying said composition to a wound. Alternatively, the method may comprise providing a composition for use as a medicament for tinea pedis, and applying said composition to a portion of skin presenting with tinea pedis. Alternatively, the method
5 may comprise providing a composition for use as a medicament for blisters, and applying said composition to at least part of the surface of the blister. Alternatively, the method may comprise providing a composition for use as a medicament for insect bites, and applying said composition to the insect bite. Alternatively, the method may comprise providing a composition for use as a medicament for inflammation, and
10 applying said composition to a portion of inflamed skin.

The composition of the present invention may alternatively or additionally be a cosmetic composition. For instance, compositions may be applied to the skin to reduce the size and/or pigmentation of nevi (also known as moles), liver spots, areas of
15 hyperpigmentation, reddened skin, and rashes.

The present invention also relates to non-therapeutic cosmetic methods. Said method comprises providing a composition as described herein, selecting an area of the body to which the composition should be applied, and topically applying the composition to the
20 selected area. For example, the method may comprise providing a composition, selecting a nevi, and applying said composition to at least part of the surface of the nevi. Alternatively, the method may comprise providing a composition, selecting a liver spot, and applying said composition to at least part of the surface of the liver spot. Alternatively, the method may comprise providing a composition, selecting an area of
25 hyperpigmentation, and applying said composition to at least part of the surface of the area of hyperpigmentation. Alternatively, the method may comprise providing a composition, selecting an area of reddened skin, and applying said composition to at least part of the surface of the reddened skin. Alternatively, the method may comprise providing a composition, selecting a rash, and applying said composition to at least part
30 of the surface of the rash.

Compositions may also be used for providing the user with a sensation at the skin and/or mucous membrane. For example, application of the composition to the skin and/or mucous membrane may result in the user perceiving a cooling and/or soothing sensation.

5

The composition of the present invention may alternatively or additionally be used on the skin and/or hair of household pets such as cats, dogs etc.

The compositions of the present invention may be provided to the user in any form. For example, the composition may be provided alone, or in combination with a form of application. For example, in one embodiment, there is provided a dressing comprising a composition as described herein. In another embodiment, there is provided a porous sheet impregnated with a composition as described herein. Such an impregnated porous sheet may be referred to as a “wet wipe”.

15

Specific combinations of components

In a particular embodiment, the composition comprises water in an amount of from 80 to 99.9% (v/v). The composition also comprises elemental silver as a colloidal suspension in an amount of from 0.1 ppm to 100 ppm (that is, 1×10^{-5} to 1×10^{-2} % (w/v)). The composition also comprises plant extract from at least 6 plants selected from the group consisting of *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. The plant extract is present in the composition in an amount of from 0.1% to 10% (w/v).

25

In a further embodiment, the composition consists of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

30

In one embodiment, the plant extract may comprise or consist of extracts from *Equisetum arvense*, *Symphytum officinale*, *Mahonia aquifolium*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be particularly suitable for use as a medicament for blisters.

5

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Equisetum arvense*, *Symphytum officinale*, *Mahonia aquifolium*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

10

In one embodiment, the plant extract may comprise or consist of extracts from *Arctium lappa*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*. This embodiment may be particularly suitable for use as a cosmetic composition to be applied to skin blemishes.

15

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Arctium lappa*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

20

In one embodiment, the plant extract may comprise or consist of extracts from *Symphytum officinale*, *Echinacea purpurea*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be particularly suitable for use as a medicament for burns.

25

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Symphytum officinale*, *Echinacea purpurea*, *Aloe vera*, *Matricaria recutita*,

30

Hypericum perforatum, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

5 In one embodiment, the plant extract may comprise or consist of extracts from *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be particularly suitable for use as a medicament for sunburn.

10 In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

15 In one embodiment, the plant extract may comprise extracts from *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be
20 particularly suitable for use as a medicament for eczema.

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*,
25 *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be particularly suitable for use as a medicament for open wounds.

In a further embodiment, the composition may consist of elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ (w/v), plant extract comprising or consisting of extracts from *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Chelidonium majus*, *Thuja occidentalis*, *Echinacea purpurea*, *Galium aparine*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*. This embodiment may be particularly suitable for use as a medicament for tinea pedis.

In a further embodiment, the composition may consist of elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ (w/v), plant extract comprising or consisting of extracts from *Chelidonium majus*, *Thuja occidentalis*, *Echinacea purpurea*, *Galium aparine*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Chelidonium majus*, *Thuja occidentalis*, *Mahonia aquifolium*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*. This embodiment may be particularly suitable for use as a cosmetic composition to be applied to nevi.

In a further embodiment, the composition may consist of elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ (w/v), plant extract comprising or consisting of extracts

from *Chelidonium majus*, *Thuja occidentalis*, *Mahonia aquifolium*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

5 In one embodiment, the plant extract may comprise or consist of extracts from *Arctium lappa*, *Urtica dioica*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*. This embodiment may be particularly suitable for use as a medicament for insect bites.

10 In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Arctium lappa*, *Urtica dioica*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

15 In one embodiment, the plant extract may comprise or consist of extracts from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*. This embodiment may be particularly suitable for use as a medicament for irritant diaper dermatitis.

20 In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

25 In one embodiment, the plant extract may comprise or consist of extracts from *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*,
30 and *Panax ginseng*. This embodiment may be particularly suitable for use as a medicament for contact dermatitis deriving from incontinence.

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*,
5 *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. This embodiment may be particularly suitable for application to the skin and hair of household pets.

15 In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in an amount of from 0.1% to 10% (w/v), and water as the remainder
20 of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*. This embodiment may be particularly
25 suitable for including in a porous sheet (also known as a wet wipe).

In a further embodiment, the composition may consist of elemental silver in an amount of from 1×10^{-5} % to 1×10^{-2} % (w/v), plant extract comprising or consisting of extracts from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis* in an amount of from 0.1% to 10%
30 (w/v), and water as the remainder of the composition up to 100%.

In one embodiment, the plant extract may comprise or consist of extracts from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, and *Calendula officinalis*. This
5 embodiment may be particularly suitable for including in a porous sheet (also known as a wet wipe), for application to the skin and hair of household pets.

In a further embodiment, the composition may consist of elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ (w/v), plant extract comprising or consisting of extracts
10 from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, and *Calendula officinalis* in an amount of from 0.1% to 10% (w/v), and water as the remainder of the composition up to 100%.

15 **Methods of preparing compositions**

Compositions according to the present invention may be manufactured according to any suitable method. One appropriate method is to provide plant extract from at least 6 plants selected from the group consisting of *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica*
20 *dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*. The method further comprises providing a colloidal suspension of silver in a base, the silver content of the colloidal suspension being of from 0.1 ppm to 100 ppm (that is, $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ (w/v) of the base).
25 The plant extract and colloidal suspension are then combined to provide a composition for topical application.

The base may be any of those described herein. In one embodiment, the base is water.

30 The plant extract and colloidal suspension of silver may be combined in any appropriate ratio. In one embodiment, they are combined in a ratio of from 1:20 to 1:4 (w/v) plant

extract to colloidal suspension of silver, or from 1:15 to 1:6, or from 1:12 to 1:8. In one embodiment, they are combined in a ratio of around 1:9 (w/v) plant extract to colloidal suspension of silver.

5 Examples

Examples 1 to 14 provide topically applicable compositions comprising plant extracts prepared by extracting plant material with aqueous ethanol in a ratio of 1:1 weight/volume plant material to aqueous ethanol. The proportion of ethanol to water in the aqueous ethanol (i.e. the ethanol content) for each extract is as follows:

Table 1

Plant material	% ethanol to water (v/v)
<i>Equisetum arvense</i>	25%
<i>Rumex crispus</i>	45%
<i>Arctium lappa</i>	45%
<i>Trifolium pratense</i>	45%
<i>Chelidonium majus</i>	45%
<i>Thuja occidentalis</i>	45%
<i>Urtica dioica</i>	45%
<i>Symphytum officinale</i>	45%
<i>Mahonia aquifolium</i>	45%
<i>Echinacea purpurea</i>	45%
<i>Stellaria media</i>	45%
<i>Galium aparine</i>	45%
<i>Aloe vera</i>	60%
<i>Matricaria recutita</i>	45%
<i>Hypericum perforatum</i>	60%
<i>Calendula officinalis</i>	90%
<i>Panax ginseng</i>	45%

Example 1

This is an example of a composition for applying to blisters.

- 5 Aqueous extracts of *Equisetum arvense* (0.20 mL), *Symphytum officinale* (0.40 mL), *Mahonia aquifolium* (0.20 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.12 mL), *Calendula officinalis* (0.20 mL), and *Panax ginseng* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable
- 10 composition.

Example 2

This is an example of a composition for applying to skin blemishes.

Aqueous extracts of *Arctium lappa* (0.20 mL), *Echinacea purpurea* (0.40 mL), *Stellaria media* (0.20 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.12 mL), and *Calendula officinalis* (0.12 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 3

10 This is an example of a composition for applying to skin burns.

Aqueous extracts of *Symphytum officinale* (0.28 mL), *Echinacea purpurea* (0.12 mL), *Aloe vera* (0.20 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.20 mL), *Calendula officinalis* (0.20 mL), and *Panax ginseng* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 4

This is an example of a composition for applying to eczema.

Aqueous extracts of *Equisetum arvense* (0.20 mL), *Rumex crispus* (0.20 mL), *Arctium lappa* (0.08 mL), *Trifolium pratense* (0.28 mL), *Chelidonium majus* (0.08 mL), *Thuja occidentalis* (0.08 mL), *Urtica dioica* (0.12 mL), *Symphytum officinale* (0.20 mL), *Mahonia aquifolium* (0.20 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.28 mL), *Galium aparine* (0.28 mL), *Aloe vera* (0.20 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.12 mL), *Calendula officinalis* (0.12 mL), and *Panax ginseng* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 5

30 This is an example of a composition for applying to cuts.

Aqueous extracts of *Equisetum arvense* (0.20 mL), *Symphytum officinale* (0.28 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.12 mL), *Aloe vera* (0.20 mL), *Matricaria recutita* (0.12 mL), *Hypericum perforatum* (0.12 mL), *Calendula officinalis* (0.28 mL), and *Panax ginseng* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 6

This is an example of a composition for applying to Tinea pedis (Athlete's Foot).

Aqueous extracts of *Chelidonium majus* (0.04 mL), *Thuja occidentalis* (0.20 mL), *Echinacea purpurea* (0.12 mL), *Galium aparine* (0.40 mL), *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.12 mL), and *Calendula officinalis* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 7

This is an example of a composition for applying to sunburn.

Aqueous extracts of *Galium aparine* (0.20 mL), *Aloe vera* (0.28 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.28 mL), *Calendula officinalis* (0.28 mL), and *Panax ginseng* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 8

This is an example of a composition for applying to moles.

Aqueous extracts of *Chelidonium majus* (0.12 mL), *Thuja occidentalis* (0.28 mL), *Mahonia aquifolium* (0.20 mL), *Aloe vera* (0.30 mL), *Matricaria recutita* (0.20 mL), and *Calendula officinalis* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 9

This is an example of a composition for applying to insect bites.

Aqueous extracts of *Arctium lappa* (0.20 mL), *Urtica dioica* (0.28 mL), *Echinacea purpurea* (0.28 mL), *Stellaria media* (0.20 mL), *Matricaria recutita* (0.12 mL), *Hypericum perforatum* (0.20 mL), and *Calendula officinalis* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 10

This is an example of a composition for applying to irritant diaper dermatitis.

Aqueous extracts of *Urtica dioica* (0.20 mL), *Symphytum officinale* (0.28 mL), *Echinacea purpurea* (0.28 mL), *Stellaria media* (0.20 mL), *Aloe vera* (0.28 mL), *Matricaria recutita* (0.28 mL), and *Calendula officinalis* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 11

This is an example of a composition for use in the treatment of contact dermatitis related to incontinence.

Aqueous extracts of *Urtica dioica* (0.20 mL), *Symphytum officinale* (0.28 mL), *Echinacea purpurea* (0.20 mL), *Mahonia aquifolium* (0.12 mL), *Stellaria media* (0.20 mL), *Aloe vera* (0.28 mL), *Matricaria recutita* (0.28 mL), *Hypericum perforatum* (0.20 mL), *Calendula officinalis* (0.28 mL), and *Panax ginseng* (0.28 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 12

This is an example of a composition for applying to the skin of household pets.

Aqueous extracts of *Rumex crispus* (0.20 mL), *Arctium lappa* (0.20 mL), *Trifolium pratense* (0.20 mL), *Symphytum officinale* (0.12 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.12 mL), *Hypericum perforatum* (0.12 mL), *Calendula officinalis* (0.12 mL), and *Panax ginseng* (0.12 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 13

This is an example of a composition suitable to be provided in combination with a form of application, such as a porous sheet impregnated with the composition.

Aqueous extracts of *Urtica dioica* (0.20 mL), *Symphytum officinale* (0.28 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.20 mL), *Aloe vera* (0.28 mL), *Matricaria recutita* (0.28 mL), and *Calendula officinalis* (0.28 mL), were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet impregnated with the composition (a “wet wipe”).

Example 14

This is an example of a composition suitable to be provided in combination with a form of application, such as a porous sheet impregnated with the composition, wherein the composition is suitable for use with household pets.

Aqueous extracts of *Rumex crispus* (0.20 mL), *Arctium lappa* (0.20 mL), *Trifolium pratense* (0.20 mL), *Symphytum officinale* (0.12 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.12 mL), *Matricaria recutita* (0.12 mL), and *Calendula officinalis* (0.12 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet impregnated with the composition (a “wet wipe”).

Examples 15 to 28 provide topically applicable compositions comprising mother tinctures prepared by extracting plant material with aqueous ethanol in a ratio of 1:9 weight/volume plant material to aqueous ethanol. The proportion of ethanol to water in the aqueous ethanol (i.e. the ethanol content) for each extract is as follows:

5

Table 2

Plant material	% ethanol to water (v/v)
<i>Equisetum arvense</i>	52%
<i>Rumex crispus</i>	49%
<i>Arctium lappa</i>	53%
<i>Trifolium pratense</i>	63%
<i>Chelidonium majus</i>	62%
<i>Thuja occidentalis</i>	64%
<i>Urtica dioica</i>	49%
<i>Symphytum officinale</i>	41%
<i>Mahonia aquifolium</i>	69%
<i>Echinacea purpurea</i>	64%
<i>Stellaria media</i>	50%
<i>Galium aparine</i>	49%
<i>Aloe vera</i>	0%
<i>Matricaria recutita</i>	65%
<i>Hypericum perforatum</i>	65%
<i>Calendula officinalis</i>	64%
<i>Panax ginseng</i>	89%

Example 15

10 This is an example of a composition for applying to blisters.

Aqueous ethanol extracts of *Equisetum arvense* (0.32 mL), *Symphytum officinale* (0.40 mL), *Mahonia aquifolium* (0.32 mL), *Matricaria recutita* (0.32 mL), *Hypericum*

perforatum (0.20 mL), *Calendula officinalis* (0.32 mL), and *Panax ginseng* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

5

Example 16

This is an example of a composition for applying to skin blemishes.

Aqueous ethanol extracts of *Arctium lappa* (0.32 mL), *Echinacea purpurea* (0.40 mL), *Stellaria media* (0.32 mL), *Matricaria recutita* (0.32 mL), *Hypericum perforatum* (0.20 mL), and *Calendula officinalis* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

10

Example 17

This is an example of a composition for applying to skin burns.

Aqueous ethanol extracts of *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.20 mL), *Aloe vera* (0.32 mL), *Matricaria recutita* (0.32 mL), *Hypericum perforatum* (0.32 mL), *Calendula officinalis* (0.32 mL), and *Panax ginseng* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

20

Example 18

This is an example of a composition for applying to eczema.

Aqueous ethanol extracts of *Equisetum arvense* (0.32 mL), *Rumex crispus* (0.32 mL), *Arctium lappa* (0.08 mL), *Trifolium pratense* (0.40 mL), *Chelidonium majus* (0.08 mL), *Thuja occidentalis* (0.08 mL), *Urtica dioica* (0.20 mL), *Symphytum officinale* (0.32 mL), *Mahonia aquifolium* (0.32 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media* (0.40 mL), *Galium aparine* (0.40 mL), *Aloe vera* (0.32 mL), *Matricaria recutita* (0.32 mL), *Hypericum perforatum* (0.20 mL), *Calendula officinalis* (0.20 mL), and *Panax ginseng* (0.32 mL) were combined in a vessel. The extract combination was

25

30

then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 19

5 This is an example of a composition for applying to cuts.

Aqueous ethanol extracts of *Equisetum arvense* (0.32 mL), *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.32 mL), *Stellaria media* (0.20 mL), *Aloe vera* (0.32 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.20 mL), *Calendula officinalis* (0.40 mL), and *Panax ginseng* (0.40 mL) were combined in a
10 vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 20

This is an example of a composition for applying to Tinea pedis (Athlete's Foot).

15 Aqueous ethanol extracts of *Chelidonium majus* (0.12 mL), *Thuja occidentalis* (0.32 mL), *Echinacea purpurea* (0.20 mL), *Galium aparine* (0.40 mL), *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.20 mL), and *Calendula officinalis* (0.40 mL), were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically
20 applicable composition.

Example 21

This is an example of a composition for applying to sunburn.

Aqueous ethanol extracts of *Galium aparine* (0.32 mL), *Aloe vera* (0.40 mL),
25 *Matricaria recutita* (0.32 mL), *Hypericum perforatum* (0.40 mL), *Calendula officinalis* (0.40 mL), and *Panax ginseng* (0.32 mL), were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

30

Example 22

This is an example of a composition for applying to moles.

Aqueous ethanol extracts of *Chelidonium majus* (0.20 mL), *Thuja occidentalis* (0.40 mL), *Mahonia aquifolium* (0.20 mL), *Aloe vera* (0.20 mL), *Matricaria recutita* (0.20 mL), and *Calendula officinalis* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 23

This is an example of a composition for applying to insect bites.

Aqueous ethanol extracts of *Arctium lappa* (0.32 mL), *Urtica dioica* (0.40 mL), *Echinacea purpurea* (0.40 mL), *Stellaria media* (0.32 mL), *Matricaria recutita* (0.20 mL), *Hypericum perforatum* (0.32 mL), and *Calendula officinalis* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

15

Example 24

This is an example of a composition for applying to irritant diaper dermatitis.

Aqueous ethanol extracts of *Urtica dioica* (0.32 mL), *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.32 mL), *Stellaria media* (0.32 mL), *Aloe vera* (0.40 mL), *Matricaria recutita* (0.40 mL), and *Calendula officinalis* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

20

Example 25

This is an example of a composition for use in the treatment of contact dermatitis related to incontinence.

Aqueous ethanol extracts of *Urtica dioica* (0.32 mL), *Symphytum officinale* (0.40 mL), *Mahonia aquifolium* (0.20 mL), *Echinacea purpurea* (0.40 mL), *Stellaria media* (0.32 mL), *Aloe vera* (0.32 mL), *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.32 mL), *Calendula officinalis* (0.32 mL), and *Panax ginseng* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the

30

addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 26

- 5 This is an example of a composition for applying to the skin of household pets. Aqueous ethanol extracts of *Rumex crispus* (0.32 mL), *Arctium lappa* (0.32 mL), *Trifolium pratense* (0.32 mL), *Symphytum officinale* (0.20 mL), *Echinacea purpurea* (0.32 mL), *Stellaria media* (0.20 mL), *Hypericum perforatum* (0.20 mL), *Calendula officinalis* (0.20 mL), and *Panax ginseng* (0.20 mL) were combined in a vessel. The
10 extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 27

- This is an example of a composition suitable to be provided in combination with a form
15 of application, such as a porous sheet impregnated with the composition. Aqueous ethanol extracts of *Urtica dioica* (0.20 mL), *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.32 mL), *Stellaria media* (0.32 mL), *Aloe vera* (0.40 mL), *Matricaria recutita* (0.40 mL), and *Calendula officinalis* (0.40 mL), were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5
20 ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet impregnated with the composition (a “wet wipe”).

Example 28

- 25 This is an example of a composition suitable to be provided in combination with a form of application, such as a porous sheet impregnated with the composition, wherein the composition is suitable for use with household pets. *Rumex crispus* (0.32 mL), *Arctium lappa* (0.32 mL), *Trifolium pratense* (0.32 mL), *Symphytum officinale* (0.20 mL), *Echinacea purpurea* (0.32 mL), *Stellaria media*
30 (0.20 mL), *Matricaria recutita* (0.20 mL), and *Calendula officinalis* (0.20 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the

addition of 5 ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet impregnated with the composition (a “wet wipe”).

- 5 Examples 29 to 42 provide topically applicable compositions comprising plant tinctures prepared by extracting plant material with aqueous ethanol in a ratio of 1:3 weight/volume plant material to aqueous ethanol. The proportion of ethanol to water in the aqueous ethanol (i.e. the ethanol content) for each extract is that as given in Table 1 hereinabove.

10 Example 29

This is an example of a composition for applying to blisters.

- Aqueous extracts of *Equisetum arvense* (0.60 mL), *Symphytum officinale* (0.40 mL), *Mahonia aquifolium* (0.60 mL), *Matricaria recutita* (0.60 mL), *Hypericum perforatum* (0.40 mL), *Calendula officinalis* (0.60 mL), and *Panax ginseng* (0.20 mL) were
15 combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 30

- 20 This is an example of a composition for applying to skin blemishes.

- Aqueous extracts of *Arctium lappa* (0.60 mL), *Echinacea purpurea* (0.40 mL), *Stellaria media* (0.60 mL), *Matricaria recutita* (0.60 mL), *Hypericum perforatum* (0.40 mL), and *Calendula officinalis* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal
25 silver to provide a topically applicable composition.

Example 31

This is an example of a composition for applying to skin burns.

- Aqueous extracts of *Symphytum officinale* (0.80 mL), *Echinacea purpurea* (0.40 mL),
30 *Aloe vera* (0.60 mL), *Matricaria recutita* (0.60 mL), *Hypericum perforatum* (0.60 mL), *Calendula officinalis* (0.60 mL), and *Panax ginseng* (0.80 mL) were combined in a

vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 32

5 This is an example of a composition for applying to eczema.

Aqueous extracts of *Equisetum arvense* (0.60 mL), *Rumex crispus* (0.60 mL), *Arctium lappa* (0.04 mL), *Trifolium pratense* (0.80 mL), *Chelidonium majus* (0.04 mL), *Thuja occidentalis* (0.04 mL), *Urtica dioica* (0.40 mL), *Symphytum officinale* (0.60 mL), *Mahonia aquifolium* (0.60 mL), *Echinacea purpurea* (0.20 mL), *Stellaria media*
10 (0.80 mL), *Galium aparine* (0.80 mL), *Aloe vera* (0.60 mL), *Matricaria recutita* (0.60 mL), *Hypericum perforatum* (0.40 mL), *Calendula officinalis* (0.40 mL), and *Panax ginseng* (0.60 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

15

Example 33

This is an example of a composition for applying to cuts.

Aqueous extracts of *Equisetum arvense* (0.60 mL), *Symphytum officinale* (0.80 mL), *Echinacea purpurea* (0.60 mL), *Stellaria media* (0.40 mL), *Aloe vera* (0.60 mL),
20 *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.40 mL), *Calendula officinalis* (0.80 mL), and *Panax ginseng* (0.80 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

25

Example 34

This is an example of a composition for applying to Tinea pedis (Athlete's Foot).

Aqueous extracts of *Chelidonium majus* (0.20 mL), *Thuja occidentalis* (0.60 mL), *Echinacea purpurea* (0.40 mL), *Galium aparine* (0.40 mL), *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.40 mL), and *Calendula officinalis* (0.80 mL)
30 were combined in a vessel. The extract combination was then made up to 100 mL by

the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 35

- 5 This is an example of a composition for applying to sunburn.
Aqueous extracts of *Galium aparine* (0.60 mL), *Aloe vera* (0.80 mL), *Matricaria recutita* (0.60 mL), *Hypericum perforatum* (0.80 mL), *Calendula officinalis* (0.80 mL), and *Panax ginseng* (0.60 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a
10 topically applicable composition.

Example 36

- This is an example of a composition for applying to moles.
Aqueous extracts of *Chelidonium majus* (0.40 mL), *Thuja occidentalis* (0.80 mL),
15 *Mahonia aquifolium* (0.20 mL), *Aloe vera* (0.20 mL), *Matricaria recutita* (0.20 mL), and *Calendula officinalis* (0.80 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 37

- 20 This is an example of a composition for applying to insect bites.
Aqueous extracts of *Arctium lappa* (0.60 mL), *Urtica dioica* (0.80 mL), *Echinacea purpurea* (0.80 mL), *Stellaria media* (0.60 mL), *Matricaria recutita* (0.40 mL), *Hypericum perforatum* (0.60 mL), and *Calendula officinalis* (0.60 mL) were combined
25 in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

Example 38

- This is an example of a composition for applying to Irritant diaper dermatitis.
30 Aqueous extracts of *Urtica dioica* (0.60 mL), *Symphytum officinale* (0.80 mL), *Echinacea purpurea* (0.60 mL), *Stellaria media* (0.60 mL), *Aloe vera* (0.80 mL),

Matricaria recutita (0.80 mL), and *Calendula officinalis* (0.80 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

5

Example 39

This is an example of a composition for use in the treatment of contact dermatitis related to incontinence.

Aqueous extracts of *Urtica dioica* (0.60 mL), *Symphytum officinale* (0.80 mL), *Mahonia aquifolium* (0.40 mL), *Echinacea purpurea* (0.80 mL), *Stellaria media* (0.60 mL), *Aloe vera* (0.60 mL), *Matricaria recutita* (0.80 mL), *Hypericum perforatum* (0.60 mL), *Calendula officinalis* (0.60 mL), and *Panax ginseng* (0.80 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

15

Example 40

This is an example of a composition for applying to the skin of household pets.

Aqueous extracts of *Rumex crispus* (0.60 mL), *Arctium lappa* (0.60 mL), *Trifolium pratense* (0.60 mL), *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.60 mL), *Stellaria media* (0.40 mL), *Hypericum perforatum* (0.40 mL), *Calendula officinalis* (0.40 mL), and *Panax ginseng* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a topically applicable composition.

25

Example 41

This is an example of a composition suitable to be provided in combination with a form of application, such as a porous sheet impregnated with the composition.

Aqueous extracts of *Urtica dioica* (0.60 mL), *Symphytum officinale* (0.80 mL), *Echinacea purpurea* (0.60 mL), *Stellaria media* (0.60 mL), *Aloe vera* (0.80 mL), *Matricaria recutita* (0.80 mL), and *Calendula officinalis* (0.80 mL), were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5

30

ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet impregnated with the composition (a “wet wipe”).

5

Example 42

This is an example of a composition suitable to be provided in combination with a form of application, such as a porous sheet impregnated with the composition, wherein the composition is suitable for use with household pets.

Aqueous extracts of *Rumex crispus* (0.60 mL), *Arctium lappa* (0.60 mL), *Trifolium pratense* (0.60 mL), *Symphytum officinale* (0.40 mL), *Echinacea purpurea* (0.60 mL),
10 *Stellaria media* (0.40 mL), *Matricaria recutita* (0.40 mL), and *Calendula officinalis* (0.40 mL) were combined in a vessel. The extract combination was then made up to 100 mL by the addition of 5 ppm aqueous colloidal silver to provide a composition. A porous sheet was treated with the resulting composition to provide a porous sheet
15 impregnated with the composition (a “wet wipe”).

Description of embodiments with reference to the Figures

Figure 1 shows a subject 100 with an area of unbroken skin 110. A localised area of skin 120 presents with an open wound 122. This picture of subject 100 was taken at
20 15:00 on Day 0 of a test period, prior to application of any composition described herein.

A composition consisting of aqueous extracts of *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*,
25 *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in a ratio of 3:4:3:2:3:2:4:4 (w/w) in 5 ppm aqueous colloidal silver was prepared (corresponding to Example 33). The area of skin 120 containing the wound was selected, and the composition topically applied to the selected area. Said composition was regularly applied to the area 120 through a test period of 8 days.

30

Figure 2 is a picture of subject 100 taken at 10:22 on Day 3 of the test period. The open wound 102 in area 120 has scabbed to provide a scabbed wound 124.

Figure 3 is a picture of the subject 100 taken at 16:26 on Day 6 of the test period, showing a decrease in the size of scabbed wound 124 in area 120 to provide scabbed wound 126.

Figure 4 is a picture of the subject 100 representing the appearance of Subject 100 on Day 8 of the test period. Scabbed wound 124 in area 120 has healed completely to provide new skin 128. New skin 128 in the area of skin 120 on Day 8 resembles the unbroken 110. Thus, Figures 1 to 4 demonstrate that regular application of a composition of the present invention to an open wound may result in an increased rate of healing of the wound.

In another example, a subject 200 presented with a nevi (mole) in a localised area of skin 220. The nevi was roughly circular in shape, had a darker pigmentation compared with the surrounding normal skin tone, and protruded from the surface of the skin. A composition consisting of aqueous extracts of *Chelidonium majus*, *Thuja occidentalis*, *Mahonia aquifolium*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis* in a ratio of 2:4:1:1:1:4 (w/w) in 5 ppm aqueous colloidal silver was prepared (corresponding to Example 36). The area of skin 220 containing the nevi was selected, and the composition topically applied to the selected area. Said composition was regularly applied to the selected area through a test period of 95 days.

Figure 5 is a picture of subject 200 taken on Day 80 of the test period. Subject 200 has an area of normal skin tone 210. The localised area of the skin 220 presents with a portion of darker pigmentation 222 compared to the normal skin tone 210. The darker pigmentation 222 also protrudes from the surrounding normal skin 210. The size of the nevi, the pigmentation of the area 220, and the protrusion of the nevi from the skin are reduced compared with the appearance of the nevi prior to application of the

composition. In particular, localised area of skin 220 now contains normal skin tone 212, whereas on Day 0 this area was part of the nevi.

Figure 6 is a picture representing the appearance of Subject 200 on Day 95 of the test period. The pigmentation of the skin 222 has substantially decreased; the pigmentation of the skin 222 in the area of the body 220 now resembles the normal skin tone 210, and area of skin 220 now contains only skin of normal skin tone 214. Further, there is now no protrusion from the body. Thus, Figures 5 and 6 demonstrate that regular application of a composition of the present invention to a nevi may result in a reduction in the pigmentation and physical protrusion of the nevi.

Figure 7 shows a subject 300 with a normal skin tone 310. Localised areas of skin 320 and 330 present with recently inflicted mosquito bites, resulting in reddened and raised skin 322 and 332. This picture of subject 300 was taken prior to topical application of any composition described herein.

A composition consisting of aqueous extracts of *Arctium lappa*, *Urtica dioica*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*, in a ratio of 3:4:4:3:2:3:3 (w/w) in 5 ppm aqueous colloidal silver was prepared (corresponding to Example 37). The areas of skin 320 and 330 presenting with insect bites were selected, and the composition topically applied to the selected areas. Said composition was regularly applied (three times-daily) to the areas 320 and 330 through a test period of 24 hours.

Figure 8 is a picture of the subject 300 taken after 6 hours of the test period. Reddened and raised skin 324 and 334 is decreased in size compared with reddened and raised skin 322 and 332. Reddened and raised skin 324 and 334 is of a skin tone more similar to normal skin tone 310 than reddened and raised skin 322 and 332.

Figure 9 is a picture of the subject 300 taken after 24 hours of the test period. Reddened and raised skin 326 and 336 is yet further decreased in size compared with reddened

and raised skin 324 and 334. Reddened and raised skin 326 and 336 is of a skin tone even more similar to normal skin tone 310 than reddened and raised skin 324 and 334. Subject 300 also reported a perceived reduction in the itching sensation provided by the mosquito bite after application of the composition.

5

Thus, Figures 7 to 9 demonstrate that regular application of a composition of the present invention to an insect bite, in particular a mosquito bite, may result in a reduction in the size, reddening and itchiness of an insect bite.

10 Figure 10 shows a subject 400 with unbroken skin 410. Localised area of skin 420 contains a recently inflicted open wound. Localised area of skin 420 contains two smaller areas 430 and 440. Area 430 contains a portion of open wound 432; area 440 contains a portion of scabbed wound 442. This picture of subject 400 was taken at 09:30 on Day 0 of a test period prior to topical application of any composition described
15 herein.

A composition consisting of aqueous extracts of *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng* in a ratio of
20 3:4:3:2:3:2:2:4:4 (w/w) in 5 ppm aqueous colloidal silver was prepared (corresponding to Example 33). The area of skin 420 containing the wound was selected, and the composition topically applied to the selected area. Said composition was regularly applied to the areas 420 for the first 5 days of a test period of 10 days.

25 Figure 11 is a picture of the subject 400 taken at 10:12 on Day 3 of the test period. The portion of open wound 432 in area 430 has scabbed to provide portion of scabbed wound 434 in area 430. The portion of scabbed wound 442 in area 440 has reduced in size to provide portion of scabbed wound 444 in area 440. The inflammation surrounding portions of wound 434 and 444 in area 420 is reduced compared with that
30 shown in Figure 10.

Figure 12 is a picture representing the appearance of Subject 400 at 11:17 on Day 10 of the test period. The portion of scabbed wound 434 in area 430 has healed to provide new skin 436 in area 430. The portion of scabbed wound 444 in area 440 has healed to provide new skin 446 in area 440. A wound of this magnitude inflicted on subject 400 would not usually be expected to heal within 10 days. Thus, Figures 10 to 12 demonstrate that regular application of a composition of the present invention to a wound may result in an increased rate of healing of the wound.

The above embodiments are to be understood as illustrative examples of the invention. Further embodiments of the invention are envisaged. It is to be understood that any feature described in relation to any one embodiment may be used alone, or in combination with other features described, and may also be used in combination with one or more features of any other of the embodiments, or any combination of any other of the embodiments. Furthermore, equivalents and modifications not described above may also be employed without departing from the scope of the invention, which is defined in the accompanying claims.

CLAIMS

1. A composition for topical application comprising a base, and dispersed within the base:
 - 5 elemental silver as a colloidal suspension; and
 - plant extract from at least 6 plants selected from the group consisting of *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, *Equisetum arvense*, *Symphytum officinale*, *Panax ginseng*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja*
 - 10 *occidentalis*, *Urtica dioica*, *Mahonia aquifolium*, and *Galium aparine*.
2. A composition according to claim 1, wherein the composition comprises:
 - base in an amount of from 80% to 99.9% by volume/volume of the composition;
 - elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by
 - 15 weight/volume of the composition; and
 - plant extract in an amount of from 0.1% to 10% by weight/volume of the composition.
3. A composition according to claim 1 or claim 2, wherein the composition is an
- 20 aqueous suspension, a cream, an ointment, a spray, a gel, or an aerosol.
4. A composition according to claim 3, wherein the composition is an aqueous suspension, and the base is water.
- 25 5. A composition according to claim 4, wherein the composition consists essentially of:
 - elemental silver in an amount of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by
 - weight/volume of the composition;
 - plant extract in an amount of from 0.1% to 10% by weight/volume of the
 - 30 composition; and
 - water as the remainder of the composition up to 100%.

6. A composition according to any of the preceding claims, wherein the plant extract is an extract in an extractant selected from C₁-C₄ alkanol, acetic acid, glycerine, propylene glycol, honey, water, and mixtures thereof.

5

7. A composition according to claim 6, wherein the extractant consists of aqueous ethanol.

8. A composition according to claim 6 or claim 7, wherein the plant extract is an
10 extract of plant material in a ratio of from 1:1 to 1: 12 weight/volume of plant material to extractant.

9. A composition according to any of claims 1 to 5, wherein the extract is an extractant-free extract.

15

10. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*,
20 *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

11. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Equisetum arvense*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and
25 *Panax ginseng*.

12. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Symphytum officinale*, *Echinacea purpurea*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

30

13. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 5 14. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Chelidonium majus*, *Thuja occidentalis*, *Echinacea purpurea*, *Galium aparine*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 10 15. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Equisetum arvense*, *Symphytum officinale*, *Mahonia aquifolium*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.
- 15 16. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Arctium lappa*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 20 17. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Chelidonium majus*, *Thuja occidentalis*, *Mahonia aquifolium*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*.
- 25 18. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Arctium lappa*, *Urtica dioica*, *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, *Hypericum perforatum*, and *Calendula officinalis*.
- 30 19. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*, *Matricaria recutita*, and *Calendula officinalis*.
20. A composition according to any of claims 1 to 9, wherein the plant extracts are from *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*,

Stellaria media, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*.

21. A composition according to any of claims 1 to 9, wherein the plant extracts are
5 from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*,
Echinacea purpurea, *Stellaria media*, *Hypericum perforatum*, *Calendula officinalis*,
and *Panax ginseng*.

22. A composition according to any of claims 1 to 9, wherein the plant extracts are
10 from *Urtica dioica*, *Symphytum officinale*, *Echinacea purpurea*, *Stellaria media*, *Aloe vera*,
Matricaria recutita, and *Calendula officinalis*.

23. A composition according to any of claims 1 to 9, wherein the plant extracts are
from *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Symphytum officinale*,
15 *Echinacea purpurea*, *Stellaria media*, *Matricaria recutita*, and *Calendula officinalis*.

24. A composition according to any of the preceding claims which is a cosmetic composition.

20 25. A composition according to any of claims 1 to 23 for use as a medicament.

26. A composition according to any of claims 1 to 10 for use in the treatment of dermatitis.

25 27. A composition according to any of claims 1 to 9 or claim 11 for use in the treatment of severed skin.

28. A composition according to any of claims 1 to 9 or claim 12 for use in the treatment of burns.

29. A composition according to any of claims 1 to 9 or claim 13 for use in the treatment of sunburn.
30. A composition according to any of claims 1 to 9 or claim 14 for use in the treatment of tinea pedis.
31. A composition according to any of claims 1 to 9 or claim 15 for use in the treatment of blisters.
32. A composition according to any of claims 1 to 9 or claim 18 for use in the treatment of insect bites.
33. A composition according to any of claims 1 to 9 or claim 19 for use in the treatment of irritant diaper dermatitis.
34. A composition according to any of claims 1 to 9 or claim 20 for use in the treatment of contact dermatitis related to incontinence.
35. A method of manufacturing a composition for topical application, the method comprising:
- providing plant extract from at least 6 plants selected from the group consisting of *Equisetum arvense*, *Rumex crispus*, *Arctium lappa*, *Trifolium pratense*, *Chelidonium majus*, *Thuja occidentalis*, *Urtica dioica*, *Symphytum officinale*, *Mahonia aquifolium*, *Echinacea purpurea*, *Stellaria media*, *Galium aparine*, *Aloe vera*, *Matricaria recutita*, *Hypericum perforatum*, *Calendula officinalis*, and *Panax ginseng*;
- providing a colloidal suspension of elemental silver in a base, the elemental silver content of the colloidal suspension being of from $1 \times 10^{-5} \%$ to $1 \times 10^{-2} \%$ by weight/volume of the base; and
- combining the plant extract and the colloidal suspension of elemental silver to provide a composition for topical application.

36. A method according to claim 35, wherein the base is water.
37. A method according to claim 35 or 36, wherein the plant extract and colloidal suspension of elemental silver are combined in a ratio of from 1:20 to 1:4 weight/volume plant extract to colloidal suspension of elemental silver.
38. A dressing comprising a composition according to any of claims 1 to 34.
39. A porous sheet impregnated with a composition according to any of claims 1 to 34.
40. A method of treating an area affected by a dermatological condition comprising: providing a composition according to any of claims 1 to 34; and topically applying the composition to the area affected by the dermatological condition.
41. A non-therapeutic cosmetic method comprising: providing a composition according to any of claims 1 to 34; selecting an area of the body to which the composition should be applied; and topically applying the composition to the selected area.

1/6




FIG 1




FIG 2

2/6




FIG 3




FIG 4

3/6




FIG 5




FIG 6

4/6




FIG 7




FIG 8

5/6




FIG 9




FIG 10

6/6




FIG 11




FIG 12

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2018/050869

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K33/38 A61K36/00 A61P17/02
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, WPI Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	RU 2 432 146 C1 (POLEZHAEV VIKTOR FEDOROVICH [RU]) 27 October 2011 (2011-10-27) claim 1	1-41
Y	DE 10 2006 049929 A1 (BEIERSDORF AG [DE]) 24 April 2008 (2008-04-24) paragraphs [0069], [0071]	1-41
Y	US 2004/185123 A1 (MAZZIO ELIZABETH A [US] ET AL) 23 September 2004 (2004-09-23) paragraph [0011]	1-41
Y	WO 2016/070194 A1 (POMEGA INC [US]) 6 May 2016 (2016-05-06) paragraph [0034]	1-41
	- / - -	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 June 2018

Date of mailing of the international search report

18/07/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Hars, Jesko

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2018/050869

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2005/120173 A2 (PAKNIKAR KISHORE MADHUKAR [IN]) 22 December 2005 (2005-12-22) page 6 - page 7 -----	1-41
Y	US 6 103 272 A (KEENEY JOSEPH A [US]) 15 August 2000 (2000-08-15) claim 4 -----	1-41
Y	DE 20 2009 014065 U1 (VOCK BRITTA [DE]) 11 February 2010 (2010-02-11) claim 1 -----	1-41
Y	DE 20 2015 006527 U1 (ALTPETER PATRICK [DE]) 4 December 2015 (2015-12-04) claims 1-2 -----	1-41
Y	WO 2005/002608 A1 (HILTERMAN KARINA ANNA [AU]) 13 January 2005 (2005-01-13) page 10 -----	1-41
Y	US 2011/262558 A1 (HUCKFELDT ROGER [US] ET AL) 27 October 2011 (2011-10-27) paragraph [0053] -----	1-41
Y	DATABASE GNPD [Online] MINTEL; May 2015 (2015-05), anonymous: "Balcann Gel", XP002782499, retrieved from www.gnpd.com accession no. 3194263 Database accession no. 3194263 page 1 -----	1-41
Y	DATABASE GNPD [Online] MINTEL; April 2014 (2014-04), anonymous: "Ionic+ Colloidal Silver Healing Cream", XP002782500, retrieved from www.gnpd.com accession no. 2370062 Database accession no. 2370062 page 1 -----	1-41
Y	DATABASE GNPD [Online] MINTEL; September 2012 (2012-09), anonymous: "Ionic+ Colloidal Silver Healing Gel", XP002782501, retrieved from www.gnpd.com accession no. 1880744 Database accession no. 1880744 page 1 -----	1-41

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2018/050869

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
RU 2432146 C1	27-10-2011	-----	
DE 102006049929 A1	24-04-2008	AT 536894 T	15-12-2011
		CN 101594891 A	02-12-2009
		DE 102006049929 A1	24-04-2008
		EP 2104519 A2	30-09-2009
		ES 2378408 T3	12-04-2012
		WO 2008046541 A2	24-04-2008

US 2004185123 A1	23-09-2004	NONE	

WO 2016070194 A1	06-05-2016	CA 2966325 A1	06-05-2016
		EP 3212288 A1	06-09-2017
		JP 2017533960 A	16-11-2017
		US 2017333504 A1	23-11-2017
		WO 2016070194 A1	06-05-2016

WO 2005120173 A2	22-12-2005	AU 2005251570 A1	22-12-2005
		CN 1953664 A	25-04-2007
		EA 200602094 A1	27-04-2007
		EP 1753293 A2	21-02-2007
		US 2007218555 A1	20-09-2007
		WO 2005120173 A2	22-12-2005
		ZA 200608552 B	28-05-2008

US 6103272 A	15-08-2000	NONE	

DE 202009014065 U1	11-02-2010	NONE	

DE 202015006527 U1	04-12-2015	NONE	

WO 2005002608 A1	13-01-2005	AU 2004253192 A1	13-01-2005
		BR PI0418854 A	20-11-2007
		CA 2572578 A1	13-01-2005
		CN 101022819 A	22-08-2007
		EP 1781376 A1	09-05-2007
		US 2008138446 A1	12-06-2008
		WO 2005002608 A1	13-01-2005

US 2011262558 A1	27-10-2011	CA 2797847 A1	10-11-2011
		EP 2563356 A2	06-03-2013
		US 2011262558 A1	27-10-2011
		WO 2011139794 A2	10-11-2011
