



US00PP17259P2

(12) **United States Plant Patent**  
**Schoenmakers**

(10) **Patent No.:** **US PP17,259 P2**

(45) **Date of Patent:** **Dec. 5, 2006**

(54) **FITTONIA PLANT NAMED 'WHITE STAR'**

(56) **References Cited**

(50) Latin Name: *Fittonia verschaffeltii*  
Varietal Denomination: **White Star**

U.S. PATENT DOCUMENTS

PP4,516 P \* 4/1980 Engelmann ..... Plt./373  
PP4,602 P \* 10/1980 Cobia ..... Plt./373

(75) Inventor: **Kees Schoenmakers**, Haaren (NL)

(73) Assignee: **Schoenmakers Tropische Potcultures**,  
Haaren (NL)

OTHER PUBLICATIONS

UPOV ROM GTITM Computer Database, GTI Jouve  
Retrieval Software 2005/05 Citations for 'White Star'.\*

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 4 days.

\* cited by examiner

(21) Appl. No.: **11/172,361**

*Primary Examiner*—Wendy Haas

(22) Filed: **Jun. 30, 2005**

(57) **ABSTRACT**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

A new cultivar of *Fittonia* plant named 'White Star' that is  
characterized by small light green to white leaves with dark  
green margins.

(52) **U.S. Cl.** ..... **Plt./373**

(58) **Field of Classification Search** ..... Plt./373  
See application file for complete search history.

**1 Drawing Sheet**

**1**

**2**

Botanical classification: *Fittonia verschaffeltii*.  
Variety denomination: 'White Star'.

BRIEF DESCRIPTION OF THE DRAWING

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar  
of *Fittonia* plant botanically known as *Fittonia verschaffeltii*  
and hereinafter referred to by the cultivar name 'White Star'.

The accompanying photograph illustrates the distinguish-  
ing traits of *Fittonia* 'White Star'. The plant in the photo-  
graph shows an overall view of a 12 week old plant. The  
photograph was taken using conventional techniques and  
although colors may appear different from actual colors due  
to light reflectance it is as accurate as possible by conven-  
tional photographic techniques.

The new cultivar was discovered by the inventor in a  
cultivated area of Haaren, The Netherlands in 2003. 'White  
Star' was discovered as a naturally occurring whole plant  
mutation of *Fittonia verschaffeltii* 'White Anne' (not  
patented).

BOTANICAL DESCRIPTION OF THE PLANT

Asexual reproduction by terminal cuttings of the new  
cultivar 'White Star' was first done in 2003 in Haaren, The  
Netherlands. Since that time, under careful observation, the  
unique characteristics of the new cultivar have been  
uniform, stable and reproduced true to type in successive  
generations of asexual reproduction.

The following is a detailed description of the new *Fittonia*  
cultivar named 'White Star'. Data was collected in Haaren,  
The Netherlands from 12 week old greenhouse grown plants  
in 8.5 cm. containers. The time of year was Spring and the  
average temperature was 24 degrees Centigrade during the  
day and 22 degrees Centigrade at night. No photoperiodic  
treatments were used. Color determinations are in accord-  
ance with The Royal Horticultural Society Colour Chart  
2001 edition, except where general color terms of ordinary  
dictionary significance are used. The growing requirements  
are similar to the species. 'White Star' has not been tested  
under all possible conditions and phenotypic differences  
may be observed with variations in environmental, climatic,  
and cultural conditions, however, without any variance in  
genotype.

SUMMARY OF THE INVENTION

The following represent the distinguishing characteristics  
of the new *Fittonia* cultivar 'White Star'.

1. *Fittonia* 'White Star' exhibits small leaves.
2. *Fittonia* 'White Star' exhibits light green to white  
leaves with dark green margins.

The closest comparison cultivars are *Fittonia* 'White  
Anne' (unpatented) and *Fittonia* 'Pink Star' (unpatented).  
The new cultivar 'White Star' is distinguishable from the  
parent plant 'White Anne' by the following characteristics:

1. 'White Star' has a more compact habit.
2. 'White Star' has crenate leaf margins.

The new cultivar 'White Star' is distinguishable from  
'Pink Star' by the following characteristics:

1. 'White Star' has a white veins. 'Pink Star' has pink  
veins.

Botanical classification: *Fittonia verschaffeltii* 'White Star'.  
Use: Ornamental.

Parentage: 'White Star' is a naturally occurring whole plant  
mutation of *Fittonia* 'White Anne'.

Vigor: Moderate.

Growth rate: Moderate.

Growth habit: Broad spreading, upright.

Plant shape: Flattened globular.

Suitable container size: 8.5 cm. diameter container.

Height: Average 4.4 cm.

Width: Average 8.9 cm.  
 Hardiness: USDA Zone 10.  
 Propagation: Terminal cuttings.  
 Time to initiate roots: Approximately 21 days to produce roots on an initial cutting at 24 degrees Centigrade.  
 Time to produce a rooted cutting: Approximately 35 days to produce a rooted cutting at 24 degrees Centigrade.  
 Crop time: 12 weeks.  
 Root system: Fine and fibrous.

## Stem:

*Branching habit.*—Moderately branching.  
*Average number of lateral branches.*—2.  
*Pinching.*—Not needed.  
*Lateral branch diameter.*—3 mm. in diameter.  
*Lateral branch length.*—3.2 cm. in length.  
*Lateral branch strength.*—Moderate.  
*Stem color.*—138B.  
*Pubescence.*—Dense, length 1 mm, color N155A.  
*Internode length.*—8 mm. between nodes.  
*Internode color.*—144B.  
*Shape.*—Rounded, dull.  
*Surface.*—Dull.  
*Stem strength.*—Moderate.

## Foliage:

*Leaf arrangement.*—Opposite.  
*Compound or single.*—Single.  
*Leaf shape.*—Oblanceolate to broad obovate, slightly convexed.  
*Leaf apex.*—Acute.  
*Leaf base.*—Attenuate.  
*Leaf texture.*—Dull, rugose.

*Leaf length.*—Average 4.5 cm. in length.  
*Leaf width.*—2 mm. in width.  
*Quantity of leaves per lateral branch.*—Average 8.  
*Pubescence.*—Short hairs on margins, 0.5 mm. in length, N155A.  
*Leaf margin.*—Repand and considerably wavy.  
*Vein pattern.*—Pinnate.  
*Young leaf color, (upper surface).*—145D with spots 145A to 145B, margins colored 137A to 139A.  
*Young leaf color, (lower surface).*—144B to 144C, venation colored 145D, margins colored 137A to 139A.  
*Mature leaf color, (upper surface).*—191C to 192C, margins colored 139A.  
*Mature leaf color, (lower surface).*—Between 137C to 138A, margins colored between 137A and 139A.  
*Vein color (upper surface).*—Between 155C to 157A.  
*Vein color (lower surface).*—145B to 145C.  
*Leaf attachment.*—Petiolate.  
*Petiole dimensions.*—Average 1.4 cm. in length, 2.2 mm. in diameter, 1.5 mm. in height.  
*Petiole color.*—144B.  
*Durability of foliage to stress.*—High.

Flowers: Flowers have not been observed.

Disease and insect resistance: Plants of the new *Fittonia* have not been observed for disease or insect resistance. It is claimed:

1. A new and distinct variety of *Fittonia* plant named 'White Star' as described and illustrated.

\* \* \* \* \*

