

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2017/219948 A1

(43) International Publication Date
28 December 2017 (28.12.2017)

(51) International Patent Classification:
C07D 487/04 (2006.01) *A61P 35/00* (2006.01)
A61K 31/519 (2006.01)

(21) International Application Number:
PCT/CN2017/089003

(22) International Filing Date:
19 June 2017 (19.06.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PCT/CN2016/086350
20 June 2016 (20.06.2016) CN

(71) Applicant: NOVARTIS AG [CH/CH]; Lichtstrasse 35,
4056 Basel (CH).

(72) Inventors; and

(71) Applicants (for US only): **LIU, Bo** [CN/CN]; Beijing Novartis Pharma Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN). **HUANG, Ying** [US/CN]; China Novartis Institutes for Bio-Medical Research Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN). **MAO, Liang** [CN/CN]; Beijing Novartis Pharma Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN). **WANG, Long** [CN/CN]; China Novartis Institutes for BioMedical Research Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN). **WAYKOLE, Liladhar Murlidhar** [US/CN]; Beijing Novartis Pharma Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN). **ZHANG, Lijun** [CN/CN]; Beijing Novartis Pharma Co., Ltd., 4218 Jinke Road, Zhangjiang Hi-Tech Park, Pudong New Area, Shanghai 201203 (CN).

(74) Agent: ZHONGZI LAW OFFICE; 7F, New Era Building,
26 Pinganli Xidajie, Xicheng District, Beijing 100034 (CN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: CRYSTALLINE FORMS OF TRIAZOLOPYRIMIDINE COMPOUND

(57) Abstract: Provided herein are crystalline forms of a triazolopyrimidine compound, which is useful for treating a PRC2-mediated disease or disorder.

CRYSTALLINE FORMS OF TRIAZOLOPYRIMIDINE COMPOUND

FIELD OF THE INVENTION

[0001] This invention relates to crystalline forms of *N*-(5-Fluoro-2,3-

5 dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, pharmaceutical compositions comprising the same, and methods of treatment using the crystalline forms and processes for preparing the crystalline forms.

BACKGROUND

10 [0002] Polycomb group (PcG) proteins are chromatin modifying enzymes that are dysregulated in many human cancers. The Polycomb Repressive Complex 2 (PRC2), which includes SUZ12 (suppressor of zeste 12), EED (embryonic ectoderm development) and the catalytic subunit, EZH2 (enhancer of zeste homolog 2), represses genes by methylating the core histone H3 lysine 27 (H3K27me3) at and around the promoter
15 regions of target genes. PRC2 is the critical component of cellular machinery involved in the epigenetic regulation of gene transcription and plays critical function in development and tissue differentiation and regeneration. Although EZH2 is the catalytic subunit, PRC2 requires at least EED and SUZ12 for its methyltransferase activity. EED, SUZ12 and EZH2 are overexpressed in many cancers, including but not limited to breast cancer,
20 prostate cancer, hepatocellular carcinoma and etc. EZH2 activating mutations have been identified in DLBCL (diffused large B cell lymphoma) patients and FL (follicular lymphoma) patients. Inhibition of PRC2 methyltransferase activity by compounds competing with the cofactor S-adenosyl methionine (SAM) in DLBCL reverses H3K27 methylation, re-activates expression of target genes and inhibits tumor
25 growth/proliferation. Therefore, PRC2 provides a pharmacological target for DLBCL and other cancers.

[0003] In preparing a pharmaceutical composition, a form of the therapeutic agent is sought that has a balance of desired properties, such as, for example, dissolution rate, solubility, bioavailability, and/or storage stability. The existence of multiple solid forms, 30 often referred to as polymorphs, is well known for solid pharmaceutical compounds, and the chemical and physical stability as well as handling properties of such compounds often depend on which solid form is used. Accordingly, the selection of a particular solid form of the active drug substance (e.g., a salt form, hydrated or solvated form, or polymorphic form) is often very important in the design of a reliable and reproducible

production process, and in storage, handling and distribution of a safe and effective form of the drug substance.

[0004] "Impurities: Guideline for Residual Solvents", is a guideline on amounts of residual solvents tolerated in pharmaceuticals, which was produced by the the International Council for Harmonisation of of Technical Requirements for Pharmaceuticals for Human Use (ICH). This Guideline recommends the use of less toxic solvents in the manufacture of drug substances and dosage forms, and sets pharmaceutical limits for residual solvents (organic volatile impurities) in drug products. This guidance divides solvents into three risk-based classes: Class 1 solvents, which are known to cause unacceptable toxicities; Class 2 solvents, which are associated with less severe toxicity; and Class 3 solvents, which have low toxic potential.


[0005] According to the ICH Guidance, residual amounts of Class 3 solvents of 50 mg per day or less (corresponding to 5000 ppm or 0.5%) would be acceptable without justification. The concentration limits are calculated using the equation below by assuming a product mass of 10 g administered daily:

$$\text{Concentration (ppm)} = (1000 \times \text{PDE})/\text{dose}$$

PDE (permitted daily exposure) is given in terms of mg/day and dose is given in g/day.


Higher amounts of Class 3 solvents may also be acceptable provided they are realistic in relation to manufacturing capability and good manufacturing practice.

[0006] Patent application number US Serial No. 14/977,273, which is herein incorporated by reference, discloses certain triazolopyrimidine compounds for treating PRC2-mediated diseases or disorders. This patent application was published on 23 June 2016 as US 2016-0176882. One compound disclosed in that application is *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (free form) and its hydrochloride salt. The free form (referred to herein as Compound A) has the following structure:



Compound A.

[0007] The hydrochloride salt (referred to herein as Compound X) has the following structure:



[0008] The free form, Compound A, displays low aqueous solubilities both at physiologically relevant pH level of 7.4 and in simulated gastric fluid (SGF). It was found that exposures of Compound A were under-proportional at higher doses in rodent 5 pharmacokinetic studies, which may lead to insufficient and variable exposures in clinical settings. Experiments were carried out to find an appropriate salt to improve aqueous solubility. It was discovered that only the hydrochloric salt provides the necessary improvement in solubility, and demonstrates the dose-proportional exposure in rodent after testing a number of salts that also included phosphate, fumarate, tartarate, 10 succinate, maleate and mesylate.

[0009] The procedure described in US 14/977,273 to produce Compound X uses isopropanol (IPA, a Class 3 solvent) as the medium. Variable amounts of IPA were observed in different batches of Compound X obtained by the procedure described in US 14/977,273 as determined by ¹H NMR (exemplified by entry 1 in table 2 below), with the 15 amount of IPA in some batches being more than 0.5% by weight. It has also been very difficult to remove the residual IPA from the final product. Additionally, Compound X thus obtained showed low and variable crystallinities, as judged from the XRPD spectra. Efforts were made to remove the residual IPA and improve crystallinity without much 20 success. Therefore, the need exists for methods that can consistently produce Compound X in the desired form, i.e., a form that is substantially free of residual organic solvent and which shows improved and reliable crystallinity.

SUMMARY

[0010] The present invention provides crystalline forms of Compound X (N-((5- 25 Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride).

[0011] Embodiments of these crystalline forms include those characterized herein as Forms A, H_A and H_B as well as combinations or mixtures of these crystalline forms.

Table 1

Compound X	Form
Anhydrous	A
monohydrate	H _A
dihydrate	H _B

[0012] The names used herein to characterize a specific form, e.g. "H_A" etc., should not be considered limiting with respect to any other substance possessing similar or

5 identical physical and chemical characteristics, but rather it should be understood that these designations are mere identifiers that should be interpreted according to the characterization information also presented herein.

[0013] The present invention provides methods of preparation of crystalline forms of Compound X.

10 **[0014]** The present invention also provides pharmaceutical compositions comprising crystalline form(s) of Compound X and at least one pharmaceutically acceptable carrier, diluent or excipient. The pharmaceutical composition may further comprise at least one additional therapeutic agent. Of particular interest are additional therapeutic agents selected from: other anti-cancer agents, immunomodulators, anti-allergic agents, anti-nausea agents (or anti-emetics), pain relievers, cytoprotective agents, and combinations thereof.

15 **[0015]** Crystalline form(s) of Compound X of the present invention may be used in the treatment of diseases or disorders mediated by EED and/or PRC2.

20 **[0016]** Crystalline form(s) of Compound X of the present invention may be used in therapy.

[0017] Crystalline form(s) of Compound X of the present invention may be used for the manufacture of a medicament for the treatment of diseases or disorders mediated by EED and/or PRC2.

25 **[0018]** Other features and advantages of the present disclosure will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019] FIG. 1 depicts a representative X-ray powder diffraction pattern of Form A (anhydrous form).

30 **[0020]** FIG. 2 depicts differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form A.

[0021] FIG. 3 depicts a dynamic vapor sorption (DVS) plot of Form A at 25°C.

[0022] FIG. 4 depicts a representative X-ray powder diffraction pattern of Form H_A (monohydrate form).

5 [0023] FIG. 5 depicts differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form H_A.

[0024] FIG. 6 depicts a dynamic vapor sorption plot of Form H_A at 25°C.

[0025] FIG. 7 depicts a dynamic vapor sorption plot of Form H_A at 50°C.

[0026] FIG. 8 depicts a representative X-ray powder diffraction pattern of Form H_B (dihydrate form).

10 [0027] FIG. 9 depicts differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form H_B.

[0028] FIG. 10 depicts a dynamic vapor sorption plot of Form H_B at 25°C.

[0029] FIG. 11 depicts comparison of Form A obtained by the procedure (A) described in Example 3 below vs. obtained by the procedure (B) described in Example 2 below (also described in US 14/977,273).

15

DETAILED DESCRIPTION

[0030] The following enumerated embodiments of the invention are representative:

20 1. Crystalline Form A of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, wherein said Form A contains less than 0.5% by weight of residual organic solvent.

25 2. Crystalline Form A of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, wherein said Form A contains less than 0.2% by weight of residual organic solvent.

30 3. Crystalline Form A of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, wherein said Form A contains less than 0.1% by weight of residual organic solvent.

35 4. Crystalline Form A of Compound X obtained by a process comprising the steps:

1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;

2) heating the resulting suspension to about or over 50 °C;

3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and

4) lowering the temperature of the resulting solution to to obtain Form A of Compound X.

5. Crystalline Form A of Compound X of any one of embodiments 1 to 3, obtained by a process comprising the steps:

10 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;

2) heating the resulting suspension to about or over 50 °C;

3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and

15 4) lowering the temperature of the resulting solution to to obtain Form A of Compound X.

6. Crystalline Form A of Compound X of embodiment 5, obtained by a process wherein:

20 in step 1) the water-miscible organic solvent is selected from a group consisting of ethanol and acetone;

in step 2) heating the resulting suspension to about 50 to 75 °C;

in step 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible

25 organic solvent, while maintaining the temperature at about 50 to 75 °C; and

in step 4) lowering the temperature of the resulting solution to to obtain Form A of Compound X.

7. A hydrated solid form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride.

30 8. The hydrated solid form according to embodiment 7, wherein said hydrated form is monohydrate Form H_A.

9. The hydrated solid form according to embodiment 7, wherein said hydrated form is dihydrate Form H_B.

10. A crystalline form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride (Form H_A) having a X-ray powder diffraction pattern comprising the following 20 values (CuK α λ =1.5418 Å): 13.8±0.1, 20.8±0.1, 26.2±0.1, 26.7±0.1, and 28.2 ±0.1. In some embodiments, Form H_A has one or more (e.g., 2, 3, 4 or 5) additional 20 values selected from those in List 2.

11. A crystalline form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride (Form H_B) having a X-ray powder diffraction pattern comprising the following 20 values (CuK α λ =1.5418 Å): 9.0±0.1, 17.1±0.1, 22.8±0.1, 26.9±0.1, and 35.3±0.1. In some embodiments, Form H_B has one or more (e.g., 2, 3, 4 or 5) additional 20 values selected from those in List 3.

12. A pharmaceutical composition, comprising one or more pharmaceutically acceptable carriers and Form A described in any one of embodiments 1 to 6.

13. The pharmaceutical composition of embodiment 12 further comprising at least one additional therapeutic agent.

14. The pharmaceutical composition of embodiment 13 where said additional therapeutic agent is selected from other anti-cancer agents, immunomodulators, anti-allergic agents, anti-emetics, pain relievers, cytoprotective agents, and combinations thereof.

15. Crystalline Form A described in any one of embodiments 1 to 6 for use in therapy.

16. Use of Form A described in any one of embodiments 1 to 6, in the manufacture of a medicament for treating a disease or disorder mediated by EED and/or PRC2.

17. The use of embodiment 16, wherein said disease or disorder is selected from diffused large B cell lymphoma, follicular lymphoma, other lymphomas, leukemia,

multiple myeloma, mesothelioma, gastric cancer, malignant rhabdoid tumor, hepatocellular carcinoma, prostate cancer, breast carcinoma, bile duct and gallbladder cancers, bladder carcinoma, brain tumors including neuroblastoma, schwannoma, glioma, glioblastoma and astrocytoma, cervical cancer, colon cancer, melanoma, endometrial cancer, esophageal cancer, head and neck cancer, lung cancer, nasopharyngeal carcinoma, ovarian cancer, pancreatic cancer, renal cell carcinoma, rectal cancer, thyroid cancers, parathyroid tumors, uterine tumors, and soft tissue sarcomas.

10 18. A process for preparing Crystalline Form A of Compound X comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

20 19. A process for preparing Crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

20. The process of embodiment 19 wherein:

35 in step 1) the water-miscible organic solvent is selected from a group consisting of ethanol and acetone;

in step 2) heating the resulting suspension to about 50 to 75 °C;
 in step 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature at about 50 to 75 °C; and
 5 in step 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

10 [0031] It was postulated that IPA may be trapped in the crystalline lattice when Compound X is formed in a medium of isopropanol, since during the entire salt formation process the material remained as a “slurry” and was never fully dissolved. Table 2 summarizes the results obtained from the experiments carried out with selected Class 3 solvents and combinations thereof. It was unexpectedly discovered that EtOH/water and acetone/water (Entries 11 and 12) enabled complete dissolution of Compound X and also sustained the supersaturation state for optimal duration, especially at elevated 15 temperature. These combinations provide crystalline forms of Compound X that are substantially free of organic solvents, e.g. containing not more than 0.3% and typically not more than 0.1% organic solvent by weight after normal drying.

Table 2

No.	Solvent	Compound A solubility (mg/mL)		Compound X solubility (mg/mL)	
		50 °C	rt	50 °C	rt
1	Acetone	<10	<10	<10	<10
2	ACN	<10	<10	<10	<10
3	IPA	<10	<10	<10	<10
4	MeOH	<10	<10	>10	>10
5	EtOH	<10	<10	<10	<10
6	EtOAc	<10	<10	<10	<10
7	THF	<10	<10	<10	<10
8	MTBE	<10	<10	<10	<10
9	Heptane	<10	<10	<10	<10
10	Water	<10	<10	<10	<10
11	EtOH/H ₂ O (1/1)	<10	<10	>50	<50

12	Acetone/H ₂ O (1/1)	<10	<10	>50	<50
----	-----------------------------------	-----	-----	-----	-----

[0032] With the selected solvent pair, the crystallization conditions were further optimized, as listed in Table 3. The conditions as shown in Entries 5 and 6 in Table 3 were found to produce the material with much lower content of the residual organic solvent and higher crystallinity, while better yield was obtained with the conditions in Entry 5. The conditions of Entry 5 could also be reproduced in multi-gram scale, as indicated as Entry 7 in Table 3.

Table 3

Entry	Amount	Temp.	Conditions			Amount of		
			Solvent, amount	Solvent multiples (fold)	Apprearence	Solvent residue	Solvent residue* (%)	Crystallinity
1	6 g	50 °C	IPA, 90 mL	15 x	slurry	IPA	0.80	Low
2	0.100 g	75 °C	EtOH, 5 mL	50 x	homogenous	EtOH	0.80	Medium
3	0.050 g	75 °C	EtOH, 4 mL	80 x	homogenous	EtOH	0.36	High
4	0.050 g	75 °C	EtOH, 5 mL	100 x	homogenous	EtOH	0.24	High
5	0.100 g	75 °C	95% EtOH / H ₂ O, 5 mL	50 x	homogenous	EtOH	0.16	High
6	0.100 g	75 °C	90% EtOH / H ₂ O, 5 mL	50 x	homogenous	EtOH	0.08	High
7	4 g	75 °C	95% EtOH / H ₂ O, 200mL	50 x	homogenous	EtOH	0.04	High

* The amount of the solvent residue was determined by ¹H NMR

Form A of Compound X (Anhydrous Form)

[0033] In one embodiment, Compound X is provided as a crystalline material comprising Form A. The crystalline form of Compound X comprises a crystalline form referred to herein as "Form A" of Compound X.

5 [0034] In one embodiment, Form A of Compound X is characterized by unit cell parameters approximately equal to the following:

Unit cell dimensions:	$a = 8.468(5) \text{ \AA}$	$\alpha = 102.58(2)^\circ$
	$b = 9.615(5) \text{ \AA}$	$\beta = 92.45(3)^\circ$
	$c = 12.102(7) \text{ \AA}$	$\gamma = 99.26(3)^\circ$

10

Space group: P-1

Molecules of Compound X/asymmetric unit: 2

Volume/Number of molecules in the unit cell = $946.0(9) \text{ \AA}^3$

Density (calculated) = 1.449 g/cm^3

15 wherein the unit cell parameters of Form A of Compound X are measured at a temperature of about $100(\pm 2) \text{ K}$ (-173°C).

20 [0035] In one embodiment, the invention provides Form A of Compound X which characterized by an X-ray powder diffraction pattern comprising the following 2θ values (CuK α $\lambda=1.5418 \text{ \AA}$): 12.5 ± 0.1 , 13.0 ± 0.1 , 25.2 ± 0.1 , and 30.8 ± 0.1 . In another embodiment, in addition to the above specified 2θ values, Form A exhibits one or more (e.g., 2, 3, 4 or 5) additional 2θ values selected from those in List 1. In yet another embodiment, Form A exhibits an X-ray powder diffraction pattern substantially the same as that shown in Figure 1.

25

List 1: XRPD peak listing for Form A of Compound X (2θ: most intense peaks are underlined)

Angle
$^{\circ}2\theta$
7.3
9.5
10.6
12.5
<u>13.0</u>

14.2
14.7
15.8
18.7
20.2
20.7
22.1
22.9
24.5
25.2
26.0
26.9
28.7
30.8
31.9
32.6
35.1

[0036] In another embodiment, Form A of Compound X exhibits a strong endotherm during DSC at about 264 ± 1 °C.

[0037] In another embodiment, Form A of Compound X has gradual loss of mass via

5 TGA amounting to about 0.6% loss by 200 °C.

Form H_A of Compound X (Monohydrate)

[0038] In one embodiment, Compound X is provided as a monohydrate in a crystalline form referred to herein as "Form H_A" of Compound X. Form H_A of Compound

10 X has a stoichiometry of one molecule of water for each molecule of Compound X.

[0039] In one aspect, the invention provides Form H_A of Compound X which characterized by an X-ray powder diffraction pattern comprising the following 20 values (CuK α $\lambda=1.5418$ Å): 13.8 ± 0.1 , 20.8 ± 0.1 , 26.2 ± 0.1 , 26.7 ± 0.1 , and 28.2 ± 0.1 . In another embodiment, in addition to the above specified 20 values, Form H_A exhibits one or more

15 (e.g., 2, 3, 4 or 5) additional 20 values selected from those in List 2. In yet another embodiment, Form H_A exhibits an X-ray powder diffraction pattern substantially the same as that shown in Figure 4.

List 2: XRPD peak listing for Form H_A of Compound X (2θ: most intense peaks are underlined)

Angle
°2θ
9.7
12.3
<u>13.4</u>
13.8
14.0
18.2
<u>18.9</u>
20.8
22.9
23.1
23.8
24.9
<u>26.2</u>
<u>26.7</u>
27.1
27.6
27.7
28.2
30.4
32.5

5 [0040] It has been discovered that monohydrate is easily formed in water/solvent mixtures with water activity above 0.5~0.6 and it shows acceptable stability to humidity and temperature.

[0041] In another embodiment, Form H_A of Compound X exhibits a strong endotherm during DSC at about 256 ± 1 °C.

10 [0042] In another embodiment, Form H_A of Compound X has gradual loss of mass via TGA amounting to about 4.2% loss by 111 °C.

Form H_B of Compound X (Dihydrate)

[0043] In one embodiment, Compound X is provided as a dihydrate in a crystalline form referred to herein as "Form H_B" of Compound X. Form H_B of Compound X has a stoichiometry of two molecules of water for each molecule of Compound X.

5 [0044] In one aspect, the invention provides Form H_B of Compound X which characterized by an X-ray powder diffraction pattern comprising the following 2θ values (CuK α $\lambda=1.5418$ Å): 9.0±0.1, 17.1±0.1, 22.8±0.1, 26.9±0.1, and 35.3±0.1. In another embodiment, in addition to the above specified 2θ values, Form H_B exhibits one or more (e.g., 2, 3, 4 or 5) additional 2θ values selected from those in List 3. In yet another 10 embodiment, Form H_B exhibits an X-ray powder diffraction pattern substantially the same as that shown in Figure 8.

List 3: XRPD peak listing for Form H_B of Compound X (2θ: most intense peaks are underlined)

15

Angle
°2θ
<u>9.0</u>
14.0
<u>17.1</u>
18.1
18.5
21.9
22.8
24.0
25.2
26.9
28.7
29.6
30.5
<u>35.3</u>
35.9
41.2

[0045] In another embodiment, Form H_B of Compound X exhibits a strong endotherm during DSC at about 244 ± 1 °C.

[0046] In another embodiment, Form H_B of Compound X has gradual loss of mass 5 via TGA amounting to about 8.5% loss by 122 °C.

[0047] The presence of reaction impurities and/or processing impurities may be determined by analytical techniques known in the art, such as, for example, chromatography, nuclear magnetic resonance spectroscopy, mass spectrometry, or 10 infrared spectroscopy.

[0048] In one aspect of the invention, the polymorphs of the invention have crystalline properties and are preferably at least 50% crystalline, more preferably at least 60% crystalline, still more preferably at least 70% crystalline and most preferably at least 80% crystalline. Crystallinity can be estimated by conventional X-ray diffractometry 15 techniques.

[0049] In some embodiments, the solid form of Compound X comprises one or more of the Forms described herein. A solid form of Compound X can include two or more of these Forms, i.e., it can be a mixture of two or more Forms. In some embodiments, a sample of the solid form mainly consists of a single Form selected from Forms A, H_A, and 20 H_B meaning that 50% or more of the material is of one solid Form. Relative amounts of various Forms in a mixture can be determined from XRPD data. As described herein, some of the Forms can evolve or interconvert under suitable conditions, such as Forms A and H_A, which can occur as a mixture, and can interconvert depending on the relative humidity and temperature at which the material is maintained.

[0050] In one aspect of the invention, the polymorphs of the invention are from 50%, 60%, 70%, 80% or 90% to 95%, 96%, 97%, 98%, 99% or 100% crystalline.

[0051] In the present specification, X-ray powder diffraction peaks (expressed in degrees 2θ) are measured using copper X-rays with a wavelength of 1.5406 Å (alpha1) and 1.5444 Å (alpha2).

[0052] The crystalline forms of the present invention can exist in either unsolvated or solvated forms. The term 'solvate' is used herein to describe a molecular complex comprising the compound of the invention and an amount of one or more pharmaceutically acceptable solvents. Examples of pharmaceutically acceptable solvents include ethanol and water. The term 'hydrate' is employed when the solvent is 30 water. Two polymorphs of Compound X as described herein are hydrates.

[0053] In another embodiment, the crystalline form is Form A of Compound X, wherein said Form A contains less than 0.5% by weight of residual organic solvent.

[0054] In another embodiment, the crystalline form is Form A of Compound X, wherein said Form A contains less than 0.4% by weight of residual organic solvent.

5 [0055] In another embodiment, the crystalline form is Form A of Compound X, wherein said Form A contains less than 0.3% by weight of residual organic solvent.

[0056] In another embodiment, the crystalline form is Form A of Compound X, wherein said Form A contains less than 0.2% by weight of residual organic solvent.

[0057] In another embodiment, the crystalline form is Form A of Compound X, 10 wherein said Form A contains less than 0.1% by weight of residual organic solvent.

[0058] In another embodiment, the crystalline form is Form H_A of Compound X.

[0059] In another embodiment, the crystalline form is Form H_B of Compound X.

[0060] In another embodiment, the crystalline form is Form A of Compound X in combination with Form H_A of Compound X.

15 [0061] In another embodiment, the crystalline form is Form A of Compound X in combination with Form H_A of Compound X.

[0062] In another embodiment, the crystalline form is Form A of Compound X in combination with Form H_B of Compound X.

[0063] In another embodiment, the crystalline form is Form A of Compound X in 20 combination with Form H_A of Compound X and Form H_B of Compound X.

[0064] In another embodiment, the present invention provides a composition comprising one solid form selected from any of the embodiment described above.

[0065] In another embodiment, the present invention provides a pharmaceutical composition comprising one solid form selected from any of the embodiment described 25 above and at least one pharmaceutically acceptable carrier, diluent or excipient.

[0066] In another embodiment, the present invention provides a pharmaceutical composition, comprising a therapeutically effective amount of one solid form selected from any of the embodiment described above and at least one pharmaceutically acceptable carrier, diluent or excipient.

30 [0067] The pharmaceutical composition is useful in the treatment of diseases or disorders mediated by EED and/or PRC2.

[0068] In another embodiment, the present invention provides a pharmaceutical composition as defined above further comprising additional therapeutic agent(s).

[0069] In another embodiment, the present invention provides one solid form 35 selected from any of the embodiment described above, for use in therapy, alone, or

optionally in combination with another compound of the present invention and/or at least one other type of therapeutic agent.

[0070] In another embodiment, the present invention provides one solid form selected from any of the embodiments described above, for use in therapy, for the 5 treatment of diseases or disorders mediated by EED and/or PRC2, alone, and/or at least one other type of therapeutic agent.

[0071] In another embodiment, the present invention provides a method for the treatment of diseases or disorders mediated by EED and/or PRC2, comprising 10 administering to a patient in need of such treatment a therapeutically effective amount of one solid form selected from any of the embodiments described above, alone, and/or at least one other type of therapeutic agent.

[0072] In another embodiment, the present invention provides a method for the treatment of diseases or disorders mediated by EED and/or PRC2, comprising 15 administering to a patient in need thereof a therapeutically effective amount of a first and second therapeutic agent, wherein the first therapeutic agent is one solid form selected from any of the embodiments described above, and the second therapeutic agent is one other type of therapeutic agent.

[0073] In another embodiment, the present invention also provides the use of one 20 solid form selected from any of the embodiments described above, for the manufacture of a medicament for the treatment of diseases or disorders mediated by EED and/or PRC2, alone, or optionally in combination with at least one other type of therapeutic agent.

[0074] In another embodiment, the present invention provides a combined 25 preparation of one solid form selected from any of the embodiments described above and additional therapeutic agent(s) for use in therapy.

[0075] In another embodiment, the present invention provides a combination of one solid form selected from any of the embodiments described above and additional therapeutic agent(s) for simultaneous or separate use in therapy.

[0076] In another embodiment, the present invention provides a combined 30 preparation of one solid form selected from any of the embodiments described above and additional therapeutic agent(s) for simultaneous, separate or sequential use in the treatment of diseases or disorders mediated by EED and/or PRC2. Said solid form may be administered as a pharmaceutical composition described herein.

[0077] Examples of diseases or disorders mediated by EED and/or PRC2 include, 35 but are not limited to, diffused large B cell lymphoma (DLBCL), follicular lymphoma, other

lymphomas, leukemia, multiple myeloma, mesothelioma, gastric cancer, malignant rhabdoid tumor, hepatocellular carcinoma, prostate cancer, breast carcinoma, bile duct and gallbladder cancers, bladder carcinoma, brain tumors including neuroblastoma, glioma, glioblastoma and astrocytoma, cervical cancer, colon cancer, melanoma, 5 endometrial cancer, esophageal cancer, head and neck cancer, lung cancer, nasopharyngeal carcinoma, ovarian cancer, pancreatic cancer, renal cell carcinoma, rectal cancer, thyroid cancers, parathyroid tumors, uterine tumors, and soft tissue sarcomas selected from rhabdomyosarcoma (RMS), Kaposi sarcoma, synovial sarcoma, osteosarcoma and Ewing's sarcoma.

10 **[0078]** In another embodiment, the present invention provides a method for the treatment of diseases or disorders mediated by EED and/or PRC2, comprising administering to a patient in need thereof a therapeutically effective amount of a first therapeutic agent optionally with a second therapeutic agent, wherein the first therapeutic agent is one solid form selected from any of the embodiments described 15 above and the second therapeutic agent is one other type of therapeutic agent; wherein the diseases or disorders are selected from diffused large B cell lymphoma (DLBCL), follicular lymphoma, other lymphomas, leukemia, multiple myeloma, gastric cancer, malignant rhabdoid tumor, and hepatocellular carcinoma.

20 **[0079]** In another embodiment, additional therapeutic agent(s) used in combined pharmaceutical compositions or combined methods or combined uses, are selected from one or more, preferably one to three, of the following therapeutic agents: other anti-cancer agents, immunomodulators, anti-allergic agents, anti-nausea agents (or anti-emetics), pain relievers, cytoprotective agents, and combinations thereof.

25 **[0080]** In another embodiment, the present invention provides a process of preparing crystalline Form A of Compound X, wherein said Form A contains less than 0.5% by weight of residual organic solvent, preferably less than 0.2% by weight of residual organic solvent, more preferably less than 0.1% by weight of residual organic solvent.

30 **[0081]** In another embodiment, the present invention provides a process of preparing crystalline Form A of Compound X, comprising the steps:

- 35 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to at least about 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at least about 50 °C; and

4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0082] In another embodiment, the present invention a process of preparing crystalline Form A of Compound X that contains less than 0.5% by weight of residual 5 organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 10 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

15 **[0083]** In another embodiment, the present invention provides a process of preparing crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, comprising the steps:

- 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 20 2) heating the resulting suspension to about 50 to 75 °C;
- 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature about 50 to 75 °C; and
- 4) lowering the temperature of the solution to prompt formation of crystals to obtain Form A of Compound X.

[0084] In another embodiment, the present invention provides crystalline Form A of Compound X obtained by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 30 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0085] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 5 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 10 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0086] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 15 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 2) heating the resulting suspension to about 50 to 75 °C;
- 20 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature at about 50 to 75 °C; and
- 4) lowering the temperature of the solution to prompt formation of crystals to obtain Form A of Compound X.

[0087] In another embodiment, the present invention provides crystalline Form A of Compound X obtainable by a process comprising the steps:

- 25 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 30 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0088] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 5 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while 10 maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0089] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, 15 preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtainable by a process comprising the steps steps:

- 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 20 2) heating the resulting suspension to about 50 to 75 °C;
- 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature about 50 to 75 °C; and
- 4) lowering the temperature of the solution to prompt formation of crystals to 25 obtain Form A of Compound X.

[0090] In another embodiment, the present invention provides a process of preparing crystalline Form A of Compound X, comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a 30 mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to at least about 50 °C;
- 3) acidifying the resulting suspension to form a clear solution with hydrochloric acid while maintaining the temperature at least about 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of 35 crystals to obtain Form A of Compound X.

[0091] In another embodiment, the present invention a process of preparing crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, comprising the steps:

- 5 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear 10 solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0092] In another embodiment, the present invention provides a process of preparing crystalline Form A of Compound X that contains less than 0.5% by weight of residual 15 organic solvent, comprising the steps:

- 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 2) heating the resulting suspension to about 50 to 75 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear 20 solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature about 50 to 75 °C; and
- 4) lowering the temperature of the solution to prompt formation of crystals to obtain Form A of Compound X.

[0093] In another embodiment, the present invention provides crystalline Form A of Compound X obtained by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear 30 solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0094] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 5 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear 10 solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0095] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, 15 preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 2) heating the resulting suspension to about 50 to 75 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear 20 solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature at about 50 to 75 °C; and
- 4) lowering the temperature of the solution to prompt formation of crystals to obtain Form A of Compound X.

[0096] In another embodiment, the present invention provides crystalline Form A of Compound X obtainable by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a 30 mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of 35 crystals to obtain Form A of Compound X.

[0097] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtained by a process comprising the steps:

- 5 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension to form a clear solution while 10 maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to prompt formation of crystals to obtain Form A of Compound X.

[0098] In another embodiment, the present invention provides crystalline Form A of Compound X that contains less than 0.5% by weight of residual organic solvent, 15 preferably less than 0.3% weight of residual organic solvent, more preferably less than 0.1% weight of residual organic solvent, obtainable by a process comprising the steps steps:

- 1) suspending Compound A in a mixture of water and water-miscible organic solvent selected from a group consisting of ethanol and acetone;
- 20 2) heating the resulting suspension to about 50 to 75 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature about 50 to 75 °C; and
- 25 4) lowering the temperature of the solution to prompt formation of crystals to obtain Form A of Compound X.

[0099] In further embodiments to the above processes of preparing crystalline Form A of Compound X, the present invention provides the following preferred ratios of the solvent mixture:

30 **[00100]** A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 80 to 100% by weight.

[00101] A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 85 to 100% by weight.

[00102] A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 90 to 100% by 35 weight.

[00103] A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 95 to 100% by weight.

[00104] A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 85 to 95% by weight.

5 [00105] A ratio of solvent mixture of H₂O and EtOH wherein EtOH is 90 to 95% by weight.

[00106] In further embodiments to the above processes of preparing crystalline Form A of Compound X, the present invention provides the following preferred temperature ranges for step 2:

10 [00107] A temperature range is 50 to 75 °C.

[00108] A temperature range is 60 to 75 °C.

[00109] A temperature range is 70 to 75 °C.

[00110] Various (enumerated) embodiments of the invention are described herein. It will be recognized that features specified in each embodiment may be combined with other specified features to provide further embodiments of the present invention. It is also understood that each individual element of the embodiments is its own independent embodiment.

15 [00111] Other features of the present invention should become apparent in the course of the above descriptions of exemplary embodiments that are given for illustration of the invention and are not intended to be limiting thereof.

20

III. DEFINITIONS

[00112] The general terms used hereinbefore and hereinafter preferably have within the context of this invention the following meanings, unless otherwise indicated, where more general terms wherever used may, independently of each other, be replaced by more specific definitions or remain, thus defining more detailed embodiments of the invention.

[00113] The features and advantages of the invention may be more readily understood by those of ordinary skill in the art upon reading the following detailed description. It is to be appreciated that certain features of the invention that are, for clarity reasons, described above and below in the context of separate embodiments, may also be combined to form a single embodiment. Conversely, various features of the invention that are, for brevity reasons, described in the context of a single embodiment, may also be combined so as to form sub-combinations thereof.

30

[00114] The term "a," "an," "the" and similar terms used in the context of the present invention (especially in the context of the claims) are to be construed to cover both the singular and plural unless otherwise indicated herein or clearly contradicted by the context.

5 **[00115]** All numbers expressing quantities of ingredients, weight percentages, temperatures, and so forth that are preceded by the word "about" are to be understood as only approximations so that slight variations above and below the stated number may be used to achieve substantially the same results as the stated number. Accordingly, unless indicated to the contrary, numerical parameters preceded by the word "about" are
10 approximations that may vary depending upon the desired properties sought to be obtained. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

15 **[00116]** The names used herein to characterize a specific form, e.g., "H_A" etc., are merely identifiers that are to be interpreted in accordance with the characterization information presented herein and are not to be limited so as to exclude any other substance possessing similar or identical physical and chemical characteristics.

20 **[00117]** All measurements are subject to experimental error and are within the spirit of the invention.

25 **[00118]** All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g. "such as") provided herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed.

[00119] The phrase "pharmaceutically acceptable" indicates that the substance or composition must be compatible chemically and/or toxicologically, with the other ingredients comprising a formulation, and/or the mammal being treated therewith.

30 **[00120]** Depending on the process conditions the end products of the present invention are obtained either in free (neutral) or salt form. Both the free form and the salts of these end products are within the scope of the invention. If so desired, one form of a compound may be converted into another form. A free base or acid may be converted into a salt; a salt may be converted into the free compound or another salt; a mixture of isomeric compounds of the present invention may be separated into the
35 individual isomers.

[00121] The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Allen, L.V., Jr., ed., *Remington: The Science and Practice of Pharmacy*, 22nd Edition, Pharmaceutical Press, London, UK (2012), the invention of which is hereby incorporated by reference.

10 [00122] As used herein, "polymorph(s)" refer to crystalline form(s) having the same chemical structure/composition but different spatial arrangements of the molecules and/or ions forming the crystals.

[00123] As used herein "amorphous" refers to a solid form of a molecule, atom, and/or ions that is not crystalline. An amorphous solid does not display a definitive X-ray diffraction pattern.

15 [00124] For pharmacopeial purposes, "residual organic solvents" in pharmaceuticals are defined as organic volatile chemicals that are used or produced in the manufacture of drug substances or excipients, or in the preparation of drug products. The residual organic solvents are not completely removed by practical manufacturing techniques.

20 [00125] As used herein, "water-miscible organic solvent" refers to an organic solvent that is liquid at room temperature and is completely miscible with water, preferably is selected from ethanol and acetone.

25 [00126] As used herein, a XRPD pattern "comprising" a number of peaks selected from a specified group of peaks, is intended to include XRPD patterns having additional peaks that are not included in the specified group of peaks.

[00127] "EED" refers to the protein product of the gene embryonic ectoderm development.

[00128] "PRC2" refers to Polycomb Repressive Complex 2.

30 [00129] The term "PRC2-mediated disease or disorder" refers to any disease or disorder which is directly or indirectly regulated by PRC2. This includes, but is not limited to, any disease or disorder which is directly or indirectly regulated by EED.

[00130] The term "diseases or disorders mediated by EED and/or PRC2" refers to diseases or disorders which are directly or indirectly regulated by EED and/or PRC2.

[00131] As used herein, the term "patient" encompasses all mammalian species.

[00132] As used herein, the term "subject" refers to an animal. Typically the animal is a mammal. A "subject" also refers to any human or non-human organism that could potentially benefit from treatment with an EED inhibitor. A subject also refers to for example, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, fish, birds and the like. In certain embodiments, the subject is a primate. In yet other embodiments, the subject is a human. Exemplary subjects include human beings of any age with risk factors for cancer disease.

[00133] As used herein, a subject is "in need of" a treatment if such subject would benefit biologically, medically or in quality of life from such treatment (preferably, a human).

[00134] As used herein, the term "inhibit", "inhibition" or "inhibiting" refers to the reduction or suppression of a given condition, symptom, or disorder, or disease, or a significant decrease in the baseline activity of a biological activity or process.

[00135] As used herein, the term "treat", "treating" or "treatment" of any disease/disorder refers to the treatment of the disease/disorder in a mammal, particularly in a human, and includes: (a) ameliorating the disease/disorder, (i.e., slowing or arresting or reducing the development of the disease/disorder, or at least one of the clinical symptoms thereof); (b) relieving or modulating the disease/disorder, (i.e., causing regression of the disease/disorder, either physically, (e.g., stabilization of a discernible symptom), physiologically, (e.g., stabilization of a physical parameter), or both); (c) alleviating or ameliorating at least one physical parameter including those which may not be discernible by the subject; and/or (d) preventing or delaying the onset or development or progression of the disease or disorder from occurring in a mammal, in particular, when such mammal is predisposed to the disease or disorder but has not yet been diagnosed as having it.

[00136] As used herein, "preventing" or "prevention" cover the preventive treatment (i.e., prophylaxis and/or risk reduction) of a subclinical disease-state in a mammal, particularly in a human, aimed at reducing the probability of the occurrence of a clinical disease-state. Patients are selected for preventative therapy based on factors that are known to increase risk of suffering a clinical disease state compared to the general population. "Prophylaxis" therapies can be divided into (a) primary prevention and (b) secondary prevention. Primary prevention is defined as treatment in a subject that has not yet presented with a clinical disease state, whereas secondary prevention is defined as preventing a second occurrence of the same or similar clinical disease state.

[00137] As used herein, "risk reduction" or "reducing risk" covers therapies that lower the incidence of development of a clinical disease state. As such, primary and secondary prevention therapies are examples of risk reduction.

[00138] "Therapeutically effective amount" is intended to include an amount of a compound of the present invention that will elicit the biological or medical response of a subject, for example, reduction or inhibition of EED and/or PRC2, or ameliorate symptoms, alleviate conditions, slow or delay disease progression, or prevent a disease or disorder mediated by PRC2. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the preventive or therapeutic effect, whether administered in combination, serially, or simultaneously.

[00139] Abbreviations as used herein, are defined as follows: "EtOH" for ethanol, "1x" for once, "2x" for twice, "3x" for thrice, " $^{\circ}\text{C}$ " for degrees Celsius, "aq" for aqueous, "Col" for column, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "L" for liter or liters, "mL" for milliliter or milliliters, " μL " for microliter or microliters, "N" for normal, "M" for molar, "nM" for nanomolar, "mol" for mole or moles, "mmol" for millimole or millimoles, "min" for minute or minutes, "h" for hour or hours, "rt" for room temperature, "RT" for retention time, "ON" for overnight, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "aq" for aqueous, "sat" or "sat'd" for saturated, "MW" for molecular weight, "mw" or " μwave " for microwave, "mp" for melting point, "Wt" for weight, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LC-MS" for liquid chromatography mass spectrometry, "HPLC" for high pressure liquid chromatography, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "nOe" for nuclear Overhauser effect spectroscopy, " ^1H " for proton, " δ " for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, "ee" for "enantiomeric excess" and " α ", " β ", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

30 GENERAL METHODS

[00140] The following methods were used in the exemplified Examples, except where noted otherwise.

[00141] Crystalline forms may be prepared by a variety of methods, including for example, crystallization or recrystallization from a suitable solvent, sublimation, growth from a melt, solid state transformation from another phase, crystallization from a

supercritical fluid, and jet spraying. Techniques for crystallization or recrystallization of crystalline forms from a solvent mixture include, for example, evaporation of the solvent, decreasing the temperature of the solvent mixture, crystal seeding a supersaturated solvent mixture of the molecule and/or salt, freeze drying the solvent mixture, and 5 addition of antisolvents (countersolvents) to the solvent mixture. High throughput crystallization techniques may be employed to prepare crystalline forms including polymorphs.

[00142] For crystallization techniques that employ solvent, the choice of solvent or solvents is typically dependent upon one or more factors, such as solubility of the 10 compound, crystallization technique, and vapor pressure of the solvent. Combinations of solvents may be employed, for example, the compound may be solubilized into a first solvent to afford a solution, followed by the addition of an antisolvent to decrease the solubility of the compound in the solution and to afford the formation of crystals. An antisolvent is a solvent in which the compound has low solubility.

15 **[00143]** In one method to prepare crystals, a compound is suspended and/or stirred in a suitable solvent to afford a slurry, which may be heated to promote dissolution. The term "slurry", as used herein, means a saturated solution of the compound, which may also contain an additional amount of the compound to afford a heterogeneous mixture of the compound and a solvent at a given temperature.

20 **[00144]** Seed crystals may be added to any crystallization mixture to promote crystallization. Seeding may be employed to control growth of a particular polymorph or to control the particle size distribution of the crystalline product. In general, seeds of small size are needed to control effectively the growth of crystals in the batch. Seed of small size may be generated by sieving, milling, or micronizing of large crystals, or by 25 micro-crystallization of solutions. Care should be taken that milling or micronizing of crystals does not result in any change in crystallinity from the desired crystal form (i.e., change to amorphous or to another polymorph).

[00145] A cooled crystallization mixture may be filtered under vacuum, and the isolated solids may be washed with a suitable solvent, such as cold recrystallization 30 solvent, and dried under a nitrogen purge to afford the desired crystalline form. The isolated solids may be analyzed by a suitable spectroscopic or analytical technique, such as solid state nuclear magnetic resonance, differential scanning calorimetry, x-ray powder diffraction, or the like, to assure formation of the preferred crystalline form of the product. The resulting crystalline form is typically produced in an amount of greater than 35 about 70 weight % isolated yield, preferably greater than 90 weight % isolated yield,

based on the weight of the compound originally employed in the crystallization procedure. The product may be comilled or passed through a mesh screen to delump the product, if necessary.

[00146] Crystalline forms may be prepared directly from the reaction medium of the final process for preparing Compound X. This may be achieved, for example, by employing in the final process step a solvent or a mixture of solvents from which Compound (I) may be crystallized. Alternatively, crystalline forms may be obtained by distillation or solvent addition techniques. Suitable solvents for this purpose include, for example, the aforementioned nonpolar solvents and polar solvents, including protic polar solvents such as alcohols, and aprotic polar solvents such as ketones.

X-ray Powder Diffraction (XRPD)

[00147] X-ray powder diffraction (XRPD) data were obtained using a Bruker D8 Discover with CuK α anode. Powder samples were placed on a glass slide and centered in the X-ray beam. The sample-detector distance was around 30 cm, three frames merged. The radiation was CuK α ($\lambda = 1.5418 \text{ \AA}$). Data were collected for $2 < 2\theta < 45^\circ$ with a sample exposure time of at least 270 seconds.

[00148] The derived atomic parameters (coordinates and temperature factors) were refined through full matrix least-squares. The function minimized in the refinements was $\Sigma_w (|F_{\text{O}}| - |F_{\text{C}}|)^2$. R is defined as $\Sigma ||F| - |F|| / \Sigma |F_{\text{O}}|$ while $R_w = [\Sigma_w (|F_{\text{O}}| - |F_{\text{C}}|)^2 / \Sigma_w |F_{\text{O}}|^2]^{1/2}$ where w is an appropriate weighting function based on errors in the observed intensities. Difference maps were examined at all stages of refinement. Hydrogen atoms were introduced in idealized positions with isotropic temperature factors, but no hydrogen parameters were varied.

25

Instrument	Bruker AXS DiscoverD8
Detector	PXC-VANTEC-500
Radiation	CuK α (0.15406 nm)
X-ray generator power	40 kV, 40 mA
Scan range	2° to 45° (2 theta value)
Scan time	90s/frame, 3 frames
X-ray optics	Monochromator
Detector distance	~30 cm

Thermogravimetric Analysis

[00149] Thermogravimetric analysis was conducted for each crystalline form using a TA Discovery TGA instrument. For each analysis, the TGA cell/sample chamber was purged with 20 ml/min of ultra-high purity nitrogen gas. A weight calibration was

5 performed using standard weights under nitrogen purge. The heating rate was 10°C per minute in the temperature range between rt and 300°C. The weight percentage change (wt%) was plotted versus the measured sample temperature.

Differential Scanning Calorimetry

10 **[00150]** Differential scanning calorimetry was conducted for each crystalline form using a TA Discovery DSC. For each analysis, the DSC cell/sample chamber was purged with 50 ml/min of ultra-high purity nitrogen gas. The instrument was calibrated with high purity indium. The heating rate was 10°C per minute in the temperature range between 30 and 300°C. The heat flow, which was normalized by sample weight, was plotted versus the measured sample temperature. The data were reported in units of watts/gram ("W/g"). The plot was made with the endothermic peaks pointing down. The endothermic melt peak (melting point) was evaluated for extrapolated onset temperature.

Dynamic Vapor Sorption

20 **[00151]** Dynamic vapor sorption was performed on a Surface Measurement Systems DVS Advantage instrument. Approximately 10 mg of material was loaded into a sample pan. The samples were exposed to sorption/desorption cycles in 10% relative humidity (RH) steps over the range of 50%-90%-0%-90%-50% RH at 25°C and 50°C. A target equilibrium condition of mass change less than 0.002% over 5min was set, with 25 minimum and maximum equilibrium periods of 10 and 360 min, respectively. The carrier gas was nitrogen with a flow rate of 100 mL/min. The system is calibrated with saturated salt solutions. The weight percentage change (wt%) of the sample at each stage was measured and plotted versus the target partial pressure.

TGA

Instrument	TA Discovery TGA
Temperature range	30-300 °C
Scan rate	10 K/min
Nitrogen flow	25mL/min

DSC

Instrument	TA Discovery DSC
Temperature range	30 °C-300 °C
Scan rate	10 K/min
Nitrogen flow	50 mL/min

DVS

Instrument	SMS DVS Advantage
Sample weight	About 10 mg
temperature	25 °C and 50°C
dm/dt	0.002 %/min

NMR Employed in Characterization of Examples

[00152] ¹H NMR spectra were obtained with Bruker Fourier transform spectrometers

5 operating at frequencies as follows: ¹H NMR: 400 MHz (Bruker). ¹³C NMR: 100 MHz (Bruker). Spectra data are reported in the format: chemical shift (multiplicity, number of hydrogens). Chemical shifts are specified in ppm downfield of a tetramethylsilane internal standard (δ units, tetramethylsilane = 0 ppm) and/or referenced to solvent peaks, which in ¹H NMR spectra appear at 2.49 ppm for CD₂HSOCD₃, 3.30 ppm for CD₂HOD, 10 1.94 for CD₃CN, and 7.24 ppm for CDCl₃, and which in ¹³C NMR spectra appear at 39.7 ppm for CD₃SOCD₃, 49.0 ppm for CD₃OD, and 77.0 ppm for CDCl₃. All ¹³C NMR spectra were proton decoupled. The measurements were carried out at room temperature or otherwise specified. The amount of solvent residue was calculated based on intergrations of most representative hydrogens in the corresponding ¹HNMR spectra.

15


V. EXAMPLES

[00153] The following Examples have been prepared, isolated and characterized using the methods disclosed herein. The following examples are not meant to be limiting of the scope of the invention.

20

Example 1: Preparation of Compound A (*N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine)

Intermediate 3: 8-bromo-5-(methylthio)-[1,2,4]triazolo[4,3-c]pyrimidine




5

[00154] **5-Bromo-4-hydrazinyl-2-(methylthio)pyrimidine (2):** To a solution of 5-bromo-4-chloro-2-(methylthio)pyrimidine (**1**, 49.0 g, 0.205 mol) in ethanol (1000 mL) was added hydrazine (21.5 g, 0.430 mol). The reaction was stirred at rt for 4 h. The resulting suspension was filtered, washed with hexane and dried in vacuum to give the title compound (44.1 g, 92%) as a white solid. $^1\text{H-NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ ppm 2.42 (s, 3H), 8.08 (s, 1H). LC-MS: $[\text{M}+\text{H}]^+ = 234.9$; 236.9.

[00155] *Intermediate 3: 5-bromo-4-hydrazinyl-2-(methylthio)pyrimidine (2)* (40.0 g, 0.17 mol) was dissolved in 200 mL triethoxymethane. The mixture was heated at reflux and stirred for 3 h. The reaction mixture was concentrated under reduced pressure, the residue was purified by flash chromatography (EA: PE=1:15~1:1) to give the title compound (38.3 g, 92%) as a white solid. $^1\text{H-NMR}$ (400 MHz, methanol-d₄) δ ppm 2.82 (s, 3H), 8.03 (s, 1H), 8.87 (s, 1H). LC-MS: [M+H]⁺ = 245.0; 247.0.

Intermediate A1: 8-bromo-N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-1,2,4,6-tetraazolo[4,3-c]pyrimidin-5-amine



[00156] 2-Bromo-4-(2,2-diethoxyethoxy)-1-fluorobenzene (A1.1): To a solution of 3-bromo-4-fluorophenol (500 g, 2.62 mol) and 2-bromo-1,1-diethoxyethane (670 g, 3.4 mol)

5 in 2.0 L DMF was added K_2CO_3 (1085 g, 7.86 mol) in one portion. The suspension was heated at 110 °C and stirred overnight under N_2 . After cooling to rt, the reaction was diluted with 10.0 L H_2O , and extracted with EtOAc (2.0 L \times 3). The combined organic phase was washed with brine twice, dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified on silica gel (EtOAc/hexane = 0:100 to 5:100) to give the title compound (810 g, 80%) as a yellow oil. $^1\text{H-NMR}$ (400 MHz, methanol-d4) δ ppm 1.27 (t, 6 H), 3.65 (q, 2 H), 3.78 (q, 2 H), 3.97 (d, 2 H), 4.82 (t, 1 H), 3.97 (d, 2 H), 6.84 (dd, 1 H), 7.04 (dd, 1 H), 7.13 (d, 1 H).


[00157] 4-Bromo-5-fluorobenzofuran (A1.2a along with regioisomer A1.2b): To a solution of PPA (1324 g, 3.93 mol) in toluene (2.0 L) was added A1.1 (810 g, 2.62 mol) over 30 min at 95 °C. The reaction mixture was stirred at 95 °C for 2 h. After cooling to rt, 4.0 L ice-water was added slowly. The mixture was extracted with PE (2.0 L \times 2), the combined organic phase was washed with brine (2.0 L \times 2), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified on silica gel (EtOAc/PE = 0:100 to 5:100) to give a mixture of A1.2a and A1.2b (A1.2a : A1.2b = 1:0.7, 310 g, 55% yield) as a yellow oil.

[00158] 5-Fluorobenzofuran-4-carbonitrile (A1.3): To a mixture of A1.2a and A1.2b (310 g, 1.44 mol) and $\text{Zn}(\text{CN})_2$ (253 g, 2.16 mol) in 1.0 L DMF was added $\text{Pd}(\text{PPh}_3)_4$ (162 g, 0.14 mol) under N_2 . The reaction mixture was heated at 100 °C and stirred for 18 h. After cooling to rt, the mixture was diluted with 5.0 L of water, and extracted with EtOAc (1.0 L \times 2). The combined organic phase was washed with brine (1 L), dried over


Na_2SO_4 (anhydrous), filtered and concentration under reduced pressure. The residue was purified by flash column (mobile phase: $\text{EtOAc/PE} = 1:70$ in 30 min, Ret. Time=11min, flow rate:120 mL/min) to give the title compound (92 g, 40%) as a white solid. $^1\text{H-NMR}$ (400 MHz, methanol- d_4) δ ppm 7.07 (d, 1H), 7.30 (dd, 1H), 7.89 (dd, 1H), 8.10 (dd, 1H).

5 **[00159] *tert-Butyl ((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)carbamate (A1.4):*** To a solution of **A1.3** (44.5 g, 276.4 mmol) and Boc_2O (90.0 g, 414.6 mmol) in 1.0 L MeOH was added Pd/C (5 g, 10% wt). The reaction mixture was degassed with H_2 and stirred under H_2 overnight. The mixture was filtered through celite, washed with MeOH (300 mL $\times 2$), the filtrate was concentrated under reduced pressure. The residue was 10 recrystallized from PE to give the title compound (61.0 g, 93%) as a white solid. $^1\text{H-NMR}$ (400 MHz, DMSO-d_6) δ ppm 1.38 (s, 9H), 3.21 (t, 2H), 4.12 (d, 2H), 4.53 (t, 2H), 6.63 (dd, 1H), 6.86 (dd, 1H), 7.25 (br s, 1H). LC-MS: $[\text{M}-^1\text{Bu}+\text{H}]^+ = 212.1$.

15 **[00160] *(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methanamine (A1.5):*** A solution of **A1.4** (18.3 g, 68.5 mmol) in 50 mL HCl/Dioxane (4 mol/L) was stirred at rt for 4 h. The mixture 20 was concentrated under reduced pressure. The residue was diluted with a mixture solvent ($\text{MeOH: MeCN} = 1:10$, 500 mL), then K_2CO_3 (18.0 g, 342.5 mmol) was added. The mixture was heated at 60 °C and stirred for 3 h, cooled to rt, filtered, and concentrated under reduced pressure. The crude product was purified on silica gel ($\text{MeOH: EtOAc} = 0:100$ to 1:4) to give the title compound (9.2 g, 80%) as a yellow oil. $^1\text{H-NMR}$ (400 MHz, methanol- d_4) δ ppm 3.27 (t, 2H), 3.77 (s, 2H), 4.56 (t, 2H), 6.59 (dd, 1H), 6.81 (dd, 1H). LC-MS: $[\text{M}+\text{H}]^+ = 168.1$.



25 **[00161] *Intermediate A1:*** A mixture of **A1.5** (1.41 g, 8.2 mmol) and 8-bromo-5-(methylthio)-[1,2,4]triazolo[4,3-c]pyrimidine (**3**) (1.0 g, 4.1 mmol) was heated at 40 °C and stirred for 16 h. After cooling to the rt, the mixture was diluted with EtOAc (35 mL). The precipitate was filtered and washed with EtOAc (3 mL $\times 3$), dried in vacuum to give the title compound (1.0 g, 67%) as a white solid. $^1\text{H NMR}$ (500 MHz, DMSO) δ ppm 3.27 (t, 2H), 4.53 (t, 2H), 4.66 (d, 2H), 6.71 (dd, 1H), 6.95 (t, 1H), 7.85 (s, 1H), 8.75 (t, 1H), 9.48 (s, 1H). LC-MS: $[\text{M}+\text{H}]^+ = 363.7; 365.7$.



[00162] To a mixture of **A1** (40 mg, 0.110 mmol) in 1,4-dioxane (3 mL), MeCN (0.30 mL) and water (0.30 mL) was added (2-methylpyridin-3-yl)boronic acid (30.1 mg, 0.220 mmol), potassium carbonate (45.5 mg, 0.330 mmol) and $\text{Pd}(\text{Ph}_3\text{P})_4$ (12.69 mg, 10.98

5 μmol). The resulting mixture was stirred under N_2 at 110 °C for 3 h, cooled to rt, and evaporated under vaccum. The residue was purified on flash chromatography (DCM: MeOH = 10:1) to afford *N*-(5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine as a white solid (20 mg, 46.0%).

10 [00163] Alternatively, *N*-(5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine was prepared as follows. To a suspension of **A1** (25.5 g, 70 mmol), 2-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (30.6 g, 140 mmol) and NaHCO_3 (35.3 g, 420 mmol) in a mixture solution of 1,4-dioxane (300 mL) and H_2O (100 mL) was added $\text{PdCl}_2(\text{dppf})$ (5.94 g, 612 mmol). The 15 mixture was degassed with N_2 , heated at 110 °C for 1 h. The resulting mixture was cooled to rt and conenetrated under reduced pressure. The residue was purified over column chromatography (EtOAc: MeOH = 20:1) to give 14 g of the desired product. 200 mL of acetone was added to the product, and the resulting suspension was heated at 50 °C for 2 h. The white solid was collected by filtration and dried under vacuum to give *N*- 20 ((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (13.6 g, 52%) $^1\text{H-NMR}$ (500 MHz, $\text{DMSO}-d_6$) δ ppm 2.40 (s, 3H), 3.33 (t, 2H), 4.56 (t, 2H), 4.72 (s, 2H), 6.72 (dd, 1H), 6.96 (dd, 1H), 7.31 (dd, 1H), 7.66 (s, 1H), 7.74 (d, 1H), 8.51 (d, 1H), 8.72 (t, 1H), 9.49 (s, 1H). LC-MS: $[\text{M}+\text{H}]^+ = 376.9$.

25 Example 2: Preparation of Form A of Compound X (*N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride) using isopropanol (IPA) (Entry 1 in Table 2)

[00164] To a suspension of *N*-(5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (6.0 g, 15.94 mmol) in 100 mL 30 of IPA, a solution of 0.5 N HCl in IPA (33.0 mL, 16.50 mmol) was added dropwise at rt. The suspension was stirred at 50 °C for 12 h, then cooled to rt and stirred for 5 h. The

resulting solid was collected by filtration, and dried at 40 °C under vacuum for 2 days to afford the hydrochloride salt of form A of compound X as a white solid (6.5 g, 98 %) ¹H NMR (DMSO-d₆) δ ppm 2.65 (s, 3H), 3.35 (t, 2H), 4.57 (t, 2H), 4.74 (d, 2H), 6.73 (dd, 1H), 6.97 (dd, 1H), 7.83 (s, 1H), 7.85 - 7.94 (m, 1H), 8.46 (d, 1H), 8.80 (dd, 1H), 9.07 (t, 1H), 9.58 (s, 1H). LC-MS: [M+H]⁺ = 376.9. The amount of solvent residue was calculated based on intergrations of most representative hydrogens in the corresponding ¹HNMR spectrum. Specifically, intregration at δ 1.04 accounting for 6 hydrogens of the methyl groupds of IPA is 0.31, while that of 3 hydrogens of the methyl group at 2-methyl-pyridine at δ 2.64 as 3. Therefore, mole percentage of IPA is calculated as follows: 0.31/ 6 (1+ 0.31 /6)= 4.9 %; while IPA percentage in weight is calculated as follows: 60 x (0.31/6) /{(376.38 + 36.46 +60 x (0.31/6)} = 0.74% {MW(IPA) =60, MW (compound A) = 376.40 and MW (HCl) = 36.46}.

Example 3: Preparation of Form A of Compound X (N-((5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride) using EtOH/ H₂O (Entry 7 in Table 2)

[00165] To 4.0 g of N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine in a 500 mL flask was added 200 mL of EtOH/H₂O (95/5, v/v), and the mixture was stirred by mechanical stirring. The resulting suspension was heated to 75 °C by oil bath and maintained at that temperature for 1 h. To the suspension maintained at 75 °C, 23.38 mL of 0.5 N HCl in EtOH (1.1eq) was added dropwisely. The mixture turned clear after the addition of hydrochloric acid. The resulting solution was stirred at 75 °C for 2 h. The mixture was then cooled down to rt in 3 h, and the resulting Compound X as a white solid (3.5 g, 79.7%) was collected by filtration and dried for 6 h under vacuum. The amount of EtOH residue was calculated based on the intergration of the hydrogen in the ¹H NMR spectrum. Specifically, intregratinon at δ 1.06 accounting for 3 H of the methyl group of EtOH is 0.01, while that of 3 hydrogens of the methyl group at 2-methyl-pyridine at δ 2.65 as 3. Therefore, mole percentage of EtOH was calculated as follows: 0.01/ 3 (1+ 0.01/3) = 0.33 %; while EtOH percentage in weight is calculated as follows: 46.07 x (0.01/3) /{(376.38 + 36.46 + x (0.01/3)} = 0.04% {MW(EtOH) =46.07, MW (compound A) = 376.40 and MW (HCl) = 36.46}.

Example 4: Preparation of Form H_A of Compound X

[00166] 1.0 g of Form A of Compound X was added to 10 mL of ethanol/water (3:1) mixture to obtain a suspension. The suspension was stirred at rt for 3 days. The resulting solid was collected with vacuum filtration and dried at rt overnight. Form H_A of Compound X (0.81g) was obtained in 77% yield.

Example 5: Preparation of Form H_B of Compound X

200 mg of Form A of Compound X was dissolved in minimum amount of acetone/water (1:1) mixtures at 60 °C, to obtain a clear solution. The solution was evaporated at rt over 10 3 to 5 days. The resulting solid was collected with vacuum filtration and dried at rt overnight. Form H_B of Compound X was obtained.

Preparation of 0.5 N HCl in EtOH or acetone:

[00167] 4.0 mL of the commercially available aqueous concentrated HCl (36.5%, w/w, in water) was added to 96.0 mL of EtOH or acetone (optionally at reduced temperature), 15 and the solution was mixed well to get 0.5 N HCl in EtOH or acetone.

Procedure for the experiments summarized in Table 2:

[00168] To 10 mg of compound X in a sample vial at 50 °C, selected solvent was added dropwise. The addition of solvent was stopped when a clear solution was 20 reached or when the amount of solvent reached 1 mL. The resulting mixture was stirred continuously at 50 °C for 2 h, then cooled to rt, and stirred overnight. The solubilities at 50 °C and rt were determined by the homogeneity of the resulting mixture.

Procedure for the experiments summarized in Table 3:

25 [00169] The experiments as Entries 1 to 6 in Table 3 were carried out according to the procedure described in the salt formation in Example 3 above, but with different parameters listed in the Table 3.

VI. PHARMACOLOGY AND UTILITY

30 [00170] As a key component of PRC2 complex, EED has no intrinsic enzymatic activity. However, it is critical for proper PRC2 function. EED directly binds to H3K27me3 and this binding event localizes the PRC2 complex to the chromatin substrate and allosterically activates the methyltransferase activity. Targeting the allosteric site within the regulatory EED subunit of PRC2, may offer a novel and unique angle to be 35 advantageous to, or complementary to, directly targeting the SAM competition

mechanism of EZH2 or PRC2. Therefore, targeting EED represents a highly attractive strategy for the development of a novel therapy for the treatment of many forms of cancers. In particular, the need exists for small molecules that inhibit the activity of PRC2 through targeting EED. It has now been found that triazolopyrimidine derivatives as presently disclosed are useful to target EED for the treatment of EED or PRC2-mediated diseases or disorders, especially cancers.

5 [00171] The utility of Compound X of the present invention may be demonstrated using any one of the following test procedures. Compound X was assessed for its ability to inhibit PRC2 activity in a pentameric complex of EZH2, SUZ12, EED, Rbap48 and

10 AEBP in biochemical assays. The ability of compounds of the present invention to inhibit cellular activity of PRC2 was assessed by analysing histone H3 lysine 27 methylation in human cell lines. The ability of Compound X to inhibit cancers was derived from their ability to modulate activity in human cancer cell lines bearing specific dependence to PRC2 activity to maintain cancerous growth.

15

EED-H3K27Me3 peptide competition binding assay by AlphaScreen (α -screen)

[00172] To assess the compounds' potency in the EED-H3K27Me3 competition binding assay, compounds were serially diluted 3-fold in DMSO to obtain a total of twelve concentrations. Then compounds at each concentration (75 nL of each) were transferred

20 by Mosquito into a 384-well Perkin Elmer ProxiPlate 384 plus plates. 8 μ L of solutions containing 30 nM EED (1-441)-His protein and 15 nM biotin-H3K27Me3 (19-33) peptide in the buffer (25 mM HEPES, pH 8, 0.02% Tween-20, 0.5% BSA) were added to the wells and then incubated with compound for 20 min. AlphaScreen detection beads mix was prepared immediately before use by mixing nickel chelate acceptor beads and

25 streptavidin donor beads in a 1:1 ratio (Perkin Elmer, Product No.6760619C/M/R) into the buffer described above. Then 4 μ L of detection beads mix was added to the plate and incubate in the dark at the rt for 1 h. The final concentration of donor and acceptor beads was 10 μ g/mL for each. Plates were read on EnVision (PerkinElmer) using the AlphaScreen setting adapted for optimal signal detection with a 615 nm filter, after

30 sample excitation at 680 nm. The emission signal at 615 nm was used to quantify compounds inhibition. AlphaScreen signals were normalized based on the reading coming from the positive (maximum signal control) and negative controls (minimum signal control) to give percentage of activities left. The data were then fit to a dose response equation using the program Helios (Novartis) to get the IC50 values. Helios is 35 a Novartis in-house assay data analysis software using the methods described by

Normolle, D. P., *Statistics in Medicine*, 12:2025-2042 (1993); Formenko, I. et al., *Computer Methods and Programs in Biomedicine*, 82, 31-37 (2006); Sebaugh, J. L., *Pharmaceutical Statistics*, 10:128-134 (2011); Kelly, C. et al., *Biometrics*, 46(4):1071-1085 (1990); and Kahm, M. et al., *Journal of Statistical Software*, 33(7): (2010) (gprof: 5 Fitting Biological Growth Curves with R, pages 1-21, available at <http://www.jstatsoft.org/>).

[00173] Each compound was counterscreened to determine if it interfered with the AlphaScreen beads. Compounds were diluted as described in the preceding section, and the assay was performed by adding 12 μ L of 10 nM biotin-miniPEG-His6 peptide in the 10 above buffer and incubating for 20 min at rt prior to addition of the beads to 10 μ g/mL each. The plates were then incubated for 1 h at rt in dark before being read on EnVision.

EED LC-MS Assay

[00174] Representative compounds of the present invention were serially and 15 separately diluted 3-fold in DMSO to obtain a total of eight or twelve concentrations. Then the test compounds at each concentration (120 nL of each) were transferred by Mosquito into a 384-well Perkin Elmer ProxiPlate 384 plus plates. Solutions (6 μ L) of 24 nM the wild type PRC2 (wtPRC2) complex and 2 μ M SAM in reaction buffer (20 mM Tris, pH 8.0, 0.1% BSA, 0.01% Triton, 0.5 mM DTT) were added to the wells that were then 20 incubated with the test compound for 20 min. A 6 μ L solution of 3 μ M of the peptide substrate H3K27Me0 (histone H3[21-44]-biotin) in reaction buffer was added to initiate each reaction. The final components in the reaction solution include 12 nM wtPRC2 complex, 1 μ M SAM, and 1.5 μ M H3K27me0 peptide with varying concentration of the 25 compounds. A positive control consisted of the enzyme, 1 μ M SAM and 1.5 μ M substrate in the absence of the test compound, and a negative control consisted of 1 μ M SAM and 1.5 μ M substrate only. Each reaction was incubated at rt for 120 min, then stopped by 30 addition of 3 μ L per of quench solution (2.5% TFA with 320 nM d4-SAH). The reaction mixture was centrifuged (Eppendorf centrifuge 5810, Rotor A-4-62) for 2 min at 2000 rpm and read on an API 4000 triple quadrupole mass spec with Turbulon Spray (Applied Biosystem) coupled with Prominence UFC (Shimadzu). The levels of SAH production were then normalized based on the values coming from the positive and negative controls to give percent enzyme activities. The data were then fit to a dose response equation using the program Helios to get the IC₅₀ values of the test compound.

35 ELISA (H3K27 methylation) assay

[00175] Representative compounds of the present invention were serially and separately diluted 3-fold in DMSO to obtain a total of eight or twelve concentrations. Then the compounds were added to G401 cell cultured in 384-well plate at 1:500 dilution to obtain the highest concentration of 20 μ M. The cells were further cultured for 48 h 5 before ELISA procedure.

[00176] Histone extraction: Cells, in 384-well plate, were washed with PBS (10 x PBS buffer (80 g NaCl (Sigma, S3014), 2 g KCl (Sigma, 60128), 14.4 g Na₂HPO₄ (Sigma, S5136), 2.4 g KH₂PO₄ (Sigma, P9791) to 1 L water, pH to 7.4) and lysed with the 10 addition of lysis buffer (0.4N HCl; 45 μ L per well). The plate was gently agitated at 4°C for 30 min. The cell lysate was neutralized with neutralization buffer (0.5 M sodium phosphate dibasic, pH 12.5, 1 mM DTT; 36 μ L per well). The plate was agitated to ensure the lysates were well mixed prior to the ELISA protocol.

[00177] ELISA protocol: Cell lysates were transferred to the wells of a 384-well plate and the final volume was adjusted to 50 μ L per well with PBS. The plate was sealed, centrifuged at 2,000 rpm for 2 min and incubated at 4°C for about 16 h. The plate was washed with TBST buffer (1 x TBS (10x TBS: 24.2 g Tris (Sigma, T6066), 80 g NaCl (Sigma, S3014) to 1 L of water and adjust pH to 7.6 with HCl) with 0.1% Tween-20). 15 Blocking buffer (TBST, 5% BSA; 50 μ L per well) was added and the plate was incubated for 1 h at rt. The blocking buffer was removed and primary antibody was added (30 μ L per well). The following dilutions were performed with blocking buffer: for anti-H3K27me3 antibody (Cell Signaling Technology, #9733), dilution was 1:1000; for anti-H3K27me2 antibody (Cell Signaling Technology, #9288), dilution was 1:100; for anti-H3 20 antibody (Abcam, Cat#24834), dilution was 1:1000. The primary antibody was incubated in the plate at rt for 1 h. The wells were washed with TBST and incubated with secondary antibody for 1 h at rt. For secondary antibodies, the following dilutions were carried out with blocking buffer: anti-rabbit antibody (Jackson ImmunoResearch, #111-035-003), dilution was 1:2000; and anti-mouse antibody (Cell signaling technology, 25 #7076), dilution was 1:1000. After 1 h of incubation at rt, the wells were washed with TBST. ECL substrate (Pierce, #34080) was added at 30 μ L per well and the plates were centrifuged at 2,000 rpm for 2 min. The signal was read using a PerkinElmer Envision Reader. The H3K27 methylation readouts were normalized using H3 signal and then 30 percentage inhibition was calculated against the samples treated with DMSO. The data

were then fit to a dose response curve using the program Helios to get the IC₅₀ values of the test compound.

Western Blot Analysis

5 [00178] Representative compounds of the present invention were analyzed for their ability to selectively inhibit PRC2. Western blot was performed using standard molecular biology techniques. Cell was lysed in SDS lysis buffer (Millipore, Cat#20-163) and protein concentration was measured by BCA protein assay (Pierce, Cat# PI-23221). Antibodies for western blots: anti-EZH2 (#3147), anti- H3 (#9715), anti-H3K4me1 (#9723), anti-H3K4me2 (#9725), anti-H3K4me3 (#9727), anti-H3K9me2 (#9753), anti-H3K36me2 (#9758), anti-H3K27me2 (#9755), and anti-H3K27me3 (#9756) were purchased from Cell Signaling Technology (Danvers, MA, USA). Anti-H3K9me1 (#07-395), anti-H3K27me1 (#07-448), and anti-H3K36me1 (#07-548) were purchased from Millipore (Billerica, MA, USA). Anti-H3K36me3 (ab9050-100) was purchased from Abcam (Cambridge, UK). Anti-H3K9me3 (#39161) was purchased from Active Motif (Carlsbad, CA, USA).

10 [00179] Compounds of the present invention specifically inhibit the methylation of the PRC2 substrate H3K27. This can be demonstrated by their ability to inhibit H3K27me2 and H3K27me3 in a number of human cancer cell lines, examples include rhabdoid cells (G401) and lymphoma cells (WSU-DLCL2, KARPAS422, SU-DHL4). Selectivity is 15 profiled against a number of other methylation marks, for example: H3K4me2; H3K9me2; H3K36me3; and H3K79me3.

Analysis of Cell Proliferation

20 [00180] B cell lymphoma cell KARPAS422 was cultured using standard cell culture conditions in RPMI-1640 (Invitrogen, cat #11875) supplemented with 15% FBS (Invitrogen, cat #10099-141) in humidified incubator at 37°C, 5% CO₂. To assess the effect of PRC2 inhibition on cell proliferation, exponentially growing cells were seeded at a density of 1 × 10⁵ cells/mL in 12-well plate (Corning, cat #CLS3513). After cell seeding, 25 a compound of the present invention was added to the cell media (in concentrations ranging from 0 to 100 µM, 3x dilution series). Viable cell numbers were determined every 3–4 days for up to 14 days using Vi-CELL (Beckman Coulter). On days of cell counting, fresh growth media and compound were replenished and cells split back to a density of 1 × 10⁵ cells/mL. Total cell number is expressed as split-adjusted viable cells 30 per mL. The dose response curves and IC₅₀ values were generated using Prism.

Analysis of Pharmacokinetic Properties

[00181] Pharmacokinetic properties of the compounds as presently disclosed can be determined by using the below described protocol.

5 **[00182]** A representative compound of the present invention was dissolved in 10% PEG300, 10% Solutol HS 15 and 80% pH 4.65 Acetate buffer to yield a final concentration of 0.2 mg/mL for intravenous (IV) and oral administration (PO).

10 **[00183]** For rat PK studies, a total of three male Sprague Dawley rats each were used for rat IV and PO PK study, respectively. The formulation solution was administered via a single bolus IV at 1 mg/kg and a single oral gavage (PO) at 2 mg/kg, respectively. Blood samples (approximately 150 μ L) were collected via jugular cannula at appropriate time points.

15 **[00184]** For mouse PK study, a total of twelve male ICR mice were used for IV and PO study, respectively. The formulation solution was administered via a single bolus IV at 1 mg/kg and a single oral gavage (PO) at 2 mg/kg, respectively. Blood samples (approximately 150 μ L) were collected via retro-orbital puncture (~150 μ L/mouse) after anesthetized by isoflurane or via cardiac puncture (terminal collection) at appropriate time points (n=3).

20 **[00185]** Samples were collected in tubes containing K3-EDTA and stored on ice until centrifuged. The blood samples were centrifuged at approximately 8000 rpm for 6 min at 2-8°C and the resulting plasma was separated and stored frozen at approximately -80°C. After adding the internal standard, the plasma samples were quantified by LC-MS/MS using the calibration curve. PK parameters including area under concentration curve (AUC), mean residence time (MRT), plasma clearance (Cl), steady state volume of distribution (Vdss), elimination half-life ($t_{1/2}$), maximum concentration (Cmax), time of maximum concentration (Tmax) and oral bioavailability (F %) were calculated using the following equations:

$$\text{AUC} = \int_0^{\infty} C \, dt$$

$$\text{MRT} = \frac{\int_0^{\infty} tC \, dt}{\int_0^{\infty} C \, dt} = \frac{\text{AUMC}}{\text{AUC}}$$

t is time and C is plasma concentration at the time (t);

30 Dose_{IV} is the dose for intravenous administration; and Dose_{oral} is the dose for oral administration.

$$Cl = \text{Dose}_{\text{iv}} / \text{AUC}$$

$$t_{1/2} = 0.693 \times \text{MRT}$$

$$Vdss = Cl * \text{MRT}$$

$$F \% = (\text{Dose}_{\text{iv}} \times \text{AUC}_{\text{oral}}) / (\text{Dose}_{\text{oral}} \times \text{AUC}_{\text{iv}}) \times 100\%$$

5

Protocol for High-Throughput Equilibrium Solubility Assay

[00186] Compounds of the present invention were first solubilized at 10 mM in pure DMSO. 20 μ L each of the DMSO stock solution was then transferred into 6 wells on 96-well plate. The DMSO solvent was dried with GeneVac solvent evaporator at 30 °C, 1

10 mbar vacuum for 1 h. After the addition of 200 μ L of buffer solutions (pH 6.8, or FaSSIF), the plate was sealed and shaken at 160 rpm for 24 h at rt. The plate was centrifuged at 3750 rpm for 20 min, 5 μ L of supernatant is mixed with 495 μ L of MeOH/H₂O (1:1). 0.01 μ M, 0.1 μ M, 1 μ M, 10 μ M stock solutions were prepared by series of dilution for the calibration curves. The supernatant was quantified by HPLC or LC/MS using the 15 calibration curve. High-Throughput equilibrium solubility was determined based on the concentration of the supernatant.

Efficacy studies in mouse xenograft model

[00187] All experiments conducted were performed in female athymic Nude-nu mice 20 in an AAALAC certificated facility. The animals were kept under SPF conditions in individual ventilation cages at constant temperature and humidity (i.e., 20-26 °C; 40-70%) with 5 or less animals in each cage. Animals had free access to irradiation sterilized dry granule food and sterile drinking water. All procedures and protocols were approved by the Institutional Animal Care and Use.

[00188] The cells Karpas 422 human B cell lymphoma were cultured in RPMI-1640 medium (Gibco; 11875-093) supplemented with 15% FBS (Gibco; 10099-141) and 1% Pen Strep (Gibco; 15140-122) at 37°C in an atmosphere of 5% CO₂ in air. Cells were maintained in suspension cultures at concentrations between 0.5 - 2 \times 10⁶ cells/ml. Cells were split at 1:3 every 2-4 days. To establish xenograft tumor models the cells were 30 collected, suspended in PBS, mixed with Matrigel (BD Bioscience) at a volume ratio of 1:1 at a concentration of 1 \times 10⁸ cells/mL and then injected subcutaneously into the right flank of balb/c nude mice (Vital River) at a concentration of 5 \times 10⁶ cells per animal.

[00189] The compound was formulated as a suspension in 0.5% methyl cellulose (MC) and 0.5% Tween 80 in 50 mM pH6.8 buffer (prepared in house according to the USP) 35 and administered orally by gavage at specific doses.

[00190] Treatment was initiated when the average tumor volume reached 100-300 mm³. Tumor growth and body weights were monitored at regular intervals. The two largest diameters, width (W) and length (L), of the xenograft tumors were measured manually with calipers and the tumor volume was estimated using the formula: 0.5 x L x 5 W².

[00191] When applicable, results are presented as mean \pm SEM. Graphing and statistical analysis was performed using GraphPad Prism 5.00 (GraphPad Software). Tumor and body weight change data were analyzed statistically. If the variances in the data were normally distributed (Bartlett's test for equal variances), the data were 10 analyzed using one-way ANOVA with *post hoc* Dunnet's test for comparison of treatment versus control group. The *post hoc* Tukey test was used for intragroup comparison. Otherwise, the Kruskal-Wallis ranked test *post hoc* Dunn's was used.

[00192] As a measure of efficacy the %T/C value is calculated at the end of the experiment according to:

$$15 \quad (\Delta\text{tumor volume}^{\text{treated}}/\Delta\text{tumor volume}^{\text{control}}) * 100$$

Tumor regression was calculated according to:

$$-(\Delta\text{tumor volume}^{\text{treated}}/\text{tumor volume}^{\text{treated at start}}) * 100$$

[00193] Where Δ tumor volumes represent the mean tumor volume on the evaluation day minus the mean tumor volume at the start of the experiment.

[00194] N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine was tested in the EED (a) Alphascreen binding Qualified, (b) LC-MS Qualified and (c) ELISA Qualified assays described above and found to have EED inhibitory activity.

IUPAC name	(a) IC ₅₀ (μ M)	(b) IC ₅₀ (μ M)	(c) IC ₅₀ (μ M)
N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine	0.0059	0.0089	0.0026

[00195] The antiproliferative activities (IC₅₀ values) in B cell lymphoma cell KARPAS422 after 14 days of treatment for N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine is 0.0030 μ M.

[00196] Accordingly, N-((5-fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine has been found to inhibit EED and is therefore useful in the treatment of diseases or disorders associated with EED and PRC2, which include, but are not limited to, diffused large B cell lymphoma (DLBCL),
5 follicular lymphoma, other lymphomas, leukemia, multiple myeloma, mesothelioma, gastric cancer, malignant rhabdoid tumor, hepatocellular carcinoma, prostate cancer, breast carcinoma, bile duct and gallbladder cancers, bladder carcinoma, brain tumors including neuroblastoma, glioma, glioblastoma and astrocytoma, cervical cancer, colon cancer, melanoma, endometrial cancer, esophageal cancer, head and neck cancer, lung
10 cancer, nasopharyngeal carcinoma, ovarian cancer, pancreatic cancer, renal cell carcinoma, rectal cancer, thyroid cancers, parathyroid tumors, uterine tumors, and soft tissue sarcomas selected from rhabdomyosarcoma (RMS), Kaposi sarcoma, synovial sarcoma, osteosarcoma and Ewing's sarcoma.

15 V. PHARMACEUTICAL COMPOSITIONS AND COMBINATIONS

[00197] A "pharmaceutically acceptable carrier (diluent or excipient)" refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals, including, generally recognized as safe (GRAS) solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drug stabilizers, binders, buffering agents (e.g., maleic acid, tartaric acid, lactic acid, citric acid, acetic acid, sodium bicarbonate, sodium phosphate, and the like), disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, and the like and combinations thereof, as would be known to those skilled in the art (see, for example, 20 Allen, L.V., Jr. et al., *Remington: The Science and Practice of Pharmacy* (2 Volumes), 22nd Edition, Pharmaceutical Press (2012). The formulations may be prepared using conventional dissolution and mixing procedures. For example, the bulk drug substance (i.e., compound of the present invention or stabilized form of the compound (e.g., complex with a cyclodextrin derivative or other known complexation agent)) is dissolved 25 in a suitable solvent in the presence of one or more of the excipients described above.

[00198] Compound X can be administered for any of the uses described herein by any suitable means, for example, orally, such as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions (including nanosuspensions, microsuspensions, 30 spray-dried dispersions), syrups, and emulsions; sublingually; buccally; parenterally, such

as by subcutaneous, intravenous, intramuscular, or intrasternal injection, or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally, including administration to the nasal membranes, such as by inhalation spray; 5 topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

[00199] Compound X is typically formulated into pharmaceutical dosage forms to provide an easily controllable dosage of the drug and to give the patient an elegant and 10 easily handleable product. The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the 15 frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. Compound X may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

[00200] In certain instances, it may be advantageous to administer Compound X in 20 combination with at least one additional pharmaceutical (or therapeutic) agent, such as other anti-cancer agents, immunomodulators, anti-allergic agents, anti-nausea agents (or anti-emetics), pain relievers, cytoprotective agents, and combinations thereof.

[00201] The term "combination therapy" refers to the administration of two or more therapeutic agents to treat a therapeutic disease, disorder or condition described in the 25 present invention. Such administration encompasses co-administration of these therapeutic agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of active ingredients. Alternatively, such administration encompasses co-administration in multiple, or in separate containers (e.g., capsules, powders, and liquids) for each active ingredient. Compound X and additional therapeutic 30 agents can be administered via the same administration route or via different administration routes. Powders and/or liquids may be reconstituted or diluted to a desired dose prior to administration. In addition, such administration also encompasses use of each type of therapeutic agent in a sequential manner, either at approximately the same time or at different times. In either case, the treatment regimen will provide

beneficial effects of the drug combination in treating the conditions or disorders described herein.

[00202] General chemotherapeutic agents considered for use in combination therapies include anastrozole (Arimidex®), bicalutamide (Casodex®), bleomycin sulfate (Blenoxane®), busulfan (Myleran®), busulfan injection (Busulfex®), capecitabine (Xeloda®), N4-pentoxy carbonyl-5-deoxy-5-fluorocytidine, carboplatin (Paraplatin®), carmustine (BiCNU®), chlorambucil (Leukeran®), cisplatin (Platinol®), cladribine (Leustatin®), cyclophosphamide (Cytoxan® or Neosar®), cytarabine, cytosine arabinoside (Cytosar-U®), cytarabine liposome injection (DepoCyt®), dacarbazine (DTIC-Dome®), dactinomycin (Actinomycin D, Cosmegan), daunorubicin hydrochloride (Cerubidine®), daunorubicin citrate liposome injection (DaunoXome®), dexamethasone, docetaxel (Taxotere®), doxorubicin hydrochloride (Adriamycin®, Rubex®), etoposide (Vepesid®), fludarabine phosphate (Fludara®), 5-fluorouracil (Adrucil®, Efudex®), flutamide (Eulexin®), tezacitibine, Gemcitabine (difluorodeoxycytidine), hydroxyurea (Hydrea®), Idarubicin (Idamycin®), ifosfamide (IFEX®), irinotecan (Camptosar®), L-asparaginase (ELSPAR®), leucovorin calcium, melphalan (Alkeran®), 6-mercaptopurine (Purinethol®), methotrexate (Folex®), mitoxantrone (Novantrone®), mylotarg, paclitaxel (Taxol®), nab-paclitaxel (Abraxane®), phoenix (Yttrium90/MX-DTPA), pentostatin, polifeprosan 20 with carmustine implant (Gliadel®), tamoxifen citrate (Nolvadex®), teniposide (Vumon®), 6-thioguanine, thiotapec, tirapazamine (Tirazone®), topotecan hydrochloride for injection (Hycamtin®), vinblastine (Velban®), vincristine (Oncovin®), and vinorelbine (Navelbine®).

[00203] Anti-cancer agents of particular interest for combinations with Compound X may include: cyclin-dependent kinase (CDK) inhibitors, checkpoint kinase (CHK) inhibitors, C-RAF inhibitors, phosphoinositide 3-kinase (PI3K) inhibitors, BCL-2 inhibitors, mitogen-activated protein kinase (MEK) inhibitors, topoisomerase II inhibitors, SRC inhibitors, histone deacetylase (HDAC) inhibitors, anti-tumor antibiotics, demethylating agents, and anti-estrogens.

[00204] Some patients may experience allergic reactions to the compounds of the present invention and/or other anti-cancer agent(s) during or after administration; therefore, anti-allergic agents are often administered to minimize the risk of an allergic reaction. Suitable anti-allergic agents include corticosteroids (Knutson, S., et al., *PLoS*

One, DOI:10.1371/journal.pone.0111840 (2014)), such as dexamethasone (e.g., Decadron®), beclomethasone (e.g., Beclovent®), hydrocortisone (also known as cortisone, hydrocortisone sodium succinate, hydrocortisone sodium phosphate, and sold under the tradenames Ala-Cort®, hydrocortisone phosphate, Solu-Cortef®, Hydrocort

5 Acetate® and Lanacort®), prednisolone (sold under the tradenames Delta-Cortel®, Orapred®, Pediapred® and Prelone®), prednisone (sold under the tradenames Deltasone®, Liquid Red®, Meticorten® and Orasone®), methylprednisolone (also known as 6-methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, sold under the tradenames Duralone®, Medralone®, Medrol®, M-Prednisol® 10 and Solu-Medrol®); antihistamines, such as diphenhydramine (e.g., Benadryl®), hydroxyzine, and cyproheptadine; and bronchodilators, such as the beta-adrenergic receptor agonists, albuterol (e.g., Proventil®), and terbutaline (Brethine®).

15 **[00205]** Immunomodulators of particular interest for combinations with the compounds of the present invention include one or more of: an activator of a costimulatory molecule or an inhibitor of an immune checkpoint molecule (e.g., one or more inhibitors of PD-1, 20 PD-L1, LAG-3, TIM-3 or CTLA4) or any combination thereof.

25 **[00206]** In certain embodiments, the immunomodulator is an activator of a costimulatory molecule. In one embodiment, the agonist of the costimulatory molecule is chosen from an agonist (e.g., an agonistic antibody or antigen-binding fragment thereof, 30 or a soluble fusion) of OX40, CD2, CD27, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB (CD137), GITR, CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3 or CD83 ligand.

35 **[00207]** In certain embodiments, the immunomodulator is an inhibitor of an immune checkpoint molecule. In one embodiment, the immunomodulator is an inhibitor of PD-1, PD-L1, PD-L2, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and/or TGFR beta. In one embodiment, the inhibitor of an immune checkpoint molecule inhibits PD-1, PD-L1, LAG-3, TIM-3 or CTLA4, or any combination thereof. The term “inhibition” or “inhibitor” includes a reduction in a certain parameter, e.g., an activity, of a given molecule, e.g., an immune checkpoint inhibitor. For example, inhibition of an activity, e.g., a PD-1 or PD-L1 activity, of at least 5%, 10%, 20%, 30%, 40% or more is included by this term. Thus, inhibition need not be 100%.

40 **[00208]** Some patients may experience nausea during and after administration of the compound of the present invention and/or other anti-cancer agent(s); therefore, anti-emetics are used in preventing nausea (upper stomach) and vomiting. Suitable anti-emetics include aprepitant (Emend®), ondansetron (Zofran®), granisetron HCl (Kytril®),

lorazepam (Ativan®), dexamethasone (Decadron®), prochlorperazine (Compazine®), casopitant (Rezonic® and Zunrisa®), and combinations thereof.

[00209] Medication to alleviate the pain experienced during the treatment period is often prescribed to make the patient more comfortable. Common over-the-counter

5 analgesics, such Tylenol®, are often used. However, opioid analgesic drugs such as hydrocodone/paracetamol or hydrocodone/acetaminophen (e.g., Vicodin®), morphine (e.g., Atramorph® or Avinza®), oxycodone (e.g., OxyContin® or Percocet®), oxymorphone hydrochloride (Opana®), and fentanyl (e.g., Duragesic®) are also useful for moderate or severe pain.

10 **[00210]** In an effort to protect normal cells from treatment toxicity and to limit organ toxicities, cytoprotective agents (such as neuroprotectants, free-radical scavengers, cardioprotectors, anthracycline extravasation neutralizers, nutrients and the like) may be used as an adjunct therapy. Suitable cytoprotective agents include Amifostine (Ethyol®), glutamine, dimesna (Tavocept®), mesna (Mesnex®), dextrazoxane (Zinecard® or

15 Totect®), xaliproden (Xapriila®), and leucovorin (also known as calcium leucovorin, citrovorum factor and folic acid).

[00211] The structure of the active compounds identified by code numbers, generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications).

20 **[00212]** In one embodiment, the present invention provides pharmaceutical compositions comprising Compound X together with a pharmaceutically acceptable carrier suitable for administration to a human or animal subject, either alone or together with other anti-cancer agents.

25 **[00213]** In one embodiment, the present invention provides methods of treating human or animal subjects suffering from a cellular proliferative disease, such as cancer. The present invention provides methods of treating a human or animal subject in need of such treatment, comprising administering to the subject a therapeutically effective amount of a compound of the present invention (e.g., a compound of the present invention) or a pharmaceutically acceptable salt thereof, either alone or in combination

30 with other anti-cancer agents.

[00214] In particular, compositions will either be formulated together as a combination therapeutic or administered separately.

[00215] In combination therapy for treatment of a malignancy, Compound X and other anti-cancer agent(s) may be administered simultaneously, concurrently or sequentially

with no specific time limits, wherein such administration provides therapeutically effective levels of the two compounds in the body of the subject.

[00216] In a preferred embodiment, Compound X and the other anti-cancer agent(s) is generally administered sequentially in any order by infusion or orally. The dosing

5 regimen may vary depending upon the stage of the disease, physical fitness of the patient, safety profiles of the individual drugs, and tolerance of the individual drugs, as well as other criteria well-known to the attending physician and medical practitioner(s) administering the combination. The compound of the present invention and other anti-cancer agent(s) may be administered within minutes of each other, hours, days, or even
10 weeks apart depending upon the particular cycle being used for treatment. In addition, the cycle could include administration of one drug more often than the other during the treatment cycle and at different doses per administration of the drug.

[00217] In another aspect of the present invention, kits that include Compound X and a combination partner as disclosed herein are provided. Representative kits include (a) 15 Compound X, (b) at least one combination partner, e.g., as indicated above, whereby such kit may comprise a package insert or other labeling including directions for administration.

[00218] Compound X may also be used to advantage in combination with known therapeutic processes, for example, the administration of hormones or especially 20 radiation. Compound X may in particular be used as a radiosensitizer, especially for the treatment of tumors which exhibit poor sensitivity to radiotherapy.

[00219] In another aspect of the present invention, kits that include Compound X and a combination partner as disclosed herein are provided. Representative kits include (a) 25 Compound X, (b) at least one combination partner, e.g., as indicated above, whereby such kit may comprise a package insert or other labeling including directions for administration.

[00220] In the combination therapies of the invention, Compound X and the other therapeutic agent may be manufactured and/or formulated by the same or different manufacturers. Moreover, Compound X and the other therapeutic (or pharmaceutical 30 agent) may be brought together into a combination therapy: (i) prior to release of the combination product to physicians (e.g. in the case of a kit comprising Compound X and the other therapeutic agent); (ii) by the physician themselves (or under the guidance of the physician) shortly before administration; (iii) in the patient themselves, e.g. during sequential administration of Compound X and the other therapeutic agent.

[00221] The pharmaceutical composition (or formulation) for application may be packaged in a variety of ways depending upon the method used for administering the drug. Generally, an article for distribution includes a container having deposited therein the pharmaceutical formulation in an appropriate form. Suitable containers are well-known to those skilled in the art and include materials such as bottles (plastic and glass), sachets, ampoules, plastic bags, metal cylinders, and the like. The container may also include a tamper-proof assemblage to prevent indiscreet access to the contents of the package. In addition, the container has deposited thereon a label that describes the contents of the container. The label may also include appropriate warnings.

10

WHAT IS CLAIMED IS:

1. A crystalline form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, wherein said crystalline form contains less than 0.5% by weight of residual organic solvent.
5
2. The crystalline form of claim 1, wherein said crystalline form contains less than 0.2% by weight of residual organic solvent.
10
3. The crystalline form of claim 1, wherein said crystalline form contains less than 0.1% by weight of residual organic solvent.
15
4. A crystalline form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride, having a x-ray powder diffraction pattern comprising 2θ values (CuK α $\lambda=1.5418\text{ \AA}$) of 12.5 ± 0.1 , 13.0 ± 0.1 , 25.2 ± 0.1 , and 30.8 ± 0.1 .
20
5. The crystalline form of claim 4, having a x-ray powder diffraction pattern comprising at least two additional 2θ values (CuK α $\lambda=1.5418\text{ \AA}$) selected from the group consisting of 13.8 ± 0.1 , 20.8 ± 0.1 , 26.2 ± 0.1 , 26.7 ± 0.1 , and 28.2 ± 0.1 .
25
6. The crystalline form of claim 4, which contains less than 0.5% by weight of residual organic solvent.
30
7. Crystalline *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride (Compound X) obtained by a process comprising the steps:
 - 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
 - 2) heating the resulting suspension to about or over 50 °C;
 - 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution while maintaining the temperature at about or over 50 °C; and
 - 4) lowering the temperature of the resulting solution to obtain crystalline Compound X.

8. The crystalline Compound X of any one of claims 1 to 7 obtained by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 5 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain crystalline

10 Compound X.

9. The crystalline Compound X of claim 8, obtained by the process wherein:

- in step 1) the water-miscible organic solvent is selected from a group consisting of ethanol and acetone;
- 15 in step 2) heating the resulting suspension to about 50 to 75 °C;
- in step 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature at about 50 to 75 °C; and
- 20 in step 4) lowering the temperature of the resulting solution to obtain crystalline Compound X.

10. Crystalline Form A of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride (Compound X) obtainable by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 30 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution while maintaining the temperature about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

11. The crystalline form of any one of claims 1 to 3 obtainable by a process comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 5 2) heating the resulting suspension to about or over 50 °C;
- 3) acidifying the resulting suspension with hydrochloric acid to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain Form A of

10 Compound X.

12. Crystalline Form A of claim 10, obtained by the process wherein:

- in step 1) the water-miscible organic solvent is selected from a group consisting of ethanol and acetone;
- 15 in step 2) heating the resulting suspension to about 50 to 75 °C;
- in step 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature about 50 to 75 °C; and
- 20 in step 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

13. A hydrated solid form of *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine hydrochloride.

25 14. The hydrated solid form according to claim 13, wherein said hydrated form is monohydrate Form H_A.

15. The hydrated solid form according to claim 13, wherein said hydrated form is monohydrate Form H_B.

30 16. The hydrated solid form according to claim 14, having a X-ray powder diffraction pattern comprising the following 2θ values (CuK α $\lambda=1.5418 \text{ \AA}$): 13.8±0.1, 20.8±0.1, 26.2±0.1, 26.7±0.1, and 28.2 ±0.1.

17. The hydrated solid form according to claim 15, having a X-ray powder diffraction pattern comprising the following 2θ values (CuK α $\lambda=1.5418\text{ \AA}$): 9.0±0.1, 17.1±0.1, 22.8±0.1, 26.9±0.1, and 35.3±0.1.
- 5 18. A pharmaceutical composition, comprising one or more pharmaceutically acceptable carriers and crystalline Compound X according to any one of claims 1 to 11.
- 10 19. The pharmaceutical composition of claim 17 further comprising at least one additional therapeutic agent.
- 15 20. The pharmaceutical composition of claim 18 where said at least one additional therapeutic agent is selected from other anti-cancer agents, immunomodulators, anti-allergic agents, anti-emetics, pain relievers, cytoprotective agents, and combinations thereof.
21. Crystalline Compound X according to any one of claims 1 to 11 for use in therapy.
22. Use of crystalline Compound X according to any one of claims 1 to 11, in the manufacture of a medicament for treating a disease or disorder mediated by EED and/or PRC2.
- 25 23. The use of claim 22, wherein said disease or disorder is selected from diffused large B cell lymphoma, follicular lymphoma, other lymphomas, leukemia, multiple myeloma, mesothelioma, gastric cancer, malignant rhabdoid tumor, hepatocellular carcinoma, prostate cancer, breast carcinoma, bile duct and gallbladder cancers, bladder carcinoma, brain tumors including neuroblastoma, schwannoma, glioma, glioblastoma and astrocytoma, cervical cancer, colon cancer, melanoma, endometrial cancer, esophageal cancer, head and neck cancer, lung cancer, nasopharyngeal carcinoma, ovarian cancer, pancreatic cancer, renal cell carcinoma, rectal cancer, thyroid cancers, parathyroid tumors, uterine tumors, and soft tissue sarcomas.
- 30 24. A process for preparing Crystalline Form A of Compound X comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 5 3) acidifying the resulting suspension to form a clear solution with hydrochloric acid while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

10 25. A process for preparing Crystalline Form of any one of claims 1 to 3, comprising the steps:

- 1) suspending *N*-(5-Fluoro-2,3-dihydrobenzofuran-4-yl)methyl)-8-(2-methylpyridin-3-yl)-[1,2,4]triazolo[4,3-c]pyrimidin-5-amine (Compound A) in a mixture of water and water-miscible organic solvent;
- 2) heating the resulting suspension to about or over 50 °C;
- 15 3) acidifying the resulting suspension to form a clear solution while maintaining the temperature at about or over 50 °C; and
- 4) lowering the temperature of the resulting solution to obtain crystalline Compound X.

20

26. The process of claim 24 wherein:

- in step 1) the water-miscible organic solvent is selected from a group consisting of ethanol and acetone;
- in step 2) heating the resulting suspension to about 50 to 75 °C;
- 25 in step 3) acidifying the resulting suspension to form a clear solution by adding a solution of 0.5 N HCl in a mixture of water and said water-miscible organic solvent, while maintaining the temperature at about 50 to 75 °C; and
- in step 4) lowering the temperature of the resulting solution to obtain Form A of Compound X.

30




FIG. 1 - Representative X-ray powder diffraction pattern of Form A (anhydrous form)

FIG. 2 - Differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form A




FIG. 3 - Dynamic vapor sorption (DVS) plot of Form A at 25°C





FIG. 4 - Representative X-ray powder diffraction pattern of Form H_A (monohydrate form)


FIG. 5 - Differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form H_A


FIG. 6 - Dynamic vapor sorption plot of Form H_A at 25°C


FIG. 7 - Dynamic vapor sorption plot of Form H_A at 50°C



FIG. 8 - Representative X-ray powder diffraction pattern of Form H₈ (dihydrate form)

FIG. 9 - Differential scanning calorimetry (DSC) / thermogravimetric analysis (TGA) thermograms of Form H_B



FIG. 10 - Dynamic vapor sorption plot of Form H_B at 25°C

FIG. 11 - Comparison of Form A obtained by the procedure (A) described in Example 3 vs. obtained by the procedure (B) described in Example 2 (also described in US 14/977,273)



INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2017/089003

A. CLASSIFICATION OF SUBJECT MATTER

C07D 487/04(2006.01)i; A61K 31/519(2006.01)i; A61P 35/00(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D; A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CNPAT;EPODOC;WPI;CNKI;STN:三唑并嘧啶, 二氢苯并呋喃, triazolo, pyrimidine, dihydrobenzofurane, PRC2, EED, EZH2, SUZ12, crystalline

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2012118812 A2 (EPIZYME, INC.ET AL.) 07 September 2012 (2012-09-07) claim 1 and paragraphs [01273]-[01285]	1-26
A	WO 03044021 A2 (AMGEN INC.ET AL.) 30 May 2003 (2003-05-30) pages 33-35 and abstract	1-26
A	WO 2014144747 A1 (EPIZYME, INC.) 18 September 2014 (2014-09-18) claim 1 and page 1	1-26
PX	WO 2016103155 A1 (NOVARTIS AG ET AL.) 30 June 2016 (2016-06-30) Example 2	1-26

 Further documents are listed in the continuation of Box C. See patent family annex.

- * Special categories of cited documents:
- “A” document defining the general state of the art which is not considered to be of particular relevance
- “E” earlier application or patent but published on or after the international filing date
- “L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- “O” document referring to an oral disclosure, use, exhibition or other means
- “P” document published prior to the international filing date but later than the priority date claimed

- “T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- “X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- “Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- “&” document member of the same patent family

Date of the actual completion of the international search 30 August 2017	Date of mailing of the international search report 21 September 2017
--	--

Name and mailing address of the ISA/CN STATE INTELLECTUAL PROPERTY OFFICE OF THE P.R.CHINA 6, Xitucheng Rd., Jimen Bridge, Haidian District, Beijing 100088 China Facsimile No. (86-10)62019451	Authorized officer CUI, Yongtao Telephone No. (86-10)61648259
---	---

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/CN2017/089003

Patent document cited in search report				Publication date (day/month/year)		Patent family member(s)		Publication date (day/month/year)	
WO	2012118812	A2	07 September 2012	US	9637472	B2		02 May 2017	
				AU	2012223448	B2		16 March 2017	
				US	2015065483	A1		05 March 2015	
				US	2015266854	A1		24 September 2015	
				US	2014057891	A1		27 February 2014	
				US	2013317026	A1		28 November 2013	
				US	8598167	B1		03 December 2013	
				US	9206157	B2		08 December 2015	
				US	8962620	B2		24 February 2015	
				JP	2014511389	A		15 May 2014	
				EP	2681216	A2		08 January 2014	
				AU	2012223448	A1		05 September 2013	
				AU	2017204062	A1		06 July 2017	
				EP	2681216	A4		01 October 2014	
				WO	2012118812	A3		08 November 2012	
WO	03044021	A2	30 May 2003	WO	03044021	A3		18 December 2003	
				PT	1448564	T		30 June 2006	
				AU	2002352722	B2		12 October 2006	
				DE	60210805	T2		14 September 2006	
				JP	2005518358	A		23 June 2005	
				PL	371428	A1		13 June 2005	
				MX	PA04004552	A		13 August 2004	
				US	2003195221	A1		16 October 2003	
				PT	1448564	E		30 June 2006	
				EP	1448564	B1		19 April 2006	
				US	6921762	B2		26 July 2005	
				ES	2262879	T3		01 December 2006	
				AT	323705	T		15 May 2006	
				DK	1448564	T3		21 August 2006	
				AU	2002352722	A1		10 June 2003	
				CA	2466072	A1		30 May 2003	
				EP	1448564	A2		25 August 2004	
				DE	60210805	D1		24 May 2006	
WO	2014144747	A1	18 September 2014	EP	2970306	A4		03 August 2016	
				US	2016024081	A1		28 January 2016	
				EP	2970306	A1		20 January 2016	
WO	2016103155	A1	30 June 2016	UY	36462	A		29 July 2016	
				US	2016176882	A1		23 June 2016	
				US	9580437	B2		28 February 2017	
				AU	2015370524	A1		01 June 2017	
				TW	201629065	A		16 August 2016	
				SG	11201703880V	A		28 July 2017	
				DO	P2017000149	A		15 July 2017	