




(19)

Europäisches
Patentamt
European
Patent Office
Office européen
des brevets



(11)

EP 3 064 198 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
07.09.2016 Bulletin 2016/36

(51) Int Cl.:
A61K 9/14 (2006.01) A61K 9/50 (2006.01)
A61K 31/485 (2006.01) A61K 31/135 (2006.01)
A61K 31/4468 (2006.01)

(21) Application number: 16163685.7

(22) Date of filing: 02.09.2008

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR

(30) Priority: 03.09.2007 US 969672 P

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
08829074.7 / 2 197 429

(71) Applicant: Nanotherapeutics, Inc.
Alachua, FL 32615 (US)

(72) Inventor: TALTON, James, D.
Alachua, FL Florida 32615 (US)

(74) Representative: Pohlman, Sandra M.
df-mp Dörries Frank-Molnia & Pohlman
Patentanwälte Rechtsanwälte PartG mbB
Theatinerstrasse 16
80333 München (DE)

Remarks:

This application was filed on 04-04-2016 as a
divisional application to the application mentioned
under INID code 62.

(54) PARTICULATE COMPOSITIONS FOR DELIVERY OF POORLY SOLUBLE DRUGS

(57) Described herein are compositions comprising
particles of poorly soluble drugs, in particular opioids, en-
capsulated by stabilizers, in particular polyethylene gly-
col (PEG). Further described are pharmaceutical com-
positions comprising such encapsulated compositions.
Also described are methods of making such encapsulat-

ed particle compositions, and methods of making the cor-
responding pharmaceutical compositions. The encapsu-
lated particle compositions described herein allow poorly
soluble drugs to be administered with good bioavailability
by routes that are non-invasive to patients, such as by
oral administration.

Description**FIELD OF THE INVENTION**

5 [0001] The present invention relates generally to compositions comprising particles of poorly soluble drugs encapsulated by stabilizers, and to pharmaceutical compositions comprising such encapsulated compositions. It further relates to methods of making such encapsulated particle compositions, and to methods of making the corresponding pharmaceutical compositions. The encapsulated particle compositions described herein allow poorly soluble drugs to be administered with good bioavailability by routes that are non-invasive to patients, such as by oral administration.

10

BACKGROUND

15 [0002] Oral administration of drugs is generally preferred for reasons of patient comfort and compliance. However, many drugs, including many opioids, are poorly soluble at neutral pH, and are thus poorly or variably absorbed when delivered orally. Consequently, many such drugs are administered through more invasive routes, such as by sublingual, buccal, subcutaneous, or intravenous routes.

20 [0003] Several approaches for improving the oral delivery of poorly soluble drugs have demonstrated some promise. For example, poorly soluble drugs have been administered as dispersions in large amounts of fatty acids, and have been wet-milled to yield nanoparticles. However, each of those approaches suffers from certain drawbacks, such as, e.g., inadequate stability, difficulty of manufacture, adverse interactions with the drug to be delivered, or the use of toxic amounts of adjuvants or inhibitors. Thus, there remains a need for compositions and methods for the non-invasive delivery of poorly soluble drugs.

SUMMARY

25

[0004] In some embodiments, the invention provides a composition comprising particles of a poorly soluble drug (such as, e.g., an opioid) encapsulated by a stabilizer. The stabilizer may be, e.g., a polymer, including, e.g., a water-soluble polymer of neutral charge. For example, in some embodiments, the invention provides a composition comprising particles of buprenorphine encapsulated by PEG, wherein the buprenorphine content ranges from about 0.2% to about 4%. In other embodiments, the composition further comprises a surfactant, such as, e.g., a polysorbate surfactant.

30 [0005] In other embodiments, the invention provides pharmaceutical compositions comprising such encapsulated compositions. Such pharmaceutical compositions may, in some embodiments, further comprise at least one excipient. In other embodiments, such pharmaceutical compositions may further comprise a second compound such as, e.g., a second drug, including, e.g., an opioid receptor antagonist, an anti-inflammatory drug, or an analgesic. For example, in some embodiments, the invention provides a pharmaceutical composition comprising a composition comprising the PEG-encapsulated buprenorphine composition described above. In certain embodiments, the pharmaceutical composition further comprises naloxone.

35 [0006] In some embodiments, the invention provides a first method of making a composition comprising particles of a poorly soluble drug encapsulated by a stabilizer, the method comprising:

40

blending a poorly soluble drug together with a stabilizer to form a mixture;

processing said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm; and

45

processing said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

50 [0007] In some embodiments, the invention provides a second method of making a composition comprising particles of a poorly soluble drug encapsulated by a stabilizer, the method comprising:

blending a poorly soluble drug together with a stabilizer to form a mixture;

heating said mixture to a temperature sufficient for extrusion of the mixture;

55

extruding said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm;

cooling said coarse particles; and

processing said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

5 [0008] In further embodiments, the invention provides a method of making a pharmaceutical composition, comprising the first or second method described above, and further comprising:

formulating the fine particles.

10 [0009] In further embodiments, the invention provides a method of making a pharmaceutical composition, comprising the first or second method described above, and further comprising:

mixing the fine particles with at least one excipient to form a second mixture; and

15 formulating the second mixture.

20 [0010] In yet other embodiments, the invention provides a method of treating pain, comprising administering the pharmaceutical composition described above to a patient in need thereof. In other embodiments, the invention provides a method of treating opiate addiction, comprising administering the pharmaceutical composition described above to a patient in need thereof.

DETAILED DESCRIPTION

I. PARTICULATE DELIVERY SYSTEMS

25 [0011] In some embodiments, the invention provides a composition (also referred to as a particulate delivery system or PDS) comprising particles of a poorly soluble drug encapsulated by a stabilizer. In some embodiments, those particles are fine particles, and have a diameter of less than 3 mm, less than 2 mm, less than 600 μm , less than 500 μm , or less than 300 μm . In some embodiments, the fine particles have an average diameter ranging from about 0.1 mm (100 μm) to about 3 mm. For example, the particles may have a diameter of less than 2.06 mm (corresponding to a 10 mesh sieve), less than 1.68 mm (corresponding to a 12 mesh sieve), less than 1.40 mm (corresponding to a 14 mesh sieve), less than 1.20 mm (corresponding to a 16 mesh sieve), less than 1.00 mm (corresponding to an 18 mesh sieve), less than 0.853 mm (corresponding to a 20 mesh sieve), less than 0.710 mm (corresponding to a 25 mesh sieve), less than 0.599 mm (corresponding to a 30 mesh sieve), or less than 0.500 mm (corresponding to a 35 mesh sieve). In other embodiments, the particles may have a diameter of less than 300 μm , and may be able to pass through a 50 mesh sieve.

30 [0012] As used herein, the term drug encompasses the corresponding salts, hydrates, solvates, prodrugs, and complexes of the drug. Thus, the drug may be present as, e.g., a free base, a salt, a hydrate, a prodrug, a solvate (including a mixed solvate), or a complex (such as a pharmaceutically acceptable complex, and/or a complex with a polymer).

35 [0013] As used herein, the terms poorly soluble drug, drug having poor solubility, and the like refer to a drug (in its neutral (i.e., uncharged) state) having a water solubility at neutral pH of less than 10 mg/ml. In some embodiments, the drug (in its neutral state) has a water solubility at neutral pH of less than 5 mg/ml. In other embodiments, the drug (in its neutral state) has a water solubility at neutral pH of less than 1 mg/ml. For example, buprenorphine base (i.e., uncharged buprenorphine) has a solubility at neutral pH of <1 mg/ml (whereas the corresponding hydrochloride salt has a solubility at neutral pH of 17 mg/ml). Thus, as used herein, buprenorphine (including buprenorphine base and its salts, hydrates, solvates, complexes, etc.) is a poorly soluble drug. Similarly, morphine base (i.e., uncharged morphine) has a solubility at neutral pH of <1 mg/ml (whereas the corresponding sulfate has a solubility at neutral pH of 64 mg/ml). Thus, as used herein, morphine (including morphine base and its salts, hydrates, solvates, complexes, etc.) is a poorly soluble drug. In a third example, oxycodone base (i.e., uncharged oxycodone) has a solubility at neutral pH of <1 mg/ml (whereas the corresponding hydrochloride has a solubility at neutral pH of 100 mg/ml). Thus, as used herein, oxycodone (including oxycodone base and its salts, hydrates, solvates, complexes, etc.) is a poorly soluble drug.

40 [0014] In certain embodiments, the poorly soluble drug is chosen from opioids (including opiates). Opioids include naturally-occurring, synthetic, and semisynthetic opioids, such as, e.g., alfentanil, allylprodine, alphaprodine, anileridine, benzylmorphine, bezitramide, buprenorphine, butorphanol, clonidine, clonitazene, codeine, cyclazocine, desomorphine, dextromoramide, dezocine, diamorphine, dihydrocodeine, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, eptazocine, ethoheptazine, ethylmethylthiambutene, ethylmorphine, etonitazine, fentanyl, heroin, hydrocodone, hydromorphone, hydroxypethidine, isomethadone, ketobemidone, levallorphan, levomethadyl, levorphanol, levophenacylmorphan, lofentanil, meperidine, meptazinol, metazocine, methadone, metopon, morphine, myrophine, nalbuphine, naloxone, naltrexone, narceine, nicomorphine, norlevorphanol, normethadone, nalorphine, normorphine, norpipanone, opium, oxycodone, oxymorphone, papaveretum, pentazocine, phena-

doxone, phenomorphan, phenazocine, phenoperidine, piminodine, piritramide, proheptazine, promedol, properidine, propiram, propoxyphene, remifentanil, sulfentanil, tramadol, and tilidine. For example, in some embodiments, the opioid may be chosen from, e.g., buprenorphine, codeine, fentanyl, hydrocodone, hydromorphone, morphine, methylnaltrexone, nalbuphine, nalmefene, oxymorphone, oxycodone, pethidine, and tramadol.

5 [0015] In some embodiments, the PDS may further comprise an additional compound, such as an additional drug. The additional drug may be chosen from, e.g., opioid receptor antagonists (including μ -receptor antagonists), opioid receptor agonists (including μ -receptor agonists), mixed μ -agonists/ μ -antagonists, anti-inflammatory drugs, and analgesics. In some embodiments, the second drug is an opioid receptor antagonist, such as, e.g., the μ -opioid receptor antagonist naloxone, including naloxone·HCl (naloxone hydrochloride). In some embodiments where the poorly soluble 10 drug is an opioid analgesic, the opioid receptor antagonist is added to deter abuse of the opioid analgesic.

10 [0016] The poorly soluble drug may be present in an amount ranging from about <1% to about 90% of the PDS by mass. For example, the poorly soluble drug may be present in an amount ranging from about 0.01% to about 90%, about 0.01% to about 10%, about 0.2 to about 5%, about <1% to about 10%, about 0.01% to about 10%, about 0.1 % to about 10%, about 0.01% to about 5%, about 0.1 % to about 5%, about 0.1 % to about 3%, about <1% to about 50%, about 15 <1% to about 30%, about <1% to about 80%, about 5% to about 90%, about 10% to about 95%, or about 0.1 to about 5% of the PDS, by mass. In some embodiments, the poorly soluble drug content may be about 0.5% by mass.

15 [0017] In some embodiments, the stabilizer is a polymer, such as, e.g., a water-soluble polymer, a polymer of neutral charge, or a water-soluble polymer of neutral charge. In some embodiments, the stabilizer is biodegradable. In some 20 embodiments, the stabilizer is bioerodable. In some embodiments, the stabilizer may be considered by the FDA to be generally regarded as safe (GRAS).

20 [0018] In some embodiments, the stabilizer is a polymer chosen from polyethylene oxide (also known as polyethylene glycol or PEG), polypropylene oxide, or copolymers thereof. In some embodiments, the stabilizer is a water-soluble 25 polymer of neutral charge chosen from polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone (PVP), block copolymers of ethylene oxide and propylene oxide such as, e.g., poloxamers, and tetrafunctional block copolymers derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine.

30 [0019] In some embodiments, the stabilizer may have an average molecular weight of about, e.g., 500, 1000, 2000, 3000, 3350, 3500, 4000, 4500, 5000, 6000, 8000, 10,000, or 100,000 Da, or an average molecular weight ranging from, e.g., about 100 Da to about 100,000 Da, about 100 Da to about 6,000 Da, about 500 Da to about 5000 Da, about 1000 Da to about 4000 Da, about 2000 Da to about 4000 Da, about 2000 Da to about 6000 Da, about 1000 Da to about 10,000 Da, or about 3000 Da to about 4000 Da.

35 [0020] For example, the stabilizer may be a PEG, such as a PEG of average molecular weight of about, e.g., 500, 1000, 2000, 3000, 3350, 3500, 4000, 4500, 5000, 6000, 8000, 10,000 or 100,000 Daltons. In exemplary embodiments, the stabilizer is PEG having an average molecular weight of 3350 Daltons (i.e., PEG 3350). In other embodiments, the 40 stabilizer may be a PEG having an average molecular weight ranging from, e.g., about 100 Da to about 100,000 Da, about 100 Da to about 6,000 Da, about 500 Da to about 5000 Da, about 1000 Da to about 4000 Da, about 2000 Da to about 6000 Da, about 1000 Da to about 10,000 Da, or about 3000 Da to about 4000 Da.

45 [0021] In some embodiments, the composition further comprises at least one excipient, such as, e.g., a surfactant, surface stabilizer, or other enhancer. For example, in some embodiments, the composition further comprises at least one surfactant, which may be a nonionic surfactant such as, e.g., a polysorbate surfactant. In exemplary embodiments, the composition further comprises polysorbate 20 (Tween 20). In other embodiments, the composition further comprises an emulsifier, such as, e.g., a phospholipid, propylene glycol, polysorbate, poloxamer, glyceryl monostearate, or other 50 pharmaceutical emulsifier.

50 [0022] In certain embodiments, the composition comprises an opioid, a PEG, and a surfactant. In some embodiments, the invention provides a composition comprising particles of buprenorphine (such as, e.g., buprenorphine·HCl) encapsulated by PEG (which, in some embodiments, may have an average molecular weight ranging from about 500 Daltons to about 10,000 Daltons), wherein the buprenorphine content ranges from about 0.2% to about 4%. In further embodiments, the particles have an average diameter of less than 2 mm. In some embodiments, the composition further comprises a surfactant. In certain embodiments, the PDS comprises buprenorphine, PEG 3350, and Tween 20.

55 II. METHODS OF MAKING PDS

[0023] In some embodiments, the invention provides a first method of making a composition (such as those described in Section I) comprising particles of a poorly soluble drug encapsulated by a stabilizer, the method comprising:

55 blending a poorly soluble drug together with a stabilizer to form a mixture;

processing (e.g., by mixing or granulating) said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm; and

processing (e.g., by milling, grinding, or crushing) said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

5 [0024] In certain embodiments, the fine particles have an average diameter ranging from about 0.1 mm (100 μm) to about 3 mm. For example, the particles may have a diameter of less than 2.06 mm (corresponding to a 10 mesh sieve), less than 1.68 mm (corresponding to a 12 mesh sieve), less than 1.40 mm (corresponding to a 14 mesh sieve), less than 1.20 mm (corresponding to a 16 mesh sieve), less than 1.00 mm (corresponding to an 18 mesh sieve), less than 0.853 mm (corresponding to a 20 mesh sieve), less than 0.710 mm (corresponding to a 25 mesh sieve), less than 0.599 mm (corresponding to a 30 mesh sieve), or less than 0.500 mm (corresponding to a 35 mesh sieve). In other embodiments, 10 the particles may have a diameter of less than 300 μm , and may be able to pass through a 50 mesh sieve. In certain embodiments, the particles have a diameter of 0.6 mm or less.

[0025] In certain embodiments, the stabilizer is heated prior to blending with the poorly soluble drug.

15 [0026] In other embodiments, the invention provides a second method of making a composition (such as those described in Section I) comprising particles of a poorly soluble drug encapsulated by a stabilizer. The second method uses a relatively continuous manufacturing process, which may be advantageous in certain circumstances. The second method comprises:

blending a poorly soluble drug together with a stabilizer to form a mixture;

20 heating said mixture to a temperature sufficient for extrusion of the mixture;

extruding said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm;

cooling said coarse particles; and

25 processing (e.g., by milling, grinding, or crushing) said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

30 [0027] In certain embodiments, the fine particles have an average diameter ranging from about 0.1 mm (100 μm) to about 3 mm. For example, the particles may have a diameter of less than 2.06 mm (corresponding to a 10 mesh sieve), less than 1.68 mm (corresponding to a 12 mesh sieve), less than 1.40 mm (corresponding to a 14 mesh sieve), less than 1.20 mm (corresponding to a 16 mesh sieve), less than 1.00 mm (corresponding to an 18 mesh sieve), less than 0.853 mm (corresponding to a 20 mesh sieve), less than 0.710 mm (corresponding to a 25 mesh sieve), less than 0.599 mm (corresponding to a 30 mesh sieve), or less than 0.500 mm (corresponding to a 35 mesh sieve). In other embodiments, 35 the particles may have a diameter of less than 300 μm , and may be able to pass through a 50 mesh sieve. In certain embodiments, the particles have a diameter of 0.6 mm or less.

[0028] In certain embodiments, the stabilizer is heated prior to blending with the poorly soluble drug.

III. PHARMACEUTICAL COMPOSITIONS (FINAL DOSAGE FORMS)

40 [0029] In some embodiments, the invention provides pharmaceutical compositions (sometimes referred to as final dosage forms or FDF) comprising the compositions described in Section I above.

[0030] The poorly soluble drug may be present in the pharmaceutical composition in an amount ranging from about <1% to about 90% by mass. For example, the poorly soluble drug may be present in an amount ranging from about 45 0.01% to about 90%, about 0.01 % to about 10%, about 0.2 to about 5%, about <1% to about 10%, about 0.01% to about 10%, about 0.1% to about 10%, about 0.01% to about 5%, about 0.1 % to about 5%, about 0.1 % to about 3%, about <1% to about 50%, about <1% to about 30%, about <1% to about 80%, about 5% to about 90%, about 10% to about 95%, or about 0.1 to about 5% of the pharmaceutical composition by mass. In some embodiments, the poorly soluble drug content may be about 0.5% by mass.

50 [0031] In some embodiments, the pharmaceutical compositions further comprise a second compound, such as a second drug. The second drug may be chosen from, e.g., opioid receptor antagonists (including μ -receptor antagonists), opioid receptor agonists (including μ -receptor agonists), mixed μ -agonists/ μ -antagonists, anti-inflammatory drugs, and analgesics. In some embodiments, the second drug is an opioid receptor antagonist, such as, e.g., the μ -opioid receptor antagonist naloxone, including naloxone-HCl (naloxone hydrochloride). In some embodiments where the poorly soluble 55 drug is an opioid analgesic, the opioid receptor antagonist is added to deter abuse of the opioid analgesic. The resulting compositions may have reduced potential for abuse of the opioid, relative to compositions that do not comprise an opioid receptor antagonist.

[0032] In some embodiments, the pharmaceutical compositions further comprise at least one excipient (such as, e.g.,

a water-soluble polymer, surfactant, and/or enhancer), such as a pharmaceutically acceptable excipient. Examples of pharmaceutically acceptable excipients are described in *Remington's Pharmaceutical Sciences* by E.W. Martin, and include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. In some embodiments, the pharmaceutical compositions also contain pH buffering reagents, and wetting or emulsifying agents.

[0033] The pharmaceutical compositions may, in some embodiments, be formulated for oral administration, for example as tablets, capsules, or other oral dosage forms. Such oral dosage forms may be prepared by conventional means. The pharmaceutical composition can also be prepared as a liquid, for example as a syrup or a suspension. The liquid can include suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats), emulsifying agents (lecithin or acacia), non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils), and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations can also include flavoring, coloring and sweetening agents. Alternatively, the composition can be presented as a dry product for constitution with water or another suitable vehicle.

[0034] For buccal and sublingual administration, the composition may take the form of tablets or lozenges according to conventional protocols.

[0035] For administration by oral inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray from a pressurized pack or nebulizer (e.g., in PBS), with a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoromethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount.

[0036] Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0036] The pharmaceutical composition can also be formulated for parenteral administration (including, e.g., intravenous or intramuscular administration) by bolus injection. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multidose containers with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, such as, e.g., pyrogen free water.

[0037] The pharmaceutical composition can also be formulated for rectal administration as a suppository or retention enema, e.g., containing conventional suppository bases such as PEG, cocoa butter or other glycerides.

[0038] In some embodiments, the pharmaceutical compositions described herein provide improved dissolution of the poorly soluble drug, relative to the unencapsulated poorly soluble drug, and/or to another dosage form (such as, e.g., a more invasive dosage form). For example, dissolution may be increased by, e.g., at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 60%, 70%, 80%, 90%, 93%, 95%, 96%, 97%, 98%, 99%, 100%, 110%, 120%, 130%, 140%, 150%, or 200%, or by, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100, or 1000 fold, as measured by a Vankel tablet dissolution apparatus approved by the United States Pharmacopeia.

[0039] In some embodiments, the pharmaceutical compositions described herein provide improved oral bioavailability of the poorly soluble drug, relative to the unencapsulated poorly soluble drug, and/or to another dosage form (such as, e.g., a more invasive dosage form). For example, absorption may be increased by, e.g., at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 60%, 70%, 80%, 90%, 93%, 95%, 96%, 97%, 98%, 99%, 100%, 110%, 120%, 130%, 140%, 150%, or 200%, or by, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100, or 1000 fold, as measured by, e.g., *in vivo* pharmacokinetic studies in a preclinical animal model or human clinical evaluation.

[0040] In some embodiments, the pharmaceutical compositions described herein are immediate-release formulations. In such embodiments, the pharmaceutical compositions provide a more rapid onset of action of the poorly soluble drug, relative to the unencapsulated poorly soluble drug, and/or to another dosage form (such as, e.g., a more invasive dosage form). For example, the onset of action may be shortened by, e.g., at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 60%, 70%, 80%, 90%, 93%, 95%, 96%, 97%, 98%, 99%, 100%, 110%, 120%, 130%, 140%, 150%, or 200%, or by, e.g., at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 100, or 1000 fold, as measured by, e.g., *in vivo* pharmacokinetic studies in a preclinical animal model or human clinical evaluation.

[0041] In other embodiments, the pharmaceutical compositions described herein are controlled-release formulations. In such embodiments, the pharmaceutical compositions described herein provide a more rapid onset of action of the poorly soluble drug.

[0042] In some embodiments, the pharmaceutical compositions described herein have reduced absorption variability, relative to the unencapsulated poorly soluble drug, and/or to another dosage form (such as, e.g., a more invasive dosage form).

[0043] In some embodiments, the pharmaceutical compositions described herein are associated with improved patient compliance, relative to another pharmaceutical composition comprising the same poorly soluble drug (which may be in another dosage form, such as, e.g., a more invasive dosage form).

IV. METHODS OF MAKING PHARMACEUTICAL COMPOSITIONS

[0044] In further embodiments, the invention provides a method of making a pharmaceutical composition, comprising the first, second, or third method described in Section II above, and further comprising:

5

formulating the fine particles.

[0045] In certain embodiments, the fine particles are formulated into unit doses.

[0046] In embodiments in which the pharmaceutical compositions comprise at least one excipient, the invention also provides a method of making a pharmaceutical composition, comprising the first, second, or third method described in Section II above, and further comprising:

mixing the fine particles with at least one excipient to form a second mixture; and

15

formulating the second mixture.

[0047] In certain embodiments, the fine particles are formulated into unit doses.

V. METHODS OF TREATMENT

20

[0048] The pharmaceutical compositions described herein are useful to treat any disease or condition for which administration of a corresponding hydrophobic drug is desirable. For example, compositions comprising opioids are useful for the treatment of pain. The terms "treat," "treatment," and "treating" refer to (1) a reduction in severity or duration of a disease or condition, (2) the amelioration of one or more symptoms associated with a disease or condition without necessarily curing the disease or condition, or (3) the prevention of a disease or condition. Suitable subjects include, e.g., humans and other mammals, such as, e.g., mice, rats, dogs, and non-human primates.

25

[0049] In certain embodiments, the invention provides a method of treating pain, comprising administering a pharmaceutical composition described in Section III to a subject in need thereof. For example, in such embodiments, the poorly soluble drug may be chosen from, e.g., opioids, including, e.g., buprenorphine, codeine, fentanyl, hydrocodone, hydro-morphine, morphine, methylnaltrexone, nalbuphine, nalmefene, oxymorphone, oxycodone, pethidine, and tramadol. Moreover, in such embodiments, the pharmaceutical composition may further comprise an opioid receptor antagonist, such as, e.g., naloxone or naltrexone, to deter abuse of the opioid analgesic.

30

[0050] In other embodiments, the invention provides a method of treating opiate addiction, comprising administering a pharmaceutical composition described in Section III to a subject in need thereof. For example, in such embodiments, the poorly soluble drug may be chosen from, e.g., methadone, naloxone, and naltrexone.

[0051] The following examples are intended to be purely exemplary of the present invention.

EXAMPLES

40

Example 1: Preparation of Buprenorphine-HCl PDS

[0052] A solution of 1 kg of autoclaved polyethylene glycol (Dow, PEG 3350, NF, FCC) and polysorbate (Tween 20, FCC), at a 10:1 mass ratio, cooled to less than 60 °C, is aseptically mixed with 20 g of buprenorphine-HCl (Johnson Matthey, milled USP grade) powder. The suspension (which contains 2% buprenorphine) is then blended in a paddle mixer and further agitated using low-frequency sonication (as described in U.S. Pat. Appl. Pub. No. 2005/0175707) to form buprenorphine-HCl particles encapsulated in a PEG/Tween matrix. The powder is then frozen at -20°C for at least 2 hours and then ground into a fine powder using a Retsch knife mill. Particles smaller than 600 µm are separated by sieving (30 mesh). Buprenorphine content is confirmed using an HPLC method.

50

Example 2: Preparation of Immediate-Release Oral Buprenorphine-HCl Capsules

[0053] An immediate-release oral dosage form (gelatin capsules) containing the buprenorphine-HCl particles prepared in Example 1, as well as naloxone-HCl as an abuse-deterrant second drug, is prepared as described below. The PDS prepared in Example 1 is dry mixed with naloxone (as naloxone-HCl dihydrate, USP) and additional PEG 3350 for bulking to achieve the correct capsule fill weight (400-500 mg) to achieve the desired dose. Clear gelatin #1 capsules are then filled with the mixture in a Fast-CAP Filling machine to yield capsules containing 2.0 ± 0.2 mg buprenorphine (with respect to the free base) and 0.5 ± 0.05 mg naloxone (with respect to the free base). The capsules thus contain buprenorphine and naloxone at a ratio of 4:1, with respect to the free bases.

Example 3: Dissolution of Immediate-Release Oral Buprenorphine-HCl Capsules

5 [0054] The in vivo dissolution rate of the fast-release formulation prepared in Example 2 was measured by USP Paddle Method 2 at 50 or 100 rpm in 900 ml of 0.1 N HCl acidic buffer (pH between 1.6 and 3) at 37°C. It was found that greater than 75% (by weight) of the therapeutically active agent was released after 45 minutes.

Example 4: Single-Dose Pharmacokinetic Study in Rats

10 [0055] The composition prepared in Example 2 (prior to loading into gelatin capsules) was administered via oral gavage once daily for 14 consecutive days to male and female F-344 rats at a low (0.03 mg/kg) or high (0.5 mg/kg) dose (n=10 each). Plasma samples collected up to 8 hours from single-dose pharmacokinetic (PK) studies in male rats (non-GLP) were compared to 14-day samples using a validated extraction and LC-MS-MS analytical method.

15 [0056] Values for the single-dose PK study were 0.53 and 4.07 ng/ml for male rats administered 0.03 and 0.5 mg/kg, respectively (n=3-5), observed at the first 15 minute collection time. The concentration-time profiles declined from C_{max} in mono-or multi-exponential relationships. AUC values for the single-dose PK study were 2.51 and 10.33 ng·hr/ml for male rats administered 0.03 and 0.5 mg/kg, respectively (n=3-5). At day 14 of the repeat dose toxicity study for rats administered 0.03 and 0.5 mg/kg, respectively (n=4 each), C_{max} values were 1.57 and 4.11 ng/ml for males and 1.85 and 4.12 ng/ml for females, also observed at the 30 minute collection time. C_{max} values were similar for males and females based on 30 minute and 8 hour time points from 14-day plasma samples analyzed. Norbuprenorphine, the main metabolite of buprenorphine, and naloxone plasma concentrations were below the limit of quantitation.

Example 5: Preparation of Pharmaceutical Compositions Comprising Oxycodone

25 [0057] Pharmaceutical compositions comprising oxycodone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Oxycodone content	0.1-200 mg
PEG 1500 content	100-1000 mg

30

Example 6: Preparation of Controlled-Release Pharmaceutical Compositions Comprising Oxycodone**Comprising Oxycodone**

35 [0058] Controlled-release pharmaceutical compositions comprising oxycodone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Oxycodone content	0.1-200 mg
PVP K-30 content	100-1000 mg

40

Example 7: Preparation of Pharmaceutical Compositions Comprising Morphine

45 [0059] Pharmaceutical compositions comprising morphine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Morphine content	0.1-100 mg
Poloxamer 407 content	100-1000 mg

50

Example 8: Preparation of Controlled-Release Pharmaceutical Compositions Comprising Morphine

55 [0060] Controlled-release pharmaceutical compositions comprising morphine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Morphine content	0.1-100 mg
------------------	------------

(continued)

PEG 8000 content	100-1000 mg
------------------	-------------

5 **Example 9: Preparation of Pharmaceutical Compositions Comprising Hydrocodone**

[0061] Pharmaceutical compositions comprising hydrocodone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

10	Hydrocodone content	0.1-100 mg
	PEG 3350 content	100-1000 mg

15 **Example 10: Preparation of Pharmaceutical Compositions Comprising Dihydrocodone**

[0062] Pharmaceutical compositions comprising dihydrocodone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

20	Dihydrocodone content	0.1-100 mg
	PEG 3350 content	100-1000 mg

25 **Example 11: Preparation of Pharmaceutical Compositions Comprising Codeine**

[0063] Pharmaceutical compositions comprising codeine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

30	Codeine content	0.1-100 mg
	Poloxamer 407 content	100-1000 mg

Example 12: Preparation of Pharmaceutical Compositions Comprising Meperidine

35 [0064] Pharmaceutical compositions comprising meperidine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

40	Meperidine content	0.1-500 mg
	PEG 3350 content	100-1000 mg

Example 13: Preparation of Pharmaceutical Compositions Comprising Propoxyphene

45 [0065] Pharmaceutical compositions comprising propoxyphene may be prepared according to the methods of the invention. The compositions may have the following characteristics:

50	Propoxyphene content	0.1-500 mg
	PEG 3350 content	100-1000 mg

Example 14: Preparation of Pharmaceutical Compositions Comprising Levorphanol

55 [0066] Pharmaceutical compositions comprising levorphanol may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Levorphanol content	0.1-100 mg
---------------------	------------

(continued)

Poloxamer 407 content	100-1000 mg
-----------------------	-------------

5 **Example 15: Preparation of Pharmaceutical Compositions Comprising Oxymorphone**

[0067] Pharmaceutical compositions comprising oxymorphone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Oxymorphone content	0.1-200 mg
PEG 3350 content	100-1000 mg

15 **Example 16: Preparation of Pharmaceutical Compositions Comprising Hydromorphone**

[0068] Pharmaceutical compositions comprising hydromorphone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Hydromorphone content	0.1-100 mg
Poloxamer 407 content	100-1000 mg

25 **Example 17: Preparation of Pharmaceutical Compositions Comprising Fentanyl**

[0069] Pharmaceutical compositions comprising hydromorphone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Fentanyl content	0.1-500 mg
PEG 3350 content	100-1000 mg

30 **Example 18: Preparation of Pharmaceutical Compositions Comprising Alfentanyl**

[0070] Pharmaceutical compositions comprising alfentanyl may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Alfentanyl content	0.01-50 µg
PEG 3350 content	100-1000 mg

35 **Example 19: Preparation of Pharmaceutical Compositions Comprising Sufentanyl**

[0071] Pharmaceutical compositions comprising sufentanyl may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Sufentanyl content	0.01-500 µg
PEG 3350 content	100-1000 mg

40 **Example 20: Preparation of Pharmaceutical Compositions Comprising Remifentanyl**

[0072] Pharmaceutical compositions comprising remifentanyl may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Remifentanyl content	0.01-500 µg
----------------------	-------------

(continued)

PEG 3350 content	100-1000 mg
------------------	-------------

5 **Example 21: Preparation of Pharmaceutical Compositions Comprising Levomethadyl**

[0073] Pharmaceutical compositions comprising levomethadyl may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Levomethadyl content	0.01-200 mg
Poloxamer 407 content	100-1000 mg

15 **Example 22: Preparation of Pharmaceutical Compositions Comprising Methadone**

[0074] Pharmaceutical compositions comprising methadone may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Methadone content	0.01-200 mg
PEG 3350 content	100-1000 mg

25 **Example 23: Preparation of Pharmaceutical Compositions Comprising Butorphanol**

[0075] Pharmaceutical compositions comprising butorphanol may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Butorphanol content	0.1-200 mg
PEG 3350 content	100-1000 mg

30 **Example 24: Preparation of Pharmaceutical Compositions Comprising Dezocine**

[0076] Pharmaceutical compositions comprising dezocine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Dezocine content	0.1-200 mg
Poloxamer 407 content	100-1000 mg

35 **Example 25: Preparation of Pharmaceutical Compositions Comprising Nalbuphine**

[0077] Pharmaceutical compositions comprising nalbuphine may be prepared according to the methods of the invention. The compositions may have the following characteristics:

Nalbuphine content	0.1-200 mg
PEG 3350 content	100-1000 mg

[0078] Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following items.

55 1. A composition comprising particles of a poorly soluble drug encapsulated by a stabilizer.

2. The composition of item 1, wherein the particles have an average diameter of less than 2 mm.

3. The composition of item 2, wherein the particles have an average diameter of less than 500 μm .
4. The composition of item 3, wherein the particles have an average diameter of less than 300 μm .
5. The composition of item 1, wherein the poorly soluble drug is an opioid.
6. The composition of item 5, wherein the opioid is chosen from buprenorphine, codeine, fentanyl, hydrocodone, hydromorphone, morphine, methylnaltrexone, nalbuphine, nalmefene, oxymorphone, oxycodone, pethidine, and tramadol.
- 10 7. The composition of **item 6**, wherein the opioid is buprenorphine.
8. The composition of **item 7**, wherein the buprenorphine is a buprenorphine salt.
- 15 9. The composition of **item 8**, wherein the buprenorphine salt is buprenorphine-HCl.
10. The composition of **item 1**, wherein the stabilizer is a polymer.
11. The composition of **item 10**, wherein the stabilizer is a water-soluble polymer.
- 20 12. The composition of **item 10**, wherein the stabilizer is a polymer of neutral charge.
13. The composition of **item 10**, wherein the stabilizer is a water-soluble polymer of neutral charge.
- 25 14. The composition of **item 13**, wherein the water-soluble polymer of neutral charge is chosen from polyethylene glycol (PEG), polyvinyl alcohol, polyvinylpyrrolidone, block copolymers of ethylene oxide and propylene oxide, and tetrafunctional block copolymers derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine.
- 30 15. The composition of **item 14**, wherein the polymer is PEG.
16. The composition of **item 15**, wherein the PEG has an average molecular weight ranging from about 100 Daltons to about 100,000 Daltons.
- 35 17. The composition of **item 16**, wherein the PEG is PEG 3350.
18. The composition of **item 1**, wherein the poorly soluble drug is present in an amount ranging from about 0.01 % to about 90% by mass.
- 40 19. The composition of **item 18**, wherein the poorly soluble drug is present in an amount ranging from about 0.01 % to about 10% by mass.
20. The composition of **item 19**, wherein the poorly soluble drug is present in an amount ranging from about 0.2% to about 5% by mass.
- 45 21. The composition of **item 20**, wherein the poorly soluble drug content is about 0.5% by mass.
22. The composition of **item 1**, further comprising at least one excipient.
- 50 23. The composition of **item 22**, wherein the at least one excipient is a surfactant.
24. The composition of **item 23**, wherein the surfactant is a nonionic surfactant.
25. The composition of **item 24**, wherein the nonionic surfactant is a polysorbate surfactant.
- 55 26. The composition of **item 25**, wherein the polysorbate surfactant is Tween 20.
27. A pharmaceutical composition comprising the composition of item 1.

28. The pharmaceutical composition of item 27, further comprising a second compound.

29. The pharmaceutical composition of **item 28**, wherein the second compound is a second drug.

5 30. The pharmaceutical composition of **item 29**, wherein the second drug is chosen from opioid receptor antagonists, anti-inflammatory drugs, and analgesics.

31. The pharmaceutical composition of **item 30**, wherein the second drug is an opioid receptor antagonist.

10 32. The pharmaceutical composition of **item 31**, wherein the second drug is naloxone.

33. The pharmaceutical composition of **item 32**, wherein the naloxone is a naloxone salt.

15 34. The pharmaceutical composition of **item 33**, wherein the naloxone salt is naloxone-HCl.

35. The pharmaceutical composition of **item 27**, wherein the pharmaceutical composition further comprises at least one excipient.

20 36. The pharmaceutical composition of **item 35**, wherein the at least one excipient is chosen from polyethylene glycol (PEG), polyvinyl alcohol, polyvinylpyrrolidone, block copolymers of ethylene oxide and propylene oxide, and tetrafunctional block copolymers derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine.

25 37. A pharmaceutical composition according to item 27, wherein the pharmaceutical composition is formulated for oral administration.

38. A method of making the composition of **item 1**, the method comprising:

30 blending an opioid together with a stabilizer to form a mixture;

processing said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm; and

35 processing said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

39. A method of making the composition of **item 1**, the method comprising:

40 blending an opioid together with a stabilizer to form a mixture;

heating said mixture to a temperature sufficient for extrusion of the mixture;

45 extruding said mixture to form coarse particles having an average diameter ranging from about 0.1 mm to about 5 mm;

cooling said coarse particles; and

processing said coarse particles to form fine particles having an average diameter ranging from about 0.1 mm to about 3 mm.

50 40. A method of making a pharmaceutical composition, comprising the method of item 38, and further comprising:

formulating the fine particles.

55 41. A method of making a pharmaceutical composition, comprising the method of item 39, and further comprising:

formulating the fine particles.

42. A method of treating pain, comprising administering the pharmaceutical composition of **item 27** to a patient in need thereof.

43. The method of item 42, wherein the poorly soluble drug is an opioid.

5 44. A method of treating opiate addiction, comprising administering the pharmaceutical composition of item 27 to a patient in need thereof.

10 45. The method of **item 44**, wherein the poorly soluble drug is chosen from methadone, naloxone, and naltrexone.

15 46. A composition comprising particles of buprenorphine encapsulated by PEG, wherein the buprenorphine content ranges from about 0.2% to about 4%.

47. The composition of **item 46**, wherein the particles have an average diameter of less than 2 mm.

15 48. The composition of **item 47**, wherein the buprenorphine is buprenorphine·HCl.

20 49. The composition of **item 46**, wherein the PEG has an average molecular weight ranging from about 500 Daltons to about 10,000 Daltons.

50. The composition of **item 49**, wherein the PEG is PEG 3350.

51. The composition of **item 46**, further comprising a surfactant.

25 52. A pharmaceutical composition comprising the composition of item 46 and naloxone.

53. The pharmaceutical composition of **item 52**, wherein the composition is formulated for oral administration.

30 **Claims**

1. A particle delivery system (PDS) comprising dry particles having an average diameter of less than about 2 mm, wherein the dry particles comprise a poorly soluble drug and a polyethylene glycol (PEG), wherein the amount of the poorly soluble drug present in the dry particles ranges from about 0.2% to about 5% by mass of the dry particles,

35 the poorly soluble drug is encapsulated by the PEG, and

the poorly soluble drug comprises at least one opioid selected from codeine, fentanyl, hydrocodone, hydromorphone, morphine, methylnaltrexone, nalbuphine, nalmefene, oxymorphone, oxycodone, pethidine, and tramadol.

40 2. The PDS of claim 1, wherein the PEG has an average molecular weight ranging from about 100 Daltons to about 100,000 Daltons.

3. The PDS of claim 2, wherein the PEG is PEG 3350.

45 4. The PDS of claim 1, further comprising a surfactant.

5. The PDS of claim 4, wherein the surfactant is a nonionic surfactant.

6. A pharmaceutical composition comprising the PDS of claim 1 and at least one second compound.

50 7. The pharmaceutical composition of claim 6, wherein the composition is formulated for oral administration.

8. The pharmaceutical composition of claim 7, wherein the at least one second compound is a drug chosen from opioid receptor antagonists, anti-inflammatory drugs, and analgesics.

55 9. The pharmaceutical composition of claim 8, wherein the drug comprises an opioid receptor antagonist.

10. The pharmaceutical composition of claim 9, wherein the drug comprises naloxone or a pharmaceutically acceptable

salt thereof.

11. The pharmaceutical composition of claim 10, wherein a pharmaceutically acceptable salt of naloxone is naloxone HCl.

5

10

15

20

25

30

35

40

45

50

55



EUROPEAN SEARCH REPORT

Application Number

EP 16 16 3685

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	X LEUNER C ET AL: "Improving drug solubility for oral delivery using solid dispersions", EUROPEAN JOURNAL OF PHARMACEUTICS AND BIOPHARMACEUTICS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 50, no. 1, 3 July 2000 (2000-07-03), pages 47-60, XP004257179, ISSN: 0939-6411, DOI: 10.1016/S0939-6411(00)00076-X * abstract * * page 52, column 1, paragraph 1 - page 53, column 1, paragraph 3 * -----	1-11	INV. A61K9/14 A61K9/50 A61K31/485 A61K31/135 A61K31/4468
15	X CA 987 588 A (UNIV CALIFORNIA) 20 April 1976 (1976-04-20) * claims 1-6 * * page 1, paragraph 1-2 * * examples I, VI-IX * -----	1-11	
20	X WO 2005/074890 A1 (PHARMACIA ITALIA SPA [IT]; CIOCCA CRISTINA [IT]; MARTINI ALESSANDRO [I]) 18 August 2005 (2005-08-18) * claims 1, 4, 12 * * page 1, line 19 - page 2, line 2 * * page 6, lines 5-11 * -----	1-11	TECHNICAL FIELDS SEARCHED (IPC)
25			A61K
30			
35			
40			
45			
50	1 The present search report has been drawn up for all claims		
55	Place of search	Date of completion of the search	Examiner
	Munich	25 May 2016	Peris Antoli, Berta
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 16 16 3685

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

25-05-2016

10

	Patent document cited in search report		Publication date		Patent family member(s)		Publication date
	CA 987588	A	20-04-1976	NONE			
15	WO 2005074890	A1	18-08-2005	BR	PI0418427 A	12-06-2007	
				CA	2552925 A1	18-08-2005	
				EP	1713442 A1	25-10-2006	
				WO	2005074890 A1	18-08-2005	

20

25

30

35

40

45

50

55

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20050175707 A [0052]

摘要

本文公开了包含难溶性药物的微粒的组合物，该难溶性药物特别是由具体为聚乙二醇(PEG)的稳定剂封装的阿片类。进一步描述了包含这类封装组合物的药物组合物。还描述了制备这类封装的微粒组合物的方法，以及制备相应的药物组合物的方法。在本文中描述的封装的微粒组合物允许通过对患者无侵害的途径，如通过口服，将难溶性药物给予有生物可利用性。