



- (51) International Patent Classification:
G01N 33/66 (2006.01) *G01N 33/68* (2006.01)
C12Q 1/68 (2006.01)
- (21) International Application Number:
PCT/US2013/068925
- (22) International Filing Date:
7 November 2013 (07.11.2013)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/723,473 7 November 2012 (07.11.2012) US
- (71) Applicant: OHIO UNIVERSITY [US/US]; 340 West State Street, Unit 14, Athens, OH 45701 (US).
- (72) Inventors: CHEN, Hao; 7972 Floyd Drive, The Plains, OH 45780 (US). PAN, Ning; 393 Clippinger Laboratories, Department of Chemistry and Biochemistry, Athens, OH 45701 (US). LIU, Pengyuan; 393 Clippinger Laboratories, Department of Chemistry and Biochemistry, Athens, OH 45701 (US).
- (74) Agents: NOLAN, Lisa, M.A. et al.; Wood, Herron & Evans, LLP, 2700 Carew Tower, 441 Vine Street, Cincinnati, OH 45202-2917 (US).
- (81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

[Continued on next page]

(54) Title: IONIZATION OF CHEMICALS IN MIXTURE AT LOW PH BY AMBIENT IONIZATION / MASS SPECTROMETRY




FIG. 1

(57) Abstract: A mass spectrometry-based method for analyzing an acidic organic target compound includes directing a charged solvent (44) toward a pre-acidified sample (12) comprising the target compound, to thereby ionize the pre-acidified sample (12). The method further includes directing the ionized pre-acidified sample (54) to a mass spectrometer (18), the mass spectrometer (18) being configured to identify and quantify the target compound.



TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, —
KM, ML, MR, NE, SN, TD, TG).

*before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

Published:

— *with international search report (Art. 21(3))*

IONIZATION OF CHEMICALS IN MIXTURE AT LOW pH BY
AMBIENT IONIZATION / MASS SPECTROMETRY

Cross-Reference to Related Application

[0001] This application claims the benefit of U.S. Provisional Patent Application Serial No. 61/723,473, filed on November 7, 2012, the disclosure of which is incorporated herein by reference in its entirety.

Technical Field

[0002] The present invention is generally related to a method for ionization of chemicals in a mixture. In particular, the present invention is related to ionization of phosphopeptides in mixtures by ambient ionization / mass spectrometry.

Background

[0003] Phosphorylation is one of the most common post-translational modifications (PTM) of proteins (approximately 30% of cellular proteins are phosphorylated), and it plays an important role in a wide range of biological processes, such as signal transduction. Phosphorylation is the addition of a phosphate (PO_4^{3-}) group to a protein or other organic molecule. Phosphorylation turns many protein enzymes on and off, thereby altering their function and activity. Protein phosphorylation in particular plays a significant role in a wide range of cellular processes. Its prominent role in biochemistry is the subject of a very large body of research.

[0004] Mass spectrometry (MS) has become an increasingly viable technology for phosphoprotein analysis. However, a major challenge in this regard is that phosphopeptide signal in the positive ion mode is severely suppressed by non-phosphorylated peptides when a phosphoprotein digest is ionized by traditional ionization methods such as electrospray ionization (ESI) in a commonly used "bottom-up" approach. Preliminary purification of phosphopeptides prior to MS analysis is often indispensable to solve the problem, using antibodies, affinity chromatography, metal oxides, nanopolymers, or nanoparticles to enrich phosphopeptides. As separation and purification could be time-consuming, a direct, rapid and sensitive method for ionizing phosphopeptides in mixtures would be instrumental to facilitating their analysis and characterization.

[0005] Desorption electrospray ionization (DESI) is a recent advance in the field of MS. DESI provides direct ionization of analytes with little or no sample preparation. Sample ionization by DESI occurs via the interactions with charged microdroplets generated in a pneumatically assisted electrospray of an appropriate solvent. In addition

to analysis of solid samples, DESI has been extended to directly ionize liquid samples, and its demonstrated applications include the coupling MS with chromatography, microfluidics, and electrochemistry, probing protein conformation, and developing submilli-second time-resolved MS.

Summary

[0006] The present invention is premised on the realization that an organic compound can be analyzed at low pH (i.e., in strong acidic media) using a mass spectrometer at very low pH by DESI. DESI utilizes a regular solvent as an ionizing electrospray, which, in turn, directs analyte compound to a mass spectrometer configured to analyze the analyte. Potential damage to the mass spectrometer caused by strong acid in sample mixture can be prevented due to the use of regular spray solvent, which is neutral or slightly acidic and can dilute the sample during the DESI ionization process.

[0007] The present invention is further premised on the realization that phosphoproteins can be successfully analyzed utilizing DESI coupled with MS. More particularly, the phosphopeptide can be acidified with a strong acid, such as hydrochloric acid, which suppresses the deprotonization of the peptide phosphate groups in solution. The acidified phosphopeptide can then be ionized using DESI, whereby the ionizing electrospray effectively dilutes the acidified phosphopeptide, allowing it to be analyzed using MS without damaging the mass spectrometer.

[0008] The objects and advantages of the present invention will be further appreciated in light of the following detailed description and drawings provided herein.

Brief Description of the Drawings

[0009] The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate embodiments of the invention and, together with a general description of the invention given above and the detailed description given below, serve to explain the principles of the invention.

[0010] FIG. 1 is a diagrammatic cross sectional view of an embodiment of the present invention.

[0011] FIG. 2A is an ESSI-MS spectrum according to Experiment 1.

[0012] FIG. 2B is a DESI-MS spectrum according to Experiment 1.

[0013] FIG. 3A is an α -casein sequence according to Experiment 2.

[0014] FIGS. 3B– 3E are α -casein digests according to Experiment 2.

[0015] FIGS. 4A–4B are CID MS/MS spectra according to Experiment 2.

[0016] FIG. 5 is a DESI-MS spectrum according to Experiment 4.

[0017] FIG. 6A is a β -casein sequence according to Experiment 5.

- [0018] FIG. 6B is an ESSI-MS spectrum according to Experiment 5.
- [0019] FIG. 6C is a DESI-MS spectrum according to Experiment 5.
- [0020] FIGS. 7A-7B are ECD MS/MS spectra according to Experiment 5.
- [0021] FIGS. 8A-8B are ESSI-MS spectrum according to Experiment 5.

Detailed Description

[0022] Although the present invention can be used with any acidic compound that suffers ion suppression in the presence of other compounds by traditional ionization method, such as ESI, it is particularly useful in the analysis of phosphoproteins and sulfated proteins. For purpose of description, a target compound, such as phosphopeptides, is made very acidic by addition of an appropriate strong acid to prevent suppression of the phosphopeptide signal by non-phosphorylated peptides. However, in general, this method is applicable to acidic compounds.

[0023] When the present invention is used for analysis of phosphopeptides, the phosphopeptide is combined with a strong acid that does not interfere with subsequent analysis utilizing a mass spectrometer. The phosphopeptide combined with the acid is referred to as a pre-acidified sample. In particular, haloacids, such as hydrochloric acid and hydrobromic acid can be combined with the phosphopeptide to achieve a desired pH of approximately 0 to 2.0. The acids must have a pKa1 smaller than the pKa1 of phosphoric acid, which is 2.12. Further, the strong acid must not provide a counter ion that interferes with the detection of the group of the phosphopeptides. Therefore, acids such as sulfuric acid and phosphoric acid are unsuitable for use in the present invention.

[0024] Briefly, the present invention provides a method for phosphoprotein analysis using DESI-MS. An apparatus 10 suitable for practicing the present invention includes a conduit 14 for the pre-acidified sample 12, an ambient ionizer 16, and the mass spectrometer 18. The pre-acidified sample 12 comprises an acidic organic target compound. The pre-acidified sample 12 is delivered through conduit 14 having an outlet 22. The conduit 14 may comprise, for example, a fused silica transfer capillary having an inner diameter of approximately 0.05 mm to approximately 0.2 mm. More specifically, the capillary may have an inner diameter of approximately 100 μm . The pre-acidified sample 12 may flow through the conduit 14 at a rate of approximately 1 $\mu\text{L}/\text{min}$ to approximately 5 $\mu\text{L}/\text{min}$.

[0025] The ambient ionizer 16 includes a spray probe 42, which generates microdroplets of a charged solvent 44 and directs the charged solvent 44 toward the pre-acidified sample 12 that emerges from the conduit outlet 22. The spray probe 42 may be positioned approximately 0.5 mm to approximately 5 mm from the conduit outlet 22, for

example. More specifically, the spray probe 42 may be positioned approximately 1 mm from the conduit outlet 22. The charged solvent 44 ionizes the pre-acidified sample 12 that emerges from the conduit outlet 22.

[0026] As shown, the ambient ionizer 16 is a DESI apparatus that includes a housing 46 having a solvent conduit 48 for solvent 49 surrounded by a gas conduit 50. The solvent 49 that is supplied to the ambient ionizer 16 may comprise, for example, methanol/water (1:1 by volume) containing 1% acetic acid. A voltage generator 52 is attached to the housing 46 and is operable to charge the solvent 49 within the solvent conduit 48. A high voltage of approximately 4 kV to approximately 5.5 kV may be applied to the solvent 49. More specifically, a voltage of approximately 5 kV may be applied to the solvent 49. The DESI apparatus generates the nebulized, charged solvent 44 that ionizes the pre-acidified sample 12 by desorption, forming an ionized sample 54. The DESI solvent flow rate may be from about 0.05 $\mu\text{L}/\text{min}$ to approximately 50 $\mu\text{L}/\text{min}$. More specifically, the solvent 49 may be injected into the DESI apparatus at a rate of approximately 10 $\mu\text{L}/\text{min}$.

[0027] An LS-DESI-MS system is described in further detail in U.S. Patent Nos. 7,915,579 and 8,330,119, the disclosures of which are incorporated in their entireties herein by reference.

[0028] The spray impact of the microdroplets of charged solvent 44 from the spray probe 42 with the pre-acidified sample 12 ionizes and deflects an ionized portion of the sample 54 into a mass spectrometer 18, such as a Thermo Finnigan LCQ DECA ion trap mass spectrometer 18 (Thermo Scientific, San Jose, CA). The mass spectrometer 18 has a sample entrance or opening 62, such as a heated capillary, which is also positioned near the conduit outlet 22 and the spray probe 42 of the ambient ionizer 16. The opening 62 may be positioned approximately 10 mm from the conduit outlet 22. The ionized sample 54 enters the opening 62, where a pump (not shown) maintains the atmosphere in the mass spectrometer 18 as a vacuum. The mass spectrometer 18 analyzes a mass-to-charge ratio of the ionized sample 54, as described in U.S. Patent Nos. 7,915,579 and 8,330,119. In this way, the mass spectrometer 18 is configured to identify and quantify an amount of the target compound.

[0029] In use, a pre-acidified sample 12 of, for example, acidified phosphopeptide solution that has been acidified by a strong acid, such as HCl, is provided. The introduction of the strong acid lowers the pH of the phosphopeptide solution and suppresses the deprotonation of peptide phosphate groups in the solution. The pre-acidified sample 12 is delivered through conduit 14 and emerges through a

conduit outlet 22, which is positioned proximate a spray probe 42 of the ambient ionizer 16. Solvent 49 is directed through an ambient ionizer 16, such as a DESI ionizer, which contacts the pre-acidified sample 12 and ionizes the pre-acidified sample 12 emerging from the conduit outlet 22. The ionization deflects an ionized portion of the sample 54 into an opening 62 of a mass spectrometer 18. The mass spectrometer 18 analyzes a mass-to-charge ratio of the ionized sample 54. In this way, the mass spectrometer 18 can identify and quantify an amount of the phosphopeptide target compound.

[0030] The method of the present invention helps prevent suppression of phosphopeptide signals during phosphoprotein analysis by MS. The intrinsic cause for the suppression of phosphopeptide ionization is that the phosphate groups of phosphopeptides tend to lose protons to carry negative charges. By adding a stronger acid than phosphoric acid to the target compound, phosphate deprotonation can be inhibited. Because the strong acid is added to the target compound prior to ionization, regular DESI-like solvent may be used for ionization.

[0031] As such, when the pre-acidified sample 12 is impinged by DESI spray 44 during ionization, the pre-acidified sample 12 is diluted by the DESI spray solvent 49, which reduces or prevents corrosion of the instruments in the apparatus 10. With the strong acid additives in the pre-acidified sample 12, phosphopeptides are charged as positive ions in the pre-acidified sample 12, which enables the pre-acidified sample 12 to be quickly converted from liquid phase to gas phase by DESI. Indeed, a 100% coverage of phosphorylated peptides is enabled by DESI-MS using only a few pmol of proteins. In addition, increased charges of phosphopeptide ions are also noted, which is valuable for tandem MS analysis.

[0032] This DESI-MS methodology may also be used for analysis of other acidic biomolecules, such as a sialic acid, a sialylated glycan, a phosphorylated protein, a sulphated protein or peptide, or a nucleic acid.

[0033] The present invention will be further appreciated in light of the following examples.

Experiment 1

[0034] With reference to FIGS. 2A-2B, O-phospho-L-tyrosine, a phosphorylated amino acid, was chosen as a test sample for a proof-of-principle experiment. When an amino acid mixture containing O-phospho-L-tyrosine and L-tyrosine (molar ratio 3:1) in MeOH/H₂O/HOAc (pH=3.3) was ionized by electronsonic spray ionization (ESSI, a variant form of ESI), the corresponding protonated molecules were observed at m/z 262 and 182, respectively (FIG. 2A). However, the intensity ratio of m/z 262 vs. m/z 182 is

0.3 although O-phospho-L-tyrosine is 3 times more concentrated than L-tyrosine, which shows the suppression of the phosphoamino acid in comparison to non-phosphorylated amino acids. Interestingly, when the amino acid mixture was acidified with HCl to pH 2.0 and ionized by DESI with the spray solvent 49 of MeOH/H₂O/HOAc, the signal intensity of m/z 262 exceeded m/z 182 (FIG. 2B). In addition, the absolute intensity of m/z , 262 in DESI-MS spectrum (2.0E7, arbitrary units, FIG. 2B) was higher than that in ESSI-MS spectrum (4.9E6, FIG. 2A). Evidently, the addition of HCl suppresses the deprotonation of phosphate group of O-phospho-L-tyrosine so that it can be more easily ionized by DESI.

Experiment 2


[0035] With reference to FIGS. 3A-4B, phosphoprotein digests were also examined with this DESI approach. α -Casein (sequence shown in FIG. 3A; the subscript "p" indicates that the residue serine is phosphorylated), a phosphoprotein carrying eight phosphate groups, was digested using trypsin following the reported procedure. In a comparison experiment using ESSI, among 16 peptide ions identified based on acquired MS spectrum (FIG. 3B; the figure inset shows the list of identified peptide ions), three phosphopeptide ions [VPQLEIVNP_pSAEER+2H]²⁺ (m/z 832), [YKVPQLEIVNP_pSAEER+2H]²⁺ (m/z 977), and [YKVPQLEIVNP_pSAEER+H]⁺ (m/z 1953) were detected, covering only one phosphorylation site of the protein. Other phosphorylated peptides are missing in the spectrum, emphasizing the well-known ion suppression effect mentioned above. In a stark contrast, when the sample was acidified by HCl to pH 2.0 and analyzed by DESI, besides the 16 peptide ions seen in the ESSI-MS spectrum, two additional doubly charged ions, [DIG_pSE_pSTEDQAMEDIK+2H]²⁺ (m/z 965) and [QMEAE_pSI_pSP_pSP_pSEEIVNP_pSVEQK+2H]²⁺ (m/z 1361), were detected (FIG. 3C). The successful visualization of the highly acidic phosphopeptide QMEAE_pSI_pSP_pSP_pSEEIVNP_pSVEQK is remarkable as it carries 5 phosphates. Upon collision-induced dissociation (CID), m/z 965 gives rise to fragment ions y_4, y_5, y_6, y_8 along with losses of one and two molecules of H₃PO₄ (FIG. 4A), and m/z 1361 dissociates into $b_5, b_7, b_{13}, b_{14}, y_5, y_7, y_8, y_9, y_{10}, y_{14}$ along with loss of one H₃PO₄ molecule (FIG. 4B), confirming their structures. The detection of these two additional phosphopeptides, DIG_pSE_pSTEDQAMEDIK and QMEAE_pSI_pSP_pSP_pSEEIVNP_pSVEQK, allows to cover all eight phosphorylation sites of the protein. In this experiment, only 5 pmol of α -casein digest was injected for ionization to acquire the DESI-MS spectrum, suggesting the high sensitivity of the method.

[0036] The feasibility of this DESI method was further confirmed with another sample, the α -casein Glu-C digest. Unlike trypsin digestion producing peptides carry a basic residue at the C-terminal, Glu-C selectively cleaves peptide bonds C-terminal to glutamic acid residues, which may provide phosphopeptides without any basic residues that would be more sensitive to the signal suppression effect. Indeed, only one phosphopeptide ion $[\text{IVPN}_\text{p}\text{SAEE}+\text{H}]^+$ (m/z 938) appears in the ESSI-MS spectrum (FIG. 3D). After acidification with HCl and being ionized by DESI, two more phosphopeptide ions, $[\text{LSKDIG}_\text{p}\text{SE}_\text{p}\text{STE}+\text{H}]^+$ (m/z 1326) and $[\text{AE}_\text{p}\text{SI}_\text{p}\text{S}_\text{p}\text{S}_\text{p}\text{SEEIVPN}_\text{p}\text{SVE}+2\text{H}]^{2+}$ (m/z 1039) were detected (FIG. 3E), again covering all phosphate-carrying residues in the protein.

Experiment 3

[0037] The corrosion effect of HCl used in DESI to the mass spectrometer 18 instrument was evaluated in a separate experiment. A piece of stainless steel (316 S.S) was chosen to block the opening 62 of the mass spectrometer 18 to receive the sprayed liquids from DESI. When a sample of MeOH/H₂O with pH adjusted to 2.0 by HCl was injected to undergo DESI ionization for 30 min (both the flow rates and the DESI spray solvent were kept the same as in the analysis of phosphopeptide samples), no detectable damage was noted. While the same sample (pH 2.0) was sprayed by ESSI for 30 min, some traces of black marks on the stainless steel surface were seen, indicating the occurrence of some corrosion. This result shows the strength of DESI in analyzing low pH samples. Sulfuric acid (H₂SO₄), another strong acid, was chosen to replace HCl for the DESI experiments; however, the ionization of the phosphopeptides failed, probably because SO₄²⁻ anions form adducts with positively charged peptide ions. HCl, a volatile acid, appears to be a better choice for such a DESI experiment. Recently HCl was also reported to be employed as an additive to the counter-flow gas in ESI to change the charge numbers of resulting protein ions.

Experiment 4



Scheme 1

[0038] A proposed mechanism for the efficient ionization of phosphopeptides from protein digests by DESI is depicted in Scheme 1. In a sample solution acidified with weak acids like acetic acid, phosphate groups carry negative charges due to deprotonation, which makes the phosphopeptides to be negatively charged. Once acidified with HCl to a low pH that suppresses the phosphate deprotonation, the phosphopeptides are positively charged in solution and therefore are able to compete with non-phosphorylated peptides to produce abundant positive ions by DESI. According to the Henderson-Hasselbalch equation $\text{pH} = \text{p}K_{a1} + \log([A^-]/[HA])$ (HA and A^- represent an acid and its conjugated base, respectively), the population of HA species (e.g., the intact phosphate group of phosphopeptides) can be further increased if the pH is lowered. This accounts for the contrast between the ionization of phosphopeptides at different pHs. Indeed, when the α -casein tryptic digest was acidified by HCl to pH 1.2, the signal intensity for $[\text{QMEAEpSlpSpSpSEIIVPNpSVEQK} + 2\text{H}]^{2+}$ (m/z 1361) was further enhanced to 1.8E4 (FIG. 5) in comparison to that obtained via ionization at pH 2.0 (1.1E4, FIG. 3C). In the experiment of the ionization of pH 1.2 sample by DESI, the sample injection rate was reduced to 2 $\mu\text{L}/\text{min}$ to avoid instrument corrosion. As shown in Scheme 1, these phosphopeptides positively charged in solution can be rapidly transferred into the gas phase by DESI for MS detection due to the direct sampling capability of DESI. An evidence to support this hypothesis is that, when the α -casein tryptic digest sample was mixed with the DESI spray solvent (1:2 by volume, mimicking the mixing in the DESI ionization process) and then was analyzed by ESSI, the m/z 1361 was not observed (data not shown). Presumably, the m/z 1361 detected in DESI-MS (FIG. 3E) survived the spray/sample solution mixing process in DESI because of the very short DESI ionization time scale (approximately 1-2 milliseconds) that favors the preservation of positively charged phosphopeptides during the ionization.

Experiment 5

[0039] Besides the efficient ionization of highly acidic phosphopeptides from protein digests, charges of ionized phosphopeptides can also be enhanced using this DESI approach. For instance, in the case of β -casein tryptic digest (protein sequence shown in FIG. 6A), the intensity of the doubly charged ion $[\text{RELEELNVPGEIVepSLpSpSpSEESITR} + 2\text{H}]^{2+}$ (m/z 1562) increased from 2.9E5 in ESSI-MS spectrum (FIG. 6B) to 5.1E5 in DESI-MS spectrum (FIG. 6C). More importantly, abundant triply charged ion $[\text{RELEELNVPGEIVepSLpSpSpSpSEESITR} + 3\text{H}]^{3+}$ (m/z 1042) was generated (FIG. 6C). The enhanced charges would be valuable in providing increased sequence coverage via

electron-based tandem MS analysis, such as electron-capture dissociation (ECD).

Indeed, ECD of m/z 1042 gives rise to fragment ions of $z_2, z_3, z_4, z_5, z_6, z_7, c_{17}, c_{21}, c_{22}, c_{23}, c_{24}$, and c_{25} (FIG. 7A) from which the locations of pSer¹⁸ and pSer¹⁹ of the protein can be clearly pinpointed. In contrast, ECD of m/z 1562 only gives rise to two fragment ions, c_{25} and z_{25} (FIG. 7B).

[0040] This DESI method is also applicable to the analysis of acidic sulfated peptides. Using a mixture containing sulfated hirudin (sequence: DFEEIPEE-Y(SO₃H)-LQ), bradykinin and angiotensin II as a test sample (1:10:10 by moles), the sulfated hirudin was suppressed and no corresponding peptide ion was observed in ESSI-MS (FIG. 8A). In contrast, the doubly charged ions [DFEEIPEE-Y(SO₃H)-LQ+2H]²⁺ (m/z 746) arose via DESI-MS analysis (FIG. 8B), and its structure was confirmed by CID (the inset of FIG. 8B).

[0041] This invention presents a novel approach for solving the ion signal suppression problem that occurs with phosphopeptide ionization in protein digests by acidifying the target compound with strong acid HCl followed by direct DESI-MS analysis. The methodology is general, fast and sensitive. Efficient ionization can be achieved with the coverage of all phosphate carrying residues without laborious separation or enrichment of phosphopeptides. Thus, it reduces the cost of phosphoprotein analysis and saves time. The methodology of this invention is applicable for improving MS analysis of other important biomolecules with high acidity such as sialic acids, sialylated glycans, sulphated proteins, and nucleic acids.

[0042] This has been a description of the present invention along with the various methods of practicing the present invention. However, the invention itself should only be defined by the appended claims. What is claimed is:

Claims

1. A mass spectrometry-based method for analyzing an acidic organic target compound, the method comprising:
directing a charged solvent toward a pre-acidified sample comprising the target compound, thereby ionizing the pre-acidified sample; and
directing the ionized pre-acidified sample to a mass spectrometer, the mass spectrometer being configured to identify and quantify the target compound.
2. The method of claim 1, wherein the pre-acidified sample comprises a blend of biomolecules and a strong acid.
3. The method of claim 2, wherein the biomolecules comprise a phosphopeptide.
4. The method of claim 2, wherein the biomolecules comprise one of a sialic acid, a sialylated glycan, a phosphorylated protein, a sulphated protein or peptide, and a nucleic acid.
5. The method of claim 2, wherein the strong acid comprises hydrochloric acid.
6. The method of claim 1, wherein the charged solvent is generated by desorption electrospray ionization.
7. The method of claim 1, wherein the charged solvent is generated by one of laser beam ionization, high energy particles ionization, plasma ionization, and ion beam ionization.
8. The method of claim 1, wherein the pre-acidified sample has a pH less than or equal to approximately 2.0.
9. The method of claim 8, wherein the pre-acidified sample has a pH less than or equal to approximately 1.2.

10. The method of claim 1, wherein the addition of the strong acid suppresses deprotonation of phosphate groups in the target compound.

11. The method of claim 1, wherein the pre-acidified sample flows at a rate of approximately 1 $\mu\text{L}/\text{min}$ to approximately 5 $\mu\text{L}/\text{min}$.

12. The method of claim 1, wherein ionization of the pre-acidified sample does not suppress ion signals in the pre-acidified sample.

13. The method of claim 1, further comprising:
flowing the pre-acidified sample through a conduit, the conduit having an outlet positioned approximately 10 mm from an opening of the mass spectrometer.

14. The method of claim 1, wherein the charged solvent is created by applying a voltage of approximately 5 kV to a solvent.

15. A mass spectrometry-based method for effectively analyzing an acidic organic/biological compound in the presence of complicated matrix and for effectively removing/reducing the ion signal suppression effect, the method comprising:

directing a charged solvent toward the pre-acidified sample comprising the target compound, thereby overcoming the ion suppression effect and ionizing the low pH sample with high ionization efficiency; and

directing the ionized low pH sample to a mass spectrometer, the mass spectrometer being configured to identify and quantify the target compound.




FIG. 1

2/18




FIG. 2A




FIG. 2B

4/18

RPKHPIKHQGLPQEVLNENLLRFFVAPFFPEVFGKEKVNELSKDIG_pSE_pSTEDQAMEDIKQMEAE_pSI_pS_pS_pSEEIV
PN_pSVEQKHQKEDVPSERYLGYLEQLRLKKYKVPQLEIVPN_pSAEERLHSMKEGIHAQQKEPMGIVNQELA
YFYPELFRQFYQLDAYPSGAWYYVPLGTQYTDAPSFSDIPNPIGSENSEKTTMPLW (SEQ. ID. NO.10)

FIG. 3A

5/18




FIG. 3B




FIG. 3C

7/18




FIG. 3D




FIG. 3E




FIG. 4A




FIG. 4B




FIG. 5

12/18

RELEELNVPGEIVE_{pSL_pS_p}SEESITRINKKIEKFQ_pSEEQQQTEDELQDK
IHPFAQTQSLVYPFPGPIPN_{SLPQNIPPLTQTPVVPPFLQPEVMGVSKV}
KEAMAPKHKEMPFPKVPVEPFTESQSLTLDVENLHPLPLQSWMH
QPHQPLPPTVMFPPQSVLSLSQSKVLPVPQKAVPYPQRDMPIQAFLLYQ
EPVLGPVRGPFPIIV(SEQ. ID. NO.26)

FIG. 6A




FIG. 6B




FIG. 6C

15/18




FIG. 7A

16/18




FIG. 7B

17/18




FIG. 8A




FIG. 8B

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/068925

A. CLASSIFICATION OF SUBJECT MATTER
INV. G01N33/66 C12Q1/68 G01N33/68
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, COMPENDEX, EMBASE, INSPEC, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LUCIE HARTMANOVA ET AL: "Fast profiling of anthocyanins in wine by desorption nano-electrospray ionization mass spectrometry", JOURNAL OF CHROMATOGRAPHY A, vol. 1217, no. 25, 1 June 2010 (2010-06-01), pages 4223-4228, XP055104613, ISSN: 0021-9673, DOI: 10.1016/j.chroma.2010.03.018 the whole document in particular: par. 2. Experimental; page 4224 page 4224; figure 2 par. 4. Conclusion; page 4228 ----- -/-	1,6,7, 11-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 March 2014

Date of mailing of the international search report

19/03/2014

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Tuynman, Antonin

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/068925

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SARA BOGIALLI ET AL: "Confirmatory analysis of sulfonamide antibacterials in bovine liver and kidney: extraction with hot water and liquid chromatography coupled to a single- or triple-quadrupole mass spectrometer", RAPID COMMUNICATIONS IN MASS SPECTROMETRY, vol. 17, no. 11, 15 June 2003 (2003-06-15), pages 1146-1156, XP055104657, ISSN: 0951-4198, DOI: 10.1002/rcm.1031 abstract par. "Sample preparation and extraction"; page 1148	1,7
A	----- LUKÁŠ KRÁSNÝ ET AL: "In-situ enrichment of phosphopeptides on MALDI plates modified by ambient ion landing", JOURNAL OF MASS SPECTROMETRY, vol. 47, no. 10, 27 October 2012 (2012-10-27), pages 1294-1302, XP055104655, ISSN: 1076-5174, DOI: 10.1002/jms.3081	1-15
A	----- BO QIU ET AL: "Desorption electrospray ionization mass spectrometry of DNA nucleobases: implications for a liquid film model", JOURNAL OF MASS SPECTROMETRY, vol. 44, no. 5, 1 May 2009 (2009-05-01), pages 772-779, XP055104658, ISSN: 1076-5174, DOI: 10.1002/jms.1554	1-15
X,P	----- NING PAN ET AL: "Highly efficient ionization of phosphopeptides at low pH by desorption electrospray ionization mass spectrometry", THE ANALYST, vol. 138, no. 5, 1 January 2013 (2013-01-01), page 1321, XP055104670, ISSN: 0003-2654, DOI: 10.1039/c3an36737a the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2013/068925

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. (means)

☒

 on paper

☒

 in electronic form
 - b. (time)

☐

 in the international application as filed

☐

 together with the international application in electronic form

☒

 subsequently to this Authority for the purpose of search
2.

☒

 In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments: