




US 20080139791A1

(19) **United States**(12) **Patent Application Publication****Lipovsek et al.**(10) **Pub. No.: US 2008/0139791 A1**(43) **Pub. Date: Jun. 12, 2008**(54) **PHARMACEUTICALLY ACCEPTABLE FN3
POLYPEPTIDES FOR HUMAN TREATMENTS**(75) Inventors: **Dasa Lipovsek**, Cambridge, MA
(US); **Richard W. Wagner**,
Concord, MA (US); **Robert G.**
Kuimelis, Brighton, MA (US)

Correspondence Address:

ROPES & GRAY LLP**PATENT DOCKETING Floor 39, One Interna-
tional Place
Boston, MA 02110-2624**(73) Assignee: **Adnexus Therapeutics, Inc.**,
Waltham, MA (US)(21) Appl. No.: **11/894,732**(22) Filed: **Aug. 20, 2007****Related U.S. Application Data**(63) Continuation of application No. 11/483,918, filed on
Jul. 7, 2006, which is a continuation of application No.10/728,078, filed on Dec. 3, 2003, now Pat. No. 7,115,
396, which is a continuation of application No. 09/688,
566, filed on Oct. 16, 2000, now abandoned, which is
a continuation-in-part of application No. 09/515,260,
filed on Feb. 29, 2000, now Pat. No. 6,818,418, which
is a continuation-in-part of application No. 09/456,
693, filed on Dec. 9, 1999, now abandoned.(60) Provisional application No. 60/111,737, filed on Dec.
10, 1998.**Publication Classification**(51) **Int. Cl.**
C07K 16/00 (2006.01)(52) **U.S. Cl.** **530/387.3**(57) **ABSTRACT**Disclosed herein are proteins that include an immunoglobulin
fold and that can be used as scaffolds. Also disclosed herein
are nucleic acids encoding such proteins and the use of such
proteins in diagnostic methods and in methods for evolving
novel compound-binding species and their ligands.




Fig. 1




Fig. 2




Fig. 3

[illegible]

Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11




Fig. 12

FIG. 13




FIG. 14




FIG. 15




FIG. 16





FIG. 17






Fig. 18




Fig. 19




Fig. 20




Fig. 21





Fig. 22






Fig. 24A




Fig. 24B




Fig. 24C




Fig. 24D

FIG. 25A

ROUND 9 (32)

WT	BC				DE				FG			
	15	16	DATAVTV3031	45	46	QSKS	60	61	74	75	GRGDS	PASSK
VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	90					
T09.08 VSEIPRDLVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	33					
T09.26 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	34					
T09.20 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	35					
T09.25 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	36					
T09.30 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	37					
T09.02 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	38					
T09.34 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	39					
T09.03* VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	40					
T09.13 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	41					
T09.29 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	42					
T09.01 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	43					
T09.14 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	44					
T09.24 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	45					
T09.19 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	46					
T09.31 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	47					
T09.07 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	48					
T09.36 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	49					
T09.16 -NTTHYKXNYXATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	50					
T09.23 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	51					
T09.21 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	52					
T09.33* VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	53					
T09.10 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	54					
T09.28 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	55					
T09.15 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	56					
T09.09 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	57					
T09.35 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	58					
T09.06 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	59					
T09.11 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	60					
T09.12* VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	61					
T09.17 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	62					
T09.04 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	63					
T09.05 VSDVPRDLEVVAAATP	TSLLISWDABAVTVR	YRITYYKGTGNSFV	QRETVGSKSTATIS	GLKRGVDY-TITVYA	VTGGRDSDPASKPIS	INVRT	64					

FIG. 25B

ROUND 10 (29)

		BC		DE		FG				
WT		15	16	45	46	60	61	74	75	90
T10.37	VSDVPRDLEVVAAATP	TSLLISWDAEAVTVR	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTORGDSFASPKPIS	INVRT	33		
T10.12	VSDVPRDLEVVAAATP	TSLLSIREB-----RR	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTPHGHFDLELPIS	INVRT	66		
T10.21	VSDVPRDLEVVAAATP	TSLLSIRKURVBSRR	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTPHGHFDLELPIS	INVRT	67		
T10.40	VSDVPRDLEVVAAATP	TSLLSIRKURVBSRR	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTSVNAFFYEGMPIS	INVRT	68		
T10.31a	VSDVPRDLEVVAAATP	TSLLSWILCTGNNR	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	LSLASLSTRNKPIS	INVRT	69		
T10.41	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTPTMLKPLSMPIS	INVRT	70		
T10.13	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VNQPTVBAHNHAPIS	INVRT	71		
T10.03	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTSHRDYHSTGRPIS	INVRT	72		
T10.33	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	73		
T10.27	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	74		
T10.05	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	75		
T10.01	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	76		
T10.34	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	77		
T10.36	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	78		
T10.30	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	79		
T10.06	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	80		
T10.17	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	81		
T10.18	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	82		
T10.39	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	83		
T10.02	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	84		
T10.08	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	85		
T10.20	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	86		
T10.29	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	87		
T10.15	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	88		
T10.15	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	89		
T10.35	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	90		
T10.26	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	91		
T10.14	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	92		
S08.02	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	94		
S08.03	VSDVPRDLEVVAAATP	TSLLSWIRWASPFG	YRITTYGEGNSPV	QEFVPSKSTATIS	GLKRGVDY-TITVYA	VTQSTNGNRNDPIS	INVRT	95		

FIG. 25D

ROUND M12 (24)

			BC			DE			FG		
			15	16	DAPAVTV3031	45	46	GSKS	60	61	75
WT											76GRGDSPASSK
M12.05	VSDVPRDLEVVAAATP	TSLLISWDAPAVTVR	YRITYGETGNSPV	QFTVPGSKSTATIS	GLKPGVDY-TITVYA	VTGRGDSPASSKPIIS	INVRT				33
M12.17	VSDVPRDLEVVAAATP	TSLLISWSPENDARR	YRITYGKTGDSFP	QFTVPGSKSTATIS	GLKPGVDY-TSVVYA	VTDQSYTYSNPIIS	INVRT				117
M12.12	VSDVPRDLEVVAAATP	TSLLISWBSQSPKGR	YRITYGETGNSPV	QFTVPGSKSTATIS	GRKPGADY-TITVYA	VTIEKDRIFLFGPIIS	ISVRT				118
M12.17	VSDVPRDLEVVAAATP	TSLLISWBSQSPKGR	YRITYGETGNSPV	QFTVPGSKSTATIS	GRKPGADY-TITVYA	VTIEKDRIFLFGPIIS	ISVRT				119
M12.06	VSDVPRDLEVVAAATP	TSLLISWBSQSPKGR	YRITYGETGNSPV	QFTVPGSKSTATIS	GRKPGADY-TITVYA	VTIEKDRIFLFGPIIS	ISVRT				120
M12.01	VSDVPRDLEVVAAATP	TSLLISWTHNAYNG	YRITYGETGNSPA	QFTVPRPEVTATIS	GLKPGVDY-TITVYA	VTIHRMFLRLFGPIIS	INVRT				121
M12.01	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSR	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTDESDTYKYDDPVS	INVRT				122
M12.01	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSR	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTDESDTYKYDDPVS	INVRT				123
M12.24	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTENUGRLDTKYPIS	IDCRT				124
M12.25	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTENUGRLDTKYPIS	INVRT				125
M12.23	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	XTSATPSRPNVHPIS	INLTT				126
M12.04	VSDVPRDLEVVAAATP	TSLLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				127
M12.03	VSDVPRDLEVVAAATP	TSLLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	ATHGKPPHHT--	INVRT				128
M12.07	VSDVPRDLEVVAAATP	TSLLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				129
M12.13	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				130
M12.09	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				131
M12.16	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				132
M12.22	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				133
M12.26	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				134
M12.15	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				135
M12.18	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				136
M12.14	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				137
M12.21	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				138
M12.19	VSDVPRDLEVVAAATP	TSRLISWNRSGLOSG	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYA	VTNRDT-----PIS	INVRT				139
M12.08	VSDVPRDLEVVAAATP	TSLLANP-----IIR	YRITYGETGNSPV	QFTVPPWASIAATIS	GLKPGVDY-TITVYG	VTNRDT-----PIS	INVRT				140

**PHARMACEUTICALLY ACCEPTABLE FN3
POLYPEPTIDES FOR HUMAN TREATMENTS****CROSS REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a continuation of U.S. patent application Ser. No. 11/483,918, filed Jul. 7, 2006, which is a continuation of U.S. patent application Ser. No. 10/728,078, filed Dec. 3, 2003, which is a continuation of U.S. patent application Ser. No. 09/688,566, filed Oct. 16, 2000, which is a continuation-in-part of U.S. patent application Ser. No. 09/515,260, filed Feb. 29, 2000, now U.S. Pat. No. 6,818,418, which is a continuation-in-part of U.S. patent application Ser. No. 09/456,693, filed Dec. 9, 1999, now abandoned, which claims the benefit of U.S. Provisional application 60/111,737, filed Dec. 10, 1998.

BACKGROUND OF THE INVENTION

[0002] This invention relates to protein scaffolds useful, for example, for the generation of products having novel binding characteristics.

[0003] Proteins having relatively defined three-dimensional structures, commonly referred to as protein scaffolds, may be used as reagents for the design of engineered products. These scaffolds typically contain one or more regions which are amenable to specific or random sequence variation, and such sequence randomization is often carried out to produce libraries of proteins from which desired products may be selected. One particular area in which such scaffolds are useful is the field of antibody design.

[0004] A number of previous approaches to the manipulation of the mammalian immune system to obtain reagents or drugs have been attempted. These have included injecting animals with antigens of interest to obtain mixtures of polyclonal antibodies reactive against specific antigens, production of monoclonal antibodies in hybridoma cell culture (Koehler and Milstein, *Nature* 256:495, 1975), modification of existing monoclonal antibodies to obtain new or optimized recognition properties, creation of novel antibody fragments with desirable binding characteristics, and randomization of single chain antibodies (created by connecting the variable regions of the heavy and light chains of antibody molecules with a flexible peptide linker) followed by selection for antigen binding by phage display (Clackson et al., *Nature* 352: 624, 1991).

[0005] In addition, several non-immunoglobulin protein scaffolds have been proposed for obtaining proteins with novel binding properties. For example, a "minibody" scaffold, which is related to the immunoglobulin fold, has been designed by deleting three beta strands from a heavy chain variable domain of a monoclonal antibody (Tramontano et al., *J. Mol. Recognit.* 7:9, 1994). This protein includes 61 residues and can be used to present two hypervariable loops. These two loops have been randomized and products selected for antigen binding, but thus far the framework appears to have somewhat limited utility due to solubility problems. Another framework used to display loops has been tenascin, a 74 residue, six-strand beta sheet sandwich held together by two disulfide bonds (McConnell and Hoess, *J. Mol. Biol.* 250:460, 1995). This scaffold includes three loops, but, to date, only two of these loops have been examined for randomization potential.

[0006] Other proteins have been tested as frameworks and have been used to display randomized residues on alpha helical surfaces (Nord et al., *Nat. Biotechnol.* 15:772, 1997; Nord et al., *Protein Eng.* 8:601, 1995), loops between alpha helices in alpha helix bundles (Ku and Schultz, *Proc. Natl. Acad. Sci. USA* 92:6552, 1995), and loops constrained by disulfide bridges, such as those of the small protease inhibitors (Markland et al., *Biochemistry* 35:8045, 1996; Markland et al., *Biochemistry* 35:8058, 1996; Rottgen and Collins, *Gene* 164: 243, 1995; Wan-g et al., *J. Biol. Chem.* 270:12250, 1995).

SUMMARY OF THE INVENTION

[0007] The present invention provides a new family of proteins capable of evolving to bind any compound of interest. These proteins, which generally make use of a scaffold derived from a fibronectin type III (Fn3) or Fn3-like domain, function in a manner characteristic of natural or engineered antibodies (that is, polyclonal, monoclonal, or single-chain antibodies) and, in addition, possess structural advantages. Specifically, the structure of these antibody mimics has been designed for optimal folding, stability, and solubility, even under conditions that normally lead to the loss of structure and function in antibodies.

[0008] These antibody mimics may be utilized for the purpose of designing proteins which are capable of binding to virtually any compound (for example, any protein) of interest. For example, the ¹⁰Fn3-based molecules described herein may be used as scaffolds which are subjected to directed evolution to form a population with one or more randomized Fn3 loops that are analogous by position and structure to the complementarity-determining regions (CDRs) of an antibody variable region, and/or to randomize Fn3's other three solvent exposed loops. Such a directed evolution approach results in the production of antibody-like molecules with high affinities for antigens of interest. In addition, the scaffolds described herein may be used to display defined exposed loops (for example, loops previously randomized and selected on the basis of antigen binding) in order to direct the evolution of molecules that bind to such introduced loops. A selection of this type may be carried out to identify recognition molecules for any individual CDR-like loop or, alternatively, for the recognition of two or all three CDR-like loops combined into a non-linear epitope.

[0009] Accordingly, in a first aspect, the present invention features randomized or mutated scaffold proteins. In particular, the invention features a non-antibody protein including a domain having an immunoglobulin-like fold, the non-antibody protein deriving from a reference protein by having a mutated amino acid sequence, wherein the non-antibody protein binds with a K_d at least as tight as 1 μ M to a compound that is not bound as tightly by the reference protein.

[0010] In addition, the invention features a non-antibody protein deriving from a scaffold protein including a domain having an immunoglobulin-like fold, wherein the amino acid sequence of the domain in the derived protein is more than 50% identical to the amino acid sequence of the domain in the scaffold protein.

[0011] In yet another embodiment, the invention features a protein that includes a fibronectin type III domain having at least one randomized loop, the protein being characterized by the ability of the Fn3 domain to bind to a compound that is not bound by the corresponding naturally-occurring Fn3 domain.

[0012] In various preferred embodiments, any of these proteins of the invention bind to their target compounds with a K_d

at least as tight as 500 nM, preferably, with a K_d at least as tight as 100 nM or 10 nM, and, more preferably, with a K_d at least as tight as 1 nM, 500 pM, 100 pM, or even 20 pM. The protein preferably contains one, two, or three mutated loops and at least one of the loops, and preferably two or all three of the loops, contributes to the binding of the protein to the compound. Additionally, the reference protein preferably lacks disulfide bonds, and the derivative protein may have at least one disulfide bond.

[0013] With respect to certain embodiments, the domain having an immunoglobulin-like fold preferably has a molecular mass less than 10 kD or greater than 7.5 kD, and, more preferably, has a molecular mass between 7.5-10 kD. The proteins of the invention may be monomers under physiological conditions or may be multimers, for example, dimers. In other preferred embodiments, the reference protein used to derive a mutated protein of the invention is a naturally-occurring mammalian protein (for example, a human protein); and the domain having an immunoglobulin-like fold is mutated and includes up to 50%, and preferably up to 34%, mutated amino acids as compared to the immunoglobulin-like fold of the reference protein. In addition, the domain having the immunoglobulin-like fold preferably consists of approximately 50-150 amino acids, and more preferably consists of approximately 50 amino acids.

[0014] Derivative proteins of the invention may be derived from any appropriate reference protein including, but not limited to, the preferred proteins, fibronectin or a fibronectin dimer, tenascin, N-cadherin, E-cadherin, ICAM, titin, GCSF-receptor, cytokine receptor, glycosidase inhibitor, antibiotic chromoprotein, myelin membrane adhesion molecule P0, CD8, CD4, CD2, class I MHC, T-cell antigen receptor, CD1, C2 and I-set domains of VCAM-1, I-set immunoglobulin domain of myosin-binding protein C, I-set immunoglobulin domain of myosin-binding protein H, I-set immunoglobulin domain of telokin, NCAM, twitchin, neuroglian, growth hormone receptor, erythropoietin receptor, prolactin receptor, interferon-gamma receptor, β -galactosidase/glucuronidase, β -glucuronidase, transglutaminase, T-cell antigen receptor, superoxide dismutase, tissue factor domain, cytochrome F_c , green fluorescent protein, GroEL, and thaumatin.

[0015] In further preferred embodiments of Fn3 domain-containing proteins, the fibronectin type III domain is a mammalian (for example, a human) fibronectin type III domain; and the protein includes the tenth module of the fibronectin type III (^{10} Fn3) domain. In such proteins, compound binding is preferably mediated by either one, two, or three ^{10} Fn3 loops. In other preferred embodiments, the second (DE) loop of ^{10} Fn3 may be extended in length relative to the naturally-occurring module, or the ^{10} Fn3 may lack, an integrin-binding motif. In these molecules, the integrin-binding motif may be replaced by an amino acid sequence in which a polar amino acid-neutral amino acid-acidic amino acid sequence (in the N-terminal to C-terminal direction) replaces the integrin-binding motif; alternatively, one preferred sequence is serine-glycine-glutamate. In another preferred embodiment, the fibronectin type III domain-containing proteins of the invention lack disulfide bonds.

[0016] Any of the proteins of the invention (for example, the fibronectin type III domain-containing proteins) may be formulated as part of a fusion protein. If the fusion protein is to be used for compound binding or compound binding selections, the fusion protein includes a heterologous protein that does not itself bind to the compound of interest. The heter-

ologous protein may, for example, be an antibody or antibody domain (such as an immunoglobulin F_c domain), a complement protein, a toxin protein, or an albumin protein. In addition, any of the proteins of the invention (for example, the fibronectin type III domain proteins) may be covalently bound to a nucleic acid (for example, an RNA), and the nucleic acid may encode the protein. Moreover, the protein may be a multimer, or, particularly if it lacks an integrin-binding motif, it may be formulated in a physiologically-acceptable carrier.

[0017] The present invention also features proteins that include a fibronectin type III domain having at least one mutation in a β -sheet sequence. Again, these proteins are characterized by their ability to bind to compounds that are not bound or are not bound as tightly by the corresponding naturally-occurring fibronectin domain.

[0018] Any of the proteins of the invention may be immobilized on a solid support (for example, a bead or chip), and these proteins may be arranged in any configuration on the solid support, including an array.

[0019] In a related aspect, the invention further features nucleic acids encoding any of the proteins of the invention. In preferred embodiments, the nucleic acid is DNA or RNA.

[0020] In another related aspect, the invention also features a method for generating a protein which includes a fibronectin type III domain and which is pharmaceutically acceptable to a mammal, involving removing the integrin-binding domain of said fibronectin type III domain. This method may be applied to any of the fibronectin type III domain-containing proteins described above and is particularly useful for generating proteins for human therapeutic applications. The invention also features such fibronectin type III domain-containing proteins which lack integrin-binding domains.

[0021] In yet another related aspect, the invention features methods of obtaining derivative non-antibody proteins which bind to compounds of interest. One such method involves: (a) providing a non-antibody scaffold protein including an immunoglobulin-like fold, wherein the scaffold protein does not bind to the compound with a K_d as tight as 1 μ M; (b) generating mutated derivatives of the non-antibody scaffold protein, thereby producing a library of mutated proteins; (c) contacting the library with the compound; (d) selecting from the library at least one derivative protein which binds to the compound with a K_d at least as tight as 1 μ M; and (e) optionally repeating steps (b)-(d) substituting for the non-antibody scaffold protein in repeated step (b) the product from the previous step (d). This technique may also be carried out with any of the proteins of the invention (for example, any of the fibronectin type III domain-containing proteins).

[0022] In yet other related aspects, the invention features screening methods which may be used to obtain or evolve randomized or mutated proteins of the invention capable of binding to compounds of interest, or to obtain or evolve compounds (for example, proteins) capable of binding to a particular protein containing a randomized or mutated motif. In addition, the invention features screening procedures which combine these two methods, in any order, to obtain either compounds or proteins of interest.

[0023] In particular, a first screening method, useful for the isolation or identification of randomized or mutated proteins of interest, involves: (a) contacting a compound of interest with a candidate protein, the candidate protein being a derivative non-antibody protein including a domain having an immunoglobulin-like fold, the non-antibody protein deriving

from a reference protein by having a mutated amino acid sequence wherein the non-antibody protein binds with a K_d at least as tight as 1 μ M to a compound that is not bound as tightly by the reference protein, wherein the contacting is carried out under conditions that allow compound-protein complex formation; and (b) obtaining, from the complex, the derivative protein that binds to the compound. This general technique may also be carried out with a fibronectin type III domain protein having at least one randomized or mutated loop.

[0024] The second screening method is for isolating or identifying a compound which binds to a protein of the invention. This method begins with a non-antibody protein including a domain having an immunoglobulin-like fold and deriving from a reference protein by having a mutated amino acid sequence, wherein the non-antibody protein binds with a K_d at least as tight as 1 μ M to a compound that is not bound as tightly by the reference protein. This derivative protein is then contacted with a candidate compound, wherein the contacting is carried out under conditions that allow compound-protein complex formation, and the compound which binds to the derivative protein is obtained from the complex. Again, this general technique may be carried out with any protein of the invention, for example, a protein with a mutated fibronectin type III domain.

[0025] In addition, the invention features diagnostic methods which employ the proteins of the invention (for example, fibronectin type III scaffold proteins and their derivatives). Such diagnostic methods may be carried out on a sample (for example, a biological sample) to detect one analyte or to simultaneously detect many different analytes in the sample. The method may employ any of the scaffold molecules described herein. Preferably, the method involves (a) contacting the sample with a protein of the invention that binds to the compound analyte, the contacting being carried out under conditions that allow compound-protein complex formation; and (b) detecting the complex, and therefore the compound in the sample. In addition, this method may be used to quantify, as well as detect, compound levels in a sample.

[0026] In preferred embodiments of any of the selection or diagnostic methods described herein, the protein of the invention binds to its target compound with a K_d at least as tight as 1 μ M or 500 nM, preferably, with a K_d at least as tight as 100 nM or 10 nM, and, more preferably, with a K_d at least as tight as 1 nM, 500 pM, 100 pM, or even 20 pM. The protein preferably contains one, two, or three mutated loops and at least one of the loops, and preferably two or all three of the loops contributes to the binding of the protein to the compound. Additionally, the reference protein preferably lacks disulfide bonds, and the derivative protein may have at least one disulfide bond.

[0027] With respect to certain embodiments of the methods, the domain having an immunoglobulin-like fold preferably has a molecular mass less than 10 kD or greater than 7.5 kD, and, more preferably, has a molecular mass between 7.5-10 kD. The proteins of the invention may be monomers under physiological conditions or may be multimers, for example, dimers. In other preferred embodiments, the reference protein used to derive a mutated protein of the invention is a naturally-occurring mammalian protein (for example, a human protein); and the domain having an immunoglobulin-like fold is mutated and includes up to 50%, and preferably up to 34%, mutated amino acids as compared to the immunoglobulin-like fold of the reference protein. In addition, the

domain having an immunoglobulin-like fold preferably consists of approximately 50-150 amino acids, and more preferably consists of approximately 50 amino acids.

[0028] Derivative proteins used in the methods of the invention may be derived from any appropriate reference protein including, but not limited to, the preferred proteins, fibronectin or a fibronectin dimer, tenascin, N-cadherin, E-cadherin, ICAM, titin, GCSF-receptor, cytokine receptor, glycosidase inhibitor, antibiotic chromoprotein, myelin membrane adhesion molecule P0, CD8, CD4, CD2, class I MHC, T-cell antigen receptor, CD1, C2 and I-set domains of VCAM-1, I-set immunoglobulin domain of myosin-binding protein C, I-set immunoglobulin domain of myosin-binding protein H, I-set immunoglobulin domain of telokin, NCAM, twitchin, neuroglian, growth hormone receptor, erythropoietin receptor, prolactin receptor, interferon-gamma receptor, β -galactosidase/glucuronidase, β -glucuronidase, transglutaminase, T-cell antigen receptor, superoxide dismutase, tissue factor domain, cytochrome F, green fluorescent protein, GroEL, and thaumatin.

[0029] In addition, the steps of the selection methods described herein may be repeated with further mutation or randomization being carried out between cycles. For example, for the methods involving a protein having a mutated or randomized fibronectin type III domain, at least one loop of the fibronectin type III domain of the protein obtained in step (b) may be mutated and steps (a) and (b) repeated using the further randomized protein, or the compound obtained in step (b) may be modified and steps (a) and (b) repeated using the further modified compound. In these methods, the compound is preferably a protein, and the fibronectin type III domain is preferably a mammalian (for example, a human) fibronectin type III domain. In other preferred embodiments, the protein includes the tenth module of the fibronectin type III domain (^{10} Fn3), and binding is mediated by one, two, or three ^{10} Fn3 loops. In addition, the second (DE) loop of ^{10} Fn3 may be extended in length relative to the naturally-occurring module, or ^{10} Fn3 may lack an integrin-binding motif. Again, as described above, the integrin-binding motif may be replaced by an amino acid sequence in which a basic amino acid-neutral amino acid-acidic amino acid sequence (in the N-terminal to C-terminal direction)-replaces the integrin-binding motif; alternatively, one preferred replacement sequence is serine-glycine-glutamate.

[0030] The selection and diagnostic methods described herein may be carried out using any of the proteins of the invention (for example, a fibronectin type III domain-containing protein). In addition, any of these proteins may be formulated as part of a fusion protein with a heterologous protein (for example, an antibody or antibody domain (including an immunoglobulin F_c domain) that does not itself bind the compound of interest, or a complement protein, toxin protein, or albumin protein). In addition, selections and diagnostic methods may be carried out using the proteins of the invention (for example, the fibronectin type III domain proteins) covalently bound to nucleic acids (for example, RNAs or any nucleic acid which encodes the protein). Moreover, the selections and diagnostic methods may be carried out using these proteins (for example, the fibronectin domain-containing proteins) as monomers or as multimers, such as dimers.

[0031] Preferably, the selections and diagnostic methods involve the immobilization of the binding target on a solid support. Preferred solid supports include columns (for example, affinity columns, such as agarose-based affinity col-

umns), microchips, or beads. Alternatively, the proteins (for example, the Fn3 domain-containing proteins) may be immobilized and contacted with one or more potential binding targets.

[0032] For the diagnostic methods, the compound is often a protein, but may also be any other analyte in a sample. Detection may be accomplished by any standard technique including, without limitation, radiography, fluorescence detection, mass spectroscopy, or surface plasmon resonance.

[0033] In a final aspect, the invention features a non-antibody protein that binds tumor necrosis factor- α (TNF- α) with a K_d at least as tight as 1 μ M, the protein having a sequence that is less than 20% identical to TNF- α receptor (for example, a naturally-occurring TNF- α receptor, such as a mammalian or human TNF- α receptor).

[0034] In preferred embodiments, this protein includes a mutated fibronectin type III domain and the protein is mutated in the DE, BC, and FG loops. Preferably, the mutated FG loop is the same length as the wild-type FG loop. In other preferred embodiments, the protein includes an immunoglobulin-like fold (preferably, having a molecular mass less than 10 kD, greater than 7.5 kD, or between 7.5-10 kD) that consists of approximately 50-150 amino acids, and preferably, approximately 50 amino acids.

[0035] The TNF- α binders according to the invention bind TNF- α with a K_d at least as tight as 1 μ M, preferably, at least as tight as 500 nM, 100 nM, or 10 nM, more preferably, at least as tight as 1 nM or 500 pM, and, most preferably, at least as tight as 100 pM or even 20 pM. Preferably, these proteins contain one, two, or three mutated loops, and at least one, and preferably two or all three of the loops, contribute to the binding of the non-antibody protein to TNF- α . In other preferred embodiments, the non-antibody protein has at least one disulfide bond, and the non-antibody protein is a monomer or dimer under physiological conditions.

[0036] The TNF- α binders may be immobilized on a solid support (for example, a chip or bead), and may be part of an array. In addition, any of the TNF- α binders may be joined to a heterologous protein (for example, a heterologous protein that is an antibody or an antibody domain that does not bind TNF- α , an immunoglobulin F_c domain, a complement protein, or an albumin protein).

[0037] If desired, the protein may include a mutated fibronectin type III domain (for example, one derived from a human fibronectin type III domain, such as a mutated tenth module of the fibronectin type III domain (¹⁰Fn3)). In addition, the protein may lack an ¹⁰Fn3 integrin-binding motif. TNF- α binders preferably include a non-naturally occurring sequence in a loop of ¹⁰Fn3 (for example, the loop sequence PW(A/G), and may include a non-naturally occurring sequence in a β -sheet of ¹⁰Fn3. Particularly preferred TNF- α binders of the invention are shown in FIG. 25 (SEQ ID NOS: 34-140).

[0038] In addition, in related aspects, the invention features nucleic acids encoding any of the TNF- α binding proteins of the invention, as well as a loop structure on any protein that includes any one of the amino acid sequences of FIG. 25 (SEQ ID NOS: 34-140).

[0039] As used herein, by "non-antibody protein" is meant a protein that is not produced by the B cells of a mammal either naturally or following immunization of a mammal. This term also excludes antibody fragments of more than 100 amino acids, preferably, more than 80 amino acids, and, most preferably, more than 50 amino acids in length.

[0040] By "immunoglobulin-like fold" is meant a protein domain of between about 80-150 amino acid residues that includes two layers of antiparallel beta-sheets, and in which the flat, hydrophobic faces of the two beta-sheets are packed against each other. Proteins according to the invention may include several immunoglobulin-like folds covalently bound or associated non-covalently into larger structures.

[0041] By "scaffold" is meant a protein used to select or design a protein framework with specific and favorable properties, such as binding. When designing proteins from the scaffold, amino acid residues that are important for the framework's favorable properties are retained, while others residues may be varied. Such a scaffold has less than 50% of the amino acid residues that vary between protein derivatives having different properties and greater than or equal to 50% of the residues that are constant between such derivatives. Most commonly, these constant residues confer the same overall three-dimensional fold to all the variant domains, regardless of their properties.

[0042] By "fibronectin type III domain" is meant a domain having 7 or 8 beta strands which are distributed between two beta sheets, which themselves pack against each other to form the core of the protein, and further containing loops which connect the beta strands to each other and are solvent exposed. There are at least three such loops at each edge of the beta sheet sandwich, where the edge is the boundary of the protein perpendicular to the direction of the beta strands. Preferably, a fibronectin type III domain includes a sequence which exhibits at least 30% amino acid identity, and preferably at least 50% amino acid identity, to the sequence encoding the structure of the ¹⁰Fn3 domain referred to as "1ttg" (ID="1ttg" (one ttg)) available from the RCSB (Research Collaboratory for Structural Bioinformatics) Protein Data Base. Sequence identity referred to in this definition is determined by the Homology program, available from Molecular Simulation (San Diego, Calif.). The invention further includes polymers of ¹⁰Fn3-related molecules, which are an extension of the use of the monomer structure, whether or not the subunits of the polypeptide are identical.

[0043] By "naturally occurring" is meant any protein that is encoded by a living organism.

[0044] By "randomized" or "mutated" is meant including one or more amino acid alterations relative to a template sequence. By "randomizing" or "mutating" is meant the process of introducing, into a sequence, such an amino acid alteration. Randomization or mutation may be accomplished through intentional, blind, or spontaneous sequence variation, generally of a nucleic acid coding sequence, and may occur by any technique, for example, PCR, error-prone PCR, or chemical DNA synthesis. By a "corresponding, non-mutated protein" is meant a protein that is identical in sequence, except for the introduced-amino acid mutations.

[0045] By a "protein" is meant any sequence of two or more amino acids, regardless of length, post-translation modification, or function. "Protein" and "peptide" are used interchangeably herein.

[0046] By "RNA" is meant a sequence of two or more covalently bonded, naturally occurring or modified ribonucleotides. One example of a modified RNA included within this term is phosphorothioate RNA.

[0047] By "DNA" is meant a sequence of two or more covalently bonded, naturally occurring or modified deoxyribonucleotides.

[0048] By a “nucleic acid” is meant any two or more covalently bonded nucleotides or nucleotide analogs or derivatives. As used herein, this term includes, without limitation, DNA, RNA, and PNA.

[0049] By “pharmaceutically acceptable” is meant a compound or protein that may be administered to an animal (for example, a mammal) without significant adverse medical consequences.

[0050] By “physiologically acceptable carrier” is meant a carrier which does not have a significant detrimental impact on the treated host and which retains the therapeutic properties of the compound with which it is administered. One exemplary physiologically acceptable carrier is physiological saline. Other physiologically acceptable carriers and their formulations are known to one skilled in the art and are described, for example, in *Remington's Pharmaceutical Sciences*, (18th edition), ed. A. Gennaro, 1990, Mack Publishing Company, Easton, Pa., incorporated herein by reference.

[0051] By a “fusion protein” is meant a protein that includes (i) a scaffold protein of the invention joined to (ii) a second, different (i.e., “heterologous”) protein. “Fusion proteins” are distinguished from “nucleic acid-protein fusions” and “RNA-protein fusions” in that a “fusion protein” is composed entirely of amino acids, while both a “nucleic acid-protein fusion” and an “RNA-protein fusion” include a stretch of nucleic acids (the nucleic acid or RNA component) joined to a stretch of amino acids (the protein component).

[0052] By “selecting” is meant substantially partitioning a molecule from other molecules in a population. As used herein, a “selecting” step provides at least a 2-fold, preferably, at least a 30-fold, more preferably, at least a 100-fold, and, most preferably, at least a 1000-fold enrichment of a desired molecule relative to undesired molecules in a population following the selection step. A selection step may be repeated any number of times, and different types of selection steps may be combined in a given approach.

[0053] By “binding partner,” as used herein, is meant any molecule which has a specific, covalent or non-covalent affinity for a portion of a desired compound (for example, protein) of interest. Examples of binding partners include, without limitation, members of antigen/antibody pairs, protein/inhibitor pairs, receptor/ligand pairs (for example cell surface receptor/ligand pairs, such as hormone receptor/peptide hormone pairs), enzyme/substrate pairs (for example, kinase/substrate pairs), lectin/carbohydrate pairs, oligomeric or heterooligomeric protein aggregates, DNA binding protein/DNA binding site pairs, RNA/protein pairs, and nucleic acid duplexes, heteroduplexes, or ligated strands, as well as any molecule which is capable of forming one or more covalent or non-covalent bonds (for example, disulfide bonds) with any portion of another molecule (for example, a compound or protein).

[0054] By a “solid support” is meant, without limitation, any column (or column material), bead, test tube, microtiter dish, solid particle (for example, agarose or sepharose), microchip (for example, silicon, silicon-glass, or gold chip), or membrane (for example, an inorganic membrane, nitrocellulose, or the membrane of a liposome or vesicle) to which an antibody mimic or an affinity complex may be bound, either directly or indirectly (for example, through other binding partner intermediates such as other antibodies or Protein A), or in which an antibody mimic or an affinity complex may be embedded (for example, through a receptor or channel).

[0055] The present invention provides a number of advantages. For example, as described in more detail below, the present antibody mimics exhibit improved biophysical properties, such as stability under reducing conditions and solubility at high concentrations. In addition, these molecules may be readily expressed and folded in prokaryotic systems, such as *E. coli*, in eukaryotic systems, such as yeast, and in in vitro translation systems, such as the rabbit reticulocyte lysate system. Moreover, these molecules are extremely amenable to affinity maturation techniques involving multiple cycles of selection, including in vitro selection using RNA-protein fusion technology (Roberts and Szostak, Proc. Natl. Acad. Sci USA 94:12297, 1997; Szostak et al., U.S. Ser. No. 09/007,005 and U.S. Ser. No. 09/247,190; Szostak et al. WO98/31700), phage display (see, for example, Smith and Petrenko, Chem. Rev. 97:317, 1997), and yeast display systems (see, for example, Boder and Wittrup, Nature Biotech. 15:553, 1997).

[0056] Other features and advantages of the present invention will be apparent from the following detailed description thereof, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0057] FIG. 1 is a photograph showing a comparison between the structures of antibody heavy chain variable regions from camel (dark blue) and llama (light blue), in each of two orientations.

[0058] FIG. 2 is a photograph showing a comparison between the structures of the camel antibody heavy chain variable region (dark blue), the llama antibody heavy chain variable region (light blue), and a fibronectin type III module number 10 (¹⁰Fn3) (yellow).

[0059] FIG. 3 is a photograph showing a fibronectin type III module number 10 (¹⁰Fn3), with the loops corresponding to the antigen-binding loops in IgG heavy chains highlighted in red.

[0060] FIG. 4 is a graph illustrating a sequence alignment between a fibronectin type III protein domain and related protein domains.

[0061] FIG. 5 is a photograph showing the structural similarities between a ¹⁰Fn3 domain and 15 related proteins, including fibronectins, terascins, collagens, and undulin. In this photograph, the regions are labeled as follows: constant, dark blue; conserved, light blue; neutral, white; variable, red; and RGD integrin-binding motif (variable), yellow.

[0062] FIG. 6 is a photograph showing space filling models of fibronectin III modules 9 and 10, in each of two different orientations. The two modules and the integrin binding loop (RGD) are labeled. In this figure, blue indicates positively charged residues, red indicates negatively charged residues, and white indicates uncharged residues.

[0063] FIG. 7 is a photograph showing space filling models of fibronectin III modules 7-10, in each of three different orientations. The four modules are labeled. In this figure, blue indicates positively charged residues, red indicates negatively charged residues, and white indicates uncharged residues.

[0064] FIG. 8 is a photograph illustrating the formation, under different salt conditions, of RNA-protein fusions which include fibronectin type III domains.

[0065] FIG. 9 is a series of photographs illustrating the selection of fibronectin type III domain-containing RNA-protein fusions, as measured by PCR signal analysis.

[0066] FIG. 10 is a graph illustrating an increase in the percent TNF- α binding during the selections described herein, as well as a comparison between RNA-protein fusion and free protein selections.

[0067] FIG. 11 is a series of schematic representations showing IgG, $^{10}\text{Fn3}$, $\text{Fn-CH}_1\text{—CH}_2\text{—CH}_3$, and $\text{Fn-CH}_2\text{—CH}_3$ (clockwise from top left).

[0068] FIG. 12 is a photograph showing a molecular model of $\text{Fn-CH}_1\text{—CH}_2\text{—CH}_3$ based on known three-dimensional structures of IgG (X-ray crystallography) and $^{10}\text{Fn3}$ (NMR and X-ray crystallography).

[0069] FIG. 13 is a graph showing the time course of an exemplary $^{10}\text{Fn3}$ -based nucleic acid-protein fusion selection of TNF- α binders. The proportion of nucleic acid-protein fusion pool (open diamonds) and free protein pool (open circles) that bound to TNF- α -Sephacrose, and the proportion of free protein pool (full circles) that bound to underivatized Sephacrose, are shown.

[0070] FIGS. 14 and 15 are graphs illustrating TNF- α binding by TNF- α Fn-binders. In particular, these figures show mass spectra data obtained from a $^{10}\text{Fn3}$ fusion chip and non-fusion chip, respectively.

[0071] FIGS. 16 and 17 are the phosphorimage and fluorescence scan, respectively, of an $^{10}\text{Fn3}$ array, illustrating TNF- α binding.

[0072] FIG. 18 is a graph showing an alignment of the primary sequences of the llama V_H domain and the wild-type human $^{10}\text{Fn3}$ domain. Homologous residues between the two sequences are indicated. The $^{10}\text{Fn3}$ residues outside the randomized loops that were found to have mutated in approximately 45% of the selected clones are marked with arrows under the wild-type $^{10}\text{Fn3}$ sequence and with the letter that identifies the selected residue.

[0073] FIG. 19 shows schematic representations of the llama V_H domain and the wild-type human $^{10}\text{Fn3}$ domain. The locations of the mutated framework residues are indicated.

[0074] FIG. 20 is a graph illustrating the efficiency and specificity of binding of a free-protein pool translated from the original library (R0) and after ten rounds of selection with TNF- α (R10). Protein pool binding to underivatized Sephacrose, to TNF- α -Sephacrose, to IL-1 α -Sephacrose, and to IL-13-Sephacrose is compared.

[0075] FIG. 21 is a series of IgG-like scaffolds for the display of up to three loops.

[0076] FIG. 22 is a series of IgG-like scaffolds for the display of up to four, or even six, loops.

[0077] FIG. 23 is a series of scaffolds, unrelated to IgG, for the display of loop structures.

[0078] FIGS. 24A-24D are photographic and graphic illustrations demonstrating the specific capture of a target (TNF- α) by a mimic immobilized on a solid surface.

[0079] FIG. 25 is a graph listing exemplary TNF- α binders (SEQ ID NOS: 33-140) according to the invention.

DETAILED DESCRIPTION

[0080] The novel antibody mimics described herein have been designed to be superior both to antibody-derived fragments and to non-antibody frameworks, for example, those frameworks cited above.

[0081] The major advantage of these antibody mimics over antibody fragments is structural. These antibody mimics are derived from whole, stable, and soluble structural scaffolds. For example, the Fn3 scaffold is found in the human body.

Consequently, they exhibit better folding and thermostability properties than antibody fragments, whose creation involves the removal of parts of the antibody native fold, often exposing amino acid residues that, in an intact antibody, would be buried in a hydrophobic environment, such as an interface between variable and constant domains. Exposure of such hydrophobic residues to solvent increases the likelihood of aggregation of the antibody fragments.

[0082] In addition, the scaffolds described herein have no disulfide bonds, which have been reported to retard or prevent proper folding of antibody fragments under certain conditions. Since the present scaffolds do not rely on disulfides for native fold stability, they are stable under reducing conditions, unlike antibodies and their fragments which unravel upon disulfide bond reduction.

[0083] Moreover, these scaffolds provide the functional advantages of antibody molecules. In particular, despite the fact that the $^{10}\text{Fn3}$ module is not an immunoglobulin, its overall fold is close to that of the variable region of the IgG heavy chain (FIG. 2), making it possible to display the three fibronectin loops analogous to CDRs in relative orientations similar to those of native antibodies. Because of this structure, the present antibody mimics possess antigen binding properties that are similar in nature and affinity to those of antibodies, and a loop randomization and shuffling strategy may be employed in vitro that is similar to the process of affinity maturation of antibodies in vivo.

[0084] There are now described below exemplary scaffolds, for example, fibronectin-based scaffolds, and their use for identifying, selecting, and evolving novel binding proteins as well as their target ligands. These examples are provided for the purpose of illustrating, and not limiting, the invention.

$^{10}\text{Fn3}$ Structural Motif

[0085] Preferred antibody mimics of the present invention are based on the structure of a fibronectin module of type III (Fn3), a common domain found in mammalian blood and structural proteins. This domain occurs more than 400 times in the protein sequence database and has been estimated to occur in 2% of the proteins sequenced to date, including fibronectins, tenascin, intracellular cytoskeletal proteins, and prokaryotic enzymes (Bork and Doolittle, *Proc. Natl. Acad. Sci. USA* 89:8990, 1992; Bork et al., *Nature Biotech.* 15:553, 1997; Meinke et al., *J. Bacteriol.* 175:1910, 1993; Watanabe et al., *J. Biol. Chem.* 265:15659, 1990). A particular scaffold is the tenth module of human Fn3 ($^{10}\text{Fn3}$), which comprises 94 amino acid residues. The overall fold of this domain is closely related to that of the smallest functional antibody fragment, the variable region of the heavy chain, which comprises the entire antigen recognition unit in camel and llama IgG (FIGS. 1, 2). The major differences between camel and llama domains and the $^{10}\text{Fn3}$ domain are that (i) $^{10}\text{Fn3}$ has fewer beta strands (seven vs. nine) and (ii) the two beta sheets packed against each other are connected by a disulfide bridge in the camel and llama domains, but not in $^{10}\text{Fn3}$.

[0086] The three loops of $^{10}\text{Fn3}$ corresponding to the antigen-binding loops of the IgG heavy chain run between amino acid residues 21-31 (BC), 51-56 (DE), and 76-88 (FG) (FIG. 3). The length of the BC and DE loop, 10 and 6 residues, respectively, fall within the narrow range of the corresponding antigen-recognition loops found in antibody heavy chains, that is, 7-10 and 4-8 residues, respectively. Accordingly, once randomized and selected for high antigen affinity,

these two loops may make contacts with antigens equivalent to the contacts of the corresponding loops in antibodies.

[0087] In contrast, the FG loop of $^{10}\text{Fn3}$ is 12 residues long, whereas the corresponding loop in antibody heavy chains ranges from 4-28 residues. To optimize antigen binding, therefore, the length of the FG loop of $^{10}\text{Fn3}$ is preferably randomized in length as well as in sequence to cover the CDR3 range of 4-28 residues to obtain the greatest possible flexibility and affinity in antigen binding. Indeed, in general, the lengths as well as the sequences of the CDR-like loops of the antibody mimics may be randomized during in vitro or in vivo affinity maturation (as described in more detail below).

[0088] The tenth human fibronectin type III domain, $^{10}\text{Fn3}$, refolds rapidly even at low temperature; its backbone conformation has been recovered within 1 second at 5° C. Thermodynamic stability of $^{10}\text{Fn3}$ is high ($\Delta G_{\text{fold}}=24 \text{ kJ/mol}=5.7 \text{ kcal/mol}$), correlating with its high melting temperature of 110° C.

[0089] One of the physiological roles of $^{10}\text{Fn3}$ is as a subunit of fibronectin, a glycoprotein that exists in a soluble form in body fluids and in an insoluble form in the extracellular matrix (Dickinson et al., *J. Mol. Biol.* 236:1079, 1994). A fibronectin monomer of 220-250 kD contains 12 type I modules, two type II modules, and 17 fibronectin type III modules (Potts and Campbell, *Curr. Opin. Cell Biol.* 6:648, 1994). Different type III modules are involved in the binding of fibronectin to integrins, heparin, and chondroitin sulfate. $^{10}\text{Fn3}$ was found to mediate cell adhesion through an integrin-binding Arg-Gly-Asp (RGD) motif on one of its exposed loops. Similar RGD motifs have been shown to be involved in integrin binding by other proteins, such as fibrinogen, von Willebrand factor, and vitronectin (Hynes et al., *Cell* 69:11, 1992). No other matrix- or cell-binding roles have been described for $^{10}\text{Fn3}$.

[0090] The observation that $^{10}\text{Fn3}$ has only slightly more adhesive activity than a short peptide containing RGD is consistent with the conclusion that the cell-binding activity of $^{10}\text{Fn3}$ is localized in the RGD peptide rather than distributed throughout the $^{10}\text{Fn3}$ structure (Baron et al., *Biochemistry* 31:2068, 1992). The fact that $^{10}\text{Fn3}$ without the RGD motif is unlikely to bind to other plasma proteins or extracellular matrix makes $^{10}\text{Fn3}$ a useful scaffold to replace antibodies. In addition, the presence of $^{10}\text{Fn3}$ in natural fibrinogen in the bloodstream suggests that $^{10}\text{Fn3}$ itself is unlikely to be immunogenic in the organism of origin.

[0091] In addition, we have determined that the $^{10}\text{Fn3}$ framework possesses exposed loop sequences tolerant of randomization, facilitating the generation of diverse pools of antibody mimics. This determination was made by examining the flexibility of the $^{10}\text{Fn3}$ sequence. In particular, the human $^{10}\text{Fn3}$ sequence was aligned with the sequences of fibronectins from other sources as well as sequences of related proteins (FIG. 4), and the results of this alignment were mapped onto the three-dimensional structure of the human $^{10}\text{Fn3}$ domain (FIG. 5). This alignment revealed that the majority of conserved residues are found in the core of the beta sheet sandwich, whereas the highly variable residues are located along the edges of the beta sheets, including the N- and C-termini, on the solvent-accessible faces of both beta sheets, and on three solvent-accessible loops that serve as the hyper-variable loops for affinity maturation of the antibody mimics. In view of these results, the randomization of these three loops are unlikely to have an adverse effect on the overall fold or stability of the $^{10}\text{Fn3}$ framework itself.

[0092] For the human $^{10}\text{Fn3}$ sequence, this analysis indicates that, at a minimum, amino acids 1-9, 44-50, 61-54, 82-94 (edges of beta sheets); 19, 21, 3046 (even), 79-65 (odd) (solvent-accessible faces of both beta sheets); 21-31, 51-56, 76-88 (CDR-like solvent-accessible loops); and 14-16 and 36-45 (other solvent-accessible loops and beta turns) may be randomized to evolve new or improved compound-binding proteins. In addition, as discussed above, alterations in the lengths of one or more solvent exposed loops may also be included in such directed evolution methods.

[0093] Alternatively, changes in the β -sheet sequences may also be used to evolve new proteins. These mutations change the scaffold and thereby indirectly alter loop structure(s). If this approach is taken, mutations should not saturate the sequence, but rather few mutations should be introduced. Preferably, no more than between 3-20 changes should be introduced to the β -sheet sequences by this approach.

[0094] Sequence variation may be introduced by any technique including, for example, mutagenesis by Taq polymerase (Tindall and Kunkel, *Biochemistry* 27:6008 (1988)), fragment recombination, or a combination thereof. Similarly, an increase of the structural diversity of libraries, for example, by varying the length as well as the sequence of the CDR-like loops, or by structural redesign based on the advantageous framework mutations found in selected pools, may be used to introduce further improvements in antibody mimic scaffolds.

Antibody Mimic Fusions

[0095] The antibody mimics described herein may be fused to other protein domains. For example, these mimics may be integrated with the human immune response by fusing the constant region of an IgG (F_c) with an antibody mimic, such as an $^{10}\text{Fn3}$ module, preferably through the C-terminus of $^{10}\text{Fn3}$. The F_c in such a $^{10}\text{Fn3}$ - F_c fusion molecule activates the complement component of the immune response and increases the therapeutic value of the antibody mimic. Similarly, a fusion between an antibody mimic, such as $^{10}\text{Fn3}$, and a complement protein, such as C1q, may be used to target cells, and a fusion between an antibody mimic, such as $^{10}\text{Fn3}$, and a toxin may be used to specifically destroy cells that carry a particular antigen. In addition, an antibody scaffold, such as $^{10}\text{Fn3}$, in any form may be fused with albumin to increase its half-life in the bloodstream and its tissue penetration. Any of these fusions may be generated by standard techniques, for example, by expression of the fusion protein from a recombinant fusion gene constructed using publically available gene sequences.

Scaffold Multimers

[0096] In addition to monomers, any of the scaffold constructs described herein may be generated as dimers or multimers of antibody mimics as a means to increase the valency and thus the avidity of antigen binding. Such multimers may be generated through covalent binding. For example, individual $^{10}\text{Fn3}$ modules may be bound by imitating the natural $^8\text{Fn3}$ - $^9\text{Fn3}$ - $^{10}\text{Fn3}$ C-to-N-terminus binding or by imitating antibody dimers that are held together through their constant regions. A $^{10}\text{Fn3}$ -Fc construct may be exploited to design dimers of the general scheme of $^{10}\text{Fn3}$ -Fc::Fc- $^{10}\text{Fn3}$. The bonds engineered into the Fc::Fc interface may be covalent or non-covalent. In addition, dimerizing or multimerizing part-

ners other than Fc can be used in hybrids, such as $^{10}\text{Fn3}$ hybrids, to create such higher order structures.

[0097] In particular examples, covalently bonded multimers may be generated by constructing fusion genes that encode the multimer or, alternatively, by engineering codons for cysteine residues into monomer sequences and allowing disulfide bond formation to occur between the expression products. Non-covalently bonded multimers may also be generated by a variety of techniques. These include the introduction, into monomer sequences, of codons corresponding to positively and/or negatively charged residues and allowing interactions between these residues in the expression products (and therefore between the monomers) to occur. This approach may be simplified by taking advantage of charged residues naturally present in a monomer subunit, for example, the negatively charged residues of fibronectin. Another means for generating non-covalently bonded antibody mimics is to introduce, into the monomer gene (for example, at the amino- or carboxy-termini), the coding sequences for proteins or protein domains known to interact. Such proteins or protein domains include coil-coil motifs, leucine zipper motifs, and any of the numerous protein subunits (or fragments thereof) known to direct formation of dimers or higher order multimers.

Fibronectin-Like Molecules

[0098] Although $^{10}\text{Fn3}$ represents a preferred scaffold for the generation of antibody mimics, other molecules may be substituted for $^{10}\text{Fn3}$ in the molecules described herein. These include, without limitation, human fibronectin modules $^1\text{Fn3}$ - $^9\text{Fn3}$ and $^{11}\text{Fn3}$ - $^{17}\text{Fn3}$ as well as related Fn3 modules from non-human animals and prokaryotes. In addition, Fn3 modules from other proteins with sequence homology to $^{10}\text{Fn3}$, such as tenascins and undulins, may also be used. Other exemplary scaffolds having immunoglobulin-like folds (but with sequences that are unrelated to the V_H domain) are shown in FIG. 21 and include N-cadherin, ICAM-2, titin, GCSF receptor, cytokine receptor, glycosidase inhibitor, E-cadherin, and antibiotic chromoprotein. Yet further domains with related structures may be derived from myelin membrane adhesion molecule P0, CD8, CD4, CD2, class I MHC, T-cell antigen receptor, CD1, C2 and I-set domains of VCAM-1, I-set immunoglobulin domain of myosin-binding protein C, I-set immunoglobulin domain of myosin-binding protein H, I-set immunoglobulin domain of telokin, telikin, NCAM, twitchin, neuroglian, growth hormone receptor, erythropoietin receptor, prolactin receptor, GC-SF receptor, interferon-gamma receptor, β -galactosidase/glucuronidase, β -glucuronidase, and transglutaminase. Alternatively, any other protein that includes one or more immunoglobulin-like folds may be utilized. Such proteins may be identified, for example, using the program SCOP (Murzin et al., *J. Mol. Biol.* 247:536 (1995); Lo Conte et al., *Nucleic Acids Res.* 25:257 (2000)).

[0099] Generally, any molecule that exhibits a structural relatedness to the V_H domain (as identified, for example, using the computer program above) may be utilized as an antibody mimic. Such molecules may, like fibronectin, include three loops at the N-terminal pole of the molecule and three loops at the C-terminal pole, each of which may be randomized to create diverse libraries; alternatively, larger domains may be utilized, having larger numbers of loops, as long as a number of such surface randomizable loops are positioned closely enough in space that they can participate in

antigen binding. FIG. 22 shows examples of useful domains having more than three loops positioned close to each other. These examples include T-cell antigen receptor and superoxide dismutase, which each have four loops that can be randomized; and an Fn3 dimer, tissue factor domains, and cytokine receptor domains, each of which have three sets of two similar domains where three randomizable loops are part of the two domains (bringing the total number of loops to six).

[0100] In yet another alternative, any protein having variable loops positioned close enough in space may be utilized for candidate binding protein production. For example, large proteins having spatially related, solvent accessible loops may be used, even if unrelated structurally to an immunoglobulin-like fold. Exemplary proteins include, without limitation, cytochrome F, green fluorescent protein, GroEL, and thaumatin (FIG. 23). The loops displayed by these proteins may be randomized and superior binders selected from a randomized library as described herein. Because of their size, molecules may be obtained that exhibit an antigen binding surface considerably larger than that found in an antibody-antigen interaction. Other useful scaffolds of this type may also be identified using the program SCOP (Murzin et al., *J. Mol. Biol.* 247:536 (1995)) to browse among candidate proteins having numerous loops, particularly loops positioned among parallel beta sheets or a number of alpha-helices.

[0101] Modules from different organisms and parent proteins may be most appropriate for different applications. For example, in designing an antibody mimic, it may be most desirable to generate that protein from a fibronectin or fibronectin-like molecule native to the organism for which a therapeutic is intended. In contrast, the organism of origin is less important or even irrelevant for antibody mimics that are to be used for in vitro applications, such as diagnostics, or as research reagents.

[0102] For any of these molecules, libraries may be generated and used to select binding proteins by any of the methods described herein.

Directed Evolution of Scaffold-Based Binding Proteins

[0103] The antibody mimics described herein may be used in any technique for evolving new or improved binding proteins. In one particular example, the target of binding is immobilized on a solid support, such as a column resin or microtiter plate well, and the target contacted with a library of candidate scaffold-based binding proteins. Such a library may consist of antibody mimic clones, such as $^{10}\text{Fn3}$ clones constructed from the wild type $^{10}\text{Fn3}$ scaffold through randomization of the sequence and/or the length of the $^{10}\text{Fn3}$ CDR-like loops. If desired, this library may be an RNA-protein fusion library generated, for example, by the techniques described in Szostak et al., U.S. Ser. No. 09/007,005 and Ser. No. 09/247,190; Szostak et al., WO98/31700; and Roberts & Szostak, *Proc. Natl. Acad. Sci. USA* (1997) vol. 94, p. 12297-12302. Alternatively, it may be a DNA-protein library (for example, as described in Lohse, *DNA-Protein Fusions and Uses Thereof*, U.S. Ser. No. 60/110,549, U.S. Ser. No. 09/459,190, and WO00/32823). The fusion library is incubated with the immobilized target, the support is washed to remove non-specific binders, and the tightest binders are eluted under very stringent conditions and subjected to PCR to recover the sequence information or to create a new library of binders which may be used to repeat the selection process, with or without further mutagenesis of the sequence. A num-

ber of rounds of selection may be performed until binders of sufficient affinity for the antigen are obtained.

[0104] In one particular example, the ¹⁰F_n3 scaffold may be used as the selection target. For example, if a protein is required that binds a specific peptide sequence presented in a ten residue loop, a single ¹⁰F_n3 clone is constructed in which one of its loops has been set to the length of ten and to the desired sequence. The new clone is expressed in vivo and purified, and then immobilized on a solid support. An RNA-protein fusion library based on an appropriate scaffold is then allowed to interact with the support, which is then washed, and desired molecules eluted and re-selected as described above.

[0105] Similarly, the scaffolds described herein, for example, the ¹⁰F_n3 scaffold, may be used to find natural proteins that interact with the peptide sequence displayed by the scaffold, for example, in an ¹⁰F_n3 loop. The scaffold protein, such as the ¹⁰F_n3 protein, is immobilized as described above, and an RNA-protein fusion library is screened for binders to the displayed loop. The binders are enriched through multiple rounds of selection and identified by DNA sequencing.

[0106] In addition, in the above approaches, although RNA-protein libraries represent exemplary libraries for directed evolution, any type of scaffold-based library may be used in the selection methods of the invention.

Use

[0107] The antibody mimics described herein may be evolved to bind any antigen of interest. These proteins have thermodynamic properties superior to those of natural antibodies and can be evolved rapidly in vitro. Accordingly, these antibody mimics may be employed in place of antibodies in all areas in which antibodies are used, including in the research, therapeutic, and diagnostic fields. In addition, because these scaffolds possess solubility and stability properties superior to antibodies the antibody mimics described herein may also be used under conditions which would destroy or inactivate antibody molecules. Finally, because the scaffolds of the present invention may be evolved to bind virtually any compound, these molecules provide completely novel binding proteins which also find use in the research, diagnostic, and therapeutic areas.

Experimental Results

[0108] Exemplary scaffold molecules described above were generated and tested, for example, in selection protocols, as follows.

Library Construction

[0109] A complex library was constructed from three fragments, each of which contained one randomized area corresponding to a CDR-like loop. The randomized residues are indicated in FIG. 18 as underlined sequences, specifically, residues 23-29 of the ¹⁰F_n3 BC loop (corresponding to CDR-H1 of the llama V_H); residues 52-55 of the ¹⁰F_n3 DE loop (corresponding to CDR-H2 of the llama V_H); and residues 78-87 of the ¹⁰F_n3 FG loop (corresponding to CDR-H3 of the llama V_H). The fragments were named BC, DE, and FG based on the names of the CDR-H-like loops contained within them; in addition to ¹⁰F_n3 and a randomized sequence, each of the

fragments contained stretches encoding an N-terminal HiS₆ domain or a C-terminal FLAG peptide tag. At each junction between two fragments (i.e., between the BC and DE fragments or between the DE and FG fragments), each DNA fragment contained recognition sequences for the EarI Type IIS restriction endonuclease. This restriction enzyme allowed the splicing together of adjacent fragments while removing all foreign, non-¹⁰F_n3, sequences. It also allowed for a recombination-like mixing of the three ¹⁰F_n3 fragments between cycles of mutagenesis and selection.

[0110] The wild-type, human ¹⁰F_n3 gene was cloned from a human liver library (Maxim Biotech, South San Francisco, Calif.) using the primers Hu5PCR-NdeI 5'CATATGGTCTGATGTTCCGAGG3'; SEQ ID NO: 28) and Hu3PCR-EcoRI (5'GAATTCCTATGTTCCGTAATTAATGGAATTG3'; SEQ ID NO: 29). Three different libraries were constructed from the wild-type segments obtained by the PCR of the ¹⁰F_n3 clone and from randomized segments obtained by oligonucleotide synthesis. The BC_r-DE_r-FG_r library was obtained by randomizing the selected residues in BC, DE, and FG loops; the BC_r-DE_{wt}-FG_r library was obtained by randomizing the selected residues in BC and FG loops, leaving the DE loop sequence wild-type; and the BC_{wt}-DE_{wt}-FG_r library was obtained by randomizing the selected residues in the FG loop only.

[0111] The BC_r, DE_r, and FG_r fragments were made synthetically. Each fragment was assembled from two overlapping oligonucleotides, which were first annealed, then extended to form the double-stranded DNA form of the fragment. The oligonucleotides that were used to construct and process the three fragments are listed below; the "Top" and "Bottom" species for each fragment are the oligonucleotides that contained the entire ¹⁰F_n3 encoding sequence. In these oligonucleotide designations, "N" indicates A, T, C, or G; and "S" indicates C or G.

Hf_nLbcTop (His) :
(SEQ ID NO: 1)
5'-GG AAT TCC TAA TAC GAC TCA CTA TAG GGA CAA
TTA CTA TTT ACA ATT ACA ATG CAT CAC CAT CAC CAT
CAC GTT TCT GAT GTT CCG AGG GAC CTG GAA GTT GTT
GCT GCG ACC CCC ACC AGC-3'

Hf_nLbcTop (an alternative N-terminus):
(SEQ ID NO: 2)
5'-GG AAT TCC TAA TAC GAC TCA CTA TAG GGA CAA
TTA CTA TTT ACA ATT ACA ATG GTT TCT GAT GTT CCG
AGG GAC CTG GAA GTT GTT GCT GCG ACC CCC ACC AGC-3'

Hf_nLBCBot-flag8:
(SEQ ID NO: 3)
5'-AGC GGA TGC CTT GTC GTC GTC GTC CTT GTA GTC
GCT CTT CCC TGT TTC TCC GTA AGT GAT CCT GTA ATA
TCT (SNN)₇ CCA GCT GAT CAG TAG GCT GGT GGG GGT CGC
AGC-3'

Hf_nBC3'-flag8:
(SEQ ID NO: 4)
5'-AGC GGA TGC CTT GTC GTC GTC GTC CTT GTA GTC GCT
CTT CCC TGT TTC TCC GTA AGT GAT CC-3'

-continued

HFnlDETop: (SEQ ID NO: 5)
 5'-GG AAT TCC TAA TAC GAC TCA CTA TAG GGA CAA
 TTA CTA TTT ACA ATT ACA ATG CAT CAC CAT CAC CAT
 CAC CTC TTC ACA GGA GGA AAT AGC CCT GTC C-3'

HFnlDEBot-flag8: (SEQ ID NO: 6)
 5'-AGC GGA TGC CTT GTC GTC GTC CTT GTA GTC GCT
 CTT CGT ATA ATC AAC TCC AGG TTT AAG GCC GCT GAT
 GGT AGC TGT (SNN)₄ AGG CAC AGT GAA CTC CTG GAC AGG
 GCT ATT TCC TCC TGT-3'

HFnlDE3'-flag8: (SEQ ID NO: 7)
 5'-AGC GGA TGC CTT GTC GTC GTC GTC CTT GTA GTC GCT
 CTT CGT ATA ATC AAC TCC AGG TTT AAG G-3'

HFnlFGTop: (SEQ ID NO: 8)
 5'-GG AAT TCC TAA TAC GAC TCA CTA TAG GGA CAA TTA
 CTA TTT ACA ATT ACA ATG CAT CAC CAT CAC CAT CAC
 CTC TTC TAT ACC ATC ACT GTG TAT GCT GTC-3'

HFnlFGBot-flag8: (SEQ ID NO: 9)
 5'-AGC GGA TGC CTT GTC GTC GTC CTT GTA GTC TGT
 TCG GTA ATT AAT GGA AAT TGG (SNN)₁₀ AGT GAC AGC
 ATA CAC AGT GAT GGT ATA-3'

HFnlFG3'-flag8: (SEQ ID NO: 10)
 5'-AGC GGA TGC CTT GTC GTC GTC CTT GTA GTC TGT
 TCG GTA ATT AAT GGA AAT TGG-3'

T7Tmv (introduces T7 promoter and TMV untranslated region needed for in vitro translation): (SEQ ID NO: 11)
 5'-GCG TAA TAC GAC TCA CTA TAG GGA CAA TTA CTA TTT
 ACA ATT ACA-3'

ASaflag8: (SEQ ID NO: 12)
 5'-AGC GGA TGC CTT GTC GTC GTC CTT GTA GTC-3'

Unispl-s (spint oligonucleotide used to ligate mRNA to the puromycin-containing linker, described by Roberts et al, 1997, supra): (SEQ ID NO: 13)
 5'-TTTTTTTTTAGCGGATGC-3'

A18--2PEG (DNA-puromycin linker): (SEQ ID NO: 14)
 5'-(A)₁₈(PEG)₂CCPur

[0112] The oligonucleotide pair BC_{Top} and BC_{Bot-flag8} was used to construct the fragment which contains the randomized BC loop; the pair DE_{Top} and DE_{Bot-flag8} was used to construct the fragment which contains the randomized DE loop; the pair BC_{Top} and DE_{3-Flag8} was used to PCR-amplify the BC_{wt}-DE_{wt} fragments; and the pair FG_{Top} and FG_{Bot-Flag8} was used to construct the fragment which contains the randomized FG loop. The pairs of oligonucleotides (500 pmol of

each) were annealed in 100 μ L of 10 mM Tris 7.5, 50 mM NaCl for 10 minutes at 85° C., followed by a slow (0.5-1 hour) cooling to room temperature. The annealed fragments with single-stranded overhangs were then extended using 100 U Klenow (New England Biolabs, Beverly, Mass.) for each 100 μ L aliquot of annealed oligos, and the buffer made of 838.5 μ L H₂O, 9 μ L 1 M Tris 7.5, 5 μ L 1M MgCl₂, 20 μ L 10 mM dNTPs, and 7.5 μ L 1M DTT. The extension reactions proceeded for 1 hour at 25° C.

[0113] In order to reduce the frequency of stop codons introduced by the random sequences, the randomized residues were encoded by (NNS)_n, where N stands for any nucleotide and S for an equimolar mixture of C and G; only one of the three stop codons (TAG) conforms to the NNS restriction. In addition to the sequence encoding ¹⁰F_n3, the gene fragments contained the 5' Tobacco Mosaic Virus (TMV) untranslated region and the T7 promoter, as well as the sequences encoding a 5' hexahistidine protein purification tag and a 3'FLAG epitope purification tag. In addition, as noted above, Ear I restriction endonuclease recognition sites were engineered into the overlaps between adjacent fragments in order to facilitate the assembly of the three fragments.

[0114] Next, each of the double-stranded fragments was transformed into an RNA-protein fusion (PROfusion™) using the technique developed by Szostak et al., U.S. Ser. No. 09/007,005 and U.S. Ser. No. 09/247,190; Szostak et al., WO98/31700; and Roberts & Szostak, Proc. Natl. Acad. Sci. USA (1997) vol. 94, p. 12297-12302. Briefly, the fragments were transcribed using an Ambion in vitro transcription kit, T7-MEGAshortscript™ (Ambion, Austin, Tex.), and the resulting mRNA was gel-purified and ligated to a 5'-phosphorylated DNA-puromycin linker, preferably, 5' dA₁₈PEG₂dCdCPur using DNA ligase (Promega, Madison, Wis.); the mRNA was aligned with the DNA linker using a DNA splint oligonucleotide (5'TTTTTTTTINAGCG-GATGC 3'; SEQ ID NO: 30) as described in Szostak (supra). The mRNA-DNA-puromycin molecule was then translated using the Ambion rabbit reticulocyte lysate-based translation kit in the presence of ³⁵S-methionine. The resulting mRNA-DNA-puromycin-protein fusion was purified using Oligo (dT) cellulose, (Type 7, Amersham Pharmacia, Piscataway, N.J.) and a complementary DNA strand was, synthesized using reverse transcriptase (Superscript™II, Gibco, Life Technologies, Rockville, Md.) and the RT primers described above (Unisplint-S or flagASA), following the manufacturer's instructions (preferably, a two-minute annealing at 70° C. and a 40 minute reaction at 42° C.).

[0115] The RNA-protein fusion with annealing cDNA obtained for each fragment was next purified on the resin appropriate to its peptide purification tag, i.e., on Ni-NTA agarose (Qiagen, Valencia, Calif.) for the His₆-tag and M2 Anti-Flag Agarose (Sigma, St. Louis, Mo.) for the FLAG-tag, following the procedures recommended by the manufacturers. The fragment-encoding genetic information recovered by KOH elution was amplified by PCR using Pharmacia Ready-to-Go PCR Beads, 10 pmol of 5' and 3' PCR primers, and the following PCR program (Pharmacia, Piscataway, N.J.): Step 1: 95° C. for 3 minutes; Step 2: 95° C. for 30 seconds, 58/62° C. for 30 seconds, 72° C. for 1 minute, 20/25/30 cycles, as required; Step 3: 72° C. for 5 minutes; Step 4: 4° C. until end (typically, 25 cycles).

[0116] The resulting DNA was cleaved by 5-6 U EarI (New England Biolabs) per μ g DNA; the reaction took place in T4 DNA Ligase Buffer (New England Biolabs) at 37° C., for 1

hour, and was followed by an optional incubation at 70° C. for 15 minutes to inactivate Ear I. Equal amounts of the BC, DE, and FG fragments were combined and ligated to form a full-length ¹⁰Fn3 gene with randomized loops. The ligation required 10 U of fresh EarI (New England Biolabs) and 20 U of T4 DNA Ligase (Promega, Madison, Wis.), and took 1 hour at 37° C. EarI and ligase were then inactivated by a 15 minute incubation at 65° C.

[0117] Three different libraries, BC_{wt}-DE_{wt}-FG_r, BC_r-DE_{wt}-FG_r, and BC_r-DE_r-FG_r, were made in the manner described above. Each contained the form of the FG loop with 10 randomized residues. The BC and the DE loops of the first library bore the wild type ¹⁰Fn3 sequence; a BC loop with 7 randomized residues and a wild type DE loop made up the second library; and a BC loop with 7 randomized residues and a DE loop with 4 randomized residues made up the third library. The complexity of the FG loop in each of these three libraries was 10¹³; the further two randomized loops provided the potential for a complexity too large to be sampled in a laboratory. The combination of these libraries provided a master library having 10¹² unique clones.

[0118] The sequences of 76 randomly picked clones from the original, randomized, BC_r-DE_r-FG_r library showed no pattern in the randomized loops (data not shown); the amino acid frequency in the library varied in proportion to the number of codons available that encoded each residue, between 1% per position (glutamic acid, methionine, tryptophan) and 14% per position (proline). In contrast, the average probability for a residue in the preserved, beta-sheet framework to have remained as wild type was found to be 99%.

[0119] Equimolar amounts of the three libraries (2 pmoles of DNA each) were combined into one master library in order to simplify the selection process; target binding itself was expected to select the most suitable library for a particular challenge. RNA-protein fusions were obtained from the master library following the general procedure described in Szostak et al., U.S. Ser. No. 09/007,005 and Ser. No. 09/247,190; Szostak et al., WO98/31700; and Roberts & Szostak, Proc. Natl. Acad. Sci. USA (1997) vol. 94, p. 12297-12302 (FIG. 8), except that affinity purification performed in rounds three to ten used only M2-Sepharose (see below).

Fusion Selections

[0120] The master library in the RNA-protein fusion form was subjected to selection for binding to TNF-α (Pepro Tech, Rocky Hill, N.J.). Two initial protocols were employed: one in which the target was immobilized on an agarose column and one in which the target was immobilized on a BIACORE chip. First, an extensive optimization of conditions to minimize background binders to the agarose column yielded the favorable buffer conditions of 50 mM HEPES pH 7.4, 0.02% Triton, 100 μg/ml sheared salmon sperm DNA. In this buffer, the non-specific binding of the ¹⁰Fn3-RNA fusion to TNF-α Sepharose was 0.3%. The non-specific binding background of the ¹⁰Fn3-RNA/cDNA library to TNF-α Sepharose was found to be 0.1%.

[0121] During each round of selection on TNF-α Sepharose, the library was first preincubated for an hour with underivatized Sepharose to remove any remaining non-specific binders; the flow-through from this pre-clearing was incubated for another hour with TNF-α Sepharose. The TNF-α Sepharose was washed for 3-30 minutes.

[0122] After each selection, the cDNA component of the complex that had been eluted from the solid support with 0.3

M NaOH or 0.1M KOH was amplified by PCR; a DNA band of the expected size persisted through multiple rounds of selection (FIG. 9); similar results were observed in the two alternative selection protocols, and only the data from the agarose column selection is shown in FIG. 9.

[0123] In this selection, in the first seven rounds, the binding of Fn3-RNA/cDNA molecules to the target remained low; in contrast, when free protein was translated from DNA pools at different stages of the selection, the proportion of the column binding species increased significantly between rounds (FIG. 10).

[0124] In later selections, the fusion pools selected in the first eight rounds of selection (R1-8) bound to TNF-α-Sepharose at levels close to the background (<0.25%) (FIG. 13). After nine rounds of selection (R9), the binding of fusion to TNF-α-Sepharose increased sharply to 0.7%, and, after ten rounds of selection (R10), the binding increased further to 7% (FIG. 13). These selections were carried out using TNF-α immobilized on Epoxy-Activated Sepharose™ 6B (Amersham Pharmacia) at 10 mg TNF/g Sepharose in 10 mL. Before use, the TNF-α-derivatized Sepharose was blocked in Binding Buffer (50 mM HEPES, pH 7.4, 0.02% Triton, 0.1 mg/mL sheared salmon sperm DNA (Ambion)), overnight, at 4° C.

[0125] The ¹⁰Fn3-based master library was transcribed, ligated to the puromycin-bearing linker, translated into an mRNA-protein library in the presence of 5-10 μL/300 μL ³⁵S-methionine, affinity purified on Oligo(dT) Cellulose, reverse-transcribed into a DNA/mRNA-protein library, and affinity-purified on M2-Sepharose (for rounds 3-10), as described above. Forty pmol of DNA/mRNA-protein fusion library molecules, the equivalent of 20 copies of 4×10¹² different sequences, were recovered, then subjected to the first round (R1) of the selection.

[0126] In the first step of the selection, 40 pmoles of the DNA/mRNA-protein library was incubated for 1 hour at 4° C., with tumbling, in 300 μL of Binding Buffer with 30 μL of Epoxy-Sepharose that had been subjected to the derivatization procedure in the absence of TNF-α. In the second round, 24 pmol of the library was added, and in the remaining eight rounds, 0.1-2 pmol of the library was added. The supernatant was recovered by microcentrifugation through a Micro Bio-Spin® chromatography column (BIO-RAD, Hercules, Calif.), then incubated with 30 μL of TNF-α-Sepharose (6 μM) in 300 μL of the Binding Buffer for 1 hour at 4° C. (during Rounds 7-10, the Binding Buffer contained an additional 1 mg/ml of BSA). The TNF-α-Sepharose was recovered on a spin column, then washed with 3×300 μL of Binding Buffer, eluted with 100 μL of 0.1 M KOH, and finally neutralized with 1 μL of 1 M Tris 8.0, 8 μL of 1 M HCl. Samples of the library, of the TNF-α-Sepharose before and after the elution, of the washes, and of the elutions were quantified by counting ³⁵S-methionine in the sample in a scintillation counter. The next round of selection began with the formation of a new DNA/mRNA-protein pool by PCR amplification, which was transcribed, translated, and reverse-transcribed from the PCR product.

[0127] The DNA pools obtained from the elution after nine and after ten rounds were cloned into the TOPO™ TA, pCR2.1 cloning vector (Invitrogen, Carlsbad, Calif.) and transformed into *E. coli*. Between 30 and 100 clones were picked and grown into plasmid minipreps (Qiagen). Thirty-eight clones from R9 and 29 clones from R10 were picked at random and sequenced (DNA Sequencing Core Facility,

Massachusetts General Hospital, Dept. of Molecular Biology, Boston, Mass.). The program ClustalW⁶⁰ was used to align the resulting protein sequences.

Amino Acid Residue Sequences of the TNF- α Binding Clones

[0128] Thirty-eight of the 61 clones derived from R9 and from R10 had unique amino acid sequences, a surprising diversity. The ten clones that were isolated more than once, presumably because of their superior binding to TNF- α , are listed in Table 1 (full sequences in FIG. 25).

[0129] Of the 61 clones picked randomly from the winning pool, only one (clone T09.08, sequence not shown) had its origin in the BC_{wt}-DE_{wt}-FG_r library, with another six from the BC_r-DE_{wt}-FG_r library. The observation that the remaining 54 (88% of the winners) were selected from the BC_r-DE_r-FG_r library points out the importance for TNF- α cooperative binding of the target by several loops.

[0130] The most common motif found in the selected loop sequences is PWA(S/T), which is found in the DE loop of 33 of the 61 clones; the more loosely defined sequence of PW(A/G) is seen in 41/61 clones. Such a strong selection for a specific DE sequence is surprising since the analogous CDR-H2 loops of antibody V_H domains generally make only a small contribution to antigen binding. On the other hand, the short length of the DE loop, which means that 10⁷ copies of each possible tetrapeptide sequence would be expected to be present in the library, would facilitate the optimization of any contribution of the DE loop to the selected properties. A survey of other Fn3 domains (Dickenson et al., J. Mol. Biol. 236:1079-1092 (1994)) shows that proline is found at positions equivalent to the ¹⁰Fn3 residue 52 as frequently as is the wild-type glycine; similarly, alanine, glycine, and the wild-type lysine are all common at positions equivalent to the ¹⁰Fn3 position 54. In consequence, it appears likely that the selected residues at positions 52 and 54 are at least consistent with favorable biophysical properties. In contrast, no tryptophan is found at the position equivalent to the ¹⁰Fn3 residue 53, which suggests that Tryptophan 53 may have been selected for a reason specific to the present selection, such as due to a contribution to TNF- α binding. This is consistent with the absence of this motif in later selections against other antigens, again suggesting that the PWA/G motif is more likely to contribute to TNF- α binding directly than through stability or solubility of the ¹⁰Fn3 domain. The preference for the PWA/G motif on loop DE suggests another possible reason for the preference for the BC_r-DE_r-FG_r library during the selection: the BC_r-DE_r-FG_r library alone contained the randomized DE loop, and would be expected to outcompete the other two libraries if the PWA/G sequence were important to target binding.

[0131] The sequences selected most frequently in the BC loop is NRSGLQS (12/61) (SEQ ID NO: 31), whereas the sequence selected most commonly in the FG loop is AQT-GHHLHDK (6/61) (SEQ ID NO: 32). An NRSGLQS BC loop and an AQTGHHLHDK FG loop have not been found in the same molecule, but two clones were found which contain the most frequently isolated sequences on two of the three randomized loops. These clones, T10.06 (BC: NRSGLQS, DE: PWA) and T09.12 (DE: PWA, FG: AQTGHHLHDK), have two of the lowest four dissociation constants from TNF- α of the clones examined (Table 1).

[0132] Due to the use of a Taq polymerase that contains no proofreading activity, every round of PCR introduced addi-

tional random mutations into both the CDR-like loops and the beta-sheet scaffold of the ¹⁰Fn3 sequence, at the estimated rate of 0.01% per base pair, i.e., 3% per ¹⁰Fn3 gene per round of PCR and approximately 75% per round of selection. Consequently, it is likely that the residues preserved as wild-type and those preserved in a non-wild-type stable sequence indicate that such sequences were selected due to their superior properties. In the mutated loops, it is impossible to distinguish between the mutations introduced by oligonucleotide synthesis or by PCR mutagenesis, but in the beta-strand scaffold, most of the mutations selected originate from Taq errors. The selected clones showed several conserved changes in the scaffold of the protein, which had not been randomized intentionally. FIG. 18 indicates the residues in the ¹⁰Fn3 beta sheet that had not been randomized, but nevertheless mutated during selection. This mutagenesis occurred at the frequency of 26-28 of the 61 clones; these mutations are marked with arrows under the wild-type ¹⁰Fn3 sequence and with the letter that identifies the selected residue. In particular, 28 of the 61 clones mutated from Leucine 18 to Arginine or to Glutamine, and 26 clones mutated from Threonine 56 to Isoleucine. FIG. 19 shows the location of these scaffold mutations. Whereas position 56 is at the stem loop DE and thus would be expected to affect the conformation and the target-binding properties of this loop, the distance of position 18 from the presumed TNF- α -binding loops suggests that the selective advantage of this mutation may arise from an indirect effect on the conformation of loop BC or from an effect on the stability of the protein (FIG. 19). This is supported by an experiment in which clone T10.06, which contains the frequently seen L18R and T56I changes from the wild-type, was mutagenized to reverse position 18 back to the wild-type leucine. This change caused an increase of the K_d of the variant by approximately 10-fold. The weaker binding of the T10.06(L18) protein to TNF- α suggests that the residue at position 18 has an effect on the binding of the target by the CDR-like loops, possibly by a minor structural change that is transmitted through the beta-strand to loop BC.

Affinity and Specificity of the Selected TNF- α Binding Pools

[0133] The apparent average K_d values of free protein pools for TNF- α after nine and after ten rounds of selection were found to be indistinguishable (4 and 6 nM, respectively; Table 1); this similarity in affinity is consistent with the relatively low (10 fold) level of enrichment observed in the last round of selection and with the similarity in the sequence composition of the two pools. The apparent average K_d values of free protein pool after four further rounds of selection was 3 nM, also indistinguishable from those of R9 and R10 pools.

[0134] In order to assess the specificity of the binding of the pool selected after ten rounds of selection, we compared the binding of two different free protein pools to three cytokines immobilized on Sepharose to TNF- α , the target of the selection, and to IL-1 α and IL-13, which were irrelevant to the selection. The first pool had been transcribed and translated from the initial, randomized DNA library before the selection (R0), and the second pool, from the library after ten rounds of selection (R10).

[0135] To carry out these experiments the PCR product of the elution after the tenth round of selection was transcribed and translated in vitro, in the presence of ³⁵S-methionine but without forming the mRNA-protein fusion. The resulting fraction of the free protein bound to TNF- α -Sepharose, to IL-1 α -Sepharose, to IL-13-Sepharose at approximately 10

μM , 30 μM , and 50 μM , respectively, and to underivatized Sepharose was compared (FIG. 20), using the procedure described above for DNA/mRNA-protein fusion binding to TNF- α -Sepharose. The amount of the selected pool bound to each of the targets was measured by scintillation counting of the washed beads.

[0136] FIG. 20 shows that, whereas the binding of R0 to TNF- α , IL-1 α , and IL-13 was similar (2%, 4%, and 3%, respectively); the ten rounds of selection resulted in 32% binding to the targeted TNF- α , in 3% binding to IL-1 α , and in 1% binding to IL-13. The absolute and the relative increase of protein binding to TNF- α demonstrates the ability of the $^{10}\text{Fn3}$ scaffold and of the DNA/mRNA-protein fusion-based selection system to select target-specific binders.

[0137] To examine the specificity of binding further, clone T09.12 was immobilized in a microarray format (as generally described below) and was tested for binding to soluble TNF- α . Specific binding of TNF- α to this clone was detected using fluorescence detection (FIG. 24A) and mass spectroscopy (FIG. 24B). For the mass spectroscopy results, binding assays were carried out in the presence of fetal bovine serum, an exemplary complex biological fluid containing a variety of potential interfering proteins. For fluorescence detection (FIG. 24A), a mixture of RNA- $^{10}\text{Fn3}$ fusion of wild-type $^{10}\text{Fn3}$ and of the T09.12 variant (Table 1) was hybridized onto a DNA microarray on which oligonucleotides complementary to the RNA portion of the fusion molecules had been immobilized at 600 micron pitch, with 24 replicate features. After removal of unhybridized fusion by washing, the surface was exposed to biotin-TNF- α (2.6 $\mu\text{g}/\text{mL}$ in TBS, 0.02% Tween-20, 0.2% BSA), washed, and air-dried. The captured biotin-TNF- α was detected by Cy3-labeled anti-biotin monoclonal antibody (Sigma) using a ScanArray 5000 system (GSI Lumonics). For mass spectroscopy detection, RNA- $^{10}\text{Fn3}$ fusion of the T09.12 variant (FIG. 24B) and wild-type $^{10}\text{Fn3}$ (FIG. 24C) was treated with RNase A to generate a fusion between the protein and the DNA linker. The resulting DNA-linked protein was hybridized to a glass coverslip arrayed with an immobilized oligonucleotide complementary to the DNA linker (FIGS. 24B and 24C; no fusion was applied in FIG. 24D). After washing, the above surfaces were exposed to TNF- α (1.5 mg/mL in 90% v/v PBS/10% fetal bovine serum). The dried chip was spotted with MALDI matrix and analyzed with a Voyager DE MALDI-TOF mass spectrometer (PerSeptive Biosystems). A signal at 17.4 kD, which corresponded to the expected molecular mass of monomeric TNF- α , was detected on the 200 μm features that contained T09.12 protein (FIG. 24A), but not on the features that contained wild-type $^{10}\text{Fn3}$ (FIG. 24B) nor on the features that did not contain DNA-protein fusion (FIG. 24C).

K_d of the Selected TNF- α Binding Clones

[0138] Dissociation constants were determined for all the clones that were represented more than once in the two pools generated after nine and after ten rounds of selection, as well as for the only clone that originated from the BC_{wr}-DE_{wr}FG_r library (clone T09.08).

[0139] To determine these binding constants, biotinylated TNF- α was prepared using the NHS-LC-LC-Biotin reagent supplied by Pierce (Rockford, Ill.). MALDI-TOF mass spectrometry was used to estimate that more than 80% of the monomeric TNF- α , and hence more than 99% of the trimer, was biotinylated.

[0140] For the R9 and R10 pools (and the R14 and M12 pools discussed below), as well as for the characterized clones derived from these two pools, eleven samples of 0.25 nM, in vitro-translated, ^{35}S -methionine-labeled free protein were incubated with the biotinylated TNF- α at a concentration between 17 pM and 23 nM, in 200 μL 10 μM HEPES, pH 7.4, 150 mM NaCl, 1% BSA, 0.02% Triton, for one hour at room temperature. Subsequently, each sample was loaded on a pre-soaked, SAM^{2R} Biotin Capture Membrane (Promega, Madison, Wis.) using a 96 well, Easy-Titer™ ELISA system (Pierce). Under vacuum, each spot was washed with 200 μL of HBS pH 7.4, 1% BSA, 0.05% Triton; next the entire membrane was rinsed in the buffer and air-dried. The membrane was exposed with a Storage Phosphor Screen (Molecular Dynamics, Sunnyvale, Calif.) overnight, and the intensities of the resulting individual spots were quantified using a STORM 860 phosphorimager with the ImageQuaNT densitometry program (Molecular Dynamics). The K_d of the binding was determined by fitting the equilibrium equation to the resulting binding curve (KaleidaGraph, Synergy Software)? the error of the experiment was estimated from 24 independent experiments.

[0141] In these studies, the K_d values were found to be in the narrow range of 1-24 nM (Table 1). The T09.12 and T10.06 clones, which contained the most commonly isolated sequences in two loops each, have the low K_d of 4 and 2 nM, respectively; on the other hand, a number of clones with less frequently seen loops, such as clones T09.07 and T10.15, showed similarly tight binding.

[0142] A sample comparison of TNF- α binding between free protein and the cDNA/mRNA-protein complex derived from the same sequence showed that the two dissociation constants were within experimental error of each other, a property of the system that makes it possible to use the cDNA/mRNA-protein complex to select for target-binding properties of the protein itself.

High-Stringency Selection of TNF- α Binding Clones

[0143] Despite the duplicate clones isolated, the TNF- α -binding pools after nine and after ten rounds of selection contained numerous different clones, i.e., 38 different sequences in 61 clones sampled. Therefore, further selection, with more stringent binding requirements, was undertaken to recover a subset of these clones with superior TNF- α binding properties. Consequently, four further rounds of selections (R11-R14) were conducted in solution, where the concentration of the target was controlled more easily. The concentration of TNF- α was limited to 0.5 nM and the concentration of DNA/mRNA- $^{10}\text{Fn3}$ pool to 0.1 nM; in addition, the length and the temperature of the washes of the $^{10}\text{Fn3}$ /TNF- α complex bound to streptavidin-coated paramagnetic beads were increased.

[0144] Specifically, these selections were carried out as follows. For rounds 11-13, 0.1 nM DNA/mRNA- $^{10}\text{Fn3}$ fusion library, which had been made as described above, was pre-cleared by tumbling for 1 hour at 4° C. with 100 μL of Dynabeads® M-280 (streptavidin-coated paramagnetic beads; Dynal, Lake Success, N.Y.) that had been pre-blocked in Binding Buffer. The resulting pre-cleared fusion mixture was combined with 0.5 nM biotinylated TNF- α in 300 μL of the above Binding Buffer, and the complex incubated at 4° C. for 1 hour. Next, 100, of resuspended Dynabeads® M-280 Streptavidin at 1.3 g/cm^3 , which had been blocked by overnight incubation in Binding Buffer, were added to the mixture

and incubated at 4° C., with tumbling, for 45 additional minutes. The paramagnetic beads were separated from the supernatant on a Dynal MPC-S rack, the supernatant was removed, and the beads were washed with the Binding Buffer for 1, 15, and 30 minutes in the case of R11 and R12, or for 1 minute, followed by nine ten-minute washes in the case of R13-R14. DNA was eluted from the washed DNA/mRNA-¹⁰F_n3:TNF- α -biotin:streptavidin-bead complexes with two washes of 100 μ L 0.1 M KOH, and treated as described above for the column-based selection to produce the next generation DNA/mRNA-¹⁰F_n3 fusion library. Round 14 differed from R11-R13 in that the selection was performed at 30° C. and in the presence of an additional 150 mM NaCl. Except for the elevated temperature, the sequence of washes was the same for R14 as for R13.

[0145] Twenty-two clones derived from the DNA eluted after four further rounds of selection (R14) were picked at random and found to represent 15 different loop sequences (Table 2; full sequences in FIG. 25). The clone T10.06, isolated previously from R10 as described above, was picked eight separate times, whereas the remaining sequences, including T09.31, which had been isolated from the R9 pool, were found in one isolate each. Similar to the isolates from rounds nine and ten, the R14 clones examined showed a preference (18 of 22 clones) for the PWA/G sequence in the DE loop, and four new, non-wild-type DE sequences were revealed.

[0146] Whereas the apparent average K_d values of the R14 free protein pool, 3 nM, is similar to those measured for the pools after nine and ten rounds (4 and 6 nM, respectively), several K_d values of the clones isolated from the R14 pool were an order of magnitude lower than the lowest values observed in the R9 and R10 pools (Table 2). The clones that bound TNF- α most tightly, T14.07 and T14.25, had a K_d of 90 pmol. Thus, the conditions used in the last four rounds of selections were stringent enough to favor ¹⁰F_n3 molecules with subnanomolar K_d , but not so stringent as to eliminate such molecules.

Mutagenic Affinity Maturation

[0147] As discussed above, the selections described herein may also be combined with mutagenesis after all or a subset of the selection steps to further increase library diversity. In one parallel selection strategy, error-prone PCR was incorporated into the amplification of DNA between rounds (Cadwell and Joyce, PCR Methods Appl 2:28 (1992)). This technique was carried out beginning with the diverse DNA pool eluted after R8 above. This pool was amplified using error-prone PCR, with the pool divided into seven equal parts and mutagenized at the target frequency of 0.8%, 1.6%, 2.4%, 3.2%, 4.0%, 4.8%, and 5.6%. The seven PCR reactions were combined, and cDNA/RNA-protein fusion was made from the mixture and subjected to a round of selection in solution. Before the second mutagenic round, M10, error-prone PCR was performed in three separate reactions, at 0.8%, 1.6%, and 2.4%. The two remaining rounds, M11 and M12, were performed using standard Taq PCR. Except for mutagenesis, the selection conditions for M9-M12 were the same as for R11-R14. The twenty M12 clones tested showed tighter binding to TNF- α than the clones selected using the two earlier selection protocols (Table 3; full sequences in FIG. 25); the tightest binding of TNF- α was seen in M12.04, and had the observed K_d of 20 pM. These results demonstrated that low-level, random mutagenesis late in a selection can improve both the

binding affinity of selected antibody mimics (20 pM vs. 90 pM) and the speed with which they can be selected (12 rounds vs. 14 rounds). In addition, the frequency of tight binders in this mutagenesis approach was observed to be about 5%, whereas the frequency is approximately 3% in other selections.

Superiority of F_n Binders

[0148] The selection of ¹⁰F_n3 variants capable of binding to TNF- α , performed using covalent mRNA-protein fusion as the unit of selection, was won by molecules with dissociation constants as low as 20 pM. These K_d values compared favorably against the standards of selection of others that used other antibody mimic scaffolds and selection methods. Consequently, the ¹⁰F_n3-based scaffold and covalent mRNA-protein fusion-based in vitro selection method may be utilized for the development of antibody mimics against a broad range of antigens. In addition, the subnanomolar, TNF- α -binding ¹⁰F_n3 variants described herein represent potential therapeutic, research, and diagnostic agents. Moreover, since this in vitro selection method can be automated, such a combination of scaffold and selection methods have applications on the genomic scale.

[0149] One of the factors that contributed to the success of the present selection was the randomization of all three CDR-like loops of ¹⁰F_n3; similar libraries which contained only one or two randomized loops were less likely to include tight binders than the library with three randomized, CDR-like loops.

[0150] In the selection reported above, the randomized loops remained the length of the corresponding, wild-type ¹⁰F_n3 loops. To generate further library diversity, the length of the loops as well as their sequences may be varied, to incorporate favorable mutations in the ¹⁰F_n3 beta-sheet into the wild-type scaffold used for library construction, and to create libraries with randomized beta-sheet scaffolds which will allow selection of structures even more successful at mimicking antibodies.

[0151] Selections similar to those described herein may be carried out with any other binding species target (for example, IL-1 or IL-13).

Animal Studies

[0152] Wild-type ¹⁰F_n3 contains an integrin-binding tripeptide motif, Arginine 78-Glycine 79-Aspartate 80 (the "RGD motif") at the tip of the FG loop. In order to avoid integrin binding and a potential inflammatory response based on this tripeptide in vivo, a mutant form of ¹⁰F_n3 was generated that contained an inert sequence, Serine 78-Glycine 79-Glutamate 80 (the "SGE mutant"), a sequence which is found in the closely related, wild-type ¹⁰F_n3 domain. This SGE mutant was expressed as an N-terminally His₆-tagged, free protein in *E. coli*, and purified to homogeneity on a metal chelate column followed by a size exclusion column.

[0153] In particular, the DNA sequence encoding His₆-¹⁰F_n3(SGE) was cloned into the pET9a expression vector and transformed into BL21 DE3 pLysS cells. The culture was then grown in LB broth containing 50 μ g/mL kanamycin at 37° C., with shaking, to A₅₆₀=1.0, and was then induced with 0.4 mM IPTG. The induced culture was further incubated, under the same conditions, overnight (14-18 hours); the bacteria were recovered by standard, low speed centrifugation. The cell pellet was resuspended in 1/50 of the original culture

volume of lysis buffer (50 mM Tris 8.0, 0.5 M NaCl, 5% glycerol, 0.05% Triton X-100, and 1 mM PMSF), and the cells were lysed by passing the resulting paste through a Microfluidics Corporation Microfluidizer M110-EH, three times. The lysate was clarified by centrifugation, and the supernatant was filtered through a 0.45 μ m filter followed by filtration through a 0.2 μ m filter. 100 mL of the clarified lysate was loaded onto a 5 mL Talon cobalt column (Clontech, Palo Alto, Calif.), washed by 70 mL of lysis buffer, and eluted with a linear gradient of 0-30 mM imidazole in lysis buffer. The flow rate through the column through all the steps was 1 mL/min. The eluted protein was concentrated 10-fold by dialysis (MW cutoff=3,500) against 15,000-20,000 PEG. The resulting sample was dialysed into buffer 1 (lysis buffer without the glycerol), then loaded, 5 mL at a time, onto a 16x60 mm Sephacryl 100 size exclusion column equilibrated in buffer 1. The column was run at 0.8 mL/min, in buffer 1; all fractions that contained a protein of the expected MW were pooled, concentrated 10 $\times$ as described above, then dialyzed into PBS. Endotoxin screens and animal studies were performed on the resulting sample (Toxikon; MA).

[0154] The endotoxin levels in the samples examined to date have been below the detection level of the assay. In a preliminary animal toxicology study, this protein was injected into two mice at the estimated 100 $\times$ therapeutic dose of 2.6 mg/mouse. The animals survived the two weeks of the study with no apparent ill effects. These safety results support the use of ¹⁰F_n3 incorporated into an IV drug.

Alternative Constructs for In Vivo Use

[0155] To extend the half life of the 8 kD ¹⁰F_n3 domain, a larger molecule has also been constructed that mimics natural antibodies. This ¹⁰F_n3-F_c molecule contains the —CH₂—CH₂—CH₃ (FIG. 11) or —CH₂—CH₃ domains of the IgG constant region of the host; in these constructs, the ¹⁰F_n3 domain is grafted onto the N-terminus in place of the IgG V_H domain (FIGS. 11 and 12). Such antibody-like constructs are to improve the pharmacokinetics of the protein as well as its ability to harness the natural immune response.

[0156] In order to construct the murine form of the ¹⁰F_n3-CH₁—CH₂—CH₃ clone, the —CH₁—CH₂—CH₃ region was first amplified from a mouse liver spleen cDNA library (Clontech), then ligated into the pET25b vector. The primers used in the cloning were 5' Fc Nest and 3' 5 Fc Nest, and the primers used to graft the appropriate restriction sites onto the ends of the recovered insert were 5'Fc HIII and 3'Fc Nhe:

```

5' Fc Nest                                     (SEQ ID NO: 15)
5' GCG GCA GGG TTT GCT TAC TGG GGC CAA GGC 3';

3' Fc Nest                                     (SEQ ID NO: 16)
5' GGG AGG GGT GGA GGT AGG TCA CAG TCC 3';

3' Fc Nhe                                     (SEQ ID NO: 17)
5' TTT GCT AGC TTT ACC AGG AGA GTG GGA GGC 3';
and
5' Fc HIII                                     (SEQ ID NO: 18)
5' AAA AAG CTT GCC AAA ACG ACA CCC CCA TCT GTC 3'.
```

[0157] Further PCR was used to remove the CH₁ region from this clone and to create the Fc part of the shorter, ¹⁰F_n3-

CH₂—CH₃ clone. The sequence encoding ¹⁰F_n3 was spliced onto the 5' end of each, clone; either the wild type ¹⁰F_n3 cloned from the same mouse spleen cDNA library or a modified ¹⁰F_n3 obtained by mutagenesis or randomization of the molecules can be used. The oligonucleotides used in the cloning of murine wild-type ¹⁰F_n3 were:

```

Mo 5PCR-NdeI:                               (SEQ ID NO: 19)
5' CATATGGTTTCTGATATTCGAGAGATCTGGAG 3';

Mo5PCR-His-NdeI (for an alternative N-terminus
with the His6 purification tag):             (SEQ ID NO: 20)
5' CAT ATG CAT CAC CAT CAC CAT CAC GTT TCT GAT
ATT CCG AGA G 3';

Mo3PCR-EcoRI:                               (SEQ ID NO: 21)
5' GAATTCCTATGTTTATAATTGATGAAAC 3'.
```

[0158] The human equivalents of the clones are constructed using the same strategy with human oligonucleotide sequences.

Antibody Mimics in Protein Chip Applications

[0159] Any of the antibody mimics described herein may be immobilized on a solid support, such as a microchip. The suitability of the present scaffolds, for example, the ¹⁰F_n3 scaffold, for protein chip applications is the consequence of (1) their ability to support many binding functions which can be selected rapidly on the bench or in an automated setup, and (2) their superior biophysical properties.

[0160] The versatile binding properties of ¹⁰F_n3 are a function of the loops displayed by the F_n3 immunoglobulin-like, beta sandwich fold. As discussed above, these loops are similar to the complementarity determining regions of antibody variable domains and can cooperate in a way similar to those antibody loops in order to bind antigens. In our system, ¹⁰F_n3 loops BC (for example, residues 21-30), DE (for example, residues 51-56), and FG (for example, residues 76-87) are randomized either in sequence, in length, or in both sequence and length in order to generate diverse libraries of mRNA-¹⁰F_n3 fusions. The binders in such libraries are then enriched based on their affinity for an immobilized or tagged target, until a small population of high affinity binders are generated. Also, error-prone PCR and recombination can be employed to facilitate affinity maturation of selected binders. Due to the rapid and efficient selection and affinity maturation protocols, binders to a large number of targets can be selected in a short time.

[0161] As a scaffold for binders to be immobilized on protein chips, the ¹⁰F_n3 domain has the advantage over antibody fragments and single-chain antibodies-of being smaller and easier to handle. For example, unlike single-chain scaffolds or isolated variable domains of antibodies, which vary widely in their stability and solubility, and which require an oxidizing environment to preserve their structurally essential disulfide bonds, ¹⁰F_n3 is extremely stable, with a melting temperature of 110° C., and solubility at a concentration >16 mg/mL. The ¹⁰F_n3 scaffold also contains no disulfides or free cysteines; consequently, it is insensitive to the redox potential of

its environment. A further advantage of $^{10}\text{Fn3}$ is that its antigen-binding loops and N-terminus are on the edge of the beta-sandwich opposite to the C-terminus; thus the attachment of a $^{10}\text{Fn3}$ scaffold to a chip by its C-terminus aligns the antigen-binding loops, allowing for their greatest accessibility to the solution being assayed. Since $^{10}\text{Fn3}$ is a single domain of only 94 amino acid residues, it is also possible to immobilize it onto a chip surface at a higher density than is used for single-chain antibodies, with their approximately 250 residues. In addition, the hydrophilicity of the $^{10}\text{Fn3}$ scaffold, which is reflected in the high solubility of this domain, minimizes unwanted binding of $^{10}\text{Fn3}$ to a chip surface.

[0162] The stability of the $^{10}\text{Fn3}$ scaffold as well as its suitability for library formation and selection of binders are likely to be shared by the large, Fn3-like class of protein domains with an immunoglobulin-like fold, such as the domains of tenascin, N-cadherin, E-cadherin, ICAM, titin, GCSF-R, cytokine receptor, glycosidase inhibitor, and antibiotic chromoprotein. The key features shared by all such domains are a stable framework provided by two beta-sheets, which are packed against each other and which are connected by at least three solvent-accessible loops per edge of the sheet; such loops can be randomized to generate a library of potential binders without disrupting the structure of the framework (as described above). In addition, as with $^{10}\text{Fn3}$, any of these loops (or similar loops from other proteins) may be immobilized alone or in combination with other loops onto a solid support surface.

Immobilization of Fn3-Based Antibody Mimics

[0163] To immobilize antibody mimics, such as Fn3-based antibody mimics, to a chip surface, a number of exemplary techniques may be utilized. For example, such antibody mimics may be immobilized as RNA-protein fusions by Watson-Crick hybridization of the RNA moiety of the fusion to a base complementary DNA immobilized on the chip surface (as described, for example, in Addressable Protein Arrays, U.S. Ser. No. 60/080,686; U.S. Ser. No. 09/282,734; and WO 99/51773; and Methods for Encoding and Sorting In Vitro Translated Proteins, U.S. Ser. No. 60/151,261 and U.S. Ser. No. 09/648,040). Alternatively, antibody mimics can be immobilized as free proteins directly on a chip surface. Manual as well as robotic devices may be used for deposition of the antibody mimics on the chip surface. Spotting robots can be used for deposition of antibody mimics with high density in an array format (for example, by the method of Lueking et al., *Anal Biochem.* 1999 May 15; 270(1):103-11). Different methods may also be utilized for anchoring the antibody mimic on the chip surface. A number of standard immobilization procedures may be used including those described in *Methods in Enzymology* (K. Mosbach and B. Danielsson, eds.), vols. 135 and 136, Academic Press, Orlando, Fla., 1987; Nilsson et al., *Protein Expr. Purif.* 1997 October; 11(1):1-16; and references therein. Oriented immobilization of antibody mimics can help to increase the binding capacity of chip-bound antibody mimics. Exemplary approaches for achieving oriented coupling are described in Lu et al., *The Analyst* (1996), vol. 121, p. 29R-32R; and Turkova, *J Chromatogr B Biomed Sci App.* 1999 Feb. 5; 722(1-2): 11-31. In addition, any of the methods described

herein for anchoring antibody mimics to chip surfaces can also be applied to the immobilization of antibody mimics on beads, or other supports.

Target Protein Capture and Detection

[0164] Selected populations of scaffold-binders may be used for detection and/or quantitation of analyte targets, for example, in samples such as biological samples. To carry out this type of diagnostic assay, selected scaffold-binders to targets of interest are immobilized on an appropriate support to form multi-featured protein chips. Next, a sample is applied to the chip, and the components of the sample that associate with the binders are identified based on the target-specificity of the immobilized binders. Using this technique, one or more components may be simultaneously identified or quantitated in a sample (for example, as a means to carry out sample profiling).

[0165] Methods for target detection allow measuring the levels of bound protein targets and include, without limitation, radiography, fluorescence scanning, mass spectroscopy (MS), and surface plasmon resonance (SPR). Autoradiography using a phosphorimager system (Molecular Dynamics, Sunnyvale, Calif.) can be used for detection and quantification of target protein which has been radioactively labeled, e.g., using ^{35}S methionine. Fluorescence scanning using a laser scanner (see below) may be used for detection and quantification of fluorescently labeled targets. Alternatively, fluorescence scanning may be used for the detection of fluorescently labeled ligands which themselves bind to the target protein (e.g., fluorescently labeled target-specific antibodies or fluorescently labeled streptavidin binding to target-biotin, as described below).

[0166] Mass spectroscopy can be used to detect and identify bound targets based on their molecular mass. Desorption of bound target protein can be achieved with laser assistance directly from the chip surface as described below. Mass detection also allows determinations, based on molecular mass, of target modifications including post-translational modifications like phosphorylation or glycosylation. Surface plasmon resonance can be used for quantification of bound protein targets where the scaffold-binder(s) are immobilized on a suitable gold-surface (for example, as obtained from Biacore, Sweden).

[0167] Described below are exemplary schemes for selecting binders (in this case, Fn-binders specific for the protein, TNF- α) and the use of those selected populations for detection on chips. This example is provided for the purpose of illustrating the invention, and should not be construed as limiting.

Selection of TNF- α Binders Based on $^{10}\text{Fn3}$ Scaffold

[0168] In one exemplary use for scaffold selection on chips, an $^{10}\text{Fn3}$ -based selection was performed against TNF- α , using a library of human $^{10}\text{Fn3}$ variants with randomized loops BC, DE, and FG. The library was constructed from three DNA fragments, each of which contained nucleotide sequences that encoded approximately one third of human $^{10}\text{Fn3}$, including one of the randomized loops. The DNA sequences that encoded the loop residues listed above were rebuilt by oligonucleotide synthesis, so that the codons for the residues of interest were replaced by (NNS)_{*m*}, where N represents any of the four deoxyribonucleotides (A, C, G, or T),

and S represents either C or G. The C-terminus of each fragment contained the sequence for the FLAG purification tag.

[0169] Once extended by Klenow, each DNA fragment was transcribed, ligated to a puromycin-containing DNA linker, and translated *in vitro*, as described by Szostak et al. (Roberts and Szostak, *Proc. Natl. Acad. Sci. USA* 94:12297, 1997; Szostak et al., U.S. Ser. No. 09/007,005 and U.S. Ser. No. 09/247,190; Szostak et al., WO98/31700), to generate an mRNA-peptide fusion, which was then reverse-transcribed into a DNA-mRNA-peptide fusion. The binding of the FLAG-tagged peptide to M2 agarose separated full-length fusion molecules from those containing frameshifts or superfluous stop codons; the DNA associated with the purified full-length fusion was amplified by PCR, then the three DNA fragments were cut by *Ear I* restriction endonuclease and ligated to form the full length template. The template was transcribed, ligated to puromycin-containing DNA linkers, and translated to generate a ¹⁰F_n3-RNA/cDNA library, which was then reverse-transcribed to yield the DNA-mRNA-peptide fusion library which was subsequently used in the selection.

[0170] Selection for TNF- α binders took place in 50 mM HEPES, pH 7.4, 0.02% Triton-X, 0.1 mg/mL salmon sperm DNA. The PROfusion™ library was incubated with Sepharose-immobilized TNF- α ; after washing, the DNA associated with the tightest binders was eluted with 0.1 M KOH, amplified by PCR, and transcribed, ligated, translated, and reverse-transcribed into the starting material for the next round of selection.

[0171] Ten rounds of such selection were performed (as shown in FIG. 13); they resulted in a PROfusion™ pool that bound to TNF- α -Sepharose with the apparent average K_d of 120 nM. Specific clonal components of the pool that were characterized showed TNF- α binding in the range of 50-500 nM.

Immobilization, Target Protein Capture, and MALDI-TOF Detection

[0172] As a first step toward immobilizing F_n3 fusions to a chip surface, an oligonucleotide capture probe was prepared with an automated DNA synthesizer (PE BioSystems Expedite 8909) using the solid-support phosphoramidite approach. All reagents were obtained from Glen Research. Synthesis was initiated with a solid support containing a disulfide bond to eventually provide a 3'-terminal thiol functionality. The first four monomers to be added were hexaethylene oxide units, followed by 20 T monomers. The 5'-terminal DMT group was not removed. The capture probe was cleaved from the solid, support and deprotected with ammonium hydroxide, concentrated to dryness in a vacuum centrifuge, and purified by reverse-phase HPLC using an acetonitrile gradient in triethylammonium acetate buffer. Appropriate fractions from the HPLC were collected, evaporated to dryness in a vacuum centrifuge, and the 5'-terminal DMT group was removed by treatment with 80% AcOH for 30 minutes. The acid was removed by evaporation, and the oligonucleotide was then treated with 100 mM DTT for 30 minutes to cleave the disulfide bond. DTT was removed by repeated extraction with EtOAc. The oligonucleotide was ethanol precipitated from the remaining aqueous layer and checked for purity by reverse-phase HPLC.

[0173] The 3'-thiol capture probe was adjusted to 250 μ M in degassed 1 $\times$ PBS buffer and applied as a single droplet (75 μ L) to a 9 $\times$ 9 mm gold-coated chip (Biacore) in an argon-flushed

chamber containing a small amount of water. After 18-hours at room temperature, the capture probe solution was removed, and the functionalized chip was washed with 50 mL 1 $\times$ PBS buffer (2 $\times$ for 15 minutes each) with gentle agitation, and then rinsed with 50 mL water (2 $\times$ for 15 minutes each) in the same fashion. Remaining liquid was carefully removed and the functionalized chips were either used immediately or stored at 4° C. under argon.

[0174] About 1 pmol of ¹⁰F_n3 fusion pool from the Round 10 TNF- α selection (above) was treated with RNase A for several hours, adjusted to 5 $\times$ SSC in 70 μ L, and applied to a functionalized gold chip from above as a single droplet. A 50 μ L volume gasket device was used to seal the fusion mixture with the functionalized chip, and the apparatus was continuously rotated at 4° C. After 18 hours the apparatus was disassembled, and the gold chip was washed with 50 mL 5 $\times$ SSC for 10 minutes with gentle agitation. Excess liquid was carefully removed from the chip surface, and the chip was passivated with a blocking solution (1 $\times$ TBS+0.02% Tween-20+0.25% BSA) for 10 minutes at 4° C. Excess liquid was carefully removed, and a solution containing 500 μ g/mL TNF- α in the same composition blocking solution was applied to the chip as a single droplet and incubated at 4° C. for two hours with occasional mixing of the droplet via Pipetman. After removal of the binding solution, the chip was washed for 5 minutes at 4° C. with gentle agitation (50 mL 1 $\times$ TBS+0.02% Tween-20) and then dried at room temperature. A second chip was prepared exactly as described above, except fusion was not added to the hybridization mix.

[0175] Next, MALDI-TOF matrix (15 mg/mL 3,5-dimethoxy-4-hydroxycinnamic acid in 1:1 ethanol/10% formic acid in water) was uniformly applied to the gold chips with a high-precision 3-axis robot (MicroGrid, BioRobotics). A 16-pin tool was used to transfer the matrix from a 384-well microtiter plate to the chips, producing 200 micron diameter features with a 600 micron pitch. The MALDI-TOF mass spectrometer (Voyager DE, PerSeptive-Biosystems) instrument settings were as follows: Accelerating Voltage=25k, Grid Voltage=92%, Guide Wire Voltage=0.05%, Delay=200 on, Laser Power=2400, Low Mass Gate=1500, Negative Ions=off. The gold chips were individually placed on a MALDI sample stage modified to keep the level of the chip the same as the level of the stage, thus allowing proper flight distance. The instrument's video monitor and motion control system were used to direct the laser beam to individual matrix features.

[0176] FIGS. 14 and 15 show the mass spectra from the ¹⁰F_n3 fusion chip and the non-fusion chip, respectively. In each case, a small number of 200 micron features were analyzed to collect the spectra, but FIG. 15 required significantly more acquisitions. The signal at 17.4 kDa corresponds to TNF- α monomer.

Immobilization, Target Protein Capture, and Fluorescence Detection

[0177] Pre-cleaned 1 $\times$ 3 inch glass microscope slides (Goldseal, #3010) were treated with Nanostrip (Cyantek) for 15 minutes, 10% aqueous NaOH at 70° C. for 3 minutes, and 1% aqueous HCl for 1 minute, thoroughly rinsing with deionized water after each reagent. The slides were then dried in a vacuum desiccator over anhydrous calcium sulfate for several hours. A 1% solution of aminopropyltrimethoxysilane in 95% acetone/5% water was prepared and allowed to hydrolyze for 20 minutes. The glass slides were immersed in the hydrolyzed

silane solution for 5 minutes with gentle agitation. Excess silane was removed by subjecting the slides to ten 5-minute washes, using fresh portions of 95% acetone/5% water for each wash, with gentle agitation. The slides were then cured by heating at 110° C. for 20 minutes. The silane treated slides were immersed in a freshly prepared 0.2% solution of phenylene 1,4-diisothiocyanate in 90% DMF/10% pyridine for two hours, with gentle agitation. The slides were washed sequentially with 90% DMF/10% pyridine, methanol, and acetone. After air drying, the functionalized slides were stored at 0° C. in a vacuum desiccator over anhydrous calcium sulfate. Similar results were obtained with commercial amine-reactive slides (3-D Link, Surmodics).

[0178] Oligonucleotide capture probes were prepared with an automated DNA synthesizer (PE BioSystems Expedite 8909) using conventional phosphoramidite chemistry. All reagents were from Glen Research. Synthesis was initiated with a solid support bearing an orthogonally protected amino functionality, whereby the 3'-terminal amine is not unmasked until final deprotection step. The first four monomers to be added were hexaethylene oxide units, followed by the standard A, G, C and T monomers. All capture oligo sequences were cleaved from the solid support and deprotected with ammonium hydroxide, concentrated to dryness, precipitated in ethanol, and purified by reverse-phase HPLC using an acetonitrile gradient in triethylammonium acetate buffer. Appropriate fractions from the HPLC were collected, evaporated to dryness in a vacuum centrifuge, and then coevaporated with a portion of water.

[0179] The purified, amine-labeled capture origos were adjusted to a concentration of 250 μ M in 50 mM sodium carbonate buffer (pH 9.0) containing 10% glycerol. The probes were spotted onto the amine-reactive glass surface at defined positions in a 5 $\times$ 5 $\times$ 6 array pattern with a 3-axis robot (MicroGrid, BioRobotics). A 16-pin tool was used to transfer the liquid from 384-well microtiter plates, producing 200 micron features with a 600 micron pitch. Each sub-grid of 24 features represents a single capture probe (i.e., 24 duplicate spots). The arrays were incubated at room temperature in a moisture-saturated environment for 12-18 hours. The attachment reaction was terminated by immersing the chips in 2% aqueous ammonium hydroxide for five minutes with gentle agitation, followed by rinsing with distilled water (3 $\times$ for 5 minutes each). The array was finally soaked in 10 $\times$ PBS solution for 30 minutes at room temperature, and then rinsed again for 5 minutes in distilled water.

[0180] Specific and thermodynamically isoenergetic sequences along the ¹⁰F_n3 mRNA were identified to serve as capture points to self-assemble and anchor the ¹⁰F_n3 protein. The software program HybSimulator v4.0 (Advanced Gene Computing Technology, Inc.) facilitated the identification and analysis of potential capture probes. Six unique capture probes were chosen and printed onto the chip, three of which are complementary to common regions of the ¹⁰F_n3 fusion pool's mRNA (CP3', CP5', and CPflag). The remaining three sequences (CPneg1, CPneg2, and CPneg3) are not complementary and function in part as negative controls. Each of the capture probes possesses a 3'-amino terminus and four hexaethylene oxide spacer units, as described above. The following is a list of the capture probe sequences that were employed (5'→3'):

CP3' :	TGTAATAGTAATTGTCCC	(SEQ ID NO: 22)
CP5' :	TTTTTTTTTTTTTTTTTTT	(SEQ ID NO: 23)
CPneg1 :	CCTGTAGGTGTCCAT	(SEQ ID NO: 24)
CPflag :	CATCGTCCTTGTAGTC	(SEQ ID NO: 25)
CPneg2 :	CGTCGTAGGGTA	(SEQ ID NO: 26)
CPneg3 :	CAGGTCTTCTTCAGAGA	(SEQ ID NO: 27)

About 1 pmol of ¹⁰F_n3 fusion pool from the Round 10 TNF- α selection was adjusted to 5 $\times$ SSC containing 0.02% Tween-20 and 2 mM vanadyl ribonucleotide complex in a total volume of 350 μ L. The entire volume was applied to the microarray under a 400 μ L gasket device and the assembly was continuously rotated for 18 hours at room temperature. After hybridization the slide was washed sequentially with stirred 500 mL portions of 5 $\times$ SSC, 2.5 $\times$ SSC, and 1 $\times$ SSC for 5 minutes each. Traces of liquid were removed by centrifugation and the slide was allowed to air-dry.

[0181] Recombinant human TNF- α (500 μ g, lyophilized, from PreproTech) was taken up in 230 μ L 1 $\times$ PBS and dialyzed against 700 mL stirred 1 $\times$ PBS at 4° C. for 18 hours in a Microdialyzer unit (3,500 MWCO, Pierce). The dialyzed TNF- α was treated with EZ-Link NHS-LC-LC biotinylation reagent (20 μ g, Pierce) for 2 hours at 0° C., and again dialyzed against 700 mL stirred 1 $\times$ PBS at 4° C. for 18 hours in a Microdialyzer unit (3,500 MWCO, Pierce). The resulting conjugate was analyzed by MAIDI-TOF mass spectrometry and was found to be almost completely functionalized with a single biotin moiety.

[0182] Each of the following processes was conducted at 4° C. with continuous rotation or mixing. The protein microarray surface was passivated by treatment with 1 $\times$ TBS-containing 0.02% Tween-20 and 0.2% BSA (200 μ L) for 60 minutes. Biotinylated TNF- α (100 nM concentration made up in the passivation buffer) was contacted with the microarray for 120 minutes. The microarray was washed with 1 $\times$ TBS containing 0.02% Tween-20 (3 $\times$ 50 mL, 5 minutes each wash). Fluorescently labeled streptavidin (2.5 μ g/mL Alexa 546-streptavidin conjugate from Molecular Probes, made up in the passivation buffer) was contacted with the microarray for 60 minutes. The microarray was washed with 1 $\times$ TBS containing 0.02% Tween-20 (2 $\times$ 50 mL, 5 minutes each wash) followed by a 3 minute rinse with 1 $\times$ TBS. Traces of liquid were removed by centrifugation, and the slide was allowed to air-dry at room temperature.

[0183] Fluorescence laser scanning was performed with a GSI Lumonics ScanArray 5000 system using 101 pixel resolution and preset excitation and emission wavelengths for Alexa 546 dye. Phosphorimage analysis was performed with a Molecular Dynamics Storm system. Exposure time was 48 hours with direct contact between the microarray and the phosphor storage screen. Phosphorimage scanning was performed at the 50 micron resolution setting, and data was extracted with ImageQuant v.4.3 software.

[0184] FIGS. 16 and 17 are the phosphorimage and fluorescence scan, respectively, of the same array. The phosphorimage shows where the ¹⁰F_n3 fusion hybridized based on the ³⁵S methionine signal. The fluorescence scan shows where the labeled TNF- α bound.

OTHER EMBODIMENTS

[0185] Other embodiments are within the claims.

[0186] All publications, patents, and patent applications mentioned herein are hereby incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 202

<210> SEQ ID NO 1

<211> LENGTH: 122

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
ggaattccta atacgactca ctataggac aattactatt tacaattaca atgcatcacc    60
atcaccatca cgtttctgat gttccgaggg acctggaagt tgtgtgtgcg acccccacca    120
gc                                                                    122
```

<210> SEQ ID NO 2

<211> LENGTH: 104

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```
ggaattccta atacgactca ctataggac aattactatt tacaattaca atgggttctg    60
atgttccgag ggacctggaa gttgtgtgctg cgacccccac cagc                  104
```

<210> SEQ ID NO 3

<211> LENGTH: 126

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)...(126)

<223> OTHER INFORMATION: n = A,T,C or G

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)...(126)

<223> OTHER INFORMATION: s = C or G

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: 74, 75, 77, 78, 80, 81, 83, 84, 86, 87, 89, 90, 92, 93

<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 3

```
agcggatgcc ttgtcgctgt cgtccttgta gtcgtctctc cctgtttctc cgtaagtgtat    60
cctgtaatat ctsnnsnnsn nsnnnsnnnn snnccagctg atcagtaggc tggtagggggt    120
cgcagc                                                                    126
```

<210> SEQ ID NO 4

<211> LENGTH: 62

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```
agcggatgcc ttgtcgctgt cgtccttgta gtcgtctctc cctgtttctc cgtaagtgtat    60
cc                                                                    62
```

<210> SEQ ID NO 5

-continued

```

<211> LENGTH: 99
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5
ggaattccta atacgactca ctataggac aattactatt tacaattaca atgcatcacc      60
atcaccatca cctcttcaca ggaggaaata gccctgtcc                             99

<210> SEQ ID NO 6
<211> LENGTH: 132
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(132)
<223> OTHER INFORMATION: n = A,T,C or G
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(132)
<223> OTHER INFORMATION: s = C or G
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 83, 84, 86, 87, 89, 90, 92, 93
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 6
agcggatgcc ttgtcgctgt cgtccttgta gtcgctcttc gtataatcaa ctccagggtt      60
aaggccgctg atggtagctg tsnnnsnnnn snnaggcaca gtgaactcct ggacagggct      120
atttcctcct gt                                                         132

<210> SEQ ID NO 7
<211> LENGTH: 64
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7
agcggatgcc ttgtcgctgt cgtccttgta gtcgctcttc gtataatcaa ctccagggtt      60
aagg                                                         64

<210> SEQ ID NO 8
<211> LENGTH: 101
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8
ggaattccta atacgactca ctataggac aattactatt tacaattaca atgcatcacc      60
atcaccatca cctcttctat accatcactg tgtatgctgt c                             101

<210> SEQ ID NO 9
<211> LENGTH: 114
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(114)
<223> OTHER INFORMATION: n = A,T,C or G
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(114)
<223> OTHER INFORMATION: s = C or G
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 59, 60, 62, 63, 65, 66, 68, 69, 71, 72, 74, 75, 77, 78,

```

-continued

80, 81, 83, 84, 86, 87
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 9

agcggatgcc ttgtcgtcgt cgtccttgta gtctgttcgg taattaatgg aaattggsnn 60

snnsnnsnns nnsnnsnnsn nsnnnsnagt gacagcatatg acagtgatgg tata 114

<210> SEQ ID NO 10
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

agcggatgcc ttgtcgtcgt cgtccttgta gtctgttcgg taattaatgg aaattgg 57

<210> SEQ ID NO 11
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: T7 phage and tobacco mosaic virus

<400> SEQUENCE: 11

gcgtaatacgt actcactata gggacaatta ctatttataa ttaca 45

<210> SEQ ID NO 12
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Flag sequence

<400> SEQUENCE: 12

agcggatgcc ttgtcgtcgt cgtccttgta gtc 33

<210> SEQ ID NO 13
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Splint oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)...(19)
<223> OTHER INFORMATION: n = A,T,C or G
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 10
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 13

tttttttttn agcggatgc 19

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Puromycin linker oligonucleotide

<400> SEQUENCE: 14

aaaaaaaaa aaaaaaaacc 20

<210> SEQ ID NO 15
<211> LENGTH: 30

-continued

<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 15
gcggcagggt ttgcttactg gggccaaggg 30

<210> SEQ ID NO 16
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 16
gggaggggtg gaggtaggtc acagtcc 27

<210> SEQ ID NO 17
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 17
tttgctagct ttaccaggag agtgggaggc 30

<210> SEQ ID NO 18
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 18
aaaaagcttg ccaaaacgac acccccatct gtc 33

<210> SEQ ID NO 19
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 19
catatggttt ctgatattcc gagagatctg gag 33

<210> SEQ ID NO 20
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 20
catatgcac accatcacca tcacgtttct gatattccga gag 43

<210> SEQ ID NO 21
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 21
gaattcctat gttttataat tgatggaaac 30

<210> SEQ ID NO 22
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22
tgtaaatagt aattgtccc 19

-continued

<210> SEQ ID NO 23
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

tttttttttt tttttttttt

20

<210> SEQ ID NO 24
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 24

cctgtaggtg tccat

15

<210> SEQ ID NO 25
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

catcgtcctt gtagtc

16

<210> SEQ ID NO 26
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 26

cgtcgtaggg gta

13

<210> SEQ ID NO 27
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 27

caggtcttct tcagaga

17

<210> SEQ ID NO 28
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

catatggttt ctgatgttcc gagg

24

<210> SEQ ID NO 29
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

gaattcctat gttcggtaat taatggaaat tg

32

-continued

<210> SEQ ID NO 30
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: DNA splint Oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 10
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 30

tttttttttn agcggatgc

19

<210> SEQ ID NO 31
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

<400> SEQUENCE: 31

Asn Arg Ser Gly Leu Gln Ser
1 5

<210> SEQ ID NO 32
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

<400> SEQUENCE: 32

Ala Gln Thr Gly His His Leu His Asp Lys
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

<400> SEQUENCE: 33

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly Asp
65 70 75 80

Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 34
<211> LENGTH: 95
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

Val Ser Glu Ile Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

-continued

```

Ser Leu Leu Phe Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr
      20              25              30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Leu Val Gln Glu Phe
      35              40              45

Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Asn Thr Ile Thr Gly Tyr Ala Val Thr Thr Thr Tyr
      65              70              75              80

Arg Thr Arg Ile Asp Lys Gln Pro Ile Ser Ile Asn Tyr Arg Thr
      85              90              95

```

```

<210> SEQ ID NO 35
<211> LENGTH: 90
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 35

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
  1              5              10              15

Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr
      20              25              30

Arg Ile Thr Tyr Gly Glu Lys Gly Gly Asn Ser Pro Val Gln Glu Phe
      35              40              45

Thr Val Pro Glu Leu Asn Pro Thr Ala Thr Ile Ser Arg Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gln Asn Gly Thr
      65              70              75              80

Pro Arg Arg His Leu Arg Pro Asn Phe His
      85              90

```

```

<210> SEQ ID NO 36
<211> LENGTH: 95
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 36

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
  1              5              10              15

Gly Leu Leu Ile Ser Trp Asn Lys Ser Arg Met Thr Thr Arg Tyr Tyr
      20              25              30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35              40              45

Thr Val Pro Val Thr Asp Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Asn Thr Ile Ile Val His Ala Val Thr Leu Thr Asn
      65              70              75              80

Gln Asn Ser Asp His Thr Tyr Pro Ile Ser Ile Asn Tyr Arg Thr
      85              90              95

```

```

<210> SEQ ID NO 37
<211> LENGTH: 91
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 37

```

```

Val Ser Asp Val Pro Arg Asp Leu Asp Val Val Ala Ala Thr Pro Thr

```

-continued

1	5	10	15
Ser Leu Leu Ile Ser Trp Asp Ser Ser His Arg Tyr Tyr Arg Ile Thr	20	25	30
Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr Ala Pro	35	40	45
Asn Asn Pro Pro Thr Ala Thr Ile Ser Gly Leu Lys Pro Gly Val Asp	50	55	60
Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Asp Gly Ser Arg His Met	65	70	75
Leu Thr Lys Pro Ile Ser Ile Asn Tyr Arg Thr	85	90	

<210> SEQ ID NO 38
 <211> LENGTH: 88
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr	1	5	10	15
Ser Leu Leu Ile Ser Trp His Asn Asn His Ile Asp Met Arg Tyr Tyr	20	25	30	
Arg Ser Ala Asn Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Val Phe	35	40	45	
Thr Val Pro Gln Arg Arg Gln Thr Ala Thr Ile Ser Gly Leu Lys Pro	50	55	60	
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Lys Asn Gln	65	70	75	80
Gly Arg Arg Arg Gln Gly Ile Arg	85			

<210> SEQ ID NO 39
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ser Thr	1	5	10	15
Ser Leu Leu Ile Ser Trp Arg Thr Pro Ala Ser Pro His Gly Tyr Tyr	20	25	30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Glu Glu Phe	35	40	45	
Thr Val Pro Leu Leu Trp Pro Thr Ala Thr Ile Ser Gly Leu Lys Pro	50	55	60	
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Thr His Met	65	70	75	80
Leu Lys Pro Gln Ser Met Pro Ile Ser Ile Asn Tyr Arg Thr	85	90		

<210> SEQ ID NO 40
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 40

-continued

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10          15
Ser Leu Leu Ile Ser Trp Arg Thr Pro Ala Ser Pro His Gly Tyr Tyr
          20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Glu Glu Phe
          35           40           45
Thr Val Pro Leu Leu Trp Pro Thr Ala Thr Ile Ser Gly Leu Lys Pro
          50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Thr His Met
65           70           75           80
Leu Lys Pro Gln Ser Met Pro Ile Ser Ile Asn Tyr Arg Thr
          85           90

```

```

<210> SEQ ID NO 41
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 41

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Ala Ser Thr
 1           5           10          15
Ser Leu Leu Ile Ser Trp Arg Pro Asn Pro Arg Leu Ser Arg Tyr Tyr
          20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35           40           45
Thr Val Pro Gly Leu Phe Ser Thr Ala Thr Ile Ser Gly Leu Asn Pro
          50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Lys Glu Thr
65           70           75           80
Ser Asn Ile Phe Ile Ala Pro Ile Ser Ile Asn Tyr Arg Thr
          85           90

```

```

<210> SEQ ID NO 42
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 42

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ser Thr
 1           5           10          15
Cys Leu Leu Ile Ser Trp Arg Pro Asn Pro Arg Leu Ser Arg Tyr Tyr
          20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35           40           45
Thr Val Pro Gly Leu Phe Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
          50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Lys Glu Thr
65           70           75           80
Ser Asn Ile Phe Ile Ala Pro Ile Ser Ile Asn Tyr Arg Thr
          85           90

```

```

<210> SEQ ID NO 43
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

-continued

<400> SEQUENCE: 43

```

Val Ser Asp Val Pro Arg Asp Pro Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Ile Ser Trp Asp Pro Asn Ile Arg Leu Arg Arg Tyr Tyr
      20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45
Thr Val Pro Gly Phe Phe Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Ser Arg Asn
65             70             75             80
Glu Asp Thr Arg Phe Gly Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

<210> SEQ ID NO 44

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Ile Ser Trp Phe Arg Ser Leu Gln Arg Asp Arg Asp Tyr
      20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45
Thr Val Pro Phe Arg Met Lys Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Ile Thr Pro Pro Asp Lys
65             70             75             80
Met Glu Pro Pro Lys Gly Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

<210> SEQ ID NO 45

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Ile Ser Trp Tyr Arg His Thr Tyr Arg Asp Arg Tyr Tyr
      20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Ser
      35             40             45
Thr Val Pro Pro Trp Ala Thr Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60
Gly Val Asp Tyr Thr Ile Ala Val Tyr Ala Val Thr Asp Thr Gly Tyr
65             70             75             80
Asp Val His Thr Lys Arg Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

<210> SEQ ID NO 46

<211> LENGTH: 94

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Ala Pro Thr
 1             5             10             15

Ser Leu Leu Ile Ser Trp Tyr Arg His Thr Tyr Arg Asp Arg Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45

Thr Val Pro Pro Trp Ala Thr Thr Ala Ala Ile Ser Gly Leu Lys Pro
      50             55             60

Gly Val Asp Tyr Ala Ile Ala Val Tyr Ala Val Thr Asp Thr Gly Tyr
      65             70             75             80

Asp Val His Thr Lys Arg Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 47
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15

Ser Gln Leu Ile Ser Trp Pro Phe Gly Trp Tyr Pro Ser Arg Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45

Thr Val Pro Pro Trp Ala Arg Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Tyr Ser Asp
      65             70             75             80

Phe Ser Gln Val His Thr Pro Asn Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 48
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15

Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Ser Arg Tyr Arg
      20             25             30

Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr
      35             40             45

Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro Gly
      50             55             60

Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Leu Pro Ile Pro
      65             70             75             80

Thr Leu Val His Gly Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

-continued

```

<210> SEQ ID NO 49
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49
Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Arg Leu Ile Ser Trp Ala Ser Pro Pro Met Trp Cys Arg Tyr Tyr
      20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Asn Gly Pro Val Gln Glu Phe
 35             40             45
Thr Val Pro Pro Trp Ala Thr Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Glu Tyr Leu Pro
65             70             75             80
Glu Trp Asn Met Thr Gln Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 50
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50
Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Arg Leu Ile Ser Trp Ala Ser Pro Pro Met Trp Cys Arg Tyr Tyr
      20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35             40             45
Thr Val Pro Pro Trp Ala Thr Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50             55             60
Gly Val Asp Tyr Thr Ile Thr Met Tyr Ala Val Thr Glu Tyr Leu Pro
65             70             75             80
Glu Trp Asn Met Thr Gln Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 51
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51
Asn Thr Thr His Tyr Arg Lys Asn Asn Tyr Tyr Ala Thr Pro Thr Ser
 1             5             10             15
Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr Arg
      20             25             30
Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr
 35             40             45
Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro Gly
 50             55             60
Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp Thr
65             70             75             80
Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

-continued

<210> SEQ ID NO 52
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 52

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser Ala Thr Arg
65 70 75 80
Thr Val Lys Arg Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 53
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

Val Ser Asp Ala Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Met Tyr Ala Val Thr Ser Asn Val Gly
65 70 75 80
Arg Leu Asp Thr Arg Tyr Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 54
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

Val Ser Asp Val Pro Arg Asp Leu Asp Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Glu Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser Asn Val Gly
65 70 75 80

-continued

Arg Leu Asp Thr Arg Tyr Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 55
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Lys Glu Pro Gln
65 70 75 80

Arg His Ala Leu Val Thr Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 56
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 56

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Glu Thr Pro Ser
65 70 75 80

Thr Lys Pro His Asn Val Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 57
<211> LENGTH: 74
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Arg Leu Ile Ser Trp Asn His Pro Gly Pro Phe Ser Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Thr Val Pro Pro Trp Ala Arg Thr Ala Ile Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala

-continued

```

<210> SEQ ID NO 59
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ser Thr
 1             5             10             15

Ser Leu Leu Ile Ser Trp His Tyr Leu Arg Arg Gln Pro Arg Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45

Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Gly Gly Leu Lys Pro
  50             55             60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Gln Thr Gly
65             70             75             80

His His Leu His Asp Glu Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 60
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Val Ser Asp Val Pro Arg Asp Leu Gln Ile Val Ala Ala Thr Pro Thr
 1          5          10          15
Ser Leu Leu Ile Ser Trp Asp Ile Ser Arg Tyr Lys His Arg Tyr Tyr
 20          25          30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asp Ser Pro Val Gln Glu Phe
 35          40          45
Thr Ala Pro Pro Trp Ala Ser Ile Ala Thr Ile Gly Leu Lys Pro
 50          55          60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Gln Thr Gly
 65          70          75          80

```

-continued

Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro	Thr
1				5					10					15	
Ser	Arg	Leu	Ile	Ser	Trp	Arg	Pro	Thr	Ser	Asn	Pro	Pro	Arg	Tyr	Tyr
		20						25					30		
Arg	Ile	Ser	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Phe
		35					40					45			
Thr	Val	Pro	Pro	Trp	Ala	Ser	Ile	Ala	Thr	Ile	Gly	Gly	Leu	Lys	Pro
	50					55					60				

-continued

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Gln Thr Gly
65 70 75 80

His His Leu His Asp Lys Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 64
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Gln Leu Ile Ser Trp Lys Thr Thr Asn Pro Thr Ala Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Thr Val Pro Pro Trp Ala Thr Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asn Leu Thr Thr
65 70 75 80

Arg Arg Arg His Arg Ala Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 65
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Arg Leu Ile Ser Trp Thr Thr Arg His Ser Pro Val Arg Tyr Tyr
20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45

Ile Val Pro Pro Trp Ala Thr Thr Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Met Pro Thr Asn
65 70 75 80

Trp Arg Phe Pro His Arg Pro Ile Ser Ile Asp Tyr Arg Thr
85 90

<210> SEQ ID NO 66
<211> LENGTH: 90
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

Val Ser Asp Val Pro Arg Asp Leu Glu Ala Val Ala Ala Thr Pro Thr
1 5 10 15

Ser Leu Leu Ile Arg Glu Arg Glu Arg Arg Tyr Tyr Arg Ile Thr Tyr
20 25 30

Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr Val Pro Gly
35 40 45

Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Glu Pro Gly Val Asp Tyr

-continued

50	55	60
Thr Ile Thr Val Tyr	Ala Val Thr Pro His His	Gly His Phe Asp Leu
65	70	75 80
Glu Leu Pro Ile Ser	Ile Asn Tyr Arg Thr	
	85	90

<210> SEQ ID NO 67
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 67

Val Ser Asp Val Pro Arg Asp Leu Glu Gly Val Ala Ala Thr Pro Thr	
1 5 10 15	
Ser Leu Leu Ile Ser Arg Lys Asp Arg Val Ser Ser Arg Arg Tyr Tyr	
20 25 30	
Arg Ile Thr Tyr Gly Glu Thr Gly Asn Ser Pro Val Gln Glu Phe	
35 40 45	
Thr Val Pro Gly Ser Lys Ser Thr Ala Ile Ile Ser Gly Leu Lys Pro	
50 55 60	
Gly Val Asp Tyr Thr Ile Thr Ala Tyr Val Val Thr Pro His His Gly	
65 70 75 80	
His Phe Asp Leu Glu Leu Pro Ile Ser Ile Asn Tyr Arg Thr	
85 90	

<210> SEQ ID NO 68
 <211> LENGTH: 95
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 68

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr	
1 5 10 15	
Ser Leu Leu Ile Ser Trp His Met Ala Thr Pro Asn Thr Arg Tyr Tyr	
20 25 30	
Arg Thr Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	
35 40 45	
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro	
50 55 60	
Gly Val Asp Tyr Asn Thr Asn Thr Val Tyr Ala Val Thr Ser Val Asn	
65 70 75 80	
Ala Phe Pro Tyr Glu Gly Met Pro Ile Ser Ile Asn Tyr Arg Thr	
85 90 95	

<210> SEQ ID NO 69
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 69

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ala Thr	
1 5 10 15	
Ser Leu Leu Ser Ser Trp Tyr Leu Cys Thr Gly Asn Asn Arg Asp Tyr	
20 25 30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Ala Phe	
35 40 45	

-continued

```

Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50                      55                      60

Gly Val Asp Tyr Ile Pro Ser Arg Cys Met Leu Ser Leu Ala Ser Leu
65                      70                      75                      80

Met Ser Thr Arg Asn Lys Pro Ile Ser Ile Asn Tyr Arg Thr
                      85                      90

```

```

<210> SEQ ID NO 70
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 70

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1                      5                      10                      15

Ser Leu Leu Ile Ser Trp Arg Thr Pro Ala Ser Pro His Gly Tyr Tyr
                20                      25                      30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Glu Glu Phe
    35                      40                      45

Thr Val Pro Leu Leu Trp Pro Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50                      55                      60

Gly Val Asp Tyr Ala Ile Thr Val Tyr Ala Val Thr Pro Thr His Met
65                      70                      75                      80

Leu Lys Pro Leu Ser Met Pro Ile Ser Ile Asn Tyr Arg Thr
                85                      90

```

```

<210> SEQ ID NO 71
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 71

```

```

Ile Ser Asp Val Pro Arg Asp Met Glu Val Val Ala Ala Thr Pro Thr
 1                      5                      10                      15

Ser Leu Leu Ile Ser Trp Asn Met Ala His Pro His Asp Arg Asn Tyr
    20                      25                      30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Ser
    35                      40                      45

Thr Val Pro Arg Tyr Leu Ser Thr Ala Thr Ile Ser Gly Pro Lys Arg
 50                      55                      60

Val Asp Tyr Thr Ile Ile Val Tyr Ala Val Asn Gln Pro Thr Val Ser
65                      70                      75                      80

Ala His Asn His Ala Pro Ile Ser Ile Asn Tyr Arg Thr
                85                      90

```

```

<210> SEQ ID NO 72
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 72

```

```

Val Ser Asp Val Pro Arg Asp Leu Lys Val Val Ala Ala Thr Pro Thr
 1                      5                      10                      15

Ser Leu Leu Ile Ser Trp Phe Pro Asp Asn Ala Thr Pro Arg Tyr Tyr
    20                      25                      30

```

-continued

```

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
   35                               40               45

Thr Val Pro Leu Phe Thr Thr Ala Thr Ile Ser Gly Leu Lys Pro Gly
   50                               55               60

Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser His Arg Asp Tyr
   65                               70               75               80

His Ser Thr Gly Arg Pro Ile Ser Ile Asn Tyr Arg Thr
           85                               90

```

```

<210> SEQ ID NO 73
<211> LENGTH: 95
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 73

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
   1           5           10           15

Ser Leu Leu Ile Ser Trp Met Leu Leu Arg Asp Asp Arg Arg Tyr Tyr
           20           25           30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
   35                               40               45

Thr Val Pro Thr Phe His Pro Thr Ala Thr Ile Ser Gly Arg Lys Pro
   50                               55               60

Gly Val Asp Tyr Asn Thr Ile Thr Val Tyr Ala Val Thr Gln Ser Thr
   65                               70               75               80

Asn Gly Asn Arg Asn Asp Phe Pro Ile Ser Ile Asn Tyr Arg Thr
           85                               90           95

```

```

<210> SEQ ID NO 74
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 74

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
   1           5           10           15

Ser Leu Leu Ile Ser Trp Ser Pro Pro Asn Asp Ala His Arg Tyr Tyr
           20           25           30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
   35                               40               45

Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
   50                               55               60

Gly Val Asp Tyr Thr Asn Thr Val Tyr Ala Val Thr Asp Gln Gln Ser
   65                               70               75               80

Tyr Thr Tyr Tyr Ser Asn Pro Ile Ser Ile Asn Tyr Arg Thr
           85                               90

```

```

<210> SEQ ID NO 75
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 75

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
   1           5           10           15

Ser Leu Val Ile Ser Trp Ser Pro Pro Asn Asp Ala His Arg Tyr Tyr

```

-continued

```
<210> SEQ ID NO 76
<211> LENGTH: 92
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

Val 1	Ser	Asp	Val 5	Pro	Arg	Asp	Leu	Glu	Val 10	Val	Ala	Ala	Thr	Pro 15	Thr
Ser	Leu	Leu	Ile 20	Ser	Trp	Ser	Pro	Pro 25	Asn	Asp	Ala	His	Arg 30	Tyr	Tyr
Arg	Ile	Thr 35	Tyr	Gly	Glu	Thr	Gly 40	Gly	Asn	Ser	Pro	Val 45	Gln	Glu	Phe
Thr	Val 50	Pro	Pro	Trp	Ala	Thr 55	Thr	Ala	Thr	Ile	Ser 60	Gly	Leu	Lys	Pro
Gly 65	Val	Asp	Tyr	Thr	Ile 70	Thr	Val	Tyr	Thr	Met 75	Pro	Thr	Asn	Trp	Arg 80
Phe	Pro	His	Arg 85	Pro	Ile	Ser	Ile	Asn	Tyr 90	Arg	Thr				

```
<210> SEQ ID NO 77
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

Val 1	Ser	Asp	Val 5	Pro	Arg	Asp	Leu	Glu	Val 10	Val	Ala	Ala	Thr	Pro 15	Thr
Ser	Gln	Leu	Ile 20	Ser	Trp	Thr	Thr	Arg 25	His	Ser	Pro	Val	Arg 30	Tyr	Tyr
Arg	Ile	Thr 35	Tyr	Gly	Glu	Thr	Gly 40	Gly	Asn	Ser	Pro	Val 45	Gln	Glu	Phe
Thr	Val 50	Pro	Pro	Trp	Ala	Thr 55	Thr	Ala	Thr	Ile	Ser 60	Gly	Leu	Lys	Pro
Gly 65	Val	Asp	Tyr	Thr	Ile 70	Thr	Val	Tyr	Ala	Val 75	Thr	Met	Pro	Thr	Asn 80
Trp	Arg	Phe	Pro 85	His	Arg	Pro	Ile	Ser 90	Ile	Asn	Tyr	Arg	Thr		

```
<210> SEQ ID NO 78
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

Val Ser Asp Val Pro Arg Asp Leu Glu Ile Val Ala Ala Thr Pro Thr
1 5 10 15

-continued

Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Gly Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Thr Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser Asn Val Gly
65 70 75 80
Arg Leu Asp Thr Arg Tyr Pro Ile Ser Thr Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 79
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 17, 30, 34
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 79

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Xaa Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Xaa Tyr Tyr
20 25 30
Arg Xaa Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Ile Ser Gly Leu Lys Pro Gly
50 55 60
Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser Asn Val Gly Arg
65 70 75 80
Leu Asp Thr Arg Tyr Pro Ile Phe Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 80
<211> LENGTH: 76
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr
65 70 75

<210> SEQ ID NO 81
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

-continued

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10          15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
 20          25          30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35          40          45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
 50          55          60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
 65          70          75          80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
 85          90

```

```

<210> SEQ ID NO 82
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 82

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10          15
Ser Arg Leu Ile Ser Cys Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
 20          25          30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35          40          45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
 50          55          60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
 65          70          75          80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
 85          90

```

```

<210> SEQ ID NO 83
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 83

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10          15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
 20          25          30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Ser Ser Pro Val Gln Glu Phe
 35          40          45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
 50          55          60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Thr His Asn
 65          70          75          80
Trp Asn Asp Gln Thr Arg Ser Ile Ser Ile Asn Tyr Arg Thr
 85          90

```

```

<210> SEQ ID NO 84
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 84

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10           15
Ser Arg Leu Ile Ser Trp Arg Pro Thr Ser Asn Pro Pro Arg Tyr Tyr
          20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35           40           45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Gly Gly Leu Lys Pro
          50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Gln Thr Gly
65           70           75           80
Tyr His Leu His Asp Lys Pro Ile Ser Ile Asn Tyr Arg Thr
          85           90

```

<210> SEQ ID NO 85

<211> LENGTH: 93

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: VARIANT

<222> LOCATION: 28, 51, 82

<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 85

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Pro Thr Ser
 1           5           10           15
Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Xaa Arg Tyr Tyr Arg
          20           25           30
Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr
          35           40           45
Val Pro Xaa Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro Gly
          50           55           60
Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Phe Pro Pro Gly Tyr
65           70           75           80
Pro Xaa Thr Glu Met Pro Ile Ser Ile Asn Tyr Arg Thr
          85           90

```

<210> SEQ ID NO 86

<211> LENGTH: 92

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

```

Ile Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1           5           10           15
Ser Leu Leu Ile Ser Trp Arg Arg Trp Pro His Phe Asp Arg Tyr Tyr
          20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35           40           45
Thr Val Pro Pro Trp Ala Thr Ile Ala Thr Ile Ser Gly Leu Lys Pro
          50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asn Pro Leu Ser
65           70           75           80
Pro Thr Thr Leu His Pro Pro Ile Asn Tyr Arg Thr
          85           90

```

-continued

<210> SEQ ID NO 87
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Lys Pro Arg Arg Thr Asn Thr Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Gly Thr Ile Ala Thr Ile Asn Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Leu Gly Thr Gly
65 70 75 80
Val Tyr Thr Arg Ala Gln Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 88
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 88

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Gln Leu Ile Ser Trp Pro Phe Gly Trp Tyr Pro Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Arg Thr Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr His Phe Pro Glu
65 70 75 80
Ser Arg Arg Pro Ala Lys Pro Met Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 89
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Leu Leu Ile Ser Trp His Thr Glu Arg Ser Phe Pro Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Gly Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Glu His Tyr Arg
65 70 75 80

-continued

Asp Thr Gly Thr Gly His Pro Ile Pro Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 90
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
 Ser Leu Leu Ile Ser Trp His Thr Glu Arg Ser Phe Pro Arg Tyr Tyr
20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
 Thr Val Pro Pro Trp Gly Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
 Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Glu His Tyr Arg
65 70 75 80
 Asp Thr Gly Thr Gly His Pro Ile Pro Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 91
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
 Ser Gln Leu Ile Ser Trp Lys Ser His Thr Phe His Pro Arg Tyr Tyr
20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
 Thr Val Pro Pro Trp Ala Ser Thr Ala Ala Ile Ser Gly Leu Lys Pro
50 55 60
 Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Leu Asn Arg Ser
65 70 75 80
 Ser Pro Asn Ser Ala Arg Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 92
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
 Ser Leu Leu Ile Ser Trp Arg Pro Gln Val Val Ser Thr Arg Tyr Tyr
20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
 Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
 Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asn His Lys Ala

-continued

65		70		75		80
----	--	----	--	----	--	----

Asn His His Asp Ala Glu Pro Ile Ser Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 93
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Arg Pro Thr Ser Asn His Pro Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Gly Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Thr Thr Asn Glu
65 70 75 80
Asp His Val Tyr Ala Leu Pro Ile Ser Ile Asn Tyr Arg Ile
85 90

<210> SEQ ID NO 94
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Leu
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 95
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Gly Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60

-continued

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Met Ser Asp
 65 70 75 80

Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 96
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 40
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 96

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1 5 10 15

Ser Leu Leu Ile Ser Trp Asp Thr His Asn Ala Tyr Asn Gly Tyr Tyr
 20 25 30

Arg Ile Thr Tyr Gly Glu Thr Xaa Gly Asn Ser Pro Val Arg Glu Phe
 35 40 45

Thr Val Pro His Pro Glu Val Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50 55 60

Gly Val Asp Asp Thr Ile Thr Val Tyr Ala Val Thr Asn His His Met
 65 70 75 80

Pro Leu Arg Ile Phe Gly Pro Ile Ser Ile Asn His Arg Thr
 85 90

<210> SEQ ID NO 97
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 13, 21
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 97

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Xaa Thr Pro Thr
 1 5 10 15

Ser Leu Leu Ile Xaa Trp Thr Arg Thr Asn Ala Asn Thr Arg Tyr Tyr
 20 25 30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35 40 45

Thr Ala Pro Asn Asn Pro Pro Thr Ala Thr Ile Gly Gly Leu Lys Pro
 50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Asp Gly Ser
 65 70 75 80

Arg His Met Leu Thr Lys Pro Ile Ser Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 98
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

Leu Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr

-continued

1	5	10	15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr	20	25	30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	35	40	45
Thr Val Pro Pro Trp Ala Ser Ile Ala Ala Ile Ser Gly Leu Lys Pro	50	55	60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp	65	70	75
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr	85	90	

<210> SEQ ID NO 99
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr	1	5	10	15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr	20	25	30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Leu	35	40	45	
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro	50	55	60	
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp	65	70	75	80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn His Arg Thr	85	90		

<210> SEQ ID NO 100
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 100

Val Ser Asp Val Pro Arg Gly Leu Glu Val Val Ala Ala Thr Pro Thr	1	5	10	15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr	20	25	30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	35	40	45	
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys His	50	55	60	
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp	65	70	75	80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr	85	90		

<210> SEQ ID NO 101
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

-continued

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Glu Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Met Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 102
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Thr Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 103
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Ala Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 104
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 104

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 105

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Cys Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Gln Arg Asp
65 70 75 80
Thr Tyr Arg Tyr Asp Asp Pro Ile Ser Thr Asn Cys Arg Thr
85 90

<210> SEQ ID NO 106

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 106

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
1 5 10 15
Ser Arg Leu Ile Ser Trp Arg Asn Ile Tyr Pro Ile Ala Arg Tyr Tyr
20 25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
35 40 45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
50 55 60
Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Asp
65 70 75 80
Thr Tyr Lys Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 107

<211> LENGTH: 92

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 27
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 107
Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ala Thr
 1             5             10             15
Ser Gln Leu Ile Ser Trp Pro Trp Pro Ser Xaa Pro Thr Arg Tyr Tyr
          20             25             30
Arg Ile Thr Tyr Gly Glu Thr Glu Gly Asn Ser Pro Val Gln Glu Phe
          35             40             45
Thr Val Pro Pro Trp Ala Ser Thr Ala Thr Ile Ser Gly Ile Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Ala Val Tyr Ala Val Thr Met Pro Glu Arg
65             70             75             80
Lys Tyr Asp Lys Pro Ile Ser Ile Asn Tyr Arg Thr
          85             90

```

```

<210> SEQ ID NO 108
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108
Val Ser Asp Val Ser Arg Asp Leu Glu Ala Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Ile Ser Trp Asn Pro Asn Arg Ser Phe Ala Arg Tyr Tyr
          20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35             40             45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Gly Gly Leu Lys Pro
          50             55             60
Arg Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Gln Thr Gly
65             70             75             80
His His Leu His Asp Lys Ser Ile Pro Ile Asn Tyr Arg Thr
          85             90

```

```

<210> SEQ ID NO 109
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 109
Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Ser Arg Tyr Tyr
          20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Ser
          35             40             45
Thr Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Phe Pro Pro Gly
65             70             75             80

```

-continued

Tyr Pro Leu Thr Glu Met Pro Ile Ser Ile Asn Tyr Arg Thr
85 90

<210> SEQ ID NO 110
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 110

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Ser
 1 5 10 15
 Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Ser Arg Tyr Tyr
 20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35 40 45
 Thr Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50 55 60
 Gly Val Asp Tyr Thr Ile Ala Val Tyr Ala Val Thr Phe Pro Thr Gly
 65 70 75 80
 Tyr Pro Leu Thr Glu Met Pro Ile Ser Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 111
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 111

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1 5 10 15
 Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Ser Arg Tyr Tyr
 20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35 40 45
 Thr Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro
 50 55 60
 Gly Val Asp Tyr Thr Ile Thr Ala Tyr Ala Val Thr Tyr Thr His Ser
 65 70 75 80
 Thr Pro Met Gln Asp Glu Pro Ile Ser Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 112
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 112

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1 5 10 15
 Ser Arg Leu Ile Ser Trp Asp Asn Ser Arg Pro Asn Thr Arg Tyr Tyr
 20 25 30
 Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
 35 40 45
 Thr Val Pro Pro Trp Gly Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
 50 55 60
 Gly Val Lys Tyr Thr Ile Thr Val Tyr Ala Val Thr Thr Ser Glu Cys

-continued

65		70		75		80							
His	Lys	Leu	Ser	Ser	Thr	Ser	Ile	Ser	Ile	Asn	Tyr	Arg	Thr
				85					90				

<210> SEQ ID NO 113
 <211> LENGTH: 91
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 113

Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro	Thr
1				5					10					15	
Ser	Leu	Leu	Ile	Ser	Trp	Thr	Arg	Thr	Asn	Ala	Ser	Thr	Arg	Tyr	Tyr
			20					25					30		
Arg	Ile	Thr	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Phe
	35					40						45			
Thr	Val	Pro	Asn	Phe	Trp	Trp	Ile	Ser	Gly	Leu	Lys	Pro	Gly	Val	Asp
	50				55						60				
Tyr	Thr	Ile	Thr	Val	Tyr	Ala	Val	Ala	Ser	Pro	Asp	Glu	Thr	Ser	Ala
65				70					75					80	
Tyr	Ser	Glu	Pro	Ile	Ser	Ile	Asn	Tyr	Arg	Thr					
			85					90							

<210> SEQ ID NO 114
 <211> LENGTH: 92
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 3, 18, 23
 <223> OTHER INFORMATION: Xaa = Any Amino Acid
 <400> SEQUENCE: 114

Val	Ser	Xaa	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Ala	Ala	Thr	Pro	Thr
1				5					10					15	
Ser	Xaa	Leu	Ile	Ser	Trp	Xaa	Pro	Arg	Ser	His	His	Asp	Arg	Tyr	Tyr
		20					25						30		
Arg	Ile	Thr	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Phe
	35					40						45			
Thr	Val	Pro	Pro	Trp	Gly	Thr	Ile	Ala	Thr	Ile	Asp	Gly	Leu	Lys	Pro
	50				55					60					
Gly	Val	Gly	Tyr	Thr	Val	Thr	Val	Tyr	Ala	Val	Thr	Asp	Asn	Pro	Asn
65				70					75					80	
Ser	Ala	Lys	Ala	Gln	His	Pro	Ile	Asn	Ser	Arg	Thr				
			85					90							

<210> SEQ ID NO 115
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 115

Val	Ser	Asp	Val	Pro	Arg	Asp	Leu	Glu	Val	Val	Val	Ala	Thr	Pro	Thr
1				5					10					15	
Ser	Gln	Leu	Ile	Ser	Trp	Met	Thr	Pro	His	Asn	His	Val	Arg	Tyr	Tyr
		20					25						30		
Gly	Ile	Thr	Tyr	Gly	Glu	Thr	Gly	Gly	Asn	Ser	Pro	Val	Gln	Glu	Ser

-continued

35	40	45
Thr Val Pro Thr Gly Asn Ala Thr Ala Thr Ile Ser Gly Leu Lys Pro		
50	55	60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro His His Gly		
65	70	75 80
His Phe Asp Leu Glu Pro Pro Ile Ser Ile Asn Tyr Arg Thr		
85	90	

<210> SEQ ID NO 116
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 116

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Ser Thr		
1	5	10 15
Gly Leu Leu Ile Ser Trp Arg Thr Pro Ala Ser Pro His Gly Tyr Tyr		
20	25	30
Arg Ile Thr Tyr Gly Glu Thr Gly Asn Ser Pro Val Glu Glu Phe		
35	40	45
Thr Val Pro Leu Leu Trp Pro Thr Ala Thr Ile Ser Gly Leu Lys Pro		
50	55	60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Thr His Met		
65	70	75 80
Leu Lys Pro Gln Ser Met Pro Ile Ser Ile Asn Tyr Arg Thr		
85	90	

<210> SEQ ID NO 117
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 117

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr		
1	5	10 15
Ser Leu Leu Ile Ser Trp Ser Pro Pro Asn Asp Ala His Arg Tyr Tyr		
20	25	30
Arg Ile Thr Tyr Gly Lys Thr Gly Gly Asp Ser Pro Val Gln Glu Phe		
35	40	45
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro		
50	55	60
Gly Val Asp Tyr Thr Ser Val Val Tyr Ala Val Thr Asp Gln Gln Ser		
65	70	75 80
Tyr Thr Tyr Tyr Ser Asn Pro Ile Ser Ile Asn Tyr Arg Thr		
85	90	

<210> SEQ ID NO 118
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 28
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 118

Val Ser Asp Val Pro Ser Asp Leu Glu Val Val Ala Ala Thr Pro Thr

-continued

1	5	10	15
Ser Leu Leu Ile Ser Trp Glu Gln Ser Pro Thr Xaa Gly Arg Tyr Tyr	20	25	30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	35	40	45
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Arg Lys Pro	50	55	60
Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ile Glu Lys Asp	65	70	75
Arg Ile Pro Leu Phe Gly Pro Ile Ser Ile Ser Tyr Arg Thr	85	90	

<210> SEQ ID NO 119
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 119

Val Ser Asp Val Pro Ser Asp Leu Glu Val Val Ala Ala Thr Pro Thr	1	5	10	15
Ser Leu Leu Ile Ser Trp Glu Gln Ser Pro Thr Tyr Gly Arg Tyr Tyr	20	25	30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	35	40	45	
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Arg Lys Pro	50	55	60	
Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ile Glu Lys Asp	65	70	75	80
Arg Ile Pro Leu Phe Gly Pro Ile Ser Ile Ser Tyr Arg Thr	85	90		

<210> SEQ ID NO 120
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 28
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 120

Val Ser Asp Val Pro Ser Asp Leu Glu Val Val Ala Ala Thr Pro Thr	1	5	10	15
Ser Leu Leu Ile Ser Trp Glu Gln Ser Pro Thr Xaa Gly Arg Tyr Tyr	20	25	30	
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	35	40	45	
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Arg Lys Pro	50	55	60	
Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ile Glu Lys Asp	65	70	75	80
Arg Ile Pro Leu Phe Gly Pro Ile Ser Ile Ser Tyr Arg Thr	85	90		

<210> SEQ ID NO 121
 <211> LENGTH: 94

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 121

Val Pro Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15

Ser Leu Leu Ile Ser Trp Asp Thr His Asn Ala Tyr Asn Gly Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Ser Ser Pro Ala Gln Glu Phe
      35             40             45

Thr Val Pro His Pro Glu Val Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60

Gly Val Asp Asp Thr Ile Thr Val Tyr Ala Val Thr Ile His His Met
65             70             75             80

Pro Leu Arg Ile Phe Gly Pro Ile Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 122
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 122

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15

Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45

Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Glu Ser Asp
65             70             75             80

Thr Tyr Lys Tyr Asp Asp Pro Val Ser Ile Asn Tyr Arg Thr
      85             90

```

```

<210> SEQ ID NO 123
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 123

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15

Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
      20             25             30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35             40             45

Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
      50             55             60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Glu Ser Asp
65             70             75             80

Thr Tyr Lys Tyr Asp Asp Pro Val Ser Thr Asn Tyr Arg Thr
      85             90

```

-continued

```

<210> SEQ ID NO 124
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 124

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10            15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Gly Tyr Tyr
          20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35             40             45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Pro Asn Val Gly
65             70             75             80
Arg Leu Asp Thr Arg Tyr Pro Ile Ser Ile Asp Cys Arg Thr
          85             90

```

```

<210> SEQ ID NO 125
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 86
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 125

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10            15
Ser Arg Leu Ile Ser Trp Asn Arg Ser Gly Leu Gln Ser Arg Tyr Tyr
          20             25             30
Arg Thr Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
          35             40             45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ser Asn Val Gly
65             70             75             80
Arg Leu Asp Thr Arg Xaa Pro Ile Ser Ile Asn Tyr Arg Thr
          85             90

```

```

<210> SEQ ID NO 126
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 75
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 126

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10            15
Ser Arg Leu Ile Ser Trp Arg Thr Met Pro Val Thr Ala Arg Tyr Tyr
          20             25             30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asp Ser Pro Val Gln Glu Phe
          35             40             45

```

-continued

Thr Val Pro Pro Trp Ala Ser Ile Ala Ala Ile Ser Gly Leu Lys Pro
 50 55 60

Gly Ala Asp Tyr Thr Ile Thr Val Tyr Ala Xaa Thr Ser Ala Thr Pro
 65 70 75 80

Ser Arg Pro Asn Val His Pro Ile Ser Ile Asn Leu Thr Thr
 85 90

<210> SEQ ID NO 127
 <211> LENGTH: 88
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 127

Val Ser Asp Val Pro Gly Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1 5 10 15

Ser Leu Leu Ile Gly Trp Ser Met Thr Pro Asn Trp Pro Arg Tyr Tyr
 20 25 30

Arg Ile Ala Tyr Gly Glu Thr Gly Gly Asp Ser Pro Val Gln Glu Phe
 35 40 45

Thr Val Pro Pro Trp Ala Ser Ile Ala Ile Ile Gly Gly Leu Lys Pro
 50 55 60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr His Arg Asp Thr
 65 70 75 80

Pro Ile Ser Ile Asn Tyr Arg Thr
 85

<210> SEQ ID NO 128
 <211> LENGTH: 90
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 128

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Ile
 1 5 10 15

Ser Gln Leu Thr Ser Trp Gln Pro Gln Pro Asn Gly Ser Arg Tyr Tyr
 20 25 30

Arg Ile Ala Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Arg Glu Phe
 35 40 45

Thr Val Pro Ala Arg Glu Gln Thr Ala Thr Ser Gly Leu Lys Pro Gly
 50 55 60

Val Asp Tyr Ala Ile Thr Val Tyr Ala Ala Thr His Gly Lys Pro Pro
 65 70 75 80

His Ile His Phe Thr Ile Asn Tyr Arg Thr
 85 90

<210> SEQ ID NO 129
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 2, 9, 12, 22
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 129

Val Xaa Asp Val Pro Arg Asp Leu Xaa Val Val Xaa Ala Thr Pro Thr
 1 5 10 15

-continued

```

Ser Leu Leu Ile Ser Xaa Arg Ser Gly Asn Arg Thr Thr Arg Tyr Tyr
      20              25              30

Arg Ile Thr Tyr Gly Asp Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35              40              45

Thr Met Pro Pro Trp Ala Thr Val Ala Ala Ile Ser Gly Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Thr His Asn Ser
      65              70              75              80

Thr Ala Gln Pro Glu Tyr Pro Ile Pro Phe Asn Arg Arg Thr
      85              90

```

```

<210> SEQ ID NO 130
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 130

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
  1              5              10              15

Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Thr Tyr Ser Arg Tyr Tyr
      20              25              30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35              40              45

Thr Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Cys Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Ala Phe Pro Pro Gly
      65              70              75              80

Tyr Pro Leu Thr Glu Met Pro Ile Ser Ile Asn Tyr Arg Thr
      85              90

```

```

<210> SEQ ID NO 131
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 131

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
  1              5              10              15

Ser Arg Leu Ile Ser Trp Arg Pro Gly Arg Ala Tyr Ser Arg Tyr Phe
      20              25              30

Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe
      35              40              45

Thr Val Pro Pro Trp Ala Asn Thr Ala Thr Ile Ser Gly Leu Lys Pro
      50              55              60

Gly Val Asp Tyr Thr Ile Ala Val Tyr Ala Val Thr Phe Pro Pro Arg
      65              70              75              80

Tyr Pro Leu Thr Glu Met Pro Ile Ser Ile Asn Tyr Arg Ala
      85              90

```

```

<210> SEQ ID NO 132
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 132

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr

```

-continued

1	5	10	15
Ser Arg Leu Ile	Ser Trp Arg Pro Gly Arg Thr Tyr	Ser Arg Tyr Tyr	
20	25	30	
Arg Ile Thr Tyr Gly Glu Ala Gly Gly Asn Ser Pro Val Gln Glu Phe			
35	40	45	
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro			
50	55	60	
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Asp Lys Ser Gly			
65	70	75	80
Thr Tyr Arg Tyr Asp Asp Pro Ile Ser Ile Asn Tyr Arg Thr			
85	90		

<210> SEQ ID NO 133
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 133

Val Ser Asp Val Pro Arg Asp Leu Arg Val Val Ala Ala Thr Pro Thr	
1	5 10 15
Ser Arg Leu Ile Ser Trp Arg Pro Ala Ser Asn Pro Ala Arg Tyr Tyr	
20	25 30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe	
35	40 45
Thr Val Pro Pro Trp Ala Ser Val Ala Thr Ile Gly Gly Leu Lys Pro	
50	55 60
Gly Val Asp Tyr Thr Val Thr Val Tyr Ala Val Thr Ala Gln Thr Gly	
65	70 75 80
His Arg Leu His Asp Lys Pro Ile Ser Ile Asn Tyr Arg Thr	
85	90

<210> SEQ ID NO 134
 <211> LENGTH: 87
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 134

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Ser	
1	5 10 15
Leu Leu Ile Ser Trp Arg Pro Pro Ala Asp Leu Asn Arg Tyr Tyr Arg	
20	25 30
Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr	
35	40 45
Val Pro Pro Trp Gly Thr Val Ala Thr Val Asn Gly Leu Lys Pro Gly	
50	55 60
Val Gly Tyr Thr Ile Thr Val Tyr Ala Val Thr His Arg Asp Thr Pro	
65	70 75 80
Ile Ser Ile Asn Tyr Arg Ala	
85	

<210> SEQ ID NO 135
 <211> LENGTH: 93
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 135

-continued

```

Val Thr Asp Val Pro Arg Gly Leu Lys Ile Val Ala Ala Thr Pro Ser
 1          5          10          15
Leu Leu Ile Ser Trp Arg Asn Ala Lys Asp Pro Gly Arg Tyr Tyr Arg
          20          25          30
Ile Thr Tyr Gly Glu Thr Gly Gly Ser Ser Pro Val Gln Glu Phe Thr
          35          40          45
Val Pro Pro Trp Gly Thr Ile Ala Ala Ile Asn Gly Leu Lys Pro Gly
          50          55          60
Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Ala Thr Asn Pro Gly
65          70          75          80
Pro Thr Gln His Arg Pro Ile Pro Ile Asn Tyr Arg Thr
          85          90

```

```

<210> SEQ ID NO 136
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 136

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Asp Pro His
 1          5          10          15
Gln Pro Leu Ile Cys Trp Ala Ser Pro Pro Met Trp Cys Arg Tyr Tyr
          20          25          30
Arg Ile Thr Tyr Gly Glu Ser Gly Gly Asn Ser Pro Val Gln Glu Phe
          35          40          45
Thr Val Pro Pro Trp Ala Thr Ala Ala Ala Ile Ser Gly Leu Lys Pro
          50          55          60
Gly Val Asp Tyr Thr Ile Thr Val His Ala Val Thr Asp Glu Ser Trp
65          70          75          80
Ser Asp Arg Ser Met Asp Pro Ile Ser Ile Asn Cys Arg Thr
          85          90

```

```

<210> SEQ ID NO 137
<211> LENGTH: 94
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 137

```

```

Val Ser Asp Val Pro Arg Asp Leu Lys Val Val Ala Ala Thr Pro Thr
 1          5          10          15
Ser Arg Leu Ile Ser Trp Thr His Asp Asn Val Pro Ala Arg Tyr Tyr
          20          25          30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Leu
          35          40          45
Thr Val Pro Pro Trp Ala Ser Ile Ala Thr Ile Ser Gly Leu Lys Pro
          50          55          60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Leu Tyr Thr Gly
65          70          75          80
Asn His Arg Pro Glu His Pro Ile Ser Ile Asn Tyr Arg Thr
          85          90

```

```

<210> SEQ ID NO 138
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 138

```

Val Ser Asp Val Pro Arg Asp Pro Val Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Ile Ser Trp Tyr Arg His Thr Tyr Arg Asp Arg Tyr Tyr
          20             25             30
Arg Val Thr Tyr Gly Glu Thr Arg Gly Asn Ser Pro Ile Arg Glu Phe
          35             40             45
Thr Val Pro Pro Trp Ala Thr Ile Ala Thr Ile Ser Gly Leu Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Ala Val Tyr Ala Val Thr Asp Ala Gly Tyr
65             70             75             80
Asp Val His Thr Lys Arg Pro Ile Ser Ile Asn Arg Thr
          85             90

```

<210> SEQ ID NO 139

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 139

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Gly Leu Leu Ile Ser Trp Arg Asn Asn Gln Tyr Thr Pro Arg His Tyr
          20             25             30
Gly Ile Thr Tyr Gly Glu Thr Gly Gly Lys Ser Pro Val Gln Glu Phe
          35             40             45
Thr Val Pro Glu Leu Asn Pro Thr Ala Thr Ile Ser Arg Leu Lys Pro
          50             55             60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gln Asn Gly Thr
65             70             75             80
Pro Arg Val Ile Tyr Gly Pro Ile Ser Ile Asn Tyr Arg Thr
          85             90

```

<210> SEQ ID NO 140

<211> LENGTH: 89

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 140

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr
 1             5             10             15
Ser Leu Leu Asn Val Pro Ile Ile Arg Tyr Tyr Arg Ile Thr Tyr Gly
          20             25             30
Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr Val Pro Ala Pro
          35             40             45
Lys Ala Ile Ala Thr Thr Ser Gly Leu Lys Pro Gly Val Asp Tyr Thr
          50             55             60
Ile Thr Val Tyr Gly Val Thr Ser His Arg Asn His Phe His Val Glu
65             70             75             80
Thr Pro Ile Ser Ile Asn Tyr Gln Ala
          85

```

<210> SEQ ID NO 141

<211> LENGTH: 21

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 141
Asp Ala Pro Ala Val Thr Val Gly Ser Lys Ser Gly Arg Gly Asp Ser
1 5 10 15
Pro Ala Ser Ser Lys
20
<210> SEQ ID NO 142
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 142
Ala Ser Pro Pro Met Trp Cys Pro Trp Ala Thr Glu Tyr Leu Pro Glu
1 5 10 15
Trp Asn Met Thr Gln
20
<210> SEQ ID NO 143
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 143
Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Asp Lys Ser Asp Thr
1 5 10 15
Tyr Lys Tyr Asp Asp
20
<210> SEQ ID NO 144
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 144
Arg Pro Thr Ser Asn Pro Pro Pro Trp Ala Ser Ala Gln Thr Gly His
1 5 10 15
His Leu His Asp Lys
20
<210> SEQ ID NO 145
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 145
His Thr Glu Arg Ser Phe Pro Pro Trp Gly Ser Glu His Tyr Arg Asp
1 5 10 15
Thr Gly Thr Gly His
20
<210> SEQ ID NO 146
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 146
Thr Thr Arg His Ser Pro Val Pro Trp Ala Thr Met Pro Thr Asn Trp

-continued

1	5	10	15
---	---	----	----

Arg Phe Pro His Arg
20

<210> SEQ ID NO 147
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 147

Arg Pro Asn Pro Arg Leu Ser Gly Leu Phe Ser Pro Lys Glu Thr Ser
1 5 10 15

Asn Ile Phe Ile Ala
20

<210> SEQ ID NO 148
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 148

Ser Pro Pro Asn Asp Ala His Gly Ser Lys Ser Asp Gln Gln Ser Tyr
1 5 10 15

Thr Tyr Tyr Ser Asn
20

<210> SEQ ID NO 149
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 149

Arg Thr Pro Ala Ser Pro His Leu Leu Trp Pro Pro Thr His Met Leu
1 5 10 15

Lys Pro Gln Ser Met
20

<210> SEQ ID NO 150
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 150

Tyr Arg His Thr Tyr Arg Asp Pro Trp Ala Thr Asp Thr Gly Tyr Asp
1 5 10 15

Val His Thr Lys Arg
20

<210> SEQ ID NO 151
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 151

Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Ser Asn Val Gly Arg
1 5 10 15

Leu Asp Thr Arg Tyr
20

-continued

<210> SEQ ID NO 152
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 152

Asp Thr His Asn Ala Tyr Asn His Pro Glu Val Asn His His Met Pro
1 5 10 15

Leu Arg Ile Phe Gly
20

<210> SEQ ID NO 153
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 153

Arg Pro Thr Ser Asn Pro Pro Pro Trp Ala Ser Pro Val Tyr Pro Met
1 5 10 15

His Ser Met Leu Ser
20

<210> SEQ ID NO 154
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 154

Arg Asn Ile Tyr Pro Ile Ala Pro Trp Ala Ser Asp Lys Ser Asp Thr
1 5 10 15

Tyr Lys Tyr Asp Asp
20

<210> SEQ ID NO 155
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 155

Asn Arg Ser Gly Leu Gln Cys Pro Trp Ala Ser Asp Gln Arg Asp Thr
1 5 10 15

Tyr Lys Tyr Asp Asp
20

<210> SEQ ID NO 156
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 156

Arg Pro Gly Arg Thr Tyr Ser Pro Trp Ala Asn Phe Pro Thr Gly Tyr
1 5 10 15

Pro Leu Thr Glu Met
20

<210> SEQ ID NO 157
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 157

-continued

Arg Pro Gly Arg Thr Tyr Ser Pro Trp Ala Asn Phe Pro Pro Gly Tyr
1 5 10 15

Pro Leu Thr Glu Met
20

<210> SEQ ID NO 158
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 158

Met Thr Pro His Asn His Val Thr Gly Asn Ala Pro His His Gly His
1 5 10 15

Phe Asp Leu Glu Pro
20

<210> SEQ ID NO 159
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 159

Thr Arg Thr Asn Ala Ser Thr Asn Phe Trp Trp Ser Pro Asp Glu Thr
1 5 10 15

Ser Ala Tyr Ser Glu
20

<210> SEQ ID NO 160
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 160

Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Asp Lys Ser Asp Thr
1 5 10 15

Tyr Lys Tyr Asp Asp
20

<210> SEQ ID NO 161
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 161

Arg Pro Gly Arg Thr Tyr Ser Pro Trp Ala Asn Tyr Thr His Ser Thr
1 5 10 15

Pro Met Gln Asp Glu
20

<210> SEQ ID NO 162
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 162

Arg Thr Pro Ala Ser Pro His Leu Leu Trp Pro Pro Thr His Met Leu
1 5 10 15

Lys Pro Gln Ser Met
20

-continued

<210> SEQ ID NO 163
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 163

Thr Arg Thr Asn Ala Asn Thr Asn Asn Pro Pro Pro Asp Gly Ser Arg
1 5 10 15

His Met Leu Thr Lys
20

<210> SEQ ID NO 164
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 164

Asp Asn Ser Arg Pro Asn Thr Pro Trp Gly Ser Thr Ser Glu Cys His
1 5 10 15

Lys Leu Ser Ser Thr
20

<210> SEQ ID NO 165
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 165

Asn Pro Asn Arg Ser Phe Ala Pro Trp Ala Ser Ala Gln Thr Gly His
1 5 10 15

His Leu His Asp Lys
20

<210> SEQ ID NO 166
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 166

Ser Met Thr Pro Asn Trp Pro Pro Trp Ala Ser His Arg Asp Thr
1 5 10 15

<210> SEQ ID NO 167
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 167

Asp Thr His Asn Ala Tyr Asn His Pro Glu Val Ile His His Met Pro
1 5 10 15

Leu Arg Ile Phe Gly
20

<210> SEQ ID NO 168
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 168

-continued

Ala	Ser	Pro	Pro	Met	Trp	Pro	Trp	Ala	Thr	Asp	Glu	Ser	Trp	Ser	Asp
1				5					10					15	

Arg	Ser	Met	Asp
			20

<210> SEQ ID NO 169
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 169

Arg	Pro	Pro	Ala	Asp	Leu	Asn	Pro	Trp	Gly	Thr	His	Arg	Asp	Thr
1				5					10				15	

<210> SEQ ID NO 170
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 170

Glu	Gln	Ser	Pro	Thr	Tyr	Gly	Gly	Ser	Lys	Ser	Ile	Glu	Lys	Asp	Arg
1				5					10					15	

Ile	Pro	Leu	Phe	Gly
				20

<210> SEQ ID NO 171
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 171

Arg	Pro	Gly	Arg	Thr	Tyr	Ser	Pro	Trp	Ala	Asn	Phe	Pro	Pro	Gly	Tyr
1				5					10					15	

Pro	Leu	Thr	Glu	Met
				20

<210> SEQ ID NO 172
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 172

Arg	Pro	Gly	Arg	Thr	Tyr	Ser	Pro	Trp	Ala	Ser	Asp	Lys	Ser	Gly	Thr
1				5					10					15	

Tyr	Arg	Tyr	Asp	Asp
				20

<210> SEQ ID NO 173
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 173

Tyr	Arg	His	Thr	Tyr	Arg	Asp	Pro	Trp	Ala	Thr	Asp	Ala	Gly	Tyr	Asp
1				5					10					15	

Val	His	Thr	Lys	Arg
				20

<210> SEQ ID NO 174
 <211> LENGTH: 21

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 174
Arg Thr Met Pro Val Thr Ala Pro Trp Ala Ser Ser Ala Thr Pro Ser
1 5 10 15
Arg Pro Asn Val His
20

<210> SEQ ID NO 175
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 175
Arg Pro Gly Arg Ala Tyr Ser Pro Trp Ala Asn Phe Pro Pro Arg Tyr
1 5 10 15
Pro Leu Thr Glu Met
20

<210> SEQ ID NO 176
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 176
Ser Pro Pro Asn Asp Ala His Gly Ser Lys Ser Asp Gln Gln Ser Tyr
1 5 10 15
Thr Tyr Tyr Ser Asn
20

<210> SEQ ID NO 177
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 177
Ile Ile Ala Pro Lys Ala Ser His Arg Asn His Phe His Val Glu Thr
1 5 10 15

<210> SEQ ID NO 178
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 178
Arg Asn Asn Gln Tyr Thr Pro Glu Leu Asn Pro Gln Asn Gly Thr Pro
1 5 10 15
Arg Val Ile Tyr Gly
20

<210> SEQ ID NO 179
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<400> SEQUENCE: 179
Arg Pro Ala Ser Asn Pro Ala Pro Trp Ala Ser Ala Gln Thr Gly His
1 5 10 15
Arg Leu His Asp Lys

-continued

20

<210> SEQ ID NO 180
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 180

Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Pro Asn Val Gly Arg
1 5 10 15
Leu Asp Thr Arg Tyr
20

<210> SEQ ID NO 181
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 181

Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Asp Glu Ser Asp Thr
1 5 10 15
Tyr Lys Tyr Asp Asp
20

<210> SEQ ID NO 182
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 182

Thr His Asp Asn Val Pro Ala Pro Trp Ala Ser Leu Tyr Thr Gly Asn
1 5 10 15
His Arg Pro Glu His
20

<210> SEQ ID NO 183
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 183

Arg Ser Gly Asn Arg Thr Thr Pro Trp Ala Thr Thr His Asn Ser Thr
1 5 10 15
Ala Gln Pro Glu Tyr
20

<210> SEQ ID NO 184
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 184

Asn Arg Ser Gly Leu Gln Ser Pro Trp Ala Ser Ser Asn Val Gly Arg
1 5 10 15
Leu Asp Thr Arg Tyr
20

<210> SEQ ID NO 185
<211> LENGTH: 21
<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 185

```

Arg Asn Ala Lys Asp Pro Gly Pro Trp Gly Thr Ala Thr Asn Pro Gly
 1           5           10           15
Pro Thr Gln His Arg
           20

```

<210> SEQ ID NO 186

<211> LENGTH: 94

<212> TYPE: PRT

<213> ORGANISM: Bovis taurus

<400> SEQUENCE: 186

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Ile Ala Ala Thr Pro Thr
 1           5           10           15
Ser Leu Leu Ile Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr
           20           25           30
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Ser Ser Pro Val Gln Glu Phe
           35           40           45
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro
           50           55           60
Gly Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly Asp
65           70           75           80
Ser Pro Ala Ser Ser Lys Pro Val Ser Ile Asn Tyr Arg Thr
           85           90

```

<210> SEQ ID NO 187

<211> LENGTH: 92

<212> TYPE: PRT

<213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 187

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Ile Ala Ser Thr Pro Thr
 1           5           10           15
Ser Leu Leu Ile Ser Trp Glu Pro Ala Val Ser Val Arg Tyr Tyr Arg
           20           25           30
Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr
           35           40           45
Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Asn Ile Lys Pro Gly Ala
           50           55           60
Asp Tyr Thr Ile Thr Leu Tyr Ala Val Thr Gly Arg Gly Asp Ser Pro
65           70           75           80
Ala Ser Ser Lys Pro Val Ser Ile Asn Tyr Gln Thr
           85           90

```

<210> SEQ ID NO 188

<211> LENGTH: 92

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 188

```

Val Ser Asp Ile Pro Arg Asp Leu Glu Val Ile Ala Ser Thr Pro Thr
 1           5           10           15
Ser Leu Leu Ile Ser Trp Glu Pro Ala Val Ser Val Arg Tyr Tyr Arg
           20           25           30

```

-continued

```

Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val Gln Glu Phe Thr
    35                      40                      45

Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Asn Ile Lys Pro Gly Ala
    50                      55                      60

Asp Tyr Thr Ile Thr Leu Tyr Ala Val Thr Gly Arg Gly Asp Ser Pro
    65                      70                      75                      80

Ala Ser Ser Lys Pro Val Ser Ile Asn Tyr Lys Thr
    85                      90

```

```

<210> SEQ ID NO 189
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculuc
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 24
<223> OTHER INFORMATION: Xaa = Any Amino Acid

```

```

<400> SEQUENCE: 189

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Ile Ala Ser Thr Pro Thr
  1          5          10          15

Ser Leu Leu Ile Ser Trp Glu Xaa Pro Ala Val Thr Val Arg Tyr Tyr
    20          25          30

Arg Ile Thr Tyr Gly Glu Thr Asn
  35          40

```

```

<210> SEQ ID NO 190
<211> LENGTH: 92
<212> TYPE: PRT
<213> ORGANISM: Gallus gallus

```

```

<400> SEQUENCE: 190

```

```

Val Ser Asp Val Pro Arg Asp Leu Glu Val Asn Thr Ser Pro Thr Ser
  1          5          10          15

Leu Glu Ile Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr Arg
    20          25          30

Ile Thr Tyr Gly Glu Thr Gly Gly Ser Ser Pro Val Gln Glu Phe Thr
    35          40          45

Val Pro Gly Thr Met Ser Ala Thr Ile Thr Gly Leu Lys Pro Gly Val
    50          55          60

Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly Asp Ser Pro
    65          70          75          80

Ala Ser Ser Lys Pro Val Thr Val Thr Tyr Lys Thr
    85          90

```

```

<210> SEQ ID NO 191
<211> LENGTH: 93
<212> TYPE: PRT
<213> ORGANISM: Xenopus laevis

```

```

<400> SEQUENCE: 191

```

```

Val Ser Asp Val Pro Thr Asp Leu Glu Val Thr Ser Ser Ser Pro Asn
  1          5          10          15

Thr Leu Thr Ile Ser Trp Glu Ala Pro Ala Val Ser Val Arg Tyr Tyr
    20          25          30

Arg Ile Thr Tyr Ser Gln Thr Gly Gly Gly Pro Glu Lys Glu Phe Thr
    35          40          45

```

-continued

Val Pro Gly Thr Ser Asn Thr Ala Thr Ile Arg Gly Leu Asn Pro Gly
 50 55 60

Val Ser Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly Asp Ser
 65 70 75 80

Pro Ala Ser Ser Lys Pro Leu Thr Ile Ile His Lys Thr
 85 90

<210> SEQ ID NO 192
 <211> LENGTH: 68
 <212> TYPE: PRT
 <213> ORGANISM: Canis familiaris

<400> SEQUENCE: 192

Ala Asp Ala Pro Ser Leu Phe Leu Ala Thr Thr Pro Ser Leu Leu Val
 1 5 10 15

Ser Trp Gln Pro Ala Ile Thr Gly Tyr Ile Ile Lys Tyr Gly Ser Glu
 20 25 30

Val Val Pro Gly Val Thr Ala Thr Ile Thr Gly Leu Pro Gly Thr Glu
 35 40 45

Tyr Thr Ile Gln Val Ile Ala Leu Lys Asn Gln Lys Ser Leu Ile Gly
 50 55 60

Lys Thr Glu Leu
 65

<210> SEQ ID NO 193
 <211> LENGTH: 67
 <212> TYPE: PRT
 <213> ORGANISM: Equus caballis

<400> SEQUENCE: 193

Ala Asp Ala Pro Ser Leu Phe Leu Ala Thr Thr Pro Ser Leu Leu Ile
 1 5 10 15

Ser Trp Gln Pro Ala Ile Thr Gly Tyr Ile Ile Lys Tyr Gly Ser Glu
 20 25 30

Val Val Pro Gly Val Thr Ala Thr Ile Thr Gly Leu Pro Gly Thr Glu
 35 40 45

Tyr Thr Ile Gln Val Ile Ala Ile Lys Asn Gln Lys Ser Leu Ile Gly
 50 55 60

Lys Thr Glu
 65

<210> SEQ ID NO 194
 <211> LENGTH: 70
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 194

Val Ser Pro Pro Lys Asp Leu Val Thr Val Thr Thr Val Asn Leu Ala
 1 5 10 15

Trp Asp Met Val Thr Tyr Leu Val Val Tyr Thr Pro Thr His Glu Gly
 20 25 30

Gly Glu Met Gln Phe Val Pro Gly Asp Gln Thr Ser Thr Ile Ile Gln
 35 40 45

Leu Pro Gly Val Glu Tyr Phe Ile Arg Val Phe Ala Ile Leu Asn Lys
 50 55 60

Lys Ser Val Ser Ala Val

-continued

```

65              70

<210> SEQ ID NO 195
<211> LENGTH: 69
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 195

Val Ser Pro Pro Lys Asp Leu Val Thr Val Thr Thr Val Asn Leu Ala
 1          5          10          15

Trp Asp Met Val Thr Tyr Leu Ile Val Tyr Thr Pro Thr His Glu Gly
      20          25          30

Glu Met Gln Phe Val Pro Gly Asp Gln Thr Ser Thr Thr Ile Arg Leu
 35          40          45

Pro Gly Val Glu Tyr Phe Ile Arg Val Phe Ala Ile Leu Asn Lys Lys
 50          55          60

Ser Val Ser Ala Val
65

```

```

<210> SEQ ID NO 196
<211> LENGTH: 75
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 196

Met Asp Gly Pro Gln Asp Leu Val Val Ala Val Thr Pro Thr Thr Leu
 1          5          10          15

Asp Leu Ser Trp Pro Gln Ala Val Asp Phe Val Val Ser Tyr Val Ser
      20          25          30

Ala Gly Asn Arg Val Leu Val Pro Pro Glu Ala Asp Thr Gln Leu Thr
      35          40          45

Leu Met Pro Gly Val Glu Tyr Val Val Thr Val Thr Ala Glu Arg Gly
 50          55          60

His Ala Val Ser Ala Ser Ile Ala Asn Thr Gly
65          70          75

```

```

<210> SEQ ID NO 197
<211> LENGTH: 69
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 197

Thr Val Pro Ser Leu Ile Tyr Val Gly Pro Thr Thr Met His Val Gln
 1          5          10          15

Trp Gln Val Gly Gly Ala Thr Gly Tyr Ile Leu Ser Tyr Pro Val Asp
      20          25          30

Thr Glu Thr Lys Glu Val Leu Gly Pro Thr Val Asn Met Gln Leu Thr
      35          40          45

Leu Val Pro Asn Thr Glu Tyr Ala Val Thr Val Gln Ala Val Leu Leu
 50          55          60

Thr Ser Val Thr Val
65

```

```

<210> SEQ ID NO 198
<211> LENGTH: 44
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

-continued

<400> SEQUENCE: 198

Thr Val Pro Ser Leu Asn Ile Tyr Val Gly Pro Thr Thr Met His Val
 1 5 10 15
 Gln Trp Gln Val Gly Gly Ala Thr Gly Tyr Ile Leu Ser Tyr Pro Val
 20 25 30
 Asp Thr Glu Thr Lys Gln Val Leu Arg Val Thr His
 35 40

<210> SEQ ID NO 199

<211> LENGTH: 66

<212> TYPE: PRT

<213> ORGANISM: Gallus gallus

<400> SEQUENCE: 199

Leu Ser Asp Leu Leu Tyr Val Ser Ser Ser Met Arg Ala Lys Trp Gly
 1 5 10 15
 Val Ala Gly Ala Thr Gly Tyr Met Ile Leu Tyr Ala Pro Leu Thr Glu
 20 25 30
 Gly Leu Ala Ala Glu Lys Glu Ile Ile Gly Glu Ala Ser Thr Leu Glu
 35 40 45
 Leu Asp Gly Leu Leu Pro Asn Thr Glu Tyr Thr Val Thr Val Tyr Ala
 50 55 60
 Met Phe
 65

<210> SEQ ID NO 200

<211> LENGTH: 72

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 200

Leu Ser Asp Leu Leu Tyr Val Thr Ser Met Arg Val Lys Trp Asp Ala
 1 5 10 15
 Val Gly Ala Ser Gly Tyr Leu Ile Leu Tyr Ala Pro Leu Thr Glu Gly
 20 25 30
 Leu Ala Gly Glu Lys Glu Met Ile Gly Glu Thr His Thr Ile Glu Leu
 35 40 45
 Ser Gly Leu Leu Pro Asn Thr Glu Tyr Thr Val Thr Val Tyr Ala Met
 50 55 60
 Phe Gly Ala Ser Asp Val Thr Gly
 65 70

<210> SEQ ID NO 201

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Lama glama

<400> SEQUENCE: 201

Asp Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Arg Thr Gly Ser Thr Tyr
 20 25 30
 Asp Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
 35 40 45
 Ala Ala Ile Asn Trp Asp Ser Ala Arg Thr Tyr Tyr Ala Ser Ser Val

-continued

50	55	60																	
Arg Gly Arg Phe Thr	Ile Ser Arg Asp Asn Ala Lys Lys Thr Val Tyr																		
65	70	75	80																
Leu Gln Met Asn Ser	Leu Lys Pro Glu Asp Thr Ala Val Tyr Thr Cys																		
	85	90	95																
Gly Ala Gly Glu Gly	Gly Thr Trp Asp Ser Trp Gly Gln Gly Thr Gln																		
	100	105	110																
Val Thr Val Ser Ser																			
	115																		

<210> SEQ ID NO 202
 <211> LENGTH: 94
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 202

Val Ser Asp Val Pro Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr																			
1	5	10	15																
Ser Leu Leu Phe Ser Trp Asp Ala Pro Ala Val Thr Val Arg Tyr Tyr																			
	20	25	30																
Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Leu Val Gln Glu Phe																			
	35	40	45																
Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro																			
	50	55	60																
Gly Val Asp Tyr Thr Ile Thr Gly Tyr Ala Val Thr Gly Arg Gly Asp																			
	65	70	75	80															
Ser Pro Ala Ser Ser Lys Pro Ile Ser Ile Asn Tyr Arg Thr																			
	85	90																	

What is claimed is:

1. A non-antibody protein comprising a domain having an immunoglobulin-like fold, said non-antibody protein deriving from a reference protein by having a mutated amino acid sequence wherein said non-antibody protein binds with a Kd

at least as tight as 1 μ M to a compound that is not bound as tightly by said reference protein.

2-43. (canceled)

* * * * *