



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/54, 9/12, 5/10, A01K 67/027, C07K 16/40, A61K 48/00, G01N 33/50, C12Q 1/48		A2	(11) International Publication Number: WO 99/25843
			(43) International Publication Date: 27 May 1999 (27.05.99)
(21) International Application Number: PCT/EP98/06981	(22) International Filing Date: 21 October 1998 (21.10.98)		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(30) Priority Data: 9722320.0 22 October 1997 (22.10.97) GB			
(71) Applicants (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; Mail Drop TPC-8, 10550 North Torrey Pines Road, La Jolla, CA 92037 (US). JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): LUYTEN, Walter, H., M., L. [BE/BE]; Janssen Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse (BE). PARKER, Andrew, E. [GB/GB]; Zeneca Pharmaceuticals, 8AF22 Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). MCGOWAN, Clare [GB/US]; 12728 Via Felino, Del Mar, CA 92014 (US). BLASINA, Alessandra [IT/US]; 12530 Carmel Creek Road, 144, San Diego, CA 92130 (US).			
(74) Agent: VOSSIUS, Volker; Holbeinstrasse 5, D-81679 München (DE).			

(54) Title: HUMAN CHECKPOINT KINASE, HCDS1, COMPOSITIONS AND METHODS

(57) Abstract

The invention provides for a novel human checkpoint kinase gene, hCDS1, translated protein, compositions, methods, and kits.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon			PT	Portugal		
CN	China	KR	Republic of Korea	RO	Romania		
CU	Cuba	KZ	Kazakhstan	RU	Russian Federation		
CZ	Czech Republic	LC	Saint Lucia	SD	Sudan		
DE	Germany	LI	Liechtenstein	SE	Sweden		
DK	Denmark	LK	Sri Lanka	SG	Singapore		
EE	Estonia	LR	Liberia				

HUMAN CHECKPOINT KINASE, HCDS1, COMPOSITIONS AND METHODS

BACKGROUND OF THE INVENTION

The integrity of the genome is of prime importance to a dividing cell. In response to DNA damage, eukaryotic cells rely upon a complex system of checkpoint controls to delay cell-cycle progression. The normal eukaryotic cell-cycle is divided into 4 phases (sequentially G1, S, G2, M) which correlate with distinct cell morphology and biochemical activity, and cells withdrawn from the cell-cycle are said to be in G0, or non-cycling state. When cells within the cell-cycle are actively replicating, duplication of DNA occurs in the S phase, and active division of the cell occurs in M phase. *See generally* Benjamin Lewin, GENES VI (Oxford University Press, Oxford, GB, Chapter 36, 1997). DNA is organized in the eukaryotic cell into successively higher levels of organization that result in the formation of chromosomes. Non-sex chromosomes are normally present in pairs, and during cell division, the DNA of each chromosome replicates resulting in paired chromatids. (*See generally* Benjamin Lewin, GENES VI (Oxford University Press, Oxford, GB, Chapter 5, 1997)).

Checkpoint delays provide time for repair of damaged DNA prior to its replication in S-phase and prior to segregation of chromatids in M-phase (Hartwell and Weinert, 1989, Science, 246: 629-634). In many cases the DNA-damage response pathways cause arrest by inhibiting the activity of the cyclin-dependent kinases (Elledge, 1997, Science, 274: 1664-1671). In human cells the DNA-damage induced G2 delay is largely dependent on inhibitory phosphorylation of Cdc2 (Blasina et al., 1997, Mol. Cell Biol., 8: 1-11; Jin et al., 1996, J. Cell Biol., 134: 963-970), and is therefore likely to result from a change in the activity of the opposing kinases and phosphatases that act on Cdc2. However, evidence that the activity of these enzymes is substantially altered in response to DNA damage is lacking (Poon et al., 1997, Cancer Res., 57: 5168-5178).

Three distinct Cdc25 proteins are expressed in human cells. Cdc25A is specifically required for the G1-S transition (Hoffmann et al., 1994, EMBO J., 13: 4302-4310; Jinno et al., 1994, EMBO J., 13: 1549-1556), whereas Cdc25B and Cdc25C are required for the G2-M transition (Gabrielli et al., 1996, J. Cell Sci., 7: 1081-1093; Galaktionov et al., 1991, Cell, 67: 1181-1194; Millar et al., 1991, Proc. Natl. Acad. Sci. USA, 88: 10500-10504; Nishijima et al., 1997, J. Cell Biol., 138:

1105-1116). The exact contribution of Cdc25B and Cdc25C to M-phase progression is not known.

Much of our current knowledge about checkpoint control has been obtained from studies using budding (*Saccharomyces cerevisiae*) and fission 5 (*Schizosaccharomyces pombe*) yeast. A number of reviews of our current understanding of cell cycle checkpoints in yeast and higher eukaryotes have recently been published (Hartwell & Kastan, 1994, *Science*, 266: 1821-1828; Murray, 1994, *Current Biology*, 6: 872-876; Elledge, 1996, *Science*, 274: 1664-1672; Kaufmann & Paules, 1996, *FASEB J.*, 10: 238-247). In the fission yeast six gene products, *rad1*⁺, 10 *rad3*⁺, *rad9*⁺, *rad17*⁺, *rad26*⁺, and *hus1*⁺ have been identified as components of both the DNA-damage dependent and DNA-replication dependent checkpoint pathways. In addition *cds1*⁺ has been identified as being required for the DNA-replication dependent checkpoint and *rad27*⁺/*chk1*⁺ has been identified as required for the DNA-damage dependent checkpoint in yeast.

15 Several of these genes have structural homologues in the budding yeast and further conservation across eukaryotes has recently been suggested with the cloning of two human homologues of *S. pombe rad3*⁺: ATM (ataxia telangiectasia mutated) (Savitsky *et al.*, 1995, *Science*, 268: 1749-1753) and ATR (ataxia telangiectasia and *rad3*⁺ related)(Bentley *et al.*, 1996, *EMBO J.*, 15: 6641-6651; Cimprich *et al.*, 1996, *Proc. Natl. Acad. Sci. USA*, 93: 2850-2855) and of a human homologue of *S. pombe rad9*⁺ (Lieberman *et al.*, 1996, *Proc. Natl. Acad. Sci. USA*, 93: 13890-13885).

20 While much is known about yeast checkpoint proteins and genes, this knowledge is not fully predictive of the existence of corresponding human genes or proteins, or their effector role in human cell-cycle control and regulation.

25 In order to develop new and more effective treatments and therapeutics for the amelioration of the effects of cancer, it is important to identify and characterize human checkpoint proteins and to identify mediators of their activity.

SUMMARY OF THE INVENTION

30 The present invention is directed to the discovery of a novel human checkpoint kinase gene hCDS1, protein and constructs and methods for the production and use of hCDS1.

In particular, the present invention encompasses a nucleic acid sequence which encodes for hCDS1, consisting of the nucleic acid sequence of SEQ ID NO.: 1. In

particular, the invention encompasses the nucleic acid sequence from position 66 to 1695 of the nucleic acid sequence of SEQ ID NO.: 1, which translates into the hCDS1 protein. The present invention also encompasses nucleic acid constructs, vectors, plasmids, cosmids and the like which contain the nucleic acid sequence of SEQ ID 5 NO.: 1. In particular, the present invention provides for nucleic acid vector constructs which contain the nucleic acid sequence of SEQ ID NO.: 1 and are capable of expressing protein from this nucleic acid sequence. The present invention encompasses nucleic acid vectors that are suitable for the transformation of host cells, whether eukaryotic or prokaryotic, suitable for incorporation into viral vectors, or suitable for 10 *in vitro* protein expression. The present invention further embodies the nucleic acid sequence of SEQ ID NO.: 1 in tandem with, or otherwise in conjunction with additional nucleic acids for the generation of fusion protein products containing at least the functional segment of the protein encoded for by the nucleic acid of SEQ ID NO.: 1. The present invention also encompasses the nucleic acid of SEQ ID NO.: 1 adapted for use as a naked DNA transformant for incorporation and expression in 15 target cells. The present invention also provides for anti-sense DNA molecule formulations which are the complement to nucleic acid sequence of SEQ ID NO.: 1, and fragments thereof, whether complementary to contiguous or discontinuous portions of the nucleic acid sequence of SEQ ID NO.: 1. The present invention also 20 provides for compositions incorporating modified nucleotides or backbone components which encode for the nucleic acid sequence of SEQ ID NO.: 1, its complement, or fragments thereof. Such modified nucleotides and nucleic acids are known in the art (see for example Verma *et al.*, Ann. Rev. Biochem. 67: 99-134 (1998)). Thus the present invention encompasses modified nucleic acids which incorporate, for example, 25 internucleotide linkage modification, base modifications, sugar modification, nonradioactive labels, nucleic acid cross-linking, and altered backbones including PNAs (polypeptide nucleic acids).

The present invention provides for the novel human checkpoint kinase protein hCDS1, which consists of the amino acid sequence of SEQ ID NO.: 2. The invention 30 encompasses hCDS1 protein produced by recombinant DNA technology and expressed *in vivo* or *in vitro*. The invention thus encompasses hCDS1 protein produced by transformed host cells in small-scale or large-scale production. The invention encompasses complete hCDS1 protein, in either glycosylated or unglycosylated forms,

produced by either eukaryotic or prokaryotic cellr. The present invention provides for hCDS1 protein expressed from mammalian, insect, plant, bacterial, fungal, or any other suitable host cell. The present invention encompasses hCDS1 protein that is produced as a fusion protein product, conjugated to a solid support, or hCDS1 protein
5 which is labeled with any chemical, radioactive, fluorescent, chemiluminescent or otherwise detectable marker. The present invention also provides for hCDS1 protein isolated from natural sources and enriched in purity over that found in nature. The present invention also provides for pharmaceutical formulations of hCDS1 protein and formulations of the hCDS1 protein in pharmaceutically acceptable carriers or
10 excipients.

The present invention encompasses any nucleic acid sequence which would encode for the amino acid sequence of SEQ ID NO.: 2, and the embodiments of these nucleic acid sequences as described for SEQ ID NO.: 1, as the nucleic acid code for generating any nucleic acid sequence which will encode for a protein having the
15 amino acid sequence of SEQ ID NO.: 2 is predictable to one of skill in the art.

The present invention encompasses antibodies which bind specifically to the hCDS1 protein, either polyclonal or monoclonal, as generated by the immunization of a mammal with protein having the amino acid sequence of SEQ ID NO.: 2, or fragments thereof.

20 The present invention also encompasses equivalent proteins where substitutions of amino acids in the sequence of SEQ ID NO.: 2 that are reasonably predictable as being equivalent, and the embodiments thereof as described for SEQ ID NO.: 2. For example, non-polar (hydrophobic side-chain) amino acids alanine, valine, leucine, isoleucine, proline, phenylalanine, tryptophan, methionine; uncharged polar amino acids glycine, serine, threonine, cysteine, tyrosine, asparagine, glutamine; charged polar amino acids aspartic acid, glutamic acid; basic amino acids lysine, arginine, and histidine are understood by those in the art to have functionally predictable effects when substituted. Thus the present invention also encompasses equivalent nucleic acids which encode for such equivalent proteins and the embodiments thereof as
25 described for SEQ ID NO.: 1.

30 The invention also provides for methods of generating hCDS1 protein, by using recombinant DNA technology and the appropriate nucleic acid encoding for hCDS1 protein, fusion protein, or fragments thereof. The invention provides for

incorporating an appropriate nucleic acid sequence into a suitable expression vector, the incorporation of any suitable control elements such as promoter, enhancer, either inducible or constitutively expressed. The invention provides for the use of expression vectors with or without at least one additional selectable marker or expressible protein.

5 The invention provides for methods wherein a suitably constructed expression vector is transformed or otherwise introduced into a suitable host cell, and protein is expressed by such a host cell. Thus the present invention also provides for the transformed host cells, which are capable of producing hCDS1 protein, fusion protein, or fragments thereof.

10 The discovery that hCDS1 acts in coordination with Cdc25 in the DNA damage checkpoint allows for the use of the compounds of the invention in methods for therapeutic treatment of diseases which involve abnormal DNA damage checkpoint function. The present invention further provides for the use of the compounds of the present invention as therapeutics for the treatment of cancer. In particular, the present 15 invention allows for the specific modification of the hCDS1-Cdc25 DNA damage checkpoint in cells.

The present invention also encompasses methods for screening test compounds for efficacy in effecting the hCDS1 mediated checkpoint function of eukaryotic cells, said method comprising contacting a test compound to eukaryotic cells, and detecting 20 any change in hCDS1 expression or function. Thus the invention further encompasses the method of screening wherein said detection of change in hCDS1 expression or function is accomplished by assaying for hCDS1 mRNA production, or by assaying for hCDS1 protein expression. In particular, the present invention allows for the screening of candidate substances for efficacy in modifying the DNA damage 25 checkpoint by screening for any change in Cdc25 phosphorylation, or kinase activity. The compounds or substances identified by the assays of the invention, or compounds corresponding to such compounds or substances, can be used for the manufacture of pharmaceutical therapeutics.

Thus, in one embodiment the present invention provides for pharmaceutical 30 compositions which include the hCDS1 protein, hCDS1 nucleic acid, hCDS1 anti-sense nucleic acids. In another embodiment, the present invention provides for compounds or substances identified as suitable for use as a therapeutic by the assays of the invention, in pharmaceutical formulations. These pharmaceutical compositions

can further include chemotherapeutic agents for the use in treating cancer, or be administered in a regimen coordinated with the administration of other anti-cancer therapies. The present invention, in one embodiment thus encompasses methods for combined chemotherapy using the hCDS1 derived pharmaceuticals independently, and 5 in combination with other chemotherapeutic agents, and in a second embodiment as admixtures with other anti-cancer therapeutics for single dose administration.

According to one aspect of the present invention, there is provided a nucleic acid encoding hCDS1 protein having the amino acid sequence illustrated in Figure 2 (SEQ ID NO.: 2), or encoding a functional equivalent fragment, or bioprecursor of 10 said protein. Preferably, the nucleic acid may be a DNA molecule such as a genomic DNA molecule and even more preferably a cDNA molecule, however it may also be RNA.

In a preferred embodiment, a nucleic acid encoding hCDS1 protein comprises the nucleic acid sequence represented by position 66 to 1694 of the sequence 15 illustrated in Figure 1 (SEQ ID NO.: 1), the complement thereof, or a nucleic acid sequence capable of hybridizing to either under high stringency conditions.

The nucleic acid sequences defined herein may, advantageously, be capable of hybridizing under low stringency conditions to nucleic acid sequences derived from family members to identify homologs therefrom or alternatively to identify nucleic 20 acid sequences from other species.

As would be well known to those skilled in the art due to the degeneracy of the genetic code the nucleic acid sequences according to the invention may include substitutions therein yet which still encode the same amino acid sequence.

Advantageously, the nucleic acids according to the invention may be 25 incorporated into an expression vector and may be subsequently used to transform, transfect or infect a suitable host cell. In such an expression vector the nucleic acid according to the invention is operably linked to a control sequence, such as a suitable promoter or the like, ensuring expression of the proteins according to the invention in a suitable host cell. The expression vector may, advantageously be a plasmid, cosmid, virus or other suitable vector. The expression vector and the host cell transfected, 30 transformed or infected with the vector also form part of the present invention. Preferably, the host cell is a eukaryotic cell or a bacterial cell and even more preferably a mammalian cell or insect cell. Mammalian host cells are particularly

advantageous because they provide the necessary post-translational modifications to the expressed proteins according to the invention, such as glycosylation or the like, which modifications confer optimal biological activity on said proteins, which when isolated may advantageously be used in diagnostic kits or the like.

5 The expression vector including said nucleic acid according to the invention may advantageously be used *in vivo*, such as in, for example, gene therapy.

According to a further aspect of the invention there is also provided a transgenic cell, tissue or organism comprising a transgene capable of expressing hCDS1 protein, which protein comprises the amino acid sequence illustrated in Figure 10 10 2 (SEQ ID NO.: 2), or the amino acid sequence of a functional equivalent or bioprecursor or fragment therefor. The term "transgene capable of expression" as used herein means a suitable nucleic acid sequence which leads to expression of hCDS1 or proteins, having the same function and/or activity. The transgene may include, for example, genomic nucleic acid isolated from human cells or synthetic 15 nucleic acid, including DNA integrated into the genome or in an extrachromosomal state. Preferably, the transgene comprises the nucleic acid sequence encoding the proteins according to the invention as described herein, or a functional fragment of said nucleic acid. A functional fragment of said nucleic acid should be taken to mean a fragment of the gene comprising said nucleic acid coding for the proteins according 20 to the invention or a functional equivalent, derivative or a non-functional derivative such as a dominant negative mutant, or bioprecursor of said proteins. For example, it would be readily apparent to persons skilled in the art that nucleotide substitutions or deletions may be used using routine techniques, which do not affect the protein sequence encoded by said nucleic acid, or which encode a functional protein according 25 to the invention.

The hCDS1 protein expressed by said transgenic cell, tissue or organism or a functional equivalent or bioprecursor of said protein also form part of the present invention.

Further provided by the present invention is an antisense molecule which is 30 capable of hybridizing to the nucleic acid according to the invention. Advantageously, the antisense molecule according to the invention may be used as a medicament, or in the preparation of a medicament for the treatment of cancer and other proliferative diseases.

The present invention also advantageously provides nucleic acid sequences of at least approximately 15 nucleotides of a nucleic acid according to the invention and preferably from 15 to 50 nucleotides. These sequences may advantageously be used as probes or primers to initiate replication, or the like. Such nucleic acid sequences 5 may be produced according to techniques well known in the art, such as by recombinant or synthetic means. They may also be used in diagnostic kits or the like for detecting the presence of a nucleic acid according to the invention. These tests generally comprise contacting the probe with the sample under hybridizing conditions and detecting for the presence of any duplex or triplex formation between the probe 10 and any nucleic acid in the sample.

Advantageously, the nucleic acid sequences, according to the invention may be produced using such recombinant or synthetic means, such as for example using PCR cloning mechanisms which generally involve making a pair of primers, which may be from approximately 15 to 50 nucleotides to a region of the gene which is desired to 15 be cloned, bringing the primers into contact with mRNA, cDNA, or genomic DNA from a human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region (and where necessary first performing a reverse transcription step), isolating the amplified region or fragment and recovering the amplified DNA. Generally, such techniques as defined herein are well known in 20 the art, such as described in Sambrook *et. al.*, (Molecular Cloning; a Laboratory Manual, 1989). Advantageously, human allelic variants of the nucleic acid according to the invention may be obtained by for example, probing genomic DNA libraries from a range of individuals for example from different populations, and other genotyping techniques. Furthermore, nucleic acids and probes according to the 25 invention may be used to sequence genomic DNA from patients, using techniques well known in the art, for example, the Sanger dideoxy chain termination method, which may advantageously ascertain any predisposition of a patient to certain proliferative disorders.

Further provided by the present invention are isolated proteins having the 30 amino acid sequences as illustrated in Figure 2 (SEQ ID NO.: 2) or the amino acid sequence of a functional equivalent functional fragment or bioprecursor of said protein in addition to antibodies, monoclonal or polyclonal capable of binding to the amino acid sequences of these proteins or fragments thereof. As would be well known to

those skilled in the art, the proteins according to the invention may comprise conservative substitutions, deletions or insertions wherein the protein comprises different amino acids than those disclosed in Figure 2, yet which substitutions, deletions or insertions do not affect the activity of the proteins according to the 5 invention or their ability to interact in the human cell cycle checkpoint pathway.

Preferred fragments include those comprising an epitope of the proteins according to the invention. The epitopes may be determined using, for example, peptide scanning techniques as described in Geysen *et. al.*, Mol. Immunol., 23; 709-715 (1986).

10 The antibodies according to the invention may be produced according to techniques which are known to those skilled in the art. Monoclonal antibodies may be prepared using conventional hybridoma technology as described in Kohler F and Milstein C (1985), Nature 256, 495-497. Polyclonal antibodies may also be prepared using conventional technology well known to those skilled in the art, and which 15 comprises inoculating a host animal, such as a mouse, with a protein or epitope according to the invention and recovering the immune serum. The present invention also includes fragments of whole antibodies which maintain their binding activity, such as for example, Fv, F(ab') and F(ab')₂ fragments as well as single chain antibodies.

20 Advantageously, the nucleic acid and/or the proteins according to the invention may be included in a pharmaceutical composition together with a pharmaceutically acceptable carrier, diluent or excipient therefor. The pharmaceutical composition containing said nucleic acids according to the invention may, for example, be used in gene therapy. Such nucleic acids, according to the invention, may be administered 25 naked, or packaged in protein capsules, lipid capsules, liposomes, membrane based capsules, virus protein, whole virus, cell vectors, bacterial cell hosts, altered mammalian cell hosts, or such suitable means for administration.

30 There is further provided by the present invention a method for detecting for the presence or absence of a nucleic acid according to the invention, in a biological sample, which method comprises, a) bringing said sample into contact with a probe comprising a nucleic acid or probe according to the invention under hybridizing conditions, and b) detecting for the presence of hybridization, for example, by the presence of any duplex or triplex formation between said probe and any nucleic acid

present in said sample. Proteins according to the invention may also be detected by a) contacting said sample with an antibody to an epitope of a protein according to the invention under conditions which allow for the formation of an antibody-antigen complex, b) monitoring for the presence of any antigen-antibody complex.

5 Kits for detecting said nucleic acids and proteins are also provided by the present invention. A kit for detecting for the presence of a nucleic acid according to the invention in a biological sample may comprise (a) means for contacting the sample with a probe comprising a nucleic acid or a probe according to the invention and means for detecting for the presence of any duplex or triplex formation between 10 said probe and any nucleic acid present in the sample.

Likewise, a kit for detecting for the presence of a protein according to the invention in a biological sample may comprise (a) means for contacting said sample with an antibody to an epitope of a protein according to the invention under conditions which allow for the formation of an antibody - protein complex, and means 15 for monitoring said sample for the presence of any protein - antibody complex.

A further aspect of the present invention provides a method of determining whether a compound is an inhibitor or an activator of expression or activity of the proteins of the human cell cycle checkpoint pathway which method comprises contacting a cell expressing the proteins in said pathway with said compound and 20 comparing the level of expression of any of the proteins of the checkpoint pathway of said cell against a cell which has not been contacted with said compound. Any compounds identified may then advantageously be used as a medicament or in the preparation of a medicament for treating cancer or proliferative disorders.

Alternatively, the compounds may be included in a pharmaceutical composition 25 together with a pharmaceutically acceptable carrier, diluent or excipient therefor. Advantageously, any compounds identified as an inhibitor of the cell checkpoint pathway may be included in a pharmaceutical composition according to the invention together with a cytotoxic agent, such as a DNA damaging chemotherapeutic agent, and a pharmaceutically acceptable carrier diluent or excipient therefor. Thus, the human 30 cell cycle checkpoint inhibitor may enhance the chemotherapeutic effect of cytotoxic agents used in, for example, anti-cancer therapy.

There is also provided by the present invention a method for screening candidate substances for anti-cancer therapy, which method comprises a) providing a

protein according to the present invention exhibiting kinase activity together with a substrate for said protein under conditions such that the kinase will act upon the substrate, b) bringing the protein and substrate into contact with a candidate substance, c) measuring the degree of any increase or decrease in the kinase activity of the 5 protein, d) selecting a candidate substance which provides a decrease or increase in activity. Such a candidate substance may also be used as a medicament, or in the preparation of a medicament for the treatment of cancer or other such proliferative cell disorders.

The present invention also comprises a method of identifying other proteins 10 active in the cell checkpoint pathway, which method comprises a) contacting a cell extract with an antibody to an epitope of a protein according to the invention, under appropriate binding conditions, b) identifying any antibody-protein complex and c) analyzing the complex to identify any protein bound to the antibody or protein which is other than the protein according to the invention.

15 Another method for identifying proteins involved in the cell checkpoint pathway utilizes a two-hybrid system developed in yeast by Chien *et. al.*, *supra* (1991). This technique is based on functional *in vivo* reconstitution of a transcription factor which activates a reporter gene. More particularly the technique comprises providing an appropriate host cell with a DNA construct comprising a reporter gene 20 under the control of a promoter regulated by a transcription factor having a DNA binding domain and an activating domain, expressing in the host cell a first hybrid DNA sequence encoding a first fusion of a fragment or all of a nucleic acid sequence according to the invention and either said DNA binding domain or the activating domain of the transcription factor, expressing in the host cell at least one second hybrid DNA sequence encoding putative binding proteins to be investigated together 25 with the DNA binding domain or activating domain of the transcription factor which is not incorporated in the first fusion; detecting any binding of the protein being investigated with a protein according to the invention by detecting for the production of any reporter gene product in the host cell; optionally isolating second hybrid DNA 30 sequences encoding the binding protein. In one embodiment of this aspect of the invention the method may comprise:

(a) constructing at least two nucleotide vectors, the first of which comprises a nucleotide segment encoding for a DNA binding domain of GAL4 protein operably

linked to a nucleic acid sequence encoding a protein according to the present invention, the second vector comprising a nucleotide sequence encoding a protein binding domain of GAL4 operably linked to a nucleotide sequence encoding a protein to be tested,

5 (b) co-transforming each of said vectors into a yeast cell being deficient for transcription of genes encoding galactose metabolizing proteins, wherein interaction between said test protein and the protein according to the invention leads to transcription of galactose metabolic genes.

BRIEF DESCRIPTION OF THE DRAWINGS

10 The invention may be more clearly understood from the following examples which are given by way of example only, with reference to the accompanying drawings, wherein:

15 FIGURE 1 illustrates the nucleotide sequence of hCDS1 (SEQ ID NO.: 1) wherein residues 66-1694 is the coding region and 3' and 5' untranslated regions(UTRs). The initiation and termination codons are shown in bold,

FIGURE 2 illustrates the deduced amino acid sequence of hCDS1 (SEQ ID NO.: 2),

20 FIGURE 3 illustrates the amino acid sequence alignment of hCDS1 and *S. pombe* *cds1* performed using the CLUSTALW alignment program and annotated using the GENEDOC program. Residues shaded black are identical between the two proteins and residues shaded grey are similar.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

25 The present invention encompasses the isolation and characterization of a novel human checkpoint kinase gene and protein which is called hCDS1. The hCDS1 gene and protein show some similarity to a homologous gene and protein found in *S. pombe*.

30 The *S. pombe* *cds1*⁺ gene was identified by its ability to complement a DNA polymerase α mutant (Murakami & Okayama, 1995, *Nature*, 374: 817-819). *S. pombe* *cds1* was also able to suppress the hydroxyurea sensitivity (DNA replication-dependent checkpoint) of *rad1*, *rad3* and *rad9* mutant *S. pombe* strains but not the UV sensitivity (DNA damage-dependent checkpoint). This shows that *S. pombe* *cds1* executes its checkpoint function during DNA synthesis.

S. pombe *cds1* is a putative protein kinase that is 70% similar to the *S. cerevisiae* checkpoint gene *RAD53*. In *S. cerevisiae* the DNA damage- and DNA replication-dependent checkpoints are genetically separate at the level of detection of DNA lesions. The two pathways then converge on the *Rad53* protein kinase which 5 potentially acts as an amplifier in the signal transduction pathway. This appears not to be the case in *S. pombe* where the same proteins are involved in detection of all types of lesion but the transduction of the signal follows separate pathways involving different protein kinases; *S. pombe* *cds1* for the DNA replication-dependent checkpoint and *Chk1/Rad27* for the DNA damage-dependent pathway. It has been suggested that 10 S-phase-specific activation of *cds1* kinase may define a subpathway of the checkpoint response in *S. pombe* (Lindsay et al., 1998, *Genes and Development*, 12: 382-395).

S. pombe *cds1* may act via an interaction with DNA polymerase α to monitor the progress of DNA replication or the integrity of replication complexes. There is evidence in Drosophila for a kinase of the appropriate molecular weight associating 15 with DNA polymerase α (Peck et al., 1993, *B.B.R.C.*, 190: 325-331). Alternatively it may act via phosphorylation of *p107^{wee1}* in a manner analogous to *Chk1* ultimately affecting the activity of the G1/S phase cyclin dependent kinases.

Many of the methods and materials for carrying out the basic molecular biology manipulations as described in the examples below are known in the art, and 20 can be found in such references as Sambrook et al., *Molecular Cloning*, 2nd edition, Cold Spring Harbor Laboratory Press (1989); Berger et al., *Guide to Molecular Cloning Techniques, Methods in Enzymology*, Vol. 152, Academic Press, Inc., (1987); Davis et al., *Basic Methods in Molecular Biology*, Elsevier Science Publishing Co., Inc. (1986); Ausubel et al., *Short Protocols in Molecular Biology*, 2nd ed., John Wiley & Sons, (1992); Goeddel *Gene Expression Technology, Methods in Enzymology*, Vol. 25 185, Academic Press, Inc., (1991); Guthrie et al., *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Vol. 194, Academic Press, Inc., (1991); McPherson et al., *PCR Volume 1*, Oxford University Press, (1991).

The invention in its several aspects can be more readily understood by 30 reviewing the following examples.

Example 1 Isolation of hCDS1

Isolation of hCDS1 began with a search for sequences similar to *S. pombe* *cds1*+ using the TBLASTN program. A human expressed sequence tag (EST No. 864164) was identified in the proprietary LifeSeq® database (Incyte Pharmaceuticals Inc., Palo Alto, CA, USA). Sequence analysis of the 1.3 kb insert revealed an incomplete open reading frame which was similar to *S. pombe* *cds1*. Approximately 650 nucleotides of novel 5' DNA sequence was obtained by 5'RACE (rapid amplification of cDNA ends) using a Marathon Ready human placental cDNA (Clontech), following the manufacturer's instructions.

10 Briefly, the two hCDS1 gene specific primers used for nested PCR (Polymerase chain reaction) reactions were GSP3 5'-
TTTGCTGATGATCTTATGGCTAC-3' (SEQ ID NO.: 3) and GSP4 5'-
CACAGGCACCACTCCAAGAGTTT-3' (SEQ ID NO.: 4). Subsequently, a
complete ORF for hCDS1 was amplified from a human SK-N-MC neuroblastoma
15 cDNA library using the PCR primers
5'-GGGCTCGAGAGCAGCGATGTCTCGGGAGTCGGATGT-3' (SEQ ID NO.: 5)
and
5'-GGCGGATCCTCGAGTCACAACACAGCAGCACACAC-3' (SEQ ID NO.: 6).
The amplification product was then cloned into pCR2.1 vector (Invitrogen) and the
20 DNA sequence determined.

25 The nucleic acid sequence of hCDS1 was found to show 47.8% identity to the *S. pombe* *cds1*+ at the DNA level. Termination codons were present in all three reading frames in the 120 nucleotides immediately 5' to the putative hCDS1 initiation codon, indicating that the complete coding region has been isolated. Parts of the sequence is also found to match partial sequences found in the NCBI databases, EST AA285249, genomic sequence H55451, and the 54 base pair fragment H55698.

30 The identified human gene and vectors encoding the hCDS1 nucleic acid sequence were deposited as plasmid HCDS1 ORF/pCR-Blunt deposited under Accession No. LMBP 3708; plasmid HCDS1 5'RACE fragment/pGEM-Easy deposited under Accession No. LMBP 3710; and plasmid HCDS1 3'fragment Incyte clone 864164/pSPORT deposited under Accession No. LMBP 3709 with the Belgian Co-ordinated Collections of Micro-organisms (BCCM) at Laboratorium Voor Moleculaire

Biologics-Plasmidencollecte (LMBP) 35, B-9000 Gent, Belgium, in accordance with the provisions of the Budapest Treaty, 28 April 1997.

The tissue expression profile of hCDS1 was examined on multiple tissue Northern blots (Clontech) and a cancer cell line Northern blot (Clontech), which were 5 probed with the hCDS1 ORF. A single transcript of approximately 2.1 kb was observed. Expression was undetectable by conventional Northern blot hybridization conditions in all normal human tissues examined. However, expression was found to be greatly elevated in all of the cancer cell lines examined.

The hCDS1 gene was localized to chromosome 22q11.2-q12, as determined 10 using the complete ORF as a probe for FISH (Fluorescent *in situ* Hybridization) analysis. The hybridization efficiency was approximately 62%, and no other loci were detected under the conditions used.

Briefly, lymphocytes isolated from human blood were cultured in α -minimal 15 essential media (MEM) supplemented with 10% fetal calf serum and phytohaemagglutinin (PHA) at 37°C for 68-72 hours. The lymphocyte cultures were treated with BrdU (0.18 mg/ml, Sigma) to synchronize the cell population. The synchronized cells were washed three times with serum-free medium to release the block and re-cultured at 37°C for 6 hours in α -MEM with thymidine (2.5 μ g/ml Sigma). Cells were harvested and slides were prepared using standard procedures 20 including hypotonic treatment, fixation and air-drying. DNA fragments containing the hCDS1 complete ORF were gel purified and biotinylated with dATP using the BRL BioNick labeling kit (15°C, 1 hour) (Heng et al., 1992, Proc. Natl. Acad. Sci. USA. 89: 9509-9513).

Slides were then baked at 55°C for 1 hour, and after RNase treatment, the 25 slides were denatured in 70% formamide in 2x SSC for 2 minutes at 70°C followed by dehydration with ethanol. Probes were denatured at 75°C for 5 minutes in a hybridization mix consisting of 50% formamide and 10% dextran sulphate. Probes were loaded on the denatured chromosomal slides. After overnight hybridization, slides were washed and detected. FISH signals and the DAPI-banding pattern were 30 recorded separately by taking photographs, and the assignment of the FISH mapping data with chromosomal bands was achieved by superimposing FISH signals with DAPI banded chromosomes (Heng & Tsui, 1994, Methods in Mol. Biol., 33: 35-49).

Example 2 Characterization of hCDS1 protein

The hCDS1 nucleic acid sequence cDNA predicts a translation product of 543 amino acids with an approximate molecular weight of 61kDa. This is close to the apparent molecular weight of endogenous Cds1 protein in HeLa cells. The predicted 5 hCDS1 protein, is 28% identical to the cds1 protein of *S. pombe*, 28% identical to RAD53 and 27% identical to the DUN1 kinase of *S. cerevisiae*. Sequence alignment of these apparent homologs shows several regions of sequence similarity outside the kinase domain, including conservation of the Fork Head Associated domain (Hoffmann et al., 1995, *Trends Biochem. Sci.*, 20: 347-9). The human protein shows 10 the same overall structure as *S. pombe* CDS1 and *S. cerevisiae* DUN1 in that it lacks the long C-terminal extension found in RAD53. Northern blot analysis with hCDS1 identified a single transcript of about 2.2 kb expressed in testis and in 8 human cancer samples examined.

Briefly, two multiple tissue Northern blots (Clontech) and a Cancer Cell line 15 Northern blot (Clontech) were hybridized with a cDNA probe for hCDS1. The probe corresponds to the complete ORF as described above. The blots were washed at high stringency (0.1 x SSC, 0.1% SDS, 50°C, 2 x 20 min) and exposed using Kodak X-OMAT autoradiography film with intensifying screens at -70°C.

20 Example 3 Cdc25 total activity assay

The possibility that dephosphorylation of Cdc2 is down-regulated in the presence of DNA damage required an assay to allow for the analysis of the total activity of Cdc25. In the presence of EDTA, Cdc2/Cyclin B from asynchronous HeLa cell extracts was found to inactivate spontaneously.

Briefly, cells were lysed in ice-cold lysis buffer (50 mM Tris pH 7.4 25 containing 2 mM magnesium chloride, 1 mM phenylmethylsulphonyl fluoride, and 5 µg/ml leupeptin, pepstatin and aprotinin). Lysates were cleared by centrifugation at 10,000 xg for 10 minutes and the protein concentration of the supernatants determined using the Lowry assay. 10 mM EDTA was added to the supernatants (100 µg in 60 30 µL) and the reaction initiated by incubation at 30°C. At assay intervals the activity of Cdc2/Cyclin B was assayed by measuring the histone-H1 kinase activity present in anti-Cyclin B immune-precipitates (Blasina et al., *supra*). For immunoblots 400 µg of cell lysate was immune-precipitated using anti-Cyclin B antibody, resolved on an 11%

acrylamide-SDS gel. Monoclonal antibody against the PSTAIRE motif of Cdc2 was used to detect the different phospho-forms of Cdc2.

Activation correlates with loss of the inhibited-phosphorylated form of Cdc2, visualized as the slower migrating species on SDS-PAGE gels. Activation was 5 prevented by vanadate, an inhibitor of Cdc25 and other tyrosine phosphatase. Furthermore, immune-depletion with Cdc25C-specific anti-sera greatly reduced activation of Cdc2/Cyclin B. There was no increase in the levels of Cdc2 or Cyclin B protein, phosphorylation by WEE1 and Myt1 was blocked by the presence of 10 mM EDTA. Thus, these result demonstrate that the activation of Cdc2 was the result of 10 dephosphorylation. In lysates of asynchronous HeLa cells, the endogenous Cdc25 phosphatase activity is sufficient to dephosphorylate and activate more than 80% of the available Cyclin B/Cdc2 in 30 minutes. Analysis of lysates of HeLa cells in which the DNA had been damaged by exposure to 10 Gy of τ -irradiation one hour before harvesting showed a significant reduction in the rate of activation of Cdc2, such that 15 less than 25% of the available Cdc2/Cyclin B was activated during the 30 minutes incubation. The amount of Cdc2/Cyclin B in complex was not significantly altered and it was activated to the same extent as control Cdc2/Cyclin B by addition of exogenous GST-Cdc25. Irradiation with 10 Gy led to more than 3-fold reduction in the rate of 20 Cdc2 dephosphorylation in the 10 time courses examined. If the inactivation of Cdc25 measured above is part of the DNA-damage checkpoint response in human cells, then experimental conditions that over-ride the DNA damage checkpoint might be expected to block the radiation-induced inhibition of Cdc25.

Example 4 DNA Damage Checkpoint Effect of hCDS1

25 DNA damage response in a variety of cells is known to require various related kinases which structurally are related to PI-3 kinases. At least one member of the family, DNA-Protein Kinase, has been shown to be sensitive to wortmannin *in vitro* (Hawley et al., 1996, Genes and Dev., 10: 2383-8; Hartley et al., 1995, Cell, 82: 849-856). Thus the possibility that a wortmannin-sensitive kinase acted upstream of the 30 radiation induced delay in M-phase entry was tested (Price et al., 1996, Cancer Research, 56: 246-250). HeLa cells can be arrested in M-phase by nocodazole, irradiation causes cells to delay in G2 prior to the nocodazole-sensitive M-phase block point. Thus, by scoring the mitotic index of cells that are cultured in nocodazole, it is

possible to determine whether entry into mitosis has been delayed. Control cells cultured in the presence of nocodazole for 14 hours contained 60% mitotic cells, the presence of wortmannin had little effect on this number. However, irradiation reduced the number of cells that reach the nocodazole block point to only 10%. In contrast, 5 irradiation in the presence of wortmannin had only a modest effect. These results demonstrate that wortmannin over-rides the DNA damage G2 checkpoint in HeLa cells.

The effects of wortmannin on the radiation-induced inactivation of Cdc25 was then tested. Wortmannin had little effect on the activation of Cdc2/Cyclin B in 10 extracts prepared from unirradiated cultures, however it did greatly diminish the irradiation-induced decrease in Cdc25 activity.

Radiation-induced G2 checkpoint is also over-ridden in cell-lines derived from patients with the genetic disorder ataxia telangiectasia. Ataxia Telangiectasia mutant cells (ATM) are defective in both the G1 and G2 checkpoints following exposure to 15 many, but not all, agents that damage DNA (Canman et al., 1994, Cancer Research, 54: 5054-5058). The failure of AT-deficient cells to delay G1 correlates with a failure to up-regulate p53 (Kastan et al., 1992, Cell, 71: 587-589), and with failure to phosphorylate and activate cAb1 (Baskaran et al., 1997, Nature, 387: 516-519; Shafman et al., 1997, Nature, 387: 520-523). The molecular basis for the failure to 20 delay G2 is unknown. AT-deficient cells show greatly reduced responses to agents that generate chromosomal breaks, such as ionizing τ -rays. Remarkably, AT-deficient cells have near normal responses following the base damage that is generated by irradiation with a UV source (Canman et al., 1994, Cancer Research, 54: 5054-5058; Painter et al., 1980, Proc. Natl. Acad. Sci. USA, 77: 7315-7317; Zampetti-Bosseler et al., 1981, Int. J. Radiat. Biol., 39: 547-558). The effects of UV and τ -irradiation on the Cdc25 25 activity of AT-plus and AT-minus SV40-transformed human fibroblast cell-lines was tested. AT-minus cells respond to UV-irradiation with a robust reduction in the rate at which Cdc2 is dephosphorylated. In contrast, τ -irradiation had only a modest effect on the rate of dephosphorylation of Cdc2. In AT-plus cells the rate of dephosphorylation 30 of Cdc2 was significantly reduced following either ionizing-radiation or UV-radiation. These data indicate that the ATM gene product is required for the efficient inactivation of Cdc25 following τ -irradiation and demonstrate a correlation between inactivation of Cdc25 and delayed entry into M-phase following DNA damage.

Mediators of the checkpoint-dependent inactivation of Cdc25 in human cells are excellent targets for generating therapeutics or therapeutic regimens that will enhance anti-cancer treatment, and reduce side-effects on normal cells.

5 To facilitate biochemical characterization of hCDS1, 6his-hCDS1 was expressed in insect cells, affinity purified and incubated in extracts of HeLa cells in the presence of an ATP-regenerating system. EDTA was added to inhibit kinase in the extract, and the rate of dephosphorylation and activation of Cdc2/CyclinB was monitored.

Briefly, recombinant viruses encoding for 6his-hCDS1, 6his-Chk1, 6his-Cdc2 10 and GST-Cdc25C were generated using the Bac-to-Bac expression system from Gibco/BRL. 6his-fusion proteins were purified following the procedure described in Kumagai et al., (1995), Mol. Biol. Cell, 6: 199-213. GSH-sepharose beads were incubated for 15 minutes in Sf9 extracts; beads were collected by centrifugation and washed three-times with lysis buffer (50 mM Tris pH 8.0, 5 mM EDTA, 150 mM 15 NaCl, 0.1% NP40, 5% glycerol, 0.1% β -mercaptoethanol and protease inhibitors). Beads were washed three-times with kinase assay buffer (50 mM Tris pH 7.4, 10 mM MgCl₂) prior to phosphorylation reactions or three-times with phosphatase assay buffer (50 mM imidazole pH 7.4, 5 mM EDTA and 0.1% β -mercaptoethanol) prior to phosphatase assays.

20 Both 6his-Chk1 and 6his-hCDS1 were found to significantly reduce the activation of Cdc2/Cyclin B in these assays. The reduced activation of Cdc2 was dose dependent and required ATP. Confirmation that Cdc2 was not irreversibly inhibited by 6his-Chk1 or 6his-hCDS1 was shown by the activation that resulted when excess GST-Cdc25C was added after kinase treatment. Thus, both 6his-hCDS1 and 6his-Chk1 25 can mimic the radiation-induced down-regulation of Cdc25 seen in extracts. These experiments used HeLa cell lysates that had been clarified by centrifugation, therefore it is unlikely that changes in sub-cellular locale could account for inactivation of Cdc25 (Peng et al., 1997, Science, 277: 1501-1505).

30 **Example 5 Direct Effect of hCDS1 on Cdc25**

Indirect mechanisms of inhibition of Cdc25 by hCDS1 could not be excluded by the cell lysate assays, therefore, affinity-purified reagents were used to determine direct phosphorylation and inhibition of GST-Cdc25 activity by hCDS1.

GST-Cdc25 was incubated with either 6his-hCDS1, mock beads, or 6his-Chk1 in the presence of γ -32P ATP for 15 minutes at 30°C. Proteins were resolved by SDS-PAGE and visualized by autoradiography. GST-Cdc25 was phosphorylated by 6his-Chk1 and by 6his-hCDS1. Assays were performed to determine if Cdc25 phosphatase activity was effected by this phosphorylation.

GST-Cdc25 was assayed for its ability to activate the histone-H1 kinase activity of Cdc2/Cyclin B immune-precipitates. It was found that phosphorylation of GST-Cdc25 by 6his-hCDS1 inhibited the ability of GST-Cdc25 to activate Cdc2/Cyclin B. Thus, these data demonstrate that 6his-hCDS1 inactivated Cdc25 *in vitro*, and that Cdc25 is inactivated *in vivo* following DNA damage.

Since 6his-Chk1 associates with GST-Cdc25 and has histone-H1 kinase activity *in vitro* (Sanchez et al., 1997, Science, 277: 1497-1501), analysis of Cdc2/Cyclin B kinase activity was obscured. In order to test GST-Chk1 effects, an assay was used in which Cdc2 dephosphorylation was monitored by the disappearance of the slower migrating species of Cdc2 on gel-mobility analysis.

Briefly, phosphorylated Cdc2 was purified from Sf9 cells that had been simultaneously infected with recombinant baculoviruses encoding 6his-Cdc2, 6his-Wee1, 6his-Myt1 and GST-Cyclin B (Parker et al., 1992, Science, 257: 1955-1957). The 6his-Cdc2 complexed to Cyclin B was purified using GSH beads under the conditions for GST-Cdc25 except that 1 mM VO₄ was included in the lysis buffer. Western Blot analysis showed that quadruple infection resulted in phosphorylation of the majority of Cdc2/GST-Cyclin B at one or both inhibitory sites. These phosphatase assays were carried out in the presence of 10 mM EDTA, and the absence of ATP, conditions that eliminate the possibility of 6his-Chk1 phosphorylating Cdc2 or Cyclin B directly. GST-Cdc25 catalyses a reduction in the slower migrating phosphorylated forms of Cdc2. Prior phosphorylation of GST-Cdc25 by 6his-Chk1 leads to a dose-dependent reduction in GST-Cdc25 activity. These data confirm that Chk1 negatively regulated Cdc25 activity (Furnari et al., 1997, Science, 277: 1495-1497; Weinert, 1997, Science, 277: 1450), and extend them by demonstrating that the negative regulation involves inactivation of the phosphatase activity.

Example 6 DNA Damage and Modification of hCDS1

As the previous data had shown that 6his-hCDS1 inactivates Cdc25, and that DNA damage is associated with Cdc25 inactivation, an assay was performed to determine if DNA damage leads to any modification or activation of hCDS1. Antisera raised against 6his-hCDS1 was used in immune-complex kinase assays using HeLa cell lysates. A weak signal corresponding to hCDS1 was detected in the sample from asynchronous HeLa cells; increased phosphorylation of hCDS1 was seen following irradiation.

Briefly, antibodies to hCDS1 were generated by immunizing a rabbit with 6his-hCDS1 purified from Sf9 cells (Harlow et al., Antibodies (Cold Spring Harbor Laboratory Press, NY, 1988). The resulting antisera immune-precipitates an active kinase of the expected molecular weight from Sf9 cells infected with 6his-hCDS1 virus, but not from uninfected Sf9 cells, or from other cells infected with 6his-Chk1 virus.

The results were confirmed as being due to hCDS1 by re-precipitation of the protein band following denaturation in 4% SDS. The *in vitro* phosphorylation is most likely due to autophosphorylation, and the increased signal reflects an increase in activity following irradiation. The increase *in vitro* phosphorylation of p64^{Cds1} suggests that, like RAD53 and DUN1, hCDS1 is modified in response to DNA damage.

The effect of arresting DNA synthesis on phosphorylation of p64^{Cds1} was examined by further assay. The hCDS1 from replication arrested cells behaved exactly like the protein from asynchronous cultures; no significant increase in phosphorylation was seen in response to thymidine or other agents that block DNA replication. The increased phosphorylation of p64^{Cds1} was detected following irradiation of thymidine-arrested cells. The effect of damaging DNA in cells that are predominantly arrested outside S-phase was also tested. Cells were cultured in the presence of nocodazole for 20 hours prior to irradiation. Again, a weak but detectable signal was seen in the unirradiated sample. However, irradiation of nocodazole arrested cells lead to increased phosphorylation.

These findings surprisingly contrast with the results found in yeast, where fission yeast Cds1 has been found to be activated in response to incompletely replicated DNA (Boddy et al., 1998, Science, 280: 909-12; Lindsay et al., 1998, Genes and Dev., 12: 382-95). The results here show a role for human Cds1 in the

DNA damage checkpoint rather than the replication checkpoint as previously found in yeast.

Example 7 Drug Identification

5 The Cdc25 assays described above are suitable for use in the identification of chemical agents that would modify the DNA damage checkpoint mediated by hCDS1 and Cdc25, either by enhanced or inhibited activity. Thus a typical screening assay would use similar conditions as described above, plus addition of a reagent to be tested. Monitoring of the activity of the assay components, i.e. detection of 10 phosphorylation as described above, can be conducted in comparison to control reactions to detect both enhanced and inhibited activity.

Clearly such assays are readily adaptable to mechanical/automated apparatus and detection. With the fundamental elements of the assay reactions being known, the assay is clearly suited for use in conjunction with automated high-throughput 15 low-signal apparatus which may incorporate microscopic slide array, or cell-biochip arrays in conjunction with CCD detection devices and the use of a visible signal triggered by phosphorylation or other reaction to kinase activity.

Example 8 Therapeutic Use

20 The characterization of hCDS1 and the elucidation that the role for human Cds1 is in the DNA damage checkpoint rather than the replication checkpoint as found in yeast, allows for the adaptation of this knowledge to the preparation of pharmaceuticals, and therapeutic methods for acting as an adjunct to chemotherapy of cancer.

25 In particular, pharmaceutical formulations of the present invention incorporating cDNA, RNA, antisense molecules, hCDS1 protein, antibodies against hCDS1 protein, or other therapeutics corresponding to those identified in the assays of the invention, can be administered in conjunction with any suitable chemotherapy agent in order to act as an adjunct to the main action of the chemotherapy agent. For 30 example, the use of anticancer drugs such as antimetabolite, antibiotics, alkylating agents, microtubule inhibitors, steroid hormones and their antagonists, and others, is generally directed against metabolic sites essential to cell replication. While ideally these drugs should intervene only with the cellular processes unique to malignant

cells, the currently available anticancer drugs affect all proliferating cells, both normal and malignant. Thus, current chemotherapy is hampered by a steep dose-response curve for both toxic and therapeutic effects. Therefore, co-administration of the hCDS1-based drugs of the present invention, and drugs identified by the hCDS1 assays of the present invention, with chemotherapeutic agents will allow for enhanced killing of malignant cells.

One mechanism for enhanced killing is effected by disabling the DNA damage checkpoint control of malignant cells, thus making the administration of DNA damaging chemotherapeutic agents more effective. The disabling of the DNA damage control checkpoint can be effected by modifying the hCDS1 response, as demonstrated by the data above.

Thus, the co-administration of novel hCDS1 based therapeutics in combination with any one or more anticancer agent is contemplated by the present invention. For example, normal dosages of the anticancer drugs Cytarabine, Fludarabine, 5-Fluorouracil, 6-Mercaptopurine, Methotrexate, 6-Thioguanine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Idarubicin, Plicamycin, Carmustine, Iomustine, Cyclophosphamide, Ifosfamide, Mechloroethamine, Streptozotocin, Navelbine, Paclitaxel, Vinblastine, Vincristine, Asparaginase, Cisplatin, Carboplatin, Etoposide, Interferons, Procarbazine etc., can be administered with the appropriate amount of hCDS1 based drug so as to a) alter the length of time of administration, b) alter the time between administrations, c) alter the efficacy of the chemotherapeutic agent on malignant cells, or d) alter the side-effects of the chemotherapeutic agent on normal cells. The effects of the co-administration of hCDS1 based drugs can be any one or combination of these effects in addition to others.

Typically, destruction of cancer cells by chemotherapeutic agents follows first-order kinetics, for a log kill effect. Thus, the co-administration of hCDS1-based therapeutics would be designed to enhance the log kill effect. Typically, chemotherapeutic treatment protocols call for a combination of drugs which act at different steps in the metabolic pathway, thus enhancing killing while staying below toxic levels. Thus, the co-administration of hCDS1 based therapeutics would ideally be in combination with such protocols, and improve efficacy thereof.

Ultimately, the most effective therapeutic methods would combine targeted administration of chemotherapeutic drugs and/or MDR (multidrug resistance)

inhibiting agents, with hCDS1 based therapeutics, to specifically target and eliminate malignant cells via the cells' own uncontrolled replication without DNA damage repair, and thus eventual cell death.

The foregoing discussion and examples are intended as illustrative of the present invention and are not to be taken as limiting. Still other variants within the spirit and scope of this invention are possible and will readily present themselves to those of skill in the art.

What is claimed is:

1. A DNA expression vector comprising an isolated DNA molecule having the nucleic acid sequence as shown in Figure 1 (SEQ ID NO.:1).
2. A DNA expression vector comprising an isolated DNA molecule having 5 the nucleic acid sequence represented by residues 66-1694 of the nucleic acid sequence shown in Figure 4 (SEQ ID NO.:1).
3. A host cell transformed, transfected or infected with the vector of claim 1 or 2.
4. A host cell according to claim 3 which cell is a eukaryotic or prokaryotic 10 cell.
5. A host cell according to claim 4 wherein said eukaryotic cell is a mammalian cell.
6. A host cell according to claim 5 wherein said eukaryotic cell is an insect cell.
- 15 7. A transgenic cell, tissue or organism comprising a transgene capable of expressing a HCDS1 protein, having the amino acid sequence of Figure 2 (SEQ ID NO.: 2), or the amino acid sequence of a fragment of said protein, a functional equivalent, of said protein, or a bioprecursor of said protein.
8. A HCDS1 protein, fragment or functional equivalent or bioprecursor 20 thereof, expressed from a cell according to any of claims 1 to 7.
9. An isolated protein having the amino acid sequence of Figure 2 (SEQ ID NO.: 2) or the amino acid sequence of the functional equivalent or bioprecursor of said protein.
10. An antibody which specifically binds to a protein of claims 8 and 9.
11. An antibody which specifically binds to a protein having the amino acid 25 sequence of Figure 2 (SEQ ID NO.: 2).
12. An antibody of claim 10 that is a monoclonal antibody.
13. A protein according to claims 8 or 9, for use in treating cancer or proliferative disease.
- 30 14. A protein according to claims 8 or 9, for use in the preparation of a medicament.
15. The use of a protein according to claims 8 or 9, for the preparation of a medicament.

16. A medicament comprising a protein according to claims 8 or 9.
17. The use of a medicament comprising a protein of claims 8 or 9, for the treatment of cancer or proliferative disease.
18. The method of treating cancer or proliferative disease in a patient, comprising administering a compound according to claims 8 or 9, to a patient in need of treatment.
19. The method of treating cancer or proliferative disease in a patient, comprising administering a medicament comprising a protein according to claims 8 or 9, to a patient in need of treatment.
20. A protein according to claims 10, 11 or 12, for use in treating cancer or proliferative disease.
21. A protein according to claims 10, 11 or 12, for use in the preparation of a medicament.
22. The use of a protein according to claims 10, 11 or 12, for the preparation of a medicament.
23. A medicament comprising a protein according to claims 10, 11 or 12.
24. The use of a medicament comprising a protein of claims 10, 11 or 12, for the treatment of cancer or proliferative disease.
25. The method of treating cancer or proliferative disease in a patient, comprising administering a compound according to claims 10, 11 or 12, to a patient in need of treatment.
26. The method of treating cancer or proliferative disease in a patient, comprising administering a medicament comprising a protein according to claims 10, 11 or 12, to a patient in need of treatment.
27. A method for identifying a compound as an inhibitor or activator of expression of the CDS1 human cell cycle checkpoint pathway protein which method comprises, contacting a cell expressing the proteins in said pathway with said compound and comparing the level of expression of the CDS1 human cell cycle checkpoint pathway protein of said cell against a cell which has not been contacted with said compound.
28. A compound selected by the method of claim 26.
29. A compound identifiable by the method of claim 26, useful for treating cancer or other proliferative diseases.

30. The use of a compound of claim 28 or 29, in the preparation of a medicament for the treatment of cancer or proliferative diseases.

31. A pharmaceutical composition comprising a compound according to claim 28 or 29, together with a pharmaceutically acceptable carrier, diluent or 5 excipient therefor.

32. A pharmaceutical composition comprising a compound of claim 28 or 29 together with a DNA damaging chemotherapeutic agent and a pharmaceutically acceptable carrier, diluent or excipient therefor.

33. Use of a medicament comprising a compound of claim 28 or 29, for the 10 treatment of cancer or proliferative disease.

34. The method of treating cancer or proliferative disease in a patient, comprising administering a compound of claim 28 or 29, to a patient in need of treatment.

35. The method of treating cancer or proliferative disease in a patient, 15 comprising administering a medicament containing a compound of claim 28 or 29, to a patient in need of treatment.

36. A method for screening candidate substances for anti-cancer therapy, which method comprises, providing a protein according to claims 8 or 9, exhibiting kinase activity together with a substrate for said protein under 20 conditions such that the kinase will act upon the substrate, bringing the protein and substrate into contact with a candidate substance, measuring the degree of any increase or decrease in the kinase activity of the protein, selecting a candidate substance which provides a decrease or increase in said kinase activity.

37. A method of increasing susceptibility of cancer cells to chemotherapy 25 and/or radiotherapy, which method comprises administering to a patient a therapeutically effective amount of a candidate substance identifiable by the method of claim 36.

38. A candidate substance selected according to the method of claim 36.

39. The use of a compound of claim 37 or 38, in the preparation of a 30 medicament for the treatment of cancer or proliferative diseases.

40. A pharmaceutical composition comprising a compound according to claim 37 or 38, together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

41. A pharmaceutical composition comprising a compound of claim 37 or 38 together with a DNA damaging chemotherapeutic agent and a pharmaceutically acceptable carrier, diluent or excipient therefor.
- 5 42. Use of a medicament comprising a compound of claim 37 or 38, for the treatment of cancer or proliferative disease.
43. The method of treating cancer or proliferative disease in a patient, comprising administering a compound of claim 37 or 38, to a patient in need of treatment.
- 10 44. The method of treating cancer or proliferative disease in a patient, comprising administering a medicament containing a compound of claim 37 or 38, to a patient in need of treatment.
- 15 45. A method for determining whether a compound modifies the activity of CDS1 human cell cycle checkpoint pathway protein said method comprising the step of combining Cdc25 protein with hCds1 protein in the presence of ATP and a compound to be tested, and detecting Cdc25 phosphatase activity.
46. A compound selected by the method of claim 45.
47. A compound corresponding to a compound selected by the method of claim 45.
- 20 48. A compound identifiable by the method of claim 45, useful for treating cancer or other proliferative diseases.
49. The use of a compound of claim 46, 47 or 48, in the preparation of a medicament for the treatment of cancer or proliferative diseases.
- 25 50. A pharmaceutical composition comprising a compound according to claim 46, 47 or 48, together with a pharmaceutically acceptable carrier, diluent or excipient therefor.
51. A pharmaceutical composition comprising a compound of claim 46, 47 or 48 together with a DNA damaging chemotherapeutic agent and a pharmaceutically acceptable carrier, diluent or excipient therefor.
- 30 52. Use of a medicament comprising a compound of claim 46, 47 or 48, for the treatment of cancer or proliferative disease.

53. The method of treating cancer or proliferative disease in a patient, comprising administering a compound of claim 46, 47 or 48, to a patient in need of treatment.

54. The method of treating cancer or proliferative disease in a patient, comprising administering a medicament containing a compound of claim 46, 47 or 48, to a patient in need of treatment.

55. A method for screening candidate substances suitable for anti-cancer therapy, which method comprises, combining a protein having the amino acid residue sequence shown in Figure 2 (SEQ ID NO.: 2) and Cdc25 protein under 10 phosphorylating conditions, contacting the resulting combination with a candidate substance, determining any change in Cdc25 protein phosphorylation activity, selecting a candidate substance which modulates said phosphorylation activity.

56. A compound selected by the method of claim 55.

57. A compound corresponding to a compound selected by the method of 15 claim 55.

58. A compound identifiable by the method of claim 55, useful for treating cancer or other proliferative diseases.

59. The use of a compound of claim 56, 57 or 58, in the preparation of a medicament for the treatment of cancer or proliferative diseases.

60. A pharmaceutical composition comprising a compound according to 20 claim 56, 57 or 58, together with a pharmaceutically acceptable carrier, diluent or excipient therefor.

61. A pharmaceutical composition comprising a compound of claim 56, 57 or 58 together with a DNA damaging chemotherapeutic agent and a 25 pharmaceutically acceptable carrier, diluent or excipient therefor.

62. Use of a medicament comprising a compound of claim 56, 57 or 58, for the treatment of cancer or proliferative disease.

63. The method of treating cancer or proliferative disease in a patient, comprising administering a compound of claim 56, 57 or 58, to a patient in need 30 of treatment.

64. The method of treating cancer or proliferative disease in a patient, comprising administering a medicament containing a compound of claim 56, 57 or 58, to a patient in need of treatment.

65. A method for modifying DNA damage checkpoint activity of a cell, comprising administering an effective amount of a protein having the amino acid sequence shown in Figure 2 (SEQ ID NO.: 2).

66. A method as in claim 65 wherein said protein is an isolated recombinant protein.

67. A method as in claim 65 wherein said protein is administered in conjunction with at least one other chemotherapy agent.

68. A method as in claim 65 wherein said cell is a cancer cell.

1/3

FIG. 1

HCDS1 cDNA nucleotide sequence

1	actagtgatt	actcacaggg	ctcgagcggc	cggccgggca	ggtcaggtgg	50
51	gctcacgcgg	tcgtgatgtc	tcgggagtcg	gatgttgagg	ctcagcagtc	100
101	tcatggcagc	agtgcctgtt	cacagccccca	tggcagcgtt	acccagtccc	150
151	aaggctcctc	ctcacagtcc	cagggcatat	ccagctcctc	taccagcacg	200
201	atgccaaact	ccagccagtc	ctctcaactcc	agctctggga	cactgagctc	250
251	cttagagaca	gtgtccactc	aggaactcta	ttcttattccct	gaggaccaag	300
301	aacctgagga	ccaagaaccc	gaggagccta	cccctgcccc	ctgggctcga	350
351	ttatgggccc	ttcaggatgg	atttgcaat	cttgaatgtg	tgaatgacaa	400
401	ctactgggtt	gggaggggaca	aaagctgtga	atattgtctt	gatgaaccac	450
451	tgctgaaaag	aacagataaa	tacccaaacat	acagcaagaa	acactttcgg	500
501	attttcaggg	aagtgggtcc	taaaaactct	tacattgcatt	acatagaaga	550
551	tcacagtggc	aatggaaacct	ttgttaatac	agagcttgc	gggaaaggaa	600
601	aacgcctc	tttgaataac	aattctgaaa	ttgcactgtc	actaagcaga	650
651	aataaagttt	ttgtcttttt	tgatctgact	gttagatgatc	agtcagttt	700
701	tcctaaggca	ttaagagatg	aatacatcat	gtcaaaaact	cttggaaagtg	750
751	gtgcctgtgg	agaggtaaag	ctggctttcg	agagggaaac	atgtaaagaaa	800
801	gtagccataa	agatcatcag	caaaaaggaa	tttgctattt	gttcagcaag	850
851	agaggcagac	ccagctctca	atgttggaaac	agaaatagaa	attttggaaaa	900
901	agctaaatca	tccttgcatc	atcaagatta	aaaacttttt	tgatgcagaa	950
951	gattattata	ttgttttggg	attgtggaa	ggggggagagc	tgtttgacaa	1000
1001	agtgggggg	aataaaacgcc	tgaaaagaagc	tacctgcacg	ctcttatttt	1050
1051	accatgtct	cttggctgtg	cagtacccctc	atggaaacgg	tattatacac	1100
1101	cgtacttaa	agccagagaaa	tgttttactg	tcatctcaag	aagaggactg	1150
1151	tcttataaaag	attactgatt	ttgggcactc	caagattttt	ggagagacct	1200
1201	ctctcatgag	aaccttatgt	ggaacccccca	cctacttggc	gcctgaagtt	1250
1251	cttgggtctg	ttgggactgc	tgggtataac	cgtgctgtgg	actgctggag	1300
1301	ttttaggagtt	attctttttt	tctgccttag	tgggtatcca	cctttctctg	1350
1351	agcataggac	tcaagtgtca	ctgaaggatc	agatcaccag	tggaaaatac	1400
1401	aacttcattc	ctgaagtctg	ggcagaagtc	ttagagaaaag	ctctggacct	1450
1451	tgtcaagaag	ttgttggtag	tggatccaaa	ggcacgtttt	acgacagaag	1500
1501	aaggcctaag	acacccgtgg	cttcaggatg	aagacatgaa	gagaaaagttt	1550
1551	caagatcttc	tgtctgagga	aaatgaatcc	acagctctac	cccaggttct	1600
1601	agcccagcct	tctactatgc	gaaaggccgc	ccgtgaagggg	gaagccgagg	1650
1651	gtgccgagac	cacaaagcgc	ccagctgtgt	gtgctgtgt	gttgtgaact	1700
1701	ccgtggtttg	aacacgaaaag	aaatgtacct	tctttcaactc	tgtcatctt	1750
1751	cttttctttt	agtctgtttt	tttatagtttt	gtattttaat	tatgggaaata	1800
1801	attgtttttt	cacagtcact	gatgtacaat	taaaaacctg	atggAACCTG	1850
1851	aaaaaaa					

2/3

FIG. 2

Predicted hCDS1 Amino acid sequence

1 MSRESDVEAQ QSHGSSACSQ PHGSVTQSQG SSSQSQGISS 40
41 SSTSTMPNSS QSSHSSSGTL SSLETVSTQE LYSIPEDQEP 80
81 EDQEPEEPTP APWARLWALQ DGFANLECVN DNYWFGRDKS 120
121 CEYCFDEPLL KRTDKYRTYS KKHFRIFREV GPKNSYIAYI 160
161 EDHSGNGTFV NTELVGKGKR RPLNNNSEIA LSLSRNKVFV 200
201 FFDLTVDDQS VYPKALRDEY IMSKTLGSGA CGEVKLAER 240
241 KTCKKVAIKI ISKRKFAIGS AREADPALNV ETEIEILKKL 280
281 NHPCIIKIKN FFDAEDYYIV LELMEGGELF DKVVGNKRLK 320
321 EATCKLYFYQ MLLAVQYLHE NGIIHRDLKP ENVLLSSQEE 360
361 DCLIKITDFG HSKILGETSL MRTLCGTPTY LAPEVLVSVG 400
401 TAGYNRAVDC WSLGVILFIC LSGYPPFSEH RTQVSLKDQI 440
441 TSGKYNFIPE VVVAEVSEKAL DLVKKLLVVD PKARFTTEEA 480
481 LRHPWLQDED MKRKFQDLLS EENESTALPQ VLAQPSTSRK 520
521 RPREGAEAGA ETTKRPAVCA AVL

3/3

HCDS1	:	MPNSSSSSSGTLSLETSTQELYSIPEPEDQEPEEPTPAPWARIWALQDGFANLECVNDNNT-FGRDKSCEY	:	78
S.p. CDS1	:	ME---EPEEATQAPQEAAPLHVSQNIAKQVNNT-----ENVFMKLVMTRMLDGKTEVPLTLDVHNGFGRHKSCEV	:	70
HCDS1	:	CFDEPLKRTDKYRTYSKKHFRIFREVGPKNSYIAYTEHDHSINGTFVNTELVGKGRPLNNSETAISLSRNK-----	:	152
S.p. CDS1	:	VLNCE---RVSNFHFEIYQGHENDSDE-----EN-----WVFLHDHSINGTFLNFERLAKNSRTTISNGDEIRIGLGVPKDEISF	:	143
HCDS1	:	-VFVFFFDLTVDDQSVYPKALRDEYIMSKTLGSGACGEVKLAFERKTCKVAIKTISKRKFAFGSAREADPALNVEIE	:	230
S.p. CDS1	:	ILKKLNHPCLIKIKNNFFDAED-YYIVLLELMEGGELFDKUVGAKNKRRIKEATCKLYFYQMLAVQYLHENGIITHRLDKPENV	:	221
HCDS1	:	ILKSLHHPGVVQCHEICENDDELFIVMEYVEGGDLMDFFLIANGSIDEQDCKPLLKQLLETLLHLHKQGVITHRDLKPENI	:	308
S.p. CDS1	:	LLSQEEDCLIKITDGFHSKTLGET-SILMRTTICGTPTYLAPEVILVSGT---ACYNRAVDGMSLCVILFICLSGYPPFS	:	383
HCDS1	:	LLSQEEDCLIKITDGFHSKTLGET-SILMRTTICGTPTYLAPEVILVSGT---ACYNRAVDGMSLCVILFICLSGYPPFS	:	376
S.p. CDS1	:	LITN---DFHLLKISDFGLAKVVIHGTCTFLEIFCGTMGYLAPEVLRSKAVNVLDDGYYDDKVDIISLCCVLYVMLTASIPFA	:	376
HCDS1	:	EHRTOVSLKDQITSKCKYNFIPPEVMAEVSEKAIDLIVKLLVVDPK2RFITTEALRHPLQDDEDMKRKFQDLLEEEESTA	:	462
S.p. CDS1	:	SS-SQAKCIELISKAYPIEPLLENEISEGIDLINRMLEINPEKRRISESEALQHPMFYT-----	:	435
HCDS1	:	LPQVLAQPSRSRKPREGEAEGAETTKRPAVCAAVL	:	498
S.p. CDS1	:	-----VSTHEHRTPESSSEHEATEQLNSSS-----	:	460

SEQUENCE LISTING

<110> The Scripps Research Institute et al.
 <120> A Novel Human Checkpoint Kinase, hCDS1, Compositions
 and Methods
 <130> TSRI649.SP8
 <140>
 <141>
 <150> GB 9722320.0
 <151> 1997-10-22
 <160> 6
 <170> PatentIn Ver. 2.0
 <210> 1
 <211> 1858
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 <222> (66)..(1694)
 <400> 1
 actagtatt actcacaggg ctcgagcggc cgccgggca ggtcaggtgg gtcacgcgg 60
 tcgtg atg tct cgg gag tcg gat gtt gag gct cag cag tct cat ggc agc 110
 Met Ser Arg Glu Ser Asp Val Glu Ala Gln Gln Ser His Gly Ser
 1 5 10 15
 agt gcc tgt tca cag ccc cat ggc agc gtt acc cag tcc caa ggc tcc 158
 Ser Ala Cys Ser Gln Pro His Gly Ser Val Thr Gln Ser Gln Gly Ser
 20 25 30
 tcc tca cag tcc cag ggc ata tcc agc tcc tct acc agc acg atg cca 206
 Ser Ser Gln Ser Gln Gly Ile Ser Ser Ser Thr Ser Thr Met Pro
 35 40 45
 aac tcc agc cag tcc tct cac tcc agc tct ggg aca ctg agc tcc tta 254
 Asn Ser Ser Gln Ser Ser His Ser Ser Ser Gly Thr Leu Ser Ser Leu
 50 55 60
 gag aca gtg tcc act cag gaa ctc tat tct att cct gag gac caa gaa 302
 Glu Thr Val Ser Thr Gln Glu Leu Tyr Ser Ile Pro Glu Asp Gln Glu
 65 70 75
 cct gag gac caa gaa cct gag gag cct acc cct gcc ccc tgg gct cga 350
 Pro Glu Asp Gln Glu Pro Glu Glu Pro Thr Pro Ala Pro Trp Ala Arg
 80 85 90 95
 tta tgg gcc ctt cag gat gga ttt gcc aat ctt gaa tgt gtg aat gac 398
 Leu Trp Ala Leu Gln Asp Gly Phe Ala Asn Leu Glu Cys Val Asn Asp
 100 105 110
 aac tac tgg ttt ggg agg gac aaa agc tgt gaa tat tgc ttt gat gaa 446
 Asn Tyr Trp Phe Gly Arg Asp Lys Ser Cys Glu Tyr Cys Phe Asp Glu
 115 120 125
 cca ctg ctg aaa aga aca gat aaa tac cga aca tac agc aag aaa cac 494
 Pro Leu Leu Lys Arg Thr Asp Lys Tyr Arg Thr Tyr Ser Lys Lys His
 130 135 140
 ttt cgg att ttc agg gaa gtg ggt cct aaa aac tct tac att gca tac 542
 Phe Arg Ile Phe Arg Glu Val Gly Pro Lys Asn Ser Tyr Ile Ala Tyr
 145 150 155
 ata gaa gat cac agt ggc aat gga acc ttt gta aat aca gag ctt gta 590
 Ile Glu Asp His Ser Gly Asn Gly Thr Phe Val Asn Thr Glu Leu Val
 160 165 170 175
 ggg aaa gga aaa cgc cgt cct ttg aat aac aat tct gaa att gca ctg 638
 Gly Lys Gly Lys Arg Arg Pro Leu Asn Asn Ser Glu Ile Ala Leu
 180 185 190
 tca cta agc aga aat aaa gtt ttt gtc ttt gat ctg act gta gat 686
 Ser Leu Ser Arg Asn Lys Val Phe Val Phe Asp Leu Thr Val Asp
 195 200 205
 gat cag tca gtt tat cct aag gca tta aga gat gaa tac atc atg tca 734
 Asp Gln Ser Val Tyr Pro Lys Ala Leu Arg Asp Glu Tyr Ile Met Ser
 210 215 220
 aaa act ctt gga agt ggt gcc tgt gga gag gta aag ctg gct ttc gag 782
 Lys Thr Leu Gly Ser Gly Ala Cys Gly Glu Val Lys Leu Ala Phe Glu
 225 230 235
 agg aaa aca tgt aag aaa gta gcc ata aag atc atc agc aaa agg aag 830
 Arg Lys Thr Cys Lys Lys Val Ala Ile Lys Ile Ser Lys Arg Lys

240	245	250	255
ttt gct att ggt tca gca aga gag gca gac cca gct ctc aat gtt gaa			
Phe Ala Ile Gly Ser Ala Arg Glu Ala Asp Pro Ala Leu Asn Val Glu			
260	265	270	
aca gaa ata gaa att ttg aaa aag cta aat cat cct tgc atc atc aag			
Thr Glu Ile Glu Ile Leu Lys Lys Leu Asn His Pro Cys Ile Ile Lys			
275	280	285	
att aaa aac ttt ttt gat gca gaa gat tat tat att gtt ttg gaa ttg			
Ile Lys Asn Phe Asp Ala Glu Asp Tyr Tyr Ile Val Leu Glu Leu			
290	295	300	
atg gaa ggg gga gag ctg ttt gac aaa gtg gtg ggg aat aaa cgc ctg			
Met Glu Gly Gly Glu Leu Phe Asp Lys Val Val Gly Asn Lys Arg Leu			
305	310	315	
aaa gaa gct acc tgc aag ctc tat ttt tac cag atg ctc ttg gct gtg			
Lys Glu Ala Thr Cys Lys Leu Tyr Phe Tyr Gln Met Leu Leu Ala Val			
320	325	330	335
cag tac ctt cat gaa aac ggt att ata cac cgt gac tta aag cca gag			
Gln Tyr Leu His Glu Asn Gly Ile Ile His Arg Asp Leu Lys Pro Glu			
340	345	350	
aat gtt tta ctg tca tct caa gaa gag gac tgt ctt ata aag att act			
Asn Val Leu Leu Ser Ser Gln Glu Asp Cys Leu Ile Lys Ile Thr			
355	360	365	
gat ttt ggg cac tcc aag att ttg gga gag acc tct ctc atg aga acc			
Asp Phe Gly His Ser Lys Ile Leu Gly Glu Thr Ser Leu Met Arg Thr			
370	375	380	
tta tgt gga acc ccc acc tac ttg gcg cct gaa gtt ctt gtt tct gtt			
Leu Cys Gly Thr Pro Thr Tyr Leu Ala Pro Glu Val Leu Val Ser Val			
385	390	395	
400	405	410	415
405	410	415	
att ctt ttt atc tgc ctt agt ggg tat cca cct ttc tct gag cat agg			
Ile Leu Phe Ile Cys Leu Ser Gly Tyr Pro Pro Phe Ser Glu His Arg			
420	425	430	
act caa gtg tca ctg aag gat cag atc acc agt gga aaa tac aac ttc			
Thr Gln Val Ser Leu Lys Asp Gln Ile Thr Ser Gly Lys Tyr Asn Phe			
435	440	445	
440	445	450	
att cct gaa gtc tgg gca gaa gtc tca gag aaa gct ctg gac ctt gtc			
Ile Pro Glu Val Trp Ala Glu Val Ser Glu Lys Ala Leu Asp Leu Val			
450	455	460	
455	460	465	
aag aag ttg ttg gta gtg gat cca aag gca cgt ttt acg aca gaa gaa			
Lys Lys Leu Leu Val Val Asp Pro Lys Ala Arg Phe Thr Thr Glu Glu			
465	470	475	
470	475	480	
gcc tta aga cac ccg tgg ctt cag gat gaa gac atg aag aga aag ttt			
Ala Leu Arg His Pro Trp Leu Gln Asp Glu Asp Met Lys Arg Lys Phe			
480	485	490	495
485	490	495	
caa gat ctt ctg tct gag gaa aat gaa tcc aca gct cta ccc cag gtt			
Gln Asp Leu Leu Ser Glu Glu Asn Glu Ser Thr Ala Leu Pro Gln Val			
500	505	510	
505	510	515	
cta gcc cag cct tct act agt cga aag cgg ccc cgt gaa ggg gaa gcc			
Leu Ala Gln Pro Ser Thr Ser Arg Lys Arg Pro Arg Glu Gly Glu Ala			
515	520	525	
520	525	530	
530	535	540	
535	540	545	
tgaactccgt ggtttgaaca cggaaagaaat gtaccttctt tcactctgtc atctttctt			1754
tctttgagtc ttgtttttta tagtttgtat tttaattatg ggaataattg ctttttcaca			1814
gtcactgtat tacaattaaa aacctgatgg aacctggaaa aaaa			1858
<210> 2			
<211> 543			
<212> PRT			
<213> Homo sapiens			
<400> 2			
Met Ser Arg Glu Ser Asp Val Glu Ala Gln Ser His Gly Ser Ser			
1	5	10	15
Ala Cys Ser Gln Pro His Gly Ser Val Thr Gln Ser Gln Gly Ser Ser			
20	25	30	
Ser Gln Ser Gln Gly Ile Ser Ser Ser Ser Thr Ser Thr Met Pro Asn			
35	40	45	

Ser Ser Gln Ser Ser His Ser Ser Ser Gly Thr Leu Ser Ser Leu Glu
 50 55 60
 Thr Val Ser Thr Gln Glu Leu Tyr Ser Ile Pro Glu Asp Gln Glu Pro
 65 70 75 80
 Glu Asp Gln Glu Pro Glu Glu Pro Thr Pro Ala Pro Trp Ala Arg Leu
 85 90 95
 Trp Ala Leu Gln Asp Gly Phe Ala Asn Leu Glu Cys Val Asn Asp Asn
 100 105 110
 Tyr Trp Phe Gly Arg Asp Lys Ser Cys Glu Tyr Cys Phe Asp Glu Pro
 115 120 125
 Leu Leu Lys Arg Thr Asp Lys Tyr Arg Thr Tyr Ser Lys Lys His Phe
 130 135 140
 Arg Ile Phe Arg Glu Val Gly Pro Lys Asn Ser Tyr Ile Ala Tyr Ile
 145 150 155 160
 Glu Asp His Ser Gly Asn Gly Thr Phe Val Asn Thr Glu Leu Val Gly
 165 170 175
 Lys Gly Lys Arg Arg Pro Leu Asn Asn Ser Glu Ile Ala Leu Ser
 180 185 190
 Leu Ser Arg Asn Lys Val Phe Val Phe Asp Leu Thr Val Asp Asp
 195 200 205
 Gln Ser Val Tyr Pro Lys Ala Leu Arg Asp Glu Tyr Ile Met Ser Lys
 210 215 220
 Thr Leu Gly Ser Gly Ala Cys Gly Glu Val Lys Leu Ala Phe Glu Arg
 225 230 235 240
 Lys Thr Cys Lys Lys Val Ala Ile Lys Ile Ile Ser Lys Arg Lys Phe
 245 250 255
 Ala Ile Gly Ser Ala Arg Glu Ala Asp Pro Ala Leu Asn Val Glu Thr
 260 265 270
 Glu Ile Glu Ile Leu Lys Lys Leu Asn His Pro Cys Ile Ile Lys Ile
 275 280 285
 Lys Asn Phe Phe Asp Ala Glu Asp Tyr Tyr Ile Val Leu Glu Leu Met
 290 295 300
 Glu Gly Gly Glu Leu Phe Asp Lys Val Val Gly Asn Lys Arg Leu Lys
 305 310 315 320
 Glu Ala Thr Cys Lys Leu Tyr Phe Tyr Gln Met Leu Leu Ala Val Gln
 325 330 335
 Tyr Leu His Glu Asn Gly Ile Ile His Arg Asp Leu Lys Pro Glu Asn
 340 345 350
 Val Leu Leu Ser Ser Gln Glu Glu Asp Cys Leu Ile Lys Ile Thr Asp
 355 360 365
 Phe Gly His Ser Lys Ile Leu Gly Glu Thr Ser Leu Met Arg Thr Leu
 370 375 380
 Cys Gly Thr Pro Thr Tyr Leu Ala Pro Glu Val Leu Val Ser Val Gly
 385 390 395 400
 Thr Ala Gly Tyr Asn Arg Ala Val Asp Cys Trp Ser Leu Gly Val Ile
 405 410 415
 Leu Phe Ile Cys Leu Ser Gly Tyr Pro Pro Phe Ser Glu His Arg Thr
 420 425 430
 Gln Val Ser Leu Lys Asp Gln Ile Thr Ser Gly Lys Tyr Asn Phe Ile
 435 440 445
 Pro Glu Val Trp Ala Glu Val Ser Glu Lys Ala Leu Asp Leu Val Lys
 450 455 460
 Lys Leu Leu Val Val Asp Pro Lys Ala Arg Phe Thr Thr Glu Glu Ala
 465 470 475 480
 Leu Arg His Pro Trp Leu Gln Asp Glu Asp Met Lys Arg Lys Phe Gln
 485 490 495
 Asp Leu Leu Ser Glu Glu Asn Glu Ser Thr Ala Leu Pro Gln Val Leu
 500 505 510
 Ala Gln Pro Ser Thr Ser Arg Lys Arg Pro Arg Glu Gly Glu Ala Glu
 515 520 525
 Gly Ala Glu Thr Thr Lys Arg Pro Ala Val Cys Ala Ala Val Leu
 530 535 540

<210> 3

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer GSP3

<400> 3

ttttgctgat gatctttatg gctac	25
<210> 4	
<211> 25	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: PCR Primer	
GSP4	
<400> 4	
cacaggcacc acttccaaga gtttt	25
<210> 5	
<211> 36	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: PCR Primer	
<400> 5	
gggctcgaga gcagcgatgt ctcgggagtc ggatgt	36
<210> 6	
<211> 35	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> Description of Artificial Sequence: PCR Primer	
<400> 6	
ggcgaggatcct cgagtcacaa cacagcagca cacac	35