



(51) International Patent Classification:

C12N 15/09 (2006.01) C12N 15/113 (2010.01)

C12N 15/10 (2006.01) C12N 15/115 (2010.01)

C12N 15/11 (2006.01)

(21) International Application Number:

PCT/US2022/017582

(22) International Filing Date:

23 February 2022 (23.02.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/153,847 25 February 2021 (25.02.2021) US

63/285,851 03 December 2021 (03.12.2021) US

(71) Applicant: ARTISAN DEVELOPMENT LABS, INC.

[US/US]; 460 Oneida Street, Boulder, Colorado 80303 (US).

(72) Inventors: BARGHETTI, Andrea; 460 Oneida Street,

Boulder, Colorado 80303 (US). BAUMGARTNER,

Roland; 460 Oneida Street, Boulder, Colorado 80303 (US).

WARNECKE, Tanya; 460 Oneida Street, Boulder, Colorado 80303 (US). MARSHALL, Kurt; 460 Oneida Street, Boulder, Colorado 80303 (US). SCHIEL, John; 460 Oneida Street, Boulder, Colorado 80303 (US).

(74) Agent: WITT, Eric et al.; Storella & Witt, LLP, 2625 Alcatraz Avenue #197, Berkeley, California 94705 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) Title: COMPOSITIONS AND METHODS FOR TARGETING, EDITING, OR MODIFYING GENES




Figure 1A

(57) Abstract: Provided herein are nucleic acids useful as guide nucleic acids (gNAs), e.g., guide ribonucleic acids (gRNAs), in a CRISPR system wherein the guide nucleic acids contain one or more modifications to one or more nucleotides, use of such guide nucleic acids in modifying cells, and other uses wherein CRISPR Cas proteins are utilized.



UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

COMPOSITIONS AND METHODS FOR TARGETING, EDITING, OR MODIFYING GENES

CROSS-REFERENCE

- 5 **[0001]** This application claims the benefit of U.S. Provisional Application No. 63/153,847, filed February 25, 2021, and U.S. Provisional Application No. 63/285,851 filed December 3, 2021, which applications are incorporated herein by reference.

SEQUENCE LISTING

- 10 **[0002]** The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on February 12, 2021, is named ARTN-005_PRO_SL.txt and is 2,203,951 bytes in size.

BACKGROUND OF THE INVENTION

- 15 **[0003]** Recent advances have been made in precise genome targeting technologies. For example, specific loci in genomic DNA can be targeted, edited, or otherwise modified by designer meganucleases, zinc finger nucleases, or transcription activator-like effectors (TALEs). Furthermore, the CRISPR-Cas systems of bacterial and archaeal adaptive immunity have been adapted for precise targeting of genomic DNA in eukaryotic cells. Compared to the earlier
20 generations of genome editing tools, the CRISPR-Cas systems are easy to set up, scalable, and amenable to targeting multiple positions within the eukaryotic genome, thereby providing a major resource for new applications in genome engineering. Two distinct classes of CRISPR-Cas systems have been identified. Class 1 CRISPR-Cas systems utilize multi-protein effector complexes, whereas class 2 CRISPR-Cas systems utilize single-protein effectors. Among the
25 three types of class 2 CRISPR-Cas systems, type II and type V systems typically target DNA and type VI systems typically target RNA. Naturally occurring type II effector complexes consist of Cas9, CRISPR RNA (crRNA), and trans-activating CRISPR RNA (tracrRNA), but the crRNA and tracrRNA can be fused as a single guide RNA in an engineered system for simplicity. Certain naturally occurring type V systems, such as type V-A, type V-C, and type V-D systems,
30 do not require tracrRNA and use crRNA alone as the guide for cleavage of target DNA.

[0004] The CRISPR-Cas systems have been engineered for various purposes, such as genomic DNA cleavage, base editing, epigenome editing, and genomic imaging. Although significant developments have been made, there still remains a need for new and useful CRISPR-Cas systems as powerful precise genome targeting tools.

SUMMARY OF THE INVENTION

[0005] In one aspect, provided herein are compositions.

[0006] In certain embodiments, provided herein is a composition comprising a synthetic guide RNA (gRNA) comprising (i) a targeter nucleic acid with a 3' end and a 5' end, comprising: (a) a spacer sequence comprising the 3' end, configured to hybridize with a target nucleotide sequence, and (b) a targeter stem sequence comprising the 5' end; and (ii) a modulator nucleic acid with a 3' end and a 5' end, comprising (a) a modulator stem sequence comprising the 3' end, complementary to the targeter stem sequence, and (b) a 5' sequence, *e.g.*, tail sequence, comprising the 5' end; wherein the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids; and either the targeter nucleic acid or the modulator nucleic acid, or both, is modified at one or more nucleotides or internucleotide linkages at or near its 3' end, at or near its 5' end, or at or near both, and a complex comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating a CRISPR Associated (Cas) nuclease that, in a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. In certain embodiments, the modification is a chemical modification. In certain embodiments, the Cas nuclease is a Type V Cas nuclease, such as a type V-A, type V-C, or type V-D Cas nuclease, for example a type V-A Cas nuclease. In certain embodiments the Type V-A Cas nuclease is a Cpf1, MAD, Csm1, ART, or ABW nuclease, or derivative or variant thereof. In certain embodiments the composition further comprises the Cas nuclease that, in a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. In certain embodiments the composition further comprises the Cas nuclease. In certain embodiments the targeter nucleic acid, the modulator nucleic acid, and the Cas nuclease are present in a ribonucleoprotein (RNP) complex. In certain embodiments some or all of the nucleic acid is RNA. In certain embodiments the modification, *e.g.*, the chemical modification comprises a chemical modification at or near the 3' end of the targeter nucleic acid. In certain embodiments the chemical modification comprises a chemical modification at a nucleotide or internucleotide linkage within 10 nucleotides of the 3' end. In certain embodiments the chemical modification comprises a chemical modification to the 3' terminal nucleotide or internucleotide linkage. In certain embodiments comprising a modification at or near the 3' end of the targeter nucleic acid the composition further comprises a chemical modification at or near the 5' end of the targeter nucleic acid. In certain embodiments the chemical modifications at or near the 3' and 5' ends are the same. In certain embodiments the chemical modifications at or near the 3' and 5' ends are different. In certain embodiments comprising a modification at or near the 3' end, optionally, at or near the 5' end of the targeter nucleic acid the composition further comprises a chemical modification at or near the 3' end of the modulator nucleic acid. In certain embodiments the

chemical modification at or near the 3' end of the modulator nucleic acid is the same as the chemical modification at or near the 3' end of the targeter nucleic acid; different from the chemical modification at or near the 3' end of the targeter nucleic acid; the same as the chemical modification at or near the 5' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 5' end of the targeter nucleic acid, if present; or a combination thereof. In certain embodiments comprising a modification at or near the 3' and, optionally, at or near the 5' end of the targeter nucleic acid and a chemical modification at or near the 3' end of the modulator nucleic acid the composition further comprises a chemical modification at or near the 5' end of the modulator nucleic acid. In certain embodiments the chemical modification at or near the 5' end of the modulator nucleic acid is the same as the chemical modification at or near the 3' end of the targeter nucleic acid; different from the chemical modification at or near the 3' end of the targeter nucleic acid; the same as the chemical modification at or near the 5' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 5' end of the targeter nucleic acid, if present; the same as the chemical modification at or near the 3' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 3' end of the modulator nucleic acid, if present; or a combination thereof. In certain embodiments the chemical modification comprises a chemical modification at or near the 5' end of the targeter nucleic acid. In certain embodiments comprising a modification at or near the 5' end of the targeter nucleic acid the composition further comprises a chemical modification at or near the 3' end of the targeter nucleic acid. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are the same. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are different. In certain embodiments comprising a modification at or near the 5' end of the targeter nucleic acid and, optionally, a chemical modification at or near the 3' end of the targeter nucleic acid, the composition further comprises a chemical modification at or near the 3' end of the modulator nucleic acid. In certain embodiments the chemical modification at or near the 3' end of the modulator nucleic acid is the same as the chemical modification at or near the 5' end of the targeter nucleic acid; different from the chemical modification at or near the 5' end of the targeter nucleic acid; the same as the chemical modification at or near the 3' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 3' end of the targeter nucleic acid, if present; or a combination thereof. In certain embodiments comprising a modification at or near the 5' end of the targeter nucleic acid and, optionally, a chemical modification at or near the 3' end of the targeter nucleic acid and/or a chemical modification at or near the 3' end of the modulator nucleic acid the composition further comprises a chemical modification at or near the 5' end of the modulator nucleic acid. In certain embodiments the

chemical modification at or near the 5' end of the modulator nucleic acid is the same as the chemical modification at or near the 5' end of the targeter nucleic acid; different from the chemical modification at or near the 5' end of the targeter nucleic acid; the same as the chemical modification at or near the 3' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 3' end of the targeter nucleic acid, if present; the same as the chemical modification at or near the 3' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 3' end of the modulator nucleic acid, if present; or a combination thereof. In certain embodiments the chemical modification comprises a chemical modification at or near the 3' end of the modulator nucleic acid. In certain embodiments the composition further comprises a chemical modification at or near the 5' end of the modulator nucleic acid. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are the same. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are different. In certain embodiments comprising a modification at or near the 3' end of the modulator nucleic acid and, optionally, a modification at or near the 5' end of the modulator nucleic acid, the composition further comprises a chemical modification at or near the 3' end of the targeter nucleic acid. In certain embodiments the chemical modification at or near the 3' end of the targeter nucleic acid is the same as the chemical modification at or near the 3' end of the modulator nucleic acid; different from the chemical modification at or near the 3' end of the modulator nucleic acid; the same as the chemical modification at or near the 5' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 5' end of the modulator nucleic acid, if present; or a combination thereof. In certain embodiments comprising a modification at or near the 3' end of the modulator nucleic acid and, optionally, a modification at or near the 5' end of the modulator nucleic acid and/or a chemical modification at or near the 3' end of the targeter nucleic acid the composition further comprises a chemical modification at or near the 5' end of the targeter nucleic acid. In certain embodiments the chemical modification at or near the 5' end of the targeter nucleic acid is the same as the chemical modification at or near the 3' end of the modulator nucleic acid; different from the chemical modification at or near the 3' end of the modulator nucleic acid; the same as the chemical modification at or near the 5' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 5' end of the modulator nucleic acid, if present; the same as the chemical modification at or near the 3' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 3' end of the targeter nucleic acid, if present; or a combination thereof. In certain embodiments the chemical modification comprises a chemical modification at or near the 5' end of the modulator nucleic acid. In certain embodiments comprising a chemical modification at or near the 5' end of the modulator nucleic acid

acid the composition further comprises a chemical modification at or near the 3' end of the modulator nucleic acid. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are the same. In certain embodiments the chemical modifications at or near the 3' and at or near the 5' ends are different. In certain embodiments comprising a chemical

5 modification at or near the 5' end of the modulator nucleic acid and, optionally, a chemical modification at or near the 3' end of the modulator nucleic acid, the composition further comprises a chemical modification at or near the 3' end of the targeter nucleic acid. In certain

10 embodiments the chemical modification at or near the 3' end of the targeter nucleic acid is the same as the chemical modification at or near the 5' end of the modulator nucleic acid; different from the chemical modification at or near the 5' end of the modulator nucleic acid; the same as the chemical modification at or near the 3' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 3' end of the modulator nucleic acid, if present; or a combination thereof. In certain embodiments comprising a chemical modification at or near the

15 5' end of the modulator nucleic acid and, optionally, a chemical modification at or near the 3' end of the modulator nucleic acid and/or a chemical modification at or near the 3' end of the targeter nucleic acid the composition further comprises a chemical modification at or near the 5' end of the targeter nucleic acid. In certain embodiments the chemical modification at or near the 5' end of the targeter nucleic acid is the same as the chemical modification at or near the 5' end of the modulator nucleic acid; different from the chemical modification at or near the 5' end of the modulator nucleic acid; the same as the chemical modification at or near the 3' end of the modulator nucleic acid, if present; different from the chemical modification at or near the 3' end of the modulator nucleic acid, if present; the same as the chemical modification at or near the 3' end of the targeter nucleic acid, if present; different from the chemical modification at or near the 3' end of the targeter nucleic acid, if present; or a combination thereof. In any of the previous

25 **embodiments, the chemical modification can be selected from the group consisting of 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), and a combination thereof.** In any of the previous embodiments,

30 the spacer sequence can comprise a sequence capable of hybridizing with a human ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, CARD11, CD247, IL7R, LCK, PLCG1, ALP NR, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8,

35 PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2,

TAPBP, TWF1, CD3D, or NLRC5 gene. Any of the previous embodiments may further comprise a Cas protein, for example a Cas nuclease. In certain embodiments, provided is eukaryotic cell comprising the gRNA of any of the previous embodiments, in some cases further comprising a Cas nuclease to which the gRNA binds. In certain embodiments the cell is an immune cell such as a human immune cell. In certain embodiments the immune cell is a T cell. In certain embodiments, the immune cell is a CAR-T cell. In certain embodiments, the gNA-Cas complex is introduced into host cell, *e.g.*, an immune cell, *e.g.*, a T cell, along with an exogenous donor template, *e.g.*, a CAR cassette, where the the exogenous donor template is introduced into the genome of the host cell through the activity of the gNA-Cas complex resulting in a modified cell, *e.g.*, a CAR-T cell. In certain embodiments provided herein is a composition comprising any of the preceding composition and further comprising a Cas protein. In certain embodiments the Cas protein comprises a Cas nuclease. In certain embodiments the Cas nuclease is a Type I, II, III, IV, V, or VI Cas nuclease. In certain embodiments the Cas nuclease is a Type V Cas nuclease. In certain embodiments the Cas nuclease is a Type V-A, V-C, or V-D nuclease. In certain embodiments the Cas nuclease is a Type V-A Cas nuclease. In certain embodiments the Type V-A Cas nuclease is a Cpf1, MAD, Csm1, ART, or ABW Cas nuclease, or a derivative or variant thereof. In certain embodiments provided herein is a pharmaceutical composition comprising any of the preceding compositions and a pharmaceutically acceptable carrier.

[0007] In one aspect, provided herein are methods.

[0008] In certain embodiments, provided herein is method of cleaving a target DNA having a target nucleotide sequence, the method comprising contacting the target DNA with a composition of the preceding paragraph, thereby resulting in cleavage of the target DNA. In certain embodiments the contacting occurs *in vitro*. In certain embodiments the contacting occurs in a cell *ex vivo*. In certain embodiments the target DNA is genomic DNA of the cell. In certain embodiments the system is delivered into the cell as a pre-formed RNP complex. In certain embodiments the pre-formed RNP complex is delivered into the cell by electroporation, lipofection, or a viral method. In certain embodiments the pre-formed RNP complex is delivered into the cell by electroporation.

[0009] In certain embodiments provided herein is a method of editing the genome of a eukaryotic cell, the method comprising delivering the engineered, non-naturally occurring system of any of the embodiments of the first paragraph of this section, thereby resulting in editing of the genome of the eukaryotic cell. In certain embodiments the system is delivered into the cell as a pre-formed RNP complex. In certain embodiments the system is delivered into the cell by electroporation, lipofection, or a viral method. In certain embodiments the system is delivered into the cell by electroporation. In certain embodiments the cell is an immune cell. In

certain embodiments the immune cell is a T lymphocyte. In certain embodiments the engineered, non-naturally occurring system is delivered to a plurality of eukaryotic cells, and wherein the system comprises a guide nucleic acid comprising one or modifications as described herein, wherein the editing efficiency of the genomes of the plurality of cells is increased by at least 5% compared to the editing efficiency when the same system but without the modification or modifications is used.

[0010] In certain embodiments, provided herein is a method of treating a disease or a disorder comprising administering to a subject in need thereof an effective amount of a composition of the first paragraph of this section. In certain embodiments the method comprises administering to a subject in need thereof of cells modified by treatment with a composition of the first paragraph of this section. In certain embodiments the cells are cells that are removed from an individual and treated *ex vivo*. In certain embodiments the subject in need of treatment and the individual whose cells are treated *ex vivo* are the same.

INCORPORATION BY REFERENCE

[0011] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0013] **Figure 1A** shows a schematic representation showing the structure of an exemplary dual single guide CRISPR-Cas system.

[0014] **Figure 1B** shows a schematic representation showing the structure of an exemplary dual guide CRISPR-Cas system.

[0015] **Figure 2A-2C** show a series of schematic representations of exemplary modifications to dual guide gRNA. 2A: protecting group at 5' end of modulator nucleic acid; 2B: donor template recruiting sequence at 5' end of modulator nucleic acid; 2C: editing enhancer at 5' end of modulator nucleic acid.

[0016] **Figure 3** shows data for editing efficiency (as measured by # of reads modified/ total # of reads) in primary T cells in an exon of an exemplary gene 1. Shown are editing results relative to the single gRNA design (left bar) vs. the negative control (far right bar).

[0017] **Figure 4A** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 1. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0018] **Figure 4B** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 1. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0019] **Figure 5A** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 2. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0020] **Figure 5B** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 2. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0021] **Figure 6A** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 3. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0022] **Figure 6B** shows a series of data for editing efficiency in primary T cells in an exon of an exemplary gene 3. Shown are editing results of multiple modified gRNA designs over the single gRNA design (2 right bars).

[0023] **Figure 7** shows chemical structures for a 5' (panel A), 3' (panel B), and internal (panel C) propanediol modification.

[0024] **Figure 8** shows editing efficiency in primary T cells in a B2M gene. Shown are the editing results of multiple modified gRNA designs over the single gRNA design.

[0025] **Figure 9** shows HLA-1 surface expression knock down after treatment with RNP comprising multiple modified gRNA design compared to the single gRNA design targeting the B2M gene as measured by flow cytometry.

DETAILED DESCRIPTION OF THE INVENTION

I. Engineered, non-naturally occurring modified guide, *e.g.*, dual guide CRISPR-Cas Systems

A. Nucleic Acid Modifications to Guide Nucleic Acids

1. Specific Modifications to Targeter and/or Modulator Nucleic Acids

B. Targeter and Modulator Nucleic Acids

C. Cas Proteins

II. Methods of Targeting, Editing, and/or Modifying Genomic DNA

A. Ribonucleoprotein (RNP) Delivery and "Cas RNA" Delivery

- B. CRISPR Expression Systems
- C. Donor Templates
- D. Efficiency and Specificity
- E. Multiplex Methods
- 5 III. Pharmaceutical Compositions
- IV. Therapeutic Uses
 - A. Gene Therapies
 - B. Immune Cell Engineering
- V. Kits
- 10 VI. Embodiments
- VII. Examples

[0026] The invention is based, in part, upon the design of a modified guide CRISPR-Cas system, such as a modified dual guide CRISPR-Cas system in which a targeter nucleic acid and a modulator nucleic acid, when hybridized to form a complex, can, *e.g.*, activate a Cas nuclease that, in a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. The engineered modified dual guide CRISPR-Cas systems described herein can be used to target, edit, or modify a target nucleic acid such as genomic DNA. Modifications include a chemical modification to one or more nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid, at or near the 5' end of the targeter nucleic acid (if a dual guide system), at or near the 3' end of the modulator nucleic acid (if a dual guide system), at or near the 5' end of the modulator nucleic acid, or combinations thereof. In cases where more than one locus is modified, the chemical modification at each locus can be the same or different. In certain embodiments the modified guide nucleic acid (gNA) can be a single guide nucleic acid such as a single guide RNA, wherein the targeter and modulator nucleic acid are joined by a plurality of nucleotides; while embodiments are described in terms of dual guide nucleic acids it is understood that the same can be applied to single guide nucleic acids, where appropriate.

[0027] A CRISPR-Cas system generally comprises a Cas protein and one or more guide nucleic acids, *e.g.*, gRNAs. The Cas protein can be directed to a specific location in a double-stranded DNA target by recognizing a protospacer adjacent motif (PAM) in the non-target strand of the DNA, and the one or more guide nucleic acids can be directed to a specific location by hybridizing with a target nucleotide sequence in the target strand of the DNA. Both PAM recognition and target nucleotide sequence hybridization are required for stable binding of a CRISPR-Cas complex to the DNA target and, if the Cas protein has an effector function, *e.g.*, nuclease activity, activation of the effector function. As a result, when creating a CRISPR-Cas system, a guide nucleic acid can be designed to comprise a nucleotide sequence called spacer sequence that hybridizes with a target nucleotide sequence, where target nucleotide sequence is located adjacent to a PAM in an orientation operable with the Cas protein. It has been observed that not all CRISPR-Cas systems designed by these criteria are equally effective.

[0028] Type V-A, type V-C, and type V-D CRISPR-Cas systems naturally include a Cas nuclease and a single guide RNA (*i.e.*, crRNA) while lacking a tracrRNA. By splitting the single guide RNA into two different nucleic acids, where at least one end of one of the nucleic acids is chemically modified, the engineered system described herein provides better flexibility and tunability. For example, the efficiency of nucleic acid cleavage can be increased or decreased by adjusting the hybridization length and/or affinity of the targeter nucleic acid and the modulator nucleic acid. Furthermore, given the length limitation of nucleic acids that can be synthesized with high yield and accuracy, the use of modified dual guide nucleic acids allows incorporation of more polynucleotide elements that can improve editing efficacy and/or specificity.

[0029] In particular, the modified dual guide system can be engineered as a tunable system to decrease off-target editing, and thus can be used to edit a nucleic acid with high specificity. The system can be employed in a number of applications, for example, editing cells such as mammalian cells for use in therapy. A decrease in off-target editing is particularly desirable when creating genetically engineered proliferating cells, such as stem cells, progenitor cells, and immune memory cells, to be administered to a subject in need of the therapy. High specificity can be accomplished using the modified dual guide systems described herein, which optionally further include, for example, one or more chemical modifications to the targeter nucleic acid and/or modulator nucleic acid, an editing enhancer sequence, and/or a donor template-recruiting sequence. The nature and/or location of the chemical modifications can modulate editing efficiency in the CRISPR system. For example, in certain embodiments a modification at or near the 5' end, at or near the 3' end, and/or at or near both of a targeter and/or modulator nucleic acid, *e.g.*, one or more modifications to one or more nucleotides, as described elsewhere herein, can result in at least a 1, 2, 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, or 90% increase in editing efficiency; in some cases a similar decrease in editing efficiency may be achieved, compared to non-modulated nucleic acids.

[0030] Thus, provided herein are guide nucleic acids, such as RNAs, comprising a targeter nucleic acid and a modulator nucleic acid; see, *e.g.*, **Figure 1A**, showing a single guide nucleic acid, and **Figure 1B**, showing a dual guide nucleic acid. One or more nucleotides or internucleotide linkages at or near the 5' end (of the modulator nucleic acid in a sgNA, of either or both of modulator nucleic acid and/or targeter nucleic acid in dual gNA), at or near the 3' end (of the targeter nucleic acid in a sgNA, of either or both of modulator nucleic acid and/or targeter nucleic acid in dual gNA), or both of the targeter and/or modulator nucleic acids comprise one or more modified nucleotides or internucleotide linkages, *e.g.*, chemically modified nucleotides. Specific embodiments are as described herein, and include embodiments in which a specific gene is targeted by the modified guide nucleic acid, *e.g.*, modified single guide nucleic acid such as

modified single guide RNA, or modified dual guide nucleic acid such as a modified dual guide RNA. In certain embodiments, provided are compositions comprising a modified guide nucleic acid as described herein and a Cas protein, such as a Cas nuclease. The protein, *e.g.*, nuclease can be any suitable protein, *e.g.*, nuclease; in certain embodiments, the nuclease is a Type I, II, III, IV, V, or VI Cas nuclease; in certain embodiments the nuclease is Type V Cas nuclease, such as a Type V-A, V-C, or V-D nuclease, for example, a Type V-A nuclease. Specific nucleases are as described herein. In certain embodiments, the composition can also comprise a donor template. In certain embodiments, provided are CRISPR expression systems for expressing one or more of the nuclease, the targeter nucleic acid, the modulator nucleic acid, and/or, optionally, a donor template; it will be appreciated that, in general, the modified nucleic acids cannot be expressed by such a system. Also provided are cells, such as an immune cell, *e.g.*, a T cell, comprising one or more of the modified guide nucleic acids described herein, Cas nucleases as described herein, and/or donor template. In certain embodiments, provided are pharmaceutical compositions comprising compositions comprising modified guide nucleic acids, as described herein. In certain embodiments, provided are methods for targeting, editing, and/or modifying genomic DNA using the modified guide nucleic acid compositions described herein. In certain embodiments, provided herein are methods of gene therapy utilizing the modified guide nucleic acid compositions described herein. In certain embodiments, provided herein are methods of immune cell engineering utilizing the modified guide nucleic acid compositions described herein. In certain embodiments, provided are kits comprising the modified guide nucleic acids described herein.

[0031] The features and uses of the modified single and dual guide CRISPR-Cas system are discussed in detail in the following sections.

I. Engineered, Non-Naturally Occurring Modified Guide Nucleic Acid-CRISPR-Cas Systems

[0032] In certain embodiments, the engineered, non-naturally occurring system of the present invention comprises a targeter nucleic acid comprising a spacer sequence designed to hybridize with a target nucleotide sequence and a targeter stem sequence; and a modulator nucleic acid comprising a modulator stem sequence complementary to the targeter stem sequence and, optionally, a 5' sequence, *e.g.*, a tail sequence, wherein, in the case of a single guide nucleic acid the guide nucleic acid is a single polynucleotide, and in the case of a dual guide nucleic acid, the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids, and wherein a guide nucleic acid comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating a Cas nuclease; in certain cases of dual gNAs, the nuclease is one that, in

a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. See **Figures 1A and 1B**. One or both of the targeter nucleic acid and/or the modulator nucleic acid includes one or more modified nucleotides or internucleotide linkages at or near the 3' end, at or near the 5' end, or at or near both.

5 **[0033]** The terms “targeter stem sequence” and “modulator stem sequence,” as used herein, can include a pair of nucleotide sequences in one or more guide nucleic acids that hybridize with each other. When a targeter stem sequence and a modulator stem sequence are contained in a single guide nucleic acid, the targeter stem sequence is proximal to a spacer sequence designed to hybridize with a target nucleotide sequence, and the modulator stem
10 sequence is proximal to the targeter stem sequence. When a targeter stem sequence and a modulator stem sequence are in separate nucleic acids, i.e. in a dual guide nucleic acid, the targeter stem sequence is in the same nucleic acid as a spacer sequence designed to hybridize with a target nucleotide sequence. In a CRISPR-Cas system that naturally includes separate crRNA and tracrRNA (*e.g.*, a type II system), the duplex formed between the targeter stem
15 sequence and the modulator stem sequence corresponds to the duplex formed between the crRNA and the tracrRNA. In a CRISPR-Cas system that naturally includes a single crRNA but no tracrRNA (*e.g.*, a type V-A system), the duplex formed between the targeter stem sequence and the modulator stem sequence corresponds to the stem portion of a stem-loop structure in the scaffold sequence (also called direct repeat sequence) of the crRNA. It is understood that 100%
20 complementarity is not required between the targeter stem sequence and the modulator stem sequence. In a type V-A CRISPR-Cas system, however, the targeter stem sequence is typically 100% complementary to the modulator stem sequence.

[0034] In certain embodiments wherein the target nucleic acid and the modulator nucleic acid comprise a single polynucleotide, a loop motif may exist between the 3' stem sequence of
25 the targeter nucleic acid and the 5' stem sequence of the modulator nucleic acid, *e.g.*, a stem loop. In certain embodiments, the loop motif is between 1-11, 2-11, 3-11, 4-11, 5-11, 3-10, 3-9, 3-8, 3-7, 3-6, 1-11, 2-10, 3-9, 4-8, 5-7, 4-6, 1-7, 2-6, 3-5 nucleotides in length. In a preferred embodiment, the loop motif is between 3-5 nucleotides in length. In a separate preferred embodiment, the loop motif is four nucleotides in length. In certain embodiments, the loop motif
30 is 5'-TCTT-3' or 5'-TATT-3'.

[0035] The term “targeter nucleic acid,” as used herein in the context of a dual guide nucleic acid CRISPR-Cas system, can include a nucleic acid comprising (i) a spacer sequence designed to hybridize with a target nucleotide sequence; and (ii) a targeter stem sequence capable of hybridizing with an additional nucleic acid to form a complex, wherein the complex is capable
35 of activating a Cas nuclease (*e.g.*, a type II or type V-A Cas nuclease) under suitable conditions,

and wherein the targeter nucleic acid alone, in the absence of the additional nucleic acid, is not capable of activating the Cas nuclease under the same conditions. The term “targeter nucleic acid,” as used herein in the context of a single guide nucleic acid CRISPR-Cas system, can include a nucleic acid comprising (i) a spacer sequence designed to hybridize with a target nucleotide sequence; and (ii) a targeter stem sequence capable of hybridizing with a complementary stem sequence in a modulator nucleic acid that is 5' to the targeter nucleic acid in the single polynucleotide of the sgNA, wherein the sgNA is capable of activating a Cas nuclease (*e.g.*, a type II or type V-A Cas nuclease).

[0036] The term “modulator nucleic acid,” as used herein in connection with a given targeter nucleic acid and its corresponding Cas nuclease, can include a nucleic acid capable of hybridizing with the targeter nucleic acid, to form an intra-polynucleotide hybridized portion in the case of a sgNA, and to form a complex in the case of a dual gNA, wherein the sgNA or complex, but not the modulator nucleic acid alone, is capable of activating the type Cas nuclease under suitable conditions.

[0037] The term “suitable conditions,” as used in connection with the definitions of “targeter nucleic acid” and “modulator nucleic acid,” can include the conditions under which a naturally occurring CRISPR-Cas system is operative, such as in a prokaryotic cell, in a eukaryotic (*e.g.*, mammalian or human) cell, or in an *in vitro* assay.

[0038] Type V-A, type V-C, and type V-D CRISPR-Cas systems are distinctive subtypes of CRISPR-Cas systems under the classification described in Makarova *et al.* (2017) CELL, 168: 328. Naturally occurring CRISPR-Cas systems of these subtypes lack a tracrRNA and rely on a single crRNA to guide the CRISPR-Cas complex to the target DNA. Naturally occurring type V-A Cas proteins comprise a RuvC-like nuclease domain but lack an HNH endonuclease domain, and recognize a 5' T-rich protospacer adjacent motif (PAM), the 5' orientation determined using the non-target strand (*i.e.* the strand not hybridized with the spacer sequence) as the coordinate.

[0039] Naturally occurring type V-A CRISPR-Cas systems cleave a double-stranded DNA to generate a staggered double-stranded break rather than a blunt end. The cleavage site is distant from the PAM site (*e.g.*, separated by at least 10, 11, 12, 13, 14, or 15 nucleotides from the PAM on the non-target strand and/or separated by at least 15, 16, 17, 18, or 19 nucleotides from the sequence complementary to PAM on the target strand).

[0040] The instant disclosure provides an engineered, non-naturally occurring system comprising a targeter nucleic acid comprising: a spacer sequence designed to hybridize with a target nucleotide sequence; and a targeter stem sequence; and a modulator nucleic acid comprising a modulator stem sequence complementary to the targeter stem sequence, and, optionally, a 5' sequence, *e.g.*, a tail sequence, wherein, in the case of a single guide nucleic acid

the targeter nucleic acid and the modulator nucleic acid are part of a single polynucleotide, and in the case of a dual guide nucleic acid, the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids; modifications can include one or more chemical modifications to one or more nucleotides at or near the 3' end of the targeter nucleic acid (dual and single gNA), at or near the 5' end of the targeter nucleic acid (dual gNA), at or near the 3' end of the modulator nucleic acid (dual gNA), at or near the 5' end of the modulator nucleic acid (single and dual gNA), or combinations thereof, and wherein the gNA comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating a Cas nuclease, such as a Type I, II, III, IV, V, or VI Cas nuclease, such as a Type V Cas nuclease, for example, a type V-A, type V-C, or type V-D Cas nuclease. In certain embodiments, the Cas nuclease is a type V-A Cas nuclease. In certain embodiments the targeter sequence comprises, from 5' to 3', a targeter stem sequence and a spacer sequence and the modulator sequence comprises, from 5' to 3', a 5' sequence, *e.g.*, a tail sequence, and a modulator stem sequence. In certain embodiments, the system also comprises a Cas nuclease, such as type V-A, type V-C, or type V-D Cas nuclease, for example, a Type V-A Cas nuclease.

A. Nucleic Acid Modifications in Guide Nucleic Acids

[0041] Provided herein are engineered, non-naturally occurring systems comprising a targeter nucleic acid comprising: a spacer sequence designed to hybridize with a target nucleotide sequence and a targeter stem sequence; and a modulator nucleic acid comprising a modulator stem sequence complementary to the targeter stem sequence, and, optionally, a 5' sequence, *e.g.*, a tail sequence, wherein, in a single guide nucleic acid the targeter nucleic acid and the modulator nucleic acid are part of a single polynucleotide, and in a dual guide nucleic acid, the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids; modifications can include one or more chemical modifications to one or more nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid (dual and single gNA), at or near the 5' end of the targeter nucleic acid (dual gNA), at or near the 3' end of the modulator nucleic acid (dual gNA), at or near the 5' end of the modulator nucleic acid (single and dual gNA), or combinations thereof as appropriate for single or dual gNA. In certain embodiments, the Cas nuclease is a type V-A Cas nuclease. Modulator and/or targeter nucleic sequences can include further sequences, as detailed in **Section IB**, and modifications can be in these further sequences, as appropriate and apparent to one of skill in the art. In embodiments described in this section, below, in certain embodiments, guide nucleic acid is oriented from 5' at the modulator nucleic acid to 3' at the modulator stem sequence, and 5' at the targeter stem sequence to 3' at the targeter sequence (see, *e.g.*, **Figure 1A and 1B**); in certain embodiments, as

appropriate, guide nucleic acid is oriented from 3' at the modulator nucleic acid to 5' at the modulator stem sequence, and 3' at the targeter stem sequence to 5' at the targeter sequence.

[0042] The targeter nucleic acid may comprise a DNA (*e.g.*, modified DNA), an RNA (*e.g.*, modified RNA), or a combination thereof. The modulator nucleic acid may comprise a DNA (*e.g.*, modified DNA), an RNA (*e.g.*, modified RNA), or a combination thereof. In certain embodiments, the targeter nucleic acid is an RNA and the modulator nucleic acid is an RNA. A targeter nucleic acid in the form of an RNA is also called targeter RNA, and a modulator nucleic acid in the form of an RNA is also called modulator RNA. The nucleotide sequences disclosed herein are presented as DNA sequences by including thymidines (T) and/or RNA sequences including uridines (U). It is understood that corresponding DNA sequences, RNA sequences, and DNA/RNA chimeric sequences are also contemplated. For example, where a spacer sequence is presented as a DNA sequence, a nucleic acid comprising this spacer sequence as an RNA can be derived from the DNA sequence disclosed herein by replacing each T with U. As a result, for the purpose of describing a nucleotide sequence, T and U are used interchangeably herein.

[0043] In certain embodiments some or all of the gNA is RNA, *e.g.*, a gRNA. In certain embodiments, 5-100%, 10-100%, 20-100%, 30-100%, 40-100%, 50-100%, 60-100%, 70-100%, 80-100%, 90-100%, 95-100%, 99-100%, 99.5-100% of the gNA is gRNA. In certain embodiments, 20%-80%, 20%-70%, 20%-60%, 20%-50%, 20%-40%, 20%-30%, 30%-80%, 30%-70%, 30%-60%, 30%-50%, 30%-40%, 40%-80%, 40%-70%, 40%-60%, 40%-50%, 50%-80%, 50%-70%, 50%-60%, 60%-80%, 60%-70%, or 70%-80% of gNA is RNA. In certain embodiments, 50% of the gNA is RNA. In certain embodiments, 70% of the gNA is RNA. In certain embodiments, 90% of the gNA is RNA. In certain embodiments, 100% of the gNA is RNA, *e.g.*, a gRNA.

[0044] In certain embodiments the stem sequences are 1-20, 2-19, 3-18, 4-17, 5-16, 6-15, 7-14, 8-13, 9-12, 10-11, 1-9, 2-8, 3-7, 4-6, or 2-9 nucleotides in length. In a preferred embodiment, the stem sequences are 4-6 nucleotides in length. In certain embodiments, the stem sequence of the modulator and targeter nucleic acids share 5%-100%, 10%-100%, 20%-100%, 30%-100%, 40%-100%, 50%-100%, 60%-100%, 70%-100%, 80%-100%, 90%-100%, 95%-100%, 99%-100%, 99.5%-100% of the gNA is gRNA. In certain embodiments, 20%-80%, 20%-70%, 20%-60%, 20%-50%, 20%-40%, 20%-30%, 30%-80%, 30%-70%, 30%-60%, 30%-50%, 30%-40%, 40%-80%, 40%-70%, 40%-60%, 40%-50%, 50%-80%, 50%-70%, 50%-60%, 60%-80%, 60%-70%, or 70%-80% sequence complementarity. In certain embodiments, the stem sequence of the modulator and targeter nucleic acids share 80%, 90%, 95%, or 100% sequence complementarity. In a preferred embodiment, the stem sequence of the modulator and targeter nucleic acids share 80%-100% sequence complementarity.

[0045] In certain embodiments, the targeter nucleic acid and/or the modulator nucleic acid are RNAs with one or more modifications in a ribose group, one or more modifications in a phosphate group, one or more modifications in a nucleobase, one or more terminal modifications, or a combination thereof. Exemplary modifications are disclosed in U.S. Patent Nos. 10,900,034 and 10,767,175, U.S. Patent Application Publication No. 2018/0119140, Watts *et al.* (2008) *Drug Discov. Today* 13: 842-55, and Hendel *et al.* (2015) *NAT. BIOTECHNOL.* 33: 985.

[0046] Modifications in a ribose group include but are not limited to modifications at the 2' position or modifications at the 4' position. For example, in certain embodiments, the ribose comprises 2'-O-C1-4alkyl, such as 2'-O-methyl (2'-OMe, or M). In certain embodiments, the ribose comprises 2'-O-C1-3alkyl-O-C1-3alkyl, such as 2'-methoxyethoxy (2'-O—CH₂CH₂OCH₃) also known as 2'-O-(2-methoxyethyl) or 2'-MOE. In certain embodiments, the ribose comprises 2'-O-allyl. In certain embodiments, the ribose comprises 2'-O-2,4-Dinitrophenol (DNP). In certain embodiments, the ribose comprises 2'-halo, such as 2'-F, 2'-Br, 2'-Cl, or 2'-I. In certain embodiments, the ribose comprises 2'-NH₂. In certain embodiments, the ribose comprises 2'-H (*e.g.*, a deoxynucleotide). In certain embodiments, the ribose comprises 2'-arabino or 2'-F-arabino. In certain embodiments, the ribose comprises 2'-LNA or 2'-ULNA. In certain embodiments, the ribose comprises a 4'-thioribosyl.

[0047] Modifications can also include a deoxy group, for example a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP).

[0048] Internucleotide linkage modifications in a phosphate group include but are not limited to a phosphorothioate (S), a chiral phosphorothioate, a phosphorodithioate, a boranophosphonate, a C₁₋₄alkyl phosphonate such as a methylphosphonate, a boranophosphonate, a phosphonocarboxylate such as a phosphonoacetate (P), a phosphonocarboxylate ester such as a phosphonoacetate ester, an amide, a thiophosphonocarboxylate such as a thiophosphonoacetate (SP), a thiophosphonocarboxylate ester such as a thiophosphonoacetate ester, and a 2',5'-linkage having a phosphodiester or any of the modified phosphates above. Various salts, mixed salts and free acid forms are also included.

[0049] Modifications in a nucleobase include but are not limited to 2-thiouracil, 2-thiocytosine, 4-thiouracil, 6-thioguanine, 2-aminoadenine, 2-aminopurine, pseudouracil, hypoxanthine, 7-deazaguanine, 7-deaza-8-azaguanine, 7-deazaadenine, 7-deaza-8-azaadenine, 5-methylcytosine, 5-methyluracil, 5-hydroxymethylcytosine, 5-hydroxymethyluracil, 5,6-dehydrouracil, 5-propynylcytosine, 5-propynyluracil, 5-ethynylcytosine, 5-ethynyluracil, 5-allyluracil, 5-allylcytosine, 5-aminoallyluracil, 5-aminoallyl-cytosine, 5-bromouracil, 5-iodouracil, diaminopurine, difluorotoluene, dihydrouracil, an abasic nucleotide, Z base, P base,

Unstructured Nucleic Acid, isoguanine, isocytosine (see, Piccirilli *et al.* (1990) NATURE, 343: 33), 5-methyl-2-pyrimidine (see, Rappaport (1993) BIOCHEMISTRY, 32: 3047), x(A,G,C,T), and y(A,G,C,T).

[0050] Terminal modifications include but are not limited to polyethyleneglycol (PEG), hydrocarbon linkers (such as heteroatom (O,S,N)-substituted hydrocarbon spacers; halo-substituted hydrocarbon spacers; keto-, carboxyl-, amido-, thionyl-, carbamoyl-, thionocarbamoyl-containing hydrocarbon spacers, propanediol), spermine linkers, dyes such as fluorescent dyes (for example, fluoresceins, rhodamines, cyanines), quenchers (for example, dabcy1, BHQ), and other labels (for example biotin, digoxigenin, acridine, streptavidin, avidin, peptides and/or proteins). In certain embodiments, a terminal modification comprises a conjugation (or ligation) of the RNA to another molecule comprising an oligonucleotide (such as deoxyribonucleotides and/or ribonucleotides), a peptide, a protein, a sugar, an oligosaccharide, a steroid, a lipid, a folic acid, a vitamin and/or other molecule. In certain embodiments, a terminal modification incorporated into the RNA is located internally in the RNA sequence via a linker such as 2-(4-butylamidofluorescein)propane-1,3-diol bis(phosphodiester) linker, which is incorporated as a phosphodiester linkage and can be incorporated anywhere between two nucleotides in the RNA.

[0051] The modifications disclosed above can be combined in the targeter nucleic acid and/or the modulator nucleic acid that are in the form of RNA. In certain embodiments, the modification in the RNA is selected from the group consisting of incorporation of 2'-O-methyl-3'phosphorothioate (MS), 2'-O-methyl-3'-phosphonoacetate (MP), 2'-O-methyl-3'-thiophosphonoacetate (MSP), 2'-halo-3'-phosphorothioate (*e.g.*, 2'-fluoro-3'-phosphorothioate), 2'-halo-3'-phosphonoacetate (*e.g.*, 2'-fluoro-3'-phosphonoacetate), and 2'-halo-3'-thiophosphonoacetate (*e.g.*, 2'-fluoro-3'-thiophosphonoacetate).

[0052] In certain embodiments, modifications can include 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), or a combination thereof, at or near either the 3' or 5' end of either the targeter or modulator nucleic acid, as appropriate for single or dual gNA.

[0053] In certain embodiments, modifications can include either a 5' or a 3' propanediol or C3 linker modification as depicted in **Figure 7A** or **B** respectively.

[0054] In certain embodiments, the modification alters the stability of the RNA. In certain embodiments, the modification enhances the stability of the RNA, *e.g.*, by increasing nuclease resistance of the RNA relative to a corresponding RNA without the modification.

Stability-enhancing modifications include but are not limited to incorporation of 2'-O-methyl, a 2'-O-C₁₋₄alkyl, 2'-halo (*e.g.*, 2'-F, 2'-Br, 2'-Cl, or 2'-I), 2'MOE, a 2'-O-C₁₋₃alkyl-O-C₁₋₃alkyl, 2'-NH₂, 2'-H (or 2'-deoxy), 2'-arabino, 2'-F-arabino, 4'-thioribosyl sugar moiety, 3'-phosphorothioate, 3'-phosphonoacetate, 3'-thiophosphonoacetate, 3'-methylphosphonate, 3'-boranophosphate, 3'-phosphorodithioate, locked nucleic acid ("LNA") nucleotide which comprises a methylene bridge between the 2' and 4' carbons of the ribose ring, and unlocked nucleic acid ("ULNA") nucleotide. Such modifications are suitable for use as a protecting group to prevent or reduce degradation of the 5' sequence, *e.g.*, a tail sequence, modulator stem sequence (dual guide nucleic acids), targeter stem sequence (dual guide nucleic acids), and/or spacer sequence (see, the "Targeter and Modulator nucleic acids" subsection).

1. Specific Modifications to Targeter and/or Modulator Nucleic Acids

[0055] In certain embodiments, a targeter nucleic acid, *e.g.*, RNA, comprises at least one nucleotide at or near the 3' end comprising a modification to a ribose, phosphate group, nucleobase, or terminal modification. In certain embodiments, the 3' end of the targeter nucleic acid comprises the spacer sequence. In certain embodiments, the 3' end of the targeter nucleic acid comprises the targeter stem sequence. Exemplary modifications are disclosed in Dang *et al.* (2015) *Genome Biol.* 16: 280, Kocaz *et al.* (2019) *Nature Biotech.* 37: 657-66, Liu *et al.* (2019) *Nucleic Acids Res.* 47(8): 4169-4180, Schubert *et al.* (2018) *J. Cytokine Biol.* 3(1): 121, Teng *et al.* (2019) *Genome Biol.* 20(1): 15, Watts *et al.* (2008) *Drug Discov. Today* 13(19-20): 842-55, and Wu *et al.* (2018) *Cell Mol. Life. Sci.* 75(19): 3593-607.

[0056] In certain embodiments, one or more nucleotides or internucleotide linkages within 15, 10, 5, 4, 3, 2, or 1 nucleotides of the 3' end of the targeter nucleic acid is modified. In certain embodiments, the nucleotide or internucleotide linkage at or near the 3' end of the targeter nucleic acid is modified. In certain embodiments, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 10, 5, 4, 3, or 2, or nucleotides of the 3' end of the targeter nucleic acid (as appropriate for total number of nucleotides or internucleotide linkages modified) of the targeter nucleic acid are modified, wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, modifications can include 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), or a combination thereof, at or near the 3' end of the targeter nucleic acid. In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides

of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 3' end of the targeter sequence, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides of the 3' end, for example the 3' end nucleotide, is modified, one or more other nucleotides at or near the 3' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides are also modified, for example with one or more of the modifications just described.

[0057] In certain embodiments, a targeter nucleic acid in a dual gNA, *e.g.*, dual gRNA, comprises at least one nucleotide at or near the 5' end comprising a modification to a ribose, phosphate internucleotide linkage, nucleobase, or terminal modification. In certain embodiments, the 5' end of the targeter nucleic acid comprises the spacer sequence. In certain embodiments, the 5' end of the targeter nucleic acid comprises the targeter stem sequence.

[0058] In certain embodiments, a nucleotide or internucleotide linkage within 15, 10, 5, 4, 3, 2, or 1 nucleotides of the 5' end of the targeter nucleic acid is modified. In certain embodiments, the nucleotide or internucleotide linkage at or near the 5' end of the targeter

nucleic acid is modified. In certain embodiments, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 10, 5, 4, 3, or 2, or nucleotides of the 5' end of the targeter nucleic acid (as appropriate for total number of nucleotides or internucleotide linkages modified) of the targeter nucleic acid are modified, wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, modifications can include 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), or a combination thereof, at or near the 5' end of the targeter nucleic acid. In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end nucleotide, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide or internucleotide

linkage, is modified, one or more other nucleotides or internucleotide linkages at or near the 5' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

5 **[0059]** In certain embodiments, a modulator nucleic acid in a dual gNA, *e.g.*, dual gRNA, comprises at least one nucleotide or internucleotide linkage at or near the 3' end comprising a modification to a ribose, phosphate internucleotide linkage, nucleobase, or terminal modification. In certain embodiments, the 3' end of the modulator nucleic acid comprises a modulator stem sequence. In certain embodiments, the 5' end of the modulator nucleic acid
10 includes a 5' sequence, *e.g.*, a tail sequence. In certain embodiments, one or more nucleotides or internucleotide linkage within 15, 10, 5, 4, 3, 2, or 1 nucleotides of the 3' end of the modulator nucleic acid is modified. In certain embodiments, the nucleotide or internucleotide linkage at or near the 3' end of the modulator nucleic acid is modified. In certain embodiments, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkage within 15, 10, 5, 4, 3, or 2, or nucleotides of the
15 3' end of the modulator nucleic acid (as appropriate for total number of nucleotides modified) of the modulator nucleic acid are modified, wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, modifications can include 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a
20 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), or a combination thereof, at or near the 3' end of the modulator nucleic acid. In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl (M). In
25 certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkages of the 3' end, for example the 3' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide
30 linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for
35 **example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain**

embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, one or more other nucleotides or internucleotide linkages at or near the 3' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

[0060] In certain embodiments, a modulator nucleic acid, *e.g.*, RNA, such as a single or dual gRNA, *e.g.*, single or dual gRNA, comprises at least one nucleotide at or near the 5' end comprising a modification to a ribose, phosphate group, nucleobase, or terminal modification. In certain embodiments, the 3' end of the modulator nucleic acid of a dual gRNA comprises the modulator stem sequence. In certain embodiments, the 5' end of the modulator nucleic acid comprises a 5' sequence, *e.g.*, a tail sequence. In certain embodiments, a nucleotide or internucleotide linkage within 15, 10, 5, 4, 3, 2, or 1 nucleotides of the 5' end of the modulator nucleic acid is modified. In certain embodiments, the nucleotide or internucleotide linkage at or near the 5' end of the modulator nucleic acid is modified. In certain embodiments, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 10, 5, 4, 3, or 2, or nucleotides of the 5' end of the modulator nucleic acid (as appropriate for total number of nucleotides modified) of the modulator nucleic acid are modified, wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, modifications can include 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), or a combination thereof, at or near the 5' end of the modulator nucleic acid. In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic

acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is modified, one or more other nucleotides or internucleotide linkages at or near the 5' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

[0061] In embodiments in which one or more nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid in a dual gNA, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, one or more nucleotides or internucleotide linkages at or near the 5' end of the targeter nucleic acid, for example, within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is also modified. In certain embodiments, a nucleotide

at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or internucleotide linkage at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, and a nucleotide or internucleotide linkage at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is modified, one or more other nucleotides or internucleotide linkage at or near the 5' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

[0062] In embodiments in which one or more nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, and/or one or more nucleotides or internucleotide linkages at or near the 5' end of the targeter nucleic

acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is modified, a nucleotide or internucleotide linkage at or near the 3' end of a modulator nucleic acid, for example, within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is also modified. In

5 certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a

10 phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the 3' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 3' end, for example the

15 3' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for

20 **example the 3' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain** embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a **2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the**

25 **3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In** certain embodiments, a nucleotide at or near the 3' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, **comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or**

30 **internucleotide linkage at or near the 3' end of the modulator nucleic acid, for example within** 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, one or more other nucleotides or internucleotide linkages at or near the 3' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

[0063] In embodiments in which one or more nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, and/or one or more nucleotides or internucleotide linkages at or near the 5' end of the targeter nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotides of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is modified, and/or one or more nucleotides or internucleotide linkages at or near the 3' end of a modulator nucleic acid, for example, within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide or internucleotide linkage, is modified, one or more nucleotides or internucleotide linkages at or near the 5' end of a modulator nucleic acid, for example, within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide or internucleotide linkage, is modified. In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl (M). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphorothioate (S). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a phosphonoacetate (P). In certain embodiments, an internucleotide linkage at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 internucleotide linkage of the 5' end, for example the 5' end internucleotide linkage, comprises a thiophosphonoacetate (SP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphorothioate (MS). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-phosphonoacetate (MP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-O-methyl-3'-thiophosphonoacetate (MSP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-phosphonoacetate (DP). In certain embodiments, a nucleotide at or near the 5' end of the modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 5' end nucleotide, comprises a 2'-deoxy-3'-thiophosphonoacetate (DSP). In embodiments in which a nucleotide or internucleotide linkage at or near the 5' end of the

modulator nucleic acid, for example within 10, 5, 4, 3, 2, or 1 nucleotide of the 5' end, for example the 3' end nucleotide or internucleotide linkage, is modified, one or more other nucleotides or internucleotide linkages at or near the 5' end are also modified, for example, an additional 1, 2, 3, 4, 5, 6, 7, 8, or 9 nucleotides or internucleotide linkages are modified, for example with one or more of the modifications just described.

[0064] In certain embodiments in which the 3' end of the targeter nucleic acid is unmodified, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator may be modified. In certain embodiments, the modifications comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides or internucleotide linkages of the 5' end of the modulator (as appropriate for total number of nucleotides modified), wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, the modifications comprise 1-5, 1-4, 1-3, 1-2, 2-4, 2-3, for example 1, 2, 3, 4, or 5, 2'-O-methoxy-3'-phosphorothioate modifications within 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) nucleotides of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 1-3, for example 1, 2, or 3, 2'-O-methoxy-3'-phosphorothioate modifications within the first 3 nucleotides of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 2 nucleotides of the 5' end of the modulator nucleic acid are 2'-O-methoxy-3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0065] In certain embodiments in which the 3' end of the targeter nucleic acid is unmodified, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator may be modified. In certain embodiments, the modifications comprise 1-17, 2-16, 3-15, 4-14, 5-13, 6-12, 7-11, or 8-10, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, phosphorothioate modification within the first 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) internucleotide linkages of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 8-10, for example 8, 9 or 10, phosphorothioate modifications within the first 10, 9 or 8 internucleotide linkages (as appropriate for total number of internucleotide linkages modified) of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 9 internucleotide linkages of the 5' end of the modulator nucleic acid are phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0066] In certain embodiments in which the 3' end of the targeter nucleic acid is unmodified, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator may be modified. In certain embodiments, modifications comprise 1-3, for example, 1, 2, or 3, 2'-O-methoxy and/or 1-3, for example, 1, 2, or 3, 3'-phosphorothioate modifications within the first 3 nucleotides (as appropriate for the total number of nucleotides modified) of the 5' end of the modulator nucleic acid. In this embodiment, any combination of 2'-O-methoxy nucleotide modifications and phosphorothioate internucleotide modifications may be used. In a preferred embodiment, the first nucleotide from the 5' end of the modulator nucleic acid is 2'-O-methoxy-3'-phosphorothioate modified and the second internucleotide linkage is phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0067] In certain embodiments in which the 3' end of the targeter nucleic acid is unmodified, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the 5' end of the modulator nucleic acid is modified with a terminal propanediol. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0068] In certain embodiments in which the 3' end of the targeter nucleic acid comprises a terminal propanediol modification, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides or internucleotide linkages of the 5' end of the modulator (as appropriate for total number of nucleotides modified), wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, the modifications comprise 1-5, 1-4, 1-3, 1-2, 2-4, 2-3, for example 1, 2, 3, 4, or 5, 2'-O-methoxy-3'-phosphorothioate modifications within 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) nucleotides of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 1-3, for example 1, 2, or 3, 2'-O-methoxy-3'-phosphorothioate modifications within the first 3 nucleotides of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 2 nucleotides of the 5' end of the modulator nucleic acid are 2'-O-methoxy-3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0069] In certain embodiments in which the 3' end of the targeter nucleic acid comprises a terminal propanediol modification, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications comprise 1-17, 2-16, 3-15, 4-14, 5-13, 6-12, 7-11, or 8-10, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, phosphorothioate modification within the first 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) internucleotide linkages of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 8-10, for example 8, 9 or 10, 3 phosphorothioate modifications within the first 10, 9 or 8 internucleotide linkages (as appropriate for total number of internucleotide linkages modified) of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 9 internucleotide linkages of the 5' end of the modulator nucleic acid are phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0070] In certain embodiments in which the 3' end of the targeter nucleic acid comprises a terminal propanediol modification, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, modifications comprise 1-3, for example, 1, 2, or 3, 2'-O-methoxy and /or 1-3, for example, 1, 2, or 3, 3'-phosphorothioate modifications within the first 3 nucleotide (as appropriate for the total number of nucleotides modified) of the 5' end of the modulator nucleic acid. In this embodiment, any combination of 2'-O-methoxy and 3'-phosphorothioate modifications may be used. In a preferred embodiment, the first nucleotide from the 5' end of the modulator nucleic acid is 2'-O-methoxy-3'-phosphorothioate modified and the second nucleotide is 3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0071] In certain embodiments in which the 3' end of the targeter nucleic acid comprises a terminal propanediol modification, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the 5' end of the modulator nucleic acid is modified with a terminal propanediol. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0072] In certain embodiments in which the 3' end of the targeter nucleic acid comprises two 2'-O-methoxy-3'-phosphorothioate modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or

internucleotide linkages within 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides or internucleotide linkages of the 5' end of the modulator (as appropriate for total number of nucleotides modified), wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, the modifications comprise 1-5, 1-4, 1-3, 1-2, 2-4, 2-3, for example 1, 2, 3, 4, or 5, 2'-O-methoxy-3'-phosphorothioate modifications within 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) nucleotides of the 5' end of the modulator nucleic acid. In certain embodiments, the the modifications comprise 1-3, for example 1, 2, or 3, 2'-O-methoxy-3'-phosphorothioate modifications within the first 3 nucleotides of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 2 nucleotides of the 5' end of the modulator nucleic acid are 2'-O-methoxy-3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0073] In certain embodiments in which the 3' end of the targeter nucleic acid comprises two 2'-O-methoxy-3'-phosphorothioate modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications comprise 1-17, 2-16, 3-15, 4-14, 5-13, 6-12, 7-11, or 8-10, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, phosphorothioate modifications within the first 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) internucleotide linkages of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 8-10, for example 8, 9 or 10, phosphorothioate modifications within the first 10, 9 or 8 internucleotide linkages (as appropriate for total number of nucleotides modified) of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 9 nucleotides of the 5' end of the modulator nucleic acid are 3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0074] In certain embodiments in which the 3' end of the targeter nucleic acid comprises two 2'-O-methoxy-3'-phosphorothioate modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, modifications comprise 1-3, for example, 1, 2, or 3, 2'-O-methoxy and /or 1-3, for example, 1, 2, or 3, 3'-phosphorothioate modifications within the first 3 nucleotide (as appropriate for the total number of nucleotides modified) of the 5' end of the modulator nucleic acid. In this embodiment, any combination of 2'-O-methoxy and 3'-phosphorothioate modifications may be used. In a preferred embodiment, the first nucleotide from the 5' end of the

modulator nucleic acid is 2'-O-methoxy-3'-phosphorothioate modified and the second nucleotide is 3'-phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

5 **[0075]** In certain embodiments in which the 3' end of the targeter nucleic acid comprises two 2'-O-methoxy-3'-phosphorothioate modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the 5' end of the modulator nucleic acid is modified with a terminal propanediol. For each of the described embodiments, the 3' nucleotide of the modulator nucleic
10 acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0076] In certain embodiments in 3' end of the targeter nucleic acid comprises five 2'-fluoro modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides or internucleotide linkages within 15, 14, 13,
15 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides or internucleotide linkages of the 5' end of the modulator (as appropriate for total number of nucleotides modified), wherein the modified nucleotides or internucleotide linkages can have the same modification, different modification, or any combination thereof. In certain embodiments, the modifications comprise 1-5, 1-4, 1-3, 1-2, 2-4, 2-3, for example 1, 2, 3, 4, or 5, 2'-O-methoxy-3'-phosphorothioate modifications within
20 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) nucleotides of the 5' end of the modulator nucleic acid. In certain embodiments, the the modifications comprise 1-3, for example 1, 2, or 3, 2'-O-methoxy-3'-phosphorothioate modifications within the first 3 nucleotides of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 2 nucleotides of the 5' end of the modulator nucleic acid are 2'-O-methoxy-3'-phosphorothioate modified. For
25 each of the described embodiments, the 3' nucleotide of the modulator nucleic acid nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0077] In certain embodiments in 3' end of the targeter nucleic acid comprises five 2'-fluoro modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the modifications
30 comprise 1-17, 2-16, 3-15, 4-14, 5-13, 6-12, 7-11, or 8-10, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17, phosphorothioate modification within the first 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 (as appropriate for total number of nucleotides modified) internucleotide linkages of the 5' end of the modulator nucleic acid. In certain embodiments, the modifications comprise 8-10, for example 8, 9 or 10, phosphorothioate modifications within the
35 first 10, 9 or 8 internucleotide linkages (as appropriate for total number of nucleotides modified)

of the 5' end of the modulator nucleic acid. In a preferred embodiment, the first 9 internucleotide linkages of the 5' end of the modulator nucleic acid are phosphorothioate modified. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

5 **[0078]** In certain embodiments in 3' end of the targeter nucleic acid comprises five 2'-fluoro modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, modifications comprise 1-3, for example, 1, 2, or 3, 2'-O-methoxy and /or 1-3, for example, 1, 2, or 3, 3'-phosphorothioate modifications within the first 3 nucleotides or internucleotide linkages (as appropriate for the
10 total number of nucleotides or internucleotide linkages modified) of the 5' end of the modulator nucleic acid. In this embodiment, any combination of 2'-O-methoxy and phosphorothioate modifications may be used. In a preferred embodiment, the first nucleotide from the 5' end of the modulator nucleic acid is 2'-O-methoxy-3'-phosphorothioate modified and the second internucleotide linkage is phosphorothioate modified. For each of the described embodiments,
15 the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0079] In certain embodiments in 3' end of the targeter nucleic acid comprises five 2'-fluoro modifications, one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid may be modified. In certain embodiments, the 5' end of the
20 modulator nucleic acid is modified with a terminal propanediol. For each of the described embodiments, the 3' nucleotide of the modulator nucleic acid is may be either an A, T, G, C with a preferred terminal nucleotide of either A or C.

[0080] In certain embodiments the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025,
25 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037. In certain embodiments, the targeter nucleic acid comprises a stem sequence complementary to a modulator stem sequence of any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037 and a spacer sequence that may hybridize with a target
30 nucleotide sequence in a target DNA of interest to a skilled artisan. Such a targeter nucleic acid may comprise any composition of modifications as described herein.

[0081] In certain embodiments, modulator and targeter nucleic acids comprise a single polynucleotide wherein the stem sequence of the modulator is in physically connection to the stem sequence of the targeter nucleic acid through a linker, *e.g.*, a loop sequence or a chemical
35 spacer sequence, *e.g.*, a propanediol linker and the like. In certain embodiments, modulator and

target nucleic acids are separate polynucleotides. In either of these embodiments, the gNA may comprise any combination of chemical modification as desired by one skilled in the art.

[0082] In certain embodiments, the modification alters the specificity of the engineered, non-naturally occurring system. In certain embodiments, the modification enhances the specificity of the engineered, non-naturally occurring system, *e.g.*, by enhancing on-target binding and/or cleavage, or reducing off-target binding and/or cleavage, or a combination thereof. Specificity-enhancing modifications include but are not limited to 2-thiouracil, 2-thiocytosine, 4-thiouracil, 6-thioguanine, 2-aminoadenine, and pseudouracil. Within 10, 5, 4, 3, 2, or 1 nucleotide of the 3' end, for example the 3' end nucleotide, is modified

[0083] In certain embodiments, the modification alters the immunostimulatory effect of the RNA relative to a corresponding RNA without the modification. For example, in certain embodiments, the modification reduces the ability of the RNA to activate TLR7, TLR8, TLR9, TLR3, RIG-I, and/or MDA5.

[0084] In certain embodiments, the targeter nucleic acid and/or the modulator nucleic acid comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 modified nucleotides or internucleotide linkages. The modification can be made at one or more positions in the targeter nucleic acid and/or the modulator nucleic acid such that these nucleic acids retain functionality. For example, the modified nucleic acids can still direct the Cas protein to the target nucleotide sequence and allow the Cas protein to exert its effector function. It is understood that the particular modification(s) at a position may be selected based on the functionality of the nucleotide or internucleotide linkage at the position. For example, a specificity-enhancing modification may be suitable for a nucleotide or internucleotide linkage in the spacer sequence, the targeter stem sequence, or the modulator stem sequence. A stability-enhancing modification may be suitable for one or more terminal nucleotides or internucleotide linkages in the targeter nucleic acid and/or the modulator nucleic acid. In certain embodiments, at least 1 (*e.g.*, at least 2, at least 3, at least 4, or at least 5) terminal nucleotides or internucleotide linkages at or near the 5' end and/or at least 1 (*e.g.*, at least 2, at least 3, at least 4, or at least 5) terminal nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid are modified. In certain embodiments, 5 or fewer (*e.g.*, 1 or fewer, 2 or fewer, 3 or fewer, or 4 or fewer) terminal nucleotides or internucleotide linkages at or near the 5' end and/or 5 or fewer (*e.g.*, 1 or fewer, 2 or fewer, 3 or fewer, or 4 or fewer) terminal nucleotides or internucleotide linkages at or near the 3' end of the targeter nucleic acid are modified. In certain embodiments, at least 1 (*e.g.*, at least 2, at least 3, at least 4, or at least 5) terminal nucleotides or internucleotide linkages at or near the 5' end and/or at least 1 (*e.g.*, at least 2, at least 3, at least 4, or at least 5) terminal nucleotides or

internucleotide linkages at or near the 3' end of the modulator nucleic acid are modified. In certain embodiments, 5 or fewer (*e.g.*, 1 or fewer, 2 or fewer, 3 or fewer, or 4 or fewer) terminal nucleotides or internucleotide linkages at or near the 5' end and/or 5 or fewer (*e.g.*, 1 or fewer, 2 or fewer, 3 or fewer, or 4 or fewer) terminal nucleotides or internucleotide linkages at or near the 3' end of the modulator nucleic acid are modified. Selection of positions for modifications is described in U.S. Patent Nos. 10,900,034 and 10,767,175. As used in this paragraph, where the targeter or modulator nucleic acid is a combination of DNA and RNA, the nucleic acid as a whole is considered as an RNA, and the DNA nucleotide(s) are considered as modification(s) of the RNA, including a 2'-H modification of the ribose and optionally a modification of the nucleobase.

[0085] It is understood that, in dual guide nucleic acid systems the targeter nucleic acid and the modulator nucleic acid, while not in the same nucleic acids, *i.e.*, not linked end-to-end through a traditional internucleotide bond, can be covalently conjugated to each other through one or more chemical modifications introduced into these nucleic acids, thereby increasing the stability of the double-stranded complex and/or improving other characteristics of the system.

B. Targeter and Modulator Nucleic Acids

[0086] The engineered, non-naturally occurring systems provided herein comprise a targeter nucleic acid and a modulator nucleic acid, one or both of which contains a modification of one or more nucleotides or internucleotide linkages at or near 3' end, at or near the 5', or at or near both ends, that, when hybridized to form a complex, are capable of activating a Cas nuclease disclosed herein. In certain embodiments, the Cas nuclease is activated by a single crRNA in the absence of a tracrRNA in a naturally occurring system. In certain embodiments, the Cas nuclease is a Type I, II, III, IV, V, or VI nuclease. In certain embodiments, the Cas nuclease is a Type V nuclease. In certain embodiments, the Cas nuclease is a type V-A, type V-C, or type V-D nuclease. In certain embodiments, the Cas nuclease is a Type V-A nuclease.

[0087] The term "targeter nucleic acid," as used herein, includes a nucleic acid comprising (i) a spacer sequence designed to hybridize with a target nucleotide sequence; and (ii) a targeter stem sequence capable of hybridizing with an additional nucleic acid to form a complex, wherein the complex is capable of activating a Cas nuclease (*e.g.*, a type V-A Cas nuclease) under suitable conditions, and wherein the targeter nucleic acid alone, in the absence of the additional nucleic acid, is not capable of activating the Cas nuclease under the same conditions.

[0088] The term "modulator nucleic acid," as used herein in connection with a given targeter nucleic acid and its corresponding Cas nuclease, includes a nucleic acid capable of

hybridizing with the targeter nucleic acid to form a complex, wherein the complex, but not the modulator nucleic acid alone, is capable of activating the type Cas nuclease under suitable conditions.

[0089] The term “suitable conditions,” as used in the definitions of “targeter nucleic acid” and “modulator nucleic acid,” includes the conditions under which a naturally occurring CRISPR-Cas system is operative, such as in a prokaryotic cell, in a eukaryotic (*e.g.*, mammalian or human) cell, or in an *in vitro* assay.

[0090] The targeter nucleic acid and/or the modulator nucleic acid can be synthesized chemically or produced in a biological process (*e.g.*, catalyzed by an RNA polymerase in an *in vitro* reaction). Such reaction or process may limit the lengths of the targeter and modulator nucleic acids. In certain embodiments, the targeter nucleic acid is no more than 100, 90, 80, 70, 60, 50, 40, 30, or 25 nucleotides in length. In certain embodiments, the targeter nucleic acid is at least 20, 25, 30, 40, 50, 60, 70, 80, or 90 nucleotides in length. In certain embodiments, the targeter nucleic acid is 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, 20-30, 20-25, 25-100, 25-90, 25-80, 25-70, 25-60, 25-50, 25-40, 25-30, 30-100, 30-90, 30-80, 30-70, 30-60, 30-50, 30-40, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-100, 50-90, 50-80, 50-70, 50-60, 60-100, 60-90, 60-80, 60-70, 70-100, 70-90, 70-80, 80-100, 80-90, or 90-100 nucleotides in length. In certain embodiments, the modulator nucleic acid is no more than 100, 90, 80, 70, 60, 50, 40, 30, or 20 nucleotides in length. In certain embodiments, the modulator nucleic acid is at least 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, or 90 nucleotides in length. In certain embodiments, the modulator nucleic acid is 10-100, 10-90, 10-80, 10-70, 10-60, 10-50, 10-40, 10-30, 10-20, 15-100, 15-90, 15-80, 15-70, 15-60, 15-50, 15-40, 15-30, 15-20, 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, 20-30, 25-100, 25-90, 25-80, 25-70, 25-60, 25-50, 25-40, 25-30, 30-100, 30-90, 30-80, 30-70, 30-60, 30-50, 30-40, 40-100, 40-90, 40-80, 40-70, 40-60, 40-50, 50-100, 50-90, 50-80, 50-70, 50-60, 60-100, 60-90, 60-80, 60-70, 70-100, 70-90, 70-80, 80-100, 80-90, or 90-100 nucleotides in length.

[0091] In naturally occurring type V-A CRISPR-Cas systems, the crRNA comprises a scaffold sequence (also called direct repeat sequence) and a spacer sequence that hybridizes with the target nucleotide sequence. In certain naturally occurring type V-A CRISPR-Cas systems, the scaffold sequence forms a stem-loop structure in which the stem consists of five consecutive base pairs. A dual guide type V-A CRISPR-Cas system may be derived from a naturally occurring type V-A CRISPR-Cas system, or a variant thereof in which the Cas protein is guided to the target nucleotide sequence by a crRNA alone, such system referred to herein as a “single guide type V-A CRISPR-Cas system.” In certain modified dual guide type V-A CRISPR-Cas systems disclosed herein, the targeter nucleic acid comprises the chain of the stem sequence

between the spacer and the loop (the “targeter stem sequence”) and the spacer sequence, and the modulator nucleic acid comprises the other chain of the stem sequence (the “modulator stem sequence”) and the 5’ sequence, *e.g.*, a tail sequence, positioned 5’ to the modulator stem sequence. The targeter stem sequence is 100% complementary to the modulator stem sequence.

- 5 As such, the double-stranded complex of the targeter nucleic acid and the modulator nucleic acid retains the orientation of the 5’ sequence, *e.g.*, a tail sequence, the modulator stem sequence, the targeter stem sequence, and the spacer sequence of a single guide type V-A CRISPR-Cas system but lacks the loop structure between the modulator stem sequence and the targeter stem sequence. A schematic representation of an exemplary double-stranded complex is shown in
- 10 **Figure 1.**

- [0092] Notwithstanding the general structural similarity, it has been discovered that the stem-loop structure of the crRNA in a naturally occurring type V-A CRISPR complex is dispensable for the functionality of the CRISPR system. This discovery is surprising because the prior art has suggested that the stem-loop structure is critical (see, Zetsche *et al.* (2015) CELL, 163: 759) and that removal of the loop structure by “splitting” the crRNA abrogated the activity
- 15 of a AsCpf1 CRISPR system (see, Li *et al.* (2017) NAT. BIOMED. ENG., 1: 0066).

- [0093] It is contemplated that the length of the duplex may be a factor in providing an operative modified dual guide CRISPR system. In certain embodiments, the targeter stem sequence and the modulator stem sequence each consist of 4-10 nucleotides that base pair with each other. In certain embodiments, the targeter stem sequence and the modulator stem sequence each consist of 4-9, 4-8, 4-7, 4-6, 4-5, 5-10, 5-9, 5-8, 5-7, or 5-6 nucleotides that base pair with each other. In certain embodiments, the targeter stem sequence and the modulator stem sequence each consist of 4, 5, 6, 7, 8, 9, or 10 nucleotides. It is understood that the composition of the nucleotides in each sequence affects the stability of the duplex, and a C-G base pair confers
- 25 greater stability than an A-U base pair. In certain embodiments, 20%-80%, 20%-70%, 20%-60%, 20%-50%, 20%-40%, 20%-30%, 30%-80%, 30%-70%, 30%-60%, 30%-50%, 30%-40%, 40%-80%, 40%-70%, 40%-60%, 40%-50%, 50%-80%, 50%-70%, 50%-60%, 60%-80%, 60%-70%, or 70%-80% of the base pairs are C-G base pairs. In certain embodiments, the targeter stem sequence and the modulator stem share at least 80%, 85%, 90%, 95%, 99%, 99.5%, or 100%
- 30 sequence complementarity. In a preferred embodiment, the target stem sequence and the modulator stem sequence share at 80-100% sequence complementarity.

- [0094] In certain embodiments, the targeter stem sequence and the modulator stem sequence each consist of 5 nucleotides. As such, the targeter stem sequence and the modulator stem sequence form a duplex of 5 base pairs. In certain embodiments, 0-4, 0-3, 0-2, 0-1, 1-5, 1-4, 1-3, 1-2, 2-5, 2-4, 2-3, 3-5, 3-4, or 4-5 out of the 5 base pairs are C-G base pairs. In certain
- 35

embodiments, 0, 1, 2, 3, 4, or 5 out of the 5 base pairs are C-G base pairs. In certain embodiments, the targeter stem sequence consists of 5'-GUAGA-3' (SEQ ID NO: 21) and the modulator stem sequence consists of 5'-UCUAC-3'. In certain embodiments, the targeter stem sequence consists of 5'-GUGGG-3' (SEQ ID NO: 22) and the modulator stem sequence consists of 5'-CCCAC-3'.

[0095] It is also contemplated that the compatibility of the duplex for a given Cas nuclease may be a factor in providing an operative modified dual guide CRISPR system. For example, the targeter stem sequence and the modulator stem sequence can be derived from a naturally occurring crRNA capable of activating a Cas nuclease in the absence of a tracrRNA. In certain embodiments, the nucleotide sequences of the targeter stem sequence and the modulator stem sequence are identical to the corresponding stem sequences of a stem-loop structure in such naturally occurring crRNA.

[0096] In certain embodiments, the targeter nucleic acid comprises, from 5' to 3', a targeter stem sequence and a spacer sequence. The spacer sequence is designed to hybridize with the target nucleotide sequence. To provide sufficient targeting to the target nucleotide sequence, the spacer sequence is generally 16 or more nucleotides in length. In certain embodiments, the spacer sequence is at least 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides in length. In certain embodiments, the spacer sequence is shorter than or equal to 75, 50, 45, 40, 35, 30, 25, or 20 nucleotides in length. Shorter spacer sequence may be desirable for reducing off-target events. Accordingly, in certain embodiments, the spacer sequence is shorter than or equal to 19, 18, or 17 nucleotides. In certain embodiments, the spacer sequence is 17-30 nucleotides in length, *e.g.*, 20-30 nucleotides, 20-25 nucleotides, 20-24 nucleotides, 20-23 nucleotides, 23-25 nucleotides, 20-22 nucleotides, such as 20 or 21 nucleotides in length. In certain embodiments, the spacer sequence is 21 nucleotides in length. In certain embodiments, the spacer sequence is 20 nucleotides in length. In certain embodiments, the spacer sequence is at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% complementary to the target nucleotide sequence. In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence in the seed region (5-10 base pairs proximal to the PAM). In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence. It has been reported that compared to DNA binding, DNA cleavage is less tolerant to mismatches between the spacer sequence and the target nucleotide sequence (see, Klein *et al.* (2018) CELL REPORTS, 22: 1413). Accordingly, in specific embodiments, when the engineered, non-naturally occurring system comprises a Cas nuclease, the spacer sequence is 100% complementary to the target nucleotide sequence.

[0097] Proper design of the spacer sequence is dependent upon the selection of target nucleotide sequence. For example, to select a target nucleotide sequence in a specific gene in a given genome, sequence analysis can be conducted to minimize potential hybridization of the spacer sequence with any other loci in the genome. The association of the target nucleotide sequence with a PAM recognized by the Cas protein is also considered by many design methods. In a type V-A CRISPR-Cas system, the PAM is immediately upstream from the target nucleotide sequence when using the non-target strand (*i.e.*, the strand not hybridized with the spacer sequence) as the coordinate. Computational models have been developed to assess the targetability of the target nucleotide sequence as well as any potential off-target effect, for example, as disclosed in Doench *et al.* (2016) NAT. BIOTECHNOL., 34: 184; Chuai *et al.* (2018) GENOME BIOLOGY, 19: 80; and Klein *et al.* (2018) CELL REPORTS, 22: 1413. Although computational methods are useful for selection of spacer sequences, it is generally advisable to design multiple spacer sequences and select one or more with high efficiency and specificity based upon the results of *in vitro* and/or *in vivo* experiments.

[0098] In certain embodiments, the engineered CRISPR-Cas systems (*e.g.*, type V-A CRISPR-Cas systems) provided can be used to target, edit, or otherwise modify specific target nucleotide sequences in human ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALPNR, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTCD2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWFI1, CD3D, or NLRC5 gene. In particular, the modified guide nucleic acids, such as single guide nucleic acids and dual guide nucleic acids, can be designed to hybridize with the selected target nucleotide sequence and, *e.g.*, activate a Cas nuclease to edit the human genes. CRISPR-Cas systems comprising such guide nucleic acids are also useful for targeting or modifying the human genes.

[0099] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 51 and 131-137, wherein the spacer sequence is capable of hybridizing with the human ADORA2A gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the ADORA2A gene locus is edited in at least 1.5% of the cells.

[0100] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID

NOs: 52, 64-66, 138-145, 622, 625-626, and 634-635, wherein the spacer sequence is capable of hybridizing with the human B2M gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the B2M gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*,

5 modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human ADORA2A gene, for example one of the spacer sequences above.

[0101] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
10 NOs: 724, 726-727, 730-732, 735-738, 741-742, and 744-745, wherein the spacer sequence is capable of hybridizing with the human CD247 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD247 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA,
15 for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human CD247 gene, for example one of the spacer sequences above.

[0102] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
20 NOs: 53 and 146, wherein the spacer sequence is capable of hybridizing with the human CD52 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD52 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as
25 described in section IA1., and a spacer sequence capable of hybridizing with the human CD52 gene, for example one of the spacer sequences above.

[0103] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
30 NOs: 54, 147-148, 636-640, 642, 644-648, 650-652, 655-656, 660-663, 666, 668, 670-671, 673-676, 678-679, and 682-685, wherein the spacer sequence is capable of hybridizing with the human CIITA gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CIITA gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more

modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human CIITA gene, for example one of the spacer sequences above.

[0104] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
5 NOs: 795, 67, 797, 798, 70, and 149-155, wherein the spacer sequence is capable of hybridizing with the human CTLA4 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CTLA4 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified
10 guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human CTLA4 gene, for example one of the spacer sequences above.

[0105] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
15 NOs: 796, 71-74, and 156-159, wherein the spacer sequence is capable of hybridizing with the human DCK gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the DCK gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more
20 modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human DCK gene, for example one of the spacer sequences above.

[0106] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
NOs: 57, 75-79, and 160-173, wherein the spacer sequence is capable of hybridizing with the human FAS gene. In certain embodiments, when the system is delivered into a population of
25 human cells *ex vivo*, the genomic sequence at the FAS gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human FAS gene, for example one of the spacer sequences above.

[0107] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
30 NOs: 58, 80, 799, 800, 83-86, and 174-187, wherein the spacer sequence is capable of hybridizing with the human HAVCR2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the HAVCR2 gene
35 locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic

acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human HAVCR2 gene, for example one of the spacer sequences above.

5 **[0108]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 748-749 and 753-754, wherein the spacer sequence is capable of hybridizing with the human IL7R gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the IL7R gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human IL7R gene, for example one of the spacer sequences above.

15 **[0109]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 59, 87, 88, and 188-198, wherein the spacer sequence is capable of hybridizing with the human LAG3 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the LAG3 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human LAG3 gene, for example one of the spacer sequences above.

25 **[0110]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises the nucleotide sequence of SEQ ID NO: 757, wherein the spacer sequence is capable of hybridizing with the human LCK gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the LCK gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human LCK gene, for example one of the spacer sequences above.

35 **[0111]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 60, 89-92, and 199-201, wherein the spacer sequence is capable of hybridizing with the human PDCD1 gene. In certain embodiments, when the system is delivered into a population of

human cells *ex vivo*, the genomic sequence at the PDCD1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PDCD1 gene, for example one of the spacer sequences above.

[0112] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 759 and 761-762, wherein the spacer sequence is capable of hybridizing with the human PLCG1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PLCG1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PLCG1 gene, for example one of the spacer sequences above.

[0113] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 61, 93, 801, 802, 96-101, 803, 103, 104, and 202-213, wherein the spacer sequence is capable of hybridizing with the human PTPN6 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PTPN6 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PTPN6 gene, for example one of the spacer sequences above.

[0114] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 62, 105, and 214-217, wherein the spacer sequence is capable of hybridizing with the human TIGIT gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TIGIT gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human TIGIT gene, for example one of the spacer sequences above.

[0115] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID

NOs: 63, 106, 804, 805, 109-130, and 218-241, wherein the spacer sequence is capable of hybridizing with the human TRAC gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TRAC gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*,

5 modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human TRAC gene, for example one of the spacer sequences above.

[0116] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID

10 NOs: 705-706, 711-712, 714-715, 717, and 719-720, wherein the spacer sequence is capable of hybridizing with the human TRBC2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TRBC2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human TRBC2 gene, for example one of the spacer sequences above.

[0117] In certain embodiments, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 705-706, wherein the spacer sequence is capable of hybridizing with both the human TRBC1 gene and the human TRBC2 gene. In certain
20 embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TRBC1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human TRBC1 gene and the
25 human TRBC2 gene, for example one of the spacer sequences above.

[0118] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 721-723, wherein the spacer sequence is capable of hybridizing with the human CARD11 gene. In certain embodiments, when the system is delivered into a population of human cells *ex*
30 *vivo*, the genomic sequence at the CARD11 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human CARD11 gene, for example one of the spacer sequences above.

- [0119] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1255-1291, wherein the spacer sequence is capable of hybridizing with the human CD38 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD38 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human CD38 gene, for example one of the spacer sequences above.
- 10 [0120] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1213-1254, wherein the spacer sequence is capable of hybridizing with the human CD3E gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD3E gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human CD3E gene, for example one of the spacer sequences above.
- 15 [0121] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1153-1212, wherein the spacer sequence is capable of hybridizing with the human CD40LG gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD40LG gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human CD40LG gene, for example one of the spacer sequences above.
- 20 [0122] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1100-1152, wherein the spacer sequence is capable of hybridizing with the human CSF2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CSF2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as
- 25 30

described in section IA 1., and a spacer sequence capable of hybridizing with the human CSF2 gene, for example one of the spacer sequences above.

[0123] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
5 NOs: 1292-1301, wherein the spacer sequence is capable of hybridizing with the human APLNR gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the APLNR gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as
10 described in section IA 1., and a spacer sequence capable of hybridizing with the human APLNR gene, for example one of the spacer sequences above.

[0124] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
15 NOs: 1302-1311, wherein the spacer sequence is capable of hybridizing with the human BBS1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the BBS1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as
described in section IA 1., and a spacer sequence capable of hybridizing with the human BBS1
20 gene, for example one of the spacer sequences above.

[0125] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
NOs: 1312-1321, wherein the spacer sequence is capable of hybridizing with the human CALR gene. In certain embodiments, when the system is delivered into a population of human cells *ex*
25 *vivo*, the genomic sequence at the CALR gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as
described in section IA 1., and a spacer sequence capable of hybridizing with the human CALR gene, for example one of the spacer sequences above.

[0126] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID
NOs: 1322-1331, wherein the spacer sequence is capable of hybridizing with the human CD3G gene. In certain embodiments, when the system is delivered into a population of human cells *ex*
35 *vivo*, the genomic sequence at the CD3G gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or

more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human CD3G gene, for example one of the spacer sequences above.

[0127] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1332-1341, wherein the spacer sequence is capable of hybridizing with the human CD58 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD58 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human CD58 gene, for example one of the spacer sequences above.

[0128] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1342-1351, wherein the spacer sequence is capable of hybridizing with the human COL17A1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the COL17A1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human COL17A1 gene, for example one of the spacer sequences above.

[0129] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1352-1361, wherein the spacer sequence is capable of hybridizing with the human DEFB134 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the DEFB134 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human DEFB134 gene, for example one of the spacer sequences above.

[0130] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1362-1371, wherein the spacer sequence is capable of hybridizing with the human ERAP1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the ERAP1 gene locus is edited in at least 1.5% of the cells. In

certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human ERAP1 gene, for example one of the spacer sequences above.

5 **[0131]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1372-1381, wherein the spacer sequence is capable of hybridizing with the human ERAP2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the ERAP2 gene locus is edited in at least 1.5% of the cells. In

10 certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human ERAP2 gene, for example one of the spacer sequences above.

15 **[0132]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1382-1391, wherein the spacer sequence is capable of hybridizing with the human IFNGR1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the IFNGR1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA,

20 comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human IFNGR1 gene, for example one of the spacer sequences above.

25 **[0133]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1392-1401, wherein the spacer sequence is capable of hybridizing with the human IFNGR2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the IFNGR2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA,

30 comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human IFNGR2 gene, for example one of the spacer sequences above.

35 **[0134]** In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1402-1411, wherein the spacer sequence is capable of hybridizing with the human JAK1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex*

vivo, the genomic sequence at the JAK1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human JAK1 gene, for example one of the spacer sequences above.

5 [0135] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1412-1421, wherein the spacer sequence is capable of hybridizing with the human JAK2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the JAK2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human JAK2 gene, for example one of the spacer sequences above.

15 [0136] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1422-1424, wherein the spacer sequence is capable of hybridizing with the human mir-101-2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the mir-101-2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human mir-101-2 gene, for example one of the spacer sequences above.

25 [0137] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1425-1434, wherein the spacer sequence is capable of hybridizing with the human MLANA gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the MLANA gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human MLANA gene, for example one of the spacer sequences above.

30 [0138] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1435-1444, wherein the spacer sequence is capable of hybridizing with the human PSMB5

gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PSMB5 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PSMB5 gene, for example one of the spacer sequences above.

[0139] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1445-1454, wherein the spacer sequence is capable of hybridizing with the human PSMB8 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PSMB8 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PSMB8 gene, for example one of the spacer sequences above.

[0140] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1455-1464, wherein the spacer sequence is capable of hybridizing with the human PSMB9 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PSMB9 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PSMB9 gene, for example one of the spacer sequences above.

[0141] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1465-1474, wherein the spacer sequence is capable of hybridizing with the human PTCD2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the PTCD2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human PTCD2 gene, for example one of the spacer sequences above.

[0142] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID

NOs: 1475-1484, wherein the spacer sequence is capable of hybridizing with the human RFX5 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the RFX5 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human RFX5 gene, for example one of the spacer sequences above.

[0143] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID

NOs: 1485-1494, wherein the spacer sequence is capable of hybridizing with the human RFXANK gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the RFXANK gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human RFXANK gene, for example one of the spacer sequences above.

[0144] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1495-1504, wherein the spacer sequence is capable of hybridizing with the human RFXAP gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the RFXAP gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human RFXAP gene, for example one of the spacer sequences above.

[0145] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1505-1514, wherein the spacer sequence is capable of hybridizing with the human RPL23 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the RPL23 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human RPL23 gene, for example one of the spacer sequences above.

[0146] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1515-1520, wherein the spacer sequence is capable of hybridizing with the human SOX10 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the SOX10 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human SOX10 gene, for example one of the spacer sequences above.

[0147] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1521-1531, wherein the spacer sequence is capable of hybridizing with the human SRP54 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the SRP54 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human SRP54 gene, for example one of the spacer sequences above.

[0148] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1532-1541, wherein the spacer sequence is capable of hybridizing with the human STAT1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the STAT1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human STAT1 gene, for example one of the spacer sequences above.

[0149] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1542-1551, wherein the spacer sequence is capable of hybridizing with the human Tap1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the Tap1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as

described in section IA 1., and a spacer sequence capable of hybridizing with the human Tap1 gene, for example one of the spacer sequences above.

[0150] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1552-1561, wherein the spacer sequence is capable of hybridizing with the human TAP2 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TAP2 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human TAP2 gene, for example one of the spacer sequences above.

[0151] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1562-1571, wherein the spacer sequence is capable of hybridizing with the human TAPBP gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TAPBP gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human TAPBP gene, for example one of the spacer sequences above.

[0152] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1572-1581, wherein the spacer sequence is capable of hybridizing with the human TWF1 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the TWF1 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA 1., and a spacer sequence capable of hybridizing with the human TWF1 gene, for example one of the spacer sequences above.

[0153] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1582-1591, wherein the spacer sequence is capable of hybridizing with the human CD3D gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the CD3D gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or

more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human CD3D gene, for example one of the spacer sequences above.

[0154] In certain embodiments of the engineered, non-naturally occurring system, the spacer sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1592-1610, wherein the spacer sequence is capable of hybridizing with the human NLRC5 gene. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the NLRC5 gene locus is edited in at least 1.5% of the cells. In certain embodiments, a modified guide nucleic acid, *e.g.*, modified guide RNA, comprises one or more modifications as described in section IA, for example, one or more modifications as described in section IA1., and a spacer sequence capable of hybridizing with the human NLRC5 gene, for example one of the spacer sequences above.

[0155] In certain embodiments of the engineered, non-naturally occurring system, the modulator-targeter complex comprises any one of SEQ ID NOs: 1004 and 1012, 1004 and 1013, 1004 and 1014, 1004 and 1015, 1004 and 1016, 1004 and 1017, 1004 and 1018, 1004 and 1019, 1004 and 1020, 1004 and 1021, 1004 and 1022, 1004 and 1023, 1004 and 1024, 1004 and 1025, 1004 and 1026, 1004 and 1027, 1004 and 1028, 1004 and 1029, 1004 and 1030, 1004 and 1031, 1004 and 1032, 1004 and 1033, 1004 and 1034, 1004 and 1035, 1004 and 1036, 1004 and 1037, 1005 and 1012, 1005 and 1013, 1005 and 1014, 1005 and 1015, 1005 and 1016, 1005 and 1017, 1005 and 1018, 1005 and 1019, 1005 and 1020, 1005 and 1021, 1005 and 1022, 1005 and 1023, 1005 and 1024, 1005 and 1025, 1005 and 1026, 1005 and 1027, 1005 and 1028, 1005 and 1029, 1005 and 1030, 1005 and 1031, 1005 and 1032, 1005 and 1033, 1005 and 1034, 1005 and 1035, 1005 and 1036, 1005 and 1037, 1006 and 1012, 1006 and 1013, 1006 and 1014, 1006 and 1015, 1006 and 1016, 1006 and 1017, 1006 and 1018, 1006 and 1019, 1006 and 1020, 1006 and 1021, 1006 and 1022, 1006 and 1023, 1006 and 1024, 1006 and 1025, 1006 and 1026, 1006 and 1027, 1006 and 1028, 1006 and 1029, 1006 and 1030, 1006 and 1031, 1006 and 1032, 1006 and 1033, 1006 and 1034, 1006 and 1035, 1006 and 1036, 1006 and 1037, 1007 and 1012, 1007 and 1013, 1007 and 1014, 1007 and 1015, 1007 and 1016, 1007 and 1017, 1007 and 1018, 1007 and 1019, 1007 and 1020, 1007 and 1021, 1007 and 1022, 1007 and 1023, 1007 and 1024, 1007 and 1025, 1007 and 1026, 1007 and 1027, 1007 and 1028, 1007 and 1029, 1007 and 1030, 1007 and 1031, 1007 and 1032, 1007 and 1033, 1007 and 1034, 1007 and 1035, 1007 and 1036, or 1007 and 1037 wherein the spacer sequence of the complex is capable of hybridizing with a target nucleotide sequence within a first human gene as referred to herein as “gene 1”. Exemplary data for these complexes may be found in Example 2, **Figure 4**. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 1

locus is edited at least 70% as efficiently as a single crRNA such as with modulator-targeter complexes comprising any one of SEQ ID NOs: 1005 and 1014, 1005 and 1027, 1006 and 1014, or 1006 and 1027. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 1 locus is edited at least as efficiently as a single crRNA. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 1 locus is edited more efficiently than a single crRNA.

[0156] In certain embodiments of the engineered, non-naturally occurring system, the modulator-targeter complex comprises any one of SEQ ID NOs: 1000 and 1012, 1000 and 1013, 1000 and 1014, 1000 and 1015, 1000 and 1016, 1000 and 1017, 1000 and 1018, 1000 and 1019, 1000 and 1020, 1000 and 1021, 1000 and 1022, 1000 and 1023, 1000 and 1024, 1000 and 1025, 1000 and 1026, 1000 and 1027, 1000 and 1028, 1000 and 1029, 1000 and 1030, 1000 and 1031, 1000 and 1032, 1000 and 1033, 1000 and 1034, 1000 and 1035, 1000 and 1036, 1000 and 1037, 1001 and 1012, 1001 and 1013, 1001 and 1014, 1001 and 1015, 1001 and 1016, 1001 and 1017, 1001 and 1018, 1001 and 1019, 1001 and 1020, 1001 and 1021, 1001 and 1022, 1001 and 1023, 1001 and 1024, 1001 and 1025, 1001 and 1026, 1001 and 1027, 1001 and 1028, 1001 and 1029, 1001 and 1030, 1001 and 1031, 1001 and 1032, 1001 and 1033, 1001 and 1034, 1001 and 1035, 1001 and 1036, 1001 and 1037, 1002 and 1012, 1002 and 1013, 1002 and 1014, 1002 and 1015, 1002 and 1016, 1002 and 1017, 1002 and 1018, 1002 and 1019, 1002 and 1020, 1002 and 1021, 1002 and 1022, 1002 and 1023, 1002 and 1024, 1002 and 1025, 1002 and 1026, 1002 and 1027, 1002 and 1028, 1002 and 1029, 1002 and 1030, 1002 and 1031, 1002 and 1032, 1002 and 1033, 1002 and 1034, 1002 and 1035, 1002 and 1036, 1002 and 1037, 1003 and 1012, 1003 and 1013, 1003 and 1014, 1003 and 1015, 1003 and 1016, 1003 and 1017, 1003 and 1018, 1003 and 1019, 1003 and 1020, 1003 and 1021, 1003 and 1022, 1003 and 1023, 1003 and 1024, 1003 and 1025, 1003 and 1026, 1003 and 1027, 1003 and 1028, 1003 and 1029, 1003 and 1030, 1003 and 1031, 1003 and 1032, 1003 and 1033, 1003 and 1034, 1003 and 1035, 1003 and 1036, or 1003 and 1037 wherein the spacer sequence of the complex is capable of hybridizing with a target nucleotide sequence within a second human gene as referred to herein as “gene 2”. Exemplary data for these complexes may be found in Example 2, **Figure 5**. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 2 locus is edited at least 70% as efficiently as a single crRNA such as with modulator-targeter complexes comprising any one of SEQ ID NOs: 1000 and 1012, 1000 and 1012, 1000 and 1013, 1000 and 1016, 1000 and 1017, 1000 and 1018, 1000 and 1019, 1000 and 1020, 1000 and 1025, 1000 and 1025, 1000 and 1025, 1000 and 1025, 1000 and 1026, 1000 and 1031, 1000 and 1032, 1000 and 1033, 1001 and 1012, 1001 and 1012, 1001 and 1018, 1001 and 1019, 1001

and 1025, 1001 and 1025, 1001 and 1026, 1001 and 1032, 1001 and 1033, 1003 and 1013, 1003 and 1014, 1003 and 1025, 1003 and 1025, 1003 and 1025, 1003 and 1027, 1003 and 1032, 1000 and 1014, 1000 and 1027, 1001 and 1014, or 1001 and 1027. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 2 locus is edited at least as efficiently as a single crRNA such as with modulator-targeter complexes comprising any one of SEQ ID NOs: 1000 and 1014, 1000 and 1027, 1001 and 1014, or 1001 and 1027. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 2 locus is edited more efficiently than a single crRNA such as with modulator-targeter complexes comprising SEQ ID NOs: 1000 and 1014, 1000 and 1027, 1001 and 1014, or 1001 and 1027.

[0157] In certain embodiments of the engineered, non-naturally occurring system, the modulator-targeter complex comprises any one of SEQ ID NOs: 1008 and 1012, 1008 and 1013, 1008 and 1014, 1008 and 1015, 1008 and 1016, 1008 and 1017, 1008 and 1018, 1008 and 1019, 1008 and 1020, 1008 and 1021, 1008 and 1022, 1008 and 1023, 1008 and 1024, 1008 and 1025, 1008 and 1026, 1008 and 1027, 1008 and 1028, 1008 and 1029, 1008 and 1030, 1008 and 1031, 1008 and 1032, 1008 and 1033, 1008 and 1034, 1008 and 1035, 1008 and 1036, 1008 and 1037, 1009 and 1012, 1009 and 1013, 1009 and 1014, 1009 and 1015, 1009 and 1016, 1009 and 1017, 1009 and 1018, 1009 and 1019, 1009 and 1020, 1009 and 1021, 1009 and 1022, 1009 and 1023, 1009 and 1024, 1009 and 1025, 1009 and 1026, 1009 and 1027, 1009 and 1028, 1009 and 1029, 1009 and 1030, 1009 and 1031, 1009 and 1032, 1009 and 1033, 1009 and 1034, 1009 and 1035, 1009 and 1036, 1009 and 1037, 1010 and 1012, 1010 and 1013, 1010 and 1014, 1010 and 1015, 1010 and 1016, 1010 and 1017, 1010 and 1018, 1010 and 1019, 1010 and 1020, 1010 and 1021, 1010 and 1022, 1010 and 1023, 1010 and 1024, 1010 and 1025, 1010 and 1026, 1010 and 1027, 1010 and 1028, 1010 and 1029, 1010 and 1030, 1010 and 1031, 1010 and 1032, 1010 and 1033, 1010 and 1034, 1010 and 1035, 1010 and 1036, 1010 and 1037, 1011 and 1012, 1011 and 1013, 1011 and 1014, 1011 and 1015, 1011 and 1016, 1011 and 1017, 1011 and 1018, 1011 and 1019, 1011 and 1020, 1011 and 1021, 1011 and 1022, 1011 and 1023, 1011 and 1024, 1011 and 1025, 1011 and 1026, 1011 and 1027, 1011 and 1028, 1011 and 1029, 1011 and 1030, 1011 and 1031, 1011 and 1032, 1011 and 1033, 1011 and 1034, 1011 and 1035, 1011 and 1036, or 1011 and 1037 wherein the spacer sequence of the complex is capable of hybridizing with a target nucleotide sequence within a third human gene as referred to herein as “gene 3”. Exemplary data for these complexes may be found in Example 2, **Figure 6**. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 3 locus is edited at least 70% as efficiently as a single crRNA such as with modulator-targeter complexes comprising any one of SEQ ID NOs: 1008 and 1012, 1008 and 1013, 1008 and 1014,

1008 and 1016, 1008 and 1017, 1008 and 1023, 1008 and 1024, 1008 and 1026, 1008 and 1027, 1008 and 1029, 1008 and 1030, 1008 and 1032, 1008 and 1033, 1008 and 1034, 1008 and 1035, 1008 and 1037, 1009 and 1012, 1009 and 1013, 1009 and 1014, 1009 and 1016, 1009 and 1017, 1009 and 1018, 1009 and 1019, 1009 and 1020, 1009 and 1021, 1009 and 1022, 1009 and 1023, 5 1009 and 1024, 1010 and 1012, 1010 and 1013, 1010 and 1014, 1010 and 1016, 1010 and 1017, 1010 and 1018, 1010 and 1020, 1010 and 1021, 1010 and 1022, 1010 and 1023, 1010 and 1024, 1010 and 1027, 1010 and 1030, 1008 and 1021, 1008 and 1031, 1008 and 1018, 1008 and 1019, 1008 and 1020, 1008 and 1022, or 1010 and 1019. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 3 locus is edited at least as efficiently as a single crRNA such as with modulator-targeter complexes 10 comprising any one of SEQ ID NOs: 1008 and 1021, 1008 and 1031, 1008 and 1018, 1008 and 1019, 1008 and 1020, 1008 and 1022, or 1010 and 1019. In certain embodiments, when the system is delivered into a population of human cells *ex vivo*, the genomic sequence at the gene 3 locus is edited more efficiently than a single crRNA such as with modulator-targeter complexes 15 comprising any one of SEQ ID NOs: 1008 and 1018, 1008 and 1019, 1008 and 1020, 1008 and 1022, or 1010 and 1019.

[0158] In certain embodiments of the engineered, non-naturally occurring system, genomic mutations are detected in no more than 2% of the cells at any off-target loci by CIRCLE-Seq. In certain embodiments, genomic mutations are detected in no more than 1% of 20 the cells at any off-target loci by CIRCLE-Seq.

[0159] In certain embodiments, provided is a guide nucleic acid a modified guide nucleic acid, *e.g.*, modified guide RNA, comprising one or more modifications as described in section IA, for example, one or more modifications as described in section IA1, and comprising a targeter stem sequence and a spacer sequence, wherein the spacer sequence comprises a 25 nucleotide sequence listed **Table 1**, **2**, or **3**, or a portion thereof sufficient to hybridize with the corresponding target gene listed in the table. In particular, **Table 1** lists the guide nucleic acid that showed the best editing efficiency for each target gene. **Table 2** lists the guide nucleic acids that showed at least 10% editing efficiency. **Table 3** lists the guide nucleic acids that showed at least 1.5% and lower than 10% editing efficiency.

[0160] In certain embodiments, a guide nucleic acid of the present invention is capable of binding the genomic locus of the corresponding target gene in the human genome. In certain 30 embodiments, a guide nucleic acid of the present invention, alone or in combination with a modulator nucleic acid, is capable of directing a Cas protein to the genomic locus of the corresponding target gene in the human genome. In certain embodiments, a guide nucleic acid of the present invention, alone or in combination with a modulator nucleic acid, is capable of 35

directing a Cas nuclease to the genomic locus of the corresponding target gene in the human genome, thereby resulting in cleavage of the genomic DNA at the genomic locus.

Table 1 Selected Spacer Sequences Targeting Human Genes

Target Gene	crRNA	Spacer Sequence	SEQ ID NO
ADORA2A	gADORA2A_12	AGGATGTGGTCCCCATGAACT	51
B2M	gB2M_41	ATAGATCGAGACATGTAAGCA	635
CARD11	gCARD11_1	TAGTACCGCTCCTGGAAGGTT	721
CD247	gCD247_12	CTAGCAGAGAAGGAAGAACCC	735
CD52	gCD52_1	CTCTTCCTCCTACTCACCATC	53
CIITA	gCIITA_32	CCTTGGGGCTCTGACAGGTAG	636
CTLA4	gCTLA4_4	AGCGGCACAAGGCTCAGCTGA	795
DCK	gDCK_6	CGGAGGCTCCTTACCGATGTT	796
FAS	gFAS_36	GTGTTGCTGGTGAGTGTGCAT	57
HAVCR2	gTIM3_6	CTTGTAAGTAGTAGCAGCAGC	58
IL7R	gIL7R_3	CAGGGGAGATGGATCCTATCT	749
LAG3	gLAG3_6	GGGTGCATACCTGTCTGGCTG	59
LCK	gLCK1_3	ACCCATCAACCCGTAGGGATG	757
PDCD1	gPD_23	TCTGCAGGGACAATAGGAGCC	60
PLCG1	gPLCG1_2	CCTTTCTGCGCTTCGTGGTGT	759
PTPN6	gPTPN6_6	TATGACCTGTATGGAGGGGAG	61
TIGIT	gTIGIT_2	AGGCCTTACCTGAGGCGAGGG	62
TRAC	gTRAC006	TGAGGGTGAAGGATAGACGCT	63
TRBC1+2	gTRBC1+2_3	CGCTGTCAAGTCCAGTTCTAC	706
TRBC2	gTRBC2_12	CCGAGGTTGAAGCCACAGTCT	712

5

Table 2 Selected Spacer Sequences Targeting Human Genes

Target Gene	crRNA	Spacer sequence	SEQ ID NO
ADORA2A	gADORA2A_12	AGGATGTGGTCCCCATGAACT	51
B2M	gB2M_4	CTCACGTCATCCAGCAGAGAA	52
B2M	gB2M_7	ACTTTCCATTCTCTGCTGGAT	64
B2M	gB2M_2	TGGCCTGGAGGCTATCCAGCG	65
B2M	gB2M_17	TATCTCTTGTACTACACTGAA	66
B2M	gB2M_30	AGTGGGGTGAATTCAGTGTA	625
B2M	gB2M_41	ATAGATCGAGACATGTAAGCA	635
CIITA	gCIITA_32	CCTTGGGGCTCTGACAGGTAG	636
CIITA	gCIITA_33	ACCTTGGGGCTCTGACAGGTA	637
CIITA	gCIITA_35	CTCCCAGAACCCGACACAGAC	639
CIITA	gCIITA_36	TGGGCTCAGGTGCTTCCTCAC	640
CIITA	gCIITA_38	CTTGTCTGGGCAGCGGAACTG	642
CIITA	gCIITA_40	TCAAAGTAGAGCACATAGGAC	644
CIITA	gCIITA_41	TGCCCCAACTTCTGCTGGCATC	645
CIITA	gCIITA_43	TCTGCAGCCTTCCCAGAGGAG	647
CIITA	gCIITA_44	TCCAGGCGCATCTGGCCGGAG	648
CIITA	gCIITA_48	CTCGGGAGGTCAGGGCAGGTT	652
CIITA	gCIITA_57	CAGAAGAAGCTGCTCCGAGGT	660

Target Gene	crRNA	Spacer sequence	SEQ ID NO
CIITA	gCIITA_59	AGAGCTCAGGGATGACAGAGC	662
CIITA	gCIITA_60	TGCCGGGCAGTGTGCCAGCTC	663
CIITA	gCIITA_63	GCCACTCAGAGCCAGCCACAG	666
CIITA	gCIITA_65	GCAGCACGTGGTACAGGAGCT	668
CIITA	gCIITA_67	TGGGCACCCGCCTCACGCCTC	670
CIITA	gCIITA_70	CCAGGTCTTCCACATCCTTCA	673
CIITA	gCIITA_71	AAAGCCAAGTCCCTGAAGGAT	674
CIITA	gCIITA_72	GGTCCCGAACAGCAGGGAGCT	675
CIITA	gCIITA_73	TTTAGGTCCCGAACAGCAGGG	676
CIITA	gCIITA_76	GGGAAAGCCTGGGGGCCTGAG	679
CIITA	gCIITA_80	CAAGGACTTCAGCTGGGGGAA	682
CIITA	gCIITA_81	TAGGCACCCAGGTCAGTGATG	683
CIITA	gCIITA_82	CGACAGCTTGTACAATAACTG	684
CD247	gCD247_1	TGTGTTGCAGTTCAGCAGGAG	724
CD247	gCD247_3	CGGAGGGTCTACGGCGAGGCT	726
CD247	gCD247_4	TTATCTGTTATAGGAGCTCAA	727
CD247	gCD247_8	GACAAGAGACGTGGCCGGGAC	731
CD247	gCD247_12	CTAGCAGAGAAGGAAGAACCC	735
CD247	gCD247_15	ATCCCAATCTCACTGTAGGCC	738
CD247	gCD247_18	TCATTTCACTCCCAAACAACC	741
CD247	gCD247_19	ACTCCCAAACAACCAGCGCCG	742
CD52	gCD52_1	CTCTTCCTCCTACTCACCATC	53
CIITA	gCIITA_4	TAGGGGCCCCAACTCCATGGT	54
CTLA4	gCTLA4_4	AGCGGCACAAGGCTCAGCTGA	795
CTLA4	gCTLA4_14	CCTGGAGATGCATACTCACAC	67
CTLA4	gCTLA4_6	CAGAAGACAGGGATGAAGAGA	797
CTLA4	gCTLA4_19	CACTGGAGGTGCCCGTGCAGA	798
CTLA4	gCTLA4_13	TGTGTGAGTATGCATCTCCAG	70
DCK	gDCK_6	CGGAGGCTCCTTACCGATGTT	796
DCK	gDCK_2	TCAGCCAGCTCTGAGGGGACC	71
DCK	gDCK_8	CTCACAACAGCTGCAGGGAAG	72
DCK	gDCK_26	AGCTTGCCATTTCAGAGAGGCA	73
DCK	gDCK_30	TACATACCTGTCACTATACAC	74
FAS	gFAS_36	GTGTTGCTGGTGAGTGTGCAT	57
FAS	gFAS_34	TTTTTCTAGATGTGAACATGG	75
FAS	gFAS_35	ATGATTCCATGTTTACATCTA	76
FAS	gFAS_12	GTGTAACATACCTGGAGGACA	77
FAS	gFAS_1	GGAGGATTGCTCAACAACCAT	78
FAS	gFAS_59	TAGGAAACAGTGGCAATAAAT	79
HAVCR2	gTIM3_6	CTTGTAAGTAGTAGCAGCAGC	58
HAVCR2	gTIM3_29	CAAGGATGCTTACCACCAGGG	80
HAVCR2	gTIM3_6	TAAGTAGTAGCAGCAGCAGCA	799
HAVCR2	gTIM3_32	TATCAGGGAGGCTCCCCAGTG	800
HAVCR2	gTIM3_30	CCACCAGGGGACATGGCCAG	83
HAVCR2	gTIM3_12	AATGTGGCAACGTGGTGCTCA	84
HAVCR2	gTIM3_25	TGACATTAGCCAAGGTCACCC	85
HAVCR2	gTIM3_18	CGCAAAGGAGATGTGTCCCTG	86
IL7R	gIL7R_3	CAGGGGAGATGGATCCTATCT	749
IL7R	gIL7R_8	CATAACACACAGGCCAAGATG	754

Target Gene	crRNA	Spacer sequence	SEQ ID NO
LAG3	gLAG3_6	GGGTGCATACCTGTCTGGCTG	59
LAG3	gLAG3_38	TCAGGACCTTGGCTGGAGGCA	87
LAG3	gLAG3_33	GGTCACCTGGATCCCTGGGGA	88
LCK	gLCK1_3	ACCCATCAACCCGTAGGGATG	757
PDCD1	gPD_23	TCTGCAGGGACAATAGGAGCC	60
PDCD1	gPD_2	CCTTCCGCTCACCTCCGCCTG	89
PDCD1	gPD_8	GCACGAAGCTCTCCGATGTGT	90
PDCD1	gPD_29	CTAGCGGAATGGGCACCTCAT	91
PDCD1	gPD_27	CAGTGGCGAGAGAAGACCCCG	92
PTPN6	gPTPN6_6	TATGACCTGTATGGAGGGGAG	61
PTPN6	gPTPN6_46	ACTGCCCCCACCAGGCCTG	93
PTPN6	gPTPN6_7	CGACTCTGACAGAGCTGGTGG	801
PTPN6	gPTPN6_26	CAGAAGCAGGAGGTGAAGAAC	802
PTPN6	gPTPN6_1	ACCGAGACCTCAGTGGGCTGG	96
PTPN6	gPTPN6_37	TGGGCCCTACTCTGTGACCAA	97
PTPN6	gPTPN6_16	TGTGCTCAGTGACCAGCCCAA	98
PTPN6	gPTPN6_25	CCCACCCACATCTCAGAGTTT	99
PTPN6	gPTPN6_12	TTGTGCGTGAGAGCCTCAGCC	100
PTPN6	gPTPN6_22	AAGAAGACGGGGATTGAGGAG	101
PTPN6	gPTPN6_5	TCCCCTCCATACAGGTCATAG	803
PTPN6	gPTPN6_19	GCTCCCCCAGGGTGGACGCT	103
PTPN6	gPTPN6_14	GGCTGGTCACTGAGCACAGAA	104
TIGIT	gTIGIT_2	AGGCCTTACCTGAGGCGAGGG	62
TIGIT	gTIGIT_18	GTCCTCCCTCTAGTGGCTGAG	105
TRAC	gTRAC006	TGAGGGTGAAGGATAGACGCT	63
TRAC	gTRAC073	GCAGACAGGGAGAAATAAGGA	106
TRAC	gTRAC017	CAGGTGAAATTCCTGAGATGT	804
TRAC	gTRAC059	GACATCATTGACCAGAGCTCT	805
TRAC	gTRAC078	CCAGCTCACTAAGTCAGTCTC	109
TRAC	gTRAC012	TATGGAGAAGCTCTCATTTCT	110
TRAC	gTRAC039	TAAGATGCTATTTCCCGTATA	111
TRAC	gTRAC067	CCGTGTCAATTCTCTGGACTGC	112
TRAC	gTRAC079	ATTCCTCCACTTCAACACCTG	113
TRAC	gTRAC038	TACGGGAAATAGCATCTTAGA	114
TRAC	gTRAC061	GTGGCAATGGATAAGGCCGAG	115
TRAC	gTRAC058	CTTGCTTCAGGAATGGCCAGG	116
TRAC	gTRAC021	TAGTTCAAAACCTCTATCAAT	117
TRAC	gTRAC049	TCTGTGATATACACATCAGAA	118
TRAC	gTRAC074	GGCAGACAGGGAGAAATAAGG	119
TRAC	gTRAC018	CTCGATATAAGGCCTTGAGCA	120
TRAC	gTRAC043	GAGTCTCTCAGCTGGTACACG	121
TRAC	gTRAC075	TGGCAGACAGGGAGAAATAAG	122
TRAC	gTRAC082	CCAGCTGACAGATGGGCTCCC	123
TRAC	gTRAC040	CCGTATAAAGCATGAGACCGT	124
TRAC	gTRAC041	CCCCAACCCAGGCTGGAGTCC	125
TRAC	gTRAC076	TTGGCAGACAGGGAGAAATAA	126
TRAC	gTRAC014	TCAGAAGAGCCTGGCTAGGAA	127
TRAC	gTRAC029	CTCTGCCAGAGTTATATTGCT	128
TRAC	gTRAC028	CCATGCCTGCCTTTACTCTGC	129

Target Gene	crRNA	Spacer sequence	SEQ ID NO
TRAC	gTRAC050	GTCTGTGATATACACATCAGA	130
TRBC1+2	gTRBC1+2_1	AGCCATCAGAAGCAGAGATCT	705
TRBC1+2	gTRBC1+2_3	CGCTGTCAAGTCCAGTTCTAC	706
TRBC2	gTRBC2_11	AGACTGTGGCTTCACCTCCGG	711
TRBC2	gTRBC2_12	CCGGAGGTGAAGCCACAGTCT	712
TRBC2	gTRBC2_15	CTAGGGAAGGCCACCTTGTAT	715
TRBC2	gTRBC2_21	GAGCTAGCCTCTGGAATCCTT	720

Table 3 Selected Spacer Sequences Targeting Human Genes

Target Gene	crRNA	Spacer sequence	SEQ ID NO
ADORA2A	gADORA2A_16	CGGATCTTCCTGGCGGCGCGA	131
ADORA2A	gADORA2A_28	AAGGCAGCTGGCACCAGTGCC	132
ADORA2A	gADORA2A_2	TGGTGTCACTGGCGGCGGCCG	133
ADORA2A	gADORA2A_23	TTCTGCCCCGACTGCAGCCAC	134
ADORA2A	gADORA2A_7	GTGACCGGCACGAGGGCTAAG	135
ADORA2A	gADORA2A_8	CCATCGGCCTGACTCCCATGC	136
ADORA2A	gADORA2A_4	CCATCACCATCAGCACCGGGT	137
B2M	gB2M_21	TCACAGCCCAAGATAGTTAAG	138
B2M	gB2M_8	CTGAATTGCTATGTGTCTGGG	139
B2M	gB2M_11	CTGAAGAATGGAGAGAGAATT	140
B2M	gB2M_18	TCAGTGGGGTGAATTCAGTG	141
B2M	gB2M_5	CATTCTCTGCTGGATGACGTG	142
B2M	gB2M_10	ATCCATCCGACATTGAAGTTG	143
B2M	gB2M_22	CCCCACTTAACATCTTGGGC	144
B2M	gB2M_1	GCTGTGCTCGCGCTACTCTCT	145
B2M	gB2M_27	AATTCTCTCTCCATTCTTCAG	622
B2M	gB2M_31	CAGTGGGGTGAATTCAGTGT	626
B2M	gB2M_40	CATAGATCGAGACATGTAAGC	634
CD247	gCD247_7	CCCCCATCTCAGGGTCCCGGC	730
CD247	gCD247_9	TCTCCCTCTAACGTCTTCCCG	732
CD247	gCD247_13	TGCAGTTCCTGCAGAAAGAGG	736
CD247	gCD247_14	TGCAGGAAGTGCAGAAAGATA	737
CD247	gCD247_21	TGATTTGCTTTCACGCCAGGG	744
CD247	gCD247_22	CTTTCACGCCAGGGTCTCAGT	745
CD52	gCD52_4	GCTGGTGTCTGTTTTGTCCTGA	146
CIITA	gCIITA_18	TGCTGGCATCTCCATACTCTC	147
CIITA	gCIITA_29	GTCTCTTGCAAGTGCCTTTCTC	148
CIITA	gCIITA_34	CCGGCCTTTTTTACCTTGGGGC	638
CIITA	gCIITA_42	TGACTTTTCTGCCCCAAGTTCT	646
CIITA	gCIITA_46	CCAGAGCCCATGGGGCAGAGT	650
CIITA	gCIITA_47	TCCCCACCATCTCCACTCTGC	651
CIITA	gCIITA_51	CAGAGCCGGTGGAGCAGTTCT	655
CIITA	gCIITA_52	CCCAGCACAGCAATCACTCGT	656
CIITA	gCIITA_55	AGCCACATCTTGAAGAGACCT	658
CIITA	gCIITA_58	AGCTGTCCGGCTTCTCCATGG	661
CIITA	gCIITA_68	CCCCTCTGGATTGGGGAGCCT	671
CIITA	gCIITA_75	CCTCCTAGGCTGGGCCCTGTC	678
CIITA	gCIITA_83	TCTTGCCAGCGTCCAGTACAA	685

Target Gene	crRNA	Spacer sequence	SEQ ID NO
CTLA4	gCTLA4_27	CTGTTGCAGATCCAGAACCGT	149
CTLA4	gCTLA4_36	ACAGCTAAAGAAAAGAAGCCC	150
CTLA4	gCTLA4_41	TCAATTGATGGGAATAAAATA	151
CTLA4	gCTLA4_28	CTCCTCTGGATCCTTGCAGCA	152
CTLA4	gCTLA4_37	CACATAGACCCCTGTTGTAAG	153
CTLA4	gCTLA4_18	CTAGATGATTCCATCTGCACG	154
CTLA4	gCTLA4_5	TTCTTCTCTTCATCCCTGTCT	155
DCK	gDCK_9	AGGATATTCACAAATGTTGAC	156
DCK	gDCK_22	GAAGGTAAAAGACCATCGTTC	157
DCK	gDCK_21	TCATACATCATCTGAAGAACA	158
DCK	gDCK_7	ATCTTTCCTCACAACAGCTGC	159
FAS	gFAS_47	AGTGAAGAGAAAGGAAGTACA	160
FAS	gFAS_45	TTTGTCTTTTCAGTGAAGAGA	161
FAS	gFAS_25	CTAGGCTTAGAAGTGGAATA	162
FAS	gFAS_10	GAAGGCCTGCATCATGATGGC	163
FAS	gFAS_32	GTGCAAGGGTCACAGTGTTC	164
FAS	gFAS_5	GGACGATAATCTAGCAACAGA	165
FAS	gFAS_14	TTCTTGGGCAGGTGAAAGGA	166
FAS	gFAS_29	GTTTACATCTGCACTTGGTAT	167
FAS	gFAS_33	CTTGGTGCAAGGGTCACAGTG	168
FAS	gFAS_71	CTGTTCTGCTGTGTCTTGGAC	169
FAS	gFAS_38	CTCTTTCGCACTTGGTGTGCT	170
FAS	gFAS_70	TGTTCTGCTGTGTCTTGGACA	171
FAS	gFAS_4	ACAGGTTCTTACGTCTGTTGC	172
FAS	gFAS_15	GGCAGGTGAAAGGAAAGCTAG	173
HAVCR2	gTIM3_42	CTAGGGTATTCTCATAGCAA	174
HAVCR2	gTIM3_10	CCCCAGCAGACGGGCACGAGG	175
HAVCR2	gTIM3_47	GCCAACCTCCCTCCCTCAGGA	176
HAVCR2	gTIM3_34	TGTTTCCATAGCAAATATCCA	177
HAVCR2	gTIM3_19	GATCCGGCAGCAGTAGATCCC	178
HAVCR2	gTIM3_48	CCAATCCTGAGGGAGGGAGGT	179
HAVCR2	gTIM3_36	CGGGACTCTGGAGCAACCATC	180
HAVCR2	gTIM3_15	GCCAGTATCTGGATGTCCAAT	181
HAVCR2	gTIM3_27	ACTGCAGCCTTTCCAAGGATG	182
HAVCR2	gTIM3_41	CCCCTTACTAGGGTATTCTCA	183
HAVCR2	gTIM3_23	ACCTGAAGTTGGTCATCAAAC	184
HAVCR2	gTIM3_28	CCAAGGATGCTTACCACCAGG	185
HAVCR2	gTIM3_40	GTTTCCCCCTTACTAGGGTAT	186
HAVCR2	gTIM3_13	ATCAGTCCTGAGCACCACGTT	187
IL7R	gIL7R_2	CCAGGGGAGATGGATCCTATC	748
IL7R	gIL7R_7	TCTGTCGCTCTGTTGGTCATC	753
LAG3	gLAG3_35	TGAGGTGACTCCAGTATCTGG	188
LAG3	gLAG3_41	CCAGCCTTGGCAATGCCAGCT	189
LAG3	gLAG3_37	TGTGGAGCTCTCTGGACACCC	190
LAG3	gLAG3_16	GGGCAGGAAGAGGAAGCTTTC	191
LAG3	gLAG3_46	TCCATAGGTGCCCAACGCTCT	192
LAG3	gLAG3_27	CCACCTGAGGCTGACCTGTGA	193
LAG3	gLAG3_31	CCCAGGGATCCAGGTGACCCA	194
LAG3	gLAG3_3	ACCTGGAGCCACCCAAAGCGG	195
LAG3	gLAG3_25	CCCTTCGACTAGAGGATGTGA	196

Target Gene	crRNA	Spacer sequence	SEQ ID NO
LAG3	gLAG3_13	CGCTAAGTGGTGATGGGGGGA	197
LAG3	gLAG3_22	GCAGTGAGGAAAGACCGGGTC	198
PDCD1	gPD_20	CAGAGAGAAGGGCAGAAAGTGC	199
PDCD1	gPD_22	GAAGTGGCCGGCTGGCCTGGG	200
PDCD1	gPD_18	GTGCCCTTCCAGAGAGAAGGG	201
PLCG1	gPLCG1_2	CCTTTCTGCGCTTCGTGGTGT	759
PLCG1	gPLCG1_4	TGCGCTTCGTGGTGTATGAGG	761
PLCG1	gPLCG1_5	GTGGTGTATGAGGAAGACATG	762
PTPN6	gPTPN6_20	GAGACCTTCGACAGCCTCACG	202
PTPN6	gPTPN6_41	CTGGACCAGATCAACCAGCGG	203
PTPN6	gPTPN6_53	CCCCCTGCACCCGGCTGCAG	204
PTPN6	gPTPN6_28	CACCAGCGTCTGGAAGGGCAG	205
PTPN6	gPTPN6_42	CTGCCGCTGGTTGATCTGGTC	206
PTPN6	gPTPN6_32	TGGCAGATGGCGTGGCAGGAG	207
PTPN6	gPTPN6_4	CTGGCTCGGCCAGTCGCAAG	208
PTPN6	gPTPN6_8	AGGTGGATGATGGTGGCGTCG	209
PTPN6	gPTPN6_40	GGGAGACCTGATTCTGGGAGAT	210
PTPN6	gPTPN6_48	AATGAACTGGGCGATGGCCAC	211
PTPN6	gPTPN6_10	TCTAGGTGGTACCATGGCCAC	212
PTPN6	gPTPN6_39	CAGGTCTCCCCGCTGGACAAT	213
TIGIT	gTIGIT_11	GGGTGGCACATCTCCCATCC	214
TIGIT	gTIGIT_7	TGCAGAGAAAGGTGGCTCTAT	215
TIGIT	gTIGIT_10	TAATGCTGACTTGGGGTGGCA	216
TIGIT	gTIGIT_27	CTCCTGAGGTCACCTTCCACA	217
TRAC	gTRAC066	CTAAGAAACAGTGAGCCTTGT	218
TRAC	gTRAC042	CCTCTTTGCCCAACCCAGGC	219
TRAC	gTRAC035	AGGTTTCCTTGAGTGGCAGGC	220
TRAC	gTRAC044	AGAATCAAAATCGGTGAATAG	221
TRAC	gTRAC072	CCCCTTACTGCTCTCTAGGC	222
TRAC	gTRAC062	GGTGGCAATGGATAAGGCCGA	223
TRAC	gTRAC020	GAATATAAATCAGAACACCT	224
TRAC	gTRAC013	TTTCTCAGAAGAGCCTGGCTA	225
TRAC	gTRAC068	CCCGTGTCATTCTCTGGACTG	226
TRAC	gTRAC025	CTGGGCCTTTTTCCCATGCCT	227
TRAC	gTRAC019	AATATAAATCAGAACACCTG	228
TRAC	gTRAC048	ATTCTCAAACAAATGTGTCAC	229
TRAC	gTRAC036	CTTGAGTGGCAGGCCAGGCCT	230
TRAC	gTRAC056	CATGTGCAAACGCCTTCAACA	231
TRAC	gTRAC064	TACTAAGAAACAGTGAGCCTT	232
TRAC	gTRAC071	CTCAGACTGTTTGCCCCTTAC	233
TRAC	gTRAC081	TAATTCCTCCACTTCAACACC	234
TRAC	gTRAC030	ATAGGATCTTCTTCAAAACCC	235
TRAC	gTRAC033	GAAGAAGATCCTATTAAATAA	236
TRAC	gTRAC001	TGTTTTTAATGTGACTCTCAT	237
TRAC	gTRAC009	GTACTTTACAGTTTATTAAAT	238
TRAC	gTRAC007	ATAAACTGTAAAGTACCAAAC	239
TRAC	gTRAC084	GACTTTTCCCAGCTGACAGAT	240
TRAC	gTRAC083	CCCAGCTGACAGATGGGCTCC	241
TRBC2	gTRBC2_14	CCAGCAAGGGGTCCCTGTCTGC	714
TRBC2	gTRBC2_17	CCATGGCCATCAGCACGAGGG	717

Target Gene	crRNA	Spacer sequence	SEQ ID NO
TRBC2	gTRBC2_19	CACAGGTCAAGAGAAAGGATT	719
CSF2	gCSF2_001	TGAGATGACTTCTACTGTTTC	1100
CSF2	gCSF2_002	CCTTTTCTACAGAATGAAACA	1101
CSF2	gCSF2_003	CTTTTCTACAGAATGAAACAG	1102
CSF2	gCSF2_004	CTACAGAATGAAACAGTAGAA	1103
CSF2	gCSF2_005	TACAGAATGAAACAGTAGAAG	1104
CSF2	gCSF2_006	CCACAGGAGCCGACCTGCCTA	1105
CSF2	gCSF2_007	CACAGGAGCCGACCTGCCTAC	1106
CSF2	gCSF2_008	ttatTTTTcTTTTTTAAAGG	1107
CSF2	gCSF2_009	tatTTTTcTTTTTTAAAGGA	1108
CSF2	gCSF2_010	atTTTTcTTTTTTAAAGGAA	1109
CSF2	gCSF2_011	ttTTTTcTTTTTTAAAGGAAA	1110
CSF2	gCSF2_012	tcTTTTTTAAAGGAAACTTC	1111
CSF2	gCSF2_013	ctTTTTTTAAAGGAAACTTCC	1112
CSF2	gCSF2_014	ttTTTTTTAAAGGAAACTTCCT	1113
CSF2	gCSF2_015	tttAAAGGAAACTTCCTGTGC	1114
CSF2	gCSF2_016	ttAAAGGAAACTTCCTGTGCA	1115
CSF2	gCSF2_017	tAAAGGAAACTTCCTGTGCAA	1116
CSF2	gCSF2_018	AAAGGTGATAATCTGGGTTGC	1117
CSF2	gCSF2_019	AAAGGAAACTTCCTGTGCAAC	1118
CSF2	gCSF2_020	AAGGAAACTTCCTGTGCAACC	1119
CSF2	gCSF2_021	AAACTTTCAAAGGTGATAATC	1120
CSF2	gCSF2_022	AAAGTTTCAAAGAGAACCTGA	1121
CSF2	gCSF2_023	AAAGAGAACCTGAAGGACTTT	1122
CSF2	gCSF2_024	TGCTTGTCATCCCCCTTGACT	1123
CSF2	gCSF2_025	ACTGCTGGGAGCCAGTCCAGG	1124
CSF2	gCSF2_026	CCTAGGTGGTCAGGCTTGCGG	1125
CSF2	gCSF2_027	TGGTCACCATTAATCATTTCC	1126
CSF2	gCSF2_028	CTCTGTGTATTTAAGAGCTCT	1127
CSF2	gCSF2_029	AGAGCTCTTTTGCCAGTGAGC	1128
CSF2	gCSF2_030	ATTCTGTAGAAAAGGAAAATG	1129
CSF2	gCSF2_031	ACCTCCAGGTAAGATGCTTCT	1130
CSF2	gCSF2_032	CAGAAGCCCCTGCCCTGGGGT	1131
CSF2	gCSF2_033	GATGGCACCACACAGGGTTGT	1132
CSF2	gCSF2_034	TCTCCAGTCAGCTGGCTGCAG	1133
CSF2	gCSF2_035	TCAGCTGAGCGGCCATGGGCA	1134
CSF2	gCSF2_036	CCACCTGTCCCCTGGTGAATC	1135
CSF2	gCSF2_037	GGGCGCTCACTGTGCCCCGAG	1136
CSF2	gCSF2_038	AGGAACAACCCTTGCCCACCC	1137
CSF2	gCSF2_039	CTGCTGCCCCCAGCCCCCAGG	1138
CSF2	gCSF2_040	TGTGCCAACAGTTATGTAATG	1139
CSF2	gCSF2_041	ATCCCAAGGAGTCAGAGCCAC	1140
CSF2	gCSF2_042	CCCTCACCTCTGACCTCATTA	1141
CSF2	gCSF2_043	CTTGGGTTTGCCCTCACCTCT	1142
CSF2	gCSF2_044	CTCTGGCCCCACATGGGGTGC	1143
CSF2	gCSF2_045	CTCCCTTCCCGCAGGAAGGAG	1144
CSF2	gCSF2_046	TGGCCTTGACTCCACTCCTTC	1145
CSF2	gCSF2_047	GTCCCAGGGCAGAGCAGGGCA	1146
CSF2	gCSF2_048	ACTGCCCAGAAGGCCAACCTC	1147
CSF2	gCSF2_049	TCTACTGCCTCTTAGAACTCA	1148

Target Gene	crRNA	Spacer sequence	SEQ ID NO
CSF2	gCSF2_050	AAAGGAAACTTCCTGTGCAAt	1149
CSF2	gCSF2_051	AAGGAAACTTCCTGTGCAAtC	1150
CSF2	gCSF2_052	AAAGGTGATAgTCTGGaTTGC	1151
CSF2	gCSF2_053	AAACTTTCAAAGGTGATAgTC	1152
CD40LG	gCD40LG_001	GTTGTATGTTTCGATCATGCT	1153
CD40LG	gCD40LG_002	AACTTTAACACAGCATGATCG	1154
CD40LG	gCD40LG_003	ACACAGCATGATCGAAACATA	1155
CD40LG	gCD40LG_004	ATGCTGATGGGCAGTCCAGTG	1156
CD40LG	gCD40LG_005	CATGCTGATGGGCAGTCCAGT	1157
CD40LG	gCD40LG_006	TATGTATTTACTTACTGTTTT	1158
CD40LG	gCD40LG_007	ATGTATTTACTTACTGTTTTT	1159
CD40LG	gCD40LG_008	TGTATTTACTTACTGTTTTTC	1160
CD40LG	gCD40LG_009	CTTACTGTTTTTCTTATCACC	1161
CD40LG	gCD40LG_010	TCTTATCACCCAGATGATTGG	1162
CD40LG	gCD40LG_011	CTTATCACCCAGATGATTGGG	1163
CD40LG	gCD40LG_012	TTATCACCCAGATGATTGGGT	1164
CD40LG	gCD40LG_013	TGCTGTGTATCTTCATAGAAG	1165
CD40LG	gCD40LG_014	GCTGTGTATCTTCATAGAAGG	1166
CD40LG	gCD40LG_015	CTGTGTATCTTCATAGAAGGT	1167
CD40LG	gCD40LG_016	ATGAATACAAAATCTTCATGA	1168
CD40LG	gCD40LG_017	CATGAATACAAAATCTTCATG	1169
CD40LG	gCD40LG_018	TCCTGTGTTGCATCTCTGTAT	1170
CD40LG	gCD40LG_019	GTATTCATGAAAACGATACAG	1171
CD40LG	gCD40LG_020	TATTCATGAAAACGATACAGA	1172
CD40LG	gCD40LG_021	ATCTCCTCACAGTTCAGTAAG	1173
CD40LG	gCD40LG_022	AATCTCCTCACAGTTCAGTAA	1174
CD40LG	gCD40LG_023	CCAGTAATTAAGCTGCTTACC	1175
CD40LG	gCD40LG_024	ACCAGTAATTAAGCTGCTTAC	1176
CD40LG	gCD40LG_025	AAGGCTTTGTGAAGGTAAGCA	1177
CD40LG	gCD40LG_026	TTCGTCTCCTCTTTGTTTAAAC	1178
CD40LG	gCD40LG_027	TTTCTTCGTCTCCTCTTTGTT	1179
CD40LG	gCD40LG_028	CTTTCTTCGTCTCCTCTTTGT	1180
CD40LG	gCD40LG_029	AGGATATAATGTTAAACAAAG	1181
CD40LG	gCD40LG_030	GGATATAATGTTAAACAAAGA	1182
CD40LG	gCD40LG_031	AAAGCTGTTTTCTTTCTTCGT	1183
CD40LG	gCD40LG_032	CATTTCAAAGCTGTTTTCTTT	1184
CD40LG	gCD40LG_033	GCATTTCAAAGCTGTTTTCTT	1185
CD40LG	gCD40LG_034	TGCATTTCAAAGCTGTTTTCT	1186
CD40LG	gCD40LG_035	AGGATTCTGATCACCTGAAAT	1187
CD40LG	gCD40LG_036	TGGTTCCATTTTCAGGTGATCA	1188
CD40LG	gCD40LG_037	GGTTCCATTTTCAGGTGATCAG	1189
CD40LG	gCD40LG_038	GTTCCATTTTCAGGTGATCAGA	1190
CD40LG	gCD40LG_039	AGGTGATCAGAATCCTCAAAT	1191
CD40LG	gCD40LG_040	CTGCTGGCCTCACTTATGACA	1192
CD40LG	gCD40LG_041	AGCCCACTGTAACACTGTTAC	1193
CD40LG	gCD40LG_042	CAGCCCACTGTAACACTGTTA	1194
CD40LG	gCD40LG_043	TCAGCCCACTGTAACACTGTT	1195
CD40LG	gCD40LG_044	CCTTTCTTTGTAACAGTGTTA	1196
CD40LG	gCD40LG_045	TTTGTAACAGTGTTACAGTGG	1197
CD40LG	gCD40LG_046	TAACAGTGTTACAGTGGGCTG	1198

Target Gene	crRNA	Spacer sequence	SEQ ID NO
CD40LG	gCD40LG_047	CAGGGTTACCAAGTTGTTGCT	1199
CD40LG	gCD40LG_048	CCAGGGTTACCAAGTTGTTGC	1200
CD40LG	gCD40LG_049	CCATTTTCCAGGGTTACCAAG	1201
CD40LG	gCD40LG_050	ACGGTCAGCTGTTTCCCATT	1202
CD40LG	gCD40LG_051	AACGGTCAGCTGTTTCCCATT	1203
CD40LG	gCD40LG_052	GGCAGAGGCTGGCTATAAATG	1204
CD40LG	gCD40LG_053	TAGCCAGCCTCTGCCTAAAGT	1205
CD40LG	gCD40LG_054	CAGCTCTGAGTAAGATTCTCT	1206
CD40LG	gCD40LG_055	GCGGAAGTGTGGGTATTTGCA	1207
CD40LG	gCD40LG_056	AATTGCAACCAGGTGCTTCGG	1208
CD40LG	gCD40LG_057	TCAATGTGACTGATCCAAGCC	1209
CD40LG	gCD40LG_058	AGTAAGCCAAAGGACGTGAAG	1210
CD40LG	gCD40LG_059	GCTTACTCAAACCTCTGAACAG	1211
CD40LG	gCD40LG_060	ACTGCTGGCCTCACTTATGAC	1212
CD3E	gCD3E_1	CACTCCATCCTACTCACCTGA	1213
CD3E	gCD3E_2	tttttCTTATTTATTTCTAG	1214
CD3E	gCD3E_3	tttttCTTATTTATTTCTAGT	1215
CD3E	gCD3E_4	tttCTTATTTATTTTCTAGTT	1216
CD3E	gCD3E_5	ttCTTATTTATTTTCTAGTTG	1217
CD3E	gCD3E_6	tCTTATTTATTTTCTAGTTGG	1218
CD3E	gCD3E_7	CTTATTTATTTTCTAGTTGGC	1219
CD3E	gCD3E_8	TTATTTATTTTCTAGTTGGCG	1220
CD3E	gCD3E_9	TTTTCTAGTTGGCGTTTGGGG	1221
CD3E	gCD3E_10	CTAGTTGGCGTTTGGGGGCAA	1222
CD3E	gCD3E_11	TAGTTGGCGTTTGGGGGCAAG	1223
CD3E	gCD3E_12	CTTTTCAGGTAATGAAGAAAT	1224
CD3E	gCD3E_13	CAGGTAATGAAGAAATGGGTA	1225
CD3E	gCD3E_14	AGGTAATGAAGAAATGGGTAA	1226
CD3E	gCD3E_15	CTTTTTTCATTTTCAGGTGGT	1227
CD3E	gCD3E_16	TTCATTTTCAGGTGGTATTAC	1228
CD3E	gCD3E_17	TCATTTTCAGGTGGTATTACA	1229
CD3E	gCD3E_18	CATTTTCAGGTGGTATTACAC	1230
CD3E	gCD3E_19	ATTTTCAGGTGGTATTACACA	1231
CD3E	gCD3E_20	CAGGTGGTATTACACAGACAC	1232
CD3E	gCD3E_21	AGGTGGTATTACACAGACACG	1233
CD3E	gCD3E_22	CCTTCTTTCTCCCCAGCATAT	1234
CD3E	gCD3E_23	TCCCCAGCATATAAAGTCTCC	1235
CD3E	gCD3E_24	AGATCCAGGATACTGAGGGCA	1236
CD3E	gCD3E_25	tcatTGTGTTGCCATAGTATT	1237
CD3E	gCD3E_26	atcatTGTGTTGCCATAGTAT	1238
CD3E	gCD3E_27	tatcatTGTGTTGCCATAGTA	1239
CD3E	gCD3E_28	tcatcctcatcaccgcctatg	1240
CD3E	gCD3E_29	atcatcctcatcaccgcctat	1241
CD3E	gCD3E_30	tatcatcctcatcaccgccta	1242
CD3E	gCD3E_31	CTCCAATTCTGAAAATTCCTT	1243
CD3E	gCD3E_32	CAGAAATTGGAGCAAAGTGTT	1244
CD3E	gCD3E_33	AGAATTGGAGCAAAGTGTTA	1245
CD3E	gCD3E_34	CTTCTCTGGGGTAGCAGACA	1246
CD3E	gCD3E_35	ATCTCTACCTGAGGGCAAGAG	1247
CD3E	gCD3E_36	TCTCTACCTGAGGGCAAGAGG	1248

Target Gene	crRNA	Spacer sequence	SEQ ID NO
CD3E	gCD3E_37	TATTCTTGCTCCAGTAGTAAA	1249
CD3E	gCD3E_38	CTACTGGAGCAAGAATAGAAA	1250
CD3E	gCD3E_39	CCTGCCGCCAGCACCCGCTCC	1251
CD3E	gCD3E_40	CCCTCCTTCCTCCGCAGGACA	1252
CD3E	gCD3E_41	TATCCACGTTACCTCATAGT	1253
CD3E	gCD3E_42	ACCCCCAGCCCATCCGGAAAG	1254
CD38	gCD38_001	TCCCCGGACACCGGGCTGAAC	1255
CD38	gCD38_002	AGTGTACTTGACGCATCGCGC	1256
CD38	gCD38_003	CCGAGACCGTCCTGGCGCGAT	1257
CD38	gCD38_004	GCAGTCTACATGTCTGAGATA	1258
CD38	gCD38_005	TGTGTTTTATCTCAGACATGT	1259
CD38	gCD38_006	TCTCAGACATGTAGACTGCCA	1260
CD38	gCD38_007	AAATAAATGCACCCCTGAAAG	1261
CD38	gCD38_008	AAGGGTGCATTTATTTCAAAA	1262
CD38	gCD38_009	TTTCAAAACATCCTTGCAACA	1263
CD38	gCD38_010	AAAACATCCTTGCAACATTAC	1264
CD38	gCD38_011	TTCTGCTCCAAAGAAGAATCT	1265
CD38	gCD38_012	TTCTTCCTTAGATTCTTCTTT	1266
CD38	gCD38_013	GAGCAGAATAAAAAGATCTGGC	1267
CD38	gCD38_014	TACAAACTATGTCTTTTAGAA	1268
CD38	gCD38_015	TCCAGTCTGGGCAAGATTGAT	1269
CD38	gCD38_016	GAAATAAACTATCAATCTTGC	1270
CD38	gCD38_017	CAGAATACTGAAACAGGGTTG	1271
CD38	gCD38_018	AGTATTCTGGAAAACGGTTTC	1272
CD38	gCD38_019	ACTACTTGGTACTTACCCTGC	1273
CD38	gCD38_020	AGTTTGCAGAAGCTGCCTGTG	1274
CD38	gCD38_021	CAGAAGCTGCCTGTGATGTGG	1275
CD38	gCD38_022	CTGCGGGATCCATTGAGCATC	1276
CD38	gCD38_023	TCAAAGATTTTACTGCGGGAT	1277
CD38	gCD38_024	GGGTTCTTTGTTTCTTCTATT	1278
CD38	gCD38_025	TTTCTTCTATTTTAGCACTTT	1279
CD38	gCD38_026	TTCTATTTTAGCACTTTTGGG	1280
CD38	gCD38_027	GCACTTTTGGGAGTGTGGAAG	1281
CD38	gCD38_028	GGAGTGTGGAAGTCCATAATT	1282
CD38	gCD38_029	CAACCAGAGAAGGTTTCAGACA	1283
CD38	gCD38_030	TGGTGGGATCCTGGCATAAGT	1284
CD38	gCD38_031	TTCCCCAGAGACTTATGCCAG	1285
CD38	gCD38_032	CTTATAATCGATTCCAGCTCT	1286
CD38	gCD38_033	CTTTTTTGCTTTCTTGTCATA	1287
CD38	gCD38_034	CTTTCTTGTCATAGACCTGAC	1288
CD38	gCD38_035	ACACACTGAAGAACTTGTCA	1289
CD38	gCD38_036	TTGTCATAGACCTGACAAGTT	1290
CD38	gCD38_037	TTCAAGTGTGTGAAAAATCCTG	1291
ALP NR	gAPLNR_001	ACAACACTATGGGGCAGACA	1292
ALP NR	gAPLNR_002	CAGTCTGTGTACTCACACTCA	1293
ALP NR	gAPLNR_003	GGAGCAGCCGGGAGAGAGGC	1294
ALP NR	gAPLNR_004	GGACCTTCTTCTGCAAGCTCA	1295
ALP NR	gAPLNR_006	TGGTGCCCTTCACCATCATGC	1296
ALP NR	gAPLNR_007	GGCGATGAAGAAGTAACAGGT	1297
ALP NR	gAPLNR_008	CCCTGTGCTGGATGCCCTACC	1298

Target Gene	crRNA	Spacer sequence	SEQ ID NO
ALPNR	gAPLNR_009	ACCTCTTCCTCATGAACATCT	1299
ALPNR	gAPLNR_010	GACCCCCGCTTCCGCCAGGCC	1300
ALPNR	gAPLNR_011	TCGTGCATCTGTTCTCCACCC	1301
BBS1	gBBS1_005	CATGGGGATGGGGAATACAAG	1302
BBS1	gBBS1_007	GGTCATCACCAGTGGTCCTTT	1303
BBS1	gBBS1_009	GCCTGGTTCCAAAGGTCTTGT	1304
BBS1	gBBS1_015	ACTTAGCTCCAGCTGCAGAAA	1305
BBS1	gBBS1_016	CAAATGCCTCCATTTCACTTA	1306
BBS1	gBBS1_017	TGCAGCTGGAGCTAAGTGAAA	1307
BBS1	gBBS1_018	TAAACCAACACAAGTCCAAC	1308
BBS1	gBBS1_028	CACTGTCCACTTCCCTAGGTG	1309
BBS1	gBBS1_032	CGTGGATCAGACACTGCGAGA	1310
BBS1	gBBS1_033	TCCACCCACCCTCTCCATAGG	1311
CALR	gCALR_001	GATTTCGATCCAGCGGGAAGTC	1312
CALR	gCALR_006	CAGACAAGCCAGGATGCACGC	1313
CALR	gCALR_011	ACCGTGAAGTGCACCACCAGC	1314
CALR	gCALR_012	CTAATAGTTTGGACCAGACAG	1315
CALR	gCALR_013	GACCAGACAGACATGCACGGA	1316
CALR	gCALR_014	CCACCACCCCCAGGCACACCT	1317
CALR	gCALR_015	CACACCTGTACACACTGATTG	1318
CALR	gCALR_017	AAGCATCAGGATCCTTTATCT	1319
CALR	gCALR_019	TGGGTGGATCCAAGTGCCCTT	1320
CALR	gCALR_021	CTCCAAGTCTCACCTGCCAGA	1321
CD3G	gCD3G_001	CCGGAGGACAGAGACTGACAT	1322
CD3G	gCD3G_004	GCTTCTGCATCACAAGTCAGA	1323
CD3G	gCD3G_006	TCTTCAGTTAGGAAGCCGATC	1324
CD3G	gCD3G_007	AAGATGGGAAGATGATCGGCT	1325
CD3G	gCD3G_008	CACTGATACATCCCTCGAGGG	1326
CD3G	gCD3G_011	GTTCAATGCAGTTCTGACACA	1327
CD3G	gCD3G_012	CCTACAGTGTGTCAGAACTGC	1328
CD3G	gCD3G_017	CCTCTCGACTGGCGAACTCCA	1329
CD3G	gCD3G_022	CTTGAAGGTGGCTGTACTGGT	1330
CD3G	gCD3G_023	CAGGTACTTTGGCCCAGTCAA	1331
CD58	gCD58_004	CCAACAAATATATGGTGTGT	1332
CD58	gCD58_005	AAGGCACATTGCTTGGTACAT	1333
CD58	gCD58_010	AAAGAGGTCCTATGGAAAAAA	1334
CD58	gCD58_012	AAAGATGAGAAAGCTCTGAAT	1335
CD58	gCD58_018	GCGATTCCATTTTCATACTCAT	1336
CD58	gCD58_019	CAGAGTCTCTTCCATCTCCCA	1337
CD58	gCD58_020	CATTGCTCCATAGGACAATCC	1338
CD58	gCD58_023	AGATGGAAAATGATCTTCCAC	1339
CD58	gCD58_028	TAGGTCATTCAAGACACAGAT	1340
CD58	gCD58_033	GGTATTCTGAAATGTGACAGA	1341
COL17A1	gCOL17A1_005	TAGTTGTCACTGAAACAGTAA	1342
COL17A1	gCOL17A1_006	GCATAGCCATTGCTGGTCCCG	1343
COL17A1	gCOL17A1_017	ACTCCGTCCTCTGGTTGAAGA	1344
COL17A1	gCOL17A1_024	CAGTGTCAGGCACCTACGATG	1345
COL17A1	gCOL17A1_047	CTGTTCCATCATTAGCTTCTT	1346
COL17A1	gCOL17A1_054	AGGTGACATGGGAAGTCCAGG	1347
COL17A1	gCOL17A1_065	CAAGAAGCAGCAAACCTGACCT	1348

Target Gene	crRNA	Spacer sequence	SEQ ID NO
COL17A1	gCOL17A1_070	GGTGACAAAGGACCAATGGGA	1349
COL17A1	gCOL17A1_084	AGAGGGGGTCATCGATGCTCAC	1350
COL17A1	gCOL17A1_094	ATGCCGGCTCTACTGTACCTT	1351
DEFB134	gDEFB134_001	CCTGCCAGCACTGGATCCCAA	1352
DEFB134	gDEFB134_004	CTTTGGGATCCAGTGCTGGCA	1353
DEFB134	gDEFB134_007	CTTCCAGGTATAAAATTCATTA	1354
DEFB134	gDEFB134_008	TTGTGCATTTCTGATGATAAT	1355
DEFB134	gDEFB134_009	TAGCATTTCTTGTGCATTTCT	1356
DEFB134	gDEFB134_010	ACTCTCATAGCATTCAAGTCT	1357
DEFB134	gDEFB134_011	ACACAGCACTCCAGCTGAAAC	1358
DEFB134	gDEFB134_012	CTTTGACACAGCACTCCAGCT	1359
DEFB134	gDEFB134_013	AGCTGGAGTGCTGTGTCAAAG	1360
DEFB134	gDEFB134_014	TTATGTTCAGGGTGCAGGATTT	1361
ERAP1	gERAP1_008	CATGGATCAAGAGATCATAAT	1362
ERAP1	gERAP1_015	CAAAAGCACCTACAGAACCAA	1363
ERAP1	gERAP1_029	AGTCTGTCAGCAAGATAACCA	1364
ERAP1	gERAP1_035	GGTAGGGGATACGGTATGCTG	1365
ERAP1	gERAP1_037	AGCATACCGTATCCCCTACCC	1366
ERAP1	gERAP1_039	CATAGCACCAGACTGAAAGTC	1367
ERAP1	gERAP1_061	CCTTATCATAAGAAACATCAT	1368
ERAP1	gERAP1_065	AATGCGTCAGCACTAAGATAC	1369
ERAP1	gERAP1_077	CCCTAATAACCATCACAGTGA	1370
ERAP1	gERAP1_078	CTCTAGGAGCATTACCCAGTG	1371
ERAP2	gERAP2_001	TGTGTGAATTAACCATTGCAG	1372
ERAP2	gERAP2_014	ATGTATCTTGAATCTTCCTCT	1373
ERAP2	gERAP2_018	AGTTACCCTGCTCATGAACAA	1374
ERAP2	gERAP2_046	GAGAGTGGATAGTAGATATCA	1375
ERAP2	gERAP2_048	ATATCTACTATCCACTCTCCA	1376
ERAP2	gERAP2_099	ATGTGGACTCAAATGGTTACT	1377
ERAP2	gERAP2_108	CCTGTCAATCACTGGCTTAAA	1378
ERAP2	gERAP2_118	GAGCAATATGAACTGTCAATG	1379
ERAP2	gERAP2_134	ACTTGGGCTCATATGACATAA	1380
ERAP2	gERAP2_261	TCCTTACCATGTTACTTGTCA	1381
IFNGR1	gIFNGR1_004	TTACAGTGCCTACACCAACTA	1382
IFNGR1	gIFNGR1_006	CCGTAGAGGTAAAGAACTATG	1383
IFNGR1	gIFNGR1_008	GTGTTAAGAATTCAGAATGGA	1384
IFNGR1	gIFNGR1_010	ATGGATCACCAACATGATCAG	1385
IFNGR1	gIFNGR1_012	ACTCTGACCCAAAGAGAATTT	1386
IFNGR1	gIFNGR1_021	GGGATCATAATCGACTTCCTG	1387
IFNGR1	gIFNGR1_025	AGTTGTAACACCCACACATG	1388
IFNGR1	gIFNGR1_042	GAGACAAAACCTGAATCAAAA	1389
IFNGR1	gIFNGR1_049	AGTAGTAACCAGTCTGAACCT	1390
IFNGR1	gIFNGR1_052	TGGAGTGATCACTCTCAGAAC	1391
IFNGR2	gIFNGR2_001	TCTGTCCCCCTCAAGACCCTC	1392
IFNGR2	gIFNGR2_003	AACTGCACTTGGTAGACAACA	1393
IFNGR2	gIFNGR2_005	CTTCCCAGCACCGACAGTAAA	1394
IFNGR2	gIFNGR2_006	AATGTCACTCTACGCCTTCGA	1395
IFNGR2	gIFNGR2_012	CCAGTAATGGACATAATAACA	1396
IFNGR2	gIFNGR2_015	AGTTATCCAATGAAATGGAGT	1397
IFNGR2	gIFNGR2_017	ATTGGATAACTTAAACCCTC	1398

Target Gene	crRNA	Spacer sequence	SEQ ID NO
IFNGR2	gIFNGR2_021	GTAGCAAGATATGTTGCTTAA	1399
IFNGR2	gIFNGR2_026	GCCTCCACTGAGCTTCAGCAA	1400
IFNGR2	gIFNGR2_031	ACACTCCACCAAGCATCCCAT	1401
JAK1	gJAK1_002	CTTCCACAACAGTATCTAAAT	1402
JAK1	gJAK1_021	GCTACAAGCGATATATTTCCAG	1403
JAK1	gJAK1_037	ATTCGAATGACGGTGGAAACG	1404
JAK1	gJAK1_059	GCATGAAGCTGATGTTATCCG	1405
JAK1	gJAK1_074	GTACACACATTTCCATGGACC	1406
JAK1	gJAK1_075	CCAGAGCGTGGTTCCAAAGCT	1407
JAK1	gJAK1_090	AGATCAGCTATGTGGTTACCT	1408
JAK1	gJAK1_100	CCTTACAAATCTGAACGGCAT	1409
JAK1	gJAK1_108	ACCAAAGCAATTGAAACCGAT	1410
JAK1	gJAK1_111	GATTGCATTAAACATTCTGGA	1411
JAK2	gJAK2_009	GAAGCAGCAATACAGATTTCT	1412
JAK2	gJAK2_101	AAGGCGTACGAAGAGAAGTAG	1413
JAK2	gJAK2_118	AGATATGTATCTAGTGATCCA	1414
JAK2	gJAK2_121	GATCACTAGATACATATCTGA	1415
JAK2	gJAK2_126	GCACATACATTCCCATGAATA	1416
JAK2	gJAK2_132	AATGCATTTCAGGTGGTACCCA	1417
JAK2	gJAK2_137	CCACAAAGTGGTACCAAACT	1418
JAK2	gJAK2_175	AAGATAGTCTCGTAACTTCC	1419
JAK2	gJAK2_187	GGTTAACCAGTCTTGCCAC	1420
JAK2	gJAK2_191	CAGGTATGCTCCAGAACTACT	1421
mir-101-2	gmir-101-2_001	GGTTATCATGGTACCGATGCT	1422
mir-101-2	gmir-101-2_002	AGATATACAGCATCGGTACCA	1423
mir-101-2	gmir-101-2_003	TCAATGTGATGGCACCACCAT	1424
MLANA	gMLANA_001	AACTTACTCTTCAGCCGTGGT	1425
MLANA	gMLANA_002	TCTATCTCTTGGGCCAGGGCC	1426
MLANA	gMLANA_003	GTCTTCTACAATACCAACAGC	1427
MLANA	gMLANA_004	CCAACCATCAAGGCTCTGTAT	1428
MLANA	gMLANA_008	CATTTTCAGGATAAAAAGTCTTC	1429
MLANA	gMLANA_009	AGGATAAAAGTCTTCATGTTG	1430
MLANA	gMLANA_010	CTGTCCCGATGATCAAACCCT	1431
MLANA	gMLANA_011	TCTTGAAGAGACACTTTGCTG	1432
MLANA	gMLANA_012	ATCATCGGGACAGCAAAGTGT	1433
MLANA	gMLANA_020	TCATAAGCAGGTGGAGCATTG	1434
PSMB5	gPSMB5_001	TGCCCACACTAGACATGGCGC	1435
PSMB5	gPSMB5_002	GGACTTGGGGGTCGTGCAGAT	1436
PSMB5	gPSMB5_003	GATTCCCTGGCTCTTCTGGGAC	1437
PSMB5	gPSMB5_005	CTCTGATCTTAACAGTTCCGC	1438
PSMB5	gPSMB5_006	GAAGCTCATAGATTTCGACATT	1439
PSMB5	gPSMB5_007	GAGGCAGCTGCTACAGAGATG	1440
PSMB5	gPSMB5_008	TACTGATACACCATGTTGGCA	1441
PSMB5	gPSMB5_010	CAGGCCTCTACTACGTGGACA	1442
PSMB5	gPSMB5_011	AGGGGCCACCTTCTCTGTAGG	1443
PSMB5	gPSMB5_012	AGGGGGTAGAGCCACTATACT	1444
PSMB8	gPSMB8_001	TCTATGCGATCTCCAGAGCTC	1445
PSMB8	gPSMB8_004	TCTTATCAGCCCACAGAATTC	1446
PSMB8	gPSMB8_005	TCCGTCCCCACCCAGGGACTG	1447
PSMB8	gPSMB8_008	AGTGTGCGCAGCCTCCAAGCT	1448

Target Gene	crRNA	Spacer sequence	SEQ ID NO
PSMB8	gPSMB8_010	ATCTTATAGGGTCCTGGACTC	1449
PSMB8	gPSMB8_011	CTGAGAGCCGAGTCCCATGTT	1450
PSMB8	gPSMB8_012	TCATTTGTCCACAGTGTACCA	1451
PSMB8	gPSMB8_013	ACCCAACCATCTTCCTTCATG	1452
PSMB8	gPSMB8_014	TCCACAGTGTACCACATGAAG	1453
PSMB8	gPSMB8_015	TACTTTCACCCAACCATCTTC	1454
PSMB9	gPSMB9_001	ACGGGGGCGTTGTGATGGGTT	1455
PSMB9	gPSMB9_002	CTCACCTGCAGACACTCGGG	1456
PSMB9	gPSMB9_005	CCTCAGGATAGAAGTGGAGGA	1457
PSMB9	gPSMB9_007	TCACCACATTTGCAGCAGCCA	1458
PSMB9	gPSMB9_009	GCTGCTGCAAATGTGGTGAGA	1459
PSMB9	gPSMB9_010	GGAGAACTCACCTGACCTCC	1460
PSMB9	gPSMB9_011	ACCTGAGGATCCCTTTCCCAG	1461
PSMB9	gPSMB9_012	CCAGGTATATGGAACCCTGGG	1462
PSMB9	gPSMB9_014	TCTATGGTTATGTGGATGCAG	1463
PSMB9	gPSMB9_015	GCAGTTCATTGCCCAAGATGA	1464
PTCD2	gPTCD2_005	ACCACATTATCTGTAAGTAGG	1465
PTCD2	gPTCD2_007	GCTAAAAGATACCTACTTACA	1466
PTCD2	gPTCD2_011	GTGCCAGAAAGATTACATGCA	1467
PTCD2	gPTCD2_018	ATTACCAGGTACCATGCAGAG	1468
PTCD2	gPTCD2_026	TTCTCAGACTCCACATCATTC	1469
PTCD2	gPTCD2_032	ATCTCTATCAATACTTGCAAA	1470
PTCD2	gPTCD2_033	GCAGGTGCTTTGCAAGTATTG	1471
PTCD2	gPTCD2_042	CCTGATTTCAGAGCTAATGCCA	1472
PTCD2	gPTCD2_043	GCTGTGGCATTAGCTCTGAAT	1473
PTCD2	gPTCD2_064	ATAGCAACGTGTGAGATTTCC	1474
RFX5	gRFX5_008	TGTAGCTCAGAGCCAAGTACA	1475
RFX5	gRFX5_012	GCAAGATCATCAGAGAGATCT	1476
RFX5	gRFX5_013	ACTTGCATCAGATATTGCTAC	1477
RFX5	gRFX5_015	GTACTTACACTCTCAGAACCC	1478
RFX5	gRFX5_016	AGGATCCGCTCTGCCCAGTCA	1479
RFX5	gRFX5_017	GTACCTCTGCAGAAAGAGGACG	1480
RFX5	gRFX5_018	GATGACCGTTCCCGAGGTGCA	1481
RFX5	gRFX5_026	GCTGGTGGAGCCTGCCCACTG	1482
RFX5	gRFX5_028	GCATCACTTGCTGTATCCTCT	1483
RFX5	gRFX5_038	GCTTCTGCTGCCCTTGATGAC	1484
RFXANK	gRFXANK_001	CCCATGGAGCTTACCCAGCCT	1485
RFXANK	gRFXANK_002	CCTGCACCCCTGAGCCTGTGA	1486
RFXANK	gRFXANK_003	CCAGCAGGCAGCTCCCTGAAG	1487
RFXANK	gRFXANK_005	GAGAGATTGAGACCGTTCGCT	1488
RFXANK	gRFXANK_006	CCAGGATGTGGGGGTCGGCAC	1489
RFXANK	gRFXANK_007	TCCTGCCCTTACCCACGACAG	1490
RFXANK	gRFXANK_008	ACGTGGTTCCCGCGCACAGCG	1491
RFXANK	gRFXANK_009	CAGCCCGAGGCGCTGACCTCA	1492
RFXANK	gRFXANK_010	CGGTATCCCAGGGCCACGGCA	1493
RFXANK	gRFXANK_011	CCTGCCCCATCTCAGTGCAAC	1494
RFXAP	gRFXAP_001	GAGGATCTAGAGGACGAGGAG	1495
RFXAP	gRFXAP_004	TACTTGTCTTGTACATCTTG	1496
RFXAP	gRFXAP_005	CCGCGCTGCCAGTCGAGGCAG	1497
RFXAP	gRFXAP_009	ACAATGGAGAGTATGTTATCT	1498

Target Gene	crRNA	Spacer sequence	SEQ ID NO
RFXAP	gRFXAP_012	GGGATCGTCCTGCAAGACCTA	1499
RFXAP	gRFXAP_016	GAACAAGTGTTAAATCAAAAA	1500
RFXAP	gRFXAP_020	TAAGTCGTTACTAAGAAGTCC	1501
RFXAP	gRFXAP_021	TGTAAAAATTGCACTACTTCT	1502
RFXAP	gRFXAP_023	CAGAAACAGCAACAGCTATTA	1503
RFXAP	gRFXAP_025	GAGCAAAGACAACAGCAGTTT	1504
RPL23	gRPL23_003	GCACCAGAGGACCCACCACGT	1505
RPL23	gRPL23_004	TATCCACAGGACGTGGTGGGT	1506
RPL23	gRPL23_008	TAGGAGCCAAAAACCTGTATA	1507
RPL23	gRPL23_013	GTTGTGCAATGACCACTGCTG	1508
RPL23	gRPL23_014	TTCTCTCAGTACATCCAGCAG	1509
RPL23	gRPL23_019	AAGATAATGCAGGAGTCATAG	1510
RPL23	gRPL23_021	CTACCTTTCATCTCGCCTTTA	1511
RPL23	gRPL23_025	ATGCAGGTTCTGCCATTACAG	1512
RPL23	gRPL23_026	CAAATATACTGGAGAATCATG	1513
RPL23	gRPL23_027	CCTTCCCTTTATATCCACAGG	1514
SOX10	gSOX10_001	CTGGCGCCGTTGACGCGCACG	1515
SOX10	gSOX10_002	TTGTGCTGCATACGGAGCCGC	1516
SOX10	gSOX10_003	ATGTGGCTGAGTTGGACCAGT	1517
SOX10	gSOX10_004	GCATCCACACCAGGTGGTGAG	1518
SOX10	gSOX10_005	ACTACTCTGACCATCAGCCCT	1519
SOX10	gSOX10_006	GGGCCGGGACAGTGTCGTATA	1520
SRP54	gSRP54_011	TCTTAGTTGCTTCACTAGTTT	1521
SRP54	gSRP54_020	GTGGGTGTCCATGCCTTAACT	1522
SRP54	gSRP54_021	GCTTGTAGACCCTGGAGTTAA	1523
SRP54	gSRP54_024	CCACTCCCTTGCAATCCAACA	1524
SRP54	gSRP54_029	TCACCCAGCTAGCATATTATT	1525
SRP54	gSRP54_030	ATATGTGCAGACACATTCAGA	1526
SRP54	gSRP54_064	ATTGGTACAGGGGAACATATA	1527
SRP54	gSRP54_087	GCACCATCCGTACTGTCTAGT	1528
SRP54	gSRP54_090	GTAAACAACCAGGAAGAATCC	1529
SRP54	gSRP54_096	CCCTCAGGTGGCGACATGTCT	1530
SRP54	gSRP54_139	AGGATAACTAACCAAGATCTG	1531
STAT1	gSTAT1_003	CATGGGAAAACCTGTCATCATA	1532
STAT1	gSTAT1_005	TAACCACTGTGCCAGGTACTG	1533
STAT1	gSTAT1_009	ATGACCTCCTGTCACAGCTGG	1534
STAT1	gSTAT1_013	TTCTAACCACTCAAATCTAGG	1535
STAT1	gSTAT1_014	AGGAAGACCCAATCCAGATGT	1536
STAT1	gSTAT1_026	TAGTGTATAGAGCATGAAATC	1537
STAT1	gSTAT1_032	TGATCACTCTTTGCCACACCA	1538
STAT1	gSTAT1_102	CCTGACATCATTCGCAATTAC	1539
STAT1	gSTAT1_103	GATACAGATACTTCAGGGGAT	1540
STAT1	gSTAT1_113	GTCACCCTTCTAGACTTCAGA	1541
Tap1	gTap1_011	GAGTGAAGGTATCGGCTGAGC	1542
Tap1	gTap1_012	AGCCCCCAGACCTGGCTATGG	1543
Tap1	gTap1_016	AGGAGAAACCTGTCTGGTTCT	1544
Tap1	gTap1_020	CTTCTGCCCAAGAAGGTGGGA	1545
Tap1	gTap1_026	GGGAAAAGCTGCAAGAAATAA	1546
Tap1	gTap1_030	AGGTATGCTGCTGAAAGTGGG	1547
Tap1	gTap1_033	TCTGAGGAGCCCACAGCCTTC	1548

Target Gene	crRNA	Spacer sequence	SEQ ID NO
Tap1	gTap1_035	GGTAGGCAAAGGAGACATCTT	1549
Tap1	gTap1_036	CCTACCCAAACCGCCCAGATG	1550
Tap1	gTap1_039	GAAGAAGTCTTCAAGAAAATA	1551
TAP2	gTAP2_004	GCAGCCCCCACAGCCCTCCCA	1552
TAP2	gTAP2_008	AGGTGAGACATTAATCCCTCA	1553
TAP2	gTAP2_014	AAGGAAGCCAGTTACTCATCA	1554
TAP2	gTAP2_027	CAGACCCTGGTATACATATAT	1555
TAP2	gTAP2_028	GCTGTCTGGTCCATGTAGGAGA	1556
TAP2	gTAP2_029	TCCTACATGGACCGACAGCCA	1557
TAP2	gTAP2_030	ACAACCCCTGCAGAGTGGTG	1558
TAP2	gTAP2_037	ATCCAGCAGCACCTGTCCCCC	1559
TAP2	gTAP2_038	AGTTGGGCAGGAGCCTGTGCT	1560
TAP2	gTAP2_040	TAGAAGATACCTGTGTATATT	1561
TAPBP	gTAPBP_001	CGCTCGCATCCTCCACGAACC	1562
TAPBP	gTAPBP_002	GCAGAGGCGGGGAGAGGCACG	1563
TAPBP	gTAPBP_003	CCTACATGCCCCCACCTCCG	1564
TAPBP	gTAPBP_004	GGCTAGAGTGGCGACGCCAGC	1565
TAPBP	gTAPBP_007	AGGAGGGCACCTATCTGGCCA	1566
TAPBP	gTAPBP_010	GTCTCTTTTCCCCAGAACCCC	1567
TAPBP	gTAPBP_011	CCCAGAACCCCCCAAAGTGTC	1568
TAPBP	gTAPBP_012	AGGGCCCTCCCTTGAGGACAG	1569
TAPBP	gTAPBP_013	CTGTCTGCCTTTCTTCTGCTT	1570
TAPBP	gTAPBP_016	CCCACAGCTGTCTACCTGTCC	1571
TWF1	gTWF1_005	CACAGCAAGTGAAGATGTAA	1572
TWF1	gTWF1_012	ATAGAGCAACTTGTGATTGGA	1573
TWF1	gTWF1_015	CCCCTGTTGGAGGACAAACAA	1574
TWF1	gTWF1_018	ATGTGGCCACCTCCAAATTCC	1575
TWF1	gTWF1_020	GAGGTGGCCACATTAAAGATG	1576
TWF1	gTWF1_022	ATCTGTCTAGTTCTTCCTCA	1577
TWF1	gTWF1_051	CAGATCGAGATAGACAATGGG	1578
TWF1	gTWF1_053	TGAAGAAGTACATCCCAAGCA	1579
TWF1	gTWF1_060	ATGTGATGACTTTAATCAGTA	1580
TWF1	gTWF1_101	AAATAGGTGGGCTACCTTTCT	1581
CD3D	gCD3D_001	TCTCTGGCCTGGTACTGGCTA	1582
CD3D	gCD3D_002	CCCTTTAGTGAGCCCCTTCAA	1583
CD3D	gCD3D_003	GTGAGCCCCTTCAAGATACCT	1584
CD3D	gCD3D_004	TGAATTGCAATACCAGCATCA	1585
CD3D	gCD3D_005	CCAGGTCCAGTCTTGTAATGT	1586
CD3D	gCD3D_006	TCCTTGTATATATCTGTCCCA	1587
CD3D	gCD3D_007	GGAGTCTTCTGCTTTGCTGGA	1588
CD3D	gCD3D_008	CTGGACATGAGACTGGAAGGC	1589
CD3D	gCD3D_009	TCTTCTCCTCTCTTAGCCCCT	1590
CD3D	gCD3D_010	CTCCAAGGTGGCTGTACTGAG	1591
NLRC5	gNLRC5_001	GCTCCTGTAGCGCTGCTGGGC	1592
NLRC5	gNLRC5_002	GGGAAGGCTGGCATGGGCAAG	1593
NLRC5	gNLRC5_003	CAGGCCCTGTTCCCTTTTGA	1594
NLRC5	gNLRC5_004	AATTCCGCCAGCTCAACTTGA	1595
NLRC5	gNLRC5_005	ATCTGTACCTGAGCCCTGAAT	1596
NLRC5	gNLRC5_006	ATGGGCTAGATGAGGCCCTCC	1597
NLRC5	gNLRC5_007	TCCCATCTCTGCAATGGGACC	1598

Target Gene	crRNA	Spacer sequence	SEQ ID NO
NLRC5	gNLRC5_008	ATGGGCCACGGGTGGAAGAAT	1599
NLRC5	gNLRC5_009	TCTGTAACCTCCACCAGGGCCC	1600
NLRC5	gNLRC5_010	CATAGAAGATAACCTTCCCTG	1601
NLRC5	gNLRC5_011	GGGCCACTCACAGCCTGCTGA	1602
NLRC5	gNLRC5_012	ACCCACCTCAGCCTGCAGGAG	1603
NLRC5	gNLRC5_013	TTCACCTTGGGGCTGGCCATC	1604
NLRC5	gNLRC5_014	TTGCTGCCCTGCACCTGATGG	1605
NLRC5	gNLRC5_015	GTCCGCTGTACCCAGCGGGAA	1606
NLRC5	gNLRC5_016	GCCCTGTGAGCTTGCGGGTGG	1607
NLRC5	gNLRC5_017	TGCGGTGAGACTGGCCAGCTC	1608
NLRC5	gNLRC5_018	CCACTGACCTGCACCGACCTG	1609
NLRC5	gNLRC5_019	ATGGCTGTCCCCTGGAGCCCC	1610

[0161] The spacer sequences provided in **Tables 1-3** are designed based upon identification of target nucleotide sequences associated with a PAM in a given target gene locus, and are selected based upon the editing efficiency detected in human cells.

- 5 [0162] Further exemplary spacer sequences useful in embodiments of the methods and compositions disclosed herein are shown in **Tables 4-23**.

Table 4 Tested crRNAs Targeting Human ADORA2A Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gADORA2A_1	GTGGTGTCACTGGCGGCGGCC	242	0.3
gADORA2A_2	TGGTGTCACTGGCGGCGGCCG	133	3.9
gADORA2A_3	GCCATCACCATCAGCACCGGG	243	0.5
gADORA2A_4	CCATCACCATCAGCACCGGGT	137	2.1
gADORA2A_5	GTCCTGGTCCTCACGCAGAGC	244	0.1
gADORA2A_6	GCCCTCGTGCCGGTCACCAAG	245	0.9
gADORA2A_7	GTGACCGGCACGAGGGCTAAG	135	2.8
gADORA2A_8	CCATCGGCCTGACTCCCATGC	136	2.2
gADORA2A_9	GCTGACCGCAGTTGTTCCAAC	246	1.1
gADORA2A_10	GGCTGACCGCAGTTGTTCCAA	247	0.5
gADORA2A_11	GCCCTCCCCGCAGCCCTGGGA	248	1.3
gADORA2A_12	AGGATGTGGTCCCCATGAACT	51	18.2
gADORA2A_13	AACTTCTTTGCCTGTGTGCTG	249	0.1
gADORA2A_14	TTTGCCTGTGTGCTGGTGCCC	250	0.2
gADORA2A_15	CCTGTGTGCTGGTGCCCCTGC	251	1.1
gADORA2A_16	CGGATCTTCCTGGCGGCGCGA	131	7.8
gADORA2A_17	AGCTGTCGTGCGCCGCCAGG	252	0.1
gADORA2A_18	TGCAGTGTGGACCGTGCCCGC	253	0.2
gADORA2A_19	GCAGCATGGACCTCCTTCTGC	254	0.4
gADORA2A_20	CCCTCTGCTGGCTGCCCCTAC	255	0.6
gADORA2A_21	ACTTTCTTCTGCCCCGACTGC	256	0.6
gADORA2A_22	CTTCTGCCCCGACTGCAGCCA	257	1.0

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gADORA2A_23	TTCTGCCCCGACTGCAGCCAC	134	2.8
gADORA2A_24	ATCTACGCCTACCGTATCCGC	258	0.0
gADORA2A_25	CGCAAGATCATTCGCAGCCAC	259	0.1
gADORA2A_26	AAAGGTTCTTGCTGCCTCAGG	260	0.1
gADORA2A_27	CAAGGCAGCTGGCACCAGTGC	261	0.1
gADORA2A_28	AAGGCAGCTGGCACCAGTGCC	132	5.8
gADORA2A_29	AGCTCATGGCTAAGGAGCTCC	262	0.2
gADORA2A_30	GCCATGAGCTCAAGGGAGTGT	263	0.5

Table 5 Tested crRNAs Targeting Human B2M Gene

crRNA Name	Spacer Sequence	SEQ ID NO	% Indel
gB2M_1	GCTGTGCTCGCGCTACTCTCT	145	1.8
gB2M_2	TGGCCTGGAGGCTATCCAGCG	65	17.4
gB2M_3	CCCGATATTCCTCAGGTACTC	264	0.1
gB2M_4	CTCACGTCATCCAGCAGAGAA	52	74.1
gB2M_5	CATTCTCTGCTGGATGACGTG	142	2.2
gB2M_6	CCATTCTCTGCTGGATGACGT	265	1.0
gB2M_7	ACTTTCATTCTCTGCTGGAT	64	17.9
gB2M_8	CTGAATTGCTATGTGTCTGGG	139	3.5
gB2M_9	AATGTCGGATGGATGAAACCC	266	0.5
gB2M_10	ATCCATCCGACATTGAAGTTG	143	2.0
gB2M_11	CTGAAGAATGGAGAGAGAATT	140	3.4
gB2M_12	TCAATTCTCTCTCCATTCTTC	267	0.7
gB2M_13	TTCAATTCTCTCTCCATTCTT	268	0.7
gB2M_14	CTGAAAGACAAGTCTGAATGC	269	0.4
gB2M_15	TCTTTCAGCAAGGACTGGTCT	270	0.9
gB2M_16	AGCAAGGACTGGTCTTTCTAT	271	0.3
gB2M_17	TATCTCTTGTAACACTGAA	66	15.3
gB2M_18	TCAGTGGGGGTGAATTCAGTG	141	3.0
gB2M_19	ACTATCTTGGGCTGTGACAAA	272	0.1
gB2M_20	GTCACAGCCCAAGATAGTTAA	273	0.8
gB2M_21	TCACAGCCCAAGATAGTTAAG	138	5.3
gB2M_22	CCCCACTTAACATCTTGGGC	144	2.0
gB2M_23	CTGGCCTGGAGGCTATCCAGC	618	0.77
gB2M_24	TCCCGATATTCCTCAGGTACT	619	0.54
gB2M_25	CCGATATTCCTCAGGTACTCC	620	0.14
gB2M_26	AGTAAGTCAACTTCAATGTCG	621	0.11
gB2M_27	AATTCTCTCTCCATTCTTCAG	622	2.70
gB2M_28	CAATTCTCTCTCCATTCTTCA	623	0.26
gB2M_29	CAGCAAGGACTGGTCTTTCTA	624	0.19
gB2M_30	AGTGGGGGTGAATTCAGTGTA	625	91.96
gB2M_31	CAGTGGGGGTGAATTCAGTGT	626	8.10

crRNA Name	Spacer Sequence	SEQ ID NO	% Indel
gB2M_33	CTATCTCTTGTA TACTACTGA	627	0.21
gB2M_34	TACTACTGAATTCACCCCC	628	0.80
gB2M_35	GGCTGTGACAAAGTCACATGG	629	0.18
gB2M_36	CAAAAGAATGTAAGACTTACC	630	0.13
gB2M_37	CCTCCATGATGCTGCTTACAT	631	0.81
gB2M_38	TCATAGATCGAGACATGTAA	632	0.18
gB2M_39	TCATAGATCGAGACATGTAAAG	633	0.20
gB2M_40	CATAGATCGAGACATGTAAAGC	634	4.25
gB2M_41	ATAGATCGAGACATGTAAAGCA	635	93.92

Table 6 Tested crRNAs Targeting Human CD52 Gene

crRNA Name	Spacer Sequence	SEQ ID NO	% Indel
gCD52_1	CTCTTCCTCCTACTCACCATC	53	28.4
gCD52_2	TCCTCCTACAGATACAACTG	274	N.D.
gCD52_3	GTCCTGAGAGTCCAGTTTGTA	275	N.D.
gCD52_4	GCTGGTGTCGTTTTGTCCTGA	146	4.1
gCD52_5	TGTTGCTGGATGCTGAGGGGC	276	1.1
gCD52_6	CCTTTTCTTCGTGGCCAATGC	277	0.2
gCD52_7	TCTTCGTGGCCAATGCCATAA	278	0.2
gCD52_8	CTTCGTGGCCAATGCCATAAT	279	0.15

Table 7 Tested crRNAs Targeting Human CIITA Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCIITA_1	GGGCTCTGACAGGTAGGACCC	280	0.5
gCIITA_2	TACCTTGGGGCTCTGACAGGT	281	0.0
gCIITA_3	TTACCTTGGGGCTCTGACAGG	282	0.0
gCIITA_4	TAGGGGCCCAACTCCATGGT	54	13.5
gCIITA_5	TTAACAGCGATGCTGACCCCC	284	0.1
gCIITA_6	TATGACCAGATGGACCTGGCT	285	0.2
gCIITA_7	TCCTCCCAGAACCCGACACAG	286	0.1
gCIITA_8	CCTCCCAGAACCCGACACAGA	287	0.1
gCIITA_9	CATGTCACACAACAGCCTGCT	288	0.1
gCIITA_10	CTCACCAGATATTGGCATAAGC	289	0.1
gCIITA_11	TCCTTGTCTGGGCAGCGGAAC	290	0.1
gCIITA_12	CCTTGTCTGGGCAGCGGAAC	291	0.4
gCIITA_13	TCTGGGCAGCGGAACCTGGACC	292	0.1
gCIITA_14	CTCAGGCCCTCCAGCTGGGAG	293	0.2
gCIITA_15	CTGAAAATGTCCTTGCTCAGG	294	0.2
gCIITA_16	TCTCAAAGTAGAGCACATAGG	295	0.1

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCIITA_17	ATCTGGTCCTATGTGCTCTAC	296	0.2
gCIITA_18	TGCTGGCATCTCCATACTCTC	147	4.8
gCIITA_19	CTGCCCCAACTTCTGCTGGCAT	297	0.5
gCIITA_20	TCTGCCCCAACTTCTGCTGGCA	298	0.1
gCIITA_21	CTGACTTTTCTGCCCCAACTTC	299	0.1
gCIITA_22	CTCTGCAGCCTTCCCAGAGGA	300	0.6
gCIITA_23	CCAGAGGAGCTTCCGGCAGAC	301	0.9
gCIITA_24	AGGTCTGCCGGAAGCTCCTCT	302	0.1
gCIITA_25	CAGTGCTTCAGGTCTGCCGGA	303	0.2
gCIITA_26	CGGCAGACCTGAAGCACTGGA	304	0.3
gCIITA_27	CTCACAGCTGAGCCCCCACT	305	0.4
gCIITA_28	CTCCAGGCGCATCTGGCCGGA	306	0.7
gCIITA_29	GTCTCTTGACAGTGCCTTTCTC	148	2.4
gCIITA_30	TCTCTTGACAGTGCCTTTCTCC	307	0.1
gCIITA_31	CTCCAGTTCCTCGTTGAGCTG	308	0.1
gCIITA_32	CCTTGGGGCTCTGACAGGTAG	636	93.85
gCIITA_33	ACCTTGGGGCTCTGACAGGTA	637	11.83
gCIITA_34	CCGGCCTTTTTACCTTGGGGC	638	2.26
gCIITA_35	CTCCCAGAACCCGACACAGAC	639	48.70
gCIITA_36	TGGGCTCAGGTGCTTCCTCAC	640	85.46
gCIITA_37	CTGGGCTCAGGTGCTTCCTCA	641	0.45
gCIITA_38	CTTGTCTGGGCAGCGGAACTG	642	38.38
gCIITA_39	CTCAAAGTAGAGCACATAGGA	643	0.25
gCIITA_40	TCAAAGTAGAGCACATAGGAC	644	15.68
gCIITA_41	TGCCCCAACTTCTGCTGGCATC	645	46.21
gCIITA_42	TGACTTTTCTGCCCCAACTTCT	646	2.72
gCIITA_43	TCTGCAGCCTTCCCAGAGGAG	647	55.09
gCIITA_44	TCCAGGCGCATCTGGCCGGAG	648	39.16
gCIITA_45	TCCAGTTCCTCGTTGAGCTGC	649	0.22
gCIITA_46	CCAGAGCCCATGGGGCAGAGT	650	1.51
gCIITA_47	TCCCCACCATCTCCACTCTGC	651	2.05
gCIITA_48	CTCGGGAGGTCAGGGCAGGTT	652	61.63
gCIITA_49	GAAGCTTGTTGGAGACCTCTC	653	0.67
gCIITA_50	GGAAGCTTGTTGGAGACCTCT	654	0.57
gCIITA_51	CAGAGCCGGTGGAGCAGTTCT	655	8.94
gCIITA_52	CCCAGCACAGCAATCACTCGT	656	2.63
gCIITA_53	TCTTCTCTGTCCCCTGCCATT	657	0.28
gCIITA_55	AGCCACATCTTGAAGAGACCT	658	5.71
gCIITA_56	CCAGAAGAAGCTGCTCCGAGG	659	0.52
gCIITA_57	CAGAAGAAGCTGCTCCGAGGT	660	12.02
gCIITA_58	AGCTGTCCGGCTTCTCCATGG	661	3.25
gCIITA_59	AGAGCTCAGGGATGACAGAGC	662	16.35
gCIITA_60	TGCCGGGCAGTGTGCCAGCTC	663	11.98
gCIITA_61	ATGTCTGCGGCCAGCTCCCA	664	1.25

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCIITA_62	GCCATCGCCCAGGTCCTCACG	665	1.29
gCIITA_63	GCCACTCAGAGCCAGCCACAG	666	35.47
gCIITA_64	TGGCTGGGCTGATCTTCCAGC	667	0.50
gCIITA_65	GCAGCACGTGGTACAGGAGCT	668	70.73
gCIITA_66	CTGGGCACCCGCCTCACGCCT	669	0.31
gCIITA_67	TGGGCACCCGCCTCACGCCTC	670	12.57
gCIITA_68	CCCCTCTGGATTGGGGAGCCT	671	4.61
gCIITA_69	AAAGGCTCGATGGTGAAGTTC	672	1.17
gCIITA_70	CCAGGTCTTCCACATCCTTCA	673	38.98
gCIITA_71	AAAGCCAAGTCCCTGAAGGAT	674	39.50
gCIITA_72	GGTCCCGAACAGCAGGGAGCT	675	89.25
gCIITA_73	TTAGGTCCCGAACAGCAGGG	676	10.88
gCIITA_74	CTTACGCAAACTCCAGTTTCT	677	0.79
gCIITA_75	CCTCCTAGGCTGGGCCCTGTC	678	2.78
gCIITA_76	GGGAAAGCCTGGGGGCCTGAG	679	68.93
gCIITA_77	CCCAAAGTGGTGCGGATCCTC	680	0.57
gCIITA_79	CTCCCTGCAGCATCTGGAGTG	681	1.12
gCIITA_80	CAAGGACTTCAGCTGGGGGAA	682	87.87
gCIITA_81	TAGGCACCCAGGTCAGTGATG	683	44.56
gCIITA_82	CGACAGCTTGTACAATAACTG	684	34.37
gCIITA_83	TCTTGCCAGCGTCCAGTACAA	685	5.62
gCIITA_84	CCCGGCCTTTTACCTTGGGG	686	0.38
gCIITA_85	CCTCCCAGGCAGCTCACAGTG	687	0.74
gCIITA_87	TCCAGCCAGGTCCATCTGGTC	688	0.15
gCIITA_88	TTCTCCAGCCAGGTCCATCTG	689	0.21
gCIITA_89	ATCACCTTCCATGTACACAA	690	0.31
gCIITA_90	TCTGGGCTCAGGTGCTTCCTC	691	0.25
gCIITA_91	TGCCAATATCGGTGAGGAAGC	692	0.17
gCIITA_92	CAGGACTCCCAGCTGGAGGGC	693	0.61
gCIITA_93	TCTGACTTTTCTGCCCAACTT	694	0.21
gCIITA_94	CAGTGCCTTTCTCCAGTTCTT	695	0.25
gCIITA_95	GCTGGCCTGGGGCACCTCACC	696	0.59
gCIITA_96	GCTCCATCAGCCACTGACCTG	697	0.29
gCIITA_97	CCTGTCATGTTTGCTCGGGAG	698	0.27
gCIITA_98	TCCATCTCCAGAGCACAAAGAC	699	0.23
gCIITA_99	TTGGAGACCTCTCCAGCTGCC	700	0.99
gCIITA_100	GCAGAGCCGGTGGAGCAGTTC	701	0.46
gCIITA_101	CTGCTGCTCCTCTCCAGCCTG	702	0.23
gCIITA_103	GCAGCCAACAGCACCTCAGCC	703	0.22
gCIITA_104	GCCCAGCACAGCAATCACTCG	704	0.07

Table 8 Tested crRNAs Targeting Human CTLA4 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCTLA4_1	TGCCGCTGAAATCCAAGGCAA	309	1.3
gCTLA4_2	CCTTGGATTTTCAGCGGCACAA	310	0.8
gCTLA4_3	GATTTTCAGCGGCACAAGGCTC	311	0.6
gCTLA4_4	AGCGGCACAAGGCTCAGCTGA	795	58.4
gCTLA4_5	TTCTTCTCTTCATCCCTGTCT	155	1.7
gCTLA4_6	CAGAAGACAGGGATGAAGAGA	797	44.6
gCTLA4_7	GCAGAAGACAGGGATGAAGAG	312	0.2
gCTLA4_8	GGCTTTTCCATGCTAGCAATG	313	0.1
gCTLA4_9	GCTTTTCCATGCTAGCAATGC	314	0.2
gCTLA4_10	TCCATGCTAGCAATGCACGTG	315	0.1
gCTLA4_11	CCATGCTAGCAATGCACGTGG	316	0.1
gCTLA4_12	GTGTGTGAGTATGCATCTCCA	317	0.8
gCTLA4_13	TGTGTGAGTATGCATCTCCAG	70	12.6
gCTLA4_14	CCTGGAGATGCATACTCACAC	67	47.4
gCTLA4_15	GCCTGGAGATGCATACTCACA	318	0.2
gCTLA4_16	GGCAGGCTGACAGCCAGGTGA	319	1.2
gCTLA4_17	AGTCACCTGGCTGTCAGCCTG	320	0.4
gCTLA4_18	CTAGATGATTCCATCTGCACG	154	2.0
gCTLA4_19	CACTGGAGGTGCCCGTGCAGA	798	42.5
gCTLA4_20	ATTTCCACTGGAGGTGCCCGT	321	0.1
gCTLA4_21	GATAGTGAGGTTCACCTTGATT	322	0.6
gCTLA4_22	CAGATGTAGAGTCCCGTGTCC	323	0.6
gCTLA4_23	CTCACCAATTACATAAATCTG	324	0.8
gCTLA4_24	GCTCACCAATTACATAAATCT	325	1.0
gCTLA4_25	GTTTTCTGTTGCAGATCCAGA	326	0.1
gCTLA4_26	TTTTCTGTTGCAGATCCAGAA	327	0.1
gCTLA4_27	CTGTTGCAGATCCAGAACCGT	149	5.0
gCTLA4_28	CTCCTCTGGATCCTTGACAGCA	152	3.0
gCTLA4_29	CAGCAGTTAGTTCGGGGTTGT	328	0.7
gCTLA4_30	TTTATAGCTTTCTCCTCACAG	329	0.6
gCTLA4_31	CTCCTCACAGCTGTTTCTTTG	330	1.0
gCTLA4_32	TCCTCACAGCTGTTTCTTTGA	331	0.7
gCTLA4_33	GCTCAAAGAAACAGCTGTGAG	332	0.8
gCTLA4_34	TTTTTGTGTTTGACAGCTAAA	333	0.5
gCTLA4_35	TGTGTTTGACAGCTAAAGAAA	334	0.1
gCTLA4_36	ACAGCTAAAGAAAAGAAGCCC	150	3.9
gCTLA4_37	CACATAGACCCCTGTTGTAAG	153	2.9
gCTLA4_38	CACATTCTGGCTCTGTTGGGG	335	0.2
gCTLA4_39	TCACATTCTGGCTCTGTTGGG	336	0.3
gCTLA4_40	AGCCTTATTTTATTCCCATCA	337	0.3

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCTLA4_41	TCAATTGATGGGAATAAAATA	151	3.0

Table 9 Tested crRNAs Targeting Human DCK Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gDCK_1	TCTTGGGCGGGTGGCCATTC	338	0.1
gDCK_2	TCAGCCAGCTCTGAGGGGACC	71	50.4
gDCK_3	CTTGATGCGGGTCCCCTCAGA	339	0.3
gDCK_4	GATGGAGATTTTCTTGATGCG	340	0.3
gDCK_5	CCGATGTTCCCTTCGATGGAG	341	0.5
gDCK_6	CGGAGGCTCCTTACCGATGTT	796	85.1
gDCK_7	ATCTTTCCTCACAACAGCTGC	159	1.5
gDCK_8	CTCACAACAGCTGCAGGGAAG	72	31.7
gDCK_9	AGGATATTCACAAATGTTGAC	156	8.1
gDCK_10	TGAATATCCTTAAACAATTGT	342	1.0
gDCK_11	CCAATCTTCACACAATTGTTT	343	0.1
gDCK_12	AACAATTGTGTGAAGATTGGG	344	0.8
gDCK_13	AACATTGCACCATCTGGCAAC	345	1.2
gDCK_14	GAACATTGCACCATCTGGCAA	346	0.6
gDCK_15	CATACCTCAAATTCATCTTGA	347	0.3
gDCK_16	ATTTTCATACCTCAAATTCAT	348	0.1
gDCK_17	AATTTTATTTTCATACCTCAA	349	0.0
gDCK_18	TGCACATTCAAATAGGAACT	350	0.4
gDCK_19	TCTGAGACATTGTAAGTTCCT	351	0.7
gDCK_20	CAATGTCTCAGAAAAATGGTG	352	0.6
gDCK_21	TCATACATCATCTGAAGAACA	158	3.6
gDCK_22	GAAGGTAAAAGACCATCGTTC	157	5.6
gDCK_23	ACCTTCCAAACATATGCCTGT	353	1.2
gDCK_24	CAAACATATGCCTGTCTCAGT	354	1.1
gDCK_25	CCATTCAGAGAGGCAAGCTGA	355	0.9
gDCK_26	AGCTTGCCATTCAGAGAGGCA	73	13.3
gDCK_27	CCTCTCTGAATGGCAAGCTCA	356	1.1
gDCK_28	TCTGCATCTTTGAGCTTGCCA	357	0.1
gDCK_29	TTGAACGATCTGTGTATAGTG	358	0.2
gDCK_30	TACATACCTGTCACATACAC	74	12.8
gDCK_31	AGGTATATTTTGCATCTAAT	359	0.05

Table 10 Tested crRNAs Targeting Human FAS Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gFAS_1	GGAGGATTGCTCAACAACCAT	78	22.6

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gFAS_2	TATTTTACAGGTTCTTACGTC	360	0.1
gFAS_3	ATTTTACAGGTTCTTACGTC	361	0.7
gFAS_4	ACAGGTTCTTACGTCTGTTGC	172	1.5
gFAS_5	GGACGATAATCTAGCAACAGA	165	1.9
gFAS_6	TGGACGATAATCTAGCAACAG	362	0.0
gFAS_7	GGCATTAACTTTTGGACGA	363	0.1
gFAS_8	GAGTTGATGTCAGTCACTTGG	364	0.1
gFAS_9	CAAGTTCTGAGTCTCAACTGT	365	0.1
gFAS_10	GAAGGCCTGCATCATGATGGC	163	2.4
gFAS_11	TGGCAGAATTGGCCATCATGA	366	0.8
gFAS_12	GTGTAACATACCTGGAGGACA	77	29.9
gFAS_13	TTTCCTTGGGCAGGTGAAAGG	367	1.1
gFAS_14	TTTCCTTGGGCAGGTGAAAGGA	166	1.7
gFAS_15	GGCAGGTGAAAGGAAAGCTAG	173	1.5
gFAS_16	TTGGCAGGGCACGCAGTCTGG	368	0.7
gFAS_17	CCTTCTTGGCAGGGCACGCAG	369	0.8
gFAS_18	TCTGTGTACTCCTTCCCTTCT	370	1.0
gFAS_19	GTCTGTGTACTCCTTCCCTTC	371	0.6
gFAS_20	GAAGAAAAATGGGCTTTGTCT	372	0.7
gFAS_21	TCTTCCAAATGCAGAAGATGT	373	0.7
gFAS_22	ATCACACAATCTACATCTTCT	374	0.5
gFAS_23	AAGACTCTTACCATGTCCTTC	375	0.6
gFAS_24	CAAACCTGATTTTCTAGGCTTA	376	0.1
gFAS_25	CTAGGCTTAGAAGTGGAATA	162	3.5
gFAS_26	GAAGTGGAAATAAACTGCACC	377	0.3
gFAS_27	GTATTCTGGGTCCGGGTGCAG	378	1.3
gFAS_28	CATCTGCACTTGGTATTCTGG	379	1.2
gFAS_29	GTTTACATCTGCACTTGGTAT	167	1.6
gFAS_30	TTTTGTAACTCTACTGTATGT	380	0.8
gFAS_31	TTTGTAACCTCTACTGTATGTG	381	1.4
gFAS_32	GTGCAAGGGTCACAGTGTTCA	164	2.4
gFAS_33	CTTGGTGCAAGGGTCACAGTG	168	1.6
gFAS_34	TTTTTCTAGATGTGAACATGG	75	59.1
gFAS_35	ATGATTCCATGTTACATCTA	76	58.5
gFAS_36	GTGTTGCTGGTGAGTGTGCAT	57	61.9
gFAS_37	CACTTGGTGTTGCTGGTGAGT	382	1.3
gFAS_38	CTCTTTGCACTTGGTGTTGCT	170	1.5
gFAS_39	GGGTGGCTTTGTCTTCTTCTT	383	0.1
gFAS_40	GTCTTCTTCTTTTGCCAATTTC	384	0.6
gFAS_41	TCTTCTTCTTTTGCCAATTCC	385	0.1
gFAS_42	GCCAATTCCACTAATTGTTTG	386	0.4
gFAS_43	CCCCAAACAATTAGTGGAATT	387	0.4
gFAS_44	AACAAAGCAAGAACTTACCCC	388	0.3
gFAS_45	TTTGTTCTTTCAGTGAAGAGA	161	6.0

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gFAS_46	TTCTTTCAGTGAAGAGAAAGG	389	0.9
gFAS_47	AGTGAAGAGAAAGGAAGTACA	160	9.8
gFAS_48	CTGTACTTCCTTTCTCTTCAC	390	0.8
gFAS_49	TGCATGTTTTCTGTACTTCCT	391	0.6
gFAS_50	CTGCATGTTTTCTGTACTTCC	392	0.4
gFAS_51	TGTGCTTTCTGCATGTTTTCT	393	0.3
gFAS_52	CTGTGCTTTCTGCATGTTTTCT	394	0.3
gFAS_53	CCTTTCTGTGCTTTCTGCATG	395	0.3
gFAS_54	GTTTTCCCTTTCTGTGCTTTCT	396	0.4
gFAS_55	AAGTTGGAGATTCATGAGAAC	397	0.4
gFAS_56	AATACCTACAGATTTAAAGT	398	0.3
gFAS_57	TTGCTTTCTAGGAAACAGTGG	399	1.1
gFAS_58	CTAGGAAACAGTGGCAATAAA	400	1.3
gFAS_59	TAGGAAACAGTGGCAATAAAT	79	11.0
gFAS_60	CCAGATAAATTTATTGCCACT	401	0.7
gFAS_61	CTATTTTTCAGATGTTGACTT	402	0.1
gFAS_62	TCAGATGTTGACTTGAGTAAA	403	0.6
gFAS_63	AGTAAATATATCACCCTATT	404	0.8
gFAS_64	AACTTGACTTAGTGTCAATGAC	405	0.4
gFAS_65	GAACAAAGCCTTTAACTTGAC	406	0.5
gFAS_66	GTTCGAAAGAATGGTGTCAAT	407	0.9
gFAS_67	ATTGACACCATTCTTTCGAAC	408	0.5
gFAS_68	TTCGAAAGAATGGTGTCAATG	409	0.7
gFAS_69	GGCTTCATTGACACCATTCTT	410	0.4
gFAS_70	TGTTCTGCTGTGTCTTGGACA	171	1.5
gFAS_71	CTGTTCTGCTGTGTCTTGGAC	169	1.5
gFAS_72	GTAATTGGCATCAACTTCATG	411	0.3
gFAS_73	CATGAAGTTGATGCCAATTAC	412	0.8
gFAS_74	TTCCATGAAGTTGATGCCAA	413	0.4
gFAS_75	TTTCTTTCCATGAAGTTGATG	414	0.5
gFAS_76	ATGGAAAGAAAGAAGCGTATG	415	1.3
gFAS_77	ATCAATGTGTCATACGCTTCT	416	0.8
gFAS_78	TTGAGATCTTTAATCAATGTG	417	1.0
gFAS_79	TTTGAGATCTTTAATCAATGT	418	0.9
gFAS_80	CTCTGCAAGAGTACAAAGATT	419	0.2
gFAS_81	TACTCTTGCAGAGAAAATTCA	420	0.2
gFAS_82	AGGATGATAGTCTGAATTTTC	421	0.4
gFAS_83	CTGAGTCACTAGTAATGTCCT	422	0.7
gFAS_84	AATTTTCTGAGTCACTAGTAA	423	0.6
gFAS_85	TGAAGTTTGAATTTTCTGAGT	424	0.4
gFAS_86	ATTTCTGAAGTTTGAATTTTC	425	0.3
gFAS_87	GATTTTCATTTCTGAAGTTTGA	426	0.5
gFAS_88	GGATTTTCATTTCTGAAGTTTG	427	0.5
gFAS_89	AGAAATGAAATCCAAAGCTTG	428	0.5

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gFAS_90	TCACTCTAGACCAAGCTTTGG	429	0.5
gFAS_91	TTGTTTTTCACTCTAGACCAA	430	0.7
gFAS_92	GTCTAGAGTGAAAAACAACAA	431	0.5

Table 11 Tested crRNAs Targeting Human HAVCR2 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTIM3_1	TCTTCTGCAAGCTCCATGTTT	432	0.1
gTIM3_2	TCTTCTGCAAGCTCCATGTTT	433	0.07
gTIM3_3	CTTCTGCAAGCTCCATGTTTT	434	0.1
gTIM3_4	CACATCTTCCCTTTGACTGTG	435	0.8
gTIM3_5	GACTGTGTCCTGCTGCTGCTG	436	0.8
gTIM3_6	TAAGTAGTAGCAGCAGCAGCA	799	53.7
gTIM3_7	CTTGTAAGTAGTAGCAGCAGC	58	64.4
gTIM3_8	TCTCTCTATGCAGGGTCTCTCA	437	0.1
gTIM3_9	TACACCCAGCCGCCCCAGGG	438	1.0
gTIM3_10	CCCCAGCAGACGGGCACGAGG	175	7.3
gTIM3_11	GCCCCAGCAGACGGGCACGAG	439	0.6
gTIM3_12	AATGTGGCAACGTGGTGCTCA	84	21.9
gTIM3_13	ATCAGTCCTGAGCACCACGTT	187	1.5
gTIM3_14	CATCAGTCCTGAGCACCACGT	440	0.1
gTIM3_15	GCCAGTATCTGGATGTCCAAT	181	2.9
gTIM3_16	CGGAAATCCCCATTTAGCCAG	441	0.4
gTIM3_17	GCGGAAATCCCCATTTAGCCA	442	0.1
gTIM3_18	CGCAAAGGAGATGTGTCCCTG	86	14.4
gTIM3_19	GATCCGGCAGCAGTAGATCCC	178	5.1
gTIM3_20	TCATCATTCATTATGCCTGGG	443	0.1
gTIM3_21	AGGTAAATTTTTCATCATTC	444	0.1
gTIM3_22	ATGACCAACTTCAGGTTAAAT	445	0.1
gTIM3_23	ACCTGAAGTTGGTCATCAAAC	184	2.2
gTIM3_24	TGTTGTTTCTGACATTAGCCA	446	0.7
gTIM3_25	TGACATTAGCCAAGGTCACCC	85	15.7
gTIM3_26	GAAAGGCTGCAGTGAAGTCTC	447	0.1
gTIM3_27	ACTGCAGCCTTTCCAAGGATG	182	2.6
gTIM3_28	CCAAGGATGCTTACCACCAGG	185	1.9
gTIM3_29	CAAGGATGCTTACCACCAGGG	80	59.8
gTIM3_30	CCACCAGGGGACATGGCCCAG	83	22.1
gTIM3_31	TATAGCAGAGACACAGACACT	448	0.3
gTIM3_32	TATCAGGGAGGCTCCCCAGTG	800	22.4
gTIM3_33	CTGTTAGATTTATATCAGGGA	449	1.4
gTIM3_34	TGTTTCCATAGCAAATATCCA	177	5.6
gTIM3_35	CATAGCAAATATCCACATTGG	450	1.0
gTIM3_36	CGGGACTCTGGAGCAACCATC	180	3.3

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTIM3_37	AAAATTAAAGCGCCGAAGATA	451	0.2
gTIM3_38	CATTTGAAAATTAAAGCGCCG	452	0.1
gTIM3_39	TGTTTCCCCCTTACTAGGGTA	453	0.7
gTIM3_40	GTTTCCCCCTTACTAGGGTAT	186	1.7
gTIM3_41	CCCCTTACTAGGGTATTCTCA	183	2.2
gTIM3_42	CTAGGGTATTCTCATAGCAAA	174	8.5
gTIM3_43	AATTCTGTATCTTCTCTTTGC	454	0.7
gTIM3_44	ATTTCCACAGCCTCATCTCTT	455	0.4
gTIM3_45	TTTCCACAGCCTCATCTCTTT	456	1.0
gTIM3_46	CACAGCCTCATCTCTTTGGCC	457	0.5
gTIM3_47	GCCAACCTCCCTCCCTCAGGA	176	6.0
gTIM3_48	CCAATCCTGAGGGAGGGAGGT	179	4.5
gTIM3_49	CTTCTGAGCGAATTCCTCTG	458	0.7
gTIM3_50	ATATACGTTCTCTTCAATGGT	459	0.5
gTIM3_51	GGGTTGTCGCTTTGCAATGCC	460	0.5

Table 12 Tested crRNAs Targeting Human LAG3 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gLAG3_1	CTGTTTCTGCAGCCGCTTTGG	461	0.2
gLAG3_2	TGCAGCCGCTTTGGGTGGCTC	462	0.2
gLAG3_3	ACCTGGAGCCACCCAAAGCGG	195	3.1
gLAG3_4	GCTCACCTAGTGAAGCCTCTC	463	1.3
gLAG3_5	TGCGAAGAGCAGGGGTCACTT	464	0.8
gLAG3_6	GGGTGCATACCTGTCTGGCTG	59	52.4
gLAG3_7	CCGCCCAGTGGCCCGCCCGCT	465	N.D.
gLAG3_8	TCGCTATGGCTGCGCCAGCC	466	0.1
gLAG3_9	TCCTTGACAGTGACTGCCAG	467	N.D.
gLAG3_10	CACAGTGACTGCCAGCCCCC	468	N.D.
gLAG3_11	GAAGTGTCTCTTCAGCCGCCC	469	0.1
gLAG3_12	AGCCGCCCTGACCGCCAGCC	470	0.1
gLAG3_13	CGCTAAGTGGTGATGGGGGA	197	2.3
gLAG3_14	CCGCTAAGTGGTGATGGGGGG	471	0.3
gLAG3_15	GCGGAAAGCTTCTCTTCCTG	472	1.0
gLAG3_16	GGGCAGGAAGAGGAAGCTTTC	191	6.4
gLAG3_17	CTCTTCCTGCCCAAGTCAGC	473	1.3
gLAG3_18	AACGTCTCCATCATGTATAAC	474	1.1
gLAG3_19	CTTTTCTCTTCAGGTCTGGAG	475	0.2
gLAG3_20	CTCTTCAGGTCTGGAGCCCCC	476	0.2
gLAG3_21	ACAGTGTACGCTGGAGCAGGT	477	0.1
gLAG3_22	GCAGTGAGGAAAGACCGGGTC	198	2.1
gLAG3_23	CTCACTGCCAAGTGGACTCCT	478	0.4
gLAG3_24	ACCCTTCGACTAGAGGATGTG	479	0.8

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gLAG3_25	CCCTTCGACTAGAGGATGTGA	196	2.7
gLAG3_26	GACTAGAGGATGTGAGCCAGG	480	1.0
gLAG3_27	CCACCTGAGGCTGACCTGTGA	193	3.4
gLAG3_28	CCCACCTGAGGCTGACCTGTG	481	0.8
gLAG3_29	TACTCTTTTCAGTGACTCCCA	482	0.3
gLAG3_30	CAGTGACTCCCAAATCCTTTG	483	0.1
gLAG3_31	CCCAGGGATCCAGGTGACCCA	194	3.1
gLAG3_32	GGGTCACCTGGATCCCTGGGG	484	0.2
gLAG3_33	GGTCACCTGGATCCCTGGGGA	88	17.1
gLAG3_34	GTGAGGTGACTCCAGTATCTG	485	0.7
gLAG3_35	TGAGGTGACTCCAGTATCTGG	188	9.3
gLAG3_36	GTGTGGAGCTCTCTGGACACC	486	0.9
gLAG3_37	TGTGGAGCTCTCTGGACACCC	190	6.9
gLAG3_38	TCAGGACCTTGGCTGGAGGCA	87	17.7
gLAG3_39	GCTGGAGGCACAGGAGGCCCA	487	0.3
gLAG3_40	CCCAGCCTTGGCAATGCCAGC	488	0.8
gLAG3_41	CCAGCCTTGGCAATGCCAGCT	189	8.3
gLAG3_42	GCAATGCCAGCTGTACCAGGG	489	0.6
gLAG3_43	TTGGAGCAGCAGTGTACTTCA	490	0.8
gLAG3_44	ACAGAGCTGTCTAGCCCAGGT	491	0.4
gLAG3_45	CTCCATAGGTGCCCAACGCTC	492	1.3
gLAG3_46	TCCATAGGTGCCCAACGCTCT	192	4.0
gLAG3_47	TCATCCTTGGTGTCTTTCTC	493	0.4
gLAG3_48	GTGTCCTTTCTCTGCTCCTTT	494	0.1
gLAG3_49	CTCTGCTCCTTTTGGTGACTG	495	0.2
gLAG3_50	TCTGCTCCTTTTGGTGACTGG	496	0.1
gLAG3_51	TGGTGACTGGAGCCTTTGGCT	497	0.6
gLAG3_52	GGTGACTGGAGCCTTTGGCTT	498	0.2
gLAG3_53	GGCTTTCACCTTTGGAGAAGA	499	0.1
gLAG3_54	GCTTTCACCTTTGGAGAAGAC	500	0.2
gLAG3_55	CTCTAAGGCAGAAAATCGTCT	501	0.1
gLAG3_56	CTGCCTTAGAGCAAGGGATTC	502	0.1
gLAG3_57	GAGCAAGGGATTACCCCTCCG	503	0.2

Table 13 Tested crRNAs Targeting Human PDCD1 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPD_1	AACCTGACCTGGGACAGTTTC	504	0.2
gPD_2	CCTTCCGCTCACCTCCGCCTG	89	46.9
gPD_3	CGCTCACCTCCGCCTGAGCAG	505	1.0
gPD_4	TCCACTGCTCAGGCGGAGGTG	506	0.6
gPD_5	TCCCCAGCCCTGCTCGTGGTG	507	1.2
gPD_6	GGTACCACGAGCAGGGCTGG	508	0.7

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPD_7	ACCTGCAGCTTCTCCAACACA	509	0.2
gPD_8	GCACGAAGCTCTCCGATGTGT	90	41.7
gPD_9	TCCAACACATCGGAGAGCTTC	510	0.2
gPD_10	GTGCTAAACTGGTACCGCATG	511	0.2
gPD_11	TCCGTCTGGTTGCTGGGGCTC	512	0.1
gPD_12	CCCGAGGACCGCAGCCAGCCC	513	0.4
gPD_13	CGTGTCACACAAC TGCCAAC	514	0.5
gPD_14	CACATGAGCGTGGTCAGGGCC	515	0.1
gPD_15	GATCTGCGCCTTGGGGGCCAG	516	0.1
gPD_16	ATCTGCGCCTTGGGGGCCAGG	517	1.2
gPD_17	GGGGCCAGGGAGATGGCCCCA	518	0.6
gPD_18	GTGCCCTTCCAGAGAGAAGGG	201	1.7
gPD_19	TGCCCTTCCAGAGAGAAGGGC	519	0.9
gPD_20	CAGAGAGAAGGGCAGAAGTGC	199	2.5
gPD_21	TGCCCTTCTCTCTGGAAGGGC	520	1.4
gPD_22	GAACTGGCCGGCTGGCCTGGG	200	1.7
gPD_23	TCTGCAGGGACAATAGGAGCC	60	57.6
gPD_24	CTCCTCAAAGAAGGAGGACCC	521	0.1
gPD_25	TCCTCAAAGAAGGAGGACCCC	522	0.5
gPD_26	TCTCGCCACTGGAATCCAGC	523	0.2
gPD_27	CAGTGGCGAGAGAAGACCCCG	92	23.7
gPD_28	CCTAGCGGAATGGGCACCTCA	524	0.1
gPD_29	CTAGCGGAATGGGCACCTCAT	91	30.3
gPD_30	GCCCCTCTGACCGGCTTCCTT	525	0.3

Table 14 Tested crRNAs Targeting Human PTPN6 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPTPN6_1	ACCGAGACCTCAGTGGGCTGG	96	58.2
gPTPN6_2	AGCAGGGTCTCTGCATCCAGC	526	0.3
gPTPN6_4	CTGGCTCGGCCCAGTCGCAAG	208	4.3
gPTPN6_5	TCCCCCTCCATACAGGTCATAG	803	14.8
gPTPN6_6	TATGACCTGTATGGAGGGGAG	61	83.4
gPTPN6_7	CGACTCTGACAGAGCTGGTGG	801	78.1
gPTPN6_8	AGGTGGATGATGGTGGCGTCG	209	3.5
gPTPN6_9	CCTGACGCTGCCTTCTCTAGG	527	0.8
gPTPN6_10	TCTAGGTGGTACCATGGCCAC	212	2.4
gPTPN6_11	GCCTGCAGCAGCGTCTCTGCC	528	0.2
gPTPN6_12	TTGTGCGTGAGAGCCTCAGCC	100	29.4
gPTPN6_13	GTGCTTTCTGTGCTCAGTGAC	529	0.8
gPTPN6_14	GGCTGGTCACTGAGCACAGAA	104	10.4
gPTPN6_15	CTGTGCTCAGTGACCAGCCCA	530	0.5
gPTPN6_16	TGTGCTCAGTGACCAGCCCCA	98	37.5

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPTPN6_17	ATGTGGGTGACCCTGAGCGGG	531	0.9
gPTPN6_18	CCTCGCACATGACCTTGATGT	532	1.4
gPTPN6_19	GCTCCCCCAGGGTGGACGCT	103	13.5
gPTPN6_20	GAGACCTTCGACAGCCTCACG	202	9.7
gPTPN6_21	GACAGCCTCACGGACCTGGTG	533	0.5
gPTPN6_22	AAGAAAGACGGGGATTGAGGAG	101	22.3
gPTPN6_23	TTGTTCAAGTTCCAACACTCGG	534	0.1
gPTPN6_24	GCTGTATCCTCGGACTCCTGC	535	0.4
gPTPN6_25	CCCACCCACATCTCAGAGTTT	99	34.8
gPTPN6_26	CAGAAGCAGGAGGTGAAGAAC	802	77.5
gPTPN6_27	CAGACGCTGGTGCAAGTTCTT	536	0.3
gPTPN6_28	CACCAGCGTCTGGAAGGGCAG	205	5.4
gPTPN6_29	TTCTCTGGCCGCTGCCCTTCC	537	0.1
gPTPN6_30	ATGTAGTTGGCATTGATGTAG	538	0.2
gPTPN6_31	CGTCCAGAACCAGCTGCTAGG	539	0.3
gPTPN6_32	TGGCAGATGGCGTGGCAGGAG	207	4.4
gPTPN6_33	TCCACCTCTCGGGTGGTCATG	540	0.7
gPTPN6_34	CTCCACCTCTCGGGTGGTCAT	541	1.2
gPTPN6_35	CCAGAACAAATGCGTCCCATA	542	0.2
gPTPN6_36	CAGAACAAATGCGTCCCATAC	543	0.5
gPTPN6_37	TGGGCCCTACTCTGTGACCAA	97	51.3
gPTPN6_38	TATTCGGTTGTGTCATGCTCC	544	0.1
gPTPN6_39	CAGGTCTCCCCGCTGGACAAT	213	1.6
gPTPN6_40	GGGAGACCTGATTGCGGAGAT	210	3.4
gPTPN6_41	CTGGACCAGATCAACCAGCGG	203	8.4
gPTPN6_42	CTGCCGCTGGTTGATCTGGTC	206	5.3
gPTPN6_43	CCTGCCGCTGGTTGATCTGGT	545	0.3
gPTPN6_44	CCCAGCGCCGGCATCGGCCGC	546	N.D.
gPTPN6_45	GTGGAGATGTTCTCCATGAGC	547	N.D.
gPTPN6_46	ACTGCCCCCACCAGGCCTG	93	80.3
gPTPN6_47	TACTGCGCCTCCGTCTGCACC	548	0.1
gPTPN6_48	AATGAACTGGGCGATGGCCAC	211	3.3
gPTPN6_49	TTCTTAGTGGTTTCAATGAAC	549	0.1
gPTPN6_50	GCATGGGCATTCTTCATGGCT	550	N.D.
gPTPN6_51	GACGAGGTGCGGGAGGCCTTG	551	N.D.
gPTPN6_52	GAGTCTAGTGCAGGGACCGTG	552	0.1
gPTPN6_53	CCCCCCTGCACCCGGCTGCAG	204	7.0
gPTPN6_54	TGTCTGCAGCCGGGTGCAGGG	553	0.9
gPTPN6_55	TCCTCCCTCTTGTCTTAGTG	554	0.0
gPTPN6_56	CTCCTCCCTCTTGTCTTAGT	555	0.1
gPTPN6_57	TTCACTTTCTCCTCCCTCTTG	556	0.2

Table 15 Tested crRNAs Targeting Human TIGIT Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTIGIT_1	CCTGAGGCGAGGGGAGCCTGC	557	0.2
gTIGIT_2	AGGCCTTACCTGAGGCGAGGG	62	81.7
gTIGIT_3	GTCTCTTCCCTAGGAATGAT	558	1.3
gTIGIT_4	TATTGTGCCTGTCATCATTCC	559	1.0
gTIGIT_5	TCTGCAGAAATGTTCCCCGTT	560	1.1
gTIGIT_6	CTCTGCAGAAATGTTCCCCGT	561	0.1
gTIGIT_7	TGCAGAGAAAGGTGGCTCTAT	215	6.0
gTIGIT_8	TGCCGTGGTGGAGGAGAGGTG	562	0.3
gTIGIT_9	TGGCCATTTGTAATGCTGACT	563	0.8
gTIGIT_10	TAATGCTGACTTGGGGTGGCA	216	1.6
gTIGIT_11	GGGTGGCACATCTCCCCATCC	214	9.7
gTIGIT_12	AAGGATGGGGAGATGTGCCAC	564	0.4
gTIGIT_13	AAGGATCGAGTGGCCCCAGGT	565	0.2
gTIGIT_14	TGCATCTATCACACCTACCT	566	1.4
gTIGIT_15	TAGGACCTCCAGGAAGATTCT	567	0.4
gTIGIT_16	CTAGGACCTCCAGGAAGATTC	568	0.5
gTIGIT_17	CTCCAGCAGGAATACCTGAGC	569	0.8
gTIGIT_18	GTCTCCCTCTAGTGGCTGAG	105	72.4
gTIGIT_19	GAGCCATGGCCGCGACGCTGG	570	0.9
gTIGIT_20	TAGTCAACGCGACCACCACGA	571	0.1
gTIGIT_21	CTAGTCAACGCGACCACCACG	572	0.1
gTIGIT_22	TAGTTTGTTTGTTTTTAGAAG	573	0.6
gTIGIT_23	TTTGTTTTTAGAAGAAAGCCC	574	1.0
gTIGIT_24	TTTTTAGAAGAAAGCCCTCAG	575	0.4
gTIGIT_25	TAGAAGAAAGCCCTCAGAATC	576	1.2
gTIGIT_26	CACAGAATGGATTCTGAGGGC	577	0.3
gTIGIT_27	CTCCTGAGGTCACCTTCCACA	217	1.6
gTIGIT_28	CTGGGGGTGAGGGAGCACTGG	578	0.5
gTIGIT_29	TGCCTGGACACAGCTTCTGG	579	0.3
gTIGIT_30	TGTAACCTCAGGACATTGAAGT	580	0.5
gTIGIT_31	AATGTCCTGAGTTACAGAAGC	581	0.5

Table 16 Tested crRNAs Targeting Human TRAC Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTRAC001	TGTTTTTAATGTGACTCTCAT	237	1.8
gTRAC002	GTGTTTTTAATGTGACTCTCA	582	0.4
gTRAC003	CGTAGGATTTTGTGTTTTTAA	583	0.1
gTRAC004	CTTAGTGCTGAGACTCATTCT	584	0.7

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTRAC005	CCTTAGTGCTGAGACTCATTC	585	0.6
gTRAC006	TGAGGGTGAAGGATAGACGCT	63	81.8
gTRAC007	ATAAACTGTAAAGTACCAAAC	239	1.7
gTRAC008	TTTGGTACTTTACAGTTTATT	586	0.2
gTRAC009	GTACTTTACAGTTTATTAAAT	238	1.7
gTRAC010	CAGTTTATTAAATAGATGTTT	587	0.5
gTRAC011	TTAAATAGATGTTTATATGGA	588	0.0
gTRAC012	TATGGAGAAGCTCTCATTTCT	110	46.7
gTRAC013	TTTCTCAGAAGAGCCTGGCTA	225	5.8
gTRAC014	TCAGAAGAGCCTGGCTAGGAA	127	16.6
gTRAC015	ACCTGCAAAATGAATATGGTG	589	0.0
gTRAC016	GCAGGTGAAATTCCTGAGATG	590	0.2
gTRAC017	CAGGTGAAATTCCTGAGATGT	804	63.6
gTRAC018	CTCGATATAAGGCCTTGAGCA	120	26.0
gTRAC019	AACTATAAATCAGAACACCTG	228	4.5
gTRAC020	GAACATAAATCAGAACACCT	224	6.4
gTRAC021	TAGTTCAAAACCTCTATCAAT	117	27.7
gTRAC022	TGGTATGTTGGCATTAAAGTTG	591	1.0
gTRAC023	CCAACTTAATGCCAACATACC	592	1.4
gTRAC024	CTTTGCTGGGCCTTTTTTCCCA	593	1.0
gTRAC025	CTGGGCCTTTTTCCCATGCCT	227	4.6
gTRAC026	TCCCATGCCTGCCTTTACTCT	594	0.6
gTRAC027	CCCATGCCTGCCTTTACTCTG	595	0.7
gTRAC028	CCATGCCTGCCTTTACTCTGC	129	15.3
gTRAC029	CTCTGCCAGAGTTATATTGCT	128	15.8
gTRAC030	ATAGGATCTTCTTCAAAACCC	235	2.2
gTRAC031	TTTAATAGGATCTTCTTCAAA	596	0.3
gTRAC032	ATTTAATAGGATCTTCTTCAA	597	0.1
gTRAC033	GAAGAAGATCCTATTAAATAA	236	2.0
gTRAC034	AAGAAGATCCTATTAAATAAAA	598	0.1
gTRAC035	AGGTTTCCTTGAGTGGCAGGC	220	7.5
gTRAC036	CTTGAGTGGCAGGCCAGGCCT	230	4.4
gTRAC037	AGTGAACGTTACGGCCAGGC	599	0.7
gTRAC038	TACGGGAAATAGCATCTTAGA	114	40.7
gTRAC039	TAAGATGCTATTTCCCGTATA	111	45.8
gTRAC040	CCGTATAAAGCATGAGACCGT	124	21.5
gTRAC041	CCCCAACCCAGGCTGGAGTCC	125	18.7
gTRAC042	CCTCTTTGCCCCAACCCAGGC	219	7.6
gTRAC043	GAGTCTCTCAGCTGGTACACG	121	25.9
gTRAC044	AGAATCAAAATCGGTGAATAG	221	7.4
gTRAC045	TTTGAGAATCAAAATCGGTGA	600	1.3
gTRAC046	TGACACATTTGTTTGAGAATC	601	0.2
gTRAC047	GATTCTCAAACAAATGTGTCA	602	0.1
gTRAC048	ATTCTCAAACAAATGTGTCAC	229	4.5

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTRAC049	TCTGTGATATACACATCAGAA	118	27.6
gTRAC050	GTCTGTGATATACACATCAGA	130	11.4
gTRAC055	CACATGCAAAGTCAGATTTGT	603	1.0
gTRAC056	CATGTGCAAACGCCTTCAACA	231	3.9
gTRAC057	GTGCCTTCGCAGGCTGTTTCC	604	0.9
gTRAC058	CTTGCTTCAGGAATGGCCAGG	116	27.8
gTRAC059	GACATCATTGACCAGAGCTCT	805	50.1
gTRAC060	AGACATCATTGACCAGAGCTC	605	1.3
gTRAC061	GTGGCAATGGATAAGGCCGAG	115	38.8
gTRAC062	GGTGGCAATGGATAAGGCCGA	223	6.5
gTRAC063	TTAGTAAAAAGAGGGTTTTGG	606	1.4
gTRAC064	TACTAAGAAACAGTGAGCCTT	232	3.5
gTRAC065	ACTAAGAAACAGTGAGCCTTG	607	0.2
gTRAC066	CTAAGAAACAGTGAGCCTTGT	218	9.5
gTRAC067	CCGTGTCATTCTCTGGACTGC	112	45.4
gTRAC068	CCCGTGTCATTCTCTGGACTG	226	5.3
gTRAC069	TCCCGTGTCATTCTCTGGACT	608	1.0
gTRAC070	TTCCCGTGTCATTCTCTGGAC	609	0.3
gTRAC071	CTCAGACTGTTTGCCCCCTTAC	233	3.4
gTRAC072	CCCCTTACTGCTCTTCTAGGC	222	6.9
gTRAC073	GCAGACAGGGAGAAATAAGGA	106	66.9
gTRAC074	GGCAGACAGGGAGAAATAAGG	119	27.1
gTRAC075	TGGCAGACAGGGAGAAATAAG	122	25.2
gTRAC076	TTGGCAGACAGGGAGAAATAA	126	16.7
gTRAC077	TCCCTGTCTGCCAAAAAATCT	610	1.1
gTRAC078	CCAGCTCACTAAGTCAGTCTC	109	47.4
gTRAC079	ATTCCTCCACTTCAACACCTG	113	45.4
gTRAC080	AATTCCTCCACTTCAACACCT	611	0.5
gTRAC081	TAATTCCTCCACTTCAACACC	234	2.3
gTRAC082	CCAGCTGACAGATGGGCTCCC	123	21.5
gTRAC083	CCCAGCTGACAGATGGGCTCC	241	1.6
gTRAC084	GACTTTTCCCAGCTGACAGAT	240	1.6
gTRAC085	TCAACCCTGAGTTAAAACACA	612	0.5
gTRAC086	CTCAACCCTGAGTTAAAACAC	613	0.2
gTRAC087	TCCTGAAGGTAGCTGTTTTCT	614	0.2
gTRAC088	GTCTGAAGGTAGCTGTTTTTC	615	0.1
gTRAC089	AACTCAGGGTTGAGAAAACAG	616	0.7
gTRAC090	ACTCAGGGTTGAGAAAACAGC	617	0.1

Table 17 Tested crRNAs Targeting Human TRBC1/TRBC2 Genes

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gTRBC1+2_1	AGCCATCAGAAGCAGAGATCT	705	66.40 (TRBC1) ; 74.7 (TRBC2)
gTRBC1+2_3	CGCTGTCAAGTCCAGTTCTAC	706	71.28 (TRBC1)
gTRBC2_7	CCCTGTTTTCTTTCAGACTGT	707	0.09
gTRBC2_8	CTTTCAGACTGTGGCTTCACC	708	0.24
gTRBC2_9	TTTCAGACTGTGGCTTCACCT	709	0.24
gTRBC2_10	CAGACTGTGGCTTCACCTCCG	710	0.16
gTRBC2_11	AGACTGTGGCTTCACCTCCGG	711	19.97
gTRBC2_12	CCGGAGGTGAAGCCACAGTCT	712	33.14
gTRBC2_13	TCAACAGAGTCTTACCAGCAA	713	1.20
gTRBC2_14	CCAGCAAGGGGTCCTGTCTGC	714	6.69
gTRBC2_15	CTAGGGAAGGCCACCTTGAT	715	21.74
gTRBC2_16	TATGCCGTGCTGGTCAGTGCC	716	0.20
gTRBC2_17	CCATGGCCATCAGCACGAGGG	717	1.75
gTRBC2_18	CCTAGCAAGATCTCATAGAGG	718	0.37
gTRBC2_19	CACAGGTCAAGAGAAAGGATT	719	1.58
gTRBC2_21	GAGCTAGCCTCTGGAATCCTT	720	11.89

Table 18 Tested crRNAs Targeting Human CARD11 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCARD11_1	TAGTACCGCTCCTGGAAGGTT	721	1.37
gCARD11_2	ATCTTGTAGTACCGCTCCTGG	722	0.07
gCARD11_3	CTTCATCTTGTAGTACCGCTC	723	0.08

5

Table 19 Tested crRNAs Targeting Human CD247 gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCD247_1	TGTGTTGCAGTTCAGCAGGAG	724	55.77
gCD247_2	CGTTATAGAGCTGGTTCTGGC	725	0.20
gCD247_3	CGGAGGGTCTACGGCGAGGCT	726	20.79
gCD247_4	TTATCTGTTATAGGAGCTCAA	727	12.31
gCD247_5	TCTGTTATAGGAGCTCAATCT	728	0.24
gCD247_6	TCCAAAACATCGTACTCCTCT	729	0.34
gCD247_7	CCCCCATCTCAGGGTCCCGGC	730	6.43

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gCD247_8	GACAAGAGACGTGGCCGGGAC	731	40.95
gCD247_9	TCTCCCTCTAACGTCTTCCCG	732	4.13
gCD247_10	CTGAGGGTTCTTCCTTCTCTG	733	0.05
gCD247_11	CCGTTGTCTTTCCTAGCAGAG	734	1.18
gCD247_12	CTAGCAGAGAAGGAAGAACCC	735	70.64
gCD247_13	TGCAGTTCCTGCAGAAGAGGG	736	4.93
gCD247_14	TGCAGGAAGTGCAGAAAGATA	737	2.91
gCD247_15	ATCCCAATCTCACTGTAGGCC	738	31.12
gCD247_16	CATCCCAATCTCACTGTAGGC	739	0.10
gCD247_17	CTCATTTCACTCCCAAACAAC	740	0.30
gCD247_18	TCATTTCACTCCCAAACAACC	741	44.34
gCD247_19	ACTCCCAAACAACCAGCGCCG	742	43.17
gCD247_20	TTTTCTGATTTGCTTTCACGC	743	0.10
gCD247_21	TGATTTGCTTTCACGCCAGGG	744	5.23
gCD247_22	CTTTCACGCCAGGGTCTCAGT	745	8.24
gCD247_23	ACGCCAGGGTCTCAGTACAGC	746	0.30

Table 20 Tested crRNAs Targeting Human IL7R Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gIL7R_1	CTTTCCAGGGGAGATGGATCC	747	0.25
gIL7R_2	CCAGGGGAGATGGATCCTATC	748	8.35
gIL7R_3	CAGGGGAGATGGATCCTATCT	749	87.87
gIL7R_4	CTAACCATCAGCATTTTGAGT	750	0.11
gIL7R_5	GAGTTTTTTCTCTGTCGCTCT	751	0.07
gIL7R_6	AGTTTTTTCTCTGTCGCTCTG	752	0.06
gIL7R_7	TCTGTCGCTCTGTTGGTCATC	753	2.61
gIL7R_8	CATAACACACAGGCCAAGATG	754	25.83

Table 21 Tested crRNAs Targeting Human LCK Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gLCK1_1	ATGTCCTTTCACCCATCAACC	755	0.06
gLCK1_2	CACCCATCAACCCGTAGGGAT	756	0.17
gLCK1_3	ACCCATCAACCCGTAGGGATG	757	16.21

5

Table 22 Tested crRNAs Targeting Human PLCG1 Gene

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPLCG1_1	CTCATACACCACGAAGCGCAG	758	0.09

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gPLCG1_2	CCTTTCTGCGCTTCGTGGTGT	759	5.14
gPLCG1_3	CTGCGCTTCGTGGTGTATGAG	760	0.05
gPLCG1_4	TGCGCTTCGTGGTGTATGAGG	761	1.91
gPLCG1_5	GTGGTGTATGAGGAAGACATG	762	3.53

Table 23 Tested crRNAs Targeting Certain Other Human Genes

crRNA	Spacer Sequence	SEQ ID NO	% Indel
gDHODH_1	TTGCAGAAGCGGGCCAGGAT	770	0.60
gDHODH_2	TTGCAGAAGCGGGCCAGGAT	771	0.59
gDHODH_3	TATGCTGAACACCTGATGCCG	772	74.94
gPLK1_1	CCAGGGTCGGCCGGTGCCCGT	773	29.06
gPLK1_2	GCCGGTGGAGCCGCCGCCGA	774	2.01
gPLK1_3	TGGGCAAGGGCGGCTTTGCCA	775	2.26
gPLK1_4	GGGCAAGGGCGGCTTTGCCAA	776	28.24
gPLK1_5	GGCAAGGGCGGCTTTGCCAAG	777	28.41
gPLK1_6	CCAAGTGCTTCGAGATCTCGG	778	2.07
gPLK1_7	CATGGACATCTTCTCCCTCTG	779	90.07
gPLK1_8	TCGAGGACAACGACTTCGTGT	780	0.16
gPLK1_9	CGAGGACAACGACTTCGTGTT	781	6.84
gPLK1_10	GAGGACAACGACTTCGTGTTC	782	8.52
gMVD_1	CAGTTAAAAACCACCACAACA	783	1.42
gMVD_2	GCTGAATGGCCGGGAGGAGGA	784	14.06
gMVD_3	TGGAGTGGCAGATGGGAGAGC	785	63.22
gTUBB_1	AACCATGAGGGAAATCGTGCA	786	2.61
gTUBB_2	ACCATGAGGGAAATCGTGAC	787	68.40
gTUBB_3	TTCTCTGTAGGTGGCAAATAT	788	18.67
gU6_1	GTCCTTTCCACAAGATATATA	763	68.1
gU6_2	GATTTCTTGGCTTTATATATC	764	0.71
gU6_3	TTGGCTTTATATATCTTGTGG	765	2.83
gU6_4	GCTTTATATATCTTGTGGAAA	766	0.37
gU6_5	ATATATCTTGTGGAAAGGACG	767	0.39
gU6_6	TATATCTTGTGGAAAGGACGA	768	0.39
gU6_7	TGGAAAGGACGAAACACCGTG	769	0.24

- 5 [0163] To provide sufficient targeting to the target nucleotide sequence, the spacer sequence can be 16 or more nucleotides in length. In certain embodiments, the spacer sequence is at least 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or 75 nucleotides in length. In certain embodiments, the spacer sequence is shorter than or equal to 75,

50, 45, 40, 35, 30, 25, or 20 nucleotides in length. Shorter spacer sequence may be desirable for reducing off-target events. Accordingly, in certain embodiments, the spacer sequence is shorter than or equal to 21, 20, 19, 18, or 17 nucleotides. In certain embodiments, the spacer sequence is 17-30 nucleotides in length, *e.g.*, 17-21, 17-22, 17-23, 17-24, 17-25, 17-30, 20-21, 20-22, 20-23, 20-24, 20-25, or 20-30 nucleotides in length. In certain embodiments, the spacer sequence is 19-22 nucleotides in length, for example 20 to 21 nucleotides in length. In certain embodiments, the spacer sequence is 21 nucleotides in length. In certain embodiments, the spacer sequence is 20 nucleotides in length.

[0164] In certain embodiments, the spacer sequence comprises a portion of a spacer sequence listed in **Table 1, 2, or 3**, wherein the portion is 16, 17, 18, 19, or 20 nucleotides in length. In certain embodiments, the spacer sequence comprises nucleotides 1-16, 1-17, 1-18, 1-19, or 1-20 of a spacer sequence listed in **Table 1, 2, or 3**. In specific embodiments, the spacer sequence consists of nucleotides 1-16, 1-17, 1-18, 1-19, or 1-20 of a spacer sequence listed in **Table 1, 2, or 3**.

[0165] In certain embodiments, the spacer sequence comprises a portion of a spacer sequence listed in **Table 1-2 or 3**, wherein the portion is 16, 17, 18, 19, or 20 nucleotides in length. In certain embodiments, the spacer sequence comprises nucleotides 1-16, 1-17, 1-18, 1-19, or 1-20 of a spacer sequence listed in **Table 1, 2, or 3**. In specific embodiments, the spacer sequence consists of nucleotides 1-16, 1-17, 1-18, 1-19, or 1-20 of a spacer sequence listed in any one of **Tables 1-23**.

[0166] In certain embodiments, the spacer sequence is 21 nucleotides in length. In certain embodiments, the spacer sequence consists of a spacer sequence shown in **Table 1, 2, or 3**.

[0167] In certain embodiments, the spacer sequence is 21 nucleotides in length. In certain embodiments, the spacer sequence consists of a spacer sequence shown in any one of **Tables 1-23**.

[0168] In certain embodiments, the spacer sequence, where it is longer than 21 nucleotides in length, comprises a spacer sequence shown in **Table 1, 2, or 3** and one or more nucleotides. In certain embodiments, the one or more nucleotides are 3' to the spacer sequence shown in **Table 1, 2, or 3**.

[0169] In certain embodiments, the spacer sequence, where it is longer than 21 nucleotides in length, comprises a spacer sequence shown in **Table 1, 2, or 3** and one or more nucleotides. In certain embodiments, the one or more nucleotides are 3' to the spacer sequence shown in any one of **Tables 1-3**.

[0170] In certain embodiments, the spacer sequence is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 93%, at least 94%, at least 95%, at least 96%, at least

97%, at least 98%, or at least 99% complementary to the target nucleotide sequence. In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence in the seed region (5-10 base pairs proximal to the PAM). In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence. The spacer sequences listed in **Tables 1-3** are designed to be 100% complementary to the wild-type sequence of the corresponding target gene. Accordingly, it is contemplated that a spacer sequence useful for targeting a gene listed in **Table 1, 2, or 3** can be at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to a corresponding spacer sequence listed in **Table 1, 2, or 3**, or a portion thereof disclosed herein. In certain embodiments, the spacer sequence is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides different from a sequence listed in **Table 1, 2, or 3**. In certain embodiments, the spacer sequence is 100% identical to a sequence listed in **Table 1, 2, or 3** in the seed region (at least 5 base pairs proximal to the PAM). It has been reported that compared to DNA binding, DNA cleavage is less tolerant to mismatches between the spacer sequence and the target nucleotide sequence (see, Klein *et al.* (2018) Cell Reports, 22: 1413). Accordingly, in certain embodiments, a guide nucleic acid to be used with a Cas nuclease comprises a spacer sequence 100% complementary to the target nucleotide sequence. In certain embodiments, a guide nucleic acid to be used with a Cas nuclease comprises a spacer sequence listed in **Table 1, 2, or 3**, or a portion thereof disclosed herein.

[0171] In certain embodiments, the spacer sequence is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% complementary to the target nucleotide sequence. In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence in the seed region (at least 5 base pairs proximal to the PAM). In certain embodiments, the spacer sequence is 100% complementary to the target nucleotide sequence. The spacer sequences listed in any one of **Tables 1-23** are designed to be 100% complementary to the wild-type sequence of the corresponding target gene. Accordingly, it is contemplated that a spacer sequence useful for targeting a gene listed in **Table 1, 2, or 3** can be at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to a corresponding spacer sequence listed in any one of **Tables 1-23**, or a portion thereof disclosed herein. In certain embodiments, the spacer sequence is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides different from a sequence listed in any one of **Tables 1-23**. In certain embodiments, the spacer sequence is 100% identical to a sequence listed in any one of **Tables 1-23** in the seed region (at least 5 base pairs proximal to the PAM). It has been reported that compared to DNA binding, DNA cleavage is less tolerant to mismatches between the spacer

sequence and the target nucleotide sequence (see, Klein *et al.* (2018) Cell Reports, 22: 1413). Accordingly, in certain embodiments, a guide nucleic acid to be used with a Cas nuclease comprises a spacer sequence 100% complementary to the target nucleotide sequence. In certain embodiments, a guide nucleic acid to be used with a Cas nuclease comprises a spacer sequence
 5 listed in any one of **Table 1-23**, or a portion thereof disclosed herein.

[0172] The present invention also provides guide nucleic acids targeting human DHODH, PLK1, MVD, TUBB, or U6 gene comprising the spacer sequences provided below in **Table 25**. DHODH, PLK1, MVD, and TUBB are known to be essential genes. It is contemplated that the guide nucleic acids targeting these genes, particularly the ones that edit the respective
 10 genomic locus at high efficiency (*e.g.*, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%), can be used as positive controls for assessing transfection efficiency and other experimental processes. The spacer sequences targeting U6 in **Table 25** are designed to hybridize with the promoter region of human U6 gene and can be used to assess expression of an inserted gene from the endogenous U6 promoter.

[0173] In certain embodiments, the 3' end of the targeter stem sequence is linked by no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides to the 5' end of the spacer sequence. In certain embodiments, the targeter stem sequence and the spacer sequence are adjacent to each other, directly linked by an internucleotide bond. In certain embodiments, the targeter stem sequence and the spacer sequence are linked by one nucleotide, *e.g.*, a uridine. In certain
 20 embodiments, the targeter stem sequence and the spacer sequence are linked by two or more nucleotides. In certain embodiments, the targeter stem sequence and the spacer sequence are linked by 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides.

[0174] In certain embodiments, the targeter nucleic acid further comprises an additional nucleotide sequence 5' to the targeter stem sequence. In certain embodiments, the additional
 25 nucleotide sequence comprises at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50) nucleotides. In certain embodiments, the additional nucleotide sequence consists of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides. In certain embodiments, the additional nucleotide sequence consists of 2
 30 nucleotides. In certain embodiments, the additional nucleotide sequence is reminiscent to the loop or a fragment thereof (*e.g.*, one, two, three, or four nucleotides at or near the 3' end of the loop) in a crRNA of a corresponding single guide CRISPR-Cas system. It is understood that an additional nucleotide sequence 5' to the targeter stem sequence is dispensable. Accordingly, in certain embodiments, the targeter nucleic acid does not comprise any additional nucleotide 5' to
 35 the targeter stem sequence.

[0175] In certain embodiments, the targeter nucleic acid further comprises an additional nucleotide sequence containing one or more nucleotides at or near the 3' end that does not hybridize with the target nucleotide sequence. The additional nucleotide sequence may protect the targeter nucleic acid from degradation by 3'-5' exonuclease. In certain embodiments, the additional nucleotide sequence is no more than 100 nucleotides in length. In certain
 5 embodiments, the additional nucleotide sequence is no more than 90, 80, 70, 60, 50, 40, 30, 20, or 10 nucleotides in length. In certain embodiments, the additional nucleotide sequence is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides in length. In certain
 10 embodiments, the additional nucleotide sequence is 5-100, 5-50, 5-40, 5-30, 5-25, 5-20, 5-15, 5-10, 10-100, 10-50, 10-40, 10-30, 10-25, 10-20, 10-15, 15-100, 15-50, 15-40, 15-30, 15-25, 15-20, 20-100, 20-50, 20-40, 20-30, 20-25, 25-100, 25-50, 25-40, 25-30, 30-100, 30-50, 30-40, 40-100, 40-50, or 50-100 nucleotides in length.

[0176] In certain embodiments, the additional nucleotide sequence forms a hairpin with the spacer sequence. Such secondary structure may increase the specificity of the engineered,
 15 non-naturally occurring system (see, Kocak *et al.* (2019) NAT. BIOTECH. 37: 657-66). In certain embodiments, the free energy change during the hairpin formation is greater than or equal to -20 kcal/mol, -15 kcal/mol, -14 kcal/mol, -13 kcal/mol, -12 kcal/mol, -11 kcal/mol, or -10 kcal/mol. In certain embodiments, the free energy change during the hairpin formation is greater than or equal to -5 kcal/mol, -6 kcal/mol, -7 kcal/mol, -8 kcal/mol, -9 kcal/mol, -10 kcal/mol, -11
 20 kcal/mol, -12 kcal/mol, -13 kcal/mol, -14 kcal/mol, or -15 kcal/mol. In certain embodiments, the free energy change during the hairpin formation is in the range of -20 to -10 kcal/mol, -20 to -11 kcal/mol, -20 to -12 kcal/mol, -20 to -13 kcal/mol, -20 to -14 kcal/mol, -20 to -15 kcal/mol, -15 to -10 kcal/mol, -15 to -11 kcal/mol, -15 to -12 kcal/mol, -15 to -13 kcal/mol, -15 to -14 kcal/mol, -14 to -10 kcal/mol, -14 to -11 kcal/mol, -14 to -12 kcal/mol, -14 to -13 kcal/mol, -13
 25 to -10 kcal/mol, -13 to -11 kcal/mol, -13 to -12 kcal/mol, -12 to -10 kcal/mol, -12 to -11 kcal/mol, or -11 to -10 kcal/mol. In other embodiments, the targeter nucleic acid does not comprise any nucleotide 3' to the spacer sequence.

[0177] In certain embodiments, the modulator nucleic acid further comprises an additional nucleotide sequence 3' to the modulator stem sequence. In certain embodiments, the
 30 additional nucleotide sequence comprises at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50) nucleotides. In certain embodiments, the additional nucleotide sequence consists of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides. In certain embodiments, the additional nucleotide sequence consists of 1
 35 nucleotide (*e.g.*, uridine). In certain embodiments, the additional nucleotide sequence consists of

2 nucleotides. In certain embodiments, the additional nucleotide sequence is reminiscent to the loop or a fragment thereof (*e.g.*, one, two, three, or four nucleotides at or near the 5' end of the loop) in a crRNA of a corresponding single guide CRISPR-Cas system. It is understood that an additional nucleotide sequence 3' to the modulator stem sequence is dispensable. Accordingly, in certain embodiments, the modulator nucleic acid does not comprise any additional nucleotide 3' to the modulator stem sequence.

[0178] It is understood that the additional nucleotide sequence 5' to the targeter stem sequence and the additional nucleotide sequence 3' to the modulator stem sequence, if present, may interact with each other. For example, although the nucleotide immediately 5' to the targeter stem sequence and the nucleotide immediately 3' to the modulator stem sequence do not form a Watson-Crick base pair (otherwise they would constitute part of the targeter stem sequence and part of the modulator stem sequence, respectively), other nucleotides in the additional nucleotide sequence 5' to the targeter stem sequence and the additional nucleotide sequence 3' to the modulator stem sequence may form one, two, three, or more base pairs (*e.g.*, Watson-Crick base pairs). Such interaction may affect the stability of the complex comprising the targeter nucleic acid and the modulator nucleic acid.

[0179] The stability of a complex comprising a targeter nucleic acid and a modulator nucleic acid can be assessed by the Gibbs free energy change (ΔG) during the formation of the complex, either calculated or actually measured. Where all the predicted base pairing in the complex occurs between a base in the targeter nucleic acid and a base in the modulator nucleic acid, *i.e.*, there is no intra-strand secondary structure, the ΔG during the formation of the complex correlates generally with the ΔG during the formation of a secondary structure within the corresponding single guide nucleic acid. Methods of calculating or measuring the ΔG are known in the art. An exemplary method is RNAfold (rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) as disclosed in Gruber *et al.* (2008) NUCLEIC ACIDS RES., 36(Web Server issue): W70–W74. Unless indicated otherwise, the ΔG values in the present disclosure are calculated by RNAfold for the formation of a secondary structure within a corresponding single guide nucleic acid. In certain embodiments, the ΔG is lower than or equal to -1 kcal/mol, *e.g.*, lower than or equal to -2 kcal/mol, lower than or equal to -3 kcal/mol, lower than or equal to -4 kcal/mol, lower than or equal to -5 kcal/mol, lower than or equal to -6 kcal/mol, lower than or equal to -7 kcal/mol, lower than or equal to -7.5 kcal/mol, or lower than or equal to -8 kcal/mol. In certain embodiments, the ΔG is greater than or equal to -10 kcal/mol, *e.g.*, greater than or equal to -9 kcal/mol, greater than or equal to -8.5 kcal/mol, or greater than or equal to -8 kcal/mol. In certain embodiments, the ΔG is in the range of -10 to -4 kcal/mol. In certain embodiments, the ΔG is in the range of -8 to -4 kcal/mol, -7 to -4 kcal/mol, -6 to -4

kcal/mol, -5 to -4 kcal/mol, -8 to -4.5 kcal/mol, -7 to -4.5 kcal/mol, -6 to -4.5 kcal/mol, or -5 to -4.5 kcal/mol, for example -8 kcal/mol, -7 kcal/mol, -6 kcal/mol, -5 kcal/mol, -4.9 kcal/mol, -4.8 kcal/mol, -4.7 kcal/mol, -4.6 kcal/mol, -4.5 kcal/mol, -4.4 kcal/mol, -4.3 kcal/mol, -4.2 kcal/mol, -4.1 kcal/mol, or -4 kcal/mol.

5 **[0180]** It is understood that the ΔG may be affected by a sequence in the targeter nucleic acid that is not within the targeter stem sequence, and/or a sequence in the modulator nucleic acid that is not within the modulator stem sequence. For example, one or more base pairs (*e.g.*, Watson-Crick base pair) between an additional sequence 5' to the targeter stem sequence and an additional sequence 3' to the modulator stem sequence may reduce the ΔG , *i.e.*, stabilize the
10 nucleic acid complex. In certain embodiments, the nucleotide immediately 5' to the targeter stem sequence comprises a uracil or is a uridine, and the nucleotide immediately 3' to the modulator stem sequence comprises a uracil or is a uridine, thereby forming a nonconventional U-U base pair.

[0181] In certain embodiments, the modulator nucleic acid comprises a nucleotide
15 sequence referred to herein as a "5' sequence", *e.g.*, a tail sequence, positioned 5' to the modulator stem sequence. Where the CRISPR system is a type V-A CRISPR system, the 5' sequence, *e.g.*, a tail sequence, in a modified dual guide system is reminiscent of the nucleotide sequence positioned 5' to the stem-loop structure of the scaffold sequence in a crRNA (the single guide). Accordingly, the 5' sequence, *e.g.*, a tail sequence, can comprise the corresponding
20 nucleotide sequences when a modified dual guide system is engineered from a single guide system. In certain embodiments, one or more nucleotides or internucleotide linkages at or near the 5' end of the 5' sequence, *e.g.*, tail sequence, can be modified, *e.g.*, chemically modified.

[0182] Without being bound by theory, it is contemplated that or near the 5' sequence, *e.g.*, tail sequence, may participate in the formation of the CRISPR-Cas complex. For example,
25 in certain embodiments, the 5' sequence, *e.g.*, tail sequence, forms a pseudoknot structure with the modulator stem sequence, which is recognized by the Cas protein (see, Yamano *et al.* (2016) CELL, 165: 949). In certain embodiments, the 5' sequence, *e.g.*, tail sequence, is at least 3 (*e.g.*, at least 4 or at least 5) nucleotides in length. In certain embodiments, the 5' sequence, *e.g.*, tail sequence, is 3, 4, or 5 nucleotides in length. In certain embodiments, the nucleotide at or near the
30 3' end of the 5' sequence, *e.g.*, tail sequence, comprises a uracil or is a uridine. In certain embodiments, the second nucleotide in the 5' sequence, *e.g.*, tail sequence, the position counted from the 3' end, comprises a uracil or is a uridine. In certain embodiments, the third nucleotide in the 5' sequence, *e.g.*, tail sequence, the position counted from the 3' end, comprises an adenine or is an adenosine. This third nucleotide may form a base pair (*e.g.*, a Watson-Crick base
35 pair) with a nucleotide 5' to the modulator stem sequence. Accordingly, in certain embodiments,

the modulator nucleic acid comprises a uridine or a uracil-containing nucleotide 5' to the modulator stem sequence. In certain embodiments, the 5' sequence, *e.g.*, tail sequence, comprises the nucleotide sequence of 5'-AUU-3'. In certain embodiments, the 5' sequence, *e.g.*, tail sequence, comprises the nucleotide sequence of 5'-AAUU-3'. In certain embodiments, the 5' sequence, *e.g.*, tail sequence, comprises the nucleotide sequence of 5'-UAAUU-3'. In certain embodiments, the 5' sequence, *e.g.*, tail sequence, is positioned immediately 5' to the modulator stem sequence.

[0183] In certain embodiments, the targeter nucleic acid and/or the modulator nucleic acid are designed to reduce the degree of secondary structure other than the hybridization between the targeter stem sequence and the modulator stem sequence. In certain embodiments, at most 75%, 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 1%, or fewer of the nucleotides of the targeter nucleic acid and/or the modulator nucleic acid participate in self-complementary base pairing when optimally folded. Optimal folding may be determined by any suitable polynucleotide folding algorithm. Some programs are based on calculating the minimal Gibbs free energy. An example of one such algorithm is mFold, as described by Zuker and Stiegler (Nucleic Acids Res. 9 (1981), 133-148). Another example folding algorithm is the online webserver RNAfold, developed at Institute for Theoretical Chemistry at the University of Vienna, using the centroid structure prediction algorithm (see *e.g.*, A. R. Gruber et al., 2008, Cell 106(1): 23-24; and PA Carr and GM Church, 2009, Nature Biotechnology 27(12): 1151-62).

[0184] The targeter nucleic acid is directed to a specific target nucleotide sequence, and the donor template is designed to modify the target nucleotide sequence or a sequence nearby. It is understood, therefore, that association of the targeter or modulator nucleic acid with a donor template can increase editing efficiency and reduce off-targeting. In a multiplex method (*e.g.*, as disclosed in the "Multiplex Methods" subsection of section II *infra*), association of a donor template with a modulator nucleic acid allows combination of a targeter nucleic acid library with a donor template library, making designs of screening or selection assays more efficient and flexible. Accordingly, in certain embodiments, the modulator nucleic acid further comprises a donor template-recruiting sequence capable of hybridizing with a donor template (see **Figure 1C**). Donor templates are described in the "Donor Templates" subsection of section II *infra*. The donor template and donor template-recruiting sequence can be designed such that they bear sequence complementarity. In certain embodiments, the donor template-recruiting sequence is at least 90% (*e.g.*, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) complementary to at least a portion of the donor template. In certain embodiments, the donor template-recruiting sequence is 100% complementary to at least a portion of the donor template. In certain embodiments, where the

donor template comprises an engineered sequence not homologous to the sequence to be repaired, the donor template-recruiting sequence is capable of hybridizing with the engineered sequence in the donor template. In certain embodiments, the donor template-recruiting sequence is at least 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 nucleotides in length. In certain embodiments, the donor template-recruiting sequence is positioned at or near the 5' end of the modulator nucleic acid. In certain embodiments, the donor template-recruiting sequence is linked to the 5' sequence, *e.g.*, tail sequence, if present, or to the modulator stem sequence, of the modulator nucleic acid through an internucleotide bond or a nucleotide linker.

[0185] In certain embodiments, a guide nucleic acid as described herein is associated with a donor template comprising a single strand oligodeoxynucleotide (ssODN).

[0186] In certain embodiments, the modulator nucleic acid further comprises an editing enhancer sequence, which increases the efficiency of gene editing and/or homology-directed repair (HDR). Exemplary editing enhancer sequences are described in Park *et al.* (2018) NAT. COMMUN. 9: 3313. In certain embodiments, the editing enhancer sequence is positioned 5' to the 5' sequence, *e.g.*, tail sequence, if present, or 5' to the modulator stem sequence. In certain embodiments, the editing enhancer sequence is 1-50, 4-50, 9-50, 15-50, 25-50, 1-25, 4-25, 9-25, 15-25, 1-15, 4-15, 9-15, 1-9, 4-9, or 1-4 nucleotides in length. In certain embodiments, the editing enhancer sequence is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 55 nucleotides in length. The editing enhancer sequence is designed to minimize homology to the target nucleotide sequence or any other sequence that the engineered, non-naturally occurring system may be contacted to, *e.g.*, the genome sequence of a cell into which the engineered, non-naturally occurring system is delivered. In certain embodiments, the editing enhancer is designed to minimize the presence of hairpin structure. The editing enhancer can comprise one or more of the chemical modifications disclosed herein.

[0187] The modulator and/or targeter nucleic acids can further comprise a protective nucleotide sequence that prevents or reduces nucleic acid degradation. In certain embodiments, the protective nucleotide sequence is at least 5 (*e.g.*, at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50) nucleotides in length. The length of the protective nucleotide sequence increases the time for an exonuclease to reach the 5' sequence, *e.g.*, tail sequence, modulator stem sequence, targeter stem sequence, and/or spacer sequence, thereby protecting these portions of the modulator and/or targeter nucleic acids from degradation by an exonuclease. In certain embodiments, the protective nucleotide sequence forms a secondary structure, such as a hairpin or a tRNA structure, to reduce the speed of degradation by an exonuclease (see, for example, Wu *et al.* (2018) CELL. MOL. LIFE SCI., 75(19): 3593-3607). Secondary structures can be predicted by methods known in the art, such as the

online webserver RNAfold developed at University of Vienna using the centroid structure prediction algorithm (see, Gruber *et al.* (2008) NUCLEIC ACIDS RES., 36: W70). Certain chemical modifications, which may be present in the protective nucleotide sequence, can also prevent or reduce nucleic acid degradation, as disclosed in the “RNA Modifications” subsection.

5 **[0188]** A protective nucleotide sequence is typically located at or near the 5' end, at or near the 3' end, or at both ends, of the modulator or targeter nucleic acid. In certain embodiments, the modulator nucleic acid comprises a protective nucleotide sequence at or near the 5' end, optionally through a nucleotide linker. In certain embodiments, the modulator nucleic acid comprises a protective nucleotide sequence at or near the 3' end. In certain embodiments, the modulator nucleic acid comprises a protective nucleotide sequence at or near the 5' end. In certain embodiments, the modulator nucleic acid comprises a protective nucleotide sequence at or near the 3' end.

15 **[0189]** As described above, various nucleotide sequences can be present in the 5' portion of a modulator nucleic acid, including but not limited to a donor template-recruiting sequence, an editing enhancer sequence, a protective nucleotide sequence, and a linker connecting such sequence to the 5' sequence, *e.g.*, tail sequence, if present, or to the modulator stem sequence. It is understood that the functions of donor template recruitment, editing enhancement, protection against degradation, and linkage are not exclusive to each other, and one nucleotide sequence can have one or more of such functions. For example, in certain embodiments, the modulator nucleic acid comprises a nucleotide sequence that is both a donor template-recruiting sequence and an editing enhancer sequence. In certain embodiments, the modulator nucleic acid comprises a nucleotide sequence that is both a donor template-recruiting sequence and a protective sequence. In certain embodiments, the modulator nucleic acid comprises a nucleotide sequence that is both an editing enhancer sequence and a protective sequence. In certain embodiments, the modulator nucleic acid comprises a nucleotide sequence that is a donor template-recruiting sequence, an editing enhancer sequence, and a protective sequence. In certain embodiments, the nucleotide sequence 5' to the 5' sequence, *e.g.*, tail sequence, if present, or 5' to the modulator stem sequence is 1-90, 1-80, 1-70, 1-60, 1-50, 1-40, 1-30, 1-20, 1-10, 10-90, 10-80, 10-70, 10-60, 10-50, 10-40, 10-30, 10-20, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, 20-30, 30-90, 30-80, 30-70, 30-60, 30-50, 30-40, 40-90, 40-80, 40-70, 40-60, 40-50, 50-90, 50-80, 50-70, 50-60, 60-90, 60-80, 60-70, 70-90, 70-80, or 80-90 nucleotides in length.

35 **[0190]** In certain embodiments, the engineered, non-naturally occurring system further comprises one or more compounds (*e.g.*, small molecule compounds) that enhance HDR and/or inhibit NHEJ. Exemplary compounds having such functions are described in Maruyama *et al.* (2015) NAT BIOTECHNOL. 33(5): 538-42; Chu *et al.* (2015) NAT BIOTECHNOL. 33(5): 543-48; Yu

et al. (2015) CELL STEM CELL 16(2): 142-47; Pinder *et al.* (2015) NUCLEIC ACIDS RES. 43(19): 9379-92; and Yagiz *et al.* (2019) COMMUN. BIOL. 2: 198. In certain embodiments, the engineered, non-naturally occurring system further comprises one or more compounds selected from the group consisting of DNA ligase IV antagonists (*e.g.*, SCR7 compound, Ad4 E1B55K protein, and Ad4 E4orf6 protein), RAD51 agonists (*e.g.*, RS-1), DNA-dependent protein kinase (DNA-PK) antagonists (*e.g.*, NU7441 and KU0060648), β 3-adrenergic receptor agonists (*e.g.*, L755507), inhibitors of intracellular protein transport from the ER to the Golgi apparatus (*e.g.*, brefeldin A), and any combinations thereof.

[0191] The sequences of the modulator nucleic acid and the targeter nucleic acid should be compatible with the Cas protein. Exemplary sequences that are operative with certain type V-A Cas proteins are provided in **Table 24**. It is understood that these sequences are merely illustrative, and other guide nucleic acid sequences may also be used with these Cas proteins.

Table 24 Type V-A Cas Protein and Corresponding Guide Nucleic Acid Sequences

Cas Protein ¹	Modulator Sequence ²	Targeter Stem Sequence	PAM ³
MAD7 (SEQ ID NO: 1)	UAAUUUCUAC (SEQ ID NO: 15)	GUAGA (SEQ ID NO: 21)	5' TTTN
MAD7 (SEQ ID NO: 1)	AUCUAC (SEQ ID NO: 791)	GUAGA (SEQ ID NO: 21)	5' TTTN
MAD7 (SEQ ID NO: 1)	GGAAUUUCUAC (SEQ ID NO: 102)	GUAGA (SEQ ID NO: 21)	5' TTTN
MAD7 (SEQ ID NO: 1)	UAAUCCAC (SEQ ID NO: 792)	GUGGG (SEQ ID NO: 22)	5' TTTN
MAD2 (SEQ ID NO: 2)	AUCUAC (SEQ ID NO: 791)	GUAGA (SEQ ID NO: 21)	5' TTTN
AsCpf1 (SEQ ID NO: 3)	UAAUUUCUAC (SEQ ID NO: 15)	GUAGA (SEQ ID NO: 21)	5' TTTN
LbCpf1 (SEQ ID NO: 4)	UAAUUUCUAC (SEQ ID NO: 15)	GUAGA (SEQ ID NO: 21)	5' TTTN
FnCpf1 (SEQ ID NO: 5)	UAAUUUCUACU (SEQ ID NO: 18)	GUAGA (SEQ ID NO: 21)	5' TTN
<i>Prevotella bryantii</i> Cpf1 (SEQ ID NO: 6)	AAUUUCUAC (SEQ ID NO: 19)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Proteocatella sphenisci</i> Cpf1 (SEQ ID NO: 7)	AAUUUCUAC (SEQ ID NO: 19)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Anaerovibrio</i> sp. RM50 Cpf1 (SEQ ID NO: 8)	AAUUUCUAC (SEQ ID NO: 19)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Moraxella caprae</i> Cpf1 (SEQ ID NO: 9)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC

Cas Protein ¹	Modulator Sequence ²	Targeter Stem Sequence	PAM ³
<i>Lachnospiraceae</i> <i>bacterium</i> COE1 Cpf1 (SEQ ID NO: 10)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Eubacterium</i> <i>coprostanoligenes</i> Cpf1 (SEQ ID NO: 11)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Smithella</i> sp. SCADC Csm1 (SEQ ID NO: 12)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Sulfuricurvum</i> sp. Csm1 (SEQ ID NO: 13)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC
<i>Microgenomates</i> (<i>Roizmanbacteria</i>) <i>bacterium</i> Csm1 (SEQ ID NO: 14)	GAAUUUCUAC (SEQ ID NO: 20)	GUAGA (SEQ ID NO: 21)	5' TTTC

¹ The amino acid sequences of the Cas proteins are provided at the end of the specification.

² It is understood that a “modulator sequence” listed herein may constitute the nucleotide sequence of a modulator nucleic acid. Alternatively, additional nucleotide sequences can be comprised in the modulator nucleic acid 5' and/or 3' to a “modulator sequence” listed herein.

5 ³ In the consensus PAM sequences, N represents A, C, G, or T. When the PAM sequence is preceded by “5’,” it means that the PAM is immediately upstream from the target nucleotide sequence when using the non-target strand (*i.e.*, the strand not hybridized with the spacer sequence) as the coordinate.

[0192] In certain embodiments, the targeter nucleic acid of the engineered, non-naturally occurring system comprises a targeter stem sequence listed in **Table 24**. In certain embodiments, the targeter nucleic acid and the modulator nucleic acid of the engineered, non-naturally occurring system comprise, respectively, a targeter stem sequence and a modulator sequence listed in the same line of **Table 24**. It is understood that one or more 3' or 5' ends of a modulator sequence may contain chemical modifications, and/or 3' end of targeter stem sequence or 5' end of a targeter stem sequence, depending on orientation, may contain one or more chemical modifications. In certain embodiments, the engineered, non-naturally occurring system further comprises a Cas nuclease comprising the amino acid sequence set forth in the SEQ ID NO listed in the same line of **Table 24**. In certain embodiments, the engineered, non-naturally occurring system is useful for targeting, editing, or modifying a nucleic acid comprising a target nucleotide sequence close or adjacent to (*e.g.*, immediately downstream of) a PAM listed in the same line of **Table 24** when using the non-target strand (*i.e.*, the strand not hybridized with the spacer sequence) as the coordinate.

[0193] In certain embodiments, the engineered, non-naturally occurring system is tunable or inducible. For example, in certain embodiments, the targeter nucleic acid, the modulator

nucleic acid, and/or the Cas protein can be introduced to the target nucleotide sequence at different times, the system becoming active only when all components are present. In certain embodiments, the amounts of the targeter nucleic acid, the modulator nucleic acid, and/or the Cas protein can be titrated to achieve desirable efficiency and specificity. In certain

5 embodiments, excess amount of a nucleic acid comprising the targeter stem sequence or the modulator stem sequence can be added to the system, thereby dissociating the complex of the targeter nucleic acid and modulator nucleic acid and turning off the system.

C. Cas Proteins

10 **[0194]** In certain embodiments, compositions and methods provided herein include a Cas protein, *e.g.*, a Cas nuclease. The present invention also provides an engineered, non-naturally occurring system comprising a guide nucleic acid (*e.g.*, a dual guide nucleic acid) disclosed herein, for example a guide nucleic acid described in section IA, IA1, and IB. In certain

15 embodiments, the engineered, non-naturally occurring system further comprises the Cas nuclease, such as a Type I, II, III, IV, V, or VI nuclease, in some cases a Type V nuclease, for example, a Type V-A, V-C, or V-D Cas nuclease, such as a Type VA nuclease, including but not limited to a Cpf1 nuclease, derivative, or variant; a MAD nuclease, derivative, or variant; a ART nuclease, derivative, or variant; a Csm1 nuclease, derivative, or variant; or an ABW nuclease, derivative, or variant; specific examples are provided in this section. In certain embodiments, the

20 modified guide nucleic acid and the Cas nuclease are present in a ribonucleoprotein (RNP) complex. In certain embodiments, the system also includes an editing sequence (donor sequence or donor template) having a change in sequence relative to the sequence of a target region.

[0195] The terms “CRISPR-Associated protein,” “Cas protein,” and “Cas,” as used interchangeably herein, can include a naturally occurring Cas protein or an engineered Cas

25 protein. Non-limiting examples of Cas protein engineering includes but are not limited to mutations and modifications of the Cas protein that alter the activity of the Cas, alter the PAM specificity, broaden the range of recognized PAMs, and/or reduce the ability to modify one or more off-target loci as compared to a corresponding unmodified Cas. In certain embodiments, the altered activity of the engineered Cas comprises altered ability (*e.g.*, specificity or kinetics) to

30 bind the naturally occurring crRNA or engineered modified dual guide nucleic acids, altered ability (*e.g.*, specificity or kinetics) to bind the target nucleotide sequence, altered processivity of nucleic acid scanning, and/or altered effector (*e.g.*, nuclease) activity. A Cas protein having the nuclease activity is referred to as a “CRISPR-Associated nuclease” or “Cas nuclease,” as used interchangeably herein. In certain cases, as will be clear from context, a Cas nuclease lacking

35 nuclease activity can also be referred to as a Cas nuclease.

[0196] The Cas nuclease that a complex comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating can be any suitable Cas nuclease, such as a Type I, II, III, IV, V, or VI nuclease, such as a Type V nuclease. In certain embodiments, provided herein are methods and compositions that include a modified guide nucleic acid, *e.g.*, RNA, as described herein, for example in section IA or section IA1, and a Type I, II, III, IV, V, or VI nuclease. In certain embodiments, provided herein are methods and compositions that include a modified guide nucleic acid, *e.g.*, RNA, as described herein, for example in section IA, IA1, or IB and a Type V, nuclease.

[0197] In certain embodiments, the Cas nuclease that a complex comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating is a type V-A, type V-C, or type V-D Cas nuclease. In certain embodiments, the Cas nuclease is a type V-A nuclease. In certain embodiments, the Cas nuclease is a Type V-E nuclease. In certain embodiments, the Cas nuclease is a MAD, ART, or ABW nuclease, as described herein. In certain embodiments, provided herein are methods and compositions that include a modified guide nucleic acid, *e.g.*, RNA, as described herein, for example in section IA, IA1, or IB, and a Type V-A, Type V-C, Type V-E, or Type V-D Cas nuclease.

[0198] In certain embodiments, a nuclease that a complex comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating can be a Type V-A Cas nuclease. When a Type V-A Cas nuclease is used with a split gRNA as described herein, it may be considered a Type V-E Cas nuclease, and “Type V-A” may be considered equivalent to “Type V-E” herein in this context. In certain embodiments, the type V-A Cas nuclease comprises Cpf1 or a variant or derivative thereof, a MAD nuclease or a variant or derivative thereof, a Csm1 nuclease or a variant or derivative thereof, an ART nuclease or variant or derivative thereof, or an ABW nuclease or variant or derivative thereof. In certain embodiments, a composition comprises a Type V-A nuclease and a modified guide nucleic acid, *e.g.*, modified dual guide RNA, as described herein, *e.g.*, in Section IA, IA1, or IB.

[0199] In certain embodiments, the type V-A Cas nucleases comprises Cpf1 or a derivative thereof. Cpf1 proteins are known in the art and are described in U.S. Patent Nos. 9,790,490 and 10,113,179. Cpf1 orthologs can be found in various bacterial and archaeal genomes. For example, in certain embodiments, the Cpf1 protein is derived from *Francisella novicida* U112 (Fn), *Acidaminococcus* sp. BV3L6 (As), *Lachnospiraceae* bacterium ND2006 (Lb), *Lachnospiraceae* bacterium MA2020 (Lb2), *Candidatus Methanoplasma termitum* (CMT), *Moraxella bovoculi* 237 (Mb), *Porphyromonas crevioricanis* (Pc), *Prevotella disiens* (Pd), *Francisella tularensis* 1, *Francisella tularensis* subsp. *novicida*, *Prevotella albensis*, *Lachnospiraceae* bacterium MC2017 1, *Butyrivibrio proteoclasticus*, *Peregrinibacteria*

bacterium GW2011_GWA2_33_10, Parabacterium bacterium GW2011_GWC2_44_17, Smithella sp. SCADC, Eubacterium eligens, Leptospira inadai, Porphyromonas macacae, Prevotella bryantii (Pb), Proteocatella sphenisci (Ps), Anaerovibrio sp. RM50 (As2), Moraxella caprae (Mc), Lachnospiraceae bacterium COE1 (Lb3), or Eubacterium coprostanoligenes (Ec).

- 5 **[0200]** In certain embodiments, the type V-A Cas nuclease comprises AsCpf1 or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid
- 10 sequence set forth in SEQ ID NO: 3. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 3.

AsCpf1 (SEQ ID NO: 3)

MTQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQQFIEEDKARNDHYKELKPIIDRIYKTYADQ
CLQLVQLDWENLSAAIDSYRKEKTEETRNALIEEQATYRNAIHDFIGRTDNLTDANKRHAIEI
15 YKGLFKAELFNGKVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSAEIDISTAI PHR
IVQDNFPKFKENCHIFTRLITAVPSLREHFENVKKAIGIFVSTSIIEEVFSFPFYNQLLTQTQID
LYNQLLGGISREAGTEKIKGLNEVLNLAIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFIL
EEFKSDEEVIQSFCYKTLRNENVLETAELFNELNSIDLTHIFISHKKLETISSALCDHWD
LRNALYERRISELTGKITKSAKEKVQRSLKHEDINLQEIISAAGKELSEAFKQKTSEILSHAH
20 ALDQPLPTTLKKQEEKEILKSQDLSLLGLYHLLDWFVAVDESNEVDPEFSARLTGKILEMEPSLS
FYNKARNYATKKPYSVEKFKLNFQMPTLASGWDVNKEKNNGAILFVKNGLYYLGIMPQKQGRYK
ALSFEPTTEKTSEGFDMYYDYFPDAAKMIPKCSTQLKAVTAHFQTHHTPIILLSNFIETPLEITK
EIYDLNNPEKEPKKFQTAYAKKTGDQKGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQY
KDLGEYYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGL
25 FSPENLAKTSIKLNGQAEFYRPKSRMKRMAHRLGEKMLNKKLDQKTPIPDITLYQELYDYVNH
RLSHDLSDEARALLPNVITKEVSHEIIKDRRFTSDKFFHFVPITLNYQAANSPSKFNQVRNAYL
KEHPETPIIGIDRGERNLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSV
VGTIKDLKQGYLSQVIHEIVDLMIHYQAVVLENLNFQFKSKRTGIAEKAVYQQFEKMLIDKLN
CLVLKDYPAEKVGGVLPYQLTDQFTSFAKMGTSQGFIFYVPAPYTSKIDPLTGFVDPFVWKT
30 KNHESRKHFLLEGFDLHYDVKTGDFILHFKMNRNLSFQRLPGFMPAWDIVFEKNETQFDAQGT
PFIAGKRIVPVIENHRFTGRYRDLYPANELIALLEEKGIVFRDGSNILPKLLENDSDHAIDTMV
ALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNPPEWPMADANGAYHIALKGQQLLNH
LKESKDLKLQNGISNQDWLAYIQELRN

- 35 **[0201]** In certain embodiments, the type V-A Cas nuclease comprises LbCpf1 or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid
- 40 sequence set forth in SEQ ID NO: 4. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 4.

LbCpf1 (SEQ ID NO: 4)

MSKLEKFTNCYSLSKTLRFKAI PVGKTQENIDNKRLLVEDEKRAEDYKGVKKLLDRYYLSFIND
 VLHSIKLKNLNNYISLFRKKTRTEKENKELENLEINLRKEIAKAFKGNEGYKSLFKKDI IETIL
 PEFLLDDKDEIALVNSFNGFTTAFTGFFDNRENMFSEEAKSTSI AFRCINENLTRYI SNMDI FEK
 5 VDAIFDKHEVQEIKEKILNSDYDVEDFFEGEFFFNFVLTQEGIDVYNAI IGGFVTESGEKIKGLN
 EYINLYNQKTKQKLPKFKPLYKQVLSDRSLSFYGEGYTSDEEVLEVFRNTLNKNSEIFSSIKK
 LEKLFKNFDEYSSAGIFVKNGPAISTISKDIFGEWNVIRDKWNAEYDDIHLKKKAVVTEKYEDD
 RRKSFKKIGSFSLEQLQEYADADLSVVEKLKEII IQKVEIYKVYGSSEKLFDAADFVLEKSLKK
 10 NDAVVAIMKDLLDSVKS FENYIKAFFGEGKETNRDES FYGDFVLAYDILLKVDHIYDAIRNYVT
 QKPYSKDKFKLYFQNPQFMGGWDKDKETDYRATILRYGSKYYLAIMDKKYAKCLQKIDKDDVNG
 NYEKINYKLLPGPNKMLPKVFFSKWMAAYNPSEDIQKIYKNGTFKKGDMFNLNDCHKLIDFFK
 DSISRYPKWSNAYDFNFSETEKYKDIAGFYREVEEQGYKVSFESASKKEVDKLVEEGKLYMFQI
 YNKDFS DSKSHGTPNLHTMYFKLLFDENNHGQIRLSGGAELFMRRASLKKEELVVHPANSPIANK
 15 NPDNPKKTTTTLSYDVYKDKRFSEDQYELHIPIAINKCPKNIFKINTEVRVLLKHDDNPYVIGID
 RGERNLLYIVVVDGKGNIVEQYSLNEIINNENGIRIKTDYHSLLDKKEKERFEARQNWT SIENI
 KELKAGYISQVVHKICELVEKYDAVIALEDLNSGFKNSRVKVEKQVYQKFEKMLIDKLNVMVDK
 KSNPCATGGALKGYQITNKFESFKSMSTQNGFI FYIPAWLTSKIDPSTGTFVNLLKTKYTSIADS
 KKFIS SFDRIMYVPEEDLFEFALDYKNFSRTDADYIKKWKLYSYGNRIRI FRNPKKNNVFDWEE
 VCLTSAYKELFNKYGINYQQGDIRALLCEQSDKAFYSSFMALMSLMLQMRNSITGRTDVDFLIS
 20 PVKNSDGI FYDSRNYEAQENAILPKNADANGAYNIARKVLWAIGQFKKAEDEKLDKVKIAISNK
 EWLEYAQT SVKH

[0202] In certain embodiments, the type V-A Cas nuclease comprises FnCpf1 or a
 variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid
 sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at
 25 least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
 least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid
 sequence set forth in SEQ ID NO: 5.

FnCpf1 (SEQ ID NO: 5)

MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEE
 30 ILSSVCISEDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNL I
 DAKKGQESDLILWLKQSKDNGIELFKANSDITDIDEALEI IKSFKGWTTYFKGFHENRKNVYSS
 NDIPTSI IYRIVDDNLPKFLENKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQR
 VFSLDEVFEIANFNNYLNQSGITKFNTIIIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYK
 35 MSVLFKQILSDTESKSFVIDKLEDDSDVVTMQSFYEQIAAFKTVEEKSIKETLSLLFDDLKAQ
 KLDLSKIYFKNDKSLTDL SQQVFDDYSVIGTAVLEYITQQIAPKNLDNPSKKEQELIAKKEKA
 KYLSLETIKLAL EEFNKHRDIDKQCRFEEILANFAAIPMI FDEIAQNKDNLAQISIKYQNQGKK
 DLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYLVFEECYFELANIV
 PLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLLGVMNKKNNKI
 40 FDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNGSPQ
 KGYEKF EFNIEDCRKFIDFYKQSI SKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLT FEN
 ISESYIDSVVNQGKLYLFQIYNKDFSAYS SKGRPNLHTLYWKALFDERNLQDVVYKLNGEAEIFY
 RKQSI PKKITHPAKEAIANKNDNP KKESVFEYDLIKDKRFTE DKFFFHC PITINFKSSGANKF
 NDEINLLLKEKANDVHILSIDRGERHLAYYTLVDGKGNI I KQDTFNI I GNDRMKTNYHDKLAAI
 EKDRDSARKDWKKINNI KEMKEGYLSQVVHEIAKLVI EYNAIVVFEDLNF GFKRGRFKVEKQVY
 45 QKLEKMLIEKLNLYLVFDNEFDKTGGVLRAYQLTAPFETFKKMGKQTGI IYYPAGFTSKICPV
 TGFVNQLYPKYESVSKSQEFFSKFDKICYNLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLI
 NFRNSDKHNHWD TREVYPTKELEKLLKDYSIEYGHGECIKAAICGESDKKFFAKLTSVLNTILQ

MRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDADANGAYHIGLKGLMLLGRIKNNQEGK
KLNLVIKNEEYFEFVQNRNN

[0203] In certain embodiments, the type V-A Cas nuclease comprises *Prevotella bryantii*

5 Cpf1 (PbCpf1) or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 6. In certain embodiments, the type V-A Cas
10 protein comprises the amino acid sequence set forth in SEQ ID NO: 6.

Prevotella bryantii Cpf1 (PbCpf1) (SEQ ID NO: 6)

MQINNLIKIIYMKFTDFTGLYSLSKTLRFELKPIGKTLENIKKAGLLEQDQHRADSYKKVKKIID
EYHKAFIEKSLSNFELKYQSEDKLDSLEEYLMYYSMKRIEKEKDKFAKIQDNLRKQIADHLKG
DESYKTI FSKDLIRKNLPDFVKSDEERTLIKEFKDFTTYFKGFYENRENMYSAEDKSTAI SHRI
15 IHENLPKFVDNINAFSKIILIPELREKLNQIYQDFEYLNVESIDEI FHLDYFSMVMQKQIEV
YNAIIGGKSTNDKKIQGLNEYINLYNQKHKDKCLPKLKLFFKQILSDRIAISWLPDNFKDDQEA
LDSIDTCYKNLLNDGNVLGEGNLKLLLENIDTYNLKGI FIRNDLQLTDISQMYASWNVIQDAV
ILDLKKQVSRKKKESAEDYNDRLKKLYTSQESFSIQYLNDCLRAYGKTENIQDYFAKLGAVNNE
HEQTINLFAQVRNAYTSVQAILTTPYPENANLAQDKETVALIKNLLDSLKRLQRFIKPLLKGD
20 ESDKDERFYGDFTPLWETLNQITPLYNMVRNYMTRKPYSQEKIKLNFENSTLLGGWDLNKEHDN
TAIILRKNGLYYLAIMKKSANKIFDKDKLDNSGDCYEKMVYKLLPGANKMLPKVFFSKSRIDEF
KPSENIIENYKKGTHKKGANFNLDCHNLIDFFKSSISKHEDWSKFNHFHSDTSSYEDLSDFYR
EVEQQGYSSISFCDSVEYINKMVEKGDLYLFQIYNKDFSEFSKGT PNMHTLYWNSLFSKENLNN
IIYKLNQAEIFFRKKSLNYKRPHPAHQAIKNKNKCNEKKESIFDYDLVKDKRYTVDFKQFHV
25 PITMNFKSTGNTNINQQVIDYLRTEDDTHIIGIDRGERHLLYLVIDSHGKIVEQFTLNEIVNE
YGGNIYRTNYHDLDTREQNREKARESWQTIENTIKELKEGYISQVIHKITDLMQKYHAVVLED
LNMGMGRGRQKVEKQVYQKFEEMLINKLNYLVNKKADQNSAGLLHAYQLTSKFESFQKLKQKS
GFLFYIPAWNTSKIDPVTGFVNLFDTRYESIDKAKAFFGKFDSIRYNADKDWFEFAFDYNNFTT
KAEGTRTNWTICTYGSRI RTFRNQAKNSQWDNEEIDLTKAYKAFFAKHGINIYDNIKEAIAMET
30 EKSFFEDLLHLLKLTLQMRNSITGTTTDYLISPVHDSKGNFYDSRICDNSLPANADANGAYNIA
RKGLMLLIQQIKDSTSSNRKFSPITNKDWLIFAQEKPYLND

[0204] In certain embodiments, the type V-A Cas nuclease comprises *Proteocatella*

sphenisci Cpf1 (PsCpf1) or a variant thereof. In certain embodiments, the type V-A Cas protein
35 comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 7. In certain embodiments, the type V-A
Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 7.

40 *Proteocatella sphenisci* Cpf1 (PsCpf1) (SEQ ID NO: 7)

MENFNKLYPINKTLRFELRPYGKTLENFKKSGLLEKDAFKANSRRSMQAIIDEKFKETIEERLK
YTEFSECDLGNMTSKDKKITDKAATNLKKQVILSFDEIFNNYLKPKDNIDALFKNDPSNPVIS
TFKGFTTYFVNFFFEIRKHI FKGESSGSMAYRIIDENLT TYLNNIEKIKKLPEELKSQLEGIDQI
DKLNNYNEFITQSGITHYNEIIIGGISKENVKIQGINEGINLYCQKNKVKLPRLTPLYKMILSD
45 RVSNSFVLDTIENDTELIEMISDLINKTEISQDVIMSDIQNIFIKYKQLGNLPGISYSSIVNAI

CSDYDNNFGDGKRKKSSENDNRKKHLETNVYSINYISELLTDTDVSSNIKMRYKELEQNYQVCKE
 NFNATNWMNIKNIKQSEKTNLIKDLLDILKSIQRFYDLFDIVDEDNPSAEFYTWLSKNAEKLD
 FEFNSVYNKSRNYLTRKQYSDKKIKLNFDSPTLAKGWDANKEIDNSTIIMRKFNNDRGDYDYFL
 GIWNKSTPANEKIIPLEDNGLFEKMQYKLYPDPSKMLPKQFLSKIWKAKHPTTPEFDKKYKEGR
 5 HKKGPDFEKEFLHELIDCFKHGLVNHDEKYQDVFGFNLNRTEDYNSYTEFLEDVERCNYNLSFN
 KIADTSNLINDGKLYVFQIWSKDFSIDSKGTKNLNTIYFESLFSEENMIEKMFKLSGAEI FYR
 PASLNYCEDIIKKGHHHAELKDKFDYPIIKDKRYSQDKFFFHVPMVINIKSEKLSKSLNNRTN
 ENLGQFTHIIGIDRGERHLLIYLTVDVSTGEIVEQKHLDEIINTDTKGVEHKTHYLNKLEEKSK
 TRDNERKSWEAIEETIKELKEGYISHVINEIQKLQEKYNALIVMENLNYGFKNSRIKVEKQVYQK
 10 FETALIKKFNYIIDKDPETYIHGYQLTNPITTLDKIGNQSGIVLYIPAWNTSKIDPVTGFVNL
 LYADDLKYKNQEQAQSFQKIDNIYFENGFEKFDIDFSKWNRYSSISKTQWTLTSYGTRIQTFR
 NPQKNKWDSAEYDLTEEFKLIILNIDGTLKSQDVETYSKFKMSLFKLMLQLRNSVTGTDIDYMS
 PVTDKTGTHFDSRENKLNLPADADANGAYNIARKGIMAIENIMNGISDPLKISNEDYLKYIQNQ
 QE

15 **[0205]** In certain embodiments, the type V-A Cas nuclease comprises *Anaerovibrio* sp.
 RM50 Cpf1 (As2Cpf1) or a variant thereof. In certain embodiments, the type V-A Cas protein
 comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least
 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least
 20 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical
 to the amino acid sequence set forth in SEQ ID NO: 8. In certain embodiments, the type V-A
 Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 8.

Anaerovibrio sp. RM50 Cpf1 (As2Cpf1) (SEQ ID NO: 8)

MVAFIDEFVGQYPVSKTLRFEARPVPEPKKWLESDQCSVLFNDQKRNEYGYVLKELLDDYRAY
 25 IEDALTSFTLDKALLENAYDLYCNRDTNAFSSCCEKLRKDLVKAFGNLKDYLLGSDQLKDLVKL
 KAKVDAPAGKGKKKI EVD SRLINWLNNAKYS AEDREKYI KAIESFEGFVTYLTNYKQARENMF
 SSEDKSTAI AFRVIDQNMVTFGNIRIYEKIKAKYPELYSALKGFEEKFFSPTAYSEILSQSKID
 EYNYQCIGRPIDDA DFKGVNSLINEYRQKNGIKARELPVMSMLYKQILSDRDNFSMFSEVINRNE
 EAIECAKNGYKVS YALFNELLQLYKKIFTEDNYGNIYVKTPLELSQALFGDWSILRNALDNG
 30 KYDKDIINLAELEKYFSEYCKVLDAADAAKIQDKFNLKDYFIQKNALDATLPDLDKITQYKPHL
 DAMLQAIRKYKLF SMYNGRKKMDVPENGIDFSNEFNAIYDKLSEFSILYDRI RN FATKKPY SDE
 KMKLSFNMPTMLAGWDYNNETANGCFLFIKDGKYFLGVADSKSKNIFDFKKNPHLLDKYSSKDI
 YYKVYKQVSGSAKMLPKVV FAGSNEKIFGHLISKRILEIREKKLYTAAAGDRKAVA EWIDFMK
 35 SAIAIHPEWNEYFKFKFN TAEYDNANKFYEDIDKQTSLEKVEIPT EYIDEMVSQHKL YLFQL
 YTKDFS DKKKKKGTDNLHTMYWHGVFSDENLKAVTEGTQPIIKLNGEAEFMRNPSIEFQVTHE
 HNKPIANKNPLNTKKESVFNYDLIKDKRYTERKFYFHCPI TLNFRADKPIKYNEKINRFVENNP
 DVCIIIGIDRGERHLLYYTVINQTDGILEQGS LNKISGSYTNDKGEKVNKETDYHDL LDRKEK GK
 HVAQQAWETIENIKELKAGYLSQVYKLTQLMLQYNAVIVLENLNVGFKRGR TKVEKQVYQKFE
 KAMIDKLNLYLVFKDRGYEMNGSYAKGLQLTDKFESFDKIGKQTGCIYYVIPS YTS HIDPKTG FV
 40 NLLNAKLRYENITKAQDTIRKFDSISYNAKADYFEFAFDYRSFGVDMARNEWVVC TCGDLRWEY
 SAKTRET KAYSVTDR LKELFKAHGIDYVGGENLVSHITEVADKHFLSTLLFYLRVLVKMRYTVS
 GTENENDFILSPVEYAPGKFFDSREATSTEP MNADANGAYHIALKGLMTIRGIEDGKLHNYGKG
 GENAAWFKFMQONQEYKNNG

45 **[0206]** In certain embodiments, the type V-A Cas nuclease comprises *Moraxella caprae*
 Cpf1 (McCpf1) or a variant thereof. In certain embodiments, the type V-A Cas protein comprises
 an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at

least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 9. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 9.

5 *Moraxella caprae* Cpf1 (McCpf1) (SEQ ID NO: 9)

MLFQDFTHLYPLSKTMRFEKLPKIGKTLLEHIHAKNFLSQDETMADMYQKVKAILDDYHRDFIADM
MGEVKLTKLAEFYDVYLKFRKNPKDDGLQKQLKDLQAVLRKEIVKPIGNNGGKYKAGYDRLFGAK
LFDKGKELGDLAKFVIAQEGESSPKLAHLAHFEKFSTYFTGFHDNRKNMYSDEDKHTAITYRLI
HENLPRFIDNLQILATIKQKHSALYDQIINELTASGLDVSLASHLDGYHKLLTQEGITAYNTLL
10 GGISGEAGSRKIQGINELINSHHNQHCHKSERIAKLRPLHKQILSDGMGVSFLPSKFADDSEMC
QAVNEFYRHYADVFAKVQSLFDGFDHQQKDGIYVEHKNLNELSKQAFGDFALLGRVLDGYVVDV
VNPEFNERFAKAKTDNAKAKLTKEKDKFIKGVHSLASLEQAI EHYTARHDDSVQAGKLGQYFK
HGLAGVDNPIQKIHNNHSTIKGFLERERPAGERALPKIKSGKNPEMTQLRQLKELLDNALNVAH
FAKLLTTKTTLDNQDGNFYGEFGALYDELAKEIPTLYNKVRDYLKQKPFSTEKYKLNFGNPTLLN
15 GWDLNKEKDNFGIILQKDGCCYLLALLDKAHKKVFDNAPNTGKNVYQKMIYKLLPGPNKMLPKVF
FAKSNLDYYNPSAELLDKYAQGTHKKGNFNLKDCHALIDFFKAGINKHPEWQHFGFKFSPTSS
YQDLSDFYREVEFPQGYQVKFVDINADYINELVEQGGYLYLFQIYNKDFSPKAHGKPNLHTLYFKA
LFSKDNLANPIYKLNGEAQIFYRKASLDMNETTIHRAGEVLENKNPDNPKKRQFVYDIIKDKRY
TQDKFMLHVPITMNFVQGMTIKEFNKKVNQSIQQYDEVNIGIDRGERHLLYLTVINSKGEIL
20 EQRSINDITTASANGTQMTTPYHKILDKREIERLNARVWGGEIETIKELKSGYLSHVHQSLSQL
MLKYNAIVVLEDLNFQFGRGRFKVEKQIYQNFENALIKKLNHLVLKDEADDEIGSYKNALQLTN
NFTDLKSIGKQTGFLFYVPAWNTSKIDPETGFVDLLKPRYENIAQSQAFFGKFDKICYNADKDY
FEFHIDYAKFTDKAKNSRQIWKICSHGDKRYVYDKTANQNKGATKGINVNDELKSLFARHHIND
KQPNLVMIDICQNNDEKFEHKSLLIYLLKTLLALRYSNASSDEDFILSPVANDEGMFFNSALADDTQ
25 PQNADANGAYHIALKGLWVLEQIKNSDDLNVKVLKLAIDNQTLNFAQNR

[0207] In certain embodiments, the type V-A Cas nuclease comprises *Lachnospiraceae* bacterium COE1 Cpf1 (Lb3Cpf1) or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%,
30 at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 10. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 10.

Lb3Cpf1 (SEQ ID NO: 10)

35 MHENNGKIADNFIGIYPVSKTLRFELKPVGKTQEYIEKHGILDEDLKRAGDYKSVKKIIDAYHK
YFIDEALNGIQLDGLKNYYELYEKRDNNNEEKEFQKIQMSLRKQIVKRFSEHPQYKYLFKKELI
KNVLPEFTKDNAEEQTLVKSFQEFTTYFEGFHQNRKNMYSDEEKSTAIAYRVVHQNLPKYIDNM
RIFSMILNTDIRSDLTELFNNLKTMDITIVEEYFAIDGFNKVVNQKIDVYNTILGAFSTDDN
TKIKGLNEYINLYNQKNKAKLPKLKPLFKQILSDRDKISFIPEQFSDTEVLEAVDMFYNRLLQ
40 FVIENEGQITISKLLTNFSAYDLNKIYVKNDDTISAI SNDLFDDWSYISKAVRENYDSENVKN
KRAAAEYEEKKEKALSKIKMYSIEELNFFVKKYSCNECHIEGYFERRILEILDKMRYAYESKIL
HDKGLINNISLCQDRQAISELKDFLDSIKEVQWLLKPLMIGQEADKEEAFYTELLRIWEELEP
ITLLYNKVRNRYVTKKPYTLEKVKLNFYKSTLLDGDKNKEKDNLGIILLKDGQYYLGIMNRRNN
KIADDAPLAKTDNVYRKMEYKLLTKVSANLPRI FLKDKYNPSEEMLEKYEKGTHLKGENFCIDD
45 CRELIDFFKKGIKQYEDWGQDFKFSDESYYDISAFYKEVEHQGYKITFRDIDETYIDSLVNE
GKLYLFQIYNKDFSPYSGTKNLHTLYWEMLFSQQNLQNIIVYKLNAGNAEIFYRKASINQKDVVV

HKADLP IKNKDPQNSKKESMFDYDI I KDKRFTCDKYQFHVPI TMNFKALGENHFNKVNRLIHD
 AENMHII GIDRGERNLI YLCMIDMKGNIVKQISLNEI I SYDKNKLEHKRNYHQLLKTREDEKNS
 ARQSWQTIHTIKELKEGYLSQVIHVITDLMVEYNAIIVVLEDLNFQFKQGRQKFERQVYQKFEKM
 LIDKLNLYLVDKSKGMDDEDGGLLHAYQLTDEFKSKQLGKQSGFLYYI PAWNTSKLDPTTG FVNL
 5 FYTKYESVEKSKEFINNFTSILYNQEREYFEFLFDYSAFTSKAEGSRLKWTVC SKGERVETYRN
 PKKNNEWDTQKIDLT FELKKLFNDYSISLLDGDLDREQMGKIDKADFYKKFMKLFALIVQMRNSD
 EREDKLISPVLNKYGAFFETGKNERMPLDADANGAYNIARKGLWII EKIKNTDVEQLDKVKLTI
 SNKEWLQYAQEHIL

[0208] In certain embodiments, the type V-A Cas nuclease comprises *Eubacterium*

10 *coprostanoligenes* Cpf1 (EcCpf1) or a variant thereof. In certain embodiments, the type V-A Cas
 protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%,
 at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at
 least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%
 identical to the amino acid sequence set forth in SEQ ID NO: 11. In certain embodiments, the
 15 type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 11.

Eubacterium coprostanoligenes Cpf1 (EcCpf1) (SEQ ID NO: 11)

MDFFKNDMYFLCINGI I V I SKLFAYLFLMYKRGVVM I KDNFVNVYSLSKTIRMALIPWGKTEDN
 FYKKFLLEEDEERAKNY I KVKGYMDEYHKNFIESALNSVVLNGVDEYCELYFKQNKSDSEVKKI
 ESLEASMRKQISKAMKEYTV DGVKIYPLLSKKEFIRELLPEFLTQDEE IETLEQFNDSTYFQG
 20 FWENRKN IYTDEEKSTGVPYRCINDNLPKFLDNVKSFEKVILALPQKAVDELNANFNGVYNVDV
 QDVFSVDYFNFVLSQSGIEKYNNIIGGYSNSDASKVQGLNEKINLYNQQIAKSDKSKKLPLLP
 LYKQILSDRSSLSFIPEKFKDDNEVLNSINVLYDNIAESLEKANDLMSDIANYNTDNIFISSGV
 AVTDISKKVFGDWSLIRNNWNDEYESTHKKGKNEEKFEYKEDKEFKKIKSFSVSELQRLANS DL
 25 SIVDYLVDDESASLYADIKTAYNNAKDLLSNEYSHSKRLSKNDDAIELIKSFLDSIKNYEAF LKP
 LCGTGKEESKDNAFYGAFLCFFEEIRQVDVAVYNKVRNHITQKPYSDNKIKLNFQNPQFLAGWDK
 NKERAYRSVLLRNGEKY YLAIMEKGKSKLFEDFPEDESSPFEKIDYKLLPEPSKMLPKVFFATS
 NKDLFNPSDEILNIRATGSFKKGDSFNLDDCHKFIDFYKAS IENHPDWSKFD FDFSETNDYEDI
 SKFFKEVSDQGYSIGYRKIS ESYLEEMVDNGSLYMFQLYNKDFSENRRKSKGTPNLHTLYFKMLF
 DERNLEDVVYKLSGGAEMFYRKPSIDKNEMIVHPKNQPIDNKNPNNVKKTSTFEYDIVKDMRYT
 30 KPQFQLHLP IVLNFKANSKGYINDDVRNVLKNS EDTYVIGIDRGERNLVYACVVDGNGKLVEQV
 PLNVI EADNGYKTDYHKLLNDREEKRNEARKSWKTIGNIKELKEGYISQVVHKICQLVVKYDAV
 IAMEDLNSGFVNSRKKVEKQVYQKFERMLTQKLNLYLVDKKLDPNEMGGLLNAYQLTNEATKVRN
 GRQDGII FYI PAWLTSKIDPTTG FVNL LKPKYNSVSASKEFFSKFDEIRYNEKENYFEFSFN YD
 NFPKCNADFKREWTVCTYGDRI RTFRDPENNNKFNSEVVVLNDEFKNLFVEFDIDYTDNLKEQI
 35 LAMDEKSFYKKLMGLLSLT LQMRNSISKNVDVDYLI SPVKNSNGEFYDSRNYDITSSLP CDADS
 NGAYNIARKGLWAINQIKQADDETKANISIKNSEWLQYAQNCDEV

[0209] In certain embodiments, the type V-A Cas nuclease is not Cpf1. In certain
 embodiments, the type V-A Cas nuclease is not AsCpf1.

[0210] In certain embodiments, the type V-A Cas nuclease comprises MAD1, MAD2,
 40 MAD3, MAD4, MAD5, MAD6, MAD7, MAD8, MAD9, MAD10, MAD11, MAD12, MAD13,
 MAD14, MAD15, MAD16, MAD17, MAD18, MAD19, or MAD20, or derivatives or variants
 thereof. MAD1-MAD20 are known in the art and are described in U.S. Patent No. 9,982,279.

[0211] In certain embodiments, the type V-A Cas nuclease comprises MAD7 or a variant
 thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at

least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 1. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 1.

MAD7 (SEQ ID NO: 1)

MNNGTNNFQNFIGISSLQKTLRNALIP TETTQQFIVKNGIIKEDEL RGENRQILKDIMDDYYRG
 F ISETLSSIDDIDWTS LFEKMEIQ LKNGDNKD TLIKEQTEYRKAIHKKFANDDRFKNMFS AKLI
 SDILPEFVIHNNNYSASEKEEKTQVIKLF SRFATSF KDYFKNRANCFSADDISSSSCHRI VNDN
 10 AEIFFSNALVYRRIVKSLSNDDINKISGDMKDSLKEMSLEETIYSYKEYGEFITQEGISFYNDIC
 GKVNSFMNLYCQKNKENKNLYKLQKLHKQILCIADTSYEVYPYKFESDEEVYQSVNGFLDNISSK
 HIVERLRKIGDNYNGYNLDKIYIVSKFYESVSQKTYRDWETINTALEIHYNILPGNGKSKADK
 VKKAVKNDLQKSITEINELVSNYKLCSDDN IKAETIHEISHILNNFEAQELKYNPEIHLVESE
 LKASELKNVLDVIMNAFWCSVFMTEELVDKDN NFYAELEEIYDEIYPVISLYNLVRNYVTQKP
 15 YSTKKIKLNFGIPTLADGWSKSKESYNNAILMRDNLYYLGI FNAKNKPKDKIIEGNTSENKGD
 YKKMIYNLLPGPNKMIPKVFLSSKTGVET YKPSAYILEGYKQNKHIKSSKDFDITFCHDLIDYF
 KNCIAIHPEWKNFGFDFSDTSTYEDISGFY REVELQGYKIDWTYI SEKDIDLLQEKGQLYLFQI
 YNKDFSKKSTGNDNLHTMYLKNL FSEENLKDIVLKLNGEAEIFFRKSSIKNPIIHKKGSILVNR
 TYEAEKDKQFGNIQIVRKNI PENIYQELYKYFNDKSDKELSDEAAKLKNVVGHEAATNIVKDY
 20 RYTYDKYFLHMPITINFKANKTGFINDRILQYIAKEKDLH VIGIDRGERNLIYVSVIDTCGNIV
 EQKSFNIVNGYDYQIKLKQQEGARQIARKEWKEIGKIKEIKEGYLSLVIHEISKMVIKYNALIA
 MEDLSYGFKKGRFKVERQVYQKFETMLINKLNYLVFKDISITENGGLLKGYQLTYIPDKLKNVG
 HQCGCIFYPAAAYTSKIDPTTG FVNI FKFKDLTVDAKREFIKKFDSIRYDSEKNLFCFTFDYNN
 FITQNTVMSKSSWSVYTYGVRIKRRFVNGRFSNESDTIDITKMEKTLEMTDINWRDGHDLRQD
 25 IIDYEIVQHI FEIFRLTVQMRNSLSELED RDYDRLISPVLNENNI FYDSAKAGDALPKDADANG
 AYCIALKGLYEIKQITENWKEDGKFSRDKL KISNKDWDFIQNKRYL

[0212] In certain embodiments, the type V-A Cas nuclease comprises MAD2 or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 2. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 2

MAD2 (SEQ ID NO: 2)

MSSLTKFTNKYSKQLTIKNELIPVGKTL ENIKENGLIDGDEQLNENYQKAKIIVDDFLRDFINK
 ALNNTQIGNWRELADALNKEDEDNIEKLQDKIRGIIVSKFETFDLFSSYSIKKDEKIIDDDNDV
 EEEELDLGKKTSSFYIFKKNL FKLVLPSYLKTTNQDKLKIISFDNFSTYFRGFFENRKNIFT
 KKPISTSIAYRIVHDNFPKFLDNIRC FNVWQTECPQLIVKADNYLKS KNVIAKDKSLANYFTVG
 AYDYFLSQNGIDFYNNIIGGLPAFAGHEKIQGLNEFINQECQKDSSELKSKLKNRHAFKMAVLFK
 35 QILSDREKSFVIDEFESDAQVIDAVKNFYAEQCKDNNVIFNLNLINKNIAFLSDDEL DGI FIEG
 KYLSSVSQKLYSDWSKLRNDIEDSANSKQGNKELAKKIKTNKGDVEKAISKYEFSLSELNSIVH
 DNTKFSDLLSCTLHKVASEKLKVN EGDWPKHLKNNEEKQKIKEPLDALLEIYNTLLIFNCKSF
 NKNGNFYVDYDRCINELSSVVYLYNKTRNYCTKPYNTDKFKLNFNSPQLGEGFSKSKENDCLT
 40 LLFKKDDNYYVGIIRKGAKINFDDTQAIADNTDNCIFKMNYFLLKDAKKFIPKCSIQLKEVKAH
 FKKSEDDYILSDKEKFASPLVIKKSTFLLATAHVKGKKNIKKFQKEYSKENPTEYRNSLNEWI
 45

AFCKEFLKTYKAATIFDITTLKKAEEYADIVEFYKDVDNLCYKLEFCPIKTSFIENLIDNGDLY
 LFRINNKDFSSKSTGTKNLHTLYLQAI FDERNLNNPTIMLNGGAELFYRKESIEQKNRITHKAG
 SILVNKVKCDGTSDDKIRNEIYQYENKFIDTLSDEAKKVLPNVIKKEATHDITKDKRFTSDKF
 FFHCPLTINYKEGDTKQFNNEVLSFLRGNPDINIIGIDRGERNLIYVTVINQKGEILDSVSFNT
 5 VTNKSSKIEQTVDYEEKLAVREKERIEAKRSWDSISKIATLKEGYLSAIVHEICLLMIKHNAIV
 VLENLNAGFKRIRGGLSEKSVYQKFEKMLINKLNYFVSKKESDWNKPSGLNLGLQLSDQFESFE
 KLGIQSGFI FYVPAAYTSKIDPTTGFANVLNLSKVRNVDAIKSFFSNFNEISYSKKEALFKFSF
 DLDSLKKGFSFVKFSKSKWNVYTFGERI IKPKNKQGYREDKRINLT FEMKKLLNEYKVSFDL
 ENNLIPNLTSANLKDTFWKELFFI FKTTLQLRNSVTNGKEDVLISPVKNAKGEFFVSGTHNKT
 10 PQDCDANGAYHIALKGLMILERNNLVREEKDTKKIMAI SNVDWFEYVQKRRGVL

[0213] In certain embodiments, the type V-A Cas nucleases comprises Csm1. Csm1
 proteins are known in the art and are described in U.S. Patent No. 9,896,696. Csm1 orthologs
 can be found in various bacterial and archaeal genomes. For example, in certain embodiments,
 15 the Csm1 protein is derived from *Smithella* sp. SCADC (Sm), *Sulfuricurvum* sp. (Ss), or
Microgenomates (*Roizmanbacteria*) *bacterium* (Mb).

[0214] In certain embodiments, the type V-A Cas nuclease comprises SmCsm1 or a
 variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid
 sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at
 20 least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at
 least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid
 sequence set forth in SEQ ID NO: 12. In certain embodiments, the type V-A Cas protein
 comprises the amino acid sequence set forth in SEQ ID NO: 12.

Smithella sp. SCADC Csm1 (SEQ ID NO: 12)

MEKYKITKTIRFKLLPDKIQDISRQVAVLQNSTNAEKKNNLLRLVQRGQELPKLLNEYIRYSDN
 HKLKSNTVTHFRWLRLFTKDLFYNWKKDNTTEKKIKISDVVYL SHVFEAFLEWESTIERVNADC
 NKPEESKTRDAEIALSIRKLGIKHQLPFIKGFVDNSNDKNS EDTKSKLTALLSEFEAVLKICEQ
 NYLPSQSSGIAIAKASFNYTINKKQKDFEAEIVALKKQLHARYGNKKYDQLLRENLNLIPLKEL
 PLKELPLIEFYSEIKKRKSTKKSEFLEAVSNGLVFDDLKSKFPLFQTESNKYDEYLLKLSNKITQ
 30 KSTAKSLLSKDSPEAQKLQTEITKLKKNRGEYFKKAFGKYVQLCELYKEIAGKRGKLGQIKGI
 ENERIDSQRLQYWALVLEDNLKHSLILIPKEKTNELYRKVWGAKDDGASSSSSTLYYFESMTY
 RALRKL CFGINGNTFLPEIQKELPQYNQKEFGFCFHKSNDDEIDEPKLISFYQSVLKTDFVK
 NTLALPQSVFNEVAIQSFETRQDFQIALEKCCYAKKQII SE SLKKEILENYNTQIFKITSLDLQ
 RSEQKNLKGHTRIWNRFWTKQNEEINYNLRLNPEIAIVWRKAKKTRIEKY GERSVLYEPEKRN
 35 YLHEQYTLCTTVTDNALNNEITFAFEDTKKKGTEIVKYNEKINQTLKKEFNKNQLWFY GIDAGE
 IELATLALMNKDKEPQLFTVYELKKLDFFKHGYIYNKERELVIREKPYKAIQNLSYFLNEELYE
 KTFRDGKFNETYNELFKEKHVSAIDLTTAKVINGKII LN GDMITFLNLRILHAQRKIYEELIEN
 PHAELKEKDYKLYFEIEGKDKDIYISRLDFEYIKPYQEISNYLFAYFASQQINEAREEEQINQT
 KRALAGNMIGVIYYLYQKYRGIISIEDLKQTKVESDRNKFEGNIERPLEWALYRKFFQOEGYVPP
 40 ISELIKLRELEKFPLKDVKQPKYENIQQFGI IKFVSPEETSTTCPKCLRRFKDYDKNKQEGFCK
 CQCGFDTRNDLKGFEGLNDPDKVA AFNIAKRGFEDLQKYK

[0215] In certain embodiments, the type V-A Cas nuclease comprises SsCsm1 or a
 variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid
 sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at

least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 13. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 13.

5 *Sulfuricurvum* sp. Csm1 (SEQ ID NO: 13)

MLHAFTNQYQLSKTLRFGATLKEDEKKCKSHEELKGFVDISYENMKSSATIAESLNENELVKKC
 ERCYSEIVKFHNAWEKIYYRTDQIAVYKDFYRQLSRKARFDAGKNSQLITLASLCGMYQGAKL
 SRYITNYWKDNI TRQKSFLKDFSQQLHQYTRALEKSDKAHTKPNLINFNKTFMVLANLVNEIVI
 PLSNGAISFPNISKLEDGEESHLEFALNDYSQSELIGELKDAIATNGGYTPFAKVTLNHYTA
 10 EQKPHVFKNDIDAKIRELKLIGLVETLKGKSSEQIEEYFSNLDKFSTYNDRNQSVIVRTQCFKY
 KPIPFVLVHQLAKYISEPNGWDEDAVAKVLDAVGAI RSPAHDYANNQEGFDLNHYPIKVAFDYA
 WEQLANSLYTTVTFPQEMCEKYLNSIYGCEVSKEPVFKFYADLLYIRKNLAVLEHKNNLPSNQE
 EFICKINNTFENIVLPYKISQFETYKKDILAWINDGHDHKKYTDKQQLGFI RGGLKGRIKAE
 VSQKDKYGKIKSYENPYTKLTNEFKQISSYTGKTFALRDKFKEKNEITKITHFGIIIEDKNR
 15 DRYLLASELKHEQINHVSTILNKLDKSSEFITYQVKSLSKTLIKLIKHNHTTKGAI SPYADFH
 TSKTGFNKNEIEKNWDNYKREQVLVEYVKDCLTDSTMAKNQNWAEFGWNFEKNSYEDIEHEID
 QKSYLLQSDTISKQSIASLVEGGCLLLPIINQDITSKERKDNQFSKDNWHI FEKSKEFRLHPE
 FAVSYRTPIEGYPVQKRYGRLQFVCAFNHIVPQNGEFINLKKQIENFNDEDVQKRNVTETFNKK
 VNHALSDKEYVVIGIDRGLKQLATLCVLDKRGKILGDFEIKKEFVRAEKRSSEHWEHTQAETR
 20 HILDLSNLRVETTIEGKKVLVDQSLTLVKKNRDTPEEATEENKQKIKLKQLSYIRKLQHKMQT
 NEQDVLDLINNEPSDEEFKKRIEGLISSFGEGQKYADLPINTMREMISDLQGV IARGNNQTEKN
 KIIELDAADNLKQGIVANMIGIVNYIFAKYSYKAYISLEDLSRAYGGAKSGYDGRYLPSTSQDE
 DVDFKEQQNQMLAGLTGYQFFEMQLLKKLQKI QSDNTVLRFPVPAFRSADNYRNILRLEETKYKS
 KPFGVVHFIDPKFTSKKCPVCSKTNVYRDKDDILVCKECCGFRSDSQLKERENNIHYIHNGDDNG
 25 AYHIALKSVENLIQMK

[0216] In certain embodiments, the type V-A Cas nuclease comprises MbCsm1 or a variant thereof. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the type V-A Cas protein comprises the amino acid sequence set forth in SEQ ID NO: 14.

Microgenomates (Roizmanbacteria) bacterium Csm1 (SEQ ID NO: 14)

MEIQELKNLYEVKKTVR FELKPSKKKIFEGGDVIKLQKDFEKVQKFFLDI FVYKNEHTKLEFKK
 35 KREIKYTWLRTNTKNEFYNNWRGKSDTGKNYALNKIGFLAEELRWLNNEWQELTKSLKDLTQREE
 HKQERKSDIAFVLRNFLKRQNLPIKDFFNVIDIQGKQGESDDKIRKFREEIKEIEKNLNAC
 SREYLPTQSGVLLYKASFYTLNKTTPKEYEDLKKEKESELSSVLLKEIYRRKRFNRTTNQKD
 TLFECTSDWLVKIKLGKDIYEWTLDEAYQMKIWKANQKSNFIEAVAGDKLTHQNFRKQFPLFD
 ASDEDFETFYRLTKALDKNPENAKKIAQKRKGFNAPNETVQTKNYHELCELYKRIAVKRGKII
 40 AEIKGIENEEVQSQLLTHWAVIAEERDKKFIVLIPRKNNGGKLENHKNHAHFLQEKDRKEPNDIK
 VYHFKSLTLRSLEKLCFKEAKNTFAPEIKKETNPKIWFPTYKQEWNSTPERLIKFKYQVLQSNY
 AQTYLDLVDFGNLNTFLETHFTTLEEFESDLEKTCYTKVPVYFAKKELETFADEFEAEVFEITT
 RSISTESKRKENAHAEIWRDFWSRENEEENHITRLNPEVSVLYRDEIKEKSNTSRKNRKSANN
 RFSDPRFTLATITL NADKKKSNLAFKTVEDINIHIDNFNKKFSKNFSGEWVYGIDRGLKELAT
 45 LNVVKFSDVKNVFGVSQPKFAKIPYIKLRDEKAILKDENGLSLKNAKGEARKVIDNISDVLEE
 GKEPDSTLFEKREVSSIDLTRAKLIKGHII SNQGDKTYLKLKETSARKRRI FELFSTAKIDKSSQ

FHVRKTIELSGTKIYWLCEWQRQDSWRTEKVSRLRNTLKGYLQNLDLKNRFENIETIEKINHRLD
 AITANMVGILSHLQNKLEMQGVIALENLDTVREQSNNKMIDEHFEQSNEHVSRRLEWALYCKFA
 NTGEVPPQIKESI FLRDEFKVCQIGILNFIDVKGTSSNCPNCDQESRKTGSHFICNFQNNCIF
 SKENRNLLLEQNLHNSDDVAAFNI AKRGLEIVKV

- 5 **[0217]** In certain embodiments, the type V-A Cas nuclease comprises an ART nuclease or a variant thereof. In general, such nucleases sequences have < 60% AA sequence similarity to Cas12a, < 60% AA sequence similarity to a positive control nuclease, and > 80% query cover. In certain embodiments, the Type V-A nuclease comprises an ART1, ART2, ART3, ART4, ART5, ART6, ART7, ART8, ART9, ART10, ART11, ART12, ART13, ART14, ART15, ART16, ART17, ART18, ART19, ART20, ART21, ART22, ART23, ART24, ART25, ART26, ART27, ART28, ART28, ART30, ART31, ART32, ART33, ART34, ART35, or ART11* (*i.e.*, ART11_L679F, *i.e.*, ART11 wherein leucine (L) at amino acid position 679 is replaced with phenylalanine (F)) nuclease, as shown in **Table 25 and Appendix A**. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 15 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence designated for the individual ART nuclease as shown in **Table 25 and Appendix A**. In certain embodiments, provided is a nucleic acid-guided nuclease comprising a nucleic acid-guided nuclease polypeptide having at least 85% 20 identity to an amino acid sequence represented by SEQ ID NOs: 950-984 or a nucleic acid encoding a nucleic acid-guided nuclease polypeptide comprising at least 85% identity with the polynucleotide represented by SEQ ID NOs: 808-949. In certain embodiments, provided is a nucleic acid-guided nuclease comprising a polypeptide having at least 90% identity to the amino acid sequence represented by SEQ ID NOs: 950-958, 968-970, 972, 973, 976, 978-982, or 984, 25 wherein the polypeptide does not contain a peptide motif of **YLFQIYNKDF** (SEQ ID NO: 806). In certain embodiments, provided is a nucleic acid-guided nuclease comprising a nucleic acid encoding a polypeptide having at least 90% identity to nucleic acids represented by SEQ ID NOs: 808-845 wherein an encoded polypeptide does not contain a peptide motif of **YLFQIYNKDF** (SEQ ID NO: 806). In certain embodiments, provided is a nucleic acid-guided 30 nuclease wherein the polypeptide comprises at least 90% identity with the amino acid sequence represented by SEQ ID NOs: 950, 951, 954, 955, 957, or 958. In certain embodiments, provided is a nucleic acid-guided nuclease, wherein the polypeptide comprises a polypeptide comprising at least 90% identity with the amino acid sequence represented by SEQ ID NO: 951.

35 **Table 25 Exemplary ART nucleases**

ART Name	Protein Reference Number	SEQ ID NO corresponding to Amino Acid sequences	SEQ ID NO corresponding to nucleic acid sequence	% AA to Cpf1 (<80% desired)	% AA to positive control (<60% desired)
ART1	WP_118425113.1	950	808	30.838	32.54
ART2	WP_137013028.1	951	812	34.189	33.07
ART3	WP_073043853.1	952	818	35.982	36.72
ART4	WP_118734405.1	953	822	30.519	51.64
ART5	WP_146683785.1	954	826	30.114	32.31
ART6	WP_117882263.1	955	830	29.421	33.49
ART7	OYP43732.1	956	834	26.323	28.64
ART8	TSC78600.1	957	838	25.379	23.01
ART9	WP_094390816.1	958	842	26.323	28.62
ART10	WP_104505765.1	959	846	31.291	32.59
ART11	WP_151622887.1	960	850	30.654	35.55
ART12	HAW84277.1	961	854	34.872	31.33
ART13	WP_119227726.1	962	858	34.993	31.55
ART14	WP_118080156.1	963	862	32.551	35.33
ART15	WP_046700744.1	964	866	31.456	33.92
ART16	WP_115247861.1	965	870	31.136	34.25
ART17	WP_062499108.1	966	874	31.136	34.17
ART18	WP_154326953.1	967	878	31.113	33.28
ART19	WP_117747221.1	968	882	30.764	32.47
ART20	WP_118211091.1	969	886	30.986	32.29
ART21	WP_118163031.1	970	890	31.134	32.54
ART22	WP_115006085.1	971	894	30.044	31.55
ART23	HCS95801.1	972	898	30.37	51.64
ART24	WP_089541090.1	973	902	30.933	33.11

ART Name	Protein Reference Number	SEQ ID NO corresponding to Amino Acid sequences	SEQ ID NO corresponding to nucleic acid sequence	% AA to Cpf1 (<80% desired)	% AA to positive control (<60% desired)
ART25	WP_120123115.1	974	906	29.978	48.88
ART26	WP_117874294.1	975	910	29.904	48.49
ART27	WP_117951432.1	976	904	29.421	33.03
ART28	WP_108977930.1	977	918	32.099	32.69
ART29	WP_117886476.1	978	922	29.643	33.41
ART30	WP_101070975.1	979	926	29.027	32.95
ART31	WP_117949317.1	980	930	29.198	33.18
ART32	WP_118128310.1	981	934	29.198	33.18
ART33	WP_138157649.1	982	938	27.273	29.89
ART34	WP_135764749.1	983	942	27.004	25
ART35	OYP46450.1	984	946	26.709	29.51

[0218] In certain embodiments, the type V-A Cas nuclease comprises an ABW nuclease or a variant thereof. See International (PCT) Publication No. WO2021/108324. Exemplary amino acid and nucleic acid sequences are shown in **Table 26**. In certain embodiments, the Type V-A nuclease comprises an ABW1, ABW 2, ABW3, ABW4, ABW5, ABW6, ABW7, ABW8, or ABW 9 nuclease, as shown in **Table 26**. In certain embodiments, the type V-A Cas protein comprises an amino acid sequence at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to the amino acid sequence designated for the individual ABW nuclease as shown in **Table 26**.

Table 26 Sequences of exemplary engineered ABW nucleases

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
ABW1	MGHHHHHSSGLVPRGSG TMAAFDKFIHQYQVSKTL RFALIPQGKTLENTKNNV LQEDDERQKNYEKVKPIL	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGGCGGCGTTTCGAT AAGTTCATCCATCAATATCAAGTAAGCAAAACCCTC CGTTTTGCACTTATTCCGCAGGGGAAAACCTTGGAG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	DRIYKVF AEESLKDCSVD WNDLNACLDAYQKNPSAD KRQKV KAAQDALRDEIAG YFTGKQYANGKNKNAVKE KEQAELYKDI FSKKIFDG TVTNNKLPQVNL SAEETE LLGCFDKFTTYFVGFYQN RENVFSGEDIATAIPHRI VQDNF PKFRENCRIYQDL IKNEPAL KPLLQQA AAV MAQNP KGIYQPRKSLDDI FVIPFYNHLL LQDDIDYF NQILGGISGAAGQKKIQG LNETINLFMQQHPQEADK LKKKKIRHRFIPLYKQIL SDRTSFSFIPEAFSNSQE ALDGIETFKKSLKKNDF GALERLIQNLASLDLKYV YLSNKKVNEISQALYGEW HCIQDVLKQDFSLES LIQ INPQNSSNGFLATLTDEG KKRISQCRNV LGNPLPVK LADDQDKAQVKNQLDTLL AAVHYLEWFKADPDLETD PNFTVPFEKIWEELVPLL SLYSKVRNFVTKKPYSTA KFKLNFANPTLADGWDIH KESDNGALLFEKGGLYYL GIMNPKDKPNFKSYQGA PYYQKMVYRFFPDCSKTI PKCSTQRKDVKKYFEDHP QATSYQI HDSKKEKFRQD FFEIPREIYELNNTTYGT GKSKYKKFQTQYYQKTQD KSGYQKALRKWIDFSKKF LQTYVSTSI FDFKGLRPS KDYQDLGEFYKDVNSRCY RVTFEKIRVQDIHEAVKN GQLYLFQLYNKDFSPKSH GLPNLHTLYWKAVFDPEN LKDP IVKLNGQAE LFYRP KSNMQIIQHKTGEEIVNK KLKDGTPVPDDIYREISA YVQGK CQGNLSPEAEKWL PSVTIKKAAHDITKDRRF TEDKFF FHPITLNYQSS GKPTAFNSQVNDFLTEHP ETNIIGIDRGERNLIYAV VITPDGKILEQKSFNVIH DFDYHESLSQREKQRVAA	AATACAAAAAATAACGTACTCCAGGAAGATGATGAG CGTCAGAAAAATTACGAAAAAGTCAAACCTATCCTT GATCGTATTTATAAGGTATTCGCTGAGGAAAGCCTG AAAGATTGCAGCGTTGACTGGAATGACCTCAATGCA TGTCTGGATGCTTACCAAAAAAATCCTAGCGCGGAT AAGCGTCAGAAGGTGAAAGCCGCGCAGGACGCGTTG CGGGACGAAATTGCCGGTTATTTTACAGGGAAACAA TACGCGAACGGGAAGAACAAAAATGCCGTTAAGGAG AAAGAGCAGGCAGAATTGTATAAGGATATCTTTAGC AAAAAGATCTTTGATGGGACCGTAACGAACAACAAA TTGCCACAGGTCAACCTTTCAGCCGAAGAAACAGAG TTATTAGGCTGTTTTGATAAATTCACAACATATTTT GTCGGCTTTTACCAGAACCGTGAGAACGTATTTTCA GGGGAGGATATTGCTACAGCTATTCCGCATCGGATC GTCCAGGATAATTTTCTAAATTCCGGGAAAACGTGT CGGATTTATCAGGACTTAATCAAAAATGAACCTGCC CTTAAACCGCTGCTTCAGCAAGCAGCGGCCGCGGTG ATGGCCCAAGATCCAAAGGGGATCTATCAACCACGT AAGAGTCTGGACGATATTTTGTTCATTCCGTTTTAT AACCATCTCCTCTTACAGGATGATATTGATTATTTT AATCAAATCTTAGGCGGCATTTCCGGGGCAGCCGGT CAGAAAAAATCCAGGGTTTAAATGAAACAATTAAT CTGTTTATGCAACAGCACCCACAAGAAGCCGATAAG TTAAAGAAAAAAGATTTCGTCATCGGTTTTATTCCG CTGTATAAACAAATTCTCTCTGACCGTACGTCTTTT TCGTTTCATCCCTGAAGCTTTTTTCCAATTCTCAGGAA GCGTTAGACGGCATTGAGACATTCAAAAAGTCTCTT AAGAAGAATGACACATTCGGCGCGTTGGAGCGGCTG ATTCAAAAATCTTGCTTCCCTGGACCTGAAATACGTG TATTTATCGAACAAGAAGGTCAATGAGATTTCGCAG GCATTATACGGCGAATGGCACTGCATCCAAGACGTC CTCAAGCAAGATTTTACGCTTGAGAGCCTGATCCAG ATCAACCCACAAAATTTAGCAATGGTTTCTTGCC ACACTTACCGACGAAGGCAAGAAACGTATCTCCCAA TGTCGTAACGTACTGGGGAATCCTCTTCCAGTCAAG CTTGCGGATGATCAAGACAAAGCGCAAGTCAAAAAC CAATTGGATACATTACTGGCTGCTGTACACTATCTC GAGTGGTTCAAGGCAGATCCAGACCTGGAAACAGAC CCTAACTTCACTGTTCTTTTCGAAAAGATCTGGGAG GAATTGGTTCTTTTACTTTCACTGTACTCTAAAGTT CGGAATTTTGTTACAAAGAAGCCATATTCTACAGCT AAATTTAAACTGAACCTTGCTAACCCGACATTAGCG GATGGGTGGGATATTCACAAGGAAAGTGATAACGGC GCGCTCCTGTTTGAAAAGGGTGGTTTGTATTACTTG GGTATCATGAACCTTAAAGATAAGCCTAATTTTAAA TCCTATCAGGGTGCAGAGCCATACTATCAGAAGATG GTGTACCGTTTTTTTCTGACTGTTTCAAGACCATC CCAAAATGCAGCACCCAACGTAAGGATGTAAAAAAG TACTTCGAAGACCACCTCAAGCGACCTCATACCAG ATCCACGACTCAAAGAAAGAGAAGTTTCGTCAGGAT

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	RQAWTAIGRIKDLKEGYL SLVVHEIAQMMIKYQAVV VLENLNTGFKRVRGGISE KAVYQQFEKMLIEKLNFL VFKDRAINQEGGVLKAYQ LTDSFTSFAKLGNSGFL FYIPSAYTSKIDPGTGFV DPFIWSHVTASEENRNEF LKGFDSLKYDAQSSAFVL HFKMKSNNKQFQKNNVEGF MPEWDICFEKNEEKISLQ GSKYTAGKRIIFDSKKKQ YMECFPQNELMKALQDVG ITWNTGNDIWQDVLKQAS TDTGFRHRMINLIRSVLQ MRSSNGATGEDYINSPVM DLDGRFFDTRAGIRDPL DADANGAYHIALKGRMVL ERIRSQKNTAIKNTDWLY AIQEERNGAPKRPAATKK AGQAKKKKASGSGAGSPK KKRKVEDPKKKRKV (SEQ ID NO: 789)	TTTTTTGAGATCCCTCGGGAGATTTACGAGCTTAAT AACACCACATACGGCACAGGTAAGTCTAAATATAAA AAATTCCAGACCCAGTATTACCAGAAGACTCAGGAT AAGTCAGGCTATCAGAAAGCACTTCGCAAATGGATT GACTTTTCCAAAAAGTTTCTTCAAACATACGTCAGT ACTTCCATTTTTGATTTCAAAGGTCTCCGTCCTTCG AAGGATTATCAGGACTTAGGCGAGTTCTATAAAGAC GTTAATTCGCGTTGTTACCGTGTGACGTTTCGAGAAA ATTTCGCGTACAGGACATCCACGAAGCAGTCAAAAAAT GGGCAACTGTATCTCTTCCAATTATATAATAAGGAC TTCTCACCTAAAAGCCATGGGTTGCCTAATCTTCAC ACTCTCTATTGGAAAGCCGTGTTTCGATCCTGAGAAC TTGAAGGACCCTATCGTAAACTTAATGGCCAAGCT GAGTTATTCTATCGGCCGAAATCCAACATGCAAATC ATCCAACATAAGACCGGGGAGGAGATTGTGAACAAA AAGCTGAAGGACGGCACCCCGGTTCTGATGATATC TACCGCGAAATCAGTGCTTACGTCCAGGGGAAATGT CAAGGCAACTTATCCCCGGAGGCAGAGAAGTGGCTC CCAAGTGTCACAATCAAGAAAGCCGCCCATGATATC ACAAAGGATCGTCGCTTTACCGAAGATAAGTTTTTC TTTCATGTCCCTATTACACTGAACTATCAGAGTTCA GGCAAGCCGACGGCATTCAACTCGCAAGTAAACGAT TTCTTGACCGAGCACCCCTGAGACAAATATCATCGGC ATTGATCGGGGTGAACGTAACCTTGATTTATGCCGTT GTAATCACTCCAGATGGCAAGATTCTCGAACAGAAA TCTTTTAACGTGATCCACGACTTTGATTATCATGAA TCCCTGTCCCAGCGGGAAAAACAGCGGGTAGCAGCG CGTCAGGCTTGGACAGCGATTGGTCGCATCAAGGAT CTCAAGGAAGGTTACCTGTGCTTGTGGTGCACGAA ATTGCTCAAATGATGATCAAATACCAAGCAGTCGTC GTATTAGAAAACCTCAACACGGGCTTTAAGCGTGTG CGCGGTGGTATCAGTGAGAAGGCCGTCTACCAACAG TTCGAAAAAATGTTGATTGAAAAATTGAACTTCCTG GTATTTAAAGATCGGGCAATCAATCAGGAAGGCGGG GTTCTCAAAGCTTACCAGCTGACAGACTCGTTTACG TCTTTTGCAAAGTTAGGTAACAGTCCGGTTTCCTG TTCTACATCCCGTCCGCCTACACCAGCAAAATCGAC CCTGGTACGGGCTTCGTCGATCCTTTTATCTGGTCT CACGTGACCGCTTCTGAGGAAAATCGGAATGAATTT TTAAAGGGCTTTGATAGCTTGAAATATGACGCCCAA TCATCCGCCTTTGTACTGCATTTCAAGATGAAATCC AATAAGCAATTTTCAGAAGAACAATGTTGAAGGTTTC ATGCCGGAATGGGATATCTGCTTCGAGAAAAACGAG GAAAAGATTTCTTGCAGGGTAGTAAGTATACAGCC GGTAAACGCATTATTTTCGACTCCAAAAAGAAGCAA TACATGGAGTGCTTCCCGCAGAATGAGCTCATGAAA GCACTGCAGGACGTAGGCATCACCTGGAACACGGGC AACGATATCTGGCAGGATGTCCTTAAACAAGCGAGC ACAGATACAGGGTTTCGTCACCGGATGATCAACCTG ATCCGTTTCAGTGCTCCAGATGCGGTCCAGTAATGGT

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		GCGACCGGGGAGGATTACATCAATTCACCTGTGATG GATCTGGACGGCCGTTTTTTCGACACTCGGGCGGGG ATTCTGTGATCTGCCATTGGATGCCGACGCCAACGGC GCATACCACATCGCTTTAAAAGGGCGTATGGTACTC GAACGCATTTCGCTCCCAAAAGAATACCGCGATTAAG AACACTGACTGGTTATACGCAATCCAAGAGGAACGT AACGGCGCGCCAAAAAGGCCGGCGGCCACGAAAAAG GCCGGCCAGGCAAAAAAGAAAAGGCTAGCGGCAGC GGCGCCGGATCCCCAAAAGAAAAGGAAGGTTGAA GACCCCAAGAAAAGAGGAAGGTGTGATAA (SEQ ID NO: 790)
ABW2	MGHHHHHHSSGLVPRGSG TMKEFTNQYSLTKTLRFE LRPVGETAEKIEDFKSGG LKQTVKDRERTEAYKQL KEVIDSYHRDFIEQAFAR QQTLSSEDFKQTYQLYKE AQKEKDGETLTKQYEHLR KKIAAMFSKATKEWAVMG ENNELIGKNKESKLYQWL EKNYRAGRIEKEEFDHNA GLIEYFEKFSTYFVGFDK NRANMYSKEAKATAISFR TINENMVKHFDNCQRLEK IKSKYPDLAEELKDFEEF FKPSYFINCMNQSGIDYY NISAIGGKDEKDQKANMK INLFTQKNHLKGS DKPPF FAKLYKQILSDREKSVVI DEFEKDSELTEALKNVFS KDGLINEEFFTKLKSAL NFMLPEYQGQLYIRNAFL TKISANIWGSWSGIIKD AVTQAAENNFTRKSDKEK YAKKDFYSIAELQQAIDE YIPTLENGVQNASLIEYF RKMNYKPRGSEEDAGLIE EINNLRQAGIVLNQAEL GSGKQREENIEKIKNLLD SVLNLERFLKPLYLEKEK MRPKAANLNKDFCESFDP LYEKLKTFKLYNKVRNY ATKKPYSKDKFKINFDTA TLLYGWSLDKETANLSVI FRKREKFYLGIIINRYNSQ IFNYKIAGSESEKGLERK RSLQQKVLAEEGEDYFEK MVYHLLLGASKTIPKCST QLKEVKAHFQKSSEDYII QSKSFAKSLTLTKEIFDL	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGAAGGAGTTTACC AACCAATATTCCTTAACCAAGACCCTGCGGTTTCGAG TTGCGGCCAGTCGGCGAAACAGCAGAAAAGATCGAA GATTTTAAATCGGGCGGGCTCAAGCAAACAGTGGAA AAGGATCGTGAGCGTACAGAAGCGTATAAGCAGTTG AAAGAGGTTATTGACTCCTATCATCGTGACTTCATT GAGCAAGCTTTTGC GCGCCAGCAGACGCTGTCCGAG GAGGATTTTAAACAAACATATCAACTGTACAAAGAG GCCCAGAAAAGAGAAGGATGGGGAAACATTAACAAAG CAGTACGAGCATTTACGGAAGAAAATCGCAGCTATG TTCAGCAAGGCTACGAAGGAATGGGCCGTTATGGGG GAGAATAACGAATTGATCGGGAAAAACAAAGAGTCA AAGTTGTATCAGTGGCTGGAGAAGAACTACCGCGCA GGTCGCATCGAAAAAGAGGAATTTCGACCATAATGCG GGCTTAATCGAATACTTCGAGAAATTTTCCACATAT TTCGTAGGTTTTGACAAAAATCGTGCGAATATGTAT TCAAAGGAGGCAAAGGCGACCGCAATTTCTTCCGG ACGATTAATGAGAACATGGTCAAGCATTTTCGATAAT TGCCAGCGGCTCGAGAAGATTAAATCTAAATATCCT GATTTGGCCGAGGAGCTGAAGGATTTTGAGGAGTTT TTTAAACCTAGCTATTTTATTAATTGTATGAATCAA TCGGGTATCGACTACTACAATATCAGCGCGATCGGC GGTAAGGATGAAAAGGATCAGAAAGCGAATATGAAG ATCAACCTTTTCACGCAAAAAAATCATTTAAAGGGC AGTGATAAACCACCATTTTTTGCTAAGCTCTACAAG CAAATTTTGAGTGACCGGGAGAGTCCGTGGTAATC GACGAGTTCGAAAAGGACAGCGAATTGACAGAGGCA CTCAAAAACGTGTTTTCCAAGGACGGTTTGATCAAT GAGGAGTTTTTTACAAAGTTAAAAAGTGCATTAGAA AATTTTATGTTGCCTGAATATCAAGGTCAACTCTAC ATCCGTAACGCTTTTCTTACGAAGATCAGCGCAAAC ATTTGGGGCTCTGGTTCTTGGGGCATCATCAAGGAC GCAGTTACCCAGGCTGCGGAAAACAATTTACGCGT AAGTCTGACAAGGAAAAGTATGCCAAGAAAGACTTC TATTCCATTGCTGAACTCCAGCAGGCTATTGATGAA TACATTCTACTCTGGAGAACGGGGTTCAAACGCA TCACTCATCGAGTACTTTTCGCAAAATGAATTACAAA CCACGCGGTTCTGAAGAAGACGCAGGCTTGATCGAA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	NNLRYNTETGEISSELS TYPKKFQKGYLQTGDVS GYKTALHKWIDFCKEFLR CYRNTEIFTFHFKDTKEY ESLDEFLKEVDSSGYEIS FDKIKASYINEKVNAGEL YLFEIYNKDFSEYSKGP NLHTIYWKSLEFETQNLDD KTAKLNGKAEIFFRPRSI KHNDKI IHRAGETLKNKN PLNEKPSSRFDYDITKDR RFTKDKFFLHCPITLNF QDKPVRFNEQVNLKDN PDVNIIGIDRGERHLLYY TLINQNGEILQQGSLNRI GEEESRPTDYHRLDERE KQRQQARETWKAVEGIKD LKAGYLSRVVHKLGLMV QNNAIIVLEDLNKGFKRG RFAVEKQVYQNFKALIQ KLNLYLVFKEVNSKDAPGH YLKAYQLTAPFISFEKLG TQSGFLFYVRAWNTSKID PATGFTDQIKPKYKNQKQ AKDFMSSFDVRYNRKEN YFEFEADFEKLAQKPKGR TRWTICSYGQERYSYSPK ERKFVKHNVTONLAELFN SEGISFDSGQCFKDEILK VEDASFFKSII FNLRLLL KLRHTCKNAEIERDFIIS PVKGNNSSFFDSRIAEQE NITSIPQNADANGAYNIA LKGLMNLHNI SKDGKAKL IKDEDWIEFVQKRKFAAA KRPAATKKAGQAKKKKAS GSGAGSPKKKRVKVEDPKK KRKV (SEQ ID NO: 16)	GAAATTAATAACAACCTGCGTCAGGCTGGGATCGTC CTGAATCAAGCCGAGCTGGGGTCTGGTAAGCAGCGG GAAGAGAATATTGAAAAATTAAGAACTTATTAGAT TCGGTTTTGAATCTCGAACGTTTCTTAAAGCCACTT TACTTGGAGAAAGAGAAAAATGCGTCCAAAAGCTGCT AACCTGAATAAGGATTTTTGTGAGTCATTTGATCCA CTTTACGAGAAACTGAAAACGTTTTTCAAGCTCTAC AATAAAGTACGTAACACGCAACAAAAGAACCATAC TCAAAGGACAAATTTAAGATCAATTTTGATACCGCT ACGTTATTATATGGGTGGAGTTTGGATAAGGAAACC GCGAATCTCAGCGTCATTTTCCGTAAACGCGAAAAA TTCTATTTGGGTATCATCAACCGGTACAATAGCCAG ATTTTCAATTATAAGATTGCGGGCAGTGAGAGCGAG AAAGGGTTAGAGCGTAAGCGGTCTGCTGCAGCAAAAG GTGCTTGCAGAGGAGGGTGAAGATTATTTTGAGAAA ATGGTATACCACCTGCTGCTTGGCGCGTCGAAAACCT ATTCCGAAATGCTCGACACAGTTGAAAGAAGTAAAA GCACACTTTCAAAAGTCATCAGAAGATTATATTATC CAATCCAAATCATTTGCAAAGTCATTAACATTAACA AAAGAGATCTTTGACTTAAATAATCTGCGGTATAAC ACAGAAACGGGCGAAATTAGTTCCGAGCTTTCTGAT ACATATCCGAAGAAGTCCAGAAGGGGTATCTCACA CAAACAGGCGACGTTTTCGGGTACAAAACCTGCTCTG CATAAGTGGATTGATTTCTGCAAAGAGTTCTTGCGT TGCTATCGTAATACGGAGATCTTCACGTTCCATTTT AAGGACACGAAGGAGTACGAGTCGTTAGATGAGTTC TTGAAAGAAGTGGATAGTTTACGTTATGAGATTTCA TTCGATAAGATCAAAGCCTCTTATATCAACGAGAAG GTTAATGCAGGCGAGCTGTACTTGTTCGAGATCTAT AATAAAGATTTCTCCGAGTATTCCAAAGGTAAGCCA AATCTGCATACCATTTATTGGAAAAGTCTCTTCGAG ACTCAAAACTTGCTGGATAAAACAGCGAAACTCAAC GGCAAGGCAGAGATCTTCTTCCGGCCACGTTTCGATC AAACACAACGACAAAATCATCCACCGTGCGGGCGAA ACACTTAAGAATAAAAAACCGCTCAATGAAAAGCCT AGTTCGCGTTTCGATTACGATATTACGAAAGATCGT CGTTTTACGAAAGACAAATTTTTTTTACACTGCCCT ATTACGTTAAACTTTAAGCAGGACAAAGCCTGTTTCGC TTTAATGAACAAGTCAACTTATACTTAAAAGACAAT CCAGACGTGAATATTATCGGTATCGATCGTGGTGAG CGTCACTTGCTTTATTACACTTTGATCAATCAGAAT GGTGAGATCTTACAGCAGGGTTCACTTAATCGCATT GGTGAGGAAGAATCTCGGCCTACGGACTACCATCGG TTACTCGATGAGCGTGAAAAGCAGCGTCAACAAGCA CGGGAGACGTGGAAAGCAGTAGAAGGGATTAAAGGAC TTAAAAGCTGGGTATCTTTCACGGGTGTACATAAA CTTGCAAGTTTAAATGGTACAAAACAACGCAATTGTC GTTCTGGAAGATCTTAACAAGGGTTTTAAGCGCGGT CGTTTCGCTGTTGAGAAACAGGTGTACCAGAACTTC GAAAAGCACTTATTCAAAGCTTAACTATTTAGTG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		TTCAAGGAGGTCAACTCTAAAGACGCCCCTGGCCAC TATTTGAAGGCATATCAGCTTACGGCCCCTTTCATC TCGTTTCGAAAAATTGGGTACTCAGAGCGGTTTCCTT TTTTATGTGCGCGCATGGAATACCTCGAAGATCGAC CCGGCGACGGGTTTTACCGACCAAATCAAACCAAAG TATAAAAACCAAAAACAAGCTAAAGACTTCATGTCA AGCTTCGACTCTGTCCGGTACAACCGCAAGGAAAAAT TATTTTGAATTCGAGGCGGACTTTGAAAACTGGCA CAGAAACCTAAGGGGCGCACCCGCTGGACGATTTGT TCCTATGGCCAGGAACGGTACTCTTACTCCCCAAAA GAACGGAAGTTTGTAAAGCACAAACGTTACACAAAAAT CTTGCTGAGCTTTTTAATTCAGAGGGTATCTCGTTC GACTCCGGGCAGTGTTCAGGATGAGATCCTGAAG GTCGAGGATGCCAGTTTCTTTAAGTCTATTATTTTC AATCTTCGCCTCCTTCTCAAGCTTCGTCACACTTGC AAGAACGCCGAGATCGAACGTGATTTTCATCATTTCT CCTGTCAAGGGGAACAATTCGTCCTTTTTTGGACTCC CGTATTGCCGAACAAGAAAATATCACCAGCATTCCA CAGAATGCTGATGCAAACGGTGCATACAACATCGCG CTGAAGGGCCTGATGAACCTCCATAATATCTCTAAG GACGGCAAGGCAAAATTAATTAAGGATGAAGATTGG ATCGAATTTGTCCAAAAACGCAAGTTCGCGGCCGCA AAAAGGCCGGCGGCCACGAAAAAGGCCGGCCAGGCA AAAAAGAAAAAGGCTAGCGGCAGCGGCGCCGATCC CCAAAGAAGAAAAGGAAGGTTGAAGACCCCAAGAAA AAGAGGAAGGTGTGATAA (SEQ ID NO: 17)
ABW3	MGHHHHHHSSGLVPRGSL QMKTLSDFTNLFPLSKTL RFKLIPIGNTLKNIEASG ILDEDRHRAESYVKVKAI IDEYHKAIFIDRVLSDTCL QTESIGKHNSLEEFFFY QIGAKSEQQKTFKKIQD ALRKQIADSLTKDKHFSR IDKKELIQEDLIQFVRDG EDAAEKTSLISEFQNFV YFTGFHENRQNMYSPEDEK STAIAYRLINENLPKFVD NMKVFDRIAASELASCFD ELYHNFEEYLQVERLHDI FSLDYFNLLLTQKHIDVY NALIGGKATETGEKIKGL NEYINLYNQRHQEKLPK FKMLFKQILTDREAI SWL PRQFDDNSQLLSAIEQCY NHLSTYTLKDGSLKYLLE NLHTYDTEKIFIRNDSL TEISQRHYGSWSILPEAI KRHLERANPQKRRETYEA YQSRIEKAFKAYPGFSIA	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCCTGCAGATGAAGACCTTGTCT GATTTTACCAATCTGTTCCCTTTATCTAAGACTCTC CGTTTCAAGCTGATTCCAATCGGCAACACGCTCAAG AACATTGAAGCTAGTGGCATCCTTGACGAGGATCGC CACCGCGCGGAGTCCTATGTCAAGGTCAAGGCCATC ATCGACGAATATCATAAAGCTTTCATCGATCGGGTC CTGTGCGGATACTTGCCTCCAGACGGAATCTATCGGC AAACACAACAGTCTCGAGGAATTCTTTTTCTACTAC CAAATTGGTGCAAAAAGTGAACAGCAGAAAAAGACG TTTAAAAAGATTCAAGACGCCTTGCGCAAACAAATC GCAGATAGCCTCACCAAGGACAAACATTTTTTCACGG ATTGATAAAAAAGAATTGATCCAAGAGGATTTGATC CAGTTTGTGCGCGATGGGGAGGATGCCGCTGAAAAG ACGTCTCTGATTTCCGAATTTCAAAATTTACAGTT TATTTTACCGGGTTTCATGAGAATCGCCAGAACATG TACAGTCCGGACGAGAAGTCCACGGCCATCGCATAT CGCTTAATTAACGAGAATCTCCCAAAATTCGTAGAC AACATGAAAGTTTTTGACCGTATCGCGGCGTCCGAA TTGGCATCGTGTTCGACGAATTATACCACAACCTTC GAGGAATACCTCCAAGTGGAGCGGTTACATGATATC TTTAGTTTGGACTATTTCAATCTGCTTCTCACGCAG AAACATATCGACGTCTATAATGCTCTGATCGGTGGG AAGGCAACCGAAACCGGGGAAAAGATCAAGGGCTTA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	FLNGCLTETGKESPSIES YFESLGAVETETSQQENW FARIANAYTDFREMQRNL HATDVPLAQDAEAVARIK KLLDALKGLQLFIKPLLD TGEAEKDERFYGDFTEF WNELDTITPLYNMVRNYL TRKPYSEEKIKLNFQNP LLNGWDLNKEVDNTSVIL RRNGRYYLAIMHRNHRRV FSQYPGTERGDCYEKMEY KLLPGANKMLPKVFFSKS RIDEFNPSEELLARYQQG THKKGENFNLHDCHALID FFKDSIEKHEEWRNFHFK FSDTSSYTDMSGFYREIE TQGYKLSFVPVACEYIDE LVRDGKIFLFQIYNKDFS TYSKGKPNMHTLYWEMLF DERNLMNVVYKLNQAEI FFRKASLSARHPEHPAGL PIKKKQAPTEESCFPYDL IKNKRYTVDQFQFHVPIIT INFKATGTSNINPSVTDY IRTADDLHIIGIDRGERH LLYLVVIDSQGRICEQFS LNEIVTQYQGHQYRTDYH ALLQKKEDERQKARQSWQ SIENIKELKEGYLSQVVH KVSELMIKYKAIVVLEDL NAGFKRSRQKVEKQVYQK FEKMLIDKLNILVFKTAE ADQPGLLHAYQLTNKFE SFKKMGKQSGFLFYIPAW NTSKIDPTTGfVNLFDTR YENVDKSRAFFGKFDSIR YRADKGTFEWTFDYNFHH KKAEGTRSSWCLSSHGHR VRTFRNPAKNNQWDNEEI DLTQAFRDLFEAWGIEIT SNLKEAICNQSEKKFFSE LFELFKLMIQLRNSVTGT NIDYMVSPVENHYGTFFD SRTCDSSLPANADANGAY NIARKGLMLARRIQATPE NDPISLTLNKEWLRFAQ GLDETTTYEAAKRPAAT KKAGQAKKKKASGSGAGS PKKKRKVEDPKKKRKV (SEQ ID NO: 29)	AATGAATACATCAATCTCTACAATCAACGTCACAAG CAGGAAAAACTGCCAAAATTCAAGATGTTATTCAAG CAAATTCTTACCGACCGTGAGGCAATCAGCTGGTTG CCACGCCAATTTGACGATAATAGTCAGTTACTCTCA GCCATTGAACAGTGTTATAACCACTTTTCGACCTAC AACTCAAGGATGGGTCACTCAAATACCTGTTAGAA AACCTGCATACATACGATACTGAAAAGATCTTCATC CGCAATGACAGTTTACTTACGGAAATCTCCCAACGG CATTACGGTTCGTGGTCGATTTTACCAGAAGCTATC AAACGTCATCTCGAGCGCGCAACCCGCAAAAACGG CGCGAAACATACGAGGCCTATCAATCTCGCATTGAG AAGGCCTTTAAGGCATATCCGGGGTTTTCAATTGCT TTCCTCAATGGGTGTTTAACAGAGACAGGTAAGGAG TCGCCATCCATCGAAAAGCTATTTTGAAAGTCTGGGT GCTGTCGAAACAGAGACCTCTCAGCAGGAAAACCTGG TTTGCCCCGCATCGAAACGCTTATACGGACTTTTCGT GAAATGCAAAATCGGCTGCACGCCACTGACGTGCCG TTGGCTCAAGACGCTGAGGCAGTGGCCCGGATCAAG AAGCTGTTAGATGCACTGAAAGGCCTGCAATTATTC ATTAAGCCTCTTTTGGTACTGGCGAAGAAGCAGAG AAAGATGAACGGTTCTATGGGGACTTTACCGAATTC TGGAACGAGTTAGACACTATCACGCCATTGTACAAT ATGGTACGGAACCTATCTCACGCGTAAGCCTTATAGT GAAGAAAAAATCAAGCTCAATTTCCAGAATCCGACA TTACTGAACGGTTGGGATTTGAACAAAGAGGTAGAT AATACATCTGTCATCCTCCGCCGAATGGTCGTTAT TATCTTGCCATCATGCACCGCAACCACCGGCGTGTA TTTTCACAGTATCCAGGCACAGAACGTGGCGATTGT TATGAGAAAATGGAATATAAACTGCTTCCGGGCGCC AACAAGATGCTCCCAAAAGTCTTCTCTCTAAATCA CGCATCGATGAATTCAACCCTAGCGAAGAATTATTA GCACGTTACCAGCAAGGTACCCACAAGAAGGGTGAG AATTTTAATTTACACGACTGCCATGCCTTGATTGAT TTTTTTAAAGACTCTATTGAGAAACATGAAGAATGG CGTAACTTTTCAATTTTAAATTTAGTGATACGTCCAGT TACACCGACATGAGCGGCTTTTATCGTGAAATCGAA ACACAGGGTTACAAGTTGTCATTTGTGCCAGTGGCG TGTGAATACATCGATGAGTTGGTACGTGATGGCAAA ATCTTTTTTGTTCAGATCTATAATAAGGACTTTTTCG ACCTACTCTAAGGGCAAGCCAAATATGCACACTCTT TATTGGGAAATGCTTTTTCGACGAGCGGAACCTGATG AACGTGGTGTATAAACTCAATGGCCAAGCAGAGATC TTTTTTCGTAAAGCATCACTGAGCGCACGTCACCCCT GAGCACCCGGCAGGGTTGCCAATTAAAAAAAACAG GCCCCGACGGAAGAATCTTGTTCCTCATATGATCTC ATTAAGAATAAGCGGTATACAGTTGACCAGTTTCAG TTTTACGTGCCAATTACTATTAATTTTAAAGCAACT GGGACTTCAAATATCAACCCGTCGGTCACTGATTAT ATTCGTACGGCCGATGACCTCCATATCATTGGCATT GATCGCGGTGAGCGCCATTTACTTTTATTAGTGGTG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		ATTGACTCACAAGGGCGCATCTGTGAACAGTTTTCC TTAAACGAGATCGTAACGCAATACCAAGGTCACCAG TACCGTACAGATTATCATGCTCTCTTGCAGAAAAAA GAGGATGAACGGCAAAAAGCTCGCCAGTCTTGGCAA TCGATCGAAAACATCAAGGAATTAAAAGAGGGGTAT CTGAGCCAAGTAGTGCACAAGGTTTCTGAACTGATG ATCAAATATAAAGCAATTGTGGTGTTGGAAGATTTA AATGCTGGGTTCAAGCGGAGTCGGCAGAAGGTTGAA AAGCAAGTGTATCAAAAATTTGAGAAGATGCTGATC GACAACTTAACTATCTTGTGTTCAAGACCGCAGAA GCTGACCAACCTGGCGGCCTCCTGCACGCATACCAA TTAACAAATAAATTTGAGTCATTCAAGAAAATGGGG AAGCAAAGTGGCTTCCTCTTCTACATTCCTGCATGG AACACGTCTAAAATCGACCCGACCACGGGCTTTGTC AACCTTTTTGATACCCGGTATGAGAACGTAGACAAA TCCCGTGCCTTCTTCGGCAAATTCGATAGCATCCGC TACCGTGCGGACAAGGGCACGTTTCGAGTGGACGTTT GATTATAATAACTTTACAAAAAGGCCGAAGGTACG CGGTGAGCTGGTGTTTGTCTTCTCATGGTAACCGG GTCCGTACTTTCCGCAATCCTGCGAAAAACAACCAA TGGGACAACGAAGAGATCGACTTAACACAAGCGTTC CGCGATCTGTTTGAAGCTTGGGGGATCGAGATCACT TCGAACTTAAAAGAGGCCATTTGCAACCAGTCTGAG AAGAAATTCTTTTCTGAGCTTTTTCGAACTGTTCAAA CTTATGATCCAGCTGCGGAACCTCAGTGACAGGCACG AATATCGACTATATGGTGAGCCAGTCGAGAATCAC TACGGCACGTTCTTCGATTTCGCGCACATGCGATTCT TCTCTGCCGGCTAACGCTGACGCTAATGGTGCTTAT AATATTGCCCGTAAGGGGTTAATGCTGGCTCGCCGC ATTCAGGCTACCCCTGAGAATGATCCGATCTCCTTA ACATTGAGCAACAAAGAGTGGTTACGCTTTGCACAG GGGCTCGATGAGACAACAACCTACGAGGCGGCCGCA AAAAGGCCGGCGGCCACGAAAAAGGCCGGCCAGGCA AAAAAGAAAAAGGCTAGCGGCAGCGGCGCCGATCC CCAAAGAGAAAAAGGAAGTTGAAGACCCCAAGAAA AAGAGGAAGGTGTGATAA (SEQ ID NO: 30)
ABW4	MGHHHHHHSSGLVPRGSG TMKNMESFINLYPVSKTL RFELKPIGKTLETFSRWI EELKEKEAIELKETGNLL AQDEHRAESYKKVKKILD EYHKWFITESLQNTKLNG LDV FYHNYMLPKKEDHEK KAFASCQDNLRKQIVNAF RQETGLFNKLSGKELFKD SKEEVALLKAIVPYFDNK TLENIGVKSNEGALLLIE EFKDFTTYFGGFHENRKN MYSDEAKSTAVAFRLIHE NLPRFIDNKKVFEEKIMN	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGAAGAACATGGAG TCTTTTATTAATTTATATCCGGTTTCGAAAACCTTTA CGTTTTGAGTTAAAGCCTATTGGCAAAACACTCGAA ACTTTCTCCCGCTGGATCGAAGAGTTGAAAGAGAAA GAGGCTATTGAGCTGAAAGAAACTGGCAACCTGTTG GCGCAGGATGAGCATCGGGCCGAGTCTTATAAGAAG GTCAAAAAAATTCTTGACGAATATCATAAATGGTTC ATCACTGAAAGCCTCCAGAACACAAAGTTAAATGGG TTGGACGTTTTTTATCATAACTATATGCTCCCGAAG AAAGAGGACCATGAGAAGAAAGCTTTTGCTTCGTGT CAAGATAATCTCCGTAAGCAAATTGTAAACGCGTTT CGTCAAGAAACCGGTTTATTTAACAACTGTCAGGC AAAGAACTGTTTAAAGATTCTGAAGGAAGAGGTTGCA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	SELKDKFPEILKELEQIL QVNEIEEMFQLDYFNDTL IQNGIDVYNHLIGGYAEE GKKKIQGLNEHINLYNQI QKEKNKRIPRLKPLYKQI LSDRETASFVIEAFENDG ELLESLEKSYRLLQQEVF TPEGKEGLANLLAAIAES ETHKIFLKNLGLTEISQ QIYESWSLIEEAWNKKQYD NKQKQVETETETYVDNRKK AFKSIKSFSAIEVEEWVK ALGNEKHKGKSVATYFKS LGKTDEKVSLEQVENNY NIIKDLNTPYPYPSKDLA QQKDDVEKIKNYLDSLKA LQRFIKPLLGSGEESDKD AHFYGEFTAFWDVLDKVT PLYNKVRNYMTKKPYSTE KFKLNFENSYFLNGWAQD YETKAGLIFLKDGNVFLA INNKKLDEKEKKQLKTNY EKNPAKRIILDFFQKPDNK NIPRLFIRSKGDNFAPAV EKYNLPISDVIDIYDEGK FKTEYRKINEPEYLSLH KLIDYFKLGFSEKESYKH YSFSWKKTHEYENIAQFY HDVEVSCYQVLDENINWD SLMEYVEQNKLYLFQIYN KDFSPNSKGTPNMHTLYW KMLFNPDLNKDVVYKLNG QAEVFYRKASIKKENKIV HKANDPIDNKNELNKKKQ NTFEYDIVKDKRYTVDFK QFHVPIITLNFKAEGNLNL NSKVNEYIKECDDLHIIG IDRGERHLLYLSLIDMKG NIVKQFSLNEIVNEHKGN TYRTNYHNLDDKREKERE KERESWKT IETIKELKEG YISQVVKITQLMIEYNA IVVLEDLNFGRGRFRKV EKQVYQKFEKMLIDKLN LVDKKKEANESGGTLKAY QLTDSYADFMKYKKKQCG FLFYVPAWNTSKIDPTTG FVNLFDTYHVNVSQAQEF FSKFKSIRYNAANNYFEF EVTDYFSFSGKAEGTKQN	CTGTTGAAAGCCATTGTACCGTATTTTCGATAACAAG ACTCTGGAAAACATTGGTGTTAAGAGTAATGAAGGG GCTCTCCTTTTAATTGAAGAGTTCAAGGATTTTACC ACGTATTTCCGGTGGCTTCCATGAGAATCGCAAAAAT ATGTATAGCGACGAAGCAAAATCAACAGCGGTTGCC TTTCGTCTTATTCACGAAAATTTGCCGCGCTTCATT GACAATAAGAAGGTCTTCGAAGAGAAAATCATGAAT AGTGAATTAAAGGATAAAATTTCCAGAGATTTTGAAG GAGCTGGAACAGATTCTGCAAGTCAACGAGATTGAA GAGATGTTTCAGCTCGACTATTTTAACGACACATTG ATCCAGAATGGCATCGATGTCTATAACCATTGATC GCGCGCTACGCCGAGGAAGGCAAGAAAAAAATTCAA GGGCTTAACGAGCATATTAACCTCTATAACCAGATC CAGAAGGAGAAGAATAAGCGTATCCCGCGGCTGAAA CCACTCTATAAGCAAATTTTGAGTGATCGCGAAACC GCCTCATTTGTTATCGAGGCGTTTGAGAACGATGGC GAGTTATTAGAATCATTGGAGAAGTCATATCGCTTA CTGCAGCAGGAGGTCTTTACGCCTGAAGGTAAAGAA GGTCTGGCGAATTTACTCGCAGCAATCGCTGAAAGC GAGACACACAAGATCTTTCTGAAGAACGACTTGGGT CTCACCGAGATCTCTCAACAAATTTATGAATCATGG TCGCTGATTGAAGAGGCATGGAATAAACAAATATGAC AACAAACAGAAGAAAGTTACGGAGACAGAGACATAT GTGGACAATCGGAAAAAGGCTTTCAAGTCCATCAAG AGCTTTAGCATCGCAGAGGTTGAGGAATGGGTGAAA GCACTTGGGAATGAGAAACACAAGGGCAAAAGCGTG GCAACCTATTTTAAAAGTCTCGGGAAGACTGACGAA AAAGTTAGCCTTATTGAACAGGTAGAGAACAATTAT AATATCATCAAGGACCTTTTGAACACACCGGTATCCT CCTTCGAAGGACTTGGCCCAGCAAAAAGATGACGTT GAAAAAATCAAAAATTATTTGGACTCTCTGAAGGCC CTCCAGCGGTTCATTAAGCCATTGTTGGGTAGCGGG GAGGAATCCGATAAAGATGCGCACTTTTATGGTGAG TTTACCGCTTTCTGGGATGTGCTCGACAAAGTAACC CCACTCTACAATAAAGTCCGCACTATATGACTAAG AAACCTTATAGCACAGAGAAATTTAAGCTGAATTTT GAAAATAGTTACTTTTTGAATGGTTGGGCACAGGAC TACGAGACAAAAGCGGGGCTTATCTTCTTGAAGGAC GGCAATTACTTCCTTGCCATCAATAATAAGAAATTA GATGAAAAGGAGAAAAAACAGCTCAAGACTAATTAT GAGAAGAATCCTGCGAAGCGTATCATCTTAGACTTT CAGAAGCCAGACAATAAGAACATTCTCTGCTTGTTT ATTCTGCAGTAAAGGCGACAATTTCTGCTCCTGCAGTA GAAAAGTATAATCTTCCGATCTCTGACGTTATTGAC ATCTATGACGAGGGGAAGTTTAAGACTGAGTATCGC AAAATTAACGAGCCGGAATATCTCAAATCTCTCCAT AAGCTGATTGACTACTTCAAACCTTGGGTTCTCCAAG CATGAATCCTACAAGCATTATTCTTTTTTCATGGAAG AAAACACATGAGTATGAGAACATCGCCCAGTTTTAC CACGACGTGGAGGTCTCTTGCTATCAGGTGCTCGAC

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	WIICTHGTRIINFNRNPEK NSQWDNKEVVITDEFKKL FEKHGIDYKNSSDLKGQI ASQSEKAFFHNEKKDTKD PDGLLQLFKLALQMRNSF IKSEEDYLVSPVMNDEGE FFDSRKAQPNQPENADAN GAYNIAMKGKWVVKQIRE SEDLDKLLAISNKEWLN FAQRSAAAKRPAATKKAG QAKKKKASGSGAGSPKKK RKVEDPKKKRKV (SEQ ID NO: 42)	GAAAATATTAAGTGGGATTCCTCATGGAGTATGTA GAACAGAACAAATTGTACTTGTTCCAGATTTATAAC AAAGACTTCTCCCCAAACTCGAAAGGCACTCCGAAT ATGCACACTTTGTACTGGAAGATGTTGTTTAATCCG GATAATCTTAAGGACGTGGTCTATAAGCTGAACGGT CAGGCTGAAGTATTCTACCGGAAGGCGAGTATTAAG AAAGAAAAACAAGATTGTCCACAAGGCGAACGACCTT ATTGACAATAAAAAACGAGTTGAATAAGAAAAAGCAA AATACATTTGAATACGACATCGTCAAAGATAAACGG TATACAGTGGATAAGTTTCAATTCCATGTTCCCTATC ACGCTCAACTTTAAAGCTGAAGGCCTGAATAACTTG AATAGCAAAGTTAACGAATACATCAAAGAGTGTGAC GACCTTCACATTATTGGCATCGACCGGGGTGAACGG CACCTCTTGATCTGAGCCTCATCGATATGAAAGGT AACATTGTAAAGCAATTTAGTCTTAACGAGATCGTT AATGAGCACAAGGGGAACACGTACCGCACGAACAT CATAACCTCTTGGACAAACGTGAAAAGGAACGTGAA AAAGAGCGCGAGTCATGGAAAACCATTGAGACCATC AAAGAGCTGAAAGAAGGCTATATTAGTCAAGTAGTA CATAAAATCACTCAGTTAATGATCGAATATAATGCG ATCGTTGTACTCGAAGACCTGAATTTCTGGCTTCAAA CGCGGCCGGTTCAAGGTGGAGAAGCAAGTGTATCAA AAATTTGAGAAGATGTTAATTGATAAACTGAACCTAC TTGGTCGATAAGAAGAAGGAAGCCAATGAGAGTGGC GGGACACTCAAAGCCTACCAGCTTACCGATAGTTAC GCTGACTTCATGAAGTACAAGAAAAAGCAATGCGGC TTCCTGTTTTATGTCCCGGCCTGGAACACTTCCAAA ATCGATCCTACTACTGGGTTCTGTGAATCTGTTTGAC ACACATTATGTCAATGTTAGTAAGGCCCAGGAATTT TTCTCGAAATTCAGTCAATTCGCTACAACGCGGCC AACAACTATTTGAGTTTGAAGTAACAGATTATTTT TCCTTCAGTGGTAAAGCTGAGGGCACCAAGCAGAA TGGATCATTTGCACCCATGGCACCCGCATTATCAAT TTTCGTAACCCGGAAAAAAATTCGAGTGGGATAAT AAGGAAGTAGTGATCACAGATGAATTCAGAAACTG TTTGAGAAGCACGGCATTGACTACAAAAATAGTTCC GACCTCAAGGGGCAGATCGCCTCTCAATCGGAGAAG GCGTTTTTTTCATAACGAAAAAAAAGATACAAAGGAC CCAGATGGCCTTCTGCAGCTTTTTAACTGGCGCTG CAGATGCGGAACTCTTTCATTAAGAGCGAAGAGGAC TACTTAGTATCTCCTGTGATGAACGACGAAGGTGAA TTCTTTGACTCGCGCAAAGCCCAGCCTAATCAGCCA GAGAACGCTGATGCTAATGGGGCGTACAATATTGCA ATGAAAGGGAAATGGGTGTTAAGCAAATCCGCGAA TCGGAGGACCTCGACAAGCTGAACTGGCAATCTCA AATAAAGAATGGTTGAACTTCGCCCAGCGCTCCGCG GCCGCAAAAAGGCCGGCGGCCACGAAAAAGGCCGGC CAGGCAAAAAGAAAAAGGCTAGCGGCAGCGCGCC GGATCCCCAAAGAAGAAAAGGAAGGTGAAGACCCC

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		AAGAAAAAGAGGAAGGTGTGATAA (SEQ ID NO: 43)
ABW5	MGHHHHHSSGLVPRGSG TMKNILEQFVGLYPLSKT LRFELKPLGKTLEHIEKK GLIAQDEQRAEEYKLVKD IIDRYHKAFIHMCLKHFK LKMYSQGYDSLEEYRKL ASISKRNEKEEQQFDKVK ENLRKQIVDAFKNGGSYD DLFKKELIQKHLPRFIEG EEEKRIVDNFNKFTTYFT GFHENRKNMYSDEKESTA IAYRLIHENLPLFLDNMK SFAKIAESEVAARFTEIE TAYRTYLNVEHISELFTL DYFSTVLTQEQIEVYNNI IGGRVDDDNVKIQGLNEY VNLYNQQKDRSKRLPLL KSLYKMILSDRIAISWLP EEFKSDKEMIEAINNMHD DLKDILAGDNEDSLKSL QHIGQYDLSKIYIANNPG LTDISQQMFGCYDVFTNG IKQELRNSITPSKKEKAD NEIYEERINKMFKSEKSF SIAYLNSLPHPKTDAPQK NVEDYFALLGTCNQNDQ PINLFAQIEMARLVASDI LAGRNVNLNQSENDIKLI KDLLDAYKALQHFVKPLL GSGDEAEKDNEFDARLRA AWNALDIVTPLYNKVRNW LTRKPYSTEKIKLNFENA QLLGWDQNKEPDCTSVL LRKDGMYYLAIMDKKANH AFDCDCLPSDGACFEKID YKLLPGANKMLPKVFFSK SRIKEFSPSESIIAAYKK GTHKKGPNFSLSDCHRLI DFFKASIDKHEDWSKFRF RFSDTKTYEDISGFYREV EQQGYMLGFRKVSEAFVN KLVDEGKLYLFHIWNKDF SKHSKGTPNLHTIYWKML FDEKNLTDVIYKLNGQAE VFYRKSLDLNKTTHKA HAPITNKNTQNAKKGSVF DYDIIKNRRYTVDKFQFH VPITLNFKATGRNYINEH	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGAAGAACATCTTA GAGCAGTTTGTCTGGCTTATATCCGTTGTCTAAAACA CTTCGGTTTGAGCTTAAACCTTTGGGTAAGACGTTG GAACATATTGAGAAAAAAGGCTTGATTGCCCAAGAC GAACAGCGGGCGGAGGAGTACAAATTGGTTAAAGAT ATTATTGATCGCTACCACAAGGCTTTTATTTCATATG TGCTTAAAACATTTTAAAGCTCAAGATGTACAGTGAA CAAGGGTATGATAGCTTGGAGGAGTACCGCAAGCTT GCGTCAATTTCCAAACGCAACGAGAAAGAGGAGCAG CAATTTGACAAAGTCAAGGAAAATCTTCGTAAGCAA ATTGTGCGACGCGTTTAAAAATGGCGGGAGTTATGAT GATCTGTTTAAAGAAAGAATTGATCCAGAAACACCTC CCACGTTTTATTGAGGGTGAAGAAGAAAAACGTATC GTTGACAACCTTCAACAAGTTCACGACCTATTTTACT GGTTTTTCATGAAAATCGCAAGAATATGTATAGTGAC GAAAAGGAATCGACGGCTATTGCTTATCGTCTCATT CACGAAAACCTTGCCATTGTTTTTGGATAACATGAAG AGCTTCGCTAAGATCGCCGAATCGGAAGTGGCTGCT CGTTTTTACCGAAATCGAAACCGCTTACCGGACATAC TTGAACGTAGAACACATTAGTGAAGTGTTCACCCTC GACTATTTTAGCACGGTTTTTGACGCAAGAACAAATC GAAGTATATAATAACATTATCGGCGGGCGCGTCGAC GACGACAACGTAAAGATCCAAGGGTTGAATGAGTAC GTAAATTTATATAATCAGCAGCAGAAGGACCGGTCT AAGCGCTTACCGCTTCTTAAGTCCCTCTACAAAATG ATCTTATCCGATCGTATTGCAATTTCTGTTGTTACCT GAGGAGTTCAAATCCGATAAGGAGATGATTGAAGCA ATTAACAACATGCATGACGACCTGAAGGACATTCTG GCAGGCGACAACGAAGACTCGCTTAAGTCCTTACTG CAGCATATTGGCCAATACGATCTCTCGAAAATCTAC ATTGCGAACAATCCGGGCTGACAGATATCTCACAA CAAATGTTCTGGGTGTTATGACGTCTTTACTAATGGG ATCAAGCAGGAGCTCCGGAACAGTATTACCCCTTCA AAAAAGGAGAGAGCCGATAACGAAATCTACGAGGAG CGGATTAACAAAATGTTTAAAGTGAGAAGAGTTTC TCAATTGCCTACCTGAATTCGTTGCCGCACCCAAAG ACGGATGCGCCTCAAAAAAATGTTGAGGATTATTTT GCTCTCCTGGGGACTTGCAATCAAAACGATGAACAG CCGATTAATTTGTTTGCCCAAATTGAGATGGCACGC TTAGTCGCCTCTGATATTCTCGCAGGCCGGCACGTT AATTTGAACCAATCTGAGAATGATATCAAGTTAATC AAGGATCTGTTAGATGCTTACAAGGCTCTGCAGCAT TTCGTCAAACCACTCCTTGGCTCGGGTGACGAGGCT GAGAAAGATAACGAGTTCGATGCACGCCCTCCGTGCG GCTTGGAATGCGTTGGACATTGTTACACCACTCTAT AACAAGGTTTCGGAAGTGGCTGACCCGCAAACCATAT TCTACAGAAAAAATCAAGCTTAATTTGAAAACGCC

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	TQEAIRNNGIEHIIGIDR GERHLLYLSLIDLKGNIV KQMTLNDIVNEYNGRTYA TNYKDLLATREGERTDAR RNWQKIENIKEIKEGYLS QVVHILSKMMVDYKAIVV LEDLNTGFMNRNRQKIERQ VYEKFEKMLIDKLNCYVD KQKDADETGGALHPLQLT NKFESFRKLKQSGWLFY IPAWNTSKIDPVTGFVNM LDTRYENADKARCFFSKF DSIRYNADKDWFEFAMDY SKFTDKAKDITYTWWTLCS YGTRIKTFRNPAKNNLWD NEEVVLTFDEFKKVFAAAG IDVHENLKEAICALTDKK YLEPLMRLMTLLVQMRNS ATNSETDYLLSPVADESG MFYDSREGKETLPKDADA NGAYNIARKGLWTIRRIQ ATNCEEKVNVLVLSNREWL QFAQQKPYLNDAAKRPA ATKKAGQAKKKKASGSGA GSPKKKRKVEDPKKKRKV (SEQ ID NO: 55)	CAACTTCTGGGGGGTTGGGATCAGAACAAAGAACCG GATTGCACATCAGTCCTCCTTCGGAAGGATGGGATG TACTATTTAGCGATCATGGATAAAAAGGCGAATCAC GCCTTTGACTGTGACTGCTTACCGTCTGACGGGGCC TGTTTCGAGAAAATTGACTACAAGCTGCTCCCGGGC GCGAATAAAATGTTGCCGAAAGTTTTTTTTTCTAAA AGCCGCATCAAAGAATTTTCCCCTTCGGAATCGATC ATCGCTGCTTATAAAAAGGGGACTCATAAAAAGGG CCGAATTTTCAGTCTCTCTGATTGTCTATCGCTTGATT GACTTTTTTTAAGGCTAGCATTGATAAGCACGAAGAT TGGTCAAAATTTTCGTTTTTCGCTTCTCAGATACCAAA ACGTATGAAGACATCAGTGGTTTTCTACCGTGAAGTA GAACAGCAAGGCTATATGCTGGGTTTTTCGTAAAGTC TCTGAGGCCTTTGTGAATAAACTCGTTGATGAAGGT AAGTTATACTTATTCCATATCTGGAACAAAGACTTT AGTAAGCACTCCAAAGGTACACCTAATCTCCACACT ATTTATTGGAAAATGCTCTTCGATGAGAAAAATCTC ACTGACGTCATCTACAACTGAATGGGCAGGCTGAA GTATTCTACCGTAAAAAAGTCTGGATCTTAATAAG ACAACTACTCACAAGGCACATGCCCCAATCACCAAT AAAAATACCCAAAACGCAAAGAAGGGTAGTGTTTTT GATTACGATATCATCAAAAATCGTCGCTACACAGTG GACAAATTCCAGTTCCACGTCCCTATCACCTTAAAT TTTAAGGCAACAGGTCGTAATTACATTAATGAGCAC ACTCAAGAGGCAATCCGTAATAATGGCATCGAACAT ATCATTGGCATCGACCGTGGGGAGCGTCACTTGCTT TACTTGTCGCTCATTGATCTGAAGGGTAATATCGTC AAGCAGATGACCCTTAATGATATTGTCAATGAATAT AATGGTCGGACTTATGCGACGAACTACAAGGACTTG CTGGCAACACGGGAGGGTGAGCGTACGGACGCTCGG CGCAACTGGCAGAAGATTGAAAATATTAAAGAAATC AAGGAAGGTTACCTTAGCCAGGTGGTGCACATCTTG AGTAAAATGATGGTCGACTACAAGGCTATCGTTGTT CTGGAAGACTTGAATACAGGCTTCATGCGGAATCGT CAAAAAATCGAACGTCAAGTATATGAGAAGTTTCGAA AAAATGTTAATTGACAAGCTGAAGTCTATGTTGAC AAACAAAAGGATGCTGACGAGACGGGCGGTGCCCTC CACCCGCTGCAGCTGACAAACAAATTTGAGTCGTTT CGTAAGTTAGGTAAGCAGAGTGTTGGCTTTTTTAC ATCCCAGCATGGAACACTTCGAAAATCGACCCAGTT ACTGGGTTCTGTAACATGTTAGACACGCGCTACGAG AACGCCGATAAGGCGCGGTGTTTCTTCTCGAAATTC GATTCCATCCGGTATAACGCTGACAAAGATTGGTTT GAGTTTGCTATGGATTACAGTAAGTTCACTGATAAA GCGAAAGATACTTACACGTGGTGGACTCTGTGTTCC TATGGGACGCGTATTAAAACTTTTCGTAATCCGGCT AAGAATAATTTGTGGGATAATGAGGAGGTTGTCCTT ACTGATGAGTTCAAGAAAGTTTTTCGCAGCGGCAGGT ATTGATGTCCATGAGAACCTTAAGGAAGCGATCTGT GCTCTGACAGATAAAAAAGTATCTTGAACCACTCATG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		CGTCTCATGACCCTGCTCGTTCAAATGCGGAACTCT GCTACTAACTCCGAAACAGACTATTTACTTTCACCA GTTGCTGACGAGTCAGGGATGTTCTATGACTCCCGC GAAGGGAAGGAAACACTGCCAAAAGATGCGGACGCC AACGGTGCATATAACATTGCCCCGTAAGGGCCTCTGG ACCATCCGGCGGATTCAAGCCACCAACTGTGAGGAG AAAGTTAACTTAGTCCTCAGTAATCGTGAATGGTTG CAGTTTGCCCAGCAGAAACCATATCTGAATGATGCG GCCGCAAAAAGGCCGGCGGCCACGAAAAAGGCCGGC CAGGCAAAAAGAAAAAGGCTAGCGGCAGCGGCCGCC GGATCCCCAAAGAAGAAAAGGAAGGTTGAAGACCCC AAGAAAAAGAGGAAGGTGTGATAA (SEQ ID NO: 56)
ABW 6	MGHHHHHHSSGLVPRGSG TMIYRENFKRKKEKIEMN TGFNDFTNLSSVTKTLCN RLIPTEITAKYIKEHGV EADQERNMMSQELKNILN DFYRSFLNENLVKVHED FKPLFTEMKKYLETKDNK EALEKAQDDMRKAIHDI ESDDRYKKMFKAELITASI LPEFILHNGAYSAAEKEE KMQVVKMFNGFMTSFSAF FTNRENCFSKEKISSAC YRIVDDNAKIHFDNIRIY KNIANKFDEIEMIEKIE EAAGGADIRNIFSYNFDH FAFNHFVSQDDISFYNYV VGGINKFMNLYCQATKEK LSPYKLRHLHKQILCIEE SLYDVPKFNCDDEVYAA VNDFLNNVRTKSVIERLQ MLGKNADSVDLDKIYISK KHFTNISQTLYRDFSVIN TALTMYSYIDTLPKGKGTK EKKAASMAKNTELISLGE IDKLVDKYNLCPDKAAST RSLIRSISDIVADYKANP LTMNSGIPLAENETEIAV LKEAIEPFMDIFRWCAKF KTDEPVDKDTDFYTELED INDEIHSIVSLYNRTRNY VTKKPYNTDKFGLYFGTS SFASGWSESKEFTNNAIL LAKDDKFYLGVFNAKNKP AKSIIKGHDTIQDGDYKK MVYSLLTGPNKMLPHMFI SSSKAVPVYGLTDELLSD YKKGRHLKTSKNFDIDYC	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGATCTACCGTGAG AATTTTAAGCGGAAAAAGGAGAAGATTGAAATGAAC ACTGGGTTTAATGACTTCACTAATTTGAGTTCCGTG ACCAAGACGTTATGCAACCGGTTGATCCCAACAGAA ATTACCGCAAAGTACATTAAGGAGCATGGGGTAATT GAGGCGGACCAAGAACGGAACATGATGAGTCAAGAG CTGAAAAATATCTTGAATGACTTTTACCGGAGTTTC CTGAACGAGAACCTTGTGAAGGTGCACGAACTTGAT TTCAAGCCGTTATTACCGAGATGAAAAAGTACCTC GAAACAAAAGATAACAAGGAAGCACTCGAAAAGGCC CAGGACGACATGCGGAAGGCAATCCATGATATCTTT GAAAGTGATGACCGCTACAAAAAATGTTCAAGGCT GAGATCACGGCGTCGATTTTGCTGAATTCATTCTT CATAACGGGGCATATTACGCCGAAGAAAAGGAGGAG AAAATGCAAGTAGTCAAGATGTTCAATGGCCTTTATG ACGTCTTTCTCAGCATTCCTTACGAATCGTGAGAAT TGTTTCTCCAAAGAAAAGATCAGCTCCTCCGCATGT TACCGTATTGTTGATGACAACGCGAAAAATCCATTC GATAACATTCGTATTTATAAAAATATCGCCAACAAG TTCGATTATGAAATTGAAATGATCGAGAAGATCGAA GAGGCGGCGGGGGGTGCCGACATTCGTAATATCTTC TCGTACAACCTTTGACCACTTTGCATTCAATCATTTT GTTAGTCAAGATGATATCTCATTCTACAATTATGTT GTTGGTGGTATTAACAAGTTTATGAACTTGTATTGT CAAGCCACCAAAGAGAAATTATCGCCTTATAAACTG CGTCACCTTCACAAACAGATTCTGTGTATTGAGGAA AGCCTCTATGACGTGCCAGCGAAGTTTAATTGTGAT GAGGACGTATATGCAGCTGTCAACGATTTTCTTAAT AACGTTTCGGACGAAATCAGTAATTGAACGCTTGCAA ATGCTCGGCAAAAATGCAGACAGTTACGACCTGGAT AAAATTTATATCTCTAAAAAGCACTTCACCAATATC TCTCAAACCTTTATATCGCGACTTCTCTGTGATCAAC ACTGCCCTCACTATGTCTTATATCGATACTCTTCCG GGTAAGGGGAAAACCAAGGAAAAAAGGCAGCATCG ATGGCCAAAAACACCGAACTTATTTTCGTTAGGCGAA ATTGATAAGTTGGTGGATAAATATAACCTCTGTCCA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	HKLIDYFKHCLALYTDWD CFNFKFSDTESYNDIGEF YKEVAEQGYMNTYIGS DDIDSLQENGQLYLFQIY NKDFSEKSF GKPSKHTAI LRSLSFDENVADPVIKLC GGTEVFFRPKSIKTPVVH KKGSILVSKTYNAQEMDE NGNIITVRKCVDDVYME LYGYNNSGTPLSAEALK YKDIVDHRTPYDIIKDR RYTEDEFFINMPVSLNYK AENRRVNVNEMALKYIAQ TKDTYIIIGIDRGERNLLY VSVIDTDGNIVEQKSLNI INNVDYQAKLKQVEIMRK LARQNWQGVKIADLKKG YLSQAVHEVAELVIKYN IVVMEDLNSRFKEKRSKI ERGVYQQFETSLIKTLNY LTFKDRKPLEAGGIANGY QLTYPESLKNVGSQCGC ILYVPAAYTSKIDPTTGF VTLFKFKDISSEKAKTDF IGRFDCIRYDAEKDLFAF EFDYDNFETYETCARTKW CAYTYGTRVKKTFRNRKF VSEVIIDITEEIKKTLAA TDINWIDSHDIKQEIIDY ALSSHI FEMFKLTVQMRN SLCESKDREYDKFVSPIL NASGKFFD TDAADKSLPI EADANDAYGIAMKGLYNV LQVKNNWAEAGEKFKFSRL SNEDWFNF MQRAAKRP AATKKAGQAKKKKASGSG AGSPKKKRKVEDPKKKRK V (SEQ ID NO: 68)	GATAAGGCAGCTAGCACTCGTAGCCTCATTCCGGTCT ATTAGCGACATCGTCGCTGACTACAAGGCAAACCCCT CTTACAATGAATAGTGGGATTCCGTTGGCAGAGAAC GAGACAGAAATCGCGGTGTTAAAAGAGGCGATCGAG CCTTTTATGGATATCTTCCGGTGGTGTGCTAAGTTT AAAACCGACGAGCCTGTCGATAAGGATACAGATTTT TACACGGAGTTAGAAGACATTAACGATGAAATCCAT AGTATTGT CAGTCTTTATAACCGGACCCGGAATTAT GTCATAAAAAGCCGTACAACACAGATAAGTTCCGGT CTGTATTTTGGCACTTCGTCGTTCCGCATCGGGTTGG AGCGAGAGCAAAGAGTTTACTAACAACGCAATTTTG TTAGCCAAGGATGACAAGTTTTACCTCGGCGTGTTT AACGCAAAAAACAAGCCAGCAAAATCGATTATCAAA GGGCATGACACAATCCAAGATGGTGATTATAAGAAA ATGGTGATTCTACTGCTCACC GGCCCAAATAAGATG CTTCTC CATGTTTATCTCGAGCAGTAAAGCGGTT CCTGTTTACGGGCTCACTGACGAGCTTCTCAGCGAC TATAAGAAAGGTGCGCACCTTAAGACATCCAAGAAT TTCGACATTGATTACTGTCACAACTTATCGATTAC TTCAAACATTGTCTCGCTTTGTATACTGATTGGGAT TGCTTCAACTTCAAATTCTCTGATACGGAGTCTTAC AATGATATCGGCGAGTTCTACAAAGAGGTTGCCGAG CAAGGCTACTACATGAACTGGACATATATCGGGTCG GACGATATCGATTTCGCTGCAGGAAAACGGCCAGCTC TATCTTTTTTCAAATTTTATAACAAAGATTTTCAGCGAA AAGTCATTCCGGTAAACCGTCTAAACATACGGCCATC CTGCGTAGCTTATTTCAGCGATGAAAACGTGGCCGAC CCAGTCATTAAACTGTGTGGGGGGACCGAAGTTTTT TTCCGGCCGAAGTCTATTAAAGACACCAGTAGTACAT AAAAAAGGCAGCATCCTCGTATCCAAAACCTATAAC GCACAAGAAATGGACGAGAATGGTAATATCATCACC GTGCGGAAGTGTGTTCCAGACGACGTCTATATGGAG CTCTACGGCTATTACAACAACCTCTGGGACGCCTCTG TCCGCCGAAGCTTTGAAATACAAGGATATTGTGGAC CACCGCACGGCTCCGTACGACATTATCAAGGACCGG CGTTACACCGAAGACGAATTTTTTCATCAACATGCCG GTGTCATTGAATTATAAAGCGGAAAACCGCCGTGTT AATGTGAACGAAATGGCCTTAAATACATCGCACAG ACCAAGGACACCTACATCATTGGCATCGATCGGGGC GAACGTAATCTGTTGTATGTGAGCGTTATCGATACT GACGGCAATATCGTTGAGCAAAGAGTCTCAATATC ATCAATAACGTGGATTATCAAGCCAAATTAAAGCAA GTGGAAATCATGCGTAAACTGGCCCGTCAGAATTGG AAGCAGGGGGTAAAGATTGCAGACCTGAAAAAGGGC TACCTGTCACAAGCGGTACATGAAGTCGCGGAACCT GTAATTAAATACAACGGGATTGTTGTAATGGAGGAC TTAAACTCCCGCTTCAAAGAGAAGCGTTCTAAAATT GAACGCGGCGTCTACCAACAGTTTGAGACATCATTA ATCAAGACATTGAATTATTTGACGTTCAAAGATCGC AAACCGTTAGAAGCCGGGGGCATTGCGAATGGTTAT

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		CAATTAACCTTATATTCCGGAGTCTCTTAAAAATGTG GGCTCTCAGTGCGGCTGTATCTTGTATGTGCCAGCA GCCTACACCTCGAAGATCGACCCTACCACTGGTTTC GTCACCTTGTTCAAATTCAAAGACATTCGAGCGAG AAAGCTAAAACGGATTTTATTGGTCGGTTCGACTGC ATCCGTTATGATGCAGAAAAGGACCTTTTCGCATTT GAATTTCGATTATGACAACCTTTGAGACTTATGAGACT TGTGCGCGTACCAAATGGTGTGCATATACATACGGG ACTCGGGTGAAGAAAACCTTTCCGGAATCGGAAATTC GTGTGAGAGGTGATCATCGACATCACTGAAGAGATC AAGAAGACCCTTGCAGCGACCGATATTAATTGGATT GACAGTCACGACATCAAACAAGAGATCATCGACTAT GCCCTTAGCAGCCATATTTTTGAAATGTTCAAATTA ACGGTACAGATGCGTAACAGCCTTTGCGAGAGTAAA GATCGCGAGTACGACAAGTTCGTCTCACCTATTCTC AACGCGTCGGGCAAATTTTTCGACACCGATGCCGCT GATAAAAGTCTGCCTATTGAAGCTGATGCGAACGAT GCGTATGGTATTGCTATGAAAGGGTTGTATAATGTT TTACAAGTCAAAAACAACCTGGGCGGAGGGCGAGAAA TTTAAGTTCTCCCGTTTAAGCAACGAAGATTGGTTC AACTTCATGCAAAAGCGGGCGGCCGCAAAAAGGCCG GCGGCCACGAAAAAGGCCGGCCAGGCCAAAAAGAAA AAGGCTAGCGGCAGCGCGCCGGATCCCCAAAGAAG AAAAGGAAGGTTGAAGACCCCAAGAAAAAGAGGAAG GTGTGATAA (SEQ ID NO: 69)
ABW 7	MGHHHHHSSGLVPRGSL QMTMDYGNQFERRAPLT KTITLRLKPIGETRETIR EQKLLAQDAEFRKLVETV TPIVDDCIRKIADNALCH FGTEYDFSCLGNAISKND SKAIKKETEKVEKLLAKV LTENLPDGLRKVNDINSA AFIQDTLTSFVQDDADKR VLIQELKGKTVLMQRFLT TRITALTVWLPDRVFENF NIFIENAEKMRILLDSPL NEKIMKFDPDAEQYASLE FYGQCLSQKDIDSYNLII SGIYADDEVKNPGINEIV KEYNQQIRGDKDESPLPK LKKLHKQILMPVEKAFFV RVLSNDSARSILEKILK DTEMLPSKIIIEAMKEADA GDIYVGSRLHELHSHVIY GDHGKLSQIIYDKESKRI SELMETLSPKERKESKKR LEGLEEHIRKSTYTFDEL NRYAEKNVMAAYIAAVEE SCAEIMRKEKDLRTLISK	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCCTGCAGATGACAATGGATTAC GGTAACGGTCAATTTGAGCGGCGCGCCCCGCTCACC AAGACAATCACTCTCCGGTTGAAACCGATCGGGGAG ACCCGTGAGACGATTTCGCGAGCAAAAGCTCCTCGAA CAAGATGCTGCATTCCGTAACTTGTTGAAACTGTC ACCCCTATCGTGGATGATTGTATCCGGAAAATTGCT GACAACGCTTTGTGTCAATTTGGCACGGAATATGAT TTCTCCTGTTTAGGTAATGCCATCTCAAAAAATGAC AGCAAAGCGATTAAAGAAAGAGACCGAAAAAGTAGAG AAGCTGTTGGCCAAGGTTCTGACAGAGAACTTGCCA GACGGTCTGCGTAAAGTCAACGATATTAACAGCGCG GCTTTTATTTCAGGACACACTGACATCATTCGTCCAG GACGATGCTGACAAACGTGTGTTAATTCAAGAGTTA AAGGGCAAACTGTGTTAATGCAACGCTTTTTAACA ACCCGGATTACTGCATTGACTGTATGGCTCCCTGAC CGGGTGTTTGAGAACTTCAACATTTTTATCGAAAAT GCTGAAAAGATGCGCATCTTGCTCGACTCACCATTG AATGAAAAGATCATGAAGTTCGATCCGGATGCTGAA CAATACGCGAGTTTGGAATCTATGGTCAATGTCTG TCCCAGAAGGATATTGATTCGTACAACCTCATCATT TCCGGGATTTATGCCGATGATGAGGTCAAGAACCCA GGTATCAATGAAATTGTTAAGGAATACAACCAGCAA ATTCGCGGGGATAAGGATGAGTCACCTTTACCTAAA CTGAAAAAGTTGCATAAACAAATTTTGATGCCTGTC

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	EDVKIRGNRHNTLIVKNY FNAWTVFRNLIRILRRKS EAEIDSDFYDVLDDSVFV LSLTYKGENLCRSYITKK IGSDLKPEIATYGSALRP NSRWSPGEKFNVKFHTI VRRDGRLLYYFILPKGAKP VELEDMDGDI ECLQMRKI PNPTIFLPLVFKDPEAF FRDNPEADEFVFLSGMKA PVTITRETYEAYRYKLYT VGKLRDGEVSEEEYKRAL LQVLTAYKEFLENRMIYA DLNFGFKDLEEYKDSSEF IKQVETHNTFMCWAKVSS SQLDDLKSGNGLLFEIW SERLESYYKYGNEKVLRG YEGVLLSILKDENLVSMR TLLNSRPM LVYRPKESSK PMVVHRDGSRVVDRFDKD GKYIPPEVHDELYRFFNN LLIKEKLGEKARKILDNK KVKVKVLESERVKWSKFY DEQFAVTF SVKKNADCLD TTKDLNAEVMEQYSESNR LILIRNTTDILYYLVLDK NGKVLKQSRSLNIINDGAR DVDWKERFRQVTKDRNEG YNEWDSRTSNDLKEVYL NYALKEIAEAVIEYNAIL IIEKMSNAFKDKYSFLDD VTFKGFETKLLAKLSDLH FRGIKDGEPCSF TNPLQL CQND SNKILQDGVIFMVP NSMTRSLDPDTGFIFAIN DHNIRTKKAKLNFLSKFD QLKVSSEGCLIMKYSGDS LPTHNTDN RVWNCCCNHP ITNYDRET KKVEFIEEPV EELSRVLEENGIETDTEL NKLNERENVP GKVVDAIY SLVLNLYLRGT VSGVAGQR AVYYSPVTGKKYDISFIQ AMNLNRKCDYYRIGSKER GEWTD FVAQLINAAAKRP AATKKAGQAKKKKASGSG AGSPKKKRKVEDPKKKRK V (SEQ ID NO: 81)	GAGAAGGCATTTTTTCGTTCCGGTACTCAGTAATGAT TCTGATGCTCGTTCAATTTTAGAAAAAATCTTGAAG GATACTGAGATGTTGCCTTCTAAGATCATTGAAGCG ATGAAAGAAGCAGACGCTGGGGACATCGCTGTATAT GGTTCACGTTTGCACGAGTTAAGCCACGTAATCTAT GGCGATCACGGGAAGCTCTCTCAGATTATCTATGAT AAGGAGTCGAAACGCATCAGCGAGCTCATGGAAACG TTATCGCCTAAGGAGCGCAAAGAGTCAAAGAAACGC TTGGAGGGTCTGGAAGAACATATCCGGAAAGTCGACA TATACCTTCGACGAGCTTAATCGTTATGCGGAAAAG AACGTCATGGCTGCCTACATCGCGGCCGTGGAGGAA AGCTGCGCCGAAATTATGCGTAAGGAGAAGGACTTA CGCACGCTTCTTAGTAAGGAGGATGTCAAGATTCTGT GGTAATCGCCACAATACGTTAATTGTTAAGAACTAC TTCAATGCCTGGACTGTCTTCCGGAATTTGATCCGC ATCCTCCGGCGGAAATCCGAGGCGGAGATCGACTCA GATTTCTATGACGTCTTGGATGACTCTGTGGAAGTT TTATCGCTCACATATAAAGGTGAAAACCTTGTCGCCG TCTTACATTACGAAGAAGATCGGGAGCGATTTAAAG CCAGAGATTGCTACCTATGGTTCCGCCTTGCGCCCT AATTCACGGTGGTGGTCACCGGGCGAGAAGTTTAAC GTAAAGTTCCACACCATTGTTCCCGGGACGGTCGC CTTTATTATTTTCATCTTGCCGAAAGGTGCCAAACCT GTCGAGCTCGAAGATATGGATGGGGACATCGAATGC TTGCAAATGCGCAAGATTCCGAATCCGACTATTTTC CTTCCAAAATTGGTTTTCAAGGACCCAGAGGCCCTTC TTCCGCGACAATCCAGAGGCAGATGAATTCGTTTTT CTTTCGGGTATGAAAGCTCCAGTGACCATCACGCGT GAAACCTATGAGGCGTATCGCTACAACTTTATACA GTTGGGAAGTTACGCGACGGTGAAGTGAGCGAAGAA GAGTATAAACGTGCGTTGTTACAAGTATTGACCGCC TATAAGGAATTCTTAGAGAATCGGATGATCTACGCA GATCTGAACTTTGGCTTTAAAGATCTCGAAGAATAC AAAGACTCGTCAGAATTTATCAAACAAGTCGAAACT CACAACTTTTATGTGCTGGGCTAAGGTCAGTAGC AGTCAGCTCGACGACCTGGTCAAGAGCGGGAACGGG TACTGTTCGAAATCTGGTCAAGACGGTTGGAGTCC TATTACAAATATGGCAACGAGAAGGTGCTGCGTGGG TACGAGGGCGTTCTTTTGAGTATCCTTAAGGACGAG AACCTCGTGAGCATGCGGACGCTGCTTAATTCTCGG CCGATGCTCGTCTACCGCCCTAAAGAATCATCCAAG CCGATGGTTCGTTACCGGGACGGTAGCCGCGTCGTT GATCGGTTTCGATAAGGATGGGAAGTATATCCACCA GAGGTACACGACGAATTATACCGGTTCTTTAACAAT TTGCTTATTAAGGAAAAGCTCGGCGAGAAAGCGCGC AAAATTTTAGACAACAAAAAAGTAAAAGTAAAGGTA TTGGAATCTGAACGTGTAAAGTGGTCAAAGTTTTAT GATGAACAGTTTGCAGTTACATTCTCTGTTAAAAAG AATGCAGACTGTCTGGATACCACGAAAGATCTCAAT GCCGAAGTTATGGAGCAGTATTCGAATCGAACCGG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		CTTATCCTGATCCGCAATACCACTGACATCTTGTAT TATCTTGTACTTGATAAGAATGGGAAAGTGCTGAAA CAACGCTCATTGAATATCATTAACGACGGGGCTCGC GACGTTGATTGGAAAGAGCGTTTTTCGGCAGGTAACA AAAGATCGTAACGAAGGCTATAACGAGTGGGACTAC TCGCGGACTAGCAACGATTTGAAAGAGGTCTATCTG AATTATGCATTGAAGGAGATTGCCGAAGCGGTAATC GAATACAACGCAATTTTTGATTATTGAAAAAATGTCTG AATGCCTTCAAGGATAAGTACTCCTTTTTTGGATGAT GTTACCTTCAAAGGTTTTGAGACCAAACCTTCTTGCG AAGCTCTCTGACTTGCAATTTCCGGGGTATTAAAGAT GGGGAGCCATGTTTCGTTTACGAACCCGTTACAGTTA TGTGAGAACGACTCAAACAAAATTTTACAAGACGGT GTGATTTTTCATGGTCCCTAACAGCATGACGCGCAGT CTGGACCCTGACACTGGGTTTCATTTTTTTCGATTAAAC GATCACAACATCCGCACTAAGAAAGCGAAGTTAAAC TTCCTTAGTAAATTCGATCAGCTGAAAGTGTCATCA GAGGGCTGTTTAATCATGAAATATTCGGGGGACTCC CTTCTACACACAACACAGATAATCGTGTATGGAAC TGTGTTGCAATCACCCGATCACCAACTACGACCGC GAGACGAAAAAGGTCGAATTCATCGAGGAGCCAGTG GAAGAGTTGAGTCGCGTCTTAGAAGAGAATGGGATT GAGACAGATACGGAACCTTAACAAGCTTAACGAGCGC GAGAATGTTCCGGGCAAGGTAGTAGATGCCATCTAT TCTCTGGTGTGTAATTACTTGCGTGGTACCGTGTCC GCGGTTGCAGGCCAACGGGCGGTCTACTATTTCCCT GTGACGGGGAAAAAATATGATATTTTCGTTTATCCAA GCAATGAATCTGAATCGTAAGTGCATTACTACCGG ATCGGGAGCAAAGAACGCGGCGAATGGACGGATTTT GTAGCGCAGTTAATTAACGCGGCCGCAAAAAGGCCG GCGGCCACGAAAAAGGCCGCCAGGCAAAAAGAGAA AAGGCTAGCGGCAGCGGCGCCGGATCCCCAAAGAAG AAAAGGAAGGTTGAAGACCCCAAGAAAAAGAGGAAG GTGTGATAA (SEQ ID NO: 82)
ABW8	MGHHHHHHSSGLVPRGSG TMCYDLNNIKTKLREREV ETMGNNMDNSFEPFIGGN SVSKTLRNELRVGSEYTG KHIKECAIIAEDAVKAEN QYIVKEMMDDFYRDFINR KLDALQGINWEQLFDIMK KAKLDSNKNVSKELDKIQ ESTRKEIGKIFSSDPIYK DMLKADMISKILPEYIVD KYGDAASRIEAVKVFGF SGYFIDFWASRKNVFSK NIASAIPHRIVNVNARIH LDNITAFNRIAEIAGDEV AGIAEDACAYLQNSLED VFTGACYGEFICQKDIDR	ATGGGCCACCATCATCATCATAGCAGCGGCCTG GTGCCGCGCGGCAGCGGTACCATGTGCTACGACTTA AACAACATCAAGACAAAGTTACGTGAACGCGAAGTC GAAACTATGGGCAATAACATGGATAATAGCTTCGAG CCTTTTATTGGCGGTAATAGTGTCTCTAAAACACTT CGGAATGAGCTGCGTGTAGGTTCCGAATATACTGGT AAACACATTAAAGAGTGCGCGATCATTCGAGAGGAC GCCGTGAAGGCGGAGAACCAGTACATCGTAAAAGAG ATGATGGACGACTTTTACCGTGACTTCATTAATCGC AAACCTTGACGCCTTGCAAGGTATTAATTGGGAGCAG CTTTTTTCGATTATGAAGAAGGCGAAATTGGATAAG TCGAATAAAGTCAGCAAAGAGTTAGACAAGATTCAA GAGTCTACGCGGAAAGAAATCGGGAAAATCTTCTCA TCCGATCCAATCTATAAAGACATGCTCAAAGCGGAC ATGATCAGCAAATTTCTGCCAGAGTATATTGTGCGAC AAATACGGTGATGCAGCCTCGCGGATCGAAGCTGTA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	YNNICGVINQHNMNQYCQN KKISRSKFKMERLHKQIL CRSESGFEIPIGFQTDGE VIDAINSFSSTILEEKDIL DRLRTLSEQEVTGYDMERI YVSSKAFESVSKYIDHKW DVIASSMYNYFSGAVRGK DDKKDVKIQTTEIKKIKSC SLLDLKKLVDMYYKMDGM CLEHEATEYVAGITEILV DFNYKTFDMDDSVKMIQN EHMINEIKEYLDITYMSIY HWAKDFMIDELVDRDMEF YSELDEIYYDLSDIVPLY NKVRNYVTQKPYSQDKIK LNFGSPTLANGWSKSKEF DNNVVVLLRDEKIYLAIL NVGNKPSKDIMAGEDRRR SDTDYKKMNYLLPGASK TLPHVFISNAWKSHGI PDEIMYGYNQNKHLKSSP NFDLEFCRKLIDYYKECI DSYPNYQIFNFKFAATET YNDISEFYKDVERQGYKI EWSYISEDINQMDRDGQ IYLFQIYNKDFAPNSKGM QNLHTLYLKNIFSEENLS DVVIKLNGEAELFFRKSS IQHKRGHKKGSVLVNKTY KTTEKTENGQGEIEVIES VPDQCYLELVKYWSEGGV GQLSEEASKYKDKVSHYA ATMDIVKDRRYTEDKFFI HMPITINFKADNRNNVNE KVLKFIAENDDLHVIGID RGERNLLYVSVIDSRGRI VEQKSFNIVENYESSKNV IRRHDYRGKLVNKEHYRN EARKSWKEIGKIKEIKEG YLSQVIHEISKLVLYNA IIVMEDLNYGFKRGRFKV ERQVYQKFETMLINKLAY LVDKSRVDEPGGLLKGY QLTYVPDNLGELGSQCGI IFYVPAAYTSKIDPVTGF VDVDFDKAYSNAEARLDF INKLDCIRYDAPRNKFEI AFDYGNFRTHTTLAKTS WTIFIHGDRIKKERGSYG WKDEIIDIEARIRKLFED	AAGGTGTTTTACGGCTTTTCGGGTATTTTATCGAC TTCTGGGCATCGCGCAAGAACGTCTTCTCAGATAAG AACATCGCGTCGGCCATTCCGCACCGGATTGTCAAT GTGAACGCTCGGATCCATCTGGACAACATCACGGCC TTCAACCGTATCGCAGAAATTGCAGGGGATGAAGTC GCCGGCATTGCTGAAGATGCTTGTGCTTACCTGCAG AATATGAGCTTAGAGGATGTATTACGGGGGCTGC TACGGTGAGTTCATCTGTCAGAAGGATATTGATCGT TACAATAACATTTGCGGTGTTATCAACCAGCACATG AATCAATACTGCCAAAACAAAAGATCTCACGCTCA AAATTTAAGATGGAACGTCTGCACAAACAGATCTTA TGTCGCTCTGAGAGTGCTTTTGAGATCCCGATTGGG TTTCAAACCGACGGGGAGGTAATCGATGCTATCAAC TCCTTTTCTACGATTCTTGAAGAGAAAGATATCTTG GATCGTCTGCGCACTTTGTCGCAGGAGGTAACAGGT TATGACATGGAGCGTATCTATGTAAGTTCCAAGGCG TTTGAGTCTGTATCAAAGTACATCGATCACAATGG GACGTAATTGCTTCTTCCATGTACAATTACTTTTCT GGGGCTGTTCGTGGGAAGGACGACAAGAAAGATGTC AAGATTCAGACGGAATTAAGAAAGATTAAGTCATGT TCGTTATTGGACCTCAAAAAGCTGGTAGATATGTAT TATAAAATGGATGGGATGTGTTTAGAGCACGAAGCG ACGGAGTACGTGGCAGGTATTACGGAGATCCTGGTT GACTTTAACTATAAGACCTTCGACATGGATGATTCC GTTAAGATGATTCAAAATGAGCACATGATTAATGAA ATTAAAGAATATTTAGATACCTATATGTCTATCTAT CATTGGGCGAAGGACTTTATGATCGATGAGCTCGTA GATCGCGACATGGAATCTACAGTGAGCTCGATGAA ATCTATTATGATTTGTCCGACATCGTACCACTGTAT AATAAAGTCCGCAACTACGTCACGCAAAAACCGTAT TCCCAGGATAAAATCAAGTTAACTTTGGCAGCCCCA ACCTTAGCAAACGGTTGGAGCAAGTCGAAAGAATTT GATAACAACGTTGTAGTATTGTTGCGTGACGAAAAG ATTTATCTGGCCATCTTAAATGTGGGGAATAAACCG TCAAAGGATATCATGGCGGGCGAAGACCGTCGTCGC TCCGATACTGATTACAAGAAAATGAATTACTATCTG CTCCCTGGGGCAAGCAAAAACCTGCCACACGTTTTT ATCTCTTCAAATGCATGGAAGAAATCCCACGGTATC CCTGACGAGATTATGTACGGCTATAACCAAATAAG CATTTAAAATCTTCGCCAAACTTCGACTTAGAGTTT TGTCGCAAGCTGATCGATTATTACAAAGAATGTATT GACAGCTATCCTAACTATCAGATCTTCAATTTCAA TTCGCCGCTACGGAACTTACAACGATATTTTCGGAG TTCTACAAAGATGTTGAACGTCAGGGGTACAAGATT GAATGGTCGTACATTTCCGAGGACGATATTAATCAG ATGGATCGTGACGGCCAGATTTATCTTTTTCAAATC TACAACAAGGATTTTGCCCCAACTCTAAGGGCATG CAGAATTTACATACACTCTATTTAAAAAATATTTTT TCAGAGGAAAACCTCTCTGATGTGCTCATTAACTG AATGGCGAGGCTGAGCTCTTCTTCCGCAAGAGCTCG

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	TDIEYADGHNLIQDINEL ESPIQKKFVGELEFDIIRF TVQLRNSKSEKYDGTETE YDKIISPVMDEEGVFFTT DSYIRADGTELPKDADAN GAYCIALKGLYDVLAVKK YWKEGEKFDRLKLLAITNY NWFDFIQNRRFAAAKRPA ATKKAGQAKKKKASGSGA GSPKKKRKVEDPKKKRKV (SEQ ID NO: 94)	ATCCAACATAAACGCGGTCATAAGAAGGGTAGTGTG TTGGTAAATAAGACCTATAAAACCACAGAAAAAACT GAAAATGGTCAAGGCGAAATTGAAGTAATCGAGAGC GTGCCGGACCAAGTGTACCTGGAGCTTGTTAAGTAC TGGTCAGAGGGTGGTGTAGGTCAAGTGTGTCAGAAGAG GCTTCCAAATACAAAGATAAAGTCAGCCACTACGCT GCAACAATGGATATTGTCAAGGACCGGCGGTACACG GAGGATAAGTTCTTTATTACATGCCGATTACGATT AATTTTAAAGCTGATAACCGGAACAATGTCAACGAG AAAGTGCTGAAGTTTATTGCAGAAAACGATGATCTC CACGTTATTGGTATTGACCGTGGGGAACGTAATCTC CTGTACGTCTCAGTAATTGATTACGTGGGCGTATT GTTGAGCAGAAGTCGTTTAATATTGTTGAGAATTAC GAGAGCAGTAAAAATGTGATCCGCCGCCATGATTAT CGTGGGAAATTAGTAAATAAAGAGCACTATCGTAAT GAGGCACGTAAGAGCTGGAAAGAAATCGGCAAAATC AAGGAGATCAAAGAAGGTTATCTCAGTCAAGTTATC CATGAGATTAGTAAGTTGGTATTAAAGTATAACGCC ATCATCGTGATGGAAGATCTTAATTATGGCTTCAAA CGCGGGCGGTTTAAAGTCGAGCGGCAGGTATACCAG AAGTTCGAGACCATGCTTATTAACAAATTAGCCTAC TTAGTGGACAAATCACGCGCGGTAGACGAACCGGGT GGGTTATTAAAAGGCTACCAGCTGACATACGTGCCA GATAACTTGGGTGAACTGGGGTCCAGTGCGGGATC ATTTTTTATGTGCCAGCAGCATACACTTCGAAAATC GATCCTGTTACGGGCTTTGTAGACGTGTTTGATTTT AAGGCATACTCCAATGCCGAAGCACGTTTAGATTTT ATCAATAAACTGGACTGCATCCGGTATGACGCGCCG CGTAACAAGTTTGAAATTGCTTTGACTACGGTAAC TTCCGGACTCATCATAACAACCTTGCAAAGACTAGC TGGACTATTTTTATTACGGCGACCGTATTAAAAAG GAGCGCGGTTCTTACGGCTGGAAGGACGAAATTATC GATATCGAGGCCCCTATTCTGAAGCTGTTTGAAGAC ACAGACATCGAATACGCCGATGGTCACAATTTGATC GGTGACATTAACGAGCTCGAGAGTCCAATTCAAAG AAATTCTGTTGGTGAGCTGTTTCGACATTATCCGTTTC ACTGTCCAACGCGCAACAGCAAAAGTGAGAAATAT GACGGCACCGAAAAGGAGTATGACAAAATTATTTTCG CCGGTAATGGACGAGGAGGGGGTCTTTTACAACC GACAGTTATATCCGCGCAGATGGTACTGAATTACCT AAAGATGCTGATGCTAACGGGGCCTATTGTATCGCG CTGAAGGGTCTTTACGACGTGCTCGCGGTAAAGAAA TATTGGAAGGAGGGGGAGAAGTTCGATCGGAAGTTA CTTGCCATCACCATTACAAGTGGTTTGATTTTCAAT CAGAATCGTCGTTTCGCGGCCGCAAAAAGGCCGGCG GCCACGAAAAAGGCCGGCCAGGCAAAAAGAAAAAG GCTAGCGGCAGCGCGCCGGATCCCCAAGAAGAAA AGGAAGGTTGAAGACCCCAAGAAAAAGAGGAAGGTG TGATAA (SEQ ID NO: 95)

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
ABW9	MGHSHHHHSSGLVPRGSG TMSDRLDVLTNQYPLSKT LRFELKPVGATADWIRKH NVIRYHNGKLVGKDAIRF QNYKYLKKMLDEMHRFL QQALVLEPNSNQAQELTA LLRAIENNYCNNNDLLAG DYPSLSTDKTIKISNGLS KLTTDLFDKFKFEDWAYQY KEDMPNFWRQDIAELEQK LQVSANAKDQKFYKGIK KLKNKIQKSELKAETHKG LYSPTESLQLEWLVRRG DIKLTYLEIGKENEKLNE LVPLVELKDIHRNFNNFA TYLSGFSKNRENVYSTKF DRRSGYKATSVIARTFEQ NLMFCLGNIAKWHKVTEF INQANNYELLQEHGIDWN KQIAALEHKLVDCLAEFF ALNDFSQTLAQGGIEKYN QVLGIAEIAQGPQTQGL NELINLARQKLSAKRSQ PTLQLLYKQILSKGDKPF IDDFKSDQELIAELNEFV SSQIHGEHGAIKLINHEL ESFINEARAAQQQIYVPK DKLTLSLLLTGSWQAIN QWRYKLFQKQLDKQKQ YSFLAQVERWLATEVEQ QNFYQTEKERQQHKDTQP ANVTSSSDGHSILTAFEQ QVQTLTNTICVAAEKYRQ LSDNLTAIDKQRESESSK GFEQIAVIKTLLDACNEL NHFLARFTVNKKDKLPED RAEFWYKQLQAYIDAFPI YELYNKVRNYLSKKPFST EKVKINFDNSHFLSGWTA DYERHSALLFKFNENYLL GVVNENLSSEEEKLLKV GGEHAKRFIYDFQKIDN SNPPRVFIRSKGSSFAPA VEKYQLPIGDIIDIYDQG KFKTEHKKKNEAEFKDSL VRLIDYFKLGFSRHDSYK HYPFKWKASHQYSDIAEF YAHTASFCYTLKEENINF NVLRELSSAGKVYLFYIY NKDFSKNKRGQGRDNLHT	ATGGGGCATCACCACCACCACCTCGTCGGGTCTT GTTCCACGTGGTTCTGGTACCATGTCTGATCGCCTG GACGTGCTTACTAACCAATACCCATTATCGAAAAC TTGCGCTTCGAATTGAAGCCGGTTGGAGCCACAGCT GACTGGATTTCGCAAACACAACGTTATCCGCTATCAT AATGGTAAACTGGTTGGAAAGGATGCGATCCGTTTT CAAAATTATAAGTATCTGAAGAAAATGCTTGATGAG ATGCATCGCTTATTTCTTCAGCAAGCACTGGTGTTG GAGCCAAAATAGCAACCAGGCGCAGGAGTTGACCGCA CTGCTGCGTGCTATTGAGAATAATTATTGCAACAAC AACGACCTGCTGGCGGGCGATTATCCAGCCTCTCT ACCGATAAGACCATTAAAATCAGCAACGGCCTTAGC AAGCTGACCACGGATCTGTTTCGATAAGAAGTTTCGAA GACTGGGCATACCAATACAAAGAAGATATGCCCAAT TTCTGGCGTCAAGATATTGCGGAATTAGAGCAAAAG CTTCAGGTGAGTGCGAACGCAAAAGATCAAAAGTTC TACAAAGGGATCATCAAGAAGCTGAAGAATAAGATC CAGAAGTCTGAACTGAAAGCGGAAACGCACAAGGGC TTATACTCACCTACGGAGTCACTGCAACTGCTGGAG TGGCTGGTACGTCGTGGCGATATTAACTGACTTAC TTAGAGATTGGTAAAGAGAACGAGAAAATTAAAGAA CTGGTCCCGCTGGTTCGAACTTAAGGACATTCATCGC AATTTCAATAATTTTCGCCACATATCTTTCTGGCTTC AGCAAGAATCGTGAGAAATGTGTACTCAACCAATTT GATCGTCGTTTCGGGTTATAAAGCCACCAGTGTAATC GCACGCACGTTTCGAACAGAATTTAATGTTCTGTCTT GGTAACATTGCCAAGTGGCACAAGGTGACAGAATTC ATCAACCAGGCGAACAATTACGAGCTCCTGCAGGAG CACGGCATCGATTGGAATAAGCAAATTGCCGCGCTG GAACACAACTGGACGTGTGTCTCGCAGAGTTCTTC GCGCTTAATAACTTCTCACAAACCCTTGCAACAACAG GGTATCGAAAAGTATAACCAGGTCTTGGCCGGCATC GCCGAGATTGCAGGCCAACCACAGACCCAGGGCCTG AACGAACTCATTAACCTGGCCCGTCAGAAATTGTCT GCCAAACGCTCACAACTGCCTACGTTGCAACTCCTT TACAAACAAATCTTAAGCAAGGGTGATAAGCCATTC ATCGACGATTTTAAAAGCGACCAAGAGTTGATCGCC GAATTAAATGAGTTTGTAGCAGCCAGATTACCGGA GAGCATGGTGCAATCAAATTAATTAATCACGAACTT GAAAGCTTTATCAATGAAGCCCGTGCAGCGCAGCAA CAGATTTATGTGCCCAAGGACAAGCTTACCGAATTA AGTCTTCTCTTAACGGGCAGTTGGCAAGCTATTAAT CAATGGCGTTACAACTGTTTCGACCAGAAACAGCTG GATAAACAACAGAAACAATATTCAATTAGCCTGGCC CAGGTTGAACGCTGGCTGGCAACTGAGGTTGAGCAA CAAAACCTTCTACCAAACCGAAAAGGAGCGCCAGCAG CATAAAGATACGCAGCCGGCGAACGTCAACCACAGC AGCGATGGACACAGCATTTTAACAGCATTTGAGCAA CAGGTGCAGACCTTATTAACCAACATCTGTGTTGCT GCCGAGAAATATCGCCAATTAAGTGATAATCTCACA

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
	SYWKLLFSAENLKDVVLK LNGQAEIFYRPASLAETK AYTHKKGEVLKHKAYSKV WEALDSPIGTRLSWDDAL KIPSITEKTNHNNQRVVQ YNGQEIGRKAFAIIKNR RYSVDKFLFHCPIITLNF ANGQDNINARVNQFLANN KKINIIIGIDRGEKHLLEY SVINQQGEVLHQESFNTI TNSYQTANGEKRQVVTDY HQKLDMSSEDKRDKARKSW STIENIKELKAGYLSHV HRLAQLIIEFNAIVALE LNHGFKRGRFKIEKQVYQ KFEKALIDKLSYLAFKDR TSCLETGHYLNALFQLTSK FKGFNILGKQSGILFYVN ADYTSTTDPLTGYIKNVY KTYSSVKDSTEFWQRFNS IRYIASENRFEFSYDLAD LKQKSLESKTKQTPAKT QWTVSSHVTRSYNQQT QHELFEVTARIQQLLSKA EISYQHQNLDLIPALASCQ SKALHKELIWLFNISILTM RVTDSKSPSATSSENDLIL SPVAPYFDSRNLNKQLPE NGDANGAYNIARKGIMLL ERIGDFVPEGNKKYPDLL IRNNDWQNFVQRPEMVNK QKKKLVLKLTKEYSNGSLF NDLAFAKAAKRPAATKKA GQAKKKKASGSGAGSPKK KRKVEDPKKKRKV (SEQ ID NO: 107)	GCCATCGATAAACAACGCGAGAGCGAATCAAGTAAG GGATTTCGAGCAAATCGCGGTGATTAAACCTTGCTG GACGCGTGTAACGAGCTGAATCACTTTCTGGCACGC TTCACGGTCAACAAGAAGGACAACTCCCCGAAGAT CGCGCAGAATTTTGGTATGAAAAGTTACAAGCGTAC ATTGACGCGTTTCCGATCTACGAGCTGTATAATAAA GTGCGTAATTACTTAAGCAAGAAGCCGTTTAGCACT GAGAAAGTCAAAATTAATTTTGACAATTCCCATTTC CTGTCCGGTTGGACGGCGGACTATGAGCGTCACAGC GCCTTATTATTCAAATTTAATGAAAATTACCTGCTG GGTGTAGTGAATGAGAACTTAAGCAGCGAGGAAGAA GAAAAGCTGAAGCTCGTGGGCGGCGAAGAATGCGC AAGCGCTTCATTTATGATTTTTCAGAAAATCGACAAC TCAAACCCACCGCGCGTTTTTCATTCTGTAGCAAGGGG TCATCGTTCGCACCTGCGGTGCGAAAAGTATCAGTTA CCGATTGGCGATATCATTGACATTTACGATCAGGGT AAATTTAAGACAGAACAAGAAGAAGATGAGGCC GAGTTTAAAGACAGTCTGGTACGTTTGATCGATTAT TTTAAGCTGGGCTTCTCTCGCCATGACAGCTATAAG CACTACCCATTCAAGTGGAAAGCCAGTCATCAATAT AGCGACATTGCGGAATTTTACGCTCATACCGCCTCA TTTTGTTACACGCTTAAGGAAGAAAACATCAATTTT AACGTTCTGCGTGAGTTGTCGTCGGCGGGCAAAGTA TATCTCTTCGAAATTTACAATAAGGATTTCTCAAAG AACAAGCGCGGCCAAGGACGCGACAACCTTGCATACC AGTTATTGGAAGTTGCTGTTCTCGGCTGAGAACCTG AAGGATGTTGTGCTGAAATTAACGGCCAAGCGGAG ATCTTTTACCGCCCGAGCGTCTTTGGCCGAAACCAAG GCCTACACCCATAAGAAAGGGGAAGTACTGAAACAT AAGGCTTATAGCAAAGTGTGGGAAGCCCTGGATTCT CCCATTGGCACCCGCCTGAGCTGGGACGATGCTTTA AAGATCCCGTCTATTACCGAGAAGACCAATCACAA AATCAGCGTGTTGTCCAGTACAACGGCCAAGAAATT GGCCGCAAAGCGGAGTTCGCTATTATCAAGAACCGC CGTTATTCCGTCGATAAATTCCTCTTTCACTGCCCC ATTACACTCAACTTCAAGGCGAACGGCCAGGACAAC ATTAACGCACGCGTTAATCAATTCCTGGCAAATAAC AAGAAGATCAACATTATTGGAATTGACCGTGGTGAA AAGCATTTACTGTATATCAGCGTGATTAATCAACAA GGCGAAGTCCTGCATCAGGAAAGCTTCAATACAATC ACGAATTCATATCAGACCGCCAATGGCGAGAAACGC CAAGTAGTCACTGACTATCACCAGAAGTTGGACATG AGCGAGGACAAACGCGATAAAGCACGTAAGAGCTGG AGTACAATCGAAAATATCAAAGAGCTGAAGGCGGGG TATCTGAGCCACGTTGTACATCGCCTCGCGCAACTG ATTATCGAATTTAATGCCATTGTTGCGTTGGAAGAT CTTAACCACGGGTTCAAACGCGGACGTTTTAAATC GAAAAGCAAGTGTATCAGAAGTTCGAAAAGGCGCTG ATCGACAAATTGAGCTACTTAGCGTTTAAGGATCGC ACGTCGTGTCTGGAACTGGACATTACTTGAATGCC

	Engineered Amino Acid Sequence	Engineered Nucleotide Sequence
		TTTCAATTAACCTCAAAGTTCAAAGGCTTTAACAAC CTTGGAAGCAATCCGGGATTTTGTCTACGTAAAC GCCGATTACACGAGCACCACGGATCCCTTAACAGGC TATATTAAGAACGTATACAAAACCTACTCCTCGGTG AAGGATTCGACCGAATTTTGGCAGCGCTTTAACTCT ATCCGCTATATTGCGAGCGAGAACCGTTTGAATTT AGCTACGACTTAGCGGACCTGAAACAGAAGTCGCTC GAGAGTAAAACCAAACAGACCCCTCTCGCCAAGACC CAATGGACGGTCTCTAGCCACGTTACCCGTTCTTAT TACAACCAGCAGACGAAGCAACATGAGTTATTTCGAA GTGACAGCGCGCATTTCAGCAATTGCTTAGCAAAGCA GAAATCAGCTATCAACATCAAACGACTTGATCCCT GCGTTAGCATCATGTCAAAGTAAGGCGTTACACAAG GAGTTGATTTGGCTGTTCAACAGCATCCTGACTATG CGCGTCACGGACTCAAGCAAACCGTCCGCGACCTCG GAGAATGATTTTATCCTGAGCCCGGTAGCGCCGTAC TTCGACTCCCGCAATCTGAATAAGCAGCTGCCGGAA AACGGCGACGCGAACGGCGCATACAATATCGCTCGT AAAGGTATCATGCTTCTGGAACGTATCGGGGACTTC GTCCCGGAAGGTAACAAGAAGTACCCCGATTTACTG ATCCGCAATAATGACTGGCAGAATTTTGTACAACGC CCGGAGATGGTGAACAAGCAGAAGAAGAACTCGTG AAGTTGAAAACGGAATACTCTAATGGCAGCCTCTTC AATGATTTGGCGTTTAAAGCCGCAGCTAAGCGCCCC GCCGCGACTAAGAAAGCGGGTCAAGCGAAGAAGAAG AAAGCGTCGGGGTCGGGAGCGGGCAGTCCGAAGAAG AAGCGTAAAGTAGAGGATCCGAAGAAGAAACGCAAA GTATAATAA (SEQ ID NO: 108)

[0219] In some embodiments, nuclease constructs disclosed herein can have a polypeptide sequence having at least 85% homology to the polypeptide represented by SEQ ID NO: 94 (ABW8), 29 (ABW3), 81 (ABW7), 107 (ABW9), 3 (ABW1), 16 (ABW2), 42 (ABW4), 55 (ABW5), and/or 68 (ABW6) of International (PCT) Publication No. WO2021/108324. In some embodiments, nuclease constructs herein can have a polynucleotide sequence at least 85% homologous to the polynucleotide encoding the polypeptide having a polynucleotide represented by SEQ ID NO: 95-104 (ABW8 variants 1-10), 30-39 (ABW3 variants 1-10), 82-91 (ABW7 variants 1-10), 108-117 (ABW9 variants 1-10), 4-13 (ABW1 variants 1-10), 17-26 (ABW2 variants 1-10), 43-52 (ABW4 variants 1-10), 56-65 (ABW5 variants 1-10), and/or 69-78 (ABW6 variants 1-10)) of International (PCT) Publication No. WO2021/108324.

[0220] In some embodiments, nuclease constructs herein having a polypeptide of at least 85% homology to the polypeptide represented SEQ ID NO: 94) of International (PCT) Publication No. WO2021/108324. (ABW8) can have increased activity and/or editing accuracy compared to other nuclease constructs. In some embodiments, nuclease constructs herein having

a polypeptide of at least 85% homology to the polypeptide represented by SEQ ID NO: 94 (ABW8), 29 (ABW3), 81 (ABW7) and/or 107 (ABW9) can have increased enzymatic activity and/or editing efficiency and/or accuracy compared to other nuclease constructs such as control nuclease constructs or native sequence-containing nucleases.

5 **[0221]** In some embodiments, nuclease constructs disclosed herein having a polynucleotide encoding a polypeptide having a polynucleotide of at least 85% homology to a polynucleotide represented by SEQ ID NO: 95-104 (ABW8 variants 1-10) can have increased enzymatic activity and/or editing efficiency and/or accuracy compared to control nuclease constructs or nuclease constructs having native sequences. In some embodiments, nuclease
10 constructs disclosed herein having a polynucleotide encoding a polypeptide of at least 85% homology to a polynucleotide represented by SEQ ID NO: 95-104 (ABW8 variants 1-10), 30-39 (ABW3 variants 1-10) or 82-91 (ABW7 variants 1-10) can have increased activity (*e.g.*, editing and/or efficiency) compared to control nuclease constructs or other nuclease constructs.

[0222] More type V-A Cas nucleases and their corresponding naturally occurring
15 CRISPR-Cas systems can be identified by computational and experimental methods known in the art, *e.g.*, as described in U.S. Patent No. 9,790,490 and Shmakov *et al.* (2015) MOL. CELL, 60: 385. Exemplary computational methods include analysis of putative Cas proteins by homology modeling, structural BLAST, PSI-BLAST, or HHPred, and analysis of putative CRISPR loci by identification of CRISPR arrays. Exemplary experimental methods include *in*
20 *vitro* cleavage assays and in-cell nuclease assays (*e.g.*, the Surveyor assay) as described in Zetsche *et al.* (2015) CELL, 163: 759.

[0223] In certain embodiments, the Cas nuclease directs cleavage of one or both strands at the target locus, such as the target strand (*i.e.*, the strand having the target nucleotide sequence that hybridizes with a single guide nucleic acid or modified dual guide nucleic acids) and/or the
25 non-target strand. In certain embodiments, the Cas nuclease directs cleavage of one or both strands within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 100, 200, 500, or more nucleotides from the first or last nucleotide of the target nucleotide sequence or its complementary sequence. In certain embodiments, the cleavage is staggered, *i.e.* generating sticky ends. In certain
embodiments, the cleavage generates a staggered cut with a 5' overhang. In certain embodiments,
30 the cleavage generates a staggered cut with a 5' overhang of 1 to 5 nucleotides, *e.g.*, of 4 or 5 nucleotides. In certain embodiments, the cleavage site is distant from the PAM, *e.g.*, the cleavage occurs after the 18th nucleotide on the non-target strand and after the 23rd nucleotide on the target strand.

[0224] In certain embodiments, the engineered, non-naturally occurring system of the
35 present invention further comprises the Cas nuclease that a complex comprising the targeter

nucleic acid and the modulator nucleic acid is capable of activating. In other embodiments, the engineered, non-naturally occurring system of the present invention further comprises a Cas protein that is related to the Cas nuclease that a complex comprising the targeter nucleic acid and the modulator nucleic acid is capable of activating. For example, in certain embodiments, the Cas protein comprises an amino acid sequence at least 80% (*e.g.*, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to the Cas nuclease. In certain embodiments, the Cas protein comprises a nuclease-inactive mutant of the Cas nuclease. In certain embodiments, the Cas protein further comprises an effector domain.

10 **[0225]** In certain embodiments, the Cas protein lacks substantially all DNA cleavage activity. Such a Cas protein can be generated by introducing one or more mutations to an active Cas nuclease (*e.g.*, a naturally occurring Cas nuclease). A mutated Cas protein is considered to lack substantially all DNA cleavage activity when the DNA cleavage activity of the protein has at most 25%, at most 10%, at most 5%, at most 1%, at most 0.1%, at most 0.01%, or less of the DNA cleavage activity of the corresponding non-mutated form, for example, nil or negligible as compared with the non-mutated form. Thus, the Cas protein may comprise one or more mutations (*e.g.*, a mutation in the RuvC domain of a type V-A Cas protein) and be used as a generic DNA binding protein with or without fusion to an effector domain. Exemplary mutations include D908A, E993A, and D1263A with reference to the amino acid positions in AsCpf1; 20 D832A, E925A, and D1180A with reference to the amino acid positions in LbCpf1; and D917A, E1006A, and D1255A with reference to the amino acid position numbering of the FnCpf1. More mutations can be designed and generated according to the crystal structure described in Yamano *et al.* (2016) CELL, 165: 949.

[0226] It is understood that the Cas protein, rather than losing nuclease activity to cleave all DNA, may lose the ability to cleave only the target strand or only the non-target strand of a double-stranded DNA, thereby being functional as a nickase (see, Gao *et al.* (2016) CELL RES., 26: 901). Accordingly, in certain embodiments, the Cas nuclease is a Cas nickase. In certain embodiments, the Cas nuclease has the activity to cleave the non-target strand but lacks substantially the activity to cleave the target strand, *e.g.*, by a mutation in the Nuc domain. In 30 certain embodiments, the Cas nuclease has the cleavage activity to cleave the target strand but lacks substantially the activity to cleave the non-target strand.

[0227] In other embodiments, the Cas nuclease has the activity to cleave a double-stranded DNA and result in a double-strand break.

[0228] Cas proteins that lack substantially all DNA cleavage activity or have the ability to cleave only one strand may also be identified from naturally occurring systems. For example, 35

certain naturally occurring CRISPR-Cas systems may retain the ability to bind the target nucleotide sequence but lose entire or partial DNA cleavage activity in eukaryotic (*e.g.*, mammalian or human) cells. Such type V-A proteins are disclosed, for example, in Kim *et al.* (2017) ACS SYNTH. BIOL. 6(7): 1273-82 and Zhang *et al.* (2017) CELL DISCOV. 3:17018.

- 5 **[0229]** The activity of the Cas protein (*e.g.*, Cas nuclease) can be altered, thereby creating an engineered Cas protein. In certain embodiments, the altered activity of the engineered Cas protein comprises increased targeting efficiency and/or decreased off-target binding. While not wishing to be bound by theory, it is hypothesized that off-target binding can be recognized by the Cas protein, for example, by the presence of one or more mismatches between the spacer
- 10 sequence and the target nucleotide sequence, which may affect the stability and/or conformation of the CRISPR-Cas complex. In certain embodiments, the altered activity comprises modified binding, *e.g.*, increased binding to the target locus (*e.g.*, the target strand or the non-target strand) and/or decreased binding to off-target loci. In certain embodiments, the altered activity comprises altered charge in a region of the protein that associates with a single guide nucleic acid
- 15 or dual guide nucleic acids, *e.g.*, modified dual guide nucleic acids as described herein. In certain embodiments, the altered activity of the engineered Cas protein comprises altered charge in a region of the protein that associates with the target strand and/or the non-target strand. In certain embodiments, the altered activity of the engineered Cas protein comprises altered charge in a region of the protein that associates with an off-target locus. The altered charge can include
- 20 decreased positive charge, decreased negative charge, increased positive charge, and increased negative charge. For example, decreased negative charge and increased positive charge may generally strengthen the binding to the nucleic acid(s) whereas decreased positive charge and increased negative charge may weaken the binding to the nucleic acid(s). In certain
- 25 embodiments, the altered activity comprises increased or decreased steric hindrance between the protein and a single guide nucleic acid or dual guide nucleic acids, *e.g.*, modified dual guide nucleic acids as described herein. In certain embodiments, the altered activity comprises increased or decreased steric hindrance between the protein and the target strand and/or the non-target strand. In certain embodiments, the altered activity comprises increased or decreased steric hindrance between the protein and an off-target locus. In certain embodiments, the modification
- 30 or mutation comprises a substitution of Lys, His, Arg, Glu, Asp, Ser, Gly, or Thr. In certain embodiments, the modification or mutation comprises a substitution with Gly, Ala, Ile, Glu, or Asp. In certain embodiments, the modification or mutation comprises an amino acid substitution in the groove between the WED and RuvC domain of the Cas protein (*e.g.*, a type V-A Cas protein).

[0230] In certain embodiments, the altered activity of the engineered Cas protein comprises increased nuclease activity to cleave the target locus. In certain embodiments, the altered activity of the engineered Cas protein comprises decreased nuclease activity to cleave an off-target locus. In certain embodiments, the altered activity of the engineered Cas protein comprises altered helicase kinetics. In certain embodiments, the engineered Cas protein comprises a modification that alters formation of the CRISPR complex.

[0231] In certain embodiments, a protospacer adjacent motif (PAM) or PAM-like motif directs binding of the Cas protein complex to the target locus. Many Cas proteins have PAM specificity. The precise sequence and length requirements for the PAM differ depending on the Cas protein used. PAM sequences are typically 2-5 base pairs in length and are adjacent to (but located on a different strand of target DNA from) the target nucleotide sequence. PAM sequences can be identified using any suitable method, *e.g.*, a method known in the art, such as testing cleavage, targeting, or modification of oligonucleotides having the target nucleotide sequence and different PAM sequences.

[0232] In one embodiment, the Cas protein is MAD7 and the PAM is TTTN, wherein N is A, C, G, or T. In one embodiment, the Cas protein is MAD7 and the PAM is CTTN, wherein N is A, C, G, or T. In another embodiment, the Cas protein is AsCpf1 and the PAM is TTTN, wherein N is A, C, G, or T. In another embodiment, the Cas protein is FnCpf1 and the PAM is 5' TTN, wherein N is A, C, G, or T. PAM sequences for certain other type V-A Cas proteins are disclosed in Zetsche *et al.* (2015) CELL, 163: 759 and U.S. Patent No. 9,982,279. Further, engineering of the PAM Interacting (PI) domain of a Cas protein may allow programming of PAM specificity, improve target site recognition fidelity, and increase the versatility of the engineered, non-naturally occurring system. Exemplary approaches to alter the PAM specificity of Cpf1 is described in Gao *et al.* (2017) NAT. BIOTECHNOL., 35: 789.

[0233] In certain embodiments, the engineered Cas protein comprises a modification that alters the Cas protein specificity in concert with modification to targeting range. Cas mutants can be designed to have increased target specificity as well as accommodating modifications in PAM recognition, for example by choosing mutations that alter PAM specificity (*e.g.*, in the PI domain) and combining those mutations with groove mutations that increase (or if desired, decrease) specificity for the on-target locus versus off-target loci. The Cas modifications described herein can be used to counter loss of specificity resulting from alteration of PAM recognition, enhance gain of specificity resulting from alteration of PAM recognition, counter gain of specificity resulting from alteration of PAM recognition, or enhance loss of specificity resulting from alteration of PAM recognition.

[0234] In certain embodiments, the engineered Cas protein comprises one or more nuclear localization signal (NLS) motifs. In certain embodiments, the engineered Cas protein comprises at least 2 (*e.g.*, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10) NLS motifs. Non-limiting examples of NLS motifs include: the NLS of SV40 large T-antigen, having the amino acid sequence of **PKKKRKV** (SEQ ID NO: 23); the NLS from nucleoplasmin, *e.g.*, the nucleoplasmin bipartite NLS having the amino acid sequence of **KRPAATKKAGOAKKKK** (SEQ ID NO: 24); the c-myc NLS, having the amino acid sequence of **PAAKRVKLD** (SEQ ID NO: 25) or **RQRRNELKRSP** (SEQ ID NO: 26); the hRNPA1 M9 NLS, having the amino acid sequence of **NOSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNOGGY** (SEQ ID NO: 27); the importin- α IBB domain NLS, having the amino acid sequence of **RMRIZFKNKGKDTAELRRRRVEVSVELRKAKKDEQILKRRNV** (SEQ ID NO: 28); the myoma T protein NLS, having the amino acid sequence of **VSRKRPRP** (SEQ ID NO: 793) or **PPKKARED** (SEQ ID NO: 794); the human p53 NLS, having the amino acid sequence of **POPKKKPL** (SEQ ID NO: 31); the mouse c-abl IV NLS, having the amino acid sequence of **SALIKKKKKMAP** (SEQ ID NO: 32); the influenza virus NS1 NLS, having the amino acid sequence of **DRLRR** (SEQ ID NO: 33) or **PKQKKRK** (SEQ ID NO: 34); the hepatitis virus δ antigen NLS, having the amino acid sequence of **RKLKKKIKKL** (SEQ ID NO: 35); the mouse Mx1 protein NLS, having the amino acid sequence of **REKKKFLKRR** (SEQ ID NO: 36); the human poly(ADP-ribose) polymerase NLS, having the amino acid sequence of **KRKGDEVVDGVDEVAKKKSKK** (SEQ ID NO: 37); the human glucocorticoid receptor NLS, having the amino acid sequence of **RKCLOAGMNLEARKTKK** (SEQ ID NO: 38), and synthetic NLS motifs such as **PAAKKKKLD** (SEQ ID NO: 39).

[0235] In general, the one or more NLS motifs are of sufficient strength to drive accumulation of the Cas protein in a detectable amount in the nucleus of a eukaryotic cell. The strength of nuclear localization activity may derive from the number of NLS motif(s) in the Cas protein, the particular NLS motif(s) used, the position(s) of the NLS motif(s), or a combination of these factors. In certain embodiments, the engineered Cas protein comprises at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10) NLS motif(s) at or near the N-terminus (*e.g.*, within at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50, or more amino acids along the polypeptide chain from the N-terminus). In certain embodiments, the engineered Cas protein comprises at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10) NLS motif(s) at or near the C-terminus (*e.g.*, within at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50, or more amino acids along the polypeptide chain from the C-terminus). In certain embodiments, the engineered Cas

protein comprises at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10) NLS motif(s) at or near the C-terminus and at least 1 (*e.g.*, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10) NLS motif(s) at or near the N-terminus. In certain embodiments, the engineered Cas protein comprises one, two, or three NLS motifs at or near the C-terminus. In certain embodiments, the engineered Cas protein comprises one NLS motif at or near the N-terminus and one, two, or three NLS motifs at or near the C-terminus. In certain embodiments, the engineered Cas protein comprises a nucleoplasmin NLS at or near the C-terminus.

[0236] Detection of accumulation in the nucleus may be performed by any suitable technique. For example, a detectable marker may be fused to the nucleic acid-targeting protein, such that location within a cell may be visualized. Cell nuclei may also be isolated from cells, the contents of which may then be analyzed by any suitable process for detecting the protein, such as immunohistochemistry, Western blot, or enzyme activity assay. Accumulation in the nucleus may also be determined indirectly, such as by an assay that detects the effect of the nuclear import of a Cas protein complex (*e.g.*, assay for DNA cleavage or mutation at the target locus, or assay for altered gene expression activity) as compared to a control not exposed to the Cas protein or exposed to a Cas protein lacking one or more of the NLS motifs.

[0237] The Cas protein may comprise a chimeric Cas protein, *e.g.*, a Cas protein having enhanced function by being a chimera. Chimeric Cas proteins may be new Cas proteins containing fragments from more than one naturally occurring Cas proteins or variants thereof. For example, fragments of multiple type V-A Cas homologs (*e.g.*, orthologs) may be fused to form a chimeric Cas protein. In certain embodiments, the chimeric Cas protein comprises fragments of Cpf1 orthologs from multiple species and/or strains.

[0238] In certain embodiments, the Cas protein comprises one or more effector domains. The one or more effector domains may be located at or near the N-terminus of the Cas protein and/or at or near the C-terminus of the Cas protein. In certain embodiments, an effector domain comprised in the Cas protein is a transcriptional activation domain (*e.g.*, VP64), a transcriptional repression domain (*e.g.*, a KRAB domain or an SID domain), an exogenous nuclease domain (*e.g.*, FokI), a deaminase domain (*e.g.*, cytidine deaminase or adenine deaminase), or a reverse transcriptase domain (*e.g.*, a high fidelity reverse transcriptase domain). Other activities of effector domains include but are not limited to methylase activity, demethylase activity, transcription release factor activity, translational initiation activity, translational activation activity, translational repression activity, histone modification (*e.g.*, acetylation or demethylation) activity, single-stranded RNA cleavage activity, double-strand RNA cleavage

activity, single-strand DNA cleavage activity, double-strand DNA cleavage activity, and nucleic acid binding activity.

[0239] In certain embodiments, the Cas protein comprises one or more protein domains that enhance homology-directed repair (HDR) and/or inhibit non-homologous end joining (NHEJ). Exemplary protein domains having such functions are described in Jayavaradhan *et al.* (2019) NAT. COMMUN. 10(1): 2866 and Janssen *et al.* (2019) MOL. THER. NUCLEIC ACIDS 16: 141-54. In certain embodiments, the Cas protein comprises a dominant negative version of p53-binding protein 1 (53BP1), for example, a fragment of 53BP1 comprising a minimum focus forming region (*e.g.*, amino acids 1231-1644 of human 53BP1). In certain embodiments, the Cas protein comprises a motif that is targeted by APC-Cdh1, such as amino acids 1-110 of human Geminin, thereby resulting in degradation of the fusion protein during the HDR non-permissive G1 phase of the cell cycle.

[0240] In certain embodiments, the Cas protein comprises an inducible or controllable domain. Non-limiting examples of inducers or controllers include light, hormones, and small molecule drugs. In certain embodiments, the Cas protein comprises a light inducible or controllable domain. In certain embodiments, the Cas protein comprises a chemically inducible or controllable domain.

[0241] In certain embodiments, the Cas protein comprises a tag protein or peptide for ease of tracking or purification. Non-limiting examples of tag proteins and peptides include fluorescent proteins (*e.g.*, green fluorescent protein (GFP), YFP, RFP, CFP, mCherry, tdTomato), HIS tags (*e.g.*, 6×His tag), hemagglutinin (HA) tag, FLAG tag, and Myc tag.

[0242] In certain embodiments, the Cas protein is conjugated to a non-protein moiety, such as a fluorophore useful for genomic imaging. In certain embodiments, the Cas protein is covalently conjugated to the non-protein moiety. The terms “CRISPR-Associated protein,” “Cas protein,” “Cas,” “CRISPR-Associated nuclease,” and “Cas nuclease” as used herein can include such conjugates despite the presence of one or more non-protein moieties.

II. Methods of Targeting, Editing, and/or Modifying Genomic DNA

[0243] The engineered, non-naturally occurring systems disclosed herein are useful for targeting, editing, and/or modifying a target nucleic acid, such as a DNA (*e.g.*, genomic DNA) in a cell or organism. Accordingly, in one aspect, the present invention provides a method of modifying a target nucleic acid (*e.g.*, DNA) having a target nucleotide sequence, the method comprising contacting the target nucleic acid with the engineered, non-naturally occurring system disclosed herein, for example in sections IA, IA1, IB, or IC, thereby resulting in modification of the target nucleic acid.

[0244] The engineered, non-naturally occurring system can be contacted with the target nucleic acid as a complex. Accordingly, in certain embodiments, the method comprises contacting the target nucleic acid with a modified guide CRISPR-Cas complex, such as a modified dual guide CRISPR-Cas complex comprising (a) a targeter nucleic acid comprising

5 (i) a spacer sequence designed to hybridize with the target nucleotide sequence with a 3' end and (ii) a targeter stem sequence with, optionally, a 5' end; (b) a modulator nucleic acid comprising a modulator stem sequence complementary to the targeter stem sequence with, optionally, a 3' end and a 5' end with, optionally, a 5' sequence, *e.g.*, tail sequence,; and (c) a Cas protein, *e.g.*, a Cas nuclease, wherein the targeter nucleic acid and the modulator nucleic acid are joined in certain

10 embodiments and separate nucleic acids in other embodiments, and wherein one or more nucleotides or internucleotide linkages, such as 1-5 or more than 5 nucleotides or internucleotide linkages, *e.g.*, 1, 2, 3, 4, 5 or more than 5 nucleotides or internucleotide linkages, at or near the 3' end of the targeter nucleotide, at or near the 5' end of the targeter stem sequence (if present), at or near the 3' end of the modulator nucleic acid (if present), at or near the 5' end of the 5'

15 sequence, *e.g.*, tail sequence, or a combination thereof, are modified; thereby resulting in modification of the target nucleic acid. Exemplary modified guide nucleic acid CRISPR-Cas complexes are described in Section IC, using modified guide nucleic acids described in sections IA, IA1, and IB, and nucleases described in section IC, and any of these may be used under suitable conditions. In some cases, the targeter nucleic acid and the modulator nucleic acid form

20 a complex that is capable of activating a Cas nuclease that, in a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. In certain embodiments, the Cas protein comprises an amino acid sequence at least 80% (*e.g.*, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to the Cas nuclease.

25 **[0245]** In certain embodiments, the Cas nuclease is a Type I, II, III, IV, V, or V Cas nuclease. In certain embodiments, the Cas nuclease is a type V-A, type V-C, or type V-D Cas nuclease. In certain embodiments, the Cas nuclease is a type V-A Cas nuclease. In certain embodiments, the target nucleic acid further comprises a cognate PAM positioned relative to the target nucleotide sequence such that (a) the modified dual guide CRISPR-Cas complex binds the

30 target nucleic acid; or (b) the Cas nuclease is activated when the modified guide CRISPR-Cas complex, *e.g.*, modified dual guide CRISPR-Cas complex, binds the target nucleic acid.

[0246] The modified guide CRISPR-Cas complex, *e.g.*, modified dual guide CRISPR-Cas complex may be delivered to a cell by introducing a pre-formed ribonucleoprotein (RNP) complex into the cell. Alternatively, one or more components of the modified guide CRISPR-

35 Cas complex, *e.g.*, modified dual guide CRISPR-Cas complex may be expressed in the cell; it

will be appreciated that segments containing modified nucleotides should be introduced into the cells, but unmodified segments can be expressed in the cell. Exemplary methods of delivery are known in the art and described in, for example, U.S. Patent Nos. 10,113,167, 8,697,359, 10,570,418, 11,125,739, 10,829,787, and 11,118,194, and U.S. Patent Application Publication
5 Nos. 2015/0344912, 2018/0119140, and 2018/0282763.

[0247] It is understood that contacting a DNA (*e.g.*, genomic DNA) in a cell with a modified dual guide CRISPR-Cas complex does not require delivery of all components of the complex into the cell. For examples, one or more of the components may be pre-existing in the cell. In certain embodiments, the cell (or a parental/ancestral cell thereof) has been engineered to
10 express the Cas protein, and the targeter nucleic acid (or a nucleic acid comprising a regulatory element operably linked to a nucleotide sequence encoding the targeter nucleic acid) and the modulator nucleic acid (or a nucleic acid comprising a regulatory element operably linked to a nucleotide sequence encoding the modulator nucleic acid) are delivered into the cell, *e.g.*, where one or the other, or both, contains one or more modified nucleotides or internucleotide linkages
15 at or near the 3' ends, at or near the 5' ends, or at or near both ends. In certain embodiments, the cell (or a parental/ancestral cell thereof) has been engineered to express the modulator nucleic acid, and the Cas protein (or a nucleic acid comprising a regulatory element operably linked to a nucleotide sequence encoding the Cas protein) and the targeter nucleic acid (or a nucleic acid comprising a regulatory element operably linked to a nucleotide sequence encoding the targeter
20 nucleic acid) are delivered into the cell, where the targeter nucleic acid contains one or more modified nucleotides at or near the 3' and/or 5' ends. In certain embodiments, the cell (or a parental/ancestral cell thereof) has been engineered to express the Cas protein and the targeter nucleic acid, and the modulator nucleic acid (or a nucleic acid comprising a regulatory element operably linked to a nucleotide sequence encoding the modulator nucleic acid) is delivered into
25 the cell, where the modulator nucleic acid contains one or more modified nucleotides at or near the 3' and/or 5' ends.

[0248] In certain embodiments, the target DNA is in the genome of a target cell. Accordingly, in another aspect, the present invention provides a cell comprising the non-naturally occurring system or a CRISPR expression system described herein. In certain
30 embodiments, the cell is an immune cell. In certain embodiments, the cell is a T cell. See following paragraph and section IVB for more potential cells to be modified. In addition, the present invention provides a cell whose genome has been modified by the modified dual guide CRISPR-Cas system or complex disclosed herein.

[0249] The target cells can be mitotic or post-mitotic cells from any organism, such as a
35 bacterial cell, an archaeal cell, a cell of a single-cell eukaryotic organism, a plant cell, an algal

cell, *e.g.*, *Botryococcus braunii*, *Chlamydomonas reinhardtii*, *Nannochloropsis gaditana*, *Chlorella pyrenoidosa*, *Sargassum patens* C. Agardh, and the like, a fungal cell (*e.g.*, a yeast cell), an animal cell, a cell from an invertebrate animal (*e.g.*, fruit fly, enidarian, echinoderm, nematode, etc.), a cell from a vertebrate animal (*e.g.*, fish, amphibian, reptile, bird, mammal), a cell from a mammal, a cell from a rodent, or a cell from a human. The types of target cells include but are not limited to a stem cell (*e.g.*, an embryonic stem (ES) cell, an induced pluripotent stem (iPS) cell, a germ cell), a somatic cell (*e.g.*, a fibroblast, a hematopoietic cell, a T lymphocyte (*e.g.*, CD8⁺ T lymphocyte), an NK cell, a neuron, a muscle cell, a bone cell, a hepatocyte, a pancreatic cell), an in vitro or in vivo embryonic cell of an embryo at any stage (*e.g.*, a 1-cell, 2-cell, 4-cell, 8-cell; stage zebrafish embryo). Cells may be from established cell lines or may be primary cells (*i.e.*, cells and cells cultures that have been derived from a subject and allowed to grow in vitro for a limited number of passages of the culture). For example, primary cultures are cultures that may have been passaged within 0 times, 1 time, 2 times, 4 times, 5 times, 10 times, or 15 times, but not enough times to go through the crisis stage. Typically, the primary cell lines of the present invention are maintained for fewer than 10 passages in vitro. If the cells are primary cells, they may be harvested from an individual by any suitable method. For example, leukocytes may be harvested by apheresis, leukocytapheresis, or density gradient separation, while cells from tissues such as skin, muscle, bone marrow, spleen, liver, pancreas, lung, intestine, or stomach can be harvested by biopsy. The harvested cells may be used immediately, or may be stored under frozen conditions with a cryopreservative and thawed at a later time in a manner as commonly known in the art.

A. Ribonucleoprotein (RNP) Delivery and “Cas RNA” Delivery

[0250] The engineered, non-naturally occurring system disclosed herein can be delivered into a cell by suitable methods known in the art, including but not limited to ribonucleoprotein (RNP) delivery and “Cas RNA” delivery described below.

[0251] In certain embodiments, a modified guide CRISPR-Cas complex, *e.g.*, modified dual guide CRISPR-Cas system including a targeter nucleic acid, a modulator nucleic acid, where the 5' and/or 3' ends of one or both of the targeter and modulator nucleic acids contain one or more, *e.g.*, 1-5 or more than 5, such as 1, 2, 3, 4, 5, or more than 5 modified nucleotides or internucleotide linkages, and a Cas protein can be combined into a RNP complex and then delivered into the cell as a pre-formed complex. Exemplary modified guide nucleic acid CRISPR-Cas complexes are described in Section IC, using modified guide nucleic acids described in sections IA, IA1, and IB, and nucleases described in section IC, and any of these may be used under suitable conditions. This method is suitable for active modification of the

genetic or epigenetic information in a cell during a limited time period. For example, where the Cas protein has nuclease activity to modify the genomic DNA of the cell, the nuclease activity only needs to be retained for a period of time to allow DNA cleavage, and prolonged nuclease activity may increase off-targeting. Similarly, certain epigenetic modifications can be maintained in a cell once established and can be inherited by daughter cells.

[0252] A “ribonucleoprotein” or “RNP,” as used herein, includes a complex comprising a nucleoprotein and a ribonucleic acid. A “nucleoprotein” as used herein includes a protein capable of binding a nucleic acid (*e.g.*, RNA, DNA). Where the nucleoprotein binds a ribonucleic acid it is referred to as “ribonucleoprotein.” The interaction between the ribonucleoprotein and the ribonucleic acid may be direct, *e.g.*, by covalent bond, or indirect, *e.g.*, by non-covalent bond (*e.g.*, electrostatic interactions (*e.g.*, ionic bond, hydrogen bond, halogen bond), van der Waals interactions (*e.g.*, dipole-dipole, dipole-induced dipole, London dispersion), ring stacking (pi effects), hydrophobic interactions, and the like). In certain embodiments, the ribonucleoprotein includes an RNA-binding motif non-covalently bound to the ribonucleic acid. For example, positively charged aromatic amino acid residues (*e.g.*, lysine residues) in the RNA-binding motif may form electrostatic interactions with the negative nucleic acid phosphate backbones of the RNA.

[0253] To ensure efficient loading of the Cas protein, the targeter nucleic acid and the modulator nucleic acid can be provided in excess molar amount (*e.g.*, at least 1 fold, at least 1.5 fold, at least 2 fold, at least 3 fold, at least 4 fold, or at least 5 fold) relative to the Cas protein. In certain embodiments, the targeter nucleic acid and the modulator nucleic acid are annealed under suitable conditions prior to complexing with the Cas protein. In other embodiments, the targeter nucleic acid, the modulator nucleic acid, and the Cas protein are directly mixed together to form an RNP.

[0254] A variety of delivery methods can be used to introduce an RNP disclosed herein into a cell. Exemplary delivery methods or vehicles include but are not limited to microinjection, liposomes (see, *e.g.*, U.S. Patent No. 10,829,787) such as molecular trojan horses liposomes that delivers molecules across the blood brain barrier (see, Pardridge *et al.* (2010) COLD SPRING HARB. PROTOC., doi:10.1101/pdb.prot5407), immunoliposomes, virosomes, microvesicles (*e.g.*, exosomes and ARMMs), polycations, lipid:nucleic acid conjugates, electroporation, cell permeable peptides (see, U.S. Patent No. 11,118,194), nanoparticles, nanowires (see, Shalek *et al.* (2012) NANO LETTERS, 12: 6498), exosomes, and perturbation of cell membrane (*e.g.*, by passing cells through a constriction in a microfluidic system, see, U.S. Patent No. 11,125,739). In certain embodiments the delivery method is electroporation. Where the target cell is a

proliferating cell, the efficiency of RNP delivery can be enhanced by cell cycle synchronization (see, U.S. Patent No. 10,570,418).

[0255] In other embodiments, the modified guide CRISPR-CAS system, *e.g.*, modified dual guide CRISPR-Cas system is delivered into a cell in a “Cas RNA” approach, *i.e.*, delivering
5 a targeter nucleic acid, a modulator nucleic acid, where the 5’ and/or 3’ ends of one or both of the targeter and modulator nucleic acids contain one or more, *e.g.*, 1-5 or more than 5, such as 1, 2, 3, 4, 5, or more than 5 modified nucleotides, and an RNA (*e.g.*, messenger RNA (mRNA)) encoding a Cas protein. The RNA encoding the Cas protein can be translated in the cell and form a complex with the targeter nucleic acid and the modulator nucleic acid intracellularly. Similar to
10 the RNP approach, RNAs have limited half-lives in cells, even though stability-increasing modification(s) can be made in one or more of the RNAs. Accordingly, the “Cas RNA” approach is suitable for active modification of the genetic or epigenetic information in a cell during a limited time period, such as DNA cleavage, and has the advantage of reducing off-targeting.

[0256] The mRNA can be produced by transcription of a DNA comprising a regulatory
15 element operably linked to a Cas coding sequence. Given that multiple copies of Cas protein can be generated from one mRNA, the targeter nucleic acid and the modulator nucleic acid are generally provided in excess molar amount (*e.g.*, at least 5 fold, at least 10 fold, at least 20 fold, at least 30 fold, at least 50 fold, or at least 100 fold) relative to the mRNA. In certain
20 embodiments, the targeter nucleic acid and the modulator nucleic acid are annealed under suitable conditions prior to delivery into the cells. In other embodiments, the targeter nucleic acid and the modulator nucleic acid are delivered into the cells without annealing *in vitro*. In certain embodiments, a modified dual guide nucleic acid system is used. In certain embodiments, a modified single guide nucleic acid system is used.

[0257] A variety of delivery systems can be used to introduce an “Cas RNA” system into
25 a cell. Non-limiting examples of delivery methods or vehicles include microinjection, biolistic particles, liposomes (see, *e.g.*, U.S. Patent No. 10,829,787) such as molecular trojan horses liposomes that delivers molecules across the blood brain barrier (see, Pardridge *et al.* (2010) COLD SPRING HARB. PROTOC., doi:10.1101/pdb.prot5407), immunoliposomes, virosomes, polycations, lipid:nucleic acid conjugates, electroporation, nanoparticles, nanowires (see, Shalek
30 *et al.* (2012) NANO LETTERS, 12: 6498), exosomes, and perturbation of cell membrane (*e.g.*, by passing cells through a constriction in a microfluidic system, see, U.S. Patent No. 11,125,739). Specific examples of the “nucleic acid only” approach by electroporation are described in International (PCT) Publication No. WO2016/164356.

[0258] In other embodiments, the modified guide CRISPR-Cas system, *e.g.*, modified
35 dual guide CRISPR-Cas system is delivered into a cell in the form of a targeter nucleic acid, a

modulator nucleic acid, where the 5' and/or 3' ends of one or both of the targeter and modulator nucleic acids contain one or more, *e.g.*, 1-5 or more than 5, such as 1, 2, 3, 4, 5, or more than 5 modified nucleotides, and a DNA comprising a regulatory element operably linked to a Cas coding sequence. The DNA can be provided in a plasmid, viral vector, or any other form described in the "CRISPR Expression Systems" subsection. Such delivery method may result in constitutive expression of Cas protein in the target cell (*e.g.*, if the DNA is maintained in the cell in an episomal vector or is integrated into the genome), and may increase the risk of off-targeting which is undesirable when the Cas protein has nuclease activity. Notwithstanding, this approach is useful when the Cas protein comprises a non-nuclease effector (*e.g.*, a transcriptional activator or repressor). It is also useful for research purposes and for genome editing of plants.

B. CRISPR Expression Systems

[0259] In another aspect, the present invention provides a CRISPR expression system comprising: either (a) a nucleic acid comprising a first regulatory element operably linked to a nucleotide sequence encoding a targeter nucleic acid disclosed herein comprising (i) a spacer sequence designed to hybridize with a target nucleotide sequence and (ii) a targeter stem sequence; or (b) a nucleic acid comprising a second regulatory element operably linked to a nucleotide sequence encoding a modulator nucleic acid disclosed herein comprising a modulator stem sequence complementary to the targeter stem sequence, wherein a complex comprising the targeter nucleic acid and a modified modulator nucleic acid, as described herein, or a complex comprising a modified targeter nucleic acid, as described herein, and the modulator nucleic acid is capable of activating a Cas nuclease that, in a naturally occurring system, is activated by a single crRNA in the absence of a tracrRNA. Exemplary targeter and modulator nucleic acids are described in sections IA, IA1, and IB.

[0260] In certain embodiments, the CRISPR expression system further comprises (c) a nucleic acid comprising a third regulatory element operably linked to a nucleotide sequence encoding a Cas protein disclosed herein. In certain embodiments, the Cas protein comprises an amino acid sequence at least 80% (*e.g.*, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to the Cas nuclease, thereby resulting in modification of the target nucleic acid (*e.g.*, DNA). In certain embodiments, the Cas protein and the Cas nuclease are identical, and the method results in cleavage of the target nucleic acid. In certain embodiments, the Cas nuclease is a Type I, II, III, IV, V, or VI nuclease. In certain embodiments the Cas nuclease is a Type V nuclease. In certain embodiments, the Cas nuclease is a type V-A, type V-C, or type V-D Cas

nuclease. In certain embodiments, the Cas nuclease is a type V-A Cas nuclease. Exemplary Cas Nucleases are described in section IC.

[0261] As used in this context, the term “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell).

[0262] The form of elements (a), (b), and (c) of the CRISPR expression system described above may be independently selected from various nucleic acids such as DNA (*e.g.*, modified DNA) and RNA (*e.g.*, modified RNA). In certain embodiments, elements (a) and (b) are each in the form of DNA. In certain embodiments, the CRISPR expression system further comprises element (c) in the form of DNA. The third regulatory element can be a constitutive or inducible promoter that drives the expression of the Cas protein. In other embodiments, the CRISPR expression system further comprises element (c) in the form of RNA (*e.g.*, mRNA).

[0263] Elements (a), (b), and/or (c) can be provided in one or more vectors. The term “vector,” as used herein, can include a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids in cells, such as prokaryotic cells, eukaryotic cells, mammalian cells, or target tissues. Non-viral vector delivery systems include DNA plasmids, RNA (*e.g.*, a transcript of a vector described herein), naked nucleic acid, and nucleic acid complexed with a delivery vehicle, such as a liposome. Viral vector delivery systems include DNA and RNA viruses, which have either episomal or integrated genomes after delivery to the cell. Gene therapy procedures are known in the art and disclosed in Van Brunt (1988) BIOTECHNOLOGY, 6: 1149; Anderson (1992) SCIENCE, 256: 808; Nabel & Feigner (1993) TIBTECH, 11: 211; Mitani & Caskey (1993) TIBTECH, 11: 162; Dillon (1993) TIBTECH, 11: 167; Miller (1992) NATURE, 357: 455; Vigne, (1995) RESTORATIVE NEUROLOGY AND NEUROSCIENCE, 8: 35; Kremer & Perricaudet (1995) BRITISH MEDICAL BULLETIN, 51: 31; Haddada *et al.* (1995) CURRENT TOPICS IN MICROBIOLOGY AND IMMUNOLOGY, 199: 297; Yu *et al.* (1994) GENE THERAPY, 1: 13; and Doerfler and Bohm (Eds.) (2012) The Molecular Repertoire of Adenoviruses II: Molecular Biology of Virus-Cell Interactions. In certain embodiments, at least one of the vectors is a DNA plasmid. In certain embodiments, at least one of the vectors is a viral vector (*e.g.*, retrovirus, adenovirus, or adeno-associated virus).

[0264] Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors and replication defective viral vectors) do not autonomously replicate in the host cell. Certain vectors, however,

may be integrated into the genome of the host cell and thereby are replicated along with the host genome. A skilled person in the art will appreciate that different vectors may be suitable for different delivery methods and have different host tropism, and will be able to select one or more vectors suitable for the use.

- 5 **[0265]** The term “regulatory element,” as used herein, can include a transcriptional and/or translational control sequence, such as a promoter, enhancer, transcription termination signal (*e.g.*, polyadenylation signal), internal ribosomal entry sites (IRES), protein degradation signal, and the like, that provide for and/or regulate transcription of a non-coding sequence (*e.g.*, a targeter nucleic acid or a modulator nucleic acid) or a coding sequence (*e.g.*, a Cas protein)
- 10 and/or regulate translation of an encoded polypeptide. Such regulatory elements are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY, 185, Academic Press, San Diego, Calif. (1990). Regulatory elements include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory
- 15 sequences). A tissue-specific promoter may direct expression primarily in a desired tissue of interest, such as muscle, neuron, bone, skin, blood, specific organs (*e.g.*, liver, pancreas), or particular cell types (*e.g.*, lymphocytes). Regulatory elements may also direct expression in a temporal-dependent manner, such as in a cell-cycle dependent or developmental stage-dependent manner, which may or may not also be tissue or cell-type specific. In certain embodiments, a
- 20 vector comprises one or more pol III promoter (*e.g.*, 1, 2, 3, 4, 5, or more pol III promoters), one or more pol II promoters (*e.g.*, 1, 2, 3, 4, 5, or more pol II promoters), one or more pol I promoters (*e.g.*, 1, 2, 3, 4, 5, or more pol I promoters), or combinations thereof. Examples of pol III promoters include, but are not limited to, U6 and H1 promoters. Examples of pol II promoters include, but are not limited to, the retroviral Rous sarcoma virus (RSV) LTR promoter
- 25 (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer), the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter. Also encompassed by the term “regulatory element” are enhancer elements, such as WPRE; CMV enhancers; the R-U5' segment in LTR of HTLV-I (see, Takebe *et al.* (1988) MOL. CELL. BIOL., 8: 466); SV40
- 30 enhancer; and the intron sequence between exons 2 and 3 of rabbit β -globin (see, O'Hare *et al.* (1981) PROC. NATL. ACAD. SCI. USA., 78: 1527). It will be appreciated by those skilled in the art that the design of the expression vector can depend on factors such as the choice of the host cell to be transformed, the level of expression desired, etc. A vector can be introduced into host cells to produce transcripts, proteins, or peptides, including fusion proteins or peptides, encoded by

nucleic acids as described herein (*e.g.*, CRISPR transcripts, proteins, enzymes, mutant forms thereof, or fusion proteins thereof).

[0266] In certain embodiments, the nucleotide sequence encoding the Cas protein is codon optimized for expression in a eukaryotic host cell, *e.g.*, a yeast cell, a mammalian cell (5 *e.g.*, a mouse cell, a rat cell, or a human cell), or a plant cell. Various species exhibit particular bias for certain codons of a particular amino acid. Codon bias (differences in codon usage between organisms) often correlates with the efficiency of translation of messenger RNA (mRNA), which is in turn believed to be dependent on, among other things, the properties of the codons being translated and the availability of particular transfer RNA (tRNA) molecules. The (10 predominance of selected tRNAs in a cell is generally a reflection of the codons used most frequently in peptide synthesis. Accordingly, genes can be tailored for optimal gene expression in a given organism based on codon optimization. Codon usage tables are readily available, for example, at the “Codon Usage Database” available at kazusa.or.jp/codon/ and these tables can be adapted in a number of ways (see, Nakamura *et al.* (2000) NUCL. ACIDS RES., 28: 292).

15 Computer algorithms for codon optimizing a particular sequence for expression in a particular host cell, such as Gene Forge (Aptagen; Jacobus, Pa.), are also available. In certain embodiments, the codon optimization facilitates or improves expression of the Cas protein in the host cell.

20 **C. Donor Templates**

[0267] Cleavage of a target nucleotide sequence in the genome of a cell by the modified dual guide CRISPR-Cas system or complex disclosed herein can activate the DNA damage pathways, which may rejoin the cleaved DNA fragments by NHEJ or HDR. HDR requires a repair template, either endogenous or exogenous, to transfer the sequence information from the (25 repair template to the target.

[0268] In certain embodiments, the engineered, non-naturally occurring system or CRISPR expression system further comprises a donor template. As used herein, the term “donor template” can include a nucleic acid designed to serve as a repair template at or near the target nucleotide sequence upon introduction into a cell or organism. In certain embodiments, the donor (30 template is complementary to a polynucleotide comprising the target nucleotide sequence or a portion thereof. When optimally aligned, a donor template may overlap with one or more nucleotides of a target nucleotide sequences (*e.g.*, at least 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 100, 500 or more nucleotides). The nucleotide sequence of the donor template is typically not identical to the genomic sequence that it replaces. Rather, the donor template may contain one or (35 more substitutions, insertions, deletions, inversions or rearrangements with respect to the

genomic sequence, so long as sufficient homology is present to support homology-directed repair. In certain embodiments, the donor template comprises a non-homologous sequence flanked by two regions of homology (*i.e.*, homology arms), such that homology-directed repair between the target DNA region and the two flanking sequences results in insertion of the non-homologous sequence at the target region. In certain embodiments, the donor template comprises a non-homologous sequence of 10-100 nucleotides, 50-500 nucleotides, 100-1,000 nucleotides, 200-2,000 nucleotides, or 500-5,000 nucleotides in length positioned between two homology arms. In certain embodiments the donor template comprises a single stranded oligodeoxynucleotide (ssODN).

[0269] Generally, the homologous region(s) of a donor template has at least 50% sequence identity to a genomic sequence with which recombination is desired. The homology arms are designed or selected such that they are capable of recombining with the nucleotide sequences flanking the target nucleotide sequence under intracellular conditions. In certain embodiments, where HDR of the non-target strand is desired, the donor template comprises a first homology arm homologous to a sequence 5' to the target nucleotide sequence and a second homology arm homologous to a sequence 3' to the target nucleotide sequence. In certain embodiments, the first homology arm is at least 50% (*e.g.*, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) identical to a sequence 5' to the target nucleotide sequence. In certain embodiments, the second homology arm is at least 50% (*e.g.*, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) identical to a sequence 3' to the target nucleotide sequence. In certain embodiments, when the donor template sequence and a polynucleotide comprising a target nucleotide sequence are optimally aligned, the nearest nucleotide of the donor template is within at least 5, 10, 15, 20, 25, 50, 75, 100, 200, 300, 400, 500, 1000, 2000, 3000, 4000, or more nucleotides from the target nucleotide sequence.

[0270] In certain embodiments, the donor template further comprises an engineered sequence not homologous to the sequence to be repaired. Such engineered sequence can harbor a barcode and/or a sequence capable of hybridizing with a donor template-recruiting sequence disclosed herein.

[0271] In certain embodiments, the donor template further comprises one or more mutations relative to the genomic sequence, wherein the one or more mutations reduce or prevent cleavage, by the same CRISPR-Cas system, of the donor template or of a modified genomic sequence with at least a portion of the donor template sequence incorporated. In certain

embodiments, in the donor template, the PAM adjacent to the target nucleotide sequence and recognized by the Cas nuclease is mutated to a sequence not recognized by the same Cas nuclease. In certain embodiments, in the donor template, the target nucleotide sequence (*e.g.*, the seed region) is mutated. In certain embodiments, the one or more mutations are silent with respect to the reading frame of a protein-coding sequence encompassing the mutated sites.

[0272] The donor template can be provided to the cell as single-stranded DNA, single-stranded RNA, double-stranded DNA, or double-stranded RNA. It is understood that the modified dual guide CRISPR-Cas system disclosed herein may possess nuclease activity to cleave the target strand, the non-target strand, or both. When HDR of the target strand is desired, a donor template having a nucleic acid sequence complementary to the target strand is also contemplated.

[0273] The donor template can be introduced into a cell in linear or circular form. If introduced in linear form, the ends of the donor template may be protected (*e.g.*, from exonucleolytic degradation) by methods known to those of skill in the art. For example, one or more dideoxynucleotide residues are added to the 3' terminus of a linear molecule and/or self-complementary oligonucleotides are ligated to one or both ends (see, for example, Chang *et al.* (1987) PROC. NATL. ACAD. SCI. USA, 84: 4959; Nehls *et al.* (1996) SCIENCE, 272: 886; see also the chemical modifications for increasing stability and/or specificity of RNA disclosed *supra*). Additional methods for protecting exogenous polynucleotides from degradation include, but are not limited to, addition of terminal amino group(s) and the use of modified nucleotides such as, for example, phosphorothioates, phosphoramidates, and O-methyl ribose or deoxyribose residues. As an alternative to protecting the termini of a linear donor template, additional lengths of sequence may be included outside of the regions of homology that can be degraded without impacting recombination.

[0274] A donor template can be a component of a vector as described herein, contained in a separate vector, or provided as a separate polynucleotide, such as an oligonucleotide, linear polynucleotide, or synthetic polynucleotide. In certain embodiments, the donor template is a DNA. In certain embodiments, a donor template is in the same nucleic acid as a sequence encoding the targeter nucleic acid, a sequence encoding the modulator nucleic acid, and/or a sequence encoding the Cas protein, where applicable. In certain embodiments, a donor template is provided in a separate nucleic acid. A donor template polynucleotide may be of any suitable length, such as at least 50, 75, 100, 150, 200, 500, 1000, 2000, 3000, 4000, or more nucleotides in length, for example 50 to 500 nucleotides in length, such as 500 nucleotides in length.

[0275] A donor template can be introduced into a cell as an isolated nucleic acid.

Alternatively, a donor template can be introduced into a cell as part of a vector (*e.g.*, a plasmid)

having additional sequences such as, for example, replication origins, promoters and genes encoding antibiotic resistance, that are not intended for insertion into the DNA region of interest. Alternatively, a donor template can be delivered by viruses (*e.g.*, adenovirus, adeno-associated virus (AAV)). In certain embodiments, the donor template is introduced as an AAV, *e.g.*, a pseudotyped AAV. The capsid proteins of the AAV can be selected by a person skilled in the art based upon the tropism of the AAV and the target cell type. For example, in certain embodiments, the donor template is introduced into a hepatocyte as AAV8 or AAV9. In certain embodiments, the donor template is introduced into a hematopoietic stem cell, a hematopoietic progenitor cell, or a T lymphocyte (*e.g.*, CD8⁺ T lymphocyte) as AAV6 or an AAVHSC (see, U.S. Patent No. 9,890,396). It is understood that the sequence of a capsid protein (VP1, VP2, or VP3) may be modified from a wild-type AAV capsid protein, for example, having at least 50% (*e.g.*, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) sequence identity to a wild-type AAV capsid sequence.

[0276] The donor template can be delivered to a cell (*e.g.*, a primary cell) by various delivery methods, such as a viral or non-viral method disclosed herein. In certain embodiments, a non-viral donor template is introduced into the target cell as a naked nucleic acid or in complex with a liposome or poloxamer. In certain embodiments, a non-viral donor template is introduced into the target cell by electroporation. In other embodiments, a viral donor template is introduced into the target cell by infection. The engineered, non-naturally occurring system can be delivered before, after, or simultaneously with the donor template (see, International (PCT) Application Publication No. WO2017/053729). A skilled person in the art can choose proper timing based upon the form of delivery (consider, for example, the time needed for transcription and translation of RNA and protein components) and the half-life of the molecule(s) in the cell. In particular embodiments, where the modified guide CRISPR-Cas system, *e.g.*, modified dual guide CRISPR-Cas system including the Cas protein is delivered by electroporation (*e.g.*, as an RNP), the donor template (*e.g.*, as an AAV) is introduced into the cell within 4 hours (*e.g.*, within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 90, 120, 150, 180, 210, or 240 minutes) after the introduction of the engineered, non-naturally occurring system.

[0277] In certain embodiments, the donor template is conjugated covalently to the modulator nucleic acid. Covalent linkages suitable for this conjugation are known in the art and are described, for example, in U.S. Patent No. 9,982,278 and Savic *et al.* (2018) *eLIFE* 7:e33761. In certain embodiments, the donor template is covalently linked to the modulator nucleic acid (*e.g.*, the 5' end of the modulator nucleic acid) through an internucleotide bond. In certain

embodiments, the donor template is covalently linked to the modulator nucleic acid (*e.g.*, the 5' end of the modulator nucleic acid) through a linker.

[0278] In certain embodiments, the donor template contains an exogenous gene, *e.g.*, CAR cassette, that is integrated at a targeted site in a host cell, *e.g.*, an immune cell, *e.g.*, a T cell, mediated through the action of a nucleic acid-guided nuclease complex, that results in a modified host cell, *e.g.*, a CAR-T cell.

D. Efficiency and Specificity

[0279] The engineered, non-naturally occurring system of the present invention can have the advantage that the efficiency of nucleic acid targeting, cleavage, or modification can be increased or decreased by, for example, adjusting the hybridization of modified guide nucleic acids, *e.g.*, modified dual guide nucleic acids, and the length of the spacer sequence.

[0280] In certain embodiments, the engineered, non-naturally occurring system has high efficiency. For example, in certain embodiments, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% of a population of nucleic acids having the target nucleotide sequence and a cognate PAM, when contacted with the engineered, non-naturally occurring system, is targeted, cleaved, or modified. In certain embodiments, the genomes of at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% of a population of cells, when contacted with the engineered, non-naturally occurring system, are targeted, cleaved, or modified.

[0281] It has been observed that the occurrence of on-target events and the occurrence of off-target events are generally correlated. For certain therapeutic purposes, low on-target efficiency can be tolerated and low off-target frequency is more desirable. For example, when editing or modifying a proliferating cell that will be delivered to a subject and proliferate *in vivo*, tolerance to off-target events is low. Prior to delivery, however, it is possible to assess the on-target and off-target events, thereby selecting one or more colonies that have the desired edit or modification and lack any undesired edit or modification.

[0282] The method disclosed herein is suitable for such use. In certain embodiments, when a population of nucleic acids having the target nucleotide sequence and a cognate PAM is contacted with the engineered, non-naturally occurring system disclosed herein, the frequency of off-target events (*e.g.*, targeting, cleavage, or modification, depending on the function of the

CRISPR-Cas system) is reduced by at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% relative to the frequency of off-target events when using the corresponding CRISPR system containing an unmodified guide nucleic acid, *e.g.*, a guide nucleic acid consisting of the sequences of the targeter and modulator nucleic acids, but without modified nucleotides or internucleotide linkages under the same conditions. In certain cases, the frequency is relative to the frequency of off-target events when using the corresponding CRISPR system containing a single guide nucleic acid (*e.g.*, a single crRNA consisting of the sequences of the targeter and modulator nucleic acids) under the same conditions. In certain embodiments, when genomic DNA having the target nucleotide sequence and a cognate PAM is contacted with the engineered, non-naturally occurring system disclosed herein in a population of cells, the frequency of off-target events (*e.g.*, targeting, cleavage, or modification, depending on the function of the CRISPR-Cas system) is reduced by at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% % relative to the frequency of off-target events when using the corresponding CRISPR system containing an unmodified guide nucleic acid, *e.g.*, a guide nucleic acid consisting of the sequences of the targeter and modulator nucleic acids, but without modified nucleotides or internucleotide linkages under the same conditions. In certain cases, the frequency is relative to the frequency of off-target events when using the corresponding CRISPR system containing a single guide nucleic acid (*e.g.*, a single crRNA consisting of the sequences of the targeter and modulator nucleic acids) under the same conditions. In certain embodiments, when delivered into a population of cells comprising genomic DNA having the target nucleotide sequence and a cognate PAM, the frequency of off-target events (*e.g.*, targeting, cleavage, or modification, depending on the function of the CRISPR-Cas system) in the cells receiving the engineered, non-naturally occurring system disclosed herein is reduced by at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% % relative to the frequency of off-target events when using the corresponding CRISPR system containing an unmodified guide nucleic acid, *e.g.*, a guide nucleic acid consisting of the sequences of the targeter and modulator nucleic acids, but without modified nucleotides or internucleotide linkages under the same conditions. In certain cases, the frequency is relative to the frequency of off-target events when using the corresponding CRISPR system containing a single guide nucleic acid (*e.g.*, a single crRNA consisting of the sequences of the targeter and modulator nucleic acids) under the same conditions. Methods of assessing off-target events were

summarized in Lazzarotto *et al.* (2018) NAT. PROTOC. 13(11): 2615-42, and include discovery of *in situ* Cas off-targets and verification by sequencing (DISCOVER-seq) as disclosed in Wienert *et al.* (2019) SCIENCE 364(6437): 286-89; genome-wide unbiased identification of double-stranded breaks (DSBs) enabled by sequencing (GUIDE-seq) as disclosed in Kleinstiver *et al.* (2016) NAT. BIOTECH. 34: 869-74; circularization for *in vitro* reporting of cleavage effects by sequencing (CIRCLE-seq) as described in Kocak *et al.* (2019) NAT. BIOTECH. 37: 657-66. In certain embodiments, the off-target events include targeting, cleavage, or modification at a given off-target locus (*e.g.*, the locus with the highest occurrence of off-target events detected). In certain embodiments, the off-target events include targeting, cleavage, or modification at all the loci with detectable off-target events, collectively.

E. Multiplex Methods

[0283] The methods of targeting, editing, and/or modifying a genomic DNA disclosed herein can be conducted in multiplicity. For example, a library of targeter nucleic acids can be used to target multiple genomic loci; a library of donor templates can also be used to generate multiple insertions, deletions, and/or substitutions. The multiplex assay can be conducted in a screening method wherein each separate cell culture (*e.g.*, in a well of a 96-well plate or a 384-well plate) is exposed to a different targeter nucleic acid or a different combination of targeter nucleic acid and donor template. The multiplex assay can also be conducted in a selection method wherein a cell culture is exposed to a mixed population of different targeter nucleic acids and/or donor templates, and the cells with desired characteristics (*e.g.*, functionality) are enriched or selected by advantageous survival or growth, resistance to a certain agent, expression of a detectable protein (*e.g.*, a fluorescent protein that is detectable by flow cytometry), *etc.*

[0284] In certain embodiments, the multiplex method employs a plurality of targeter nucleic acids that are capable of hybridizing with different target nucleotide sequences. In certain embodiments, the plurality of targeter nucleic acids comprise a common targeter stem sequence. In certain embodiments, the multiplex method employs a single modulator nucleic acid capable of hybridizing with the plurality of targeter nucleic acids. In certain embodiments, the multiplex method employs a single Cas protein (*e.g.*, Cas nuclease) disclosed herein.

[0285] In certain embodiments, the multiplex method employs a plurality of targeter nucleic acids that are capable of hybridizing with different target nucleotide sequences that are close to or adjacent to different PAMs. In certain embodiments, the plurality of targeter nucleic acids comprise different targeter stem sequences. In certain embodiments, the multiplex method employs a plurality of modulator nucleic acids each capable of hybridizing with a different

targeter nucleic acid. In certain embodiments, the multiplex method employs a plurality of Cas proteins (*e.g.*, Cas nucleases) disclosed herein that have different PAM specificity.

[0286] In certain embodiments, the multiplex method further comprises introducing one or more donor templates into the population of cells. In certain embodiments, the multiplex method employs a plurality of modulator nucleic acids each comprising a different donor template-recruiting sequence, wherein each donor template-recruiting sequence is capable of hybridizing with a different donor template.

[0287] In certain embodiments, the plurality of targeter nucleic acids and/or the plurality of donor templates are designed for saturation editing. For example, in certain embodiments, each nucleotide position in a sequence of interest is systematically modified with each of all four traditional bases, A, T, G and C. In other embodiments, at least one sequence in each gene from a pool of genes of interest is modified, for example, according to a CRISPR design algorithm. In certain embodiments, each sequence from a pool of exogenous elements of interest (*e.g.*, protein coding sequences, non-protein coding genes, regulatory elements) is inserted into one or more given loci of the genome.

[0288] It is understood that the multiplex methods suitable for the purpose of carrying out a screening or selection method, which is typically conducted for research purposes, may be different from the methods suitable for therapeutic purposes. For example, constitutive expression of certain elements (*e.g.*, a Cas nuclease and/or a modulator nucleic acid) may be undesirable for therapeutic purposes due to the potential of increased off-targeting. Conversely, for research purposes, constitutive expression of a Cas nuclease and/or a modulator nucleic acid may be desirable. For example, the constitutive expression provides a large window during which other elements can be introduced. When a stable cell line is established for the constitutive expression, the number of exogenous elements that need to be co-delivered into a single cell is also reduced. Therefore, constitutive expression of certain elements can increase the efficiency and reduce the complexity of a screening or selection process. Inducible expression of certain elements of the system disclosed herein may also be used for research purposes given similar advantages. Expression may be induced by an exogenous agent (*e.g.*, a small molecule) or by an endogenous molecule or complex present in a particular cell type (*e.g.*, at a particular stage of differentiation). Methods known in the art, such as those described in the “CRISPR Expression Systems” subsection *supra*, can be used for constitutively or inducibly expressing one or more elements.

[0289] It is further understood that despite the need to introduce at least three elements—the targeter nucleic acid, the modulator nucleic acid, and the Cas protein—these three elements can be delivered into the cell as a single complex of pre-formed RNP. Therefore, the efficiency

of the screening or selection process can also be achieved by pre-assembling a plurality of RNP complexes in a multiplex manner.

[0290] In certain embodiments, the method disclosed herein further comprises a step of identifying a targeter nucleic acid, a modulator nucleic acid, a Cas protein, a donor template, or a combination of two or more of these elements from the screening or selection process. A set of barcodes may be used, for example, in the donor template between two homology arms, to facilitate the identification. In specific embodiments, the method further comprises harvesting the population of cells; selectively amplifying a genomic DNA or RNA sample including the target nucleotide sequence(s) and/or the barcodes; and/or sequencing the genomic DNA or RNA sample and/or the barcodes that has been selectively amplified.

[0291] In another aspect, the present invention provides a library comprising a plurality of targeter nucleic acids disclosed herein, optionally further comprising one or more modulator nucleic acids disclosed herein. In another aspect, the present invention provides a library comprising a plurality of nucleic acids each comprising a regulatory element operably linked to a different targeter nucleic acid disclosed herein, optionally further comprising a regulatory element operably linked to a modulator nucleic acid disclosed herein. These libraries can be used in combination with one or more Cas proteins or Cas-coding nucleic acids disclosed herein, and/or one or more donor templates as disclosed herein for a screening or selection method.

20 **III. Pharmaceutical Compositions**

[0292] The present invention provides a composition (*e.g.*, pharmaceutical composition) comprising an engineered, non-naturally occurring system or a eukaryotic cell disclosed herein. In certain embodiments, the composition comprises a complex of the targeter nucleic acid and the modulator nucleic acid. In certain embodiments, the composition comprises an RNP comprising the targeter nucleic acid, the modulator nucleic acid, and a Cas protein (*e.g.*, the Cas nuclease that the targeter nucleic acid and the modulator nucleic acid are capable of activating or a related Cas protein). Targeter and modulator nucleic acids, and modifications thereto, are described in sections IA, IA 1, and IB; Cas proteins, *e.g.*, Cas nucleases, are described in section IC; RNPs are described in section IIA.

[0293] In addition, the present invention provides a method of producing a composition, the method comprising incubating the targeter nucleic acid and the modulator nucleic acid of an engineered, non-naturally occurring system disclosed herein under suitable conditions, thereby producing a composition (*e.g.*, pharmaceutical composition) comprising a complex of the targeter nucleic acid and the modulator nucleic acid. In certain embodiments, the method further comprises incubating the targeter nucleic acid and the modulator nucleic acid with a Cas protein

(*e.g.*, the Cas nuclease that the targeter nucleic acid and the modulator nucleic acid are capable of activating or a related Cas protein), thereby producing a complex of the targeter nucleic acid, the modulator nucleic acid, and the Cas protein (*e.g.*, an RNP). In certain embodiments, the method further comprises purifying the complex (*e.g.*, the RNP).

5 [0294] For therapeutic use, an engineered, non-naturally occurring system, a CRISPR expression system, or a cell comprising such system or modified by such system disclosed herein is combined with a pharmaceutically acceptable carrier. The term “pharmaceutically acceptable” as used herein can include those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the
10 tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit-to-risk ratio.

[0295] The term “pharmaceutically acceptable carrier” as used herein can include buffers, carriers, and excipients suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or
15 complication, commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable carriers include any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, emulsions (*e.g.*, such as an oil/water or water/oil emulsions), and various types of wetting agents. The compositions also can include stabilizers and preservatives. For examples of carriers, stabilizers and adjuvants, see, *e.g.*, Martin, Remington’s Pharmaceutical Sciences,
20 15th Ed., Mack Publ. Co., Easton, PA (1975). Pharmaceutically acceptable carriers include buffers, solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is known in the art.

[0296] In certain embodiments, a pharmaceutical composition disclosed herein comprises
25 a salt, *e.g.*, NaCl, MgCl₂, KCl, MgSO₄, etc.; a buffering agent, *e.g.*, a Tris buffer, N-(2-Hydroxyethyl)piperazine-N’-(2-ethanesulfonic acid) (HEPES), 2-(N-Morpholino)ethanesulfonic acid (MES), MES sodium salt, 3-(N-Morpholino)propanesulfonic acid (MOPS), N-tris[Hydroxymethyl]methyl-3-aminopropanesulfonic acid (TAPS), etc.; a solubilizing agent; a detergent, *e.g.*, a non-ionic detergent such as Tween-20, etc.; a nuclease inhibitor; and the like.
30 For example, in certain embodiments, a subject composition comprises a subject DNA-targeting RNA and a buffer for stabilizing nucleic acids.

[0297] In certain embodiments, a pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or
35 penetration of the composition. In such embodiments, suitable formulation materials include, but

are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants (see, *Remington's Pharmaceutical Sciences*, 18th ed. (Mack Publishing Company, 1990).

[0298] In certain embodiments, a pharmaceutical composition may contain nanoparticles, *e.g.*, polymeric nanoparticles, liposomes, or micelles (See Anselmo *et al.* (2016) *BIOENG. TRANSL. MED.* 1: 10-29). In certain embodiment, the pharmaceutical composition comprises an inorganic nanoparticle. Exemplary inorganic nanoparticles include, *e.g.*, magnetic nanoparticles (*e.g.*, Fe₃MnO₂) or silica. The outer surface of the nanoparticle can be conjugated with a positively charged polymer (*e.g.*, polyethylenimine, polylysine, polyserine) which allows for attachment (*e.g.*, conjugation or entrapment) of payload. In certain embodiment, the pharmaceutical composition comprises an organic nanoparticle (*e.g.*, entrapment of the payload inside the nanoparticle). Exemplary organic nanoparticles include, *e.g.*, SNALP liposomes that contain cationic lipids together with neutral helper lipids which are coated with polyethylene glycol (PEG) and protamine and nucleic acid complex coated with lipid coating. In certain embodiment, the pharmaceutical composition comprises a liposome, for example, a liposome disclosed in International (PCT) Publication No. WO2015/148863.

[0299] In certain embodiments, the pharmaceutical composition comprises a targeting moiety to increase target cell binding or uptake of nanoparticles and liposomes. Exemplary targeting moieties include cell specific antigens, monoclonal antibodies, single chain antibodies,

aptamers, polymers, sugars, and cell penetrating peptides. In certain embodiments, the pharmaceutical composition comprises a fusogenic or endosome-destabilizing peptide or polymer.

[0300] In certain embodiments, a pharmaceutical composition may contain a sustained-
5 or controlled-delivery formulation. Techniques for formulating sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. Sustained-release preparations may include, *e.g.*, porous polymeric microparticles or semipermeable polymer matrices in the form of shaped articles, *e.g.*, films, or microcapsules. Sustained release matrices may include polyesters,
10 hydrogels, polylactides, copolymers of L-glutamic acid and gamma ethyl-L-glutamate, poly (2-hydroxyethyl-inethacrylate), ethylene vinyl acetate, or poly-D(-)-3-hydroxybutyric acid. Sustained release compositions may also include liposomes that can be prepared by any of several methods known in the art.

[0301] A pharmaceutical composition of the invention can be administered by a variety
15 of methods known in the art. The route and/or mode of administration vary depending upon the desired results. Administration can be intravenous, intramuscular, intraperitoneal, or subcutaneous, or administered proximal to the site of the target. The pharmaceutically acceptable carrier should be suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (*e.g.*, by injection or infusion). Depending on the route of
20 administration, the active compound, *i.e.*, the multispecific antibody of the invention, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

[0302] Formulation components suitable for parenteral administration include a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin,
25 propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as EDTA; buffers such as acetates, citrates or phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose.

[0303] For intravenous administration, suitable carriers include physiological saline,
30 bacteriostatic water, Cremophor ELTM (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). The carrier should be stable under the conditions of manufacture and storage, and should be preserved against microorganisms. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof.

[0304] Pharmaceutical formulations preferably are sterile. Sterilization can be accomplished by any suitable method, *e.g.*, filtration through sterile filtration membranes. Where the composition is lyophilized, filter sterilization can be conducted prior to or following lyophilization and reconstitution. In certain embodiments, a multispecific antibody is

5 lyophilized, and then reconstituted in buffered saline, at the time of administration.

[0305] Pharmaceutical compositions of the invention can be prepared in accordance with methods well known and routinely practiced in the art. See, *e.g.*, Remington: The Science and Practice of Pharmacy, Mack Publishing Co., 20th ed., 2000; and Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

10 Pharmaceutical compositions are preferably manufactured under GMP conditions. Typically, a therapeutically effective dose or efficacious dose of the multispecific antibody of the invention is employed in the pharmaceutical compositions of the invention. The multispecific antibodies of the invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. Dosage regimens are adjusted to provide the optimum
15 desired response (*e.g.*, a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically
20 discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0306] Actual dosage levels of the active ingredients in the pharmaceutical compositions of the invention can be varied so as to obtain an amount of the active ingredient which is
25 effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level depends upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the
30 duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors.

IV. Therapeutic Uses

[0307] The engineered, non-naturally occurring system and CRISPR expression system disclosed herein are useful for targeting, editing, and/or modifying the genomic DNA in a cell or organism. These systems, as well as a cell comprising one of the systems or a cell whose genome has been modified by the engineered, non-naturally occurring system, can be used to treat a disease or disorder in which modification of genetic or epigenetic information is desirable. Accordingly, in another aspect, the present invention provides a method of treating a disease or disorder, the method comprising administering to a subject in need thereof an effective amount of a non-naturally occurring system, a CRISPR expression system, or a cell disclosed herein.

[0308] The term “subject” can include human and non-human animals. Non-human animals include all vertebrates, *e.g.*, mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, and reptiles. Except when noted, the terms “patient” or “subject” are used herein interchangeably.

[0309] The terms “treatment”, “treating”, “treat”, “treated”, and the like, as used herein, can include obtaining a desired pharmacologic and/or physiologic effect. The effect may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease or delaying the disease progression. “Treatment”, as used herein, covers any treatment of a disease in a mammal, *e.g.*, in a human, and includes: (a) inhibiting the disease, *i.e.*, arresting its development; and (b) relieving the disease, *i.e.*, causing regression of the disease. It is understood that a disease or disorder may be identified by genetic methods and treated prior to manifestation of any medical symptom.

[0310] For therapeutic purposes, the method disclosed herein is particularly suitable for editing or modifying a proliferating cell, such as a stem cell (*e.g.*, a hematopoietic stem cell), a progenitor cell (*e.g.*, a hematopoietic progenitor cell or a lymphoid progenitor cell), or a memory cell (*e.g.*, a memory T cell). Given that such cell is delivered to a subject and will proliferate *in vivo*, tolerance to off-target events is low. Prior to delivery, however, it is possible to assess the on-target and off-target events, thereby selecting one or more colonies that have the desired edit or modification and lack any undesired edit or modification. Therefore, lower editing or modifying efficiency can be tolerated for such cell. The engineered, non-naturally occurring system of the present invention has the advantage of increasing or decreasing the efficiency of nucleic acid cleavage by, for example, adjusting the hybridization of modified dual guide nucleic acids. As a result, it can be used to minimize off-target events when creating genetically engineered proliferating cells.

[0311] For minimization of toxicity and off-target effect, it is important to control the concentration of the modified guide CRISPR-Cas system, *e.g.*, modified dual guide CRISPR-Cas

system, delivered. Optimal concentrations can be determined by testing different concentrations in a cellular, tissue, or non-human eukaryote animal model and using deep sequencing to analyze the extent of modification at potential off-target genomic loci. The concentration that gives the highest level of on-target modification while minimizing the level of off-target modification can be selected for *ex vivo* or *in vivo* delivery.

B. Gene Therapies

[0312] It is understood that the engineered, non-naturally occurring system and CRISPR expression system disclosed herein can be used to treat a genetic disease or disorder, *i.e.*, a disease or disorder associated with or otherwise mediated by an undesirable mutation in the genome of a subject.

[0313] Exemplary genetic diseases or disorders include age-related macular degeneration, adrenoleukodystrophy (ALD), Alagille syndrome, alpha-1-antitrypsin deficiency, argininemia, argininosuccinic aciduria, ataxia (*e.g.*, Friedreich ataxia, spinocerebellar ataxias, ataxia telangiectasia, essential tremor, spastic paraplegia), autism, biliary atresia, biotinidase deficiency, carbamoyl phosphate synthetase I deficiency, carbohydrate deficient glycoprotein syndrome (CDGS), a central nervous system (CNS)-related disorder (*e.g.*, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), canavan disease (CD), ischemia, multiple sclerosis (MS), neuropathic pain, Parkinson's disease), Bloom's syndrome, cancer, Charcot-Marie-Tooth disease (*e.g.*, peroneal muscular atrophy, hereditary motor sensory neuropathy), congenital hepatic porphyria, citrullinemia, Crigler-Najjar syndrome, cystic fibrosis (CF), Dentatorubro-Pallidoluysian Atrophy (DRPLA), Diabetes insipidus, Fabry, familial hypercholesterolemia (LDL receptor defect), Fanconi's anemia, fragile X syndrome, a fatty acid oxidation disorder, galactosemia, glucose-6-phosphate dehydrogenase (G6PD), glycogen storage diseases (*e.g.*, type I (glucose-6-phosphatase deficiency, Von Gierke II (alpha glucosidase deficiency, Pompe), III (debrancher enzyme deficiency, Cori), IV (brancher enzyme deficiency, Anderson), V (muscle glycogen phosphorylase deficiency, McArdle), VII (muscle phosphofructokinase deficiency, Tauri), VI (liver phosphorylase deficiency, Hers), IX (liver glycogen phosphorylase kinase deficiency)), hemophilia A (associated with defective factor VIII), hemophilia B (associated with defective factor IX), Huntington's disease, glutaric aciduria, hypophosphatemia, Krabbe, lactic acidosis, Lafora disease, Leber's Congenital Amaurosis, Lesch Nyhan syndrome, a lysosomal storage disease, metachromatic leukodystrophy disease (MLD), mucopolysaccharidosis (MPS) (*e.g.*, Hunter syndrome, Hurler syndrome, Maroteaux-Lamy syndrome, Sanfilippo syndrome, Scheie syndrome, Morquio syndrome, other, MPSI, MPSII, MPSIII, MSIV, MPS 7), a

muscular/skeletal disorder (e.g., muscular dystrophy, Duchenne muscular dystrophy), myotonic Dystrophy (DM), neoplasia, N-acetylglutamate synthase deficiency, ornithine transcarbamylase deficiency, phenylketonuria, primary open angle glaucoma, retinitis pigmentosa, schizophrenia, Severe Combined Immune Deficiency (SCID), Spinobulbar Muscular Atrophy (SBMA), sickle cell anemia, Usher syndrome, Tay-Sachs disease, thalassemia (e.g., β -Thalassemia), trinucleotide repeat disorders, tyrosinemia, Wilson's disease, Wiskott-Aldrich syndrome, X-linked chronic granulomatous disease (CGD), X-linked severe combined immune deficiency, and xeroderma pigmentosum.

[0314] Genetic Disorders contemplated herein can include, but are not limited to:

[0315] **Neoplasia:** Genes linked to this disorder: PTEN; ATM; ATR; EGFR; ERBB2; ERBB3; ERBB4; Notch1; Notch2; Notch3; Notch4; AKT; AKT2; AKT3; HIF; HIF1 α ; HIF3 α ; Met; HRG; Bcl2; PPAR α ; PPAR γ ; WT1 (Wilms Tumor); FGF Receptor Family members (5 members: 1, 2, 3, 4, 5); CDKN2a; APC; RB (retinoblastoma); MEN1; VHL; BRCA1; BRCA2; AR (Androgen Receptor); TSG101; IGF; IGF Receptor; Igf1 (4 variants); Igf2 (3 variants); Igf 1 Receptor; Igf 2 Receptor; Bax; Bcl2; caspases family (9 members: 1, 2, 3, 4, 6, 7, 8, 9, 12); Kras; Apc;

[0316] **Age-related Macular Degeneration:** Genes linked to these disorders Abcr; Cc12; Cc2; cp (cemloplasmin); Timp3; cathepsinD; Vidlr; Ccr2;

[0317] **Schizophrenia Disorders:** Genes linked to this disorder: Neuregulin1 (Nrg1); Erb4 (receptor for Neuregulin); Complexin1 (Cplx1); Tph1 Tryptophan hydroxylase; Tph2 Tryptophan hydroxylase 2; Neurexin 1; GSK3; GSK3 α ; GSK3 β ;

[0318] **Trinucleotide Repeat Disorders:** Genes linked to this disorder: 5 HTT (Huntington's Dx); SBMA/SMAX1/AR (Kennedy's Dx); FXN/X25 (Friedrich's Ataxia); ATX3 (Machado-Joseph's Dx); ATXN1 and ATXN2 (spinocerebellar ataxias); DMPK (myotonic dystrophy); Atrophin-1 and Atnl (DRPLA Dx); CBP (Creb-BP – global instability); VLDLR (Alzheimer's); Atxn7; Atxn10;

[0319] **Fragile X Syndrome:** Genes linked to this disorder: FMR2; FXR1; FXR2; mGLURS;

[0320] **Secretase Related Disorders:** Genes linked to this disorder: A β -1 (alpha and beta); Presenil n (Psen1); nicastrin (Ncstn); PEN-2;

[0321] **Others:** Genes linked to this disorder: Nos1; Paip1; Nati; Nat2;

[0322] **Prion – related disorders:** Gene linked to this disorder: Prp;

[0323] **ALS:** Genes linked to this disorder: SOD1; ALS2; STEX; FUS; TARDBP; VEGF (VEGF-a; VEGF-b; VEGF-c);

- [0324] Drug addiction:** Genes linked to this disorder: Prkce (alcohol); Drd2; Drd4; ABAT (alcohol); GRIA2; GrmS; Grinl; Htrlb; Grin2a; Drd3; Pdyn; Grial (alcohol);
- [0325] Autism:** Genes linked to this disorder: Meep2; BZRAP1; MDGA2; SemaSA; Neurexin 1; Fragile X (FMR2 (AFF2); FXR1; FXR2; MglurS);
- 5 **[0326] Alzheimer's Disease** Genes linked to this disorder: El; CHIP; UCH; UBB; Tau; LRP; PICALM; Clusterin; PS1; SORL1; CR1; Vidlr; Ubal; Uba3; CHIP28 (Aqp1, Aquaporin 1); Uchl1; Uchl3; APP;
- [0327] Inflammation and Immune-related disorders** Genes linked to this disorder: IL-10; IL-1 (IL-1a; IL-1b); IL-13; IL-17 (IL-17a (CTLA8); IL-17b; IL-17c; IL-17d; IL-17f); 11-23; Cx3crl; ptpn22; TNFa; NOD2/CARD15 for IBD; IL-6; IL-12 (IL-12a; IL-12b); CTLA4; Cx3c11, AAT deficiency/mutations, AIDS (KIR3DL1, NKAT3, NKB1, ANIB11, KIR3DS1, IFNG, CXCL12, SDF1); Autoimmune lymphoproliferative syndrome (TNFRSF6, APT1, FAS, CD95, ALPS1A); Combined immunodeficiency, (IL2RG, SCIDX1, SCIDX, IMD4); HIV-1 (CCL5, SCYA5, D17S136E, TCP228), HIV susceptibility or infection (IL10, CSIF, CMKBR2, CCR2, CMKBR5, CCCR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFSF5, CD4OLG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TACI); Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-17f), 11-23, Cx3crl, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b), CTLA4, Cx3c11); Severe combined
- 10 Cx3crl; ptpn22; TNFa; NOD2/CARD15 for IBD; IL-6; IL-12 (IL-12a; IL-12b); CTLA4; Cx3c11, AAT deficiency/mutations, AIDS (KIR3DL1, NKAT3, NKB1, ANIB11, KIR3DS1, IFNG, CXCL12, SDF1); Autoimmune lymphoproliferative syndrome (TNFRSF6, APT1, FAS, CD95, ALPS1A); Combined immunodeficiency, (IL2RG, SCIDX1, SCIDX, IMD4); HIV-1 (CCL5, SCYA5, D17S136E, TCP228), HIV susceptibility or infection (IL10, CSIF, CMKBR2, CCR2, CMKBR5, CCCR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFSF5, CD4OLG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TACI); Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-17f), 11-23, Cx3crl, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b), CTLA4, Cx3c11); Severe combined
- 15 CCR2, CMKBR5, CCCR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFSF5, CD4OLG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TACI); Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-17f), 11-23, Cx3crl, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b), CTLA4, Cx3c11); Severe combined
- 20 immunodeficiencies (SCIDs)(JAK3, JAKL, DCLRE1C, ARTEMIS, SCIDA, RAG1, RAG2, ADA, PTPRC, CD45, LCA, IL7R, CD3D, T3D, IL2RG, SCIDX1, SCIDX, IMD4);
- [0328] Parkinson's,** Genes linked to this disorder: x-Synuclein; DJ-1; LRRK2; Parkin; PINK1;
- [0329] Blood and coagulation disorders:** Genes linked to these disorders: Anemia (CDAN1, CDA1, RPS19, DBA, PKLR, PK1, NT5C3, UMPH I, PSN1, RHAG, RH50A, NRAMP2, SPTB, ALAS2, ANH I, ASB, ABCB7, ABC7, ASAT); Bare lymphocyte syndrome (TAPBP, TPSN, TAP2, ABCB3, PSF2, RINGI 1, MHC2TA, C2TA, RFX5, RFXAP, RFX5), Bleeding disorders (TBXA2R, P2RX I, P2X I); Factor H and factor H-like 1 (HF1, CFH, HUS); Factor V and factor VIII (MCFD2); Factor VII deficiency (F7); Factor X deficiency (F10); Factor XI deficiency (F11); Factor XII deficiency (F12, HAF); Factor XIIIa deficiency (F13A1, F13A); Factor XIIIb deficiency (F13B); Fanconi anemia (FANCA, FACA, FA1, FA, FAA, FAAP95, FAAP90, FLJ34064, FANCB, FANCC, FACC, BRCA2, FANCD1, FANCD2, FANCD, FACD, FAD, FANCE, FACE, FANCF, XRCC9, FANCG, BRIP1, BACH1, FANCI, PHF9, FANCL, FANCM, ICIAA1596); Hemophagocytic lymphohistiocytosis disorders (PRF1, HPLH2, UNC13D, MUNC13-4, HPLH3, HLH3, FHL3); Hemophilia A (F8, F8C, HEMA);
- 35 HPLH2, UNC13D, MUNC13-4, HPLH3, HLH3, FHL3); Hemophilia A (F8, F8C, HEMA);

Hemophilia B (F9, HEMB), Hemorrhagic disorders (PI, ATT, F5); Leukocyte deficiencies and disorders (ITGB2, CD18, LCAMB, LAD, EIF2B1, EIF2BA, EIF2B2, EIF2B3, EIF2B5, LVWM, CACH, CLE, EIF2B4); Sickle cell anemia (HBB); Thalassemia (HBA2, HBB, HBD, LCRB, HBA1);

- 5 **[0330] Cell dysregulation and oncology disorders:** Genes linked to these disorders: B-cell non-Hodgkin lymphoma (BCL7A, BCL7); Leukemia (TALI, TCL5, SCL, TAL2, FLT3, NBS 1, NBS, ZNFNIA1, IK1, LYF1, HOXD4, HOX4B, BCR, CML, PHL, ALL, ARNT, KRAS2, RASK2, GMPS, AFIO, ARHGEF12, LARG, KIAA0382, CALM, CLTH, CEBPA, CEBP, CHIC2, BTL, FLT3, KIT, PBT, LPP, NPM1, NUP214, D9S46E, CAN, CAIN, RUNX 1, 10 CBFA2, AML1, WHSC 1 LI, NSD3, FLT3, AF1Q, NPM 1, NUMA1, ZNF145, PLZF, PML, MYL, STAT5B, AFI 0, CALM, CLTH, ARL1 1, ARLTS1, P2RX7, P2X7, BCR, CML, PHL, ALL, GRAF, NFI, VRNF, WSS, NFNS, PTPNI 1, PTP2C, SHP2, NS 1, BCL2, CCND1, PRAD1, BCL1, TCRA, GATA1, GF1, ERYF1, NFE1, ABL1, NQO1, DIA4, NMOR1, NUP214, D9S46E, CAN, CAIN);
- 15 **[0331] Metabolic, liver, kidney disorders:** Genes linked to these disorders: Amyloid neuropathy (TTR, PALS); Amyloidosis (APOA1, APP, AAA, CVAP, AD1, GSN, FGA, LYZ, UR, PALS); Cirrhosis (KATI 8, KRT8, CaHIA, NAIC, TEX292, KIAA1988); Cystic fibrosis (CFTR, ABCC7, CF, MRP7); Glycogen storage diseases (SLC2A2, GLUT2, G6PC, G6PT, G6PT1, GAA, LAMP2, LAMPS, AGL, GDE, GBE1, GYS2, PYGL, PFKM); Hepatic adenoma, 20 142330 (TCF1, HNF1A, MODY3), Hepatic failure, early onset, and neurologic disorder (SCOD1, SCO1), Hepatic lipase deficiency (LIPC), Hepatoblastoma, cancer and carcinomas (CTNNB1, PDGFRL, PDGRL, PRLTS, AXIN1, AXIN, CTNNB1, TP53, P53, LFS1, IGF2R, MPRI, MET, CASP8, MCH5; Medullary cystic kidney disease (UMOD, HNFJ, FJHN, MCKD2, ADMCKD2); Phenylketonuria (PAH, PKU1, QDPR, DHPR, PTS); Polycystic kidney and 25 hepatic disease (FCYT, PKHD1, ARPKD, PKD2, PKD4, PKDTS, PRKCSH, G19P1, PCLD, SEC63);

- [0332] Muscular/Skeletal Disorders:** Genes linked to these disorders: Becker muscular dystrophy (DMD, BMD, MYF6), Duchenne Muscular Dystrophy (DMD, BMD); Emery-Dreifuss muscular dystrophy (LMNA, LMN1, EMD2, FPLD, CMD1A, HGPS, LGMD1B, 30 LMNA, LMN1, EMD2, FPLD, CMD1A); Facioscapulohumeral muscular dystrophy (FSHMD1A, FSHD1A); Muscular dystrophy (FKRP, MDC1C, LGMD2I, LAMA2, LAMM, LARGE, KIAA0609, MDC1D, FCMD, TTID, MYOT, CAPN3, CANP3, DYSF, LGMD2B, SGCG, LGMD2C, DMDA1, SCG3, SGCA, ADL, DAG2, LGMD2D, DMDA2, SGCB, LGMD2E, SGCD, SGD, LGMD2F, CMD1L, TCAP, LGMD2G, CMD1N, TRIM32, HT2A, 35 LGMD2H, FKRP, MDC1C, LGMD2I, TTN, CMD1G, TMD, LGMD2J, POMT1, CAV3,

- LGMD1C, SEPN1, SELN, RSMD1, PLEC1, PLTN, EBS1); Osteopetrosis (LAPS, BMND1, LRP7, LR3, OPPG, VBCH2, CLCN7, CLC7, OPTA2, OSTM1, GL, TCIRG1, TIRC7, OC116, OPTB1); Muscular atrophy (VAPB, VAPC, ALS8, SMN1, SMA1, SMA2, SMA3, SMA4, BSCL2, SPG17, GARS, SMAD1, CMT2D, HEXB, IGHMBP2, SMUBP2, CATF1, SMARD1);
- 5 **[0333] Neurological and Neuronal disorders:** Genes linked to these disorders: ALS (SOD1, ALS2, STEX, FUS, TARDBP, VEGF (VEGF-a, VEGF-b, VEGF-c); Alzheimer disease (APP, AAA, CVAP, AD1, APOE, AD2, PSEN2, AD4, STM2, APBB2, FE65L1, NOS3, PLA2, URK, ACE, DCPI, ACEI, MPO, PACIP1, PAXIPIL, PTIP, A2M, BLMH, BMH, PSEN1, AD3); Autism (Mecp2, BZRAP I, MDGA2, Sema5A, Neurex 1, GLO1, MECP2, RTT, PPMX,
- 10 MRX16, MRX79, NLGN3, NLGN4, KIAA1260, AUTSX2); Fragile X Syndrome (FMR2, FXR1, FXR2, mGLUR5); Huntington's disease and disease like disorders (HD, IT15, PRNP, PRIP, JPH3, JP3, HDL2, TBP, SCA17); Parkinson disease (NR4A2, NURR1, NOT, TINUR, SNCAIP, TBP, SCA17, SNCA, NACP, PARK1, PARK4, DJ1, PARK7, LRRK2, PARKS, PINK1, PARK6, UCHL1, PARKS, SNCA, NACP, PARK1, PARK4, PRKN, PARK-2, PDJ,
- 15 DBH, NDUFV2); Rett syndrome (MECP2, RTT, PPMX, MRX16, MRX79, CDKL5, STK9, MECP2, RTT, PPMX, MRX16, MRX79, x-Synuclein, DJ-1); Schizophrenia (Neuregulin1 (Nrg1), Erb4 (receptor for Neuregulin), Complexin1 (Cp1x1), Tph1 Tryptophan hydroxylase, Tph2, Tryptophan hydroxylase 2, Neurexin 1, GSK3, GSK3a, GSK3b, 5-HTT (S1c6a4), COMT, DRD (Drd 1a), SLC6A3, DAOA, DTNBP1, Dao (Daol)); Secretase Related Disorders (APH-1 (alpha and beta), Presenilin 1 (Psen1), nicastrin, (Ncstn), PEN-2, Nosl, Parpl, Natl, Nat2); Trinucleotide Repeat Disorders (HTT (Huntington's Dx), SBMA/SMAX1/AR (Kennedy's Dx), FXN/X25 (Friedrich's Ataxia), ATX3 (Machado-Joseph's Dx), ATXN1 and ATXN2 (spinocerebellar ataxias), DMPK (myotonic dystrophy), Atrophin-1 and Atn1 (DRPLA Dx), CBP (Creb-BP – global instability), VLDLR (Alzheimer's), Atxn7, Atxn10);
- 25 **[0334] Ocular-related disorders:** Genes linked to these disorders: Age-related macular degeneration (Aber, Cc12, Cc2, cp (ceruloplasmin), Timp3, cathepsinD, Vldlr, Ccr2); Cataract (CRYAA, CRYA1, CRYBB2, CRYB2, PITX3, BFSP2, CP49, CP47, CRYAA, CRYA1, PAX6, AN2, MGDA, CRYBA1, CRYB1, CRYGC, CRYG3, CCL, LIM2, MP19, CRYGD, CRYG4, BFSP2, CP49, CP47, HSF4, CTM, HSF4, CTM, MIP, AQPO, CRYAB, CRYA2, CTPP2,
- 30 CRYBB1, CRYGD, CRYG4, CRYBB2, CRYB2, CRYGC, CRYG3, CCL, CRYAA, CRYA1, GJA8, CX50, CAE1, GJA3, CX46, CZP3, CAE3, CCM1, CAM, KRIT1); Corneal clouding and dystrophy (APOA1, TGFBI, CSD2, CDGG1, CSD, BIGH3, CDG2, TACSTD2, TROP2, M1S1, VSX1, RINX, PPCD, PPD, KTCN, COL8A2, FECD, PPCD2, PIP5K3, CFD); Cornea plana congenital (KERA, CNA2); Glaucoma (MYOC, TIGR, GLC1A, JOAG, GPOA, OPTN, GLC1E,
- 35 FIP2, HYPL, NRP, CYP1B1, GLC3A, OPAL, NTG, NPG, CYP1B1, GLC3A); Leber congenital

amaurosis (CRB1, RP12, CRX, CORD2, CRD, RPGRIP1, LCA6, CORD9, RPE65, RP20, AIPL1, LCA4, GUCY2D, GUC2D, LCA1, CORD6, RDH12, LCA3); Macular dystrophy (ELOVL4, ADMD, STGD2, STGD3, RDS, RP7, PRPH2, PRPH, AVMD, AOFMD, VMD2);

[0335] P13K/AKT Cellular Signaling disorders: Genes linked to these disorders:

5 PRKCE; ITGAM; ITGA5; IRAK1; PRKAA2; EIF2AK2; PTEN; EIF4E; PRKCZ; GRK6; MAPK1; TSC1; PLK1; AKT2; IKBKB; PIK3CA; CDK8; CDKN1B; NFKB2; BCL2; PIK3CB; PPP2R1A; MAPK8; BCL2L1; MAPK3; TSC2; ITGA1; KRAS; EIF4EBP1; RELA; PRKCD; NOS3; PRKAA1; MAPK9; CDK2; PPP2CA; PIM1; ITGB7; YWHAZ; ILK; TP53; RAF1; IKBKG; RELB; DYRK1A; CDKN1A; ITGB1; MAP2K2; JAK1; AKT1; JAK2; PIK3R1;
10 CHUK; PDPK1; PPP2R5C; CTNNB1; MAP2K1; NFKB1; PAK3; ITGB3; CCND1; GSK3A; FRAP1; SFN; ITGA2; TTK; CSNK1A1; BRAF; GSK3B; AKT3; FOXO1; SOK; HS P9OAA1; RP S 6KB1;

[0336] ERK/MAPK Cellular Signaling disorders: Genes linked to these disorders:

PRKCE; ITGAM; ITGA5; HSPB1; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A; TLN1;
15 EIF4E; ELK1; GRK6; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; CREB1; PRKCI; PTK2; FOS; RPS6KA4; PIK3CB; PPP2R1A; PIK3C3; MAPK8; MAPK3; ITGA1; ETS1; KRAS; MYCN; EIF4EBP1; PPARG; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PPP2CA; PIM1; PIK3C2A; ITGB7; YWHAZ; PPP1CC; KSR1; PXN; RAF1; FYN; DYRK1A; ITGB1; MAP2K2; PAK4; PIK3R1; STAT3; PPP2R5C; MAP2K1; PAK3; ITGB3; ESR1; ITGA2; MYC;
20 TTK; CSNK1A1; CRKL; BRAE; ATF4; PRKCA; SRF; STAT1; SGK;

[0337] Glucocorticoid Receptor Cellular Signaling disorders: Genes linked to these

disorders: RAC1; TAF4B; EP300; SMAD2; TRAF6; PCAF; ELK1; MAPK1; SMAD3; AKT2; IKBKB; NCOR2; UBE2I; PIK3CA; CREB1; FOS; HSPA5; NFKB2; BCL2; MAP3K14; STAT5B; PIK3CB; PIK3C3; MAPK8; BCL2L1; MAPK3; TSC22D3; MAPK10; NRIP1;
25 KRAS; MAPK13; RELA; STAT5A; MAPK9; NOS2A; PBX1; NR3C1; PIK3C2A; CDKN1C; TRAF2; SERPINE1; NCOA3; MAPK14; TNF; RAF1; IKBKG; MAP3K7; CREBBP; CDKN1A; MAP2K2; JAK1; IL8; NCOA2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; TGFBR1; ESR1; SMAD4; CEBPB; JUN; AR; AKT3; CCL2; MMP 1; STAT1; IL6; HSP9OAA1;

[0338] Axonal Guidance Cellular Signaling disorders: Genes linked to these

disorders: PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; ADAM12; IGF1; RAC1; RAP1A; EIF4E; PRKCZ; NRPI; NTRK2; ARHGEF7; SMO; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; AKT2; PIK3CA; ERBB2; PRKCI; PTK2; CFL1; GNAQ; PIK3CB; CXCL12; PIK3C3; WNT11; PRKD1; GNB2L1; ABL1 ; MAPK3; ITGA1; KRAS; RHOA; PRKCD; PIK3C2A;
35 ITGB7; GLI2; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; ADAM17; AKT1;

PIK3R1; GUI; WNT5A; ADAM10; MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; CRKL; RND1; GSK3B; AKT3; PRKCA;

[0339] Ephrin Receptor Cellular Signaling disorders: Genes linked to these disorders:

PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A;
 5 GRK6; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; PLK1; AKT2; DOK1; CDK8;
 CREB1; PTK2; CFL1; GNAQ; MAP3K14; CXCL12; MAPK8; GNB2L1; ABL1; MAPK3;
 ITGA1; KRAS; RHOA; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PIM1; ITGB7; PXN;
 RAF1; FYN; DYRK1A; ITGB1; MAP2K2; PAK4; AKT1; JAK2; STAT3; ADAM10;
 MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; TTK; CSNK1A1; CRKL; BRAF;
 10 PTPN13; ATF4; AKT3; SGK;

[0340] Actin Cytoskeleton Cellular Signaling disorders: Genes linked to these

disorders: ACTN4; PRKCE; ITGAM; ROCK1; ITGA5; IRAK1; PRKAA2; EIF2AK2; RAC1;
 INS; ARHGEF7; GRK6; ROCK2; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; PTK2;
 CFL1; PIK3CB; MYH9; DIAPH1; PIK3C3; MAPK8; F2R; MAPK3; SLC9A1; ITGA1; KRAS;
 15 RHOA; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; ITGB7; PPP1CC; PXN; VIL2;
 RAF1; GSN; DYRK1A; ITGB1; MAP2K2; PAK4; PIP5K1A; PIK3R1; MAP2K1; PAK3;
 ITGB3; CDC42; APC; ITGA2; TTK; CSNK1A1; CRKL; BRAF; VAV3; SGK;

[0341] Huntington's Disease Cellular Signaling disorders: Genes linked to these

disorders: PRKCE; IGF1; EP300; RCOR1; PRKCZ; HDAC4; TGM2; MAPK1; CAPNS1;
 20 AKT2; EGFR; NCOR2; SP1; CAPN2; PIK3CA; HDAC5; CREB1; PRKC1; HS PA5 ; REST;
 GNAQ; PIK3CB; PIK3C3; MAPK8; IGF1R; PRKD1; GNB2L1; BCL2L1; CAPN1; MAPK3;
 CASP8; HDAC2; HDAC7A; PRKCD; HDAC11; MAPK9; HDAC9; PIK3C2A; HDAC3; TP53;
 CASP9; CREBBP; AKT1; PIK3R1; PDPK1; CASP1; APAF1; FRAP1; CASP2; JUN; BAX;
 ATF4; AKT3; PRKCA; CLTC; SGK; HDAC6; CASP3;

[0342] Apoptosis Cellular Signaling disorders: Genes linked to these disorders:

PRKCE; ROCK1; BID; IRAK1; PRKAA2; EIF2AK2; BAK1; BIRC4; GRK6; MAPK1;
 CAPNS1; PLK1; AKT2; IKBKB; CAPN2; CDK8; FAS; NFKB2; BCL2; MAP3K14; MAPK8;
 BCL2L1; CAPN1; MAPK3; CASP8; KRAS; RELA; PRKCD; PRKAA1; MAPK9; CDK2;
 PIM1; TP53; TNF; RAF1; IKBKG; RELB; CASP9; DYRK1A; MAP2K2; CHUK; APAF1;
 30 MAP2K1; NFKB1; PAK3; LMNA; CASP2; BIRC2; TTK; CSNK1A1; BRAF; BAX; PRKCA;
 SGK; CASP3; BTRC3; PARP1;

[0343] B Cell Receptor Cellular Signaling disorders: Genes linked to these disorders:

RAC1; PTEN; LYN; ELK1; MAPK1; RAC2; PTPN11; AKT2; IKBKB; PIK3CA; CREB1;
 SYK; NFKB2; CAMK2A; MAP3K14; PIK3CB; PIK3C3; MAPK8; BCL2L1; ABL1; MAPK3;
 35 ETS1; KRAS; MAPK13; RELA; PTPN6; MAPK9; EGR1; PIK3C2A; BTK; MAPK14; RAF1;

IKBKG; RELB; MAP3K7; MAP2K2; AKT1; PIK3R1; CHUK; MAP2K1; NFKB1; CDC42; GSK3A; FRAP1; BCL6; BCL10; JUN; GSK3B; ATF4; AKT3; VAV3; RPS6KB1;

[0344] Leukocyte Extravasation Cellular Signaling disorders: Genes linked to these disorders: ACTN4; CD44; PRKCE; ITGAM; ROCK1; CXCR4; CYBA; RAC1; RAP1A;

5 PRKCZ; ROCK2; RAC2; PTPN11; MMP14; PIK3CA; PRKCI; PTK2; PIK3CB; CXCL12; PIK3C3; MAPK8; PRKD1; ABL1; MAPK10; CYBB; MAPK13; RHOA; PRKCD; MAPK9; SRC; PIK3C2A; BTK; MAPK14; NOX1; PXN; VIL2; VASP; ITGB1; MAP2K2; CTNND1; PIK3R1; CTNNB1; CLDN1; CDC42; FUR; ITK; CRKL; VAV3; CTTN; PRKCA; MMP1; MMP9;

10 **[0345] Integrin Cellular Signaling disorders:** Genes linked to these disorders: ACTN4; ITGAM; ROCK1; ITGA5; RAC1; PTEN; RAP1A; TLN1; ARHGEF7; MAPK1; RAC2; CAPNS1; AKT2; CAPN2; PIK3CA; PTK2; PIK3CB; PIK3C3; MAPK8; CAV1; CAPN1; ABL1; MAPK3; ITGA1; KRAS; RHOA; SRC; PIK3C2A; ITGB7; PPP1CC; ILK; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; AKT1; PIK3R1; TNK2; MAP2K1; PAK3; ITGB3;

15 CDC42; RND3; ITGA2; CRKL; BRAF; GSK3B; AKT3;

[0346] Acute Phase Response Cellular Signaling disorders: Genes linked to these disorders: IRAK1; SOD2; MYD88; TRAF6; ELK1; MAPK1; PTPN11; AKT2; IKBKB; PIK3CA; FOS; NFKB2; MAP3K14; PIK3CB; MAPK8; RIPK1; MAPK3; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; FTL; NR3C1; TRAF2; SERPINE1; MAPK14; TNF;

20 RAF1; PDK1; IKBKG; RELB; MAP3K7; MAP2K2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; FRAP1; CEBPB; JUN; AKT3; IL1R1; IL6;

[0347] PTEN Cellular Signaling disorders: Genes linked to these disorders: ITGAM; ITGA5; RAC1; PTEN; PRKCZ; BCL2L11; MAPK1; RAC2; AKT2; EGFR; IKBKB; CBL; PIK3CA; CDKN1B; PTK2; NFKB2; BCL2; PIK3CB; BCL2L1; MAPK3; ITGA1; KRAS;

25 ITGB7; ILK; PDGFRB; INSR; RAF1; IKBKG; CASP9; CDKN1A; ITGB1; MAP2K2; AKT1; PIK3R1; CHUK; PDGFRA; PDPK1; MAP2K1; NFKB1; ITGB3; CDC42; CCND1; GSK3A; ITGA2; GSK3B; AKT3; FOXO1; CASP3;

[0348] p53 Cellular Signaling disorders: Genes linked to these disorders: RPS6KB1 PTEN; EP300; BBC3; PCAF; FASN; BRCA1; GADD45A; BIRC5; AKT2; PIK3CA; CHEK1;

30 TP53INP1; BCL2; PIK3CB; PIK3C3; MAPK8; THBS 1; ATR; BCL2L1; E2F1; PMAIP1; CHEK2; TNFASF10B; TP73; RB1; HDAC9; CDK2; PIK3C2A; MAPK14; TP53; LRDD; CDKN1A; HIPK2; AKT1; PIK3R1; RAM2B; APAF1; CTNNB1; SIRT1; CCND1; PRKDC; ATM; SFN; CDKN2A; JUN; SNAI2; GSK3B; BAX; AKT3;

[0349] Aryl Hydrocarbon Receptor Cellular Signaling disorders: Genes linked to these disorders: HSPB1; EP300; FASN; TGM2; RXRA; MAPK1; NQO1; NCOR2; SP1; ARNT;

35

CDKN1B; FOS; CHEK1; SMARCA4; NFKB2; MAPK8; ALDH1A1; ATR; E2F1; MAPK3; NRIP1; CHEK2; RELA; TP73; GSTP1; RB1; SRC; CDK2; AHR; NFE2L2; NCOA3; TP53; TNF; CDKN1A; NCOA2; APAF1; NFKB1; CCND1; ATM; ESR1; CDKN2A; MYC; JUN; ESR2; BAX; IL6; CYP1B1; HSP90AA1;

- 5 **[0350] Xenobiotic Metabolism Cellular Signaling disorders:** Genes linked to these disorders: PRKCE; EP300; PRKCZ; RXRA; MAPK1; NQO1; NCOR2; PIK3CA; ARNT; PRKCI; NFKB2; CAMK2A; PIK3CB; PPP2R1A; PIK3C3; MAPK8; PRKD1; ALDH1A1; MAPK3; NRIP1; KRAS; MAPK13; PRKCD; GSTP1; MAPK9; NOS2A; ABCB1; AHR; PPP2CA; FTL; NFE2L2; PIK3C2A; PPARGC1A; MAPK14; TNF; RAF1; CREBBP; MAP2K2;
- 10 PIK3R1; PPP2R5C; MAP2K1; NFKB1; KEAP1; PRKCA; EIF2AK3; IL6; CYP1B1; HSP90AA1;

- [0351] SAPL/JNK Cellular Signaling disorders:** Genes linked to these disorders: PRKCE; IRAK1; PRKAA2; EIF2AK2; RAC1; ELK1; GRK6; MAPK1; GADD45A; RAC2; PLK1; AKT2; PIK3CA; FADD; CDK8; PIK3CB; PIK3C3; MAPK8; RIPK1; GNB2L1; IRS1;
- 15 MAPK3; MAPK10; DAXX; KRAS; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; TRAF2; TP53; LCK; MAP3K7; DYRK1A; MAP2K2; PIK3R1; MAP2K1; PAK3; CDC42; JUN; TTK; CSNK1A1; CRKL; BRAF; SGK;

- [0352] PPAR/RXR Cellular Signaling disorders:** Genes linked to these disorders: PRKAA2; EP300; INS; SMAD2; TRAF6; PPARA; FASN; RXRA; MAPK1; SMAD3; GNAS;
- 20 IKBKB; NCOR2; ABCA1; GNAQ; NFKB2; MAP3K14; STAT5B; MAPK8; IAS1; MAPK3; KRAS; RELA; PRKAA1; PPARGC1A; NCOA3; MAPK14; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; JAK2; CHUK; MAP2K1; NFKB1; TGFBA1; SMAD4; JUN; IL1R1; PRKCA; IL6; HSP90AA1; ADIPOO;

- [0353] NF-KB Cellular Signaling disorders:** Genes linked to these disorders: IRAK1;
- 25 EIF2AK2; EP300; INS; MYD88; PRKCZ; TRAF6; TBK1; AKT2; EGFR; IKBKB; PIK3CA; BTRC; NFKB2; MAP3K14; PIK3CB; PIK3C3; MAPK8; RIPK1; HDAC2; KRAS; RELA; PIK3C2A; TRAF2; TLR4; PDGFRB; TNF; INSR; LCK; IKBKG; RELB; MAP3K7; CREBBP; AKT1; PIK3R1; CHUK; PDGFRA; NFKB1; TLR2; BCL10; GSK3B; AKT3; TNFAIP3; IL1R1;

- 30 **[0354] Neuregulin Cellular Signaling disorders:** Genes linked to these disorders: ERBB4; PRKCE; ITGAM; ITGA5; PTEN; PRKCZ; ELK1; MAPK1; PTPN11; AKT2; EGFR; ERBB2; PRKCI; CDKN1B; STAT5B; PRKD1; MAPK3; ITGA1; KRAS; PRKCD; STAT5A; SRC; ITGB7; RAF1; ITGB1; MAP2K2; ADAM17; AKT1; PIK3R1; PDPK1; MAP2K1; ITGB3; EREG; FRAP1; PSEN1; ITGA2; MYC; NRG1; CRKL; AKT3; PRKCA; HS P90AA1;
- 35 RPS6KB1;

[0355] Wnt and Beta catenin Cellular Signaling disorders: Genes linked to these disorders: CD44; EP300; LRP6; DVL3; CSNK1E; GJA1; SMO; AKT2; PIN1; CDH1; BTRC; GNAQ; MARK2; PPP2R1A; WNT11; SRC; DKK1; PPP2CA; SOX6; SFRP2; ILK; LEF1; SOX9; TP53; MAP3K7; CREBBP; TCF7L2; AKT1; PPP2R5C; WNT5A; LAPS; CTNBNB1; 5 TGFBRI; CCND1; GSK3A; DVL1; APC; CDKN2A; MYC; CSNK1A1; GSK3B; AKT3; SOX2;

[0356] Insulin Receptor Signaling disorders: Genes linked to these disorders: PTEN; INS; EIF4E; PTPN1; PRKCZ; MAPK1; TSC1; PTPN11; AKT2; CBL; PIK3CA; PRKCI; PIK3CB; PIK3C3; MAPK8; IASI; MAPK3; TSC2; KRAS; EIF4EBP1; SLC2A4; PIK3C2A; 10 PPP1CC; INSR; RAF1; FYN; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; PDPK1; MAP2K1; GSK3A; FRAP1; CRKL; GSK3B; AKT3; FOXO1; SGK; RPS6KB1;

[0357] IL-6 Cellular Signaling disorders: Genes linked to these disorders: HSPB1; TRAF6; MAPKAPK2; ELK1; MAPK1; PTPN11; IKBKB; FOS; NFKB2; MAP3K14; MAPK8; MAPK3; MAPK10; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; ABCB1; 15 TRAF2; MAPK14; TNF; RAF1; IKBKG; RELB; MAP3K7; MAP2K2; IL8; JAK2; CHUK; STAT3; MAP2K1; NFKB1; CEBPB; JUN; IL1R1; SRF; IL6;

[0358] Hepatic Cholestasis Cellular Signaling disorders: Genes linked to these disorders: PRKCE; IRAK1; INS; MYD88; PRKCZ; TRAF6; PPARA; RXRA; IKBKB; PRKCI; NFKB2; MAP3K14; MAPK8; PRKD1; MAPK10; RELA; PRKCD; MAPK9; ABCB1; TRAF2; 20 TLR4; TNF; INSR; IKBKG; RELB; MAP3K7; IL8; CHUK; NR1H2; TJP2; NFKB1; ESR1; SREBF1; FGFR4; JUN; IL1R1; PRKCA; IL6;

[0359] IGF-1 Cellular Signaling disorders: Genes linked to these disorders: IGF1; PRKCZ; ELK1; MAPK1; PTPN11; NEDD4; AKT2; PIK3CA; PRKCI; PTK2; FOS; PIK3CB; PIK3C3; MAPK8; IGF1R; IRS1; MAPK3; IGFBP7; KRAS; PIK3C2A; YWHAZ; PXN; RAF1; 25 CASP9; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; IGFBP2; SFN; JUN; CYR61; AKT3; FOXO1; SRF; CTGF; RPS6KB1;

[0360] NRF2-mediated Oxidative Stress Response Signaling disorders: Genes linked to these disorders: PRKCE; EP300; SOD2; PRKCZ; MAPK1; SQSTM1; NQO1; PIK3CA; PRKCI; FOS; PIK3CB; PIK3C3; MAPK8; PRKD1; MAPK3; KRAS; PRKCD; GSTP1; 30 MAPK9; FTL; NFE2L2; PIK3C2A; MAPK14; RAF1; MAP3K7; CREBBP; MAP2K2; AKT1; PIK3R1; MAP2K1; PPIB; JUN; KEAP1; GSK3B; ATF4; PRKCA; EIF2AK3; HSP90AA1;

[0361] Hepatic Fibrosis/Hepatic Stellate Cell Activation Signaling disorders: Genes linked to these disorders: EDN1; IGF1; KDR; FLT1; SMAD2; FGFR1; MET; PGF; SMAD3; EGFR; FAS; CSF1; NFKB2; BCL2; MYH9; IGF1R; IL6R; RELA; TLR4; PDGFRB; TNF;

RELB; IL8; PDGFRA; NFKB1; TGFB1; SMAD4; VEGFA; BAX; IL1R1; CCL2; HGF; MMP1; STAT1; IL6; CTGF; MMP9;

[0362] PPAR Signaling disorders: Genes linked to these disorders: EP300; INS;

TRAF6; PPARA; RXRA; MAPK1; IKBKB; NCOR2; FOS; NFKB2; MAP3K14; STAT5B;

5 MAPK3; NRIP1; KRAS; PPARG; RELA; STAT5A; TRAF2; PPARGC1A; PDGFRB; TNF; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; CHUK; PDGFRA; MAP2K1; NFKB1; JUN; IL1R1; HSP90AA1;

[0363] Fc Epsilon RI Signaling disorders: Genes linked to these disorders: PRKCE;

RAC1; PRKCZ; LYN; MAPK1; RAC2; PTPN11; AKT2; PIK3CA; SYK; PRKCI; PIK3CB;

10 PIK3C3; MAPK8; PRKD1; MAPK3; MAPK10; KRAS; MAPK13; PRKCD; MAPK9;

PIK3C2A; BTK; MAPK14; TNF; RAF1; FYN; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; AKT3; VAV3; PRKCA;

[0364] G-Protein Coupled Receptor Signaling disorders: Genes linked to these

disorders: PRKCE; RAP1A; RGS16; MAPK1; GNAS; AKT2; IKBKB; PIK3CA; CREB1;

15 GNAQ; NFKB2; CAMK2A; PIK3CB; PIK3C3; MAPK3; KRAS; RELA; SRC; PIK3C2A;

RAF1; IKBKG; RELB; FYN; MAP2K2; AKT1; PIK3R1; CHUK; PDPK1; S TAT3 ; MAP2K1; NFKB1; BRAF; ATF4; AKT3; PRKCA;

[0365] Inositol Phosphate Metabolism Signaling disorders: Genes linked to these

disorders: PRKCE; IRAK1; PRKAA2; EIF2AK2; PTEN; GRK6; MAPK1; PLK1; AKT2;

20 PIK3CA; CDK8; PIK3CB; PIK3C3; MAPK8; MAPK3; PRKCD; PRKAA1; MAPK9; CDK2;

PIM1; PIK3C2A; DYRK1A; MAP2K2; PIP5K1A; PIK3R1; MAP2K1; PAK3; ATM; TTK; CSNK1A1; BRAF; SGK;

[0366] PDGF Signaling disorders: Genes linked to these disorders: EIF2AK2; ELK1;

ABL2; MAPK1; PIK3CA; FOS; PIK3CB; P IK3 C3 ; MAPK8; CAV1; ABL1 ; MAPK3;

25 KRAS; SRC; PIK3C2A; PDGFRB; RAF1; MAP2K2; JAK1; JAK2; PIK3R1; PDGFRA;

STAT3; SPHK1; MAP2K1; MYC; JUN; CRKL; PRKCA; SRF; STAT1; SPHK2 **VEGF**

Signaling disorders: Genes linked to these disorders: ACTN4; ROCK1; KDR; FLT1; ROCK2;

MAPK1; PGF; AKT2; PIK3CA; ARNT; PTK2; BCL2; PIK3CB; PIK3C3; BCL2L1; MAPK3;

KRAS; HIF1A; NOS3; PIK3C2A; PXN; RAF1; MAP2K2; ELAVL1; AKT1; PIK3R1;

30 MAP2K1; SFN; VEGFA; AKT3; FOXO1; PRKCA;

[0367] Natural Killer Cell Signaling disorders: Genes linked to these disorders:

PRKCE; RAC1; PRKCZ; MAPK1; RAC2; PTPN11; KIR2DL3; AKT2; PIK3CA; SYK;

PRKCI; PIK3CB; PIK3C3; PRKD1; MAPK3; KRAS; PRKCD; PTPN6; PIK3C2A; LCK;

RAF1; FYN; MAP2K2; PAK4; AKT1; PIK3R1; MAP2K1; PAK3; AKT3; VAV3; PRKCA;

- [0368] **Cell Cycle: G1/S Checkpoint Regulation Signaling disorders:** Genes linked to these disorders: HDAC4; SMAD3; SUV39H1; HDAC5; CDKN1B; BTRC; ATR; ABL1 ; E2F1; HDAC2; HDAC7A; RB1; HDAC11 ; HDAC9; CDK2; E2F2; HDAC3; TP53; CDKN1A; CCND1; E2F4; ATM; RBL2; SMAD4; CDKN2A; MYC; NRG1; GSK3B; RBL1; HDAC6;
- 5 [0369] **T Cell Receptor Signaling disorders:** Genes linked to these disorders: RAC1; ELK1; MAPK1; IKBKB; CBL; PIK3CA; FOS; NFKB2; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; RELA; PIK3C2A; BTK; LCK; RAF1; IKBKG; RELB; FYN; MAP2K2; PIK3R1; CHUK; MAP2K1; NFKB1; ITK; BCL10; JUN; VAV3;
- [0370] **Death Receptor disorders:** Genes linked to these disorders: CRADD; HSPB1; 10 BID; BIRC4; TBK1; IKBKB; FADD; FAS; NFKB2; BCL2; MAP3K14; MAPK8; RIPK1; CASP8; DAXX; TNFRSF10B; RELA; TRAF2; TNF; IKBKG; RELB; CASP9; CHUK; APAF1; NFKB1; CASP2; BIRC2; CASP3; BIRC3;
- [0371] **FGF Cell Signaling disorders:** Genes linked to these disorders: RAC1; FGFR1; MET; MAPKAPK2; MAPK1; PTPN11; AKT2; PIK3CA; CREB1; PIK3CB; PIK3C3; MAPK8; 15 MAPK3; MAPK13; PTPN6; PIK3C2A; MAPK14; RAF1; AKT1; PIK3R1; STAT3; MAP2K1; FGFR4; CRKL; ATF4; AKT3; PRKCA; HGF;
- [0372] **GM-CSF Cell Signaling disorders:** Genes linked to these disorders: LYN; ELK1; MAPK1; PTPN11; AKT2; PIK3CA; CAMK2A; STAT5B; PIK3CB; PIK3C3; GNB2L1; BCL2L1; MAPK3; ETS1; KRAS; RUNX1; PIM1; PIK3C2A; RAF1; MAP2K2; AKT1; JAK2; 20 PIK3R1; STAT3; MAP2K1; CCND1; AKT3; STAT1;
- [0373] **Amyotrophic Lateral Sclerosis Cell Signaling disorders:** Genes linked to these disorders: BID; IGF1; RAC1; BIRC4; PGF; CAPNS1; CAPN2; PIK3CA; BCL2; PIK3CB; PIK3C3; BCL2L1; CAPN1; PIK3C2A; TP53; CASP9; PIK3R1; RAB5A; CASP1; APAF1; VEGFA; BIRC2; BAX; AKT3; CASP3; BIRC3 PTPN1; MAPK1; PTPN11; AKT2; PIK3CA; 25 STAT5B; PIK3CB; PIK3C3; MAPK3; KRAS; SOCS1; STAT5A; PTPN6; PIK3C2A; RAF1; CDKN1A; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; FRAP1; AKT3; STAT1;
- [0374] **JAK/Stat Cell Signaling disorders:** Genes linked to these disorders: PTPN1; MAPK1; PTPN11; AKT2; PIK3CA; STAT5B; PIK3CB; PIK3C3; MAPK3; KRAS; SOCS1; 30 STAT5A; PTPN6; PIK3C2A; RAF1; CDKN1A; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; FRAP1; AKT3; STAT1;
- [0375] **Nicotinate and Nicotinamide Metabolism Cell Signaling disorders:** Genes linked to these disorders: PRKCE; IRAK1; PRKAA2; EIF2AK2; GRK6; MAPK1; PLK1; AKT2; CDK8; MAPK8; MAPK3; PRKCD; PRKAA1; PBEF1; MAPK9; CDK2; PIM1; 35 DYRK1A; MAP2K2; MAP2K1; PAK3; NT5E; TTK; CSNK1A1; BRAF; SGK;

- [0376] **Chemokine Cell Signaling disorders:** Genes linked to these disorders: CXCR4; ROCK2; MAPK1; PTK2; FOS; CFL1; GNAQ; CAMK2A; CXCL12; MAPK8; MAPK3; KRAS; MAPK13; RHOA; CCR3; SRC; PPP1CC; MAPK14; NOX1; RAF1; MAP2K2; MAP2K1; JUN; CCL2; PRKCA;
- 5 [0377] **IL-2 Cell Signaling disorders:** Genes linked to these disorders: ELK1; MAPK1; PTPN11; AKT2; PIK3CA; SYK; FOS; STAT5B; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; SOCS1; STAT5A; PIK3C2A; LCK; RAF1; MAP2K2; JAK1; AKT1; PIK3R1; MAP2K1; JUN; AKT3;
- [0378] **Synaptic Long Term Depression Signaling disorders:** Genes linked to these disorders: PRKCE; IGF1; PRKCZ; PRDX6; LYN; MAPK1; GNAS; PRKCI; GNAQ; PPP2R1A; IGF1R; PRKD1; MAPK3; KRAS; GRN; PRKCD; NOS3; NOS2A; PPP2CA; YWHAZ; RAF1; MAP2K2; PPP2R5C; MAP2K1; PRKCA;
- 10 [0379] **Estrogen Receptor Cell Signaling disorders:** Genes linked to these disorders: TAF4B; EP300; CARM1; PCAF; MAPK1; NCOR2; SMARCA4; MAPK3; NRIP1; KRAS; SRC; NR3C1; HDAC3; PPARGC1A; RBM9; NCOA3; RAF1; CREBBP; MAP2K2; NCOA2; MAP2K1; PRKDC; ESR1; ESR2;
- [0380] **Protein Ubiquitination Pathway Cell Signaling disorders:** Genes linked to these disorders: TRAF6; SMURF1; BIRC4; BRCA1; UCHL1; NEDD4; CBL; UBE2I; BTRC; HSPA5; USP7; USP10; FBXW7; USP9X; STUB1; USP22; B2M; BIRC2; PARK2; USP8;
- 20 [0381] **IL-10 Cell Signaling disorders:** Genes linked to these disorders: TRAF6; CCR1; ELK1; IKBKB; SP1; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; MAPK14; TNF; IKBKG; RELB; MAP3K7; JAK1; CHUK; STAT3; NFKB1; JUN; IL1R1; IL6;
- [0382] **VDR/RXR Activation Signaling disorders:** Genes linked to these disorders: PRKCE; EP300; PRKCZ; RXRA; GADD45A; HES1; NCOR2; SP1; PRKCI; CDKN1B; PRKD1; PRKCD; RUNX2; KLF4; YY1; NCOA3; CDKN1A; NCOA2; SPP1; LAPS; CEBPB; FOXO1; PRKCA;
- 25 [0383] **TGF-beta Cell Signaling disorders:** Genes linked to these disorders: EP300; SMAD2; SMURF1; MAPK1; SMAD3; SMAD1; FOS; MAPK8; MAPK3; KRAS; MAPK9; RUNX2; SERPINE1; RAF1; MAP3K7; CREBBP; MAP2K2; MAP2K1; TGFB1; SMAD4; JUN; SMAD5;
- [0384] **Toll-like Receptor Cell Signaling disorders:** Genes linked to these disorders: IRAK1; EIF2AK2; MYD88; TRAF6; PPARA; ELK1; IKBKB; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; TLR4; MAPK14; IKBKG; RELB; MAP3K7; CHUK; NFKB1;
- 35 [0385] **Toll-like Receptor Cell Signaling disorders:** Genes linked to these disorders: TLR2; JUN;

[0385] p38 MAPK Cell Signaling disorders: Genes linked to these disorders: HSPB1; IRAK1; TRAF6; MAPKAPK2; ELK1; FADD; FAS; CREB1; DDIT3; RPS6KA4; DAXX; MAPK13; TRAF2; MAPK14; TNF; MAP3K7; TGFBR1; MYC; ATF4; IL1R1; SRF; STAT1; and

5 **[0386] Neurolrophin/TRK Cell Signaling disorders:** Genes linked to these disorders: NTRK2; MAPK1; PTPN11; PIK3CA; CREB1; FOS; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; PIK3C2A; RAF1; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; CDC42; JUN; ATF4.

[0387] Other cellular dysfunction disorders linked to a genetic modification are contemplated herein for example, FXR/RXR Activation, Synaptic Long Term Potentiation, Calcium Signaling EGF Signaling, Hypoxia Signaling in the Cardiovascular System, LPS/IL-1 Mediated Inhibition of RXR Function LXR/RXR Activation, Amyloid Processing, IL-4 Signaling, Cell Cycle: G2/M DNA Damage Checkpoint Regulation, Nitric Oxide Signaling in the Cardiovascular System Purine Metabolism, cAMP-mediated Signaling, Mitochondrial Dysfunction Notch Signaling Endoplasmic Reticulum Stress Pathway Pyrimidine Metabolism, Parkinson's Signaling Cardiac & Beta Adrenergic Signaling Glycolysis/ Gluconeogenesis Interferon Signaling Sonic Hedgehog Signaling Glycerophospholipid Metabolism, Phospholipid Degradation, Tryptophan Metabolism Lysine Degradation Nucleotide Excision Repair Pathway, Starch and Sucrose Metabolism, Aminosugars Metabolism Arachidonic Acid Metabolism, Circadian Rhythm Signaling, Coagulation System Dopamine Receptor Signaling, Glutathione Metabolism Glycerolipid Metabolism Linoleic Acid Metabolism Methionine Metabolism Pyruvate Metabolism Arginine and Praline Metabolism, Eicosanoid Signaling Fructose and Mannose Metabolism, Galactose Metabolism Stilbene, Coumarine and Lignin Biosynthesis Antigen Presentation Pathway, Biosynthesis of Steroids Butanoate Metabolism Citrate Cycle Fatty Acid Metabolism Glycerophospholipid Metabolism, Histidine Metabolism Inositol Metabolism Metabolism of Xenobiotics by Cytochrome p450, Methane Metabolism, Phenylalanine Metabolism, Propanoate Metabolism Selenoamino Acid Metabolism Sphingolipid Metabolism Aminophosphonate Metabolism, Androgen and Estrogen Metabolism Ascorbate and Aldarate Metabolism, Bile Acid Biosynthesis Cysteine Metabolism Fatty Acid Biosynthesis Glutamate Receptor Signaling, NRF2-mediated, Oxidative Stress Response Pentose Phosphate Pathway, Pentose and Glucuronate Interconversions, Retinol Metabolism Riboflavin Metabolism Tyrosine Metabolism Ubiquinone Biosynthesis Valine, Leucine and Isoleucine Degradation Glycine, Serine and Threonine Metabolism Lysine Degradation Pain/Taste, or Mitochondrial Function Developmental Neurology or combinations thereof.

[0388] Additional exemplary genetic diseases or disorders and associated information are available on the world wide web at kumc.edu/gec/support, genome.gov/10001200, and

ncbi.nlm.nih.gov/books/NBK22183/. Additional exemplary genetic diseases or disorders, associated genetic mutations, and gene therapy approaches to treat genetic diseases or disorders are described in International (PCT) Publication Nos. WO2013/126794, WO2013/163628, WO2015/048577, WO2015/070083, WO2015/089354, WO2015/134812, WO2015/138510, WO2015/148670, WO2015/148860, WO2015/148863, WO2015/153780, WO2015/153789, and WO2015/153791, U.S. Patent Nos. 89,568,28, 83,83,604, 9,273,296, 8,859,597, and 9,255,130, and U.S. Patent Publication Nos. 2009/0222937, 2009/0271881, 2010/0229252, 2010/0311124, 2011/0016540, 2011/0023139, 2011/0023144, 2011/0023145, 2011/0023146, 2011/0023153, 2011/0091441, 2012/0159653, and 2013/0145487.

B. Immune Cell Engineering

[0389] It is understood that the engineered, non-naturally occurring system and CRISPR expression system disclosed herein can be used to engineer an immune cell. Immune cells include but are not limited to lymphocytes (*e.g.*, B lymphocytes or B cells, T lymphocytes or T cells, and natural killer cells), myeloid cells (*e.g.*, monocytes, macrophages, eosinophils, mast cells, basophils, and granulocytes), and the stem and progenitor cells that can differentiate into these cell types (*e.g.*, hematopoietic stem cells, hematopoietic progenitor cells, and lymphoid progenitor cells). The cells can include autologous cells derived from a subject to be treated, or alternatively allogenic cells derived from a donor. Cells are treated with a non-naturally occurring CRISPR-Cas system can be treated with a system that includes a modified targeter nucleic acid, a modified modulator nucleic acid, or both, as described in sections IA, IA1, and IB. Systems that include Cas proteins such as Cas nucleases are described in section IC. RNPs, Expression systems, donor templates, efficiency and specificity, and multiplex methods are described in sections IIA, B, C, D, and E, respectively.

[0390] It is understood that the guide nucleic acid, the engineered, non-naturally occurring system, and the CRISPR expression system disclosed herein can be used to treat any disease or disorder that can be improved by editing or modifying a target sequence; exemplary genes containing target sequences to be modified for therapeutic purposes include ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALP NR, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWF1, CD3D, or NLRC5 gene in a cell.

[0391] In certain embodiments, the immune cell is a T cell, which can be, for example, a cultured T cell, a primary T cell, a T cell from a cultured T cell line (*e.g.*, Jurkat, SupTi), or a T cell obtained from a mammal, for example, from a subject to be treated. If obtained from a mammal, the T cell can be obtained from numerous sources, including but not limited to blood, bone marrow, lymph node, the thymus, or other tissues or fluids. T cells can also be enriched or purified. The T cell can be any type of T cell and can be of any developmental stage, including but not limited to, CD4⁺/CD8⁺ double positive T cells, CD4⁺ helper T cells (*e.g.*, Th1 and Th2 cells), CD8⁺ T cells (*e.g.*, cytotoxic T cells), tumor infiltrating lymphocytes (TILs), memory T cells (*e.g.*, central memory T cells and effector memory T cells), regulatory T cells, naive T cells, and the like.

[0392] In certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to express an exogenous gene. For example, in certain embodiments, an engineered CRISPR system disclosed herein may be used to engineer an immune cell to express an exogenous gene. For example, in certain embodiments, the guide nucleic acid, the engineered, non-naturally occurring system, and the CRISPR expression system disclosed herein may be used to engineer an immune cell to express an exogenous gene at the locus of a ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALPNR, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWIF1, CD3D, or NLRC5 gene. For example, in certain embodiments, an engineered CRISPR system disclosed herein may catalyze DNA cleavage at a gene locus, allowing for site-specific integration of the exogenous gene at the gene locus by HDR.

[0393] In certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to express a chimeric antigen receptor (CAR), *i.e.*, the T cell comprises an exogenous nucleotide sequence encoding a CAR. As used herein, the term “chimeric antigen receptor” or “CAR” can include any artificial receptor including an antigen-specific binding moiety and one or more signaling chains derived from an immune receptor. CARs can comprise a single chain fragment variable (scFv) of an antibody specific for an antigen coupled via hinge and transmembrane regions to cytoplasmic domains of T cell signaling molecules, *e.g.*, a T cell costimulatory domain (*e.g.*, from CD28, CD137, OX40, ICOS, or CD27) in tandem with a T cell triggering domain (*e.g.*, from CD3 ζ). **A T cell expressing a chimeric antigen receptor is referred to as a CAR T cell.** Exemplary CAR T cells include CD19 targeted CTL019 cells (see, Grupp *et al.* (2015) BLOOD, 126: 4983), 19-28z cells (see, Park *et al.* (2015) J. CLIN. ONCOL., 33: 7010), and KTE-C19 cells

(see, Locke *et al.* (2015) BLOOD, 126: 3991). Additional exemplary CAR T cells are described in U.S. Patent Nos. 8,399,645, 8,906,682, 7,446,190, 9,181,527, 9,272,002, 9,266,960, 10,253,086, 10,808,035, and 10,640,569, and International (PCT) Publication Nos. WO2013/142034, WO2015/120180, WO2015/188141, WO2016/120220, and WO2017/040945. Exemplary approaches to express CARs using CRISPR systems are described in Hale *et al.* (2017) MOL THER METHODS CLIN DEV., 4: 192, MacLeod *et al.* (2017) MOL THER, 25: 949, and Eyquem *et al.* (2017) NATURE, 543: 113.

[0394] In certain embodiments, an immune cell, *e.g.*, a T cell, binds an antigen, *e.g.*, a cancer antigen, through an endogenous T cell receptor (TCR). In certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to express an exogenous TCR, *e.g.*, an exogenous naturally occurring TCR or an exogenous engineered TCR. T cell receptors comprise two chains referred to as the α - and β -chains, that combine on the surface of a T cell to form a heterodimeric receptor that can recognize MHC-restricted antigens. Each of α - and β - chain comprises a constant region and a variable region. Each variable region of the α - and β - chains defines three loops, referred to as complementary determining regions (CDRs) known as CDR₁, CDR₂, and CDR₃ that confer the T cell receptor with antigen binding activity and binding specificity.

[0395] In certain embodiments, a CAR or TCR binds a cancer antigen selected from B-cell maturation antigen (BCMA), mesothelin, prostate specific membrane antigen (PSMA), prostate stem cell antigen (PCSA), carbonic anhydrase IX (CAIX), carcinoembryonic antigen (CEA), CD5, CD7, CD10, CD19, CD20, CD22, CD30, CD33, CD34, CD38, CD41, CD44, CD49f, CD56, CD70, CD74, CD123, CD133, CD138, epithelial glycoprotein2 (EGP 2), epithelial glycoprotein-40 (EGP-40), epithelial cell adhesion molecule (EpCAM), receptor-type tyrosine-protein kinase (FLT3), folate-binding protein (FBP), fetal acetylcholine receptor (AChR), folate receptor- α and β (FR α and β), Ganglioside G2 (GD2), Ganglioside G3 (GD3), epidermal growth factor receptor 2 (HER-2/ERB2), epidermal growth factor receptor VIII (EGFRvIII), ERB3, ERB4, human telomerase reverse transcriptase (hTERT), Interleukin-13 receptor subunit α -2 (IL-13R α 2), K-light chain, kinase insert domain receptor (KDR), Lewis A (CA19.9), Lewis Y (LeY), LI cell adhesion molecule (LICAM), melanoma-associated antigen 1 (melanoma antigen family A1, MAGE-A1), Mucin 16 (MUC-16), Mucin 1 (MUC-1; *e.g.*, a truncated MUC-1), KG2D ligands, cancer-testis antigen NY-ESO-1, oncofetal antigen (h5T4), tumor-associated glycoprotein 72 (TAG-72), vascular endothelial growth factor R2 (VEGF-R2), Wilms tumor protein (WT-1), type 1 tyrosine-protein kinase transmembrane receptor (ROR1), B7-H3 (CD276), B7-H6 (Nkp30), Chondroitin sulfate proteoglycan-4 (CSPG4), DNAX Accessory Molecule (DNAM-1), Ephrin type A Receptor 2 (EpHA2), Fibroblast Associated Protein (FAP), Gp100/HLA-A2, Glypican 3 (GPC3), HA-IH, HERK-V, IL-1 IR α , Latent

Membrane Protein 1 (LMP1), Neural cell-adhesion molecule (N-CAM/CD56), and Trail Receptor (TRAIL-R).

[0396] In certain embodiments, a genetic loci suitable for insertion of a exogenous gene is intergenic. In certain embodiments, introduction of an exogenous DNA into the genetic loci mediated by the action on a nucleic acid-guided nucleases, *e.g.*, a Cas protein bound to a suitable gRNA, results in no phenotypic side effects to the host cell. In certain embodiments, the exogenous gene and or genes inserted into the genetic loci demonstrate robust transcription. In certain embodiments, the transcription of the exogenous gene or genes doesn't affect the expression of neighboring upstream or down-stream genes. In certain embodiments, robust transcription of the exogenous gene or genes in the genetic loci is maintained in progeny after proliferation of the host cell.

[0397] Genetic loci suitable for insertion of a CAR- or exogenous TCR-encoding sequence include but are not limited to safe harbor loci (*e.g.*, the AAVS1 locus), TCR subunit loci (*e.g.*, the **TCR α constant (TRAC) locus**), and other loci associated with certain advantages (*e.g.*, the CCR5 locus, the inactivation of which may prevent or reduce HIV infection). It is understood that insertion in the TRAC locus reduces tonic CAR signaling and enhances T cell potency (see, Eyquem *et al.* (2017) NATURE, 543: 113). Furthermore, inactivation of the endogenous TRAC gene may reduce a graft-versus-host disease (GVHD) response, thereby allowing use of allogeneic T cells as starting materials for preparation of CAR-T cells. Accordingly, in certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to have reduced expression of an endogenous TCR or TCR subunit, *e.g.*, **TCR α subunit constant (TRAC)**. The cell may be engineered to have partially reduced or no expression of the endogenous TCR or TCR subunit. For example, in certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have less than 80% (*e.g.*, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, or less than 5%) of the expression of the endogenous TCR or TCR subunit relative to a corresponding unmodified or parental cell. In certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have no detectable expression of the endogenous TCR or TCR subunit. Exemplary approaches to reduce expression of TCRs using CRISPR systems are described in U.S. Patent No. 9,181,527, Liu *et al.* (2017) CELL RES, 27: 154, Ren *et al.* (2017) CLIN CANCER RES, 23: 2255, Cooper *et al.* (2018) LEUKEMIA, 32: 1970, and Ren *et al.* (2017) ONCOTARGET, 8: 17002.

[0398] It is understood that certain immune cells, such as T cells, also express major histocompatibility complex (MHC) or human leukocyte antigen (HLA) genes, and inactivation of these endogenous gene may reduce a GVHD response, thereby allowing use of allogeneic T cells as starting materials for preparation of CAR-T cells. Accordingly, in certain embodiments,

an immune cell, *e.g.*, a T cell, is engineered to have reduced expression of one or more endogenous class I or class II MHCs or HLAs (*e.g.*, beta 2-microglobulin (B2M), class II major histocompatibility complex transactivator (CIITA), HLA-E, and/or HLA-G). The cell may be engineered to have partially reduced or no expression of an endogenous MHC or HLA. For example, in certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have less than less than 80% (*e.g.*, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, or less than 5%) of the expression of endogenous MHC (*e.g.*, B2M, CIITA, HLA-E, or HLA-G) relative to a corresponding unmodified or parental cell. In certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have no detectable expression of an endogenous MHC (*e.g.*, B2M, CIITA, HLA-E, or HLA-G). Exemplary approaches to reduce expression of MHCs using CRISPR systems are described in Liu *et al.* (2017) CELL RES, 27: 154, Ren *et al.* (2017) CLIN CANCER RES, 23: 2255, and Ren *et al.* (2017) ONCOTARGET, 8: 17002.

[0399] Other genes that may be inactivated to reduce a GVHD response include but are not limited to CD3, CD52, and deoxycytidine kinase (DCK). For example, inactivation of DCK may render the immune cells (*e.g.*, T cells) resistant to purine nucleotide analogue (PNA) compounds, which are often used to compromise the host immune system in order to reduce a GVHD response during an immune cell therapy. In certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have less than less than 80% (*e.g.*, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, or less than 5%) of the expression of endogenous CD52 or DCK relative to a corresponding unmodified or parental cell.

[0400] In certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to have reduced expression of an endogenous gene. For example, in certain embodiments, an engineered CRISPR system disclosed herein may be used to engineer an immune cell to have reduced expression of an endogenous gene. For example, in certain embodiments, an engineered CRISPR system disclosed herein may result in DNA cleavage at a gene locus, thereby inactivating the targeted gene. In other embodiments, an engineered CRISPR system disclosed herein may be fused to an effector domain (*e.g.*, a transcriptional repressor or histone methylase) to reduce the expression of the target gene.

[0401] It is understood that the activity of an immune cell (*e.g.*, T cell) may be enhanced by inactivating or reducing the expression of an immune suppressor such as an immune checkpoint protein. Accordingly, in certain embodiments, an immune cell, *e.g.*, a T cell, is engineered to have reduced expression of an immune checkpoint protein. Exemplary immune checkpoint proteins expressed by wild-type T cells include but are not limited to PDCD1 (PD-1), CSF2, CTLA4, ADORA2A (A2AR), B7-H3, B7-H4, BTLA, KIR, LAG3, HAVCR2 (TIM3),

TIGIT, VISTA, PTPN6 (SHP-1), and FAS. The cell may be modified to have partially reduced or no expression of the immune checkpoint protein. For example, in certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have less than 80% (*e.g.*, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, or less than 5%) of the expression of the immune checkpoint protein relative to a corresponding unmodified or parental cell. In certain embodiments, the immune cell, *e.g.*, a T cell, is engineered to have no detectable expression of the immune checkpoint protein. Exemplary approaches to reduce expression of immune checkpoint proteins using CRISPR systems are described in International (PCT) Publication No. WO2017/017184, Cooper *et al.* (2018) LEUKEMIA, 32: 1970, Su *et al.* (2016) ONCOIMMUNOLOGY, 6: e1249558, and Zhang *et al.* (2017) FRONT MED, 11: 554.

[0402] The immune cell can also be engineered to express an exogenous protein (besides an antigen-binding protein described above) at the locus of a human ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALPNI, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWF1, CD3D, or NLRC5 gene.

[0403] In certain embodiments, an immune cell, *e.g.*, a T cell, is modified to express a dominant-negative form of an immune checkpoint protein. In certain embodiments, the dominant-negative form of the checkpoint inhibitor can act as a decoy receptor to bind or otherwise sequester the natural ligand that would otherwise bind and activate the wild-type immune checkpoint protein. Examples of engineered immune cells, for example, T cells containing dominant-negative forms of an immune suppressor are described, for example, in International (PCT) Publication No. WO2017/040945.

[0404] In certain embodiments, an immune cell, *e.g.*, a T cell, is modified to express a gene (*e.g.*, a transcription factor, a cytokine, or an enzyme) that regulates the survival, proliferation, activity, or differentiation (*e.g.*, into a memory cell) of the immune cell. In certain embodiments, the immune cell is modified to express TET2, FOXO1, IL-12, IL-15, IL-18, IL-21, IL-7, GLUT1, GLUT3, HK1, HK2, GAPDH, LDHA, PDK1, PKM2, PFKFB3, PGK1, ENO1, GYS1, and/or ALDOA. In certain embodiments, the modification is an insertion of a nucleotide sequence encoding the protein operably linked to a regulatory element. In certain embodiments, the modification is a substitution of a single nucleotide polymorphism (SNP) site in the endogenous gene. In certain embodiments, an immune cell, *e.g.*, a T cell, is modified to express a variant of a gene, for example, a variant that has greater activity than the respective

wild-type gene. In certain embodiments, the immune cell is modified to express a variant of CARD11, CD247, IL7R, LCK, or PLCG1. For example, certain gain-of-function variants of IL7R were disclosed in Zenatti *et al.*, (2011) NAT. GENET. 43(10):932-39. The variant can be expressed from the native locus of the respective wild-type gene by delivering an engineered system described herein for targeting the native locus in combination with a donor template that carries the variant or a portion thereof.

[0405] In certain embodiments, an immune cell, *e.g.*, a T cell, is modified to express a protein (*e.g.*, a cytokine or an enzyme) that regulates the microenvironment that the immune cell is designed to migrate to (*e.g.*, a tumor microenvironment). In certain embodiments, the immune cell is modified to express CA9, CA12, a V-ATPase subunit, NHE1, and/or MCT-1.

[0406] In certain embodiments, provided is a method for treatment of a disease, *e.g.*, a cancer, by administering to a subject suffering from the disease an effective amount of T cells modified to express a CAR specific to the disease using the modified guide nucleic acids and CRISPR-Cas systems described herein, *e.g.*, in sections IA, IA1, IB, IC, and IVB. In certain embodiments, the T cells are autologous cells removed from the subject, treated to modify genomic DNA to express CAR, expanded, and administered to the subject; in certain embodiments, the T cells are allogeneic T cells that have been treated to modify genomic DNA to express CAR. In certain embodiments, the disease is a blood cancer, such as leukemia or lymphoma; in certain embodiments the disease is a solid tumor cancer.

V. Kits

[0407] It is understood that the engineered, non-naturally occurring system, the CRISPR expression system, and/or the library disclosed herein can be packaged in a kit suitable for use by a medical provider, researcher, and the like. Accordingly, in another aspect, the invention provides kits containing any one or more of the elements disclosed in the above systems, libraries, methods, and compositions. In certain embodiments, the kit comprises an engineered, non-naturally occurring system as disclosed herein and instructions for using the kit. The instructions may be specific to the applications and methods described herein. In certain embodiments, one or more of the elements of the system are provided in a solution. In certain embodiments, one or more of the elements of the system are provided in lyophilized form, and the kit further comprises a diluent. Elements may be provided individually or in combinations, and may be provided in any suitable container, such as a vial, a bottle, a tube, or immobilized on the surface of a solid base (*e.g.*, chip or microarray). In certain embodiments, the kit comprises one or more of the nucleic acids and/or proteins described herein. In certain embodiments, the kit provides all elements of the systems of the invention.

[0408] In certain embodiments of a kit comprising the engineered, non-naturally occurring system, the targeter nucleic acid and the modulator nucleic acid, where one or more nucleotides at or near the 3' and/or 5' end of either or both, is modified, are provided in separate containers. In other embodiments, the targeter nucleic acid and the modulator nucleic acid are pre-complexed, and the complex is provided in a single container. In certain embodiments, the kit comprises a Cas protein or a nucleic acid comprising a regulatory element operably linked to a nucleic acid encoding a Cas protein provided in a separate container. In other embodiments, the kit comprises a Cas protein pre-complexed with the targeter nucleic acid and the modulator nucleic acid, and the complex is provided in a single container.

[0409] In order to target multiple target nucleotide sequences, *e.g.*, for use in a screening or selection process, a kit may be provided comprising multiple targeter nucleic acids. Accordingly, in certain embodiments, the kit comprises a plurality of targeter nucleic acids as disclosed herein (*e.g.*, in separate tubes or immobilized on the surface of a solid base such as a chip or a microarray), optionally one or more modulator nucleic acids as disclosed herein, and optionally a Cas protein or a regulatory element operably linked to a nucleic acid encoding a Cas protein as disclosed herein. Such kits are useful for identifying a targeter nucleic acid with the highest efficiency and/or specificity to target a given gene, for identifying a gene implicated in a physiological or pathological pathway, or for engineering a cell to achieve desired functionality in a multiplex assay. In certain embodiments, the kit further comprises one or more donor templates provided in one or more separate containers. In certain embodiments, the kit comprises a plurality of donor templates as disclosed herein (*e.g.*, in separate tubes or immobilized on the surface of a solid base such as a chip or a microarray), one or more targeter nucleic acids disclosed herein, and one or more modulator nucleic acids as disclosed herein, and optionally a Cas protein or a regulatory element operably linked to a nucleic acid encoding a Cas protein as disclosed herein. Such kits are useful for identifying a donor template that introduces optimal genetic modification in a multiplex assay. The CRISPR expression systems as disclosed herein are also suitable for use in a kit.

[0410] In certain embodiments, a kit further comprises one or more reagents and/or buffers for use in a process utilizing one or more of the elements described herein. Reagents may be provided in any suitable container and may be provided in a form that is usable in a particular assay, or in a form that requires addition of one or more other components before use (*e.g.*, in concentrate or lyophilized form). A buffer may be a reaction or storage buffer, including but not limited to a sodium carbonate buffer, a sodium bicarbonate buffer, a borate buffer, a Tris buffer, a MOPS buffer, a HEPES buffer, and combinations thereof. In some embodiments, the buffer is alkaline. In certain embodiments, the buffer has a pH from 6-9, 6.5-8.5, 7-8, 6.5-7.5, 6-8, 7.5-8.5,

7-9, 6.5-9.5, 6-10, 8-9, 7.5-9.5, 7-10, for example 7-8, such as 7.5. In certain embodiments, the kit further comprises a pharmaceutically acceptable carrier. In certain embodiments, the kit further comprises one or more devices or other materials for administration to a subject.

[0411] Throughout the description, where compositions are described as having,

5 including, or comprising specific components, or where processes and methods are described as having, including, or comprising specific steps, it is contemplated that, additionally, there are compositions of the present invention that consist essentially of, or consist of, the recited components, and that there are processes and methods according to the present invention that consist essentially of, or consist of, the recited processing steps.

10 [0412] In the application, where an element or component is said to be included in and/or selected from a list of recited elements or components, it should be understood that the element or component can be any one of the recited elements or components, or the element or component can be selected from a group consisting of two or more of the recited elements or components.

15 [0413] Further, it should be understood that elements and/or features of a composition or a method described herein can be combined in a variety of ways without departing from the spirit and scope of the present invention, whether explicit or implicit herein. For example, where reference is made to a particular compound, that compound can be used in various embodiments of compositions of the present invention and/or in methods of the present invention, unless
20 otherwise understood from the context. In other words, within this application, embodiments have been described and depicted in a way that enables a clear and concise application to be written and drawn, but it is intended and will be appreciated that embodiments may be variously combined or separated without parting from the present teachings and invention(s). For example, it will be appreciated that all features described and depicted herein can be applicable to all
25 aspects of the invention(s) described and depicted herein.

[0414] The terms “a” and “an” and “the” and similar references in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. For example, the term “a cell” can include a plurality of cells, including mixtures
30 thereof. Where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

[0415] It should be understood that the expression “at least one of” includes individually each of the recited objects after the expression and the various combinations of two or more of the recited objects unless otherwise understood from the context and use. The expression

“and/or” in connection with three or more recited objects should be understood to have the same meaning unless otherwise understood from the context.

[0416] The use of the term “include,” “includes,” “including,” “have,” “has,” “having,” “contain,” “contains,” or “containing,” including grammatical equivalents thereof, should be understood generally as open-ended and non-limiting, for example, not excluding additional unrecited elements or steps, unless otherwise specifically stated or understood from the context.

[0417] Where the use of the term “about” is before a quantitative value, the present invention also includes the specific quantitative value itself, unless specifically stated otherwise. As used herein, the term “about” refers to a $\pm 10\%$ variation from the nominal value unless otherwise indicated or inferred.

[0418] It should be understood that the order of steps or order for performing certain actions is immaterial so long as the present invention remain operable. Moreover, two or more steps or actions may be conducted simultaneously.

[0419] The use of any and all examples, or exemplary language herein, for example, “such as” or “including,” is intended merely to illustrate better the present invention and does not pose a limitation on the scope of the invention unless claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the present invention.

VI. Embodiments

[0420] In embodiment 1 provided herein is a composition comprising a synthetic guide nucleic acid (gNA) comprising (i) a targeter nucleic acid comprising: (a) a spacer sequence configured to hybridize with a target nucleotide sequence, and (b) a targeter stem sequence; and (ii) a modulator nucleic acid comprising (a) a modulator stem sequence complementary to the targeter stem sequence, and (b) a 5' sequence; wherein either the targeter nucleic acid or the modulator nucleic acid, or both, comprise one or more modified nucleotides at or near its 3' end, if present, at or near its 5' end, if present, or both. In embodiment 2 provided herein is the composition of embodiment 1 wherein the synthetic gNA is a single polynucleotide. In embodiment 3 provided herein is the composition of embodiment 1 wherein the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids. In embodiment 4 provided herein is the composition of any one of embodiments 1 through 3 wherein some or all of the gNA is RNA, e.g provided herein is a gRNA. In embodiment 5 provided herein is the composition of embodiment 4 wherein at least 50% of the nucleic acid is RNA. In embodiment 6 provided herein is the composition of embodiment 4 wherein at least 70% of the nucleic acid is RNA. In embodiment 7 provided herein is the composition of embodiment 4 wherein at least 90% of the

nucleic acid is RNA. In embodiment 8 provided herein is the composition of embodiment 4 wherein at least 100% of the nucleic acid is RNA. In embodiment 9 provided herein is the composition of any one of embodiments 1 through 8 wherein the modified nucleotide comprises a chemical modification. In embodiment 10 provided herein is the composition of any one of
5 embodiments 1 through 9 wherein the stem sequences are between one and ten nucleotides in length. In embodiment 11 provided herein is the composition of any one of embodiments 1 through 9 wherein the stem sequences are between two and nice nucleotides in length. In embodiment 12 provided herein is the composition of any one of embodiments 1 through 9 wherein the stem sequences are between four and six nucleotides in length. In embodiment 13
10 provided herein is the composition of any one of embodiments 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 80% sequence complementarity. In embodiment 14 provided herein is the composition of any one of embodiments 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 90% sequence complementarity. In embodiment 15 provided herein is the
15 composition of any one of embodiments 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 95% sequence complementarity. In embodiment 16 provided herein is the composition of any one of embodiments 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share 100% sequence complementarity. In embodiment 17 provided herein is the composition of embodiment 2
20 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 5' end of the gNA. In embodiment 18 provided herein is the composition of embodiment 17 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 5' end. In embodiment 19 provided herein is the composition of embodiment 17 wherein the
25 chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 5' end. In embodiment 20 provided herein is the composition of embodiment 19 wherein the chemical modification comprises one to ten phopshorothioate modifications within ten internucleotide linkages of the 5' end. In embodiment 21 provided herein is the composition of embodiment 19 wherein the chemical modification comprises nine
30 phosphorothioate modifications to the final nine internucleotide linkages of the 5' end. In embodiment 22 provided herein is the composition of embodiment 17 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 5' end. In embodiment 23 provided herein is the composition of any one of embodiments 17 through 21 wherein the chemical modification comprises two 2'-O-
35 methoxy modifications to the final two nucleotides of the 5' end. In embodiment 24 provided

herein is the composition of any one of embodiments 17 through 21 wherein the chemical modification comprises one 2'-O-methoxy modification to the 5' terminal nucleotide. In embodiment 25 provided herein is the composition of any one of embodiments 17 through 24 wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide. In embodiment 26 provided herein is the composition of embodiment 17 wherein the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037. In embodiment 27 provided herein is the composition of embodiment 2 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 3' end of the gNA. In embodiment 28 provided herein is the composition of any one of embodiments 17 through 25 wherein the chemical modification further comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 3' end of the gNA. In embodiment 29 provided herein is the composition of embodiment 27 or embodiment 28 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3' end. In embodiment 30 provided herein is the composition of embodiment 27 or embodiment 28 wherein the chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 3' end. In embodiment 31 provided herein is the composition of embodiment 30 wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 3' end. In embodiment 32 provided herein is the composition of embodiment 30 wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 3' end. In embodiment 33 provided herein is the composition of any one of embodiments 27 through 32 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 3' end. In embodiment 34 provided herein is the composition of any one of embodiments 27 through 32 wherein the chemical modification comprises two 2'-O-methoxy modifications to the final two nucleotides of the 3' end. In embodiment 35 provided herein is the composition of any one of embodiments 27 through 32 wherein the chemical modification comprises one 2'-O-methoxy modification to the last nucleotide of the 3' end. In embodiment 36 provided herein is the composition of any one of embodiments 27 through 35 wherein the chemical modification comprises a propanediol modification to the 3' terminal nucleotide. In embodiment 37 provided herein is the composition of embodiment 3 wherein the modulator nucleic acid is unmodified. In embodiment 38 provided herein is the composition of embodiment 3 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or

near the 5' end of the modulator nucleic acid. In embodiment 39 provided herein is the composition of embodiment 38 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 5' end. In embodiment 40 provided herein is the composition of embodiment 38 wherein the chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 5' end. In embodiment 41 provided herein is the composition of embodiment 40 wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 5' end. In embodiment 42 provided herein is the composition of embodiment 40 wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 5' end. In embodiment 43 provided herein is the composition of any one of embodiments 38 through 42 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 5' end. In embodiment 44 provided herein is the composition of any one of embodiments 38 through 42 wherein the chemical modification comprises two 2'-O-methoxy modifications to the final two nucleotides of the 5' end. In embodiment 45 provided herein is the composition of any one of embodiments 38 through 42 wherein the chemical modification comprises one 2'-O-methoxy modification to the 5' end nucleotide. In embodiment 46 provided herein is the composition of any one of embodiments 38 through 45 wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide. In embodiment 47 provided herein is the composition of embodiment 38 wherein the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037. In embodiment 48 provided herein is the composition of any one of embodiments 3 and embodiments 38 through 46 wherein the modification comprises a chemical modification to one or more nucleotides or internucleotide linkages at or near the 3' end of the modulator nucleic acid. In embodiment 49 provided herein is the composition of embodiment 48 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3' end. In embodiment 50 provided herein is the composition of embodiment 48 wherein the chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 3' end. In embodiment 51 provided herein is the composition of embodiment 50 wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 3' end. In embodiment 52 provided herein is the composition of embodiment 50 wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 3' end. In

embodiment 53 provided herein is the composition of any one of embodiments 3 and
embodiments 38 through 52 wherein the chemical modification comprises one to five 2'-O-
methoxy modifications to one or more nucleotides within five nucleotides of the 3' end. In
embodiment 54 provided herein is the composition of any one of embodiments 3 and
5 embodiments 38 through 52 wherein the chemical modification comprises two 2'-O-methoxy
modifications to the final two nucleotides of the 3' end. In embodiment 55 provided herein is the
composition of any one of embodiments 3 and embodiments 38 through 52 wherein the chemical
modification comprises one 2'-O-methoxy modification to the last nucleotide of the 3' end. In
embodiment 56 provided herein is the composition of any one of embodiments 48 through 55
10 wherein the chemical modification comprises a propanediol modification to the 3' terminal
nucleotide. In embodiment 57 provided herein is the composition of any one of embodiments 3
and 38 through 56 wherein the targeter nucleic acid is unmodified. In embodiment 58 provided
herein is the composition of any one of embodiments 37 through 56 wherein the chemical
modification comprises a chemical modification in one or more nucleotides or internucleotide
15 linkages at or near the 3' end of the targeter nucleic acid. In embodiment 59 provided herein is
the composition of embodiment 58 wherein the chemical modification comprises a chemical
modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3'
end. In embodiment 60 provided herein is the composition of embodiment 58 wherein the
chemical modification comprises one to five 2'-O-methoxy-3'-phosphorothioate modifications
20 within five nucleotides of the 3' end. In embodiment 61 provided herein is the composition of
embodiment 58 wherein the chemical modification comprises two 2'-O-methoxy-3'-
phosphorothioate modifications to the final two nucleotides of the 3' end. In embodiment 62
provided herein is the composition of embodiment 58 wherein the chemical modification
comprises one through nine 2'-fluoro modifications to the final nine nucleotides of the 3' end. In
25 embodiment 63 provided herein is the composition of embodiment 58 wherein the chemical
modification comprises five 2'-fluoro modifications to the final five nucleotides of the 3' end. In
embodiment 64 provided herein is the composition of any one of embodiments 58 through 63
wherein the chemical modification comprises a propanediol modification to the 3' terminal
nucleotide. In embodiment 65 provided herein is the composition of any one of embodiments 3,
30 embodiments 37 through 56, and embodiments 58 through 64 wherein the modification
comprises a chemical modification at or near the 5' end of the targeter nucleic acid. In
embodiment 66 provided herein is the composition of embodiment 65 wherein the chemical
modification comprises a chemical modification in at least one nucleotide or internucleotide
linkage within ten nucleotides of the 5' end. In embodiment 67 provided herein is the
35 composition of embodiment 65 wherein the chemical modification comprises one to five 2'-O-

methoxy-3'-phosphorothioate modifications within five nucleotides of the 5' end. In embodiment 68 provided herein is the composition of embodiment 65 wherein the chemical modification comprises two 2'-O-methoxy-3'-phosphorothioate modifications to the final two nucleotides of the 5' end. In embodiment 69 provided herein is the composition of embodiment 65 wherein the chemical modification comprises one through nine 2'-fluoro modifications to one or more nucleotides within nine nucleotides of the 5' end. In embodiment 70 provided herein is the composition of embodiment 65 wherein the chemical modification comprises five 2'-fluoro modification to the final five nucleotides of the 5' end. In embodiment 71 provided herein is the composition of any one of embodiments 65 through 70 wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide. In embodiment 72 provided herein is the composition of embodiment 1, 9, 17, 18, 27, 28, 29, 38, 39, 48, 49, 58, 59, or 65 wherein the chemical modification is 2'-O-alkyl, 2'-O-methyl, a phosphorothioate, a phosphonoacetate, a thiophosphonoacetate, a 2'-O-methyl-3'-phosphorothioate, a 2'-O-methyl-3'-phosphonoacetate, a 2'-O-methyl-3'-thiophosphonoacetate, a 2'-deoxy-3'-phosphonoacetate, a 2'-deoxy-3'-thiophosphonoacetate, or a combination thereof. In embodiment 73 provided herein is a composition comprising a Cas nuclease complexed with a gNA of any of the previous embodiments. In embodiment 74 provided herein is the composition of embodiment 73 wherein the Cas nuclease comprises a Type V Cas nuclease. In embodiment 75 provided herein is the composition of embodiment 74, wherein the Cas nuclease comprises a type V-A, type V-C, or type V-D Cas nuclease. In embodiment 76 provided herein is the composition of embodiment 75, wherein the Cas nuclease comprises a type V-A Cas nuclease. In embodiment 77 provided herein is the composition of embodiment 76 wherein the Type V-A Cas nuclease comprises a MAD, Csm1, ART, ABW, or other Cpf1 nuclease, or derivative or variant thereof. In embodiment 78 provided herein is the composition of embodiment 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 3, 4, 5, 6, 7, 8, 9, 10, or 11. In embodiment 79 provided herein is the composition of embodiment 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 1 or 2. In embodiment 80 provided herein is the composition of embodiment 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, or 984. In embodiment 81 provided herein is the composition of embodiment 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 789, 16, 29, 42, 55, 68, 81, 94, or 107. In embodiment 82 provided herein is the composition of any one of embodiments 78 through 81 wherein the nuclease sequence is at least 85% identical to the sequence. In embodiment 83 provided herein is

the composition of any one of embodiments 78 through 81 wherein the nuclease sequence is at least 90% identical to the sequence. In embodiment 84 provided herein is the composition of any one of embodiments 78 through 81 wherein the nuclease sequence is at least 95% identical to the sequence. In embodiment 85 provided herein is the composition of any one of embodiments 78 through 81 wherein the nuclease sequence is at least 97.5% identical to the sequence. In embodiment 86 provided herein is the composition of any one of embodiments 78 through 81 wherein the nuclease sequence is 100% identical to the sequence. In embodiment 87 provided herein is the composition of any one of embodiments 1 through 71 wherein the spacer sequence comprises a sequence capable of hybridizing with a target nucleotide sequence within a human ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALPNR, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWIF1, CD3D, or NLRC5 gene. In embodiment 88 provided herein is the composition of any one of embodiments 1 through 71 wherein the spacer sequence comprises a sequence capable of hybridizing with a target nucleotide sequence within a safe harbor site. In embodiment 89 provided herein is the composition of any one of embodiments 3 and 37 through 71 wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 70% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA. In embodiment 90 provided herein is the composition of any one of embodiments 3 and 37 through 71 wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 80% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA. In embodiment 91 provided herein is the composition of any one of embodiments 3 and 37 through 71 wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 90% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA. In embodiment 92 provided herein is the composition of any one of embodiments 3 and 37 through 71 wherein the dual gNA when complexed with a Cas nuclease as an RNP results in the same editing efficiency as compared to a single 5' and 3' propanediol modified gNA. In embodiment 93 provided herein is the composition of any one of embodiments 3 and 37 through 71 wherein the modified gNA when complexed with a Cas nuclease as an RNP results in higher editing efficiency as compared to a single gNA. In embodiment 94 provided herein is a eukaryotic cell comprising the gRNA of any of embodiments 1 through 71. In embodiment 95 provided herein is a eukaryotic cell comprising the nucleic acid-guided nuclease complex of any of embodiments 73 through 86. In embodiment

96 provided herein is the composition of embodiment 94 or embodiment 95 wherein the eukaryotic cell is an allogeneic cell. In embodiment 97 provided herein is the composition of any one of embodiments 94 through 96 wherein the eukaryotic cell is an immune cell. In embodiment 98 provided herein is the composition of embodiment 97 wherein the immune cell is a human immune cell. In embodiment 99 provided herein is the composition of embodiment 97 wherein the immune cell comprises a neutrophil, an eosinophil, a basophil, a mast cell, a monocyte, a macrophage, a dendritic cell, a natural killer cell, or a lymphocyte. In embodiment 100 provided herein is the composition of embodiment 97 wherein the immune cell is a T cell. In embodiment 101 provided herein is the composition of embodiment 97 wherein the immune cell is a CAR T cell. In embodiment 102 provided herein is the composition of any one of embodiments 94 through 96 wherein the eukaryotic cell is a stem cell. In embodiment 103 provided herein is the composition of embodiment 102 wherein the stem cell is a human pluripotent stem cell. In embodiment 104 provided herein is the composition of embodiment 102 wherein the stem cell is an embryonic stem cell or an induced pluripotent stem cell. In embodiment 105 provided herein is the composition of embodiment 102 wherein the stem cell is a hematopoietic stem cell. In embodiment 106 provided herein is a method of cleaving a target DNA having a target nucleotide sequence, the method comprising contacting the target DNA with the composition of any one of embodiments 73 through 86, thereby resulting in cleavage of the target DNA. In embodiment 107 provided herein is the method of embodiment 106, wherein the contacting occurs in vitro. In embodiment 108 provided herein is the method of embodiment 106, wherein the contacting occurs in a cell ex vivo. In embodiment 109 provided herein is the method of embodiment 108, wherein the target DNA is genomic DNA of the cell. In embodiment 110 provided herein is the method of embodiment 108, wherein the system is delivered into the cell as a pre-formed RNP complex. In embodiment 111 provided herein is the method of embodiment 110, wherein the pre-formed RNP complex is delivered into the cell by electroporation, lipofection, or a viral method. In embodiment 112 provided herein is the method of embodiment 111 wherein the pre-formed RNP complex is delivered into the cell by electroporation. In embodiment 113 provided herein is a method of editing the genome of a eukaryotic cell, the method comprising delivering the engineered, non-naturally occurring system of any one of embodiments 73 through 86, or components thereof and/or one or more polynucleotides coding for one or more components thereof, into the eukaryotic cell, thereby resulting in editing of the genome of the eukaryotic cell. In embodiment 114 provided herein is the method of embodiment 113, wherein the system is delivered into the cell as a pre-formed RNP complex. In embodiment 115 provided herein is the method of embodiment 113 or 114, wherein the system is delivered into the cell by electroporation, lipofection, or a viral method. In

embodiment 116 provided herein is the method of embodiment 113 or 114, wherein the system is delivered into the cell by electroporation. In embodiment 117 provided herein is the method of any one of embodiments 113 through 116, wherein the cell is an immune cell. In embodiment 118 provided herein is the method of embodiment 117, wherein the immune cell is a neutrophil, eosinophil, basophil, mast cell, monocyte, macrophage, dendritic cell, natural killer cell, or a lymphocyte. In embodiment 119 provided herein is the method of embodiment 117, wherein the immune cell is a T lymphocyte. In embodiment 120 provided herein is the method of embodiment 119, wherein an exogenous donor template is delivered in addition to the RNP wherein the donor template is used to repair the cleaved target DNA. In embodiment 121 provided herein is the method of embodiment 120 wherein the exogenous donor template is CAR cassette. In embodiment 122 provided herein is the composition of a CAR T cell resulting from the method of embodiment 121. In embodiment 123 provided herein is the method of embodiment 113 wherein the engineered, non-naturally occurring system is delivered to a plurality of eukaryotic cells, and wherein the system comprises a guide nucleic acid comprising one or modifications as described herein, wherein the editing efficiency of the genomes of the plurality of cells is increased by at least 5% compared to the editing efficiency when the same system but without the modification or modifications is used. In embodiment 124 provided herein is a pharmaceutical composition comprising the composition of any one of embodiments 1 through 93 and a pharmaceutically acceptable carrier. In embodiment 125 provided herein is a method of treating a disease or a disorder comprising administering to a subject in need thereof an effective amount of a composition of any one of embodiments 1 through 93, or an effective amount of cells modified by treatment with a composition of any one of embodiments 1 through 93. In embodiment 126 provided herein is the method of embodiment 125 comprising administering to a subject in need thereof of cells modified by treatment with a composition of any one of embodiment 1 to 93. In embodiment 127 provided herein is the method of embodiment 125 wherein the cells are cells that are removed from an individual and treated ex vivo with a composition of any one of embodiment 1 to 93. In embodiment 128 provided herein is the method of embodiment 127 wherein the subject in need of treatment and the individual whose cells are treated ex vivo are the same. In embodiment 129 provided herein is a composition comprising the composition of any one of embodiments 1 through 72 further comprising a Cas protein. In embodiment 130 provided herein is the composition of embodiment 128 wherein the Cas protein is a Cas nuclease. In embodiment 131 provided herein is the composition of embodiment 130 wherein the Cas nuclease is a Type I, II, III, IV, V, or VI Cas nuclease. In embodiment 132 provided herein is the composition of embodiment 131 wherein the Cas nuclease is a Type II Cas nuclease.

VII. Examples

Example 1

5 [0421] In this Example various combinations of modified and unmodified modulator and targeter nucleic acids were used with a Cas nuclease and the editing efficiency for each combination was evaluated.

[0422] Modifications included at least one of 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a
10 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), propanediol (Alt-R) or a combination thereof at either the 3' or 5' end. The modified/unmodified crRNA and tracrRNA (sequences shown in Table 27) were mixed in the molar ratio 1:1 (200pmol + 200pmol) at room temperature and incubated for 15min. For the formation of the RNPs 100
15 pmol unmodified/modified gRNA were mixed with 100 pmol Mad7 nuclease and transfected into primary Pan T cells using the Lonza nucleofector system. Combinations of modifications tested are shown in Figure 2.

[0423] Figure 3 shows data for editing efficiency (as measured by # of reads modified/total # of reads) in primary T cells in an exon at a gene locus. The gRNA designs are shown and
20 show the various combination of modifications tested at either the 3' and/or the 5' end of the crRNA or tracrRNA. Sequences for each tested modification are shown in Table 27. The Figure shows the editing results relative to the single gRNA design (left bar) vs. the negative control (far right bar). It is observed that the modifications made to the crRNA make a much larger difference than the tracrRNA although the combination of dual modified crRNA and tracrRNA
25 make the greatest increase in editing efficiency.

Table 27 Modified Single and Dual gRNAs

Name	Description	SEQ ID NO	Sequence ¹
sgRNA	-	1038	/AltR1/TAATTTCTACTCTTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
STAR	Modulator	1039	TAATTTCTACTC
	Targeter	1040	TTGTAGATGAGTCTCTCAGCTGGTACACG
STAR MOD1	Modulator	1041	/AltR1/TAATTTCTACTC
	Targeter	1042	TTGTAGATGAGTCTCTCAGCTGGTACACG
STAR MOD2	Modulator	1043	TAATTTCTACTC
	Targeter	1044	TTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/

STAR MOD3	Modulator	1045	/AltR1/TAATTTCTACTC/AltR2/
	Targeter	1046	TTGTAGATGAGTCTCTCAGCTGGTACACG
STAR MOD4	Modulator	1047	TAATTTCTACTC
	Targeter	1048	/AltR1/TTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
STAR MOD5	Modulator	1049	/AltR1/TAATTTCTACTC
	Targeter	1050	TTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
STAR MOD6	Modulator	1051	/AltR1/TAATTTCTACTC/AltR2/
	Targeter	1052	/AltR1/TTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/

¹/AltR1/ indicated a sequence modified with a 5' Alt-R (propanediol) modification. AltR2/ indicates a sequence modified with a 3' Alt-R (propanediol) modification.

Example 2

5 **[0424]** In this Example various combinations of modified and unmodified modulator and targeter nucleic acids were used with a Cas nuclease and the editing efficiency for each combination was evaluated for three gene targets (Gene 1, Gene 2, Gene 3).

[0425] Modifications included at least one of 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), a 2'-fluoro-ribonucleotide, propanediol (Alt-R) or a combination thereof at either the 3' or 5' end. The modified/unmodified crRNA and tracrRNA were mixed in the molar ratio 1:1 (200pmol + 200pmol) at room temperature and incubated for 15min. For the formation of the RNPs 100 pmol unmodified/modified gRNA were mixed with 100 pmol Mad7 nuclease and transfected into primary Pan T cells using the Lonza nucleofector system. Sequences for each modulator nucleic acid and targeter nucleic acid used in the study with their respective modifications are shown in **Table 28 and Table 29**.

[0426] The specific sequence for each modulator and targeter complex in **Figures 4-6** may be deduced using the naming convention (modulator description) x (targeter description) using **Table 28 and Table 29**. For example, an experimental test using complex Mod_A_7_x_Tar_1 for gene 3 would comprise a modulator of SEQ ID NO: 1031 and a targeter of SEQ ID NO: 1008. It is observed that some combinations made to the modulator-targeter complex demonstrate activity greater than the sgRNA. It is also observed that some combinations made to the modulator-targeter complex demonstrate lower but sufficient (greater than or equal to 70% activity) activity compared to the sgRNA.

[0427] **Figure 4A and B** show data for the editing efficiency (as measured by # of reads modified/ total # of reads) in primary T cells in an exon at a gene locus for gene 1 for dual

gRNAs. Each tested gNA is shown on the x-axis with the first two representing single gRNA controls. The editing efficiency for each tested dual gRNA is normalized to the activity of the single gRNA control. The dotted line represents an editing efficiency of 70% relative to the single gRNA control. The dashed line represents an editing efficiency equal to the single gRNA control. **Figure 4A** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' A nucleotide. Control gRNAs for **Figure 4A** are SEQ ID NO: 1056. **Figure 4B** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' C nucleotide. Control RNAs for **Figure 4B** are SEQ ID NO: 1055.

[0428] **Figure 5A and B** show data for the editing efficiency (as measured by # of reads modified/ total # of reads) in primary T cells in an exon at a gene locus for gene 2 for dual gRNAs. Each tested gNA is shown on the x-axis with the first two representing single gRNA controls. The editing efficiency for each tested dual gRNA is normalized to the activity of the single gRNA control. The dotted line represents an editing efficiency of 70% relative to the single gRNA control. The dashed line represents an editing efficiency equal to the single gRNA control. **Figure 5A** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' A nucleotide. Control RNAs for **Figure 5A** are SEQ ID NO: 1058. **Figure 5B** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' C nucleotide. Control RNAs for **Figure 5B** are SEQ ID NO: 1057.

[0429] **Figure 6A and B** show data for the editing efficiency (as measured by # of reads modified/ total # of reads) in primary T cells in an exon at a gene locus for gene 3 for dual gRNAs. Each tested gNA is shown on the x-axis with the first two representing single gRNA controls. The editing efficiency for each tested dual gRNA is normalized to the activity of the single gRNA control. The dotted line represents an editing efficiency of 70% relative to the single gRNA control. The dashed line represents an editing efficiency equal to the single gRNA control. **Figure 6A** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' A nucleotide. Control RNAs for **Figure 6A** are SEQ ID NO: 1054. **Figure 6B** shows normalized editing efficiencies for dual guide RNAs comprising modulator nucleic acids with a terminal 3' C nucleotide. Control RNAs for **Figure 6B** are SEQ ID NO: 1053.

Table 28 Targeter Sequences

SEQ ID NO	Description	Sequence ¹
1000	Tar_1_gene 2	TTGTAGATCACAGGAGCCGACCTGCCTAC
1001	Tar_2_gene 2	TTGTAGATCACAGGAGCCGACCTGCCTAC/AltR2/
1002	Tar_3_gene 2	TTGTAGATCACAGGAGCCGACCTGCCT*mA*mC
1003	Tar_4_gene 2	TTGTAGATCACAGGAGCCGACCTG/i2FC//i2FC//i2FT//i2FA//i2FC/
1004	Tar_1_gene 1	TTGTAGATCAAGGACTTCAGCTGGGGGAA
1005	Tar_2_gene 1	TTGTAGATCAAGGACTTCAGCTGGGGGAA/AltR2/
1006	Tar_3_gene 1	TTGTAGATCAAGGACTTCAGCTGGGGG*mA*mA
1007	Tar_4_gene 1	TTGTAGATCAAGGACTTCAGCTGG/i2FG//i2FG//i2FG//i2FA//i2FA/
1008	Tar_1_gene 3	TTGTAGATGAGTCTCTCAGCTGGTACACG
1009	Tar_2_gene 3	TTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
1010	Tar_3_gene 3	TTGTAGATGAGTCTCTCAGCTGGTACA*mC*mG
1011	Tar_4_gene 3	TTGTAGATGAGTCTCTCAGCTGGT/i2FA//i2FC//i2FA//i2FC//i2FG/

¹m indicates a 2-O-methoxy phosphorothioate nucleotide modification. * indicates a phosphorothioate internucleotide linkage modification. /i2FN/ indicates a 2'-fluoro nucleotide modification where the N may comprise any base (A, T, G, or C). /AltR2/ indicates a sequence modified with a 3' Alt-R (propanediol) modification as shown in **Figure 7B**.

5

Table 29 Modulator Sequences

SEQ ID NO	Description	Sequences ¹
1012	STAR_MOD_C_1	TAATTTCTACTC
1013	STAR_MOD_C_2	/Alt1/TAATTTCTACTC
1014	STAR_MOD_C_3	mT*A*TAATTTCTACTC
1015	STAR_MOD_C_4	mT*mA*TAATTTCTACTC
1016	STAR_MOD_C_5	AGACCTTTTTTAATTTCTACTC
1017	STAR_MOD_C_6	/Alt1/AGACCTTTTTTAATTTCTACTC
1018	STAR_MOD_C_7	A*G*A*C*C*T*T*T*T*TAATTTCTACTC
1019	STAR_MOD_C_8	mA*mG*ACCTTTTTTAATTTCTACTC
1020	STAR_MOD_C_9	mA*G*ACCTTTTTTAATTTCTACTC
1021	STAR_MOD_C_10	GTCAAAAGACCTTTTTTAATTTCTACTC
1022	STAR_MOD_C_11	/Alt1/GTCAAAAGACCTTTTTTAATTTCTACTC
1023	STAR_MOD_C_12	mG*mT*CAAAAGACCTTTTTTAATTTCTACTC
1024	STAR_MOD_C_13	mG*T*CAAAAGACCTTTTTTAATTTCTACTC
1025	STAR_MOD_A_1	TAATTTCTACTA

SEQ ID NO	Description	Sequences ¹
1026	STAR_MOD_A_2	/Alt1/TAATTTCTACTA
1027	STAR_MOD_A_3	mT*A*TAATTTCTACTA
1028	STAR_MOD_A_4	mT*mA*TAATTTCTACTA
1029	STAR_MOD_A_5	AGACCTTTTAAATTTCTACTA
1030	STAR_MOD_A_6	/Alt1/AGACCTTTTAAATTTCTACTA
1031	STAR_MOD_A_7	A*G*A*C*C*T*T*T*T*TAATTTCTACTA
1032	STAR_MOD_A_8	mA*mG*ACCTTTTAAATTTCTACTA
1033	STAR_MOD_A_9	mA*G*ACCTTTTAAATTTCTACTA
1034	STAR_MOD_A_10	GTCAAAAGACCTTTTAAATTTCTACTA
1035	STAR_MOD_A_11	/Alt1/GTCAAAAGACCTTTTAAATTTCTACTA
1036	STAR_MOD_A_12	mG*mT*CAAAAGACCTTTTAAATTTCTACTA
1037	STAR_MOD_A_13	mG*T*CAAAAGACCTTTTAAATTTCTACTA

¹m indicates a 2-O-methoxy phosphorothioate nucleotide modification. * indicates a phosphorothioate internucleotide linkage modification. /Alt1/ indicates a sequence modified a 5' Alt-R (propanediol) modification as shown in **Figure 7A**.

5

Table 30 Control Sequences

SEQ ID NO	Description	Sequences ¹
1053	gGENE3_plate_1 and gGENE3_plate_2	/AltR1/TAATTTCTACTCTTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
1054	gGENE3_plate_3 and gGENE3_plate_4	/AltR1/TAATTTCTACTATTGTAGATGAGTCTCTCAGCTGGTACACG/AltR2/
1055	gGENE1_plate_2	/AltR1/TAATTTCTACTCTTGTAGATCAAGGACTTCAGCTGGGGAA/AltR2/
1056	gGENE1_plate_3 and gGENE1_plate_4	/AltR1/TAATTTCTACTATTGTAGATCAAGGACTTCAGCTGGGGAA/AltR2/
1057	gGENE2_plate_1 and gGENE2_plate_2	/AltR1/TAATTTCTACTCTTGTAGATCACAGGAGCCGACCTGCCTAC/AltR2/
1058	gGENE2_plate_3 and gGENE2_plate_4	/AltR1/TAATTTCTACTATTGTAGATCACAGGAGCCGACCTGCCTAC/AltR2/

Example 3

[0430] In this example, various combinations of modified and unmodified modulator and targeter nucleic acids targeting the B2M gene were used with a Cas nuclease. For each combination, the editing efficiency and surface expression of HLA-1 was evaluated.

5 **[0431]** Modifications included at least one of 2'-O-methyl (M), a phosphorothioate (S), a phosphonoacetate (P), a thiophosphonoacetate (SP), a 2'-O-methyl-3'-phosphorothioate (MS), a 2'-O-methyl-3'-phosphonoacetate (MP), a 2'-O-methyl-3'-thiophosphonoacetate (MSP), a 2'-deoxy-3'-phosphonoacetate (DP), a 2'-deoxy-3'-thiophosphonoacetate (DSP), a 2'-fluoro-ribonucleotide, propanediol (Alt-R) or a combination thereof at either the 3' or 5' end. The modified/unmodified crRNA and tracrRNA were mixed in the molar ratio 1:1 (200pmol + 200pmol) at room temperature and incubated for 15min. For the formation of the RNPs 100 pmol unmodified/modified gRNA were mixed with 100 pmol Mad7 nuclease and transfected into primary Pan T cells using the Lonza nucleofector system. Sequences for each modulator nucleic acid and targeter nucleic acid used in the study with their respective modifications are shown in **Table 31** and **Table 29**.

[0432] The specific sequence for each modulator and targeter complex in **Figures 7-8** may be deduced using the naming convention (modulator description) (targeter description) using **Table 31** and **Table 29**. For example, an experimental test using complex Mod C2 Tar 2 would comprise a modulator comprising SEQ ID NO: 1013 and a targeter comprising SEQ ID NO: 999. It is observed that some combinations made to the modulator-targeter complex demonstrate activity greater than the sgRNA. It is observed that some combinations made to the modulator-targeter complex demonstrate activity similar to the sgRNA. It is also observed that some combinations made to the modulator-targeter complex demonstrate lower but sufficient activity compared to the sgRNA.

25 **[0433]** **Figure 7** show data for the editing efficiency at the B2M gene (as measured by % INDELs formed) in primary T cells after treatment with various combinations of modified and unmodified modulator and targeter nucleic acids complexed with MAD7. INDEL formation was measured by amplicon sequencing of the gene post treatment. Each tested gNA is shown on the x-axis with the first two representing a no gRNA and single gRNA controls, respectively. INDEL formation as a percentage of total reads is shown on the y-axis.

[0434] **Figure 8** show data for the editing efficiency and resultant reduction in HLA-1 surface expression in primary T cells as measured by flow cytometry (proportion of cells lacking HLA-1 expression in the population) after treatment with various combinations of modified and unmodified modulator and targeter nucleic acids complexed with MAD7. Each tested gNA is

shown on the x-axis with the first two representing a no gRNA (cell only control) and a single gRNA control, respectively.

Table 31 B2M Targeter Sequences

SEQ ID NO	Description	Sequence
998	Tar_1_B2M	TTGTAGATAGTGGGGGTGAATTCAGTGTA
999	Tar_2_B2M	TTGTAGATAGTGGGGGTGAATTCAGTGTA/AltR2/

5

[0435] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

10

CLAIMS

WHAT IS CLAIMED IS:

1. A composition comprising a synthetic guide nucleic acid (gNA) comprising
 - (i) a targeter nucleic acid comprising:
 - (a) a spacer sequence configured to hybridize with a target nucleotide sequence, and
 - (b) a targeter stem sequence; and
 - (ii) a modulator nucleic acid comprising
 - (a) a modulator stem sequence complementary to the targeter stem sequence, and
 - (b) a 5' sequence;wherein either the targeter nucleic acid or the modulator nucleic acid, or both, comprise one or more modified nucleotides at or near its 3' end, if present, at or near its 5' end, if present, or both.
2. The composition of claim 1 wherein the synthetic gNA is a single polynucleotide.
3. The composition of claim 1 wherein the targeter nucleic acid and the modulator nucleic acid are separate nucleic acids.
4. The composition of any one of claims 1 through 3 wherein some or all of the gNA is RNA, *e.g.*, a gRNA.
5. The composition of claim 4 wherein at least 50% of the nucleic acid is RNA.
6. The composition of claim 4 wherein at least 70% of the nucleic acid is RNA.
7. The composition of claim 4 wherein at least 90% of the nucleic acid is RNA.
8. The composition of claim 4 wherein at least 100% of the nucleic acid is RNA.
9. The composition of any one of claims 1 through 8 wherein the modified nucleotide comprises a chemical modification.

10. The composition of any one of claims 1 through 9 wherein the stem sequences are between one and ten nucleotides in length.
11. The composition of any one of claims 1 through 9 wherein the stem sequences are between
5 two and nice nucleotides in length.
12. The composition of any one of claims 1 through 9 wherein the stem sequences are between four and six nucleotides in length.
- 10 13. The composition of any one of claims 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 80% sequence complementarity.
14. The composition of any one of claims 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 90% sequence complementarity.
- 15 15. The composition of any one of claims 1 through 12 wherein the stem sequences of the modulator and targeter nucleic acids share at least 95% sequence complementarity.
16. The composition of any one of claims 1 through 12 wherein the stem sequences of the
20 modulator and targeter nucleic acids share 100% sequence complementarity.
17. The composition of claim 2 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 5' end of the gNA.
- 25 18. The composition of claim 17 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 5' end.
19. The composition of claim 17 wherein the chemical modification comprises one to 17
30 phosphorothioate modifications within 17 internucleotide linkages of the 5' end.
20. The composition of claim 19 wherein the chemical modification comprises one to ten phopshorothioate modifications within ten internucleotide linkages of the 5' end.

21. The composition of claim 19 wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 5' end.
22. The composition of claim 17 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 5' end.
23. The composition of any one of claims 17 through 21 wherein the chemical modification comprises two 2'-O-methoxy modifications to the final two nucleotides of the 5' end.
24. The composition of any one of claims 17 through 21 wherein the chemical modification comprises one 2'-O-methoxy modification to the 5' terminal nucleotide.
25. The composition of any one of claims 17 through 24 wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide.
26. The composition of claim 17 wherein the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037.
27. The composition of claim 2 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 3' end of the gNA.
28. The composition of any one of claims 17 through 25 wherein the chemical modification further comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 3' end of the gNA.
29. The composition of claim 27 or claim 28 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3' end.
30. The composition of claim 27 or claim 28 wherein the chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 3' end.
31. The composition of claim 30 wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 3' end.

32. The composition of claim 30 wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 3' end.
- 5 33. The composition of any one of claims 27 through 32 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 3' end.
34. The composition of any one of claims 27 through 32 wherein the chemical modification
10 comprises two 2'-O-methoxy modifications to the final two nucleotides of the 3' end.
35. The composition of any one of claims 27 through 32 wherein the chemical modification comprises one 2'-O-methoxy modification to the last nucleotide of the 3' end.
- 15 36. The composition of any one of claims 27 through 35 wherein the chemical modification comprises a propanediol modification to the 3' terminal nucleotide.
37. The composition of claim 3 wherein the modulator nucleic acid is unmodified.
- 20 38. The composition of claim 3 wherein the modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or near the 5' end of the modulator nucleic acid.
39. The composition of claim 38 wherein the chemical modification comprises a chemical
25 modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 5' end.
40. The composition of claim 38 wherein the chemical modification comprises one to 17
phosphorothioate modifications within 17 internucleotide linkages of the 5' end.
30
41. The composition of claim 40 wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 5' end.
42. The composition of claim 40 wherein the chemical modification comprises nine
35 phosphorothioate modifications to the final nine internucleotide linkages of the 5' end.

- 5 **43.** The composition of any one of claims **38** through **42** wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 5' end.
- 44.** The composition of any one of claims **38** through **42** wherein the chemical modification comprises two 2'-O-methoxy modifications to the final two nucleotides of the 5' end.
- 10 **45.** The composition of any one of claims **38** through **42** wherein the chemical modification comprises one 2'-O-methoxy modification to the 5' end nucleotide.
- 46.** The composition of any one of claims **38** through **45** wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide.
- 15 **47.** The composition of claim **38** wherein the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037.
- 20 **48.** The composition of any one of claims **3** and claims **38** through **46** wherein the modification comprises a chemical modification to one or more nucleotides or internucleotide linkages at or near the 3' end of the modulator nucleic acid.
- 49.** The composition of claim **48** wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3' end.
- 25 **50.** The composition of claim **48** wherein the chemical modification comprises one to 17 phosphorothioate modifications within 17 internucleotide linkages of the 3' end.
- 30 **51.** The composition of claim **50** wherein the chemical modification comprises one to ten phosphorothioate modifications within ten internucleotide linkages of the 3' end.
- 52.** The composition of claim **50** wherein the chemical modification comprises nine phosphorothioate modifications to the final nine internucleotide linkages of the 3' end.

35

53. The composition of any one of claims 3 and claims 38 through 52 wherein the chemical modification comprises one to five 2'-O-methoxy modifications to one or more nucleotides within five nucleotides of the 3' end.
- 5 54. The composition of any one of claims 3 and claims 38 through 52 wherein the chemical modification comprises two 2'-O-methoxy modifications to the final two nucleotides of the 3' end.
55. The composition of any one of claims 3 and claims 38 through 52 wherein the chemical
10 modification comprises one 2'-O-methoxy modification to the last nucleotide of the 3' end.
56. The composition of any one of claims 48 through 55 wherein the chemical modification comprises a propanediol modification to the 3' terminal nucleotide.
- 15 57. The composition of any one of claims 3 and 38 through 56 wherein the targeter nucleic acid is unmodified.
58. The composition of any one of claims 37 through 56 wherein the chemical modification comprises a chemical modification in one or more nucleotides or internucleotide linkages at or
20 near the 3' end of the targeter nucleic acid.
59. The composition of claim 58 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 3' end.
- 25 60. The composition of claim 58 wherein the chemical modification comprises one to five 2'-O-methoxy-3'-phosphorothioate modifications within five nucleotides of the 3' end.
61. The composition of claim 58 wherein the chemical modification comprises two 2'-O-methoxy-3'-phosphorothioate modifications to the final two nucleotides of the 3' end.
- 30 62. The composition of claim 58 wherein the chemical modification comprises one through nine 2'-fluoro modifications to the final nine nucleotides of the 3' end.

63. The composition of claim 58 wherein the chemical modification comprises five 2'-fluoro modifications to the final five nucleotides of the 3' end.

64. The composition of any one of claims 58 through 63 wherein the chemical modification
5 comprises a propanediol modification to the 3' terminal nucleotide.

65. The composition of any one of claims 3, claims 37 through 56, and claims 58 through 64 wherein the modification comprises a chemical modification at or near the 5' end of the targeter nucleic acid.

10

66. The composition of claim 65 wherein the chemical modification comprises a chemical modification in at least one nucleotide or internucleotide linkage within ten nucleotides of the 5' end.

15 67. The composition of claim 65 wherein the chemical modification comprises one to five 2'-O-methoxy-3'-phosphorothioate modifications within five nucleotides of the 5' end.

68. The composition of claim 65 wherein the chemical modification comprises two 2'-O-methoxy-3'-phosphorothioate modifications to the final two nucleotides of the 5' end.

20

69. The composition of claim 65 wherein the chemical modification comprises one through nine 2'-fluoro modifications to one or more nucleotides within nine nucleotides of the 5' end.

70. The composition of claim 65 wherein the chemical modification comprises five 2'-fluoro
25 modification to the final five nucleotides of the 5' end.

71. The composition of any one of claims 65 through 70 wherein the chemical modification comprises a propanediol modification to the 5' terminal nucleotide.

30 72. The composition of claim 1, 9, 17, 18, 27, 28, 29, 38, 39, 48, 49, 58, 59, or 65 wherein the chemical modification is 2'-O-alkyl, 2'-O-methyl, a phosphorothioate, a phosphonoacetate, a thiophosphonoacetate, a 2'-O-methyl-3'-phosphorothioate, a 2'-O-methyl-3'-phosphonoacetate, a 2'-O-methyl-3'-thiophosphonoacetate, a 2'-deoxy-3'-phosphonoacetate, a 2'-deoxy-3'-thiophosphonoacetate, or a combination thereof.

35

73. A composition comprising a Cas nuclease complexed with a gNA of any of the previous claims.

74. The composition of claim 73 wherein the Cas nuclease comprises a Type V Cas nuclease.

75. The composition of claim 74, wherein the Cas nuclease comprises a type V-A, type V-C, or type V-D Cas nuclease.

76. The composition of claim 75, wherein the Cas nuclease comprises a type V-A Cas nuclease.

77. The composition of claim 76 wherein the Type V-A Cas nuclease comprises a MAD, Csm1, ART, ABW, or other Cpf1 nuclease, or derivative or variant thereof.

78. The composition of claim 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 3, 4, 5, 6, 7, 8, 9, 10, or 11.

79. The composition of claim 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 1 or 2.

80. The composition of claim 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, or 984.

81. The composition of claim 77 wherein the nuclease comprises a sequence with at least 80% identity to any one of SEQ ID NOs: 789, 16, 29, 42, 55, 68, 81, 94, or 107.

82. The composition of any one of claims 78 through 81 wherein the nuclease sequence is at least 85% identical to the sequence.

83. The composition of any one of claims 78 through 81 wherein the nuclease sequence is at least 90% identical to the sequence.

84. The composition of any one of claims 78 through 81 wherein the nuclease sequence is at least 95% identical to the sequence.

85. The composition of any one of claims **78** through **81** wherein the nuclease sequence is at least 97.5% identical to the sequence.

5 **86.** The composition of any one of claims **78** through **81** wherein the nuclease sequence is 100% identical to the sequence.

87. The composition of any one of claims **1** through **71** wherein the spacer sequence comprises a sequence capable of hybridizing with a target nucleotide sequence within a human ADORA2A, B2M, CD3E, CD38, CD40LG, CD52, CIITA, CSF2, CTLA4, DCK, FAS, HAVCR2 (also called TIM3), LAG3, PDCD1 (also called PD-1), PTPN6, TIGIT, TRAC, TRBC1, TRBC2, TRBC1_2 (or TRBC1+2), CARD11, CD247, IL7R, LCK, PLCG1, ALPNI, BBS1, CALR, CD3G, CD58, COL17A1, DEFB134, ERAP1, ERAP2, IFNGR1, IFNGR2, JAK1, JAK2, mir-101-2, MLANA, PSMB5, PSMB8, PSMB9, PTC2, RFX5, RFXANK, RFXAP, RPL23, SOX10, SRP54, STAT1, Tap1, TAP2, TAPBP, TWF1, CD3D, or NLRC5 gene.

88. The composition of any one of claims **1** through **71** wherein the spacer sequence comprises a sequence capable of hybridizing with a target nucleotide sequence within a safe harbor site.

20 **89.** The composition of any one of claims **3** and **37** through **71** wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 70% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA.

90. The composition of any one of claims **3** and **37** through **71** wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 80% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA.

91. The composition of any one of claims **3** and **37** through **71** wherein the dual gNA when complexed with a Cas nuclease as an RNP results in at least 90% of the editing efficiency as compared to a 5' and 3' propanediol modified single gNA.

92. The composition of any one of claims **3** and **37** through **71** wherein the dual gNA when complexed with a Cas nuclease as an RNP results in the same editing efficiency as compared to a single 5' and 3' propanediol modified gNA.

35

93. The composition of any one of claims 3 and 37 through 71 wherein the modified gNA when complexed with a Cas nuclease as an RNP results in higher editing efficiency as compared to a single gNA.
- 5 94. A eukaryotic cell comprising the gRNA of any of claims 1 through 71.
95. A eukaryotic cell comprising the nucleic acid-guided nuclease complex of any of claims 73 through 86.
- 10 96. The composition of claim 94 or claim 95 wherein the eukaryotic cell is an allogeneic cell.
97. The composition of any one of claims 94 through 96 wherein the eukaryotic cell is an immune cell.
- 15 98. The composition of claim 97 wherein the immune cell is a human immune cell.
99. The composition of claim 97 wherein the immune cell comprises a neutrophil, an eosinophil, a basophil, a mast cell, a monocyte, a macrophage, a dendritic cell, a natural killer cell, or a lymphocyte.
- 20 100. The composition of claim 97 wherein the immune cell is a T cell.
101. The composition of claim 97 wherein the immune cell is a CAR T cell.
- 25 102. The composition of any one of claims 94 through 96 wherein the eukaryotic cell is a stem cell.
103. The composition of claim 102 wherein the stem cell is a human pluripotent stem cell.
- 30 104. The composition of claim 102 wherein the stem cell is an embryonic stem cell or an induced pluripotent stem cell.
105. The composition of claim 102 wherein the stem cell is a hematopoietic stem cell.

106. A method of cleaving a target DNA having a target nucleotide sequence, the method comprising contacting the target DNA with the composition of any one of claims **73** through **86**, thereby resulting in cleavage of the target DNA.

5 **107.** The method of claim **106**, wherein the contacting occurs *in vitro*.

108. The method of claim **106**, wherein the contacting occurs in a cell *ex vivo*.

109. The method of claim **108**, wherein the target DNA is genomic DNA of the cell.

10

110. The method of claim **108**, wherein the system is delivered into the cell as a pre-formed RNP complex.

15 **111.** The method of claim **110**, wherein the pre-formed RNP complex is delivered into the cell by electroporation, lipofection, or a viral method.

112. The method of claim **111** wherein the pre-formed RNP complex is delivered into the cell by electroporation.

20 **113.** A method of editing the genome of a eukaryotic cell, the method comprising delivering the engineered, non-naturally occurring system of any one of claims **73** through **86**, or components thereof and/or one or more polynucleotides coding for one or more components thereof, into the eukaryotic cell, thereby resulting in editing of the genome of the eukaryotic cell.

25 **114.** The method of claim **113**, wherein the system is delivered into the cell as a pre-formed RNP complex.

115. The method of claim **113** or **114**, wherein the system is delivered into the cell by electroporation, lipofection, or a viral method.

30

116. The method of claim **113** or **114**, wherein the system is delivered into the cell by electroporation.

117. The method of any one of claims **113** through **116**, wherein the cell is an immune cell.

35

118. The method of claim 117, wherein the immune cell is a neutrophil, eosinophil, basophil, mast cell, monocyte, macrophage, dendritic cell, natural killer cell, or a lymphocyte.

119. The method of claim 117, wherein the immune cell is a T lymphocyte.

5

120. The method of claim 119, wherein an exogenous donor template is delivered in addition to the RNP wherein the donor template is used to repair the cleaved target DNA.

121. The method of claim 120 wherein the exogenous donor template is CAR cassette.

10

122. The composition of a CAR T cell resulting from the method of claim 121.

15

123. The method of claim 113 wherein the engineered, non-naturally occurring system is delivered to a plurality of eukaryotic cells, and wherein the system comprises a guide nucleic acid comprising one or modifications as described herein, wherein the editing efficiency of the genomes of the plurality of cells is increased by at least 5% compared to the editing efficiency when the same system but without the modification or modifications is used.

20

124. A pharmaceutical composition comprising the composition of any one of claims 1 through 93 and a pharmaceutically acceptable carrier.

25

125. A method of treating a disease or a disorder comprising administering to a subject in need thereof an effective amount of a composition of any one of embodiments 1 through 93, or an effective amount of cells modified by treatment with a composition of any one of embodiments 1 through 93

30

126. The method of claim 125 comprising administering to a subject in need thereof of cells modified by treatment with a composition of any one of claim 1 to 93.

127. The method of claim 125 wherein the cells are cells that are removed from an individual and treated *ex vivo* with a composition of any one of claim 1 to 93.

35

128. The method of claim 127 wherein the subject in need of treatment and the individual whose cells are treated *ex vivo* are the same.

129. A composition comprising the composition of any one of claims **1** through **72** further comprising a Cas protein.

130. The composition of claim **128** wherein the Cas protein is a Cas nuclease.

5

131. The composition of claim **130** wherein the Cas nuclease is a Type I, II, III, IV, V, or VI Cas nuclease.

132. The composition of claim **131** wherein the Cas nuclease is a Type II Cas nuclease.

10




Figure 1A




Figure 1B




Figure 2A




Figure 2B




Figure 2C




Figure 3






Figure 4B



Figure 5A




Figure 5B




Figure 6A




Figure 6B




Figure 7




Figure 8




Figure 9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2022/017582

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12N 15/09; C12N 15/10; C12N 15/11; C12N 15/113; C12N 15/115 (2022.01)

CPC - C12N 15/09; C12N 15/10; C12N 15/102; C12N 15/11; C12N 15/113 (2022.05)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

see Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

see Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

see Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2019/0017050 A1 (ACTYM THERAPEUTICS, INC.) 17 January 2019 (17.01.2019) entire document	1-4, 8, 37
A	WO 2019/126037 A1 (CITY OF HOPE) 27 June 2019 (27.06.2019) entire document	1-4, 8, 37
A	US 2018/0371497 A1 (INSCRIPTA INC.) 27 December 2018 (27.12.2018) entire document	1-4, 8, 37
A	US 2018/0028613 A1 (UNIVERSITY OF LEICESTER et al) 01 February 2018 (01.02.2018) entire document	1-4, 8, 37

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 June 2022

Date of mailing of the international search report

JUL 13 2022

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Taina Matos

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/017582

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

a. ☒ forming part of the international application as filed:

☒ in the form of an Annex C/ST.25 text file.

☐ on paper or in the form of an image file.

b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.

c. ☐ furnished subsequent to the international filing date for the purposes of international search only:

☐ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).

☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

SEQ ID NOs: 51, 1000, and 1012 were searched.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/017582

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 9-16, 25, 28-36, 46, 48-132
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See extra sheet(s).

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-4, 8, 37

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2022/017582

Continued from Box No. III Observations where unity of invention is lacking

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees need to be paid.

Group I+: claims 1-8, 17-24, 26, 27, 37-45, and 47 are drawn to synthetic guide nucleic acids.

The first invention of Group I+ is restricted to a composition comprising a synthetic guide nucleic acid (gNA) comprising (i) a targeter nucleic acid comprising: (a) a spacer sequence configured to hybridize with a target nucleotide sequence, and (b) a targeter stem sequence; and (ii) a modulator nucleic acid comprising (a) a modulator stem sequence complementary to the targeter stem sequence, where the spacer sequence is selected to be SEQ ID NO: 51, the targeter stem sequence is selected to be SEQ ID NO: 1000, and the modulator stem sequence is selected to be SEQ ID NO: 1012. It is believed that claims 1-4, 8, and 37 read on this first named invention and thus these claims will be searched without fee to the extent that they read on SEQ ID NOs: 51, 1000, and 1012.

Applicant is invited to elect additional synthetic guide nucleic acids (gNAs), with their respective, corresponding, SEQ ID NOs to be searched in a specific combination by paying additional fee for each set of election. An exemplary election would be a composition comprising a synthetic guide nucleic acid (gNA) comprising (i) a targeter nucleic acid comprising: (a) a spacer sequence configured to hybridize with a target nucleotide sequence, and (b) a targeter stem sequence; and (ii) a modulator nucleic acid comprising (a) a modulator stem sequence complementary to the targeter stem sequence, where the spacer sequence is selected to be SEQ ID NO: 52, the targeter stem sequence is selected to be SEQ ID NO: 1001, and the modulator stem sequence is selected to be SEQ ID NO: 1013. Additional synthetic guide nucleic acids (gNAs), with their respective, corresponding, SEQ ID NOs will be searched upon the payment of additional fees. Applicants must specify the claims that read on any additional elected inventions. Applicants must further indicate, if applicable, the claims which read on the first named invention if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched/examined.

The inventions listed in Groups I+ do not relate to a single general inventive concept under PCT Rule 13.1, because under PCT Rule 13.2 they lack the same or corresponding special technical features for the following reasons:

The Groups I+ formulas do not share a significant structural element responsible for genomic DNA editing, requiring the selection of alternative synthetic guide nucleic acids where "the modulator nucleic acid comprises any one of SEQ ID NOs: 1012, 1013, 1014, 1015, 1016, 1017, 1018, 1019, 1020, 1021, 1022, 1023, 1024, 1025, 1026, 1027, 1028, 1029, 1030, 1031, 1032, 1033, 1034, 1035, 1036, or 1037."

Additionally, even if Groups I+ were considered to share the technical features of a composition comprising a synthetic guide nucleic acid (gNA) comprising (i) a targeter nucleic acid comprising: (a) a spacer sequence configured to hybridize with a target nucleotide sequence, and (b) a targeter stem sequence; and (ii) a modulator nucleic acid comprising (a) a modulator stem sequence complementary to the targeter stem sequence, and (b) a 5' sequence; wherein either the targeter nucleic acid or the modulator nucleic acid, or both, comprise one or more modified nucleotides at or near its 3' end, if present, at or near its 5' end, if present, or both. However, these shared technical features do not represent a contribution over the prior art.

Specifically, WO 2019/126037 A1 to City of Hope discloses a composition (provided herein is a composition, Para. [0007]) comprising a synthetic guide nucleic acid (gNA) (the nucleic acid further includes a guide RNA (gRNA) sequence, wherein one or more cytosine nucleotides and/or one or more uracil nucleotides of said gRNA sequence are modified nucleotides, Para. [0004]; engineered through one or more rounds of in vitro selection techniques, Para. [0171]) comprising (i) a targeter nucleic acid (gRNAs that target the TAR element, Para. [0038]) comprising: (a) a spacer sequence configured to hybridize with a target nucleotide sequence (containing individual spacer sequences, Para. [0104]), And (b) a targeter stem sequence (the gRNA sequence or tracrRNA sequence of the disclosure has one or more loops in the sequence. The loop (or loop region) used herein is according to its plain and ordinary meaning in the art and refers to a stem-loop, Para. [0165]); and (ii) a modulator nucleic acid comprising (a) a modulator stem sequence complementary to the targeter stem sequence (gDNA includes a nucleotide sequence complementary to a target site (e.g., a modulator binding sequence), Para. [0099]), and (b) a 5' sequence (a requisite 5'-NNGRRT protospacer adjacent motif (PAM), Para. [0101]); wherein either the targeter nucleic acid or the modulator nucleic acid, or both, comprise one or more modified nucleotides at or near its 3' end, if present, at or near its 5' end, if present, or both (gDNA includes a nucleotide sequence complementary to a target site (e.g., a modulator binding sequence), Para. [0099]), and (b) a 5' sequence (a requisite 5'-NNGRRT protospacer adjacent motif (PAM), Para. [0101]).

The inventions listed in Groups I+ therefore lack unity under Rule 13 because they do not share a same or corresponding special technical features.