

**(12) STANDARD PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. **AU 2014350197 B2**

(54) Title  
**Selective PYY compounds and uses thereof**

(51) International Patent Classification(s)  
**C07K 14/575** (2006.01) **A61K 38/22** (2006.01)

(21) Application No: **2014350197** (22) Date of Filing: **2014.11.13**

(87) WIPO No: **WO15/071355**

(30) Priority Data

(31) Number	(32) Date	(33) Country
<b>13193082.8</b>	<b>2013.11.15</b>	<b>EP</b>
<b>14176725.1</b>	<b>2014.07.11</b>	<b>EP</b>
<b>61/906,020</b>	<b>2013.11.19</b>	<b>US</b>

(43) Publication Date: **2015.05.21**

(44) Accepted Journal Date: **2018.10.04**

(71) Applicant(s)  
**Novo Nordisk A/S**

(72) Inventor(s)  
**Ostergaard, Soren;Frieboes, Kilian Waldemar Conde;Wieczorek, Birgit;Thomsen, Jens  
Kaalby;Wulff, Birgitte Schjellerup;Jessen, Carsten**

(74) Agent / Attorney  
**Shelston IP Pty Ltd., Level 21, 60 Margaret Street, Sydney, NSW, 2000, AU**

(56) Related Art  
**WO 2011033068 A1**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 May 2015 (21.05.2015)

(10) International Publication Number  
**WO 2015/071355 A1**

- (51) **International Patent Classification:**  
*A61K 38/22* (2006.01) *C07K 14/575* (2006.01)
- (21) **International Application Number:**  
PCT/EP2014/074476
- (22) **International Filing Date:**  
13 November 2014 (13.11.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
13193082.8 15 November 2013 (15.11.2013) EP  
61/906,020 19 November 2013 (19.11.2013) US  
14176725.1 11 July 2014 (11.07.2014) EP
- (71) **Applicant:** NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).
- (72) **Inventors:** ØSTERGAARD, Søren; Novo Allé, DK-2880 Bagsværd (DK). FRIEBOES, Kilian Waldemar Conde; Novo Allé, DK-2880 Bagsværd (DK). WIECZOREK, Birgit; Novo Allé, DK-2880 Bagsværd (DK). THOMSEN, Jens Kaalby; Novo Allé, DK-2880 Bagsværd (DK). WULFF, Birgitte Schjellerup; Novo Allé, DK-2880 Bagsværd (DK). JESSEN, Carsten; Novo Allé, DK-2880 Bagsværd (DK).

- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** SELECTIVE PYY COMPOUNDS AND USES THEREOF

(57) **Abstract:** The invention relates to PYY compounds having the amino acid in the position corresponding to position 30 of hPYY(1-36) substituted with tryptophan and derivatives thereof with a modifying group attached to the position corresponding to position 7 of hPYY(1-36). The compounds of the invention are selective Y2 receptor agonists. The invention also relates to pharmaceutical compositions comprising such PYY compounds and pharmaceutically acceptable excipients, as well as the medical use of the PYY compounds.



WO 2015/071355 A1

**DESCRIPTION**

**TITLE:** Selective PYY compounds and uses thereof

**TECHNICAL FIELD**

The present invention relates to analogues and/or derivatives of Peptide YY (PYY), and their pharmaceutical use.

**INCORPORATION-BY-REFERENCE OF THE SEQUENCE LISTING**

The Sequence Listing, entitled "SEQUENCE LISTING", is 35329 bytes, was created on 22 October 2014 and is incorporated herein by reference.

**BACKGROUND OF THE INVENTION**

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

PYY is released during a meal from L-cells in the distal small intestine and the colon. PYY is known to have peripheral effects in the gastrointestinal (GI) tract and also act centrally as a satiety signal. PYY is naturally secreted as a 36 amino acid peptide (PYY(1-36)) with a C-terminal amide but is cleaved to PYY(3-36) which constitutes approximately 50% of the circulating PYY. The enzyme responsible for the degradation is dipeptidyl peptidase IV (DPP IV). PYY(3-36) is rapidly eliminated by proteases and other clearance mechanisms. The half-life of PYY(3-36) has been reported to be <30 minutes in pigs. Thus, PYY displays suboptimal pharmacokinetic properties, meaning that the peptide has to be administered at least twice daily.

Whereas PYY(1-36) activates Y1, Y2 and Y5 receptors with very little selectivity and the Y4 receptor slightly less, the DPP IV processed PYY(3-36) displays increased selectivity for the Y2 receptor over Y1, Y4 and Y5 receptors, albeit some Y1 and Y5 affinity is retained. Y2 receptor activation is known to decrease appetite and food intake whereas Y1 and Y5 receptor activation leads to an increase in appetite and food intake.

Furthermore, Y1 and Y5 receptor activation may lead to an increase in blood pressure.

PYY(3-36) has been suggested for use in the treatment of obesity and associated diseases based on the demonstrated effects of certain of these peptides in animal models and in man, and on the fact that obese people have low basal levels of PYY as well as lower meal responses of this peptide. Furthermore, Y2 agonists have been demonstrated to have anti-secretory and pro-absorptive effects in the gastro-intestinal (GI) tract. The potential use of Y2 agonists in the treatment of a number of gastro-intestinal disorders has been suggested.

Based on demonstrated effects in e.g. Zucker rats and Diet-Induced Obese (DIO) mice Y2 selective PYY(3-36) analogues have a positive effect on glucose metabolism and are thus suggested to be used for the treatment of diabetes.

WO 2009/138511 A1 relates to long-acting Y2 and/or Y4 receptor agonists. WO 2011/033068 A1 relates to PYY analogues stabilised against C-terminal proteolytic breakdown. WO 2011/058165 A1 relates to Y2 receptor agonists with protracted pharmacokinetic properties.

For the treatment of conditions responsive to Y receptor modulation such as obesity and diabetes it would be attractive to use PYY analogues which are specific for the Y receptor subtype Y2 and importantly also display protracted pharmacokinetic properties and as such can be used in a dosing regimen with lower frequency of administration than PYY or PYY(3-36).

## SUMMARY

The invention relates to PYY compounds. The PYY compounds of the present invention may comprise up to 10 amino acid modifications as compared to human PYY(3-36) (hPYY(3-36), SEQ ID NO:2), and have the amino acid in the position corresponding to position 30 of human PYY(1-36) (hPYY(1-36), SEQ ID NO:1) substituted with tryptophan.

In one aspect, the PYY compounds further comprise N(alpha)-methylarginine in the position corresponding to position 35 of hPYY(1-36).

Also or alternatively, in one aspect, the PYY compounds further comprise a lysine in the position corresponding to position 7 of hPYY(1-36), and a modifying group attached to the epsilon amino group of this lysine.

In one aspect, the invention also relates to pharmaceutical compositions comprising such PYY compounds and pharmaceutically acceptable excipients, as well as the medical use of the PYY compounds.

Also or alternatively, in one aspect, the invention relates to PYY compounds being Y2 receptor agonists.

Also or alternatively, in one aspect, the invention relates to PYY compounds displaying selectivity towards the Y receptor subtype Y2 as compared to Y receptor subtypes Y1, Y4 and Y5.

Also or alternatively, in one aspect, the invention relates to PYY compounds with longer half-life than the half-life of hPYY(3-36). Also or alternatively, in one aspect, the invention relates to PYY compounds with longer half-life than the half-life of hPYY(1-36).

**DESCRIPTION OF THE INVENTION**

According to a first aspect, the present invention provides a PYY compound comprising:

i) tryptophan at the position corresponding to position 30 of hPYY(1-36) (SEQ ID NO:1),


ii) N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36)

(SEQ ID NO:1),

iii) a lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1) and a


modifying group attached to the epsilon amino group of said lysine, wherein said modifying group is defined by **A-B-C-**,

wherein **A-** is selected from




wherein a is an integer from 12 to 19, b is an integer from 10 to 16, and c is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-,

wherein **B-** is selected from



wherein d is 1 or 2; e is 1 or 2; and f is 2, 3 or 4; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-,

wherein **-C-** is



wherein g is an integer in the range of 1-5, h is an integer in the range of 1-5, i is an integer in the range of 2 to 6, j is an integer in the range of 1 to 6, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36), and

iv) a maximum of 8 amino acid modifications as compared to hPYY(3-36), or a pharmaceutically acceptable salt thereof.

According to a second aspect, the present invention provides a pharmaceutical composition comprising a PYY compound according to the invention, and at least one pharmaceutically acceptable excipient.

According to a third aspect, the present invention provides a PYY compound according to the invention, when used as a medicament.

According to a fourth aspect, the present invention provides a method of treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression and/or obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence which comprises administering to a subject in need thereof an effective amount of a PYY compound according to the invention.

According to a fifth aspect, the present invention provides a pharmaceutical composition according to the invention, when used as a medicament.

According to a sixth aspect, the present invention provides a method of treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression and/or eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence which comprises administering to a subject in need thereof an effective amount of a pharmaceutical composition according to the invention.

According to a seventh aspect, the present invention provides use of a PYY compound according to the invention in the manufacture of a medicament for the treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression and/or obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence.

According to an eighth aspect, the present invention provides use of a pharmaceutical composition according to the invention in the manufacture of a medicament for the treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression and/or obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence.

Unless the context clearly requires otherwise, throughout the description and the claims, the words "comprise", "comprising", and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".

The invention relates to PYY compounds. The PYY compounds of the present invention may comprise up to 10 amino acid modifications as compared to hPYY(3-36), and have the amino acid in the position that corresponds to position 30 of hPYY(1-36) substituted with tryptophan.

Also, in one aspect, the arginine residue in the position corresponding to position 35 of hPYY(1-36) has a methyl group added to the  $\alpha$ -amino group, meaning that the residue in the position corresponding to position 35 of hPYY(1-36) is thus N(alpha)-methylarginine.

Also or alternatively, in one aspect, the PYY compounds further comprise a lysine in the position corresponding to position 7 of hPYY(1-36), and a modifying group attached to the epsilon amino group of this lysine.

Also or alternatively, in one aspect, the invention relates to PYY compounds being Y receptor subtype Y2 agonists.

Also or alternatively, in one aspect, the invention relates to PYY compounds displaying selectivity towards the Y receptor subtype Y2 as compared to Y receptor subtypes Y1, Y4 and Y5.


In one aspect peptides being "selective" for specific receptors over other receptors refers to peptides that display at least 10 fold, such as at least 20 fold, at least 50 fold, or at least 100 fold higher potency for one Y receptor over other Y receptors as measured *in vitro* in an assay for receptor function, such as an Actone functional potency assay, and compared by EC50 values, or a Scintillation Proximity Assay (SPA) measuring receptor binding affinity, and compared by Ki values.

In what follows, Greek letters may be represented by their symbol or the corresponding written name, for example:  $\alpha$  = alpha;  $\beta$  = beta;  $\varepsilon$  = epsilon;  $\gamma$  = gamma;  $\omega$  = omega; etc.

### **PYY compounds**

The term "hPYY(1-36)" as used herein refers to the human Peptide YY, the sequence of which is included in the sequence listing as SEQ ID NO:1. The peptide having the sequence of SEQ ID NO:1 may also be designated native hPYY.

The term "PYY compound" as used herein refers to a peptide, or a compound, which is a variant of hPYY(1-36). The term "PYY compound" as used herein may also refer to a peptide, or a compound, which is a variant of hPYY(3-36) (SEQ ID NO:2).





The term "PYY compound" as used herein may also refer to a peptide, or a compound, which is a variant of hPYY(4-36).

The C-terminal of the PYY compounds of the present invention is an amide, as is the C-terminal of native hPYY(1-36) (SEQ ID NO:1) and hPYY(3-36) (SEQ ID NO:2),  
5 respectively.

The PYY compounds of the present invention can be PYY analogues and/or derivatives thereof.

The term "PYY analogue" is used for PYY compounds, where at least one amino acid modification in the backbone is present.

10 The term "PYY derivative" is used for PYY compounds comprising at least one non-amino acid substituent covalently attached.

A derivative of a PYY analogue is thus a PYY compound comprising at least one amino acid modification and at least one non-amino acid substituent covalently attached.

15 The PYY compounds of the present invention may comprise up to 10 amino acid modifications as compared to hPYY(3-36).

The term "amino acid modification" used throughout this application is used in the meaning of a modification to an amino acid as compared to hPYY(3-36). This modification can be the result of a deletion of an amino acid, addition of an amino acid,  
20 substitution of one amino acid with another or a substituent covalently attached to an amino acid of the peptide.

The PYY compounds of the invention comprises a tryptophan at the position corresponding to position 30 of hPYY(1-36), meaning that the PYY compounds of the invention may comprise up to 9 amino acid modifications as compared to hPYY(3-36) in  
25 addition to this modification in the position corresponding to position 30 of hPYY(1-36).

As an example, [Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) comprises 4 amino acid substitutions as compared to hPYY(3-36).

As another example, [Arg4, Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) comprises 5 amino acid substitutions and 1 deletion as compared to hPYY(3-36), meaning that this  
30 compound has 6 amino acid modifications as compared to hPYY(3-36).

In yet another aspect, the PYY peptides of the invention may exhibit at least 70%, 75% or 80% sequence identity to hPYY(3-36). As an example of a method for determination of the sequence identity between two analogues the two peptides [NMeArg35]hPYY(3-36) and hPYY(3-36) are aligned. The sequence identity of the  
35 [NMeArg35]hPYY(3-36) analogue relative to hPYY(3-36) is given by the total number of aligned residues minus the number of different residues (i.e. the number of aligned

identical residues) divided by the total number of residues in hPYY(3-36). Accordingly, in said example the sequence identity is (34-1)/34.

PYY compounds or PYY analogues of the invention may be described by reference to i) the number of the amino acid residue in hPYY(1-36) which corresponds to the amino acid residue which is changed (i.e., the corresponding position in hPYY(1-36), and to ii) the actual change.

The following is a non-limiting example of suitable analogue nomenclature. [NMeArg35]hPYY(3-36) designates an analogue of the human PYY(1-36), wherein the naturally occurring arginine in position 35 has been substituted with N(alpha)-methylarginine (the arginine residue in the position corresponding to position 35 of hPYY(1-36) has a methyl group added to the alpha-amino group) and the naturally occurring tyrosine and proline in position 1 and 2, respectively, have been deleted.

Likewise, [Trp30]hPYY(3-36) designates an analogue of human PYY(3-36), wherein the naturally occurring leucine in position 30 of human PYY(1-36) has been substituted with tryptophan.

The following is a non-limiting example of suitable nomenclature for a derivative of a PYY analogue. 7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) designates a derivative of an analogue of hPYY(3-36) (SEQ ID NO:2), wherein [Lys7,Gln18,Trp30,NMeArg35] designate the amino acid changes as compared to human PYY(3-36) with the numbers referring to the corresponding positions of PYY(1-36), and wherein the substituent [2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]- is attached to the epsilon amino group of the lysine in the position corresponding to position 7 in hPYY(1-36).

The expressions "a position equivalent to" or "corresponding position" are used to characterise the site of change in a variant PYY sequence by reference to hPYY(1-36).

In general throughout the application, when referring to a particular position of a PYY analogue, the position referred to is the position of the PYY analogue corresponding to that particular position of hPYY(1-36).

The expression used throughout this application, that a PYY compound comprises a particular amino acid at a position corresponding to a certain position of hPYY(1-36), means that the native amino acid in that position has been replaced with that particular amino acid.

Amino acid residues may be identified by their full name, their one-letter code, and/or their three-letter code. These three ways are fully equivalent.

Analogues "comprising" certain specified changes may comprise further changes, when compared to hPYY(1-36). In one aspect, the analogue "has" the specified changes.

5

### **PYY analogues**

A PYY analogue is a PYY peptide in which a number of amino acid residues have been modified when compared to hPYY(1-36). These modifications include substitutions, insertions, and/or deletions, alone or in combination.

10

In a specific aspect, the PYY analogues of the invention include one or more modifications of a "non-essential" amino acid residue. In the context of the invention, a "non-essential" amino acid residue is a residue that can be altered, i.e., deleted or substituted in the human PYY amino acid sequence without abolishing or substantially reducing the activity of the PYY analogue towards the Y2 receptor.

15

Substitutions. In one aspect amino acids may be substituted by conservative substitution. The term "conservative substitution" as used herein denotes that one or more amino acids are replaced by another, biologically similar residue. Examples include substitution of amino acid residues with similar characteristics, e.g. small amino acids, acidic amino acids, polar amino acids, basic amino acids, hydrophobic amino acids and aromatic amino acids.

20

In one aspect, the PYY analogues of the invention may comprise substitutions of one or more unnatural and/or non-amino acids, e.g., amino acid mimetics, into the sequence of PYY.

25

Deletions and Truncations. In one aspect, the PYY analogues of the invention may have one or more amino acid residues deleted from the amino acid sequence of human PYY, alone or in combination with one or more insertions or substitutions.

Insertions. In one aspect, the PYY analogues of the invention may have one or more amino acid residues inserted into the amino acid sequence of human PYY, alone or in combination with one or more deletions and/or substitutions.

30

In one aspect, the PYY analogues of the invention may include insertions of one or more unnatural amino acids and/or non-amino acids into the sequence of PYY.

35

The PYY peptide may be derived from vertebrates, such as human, mouse, sheep, goat, cow, or horse. The term "vertebrate" means members of the subphylum Vertebrata, a primary division of the phylum Chordata that includes the fish, amphibians, reptiles, birds, and mammals, all of which are characterized by a segmented spinal

column and a distinct well-differentiated head. The term "mammal" means humans as well as all other warm-blooded members of the animal kingdom possessed of a homeostatic mechanism in the class Mammalia, e.g., companion mammals, zoo mammals, and food-source mammals. Some examples of companion mammals are  
5 canines (e.g., dogs), felines (e.g., cats) and horses; some examples of food-source mammals are pigs, cattle, sheep, and the like. In one aspect the mammal is a human or a companion mammal. In one aspect the mammal is a human, male or female.

The term "peptide", as e.g. used in the context of the PYY compounds of the  
10 invention, refers to a compound which comprises a series of amino acids interconnected by amide (or peptide) bonds.

The PYY peptides of the invention comprise at least 24 constituent amino acids connected by peptide bonds. In particular embodiments the PYY peptides comprise at least 33 amino acids. In particular embodiments the PYY peptides comprise  
15 at least 34 amino acids.

Amino acids are molecules containing an amine group and a carboxylic acid group, and, optionally, one or more additional groups, often referred to as a side chain.

The term "amino acid" includes proteinogenic (or coded or natural) amino acids  
20 (amongst the 20 standard amino acids), as well as non-proteinogenic (or non-coded or non-natural) amino acids. Proteinogenic amino acids are those which are naturally incorporated into proteins. The standard amino acids are those encoded by the genetic code. Non-proteinogenic amino acids are either not found in proteins, or not produced by standard cellular machinery (e.g., they may have been subject to post-translational  
25 modification). Non-limiting examples of non-proteinogenic amino acids are Aib ( $\alpha$ -aminoisobutyric acid), N( $\alpha$ )-methylarginine, as well as the D-isomers of the proteinogenic amino acids. One example of a D-isomer of a proteinogenic amino acid is the D-isomer of aspartic acid, which can also be written as D-Asp.

In what follows, all amino acids of the PYY compound for which the optical  
30 isomer is not stated is to be understood to mean the L-isomer (unless otherwise specified).

### **PYY derivatives**

The term "derivative" as used herein in the context of a PYY peptide or analogue  
35 means a chemically modified PYY peptide, in which one or more substituents have been covalently attached to the peptide.

In one aspect of the invention, the substituent may be an N-terminal substituent.

Also or alternatively, in one aspect, the substituent may be a modifying group or alternatively, referred to as a protracting moiety.

5

***N-terminal substituent***

In one aspect of the invention, the PYY compound comprises a substituent covalently attached to the alpha-amino group in the amino acid residue in the N-terminus of the PYY compound. In one aspect, the amino acid residues in the positions  
10 corresponding to positions 1-3 of hPYY(1-36) are absent, and the N-terminal substituent is covalently attached to the amino acid residue in the position corresponding to position 4 of hPYY(1-36).

In one aspect, the N-terminal substituent is an alkoxy group. In one aspect, the N-terminal substituent is an alkoxy group comprising up to 12 carbon atoms. In another  
15 aspect, the N-terminal substituent is an alkoxy group comprising up to 6 carbon atoms.

***Modifying group / Protracting moiety***

In one aspect, the PYY compound comprises a substituent or modifying group covalently attached to the amino acid residue in the position corresponding to position 7  
20 of hPYY(1-36). In one further aspect, the substituent or modifying group is capable of forming non-covalent conjugates with proteins, thereby promoting the circulation of the derivative with the blood stream, and also having the effect of protracting the time of action of the derivative, due to the fact that the conjugate of the PYY derivative and albumin is only slowly removed by renal clearance. Thus, the substituent, or modifying  
25 group, as a whole may also be referred to as a protracting moiety.

The modifying group may be covalently attached to a lysine residue of the PYY peptide by acylation, i.e., via an amide bond formed between a carboxylic acid group of the modifying group and the epsilon amino group of the lysine residue. The amino group of lysine could also be coupled to an aldehyde of the modifying group by reductive  
30 amination. In another aspect the thiol group of cysteine could be coupled to a maleimido group of the modifying group by Michael addition or coupled to the chloro- or iodoacetyl group of the modifying group by nucleophilic substitution.

In one aspect, the modifying group may be covalently attached to a lysine residue in a position corresponding to position 7 of hPYY(1-36) by acylation, i.e., via an  
35 amide bond formed between a carboxylic acid group of the modifying group and the epsilon amino group of the lysine residue.

The derivatives of the invention may exist in different stereoisomeric forms having the same molecular formula and sequence of bonded atoms, but differing only in the three-dimensional orientation of their atoms in space. The stereoisomerism of the exemplified derivatives of the invention is indicated in the experimental section, in the names as well as the structures, using standard nomenclature. Unless otherwise stated the invention relates to all stereoisomeric forms of the claimed derivative.

Herein, all amino acids of the PYY compound for which the optical isomer is not stated are to be understood to mean the L-isomer (unless otherwise specified).

### **Pharmaceutically acceptable salts**

The PYY compounds of the invention may be in the form of a pharmaceutically acceptable salt.

Salts are e.g. formed by a chemical reaction between a base and an acid, e.g.:  
$$2\text{NH}_3 + \text{H}_2\text{SO}_4 \rightarrow (\text{NH}_4)_2\text{SO}_4.$$

The salt may be a basic salt, an acidic salt, or it may be neither nor (i.e. a neutral salt). Basic salts produce hydroxide ions and acid salts hydronium ions in water.

The salts of the derivatives of the invention may be formed with added cations or anions between anionic or cationic groups, respectively. These groups may be situated in the peptide moiety, and/or in the side chain of the derivatives of the invention.

Non-limiting examples of anionic groups of the derivatives of the invention include free carboxylic groups in the side chain, if any, as well as in the peptide moiety. The peptide moiety often includes free carboxylic groups at internal acid amino acid residues such as Asp and Glu.

Non-limiting examples of cationic groups in the peptide moiety include the free amino group at the N-terminus, if present, as well as any free amino group of internal basic amino acid residues such as His, Arg, and Lys.

### **FUNCTIONAL PROPERTIES**

In a first functional aspect, the PYY compounds of the invention have a good Y2 receptor potency. Also, or alternatively, in a second aspect, they bind very well to the Y2 receptor. Preferably they are full Y2 receptor agonists as is reflected by their ability to bind strongly to the Y2 receptor combined with the capacity to fully activate the receptor compared to hPYY(1-36) and hPYY(3-36).

Also or alternatively, in a second functional aspect, the invention relates to PYY compounds displaying selectivity towards the Y receptor subtype Y2 as compared to Y receptor subtypes Y1, Y4 and Y5.

Also, or alternatively, in a third functional aspect, the PYY compounds of the invention have improved pharmacokinetic properties. Also, or alternatively, in a fourth functional aspect, the PYY compounds of the invention have increased half-life and/or a decreased clearance. Also, or alternatively, in a fifth functional aspect, they have the effect in vivo of decreasing the blood glucose. Also, or alternatively, in a sixth functional aspect, they have the effect in vivo of decreasing food intake. Also, or alternatively, in a seventh functional aspect, they have the effect in vivo of decreasing body weight.

### **Biological activity – in vitro potency**

According to the first functional aspect, the PYY compounds of the invention are biologically active, or potent.

In a particular embodiment, potency and/or activity refers to in vitro potency, i.e. performance in a functional Y2 receptor assay, more in particular to the capability of activating the human Y2 receptor.

The term half maximal effective concentration ( $EC_{50}$ ) generally refers to the concentration which induces a response halfway between the baseline and maximum, by reference to the dose response curve.  $EC_{50}$  is used as a measure of the potency of a compound and represents the concentration where 50% of its maximal effect is observed.

The in vitro potency of the derivatives of the invention may be determined as described in Example 39, and the  $EC_{50}$  of the derivative in question determined. The lower the  $EC_{50}$  value, the better the potency.

In one aspect of the invention, the derivative of the invention has an in vitro potency determined using the method of Example 39 corresponding to an  $EC_{50}$  at or below 100 nM. In one aspect, the derivative of the invention has an in vitro potency determined using the method of Example 39 corresponding to an  $EC_{50}$  at or below 50 nM. In one aspect, the derivative of the invention has an in vitro potency determined using the method of Example 39 corresponding to an  $EC_{50}$  at or below 25 nM.

### **Biological activity – in vitro receptor binding**

According to the second functional aspect, the PYY compounds of the invention bind very well to the Y2 receptor. This may be determined as described in Example 40.

Generally, the binding to the Y2 receptor should be as good as possible, corresponding to a low  $K_i$  value. The  $K_i$  value is determined by the Cheng-Prusoff equation  $K_i = IC_{50} / (1 + [L] / K_d)$ , wherein  $IC_{50}$  is the half maximal inhibitory concentration of the agonist,  $[L]$  is the concentration of the radioligand and  $K_d$  is the dissociation constant for binding.

As an example, in a particular aspect, the Y2 receptor binding affinity ( $K_i$ ) is below 100 nM. In one aspect of the invention, the Y2 receptor binding affinity ( $K_i$ ) is below 50 nM. In one aspect of the invention, the Y2 receptor binding affinity ( $K_i$ ) is below 10 nM.

### **Biological activity – in vivo pharmacology**

In another particular embodiment the PYY compounds of the invention are potent in vivo, which may be determined as is known in the art in any suitable animal model, as well as in clinical trials.

The diabetic db/db mouse is one example of a suitable animal model, and the blood glucose lowering effect may be determined in such mice in vivo, e.g. as described in Example 42.

In addition, inhibition of food intake in the db/db mice is a suitable model for determination of effect on food intake and body weight as also described in Example 42.

Generally, the glucose lowering effect of a 1  $\mu\text{mol/kg}$  dose should be as good as possible corresponding to a low relative % glucose level.

As an example, in a particular aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 80%. In one aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 70%. In one aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 60%.

As an example, in a particular aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 40%. In one aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 30%. In one aspect of the invention, 16 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 20%.

As an example, in a particular aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 80%. In one aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 70%. In one aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the relative % glucose level is below 60%.



As an example, in a particular aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 40%. In one aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 30%. In one aspect of the invention, 4 hours after dosing (1  $\mu\text{mol/kg}$ ) the % relative food intake is below 20%.

5

### **Pharmacokinetics profile**

According to the third functional aspect, the PYY compounds of the invention have improved pharmacokinetic properties such as increased terminal half-life and/or decreased clearance.

10

Increasing terminal half-life and/or decreasing of the clearance means that the compound in question is eliminated slower from the body. For the compounds of the invention this entails an extended duration of pharmacological effect.

The pharmacokinetic properties of the derivatives of the invention may suitably be determined in-vivo in pharmacokinetic (PK) studies. Such studies are conducted to evaluate how pharmaceutical compounds are absorbed, distributed, and eliminated in the body, and how these processes affect the concentration of the compound in the body, over the course of time.

15

In the discovery and preclinical phase of pharmaceutical drug development, animal models such as the mouse, rat, monkey, dog, or pig, may be used to perform this characterisation. Any of these models can be used to test the pharmacokinetic properties of the derivatives of the invention.

20

The estimate of terminal half-life and/or clearance is relevant for evaluation of dosing regimens and an important parameter in drug development, in the evaluation of new drug compounds.

25

### **Pharmacokinetics profile – half life in vivo in minipigs**

According to the third functional aspect, the derivatives of the invention have improved pharmacokinetic properties.

In a particular embodiment, the pharmacokinetic properties may be determined as terminal half-life ( $T_{1/2}$ ) in vivo in minipigs after i.v. administration, e.g. as described in Example 41 herein.

30

In one aspect of the invention, the terminal half-life in minipigs is at least 10 hours. In one aspect of the invention, the terminal half-life in minipigs is at least 20 hours. In yet another aspect of the invention, the terminal half-life in minipigs is at least 40 hours.

35

**PRODUCTION OF PYY COMPOUNDS**

The production of peptides like the PYY compounds of the present invention is well known in the art.

5           The PYY moiety of the derivatives of the invention may for instance be produced by classical peptide synthesis, e.g., solid phase peptide synthesis using t-Boc or Fmoc chemistry or other well established techniques, see, e.g., Greene and Wuts, "Protective Groups in Organic Synthesis", John Wiley & Sons, 1999, Florencio Zaragoza Dörwald, "Organic Synthesis on solid Phase", Wiley-VCH Verlag GmbH, 2000, and "Fmoc Solid  
10   Phase Peptide Synthesis", Edited by W.C. Chan and P.D. White, Oxford University Press, 2000.

Also, or alternatively, they may be produced by recombinant methods, viz. by culturing a host cell containing a DNA sequence encoding the analogue and capable of expressing the peptide in a suitable nutrient medium under conditions permitting the  
15   expression of the peptide. Non-limiting examples of host cells suitable for expression of these peptides are: Escherichia coli, Saccharomyces cerevisiae, as well as mammalian BHK or CHO cell lines.

The PYY compounds of the invention which include non-natural amino acids and/or covalently attached substituents may e.g. be produced as described in the  
20   experimental part.

Specific examples of methods of preparing a number of the PYY compounds of the invention are included in the experimental part.

**PROTEIN PURIFICATION**

25           The PYY compounds of the present invention may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, and reverse-phase high performance liquid chromatography (RP-HPLC)), electrophoretic procedures, or extraction (see, e.g., Protein Purification, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).  
30

**MODE OF ADMINISTRATION**

The term "treatment" is meant to include both the prevention and minimization of the referenced disease, disorder, or condition (i.e., "treatment" refers to both prophylactic and therapeutic administration of the PYY compounds of the invention or  
35   composition comprising the PYY compounds of the invention) unless otherwise indicated or clearly contradicted by context.

The route of administration may be any route which effectively transports a compound of this invention to the desired or appropriate place in the body, such as parenterally, for example, subcutaneously, intramuscularly or intravenously.

Alternatively, a compound of this invention can be administered orally, pulmonary, rectally, transdermally, buccally, sublingually, or nasally.

## PHARMACEUTICAL COMPOSITIONS

Injectable compositions comprising PYY compounds of the present invention can be prepared using the conventional techniques of the pharmaceutical industry which involve dissolving and mixing the ingredients as appropriate to give the desired end product. Thus, according to one procedure, a PYY compound of this invention is dissolved in a suitable buffer at a suitable pH so precipitation is minimised or avoided. The injectable composition is made sterile, for example, by sterile filtration.

A composition may be a stabilised formulation. The term "stabilised formulation" refers to a formulation with increased physical and/or chemical stability, preferably both. In general, a formulation must be stable during use and storage (in compliance with recommended use and storage conditions) until the expiration date is reached.

The term "physical stability" refers to the tendency of the polypeptide to form biologically inactive and/or insoluble aggregates as a result of exposure to thermo-mechanical stress, and/or interaction with destabilising interfaces and surfaces (such as hydrophobic surfaces). The physical stability of an aqueous polypeptide formulation may be evaluated by means of visual inspection, and/or by turbidity measurements after exposure to mechanical/physical stress (e.g. agitation) at different temperatures for various time periods. Alternatively, the physical stability may be evaluated using a spectroscopic agent or probe of the conformational status of the polypeptide such as e.g. Thioflavin T or "hydrophobic patch" probes.

The term "chemical stability" refers to chemical (in particular covalent) changes in the polypeptide structure leading to formation of chemical degradation products potentially having a reduced biological potency, and/or increased immunogenic effect as compared to the intact polypeptide. The chemical stability can be evaluated by measuring the amount of chemical degradation products at various time-points after exposure to different environmental conditions, e.g. by SEC-HPLC, and/or RP-HPLC.

In one aspect, the invention provides PYY compounds with improved physical stability. In one aspect, the invention provides PYY compounds with improved chemical stability.

**COMBINATION TREATMENT**

The treatment with a PYY compound according to the present invention may also be combined with one or more additional pharmacologically active substances, e.g. selected from antidiabetic agents, antiobesity agents, appetite regulating agents, antihypertensive agents, agents for the treatment and/or prevention of complications resulting from or associated with diabetes and agents for the treatment and/or prevention of complications and disorders resulting from or associated with obesity.

Examples of these pharmacologically active substances are: GLP-1 receptor agonists, insulin, DPP-IV (dipeptidyl peptidase-IV) inhibitors, amylin agonists and leptin receptor agonists.

In one aspect of the invention, a PYY compound according to the present invention is combined with a GLP-1 agonist. The compounds may be supplied in a single-dosage form wherein the single-dosage form contains both compounds, or in the form of a kit-of-parts comprising a preparation of the PYY compound as a first unit dosage form and a preparation of the GLP-1 agonist as a second unit dosage form.

Non-limiting examples of GLP-1 agonists to be combined with the PYY compounds of the present invention are liraglutide, semaglutide, exenatide, dulaglutide, lixisenatide, taspoglutide, and albiglutide.

Liraglutide, a mono-acylated GLP-1 derivative for once daily administration which is marketed as of 2009 by Novo Nordisk A/S, is disclosed in WO 98/08871, Example 37.

WO 2006/097537 discloses additional GLP-1 derivatives including semaglutide (Example 4), a mono-acylated GLP-1 derivative for once weekly administration which is under development by Novo Nordisk A/S.

Exenatide is a synthetic version of exendin-4, a hormone found in the saliva of the Gila monster. It displays biological properties similar to GLP-1.

Dulaglutide is a GLP-1-Fc construct (GLP-1 - linker - Fc from IgG4).

Lixisenatide is based on exendin-4(1-39) modified C-terminally with six Lys residues.

Taspoglutide is the 8-(2-methylalanine)-35-(2-methylalanine)-36-L-argininamide derivative of the amino acid sequence 7-36 of human GLP-1.

Albiglutide is a recombinant human serum albumin (HSA)-GLP-1 hybrid protein, likely a GLP-1 dimer fused to HSA. The constituent GLP-1 peptide is an analogue, in which Ala at position 8 has been substituted by Gly.

**PHARMACEUTICAL INDICATIONS**

The present invention also relates to a PYY compound of the invention for use as

a medicament.

In particular aspects of the invention, the PYY compounds of the invention may be used for the following medical treatments:

(i) prevention and/or treatment of all forms of diabetes, such as hyperglycemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, non-insulin dependent diabetes, MODY (maturity onset diabetes of the young), gestational diabetes, and/or for reduction of HbA1C;

(ii) delaying or preventing diabetic disease progression, such as progression in type 2 diabetes, delaying the progression of impaired glucose tolerance (IGT) to insulin requiring type 2 diabetes, delaying or preventing insulin resistance, and/or delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes;

(iii) improving  $\beta$ -cell function, such as decreasing  $\beta$ -cell apoptosis, increasing  $\beta$ -cell function and/or  $\beta$ -cell mass, and/or for restoring glucose sensitivity to  $\beta$ -cells;

(iv) prevention and/or treatment of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence;

(v) prevention and/or treatment of diabetic complications, such as angiopathy; neuropathy, including peripheral neuropathy; nephropathy; and/or retinopathy;

(vi) improving lipid parameters, such as prevention and/or treatment of dyslipidemia, lowering total serum lipids; increasing HDL; lowering small, dense LDL; lowering VLDL; lowering triglycerides; lowering cholesterol; lowering plasma levels of lipoprotein a (Lp(a)) in a human; inhibiting generation of apolipoprotein a (apo(a)) in vitro and/or in vivo;

(vii) prevention and/or treatment of cardiovascular diseases; and/or

(viii) prevention and/or treatment of sleep apnoea.

The following indications are particularly preferred: Type 2 diabetes, and/or obesity.

In one aspect, a method is disclosed herein for altering energy metabolism in a subject. The method includes administering a therapeutically effective amount of a PYY compound of the invention to the subject, thereby altering energy expenditure. Energy is burned in all physiological processes. The body can alter the rate of energy expenditure

directly, by modulating the efficiency of those processes, or changing the number and nature of processes that are occurring. For example, during digestion the body expends energy moving food through the bowel, and digesting food, and within cells, the efficiency of cellular metabolism can be altered to produce more or less heat.

5 In one aspect a method is disclosed herein for any and all manipulations of the accurate circuitry described in this application, which alter food intake co-ordinately and reciprocally alter energy expenditure. Energy expenditure is a result of cellular metabolism, protein synthesis, metabolic rate, and calorie utilization. Thus, in this embodiment, peripheral administration results in increased energy expenditure, and  
10 decreased efficiency of calorie utilization. In one aspect, a therapeutically effective amount of a PYY compound according to the invention is administered to a subject, thereby increasing energy expenditure.

While "obesity" is generally defined as a body mass index over 30, for purposes  
15 of this disclosure, any subject, including those with a body mass index of less than 30, who needs or wishes to reduce body weight is included in the scope of "obese." Without intending to be limited by theory, it is believed that the effects of peripherally administered PYY compounds of the present invention in the reduction of food intake, in the delay of gastric emptying, in the reduction of nutrient availability, and in the  
20 causation of weight loss are determined by interactions with one or more unique receptor classes in, or similar to, those in the PP family. More particularly, it appears that a receptor or receptors similar to the PYY-preferring (or Y2) receptors are involved.

#### **PARTICULAR EMBODIMENTS**

25 The invention is further described by the following non-limiting embodiments of the invention:

1. A PYY compound comprising a tryptophan at a position corresponding to position  
30 of hPYY(1-36) (SEQ ID NO:1), and a maximum of 10 amino acid modifications as compared to hPYY(3-36), and a pharmaceutically acceptable salt thereof.

2. A PYY compound comprising tryptophan at the position corresponding to position  
30 of hPYY(1-36) (SEQ ID NO:1), and a maximum of 10 amino acid modifications as compared to hPYY(3-36), or a pharmaceutically acceptable salt thereof.

3. A PYY compound according to any one of the preceding embodiments, further comprising N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36) (SEQ ID NO:1).

4. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound comprises N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36) (SEQ ID NO:1).

5. A PYY compound according to any one of the preceding embodiments, further comprising glutamine at a position corresponding to position 18 of hPYY(1-36) (SEQ ID NO:1).

6. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound comprises glutamine at a position corresponding to position 18 of hPYY(1-36) (SEQ ID NO:1).

7. A PYY compound according to any one of the preceding embodiments, further comprising arginine at a position corresponding to position 4 of hPYY(1-36) (SEQ ID NO:1).

8. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound comprises arginine at a position corresponding to position 4 of hPYY(1-36) (SEQ ID NO:1).

9. A PYY compound according to any one of the preceding embodiments, further comprising Aib at a position corresponding to position 28 of hPYY(1-36) (SEQ ID NO:1).

10. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound comprises Aib at a position corresponding to position 28 of hPYY(1-36) (SEQ ID NO:1).

11. A PYY compound according to any one of the preceding embodiments, wherein the positions corresponding to positions 1 and 2 of hPYY(1-36) (SEQ ID NO:1) are absent.

12. A PYY compound according to any one of the preceding embodiments, wherein the positions corresponding to position 1-3 of hPYY(1-36) (SEQ ID NO:1) are absent.

13. A PYY compound according to any one of the preceding embodiments, wherein the positions corresponding to positions 1-3 of hPYY(1-36) (SEQ ID NO:1) are absent, and wherein the PYY compound further comprises an N-terminal substituent, wherein the N-terminal substituent is an alkoxy group comprising up to 12 carbon atoms.

14. A PYY compound according to embodiment 13, wherein the N-terminal substituent is an alkoxy group comprising up to 10 carbon atoms.

15. A PYY compound according to embodiment 13, wherein the N-terminal substituent is an alkoxy group comprising up to 8 carbon atoms.

16. A PYY compound according to embodiment 13, wherein the N-terminal substituent is an alkoxy group comprising up to 6 carbon atoms.

17. A PYY compound according to embodiment 13, wherein the N-terminal substituent is an alkoxy group comprising 6 carbon atoms.

18. A PYY compound according to embodiment 13, wherein the N-terminal substituent is selected from 3-methylbutanoyl, 3-methylpentanoyl or hexanoyl.

19. A PYY compound according to embodiment 13, wherein the N-terminal substituent is 3-methylbutanoyl.

20. A PYY compound according to embodiment 13, wherein the N-terminal substituent is 3-methylpentanoyl.

21. A PYY compound according to embodiment 13, wherein the N-terminal substituent is hexanoyl.

22. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has a maximum of 8 amino acid modifications as compared to hPYY(3-36).



23. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has a maximum of 6 amino acid modifications as compared to hPYY(3-36).

5 24. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has a minimum of 4 amino acid modifications as compared to hPYY(3-36).

10 25. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has a minimum of 6 amino acid modifications as compared to hPYY(3-36).

15 26. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has a minimum of 8 amino acid modifications as compared to hPYY(3-36).

20 27. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has in the range of 4 to 10 amino acid modifications as compared to hPYY(3-36).

28. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has in the range of 6 to 8 amino acid modifications as compared to hPYY(3-36).

25 29. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has in the range of 4 to 6 amino acid modifications as compared to hPYY(3-36).

30 30. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has 4 amino acid modifications as compared to hPYY(3-36).

31. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has 6 amino acid modifications as compared to hPYY(3-36).

35 32. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound has 8 amino acid modifications as compared to hPYY(3-36).

33. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound exhibit at least 70% sequence identity to hPYY(3-36).


34. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound exhibit at least 75% sequence identity to hPYY(3-36).

35. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound exhibit at least 80% sequence identity to hPYY(3-36).

36. A PYY compound according to any one of the preceding embodiments, further comprising a Lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1) and a modifying group attached to the epsilon amino group of the Lysine residue in position 7, wherein said modifying group is defined by **A-B-C-**, wherein **A-** comprises a carboxylic acid, a tetrazole or a sulfonic acid.

37. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound comprises lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1) and a modifying group attached to the epsilon amino group of said lysine, wherein said modifying group is defined by **A-B-C-**.

38. A PYY compound according to any one of embodiments 36-37, wherein **A-** is selected from




wherein a is an integer from 12 to 19, b is an integer from 10 to 16, and c is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-.

39. A PYY compound according to embodiment 38, wherein a is 15, b is 13, or c is 13.

40. A PYY compound according to any one of embodiments 36-37, wherein **A-** is selected from


22



wherein a is an integer from 12 to 19, c is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-.

5    41.    A PYY compound according to embodiment 40, wherein a is 15, or c is 13.


42.    A PYY compound according to any one of embodiments 36-37, wherein **A-** is



wherein a is an integer from 12 to 19, and wherein \* denotes the attachment point to -B-.

43.    A PYY compound according to embodiment 42, wherein a is 15.


44.    A PYY compound according to any one of embodiments 36-37, wherein **A-** is



wherein b is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-.

45.    A PYY compound according to embodiment 44, wherein b is 13.

46.    A PYY compound according to any one of embodiments 36-37, wherein **A-** is




wherein c is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-.

47. A PYY compound according to embodiment 46, wherein c is 13.

48. A PYY compound according to any one of embodiments 36-47, wherein **B-** is selected from


5



wherein d is 1 or 2; e is 1 or 2; and f is 2, 3 or 4; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-.

10


49. A PYY compound according to embodiment 48, wherein **B-** is selected from



15 wherein d is 1 or 2; and f is 2, 3 or 4; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-.


50. A PYY compound according to embodiment 49, wherein d is 1 or 2; and f is 3.

20 51. A PYY compound according to embodiment 48, wherein **B-** is



wherein d is 1 or 2; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-.

52. A PYY compound according to embodiment 48, wherein **B-** is




wherein f is 2, 3 or 4; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-.


5

53. A PYY compound according to embodiment 52, wherein f is 3.

54. A PYY compound according to any one of embodiments 36-53, wherein **-C-** is




or




wherein g is an integer in the range of 1-5, h is an integer in the range of 1-5, i is an integer in the range of 2 to 6, j is an integer in the range of 2 to 6, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36).

55. A PYY compound according to any one of embodiments 36-53, wherein **-C-** is



or




wherein g is an integer in the range of 1-5, h is an integer in the range of 1-5, i is an integer in the range of 2 to 6, j is an integer in the range of 1 to 6, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36).

56. A PYY compound according to any one of embodiments 54-55, wherein g and h are each 1.

57. A PYY compound according to any one of embodiments 54-56, wherein i is selected from 2, 3, 4 or 5 and j is selected from 1, 2 or 3.

58. A PYY compound according to any one of embodiments 36-53, wherein -C- is



VII

wherein g is an integer in the range of 1-5, h is an integer in the range of 1-5, i is an integer in the range of 2 to 6, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36).

59. A PYY compound according to embodiment 58, wherein g and h are each 1.

60. A PYY compound according to any one of embodiments 58-59, wherein i is selected from 2, 4 or 6.

61. A PYY compound according to embodiment 60, wherein i is 2.

62. A PYY compound according to embodiment 60, wherein i is 4.


63. A PYY compound according to embodiment 60, wherein i is 6.

64. A PYY compound according to embodiment 58, wherein g and h are each 1, and i is 2.

65. A PYY compound, wherein the PYY compound has a maximum of 10 amino acid modifications as compared to hPYY(3-36), and wherein the PYY compound comprises


- (i) tryptophan at the position corresponding to position 30 of hPYY(1-36) (SEQ ID NO:1);
- (ii) N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36) (SEQ ID NO:1);
- (iii) glutamine at a position corresponding to position 18 of hPYY(1-36) (SEQ ID NO:1);
- (iv) lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1); and
- (v) a modifying group attached to the epsilon amino group of said lysine, wherein said modifying group is defined by **A-B-C-**, wherein

**A-** is selected from




wherein a is 15, c is 13, and wherein \* denotes the attachment point to -B-;

**B-** is



wherein d is 1 or 2; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-; and

**-C-** is



wherein g and h is each 1, i is selected from 2 or 4, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36); or a pharmaceutically acceptable salt thereof.

66. A PYY compound according embodiment 65, wherein the PYY compound comprises arginine at a position corresponding to position 4 of hPYY(1-36) (SEQ ID NO:1).

67. A PYY compound according to any one of embodiments 65-66, wherein the positions corresponding to positions 1 and 2 of hPYY(1-36) (SEQ ID NO:1) are absent.

68. A PYY compound according to any one of embodiments 65-67, wherein the positions corresponding to positions 1-3 of hPYY(1-36) (SEQ ID NO:1) are absent, and wherein the PYY compound further comprises an N-terminal substituent, wherein the N-terminal substituent is selected from 3-methylbutanoyl, 3-methylpentanoyl or hexanoyl.

69. A PYY compound according to any one of embodiments 65-68, wherein the PYY compound has a maximum of 8 amino acid modifications as compared to hPYY(3-36).

70. A PYY compound, wherein the PYY compound has a maximum of 10 amino acid modifications as compared to hPYY(3-36), wherein the positions corresponding to positions 1-3 of hPYY(1-36) (SEQ ID NO:1) are absent, and wherein the PYY compound comprises

(i) tryptophan at the position corresponding to position 30 of hPYY(1-36) (SEQ ID NO:1);


(ii) N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36) (SEQ ID NO:1);

(iii) glutamine at a position corresponding to position 18 of hPYY(1-36) (SEQ ID NO:1);

(iv) lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1);

(v) a modifying group attached to the epsilon amino group of said lysine, wherein said modifying group is defined by **A-B-C-**, wherein


**A-** is



wherein c is 13, and wherein \* denotes the attachment point to -B-;


**B-** is





wherein d is 1; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-; and

5 -C- is



wherein g and h is each 1, i is 2, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36);

- 10 (vi) arginine at a position corresponding to position 4 of hPYY(1-36) (SEQ ID NO:1); and  
 (vii) an N-terminal substituent, wherein the N-terminal substituent is selected from 3-methylbutanoyl, 3-methylpentanoyl or hexanoyl;  
 or a pharmaceutically acceptable salt thereof.

15 71. A PYY compound according to embodiment 70, wherein the PYY compound has a maximum of 8 amino acid modifications as compared to hPYY(3-36).

71a. A PYY compound according to embodiment 70, wherein the PYY compound has 6 amino acid modifications as compared to hPYY(3-36).

20

72. A PYY compound according to any one of the preceding embodiments, wherein the PYY compound is not a salt.

73. A PYY compound according to any one of the preceding embodiments selected from the following:


25

[Trp30]hPYY(3-36) (SEQ ID NO:3)




; [Trp30,NMeArg35]hPYY3-36 (SEQ ID NO:4)

30



5 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[[4S]-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:7)




10 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[[4S]-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:8)

[illegible]

10 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-  
4-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]butanoyl]-  
amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Aib28,Trp30,NMeArg35]  
hPYY(3-36) (SEQ ID NO:10)

[illegible]


5 Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:11)




10 ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:12)

CCCCCC(=O)RPE[C@@H](CCCCNC(=O)COCCOC(=O)NCCCC(C(=O)O)[C@H](CCCNC(=O)N=[NH2+]=[NH2+])C(=O)N[C@@H](C)C(=O)N[C@@H](Cc1ccc(O)cc1)C(=O)N)C(=O)PGEDASPEELQRYYASLRHYLNWVTRQN(CCCCCCCCCCCCCCCCCCCCCCCCCCS(=O)(=O)O)C(=O)N


5 ; 4-N{alpha}-(3-methyl-pentanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-  
4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:13)



10 ; 4-N{alpha}-(3-methyl-pentanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-  
4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:14)




; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-  
 [[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]butanoyl]amino]-  
 ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]  
 hPYY(4-36) (SEQ ID NO:15)



5


; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-  
 (16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-  
 10 ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:16)




; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-  
 15 (16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]-  
 amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(4-36) (SEQ ID NO:17)

C[C@H](C)CC(=O)R[PE-NH-CH(C(=O)PGEDASPEELQRYRYASLRHYLNWVTRQNH-CH(C(=O)N[C@@H](CCCNC(N)=N)C(=O)N[C@@H](CCCCNC(=O)COCCOCNC(=O)COCCOCNC(=O)COCCOCNC(=O)CCCCCS(=O)(=O)O)]C(=O)N[C@@H](Cc1ccc(O)cc1)C(=O)N]

5 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-  
(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:18)




10 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-  
(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:19)





36

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(14-sulfotetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:20)



5

74. A PYY compound according to any one of the preceding embodiments selected from the following:

10 [Trp30]hPYY(3-36) (SEQ ID NO:3)



; [Trp30,NMeArg35]hPYY3-36 (SEQ ID NO:4)



15


; 7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:5)

[illegible][illegible]

10 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-4-(17-  
carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-  
[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:7)

[illegible]


; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:8)



; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[4-[16-(1H-tetrazol-5-yl)-  
hexadecanoylsulfamoyl]butanoylamino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,  
Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:9)

[illegible]


5 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-  
4-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]butanoyl]-  
amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Aib28,Trp30,NMeArg35]  
hPYY(3-36) (SEQ ID NO:10)



10 ; 7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-  
butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,  
Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:11)

[illegible]

5



10

[illegible]

10 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-  
[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]butanoyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]  
hPYY(4-36) (SEQ ID NO:15)

[illegible][illegible]


10 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-  
(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]-  
amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(4-36) (SEQ ID NO:17)

C[C@H](C)CC(=O)[R]P[NH-][CH(C(=O)[PGEDASPEELQRYYASLRHYLNWVTRQNHC(=O)N[C@@H](Cc1ccc(O)cc1))C(N)=N]CCCNC(=O)COCCOCNC(=O)COCCOCNC(=O)CC[C@H](NC(=O)CCCCS(=O)(=O)O)C(=O)OC[C@H](C)CC(=O)RPE-N[C@@H](CCCCNC(=O)COCCOCN(C(=O)OCCCCCS(=O)(=O)O)CCCNC(=O)COCCOCN(C(=O)OCC[C@H](N)C(=O)O)CC[C@H](N)C(=O)NCC[C@H](N)C(=O)N[C@@H](Cc1ccc(O)cc1)C(=O)N[illegible]




44

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(14-sulfotetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:20)



5

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-carboxy-hexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:21)




10

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(14-carboxy-tetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:22)


[illegible]

5 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-  
(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]-[Arg4,Lys7,Gln18,Ile28,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:23)




; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:24)

46




5 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-  
 [[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]-  
 ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]et  
 hoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:25)




[illegible]

10




15

48




5 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:29)




10

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[(2S)-2-amino-6-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-4-oxobutanoyl]amino]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:29)


amino]hexanoyl]]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:30)




; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(15-carboxypentadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:31)



; 7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Val3,Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:32)




5 ; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,D-Asp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:33)




51

; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[2-(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:34)



5


; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[2-(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,D-isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:35)




10



; 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(4-36) (SEQ ID NO:36)



5 ; 7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(3-36) (SEQ ID NO:37)



10

75. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:3.

76. A PYY compound according to any one of embodiments 1-2, wherein the PYY  
15 compound is SEQ ID NO:4.

77. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:5.

78. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:6.

5 79. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:7.

80. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:8.

10 81. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:9.

82. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:10.

15 83. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:11.

84. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:12.

20

85. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:13.

25 86. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:14.

87. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:15.

30 88. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:16.

89. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:17.

35

90. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:18.

91. A PYY compound according to any one of embodiments 1-2, wherein the PYY  
5 compound is SEQ ID NO:19.

92. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:20.

10 93. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:21.

94. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:22.

15 95. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:23.

96. A PYY compound according to any one of embodiments 1-2, wherein the PYY  
20 compound is SEQ ID NO:24.

97. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:25.

25 98. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:26.

99. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:27.

30 100. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:28.

101. A PYY compound according to any one of embodiments 1-2, wherein the PYY  
35 compound is SEQ ID NO:29.

102. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:30.

5 103. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:31.

104. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:32.

10 105. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:33.

106. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:34.

15

107. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:35.

20 108. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:36.

109. A PYY compound according to any one of embodiments 1-2, wherein the PYY compound is SEQ ID NO:37.

25 110. A PYY compound according to any one of the preceding embodiments which is a human Y2 receptor agonist.

111. A PYY compound according to any one of the preceding embodiments which is a full human Y2 receptor agonist.

30

112. A PYY compound according to any one of the preceding embodiments which is a selective human Y2 receptor agonist.

35 113. A PYY compound according to any one of the preceding embodiments which is a selective full human Y2 receptor agonist.

114. A PYY compound according to any one of the preceding embodiments which is capable of activating the human Y2 receptor.

5 115. A PYY compound according to any one of the preceding embodiments which is capable of activating the human Y2 receptor in an assay with whole cells expressing the human Y2 receptor.

10 116. A PYY compound according to any one of the preceding embodiments which is capable of activating the human Y2 receptor in the Actone functional potency assay of example 39.

117. A PYY compound according to any one of the preceding embodiments which is capable of binding to the human Y2 receptor.

15 118. A PYY compound according to any one of the preceding embodiments which is capable of binding to the human Y2 receptor, wherein the binding to the human Y2 receptor is measured in a competitive binding assay, such as the assay of example 40.

20 119. A PYY compound according to any one of the preceding embodiments which has improved pharmacokinetic properties.

120. A PYY compound according to any one of the preceding embodiments which has an increased half-life and/or a decreased clearance.

25 121. A PYY compound according to any one of the preceding embodiments which has the effect in vivo of decreasing the blood glucose determined in a single-dose study in a db/db mouse model.

30 122. A PYY compound according to any one of the preceding embodiments which has the effect in vivo of decreasing food intake determined in a single-dose study in a db/db mouse model.

123. A pharmaceutical composition comprising a PYY compound according to any one of embodiments 1-122, and at least one pharmaceutically acceptable excipient.

124. A PYY compound according to any one of embodiments 1-122, for use as a medicament.

125. A PYY compound according to any one of embodiments 1-122, for use in the treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression.

126. A PYY compound according to any one of embodiments 1-122, for use in the treatment and/or prevention of diabetes.

127. A PYY compound according to any one of embodiments 1-122, for use in the treatment and/or prevention of type 2 diabetes.

128. Use of a PYY compound according to any one of embodiments 1-122, for the manufacture of a medicament for the treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression.

129. Use of a PYY compound according to any one of embodiments 1-122, for the manufacture of a medicament for the treatment and/or prevention of diabetes.

130. Use of a PYY compound according to any one of embodiments 1-122, for the manufacture of a medicament for the treatment and/or prevention of type 2 diabetes.

131. A method of treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or  
5 for delaying or preventing diabetic disease progression by administering a pharmaceutically active amount of a PYY compound according to any one of embodiments 1-122.

132. A method of treatment and/or prevention of diabetes by administering a  
10 pharmaceutically active amount of a PYY compound according to any one of embodiments 1-122.

133. A method of treatment and/or prevention of type 2 diabetes by administering a pharmaceutically active amount of a PYY compound according to any one of  
15 embodiments 1-122.

134. A PYY compound according to any one of embodiments 1-122, for use in the treatment and/or prevention of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or  
20 preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence.

25 135. A PYY compound according to any one of embodiments 1-122, for use in the treatment and/or prevention of obesity.

136. Use of a PYY compound according to any one of embodiments 1-122, in the manufacture of a medicament for the treatment and/or prevention of eating disorders,  
30 such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or  
35 urine incontinence.

137. Use of a PYY compound according to any one of embodiments 1-122, in the manufacture of a medicament for the treatment and/or prevention of obesity.

138. A method of treatment and/or prevention of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence by administering a pharmaceutically active amount of a PYY compound according to any one of embodiments 1-122.

139. A method of treatment and/or prevention of obesity, by administering a pharmaceutically active amount of a PYY compound according to any one of embodiments 1-122.

140. A pharmaceutical composition comprising a PYY compound according to any one of embodiments 1-122, a GLP-1 agonist, and at least one pharmaceutically acceptable excipient.

141. A pharmaceutical composition according to embodiment 140, wherein the GLP-1 agonist is liraglutide.

142. A pharmaceutical composition according to embodiment 140, wherein the GLP-1 agonist is semaglutide.

143. A pharmaceutical composition according to any one of embodiments 140-142, wherein the PYY compound is SEQ ID NO.18.

144. A pharmaceutical composition according to any one of embodiments 140-143 for use as a medicament.

145. A pharmaceutical composition according to any one of embodiments 140-143, for use in the treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for



delaying or preventing diabetic disease progression and/or eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence.

146. A pharmaceutical composition according to any one of embodiments 140-143, for use in the treatment and/or prevention of diabetes.

147. A pharmaceutical composition according to any one of embodiments 140-143, for use in the treatment and/or prevention of type 2 diabetes.

148. A pharmaceutical composition according to any one of embodiments 140-143, for use in the treatment and/or prevention of obesity.

149. Use of a pharmaceutical composition according to any one of embodiments 140-143, for the manufacture of a medicament for the treatment and/or prevention diabetes and/or obesity.

150. A method of treatment and/or prevention of all forms of diabetes and related diseases, such as eating disorders, diabetic complications, cardiovascular diseases and/or sleep apnoea; and/or for improving lipid parameters, improving  $\beta$ -cell function, and/or for delaying or preventing diabetic disease progression by administering a pharmaceutically active amount of the pharmaceutical composition according to any one of embodiments 140-143.

151. A method of treatment and/or prevention of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; delaying gastric emptying; increasing physical mobility; and/or prevention and/or treatment of comorbidities to obesity, such as osteoarthritis and/or urine incontinence by

administering a pharmaceutically active amount the pharmaceutical composition according to any one of embodiments 140-143.

## EXAMPLES

- 5            This experimental part starts with a list of abbreviations, and is followed by a section including general methods for synthesising and characterising compounds of the invention. Then follows a number of examples which relate to the preparation of specific PYY compounds, and at the end a number of examples have been included relating to the activity and properties of these compounds (section headed pharmacological methods).
- 10           The examples serve to illustrate the invention.

### List of Abbreviations

- ACN: acetonitrile  
Aib:  $\alpha$ -aminoisobutanoic acid  
15    Boc: tert butyloxycarbonyl  
CH<sub>3</sub>CN: acetonitrile  
cpm: counts per minute  
DCM: dichloromethane  
DIC: Diisopropylcarbodiimide  
20    DIPEA: diisopropylethylamine  
DMF: N,N-dimethylformamide  
Et<sub>2</sub>O: diethyl ether  
Fmoc: 9 H-fluoren-9-ylmethoxycarbonyl  
HFIP: Hexafluoroisopropanol  
25    HMWP: High molecular weight proteins  
h: hours  
H<sub>2</sub>O: water  
HOAc: acetic acid  
HOAt: 1-Hydroxy-7-azabenzotriazole  
30    HOBt: 1-Hydroxybenzotriazole  
Min: minutes  
Mtt: 4-methyltrityl  
MW: Molecular weight  
NMeArg: N( $\alpha$ )-methyl-L-arginine  
35    NMF: 1-Methyl-formamide

NMP: 1-Methyl-pyrrolidin-2-one

OtBu: tert butyl ester

Pbf: 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl

PyBOP: benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate

5 rpm: rounds per minute

r.t: Room temperature

tBu: tert butyl

TFA: trifluoroacetic acid

TIPS: triisopropylsilane

10 Trt: triphenylmethyl

## Materials and Methods

### General Methods of Preparation

15 This section relates to methods for solid phase synthesis of peptide backbone and synthesis of side chain attached to backbone (SPPS methods, including methods for the coupling of amino acids, the de-protection of Fmoc-amino acids, methods for cleaving the peptide from the resin, and for its purification).

#### 20 1. Synthesis of resin bound protected peptide backbone

Procedure for the automatic step-wise assembly of peptide backbone. The protected peptidyl resin was synthesized according to the Fmoc strategy on a solid phase peptide synthesiser Prelude (Protein Technologies, Tucson, USA) either 0.25 mmol scale or 0.4 mmol scale using the manufacturer supplied machine protocols. The Fmoc-protected amino acid derivatives used were the standard recommended: Fmoc-Ala-OH, Fmoc-Arg(Pbf)-OH, Fmoc-Asn(Trt)-OH, Fmoc-Asp(OtBu)-OH, Fmoc-Cys(Trt)-OH, Fmoc-Gln(Trt)-OH, Fmoc-Glu(OtBu)-OH, Fmoc-Gly-OH, Fmoc-His(Trt)-OH, Fmoc-Ile-OH, Fmoc-Leu-OH, Fmoc-Lys(Boc)-OH, Fmoc-Lys(Mtt)-OH, Fmoc-Met-OH, Fmoc-Phe-OH, Fmoc-Pro-OH, Fmoc-Ser(tBu)-OH, Fmoc-Thr(tBu)-OH, Fmoc-Trp(Boc)-OH, Fmoc-Tyr(tBu)-OH, or, 25 Fmoc-Val-OH etc. supplied from e.g., Bachem, Iris Biotech, Protein Technologies or Novabiochem. If nothing else is specified the natural L-form of the amino acids are used. Coupling was done by the use of DIC (dicyclohexylcarbodiimide) and Ozyma Pure (ethyl 2-cyano-2-(hydroxyimino)-acetate, Merck, Novabiochem, Switzerland) mediated couplings in NMP (N-methyl pyrrolidone). The coupling of the Fmoc-amino acid was done 30 as described above using 4-8 time excess of amino acid relative to resin substitution (4-8 eq). Coupling time ranged from 1 hour up to 4 hours. The Fmoc-Arg(pbf)-OH was

coupled using a double coupling procedure (1 hour + 1 hour). The resin used for the synthesis of the peptide amides can be Tentagel RAM (Rapp Polymere, Germany), Rink amid ChemMatrix resin (Matrix Innovation, Canada) Rink-Amide resin (Merck/Novabiochem). The protected amino acid derivatives used were standard Fmoc-

amino acids (supplied from e.g. Protein Technologies, or Novabiochem. The epsilon

amino group of lysine to be derivatised was protected with Mtt. The N-terminal amino acid or building was coupled as a Boc-protected amino acid, e.g., Boc-Ile. Alternatively

isovaleric acid was coupled according to the above described coupling procedure for the Fmoc-amino acids. The step-wise solid phase assembly on the Prelude was done using

the following steps: 1) deprotection (removal of Fmoc) by the use of 25% piperidine in NMP for 2x4 min., step 2) Wash (removal of piperidine) with NMP and DCM, step 3)

Coupling of Fmoc-amino acid (0.3M Fmoc-amino acid in 0.3M Oxyma Pure in NMP) 4-8 eq excess for 1-4 hours coupling initiated by adding 1/10 volume of 3M DIC in NMP and

1/10 volume collidine in NMP. Mixing was done by occasional bubbling with nitrogen, step

4) Wash (removal of excess amino acid and reagents by the use of NMP and DCM). Last step included washing with DCM which made the resin ready for attachment of albumin binding moiety on lysine side chain.

## 2. Attachment of modifying groups to resin bound protected peptide backbone

*Procedure for manual removal of Mtt-protection (lysine(Mtt)):* Before synthesis of the modifying group, the Mtt group on the site of attachment (lysine) must be removed. The resin was placed in a syringe or reaction flask and treated with 75% hexafluoroisopropanol (HFIP) + 25% DCM for 2 X 30 minutes to remove the Mtt group. The resin was then washed with DCM and NMP as described above and neutralized with 5% DIPEA

(neutralisation step) in NMP or 25% piperidine in NMP followed by NMP washing before coupling the albumin moiety. Alternatively, the neutralisation step was omitted.

*Procedure for Prelude removal of Mtt-protection (Lysine(Mtt)):* On the Prelude the resin was treated with 75% hexafluoroisopropanol (HFIP) + 25% DCM for 2 x 2 minutes

followed 2 X 30 minutes to remove the Mtt group on the lysine. The resin was then washed with DCM and NMP followed by a neutralisation step using 25% piperidine in NMP by 4 minutes, and was then ready for the synthesis of the modifying group.

## *Procedure for manual synthesis of modifying groups onto a lysine residue:*

The building blocks Fmoc-8-amino-3,6-dioxaoctanoic acid (CAS No. 166108-71-0), Fmoc-TTDS-OH (CAS No. 172089-14-4, IRIS Biotech GmbH), Fmoc-L-Glu-OtBu (84793-

07-7), and eicosanedioic acid mono-tert-butyl ester (CAS No. 843666-40-0) were coupled using DIC and Oxyma Pure in 4-8 eq relative to resin substitution. The coupling time was 2-16 hours usually followed by a capping step using 1 M acetic anhydride for 15-60 min. The Fmoc-group was removed by 25% piperidine in NMP for 10-30 min.

5 followed by washing.

The 16-sulfonic hexadecanoic acid was solubilised in NMP or N-methylformamid (NMF) at 60 degree Celsius or above and activated by PyBOP 1 eq relative to the sulfonic hexadecanoic acid and 2eq of diisopropylethylamine (DIPEA) relative to sulfonic hexadecanoic acid was also added. The peptidyl resin was washed with hot NMP or NMF  
10 just prior to the addition of activated sulfonic hexadecanoic acid. An excess of 3-4 of the sulfonic building block was used and coupling allowed to proceed > 16 hours.

*Procedure for automated synthesis of modifying groups onto a lysine residue:*

For the synthesis of the modifying groups the following building blocks were used: Fmoc-  
15 8-amino-3,6-dioxaoctanoic acid, Fmoc-TTDS-OH, Fmoc-Glu-OtBu, and eicosanedioic acid mono-tert-butyl ester (CAS No. 843666-40-0). Modifying groups were coupled using DIC and Oxyma Pure in 4-8 eq relative to resin substitution. The coupling time was 2-16 hours usually followed by a capping step using 1 M acetic anhydride for 20 min. The Fmoc-group was removed by 25% piperidine in NMP for 2x4 min. followed by washing as  
20 described in the SPPS of the peptide backbone. All other synthesis steps were also the same as described above with the backbone synthesis. The coupling of 16-sulfonic hexadecanoic acid was done by the manual procedure as described above using pyBOP as coupling reagent.

25 3. Cleavage of resin bound peptide with or without attached modifying groups and purification

Prior to TFA deprotection the peptidyl resin was washed with DCM or diethyl ether and dried. The peptide and side chain protection groups were removed by addition of 20-40 ml (0.25 mmol scale) 30-60 (0.4 mmol scale) ml 92% TFA, 5% TIPS and 3% H<sub>2</sub>O for 2-4  
30 hours. Then TFA was filtered and in some cases concentrated by a stream of argon and diethyl ether was added to precipitate the peptide. The peptide was washed three-five times with diethyl ether and dried.

**General Methods of Detection and Characterisation**

35 This section relates to methods for detection and characterisation of the resulting peptides, including LCMS, MALDI and UPLC methods.

1. LC-MS method (LCMS1)

Agilent Technologies LC/MSD TOF (G1969A) mass spectrometer was used to identify the molecular weight of the peptide after elution from an Agilent 1200 series HPLC system.

- 5 The de-convolution of the mass data was calculated using the Agilents software.

Eluents:

Buffer A: 0.1% TFA in water

Buffer B: 0.1% TFA in CH<sub>3</sub>CN

- 10 LC-MS Waters Acquity (LCMS2)

LC-system: Waters Acquity UPLC

Column: Waters Acquity UPLC BEH, C-18, 1.7µm, 2.1mm x 50mm

Detector: Waters (Micromass) LCT Premier XE

Linear gradient: 5 % to 95 % B

- 15 Gradient run-time: 4.0 minutes

Total run-time: 7.0 minutes

Flow rate: 0.4 ml/minutes

Column temperature: 40°C

Solvent A: 99.90 % MQ-water, 0.1% formic acid

- 20 Solvent B: 99.90 % acetonitrile, 0.1 % formic acid

2. UPLC methods

*Method UPLC2*

Buffer A: 0.05% TFA

- 25 Buffer B: CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.4ml/min

Gradient: 5-95% B, (16min),

Column: Acquity UPLC BEH C18, 1.7µm, 2.1 x 150 mm column

Column temp: 40 °C

30

*Method UPLC26v01*

Buffer A: 0.05% TFA

Buffer B: CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.45 ml/min

- 35 Gradient: 5-60% Buffer B (0,5 – 4 min.)

Column: Acquity UPLC BEH C18 1.7µm, 2.1 x 50 mm

Column temperature: 40 °C

*Method UPLC29v01*

Buffer A: 0.05% TFA

5 Buffer B CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.45 ml/min

Gradient: 15-35% Buffer B (0,5 – 4 min.)

Column: Acquity UPLC BEH C18 1.7µm, 2.1 x 50 mm

Column temperature: 40 °C

10

*Method UPLC30v01*

Buffer A: 0.05% TFA

Buffer B: CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.45 ml/min

15 Gradient: 20-40% Buffer B (0.5 – 4 min.)

Column: Acquity UPLC BEH C18 1.7µm, 2.1 x 50 mm

Column temperature: 40 °C

*Method UPLC31v01*

20 Buffer A: 0.05% TFA

Buffer B: CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.45 ml/min

Gradient: 25-45% Buffer B (0.5 – 4 min.)

Column: Acquity UPLC BEH C18 1.7µm, 2.1 x 50 mm

25 Column temperature: 40 °C

*Method UPLC02v01*

System: Waters Acquity UPLC system

Buffer A: 0.05% TFA in H<sub>2</sub>O

30 Buffer B: CH<sub>3</sub>CN + 0.05% TFA

Flow: 0.40 ml/min

Gradient: 5-95 % Buffer B (16 min.)

Column: Acquity UPLC BEH C18 1.7µm, 2.1 x 150 mm

Column temperature: 40 °C

35

*Method UPLC07v01*

System: Waters Acquity UPLC system

Buffer A: 0.09 M di-Ammonium Hydrogen Phosphate (aq) and 10% Acetonitrile, pH 3.6

Buffer B: 20% Isopropanole, 20% Water and 60% Acetonitrile

Flow: 0.50 ml/min

5 Gradient: 35-65 % Buffer B (2-17 min.)

Column: Phenomenex Kinetex C18, 1.7µm, 2.1 mm x 150 mm column

Column temperature: 60 °C

*Method UPLC16v01*

10 System: Waters Acquity UPLC system

Buffer A: 0.2 M Sodium Sulfate, 0.02 M di-Sodium Hydrogen Phosphate, 0.02 M Sodium di-Hydrogen Phosphate, 90% Water and 10% Acetonitrile, pH 7.2

Buffer B: 70% Acetonitrile, 30% Water

Flow: 0.40 ml/min

15 Gradient: 20-50 % Buffer B (3-20 min.)

Column: ACQUITY UPLC BEH Shield RP18, 1.7µm, 2.1 mm x 150 mm column

Column temperature: 60 °C

*Method UPLC60*

20 System: Waters Acquity UPLC system

Buffer A: 0.02 M Sodium Sulfate, 0.02 M di-Sodium Hydrogen Phosphate, 0.02 M Sodium di-Hydrogen Phosphate, 90% Water and 10% Acetonitrile, pH 7.2

Buffer B: 70% Acetonitrile, 30% Water

Flow: 0.40 ml/min

25 Gradient: 20-50 % Buffer B (3-20 min.)

Column: ACQUITY UPLC BEH Shield RP18, 1.7µm, 2.1 mm x 150 mm column

Column temperature: 60 °C

*Method UPLC17*

30 System: Waters Acquity UPLC system

Buffer A: 0.2 M Sodium Sulfate, 0.02 M di-Sodium Hydrogen Phosphate, 0.02 M Sodium di-Hydrogen Phosphate, 90% Water and 10% Acetonitrile, pH 7.2

Buffer B: 70% Acetonitrile, 30% Water

Flow: 0.40 ml/min

35 Step gradient: 10-20% B over 3 minutes, then 20-80% B over 17 minutes, then 80-90% B over 1 minute



Column: ACQUITY UPLC BEH Shield RP18, 1.7 $\mu$ m, 2.1 mm x 150 mm column

Column temperature: 60 °C

*Method UPLC61*

- 5 System: Waters Acquity UPLC system  
Buffer A: 0.02 M Sodium Sulfate, 0.02 M di-Sodium Hydrogen Phosphate, 0.02 M Sodium di-Hydrogen Phosphate, 90% Water and 10% Acetonitrile, pH 7.2  
Buffer B: 70% Acetonitrile, 30% Water  
Flow: 0.40 ml/min
- 10 Gradient: 10-20% buffer B (0-3 min) 20-80 % Buffer B (3-20 min.)  
Column: ACQUITY UPLC BEH Shield RP18, 1.7 $\mu$ m, 2.1 mm x 150 mm column  
Column temperature: 60 °C

*Method UPLC- AP-01*

- 15 Buffer A: 0,1% TFA in H<sub>2</sub>O  
Buffer B: CH<sub>3</sub>CN + 0,1% TFA  
Flow: 0,40 ml/min  
Gradient: 5-95 % Buffer B (16 min.)  
Column; Acquity UPLC BEH130; 150x 2.1; 1.7 $\mu$ m
- 20 Column temperature: 40 °C

*Method UPLC- AP-02*

- Buffer A: 20mM Na<sub>2</sub>HPO<sub>4</sub> , 20mM NaH<sub>2</sub>PO<sub>4</sub>, 200mM Na<sub>2</sub>SO<sub>4</sub> in 90% water / 10% acetonitrile , pH 7.20
- 25 Buffer B: 70% acetonitrile/ 30% water  
Flow: 0,40 ml/min  
Gradient: 10-20 % Buffer B (0-3min.); 20-50% buffer B (3-20 min); 50-80% (20-21 min)  
Column; Acquity UPLC BEH Shield, RP18 1.7  $\mu$ m, 2.1 x 150 mm
- 30 Column temperature: 40 °C

3. MALDI-MS method

- Molecular weights of the peptides were determined using matrix-assisted laser desorption time of flight mass spectroscopy (MALDI-MS), recorded on a Microflex (Bruker). A matrix
- 35 of alpha-cyano-4-hydroxy cinnamic acid was used. The molecular weight of the product was calculated based on the result of MALDI-MS analysis using the software supplied from the manufacturer.

### Synthesis of Intermediates

#### Synthesis of 16-sulfo-hexadecanoic acid

16-Hexadecanolide (997 g, 3.92 mol) was dissolved in methanol (15.1 L) and  
5 toluene-4-sulfonic acid monohydrate (90.0 g, 0.473 mol) was added. Reaction mixture  
was heated in 50 L reactor at 55 °C for 16 hours. After cooling down sodium hydrogen  
carbonate (56.0 g, 0.67 mol) was added and the reaction mixture was stirred for 15 min.  
Solvent was evaporated on Heidolph 20 L rotary evaporator. Ethyl acetate (12 L) was  
added and the mixture was extracted with 5% solution of sodium hydrogen carbonate  
10 (10 L). Organic layer was separated; emulsion layer was extracted with ethyl acetate (3  
x 3 L), white insoluble muddy material was separated and ethyl acetate layer was  
washed again with 5% solution of sodium hydrogen carbonate (5 L). Organic layers were  
combined and washed with saturated solution of sodium hydrogen carbonate (5 L) and  
brine (10 L). Solvent was evaporated on Heidolph 20 L rotary evaporator. Crude product  
15 was crystallized from hexanes (8 L). Hot solution in hexanes was decanted and then let  
to crystallize in ice bath. The material was filtered on large frit and washed with cold  
hexanes (2 L). Pure material was dried in *vacuo*.

Yield: 1062.2 g (95%).  $R_F$  (SiO<sub>2</sub>, dichloromethane/methanol 95:5): 0.65.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>,  $\delta_H$ ): 3.67 (s, 3 H); 3.67-3.60 (m, 2 H); 2.30 (t,  
20 J=7.5 Hz, 2 H); 1.67-1.53 (m, 4 H); 1.25 (s, 22 H).

The above ester (957 g, 3.34 mol) was dissolved in dichloromethane (7 L) on  
Heidolph 20 L rotary evaporator. Triethylamine (695 mL, 4.98 mol) was added, reaction  
mixture was cooled to 0 °C (by putting ice into evaporator bath) and methanesulfonyl  
25 chloride (325 mL, 4.19 mol) in dichloromethane (200 mL) was added slowly during 10  
minutes by external tubing using small *vacuum*. Then the reaction mixture was heated to  
35 °C for 1 hour. NMR analysis showed complete conversion. Water was added (690 mL)  
and solvents were evaporated. Ethyl acetate (8 L) was added and the mixture was  
washed with 1 M hydrochloric acid (4 L) and 5% solution of sodium carbonate (4 L).  
30 Since sodium carbonate extraction formed an emulsion this layer was extracted with  
ethyl acetate (4 L) and added to main portion. Combined ethyl acetate layer was washed  
with brine (4 L), dried over anhydrous sodium sulfate and filtered. Solvent was  
evaporated giving 16-methanesulfonyloxy-hexadecanoic acid methyl ester as white solid.  
Yield: 1225.4 g (100%).

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, δ<sub>H</sub>): 4.22 (t, J=6.6 Hz, 2 H); 3.66 (s, 3 H); 3.00 (s, 3 H); 2.30 (t, J=7.5 Hz, 2 H) 1.82-1.67 (m, 2 H); 1.68-1.54 (m, 2 H); 1.36-1.17 (m, 22 H).

The above mesylate (1.23 kg, 3.34 mol) was dissolved in acetone (8 L) and lithium bromide (585 g, 6.73 mol) was added and the reaction mixture was heated on Heidolph 20 L rotary evaporator at 50 °C for 12 hours. After cooling down solvent was evaporated, ethyl acetate (10 L) was added and the mixture was washed with 5% solution of sodium hydrogencarbonate (3 x 15 L) and brine (8 L). Solvent was evaporated to dryness to yield 16-bromo-hexadecanoic acid methyl ester as pale yellow oil which started to crystallize.

Yield: 1219 g (105%); contains acetone and product of acetone aldolization.

R<sub>F</sub> (SiO<sub>2</sub>, hexanes/ethyl acetate 9:1): 0.90.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, δ<sub>H</sub>): 3.65 (s, 3 H); 3.42 (t, J=6.9 Hz, 2 H); 2.32 (t, J=7.5 Hz, 2 H); 1.92-1.77 (m, 2 H) 1.69-1.53 (m, 2 H); 1.50-1.35 (m, 2 H); 1.25 (bs, 10 H).

Solutions of sodium sulfite (327 g, 2.60 mol) in water (1.26 L) and 16-bromo-hexadecanoic acid methyl ester (728 g, 2.00 mol, 96% purity) in 1-propanol (945 mL) and methanol (420 mL) were heated to reflux in 6 L reactor equipped with mechanical stirrer for 48 hours. The reaction mixture was cooled to 27 °C and diluted with tetrahydrofuran (2 L). Reaction mixture was filtered and solid material was washed with tetrahydrofuran (3 x 700 mL). Filtrate was cooled to 0 °C and another portion of material was precipitated. This precipitate was filtered and washed with tetrahydrofuran (2 x 200 mL). Solids were combined and mixed with water (8.4 L) in 20 L pot. Solution of sodium hydroxide (120 g, 3.00 mol) was added. The mixture was heated to boiling for about 5 hours. Solution of sulfuric acid (430 mL, 8.00 mol) in water (500 mL) was slowly added into the reaction mixture (sulfur dioxide is formed). Reaction mixture was heated to boiling for 10 minutes and then let to cool to 15 °C (ice bath). The mixture was filtered on Büchner funnel through filter paper Seitz (several layers filter) applying *vacuo*. This procedure was very slow and took two days. Solid material was several times washed with distilled water until pH of filtrate was between 2 and 3. This procedure took about three days. Muddy white material was dried in oven at 80 °C giving desired product.

Yield: 510 g (76%).

<sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>, δ<sub>H</sub>): 2.45-2.33 (m, 2 H); 2.18 (t, J=7.3 Hz, 2 H); 1.60-1.40 (m, 4 H); 1.24 (s, 22 H).

MS-ESI (neg, sample in H<sub>2</sub>O/MeCN + NaHCO<sub>3</sub>; m/z) : 335.5 (M-H)<sup>-</sup>, 357.5 (M-2H+Na)<sup>-</sup>, 167.3 (M-2H)<sup>2-</sup>

Synthesis of 14-sulfo-hexadecanoic acid

1 M Solution of borane-tetrahydrofuran complex in tetrahydrofuran (52 mL, 52,0 mmol) was added dropwise to a solution of tetradecanedioic acid mono-tert-butyl ester (10.0 g, 31.9 mmol) in dry tetrahydrofuran (75 mL) at 0 C under argon. The resulting solution was stirred at 0 C for 2 hrs, then the cooling bath was removed and the mixture stirred at room temperature overnight. Saturated aqueous solution of sodium hydrogencarbonate (150 mL) was added and the resulting mixture was extracted with dichloromethane (3 x 100 mL). Combined organic extracts were extracted with 5% aqueous solution of sodium carbonate (1 x 150 mL) and 10% aqueous solution citric acid (1 x 100 mL). Combined organic extracts were dried over anhydrous magnesium sulfate and evaporated to dryness. The residue was subjected to column chromatography (Silicagel 60A, 0.060-0.200 mm; eluent: dichloromethane/methanol 100:0-99:1) to give 14-hydroxy-tetradecanoic acid tert-butyl ester as yellowish oil.

Yield: 9.19 g (96%).

RF (SiO<sub>2</sub>, chloroform/methanol 9:1): 0.60.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, dH): 3.60 (t, J=6.4 Hz, 2 H); 2.18 (t, J=7.2 Hz, 2 H); 1.61-1.48 (m, 4 H); 1.42 (s, 9 H); 1.24 (bs, 18 H).

A solution of potassium hydroxide (8.60 g, 153 mmol) in water (100 mL) was added to a solution of the above ester (9.20 g, 30.6 mmol) in methanol (100 mL) and the resulting mixture was heated at 60 C for 2 days. The mixture was cooled to room temperature; then it was washed with hexanes (2 x 70 mL) and concentrated in vacuo. Concentrated hydrochloric acid (32%, 20 mL, 0.65 mol) was added dropwise and the mixture was extracted with ethyl acetate (2 x 150 mL). The combined organic layers were dried over anhydrous magnesium sulfate and evaporated to dryness to yield 14-hydroxy-tetradecanoic acid as white solid.

Yield: 7.10 g (95%).

RF (SiO<sub>2</sub>, chloroform/methanol 85:15): 0.50.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, dH): 3.66 (t, J=6.6 Hz, 2 H); 2.36 (t, J=7.4 Hz, 2 H); 1.72-1.52 (m, 4 H); 1.29 (bs, 18 H).

p-Toluenesulfonic acid monohydrate (0.28 g, 1.45 mmol) was added to a solution of the above 14-hydroxy-tetradecanoic acid (7.10 g, 29.1 mmol) in methanol (150 mL) and the resulting solution was stirred at room temperature for 3 days. Methanol was removed under reduced pressure; the residue was dissolved in ethyl acetate (170 mL); washed

with 5% aqueous solution of sodium carbonate (3 x 50 mL) and brine (25 mL); dried over anhydrous magnesium sulfate and evaporated in vacuo to give methyl 14-hydroxy-tetradecanoate as white solid.

Yield: 7.28 g (97%).

5 RF (SiO<sub>2</sub>, dichloromethane/methanol 95:5): 0.45.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, dH): 3.61-3.71 (m, 5 H); 2.32 (t, J=7.5 Hz, 2 H); 1.70-1.52 (m, 4 H); 1.44 (s, 1 H); 1.28 (bs, 18 H).

Above prepared ester (7.28 g, 28.2 mmol) was dissolved in DCM (60 mL). Triethylamine  
10 was added (5.30 mL, 52.3 mmol), reaction mixture was cooled to 0 °C and mesylchloride (2.45 mL, 31.7 mmol) was added slowly during 10 minutes. After one hour the reaction mixture was allowed to warm to room temperature and has been stirred overnight. After 16 hrs water was added (1 mL) and the mixture was stirred 30 minutes. Solvents were  
15 evaporated, ethyl acetate was added (80 mL) and the mixture was extracted with 1M hydrochloric acid (2 x 30 mL), 5% solution of sodium carbonate (2 x 20 mL) and water (20 mL). After drying with anhydrous magnesium sulfate, filtration and evaporation of solvents 16-mesylhexadecanoic acid methyl ester was obtained as white solid.

Yield: 9.15 g (92%).

RF (SiO<sub>2</sub>, dichloromethane/methanol 95:5): 0.70.

20 <sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, dH): 4.24 (t, J=6.6 Hz, 2 H); 3.68 (s, 3 H); 3.01 (s, 3 H); 2.32 (t, J=7.5 Hz, 2 H) 1.83-1.70 (m, 2 H); 1.70-1.56 (m, 2 H); 1.47-1.20 (m, 22 H).

Above prepared mesylate (9.15 g, 26.0 mmol) was dissolved in acetone (230 mL) and  
25 lithium bromide (4.50 g, 51.8 mmol) was added and the reaction mixture was refluxed overnight. After cooling down solvent was evaporated, ethyl acetate (530 mL) was added and the mixture was extracted with 5% solution of sodium hydrogencarbonate (3 x 230 mL). Combined organic extracts were dried over anhydrous magnesium sulfate and evaporated to dryness to yield 14-bromo-tetradecanoic acid methyl ester as orange oil.

30 Yield: 8.34 g (100%).

RF (SiO<sub>2</sub>, dichloromethane/methanol 95:5): 0.90.

<sup>1</sup>H NMR spectrum (300 MHz, CDCl<sub>3</sub>, dH): 3.66 (s, 3 H); 3.42 (t, J=6.9 Hz, 2 H); 2.32 (t, J=7.5 Hz, 2 H); 1.93-1.80 (m, 2 H) 1.69-1.56 (m, 2 H); 1.50-1.38 (m, 2 H); 1.28 (bs, 16 H).

35

Above prepared 14-bromo-tetradecanoic acid (8.34 g, 26.0 mmol) was dissolved in n-propanol (10 mL), water (25 mL) and 1M aqueous solution of sodium hydroxide (32 mL) and sodium sulfite (5.00 g, 39.7 mmol) was added. Reaction mixture was heated to reflux for 22 hrs. After cooling down white precipitate was filtered off. Conc. hydrochloric acid was added to give acidic pH and precipitate was centrifuged and decanted two times with water (2 x 50 ml). After lyophilization 14-sulfo-tetradecanoic acid sodium salt was obtained as a white solid.

Yield: 7.3 g (85%).

RF (SiO<sub>2</sub>, dichloromethane/methanol 95:5): 0.60.

<sup>1</sup>H NMR spectrum (300 MHz, DMSO-d<sub>6</sub>, dH): 2.40 (m, 2 H); 2.18 (t, J=7.2 Hz, 2 H); 1.61–1.40 (m, 4 H); 1.23 (bs, 18 H).

### Synthesis of compounds of the invention

#### Example 1:

SEQ ID NO:1

hPYY(1-36)

YPIKPEAPGEDASPEELNRYYYASLRHYLNLVTRQRY-NH<sub>2</sub>

#### Example 2:

SEQ ID NO:2

hPYY(3-36)

IKPEAPGEDASPEELNRYYYASLRHYLNLVTRQRY-NH<sub>2</sub>

Retention time HPLC method *UPLC16v01*: 3.37 minutes (91.4%)

Retention time HPLC method *UPLC29v01*: 10.07 minutes (85.6%)

MW calculated: 4049.6 g/mol

MALDI MS: 4048.2 g/mol

#### Example 3:

SEQ ID NO:3

[Trp30]hPYY(3-36)



Retention time HPLC method *UPLC- AP-01*: (96.6%)

MW calculated: 4123.6 g/mol


LCMS1: ((M/3)+3) 1374,8; ((M/4)+4) 1031,3

5

Example 4:

SEQ ID NO:4

[Trp30,NMeArg35]hPYY3-36



10 Retention time UPLC29v01: 3.43 minutes (100%)

Retention time UPLC16v01: 10.93 (90.9%)

MW calculated: 4136,6 g/mol

LCMS2: ((M/1)+1) 4136.03; ((M/2)+2) 2069,02 ((M/3)+3) 1379,68; ((M/4)+4) 1035,02

Example 5:

15 SEQ ID NO:5


7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36)

[illegible]

LCMS: ((M/3)+3) 1641.9; ((M/4)+4) 1231.5; ((M/5)+5) 985.6

SEQ ID NO:6

7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[2-[2-[2-[2-[2-[2-[[(4S)-4-carboxy-4-  
[[ (4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]butanoyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36)




LCMS2: ((M/3+3) 1782.6; ((M/4)+4) 1336.6; ((M/5+5) 1069.6

15 LCMS2: ((M/3+3) 1782.6; ((M/4)+4) 1336.6; ((M/5+5) 1069.6





77



Retention time UPLC02v02: 6.63 minutes (97%)

Retention time UPLC16v01: 11.94 minutes (93.2%)


5 MW calculated: 4945.67 g/mol

LCMS2: ((M/4)+4) 1237.47; ((M/3)+3) 1649.61; ((M/2)+2) 2473.87

### Example 9:

SEQ ID NO:9

7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[4-[16-(1H-tetrazol-5-yl)-  
hexadecanoylsulfamoyl]butanoylamino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-  
[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36)



15

Retention time HPLC method *UPLC30v01*: 3.78 minutes (100%)



[illegible]C(CCCC(=O)R)P[C@@H](CCCCNC(=O)COCOCOCNC(=O)COCOCNC(=O)CC[C@H](N)C(=O)O)[C@@H](C)C(=O)N[C@@H](Cc1ccc(O)cc1)C(N)=N

LCMS2: ((M/4)+4) 1240.64



Retention time UPLC02v01: 6.96 minutes (99%)

Retention time UPLC07v01: 5.04 minutes (89%)

Retention time UPLC16v01: 13.01 minutes (89.6%)


MW calculated: 4930.65 g/mol

5 LCMS2: ((M/1)+1) 4931.6; ((M/3)+3) 1644.5; ((M/4)+4) 1233.4

#### Example 15:

SEQ ID NO:15

10 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-  
[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]butanoyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]  
hPYY(4-36)



15 Retention time UPLC30v01: 3.37 minutes (91.7%)

Retention time UPLC16v01: 12.18 minutes (91.35%)

MW calculated: 5045.7 g/mol

MALDI (Found): 5045 g/mol

#### Example 16:


20 SEQ ID NO:16

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-  
(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)

[illegible]

MALDI (Found): 4915 g/mol


4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(4-36)



MALDI-MS (Found): 4929.8 g/mol

SEQ ID NO:18

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



5

Retention time UPLC16v01: 12.23 minutes (95.2%)

MW calculated: 4944.6 g/mol

Mass found 4945.34


10 LCMS2: M((/4)+4) 1237.20

#### Example 19:

SEQ ID NO:19

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)

15



Retention time UPLC02v01: 7.69 minutes (96.7%)

Retention time UPLC17v01: 9.50 minutes (92.5%)

MW calculated: 4922.62 g/mol


20 LCMS (LCMS1): m/z: 1231.7.0 ((M/4)+4); 985.3 ((M/5+5); 821.3 ((M/6+6)



Example 20:

SEQ ID NO:20

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(14-sulfotetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC30v01: 3.32 minutes (90.4%)

Retention time HPLC method UPLC60 : 17.1 minutes (85.2%)


MW calculated: 4916,48g/mol

10 LCMS (LCMS1): m/z: 1229.9 ((M/4)+4); 984.33 ((M/5)+5)

Example 21:

SEQ ID NO:21

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-carboxy-hexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC31v01: 13.0 minutes (97.4%)

20 Retention time HPLC method UPLC16v01: 3.3 minutes (95.8%)


MW calculated: 4908.6 g/mol

LCMS (LCMS1): m/z: 1637.20 ((M/3)+3); 1227.91 ((M/4)+4); 982.52 ((M/5)+5)

Example 22:

SEQ ID NO:22:

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(14-carboxy-tetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



10 Retention time HPLC method *UPLC30v01*: 3.8 minutes (97%)

Retention time HPLC method *UPLC16v01*: 10.7 minutes (100%)

MW calculated: 4880.43 g/mol


LCMS (LCMS1): m/z: 1627.64 ((M/3)+3); 1220.98 ((M/4)+4); 977.19 ((M/5)+5)

Example 23:

15 SEQ ID NO:23:

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Ile28,Trp30,NMeArg35]hPYY(4-36)

86

Retention time HPLC method *UPLC2*: 6.8 minutes (100%)5 Retention time HPLC method *UPLC17*: 8.4 minutes (91.8%)


MW calculated: 4944.53 g/mol

LCMS (LCMS1): m/z: 1648.2 ((M/3)+3); 1236.2 ((M/4)+4); 988.7 ((M/5)+5)

**Example 24:**


10 SEQ ID NO:24

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)





88

Retention time HPLC method *UPLC2:6.7 minutes (100%)*

Retention time HPLC method *UPLC16v01*: 8.5 minutes (88.9%)


5 MW calculated: .5380.00g/mol

LCMS (LCMS1): m/z: 1793 ((M/3)+3); 1344.9 ((M/4)+4); 1075.8 ((M/5)+5)

Example 27:

SEQ ID NO:27

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[ (4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC30v01:3.69 minutes (93.2%)

15 Retention time HPLC method UPLC16v01: 12.21 minutes (84.6%)

MW calculated: 4956.59.g/mol


MW(found): 4957.4 g/mol

LCMS (LCMS1): m/z: 1653.23 ((M/3)+3); 1240.19 ((M/4)+4); 992.36 ((M/5)+5)

Example 28:

SEQ ID NO:28

- 5 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]-ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



10

Retention time HPLC method UPLC30v01:3.66 minutes (92.4%)

Retention time HPLC method UPLC16v01: 12.36 minutes (65.6%)

MW calculated: 5258.96 .g/mol

MW(found): 5258.96 g/mol


- 15 LCMS (LCMS1): m/z: 1753.96 ((M/3)+3); 1315.80 ((M/4)+4); 1052.80 ((M/5)+5)

Example 29:

SEQ ID NO:29

- 20 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]-ethoxy]ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]-ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)

90



Retention time HPLC method UPLC30v01: 3.68 minutes (94.5%)

Retention time HPLC method UPLC16v01: 11.89 minutes (85.8%)

MW calculated: 5561.32 .g/mol


5 MW(found): 5562.1 g/mol

LCMS (LCMS1): m/z: 1391.31 ((M/4)+4); 1113.26 ((M/5)+5)

### Example 30:

SEQ ID NO:30

10 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[(2S)-2-amino-6-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-amino]hexanoyl]]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC16v01: 11.8 minutes (93.29%)

MW calculated: 4927.55 .g/mol


MW(found): 4928.36 g/mol

LCMS (LCMS1): m/z: 1232.92 ((M/4)+4); 986.54 ((M/5+5)

### 5 Example 31:

SEQ ID NO:31

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(15-carboxypentadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36)



10

Retention time HPLC method UPLC31v01:3.21 minutes (95.8%)

Retention time HPLC method UPLC61v01: 12.36 minutes (85.9%)

MW calculated: 4894.46 .g/mol

15 MW(found): 4894.56 g/mol

LCMS (LCMS1): m/z: 1632.52 ((M/3)+3); 1224.64 ((M/4+4)

### Example 32:


SEQ ID NO:32

7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Val3,Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36)

20



92



Retention time HPLC method UPLC16v01: 12.83 minutes (88.6%)

MW calculated: 4859.55 .g/mol


5 MW(found): 4960.21 g/mol

LCMS (LCMS1): m/z: 1240-64 ((M/4)+4); 992.91 ((M/5)+5)

### Example 33:

SEQ ID NO:33

10 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,D-Asp11,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC30v01:3.82 minutes (95.3%)

Retention time HPLC method UPLC17v01: 9.07 minutes (90.5%)

MW calculated: 4944.53 .g/mol


MW(found): 4944.60 g/mol

5 LCMS (LCMS1): m/z: 1237.15 ((M/4)+4); 989.68 ((M/5)+5)

Example 34:

SEQ ID NO:34

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36)



Retention time HPLC method UPLC30v01:1.87 minutes (89.0%)

Retention time HPLC method UPLC17v01: 8.65 minutes (74.3%)

15 MW calculated: 4944.53 .g/mol

MW(found): 4943.36 g/mol


LCMS (LCMS1): m/z: 1236.84 ((M/4)+4); 989.68 ((M/5)+5)

Example 35:

SEQ ID NO:35

20 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-  
(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]-[Arg4,Lys7,D-isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36)

94



Retention time HPLC method UPLC30v01: 3.72 minutes (98.1%)

Retention time HPLC method UPLC61: 12.78 minutes (96.2%)


5 MW calculated: 4944.53 .g/mol

#### Example 36:

SEQ ID NO:36

4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(4-36)

10



Retention time UPLC- AP-01: 7.78 minutes (96%)

MW calculated: 4909.48 .g/mol


15 MW(found): 4909.8 g/mol

LCMS (LCMS1): m/z: 1637.6 ((M/3)+3); 1228.2 ((M/4)+4); 982.9 ((M/5)+5)

#### Example 37:

SEQ ID NO:37

7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30]hPYY(3-36)



5

Retention time UPLC- AP-01: 6.51 minutes (95.3%)

MW calculated: 4960.7 .g/mol

MW(found): 4960.2 g/mol

LCMS (LCMS1): m/z: 1654.46 ((M/3)+3); 1241.1 ((M/4)+4); 992.7 ((M/5)+5)

10

#### Example 38:

SEQ ID NO:38

4-N{alpha}-(3-Methylbutanoyl)-[Arg4,Gln18,Trp30,NMeArg35]hPYY(4-36)



15

Retention time UPLC16: 13.18 minutes (62.5%)

Retention time UPLC29: 3.80 minutes (92.3%)

MW calculated: 4149.6 g/mol

MW(found): 4149.9.2 g/mol

20 LCMS (LCMSv27): m/z: 1384.04 ((M/3)+3); 1038.28 ((M/4)+4); 830.83 ((M/5)+5)

### Pharmacological methods

The utility of PYY peptide derivatives or analogues thereof of the present invention as pharmaceutically active agents in the reduction of weight gain and treatment of obesity in mammals (such as humans), and for treatment of diabetes may be demonstrated by the activity of the agonists in conventional assays and in the *in vitro* and *in vivo* assays described below.

Such assays also provide a means whereby the activities of the PYY compounds of this invention can be compared with the activities of known compounds.

#### Example 39: Receptor potency of PYY compounds

The purpose of this example is to test the activity, or potency, of the PYY compounds *in vitro*. The *in vitro* potency is the measure of the activation of the human Y1, Y2, Y4 and Y5 receptor subtypes, respectively, in a whole cell assay.

The potencies of the PYY compounds of examples 3-38 were determined using the Actone functional potency assay as described below. hPYY(3-36) (Example 2, SEQ ID NO:2) was included as a reference.

#### *Actone functional potency assay*

The Neuropeptide Y (NPY) receptors are G<sub>i</sub>-coupled seven trans-membrane receptors that mainly signal through the cAMP dependent pathway by inhibiting adenylate cyclase activity which results in a decrease of cAMP production from ATP. The Actone assay is based on a modified calcium channel that has a selective binding for cAMP, resulting in cellular calcium influx, detected by a calcium responsive dye. In order to measure decreased levels of cAMP, as result of NPY receptor activation, the  $\beta$ 1/ $\beta$ 2-adrenoreceptor agonist, isoproterenol, is added to activate adenylate cyclase and increases cAMP levels in the cell. Decreased cellular calcium concentrations, reflecting a decrease of cAMP levels due to NPY receptor activation, is detected as a decrease in fluorescence from the calcium sensitive dye.

HEK-293 cells expressing the cAMP sensitive calcium channel and one of the human NPY receptors Y1, Y2, Y4 or Y5 (CodexBiosolution, Gaithersburg, MD, USA) were seeded into poly lysine coated 384 well plates at a density of 14.000 cells/well in a volume of 25  $\mu$ l in DMEM medium containing 10% Fetal calf serum (FCS), 1% Penicillin-Streptomycin, 250  $\mu$ g/ml aminoglycoside antibiotic G418 and 1  $\mu$ g/ml aminonucleoside antibiotic puromycin. The cells were incubated over night at +37 °C in a humidified milieu in 5% CO<sub>2</sub> followed by addition of 25  $\mu$ l calcium dye buffer containing: 1 vial Calcium 5 dye (Molecular Devices, Sunnyvale, CA, USA) solved in 100 ml buffer

containing 20 mM Hepes, 0.1% Ovalbumin, 0.005% Tween 20, 1.5 mM probenecid, 250  $\mu$ M PDE-inhibitor 4-(3-Butoxy-4-methoxybenzyl)imidazolidin-2-one and 8 mM  $\text{CaCl}_2$  and pH was adjusted to 7.40. Cells were incubated for 1 hour with the calcium dye buffer and then placed in a FLIPR Tetra System (Molecular Devices) where the liquid handling system added PYY compound (1000-1 nM final concentrations) and isoproterenol (0.05  $\mu$ M final concentration) simultaneously directly followed by fluorescence signal measurement (Ex540/Em590) for 360 seconds with 30 seconds intervals. All measurements were performed in duplicates and  $\text{EC}_{50}$  values were calculated by nonlinear regression analysis of sigmoidal dose response curves using the GraphPad Prism v 5.02 (Graph Pad software, La Jolla, CA, USA). The  $\text{EC}_{50}$  values are shown in table 1.

Table 1. In vitro potency

Example	Compound	Y2 EC <sub>50</sub> (nM)	Y1 EC <sub>50</sub> (nM)	Y4 EC <sub>50</sub> (nM)	Y5 EC <sub>50</sub> (nM)
2	hPYY(3-36) SEQ ID NO:2	0.80	6.6	651	8.4
3	SEQ ID NO:3	0.57	335	403	16
4	SEQ ID NO:4	0.64	>1000	>1000	201
5	SEQ ID NO:5	2.2	>1000	>1000	260
6	SEQ ID NO:6	8.5	628	>1000	280
7	SEQ ID NO:7	3.2	977	>1000	540
8	SEQ ID NO:8	1.1	>1000	>1000	345
9	SEQ ID NO:9	2.9	>1000	>1000	219
10	SEQ ID NO:10	21	>1000	>1000	>1000
11	SEQ ID NO:11	1.9	>1000	>1000	189
12	SEQ ID NO:12	3.1	>1000	999	188
13	SEQ ID NO:13	3.6	>1000	>1000	235
14	SEQ ID NO:14	1.6	>1000	>1000	246
15	SEQ ID NO:15	4.2	599	>1000	860
16	SEQ ID NO:16	1.9	>1000	>1000	389
17	SEQ ID NO:17	1.7	>1000	>1000	81
18	SEQ ID NO:18	1.9	>1000	971	311
19	SEQ ID NO:19	4.5	>1000	>1000	>1000
20	SEQ ID NO:20	7.8	>1000	>1000	>1000

21	SEQ ID NO:21	5.6	>1000	>1000	876
22	SEQ ID NO:22	5.7	>1000	>1000	>1000
23	SEQ ID NO:23	2.8	>1000	>1000	526
24	SEQ ID NO:24	5.0	>1000	>1000	803
25	SEQ ID NO:25	6.8	>1000	>1000	>1000
26	SEQ ID NO:26	9.3	>1000	>1000	>1000
27	SEQ ID NO:27	2.9	>1000	>1000	>1000
28	SEQ ID NO:28	6.5	>1000	>1000	>1000
29	SEQ ID NO:29	19	>1000	>1000	>1000
30	SEQ ID NO:30	1.7	>1000	>1000	292
31	SEQ ID NO:31	3.5	>1000	>1000	976
32	SEQ ID NO:32	1.4	>1000	>1000	167
33	SEQ ID NO:33	3.2	>1000	>1000	923
34	SEQ ID NO:34	3.2	>1000	>1000	420
35	SEQ ID NO:35	12	>1000	>1000	626
36	SEQ ID NO:36	1.5	>1000	>1000	61
37	SEQ ID NO:37	1.0	>1000	>1000	27
38	SEQ ID NO:38	1.0	>1000	>1000	>1000

The PYY compounds of the inventions all display good Y2 potency, whereas the potency on the receptors Y1, Y4 and Y5 is strongly reduced.

#### 5 Example 40: Y1, Y2, Y4 and Y5 receptor subtype binding

The purpose of this example is to test the in vitro binding of the PYY compounds to the Y1, Y2, Y4 and Y5 receptor subtypes, respectively. The receptor binding affinity is a measure of affinity of a compound for the human Y1, Y2, Y4 and Y5 receptor subtypes, respectively.

10 The in vitro binding of the PYY compounds of examples 3-38 were determined in a scintillation proximity assay (SPA) as described below. hPYY(3-36) (Example 2, SEQ ID NO:2) was included as a reference.

#### *Scintillation Proximity Assay (SPA)*

15 NPY-receptor expressing cell lines. All cells were cultured at +37 °C in a humidified atmosphere with 5% CO<sub>2</sub>. BHK-482-8 cells with inducible expression of the human Y1 receptor (P25929, NPY1R\_HUMAN, Uniprot) were cultured in Dulbecco's Modified Eagle Medium (DMEM) with 10% Fetal bovine serum (FBS), 1% Penicillin-

Streptomycin (P/S), 1 mg/ml G418 antibiotic, 1mg/ml Hygromycin B antibiotic and 1% Non-essential amino acids. 1 mM Isopropyl  $\beta$ -D-1-thiogalactopyranoside (IPTG) was added 24 hours prior to harvesting cells for induction of NPY-Y1 receptor expression.

CHO-K1 cells stably expressing the human Y2 receptor (P49146, NPY2R\_HUMAN,

- 5 Uniprot) were cultured in DMEM F-12 with 10% FBS, 1% P/S, 150  $\mu$ g/ml Hygromycin B and 10  $\mu$ g/ml Puromycin antibiotic. CHO-K1 cells stably expressing the human Y4 receptor (P50391, NPY4R\_HUMAN, Uniprot) were cultured in DMEM F-12 with 10% FBS, 1% P/S, 10  $\mu$ g/ml Puromycin. HEK-293 cells stably expressing the human Y5 receptor (Q15761, NPY5R\_HUMAN, Uniprot) were cultured in DMEM F-12 medium containing 10%  
10 FBS, 1% Penicillin-Streptomycin, 250  $\mu$ g/ml G418 and 1  $\mu$ g/ml puromycin.

Membrane preparation. Cultured cells were detached mechanically by scraping and washed in ice cold PBS (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.47 mM KH<sub>2</sub>PO<sub>4</sub> pH adjusted to 7.4) and transferred to tubes and centrifuged for 5 minutes at  
15 1000 g at +4 °C. Pellets were resuspended in ice cold homogenization buffer; Y1: 20 mM Hepes, 10mM EDTA, with 2 complete EDTA-free protease inhibitor cocktail tablets/50 ml (Roche, Mannheim, Germany) pH 7.4); Y2, Y4: 20 mM Hepes, 5mM MgCl<sub>2</sub>, 1 mg/ml Bacitracin, pH 7.1; Y5: 10 mM NaCl, 20 mM Hepes, 0.22 mM KH<sub>2</sub>PO<sub>4</sub>, 1.26 mM CaCl<sub>2</sub>, 0.81 mM MgSO<sub>4</sub>, pH 7.4 and then homogenized for 30 seconds using a tissue  
20 homogenizer at medium speed. The homogenate was centrifuged at 35000g using an ultracentrifuge for 10 minutes at +4°C and the supernatant was discarded and fresh homogenization buffer added. Homogenization of the pellet was repeated a total of three times. The final pellet was resuspended in a few millilitres of homogenization buffer and protein concentration was determined using the Bradford method and measured at 595  
25 nm in a microplate reader. Protein concentration were adjusted to 1mg/ml and transferred to cryotubes and stored at -80°C. 250 mM sucrose was added to Y5 membranes prior to freezing.

Assay. Human Y receptor SPA binding assay were performed in white 96-well  
30 plates in a total volume of 200  $\mu$ l per well. Wheat germ agglutinin coated beads containing scintillation liquid (PerkinElmer, Waltham, MA, USA) were reconstituted in binding buffer; Y1, Y2: 50 mM Hepes, 1 mM CaCl<sub>2</sub>, 5 mM MgCl<sub>2</sub>, 0.02% tween 20, 0.25% ovalbumin pH 7.4; Y4, Y5: 20 mM Hepes, 10mM NaCl, 0.22 mM KH<sub>2</sub>PO<sub>4</sub>, 1.26 mM CaCl<sub>2</sub>, 0.81mM MgSO<sub>4</sub>, 0.1% bacitracin and 0.25% ovalbumin pH 7.4 and mixed with  
35 membrane preparation to give final concentration of 1mg beads and 3  $\mu$ g of Y1 membranes/well, 3  $\mu$ g of Y2 membranes/well, 1 $\mu$ g of Y4 membranes/well or 20  $\mu$ g of Y5



membranes/well. 50000 cpm per well of radio ligand human [ $^{125}$ I]-PYY was added corresponding to a concentration of 100 pM in Y1, Y2 and Y5 binding assays. 50000 cpm per well of radio ligand human [ $^{125}$ I]-Pancreatic Polypeptide (PP) corresponding to a concentration of 100 pM was used in Y4 binding assay.

- 5 Freeze dried analogues were dissolved in 80% dimethyl sulfoxide (DMSO), 19% H<sub>2</sub>O and 1% acetic acid (CH<sub>3</sub>COOH) to stock solutions of 2000  $\mu$ M (Y1,Y4 and Y5) and 200  $\mu$ M (Y2) and serial dilutions (1:10) were performed in binding buffer to final concentrations ranging from 10000nM to 1 pM in the Y1, Y4 and Y5 assays and 1000 nM to 0.1 pM in the Y2 assay. Plates were sealed and incubated at +25 °C for 2 hours in a plate shaker set at 400 rpm and thereafter centrifuged at 1500 rpm for 10 minutes prior to reading of luminescence on a microplate scintillation and luminescence counter. Y1 SPA plates were let to stand in room temperature for 16 hours prior to reading. Displacement of radioligand was measured as reduction in luminescence and IC<sub>50</sub> values were calculated by nonlinear regression analysis of sigmoidal dose-response curves. Ki values for binding affinity were acquired by the Cheng-Prusoff equation (Ki=IC<sub>50</sub>/(1+[L]/Kd) including receptor specific Kd values (Y1=0.556 nM; Y2=0.275 nM; Y4=0.111 nM; Y5=0.345 nM), radioligand concentration and IC<sub>50</sub> values.

Table 2: Y receptor binding affinity

Example	Compound	Y2 Ki (nM)	Y1 Ki (nM)	Y4 Ki (nM)	Y5 Ki (nM)
2	hPYY(3-36) SEQ ID NO:2	0.25	45.5	83	4.5
3	SEQ ID NO:3	0.21	443	51	15
4	SEQ ID NO:4	0.7	5089	788	379
5	SEQ ID NO:5	1.3	8021	9866	252
6	SEQ ID NO:6	5.7	6388	>10000	894
7	SEQ ID NO:7	1.5	6370	7337	713
8	SEQ ID NO:8	0.87	>10000	8881	427
9	SEQ ID NO:9	3.2	>10000	9068	495
10	SEQ ID NO:10	48	>10000	>10000	5534
11	SEQ ID NO:11	1.1	>10000	6077	246
12	SEQ ID NO:12	3.3	>10000	9807	665
13	SEQ ID NO:13	1.4	1666	7984	275
14	SEQ ID NO:14	4.8	>10000	9968	803
15	SEQ ID NO:15	6.1	>10000	>10000	2027

16	SEQ ID NO:16	3.0	>10000	>10000	1152
17	SEQ ID NO:17	0.93	8853	371	166
18	SEQ ID NO:18	4.0	>10000	9584	621
19	SEQ ID NO:19	2.1	>10000	6188	760
20	SEQ ID NO:20	7.0	>10000	6730	2540
21	SEQ ID NO:21	3.3	>10000	6587	1584
22	SEQ ID NO:22	5.0	>10000	8601	2274
23	SEQ ID NO:23	3.7	>10000	3115	1224
24	SEQ ID NO:24	4.8	>10000	6276	1848
25	SEQ ID NO:25	5.6	9200	7196	1947
26	SEQ ID NO:26	6.1	>10000	8852	3052
27	SEQ ID NO:27	2.7	>10000	5411	1437
28	SEQ ID NO:28	6.5	>10000	4912	2623
29	SEQ ID NO:29	8.3	>10000	6427	2326
30	SEQ ID NO:30	1.2	>10000	6826	690
31	SEQ ID NO:31	4.0	>10000	3040	2000
32	SEQ ID NO:32	0.84	>10000	4138	305
33	SEQ ID NO:33	2.6	>10000	5182	1622
34	SEQ ID NO:34	4.3	>10000	7692	1436
35	SEQ ID NO:35	7.5	>10000	>10000	1573
36	SEQ ID NO:36	1.5	6972	321	75
37	SEQ ID NO:37	0.37	2591	177	17
38	SEQ ID NO:38	0.87	3767	403	935

The PYY compounds of the invention all display good Y2 binding while the binding affinity on the receptors Y1, Y4 and Y5 is strongly reduced.

#### 5 Example 41: Pharmacokinetic study in minipigs

The purpose of this study is to determine the half-life in vivo of the PYY compounds after i.v. administration to minipigs, i.e. the prolongation of their time in the body and thereby their time of action. This is done in a pharmacokinetic (PK) study, where the terminal half-life of the derivative in question is determined. By terminal half-life is generally meant the period of time it takes to halve a certain plasma concentration, measured after the initial distribution phase.

*In vivo studies on pharmacokinetic evaluation in Göttingen minipigs after intravenous administration.*

Animals. Göttingen minipigs female, 15-25 kg, purchased from Ellegaard Minipigs, Denmark. The animals were housed in the Animal Unit, Novo Nordisk A/S and were kept and handled according to normal procedure in the Animal Unit. After minimum 2 weeks of acclimatization two permanent central venous catheters were implemented in vena cava caudalis in each animal. After surgery the animals were in their normal individual pens during the pharmacokinetic experiments.

Body Weight. The animals were weighed weekly. The animals were fasted on the morning prior to dosing but had ad libitum access to water; food was supplied during dosing.

Administration of peptides and dosing solutions. Intravenous injections were given through the central short catheter, which was flushed with minimum 10 ml of sterile saline post administration. The test substance was dosed at 15 nmol/kg,  $n = 3$ , in a volume of 0.05 ml/kg. Buffer: 50 mM sodium phosphate, 70 mM sodium chloride, 0.05% tween 80, pH 7.4 or 20 mM HEPES, 2.2% glycerol, 0.05% Polysorbate 80, pH 6.5.

Blood samples and analysis. Blood samples were taken through the central catheter according to the following schedule: Predose, 5, 15, 30, 45 min, 1 h, 1.5 h, 2 h, 3 h, 4 h, 6 h, 8 h, 10 h, 24 h, 48 h, 72 h, 96 h, 120 h, 168 h, 192 h, 216 h, 240 h, 264 h and 288 h. On day 1 the catheters are coupled to extension tubes, which will be removed at the end of day 1. Samples (0.8 ml) were taken through the catheter. Blood was collected in test tubes containing EDTA buffer (8 mM) and 50  $\mu$ l Val-Pyr buffer (Stabilization buffer containing 3.097g K3EDTA dissolved in 50 ml Trasylol and 0.5 ml 20 mM Val-Pyr was added. The pH was regulated to 7.4). After each blood sample the catheter was flushed with minimum 5 ml of sterile 0.9 % NaCl and 10 IE/ml heparin. Aseptic technique was demanded to avoid bacterial growth in the catheter that increases the risk of clot formation in the catheter. Samples were kept on wet ice until centrifugation (10 min, 4 °C, 1942g). Afterwards, plasma (min. 200  $\mu$ l) was transferred immediately to Micronic tubes and kept at -20 °C until analysis. The plasma samples were analysed by LC/MS as described below.

Data and results. Plasma concentration-time profiles was analysed by a non-compartmental pharmacokinetics analysis using Phoenix (Pharsight Inc., Mountain View, CA, USA). Calculations were performed using individual concentration-time values from each animal.

#### *Sample analysis*

Quantitative Assay for Plasma Samples. The test substances were assayed in plasma by Turbulent Flow Chromatography coupled to Liquid Chromatography with

subsequent Mass Spectrometric Detection (TFC/LC/MS). The selectivity of the method allowed various compounds to be quantitated in one sample, e.g. cassette dosing of four compounds per animal. The concentrations of the test substance in unknown samples were calculated using the peak area as a function of amount. Calibration graphs based on plasma samples spiked with the analyte were constructed by regression analysis. Typical dynamic range for the assay was 1 – 2,000 nmol/l. The method performance was assured by co-assaying quality control (QC) samples in duplicate at three concentration levels. Stock and working solutions of analytes were prepared in plasma and incubated by 37 °C for 1 hour.

Sample Preparation. 40.0 µl EDTA-plasma was added 160 µl 50% methanol, 1% formic acid, then vortexed and centrifuged at 14300 rpm (16457 g) at 4 °C for 20 minutes. The supernatant was transferred to a 96 well plate, (the plates have been preincubated with 0.4% BSA, 37 °C for ½ hour). Injection volume was 25 µl.

For sample clean up a TurboFlow Cyclone column (0.5 x 50 mm) both from Thermo Scientific, Franklin, MA, USA, was used and the LC separation was done either on an Onyx C18 column (2.0 x 50 mm) from Phenomenex, Torrance, CA, USA. Eluents were isocratic and gradient combinations of methanol, acetonitril, Milli-Q water and formic acid. Selective detection was done by mass spectrometry operated in positive mode ionisation.

Data handling. Plasma concentration-time profiles was analysed by a non-compartmental pharmacokinetics analysis using Phoenix (Pharsight Inc., Mountain View, CA, USA). Calculations were performed using individual concentration-time values from each animal.

Table 3: Half-life ( $t_{1/2}$ )

Example	Compound	$t_{1/2}$ (hours)
2	hPYY(3-36) SEQ ID NO:2	0.33
7	SEQ ID NO:7	63
9	SEQ ID NO:9	72
11	SEQ ID NO:11	28
12	SEQ ID NO:12	65
13	SEQ ID NO:13	55
14	SEQ ID NO:14	52
15	SEQ ID NO:15	29

16	SEQ ID NO:16	50
17	SEQ ID NO:17	58
18	SEQ ID NO:18	66
19	SEQ ID NO:19	89
20	SEQ ID NO:20	14
21	SEQ ID NO:21	52
22	SEQ ID NO:22	11
25	SEQ ID NO:25	53
26	SEQ ID NO:26	52
27	SEQ ID NO:27	59
28	SEQ ID NO:28	50
30	SEQ ID NO:30	73
31	SEQ ID NO:31	30
33	SEQ ID NO:33	58
34	SEQ ID NO:34	62
35	SEQ ID NO:35	61
36	SEQ ID NO:36	83

The tested PYY compounds of the invention have very long half-lives as compared to the half-life of hPYY(3-36).

#### 5 Example 42: Pharmacodynamic studies in db/db mice

In order to determine the in vivo effects of the PYY compounds on blood glucose and food intake in a diabetic setting, the compounds were tested in an obese, diabetic mouse model (db/db mice) as described below.

Male db/db mice are housed in a normal daily rhythm (6 pm to 6 am dark cycle) and provided ad libitum access to Altromin diet. At 11-13 weeks of age the mice are matched for blood glucose as well as body weight and divided into matching groups of 9 mice and housed 3 per cage. Mice are dosed subcutaneously with the indicated compound or vehicle (50 mM Na<sub>2</sub>HPO<sub>4</sub>, pH 7.4, 70 mM NaCl, 0.05% Tween 80) at a volume of 2.5 ml/kg at the indicated doses at 4 pm (time =0) and in some experiments a second injection was given at time=23 hours. Blood glucose and food intake are measured at the indicated time points post injection, e.g at 4 hours (4 h), 16 hours (16 h), 23 hours (23 h) and 40 hours (40 h) post injection. Blood samples for blood glucose are taken from the tail vein, into a 5 µl heparin coated capillary tube which is placed in

an eppendorf tube with Biosen<sup>®</sup> system solution (250 µl). The samples are analysed on a Biosen<sup>®</sup> instrument immediately.

Blood glucose (BG) measurements are reported as mean ± SEM of vehicle adjusted % BG relative to pre-treatment and calculated as follows:

- 5        100-[%BG(vehicle,average)-%BG] where,  
           %BG=100\*[BG(time=t)/BG(pre-treatment)]  
           and %BG(vehicle,average)=average of %BG values for the vehicle group at  
           time=t relative to vehicle pre-treatment.

- 10        Food intake is reported as mean ± SEM food intake per cage as a percentage of  
           average food intake of the vehicle group for the indicated interval.

Table 4: Effect on blood glucose in db/db mice. Blood glucose (BG) measurements are reported as mean ± SEM of vehicle adjusted % BG relative to pre-treatment

Example	Compound	Dose (µmol/kg)	% relative change in blood glucose			
			4 h	16 h	23 h	40 h
7	SEQ ID NO:7	0.3	77	88	83	86
8	SEQ ID NO:8	1	nd	68	55	nd
9	SEQ ID NO:9	1	73	70	66	66
11	SEQ ID NO:11	1	nd	62	48	nd
12	SEQ ID NO:12	1	70	68	73	67
14	SEQ ID NO:14	1	47	47	35	37
15	SEQ ID NO:15	1	nd	78	68	nd
16	SEQ ID NO:16	0.3	67	59	70	68
16	SEQ ID NO:16	1	64	47	44	40
17	SEQ ID NO:17	1	nd	58	42	nd
18	SEQ ID NO:18	0.3	66	53	56	64
18	SEQ ID NO:18	1	67	56	49	62
19	SEQ ID NO:19	1	nd	47	30	nd
20	SEQ ID NO:20	1	58	86	83	nd

Table 5: Effect on food intake in db/db mice. Food intake is reported as mean  $\pm$  SEM food intake per cage as a percentage of average food intake of the vehicle group for the indicated interval.

Example	Compound	Dose ( $\mu$ mol/kg)	% relative food intake			
			4 h	16 h	23 h	40 h
7	SEQ ID NO:7	0.3	27	34	45	54
8	SEQ ID NO:8	1	nd	33	24	nd
9	SEQ ID NO:9	1	16	14	16	16
11	SEQ ID NO:11	1	nd	24	20	nd
12	SEQ ID NO:12	1	20	23	22	22
14	SEQ ID NO:14	1	11	14	13	15
15	SEQ ID NO:15	1	nd	13	26	nd
16	SEQ ID NO:16	0.3	6.5	12	19	27
16	SEQ ID NO:16	1	6.5	8.0	7.1	12
17	SEQ ID NO:17	1	nd	18	23	nd
18	SEQ ID NO:18	0.3	10	17	24	32
18	SEQ ID NO:18	1	12	12	20	21
19	SEQ ID NO:19	1	nd	7.6	6.7	nd
20	SEQ ID NO:20	1	0	56	61	nd

- 5 These data strongly support the blood glucose lowering effect and the inhibition of food intake of the PYY compounds of the invention.

#### Example 43: Physical stability

10 The aim of this study is to determine the stability of peptide formulations in the presence and absence of phenol. As a measure of stability of the peptide formulation, the formation of high molecular weight peptide formation (%HMWP) as a function of time was analysed by size-exclusion HPLC (SE-HPLC).

15 Formulations: The analogues were solubilised to 1 mM in 30 mM Na-phosphate buffer, 10 mM NaCl pH 8.2. If necessary the pH of the stock solution was adjusted with NaOH. These stock solutions were further diluted in in the above mentioned buffer or in a corresponding buffer containing phenol, giving the final formulations with the compositions as shown in table 6.

20 Incubation: The samples were divided into UPLC vials with tight screw caps and stored incubated at 37°C and 5°C, samples at 37°C were moved to 5°C or analysed after 1, 2 and 4 weeks. The 5°C sample is used as start value.

Analysis method: The HMWP content was analysed using a dissociating SE-HPLC method with a Waters Insulin HMWP SEC column (7.8x300 mm) with a column temperature of 50°C. The samples were eluted by isocratic elution with an eluent of 500 mM NaCl, 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 5 mM H<sub>3</sub>PO<sub>4</sub>, 50% (v/v) 2-propanol at a flow rate of 0.5 ml/min. Detection was carried out at 215 nm.

As can be seen from table 6, adding a N-terminal substituent such as 3-methylbutanoyl reduces the formation of aggregates (high molecular weight proteins – HMWP). Changing lysine in position 4 to an arginine also reduces the formation of aggregates. Combining the addition of a N-terminal substituent and the change of lysine in position 4 to an arginine reduces the formation of aggregates remarkably.

Table 6. HMWP formation

Formulation No.	Compound (0.2 mM) of Example	Phosphate (mM)	NaCl (mM)	Phenol (mM)	pH	HMWP formation (%/month)
1A	18	30	10	-	8.2	0.5
1B	18	27	9	58	8.2	1.6
2A	16	30	10	-	8.2	1.0
2B	16	27	9	58	8.2	13.0
3A	8	29	10	-	8.2	1.9
3B	8	26	8	55	8.2	19.1
4A	11	29	10	-	8.2	0.7
4B	11	26	8	55	8.2	7.5

Example 44: Pharmacodynamic effect of combination treatment with a PYY compound and a GLP-1 agonist in female Landrace Yorkshire Duroc (LYD) pigs

In order to determine the in vivo effects of the combination treatment with a PYY compound and a GLP-1 agonist on food intake in pigs, the effect of either the GLP-1 agonist liraglutide alone or a combination of liraglutide and the PYY compound of Example 18 (SEQ ID NO:18) were measured as described below.

**Materials and methods.**

WO 98/08871 discloses liraglutide (example 37), including a method of preparation liraglutide.



Effect on food intake was studied in female Landrace Yorkshire Duroc (LYD) pigs (Askelygaard, Roskilde, Denmark), approximately 3 months of age, weighing 30-43 kg (n=4). The animals were acclimatised for 1-2 weeks before the study and were fed *ad libitum* with pig chow (Svine 5, Brogaarden, Denmark) at all times both during the acclimatisation and the experimental period. During the experimental period the animals were placed in individual pens for measurement of individual food intake using the system Mpigwin (Ellegaard Systems, Faaborg, Denmark). Any spillage of food (food waste) was collected and weighed, and the measured food intake was manually corrected for this amount (corrected food intake).

The PYY compound of example 18 (SEQ ID NO:18) was tested as a single s.c. administration of 2 different doses, 25 and 50 nmol/kg, in combination with 3 nmol/kg liraglutide. Liraglutide was dosed s.c. once daily from day -4 to day 3 (0.017 mL/kg corresponding to 2 nmol/kg the first 2 days, and then 0.025 mL/kg corresponding to 3 nmol/kg the remaining 6 days). The PYY analogue was dosed once in the morning of day 0 (dose volume was 0.025 mL/kg). The PYY analogue was formulated in the following buffer that also served as a vehicle for the PYY injections: 50 mM sodium phosphate 70 mM sodium chloride, 0.05% tween 80, pH 7.4. Liraglutide was formulated in the following buffer that also served as vehicle for the daily liraglutide injections: 8 mM phosphate, 184 mM propylenglycol, 58 mM phenol, pH 8.15.

The individual food intake in percent of mean vehicle food intake was calculated on day 0-4 and evaluated statistically with two-way ANOVA followed by Bonferroni's post test. Body weight change from day 0 to day 4 was calculated and evaluated statistically with one-way ANOVA followed by Bonferroni's post test.

Table 7. Percent food intake of vehicle on day 0-1 to day 3-4 after dosing of vehicle, liraglutide 3 nmol/kg, PYY analogue 25 nmol/kg in combination with liraglutide 3 nmol/kg and PYY analogue 50 nmol/kg in combination with liraglutide 3 nmol/kg in young, growing LYD pigs. Dosing of liraglutide was done once daily in the morning of day -4 to day 3, whereas the PYY compound of example 18 (SEQ ID NO:18) was given as a single s.c. dose in the morning of day 0. N=4. Data presented as mean. Two-way ANOVA followed by Bonferroni's post test.

<b>Day</b>	<b>Vehicle</b>	<b>Liraglutide 3 nmol/kg</b>	<b>Liraglutide 3 nmol/kg + SEQ ID NO:18 25 nmol/kg</b>	<b>Liraglutide 3 nmol/kg + SEQ ID NO:18 50 nmol/kg</b>
<b>0-1</b>	100.0	39.3	4.5	1.2
<b>1-2</b>	100.0	57.7	7.1	1.8
<b>2-3</b>	100.0	54.8	22.0	7.4
<b>3-4</b>	100.0	53.0	28.9	21.5

Table 8. Body weight change from day 0 to day 4 after dosing of vehicle, liraglutide 3 nmol/kg, the PYY compound of example 18 (SEQ ID NO:18) 25 nmol/kg in combination with liraglutide 3 nmol/kg and the PYY compound of example 18 (SEQ ID NO:18) 50 nmol/kg in combination with liraglutide 3 nmol/kg in young, growing LYD pigs. Dosing of liraglutide was done once daily in the morning of day -4 to day 3, whereas the PYY compound of example 18 (SEQ ID NO:18) was given as a single s.c. dose in the morning of day 0. N=4, data presented as mean. One-way ANOVA followed by Bonferroni's post test.

	<b>Vehicle</b>	<b>Liraglutide 3 nmol/kg</b>	<b>Liraglutide 3 nmol/kg + SEQ ID NO:18 25 nmol/kg</b>	<b>Liraglutide 3 nmol/kg + SEQ ID NO:18 50 nmol/kg</b>
<b>Delta BW day 0-4 (kg)</b>	4.5	3.1	0.4	-2.3

In conclusion, these data strongly support the added benefits on inhibition of food intake and body weight lowering effect of combining PYY analogues with a GLP-1 analogue.

Example 45. Pharmacodynamic effect of combination treatment with a PYY analogue and a GLP-1 analogue in db/db mice

In order to determine the in vivo effects of the combination treatment with a PYY compound and a GLP-1 agonist on blood glucose and body weight, the effect of either PYY compound, liraglutide alone or a combination were measured in an obese, diabetic mouse model (db/db mice) as described below.

Male db/db mice (9-10 weeks old) were used to measure the effect on blood glucose and body weight following treatment with varying doses of a PYY-derived analogue, the PYY compound of example 18 (SEQ ID NO:18), liraglutide (Lira), or both (combination). Prior to study mice were randomly assigned into groups that were matched for body weight, non-fasting blood glucose and HbA1c. Differences in blood glucose among groups were studied according to the dosing-schedule outlined in Table 9. Up-titration of SEQ ID NO:18 and liraglutide was used to prevent sudden appetite loss on days -9 to 0 prior the first dose (data not shown).

Table 9. Dosing-schedule outline

<b>2 x daily treatment</b>	<b>Sample, n=</b>	<b>Dose given per injection days 1-8 (nmol/kg)</b>	<b>Dose given per injection days 9-16</b>	<b>Dose given per injection days 17-24</b>
<b>Vehicle + Vehicle</b>	9	-	-	-
<b>SEQ ID NO:18</b>	9	30	100	1000
<b>Liraglutide</b>	8	10	10	10
<b>SEQ ID NO:18 + liraglutide</b>	9	15 + 5	50 + 5	1000 + 5
<b>SEQ ID NO:18 + liraglutide</b>	8	30 + 10	100 + 10	1000 + 10

Briefly, compounds were administered subcutaneously (2.5 ml/kg) at 11:00 and 17:00 hour each day. The PYY compound of example 18 (SEQ ID NO:18) was given in 50 mM phosphate; 70 mM sodium chloride; 0.05 % polysorbate 80 buffer (pH=8.0). Liraglutide was given in 50 mM phosphate; 70 mM sodium chloride; 0.05 % polysorbate 80 buffer (pH=7.4). Blood glucose was measured at a single time point (9 am) after 3-5 days of treatment at each dose based on the glucose oxidase method, using a glucose analyzer (Biosen 5040). Body weight was measured the same day. The results are shown in table 10 below.

Table 10. The effect of varying doses of the PYY compound of example 18 (SEQ ID NO:18) or liraglutide alone, or, in combination, on blood glucose and body weight. Data presented as mean. Differences were analyzed via One-way ANOVA and Tukey's multiple comparisons test. Sample, n=9.

	<b>2 x daily treatment</b>	<b>Dose given per injection</b>	<b>Blood glucose (mmol/L)</b>	<b>Body weight (g)</b>
<b>Injection days 1-8</b>	vehicle + vehicle	Vehicle	23.1	47.6
	SEQ ID NO:18	30 nmol/kg	20.5	48.6
	liraglutide	10 nmol/kg	15.4	47.6
	SEQ ID NO:18 + liraglutide	15+5 nmol/kg	9.9	46.4
	SEQ ID NO:18 + liraglutide	30+10 nmol/kg	6.6	44.0
<b>Injection days 9-16</b>	vehicle + vehicle	Vehicle	25.3	46.8
	SEQ ID NO:18	100 nmol/kg	20.2	47.7
	liraglutide	10 nmol/kg	17.4	47.7
	SEQ ID NO:18 + liraglutide	50+5 nmol/kg	10.3	45.5
	SEQ ID NO:18 + liraglutide	100+10 nmol/kg	7.6	42.8
<b>Injection days 17-24</b>	vehicle + vehicle	Vehicle	23.4	45.8
	SEQ ID NO:18	1000 nmol/kg	13.7	45.3
	liraglutide	10 nmol/kg	19.8	48.1
	SEQ ID NO:18 + liraglutide	1000+5 nmol/kg	6.2	41.9
	SEQ ID NO:18 + liraglutide	1000+10 nmol/kg	5.8	38.8

The glucose-lowering effect of the combination of the PYY compound of example 18 (SEQ ID NO:18) and liraglutide was significantly greater than the benefit produced by either compound in isolation. Glycemic levels were normalized in mice treated with the combination in the majority of the doses that were tested. Importantly, no instances of hypoglycemia were observed. The synergistic effect of the combination of the PYY compound of example 18 (SEQ ID NO:18) and liraglutide was also evident with respect to reductions in body weight. In conclusion, a combination of the PYY compound of example 18 (SEQ ID NO:18) and liraglutide synergistically lowers resting blood glucose and body weight, greatly reducing the amount of drug necessary to achieve the same, or similar, benefit produced by monotherapy.

While certain features of the invention have been illustrated and described herein, many modifications, substitutions, changes, and equivalents will now occur to those of ordinary skill in the art. It is, therefore, to be understood that the appended


claims are intended to cover all such modifications and changes as fall within the true spirit of the invention.

**CLAIMS**


1. A PYY compound comprising:

- i) tryptophan at the position corresponding to position 30 of hPYY(1-36) (SEQ ID NO:1),
- ii) N(alpha)-methyl-L-arginine at a position corresponding to position 35 of hPYY(1-36) (SEQ ID NO:1),
- iii) a lysine at a position corresponding to position 7 of hPYY(1-36) (SEQ ID NO:1) and a modifying group attached to the epsilon amino group of said lysine, wherein said modifying group is defined by **A-B-C-**,


wherein **A-** is selected from




wherein a is an integer from 12 to 19, b is an integer from 10 to 16, and c is an integer from 10 to 16, and wherein \* denotes the attachment point to -B-,  
wherein **B-** is selected from



wherein d is 1 or 2; e is 1 or 2; and f is 2, 3 or 4; and wherein \*\*\* denotes the attachment point to A-, and \*\* denotes the attachment point to -C-,  
wherein **-C-** is






VIII


wherein g is an integer in the range of 1-5, h is an integer in the range of 1-5, i is an integer in the range of 2 to 6, j is an integer in the range of 1 to 6, and wherein \*\*\*\* denotes the attachment point to -B-, and \*\*\*\*\* denotes the attachment point to the epsilon amino group of the Lysine residue in the position corresponding to position 7 of hPYY(1-36), and

iv) a maximum of 8 amino acid modifications as compared to hPYY(3-36), or a pharmaceutically acceptable salt thereof.


2. A PYY compound according to claim 1 selected from the group consisting of:  
7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[4S]-4-carboxy-4-(17-carboxyheptadecanoylamino)-  
butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,  
Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:5)



7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[[(4S)-4-carboxy-4-  
[[ (4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]butanoyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:6)




7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[2-[2-[2-[2-[2-[2-[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:7)




7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:8)







7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[2-[2-[2-[2-[4-[16-(1H-tetrazol-5-yl)-  
hexadecanoylsulfamoyl]butanoylamino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,  
Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:9)



7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-  
4-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]butanoyl]-  
amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Aib28,Trp30,NMeArg35]  
hPYY(3-36) (SEQ ID NO:10)




7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:11)




4-N{alpha}-(hexanoyl)-7-N{Epsilon}-2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:12)


- 118 -




4-N{alpha}-(3-methyl-pentanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:13)




4-N{alpha}-(3-methyl-pentanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:14)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]butanoyl]amino]-ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:15)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:16)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:18)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[4S]-4-carboxy-4-(17-carboxyheptadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:19)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[4S]-4-carboxy-4-(14-sulfotetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:20)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-carboxy-hexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:21)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(14-carboxy-tetradecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:22)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Ile28,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:23)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:24)



4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-  
[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]-  
ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]et  
hoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:25)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-  
[[2-[2-[2-[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]-  
ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-  
[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:26)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[ (4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-




oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:27)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[4-[3-[2-[2-[3-[[4-(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]-ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:28)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[4-(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]-ethoxy]ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]-ethoxy]propylamino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:29)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[(2S)-2-amino-6-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]-amino]hexanoyl]]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:30)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(15-carboxypentadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]-ethoxy]ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:31)




7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Val3,Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(3-36) (SEQ ID NO:32)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,D-Asp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:33)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:34)




4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,D-isoAsp11,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:35)



3. The PYY compound according to claim 1 or claim 2, wherein the PYY compound is 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]butanoyl]amino]-ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-[Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:15)




4. The PYY compound according to claim 1 or claim 2, wherein the PYY compound is 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[2-[2-[2-[2-[2-[2-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)-butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]-ethoxy]acetyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:18)



5. The PYY compound according to claim 1 or claim 2, wherein the PYY compound is 4-N{alpha}-(3-Methylbutanoyl)-7-N{Epsilon}-[4-[3-[2-[2-[3-[[[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]propoxy]ethoxy]ethoxy]propylamino]-4-

oxobutanoyl]-[Arg4,Lys7,Gln18,Trp30,NMeArg35]hPYY(4-36) (SEQ ID NO:27)



6. A pharmaceutical composition comprising a PYY compound according to any one of claims 1-5, and at least one pharmaceutically acceptable excipient.
7. A PYY compound according to any one of claims 1-5, suitable for use as a medicament.
8. A method of treatment and/or prevention of diabetes, type 2 diabetes or obesity, which comprises administering to a subject in need thereof an effective amount of a PYY compound according to any one of claims 1-5, or composition of claim 6.
9. A pharmaceutical composition comprising a PYY compound according to any one of claims 1-5, a GLP-1 agonist, and at least one pharmaceutically acceptable excipient.
10. A pharmaceutical composition according to claim 9, wherein the GLP-1 agonist is liraglutide or semaglutide, and wherein the PYY compound is SEQ ID NO:18.
11. A pharmaceutical composition according to claim 9 or claim 10, suitable for use as a medicament.
12. A method of treatment and/or prevention of diabetes, type 2 diabetes or obesity, which comprises administering to a subject in need thereof an effective amount of a pharmaceutical composition according to claim 9 or claim 10.
13. Use of a PYY compound according to any one of claims 1-5 in the manufacture of a medicament for the treatment and/or prevention of diabetes, types 2 diabetes or obesity.

14. Use of a pharmaceutical composition according to any one of claims 6 or 9-10 in the manufacture of a medicament for the treatment and/or prevention of diabetes, type 2 diabetes or obesity.

## SEQUENCE LISTING

<110> Novo Nordisk A/S  
 <120> Selective PYY compounds and uses thereof  
 <130> 130009W001  
 <160> 38  
 <170> Novo Nordisk A/S PatSeq 1.0.2.2  
 <210> 1  
 <211> 36  
 <212> PRT  
 <213> HOMO SAPIENS  
 <220>  
 <221> MOD\_RES  
 <222> (36)..(36)  
 <223> AMIDATION  
 <400> 1  
 Tyr Pro Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35  
 <210> 2  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL  
 <220>  
 <223> Artificial protein based on human PYY  
 <220>  
 <221> MOD\_RES  
 <222> (1)..(2)  
 <223> These positions are deleted  
 <220>  
 <221> MOD\_RES  
 <222> (36)..(36)  
 <223> AMIDATION  
 <400> 2  
 Xaa Xaa Ile Lys Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Asn Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Leu Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35  
 <210> 3  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL



&lt;220&gt;

&lt;223&gt; Artificial protein based on human PYY

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (1)..(2)

&lt;223&gt; These positions are deleted

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (36)..(36)

&lt;223&gt; AMIDATION

&lt;400&gt; 3

Xaa	Xaa	Ile	Lys	Pro	Glu	Ala	Pro	Gly	Glu	Asp	Ala	Ser	Pro	Glu	Glu
1				5				10					15		
Leu	Asn	Arg	Tyr	Tyr	Ala	Ser	Leu	Arg	His	Tyr	Leu	Asn	Trp	Val	Thr
		20					25						30		
Arg	Gln	Arg	Tyr												
		35													

&lt;210&gt; 4

&lt;211&gt; 36

&lt;212&gt; PRT

&lt;213&gt; ARTIFICIAL

&lt;220&gt;

&lt;223&gt; Artificial protein based on human PYY

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (1)..(2)

&lt;223&gt; These positions are deleted

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (35)..(35)

&lt;223&gt; METHYLATION

&lt;220&gt;

&lt;221&gt; MOD\_RES

&lt;222&gt; (36)..(36)

&lt;223&gt; AMIDATION

&lt;400&gt; 4

Xaa	Xaa	Ile	Lys	Pro	Glu	Ala	Pro	Gly	Glu	Asp	Ala	Ser	Pro	Glu	Glu
1				5				10					15		
Leu	Asn	Arg	Tyr	Tyr	Ala	Ser	Leu	Arg	His	Tyr	Leu	Asn	Trp	Val	Thr
		20					25						30		
Arg	Gln	Arg	Tyr												
		35													

&lt;210&gt; 5

&lt;211&gt; 36

&lt;212&gt; PRT

&lt;213&gt; ARTIFICIAL

&lt;220&gt;

&lt;223&gt; Artificial protein based on human PYY

&lt;220&gt;

```
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> The epsilon amino group of Lys has the following substituent:
[2-[2-[2-[[2-[2-[2-[[ (4S)-4-carboxy-4-(17-carboxyheptadecanoyl amino)butanoyl ]amino]
ethoxy]ethoxy]acetyl ]amino]ethoxy]ethoxy]acetyl ]-

```

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```
<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMINATION
```

<400> 5  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30 35  
Arg Gln Arg Tyr

<210>	6
<211>	36
<212>	PRT
<213>	ART F C AL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

<220>  
 <221> MOD\_RES  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[2-[[2-[2-[2-[[[2-[2-[2-[[[2-(4S)-4-carboxy-4-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoyl)amino]butanoyl]amino]butanoyl]aminoethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]ethoxy]acetyl]-

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```
<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMIDATION
```

<400> 6  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu

1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30 35  
Arg Gln Arg Tyr

<210>	7
<211>	36
<212>	PRT
<213>	ARTI F I C I A L

<220>  
<223> Arti fi ci al protei n based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

<220>  
 <221> MOD\_RES  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[2-[2-[2-[[[2-(4S)-4-carboxy-4-(17-carboxyheptadecanoyl)amino]butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```

<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMI DATI ON

```

<400> 7  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

<210>	8
<211>	36
<212>	PRT
<213>	ARTI F I C I A L

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

```
<220>
<221> MOD_RES
<222> (7)..(7)
```

eol f-seql . txt

<223> The epsilon amino group of Lys has the following substituent:  
[2-[2-[2-[[2-[2-[2-[[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>  
<221> MOD\_RES  
<222> (35)..(35)  
<223> METHYLATION

<220>  
<221> MOD\_RES  
<222> (36)..(36)  
<223> AMINATION

<400> 8  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

<210> 9  
<211> 36  
<212> PRT  
<213> ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

<220>  
<221> MOD\_RES  
<222> (1)..(2)  
<223> These positions are deleted

<220>  
<221> MOD\_RES  
<222> (7)..(7)  
<223> The epsilon amino group of Lys has the following substituent:  
[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[4-[16-(1H-tetrazol-5-yl)hexadecanoyl sulfamoyl]butanoyl amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>  
<221> MOD\_RES  
<222> (35)..(35)  
<223> METHYLATION

<220>  
<221> MOD\_RES  
<222> (36)..(36)  
<223> AMINATION

<400> 9  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

eol f-seql . txt

<210>	10
<211>	36
<212>	PRT
<213>	ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> The epsilon amino group of Lys has the following substituent:
[2-[2-[2-[[2-[2-[2-[[2-[2-[2-[[(4S)-4-carboxy-4-[[[(4S)-4-carboxy-4-(17-carboxyheptadecanoyl amino)butanoyl] amino]butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

```

```
<220>
<221> MOD_RES
<222> (28)..(28)
<223> The amino acid in this position is Aib (2-Aminoisobutyric acid)
```

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```
<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMINATION
```

<400> 10  
Xaa Xaa Ile Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gl n Arg Tyr Tyr Ala Ser Leu Arg Hi s Tyr Xaa Asn Trp Val Thr  
20 25 30 35  
Arg Gl n Arg Tyr  
35

<210>	11
<211>	36
<212>	PRT
<213>	ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These positions are deleted
```

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> The epsilon amino group of Lys has the following substituent:
[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]eth

```

eol f-seql . txt

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```
<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMINATION
```

<400> 11  
Xaa Xaa Ile Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

<210>	12
<211>	36
<212>	PRT
<213>	ARTICLE CIAL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(3)
<223> These positions are deleted
```

```
<220>
<221> MOD_RES
<222> (4)..(4)
<223> The alpha amino group of Arg has the following substituent: hexanoyl
```

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> The epsilon amino group of Lys has the following substituent:
2-[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoylamino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

```

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```

<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMIDATION

```

<400> 12  
Xaa Xaa Xaa Arg Pro Gl u Lys Pro Gl y Gl u Asp Al a Ser Pro Gl u Gl u  
1 5 10 15  
Leu Gl n Arg Tyr Tyr Al a Ser Leu Arg Hi s Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gl n Arg Tyr  
35

<210>	13
<211>	36
<212>	PRT
<213>	ARTICLE CIAL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(3)
<223> These positions are deleted
```

```
<220>
<221> MOD_RES
<222> (4)..(4)
<223> The alpha amino group of Lys has the following substituent:
3-methyl-pentanoyl
```

```

<220>
<221> MOD_RES
<222> (7)..(7)
<223> The epsilon amino group of Lys has the following substituent:
[2-[2-[2-[[2-[2-[2-[[ (4S)-4-carboxy-4-(17-carboxyheptadecanoyl amino)butanoyl ]amino]
ethoxy]ethoxy]acetyl ]amino]ethoxy]ethoxy]acetyl ]-

```

```
<220>
<221> MOD_RES
<222> (35)..(35)
<223> METHYLATION
```

```
<220>
<221> MOD_RES
<222> (36)..(36)
<223> AMIDATION
```

<400> 13  
Xaa Xaa Xaa Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gl n Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gl n Arg Tyr  
35

<210>	14
<211>	36
<212>	PRT
<213>	ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

```
<220>
<221> MOD_RES
<222> (1)..(3)
<223> These positions are deleted
```

```
<220>
<221> MOD_RES
<222> (4)..(4)
```

<223> The alpha amino group of Lys has the following substituent:  
3-methyl-pentanoyl

<220>

<221> MOD\_RES

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent:

[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> MOD\_RES

<222> (36)..(36)

<223> AMIDATION

<400> 14

Xaa	Xaa	Xaa	Lys	Pro	Glu	Lys	Pro	Gly	Glu	Asp	Ala	Ser	Pro	Glu	Glu
1				5				10						15	
Leu	Gln	Arg	Tyr	Tyr	Ala	Ser	Leu	Arg	His	Tyr	Leu	Asn	Trp	Val	Thr
			20					25					30		
Arg	Gln	Arg	Tyr												
			35												

<210> 15

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> These positions are deleted

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> The alpha amino group of Lys has the following substituent: 3-Methylbutanoyl

<220>

<221> MOD\_RES

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent:

[2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> MOD\_RES

<222> (36)..(36)

<223> AMIDATION



eof-seql.txt

<400> 15  
Xaa Xaa Xaa Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

<210> 16  
<211> 36  
<212> PRT  
<213> ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

<220>  
<221> MOD\_RES  
<222> (1)..(3)  
<223> These positions are deleted

<220>  
<221> MOD\_RES  
<222> (4)..(4)  
<223> The alpha amino group of Lys has the following substituent: 3-Methylbutanoyl

<220>  
<221> MOD\_RES  
<222> (7)..(7)  
<223> The epsilon amino group of Lys has the following substituent: [2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

<220>  
<221> MOD\_RES  
<222> (35)..(35)  
<223> METHYLATION

<220>  
<221> MOD\_RES  
<222> (36)..(36)  
<223> AMIDATION

<400> 16  
Xaa Xaa Xaa Lys Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
1 5 10 15  
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30  
Arg Gln Arg Tyr  
35

<210> 17  
<211> 36  
<212> PRT  
<213> ARTIFICIAL

<220>  
<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> MOD\_RES  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>  
 <221> MOD\_RES  
 <222> (36).. (36)  
 <223> AMIDATION

<400> 17  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 18  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> MOD\_RES  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>  
 <221> MOD\_RES  
 <222> (35).. (35)  
 <223> METHYLATION

<220>  
 <221> MOD\_RES  
 <222> (36).. (36)

<223> AMIDATION

<400> 18

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 19

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> These positions are deleted

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>

<221> MOD\_RES

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent: [2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(17-carboxyheptadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> MOD\_RES

<222> (36)..(36)

<223> AMIDATION

<400> 19

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 20

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> MOD\_RES  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[[ (4S)-4-carboxy-4-(14-sulfotetradecanoyl amino)butanoyl ] amino]ethoxy]ethoxy]acetyl ] ] amino]ethoxy]ethoxy]acetyl ] -

<220>  
 <221> MOD\_RES  
 <222> (35).. (35)  
 <223> METHYLATION

<220>  
 <221> MOD\_RES  
 <222> (36).. (36)  
 <223> AMINATION

<400> 20  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 21  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[[ (4S)-4-carboxy-4-(16-carboxy-hexadecanoyl amino)butanoyl ] amino]ethoxy]ethoxy]acetyl ] ] amino]ethoxy]ethoxy]acetyl ] -

<220>  
 <221> MOD\_RES  
 <222> (35).. (35)

<223> METHYLATION

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> AMIDATION

<400> 21

Xaa	Xaa	Xaa	Arg	Pro	Glu	Lys	Pro	Gly	Glu	Asp	Ala	Ser	Pro	Glu	Glu
1				5					10					15	
Leu	Gln	Arg	Tyr	Tyr	Ala	Ser	Leu	Arg	His	Tyr	Leu	Asn	Trp	Val	Thr
			20					25					30		
Arg	Gln	Arg	Tyr												
			35												

<210> 22

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> These positions are deleted

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>

<221> Mod\_Res

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent:

[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(14-carboxytetradecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]]amino]ethoxy]ethoxy]acetyl]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> AMIDATION

<400> 22

Xaa	Xaa	Xaa	Arg	Pro	Glu	Lys	Pro	Gly	Glu	Asp	Ala	Ser	Pro	Glu	Glu
1				5					10					15	
Leu	Gln	Arg	Tyr	Tyr	Ala	Ser	Leu	Arg	His	Tyr	Leu	Asn	Trp	Val	Thr
			20					25					30		
Arg	Gln	Arg	Tyr												
			35												

<210> 23

<211> 36

<212> PRT

<213> ARTI F I C I A L

<220>

<223> Arti f i c i a l p r o t e i n b a s e d o n h u m a n P Y Y

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> T h e s e p o s i t i o n s a r e d e l e t e d

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> T h e a l p h a a m i n o g r o u p o f A r g h a s t h e f o l l o w i n g s u b s t i t u e n t : 3-Methyl butanoyl

<220>

<221> Mod\_Res

<222> (7)..(7)

<223> T h e e p s i l o n a m i n o g r o u p o f L y s h a s t h e f o l l o w i n g s u b s t i t u e n t :

[2-[2-[2-[2-[2-[[ (4S)-4-carboxy-4-(16-sul fo hexadecanoyl ami no) butanoyl ] ami no] ethoxy]ethoxy]acetyl ] ami no]ethoxy]ethoxy]acetyl ]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> M E T H Y L A T I O N

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> A M I D A T I O N

<400> 23

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu

1

5

10

15

Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Ile Asn Trp Val Thr

20

25

30

Arg Gln Arg Tyr

35

<210> 24

<211> 36

<212> PRT

<213> ARTI F I C I A L

<220>

<223> Arti f i c i a l p r o t e i n b a s e d o n h u m a n P Y Y

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> T h e s e p o s i t i o n s a r e d e l e t e d

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> T h e a l p h a a m i n o g r o u p o f A r g h a s t h e f o l l o w i n g s u b s t i t u e n t : 3-Methyl butanoyl

<220>

<221> Mod\_Res

<222> (7)..(7)

<223> T h e e p s i l o n a m i n o g r o u p o f L y s h a s t h e f o l l o w i n g s u b s t i t u e n t :

[2-[2-[2-[2-[2-[[ (4S)-4-carboxy-4-(16-sul fo hexadecanoyl ami no) butanoyl ] ami no] ethoxy]ethoxy]acetyl ]-

eol f-seql . txt

```
<220>
<221> Mod_Res
<222> (36)..(36)
<223> AMINATION
```

<210>	25
<211>	36
<212>	PRT
<213>	ART F C A

```
<220>
<221> MOD_RES
<222> (1)..(3)
<223> These positions are deleted
```

<220>  
<221> Mod\_Res  
<222> (7)..(7)  
<223> The epsilon amino group of Lys has the following substituent:  
[2-[2-[2-[2-[2-[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoyl)amino]butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

```
<220>
<221> Mod_Res
<222> (36)..(36)
<223> AMI DATI ON
```

Page 16

```
<220>
<221> MOD_RES
```



<222> (4)..(4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [4-[3-[2-[2-[3-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]propoxy]  
 ]ethoxy]ethoxy]propyl amino]-4-oxobutanoyl]-

<220>  
 <221> MOD\_RES  
 <222> (35)..(35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36)..(36)  
 <223> AMIDATION

<400> 27  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 28  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1)..(3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4)..(4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butano  
 yl] amino]propoxy]ethoxy]ethoxy]propyl amino]-4-oxobutanoyl] amino]propoxy]etho  
 xy]propyl amino]-4-oxobutanoyl]-

<220>  
 <221> MOD\_RES  
 <222> (35)..(35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36)..(36)

<223> AMIDATION

<400> 28

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 29

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> These positions are deleted

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>

<221> Mod\_Res

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent:

[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[4-[3-[2-[2-[3-[[ (4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]propoxy]ethoxy]ethoxy]propyl amino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propyl amino]-4-oxobutanoyl]amino]propoxy]ethoxy]ethoxy]propyl amino]-4-oxobutanoyl]-

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> AMIDATION

<400> 29

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 30

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1)..(3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4)..(4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [(2S)-2-amino-6-[[2-[2-[2-[(4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]hexanoyl]]-

<220>  
 <221> MOD\_RES  
 <222> (35)..(35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36)..(36)  
 <223> AMIDATION

<400> 30  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 31  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1)..(3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4)..(4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[(4S)-4-carboxy-4-(15-carboxypentadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]]-

<220>  
 <221> MOD\_RES

<222> (35).. (35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36).. (36)  
 <223> AMIDATION

<400> 31  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 32  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (2)  
 <223> These positions are deleted

<220>  
 <221> Mod\_Res  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[[ (4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl] amino]ethoxy]ethoxy]acetyl] amino]ethoxy]ethoxy]acetyl]-

<220>  
 <221> MOD\_RES  
 <222> (35).. (35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36).. (36)  
 <223> AMIDATION

<400> 32  
 Xaa Xaa Val Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 33  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

<220>  
 <221> MOD\_RES  
 <222> (11).. (11)  
 <223> D-Asp

<220>  
 <221> MOD\_RES  
 <222> (35).. (35)  
 <223> METHYLATION

<220>  
 <221> Mod\_Res  
 <222> (36).. (36)  
 <223> AMINATION

<400> 33  
 Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Xaa Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 34  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1).. (3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4).. (4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> Mod\_Res  
 <222> (7).. (7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[[2-[2-[2-[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

oxy]ethoxy]acetyl ]ami no]ethoxy]ethoxy]acetyl ]-

<220>

<221> Mod\_Res

<222> (11)..(11)

<223> IsoAsp

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> AMINATION

<400> 34

Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Xaa Ala Ser Pro Glu Glu  
1 5 10 15

Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
20 25 30

Arg Gln Arg Tyr  
35

<210> 35

<211> 36

<212> PRT

<213> ARTIFICIAL

<220>

<223> Artificial protein based on human PYY

<220>

<221> MOD\_RES

<222> (1)..(3)

<223> These positions are deleted

<220>

<221> MOD\_RES

<222> (4)..(4)

<223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>

<221> Mod\_Res

<222> (7)..(7)

<223> The epsilon amino group of Lys has the following substituent:

[2-[2-[2-[[2-[2-[2-[[[4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl ]ami no]ethoxy]ethoxy]acetyl ]ami no]ethoxy]ethoxy]acetyl ]-

<220>

<221> Mod\_Res

<222> (11)..(11)

<223> D-IsoAsp

<220>

<221> MOD\_RES

<222> (35)..(35)

<223> METHYLATION

<220>

<221> Mod\_Res

<222> (36)..(36)

<223> AMINATION

eol f-seql . txt

```
<400> 35
Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Xaa Ala Ser Pro Glu Glu
 1      5      10      15
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr
      20      25      30
Arg Gln Arg Tyr
      35
```

```
<210> 36
<211> 36
<212> PRT
<213> ARTI F I C I A L
```

```
<220>
<223> Arti f i c i a l protein based on human PYY
```

```
<220>
<221> MOD_RES
<222> (1)..(3)
<223> These posi ti ons are deleted
```

```
<220>
<221> MOD_RES
<222> (4)..(4)
<223> The alpha ami no group of Arg has the fol low ing substi tuent: 3-Methyl butanoyl
```

```
<220>
<221> Mod_Res
<222> (7)..(7)
<223> The epsilon ami no group of Lys has the fol low ing substi tuent:
[2-[2-[2-[2-[2-[2-[[ (4S)-4-carboxy-4-(17-carboxyheptadecanoyl ami no)butanoyl ]ami no]
ethoxy]ethoxy]acetyl ]ami no]ethoxy]ethoxy]acetyl ]-
<220>
<221> Mod_Res
<222> (36)..(36)
<223> AMI DATION
```

```
<400> 36
Xaa Xaa Xaa Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu
 1      5      10      15
Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr
      20      25      30
Arg Gln Arg Tyr
      35
```

```
<210> 37
<211> 36
<212> PRT
<213> ARTI F I C I A L
```

```
<220>
<223> Arti f i c i a l protein based on human PYY
```

```
<220>
<221> MOD_RES
<222> (1)..(2)
<223> These posi ti ons are deleted
```

```
<220>
<221> Mod_Res
```

eof-seql.txt

<222> (7)..(7)  
 <223> The epsilon amino group of Lys has the following substituent:  
 [2-[2-[2-[2-[2-[2-[[(4S)-4-carboxy-4-(16-sulfohexadecanoyl amino)butanoyl]amino]ethoxy]ethoxy]acetyl]amino]ethoxy]ethoxy]acetyl]-

<220>  
 <221> Mod\_Res  
 <222> (36)..(36)  
 <223> AMIDATION

<400> 37  
 Xaa Xaa Ile Arg Pro Glu Lys Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35

<210> 38  
 <211> 36  
 <212> PRT  
 <213> ARTIFICIAL

<220>  
 <223> Artificial protein based on human PYY

<220>  
 <221> MOD\_RES  
 <222> (1)..(3)  
 <223> These positions are deleted

<220>  
 <221> MOD\_RES  
 <222> (4)..(4)  
 <223> The alpha amino group of Arg has the following substituent: 3-Methylbutanoyl

<220>  
 <221> MOD\_RES  
 <222> (35)..(35)  
 <223> METHYLATION

<220>  
 <221> MOD\_RES  
 <222> (36)..(36)  
 <223> AMIDATION

<400> 18  
 Xaa Xaa Xaa Arg Pro Glu Ala Pro Gly Glu Asp Ala Ser Pro Glu Glu  
 1 5 10 15  
 Leu Gln Arg Tyr Tyr Ala Ser Leu Arg His Tyr Leu Asn Trp Val Thr  
 20 25 30  
 Arg Gln Arg Tyr  
 35