

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau

(10) International Publication Number

WO 2018/222922 A1

(43) International Publication Date  
06 December 2018 (06.12.2018)

## (51) International Patent Classification:

*A61K 9/00* (2006.01)      *A61K 47/22* (2006.01)  
*A61K 47/14* (2017.01)      *A61K 9/08* (2006.01)  
*A61K 47/16* (2006.01)      *A61K 31/5513* (2006.01)  
*A61K 47/20* (2006.01)

Pharmaceuticals, Inc., 180 N. LaSalle Street, Suite 1800, Chicago, Illinois 60601 (US). **SLOAT, Brian R.**; c/o Xeris Pharmaceuticals, Inc., 180 N. LaSalle Street, Suite 1800, Chicago, Illinois 60601 (US).

## (21) International Application Number:

PCT/US2018/035473

## (22) International Filing Date:

31 May 2018 (31.05.2018)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

62/514,474      02 June 2017 (02.06.2017)      US

(71) Applicant: **XERIS PHARMACEUTICALS, INC.**  
 [US/US]; 180 N. LaSalle Street, Suite 1800, Chicago, Illinois 60601 (US).

(72) Inventors: **PRESTRELSKI, Steven J.**; c/o Xeris Pharmaceuticals, Inc., 180 N. LaSalle Street, Suite 1800, Chicago, Illinois 60601 (US). **SANDOVAL, Michael A.**; c/o Xeris

(74) Agent: **LANDRUM, Charles P.**; NORTON ROSE FUL-BRIGHT US LLP, 98 San Jacinto Boulevard, Suite 1100, Austin, Texas 78701 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

## (54) Title: PRECIPITATION RESISTANT SMALL MOLECULE DRUG FORMULATIONS




FIG. 2

(57) Abstract: Embodiments of the invention are directed to formulations that provide a solution to the problem of small molecule (e.g., diazepam) precipitation at the injection site when administered as a highly-concentrated formulation. In certain aspects the formulations include at least one surfactant in a non-aqueous formulation, which prevents small molecule precipitation and improves the bioavailability of the drug by enhancing absorption into the systemic circulation.



UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

**Published:**

- *with international search report (Art. 21(3))*

**PRECIPITATION RESISTANT SMALL MOLECULE DRUG**

**FORMULATIONS**

**CROSS REFERENCE TO RELATED APPLICATIONS**

**[0001]** This Application claims priority to U.S. Provisional Patent Application serial number 62/514,474 filed June 2, 2017, which is incorporated herein by reference in its entirety.

**BACKGROUND OF THE INVENTION**

**A. Field of the Invention**

**[0002]** The present invention relates generally to pharmaceutical formulations and, more particularly, to therapeutic formulations of small molecule drugs having improved solubility, stability, and bioavailability, and to methods of using such pharmaceutical formulations to treat various diseases, conditions, and disorders.

**B. Description of the Related Art**

**[0003]** While many small molecule drugs are orally bioavailable, parenteral injection is also used in situations where the drug has insufficient oral bioavailability, the patient is unable to accept drugs orally, or there is a need for more rapid onset of drug action. For example, administration of benzodiazepines for emergency treatment of epileptic seizures, catecholamines for allergic reactions, and “triptans” for the treatment of migraine headaches represent situations where oral administration is not as efficient or advisable and thus, the drugs must be administered via a non-oral route, often parenteral administration.

**[0004]** Standard practice for preparing formulations containing small molecule drugs has been to develop aqueous solutions for parenteral injection. The primary reason is that the majority of the human body is composed of water, including blood plasma which is an aqueous environment. Accordingly, there is a natural tendency to administer a drug formulation that is compatible with the environment that the drug is intended to reach. Several small molecule drugs, however, have limited solubility and poor stability in such aqueous environments. This problem has been solved, at least in

part, by including co-solvents and/or stabilizers into the formulation to more efficiently solubilize and stabilize the small molecule drug in an aqueous solution.

**[0005]** An example of some of the difficulties associated with parenteral injection of small molecule drugs can be seen with diazepam. This drug, which is used for emergency treatment of epileptic seizures, has been hampered by its poor aqueous solubility. Thus, the currently available emergency treatment consists of a rectal gel. An attempt has also been made to develop a large-volume (up to 3 mL) intramuscular injection based on an aqueous formulation with co-solvents (larger volumes are needed due to lower solubility of diazepam). However, the development of this drug has been limited by the difficulty in delivering a deep, large volume intramuscular injection to a convulsing patient, as well as the pain associated with such a large dosage volume.

**[0006]** Further, due to the stability issues of small molecule drugs in aqueous environments, current products are oftentimes sold as lyophilized powders that require reconstitution in an aqueous carrier prior to injection. This allows for longer shelf-life of the drug. Some products are even sold as liquids that require further dilution prior to injection with sterile water, phosphate buffer solution, or isotonic saline.

**[0007]** Due to the limited aqueous solubility of therapeutically relevant benzodiazepine drugs (*e.g.*, diazepam) there have been previous attempts to formulate these drugs as non-aqueous compositions. For example, U.S. Patent 8,946,208 (Castile et al.) describes compositions for intranasal administration wherein a small molecule drug (*e.g.*, diazepam) is dissolved in a non-aqueous vehicle comprising propylene glycol and one additional non-aqueous solvent selected from a group including N-methylpyrrolidone (NMP) and dimethylsulfoxide (DMSO). The pharmaceutical compositions for intranasal administration described by Castile et al. can be highly concentrated, with examples citing diazepam concentrations ranging up to 200 mg/mL and preferred embodiments directed to 50 mg/mL concentrations. The nasal spray formulations described by Castile et al. do not address the problem of poor water solubility encountered by the drug when it is administered to a patient, and

specifically the problem of low bioavailability that occurs when such a formulation is administered to the patient as a highly concentrated injectable formulation.

**[0008]** Another example is the non-aqueous small molecule compositions for parenteral injection as described in U.S. Patent No. 9,125,805 (Patent '805). Patent '805 disclosed compositions include concentrated diazepam formulations (*e.g.*, 50 and 100 mg/mL solutions) solubilized in non-aqueous solvents, including DMSO and NMP, with examples describing excellent long-term stability under accelerated storage conditions (40°C / 75% relative humidity). However, while these formulations may exhibit excellent long-term stability when stored in a pharmaceutically relevant container-closure system (*e.g.*, a vial, a pre-filled syringe), they exhibit poor bioavailability when injected into a patient due to the extremely low water solubility of diazepam at physiological pH, resulting in precipitation of the drug at the injection site.

**[0009]** The technology described in this application builds upon the previous discoveries by Xeris Pharmaceuticals (*e.g.*, Patent '805) that solved the problem of poor water solubility and stability of many small molecule drugs, enabling the development of injectable solutions using aprotic polar solvents. However, while these formulations possessed excellent solubility of poorly water-soluble drugs such as diazepam ( $\geq$  50 mg/mL), coupled with excellent long-term storage stability in vials and pre-filled syringes, the bioavailability of the active pharmaceutical ingredient (API) following injection was minimal due to the precipitation of the drug at the injection site.

**[0010]** Accordingly, there remains a need for a formulation that addresses the poor aqueous solubility and stability of small molecules, including benzodiazepines, while providing for improved bioavailability of highly-concentrated formulations when parenterally administered to a patient.

## SUMMARY

**[0011]** The problem of small molecule (*e.g.*, diazepam) precipitation at the injection site when administered as a highly-concentrated formulation is solved by the

inclusion of at least one surfactant in the non-aqueous formulation, which prevents small molecule precipitation and improves the bioavailability of the drug by enhancing absorption into the systemic circulation. The present invention provides a solution to the current problems facing the use of small molecule drugs in therapeutic applications as described above. The solution is premised on solubilizing and stabilizing a small molecule drug in a non-aqueous environment and then directly injecting the solubilized drug into a patient via parenteral administration, without the need for a reconstitution and/or dilution step prior to administration. The formulation can be in liquid form. Once the formulation is prepared, it can be stored for an extended period (even in an injection device) and directly injected into a subject (e.g., human) without the reconstitution or dilution steps seen in current products. Indeed, this solution goes against the prevailing industry standard. In this regard, the inventors' solution has resulted in a more stable environment for the drug and a more efficient and effective way to provide life-saving drugs to those in need of treatment. Importantly, the inventors' discovery is widely applicable for the delivery of numerous small molecule drugs that, like diazepam, that have poor or limited stability and solubility in an aqueous environment.

**[0012]** The non-aqueous vehicles described in this application are suitable for preparing pharmaceutical compositions for the intracutaneous delivery of a wide range of drug compounds. Given the current description, a person having ordinary skill in the art can determine whether a particular aprotic polar solvent system is suitable for use in combination with a particular drug. For example, this can be done by measuring the solubility of the drug compound in the vehicle. The solubility can be tested by adding an excess of the drug to the vehicle and stirring the mixture for 24 hours at room temperature. Undissolved drug is then removed by filtration or centrifugation and the solution is assayed for dissolved drug content by an appropriate analytical method, such as high-performance liquid chromatography (HPLC).

**[0013]** Without wishing to be bound by theory, it is believed that the use of surfactants in the aprotic polar solvent formulations improve bioavailability by preventing the precipitation of the highly-concentrated small molecules (e.g., benzodiazepine) formulations when injected into a patient. Specifically, it is believed

that surfactants improve solubility by entrapping drugs such as diazepam in micelles such that upon injection into an aqueous environment the surfactant facilitates diffusion of diazepam away from the injection site allowing for rapid absorption and greater bioavailability. When injecting highly-concentrated diazepam formulation into an aqueous environment at physiological pH without the use of surfactants, the diazepam partially precipitates and is slowly resolubilized and absorbed.

**[0014]** Certain embodiments of the invention are directed to stable precipitation resistant formulations for parenteral injection. The formulations can include (a) a biocompatible non-aqueous solvent; (b) a small molecule drug, or a salt thereof, solubilized within the non-aqueous solvent; and (c) 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15% to 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30% w/w (including all values and ranges there between) of a surfactant, wherein the surfactant attenuates the precipitation of the small molecule drug when injected into a subject. In certain aspects, the surfactant is present at a molar ratio of surfactant:small molecule drug of 0.5:1 to 4:1, or 1:1 to 2:1. The surfactant can be sodium deoxycholate, polysorbate 80, polysorbate 20, dodecyl maltoside, sodium dodecyl sulfate, sodium tetradecyl sulfate, alcohol ethoxylate, alkyldimethylamine oxide, or alkyl betaine. In certain aspects, the liquid formulation comprises less than 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1% by weight moisture content. The formulation of claim 1, wherein the volume of the liquid formulation to be parenterally injected is 3 mL or less. The formulation can be included within a device for dispensing the formulation. In certain aspects, the device is a syringe, a pen injection device, an auto-injector device, an external or implantable pump, or a perfusion bag. The biocompatible non-aqueous solvent can be an aprotic polar solvent, an alkyl or aryl benzoate solvent, a lipid solvent, a protic solvent, or a mixture thereof. In certain aspects, the formulation includes an aprotic polar solvent, and wherein the aprotic polar solvent is dimethylsulfoxide (DMSO), dimethylformamide (DMF), ethyl acetate, n-methyl pyrrolidone (NMP), dimethyl acetamide (DMA), propylene carbonate, or mixtures thereof. In particular aspects, the aprotic polar solvent is DMSO, NMP, or a mixture thereof. The formulation can also include an aryl or alkyl benzoate solvent, and wherein the aryl or alkyl benzoate solvent is ethyl benzoate, benzyl benzoate, or mixtures thereof. The formulation can

include from 0.5, 5, 10, 20, 40, 80, 160, 320 mg/mL to 350, 400, 450, 500, 550, 600, 650, 700, 750 mg/mL (including all values and ranges there between) of the small molecule drug. In certain aspects, the volume of the liquid formulation to be parenterally injected is from 1  $\mu$ L to 10  $\mu$ L, 10  $\mu$ L to 100  $\mu$ L, or 100  $\mu$ L to 1 mL. In certain aspects, the small molecule drug is a benzodiazepine. In a particular aspect, the benzodiazepine is diazepam. The formulation can include 25, 50, 75, 100, 125, 150 mg/mL to 175, 200, 225, 250, 275, 300 mg/mL (including all values and ranges there between) of a benzodiazepine. The solvent can be, but is not limited to DMSO, NMP, or a mixture thereof.

**[0015]** Certain embodiments are directed to methods of administering the formulation of the invention to a subject in need thereof comprising parenterally injecting the formulation to the subject. In certain aspects injecting is by parenteral injection or intracutaneous injection. In certain aspects, the formulation is not diluted prior to administration.

**[0016]** Still other embodiments of the invention are directed to methods for treating or preventing a condition by parenterally administering to a subject in need thereof a formulation of the claimed invention having an amount of a small molecule drug effective to treat or prevent the condition.

**[0017]** Certain aspects are directed to methods for treating or preventing anxiety, muscle spasms, or seizures, the method including parenterally administering to a subject in need thereof a benzodiazepine formulation. The method can include administering a formulation in an injectable volume that is within a device for dispensing the formulation. The device can be a syringe, a pen injection device, an auto-injector device, an external or implantable pump, or a perfusion bag. In certain aspects, the liquid formulation includes 25 mg/mL to 300 mg/mL benzodiazepine. In a particular aspect, the benzodiazepine is diazepam. The volume of the liquid formulation to be parenterally injected can be from 1  $\mu$ L to 10  $\mu$ L, 10  $\mu$ L to 100  $\mu$ L, or 100  $\mu$ L to 1 mL. In certain aspects, the liquid formulation is not diluted prior to administration.

**[0018]** The term “benzodiazepines” as used herein refers to the class of chemical compounds containing a bicyclic core unit in which a benzene ring is fused with a diazepine ring. Benzodiazepines are widely used lipophilic drugs that act on the central nervous system, causing sedation, decreased anxiety, muscle relaxation, and anticonvulsant actions. Non-limiting examples of benzodiazepines include diazepam, clonazepam, lorazepam, alprazolam, midazolam, and temazepam.

**[0019]** The term “surfactants” as used herein refers to surface-active agents which may adsorb onto the surfaces and/or interfaces of a system and alter the surface or interfacial free energy and the surface or interfacial tension. Non-limiting functions of surfactants include use as wetting agents, emulsifying agents, dispersing agents, and suspending agents. Surfactants are monomeric, amphipathic molecules possessing a characteristic structure that has both a non-polar (hydrophobic) region termed a “tail” and a polar (hydrophilic) region that termed the “head.” The hydrophobic tail of the surfactant molecule is generally comprised of an unsaturated or saturated hydrocarbon chain or a heterocyclic or aromatic ring system. The hydrophilic head of the surfactant molecule is polar and depending upon the charge group present in the polar head of the surfactant, they may be classified as either non-ionic, cationic, anionic, or zwitterionic (ampholytic). If the molecule does not carry a charge group on its head, the surfactant is non-ionic. If the charge group on the polar head is positive the molecule is a cationic surfactant, while an anionic surfactant has a negative charge group on the polar head of the molecule. Zwitterionic (or ampholytic) surfactants possess both positively and negatively charged groups, and depending on the pH of the solution can exist as either cationic or anionic surfactant. Examples of anionic surfactants include, but are not limited to, carboxylate, phosphate, sulfate and sulfonate ions. Non-limiting examples of anionic surfactants sodium dodecyl sulfate (SDS), and sodium tetradecyl sulfate (STS). Non-limiting examples of cationic surfactants include alkylpyridinium chloride and alkyltrimethyl ammonium bromide. Examples of non-ionic surfactants include, but are not limited to, alcohol ethoxylate, while non-limiting examples of zwitterionic surfactants include alkyltrimethylamine oxide and alkyl betaine.

**[0020]** The term “dissolution” as used herein refers to a process by which a material(s) in a gas, solid, or liquid state becomes a solute(s), a dissolved component(s), of a solvent, forming a solution of the gas, liquid, or solid in the solvent. In certain aspects a therapeutic agent or an excipient, *e.g.*, an ionization stabilizing excipient, is present in an amount up to its solubility limited or is fully solubilized. The term “dissolve” refers to a gas, liquid, or solid becoming incorporated into a solvent to form a solution.

**[0021]** The term “excipient” as used herein refers to a natural or synthetic substance formulated alongside the active or therapeutic ingredient (an ingredient that is not the active ingredient) of a medication, included for the purpose of stabilization, bulking, or to confer a therapeutic enhancement on the active ingredient in the final dosage form, such as facilitating drug absorption, reducing viscosity, enhancing solubility, adjusting tonicity, mitigating injection site discomfort, depressing the freezing point, or enhancing stability. Excipients can also be useful in the manufacturing process, to aid in the handling of the active substance concerned such as by facilitating powder flowability or non-stick properties, in addition to aiding *in vitro* stability such as prevention of denaturation or aggregation over the expected shelf life.

**[0022]** “Small molecule drugs” in the context of the present invention are biologically active compounds (and salts thereof) that can bring about a desired, beneficial, and/or pharmacological effect on a subject. These “small molecule drugs” are organic or inorganic compounds. Therefore, the small molecule drugs in the context of the present invention are not polymeric compounds. Typically, the small molecule drugs have a molecular weight of less than approximately 1000 Daltons. Certain small molecule drugs are “moisture sensitive” in that they are increasingly unstable in the presence of water. Also, salts that can be used with the small molecule drugs are known to those skilled in the art and include salts with inorganic acids, organic acids, inorganic bases, or organic bases.

**[0023]** As used herein “inhibiting” or “reducing” or any variation of these terms includes any measurable decrease or complete inhibition to achieve a desired result.

**[0024]** As used herein “effective” or “treating” or “preventing” or any variation of these terms means adequate to accomplish a desired, expected, or intended result.

**[0025]** As used herein “chemical stability,” when referring to a therapeutic agent, refers to an acceptable percentage of degradation products produced by chemical pathways such as oxidation and/or hydrolysis and/or fragmentation and/or other chemical degradation pathways. In particular, a formulation is considered chemically stable if no more than about 20% breakdown products are formed after one year of storage at the intended storage temperature of the product (*e.g.*, room temperature); or storage of the product at 25°C / 60% relative humidity (RH) for one year; or storage of the product at 40°C / 75% relative humidity for one month, and preferably three months. In some embodiments, a chemically stable formulation has less than 20%, less than 15%, less than 10%, less than 5%, less than 4%, less than 3%, less than 2%, or less than 1% breakdown products formed after an extended period of storage at the intended storage temperature of the product.

**[0026]** As used herein “physical stability,” when referring to a therapeutic agent, refers to an acceptable percentage of insoluble precipitates and/or aggregates (*e.g.*, dimers, trimers, and larger forms) being formed. In particular, a formulation is considered physically stable if no more than about 15% aggregates are formed after one year of storage at the intended storage temperature of the product (*e.g.*, room temperature); or storage of the product at 25°C / 60% relative humidity for one year; or storage of the product at 40°C / 75% relative humidity for one month, and preferably three months. In some embodiments, a physically stable formulation has less than less than 15%, less than 10%, less than 5%, less than 4%, less than 3%, less than 2%, or less than 1% aggregates formed after an extended period of storage at the intended storage temperature of the product.

**[0027]** As used herein “stable formulation” refers to a formulation where at least about 65% of the therapeutic agents (*e.g.*, peptides or salts thereof) remain chemically and physically stable after two months of storage at room temperature. Particularly preferred formulations are those in which at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% chemically and physically stable

therapeutic agent remains under these storage conditions. Especially preferred stable formulations are those which do not exhibit degradation after sterilizing irradiation (*e.g.*, gamma, beta, or electron beam).

**[0028]** As used herein, “parenteral administration” refers to administration of a therapeutic agent to a patient via a route other than the alimentary canal - any administration that is not by way of the digestive tract.

**[0029]** As used herein, “parenteral injection” refers to the administration of therapeutic agents (*e.g.*, peptides or small molecules) *via* injection under or through one or more layers of skin or mucus membranes of an animal, such as a human. Standard parenteral injections are given into the subcutaneous, intramuscular, or intradermal region of an animal, *e.g.*, a human. These deep locations are targeted because the tissue expands more easily relative to shallow dermal sites to accommodate injection volumes required to deliver most therapeutic agents, *e.g.*, 0.1 to 3.0 cc (mL).

**[0030]** The term “intracutaneous” encompasses administration into the epidermal, dermal or subcutaneous skin layer.

**[0031]** As used herein, the term “aprotic polar solvent” refers to a polar solvent which does not contain acidic hydrogen and thus does not act as a hydrogen bond donor. Polar aprotic solvents include, but are not limited to dimethylsulfoxide (DMSO), dimethylformamide (DMF), ethyl acetate, n-methyl pyrrolidone (NMP), dimethylacetamide (DMA), and propylene carbonate.

**[0032]** As used herein, the term “aprotic polar solvent system” refers to a solution wherein the solvent is a single aprotic polar solvent (for example, neat DMSO), or a mixture of two or more aprotic polar solvents (for example, a mixture of DMSO and NMP).

**[0033]** As used herein, “residual moisture” may refer to the residual moisture in the drug powder following preparation by the manufacturer/supplier. Typical powders often have residual moisture contents ranging from up to 10% (w/w). When these

powders are dissolved in an aprotic polar solvent system, the residual moisture in the powder is incorporated into the formulation. Additionally, the aprotic polar solvents may also contain a certain level of residual moisture. For example, a freshly opened bottle of USP-grade DMSO typically contains up to 0.1% (w/w) moisture. The residual moisture is different from “added moisture,” where water is intentionally added to the formulation, for example to serve as a co-solvent, or to depress the freezing point of the aprotic polar solvent system. Moisture may also be introduced into the formulation during addition of an ionization stabilizing excipient (for example, through addition of a mineral acid from an aqueous stock solution (e.g., 1 N HCl)). The “total moisture” (% w/w, unless otherwise stated) in a formulation immediately following preparation is due to the contributions from both the residual moisture and the added moisture.

**[0034]** The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

**[0035]** The words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

**[0036]** The term “about” or “approximately” or “substantially unchanged” are defined as being close to as understood by one of ordinary skill in the art, and in one non-limiting embodiment the terms are defined to be within 10%, preferably within 5%, more preferably within 1%, and most preferably within 0.5%. Further, “substantially non-aqueous” refers to less than 5%, 4%, 3%, 2%, 1%, or less by weight or volume of water.

**[0037]** As used herein “pharmaceutically acceptable” ingredient, excipient, or component is one that is suitable for use with humans and/or animals without undue adverse side effects (such as toxicity, irritation and allergic response) commensurate with a reasonable benefit/risk ratio.

**[0038]** As used herein “pharmaceutically acceptable carrier” means a pharmaceutically acceptable solvent, suspending agent, or vehicle for delivering a drug compound of the present invention to a mammal such as a human.

**[0039]** Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the examples, while indicating specific embodiments of the invention, are given by way of illustration only. Additionally, it is contemplated that changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description. Other embodiments of the invention are discussed throughout this application. Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well and vice versa. Each embodiment described herein is understood to be embodiments of the invention that are applicable to all aspects of the invention. It is contemplated that any embodiment discussed herein can be implemented with respect to any method or composition of the invention, and vice versa. Furthermore, compositions and kits of the invention can be used to achieve methods of the invention.

#### **DESCRIPTION OF THE DRAWINGS**

**[0040]** The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of the specification embodiments presented herein.

**[0041]** FIG. 1. Illustrates the stability of diazepam dissolved in DMSO through several freeze-thaw cycles.

**[0042]** FIG. 2. Illustrates pharmacokinetic profiles of diazepam in DMSO formulations compared to the Diastat® commercial formulation.

**[0043]** FIG. 3. Illustrates pharmacokinetic profiles of diazepam and nordiazepam in a DMSO formulation containing 10% (w/v) SDS compared to Hospira's IM formulation

**[0044]** FIG 4. Diazepam plasma concentrations in mini-pigs following diazepam administration.

**[0045]** FIG. 5. Percent-diazepam remaining after 20 days storage at 75 °C.

**[0046]** FIG. 6. Percent-diazepam remaining after 30 days of storage under different storage conditions.

## DESCRIPTION

**[0047]** The following discussion is directed to various embodiments of the invention. The term “invention” is not intended to refer to any particular embodiment or otherwise limit the scope of the disclosure. Although one or more of these embodiments may be preferred, the embodiments disclosed should not be interpreted, or otherwise used, as limiting the scope of the disclosure, including the claims. In addition, one skilled in the art will understand that the following description has broad application, and the discussion of any embodiment is meant only to be exemplary of that embodiment, and not intended to intimate that the scope of the disclosure, including the claims, is limited to that embodiment.

### A. Small Molecule Drugs

**[0048]** In addition to the benzodiazepines, other non-limiting small molecule drugs that can be used in the context of the present invention include epinephrine, naloxone, naltrexone, remifentanil, ganaxolone, fenfluramine, brivaracetam, apomorphine, carbidopa, levodopa, dihydroergotamine, levothyroxine, hemin, palonosetron, sumatriptan, aprepitant, novantrone, chemotherapy small molecules (e.g., mitoxantrone), corticosteroid small molecules (e.g., methylprednisolone), immunosuppressive small molecules (e.g., azathioprine, cladribine,

cyclophosphamide, methotrexate), anti-inflammatory small molecules (e.g., salicylic acid, acetylsalicylic acid, diflunisal, choline magnesium trisalicylate, salicylate, benorylate, flufenamic acid, mefenamic acid, meclofenamic acid, triflumic acid, diclofenac, fenclofenac, alclofenac, fentiazac, ibuprofen, flurbiprofen, ketoprofen, naproxen, fenoprofen, fenbufen, suprofen, indoprofen, tiaprofenic acid, benoxaprofen, pirprofen, tolmetin, zomepirac, clopinac, indomethacin, sulindac, phenylbutazone, oxyphenbutazone, azapropazone, feprazone, piroxicam, isoxicam), small molecules used to treat neurological disorders (e.g., cimetidine, ranitidine, famotidine, nizatidine, tacrine, donepezil, metrifonate, rivastigmine, selegilene, imipramine, fluoxetine, olanzapine, sertindole, risperidone, paliperidone, aripiprazole, valproate semisodium, gabapentin, carbamazepine, topiramate, phenyloin, haloperidol), and small molecules used to treat cancer (e.g., vincristine, vinblastin, paclitaxel, docetaxel, cisplatin, irinotecan, topotecan, gemcitabine, temozolomide, imatinib, bortezomib), statins (e.g., atorvastatin, amlodipine, rosuvastatin, sitagliptin, simvastatin, fluvastatin, pitavastatin, lovastatin, pravastatin, simvastatin), and other taxane derivatives. In particular embodiments, the small molecules that can be used include those that treat tuberculosis (e.g., rifampicin), small molecule anti-fungal agents (e.g., fluconazole), small molecule anti-anxiety agents and small molecule anti-convulsant agents (e.g., lorazepam), small molecule anti-cholinergic agents (e.g., atropine), small molecule  $\beta$ -agonist drugs (e.g., albuterol sulfate), small molecule mast cell stabilizers and small molecule agents used to treat allergies (e.g., cromolyn sodium), small molecule anesthetic agents and small molecule anti-arrhythmic agents (e.g., lidocaine), small molecule antibiotic agents (e.g., tobramycin, ciprofloxacin), small molecule anti-migraine agents (e.g., sumatriptan), and small molecule anti-histamine drugs (e.g., diphenhydramine). Further, the amount of the small molecule drugs in the dosage formulations can be varied depending on current acceptable amounts, subject/patient needs, and the like.

**[0049]** With respect to the biocompatible non-aqueous solvent, examples include aprotic polar solvents, alkyl or aryl benzoate solvents, lipid solvents, protic solvents, or a mixture thereof. Non-limiting examples of aprotic polar solvents include dimethylsulfoxide (DMSO), dimethylformamide (DMF), ethyl acetate, n-

methylpyrrolidone (NMP), dimethyl acetamide (DMA), propylene carbonate, or mixtures thereof. In some instances, however, the formulations of the present invention do not have to include the aforementioned solvents (*i.e.*, others can be used). In one instance, for example, the formulations do not include non-aqueous aprotic polar solvents and/or do not include non-aqueous protic solvents (*e.g.*, polyethylene glycol (PEG), propylene glycol (PG), polyvinylpyrrolidone (PVP), methoxypropylene glycol (MPEG), glycerol, glycofurool, and mixtures thereof). As noted above, the increased solubility of the small molecule drugs can result in small dosage volumes (and, in turn, small storage devices and containers), which provides for an easier and less painful administration parenterally. Non-limiting examples of aryl or alkyl benzoate solvents include methyl benzoate, ethyl benzoate, propyl benzoate, C12-C15 alkyl benzoates, in which R is a C12-15 alkyl group, C16-17 alkyl benzoate, in which R is a C16-17 fatty alcohol group, and benzyl benzoate. A non-limiting example of a lipid is triacetin, which is the triester of glycerol and acetic acid. Non-limiting examples of protic solvents include polyethylene glycol (PEG), propylene glycol (PG), polyvinylpyrrolidone (PVP), methoxypropylene glycol (MPEG), glycerol, glycofurool, or mixtures thereof. In certain aspects, the formulation does not include a co-solvent, while in other aspects it can include a co-solvent. In one instance, the formulation can include a single / only one biocompatible non-aqueous solvent (*i.e.*, in neat or pure form). In other aspects, the formulation includes a mixture of two, three, four, or more biocompatible non-aqueous solvents. In still additional aspects, the formulation can exclude co-solvents, salts, and other ingredients that can help with or increase the solubility of the small molecule drug in the non-aqueous solvent. For instance, the formulation can consist of or consist essentially of a small molecule drug and a non-aqueous solvent (or mixture of non-aqueous solvents) and still be directly injected through parenteral administration to a subject - with consist essentially of meaning in the context of this sentence exclusion of other ingredients that could increase the solubility of the drug within the non-aqueous solvent or mixture of non-aqueous solvents - *e.g.*, a preservative can be included to further preserve the injectable formulation.

**[0050]** Further, the formulation of the present invention can be non-aqueous or substantially non-aqueous (*e.g.*, less than 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5% or less of total moisture (i.e. moisture present in the formulation when it is initially prepared) by weight. In some instances, the small molecule drug has previously been dried in the presence of a buffer prior to being solubilized in the non-aqueous solvent. As explained below, this can add to the stability of the small molecule drug. In some instances, the dried small molecule drug has a pH memory that is about equal to the pH of the small molecule drug in the presence of the aqueous buffer such that the pH of the small molecule drug that is solubilized in the biocompatible non-aqueous solvent is about equal to the pH of the small molecule drug in the presence of the buffer. The memory pH can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or more or can be a range of 1 to 3, 2 to 4, 3 to 5, 4 to 6, 5 to 7, 6 to 8, 7 to 9, 8 to 10 or 9 to 11. In certain aspects, the buffer is a non-volatile buffer (non-limiting examples of which include glycine buffers, citrate buffers, or phosphate buffers, or a mixture thereof). In other instances, the buffer can be a volatile buffer. Further, the moisture content of the small molecule drug can be less than 5%, 4%, 3%, 2%, 1%, 0.5% or less w/w.

**[0051]** In certain aspects, the formulation includes from 0.5 mg/mL to about 300 mg/mL, 10 mg/mL to 50 mg/mL, 20 mg/mL to 50 mg/mL, 5 mg/mL to 15 mg/mL, or 0.5 mg/mL to 2 mg/mL of the small molecule drug. In some instances, the amount of the small molecule drug can be as high as 400, 500, 600, 700, 800, 900, 1000, 2000, or 3000 mg/mL or more. One of the unique aspects of the present formulation is that the formulation can have a high content of the drug, yet the dosage of the formulation can be relatively low (*e.g.*, 0.1  $\mu$ L, 1  $\mu$ L, 10  $\mu$ L, 20  $\mu$ L, 50  $\mu$ L, 75  $\mu$ L, 100  $\mu$ L, 200  $\mu$ L, 300  $\mu$ L, 400  $\mu$ L, 500  $\mu$ L, 600  $\mu$ L, 700  $\mu$ L, 800  $\mu$ L, 900  $\mu$ L, 1 mL, 2 mL, or 3 mL, or more as needed (*e.g.*, 4, 5, 6, 7, 8, 9, 10 mL or more). In certain instances, the volume of the liquid formulation to be parenterally injected is 3 mL or less (*e.g.*, 3, 2.5, 2, 1.5, 1, 0.5, 0.1 mL or less) or is from 0.1  $\mu$ L to 3 mL or from 0.1  $\mu$ L to 1  $\mu$ L or from 1  $\mu$ L to 10  $\mu$ L or from 10  $\mu$ L to 1 mL or from 0.1  $\mu$ L to 2.5 mL or from 0.1  $\mu$ L to 2 mL or from 0.1  $\mu$ L to 1.5 mL or from 0.1  $\mu$ L to 1 mL or from 0.1  $\mu$ L to 0.5 mL or from 0.1  $\mu$ L to 0.1 mL. Another unique aspect of the present formulation is that it can be

contained in a container or device, be stored, and be immediately ready for parenteral injection on an as-needed basis without having to reconstitute or dilute the formulation. The device can be a syringe, a pen injection device, an auto-injector device, a device that can pump or administer the formulation (*e.g.*, automatic or non-automatic external pumps, implantable pumps, etc.) or a perfusion bag. Also contemplated for use in the formulations are additional ingredients/pharmaceutical excipients, non-limiting example of which include: antioxidants (examples include ascorbic acid, cysteine, methionine, monothioglycerol, sodium thiosulfate, sulfites, BHT, BHA, ascorbyl palmitate, propyl gallate, or vitamin E); chelating agents (examples include EDTA, EGTA, tartaric acid, glycerin, or citric acid); or preservatives (examples include alkyl alcohols, benzyl alcohol, a methyl paraben, or a propyl paraben or mixtures thereof). The formulation can be in liquid form, semi-solid form, or gel form. As discussed below, the formulation can have a desired viscosity range (in one non-limiting example, such a range could be between 0.5 to 15 cps). The formulation can be such that at least 65% of the small molecule drug within the formulation remains chemically and physically stable when the formulation is stored at room temperature for two months or at least 80% of the therapeutic agent within the formulation remains chemically and physically stable when the formulation is stored at room temperature for two months.

**[0052]** In one particular aspect of the present invention, there is disclosed a stable liquid formulation for parenteral injection comprising diazepam, or a salt thereof, at least one surfactant, and a biocompatible non-aqueous solvent, wherein the diazepam and the surfactant(s) are solubilized within the non-aqueous solvent, wherein the total water content of the formulation when prepared is less than 5% w/w, wherein the volume of the formulation to be parenterally injected is between 50  $\mu$ L to 1000  $\mu$ L or any range therein (*e.g.*, 75  $\mu$ L, 100  $\mu$ L, 150  $\mu$ L, 200  $\mu$ L, 300  $\mu$ L, 400  $\mu$ L, 500  $\mu$ L, 600  $\mu$ L, 700  $\mu$ L, 800  $\mu$ L, 900  $\mu$ L, etc.). In a further aspect the formulation can include additional excipients. As explained above, such a formulation can be comprised in a container selected from a sealed syringe, a sealed pen injection device, a sealed auto-injector device, or a pump. Also, as explained above, the diazepam can be dried in the presence of a buffer prior to being solubilized in the non-aqueous solvent. This can

provide the dried diazepam with a pH memory that is about equal to the pH of diazepam in the presence of the aqueous buffer such that the pH of the diazepam that is solubilized in the biocompatible non-aqueous solvent is about equal to the pH of the diazepam in the presence of the aqueous buffer (e.g., the aforementioned non-volatile buffers such as glycine buffers, citrate buffers, or phosphate buffers, or a mixture thereof).

**[0053]** Still further, the formulations of the present invention can include one or more other excipients in addition to the surfactant(s). In some embodiments, the other excipient is selected from sugars, starches, sugar alcohols, antioxidants, chelators, polymers, and preservatives. Examples of suitable sugars excipients include, but are not limited to trehalose, glucose, sucrose, etc. Examples of suitable starches for stabilizing excipients include, but are not limited to, hydroxyethyl starch (HES). Examples of suitable sugar alcohols (also referred to as polyols) include, but are not limited to, mannitol and sorbitol. Examples of suitable antioxidants include, but are not limited to, ascorbic acid, cysteine, methionine, monothioglycerol, sodium thiosulphate, sulfites, BHT, BHA, ascorbyl palmitate, propyl gallate, N-acetyl-L-cysteine (NAC), and Vitamin E. Examples of suitable chelators include, but are not limited to, EDTA, EDTA disodium salt (edetate disodium), tartaric acid and salts thereof, glycerin, and citric acid and salts thereof. Examples of suitable inorganic salts include sodium chloride, potassium chloride, calcium chloride, magnesium chloride, calcium sulfate, and magnesium sulfate. Examples of suitable preservatives include, but are not limited to, benzyl alcohols, methyl parabens, propyl parabens, and mixtures thereof. Additional formulation components include local anesthetics, such as lidocaine or procaine. In some embodiments, the additional stabilizing excipient is present in the formulation in an amount ranging from about 0.05% (w/v) to about 60% (w/v), from about 1% (w/v) to about 50% (w/v), from about 1% (w/v) to about 40% (w/v), from about 1% (w/v) to about 30% (w/v), from about 1% (w/v) to about 20% (w/v), from about 5% (w/v) to about 60% (w/v), from about 5% (w/v) to about 50% (w/v), from about 5% (w/v) to about 40% (w/v), from about 5% (w/v) to about 30% (w/v), from about 5% (w/v) to about 20% (w/v), from about 10% (w/v) to about 60% (w/v), from about 10% (w/v) to about 50% (w/v), from about 10% (w/v) to about 40% (w/v), from about 10% (w/v) to about 30% (w/v), from about 10% (w/v) to about 20% (w/v), from about 10% (w/v) to about 10% (w/v), from about 10% (w/v) to about 5% (w/v), and from about 1% (w/v) to about 5% (w/v).

40% (w/v), from about 10% (w/v) to about 30% (w/v), or from about 10% (w/v) to about 20% (w/v). In some embodiments, the additional stabilizing excipient is present in the formulation in an amount that is about, at most, or at least 0.1, 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, or 60% (w/v).

## B. Method of Making

**[0054]** In certain embodiments, the invention also provides methods of formulating compositions comprising a small molecule drug with one or more of: a pharmaceutically acceptable solvent; a surfactant; a carrier; a solubilizer; an emulsifier; a preservative; and/or other excipient. Such compositions may contain an effective amount of at least one small molecule drug. Thus, the use of one or more small molecule drugs in the preparation of a pharmaceutical composition of a medicament is also included. Acceptable formulation components for pharmaceutical preparations are nontoxic to recipients at the dosages and concentrations employed. Formulation components are present in concentrations that are acceptable to the site of administration. The pharmaceutical composition to be used for *in vivo* administration is typically sterile. Sterilization may be accomplished by filtration through sterile filtration membranes.

**[0055]** Once the pharmaceutical composition of the invention has been formulated, it may be stored in sterile vials as a solution. Such formulations may be stored in a ready-to-use form. The components used to formulate the pharmaceutical compositions are preferably of high purity and are substantially free of potentially harmful contaminants (e.g., at least National Food (NF) grade, generally at least analytical grade, and more typically at least pharmaceutical grade). Moreover, compositions intended for *in vivo* use are usually sterile.

**[0056]** Therapeutically effective doses will be easily determined by one of skill in the art and will depend on the severity and course of the disease, the patient's health and response to treatment, the patient's age, weight, height, sex, previous medical history and the judgment of the treating physician.

**[0057]** Other embodiments of the present invention are directed to methods of stably formulating a small molecule drugs (*e.g.*, a benzodiazepine drug) comprising the steps of: dissolving the drug (*e.g.*, benzodiazepine) and surfactant (and any other additional excipients) in DMSO – the order of dissolution may or may not be of importance.

### **C. Method of Treating**

**[0058]** In a further aspect of the present invention there is disclosed a method for treating or preventing a condition, disease, disorder, etc. comprising administering to a subject in need thereof a formulation(s) of the present invention in an amount effective to treat or prevent the condition, disease, disorder, etc. Any suitable dosage of a therapeutic agent (*e.g.*, small molecule) may be administered in the methods of the present invention. Depending on the small molecule(s) comprising the therapeutic formulation, use will be governed by the indications for which they are approved.

**[0059]** As a non-limiting example, benzodiazepines such as diazepam are indicated for short-term relief for the symptoms of anxiety disorders or for the management of anxiety disorders. Indications also include the relief of symptoms of acute alcohol withdrawal, including tremor, impeding or acute delirium tremens, hallucinosis, and acute agitation, as well as use as an adjunct for the relief of skeletal muscle spasm. Further indications include status epilepticus, and severe recurrent or convulsive seizures. Injectable diazepam may also be used in the management of selected patients with epilepsy, on stable regimens of AEDs, who may require intermittent use of diazepam to control bouts of increased seizure activity.

**[0060]** The dosage administered will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular compound, salt, or combination; the age, health, or weight of the subject; the nature and extent of symptoms; the metabolic characteristics of the drug and patient, the kind of concurrent treatment; the frequency of treatment; or the effect desired. In certain aspects epileptic seizures can be treated by administering a formulation described herein comprising an effective amount of diazepam.

**[0061]** Though well-known for their use in the treatment of seizures, benzodiazepines such as diazepam can also be used for the prevention of seizures. One reported example describing the use of a benzodiazepine as a prophylactic / preventative measure for eliminating the occurrence of seizures is the use of diazepam for the prevention of febrile seizures. Shinnar et al., in *Pediatric Epilepsy: Diagnosis and Therapy, Chapter 19, Febrile Seizures*, 2007, 3:293-301 (hereinafter “Shinnar”) teaches that diazepam can be administered at the onset of fever, but prior to the onset of seizures to prevent febrile seizures. Shinnar states: “Despite febrile status epilepticus representing only 5% of febrile seizures, it accounts for approximately one quarter of all episodes of childhood status epilepticus, and more than two thirds of status epilepticus cases in the second year of life,” (Shinnar, page 293). “Two distinct approaches to the treatment of febrile seizures have developed based on the perceived immediate and long-term risks of febrile seizures. One approach is based on the old idea that febrile seizures are harmful and may lead to the development of epilepsy; this approach is aimed at *preventing febrile seizures* by using either intermittent or chronic treatment with medications. The second approach is based on the epidemiological data that febrile seizures are benign; the only concern focusing on aborting febrile seizures to prevent status epilepticus.” (Shinnar, page 297, emphasis added). “Diazepam, given generally orally or rectally at the *time of onset of a febrile illness* has demonstrated a statistically significant, yet clinically modest, ability to *reduce the probability of a febrile seizure.*” (Shinnar, page 298, emphasis added).

**[0062]** Knudsen in *Archives of Disease in Childhood*, 1985 Vol. 60, pp. 1045 – 1049. (hereinafter “Knudsen”) teaches that non-orally administered diazepam provides effective prophylactic seizure control and reduces incidence of recurrence in high and intermediate risk children. Knudsen states: “Our recent study indicated that short term diazepam prophylaxis reduced the 18 month recurrence rate for 39% to 12% and *thus prevented two thirds of all further febrile fits.* Stratification showed a remarkably wide range of recurrence rates in untreated children, a uniformly low recurrence rate in response to diazepam prophylaxis at times of fever, and an appreciable difference in the efficacy of prophylaxis, in terms of risk reduction.” (Knudson, page 1048, emphasis added).

**[0063]** As described in the prior art, the use of non-orally administered diazepam, for example, the commercially available diazepam sold under the trade name Diastat® is administered for prevention of seizures.

#### **D. Method of Administering**

**[0064]** Also disclosed is a method of administering the formulations of the present invention by parenteral administration of the formulation to a subject in need thereof. The administration can be performed without having to reconstitute and/or dilute the formulation. Further, the administration can be performed with a syringe, a pen injection device, an auto-injector device, a pump, or a perfusion bag. Also, the formulation can be stored in said syringe, pen injection device, auto-injector device, pump, or perfusion bag, which can then be immediately used (again without having to reconstitute and/or dilute the formulation). Further, and as noted above, the amount of the formulation being administered can range from 1  $\mu$ L, 10  $\mu$ L, 20  $\mu$ L, 50  $\mu$ L, 75  $\mu$ L, 100  $\mu$ L, 200  $\mu$ L, 300  $\mu$ L, 400  $\mu$ L, 500  $\mu$ L, 600  $\mu$ L, 700  $\mu$ L, 800  $\mu$ L, 900  $\mu$ L, 1 mL, 2 mL, 3 mL, 4 mL, 5 mL, 6 mL, 7 mL, 8 mL, 9 mL, or 10 mL, or more as needed. In certain aspects, the formulations are such that the small molecule drug remains stable and solubilized (i.e., no coalescence or crystallization of the small molecule drug) and when stored at room temperature (approximately 20-25°C.) for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months.

#### **E. Examples**

**[0065]** The following examples as well as the figures are included to demonstrate certain embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples or figures represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute a mode for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

**[0066]** Some embodiments of the present disclosure will be described in greater detail by way of specific examples. The following examples are offered for

illustrative purposes, and are not intended to limit any present invention in any manner. For example, those of skill in the art will readily recognize a variety of noncritical parameters that can be changed or modified to yield essentially the same results.

#### EXAMPLE 1

**[0067]** In this example, either 1% (w/v) or 2% (w/v) sodium deoxycholate (SDC)(CAS 302-95-4) was dissolved in dimethyl sulfoxide (DMSO)(CAS 67-68-5) alone, or DMSO containing 5% (v/v) benzyl alcohol (BA)(CAS 100-51-6), followed by the dissolution of diazepam to a concentration of 100 mg/mL (CAS 439-14-5). Diazepam formulations were screened for their ability to prevent diazepam precipitation using an anti-precipitation assay coupled with reversed-phase high performance liquid chromatography (RP-HPLC).

**[0068]** The anti-precipitation assay used to screen diazepam formulations consisted of diluting diazepam into a simulated extracellular fluid (SEF), such that there was a 50x dilution of the formulation solvent. The SEF consisted of 0.7 mM magnesium chloride (CAS 7786-30-3), 1.2 mM calcium chloride (CAS 10043-52-4), 2.0 mM potassium chloride (CAS 7447-40-7), 2.0 mM monopotassium phosphate (CAS 7778-77-0), 0.5 mM sodium sulfate (CAS 7757-82-6), 104 mM sodium chloride (CAS 7440-23-5), and 28.3 mM sodium bicarbonate (CAS 144-55-8). After diluting the diazepam formulation into SEF, the sample was mixed by inverting several times, and the sample was allowed to equilibrate for 10 minutes. The precipitate was collected by centrifugation and the soluble diazepam content was measured in the supernatant by RP-HPLC. Greater solubility in the supernatant is expected to translate into better dissolution of diazepam away from the injection site following subcutaneous administration.

**[0069]** The RP-HPLC method used to determine soluble diazepam was an isocratic method with a mobile phase consisting of 20% (v/v) 3.2 mM ammonium formate (CAS 540-69-2) and 80% (v/v) methanol (CAS 67-56-1). A C18 column (Luna<sup>TM</sup>, Phenomenex, 3.9 mm I.D. x 150 mm length, 5 micron particle size) was used with a

column temperature of 40°C, a 0.6 mL/min flow rate, a 5- $\mu$ L sample injection volume, and 230-nm detection wavelength.

**[0070]** Sample formulations containing SDC significantly improved diazepam recovery in the anti-precipitation assay (Table 1). Samples without SDC had 11.4% and 9.1% diazepam recovery, respectively, with and without the addition of BA to the formulation. Adding 1% (w/v) SDC to the formulation improved diazepam recovery by a factor of 2.6 and 2% (w/v) SDC improved recovery by a factor of 4.1. An improvement in the recovery of soluble diazepam measured using the anti-precipitation assay and RP-HPLC is expected to translate into improved dissolution of diazepam from the injection site and the prevention of diazepam precipitation following subcutaneous administration. Accordingly, improved dissolution is expected to yield a more desirable pharmacokinetic profile with rapid absorption and slowed elimination.

Table 1. Diazepam formulations prepared with sodium deoxycholate in DMSO prevented diazepam precipitation in SEF.

Form.	Diazepam mg/mL	% SDC (w/v)	% DMSO (v/v)	% BA (v/v)	% Diazepam Recovered
1	100	0 %	100 %	0 %	9.1 $\pm$ 2.2
2	100	0 %	95 %	5 %	11.4 $\pm$ 1.8
3	100	1 %	95 %	5 %	29.3 $\pm$ 0.4
4	100	2 %	95 %	5 %	46.8 $\pm$ 3.6

## EXAMPLE 2

**[0071]** In this example, either 1%, 10%, 20%, or 30% (v/v) Polysorbate 80 (PS80)(CAS 9005-65-5) was dispersed into DMSO with 5% (v/v) BA, followed by dissolving diazepam at a concentration of 50, 75 or 100 mg/mL. Diazepam formulations containing PS80 were screened for their ability to prevent diazepam precipitation using an anti-precipitation assay, followed by RP-HPLC (as detailed above in Example 1).

**[0072]** Sample formulations containing PS80 significantly improve diazepam recovery in the anti-precipitation assay (Table 2). When diazepam was dissolved at

100 mg/mL into DMSO with 5% BA and increasing concentrations of PS80 (up to 30% (v/v)), the highest concentration of PS80 yielded the best diazepam recovery. 30% (v/v) PS80 improved diazepam recovery by a factor of 7.9. When diazepam was dissolved at 75 mg/mL into DMSO with 5% (v/v) BA and increasing concentrations of PS80 (up to 30% (v/v)), the highest concentration of PS80 yielded the best diazepam recovery. Thirty (30) percent PS80 improved diazepam recovery by a factor of 8.4. Similarly, when diazepam was dissolved at 50 mg/mL into DMSO and 5% (v/v) BA with 20% (v/v) PS80, diazepam recovery was improved by a factor of 8.8.

**[0073]** Data in Table 2 indicates that with a fixed concentration of diazepam (either 50, 75 or 100 mg/mL), the percent of recoverable diazepam increases with the PS80 concentration. The data indicate that there is an optimal ratio between PS80 and diazepam that maximizes diazepam solubility in SEF. In the case of PS80, a ~1:1 molar ratio of diazepam and PS80 prevents precipitation in SEF.

Table 2. Diazepam formulations prepared with PS80 in DMSO prevented diazepam precipitation in SEF.

Form.	Diazepam mg/mL	% PS80 (v/v)	% DMSO (v/v)	% BA (v/v)	% Diazepam Recovered
1	100	0 %	100 %	0 %	9.1 ± 2.2
2	100	0 %	95 %	5 %	11.4 ± 1.8
3	100	1%	94 %	5 %	5.8 ± 11.0
4	100	10%	85 %	5 %	37.6 ± 7.4
5	100	20%	75 %	5 %	47.9 ± 2.5
6	100	30%	65 %	5 %	90.3 ± 1.6
7	75	10%	85 %	5 %	60.4 ± 3.2
8	75	20%	75 %	5 %	72.5 ± 4.4
9	75	30%	65 %	5 %	96.3 ± 7.6
10	50	10%	85 %	5 %	72.9 ± 2.7
11	50	20%	75 %	5 %	100.7 ± 4.7

### EXAMPLE 3

**[0074]** In this example, either 1%, 5%, 10%, or 17.7% (w/v) of dodecyl maltoside (DM)(CAS 69227-93-6) was dissolved DMSO with 5% (v/v) BA, followed by dissolving diazepam at a concentration of 50, 75 or 100 mg/mL. Diazepam

formulations containing DM were screened for their ability to prevent diazepam precipitation using an anti-precipitation assay and RP-HPLC (as detailed above in Example 1).

**[0075]** Shown in Table 3 are the percent-diazepam recovery data for formulations prepared with DM. For each concentration of diazepam (50, 75 or 100 mg/mL), increasing concentrations of DM increased the percent of diazepam that remained soluble when the formulation was diluted into SEF. For example, diazepam at 50 mg/mL dissolved into DMSO with 5% BA and 17.7% (w/v) DM had nearly 100% diazepam solubility in SEF. Like PS80, the data in Table 3 suggest that there is an ideal ratio that prevents diazepam precipitation, and in this example the ratio of DM to diazepam is ~2:1.

Table 3. Diazepam formulations prepared with DM in DMSO prevented diazepam precipitation in SEF.

Form.	Diazepam mg/mL	% DM (w/v)	% DMSO (v/v)	% BA (v/v)	% Diazepam Recovered
1	100	0 %	100 %	0 %	9.1 ± 2.2
2	100	0 %	95 %	5 %	11.4 ± 1.8
3	100	1%	95 %	5 %	22.5 ± 1.1
4	100	5%	95 %	5 %	32.8 ± 2.1
5	100	10%	95 %	5 %	46.4 ± 2.9
6	75	5%	95 %	5 %	36.2 ± 0.5
7	75	10%	95 %	5 %	51.1 ± 0.6
8	75	17.7%	95 %	5 %	75.1 ± 2.8
9	50	5%	95 %	5 %	44.3 ± 1.4
10	50	10%	95 %	5 %	68.5 ± 0.4
11	50	17.7%	95 %	5 %	97.9 ± 1.2

#### EXAMPLE 4

**[0076]** In this example, either 0% or 2 % (v/v) of polysorbate 20 (PS20)(CAS 9005-64-5) was dispersed into DMSO with 5% (v/v) BA, or DMSO with 5% (v/v) BA and 5% (v/v) NMP, followed by dissolving diazepam at a drug concentration of 100 mg/mL. Diazepam formulations containing PS20 were screened for their ability

to prevent diazepam precipitation using an anti-precipitation assay and RP-HPLC (as detailed above in example 1).

**[0077]** Diazepam formulations containing PS20 significantly improved diazepam recovery in the anti-precipitation assay (Table 4). Inclusion of 1% (v/v) PS20 was not sufficient to prevent diazepam precipitation compared to formulations without the excipient. However, inclusion of 2% (v/v) PS20 improved diazepam recovery by a factor of 2.9. Diazepam formulations containing PS20 and 5% (v/v) NMP were significantly more effective at preventing diazepam precipitation than those without NMP. 1% (v/v) PS20 and 5% (v/v) NMP in the formulation improved diazepam recovery by a factor of 4.1, and inclusion of 2% (v/v) PS20 with 5% (v/v) NMP improved recovery by a factor of 5.4.

Table 4. Diazepam formulations prepared with PS20 in DMSO prevented diazepam precipitation in SEF.

Form.	Diazepam mg/mL	% PS20 (w/v)	% DMSO (v/v)	% BA (v/v)	% NMP (v/v)	% Diazepam Recovered
1	100	0 %	100 %	0 %	0 %	9.1 ± 2.2
2	100	0 %	95 %	5 %	0 %	11.4 ± 1.8
3	100	1 %	94 %	5 %	0 %	9.9 ± 3.4
4	100	2 %	93 %	5 %	0 %	32.9 ± 6.9
5	100	1 %	89 %	5 %	5 %	46.3 ± 2.8
6	100	2 %	88 %	5 %	5 %	61.3 ± 3.9

## EXAMPLE 5

**[0078]** In this example, 10% (w/v) or less of sodium dodecyl sulfate (SDS) (CAS 151-21-3) was dissolved in DMSO with 5% (v/v) BA, followed by dissolving diazepam at a concentration of 50, 75 or 100 mg/mL. Diazepam formulations containing SDS were screened for their ability to prevent diazepam precipitation using an anti-precipitation assay and RP-HPLC (as detailed above in example 1).

**[0079]** Shown in Table 5 are the percent-diazepam recovery data after the formulations prepared with SDS were diluted into SEF. For each concentration of diazepam (50, 75, or 100 mg/mL), increasing concentrations of SDS increased the percent of diazepam that remained soluble when diluted into SEF. When 100 mg/mL

of diazepam was dissolved into DMSO with 5% (v/v) BA and 10% (w/v) SDS is diluted in SEF, the amount of recovered diazepam improved by a factor of 8.2 over diazepam in DMSO and 5% (v/v) BA without SDS. As the concentration of diazepam was decreased, the amount of SDS required to maintain diazepam solubility when diluted into SEF also decreased. The data in Table 5 suggest that there is an ideal ratio that prevents diazepam precipitation, and in this example the molar ratio of SDS to diazepam was ~1:1.

Table 5. Diazepam formulations prepared with SDS in DMSO prevented diazepam precipitation in SEF.

Form.	Diazepam mg/mL	% SDS (w/v)	% DMSO (v/v)	% BA (v/v)	% Diazepam Recovered
1	100	0 %	100 %	0 %	12.8 ± 0.2
2	100	0 %	95 %	5 %	11.4 ± 1.8
3	100	10 %	95 %	5 %	93.6 ± 4.5
4	100	8 %	95 %	5 %	81.9 ± 1.7
5	100	6 %	95 %	5 %	66.7 ± 2.1
6	100	4 %	95 %	5 %	51.9 ± 0.6
7	75	8 %	95 %	5 %	93.4 ± 1.6
8	75	6 %	95 %	5 %	68.0 ± 1.2
9	75	4 %	95 %	5 %	52.2 ± 3.2
10	50	8 %	95 %	5 %	95.5 ± 1.5
11	50	6 %	95 %	5 %	99.2 ± 3.4
12	50	4 %	95 %	5 %	92.3 ± 2.4
13	50	2 %	95 %	5 %	54.8 ± 0.9

#### EXAMPLE 6

**[0080]** Diazepam formulations (100 mg/mL) prepared in DMSO and 5% (v/v) BA (and including different surfactants) were sealed in 2 mL CZ vials (Crystal-Zenith, West Pharmaceuticals, PA, USA) with 13 mm FluroTec® stoppers (rubber stoppers coated with a fluorocarbon film, produced by West Pharmaceuticals), and stored at 40°C / 75% relative humidity for 90 days. A diazepam formulation was also prepared with 2% (v/v) PS20, 5% (v/v) BA, and 5% (v/v) NMP in DMSO. Sealed CZ vials were packaged in foil pouches for the stability study. All formulations were compared with 100 mg/mL diazepam dissolved in DMSO (containing 5% (v/v) BA). The

stability of the formulations are presented as percent diazepam remaining ( $\pm$  standard deviation) in Table 6 below.

Table 6. The stability of diazepam dissolved in DMSO with various excipients at 40°C/75% relative humidity. Stability data is given as the average ( $\pm$  standard deviation) %purity of the diazepam peak for N = 3 replicates.

Solvent System	Excipients	Day 0	Day 90
DMSO + 5% (v/v) BA	None	100	95.3 $\pm$ 0.4
DMSO + 5% (v/v) BA	10% (v/v) PS80	100	98.8 $\pm$ 0.2
DMSO + 5% (v/v) BA	10% (w/v) SDS	100	97.1 $\pm$ 0.1
DMSO + 5% (v/v) BA	2% (w/v) SDC	100	91.0 $\pm$ 0.1
DMSO + 5% (v/v) BA	2% (v/v) PS20	100	100 $\pm$ 0.2
DMSO + 5% (v/v) BA	2% (v/v) PS20 + 5% (v/v) NMP	100	95.6 $\pm$ 0.2

**[0081]** Diazepam dissolved in the DMSO-BA solvent system (with no additional excipients) experienced a 4.2% loss of diazepam purity in the first 30 days when stored at 40°C and 75% relative humidity, but remained at ~95% for the duration of the study. Diazepam formulations prepared with excipients of 10% (v/v) PS80, 10% (w/v) SDS, and 2% (v/v) PS20 exhibited no less than 97.1% diazepam peak purity remaining at the end of the 90-day study. These formulations demonstrated enhanced stability compared to diazepam in DMSO and 5% (v/v) benzyl alcohol (with no additional excipients). Diazepam formulations prepared with 2% (v/v) PS20 and 5% (v/v) NMP exhibited stability comparable to that observed the DMSO-BA solvent system without added excipients. Finally, diazepam formulations prepared with 2% (w/v) SDC had 91% diazepam peak purity remaining at the conclusion of the 90-day stability study.

#### EXAMPLE 7

**[0082]** The stability of the diazepam formulation in a DMSO-BA solvent system was studied through 40 freeze-thaw cycles. Diazepam (100 mg/mL) was dissolved into DMSO containing 5% (v/v) BA and sealed in 2 mL CZ vials with 13 mm FluroTec™ stoppers as described above. Three vials of diazepam formulation were sealed in an aluminum foil pouch and stored in a -20°C freezer for 2 hours. Following storage the pouch was placed at room temperature and allowed to thaw for 2 hours. This process was repeated at a rate of three freeze-thaw cycles per day, where the

third freeze cycle was left overnight. Following several days of freeze-thaw cycles, 2  $\mu$ L of diazepam formulation was removed for analysis by RP-HPLC. Data shown in FIG. 1 shows the percent of diazepam remaining  $\pm$  standard deviation in 5 freeze-thaw cycle intervals.

**[0083]** No significant loss of diazepam was measured following 20 freeze-thaw cycles. However, after 25 cycles the percent of diazepam remaining was significantly different from the pre-frozen sample (bar 0,  $p < 0.05$ ). With each successive interval of 5 freeze-thaw cycles, the percent of diazepam remaining decreased. After 40 cycles, there was  $81.7 \pm 0.5\%$  of diazepam remaining.

#### EXAMPLE 8

**[0084]** In this example the pharmacokinetic (PK) profile of several diazepam-DMSO formulations (with and without excipients) were evaluated in a rat model. Studies were conducted in adult male Sprague-Dawley rats (HillTop Lab Animals, Inc., Scottsdale, PA) between 8 and 10 weeks of age and between 275 to 325 grams, with at least 5 rats per group. Rats were housed individually in stainless steel cages and identified numerically via a permanent marker applied to the tail. Rats were observed twice-daily during acclimation and any abnormal health findings recorded. Animals were fed standard rodent chow and water during the acclimation period, but fasted overnight prior to study initiation. The animals were allowed to acclimatize for 1 week prior to the study.

**[0085]** Test articles for the pharmacokinetics study are described in Table 7. The study contained two commercially available diazepam formulations as comparators – Diastat® rectal diazepam (Valeant Pharmaceuticals) and Hospira’s intramuscular (IM) diazepam formulation. Rats were weighed prior to study initiation to accurately administer the study dose of 3.5 mg/kg. Group 1 was dosed rectally with Diastat and group 2 received an IM injection of diazepam in the rear thigh muscle. Groups 3 through 6 were dosed subcutaneously (SC) in the mid-scapular region with diazepam using 1 of the 4 test formulations. Whole blood samples were collected in vials with K<sub>2</sub> EDTA from each rat via the jugular vein at time 0 (pre-dose), 10, 20, 30, 45, 60, and 120 minutes. Whole blood samples were centrifuged (3000 rpm, 15 minutes) and

plasma was harvested. Plasma samples were frozen at -70°C for future diazepam and nordiazepam analysis.

Table 7. Test articles and dosages administered to rats in the pharmacokinetics study

Group	Test Article	Diazepam Concentration	Route of Administration	Study Dosage
1	Diastat® - AcuDial™ (rectal diazepam)	5 mg/mL	Rectal	~200 µL
2	Hospira Diazepam	5 mg/mL	Intramuscular	~200 µL
3	DMSO alone	100 mg/mL	Subcutaneous	~10 µL
4	2% SDC in DMSO + 5% BA	100 mg/mL	Subcutaneous	~10 µL
5	2% PS20 in DMSO + 5% BA	100 mg/mL	Subcutaneous	~10 µL
6	10% SDS in DMSO + 5% BA	100 mg/mL	Subcutaneous	~10 µL

**[0086]** Rat plasma samples were assayed for diazepam and nordiazepam concentrations by LC/MS using diazepam-d<sub>5</sub> and nordiazepam-d<sub>5</sub> as the internal standards. To each 25 µL sample, 25 µL of 25 ng/mL diazepam-d<sub>5</sub>/nordiazepam-d<sub>5</sub> solution and 0.2 mL of water was added. The samples were vortexed, extracted using Evolute® ABN SPE plates (solid phase extraction, Biotage, Charlotte, NC), eluted with methanol, and dried under nitrogen at approximately 40°C. The dried residues were reconstituted in 500 µL of 1:1 methanol:water and mixed by vortexing. A reversed-phase ultra-performance liquid chromatography (RP-UPLC) and mass spectrometry (MS) method was used to measure diazepam and nordiazepam. The gradient method utilizes a chromatographic column (Waters Atlantis dC-18, 3 µm, 2.1 x 50 mm) with 0.1% formic acid in 60:40 (% v/v) acetonitrile:water as the mobile phase. Analyte detection was performed utilizing an ultraviolet detector at 280 nm for assay and a mass spectrometer for identity confirmation with a 0.25 mL/minute flow rate. Data obtained was expressed in ng/mL. Diazepam and nordiazepam plasma concentration data was analyzed using PKSolver Software, a menu-driven add-in program for Microsoft Excel.

**[0087]** Shown in FIG. 2 are the diazepam plasma concentration profiles (mean ± SEM) for Diastat® rectal diazepam, diazepam in DMSO and 5% (v/v) BA with 2% (w/v) SDC, diazepam in DMSO alone, and diazepam in DMSO and 5% (v/v) BA with 2% (v/v) PS20. The Diastat® group had a Tmax at 10 minutes with Cmax at ~200

ng/mL, and rapid decline in plasma concentration over the remaining 110 minutes. Diazepam in DMSO alone had a  $T_{max}$  at 10 minutes with a  $C_{max}$  of ~70 ng/mL. Unlike Diastat, diazepam in DMSO alone maintained diazepam plasma concentrations at ~50 ng/mL over the remaining 110 minutes. The lack of elimination phase observed in this group (and other DMSO groups) is believed due to partial diazepam precipitation at the injection site, followed by slow absorption. Adding 2% (v/v) PS20 to the diazepam in DMSO formulation resulted in a diazepam plasma concentration spike at 10 minutes, but due to the partial dose precipitation and slow absorption, the  $T_{max}$  (110 ng/mL) occurred at 120 minutes ( $C_{max}$ ). Adding 2% (w/v) SDC to the diazepam in DMSO formulation resulted in diazepam plasma concentration spike at 10 minutes post-injection, but with a delayed  $T_{max}$  (~110 ng/mL) occurring at 60 minutes post-injection ( $C_{max}$ ). Overall the addition of excipients to the diazepam in DMSO formulations incrementally increased the diazepam plasma concentration compared to diazepam in DMSO alone at 10 minutes post-injection.

**[0088]** Shown in FIG. 3 are the combined diazepam (DZ) and nordiazepam (NDZ) PK profiles (mean  $\pm$  SEM) of the intramuscular injection of diazepam (Hospira IM) and the diazepam in DMSO and 5% (v/v) BA with 10% (w/v) SDS formulations. Both diazepam and its metabolite, nordiazepam have pharmacodynamic activity and are clinically relevant. Both the commercial comparator and the DMSO based formulation had a  $T_{max}$  at 30 minutes, and  $C_{max}$  of 424 ng/mL and 396 ng/mL, respectively. The difference in  $C_{max}$  between groups was statistically equivalent ( $p = 0.3123$ ). Together, FIG. 2 and FIG. 3 indicate that a judicious choice of excipient can modulate the diazepam PK profile, promoting either rapid or delayed/slow absorption.

#### EXAMPLE 9

**[0089]** In this example the pharmacokinetic (PK) profile of diazepam in DMSO and 30% (w/v) polysorbate 80 following subcutaneous and intramuscular injection were evaluated in Gottingen mini-pig model. Studies were conducted in female mini-pigs (Xenometrics, Stilwell, KS) weighing between 10 and 12 kilograms, with 3 mini-pigs per group (2 groups). The animals were allowed to acclimatize for 2 weeks prior

to the study. Animals were allowed a 2-week washout period between administrations. Mini-pigs were housed individually in stainless steel cages and identified numerically via a placard on the cage. Mini-pigs were observed twice-daily during the study and any abnormal health findings were recorded. Animals were fasted overnight prior to study initiation through 4 hours after dose administration.

**[0090]** Test articles for the PK study are described in Table 8. The study contained two commercially available diazepam formulation as comparators – Diastat® rectal (PR) diazepam (Valeant Pharmaceuticals) and Hospira’s intramuscular (IM) diazepam formulation. Mini-pigs were weighted prior to study initiation to accurately administer the study dose of 1.0 mg/kg. Group 1 was dosed rectally with Diastat and group 2 received an IM injection of Hospira’s diazepam in the rear thigh muscle. Group 3 received a subcutaneous dose behind the ear with the diazepam test formulation, where Group 4 received an intramuscular dose of the same formulation in the rear thigh. Animals were dosed such that the same group of animals didn’t receive both intramuscular administrations. Whole blood samples were collected in vials with K<sub>2</sub> EDTA from each mini-pig via the jugular vein at time 0 (pre-dose), 0.017, 0.083, 0.167, 0.25, 0.5, 0.75, 1.0, 2.0, 4.0, 6.0, 8.0, and 24.0 hours after dose administration. Whole blood samples were centrifuged (3000 rpm, 15 minutes) and plasma was collected. Plasma samples were frozen at -70°C for future diazepam analysis.

Table 8: Test articles and dosages administered to mini-pigs in the pharmacokinetic study

Group	Test Article	Diazepam Concentration	Route of Administration	Study Dosage
1	Diastat® Rectal Diazepam	5 mg/mL	Rectal	1 mg/kg
2	Hospira Diazepam	5 mg/mL	Intramuscular	
3	Diazepam in DMSO and 30% PS80	100 mg/mL	Subcutaneous	
4	Diazepam in DMSO and 30% PS80	100 mg/mL	Intramuscular	

**[0091]** Mini-pig plasma samples were assayed for diazepam concentrations by LC/MS using diazepam-d<sub>5</sub> as the internal standard. To each 20 µL sample or standard,

500  $\mu$ L of Trizma base buffer was added, followed by vortex-mixing for 1 minute. After mixing, 600  $\mu$ L of ethyl acetate was added to each sample and mixed by inversion 40 times. Six hundred  $\mu$ L of this solution was transferred to a new collection tube and dried under nitrogen at 60°C. The dried product is reconstituted in 100  $\mu$ L of reconstitution solution (1 mM ammonium formate and 0.1% formic acid in 65/35 (%v/v) methanol/water), followed by vortex-mixing for 1 minute. A reversed-phase high performance liquid chromatography (RP-HPLC) and mass spectrometry (MS) method was used to measure diazepam content. The gradient utilized a chromatographic column (Waters ACE® C18, 3  $\mu$ m, 4.6 x 30 mm) with mobile phase A being 1 mM ammonium formate and 0.1% formic acid in 65/35 (v/v) methanol/water and mobile phase B being 1% acetic acid in 90/10 (v/v) methanol/water. Diazepam detection was performed utilizing an ultraviolet detector at 280 nm for assay and mass spectrometer for identity confirmation with a 0.25 mL/minute flow rate. Data obtained was expressed in ng/mL.

**[0092]** Shown in FIG. 4 are the diazepam plasma concentration profiles (mean  $\pm$  SEM) for 100 mg/mL diazepam in DMSO with 30% polysorbate 80 (30% (w/v) PS80) administered by intramuscular (IM) or subcutaneous (SC) injection, Hospira's diazepam administered by intramuscular injection, and Diastat® rectal diazepam. The Diastat® group had a Tmax at 10 minutes with a Cmax of ~271 ng/mL, followed by a rapid decline in plasma concentration over the first 5 hours of the study. The Hospira diazepam group also had a Tmax at 10 minutes and a Cmax of 425.3 ng/mL, followed by a rapid decline in plasma concentration, though not as rapid as Diastat rectal gel. The diazepam formulation in DMSO with 30% PS80 administered by IM injection had a Tmax at 45 minutes, Cmax of 391.3 ng/mL, and an elimination profile that matched Hospira's IM administration. The SC administration of diazepam in DMSO and 30% (w/v) PS80 resulted in three peak plasma concentrations, with a true Cmax of 202.0 ng/mL occurring at 4.0 hours. Depending on the route of administration, the pharmacokinetic profile and peak plasma concentrations of diazepam in DMSO with 30% (w/v) polysorbate 80 can have a rapid onset or prolonged systemic circulation.

## EXAMPLE 10

**[0093]** The stability of diazepam in DMSO with 30% (w/v) polysorbate 80 was studied when stored in glass vials at 75°C for 20 days. Diazepam (100 mg/mL) was dissolved into DMSO containing 30% (w/v) polysorbate 80 and sealed into 10 mL borosilicate glass vials with 20 mm FluroTec™ stoppers and 20 mm flip-off seals. Vials were stored at 75°C and three vials were pulled on days 5, 12, and 20. Percent-diazepam remaining in each vial was determined by RP-HPLC. Data in FIG. 5 shows the percent diazepam remaining ± standard deviation.

**[0094]** After 5 and 12 days of storage at 75°C the percent-diazepam remaining was not significantly different from the initial (time zero) measurement ( $p = 0.431$ ). However, after 8 additional days of storage at 75°C (day 20), the percent diazepam decreased by approximately 6.5% (106.98% to 99.99%,  $p = 0.001$ ). Despite the reduction in the percent-diazepam remaining after storage at 75°C, the results indicate diazepam at 100 mg/mL is stable in DMSO with 30% polysorbate 80 (w/v).

## EXAMPLE 11

**[0095]** The stability of diazepam in DMSO with 30% (w/v) polysorbate 80 was studied when stored in glass vials at 4 - 8°C, 25°C and 60% relative humidity, 30°C, and 40°C and 75% relative humidity. Diazepam (100 mg/mL) was dissolved in DMSO containing 30% (w/v) polysorbate 80 and sealed in 10 mL borosilicate glass vials with 20 mm FluroTec™ stoppers and 20 mm flip-off seals. Vials were stored under the aforementioned storage conditions in triplicate. Percent-diazepam remaining in each vial was determined by RP-HPLC following 30 days of storage. Data in FIG 6 shows the percent-diazepam remaining ± standard deviation.



**[0096]** Initial diazepam content was measured by RP-HPLC at time zero and is represented in the graph by the bottom bar labeled “Initial”. The percent-diazepam remaining at any storage condition (4 - 8°C, 25°C/60% RH, 30°C, and 40°C/75% RH) was at 100%, showing no loss of diazepam after 30 days of storage.

## CLAIMS



1. A stable precipitation resistant formulation for parenteral injection comprising:
  - (a) a biocompatible non-aqueous solvent; and
  - (b) a small molecule drug, or a salt thereof, solubilized within the non-aqueous solvent; and
  - (c) 1% to 30% w/w of a surfactant, wherein the surfactant attenuates the precipitation of the small molecule drug when injected into a subject.
2. The formulation of claim 1, wherein the surfactant is present at a molar ratio of surfactant:small molecule drug of 0.5:1 to 4:1, or 1:1 to 2:1.
3. The formulation of claim 1, wherein the surfactant is sodium deoxycholate, polysorbate 80, polysorbate 20, dodecyl maltoside, sodium dodecyl sulfate, sodium tetradecyl sulfate, alcohol ethoxylate, alkyldimethylamine oxide, or alkyl betaine.
4. The formulation of claim 1, wherein the liquid formulation comprises less than 10% by weight moisture content.
5. The formulation of claim 1, wherein the volume of the liquid formulation to be parenterally injected is 3 mL or less.
6. The formulation of claim 1, further comprised within a device for dispensing the formulation.
7. The formulation of claim 6, wherein the device is a syringe, a pen injection device, an auto-injector device, an external or implantable pump, or a perfusion bag.
8. The formulation of any one of claims 1 to 7, wherein the biocompatible non-aqueous solvent is an aprotic polar solvent, an alkyl or aryl benzoate solvent, a lipid solvent, a protic solvent, or a mixture thereof.
9. The formulation of claim 8, wherein the formulation comprises the aprotic polar solvent, and wherein the aprotic polar solvent is dimethylsulfoxide (DMSO), dimethylformamide (DMF), ethyl acetate, n-methyl pyrrolidone (NMP), dimethyl acetamide (DMA), propylene carbonate, or mixtures thereof.

10. The formulation of claim 9, wherein the aprotic polar solvent is DMSO, NMP, or a mixture thereof.
11. The formulation of claim 8, wherein the formulation comprises an aryl or alkyl benzoate solvent, and wherein the aryl or alkyl benzoate solvent is ethyl benzoate, benzyl benzoate, or mixtures thereof.
12. The formulation of any one of claims 1 to 7, wherein the liquid formulation comprises less than 10% by weight moisture content.
13. The formulation of any one of claims 1 to 7, wherein the liquid formulation includes from 0.5 mg/mL to 750 mg/mL of the small molecule drug.
14. The formulation of any one of claims 1 to 7, wherein the volume of the liquid formulation to be parenterally injected is from 1  $\mu$ L to 10  $\mu$ L.
15. The formulation of any one of claims 1 to 7, wherein the volume of the liquid formulation to be parenterally injected is from 10  $\mu$ L to 100  $\mu$ L.
16. The formulation of any one of claims 1 to 7, wherein the volume of the liquid formulation to be parenterally injected is from 100  $\mu$ L to 1 mL.
17. The formulation of any one of claims 1 to 7, wherein the small molecule drug is a benzodiazepine.
18. The formulation of claim 17, wherein the benzodiazepine is diazepam.
19. The formulation of claim 17, wherein the liquid formulation comprises 25 mg/mL to 300 mg/mL of a benzodiazepine.
20. The formulation of claim 11, wherein the solvent is DMSO, NMP, or a mixture thereof.
21. A method of administering the formulation of any of claims 1 to 20 to a subject in need thereof comprising parenterally injecting the formulation to the subject.



22. The method of claim 21, wherein injecting is by parenteral injection or intracutaneous injection.
23. The method of claim 21, wherein the formulation is not diluted prior to administration.
24. A method for treating or preventing a condition comprising parenterally administering to a subject in need thereof the formulation of any of claims 1 to 20 in an amount effective to treat or prevent the condition.
25. A method for treating or preventing anxiety, muscle spasms, or seizures, the method comprising parenterally administering to a subject in need thereof a formulation of claim 1 comprising a benzodiazepine.
26. The method of claim 25, further comprising the injectable volume within a device for dispensing the liquid formulation.
27. The method of claim 26, wherein the device is a syringe, a pen injection device, an auto-injector device, an external or implantable pump, or a perfusion bag.
28. The method of claim 25, wherein the liquid formulation comprises 25 mg/mL to 300 mg/mL of a benzodiazepine.
29. The method of claim 25, wherein the benzodiazepine is diazepam.
30. The method of claim 25, wherein the volume of the liquid formulation to be parenterally injected is from 1  $\mu$ L to 10  $\mu$ L.
31. The method of claim 25, wherein the volume of the liquid formulation to be parenterally injected is from 10  $\mu$ L to 100  $\mu$ L.
32. The method of claim 25, wherein the volume of the liquid formulation to be parenterally injected is from 100  $\mu$ L to 1 mL.
33. The method of claim 25, wherein the liquid formulation is not diluted prior to administration.

**FIG. 1****FIG. 2**

2/3

**FIG. 3****FIG. 4**

3/3

**FIG. 5****FIG. 6**

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2018/035473

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>	INV. A61K9/00	A61K47/14	A61K47/16	A61K47/20	A61K47/22
	A61K9/08	A61K31/5513			

**ADD.**

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2014/004895 A1 (XERIS PHARMACEUTICALS INC [US]) 3 January 2014 (2014-01-03) paragraph [0006]; claim 1; example 1 -----	1-33
Y	WO 2009/027697 A2 (ARCHIMEDES DEV LTD [GB]; WATTS PETER JAMES [GB]; CHENG YU-HUI [GB]; SM) 5 March 2009 (2009-03-05) examples -----	1-33
X	EP 2 030 610 A1 (ARCHIMEDES DEV LTD [GB]) 4 March 2009 (2009-03-04) examples -----	1-33
Y	WO 2005/021046 A1 (DBL AUSTRALIA PTY LTD) 10 March 2005 (2005-03-10) the whole document ----- -/-	1-33

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
---	--

20 July 2018

31/07/2018

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer
--	--------------------

Giese, Hans-Hermann

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2018/035473

**C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2007/059019 A2 (WINCH PETER D [US]) 24 May 2007 (2007-05-24) claims 133-138; examples 32,33	1-33
A	WO 2006/122217 A2 (UNIV MINNESOTA [US]; CLOYD JAMES C [US]; SIEGEL ROLAND A [US]) 16 November 2006 (2006-11-16) paragraphs [0010], [0013], [0014]	1-33
A	US 6 365 637 B1 (ZIRNSTEIN MICHAEL [DE] ET AL) 2 April 2002 (2002-04-02)	1-33
A	example 18	-----
A	US 5 397 771 A (BECHGAARD ERIK [DK] ET AL) 14 March 1995 (1995-03-14)	1-33
	figure 6	-----

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/US2018/035473

Patent document cited in search report	Publication date	Patent family member(s)			Publication date
WO 2014004895	A1 03-01-2014	AU 2013284412 A1 BR 112014032695 A2 CA 2877909 A1 CN 104394848 A CN 107913248 A EP 2874599 A1 HK 1204771 A1 JP 2015523361 A KR 20150027263 A MX 345256 B US 2014005135 A1 US 2016151385 A1 WO 2014004895 A1			22-01-2015 27-06-2017 03-01-2014 04-03-2015 17-04-2018 27-05-2015 04-12-2015 13-08-2015 11-03-2015 23-01-2017 02-01-2014 02-06-2016 03-01-2014
WO 2009027697	A2 05-03-2009	AU 2008291873 A1 BR PI0815782 A2 CA 2697825 A1 CN 101842079 A EP 2194965 A2 ES 2598114 T3 HK 1148464 A1 JP 5539875 B2 JP 2010537963 A KR 20100047311 A NZ 583265 A TW 200922635 A US 2009233912 A1 WO 2009027697 A2 ZA 201001303 B			05-03-2009 24-02-2015 05-03-2009 22-09-2010 16-06-2010 25-01-2017 01-03-2013 02-07-2014 09-12-2010 07-05-2010 12-01-2012 01-06-2009 17-09-2009 05-03-2009 23-12-2015
EP 2030610	A1 04-03-2009	NONE			
WO 2005021046	A1 10-03-2005	CA 2546482 A1 EP 1667656 A1 JP 2007504256 A US 2007108405 A1 WO 2005021046 A1			10-03-2005 14-06-2006 01-03-2007 17-05-2007 10-03-2005
WO 2007059019	A2 24-05-2007	AU 2006315610 A1 CA 2635496 A1 EP 1951248 A2 JP 2009515895 A US 2009156562 A1 WO 2007059019 A2			24-05-2007 24-05-2007 06-08-2008 16-04-2009 18-06-2009 24-05-2007
WO 2006122217	A2 16-11-2006	US 2007021411 A1 US 2007208011 A1 WO 2006122217 A2			25-01-2007 06-09-2007 16-11-2006
US 6365637	B1 02-04-2002	AT 302615 T CN 1253021 A DE 19851777 A1 EP 1002546 A2 JP 4819210 B2 JP 2000143544 A US 6365637 B1			15-09-2005 17-05-2000 11-05-2000 24-05-2000 24-11-2011 23-05-2000 02-04-2002
US 5397771	A 14-03-1995	NONE			

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No  
PCT/US2018/035473

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
<hr/>			