

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2016/210224 A1

(43) International Publication Date  
29 December 2016 (29.12.2016)

WIPO | PCT

(51) International Patent Classification:  
*C12N 15/10* (2006.01)    *C12Q 1/68* (2006.01)  
*C12P 19/34* (2006.01)

(21) International Application Number:  
PCT/US2016/039167

(22) International Filing Date:  
24 June 2016 (24.06.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
62/183,854    24 June 2015 (24.06.2015)    US

(71) Applicant: DANA-FARBER CANCER INSTITUTE, INC. [US/US]; 450 Brookline Avenue, Boston, MA 02215 (US).

(72) Inventors: MAKRIGIORGOS, Gerassimos; 63 Payson Road, Chestnut Hill, MA 02467 (US). SONG, Chen; 270 Babcock Street, Apt. 5C, Boston, MA 02215 (US).

(74) Agent: COLANTONIO, Jessica R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: SELECTIVE DEGRADATION OF WILD-TYPE DNA AND ENRICHMENT OF MUTANT ALLELES USING NUCLEASE




Fig. 1A

(57) Abstract: The present disclosure provides methods for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid using nucleases that have a substantially higher activity on double stranded DNA versus single stranded DNA or RNA. The present disclosure also includes methods for enriching a target mutant nucleic acid and for preparing unmethylated/methylated nucleic acids of interest for subsequent enrichment.

WO 2016/210224 A1

## SELECTIVE DEGRADATION OF WILD-TYPE DNA AND ENRICHMENT OF MUTANT ALLELES USING NUCLEASE

### RELATED APPLICATIONS

5 This application claims priority under 35 U.S.C. § 119(e) to U.S. provisional application number 62/183,854, filed June 24, 2015, the contents of which are incorporated herein by reference in their entirety.

### GOVERNMENT SUPPORT

10 This invention was made with government support under grant number R21 CA175542 awarded by The National Institutes of Health. The government has certain rights in the invention.

### BACKGROUND OF THE INVENTION

15 A commonly encountered situation in genetic analysis entails the need to identify a low percent of variant DNA sequences ('minority alleles') in the presence of a large excess of non-variant sequences ('majority alleles'). Examples for such situations include the following: identification and sequencing of a few mutated alleles in the presence of a large excess of normal (wild-type) alleles, a commonly encountered 20 situation in cancer (for example, identification of tumor-circulating DNA in blood or in urine of cancer patients (or abnormal DNA in people suspected of having cancer) in the presence of a large excess of wild type alleles); identification of a few methylated alleles in the presence of a large excess of unmethylated alleles (or vice-versa) in epigenetic analysis; identification and genotyping of a few fetal DNA sequences circulating in the 25 maternal blood where a large excess of maternal DNA sequences are also present; identification of emerging mutated strains in infectious agents (bacteria or viruses); and variant sequence detection for crop development.

30 While reliable high throughput screening methods for germline or high-prevalence somatic mutations have been described, detection of low-prevalence somatic mutations in tumors with heterogeneity, stromal contamination, or in bodily fluids is still problematic. And yet, the clinical significance of identifying these mutations is very important in several situations. For example, in lung adenocarcinoma, low-level EGFR mutations that cannot be identified by regular sequencing can confer positive response to

tyrosine kinase inhibitors or drug resistance. Mutations in plasma useful as biomarkers for early detection of cancer or cancer response to treatment, cannot be sequenced using conventional methods due to the high excess of wild type alleles originating from normal tissues. Additionally, mutations in tumors with frequent stromal contamination, such as 5 pancreatic or prostate cancer, can be ‘masked’ by presence of wild type alleles, thus requiring laborious micro-dissection or resulting in missing mutations altogether.

Beyond cancer, low levels of target DNA in the presence of high levels of non-target DNA occurs in many other fields and applications. For example, the detection of a small amount of fetal alleles within maternal alleles is especially important for prenatal 10 diagnosis during early stages in pregnancy where fetal alleles comprise a low fraction. An especially interesting application to this end is the fact that fetal alleles are substantially hypomethylated compared to maternal alleles. Thus, there is a general need to develop techniques that allow for identification of low level minority alleles (for example, mutated or hypo/hypermethylated alleles) in the presence of high level non-15 variant majority alleles.

#### SUMMARY OF THE INVENTION

The present disclosure relates to a novel development (Nuclease-assisted Mutation Enrichment, NaME) that results to preferential cleavage of wild type nucleic acids, thereby allowing for subsequent enrichment of mutated target sequences of 20 interest. The mutation-enriched sequences can then be screened using any currently available methods for identifying mutations, such as Sanger Sequencing, high resolution melting (HRM), *etc.*

Accordingly, some aspects of the disclosure provide a method for preparing a 25 target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid. The method comprises subjecting a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a condition that destabilizes the double stranded wild type and target mutant nucleic acids; contacting the destabilized double stranded wild type and target 30 mutant nucleic acids with a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby

forming complimentary wild-type-probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; and exposing the complimentary wild-type-probe duplexes and the partially complimentary target mutant-probe duplexes to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

Some aspects of the disclosure provide a method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; and subjecting the reaction mixture to a condition that destabilizes the double stranded wild type and target mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

In some embodiments, the condition that destabilizes the double stranded wild type and mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and mutant nucleic acids comprises addition of an organic solvent and/or an increase in temperature combined with a thermostable DSN.

Some aspects of the disclosure provide a method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to

create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; reducing the temperature of the reaction mixture to permit 5 formation of complimentary wild type – probe duplexes on top and bottom strands and partially complimentary target mutant-probe duplexes; and exposing the reaction mixture to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

10 In some embodiments, the method is used to prepare an unmethylated target nucleic acid of interest for subsequent enrichment, and wherein prior to implementing the NaME protocol described herein on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite and one of the oligonucleotide probes is complimentary to top strand of the methylated nucleic acid of interest, while the other oligonucleotide 15 probe is complimentary to the bottom strand of the methylated nucleic acid of interest.

20 In some embodiments, the method is used to prepare a methylated target nucleic acid of interest for subsequent enrichment, and wherein prior to implementing the NaME protocol described herein on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite and one of the oligonucleotide probes is complimentary to top strand of the unmethylated nucleic acid of interest, while the other oligonucleotide probe is 25 complimentary to the bottom strand of the unmethylated nucleic acid of interest.

30 In some embodiments, the method is used to prepare both an unmethylated target nucleic acid of interest and a methylated target nucleic acid of interest for subsequent enrichment, wherein the method comprises: (i) a pair of oligonucleotide probes, one of which is complimentary to top strand of the methylated form of the unmethylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the methylated form of the unmethylated target nucleic acid of interest, (ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of the methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the unmethylated form of the methylated target nucleic acid of interest; and wherein prior to implementing the NaME protocol described herein on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite.

In some embodiments, the method is used to prepare multiple target nucleic acids of interest, some of which are methylated target nucleic acids of interest, and some of which are unmethylated target nucleic acids of interest, and the method comprises (i) a pair of oligonucleotide probes, one of which is complimentary to top strand of the 5 methylated form of each unmethylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the methylated form of each unmethylated target nucleic acid of interest, (ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of each methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the 10 unmethylated form of each methylated target nucleic acid of interest.

Some aspects of the disclosure provide a method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid, the method comprising the steps of: (a) exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a thermostable double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; and (c) reducing the temperature to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

In some embodiments, steps (b) and (c) are repeated for two or more cycles. In some embodiments, the reaction mixture further comprises an organic solvent. In some embodiments, the denaturing temperature is between 65-85°C.

30 Furthermore, in some embodiments step (c) of reducing the temperature is performed by applying a decreasing temperature gradient between temperatures that permit hybridization of probes having different  $T_m$  to their corresponding sequences on the target DNA. For example, a decreasing temperature gradient from 67°C to 64°C can

be applied during DSN digestion, to allow diverse probes that have distinct Tms of 64-67°C to hybridize effectively to their respective targets. ('Touch-down NaME'). The temperature gradient can preferably range between 2-20°C.

Some aspects of the disclosure provide a method for enriching a target mutant nucleic acid, the method comprising the steps of: (a) preparing an amplification reaction mixture comprising: a double-stranded wild type nucleic acid, a double-stranded target nucleic acid suspected of containing a mutation, a thermostable double strand-specific nuclease (DSN), a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation and PCR amplification components; (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; (c) reducing the temperature to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes; and (d) subjecting the reaction mixture to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid.

In some embodiments the amplification condition is such that amplification is applied to the probes rather than the hybridized nucleic acid. In some embodiments, a purification step is applied following probe binding to top-and-bottom DNA target strands, either before or after DSN cleavage, to remove excess unbound probes. Then following DSN cleavage the uncut probes are amplified (instead of amplifying the target DNA) and identified/quantified. Since probes that bind WT DNA will have been selectively digested by DSN, the presence of any given probe after amplification indicates a mutation under the region covered by this probe.

In some embodiments, steps (b) and (c) are repeated for two or more cycles before executing step (d). In some embodiments, steps (b), (c) and (d) are repeated for two or more cycles. In some embodiments, the reaction mixture further comprises an organic solvent. In some embodiments, the denaturing temperature is between 65-85°C. In some embodiments, the primers used for PCR amplification have a melting

temperature that is below the temperature applied in step (c). In some embodiments, the amplification condition is COLD-PCR.

Some aspects of the disclosure provide a method for preparing unmethylated nucleic acids of interest for subsequent enrichment relative to corresponding methylated nucleic acids comprising the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated nucleic acids while methylated nucleic acids remain double-stranded; (d) exposing the unmethylated and methylated nucleic acids to double strand-specific nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN cleaves the methylated double-stranded nucleic acids but not the unmethylated single-stranded nucleic acids.

Some aspects of the disclosure provide a method for preparing unmethylated nucleic acids of interest for subsequent enrichment relative to corresponding methylated nucleic acids comprising the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits denaturation of both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids; (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and (e) exposing the unmethylated and methylated nucleic acids to double strand-specific nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN preferentially cleaves the methylated duplexes but not the unmethylated single-stranded nucleic acids.

Some aspects of the disclosure provide a method for preparing methylated nucleic acids of interest for subsequent enrichment relative to corresponding unmethylated nucleic acids comprising the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated

nucleic acids while methylated nucleic acids remain double-stranded; and (d) exposing the unmethylated and methylated nucleic acids to an exonuclease and conditions for optimal exonuclease activity, wherein the exonuclease cleaves the unmethylated single-stranded nucleic acids but not the methylated double-stranded nucleic acids.

5 Some aspects of the disclosure provide a method for preparing methylated nucleic acids of interest for subsequent enrichment relative to corresponding unmethylated nucleic acids comprising the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to 10 form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits denaturation of both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids; (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and (e) exposing the unmethylated 15 and methylated nucleic acids to an exonuclease and conditions for optimal exonuclease activity, wherein the exonuclease preferentially cleaves the unmethylated single-stranded nucleic acids, but not the methylated duplexes.

In some embodiments, the nucleic acid amplification reaction of step (b) is selected from the group consisting of: PCR; full COLD-PCR, fast COLD-PCR; ice-COLD-PCR, temperature-tolerant COLD-PCR and limited denaturation time COLD-PCR. In some embodiments, the cleaved unmethylated single stranded nucleic acids and the uncleaved methylated duplexes are subjected to an amplification condition using the bisulfite resistant adaptors ligated in step (a). In some embodiments, the amplification condition is selected from the group consisting of: PCR, full COLD-PCR, fast COLD-PCR; ice-COLD-PCR, temperature-tolerant COLD-PCR and limited denaturation time COLD-PCR. In some embodiments, naturally AT-rich sequences are removed prior to 25 the sodium bisulfite treatment.

In any of the foregoing methods, the probes are in a molar excess of 100-fold to 1 billion-fold compared to the wild type and target nucleic acids. In any of the foregoing 30 methods, one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other. In any of the foregoing methods, each probe comprises a locked

nucleic acid (LNA), peptide nucleic acid (PNA), xeno nucleic acid (XNA), nucleic acid with any known modified base or RNA. In any of the foregoing methods, the method is used to prepare two or more different target mutant nucleic acids for subsequent enrichment relative to corresponding wild type nucleic acids, and the method further

5 comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand. In any of the foregoing methods, the nucleic acid sample comprises genomic DNA or circulating DNA in urine, plasma or other bodily fluids.

10 In any of the foregoing methods, the methods further comprise enriching the nucleic acids for regions of interest prior to implementing NaME protocol described herein as follows: contacting the nucleic acid sample with bait oligonucleotides that bind to different nucleic acids of interest on top and bottom strands, permitting binding of the bait oligonucleotides to the nucleic acids of interest on top and bottom strands, and

15 isolating the bait oligonucleotides with the nucleic acids of interest bound thereto from remaining nucleic acids. In some embodiments, the bait oligonucleotides are biotinylated at one end. In some embodiments, the bait oligonucleotides are attached to beads.

In any of the foregoing methods, prior to implementing the NaME protocol described herein, the nucleic acid sample is subjected to an amplification condition.

20 In any of the foregoing methods, the methods further comprise enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid. In some embodiments, the

25 amplification condition is selected from the group consisting of: PCR, COLD-PCR, ligation mediated PCR or COLD-PCR using common ligated adaptors, multiplex PCR, and isothermal amplification.

In any of the foregoing methods, each probe can be optionally modified at the 3' end to prevent polymerase extension.

30 In any of the foregoing methods, the methods further comprise enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type -probe duplexes and uncleaved target mutant nucleic acids to a further DNA degradation condition which hydrolyzes enzymatically the DSN-

cleaved wild type-probe duplexes, with the degradation initiated at the position of the cleavage.

In any of the foregoing methods, the methods further comprise analyzing the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids using one or more of the methods selected from the group consisting of: 5 MALDI-TOF, HR-Melting, Di-deoxy-sequencing, Single-molecule sequencing, massively parallel sequencing (MPS), pyrosequencing, SSCP, RFLP, dHPLC, CCM, digital PCR and quantitative-PCR.

In some embodiments of any one of the provided methods, the DSN enzyme is a 10 DNA guided or RNA guided enzyme. In some embodiments of any one of the provided methods, the enzyme is an RNA guided enzyme, e.g., Cas9. In some embodiments of any one of the provided methods, the enzyme is a DNA guided enzyme, e.g., an Argonaute enzyme.

Some aspects of the disclosure provide a method for preparing a target mutant 15 nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising subjecting a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a condition that destabilizes the double stranded wild type and target mutant nucleic acids; contacting the destabilized double stranded wild type and target mutant nucleic acids 20 with a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild-type-probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein at least one of the probes overlaps a sequence on the target 25 nucleic acid containing the suspected mutation, and wherein one or both probes comprise a locked nucleic acid (LNA), peptide nucleic acid (PNA), xeno nucleic acid (XNA), or a nucleic acid with any known modified base or RNA which is capable of blocking PCR amplification; and subjecting the complimentary wild-type- probe duplexes on top and 30 bottom strands, and partially complimentary target mutant-probe duplexes to an amplification condition. The probe(s) that overlap the mutation position act to block PCR amplification, e.g., acting as a clamp, for the wild-type top and bottom DNA strands, thereby inhibiting amplification of the wild-type nucleic acid. When the probe

duplexes with a partially complimentary target mutant sequence, it is less able to inhibit PCR amplification, thereby permitting selective amplification of the mutant nucleic acid as compared to the wild-type, without a need for a cleaving enzyme (e.g., DSN).

Each of the embodiments and aspects of the invention can be practiced  
5 independently or combined. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including", "comprising", or "having", "containing", "involving", and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

10 These and other aspects of the inventions, as well as various advantages and utilities will be apparent with reference to the Detailed Description. Each aspect of the invention can encompass various embodiments as will be understood.

All documents identified in this application are incorporated in their entirety herein by reference.

15

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1A is a schematic the selective degradation of wild-type double-stranded DNA using duplex specific nuclease (DSN) and sequence selective oligonucleotides ('probes') at elevated temperatures. A DNA denaturation step prior to DSN action may  
20 optionally be applied in order to generate single stranded DNA prior to probe binding and selective wild type degradation of both DNA strands.. Fig. 1B shows the validation of NaME-based mutation enrichment as described in Fig 1A via digital PCR (ddPCR) for a mutated KRAS amplicon with 5% original mutation abundance (mutation abundance = fractional ratio of mutant to wild-type DNA, expressed as a percentage). In this example, the DNA did not undergo a 95°C denaturation step; the DNA, two probes, and DSN were  
25 mixed and the temperature was elevated (67°C) to de-stabilize the duplex and enable probe binding. A range of temperatures was applied to identify the optimal temperatures for WT-specific digestion. Following DSN action, the DSN was inactivated via heating at 95°C and droplet digital PCR (ddPCR) was applied to the digested sample. ddPCR  
30 quantifies the mutation enrichment achieved by measuring fractional mutation abundance before and after DSN action.

Fig. 2 is a schematic showing the use of partially overlapping probes to provide selectivity simultaneously on both DNA strands when binding a mutation. Probes are

preferably 3'-blocked to prevent subsequent polymerase extension.

Fig. 3 is a schematic showing the use of duplex specific nuclease and partially overlapping sequence selective oligonucleotides to enable the selective degradation of double-stranded, or denatured, DNA. With the mutant sequence, there is limited complementarity between the top and bottom strand probes, which prohibits probe-to-probe interactions at higher temperatures.

Fig. 4 is a schematic showing wild-type allele degradation directly from fragmented genomic DNA (i.e. not pre-amplified DNA). Note that a multiplex approach can be taken; thousands of probes can be used simultaneously on selected DNA targets of interest.

Fig. 5 shows the results of a single-plex assay showing the major increase in mutation abundance of mutational sequences following a DSN reaction directly on fragmented genomic DNA for a selected TP53 exon 8 target sequence. The mutation abundance is quantified before and after treatment of the sample with DSN, using ddPCR. The mutation abundance increases only if both top and bottom probes are included in the reaction, in addition to DSN nuclease.

Fig. 6 shows the results of a duplex assay, showing the mutation abundance of a sample containing mutated KRAS and p53 at 5% or 0.3% original abundance, following a DSN reaction directly on fragmented genomic DNA.

Fig. 7A shows the results of a DSN reaction on genomic DNA for a single-plex assay using DNA mutated at three different positions on KRAS exon 2 in three different cell lines: H2009, A549, and HCT-15.

Fig. 7B is a table summarizing the mutation abundance found in a DSN reaction performed directly on fragmented genomic DNA via an 11-plex assay. The 11 mutated targets were formed using DNA from Horizon Dx. In the figure, column 1 shows the name of the target gene, columns 2 and 3 represent the mutation position and mutation type, respectively, column 4 shows the mutation abundance as derived via digital PCR when DSN is omitted; column 5 shows the mutation abundance when DSN is applied (NaME), column 6 represents the mutation abundance when both DSN and the probes are omitted, and column 7 shows the expected mutation abundance according to the Horizon manufacturer, without any treatment.

Fig. 8 is a schematic illustrating two approaches to selective degradation of wild-type DNA prior to targeted re-sequencing: one after multiplexed PCR (top), and one

prior to PCR (bottom) using selective binding of DNA targets to beads. In the latter situation, the probes utilized are designed to be partially overlapping each other and bind both top and bottom DNA target strands. In addition, they are biotinylated. These probes are first used without DSN to bind their targets and to immobilize selected DNA 5 targets to beads. The non-immobilized DNA is then removed from the solution. The temperature then is then adjusted accordingly, and NaME is applied as described above.

Fig. 9 is a schematic illustrating the 'nuclease chain reaction'. Wild-type degradation is combined with denaturation cycles during the DSN digestion process resulting in improved discrimination between mutant and WT and, therefore, better 10 mutation enrichment. Note that, following brief denaturation at 85°C and cooling to 65°C, the reaction cycles again at 85°C before substantial re-hybridization of the two strands can occur.

Fig. 10 is a schematic illustrating the PCR-NaME chain reaction where both PCR amplification and nuclease chain reaction operate simultaneously on the sample, in a 15 single tube. The successive cycles of PCR synthesis and wild-type-specific degradation lead to improved enrichment of mutated sequences.

Fig. 11 is a schematic illustrating COLD-PCR-NaME. The successive cycles of mutant-specific synthesis and wild-type-specific degradation enrich for mutated sequences.

20 Fig. 12A is schematic illustrating mutation scanning using two or more longer (non-overlapping) probes on opposite strands. Probes are preferentially 3'-blocked to prevent polymerase extension, and may contain modified bases, such as LNA, PNA, XNA, deoxy- inosine triphosphate (dITP), or contain dUTP, or comprise RNA. In some cases part of the probes can comprise one or more random nucleotides, so that the probe 25 can be directed against a plurality of DNA targets. The total combined sequence under the two probes is interrogated during NaME: if there is a mutation anywhere under the two probes, it will prevent strand cutting.

Fig. 12B shows that the combination of denaturation and DSN for mutation scanning as described in 12A results in the preferential cutting of wild-type DNA. A 30 mutation present anywhere under the two probes results in the amplification of the mutated DNA during subsequent PCR or digital PCR. The effects of probe length and concentration on mutation enrichment are depicted in the graphs.

Fig. 12C shows mutation scanning using two adjacent probes and DSN in NaME

as described in 12A. The graph presents the average mutation enrichment-fold of each listed mutation.

Fig. 13 is a schematic representing the application of NaME with RNA or single-stranded DNA. Multiplexed wild-type nucleic acid degradation is used to enrich mutants prior to cDNA synthesis or PCR.

Fig. 14 is a schematic showing hypomethylation enrichment from fragmented genomic DNA using probes designed to match the methylated alleles on the target sequences.

Fig. 15 shows the enrichment of unmethylated DNA sequences and methylation-sensitive temperature-tolerant-COLD-PCR following preferential DSN digestion of the sequences having a Tm higher than the Tm of choice. This scheme is a genome-wide application and does not require the use of gene-specific probes or selection of target sequences. The scheme enriches and amplifies hypo-methylated alleles on a genome-wide basis.

Fig. 16 depicts the process of using exonuclease-based temperature fractionation of DNA fragments to remove lower Tm fragments prior to bisulfite conversion.

Fig. 17 is a flowchart showing the temperature-based elution of genomic DNA fragments by utilizing binding of DNA to magnetic beads.

20

## DETAILED DESCRIPTION OF THE INVENTION

In most applications involving detection of low-prevalence somatic mutations, the mutant alleles are detected following a polymerase chain reaction (PCR) step that amplifies both mutant and wild type alleles. Methods have also been described to preferentially amplify the mutated alleles over wild-type alleles (*e.g.* co-amplification at lower denaturation temperature or COLD-PCR and improved and complete enrichment COLD PCR or ice-COLD-PCR; Li J, Wang L, Mamon H, Kulke MH, Berbeco R, Makrigiorgos GM. Replacing PCR with COLD-PCR enriches variant DNA sequences and redefines the sensitivity of genetic testing. *Nat Med* 2008;14:579-84; Milbury CA, Li J, Makrigiorgos GM. Ice-COLD-PCR enables rapid amplification and robust enrichment for low-abundance unknown DNA mutations. *Nucleic Acids Res*;39:e2). However, the enrichment that can be obtained via such PCR-based methods has a limit, since after several cycles of synthesis, the polymerase unavoidably introduces mis-incorporations (PCR errors) that are subsequently scored as mutations. Repeated

amplifications can also introduce mis-priming which results in the amplification of unwanted non-target sequences. Furthermore, there are powerful genetic analysis methods currently emerging ('third generation sequencing' Nanopore, Pac-Bio systems) that may obviate the use of PCR altogether. Therefore, mutation enrichment methods 5 that reduce the amount of PCR performed, or that can operate without PCR, or in conjunction with PCR if so required, are important.

The present disclosure is based, at least in part, on the novel development of a technique, Nuclease-assisted Mutation Enrichment (NaME) that results in the preferential cleavage of non-variant/wild type DNA or RNA, thereby allowing for 10 subsequent selective enrichment of variant (mutant) target sequences. Thus, NaME can be used before, during, or after an amplification step, such as PCR, , or without any amplification, depending on the application. Subsequently, the mutation-enriched sequences can be screened via any currently available method for identifying mutations, including Sanger Sequencing, high resolution melting (HRM), SSCP, next generation 15 massively parallel sequencing, and MALDI-TOF for known mutations; and Single Molecule Sequencing-or third generation sequencing for high-throughput sequencing of low-level mutations. NaME can also be applied to detect low levels of un-methylated alleles (Methylation-Sensitive Nuclease-assisted minor-allele Enrichment or MS-NaME) in a background of methylated alleles (or vice-versa).

20 The methods described herein greatly improve the current detection limits of mutation/methylation detection technologies, thereby enhancing reliability of patient-specific mutation screening, for example, in heterogeneous tumor samples or in circulating DNA. The methods described herein also enable high multiplexity of targets (*i.e.*, enable the simultaneous screening of a panel of DNA regions), thus enabling high- 25 throughput methods to be used for somatic mutation detection, (for example, massively parallel sequencing, MPS). NaME is particularly useful in the field of circulating biomarkers for cancer applications, pre-natal diagnostic applications, and infectious disease applications.

30 'Wild type target sequence' refers to a nucleic acid that is more prevalent in a nucleic acid sample than a corresponding target sequence (*e.g.*, same region of gene but different nucleic acid sequence). The wild type sequence makes-up over 50% of the total wild type sequence + mutant target sequence in a sample. The wild type sequence can be expressed at the RNA and/or DNA level 10X, 15X, 20X, 25X, 30X, 35X, 40X, 45X,

50X, 60X, 70X, 80X, 90X 100X, 150X, 200X or more than the target sequence. For example, a sample (*e.g.*, blood sample) may contain numerous normal cells and few cancerous cells. The normal cells contain wild-type alleles (non-mutant) sequences, while the small number of cancerous cells contain target sequences. As used herein, a  
5 ‘wild type strand’ refers to a single nucleic acid strand of a wild type sequence. The term ‘wild-type’ typically refers to the most common polynucleotide sequence or allele for a certain gene in a population. Generally, the wild-type allele will be obtained from normal cells.

10 The wild type sequence is about 13-2000 nucleotides long. In some embodiments, the wild type sequence is 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900 or more nucleotides long. Wild type sequences will share at least 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homology to the corresponding target sequence, but will differ by at least one nucleotide from the target  
15 sequence. In some embodiments, the at least one nucleotide is a methylated cytosine. In some embodiments, the at least one nucleotide is an unmethylated cytosine. Wild type sequences according to the invention can be amplified by PCR with the same pair of primers as that used for the mutant sequence.

20 A ‘target nucleic acid’ or ‘target sequence’, used interchangeably herein, refers to a nucleic acid that is less prevalent in a nucleic acid sample than a corresponding wild type sequence. The target sequence makes-up less than 50% of the total amount of wild type sequence+target sequence in a sample. Preferably the target sequence is expressed at the RNA and/or DNA level 1:10, 1:15, 1:20, 1:25 $\times$ , 1:30, 1:35, 1:40, 1:45, 1:50, 1:60, 1:70, 1:80, 1:90, 1:100, 1:150, 1:200 $\times$  or less than the wild type sequence. In some  
25 embodiments, the target sequence is a mutant allele. For example, a sample (*e.g.*, blood sample) may contain numerous normal cells and few cancerous cells. The normal cells contain wild-type (*i.e.*, non-mutant) sequences, while the small number of cancerous cells contain target mutant sequences. In some embodiments, the target sequence is repeat sequences that occur at large numbers in human genome (including but not limited  
30 to ALU elements, LINE elements, SINE elements, di-nucleotide repeats, tri-nucleotide repeats). Altered repeat sequences occur often in diseased states and application of the present invention in detecting alterations in repeat sequences are of interest. In some embodiments, the methods described herein are directed to detecting fetal DNA in a

5 nucleic acid sample obtained from a mother. In this embodiment, the fetal DNA is the target sequence while the more prevalent mother DNA is the wild type sequence. In some embodiments, the target sequence is a methylated allele. In some embodiments, the target sequence is an unmethylated allele. As used herein, a “target strand” refers to a single nucleic acid strand of a target sequence.

In some embodiments, the target sequence is 13-2000 nucleotides long. In some embodiments, the target sequence is 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900 or more nucleotides long. Target sequences share at least 50%, 60%, 70%, 80%, 85%, 90%, 10 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homology to the corresponding wild type sequence, but differs by at least one nucleotide from the wild type sequence. In some embodiments, the at least one nucleotide is a methylated cytosine. In some embodiments, the at least one nucleotide is an unmethylated cytosine. Target sequences according to the invention can be amplified via PCR with the same pair 15 of primers as those used for the wild type sequence.

‘Target mutant sequence’ or ‘mutant target sequence’ refers to a nucleic acid that is less prevalent in a nucleic acid sample than a corresponding wild type sequence. The target mutant sequence typically makes-up less than 50% of the total amount of wild type sequence + mutant sequence in a sample. The target mutant sequence may be 20 expressed at the RNA and/or DNA level 1 :10, 1 :15, 1 :20, 1 :25X, 1 :30, 1 :35, 1 :40, 1 :45, 1 :50, 1 :60, 1 :70, 1 :80, 1 :90, 1 :100, 1 :150, 1 :200X or less than the wild type sequence. For example, a sample (*e.g.*, blood sample) may contain numerous normal cells and few cancerous cells. The normal cells contain wild-type (non-mutant) alleles, while the small number of cancerous cells contain target mutant sequences. In some 25 embodiments, the invention is directed to detecting fetal DNA in a nucleic acid sample obtained from a mother. In this embodiment, the target mutant sequence is the fetal DNA while the more prevalent mother DNA is the wild type sequence. As used herein, a target mutant sequence is meant to include fetal DNA obtained from a pregnant mother. In some embodiments, the present disclosure is directed to detecting one or 30 more methylated alleles in the presence of a large excess of unmethylated alleles, or vice versa in epigenetic analysis.

The target mutant sequence is about 13-2000 nucleotides long. In some embodiments, the target mutant sequence is 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120,

130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900 or more nucleotides long. Target mutant sequences share at least 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homology to the corresponding wild type sequence, but differs by at least one nucleotide from the wild

5 type sequence. Mutant target sequences according to the invention can be amplified via PCR with the same pair of primers as those used for the wild type sequence.

The term ‘mutant’ refers to a nucleotide change (*i.e.*, a single or multiple nucleotide substitution, deletion, insertion, or methylation, or alteration in the number of poly-nucleotide repeats) in a nucleic acid sequence. A nucleic acid which bears a

10 mutation has a nucleic acid sequence (mutant allele) that is different in sequence from that of the corresponding wild- type sequence. The methods described herein are especially useful in preferentially cleaving wild type sequences, thereby allowing for selective enrichment of several or numerous mutant target sequences simultaneously. The mutant alleles can contain between 1 and 500 nucleotide sequence changes. A

15 mutant allele may have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500 nucleotide sequence changes compared to a corresponding wild-type allele. Typically, a mutant allele will contain between 1 and 10 nucleotide sequence changes, and more typically between 1 and 5 nucleotide sequence changes. The mutant allele will have 50%, 60%, 70%, 80%,

20 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homology to the wild-type allele. Generally, the mutant allele will be obtained from diseased tissues or cells and is associated with a disease state.

As used herein, a ‘region of interest’ is a sequence that will be interrogated for variations such as clinically relevant mutations, and methylation/unmethylation patterns.

25 ‘Enriching a mutant target sequence’ refers to increasing the amount of a mutant target sequence and/or increasing the ratio of mutant target sequence relative to the corresponding wild type sequence in a sample. For example, where the ratio of mutant sequence to wild type sequence is initially 5% to 95% in a sample, the mutant sequence may be preferentially amplified in an amplification reaction so as to produce a ratio of

30 70% mutant sequence to 30% wild type sequence. Thus, there is a 14 fold enrichment of the mutant sequence relative to the wild type sequence in this hypothetical example. Generally, enrichment of a mutant target sequence results in a 2X to 200X increase in the mutant target sequence relative to the wild type sequence prior to enrichment. The

enrichment of the mutant target sequence is at least a 2X, 3X, 4X, 5X, 6X, 7X, 8X, 9X, 10X, 15X, 20X, 25X, 30X, 35X, 40X, 45X, 50X, 60X, 70X, 80X, 90X 100X, 150X, 200X or more fold enrichment. Enrichment of a mutant target sequence results in a sample having 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%,

- 5 70%, 80%, 90%, 95% or more, mutant target sequence compared to wild type sequence (e.g., 10% mutant target sequence : 90% wild type sequence to 95% mutant target sequence : 5% wild type sequence).

‘Allele’ refers to alternative forms of a gene, portion thereof or non-coding region of DNA that occupy the same locus or position on homologous chromosomes that have 10 at least one difference in the nucleotide sequence. The term allele can be used to describe DNA from any organism including but not limited to bacteria, viruses, fungi, protozoa, molds, yeasts, plants, humans, non-humans, animals, and archaebacteria. The alleles may be found in a single cell (e.g., two alleles, one inherited from the father and one from the mother) or within a population of cells (e.g., a wild-type allele from normal 15 tissue and a somatic mutant allele from diseased tissue).

An allele can be 13-2000 nucleotides long. In one embodiment the allele is 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900 or more nucleotides long. Alleles will generally share 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 20 97%, 98%, 99% or more homology to each other. Alleles according to the invention can be amplified by PCR with the same pair of primers.

In one embodiment, the methods described herein are used to enrich a polymorphism. Any given gene may have none, one, or many allelic forms (polymorphism). Common mutational changes which give rise to alleles may be the 25 result of natural or artificial (e.g., chemical carcinogens) deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

As used herein the term ‘melting temperature’ or ‘Tm’ refers to the temperature at which a polynucleotide dissociates from its complementary sequence. Generally, the 30 Tm may be defined as the temperature at which one-half of the Watson-Crick base pairs in a duplex nucleic acid molecule are broken or dissociated (*i.e.*, are ‘melted’) while the other half of the Watson-Crick base pairs remain intact in a double stranded conformation. In other words, the Tm is defined as the temperature at which 50% of the

nucleotides of two complementary sequences are annealed (double strands) and 50% of the nucleotides are denatured (single strands). Tm, therefore defines a midpoint in the transition from double-stranded to single- stranded nucleic acid molecules (or, conversely, in the transition from single-stranded to double-stranded nucleic acid 5 molecules).

The Tm can be estimated by a number of methods, for example by a nearest-neighbor calculation as per Wetmur 1991 (Wetmur, J. G. 1991. DNA probes: applications of the principles of nucleic acid hybridization. Crit Rev Biochem MoI Biol 26: 227-259, hereby incorporated by reference) and by commercial programs including 10 Oligo™ Primer Design and programs available on the internet. Alternatively, the Tm can be determined though actual experimentation. For example, double-stranded DNA binding or intercalating dyes, such as ethidium bromide or SYBR-green (Molecular Probes) can be used in a melting curve assay to determine the actual Tm of the nucleic acid. Additional methods for determining the Tm of a nucleic acid are well known in 15 the art and described herein.

As used herein, ‘reaction mixture’ refers to a mixture of constituents including but not limited to a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation, and a pair of oligonucleotide probes that are complimentary to top and bottom strands of the 20 wild type nucleic acid. The reaction mixture can also include reagents, such as, but not limited to, salt(s), buffer(s), and enzyme(s) such as double strand-specific nuclease (DSN), exonuclease, and polymerase.

As used herein, a nucleic acid sample refers to any substance containing or presumed to contain a nucleic acid of interest (target and wild type sequences) or which 25 is itself a nucleic acid containing or presumed to contain a target nucleic acid of interest. The term “nucleic acid sample” thus includes a sample of nucleic acid (genomic DNA, cDNA, RNA), cell, organism, tissue, or fluid, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, synovial fluid, urine, tears, stool, external secretions of the skin, respiratory, intestinal and genitourinary tracts, saliva, blood cells, 30 tumors, organs, tissue, samples of *in vitro* cell culture constituents, natural isolates (such as drinking water, seawater, solid materials), microbial specimens, and objects or specimens that have been “marked” with nucleic acid tracer molecules. The nucleic acid sample may be obtained from mammals, viruses, bacteria or plants. In some

embodiments, the nucleic acid sample is DNA circulating in plasma, urine or other bodily fluids.

As used herein “oligonucleotide probes” refer to molecules comprising two or more deoxyribonucleotides or ribonucleotides. The methods described herein utilize a 5 pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand. By “complimentary” it is meant that the probes hybridize without any mismatches to the sequences in the top and bottom stands of the wild type nucleic acid. The oligonucleotide probes are non-identical, *i.e.*, the sequences of the two probes are 10 different from each other. In some embodiments, the probes do not overlap each other, *i.e.*, they do not bind to each other. At least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation, *i.e.*, the probe hybridizes to the sequence on the target nucleic acid containing the suspected mutation with at least one mismatch thereby forming a “partially complimentary” target mutant -probe duplex.

15 In some embodiments, one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation. Thus, the probes hybridize respectively to the top and bottom sequences on the target nucleic acid containing the suspected mutation with at least one mismatch. In such embodiments, the 20 probes partially overlap each other. However, they do not bind substantially to each other.

25 The oligonucleotide probes may be anywhere between 5 and 100 bases long. In some embodiments, the probes are 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 bases long. In some embodiments, the probes are in a molar excess of 100-fold- to 1 billion-fold compared to the wild type and target nucleic acids (*e.g.*, 100-fold, 500-fold, 1000-fold, 10,000-fold, 50,000-fold, 100,000-fold, 500,000-fold, 1 million-fold, 500-million-fold, 100 million-fold, 1 billion-fold in excess as compared to the wild type and target nucleic acids). In some embodiments the probes are in a molar concentration of 1 $\mu$ M, 10 $\mu$ M, 50 $\mu$ M, 100 $\mu$ M, 200 $\mu$ M, 300 $\mu$ M, 400 $\mu$ M, 500 $\mu$ M, 600 $\mu$ M, 700 $\mu$ M, 800 $\mu$ M, 30 900 $\mu$ M, or 1,000 $\mu$ M in the reaction.

By “selectively cleaved” or “preferentially cleaved” is meant that the subject methods preferentially cut, *i.e.*, cleave, deoxyribonucleic acid molecules present in perfectly matched double-stranded nucleic acids, *e.g.*, DNA-DNA duplexes and DNA-

RNA duplexes. Perfectly matched double-stranded nucleic acids are hybrid structures between complementary strands where no mismatches are present, as compared to partially complimentary nucleic acid duplexes of the same length. Thus, in the methods described herein complimentary DNA containing duplex nucleic acids (*i.e.*, without any 5 mismatches) are cleaved to a much greater extent than partially complimentary nucleic acid duplexes (*i.e.*, with one or more mismatches), non-DNA containing nucleic acid duplexes and/or single-stranded nucleic acids. In other words, the subject methods are able to cleave or cut perfectly matched nucleic acids duplexes in a sample at a much greater rate than other nucleic acid molecules that may be present in the sample being 10 treated, where the rate of perfectly matched nucleic acids duplex cleavage is typically at least 5 fold, at least 10 fold, at least 50 fold, or at least 100 fold greater than the rate of cleavage of other nucleic acids that may be present in the sample being treated.

15 As used herein, “primers” refers to oligonucleotides that anneal to opposite strands of a mutant target and wild type sequence so as to form an amplification product during a PCR reaction.

NaME on double stranded DNA

Accordingly, some aspects of the disclosure provide methods for preparing a 20 target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid. The subsequent enrichment can be achieved by applying amplification conditions to the produced reaction mixture.

In some embodiments, the method comprises subjecting a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a condition that destabilizes the double 25 stranded wild type and target mutant nucleic acids; contacting the destabilized double stranded wild type and target mutant nucleic acids with a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids 30 thereby forming complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; and exposing the complimentary wild-type-probe duplexes and the partially

complimentary target mutant-probe duplexes to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

In some embodiments, the method comprises exposing a nucleic acid sample 5 comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a 10 sequence on the target nucleic acid containing the suspected mutation; and subjecting the reaction mixture to a condition that destabilizes the double stranded wild type and target mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild-type- probe duplexes on top and bottom strands, and partially 15 complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

NaME utilizes nucleases (DNases) that show a strong preference for cleaving double stranded DNA versus single stranded DNA or RNA. DNases that can be used in 20 the methods described herein include, but are not limited to, native shrimp dsDNase, recombinant shrimp dsDNase, King crab nuclease (DSN) and bovine DNase I. NaME takes advantage of the DSN properties to cleave specific sequences from both top and bottom DNA strands of wild-type DNA as shown on Fig. 1A. In contrast, mutation-containing DNA is not cleaved or cleaved to a significantly less extent than wild type 25 DNA. Hence, a subsequent PCR reaction after DSN digestion amplifies preferentially the mutant alleles that remain substantially intact, and leads to enrichment of mutant versus wild type alleles.

For the purposes of the present disclosure, the term “double-strand specific nuclelease” or “DSN” includes DNA/RNA guided enzymes which have preferential 30 activity on double-stranded DNA , as compared to single stranded DNA. Examples of such enzymes that can be employed in conjunction with NaME include the RNA-guided Cas9 enzymes (Gu et al, *Depletion of Abundant Sequences by Hybridization (DASH): Using Cas9 to remove unwanted high-abundance species in sequencing libraries and*

*molecular counting applications* Genome Biology 2016; 17, 41), or the Argonaute DNA-guided enzymes (Gao et al, *DNA-guided genome editing using the Natronobacterium gregoryi Argonaute*, Nature Biotechnology May 2016 advanced online publication). These DNA/RNA guided enzymes digest DNA with high preference 5 when the probe ('guide oligonucleotide') is fully matched to the target DNA, and less so when there is a mismatch. By employing probes targeting both top and bottom DNA strands in an overlapping fashion as described in the present invention, NAME can be applied with DNA/RNA guided enzymes, in the same manner as when using other DSN nucleases described herein.

10 During NaME, (Fig. 1A) DSN and a pair of oligonucleotide probes that match the top and bottom strands of the wild type nucleic acid of interest are added to (*i.e.*, exposed or contacted with) a nucleic acid sample comprising double-stranded wild type nucleic acid and double-stranded target nucleic acid suspected of containing a mutation to create a reaction mixture. The nucleic acid sample is exposed to the DSN and the 15 oligonucleotide probes at a low temperature at which the DSN is inactive (*e.g.*, 4°C). At least one of the oligonucleotide probes overlaps sequences on the target nucleic acid that are suspected of containing clinically important mutations (*e.g.* KRAS codon 12/13 sequences; p53 sequences; tri-nucleotide repeat sequences; etc.). The second oligonucleotide binds the opposite target nucleic acid strand from the first 20 oligonucleotide probe, and can have similar length as the first oligonucleotide. In some embodiments, this second probe is designed to match a sequence on the target nucleic acid that normally does not contain mutations. In some embodiments, the probes are in a molar excess of 100-fold, 500-fold, 1000-fold, 10,000-fold, 50,000-fold, 100,000-fold, 25 500,000-fold, 1 million-fold, 500-million fold, 100 million-fold, 1 billion-fold compared to the wild type and target nucleic acids.

30 The reaction mixture is then subjected to a condition that destabilizes the double stranded wild type and mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and mutant nucleic acids thereby forming complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary mutant-probe duplexes. By "destabilizing" it is meant that the double stranded wild type and target mutant nucleic acids denature to such an extent so as to allow the probes to hybridize to their corresponding sequences, but the wild type and target mutant nucleic acids do not denature completely. A condition that destabilizes the

double stranded wild type and mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and mutant nucleic acids include addition of an organic solvent such as, but not limited to DMSO, betaine or formamide and/or an increase in temperature combined with a thermostable DSN. The increase in 5 temperature is such that it enables specific probe hybridization to its corresponding sequence. The temperature of the reaction mixture is raised to a temperature that destabilizes the double stranded structure (*e.g.*, 65°C - 80°C including 65°C, 70°C, 75°C, 80°C) but does not denature it completely. This destabilizing temperature is typically about 10-20°C below the melting temperature (Tm) of the nucleic acid sequence. At this 10 temperature, the oligonucleotide probes invade and bind to their corresponding sequences on the wild type and mutant nucleic acids. The probes fully match the sequences on the wild type nucleic acid and can, thus, form complimentary wild type-probe duplexes (*i.e.*, with no mis-matches).

If a suspected mutation is present on the target nucleic acid, the binding between 15 the probe and the target nucleic acid is inefficient and results in partially complimentary mutant-probe duplexes (*i.e.*, with mis-matches). The complimentary wild type-probe duplexes are recognized and cleaved by the DSN enzyme. In contrast, the partially complimentary mutant-probe duplexes remains substantially intact.

In some embodiments, one of the oligonucleotide probes overlaps a sequence on 20 the target nucleic acid that is suspected of containing a mutation while the second probe is designed to match a sequence at a different position on the target nucleic acid that normally does not contain mutations (Fig 1A). Hence, the approach shown on Fig 1A typically leads to cleavage of both strands for wild type nucleic acid while only a single DNA strand of the mutant nucleic acid is cleaved.

25 In some embodiments, the methods described herein are performed by first destabilizing the double-stranded wild type nucleic acid and the double-stranded target nucleic acid suspected of containing a mutation. The destabilized wild type nucleic acid and the target mutant nucleic acid are then contacted with the oligonucleotide probes to permit hybridization of the probes to their corresponding sequences on the wild type and 30 target mutant nucleic acids thereby forming complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes. By “contacting” it is meant that the probes are added to the nucleic acids and the components are mixed, or the nucleic acids are added to the probes and the components

are mixed. The duplexes are then exposed to DSN which preferentially cuts the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

Some aspects of the disclosure provide methods for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; reducing the temperature of the reaction mixture to permit formation of complimentary wild type – probe duplexes on top and bottom strands and partially complimentary target mutant-probe duplexes; and exposing the reaction mixture to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

In these methods, the double stranded wild type and target nucleic acids in the presence of the two probes are first denatured by subjecting the reaction mixture to a denaturing temperature, while also optionally including organic solvents like DMSO, betaine or formamide. The denaturing temperature should be sufficiently high so as to allow the full denaturation of the wild type and target nucleic acids (e.g., 75° C, 80° C, 85° C, 90° C, or 95° C). In some embodiments, the denaturing temperature is about 1° C to 30° C above the Tm of the wild type and nucleic acid sequence (e.g., 1° C, 5° C, 10° C, 15° C, 20° C, 25° C, 30° C above the Tm of the wild type and nucleic acid sequence).

Next the temperature of the reaction mixture is decreased allowing the wild type and target nucleic acids to hybridize with the oligonucleotide probes to form complimentary wild type – probe duplexes on top and bottom strands (*i.e.*, with no mis-matches) and partially complimentary mutant-probe duplexes (*i.e.*, with mis-matches). In some embodiments, this hybridization temperature is 40° C, 45° C, 50° C, 55° C, 60° C, 65° C, 70° C, or 75° C). At this hybridization temperature, since the two probes are in high excess relative to the target nucleic acid, they bind first to their respective targets,

5 *i.e.*, while the two parent strands of the wild type and target nucleic acids have not yet re-associated and remain substantially single-stranded. In some embodiments, the probes are in a molar excess of 100-fold, 500-fold, 1000-fold, 10,000-fold, 50,000-fold, 100,000-fold, 500,000-fold, 1 million-fold, 500-million fold, 100 million-fold, 1 billion-fold compared to the wild type and target nucleic acids.

10 In this method, DSN is, optionally, not added from the beginning in order to avoid partial or total inactivation of the DSN at the denaturing temperature. DSN is added once the temperature is reduced to allow formation of complimentary wild type – probe duplexes on top and bottom strands and partially complimentary mutant-probe duplexes. DSN then preferentially degrades the complimentary wild type – probe duplexes, while the partially complimentary mutant-probe duplexes remains substantially intact. The DSN activity can then be stopped, for example, by heating the sample to 95°C for 1-10 min to inactivate the DSN. A subsequent PCR reaction amplifies 15 preferentially the mutated alleles that remain substantially intact, while the DSN-digested wild type alleles do not amplify. Fig. 1B demonstrates quantification of the fractional mutation abundance following this PCR reaction. It can be seen that in the absence of DSN and/or both probes the mutation abundance is low (4-6%) while in the presence of DSN and both probes the resulting mutation abundance is 37-38%, *i.e.* demonstrating the enrichment of the mutated alleles using NaME.

20 In some embodiments, the subsequent PCR reaction amplifies the probes rather than the hybridized nucleic acid. In this approach, a purification step is applied following probe binding to top-and-bottom DNA target strands, either before or after DSN cleavage, to remove excess unbound probes (e.g., using a DNA affinity column, such as QIAquick® PCR purification kit commercially available from QIAGEN). Then 25 following DSN cleavage the uncut probes are amplified (instead of amplifying the target DNA) and identified/quantified. Since probes that bind WT DNA will have been selectively digested by DSN, the presence of any given probe after amplification indicates a mutation under the region covered by this probe.

30 **NAME using OVERLAPPING PROBES**

In some embodiments of the methods described herein, the probes are constructed such that one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on

the bottom strand of the target nucleic acid containing the mutation of interest and the two probes partially overlap each other (Figs. 2 and 3). The two probes overlap only partially, so that they do not bind substantially to each other in solution at the temperatures used for NaME during DSN digestion (e.g., 65°C -70°C). Accordingly, it  
5 is important during probe hybridization to their corresponding sequences to retain a temperature low enough for probe binding to the template, but high enough so that it does not allow substantial probe-to-probe binding. This approach increases the specificity of the process for mutated sequences, and the mutation enrichment becomes much more pronounced than when using only one mutation-specific probe with a second  
10 probe which matches the wild-type nucleic acid (Fig 1A).

In some embodiments, a thermostable DSN, such as but not limited to, king crab nuclease is used in the methods described herein. In some embodiments, a non-thermostable DSN, such as but not limited to, native shrimp dsDNase, recombinant shrimp dsDNase, and bovine DNase I is used in the methods described herein. If a non-thermostable DSN is used during NaME, then the overlap between the probes designed  
15 must be such that at the temperature used for probe-nucleic acid duplex formation (e.g., 37-45°C there is minimal binding of the probes to each other, while they still bind specifically to genomic DNA targets). One way to reduce the Tm of the probes to match the optimal temperature of the nuclease used is to add organic solvents (DMSO,  
20 formamide) that lower the Tm of the probes. For example, instead of probes with a Tm of 65°C matching the optimal temperature of thermostable DSN enzyme, one may use shrimp nuclease in the presence of 10% DMSO which reduces the probe Tm as well as the Tm of probe-probe overlap regions.

For the purposes of the present disclosure, the term “double-strand specific  
25 nuclease” or “DSN” includes DNA/RNA guided enzymes which have preferential activity on double-stranded DNA, as opposed to single stranded DNA. Examples of such enzymes that can be employed in conjunction with NaME include the RNA-guided Cas9 enzymes (Gu et al, *Depletion of Abundant Sequences by Hybridization (DASH): Using Cas9 to remove unwanted high-abundance species in sequencing libraries and 30 molecular counting applications* Genome Biology 2016; 17, 41), or the Argonaute DNA-guided enzymes (Gao et al, *DNA-guided genome editing using the Natronobacterium gregoryi Argonaute*, Nature Biotechnology May 2016 advanced online publication). These DNA/RNA guided enzymes digest DNA with high preference

when the probe ('guide oligonucleotide') is fully matched to the target DNA, and less so when there is a mismatch. By employing probes targeting both top and bottom DNA strands in an overlapping fashion as described in the present invention, NAME can be applied with DNA/RNA guided enzymes, in the same manner as when using other DSN nucleases described herein.

In some embodiments, prior to implementing the methods described herein, the nucleic acid sample is subjected to an amplification condition. In some embodiments, the methods described herein further comprise enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture containing 5 cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid. Any known amplification condition can be used. In some embodiments, the amplification condition is selected from the group 10 consisting of: PCR, ligation mediated PCR using common ligated adaptors, multiplex PCR, using multiple pairs of primers, PCR of repeat elements using primers specific for ALU, LINE 1, poly-nucleotide repeats, micro-satellites and other repeat elements spread over the genome, arbitrarily-primed PCR (AP-PCR) and isothermal amplification (such 15 as but not limited to displacement amplification based on phi-29 based; or Loop Mediated LAMP amplification; or any other isothermal mode of amplification). The 20 wild type nucleic acid will not amplify during this amplification step since it was cleaved selectively by DSN. The mutation-enriched amplified product can then be analyzed for mutations using any available method such as MALDI-TOF, HR-Melting, Di-deoxy-sequencing, Single-molecule sequencing, massively parallel sequencing (MPS), pyrosequencing, single strand conformational polymorphism SSCP, restriction fragment 25 length polymorphism RFLP, denaturing high precision liquid chromatography dHPLC, chemical cleavage of mismatches CCM, capillary electrophoresis, digital PCR and quantitative-PCR.

The probes used in the methods described herein preferably contain a 3'-block to 30 polymerase extension, so that if the NaME -reaction product is subsequently amplified there is no interference of the probes with the amplification reaction. A 3'-polymerase block can comprise a simple phosphate; or abasic site; or any other modification that prevents polymerase synthesis past the block. In addition, for added discrimination of wild type versus mutant sequences during NaME, in some embodiments, each probe

comprises a locked nucleic acid (LNA), peptide nucleic acid (PNA), xeno nucleic acid (XNA), nucleic acid with any known natural or modified base such as dITP or 2,6-diaminopurine dATP or RNA that increases the destabilization caused by a mutation-induced mismatch between the oligonucleotide probe and its target nucleic acid.

5 In some cases part of the probes used in the methods provided herein can comprise one or more random nucleotides, so that the probe can be directed against a plurality of DNA targets. For example, a probe can include a core region of one or more nucleotides which are complimentary to the wild-type nucleic acid sequence in a region of interest. e.g., a suspected mutation site. Any one or more of the remaining nucleotides 10 in the probe may be selected randomly from any or all possible nucleotides. The probe containing the core region plus one or more random nucleotides can form a duplex with a fully complimentary wild-type nucleic acid sequence which also contains the core region. A cleavage enzyme, e.g., DSN, can be used to cleave the complimentary wild-type probe duplexes, but not the partially complimentary target mutant-probe duplexes.

15 In some embodiments, the methods described herein are used to prepare two or more different target mutant nucleic acids for subsequent enrichment relative to corresponding wild type nucleic acids. In such embodiments, one or more additional pairs of probes directed to the different wild type nucleic acids are used. For each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand, while 20 the other is complimentary to the wild type nucleic acid bottom strand.

25 In all embodiments described above, it is also understood that the concentration of probes for the top and bottom DNA strands does not necessarily need be the same. Thus one may combine a high concentration of probe for the bottom strand and a low concentration of probe for the top strand, or vice versa. Probe concentrations can also be different for each DNA target when many targets are simultaneously enriched. Optimized concentrations depending on sequence context, local sequence Tm and mutation being targeted can be applied. Furthermore a subsequent PCR amplification reaction following application of DSN can utilize equal amounts of primers or different amounts of primers for each DNA strand (asymmetric PCR, or Linear After The 30 Exponential, L.A.T.E PCR).

NAME applied directly on genomic DNA

In some embodiments, the methods described herein are performed directly on genomic DNA. The genomic DNA can optionally be fragmented prior to application of NaME (Fig. 4). In this approach, NaME is applied by (a) fragmenting the genomic DNA 5 using enzymatic or physical means; (b) adding overlapping (or non-overlapping) probes that address both top and bottom DNA strands and optionally denaturing both the wild type and target nucleic acids (for example, at 95°C); (c) reducing the temperature to, for example, 60-70°C to enable probes to find their respective targets prior to substantial renaturation of the parent DNA strands, and keeping the temperature high enough to 10 minimize probe-to-probe interactions; (d) adding DSN to selectively cleave one or more (multiple) complimentary wild type-probe duplexes DNA targets while leaving the partially complimentary target mutant –probe duplexes substantially intact. The resulting reaction mixture with cleaved wild type-probe duplexes and uncleaved target 15 mutant nucleic acids can be amplified using methods known in the art, such as but not limited to, PCR, COLD-PCR, ligation-mediated PCR or COLD-PCR using common ligated adaptors, multiplex-PCR or isothermal amplification (such as phi-29 based; or LAMP-based; or any other isothermal mode of amplification) thereby enriching the target mutant nucleic acid as compared to the wild type. This amplified product can be 20 examined for mutations using any available method, such as but not limited to, MALDI-TOF, HR-Melting, Di-deoxy-sequencing, Single-molecule sequencing, massively parallel sequencing (MPS), pyrosequencing, SSCP, RFLP, dHPLC, CCM, digital PCR and quantitative-PCR (the wild type nucleic acid will not amplify during this amplification step since it was selectively cleaved by DSN).

Fig. 5 depicts the enrichment of an original ~1% mutation to an 83% mutation 25 and a 0.5% mutation to 14% mutation, following application of NaME directly to genomic DNA. Fig. 6 shows a duplex application of NaME on two mutated targets simultaneously, in KRAS and TP53 genes. Fig 7A shows application of NaME using two overlapping probes covering 3 different mutations in codons 12 and 13 of Kras, and indicating that any mutation under the probes will be enriched during NaME. And Fig. 30 7B demonstrates simultaneous enrichment of 11 different targets when NaME is applied directly from genomic DNA. For each target, a separate pair of overlapping probes was designed, and all probes are included in a single reaction during NaME.

In some embodiments the probes used are directed against poly-nucleotide repeats that are widespread around the genome, so that multiple targets are addressed via NaME simultaneously, using a single pair of probes. For example, if the target is a poly-A-containing sequence, two overlapping probes are used, one for bottom strand 5 containing poly-T and one for top strand containing poly-A in this case. To increase the length of the probe, one may also add an optional number of inosines that can generically bind to neighboring bases.

Combination of NaME with Massively Parallel Sequencing (MPS).

10 In some embodiments, the methods described herein may be used in conjunction with massively parallel sequencing (MPS). MPS is currently the most advanced approach for mutation identification. Sample ('library') preparation for MPS is a very important step prior to applying genome-wide or exome-wide sequencing or targeted re-sequencing. NaME provides a unique advantage for sample preparation prior to MPS, as 15 it can enrich predictably numerous targets for mutations, thereby enabling MPS to identify easily mutations that are originally at very low abundance, without requiring an excessive number of sequence reads. This enables cost reduction and increased sensitivity and simplicity. Fig. 8 provides an example of NaME - enhanced sample preparation process prior to MPS. In this approach, multiplexed NaME (using overlapping probes against numerous gene mutations simultaneously as described in previous sections) is applied to the original starting material in order to cleave selectively 20 wild type nucleic acid on all targets of interest simultaneously. The sample is then amplified to enrich preferentially the mutated DNA targets. Finally, the resulting mutation-enriched DNA is used for routine library construction prior to MPS.

25 In some embodiments, multiplexed application of NaME on numerous targets of interest can be applied directly from denatured genomic DNA as described in Fig 4. Following this, a multiplexed PCR can be applied using primers addressing the DNA targets of interest (for example, but not limited to, the primers used in the Life- 30 Technologies Ampliseq kit, or the Illumina Trueseq kit.). The multiplexed PCR products will now be enriched for mutations in view of the NaME treatment, thus the resulting library preparation will provide mutation enriched DNA for targeted re-sequencing.

In some embodiments, prior to implementing the NaME method described herein, targets of interest are captured from genomic DNA within molecular inversion

probes (MIPS); or using the 'bait' oligonucleotides approach. In some embodiments, the MIPS or the bait oligonucleotides are biotinylated (e.g., similar to those included in the Agilent SureSelect™ kit, but re-designed to capture both top and bottom target DNA strands as per previous sections). In some embodiments, the bait oligonucleotides are  
5 attached to beads. The nucleic acid sample is contacted with the bait oligonucleotides that bind to selected targets of interest and binding of the bait oligonucleotides to the targets of interest is enabled. Next the bait oligonucleotides with the regions of interest bound thereto are isolated from the remaining nucleic acids. Finally, the isolated targets are released from beads and multiplexed NaME (using overlapping probes addressing  
10 both top and bottom DNA strands and annealing to numerous targeted gene mutations simultaneously) is used to cleave the different wild type nucleic acids simultaneously. Following this, PCR and library construction using the mutation-enriched sample can be used prior to MPS.

In some embodiments, the probe oligonucleotides that can be used in the methods  
15 described herein systematically tile a genomic region of interest, for example, chromosome Y. In some embodiments, degenerate oligonucleotide probes are synthesized that cover all AT-rich regions, all GC rich regions, gene promoters; or CpG islands. Any genomic fraction of interest can be targeted for selective cleavage 'at will' using multiple overlapping probes targeting both top and bottom strands and designed as  
20 described herein.

*Mutation enrichment using NaME in the absence of subsequent amplification.*

In all embodiments described thus far, in order to produce a sample with enriched mutated target sequences, an amplification step is conducted following application of  
25 DSN digestion of the wild type DNA alleles. Alternatively, another way to enrich the mutated target sequences is to eliminate the wild type sequences (without amplification). Thus, in some embodiments, the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids is subjected to a further DNA degradation condition which hydrolyzes enzymatically the DSN-cleaved wild type-probe duplexes,  
30 with the degradation initiated at the position of the cleavage. The "DNA degradation condition" includes contacting the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an exonuclease (e.g., using exo I or exo III or Klenow fragment of E.coli DNA polymerase I that digest DNA from the 3'-

end) under conditions of optimal exonuclease activity (that is, the temperature, pH ion concentrations *etc.* are maintained to provide optimal enzyme activity).

In this approach, genomic DNA can either be not fragmented, or it may be randomly fragmented as described in preceding sections. If the DNA is fragmented, the 5 fragmented genomic DNA is first ligated to adaptors that are resistant to 3' - exonuclease digestion (e.g., by using adaptors that have a 3'-terminal phosphorothioate linkage).

Next, the DNA sample is denatured and NaME is applied as described in previous sections to generate cleavage of wild type sequences while leaving intact the mutated sequences. Next, an exonuclease digestion is applied (*e.g.*, using exo I or exo III or 10 Klenow fragment of E.coli DNA polymerase I that digest DNA from the 3'-end).

Exonuclease will digest all sequences that do not have an exonuclease resistant 3'-end, *i.e.* without 3'-terminal phosphorothioate. Since DSN -nicked fragments do not have a 3'-terminal phosphorothioate, they will be fully digested by the enzyme, thereby eliminating wild type DNA strands. Digestion will proceed from the 3'-position of the 15 DSN-induced nick all the way to the 5'end, while leaving the (un-nicked) mutated target sequences intact. Optionally, one may also digest wild-type sequences from the 5'-end of the nick all the way to the 3'-end by using E.coli DNA polymerase I that has 5' to 3' exonuclease activity. Following the complete digestion of the DSN-nicked, wild type sequences, an endpoint detection method that does not require amplification, such as a 20 single molecule sequencing approach (Nanopore system; or Pacific-Bio system) can be used to sequence the mutation-enriched DNA sample. This embodiment that does not rely on any form of nucleic acid amplification and can be particularly useful in the sequencing of small genomes (*e.g.*, bacterial or viral genomes) where low level mutations are currently difficult to detect in view of the relatively high error rate of these 25 sequencing systems. The approach can be combined with selective capture of genomic fragments on beads, to reduce the complexity of a larger genome followed by NaME to eliminate wild type DNA and enrich mutated target sequences in the absence of amplification.

30 *Mutation scanning using NaME*

Often there is a need to scan for mutations in long DNA regions (as opposed to identifying known mutations at a single hotspot position, such as KRAS). Using two longer probes that are complementary to adjacent DNA sequences (one probe on bottom

strand and the second, adjacent probe, on the top strand) one can adapt NaME for 'mutation scanning'. For example, probes of 50-70 bp can be used to cover a region of 100-140 base pairs. If there is a mutation at any position along these 140 base pairs, the corresponding strand will not be substantially cleaved by DSN. Hence, the mutant strand 5 will survive and will lead to a subsequent PCR product that can be sequenced (Fig 12A). In this way, longer regions on tumor suppressor genes like p53 can be enriched for mutations irrespective of the mutation position (Fig. 12B and 12C).

NaME application with RNA or ssDNA

10 In some embodiments, the methods described herein can be used to selectively cleave wild type cDNA, or mRNA or DNA in single stranded format. For this approach (Fig. 13), only a single probe per targeted gene would be needed, as opposed to one probe on each opposite strand used in dsDNA.

15 Methylation-sensitive NaME (with probes)

20 In some embodiments, the methods described herein are used to prepare an unmethylated target nucleic acid of interest for subsequent enrichment. In such embodiments, prior to implementing NaME on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite (bisulfite converts all cytosines to uracils, unless the cytosines are methylated at CpG dinucleotide positions). The pair of oligonucleotide probes used in these methods are complimentary to top and bottom strands of the 25 methylated nucleic acid of interest, that is, one of the oligonucleotide probes is complimentary to top strand of the methylated nucleic acid of interest, while the other oligonucleotide probe is complimentary to the bottom strand of the methylated nucleic acid of interest following bisulfite conversion. The probes will, thus, form complimentary (*i.e.*, without any mismatches) duplexes with the top and bottom strands in alleles that contain fully methylated DNA. In contrast, the probes will form partially complimentary duplexes with the alleles containing unmethylated cytosines because of mismatches at the positions of uracils (which used to be cytosines before bisulfite 30 conversion). As a result, the methylated duplexes will be cleaved by DSN, while the unmethylated duplexes will remain substantially intact for subsequent amplification (Fig. 14).

Since NaME works well with mismatches caused by single point mutations, it can be expected that presence of several mismatches on a sequence due to conversion of multiple cytosines makes DSN match/mismatch discrimination work even better. Thus, one can optionally also use probes that are longer, (for example, 50-200 bp or longer) 5 with this approach. Furthermore, in contrast to regular double stranded DNA, bisulfite converted DNA remains single stranded after chemical treatment, and the two DNA strands are not complementary to each other any longer. Accordingly, one may optionally use probes matching only the top DNA strand, or matching only the bottom DNA strand following bisulfite conversion of DNA.

10 One can use thousands of probes covering all promoters and tiling entire genomic regions. For example, genomic DNA is digested into smaller fragments, using physical shearing for random fragmentation or restriction enzyme fragmentation (using enzymes that are methylation – independent). DNA is randomly fragmented, end repaired, and ligated to methylated adaptors that are resistant to bisulfite conversion. This is a standard 15 first step in whole genome bisulfite sequencing preparations. Next, the sample is treated with sodium bisulfite, to convert unmethylated C to U (Fig. 14). The DNA at this point comprises mostly single stranded sequences. Next, the NaME procedure is applied, by adding DSN plus a large set of synthesized oligonucleotide probes designed to match the 20 methylated bisulfite-converted version of the regions of interest (for example, an entire tumor suppressor gene like TP53 or BRCA1; or a large portion of chromosome 21 if trisomy 21 is under examination for pre-natal diagnostics; or all promoters in oncogenes; or regions that are differentially methylated among various tissues in order to assist 25 definition of the tissue of origin when examining circulating DNA or other liquid biopsies). The probes will form perfectly double stranded DNA (i.e., complimentary duplexes) in alleles that contained fully methylated DNA. Both top and bottom strands of the original DNA need to be addressed by the oligonucleotides used, as both parent DNA strands need to be selectively digested and prevented from subsequent 30 amplification. Alleles with unmethylated DNA will remain undigested, because the probes will contain mismatches at the positions of uracils (bisulfite converted cytosines). As a result, DSN will not cut these sequences, thereby allowing their subsequent amplification (Fig. 14). Following selective cleavage of methylated targets of interest, an amplification condition, such as PCR using the common adaptors, can be applied followed by sequencing of the sample. Alternatively one can apply PCR of repeat

elements using primers specific for bisulfite-treated ALU, LINE 1 and other repeat elements spread over the genome; or arbitrarily-primed PCR (AP-PCR); or COLD-PCR. Isothermal forms of amplification may also be used in the place of PCR. This approach is of relevance to many applications such as cancer diagnosis/therapy (for detecting 5 global hypomethylation), to prenatal diagnosis (*e.g.* for detecting placental DNA which contains fetal sequences that are substantially unmethylated), and to other diseases known to result/cause hypomethylation, such as systemic lupus erythmatosis.

In some embodiments, *e.g.*, following sodium bisulfite treatment, the method is applied in the opposite manner, that is, to prepare a methylated target nucleic acid of 10 interest for subsequent enrichment. In such embodiments, the pair of oligonucleotide probes used are fully complimentary to top and bottom strands of the unmethylated nucleic acid of interest, that is, one of the oligonucleotide probes is complimentary to top strand of the unmethylated nucleic acid of interest, while the other oligonucleotide probe is complimentary to the bottom strand of the unmethylated nucleic acid of interest 15 following its bisulfite conversion. This results in the preferential removal of the unmethylated regions of the targets of interest.

In some embodiments, *e.g.*, following bisulfite treatment, the method is used to prepare both an unmethylated target nucleic acid of interest and a (different) methylated target nucleic acid of interest for subsequent enrichment. The method comprises: (i) a 20 pair of oligonucleotide probes, one of which is complimentary to top strand of the methylated form of the unmethylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the methylated form of the unmethylated target nucleic acid of interest, (ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of the methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the 25 unmethylated form of the methylated target nucleic acid of interest; and wherein prior to implementing NaME protocol described herein on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite.

In some embodiments, the method is used to prepare multiple target nucleic acids 30 of interest, some of which are methylated target nucleic acids of interest, and some of which are unmethylated target nucleic acids of interest. In such embodiments, (i) a pair of oligonucleotide probes, one of which is complimentary to top strand of the methylated form of each unmethylated target nucleic acid of interest, while the other is

complimentary to the bottom strand of the methylated form of each unmethylated target nucleic acid of interest, and (ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of each methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the unmethylated form of each methylated target nucleic acid of interest are used. Prior to implementing NaME on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite. Thus, the methods described herein allow for the simultaneously removal of (a) unmethylated promoters in tumor suppressor genes (so that it becomes easy to reveal the methylated genes of interest), and (b) methylated promoters of oncogenes (so that it becomes easy to reveal the unmethylated genes of interest). In this way one can enrich for methylated oncogene promoters, as well as for unmethylated oncogene promoters, simultaneously.

Finally, similar approaches to those described above using bisulfite treatment of DNA to selectively enrich 5-methylcytosine (5mC) -based differentially methylated/unmethylated DNA regions may also be applied to selectively enrich different epigenetic DNA modifications of interest, such as 5-hydroxy-methylation (5hmC). 5-hydroxy-methylation is an epigenetic modification functionally and biologically different from 5-methylcytosine-modification. Thus it is important to measure separately these two modifications. One way to separate DNA containing 5hmC from that containing 5mC is TAB-seq (tet-assisted bisulfite sequencing, Ito S et al, Science 2011 333(6047):1300-1303; and Yu M et al, Cell 2012: 149:1368-80). In TAB-seq, genomic 5hmC is first protected with protected by glucosylation, prior to performing bisulfite conversion. The DNA is then treated with a Tet enzyme, converting 5mC to 5caC, while leaving the glycosylated 5-hydroxy-methylation untouched. Any C's read in the resulting sequencing are thus interpreted as 5-hydroxy-methylated. Accordingly, depending on whether bisulfite treatment is applied directly, OR following glucosylation, the present invention can be directed to enriching either 5-methylcytosine-containing DNA or 5-hydroxy-methylcytosine containing DNA.

### 30 Nuclease chain reaction

Some aspects of the disclosure relate to a method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid, using a 'chain reaction approach'. The method comprises the steps of: (a) exposing a nucleic acid

sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a thermostable double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type 5 nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; and (c) reducing the temperature to permit rapid hybridization of the probes to their corresponding sequences 10 on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes and but not the partially complimentary target mutant-probe duplexes.

More specifically, in some embodiments, NaME can be applied in a temperature 15 cycling fashion, including successive brief denaturation cycles followed by DSN incubation in the presence of probes (Fig. 9). The thermostable DSN is included in the reaction mixture from the beginning despite the use of denaturing temperature. The temperature is raised such that it allows denaturation of the wild type nucleic acid and the target mutant nucleic acid without destroying the DSN enzyme which is 20 simultaneously present in the reaction mixture. In some embodiments, this denaturing temperature is 65° C, 70° C, 75° C, 80° C, or 85° C. In some embodiments, an organic solvent that can lower the Tm of the nucleic acids is included in the reaction mixture. The solvent lowers the Tm of the nucleic acids, without inhibiting the activity of DSN. Examples of such solvents include, but are not limited to DMSO, betaine or formamide. 25 In some embodiments, 10-15% of DMSO is included in the reaction mixture.

After briefly denaturing the wild type and target nucleic acids, the temperature is lowered to permit hybridization of the probes to their corresponding sequences on the wild type and mutant nucleic acids. This hybridization temperature allows for DSN activity. In some embodiments, this hybridization temperature is 50° C, 55° C, 60° C, 30 65° C, or 70° C). Since the oligonucleotide probes are in high excess as compared to the wild type and target nucleic acids, they bind to their corresponding sequences and form complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary mutant-probe duplexes. The thermostable DSN then digests the wild

type-probe complimentary duplexes, while the partially complimentary mutant-probe duplexes remain intact.

In some embodiments, steps (b) (denaturing step) and (c) (hybridization/DSN incubation step) are repeated for one or more cycles. In some embodiments, these steps 5 are repeated for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, or 50 cycles. The hybridization/DSN incubation step is only applied intermittently (for example, 2-4 min) and is interrupted by another denaturation cycle, followed by lowering the temperature again for DSN digestion, and so on. In this way it is possible to maximize the difference between wild type and mutant nucleic acid cleavage, while still preventing substantial re-10 hybridization of the two parent nucleic acid strands (which would lead to indiscriminate destruction of the nucleic acid, whether mutant or not).

In some embodiments, the method further comprises enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an 15 amplification condition such as but not limited to PCR, COLD-PCR, ligation mediated PCR or COLD-PCR using common ligated adaptors, multiplex PCR, and isothermal amplification (such as but not limited to displacement amplification based on phi-29 based; or Loop Mediated LAMP amplification; or any other isothermal mode of amplification).

20 In some embodiments, one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other.

In some embodiments, the method is used to prepare two or more different target 25 mutant nucleic acids for subsequent enrichment relative to corresponding wild type nucleic acids and the method further comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand.

30

#### NaME-PCR

Some aspects of the present disclosure provide methods that combine the NaME process with PCR amplification in a single step as shown in Fig 10. Accordingly,

aspects of the disclosure provide a method for enriching a target mutant nucleic acid, the method comprising the steps of: (a) preparing an amplification reaction mixture comprising: a double-stranded wild type nucleic acid, a double-stranded target nucleic acid suspected of containing a mutation, a thermostable double strand-specific nuclease 5 (DSN), a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation and PCR amplification components; (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the 10 wild type nucleic acid and the target mutant nucleic acid; (c) reducing the temperature to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the 15 partially complimentary target mutant-probe duplexes; and (d) subjecting the reaction mixture to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid.

In this approach, NaME and PCR are performed in a single tube. All PCR components such as but not limited to primers, dNTPs, polymerase and polymerase 20 buffer are included together with DSN and oligonucleotide probes in the reaction mixture. In this manner, the wild type nucleic acid is successively selectively destroyed by DSN while also re-synthesized by PCR, so that the total target DNA remains the same or increases, while at the same time continuously enriching the mutated DNA during the cycling process. In some embodiments, the method is performed in COLD-PCR format, 25 instead of standard PCR.

DSN is compatible with most PCR buffers used commercially; hence DSN cleavage works in a PCR environment. Since both enzymes, DSN and polymerase, are thermostable it is possible to operate a combined reaction in a common thermocycler. The amplification condition applied in step (d) permits annealing of primer pairs to the 30 wild type and target mutant nucleic acids followed by extension thereby enriching the target mutant nucleic acid relative to wild type.

In some embodiments, steps (b) (denaturing step) and (c) (hybridization/DSN incubation step) are repeated for two or more cycles before executing step (d) (PCR

amplification). In some embodiments, these steps are repeated for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, or 50 cycles. In some embodiments, steps (b) (denaturing step), (c) (hybridization/DSN incubation step) and (d) (amplification step) are repeated for two or more cycles. In some embodiments, these steps are repeated for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, or 50 cycles.

5 In some embodiments, the reaction mixture further comprises an organic solvent that can lower the Tm of the nucleic acids is included in the reaction mixture. The solvent lowers the Tm of the nucleic acids, without inhibiting the activity of DSN. Examples of such solvents include, but are not limited to DMSO, betaine or formamide.

10 The denaturing temperature used is such that it allows denaturation of the wild type nucleic acid and the target mutant nucleic acid without destroying the DSN enzyme which is simultaneously present in the reaction mixture. In some embodiments, this denaturing temperature is 65° C, 70° C, 75° C, 80° C, or 85° C applied for time periods of 1 sec, 30 sec, 1 min, 2 min, 3 min, 4 min, 5 min, 6 min, 7 min, 8 min, 9 min, or 10 15 min.

20 In some embodiments, one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other. In some embodiments, the probes are modified at the 25 3' end to prevent polymerase extension and to ensure that the probes do not act as primers during NaME -PCR. In some embodiments, the primers used for PCR amplification have a melting temperature that is below the temperature applied in step (c). This ensures that the primers do not bind to the nucleic acids during the time DSN is used to selectively cleave wild type nucleic acid. For example, the Tm of the primers can 25 be 55°C while the hybridization/DSN incubation step is performed above 65°C where the primers do not interfere.

30 In some embodiments, the method is used to enrich two or more different target mutant nucleic acids relative to wild type nucleic acids and the method further comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand.

Figure 10 depicts a process of sequential DNA synthesis via PCR (during which DSN does not have enough time for substantial cleavage of the nucleic acids, while polymerase acts within seconds to synthesize templates) followed by arresting amplification for 10min at 65°C during which DSN acts selectively on wild type nucleic acid. At this temperature, primers do not bind, and hence polymerase synthesis does not proceed. This is then followed by more DNA synthesis via PCR and more DSN digestion and so on in a sequential process. The end product is DNA comprising mainly mutant DNA. The product can be directly sequenced or analyzed for mutations using other known methods.

Figure 11 depicts combination of NaME with COLD-PCR (COLD- NaME -PCR) for even greater enrichment of mutation containing nucleic acid. Conditions for COLD-PCR are applied that enable selective amplification of mutation containing nucleic acid during PCR, followed by selective cleavage of wild type nucleic acid using DSN in sequential single tube reactions. All types of COLD-PCR can be used, including but not limited to, full COLD, fast COLD, ICE-COLD PCR or Limited Denaturation Time LDT-COLD-PCR. Methods to perform COLD-PCR are highly compatible with NaME -PCR and have been described previously (see, for example, Li J, Wang L, Mamon H, Kulke MH, Berbeco R, Makrigiorgos GM. Nat Med 2008;14:579-84; Milbury CA, Li J, Makrigiorgos GM. Nucleic Acids Res;39:e2; Murphy DM, Castellanos-Rizaldos E, Makrigiorgos GM. Clin Chem. 2014 60:1014-6; the contents of which are incorporated by reference herein).

#### *Methylation-sensitive NaME - No probes*

Some aspects of the disclosure provide methods for preparing unmethylated or methylated nucleic acids of interest using temperature-based preferential enrichment of the alleles of interest on a genome wide level using enzymes such as DSN or exonuclease. In this approach, no probes need to be used. Genomic DNA is digested to smaller fragments. Following end repair and bisulfite-resistant adaptor ligation, bisulfite treatment is applied. Now the unmethylated sequences revert to a lower Tm in view of the C>T conversion, while methylated sequences remain at high Tm. Next, PCR is applied using the ligated adaptors, in order to generate amplified double stranded DNA library. (Alternatively one can apply PCR of repeat elements using primers specific for bisulfite-treated ALU, LINE 1 and other repeat elements spread over the genome, in

order to apply the method to repeat sequences only; or arbitrarily-primed PCR (AP-PCR) to apply the method to large, arbitrary genomic fractions; or one can apply COLD-PCR. Isothermal forms of amplification may also be used in the place of PCR).

Following amplification, one may use either preferential denaturation approach,  
5 or a preferential hybridization approach to enrich unmethylated sequences. The two approaches are described below.

For preferential denaturation approach, the temperature is raised to a level of choice that generates partial or complete denaturation of lower Tm sequences. These include the originally unmethylated sequences which due to the C>U conversion resulted  
10 to a C>T conversion in the final PCR product, reverting to a lower Tm than the methylated sequences. High Tm sequences remain double stranded. These include the highly methylated GC-rich sequences. One of ordinary skill can determine the Tm of the sequences using methods known in the art and as described herein. The temperature used for this preferential denaturation of the low Tm sequences includes, but is not  
15 limited to 50° C, 55° C, 60° C, 66° C, 70° C, 75° C, 78° C, 80° C, or 85° C). Next, DSN is added and the nucleic acids are exposed to conditions optimal for DSN activity. Conditions optimal for DSN activity include the most favorable conditions that allow the enzyme to work most efficiently for cleaving complimentary duplexes. The optimum  
20 DSN activity may be affected by conditions which include temperature; pH; and salt concentrations. In some embodiments, the temperature used for optimal for DSN activity includes, but not limited to 50° C, 55° C, 60° C, 65° C, or 70° C. The originally unmethylated sequences with lower Tm, (as well as naturally AT-rich sequences with  
25 lower Tm) will be denatured (*i.e.*, will become single stranded) and therefore will escape DSN digestion, while the originally methylated sequences will be cleaved since they will remain in double stranded formation at the denaturation temperature chosen. A subsequent amplification of the remaining sequences using the ligated common adaptors will amplify preferentially the intact unmethylated sequences and will enable downstream massively parallel sequencing of unmethylated alleles. In some embodiments, the preferential denaturation of genomic DNA is performed in the  
30 presence of organic solvents that can lower the Tm of the nucleic acids. Examples include but are not limited to betaine, DMSO, or formamide. It is known that betaine generates a narrower melting peak for DNA duplexes, hence by adding betaine the

discrimination between high and low Tm sequences at a given denaturation temperature will be ‘sharper’.

Accordingly, in some embodiments, the method comprises the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) 5 subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated nucleic acids while methylated nucleic acids remain double-stranded; and (d) exposing the unmethylated and methylated nucleic acids to 10 double strand-specific nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN cleaves the methylated double-stranded nucleic acids but not the unmethylated single-stranded nucleic acids.

Alternatively, for preferential re-hybridization approach, a complete denaturation step is applied. The denaturing temperature should be sufficiently high so as to allow the 15 full denaturation of the nucleic acids (*e.g.*, 75° C, 80° C, 85° C, 90° C, or 95° C). In some embodiments, the denaturing temperature is applied for 30 seconds, 1 min, 2 min, or 3 min. In some embodiments, the denaturing temperature of 95° C is applied for 2 min. Following this, the temperature is lowered to a level that allows methylated (or other high Tm) sequences to re-hybridize rapidly (due to their higher Tm), while 20 unmethylated sequences stay substantially single stranded. Next, application of DSN enzyme at conditions for optimal DSN activity digests preferentially the methylated duplexes. In some embodiments, the re-hybridization takes place in the presence of an organic solvent such as DMSO which lowers the Tm of the nucleic acid in combination with concurrent digestion using DSN (that is, instead of adding DSN in a later step). Use 25 of organic solvents such as DMSO allows temperatures compatible with DSN action (*e.g.*, 60-75° C) to be applied during re-hybridization. Finally, a PCR amplification step can be applied to amplify preferentially the non-digested, unmethylated alleles followed by sequencing.

Accordingly, in some embodiments, the method comprises the steps of: (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) 30 subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits

denaturation of both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids; (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and (d) exposing the unmethylated and methylated nucleic acids to double strand-specific

5 nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN preferentially cleaves the methylated duplexes but not the unmethylated single-stranded nucleic acids.

Bisulfite conversion of DNA has formed the basis of identifying the methylation state of individual genes. With the advent of high throughput parallel sequencing methods, this technology has extended to the sequencing of libraries of bisulfite-treated 10 DNA. The approach involves fragmenting DNA, ligating adaptors, bisulfite treatment and then amplifying the libraries for high throughput sequencing (see, for example, US 2013/0059734, US 2008/0254453, US 2009/0148842 and US 8440404).

In a reverse approach that aims to enrich the methylated alleles, following 15 preferential denaturation of the lower Tm sequences (which includes unmethylated alleles), treatment with any enzyme with selective action against single stranded (as opposed to double stranded) DNA such as but not limited to exonuclease I, or III is applied to remove the single stranded sequences. Subsequently, the remaining sequences (including the substantially methylated alleles) can be amplified and sequenced.

Accordingly, in some embodiments, the method comprises the steps of: (a) ligating 20 bisulfite-resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated nucleic acids while methylated nucleic acids remain double-stranded; and (d) exposing the unmethylated and methylated nucleic acids to an exonuclease and conditions for optimal exonuclease activity, wherein the exonuclease 25 cleaves the unmethylated single-stranded nucleic acids but not the methylated double-stranded nucleic acids.

In some embodiments, the method comprises the steps of: (a) ligating bisulfite- 30 resistant adaptors to double stranded nucleic acids of interest; (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids; (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits denaturation of

both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids; (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and (d) exposing the unmethylated and methylated nucleic acids to an exonuclease and conditions for optimal 5 exonuclease activity, wherein the exonuclease preferentially cleaves the unmethylated single-stranded nucleic acids, but not the methylated duplexes.

In some embodiments, the nucleic acid amplification reaction used to form double-stranded bisulfite-treated nucleic acids is selected from the group consisting of: PCR; full COLD-PCR, fast COLD-PCR; ice-COLD-PCR; temperature-tolerant COLD-10 PCR; LDT-COLD-PCR; AP-PCR; and repeat element PCR (ALU, LINE1, and other repeat sequences). In some embodiments, the resultant cleaved unmethylated single stranded nucleic acids and the uncleaved methylated duplexes are subjected to an amplification condition using the bisulfite resistant ligated adaptors. In some 15 embodiments, the amplification condition is selected from the group consisting of: PCR; LDT-COLD-PCR; AP-PCR; and repeat element PCR (ALU, LINE1, and other repeat sequences).

The specific advantage of performing the genome-wide amplification in COLD-PCR format is that, by employing a desired denaturation temperature during 20 amplification, COLD-PCR provides an additional enrichment of lower-Tm sequences, as has been demonstrated previously by using COLD-PCR on unmethylated single gene sequences (Castellanos-Rizaldos, E., Milbury, C.A., Karatza, E., Chen, C.C., Makrigiorgos, G.M. and Merewood, A. (2014) COLD-PCR amplification of bisulfite-converted DNA allows the enrichment and sequencing of rare un-methylated genomic regions. PLoS One, 9, e94103.). Sequences with Tm above the selected denaturation 25 temperature do not amplify during COLD-PCR. Any of the described COLD-PCR formats can be used to amplify selected fractions of un-methylated sequences from the genome (full COLD-PCR (11); fast COLD-PCR (11); ice-COLD-PCR (12); temperature-tolerant COLD-PCR (13); and limited denaturation time LDT-COLD-PCR, Murphy DM, Castellanos-Rizaldos E, Makrigiorgos GM. Clin Chem. 2014 60:1014-6). 30 Each COLD-PCR format will achieve amplification of different genomic fractions. For example, fast COLD-PCR utilizes a single denaturation temperature, hence any sequence with Tm above this temperature will not be amplified at all. Alternatively, temperature tolerant COLD-PCR utilizes successive steps of increasing denaturation temperatures

covering a broad range of temperatures (*e.g.* a range of 10°C in 10 steps of 1°C each), hence amplifying a broader range of unmethylated sequences from the genome.

Depending on the application, different COLD-PCR programs can be employed. Fig. 15 describes the combination of DSN digestion of methylated sequences with tandem

5 temperature tolerant COLD-PCR amplification of unmethylated sequences for further enrichment of globally unmethylated alleles.

In some embodiments of the methods used for preparing unmethylated or methylated nucleic acids of interest without using oligonucleotide probes, the naturally AT-rich sequences are removed prior to the sodium bisulfite treatment. Specifically, in 10 the approaches described herein for preferential amplification of un-methylated DNA (or methylated DNA), a bisulfite step is applied that converts sequences that were originally at higher Tm (GC-rich sequences that are un-methylated) to lower Tm sequences (due to the C>T conversion). In contrast, methylated alleles retain their high Tm. In this way, a preferential denaturation step followed by DSN digestion and by subsequent 15 amplification preferentially enriches the unmethylated alleles. However, during this amplification step, sequences with naturally low Tm will also be amplified irrespective of their methylation status (for example, AT-rich sequences). In some embodiments, such sequences with melting temperature (Tm) below a temperature of choice are removed to avoid these multiple ‘normal, non-informative’ sequences with low Tm to 20 amplify in competition to the target sequences, thereby overwhelming the amplification of the sequences of interest.

Such ‘non-informative’ sequences can be substantially depleted from the sample before performing the bisulfite conversion step as follows:

(a) Removing lower – Tm sequences. A temperature-based fractionation of the 25 starting material is performed just prior to sodium bisulfite treatment (Fig. 16).

Following shearing of genomic DNA, the DNA is treated with an enzyme that creates blunt ends (*e.g.* ‘end repair’) such as but not limited to T7 DNA polymerase. Next, the temperature is raised to a desired cutoff temperature. This desired cut off temperature is the temperature at which sequences having a lower Tm need to be selectively removed.

30 As an example, let it be assumed that the desired cut off temperature is 80°C. By raising the temperature to 80°C for time periods of 1 sec-5 min (*e.g.*, 1 sec, 5 secs, 15 secs, 30 secs, 45 secs, 1 min, 2 mins, 3 mins, 4, mins, or 5 mins), DNA fragments with Tm below 80°C will be substantially denatured, while other fragments with Tm above 80°C will

remain substantially double stranded. Next, the temperature is quickly lowered, for example, by placing the sample on ice. Next, an enzyme that degrades preferentially single stranded DNA is added (for example, exonucleases I or III) and the temperature is raised to the optimal temperature for this enzyme activity (for example, 37°C) for 1 min 5 – 60 min (e.g., 1 min, 5 mins, 10 mins, 20 mins, 30 mins, 40 mins, 45 mins, 50 mins, 55 mins, or 60 mins). Due to the complexity of genomic DNA, during this time period there is no substantial re-naturation of the single stranded fragments that undergo denaturation, and these become degraded by the enzyme. Finally, the exonuclease is heat inactivated. This process yields double stranded DNA fragments with Tm higher than about 80°C. In 10 this way, fragments with Tm substantially lower than the chosen cutoff temperature are substantially depleted. In some embodiments, the process is repeated for additional rounds if more strict temperature fractionation is needed. The additional rounds can be at the same temperature (for example, 80°C) or different temperatures (for example, 82- 85°C). Following removal of sequences with Tm below the desired cut off temperature, a 15 sodium bisulfite step is applied to selectively convert C to U in unmethylated CpG positions in nucleic acids. Accordingly, these unmethylated DNA fragments will now revert to sequences with lower Tm. Because the majority of sequences with naturally lower Tm (for example, AT-rich sequences) has been removed during exonuclease-based temperature fractionation, it is now possible to remove higher Tm methylated sequences 20 by preferential DSN digestion of double-stranded sequences, as well to amplify preferentially the unmethylated sequences via an amplification step such as PCR (or COLD-PCR) without interference of the lower Tm AT-rich sequences.

(b) Solid support-based temperature fractionation

25 Another approach for removing sequences with selected Tm from a genomic DNA sample includes solid support-based temperature fractionation. In some embodiments, the solid support comprises magnetic beads. Magnetic beads may be used for immobilization of the genomic DNA sample in order to enable separation of genomic DNA fragments into discrete temperature domains prior to further treatment (Fig. 17). 30 After separating the genomic DNA fragments into discrete temperature domains, preferential amplification of unmethylated alleles can be applied separately within each domain, with minimal interference from amplification of non-desired lower Tm DNA fragments that do not provide any enrichment advantage. To apply this protocol, the

ligation step depicted in Fig. 15 is performed with a mix of biotinylated and non-biotinylated bisulfite-resistant adaptors following which the majority of genomic DNA fragments are captured on streptavidin-coated magnetic beads from one strand only (the opposite strand remaining non-biotinylated). The conditions applied (total beads relative 5 to total DNA) enable immobilization of several hundred nanograms of biotinylated DNA so that enough sequence copies are immobilized to retain low-level events in the fragment population. Following washing of unbound DNA, the temperature is ramped-up in 5 different consecutive steps differing by, for example, 3°C. During each step, the non-biotinylated DNA strands from lower-Tm fragments are gradually denaturing and 10 are eluted in the supernatant which are then collected following bead magnetization (Fig. 17). DNA transitions from double stranded to mostly single stranded within an interval of ~4-5°C. By collecting the supernatant in temperature intervals of 3°C, the genomic DNA fragments are highly enriched (for example, 10-100-fold) in sequences within pre-defined temperature domains.

15

No DSN

Some aspects of the disclosure provide a method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising subjecting a nucleic acid sample comprising a double-stranded wild type nucleic acid 20 and a double-stranded target nucleic acid suspected of containing a mutation to a condition that destabilizes the double stranded wild type and target mutant nucleic acids; contacting the destabilized double stranded wild type and target mutant nucleic acids with a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid 25 bottom strand, to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild-type-probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation, and wherein one or both probes comprise 30 a locked nucleic acid (LNA), peptide nucleic acid (PNA), xeno nucleic acid (XNA), or a nucleic acid with any known modified base or RNA which is capable of blocking PCR amplification; and subjecting the complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes to an

amplification condition. The probes that overlap the mutation position act to block PCR amplification, e.g., acting as a clamp, for the wild-type top and bottom DNA strands, thereby inhibiting amplification of the wild-type nucleic acid. When the probe duplexes with a partially complimentary target mutant sequence, it is less able to inhibit PCR 5 amplification, thereby permitting selective amplification of the mutant nucleic acid as compared to the wild-type, without a need for a cleaving enzyme (e.g., DSN).

The present invention is further illustrated by the following Example, which in no way should be construed as further limiting. The entire contents of all of the references 10 (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

### EXAMPLE

15 *NaME on double-stranded DNA*

NaME utilizes nucleases (DNases) that have a substantially higher activity on double-stranded DNA (ds DNA) versus single-stranded DNA (ss DNA). Many DNases display such activity, including native shrimp dsDNase, recombinant shrimp dsDNase, King crab nuclease (DSN) and bovine DNase I. In the following sections, NaME 20 embodiments for DSN thermostable nuclease are provided, but the same approaches can be used for all other nucleases that display substantially higher activity for ds DNA versus ss DNA. Thermostable nuclease (DSN) selectively degrades double stranded DNA (or DNA/RNA hybrids), while it has minimal or no action on single stranded DNA or RNA.

25 For the purposes of the present disclosure, the term “double-strand specific nuclease” or “DSN” includes DNA/RNA guided enzymes which have preferential activity on double-stranded DNA, as opposed to single stranded DNA. Examples of such enzymes that can be employed in conjunction with NaME include the RNA-guided Cas9 enzymes (Gu et al, *Depletion of Abundant Sequences by Hybridization (DASH)*):

30 *Using Cas9 to remove unwanted high-abundance species in sequencing libraries and molecular counting applications* Genome Biology 2016; 17, 41), or the Argonaute DNA-guided enzymes (Gao et al, *DNA-guided genome editing using the Natronobacterium gregoryi Argonaute*, Nature Biotechnology May 2016 advanced

online publication). These DNA/RNA guided enzymes digest DNA with high preference when the probe ('guide oligonucleotide') is fully matched to the target DNA, and less so when there is a mismatch. By employing probes targeting both top and bottom DNA strands in an overlapping fashion as described in the present invention, NAME can be 5 applied with DNA/RNA guided enzymes, in the same manner as when using other DSN nucleases described herein.

NaME takes advantage of the DSN properties to degrade specific sequences from both the top and bottom DNA strands of wild-type (WT) DNA (Figure 1A). In contrast, 10 mutation-containing DNA is not degraded or degraded much less than the WT DNA. Hence, a subsequent PCR reaction after DSN digestion specifically amplifies the mutant alleles that remain substantially intact.

An example of the application of this approach is demonstrated in Figure 1B: a 114bp ds KRAS PCR amplicon with a 5% mutation was subjected to the process of 15 Figure 1A. The DNA template used consisted of the KRAS PCR amplicon with a 5% mutation (1:1000, 1:10,000 (approximately 0.001nM), and 1:100,000) and wtDAN-Taqman-probe (500nM) and with or without KRAS-cutter (500nM). The samples were then either incubated with DSN (1U) or without DSN. For the 1:1000 and 1:100,000 samples, the mixture was heated to 67°C for 10 minutes and then 94° for 2 minutes. For 20 the 1:10,000 samples, the mixture was heated to a selected temperature (63°C, 67°C, 70°C, or 73°C) for 10 minutes, and then heated to 94°C for two minutes. By comparing the mutation abundance in parallel reactions without DSN versus reactions with DSN, the data in Figure 1B and Table 1 demonstrate mutation enrichments of about 10-fold for temperatures 63-67°C.

25 **Table 1:** KRAS Mutation Abundance at Different Temperatures

Sample ID	Delta Ct	Final mutation abundance	Enrichment-fold
1-10k-NO-DSN	0	4.14	1.0 (untreated)
1-10k-1U-63C	9.07	38.6	9.3
1-10k-1U-67C	<b>8.58</b>	37.4	<b>9.0</b>
1-10k-1U-70C	6.54	14	3.4
1-10k-1U-73C	7.11	19.5	4.7
1-10k-1U-67C-no-	<b>5.75</b>	6.07	<b>1.5</b>

cutter			
--------	--	--	--

*NaME applied directly from genomic DNA*

Figures 5-7 demonstrate NaME applied directly to genomic DNA for a single KRAS target sequence, or single p53 sequence (Figure 5), two targets simultaneously, 5 duplex KRAS and p53 (Figure 6) and three different KRAS mutations in a single-target reaction (Figure 7A). In Figure 5, genomic DNA with approximately a 0.5% KRAS mutation, or a p53 mutation in a separate reaction was denatured at 95°C and incubated at 65°C for 20 minutes in the presence of overlapping blocked probes and DSN. The enrichment of the KRAS or p53 mutations were detected via a subsequent digital PCR 10 that quantifies the mutation abundance. In Figure 6, the genomic DNA underwent the same protocol as in Figure 5, except that the KRAS and p53 mutated genomic DNA were mixed in a single tube. Genomic DNA from the SW480 cell line containing both KRAS and p53 mutations was mixed with wild-type DNA to create low-abundance mutations on both genes. Two mutation percentage were tested: approximately 5% and 15 approximately 0.3%. In Figure 7A, genomic DNA with 1-5% KRAS mutations from three different cell lines, in separate reactions, underwent the same protocol as described above. In Figure 7B, the DNA was denatured, the temperature was reduced to 67°C, DSN and overlapping probes were added for a 20 minute incubation. The DSN was inactivated, and PCR and digital PCR were performed on each target amplicon to derive 20 their respective mutational abundances. All mutations are enriched simultaneously via NaME. Figure 7B depicts mutation enrichment when NaME is applied on 11 targets simultaneously, directly from genomic DNA obtained from Horizon Dx, containing known low level mutations on the respective targets. It is demonstrated that all 11 targets are enriched simultaneously following application of NaME directly to genomic 25 DNA.

*Mutation scanning using NaME*

Figures 12B and 12C depict the results when using NaME for mutation enrichment of a 40-80bp region in TP53 exon 8. All of the 4 mutations tested are 30 enriched via NaME. In Figure 12B, mutation-containing DNA from the PFSK or HCC cell lines was serially diluted into WT DNA, and then a first PCR was applied to amplify the target of interest (tp53). The amplicon was denatured and then incubated at 65°C in

the presence of probes and DSN. The two probes correspond to the WT top and bottom strands, respective. The presence of a mutation inhibits DSN digestion, hence the mutated DNA is amplified during rounds of PCR. The effects of probe length and concentration on mutation enrichment are also depicted in Figure 12B. In Figure 12C, 5 mutation-containing DNA from PFSK, HCC, SW480, and MDAMB cell lines, all of which have different mutations on p53 exon 8, were serially diluted into WT DNA, and then the same protocol as described in Figure 12B was performed. The mutation enrichment-fold was calculated by performing digital PCR both before and after NaME application.

10

### References

1. Thomas, R.K., Baker, A.C., Debiasi, R.M., Winckler, W., Laframboise, T., Lin, W.M., Wang, M., Feng, W., Zander, T., Macconnail, L.E. *et al.* (2007) High-throughput oncogene mutation profiling in human cancer. *Nat Genet*, **39**, 347-351.
- 15 2. Chou, L.S., Lyon, E. and Wittwer, C.T. (2005) A comparison of high-resolution melting analysis with denaturing high-performance liquid chromatography for mutation scanning: cystic fibrosis transmembrane conductance regulator gene as a model. *Am J Clin Pathol*, **124**, 330-338.
3. Thomas, R.K., Nickerson, E., Simons, J.F., Janne, P.A., Tengs, T., Yuza, Y., 20 Garraway, L.A., Laframboise, T., Lee, J.C., Shah, K. *et al.* (2006) Sensitive mutation detection in heterogeneous cancer specimens by massively parallel picoliter reactor sequencing. *Nat Med*, **12**, 852-855.
4. Paez, J.G., Janne, P.A., Lee, J.C., Tracy, S., Greulich, H., Gabriel, S., Herman, P., Kaye, F.J., Lindeman, N., Boggon, T.J. *et al.* (2004) EGFR Mutations in Lung Cancer: 25 Correlation with Clinical Response to Gefitinib Therapy. *Science*, **304**, 1497-1500.
5. Janne, P.A., Borras, A.M., Kuang, Y., Rogers, A.M., Joshi, V.A., Liyanage, H., Lindeman, N., Lee, J.C., Halmos, B., Maher, E.A. *et al.* (2006) A rapid and sensitive enzymatic method for epidermal growth factor receptor mutation screening. *Clin Cancer Res*, **12**, 751-758.
- 30 6. Engelman, J.A., Mukohara, T., Zejnnullahu, K., Lifshits, E., Borras, A.M., Gale, C.M., Naumov, G.N., Yeap, B.Y., Jarrell, E., Sun, J. *et al.* (2006) Allelic dilution obscures detection of a biologically significant resistance mutation in EGFR -amplified lung cancer. *The Journal of clinical investigation*.

7. Diehl, F., Li, M., Dressman, D., He, Y., Shen, D., Szabo, S., Diaz, L.A., Jr., Goodman, S.N., David, K.A., Juhl, H. *et al.* (2005) Detection and quantification of mutations in the plasma of patients with colorectal tumors. *Proc Natl Acad Sci U S A*, **102**, 16368-16373.
- 5 8. Kimura, T., Holland, W.S., Kawaguchi, T., Williamson, S.K., Chansky, K., Crowley, J.J., Doroshow, J.H., Lenz, H.J., Gandara, D.R. and Gumerlock, P.H. (2004) Mutant DNA in plasma of lung cancer patients: potential for monitoring response to therapy. *Annals of the New York Academy of Sciences*, **1022**, 55-60.
9. Nilsen, I.W., Overbo, K., Jensen Havdalen, L., Elde, M., Gjellesvik, D.R. and 10 Lanes, O. (2010) The enzyme and the cDNA sequence of a thermolabile and double-strand specific DNase from Northern shrimps (*Pandalus borealis*). *PLoS One*, **5**, e10295.
10. Castellanos-Rizaldos, E., Milbury, C.A., Karatza, E., Chen, C.C., Makrigiorgos, G.M. and Merewood, A. (2014) COLD-PCR amplification of bisulfite-converted DNA allows the enrichment and sequencing of rare un-methylated genomic regions. *PLoS One*, **9**, e94103.
- 15 11. Li, J., Wang, L., Mamon, H., Kulke, M.H., Berbeco, R. and Makrigiorgos, G.M. (2008) Replacing PCR with COLD-PCR enriches variant DNA sequences and redefines the sensitivity of genetic testing. *Nat Med*, **14**, 579-584.
12. Milbury, C.A., Li, J. and Makrigiorgos, G.M. (2011) Ice-COLD-PCR enables 20 rapid amplification and robust enrichment for low-abundance unknown DNA mutations. *Nucleic Acids Res*, **39**, e2.
13. Castellanos-Rizaldos, E., Liu, P., Milbury, C.A., Guha, M., Brisci, A., Cremonesi, L., Ferrari, M., Mamon, H. and Makrigiorgos, G.M. (2012) Temperature-Tolerant COLD-PCR Reduces Temperature Stringency and Enables Robust Mutation 25 Enrichment. *Clin Chem*, **58**, 1130-1138.
14. Shagin, D.A., Rebrikov, D.V., Kozhemyako, V.B., Altshuler, I.M., Shcheglov, A.S., Zhulidov, P.A., Bogdanova, E.A., Staroverov, D.B., Rasskazov, V.A. and Lukyanov, S. (2002) A novel method for SNP detection using a new duplex-specific nuclease from crab hepatopancreas. *Genome Res*, **12**, 1935-1942.
- 30 15. Qiu, X., Zhang, H., Yu, H., Jiang, T. and Luo, Y. (2015) Duplex-specific nuclease-mediated bioanalysis. *Trends in biotechnology*, **33**, 180-188.

We claim:

CLAIMS

1. A method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising
  - subjecting a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a condition that destabilizes the double stranded wild type and target mutant nucleic acids;
  - contacting the destabilized double stranded wild type and target mutant nucleic acids with a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild-type-probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; and
  - exposing the complimentary wild-type-probe duplexes and the partially complimentary target mutant-probe duplexes to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.
2. A method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising
  - exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation; and
  - subjecting the reaction mixture to a condition that destabilizes the double stranded wild type and target mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby

forming complimentary wild-type- probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes,

wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

3. The method of any one of claims 1-2, wherein the condition that destabilizes the double stranded wild type and mutant nucleic acids to permit hybridization of the probes to their corresponding sequences on the wild type and mutant nucleic acids comprises addition of an organic solvent and/or an increase in temperature combined with a thermostable DSN.

4. A method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid comprising

exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation;

subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid;

reducing the temperature of the reaction mixture to permit formation of complimentary wild type – probe duplexes on top and bottom strands and partially complimentary target mutant-probe duplexes; and

exposing the reaction mixture to a double strand-specific nuclease (DSN), wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

5. The method of any one of claims 1 – 4, wherein the probes are in a molar excess of 100-fold to 1 billion-fold compared to the wild type and target nucleic acids.

6. The method of any one of claims 1-4, wherein one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the

other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other.

7. The method of any one of claims 1-6, wherein each probe comprises a locked nucleic acid, peptide nucleic acid, xeno nucleic acid (XNA), nucleic acid with any known modified base or RNA.

8. The method of any one of claims 1-7, wherein the method is used to prepare two or more different target mutant nucleic acids for subsequent enrichment relative to corresponding wild type nucleic acids, and the method further comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand.

9. The method of any one of claims 1-8, wherein the nucleic acid sample comprises genomic DNA or circulating DNA in urine, plasma or other bodily fluids.

10. The method of claim 9, further comprising enriching the nucleic acids for regions of interest prior to implementing the method of claim 2 or 4 as follows: contacting the nucleic acid sample with bait oligonucleotides that bind to different nucleic acids of interest on top and bottom strands, permitting binding of the bait oligonucleotides to the nucleic acids of interest on top and bottom strands, and isolating the bait oligonucleotides with the nucleic acids of interest bound thereto from remaining nucleic acids.

11. The method of claim 10, wherein the bait oligonucleotides are biotinylated at one end.

12. The method of claim 11, wherein the bait oligonucleotides are attached to beads.

13. The method of any one of claims 1-12, wherein prior to implementing the method of claim 2 or 4, the nucleic acid sample is subjected to an amplification condition.

14. The method of any one of claims 1-13, further comprising enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid.
15. The method of claim 13 or 14, wherein the amplification condition is selected from the group consisting of: PCR, COLD-PCR, ligation mediated PCR or COLD-PCR using common ligated adaptors, multiplex PCR, and isothermal amplification.
16. The method of any one of claims 13-14, wherein each probe is modified at the 3' end to prevent polymerase extension.
17. The method of any one of claims 1-16, further comprising enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type -probe duplexes and uncleaved target mutant nucleic acids to a further DNA degradation condition which hydrolyzes enzymatically the DSN-cleaved wild type-probe duplexes, with the degradation initiated at the position of the cleavage.
18. The method of any one of claims 1-16, further comprising analyzing the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids using one or more of the methods selected from the group consisting of: MALDI-TOF, HR-Melting, Di-deoxy-sequencing, Single-molecule sequencing, massively parallel sequencing (MPS), pyrosequencing, SSCP, RFLP, dHPLC, CCM, digital PCR and quantitative-PCR.
19. The method of any one of claims 1-18, wherein the method is used to prepare an unmethylated target nucleic acid of interest for subsequent enrichment, and wherein prior to implementing the method of claim 2 or 4 on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite and one of the oligonucleotide probes is complimentary to top strand of the methylated nucleic acid of interest, while the other

oligonucleotide probe is complimentary to the bottom strand of the methylated nucleic acid of interest.

20. The method of any one of claims 1-18, wherein the method is used to prepare a methylated target nucleic acid of interest for subsequent enrichment, and wherein prior to implementing the method of claim 2 or 4 on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite and one of the oligonucleotide probes is complimentary to top strand of the unmethylated nucleic acid of interest, while the other oligonucleotide probe is complimentary to the bottom strand of the unmethylated nucleic acid of interest.

21. The method of any one of claims 1-18, wherein the method is used to prepare both an unmethylated target nucleic acid of interest and a methylated target nucleic acid of interest for subsequent enrichment, wherein the method comprises:

(i) a pair of oligonucleotide probes, one of which is complimentary to top strand of the methylated form of the unmethylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the methylated form of the unmethylated target nucleic acid of interest,

(ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of the methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the unmethylated form of the methylated target nucleic acid of interest; and

wherein prior to implementing the method of claim 2 or 4 on the reaction mixture, the nucleic acid sample is treated with sodium bisulfite.

22. The method of claim 21, wherein the method is used to prepare multiple target nucleic acids of interest, some of which are methylated target nucleic acids of interest, and some of which are unmethylated target nucleic acids of interest, and the method comprises

(i) a pair of oligonucleotide probes, one of which is complimentary to top strand of the methylated form of each unmethylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the methylated form of each unmethylated target nucleic acid of interest,

(ii) a pair of oligonucleotide probes, one of which is complimentary to top strand of the unmethylated form of each methylated target nucleic acid of interest, while the other is complimentary to the bottom strand of the unmethylated form of each methylated target nucleic acid of interest.

23. A method for preparing a target mutant nucleic acid for subsequent enrichment relative to a wild type nucleic acid, the method comprising the steps of:

- (a) exposing a nucleic acid sample comprising a double-stranded wild type nucleic acid and a double-stranded target nucleic acid suspected of containing a mutation to a thermostable double strand-specific nuclease (DSN) and a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, to create a reaction mixture, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation;
- (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid; and
- (c) reducing the temperature to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes,

wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes.

24. The method of claim 23, wherein the probes are in a molar excess of 100-fold- to 1 billion-fold compared to the wild type and target nucleic acids.

25. The method of any one of claims 23-24, further comprising repeating steps (b) and (c) for two or more cycles.

26. The method of any one of claims 23-25, wherein the reaction mixture further comprises an organic solvent.

27. The method of any one of claims 23-26, wherein the denaturing temperature is between 65-85°C.
28. The method of any one of claims 23-27, wherein step (c) of reducing the temperature is performed by applying a decreasing temperature gradient between temperatures that permit hybridization of probes having different Tm to their corresponding sequences on the target DNA.
29. The method of claim 28, wherein a decreasing temperature gradient from 67°C to 64°C is applied during DSN digestion, to allow probes that have distinct Tms of 64-67°C to hybridize to their respective targets.
30. The method of claim 28 or 29, wherein the temperature gradient ranges between 2-20°C
31. The method of any one of claims 23-30, further comprising enriching the target mutant nucleic acid relative to the wild type nucleic acid by subjecting the reaction mixture with cleaved wild type-probe duplexes and uncleaved target mutant nucleic acids to an amplification condition.
32. The method of any one of claims 23-31, wherein one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other.
33. The method of any one of claims 23-32, wherein the method is used to prepare two or more different target mutant nucleic acids for subsequent enrichment relative to corresponding wild type nucleic acids and the method further comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand.

34. A method for enriching a target mutant nucleic acid, the method comprising the steps of:

- (a) preparing an amplification reaction mixture comprising:  
a double-stranded wild type nucleic acid, a double-stranded target nucleic acid suspected of containing a mutation, a thermostable double strand-specific nuclease (DSN), a pair of oligonucleotide probes, one of which is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand, wherein at least one of the probes overlaps a sequence on the target nucleic acid containing the suspected mutation and PCR amplification components;
- (b) subjecting the reaction mixture to a denaturing temperature to permit denaturation of the wild type nucleic acid and the target mutant nucleic acid;
- (c) reducing the temperature to permit hybridization of the probes to their corresponding sequences on the wild type and target mutant nucleic acids thereby forming complimentary wild type – probe duplexes on top and bottom strands, and partially complimentary target mutant-probe duplexes, wherein the DSN cleaves the complimentary wild type – probe duplexes but not the partially complimentary target mutant-probe duplexes; and
- (d) subjecting the reaction mixture to an amplification condition thereby enriching the uncleaved target mutant nucleic acid relative to the cleaved wild type nucleic acid.

35. The method of claim 34, further comprising repeating steps (b) and (c) for two or more cycles before executing step (d).

36. The method of any one of claims 34-35, further comprising repeating steps (b), (c) and (d) for two or more cycles.

37. The method of any one of claims 34-36, wherein the reaction mixture further comprises an organic solvent.

38. The method of any one of claims 34-37, wherein the denaturing temperature is between 65-85°C.

39. The method of any one of claims 34-38, wherein one of the probes overlaps a sequence on the top strand of the target nucleic acid containing the mutation, while the other probe overlaps a sequence on the bottom strand of the target nucleic acid containing the mutation and the two probes partially overlap each other.

40. The method of any one of claims 34-39, wherein each probe is modified at the 3' end to prevent polymerase extension.

41. The method of any one of claims 34-40, wherein primers used for PCR amplification have a melting temperature that is below the temperature applied in step (c).

42. The method of any one of claims 34-41, wherein the method is used to enrich two or more different target mutant nucleic acids relative to wild type nucleic acids and the method further comprises one or more additional pairs of probes directed to the different wild type nucleic acids, wherein for each pair of probes, one of the probes is complimentary to the wild type nucleic acid top strand and the other is complimentary to the wild type nucleic acid bottom strand.

43. The method of any one of claims 34-42, wherein the amplification condition is COLD-PCR.

44. A method for preparing unmethylated nucleic acids of interest for subsequent enrichment relative to corresponding methylated nucleic acids comprising the steps of:

- (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest;
- (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids;
- (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated nucleic acids while methylated nucleic acids remain double-stranded;
- (d) exposing the unmethylated and methylated nucleic acids to double strand-specific nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN cleaves the

methylated double-stranded nucleic acids but not the unmethylated single-stranded nucleic acids.

45. A method for preparing unmethylated nucleic acids of interest for subsequent enrichment relative to corresponding methylated nucleic acids comprising the steps of:

- (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest;
- (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids;
- (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits denaturation of both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids;
- (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and
- (e) exposing the unmethylated and methylated nucleic acids to double strand-specific nuclease (DSN) and conditions for optimal DSN activity, wherein the DSN preferentially cleaves the methylated duplexes but not the unmethylated single-stranded nucleic acids.

46. A method for preparing methylated nucleic acids of interest for subsequent enrichment relative to corresponding unmethylated nucleic acids comprising the steps of:

- (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest;
- (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids;
- (c) subjecting the bisulfite-treated nucleic acids to a temperature that permits preferential denaturation of unmethylated nucleic acids while methylated nucleic acids remain double-stranded; and
- (d) exposing the unmethylated and methylated nucleic acids to an exonuclease and conditions for optimal exonuclease activity, wherein the exonuclease cleaves the unmethylated single-stranded nucleic acids but not the methylated double-stranded nucleic acids.

47. A method for preparing methylated nucleic acids of interest for subsequent enrichment relative to corresponding unmethylated nucleic acids comprising the steps of:

- (a) ligating bisulfite-resistant adaptors to double stranded nucleic acids of interest;
- (b) subjecting the adaptor-linked nucleic acids to sodium bisulfite treatment and a nucleic acid amplification reaction to form double-stranded bisulfite-treated nucleic acids;
- (c) subjecting the bisulfite-treated nucleic acids to a denaturing temperature that permits denaturation of both unmethylated and methylated nucleic acids to form unmethylated and methylated single stranded nucleic acids;
- (d) reducing the temperature to permit preferential formation of methylated duplexes, but not unmethylated duplexes; and
- (e) exposing the unmethylated and methylated nucleic acids to an exonuclease and conditions for optimal exonuclease activity, wherein the exonuclease preferentially cleaves the unmethylated single-stranded nucleic acids, but not the methylated duplexes.

48. The method of any one of claims 44-47, wherein the nucleic acid amplification reaction of step (b) is selected from the group consisting of: PCR; full COLD-PCR, fast COLD-PCR; ice-COLD-PCR, temperature-tolerant COLD-PCR and limited denaturation time COLD-PCR.

49. The method of any one of claims 44-48 where the cleaved unmethylated single stranded nucleic acids and the uncleaved methylated duplexes are subjected to an amplification condition using the bisulfite resistant adaptors ligated in step (a).

50. The method of claim 49, wherein the amplification condition is selected from the group consisting of: PCR, full COLD-PCR, fast COLD-PCR; ice-COLD-PCR, temperature-tolerant COLD-PCR and limited denaturation time COLD-PCR.

51. The method of any one of claims 44-50, wherein naturally AT-rich sequences are removed prior to the sodium bisulfite treatment.





Fig. 1A

# Digital PCR result



Fractional Abundance(%)

SUBSTITUTE SHEET (RULE 26)

Fig. 1B

3/22




Fig. 2





Fig. 3



6/22




Fig. 5

7/22




Fig. 6

8/22




Fig. 7A

9/22

1	2	3	4	5	6	7
Gene	Variant	No-DSN (%)	0.2U-DSN prod (%)	5% gDNA ctrl (%)	% given from HDx	
BRAF	V600E	c.1799T>A	11.7	99.8	6.64	8%
BRAF	V600K	c.1798_1799GT>AA	6.29	97.9	2.78	4%
EGFR	T790M	c.2369C>T	7.15	69.1	3.84	4.2%
FLT3	Δ1836	c.2506_2508delATC	7.18	99.9	6.13	3%
IDH1	R132C	c.394C>T	7.75	99.3	5.62	5%
JAK2	V617F	c.1849G>T	8.33	69.4	5.21	5%
KRAS	G12A	c.35G>C	7.05	21.3	4.41	5%
KRAS	G12R	c.34G>C	7.94	16.6	1.14	5%
MEK1	P124L	c.371C>T	6.18	95.5	5.04	5%
NOTCH1	L1601P	c.4799T>C	10	64	9.55	4.80%
NRAS	Q61K	c.181C>A	8.98	78.3	5.45	5%

Fig. 7B

10/22




Fig. 8






Fig. 9





14/22




Fig. 12A





Fig. 12B

16/22

## Mutation Enrichment-Fold as a Function of Probe Length



## Mutation Enrichment-Fold as a function of Probe Conc.




Fig. 12B  
(Continued)

17/22




Fig. 12C




Fig. 13




Fig. 14




Fig. 15




Fig. 16

22/22




Fig. 17

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/039167

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12N 15/10; C12P 19/34; C12Q 1/68 (2016.01)

CPC - C12P 19/34; C12Q 1/68; C12Q 1/6827; C12Q 1/6858 (2016.08)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC - C12N 15/10; C12P 19/34; C12Q 1/68

CPC - C12P 19/34; C12Q 1/68; C12Q 1/6827; C12Q 1/6858

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC - 435/6; 435/91.1; 536/23.1 (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatBase, Google Patents, Google Scholar, PubMed

Search terms used: double strand-specific nuclease duplex-specific nuclease minor allele variant mutant methylated unmethylated

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SHAGIN et al. "A novel method for SNP detection using a new duplex-specific nuclease from crab hepatopancreas," Genome Res. December 2002, Vol. 12, Pgs. 1935-42 entire document	1-4, 23-25, 34-36, 44-48
A	CASTELLANOS-RIZALDOS et al. "COLD-PCR amplification of bisulfite-converted DNA allows the enrichment and sequencing of rare un-methylated genomic regions," PLoS One, 11 April 2014 (11.04.2014), Vol. 9, e94103, Pgs. 1-5. entire document	1-4, 23-25, 34-36, 44-48
A	WO 2015/013166 A1 (DANA-FARBER CANCER INSTITUTE, INC) 29 January 2015 (29.01.2015) entire document	1-4, 23-25, 34-36, 44-48
A	US 2014/0051087 A1 (MAKRIGIORGOS et al) 20 February 2014 (20.02.2014) entire document	1-4, 23-25, 34-36, 44-48
A	US 2009/0068652 A1 (TAYLOR et al) 12 March 2009 (12.03.2009) entire document	1-4, 23-25, 34-36, 44-48

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

30 August 2016

Date of mailing of the international search report

26 SEP 2016

Name and mailing address of the ISA/  
Mail Stop PCT, Attn: ISA/US, Commissioner for Patents  
P.O. Box 1450, Alexandria, VA 22313-1450  
Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300  
PCT OSP: 571-272-7774

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US2016/039167

**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3.  Claims Nos.. 5-22, 26-33, 37-43, 49-51  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.