



(86) Date de dépôt PCT/PCT Filing Date: 2012/09/28
(87) Date publication PCT/PCT Publication Date: 2013/04/04
(45) Date de délivrance/Issue Date: 2023/03/21
(85) Entrée phase nationale/National Entry: 2014/03/26
(86) N° demande PCT/PCT Application No.: JP 2012/075092
(87) N° publication PCT/PCT Publication No.: 2013/047752
(30) Priorités/Priorities: 2011/09/30 (JP2011-217498);
2012/02/24 (JP PCT/JP2012/054624);
2012/08/24 (JP2012-185866)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61P 29/00 (2006.01), *A61P 35/00* (2006.01),
A61P 37/02 (2006.01), *C07K 14/47* (2006.01),
C12P 21/02 (2006.01), *C12N 15/09* (2006.01)
(72) Inventeurs/Inventors:
IGAWA, TOMOYUKI, JP;
MAEDA, ATSUHIKO, JP;
HARAYA, KENTA, JP;
IWAYANAGI, YUKI, JP;
TACHIBANA, TATSUHIKO, JP
(73) Propriétaire/Owner:
CHUGAI SEIYAKU KABUSHIKI KAISHA, JP
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : MOLECULE LIANT UN ANTIGENE SERVANT A PROMOUVOIR L'ELIMINATION DES ANTIGENES
(54) Title: ANTIGEN-BINDING MOLECULE FOR PROMOTING ELIMINATION OF ANTIGENS

(57) **Abrégé/Abstract:**

The present inventors created antigen-binding molecules containing an antigen-binding domain and an Fcγ-receptor-binding domain, wherein the molecules have human-FcRn-binding activity in an acidic pH range condition, the antigen-binding domain changes the antigen-binding activity of the antigen-binding molecules depending on the ion-concentration condition, and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor in a neutral pH range condition than an Fc region of a native human IgG in which the sugar chain bound at position 297 (EU numbering) is a fucose-containing sugar chain.

ABSTRACT

5 The present inventors created antigen-binding molecules containing an antigen-binding domain and an Fc γ -receptor-binding domain, wherein the molecules have human-FcRn-binding activity in an acidic pH range condition, the antigen-binding domain changes the antigen-binding activity of the antigen-binding molecules depending on the ion-concentration condition, and the Fc γ receptor-binding domain has higher binding activity to the Fc γ receptor in a neutral pH range condition than an Fc region of a native human IgG in which the sugar chain bound at position 297 (EU numbering) is a fucose-containing sugar chain.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 254

NOTE : Pour les tomes additionnels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 254

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

DESCRIPTION

ANTIGEN-BINDING MOLECULE FOR PROMOTING ELIMINATION OF ANTIGENS

Technical Field

The present invention provides antigen-binding molecules with enhanced intracellular uptake of a bound antigen, antigen-binding molecules in which the number of antigens that can be bound per single molecule is increased, antigen-binding molecules with improved pharmacokinetics, antigen-binding molecules with enhanced intracellular dissociation of extracellularly bound antigen, antigen-binding molecules with enhanced extracellular release in the antigen-unbound state, antigen-binding molecules having the function of decreasing the total antigen concentration or the free antigen concentration in plasma, pharmaceutical compositions comprising such an antigen-binding molecules, and methods for producing them.

Background Art

Antibodies are drawing attention as pharmaceuticals as they are highly stable in plasma and have few side effects. A number of IgG-type antibody pharmaceuticals are now available on the market and many antibody pharmaceuticals are currently under development (Non-patent Documents 1 and 2). Meanwhile, various technologies applicable to second-generation antibody pharmaceuticals have been reported, including those that enhance effector function, antigen-binding ability, pharmacokinetics and stability, and those that reduce the risk of immunogenicity (Non-patent Document 3). In general, the requisite dose of an antibody pharmaceutical is very high. This in turn has led to problems such as high production cost as well as difficulty in producing subcutaneous formulations. In theory, the dose of an antibody pharmaceutical may be reduced by improving antibody pharmacokinetics or improving the affinity between antibodies and antigens.

Literature has reported methods for improving antibody pharmacokinetics using artificial substitution of amino acids in constant regions (Non-patent Documents 4 and 5). Similarly, affinity maturation has been reported as a technology for enhancing antigen-binding ability or antigen-neutralizing activity (Non-patent Document 6). This technology enables enhancement of antigen-binding activity by introduction of amino acid mutations into the CDR of a variable region or such. The enhancement of antigen-binding ability enables improvement of *in vitro* biological activity or reduction of dosage, and further enables improvement of *in vivo* efficacy (Non-patent Document 7).

The antigen-neutralizing capacity of a single antibody molecule depends on its affinity.

By increasing the affinity, an antigen can be neutralized by a smaller amount of an antibody. Various methods can be used to enhance antibody affinity (Non-patent Document 6). Furthermore, if affinity could be made infinite by covalently binding an antibody to an antigen, a single antibody molecule could neutralize one antigen molecule (a divalent antibody can
5 neutralize two antigen molecules). However, the stoichiometric neutralization of one antibody against one antigen (one divalent antibody against two antigens) is the limit of pre-existing methods, and thus it is impossible to completely neutralize an antigen with an amount of antibody smaller than the amount of antigen. In other words, the affinity enhancing effect has a limit (Non-patent Document 9). To prolong the neutralization effect of a neutralizing antibody
10 for a certain period, the antibody must be administered at a dose higher than the amount of antigen produced in the body during the same period. With just the improvement of antibody pharmacokinetics or affinity maturation technology described above, there is thus a limitation in the reduction of the required antibody dose. Accordingly, in order to sustain an antibody's antigen-neutralizing effect for a target period with an amount of antibody smaller than the
15 amount of antigen, a single antibody must neutralize multiple antigens. An antibody that binds to an antigen in a pH-dependent manner has recently been reported as a novel method for achieving the above objective (Patent Document 1). pH-dependent antigen-binding antibodies, which bind strongly to an antigen under neutral conditions in plasma and dissociate from the antigen under acidic conditions in the endosome, can dissociate from the antigen in the
20 endosome. When a pH-dependent antigen-binding antibody dissociates from the antigen is recycled to the plasma by FcRn, it can bind to another antigen again. Thus, a single pH-dependent antigen-binding antibody can bind to a number of antigens repeatedly.

In addition, plasma retention of an antigen is very short as compared to antibodies recycled via FcRn binding. When an antibody with such long plasma retention binds to the
25 antigen, the plasma retention time of the antigen-antibody complex is prolonged to the same retention time as that of the antibody. Thus, plasma retention of the antigen is prolonged by binding to the antibody, and thus the plasma antigen concentration is increased.

Accordingly, pH-dependent antigen-binding antibodies have effects that could not be accomplished by normal antibodies, since they can promote elimination of antigens from plasma
30 compared to normal antibodies, by binding of a single antibody to a plurality of antigens. However, antibody engineering methods for improving the effects of pH-dependent antigen-binding antibodies that can bind repeatedly to antigens and that promote elimination of antigens from plasma have not been reported to date.

IgG antibodies have long retentivity in plasma due to their binding to FcRn. Binding
35 between IgG and FcRn is observed only under acidic conditions (pH6.0), and the binding is hardly observed under neutral conditions (pH7.4). IgG antibodies are taken up into cells

non-specifically, but upon binding to FcRn in the endosome under an intra-endosome acidic condition, they return to the cell surface, and dissociate from FcRn under neutral conditions in plasma. When mutations are introduced into an Fc region of IgG so that binding to FcRn in an acidic pH range condition is lost, recycling of the antibody from endosome to plasma does not
5 take place and plasma retentivity of the antibodies is significantly impaired. A method for improving FcRn binding in an acidic pH range condition has been reported as a method for improving plasma retentivity of an IgG antibody. Improving binding to FcRn in an acidic pH range condition by introducing amino acid substitutions to an Fc region of IgG antibody leads to increased efficiency of antibody recycling from endosome to plasma, and as a result, plasma
10 retentivity is improved.

Many studies have been carried out so far on antibody-dependent cellular cytotoxicity (hereinafter denoted as ADCC) and complement-dependent cytotoxicity (hereinafter denoted as CDC), which are effector functions of IgG class antibodies. It has been reported that in the human IgG class, antibodies of the IgG1 subclass have the highest ADCC activity and CDC
15 activity (Non-Patent Document 13). Furthermore, antibody-dependent cell-mediated phagocytosis (ADCP), which is phagocytosis of target cells mediated by IgG class antibodies, is also suggested to be one of the antibody effector functions (Non-Patent Documents 14 and 15). Since IgG1 subclass antibodies can exert these effector functions against tumors, IgG1 subclass antibodies are used for most antibody pharmaceuticals against cancer antigens.

20 In order for IgG antibodies to mediate ADCC and ADCP activities, the Fc region of the IgG antibodies must bind to antibody receptors (hereinafter denoted as Fcγ receptor or FcγR) that are present on the surface of effector cells such as killer cells, natural killer cells, and activated macrophages. In humans, isoforms FcγRIa, FcγRIIa, FcγRIIb, FcγRIIIa, and FcγRIIIb have been reported as members of the Fcγ receptor protein family, and their respective allotypes have
25 been reported as well (Non-Patent Document 16).

Enhancement of cytotoxic effector functions such as ADCC and ADCP has been drawing attention as a promising means for enhancing the antitumor effects of anticancer antibodies. Importance of Fcγ receptor-mediated effector functions aimed for antitumor effects of antibodies has been reported using mouse models (Non-Patent Documents 17 and 18).
30 Furthermore, it was observed that clinical effects in humans correlated with the high-affinity polymorphic allotype (V158) and the low-affinity polymorphic allotype (F158) of FcγRIIIa (Non-Patent Document 19). These reports suggest that antibodies with an Fc region optimized for binding to a specific Fcγ receptor mediates stronger effector functions, and thereby exert more effective antitumor effects. The balance between the affinity of antibodies against the
35 activating receptors including FcγRIa, FcγRIIa, FcγRIIIa, and FcγRIIIb, and the inhibitory receptors including FcγRIIb is an important factor in optimizing antibody effector functions.

Enhancing the affinity to activating receptors may give antibodies a property to mediate stronger effector functions (Non-Patent Document 20), and therefore has been reported in various reports to date as an antibody engineering technique for improving or enhancing the antitumor activity of antibody pharmaceuticals against cancer antigens.

5 Regarding binding between the Fc region and Fcγ receptor, several amino acid residues in the antibody hinge region and the CH2 domain, and a sugar chain added to Asn at position 297 (EU numbering) bound to the CH2 domain have been shown as being important (Non-Patent Documents 13, 21, and 22). Focusing on this binding site, studies have so far been carried out on mutants of the Fc region having various Fcγ receptor binding properties, and Fc region
10 mutants with higher affinity to activating Fcγ receptor have been obtained (Patent Documents 2 and 3). For example, Lazar *et al.* have succeeded in increasing the binding of human IgG1 to human FcγRIIIa (V158) by approximately 370 fold by substituting Ser at position 239, Ala at position 330, and Ile at position 332 (EU numbering) of human IgG1 with Asp, Leu, and Glu, respectively (Non-Patent Document 19 and Patent Document 2). The ratio of binding to
15 FcγRIIIa and FcγRIIb (A/I ratio) for this mutant was approximately 9-fold that of the wild type. Furthermore, Shinkawa *et al.* have succeeded in increasing the binding to FcγRIIIa up to approximately 100 fold by removing fucose from the sugar chain added to Asn at position 297 (EU numbering) (Non-Patent Document 24). These methods can greatly improve the ADCC activity of human IgG1 compared to that of naturally-occurring human IgG1.

20 Thus, since Fcγ-receptor-binding activity plays an important role in cytotoxic activity in antibodies targeting membrane-type antigens, an isotype of human IgG1 with high FcγR-binding activity is used when cytotoxic activity is needed. Improvement of cytotoxic activity by enhancing Fcγ-receptor-binding activity is also widely used technique. On the other hand, the role played by Fcγ-receptor-binding activity in antibodies targeting soluble antigens is not known,
25 and it has been thought that there is no difference between human IgG1 with high Fcγ-receptor-binding activity and human IgG2 and human IgG4 with low FcγR-binding activity. Therefore, to date, enhancement of Fcγ-receptor-binding activity has not been attempted for antibodies targeting soluble antigens, and their effects have not been reported.

[Prior Art Documents]

30 Patent Documents

[Patent Document 1] WO 2009/125825

[Patent Document 2] WO 2000/042072

[Patent Document 3] WO 2006/019447

35 Non-patent Documents

[Non-patent Document 1] Janice M Reichert, Clark J Rosensweig, Laura B Faden & Matthew C

- Dewitz, Monoclonal antibody successes in the clinic., *Nat. Biotechnol.* (2005) 23, 1073-1078
- [Non-patent Document 2] Pavlou AK, Belsey MJ., The therapeutic antibodies market to 2008., *Eur J Pharm Biopharm.* (2005) 59 (3), 389-396
- [Non-patent Document 3] Kim SJ, Park Y, Hong HJ., Antibody engineering for the development
 5 of therapeutic antibodies., *Mol Cells.* (2005) 20 (1), 17-29
- [Non-patent Document 4] Hinton PR, Xiong JM, Johlfs MG, Tang MT, Keller S, Tsurushita N.,
 An engineered human IgG1 antibody with longer serum half-life., *J. Immunol.* (2006) 176 (1),
 346-356
- [Non-patent Document 5] Ghetie V, Popov S, Borvak J, Radu C, Matesoi D, Medesan C, Ober
 10 RJ, Ward ES., Increasing the serum persistence of an IgG fragment by random mutagenesis., *Nat. Biotechnol.* (1997) 15 (7), 637-640
- [Non-patent Document 6] Rajpal A, Beyaz N, Haber L, Cappuccilli G, Yee H, Bhatt RR,
 Takeuchi T, Lerner RA, Crea R., A general method for greatly improving the affinity of
 antibodies by using combinatorial libraries., *Proc. Natl. Acad. Sci. U. S. A.* (2005) 102 (24),
 15 8466-8471
- [Non-patent Document 7] Wu H, Pfarr DS, Johnson S, Brewah YA, Woods RM, Patel NK, White
 WI, Young JF, Kiener PA., Development of Motavizumab, an Ultra-potent Antibody for the
 Prevention of Respiratory Syncytial Virus Infection in the Upper and Lower Respiratory Tract., *J. Mol. Biol.* (2007) 368, 652-665
- [Non-patent Document 8] Hanson CV, Nishiyama Y, Paul S., Catalytic antibodies and their
 20 applications., *Curr Opin Biotechnol.* (2005) 16 (6), 631-636
- [Non-patent Document 9] Rathanaswami P, Roalstad S, Roskos L, Su QJ, Lackie S, Babcook J.,
 Demonstration of an in vivo generated sub-picomolar affinity fully human monoclonal antibody
 to interleukin-8., *Biochem. Biophys. Res. Commun.* (2005) 334 (4), 1004-1013
- [Non-patent Document 10] Dall'Acqua WF, Woods RM, Ward ES, Palaszynski SR, Patel NK,
 25 Brewah YA, Wu H, Kiener PA, Langermann S., Increasing the affinity of a human IgG1 for the
 neonatal Fc receptor: biological consequences., *J. Immunol.* (2002) 169 (9), 5171-5180
- [Non-patent Document 11] Yeung YA, Leabman MK, Marvin JS, Qiu J, Adams CW, Lien S,
 Starovasnik MA, Lowman HB., Engineering human IgG1 affinity to human neonatal Fc
 30 receptor: impact of affinity improvement on pharmacokinetics in primates., *J. Immunol.* (2009)
 182 (12), 7663-7671
- [Non-patent Document 12] Datta-Mannan A, Witcher DR, Tang Y, Watkins J, Wroblewski VJ.,
 Monoclonal antibody clearance. Impact of modulating the interaction of IgG with the neonatal
 Fc receptor., *J. Biol. Chem.* (2007) 282 (3), 1709-1717
- [Non-patent Document 13] Clark, M., Antibody Engineering IgG Effector Mechanisms.,
 35 *Chemical Immunology* (1997) 65, 88-110

[Non-patent Document 14] Horton HM, Bennett MJ, Pong E, Peipp M, Karki S, Chu SY, Richards JO, Vostiar I, Joyce PF, Repp R, Desjarlais JR, Zhukovsky EA., Potent in vitro and in vivo activity of an Fc-engineered anti-CD19 monoclonal antibody against lymphoma and leukemia., *Cancer Res.* (2008) 68, 8049-8057

- 5 [Non-patent Document 15] Zalevsky J, Leung IW, Karki S, Chu SY, Zhukovsky EA, Desjarlais JR, Carmichael DF, Lawrence CE., The impact of Fc engineering on an anti-CD19 antibody: increased Fcγ receptor affinity enhances B-cell clearing in nonhuman primates., *Blood* (2009) 113, 3735-3743

- [Non-patent Document 16] Jefferis R, Lund J., Interaction sites on human IgG-Fc for FcγR: current models., *Immunol. Lett.* (2002) 82, 57-65

[Non-patent Document 17] Clynes, R., Yoshizumi, T., Moroi, Y., Houghton, A.N., and Ravetch, J.V., Fc Receptors are required for passive and active immunity to melanoma., *Proc. Natl. Acad. Sci. U. S. A.* (1998) 95, 652-656

- 15 [Non-patent Document 18] Clynes RA, Towers TL, Presta LG, Ravetch JV., Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets., *Nat. Med.* (2000) 6, 443-446

[Non-patent Document 19] Cartron G, Dacheux L, Salles G, Solal-Celigny P, Bardos P, Colombat P, Watier H., Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor FcγRIIIa gene., *Blood* (2002) 99, 754-758

- [Non-patent Document 20] Nimmerjahn F, Ravetch JV., Divergent immunoglobulin g subclass activity through selective Fc receptor binding., *Science* (2005) 310, 1510-1512

[Non-patent Document 21] Greenwood J, Clark M, Waldmann H., Structural motifs involved in human IgG antibody effector functions., *Eur. J. Immunol.* (1993) 23, 1098-1104

- [Non-patent Document 22] Morgan A, Jones ND, Nesbitt AM, Chaplin L, Bodmer MW, Emtage JS., The N-terminal end of the CH2 domain of chimeric human IgG1 anti-HLA-DR is necessary for C1q, Fc γRI and Fc γRIII binding., *Immunology* (1995) 86, 319-324

25 [Non-patent Document 23] Lazar GA, Dang W, Karki S, Vafa O, Peng JS, Hyun L, Chan C, Chung HS, Eivazi A, Yoder SC, Vielmetter J, Carmichael DF, Hayes RJ, Dahiyat BI., Engineered antibody Fc variants with enhanced effector function., *Proc. Nat. Acad. Sci. U. S. A.* (2006) 103, 4005-4010

- 30 [Non-patent Document 24] Shinkawa T, Nakamura K, Yamane N, Shoji-Hosaka E, Kanda Y, Sakurada M, Uchida K, Anazawa H, Satoh M, Yamasaki M, Hanai N, Shitara K., The absence of fucose but not the presence of galactose or bisecting N-acetylglucosamine of human IgG1 complex-type oligosaccharides shows the critical role of enhancing antibody-dependent cellular cytotoxicity., *J. Biol. Chem.* (2003) 278, 3466-3473

35

Summary of the Invention

[Problems to be Solved by the Invention]

The present invention was achieved in view of the above circumstances. An objective of the present invention is to provide antigen-binding molecule with enhanced intracellular uptake of a bound antigen, antigen-binding molecules in which the number of antigens that can be bound per single molecule is increased, antigen-binding molecules with improved
 5 pharmacokinetics, antigen-binding molecules with enhanced intracellular dissociation of extracellularly bound antigen, antigen-binding molecules with enhanced extracellular release in the antigen-unbound state, antigen-binding molecules having the function of decreasing the total antigen concentration or the free antigen concentration in plasma, pharmaceutical compositions
 10 comprising such an antigen-binding molecules, and methods for producing them.

[Means for Solving the Problems]

As a result of conducting dedicated research to accomplish the above-mentioned objectives, the present inventors created an antigen-binding molecule containing an
 15 antigen-binding domain having human-FcRn-binding activity in an acidic pH range condition and in which the antigen-binding activity of an antigen-binding molecule changes depending on ion-concentration, and an Fc γ receptor-binding domain having a binding activity higher to the Fc γ receptor in a neutral pH range condition than the Fc γ -receptor-binding domain of an Fc region of a native human IgG in which the sugar chain bonded at position 297 (EU numbering) is
 20 a fucose-containing sugar chain. Furthermore, the present inventors discovered a method for enhancing intracellular uptake of bound antigens, a method for increasing the number of antigens that can bind to a single antigen-binding molecule, a method for improving pharmacokinetics of an antigen-binding molecule, a method for promoting intracellular dissociation of an antigen, which is extracellularly bound to the antigen-binding molecule, from an antigen-binding
 25 molecule, a method for promoting extracellular release of the antigen-binding molecule not bound to an antigen, and a method for decreasing total antigen concentration or free antigen concentration in plasma, wherein the methods comprises contacting the antigen-binding molecule with an Fc γ -receptor-expressing cell *in vivo* or *in vitro*. Furthermore, the inventors discovered methods for producing antigen-binding molecules having the above-mentioned
 30 properties, and also discovered the utility of pharmaceutical compositions containing, as an active ingredient, such an antigen-binding molecule or an antigen-binding molecule produced by the production method of the present invention, and thereby completed the present invention.

That is, more specifically, the present invention provides [1] to [46] below:

[1] A pharmaceutical composition which comprises an antigen-binding molecule comprising an
 35 antigen-binding domain and an Fc γ -receptor-binding domain, wherein the antigen-binding molecule has human-FcRn-binding activity in an acidic pH range condition, and wherein the

antigen-binding domain has antigen-binding activity that changes depending on an ion-concentration condition, and the Fc γ -receptor-binding domain has higher binding activity to the Fc γ receptor in a neutral pH range condition than an Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain.

[2] The pharmaceutical composition of [1], wherein the antigen is a soluble antigen.

[3] The pharmaceutical composition of [1] or [2], wherein the ion concentration is calcium ion concentration.

[4] The pharmaceutical composition of [3], wherein the antigen-binding domain is an antigen-binding domain in which binding activity to the antigen under a high-calcium-ion concentration condition is higher than that under a low-calcium-ion concentration condition.

[5] The pharmaceutical composition of [1] or [2], wherein the ion-concentration condition is a pH condition.

[6] The pharmaceutical composition of [5], wherein the antigen-binding domain is an antigen-binding domain in which binding activity to the antigen in a neutral pH range condition is higher than that in an acidic pH range condition.

[7] The pharmaceutical composition of any one of [1] to [6], wherein the antigen-binding molecule has neutralizing activity against the antigen.

[8] The pharmaceutical composition of any one of [1] to [7], wherein the Fc γ receptor-binding domain comprises an antibody Fc region.

[9] The pharmaceutical composition of [8], wherein the Fc region is an Fc region in which at least one or more amino acids selected from the group consisting of amino acids at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 in the Fc region site according to EU numbering are different from amino acids at corresponding sites in a native Fc region.

[10] The pharmaceutical composition of [9], wherein the Fc region is an Fc region which comprises at least one or more amino acids selected from the group consisting of:
either Lys or Tyr at amino acid position 221;

any one of Phe, Trp, Glu, and Tyr at amino acid position 222;

any one of Phe, Trp, Glu, and Lys at amino acid position 223;

any one of Phe, Trp, Glu, and Tyr at amino acid position 224;

- any one of Glu, Lys, and Trp at amino acid position 225;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 227;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 228;
- any one of Ala, Glu, Gly, and Tyr at amino acid position 230;
- 5 any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 232;
- any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 233;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
- 10 Tyr at amino acid position 234;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 235;
- any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 236;
- 15 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 237;
- any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 238;
- any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 239;
- 20 any one of Ala, Ile, Met, and Thr at amino acid position 240;
- any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;
- any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;
- His at amino acid position 244;
- 25 Ala at amino acid position 245;
- any one of Asp, Glu, His, and Tyr at amino acid position 246;
- any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;
- any one of Glu, His, Gln, and Tyr at amino acid position 249;
- either Glu or Gln at amino acid position 250;
- 30 Phe at amino acid position 251;
- any one of Phe, Met, and Tyr at amino acid position 254;
- any one of Glu, Leu, and Tyr at amino acid position 255;
- any one of Ala, Met, and Pro at amino acid position 256;
- any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;
- 35 any one of Asp, Glu, His, and Tyr at amino acid position 260;
- any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;

- any one of Ala, Ile, Met, and Thr at amino acid position 263;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr
 at amino acid position 264;
 any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 5 Tyr at amino acid position 265;
 any one of Ala, Ile, Met, and Thr at amino acid position 266;
 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at
 amino acid position 267;
 any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid
 10 position 268;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 269;
 any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid
 position 270;
 15 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 271;
 any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 272;
 either Phe or Ile at amino acid position 273;
 20 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at
 amino acid position 274;
 either Leu or Trp at amino acid position 275;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 276;
 25 any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at
 amino acid position 278;
 Ala at amino acid position 279;
 any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;
 any one of Asp, Lys, Pro, and Tyr at amino acid position 281;
 30 any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;
 any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;
 any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;
 any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;
 any one of Glu, Gly, Pro, and Tyr at amino acid position 286;
 35 any one of Asn, Asp, Glu, and Tyr at amino acid position 288;
 any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;

- any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;
 any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;
 any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 293;
- 5 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 294;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 295;
 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino acid position 296;
- 10 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 297;
 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 298;
- 15 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 299;
 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 300;
 any one of Asp, Glu, His, and Tyr at amino acid position 301;
- 20 Ile at amino acid position 302;
 any one of Asp, Gly, and Tyr at amino acid position 303;
 any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;
 any one of Glu, Ile, Thr, and Tyr at amino acid position 305;
 any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;
- 25 Phe at amino acid position 313;
 Leu at amino acid position 315;
 either Glu or Gln at amino acid position 317;
 any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;
 any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 320;
- 30 any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 322;
 Ile at amino acid position 323;
 any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 324;
- 35 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and

- Tyr at amino acid position 325;
 any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino acid position 326;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr
 5 at amino acid position 327;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 328;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 329;
 10 any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 330;
 any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 331;
 any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp,
 15 and Tyr at amino acid position 332;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid position 333;
 any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;
 any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid
 20 position 335;
 any one of Glu, Lys, and Tyr at amino acid position 336;
 any one of Glu, His, and Asn at amino acid position 337;
 any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;
 either Ala or Val at amino acid position 376;
 25 either Gly or Lys at amino acid position 377;
 Asp at amino acid position 378;
 Asn at amino acid position 379;
 any one of Ala, Asn, and Ser at amino acid position 380;
 either Ala or Ile at amino acid position 382;
 30 Glu at amino acid position 385;
 Thr at amino acid position 392;
 Leu at amino acid position 396;
 Lys at amino acid position 421;
 Asn at amino acid position 427;
 35 either Phe or Leu at amino acid position 428;
 Met at amino acid position 429;

Trp at amino acid position 434;

Ile at amino acid position 436; and

any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440,

in the Fc region site according to EU numbering.

- 5 [11] The pharmaceutical composition of any one of [1] to [10], wherein the Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, is an Fc region of any one of native human IgG1, native human IgG2, native human IgG3, and native human IgG4 in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain.

- 10 [12] The pharmaceutical composition of any one of [1] to [11], wherein the human Fcγ receptor is FcγRIa, FcγRIIa(R), FcγRIIa(H), FcγRIIb, FcγRIIIa(V), or FcγRIIIa(F).

[13] The pharmaceutical composition of any one of [1] to [11], wherein the human Fcγ receptor is FcγRIIb.

- [14] The pharmaceutical composition of any one of [8] to [13], wherein the Fc region is an Fc
15 region which comprises at least one or more of

Asp at amino acid position 238, and

Glu at amino acid position 328

in the Fc region site according to EU numbering.

- [15] A method comprising the step of contacting an antigen-binding molecule with an
20 Fcγ-receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human-FcRn-binding activity in an acidic pH range condition and comprises an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition and an Fcγ-receptor-binding domain that has higher binding activity to the Fcγ receptor in a neutral pH range condition compared to an Fc region of a native human IgG in which the sugar
25 chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, which is a method of any one of:

(i) a method for increasing the number of antigens that can bind to a single antigen-binding molecule;

(ii) a method for eliminating plasma antigens;

- 30 (iii) a method for improving antigen-binding molecule pharmacokinetics;

(iv) a method for promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule;

(v) a method for promoting extracellular release of an antigen-binding molecule not bound to an antigen; and

- 35 (vi) a method for decreasing a total antigen concentration or free antigen concentration in plasma.

- [16] The method of [15], wherein the antigen is a soluble antigen.
- [17] The method of [15] or [16], wherein the ion concentration is a calcium ion concentration.
- [18] The method of [17], wherein the antigen-binding domain is an antigen-binding domain in which binding activity to the antigen under a high-calcium-ion concentration condition is higher than that under a low-calcium-ion concentration condition.
- [19] The method of [15] or [16], wherein the ion concentration condition is a pH condition.
- [20] The method of [19], wherein the antigen-binding domain is an antigen-binding domain in which binding activity to the antigen in a neutral pH range condition is higher than that in an acidic pH range condition.
- [21] The method of any one of [15] to [20], wherein the antigen-binding molecule has neutralizing activity against the antigen.
- [22] The method of any one of [15] to [21], wherein the Fc γ receptor-binding domain comprises an antibody Fc region.
- [23] The method of [22], wherein the Fc region is an Fc region in which at least one or more amino acids selected from the group consisting of amino acids at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 in the Fc region site according to EU numbering are different from the amino acids at corresponding sites in the native Fc region.
- [24] The method of [23], wherein the Fc region is an Fc region which comprises at least one or more amino acids selected from the group consisting of:
- either Lys or Tyr at amino acid position 221;
 - any one of Phe, Trp, Glu, and Tyr at amino acid position 222;
 - any one of Phe, Trp, Glu, and Lys at amino acid position 223;
 - any one of Phe, Trp, Glu, and Tyr at amino acid position 224;
 - any one of Glu, Lys, and Trp at amino acid position 225;
 - any one of Glu, Gly, Lys, and Tyr at amino acid position 227;
 - any one of Glu, Gly, Lys, and Tyr at amino acid position 228;
 - any one of Ala, Glu, Gly, and Tyr at amino acid position 230;
 - any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;
 - any one of Glu, Gly, Lys, and Tyr at amino acid position 232;
 - any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr

- at amino acid position 233;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 234;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 5 Tyr at amino acid position 235;
 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 236;
 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 237;
 10 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 238;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr
 at amino acid position 239;
 any one of Ala, Ile, Met, and Thr at amino acid position 240;
 15 any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;
 any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;
 His at amino acid position 244;
 Ala at amino acid position 245;
 any one of Asp, Glu, His, and Tyr at amino acid position 246;
 20 any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;
 any one of Glu, His, Gln, and Tyr at amino acid position 249;
 either Glu or Gln at amino acid position 250;
 Phe at amino acid position 251;
 any one of Phe, Met, and Tyr at amino acid position 254;
 25 any one of Glu, Leu, and Tyr at amino acid position 255;
 any one of Ala, Met, and Pro at amino acid position 256;
 any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;
 any one of Asp, Glu, His, and Tyr at amino acid position 260;
 any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;
 30 any one of Ala, Ile, Met, and Thr at amino acid position 263;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr
 at amino acid position 264;
 any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 265;
 35 any one of Ala, Ile, Met, and Thr at amino acid position 266;
 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at

- amino acid position 267;
 any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid position 268;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 269;
 5 any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 270;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 271;
 10 any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 272;
 either Phe or Ile at amino acid position 273;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 274;
 15 either Leu or Trp at amino acid position 275;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 276;
 any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 278;
 20 Ala at amino acid position 279;
 any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;
 any one of Asp, Lys, Pro, and Tyr at amino acid position 281;
 any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;
 any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;
 25 any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;
 any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;
 any one of Glu, Gly, Pro, and Tyr at amino acid position 286;
 any one of Asn, Asp, Glu, and Tyr at amino acid position 288;
 any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;
 30 any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;
 any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;
 any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 293;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 294;
 35 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at

- amino acid position 295;
 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino acid position 296;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 5 at amino acid position 297;
 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 298;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 299;
 10 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 300;
 any one of Asp, Glu, His, and Tyr at amino acid position 301;
 Ile at amino acid position 302;
 any one of Asp, Gly, and Tyr at amino acid position 303;
 15 any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;
 any one of Glu, Ile, Thr, and Tyr at amino acid position 305;
 any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;
 Phe at amino acid position 313;
 Leu at amino acid position 315;
 20 either Glu or Gln at amino acid position 317;
 any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;
 any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 320;
 any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position
 25 322;
 Ile at amino acid position 323;
 any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 324;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 30 Tyr at amino acid position 325;
 any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino acid position 326;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 327;
 35 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 328;

- any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 329;
- any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 330;
- 5 any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 331;
- any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 332;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid
- 10 position 333;
- any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;
- any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid position 335;
- any one of Glu, Lys, and Tyr at amino acid position 336;
- 15 any one of Glu, His, and Asn at amino acid position 337;
- any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;
- either Ala or Val at amino acid position 376;
- either Gly or Lys at amino acid position 377;
- Asp at amino acid position 378;
- 20 Asn at amino acid position 379;
- any one of Ala, Asn, and Ser at amino acid position 380;
- either Ala or Ile at amino acid position 382;
- Glu at amino acid position 385;
- Thr at amino acid position 392;
- 25 Leu at amino acid position 396;
- Lys at amino acid position 421;
- Asn at amino acid position 427;
- either Phe or Leu at amino acid position 428;
- Met at amino acid position 429;
- 30 Trp at amino acid position 434;
- Ile at amino acid position 436; and
- any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440,
- in the Fc region site according to EU numbering.
- [25] The method of any one of [15] to [24], wherein the Fc region of a native human IgG in
- 35 which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, is an Fc region of any one of native human IgG1, native human IgG2, native human

IgG3, and native human IgG4 in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain.

[26] The method of any one of [15] to [25], wherein the human Fc γ receptor is Fc γ RIa, Fc γ RIIa(R), Fc γ RIIa(H), Fc γ RIIb, Fc γ RIIIa(V), or Fc γ RIIIa(F).

5 [27] The method of any one of [15] to [25], wherein the human Fc γ receptor is Fc γ RIIb.

[28] The method of any one of [22] to [27], wherein the Fc region is an Fc region which comprises at least one or more of

Asp at amino acid position 238, and

Glu at amino acid position 328

10 in the Fc region site according to EU numbering.

[29] A method comprising the step of enhancing Fc γ -receptor-binding activity in a neutral pH range condition of the Fc γ -receptor-binding domain in an antigen-binding molecule compared to that of a native human IgG Fc region in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, wherein the antigen-binding molecule has
15 human-FcRn-binding activity in an acidic pH range condition and comprises an Fc γ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition, which is a method of any one of:

(i) a method for altering an antigen-binding molecule, wherein the intracellular uptake of the antigen to which it binds is enhanced;

20 (ii) a method for increasing the number of antigens that can bind to a single molecule of antigen-binding molecule;

(iii) a method for increasing the ability of an antigen-binding molecule to eliminate plasma antigens;

(iv) a method for improving antigen-binding molecule pharmacokinetics;

25 (v) a method for promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule;

(vi) a method for promoting extracellular release of an antigen-binding molecule not bound to an antigen, wherein the antigen-binding molecule had been taken up into a cell in an antigen-bound form; and

30 (vii) a method for altering an antigen-binding molecule, which can decrease a total antigen concentration or free antigen concentration in plasma.

[30] The method of [29], wherein the antigen is a soluble antigen.

[31] The method of [29] or [30], wherein the ion concentration is a calcium ion concentration.

[32] The method of [31], wherein the antigen-binding domain is an antigen-binding domain in
35 which binding activity to the antigen under high calcium ion concentration conditions is higher than that under low calcium ion concentration conditions.

[33] The method of [29] or [30], wherein the ion concentration condition is a pH condition.

[34] The method of [33], wherein the antigen-binding domain is an antigen-binding domain in which binding activity to the antigen in a neutral pH range condition is higher than that in an acidic pH range condition.

5 [35] The method of any one of [29] to [34], wherein the antigen-binding molecule has neutralizing activity against the antigen.

[36] The method of any one of [29] to [35], wherein the Fcγ receptor-binding domain comprises an antibody Fc region.

[37] The method of [36], wherein the Fc region is an Fc region in which at least one or more
 10 amino acids selected from the group consisting of amino acids at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322,
 15 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 in the Fc region site according to EU numbering are different from the amino acids at corresponding sites in the native Fc region.

[38] The method of [33], wherein the Fc region is an Fc region comprising at least one or more
 20 amino acids selected from the group consisting of:

either Lys or Tyr at amino acid position 221;

any one of Phe, Trp, Glu, and Tyr at amino acid position 222;

any one of Phe, Trp, Glu, and Lys at amino acid position 223;

any one of Phe, Trp, Glu, and Tyr at amino acid position 224;

25 any one of Glu, Lys, and Trp at amino acid position 225;

any one of Glu, Gly, Lys, and Tyr at amino acid position 227;

any one of Glu, Gly, Lys, and Tyr at amino acid position 228;

any one of Ala, Glu, Gly, and Tyr at amino acid position 230;

any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;

30 any one of Glu, Gly, Lys, and Tyr at amino acid position 232;

any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 233;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 234;

35 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 235;

- Tyr at amino acid position 236;
 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 237;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 5 at amino acid position 238;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr
 at amino acid position 239;
 any one of Ala, Ile, Met, and Thr at amino acid position 240;
 any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;
 10 any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;
 His at amino acid position 244;
 Ala at amino acid position 245;
 any one of Asp, Glu, His, and Tyr at amino acid position 246;
 any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;
 15 any one of Glu, His, Gln, and Tyr at amino acid position 249;
 either Glu or Gln at amino acid position 250;
 Phe at amino acid position 251;
 any one of Phe, Met, and Tyr at amino acid position 254;
 any one of Glu, Leu, and Tyr at amino acid position 255;
 20 any one of Ala, Met, and Pro at amino acid position 256;
 any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;
 any one of Asp, Glu, His, and Tyr at amino acid position 260;
 any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;
 any one of Ala, Ile, Met, and Thr at amino acid position 263;
 25 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr
 at amino acid position 264;
 any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 265;
 any one of Ala, Ile, Met, and Thr at amino acid position 266;
 30 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at
 amino acid position 267;
 any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid
 position 268;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 35 acid position 269;
 any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid

- position 270;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 271;
any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 272;
5 either Phe or Ile at amino acid position 273;
any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 274;
either Leu or Trp at amino acid position 275;
10 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 276;
any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 278;
Ala at amino acid position 279;
15 any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;
any one of Asp, Lys, Pro, and Tyr at amino acid position 281;
any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;
any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;
any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;
20 any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;
any one of Glu, Gly, Pro, and Tyr at amino acid position 286;
any one of Asn, Asp, Glu, and Tyr at amino acid position 288;
any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;
any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;
25 any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;
any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 293;
any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 294;
30 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 295;
any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino acid position 296;
any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 297;
35 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino

- acid position 298;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 299;
any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp
5 at amino acid position 300;
any one of Asp, Glu, His, and Tyr at amino acid position 301;
Ile at amino acid position 302;
any one of Asp, Gly, and Tyr at amino acid position 303;
any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;
10 any one of Glu, Ile, Thr, and Tyr at amino acid position 305;
any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;
Phe at amino acid position 313;
Leu at amino acid position 315;
either Glu or Gln at amino acid position 317;
15 any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;
any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 320;
any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 322;
20 Ile at amino acid position 323;
any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 324;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 325;
25 any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino acid position 326;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 327;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 328;
30 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 329;
any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 330;
35 any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 331;

any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 332;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid position 333;

5 any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;

any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid position 335;

any one of Glu, Lys, and Tyr at amino acid position 336;

any one of Glu, His, and Asn at amino acid position 337;

10 any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;

either Ala or Val at amino acid position 376;

either Gly or Lys at amino acid position 377;

Asp at amino acid position 378;

Asn at amino acid position 379;

15 any one of Ala, Asn, and Ser at amino acid position 380;

either Ala or Ile at amino acid position 382;

Glu at amino acid position 385;

Thr at amino acid position 392;

Leu at amino acid position 396;

20 Lys at amino acid position 421;

Asn at amino acid position 427;

either Phe or Leu at amino acid position 428;

Met at amino acid position 429;

Trp at amino acid position 434;

25 Ile at amino acid position 436; and

any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440,

in the Fc region site according to EU numbering.

[39] The method of any one of [29] to [38], wherein the Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, is an Fc region of any one of native human IgG1, native human IgG2, native human
30 IgG3, and native human IgG4 in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain.

[40] The method of any one of [29] to [39], wherein the human Fcγ receptor is FcγRIa, FcγRIIa(R), FcγRIIa(H), FcγRIIb, FcγRIIIa(V), or FcγRIIIa(F).

35 [41] The method of any one of [29] to [39], wherein the human Fcγ receptor is FcγRIIb.

[42] The method of any one of [36] to [41], wherein the Fc region is an Fc region which

comprises at least one or more of:

Asp at amino acid position 238, and

Glu at amino acid position 328

in the Fc region site according to EU numbering.

- 5 [43] A method for producing an antigen-binding molecule, which comprises the steps of:
- (a) determining the antigen-binding activity of an antigen-binding domain under a high-calcium-ion concentration condition;
 - (b) determining the antigen-binding activity of an antigen-binding domain under a low-calcium-ion concentration condition;
 - 10 (c) selecting the antigen-binding domain for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
 - (d) linking a polynucleotide encoding the antigen-binding domain selected in (c) to a polynucleotide encoding an Fc γ receptor-binding domain having human-FcRn-binding activity in an acidic pH range condition and in which binding activity to the Fc γ receptor in a neutral pH
 - 15 range condition is higher than that of an Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain;
 - (e) culturing cells introduced with a vector in which the polynucleotide obtained in (d) is operably linked; and
 - (f) collecting antigen-binding molecules from the cell culture of (e).
- 20 [44] A method for producing an antigen-binding molecule, which comprises the steps of:
- (a) determining the antigen-binding activity of an antibody under a high-calcium-ion concentration condition;
 - (b) determining the antigen-binding activity of an antibody under a low-calcium-ion concentration condition;
 - 25 (c) selecting the antibody for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
 - (d) linking a polynucleotide encoding the antigen-binding domain of the antibody selected in (c) to a polynucleotide encoding an Fc γ receptor-binding domain having human-FcRn-binding activity in an acidic pH range, and in which binding activity to the Fc γ receptor in a neutral pH
 - 30 range condition is higher than that of an Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain;
 - (e) culturing cells introduced with a vector in which the polynucleotide obtained in (d) is operably linked; and
 - (f) collecting antigen-binding molecules from the cell culture of (e).
- 35 [45] A method for producing an antigen-binding molecule, which comprises the steps of:
- (a) determining the antigen-binding activity of an antigen-binding domain in a neutral pH range

condition;

(b) determining the antigen-binding activity of an antigen-binding domain in an acidic pH range condition;

- (c) selecting the antigen-binding domain for which the antigen-binding activity determined in
 5 (a) is higher than the antigen-binding activity determined in (b);
 (d) linking a polynucleotide encoding the antigen-binding domain selected in (c) to a polynucleotide encoding an Fc γ receptor-binding domain having human-FcRn-binding activity in an acidic pH range condition, and in which binding activity to the Fc γ receptor in a neutral pH range condition is higher than that of an Fc region of a native human IgG in which the sugar
 10 chain bound at position 297 according to EU numbering is a fucose-containing sugar chain;
 (e) culturing cells introduced with a vector in which the polynucleotide obtained in (d) is operably linked; and
 (f) collecting antigen-binding molecules from the cell culture of (c).

[46] A method for producing an antigen-binding molecule, which comprises the steps of:

- 15 (a) determining the antigen-binding activity of an antibody in a neutral pH range condition;
 (b) determining the antigen-binding activity of an antibody in an acidic pH range condition;
 (c) selecting the antibody for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
 (d) linking a polynucleotide encoding the antigen-binding domain of the antibody selected in
 20 (c) to a polynucleotide encoding an Fc γ receptor-binding domain having human-FcRn-binding activity in an acidic pH range condition, in which binding activity to the Fc γ receptor in a neutral pH range condition is higher than that of an Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain;
 (e) culturing cells introduced with a vector in which the polynucleotide obtained in (d) is
 25 operably linked; and
 (f) collecting antigen-binding molecules from the cell culture of (e).

[47] The production method of any one of [43] to [46], wherein the antigen is a soluble antigen.

[48] The production method of any one of [43] to [47], wherein the Fc γ receptor-binding domain comprises an antibody Fc region.

- 30 [49] The production method of [48], wherein the Fc region is an Fc region in which at least one or more amino acids selected from the group consisting of amino acids at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292,
 35 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377,

378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 in the Fc region site according to EU numbering are different from the amino acids at corresponding sites in the native Fc region.

- [50] The production method of [49], wherein the Fc region comprises at least one or more amino acids selected from the group consisting of:
- 5 either Lys or Tyr at amino acid position 221;
 - any one of Phe, Trp, Glu, and Tyr at amino acid position 222;
 - any one of Phe, Trp, Glu, and Lys at amino acid position 223;
 - any one of Phe, Trp, Glu, and Tyr at amino acid position 224;
 - 10 any one of Glu, Lys, and Trp at amino acid position 225;
 - any one of Glu, Gly, Lys, and Tyr at amino acid position 227;
 - any one of Glu, Gly, Lys, and Tyr at amino acid position 228;
 - any one of Ala, Glu, Gly, and Tyr at amino acid position 230;
 - any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;
 - 15 any one of Glu, Gly, Lys, and Tyr at amino acid position 232;
 - any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 233;
 - any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 234;
 - 20 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 235;
 - any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 236;
 - any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 237;
 - 25 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 238;
 - any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 239;
 - 30 any one of Ala, Ile, Met, and Thr at amino acid position 240;
 - any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;
 - any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;
 - His at amino acid position 244;
 - Ala at amino acid position 245;
 - 35 any one of Asp, Glu, His, and Tyr at amino acid position 246;
 - any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;

- any one of Glu, His, Gln, and Tyr at amino acid position 249;
 either Glu or Gln at amino acid position 250;
 Phe at amino acid position 251;
 any one of Phe, Met, and Tyr at amino acid position 254;
 5 any one of Glu, Leu, and Tyr at amino acid position 255;
 any one of Ala, Met, and Pro at amino acid position 256;
 any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;
 any one of Asp, Glu, His, and Tyr at amino acid position 260;
 any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;
 10 any one of Ala, Ile, Met, and Thr at amino acid position 263;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr
 at amino acid position 264;
 any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 265;
 15 any one of Ala, Ile, Met, and Thr at amino acid position 266;
 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at
 amino acid position 267;
 any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid
 position 268;
 20 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 269;
 any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid
 position 270;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and
 25 Tyr at amino acid position 271;
 any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 272;
 either Phe or Ile at amino acid position 273;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at
 30 amino acid position 274;
 either Leu or Trp at amino acid position 275;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 276;
 any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at
 35 amino acid position 278;
 Ala at amino acid position 279;

- any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;
 any one of Asp, Lys, Pro, and Tyr at amino acid position 281;
 any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;
 any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;
 5 any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;
 any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;
 any one of Glu, Gly, Pro, and Tyr at amino acid position 286;
 any one of Asn, Asp, Glu, and Tyr at amino acid position 288;
 any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;
 10 any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;
 any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;
 any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 293;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 294;
 15 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 295;
 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino acid position 296;
 20 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 297;
 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 298;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 299;
 25 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 300;
 any one of Asp, Glu, His, and Tyr at amino acid position 301;
 Ile at amino acid position 302;
 30 any one of Asp, Gly, and Tyr at amino acid position 303;
 any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;
 any one of Glu, Ile, Thr, and Tyr at amino acid position 305;
 any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;
 Phe at amino acid position 313;
 35 Leu at amino acid position 315;
 either Glu or Gln at amino acid position 317;

- any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;
any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid
position 320;
any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position
5 322;
Ile at amino acid position 323;
any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid
position 324;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
10 Tyr at amino acid position 325;
any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino
acid position 326;
any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr
at amino acid position 327;
15 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
Tyr at amino acid position 328;
any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
at amino acid position 329;
any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr
20 at amino acid position 330;
any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position
331;
any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp,
and Tyr at amino acid position 332;
25 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid
position 333;
any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;
any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid
position 335;
30 any one of Glu, Lys, and Tyr at amino acid position 336;
any one of Glu, His, and Asn at amino acid position 337;
any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;
either Ala or Val at amino acid position 376;
either Gly or Lys at amino acid position 377;
35 Asp at amino acid position 378;
Asn at amino acid position 379;

- any one of Ala, Asn, and Ser at amino acid position 380;
 either Ala or Ile at amino acid position 382;
 Glu at amino acid position 385;
 Thr at amino acid position 392;
 5 Leu at amino acid position 396;
 Lys at amino acid position 421;
 Asn at amino acid position 427;
 either Phe or Leu at amino acid position 428;
 Met at amino acid position 429;
 10 Trp at amino acid position 434;
 Ile at amino acid position 436; and
 any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440,
 in the Fc region site according to EU numbering.
- [51] The production method of any one of [43] to [50], wherein the Fc γ receptor binding domain
 15 is an Fc region of any one of native human IgG1, native human IgG2, native human IgG3, and
 native human IgG4 in which the sugar chain bound at position 297 according to EU numbering
 is a fucose-containing sugar chain.
- [52] The production method of any one of [43] to [51], wherein the human Fc γ receptor is
 Fc γ RIa, Fc γ RIIa(R), Fc γ RIIa(H), Fc γ RIIb, Fc γ RIIIa(V), or Fc γ RIIIa(F).
- 20 [53] The production method of any one of [43] to [51], wherein the human Fc γ receptor is
 Fc γ RIIb.
- [54] The production method of any one of [48] to [53], wherein the Fc region comprises at least
 one or more amino acids of:
 Asp at amino acid position 238, and
 25 Glu at amino acid position 328
 in the Fc region site according to EU numbering.

Brief Description of the Drawings

Fig. 1 shows a non-limiting action mechanism for the elimination of soluble antigen
 30 from plasma by administering an antibody that binds to an antigen in an ion
 concentration-dependent manner and whose Fc γ receptor binding is enhanced at a neutral pH as
 compared to existing neutralizing antibodies.

Fig. 2 shows a time course of human IL-6 receptor concentration in the plasma of
 human FcRn transgenic mice administered with Fv4-IgG1 which binds to human IL-6 receptor
 35 in a pH-dependent manner or H54/L28-IgG1.

Fig. 3 shows a time course of human IL-6 receptor concentration in the plasma of

human FcRn transgenic mice administered with Fv4-IgG1 which binds to human IL-6 receptor in a pH-dependent manner, Fv4-IgG1-F760 which is an Fv4-IgG1 variant that lacks mouse Fc γ R binding, Fv4-IgG1-F1022 which is an Fv4-IgG1 variant with enhanced mouse Fc γ R binding, or Fv4-IgG1-Fuc which is an Fv4-IgG1 antibody with low fucose content.

5 Fig. 4 shows a time course of human IL-6 receptor concentration in the plasma of human FcRn transgenic mice administered with Fv4-IgG1 or antigen-binding molecules comprising as the heavy chain, Fv4-IgG1-F1022 or Fv4-IgG1-F1093 which is a Fv4-IgG1-F1022 variant with improved FcRn binding in an acidic pH range.

10 Fig. 5 shows a concentration time course of the administered antigen-binding molecules in the plasma of human FcRn transgenic mice administered with Fv4-IgG1 or antigen-binding molecules comprising as the heavy chain, Fv4-IgG1-F1022 or Fv4-IgG1-F1093 which is a Fv4-IgG1-F1022 variant with improved FcRn binding in an acidic pH range.

15 Fig. 6 shows a time course of human IL-6 receptor concentration in the plasma of human FcRn transgenic mice administered with Fv4-IgG1, Fv4-IgG1-F1087 which is an Fv4-IgG1 variant with enhanced mouse Fc γ R binding (in particular, enhanced mouse Fc γ RIIb binding and mouse Fc γ RIII binding), and Fv4-IgG1-F1182 which is an Fv4-IgG1 variant with enhanced mouse Fc γ R binding (in particular, enhanced mouse Fc γ RI binding and mouse Fc γ RIV binding).

20 Fig. 7 shows a concentration time course of the administered antigen-binding molecules in the plasma of human FcRn transgenic mice administered with Fv4-IgG1, Fv4-IgG1-F1087, and Fv4-IgG1-F1180 and Fv4-IgG1-F1412 which are Fv4-IgG1-F1087 variants with improved FcRn binding in an acidic pH range.

25 Fig. 8 shows a concentration time course of the administered antigen-binding molecules in the plasma of human FcRn transgenic mice administered with Fv4-IgG1, Fv4-IgG1-F1182, and Fv4-IgG1-F1181 which is an Fv4-IgG1-F1182 variant with improved FcRn binding in an acidic pH range.

30 Fig. 9 shows a time course of human IL-6 receptor concentration in the plasma of human FcRn transgenic mice administered with Fv4-IgG1, Fv4-IgG1-F1087, and Fv4-IgG1-F1180 and Fv4-IgG1-F1412 which are Fv4-IgG1-F1087 variants with improved FcRn binding in an acidic pH range.

Fig. 10 shows a time course of human IL-6 receptor concentration in the plasma of human FcRn transgenic mice administered with Fv4-IgG1, Fv4-IgG1-F1182, and Fv4-IgG1-F1181 which is an Fv4-IgG1-F1182 variant with improved FcRn binding in an acidic pH range.

35 Fig. 11 shows the results of change in plasma concentration of Fv4-IgG1, Fv4-IgG1-F1782, or Fv4-IgG1-F1087 in a human FcRn transgenic mouse when Fv4-IgG1,

Fv4-IgG1-F1782, or Fv4-IgG1-F1087 is administered to the mouse.

Fig. 12 shows the results of change in plasma concentration of a soluble human IL-6 receptor in a human FcRn transgenic mouse when Fv4-IgG1, Fv4-IgG1-F1782, or Fv4-IgG1-F1087 is administered to the mouse.

5 Fig. 13 shows a time course of human IL-6 receptor concentration in the plasma of normal mice administered with Fv4-mIgG1, Fv4-mIgG1-mF44 which is an Fv4-mIgG1 variant with enhanced mouse FcγRIIb binding and mouse FcγRIII binding, and Fv4-mIgG1-mF46 which is an Fv4-mIgG1 variant with further enhanced mouse FcγRIIb binding and mouse FcγRIII binding.

10 Fig. 14 shows a time course of human IL-6 receptor concentration in the plasma of FcγRIII-deficient mice administered with Fv4-mIgG1, Fv4-mIgG1-mF44 which is an Fv4-mIgG1 variant with enhanced mouse FcγRIIb binding and mouse FcγRIII binding, and Fv4-mIgG1-mF46 which is an Fv4-mIgG1 variant with further enhanced mouse FcγRIIb binding and mouse FcγRIII binding.

15 Fig. 15 shows a time course of human IL-6 receptor concentration in the plasma of Fc receptor γ chain-deficient mice administered with Fv4-mIgG1, Fv4-mIgG1-mF44 which is an Fv4-mIgG1 variant with enhanced mouse FcγRIIb binding and mouse FcγRIII binding, and Fv4-mIgG1-mF46 which is an Fv4-mIgG1 variant with further enhanced mouse FcγRIIb binding and mouse FcγRIII binding.

20 Fig. 16 shows a time course of human IL-6 receptor concentration in the plasma of FcγRIIb-deficient mice administered with Fv4-mIgG1, Fv4-mIgG1-mF44 which is an Fv4-mIgG1 variant with enhanced mouse FcγRIIb binding and mouse FcγRIII binding, and Fv4-mIgG1-mF46 which is an Fv4-mIgG1 variant with further enhanced mouse FcγRIIb binding and mouse FcγRIII binding.

25 Fig. 17 shows a result of evaluating the platelet aggregation ability of the omalizumab-G1d-v3/IgE immunocomplex by platelet aggregation assay using platelets derived from donors with FcγRIIa allotype (R/H).

Fig. 18 shows a result of evaluating the platelet aggregation ability of the omalizumab-G1d-v3/IgE immunocomplex by platelet aggregation assay using platelets derived from donors with FcγRIIa allotype (H/H).

30 Fig. 19 shows a result of assessing CD62p expression on the membrane surface of washed platelets. The black-filled area in the graph indicates a result of ADP stimulation after reaction with PBS. The area that is not filled in the graph indicates a result of ADP stimulation after reaction with the immunocomplex.

35 Fig. 20 shows a result of assessing the expression of active integrin on the membrane surface of washed platelets. The black-filled area in the graph indicates a result of ADP

stimulation after reaction with PBS. The area that is not filled in the graph indicates a result of ADP stimulation after reaction with the immunocomplex.

Fig. 21 shows the results of evaluating platelet aggregation activity induced by the omalizumab-BP230/IgE immunocomplex and the omalizumab-G1d-v3/IgE immunocomplex in a platelet aggregation assay using platelets derived from a donor with an FcγRIIa polymorphism (R/H).

Fig. 22 shows the results of evaluating CD62p expression on the surface of the membrane of washed platelets. The graph shaded with grey indicates the result when stimulation by adding ADP was performed after reaction with PBS, the solid line and the dotted line indicate the results when stimulation by ADP was performed after reaction with the omalizumab-G1d-v3/IgE immunocomplex and the omalizumab-BP230/IgE immunocomplex, respectively.

Fig. 23 shows the results of evaluating activating integrin expression on the surface of the membrane of washed platelets. The graph shaded with grey indicates the result when stimulation by adding ADP was performed after reaction with PBS, the solid line and the dotted line indicate the results when stimulation by ADP was performed after reaction with the omalizumab-G1d-v3/IgE immunocomplex and the omalizumab-BP230/IgE immunocomplex, respectively.

Fig. 24 shows a graph in which the horizontal axis shows the relative value of FcγRIIb-binding activity of each PD variant, and the vertical axis shows the relative value of FcγRIIa type R-binding activity of each PD variant. The value for the amount of binding of each PD variant to each FcγR was divided by the value for the amount of binding of IL6R-F652 / IL6R-L, which is a control antibody prior to introduction of the alteration (IL6R-F652, defined by SEQ ID NO: 142, is an antibody heavy chain comprising an altered Fc with substitution of Pro at position 238 (EU numbering) with Asp), to each FcγR; and then the obtained value was multiplied by 100, and used as the relative binding activity value for each PD variant to each FcγR. The F652 plot in the figure shows the value for IL6R-F652/IL6R-L.

Fig. 25 shows a graph in which the vertical axis shows the relative value of FcγRIIb-binding activity of variants produced by introducing each alteration into GpH7-B3 (SEQ ID NO: 159) / GpL16-k0 (SEQ ID NO: 160) which does not have the P238D alteration, and the horizontal axis shows the relative value of FcγRIIb-binding activity of variants produced by introducing each alteration into IL6R-F652 (SEQ ID NO: 142) / IL6R-L which has the P238D alteration. The value for the amount of FcγRIIb binding of each variant was divided by the value for the amount of FcγRIIb binding of the pre-altered antibody; and then the obtained value was multiplied by 100, and used as the value of relative binding activity. Here, region A contains alterations that exhibit the effect of enhancing FcγRIIb binding in both cases where an

alteration is introduced into GpH7-B3 / GpL16-k0 which does not have P238D and where an alteration is introduced into IL6R-F652 / IL6R-L which has P238D. Region B contains alterations that exhibit the effect of enhancing FcγRIIb binding when introduced into GpH7-B3 / GpL16-k0 which does not have P238D, but do not exhibit the effect of enhancing FcγRIIb binding when introduced into IL6R-F652 / IL6R-L which has P238D.

Fig. 26 shows a crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex.

Fig. 27 shows an image of superimposing the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex and the model structure of the Fc(WT) / FcγRIIb extracellular region complex, with respect to the FcγRIIb extracellular region and the Fc CH2 domain A by the least squares fitting based on the Cα atom pair distances.

Fig. 28 shows comparison of the detailed structure around P238D after superimposing the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex and the model structure of the Fc(WT) / FcγRIIb extracellular region complex with respect to the only Fc CH2 domain A or the only Fc CH2 domain B by the least squares fitting based on the Cα atom pair distances.

Fig. 29 shows that a hydrogen bond can be found between the main chain of Gly at position 237 (indicated by EU numbering) in Fc CH2 domain A, and Tyr at position 160 in FcγRIIb in the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex.

Fig. 30 shows that an electrostatic interaction can be found between Asp at position 270 (indicated by EU numbering) in Fc CH2 domain B, and Arg at position 131 in FcγRIIb in the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex.

Fig. 31 shows a graph in which the horizontal axis shows the relative value of FcγRIIb-binding activity of each 2B variant, and the vertical axis shows the relative value of FcγRIIa type R-binding activity of each 2B variant. The value for the amount of binding of each 2B variant to each FcγR was divided by the value for the amount of binding of a control antibody prior to alteration (altered Fc with substitution of Pro at position 238 (indicated by EU numbering) with Asp) to each FcγR; and then the obtained value was multiplied by 100, and used as the value of relative binding activity of each 2B variant towards each FcγR.

Fig. 32 shows Glu at position 233 (indicated by EU numbering) in Fc Chain A and the surrounding residues in the extracellular region of FcγRIIb in the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex.

Fig. 33 shows Ala at position 330 (indicated by EU numbering) in Fc Chain A and the surrounding residues in the extracellular region of FcγRIIb in the crystal structure of the Fc(P238D) / FcγRIIb extracellular region complex.

Fig. 34 shows the structures of Pro at position 271 (EU numbering) of Fc Chain B after

superimposing the crystal structures of the Fc(P238D) / Fc γ RIIb extracellular region complex and the Fc(WT) / Fc γ RIIIa extracellular region complex by the least squares fitting based on the C α atom pair distances with respect to Fc Chain B.

Fig. 35 shows an image of the Fc (P208)/Fc γ RIIb extracellular region complex
 5 determined by X-ray crystal structure analysis. For each of the CH2 and CH3 domains in the Fc portion, those on the left side are referred to as domain A and those on the right side are referred to as domain B.

Fig. 36 shows comparison after superimposing the structures of Fc (P208)/Fc γ RIIb extracellular region complex and Fc (WT)/Fc γ RIIIa extracellular region complex (PDB code:
 10 3RY6) determined by X-ray crystal structure analysis with respect to the CH2 domain A of the Fc portion by the least squares fitting based on the C α atom pair distances. In the diagram, the structure drawn with heavy line shows the Fc (P208)/Fc γ RIIb extracellular region complex, while the structure drawn with thin line indicates the structure of Fc (WT)/Fc γ RIIIa extracellular region complex. Only the CH2 domain A of the Fc portion is drawn for the Fc (WT)/Fc γ RIIIa
 15 extracellular region complex.

Fig. 37 shows in the X-ray crystal structure of the Fc (P208)/Fc γ RIIb extracellular region complex, a detailed structure around Asp at position 237 (EU numbering) in the CH2 domain A of the Fc portion, which forms a hydrogen bond with Tyr at position 160 in Fc γ RIIb at the main chain moiety.

Fig. 38 shows in the X-ray crystal structure of the Fc (P208)/Fc γ RIIb extracellular region complex, the structure of amino acid residues around Asp at position 237 (EU numbering)
 20 in the CH2 domain A of the Fc portion, which forms a hydrogen bond with Tyr at position 160 in Fc γ RIIb at the main chain moiety.

Fig. 39 shows comparison around the loop at positions 266 to 271 (EU numbering) after
 25 superimposing the X-ray crystal structures of the Fc (P238D)/Fc γ RIIb extracellular region complex shown in Example 10 and the Fc (P208)/Fc γ RIIb extracellular region complex with respect to the CH2 domain B of the Fc portion by the least squares fitting based on the C α atom pair distances. When compared to Fc (P238D), Fc (P208) has the H268D alteration at position 268 (EU numbering) and the P271G alteration at position 271 (EU numbering) in the loop.

Fig. 40 is a diagram showing the structure around Ser239 in the CH2 domain B of the Fc portion in the X-ray crystal structure of the Fc (P208)/Fc γ RIIb extracellular region complex,
 30 along with the electron density determined by X-ray crystal structure analysis with 2Fo-Fc coefficient.

Fig. 41 shows comparison after superimposing the three-dimensional structures of the
 35 Fc (P208)/Fc γ RIIIaR extracellular region complex and Fc (P208)/Fc γ RIIb extracellular region complex determined by X-ray crystal structure analysis by the least squares fitting based on the

C α atom pair distances.

Fig. 42 shows comparison around Asp at position 237 (EU numbering) in the CH2 domain A of the Fc portion between the X-ray crystal structures of the Fc (P208)/Fc γ RIIaR extracellular region complex and the Fc (P208)/Fc γ RIIb extracellular region complex, along with the electron density determined by X-ray crystal structure analysis with 2Fo-Fc coefficient.

Fig. 43 shows comparison around Asp at position 237 (EU numbering) in the CH2 domain B of the Fc portion between the X-ray crystal structures of the Fc (P208)/Fc γ RIIaR extracellular region complex and the Fc (P208)/Fc γ RIIb extracellular region complex, along with the electron density determined by X-ray crystal structure analysis with 2Fo-Fc coefficient.

Fig. 44 shows comparison between the constant-region sequences of G1d and G4d. In the diagram, the amino acids boxed with thick-frame indicate positions with different amino acid residues between G1d and G4d.

Fig. 45 shows the change in plasma antibody concentration of GA2-IgG1 and GA2-F1087 in normal mice.

Fig. 46 shows the change in plasma hIgA concentration in normal mice administered with GA2-IgG1 and GA2-F1087.

Fig. 47 shows the change in plasma antibody concentration of 278-IgG1 and 278-F1087 in C57BL/6J mice.

Fig. 48 shows the change in plasma hIgE (Asp6) concentration in C57BL/6J mice administered with 278-IgG1 and 278-F1087.

Fig. 49 shows the structure of the heavy chain CDR3 of the 6RL#9 antibody Fab fragment determined by X-ray crystal structure analysis. (i) shows the crystal structure of the heavy chain CDR3 obtained under a crystallization condition in the presence of calcium ion. (ii) shows the crystal structure of the heavy chain CDR3 obtained under a crystallization condition in the absence of calcium ion.

Fig. 50 shows a time course of the plasma concentration of each antibody in normal mice administered with antibody H54/L28-IgG1, FH4-IgG1, or 6RL#9-IgG1.

Fig. 51 shows a time course of the plasma concentration of soluble human IL-6 receptor (hsIL-6R) in normal mice administered with antibody H54/L28-IgG1, FH4-IgG1, or 6RL#9-IgG1.

Fig. 52 shows ion-exchange chromatograms for an antibody having human Vk5-2 sequence and an antibody having h Vk5-2_L65 sequence which has an altered glycosylation sequence in the human Vk5-2 sequence. Solid line indicates a chromatogram for an antibody having human Vk5-2 sequence (heavy chain: CIM_H (SEQ ID NO: 67); light chain: hVk5-2 (SEQ ID NO: 4)); broken line indicates a chromatogram for an antibody having hVk5-2_L65 sequence (heavy chain: CIM_H (SEQ ID NO: 67); light chain: hVk5-2_L65 (SEQ ID NO: 70)).

Fig. 53A shows ion-exchange chromatograms for an antibody having LfVkl_Ca sequence (heavy chain: GC_H (SEQ ID NO: 51); light chain: LfVkl_Ca (SEQ ID NO: 83)) and an antibody having a sequence in which Asp (D) in the LfVkl_Ca sequence is substituted with Ala (A) after storage at 5°C (solid line) or 50°C (dotted line). After storage at 5°C, the highest peak in the chromatogram for each antibody is defined as a main peak, and the y axis of each ion-exchange chromatogram was normalized to the main peak. The graph shows a chromatogram for an antibody having LfVkl_Ca (SEQ ID NO: 83) as the light chain.

Fig. 53B shows a chromatogram for an antibody having LfVkl_Ca1 (SEQ ID NO: 85) as the light chain.

Fig. 53C shows a chromatogram for an antibody having LfVkl_Ca2 (SEQ ID NO: 86) as the light chain.

Fig. 53D shows a chromatogram for an antibody having LfVkl_Ca3 (SEQ ID NO: 87) as the light chain.

Fig. 54A shows ion-exchange chromatograms for an antibody having LfVkl_Ca sequence (heavy chain: GC_H (SEQ ID NO: 51); light chain: LfVkl_Ca (SEQ ID NO: 83)) and an antibody having LfVkl_Ca6 sequence (heavy chain: GC_H (SEQ ID NO: 51); light chain: LfVkl_Ca6 (SEQ ID NO: 88)) in which Asp (D) at position 30 (Kabat numbering) in the LfVkl_Ca sequence is substituted with Ser (S) after storage at 5°C (solid line) or 50°C (dotted line). After storage at 5°C, the highest peak in the chromatogram for each antibody is defined as a main peak, and the y axis of each ion-exchange chromatogram was normalized to the main peak. The graph shows a chromatogram for an antibody having LfVkl_Ca (SEQ ID NO: 83) as the light chain.

Fig. 54B shows a chromatogram for an antibody having LfVkl_Ca6 (SEQ ID NO: 88) as the light chain.

Fig. 55 shows the relationship between designed amino acid distribution (indicated with "Design") and amino acid distribution for sequence information on 290 clones isolated from *E. coli* introduced with a gene library of antibodies that bind to antigens in a Ca-dependent manner (indicated with "Library"). The horizontal axis indicates amino acid position (Kabat numbering). The vertical axis indicates percentage in amino acid distribution.

Fig. 56 shows sensorgrams for anti-IL-6R antibody (tocilizumab), antibody 6RC1IgG_010, antibody 6RC1IgG_012, and antibody 6RC1IgG_019 under a high calcium ion concentration (1.2 mM) condition. The horizontal axis shows time, and the vertical axis shows RU value.

Fig. 57 shows sensorgrams for anti-IL-6R antibody (tocilizumab), antibody 6RC1IgG_010, antibody 6RC1IgG_012, and antibody 6RC1IgG_019 under a low calcium ion concentration (3 μ M) condition. The horizontal axis shows time, and the vertical axis shows

RU value.

Fig. 58 shows the relationship between designed amino acid distribution (indicated with "Design") and amino acid distribution for sequence information on 132 clones isolated from *E. coli* introduced with a gene library of antibodies that bind to antigens in a pH-dependent manner (indicated with "Library"). The horizontal axis shows amino acid position (Kabat numbering). The vertical axis indicates percentage in amino acid distribution.

Fig. 59 shows sensorgrams for anti-IL-6R antibody (tocilizumab), antibody 6RpH#01, antibody 6RpH#02, and antibody 6RpH#03 at pH 7.4. The horizontal axis shows time, and the vertical axis shows RU value.

Fig. 60 shows sensorgrams for anti-IL-6R antibody (tocilizumab), antibody 6RpH#01, antibody 6RpH#02, and antibody 6RpH#03 at pH 6.0. The horizontal axis shows time, and the vertical axis shows RU value.

Fig. 61A depicts a graph of ECL responses to native Fc and altered Fc from sera isolated from 15 to 30 independent rheumatism patients. Graphs of ECL responses to native Fc (Fig. 61A), Fv4-YTE (Fig. 61B), Fv4-F1166 (= YTE + Q438R/S440E) (Fig. 61C), Fv4-F1167 (= YTE + S424N) (Fig. 61D), Fv4-LS (Fig. 61E), Fv4-F1170 (= LS + Q438R/S440E) (Fig. 61F), Fv4-F1171 (= LS + S424N) (Fig. 61G), Fv4-N434H (Fig. 61H), Fv4-F1172 (= N434H + Q438R/S440E) (Fig. 61I), Fv4-F1173 (= N434H + S424N) (Fig. 61J) are shown, respectively.

Fig. 61B is a continuation of Fig. 61A.

Fig. 61C is a continuation of Fig. 61B.

Fig. 61D is a continuation of Fig. 61C.

Fig. 61E is a continuation of Fig. 61D.

Fig. 61F is a continuation of Fig. 61E.

Fig. 61G is a continuation of Fig. 61F.

Fig. 61H is a continuation of Fig. 61G.

Fig. 61I is a continuation of Fig. 61H.

Fig. 61J is a continuation of Fig. 61I.

Fig. 62A depicts a graph of ECL responses to altered Fc from sera isolated from 30 independent rheumatism patients. Graphs of ECL responses to Fv4-LS (Fig. 62A), Fv4-F1380 (Fig. 62B), Fv4-F1384 (Fig. 62C), Fv4-F1385 (Fig. 62D), Fv4-F1386 (Fig. 62E), Fv4-F1388 (Fig. 62F), and Fv4-F1389 (Fig. 62G) are shown, respectively.

Fig. 62B is a continuation of Fig. 62A.

Fig. 62C is a continuation of Fig. 62B.

Fig. 62D is a continuation of Fig. 62C.

Fig. 62E is a continuation of Fig. 62D.

Fig. 62F is a continuation of Fig. 62E.

Fig. 62G is a continuation of Fig. 62F.

[Mode for Carrying Out the Invention]

The definitions and detailed description below are provided to help the understanding of the present invention illustrated herein.

Amino acids

Herein, amino acids are described in one- or three-letter codes or both, for example, Ala/A, Leu/L, Arg/R, Lys/K, Asn/N, Met/M, Asp/D, Phe/F, Cys/C, Pro/P, Gln/Q, Ser/S, Glu/E, Thr/T, Gly/G, Trp/W, His/H, Tyr/Y, Ile/I, or Val/V.

Alteration of amino acids

For amino acid alterations in the amino acid sequence of an antigen-binding molecule, known methods such as site-directed mutagenesis methods (Kunkel *et al.* (Proc. Natl. Acad. Sci. USA (1985) 82, 488-492)) and overlap extension PCR may be appropriately employed. Additions, deletions, and/or substitutions of an amino acid are added appropriately by these known methods. Substituting amino acid residues means substituting an amino acid residue with another amino acid residue for the purpose of altering aspects such as the following:

- (a) backbone structure of a polypeptide in a helical structure region or a sheet structure region;
- (b) charge or hydrophobicity at a target site; or
- (c) length of a side chain.

Amino acid residues are classified into the following groups based on the properties of side chains included in their structures:

- (1) hydrophobic: norleucine, Met, Ala, Val, Leu, and Ile;
- (2) neutral hydrophilic: Cys, Ser, Thr, Asn, and Gln;
- (3) acidic: Asp and Glu;
- (4) basic: His, Lys, and Arg;
- (5) residues that affect the orientation of the chain: Gly and Pro; and
- (6) aromatic: Trp, Tyr, and Phe.

Substitution between amino acid residues within each of these groups is referred to as conservative substitution. On the other hand, substitution between amino acid residues from different amino acid groups is referred to as non-conservative substitution. Substitutions in the present invention may be conservative substitutions or non-conservative substitutions, or a combination of conservative and non-conservative substitutions. Furthermore, a plurality of known methods may be employed as amino acid alteration methods for substitution to non-native amino acids (Annu. Rev. Biophys. Biomol. Struct. (2006) 35, 225-249; and Proc. Natl.

Acad. Sci. U.S.A. (2003) 100 (11), 6353-6357). For example, a cell-free translation system (Clover Direct (Protein Express)) containing a tRNA which has the non-native amino acid bound to a complementary amber suppressor tRNA of the UAG codon (amber codon), which is one of the stop codons, is suitably used.

5 Furthermore, an expression that uses one-letter amino-acid codes of the amino acid before alteration and the amino acid after the alteration before and after a number indicating a specific position, respectively, may be used appropriately as an expression for an amino acid alteration. For example, the alteration P238D, which is used when substituting an amino acid of the Fc region included in an antibody constant region, expresses substitution of Pro at position
10 238 (according to EU numbering) with Asp. That is, the number shows the position of the amino acid according to EU numbering, the one-letter amino-acid code written before the number shows the amino acid before substitution, and the one-letter amino-acid code written after the number shows the amino acid after substitution.

15 And/or

As used herein, the term "and/or" means a combination of the terms before and after the set phrase "and/or", and includes every combination where "and" and "or" are suitably combined. Specifically, for example, "the amino acids at positions 326, 328, and/or 428 are substituted" includes a variation of alterations of the following amino acids:

20 amino acid(s) at (a) position 326, (b) position 328, (c) position 428, (d) positions 326 and 328, (e) positions 326 and 428, (f) positions 328 and 428, and (g) positions 326, 328, and 428.

Antigens

25 As used herein, the structure of an "antigen" is not particularly limited to a specific structure as long as it includes an epitope which is bound by an antigen-binding domain. In another meaning, an antigen may be an inorganic matter or an organic matter, and it is preferably a soluble antigen which is present in the body fluid of an organism and which is in an embodiment which may be bound by an antigen-binding molecule of the present invention. The following molecules are examples of the antigens:

30 17-IA, 4-1BB, 4Dc, 6-keto-PGF1a, 8-iso-PGF2a, 8-oxo-dG, A1 adenosine receptor, A33, ACE, ACE-2, activin, activin A, activin AB, activin B, activin C, activin RIA, activin RIA ALK-2, activin RIB ALK-4, activin RIIA, activin RIIB, ADAM, ADAM10, ADAM12, ADAM15, ADAM17/TACE, ADAM8, ADAM9, ADAMTS, ADAMTS4, ADAMTS5, addressin, aFGF,
35 ALCAM, ALK, ALK-1, ALK-7, alpha-1-antitrypsin, alpha-V/beta-1 antagonist, ANG, Ang, APAF-1, APE, APJ, APP, APRIL, AR, ARC, ART, artemin, anti-Id, ASPARTIC, atrial natriuretic

peptide, av/b3 integrin, Ax1, b2M, B7-1, B7-2, B7-H, B-lymphocyte stimulating factor (BlyS), BACE, BACE-1, Bad, BAFF, BAFF-R, Bag-1, BAK, Bax, BCA-1, BCAM, Bcl, BCMA, BDNF, b-ECGF, bFGF, BID, Bik, BIM, BLC, BL-CAM, BLK, BMP, BMP-2 BMP-2a, BMP-3 Osteogenin, BMP-4 BMP-2b, BMP-5, BMP-6 Vgr-1, BMP-7 (OP-1), BMP-8 (BMP-8a, OP-2),

5 BMPR, BMPR-IA (ALK-3), BMPR-IB (ALK-6), BRK-2, RPK-1, BMPR-II (BRK-3), BMP, b-NGF, BOK, bombesin, bone-derived neurotrophic factor, BPDE, BPDE-DNA, BTC, complement factor 3 (C3), C3a, C4, C5, C5a, C10, CA125, CAD-8, calcitonin, cAMP, carcinoembryonic antigen (CEA), cancer associated antigen, cathepsin A, cathepsin B, cathepsin C/DPPI, cathepsin D, cathepsin E, cathepsin H, cathepsin L, cathepsin O, cathepsin S, cathepsin

10 V, cathepsin X/Z/P, CBL, CCI, CCK2, CCL, CCL1, CCL11, CCL12, CCL13, CCL14, CCL15, CCL16, CCL17, CCL18, CCL19, CCL2, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCL26, CCL27, CCL28, CCL3, CCL4, CCL5, CCL6, CCL7, CCL8, CCL9/10, CCR, CCR1, CCR10, CCR10, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CD1, CD2, CD3, CD3E, CD4, CD5, CD6, CD7, CD8, CD10, CD11a, CD11b, CD11c, CD13, CD14, CD15, CD16,

15 CD18, CD19, CD20, CD21, CD22, CD23, CD25, CD27L, CD28, CD29, CD30, CD30L, CD32, CD33 (p67 protein), CD34, CD38, CD40, CD40L, CD44, CD45, CD46, CD49a, CD52, CD54, CD55, CD56, CD61, CD64, CD66e, CD74, CD80 (B7-1), CD89, CD95, CD123, CD137, CD138, CD140a, CD146, CD147, CD148, CD152, CD164, CEACAM5, CFTR, cGMP, CINC, Botulinum toxin, Clostridium perfringens toxin, CKb8-1, CLC, CMV, CMV UL, CNTF,

20 CNTN-1, COX, C-Ret, CRG-2, CT-1, CTACK, CTGF, CTLA-4, CX3CL1, CX3CR1, CXCL, CXCL1, CXCL2, CXCL3, CXCL4, CXCL5, CXCL6, CXCL7, CXCL8, CXCL9, CXCL10, CXCL11, CXCL12, CXCL13, CXCL14, CXCL15, CXCL16, CXCR, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, cytokeratin tumor associated antigen, DAN, DCC, DcR3, DC-SIGN, complement regulatory factor (Decay accelerating factor), des (1-3)-IGF-I (brain IGF-1), Dhh,

25 digoxin, DNAM-1, Dnase, Dpp, DPPIV/CD26, Dtk, ECAD, EDA, EDA-A1, EDA-A2, EDAR, EGF, EGFR (ErbB-1), EMA, EMMPRIN, ENA, endothelin receptor, enkephalinase, eNOS, Eot, eotaxin 1, EpCAM, ephrin B2/EphB4, EPO, ERCC, E-selectin, ET-1, factor IIa, factor VII, factor VIIIc, factor IX, fibroblast activation protein (FAP), Fas, FcR1, FEN-1, ferritin, FGF, FGF-19, FGF-2, FGF3, FGF-8, FGFR, FGFR-3, fibrin, FL, FLIP, Flt-3, Flt-4, follicle

30 stimulating hormone, fractalkine, FZD1, FZD2, FZD3, FZD4, FZD5, FZD6, FZD7, FZD8, FZD9, FZD10, G250, Gas6, GCP-2, GCSF, GD2, GD3, GDF, GDF-1, GDF-3 (Vgr-2), GDF-5 (BMP-14, CDMP-1), GDF-6 (BMP-13, CDMP-2), GDF-7 (BMP-12, CDMP-3), GDF-8 (myostatin), GDF-9, GDF-15 (MIC-1), GDNF, GDNF, GFAP, GFRa-1, GFR-alpha1, GFR-alpha2, GFR-alpha3, GTR, glucagon, Glut4, glycoprotein IIb/IIIa (GPIIb/IIIa), GM-CSF,

35 gp130, gp72, GRO, growth hormone releasing hormone, hapten (NP-cap or NIP-cap), HB-EGF, HCC, HCMV gB envelope glycoprotein, HCMV gH envelope glycoprotein, HCMV UL,

hematopoietic growth factor (HGF), Hep B gp120, heparanase, Her2, Her2/neu (ErbB-2), Her3 (ErbB-3), Her4 (ErbB-4), herpes simplex virus (HSV) gB glycoprotein, HSV gD glycoprotein, HGFA, high molecular weight melanoma-associated antigen (HMW-MAA), HIV gp120, HIV IIIB gp 120 V3 loop, HLA, HLA-DR, HM1.24, HMFG PEM, HRG, Hrk, human cardiac myosin,

5 human cytomegalovirus (HCMV), human growth hormone (HGH), HVEM, I-309, IAP, ICAM, ICAM-1, ICAM-3, ICE, ICOS, IFNg, Ig, IgA receptor, IgE, IGF, IGF binding protein, IGF-1R, IGFBP, IGF-I, IGF-II, IL, IL-1, IL-1R, IL-2, IL-2R, IL-4, IL-4R, IL-5, IL-5R, IL-6, IL-6R, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-18, IL-18R, IL-23, interferon (INF)-alpha, INF-beta, INF-gamma, inhibin, iNOS, insulin A chain, insulin B chain, insulin-like growth factor1, integrin

10 alpha2, integrin alpha3, integrin alpha4, integrin alpha4/beta1, integrin alpha4/beta7, integrin alpha5 (alpha V), integrin alpha5/beta1, integrin alpha5/beta3, integrin alpha6, integrin beta1, integrin beta2, interferon gamma, IP-10, I-TAC, JE, kallikrein 2, kallikrein 5, kallikrein 6, kallikrein 11, kallikrein 12, kallikrein 14, kallikrein 15, kallikrein L1, kallikrein L2, kallikrein L3, kallikrein L4, KC, KDR, keratinocyte growth factor (KGF), laminin 5, LAMP, LAP, LAP

15 (TGF-1), latent TGF-1, latent TGF-1 bp1, LBP, LDGF, LECT2, lefty, Lewis-Y antigen, Lewis-Y associated antigen, LFA-1, LFA-3, Lfo, LIF, LIGHT, lipoprotein, LIX, LKN, Lptn, L-selectin, LT-a, LT-b, LTB4, LTBP-1, lung surface, luteinizing hormone, lymphotoxin beta receptor, Mac-1, MAdCAM, MAG, MAP2, MARC, MCAM, MCAM, MCK-2, MCP, M-CSF, MDC, Mer, METALLOPROTEASES, MGDF receptor, MGMT, MHC (HLA-DR), MIF, MIG, MIP,

20 MIP-1-alpha, MK, MMAC1, MMP, MMP-1, MMP-10, MMP-11, MMP-12, MMP-13, MMP-14, MMP-15, MMP-2, MMP-24, MMP-3, MMP-7, MMP-8, MMP-9, MPIF, Mpo, MSK, MSP, mucin (Muc1), MUC18, Mullerian-inhibiting substance, Mug, MuSK, NAIP, NAP, NCAD, N-C adherin, NCA 90, NCAM, NCAM, neprilysin, neurotrophin-3, -4, or -6, neurturin, nerve growth factor (NGF), NGFR, NGF-beta, nNOS, NO, NOS, Npn, NRG-3, NT, NTN, OB, OGG1, OPG,

25 OPN, OSM, OX40L, OX40R, p150, p95, PADPr, parathyroid hormone, PARC, PARP, PBR, PBSF, PCAD, P-cadherin, PCNA, PDGF, PDGF, PDK-1, PECAM, PEM, PF4, PGE, PGF, PGI2, PGJ2, PIN, PLA2, placental alkaline phosphatase (PLAP), PlGF, PLP, PP14, proinsulin, prorelaxin, protein C, PS, PSA, PSCA, prostate-specific membrane antigen (PSMA), PTEN, PTHrp, Ptk, PTN, R51, RANK, RANKL, RANTES, RANTES, relaxin A chain, relaxin B chain,

30 renin, respiratory syncytial virus (RSV) F, RSV Fgp, Ret, Rheumatoid factor, RLIP76, RPA2, RSK, S100, SCF/KL, SDF-1, SERINE, serum albumin, sFRP-3, Shh, SIGIRR, SK-1, SLAM, SLPI, SMAC, SMDF, SMOH, SOD, SPARC, Stat, STEAP, STEAP-II, TACE, TACI, TAG-72 (tumor-associated glycoprotein-72), TARC, TCA-3, T-cell receptor (for example, T-cell receptor alpha/beta), TdT, TECK, TEM1, TEM5, TEM7, TEM8, TERT, testis PLAP-like alkaline

35 phosphatase, TfR, TGF, TGF-alpha, TGF-beta, TGF-beta Pan Specific, TGF-betaRI (ALK-5), TGF-betaRII, TGF-betaRIIb, TGF-betaRIII, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta4,

TGF-beta5, thrombin, thymus Ck-1, thyroid-stimulating hormone, Tie, TIMP, TIQ, tissue factor, TMEFF2, Tmpo, TMPRSS2, TNF, TNF-alpha, TNF-alphabeta, TNF-beta2, TNFc, TNF-RI, TNF-RII, TNFRSF10A (TRAIL R1 Apo-2, DR4), TNFRSF10B (TRAIL R2 DR5, KILLER, TRICK-2A, TRICK-B), TNFRSF10C (TRAIL R3 DcR1, LIT, TRID), TNFRSF10D (TRAIL R4 DcR2, TRUNDD), TNFRSF11A (RANK ODF R, TRANCE R), TNFRSF11B (OPG OCIF, TR1),
 5 TNFRSF12 (TWEAK R FN14), TNFRSF13B (TACI), TNFRSF13C (BAFF R), TNFRSF14 (HVEM ATAR, HveA, LIGHT R, TR2), TNFRSF16 (NGFR p75NTR), TNFRSF17 (BCMA), TNFRSF18 (GITR AITR), TNFRSF19 (TROY TAJ, TRADE), TNFRSF19L (RELT), TNFRSF1A (TNF RI CD120a, p55-60), TNFRSF1B (TNF RII CD120b, p75-80), TNFRSF26
 10 (TNFRH3) , TNFRSF3 (LTbR TNF RIII, TNFC R), TNFRSF4 (OX40 ACT35, TXGP1 R), TNFRSF5 (CD40 p50), TNFRSF6 (Fas Apo-1, APT1, CD95), TNFRSF6B (DcR3 M68, TR6), TNFRSF7 (CD27), TNFRSF8 (CD30), TNFRSF9 (4-1BB CD137, ILA), TNFRSF21 (DR6), TNFRSF22 (DcTRAIL R2 TNFRH2), TNFRST23 (DcTRAIL R1 TNFRH1), TNFRSF25 (DR3 Apo-3, LARD, TR-3, TRAMP, WSL-1), TNFSF10 (TRAIL Apo-2 ligand, TL2), TNFSF11
 15 (TRANCE/RANK ligand ODF, OPG ligand), TNFSF12 (TWEAK Apo-3 ligand, DR3 ligand), TNFSF13 (APRIL TALL2), TNFSF13B (BAFF BLYS, TALL1, THANK, TNFSF20), TNFSF14 (LIGHT HVEM ligand, LTg), TNFSF15 (TL1A/VEGI), TNFSF18 (GITR ligand AITR ligand, TL6), TNFSF1A (TNF-a Conectin, DIF, TNFSF2), TNFSF1B (TNF-b LTa, TNFSF1), TNFSF3 (LTb TNFC, p33), TNFSF4 (OX40 ligand gp34, TXGP1), TNFSF5 (CD40 ligand CD154, gp39,
 20 HIGM1, IMD3, TRAP), TNFSF6 (Fas ligand Apo-1 ligand, APT1 ligand), TNFSF7 (CD27 ligand CD70), TNFSF8 (CD30 ligand CD153), TNFSF9 (4-1BB ligand CD137 ligand), TP-1, t-PA, Tpo, TRAIL, TRAIL R, TRAIL-R1, TRAIL-R2, TRANCE, transferrin receptor, TRF, Trk, TROP-2, TSG, TSLP, tumor associated antigen CA125, tumor associated antigen expressing Lewis-Y associated carbohydrates, TWEAK, TXB2, Ung, uPAR, uPAR-1, urokinase, VCAM,
 25 VCAM-1, VECAD, VE-Cadherin, VE-cadherin-2, VEGFR-1 (flt-1), VEGF, VEGFR, VEGFR-3 (flt-4), VEGI, VIM, virus antigen, VLA, VLA-1, VLA-4, VNR integrin, von Willebrand factor, WIF-1, WNT1, WNT2, WNT2B/13, WNT3, WNT3A, WNT4, WNT5A, WNT5B, WNT6, WNT7A, WNT7B, WNT8A, WNT8B, WNT9A, WNT9A, WNT9B, WNT10A, WNT10B, WNT11, WNT16, XCL1, XCL2, XCR1, XCR1, XEDAR, XIAP, XPD, HMGB1, IgA, A β , CD81,
 30 CD97, CD98, DDR1, DKK1, EREG, Hsp90, IL-17/IL-17R, IL-20/IL-20R, oxidized LDL, PCSK9, prekallikrein, RON, TMEM16F, SOD1, Chromogranin A, Chromogranin B, tau, VAP1, high molecular weight kininogen, IL-31, IL-31R, Nav1.1, Nav1.2, Nav1.3, Nav1.4, Nav1.5, Nav1.6, Nav1.7, Nav1.8, Nav1.9, EPCR, C1, C1q, C1r, C1s, C2, C2a, C2b, C3, C3a, C3b, C4, C4a, C4b, C5, C5a, C5b, C6, C7, C8, C9, factor B, factor D, factor H, properdin, sclerostin,
 35 fibrinogen, fibrin, prothrombin, thrombin, tissue factor, factor V, factor Va, factor VII, factor VIIa, factor VIII, factor VIIa, factor IX, factor IXa, factor X, factor Xa, factor XI, factor XIa,

factor XII, factor XIIa, factor XIII, factor XIIIa, TFPI, antithrombin III, EPCR, thrombomodulin, TAPI, tPA, plasminogen, plasmin, PAI-1, PAI-2, GPC3, Syndecan-1, Syndecan-2, Syndecan-3, Syndecan-4, LPA, S1P, Acetylcholine receptor, AdipoR1, AdipoR2, ADP ribosyl cyclase-1, alpha-4/beta-7 integrin, alpha-5/beta-1 integrin, alpha-v/beta-6 integrin, alphavbeta1 integrin, 5 Angiopoietin ligand-2, Angptl2, Anthrax, Cadherin, Carbonic anhydrase-IX, CD105, CD155, CD158a, CD37, CD49b, CD51, CD70, CD72, Claudin 18, Clostridium difficile toxin, CS1, Delta-like protein ligand 4, DHICA oxidase, Dickkopf-1 ligand, Dipeptidyl peptidase IV, EPOR, F protein of RSV, Factor Ia, FasL, Folate receptor alpha, Glucagon receptor, Glucagon-like peptide 1 receptor, Glutamate carboxypeptidase II, GMCSFR, Hepatitis C virus E2 glycoprotein, 10 Hepcidin, IL-17 receptor, IL-22 receptor, IL-23 receptor, IL-3 receptor, Kit tyrosine kinase, Leucine Rich Alpha-2-Glycoprotein 1 (LRG1), Lysosphingolipid receptor, Membrane glycoprotein OX2, Mesothelin, MET, MICA, MUC-16, Myelin associated glycoprotein, Neuropilin-1, Neuropilin-2, Nogo receptor, PLXNA1, PLXNA2, PLXNA3, PLXNA4A, PLXNA4B, PLXNB1, PLXNB2, PLXNB3, PLXNC1, PLXND1, Programmed cell death ligand 15 1, Proprotein convertase PC9, P-selectin glycoprotein ligand-1, RAGE, Reticulon 4, RF, RON-8, SEMA3A, SEMA3B, SEMA3C, SEMA3D, SEMA3E, SEMA3F, SEMA3G, SEMA4A, SEMA4B, SEMA4C, SEMA4D, SEMA4F, SEMA4G, SEMA5A, SEMA5B, SEMA6A, SEMA6B, SEMA6C, SEMA6D, SEMA7A, Shiga like toxin II, Sphingosine-1-phosphate receptor-1, ST2, Staphylococcal lipoteichoic acid, Tenascin, TG2, Thymic stromal 20 lymphoprotein receptor, TNF superfamily receptor 12A, Transmembrane glycoprotein NMB, TREM-1, TREM-2, Trophoblast glycoprotein, TSH receptor, TTR, Tubulin, and ULBP2, and receptors for growth factors and hormones, molecules that exist in their soluble form and are not anchored to cells in the body fluid of organisms. For example, among the receptors, soluble antigens present in the body fluid of an organism due to some mechanism including 25 protease-mediated digestion of receptors or such expressed on a cell surface are also suitable examples of the soluble antigens of the present invention. Examples of such molecules may include the soluble IL-6R molecule (J. Immunol. (1994) 152, 4958-4968) and CD20, as well as CD52 (Br. J. Haematol. (2003) 123 (5), 850-857), described herein. Furthermore, not only the molecules inherently expressed in a living organism, but also soluble antigens existing in the 30 body fluid of an organism, which are infectious molecules such as prions or antigens presented by infectious organisms such as viruses or presented on such organisms are also examples of the soluble antigens of the present invention. Suitable examples of the body fluid include blood, plasma, serum, urine, lymph, saliva, and tear fluid.

35 Epitope

“Epitope” means an antigenic determinant in an antigen, and refers to an antigen site to

which the antigen-binding domain of an antigen-binding molecule disclosed herein binds. Thus, for example, the epitope can be defined according to its structure. Alternatively, the epitope may be defined according to the antigen-binding activity of an antigen-binding molecule that recognizes the epitope. When the antigen is a peptide or polypeptide, the epitope can be specified by the amino acid residues forming the epitope. Alternatively, when the epitope is a sugar chain, the epitope can be specified by its specific sugar chain structure.

A linear epitope is an epitope that contains an epitope whose primary amino acid sequence is recognized. Such a linear epitope typically contains at least three and most commonly at least five, for example, about 8 to about 10 or 6 to 20 amino acids in its specific sequence.

In contrast to the linear epitope, “conformational epitope” is an epitope in which the primary amino acid sequence containing the epitope is not the only determinant of the recognized epitope (for example, the primary amino acid sequence of a conformational epitope is not necessarily recognized by an epitope-defining antibody). Conformational epitopes may contain a greater number of amino acids compared to linear epitopes. A conformational epitope-recognizing antibody recognizes the three-dimensional structure of a peptide or protein. For example, when a protein molecule folds and forms a three-dimensional structure, amino acids and/or polypeptide main chains that form a conformational epitope become aligned, and the epitope is made recognizable by the antibody. Methods for determining epitope conformations include, for example, X ray crystallography, two-dimensional nuclear magnetic resonance, site-specific spin labeling, and electron paramagnetic resonance, but are not limited thereto. See, for example, Epitope Mapping Protocols in Methods in Molecular Biology (1996), Vol. 66, Morris (ed.).

Binding Activity

Examples of a method for assessing the epitope binding by a test antigen-binding molecule containing an IL-6R antigen-binding domain are described below. According to the examples below, methods for assessing the epitope binding by a test antigen-binding molecule containing an antigen-binding domain for an antigen other than IL-6R, can also be appropriately conducted.

For example, whether a test antigen-binding molecule containing an IL-6R antigen-binding domain recognizes a linear epitope in the IL-6R molecule can be confirmed for example as mentioned below. A linear peptide comprising an amino acid sequence forming the extracellular domain of IL-6R is synthesized for the above purpose. The peptide can be synthesized chemically, or obtained by genetic engineering techniques using a region encoding the amino acid sequence corresponding to the extracellular domain in an IL-6R cDNA. Then, a

test antigen-binding molecule containing an IL-6R antigen-binding domain is assessed for its binding activity towards a linear peptide comprising the amino acid sequence forming the extracellular domain. For example, an immobilized linear peptide can be used as an antigen by ELISA to evaluate the binding activity of the antigen-binding molecule towards the peptide.

- 5 Alternatively, the binding activity towards a linear peptide can be assessed based on the level that the linear peptide inhibits the binding of the antigen-binding molecule to IL-6R-expressing cells. These tests can demonstrate the binding activity of the antigen-binding molecule towards the linear peptide.

Whether a test antigen-binding molecule containing an IL-6R antigen-binding domain
10 recognizes a conformational epitope can be assessed as follows. IL-6R-expressing cells are prepared for the above purpose. A test antigen-binding molecule containing an IL-6R antigen-binding domain can be determined to recognize a conformational epitope when it strongly binds to IL-6R-expressing cells upon contact, but does not substantially bind to an immobilized linear peptide comprising an amino acid sequence forming the extracellular domain
15 of IL-6R. Herein, “not substantially bind” means that the binding activity is 80% or less, generally 50% or less, preferably 30% or less, and particularly preferably 15% or less compared to the binding activity towards cells expressing human IL-6R.

Methods for assaying the binding activity of a test antigen-binding molecule containing an IL-6R antigen-binding domain towards IL-6R-expressing cells include, for example, the
20 methods described in *Antibodies: A Laboratory Manual* (Ed Harlow, David Lane, Cold Spring Harbor Laboratory (1988) 359-420). Specifically, the assessment can be performed based on the principle of ELISA or fluorescence activated cell sorting (FACS) using IL-6R-expressing cells as antigen.

In the ELISA format, the binding activity of a test antigen-binding molecule containing
25 an IL-6R antigen-binding domain towards IL-6R-expressing cells can be assessed quantitatively by comparing the levels of signal generated by enzymatic reaction. Specifically, a test antigen-binding molecule is added to an ELISA plate onto which IL-6R-expressing cells are immobilized. Then, the test antigen-binding molecule bound to the cells is detected using an enzyme-labeled antibody that recognizes the test antigen-binding molecule. Alternatively,
30 when FACS is used, a dilution series of a test antigen-binding molecule is prepared, and the antibody binding titer for IL-6R-expressing cells can be determined to compare the binding activity of the test antigen-binding molecule towards IL-6R-expressing cells.

The binding of a test antigen-binding molecule towards an antigen expressed on the surface of cells suspended in buffer or the like can be detected using a flow cytometer. Known
35 flow cytometers include, for example, the following devices:
FACSCanto™ II

FACSAria™

FACSArray™

FACSVantage™ SE

FACSCalibur™ (all are trade names of BD Biosciences)

5 EPICS ALTRA HyPerSort™

Cytomics FC 500

EPICS XL-MCL ADC EPICS XL ADC

Cell Lab Quanta/Cell Lab Quanta SC (all are trade names of Beckman Coulter).

Preferable methods for assaying the binding activity of a test antigen-binding molecule containing an IL-6R antigen-binding domain towards an antigen include, for example, the following method. First, IL-6R-expressing cells are reacted with a test antigen-binding molecule, and then this is stained with an FITC-labeled secondary antibody that recognizes the antigen-binding molecule. The test antigen-binding molecule is appropriately diluted with a suitable buffer to prepare the molecule at a desired concentration. For example, the molecule can be used at a concentration within the range of 10 µg/ml to 10 ng/ml. Then, the fluorescence intensity and cell count are determined using FACSCalibur (BD). The fluorescence intensity obtained by analysis using the CELL QUEST Software (BD), *i.e.*, the Geometric Mean value, reflects the quantity of antibody bound to cells. That is, the binding activity of a test antigen-binding molecule, which is represented by the quantity of the test antigen-binding molecule bound, can be determined by measuring the Geometric Mean value.

Whether a test antigen-binding molecule containing an IL-6R antigen-binding domain shares a common epitope with another antigen-binding molecule can be assessed based on the competition between the two molecules for the same epitope. The competition between antigen-binding molecules can be detected by cross-blocking assay or the like. For example, the competitive ELISA assay is a preferred cross-blocking assay.

Specifically, in cross-blocking assay, the IL-6R protein immobilized to the wells of a microtiter plate is pre-incubated in the presence or absence of a candidate competitor antigen-binding molecule, and then a test antigen-binding molecule is added thereto. The quantity of test antigen-binding molecule bound to the IL-6R protein in the wells is indirectly correlated with the binding ability of a candidate competitor antigen-binding molecule that competes for the binding to the same epitope. That is, the greater the affinity of the competitor antigen-binding molecule for the same epitope, the lower the binding activity of the test antigen-binding molecule towards the IL-6R protein-coated wells.

The quantity of the test antigen-binding molecule bound to the wells *via* the IL-6R protein can be readily determined by labeling the antigen-binding molecule in advance. For example, a biotin-labeled antigen-binding molecule is measured using an avidin/peroxidase

conjugate and appropriate substrate. In particular, cross-blocking assay that uses enzyme labels such as peroxidase is called “competitive ELISA assay”. The antigen-binding molecule can also be labeled with other labeling substances that enable detection or measurement.

Specifically, radiolabels, fluorescent labels, and such are known.

5 When the candidate competitor antigen-binding molecule can block the binding by a test antigen-binding molecule containing an IL-6R antigen-binding domain by at least 20%, preferably at least 20 to 50%, and more preferably at least 50% compared to the binding activity in a control experiment conducted in the absence of the competitor antigen-binding molecule, the test antigen-binding molecule is determined to substantially bind to the same epitope bound by
10 the competitor antigen-binding molecule, or compete for the binding to the same epitope.

 When the structure of an epitope bound by a test antigen-binding molecule containing an IL-6R antigen-binding domain has already been identified, whether the test and control antigen-binding molecules share a common epitope can be assessed by comparing the binding activities of the two antigen-binding molecules towards a peptide prepared by introducing amino
15 acid mutations into the peptide forming the epitope.

 To measure the above binding activities, for example, the binding activities of test and control antigen-binding molecules towards a linear peptide into which a mutation is introduced are compared in the above ELISA format. Besides the ELISA methods, the binding activity towards the mutant peptide bound to a column can be determined by flowing test and control
20 antigen-binding molecules in the column, and then quantifying the antigen-binding molecule eluted in the elution solution. Methods for adsorbing a mutant peptide to a column, for example, in the form of a GST fusion peptide, are known.

 Alternatively, when the identified epitope is a conformational epitope, whether test and control antigen-binding molecules share a common epitope can be assessed by the following
25 method. First, IL-6R-expressing cells and cells expressing IL-6R with a mutation introduced into the epitope are prepared. The test and control antigen-binding molecules are added to a cell suspension prepared by suspending these cells in an appropriate buffer such as PBS. Then, the cell suspensions are appropriately washed with a buffer, and an FITC-labeled antibody that recognizes the test and control antigen-binding molecules is added thereto. The fluorescence
30 intensity and number of cells stained with the labeled antibody are determined using FACSCalibur (BD). The test and control antigen-binding molecules are appropriately diluted using a suitable buffer, and used at desired concentrations. For example, they may be used at a concentration within the range of 10 µg/ml to 10 ng/ml. The fluorescence intensity determined by analysis using the CELL QUEST Software (BD), *i.e.*, the Geometric Mean value, reflects the
35 quantity of labeled antibody bound to cells. That is, the binding activities of the test and control antigen-binding molecules, which are represented by the quantity of labeled antibody bound, can

be determined by measuring the Geometric Mean value.

In the above method, whether an antigen-binding molecule does “not substantially bind to cells expressing mutant IL-6R” can be assessed, for example, by the following method. First, the test and control antigen-binding molecules bound to cells expressing mutant IL-6R are stained with a labeled antibody. Then, the fluorescence intensity of the cells is determined. When FACSCalibur is used for fluorescence detection by flow cytometry, the determined fluorescence intensity can be analyzed using the CELL QUEST Software. From the Geometric Mean values in the presence and absence of the antigen-binding molecule, the comparison value (Δ Geo-Mean) can be calculated according to the following formula to determine the ratio of increase in fluorescence intensity as a result of the binding by the antigen-binding molecule.

$$\Delta\text{Geo-Mean} = \text{Geo-Mean (in the presence of the antigen-binding molecule)} / \text{Geo-Mean (in the absence of the antigen-binding molecule)}$$

The Geometric Mean comparison value (Δ Geo-Mean value for the mutant IL-6R molecule) determined by the above analysis, which reflects the quantity of a test antigen-binding molecule bound to cells expressing mutant IL-6R, is compared to the Δ Geo-Mean comparison value that reflects the quantity of the test antigen-binding molecule bound to IL-6R-expressing cells. In this case, the concentrations of the test antigen-binding molecule used to determine the Δ Geo-Mean comparison values for IL-6R-expressing cells and cells expressing mutant IL-6R are particularly preferably adjusted to be equal or substantially equal. An antigen-binding molecule that has been confirmed to recognize an epitope in IL-6R is used as a control antigen-binding molecule.

If the Δ Geo-Mean comparison value of a test antigen-binding molecule for cells expressing mutant IL-6R is smaller than the Δ Geo-Mean comparison value of the test antigen-binding molecule for IL-6R-expressing cells by at least 80%, preferably 50%, more preferably 30%, and particularly preferably 15%, then the test antigen-binding molecule “does not substantially bind to cells expressing mutant IL-6R”. The formula for determining the Geo-Mean (Geometric Mean) value is described in the CELL QUEST Software User’s Guide (BD biosciences). When the comparison shows that the comparison values are substantially equivalent, the epitope for the test and control antigen-binding molecules can be determined to be the same.

Antigen-binding domain

Herein, an “antigen-binding domain” may be of any structure as long as it binds to an antigen of interest. Such domains preferably include, for example:

antibody heavy-chain and light-chain variable regions;

a module of about 35 amino acids called A domain which is contained in the *in vivo* cell membrane protein Avimer (WO 2004/044011, WO 2005/040229);

Adnectin containing the 10Fn3 domain which binds to the protein moiety of fibronectin, a glycoprotein expressed on cell membrane (WO 2002/032925);

Affibody which is composed of a 58-amino acid three-helix bundle based on the scaffold of the IgG-binding domain of Protein A (WO 1995/001937);

Designed Ankyrin Repeat proteins (DARPin) which are a region exposed on the molecular surface of ankyrin repeats (AR) having a structure in which a subunit consisting of a turn comprising 33 amino acid residues, two antiparallel helices, and a loop is repeatedly stacked (WO 2002/020565);

Anticalins and such, which are domains consisting of four loops that support one side of a barrel structure composed of eight circularly arranged antiparallel strands that are highly conserved among lipocalin molecules such as neutrophil gelatinase-associated lipocalin (NGAL) (WO 2003/029462); and

the concave region formed by the parallel-sheet structure inside the horseshoe-shaped structure constituted by stacked repeats of the leucine-rich-repeat (LRR) module of the variable lymphocyte receptor (VLR) which does not have the immunoglobulin structure and is used in the system of acquired immunity in jawless vertebrate such as lamprey and hagfish (WO 2008/016854). Preferred antigen-binding domains of the present invention include, for example, those having antibody heavy-chain and light-chain variable regions. Preferred examples of antigen-binding domains include "single chain Fv (scFv)", "single chain antibody", "Fv", "single chain Fv 2 (scFv2)", "Fab", and "F(ab')₂".

The antigen-binding domains of antigen-binding molecules of the present invention can bind to an identical epitope. Such epitope can be present, for example, in a protein comprising the amino acid sequence of SEQ ID NO: 1. Alternatively, the epitope can be present in the protein comprising the amino acids at positions 20 to 365 in the amino acid sequence of SEQ ID NO: 1. Alternatively, each of the antigen-binding domains of antigen-binding molecules of the present invention can bind to a different epitope. Herein, the different epitope can be present in, for example, a protein comprising the amino acid sequence of SEQ ID NO: 1. Alternatively, the epitope can be present in the protein comprising the amino acids at positions 20 to 365 in the amino acid sequence of SEQ ID NO: 1.

Specificity

"Specific" means that one of molecules that specifically binds to does not show any significant binding to molecules other than a single or a number of binding partner molecules.

Furthermore, "specific" is also used when an antigen-binding domain is specific to a particular epitope among multiple epitopes in an antigen. When an epitope bound by an antigen-binding domain is contained in multiple different antigens, antigen-binding molecules containing the antigen-binding domain can bind to various antigens that have the epitope.

5

Antibodies

Herein, "antibody" refers to a natural immunoglobulin or an immunoglobulin produced by partial or complete synthesis. Antibodies can be isolated from natural sources such as naturally-occurring plasma and serum, or culture supernatants of antibody-producing
10 hybridomas. Alternatively, antibodies can be partially or completely synthesized using techniques such as genetic recombination. Preferred antibodies include, for example, antibodies of an immunoglobulin isotype or subclass belonging thereto. Known human immunoglobulins include antibodies of the following nine classes (isotypes): IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM. Of these isotypes, antibodies of the present invention
15 include IgG1, IgG2, IgG3, and IgG4. IgG constant regions include mutants naturally formed therefrom. A number of allotype sequences due to genetic polymorphism are described in "Sequences of proteins of immunological interest", NIH Publication No.91-3242, for the constant regions of human IgG1, human IgG2, human IgG3, and human IgG4 antibodies, and any one of them may be used in the present invention. In particular for the human IgG1
20 sequence, the amino acid sequence of positions 356 to 358 (EU numbering) may be either DEL or EEM.

Methods for producing an antibody with desired binding activity are known to those skilled in the art. Below is an example that describes a method for producing an antibody that binds to IL-6R (anti-IL-6R antibody). Antibodies that bind to an antigen other than IL-6R can
25 also be produced according to the example described below.

Anti-IL-6R antibodies can be obtained as polyclonal or monoclonal antibodies using known methods. The anti-IL-6R antibodies preferably produced are monoclonal antibodies derived from mammals. Such mammal-derived monoclonal antibodies include antibodies produced by hybridomas or host cells transformed with an expression vector carrying an
30 antibody gene by genetic engineering techniques. "Humanized antibodies" or "chimeric antibodies" are included in the monoclonal antibodies of the present invention.

Monoclonal antibody-producing hybridomas can be produced using known techniques, for example, as described below. Specifically, mammals are immunized by conventional immunization methods using an IL-6R protein as a sensitizing antigen. Resulting immune cells
35 are fused with known parental cells by conventional cell fusion methods. Then, hybridomas producing an anti-IL-6R antibody can be selected by screening for monoclonal

antibody-producing cells using conventional screening methods.

Specifically, monoclonal antibodies are prepared as mentioned below. First, the IL-6R gene whose nucleotide sequence is disclosed in SEQ ID NO: 2 can be expressed to produce an IL-6R protein shown in SEQ ID NO: 1, which will be used as a sensitizing antigen for antibody preparation. That is, a gene sequence encoding IL-6R is inserted into a known expression vector, and appropriate host cells are transformed with this vector. The desired human IL-6R protein is purified from the host cells or their culture supernatants by known methods. In order to obtain soluble IL-6R from culture supernatants, for example, a protein consisting of the amino acids at positions 1 to 357 in the IL-6R polypeptide sequence of SEQ ID NO: 1, such as described in Mullberg *et al.* (J. Immunol. (1994) 152 (10), 4958-4968), is expressed as a soluble IL-6R, instead of the IL-6R protein of SEQ ID NO: 1. Purified natural IL-6R protein can also be used as a sensitizing antigen.

The purified IL-6R protein can be used as a sensitizing antigen for immunization of mammals. A partial IL-6R peptide may also be used as a sensitizing antigen. In this case, a partial peptide can be prepared by chemical synthesis based on the amino acid sequence of human IL-6R, or by inserting a partial IL-6R gene into an expression vector for expression. Alternatively, a partial peptide can be produced by degrading an IL-6R protein with a protease. The length and region of the partial IL-6R peptide are not limited to particular embodiments. A preferred region can be arbitrarily selected from the amino acid sequence at amino acid positions 20 to 357 in the amino acid sequence of SEQ ID NO: 1. The number of amino acids forming a peptide to be used as a sensitizing antigen is preferably at least five or more, six or more, or seven or more. More specifically, a peptide of 8 to 50 residues, more preferably 10 to 30 residues can be used as a sensitizing antigen.

For sensitizing antigen, alternatively it is possible to use a fusion protein prepared by fusing a desired partial polypeptide or peptide of the IL-6R protein with a different polypeptide. For example, antibody Fc fragments and peptide tags are preferably used to produce fusion proteins to be used as sensitizing antigens. Vectors for expression of such fusion proteins can be constructed by fusing in frame genes encoding two or more desired polypeptide fragments and inserting the fusion gene into an expression vector as described above. Methods for producing fusion proteins are described in Molecular Cloning 2nd ed. (Sambrook, J *et al.*, Molecular Cloning 2nd ed., 9.47-9.58 (1989) Cold Spring Harbor Lab. Press). Methods for preparing IL-6R to be used as a sensitizing antigen, and immunization methods using IL-6R are specifically described in WO 2003/000883, WO 2004/022754, WO 2006/006693, and such.

There is no particular limitation on the mammals to be immunized with the sensitizing antigen. However, it is preferable to select the mammals by considering their compatibility with the parent cells to be used for cell fusion. In general, rodents such as mice, rats, and

hamsters, rabbits, and monkeys are preferably used.

The above animals are immunized with a sensitizing antigen by known methods. Generally performed immunization methods include, for example, intraperitoneal or subcutaneous injection of a sensitizing antigen into mammals. Specifically, a sensitizing antigen is appropriately diluted with PBS (Phosphate-Buffered Saline), physiological saline, or the like. If desired, a conventional adjuvant such as Freund's complete adjuvant is mixed with the antigen, and the mixture is emulsified. Then, the sensitizing antigen is administered to a mammal several times at 4- to 21-day intervals. Appropriate carriers may be used in immunization with the sensitizing antigen. In particular, when a low-molecular-weight partial peptide is used as the sensitizing antigen, it is sometimes desirable to couple the sensitizing antigen peptide to a carrier protein such as albumin or keyhole limpet hemocyanin for immunization.

Alternatively, hybridomas producing a desired antibody can be prepared using DNA immunization as mentioned below. DNA immunization is an immunization method that confers immunostimulation by expressing a sensitizing antigen in an animal immunized as a result of administering a vector DNA constructed to allow expression of an antigen protein-encoding gene in the animal. As compared to conventional immunization methods in which a protein antigen is administered to animals to be immunized, DNA immunization is expected to be superior in that:

- immunostimulation can be provided while retaining the structure of a membrane protein such as IL-6R; and
- there is no need to purify the antigen for immunization.

In order to prepare a monoclonal antibody of the present invention using DNA immunization, first, a DNA expressing an IL-6R protein is administered to an animal to be immunized. The IL-6R-encoding DNA can be synthesized by known methods such as PCR. The obtained DNA is inserted into an appropriate expression vector, and then this is administered to an animal to be immunized. Preferably used expression vectors include, for example, commercially-available expression vectors such as pcDNA3.1. Vectors can be administered to an organism using conventional methods. For example, DNA immunization is performed by using a gene gun to introduce expression vector-coated gold particles into cells in the body of an animal to be immunized. Antibodies that recognized IL-6R can also be produced by the methods described in WO 2003/104453.

After immunizing a mammal as described above, an increase in the titer of an IL-6R-binding antibody is confirmed in the serum. Then, immune cells are collected from the mammal, and then subjected to cell fusion. In particular, splenocytes are preferably used as immune cells.

A mammalian myeloma cell is used as a cell to be fused with the above-mentioned immune cells. The myeloma cells preferably comprise a suitable selection marker for screening. A selection marker confers characteristics to cells for their survival (or death) under a specific culture condition. Hypoxanthine-guanine phosphoribosyltransferase deficiency (hereinafter abbreviated as HGPRT deficiency) and thymidine kinase deficiency (hereinafter abbreviated as TK deficiency) are known as selection markers. Cells with HGPRT or TK deficiency have hypoxanthine-aminopterin-thymidine sensitivity (hereinafter abbreviated as HAT sensitivity). HAT-sensitive cells cannot synthesize DNA in a HAT selection medium, and are thus killed. However, when the cells are fused with normal cells, they can continue DNA synthesis using the salvage pathway of the normal cells, and therefore they can grow even in the HAT selection medium.

HGPRT-deficient and TK-deficient cells can be selected in a medium containing 6-thioguanine, 8-azaguanine (hereinafter abbreviated as 8AG), or 5'-bromodeoxyuridine, respectively. Normal cells are killed because they incorporate these pyrimidine analogs into their DNA. Meanwhile, cells that are deficient in these enzymes can survive in the selection medium, since they cannot incorporate these pyrimidine analogs. In addition, a selection marker referred to as G418 resistance provided by the neomycin-resistant gene confers resistance to 2-deoxystreptamine antibiotics (gentamycin analogs). Various types of myeloma cells that are suitable for cell fusion are known.

For example, myeloma cells including the following cells can be preferably used:

P3(P3x63Ag8.653) (J. Immunol. (1979) 123 (4), 1548-1550);
 P3x63Ag8U.1 (Current Topics in Microbiology and Immunology (1978)81, 1-7);
 NS-1 (C. Eur. J. Immunol. (1976)6 (7), 511-519);
 MPC-11 (Cell (1976) 8 (3), 405-415);
 SP2/0 (Nature (1978) 276 (5685), 269-270);
 FO (J. Immunol. Methods (1980) 35 (1-2), 1-21);
 S194/5.XX0.BU.1 (J. Exp. Med. (1978) 148 (1), 313-323);
 R210 (Nature (1979) 277 (5692), 131-133), etc.

Cell fusions between the immunocytes and myeloma cells are essentially carried out using known methods, for example, a method by Kohler and Milstein *et al.* (Methods Enzymol. (1981) 73: 3-46).

More specifically, cell fusion can be carried out, for example, in a conventional culture medium in the presence of a cell fusion-promoting agent. The fusion-promoting agents include, for example, polyethylene glycol (PEG) and Sendai virus (HVJ). If required, an auxiliary substance such as dimethyl sulfoxide is also added to improve fusion efficiency.

The ratio of immune cells to myeloma cells may be determined at one's own discretion,

preferably, for example, one myeloma cell for every one to ten immunocytes. Culture media to be used for cell fusions include, for example, media that are suitable for the growth of myeloma cell lines, such as RPMI1640 medium and MEM medium, and other conventional culture medium used for this type of cell culture. In addition, serum supplements such as fetal calf serum (FCS) may be preferably added to the culture medium.

For cell fusion, predetermined amounts of the above immune cells and myeloma cells are mixed well in the above culture medium. Then, a PEG solution (for example, the average molecular weight is about 1,000 to 6,000) prewarmed to about 37°C is added thereto at a concentration of generally 30% to 60% (w/v). This is gently mixed to produce desired fusion cells (hybridomas). Then, an appropriate culture medium mentioned above is gradually added to the cells, and this is repeatedly centrifuged to remove the supernatant. Thus, cell fusion agents and such which are unfavorable to hybridoma growth can be removed.

The hybridomas thus obtained can be selected by culture using a conventional selective medium, for example, HAT medium (a culture medium containing hypoxanthine, aminopterin, and thymidine). Cells other than the desired hybridomas (non-fused cells) can be killed by continuing culture in the above HAT medium for a sufficient period of time. Typically, the period is several days to several weeks. Then, hybridomas producing the desired antibody are screened and singly cloned by conventional limiting dilution methods.

The hybridomas thus obtained can be selected using a selection medium based on the selection marker possessed by the myeloma used for cell fusion. For example, HGPRT- or TK-deficient cells can be selected by culture using the HAT medium (a culture medium containing hypoxanthine, aminopterin, and thymidine). Specifically, when HAT-sensitive myeloma cells are used for cell fusion, cells successfully fused with normal cells can selectively proliferate in the HAT medium. Cells other than the desired hybridomas (non-fused cells) can be killed by continuing culture in the above HAT medium for a sufficient period of time. Specifically, desired hybridomas can be selected by culture for generally several days to several weeks. Then, hybridomas producing the desired antibody are screened and singly cloned by conventional limiting dilution methods.

Desired antibodies can be preferably selected and singly cloned by screening methods based on known antigen/antibody reaction. For example, an IL-6R-binding monoclonal antibody can bind to IL-6R expressed on the cell surface. Such a monoclonal antibody can be screened by fluorescence activated cell sorting (FACS). FACS is a system that assesses the binding of an antibody to cell surface by analyzing cells contacted with a fluorescent antibody using laser beam, and measuring the fluorescence emitted from individual cells.

To screen for hybridomas that produce a monoclonal antibody of the present invention by FACS, IL-6R-expressing cells are first prepared. Cells preferably used for screening are

mammalian cells in which IL-6R is forcedly expressed. As control, the activity of an antibody to bind to cell-surface IL-6R can be selectively detected using non-transformed mammalian cells as host cells. Specifically, hybridomas producing an anti-IL-6R monoclonal antibody can be isolated by selecting hybridomas that produce an antibody which binds to cells forced to express IL-6R, but not to host cells.

Alternatively, the activity of an antibody to bind to immobilized IL-6R-expressing cells can be assessed based on the principle of ELISA. For example, IL-6R-expressing cells are immobilized to the wells of an ELISA plate. Culture supernatants of hybridomas are contacted with the immobilized cells in the wells, and antibodies that bind to the immobilized cells are detected. When the monoclonal antibodies are derived from mouse, antibodies bound to the cells can be detected using an anti-mouse immunoglobulin antibody. Hybridomas producing a desired antibody having the antigen-binding ability are selected by the above screening, and they can be cloned by a limiting dilution method or the like.

Monoclonal antibody-producing hybridomas thus prepared can be passaged in a conventional culture medium, and stored in liquid nitrogen for a long period.

The above hybridomas are cultured by a conventional method, and desired monoclonal antibodies can be prepared from the culture supernatants. Alternatively, the hybridomas are administered to and grown in compatible mammals, and monoclonal antibodies are prepared from the ascites. The former method is suitable for preparing antibodies with high purity.

Antibodies encoded by antibody genes that are cloned from antibody-producing cells such as the above hybridomas can also be preferably used. A cloned antibody gene is inserted into an appropriate vector, and this is introduced into a host to express the antibody encoded by the gene. Methods for isolating antibody genes, inserting the genes into vectors, and transforming host cells have already been established, for example, by Vandamme *et al.* (Eur. J. Biochem. (1990) 192(3), 767-775). Methods for producing recombinant antibodies are also known as described below.

For example, a cDNA encoding the variable region (V region) of an anti-IL-6R antibody is prepared from hybridoma cells expressing the anti-IL-6R antibody. For this purpose, total RNA is first extracted from hybridomas. Methods used for extracting mRNAs from cells include, for example:

- the guanidine ultracentrifugation method (Biochemistry (1979) 18(24), 5294-5299), and
- the AGPC method (Anal. Biochem. (1987) 162(1), 156-159)

Extracted mRNAs can be purified using the mRNA Purification Kit (GE Healthcare Bioscience) or such. Alternatively, kits for extracting total mRNA directly from cells, such as the QuickPrep mRNA Purification Kit (GE Healthcare Bioscience), are also commercially available. mRNAs can be prepared from hybridomas using such kits. cDNAs encoding the

antibody V region can be synthesized from the prepared mRNAs using a reverse transcriptase. cDNAs can be synthesized using the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit (Seikagaku Co.) or such. Furthermore, the SMART RACE cDNA amplification kitTM (Clontech) and the PCR-based 5'-RACE method (Proc. Natl. Acad. Sci. USA (1988) 85(23), 8998-9002; Nucleic Acids Res. (1989) 17(8), 2919-2932) can be appropriately used to synthesize and amplify cDNAs. In such a cDNA synthesis process, appropriate restriction enzyme sites described below may be introduced into both ends of a cDNA.

The cDNA fragment of interest is purified from the resulting PCR product, and then this is ligated to a vector DNA. A recombinant vector is thus constructed, and introduced into *E. coli* or such. After colony selection, the desired recombinant vector can be prepared from the colony-forming *E. coli*. Then, whether the recombinant vector has the cDNA nucleotide sequence of interest is tested by a known method such as the dideoxy nucleotide chain termination method.

The 5'-RACE method which uses primers to amplify the variable region gene is conveniently used for isolating the gene encoding the variable region. First, a 5'-RACE cDNA library is constructed by cDNA synthesis using RNAs extracted from hybridoma cells as a template. A commercially available kit such as the SMART RACE cDNA amplification kitTM is appropriately used to synthesize the 5'-RACE cDNA library.

The antibody gene is amplified by PCR using the prepared 5'-RACE cDNA library as a template. Primers for amplifying the mouse antibody gene can be designed based on known antibody gene sequences. The nucleotide sequences of the primers vary depending on the immunoglobulin subclass. Therefore, it is preferable that the subclass is determined in advance using a commercially available kit such as the Iso Strip mouse monoclonal antibody isotyping kit (Roche Diagnostics).

Specifically, for example, primers that allow amplification of genes encoding $\gamma 1$, $\gamma 2a$, $\gamma 2b$, and $\gamma 3$ heavy chains and κ and λ light chains are used to isolate mouse IgG-encoding genes. In general, a primer that anneals to a constant region site close to the variable region is used as a 3'-side primer to amplify an IgG variable region gene. Meanwhile, a primer attached to a 5' RACE cDNA library construction kit is used as a 5'-side primer.

PCR products thus amplified are used to reshape immunoglobulins composed of a combination of heavy and light chains. A desired antibody can be selected using the IL-6R-binding activity of a reshaped immunoglobulin as an indicator. For example, when the objective is to isolate an antibody against IL-6R, it is more preferred that the binding of the antibody to IL-6R is specific. An IL-6R-binding antibody can be screened, for example, by the following steps:

(1) contacting an IL-6R-expressing cell with an antibody comprising the V region encoded by a

cDNA isolated from a hybridoma;

- (2) detecting the binding of the antibody to the IL-6R-expressing cell; and
- (3) selecting an antibody that binds to the IL-6R-expressing cell.

Methods for detecting the binding of an antibody to IL-6R-expressing cells are known.

- 5 Specifically, the binding of an antibody to IL-6R-expressing cells can be detected by the above-described techniques such as FACS. Immobilized samples of IL-6R-expressing cells are appropriately used to assess the binding activity of an antibody.

- Preferred antibody screening methods that use the binding activity as an indicator also include panning methods using phage vectors. Screening methods using phage vectors are
 10 advantageous when the antibody genes are isolated from heavy-chain and light-chain subclass libraries from a polyclonal antibody-expressing cell population. Genes encoding the heavy-chain and light-chain variable regions can be linked by an appropriate linker sequence to form a single-chain Fv (scFv). Phages presenting scFv on their surface can be produced by inserting a gene encoding scFv into a phage vector. The phages are contacted with an antigen
 15 of interest. Then, a DNA encoding scFv having the binding activity of interest can be isolated by collecting phages bound to the antigen. This process can be repeated as necessary to enrich scFv having the binding activity of interest.

- After isolation of the cDNA encoding the V region of the anti-IL-6R antibody of interest, the cDNA is digested with restriction enzymes that recognize the restriction sites introduced into
 20 both ends of the cDNA. Preferred restriction enzymes recognize and cleave a nucleotide sequence that occurs in the nucleotide sequence of the antibody gene at a low frequency. Furthermore, a restriction site for an enzyme that produces a sticky end is preferably introduced into a vector to insert a single-copy digested fragment in the correct orientation. The cDNA encoding the V region of the anti-IL-6R antibody is digested as described above, and this is
 25 inserted into an appropriate expression vector to construct an antibody expression vector. In this case, if a gene encoding the antibody constant region (C region) and a gene encoding the above V region are fused in-frame, a chimeric antibody is obtained. Herein, "chimeric antibody" means that the origin of the constant region is different from that of the variable region. Thus, in addition to mouse/human heterochimeric antibodies, human/human allochimeric antibodies
 30 are included in the chimeric antibodies of the present invention. A chimeric antibody expression vector can be constructed by inserting the above V region gene into an expression vector that already has the constant region. Specifically, for example, a recognition sequence for a restriction enzyme that excises the above V region gene can be appropriately placed on the 5' side of an expression vector carrying a DNA encoding a desired antibody constant region (C
 35 region). A chimeric antibody expression vector is constructed by fusing in frame the two genes digested with the same combination of restriction enzymes.

To produce an anti-IL-6R monoclonal antibody, antibody genes are inserted into an expression vector so that the genes are expressed under the control of an expression regulatory region. The expression regulatory region for antibody expression includes, for example, enhancers and promoters. Furthermore, an appropriate signal sequence may be attached to the amino terminus so that the expressed antibody is secreted to the outside of cells. In the Examples described later, a peptide having the amino acid sequence MGWSCIILFLVATATGVHS (SEQ ID NO: 3) are used as a signal sequence. Meanwhile, other appropriate signal sequences may be attached. The expressed polypeptide is cleaved at the carboxyl terminus of the above sequence, and the resulting polypeptide is secreted to the outside of cells as a mature polypeptide. Then, appropriate host cells are transformed with the expression vector, and recombinant cells expressing the anti-IL-6R antibody-encoding DNA are obtained.

DNAs encoding the antibody heavy chain (H chain) and light chain (L chain) are separately inserted into different expression vectors to express the antibody gene. An antibody molecule having the H and L chains can be expressed by co-transfecting the same host cell with vectors into which the H-chain and L-chain genes are respectively inserted. Alternatively, host cells can be transformed with a single expression vector into which DNAs encoding the H and L chains are inserted (see WO 1994/011523).

There are various known host cell/expression vector combinations for antibody preparation by introducing isolated antibody genes into appropriate hosts. All of these expression systems are applicable to isolation of the antigen-binding domains of the present invention. Appropriate eukaryotic cells used as host cells include animal cells, plant cells, and fungal cells. Specifically, the animal cells include, for example, the following cells.

- (1) mammalian cells: CHO, COS, myeloma, baby hamster kidney (BHK), HeLa, Vero, human embryonic kidney (HEK) 293, FreestyleTM293, or such;
- (2) amphibian cells: *Xenopus* oocytes, or such; and
- (3) insect cells: sf9, sf21, Tn5, or such.

In addition, as a plant cell, an antibody gene expression system using cells derived from the *Nicotiana* genus such as *Nicotiana tabacum* is known. Callus cultured cells can be appropriately used to transform plant cells.

Furthermore, the following cells can be used as fungal cells:

- yeasts: the *Saccharomyces* genus such as *Saccharomyces cerevisiae*, and the *Pichia* genus such as *Pichia pastoris*; and
- filamentous fungi: the *Aspergillus* genus such as *Aspergillus niger*.

Furthermore, antibody gene expression systems that utilize prokaryotic cells are also known. For example, when using bacterial cells, *E. coli* cells, *Bacillus subtilis* cells, and such

can suitably be utilized in the present invention. Expression vectors carrying the antibody genes of interest are introduced into these cells by transfection. The transfected cells are cultured *in vitro*, and the desired antibody can be prepared from the culture of transformed cells.

In addition to the above-described host cells, transgenic animals can also be used to
5 produce a recombinant antibody. That is, the antibody can be obtained from an animal into which the gene encoding the antibody of interest is introduced. For example, the antibody gene can be constructed as a fusion gene by inserting in frame into a gene that encodes a protein produced specifically in milk. Goat β -casein or such can be used, for example, as the protein secreted in milk. DNA fragments containing the fused gene inserted with the antibody gene is
10 injected into a goat embryo, and then this embryo is introduced into a female goat. Desired antibodies can be obtained as a protein fused with the milk protein from milk produced by the transgenic goat born from the embryo-recipient goat (or progeny thereof). In addition, to increase the volume of milk containing the desired antibody produced by the transgenic goat, hormones can be administered to the transgenic goat as necessary (Ebert, K. M. *et al.*,
15 Bio/Technology (1994) 12 (7), 699-702).

When an antigen-binding molecule described herein is administered to human, an antigen-binding domain derived from a genetically recombinant antibody that has been artificially modified to reduce the heterologous antigenicity against human and such, can be appropriately used as the antigen-binding domain of the molecule. Such genetically
20 recombinant antibodies include, for example, humanized antibodies. These modified antibodies are appropriately produced by known methods.

An antibody variable region used to produce the antigen-binding domain of an antigen-binding molecule described herein is generally formed by three complementarity-determining regions (CDRs) that are separated by four framework regions
25 (FRs). CDR is a region that substantially determines the binding specificity of an antibody. The amino acid sequences of CDRs are highly diverse. On the other hand, the FR-forming amino acid sequences often have high identity even among antibodies with different binding specificities. Therefore, generally, the binding specificity of a certain antibody can be introduced to another antibody by CDR grafting.

30 A humanized antibody is also called a reshaped human antibody. Specifically, humanized antibodies prepared by grafting the CDR of a non-human animal antibody such as a mouse antibody to a human antibody and such are known. Common genetic engineering techniques for obtaining humanized antibodies are also known. Specifically, for example, overlap extension PCR is known as a method for grafting a mouse antibody CDR to a human FR.
35 In overlap extension PCR, a nucleotide sequence encoding a mouse antibody CDR to be grafted is added to primers for synthesizing a human antibody FR. Primers are prepared for each of the

four FRs. It is generally considered that when grafting a mouse CDR to a human FR, selecting a human FR that has high identity to a mouse FR is advantageous for maintaining the CDR function. That is, it is generally preferable to use a human FR comprising an amino acid sequence which has high identity to the amino acid sequence of the FR adjacent to the mouse CDR to be grafted.

Nucleotide sequences to be ligated are designed so that they will be connected to each other in frame. Human FRs are individually synthesized using the respective primers. As a result, products in which the mouse CDR-encoding DNA is attached to the individual FR-encoding DNAs are obtained. Nucleotide sequences encoding the mouse CDR of each product are designed so that they overlap with each other. Then, complementary strand synthesis reaction is conducted to anneal the overlapping CDR of the products synthesized using a human antibody gene as template. Human FRs are ligated *via* the mouse CDR sequences by this reaction.

The full length V region gene, in which three CDRs and four FRs are ultimately ligated, is amplified using primers that anneal to its 5'- or 3'-end, which are added with suitable restriction enzyme recognition sequences. An expression vector for humanized antibody can be produced by inserting the DNA obtained as described above and a DNA that encodes a human antibody C region into an expression vector so that they will ligate in frame. After the recombinant vector is transfected into a host to establish recombinant cells, the recombinant cells are cultured, and the DNA encoding the humanized antibody is expressed to produce the humanized antibody in the cell culture (see, European Patent Publication No. EP 239400 and International Patent Publication No. WO 1996/002576).

By qualitatively or quantitatively measuring and evaluating the antigen-binding activity of the humanized antibody produced as described above, one can suitably select human antibody FRs that allow CDRs to form a favorable antigen-binding site when ligated through the CDRs. Amino acid residues in FRs may be substituted as necessary, so that the CDRs of a reshaped human antibody form an appropriate antigen-binding site. For example, amino acid sequence mutations can be introduced into FRs by applying the PCR method used for grafting a mouse CDR into a human FR. More specifically, partial nucleotide sequence mutations can be introduced into primers that anneal to the FR. Nucleotide sequence mutations are introduced into the FRs synthesized by using such primers. Mutant FR sequences having the desired characteristics can be selected by measuring and evaluating the activity of the amino acid-substituted mutant antibody to bind to the antigen by the above-mentioned method (Cancer Res. (1993) 53: 851-856).

Alternatively, desired human antibodies can be obtained by immunizing transgenic animals having the entire repertoire of human antibody genes (see WO 1993/012227; WO

1992/003918; WO 1994/002602; WO 1994/025585; WO 1996/034096; WO 1996/033735) by DNA immunization.

Furthermore, techniques for preparing human antibodies by panning using human antibody libraries are also known. For example, the V region of a human antibody is expressed as a single-chain antibody (scFv) on phage surface by the phage display method. Phages expressing an scFv that binds to the antigen can be selected. The DNA sequence encoding the human antibody V region that binds to the antigen can be determined by analyzing the genes of selected phages. The DNA sequence of the scFv that binds to the antigen is determined. An expression vector is prepared by fusing the V region sequence in frame with the C region sequence of a desired human antibody, and inserting this into an appropriate expression vector. The expression vector is introduced into cells appropriate for expression such as those described above. The human antibody can be produced by expressing the human antibody-encoding gene in the cells. These methods are already known (see WO 1992/001047; WO 1992/020791; WO 1993/006213; WO 1993/011236; WO 1993/019172; WO 1995/001438; WO 1995/015388).

In addition to the techniques described above, techniques of B cell cloning (identification of each antibody-encoding sequence, cloning and its isolation; use in constructing expression vector in order to prepare each antibody (IgG1, IgG2, IgG3, or IgG4 in particular); and such) such as described in Bernasconi *et al.* (Science (2002) 298: 2199-2202) or in WO 2008/081008 can be appropriately used to isolate antibody genes.

EU numbering system and Kabat's numbering system

According to the methods used in the present invention, amino acid positions assigned to antibody CDR and FR are specified according to Kabat's numbering (Sequences of Proteins of Immunological Interest (National Institute of Health, Bethesda, Md., 1987 and 1991)). Herein, when an antigen-binding molecule is an antibody or antigen-binding fragment, variable region amino acids are indicated according to Kabat's numbering system, while constant region amino acids are indicated according to EU numbering system based on Kabat's amino acid positions.

Conditions of ion concentration

Conditions of metal ion concentration

In one embodiment of the present invention, the ion concentration refers to a metal ion concentration. "Metal ions" refer to ions of group I elements except hydrogen such as alkaline metals and copper group elements, group II elements such as alkaline earth metals and zinc group elements, group III elements except boron, group IV elements except carbon and silicon, group VIII elements such as iron group and platinum group elements, elements belonging to subgroup A of groups V, VI, and VII, and metal elements such as antimony, bismuth, and

polonium. Metal atoms have the property of releasing valence electrons to become cations. This is referred to as ionization tendency. Metals with strong ionization tendency are deemed to be chemically active.

In the present invention, preferred metal ions include, for example, calcium ion.

5 Calcium ion is involved in modulation of many biological phenomena, including contraction of muscles such as skeletal, smooth, and cardiac muscles; activation of movement, phagocytosis, and the like of leukocytes; activation of shape change, secretion, and the like of platelets; activation of lymphocytes; activation of mast cells including secretion of histamine; cell responses mediated by catecholamine α receptor or acetylcholine receptor; exocytosis; release of
10 transmitter substances from neuron terminals; and axoplasmic flow in neurons. Known intracellular calcium ion receptors include troponin C, calmodulin, parvalbumin, and myosin light chain, which have several calcium ion-binding sites and are believed to be derived from a common origin in terms of molecular evolution. There are also many known calcium-binding motifs. Such well-known motifs include, for example, cadherin domains, EF-hand of
15 calmodulin, C2 domain of Protein kinase C, Gla domain of blood coagulation protein Factor IX, C-type lectins of acyarglycoprotein receptor and mannose-binding receptor, A domains of LDL receptors, annexin, thrombospondin type 3 domain, and EGF-like domains.

In the present invention, when the metal ion is calcium ion, the conditions of calcium ion concentration include low calcium ion concentrations and high calcium ion concentrations.

20 “The binding activity varies depending on calcium ion concentrations” means that the antigen-binding activity of an antigen-binding molecule varies due to the difference in the conditions between low and high calcium ion concentrations. For example, the antigen-binding activity of an antigen-binding molecule may be higher at a high calcium ion concentration than at a low calcium ion concentration. Alternatively, the antigen-binding activity of an
25 antigen-binding molecule may be higher at a low calcium ion concentration than at a high calcium ion concentration.

Herein, the high calcium ion concentration is not particularly limited to a specific value; however, the concentration may preferably be selected between 100 μ M and 10 mM. In another embodiment, the concentration may be selected between 200 μ M and 5 mM. In an
30 alternative embodiment, the concentration may be selected between 500 μ M and 2.5 mM. In still another embodiment, the concentration may be selected between 200 μ M and 2 mM. Furthermore, the concentration may be selected between 400 μ M and 1.5 mM. In particular, a concentration selected between 500 μ M and 2.5 mM, which is close to the plasma (blood) concentration of calcium ion *in vivo*, is preferred.

35 Herein, the low calcium ion concentration is not particularly limited to a specific value; however, the concentration may preferably be selected between 0.1 μ M and 30 μ M. In another

embodiment, the concentration may be selected between 0.2 μM and 20 μM . In still another embodiment, the concentration may be selected between 0.5 μM and 10 μM . In an alternative embodiment, the concentration may be selected between 1 μM and 5 μM . Furthermore, the concentration may be selected between 2 μM and 4 μM . In particular, a concentration selected between 1 μM and 5 μM , which is close to the concentration of ionized calcium in early endosomes *in vivo*, is preferred.

Herein, "the antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration" means that the antigen-binding activity of an antigen-binding molecule is weaker at a calcium ion concentration selected between 0.1 μM and 30 μM than at a calcium ion concentration selected between 100 μM and 10 mM. Preferably, it means that the antigen-binding activity of an antigen-binding molecule is weaker at a calcium ion concentration selected between 0.5 μM and 10 μM than at a calcium ion concentration selected between 200 μM and 5 mM. It particularly preferably means that the antigen-binding activity at the calcium ion concentration in the early endosome *in vivo* is weaker than that at the *in vivo* plasma calcium ion concentration; and specifically, it means that the antigen-binding activity of an antigen-binding molecule is weaker at a calcium ion concentration selected between 1 μM and 5 μM than at a calcium ion concentration selected between 500 μM and 2.5 mM.

Whether the antigen-binding activity of an antigen-binding molecule is changed depending on metal ion concentrations can be determined, for example, by the use of known measurement methods such as those described in the section "Binding Activity" above. For example, in order to confirm that the antigen-binding activity of an antigen-binding molecule becomes higher at a high calcium ion concentration than at a low calcium ion concentration, the antigen-binding activity of the antigen-binding molecule at low and high calcium ion concentrations is compared.

In the present invention, the expression "the antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration" can also be expressed as "the antigen-binding activity of an antigen-binding molecule is higher at a high calcium ion concentration than at a low calcium ion concentration". In the present invention, "the antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration" is sometimes written as "the antigen-binding ability is weaker at a low calcium ion concentration than at a high calcium ion concentration". Also, "the antigen-binding activity at a low calcium ion concentration is reduced to be lower than that at a high calcium ion concentration" may be written as "the antigen-binding ability at a low calcium ion concentration is made weaker than that at a high calcium ion concentration".

When determining the antigen-binding activity, the conditions other than calcium ion concentration can be appropriately selected by those skilled in the art, and are not particularly

limited. For example, the activity can be determined at 37°C in HEPES buffer. For example, Biacore (GE Healthcare) or such can be used for the determination. When the antigen is a soluble antigen, the antigen-binding activity of an antigen-binding molecule can be assessed by flowing the antigen as an analyte over a chip onto which the antigen-binding molecule is
 5 immobilized. When the antigen is a membrane antigen, the binding activity of an antigen-binding molecule to the membrane antigen can be assessed by flowing the antigen-binding molecule as an analyte over a chip onto which the antigen is immobilized.

As long as the antigen-binding activity of an antigen-binding molecule of the present invention at a low calcium ion concentration is weaker than that at a high calcium ion
 10 concentration, the ratio of the antigen-binding activity between that at a low calcium ion concentration and at a high calcium ion concentration is not particularly limited; and the value of $KD(Ca\ 3\mu M) / KD(Ca\ 2\ mM)$, which is the ratio of the dissociation constant (KD) for an antigen at a low calcium ion concentration to the KD at a high calcium ion concentration, is preferably 2 or more; more preferably the value of $KD(Ca\ 3\mu M) / KD(Ca\ 2\ mM)$ is 10 or more;
 15 and still more preferably the value of $KD(Ca\ 3\mu M) / KD(Ca\ 2\ mM)$ is 40 or more. The upper limit of $KD(Ca\ 3\mu M) / KD(Ca\ 2\ mM)$ value is not particularly limited, and may be any value such as 400, 1000, or 10000, as long as the molecule can be produced by the techniques of those skilled in the art. Alternatively, the value of $KD(Ca\ 3\mu M) / KD(Ca\ 1.2\ mM)$ is specified. That is, the value of $KD(Ca\ 3\mu M) / KD(Ca\ 1.2\ mM)$ is 2 or more; more preferably the value of
 20 $KD(Ca\ 3\mu M) / KD(Ca\ 1.2\ mM)$ is 10 or more; and still more preferably the value of $KD(Ca\ 3\mu M) / KD(Ca\ 1.2\ mM)$ is 40 or more. The upper limit of $KD(Ca\ 3\mu M) / KD(Ca\ 1.2\ mM)$ value is not particularly limited, and may be any value such as 400, 1000, or 10000, as long as the molecule can be produced by the techniques of those skilled in the art.

When the antigen is a soluble antigen, KD (dissociation constant) can be used to
 25 represent the antigen-binding activity. Meanwhile, when the antigen is a membrane antigen, apparent KD (apparent dissociation constant) can be used to represent the activity. KD (dissociation constant) and apparent KD (apparent dissociation constant) can be determined by methods known to those skilled in the art, for example, using Biacore (GE healthcare), Scatchard plot, or flow cytometer.

30 Alternatively, for example, the dissociation rate constant (kd) can also be preferably used as an index to represent the ratio of the antigen-binding activity of an antigen-binding molecule of the present invention between low and high calcium concentrations. When the dissociation rate constant (kd) is used instead of the dissociation constant (KD) as an index to represent the binding activity ratio, the ratio of the dissociation rate constant (kd) between low
 35 and high calcium concentrations, i.e. the value of $kd(\text{low calcium concentration})/kd(\text{high calcium concentration})$, is preferably 2 or more, more preferably 5 or more, still more preferably

10 or more, and yet more preferably 30 or more. The upper limit of the K_d (low calcium concentration)/ k_d (high calcium concentration) value is not particularly limited, and can be any value such as 50, 100, or 200 as long as the molecule can be produced by techniques known to those skilled in the art.

5 When the antigen is a soluble antigen, k_d (dissociation rate constant) can be used to represent the antigen-binding activity. Meanwhile, when the antigen is a membrane antigen, apparent k_d (apparent dissociation rate constant) can be used to represent the antigen-binding activity. The k_d (dissociation rate constant) and apparent k_d (apparent dissociation rate constant) can be determined by methods known to those skilled in the art, for example, using
10 Biacore (GE healthcare) or flow cytometer. In the present invention, when the antigen-binding activity of an antigen-binding molecule is determined at different calcium ion concentrations, it is preferable to use the same conditions except for the calcium concentrations.

For example, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one
15 embodiment of the present invention, can be obtained via screening of antigen-binding domains or antibodies including the steps of:

- (a) determining the antigen-binding activity of an antigen-binding domain or antibody at a low calcium concentration;
- (b) determining the antigen-binding activity of an antigen-binding domain or antibody at a high
20 calcium concentration; and
- (c) selecting an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium concentration than at a high calcium concentration.

Moreover, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one
25 embodiment of the present invention, can be obtained via screening of antigen-binding domains or antibodies, or a library thereof, including the steps of:

- (a) contacting an antigen with an antigen-binding domain or antibody, or a library thereof at a high calcium concentration;
- (b) incubating at a low calcium concentration an antigen-binding domain or antibody that has
30 bound to the antigen in step (a); and
- (c) isolating an antigen-binding domain or antibody dissociated in step (b).

Furthermore, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one
embodiment of the present invention, can be obtained via screening of antigen-binding domains
35 or antibodies, or a library thereof, including the steps of:

- (a) contacting an antigen with a library of antigen-binding domains or antibodies at a low

calcium concentration;

(b) selecting an antigen-binding domain or antibody which does not bind to the antigen in step (a);

5 (c) allowing the antigen-binding domain or antibody selected in step (b) to bind to the antigen at a high calcium concentration ; and

(d) isolating an antigen-binding domain or antibody that has bound to the antigen in step (c).

10 In addition, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one embodiment of the present invention, can be obtained by a screening method comprising the steps of:

(a) contacting at a high calcium concentration a library of antigen-binding domains or antibodies with a column onto which an antigen is immobilized;

(b) eluting an antigen-binding domain or antibody that has bound to the column in step (a) from the column at a low calcium concentration; and

15 (c) isolating the antigen-binding domain or antibody eluted in step (b).

Furthermore, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one embodiment of the present invention, can be obtained by a screening method comprising the steps of:

20 (a) allowing at a low calcium concentration a library of antigen-binding domains or antibodies to pass through a column onto which an antigen is immobilized;

(b) collecting an antigen-binding domain or antibody that has been eluted without binding to the column in step (a);

25 (c) allowing the antigen-binding domain or antibody collected in step (b) to bind to the antigen at a high calcium concentration; and

(d) isolating an antigen-binding domain or antibody that has bound to the antigen in step (c).

30 Moreover, an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which is one embodiment of the present invention, can be obtained by a screening method comprising the steps of:

(a) contacting an antigen with a library of antigen-binding domains or antibodies at a high calcium concentration;

(b) obtaining an antigen-binding domain or antibody that has bound to the antigen in step (a);

35 (c) incubating at a low calcium concentration the antigen-binding domain or antibody obtained in step (b); and

(d) isolating an antigen-binding domain or antibody whose antigen-binding activity in step (c) is

weaker than the criterion for the selection of step (b).

The above-described steps may be repeated twice or more times. Thus, the present invention provides antigen-binding domains or antibodies whose antigen-binding activity is lower at a low calcium ion concentration than at a high calcium ion concentration, which are obtained by screening methods that further comprises the step of repeating twice or more times steps (a) to (c) or (a) to (d) in the above-described screening methods. The number of cycles of steps (a) to (c) or (a) to (d) is not particularly limited, but generally is 10 or less.

In the screening methods of the present invention, the antigen-binding activity of an antigen-binding domain or antibody at a low calcium concentration is not particularly limited as long as it is antigen-binding activity at an ionized calcium concentration of between 0.1 μM and 30 μM , but preferably is antigen-binding activity at an ionized calcium concentration of between 0.5 μM and 10 μM . More preferably, it is antigen-binding activity at the ionized calcium concentration in the early endosome *in vivo*, specifically, between 1 μM and 5 μM . Meanwhile, the antigen-binding activity of an antigen-binding domain or antibody at a high calcium concentration is not particularly limited, as long as it is antigen-binding activity at an ionized calcium concentration of between 100 μM and 10 mM, but preferably is antigen-binding activity at an ionized calcium concentration of between 200 μM and 5 mM. More preferably, it is antigen-binding activity at the ionized calcium concentration in plasma *in vivo*, specifically, between 0.5 mM and 2.5 mM.

The antigen-binding activity of an antigen-binding domain or antibody can be measured by methods known to those skilled in the art. Conditions other than the ionized calcium concentration can be determined by those skilled in the art. The antigen-binding activity of an antigen-binding domain or antibody can be evaluated as a dissociation constant (KD), apparent dissociation constant (apparent KD), dissociation rate constant (kd), apparent dissociation constant (apparent kd), and such. These can be determined by methods known to those skilled in the art, for example, using Biacore (GE healthcare), Scatchard plot, or FACS.

In the present invention, the step of selecting an antigen-binding domain or antibody whose antigen-binding activity is higher at a high calcium concentration than at a low calcium concentration is synonymous with the step of selecting an antigen-binding domain or antibody whose antigen-binding activity is lower at a low calcium concentration than at a high calcium concentration.

As long as the antigen-binding activity is higher at a high calcium concentration than at a low calcium concentration, the difference in the antigen-binding activity between high and low calcium concentrations is not particularly limited; however, the antigen-binding activity at a high calcium concentration is preferably twice or more, more preferably 10 times or more, and still more preferably 40 times or more than that at a low calcium concentration.

Antigen-binding domains or antibodies of the present invention to be screened by the screening methods described above may be any antigen-binding domains and antibodies. For example, it is possible to screen the above-described antigen-binding domains or antibodies. For example, antigen-binding domains or antibodies having natural sequences or substituted amino acid sequences may be screened.

Libraries

In an embodiment, an antigen-binding domain or antibody of the present invention can be obtained from a library that is mainly composed of a plurality of antigen-binding molecules whose sequences are different from one another and whose antigen-binding domains have at least one amino acid residue that alters the antigen-binding activity of the antigen-binding molecules depending on ion concentrations. The ion concentrations preferably include, for example, metal ion concentration and hydrogen ion concentration.

Herein, a "library" refers to a plurality of antigen-binding molecules or a plurality of fusion polypeptides containing antigen-binding molecules, or nucleic acids or polynucleotides encoding their sequences. The sequences of a plurality of antigen-binding molecules or a plurality of fusion polypeptides containing antigen-binding molecules in a library are not identical, but are different from one another.

Herein, the phrase "sequences are different from one another" in the expression "a plurality of antigen-binding molecules whose sequences are different from one another" means that the sequences of antigen-binding molecules in a library are different from one another. Specifically, in a library, the number of sequences different from one another reflects the number of independent clones with different sequences, and may also be referred to as "library size". The library size of a conventional phage display library ranges from 10^6 to 10^{12} . The library size can be increased up to 10^{14} by the use of known techniques such as ribosome display. However, the actual number of phage particles used in panning selection of a phage library is in general 10-10000 times greater than the library size. This excess multiplicity is also referred to as "the number of library equivalents", and means that there are 10 to 10,000 individual clones that have the same amino acid sequence. Thus, in the present invention, the phrase "sequences are different from one another" means that the sequences of independent antigen-binding molecules in a library, excluding library equivalents, are different from one another. More specifically, the above means that there are 10^6 to 10^{14} antigen-binding molecules whose sequences are different from one another, preferably 10^7 to 10^{12} molecules, more preferably 10^8 to 10^{11} molecules, and particularly preferably 10^8 to 10^{12} molecules whose sequences are different from one another.

Herein, the phrase "a plurality of" in the expression "a library mainly composed of a

plurality of antigen-binding molecules" generally refers to, in the case of, for example, antigen-binding molecules, fusion polypeptides, polynucleotide molecules, vectors, or viruses of the present invention, a group of two or more types of the substance. For example, when two or more substances are different from one another in a particular characteristic, this means that there are two or more types of the substance. Such examples may include, for example, mutant amino acids observed at specific amino acid positions in an amino acid sequence. For example, when there are two or more antigen-binding molecules of the present invention whose sequences are substantially the same or preferably the same except for flexible residues or except for particular mutant amino acids at hypervariable positions exposed on the surface, there are a plurality of antigen-binding molecules of the present invention. In another example, when there are two or more polynucleotide molecules whose sequences are substantially the same or preferably the same except for nucleotides encoding flexible residues or nucleotides encoding mutant amino acids of hypervariable positions exposed on the surface, there are a plurality of polynucleotide molecules of the present invention.

In addition, herein, the phrase "mainly composed of" in the expression "a library mainly composed of a plurality of antigen-binding molecules" reflects the number of antigen-binding molecules whose antigen-binding activity varies depending on ion concentrations, among independent clones with different sequences in a library. Specifically, it is preferable that there are at least 10^4 antigen-binding molecules having such binding activity in a library. More preferably, antigen-binding domains of the present invention can be obtained from a library containing at least 10^5 antigen-binding molecules having such binding activity. Still more preferably, antigen-binding domains of the present invention can be obtained from a library containing at least 10^6 antigen-binding molecules having such binding activity. Particularly preferably, antigen-binding domains of the present invention can be obtained from a library containing at least 10^7 antigen-binding molecules having such binding activity. Yet more preferably, antigen-binding domains of the present invention can be obtained from a library containing at least 10^8 antigen-binding molecules having such binding activity. Alternatively, this may also be preferably expressed as the ratio of the number of antigen-binding molecules whose antigen-binding activity varies depending on ion concentrations with respect to the number of independent clones having different sequences in a library. Specifically, antigen-binding domains of the present invention can be obtained from a library in which antigen-binding molecules having such binding activity account for 0.1% to 80%, preferably 0.5% to 60%, more preferably 1% to 40%, still more preferably 2% to 20%, and particularly preferably 4% to 10% of independent clones with different sequences in the library. In the case of fusion polypeptides, polynucleotide molecules, or vectors, similar expressions may be possible using the number of molecules or the ratio to the total number of molecules. In the

case of viruses, similar expressions may also be possible using the number of virions or the ratio to total number of virions.

Amino acids that alter the antigen-binding activity of antigen-binding domains depending on calcium ion concentrations

Antigen-binding domains or antibodies of the present invention to be screened by the above-described screening methods may be prepared in any manner. For example, when the metal ion is calcium ion, it is possible to use preexisting antibodies, preexisting libraries (phage library, etc.), antibodies or libraries prepared from hybridomas obtained by immunizing animals or from B cells of immunized animals, antibodies or libraries obtained by introducing amino acids capable of chelating calcium (for example, aspartic acid and glutamic acid) or unnatural amino acid mutations into the above-described antibodies or libraries (calcium-chelatable amino acids (such as aspartic acid and glutamic acid), libraries with increased content of unnatural amino acids, libraries prepared by introducing calcium-chelatable amino acids (such as aspartic acid and glutamic acid) or unnatural amino acid mutations at particular positions, or the like.

Examples of the amino acids that alter the antigen-binding activity of antigen-binding molecules depending on ion concentrations as described above may be any types of amino acids as long as the amino acids form a calcium-binding motif. Calcium-binding motifs are well known to those skilled in the art and have been described in details (for example, Springer *et al.* (Cell (2000) 102, 275-277); Kawasaki and Kretsinger (Protein Prof. (1995) 2, 305-490); Moncrief *et al.* (J. Mol. Evol. (1990) 30, 522-562); Chauvaux *et al.* (Biochem. J. (1990) 265, 261-265); Bairoch and Cox (FEBS Lett. (1990) 269, 454-456); Davis (New Biol. (1990) 2, 410-419); Schaefer *et al.* (Genomics (1995) 25, 638-643); Economou *et al.* (EMBO J. (1990) 9, 349-354); Wurzburg *et al.* (Structure. (2006) 14, 6, 1049-1058)). Specifically, any known calcium-binding motifs, including type C lectins such as ASGPR, CD23, MBR, and DC-SIGN, can be included in antigen-binding molecules of the present invention. Preferred examples of such preferred calcium-binding motifs also include, in addition to those described above, for example, the calcium-binding motif in the antigen-binding domain of SEQ ID NO: 62.

Furthermore, as amino acids that alter the antigen-binding activity of antigen-binding molecules depending on calcium ion concentrations, for example, amino acids having metal-chelating activity may also be preferably used. Examples of such metal-chelating amino acids include, for example, serine (Ser(S)), threonine (Thr(T)), asparagine (Asn(N)), glutamine (Gln(Q)), aspartic acid (Asp(D)), and glutamic acid (Glu(E)).

Positions in the antigen-binding domains at which the above-described amino acids are contained are not particularly limited to particular positions, and may be any positions within the heavy chain variable region or light chain variable region that forms an antigen-binding domain,

as long as they alter the antigen-binding activity of antigen-binding molecules depending on calcium ion concentrations. Specifically, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose heavy chain antigen-binding domains contain amino acids that alter the antigen-binding activity of the antigen-binding molecules depending on calcium ion concentrations. In another non-limiting embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose heavy chain CDR3 domains contain the above-mentioned amino acids. In still another non-limiting embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose heavy chain CDR3 domains contain the above-mentioned amino acids at positions 95, 96, 100a, and/or 101 as indicated according to the Kabat numbering system.

Meanwhile, in a non-limiting embodiment of the present invention, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain antigen-binding domains contain amino acids that alter the antigen-binding activity of antigen-binding molecules depending on calcium ion concentrations. In another embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR1 domains contain the above-mentioned amino acids. In still another embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR1 domains contain the above-mentioned amino acids at positions 30, 31, and/or 32 as indicated according to the Kabat numbering system.

In another non-limiting embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR2 domains contain the above-mentioned amino acid residues. In yet another embodiment, the present invention provides libraries mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR2 domains contain the above-mentioned amino acid residues at position 50 as indicated according to the Kabat numbering system.

In still another non-limiting embodiment of the present invention, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR3 domains contain the above-mentioned amino acid residues. In an alternative

embodiment, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chain CDR3 domains contain the above-mentioned amino acid residues at position 92 as indicated according to the Kabat numbering system.

5 Furthermore, in a different embodiment of the present invention, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and in which two or three CDRs selected from the above-described light chain CDR1, CDR2, and CDR3 contain the
10 aforementioned amino acid residues. Moreover, antigen-binding domains of the present invention can be obtained from a library mainly composed of antigen-binding molecules whose sequences are different from one another and whose light chains contain the aforementioned amino acid residues at any one or more of positions 30, 31, 32, 50, and/or 92 as indicated according to the Kabat numbering system.

In a particularly preferred embodiment, the framework sequences of the light chain
15 and/or heavy chain variable region of an antigen-binding molecule preferably contain human germ line framework sequences. Thus, in an embodiment of the present invention, when the framework sequences are completely human sequences, it is expected that when such an antigen-binding molecule of the present invention is administered to humans (for example, to treat diseases), it induces little or no immunogenic response. In the above sense, the phrase
20 "containing a germ line sequence" in the present invention means that a part of the framework sequences of the present invention is identical to a part of any human germ line framework sequences. For example, when the heavy chain FR2 sequence of an antigen-binding molecule of the present invention is a combination of heavy chain FR2 sequences of different human germ line framework sequences, such a molecule is also an antigen-binding molecule of the present
25 invention "containing a germ line sequence".

Preferred examples of the frameworks include, for example, fully human framework region sequences currently known, which are included in the website of V-Base
or others. Those framework region sequences can be appropriately used as a germ line sequence contained in an antigen-binding molecule of the
30 present invention. The germ line sequences may be categorized according to their similarity (Tomlinson *et al.* (J. Mol. Biol. (1992) 227, 776-798); Williams and Winter (Eur. J. Immunol. (1993) 23, 1456-1461); Cox *et al.* (Nat. Genetics (1994) 7, 162-168)). Appropriate germ line sequences can be selected from V κ , which is grouped into seven subgroups; V λ , which is grouped into ten subgroups; and V H , which is grouped into seven subgroups.

35 Fully human V H sequences preferably include, but are not limited to, for example, V H sequences of:

subgroup VH1 (for example, VH1-2, VH1-3, VH1-8, VH1-18, VH1-24, VH1-45, VH1-46, VH1-58, and VH1-69);

subgroup VH2 (for example, VH2-5, VH2-26, and VH2-70);

subgroup VH3 (VH3-7, VH3-9, VH3-11, VH3-13, VH3-15, VH3-16, VH3-20, VH3-21, VH3-23, 5 VH3-30, VH3-33, VH3-35, VH3-38, VH3-43, VH3-48, VH3-49, VH3-53, VH3-64, VH3-66, VH3-72, VH3-73, and VH3-74);

subgroup VH4 (VH4-4, VH4-28, VH4-31, VH4-34, VH4-39, VH4-59, and VH4-61);

subgroup VH5 (VH5-51);

subgroup VH6 (VH6-1); and

10 subgroup VH7 (VH7-4 and VH7-81).

These are also described in known documents (Matsuda *et al.* (J. Exp. Med. (1998) 188, 1973-1975)) and such, and thus persons skilled in the art can appropriately design antigen-binding molecules of the present invention based on the information of these sequences. It is also preferable to use other fully human frameworks or framework sub-regions.

15 Fully human VK sequences preferably include, but are not limited to, for example: A20, A30, L1, L4, L5, L8, L9, L11, L12, L14, L15, L18, L19, L22, L23, L24, O2, O4, O8, O12, O14, and O18, grouped into subgroup Vk1;

A1, A2, A3, A5, A7, A17, A18, A19, A23, O1, and O11, grouped into subgroup Vk2;

A11, A27, L2, L6, L10, L16, L20, and L25, grouped into subgroup Vk3;

20 B3, grouped into subgroup Vk4;

B2 (herein also referred to as Vk5-2), grouped into subgroup Vk5; and

A10, A14, and A26, grouped into subgroup VK6

(Kawasaki *et al.* (Eur. J. Immunol. (2001) 31, 1017-1028); Schable and Zachau (Biol. Chem. Hoppe Seyler (1993) 374, 1001-1022); Brensing-Kuppers *et al.* (Gene (1997) 191, 173-181)).

25 Fully human VL sequences preferably include, but are not limited to, for example:

V1-2, V1-3, V1-4, V1-5, V1-7, V1-9, V1-11, V1-13, V1-16, V1-17, V1-18, V1-19, V1-20, and V1-22, grouped into subgroup VL1;

V2-1, V2-6, V2-7, V2-8, V2-11, V2-13, V2-14, V2-15, V2-17, and V2-19, grouped into subgroup VL1;

30 V3-2, V3-3, and V3-4, grouped into subgroup VL3;

V4-1, V4-2, V4-3, V4-4, and V4-6, grouped into subgroup VL4; and

V5-1, V5-2, V5-4, and V5-6, grouped into subgroup VL5 (Kawasaki *et al.* (Genome Res. (1997) 7, 250-261)).

Normally, these framework sequences are different from one another at one or more 35 amino acid residues. These framework sequences can be used in combination with "at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule

depending on ion concentrations" of the present invention. Other examples of the fully human frameworks used in combination with "at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule depending on ion concentrations" of the present invention include, but are not limited to, for example, KOL, NEWM, REI, EU, TUR, 5 TEI, LAY, and POM (for example, Kabat *et al.* (1991) *supra*; Wu *et al.* (J. Exp. Med. (1970) 132, 211-250)).

Without being bound by a particular theory, one reason for the expectation that the use of germ line sequences precludes adverse immune responses in most individuals is believed to be as follows. As a result of the process of affinity maturation during normal immune responses, 10 somatic mutation occurs frequently in the variable regions of immunoglobulin. Such mutations mostly occur around CDRs whose sequences are hypervariable, but also affect residues of framework regions. Such framework mutations do not exist on the germ line genes, but they are less likely to be immunogenic in patients. This is because the normal human population is exposed to most of the framework sequences expressed from the germ line genes, and as a result 15 of immunotolerance, these germ line frameworks are expected to have low or no immunogenicity in patients. To maximize the possibility of immunotolerance, variable region-encoding genes may be selected from a group of commonly occurring functional germ line genes.

Known methods such as site-directed mutagenesis (Kunkel *et al.* (Proc. Natl. Acad. Sci. 20 USA (1985) 82, 488-492)) and overlap extension PCR can be appropriately employed to produce antigen-binding molecules of the present invention in which the above-described framework sequences contain amino acids that alter the antigen-binding activity of the antigen-binding molecules depending on calcium ion concentrations.

For example, a library which contains a plurality of antigen-binding molecules of the present 25 invention whose sequences are different from one another can be constructed by combining heavy chain variable regions prepared as a randomized variable region sequence library with a light chain variable region selected as a framework sequence originally containing at least one amino acid residue that alters the antigen-binding activity of the antigen-binding molecule depending on calcium ion concentrations. As a non-limiting example, when the ion 30 concentration is calcium ion concentration, such preferred libraries include, for example, those constructed by combining the light chain variable region sequence belonging to the Vk5-2 family represented by the light chain variable region sequence of SEQ ID NO: 62 (Vk5-2) and the heavy chain variable region produced as a randomized variable region sequence library.

Alternatively, a light chain variable region sequence selected as a framework region 35 originally containing at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule as mentioned above can be design to contain various amino acid

residues other than the above amino acid residues. Herein, such residues are referred to as flexible residues. The number and position of flexible residues are not particularly limited as long as the antigen-binding activity of the antigen-binding molecule of the present invention varies depending on ion concentrations. Specifically, the CDR sequences and/or FR sequences
5 of the heavy chain and/or light chain may contain one or more flexible residues. For example, when the ion concentration is calcium ion concentration, non-limiting examples of flexible residues to be introduced into the light chain variable region sequence of SEQ ID NO: 62 (Vk5-2) include the amino acid residues listed in Tables 1 or 2.

10 [Table 1]

CDR	Kabat NUMBERING	70 % OF AMINO ACID OF THE TOTAL				
CDR1	28	S : 100%				
	29	I : 100%				
	30	E : 72%	N : 14%	S : 14%		
	31	D : 100%				
	32	D : 100%				
	33	L : 100%				
	34	A : 70%	N : 30%			
CDR2	50	E : 100%				
	51	A : 100%				
	52	S : 100%				
	53	H : 5%	N : 25%	S : 45%	T : 25%	
	54	L : 100%				
	55	Q : 100%				
	56	S : 100%				
CDR3	90	Q:100%				
	91	H : 25%	S : 15%	R : 15%	Y : 45%	
	92	D : 80%	N : 10%	S : 10%		
	93	D : 5%	G : 10%	N : 25%	S : 50%	R : 10%
	94	S : 50%	Y : 50%			
	95	P : 100%				
	96	L : 50%	Y : 50%			

[Table 2]

CDR	Kabat NUMBERING	30 % OF AMINO ACID OF THE TOTAL				
CDR1	28	S : 100%				
	29	I : 100%				
	30	E : 83%	S : 17%			
	31	D : 100%				
	32	D : 100%				
	33	L : 100%				
	34	A : 70%	N : 30%			
CDR2	50	H : 100%				
	51	A : 100%				
	52	S : 100%				
	53	H : 5%	N : 25%	S : 45%	T : 25%	
	54	L : 100%				
	55	Q : 100%				
	56	S : 100%				
CDR3	90	Q : 100%				
	91	H : 25%	S : 15%	R : 15%	Y : 45%	
	92	D : 80%	N : 10%	S : 10%		
	93	D : 5%	G : 10%	N : 25%	S : 50%	R : 10%
	94	S : 50%	Y : 50%			
	95	P : 100%				
	96	L : 50%	Y : 50%			

Herein, flexible residues refer to amino acid residue variations present at hypervariable positions at which several different amino acids are present on the light chain and heavy chain variable regions when the amino acid sequences of known and/or native antibodies or antigen-binding domains are compared. Hypervariable positions are generally located in the CDR. In an embodiment, the data provided by Kabat, Sequences of Proteins of Immunological Interest (National Institute of Health Bethesda Md.) (1987 and 1991) is useful to determine hypervariable positions in known and/or native antibodies. Furthermore, databases on the Internet provide the collected sequences of many human light chains and heavy chains and their locations. The

information on the sequences and locations is useful to determine hypervariable positions in the present invention. According to the present invention, when a certain amino acid position has preferably about 2 to about 20 possible amino acid residue variations, preferably about 3 to about 19, preferably about 4 to about 18, preferably 5 to 17, preferably 6 to 16, preferably 7 to 15, preferably 8 to 14, preferably 9 to 13, and preferably 10 to 12 possible amino acid residue variations, the position is hypervariable. In some embodiments, a certain amino acid position may have preferably at least about 2, preferably at least about 4, preferably at least about 6, preferably at least about 8, preferably about 10, and preferably about 12 amino acid residue variations.

Alternatively, a library containing a plurality of antigen-binding molecules of the present invention whose sequences are different from one another can be constructed by combining heavy chain variable regions produced as a randomized variable region sequence library with light chain variable regions into which at least one amino acid residue that alters the antigen-binding activity of antigen-binding molecules depending on ion concentrations as mentioned above is introduced. When the ion concentration is calcium ion concentration, non-limiting examples of such libraries preferably include, for example, libraries in which heavy chain variable regions produced as a randomized variable region sequence library are combined with light chain variable region sequences in which a particular residue(s) in a germ line sequence such as SEQ ID NO: 5 (Vk1), SEQ ID NO: 6 (Vk2), SEQ ID NO: 7 (Vk3), or SEQ ID NO: 8 (Vk4) has been substituted with at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule depending on calcium ion concentrations. Non-limiting examples of such amino acid residues include amino acid residues in light chain CDR1. Furthermore, non-limiting examples of such amino acid residues include amino acid residues in light chain CDR2. In addition, non-limiting, other examples of such amino acid residues also include amino acid residues in light chain CDR3.

Non-limiting examples of such amino acid residues contained in light chain CDR1 include those at positions 30, 31, and/or 32 in the CDR1 of light chain variable region as indicated by Kabat numbering. Furthermore, non-limiting examples of such amino acid residues contained in light chain CDR2 include an amino acid residue at position 50 in the CDR2 of light chain variable region as indicated by Kabat numbering. Moreover, non-limiting examples of such amino acid residues contained in light chain CDR3 include an amino acid residue at position 92 in the CDR3 of light chain variable region as indicated by Kabat numbering. These amino acid residues can be contained alone or in combination as long as they form a calcium-binding motif and/or as long as the antigen-binding activity of an antigen-binding molecule varies depending on calcium ion concentrations. Meanwhile, as troponin C, calmodulin, parvalbumin, and myosin light chain, which have several calcium

ion-binding sites and are believed to be derived from a common origin in terms of molecular evolution, are known, the light chain CDR1, CDR2, and/or CDR3 can be designed to have their binding motifs. For example, it is possible to use cadherin domains, EF hand of calmodulin, C2 domain of Protein kinase C, Gla domain of blood coagulation protein Factor IX, C type lectins of acyarglycoprotein receptor and mannose-binding receptor, A domains of LDL receptors, annexin, thrombospondin type 3 domain, and EGF-like domains in an appropriate manner for the above purposes.

When heavy chain variable regions produced as a randomized variable region sequence library and light chain variable regions into which at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule depending on ion concentrations has been introduced are combined as described above, the sequences of the light chain variable regions can be designed to contain flexible residues in the same manner as described above. The number and position of such flexible residues are not particularly limited to particular embodiments as long as the antigen-binding activity of antigen-binding molecules of the present invention varies depending on ion concentrations. Specifically, the CDR sequences and/or FR sequences of heavy chain and/or light chain can contain one or more flexible residues. When the ion concentration is calcium ion concentration, non-limiting examples of flexible residues to be introduced into the sequence of light chain variable region include the amino acid residues listed in Tables 1 and 2.

The preferred heavy chain variable regions to be combined include, for example, randomized variable region libraries. Known methods are combined as appropriate to produce a randomized variable region library. In a non-limiting embodiment of the present invention, an immune library constructed based on antibody genes derived from lymphocytes of animals immunized with a specific antigen, patients with infections, persons with an elevated antibody titer in blood as a result of vaccination, cancer patients, or auto immune disease patients, may be preferably used as a randomized variable region library.

In another non-limiting embodiment of the present invention, a synthetic library produced by replacing the CDR sequences of V genes in genomic DNA or functional reshaped V genes with a set of synthetic oligonucleotides containing sequences encoding codon sets of an appropriate length can also be preferably used as a randomized variable region library. In this case, since sequence diversity is observed in the heavy chain CDR3 sequence, it is also possible to replace the CDR3 sequence only. A criterion of giving rise to diversity in amino acids in the variable region of an antigen-binding molecule is that diversity is given to amino acid residues at surface-exposed positions in the antigen-binding molecule. The surface-exposed position refers to a position that is considered to be able to be exposed on the surface and/or contacted with an antigen, based on structure, ensemble of structures, and/or modeled structure of an

antigen-binding molecule. In general, such positions are CDRs. Preferably, surface-exposed positions are determined using coordinates from a three-dimensional model of an antigen-binding molecule using a computer program such as the InsightII program (Accelrys). Surface-exposed positions can be determined using algorithms known in the art (for example, 5 Lee and Richards (J. Mol. Biol. (1971) 55, 379-400); Connolly (J. Appl. Cryst. (1983) 16, 548-558)). Determination of surface-exposed positions can be performed using software suitable for protein modeling and three-dimensional structural information obtained from an antibody. Software that can be used for these purposes preferably includes SYBYL Biopolymer Module software (Tripos Associates). Generally or preferably, when an algorithm requires a 10 user input size parameter, the "size" of a probe which is used in the calculation is set at about 1.4 Angstrom or smaller in radius. Furthermore, methods for determining surface-exposed regions and areas using software for personal computers are described by Pacios (Comput. Chem. (1994) 18 (4), 377-386; J. Mol. Model. (1995) 1, 46-53).

In another non-limiting embodiment of the present invention, a naive library, which is 15 constructed from antibody genes derived from lymphocytes of healthy persons and whose repertoire consists of naive sequences, which are antibody sequences with no bias, can also be particularly preferably used as a randomized variable region library (Gejima *et al.* (Human Antibodies (2002) 11, 121-129); Cardoso *et al.* (Scand. J. Immunol. (2000) 51, 337-344)). Herein, an amino acid sequence comprising a naive sequence refers to an amino acid sequence 20 obtained from such a naive library.

In one embodiment of the present invention, an antigen-binding domain of the present invention can be obtained from a library containing a plurality of antigen-binding molecules of the present invention whose sequences are different from one another, prepared by combining light chain variable regions constructed as a randomized variable region sequence library with a 25 heavy chain variable region selected as a framework sequence that originally contains "at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule depending on ion concentrations". When the ion concentration is calcium ion concentration, non-limiting examples of such libraries preferably include those constructed by combining light chain variable regions constructed as a randomized variable region sequence library with the 30 sequence of heavy chain variable region of SEQ ID NO: 9 (6RL#9-IgG1) or SEQ ID NO: 10 (6KC4-1#85-IgG1). Alternatively, such a library can be constructed by selecting appropriate light chain variable regions from those having germ line sequences, instead of light chain variable regions constructed as a randomized variable region sequence library. Such preferred libraries include, for example, those in which the sequence of heavy chain variable region of 35 SEQ ID NO: 9 (6RL#9-IgG1) or SEQ ID NO: 10 (6KC4-1#85-IgG1) is combined with light chain variable regions having germ line sequences.

Alternatively, the sequence of an heavy chain variable region selected as a framework sequence that originally contains "at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule" as mentioned above can be designed to contain flexible residues. The number and position of the flexible residues are not particularly limited as long as the antigen-binding activity of an antigen-binding molecule of the present invention varies depending on ion concentrations. Specifically, the CDR and/or FR sequences of heavy chain and/or light chain can contain one or more flexible residues. When the ion concentration is calcium ion concentration, non-limiting examples of flexible residues to be introduced into the sequence of heavy chain variable region of SEQ ID NO: 9 (6RL#9-IgG1) include all amino acid residues of heavy chain CDR1 and CDR2 and the amino acid residues of the heavy chain CDR3 except those at positions 95, 96, and/or 100a. Alternatively, non-limiting examples of flexible residues to be introduced into the sequence of heavy chain variable region of SEQ ID NO: 10 (6KC4-l#85-IgG1) include all amino acid residues of heavy chain CDR1 and CDR2 and the amino acid residues of the heavy chain CDR3 except those at amino acid positions 95 and/or 101.

Alternatively, a library containing a plurality of antigen-binding molecules whose sequences are different from one another can be constructed by combining light chain variable regions constructed as a randomized variable region sequence library or light chain variable regions having germ line sequences with heavy chain variable regions into which "at least one amino acid residue responsible for the ion concentration-dependent change in the antigen-binding activity of an antigen-binding molecule" has been introduced as mentioned above. When the ion concentration is calcium ion concentration, non-limiting examples of such libraries preferably include those in which light chain variable regions constructed as a randomized variable region sequence library or light chain variable regions having germ line sequences are combined with the sequence of a heavy chain variable region in which a particular residue(s) has been substituted with at least one amino acid residue that alters the antigen-binding activity of an antigen-binding molecule depending on calcium ion concentrations. Non-limiting examples of such amino acid residues include amino acid residues of the heavy chain CDR1. Further non-limiting examples of such amino acid residues include amino acid residues of the heavy chain CDR2. In addition, non-limiting examples of such amino acid residues also include amino acid residues of the heavy chain CDR3. Non-limiting examples of such amino acid residues of heavy chain CDR3 include the amino acids of positions 95, 96, 100a, and/or 101 in the CDR3 of heavy chain variable region as indicated by the Kabat numbering. Furthermore, these amino acid residues can be contained alone or in combination as long as they form a calcium-binding motif and/or the antigen-binding activity of an antigen-binding molecule varies depending on calcium ion concentrations.

When light chain variable regions constructed as a randomized variable region sequence library or light chain variable regions having germ line sequence are combined with a heavy chain variable region into which at least one amino acid residue that alter the antigen-binding activity of an antigen-binding molecule depending on ion concentrations as mentioned above has been introduced, the sequence of the heavy chain variable region can also be designed to contain flexible residues in the same manner as described above. The number and position of flexible residues are not particularly limited as long as the antigen-binding activity of an antigen-binding molecule of the present invention varies depending on ion concentrations. Specifically, the heavy chain CDR and/or FR sequences may contain one or more flexible residues.

Furthermore, randomized variable region libraries can be preferably used as amino acid sequences of CDR1, CDR2, and/or CDR3 of the heavy chain variable region other than the amino acid residues that alter the antigen-binding activity of an antigen-binding molecule. When germ line sequences are used as light chain variable regions, non-limiting examples of such sequences include those of SEQ ID NO: 5 (Vk1), SEQ ID NO: 6 (Vk2), SEQ ID NO: 7 (Vk3), and SEQ ID NO: 8 (Vk4).

Any of the above-described amino acids that alter the antigen-binding activity of an antigen-binding molecule depending on calcium ion concentrations can be preferably used, as long as they form a calcium-binding motif. Specifically, such amino acids include electron-donating amino acids. Preferred examples of such electron-donating amino acids include serine, threonine, asparagine, glutamic acid, aspartic acid, and glutamic acid.

Condition of hydrogen ion concentrations

In an embodiment of the present invention, the condition of ion concentrations refers to the condition of hydrogen ion concentrations or pH condition. In the present invention, the concentration of proton, i.e., the nucleus of hydrogen atom, is treated as synonymous with hydrogen index (pH). When the activity of hydrogen ion in an aqueous solution is represented as aH^+ , pH is defined as $-\log_{10}aH^+$. When the ionic strength of the aqueous solution is low (for example, lower than 10^{-3}), aH^+ is nearly equal to the hydrogen ion strength. For example, the ionic product of water at 25°C and 1 atmosphere is $K_w = aH^+aOH = 10^{-14}$, and therefore in pure water, $aH^+ = aOH = 10^{-7}$. In this case, pH=7 is neutral; an aqueous solution whose pH is lower than 7 is acidic or whose pH is greater than 7 is alkaline.

In the present invention, when pH condition is used as the ion concentration condition, pH conditions include high hydrogen ion concentrations or low pHs, i.e., an acidic pH range, and low hydrogen ion concentrations or high pHs, i.e., a neutral pH range. "The binding activity varies depending on pH condition" means that the antigen-binding activity of an antigen-binding molecule varies due to the difference in conditions of a high hydrogen ion concentration or low

pH (an acidic pH range) and a low hydrogen ion concentration or high pH (a neutral pH range). This includes, for example, the case where the antigen-binding activity of an antigen-binding molecule is higher in a neutral pH range than in an acidic pH range and the case where the antigen-binding activity of an antigen-binding molecule is higher in an acidic pH range than in a
5 neutral pH range.

In the present specification, neutral pH range is not limited to a specific value and is preferably selected from between pH 6.7 and pH 10.0. In another embodiment, the pH can be selected from between pH 6.7 and pH 9.5. In still another embodiment, the pH can be selected from between pH 7.0 and pH 9.0. In yet another embodiment, the pH can be selected from
10 between pH 7.0 and pH 8.0. In particular, the preferred pH includes pH 7.4, which is close to the pH of plasma (blood) *in vivo*.

In the present specification, an acidic pH range is not limited to a specific value and is preferably selected from between pH 4.0 and pH 6.5. In another embodiment, the pH can be selected from between pH 4.5 and pH 6.5. In still another embodiment, the pH can be selected
15 from between pH 5.0 and pH 6.5. In yet another embodiment, the pH can be selected from between pH 5.5 and pH 6.5. In particular, the preferred pH includes pH 5.8, which is close to the ionized calcium concentration in the early endosome *in vivo*.

In the present invention, "the antigen-binding activity of an antigen-binding molecule at a high hydrogen ion concentration or low pH (an acidic pH range) is lower than that at a low
20 hydrogen ion concentration or high pH (a neutral pH range)" means that the antigen-binding activity of an antigen-binding molecule at a pH selected from between pH 4.0 and pH 6.5 is weaker than that at a pH selected from between pH 6.7 and pH 10.0; preferably means that the antigen-binding activity of an antigen-binding molecule at a pH selected from between pH 4.5 and pH 6.5 is weaker than that at a pH selected from between pH 6.7 and pH 9.5; more
25 preferably, means that the antigen-binding activity of an antigen-binding molecule at a pH selected from between pH 5.0 and pH 6.5 is weaker than that at a pH selected from between pH 7.0 and pH 9.0; still more preferably means that the antigen-binding activity of an antigen-binding molecule at a pH selected from between pH 5.5 and pH 6.5 is weaker than that at a pH selected from between pH 7.0 and pH 8.0; particularly preferably means that the
30 antigen-binding activity at the pH in the early endosome *in vivo* is weaker than the antigen-binding activity at the pH of plasma *in vivo*; and specifically means that the antigen-binding activity of an antigen-binding molecule at pH 5.8 is weaker than the antigen-binding activity at pH 7.4.

Whether the antigen-binding activity of an antigen-binding molecule has changed by the
35 pH condition can be determined, for example, by the use of known measurement methods such as those described in the section "Binding Activity" above. Specifically, the binding activity is

measured under different pH conditions using the measurement methods described above. For example, the antigen-binding activity of an antigen-binding molecule is compared under the conditions of acidic pH range and neutral pH range to confirm that the antigen-binding activity of the antigen-binding molecule changes to be higher under the condition of neutral pH range than that under the condition of acidic pH range.

Furthermore, in the present invention, the expression "the antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range" can also be expressed as "the antigen-binding activity of an antigen-binding molecule at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is higher than that at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range". In the present invention, "the antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range" may be described as "the antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is weaker than the antigen-binding ability at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range". Alternatively, "the antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is reduced to be lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range" may be described as "the antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is reduced to be weaker than the antigen-binding ability at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range".

The conditions other than hydrogen ion concentration or pH for measuring the antigen-binding activity may be suitably selected by those skilled in the art and are not particularly limited. Measurements can be carried out, for example, at 37°C using HEPES buffer. Measurements can be carried out, for example, using Biacore (GE Healthcare). When the antigen is a soluble antigen, the antigen-binding activity of an antigen-binding molecule can be determined by assessing the binding activity to the soluble antigen by pouring the antigen as an analyte into a chip immobilized with the antigen-binding molecule. When the antigen is a membrane antigen, the binding activity to the membrane antigen can be assessed by pouring the antigen-binding molecule as an analyte into a chip immobilized with the antigen.

As long as the antigen-binding activity of an antigen-binding molecule of the present invention at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range is weaker than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, the ratio of the antigen-binding activity between that at a high hydrogen ion concentration or low pH, i.e., an acidic pH range, and at a low hydrogen ion concentration or high pH, i.e., a neutral pH range is not particularly limited, and the value of $KD(pH\ 5.8) / KD(pH\ 7.4)$, which is the ratio of the

dissociation constant (KD) for an antigen at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range to the KD at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is preferably 2 or more; more preferably the value of KD (pH 5.8) / KD (pH 7.4) is 10 or more; and still more preferably the value of KD (pH 5.8) / KD (pH 7.4) is 40 or more. The
5 upper limit of KD (pH 5.8) / KD (pH 7.4) value is not particularly limited, and may be any value such as 400, 1000, or 10000, as long as the molecule can be produced by the techniques of those skilled in the art.

When the antigen is a soluble antigen, the dissociation constant (KD) can be used as the value for antigen-binding activity. Meanwhile, when the antigen is a membrane antigen, the
10 apparent dissociation constant (KD) can be used. The dissociation constant (KD) and apparent dissociation constant (KD) can be measured by methods known to those skilled in the art, and Biacore (GE healthcare), Scatchard plot, flow cytometer, and such can be used.

Alternatively, for example, the dissociation rate constant (kd) can be suitably used as an index for indicating the ratio of the antigen-binding activity of an antigen-binding molecule of
15 the present invention between that at a high hydrogen ion concentration or low pH, i.e., an acidic pH range and a low hydrogen ion concentration or high pH, i.e., a neutral pH range. When kd (dissociation rate constant) is used as an index for indicating the binding activity ratio instead of KD (dissociation constant), the value of kd (in an acidic pH range) / kd (in a neutral pH range), which is the ratio of kd (dissociation rate constant) for the antigen at a high hydrogen ion
20 concentration or low pH, i.e., in an acidic pH range to kd (dissociation rate constant) at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is preferably 2 or more, more preferably 5 or more, still more preferably 10 or more, and yet more preferably 30 or more. The upper limit of kd (in an acidic pH range) / kd (in a neutral pH range) value is not particularly limited, and may be any value such as 50, 100, or 200, as long as the molecule can be produced
25 by the techniques of those skilled in the art.

When the antigen is a soluble antigen, the dissociation rate constant (kd) can be used as the value for antigen-binding activity and when the antigen is a membrane antigen, the apparent dissociation rate constant (kd) can be used. The dissociation rate constant (kd) and apparent dissociation rate constant (kd) can be determined by methods known to those skilled in the art,
30 and Biacore (GE healthcare), flow cytometer, and such may be used. In the present invention, when the antigen-binding activity of an antigen-binding molecule is measured at different hydrogen ion concentrations, i.e., pHs, conditions other than the hydrogen ion concentration, i.e., pH, are preferably the same.

For example, an antigen-binding domain or antibody whose antigen-binding activity at a
35 high hydrogen ion concentration or low pH, i.e., in an acidic pH range is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is one embodiment

provided by the present invention, can be obtained via screening of antigen-binding domains or antibodies, comprising the following steps (a) to (c):

- (a) obtaining the antigen-binding activity of an antigen-binding domain or antibody in an acidic pH range;
- 5 (b) obtaining the antigen-binding activity of an antigen-binding domain or antibody in a neutral pH range; and
- (c) selecting an antigen-binding domain or antibody whose antigen-binding activity in the acidic pH range is lower than that in the neutral pH range.

Alternatively, an antigen-binding domain or antibody whose antigen-binding activity at
10 a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is one embodiment provided by the present invention, can be obtained via screening of antigen-binding domains or antibodies, or a library thereof, comprising the following steps (a) to (c):

- (a) contacting an antigen-binding domain or antibody, or a library thereof, in a neutral pH range
15 with an antigen;
- (b) placing in an acidic pH range the antigen-binding domain or antibody bound to the antigen in step (a); and
- (c) isolating the antigen-binding domain or antibody dissociated in step (b).

An antigen-binding domain or antibody whose antigen-binding activity at a high
20 hydrogen ion concentration or low pH, i.e., in an acidic pH range is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is another embodiment provided by the present invention, can be obtained *via* screening of antigen-binding domains or antibodies, or a library thereof, comprising the following steps (a) to (d):

- (a) contacting in an acidic pH range an antigen with a library of antigen-binding domains or
25 antibodies;
- (b) selecting the antigen-binding domain or antibody which does not bind to the antigen in step (a);
- (c) allowing the antigen-binding domain or antibody selected in step (b) to bind with the antigen in a neutral pH range; and
- 30 (d) isolating the antigen-binding domain or antibody bound to the antigen in step (c).

An antigen-binding domain or antibody whose antigen-binding activity at a high
hydrogen ion concentration or low pH, i.e., in an acidic pH range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is even another embodiment provided by the present invention, can be obtained by a screening method
35 comprising the following steps (a) to (c):

- (a) contacting in a neutral pH range a library of antigen-binding domains or antibodies with a

column immobilized with an antigen;

(b) eluting in an acidic pH range from the column the antigen-binding domain or antibody bound to the column in step (a); and

(c) isolating the antigen-binding domain or antibody eluted in step (b).

5 An antigen-binding domain or antibody whose antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH, range is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is still another embodiment provided by the present invention, can be obtained by a screening method comprising the following steps (a) to (d):

10 (a) allowing, in an acidic pH range, a library of antigen-binding domains or antibodies to pass a column immobilized with an antigen;

(b) collecting the antigen-binding domain or antibody eluted without binding to the column in step (a);

(c) allowing the antigen-binding domain or antibody collected in step (b) to bind with the antigen
15 in a neutral pH range; and

(d) isolating the antigen-binding domain or antibody bound to the antigen in step (c).

 An antigen-binding domain or antibody whose antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, which is yet another
20 embodiment provided by the present invention, can be obtained by a screening method comprising the following steps (a) to (d):

(a) contacting an antigen with a library of antigen-binding domains or antibodies in a neutral pH range;

(b) obtaining the antigen-binding domain or antibody bound to the antigen in step (a);

25 (c) placing in an acidic pH range the antigen-binding domain or antibody obtained in step (b);
and

(d) isolating the antigen-binding domain or antibody whose antigen-binding activity in step (c) is weaker than the standard selected in step (b).

 The above-described steps may be repeated twice or more times. Thus, the present
30 invention provides antigen-binding domains and antibodies whose antigen-binding activity in an acidic pH range is lower than that in a neutral pH range, which are obtained by a screening method that further comprises the steps of repeating, twice or more times, steps (a) to (c) or (a) to (d) in the above-described screening methods. The number of times that steps (a) to (c) or (a) to (d) is repeated is not particularly limited; however, the number is 10 or less in general.

35 In the screening methods of the present invention, the antigen-binding activity of an antigen-binding domain or antibody at a high hydrogen ion concentration or low pH, i.e., in an

acidic pH range, is not particularly limited, as long as it is the antigen-binding activity at a pH of between 4.0 and 6.5, and includes the antigen-binding activity at a pH of between 4.5 and 6.6 as the preferred pH. The antigen-binding activity also includes that at a pH of between 5.0 and 6.5, and that at a pH of between 5.5 and 6.5 as another preferred pH. The antigen-binding activity
5 also includes that at the pH in the early endosome *in vivo* as the more preferred pH, and specifically, that at pH5.8. Meanwhile, the antigen-binding activity of an antigen-binding domain or antibody at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is not particularly limited, as long as it is the antigen-binding activity at a pH of between 6.7 and 10, and includes the antigen-binding activity at a pH of between 6.7 and 9.5 as the preferred pH.
10 The antigen-binding activity also includes that at a pH of between 7.0 and 9.5 and that at a pH of between 7.0 and 8.0 as another preferred pH. The antigen-binding activity also includes that at the pH of plasma *in vivo* as the more preferred pH, and specifically, that at pH 7.4.

The antigen-binding activity of an antigen-binding domain or antibody can be measured by methods known to those skilled in the art. Those skilled in the art can suitably determine
15 conditions other than ionized calcium concentration. The antigen-binding activity of an antigen-binding domain or antibody can be assessed based on the dissociation constant (KD), apparent dissociation constant (KD), dissociation rate constant (kd), apparent dissociation rate constant (kd), and such. These can be determined by methods known to those skilled in the art, for example, using Biacore (GE healthcare), Scatchard plot, or FACS.

20 Herein, the step of selecting an antigen-binding domain or antibody whose antigen-binding activity at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is higher than that at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, is synonymous with the step of selecting an antigen-binding domain or antibody whose antigen-binding activity at a high hydrogen ion concentration or low pH, i.e., in an acidic pH
25 range, is lower than that at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range.

As long as the antigen-binding activity at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is higher than that at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, the difference between the antigen-binding activity at a low hydrogen ion
30 concentration or high pH, i.e., a neutral pH range, and that at a high hydrogen ion concentration or low pH, i.e., an acidic pH range, is not particularly limited; however, the antigen-binding activity at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is preferably twice or more, more preferably 10 times or more, and still more preferably 40 times or more than that at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range.

35 The antigen binding domain or antibody of the present invention screened by the screening methods described above may be any antigen-binding domain or antibody, and the

above-mentioned antigen-binding domain or antibody may be screened. For example, antigen-binding domain or antibody having the native sequence may be screened, and antigen-binding domain or antibody in which their amino acid sequences have been substituted may be screened.

5 The antigen-binding domain or antibody of the present invention to be screened by the above-described screening methods may be prepared in any manner. For example, conventional antibodies, conventional libraries (phage library, etc.), antibodies or libraries prepared from B cells of immunized animals or from hybridomas obtained by immunizing animals, antibodies or libraries (libraries with increased content of amino acids with a side chain
10 pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids, libraries introduced with amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acid mutations at specific positions, etc.) obtained by introducing amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acid mutations into the above-described antibodies or libraries may be
15 used.

Methods for obtaining an antigen-binding domain or antibody whose antigen-binding activity at a low hydrogen ion concentration or high pH, i.e., in a neutral pH range, is higher than that at a high hydrogen ion concentration or low pH, i.e., in an acidic pH range, from an antigen-binding domains or antibodies prepared from hybridomas obtained by immunizing
20 animals or from B cells of immunized animals preferably include, for example, the antigen-binding molecule or antibody in which at least one of the amino acids of the antigen-binding domain or antibody is substituted with an amino acid with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or an unnatural amino acid mutation, or the antigen-binding domain or antibody inserted with an amino acid with a side chain pKa of 4.0-8.0
25 (for example, histidine and glutamic acid) or unnatural amino acid, such as those described in WO 2009/125825.

The sites of introducing mutations of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids are not particularly limited, and may be any position as long as the antigen-binding activity in an acidic pH range becomes
30 weaker than that in a neutral pH range (the value of K_D (in an acidic pH range) / K_D (in a neutral pH range) or k_d (in an acidic pH range) / k_d (in a neutral pH range) is increased) as compared to before substitution or insertion. For example, when the antigen-binding molecule is an antibody, antibody variable region and CDRs are suitable. Those skilled in the art can appropriately determine the number of amino acids to be substituted with or the number of
35 amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids to be inserted. It is possible to substitute with a single amino acid having

a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or a single unnatural amino acid; it is possible to insert a single amino acid having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or a single unnatural amino acid; it is possible to substitute with two or more amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or two or more unnatural amino acids; and it is possible to insert two or more amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or two or more unnatural amino acids. Alternatively, other amino acids can be deleted, added, inserted, and/or substituted concomitantly, aside from the substitution into amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids, or the insertion of amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids. Substitution into or insertion of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids can be performed randomly by methods such as histidine scanning, in which the alanine of alanine scanning known to those skilled in the art is replaced with histidine. Antigen-binding molecules exhibiting a greater value of KD (in an acidic pH range) / KD (in a neutral pH range) or kd (in an acidic pH range) / kd (in a neutral pH range) as compared to before the mutation can be selected from antigen-binding domains or antibodies introduced with random insertions or substitution mutations of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids.

Preferred examples of antigen-binding molecules containing the mutation into amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids as described above and whose antigen-binding activity in an acidic pH range is lower than that in a neutral pH range include, antigen-binding molecules whose antigen-binding activity in the neutral pH range after the mutation into amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids is comparable to that before the mutation into amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids. Herein, "an antigen-binding molecule after the mutation with amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids has an antigen-binding activity comparable to that before the mutation with amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids" means that, when taking the antigen-binding activity of an antigen-binding molecule before the mutation with amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids as 100%, the antigen-binding activity of an antigen-binding molecule after the mutation with amino acids having a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids is at least 10% or more, preferably 50% or more, more preferably 80% or more, and still

more preferably 90% or more. The antigen-binding activity after the mutation of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids at pH 7.4 may be higher than that before the mutation of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids at pH 7.4. If the antigen-binding activity of an antigen-binding molecule is decreased due to insertion of or substitution into amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids, the antigen-binding activity can be made to be comparable to that before the insertion of or substitution into amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids, by introducing a substitution, deletion, addition, and/or insertion of one or more amino acids of the antigen-binding molecule. The present invention also includes antigen-binding molecules whose binding activity has been adjusted to be comparable by substitution, deletion, addition, and/or insertion of one or more amino acids after substitution or insertion of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids.

Meanwhile, when an antigen-binding molecule is a substance containing an antibody constant region, preferred embodiments of antigen-binding molecules whose antigen-binding activity at an acidic pH range is lower than that in a neutral pH range include methods in which the antibody constant regions contained in the antigen-binding molecules have been modified. Specific examples of modified antibody constant regions preferably include the constant regions of SEQ ID NOs: 11, 12, 13, and 14.

Amino acids that alter the antigen-binding activity of antigen-binding domain depending on the hydrogen ion concentration conditions

Antigen-binding domains or antibodies of the present invention to be screened by the above-described screening methods may be prepared in any manner. For example, when ion concentration condition is hydrogen ion concentration condition or pH condition, conventional antibodies, conventional libraries (phage library, etc.), antibodies or libraries prepared from B cells of immunized animals or from hybridomas obtained by immunizing animals, antibodies or libraries (libraries with increased content of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids, libraries introduced with mutations of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids at specific positions, etc.) obtained by introducing mutations of amino acids with a side chain pKa of 4.0-8.0 (for example, histidine and glutamic acid) or unnatural amino acids into the above-described antibodies or libraries may be used.

In one non-limiting embodiment of the present invention, a library containing multiple antigen-binding molecules of the present invention whose sequences are different from one

another can also be constructed by combining heavy chain variable regions, produced as a randomized variable region sequence library, with light chain variable regions introduced with "at least one amino acid residue that changes the antigen-binding activity of an antigen-binding molecule depending on the hydrogen ion concentration condition".

5 Such amino acid residues include, but are not limited to, for example, amino acid residues contained in the light chain CDR1. The amino acid residues also include, but are not limited to, for example, amino acid residues contained in the light chain CDR2. The amino acid residues also include, but are not limited to, for example, amino acid residues contained in the light chain CDR3.

10 The above-described amino acid residues contained in the light chain CDR1 include, but are not limited to, for example, amino acid residues of positions 24, 27, 28, 31, 32, and/or 34 according to Kabat numbering in the CDR1 of light chain variable region. Meanwhile, the amino acid residues contained in the light chain CDR2 include, but are not limited to, for example, amino acid residues of positions 50, 51, 52, 53, 54, 55, and/or 56 according to Kabat
15 numbering in the CDR2 of light chain variable region. Furthermore, the amino acid residues in the light chain CDR3 include, but are not limited to, for example, amino acid residues of positions 89, 90, 91, 92, 93, 94, and/or 95A according to Kabat numbering in the CDR3 of light chain variable region. Moreover, the amino acid residues can be contained alone or can be contained in combination of two or more amino acids as long as they allow the change in the
20 antigen-binding activity of an antigen-binding molecule depending on the hydrogen ion concentration.

 Even when the heavy chain variable region produced as a randomized variable region sequence library is combined with the above-described light chain variable region introduced with "at least one amino acid residue that changes the antigen-binding activity of an
25 antigen-binding molecule depending on the hydrogen ion concentration condition", it is possible to design so that the flexible residues are contained in the sequence of the light chain variable region in the same manner as described above. The number and position of the flexible residues are not particularly limited to a specific embodiment, as long as the antigen-binding activity of an antigen-binding molecule of the present invention changes depending on the
30 hydrogen ion concentration condition. Specifically, the CDR and/or FR sequences of heavy chain and/or light chain can contain one or more flexible residues. For example, flexible residues to be introduced into the sequences of the light chain variable regions include, but are not limited to, for example, the amino acid residues listed in Tables 3 and 4. Meanwhile, amino acid sequences of light chain variable regions other than the flexible residues and amino acid
35 residues that change the antigen-binding activity of an antigen-binding molecule depending on the hydrogen ion concentration condition suitably include, but are not limited to, germ line

sequences such as Vk1 (SEQ ID NO: 5), Vk2 (SEQ ID NO: 6), Vk3 (SEQ ID NO: 7), and Vk4 (SEQ ID NO: 8).

[Table 3]

POSITION	AMINO ACID								
CDR1									
28	S:100%								
29	I:100%								
30	N:25%	S:25%	R:25%	H:25%					
31	S:100%								
32	H:100%								
33	L:100%								
34	A:50%	N:50%							
CDR2									
50	H:100%				OR	A:25%	D:25%	G:25%	K:25%
51	A:100%					A:100%			
52	S:100%					S:100%			
53	K:33.3%	N:33.3%	S:33.3%			H:100%			
54	L:100%					L:100%			
55	Q:100%					Q:100%			
56	S:100%					S:100%			
CDR3									
90	Q:100%				OR	Q:100%			
91	H:100%					S:33.3%	R:33.3%	Y:33.3%	
92	G:25%	N:25%	S:25%	Y:25%		H:100%			
93	H:33.3%	N:33.3%	S:33.3%			H:33.3%	N:33.3%	S:33.3%	
94	S:50%	Y:50%				S:50%	Y:50%		
95	P:100%					P:100%			
96	L:50%	Y:50%				L:50%	Y:50%		

5

(Position indicates Kabat numbering)

[Table 4]

CDR	POSITION	AMINO ACID				
CDR1	28	S:100%				
	29	I:100%				
	30	H:30%	N:10%	S:50%	R:10%	
	31	N:35%	S:65%			
	32	H:40%	N:20%	Y:40%		
	33	L:100%				
	34	A:70%	N:30%			
CDR2	50	A:25%	D:15%	G:25%	H:30%	K:5%
	51	A:100%				
	52	S:100%				
	53	H:30%	K:10%	N:15%	S:45%	
	54	L:100%				
	55	Q:100%				
	56	S:100%				
CDR3	90	Q:100%				
	91	H:30%	S:15%	R:10%	Y:45%	
	92	G:20%	H:30%	N:20%	S:15%	Y:15%
	93	H:30%	N:25%	S:45%		
	94	S:50%	Y:50%			
	95	P:100%				
	96	L:50%	Y:50%			

(Position indicates Kabat numbering)

Any amino acid residue may be suitably used as the above-described amino acid residues that change the antigen-binding activity of an antigen-binding molecule depending on the hydrogen ion concentration condition. Specifically, such amino acid residues include amino acids with a side chain pKa of 4.0-8.0. Such electron-releasing amino acids preferably include, for example, naturally occurring amino acids such as histidine and glutamic acid, as well as unnatural amino acids such as histidine analogs (US 20090035836), m-NO₂-Tyr (pKa 7.45), 3,5-Br₂-Tyr (pKa 7.21), and 3,5-I₂-Tyr (pKa 7.38) (Bioorg. Med. Chem. (2003) 11 (17), 3761-2768). Particularly preferred amino acid residues include, for example, amino acids with a side chain pKa of 6.0-7.0. Such electron-releasing amino acid residues preferably include, for example, histidine.

Known methods such as site-directed mutagenesis (Kunkel *et al.* (Proc. Natl. Acad. Sci. USA (1985) 82, 488-492)) and Overlap extension PCR can be appropriately employed to modify the amino acids of antigen-binding domains. Furthermore, various known methods can also be used as an amino acid modification method for substituting amino acids by those other than natural amino acids (Annu. Rev. Biophys. Biomol. Struct. (2006) 35, 225-249; Proc. Natl. Acad. Sci. U.S.A. (2003) 100 (11), 6353-6357). For example, a cell-free translation system (Clover Direct (Protein Express)) containing tRNAs in which amber suppressor tRNA, which is complementary to UAG codon (amber codon) that is a stop codon, is linked with an unnatural amino acid may be suitably used.

The preferred heavy chain variable region that is used in combination includes, for example, randomized variable region libraries. Known methods are appropriately combined as a method for producing a randomized variable region library. In a non-limiting embodiment of the present invention, an immune library constructed based on antibody genes derived from animals immunized with specific antigens, patients with infection or persons with an elevated antibody titer in blood as a result of vaccination, cancer patients, or lymphocytes of auto immune diseases may be suitably used as a randomized variable region library.

In another non-limiting embodiment of the present invention, in the same manner as described above, a synthetic library in which the CDR sequences of V genes from genomic DNA or functional reconstructed V genes are replaced with a set of synthetic oligonucleotides containing the sequences encoding codon sets of an appropriate length can also be suitably used as a randomized variable region library. In this case, the CDR3 sequence alone may be replaced because variety in the gene sequence of heavy chain CDR3 is observed. The basis for giving rise to amino acid variations in the variable region of an antigen-binding molecule is to generate variations of amino acid residues of surface-exposed positions of the antigen-binding molecule. The surface-exposed position refers to a position where an amino acid is exposed on the surface and/or contacted with an antigen based on the conformation, structural ensemble, and/or modeled structure of an antigen-binding molecule, and in general, such positions are the CDRs. The surface-exposed positions are preferably determined using the coordinates derived from a three-dimensional model of the antigen-binding molecule using computer programs such as InsightII program (Accelrys). The surface-exposed positions can be determined using algorithms known in the art (for example, Lee and Richards (J. Mol. Biol. (1971) 55, 379-400); Connolly (J. Appl. Cryst. (1983) 16, 548-558)). The surface-exposed positions can be determined based on the information on the three dimensional structure of antibodies using software suitable for protein modeling. Software which is suitably used for this purpose includes the SYBYL biopolymer module software (Tripos Associates). When the algorithm requires the input size parameter from the user, the "size" of probe for use in computation is

generally or preferably set at about 1.4 angstrom or less in radius. Furthermore, a method for determining surface-exposed region and area using personal computer software is described by Pacios (Comput. Chem. (1994) 18 (4), 377-386; and J. Mol. Model. (1995) 1, 46-53).

In still another non-limiting embodiment of the present invention, a naive library
5 constructed from antibody genes derived from lymphocytes of healthy persons and consisting of naive sequences, which are unbiased repertoire of antibody sequences, can also be particularly suitably used as a randomized variable region library (Gejima *et al.* (Human Antibodies (2002) 11, 121-129); and Cardoso *et al.* (Scand. J. Immunol. (2000) 51, 337-344)).

10 Neutralizing activity

A non-limiting embodiment of the present invention provides an antigen-binding molecule having human-FcRn-binding activity in an acidic pH range including an antigen-binding domain and an Fc γ receptor-binding domain, and having neutralizing activity against an antigen, wherein the antigen-binding domain has antigen-binding activity that changes
15 depending on the ion-concentration condition, and the Fc γ receptor-binding domain has higher binding activity to the Fc γ receptor in a neutral pH range condition than an Fc region of a native human IgG in which the sugar chain bonded at position 297 (EU numbering) is a fucose-containing sugar chain; and a pharmaceutical composition comprising the antigen-binding molecule. Generally, neutralizing activity refers to activity of inhibiting the
20 biological activity of a ligand, such as viruses and toxins, having biological activity on cells. Thus, substances having neutralizing activity refer to substances that bind to the ligand or the receptor to which the ligand binds, and inhibits the binding between the ligand and the receptor. Receptors blocked from binding with the ligand by the neutralizing activity will not be able to exhibit biological activity through this receptor. When the antigen-binding molecule is an
25 antibody, such an antibody having neutralizing activity is generally called a neutralizing antibody. Neutralizing activity of a test substance may be measured by comparing the biological activity in the presence of a ligand between when the test substance is present and absent.

For example, major possible ligands for the IL-6 receptor preferably include IL-6 as shown in SEQ ID NO: 15. The IL-6 receptor, which is an I-type membrane protein with its
30 amino terminus forming the extracellular domain, forms a hetero-tetramer with a gp130 receptor which has been induced to dimerize by IL-6 (Heinrich *et al.* (Biochem. J. (1998) 334, 297-314)). Formation of the heterotetramer activates Jak which is associated with the gp130 receptor. Jak undergoes autophosphorylation and phosphorylates the receptor. The phosphorylation site of the receptor and Jak serves as a binding site for SH2-carrying molecules belonging to the Stat
35 family such as Stat3; MAP kinase; PI3/Akt; and other SH2-carrying proteins and adapters. Next, Stat bound to the gp130 receptor is phosphorylated by Jak. The phosphorylated Stat

dimerizes and moves into the nucleus, and regulates the transcription of target genes. Jak or Stat can also be involved in signal cascades via receptors of other classes. Deregulated IL-6 signal cascades are observed in inflammation and pathological conditions of autoimmune diseases, and cancers such as prostate cancer and multiple myeloma. Stat3 which may act as an oncogene is constitutively activated in many cancers. In prostate cancer and multiple myeloma, there is a crosstalk between the signaling cascade via the IL-6 receptor and the signaling cascade via the epithelial growth factor receptor (EGFR) family members (Ishikawa *et al.* (J. Clin. Exp. Hematopathol. (2006) 46 (2), 55-66)).

Such intracellular signaling cascades are different for each cell type; therefore, appropriate target molecules can be determined for each target cell of interest, and are not limited to the above-mentioned factors. Neutralization activity can be evaluated by measuring the activation of *in vivo* signaling. Furthermore, the activation of *in vivo* signaling can be detected by using as an index the action of inducing the transcription of a target gene that exists downstream of the *in vivo* signaling cascade. Change in the transcription activity of the target gene can be detected by the principle of reporter assays. Specifically, a reporter gene such as green fluorescence protein (GFP) or luciferase is placed downstream of a promoter region or a transcription factor of the target gene, its reporter activity is measured, and thereby change in the transcription activity can be measured as the reporter activity. Commercially available kits for measuring the activation of *in vivo* signaling can be used appropriately (for example, Mercury Pathway Profiling Luciferase System (Clontech)).

Furthermore, for methods of measuring the activity of neutralizing receptors/ligands of the EGF receptor family and such, which normally act on signaling cascades that work toward promoting cell proliferation, the neutralization activity of neutralizing antibodies can be evaluated by measuring the proliferation activity of target cells. For example, when cells are promoted to proliferate by growth factors of the EGF family such as HB-EGF, the inhibitory effect on the proliferation of such cells based on the neutralizing activity of an anti-HB-EGF antibody can be suitably evaluated or measured by the following methods: For evaluating or measuring the cell proliferation inhibitory activity *in vitro*, a method of measuring the incorporation of [³H]-labeled thymidine added to the medium by viable cells as an index of DNA replication ability is used. As more convenient methods, a dye exclusion method, in which the ability of a cell to exclude a dye such as trypan blue from the cell is measured under the microscope, and the MTT method, are used. The latter method makes use of the ability of viable cells to convert MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide), which is a tetrazolium salt, to a blue formazan product. More specifically, a test antibody is added as well as a ligand to the culture solution of a test cell, and after a certain period of time, the MTT solution is added to the culture solution, and this is left to stand for a while for

incorporation of MTT into the cell. As a result, MTT, which is a yellow compound, is converted to a blue compound by the action of succinate dehydrogenase in the mitochondria of the cell. After dissolving this blue product for coloration, its absorbance is measured and used as an index for the number of viable cells. In addition to MTT, reagents such as MTS, XTT, 5 WST-1, and WST-8 are also commercially available (Nacalai Tesque, and such) and can be suitably used. For measuring the activity, a binding antibody which is of the same isotype as the anti-HB-EGF antibody but does not have the cell proliferation inhibitory activity can be used as a control antibody in the same manner as the anti-HB-EGF antibody, and the activity can be determined when the anti-HB-EGF antibody shows stronger cell proliferation inhibitory activity 10 than the control antibody.

Cells that can be preferably used for evaluating the activity include, for example, cells promoted to proliferate by HB-EGF such as the ovarian cancer cell line RMG-1, and mouse Ba/F3 cells which have been transformed by a vector for expressing a gene encoding hEGFR/mG-CSFR, which is a fusion protein in which the extracellular domain of human EGFR 15 is fused in frame with the intracellular domain of the mouse GCSF receptor. In this way, those skilled in the art can appropriately select cells to be used for evaluating the activity and use them to measure the cell proliferation activity as mentioned above.

Since the antigen-binding molecule provided by the present invention can eliminate antigens from plasma, the antigen-binding molecule itself does not necessarily have to have 20 neutralizing activity. However, it is more favorable to block the function of the antigen present in plasma by exerting neutralizing activity against the antigen until the antigen is taken up with the antigen-binding molecule into Fcγ-receptor-expressing cells by Fcγ-receptor-mediated endocytosis.

Furthermore, since the antigen-binding molecule provided by the present invention can 25 promote intracellular dissociation of an antigen, which has been extracellularly bound to the antigen-binding molecule, from an antigen-binding molecule, the antigen that dissociated from the antigen-binding molecule inside the cell is degraded in the lysosome. Therefore, the antigen-binding molecule itself does not necessarily have to have neutralizing activity. However, it is more favorable to block the function of the antigen present in plasma by exerting 30 neutralizing activity against the antigen until the antigen is taken up with the antigen-binding molecule into Fcγ-receptor-expressing cells by Fcγ-receptor-mediated endocytosis.

Furthermore, since the antigen-binding molecule provided by the present invention can decrease the total antigen concentration or free antigen concentration in plasma, the antigen-binding molecule itself does not necessarily have to have neutralizing activity. 35 However, it is more favorable to block the function of the antigen present in plasma by exerting neutralizing activity against the antigen until the antigen is taken up with the antigen-binding

molecule into Fcγ-receptor-expressing cells by Fcγ-receptor-mediated endocytosis.

Fcγ receptor

Fcγ receptor (FcγR) refers to a receptor capable of binding to the Fc region of monoclonal
 5 IgG1, IgG2, IgG3, or IgG4 antibodies, and includes all members belonging to the family of
 proteins substantially encoded by an Fcγ receptor gene. In humans, the family includes FcγRI
 (CD64) including isoforms FcγRIa, FcγRIb and FcγRIc; FcγRII (CD32) including isoforms
 FcγRIIa (including allotype H131 and R131), FcγRIIb (including FcγRIIb-1 and FcγRIIb-2), and
 FcγRIIc; and FcγRIII (CD16) including isoform FcγRIIIa (including allotype V158 and F158)
 10 and FcγRIIIb (including allotype FcγRIIIb-NA1 and FcγRIIIb-NA2); as well as all unidentified
 human FcγRs, FcγR isoforms, and allotypes thereof. However, Fcγ receptor is not limited to
 these examples. Without being limited thereto, FcγR includes those derived from humans, mice,
 rats, rabbits, and monkeys. FcγR may be derived from any organism. Mouse FcγR includes,
 without being limited to, FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16), and FcγRIII-2
 15 (FcγRIV, CD16-2), as well as all unidentified mouse FcγRs, FcγR isoforms, and allotypes thereof.
 Such preferred Fcγ receptors include, for example, human FcγRI (CD64), FcγRIIa (CD32),
 FcγRIIb (CD32), FcγRIIIa (CD16), and/or FcγRIIIb (CD16). The polynucleotide sequence and
 amino acid sequence of human FcγRI are shown in SEQ ID NOs: 16 (NM_000566.3) and 17
 (NP_000557.1), respectively; the polynucleotide sequence and amino acid sequence of human
 20 FcγRIIa (allotype H131) are shown in SEQ ID NOs: 18 (BC020823.1) and 19 (AAH20823.1)
 (allotype R131 is a sequence in which amino acid at position 166 of SEQ ID NO: 19 is
 substituted with Arg), respectively; the polynucleotide sequence and amino acid sequence of
 FcγIIB are shown in SEQ ID NOs: 20 (BC146678.1) and 21 (AAI46679.1), respectively; the
 polynucleotide sequence and amino acid sequence of FcγRIIIa are shown in SEQ ID NOs: 22
 25 (BC033678.1) and 23 (AAH33678.1), respectively; and the polynucleotide sequence and amino
 acid sequence of FcγRIIIb are shown in SEQ ID NOs: 24 (BC128562.1) and 25 (AAI28563.1),
 respectively (RefSeq accession number is shown in each parentheses). Whether an Fcγ receptor
 has binding activity to the Fc region of a monoclonal IgG1, IgG2, IgG3, or IgG4 antibody can be
 assessed by ALPHA screen (Amplified Luminescent Proximity Homogeneous Assay), surface
 30 plasmon resonance (SPR)-based BIACORE method, and others (Proc. Natl. Acad. Sci. USA
 (2006) 103(11), 4005-4010), in addition to the above-described FACS and ELISA formats.

Meanwhile, "Fc ligand" or "effector ligand" refers to a molecule and preferably a
 polypeptide that binds to an antibody Fc region, forming an Fc/Fc ligand complex. The
 molecule may be derived from any organism. The binding of an Fc ligand to Fc preferably
 35 induces one or more effector functions. Such Fc ligands include, but are not limited to, Fc
 receptors, FcγR, FcαR, FcεR, FcRn, C1q, and C3, mannan-binding lectin, mannose receptor,

Staphylococcus Protein A, *Staphylococcus* Protein G, and viral FcγRs. The Fc ligands also include Fc receptor homologs (FcRH) (Davis *et al.*, (2002) Immunological Reviews 190, 123-136) or FCRL (Annu Rev Immunol. 2007; 25: 525-60), which are a family of Fc receptors homologous to FcγR. The Fc ligands also include unidentified molecules that bind to Fc.

5 In FcγRI (CD64) including FcγRIa, FcγRIb, and FcγRIc, and FcγRIII (CD16) including isoforms FcγRIIIa (including allotypes V158 and F158) and FcγRIIIb (including allotypes FcγRIIIb-NA1 and FcγRIIIb-NA2), α chain that binds to the Fc portion of IgG is associated with common γ chain having ITAM responsible for transduction of intracellular activation signal. Meanwhile, the cytoplasmic domain of FcγRII (CD32) including isoforms FcγRIIa (including
10 allotypes H131 and R131) and FcγRIIc contains ITAM. These receptors are expressed on many immune cells such as macrophages, mast cells, and antigen-presenting cells. The activation signal transduced upon binding of these receptors to the Fc portion of IgG results in enhancement of the phagocytic activity and inflammatory cytokine production of macrophages, mast cell degranulation, and the enhanced function of antigen-presenting cells. Fcγ receptors
15 having the ability to transduce the activation signal as described above are also referred to as activating Fcγ receptors.

Meanwhile, the intracytoplasmic domain of FcγRIIb (including FcγRIIb-1 and FcγRIIb-2) contains ITIM responsible for transduction of inhibitory signals. The crosslinking between FcγRIIb and B cell receptor (BCR) on B cells suppresses the activation signal from
20 BCR, which results in suppression of antibody production via BCR. The crosslinking of FcγRIII and FcγRIIb on macrophages suppresses the phagocytic activity and inflammatory cytokine production. Fcγ receptors having the ability to transduce the inhibitory signal as described above are also referred to as inhibitory Fcγ receptors.

25 Binding activity to the Fcγ receptor

The binding activity of an FcγR-binding domain, which is included in an antigen-binding molecule of the present invention, to any of the human Fcγ receptors, FcγRI, FcγRIIa, FcγRIIb, FcγRIIIa, and/or FcγRIIIb, can be confirmed by the above-described FACS and ELISA format, as well as ALPHA Screen (Amplified Luminescent Proximity Homogeneous
30 Assay), a BIACORE method using the surface plasmon resonance (SPR) phenomena, and such (Proc. Natl. Acad. Sci. USA (2006) 103 (11), 4005-4010). The extracellular domain of a human Fcγ receptor may be used as the soluble antigen in these assays.

ALPHA screen is performed by the ALPHA technology based on the principle described below using two types of beads: donor and acceptor beads. A luminescent signal is detected
35 only when molecules linked to the donor beads interact biologically with molecules linked to the acceptor beads and when the two beads are located in close proximity. Excited by laser beam,

the photosensitizer in a donor bead converts oxygen around the bead into excited singlet oxygen. When the singlet oxygen diffuses around the donor beads and reaches the acceptor beads located in close proximity, a chemiluminescent reaction within the acceptor beads is induced. This reaction ultimately results in light emission. If molecules linked to the donor beads do not
5 interact with molecules linked to the acceptor beads, the singlet oxygen produced by donor beads do not reach the acceptor beads and chemiluminescent reaction does not occur.

For example, a biotin-labeled antigen-binding molecule comprising Fc region is immobilized to the donor beads and glutathione S-transferase (GST)-tagged Fc γ receptor is immobilized to the acceptor beads. In the absence of an antigen-binding molecule comprising a
10 competitive Fc region variant, Fc γ receptor interacts with a antigen-binding molecule comprising a native Fc region, inducing a signal of 520 to 620 nm as a result. The antigen-binding molecule having a non-tagged Fc region variant competes with the antigen-binding molecule comprising a native Fc region for the interaction with Fc γ receptor. The relative binding affinity can be determined by quantifying the reduction of fluorescence as a result of competition.
15 Methods for biotinylating the antigen-binding molecules such as antibodies using Sulfo-NHS-biotin or the like are known. Appropriate methods for adding the GST tag to an Fc γ receptor include methods that involve fusing polypeptides encoding Fc γ and GST in-frame, expressing the fused gene using cells introduced with a vector to which the gene is operably linked, and then purifying using a glutathione column. The induced signal can be preferably
20 analyzed, for example, by fitting to a one-site competition model based on nonlinear regression analysis using software such as GRAPHPAD PRISM (GraphPad; San Diego).

One of the substances for observing their interaction is immobilized as a ligand onto the gold thin layer of a sensor chip. When light is shed on the rear surface of the sensor chip so that total reflection occurs at the interface between the gold thin layer and glass, the intensity of
25 reflected light is partially reduced at a certain site (SPR signal). The other substance for observing their interaction is injected as an analyte onto the surface of the sensor chip. The mass of immobilized ligand molecule increases when the analyte binds to the ligand. This alters the refraction index of solvent on the surface of the sensor chip. The change in refraction index causes a positional shift of SPR signal (conversely, the dissociation shifts the signal back
30 to the original position). In the Biacore system, the amount of shift described above (*i.e.*, the change of mass on the sensor chip surface) is plotted on the vertical axis, and thus the change of mass over time is shown as measured data (sensorgram). Kinetic parameters (association rate constant (k_a) and dissociation rate constant (k_d)) are determined from the curve of sensorgram, and affinity (KD) is determined from the ratio between these two constants. Inhibition assay is
35 preferably used in the BIACORE methods. Examples of such inhibition assay are described in Proc. Natl. Acad. Sci. USA (2006) 103(11), 4005-4010.

Fcγ-receptor-binding domain

An Fcγ-receptor-binding domain having higher Fcγ-receptor-binding activity than an Fc region of a native human IgG in which the sugar chain bonded at position 297 (EU numbering) is a fucose-containing sugar chain, may be produced by altering the amino acid of the native human IgG Fc region. Furthermore, any structure of the antigen-binding domain described previously, which is characterized by being bound to an Fcγ receptor, may be used for the Fcγ-receptor-binding domain. In such a case, the Fcγ-receptor-binding domain may be produced without the need for introducing amino acid alterations, or affinity to the Fcγ receptor may be increased by introducing further alterations. Examples of such an Fcγ-receptor-binding domain include an Fab fragment antibody that binds to FcγRIIIa, which is described in Protein Eng Des Sel. 2009 Mar;22(3):175-88, Protein Eng Des Sel. 2008 Jan;21(1):1-10 and J Immunol. 2002 Jul 1;169(1):137-44, a camel-derived single domain antibody and a single-chain Fv antibody, and an FcγRI-binding cyclic peptide described in FASEB J. 2009 Feb;23(2):575-85. Whether or not the FcγR-binding activity of the Fcγ-receptor-binding domain is higher than that of the Fc region of a native human IgG in which the sugar chain bonded at position 297 (EU numbering) is a fucose-containing sugar chain may be determined appropriately using the method described in the above-mentioned section on binding activity.

In the present invention, a human IgG Fc region is a suitable example of a starting-material Fcγ-receptor-binding domain. In the present invention, "altering the amino acid" or "amino acid alteration" of the Fc region includes altering the amino acid sequence of the starting-material Fc region to a different amino acid sequence. As long as the modified variant of the starting-material Fc region can bind to a human Fcγ receptor in a neutral pH range, any Fc region may be used as the starting-material Fc region. Furthermore, an Fc region produced by further altering an already altered Fc region used as a starting Fc region may also be preferably used as the Fc region of the present invention. The "starting Fc region" can refer to the polypeptide itself, a composition comprising the starting Fc region, or an amino acid sequence encoding the starting Fc region. Starting Fc regions can comprise a known Fc region produced *via* recombination described briefly in section "Antibodies". The origin of starting Fc regions is not limited, and they may be obtained from human or any nonhuman organisms. Such organisms preferably include mice, rats, guinea pigs, hamsters, gerbils, cats, rabbits, dogs, goats, sheep, bovines, horses, camels and organisms selected from nonhuman primates. In another embodiment, starting Fcγ receptor binding domains can also be obtained from cynomolgus monkeys, marmosets, rhesus monkeys, chimpanzees, or humans. Starting Fc regions can be obtained preferably from human IgG1; however, they are not limited to any particular IgG class. This means that an Fc region of human IgG1, IgG2, IgG3, or IgG4 can be used appropriately as a

starting Fc region, and herein also means that an Fc region of an arbitrary IgG class or subclass derived from any organisms described above can be preferably used as a starting Fc region. Examples of naturally-occurring IgG variants or modified forms are described in published documents (Curr. Opin. Biotechnol. (2009) 20 (6): 685-91; Curr. Opin. Immunol. (2008) 20 (4),
5 460-470; Protein Eng. Des. Sel. (2010) 23 (4): 195-202; WO 2009/086320; WO 2008/092117; WO 2007/041635; and WO 2006/105338); however, they are not limited to the examples.

Examples of alterations include those with one or more mutations, for example, mutations by substitution of different amino acid residues for amino acids of starting Fc regions, by insertion of one or more amino acid residues into starting Fc regions, or by deletion of one or
10 more amino acids from starting Fc region. Preferably, the amino acid sequences of altered Fc regions comprise at least a part of the amino acid sequence of a non-native Fc region. Such variants necessarily have sequence identity or similarity less than 100% to their starting Fc region. In a preferred embodiment, the variants have amino acid sequence identity or similarity about 75% to less than 100%, more preferably about 80% to less than 100%, even more
15 preferably about 85% to less than 100%, still more preferably about 90% to less than 100%, and yet more preferably about 95% to less than 100% to the amino acid sequence of their starting Fc region. In a non-limiting embodiment of the present invention, at least one amino acid is different between a modified Fc region of the present invention and its starting Fc region. Amino acid difference between a modified Fc region of the present invention and its starting Fc
20 region can also be preferably specified based on amino acid differences at above-described particular amino acid positions according to EU numbering.

Known methods such as site-directed mutagenesis (Kunkel *et al.* (Proc. Natl. Acad. Sci. USA (1985) 82, 488-492)) and overlap extension PCR can be appropriately employed to modify the amino acids of Fc regions. Furthermore, various known methods can also be used as an
25 amino acid modification method for substituting amino acids by those other than natural amino acids (Annu. Rev. Biophys. Biomol. Struct. (2006) 35, 225-249; Proc. Natl. Acad. Sci. U.S.A. (2003) 100 (11), 6353-6357). For example, a cell-free translation system (Clover Direct (Protein Express)) containing tRNAs in which amber suppressor tRNA, which is complementary to UAG codon (amber codon) which is a stop codon, is linked with an unnatural amino acid may
30 be suitably used.

An Fc region having Fc γ receptor-binding activity in a neutral pH range that is contained in the antigen-binding molecules of the present invention may be obtained by any method, but specifically, an Fc region having Fc γ receptor-binding activity in the neutral pH range may be obtained by altering amino acids of human IgG immunoglobulin used as a starting
35 Fc region. Preferred IgG immunoglobulin Fc regions to be altered include, for example, the Fc regions of human IgG (IgG1, IgG2, IgG3, or IgG4, and their variants). IgG Fc regions include

mutants naturally formed therefrom. A number of allotype sequences due to genetic polymorphism are described in "Sequences of proteins of immunological interest", NIH Publication No.91-3242, for the Fc regions of human IgG1, human IgG2, human IgG3, and human IgG4 antibodies, and any one of them may be used in the present invention. In particular for the human IgG1 sequence, the amino acid sequence of positions 356 to 358 (EU numbering) may be either DEL or EFM.

Amino acids at any positions may be altered to other amino acids as long as the Fc region has Fc γ receptor-binding activity in a neutral pH range, or its Fc γ receptor-binding activity in a neutral range can be enhanced. When an antigen-binding molecule contains the Fc region of human IgG1, it is preferred to include alterations that result in enhancement of Fc γ receptor-binding in a neutral pH range compared to the binding activity of the starting Fc region of human IgG1. Amino acid alterations for enhancing Fc γ receptor-binding activity in a neutral pH range have been reported, for example, in WO 2007/024249, WO 2007/021841, WO 2006/031370, WO 2000/042072, WO 2004/029207, WO 2004/099249, WO 2006/105338, WO 2007/041635, WO 2008/092117, WO 2005/070963, WO 2006/020114, WO 2006/116260, and WO 2006/023403.

Examples of such amino acids that can be altered include at least one or more amino acids selected from the group consisting of those at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 according to EU numbering. Alteration of these amino acids enhances the Fc γ receptor-binding of an IgG immunoglobulin Fc region in a neutral pH range.

Particularly preferred alterations for use in the present invention include the following alterations:

- the amino acid at position 221 to either Lys or Tyr;
- the amino acid at position 222 to any one of Phe, Trp, Glu, and Tyr;
- the amino acid at position 223 to any one of Phe, Trp, Glu, and Lys;
- the amino acid at position 224 to any one of Phe, Trp, Glu, and Tyr;
- the amino acid at position 225 to any one of Glu, Lys, and Trp;
- the amino acid at position 227 to any one of Glu, Gly, Lys, and Tyr;
- the amino acid at position 228 to any one of Glu, Gly, Lys, and Tyr;
- the amino acid at position 230 to any one of Ala, Glu, Gly, and Tyr;

- the amino acid at position 231 to any one of Glu, Gly, Lys, Pro, and Tyr;
the amino acid at position 232 to any one of Glu, Gly, Lys, and Tyr;
the amino acid at position 233 to any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
- 5 the amino acid at position 234 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 235 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 236 to any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro,
- 10 Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 237 to any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 238 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
- 15 the amino acid at position 239 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr;
the amino acid at position 240 to any one of Ala, Ile, Met, and Thr;
the amino acid at position 241 to any one of Asp, Glu, Leu, Arg, Trp, and Tyr;
the amino acid at position 243 to any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr;
- 20 the amino acid at position 244 to His;
the amino acid at position 245 to Ala;
the amino acid at position 246 to any one of Asp, Glu, His, and Tyr;
the amino acid at position 247 to any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr;
the amino acid at position 249 to any one of Glu, His, Gln, and Tyr;
- 25 the amino acid at position 250 to either Glu or Gln;
the amino acid at position 251 to Phe;
the amino acid at position 254 to any one of Phe, Met, and Tyr;
the amino acid at position 255 to any one of Glu, Leu, and Tyr;
the amino acid at position 256 to any one of Ala, Met, and Pro;
- 30 the amino acid at position 258 to any one of Asp, Glu, His, Ser, and Tyr;
the amino acid at position 260 to any one of Asp, Glu, His, and Tyr;
the amino acid at position 262 to any one of Ala, Glu, Phe, Ile, and Thr;
the amino acid at position 263 to any one of Ala, Ile, Met, and Thr;
the amino acid at position 264 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro,
- 35 Gln, Arg, Ser, Thr, Trp, and Tyr;
the amino acid at position 265 to any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro,

- Gln, Arg, Ser, Thr, Val, Val, Trp, and Tyr;
the amino acid at position 266 to any one of Ala, Ile, Met, and Thr;
the amino acid at position 267 to any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr;
5 the amino acid at position 268 to any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp;
the amino acid at position 269 to any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 270 to any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser,
10 Thr, Trp, and Tyr;
the amino acid at position 271 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 272 to any one of Asp, Phe, Gly, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
15 the amino acid at position 273 to either Phe or Ile;
the amino acid at position 274 to any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 275 to either Leu or Trp;
the amino acid at position 276 to any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser,
20 Thr, Val, Trp, and Tyr;
the amino acid at position 278 to any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp;
the amino acid at position 279 to Ala;
the amino acid at position 280 to any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr;
25 the amino acid at position 281 to any one of Asp, Lys, Pro, and Tyr;
the amino acid at position 282 to any one of Glu, Gly, Lys, Pro, and Tyr;
the amino acid at position 283 to any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr;
the amino acid at position 284 to any one of Asp, Glu, Leu, Asn, Thr, and Tyr;
the amino acid at position 285 to any one of Asp, Glu, Lys, Gln, Trp, and Tyr;
30 the amino acid at position 286 to any one of Glu, Gly, Pro, and Tyr;
the amino acid at position 288 to any one of Asn, Asp, Glu, and Tyr;
the amino acid at position 290 to any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr;
the amino acid at position 291 to any one of Asp, Glu, Gly, His, Ile, Gln, and Thr;
the amino acid at position 292 to any one of Ala, Asp, Glu, Pro, Thr, and Tyr;
35 the amino acid at position 293 to any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;

- the amino acid at position 294 to any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
- the amino acid at position 295 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
- 5 the amino acid at position 296 to any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val;
- the amino acid at position 297 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
- the amino acid at position 298 to any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg,
- 10 Thr, Val, Trp, and Tyr;
- the amino acid at position 299 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr;
- the amino acid at position 300 to any one of Ala, Asp, Glu, Gly, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp;
- 15 the amino acid at position 301 to any one of Asp, Glu, His, and Tyr;
- the amino acid at position 302 to Ile;
- the amino acid at position 303 to any one of Asp, Gly, and Tyr;
- the amino acid at position 304 to any one of Asp, His, Leu, Asn, and Thr;
- the amino acid at position 305 to any one of Glu, Ile, Thr, and Tyr;
- 20 the amino acid at position 311 to any one of Ala, Asp, Asn, Thr, Val, and Tyr;
- the amino acid at position 313 to Phe;
- the amino acid at position 315 to Leu;
- the amino acid at position 317 to either Glu or Gln;
- the amino acid at position 318 to any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr;
- 25 the amino acid at position 320 to any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr;
- the amino acid at position 322 to any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr;
- the amino acid at position 323 to Ile;
- 30 the amino acid at position 324 to any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr;
- the amino acid at position 325 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
- the amino acid at position 326 to any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser,
- 35 Thr, Val, Trp, and Tyr;
- the amino acid at position 327 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn,

- Pro, Arg, Thr, Val, Trp, and Tyr;
the amino acid at position 328 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 329 to any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln,
5 Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 330 to any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 331 to any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr;
10 the amino acid at position 332 to any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr;
the amino acid at position 333 to any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr;
the amino acid at position 334 to any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr;
15 the amino acid at position 335 to any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr;
the amino acid at position 336 to any one of Glu, Lys, and Tyr;
the amino acid at position 337 to any one of Glu, His, and Asn;
the amino acid at position 339 to any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and
20 Thr;
the amino acid at position 376 to either Ala or Val;
the amino acid at position 377 to either Gly or Lys;
the amino acid at position 378 to Asp;
the amino acid at position 379 to Asn;
25 the amino acid at position 380 to any one of Ala, Asn, and Ser;
the amino acid at position 382 to either Ala or Ile;
the amino acid at position 385 to Glu;
the amino acid at position 392 to Thr;
the amino acid at position 396 to Leu;
30 the amino acid at position 421 to Lys;
the amino acid at position 427 to Asn;
the amino acid at position 428 to either Phe or Leu;
the amino acid at position 429 to Met;
the amino acid at position 434 to Trp;
35 the amino acid at position 436 to Ile; and
the amino acid at position 440 to any one of Gly, His, Ile, Leu, and Tyr,

according to EU numbering in the Fc region.

The number of amino acids that are altered is not particularly limited. An amino acid at one position only may be altered, or amino acids at two or more positions may be altered.

5 Examples of combinations of amino acid alterations at two or more positions include the combinations shown in Table 5 (Tables 5-1 to 5-3).

[Table 5-1]

COMBINATION OF AMINO ACIDS	COMBINATION OF AMINO ACIDS
K370E/P396L/D270E	S239Q/I332Q
Q419H/P396L/D270E	S267D/I332E
V240A/P396L/D270E	S267E/I332E
R255L/P396L/D270E	S267L/A327S
R255L/P396L/D270E	S267Q/A327S
R255L/P396L/D270E/R292G	S298A/I332E
R255L/P396L/D270E	S304T/I332E
R255L/P396L/D270E/Y300L	S324G/I332D
F243L/D270E/K392N/P396L	S324G/I332E
F243L/R255L/D270E/P396L	S324I/I332D
F243L/R292P/Y300L/V305I/P396L	S324I/I332E
F243L/R292P/Y300L/P396L	T260H/I332E
F243L/R292P/Y300L	T335D/I332E
F243L/R292P/P396L	V240I/V266I
F243L/R292P/V305I	V264I/I332E
F243L/R292P	D265F/N297E/I332E
S298A/E333A/K334A	D265Y/N297D/I332E
E380A/T307A	F243L/V262I/V264W
K326M/E333S	N297D/A330Y/I332E
K326A/E333A	N297D/T299E/I332E
S317A/K353A	N297D/T299F/I332E
A327D/I332E	N297D/T299H/I332E
A330L/I332E	N297D/T299I/I332E
A330Y/I332E	N297D/T299L/I332E
E258H/I332E	N297D/T299V/I332E
E272H/I332E	P230A/E233D/I332E
E272I/N276D	P244H/P245A/P247V
E272R/I332E	S239D/A330L/I332E
E283H/I332E	S239D/A330Y/I332E
E293R/I332E	S239D/H268E/A330Y
F241L/V262I	S239D/I332E/A327A
F241W/F243W	S239D/I332E/A330I

[Table 5-2]

F243L/V264I	S239D/N297D/I332E
H268D/A330Y	S239D/S298A/I332E
H268E/A330Y	S239D/V264I/I332E
K246H/I332E	S239E/N297D/I332E
L234D/I332E	S239E/V264I/I332E
L234E/I332E	S239N/A330L/I332E
L234G/I332E	S239N/A330Y/I332E
L234I/I332E	S239N/S298A/I332E
L234I/L235D	S239Q/V264I/I332E
L234Y/I332E	V264E/N297D/I332E
L235D/I332E	V264I/A330L/I332E
L235E/I332E	V264I/A330Y/I332E
L235I/I332E	V264I/S298A/I332E
L235S/I332E	Y296D/N297D/I332E
L328A/I332D	Y296E/N297D/I332E
L328D/I332D	Y296H/N297D/I332E
L328D/I332E	Y296N/N297D/I332E
L328E/I332D	Y296Q/N297D/I332E
L328E/I332E	Y296T/N297D/I332E
L328F/I332D	D265Y/N297D/T299L/I332E
L328F/I332E	F241E/F243Q/V262T/V264E
L328H/I332E	F241E/F243R/V262E/V264R
L328I/I332D	F241E/F243Y/V262T/V264R
L328I/I332E	F241L/F243L/V262I/V264I
L328M/I332D	F241R/F243Q/V262T/V264R
L328M/I332E	F241S/F243H/V262T/V264T
L328N/I332D	F241W/F243W/V262A/V264A
L328N/I332E	F241Y/F243Y/V262T/V264T
L328Q/I332D	I332E/A330Y/H268E/A327A
L328Q/I332E	N297D/I332E/S239D/A330L
L328T/I332D	N297D/S298A/A330Y/I332E
L328T/I332E	S239D/A330Y/I332E/K326E
L328V/I332D	S239D/A330Y/I332E/K326T
L328V/I332E	S239D/A330Y/I332E/L234I
L328Y/I332D	S239D/A330Y/I332E/L235D

[Table 5-3]

L328Y/I332E	S239D/A330Y/I332E/V240I
N297D/I332E	S239D/A330Y/I332E/V264T
N297E/I332E	S239D/A330Y/I332E/V266I
N297S/I332E	S239D/D265F/N297D/I332E
P227G/I332E	S239D/D265H/N297D/I332E
P230A/E233D	S239D/D265I/N297D/I332E
Q295E/I332E	S239D/D265L/N297D/I332E
R255Y/I332E	S239D/D265T/N297D/I332E
S239D/I332D	S239D/D265V/N297D/I332E
S239D/I332E	S239D/D265Y/N297D/I332E
S239D/I332N	S239D/I332E/A330Y/A327A
S239D/I332Q	S239D/I332E/H268E/A327A
S239E/D265G	S239D/I332E/H268E/A330Y
S239E/D265N	S239D/N297D/I332E/A330Y
S239E/D265Q	S239D/N297D/I332E/K326E
S239E/I332D	S239D/N297D/I332E/L235D
S239E/I332E	S239D/V264I/A330L/I332E
S239E/I332N	S239D/V264I/S298A/I332E
S239E/I332Q	S239E/V264I/A330Y/I332E
S239N/I332D	F241E/F243Q/V262T/V264E/I332E
S239N/I332E	F241E/F243R/V262E/V264R/I332E
S239N/I332N	F241E/F243Y/V262T/V264R/I332E
S239N/I332Q	F241R/F243Q/V262T/V264R/I332E
S239Q/I332D	S239D/I332E/H268E/A330Y/A327A
S239Q/I332E	S239E/V264I/S298A/A330Y/I332E
S239Q/I332N	F241Y/F243Y/V262T/V264T/N297D/I332E
S267E/L328F	G236D/S267E
S239D/S267E	

For the pH conditions to measure the binding activity of the Fcγ receptor-binding domain contained in the antigen-binding molecule of the present invention and the Fcγ receptor, conditions in an acidic pH range or in a neutral pH range may be suitably used. The neutral pH range, as a condition to measure the binding activity of the Fcγ receptor-binding domain contained in the antigen-binding molecule of the present invention and the Fcγ receptor, generally indicates pH 6.7 to pH 10.0. Preferably, it is a range indicated with arbitrary pH

values between pH 7.0 and pH8.0; and preferably, it is selected from pH 7.0, pH 7.1, pH 7.2, pH 7.3, pH 7.4, pH 7.5, pH 7.6, pH 7.7, pH 7.8, pH 7.9, and pH 8.0; and particularly preferably, it is pH 7.4, which is close to the pH of plasma (blood) *in vivo*. Herein, the acidic pH range, as a condition for having a binding activity of the Fcγ receptor-binding domain contained in the antigen-binding molecule of the present invention and the Fcγ receptor, generally indicates pH 4.0 to pH 6.5. Preferably, it indicates pH 5.5 to pH 6.5, and particularly preferably, it indicates pH 5.8 to pH 6.0, which is close to the pH in the early endosome *in vivo*. With regard to the temperature used as measurement condition, the binding affinity between the Fcγ receptor-binding domain and the human Fcγ receptor can be evaluated at any temperature between 10°C and 50°C. Preferably, a temperature between 15°C and 40°C is used to determine the binding affinity between the human Fcγ receptor-binding domain and the Fcγ receptor. More preferably, any temperature between 20°C and 35°C, such as any from 20°C, 21°C, 22°C, 23°C, 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 31°C, 32°C, 33°C, 34°C, or 35°C, can similarly be used to determine the binding affinity between the Fcγ receptor-binding domain and the Fcγ receptor. A temperature of 25°C is a non-limiting example in an embodiment of the present invention.

Herein, “the binding activity of the Fcγ receptor-binding domain toward Fcγ receptor is higher than the binding activity of the native Fc region toward activating Fcγ receptor” means that the binding activity of the Fcγ receptor-binding domain toward any of the human Fcγ receptors of FcγRI, FcγRIIa, FcγRIIb, FcγRIIIa, and/or FcγRIIIb is higher than the binding activity of the native Fcγ receptor-binding domain toward these human Fcγ receptors.

For example, it refers to the binding activity of the antigen-binding molecule including an Fcγ receptor-binding domain being 105% or more, preferably 110% or more, 115% or more, 120% or more, 125% or more, particularly preferably 130% or more, 135% or more, 140% or more, 145% or more, 150% or more, 155% or more, 160% or more, 165% or more, 170% or more, 175% or more, 180% or more, 185% or more, 190% or more, 195% or more, two-times or more, 2.5 times or more, three times or more, 3.5 times or more, four times or more, 4.5 times or more, five times or more, 7.5 times or more, ten times or more, 20 times or more, 30 times or more, 40 times or more, 50 times or more, 60 times or more, 70 times or more, 80 times or more, 90 times or more, or 100 times or more compared to the binding activity of the antigen-binding molecule including an Fc region of a native human IgG which is used as a control, based on the above-mentioned analysis method. For the native Fcγ receptor-binding domain, the starting-material Fcγ receptor-binding domain may be used, and the Fcγ receptor-binding domain of a native antibody belonging to the same subclass may also be used.

In the present invention, in particular, an Fc region of a native human IgG, in which the sugar chain bonded to an amino acid at position 297 (EU numbering) is a fucose-containing

sugar chain, is suitably used as the Fc region of a native human IgG to be used as a control. Whether or not the sugar chain bonded to an amino acid at position 297 (EU numbering) is a fucose-containing sugar chain can be determined using the technique described in Non-Patent Document 24. For example, a method such the following enables determination of whether the

5 sugar chain bonded to the native human IgG Fc region is a fucose-containing sugar chain. By reaction of *N*-glycosidase F (Roche diagnostics) with a native human IgG to be tested, a sugar chain is dissociated from the native human IgG to be tested (Weitzhandler *et al.* (J. Pharma. Sciences (1994) 83, 12, 1670-1675)). Next, a concentrated inspissated material of a reaction

10 solution from which the proteins have been removed by reaction with ethanol (Schenk *et al.* (J. Clin. Investigation (2001) 108 (11) 1687-1695)) is fluorescence labeled by 2-aminopyridine (Bigge *et al.* (Anal. Biochem. (1995) 230 (2) 229-238)). Fluorescence-labeled 2-AB-modified sugar chain produced by removal of reagent by solid-phase extraction using a cellulose cartridge, is analyzed by normal-phase chromatography. Observation of the detected chromatogram peak

15 enables determination of whether or not the sugar chain bonded to the native human IgG Fc region is a fucose-containing sugar chain. Non-limiting examples of such an Fc region of a native human IgG, in which the sugar chain bonded to an amino acid at position 297 (EU numbering) is a fucose-containing sugar chain, include Fc regions included in antibodies obtained by expressing in CHO cells, such as CHO-K1 (American Type Culture Collection, ATCC No. CRL-61) or DXB11 (American Type Culture Collection, ATCC No. CRL-11397),

20 genes encoding antibodies containing a native human IgG Fc region. Using the Fc γ -receptor-binding activities of Fc regions included in such antibodies as controls, the Fc γ -receptor-binding activities of Fc regions of the present invention were compared. This enables determination of whether the Fc γ -receptor-binding domain of the present invention has higher Fc γ -receptor-binding activity than the Fc region of a native human IgG in which the sugar

25 chain bonded at position 297 (EU numbering) is a fucose-containing sugar chain. Furthermore, the amount of fucose bonded to the sugar chain included in the compared Fc regions can be compared by determining the amount of fucose in the sugar chain bonded to the Fc regions included in these antibodies and the amount of fucose in the sugar chain bonded to the Fc regions of the present invention using methods such as those mentioned above.

30 As antigen-binding molecule containing an Fc region of the same subclass native antibody that is used as a control, antigen-binding molecules having an Fc region of a monoclonal IgG antibody may be suitably used. The structures of such Fc regions are shown in SEQ ID NO: 11 (A is added to the N terminus of RefSeq Accession No. AAC82527.1), SEQ ID NO: 12 (A is added to the N terminus of RefSeq Accession No. AAB59393.1), SEQ ID NO: 13

35 (RefSeq Accession No. CAA27268.1), and SEQ ID NO: 14 (A is added to the N terminus of RefSeq Accession No. AAB59394.1). Further, when an antigen-binding molecule containing

an Fc region of a particular antibody isotype is used as the test substance, the effect of the binding activity of the antigen-binding molecule containing a test Fc region toward the Fc γ receptor is tested by using as a control an antigen-binding molecule having an Fc region of a monoclonal IgG antibody of that particular isotype. In this way, antigen-binding molecules
 5 containing an Fc region whose binding activity toward the Fc γ receptor was demonstrated to be high are suitably selected.

Examples of Fc γ -receptor-binding domains suitable for use in the present invention include Fc γ -receptor-binding domains with the property of having higher binding activity to certain Fc γ receptors than to other Fc γ receptors (Fc γ -receptor-binding domains with selective
 10 Fc γ -receptor-binding activity). When an antibody is used as the antigen-binding molecule (when an Fc region is used as the Fc γ -receptor-binding domain), since a single antibody molecule can only bind to a single Fc γ receptor, a single antigen-binding molecule which is bound to an inhibitory Fc γ receptor cannot bind to another activating Fc γ R, and a single antigen-binding molecule which is bound to an activating Fc γ receptor cannot bind to another
 15 activating Fc γ receptor nor an inhibitory Fc γ receptor.

As described above, suitable examples of the activating Fc γ receptor are Fc γ RI (CD64) including Fc γ RIa, Fc γ RIb, and Fc γ RIc, and Fc γ RIII (CD16) including isoforms Fc γ RIIIa (including allotypes V158 and F158) and Fc γ RIIIb (including allotypes Fc γ RIIIb-NA1 and Fc γ RIIIb-NA2). Fc γ RIIb (including Fc γ RIIb-1 and Fc γ RIIb-2) is a suitable example of the
 20 inhibitory Fc γ receptor.

Fc γ R-binding domain having selective binding activity to an Fc γ receptor

Whether or not an Fc γ R-binding domain of the present invention has selective binding activity can be confirmed by comparing binding activities to the respective Fc γ receptors,
 25 determined by the method described in the above-mentioned section on binding activity to Fc γ receptors. An Fc γ R-binding domain with higher binding activity to inhibitory Fc γ receptors than to activating Fc γ receptors may be used as the selective Fc γ R-binding domain included in the antigen-binding molecule provided by the present invention. In a non-limiting embodiment, an Fc γ R-binding domain with higher binding activity to Fc γ RIIb (including Fc γ RIIb-1 and
 30 Fc γ RIIb-2) than to an activating Fc γ receptor selected from the group consisting of Fc γ RI(CD64) including Fc γ RIa, Fc γ RIb, Fc γ RIc, Fc γ RIII(CD16) including isoforms Fc γ RIIIa (including allotypes V158 and F158) and Fc γ RIIIb (including allotypes Fc γ RIIIb-NA1 and Fc γ RIIIb-NA2), and Fc γ RII(CD32) including isoforms Fc γ RIIa and Fc γ RIIc (including allotypes H131 and R131) may be used as a selective Fc γ R-binding domain included in an antigen-binding molecule
 35 provided by the present invention. Furthermore, in a non-limiting embodiment of the present invention, an Fc γ R-binding domain with higher binding activity to Fc γ RIIb-1 and/or Fc γ RIIb-2

than to FcγRIa, FcγRIb, and FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc may be used as a selective FcγR-binding domain included in an antigen-binding molecule provided by the present invention. Whether an FcγR-binding domain to be tested has selective binding activity to Fcγ receptors can be determined by comparing the value (ratio) obtained by dividing the KD values of the FcγR-binding domain for FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc by the KD values for FcγRIIb-1 and/or FcγRIIb-2, wherein the KD values are determined by the method described in the above-mentioned section on binding activity to Fcγ receptors, or more specifically, by comparing the FcγR selectivity indices shown in Equation 1.

[Equation 1]

FcγR selectivity index = KD value for activating FcγR / KD value for inhibitory FcγR

In Equation 1 mentioned above, “activating FcγR” refers to FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc, and inhibitory FcγR refers to FcγRIIb-1 and/or FcγRIIb-2. Although the activating FcγR and inhibitory FcγR used for the KD value measurements may be selected from any combination, in a non-limiting embodiment, a value (ratio) obtained by dividing the KD value for FcγRIIa including allotype H131 by the KD value for FcγRIIb-1 and/or FcγRIIb-2 may be used.

For example, the FcγR selectivity indices have values of, 1.2 or greater, 1.3 or greater, 1.4 or greater, 1.5 or greater, 1.6 or greater, 1.7 or greater, 1.8 or greater, 1.9 or greater, 2 or greater, 3 or greater, 5 or greater, 6 or greater, 7 or greater, 8 or greater, 9 or greater, 10 or greater, 15 or greater, 20 or greater, 25 or greater, 30 or greater, 35 or greater, 40 or greater, 45 or greater, 50 or greater, 55 or greater, 60 or greater, 65 or greater, 70 or greater, 75 or greater, 80 or greater, 85 or greater, 90 or greater, 95 or greater, 100 or greater, 110 or greater, 120 or greater, 130 or greater, 140 or greater, 150 or greater, 160 or greater, 170 or greater, 180 or greater, 190 or greater, 200 or greater, 210 or greater, 220 or greater, 230 or greater, 240 or greater, 250 or greater, 260 or greater, 270 or greater, 280 or greater, 290 or greater, 300 or greater, 310 or greater, 320 or greater, 330 or greater, 340 or greater, 350 or greater, 360 or greater, 370 or greater, 380 or greater, 390 or greater, 400 or greater, 410 or greater, 420 or greater, 430 or

greater, 440 or greater, 450 or greater, 460 or greater, 470 or greater, 480 or greater, 490 or greater, 500 or greater, 520 or greater, 540 or greater, 560 or greater, 580 or greater, 600 or greater, 620 or greater, 640 or greater, 660 or greater, 680 or greater, 700 or greater, 720 or greater, 740 or greater, 760 or greater, 780 or greater, 800 or greater, 820 or greater, 840 or greater, 860 or greater, 880 or greater, 900 or greater, 920 or greater, 940 or greater, 960 or greater, 980 or greater, 1000 or greater, 1500 or greater, 2000 or greater, 2500 or greater, 3000 or greater, 3500 or greater, 4000 or greater, 4500 or greater, 5000 or greater, 5500 or greater, 6000 or greater, 6500 or greater, 7000 or greater, 7500 or greater, 8000 or greater, 8500 or greater, 9000 or greater, 9500 or greater, 10000 or greater, or 100000 or greater.

10 A non-limiting embodiment of the selective FcγR-binding domain in an antigen-binding molecule of the present invention includes, for example, Fc regions produced by modifying the FcγR-binding domain included in an Fc region (an Fc region of the IgG class refers to the region from cysteine at position 226 (EU numbering) to the C terminus, or from proline at position 230 (EU numbering) to the C terminus, but is not limited thereto) constituting a portion of a constant region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17). An example of a method for producing the modified Fc regions includes the method described in the above-mentioned section on amino acid alterations. Examples of such altered Fc regions include an Fc region in which amino acid at position 238 (EU numbering) is Asp or an Fc region in which amino acid at position 328 (EU numbering) is Glu in a human IgG (IgG1, IgG2, IgG3, or IgG4). An Fc region in which amino acid at position 238 (EU numbering) is Asp or an Fc region in which amino acid at position 328 (EU numbering) is Glu in a human IgG (IgG1, IgG2, IgG3, or IgG4), and antigen-binding molecules containing such an Fc region show higher binding activity to FcγRIIb-1 and/or FcγRIIb-2 than to FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc.

Constant regions containing a selective FcγR-binding domain which are included in the antigen-binding molecules of the present invention and antigen-binding molecules containing such a constant region may also be Fc regions and antigen-binding molecules containing such an Fc region which maintains or shows reduced binding activity to activating FcγR (FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc) when compared to an Fc region (an Fc region of the IgG class refers to the region from cysteine at position 226 (EU numbering) to the C terminus, or from proline at position 230 (EU numbering) to the C terminus, but is not limited thereto) constituting a portion of human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3

(SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) (hereinafter referred to as a wild-type Fc region) and an antigen-binding molecule containing such a wild-type Fc region.

Compared to a wild-type Fc region and an antigen-binding molecule containing a wild-type Fc region, the degree of the aforementioned reduction in binding activity to activating FcγR of an Fc region containing a selective FcγR-binding domain included in an antigen-binding molecule of the present invention, and an antigen-binding molecule containing such an Fc region is, for example, 99% or less, 98% or less, 97% or less, 96% or less, 95% or less, 94% or less, 93% or less, 92% or less, 91% or less, 90% or less, 88% or less, 86% or less, 84% or less, 82% or less, 80% or less, 78% or less, 76% or less, 74% or less, 72% or less, 70% or less, 68% or less, 66% or less, 64% or less, 62% or less, 60% or less, 58% or less, 56% or less, 54% or less, 52% or less, 50% or less, 45% or less, 40% or less, 35% or less, 30% or less, 25% or less, 20% or less, 15% or less, 10% or less, 5% or less, 4% or less, 3% or less, 2% or less, 1% or less, 0.5% or less, 0.4% or less, 0.3% or less, 0.2% or less, 0.1% or less, 0.05% or less, 0.01% or less, or 0.005% or less.

The Fc regions containing a selective FcγR-binding domain and constant regions containing such an Fc region, and antigen-binding molecules containing such a constant region, which are included in the antigen-binding molecules of the present invention, may also be Fc regions and antigen-binding molecules containing such an Fc region which shows enhanced binding activity to inhibitory FcγR (FcγRIIb-1 and/or FcγRIIb-2) when compared to an Fc region (an Fc region of the IgG class refers to the region from cysteine at position 226 (EU numbering) to the C terminus, or from proline at position 230 (EU numbering) to the C terminus, but is not limited thereto) constituting a portion of a constant region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) (hereinafter referred to as a wild-type Fc region) and an antigen-binding molecule containing such a wild-type Fc region.

Compared to a wild-type Fc region and an antigen-binding molecule containing a wild-type Fc region, the degree of the aforementioned enhancement in binding activity to inhibitory FcγR of an Fc region containing a selective FcγR-binding domain included in an antigen-binding molecule of the present invention and an antigen-binding molecule containing such an Fc region is, for example, 101% or greater, 102% or greater, 103% or greater, 104% or greater, 105% or greater, 106% or greater, 107% or greater, 108% or greater, 109% or greater, 110% or greater, 112% or greater, 114% or greater, 116% or greater, 118% or greater, 120% or greater, 122% or greater, 124% or greater, 126% or greater, 128% or greater, 130% or greater, 132% or greater, 134% or greater, 136% or greater, 138% or greater, 140% or greater, 142% or greater, 144% or greater, 146% or greater, 148% or greater, 150% or greater, 155% or greater, 160% or greater, 165% or greater, 170% or greater, 175% or greater, 180% or greater, 185% or

greater, 190% or greater, 195% or greater, 2-fold or greater, 3-fold or greater, 4-fold or greater, 5-fold or greater, 6-fold or greater, 7-fold or greater, 8-fold or greater, 9-fold or greater, 10-fold or greater, 20-fold or greater, 30-fold or greater, 40-fold or greater, 50-fold or greater, 60-fold or greater, 70-fold or greater, 80-fold or greater, 90-fold or greater, 100-fold or greater, 200-fold or greater, 300-fold or greater, 400-fold or greater, 500-fold or greater, 600-fold or greater, 700-fold or greater, 800-fold or greater, 900-fold or greater, 1000-fold or greater, 10000-fold or greater, or 100000-fold or greater.

Furthermore, the Fc region containing a selective FcγR-binding domain included in an antigen-binding molecule of the present invention and the antigen-binding molecule containing such an Fc region may be an Fc region and an antigen-binding molecule containing such an Fc region which maintains or shows reduced binding activity to activating FcγR (FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc) when compared to an Fc region (an Fc region of the IgG class refers to, for example, the region from cysteine at position 226 (EU numbering) to the C terminus or from proline at position 230 (EU numbering) to the C terminus, but is not limited thereto) constituting a portion of a constant region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) (hereinafter referred to as a wild-type Fc region) and an antigen-binding molecule containing such a wild-type Fc region; and shows enhanced binding activity to inhibitory FcγR (FcγRIIb-1 and/or FcγRIIb-2) when compared to an Fc region (an Fc region of the IgG class refers to, for example, the region from cysteine at position 226 (EU numbering) to the C terminus or from proline at position 230 (EU numbering) to the C terminus, but is not limited thereto) constituting a portion of a constant region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) (hereinafter referred to as a wild-type Fc region) and an antigen-binding molecule containing such a wild-type Fc region.

Furthermore, the Fc region containing a selective FcγR-binding domain included in an antigen-binding molecule of the present invention and the antigen-binding molecule containing such an Fc region may be an Fc region and an antigen-binding molecule containing such an Fc region with higher degree of enhancement of binding activity to an inhibitory Fcγ receptor (FcγRIIb-1 and/or FcγRIIb-2) than to an activating Fcγ receptor (FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131), when compared to an Fc region (an Fc region of the IgG class refers to, for example, the region from cysteine at position 226 (EU numbering) to the C terminus or from proline at position 230 (EU numbering) to the C terminus, but is not limited

thereto) constituting a portion of a constant region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) (hereinafter referred to as a wild-type Fc region) and an antigen-binding molecule containing such a wild-type Fc region.

5 In the present invention, at least another alteration to the Fc region may be added to the Fc region in which amino acid at position 238 (EU numbering) is Asp and the Fc region in which amino acid at position 328 (EU numbering) is Glu, by the embodiments and such described in the aforementioned section on amino acid alterations. In addition to these alterations, additional alterations may also be added. The additional alterations can be selected from any of
10 substitutions, deletions, and modifications of an amino acid, and combinations thereof. For example, alterations that enhance binding activity to FcγRIIb while maintaining or reducing binding activity to FcγRIIa (H type) and FcγRIIa (R type) may be added. Addition of such alterations improves binding selectivity to FcγRIIb over FcγRIIa.

 Among these, alterations that improve binding selectivity to FcγRIIb over FcγRIIa (R
15 type) is favorable, and alterations that improve binding selectivity to FcγRIIb over FcγRIIa (H type) is more favorable. Examples of preferred amino acid substitutions for such alterations include: an alteration by substituting Gly at position 237 (EU numbering) with Trp; an alteration by substituting Gly at position 237 (EU numbering) with Phe; an alteration by substituting Pro at position 238 (EU numbering) with Phe; an alteration by substituting Asn at position 325 (EU
20 numbering) with Met; an alteration by substituting Ser at position 267 (EU numbering) with Ile; an alteration by substituting Leu at position 328 (EU numbering) with Asp; an alteration by substituting Ser at position 267 (EU numbering) with Val; an alteration by substituting Leu at position 328 (EU numbering) with Trp; an alteration by substituting Ser at position 267 (EU numbering) with Gln; an alteration by substituting Ser at position 267 (EU numbering) with Met;
25 an alteration by substituting Gly at position 236 (EU numbering) with Asp; an alteration by substituting Ala at position 327 (EU numbering) with Asn; an alteration by substituting Asn at position 325 (EU numbering) with Ser; an alteration by substituting Leu at position 235 (EU numbering) with Tyr; an alteration by substituting Val at position 266 (EU numbering) with Met; an alteration by substituting Leu at position 328 (EU numbering) with Tyr; an alteration by
30 substituting Leu at position 235 (EU numbering) with Trp; an alteration by substituting Leu at position 235 (EU numbering) with Phe; an alteration by substituting Ser at position 239 (EU numbering) with Gly; an alteration by substituting Ala at position 327 (EU numbering) with Glu; an alteration by substituting Ala at position 327 (EU numbering) with Gly; an alteration by substituting Pro at position 238 (EU numbering) with Leu; an alteration by substituting Ser at
35 position 239 (EU numbering) with Leu; an alteration by substituting Leu at position 328 (EU numbering) with Thr; an alteration by substituting Leu at position 328 (EU numbering) with Ser;

an alteration by substituting Leu at position 328 (EU numbering) with Met; an alteration by substituting Pro at position 331 (EU numbering) with Trp; an alteration by substituting Pro at position 331 (EU numbering) with Tyr; an alteration by substituting Pro at position 331 (EU numbering) with Phe; an alteration by substituting Ala at position 327 (EU numbering) with Asp;

5 an alteration by substituting Leu at position 328 (EU numbering) with Phe; an alteration by substituting Pro at position 271 (EU numbering) with Leu; an alteration by substituting Ser at position 267 (EU numbering) with Glu; an alteration by substituting Leu at position 328 (EU numbering) with Ala; an alteration by substituting Leu at position 328 (EU numbering) with Ile; an alteration by substituting Leu at position 328 (EU numbering) with Gln; an alteration by substituting Leu at position 328 (EU numbering) with Val; an alteration by substituting Lys at position 326 (EU numbering) with Trp; an alteration by substituting Lys at position 334 (EU numbering) with Arg; an alteration by substituting His at position 268 (EU numbering) with Gly; an alteration by substituting His at position 268 (EU numbering) with Asn; an alteration by substituting Ser at position 324 (EU numbering) with Val; an alteration by substituting Val at position 266 (EU numbering) with Leu; an alteration by substituting Pro at position 271 (EU numbering) with Gly; an alteration by substituting Ile at position 332 (EU numbering) with Phe; an alteration by substituting Ser at position 324 (EU numbering) with Ile; an alteration by substituting Glu at position 333 (EU numbering) with Pro; an alteration by substituting Tyr at position 300 (EU numbering) with Asp; an alteration by substituting Ser at position 337 (EU numbering) with Asp; an alteration by substituting Tyr at position 300 (EU numbering) with Gln; an alteration by substituting Thr at position 335 (EU numbering) with Asp; an alteration by substituting Ser at position 239 (EU numbering) with Asn; an alteration by substituting Lys at position 326 (EU numbering) with Leu; an alteration by substituting Lys at position 326 (EU numbering) with Ile; an alteration by substituting Ser at position 239 (EU numbering) with Glu;

25 an alteration by substituting Lys at position 326 (EU numbering) with Phe; an alteration by substituting Lys at position 326 (EU numbering) with Val; an alteration by substituting Lys at position 326 (EU numbering) with Tyr; an alteration by substituting Ser at position 267 (EU numbering) with Asp; an alteration by substituting Lys at position 326 (EU numbering) with Pro; an alteration by substituting Lys at position 326 (EU numbering) with His; an alteration by substituting Lys at position 334 (EU numbering) with Ala; an alteration by substituting Lys at position 334 (EU numbering) with Trp; an alteration by substituting His at position 268 (EU numbering) with Gln; an alteration by substituting Lys at position 326 (EU numbering) with Gln; an alteration by substituting Lys at position 326 (EU numbering) with Glu; an alteration by substituting Lys at position 326 (EU numbering) with Met; an alteration by substituting Val at position 266 (EU numbering) with Ile; an alteration by substituting Lys at position 334 (EU numbering) with Glu; an alteration by substituting Tyr at position 300 (EU numbering) with Glu;

35

an alteration by substituting Lys at position 334 (EU numbering) with Met; an alteration by substituting Lys at position 334 (EU numbering) with Val; an alteration by substituting Lys at position 334 (EU numbering) with Thr; an alteration by substituting Lys at position 334 (EU numbering) with Ser; an alteration by substituting Lys at position 334 (EU numbering) with His; 5 an alteration by substituting Lys at position 334 (EU numbering) with Phe; an alteration by substituting Lys at position 334 (EU numbering) with Gln; an alteration by substituting Lys at position 334 (EU numbering) with Pro; an alteration by substituting Lys at position 334 (EU numbering) with Tyr; an alteration by substituting Lys at position 334 (EU numbering) with Ile; an alteration by substituting Gln at position 295 (EU numbering) with Leu; an alteration by substituting Lys at position 334 (EU numbering) with Leu; an alteration by substituting Lys at position 334 (EU numbering) with Asn; an alteration by substituting His at position 268 (EU numbering) with Ala; an alteration by substituting Ser at position 239 (EU numbering) with Asp; an alteration by substituting Ser at position 267 (EU numbering) with Ala; an alteration by substituting Leu at position 234 (EU numbering) with Trp; an alteration by substituting Leu at position 234 (EU numbering) with Tyr; an alteration by substituting Gly at position 237 (EU numbering) with Ala; an alteration by substituting Gly at position 237 (EU numbering) with Asp; an alteration by substituting Gly at position 237 (EU numbering) with Glu; an alteration by substituting Gly at position 237 (EU numbering) with Leu; an alteration by substituting Gly at position 237 (EU numbering) with Met; an alteration by substituting Gly at position 237 (EU numbering) with Tyr; an alteration by substituting Ala at position 330 (EU numbering) with Lys; an alteration by substituting Ala at position 330 (EU numbering) with Arg; an alteration by substituting Glu at position 233 (EU numbering) with Asp; an alteration by substituting His at position 268 (EU numbering) with Asp; an alteration by substituting His at position 268 (EU numbering) with Glu; an alteration by substituting Lys at position 326 (EU numbering) with Asp; 25 an alteration by substituting Lys at position 326 (EU numbering) with Ser; an alteration by substituting Lys at position 326 (EU numbering) with Thr; an alteration by substituting Val at position 323 (EU numbering) with Ile; an alteration by substituting Val at position 323 (EU numbering) with Leu; an alteration by substituting Val at position 323 (EU numbering) with Met; an alteration by substituting Tyr at position 296 (EU numbering) with Asp; an alteration by substituting Lys at position 326 (EU numbering) with Ala; an alteration by substituting Lys at position 326 (EU numbering) with Asn; and an alteration by substituting Ala at position 330 (EU numbering) with Met. 30

Favorable amino acid substitutions among these alterations are, for example, an alteration by substituting Gly at position 237 (EU numbering) with Trp; an alteration by substituting Gly at position 237 (EU numbering) with Phe; an alteration by substituting Ser at position 267 (EU numbering) with Val; an alteration by substituting Ser at position 267 (EU 35

numbering) with Gln; an alteration by substituting His at position 268 (EU numbering) with Asn; an alteration by substituting Pro at position 271 (EU numbering) with Gly; an alteration by substituting Lys at position 326 (EU numbering) with Leu; an alteration by substituting Lys at position 326 (EU numbering) with Gln; an alteration by substituting Lys at position 326 (EU numbering) with Glu; an alteration by substituting Lys at position 326 (EU numbering) with Met; an alteration by substituting Ser at position 239 (EU numbering) with Asp; an alteration by substituting Ser at position 267 (EU numbering) with Ala; an alteration by substituting Leu at position 234 (EU numbering) with Trp; an alteration by substituting Leu at position 234 (EU numbering) with Tyr; an alteration by substituting Gly at position 237 (EU numbering) with Ala; an alteration by substituting Gly at position 237 (EU numbering) with Asp; an alteration by substituting Gly at position 237 (EU numbering) with Glu; an alteration by substituting Gly at position 237 (EU numbering) with Leu; an alteration by substituting Gly at position 237 (EU numbering) with Met; an alteration by substituting Gly at position 237 (EU numbering) with Tyr; an alteration by substituting Ala at position 330 (EU numbering) with Lys; an alteration by substituting Ala at position 330 (EU numbering) with Arg; an alteration by substituting Glu at position 233 (EU numbering) with Asp; an alteration by substituting His at position 268 (EU numbering) with Asp; an alteration by substituting His at position 268 (EU numbering) with Glu; an alteration by substituting Lys at position 326 (EU numbering) with Asp; an alteration by substituting Lys at position 326 (EU numbering) with Ser; an alteration by substituting Lys at position 326 (EU numbering) with Thr; an alteration by substituting Val at position 323 (EU numbering) with Ile; an alteration by substituting Val at position 323 (EU numbering) with Leu; an alteration by substituting Val at position 323 (EU numbering) with Met; an alteration by substituting Tyr at position 296 (EU numbering) with Asp; an alteration by substituting Lys at position 326 (EU numbering) with Ala; an alteration by substituting Lys at position 326 (EU numbering) with Asn; and an alteration by substituting Ala at position 330 (EU numbering) with Met.

The above-mentioned alteration may be at one position, or alterations at two or more positions may be combined. Favorable examples of such alterations are those described in Tables 14 to 15, Tables 17 to 24, and Tables 26 to 28.

Fc region produced by altering the FcγR-binding domain included in the Fc region presented as human IgG1 (SEQ ID NO: 14), IgG2 (SEQ ID NO: 15), IgG3 (SEQ ID NO: 16), or IgG4 (SEQ ID NO: 17) can be given as an example of another non-limiting embodiment of the selective FcγR-binding domain included in the antigen-binding molecules of the present invention. A method for producing the modified Fc regions is, for example, the method described in the above-mentioned section on amino acid alterations. Examples of such altered Fc regions include an Fc region in which amino acid at position 238 (EU numbering) is Asp and

amino acid at position at 271 (EU numbering) is Gly in a human IgG (IgG1, IgG2, IgG3, or IgG4). An Fc region in which amino acid at position 238 (EU numbering) is Asp and amino acid at position at 271 (EU numbering) is Gly in a human IgG (IgG1, IgG2, IgG3, or IgG4), and antigen-binding molecules containing such an Fc region show higher binding activity to

5 FcγRIIb-1 and/or FcγRIIb-2 than to FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131, and/or FcγRIIc.

In the present invention, at least another alteration to the Fc region may be added to the

10 Fc region in which amino acid at position 238 (EU numbering) is Asp and the amino acid at position 271 (EU numbering) is Gly, by the embodiments and such described in the aforementioned section on amino acid alterations. In addition to these alterations, additional alterations may also be added. The additional alterations can be selected from any of substitutions, deletions, and modifications of an amino acid, and combinations thereof. For

15 example, alterations that maintain or reduce binding activity to activating Fcγ receptors (FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131) can be added. Alterations that enhance binding activity to inhibitory Fcγ receptors (FcγRIIb-1 and/or FcγRIIb-2) while maintaining or

20 reducing binding activity to FcγRIIa (H type) and FcγRIIa (R type) may be added. Furthermore, alterations where the degree of enhancement of binding activity to inhibitory Fcγ receptors (FcγRIIb-1 and/or FcγRIIb-2) is higher than the degree of enhancement of binding activity to activating Fcγ receptors (FcγRIa, FcγRIb, FcγRIc, FcγRIIIa including allotype V158, FcγRIIIa including allotype F158, FcγRIIIb including allotype FcγRIIIb-NA1, FcγRIIIb including allotype

25 FcγRIIIb-NA2, FcγRIIa including allotype H131, FcγRIIa including allotype R131) may also be added. Addition of such alterations improves binding selectivity to FcγRIIb over FcγRIIa.

An example of a non-limiting embodiment of the altered Fc region comprising a selective FcγR-binding domain includes an altered Fc region in which any one or more of positions 233, 234, 237, 264, 265, 266, 267, 268, 269, 272, 296, 326, 327, 330, 331, 332, 333,

30 and 396 (EU numbering) are substituted in the Fc region in which amino acid at position 238 (EU numbering) is Asp and amino acid at position 271 (EU numbering) is Gly in a human IgG (IgG1, IgG2, IgG3, or IgG4).

In addition, an example of a non-limiting embodiment of the altered Fc region comprising a selective FcγR-binding domain is an altered Fc region comprising any one or more

35 of Asp at amino acid position 233,

- Tyr at amino acid position 234,
Asp at amino acid position 237,
Ile at amino acid position 264,
Glu at amino acid position 265,
5 any one of Phe, Met, and Leu at amino acid position 266,
any one of Ala, Glu, Gly, and Gln at amino acid position 267,
Asp or Glu at amino acid position 268,
Asp at amino acid position 269,
any one of Asp, Phe, Ile, Met, Asn, and Gln at amino acid position 272,
10 Asp at amino acid position 296,
Ala or Asp at amino acid position 326,
Gly at amino acid position 327,
Lys or Arg at amino acid position 330,
Ser at amino acid position 331,
15 Thr at amino acid position 332,
any one of Thr, Lys, and Arg at amino acid position 333,
any one of Asp, Glu, Phe, Ile, Lys, Leu, Met, Gln, Arg, and Tyr at amino acid position 396,
shown by EU numbering, in the Fc region in which amino acid at position 238 is Asp and amino
acid at position 271 (EU numbering) is Gly in a human IgG (IgG1, IgG2, IgG3, or IgG4).
20 Examples of a non-limiting embodiment of Fc region which further comprises at least
another alteration to the Fc region and further comprises additional alterations mentioned above
include Fc regions shown in Tables 6-1 to 6-7.
[Table 6-1]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP208	E233D/G237D/P238D/H268D/P271G/A330R
BP209	G237D/P238D/H268D/P271G/K326A/A330R
BP210	G237D/P238D/H268D/P271G/A330R
BP211	E233D/P238D/H268D/P271G/K326A/A330R
BP212	E233D/P238D/H268D/P271G/Y296D/A330R
BP213	E233D/P238D/H268D/P271G/A330R
BP214	E233D/L234Y/G237D/P238D/Y296D/K326D/A330K
BP215	G237D/P238D/H268D/P271G/Y296D/A330K
BP216	G237D/P238D/S267Q/H268D/P271G/A330K
BP217	G237D/P238D/S267Q/H268D/P271G/Y296D/A330K
BP218	G237D/P238D/H268D/P271G/K326D/A330K
BP219	L234Y/G237D/P238D/H268D/P271G/A330K
BP220	E233D/G237D/P238D/H268D/P271G/Y296D/A330K
BP221	L234Y/G237D/P238D/Y296D/K326A/A330R
BP222	L234Y/G237D/P238D/P271G/K326A/A330R
BP223	L234Y/G237D/P238D/H268D/P271G/K326A/A330R
BP224	L234Y/G237D/P238D/S267Q/H268D/P271G/K326A/A330R
BP225	L234Y/G237D/P238D/K326D/A330R
BP226	L234Y/G237D/P238D/P271G/K326D/A330R
BP227	L234Y/G237D/P238D/H268D/P271G/K326D/A330R
BP228	L234Y/G237D/P238D/S267Q/H268D/P271G/K326D/A330R
BP229	E233D/L234Y/G237D/P238D/P271G/K326A/A330R
BP230	E233D/G237D/P238D/H268D/P271G/Y296D/A330R
BP231	G237D/P238D/H268D/P271G/Y296D/A330R
BP232	L234Y/G237D/P238D/P271G/K326A/A330K
BP233	L234Y/G237D/P238D/P271G/A330K
BP234	E233D/L234Y/G237D/P238D/S267Q/H268D/P271G/Y296D/K326D/A330K
BP235	E233D/L234Y/G237D/P238D/H268D/P271G/Y296D/K326D/A330R
BP236	E233D/L234Y/G237D/P238D/S267Q/H268D/P271G/Y296D/K326D/A330R
BP237	E233D/L234Y/G237D/P238D/S267Q/H268D/P271G/Y296D/K326A/A330K

(Table 6-2 is a continuation table of Table 6-1.)

[Table 6-2]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP238	E233D/L234Y/G237D/P238D/H268D/P271G/Y296D/K326A/A330R
BP239	E233D/L234Y/G237D/P238D/S267Q/H268D/P271G/Y296D/K326A/A330R
BP240	E233D/G237D/P238D/S267Q/H268D/P271G/A330R
BP241	E233D/G237D/P238D/H268D/P271G/K326D/A330R
BP242	E233D/G237D/P238D/H268D/P271G/K326A/A330R
BP243	E233D/L234Y/G237D/P238D/H268D/P271G/A330R
BP244	E233D/G237D/P238D/S267Q/H268D/P271G/Y296D/A330R
BP245	E233D/G237D/P238D/S267Q/H268D/P271G/Y296D/K326D/A330R
BP246	E233D/G237D/P238D/S267Q/H268D/P271G/Y296D/K326A/A330R
BP247	E233D/G237D/P238D/H268D/P271G/Y296D/K326D/A330R
BP248	E233D/G237D/P238D/H268D/P271G/Y296D/K326A/A330R
BP249	E233D/L234Y/G237D/P238D/H268D/P271G/Y296D/A330R
BP262	G237D/P238D/H268E/P271G
BP264	E233D/G237D/P238D/H268E/P271G/Y296D/A330R
BP265	G237D/P238D/H268E/P271G/Y296D/A330R
BP266	E233D/G237D/P238D/H268E/P271G/A330R
BP267	E233D/G237D/P238D/H268E/P271G
BP268	E233D/G237D/P238D/H268E/P271G/Y296D
BP269	G237D/P238D/H268E/P271G/Y296D
BP300	E233D/G237D/P238D/V264I/H268E/P271G
BP313	E233D/G237D/P238D/D265E/H268E/P271G
BP333	E233D/G237D/P238D/V266F/H268E/P271G
BP338	E233D/G237D/P238D/V266L/H268E/P271G
BP339	E233D/G237D/P238D/V266M/H268E/P271G
BP348	E233D/G237D/P238D/S267A/H268E/P271G
BP350	E233D/G237D/P238D/S267E/H268E/P271G
BP352	E233D/G237D/P238D/S267G/H268E/P271G
BP367	E233D/G237D/P238D/H268E/E269D/P271G
BP384	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/K334R
BP390	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/I332S
BP391	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/I332T

(Table 6-3 is a continuation table of Table 6-2.)

[Table 6-3]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP392	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/I332K
BP393	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/I332R
BP423	E233D/G237D/P238D/S267A/H268E/P271G/A330R
BP425	E233D/G237D/P238D/V266L/S267A/H268E/P271G/A330R
BP426	E233D/G237D/P238D/S267A/H268E/E269D/P271G/A330R
BP427	E233D/G237D/P238D/S267A/H268E/E269Y/P271G/A330R
BP428	E233D/G237D/P238D/S267G/H268E/P271G/A330R
BP429	E233D/G237D/P238D/V264I/S267G/H268E/P271G/A330R
BP430	E233D/G237D/P238D/V266L/S267G/H268E/P271G/A330R
BP431	E233D/G237D/P238D/S267G/H268E/E269D/P271G/A330R
BP432	E233D/G237D/P238D/S267G/H268E/E269Y/P271G/A330R
BP433	E233D/G237D/P238D/H268D/P271G/Y296D/A330K/I332T
BP434	E233D/G237D/P238D/H268D/P271G/Y296D/K326D/A330R/I332T
BP435	E233D/G237D/P238D/H268D/P271G/Y296D/K326A/A330R/I332T
BP436	E233D/G237D/P238D/S267A/H268E/P271G/Y296D/A330R/I332T
BP437	G237D/P238D/S267A/H268E/P271G/Y296D/A330R/I332T
BP438	E233D/G237D/P238D/S267A/H268E/P271G/A330R/I332T
BP439	E233D/G237D/P238D/V264I/V266L/S267A/H268E/P271G/A330R
BP440	E233D/G237D/P238D/V264I/H268E/P271G/A330R
BP441	E233D/G237D/P238D/V266L/H268E/P271G/A330R
BP442	E233D/G237D/P238D/H268E/E269D/P271G/A330R
BP443	E233D/G237D/P238D/V266L/H268E/E269D/P271G/A330R
BP444	E233D/G237D/P238D/H268E/E269N/P271G/A330R
BP445	E233D/G237D/P238D/V264I/S267A/H268E/P271G/A330R
BP446	E233D/G237D/P238D/S267A/H268E/E269N/P271G/A330R
BP447	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396A
BP448	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396D
BP449	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396E
BP450	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396F
BP451	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396G
BP452	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396H

(Table 6-4 is a continuation table of Table 6-3.)

[Table 6-4]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP453	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396I
BP454	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396K
BP455	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396L
BP456	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396M
BP457	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396N
BP458	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396Q
BP459	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396R
BP460	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396S
BP461	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396T
BP462	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396V
BP463	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396W
BP464	E233D/G237D/P238D/S267A/H268E/P271G/A330R/P396Y
BP465	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E333K
BP466	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E333R
BP467	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E334S
BP468	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E334T
BP469	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E333S
BP470	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/E333T
BP471	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/P331S
BP472	E233D/G237D/P238D/H268D/P271G/Y296D/A330S
BP473	E233D/G237D/P238D/H268D/P271G/Y296D/A327G/A330R
BP474	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/P331S
BP475	E233D/G237D/P238D/H268D/P271G/Y296D/A327G/A330S
BP476	E233D/G237D/P238D/H268D/P271G/Y296D/A327G/A330S/P331S
BP477	E233D/G237D/P238D/H268D/P271G/Y296D/A327G/A330R/P331S
BP478	E233D/G237D/P238D/H268D/P271G/Y296D/A330R + S131C/K133R/G137E/G138S/Q196K/I199T/N203D/K214R/P217S + 219-221 DELETION + K222Y/T223G/H224P/T225P
BP479	E233D/G237D/P238D/V264I/V266L/S267A/H268E/P271G
BP480	E233D/G237D/P238D/V266L/H268E/E269D/P271G
BP481	E233D/G237D/P238D/V264I/S267A/H268E/P271G

(Table 6-5 is a continuation table of Table 6-4.)

[Table 6-5]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP482	E233D/G237D/P238D/S267A/H268E/E269N/P271G
BP483	E233D/G237D/P238D/V266L/S267A/H268E/P271G
BP484	E233D/G237D/P238D/S267A/H268E/E269D/P271G
BP485	E233D/G237D/P238D/S267A/H268E/E269Y/P271G
BP487	E233D/G237D/P238D/V264I/S267A/H268E/P271G/A330R/P396M
BP488	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A330R
BP489	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A330R/P396M
BP490	G237D/P238D/V264I/S267A/H268E/P271G/A330R
BP491	G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A330R
BP492	P238D/V264I/S267A/H268E/P271G
BP493	P238D/V264I/S267A/H268E/P271G/Y296D
BP494	G237D/P238D/S267A/H268E/P271G/Y296D/A330R
BP495	G237D/P238D/S267G/H268E/P271G/Y296D/A330R
BP496	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D
BP497	E233D/G237D/P238D/V264I/S267A/H268E/P271G/A327G/A330R
BP498	E233D/G237D/P238D/V264I/S267A/H268E/P271G/A330R/P396L
BP499	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A330R/P396L
BP500	G237D/P238D/V264I/S267A/H268E/P271G/Y296D
BP501	G237D/P238D/V264I/S267A/H268E/P271G
BP502	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A327G/A330R
BP503	E233D/G237D/P238D/V264I/S267A/H268E/P271G/Y296D/A327G/A330R/P396M
BP504	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272P
BP505	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D
BP506	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272P/Y296D/A330R
BP507	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272P/A330R
BP508	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272P/Y296D
BP509	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D/Y296D
BP510	G237D/P238D/V264I/S267A/H268E/P271G/E272P/A330R
BP511	G237D/P238D/V264I/S267A/H268E/P271G/E272P/Y296D/A330R
BP513	E233D/G237D/P238D/H268E/E272D/P271G

(Table 6-6 is a continuation table of Table 6-5.)

[Table 6-6]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP514	E233D/G237D/P238D/H268E/E272F/P271G
BP517	E233D/G237D/P238D/H268E/E272I/P271G
BP521	E233D/G237D/P238D/H268E/E272N/P271G
BP521	E233D/G237D/P238D/H268E/E272Q/P271G
BP531	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R/P396M
BP532	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A330R/P396M
BP533	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R/P396L
BP534	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A330R/P396L
BP535	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A327G/A330R/P396M
BP536	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A327G/A330R/P396M
BP537	G237D/P238D/V264I/S267G/H268E/P271G/A330R
BP538	G237D/P238D/V264I/H268E/P271G/A330R
BP539	G237D/P238D/V264I/S267G/H268E/P271G/E272P/Y296D/A330R
BP540	G237D/P238D/V264I/H268E/P271G/E272P/Y296D/A330R
BP541	E233D/G237D/P238D/H268D/P271G/K274Q/Y296D/A330R
BP542	E233D/G237D/P238D/H268D/P271G/Y296F/A330R
BP543	E233D/G237D/P238D/H268Q/P271G/Y296D/A330R
BP544	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/R355Q
BP545	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/D356E
BP546	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/L358M
BP547	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/K409R
BP548	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/Q419E
BP549	G237D/P238D/S267G/H268E/P271G/A330R
BP550	G237D/P238D/V264I/S267G/H268E/P271G/E272D/Y296D/A330R
BP551	G237D/P238D/V264I/H268E/P271G/E272D/Y296D/A330R
BP552	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D/Y296D/A330R
BP553	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D/A330R
BP554	G237D/P238D/V264I/S267A/H268E/P271G/E272D/A330R
BP555	G237D/P238D/V264I/S267A/H268E/P271G/E272D/Y296D/A330R
BP556	G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R
ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP514	E233D/G237D/P238D/H268E/E272F/P271G
BP517	E233D/G237D/P238D/H268E/E272I/P271G
BP521	E233D/G237D/P238D/H268E/E272N/P271G
BP521	E233D/G237D/P238D/H268E/E272Q/P271G
BP531	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R/P396M
BP532	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A330R/P396M
BP533	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R/P396L
BP534	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A330R/P396L
BP535	E233D/G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A327G/A330R/P396M
BP536	E233D/G237D/P238D/V264I/H268E/P271G/Y296D/A327G/A330R/P396M
BP537	G237D/P238D/V264I/S267G/H268E/P271G/A330R
BP538	G237D/P238D/V264I/H268E/P271G/A330R
BP539	G237D/P238D/V264I/S267G/H268E/P271G/E272P/Y296D/A330R
BP540	G237D/P238D/V264I/H268E/P271G/E272P/Y296D/A330R
BP541	E233D/G237D/P238D/H268D/P271G/K274Q/Y296D/A330R
BP542	E233D/G237D/P238D/H268D/P271G/Y296F/A330R
BP543	E233D/G237D/P238D/H268Q/P271G/Y296D/A330R
BP544	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/R355Q
BP545	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/D356E
BP546	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/L358M
BP547	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/K409R
BP548	E233D/G237D/P238D/H268D/P271G/Y296D/A330R/Q419E
BP549	G237D/P238D/S267G/H268E/P271G/A330R
BP550	G237D/P238D/V264I/S267G/H268E/P271G/E272D/Y296D/A330R
BP551	G237D/P238D/V264I/H268E/P271G/E272D/Y296D/A330R
BP552	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D/Y296D/A330R
BP553	E233D/G237D/P238D/V264I/S267A/H268E/P271G/E272D/A330R
BP554	G237D/P238D/V264I/S267A/H268E/P271G/E272D/A330R
BP555	G237D/P238D/V264I/S267A/H268E/P271G/E272D/Y296D/A330R
BP556	G237D/P238D/V264I/S267G/H268E/P271G/Y296D/A330R

(Table 6-7 is a continuation table of Table 6-6.)

[Table 6-7]

ALTERED Fc REGION	ALTERED AMINO ACID (EU NUMBERING)
BP558	G237D/P238D/V264I/S267G/H268E/P271G/E272D/A330R
BP559	P238D/V264I/S267A/H268E/P271G/E272D/Y296D
BP560	P238D/S267G/H268E/P271G/Y296D/A330R
BP561	E233D/G237D/P238D/H268D/P271G/E272D/Y296D/A330R
BP562	G237D/P238D/H268D/P271G/E272D/Y296D/A330R
BP563	E233D/G237D/P238D/H268E/P271G/E272D/Y296D/A330R
BP564	G237D/P238D/H268E/P271G/E272D/Y296D/A330R
BP565	E233D/G237D/P238D/S267A/H268E/P271G/Y296D/A330R

- Four types of FcγRs, FcγRI, FcγRIIb, FcγRIII, and FcγRIV, have been identified in mice. In humans as well, as corresponding FcγRs, FcγRI, FcγRIIa, FcγRIIb, FcγRIIIa, FcγRIIIa, and FcγRIIIb have been identified. FcγRIIb which is considered to be the only inhibitory type among these FcγRs is conserved in both humans and mice. The other FcγRs, except for FcγRIIIb, transmit activation signals *via* the immunoreceptor tyrosine-based activating motif (ITAM), whereas FcγRIIb transmits inhibitory signals *via* the immunoreceptor tyrosine-based inhibitory motif (ITIM) present inside the cells (Nat. Rev. Immunol. (2008) 8, 34-47).
- FcγRIIb1 and FcγRIIb2 have been reported as splicing variants of FcγRIIb. In both humans and mice, FcγRIIb1 has a longer intracellular domain than FcγRIIb2. FcγRIIb1 has been confirmed to be expressed in B cells, and FcγRIIb2 has been confirmed to be expressed in macrophages, mast cells, dendritic cells, basophils, neutrophils, and eosinophils (J. Clin. Immunol. (2005) 25 (1), 1-18).
- So far, in humans, dysfunction and decreased expression of FcγRIIb have been reported to be correlated with onset of autoimmune diseases. For example, it has been reported that in some SLE patients, binding of transcriptional activators is attenuated due to polymorphism in an expression promoter region of FcγRIIb, which results in the decreased FcγRIIb expression (Hum. Genet. (2005) 117, 220-227; J. Immunol. (2004) 172, 7192-7199; and J. Immunol. (2004) 172, 7186-7191). Furthermore, among SLE patients, two types of allotypes have been reported, where the amino acid at position 233 is Ile or Thr in FcγRIIb. This position exists in the transmembrane region of FcγRIIb, and it is reported that FcγRIIb is less likely to exist in the lipid raft when the amino acid at position 233 is Thr compared to when this amino acid is Ile, and as a result, signal transduction function of FcγRIIb decreases (Nat. Med. (2005) 11, 1056-1058; Hum. Mol. Genet., (2005) 14, 2881-2892). In mice as well, knockout mice produced by disrupting the FcγRIIb gene in C57BL/6 mice has been reported to present SLE-like symptoms such as autoantibody production and glomerulonephritis (Immunity 13 (2000) 277-285; J. Exp. Med. (2002) 195, 1167-1174). Furthermore, so far, reduced expression level of FcγRIIb and such have been reported in mice considered to be models with natural onset of SLE (Immunogenetics (2000) 51, 429-435; Int. Immunol. (1999) 11, 1685-1691; Curr. Biol. (2000) 10,

227-230; J. Immunol. (2002) 169, 4340-4346). From these reports, FcγRIIb is considered to regulate humoral immunity in mice as in humans.

When an antibody carrying an Fc of the present invention eliminates antigens *via* FcγRIIb, the endocytosis function of FcγRIIb is considered to be making the most important contribution among the functions of FcγRIIb. As described above, FcγRIIb1 and FcγRIIb2 exist as splicing variants of FcγRIIb, but it is reported that the latter is mainly involved in the endocytosis of an immunocomplex of an antibody and antigen (J. Immunol. (1994), 152 574-585; Science (1992) 256, 1808-1812; Cell (1989) 58, 317-327). So far, mouse FcγRIIb2 has been reported to be incorporated into a clathrin-coated pit and endocytosed (Cell (1989) 58, 10 317-327). Furthermore, it has been reported that a dileucine motif is necessary for FcγRIIb2-mediated endocytosis, and the dileucine motif is conserved in both humans and mice (EMBO J. (1994) 13 (13), 2963-2969). From these, FcγRIIb2 may have an endocytotic ability in humans as in mice.

On the other hand, unlike FcγRIIb2, it has been reported that FcγRIIb1 does not cause endocytosis. FcγRIIb1 has an inserted sequence in its intracellular domain that is not found in FcγRIIb2. It is considered that this sequence inhibits the uptake of FcγRIIb1 into a clathrin-coated pit, and as a result endocytosis is inhibited (J. Cell. Biol. (1992) 116, 875-888; J. Cell. Biol. (1989) 109, 3291-3302). In humans as well, FcγRIIb1 has an insertion sequence at a site similar to that of FcγRIIb2 as in mice; therefore, difference in the endocytotic ability 20 between FcγRIIb1 and FcγRIIb2 is presumed to be caused by a similar mechanism. Furthermore, in both humans and mice, approximately 40% of immunocomplexes on the cell surface is reported to be taken up into the cell in 20 minutes (Mol. Immunol. (2011) 49, 329-337; Science (1992) 256, 1808-1812). Therefore, in humans as well, FcγRIIb2 is presumed to uptake immunocomplexes into cells at rates similar to those in mice.

Since FcγRIIb is the only one that has ITIM inside the cell in both humans and mice among the FcγR family and the distribution of expressing cells are the same, it is presumed that its function in immune control is similar. Furthermore, considering the fact that immunocomplexes are taken up into cells at similar rates in humans and mice, antigen elimination effects of antibodies mediated by FcγRIIb in humans may be predictable using mice. 30 Antigen-binding molecules mF44 and mF46 have properties of binding to soluble antigens in a pH-dependent manner, and have enhanced affinity to mouse FcγRIIb and FcγRIII compared to mIgG1 which is an antigen-binding molecule having the property of binding to a soluble antigen in a pH-dependent manner. Indeed, it is shown in Example 5 that antigen clearance increased when mF44 or mF46 was administered to normal mice compared to when mIgG1 was 35 administered.

Furthermore, in the later-described Example 6, a similar experiment was carried out

using Fc receptor γ chain-deficient mice. It has been reported that Fc γ Rs other than Fc γ RIIb are expressed only in the co-presence of a gamma chain in mice. Thus, only Fc γ RIIb is expressed in the Fc receptor γ chain-deficient mice. Administration of mF44 or mF46, which are antigen-binding molecules having the property of binding to soluble antigens in a pH-dependent manner, to Fc receptor γ chain-deficient mice enables assessment of antigen elimination-acceleration effects when Fc γ RIIb-binding is selectively enhanced. From the results of Example 6, when mF44 or mF46 (which are antigen-binding molecules having the property of binding to soluble antigens in a pH-dependent manner) was administered to Fc receptor γ chain-deficient mice, antigen clearance was shown to increase compared to when mIgG1 (which is an antigen-binding molecule having the property of binding to soluble antigens in a pH-dependent manner) was administered to the mice. Furthermore, the results of Example 6 shows that when administered to Fc receptor γ chain-deficient mice, mF44 or mF46 cause similar degrees of antigen elimination as when administered to normal mice.

In Example 6, a similar experiment was performed using Fc γ RIII-deficient mice. Since mIgG1, mF44, and mF46 bind only to Fc γ RIIb and Fc γ RIII among the mFc γ Rs, administration of the antibodies to Fc γ RIII-deficient mice enables assessment of antigen elimination-accelerating effects when Fc γ RIIb-binding is selectively enhanced. The results of Example 6 indicate that when mF44 or mF46 was administered to Fc γ RIII-deficient mice, antigen clearance was increased compared to when mIgG1 was administered to the mice antigen clearance. Furthermore, the results of Example 6 showed that when administered to Fc γ RIII-deficient mice, mF44 and mF46 cause similar degrees of antigen elimination as when administered to Fc receptor γ chain-deficient mice and when administered to normal mice.

These results revealed that antigen elimination can be accelerated by enhancing selective binding to Fc γ RIIb alone without enhancing binding to active Fc γ Rs.

In addition to the reported documents discussed so far, based on the above-mentioned assessment results using mice, it is considered that uptake of immunocomplexes into cells *via* Fc γ RIIb takes place *in vivo* in humans as in mice, and as a result, antibodies that have Fc with selectively enhanced binding to human Fc γ RIIb can accelerate elimination of its antigens. Furthermore, as discussed above, since uptake of immunocomplexes into cells *via* Fc γ RIIb is considered to take place at similar rates in mice and humans, effects of accelerating antigen elimination comparable to those of antibodies having Fc with enhanced affinity to mouse Fc γ RIIb may be achieved *in vivo* in humans by using Fc in which affinity to human Fc γ RIIb is enhanced to a similar extent.

As described in WO2009/125825, Fv4-IgG1 is an antibody that results from conferring to a humanized anti-IL-6 receptor antibody H54/L28-IgG1 the activity to bind to the antigen in a pH-dependent manner, *i.e.*, altering the variable region to confer the property to bind to an

antigen at pH 7.4 and dissociate from the antigen at pH 5.8. WO2009/125825 showed that the elimination of soluble human IL-6 receptor is greatly accelerated in mice co-administered with Fv4 IgG1 and soluble human IL-6 receptor as the antigen as compared to mice co-administered with H54/L28-IgG1 and the antigen. Herein, heavy chain H54-IgG1 and light-chain L28-CK included in H54/L28-IgG1 are shown in SEQ ID NO: 36 and SEQ ID NO: 37, respectively; and heavy chain VH3-IgG1 and light-chain VL3-CK included in Fv4-IgG1 are shown in SEQ ID NO: 38 and SEQ ID NO: 39, respectively.

Soluble human IL-6 receptor bound to an antibody H54/L28-IgG1, which binds to soluble human IL-6 receptor, is recycled to the plasma along with the antibody *via* FcRn. Meanwhile, antibody Fv4-IgG1 which binds to soluble human IL-6 receptor in a pH-dependent manner dissociates from the soluble human IL-6 receptor that has been bound to the antibody under an acidic condition in the endosome. Since the dissociated soluble human IL-6 receptor is degraded in the lysosome, elimination of the soluble human IL-6 receptor can be greatly accelerated, and the antibody Fv4-IgG1 which binds to the soluble human IL-6 receptor in a pH-dependent manner is recycled to the plasma after binding to FcRn in the endosome. Since the recycled antibody can bind to a soluble human IL-6 receptor again, binding to the antigen (soluble human IL-6 receptor) and recycling to the plasma *via* FcRn are repeated. As a result, a single antibody molecule can repeatedly bind to the soluble human IL-6 receptor multiple times (WO2009/125825).

On the other hand, as disclosed in the present invention, it was found that plasma concentration of the soluble antigen can be reduced greatly by administration of an antigen-binding molecule with enhanced FcγR-binding activity of the Fcγ receptor-binding domain included in the antigen-binding molecule which comprises an antigen-binding domain in which antigen-binding activity changes depending on the ion concentration condition such as pH, an FcRn-binding domain having FcRn-binding activity under an acidic pH range condition, and an Fcγ receptor-binding domain.

While not being restricted to a particular theory, the unexpected decrease in soluble antigen concentration in plasma observed by administration of an antigen-binding molecule with enhanced binding to FcγRs, which comprises an antigen-binding domain in which antigen-binding activity changes depending on the ion-concentration condition such as pH and an FcRn-binding domain having FcRn-binding activity under an acidic pH range condition can be explained as follows.

As described above, an antigen-binding molecule such as Fv4-IgG1 comprising an antigen-binding domain in which antigen-binding activity changes depending on the ion-concentration condition may be able to bind repeatedly to the antigen multiple times, but the effect of dissociating the soluble antigen in the endosome to accelerate the antigen elimination

from plasma may be dependent on the rate of uptake of the complex of the antigen and antigen-binding molecule into the endosome. The antigen-binding molecules with enhanced binding activities to various FcγRs, which comprise an antigen-binding domain in which antigen-binding activity changes depending on the ion-concentration condition, are actively
5 taken up into cells by binding to various FcγRs expressed on the cell membrane, and can circulate in the plasma again by recycling *via* binding between FcRn and the FcRn-binding domain in the molecule having binding activity to FcRn under an acidic pH range condition. More specifically, since the aforementioned antigen-binding molecules that formed complexes with soluble antigens in plasma are taken up actively into cells *via* FcγRs expressed on the cell
10 membrane, the effect of accelerating elimination of soluble antigens in plasma may be more pronounced than that of antigen-binding molecules whose binding activities to various FcγRs are not enhanced.

FcγR-binding activities of antibodies that bind to membranous antigens have an important role in cytotoxic activity of the antibodies. Therefore, when cytotoxic activity is necessary for
15 an antibody to be used as a pharmaceutical, a human IgG1 isotype which has high FcγR-binding activity is used, and the technique of enhancing the FcγR-binding activities of the antibody to enhance the cytotoxic activity of the antibody is widely utilized. On the other hand, the role of FcγR-binding activities of antibodies that bind to soluble antigens and are used as pharmaceuticals had not been known, and differences in physiological effects on organisms
20 administered with human IgG1 with high FcγR-binding activities and human IgG2 and human IgG4 with low FcγR-binding activities, due to their differences in FcγR-binding activities, had not been fully examined so far. As described later in the Examples, it was actually confirmed that in the plasma of individuals administered with antibodies whose FcγR-binding activities have been lost, changes in soluble-antigen concentration were not affected. On the other hand,
25 in the present invention, the concentration of soluble antigens in the plasma was found to be greatly reduced in individuals administered with antigen-binding molecules with enhanced FcγR-binding activities and comprising an antigen-binding domain whose binding activity to soluble antigens changes depending on the ion concentration condition. More specifically, by combining an FcRn-binding domain having an FcRn-binding activity under an acidic pH range
30 condition and an antigen-binding domain whose binding to soluble antigens changes depending on the ion concentration condition, which are domains included in antigen-binding molecules targeting soluble antigens, an advantage of enhancing binding to FcγR was found for the first time.

35 Antigen-binding molecule

In the present invention, “an antigen-binding molecule” is used in the broadest sense to

refer to a molecule having human FcRn-binding activity at an acidic pH range and containing an antigen-binding domain and an Fc γ receptor-binding domain. Specifically, the antigen-binding molecules include various types of molecules as long as they exhibit the antigen-binding activity. Molecules in which an antigen-binding domain is linked to an Fc region include, for example, antibodies. Antibodies may include single monoclonal antibodies (including agonistic antibodies and antagonistic antibodies), human antibodies, humanized antibodies, chimeric antibodies, and such. Alternatively, when used as antibody fragments, they preferably include antigen-binding domains and antigen-binding fragments (for example, Fab, F(ab')₂, scFv, and Fv). Scaffold molecules where three dimensional structures, such as already-known stable α/β barrel protein structure, are used as a scaffold (base) and only some portions of the structures are made into libraries to construct antigen-binding domains are also included in antigen-binding molecules of the present invention.

An antigen-binding molecule of the present invention may contain at least some portions of an Fc region that mediates the binding to FcRn and Fc γ receptor. In a non-limiting embodiment, the antigen-binding molecule includes, for example, antibodies and Fc fusion proteins. A fusion protein refers to a chimeric polypeptide comprising a polypeptide having a first amino acid sequence that is linked to a polypeptide having a second amino acid sequence that would not naturally link in nature. For example, a fusion protein may comprise the amino acid sequence of at least a portion of an Fc region (for example, a portion of an Fc region responsible for the binding to FcRn or a portion of an Fc region responsible for the binding to Fc γ receptor) and a non-immunoglobulin polypeptide containing, for example, the amino acid sequence of the ligand-binding domain of a receptor or a receptor-binding domain of a ligand. The amino acid sequences may be present in separate proteins that are transported together to a fusion protein, or generally may be present in a single protein; however, they are included in a new rearrangement in a fusion polypeptide. Fusion proteins can be produced, for example, by chemical synthesis, or by genetic recombination techniques to express a polynucleotide encoding peptide regions in a desired arrangement.

Respective domains of the present invention can be linked together via linkers or directly via polypeptide binding.

The linkers comprise arbitrary peptide linkers that can be introduced by genetic engineering, synthetic linkers, and linkers disclosed in, for example, Protein Engineering (1996) 9(3), 299-305. However, peptide linkers are preferred in the present invention. The length of the peptide linkers is not particularly limited, and can be suitably selected by those skilled in the art according to the purpose. The length is preferably five amino acids or more (without particular limitation, the upper limit is generally 30 amino acids or less, preferably 20 amino acids or less), and particularly preferably 15 amino acids.

For example, such peptide linkers preferably include:

- Ser
 - Gly·Ser
 - Gly·Gly·Ser
 - 5 Ser·Gly·Gly
 - Gly·Gly·Gly·Ser (SEQ ID NO: 26)
 - Ser·Gly·Gly·Gly (SEQ ID NO: 27)
 - Gly·Gly·Gly·Gly·Ser (SEQ ID NO: 28)
 - Ser·Gly·Gly·Gly·Gly (SEQ ID NO: 29)
 - 10 Gly·Gly·Gly·Gly·Gly·Ser (SEQ ID NO: 30)
 - Ser·Gly·Gly·Gly·Gly·Gly (SEQ ID NO: 31)
 - Gly·Gly·Gly·Gly·Gly·Gly·Ser (SEQ ID NO: 32)
 - Ser·Gly·Gly·Gly·Gly·Gly·Gly (SEQ ID NO: 33)
 - (Gly·Gly·Gly·Gly·Ser (SEQ ID NO: 28))_n
 - 15 (Ser·Gly·Gly·Gly·Gly (SEQ ID NO: 29))_n
- where n is an integer of 1 or larger. The length or sequences of peptide linkers can be selected accordingly by those skilled in the art depending on the purpose.

Synthetic linkers (chemical crosslinking agents) is routinely used to crosslink peptides, and for example:

- 20 N-hydroxy succinimide (NHS),
- disuccinimidyl suberate (DSS),
- bis(sulfosuccinimidyl) suberate (BS³),
- dithiobis(succinimidyl propionate) (DSP),
- dithiobis(sulfosuccinimidyl propionate) (DTSSP),
- 25 ethylene glycol bis(succinimidyl succinate) (EGS),
- ethylene glycol bis(sulfosuccinimidyl succinate) (sulfo-EGS),
- disuccinimidyl tartrate (DST), disulfosuccinimidyl tartrate (sulfo-DST),
- bis[2-(succinimidoxycarbonyloxy)ethyl] sulfone (BSOCOES),
- and bis[2-(sulfosuccinimidoxycarbonyloxy)ethyl] sulfone (sulfo-BSOCOES). These
- 30 crosslinking agents are commercially available.

When multiple linkers for linking the respective domains are used, they may all be of the same type, or may be of different types.

- In addition to the linkers exemplified above, linkers with peptide tags such as His tag,
- HA tag, myc tag, and FLAG tag may also be suitably used. Furthermore, hydrogen bonding,
- 35 disulfide bonding, covalent bonding, ionic interaction, and properties of binding with each other
- as a result of combination thereof may be suitably used. For example, the affinity between CH1

and CL of antibody may be used, and Fc regions originating from the above-described bispecific antibodies may also be used for hetero Fc region association. Moreover, disulfide bonds formed between domains may also be suitably used.

In order to link respective domains via peptide linkage, polynucleotides encoding the domains are linked together in frame. Known methods for linking polynucleotides in frame include techniques such as ligation of restriction fragments, fusion PCR, and overlapping PCR. Such methods can be appropriately used alone or in combination to construct antigen-binding molecules of the present invention. In the present invention, the terms "linked" and "fused", or "linkage" and "fusion" are used interchangeably. These terms mean that two or more elements or components such as polypeptides are linked together to form a single structure by any means including the above-described chemical linking means and genetic recombination techniques. Fusing in frame means, when two or more elements or components are polypeptides, linking two or more units of reading frames to form a continuous longer reading frame while maintaining the correct reading frames of the polypeptides. When two molecules of Fab are used as an antigen-binding domain, an antibody, which is an antigen-binding molecule of the present invention where the antigen-binding domain is linked in frame to an Fc region via peptide bond without linker, can be used as a preferred antigen-binding molecule of the present invention.

FcRn

Unlike Fc γ receptor belonging to the immunoglobulin superfamily, FcRn, human FcRn in particular, is structurally similar to polypeptides of major histocompatibility complex (MHC) class I, exhibiting 22% to 29% sequence identity to class I MHC molecules (Ghetie *et al.*, Immunol. Today (1997) 18 (12): 592-598). FcRn is expressed as a heterodimer consisting of soluble β or light chain (β 2 microglobulin) complexed with transmembrane α or heavy chain. Like MHC, FcRn α chain comprises three extracellular domains (α 1, α 2, and α 3) and its short cytoplasmic domain anchors the protein onto the cell surface. α 1 and α 2 domains interact with the FcRn-binding domain of the antibody Fc region (Raghavan *et al.*, Immunity (1994) 1: 303-315).

FcRn is expressed in maternal placenta and yolk sac of mammals, and is involved in mother-to-fetus IgG transfer. In addition, in neonatal small intestine of rodents, where FcRn is expressed, FcRn is involved in transfer of maternal IgG across brush border epithelium from ingested colostrum or milk. FcRn is expressed in a variety of other tissues and endothelial cell systems of various species. FcRn is also expressed in adult human endothelia, muscular blood vessels, and hepatic sinusoidal capillaries. FcRn is believed to play a role in maintaining the plasma IgG concentration by mediating recycling of IgG to serum upon binding to IgG. Typically, binding of FcRn to IgG molecules is strictly pH dependent. The optimal binding is

observed in an acidic pH range below 7.0.

Human FcRn whose precursor is a polypeptide having the signal sequence of SEQ ID NO: 34 (the polypeptide with the signal sequence is shown in SEQ ID NO: 35) forms a complex with human β 2-microglobulin *in vivo*. As shown in the Reference Examples described below, soluble human FcRn complexed with β 2-microglobulin is produced by using conventional recombinant expression techniques. FcRn-binding domain of the present invention can be assessed for their binding activity to such a soluble human FcRn complexed with β 2-microglobulin. In the present invention, unless otherwise specified, human FcRn refers to a form capable of binding to an FcRn-binding domain of the present invention. Examples include a complex between human FcRn and human β 2-microglobulin.

Antigen-binding molecules of the present invention have an FcRn-binding domain. The FcRn-binding domain is not particularly limited as long as the antigen-binding molecules have an FcRn-binding activity in an acidic pH range, and it may be a domain that has direct or indirect binding activity to FcRn. Preferred examples of such a domain include the Fc region of IgG immunoglobulin, albumin, albumin domain 3, anti-FcRn antibody, anti-FcRn peptide, anti-FcRn scaffold molecule, and such which have an activity to directly bind to FcRn, or molecules that bind to IgG or albumin, and such that have an activity to indirectly bind to FcRn. In the present invention, a domain that has FcRn-binding activity in an acidic pH range and in a neutral pH range is preferred. If the domain already has FcRn-binding activity in an acidic pH range, it may preferably be used as it is. If the domain does not have or has weak FcRn-binding activity in an acidic pH range, amino acids in the antigen-binding molecule may be altered to confer FcRn-binding activity. Alternatively, amino acids may be altered in a domain already having FcRn-binding activity in an acidic pH range to increase the FcRn-binding activity. For amino acid alteration of the FcRn-binding domain, the alteration of interest can be identified by comparing the FcRn-binding activities in an acidic pH range before and after the amino acid alteration.

The preferred human FcRn-binding domain is a region that directly binds to FcRn. Such preferred FcRn-binding domains include, for example, antibody Fc regions. Meanwhile, regions capable of binding to a polypeptide such as albumin or IgG, which has FcRn-binding activity, can indirectly bind to FcRn *via* albumin, IgG, or such. Therefore, for the FcRn-binding region of the present invention, a region that binds to a polypeptide having FcRn-binding activity may be preferably used. An Fc region contains an amino acid sequence derived from the constant region of an antibody heavy chain. An Fc region is a portion of the antibody heavy chain constant region beginning from the N terminus of the hinge region at the papain cleavage site, which is on the amino acid at approximately position 216 according to EU numbering, and including the hinge, CH2 and CH3 domains.

The binding activity of an FcRn binding domain for FcRn, in particular human FcRn, or an antigen-binding molecule comprising the domain

The binding activity of an FcRn binding domain of the present invention to FcRn, human FcRn in particular, can be measured by methods known to those skilled in the art, as described in the section "Binding Activity" above. Those skilled in the art can appropriately determine the conditions other than pH. The antigen-binding activity and human FcRn-binding activity of an antigen-binding molecule can be assessed based on the dissociation constant (KD), apparent dissociation constant (KD), dissociation rate (kd), apparent dissociation rate (kd), and such. They can be measured by methods known to those skilled in the art. For example, Biacore (GE healthcare), Scatchard plot, or flow cytometer may be used.

When the FcRn-binding activity of an FcRn-binding domain is measured, conditions other than the pH are not particularly limited, and can be appropriately selected by those skilled in the art. Measurements can be carried out, for example, at 37°C using MES buffer, as described in WO 2009/125825. Alternatively, the FcRn-binding activity of an FcRn-binding domain can be measured by methods known to those skilled in the art, and may be measured by using, for example, Biacore (GE Healthcare) or such. The binding activity of an FcRn-binding domain to FcRn can be assessed by pouring, as an analyte, FcRn, an FcRn-binding domain, or an antigen-binding molecule of the present invention containing the FcRn-binding domain into a chip immobilized with an FcRn-binding domain, an antigen-binding molecule of the present invention containing the FcRn-binding domain, or FcRn.

The acidic pH range presented as the condition for having binding activity between FcRn and an FcRn-binding domain in an antigen-binding molecule of the present invention generally refers to pH 4.0 to pH 6.5. Preferably it refers to pH 5.5 to pH 6.5, and particularly preferably it refers to pH 5.8 to pH 6.0 which is close to the pH in an early endosome *in vivo*. The neutral pH range presented as the condition for having binding activity between FcRn and an antigen-binding molecule of the present invention or an FcRn-binding domain included in such a molecule generally refers to pH 6.7 to pH 10.0. Neutral pH range is preferably a range indicated by any pH value from pH 7.0 to pH 8.0, and is preferably selected from pH 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, and 8.0, and is particularly preferably pH 7.4 which is close to the pH in plasma (in blood) *in vivo*. When evaluating the binding affinity between human FcRn and a human FcRn-binding domain or an antigen-binding molecule containing that domain at pH 7.4 is difficult due to low binding affinity, pH 7.0 can be used instead of pH 7.4. As temperature to be used in assay conditions, binding affinity between an FcRn-binding domain and FcRn may be assessed at any temperature from 10°C to 50°C. Preferably, a temperature from 15°C to 40°C is used to determine the binding affinity between an FcRn-binding domain

and human FcRn. More preferably, any temperature from 20°C to 35°C such as any one of 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, and 35°C is also equally used to determine the binding affinity between an FcRn-binding domain and FcRn. A temperature of 25°C is a non-limiting example of an embodiment of the present invention.

5 According to Yeung *et al.* (J. Immunol. (2009) 182, 7663-7671), the human FcRn-binding activity of native human IgG1 in an acidic pH range (pH 6.0) is KD 1.7 μ M, but the activity can be hardly detected in a neutral pH range. Therefore, in a preferred embodiment, an antigen-binding molecule of the present invention having human FcRn-binding activity under an acidic pH range condition, which includes antigen-binding molecules whose human
10 FcRn-binding activity under an acidic pH range condition is KD 20 μ M or stronger and human FcRn-binding activity under a neutral pH range condition is equivalent to that of a native human IgG may be used. In a more preferred embodiment, an antigen-binding molecule of the present invention including antigen-binding molecules whose human FcRn-binding activity under an acidic pH range condition is KD 2.0 μ M or stronger may be used. In an even more preferred
15 embodiment, antigen-binding molecules whose human FcRn-binding activity under an acidic pH range condition is KD 0.5 μ M or stronger may be used. The above-mentioned KD values are determined by the method described in The Journal of Immunology (2009) 182: 7663-7671 (by immobilizing antigen-binding molecule onto a chip and loading human FcRn as the analyte).

In the present invention, an Fc region that has FcRn-binding activity under an acidic pH
20 range condition is preferred. If the domain is an Fc region already having FcRn-binding activity under an acidic pH range condition, it can be used as it is. If the domain does not have or has weak FcRn-binding activity under an acidic pH range condition, amino acids in the antigen-binding molecule may be altered to obtain an Fc region having the desired FcRn-binding activity. An Fc region having the desired FcRn-binding activity or enhanced FcRn-binding
25 activity under an acidic pH range condition can also be preferably obtained by altering amino acids in the Fc region. Amino acid alteration of an Fc region that confers such a desired binding activity can be identified by comparing the FcRn-binding activity under an acidic pH range condition before and after the amino acid alteration. Those skilled in the art can make appropriate amino acid alterations using a well-known method similar to the aforementioned
30 method used to alter the Fc γ receptor-binding activity.

An Fc region having an FcRn-binding activity under an acidic pH range condition, which is included in an antigen-binding molecule of the present invention, may be obtained by any method; but specifically, an FcRn-binding domain having FcRn-binding activity or having enhanced FcRn-binding activity under an acidic pH range condition can be obtained by altering
35 amino acids of a human IgG immunoglobulin used as the starting Fc region. Preferred Fc regions of IgG immunoglobulins for the alteration include Fc regions of human IgG (IgG1, IgG2,

IgG3, IgG4, and variants thereof). For alterations to other amino acids, amino acids at any position may be altered as long as the Fc region has FcRn-binding activity under an acidic pH range condition, or the binding activity to human FcRn under an acidic range condition can be increased. When an antigen-binding molecule includes an Fc region of a human IgG1 as the Fc region, it preferably includes an alteration that has the effect of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1. Preferred examples of such amino acids that can be altered include amino acids at positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 386, 388, 400, 413, 415, 424, 433, 434, 435, 436, 439, and/or 447 (EU numbering) as described in WO2000/042072. Similarly, preferred examples of amino acids that can be altered also include amino acids at positions 251, 252, 254, 255, 256, 308, 309, 311, 312, 385, 386, 387, 389, 428, 433, 434, and/or 436 (EU numbering) as described in WO2002/060919. Furthermore, such amino acids that can be altered also include amino acids at positions 250, 314, and 428 (EU numbering) as described in WO2004/092219. Such amino acids that can be altered further include amino acids at positions 251, 252, 307, 308, 378, 428, 430, 434, and/or 436 (EU numbering) as described in WO2010/045193. These amino acid alterations enhance the FcRn-binding under an acidic pH range condition of an Fc region of an IgG immunoglobulin.

In the present invention, an Fc region that has an FcRn-binding activity under an acidic pH range condition is preferred. If the domain is an Fc region already having FcRn-binding activity under an acidic pH range condition, it may be used as it is. If the domain does not have or has weak FcRn-binding activity under an acidic pH range condition, amino acids in the antigen-binding molecule may be altered to obtain an Fc region having the desired FcRn-binding activity. An Fc region having the desired FcRn-binding activity or enhanced FcRn-binding activity under an acidic pH range condition can preferably be obtained by altering amino acids in the Fc region. Amino acid alteration of an Fc region that confers such a desired binding activity can be identified by comparing the FcRn-binding activity under an acidic pH range condition before and after the amino acid alteration. Those skilled in the art can make appropriate amino acid alterations using a well-known method similar to the aforementioned method used to alter the Fcγ receptor-binding activity.

An Fc region having FcRn-binding activity under an acidic pH range condition, which is included in an antigen-binding molecule of the present invention may be obtained by any method; but specifically, an FcRn-binding domain having binding activity or having enhanced binding activity to FcRn under an acidic pH range condition can be obtained by altering amino acids of a human IgG immunoglobulin used as the starting Fc region. Preferred examples of Fc regions of IgG immunoglobulins for the alteration include Fc regions of human IgG (IgG1, IgG2,

IgG3, IgG4, and variants thereof). For alterations to other amino acids, amino acids at any position may be altered as long as an Fc region has FcRn-binding activity under an acidic pH range condition, or the binding activity to human FcRn under an acidic range condition can be increased. When an antigen-binding molecule comprises an Fc region of a human IgG1 as the

5 Fc region, it preferably includes an alteration that has effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1. Examples of such amino acids that can be altered include amino acids at positions 252, 254, 256, 309, 311, 315, 433, and/or 434, as well as amino acids that may be combined with them, which are amino acids at positions 253, 310, 435, and/or 426 (EU numbering) as described

10 in WO1997/034631. Preferred amino acids are amino acids at positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 386, 388, 400, 413, 415, 424, 433, 434, 435, 436, 439, and/or 447 (EU numbering) as described in WO2000/042072. Similarly, preferred examples of such amino acids that can be altered also include amino acids at positions 251, 252, 254, 255, 256, 308, 309, 311, 312, 385,

15 386, 387, 389, 428, 433, 434, and/or 436 (EU numbering) as described in WO2002/060919. Furthermore, such amino acids that can be altered include amino acids at positions 250, 314, and 428 (EU numbering) as described in WO2004/092219. In addition, preferred examples of such amino acids that can be altered include amino acids at positions 238, 244, 245, 249, 252, 256, 257, 258, 260, 262, 270, 272, 279, 283, 285, 286, 288, 293, 307, 311, 312, 316, 317, 318, 332,

20 339, 341, 343, 375, 376, 377, 378, 380, 382, 423, 427, 430, 431, 434, 436, 438, 440, and/or 442 as described in WO2006/020114. Other preferred examples of such amino acids that can be altered also include amino acids at positions 251, 252, 307, 308, 378, 428, 430, 434, and/or 436 (EU numbering) as described in WO2010/045193. These amino acid alterations enhance the FcRn-binding under an acidic pH range condition of an Fc region of an IgG immunoglobulin.

25 In a non-limiting embodiment of alterations that produce the effect of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 when an Fc region of human IgG1 is included as the Fc region, examples include at least one or more amino acid alterations selected from the group consisting of:

- either Arg or Leu at amino acid position 251;
- 30 any of Phe, Ser, Thr, and Tyr at amino acid position 252;
- either Ser or Thr at amino acid position 254;
- any of Arg, Gly, Ile, and Leu at amino acid position 255;
- any of Ala, Arg, Asn, Asp, Gln, Glu, and Thr at amino acid position 256;
- either Ile or Thr at amino acid position 308;
- 35 Pro at amino acid position 309;
- any of Glu, Leu, and Ser at amino acid position 311;

either Ala or Asp at amino acid position 312;
 either Ala or Leu at amino acid position 314;
 any of Ala, Arg, Asp, Gly, His, Lys, Ser, and Thr at amino acid position 385;
 any of Arg, Asp, Ile, Lys, Met, Pro, Ser, and Thr at amino acid position 386;
 5 any of Ala, Arg, His, Pro, Ser, and Thr at amino acid position 387;
 any of Asn, Pro, and Ser at amino acid position 389;
 any of Leu, Met, Phe, Ser, and Thr at amino acid position 428;
 any of Arg, Gln, His, Ile, Lys, Pro, and Ser at amino acid position 433;
 any of His, Phe, and Tyr at amino acid position 434; and
 10 any of Arg, Asn, His, Lys, Met, and Thr at amino acid position 436
 according to EU numbering. The number of amino acids to be altered is not particularly limited,
 and the amino acid may be altered at just one position or at two or more positions.

When an Fc region of human IgG1 is included as the Fc region, a non-limiting
 embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH
 15 range condition compared to the binding activity of the starting Fc region of human IgG1 can be
 an alteration(s) comprising Ile at amino acid position 308, Pro at amino acid position 309, and/or
 Glu at amino acid position 311 according to EU numbering. In another non-limiting
 embodiment of the alterations, the alterations may be Thr at amino acid position 308, Pro at
 amino acid position 309, Leu at amino acid position 311, Ala at amino acid position 312, and/or
 20 Ala at amino acid position 314. In yet another non-limiting embodiment of the alterations, the
 alterations may be Ile or Thr at amino acid position 308; Pro at amino acid position 309; Glu,
 Leu, or Ser at amino acid position 311; Ala at amino acid position 312; and/or Ala or Leu at
 amino acid position 314. In a different non-limiting embodiment of the alterations, the
 alterations can be Thr at amino acid position 308, Pro at amino acid position 309, Ser at amino
 25 acid position 311, Asp at amino acid position 312, and/or Leu at amino acid position 314.

When an Fc region of human IgG1 is included as the Fc region, a non-limiting
 embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH
 range condition compared to the binding activity of the starting Fc region of human IgG1 can be
 an alteration(s) comprising Leu at amino acid position 251, Tyr at amino acid position 252, Ser
 30 or Thr at amino acid position 254, Arg at amino acid position 255, and/or Glu at amino acid
 position 256 according to EU numbering.

When an Fc region of human IgG1 is included as the Fc region, a non-limiting
 embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH
 range condition compared to the binding activity of the starting Fc region of human IgG1 can be
 35 an alteration(s) comprising any one of Leu, Met, Phe, Ser, and Thr at amino acid position 428;
 any one of Arg, Gln, His, Ile, Lys, Pro, and Ser at amino acid position 433; any one of His, Phe,

and Tyr at amino acid position 434; and/or any one of Arg, Asn, His, Lys, Met, and Thr at amino acid position 436 indicated by EU numbering. In another non-limiting embodiment of the alteration(s), the alteration(s) may include His or Met at amino acid position 428 and/or His or Met at amino acid position 434.

- 5 When an Fc region of human IgG1 is included as the Fc region, a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 can be an alteration(s) comprising Arg at amino acid position 385, Thr at amino acid position 386, Arg at amino acid position 387, and/or Pro at amino acid position 389 according to EU numbering.
- 10 In another non-limiting embodiment of the alterations, the alteration(s) can be Asp at amino acid position 385, Pro at amino acid position 386, and/or Ser at amino acid position 389.

- Furthermore, when an Fc region of human IgG1 is included as the Fc region, a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human
- 15 IgG1, alterations include those of at least one or more amino acids selected from the group consisting of:
- either Gln or Glu at amino acid position 250; and
- either Leu or Phe at amino acid position 428
- according to EU numbering. The number of amino acids to be altered is not particularly limited,
- 20 and the amino acid may be altered at one position alone or at two or more positions.

- When an Fc region of human IgG1 is included as the Fc region, a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 can be an alteration(s) comprising Gln at amino acid position 250, and/or either Leu or Phe at amino
- 25 acid position 428 according to EU numbering. In another non-limiting embodiment of the alterations, the alterations can be those comprising Glu at amino acid position 250 and/or either Leu or Phe at amino acid position 428.

- When an Fc region of human IgG1 is included as the Fc region, a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH
- 30 range condition compared to the binding activity of the starting Fc region of human IgG1, examples include alterations of at least two or more amino acids selected from the group consisting of:
- either Asp or Glu at amino acid position 251;
- Tyr at amino acid position 252;
- 35 Gln at amino acid position 307;
- Pro at amino acid position 308;

Val at amino acid position 378;

Ala at amino acid position 380;

Leu at amino acid position 428;

either Ala or Lys at amino acid position 430;

5 any one of Ala, His, Ser, and Tyr at amino acid position 434; and

Ile at amino acid position 436

according to EU numbering. The number of amino acids to be altered is not particularly limited, and the amino acid may be altered at only two positions or at three or more positions.

When an Fc region of human IgG1 is included as the Fc region, a non-limiting
 10 embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 can be alteration(s) comprising Gln at amino acid position 307, and either Ala or Ser at amino acid position 434 according to EU numbering. In another non-limiting embodiment of the alterations, the alterations may include Pro at amino acid position 308 and Ala at amino acid
 15 position 434. In yet another non-limiting embodiment of the alterations, the alterations may include Tyr at amino acid position 252 and Ala at amino acid position 434. In a different non-limiting embodiment of the alterations, the alterations may include Val at amino acid position 378 and Ala at amino acid position 434. In another non-limiting embodiment of the alterations, the alterations may include Leu at amino acid position 428 and Ala at amino acid
 20 position 434. In yet another non-limiting embodiment of the alterations, the alterations may include Ala at amino acid position 434 and Ile at amino acid position 436. Furthermore, in another non-limiting embodiment of the alterations, the alterations may include Pro at amino acid position 308 and Tyr at amino acid position 434. Furthermore, in another different non-limiting embodiment of the alterations, the alterations may include Gln at amino acid
 25 position 307 and Ile at amino acid position 436.

When an Fc region of human IgG1 is included as the Fc region, a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 can be an alteration(s) comprising Gln at amino acid position 307, Ala at amino acid position 380, and
 30 Ser at amino acid position 434 as according to EU numbering. In another non-limiting embodiment of the alterations, the alterations may include Gln at amino acid position 307, Ala at amino acid position 380, and Ala at amino acid position 434. In yet another non-limiting embodiment of the alterations, the alterations may include Tyr at amino acid position 252, Pro at amino acid position 308, and Tyr at amino acid position 434. In a different non-limiting
 35 embodiment of the alterations, the alterations may include Asp at amino acid position 251, Gln at amino acid position 307, and His at amino acid position 434.

- When an Fc region of human IgG1 is included as the Fc region, in a non-limiting embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1, examples include alterations of at least two or more amino acids selected from the group
- 5 consisting of:
 - Leu at amino acid position 238;
 - Leu at amino acid position 244;
 - Arg at amino acid position 245;
 - Pro at amino acid position 249;
 - 10 Tyr at amino acid position 252;
 - Pro at amino acid position 256;
 - any one of Ala, Ile, Met, Asn, Ser, and Val at amino acid position 257;
 - Asp at amino acid position 258;
 - Ser at amino acid position 260;
 - 15 Leu at amino acid position 262;
 - Lys at amino acid position 270;
 - either Leu or Arg at amino acid position 272;
 - any one of Ala, Asp, Gly, His, Met, Asn, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 279;
 - 20 any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 283;
 - Asn at amino acid position 285;
 - Phe at amino acid position 286;
 - either Asn or Pro at amino acid position 288;
 - 25 Val at amino acid position 293;
 - any one of Ala, Glu, and Met at amino acid position 307;
 - any one of Ala, Ile, Lys, Leu, Met, Val, and Trp at amino acid position 311;
 - Pro at amino acid position 312;
 - Lys at amino acid position 316;
 - 30 Pro at amino acid position 317;
 - either Asn or Thr at amino acid position 318;
 - any one of Phe, His, Lys, Leu, Met, Arg, Ser, and Trp at amino acid position 332;
 - any one of Asn, Thr, and Trp at amino acid position 339;
 - Pro at amino acid position 341;
 - 35 any one of Glu, His, Lys, Gln, Arg, Thr, and Tyr at amino acid position 343;
 - Arg at amino acid position 375;

- any one of Gly, Ile, Met, Pro, Thr, and Val at amino acid position 376;
 Lys at amino acid position 377;
 either Asp or Asn at amino acid position 378;
 any one of Asn, Ser, and Thr at amino acid position 380;
 5 any one of Phe, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 382;
 Asn at amino acid position 423;
 Asn at amino acid position 427;
 any one of Ala, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, and Tyr at amino acid position 430;
 10 either His or Asn at amino acid position 431;
 any one of Phe, Gly, His, Trp, and Tyr at amino acid position 434;
 any one of Ile, Leu, and Thr at amino acid position 436;
 any one of Lys, Leu, Thr, and Trp at amino acid position 438;
 15 Lys at amino acid position 440; and
 Lys at amino acid position 442;
 according to EU numbering. The number of amino acids to be altered is not particularly limited, and the amino acid may be altered only at two positions or at three or more positions.

- When an Fc region of human IgG1 is included as the Fc region, a non-limiting
 20 embodiment of alterations that produce effects of enhancing binding to FcRn under an acidic pH range condition compared to the binding activity of the starting Fc region of human IgG1 can be alterations comprising Ile at amino acid position 257 and Ile at amino acid position 311 according to EU numbering. In another non-limiting embodiment of the alterations, the alterations may include Ile at amino acid position 257 and His at amino acid position 434. In
 25 yet another non-limiting embodiment of the alterations, the alterations may include Val at amino acid position 376 and His at amino acid position 434.

- Furthermore, as described later, for the FcRn-binding domain included in an antigen-binding molecule of the present invention, an Fc region having FcRn-binding activity under a neutral pH range condition may also be preferably used. Such Fc region can be
 30 obtained by any method according to the aforementioned method of obtaining Fc regions having FcRn-binding activity under an acidic pH range condition, but specifically, an FcRn-binding domain having a binding activity or having enhanced binding activity to FcRn under a neutral pH range condition can be obtained by altering amino acids of a human IgG immunoglobulin used as the starting Fc region. Preferred Fc regions of IgG immunoglobulins for the alteration
 35 include Fc regions of human IgG (IgG1, IgG2, IgG3, IgG4, and variants thereof). For alterations to other amino acids, amino acid at any position may be altered as long as Fc region

has FcRn-binding activity under a neutral pH range condition or the binding activity to human FcRn under an acidic pH range condition can be increased. When an antigen-binding molecule includes an Fc region of a human IgG1 as the Fc region, it preferably comprises an alteration that has effects of enhancing binding to FcRn under a neutral pH range condition compared to the binding activity of the starting Fc region of human IgG1. Preferred examples of such altered Fc regions include human FcRn-binding domains in which at least one or more amino acids selected from the group consisting of amino acids at positions 237, 238, 239, 248, 250, 252, 254, 255, 256, 257, 258, 265, 270, 286, 289, 297, 298, 303, 305, 307, 308, 309, 311, 312, 314, 315, 317, 325, 332, 334, 360, 376, 380, 382, 384, 385, 386, 387, 389, 424, 428, 433, 434, and 436 in the starting Fc region site according to EU numbering are different from the corresponding amino acids in the native Fc region.

Preferred examples of such altered Fc regions include Fc regions comprising at least one or more amino acids selected from the group consisting of:

- Met at amino acid position 237;
- Ala at amino acid position 238;
- Lys at amino acid position 239;
- Ile at amino acid position 248;
- any one of Ala, Phe, Ile, Met, Gln, Ser, Val, Trp, and Tyr at amino acid position 250;
- any one of Phe, Trp, and Tyr at amino acid position 252;
- Thr at amino acid position 254;
- Glu at amino acid position 255;
- any one of Asp, Glu, and Gln at amino acid position 256;
- any one of Ala, Gly, Ile, Leu, Met, Asn, Ser, Thr, and Val at amino acid position 257;
- His at amino acid position 258;
- Ala at amino acid position 265;
- Phe at amino acid position 270;
- either Ala or Glu at amino acid position 286;
- His at amino acid position 289;
- Ala at amino acid position 297;
- Gly at amino acid position 298;
- Ala at amino acid position 303;
- Ala at amino acid position 305;
- any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 307;
- any one of Ala, Phe, Ile, Leu, Met, Pro, Gln, and Thr at amino acid position 308;
- any one of Ala, Asp, Glu, Pro, and Arg at amino acid position 309;

- any one of Ala, His, and Ile at amino acid position 311;
 either Ala or His at amino acid position 312;
 either Lys or Arg at amino acid position 314;
 either Ala or His at amino acid position 315;
- 5 Ala at amino acid position 317;
 Gly at amino acid position 325;
 Val at amino acid position 332;
 Leu at amino acid position 334;
 His at amino acid position 360;
- 10 Ala at amino acid position 376;
 Ala at amino acid position 380;
 Ala at amino acid position 382;
 Ala at amino acid position 384;
 either Asp or His at amino acid position 385;
- 15 Pro at amino acid position 386;
 Glu at amino acid position 387;
 either Ala or Ser at amino acid position 389;
 Ala at amino acid position 424;
 any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at
 20 amino acid position 428;
 Lys at amino acid position 433;
 any one of Ala, Phe, His, Ser, Trp, and Tyr at amino acid position 434; and
 His at amino acid position 436;
 in the Fc region according to EU numbering.
- 25 For example, by using these amino acid alterations alone or multiple alterations in
 combination, binding of an IgG Fc region to FcRn in an acidic pH range and/or neutral pH range
 can be enhanced; however, the amino acid alterations to be introduced are not particularly
 limited, and as long as the effect of improving plasma retention is conferred, any amino acid
 alteration may be introduced.

30

Pharmaceutical compositions

- When a conventional neutralizing antibody against a soluble antigen is administered, the
 plasma retention of the antigen is expected to be prolonged by binding to the antibody. In
 general, antibodies have a long half-life (one week to three weeks) while the half-life of antigen
 35 is generally short (one day or less). Meanwhile, antibody-bound antigens have a significantly
 longer half-life in plasma as compared to when the antigens are present alone. For this reason,

administration of existing neutralizing antibody results in an increased antigen concentration in plasma. Such cases have been reported with various neutralizing antibodies that target soluble antigens including, for example, IL-6 (J. Immunotoxicol. (2005) 3, 131-139), amyloid beta (mAbs (2010) 2 (5), 1-13), MCP-1 (ARTHRITIS & RHEUMATISM (2006) 54, 2387-2392),
 5 hepcidin (AAPS J. (2010) 4, 646-657), and sIL-6 receptor (Blood (2008) 112 (10), 3959-64). Administration of existing neutralizing antibodies has been reported to increase the total plasma antigen concentration to about 10 to 1,000 times (the level of increase varies depending on antigen) the base line. Herein, the total plasma antigen concentration refers to a concentration as a total amount of antigen in plasma, *i.e.*, the sum of concentrations of antibody-bound and
 10 antibody-unbound antigens. An increase in the total plasma antigen concentration is undesirable for such antibody pharmaceuticals that target a soluble antigen. The reason is that the antibody concentration has to be higher than at least the total plasma antigen concentration to neutralize the soluble antigen. Specifically, "the total plasma antigen concentration is increased to 10 to 1,000 times" means that, in order to neutralize the antigen, the plasma antibody
 15 concentration (*i.e.*, antibody dose) has to be 10 to 1,000 times higher as compared to when increase in the total plasma antigen concentration does not occur. Conversely, if the total plasma antigen concentration can be reduced by 10 to 1,000 times as compared to the existing neutralizing antibody, the antibody dose can also be reduced to similar extent. Thus, antibodies capable of decreasing the total plasma antigen concentration by eliminating the soluble antigen
 20 from plasma are highly useful as compared to existing neutralizing antibodies.

While the present invention is not restricted to a particular theory, the reason for increase in the number of antigens that can bind to a single antigen-binding molecule and the reason for enhanced dissipation of plasma antigen concentration when an antigen-binding molecule is administered to a living organism which leads to increase of uptake of the
 25 antigen-binding molecule into cells *in vivo*, wherein the antigen-binding molecule comprises an antigen-binding domain whose antigen-binding activity changes depending on the ion-concentration condition so that the antigen-binding activity is lower under an acidic pH range condition than under a neutral pH range condition, and an FcRn-binding domain such as an antibody constant region having human Fc γ receptor-binding activity under a neutral pH
 30 range condition, can be explained as follows.

For example, when an antibody that binds to a membrane antigen is administered *in vivo*, after binding to an antigen, the antibody is, in a state bound to the antigen, incorporated into the endosome *via* intracellular internalization. Then, the antibody is transferred to the lysosome while remaining bound to the antigen, and is degraded together with the antigen there. The
 35 internalization-mediated elimination from plasma is referred to as antigen-dependent elimination, and has been reported for many antibody molecules (Drug Discov Today (2006) 11(1-2), 81-88).

When a single IgG antibody molecule binds to antigens in a divalent manner, the single antibody molecule is internalized while remaining bound to the two antigens, and is degraded in the lysosome. In the case of typical antibodies, thus, a single IgG antibody molecule cannot bind to three antigen molecules or more. For example, a single IgG antibody molecule having a
5 neutralizing activity cannot neutralize three antigen molecules or more.

The plasma retention of IgG molecule is relatively long (the elimination is slow) since FcRn, which is known as a salvage receptor for IgG molecule, functions. IgG molecules incorporated into endosomes by pinocytosis bind under the endosomal acidic condition to FcRn expressed in endosomes. IgG molecules that cannot bind to human FcRn are transferred to
10 lysosomes and degraded there. Meanwhile, IgG molecules bound to FcRn are transferred to cell surface. The IgG molecules are dissociated from FcRn under the neutral condition in plasma, and thus they are recycled back to plasma.

Alternatively, when antigen-binding molecules are antibodies that bind to a soluble antigen, the *in vivo* administered antibodies bind to antigens, and then the antibodies are
15 incorporated into cells while remaining bound to the antigens. Most of antibodies incorporated into cells bind to FcRn in the endosome and then are transferred to cell surface. The antibodies are dissociated from FcRn under the neutral condition in plasma and released to the outside of cells. However, antibodies having typical antigen-binding domains whose antigen-binding activity does not vary depending on ion concentration condition such as pH are released to the
20 outside of cells while remaining bound to the antigens, and thus cannot bind to an antigen again. Thus, like antibodies that bind to membrane antigens, single typical IgG antibody molecule whose antigen-binding activity does not vary depending on ion concentration condition such as pH cannot bind to three antigen molecules or more.

Antibodies that bind to antigens in a pH-dependent manner, which strongly bind to
25 antigens under the neutral pH range condition in plasma and are dissociated from antigens under the endosomal acidic pH range condition (antibodies that bind to antigens under the neutral pH range condition and are dissociated under an acidic pH range condition), and antibodies that bind to antigens in a calcium ion concentration-dependent manner, which strongly bind to antigens under a high calcium ion concentration condition in plasma and are dissociated from antigens
30 under a low calcium ion concentration condition in the endosome (antibodies that bind to antigens under a high calcium ion concentration condition and are dissociated under a low calcium ion concentration condition) can be dissociated from antigen in the endosome. Antibodies that bind to antigens in a pH-dependent manner or in a calcium ion concentration-dependent manner, when recycled to plasma by FcRn after dissociation from
35 antigens, can again bind to an antigen. Thus, such single antibody molecule can repeatedly bind to several antigen molecules. Meanwhile, antigens bound to antigen-binding molecules

are dissociated from antibody in the endosome and degraded in the lysosome without recycling to plasma. By administering such antigen-binding molecules *in vivo*, antigen uptake into cells is accelerated, and it is possible to decrease plasma antigen concentration.

Uptake of antigens bound by antigen-binding molecules into cells are further enhanced
5 by conferring or enhancing the Fcγ receptor-binding activity under the neutral pH range condition (pH 7.4) to antibodies that bind to antigens in a pH-dependent manner, which strongly bind to antigens under the neutral pH range condition in plasma and are dissociated from antigens under the endosomal acidic pH range condition (antibodies that bind to antigens under the neutral pH range condition and are dissociated under an acidic pH range condition), and
10 antibodies that bind to antigens in a calcium ion concentration-dependent manner, which strongly bind to antigens under a high calcium ion concentration condition in plasma and are dissociated from antigens under a low calcium ion concentration condition in the endosome (antibodies that bind to antigens under a high calcium ion concentration condition and are dissociated under a low calcium ion concentration condition). Specifically, by administering
15 such antigen-binding molecules *in vivo*, the antigen elimination is accelerated, and it is possible to reduce plasma antigen concentration. Typical antibodies that do not have the ability to bind to antigens in a pH-dependent manner or in a calcium ion concentration-dependent manner, and antigen-antibody complexes of such antibodies are incorporated into cells by non-specific endocytosis, and transported onto cell surface by binding to FcRn under the endosomal acidic
20 condition. They are dissociated from FcRn under the neutral condition on cell surface and recycled to plasma. Thus, when an antibody that binds to an antigen in a fully pH-dependent manner (that binds under the neutral pH range condition and is dissociated under an acidic pH range condition) or in a fully calcium ion concentration-dependent manner (that binds under a high calcium ion concentration condition and is dissociated under a low calcium ion
25 concentration condition) binds to an antigen in plasma and is dissociated from the antigen in the endosome, the rate of antigen elimination is considered to be equal to the rate of uptake into cells of the antibody or antigen/antibody complex by non-specific endocytosis. When the pH or calcium ion concentration dependency of antigen/antibody binding is insufficient, antigens that are not dissociated from antibodies in the endosome are, along with the antibodies, recycled to
30 plasma. On the other hand, when the pH dependency is sufficiently strong, the rate limiting step of antigen elimination is the cellular uptake by non-specific endocytosis. Meanwhile, FcRn transports antibodies from the endosome to the cell surface, and a fraction of FcRn is expected to be also distributed on the cell surface.

The present inventors considered that IgG immunoglobulins having binding activity or
35 having enhanced binding activity to Fcγ receptors under a neutral pH range condition can bind to Fcγ receptors present on the cell surface, and that the IgG immunoglobulins are taken up into

cells in an Fc γ receptor-dependent manner by binding to Fc γ receptors present on the cell surface. The rate of uptake into cells *via* Fc γ receptors is faster than the rate of uptake into cells by non-specific endocytosis. Therefore, it is thought that the rate of antigen elimination by antigen-binding molecules can further be accelerated by enhancing the ability to bind to Fc γ receptors under a neutral pH range condition. That is, antigen-binding molecules having the ability to bind to Fc γ receptors under a neutral pH range condition uptake antigens into cells more quickly than common (native human) IgG immunoglobulins, and after binding to FcRn in the endosome and dissociating the antigens, they are recycled again into plasma, and they bind again to antigens and are taken up into cells *via* Fc γ receptors. Since turnover of this cycle can be increased by increasing the ability to bind to Fc γ receptors under a neutral pH range condition, the rate of antigen elimination from plasma is accelerated. Furthermore, by making antigen-binding molecules that have decreased antigen-binding activity under an acidic pH range condition compared to antigen-binding activity under a neutral pH range condition, the rate of antigen elimination from plasma can further be accelerated. By increasing the number of cycles resulting from accelerating the rate of turnover of this cycle, the number of antigen molecules that can be bound by a single antigen-binding molecule may increase. Antigen-binding molecules of the present invention comprise an antigen-binding domain and an Fc γ receptor-binding domain, and since the Fc γ receptor-binding domain does not affect antigen binding, and based on the above-mentioned mechanism, it is considered that regardless of the type of antigens, antigen uptake into cells by antigen-binding molecules can be enhanced and the rate of antigen elimination can be accelerated by reducing binding activity (binding ability) of the antigen-binding molecule to antigens under an ion concentration condition such as acidic pH range or low calcium ion concentration condition compared to binding activity (binding ability) to antigens under an ion concentration condition such as neutral pH range or high calcium ion concentration condition, and/or enhancing binding activity to Fc γ receptors at pH in plasma. Therefore, antigen-binding molecules of the present invention may exhibit better effects than conventional therapeutic antibodies in terms of reduction of side effects caused by antigens, increase in antibody dose, and improvement of *in vivo* kinetics of antibodies.

Fig. 1 shows an embodiment of the mechanism for eliminating soluble antigens from plasma by administering ion concentration-dependent antigen-binding antibodies with enhanced binding to Fc γ receptors at neutral pH compared to conventional neutralizing antibodies. Herein below, hydrogen ion concentration (that is, pH)-dependent antigen-binding antibodies will be described as an example of the ion concentration-dependent antigen-binding antibodies, but the mechanisms are not limited to hydrogen ion concentration. Existing neutralizing antibodies which do not have pH-dependent antigen-binding ability may be gradually taken up mainly by non-specific interactions with cells after binding to soluble antigens in plasma. The

complexes of neutralizing antibodies and soluble antigens, which are taken up into cells, translocate to acidic endosomes and are recycled into plasma by FcRn. On the other hand, a pH-dependent antigen-binding antibody with enhanced binding to Fc γ receptors under a neutral condition binds to a soluble antigen in plasma, and thereafter, it may be quickly taken up into
5 cells expressing Fc γ receptors on the cell surface *via* interaction with an Fc γ receptor besides non-specific interaction. Here, the soluble antigens bound to pH-dependent antigen-binding antibody dissociate from the antibody due to pH-dependent binding ability in the acidic endosomes. Soluble antigens that have dissociated from the antibody then translocate to lysosomes, and they are degraded by proteolysis. Meanwhile, antibodies that released the
10 soluble antigens bind to FcRn in the acidic endosome, and are then recycled onto the cell membrane by FcRn, and are released again into plasma. In this way, the free antibodies that are recycled can bind again to other soluble antigens. Such pH-dependent antigen-binding antibodies whose binding to Fc γ receptors under a neutral condition can translocate a large amount of soluble antigens to lysosomes to reduce the total antigen concentration in plasma by
15 repeating a cycle of: uptake into cells *via* Fc γ receptors; dissociation and degradation of soluble antigens; and antibody recycling.

Specifically, the present invention also relates to pharmaceutical compositions comprising antigen-binding molecules of the present invention, antigen-binding molecules produced by alteration methods of the present invention, or antigen-binding molecules produced
20 by production methods of the present invention. Antigen-binding molecules of the present invention or antigen-binding molecules produced by production methods of the present invention are useful as pharmaceutical compositions since they, when administered, have the strong effect to reduce the plasma antigen concentration as compared to typical antigen-binding molecules, and exhibit the improved *in vivo* immune response, pharmacokinetics, and others in animals
25 administered with the molecules. The pharmaceutical compositions of the present invention may comprise pharmaceutically acceptable carriers.

In the present invention, pharmaceutical compositions generally refer to agents for treating or preventing, or testing and diagnosing diseases.

The pharmaceutical compositions of the present invention can be formulated by
30 methods known to those skilled in the art. For example, they can be used parenterally, in the form of injections of sterile solutions or suspensions including water or other pharmaceutically acceptable liquid. For example, such compositions can be formulated by mixing in the form of unit dose required in the generally approved medicine manufacturing practice, by appropriately combining with pharmacologically acceptable carriers or media, specifically with sterile water,
35 physiological saline, vegetable oil, emulsifier, suspension, surfactant, stabilizer, flavoring agent, excipient, vehicle, preservative, binder, or such. In such formulations, the amount of active

ingredient is adjusted to obtain an appropriate amount in a pre-determined range.

Sterile compositions for injection can be formulated using vehicles such as distilled water for injection, according to standard formulation practice.

Aqueous solutions for injection include, for example, physiological saline and isotonic solutions containing dextrose or other adjuvants (for example, D-sorbitol, D-mannose, D-mannitol, and sodium chloride). It is also possible to use in combination appropriate solubilizers, for example, alcohols (ethanol and such), polyalcohols (propylene glycol, polyethylene glycol, and such), non-ionic surfactants (polysorbate 80(TM), HCO-50, and such).

Oils include sesame oil and soybean oils. Benzyl benzoate and/or benzyl alcohol can be used in combination as solubilizers. It is also possible to combine buffers (for example, phosphate buffer and sodium acetate buffer), soothing agents (for example, procaine hydrochloride), stabilizers (for example, benzyl alcohol and phenol), and/or antioxidants. Appropriate ampules are filled with the prepared injections.

The pharmaceutical compositions of the present invention are preferably administered parenterally. For example, the compositions in the dosage form for injections, transnasal administration, transpulmonary administration, or transdermal administration are administered. For example, they can be administered systemically or locally by intravenous injection, intramuscular injection, intraperitoneal injection, subcutaneous injection, or such.

Administration methods can be appropriately selected in consideration of the patient's age and symptoms. The dose of a pharmaceutical composition containing an antigen-binding molecule can be, for example, from 0.0001 to 1,000 mg/kg for each administration. Alternatively, the dose can be, for example, from 0.001 to 100,000 mg per patient. However, the present invention is not limited by the numeric values described above. The doses and administration methods vary depending on the patient's weight, age, symptoms, and such. Those skilled in the art can set appropriate doses and administration methods in consideration of the factors described above.

Amino acids contained in the amino acid sequences of the present invention may be post-translationally modified (for example, the modification of an N-terminal glutamine into a pyroglutamic acid by pyroglutamylation is well-known to those skilled in the art). Naturally, such post-translationally modified amino acids are included in the amino acid sequences in the present invention.

Methods that use antigen-binding molecules of the present invention

The present invention also provides a method comprising contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an

antigen-binding domain and an Fc γ receptor-binding domain, in which antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition and the Fc γ receptor-binding domain has higher binding activity to the Fc γ receptor under a neutral pH range condition than a native Fc γ receptor-binding domain to which the sugar chain linked at position 5 297 (EU numbering) is a fucose-containing sugar chain, wherein the method is any one of the following:

- (i) a method of increasing the number of antigens to which a single antigen-binding molecule can bind;
- (ii) a method of eliminating plasma antigens;
- 10 (iii) a method of improving pharmacokinetics of the antigen-binding molecule;
- (iv) a method of promoting intracellular dissociation of an antigen from the antigen-binding molecule, wherein the antigen was extracellularly bound to the antigen-binding molecule;
- (v) a method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form; and
- 15 (vi) a method of decreasing the concentration of free antigen or the total antigen concentration in plasma.

Furthermore, the present invention provides a method comprising enhancing Fc γ receptor-binding activity under a neutral pH range condition of the Fc γ receptor-binding domain in an antigen-binding molecule compared to that of a native Fc γ receptor-binding domain to 20 which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an Fc γ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition, wherein the method is any one of the following:

- 25 (i) a method of altering an antigen-binding molecule in which intracellular uptake of the antigen to be bound is enhanced;
- (ii) a method of increasing the number of antigens to which a single antigen-binding molecule can bind;
- (iii) a method of increasing the ability of the antigen-binding molecule to eliminate plasma 30 antigens;
- (iv) a method of improving pharmacokinetics of antigen-binding molecule;
- (v) a method of promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule;
- 35 (vi) a method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form, wherein the antigen-binding molecule had been taken up into a

- cell in an antigen-bound form; and
- (vii) a method of altering an antigen-binding molecule, which can decrease total antigen concentration or free antigen concentration in plasma.

Examples of methods of contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo* include (1) the so-called *ex vivo* method where plasma containing the antigen-binding molecules and antigens that bind to the antigen-binding molecules is temporarily taken out from a living organism; contacted with cells expressing Fcγ receptors; and after a certain period of time, the plasma containing the extracellularly recycled (or also referred to as re-secreted or recirculated) antigen-binding molecules without bounded antigens is then placed back into the living organism, and (2) the method of administering the antigen-binding molecules to a living organism. In the method of (1), one may also utilize a method of: temporarily taking out from a living organism plasma containing the antigens to which the antigen-binding molecules bind; contacting with cells expressing the Fcγ receptors and antigen-binding molecules; and after a certain period of time, returning the plasma into the living organism.

In the present invention, Fcγ receptor-expressing cells are not limited to particular cells and any kind of cells may be used as long as they are cells that express the desired Fcγ receptor(s). To identify cells that express the desired Fcγ receptor(s), publicly known databases such as Human Protein Atlas may be used. Furthermore, whether the receptors are expressed in cells used for contacting antigen-binding molecules of the present invention can be confirmed by a method that confirms expression of a gene encoding the desired Fcγ receptor(s), or by an immunological method that uses antibodies that bind to the desired Fcγ receptor(s). These methods are also publicly known. Since contact of Fcγ receptor-expressing cells with antigen-binding molecules and antigens that bind to the antigen-binding molecules is carried out *in vivo* as well, contacting Fcγ receptor-expressing cells with antigen-binding molecules in the present invention include administering the antigen-binding molecules to living organisms. The duration of contact is, for example, a suitable duration from among one minute to several weeks, 30 minutes to one week, one hour to three days, and two hours to one day. More specifically, the duration necessary for uptake of the antigen-binding molecules or antigens bound to the antigen-binding molecules into cells by endocytosis mediated by Fcγ receptors is appropriately employed for the duration of contact. For example, various immune cells may be used for the Fcγ receptor-expressing cells.

A method of enhancing binding activity of the Fcγ receptor-binding domain to an Fcγ receptor under a neutral pH range condition than that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain is described in the later-described section on method for producing antigen-binding molecules of

the present invention.

Method of altering an antigen-binding molecule with enhanced intracellular uptake of antigens that bind to the antigen-binding molecule

5 The present invention provides methods of altering an antigen-binding molecule with enhanced intracellular uptake of antigens that bind to the antigen-binding molecule, wherein the method comprises enhancing, under a neutral pH range condition, Fcγ receptor-binding activity of the Fcγ receptor-binding domain in an antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a
10 fucose-containing sugar chain, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an Fcγ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

 In the present invention, "intracellular uptake of antigens" by an antigen-binding
15 molecule means that antigens are taken up into cells by Fcγ receptor-mediated endocytosis and internalization. In the present invention, "enhance uptake into cells" means that rate of uptake into cells of antigen-binding molecules bound to antigens in plasma is increased and/or the amount of antigens that were taken up which are recycled in plasma is reduced, and it is only necessary that the rate of uptake into cells is increased compared to the antigen-binding molecule
20 prior to reduction of antigen-binding activity (binding ability) of the antigen-binding molecule under an ion concentration condition such as an acidic pH range or low calcium ion concentration compared to the antigen-binding activity under an ion concentration condition such as a neutral pH range or high calcium ion concentration in addition to enhancing binding activity of the antigen-binding molecule to a Fcγ receptor in a neutral pH range, and it is
25 preferred that the rate is increased compared to a native human IgG to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, and it is particularly preferred that the rate is increased compared to any of a native human IgG1, IgG2, IgG3, or IgG4. Accordingly, in the present invention, whether intracellular uptake of antigens by antigen-binding molecules is enhanced can be determined by observing whether the rate of
30 uptake of antigens into cells increased. The rate of uptake of antigens into cells can be calculated, for example, by adding the antigen-binding molecule and antigen to a culture medium containing Fcγ receptor-expressing cells and measuring the decrease in antigen concentration in the culture medium over time or measuring the amount of antigens taken up into the Fcγ receptor-expressing cells over time. Using the method of increasing the rate by which antigens
35 are taken up into cells by the antigen-binding molecule of the present invention, for example, the rate of antigen elimination in plasma can be increased by administering the antigen-binding

molecule. Therefore, whether uptake of antigen into cells by the antigen-binding molecule is enhanced can also be confirmed, for example, by assessing whether the rate of antigen elimination in plasma is accelerated, or whether administration of antigen-binding molecule reduces the total antigen concentration in plasma.

5 In the present invention, "total antigen concentration in plasma" means the sum of the concentrations of antigens bound to the antigen-binding molecules and unbound antigens, or "free antigen concentrations in plasma" which is the concentration of antigens not bound to antigen-binding molecules. Various methods for measuring "total antigen concentration in plasma" or "free antigen concentrations in plasma" are well-known in the art as herein described
10 below.

In the present invention, "native human IgG" means unmodified human IgG, and is not limited to a particular class of IgG. Furthermore, in "native human IgG", it is desirable that the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain. This means that as long as human IgG1, IgG2, IgG3, or IgG4 can bind to human FcRn in an acidic pH
15 range, it can be used as a "native human IgG". Preferably, "native human IgG" may be human IgG1.

Method of increasing the number of antigens that can be bound by a single antigen-binding molecule

20 The present invention provides a method of increasing the number of antigens to which a single antigen-binding molecule can bind, wherein the method comprises contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fcγ receptor-binding domain, in which an
25 antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor under a neutral pH range condition compared to a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

30 Furthermore, the present invention provides a method of increasing the number of antigens to which a single antigen-binding molecule can bind, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in an antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a
35 fucose-containing sugar chain, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an Fcγ receptor-binding domain and

an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

The phrase "number of antigens to which a single antigen-binding molecule can bind" in the present invention means the number of antigens to which an antigen-binding molecule can
5 bind until the molecule is degraded and eliminated. The phrase "increasing the number of antigens to which a single antigen-binding molecule can bind" in the present invention refers to increasing the number of times an antigen molecule bound to the antigen-binding molecule dissociates and the antigen-binding molecule binds again to an antigen molecule. Antigen
10 molecules that bind to the antigen-binding molecule may be the same antigen molecule or different molecules existing in the reaction system where both molecules are present. In other words, it refers to the total number of times the antigen-binding molecule binds to an antigen in the reaction system. In another expression, when one cycle is assumed to be intracellular uptake of an antigen-binding molecule bound to an antigen, dissociation of the antigen in an endosome, and then return of the antigen-binding molecule to the outside of the cell, the phrase
15 refers to increase in the number of times this cycle can be repeated until the antigen-binding molecule is degraded and eliminated. After binding to an Fcγ receptor, antigen-binding molecules of the present invention having Fcγ receptor-binding activity in a neutral pH range is taken up into a cell expressing this Fcγ receptor by endocytosis. The antigen-binding molecule of the present invention that dissociated from the Fcγ receptor under an ion concentration
20 condition such as an acidic pH range or low calcium ion concentration is recycled to the outside of the cell again by binding to FcRn under an acidic pH range condition. The antigen-binding molecule of the present invention which is recycled to outside the cell after dissociating antigen from the antigen-binding molecule under an ion concentration condition such as an acidic pH range or low calcium ion concentration can bind again to an antigen. Therefore, whether the
25 number of cycles increased may be assessed by determining whether the aforementioned "intracellular uptake is enhanced", or whether the later described "pharmacokinetics is improved".

Method of eliminating plasma antigens or method of increasing the ability of the antigen-binding molecule to eliminate plasma antigens
30

The present invention provides a method of eliminating plasma antigens, which comprises contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fcγ receptor-binding
35 domain, in which an antigen binding activity of the antigen-binding domain changes depending on the ion concentration condition, and the Fcγ receptor-binding domain has higher binding

activity to the Fcγ receptor under a neutral pH range condition compared to a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

Furthermore, the present invention provides a method of increasing the ability of the antigen-binding molecule to eliminate plasma antigens, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in an antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an Fcγ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

In the present invention, the phrase "method of increasing the ability to eliminate plasma antigens" has the same meaning as "method of increasing the ability of an antigen-binding molecule to eliminate antigens from plasma".

In the present invention, "ability to eliminate plasma antigens" refers to the ability to eliminate from plasma the antigens present in plasma when antigen-binding molecules are administered *in vivo* or antigen-binding molecules are secreted *in vivo*. Therefore, in the present invention, all that the phrase "an ability of an antigen-binding molecule to eliminate plasma antigens increases" has to mean is that when an antigen-binding molecule is administered, rate of antigen elimination from the plasma is accelerated compared to before reducing the antigen-binding activity of the antigen-binding molecule under an ion concentration condition such as acidic pH range or low calcium ion concentration compared to the antigen-binding activity under a neutral pH range or high calcium ion concentration in addition to enhancing binding activity of the antigen-binding molecule to a Fcγ receptor in a neutral pH range. Whether the ability of antigen-binding molecules to eliminate plasma antigens increases can be determined, for example, by administering soluble antigens and antigen-binding molecules to a living organism and measuring the plasma concentration of the soluble antigens after the administration. When the concentration of soluble antigens in plasma is decreased after administration of soluble antigens and antigen-binding molecules by enhancing the binding activity of the antigen-binding molecules to Fcγ receptors under a neutral pH range condition, or by reducing the antigen-binding activity of the antigen-binding molecule under an ion concentration condition such as acidic pH range or low calcium ion concentration compared to the antigen-binding activity under an ion concentration condition such as a neutral pH range or high calcium ion concentration in addition to enhancing Fcγ receptor-binding activity, it can be determined that the ability of antigen-binding molecules to eliminate plasma antigens is

enhanced. Soluble antigens may be antigens to which antigen-binding molecules are actually bounded (antigens in the form of an antigen/antigen-binding molecule complex), or antigens to which antigen-binding molecules are not bound, and their concentrations may be determined as "concentration of antigen-binding molecule-bound antigens in plasma" and "concentration of antigen-binding molecule-unbound antigens in plasma", respectively (the latter has the same meaning as "free antigen concentrations in plasma"). Since "total antigen concentration in plasma" means the sum of the concentrations of antigen-binding molecule-bound antigens and antigen-binding molecule-unbound antigens, or "concentration of free antigens in plasma" which is the concentration of antigen-binding molecule-unbound antigens, soluble antigen concentration can be determined as "total antigen concentration in plasma". Various methods of measuring "total antigen concentration in plasma" or "free antigen concentrations in plasma" are well-known in the art as herein described below.

Method of improving the pharmacokinetics of antigen-binding molecules

The present invention provides a method of improving pharmacokinetics of an antigen-binding molecule, which comprises contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fcγ receptor-binding domain, in which an antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition, and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor under a neutral pH range condition compared to a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

Furthermore, the present invention provides a method of improving the pharmacokinetics of an antigen-binding molecule, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in an antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, wherein the antigen-binding molecule having human FcRn-binding activity under an acidic pH range condition comprises an Fcγ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

Herein, "enhancement of pharmacokinetics", "improvement of pharmacokinetics", and "superior pharmacokinetics" can be restated as "enhancement of plasma (blood) retention", "improvement of plasma (blood) retention", "superior plasma (blood) retention", and "prolonged plasma (blood) retention". These terms are synonymous.

Herein, "improvement of pharmacokinetics" means not only prolongation of the period

until elimination from the plasma (for example, until the antigen-binding molecule is degraded intracellularly or the like and cannot return to the plasma) after administration of the antigen-binding molecule to humans, or non-human animals such as mice, rats, monkeys, rabbits, and dogs, but also prolongation of the plasma retention of the antigen-binding molecule in a

5 form that allows antigen binding (for example, in an antigen-free form of the antigen-binding molecule) during the period of administration to elimination due to degradation. Native IgG can bind to FcRn from non-human animals. For example, mouse can be preferably used to be administered in order to confirm the property of the antigen-binding molecule of the invention since native human IgG can bind to mouse FcRn stronger than to human FcRn (Int Immunol.

10 (2001) 13(12): 1551-1559). As another example, mouse in which its native FcRn genes are deficient and a transgene for human FcRn gene is harbored to be expressed (Methods Mol Biol. 2010; 602: 93-104) can also be preferably used as a subject to be administered in order to confirm the property of the antigen-binding molecule of the invention described hereinafter.

Specifically, "improvement of pharmacokinetics" also includes prolongation of the period until

15 elimination due to degradation of the antigen-binding molecule not bound to antigens (the antigen-free form of antigen-binding molecule). The antigen-binding molecule in plasma cannot bind to a new antigen if the antigen-binding molecule has already bound to an antigen. Thus, the longer the period that the antigen-binding molecule is not bound to an antigen, the longer the period that it can bind to a new antigen (the higher the chance of binding to another

20 antigen). This enables reduction of the time period that an antigen is free of the antigen-binding molecule *in vivo* and prolongation of the period that an antigen is bound to the antigen-binding molecule. The plasma concentration of the antigen-free form of antigen-binding molecule can be increased and the period that the antigen is bound to the antigen-binding molecule can be prolonged by accelerating the antigen elimination from the plasma by administration of the

25 antigen-binding molecule. Specifically, herein "improvement of the pharmacokinetics of antigen-binding molecule" includes the improvement of a pharmacokinetic parameter of the antigen-free form of the antigen-binding molecule (any of prolongation of the half-life in plasma, prolongation of mean retention time in plasma, and impairment of plasma clearance),

prolongation of the period that the antigen is bound to the antigen-binding molecule after

30 administration of the antigen-binding molecule, and acceleration of antigen-binding molecule-mediated antigen elimination from the plasma. The improvement of pharmacokinetics of antigen-binding molecule can be assessed by determining any one of the parameters, half-life in plasma, mean plasma retention time, and plasma clearance for the antigen-binding molecule or the antigen-free form thereof ("Pharmacokinetics: Enshu-niyoru

35 Rikai (Understanding through practice)" Nanzando). For example, the plasma concentration of the antigen-binding molecule or antigen-free form thereof is determined after administration of

the antigen-binding molecule to mice, rats, monkeys, rabbits, dogs, or humans. Then, each parameter is determined. When the plasma half-life or mean plasma retention time is prolonged, the pharmacokinetics of the antigen-binding molecule can be judged to be improved. The parameters can be determined by methods known to those skilled in the art. The parameters
5 can be appropriately assessed, for example, by noncompartmental analysis using the pharmacokinetics analysis software WinNonlin (Pharsight) according to the appended instruction manual. The plasma concentration of antigen-free antigen-binding molecule can be determined by methods known to those skilled in the art, for example, using the assay method measured in a known method (Clin Pharmacol. 2008 Apr; 48(4): 406-417).

10 Herein, "improvement of pharmacokinetics" also includes prolongation of the period that an antigen is bound to an antigen-binding molecule after administration of the antigen-binding molecule. Whether the period that an antigen is bound to the antigen-binding molecule after administration of the antigen-binding molecule is prolonged can be assessed by determining the plasma concentration of free antigen. The prolongation can be judged based on
15 the determined plasma concentration of free antigen or the time period required for an increase in the ratio of free antigen concentration to the total antigen concentration.

The plasma concentration of free antigen not bound to the antigen-binding molecule or the ratio of free antigen concentration to the total concentration can be determined by methods known to those skilled in the art, for example, the method measured in Pharm Res. 2006 Jan; 23
20 (1): 95-103 can be used. Alternatively, when an antigen exhibits a particular function *in vivo*, whether the antigen is bound to an antigen-binding molecule that neutralizes the antigen function (antagonistic molecule) can be determined by testing whether the antigen function is neutralized. Whether the antigen function is neutralized can be assessed by assaying an *in vivo* marker that reflects the antigen function. Whether the antigen is bound to an antigen-binding molecule that
25 activates the antigen function (agonistic molecule) can be assessed by assaying an *in vivo* marker that reflects the antigen function.

Determination of the plasma concentration of free antigen and ratio of the amount of free antigen in plasma to the amount of total antigen in plasma, *in vivo* marker assay, and such measurements are not particularly limited; however, the assays are preferably carried out after a
30 certain period of time has passed after administration of the antigen-binding molecule. In the present invention, the period after administration of the antigen-binding molecule is not particularly limited; those skilled in the art can determine the appropriate period depending on the properties and the like of the administered antigen-binding molecule. Such periods include, for example, one day after administration of the antigen-binding molecule, three days after
35 administration of the antigen-binding molecule, seven days after administration of the antigen-binding molecule, 14 days after administration of the antigen-binding molecule, and 28

days after administration of the antigen-binding molecule. Herein, "plasma antigen concentration" refers to either "total antigen concentration in plasma" which is the sum of antigen-binding molecule bound antigen and non-bound antigen concentration or "free antigen concentration in plasma" which is antigen-binding molecule non-bound antigen concentration.

5 Total antigen concentration in plasma can be lowered by administration of antigen-binding molecule of the present invention by 2-fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, 1,000-fold, or even higher compared to the administration of a reference antigen-binding molecule comprising the native human IgG Fc region in which a sugar chain linked to at position 297 (EU numbering) is a sugar chain having fucose as an Fc γ receptor-binding domain or compared to when antigen-binding domain molecule of the present invention comprising the antigen-binding domain is not administered.

Molar antigen/antigen-binding molecule ratio can be calculated as shown below;
 value A: Molar antigen concentration at each time point
 value B: Molar antigen-binding molecule concentration at each time point
 15 value C: Molar antigen concentration per molar antigen-binding molecule concentration (molar antigen/antigen-binding molecule ratio) at each time point
 $C=A/B$.

Smaller value C indicates higher efficiency of antigen elimination per antigen-binding molecule whereas higher value C indicates lower efficiency of antigen elimination per
 20 antigen-binding molecule.

Molar antigen/antigen-binding molecule ratio can be calculated as described above.

Molar antigen/antigen-binding molecule ratio can be lowered by administration of antigen-binding molecule of present invention by 2-fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, 1,000-fold, or even higher as compared to the administration of a
 25 reference antigen-binding molecule comprising the wild-type human IgG Fc region as a human Fc γ receptor-binding domain.

In the present invention, a native human IgG I, IgG2, IgG3 or IgG4 is preferably used as the native human IgG which is used as a reference native human IgG to be compared with the antigen-binding molecules for their Fc γ receptor-binding activity or *in vivo* activity. Preferably,
 30 a reference antigen-binding molecule comprising the same antigen-binding domain as the antigen-binding molecule of interest and a native human IgG Fc region as the Fc γ receptor-binding domain can be appropriately used. More preferably, a native human IgG1 is used as a reference native human IgG to be compared with the antigen-binding molecules for their Fc γ receptor-binding activity or *in vivo* activity. Furthermore, in the present invention, as
 35 a reference antigen-binding molecule to be compared with the antigen-binding molecules of the present invention, an antigen-binding molecule comprising an antigen-binding domain whose

antigen-binding activity does not change depending on the ion concentration, an antigen-binding molecule comprising an FcRn-binding domain whose FcRn-binding activity under an acidic pH range condition has not been enhanced, an antigen-binding molecule comprising an Fc γ receptor-binding domain that does not have selective binding activity to Fc γ receptors, or such
 5 may also be appropriately used depending on the purpose.

Reduction of total antigen concentration in plasma or molar antigen/antibody ratio can be assessed as described in Examples 6, 8, and 13. More specifically, using human FcRn transgenic mouse line 32 or line 276 (Jackson Laboratories, Methods Mol Biol. 2010; 602: 93-104), they can be assessed by either antigen-antibody co-administration model or steady-state
 10 antigen infusion model when antigen-binding molecule do not cross-react to the mouse counterpart antigen. When an antigen-binding molecule cross-react with mouse counterpart, they can be assessed by simply administering antigen-binding molecule to human FcRn transgenic mouse line 32 or line 276 (Jackson Laboratories). In co-administration model, mixture of antigen-binding molecule and antigen is administered to the mouse. In steady-state
 15 antigen infusion model, infusion pump containing antigen solution is implanted to the mouse to achieve constant plasma antigen concentration, and then antigen-binding molecule is administered to the mouse. Test antigen-binding molecule is administered at same dosage. Total antigen concentration in plasma, free antigen concentration in plasma and plasma antigen-binding molecule concentration is measured at appropriate time point using method
 20 known to those skilled in the art.

For assessing the effects of an Fc γ receptor-binding domain having selective binding activity to Fc γ receptors, when an antigen-binding molecule does not cross-react with a mouse counterpart antigen, total antigen concentration in plasma or decrease in antigen/antibody mole ratio can be assessed by either the antigen-antibody simultaneous injection model or the
 25 steady-state antigen injection model using the conventionally used C57BL/6J mice (Charles River Japan). When an antigen-binding molecule cross-reacts with the mouse counterpart, the antigen-binding molecule can simply be injected to conventionally used C57BL/6J mice (Charles River Japan) to carry out the assessment.

Total or free antigen concentration in plasma and molar antigen/antigen-binding molecule ratio can be measured at 2, 4, 7, 14, 28, 56, or 84 days after administration to evaluate
 30 the long-term effect of the present invention. In other words, a long term plasma antigen concentration is determined by measuring total or free antigen concentration in plasma and molar antigen/ antigen-binding molecule ratio at 2, 4, 7, 14, 28, 56, or 84 days after administration of an antigen-binding molecule in order to evaluate the property of the antigen-binding molecule of
 35 the present invention. Whether the reduction of plasma antigen concentration or molar antigen/antigen-binding molecule ratio is achieved by antigen-binding molecule described in the

present invention can be determined by the evaluation of the reduction at any one or more of the time points described above.

5 Total or free antigen concentration in plasma and molar antigen/antigen-binding molecule ratio can be measured at 15 min, 1, 2, 4, 8, 12, or 24 hours after administration to evaluate the short-term effect of the present invention. In other words, a short term plasma antigen concentration is determined by measuring total or free antigen concentration in plasma and molar antigen/antigen-binding molecule ratio at 15 min, 1, 2, 4, 8, 12, or 24 hours after administration of an antigen-binding molecule in order to evaluate the property of the antigen-binding molecule of the present invention.

10 Route of administration of an antigen-binding molecule of the present invention can be selected from intradermal, intravenous, intravitreal, subcutaneous, intraperitoneal, parenteral and intramuscular injection.

In the present invention, improvement of pharmacokinetics in human is preferred. When the plasma retention in human is difficult to determine, it may be predicted based on the plasma retention in mice (for example, normal mice, human antigen-expressing transgenic mice, 15 human FcRn-expressing transgenic mice) or monkeys (for example, cynomolgus monkeys).

Method of promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule

20 The present invention provides a method of promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule, wherein the method comprises contacting the antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fcγ receptor-binding domain, in which an antigen-binding 25 activity of the antigen-binding domain changes depending on the ion concentration condition, and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor under a neutral pH range condition compared to a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

30 The present invention provides a method of promoting intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in an antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, wherein 35 the antigen-binding molecule has human FcRn-binding activity under an acidic pH range

condition and comprises an Fcγ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

In the present invention, the place where an antigen dissociates from an antigen-binding molecule may be any as long as it is in a cell, but preferably it is in an early endosome. In the present invention, "intracellular dissociation of an antigen from an antigen-binding molecule, wherein the antigen has been extracellularly bound to the antigen-binding molecule" does not have to mean that all antigens taken up into cells after being bound to antigen-binding molecules are intracellularly dissociated from the antigen-binding molecules, and the intracellular dissociation of antigens from antigen-binding molecules only needs to be higher in ratio when compared to before lowering the antigen-binding activity of the antigen-binding molecules under an ion concentration condition such as an acidic pH range or low calcium ion concentration than under an ion concentration condition such as a neutral pH range or high calcium ion concentration and increasing the Fcγ receptor-binding activity in a neutral pH range. Furthermore, the method of promoting intracellular dissociation of an antigen from an antigen-binding molecule may be referred to as a method that enhances intracellular uptake of antigen-bound antigen-binding molecules and confers to the antigen-binding molecules the property of easing promotion of intracellular dissociation of antigens from the antigen-binding molecules.

20 Method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form

The present invention provides a method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form, wherein the method comprises contacting an antigen-binding molecule with an Fcγ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fcγ receptor-binding domain, in which an antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition, and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor under a neutral pH range condition compared to a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

Furthermore, the present invention provides a method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form, wherein the antigen-binding molecule has been taken up into a cell in an antigen-bound form, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in the antigen-binding molecule compared to that of a native Fcγ

receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain, wherein the antigen-binding molecule has human-FcRn-binding activity under an acidic pH range condition and comprises an Fc γ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

In the present invention, the phrase "extracellular release of an antigen-binding molecule in an antigen-unbound form, wherein the antigen-binding molecule has been taken up into a cell in an antigen-bound form" does not have to mean that all antigen-binding molecules that has been taken up into a cell in an antigen-bound form are extracellularly released in an antigen-unbound form, and the ratio of antigen-binding molecules extracellularly released in an antigen-unbound form only needs to be higher when compared to before lowering the antigen-binding activity of the antigen-binding molecules under an ion concentration condition such as an acidic pH range or low calcium ion concentration than under an ion concentration condition such as a neutral pH range or high calcium ion concentration and increasing the Fc γ receptor-binding activity in a neutral pH range. The extracellularly released antigen-binding molecules preferably maintain antigen-binding activity. Furthermore, the method of promoting extracellular release of an antigen-binding molecule in an antigen-unbound form, wherein the antigen-binding molecule has been taken up into a cell in an antigen-bound form may be referred to as a method that enhances uptake of antigen-bound antigen-binding molecules into cells and confers to the antigen-binding molecules the property of easing promotion of extracellular release of an antigen-binding molecule in an antigen-unbound form.

Method of decreasing total antigen concentration or free antigen concentration in plasma, or a method of altering an antigen-binding molecule which can decrease total antigen concentration or free antigen concentration in plasma

The present invention provides a method of decreasing total antigen concentration or free antigen concentration in plasma, wherein the method comprises contacting the antigen-binding molecule with an Fc γ receptor-expressing cell *in vivo* or *ex vivo*, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an antigen-binding domain and an Fc γ receptor-binding domain, in which an antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition, and the Fc γ receptor-binding domain has higher binding activity to the Fc γ receptor under a neutral pH range condition compared to a native Fc γ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

Furthermore, the present invention provides method of altering an antigen-binding

molecule which can decrease total antigen concentration or free antigen concentration in plasma, wherein the method comprises enhancing Fcγ receptor-binding activity under a neutral pH range condition of the Fcγ receptor-binding domain in the antigen-binding molecule compared to that of a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU
5 numbering) is a fucose-containing sugar chain, wherein the antigen-binding molecule has human FcRn-binding activity under an acidic pH range condition and comprises an Fcγ receptor-binding domain and an antigen-binding domain whose antigen-binding activity changes depending on the ion concentration condition.

The method of assessing decrease of total antigen concentration or free antigen
10 concentration in plasma is described in the aforementioned section on the method of improving pharmacokinetics of antigen-binding-molecules.

Ex vivo method of eliminating the antigens from plasma

An example of a non-limiting embodiment of the use of an antigen-binding molecule for
15 the method of eliminating the antigens from plasma, which is provided by the present invention, includes use of the antigen-binding molecule for a so-called *ex vivo* method of eliminating the antigens from plasma, which comprises contacting the antigen-binding molecule of the present invention with plasma isolated from subjects to allow forming immunocomplexes, and allowing the immunocomplexes to contact cells expressing Fcγ receptors and FcRn. The rate of
20 elimination of plasma antigens can be promoted by replacing/combining a method of administering antigen-binding molecules *in vivo* with a so-called *ex vivo* method where plasma containing antigen-binding molecules and antigens that bind to the antigen-binding molecules is temporarily taken out from a living organism, and then contacted with cells expressing Fcγ receptors and FcRn, and the plasma containing extracellularly recycled (or also referred to as
25 re-secreted or recirculated) antigen-binding molecules without bound antigen after a certain period of time are returned into the living organism.

Furthermore, an example of a non-limiting embodiment of the use of an antigen-binding molecule for the method of eliminating the antigens from plasma, which is provided by the present invention, includes use of the antigen-binding molecule for a so-called *ex vivo* method of
30 eliminating the antigens from plasma, which comprises contacting immunocomplexes present in plasma isolated from subjects who have been administered with the antigen-binding molecules of the present invention with cells expressing FcRn and Fcγ receptors.

Whether or not the antigens are eliminated from plasma can be confirmed, for example, by evaluating whether or not the rate of elimination of plasma antigens mentioned above is
35 promoted when, instead of the antigen-binding molecule of the present invention, an antigen-binding molecule comprising an antigen-binding domain in which the antigen-binding

activity does not change depending on the ion concentration, an antigen-binding molecule comprising an FcRn-binding domain whose FcRn-binding activity under an acidic pH range condition has not been enhanced, or an antigen-binding molecule comprising an Fcγ receptor-binding domain that does not have selective binding activity to Fcγ receptors is used as
 5 a control for comparison.

Method of producing antigen-binding molecules

The present invention also provides a method of producing an antigen-binding molecule comprising an antigen-binding domain and an Fcγ receptor-binding domain, and human
 10 FcRn-binding activity under an acidic pH range condition, wherein an antigen-binding activity of the antigen-binding domain changes depending on the ion concentration condition and the Fcγ receptor-binding domain has higher binding activity to the Fcγ receptor under a neutral pH range condition than a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain.

15 Specifically, the present invention provides a method of producing an antigen-binding molecule, which comprises the steps of the following (a) to (f):

- (a) determining an antigen-binding activity of an antigen-binding domain under a condition of high calcium ion concentration;
- (b) determining an antigen-binding activity of the antigen-binding domain under a condition
 20 of low calcium ion concentration;
- (c) selecting an antigen-binding domain for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
- (d) linking a polynucleotide encoding the antigen-binding domain selected in (c) to a polynucleotide encoding an Fcγ receptor-binding domain having a human FcRn-binding
 25 activity in an acidic pH range, which has higher binding activity to the Fcγ receptor under a neutral pH range condition than a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain;
- (e) culturing cells introduced with a vector to which the polynucleotide obtained in (d) is operably linked; and
- 30 (f) collecting antigen-binding molecules from the cell culture of (e).

The present invention also provides a method of producing an antigen-binding molecule, which comprises the steps of the following (a) to (f):

- (a) determining an antigen-binding activity of an antibody under a high calcium ion concentration condition;
- 35 (b) determining an antigen-binding activity of the antibody under a low calcium ion concentration condition;

- (c) selecting an antibody for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
- (d) linking a polynucleotide encoding an antigen-binding domain of the antibody selected in (c) to a polynucleotide encoding an Fcγ receptor-binding domain having human
 5 FcRn-binding activity in an acidic pH range and having higher binding activity to the Fcγ receptor under a neutral pH range condition than a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain;
- (e) culturing cells introduced with a vector to which the polynucleotide obtained in (d) is
 10 operably linked; and
- (f) collecting antigen-binding molecules from the cell culture of (e).

In addition, the present invention provides a method of producing an antigen-binding molecule, which comprises the steps of the following (a) to (f):

- (a) determining an antigen-binding activity of an antigen-binding domain under a neutral pH
 15 range condition;
- (b) determining an antigen-binding activity of the antigen-binding domain under an acidic pH range condition;
- (c) selecting the antigen-binding domain for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);
- 20 (d) linking a polynucleotide encoding the antigen-binding domain selected in (c) to a polynucleotide encoding an Fcγ receptor-binding domain having a human FcRn-binding activity under an acidic pH range condition, which has higher binding activity to the Fcγ receptor under a neutral pH range condition than a native Fcγ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar
 25 chain;
- (e) culturing cells introduced with a vector to which the polynucleotide obtained in (d) is operably linked; and
- (f) collecting antigen-binding molecules from the cell culture of (e).

In addition, the present invention provides a method of producing an antigen-binding
 30 molecule, which comprises the steps of the following (a) to (f):

- (a) determining an antigen-binding activity of an antibody under a neutral pH range condition;
- (b) determining an antigen-binding activity of the antibody under an acidic pH range condition;
- 35 (c) selecting the antibody for which the antigen-binding activity determined in (a) is higher than the antigen-binding activity determined in (b);

- (d) linking a polynucleotide encoding the antigen-binding domain of the antibody selected in (c) to a polynucleotide encoding an Fc γ receptor-binding domain having human-FcRn-binding activity in an acidic pH range and having higher binding activity to the Fc γ receptor under a neutral pH range condition than a native Fc γ receptor-binding domain to which the sugar chain linked at position 297 (EU numbering) is a fucose-containing sugar chain;
- (e) culturing cells introduced with a vector to which the polynucleotide obtained in (d) is operably linked; and
- (f) collecting antigen-binding molecules from the cell culture of (e).

10 The terms "cell", "cell line", and "cell culture" are used synonymously herein, and such designations may include all progeny of a cell or cell line. Thus, for example, the terms "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content due to deliberate or inadvertent mutations. Mutant
15 progeny that have substantially the same function or biological activity as screened for in the originally transformed cell may also be included. Where distinct designations are intended, such intention will be clear from the context of the description.

 When referring to the expression of a coding sequence, the term "control sequences" refers to DNA nucleotide sequences that are necessary for the expression of an operably linked
20 coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes include, for example, a promoter, optionally an operator sequence, a ribosome binding site, and possibly, other sequences as yet poorly understood. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers for the expression of a coding sequence.

25 For a nucleic acid, the term "operably linked" means that the nucleic acid is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a precursor protein that participates in the secretion of the polypeptide. A promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence. A
30 ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and being in the reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at suitable restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or
35 linkers are used in accordance with conventional practice. Furthermore, linked nucleic acids may be produced by the above-mentioned overlap extension PCR technique.

"Ligation" refers to the process of forming phosphodiester bonds between two nucleic acid fragments. For ligation of the two fragments, the ends of the fragments must be compatible with each other. In some cases, the ends will be directly compatible after endonuclease digestion. However, it may be necessary first to convert the cohesive ends commonly produced after endonuclease digestion to blunt ends to make them compatible for ligation. For blunting the ends, the DNA is treated in a suitable buffer for at least 15 minutes at 15°C with about 10 units of the Klenow fragment of DNA polymerase I or T4 DNA polymerase in the presence of the four deoxyribonucleotide triphosphates. The DNA is then purified by phenol-chloroform extraction and ethanol precipitation, or by silica purification. The DNA fragments that are to be ligated together are put in solution in equimolar amounts. The solution will contain ATP, ligase buffer, and a ligase such as T4 DNA ligase at about 10 units per 0.5 µg of DNA. If the DNA is to be ligated to a vector, the vector is first linearized by digestion with appropriate restriction endonucleases. The linearized fragment is then treated with bacterial alkaline phosphatase or calf intestinal phosphatase to prevent self-ligation of the fragment during the ligation step.

In the production methods of the present invention, antigen-binding domains or antibodies having higher antigen-binding activity under a high-calcium-ion-concentration condition than under a low-calcium-ion-concentration condition, which have been selected by the method described in the above-mentioned section on "ion concentration conditions" are isolated. Furthermore, antigen-binding domains or antibodies having higher antigen-binding activity in a neutral pH range condition than in an acidic pH range condition, which have been selected by the method described in the above-mentioned section on "ion concentration conditions" are isolated. For example, when antigen-binding domains isolated in this manner are selected from a library, as described later in the Examples, polynucleotides encoding the antigen-binding domains are isolated by conventional gene amplification from viruses such as phages. Alternatively, when antigen-binding domains or antibodies isolated in this manner are those selected from cultures of cells such as hybridomas, as indicated in the aforementioned section on antibodies, antibody genes and such are isolated by conventional gene amplification from those cells.

Next, a polynucleotide encoding an antigen-binding domain isolated as described above is linked in frame to a polynucleotide encoding an Fcγ-receptor-binding domain having human-FcRn-binding activity in an acidic pH range, and has higher binding activity to the Fcγ receptor in a neutral pH range condition than a native Fcγ-receptor-binding domain in which the sugar chain bound at position 297 (EU numbering) is a fucose-containing sugar chain. Suitable example of the Fcγ-receptor-binding domain includes an antibody Fc region as described in the above-mentioned section on Fcγ-receptor-binding domain. Furthermore, examples of the Fcγ

receptor include FcγRIa, FcγRIIa(R), FcγRIIa(H), FcγRIIb, FcγRIIIa(V), or FcγRIIIa(F).

Suitable examples of the antibody Fc region include Fc regions having at least one or more amino acids, selected from the group consisting of amino acids at positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 in the Fc region site according to EU numbering, that are different from the amino acids at corresponding sites in the native Fc region. A suitable example of the native Fc region is an Fc region of any one of IgG1, IgG2, IgG3, and IgG4.

In a non-limiting embodiment, a suitable example of the antibody Fc region is an Fc region comprising at least one or more amino acids selected from the group consisting of:

- either Lys or Tyr at amino acid position 221;
- any one of Phe, Trp, Glu, and Tyr at amino acid position 222;
- any one of Phe, Trp, Glu, and Lys at amino acid position 223;
- any one of Phe, Trp, Glu, and Tyr at amino acid position 224;
- any one of Glu, Lys, and Trp at amino acid position 225;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 227;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 228;
- any one of Ala, Glu, Gly, and Tyr at amino acid position 230;
- any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;
- any one of Glu, Gly, Lys, and Tyr at amino acid position 232;
- any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 233;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 234;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 235;
- any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 236;
- any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 237;
- any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 238;

- any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 239;
- any one of Ala, Ile, Met, and Thr at amino acid position 240;
- any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;
- 5 any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;
- His at amino acid position 244;
- Ala at amino acid position 245;
- any one of Asp, Glu, His, and Tyr at amino acid position 246;
- any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;
- 10 any one of Glu, His, Gln, and Tyr at amino acid position 249;
- either Glu or Gln at amino acid position 250;
- Phe at amino acid position 251;
- any one of Phe, Met, and Tyr at amino acid position 254;
- any one of Glu, Leu, and Tyr at amino acid position 255;
- 15 any one of Ala, Met, and Pro at amino acid position 256;
- any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;
- any one of Asp, Glu, His, and Tyr at amino acid position 260;
- any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;
- any one of Ala, Ile, Met, and Thr at amino acid position 263;
- 20 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 264;
- any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 265;
- any one of Ala, Ile, Met, and Thr at amino acid position 266;
- 25 any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 267;
- any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid position 268;
- any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 269;
- 30 any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 270;
- any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 271;
- 35 any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 272;

- either Phe or Ile at amino acid position 273;
 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at
 amino acid position 274;
 either Leu or Trp at amino acid position 275;
- 5 any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 276;
 any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at
 amino acid position 278;
 Ala at amino acid position 279;
- 10 any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;
 any one of Asp, Lys, Pro, and Tyr at amino acid position 281;
 any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;
 any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;
 any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;
- 15 any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;
 any one of Glu, Gly, Pro, and Tyr at amino acid position 286;
 any one of Asn, Asp, Glu, and Tyr at amino acid position 288;
 any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;
 any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;
- 20 any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;
 any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid
 position 293;
 any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 294;
- 25 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at
 amino acid position 295;
 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino
 acid position 296;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 297;
- 30 any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino
 acid position 298;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and
 Tyr at amino acid position 299;
- 35 any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp
 at amino acid position 300;

- any one of Asp, Glu, His, and Tyr at amino acid position 301;
 Ile at amino acid position 302;
 any one of Asp, Gly, and Tyr at amino acid position 303;
 any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;
 5 any one of Glu, Ile, Thr, and Tyr at amino acid position 305;
 any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;
 Phe at amino acid position 313;
 Leu at amino acid position 315;
 either Glu or Gln at amino acid position 317;
 10 any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;
 any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid
 position 320;
 any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position
 322;
 15 Ile at amino acid position 323;
 any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid
 position 324;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 Tyr at amino acid position 325;
 20 any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino
 acid position 326;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr
 at amino acid position 327;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and
 25 Tyr at amino acid position 328;
 any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 329;
 any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr
 at amino acid position 330;
 30 any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position
 331;
 any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp,
 and Tyr at amino acid position 332;
 any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid
 35 position 333;
 any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;

- any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid position 335;
 any one of Glu, Lys, and Tyr at amino acid position 336;
 any one of Glu, His, and Asn at amino acid position 337;
 5 any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;
 either Ala or Val at amino acid position 376;
 either Gly or Lys at amino acid position 377;
 Asp at amino acid position 378;
 Asn at amino acid position 379;
 10 any one of Ala, Asn, and Ser at amino acid position 380;
 either Ala or Ile at amino acid position 382;
 Glu at amino acid position 385;
 Thr at amino acid position 392;
 Leu at amino acid position 396;
 15 Lys at amino acid position 421;
 Asn at amino acid position 427;
 either Phe or Leu at amino acid position 428;
 Met at amino acid position 429;
 Trp at amino acid position 434;
 20 Ile at amino acid position 436; and
 any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440;
 in the Fc region site according to EU numbering.

Furthermore, for Fc regions of the present invention, Fc regions which have binding activity or enhanced binding activity to FcRn in an acidic pH range condition may be suitably
 25 used. Examples of such Fc regions include Fc regions of IgG-type immunoglobulins such as Fc regions of human IgG (IgG1, IgG2, IgG3, IgG4, and variants thereof). For variants having alterations to other amino acids, Fc regions with amino acid alterations at any position may be used as long as there is FcRn-binding activity in an acidic pH range or the binding activity to human FcRn in an acidic pH range condition can be increased, and when an antigen-binding
 30 molecule includes an Fc region of a human IgG1 as the Fc region, it preferably includes an alteration that enhances binding to FcRn in an acidic pH range condition compared to the binding activity of the starting-material Fc region of human IgG1. Suitable examples of amino acids that can be altered include the amino acids at positions 238, 252, 253, 254, 255, 256, 265, 272, 286, 288, 303, 305, 307, 309, 311, 312, 317, 340, 356, 360, 362, 376, 378, 380, 382, 386,
 35 388, 400, 413, 415, 424, 433, 434, 435, 436, 439, and/or 447 (EU numbering) as described in WO2000/042072. Similarly, suitable examples of amino acids that can be altered also include

the amino acids at positions 251, 252, 254, 255, 256, 308, 309, 311, 312, 385, 386, 387, 389, 428, 433, 434, and/or 436 (EU numbering) as described in WO2002/060919. Furthermore, examples of amino acids that can be altered also include the amino acids at positions 250, 314, and 428 (EU numbering) as described in WO2004/092219. Other suitable examples of amino acids that can be altered also include the amino acids at positions 251, 252, 307, 308, 378, 428, 430, 434, and/or 436 (EU numbering) as described in WO2010/045193. Fc regions produced by enhancing the FcRn-binding in an acidic pH range of an IgG immunoglobulin Fc region by the amino acid alterations may be used in the production methods of the present invention.

Furthermore, as described later, for the Fc region included in an antigen-binding molecule of the present invention, an Fc region having FcRn-binding activity in a neutral pH range may also be suitably used. Such an Fc region may be obtained by any method according to the aforementioned method for obtaining Fc regions having FcRn-binding activity in an acidic pH range. Specifically, an Fc region containing an FcRn-binding domain having binding activity or having enhanced binding activity to FcRn in a neutral pH range due to alteration of amino acids of the Fc region of a human IgG immunoglobulin used as the starting-material Fc region may be obtained. Favorable Fc regions of IgG immunoglobulins for the alteration include Fc regions of human IgG (IgG1, IgG2, IgG3, IgG4, and variants thereof). For variants having alterations to other amino acids, Fc regions with amino acid alterations at any position may be used as long as there is FcRn-binding activity in a neutral pH range or the binding activity to human FcRn in a neutral pH range can be increased. When an antigen-binding molecule includes an Fc region of a human IgG1 as the Fc region, it preferably includes an alteration that enhances binding to FcRn in a neutral pH range compared to the binding activity of the starting-material Fc region of human IgG1. Suitable examples of such altered Fc regions include human Fc regions having at least one or more amino acids, selected from the group consisting of amino acids at positions 237, 238, 239, 248, 250, 252, 254, 255, 256, 257, 258, 265, 270, 286, 289, 297, 298, 303, 305, 307, 308, 309, 311, 312, 314, 315, 317, 325, 332, 334, 360, 376, 380, 382, 384, 385, 386, 387, 389, 424, 428, 433, 434, and 436 in the starting-material Fc region site according to EU numbering, that are different from the corresponding amino acids in the native Fc region.

Suitable examples of such altered Fc regions include Fc regions containing at least one amino acid selected from the group consisting of:

Met at amino acid position 237;

Ala at amino acid position 238;

Lys at amino acid position 239;

Ile at amino acid position 248;

any one of Ala, Phe, Ile, Met, Gln, Ser, Val, Trp, and Tyr at amino acid position 250;

- any one of Phe, Trp, and Tyr at amino acid position 252;
Thr at amino acid position 254;
Glu at amino acid position 255;
any one of Asp, Glu, and Gln at amino acid position 256;
- 5 any one of Ala, Gly, Ile, Leu, Met, Asn, Ser, Thr, and Val at amino acid position 257;
His at amino acid position 258;
Ala at amino acid position 265;
Phe at amino acid position 270;
either Ala or Glu at amino acid position 286;
- 10 His at amino acid position 289;
Ala at amino acid position 297;
Gly at amino acid position 298;
Ala at amino acid position 303;
Ala at amino acid position 305;
- 15 any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr
at amino acid position 307;
any one of Ala, Phe, Ile, Leu, Met, Pro, Gln, and Thr at amino acid position 308;
any one of Ala, Asp, Glu, Pro, and Arg at amino acid position 309;
any one of Ala, His, and Ile at amino acid position 311;
- 20 either Ala or His at amino acid position 312;
either Lys or Arg at amino acid position 314;
either Ala or His at amino acid position 315;
Ala at amino acid position 317;
Gly at amino acid position 325;
- 25 Val at amino acid position 332;
Leu at amino acid position 334;
His at amino acid position 360;
Ala at amino acid position 376;
Ala at amino acid position 380;
- 30 Ala at amino acid position 382;
Ala at amino acid position 384;
either Asp or His at amino acid position 385;
Pro at amino acid position 386;
Glu at amino acid position 387;
- 35 either Ala or Ser at amino acid position 389;
Ala at amino acid position 424;

any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino acid position 428;

Lys at amino acid position 433;

any one of Ala, Phe, His, Ser, Trp, and Tyr at amino acid position 434; and

5 His at amino acid position 436;

in the Fc region according to EU numbering.

For example, by using the amino acid alterations individually, or by using more than one of them in combination, FcRn-binding of an IgG Fc region in an acidic and/or neutral pH range can be enhanced, and the amino acid alterations that are introduced are not particularly
10 limited, and as long as the retentivity in plasma is improved, any amino acid alteration may be introduced.

An antigen-binding molecule of the present invention is isolated from the culture of cells transformed by a desired expression vector carrying an operably linked polynucleotide obtained by linking, as described above, a polynucleotide encoding the antigen-binding domain
15 to a polynucleotide encoding an Fcγ receptor-binding domain having human-FcRn-binding activity in an acidic pH range and having higher binding activity to the Fcγ receptor in a neutral pH range condition than a native Fcγ receptor-binding domain in which the sugar chain bound at position 297 (EU numbering) is a fucose-containing sugar chain. Antigen-binding molecules of the present invention are produced using a method according to the method for producing
20 antibodies described in the above-mentioned section on antibodies.

Herein below, the present invention will be specifically described with reference to the Examples, but it is not to be construed as being limited thereto.

25 [Examples]

[Example 1] Preparation of antigen-binding molecules whose mouse FcγR-binding activity under a neutral pH range condition is higher than the binding activity of native human IgG Fc region
(1-1) pH-dependent human IL-6 receptor-binding antibodies

H54/L28-IgG1 which comprises H54-IgG1 (SEQ ID NO: 36) and L28-CK (SEQ ID
30 NO: 37) described in WO2009/125825 is a humanized anti-IL-6 receptor antibody. Meanwhile, Fv4-IgG1 which comprises VH3-IgG1 (SEQ ID NO: 38) and VL3-CK (SEQ ID NO: 39) is a humanized anti-IL-6 receptor antibody resulting from conferring, to H54/L28-IgG1, the property of binding to soluble human IL-6 receptor in a pH-dependent manner (which binds at pH 7.4 and dissociates at pH 5.8). The *in vivo* mouse test described in WO2009/125825 demonstrated that,
35 in the group administered with a mixture of Fv4-IgG1 and soluble human IL-6 receptor as the antigen, the elimination of soluble human IL-6 receptor from plasma was significantly

accelerated as compared to the group administered with a mixture of H54/L28-IgG1 and soluble human IL-6 receptor as the antigen.

The soluble human IL-6 receptor bound to H54/L28-IgG1, which is an antibody that binds to a soluble human IL-6 receptor, is, together with the antibody, recycled to plasma by FcRn. Meanwhile, Fv4-IgG1, which is an antibody that binds to a soluble human IL-6 receptor in a pH dependent manner, dissociates soluble human IL-6 receptor under the acidic condition in the endosome. The dissociated soluble human IL-6 receptor is degraded in the lysosomes, thus this enables considerable acceleration of the elimination of soluble human IL-6 receptor. Furthermore, after binding to FcRn in the endosome, Fv4-IgG1, which is an antibody that binds to a soluble human IL-6 receptor in a pH dependent manner, is recycled to the plasma. Since the recycled antibody can bind to soluble human IL-6 receptor again, the antibody repeatedly binds to the antigen (soluble human IL-6 receptor) and is recycled by FcRn to the plasma. It is thought that, as a result, a single antibody molecule can bind repeatedly several times to soluble human IL-6 receptor (WO 2009/125825).

(1-2) Preparation of an anti-human IL-6 receptor antibody with enhanced mouse FcγR binding and anti-human IL-6 receptor antibody without mouse FcγR binding

VH3-IgG1-F1022 (SEQ ID NO: 40), an antigen-binding molecule with enhanced mouse FcγR binding, was prepared by substituting Asp for Lys at position 326 (EU numbering) and Tyr for Leu at position 328 (EU numbering) in VH3-IgG1. Fv4-IgG1-F1022 containing VH3-IgG1-F1022 as the heavy chain and VL3-CK as the light chain was produced using the method described in Reference Example 2.

Meanwhile, VH3-IgG1-F760 (SEQ ID NO: 41), an antigen-binding molecule without mouse FcγR binding, was prepared by substituting Arg for Leu at position 235 and Lys for Ser at position 239 (EU numbering) in VH3-IgG1. Fv4-IgG1-F760 containing VH3-IgG1-F760 as the heavy chain and VL3-CK as the light chain was produced using the method described in Reference Example 2.

(1-3) Assessment of mouse FcγR-binding activity

VH3/L(WT)-IgG1, VH3/L(WT)-IgG1-F1022, and VH3/L(WT)-IgG1-F760, which contain VH3-IgG1, VH3-IgG1-F1022, and VH3-IgG1-F760 as the heavy chain, and L(WT)-CK (SEQ ID NO: 42) as the light chain, were produced using the method described in Reference Example 2. These antibodies were kinetically analyzed for their mouse FcγR binding as described below.

(1-4) Kinetic analysis of mouse FcγR binding

The binding of antibodies to mouse FcγRI, FcγRIIb, FcγRIII, and FcγRIV (hereinafter, referred to as mouse FcγRs) (prepared by Reference Example 26) was kinetically analyzed using Biacore T100 and T200 (GE Healthcare). An appropriate amount of protein L (ACTIGEN) was immobilized onto a Sensor chip CM4 (GE Healthcare) by an amino coupling method, and antibodies of interest were captured thereto. Then, diluted solutions of mouse FcγRs and a running buffer as a blank were injected, and the mouse FcγRs were allowed to interact with antibodies captured onto the sensor chip. The running buffer used was 20 mmol/l ACES, 150 mmol/l NaCl, 0.05% (w/v) Tween™20, pH 7.4. This buffer was also used to dilute the mouse FcγRs. The sensor chip was regenerated using 10 mmol/l glycine-HCl, pH 1.5. All measurements were carried out at 25°C. The binding rate constant k_a (1/Ms) and dissociation rate constant k_d (1/s), which are kinetic parameters, were calculated from the sensorgrams obtained by the measurement. K_D (M) of each antibody for human FcγR was calculated based on the values. Each parameter was calculated using Biacore T100 or T200 Evaluation Software (GE Healthcare).

The result shown in Table 7 was obtained by the measurement. VH3/L (WT)-IgG1-F1022 was demonstrated to have increased binding activity to mFcγRI, mFcγRIIb, and mFcγRIII as compared to VH3/L (WT)-IgG1. Regarding VH3/L (WT)-IgG1-F760, the binding to the various mouse FcγRs was undetectable, demonstrating that VH3/L (WT)-IgG1-F760 lacks the binding activity to the various mouse FcγRs.

[Table 7]

VARIANT NAME	KD (M)			
	mFc γ R I	mFc γ R IIb	mFc γ R III	mFc γ R IV
IgG1	6.0E-08	5.0E-07	2.2E-07	2.4E-08
F1022	9.1E-09	8.5E-09	8.1E-09	3.8E-08
F760	NOT DETECTED	NOT DETECTED	NOT DETECTED	NOT DETECTED

(1-5) Preparation of antibodies with low fucose content

Known methods for increasing the FcγR-binding activity of antibodies include methods for making sugar chains linked to an antibody be sugar chains with low fucose content (J. Biol. Chem. (2003) 278, 3466-3473) in addition to methods for introducing an amino acid alteration into the Fc region of an antibody. An Fv4-IgG1 with low fucose content (hereinafter, abbreviated as Fv4-IgG1-Fuc) was produced by expressing Fv4-IgG1 using fucose transporter gene-deficient CHO cells (WO2006067913) as host cells according to the method described in Reference Example 4. It has been reported that, of the mFcγRs (mouse Fcγ receptors), antibodies with low fucose content have selectively increased FcγRIV-binding activity (Science,

2005, 310 (5753) 1510-1512).

[Example 2] Effect of eliminating antigens from plasma by antigen-binding molecules whose FcγR-binding activity is higher than the binding activity of native human IgG Fc region

5 (2-1) Effect of H54/L28-IgG1 and Fv4-IgG1 to eliminate antigens from plasma

H54/L28-IgG1, which is an anti-human IL-6 receptor antibody, and Fv4-IgG1 having the property of binding to human IL-6 receptor in a pH-dependent manner were produced by the method described in Reference Example 1. *In vivo* infusion tests were carried out using the produced H54/L28-IgG1 and Fv4-IgG1 by the method described below.

10

(2-1-1) *In vivo* infusion tests using human FcRn transgenic mice

An animal model in which the soluble human IL-6 receptor concentration is maintained constant in plasma was created by implanting an infusion pump (MINI-OSMOTIC PUMP MODEL2004, alzet) containing soluble human IL-6 receptor under the skin on the back of human FcRn transgenic mice (B6.mFcRn-/-, hFcRn Tg line 32 +/+ mouse, Jackson Laboratories, Methods Mol Biol. (2010) 602, 93-104). The *in vivo* dynamics after administration of an anti-human IL-6 receptor antibody was assessed in the animal model. To suppress the production of neutralizing antibodies against soluble human IL-6 receptor, an anti-mouse CD4 monoclonal antibody (prepared by a known method) was administered once at 20 mg/kg into the caudal vein. Then, an infusion pump containing 92.8 μg/ml soluble human IL-6 receptor was subcutaneously implanted on the back of the mice. Three days after implantation of the infusion pump, an anti-human IL-6 receptor antibody was administered once at 1 mg/kg into the caudal vein. The blood was collected from the mice 15 minutes, seven hours, one day, two days, four days, and seven days after administration of the anti-human IL-6 receptor antibody. Immediately, the collected blood was centrifuged at 15,000 rpm and 4°C for 15 minutes to prepare plasma. The isolated plasma was stored in a freezer set at -20°C or below until use.

(2-1-2) Determination of the human IL-6 receptor (hsIL-6R) concentration in plasma by an electrochemiluminescent method

30 The human IL-6 receptor concentrations in mouse plasma were determined by an electrochemiluminescent method. hsIL-6R standard curve samples prepared at 2000, 1000, 500, 250, 125, 62.5, and 31.25 pg/ml and assay samples of mouse plasma diluted 50 times or more were mixed with Monoclonal Anti-human IL-6R Antibody (R&D), Biotinylated Anti-human IL-6 R Antibody (R&D), Tocilizumab, which had been ruthenated with SULFO-TAG NHS Ester (Meso Scale Discovery). The mixtures were incubated at 37°C overnight. Tocilizumab was prepared at a final concentration of 333 μg/ml. Then, the reaction mixtures were aliquoted in

35

an MA400 PR Streptavidin Plate (Meso Scale Discovery). The solution reacted at room temperature for one hour was washed out, and then Read Buffer T (x4) (Meso Scale Discovery) was aliquoted. Immediately thereafter, the measurement was carried out using SECTOR PR 400TM Reader (Meso Scale Discovery). The concentration of human IL-6 receptor was
 5 determined based on the response of the standard curve using analysis software SOFTmax PRO (Molecular Devices).

A time course of the monitored human IL-6 receptor concentration is shown in Fig. 2. As compared to H54/L28-IgG1, Fv4-IgG1 that binds to human IL-6 receptor in a pH-dependent manner could reduce the human IL-6 receptor concentration, but could not reduce it below the
 10 baseline without antibody administration. That is, the administered antibody which binds to an antigen in a pH-dependent manner could not reduce the antigen concentration in plasma below the level prior to antibody administration.

(2-2) The effect of eliminating an antigen from plasma by an antibody with increased or reduced
 15 FcγR-binding activity

Whether the time course of human IL-6 receptor concentration is influenced by increasing or reducing the FcγR-binding activity of Fv4-IgG1, which is a pH-dependent human IL-6 receptor-binding antibody, was assessed by the method described below. Using Fv4-IgG1, Fv4-IgG1-F760, Fv4-IgG1-F1022, and Fv4-IgG1-Fuc prepared as described in Example 1, *in*
 20 *vivo* infusion tests were performed by the method described below.

(2-2-1) *In vivo* infusion tests using human FcRn transgenic mice

A animal model in which the soluble human IL-6 receptor concentration is maintained constant in plasma was created by implanting an infusion pump (MINI-OSMOTIC PUMPTM MODEL2004, alzet) containing soluble human IL-6 receptor under the skin on the back of
 25 human FcRn transgenic mice (B6.mFcRn-/-hFcRn Tg line 32 +/- mouse, Jackson Laboratories, Methods Mol Biol. (2010) 602, 93-104). In the animal model, an anti-human IL-6 receptor antibody was administered simultaneously with Sanglopor (CSL Behring) which is a human immunoglobulin preparation, to assess the *in vivo* dynamics of the soluble human IL-6 receptor
 30 after antibody administration. To suppress the production of neutralizing antibodies against soluble human IL-6 receptor, an anti-mouse CD4 monoclonal antibody (prepared by a known method) was administered once at 20 mg/kg into the caudal vein. Then, an infusion pump containing 92.8 μg/ml soluble human IL-6 receptor was subcutaneously implanted on the back of the mice. Three days after implantation of the infusion pump, an anti-human IL-6 receptor
 35 antibody and SangloporTM were administered once at 1 mg/kg and 1000 mg/kg, respectively, into the caudal vein. The blood was collected from the mice 15 minutes, seven hours, one day, two

days, four days, and seven days after administration of the anti-human IL-6 receptor antibody. The blood was collected from the mice 15 minutes, seven hours, one day, two days, three days, and seven days after administration of the anti-human IL-6 receptor antibody. Immediately, the collected blood was centrifuged at 15,000 rpm and 4°C for 15 minutes to prepare the plasma.

- 5 The isolated plasma was stored in a freezer set at -20°C or below until use.

(2-2-2) Determination of the soluble human IL-6 receptor (hsIL-6R) concentration in plasma by an electrochemiluminescent method

- 10 The hsIL-6R concentrations in mouse plasma were determined by the same electrochemiluminescent method as described in (2-1-2).

- 15 The result is shown in Fig. 3. The time course of human IL-6 receptor concentration in plasma of mice administered with Fv4-IgG1-F760, from which the mouse FcγR binding of Fv4-IgG1 is deleted, was demonstrated to be comparable to that in mice administered with Fv4-IgG1. The cytotoxic activity to a membrane antigen depends on the FcγR binding, and thus the cytotoxic activity is lost when eliminating the FcγR binding. On the other hand, even when administering an antibody, from which mouse FcγR binding is deleted, against human IL-6 receptor which is a soluble antigen, there was no effect on the time course of human IL-6 receptor concentration in the plasma of the administered mice. Thus, it would be thought that the FcγR binding of an antibody against the soluble antigen has no contribution to the time course of antigen concentration in the plasma of mice administered with the antibody.

- 20 Surprisingly, however, the human IL-6 receptor concentration in the plasma of mice administered with Fv4-IgG1-F1022 with enhanced mouse FcγR binding was considerably reduced as compared to the human IL-6 receptor concentration in the plasma of mice administered with Fv4-IgG1. As to the degree of reduction, the concentration was confirmed to be decreased below the baseline human IL-6 receptor concentration without antibody administration. In particular, the human IL-6 receptor concentration in the plasma of mice administered with Fv4-IgG1-F1022 was reduced down to about 1/100 three days after administration as compared to the case of Fv4-IgG1 administration. This finding demonstrates that, by administering to mice an antibody that binds to human IL-6 receptor in a pH-dependent manner and whose FcγR binding has been enhanced, the human IL-6 receptor concentration in the plasma of the mice can be significantly reduced, and as to the degree of reduction, the antigen concentration in plasma can be reduced below the level before antibody administration.

- 30 Furthermore, it was also demonstrated that, as compared to mice administered with Fv4-IgG1, the human IL-6 receptor concentration in plasma was reduced in mice administered with Fv4-IgG1-Fuc which has sugar chains with low fucose content and with increased mouse FcγR IV-binding activity. In particular, the human IL-6 receptor concentration in the plasma of

mice administered with Fv4-IgG1-Fuc was reduced down to about 1/2 seven days after administration as compared to the case of Fv4-IgG1 administration. The above finding demonstrates that, by administering to mice a pH-dependent antigen-binding molecule that binds to human IL-6 receptor in a pH-dependent manner and whose Fc γ R binding has been enhanced, the soluble antigen concentration in the plasma of the mice can be reduced. In this case, methods for enhancing the Fc γ R binding are not particularly limited to introduction of amino acid alterations. It was demonstrated that such enhancement can be achieved, for example, by using a human IgG Fc region to which a sugar chain with low fucose content is linked at position 297 (EU numbering); however, the effect of Fv4-IgG1-Fuc to reduce antigen concentration was smaller than Fv4-F1022. Based on this result, it would be thought that, of several Fc γ Rs (Fc γ RI, II, III, and IV for mouse), mFc γ IV, to which the binding of Fv4-IgG1-Fuc is enhanced, does not have a large contribution to the reduction of antigen concentration as an Fc γ R.

Thus, it was revealed that, by administering to an individual an antibody that binds to a soluble antigen in a pH-dependent manner and whose Fc γ R binding has been enhanced, the soluble antigen concentration in the plasma of the individual can be markedly reduced.

Without being bound by a particular theory, the unexpected reduction of soluble antigen concentration in plasma, which was observed when administering an antigen-binding molecule that comprises an antigen-binding domain whose Fc γ R binding has been enhanced and whose antigen-binding activity is altered depending on the ion concentration condition such as pH and an FcRn-binding domain that has FcRn-binding activity under an acidic pH range condition, can be explained as follows.

IgG antibodies that are non-specifically incorporated into cells return to the cell surface by binding to FcRn under the acidic condition in the endosome, and then dissociate from FcRn under the neutral condition in plasma. In such a case, when an antibody that neutralizes the function of a soluble antigen by binding to the antigen is administered to mice in which the concentration of the soluble antigen is maintained constant in plasma, the soluble antigen in plasma forms a complex with the antibody administered. The soluble antigen incorporated into cells while remaining as the complex is thought to be recycled, in a state bound to the antibody, to the plasma together with the antibody, because the Fc region of the antibody binds to FcRn under the acidic condition in the endosome.

Meanwhile, when the antibody against the soluble antigen is an antibody that binds to the antigen in a pH-dependent manner (*i.e.*, an antibody that dissociates the soluble antigen under the acidic condition in the endosome), the soluble antigen that is non-specifically incorporated into cells while remaining as a complex with the antibody, is dissociated from the antibody in the endosome and degraded in the lysosome; thus, the soluble antigen is not recycled to the plasma. That is, it is thought that Fv4-IgG1 incorporated as a complex with the soluble antigen into cells

can dissociate the soluble antigen in the endosome and thus accelerate the elimination of the soluble antigen.

As described above, antigen-binding molecules such as Fv4-IgG1, which contain an antigen-binding domain whose antigen-binding activity is altered depending on the ion concentration, are thought to be capable of binding to antigens repeatedly several times. The effect to accelerate the elimination of soluble antigens from the plasma by dissociating them in the endosome is thought to depend on the rate of incorporation of the antigen/antigen-binding molecule complex into the endosome. An antigen-binding molecule that contains an antigen-binding domain whose binding activity to various FcγRs has been increased and whose antigen-binding activity is altered depending on the condition of ion concentration, is actively incorporated into cells by binding to various FcγRs expressed on the cell membrane, and can be shuttled back to plasma by recycling *via* the binding between FcRn and the FcRn-binding domain comprised in the molecule, which has FcRn-binding activity under an acidic pH range condition. That is, it is thought that, since the above antigen-binding molecule which forms a complex with a soluble antigen in plasma is actively incorporated into cells *via* FcγR expressed on the cell membrane, its effect to accelerate the elimination of the soluble antigen from plasma is more markedly shown than antigen-binding molecules whose binding activity to various FcγRs has not been increased.

In the living organism, various FcγRs are expressed on the cell membrane of immune cells, and play a variety of functions. Any FcγRs may be used to incorporate antibodies into cells. Specifically, in human, the presence of inhibitory FcγRIIb, and activating FcγRs including FcγRI, FcγRIIa, and FcγRIIIa is known, and antibodies may be incorporated by any of them. Antibodies may be incorporated by all or any one of the FcγRs. Alternatively, antibodies may be incorporated in such a manner mediated by various activating FcγRs alone, or by inhibitory FcγRIIb alone.

On the other hand, to achieve the above purposes, it is possible to employ any methods for increasing the FcγR-binding activity of the FcγR-binding domain of antigen-binding molecules. For example, as shown in Example 1, amino acid mutations for increasing the FcγR-binding activity may be introduced into the FcγR-binding domain of antigen-binding molecules, or one can use low-fucose-type antibodies. Meanwhile, the effect to increase the FcγR-binding activity, which is achieved by such methods, may be an effect of augmenting the binding to any FcγR. Specifically, it is possible to increase the binding activity to any one, some, or all of the FcγRs. Furthermore, it is possible to only increase the binding activity to various activating FcγRs, or inhibitory FcγRIIb.

The FcγR-binding activity of an antibody that binds to a membrane antigen plays an

important role in the cytotoxic activity of the antibody. Thus, when it is necessary for an antibody used as a pharmaceutical agent to have cytotoxic activity, a human IgG1 isotype with strong FcγR-binding activity is used. In addition, techniques to enhance the cytotoxic activity of such antibodies by increasing the FcγR-binding activity of the antibodies are used commonly in the art.

Meanwhile, the role of the FcγR-binding activity of antibodies that bind to soluble antigens and which are used as pharmaceutical agents has not been known in the art. There has been no sufficient assessment on what difference in the effect on the living organism administered with the antibodies is caused by the difference in the FcγR-binding activity between human IgG1 with high FcγR-binding activity and human IgG2 and human IgG4 with low FcγR-binding activity. Actually, it was demonstrated in the present Example that there was no influence on the time course of soluble antigen concentration in the plasma of the individuals administered with an antibody that lacks FcγR-binding activity. Meanwhile, in the present invention, it was revealed that the soluble antigen concentration was significantly reduced in the plasma of the individuals administered with an antigen-binding molecule whose FcγR-binding activity has been increased and which contains an antigen-binding domain whose soluble antigen-binding activity is altered depending on the ion concentration condition. Specifically, it can be said that the present inventors revealed for the first time the benefit of the enhancement of FcγR binding by combining an FcRn-binding domain that has FcRn-binding activity under an acidic pH range condition with an antigen-binding domain whose soluble antigen binding is altered depending on the ion concentration condition, comprised in an antigen-binding molecule targeted to a soluble antigen.

[Example 3] Effect of eliminating antigens from plasma by antigen-binding molecules whose FcγR-binding activity is greater than that of native human IgG Fc region and whose human FcRn-binding activity has been increased under an acidic pH range condition

(3-1) Preparation of antigen-binding molecules whose FcγR-binding activity is greater than the binding activity of native human IgG Fc region and whose human FcRn-binding activity has been increased under an acidic pH range condition

A reported method for improving the retention of IgG antibody in plasma is to improve the FcRn binding under an acidic pH range condition. It is thought that, when the FcRn binding under an acidic pH range condition is improved by introducing an amino acid substitution into the Fc region of an IgG antibody, this increases the recycling efficiency from the endosome to plasma, resulting in an improvement of the plasma retention of the IgG antibody.

There are many reports on amino acid alterations to improve the plasma retention by improving the human FcRn-binding activity under an acidic pH range condition. Such

alterations include, for example:

the method for substituting Leu for Met at position 428 and Ser for Asn at position 434 (EU numbering) in an IgG antibody (Nat. Biotechnol. (2010) 28, 157-159); the method for substituting Ala for Asn at position 434 (Drug. Metab. Dispos. (2010) 38 (4), 600-605); the method for substituting Tyr for Met at position 252, Thr for Ser at position 254, and Glu for Thr at position 256 (J. Biol. Chem. (2006) 281, 23514-23524); the method for substituting Gln for Thr at position 250 and Leu for Met at position 428 (J. Immunol. (2006) 176 (1) 346-356); the method for substituting His for Asn at position 434 (Clin. Pharm. & Ther. (2011) 89 (2) 283-290.); and WO2010/106180; WO2010/045193; WO2009/086320; WO2009/058492; WO2008/022152; WO2006/050166, WO2006/053301, WO2006/031370; WO2005/123780; WO2005/047327; WO2005/037867; WO2004/035752; and WO2002/060919.

VH3-IgG1-F1093 (SEQ ID NO: 43) with a substitution of Leu for Met at position 428 and Ser for Asn at position 434 (EU numbering) in VH3-IgG1-F1022 was prepared to improve the pharmacodynamics of Fv4-IgG1-F1022 that was demonstrated to produce, when administered, the effect of significantly reducing the soluble antigen concentration in plasma, as described in Example 2. Fv4-IgG1-F1093 comprising VH3-IgG1-F1093 as the heavy chain and VL3-CK as the light chain was constructed using the method described in Reference Example 2.

(3-2) Effect of eliminating antigens from plasma by antigen-binding molecules whose FcγR-binding activity is greater than that of native human IgG Fc region and whose human FcRn-binding activity has been increased under an acidic pH range condition

An *in vivo* infusion test was carried out for Fv4-IgG1-F1093 by the same method as described in Example (2-1-1) using human FcRn transgenic mice in which the soluble human IL-6 receptor concentration is maintained constant in plasma. Soluble human IL-6 receptor concentrations in the plasma of the mice were determined by the method described in Example (2-1-2). The result is shown in Fig. 4.

(3-2-1) Determination of the anti-human IL-6 receptor antibody concentration in plasma by the ELISA method

Anti-human IL-6 receptor antibody concentrations in mouse plasma were determined by the ELISA method. First, an anti-Fv4 MABTECH idotype antibody was aliquoted in a Nunc-Immuno Plate, MaxiSoup (Nalge nunc International). The plate was allowed to stand at 4°C overnight to prepare a plate immobilized with the anti-Fv4 idotype antibody. The idotype antibody was obtained by immunizing a rabbit with Fv4-M73 (WO2009/125825). After purifying the serum using an ion-exchange resin, the antibody was affinity-purified by a

column immobilized with Fv4-M73, followed by adsorption using an immobilized column for human. Standard curve samples containing an anti-human IL-6 receptor antibody (concentration in plasma: 0.8, 0.4, 0.2, 0.1, 0.05, 0.025, and 0.0125 $\mu\text{g/ml}$) and assay samples of mouse plasma diluted 100 times or more were prepared. 100 μl each of the standard curve and assay samples were combined with 200 μl of 20 ng/ml soluble human IL-6 receptor. The resulting mixtures were allowed to stand at room temperature for one hour, and aliquoted to each well of the plate immobilized with the anti-Fv4 idotype antibody. The plate was allowed to stand at room temperature for another one hour. Then, Biotinylated Anti-human IL-6 R Antibody (R&D) was reacted thereto at room temperature for one hour. Next, Streptavidin-PolyHRP80 (Stereospecific Detection Technologies) was reacted thereto at room temperature for one hour. The chromogenic reaction of the reaction solution was performed using as a substrate TMB One Component TM HRP Microwell Substrate (BioFX Laboratories). After terminating the reaction with 1N sulfuric acid (Showa Chemical), the absorbance at 450 nm of the reaction solution of each well was measured with a microplate reader. Antibody concentrations in mouse plasma were determined based on the absorbance of the standard curve using the analysis software SOFTmax PRO (Molecular Devices).

The result is shown in Fig. 5.

(3-3) Improvement of pharmacodynamics by increasing the human FcRn-binding activity under an acidic pH range condition

As shown in Fig. 5, in the group administered with Fv4-IgG1-F1022 resulting from the enhancement of the Fc γ R-binding activity of Fv4-IgG1 under a neutral pH range condition, the plasma retention of the administered antibody was demonstrated to be reduced as compared to the group administered with Fv4-IgG1. Meanwhile, in the group administered with Fv4-IgG1-F1093 resulting from the enhancement of the human FcRn-binding activity of Fv4-IgG1-F1022 under an acidic pH range condition, the plasma retention of the administered antibody was demonstrated to be significantly improved as compared to the group administered with Fv4-IgG1-F1022.

Furthermore, as shown in Fig. 4, the time course of the soluble human IL-6 receptor concentration in the plasma of the Fv4-IgG1-F1022-administered group was equivalent to that of the Fv4-IgG1-F1093-administered group, up to three days after antibody administration. On day three after administration, as compared to the Fv4-IgG1-administered group, the soluble human IL-6 receptor concentration in plasma was reduced as much as about 100 times in both of the Fv4-IgG1-F1022 and Fv4-IgG1-F1093-administered groups. However, on day seven after antibody administration, the soluble human IL-6 receptor concentration in plasma was observed to be elevated in the Fv4-IgG1-F1022-administered group as compared to on day three after

administration. On the other hand, in the Fv4-IgG1-F1093-administered group, an increase in the plasma concentration of soluble human IL-6 receptor was not observed, showing that the effect to reduce the soluble human IL-6 receptor concentration was sustained in this administration group.

5 Specifically, Fv4-IgG1-F1093, when administered, reduced the soluble human IL-6 receptor concentration in the plasma of the administered individual down to about 1/100 as compared to Fv4-IgG1, and in addition, it sustained this condition for a long period. Thus, Fv4-IgG1-F1093 was demonstrated to be a highly excellent antigen-binding molecule. Without being bound by a particular theory, the phenomenon observed herein can be explained as follows.
10 Fv4-IgG1-F1022 in which the FcγR-binding activity of Fv4-IgG1 has been increased under a neutral pH range condition is thought to be incorporated in a large amount mainly into immune cells expressing FcγR on the cell membrane. The incorporated antibody is transferred into the endosome, and by binding to FcRn in the endosome, the antibody is recycled to the plasma. When the FcRn-binding activity of the antibody is not high enough under the condition at acidic
15 pH in the endosome, the antibody incorporated into the endosome is thought to be incapable of sufficient recycling. Specifically, a possible reason for the reduced plasma retention of Fv4-IgG1-F1022 relative to Fv4-IgG1 would be that the FcRn-binding activity under an acidic pH range condition is insufficient for sufficient recycling of the endosome-incorporated antibody to the plasma by FcRn binding, and the antibody that was not recycled was degraded in the
20 lysosome.

On the other hand, as with Fv4-IgG1-F1022, Fv4-IgG1-F1093 resulting from the enhancement of the human FcRn-binding activity of Fv4-IgG1-F1022 under an acidic pH range condition is thought to be incorporated in a large amount mainly into immune cells expressing FcγR on the cell membrane. An antibody incorporated and transferred into the endosome is
25 recycled to the plasma by binding to FcRn in the endosome. Since its human FcRn-binding activity under an acidic pH range condition is enhanced, Fv4-IgG1-F1093 is thought to have sufficient FcRn-binding activity in the endosome. Thus, after incorporation into cells, most of Fv4-IgG1-F1093 is recycled to the plasma. Thus, it would be thought that the plasma retention of Fv4-IgG1-F1093 was improved in administered individuals as compared to Fv4-IgG1-F1022.

30 On the other hand, it has been known that the plasma retention of ordinary antibodies is improved when their FcRn-binding activity is improved under an acidic pH range condition. However, it is thought that, when the antibody retention in plasma is improved, the plasma retention of antibody-bound antigens is also improved, and this results in an increase of the antigen concentration in plasma. In actual, as described in WO2010/088444, Antibody 18E
35 introduced with the alteration YTE into Antibody 18, which is a human IgG1 antibody against IL-6, to increase the FcRn-binding activity under an acidic pH range condition, showed

improved antibody retention in the plasma of cynomolgus monkeys, and at the same time, the concentration of the IL-6 antigen was also elevated in the plasma.

Surprisingly, however, when administering Fv4-IgG1-F1093 introduced with an alteration similar to YTE for increasing the FcRn-binding activity under an acidic pH range condition into Fv4-F1022 that binds to the antigen in a pH-dependent manner and has increased FcγR-binding activity, the plasma retention of the antibody was significantly improved in the administered individuals without increasing the concentration of soluble human IL-6 receptor which is the antigen. Rather, on day seven after antibody administration, the soluble human IL-6 receptor concentration remained low in the individuals administered with Fv4-IgG1-F1093 as compared to those administered with Fv4-F1022.

Without being bound by a particular theory, the phenomenon observed herein can be explained as follows. When administered to a living organism, an antibody without pH-dependent antigen binding is non-specifically incorporated into cells. Antigens that remain to be bound to the antibody are recycled to the plasma in the same extent as the antibody. Meanwhile, for an antibody with increased FcRn-binding activity under an acidic pH range condition, the extent of recycling to the plasma in a living organism administered with the antibody is higher than that of an antibody without increased FcRn-binding activity, and this results in an increased extent of recycling of antigens bound to the antigen to the plasma in the living organism. Thus, due to the improved plasma retention of the antibody administered in the living organism, the plasma concentration of the antigen to which the antibody binds is thought to be also increased in the living organism.

Meanwhile, when administered to a living organism, an antibody that binds to an antigen in a pH-dependent manner and which has increased FcγR-binding activity is mainly incorporated into immune cells expressing FcγR on the cell membrane, and this reduces the plasma retention. Furthermore, after being incorporated into the cells while bound to the antibody, the antigen is dissociated from the antibody in the endosome and then degraded in the lysosome, resulting in a decrease of the antigen concentration in plasma in the living organism. When the FcRn-binding activity is increased under an acidic pH range condition, the antibody retention in plasma, even if worsened due to increased FcγR-binding activity, is improved by an increase in the rate of recycling by FcRn. In this case, since the antigen bound to the antibody that binds to the antigen in a pH-dependent manner is dissociated from the antibody in the endosome and directly degraded in the lysosome, it is not thought that the antigen concentration is increased in the plasma. Furthermore, the improved plasma retention of the antibody administered to the living organism is thought to allow the antigen elimination effect of the antibody to be sustained, and the antigen concentration to be maintained low for a longer period.

The above findings demonstrate that the plasma retention of an administered antibody is

improved in a living organism administered with the antibody in which the human FcRn-binding activity under an acidic pH range condition is enhanced in an antigen-binding molecule whose FcγR-binding activity is higher than that of native human IgG Fc region. Furthermore, it was revealed that, in this case, the antibody retention in plasma is improved without deteriorating the antigen-elimination effect.

[Example 4] Further assessment of the effect of eliminating antigens from plasma antigen-binding molecules whose FcγR-binding activity is greater than that of native human IgG Fc region and whose human FcRn-binding activity has been increased under an acidic pH range condition

(4-1) The antigen elimination effect in the living organism administered with an antibody whose FcγR-binding activity is higher than that of the Fc region of native human IgG and which has human FcRn-binding activity increased under conditions of acidic pH range

As described in Example 2, the antigen concentration in plasma was significantly reduced in the group administered with Fv4-IgG1-F1022 with enhanced mouse FcγR binding. Meanwhile, as shown in Example 3, the reduced plasma retention observed in the Fv4-IgG1-F1022-administered group was markedly improved by increasing the human FcRn-binding activity of Fv4-IgG1-F1022 under an acidic pH range condition. Next, the effect of eliminating soluble antigens from plasma by enhancing mouse FcγR binding and the effect of improving the plasma retention of an antibody in the living organism administered with it by enhancing the human FcRn binding activity under an acidic pH range condition, were further assessed as described below.

(4-2) Preparation of an anti-human IL-6 receptor antibody with enhanced mouse FcγR binding

VH3-IgG1-F1087 (SEQ ID NO: 123) resulting from substituting Asp for Lys at position 326 (EU numbering) in VH3-IgG1, and VH3-IgG1-F1182 (SEQ ID NO: 124) resulting from substituting Asp for Ser at position 239 and Glu for Ile at position 332 (EU numbering) in VH3-IgG1, were prepared as antigen-binding molecules with enhanced mouse FcγR binding. Fv4-IgG1-F1087 that contains VH3-IgG1-F1087 as the heavy chain and VL3-CK as the light chain, and Fv4-IgG1-F1182 that contains VH3-IgG1-F1182 as the heavy chain and VL3-CK as the light chain, were produced using the method described in Reference Example 2.

(4-3) Assessment of mouse FcγR-binding activity

VH3/L (WT)-IgG1-F1087 and VH3/L (WT)-IgG1-F1182 which contain VH3-IgG1-F1087 and VH3-IgG1-F1182 as the heavy chain, respectively, and L (WT)-CK (SEQ

ID NO: 42) as the light chain, were prepared by the method described in Reference Example 2. These antibodies and VH3/L (WT)-IgG1-F1022 were assessed for their mouse Fc γ R-binding activity by the method described in Reference Example 2. The result is shown in Table 8. In addition, the ratio of the increase in the mouse Fc γ R-binding activity of each variant relative to the IgG1 before alteration is shown in Table 9.

[Table 8]

VARIANT NAME	KD (M)			
	mFc γ RI	mFc γ RIib	mFc γ RIII	mFc γ RIV
IgG1	5.3E-08	9.8E-07	2.4E-06	8.6E-08
F1022	7.6E-09	1.0E-08	5.5E-09	1.4E-07
F1087	2.9E-08	5.6E-08	5.2E-08	3.3E-07
F1182	2.4E-09	1.1E-07	4.8E-07	5.3E-10

[Table 9]

VARIANT NAME	RATIO OF BINDING TO:mIgG1			
	mFc γ RI	mFc γ RIib	mFc γ RIII	mFc γ RIV
IgG1	1.0	1.0	1.0	1.0
F1022	7.0	93.6	440.5	0.6
F1087	1.8	17.5	46.2	0.3
F1182	22.1	9.1	5.0	162.3

10

15

20

As shown in Table 9, it was demonstrated that F1087 and F1022 had increased binding activity to mouse Fc γ RI, mouse Fc γ RIib, and mouse Fc γ RIII as compared to IgG1, whereas their mouse Fc γ RIV-binding activity was not increased. Regarding the binding activity of F1087 to mouse Fc γ RI, mouse Fc γ RIib, mouse Fc γ RIII, and mouse Fc γ RIV, the extent of its increase was revealed to be smaller than that of F1022. Meanwhile, it was shown that the binding activity of F1182 to mouse Fc γ RI and mouse Fc γ RIV was considerably increased, whereas the extent of increase in its binding activity to Fc γ RIib and Fc γ RIII was smaller than those of F1022 and F1087. As mentioned above, these three types of variants showed enhanced binding to some mouse Fc γ Rs; however, it was shown that the Fc γ R to which the binding activity is selectively increased and the extent of the increase vary depending on the variant.

(4-4) The effect of eliminating antigens from the plasma in an individual administered with Fv4-IgG1-F1087 and Fv4-IgG1-F1182

By the same method as described in Example 2, *in vivo* infusion tests using human FcRn transgenic mice were carried out to determine the soluble IL-6 receptor concentrations in the plasma of the mice. The result is shown in Fig. 6.

In both of the groups administered with Fv4-IgG1-F1087 and Fv4-IgG1-F1182 *in vivo*,
 5 which have increased mouse FcγR-binding activity as compared to Fv4-IgG1, the *in vivo* plasma concentration of soluble human IL-6 receptor reduced as compared to the group administered with Fv4-IgG1. The effect to reduce the above plasma concentration of soluble human IL-6 receptor was high especially in the group administered with Fv4-IgG1-F1087 which has enhanced binding to mouse FcγRII and mouse FcγRIII. Meanwhile, the effect of F1182
 10 administration to reduce the plasma concentration of soluble human IL-6 receptor was small in the group administered with F1182 *in vivo* which has considerably increased binding activity to mouse FcγRI and mouse FcγRIV (as well as several-fold enhanced binding to mouse FcγRII and mouse FcγRIII). It was thought from these results that the mouse FcγRs that more significantly contribute by an effect that efficiently decreases the antigen concentration in the plasma of mice
 15 administered with a pH-dependent antigen-binding antibody, are mouse FcγRII and/or mouse FcγRIII. Specifically, it is thought that the plasma antigen concentration can be more efficiently reduced *in vivo* by administering into a living organism a pH-dependent antigen-binding antibody with enhanced binding to mouse FcγRII and/or mouse FcγRIII.

20 (4-5) Preparation of antigen-binding molecules whose FcγR-binding activity is greater than the binding activity of native human IgG Fc region and which have increased human FcRn-binding activity under an acidic pH range condition

As described in Example 3, when compared to human FcRn transgenic mice administered with Fv4-IgG1-F1022, the plasma retention of an administered antibody is
 25 markedly improved in human FcRn transgenic mice administered with Fv4-IgG1-F1093 resulting from increasing the human FcRn-binding activity under an acidic pH range condition of Fv4-IgG1-F1022 in which the mouse FcγR-binding activity has been increased. Whether this effect is also observed in human FcRn transgenic mice administered with Fv4-IgG1-F1087 and Fv4-IgG1-F1182, and whether the same effect is observed in mice administered with variants
 30 whose human FcRn-binding activity has been increased under an acidic pH range condition by addition of an alteration distinct from the alteration assessed in Example 3 were assessed as follows.

VH3-IgG1-F1180 (SEQ ID NO: 125) and VH3-IgG1-F1181 (SEQ ID NO: 126) were prepared by substituting Leu for Met at position 428 and Ser for Asn at position 434 (EU
 35 numbering) in the heavy chains VH3-IgG1-F1087 and VH3-IgG1-F1182, in order to increase their human FcRn-binding activity of Fv4-IgG1-F1087 and Fv4-IgG1-F1182 under an acidic pH

range condition. Furthermore, VH3-IgG1-F1412 (SEQ ID NO: 127) was prepared by substituting Ala for Asn at position 434 (EU numbering) in the heavy chain VH3-IgG1-F1087, in order to increase the human FcRn-binding activity of Fv4-IgG1-F1087 under an acidic pH range condition. Fv4-IgG1-F1180, Fv4-IgG1-F1181, and Fv4-IgG1-F1412, which contain the above heavy chains and VL3-CK as the light chain, were prepared using the method described in Reference Example 2.

(4-6) Improvement of pharmacodynamics of antibodies with increased human FcRn-binding activity under an acidic pH range condition

In vivo infusion tests were carried out by administering Fv4-IgG1-F1180, Fv4-IgG1-F1181, and Fv4-IgG1-F1412 to human FcRn transgenic mice according to the same method as described in Example 2 to determine the soluble IL-6 receptor concentrations in the plasma of the mice. The results on the antibody concentrations in the plasma of the mouse groups administered with Fv4-IgG1-F1087, Fv4-IgG1-F1180, Fv4-IgG1-F1412, and Fv4-IgG1 are shown in Fig. 7. The results on the antibody concentrations in the plasma of the mouse groups administered with Fv4-IgG1-F1182, Fv4-IgG1-F1181, and Fv4-IgG1 are shown in Fig. 8. Meanwhile, the plasma antibody concentrations in the mouse groups were measured by the method described in Example 3. The results on the plasma soluble IL-6 receptor concentrations of Fv4-IgG1-F1087, Fv4-IgG1-F1180, Fv4-IgG1-F1412, and Fv4-IgG1 in the mouse groups are shown in Fig. 9; and the results on the plasma soluble IL-6 receptor concentrations of Fv4-IgG1-F1182, Fv4-IgG1-F1181, and Fv4-IgG1 are shown in Fig. 10.

It was confirmed that, as compared to the group of mice administered with Fv4-IgG1-F1182, the plasma retention of administered antibodies was improved in the group of mice administered with Fv4-IgG1-F1181 resulting from increasing the human FcRn-binding activity of Fv4-IgG1-F1182 in an acidic pH range. Meanwhile, the soluble IL-6 receptor concentration in the plasma of the mouse groups administered with Fv4-IgG1-F1181 was comparable to that in the group of mice administered with Fv4-IgG1-F1182. When compared to the mouse groups administered with Fv4-IgG1, the soluble IL-6 receptor concentration in the plasma was decreased in both groups.

On the other hand, as compared to the group of mice administered with Fv4-IgG1-F1087, the plasma retention of administered antibodies was improved in both groups of mice administered with Fv4-IgG1-F1180 and Fv4-IgG1-F1412 resulting from increasing the human FcRn-binding activity of Fv4-IgG1-F1087 in an acidic pH range, and surprisingly, the plasma retention was improved up to a level comparable to that of the mouse groups administered with Fv4-IgG1. Furthermore, the sustainability of the effect of reducing the soluble IL-6 receptor concentration in plasma was improved by the improvement of the plasma

antibody retention in the groups of administered mice. Specifically, in the groups of administered mice, the soluble IL-6 receptor concentrations in plasma 14 days and 21 days after administration of Fv4-IgG1-F1180 and Fv4-IgG1-F1412 were significantly reduced as compared to the concentrations 14 days and 21 days after administration of Fv4-IgG1-F1087.

5 In view of the above, as for the groups of mice administered with the four examples of antibodies, Fv4-IgG1-F1093, Fv4-IgG1-F1181, Fv4-IgG1-F1180, and Fv4-IgG1-F1412, it was demonstrated that the plasma retention can be improved in a living organism administered with an antibody in which the human FcRn-binding activity under an acidic pH range condition has been enhanced in an antigen-binding molecule whose FcγR-binding activity is higher than the
10 binding activity of native human IgG Fc region. It was also demonstrated that, in the living organism administered with the antigen-binding molecule, the plasma retention is improved without deteriorating the effect of eliminating antigens from the living organism, and rather, the antigen elimination effect can be sustained.

15 (4-7) Preparation of antigen-binding molecules with increased human FcRn-binding activity under an acidic pH range condition and suppressed binding to a rheumatoid factor

In recent years, an antibody molecule resulting from substituting His for Asn at position 434 (EU numbering) in a humanized anti-CD4 antibody to improve the plasma retention by increasing its human FcRn-binding activity under an acidic pH range condition, has been
20 reported to bind to the rheumatoid factor (RF) (Clin. Pharmacol. Ther. (2011) 89 (2), 283-290). This antibody has a human IgG1 Fc region and a substitution of His for Asn at position 434 (EU numbering) in the FcRn-binding site. The rheumatoid factor has been demonstrated to recognize and bind to the substituted portion.

As described in Example (4-6), as compared to the case where Fv4-IgG1-F1087 was
25 administered to human FcRn transgenic mice, Fv4-IgG1-F1180 resulting from increasing under conditions of acidic pH range the human FcRn-binding activity of Fv4-IgG1-F1087 with increased mouse FcγR-binding activity, showed improved retention in plasma. Various alterations have been reported to increase the human FcRn-binding activity under conditions of acidic pH range. Of such modifications, a variant with a substitution of Leu for Met at position
30 428 and Ser for Asn at position 434 (EU numbering) in the heavy chain has been reported to show augmented binding to rheumatoid factors.

However, a variant that has a substitution of Thr for Tyr at position 436 (EU numbering) in addition to the above substitutions at positions 428 and 434 (EU numbering) shows significantly reduced binding to rheumatoid factors while retaining increased human
35 FcRn-binding activity under conditions of acidic pH range.

That is, antigen-binding molecules that have increased human FcRn-binding activity under an acidic pH range condition but do not have the binding to the rheumatoid factor can be produced by introducing into the site of the Fc region an alteration that reduces the rheumatoid factor-binding activity alone without reducing the FcRn-binding activity under an acidic pH range condition.

Such alterations used for reducing the rheumatoid factor-binding activity include alterations at positions 248-257, 305-314, 342-352, 380-386, 388, 414-421, 423, 425-437, 439, and 441-444 (EU numbering), preferably those at positions 387, 422, 424, 426, 433, 436, 438, and 440 (EU numbering), and particularly preferably, an alteration that substitutes Glu or Ser for Val at position 422, an alteration that substitutes Arg for Ser at position 424, an alteration that substitutes Asp for His at position 433, an alteration that substitutes Thr for Tyr at position 436, an alteration that substitutes Arg or Lys for Gln at position 438, and an alteration that substitutes Glu or Asp for Ser at position 440 (EU numbering). These alterations may be used alone or in combination.

Alternatively, it is possible to introduce N-type glycosylation sequences to reduce the rheumatoid factor-binding activity. Specifically, known N-type glycosylation sequences include Asn-Xxx-Ser/Thr (Xxx represents an arbitrary amino acid other than Pro). This sequence can be introduced into the Fc region to add an N-type sugar chain, and the binding to RF can be inhibited by the steric hindrance of the N-type sugar chain. Alterations used for adding an N-type sugar chain preferably include an alteration that substitutes Asn for Lys at position 248, an alteration that substitutes Asn for Ser at position 424, an alteration that substitutes Asn for Tyr at position 436 and Thr for Gln at position 438, and an alteration that substitutes of Asn for Qln at position 438, according to EU numbering, particularly preferably an alteration that substitutes Asn for Ser at position 424 (EU numbering).

(4-8) Assessment of the pharmacodynamics-improving effect of antigen-binding molecules with increased human FcRn-binding activity under conditions of acidic pH range and reduced rheumatoid factor binding

In order to assess the effect of antibodies with an alteration that reduces the above-mentioned rheumatoid factor-binding activity, Fv4-IgG1-F1782 with a substitution of Leu for Met at position 428, Ser for Asn at position 434, and Thr for Tyr at position 436 (EU numbering) in the heavy chain of Fv4-IgG1-F1087, was produced using the method described in Reference Example 2. As described in Example (4-7), Fv4-IgG1-F1782 is an antibody whose human FcRn-binding activity under conditions of acidic pH and mouse FcγR-binding activity have been both increased, but its rheumatoid factor-binding activity has not been increased as compared to native human IgG1. To test whether the plasma retention of antibody

Fv4-IgG1-F1782 has been improved as compared to Fv4-IgG1-F1087, the pharmacodynamics of the antibodies in the plasma of human FcRn transgenic mice administered with the antibodies was assessed by the same method as described in Example 2. The plasma concentration of soluble human IL-6 receptor was determined by the method described in Example (2-1-2), while
5 plasma antibody concentrations were determined by the method described in Example (3-2-1).

A time course of antibody concentrations in plasma is shown in Fig. 11, and a time course of plasma concentration of soluble human IL-6 receptor is shown in Fig. 12. Fv4-IgG1-F1782 showed improved plasma antibody retention as compared to Fv4-IgG1-F1087. Meanwhile, the plasma concentration of soluble human IL-6 receptor was significantly reduced
10 in the groups administered with the above antibodies as compared to the Fv4-IgG1 administration group.

Without being bound by a particular theory, the above results can be interpreted as follows. The results described in Examples 3 and 4 show that the plasma retention can be prolonged in the living organism administered with an antibody resulting from increasing under
15 conditions of acidic pH range the human FcRn-binding activity of an antigen-binding molecule whose FcγR-binding activity is higher than the binding activity of the Fc region of natural human IgG. It was also demonstrated that, in the living organism administered with such an antigen-binding molecule, the plasma retention is prolonged without deteriorating the effect of eliminating antigens from the living organism, and rather the antigen elimination effect can be
20 sustained.

However, it is concerned that antigen-binding molecules introduced with an alteration for increasing the human FcRn-binding activity under conditions of acidic pH range would have a risk of increased rheumatoid factor-binding activity. Thus, the plasma retention can be improved without increasing the rheumatoid factor-binding activity by introducing into the
25 binding molecules a mutation that reduces the rheumatoid factor-binding activity while retaining the human FcRn-binding activity under conditions of acidic pH range.

In other words, it was revealed that when administered into a living organism, antigen-binding molecules that have the property of binding to antigens in a pH-dependent manner, and have increased human FcRn-binding activity under conditions of acidic pH range, and whose FcγR-binding activity is greater than that of the Fc region of native human IgG, and
30 that have reduced rheumatoid factor-binding activity, have an excellent property in that the antigen-binding molecules show prolonged plasma retention without increasing their rheumatoid factor-binding activity, and effectively reduce the soluble antigen concentration in the living organism.

35
[Example 5] Effect of eliminating antigens from the plasma of a living organism administered

with an antigen-binding molecule whose Fc γ R-binding activity is higher than the binding activity of native mouse IgG Fc region

(5-1) The antigen elimination effect from the plasma of a living organism administered with a mouse antibody with increased Fc γ R-binding activity

5 As described in Examples 1 to 4, it was demonstrated that the elimination of soluble human IL-6 receptor from mouse plasma is accelerated in the groups of human FcRn transgenic mice administered with antigen-binding molecules resulting from increasing the mouse Fc γ R-binding activity of antigen-binding molecules that have a human antibody Fc region and the property of binding to human IL-6 receptor in a pH-dependent manner. Whether this effect
10 is also achieved in normal mice having mouse FcRn that was administered with antigen-binding molecules that have a mouse antibody Fc region and the property of binding to human IL-6 receptor in a pH-dependent manner, was assessed in normal mice having mouse FcRn as follows.

(5-2) Preparation of mouse antibodies with increased Fc γ R-binding activity

15 For a mouse IgG1 antibody having the property of binding to human IL-6 receptor in a pH-dependent manner, the heavy chain VH3-mIgG1 (SEQ ID NO: 128) and the light chain VL3-mk1 (SEQ ID NO: 129) were constructed using the method described in Reference Example 2. Meanwhile, to increase the mouse Fc γ R-binding activity of VH3-mIgG1, VH3-mIgG1-mF44 (SEQ ID NO: 130) was produced by substituting Asp for Ala at position 327
20 (EU numbering). Likewise, VH3-mIgG1-mF46 (SEQ ID NO: 131) was produced by substituting Asp for Ser at position 239 and Asp for Ala at position 327, according to EU numbering, in VH3-mIgG1. Fv4-mIgG1, Fv4-mIgG1-mF44, and Fv4-mIgG1-mF46, which contain VH3-mIgG1, VH3-mIgG1-mF44, and VH3-mIgG1-mF46, respectively, as the heavy chain, and VL3-mk1 as the light chain, were prepared using the method described in Reference
25 Example 2.

(5-3) Assessment of mouse Fc γ R-binding activity of mouse antibodies with enhanced Fc γ R-binding activity

VH3/L (WT)-mIgG1, VH3/L (WT)-mIgG1-mF44, and VH3/L (WT)-mIgG1-mF46,
30 which contain VH3-mIgG1, VH3-mIgG1-mF44, and VH3-mIgG1-mF46, respectively, as the heavy chain, and L (WT)-CK (SEQ ID NO: 42) as the light chain, were prepared by the method described in Reference Example 2. These antibodies were assessed for their mouse Fc γ R-binding activity by the method described in Reference Example 25. The result is shown in Table 10. In addition, the ratio of the increase in the mouse Fc γ R-binding activity of each
35 variant relative to the mIgG1 before alteration is shown in Table 11.

[Table 10]

VARIANT NAME	KD (M)			
	mFc γ RI	mFc γ RIIfb	mFc γ RIIf	mFc γ RIV
mIgG1	NOT DETECTED	1.1E-07	2.1E-07	NOT DETECTED
mF44	NOT DETECTED	8.9E-09	6.7E-09	NOT DETECTED
mF46	NOT DETECTED	1.2E-09	3.6E-09	NOT DETECTED

[Table 11]

VARIANT NAME	RATIO OF BINDING TO IgG1			
	mFc γ RI	mFc γ RIIfb	mFc γ RIIf	mFc γ RIV
mIgG1	NOT DETECTED	1.0	1.0	NOT DETECTED
mF44	NOT DETECTED	11.9	31.0	NOT DETECTED
mF46	NOT DETECTED	91.4	57.5	NOT DETECTED

5

The assessment result of Example 4 showing that VH3/L (WT)-mIgG1 having the Fc region of native mouse IgG1 antibody only binds to mouse Fc γ RIIfb and mouse Fc γ RIIf but not to mouse Fc γ RI and mouse Fc γ RIV, suggests that mouse Fc γ Rs important for the reduction of antigen concentration are mouse Fc γ RIIf and/or mouse Fc γ RIIf. VH3/L (WT)-mIgG-mF44 and VH3/L (WT)-mIgG1-mF46 introduced with an alteration that is thought to increase the Fc γ R-binding activity of VH3/L (WT)-mIgG1 was demonstrated to have increased binding activity to both of mouse Fc γ RIIfb and mouse Fc γ RIIf.

15 (5-4) Assessment of the effect to reduce the soluble IL-6 receptor concentration in the plasma of normal mice

The effect to eliminate soluble IL-6 receptor from the plasma of normal mice administered with the anti-human IL-6 receptor antibody Fv4-mIgG1, Fv4-mIgG1-mF44, or Fv4-mIgG1mF46 was assessed as follows.

20 An animal model where the soluble human IL-6 receptor concentration is maintained in a steady state in plasma was created by implanting an infusion pump (MINI-OSMOTIC PUMP MODEL2004, alzet) containing soluble human IL-6 receptor under the skin on the back of normal mice (C57BL/6J mouse, Charles River Japan). The *in vivo* dynamics of soluble human IL-6 receptor after administration of the anti-human IL-6 receptor antibody was assessed in the animal model. To suppress the production of antibodies against soluble human IL-6 receptor, 25 an anti-mouse CD4 monoclonal antibody was administered once at 20 mg/kg into the caudal

vein. Then, an infusion pump containing 92.8 µg/ml soluble human IL-6 receptor was subcutaneously implanted on the back of the mice. Three days after implantation of the infusion pump, the anti-human IL-6 receptor antibody was administered once at 1 mg/kg into the caudal vein. The blood was collected from the mice 15 minutes, seven hours, one day, two days, four days, seven days, 14 days (or 15 days), and 21 days (or 22 days) after administration of the anti-human IL-6 receptor antibody. Immediately thereafter, the collected blood was centrifuged at 15,000 rpm and 4°C for 15 minutes to prepare the plasma. The isolated plasma was stored in a freezer set at -20°C or below until use.

The soluble human IL-6 receptor concentrations in plasma were determined by the method described in (2-1-2). The result is shown in Fig. 13.

Surprisingly, it was demonstrated that, in mice administered with mF44 and mF46 introduced with an alteration to increase the binding activity of mIgG1 (native mouse IgG1) to mouse FcγRIIb and mouse FcγRIII, the plasma IL-6 receptor concentration was markedly reduced as compared to mice administered with mIgG1. In particular, even on day 21 after administration of mF44, the plasma IL-6 receptor concentration in the mF44-administered group was reduced by about 6 times as compared to the plasma IL-6 receptor concentration in the group without antibody administration, and about 10 times as compared to the mIgG1-administered group. On the other hand, on day seven after administration of mF46, the plasma IL-6 receptor concentration in the mF46-administered group was markedly reduced by about 30 times as compared to the plasma IL-6 receptor concentration in the group without antibody administration, and about 50 times as compared to the mIgG1-administered group.

The above findings demonstrate that the elimination of soluble IL-6 receptor from plasma was also accelerated in mice administered with antibodies in which the mouse FcγR-binding activity of an antigen-binding molecule having the Fc regions of mouse IgG1 antibody is increased, as with antibodies in which the mouse FcγR-binding activity of an antigen-binding molecule having the Fc region of human IgG1 antibody is increased. Without being bound by a particular theory, the phenomenon observed as described above can be explained as follows.

When administered to mice, antibodies that bind to a soluble antigen in a pH-dependent manner and have increased FcγR-binding activity are actively incorporated mainly into cells expressing FcγR on the cell membrane. The incorporated antibodies dissociate the soluble antigen under an acidic pH condition in the endosome, and then recycled to plasma *via* FcRn. Thus, a factor that achieves the effect of eliminating the plasma soluble antigen of such an antibody is the FcγR-binding activity level of the antibody. Specifically, as the FcγR-binding activity is greater, the incorporation into FcγR-expressing cells occurs more actively, and this makes the elimination of soluble antigens from plasma more rapid. Furthermore, as long as the

Fc γ R-binding activity has been increased, the effect can be assessed in the same manner regardless of whether the Fc region contained in an antibody originates from human or mouse IgG1. Specifically, the assessment can be achieved for an Fc region of any animal species, such as any of human IgG1, human IgG2, human IgG3, human IgG4, mouse IgG1, mouse IgG2a, mouse IgG2b, mouse IgG3, rat IgG, monkey IgG, and rabbit IgG, as long as the binding activity to the Fc γ R of the animal species to be administered has been increased.

[Example 6] The antigen elimination effect by antibodies with the binding activity increased in a Fc γ RIIb-selective manner

10 (6-1) The antigen elimination effect of antibodies in which the Fc γ RIIb-binding activity has been selectively increased

Fc γ RIII-deficient mice (B6.129P2-FcgrFc γ R3tm1Sjv/J mouse, Jackson Laboratories) express mouse Fc γ RI, mouse Fc γ RIIb, and mouse Fc γ RIV, but not mouse Fc γ RIII. Meanwhile, Fc receptor γ chain-deficient mice (Fcer1g mouse, Taconic, Cell (1994) 76, 519-529) express mouse Fc γ RIIb alone, but not mouse Fc γ RI, mouse Fc γ RIII, and mouse Fc γ RIV.

As described in Example 5, it was demonstrated that mF44 and mF46 with increased Fc γ R-binding activity of native mouse IgG1 show selectively enhanced binding to mouse Fc γ RIIb and mouse Fc γ RIII. It was conceived that, using the selectively increased binding activity of the antibodies, the condition under which an antibody with selectively enhanced mouse Fc γ RIIb binding is administered can be mimicked by administering mF44 and mF46 to mouse Fc γ RIII-deficient mice or Fc receptor γ chain-deficient mice which do not express mouse Fc γ RIII.

25 (6-2) Assessment of the antigen elimination effect in a living organism administered with an antibody with -selective enhancement of binding to mouse Fc γ RIIb using Fc γ RIII-deficient mice

The effect to eliminate soluble IL-6 receptor from plasma in Fc γ RIII-deficient mice administered with the anti-human IL-6 receptor antibody Fv4-mIgG1, Fv4-mIgG1-mF44, or Fv4-mIgG1-mF46 was assessed by the same method described in Example 5. The soluble human IL-6 receptor concentrations in the plasma of the mice were determined by the method described in Example (2-1-2). The result is shown in Fig. 14.

Surprisingly, it was demonstrated that, the plasma IL-6 receptor concentrations in Fc γ RIII-deficient mice administered with mF44 and mF46, which mimic the condition under which the mouse Fc γ RIIb-binding activity of mIgG1 (native mouse IgG1) is selectively increased, were markedly reduced as compared to the plasma IL-6 receptor concentration in mice administered with mIgG1. In particular, the plasma IL-6 receptor concentration of the mF44-administered group was reduced by about three times as compared to that of the

mIgG1-administered group and the accumulation of antibody concentration due to antibody administration was suppressed. Meanwhile, on day three after administration, the plasma IL-6 receptor concentration of the mF46-administered group was markedly reduced by about six times as compared to the plasma IL-6 receptor concentration of the group without antibody administration, and about 25 times as compared to the plasma IL-6 receptor concentration of the mIgG1-administered group. This result shows that, as the mouse FcγRIIb-binding activity of an anti-human IL-6 receptor antibody that binds to the antigen in a pH-dependent manner is greater, the IL-6 receptor concentration can be reduced more in the plasma of mice administered with the antibody.

(6-3) Assessment of the antigen elimination effect in the plasma of a living organism administered with an antibody with selective enhancement of mouse FcγRIIb binding using Fc receptor γ chain-deficient mice

The effect to eliminate soluble IL-6 receptor from the plasma of Fc receptor γ chain-deficient mice administered with the anti-human IL-6 receptor antibody Fv4-mIgG1, Fv4-mIgG1-mF44, or Fv4-mIgG1-mF46, was assessed by the same method as described in Example 5. The soluble human IL-6 receptor concentrations in the plasma of the mice were determined by the method described in Example (2-1-2). The result is shown in Fig. 15.

As with the case where mF44 and mF46 were administered to FcγRIII-deficient mice, the plasma IL-6 receptor concentration in Fc receptor γ chain-deficient mice administered with mF44 and mF46, which mimic the condition resulting from the selective increase in the mouse FcγRIIb-binding activity of mIgG1 (native mouse IgG1), was demonstrated to be markedly reduced as compared to the plasma IL-6 receptor concentration in Fc receptor γ chain-deficient mice administered with mIgG1. In particular, the plasma IL-6 receptor concentration in the mF44-administered group was reduced to about three times that in the mIgG1-administered group, and the accumulation of antigen concentration due to antibody administration was suppressed. Meanwhile, on day three after administration, the plasma IL-6 receptor concentration in the mF46-administered group was markedly reduced by about five times as compared to that in the group without antibody administration, and about 15 times as compared to that in the mIgG1-administered group.

The results described in Examples (6-2) and (6-3) show that the soluble antigen concentration in the plasma is markedly reduced in the group administered with an antibody that binds to a soluble antigen in a pH-dependent manner and has selectively increased mouse FcγRIIb-binding activity.

[Example 7] The antigen elimination effect of antibodies with selective enhancement of the

binding to Fc γ RIII

(7-1) The antigen elimination effect in the plasma of a living organism administered with antibodies with selectively enhanced Fc γ RIII binding

Fc γ RIIb-deficient mice (Fc γ R2b (Fc γ RII) mouse, Taconic) (Nature (1996) 379
 5 (6563), 346-349) express mouse Fc γ RI, mouse Fc γ RIII, and mouse Fc γ RIV, but not mouse Fc γ RIIb. As described in Example 5, it was demonstrated that mF44 and mF46 resulting from increasing the Fc γ R-binding activity of native mouse IgG1 show selectively enhanced binding to mouse Fc γ RIIb and mouse Fc γ RIII. It was conceived that, based on the use of the selectively increased binding activity of the antibodies, the condition of administration of an antibody with
 10 selectively enhanced binding to mouse Fc γ RIII can be mimicked by administering mF44 or mF46 to mouse Fc γ RIIb-deficient mice which do not express mouse Fc γ RIIb.

As described in Example 6, the soluble antigen concentration was reduced in the plasma of Fc γ RIII-deficient mice, which mimic the condition of administration of an antibody with selectively increased mouse Fc γ RIIb-binding activity. Meanwhile, whether the soluble antigen
 15 concentration is reduced in the plasma of Fc γ RIIb-deficient mice, which mimic the condition of administration of an antibody with selectively increased mouse Fc γ RIII-binding activity, was assessed by the test described below.

(7-2) Assessment of the antigen elimination effect by selective enhancement of mouse Fc γ RIII binding using Fc γ RIIb-deficient mice

The effect to eliminate soluble IL-6 receptor from the plasma of Fc γ RIIb-deficient mice administered with the anti-human IL-6 receptor antibody Fv4-mIgG1, Fv4-mIgG1-mF44, or Fv4-mIgG1-mF46, was assessed by the same method as described in Example 5. The soluble human IL-6 receptor concentrations in plasma were determined by the method described in
 25 Example (2-1-2). The result is shown in Fig. 16.

Surprisingly, in the groups administered with mF44 and mF46, which mimic selective increase of the mouse Fc γ RIII-binding activity of mIgG1 (native mouse IgG1), the plasma IL-6 receptor concentration was reduced, but the reduction was not confirmed compared to that shown in Example 6.

Without being bound by a particular theory, based on the results described in Examples 5, 6, and 7, the following discussion is possible. The elimination of soluble IL-6 receptor from plasma was found to be markedly accelerated in normal mice expressing both mouse Fc γ RIIb and mouse Fc γ RIII that were administered with mF44 and mF46 with selectively increased binding activity of mIgG1 (native mouse IgG1) to mouse Fc γ RIIb and mouse Fc γ RIII.
 35 Furthermore, it was revealed that, when mF44 and mF46 were administered to mice that express mouse Fc γ RIIb but not mouse Fc γ RIII (*i.e.*, Fc γ RIII-deficient mice and Fc receptor γ

chain-deficient mice), the elimination of soluble IL-6 receptor from plasma was also accelerated markedly in the mice. Meanwhile, when mF44 and mF46 were administered to mice that express mouse FcγRIII but not mouse FcγRIIb (*i.e.*, FcγRII-deficient mice), the elimination of soluble IL-6 receptor from plasma was not accelerated in the mice to the extent that they were administered in FcγRIII-deficient mice or Fc receptor γ chain-deficient mice.

From the above findings, it is thought that, the antibodies mF44 and mF46 in which the binding activity of mIgG1 (native mouse IgG1) to mouse FcγRIIb and mouse FcγRIII is increased, are incorporated into FcγR-expressing cells mainly by mouse FcγRIIb, and thus the soluble antigen in the plasma that binds to the antibodies is eliminated. Meanwhile, the FcγRIII-mediated incorporation of antibody/antigen complexes into FcγR-expressing cells is thought not to significantly contribute to the elimination of the soluble antigen from plasma.

Furthermore, as shown in Example 4, the plasma concentration of soluble IL-6 receptor was markedly reduced in mice administered with Fv4-IgG1-F1087 having increased binding activity to mouse FcγRIIb and mouse FcγRIII, in particular. Meanwhile, the effect to eliminate soluble IL-6 receptor from the plasma of mice administered with Fv4-IgG1-F1182 with increased binding activity to mouse FcγRI and mouse FcγRIV, in particular, was smaller than that of Fv4-IgG1-F1087.

Furthermore, as shown in Example 2, in mice administered with Fv4-IgG1-Fuc whose mouse FcγRIV-binding activity has been considerably increased by having sugar chains with low fucose content (Science (2005) 310 (5753) 1510-1512), the plasma concentration of soluble IL-6 receptor was reduced as compared to that in mice administered with Fv4-IgG1; however, the reduction effect was as small as about twice. Thus, mouse FcγRIV-mediated incorporation of antibodies into FcγR-expressing cells is thought not to significantly contribute to the elimination of soluble antigens from plasma.

In view of the above, it was demonstrate that, of several mouse FcγRs, mouse FcγRIIb and mouse FcγRIII, in particular mouse FcγRIIb, plays a major role in antibody incorporation into FcγR-expressing cells in mice. Thus, it would be thought that mutations to be introduced into the mouse FcγR-binding domain preferably include, but are not limited to, mutations that augment the binding to mouse FcγRIIb and mouse FcγRIII, in particular, the binding to mouse FcγRIIb.

The above findings demonstrate that, in mice, the binding activity to mouse FcγRIIb and mouse FcγRIII, in particular, the FcγRIIb-binding activity of the antibodies to be administered is more preferably increased to accelerate the elimination of soluble antigens from the plasma of a living organism by administering to it antigen-binding molecules that bind to soluble antigens in a pH-dependent manner and have increased FcγR binding activity. Specifically, when administered to a living organism, antigen-binding molecules that bind to soluble antigens in a

pH-dependent manner and have increased binding activity to mouse FcγRIIb and mouse FcγRIII, in particular increased FcγRIIb-binding activity, can accelerate the elimination of the soluble antigens from plasma and effectively reduce the plasma concentration of soluble antigens, and thus, the antigen-binding molecules were revealed to show a very effective action.

5

[Example 8] Assessment of the platelet aggregatory ability of antibodies containing an Fc region introduced with an existing alteration that enhances the FcγRIIb binding

(8-1) Preparation of antibodies containing an Fc region introduced with an existing alteration that enhances the FcγRIIb binding

10

As described in Reference Example 7, antigens can be efficiently eliminated from the plasma of the living organism by administering antibodies with selectively increased FcγRIIb-binding activity to the living organism. Furthermore, the administration of antibodies containing an Fc region with selectively increased FcγRIIb-binding activity is thought to be preferred from the viewpoint of safety and side effects in the living organism administered with

15

such antibodies. However, the mouse FcγRIIb binding and mouse FcγRIII binding are both enhanced in mF44 and mF46, and thus the binding enhancement is not selective for mouse FcγRIIb. Since the homology between mouse FcγRIIb and mouse FcγRIII is high, it would be difficult to find an alteration that enhances the mouse FcγRIIb-selective binding while distinguishing the two.

20

Moreover, there is no previous report on Fc regions with selectively enhanced mouse FcγRIIb binding. Also, the homology between human FcγRIIb and human FcγRIIa (the two allotypes, 131Arg and 131His) is also known to be high. Moreover, there is no report on Fc regions that contain an alteration that augments the human FcγRIIb-selective binding while distinguishing the two (Seung *et al.*, (Mol. Immunol. (2008) 45, 3926-3933); Greenwood *et al.*, (Eur. J. Immunol. (1993) 23 (5), 1098-1104)). Furthermore, the FcγRIIa binding has been reported to play an important role in the platelet aggregatory activity of an antibody (Meyer *et al.*, (J. Thromb. Haemost. (2009), 7 (1), 171-181); Robles-Carrillo *et al.*, (J. Immunol. (2010), 185 (3), 1577-1583)). In view of these reports, it is possible that an antibody with augmented FcγRIIa binding can increase the risk of developing thrombosis in a living organism administered with it.

25

Thus, whether antibodies with augmented FcγRIIa binding have an increased platelet aggregatory activity was assessed as follows.

30

(8-2) Assessment of the human FcγR-binding activity of antibodies containing an Fc region introduced with an existing alteration that enhances the FcγRIIb binding

35

Antibodies containing an Fc region introduced with an existing alteration that enhances the human FcγRIIb binding were analyzed for their affinity for human FcγRIa, R-type and

H-type Fc γ RIIa, Fc γ RIIb, and Fc γ RIIIa by the following procedure. An H chain was constructed to have, as the antibody H chain variable region, the antibody variable region IL6R-H (SEQ ID NO: 132) against human interleukin 6 receptor which is disclosed in WO2009/125825, and as the antibody H chain constant region, IL6R-G1d (SEQ ID NO: 133) that has G1d resulting from removing the C-terminal Gly and Lys from human IgG1. Then, IL6R-G1d-v3 (SEQ ID NO: 138) was constructed by altering the Fc region of IL6R-G1d by the substitution of Glu for Ser at position 267 (EU numbering) and Phe for Leu at position 328 (EU numbering), as described in Seung *et al.*, (Mol. Immunol. (2008) 45, 3926-3933). IL6R-L (SEQ ID NO: 135) which is the L chain of anti-human interleukin 6 receptor antibody was used as a common antibody L chain, and expressed in combination with respective H chains according to the method described in Reference Example 1, and the resulting antibodies were purified. Hereinafter, antibodies containing IL6R-G1d and IL6R-G1d-v3 as the heavy chain are referred to as IgG1 and IgG1-v3, respectively.

Then, the interaction between Fc γ R and the above antibodies was kinetically analyzed using Biacore T100 (GE Healthcare). The assay for the interaction was carried out at 25°C using HBS-EP+ (GE Healthcare) as a running buffer. The chip used was a Series S Sencor Chip CM5 (GE Healthcare) immobilized with Protein A by an amino coupling method. Each Fc γ R diluted with the running buffer was allowed to interact with the antibodies of interest captured onto the chip to measure the binding of the antibodies to each Fc γ R. After measurement, 10 mM glycine-HCl (pH 1.5) was reacted to the chip to wash off the captured antibodies to repeatedly use the regenerated chip. A sensorgram obtained as a result of the measurement was analyzed using 1:1 Langmuir binding model with Biacore Evaluation Software, and binding rate constant k_a (L/mol/s) and dissociation rate constant k_d (1/s) were calculated, and the dissociation constant K_D (mol/l) was calculated from these values. The K_D values of IgG1 and IgG1-v3 to each Fc γ R are shown in Table 12 (the K_D values of each antibody to each Fc γ R), while the relative K_D values of IgG1-v3, which are obtained by dividing K_D of IgG1 to each Fc γ R by K_D of IgG1-v3 to each Fc γ R, are shown in Table 13.

[Table 12]

ANTIBODY	KD (M)				
	Fc γ RIa	Fc γ RIIfaR	Fc γ RIIfaH	Fc γ RIIfb	Fc γ RIIfa
IgG1	3.4E-10	1.2E-06	7.7E-07	5.3E-06	3.1E-06
IgG1-v3	1.9E-10	2.3E-09	1.5E-06	1.3E-08	8.8E-06

[Table 13]

	Fc γ RIa	Fc γ RIIR	Fc γ RIIRH	Fc γ RIIRb	Fc γ RIIRa
KD VALUE RATIO	1.8	522	0.51	408	0.35

(8-3) Assessment of the ability to aggregate platelets

Next, whether the increased/reduced Fc γ RIIRa affinity of the antibody containing the Fc region with the substitution of Glu for Ser at position 267 and Phe for Leu at position 328 (EU numbering) in the Fc region of IgG1 changes the platelet aggregatory ability, was assessed using platelets derived from donors with H-type or R-type Fc γ RIIRa. The antibody comprising as the light chain omalizumab_VL-CK (SEQ ID NO: 137) and omalizumab_VH-G1d (SEQ ID NO: 136) that contains the heavy chain variable region of hIgG1 antibody (human IgG1 constant region) that binds to IgE and the G1d heavy chain constant region, was constructed using the method described in Reference Example 2. Furthermore, omalizumab_VH-G1d-v3 (SEQ ID NO: 138) was constructed by substituting Glu for Ser at position 267 and Phe for Leu at position 328 (EU numbering) in omalizumab_VH-G1d. Omalizumab-G1d-v3, which contains omalizumab_VH-G1d-v3 as the heavy chain and omalizumab_VL-CK as the light chain, was prepared using the method described in Reference Example 2. This antibody was assessed for the platelet aggregatory ability.

Platelet aggregation was assayed using the platelet aggregometer HEMA TRACER 712 (LMS Co.). First, about 50 ml of whole blood was collected at a fixed amount into 4.5-ml evacuated blood collection tubes containing 0.5 ml of 3.8% sodium citrate, and this was centrifuged at 200 g for 15 minutes. The resultant supernatant was collected and used as platelet-rich plasma (PRP). After PRP was washed with buffer A (137 mM NaCl, 2.7 mM KCl, 12 mM NaHCO₃, 0.42 mM NaH₂PO₄, 2 mM MgCl₂, 5 mM HEPES, 5.55 mM dextrose, 1.5 U/ml apyrase, 0.35 % BSA), the buffer was replaced with buffer B (137 mM NaCl, 2.7 mM KCl, 12 mM NaHCO₃, 0.42 mM NaH₂PO₄, 2 mM MgCl₂, 5 mM HEPES, 5.55 mM dextrose, 2 mM CaCl₂, 0.35 % BSA). This yielded washed platelets at a density of about 300,000/ μ l. 156 μ l of the washed platelets was aliquoted into assay cuvettes containing a stir bar in the platelet aggregometer. The platelets were stirred at 1000 rpm with the stir bar in the cuvettes maintained at 37.0°C in the platelet aggregometer. 44 μ l of the immune complex of omalizumab-G1d-v3 and IgE at a molar ratio of 1:1, prepared at final concentrations of 600 μ g/ml and 686 μ g/ml, respectively, was added to the cuvettes. The platelets were reacted with the immune complex for five minutes. Then, at a concentration that does not allow secondary platelet aggregation, adenosine diphosphate (ADP, SIGMA) was added to the reaction mixture to test whether the aggregation is enhanced.

The result of platelet aggregation for each donor with an Fc γ RIIRa polymorphic form

(R/H or H/H) obtained from the above assay is shown in Figs. 17 and 18, respectively. From the result in Fig. 17, platelet aggregation is observed when the immune complex is added to the platelets of a donor with the FcγRIIa polymorphic form (R/H). Meanwhile, as shown in Fig. 18, platelet aggregation was not observed when the immune complex is added to the platelets of a donor with the FcγRIIa polymorphic form (H/H).

Next, platelet activation was assessed using activation markers. Platelet activation can be measured based on the increased expression of an activation marker such as CD62p (p-selectin) or active integrin on the platelet membrane surface. 2.3 μl of the immune complex was added to 7.7 μl of the washed platelets prepared by the method described above. After five minutes of reaction at room temperature, activation was induced by adding ADP at a final concentration of 30 μM, and whether the immune complex enhances the ADP-dependent activation was assessed. A sample added with phosphate buffer (pH 7.4) (Gibco), instead of the immune complex, was used as a negative control. Staining was performed by adding, to each post-reaction sample, PE-labeled anti-CD62 antibody (BECTON DICKINSON), PerCP-labeled anti-CD61 antibody, and FITC-labeled PAC-1 antibody (BD bioscience). Fluorescence intensity for each stain was measured using a flow cytometer (FACSCanto™II, BD bioscience).

The result on CD62p expression, obtained by the above assay method, is shown in Fig. 19. The result on the activated integrin expression is shown in Fig. 20. The washed platelets used were obtained from a healthy person with the FcγRIIa polymorphic form R/H. The expression amount of both CD62p and active integrin expressed on platelet membrane surface, which is induced by ADP stimulation, was enhanced in the presence of the immune complex.

The above results demonstrate that the antibody having the Fc region introduced with an existing alteration that enhances the human FcγRIIb binding, which is the substitution of Glu for Ser at position 267 and Phe for Leu at position 328 (EU numbering) in the Fc region of IgG1, promotes the aggregation of platelets expressing FcγRIIa with the FcγRIIa allotype in which the amino acid at position 131 is R, as compared to platelets expressing FcγRIIa with the FcγRIIa polymorphic form in which the amino acid at position 131 is H. That is, it was suggested that the risk of developing thrombosis due to platelet aggregation can be increased when an antibody containing an Fc region introduced with an existing alteration that enhances the binding to existing human FcγRIIb is administered to humans having R-type FcγRIIa in at least one allele. It was shown that the antigen-binding molecules containing an Fc region of the present invention that enhances the FcγRIIb binding more selectively not only improves the antigen retention in plasma, but also possibly solves the above problems. Thus, the usefulness of the antigen-binding molecules of the present invention is obvious.

(8-4) Comparison of the platelet aggregatory ability between an antibody (BP230) comprising Fc

with selectively augmented FcγRIIb binding and an antibody comprising an existing Fc with selectively augmented FcγRIIb binding

(8-4-1) Preparation of antibodies with selectively augmented FcγRIIb binding

As described in Example (8-3), it was shown that antibodies with augmented FcγRIIa binding have increased platelet aggregatory ability and increased platelet activation ability and, they can increase the risk of developing thrombosis when administered to humans. Meanwhile, considering that of the FcγRs, FcγRIIa alone is expressed on platelets, it is thought that antibodies with selectively augmented FcγRIIb binding do not show increased platelet aggregatory ability or activation ability, or do not increase the risk of developing thrombosis when administered to humans. To confirm this, antibodies with selectively augmented FcγRIIb binding were actually tested for their platelet aggregatory ability and platelet activation ability.

Specifically, omalizumab_VH-G1d (SEQ ID NO: 136) and omalizumab_VL-CK (SEQ ID NO: 137), as the heavy chain and light chain of the hIgG1 antibody (human IgG1 constant region) that binds to IgE, respectively, were produced by the method described in Reference Example 2. Then, to selectively increase the human FcγRIIb-binding activity of omalizumab_VH-G1d, omalizumab_VH-BP230 (SEQ ID NO: 140) was produced by substituting Asp for Glu at position 233, Asp for Gly at position 237, Asp for Pro at position 238, Asp for His at position 268, Gly for Pro at position 271, Asp for Tyr at position 296, Arg for Ala at position 330, and Glu for Lys at position 439 (EU numbering). Likewise, to increase the human FcγRIIb- and FcγRIIa R-binding activity of omalizumab_VH-G1d, omalizumab_VH-G1d-v3 (SEQ ID NO: 138) was produced by substituting Glu for Ser at position 267 and Phe for Leu at position 328 (EU numbering).

Omalizumab-BP230 that comprises omalizumab_VH-BP230 (SEQ ID NO: 140) as the heavy chain and omalizumab_VL-CK (SEQ ID NO: 137) as the light chain, and omalizumab-G1d-v3 that comprises omalizumab_VH-G1d-v3 (SEQ ID NO: 138) as the heavy chain and omalizumab_VL-CK (SEQ ID NO: 137) as the light chain, were produced using the method described in Reference Example 2. These antibodies were assessed for their ability to aggregate and activate platelets.

(8-4-2) Assessment of omalizumab-BP230 and omalizumab-G1d-v3 for their human FcγR-binding activity

The analysis results on the affinity between each human FcγR and the Fc region of omalizumab-G1d-v3 resulting from augmenting the human FcγRIIb binding of omalizumab, which is a known antibody, are shown in Example (8-2). The affinity of the Fc region of omalizumab-BP230 for each human FcγR was also analyzed in the same manner, and the results are shown in Table 22.

(8-4-3) Assessment of the ability to aggregate platelets

Then, omalizumab-BP230 and omalizumab-G1d-v3 produced as described in Example (8-4-1) were assayed for their binding using platelets from a donor with the FcγRIIa polymorphic form (R/H). The assay result was used to assess whether the platelet aggregatory ability is altered depending on the level of the affinity for FcγRIIa.

Platelet aggregation assay was carried out using the platelet aggregometer HEMA TRACER 712 (LMS Co.). First, about 50 ml of whole blood was prepared by collecting fixed amounts of blood into 4.5-ml evacuated blood collection tubes containing 0.5 ml of 3.8% sodium citrate, and centrifuged at 200 g for 15 minutes. The resultant supernatant was collected and used as platelet-rich plasma (PRP). After washing PRP with buffer A (137 mM NaCl, 2.7 mM KCl, 12 mM NaHCO₃, 0.42 mM NaH₂PO₄, 2 mM MgCl₂, 5 mM HEPES, 5.55 mM dextrose, 1.5 U/mL apyrase, 0.35 % BSA), the buffer was changed with buffer B (137 mM NaCl, 2.7 mM KCl, 12 mM NaHCO₃, 0.42 mM NaH₂PO₄, 2 mM MgCl₂, 5 mM HEPES, 5.55 mM dextrose, 2 mM CaCl₂, 0.35 % BSA). This yielded washed platelets at a density of about 300,000/μl. 150.2 μl of washed platelets was aliquoted to assay cuvettes containing a stir bar in the platelet aggregometer. The platelets were stirred at 1000 rpm with the stir bars in the cuvettes maintained at 37.0°C in the platelet aggregometer. 24.3 μl of omalizumab-G1d-v3 or omalizumab-BP230 prepared at a final concentration of 600 μg/ml was added to the cuvettes. After one minute of reaction, 25.4 μl of IgE, which was prepared so that its molar ratio to the antibody is 1:1, was added thereto. The mixtures were incubated for five minutes. Then, at a concentration that does not induce secondary platelet aggregation, adenosine diphosphate (ADP, SIGMA) was added to the reaction mixture to test whether the aggregation is augmented. The washed platelets used were obtained from one healthy person with the FcγRIIa polymorphic form R/H.

The result obtained by the above-described assay method is shown in Fig. 21. The result shown in Fig. 21 revealed that platelets were strongly aggregated when adding the immune complex containing omalizumab-G1d-v3. On the other hand, there was no difference in platelet aggregation between when PBS was added and when the immune complex containing omalizumab-BP230 was added.

Next, the platelet activation by omalizumab-G1d-v3 and omalizumab-BP230 was assessed. Platelet activation can be measured based on the increased expression of an activation marker such as CD62p (p-selectin) and active integrin, on the platelet membrane surface. 1.22 μl of omalizumab-G1d-v3 or omalizumab-BP230 prepared at a final concentration of 600 μg/ml was added to 7.51 μl of washed platelets prepared by the method described above. After five minutes of reaction at room temperature, 1.27 μl of IgE, which was

prepared so that its molar ratio to the antibody is 1:1, was added thereto. Then, after five minutes of reaction at room temperature, whether platelet activation is induced was assessed. The negative control used was a sample added with phosphate buffer (pH 7.4, Gibco), instead of immune complexes. After reaction, each sample was fluorescently stained using PE-labeled
 5 anti-CD62 antibody (BECTON DICKINSON), PerCP-labeled anti-CD61 antibody, and FITC-labeled PAC-1 antibody (BD bioscience), and the fluorescence intensity was determined using a flow cytometer (FACS CantoII, BD bioscience). The washed platelets used were obtained from one healthy person with the FcγRIIa polymorphic form R/H.

The results obtained by the above-described assay method are shown in Figs. 22 and 23.

10 The expression of both CD62p and active integrin on the platelet membrane surface was increased by addition of the immune complex containing omalizumab-G1d-v3. Meanwhile, the expression of the molecules on the platelet membrane surface by addition of the immune complex with omalizumab-BP230 was comparable to the expression on the platelet membrane surface by addition of the phosphate buffer.

15 The above results demonstrate that the antibody that comprising the existing modified Fc region with substitutions of Glu for Ser at position 267 and Phe for Leu at position 328 (EU numbering) in the Fc region of IgG1, and which has augmented binding to both human FcγRIIb and human FcγRIIa R, promote the aggregation and activation of platelets in which the amino acid of position 131 is Arg in at least either of the alleles for the polymorphic FcγRIIa gene, as
 20 compared to the antibody that comprises an Fc region with substitutions of Asp for Glu at position 233, Asp for Gly at position 237, Asp for Pro at position 238, Asp for His at position 268, Gly for Pro at position 271, Asp for Tyr at position 296, Arg for Ala at position 330, and Glu for Lys at position 439 (EU numbering), and which has selectively augmented binding to human FcγRIIb. Specifically, the antibody that comprises the existing Fc region variant with
 25 augmented human FcγRIIb binding was suggested to have a problem that it can increase the risk of developing thrombosis due to platelet aggregation in humans with the FcγRIIa R allele; on the other hand, the Fc region variant with selectively augmented FcγRIIb binding was demonstrated to overcome this problem.

30 [Example 9] Comprehensive analysis of FcγRIIb binding of variants introduced with an alteration at the hinge portion in addition to the P238D alteration

In an Fc produced by substituting Pro at position 238 (EU numbering) with Asp in a naturally-occurring human IgG1, an anticipated combinatorial effect could not be obtained even by combining it with another alteration predicted to further increase FcγRIIb binding from the
 35 analysis of naturally-occurring antibodies. Therefore, in order to find variants that further enhance FcγRIIb binding, alterations were comprehensively introduced into the altered Fc

produced by substituting Pro at position 238 (EU numbering) with Asp. IL6R-F11 (SEQ ID NO: 141) was produced by introducing an alteration of substituting Met at position 252 (EU numbering) with Tyr and an alteration of substituting Asn at position 434 (EU numbering) with Tyr in IL6R-G1d (SEQ ID NO: 133) which was used as the antibody H chain. Furthermore, 5 IL6R-F652 (SEQ ID NO: 142) was prepared by introducing an alteration of substituting Pro at position 238 (EU numbering) with Asp into IL6R-F11. Expression plasmids containing an antibody H chain sequence were prepared for each of the antibody H chain sequences produced by substituting the region near the residue at position 238 (EU numbering) (positions 234 to 237, and 239 (EU numbering)) in L6R-F652 each with 18 amino acids excluding the original amino 10 acids and Cysteine. IL6R-L (SEQ ID NO: 135) was utilized as an antibody L chain. These variants were expressed and purified by the method of Reference Example 2. These Fc variants are called PD variants. Interactions of each PD variant with FcγRIIa type R and FcγRIIb were comprehensively evaluated by the method of Reference Example 25.

A figure that shows the results of analyzing the interaction with the respective FcγRs 15 was produced according to the following method. The value obtained by dividing the value for the amount of binding of each PD variant to each FcγR by the value for the amount of FcγR binding of the pre-altered antibody which is used as the control (IL6R-F652/IL6R-L, which has an alteration of substituting Pro at position 238 (EU numbering) with Asp and then multiplying the result by 100, was shown as the relative binding activity value of each PD variant to each 20 FcγR. The horizontal axis shows relative values of the FcγRIIb-binding activity of each PD variant, and the vertical axis shows relative values of the FcγRIIa type R-binding activity values of each PD variant (Fig. 24).

As a result, it was found that the FcγRIIb binding of eleven types of alterations were enhanced compared with the antibody before introducing alterations, and they have the effects of 25 maintaining or enhancing FcγRIIa type R-binding. The activities of these eleven variants to bind FcγRIIb and FcγRIIa R are summarized in Table 14. In the table, the sequence ID numbers refer to those of the H chains of the variants, and alteration refers to the alteration introduced into IL6R-F11 (SEQ ID NO: 141).

30 [Table 14]

VARIANT NAME	ALTERATION	RELATIVE BINDING ACTIVITY TO FcγRIIb	RELATIVE BINDING ACTIVITY TO FcγRIIaR
IL6R-F652/IL6R-L	P238D	100	100
IL6R-PD042/IL6R-L	P238D/L234W	106	240
IL6R-PD043/IL6R-L	P238D/L234Y	112	175
IL6R-PD079/IL6R-L	P238D/G237A	101	138
IL6R-PD080/IL6R-L	P238D/G237D	127	222
IL6R-PD081/IL6R-L	P238D/G237E	101	117
IL6R-PD082/IL6R-L	P238D/G237F	108	380
IL6R-PD086/IL6R-L	P238D/G237L	112	268
IL6R-PD087/IL6R-L	P238D/G237M	109	196
IL6R-PD094/IL6R-L	P238D/G237W	122	593
IL6R-PD095/IL6R-L	P238D/G237Y	124	543
IL6R-PD097/IL6R-L	P238D/S239D	139	844

Fig. 25 shows relative values for the FcγRIIb-binding activity obtained by additionally introducing the above eleven alterations into a variant carrying the P238D alteration, and relative values for the FcγRIIb-binding activity of a variant obtained by introducing the alterations into an Fc that does not contain the P238D. These eleven alterations enhanced the amount of FcγRIIb binding compared with before introduction when they were further introduced into the P238D variant. On the contrary, the effect of lowering FcγRIIb binding was observed for eight of those alterations except G237F, G237W, and S239D, when they were introduced into the variant that does not contain P238D (data not shown).

These results showed that, based on the effects of introducing alterations into a naturally-occurring IgG1, it is difficult to predict the effects of combining and introducing the same alterations into the variant containing the P238D alteration. In other words, it would not have been possible to discover these eight alterations identified this time without this investigation that introduces the same alterations are combined and introduced into the variant containing the P238D alteration.

The results of measuring KD values of the variants indicated in Table 14 for FcγRIa, FcγRIIaR, FcγRIIaH, FcγRIIb, and FcγRIIIaV by the method of Reference Example 25 are summarized in Table 15. In the table, alteration refers to the alteration introduced into IL6R-F11 (SEQ ID NO: 141). The template used for producing IL6R-F11, IL6R-G1d/IL6R-L, is indicated with an asterisk (*). Furthermore, KD (IIaR)/KD (IIb) and KD (IIaH)/KD (IIb) in the table respectively show the value obtained by dividing the KD value of each variant for FcγRIIaR by the KD value of each variant for FcγRIIb, and the value obtained by dividing the KD value of each variant for FcγRIIaH by the KD value of each variant for FcγRIIb. KD (IIb) of the parent polypeptide / KD (IIb) of the variant refers to a value obtained by dividing the KD value of the parent polypeptide for FcγRIIb by the KD value of each variant for FcγRIIb.

In addition, Table 15 shows KD values for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of each variant / KD values for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide. Here, parent polypeptide refers to a variant which has IL6R-F11 (SEQ ID NO: 141) as the H chain. It was determined that due to weak binding of FcγR to IgG, it was impossible to accurately analyze by kinetic analysis, and thus the gray-filled cells in Table 15 show values calculated by using Equation 2 of Reference Example 25.

[Equation 2]

$$KD = C \bullet R_{\max} / (R_{\text{eq}} - RI) - C$$

Table 15 shows that all variants improved their affinity for FcγRIIb in comparison with IL6R-F11, and the range of improvement was 1.9 fold to 5.0 fold. The ratio of KD value of each variant for FcγRIIaR / KD value of each variant for FcγRIIb, and the ratio of KD value of each variant for FcγRIIaH / KD value of each variant for FcγRIIb represent an FcγRIIb-binding activity relative to the FcγRIIaR-binding activity and FcγRIIaH-binding activity, respectively. That is, these values show the degree of binding selectivity of each variant for FcγRIIb, and a larger value indicates a higher binding selectivity for FcγRIIb. For the parent polypeptide IL6R-F11/IL6R-L, the ratio of KD value for FcγRIIaR / KD value for FcγRIIb and the ratio of KD value for FcγRIIaH / KD value for FcγRIIb are both 0.7, and accordingly all variants in Table 15 showed improvement of binding selectivity for FcγRIIb in comparison with the parent polypeptide. When the KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant / KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide is 1 or more, this means that the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant has equivalent or reduced binding compared with the binding by the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide. Since this value was 0.7 to 5.0 for the variants obtained this time, one may say that binding by the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the variants obtained this time was nearly the same or decreased in comparison with the parent polypeptide. These results showed that compared with the parent polypeptide, the variants obtained this time have maintained or decreased binding activities to FcγRIIa type R and type H and enhanced binding activity to FcγRIIb, and thus have improved selectivity for FcγRIIb. Furthermore, compared with IL6R-F11, all variants had lower affinity to FcγRIa and FcγRIIIaV.

[Table 15]

ALTERATION	KD [mol/L]				KD[IIaR]/ KD[IIb]	KD[IIaR]/ KD[IIb]	KD (IIb) OF PARENT POLYPEPTIDE/ KD (IIb) OF ALTERED POLYPEPTIDE	KD VALUE FOR THE STRONGER OF THE BINDING ACTIVITIES OF A VARIANT TO FcγRIIaR AND FcγRIIaH/ KD VALUE FOR THE STRONGER OF THE BINDING ACTIVITIES OF THE PARENT POLYPEPTIDE TO FcγRIIaR AND FcγRIIaH
	FcγRIa	FcγRIIaR	FcγRIIaH	FcγRIIb	FcγRIIaV			
*	3.2E-10	1.0E-06	6.7E-07	2.6E-06	3.5E-07	0.4	0.3	0.1
	9.0E-10	5.0E-06	5.0E-06	6.8E-06	1.5E-06	0.7	0.7	1.0
L234W/P238D	6.3E-08	1.6E-05	1.9E-05	2.0E-06	3.7E-05	8.1	9.5	3.2
L234Y/P238D	7.5E-08	2.6E-05	2.3E-05	1.6E-06	4.5E-05	15.9	14.4	4.6
G237A/P238D	1.4E-07	3.2E-05	2.1E-05	3.0E-06	3.7E-05	10.5	7.0	4.2
G237D/P238D	1.4E-07	2.1E-05	2.5E-05	2.0E-06	4.3E-05	10.7	12.8	4.2
G237E/P238D	3.4E-07	3.8E-05	2.5E-05	3.6E-06	4.1E-05	10.6	7.0	5.0
G237F/P238D	5.2E-08	1.4E-05	1.6E-05	3.4E-06	4.3E-05	4.1	4.7	2.8
G237L/P238D	1.2E-07	1.8E-05	1.8E-05	2.6E-06	4.1E-05	6.9	7.1	3.5
G237M/P238D	5.2E-08	2.2E-05	2.0E-05	2.9E-06	3.7E-05	7.7	7.0	4.0
G237W/P238D	3.6E-08	7.2E-06	1.2E-05	2.3E-06	3.8E-05	3.1	5.2	1.4
G237Y/P238D	9.3E-08	7.9E-06	1.5E-05	2.3E-06	4.2E-05	3.4	6.4	1.6
P238D/S239D	4.9E-09	3.5E-06	1.9E-05	1.4E-06	1.7E-05	2.6	14.0	0.7

[Example 10] X-ray crystal structure analysis of a complex formed between an Fc containing P238D and an extracellular region of FcγRIIb

As indicated earlier in Example 9, even though an alteration that is predicted from the analysis of naturally-occurring IgG1 antibodies to improve FcγRIIb-binding activity or selectivity for FcγRIIb is introduced into an Fc containing P238D, the FcγRIIb-binding activity was found to decrease, and the reason for this may be that the structure at the interacting interface between Fc and FcγRIIb is changed due to introduction of P238D. Therefore, to pursue the reason for this phenomena, the three-dimensional structure of the complex formed between an IgG1 Fc containing the P238D mutation (hereinafter, referred to as Fc (P238D)) and the extracellular region of FcγRIIb was elucidated by X-ray crystal structure analysis, and this was compared to the three-dimensional structure of the complex formed between the Fc of a naturally-occurring IgG1 (hereinafter, referred to as Fc (WT)) and the extracellular region of FcγRIIb, and the binding modes were compared. Multiple reports have been made on the three-dimensional structure of a complex formed between an Fc and an FcγR extracellular region; and the three-dimensional structures of the Fc (WT) / FcγRIIb extracellular region complex (Nature (2000) 400, 267-273; J. Biol. Chem. (2011) 276, 16469-16477), the Fc (WT) / FcγRIIIa extracellular region complex (Proc. Natl. Acad. Sci. USA (2011) 108, 12669-126674), and the Fc (WT) / FcγRIIa extracellular region complex (J. Immunol. (2011) 187, 3208-3217) have been analyzed. While the three-dimensional structure of the Fc (WT) / FcγRIIb extracellular region complex has not been analyzed, the three-dimensional structure of a complex formed with Fc (WT) is known for FcγRIIa and FcγRIIb, and their extracellular regions match 93% in amino acid sequence and have very high homology. Thus, the three-dimensional structure of the Fc (WT) / FcγRIIb extracellular region complex was predicted by modeling using the crystal structure of the Fc (WT) / FcγRIIa extracellular region complex.

The three-dimensional structure of the Fc (P238D) / FcγRIIb extracellular region complex was determined by X-ray crystal structure analysis at 2.6 Å resolution. The structure obtained as a result of this analysis is shown in Fig. 26. The FcγRIIb extracellular region is bound between two Fc CH2 domains, and this was similar to the three-dimensional structures of complexes formed between Fc (WT) and the respective extracellular region of FcγRIIIa, FcγRIIb, or FcγRIIa analyzed so far. Next, for detailed comparison, the crystal structure of the Fc (P238D) / FcγRIIb extracellular region complex and the model structure of the Fc (WT) / FcγRIIb extracellular region complex were superimposed by the least squares fitting based on the Cα atom pair distances with respect to the FcγRIIb extracellular region and the Fc CH2 domain A (Fig. 27). In that case, the degree of overlap between Fc CH2 domains B was not satisfactory, and conformational differences were found in this portion. Furthermore, using the crystal structure of the Fc (P238D) / FcγRIIb extracellular region complex and the model

structure of the Fc (WT) / FcγRIIb extracellular region complex, pairs of atoms that have a distance of 3.7 Å or less between the extracted FcγRIIb extracellular region and Fc CH2 domain B were extracted and compared in order to compare the interatomic interaction between FcγRIIb and Fc (WT) CH2 domain B with the interatomic interaction between FcγRIIb and Fc (P238D).

- 5 As shown in Table 16, the interatomic interactions between Fc CH2 domain B and FcγRIIb in Fc (P238D) and Fc (WT) did not match.

[Table 16]

FcγRIIb ATOM	Fc (P238D) CH2 DOMAIN B INTERACTION PARTNER (DISTANCE BETWEEN ATOMS, Å)	Fc(WT) CH2 DOMAIN B INTERACTION PARTNER (DISTANCE BETWEEN ATOMS, Å)
Val 116 CG2		Asp 265 OD2 (3.47) Gly 237 O (3.65)
Ser 126 OG	Ser 298 N (3.31) Ser 298 CB (3.32) Tyr 296 O (3.05)	
Lys 128 CA	Ser 298 OG (3.50)	
Phe 129 CB	Ser 298 O (3.36)	
Phe 129 CD2		Asn 297 CB (3.50) Asn 297 CG (3.43)
Lys 128 C	Ser 298 OG (3.47)	
Phe 129 N	Ser 298 OG (3.30)	
Phe 129 O	Ser 267 OG (3.54)	
Arg 131 CB		Val 266 O (3.02)
Arg 131 CG		Val 266 O (3.22)
Arg 131 CD		Val 266 CG1 (3.45) Val 266 C (3.55) Val 266 O (3.10)
Arg 131 NE	Ala 327 O (3.60)	Val 266 C (3.66) Val 266 O (3.01) Val 266 N (3.49)
Arg 131 CZ	Asp 270 CG (3.64) Asp 270 OD2 (3.22) Asp 270 OD1 (3.27) Ala 327 CB (3.63)	Val 266 N (3.13)
Arg 131 NH1	Asp 270 CG (3.19) Asp 270 OD2 (2.83) Asp 270 OD1 (2.99) Ser 267 CB (3.56)	Val 266 CG1 (3.47) Val 266 N (3.43) Thr 299 OG1 (3.66) Ser 298 O (3.11)
Arg 131 NH2	Asp 270 CG (3.20) Asp 270 OD2 (2.80) Asp 270 OD1 (2.87) Ala 327 CB (3.66)	Asp 265 CA (3.16) Val 266 N (3.37)
Tyr 157 CE1		Leu 234 CG (3.64) Leu 234 CD1 (3.61)
Tyr 157 OH		Gly 236 O (3.62) Leu 234 CA (3.48) Leu 234 CG (3.45)

Furthermore, the detailed structures around P238D were compared by superposing the X-ray crystal structure of Fc (P238D)/Fc γ RIIb extracellular region complex on the model structure of the Fc (WT)/Fc γ RIIb extracellular region complex using the least squares method based on the C α atomic distance between Fc CH2 domains A and B alone. As the position of the amino acid residue at position 238 (EU numbering), *i.e.*, a mutagenesis position of Fc (P238D), is altered from Fc (WT), the loop structure around the amino acid residue at position 238 following the hinge region is found to be different between Fc (P238D) and Fc (WT) (Fig. 28). Pro at position 238 (EU numbering) is originally located inside Fc (WT), forming a hydrophobic core with residues around position 238. However, if Pro at position 238 (EU numbering) is altered to highly hydrophilic and charged Asp, the presence of the altered Asp residue in a hydrophobic core is energetically disadvantageous in terms of desolvation. Therefore, in Fc (P238D), to cancel this energetically disadvantageous situation, the amino acid residue at position 238 (EU numbering) changes its orientation to face the solvent side, and this may have caused this change in the loop structure near the amino acid residue at position 238. Furthermore, since this loop is not far from the hinge region crosslinked by an S-S bond, its structural change will not be limited to a local change, and will affect the relative positioning of the FcCH2 domain A and domain B. As a result, the interatomic interactions between Fc γ RIIb and Fc CH2 domain B have been changed. Therefore, predicted effects could not be observed when alterations that improve selectivity and binding activity towards Fc γ RIIb in a naturally-occurring IgG were combined with an Fc containing the P238D alteration.

Furthermore, as a result of structural changes due to introduction of P238D in Fc CH2 domain A, a hydrogen bond has been found between the main chain of Gly at position 237 (EU numbering), which is adjacent to P238D mutated, and Tyr at position 160 in Fc γ RIIb (Fig. 29). The residue in Fc γ RIIa that corresponds to this Tyr 160 is Phe; and when the binding is to Fc γ RIIa, this hydrogen bond is not formed. Considering that the amino acid at position 160 is one of the few differences between Fc γ RIIa and Fc γ RIIb at the interface of interaction with Fc, the presence of this hydrogen bond which is specific to Fc γ RIIb is presumed to have led to improvement of Fc γ RIIb-binding activity and decrease of Fc γ RIIa-binding activity in Fc (P238D), and improvement of its selectivity. Furthermore, in Fc CH2 domain B, an electrostatic interaction is observed between Asp at position 270 (EU numbering) and Arg at position 131 in Fc γ RIIb (Fig. 30). In Fc γ RIIa type H, which is one of the allotypes of Fc γ RIIa, the residue corresponding to Arg at position 131 of Fc γ RIIb is His, and therefore cannot form this electrostatic interaction. This can explain why the Fc (P238D)-binding activity is lowered in Fc γ RIIa type H compared with Fc γ RIIa type R. Observations based on such results of X-ray crystal structure analysis showed that the change of the loop structure near P238D due to P238D

introduction and the accompanying change in the relative domain positioning causes formation of new interactions which is not found in the binding of the naturally-occurring IgG and FcγR, and this could lead to a selective binding profile of P238D variants for FcγRIIb.

5 [Expression and Purification of Fc (P238D)]

An Fc containing the P238D alteration was prepared as follows. First, Cys at position 220 (EU numbering) of hIL6R-IgG1-v1 (SEQ ID NO: 143) was substituted with Ser. Then, genetic sequence of Fc (P238D) from Glu at position 236 (EU numbering) to its C terminal was cloned by PCR. Using this cloned genetic sequence, production of expression vectors, and
 10 expression and purification of Fc (P238D) were carried out according to the method of Reference Examples 1 and 2. Cys at position 220 (EU numbering) forms a disulfide bond with Cys of the L chain in general IgG1. The L chain is not co-expressed when Fc alone is prepared, and therefore, the Cys residue was substituted with Ser to avoid formation of unnecessary disulfide bonds.

15

[Expression and purification of the FcγRIIb extracellular region]

The FcγRIIb extracellular region was prepared according to the method of Reference Example 25.

20 [Purification of the Fc (P238D) / FcγRIIb extracellular region complex]

To 2 mg of the FcγRIIb extracellular region sample obtained for use in crystallization, 0.29 mg of Endo F1 (Protein Science (1996) 5: 2617-2622) expressed and purified from *Escherichia coli* as a glutathione S-transferase fusion protein was added. This was allowed to remain at room temperature for three days in 0.1 M Bis-Tris buffer at pH 6.5, and the N-linked
 25 oligosaccharide was cleaved, except for *N*-acetylglucosamine directly bound to Asn of the FcγRIIb extracellular region. Next, the FcγRIIb extracellular region sample subjected to carbohydrate cleavage treatment, which was concentrated by ultrafiltration with 5000 MWCO, was purified by gel filtration chromatography (Superdex200 10/300) using a column equilibrated in 20 mM HEPES at pH 7.5 containing 0.05 M NaCl. Furthermore, to the obtained
 30 carbohydrate-cleaved FcγRIIb extracellular region fraction, Fc (P238D) was added so that the molar ratio of the FcγRIIb extracellular region would be present in slight excess. The mixture concentrated by ultrafiltration with 10,000 MWCO was subjected to purification by gel filtration chromatography (Superdex200 10/300) using a column equilibrated in 20 mM HEPES at pH 7.5 containing 0.05 M NaCl. Thus, a sample of the Fc (P238D) / FcγRIIb extracellular region
 35 complex was obtained.

[Crystallization of the Fc (P238D) / FcγRIIb extracellular region complex]

Using the sample of the Fc (P238D) / FcγRIIb extracellular region complex which was concentrated to approximately 10 mg/mL by ultrafiltration with 10,000 MWCO, crystallization of the complex was carried out by the sitting drop vapor diffusion method. Hydra II Plus One (MATRIX) was used for crystallization; and for a reservoir solution containing 100 mM Bis-Tris pH 6.5, 17% PEG3350, 0.2 M ammonium acetate, and 2.7% (w/v) D-Galactose, a crystallization drop was produced by mixing at a ratio of reservoir solution : crystallization sample = 0.2 μl : 0.2 μl. The crystallization drop after sealing was allowed to remain at 20°C, and thus thin plate-like crystals were obtained.

[Measurement of X-ray diffraction data from an Fc (P238D) / FcγRIIb extracellular region complex crystal]

One of the obtained single crystals of the Fc (P238D) / FcγRIIb extracellular region complex was soaked into a solution of 100 mM Bis-Tris pH 6.5, 20% PEG3350, ammonium acetate, 2.7% (w/v) D-Galactose, 22.5% (v/v) ethylene glycol. The single crystal was fished out of the solution using a pin with attached tiny nylon loop, and frozen in liquid nitrogen. Then, the X-ray diffraction data of the crystal was measured at synchrotron radiation facility Photon Factory BL-1A in High Energy Accelerator Research Organization. During the measurement, the crystal was constantly placed in a nitrogen stream at -178°C to maintain in a frozen state, and a total of 225 X ray diffraction images were collected using Quantum 270 CCD detector (ADSC) attached to a beam line with rotating the crystal 0.8° at a time. Determination of cell parameters, indexing of diffraction spots, and diffraction data processing from the obtained diffraction images were performed using the Xia2 program (CCP4 Software Suite), XDS Package (Wolfgang Kabsch) and Scala (CCP4 Software Suite); and finally, diffraction intensity data of the crystal up to 2.46 Å resolution was obtained. The crystal belongs to the space group P2₁, and has the following cell parameters; a = 48.85 Å, b = 76.01 Å, c = 115.09 Å, α = 90°, β = 100.70°, γ = 90°.

[X ray crystal structure analysis of the Fc (P238D) / FcγRIIb extracellular region complex]

Crystal structure of the Fc (P238D) / FcγRIIb extracellular region complex was determined by the molecular replacement method using the program Phaser (CCP4 Software Suite). From the size of the obtained crystal lattice and the molecular weight of the Fc (P238D) / FcγRIIb extracellular region complex, the number of complexes in the asymmetric unit was predicted to be one. From the structural coordinates of PDB code: 3SGJ which is the crystal structure of the Fc (WT) / FcγRIIIa extracellular region complex, the amino acid residue portions of the A chain positions 239-340 and the B chain positions 239-340 were taken out as separate

coordinates, and they were set respectively as models for searching the Fc CH2 domains. The amino acid residue portions of the A chain positions 341-444 and the B chain positions 341-443 were taken out as a single set of coordinates from the same structural coordinates of PDB code: 3SGJ; and this was set as a model for searching the Fc CH3 domains. Finally, from the

5 structural coordinates of PDB code: 2FCB which is a crystal structure of the FcγRIIb extracellular region, the amino acid residue portions of the A chain positions 6-178 was taken out and set as a model for searching the FcγRIIb extracellular region. The orientation and position of each search model in the crystal lattice were determined in the order of Fc CH3 domain, FcγRIIb extracellular region, and Fc CH2 domain, based on the rotation function and translation

10 function to obtain the initial model for the crystal structure of the Fc (P238D) / FcγRIIb extracellular region complex. When rigid body refinement which moves the two Fc CH2 domains, the two Fc CH3 domains, and the FcγRIIb extracellular region was performed on the obtained initial model, the crystallographic reliability factor, R value became 40.4%, and the Free R value became 41.9% to diffraction intensity data from 25 Å to 3.0 Å at this point.

15 Furthermore, structural refinement using the program Refmac5 (CCP4 Software Suite), and revision of the model to observe the electron density maps whose coefficient have 2Fo-Fc or Fo-Fc, which are calculated based on the experimentally determined structural factor Fo, the calculated structural factor Fc and the calculated phase using the model, was carried out by the Coot program (Paul Emsley). Model refinement was carried out by repeating these steps.

20 Finally, as a result of incorporation of water molecules into the model based on the electron density maps which use 2Fo-Fc or Fo-Fc as the coefficient, and the following refinement, the crystallographic reliability factor, R values and the Free R value of the model containing 4846 non-hydrogen atoms became 23.7% and 27.6% to 24291 diffraction intensity data from 25 Å to 2.6 Å resolution, respectively.

25

[Production of a model structure of the Fc (WT) / FcγRIIb extracellular region complex]

Based on the structural coordinates of PDB code: 3RY6 which is a crystal structure of the Fc (WT) / FcγRIIa extracellular region complex, the Build Mutants function of the Discovery Studio 3.1 program (Accelrys) was used to introduce mutations to match the amino acid

30 sequence of FcγRIIb into FcγRIIa in this structural coordinates. In that case, the Optimization Level was set to High, Cut Radius was set to 4.5, five models were generated, and the one with the best energy score from among them was set as the model structure for the Fc (WT)/ FcγRIIb extracellular region complex.

35 [Example 11] Analysis of FcγR binding of Fc variants whose alteration sites were determined based on crystal structures

Based on the results of X-ray crystal structure analysis on the complex formed between Fc (P238D) and the FcγRIIb extracellular region obtained in Example 10, variants were constructed by comprehensively introducing alterations into sites on the altered Fc having substitution of Pro at position 238 (EU numbering) with Asp that were predicted to affect interaction with FcγRIIb (residues of positions 233, 240, 241, 263, 265, 266, 267, 268, 271, 273, 295, 296, 298, 300, 323, 325, 326, 327, 328, 330, 332, and 334 (EU numbering)), and whether combinations of alterations that further enhance FcγRIIb binding in addition to the P238D alteration can be obtained, was examined.

IL6R-B3 (SEQ ID NO: 144) was produced by introducing into IL6R-G1d (SEQ ID NO: 134), the alteration produced by substituting Lys at position 439 (EU numbering) with Glu. Next, IL6R-BF648 was produced by introducing into IL6R-B3, the alteration produced by substituting Pro at position 238 (EU numbering) with Asp. IL6R-L (SEQ ID NO: 135) was utilized as the common antibody L chain. These antibody variants expressed were purified according to the method of Reference Example 2. The binding of these antibody variants to each of the FcγRs (FcγRIa, FcγRIIa type H, FcγRIIa type R, FcγRIIb, and FcγRIIIa type V) was comprehensively evaluated by the method of Reference Example 25.

A figure was produced according to the following method to show the results of analyzing the interactions with the respective FcγRs. The value for the amount of binding of each variant to each FcγR was divided by the value for the amount of binding of the pre-altered control antibody (IL6R-BF648/IL6R-L, alteration by substituting Pro at position 238 (EU numbering) with Asp) to each FcγR, and the obtained was then multiplied by 100 and shown as the relative binding activity value of each variant to each FcγR. The horizontal axis shows the relative binding activity value of each variant to FcγRIIb, and the vertical axis shows the relative binding activity value of each variant to FcγRIIa type R (Fig. 31).

As shown in Fig. 31, the results show that of all the alterations, 24 types of alterations were found to maintain or enhance FcγRIIb binding in comparison with the pre-altered antibody. The binding of these variants to each of the FcγRs are shown in Table 17. In the table, alteration refers to the alteration introduced into IL6R-B3 (SEQ ID NO: 144). The template used for producing IL6R-B3, IL6R-G1d/IL6R-L, is indicated with an asterisk (*).

[Table 17]

VARIANT NAME	ALTERATION	RELATIVE BINDING				
		FcyRIa	FcyRIIaR	FcyRIIaH	FcyRIIb	FcyRIIIa
IL6R-G1d/IL6R-L	*	140	650	1670	62	3348
IL6R-B3/IL6R-L		145	625	1601	58	3264
IL6R-BF648/IL6R-L	P238D	100	100	100	100	100
IL6R-2B002/IL6R-L	P238D/E233D	118	103	147	116	147
IL6R-BP100/IL6R-L	P238D/S267A	121	197	128	110	138
IL6R-BP102/IL6R-L	P238D/S267Q	104	165	66	106	86
IL6R-BP103/IL6R-L	P238D/S267V	56	163	69	107	77
IL6R-BP106/IL6R-L	P238D/H268D	127	150	110	116	127
IL6R-BP107/IL6R-L	P238D/H268E	123	147	114	118	129
IL6R-BP110/IL6R-L	P238D/H268N	105	128	127	101	127
IL6R-BP112/IL6R-L	P238D/P271G	119	340	113	157	102
IL6R-2B128/IL6R-L	P238D/Y296D	95	87	37	103	96
IL6R-2B169/IL6R-L	P238D/V323I	73	92	83	104	94
IL6R-2B171/IL6R-L	P238D/V323L	116	117	115	113	122
IL6R-2B172/IL6R-L	P238D/V323M	140	244	179	132	144
IL6R-BP136/IL6R-L	P238D/K326A	117	159	103	119	102
IL6R-BP117/IL6R-L	P238D/K326D	124	166	96	118	105
IL6R-BP120/IL6R-L	P238D/K326E	125	175	92	114	103
IL6R-BP126/IL6R-L	P238D/K326L	113	167	132	103	146
IL6R-BP119/IL6R-L	P238D/K326M	117	181	133	110	145
IL6R-BP142/IL6R-L	P238D/K326N	98	103	97	106	102
IL6R-BP121/IL6R-L	P238D/K326Q	118	155	135	113	157
IL6R-BP118/IL6R-L	P238D/K326S	101	132	128	104	144
IL6R-BP116/IL6R-L	P238D/K326T	110	126	110	108	114
IL6R-BP911/IL6R-L	P238D/A330K	52	101	108	119	120
IL6R-BP078/IL6R-L	P238D/A330M	106	101	89	105	91
IL6R-BP912/IL6R-L	P238D/A330R	60	81	93	103	97

The results of measuring KD values of the variants shown in Table 17 for FcγRIa, FcγRIIaR, FcγRIIaH, FcγRIIb, and FcγRIIIa V types by the method of Reference Example 25 are summarized in Table 18. In the table, alteration refers to the alteration introduced into IL6R-B3 (SEQ ID NO: 144). The template used for producing IL6R-B3, IL6R-G1d/IL6R-L, is indicated with an asterisk (*). Furthermore, KD (IIaR)/KD (IIb) and KD (IIaH)/KD (IIb) in the table respectively represent the value obtained by dividing the KD value of each variant for FcγRIIaR by the KD value of each variant for FcγRIIb, and the value obtained by dividing the KD value of each variant for FcγRIIaH by the KD value of each variant for FcγRIIb. KD (IIb) of the parent polypeptide / KD (IIb) of the altered polypeptide refers to the value obtained by dividing the KD value of the parent polypeptide for FcγRIIb by the KD value of each variant for FcγRIIb. In addition, the KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of each variant / KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide are shown in Table 18. Here, parent polypeptide refers to the variant which has IL6R-B3 (SEQ ID NO: 144) as the H chain. It was determined that due to weak binding of FcγR to IgG, it was impossible to accurately analyze by kinetic analysis, and thus the gray-filled cells in Table 18 show values calculated by using Equation 2 of Reference Example 25.

[Equation 2]

$$KD = C \bullet R_{\max} / (R_{\text{eq}} - RI) - C$$

[Table 18]

VARIANT NAME	ALTERATION	KD[μmol/L]					FcγRIIaH/ KD[μb]	FcγRIIb/ KD[μb]	FcγRIIaV	FcγRIIb	FcγRIIaH/ KD[μb]	KD[μb] OF THE PARENT POLYPEPTIDE / KD[μb] OF THE ALTERED POLYPEPTIDE	KD VALUE FOR THE STRONGER OF THE FcγRIIaR- AND FcγRIIaH- BINDING ACTIVITIES OF THE VARIANT / KD VALUE FOR THE STRONGER OF THE FcγRIIaR- AND FcγRIIaH-BINDING ACTIVITIES OF THE PARENT POLYPEPTIDE
		FcγRIa	FcγRIIaR	FcγRIIaH	FcγRIIb	FcγRIIaV							
IL6R-B37/IL6R-L		3.2E-10	1.0E-05	6.7E-07	2.6E-06	3.5E-07	0.4	0.3	1.2	0.9			
IL6R-B37/IL6R-L		4.2E-10	1.1E-06	7.7E-07	3.1E-06	3.3E-07	0.3	0.2	1.0	1.0			
IL6R-BF648/IL6R-L		1.1E-08	1.5E-05	4.0E-05	1.2E-05	7.1E-05	13.0	33.9	2.6	19.9			
IL6R-2B002/IL6R-L		6.4E-09	1.9E-05	8.6E-05	9.3E-07	5.3E-05	20.4	92.3	3.3	24.7			
IL6R-BF100/IL6R-L		1.1E-09	7.8E-06	4.6E-05	1.1E-06	5.9E-05	7.3	42.8	2.9	10.2			
IL6R-BF102/IL6R-L		8.2E-09	8.4E-06	6.1E-05	9.0E-07	8.2E-05	9.4	67.6	3.4	11.0			
IL6R-BF103/IL6R-L		3.5E-08	1.1E-05	8.8E-05	1.2E-06	1.1E-04	9.0	71.5	2.5	14.4			
IL6R-BF105/IL6R-L		4.0E-09	1.1E-05	3.6E-05	9.3E-07	5.5E-05	11.6	38.7	3.3	14.0			
IL6R-BF107/IL6R-L		1.5E-09	1.2E-05	5.2E-05	9.3E-07	6.3E-05	12.7	56.1	3.3	15.3			
IL6R-BF110/IL6R-L		7.3E-09	1.7E-05	4.7E-05	1.5E-06	6.4E-05	11.7	31.5	2.1	22.6			
IL6R-BF112/IL6R-L		6.3E-09	3.5E-06	3.5E-05	3.2E-07	8.9E-05	11.0	109.4	9.7	4.6			
IL6R-2B128/IL6R-L		1.3E-08	2.6E-05	3.4E-05	1.4E-06	7.2E-05	17.7	23.6	2.1	33.1			
IL6R-2B169/IL6R-L		2.5E-08	1.9E-05	4.8E-05	1.2E-06	7.5E-05	15.8	40.7	2.6	20.8			
IL6R-2B171/IL6R-L		9.1E-09	1.6E-05	3.4E-05	1.1E-06	5.7E-05	15.0	31.8	2.9	8.0			
IL6R-2B172/IL6R-L		3.0E-09	5.1E-06	2.1E-05	7.7E-07	4.8E-05	8.0	27.3	4.0	11.8			
IL6R-BF136/IL6R-L		6.8E-09	9.1E-06	3.8E-05	8.0E-07	6.9E-05	11.4	47.6	3.9	12.0			
IL6R-BF117/IL6R-L		4.1E-09	9.2E-06	4.1E-05	8.0E-07	6.7E-05	11.6	51.4	3.9	12.5			
IL6R-BF120/IL6R-L		6.6E-09	9.6E-06	6.5E-05	1.0E-06	7.9E-05	9.3	63.1	3.0	14.4			
IL6R-BF126/IL6R-L	7.4E-09	1.1E-05	4.5E-05	1.4E-06	5.6E-05	7.8	31.7	2.2	12.8				
IL6R-BF119/IL6R-L	7.0E-09	9.9E-06	4.5E-05	1.1E-06	5.6E-05	8.7	39.5	2.7	23.5				
IL6R-BF142/IL6R-L	5.3E-09	1.8E-05	9.3E-05	1.2E-06	1.1E-04	15.5	79.5	2.6	16.6				
IL6R-BF121/IL6R-L	1.1E-08	1.3E-05	4.4E-05	1.1E-06	5.2E-05	11.7	40.4	2.8	19.7				
IL6R-BF113/IL6R-L	1.2E-08	1.5E-05	4.6E-05	1.2E-06	5.6E-05	13.2	40.0	2.7	19.4				
IL6R-BF116/IL6R-L	2.6E-09	1.5E-05	5.4E-05	1.1E-06	7.2E-05	13.3	48.2	2.8	21.3				
IL6R-BF911/IL6R-L	4.9E-08	1.6E-05	3.7E-05	8.9E-07	5.8E-05	18.5	41.7	3.5	19.0				
IL6R-BF078/IL6R-L	8.2E-09	1.5E-05	4.5E-05	1.1E-06	7.8E-05	13.4	41.3	2.8	34.0				
IL6R-BF912/IL6R-L	3.8E-08	2.5E-05	3.8E-05	1.5E-06	7.8E-05	17.8	25.9	2.1					

Table 18 shows that in comparison with IL6R-B3, all variants showed improvement of affinity for FcγRIIb, and the range of improvement was 2.1 fold to 9.7 fold. The ratio of KD value of each variant for FcγRIIaR / KD value of each variant for FcγRIIb, and the ratio of KD value of each variant for FcγRIIaH / KD value of each variant for FcγRIIb represent an FcγRIIb-binding activity relative to the FcγRIIaR-binding activity and FcγRIIaH-binding activity, respectively. That is, these values show the degree of binding selectivity of each variant for FcγRIIb, and a greater value indicates a higher binding selectivity for FcγRIIb. Since the ratio of KD value for FcγRIIaR / KD value for FcγRIIb, and the ratio of KD value for FcγRIIaH / KD value for FcγRIIb in the parent polypeptide IL6R-B3/IL6R-L were 0.3 and 0.2, respectively, all variants in Table 18 showed improvement of binding selectivity for FcγRIIb in comparison with the parent polypeptide. When the KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant / KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide is 1 or more, this means that the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant has equivalent or decreased binding compared with the binding by the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide. Since this value was 4.6 to 34.0 for the variants obtained this time, one may say that in comparison with the parent polypeptide, the variants obtained this time had reduced binding by the stronger of the FcγRIIaR- and FcγRIIaH-binding activities. These results showed that compared with the parent polypeptide, the variants obtained this time have maintained or decreased FcγRIIa type R- and type H-binding activities, enhanced FcγRIIb-binding activity, and improved selectivity for FcγRIIb. Furthermore, compared with IL6R-B3, all variants had lower affinity to FcγRIa and FcγRIIIaV.

With regard to the promising variants among the obtained combination variants, the factors leading to their effects were studied using the crystal structure. Fig. 32 shows the crystal structure of the Fc (P238D) / FcγRIIb extracellular region complex. In this figure, the H chain positioned on the left side is Fc Chain A, and the H chain positioned on the right side is Fc Chain B. Here, one can see that the site at position 233 (EU numbering) in Fc Chain A is located near Lys at position 113 of FcγRIIb. However, in this crystal structure, the E233 side chain is in a condition of considerably high mobility, and its electron density is not well observed. Therefore, the alteration produced by substituting Glu at position 233 (EU numbering) with Asp leads to decrease in the degree of freedom of the side chain since the side chain becomes one carbon shorter. As a result, the entropy loss when forming an interaction with Lys at position 113 of FcγRIIb may be decreased, and consequently this is speculated to contribute to improvement of binding free energy.

Similarly, Fig. 33 shows the environment near the site at position 330 (EU numbering) in the structure of the Fc (P238D) / FcγRIIb extracellular region complex. This figure shows

that the environment around the site at position 330 (EU numbering) of Fc Chain A of Fc (P238D) is a hydrophilic environment composed of Ser at position 85, Glu at position 86, Lys at position 163, and such of FcγRIIb. Therefore, the alteration produced by substituting Ala at position 330 (EU numbering) with Lys or Arg is speculated to contribute to strengthening the interaction with Ser at position 85 or Glu at position 86 in FcγRIIb.

Fig. 34 depicts the structures of Pro at position 271 (EU numbering) of Fc Chain B after superimposing the crystal structures of the Fc (P238D) / FcγRIIb extracellular region complex and the Fc (WT) / FcγRIIIa extracellular region complex by the least squares fitting based on the Cα atom pair distances with respect to Fc Chain B. These two structures match well, but have different three-dimensional structures of Pro at position 271 (EU numbering). When the weak electron density around this area in the crystal structure of the Fc (P238D)/FcγRIIb extracellular region complex is also taken into consideration, it is suggested that there is possibility that Pro at position 271 (EU numbering) in Fc (P238D) / FcγRIIb causes a large strain on the structure, thus disturbing the loop structure to attain an optimal structure. Therefore, the alteration produced by substituting Pro at position 271 (EU numbering) with Gly gives flexibility to this loop structure, and is speculated to contribute to enhancement of binding by reducing the energetic barrier when allowing an optimum structure to form during interaction with FcγRIIb.

[Example 12] Examination of the combinatorial effect of alterations that enhance FcγRIIb binding when combined with P238D

Of the alterations obtained in Examples 9 and 11, those that enhanced FcγRIIb binding or maintained FcγRIIb binding and showed effects of suppressing binding to other FcγRs were combined with each other, and its effect was examined.

Particularly good alterations selected from Tables 14 and 18 were introduced into the antibody H chain IL6R-BF648 in a similar manner to the method of Example 11. IL6R-L was utilized as the antibody L chain, the expressed antibodies were purified according to the method of Reference Example 1. The binding to each of the FcγRs (FcγRIa, FcγRIIa H type, FcγRIIa R type, FcγRIIb, and FcγRIIIa V type) was comprehensively evaluated by the method of Reference Example 25.

According to the following method, relative binding activities were calculated for the results of analyzing interactions with the respective FcγRs. The value for the amount of binding of each variant to each FcγR was divided by the value for the amount of binding of the pre-altered control antibody (IL6R-BF648/IL6R-L with substitution of Pro at position 238 (EU numbering) with Asp to each FcγR, and multiplied by 100; and then the value was shown as the relative binding activity value of each variant to each FcγR (Table 19).

In the table, alteration refers to the alteration introduced into IL6R-B3 (SEQ ID NO:

144). The template used for producing IL6R-B3, IL6R-G1d/IL6R-L, is indicated with an asterisk (*).

[Table 19-1]

VARIANT NAME	ALTERATION	RELATIVE BINDING ACTIVITY				
		FcgRIa	FcgRIIaR	FcgRIIaH	FcgRIIb	FcgRIIaV
IL6R-G1d/IL6R-L		140	650	1670	62	3348
IL6R-E3/IL6R-L		145	625	1601	56	3264
IL6R-BF648/IL6R-L	P238D					
IL6R-2B253/IL6R-L	P233D/P238D/V323M	100	100	100	100	100
IL6R-2B261/IL6R-L	P233D/P238D/Y296D	155	268	207	156	126
IL6R-BP062/IL6R-L	P233D/P238D/A330K	100	94	91	115	87
IL6R-BP063/IL6R-L	P238D/Y296D/A330K	74	126	106	136	87
IL6R-BP064/IL6R-L	P238D/V323M/A330K	50	87	91	123	107
IL6R-BP065/IL6R-L	P238D/P238D/A330K	109	203	162	141	106
IL6R-BP066/IL6R-L	P238D/K326A/A330K	19	279	158	152	104
IL6R-BP067/IL6R-L	P238D/V323M/A330K	72	155	116	137	123
IL6R-BP068/IL6R-L	P238D/P238D/A330K	33	163	179	137	158
IL6R-BP069/IL6R-L	P238D/K326A/A330K	25	377	166	161	122
IL6R-BP129/IL6R-L	P233D/P238D/Y296D/A330K	43	222	186	147	136
IL6R-BP130/IL6R-L	P233D/P238D/V323M/A330K	66	111	98	138	95
IL6R-BP131/IL6R-L	P233D/G237D/P238D/A330K	104	272	224	160	115
IL6R-BP132/IL6R-L	P233D/P238D/K326A/A330K	33	364	253	160	118
IL6R-BP133/IL6R-L	P233D/L234Y/P238D/A330K	91	191	130	150	120
IL6R-BP143/IL6R-L	P234Y/P238D/K326A	41	174	151	137	114
IL6R-BP144/IL6R-L	P237D/P238D/K326A	86	238	143	133	114
IL6R-BP145/IL6R-L	L234Y/G237D/P238D	64	304	108	121	128
IL6R-BP146/IL6R-L	L234Y/G237D/P238D/K326A	41	350	224	152	153
IL6R-BP147/IL6R-L	L234Y/G237D/P238D/K326A/A330K	50	445	203	156	180
IL6R-BP148/IL6R-L	P233D/L234Y/G237D/P238D/K326A/A330K	24	650	582	177	309
IL6R-BP149/IL6R-L	P233D/L234Y/G237D/P238D/Y296D/K326A/A330K	33	603	462	176	227
IL6R-BP150/IL6R-L	L234Y/G237D/P238D/K326A/A330K	29	539	401	173	186
IL6R-BP151/IL6R-L	P233D/L234Y/G237D/P238D/K326A/A330K	30	757	770	183	204
IL6R-BP152/IL6R-L	P233D/L234Y/G237D/P238D/Y296D/K326A/A330K	39	705	621	180	221
IL6R-BP176/IL6R-L	P233D/P238D/K326D/A330K	34	638	548	178	146
IL6R-BP177/IL6R-L	P233D/L234Y/G237D/P238D/P271G/K326D/A330K	102	201	128	147	131
IL6R-BP178/IL6R-L	P233D/G237D/P238D/P271G/A330K	57	691	409	177	186
IL6R-BP179/IL6R-L	G237D/P238D/P271G/K326A/A330K	51	653	359	179	110
IL6R-BP180/IL6R-L	K327D/P238D/P271G/A330K	39	570	226	177	125
IL6R-BP180/IL6R-L	K327D/P238D/P271G/A330K	28	602	203	179	100

Table 19-2 is a continuation table of Table 19-1.

[Table 19-2]

U6R-BP181/U6R-L	E233D/P238D/P271G/K326A/A330K	108	362	150	170	122
U6R-BP182/U6R-L	E233D/P238D/P271G/Y296D/A330K	95	413	139	173	120
U6R-BP183/U6R-L	E233D/L234Y/P238D/P271G/K326A/A330K	83	423	191	164	113
U6R-BP184/U6R-L	E233D/P238D/P271G/A330K	96	436	131	171	106
U6R-BP185/U6R-L	E233D/L234Y/G237D/P238D/P271G/K326A/A330K	47	670	446	179	191
U6R-BP186/U6R-L	E233D/L234Y/G237D/P238D/P271G/Y296D/K326A/A330K	43	614	368	175	143
U6R-BP187/U6R-L	L234Y/P238D/P271G/K326A/A330K	68	387	206	157	124
U6R-BP188/U6R-L	E233D/G237D/P238D/H268D/P271G/A330K	74	636	234	179	131
U6R-BP189/U6R-L	G237D/P238D/H268D/P271G/K326A/A330K	56	557	163	177	141
U6R-BP190/U6R-L	G237D/P238D/H268D/P271G/A330K	50	615	234	181	155
U6R-BP191/U6R-L	E233D/P238D/H268D/P271G/K326A/A330K	125	362	145	170	142
U6R-BP192/U6R-L	E233D/P238D/H268D/P271G/Y296D/A330K	109	406	123	173	118
U6R-BP193/U6R-L	E233D/P238D/H268D/P271G/A330K	113	449	154	173	135
U6R-BP194/U6R-L	E233D/L234Y/G237D/P238D/H268D/P271G/K326A/A330K	69	672	395	178	249
U6R-BP195/U6R-L	E233D/L234Y/G237D/P238D/H268D/P271G/Y296D/K326A/A330K	68	661	344	181	221
U6R-BP196/U6R-L	L234Y/P238D/H268D/P271G/K326A/A330K	89	402	195	157	137
U6R-BP197/U6R-L	E233D/L234Y/G237D/P238D/H268D/P271G/Y296D/K326D/A330K	71	642	294	179	206
U6R-BP198/U6R-L	E233D/L234Y/P238D/H268D/P271G/K326A/A330K	104	449	188	164	157
U6R-BP199/U6R-L	E233D/P238D/K326D/A330K	112	172	116	144	103
U6R-BP200/U6R-L	E233D/L234Y/G237D/P238D/P271G/K326D/A330R	60	754	517	188	164
U6R-BP201/U6R-L	E233D/G237D/P238D/P271G/A330K	57	696	359	186	121
U6R-BP202/U6R-L	G237D/P238D/P271G/K326A/A330K	43	615	285	185	106
U6R-BP203/U6R-L	G237D/P238D/P271G/A330K	35	637	255	185	88
U6R-BP204/U6R-L	E233D/P238D/P271G/K326A/A330K	110	301	137	165	121
U6R-BP205/U6R-L	E233D/P238D/P271G/Y296D/A330K	97	335	108	167	93
U6R-BP206/U6R-L	E233D/P238D/P271G/A330K	101	362	123	166	92
U6R-BP207/U6R-L	E233D/P238D/A330K	74	103	103	124	97
U6R-BP208/U6R-L	E233D/G237D/P238D/H268D/P271G/A330K	81	690	310	188	118
U6R-BP209/U6R-L	G237D/P238D/H268D/P271G/K326A/A330K	66	625	267	186	153
U6R-BP210/U6R-L	G237D/P238D/H268D/P271G/A330K	57	661	279	187	135
U6R-BP211/U6R-L	E233D/P238D/H268D/P271G/K326A/A330K	128	312	111	165	87
U6R-BP212/U6R-L	E233D/P238D/H268D/P271G/Y296D/A330K	117	363	135	173	122
U6R-BP213/U6R-L	E233D/P238D/H268D/P271G/A330K	118	362	123	169	100
U6R-BP214/U6R-L	E233D/L234Y/G237D/P238D/Y296D/K326D/A330K	96	498	285	174	165

The results of measuring KD values of the variants shown in Table 19 for FcγRIa,

FcγRIIaR, FcγRIIaH, FcγRIIb, and FcγRIIIaV types by the method of Reference Example 25 are summarized in Tables 20-1 and 20-2. In the table, alteration refers to the alteration introduced into IL6R-B3 (SEQ ID NO: 144). The template used for producing IL6R-B3, IL6R-G1d/IL6R-L, is indicated with an asterisk (*). Furthermore, KD (IIaR)/KD (IIb) and KD (IIaH)/KD (IIb) in the table respectively represent the value obtained by dividing the KD value of the variant for FcγRIIaR by the KD value of the variant for FcγRIIb, and the value obtained by dividing the KD value of the variant for FcγRIIaH by the KD value of each variant for FcγRIIb. KD (IIb) of the parent polypeptide / KD (IIb) of the altered polypeptide refers to the value obtained by dividing the KD value of the parent polypeptide for FcγRIIb by the KD value of each variant for FcγRIIb. In addition, the KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of each variant / KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide are shown in Tables 20-1 and 20-2. Here, parent polypeptide refers to the variant which has IL6R-B3 (SEQ ID NO: 144) as the H chain. It was determined that due to weak binding of FcγR to IgG, it was impossible to accurately analyze by kinetic analysis, and thus the values of gray-filled cells in Tables 20-1 and 20-2 show values calculated by using Equation 2 of Reference Example 25.

[Equation 2]

$$KD = C \bullet R_{\max} / (R_{\text{eq}} - RI) - C$$

Tables 20-1 and 20-2 show that in comparison with IL6R-B3, all variants showed improvement of affinity for FcγRIIb, and the range of improvement was 3.0 fold to 99.0 fold. The ratio of KD value of each variant for FcγRIIaR / KD value of each variant for FcγRIIb, and the ratio of KD value of each variant for FcγRIIaH / KD value of each variant for FcγRIIb represent an FcγRIIb-binding activity relative to the FcγRIIaR-binding activity and FcγRIIaH-binding activity, respectively. That is, those values show the degree of binding selectivity of each variant for FcγRIIb, and a greater value indicates a higher binding selectivity for FcγRIIb. Since the ratio of KD value for FcγRIIaR / KD value for FcγRIIb, and the ratio of KD value for FcγRIIaH / KD value for FcγRIIb of the parent polypeptide IL6R-B3/IL6R-L were 0.3 and 0.2, respectively, all variants in Tables 20-1 and 20-2 showed improvement of binding selectivity for FcγRIIb in comparison with the parent polypeptide. When the KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant / KD value for the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide is 1 or more, this means that the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of a variant has equivalent or decreased binding compared with the binding by the stronger of the FcγRIIaR- and FcγRIIaH-binding activities of the parent polypeptide. Since this value was 0.7 to 29.9 for the variants obtained this time, one may say that binding by the stronger of the FcγRIIaR- and

FcγRIIaH-binding activities of the variants obtained this time was nearly equivalent or decreased compared with that of the parent polypeptide. These results showed that compared with the parent polypeptide, the variants obtained this time have maintained or decreased FcγRIIa type R- and type H-binding activities, enhanced FcγRIIb-binding activity, and improved selectivity for
5 FcγRIIb. Furthermore, compared with IL6R-B3, all variants had lower affinity for FcγRIa and FcγRIIIaV.

[Table 20-1]

VARIANT NAME	ALTERATION	KD (nM) (1)					KD (nM) / KD (nM)	KD (nM) OF POLYPEPTIDE / KD (nM) OF THE ALTERED POLYPEPTIDE	KD VALUE FOR THE STRONGER OF THE FcγRIIaR- AND FcγRIIaH-BINDING ACTIVITIES OF THE VARIANT / KD VALUE FOR THE STRONGER OF THE FcγRIIaR- AND FcγRIIaH-BINDING ACTIVITIES OF THE PARENT POLYPEPTIDE
		FcγRIa	FcγRIIaR	FcγRIIaH	FcγRIIb	FcγRIIaV			
IL6R-G16/IL6R-L		3.2E-10	1.0E-06	6.7E-07	2.6E-06	3.9E-07	0.4	0.3	0.9
IL6R-B3/IL6R-L		4.2E-10	1.1E-06	7.7E-07	3.1E-06	3.3E-07	0.3	0.2	1.0
IL6R-B7E49/IL6R-L	2238D	1.1E-06	1.5E-05	4.0E-05	1.2E-06	7.1E-05	15.0	33.9	2.6
IL6R-B2E63/IL6R-L	2235D/P236D/V236M	1.4E-09	5.0E-06	1.0E-05	4.3E-07	5.0E-05	11.5	30.1	7.2
IL6R-B2E61/IL6R-L	2235D/P236D/Y236D	9.0E-09	2.2E-06	3.3E-05	1.0E-04	7.3E-05	21.8	32.4	3.0
IL6R-BF063/IL6R-L	2235D/P236D/A330K	1.8E-08	1.2E-05	5.7E-05	5.4E-07	5.1E-05	22.8	69.0	5.8
IL6R-BF063/IL6R-L	2238D/Y236D/A330K	9.8E-08	2.3E-05	4.4E-05	7.9E-07	5.6E-05	29.0	56.5	3.9
IL6R-BF064/IL6R-L	2235D/V236M/A330K	7.0E-09	7.2E-06	2.4E-05	5.0E-07	5.7E-05	14.3	47.6	6.1
IL6R-BF065/IL6R-L	2237D/P236D/A330K	2.9E-07	4.2E-06	2.4E-05	3.2E-07	5.6E-05	13.1	74.5	9.6
IL6R-BF066/IL6R-L	2238D/K326A/A330K	2.7E-08	9.7E-06	3.4E-05	5.7E-07	5.7E-05	17.1	59.9	5.4
IL6R-BF068/IL6R-L	2237D/P236D/K326A/A330K	3.9E-07	2.9E-06	2.1E-05	6.1E-07	4.1E-05	16.0	34.7	5.1
IL6R-BF069/IL6R-L	2234V/P236D/K326A/A330K	5.3E-08	6.4E-06	3.0E-05	2.2E-07	5.7E-05	19.3	106.5	14.3
IL6R-BF129/IL6R-L	2235D/P236D/Y236D/A330K	2.5E-08	1.5E-05	4.0E-05	5.2E-07	7.1E-05	16.6	51.9	8.0
IL6R-BF130/IL6R-L	2235D/P236D/V236M/A330K	1.8E-09	5.2E-06	2.6E-05	3.0E-07	7.1E-05	17.5	35.5	10.2
IL6R-BF131/IL6R-L	2237D/G237D/P236D/A330K	1.2E-07	3.1E-06	1.4E-05	2.9E-07	5.3E-05	12.5	56.9	12.6
IL6R-BF132/IL6R-L	2235D/P236D/K326A/A330K	1.5E-08	9.0E-06	5.0E-05	5.7E-07	5.3E-05	21.5	81.1	8.4
IL6R-BF133/IL6R-L	2235D/L234V/P236D/A330K	1.3E-07	8.1E-06	2.6E-05	5.8E-07	5.2E-05	15.5	46.6	5.6
IL6R-BF143/IL6R-L	2235D/P236D/K326A	1.6E-08	5.7E-06	3.7E-05	5.7E-07	5.3E-05	10.0	47.1	5.4
IL6R-BF144/IL6R-L	2237D/P236D/K326A	3.7E-08	6.4E-06	1.7E-05	3.4E-07	4.5E-05	9.9	49.9	9.1
IL6R-BF145/IL6R-L	2234V/G237D/P236D	7.4E-08	2.1E-06	1.8E-05	2.9E-07	3.8E-05	9.3	80.0	15.7
IL6R-BF146/IL6R-L	2234V/G237D/P236D/K326A	1.4E-07	8.9E-07	5.1E-05	6.5E-08	3.5E-05	13.6	77.7	47.1
IL6R-BF147/IL6R-L	2234V/G237D/P236D/K326A/A330K	9.9E-08	1.1E-06	7.0E-05	7.5E-08	3.0E-05	14.5	93.8	41.4
IL6R-BF149/IL6R-L	2235D/L234V/G237D/P236D/K326A/A330K	1.2E-07	1.4E-06	8.4E-05	9.3E-08	3.7E-05	18.0	89.9	33.1
IL6R-BF150/IL6R-L	2234V/G237D/P236D/K326A/A330K	5.4E-07	5.5E-07	3.4E-05	3.1E-08	3.1E-05	17.7	109.0	99.0
IL6R-BF151/IL6R-L	2235D/L234V/G237D/P236D/K326A/A330K	6.4E-08	6.7E-07	4.7E-06	4.0E-08	3.1E-05	16.9	117.8	77.4
IL6R-BF152/IL6R-L	2235D/L234V/G237D/P236D/K326A/A330K	7.3E-08	8.1E-07	5.5E-06	4.1E-08	4.3E-05	19.5	135.9	75.0
IL6R-BF176/IL6R-L	2235D/P236D/A330K	7.3E-09	6.9E-06	3.0E-05	3.6E-07	5.4E-05	19.1	83.1	8.6
IL6R-BF177/IL6R-L	2235D/L234V/G237D/P236D/P271G/K326D/A330K	3.5E-08	7.1E-07	6.2E-05	5.2E-08	3.7E-05	13.8	159.2	60.9
IL6R-BF178/IL6R-L	2235D/G237D/P236D/P271G/A330K	4.3E-08	9.3E-07	1.4E-05	5.1E-08	6.4E-05	18.1	272.4	60.1
IL6R-BF179/IL6R-L	2237D/P236D/P271G/K326A/A330K	5.4E-08	1.4E-06	1.6E-05	8.4E-08	5.6E-05	16.7	190.9	36.9
IL6R-BF180/IL6R-L	2237D/P236D/P271G/A330K	9.8E-08	1.2E-05	1.6E-05	5.2E-08	7.0E-05	18.6	240.8	49.9
IL6R-BF181/IL6R-L	2235D/P236D/P271G/K326A/A330K	7.5E-09	3.2E-06	3.6E-05	1.6E-07	6.7E-05	20.3	162.5	19.3
IL6R-BF182/IL6R-L	2235D/P236D/P271G/P236D/A330K	1.0E-06	2.6E-06	2.8E-05	1.1E-07	5.8E-05	23.5	256.9	28.3

Table 20-2 is a continuation table of Table 20-1.

[Table 20-2]

LAR-BP163/LAR-L	P233D/L234Y/P236D/P271G/K326A/AS30K	1.7E-08	2.6E-06	1.5E-05	2.4E-07	5.6E-05	10.7	62.5	12.9	3.3
LAR-BP164/LAR-L	P233D/P236D/P271G/AS30K	1.1E-08	2.3E-06	9.0E-06	1.3E-07	6.9E-05	18.2	238.1	24.5	3.0
LAR-BP165/LAR-L	P233D/L234Y/G237D/P236D/P271G/K326A/AS30K	6.3E-08	8.8E-07	7.4E-06	6.3E-08	3.6E-05	12.6	106.2	44.5	1.1
LAR-BP166/LAR-L	P233D/L234Y/G237D/P236D/P271G/P296D/K326A/AS30K	4.5E-08	9.6E-07	9.3E-06	6.1E-06	4.9E-05	15.6	152.5	50.7	1.5
LAR-BP167/LAR-L	L234Y/P236D/P271G/K326A/AS30K	2.5E-08	2.6E-06	1.8E-05	2.9E-07	5.6E-05	9.7	62.3	10.7	3.6
LAR-BP168/LAR-L	P233D/G237D/P236D/H266D/P271G/AS30K	2.1E-08	1.0E-06	1.4E-05	4.9E-07	6.4E-05	21.9	350.1	57.6	1.3
LAR-BP169/LAR-L	G237D/P236D/H266D/P271G/K326A/AS30K	4.2E-08	1.4E-06	2.1E-05	7.4E-08	4.9E-05	18.5	263.6	41.6	1.8
LAR-BP170/LAR-L	G237D/P236D/H266D/P271G/K326A/AS30K	6.3E-08	1.1E-06	1.7E-05	5.3E-08	4.9E-05	19.3	203.6	53.2	1.5
LAR-BP171/LAR-L	G237D/P236D/H266D/P271G/K326A/AS30K	4.0E-08	3.0E-06	2.7E-05	1.9E-07	4.9E-05	20.3	184.9	21.2	3.6
LAR-BP172/LAR-L	P233D/P236D/H266D/P271G/P296D/AS30K	5.5E-09	2.6E-06	3.2E-05	1.1E-07	5.9E-05	23.1	283.2	27.3	3.4
LAR-BP173/LAR-L	P233D/P236D/H266D/P271G/AS30K	5.3E-09	2.3E-06	2.5E-05	1.2E-07	5.2E-05	18.3	206.6	25.5	2.9
LAR-BP174/LAR-L	P233D/L234Y/G237D/P236D/H266D/P271G/K326A/AS30K	2.4E-08	6.2E-07	6.9E-06	5.2E-06	2.7E-05	15.6	163.5	59.4	1.1
LAR-BP175/LAR-L	P233D/L234Y/G237D/P236D/H266D/P271G/P296D/K	2.3E-08	9.1E-07	1.0E-05	5.0E-08	3.1E-05	19.2	200.6	62.0	1.2
LAR-BP176/LAR-L	L234Y/P236D/H266D/P271G/K326A/AS30K	1.4E-08	3.0E-06	1.9E-05	2.2E-07	5.1E-05	13.4	85.2	13.9	9.9
LAR-BP177/LAR-L	P233D/L234Y/G237D/P236D/H266D/P271G/K326A/AS30K	1.9E-08	9.9E-07	1.2E-05	5.8E-08	3.3E-05	17.1	208.7	53.7	1.3
LAR-BP178/LAR-L	P233D/L234Y/P236D/H266D/P271G/K326A/AS30K	1.1E-08	2.2E-06	2.0E-05	2.0E-07	4.4E-05	11.0	101.5	15.7	2.8
LAR-BP179/LAR-L	P233D/P236D/K326D/AS30K	6.4E-09	6.9E-06	2.8E-05	4.9E-07	6.1E-05	17.5	53.0	6.3	11.1
LAR-BP180/LAR-L	P233D/L234Y/G237D/P236D/P271G/K326D/AS30K	3.3E-08	6.3E-07	4.2E-06	3.4E-08	3.8E-05	18.6	123.9	91.2	0.6
LAR-BP181/LAR-L	P233D/G237D/P236D/P271G/AS30K	5.1E-08	6.4E-07	5.9E-06	4.0E-08	5.2E-05	21.0	172.1	77.1	1.1
LAR-BP182/LAR-L	G237D/P236D/P271G/K326A/AS30K	9.5E-08	1.2E-06	9.2E-06	6.4E-08	5.9E-05	19.2	144.0	46.4	1.6
LAR-BP183/LAR-L	G237D/P236D/P271G/AS30K	1.2E-07	9.9E-07	1.1E-05	4.9E-08	7.2E-05	20.5	226.6	63.7	1.3
LAR-BP184/LAR-L	P233D/P236D/P271G/K326A/AS30K	7.6E-09	4.5E-06	2.1E-05	2.5E-07	5.2E-05	17.6	82.7	12.2	4.5
LAR-BP185/LAR-L	P233D/P236D/P271G/P296D/AS30K	7.7E-09	3.3E-06	2.8E-05	1.5E-07	6.6E-05	21.8	176.1	19.4	4.5
LAR-BP186/LAR-L	P233D/P236D/P271G/AS30K	8.2E-09	3.1E-06	2.4E-05	2.0E-07	6.9E-05	16.1	128.1	15.8	4.1
LAR-BP187/LAR-L	P233D/G237D/P236D/H266D/P271G/AS30K	2.2E-08	1.9E-06	2.9E-05	6.4E-07	6.9E-05	23.0	34.5	3.7	25.1
LAR-BP188/LAR-L	G237D/P236D/H266D/P271G/K326A/AS30K	1.9E-08	6.9E-07	9.3E-06	3.2E-08	5.3E-05	26.3	255.2	95.4	1.1
LAR-BP189/LAR-L	P233D/P236D/H266D/P271G/K326A/AS30K	3.9E-08	1.2E-06	1.0E-05	5.1E-08	4.1E-05	22.7	196.3	60.4	1.5
LAR-BP190/LAR-L	G237D/P236D/H266D/P271G/AS30K	6.5E-08	1.0E-06	9.5E-06	3.9E-08	4.5E-05	25.4	241.1	78.4	1.3
LAR-BP191/LAR-L	P233D/P236D/H266D/P271G/K326A/AS30K	4.2E-09	4.1E-06	2.7E-05	2.2E-07	7.3E-05	18.5	120.5	13.6	5.4
LAR-BP192/LAR-L	P233D/P236D/H266D/P271G/P296D/AS30K	5.2E-09	3.5E-06	2.2E-05	1.7E-07	6.2E-05	21.1	133.3	18.7	4.5
LAR-BP193/LAR-L	P233D/P236D/H266D/P271G/AS30K	4.1E-09	3.1E-06	2.4E-05	1.6E-07	6.3E-05	17.7	136.4	17.6	4.0
LAR-BP194/LAR-L	P233D/L234Y/G237D/P236D/P271G/P296D/K326D/AS30K	5.9E-08	1.7E-06	2.2E-05	1.2E-07	3.8E-05	14.5	78.0	26.2	2.2

[Example 13] Preparation of variants with enhanced FcγRIIb binding

As shown in Example 8, when enhancing the FcγRIIb binding, it is preferable that the FcγRIIb binding is enhanced while maximally suppressing the binding to other activating FcγRs. Thus, the present inventors additionally produced variants with enhanced FcγRIIb binding or improved selectivity to FcγRIIb by combining alterations that enhance the FcγRIIb binding or improving the selectivity to FcγRIIb. Specifically, the alterations described in Examples 9, 11, and 12 which were found to be effective when combined with alteration P238D, were combined with one another, on the basis of the P238D alteration which showed the excellent effect to enhance the FcγRIIb binding and to improve the selectivity to FcγRIIb.

10 Variants were produced by combining the Fc regions of IL6R-G1d (SEQ ID NO: 133) and IL6R-B3 (SEQ ID NO: 144) with alterations E233D, L234Y, G237D, S267Q, H268D, P271G, Y296D, K326D, K326A, A330R, and A330K described in Examples 9, 11, and 12 which were found to be effective when combined with alteration P238D. Using IL6R-L (SEQ ID NO: 135) as the antibody L chain, antibodies comprising the above-described variants in the heavy
15 chain were expressed and purified according to the method described in Reference Examples 1 and 2. The resulting variants were respectively assessed for the binding to each FcγR (FcγRIa, FcγRIIaH, FcγRIIaR, FcγRIIb, or FcγRIIIaV) by the method described in Reference Example 25.

The KD of each variant to each FcγR is shown in Table 21. "Alteration" refers to an alteration introduced into IL6R-B3 (SEQ ID NO: 144). IL6R-B3/IL6R-L which is used as the
20 template to produce each variant is indicated by asterisk (*). "KD (IIaR)/KD (IIb)" in the table shows the value obtained by dividing the KD of each variant for FcγRIIaR by the KD of each variant for FcγRIIb. The greater the value, the higher the selectivity to FcγRIIb. "KD (IIb) of parent polypeptide/KD (IIb) of altered polypeptide" shows the value obtained by dividing the KD value of IL6R-B3/IL6R-L for FcγRIIb by the KD value of each variant for FcγRIIb.
25 Meanwhile, KD (IIaR) of parent polypeptide/KD (IIaR) of altered polypeptide shows the value obtained by dividing the KD value of IL6R-B3/IL6R-L for FcγRIIaR by the KD value of each variant for FcγRIIaR. In Table 21, the numeral in the gray-filled cells indicates that the binding of FcγR to IgG was concluded to be too weak to analyze correctly by kinetic analysis and thus was calculated using:

30 [Equation 2]

$$KD = C \bullet R_{\max} / (R_{eq} - RI) - C$$

described in Reference Example 25.

[Table 21]

VARIANT NAME	ALTERATION	KD AGAINST FcγRIIIa (mol/L)	KD AGAINST FcγRIIIb (mol/L)	KD AGAINST FcγRIIIc (mol/L)	KD AGAINST FcγRIIIa (mol/L)	KD AGAINST FcγRIIIb (mol/L)	KD AGAINST FcγRIIIc (mol/L)	KD (I _h R) ₁ /KD (I _h)	KD (I _h R) ₁ OF PARENT POLYPEPTIDE/ KD (I _h R) ₁ OF ALTERED POLYPEPTIDE	KD (I _h R) ₂ OF PARENT POLYPEPTIDE/ KD (I _h R) ₂ OF ALTERED POLYPEPTIDE
IL6R-BP14/IL6R-L		3.40E-10	1.00E-06	6.70E-07	2.40E-06	3.50E-06	3.50E-07	0.4	1.2	1.1
IL6R-BP15/IL6R-L		4.20E-10	1.00E-06	7.70E-07	3.10E-06	3.30E-07	3.30E-07	0.3	1	1
IL6R-BP16/IL6R-L		1.10E-08	1.50E-05	4.90E-05	1.20E-06	7.10E-05	7.10E-05	12.5	2.6	0.1
IL6R-BP17/IL6R-L		4.30E-08	1.30E-05	1.30E-05	1.10E-06	6.70E-05	6.70E-05	3.7	75.6	0.8
IL6R-BP18/IL6R-L		6.20E-07	2.90E-05	4.20E-05	1.40E-07	5.10E-05	5.10E-05	20.7	22.1	0.3
IL6R-BP19/IL6R-L		2.80E-06	3.90E-06	2.50E-05	1.50E-07	5.00E-05	5.00E-05	24	20.7	0.3
IL6R-BP20/IL6R-L		3.70E-08	1.50E-06	2.20E-06	7.60E-08	3.80E-05	3.80E-05	19.7	40.6	0.7
IL6R-BP21/IL6R-L		1.60E-08	6.10E-07	2.30E-06	3.40E-08	2.90E-05	2.90E-05	17.9	91.2	1.2
IL6R-BP22/IL6R-L		2.03E-08	1.10E-06	1.20E-06	3.60E-08	5.80E-05	5.80E-05	30.6	85.1	1
IL6R-BP23/IL6R-L		1.30E-07	7.10E-07	3.30E-06	2.80E-08	4.50E-05	4.50E-05	25.4	110.7	1.3
IL6R-BP24/IL6R-L		5.10E-08	7.10E-07	2.50E-06	3.40E-08	4.70E-05	4.70E-05	20.9	91.2	1.3
IL6R-BP25/IL6R-L		2.70E-08	6.00E-07	2.80E-06	2.50E-08	2.50E-05	2.50E-05	24	124	1.8
IL6R-BP26/IL6R-L		6.20E-09	4.50E-07	9.30E-06	4.50E-07	3.50E-05	3.50E-05	12.9	88.6	2.4
IL6R-BP27/IL6R-L		9.50E-08	6.90E-07	2.80E-06	3.50E-08	3.20E-05	3.20E-05	19.7	88.6	1.6
IL6R-BP28/IL6R-L		5.20E-08	5.70E-07	2.40E-06	3.20E-08	3.60E-05	3.60E-05	17.3	53.9	1.9
IL6R-BP29/IL6R-L		2.70E-08	6.20E-07	2.90E-06	3.20E-08	2.60E-05	2.60E-05	19.4	96.9	1.8
IL6R-BP30/IL6R-L		5.30E-08	4.20E-07	1.10E-05	4.00E-08	5.20E-05	5.20E-05	10.5	77.3	2.6
IL6R-BP31/IL6R-L		5.00E-08	8.10E-07	3.30E-06	4.20E-08	3.70E-05	3.70E-05	19.3	73.8	1.4
IL6R-BP32/IL6R-L		1.40E-08	5.70E-07	3.80E-06	2.10E-08	6.70E-05	6.70E-05	27.1	147.6	1.9
IL6R-BP33/IL6R-L		9.40E-09	7.40E-07	1.10E-05	2.30E-08	4.00E-05	4.00E-05	32.2	134.8	1.5
IL6R-BP34/IL6R-L		7.60E-08	8.40E-07	3.30E-06	5.60E-08	4.50E-05	4.50E-05	15	55.4	1.3
IL6R-BP35/IL6R-L		1.00E-08	6.90E-07	2.80E-06	3.70E-08	5.10E-05	5.10E-05	18.6	83.8	1.6
IL6R-BP36/IL6R-L		6.50E-09	1.20E-06	2.00E-05	1.20E-07	3.10E-05	3.10E-05	10	25.6	0.9
IL6R-BP37/IL6R-L		3.90E-09	6.90E-07	7.30E-06	4.40E-08	2.50E-05	2.50E-05	15.5	70.5	1.6
IL6R-BP38/IL6R-L		7.70E-09	8.40E-07	1.90E-05	6.50E-08	3.50E-05	3.50E-05	12.9	47.7	1.3
IL6R-BP39/IL6R-L		4.10E-09	1.10E-06	1.90E-05	1.00E-07	3.50E-05	3.50E-05	11	31	1
IL6R-BP40/IL6R-L		2.40E-09	6.40E-07	7.00E-06	3.00E-08	2.70E-05	2.70E-05	17.8	86.1	1.7
IL6R-BP41/IL6R-L		7.60E-09	8.10E-07	1.70E-05	6.00E-08	4.80E-05	4.80E-05	13.5	51.7	1.4
IL6R-BP42/IL6R-L		1.10E-09	1.50E-06	2.40E-05	9.50E-08	5.20E-05	5.20E-05	15.1	68.9	1.6
IL6R-BP43/IL6R-L		1.40E-09	6.80E-07	8.50E-06	4.50E-08	3.10E-05	3.10E-05	14.7	60.8	1.5
IL6R-BP44/IL6R-L		3.00E-09	5.40E-07	5.00E-05	3.60E-08	2.50E-05	2.50E-05	15	86.1	1.5
IL6R-BP45/IL6R-L		7.80E-09	1.80E-06	2.60E-05	1.10E-07	4.60E-05	4.60E-05	16.4	28.2	0.6
IL6R-BP46/IL6R-L		6.30E-09	1.40E-06	2.50E-05	8.30E-08	3.80E-05	3.80E-05	16.9	37.3	0.8
IL6R-BP47/IL6R-L		8.00E-09	1.60E-06	2.30E-05	9.20E-08	4.40E-05	4.40E-05	17.1	33.7	0.7
IL6R-BP48/IL6R-L		7.30E-09	8.10E-07	1.20E-05	3.70E-08	4.40E-05	4.40E-05	21.9	83.8	1.1
IL6R-BP49/IL6R-L		1.70E-09	8.30E-07	1.00E-05	3.50E-08	4.40E-05	4.40E-05	23.4	88.6	1.3
IL6R-BP50/IL6R-L		7.00E-09	6.20E-07	7.20E-06	3.70E-08	2.80E-05	2.80E-05	16.8	85.8	1.8

When taking the binding to each FcγR by IL6R-B3/IL6R-L resulting from introducing the K439E alteration into the H chain of IL6R-G1d/IL6R-L containing the sequence of native human IgG1 as 1, the binding of IL6R-G1d/IL6R-L to FcγRIa as 1.3 times; the binding of IL6R-G1d/IL6R-L to FcγRIIaR was 1.1 times; the binding of IL6R-G1d/IL6R-L to FcγRIIaH was 1.1 times, the binding of IL6R-G1d/IL6R-L to FcγRIIb binding was 1.2 times, and the binding of IL6R-G1d/IL6R-L to FcγRIIIaV was 0.9 times. Thus, for any given FcγR type, the binding of IL6R-B3/IL6R-L to FcγR was comparable to the binding of IL6R-G1d/IL6R-L to FcγR. Thus, the comparison of the binding of each variant to each FcγR with that of IL6R-B3/IL6R-L prior to introduction of the alteration is assumed to be equivalent to the comparison of the binding of each variant to each FcγR with the binding to each FcγR by IL6R-G1d/IL6R-L containing the sequence of native human IgG1. For this reason, in the subsequent Examples below, the binding activity of each variant to each FcγR will be compared to that to each FcγR by IL6R-B3/IL6R-L prior to introduction of the alteration. Table 21 shows that all the variants have increased FcγRIIb binding activity as compared to IL6R-B3 prior to introduction of the alteration. The binding activity of IL6R-BF648/IL6R-L, which was the lowest, was increased by 2.6 times, while the binding activity of IL6R-BP230/IL6R-L, which is the highest, was increased by 147.6 times. Regarding the value of KD (IIaR)/KD (IIb) that represents the selectivity, the value for IL6R-BP234/IL6R-L, which was the lowest, was 10.0, while the value for IL6R-BP231/IL6R-L, which was the highest, was 32.2. Compared to 0.3 for IL6R-B3/IL6R-L prior to introduction of the alteration, these values imply that all the variants have improved selectivity. All the variants showed lower binding activity to FcγRIa, FcγRIIaH, and FcγRIIIaV than that of IL6R-B3/IL6R-L prior to introduction of the alteration.

[Example 14] X-ray crystal structure analysis of the complexes of FcγRIIb extracellular region or FcγRIIaR extracellular region and Fc region with enhanced FcγRIIb binding

As shown in Example 13, the FcγRIIb binding of variant IL6R-BP230/IL6R-L, whose FcγRIIb binding was enhanced most, was enhanced to about 150 times as compared to IL6R-B3/IL6R-L prior to introduction of the alteration, while the enhancement of its FcγRIIaR binding was suppressed to an extent of about 1.9 times. Thus, IL6R-BP230/IL6R-L is a variant excellent in both FcγRIIb binding and selectivity. However, the present inventors sought a possibility to create more preferable variants with further enhanced FcγRIIb binding while suppressing the FcγRIIaR binding as possible.

As shown in Fig. 30 described in Example 10, in the Fc region with alteration P238D, Asp at position 270 (EU numbering) in its CH2 domain B forms a tight electrostatic interaction with Arg at position 131 in FcγRIIb. This amino acid residue at position 131 is His in FcγRIIIa and FcγRIIaH, while it is Arg in FcγRIIaR like in FcγRIIb. Thus, there is no difference between

FcγRIIaR and FcγRIIb in terms of the interaction of the amino acid residue at position 131 with Asp at position 270 (EU numbering) in the CH2 domain B. This is assumed to be a major factor for the poor selectivity between the FcγRIIb binding and FcγRIIaR binding of the Fc region.

5 On the other hand, the extracellular regions of FcγRIIa and FcγRIIb are 93% identical in amino acid sequence, and thus they share very high homology. Based on the crystal structure analysis of the complex of the Fc region of native IgG1 (hereinafter abbreviated as Fc (WT)) and the extracellular region of FcγRIIaR (J. Immunol. (2011) 187, 3208-3217), a difference found
10 (Gln127, Leu132, Phe160) between FcγRIIaR and FcγRIIb. Thus, the present inventors predicted that it was extremely difficult to improve the selectivity of the Fc region between the FcγRIIb binding and FcγRIIaR binding.

In this context, the present inventors conceived that, in order to further enhance the FcγRIIb-binding activity of the Fc region, and to improve the selectivity of the Fc regions for the
15 binding to FcγRIIb and FcγRIIaR binding, it was important to clarify subtle differences between Fc region-FcγRIIb interaction and Fc region-FcγRIIaR interaction by analyzing not only the three-dimensional structure of the complex of the Fc region with enhanced FcγRIIb binding and the extracellular region of FcγRIIb but also the three-dimensional structure of the complex of the Fc region with enhanced FcγRIIb binding and the extracellular region of FcγRIIaR. First, the
20 present inventors analyzed the X-ray crystal structure of the complex of the extracellular region of FcγRIIb or FcγRIIaR and Fc (P208) resulting from eliminating the K439E alteration from the Fc region of IL6R-BP208/IL6R-L created as described in Example 12, which was the variant used as the basis in producing IL6R-BP230/IL6R-L.

25 (14-1) X-ray crystal structure analysis of the complex of Fc (P208) and the extracellular region of FcγRIIb

[Expression and purification of Fc (P208)]

Fc (P208) was prepared as described below. First, IL6R-P208 was produced by substituting Lys for Glu at position 439 (EU numbering) in IL6R-BP208, as is in the case of the
30 sequence of native human IgG1. Then, the gene sequence of Fc (P208), which was cloned by PCR from Glu at position 216 (EU numbering) to the C terminus using as a template a DNA encoding a variant with a substitution of Ser for Cys at position 220 (EU numbering), was cloned. Expression vector construction, expression, and purification were achieved according to the method described in Reference Examples 1 and 2. Meanwhile, Cys at position 220 (EU
35 numbering) in ordinary IgG1 forms a disulfide bond to a Cys in the L chain. When preparing the Fc region alone, the L chain is not expressed in combination. Thus, Cys at position 220 was

substituted by Ser to avoid unnecessary disulfide bond formation.

[Expression and purification of the extracellular region of FcγRIIb]

The extracellular region of FcγRIIb was prepared according to the method described in
5 Reference Example 25.

[Purification of the Fc (P208)/FcγRIIb extracellular region complex]

0.15 mg of the purified product of Endo F1 (Protein Science (1996) 5, 2617-2622)
expressed in *E. coli* as a fusion protein with glutathione S-transferase was added 1.5 mg of a
10 crystallization sample of the extracellular region of FcγRIIb. This added sample in 0.1 M
Bis-Tris buffer (pH 6.5) was allowed to stand at room temperature for three days to cleave off
N-type sugar chains except N-acetylglucosamine directly linked to the Asn in the sample of the
extracellular region of FcγRIIb. Then, the sample of the extracellular region of FcγRIIb
subjected to the sugar chain cleavage treatment was concentrated with a 5000MWCO
15 ultrafiltration filter, and purified by chromatography with a gel filtration column (Superdex200
10/300) equilibrated with 20 mM HEPES (pH 7.5)/0.1 M NaCl. Next, Fc (P208) was added in
such a way that the extracellular region of FcγRIIb is present in a slightly excessive molar ratio.
After concentrating with a 10000MWCO ultrafiltration filter, the purified fraction of the
extracellular region of FcγRIIb subjected to the sugar chain cleavage was purified by
20 chromatography with a gel filtration column (Superdex200 10/300) equilibrated with 25 mM
HEPES (pH 7.5)/0.1 M NaCl. The purified fraction prepared as described above was used as a
sample of Fc (P208)/FcγRIIb extracellular region complex in the subsequent assessment.

[Crystallization of the complex of Fc (P208)/FcγRIIb extracellular region]

25 A sample of Fc (P208)/FcγRIIb extracellular region complex concentrated to about 10
mg/ml with a 10000MWCO ultrafiltration filter was crystallized using the hanging drop vapor
diffusion method in combination with the seeding method. VDXTM plate (Hampton Research)
was used for crystallization. Using a reservoir solution of 0.1 M Bis-Tris (pH 6.5)/19%(w/v)
PEG3350/0.2 M potassium phosphate dibasic, crystallization drops were prepared at a mixing
30 ratio of reservoir solution : crystallization sample = 0.85 μl : 0.85 μl. Crystals of the complex
obtained under the same condition were crushed with Seed BeadTM (Hampton Research) to prepare
a seed crystal solution. The crystallization drops were added with 0.15 μl of a diluted solution
prepared from the seed solution and allowed to stand at 20°C in sealed reservoir wells. This
yielded plate-like crystals.

35

[X-ray diffraction data measurements from an Fc (P208)/FcγRIIb extracellular region complex]

crystal]

A single crystal of Fc (P208)/FcγRIIb extracellular region complex prepared as described above was soaked into a solution of 0.1 M Bis-Tris (pH 6.5)/24% (w/v) PEG3350/0.2 M potassium phosphate dibasic/20% (v/v) ethylene glycol. Then, the single crystal was fished
 5 out of the solution using a pin with attached tiny nylon loop, and frozen in liquid nitrogen. X-ray diffraction data of the single crystal was collected with Spring-8 BL32XU. During the measurement, the crystal was constantly placed in a nitrogen stream at -178°C to maintain in a frozen state. A total of 300 X-ray diffraction images of the single crystal were collected using CCD detector MX-225HE (RAYONIX) attached to a beam line with rotating the single crystal
 10 0.6° at a time. Based on the obtained diffraction images, lattice constant determination, diffraction spot indexing, and diffraction data processing were performed using programs Xia2 (J. Appl. Cryst. (2010) 43, 186-190), XDS Package (Acta Cryst. (2010) D66, 125-132) and Scala (Acta Cryst. (2006) D62, 72-82). Finally, the diffraction intensity data of the single crystal up to 2.81 Å resolution was obtained. The crystal belongs to the space group C222₁ with lattice
 15 constant $a = 156.69 \text{ Å}$, $b = 260.17 \text{ Å}$, $c = 56.85 \text{ Å}$, $\alpha = 90^\circ$, $\beta = 90^\circ$, and $\gamma = 90^\circ$.

[X-ray crystal structure analysis of Fc (P208)/FcγRIIb extracellular region complex]

The structure of Fc (P208)/FcγRIIb extracellular region complex was determined by a molecular replacement method using program Phaser (J. Appl. Cryst. (2007) 40, 658-674). The
 20 number of complexes in an asymmetrical unit was estimated to be one from the size of the obtained crystal lattice and the molecular weight of Fc (P208)/FcγRIIb extracellular region complex. The segments spanning the amino acid residues at positions 239-340 of the A chain and at positions 239-340 of the B chain, which were retrieved as an independent coordinate from the structural coordinate of PDB code: 3SGJ for the crystal structure of Fc (WT)/FcγRIIIa
 25 extracellular region complex, were used as a model for searching the CH2 domain of the Fc region. Likewise, the segments spanning the amino acid residues at positions 341-444 of the A chain and at positions 341-443 of the B chain, which were retrieved as a coordinate from the structural coordinate of PDB code: 3SGJ, were used as a model for searching the CH3 domain of the Fc region. Finally, the segment spanning the amino acid residues at positions 6-178 of the A
 30 chain, which was retrieved from the structural coordinate of PDB code: 2FCB for the crystal structure of the extracellular region of FcγRIIb, was used as a model for searching Fc (P208). The present inventors tried to determine the orientations and positions of the respective search models of the CH3 domain of the Fc region, the extracellular region of FcγRIIb, and the CH2 domain of the Fc region in the crystal lattices based on the rotation function and translation
 35 function, but failed to determine the position of one of the CH2 domains. Then, with reference to the crystal structure of the complex of Fc (WT)/FcγRIIIa extracellular region, the position of

the last CH2 domain A was determined from an electron density map that was calculated based on the phase determined for the remaining three parts. Thus, the present inventors obtained an initial model for the crystal structure of the complex of Fc (P208)/FcγRIIb extracellular region. The crystallographic reliability factor R value of the structural model for the data of diffracted
 5 intensity at 25 to 3.0 Å was 42.6% and Free R value was 43.7% after rigid body refinement where the two CH2 domains and two CH3 domains of the Fc region, and the extracellular region of FcγRIIb were allowed to deviate from the obtained initial structural model. Then, structural model refinement was achieved by repeating structural refinement using program REFMAC5 (Acta Cryst. (2011) D67, 355-367) followed by revision of the structural model performed using
 10 program Coot (Acta Cryst. (2010) D66, 486-501) with reference to the electron density maps where the coefficients 2Fo-Fc and Fo-Fc were calculated using experimentally determined structural factor Fo, structural factor Fc region calculated according to the structural model, and the phases calculated according to the structural model. Then, further refinement was carried out based on the electron density maps with coefficients of 2Fo-Fc and Fo-Fc by integrating
 15 water molecules into the structural model. With 27259 diffracted intensity data at 25 to 2.81 Å resolution, ultimately the crystallographic reliability factor R value was 24.5% and free R value was 28.2% for the structural model comprising 4786 non-hydrogen atoms.

The three-dimensional structure of the complex of Fc (P208)/FcγRIIb extracellular region was determined at a resolution of 2.81 Å by structure analysis. The structure obtained
 20 by the analysis is shown in Fig. 35. FcγRIIb extracellular region was revealed to be bound and sandwiched between the two CH2 domains of the Fc region, which resembles the three-dimensional structures of the previously analyzed complexes of Fc (WT), which is the Fc of native IgG, and each of the extracellular regions of FcγRIIIa (Proc. Natl. Acad. Sci. USA (2011) 108, 12669-126674), FcγRIIIb (Nature (2000) 400, 267-273; J. Biol. Chem. (2011) 276,
 25 16469-16477), and FcγRIIa (J. Immunol. (2011) 187 (6), 3208-3217).

A close observation of the complex of Fc (P208)/FcγRIIb extracellular region revealed a change in the loop structure at positions 233 to 239 (EU numbering) following the hinge region in the CH2 domain A of the Fc region due to an influence of the introduced the G237D and P238D alterations as compared to the complex of Fc (WT)/FcγRIIaR extracellular region (Fig.
 30 36). This leads to that the main chain of Asp at position 237 (EU numbering) in Fc (P208) formed a tight hydrogen bond to the side chain of Tyr at position 160 in FcγRIIb (Fig. 37). In both FcγRIIaH and FcγRIIaR, the amino acid residue at position 160 is Phe, which is incapable of forming such a hydrogen bond. This suggests that the above described hydrogen bond has important contribution to the enhancement of the FcγRIIb binding and the acquisition of the
 35 FcγRIIa binding selectivity of Fc (P208), *i.e.*, improvement of the FcγRIIb-binding activity and reduction of FcγRIIa-binding activity of Fc (P208).

On the other hand, the side chain of Asp at position 237 (EU numbering) in Fc (P208) forms neither particularly significant interaction in the Fc γ RIIb binding nor interaction with other residues within the Fc region. Ile at position 332, Glu at position 333, and Lys at position 334 (EU numbering) in the Fc region are located close to Asp at position 237 (EU numbering) (Fig. 38). When the amino acid residues of these positions are substituted by hydrophilic residues to form an interaction with the side chain of Asp at position 237 (EU numbering) in Fc (P208) and the loop structure can be stabilized by the interaction, this can lead to reduction of the entropic energy loss due to the hydrogen bonding between the Fc region and Tyr at position 160 in Fc γ RIIb and thereby to an increase in the binding free energy, *i.e.*, an increase in the binding activity.

When the X-ray crystal structure of the complex of Fc (P238D) with the P238D alteration and Fc γ RIIb extracellular region described in Example 10 is compared to the X-ray crystal structure of the complex of Fc (P208) and Fc γ RIIb extracellular region, deviations are observed at five portions in Fc (P208) as compared to Fc (P238D) and most of the changes are seen only at the side chain level. Meanwhile, a positional deviation at the main chain level due to the Pro-to-Gly alteration at position 271 (EU numbering) is also observed in the CH2 domain B of the Fc region, and in addition there is a structural change in the loop at positions 266 to 270 (EU numbering) (Fig. 39). As described in Example 11, it is suggested that, when Asp at position 270 (EU numbering) in Fc (P238D) forms a tight electrostatic interaction with Arg at position 131 in Fc γ RIIb, the interaction can induce stereochemical stress at Pro at position 271 (EU numbering). The experiment described herein suggests that the structural change observed with the alteration to Gly for the amino acid at position 271 (EU numbering) is assumed to be a result of elimination of the structural distortion accumulated at Pro prior to the alteration and the elimination results in an increase in the free energy for the Fc γ RIIb binding, *i.e.*, an increase in the binding activity.

Furthermore, it was demonstrated that, due to the change of the loop structure at positions 266 to 271 (EU numbering), Arg at position 292 (EU numbering) underwent a structural change while being in two states. In this case, the electrostatic interaction (Fig. 39) formed between Arg at position 292 (EU numbering) and Asp at position 268 (EU numbering) which is an altered residue in Fc (P208) can contribute to the stabilization of the loop structure. Since the electrostatic interaction formed between Asp at position 270 (EU numbering) in the loop and Arg at position 131 in Fc γ RIIb largely contribute to the binding activity of Fc (P208) to Fc γ RIIb, the stabilization of the loop structure in the binding conformation was likely to reduce the entropic energy loss upon binding. Thus, the alteration is expected to result in an increase in the binding free energy, *i.e.*, an increase in the binding activity.

Moreover, the possibility of alteration to increase the activity was scrutinized based on

the result of structural analysis. Ser at position 239 (EU numbering) was found as a candidate for the site to introduce alteration. As shown in Fig. 40, Ser at position 239 (EU numbering) in the CH2 domain B is present at the position toward which Lys at position 117 in FcγRIIb extends most naturally in structure. However, since the electron density was not observed for Lys at position 117 in FcγRIIb by the analysis described above, the Lys has no definite structure. In this situation, Lys117 is likely to have only a limited effect on the interaction with Fc (P208). When Ser at position 239 (EU numbering) in the CH2 domain B is substituted with negatively charged Asp or Glu, such an alteration is expected to cause an electrostatic interaction with the positively charged Lys at position 117 in FcγRIIb, thereby resulting in improved FcγRIIb-binding activity.

On the other hand, an observation of the structure of Ser at position 239 (EU numbering) in the CH2 domain A revealed that, by forming a hydrogen bond to the main chain of Gly at position 236 (EU numbering), the side chain of this Ser stabilized the loop structure at positions 233 to 239, including Asp at position 237 (EU numbering) that forms a hydrogen bond to the side chain of Tyr at position 160 in FcγRIIb, following the hinge region (Fig. 37). The stabilization of the loop structure in the binding conformation can reduce the entropic energy loss upon binding, and result in an increase in the binding free energy, *i.e.*, an improvement of the binding activity. Meanwhile, when Ser at position 239 (EU numbering) in the CH2 domain A is substituted with Asp or Glu, the loop structure can become unstable due to loss of the hydrogen bond to the main chain of Gly at position 236 (EU numbering). In addition, the alteration can result in electrostatic repulsion to Asp at position 265 (EU numbering) in close proximity, leading to further destabilization of the loop structure. The energy for the destabilization corresponds to loss of free energy for the FcγRIIb binding, which can result in reduction in the binding activity.

(14-2) X-ray crystal structure analysis of the complex of Fc (P208) and FcγRIIaR extracellular region

[Expression and purification of the extracellular region of FcγRIIaR]

The extracellular region of FcγRIIaR was prepared according to the method described in Reference Example 2.

[Purification of the complex of Fc (P208)/FcγRIIaR type extracellular region]

1.5 mg of purified sample of the extracellular region of FcγRIIaR was added with 0.15 mg of the purified product of Endo F1 (Protein Science (1996) 5, 2617-2622) expressed in *E. coli* as a fusion protein with S-transferase, 20 μl of 5 U/ml Endo F2 (QA-bio), and 20 μl of 5 U/ml Endo F3 (QA-bio). After 9 days of incubation at room temperature in 0.1 M Na acetate

buffer (pH 4.5), the sample was further added with 0.07 mg of the above-described Endo F1, 7.5 μ l of the above-described Endo F2, and 7.5 μ l of the above-described Endo F3, and was incubated for three days to cleave off N-type sugar chains except N-acetylglucosamine directly linked to the Asn in the sample of the extracellular region of Fc γ RIIa R. Then, the sample of
 5 the extracellular region of Fc γ RIIaR concentrated with a 10000MWCO ultrafiltration filter and subjected to the above-described sugar chain cleavage treatment was purified by chromatography with a gel filtration column (Superdex™200 10/300) equilibrated with 25 mM HEPES (pH 7)/0.1M NaCl. Next, Fc (P208) was added in such a way that the extracellular region of Fc γ RIIaR is present in a slightly excessive molar ratio. After concentrating with a
 10 10000MWCO ultrafiltration filter, the purified fraction of the extracellular region of Fc γ RIIaR subjected to the above-described sugar chain cleavage treatment was purified by chromatography with a gel filtration column (Superdex200 10/300) equilibrated with 25 mM HEPES (pH 7)/0.1 M NaCl. The purified fraction prepared as described above was used as a sample of Fc (P208)/Fc γ RIIaR type extracellular region complex in the subsequent assessment.

15

[Crystallization of the complex of Fc (P208)/Fc γ RIIaR type extracellular region]

A sample of Fc (P208)/Fc γ RIIa R type extracellular region complex concentrated to about 10 mg/ml with a 10000MWCO ultrafiltration filter was crystallized using the sitting drop vapor diffusion method. Using a reservoir solution of 0.1 M Bis-Tris (pH 7.5)/26% (w/v)
 20 PEG3350/ 0.2 M ammonium sulfate, crystallization drops were prepared at a mixing ratio of reservoir solution : crystallization sample = 0.8 μ l : 1.0 μ l. The drops were tight sealed and allowed to stand at 20°C. This yielded plate-like crystals.

[X-ray diffraction data measurement from Fc (P208)/Fc γ RIIaR extracellular region complex
 25 crystal]

A single crystal of Fc (P208)/Fc γ RIIaR extracellular region complex prepared as described above was soaked into a solution of 0.1 M Bis-Tris (pH 7.5)/27.5% (w/v) PEG3350/0.2 M ammonium sulfate/20% (v/v) glycerol. Then, the crystal was fished out of the solution using a pin with attached tiny nylon loop, and frozen in liquid nitrogen. X-ray
 30 diffraction data of the single crystal was collected from Photon Factory BL-17A of the synchrotron radiation institution in the High Energy Accelerator Research Organization. The crystal was constantly placed in a nitrogen stream at -178°C to maintain in a frozen state during the measurement. A total of 225 X-ray diffraction images of the single crystal were collected using CCD detector Quantum 315r (ADSC) equipped to the beam line with rotating the single
 35 crystal at 0.6° at a time. Based on the obtained diffraction images, lattice constant determination, diffraction spot indexing, and diffraction data processing were performed using

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 254

NOTE : Pour les tomes additionnels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 254

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

CLAIMS

1. A pharmaceutical composition which comprises pharmaceutically acceptable carriers and an antigen-binding molecule comprising an antigen-binding domain and an Fc region variant of human IgG, wherein the antigen-binding molecule has human-FcRn-binding activity in an acidic pH range condition, wherein the Fc region variant has higher binding activity to an Fc γ receptor at pH 7.4 than the binding activity to the Fc γ receptor of an Fc region of a native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, and wherein the antigen-binding domain is an antibody variable region and has at least one histidine and the antigen binding domain has antigen-binding activity that changes depending on pH, wherein the value of $KD(pH5.8)/KD(pH7.4)$, which is the ratio of KD for the antigen in an acidic pH range condition to the KD in a neutral pH range condition, is 2 or more.

2. The pharmaceutical composition of claim 1, wherein the antigen is a soluble antigen.

3. The pharmaceutical composition of claim 1 or 2, wherein the antigen-binding molecule has neutralizing activity against the antigen.

4. The pharmaceutical composition of any one of claims 1 to 3, wherein the Fc region variant is an Fc region with one or more amino acid substitutions at a position selected from the group consisting of positions 221, 222, 223, 224, 225, 227, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 251, 254, 255, 256, 258, 260, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 288, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 311, 313, 315, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 339, 376, 377, 378, 379, 380, 382, 385, 392, 396, 421, 427, 428, 429, 434, 436, and 440 according to EU numbering in a native Fc region of any one of native human IgG1, native human IgG2, native human IgG3, and native human IgG4.

5. The pharmaceutical composition of claim 4, wherein the Fc region variant is an Fc region which comprises one or more amino acids selected from the group consisting of:

either Lys or Tyr at amino acid position 221;

any one of Phe, Trp, Glu, and Tyr at amino acid position 222;

any one of Phe, Trp, Glu, and Lys at amino acid position 223;

any one of Phe, Trp, Glu, and Tyr at amino acid position 224;

any one of Glu, Lys, and Trp at amino acid position 225;

any one of Glu, Gly, Lys, and Tyr at amino acid position 227;

any one of Glu, Gly, Lys, and Tyr at amino acid position 228;

any one of Ala, Glu, Gly, and Tyr at amino acid position 230;

any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 231;

any one of Glu, Gly, Lys, and Tyr at amino acid position 232;

any one of Ala, Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 233;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 234;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 235;

any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 236;

any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 237;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 238;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 239;

any one of Ala, Ile, Met, and Thr at amino acid position 240;
any one of Asp, Glu, Leu, Arg, Trp, and Tyr at amino acid position 241;

any one of Leu, Glu, Leu, Gln, Arg, Trp, and Tyr at amino acid position 243;

His at amino acid position 244;

Ala at amino acid position 245;

any one of Asp, Glu, His, and Tyr at amino acid position 246;

any one of Ala, Phe, Gly, His, Ile, Leu, Met, Thr, Val, and Tyr at amino acid position 247;

any one of Glu, His, Gln, and Tyr at amino acid position 249;

either Glu or Gln at amino acid position 250;

Phe at amino acid position 251;

any one of Phe, Met, and Tyr at amino acid position 254;

any one of Glu, Leu, and Tyr at amino acid position 255;

any one of Ala, Met, and Pro at amino acid position 256;

any one of Asp, Glu, His, Ser, and Tyr at amino acid position 258;

any one of Asp, Glu, His, and Tyr at amino acid position 260;

any one of Ala, Glu, Phe, Ile, and Thr at amino acid position 262;

any one of Ala, Ile, Met, and Thr at amino acid position 263;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 264;

any one of Ala, Leu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 265;

any one of Ala, Ile, Met, and Thr at amino acid position 266;

any one of Asp, Glu, Phe, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 267;

any one of Asp, Glu, Phe, Gly, Ile, Lys, Leu, Met, Pro, Gln, Arg, Thr, Val, and Trp at amino acid position 268;

any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 269;

any one of Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Gln, Arg, Ser, Thr, Trp, and Tyr at amino acid position 270;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 271;

any one of Asp, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 272;

either Phe or Ile at amino acid position 273;

any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 274;

either Leu or Trp at amino acid position 275;

any one of Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 276;

any one of Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 278;

Ala at amino acid position 279;

any one of Ala, Gly, His, Lys, Leu, Pro, Gln, Trp, and Tyr at amino acid position 280;

any one of Asp, Lys, Pro, and Tyr at amino acid position 281;

any one of Glu, Gly, Lys, Pro, and Tyr at amino acid position 282;

any one of Ala, Gly, His, Ile, Lys, Leu, Met, Pro, Arg, and Tyr at amino acid position 283;

any one of Asp, Glu, Leu, Asn, Thr, and Tyr at amino acid position 284;

any one of Asp, Glu, Lys, Gln, Trp, and Tyr at amino acid position 285;

any one of Glu, Gly, Pro, and Tyr at amino acid position 286;

any one of Asn, Asp, Glu, and Tyr at amino acid position 288;

any one of Asp, Gly, His, Leu, Asn, Ser, Thr, Trp, and Tyr at amino acid position 290;

any one of Asp, Glu, Gly, His, Ile, Gln, and Thr at amino acid position 291;

any one of Ala, Asp, Glu, Pro, Thr, and Tyr at amino acid position 292;

any one of Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 293;

any one of Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 294;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 295;

any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, and Val at amino acid position 296;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 297;

any one of Ala, Asp, Glu, Phe, His, Ile, Lys, Met, Asn, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 298;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Val, Trp, and Tyr at amino acid position 299;

any one of Ala, Asp, Glu, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, and Trp at amino acid position 300;

any one of Asp, Glu, His, and Tyr at amino acid position 301;

Ile at amino acid position 302;

any one of Asp, Gly, and Tyr at amino acid position 303;

any one of Asp, His, Leu, Asn, and Thr at amino acid position 304;

any one of Glu, Ile, Thr, and Tyr at amino acid position 305;

any one of Ala, Asp, Asn, Thr, Val, and Tyr at amino acid position 311;

Phe at amino acid position 313;

Leu at amino acid position 315;

either Glu or Gln at amino acid position 317;

any one of His, Leu, Asn, Pro, Gln, Arg, Thr, Val, and Tyr at amino acid position 318;

any one of Asp, Phe, Gly, His, Ile, Leu, Asn, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 320;

any one of Ala, Asp, Phe, Gly, His, Ile, Pro, Ser, Thr, Val, Trp, and Tyr at amino acid position 322;

Ile at amino acid position 323;

any one of Asp, Phe, Gly, His, Ile, Leu, Met, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 324;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 325;

any one of Ala, Asp, Glu, Gly, Ile, Leu, Met, Asn, Pro, Gln, Ser, Thr, Val, Trp, and Tyr at amino acid position 326;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Thr, Val, Trp, and Tyr at amino acid position 327;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Lys, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 328;

any one of Asp, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 329;

any one of Cys, Glu, Phe, Gly, His, Ile, Lys, Leu, Met, Asn, Pro, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 330;

any one of Asp, Phe, His, Ile, Leu, Met, Gln, Arg, Thr, Val, Trp, and Tyr at amino acid position 331;

any one of Ala, Asp, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Val, Trp, and Tyr at amino acid position 332;

any one of Ala, Asp, Glu, Phe, Gly, His, Ile, Leu, Met, Pro, Ser, Thr, Val, and Tyr at amino acid position 333;

any one of Ala, Glu, Phe, Ile, Leu, Pro, and Thr at amino acid position 334;

any one of Asp, Phe, Gly, His, Ile, Leu, Met, Asn, Pro, Arg, Ser, Val, Trp, and Tyr at amino acid position 335;

any one of Glu, Lys, and Tyr at amino acid position 336;

any one of Glu, His, and Asn at amino acid position 337;

any one of Asp, Phe, Gly, Ile, Lys, Met, Asn, Gln, Arg, Ser, and Thr at amino acid position 339;

either Ala or Val at amino acid position 376;

either Gly or Lys at amino acid position 377;

Asp at amino acid position 378;

Asn at amino acid position 379;

any one of Ala, Asn, and Ser at amino acid position 380;
 either Ala or Ile at amino acid position 382;
 Glu at amino acid position 385;
 Thr at amino acid position 392;
 Leu at amino acid position 396;
 Lys at amino acid position 421;
 Asn at amino acid position 427;
 either Phe or Leu at amino acid position 428;
 Met at amino acid position 429;
 Trp at amino acid position 434;
 Ile at amino acid position 436; and
 any one of Gly, His, Ile, Leu, and Tyr at amino acid position 440,
 in the Fc region site according to EU numbering.

6. The pharmaceutical composition of any one of claims 1 to 5, wherein the Fc region of the native human IgG in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain, is an Fc region of any one of native human IgG1, native human IgG2, native human IgG3, and native human IgG4 in which the sugar chain bound at position 297 according to EU numbering is a fucose-containing sugar chain.

7. The pharmaceutical composition of any one of claims 1 to 6, wherein the Fcγ receptor is human FcγRIa, FcγRIIa(R), FcγRIIa(H), FcγRIIb, or FcγRIIIa.

8. The pharmaceutical composition of any one of claims 1 to 6, wherein the Fcγ receptor is human FcγRIIb.

9. The pharmaceutical composition of any one of claims 1 to 8, wherein the Fc region variant is an Fc region which comprises one or more of

Asp at amino acid position 238, and

Glu at amino acid position 328

in the Fc region site according to EU numbering.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37






FIG. 38



THIN LINE: X-RAY CRYSTAL STRUCTURE OF Fc(P238D)/FcγRIIb
EXTRACELLULAR REGION COMPLEX

HEAVY LINE: X-RAY CRYSTAL STRUCTURE OF Fc(P208)/FcγRIIb
EXTRACELLULAR REGION COMPLEX


FIG. 39



THIN LINE : Fc(P208)

HEAVY LINE : FcγRIIb EXTRACELLULAR REGION

FIG. 40



BLACK: X-RAY CRYSTAL STRUCTURE OF Fc(P208)/FcγRIIaR
EXTRACELLULAR REGION COMPLEX
GRAY: X-RAY CRYSTAL STRUCTURE OF Fc(P208)/FcγRIIb
EXTRACELLULAR REGION COMPLEX

FIG. 41




FIG. 42




FIG. 43

G1d	118	ASTKGP	SVFP	LAP	SS	SK	ST	SG	TA	AL	GC	LV	KD	YF	PE	PV	TV	SV	WN	SG	AL	TSG	VH	TF	PA	VL	QS	GL	YS	LS	SV	VT	VP	SS	SL	GT	Q	T	Y	T	C	N	V	N	H	K	P	S																																			
G4d	118	ASTKGP	SVFP	LAP	CS	RS	ST	SE	TA	AL	GC	LV	KD	YF	PE	PV	TV	SV	WN	SG	AL	TSG	VH	TF	PA	VL	QS	GL	YS	LS	SV	VT	VP	SS	SL	GT	Q	T	Y	T	C	N	V	N	H	K	P	S																																			
G1d	208	NTKVDK	KV	EP	KS	C	D	K	T	H	T	C	P	P	C	P	A	P	E	L	L	G	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	R	F	N	W	Y	V	D	G	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	I	N
G4d	208	NTKVDK	KV	EP	KS	C	D	K	T	H	T	C	P	P	C	P	A	P	E	L	L	G	G	P	S	V	F	L	F	P	P	K	P	K	D	T	L	M	I	S	R	T	P	E	V	T	C	V	V	V	D	V	S	H	E	D	P	E	V	R	F	N	W	Y	V	D	G	V	E	V	H	N	A	K	T	K	P	R	E	E	Q	I	N
G1d	298	STYRVV	SVLT	VL	FQ	D	W	I	N	G	K	E	Y	K	C	K	V	S	N	K	A	L	P	A	P	I	E	K	T	I	S	K	A	K	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	D	E	L	T	K	N	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E	S	N	G	Q	P		
G4d	298	STYRVV	SVLT	VL	FQ	D	W	I	N	G	K	E	Y	K	C	K	V	S	N	K	A	L	P	A	P	I	E	K	T	I	S	K	A	K	G	Q	P	R	E	P	Q	V	Y	T	L	P	P	S	R	D	E	L	T	K	N	Q	V	S	L	T	C	L	V	K	G	F	Y	P	S	D	I	A	V	E	W	E	S	N	G	Q	P		
G1d	388	ENNYKT	TPPV	LD	S	D	G	S	F	F	L	Y	S	R	L	T	V	D	K	S	R	W	Q	E	G	N	V	F	S	C	S	V	M	H	E	A	L	H	N	H	Y	T	Q	K	S	L	S	L	S	P	(	S	E	Q	I	D	N	O	:	1	6	1	)																				
G4d	388	ENNYKT	TPPV	LD	S	D	G	S	F	F	L	Y	S	R	L	T	V	D	K	S	R	W	Q	E	G	N	V	F	S	C	S	V	M	H	E	A	L	H	N	H	Y	T	Q	K	S	L	S	L	S	L	(	S	E	Q	I	D	N	O	:	1	6	2	)																				

FIG. 44




FIG. 45




FIG. 46




FIG. 47





FIG. 48

(i) CRYSTAL STRUCTURE IN THE
PRESENCE OF CALCIUM ION



(ii) CRYSTAL STRUCTURE IN THE
ABSENCE OF CALCIUM ION




FIG. 49




FIG. 50




FIG. 51




FIG. 52




FIG. 53A

(gi) LIGHT CHAIN : LfVkl_Ca1




FIG. 53B

(iii) LIGHT CHAIN : LfV_k1_Ca2




FIG. 53C

(iv) LIGHT CHAIN : LfV_k1_Ca3




FIG. 53D




FIG. 54A

(ii) LIGHT CHAIN : LfVkl_Ca6




FIG. 54B




FIG. 55




FIG. 56




FIG. 57




FIG. 58




FIG. 59









FIG. 60

**FIG. 61A**

**FIG. 61B**

F1166**FIG. 61C**

F1167**FIG. 61D**

**FIG. 61E**

F1170




FIG. 61F

71 / 81

F1171




FIG. 61G




FIG. 61H

F1172




FIG. 61I

F1173







FIG. 61J

75 / 81


**FIG. 62A**


**FIG. 62B**


77 / 81

F1384**FIG. 62C**

78 / 81

F1385**FIG. 62D**

**FIG. 62E**

F1388**FIG. 62F**




FIG. 62G