



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/05/14
(87) Date publication PCT/PCT Publication Date: 2020/11/19
(85) Entrée phase nationale/National Entry: 2021/11/09
(86) N° demande PCT/PCT Application No.: US 2020/032988
(87) N° publication PCT/PCT Publication No.: 2020/232305
(30) Priorités/Priorities: 2019/05/14 (US62/847,914);
2019/11/21 (US62/938,786)

(51) Cl.Int./Int.Cl. *C07K 19/00* (2006.01),
A61K 47/65 (2017.01), *C07K 14/00* (2006.01),
C07K 14/475 (2006.01), *C07K 14/52* (2006.01),
C07K 14/705 (2006.01), *C07K 7/06* (2006.01),
C12N 15/62 (2006.01), *C12P 21/00* (2006.01)
(71) Demandeur/Applicant:
WEREWOLF THERAPEUTICS, INC., US
(72) Inventeurs/Inventors:
EVNIN, LUKE, US;
BHASKAR, VINAY, US;
KNUDSEN, GISELLE, US;
HICKLIN, DANIEL, US;
SEIDEL-DUGAN, CYNTHIA, US; ...

(54) Titre : GROUPE CARACTERISTIQUES DE SEPARATION, PROCEDES ET UTILISATION ASSOCIES
(54) Title: SEPARATION MOIETIES AND METHODS AND USE THEREOF




FIG. 1A

(57) **Abrégé/Abstract:**

Provided herein are separation moieties that are suitable for use in conjunction with a variety of therapeutic payloads. The separation moieties serve to generate conditionally active macromolecules whereby the macromolecules have reduced or minimal biological activity until the separation moieties are modified under specific conditions.

(72) **Inventeurs(suite)/Inventors(continued):** SALMERON GARCIA, JOSE ANDRES, US; BRODKIN, HEATHER, US

(74) **Agent:** GOWLING WLG (CANADA) LLP

Date Submitted: 2021/11/09

CA App. No.: 3137512

Abstract:

Provided herein are separation moieties that are suitable for use in conjunction with a variety of therapeutic payloads. The separation moieties serve to generate conditionally active macromolecules whereby the macromolecules have reduced or minimal biological activity until the separation moieties are modified under specific conditions.

SEPARATION MOIETIES AND METHODS OF USE THEREOF

[1] The present application claims the benefit of U.S. Provisional Application No. 62/847,914 filed on May 14, 2019 and U.S. Provisional Application No. 62/938,786 filed on November 21, 2019, each of which are incorporated herein by reference in their entireties.

1. SEQUENCE LISTING

[2] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on May 14, 2020, is named 761146_000140_SL.txt and is 896,815 bytes in size.

2. BACKGROUND

[3] Recombinant fusion proteins containing two or more functional polypeptides have utility in many fields including, for use in protein purification, imaging, as therapeutics, and drug delivery. For example, protein drugs can be fused to Fc domains of antibodies or to carrier proteins (i.e., human serum albumin) for targeting, to extend their plasma half-lives and/or to achieve therapeutic effects. Chen et al., (2013), *Adv Drug Deliv Rev.* 65(10):1357-1369.

[4] Direct fusion of functional polypeptide or domains without a linker may lead to many undesirable outcomes, including misfolding of the fusion proteins (Zhao et al., (2008), *Protein Expr. Purif.*, 61:73-77), low yield in protein production (Amet et al., (2009), *Pharm. Res.* 26:523-528), or impaired bioactivity (Bai et al., (2006) *Proc. Natl Acad. Sci. USA*, 102:7292-7296). One approach to overcome these difficulties is to use linker sequences between the component polypeptides or domains in a fusion protein. However, the selection of a suitable linker to join protein domains together can be complicated and is often neglected in the design of fusion proteins. Chen et al., (2013), *Adv Drug Deliv Rev.* 65(10):1357-1369. The properties of linker sequences, such as length, hydrophobicity, amino acid composition, secondary structures, and the overall folding can affect linker suitability and need to be taken into consideration in designing and selecting an appropriate linker. In addition, linkers that can be cleaved under selected conditions or in selected biological locations (e.g., in tumor microenvironment) to deliver active therapeutic agents (e.g., therapeutic polypeptides) can provide targeted pharmacological activity of the therapeutic agents and reduce unwanted systemic effects. This introduces further complexity into designing suitable linkers.

[5] There is a need for improved linker sequences that can be used to prepare stable fusion proteins, including linkers that can be cleaved under selected conditions. Accordingly, novel separation moieties or linkers are disclosed herein. The separation moieties or linkers disclosed herein can be utilized, for example to specifically deliver prodrugs such as conditionally active

and/or targeted cytokines to target sites where the linkers are processed to activate bioactivity.

3. SUMMARY

[6] Provided herein are compositions and methods to generate and use high efficiency separation moieties and/or linkers. The linkers can confer site-selectivity with regards to biological activity of the attached payload or payloads. In some embodiments, the separation moieties and/or linkers are used in conjugation with therapeutic proteins to treat a disease or disorder, such as proliferative disease, a tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic reaction, a parasitic reaction, graft-versus-host disease and the like.

[7] Disclosed herein are recombinant polypeptides comprising a separation moiety, wherein the separation moiety comprises an amino acid sequence is a substrate for an enzyme, specifically a protease. The protease can be selected from the group consisting of Fibroblast activation protein alpha (FAP α , also known as prolyl endopeptidase FAP), Cathepsin L (CTSL1), an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 or MMP14. The cleavable moiety can also be a substrate for a cathepsin, such as cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin K and/or cathepsin L. Preferably, the cathepsin is Cathepsin L. Preferably, the protease is MMP14 or cathepsin L. The cleavable moiety can comprise an amino acid sequence that is a substrate for at least two proteases. In another embodiment, the separation moiety comprises two or more cleavable moieties, each of which is a substrate for a protease. The separation moiety can comprise a first cleavable moiety comprising a first amino acid sequence that is a substrate for a first protease and a second cleavable moiety comprising a second amino acid sequence that is a substrate for a second protease. In embodiments, the disclosure related to recombinant polypeptides that comprise a separation moiety that contains a protease cleavage motif as disclosed herein. The recombinant polypeptide can comprise a separation moiety that comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220, or an amino acid sequence that has at least 90% identity to SEQ ID NOs:195-220. Preferred separation moieties comprise the sequence GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198). The disclosure also relates to functional variants of separation moieties comprising SEQ IND NOs: 195-220. The functional variants of of SEQ ID NO: 195 can comprise any of SEQ ID NOS: 258-331. The functional variants of SEQ ID NO: 198 can comprise of SEQ ID NO:198 comprises SEQ ID NOS: 199 or any of SEQ ID NOS: 332-408. The separation moieties disclosed herein can comprise Formula I: [D1]-[L1]-[D2]. D1 is a first domain of interest. L1 is a separation moiety that connects or links D1 to D2, wherein the separation moiety comprises an amino acid sequence selected from SEQ ID NOs: 195-220 or an amino acid sequence that has at least about 90%

identity to SEQ ID NOs:195-220. D2 is a second domain of interest.

[8] In one embodiment, the recombinant polypeptide can further comprise a non-cleavable linker sequence. The recombinant polypeptide can include a therapeutic protein, such as a cytokine, chemokine, growth factor, soluble receptor, antigen-binding portion of an antibody (e.g., scFV, dAb) and the like. In another embodiment, the recombinant polypeptide comprises a cytokine, chemokine, growth factor, a soluble receptor, or any combination thereof. In another embodiment, the recombinant polypeptide comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain. In one embodiment, the recombinant polypeptide comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit. In one embodiment, the recombinant polypeptide comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.

[9] In one embodiment, the cleavable moiety is cleaved with either (a) greater catalytic efficiency or (b) greater specificity or (c) both (a) and (b), by one or more proteases than a reference polypeptide sequence. In another embodiment, the one or more proteases are selected from the group consisting of FAP α , CTSL1, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 and MMP14. The one or more proteases can also include a protease selected from cathepsins, such as cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin K and/or cathepsin L. In another embodiment, the reference polypeptide sequence is present in a naturally occurring polypeptide substrate for FAP α , CTSL1, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 and MMP14, a cathepsin, such as cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin K and/or cathepsin L or a combination thereof. In another embodiment, the cleavable moiety is cleaved with reduced catalytic efficiency by one or more proteases than a reference polypeptide sequence. In another embodiment the cleavable moiety is cleaved with reduced catalytic efficiency by one or more serum proteases. In another embodiment, the cleavable moiety is cleaved with reduced catalytic efficiency by one or more hepatic proteases. In another embodiment, the cleavable moiety is cleaved with reduced catalytic efficiency by one or more Factor Xa, hepsin, or thrombin.

[10] In one embodiment, the recombinant polypeptide comprises two or more separation moieties. In one embodiment, the recombinant polypeptide is operably linked to a moiety selected from the group consisting of a polypeptide moiety, a lipid moiety, a nucleic acid moiety, a detectable moiety, and a small molecule.

[11] Provided herein is a recombinant pro-protein comprising: a recombinant polypeptide comprising a cleavable moiety that is a substrate for a protease, wherein the cleavable moiety comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220;

and a polypeptide with biological activity, wherein the pro-protein has attenuated biological activity and wherein cleavage of the cleavable moiety by the protease produces a polypeptide with biological activity that is not attenuated. In one embodiment, the polypeptide with biological activity comprises a cytokine, chemokine, growth factor, a soluble receptor or a combination thereof. In some preferred aspects, the linkers are components of fusion proteins with therapeutic utility, and the linkers are not cleaved or are cleaved with low efficiency in the peripheral circulation but are cleaved with higher efficiency at a desired location in the body, such as a tumor microenvironment or site of inflammation. In another embodiment, the polypeptide with biological activity comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain. In another embodiment, the polypeptide with biological activity comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit. In another embodiment, the polypeptide with biological activity comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.

[12] The disclosure also relates to a fusion protein comprising: a. a signaling protein or molecule; and b. a blocking moiety selected from a steric blocking moiety, a specific blocking moiety, and the combination thereof; and c. a peptide linker that comprises a cleavable moiety, e.g., a cleavable moiety disclosed herein, having at least one protease-cleavable sequence. In one embodiment, the fusion protein, wherein (a) and (b) are operably linked by (c). In another embodiment, the fusion protein, wherein the peptide linker comprises two or more copies of the same cleavable moiety. In another embodiment, the fusion protein wherein the steric blocking moiety comprises human serum albumin (HSA) or an anti-HSA antibody. In another embodiment, the fusion protein wherein the signaling protein is an interleukin-2 amino acid sequence comprising: (i) a non-native N terminus and/or (ii) a non-native C terminus. In another embodiment, the fusion protein wherein the signaling protein is an interleukin-2 amino acid sequence. In another embodiment, the fusion protein further comprising one or more half-life extension domains that are not also a specific blocker. In another embodiment, the fusion protein wherein the non-native N and/or C termini are generated by circular permutation.

[13] The fusion polypeptide provided herein can a first polypeptide fusion partner linked to a ligand by a protease cleavable linker, wherein the cleavable linker has been optimized for catalytic efficiency, and wherein the ligand has been optionally modified, wherein the first polypeptide fusion partner is a blocking moiety which prevents binding of the modified ligand to a target receptor or a subunit of a target receptor until cleavage of the protease cleavable linker.

[14] The fusion polypeptide provided herein can also a fusion polypeptide comprising a first polypeptide fusion partner linked to a ligand by a protease cleavable linker, wherein the cleavable linker has been optimized for catalytic efficiency, and wherein the ligand has been optionally modified, including by circularly permutation to create a non-native N-terminus and a new C-

terminus as compared to a native ligand, and wherein at least one of the new N-terminus or the new C-terminus of the modified ligand is operably linked to a first polypeptide fusion partner to form a fusion polypeptide wherein the first polypeptide fusion partner is a blocking moiety which prevents binding of the modified ligand to a target receptor or a subunit of a target receptor until cleavage of the protease cleavable linker.

[15] In one embodiment, the first polypeptide fusion partner is selected from the group consisting of an antibody, an antibody fragment, and an albumin molecule. In another embodiment, the first polypeptide fusion partner further comprising a second polypeptide fusion partner comprising a second blocking moiety. In another embodiment, the second polypeptide fusion partner is a different kind of blocking moiety than the first polypeptide fusion partner. In another embodiment, the first polypeptide fusion partner is albumin and the second polypeptide fusion partner is a domain comprising a complementary amino acid sequence that blocks activity of the cytokine. In another embodiment, the first polypeptide fusion partner is a steric blocker, such as albumin, and the second polypeptide is a specific blocker, such as a cytokine receptor, portion of a cytokine receptor, a de novo affinity peptide specific for the cytokine, or an antibody or antibody fragment that specifically binds the cytokine of the fusion polypeptide. In another embodiment, the second polypeptide fusion partner is the same kind of blocking moiety as the first polypeptide fusion partner.

[16] In one embodiment, the fusion protein further comprises a tumor antigen binding component. In another embodiment, the fusion protein further comprises a serum half-life extension domain. In another embodiment, the ligand is selected from the group consisting of helix bundle proteins and cytokines (including, but not limited to, growth hormone, IL-2, IL-4, IL-5, IL-6, IL-10, IL-22, IL-23p19, IL-11, IL-13, IL-15, IL-12p35, IL-21, IL-30 (IL27p28), IL-34, IL-35, IL-35p35, IFN- α , IFN- β , IFN γ , LIF, CNTF, oncostatin M, CLCF-1, GCSF, GM-CSF, EPO, ferritin, leptin, placental lactogen, prolactin, apolipoprotein e), b-trefoil proteins (including, but not limited to, IL-1 α , IL-1 β , IL-1Ra, IL18, IL-33, IL-36Ra, IL-36a, IL-36b, IL-36g, IL-37, IL-38, IL1Hy2, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8a, FGF-8b, FGF-8e, FGF-8f, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, FGF-16, FGF-17, FGF-18, FGF-19, FGF-20, FGF-21, FGF-22, FGF-23), α/β (TIM) barrel proteins (including, but not limited to, triosephosphate isomerase), beta sandwich proteins (including, but not limited to, galectin-1, galectin-3, TNF-beta, seven β -propeller proteins, class 1 MHC $\alpha 1\alpha 2$ domain, integrin I domain, GYF domain, C1 domain, C2 domain (for example, from cPLA2, PKC, synaptotagmin), PDZ domains, C3d, C5a. In one embodiment, wherein the ligand comprises IL-2 polypeptide or a fragment or fragments thereof. In another embodiment, the protease-cleavable linker polypeptide comprises a sequence that is capable of being cleaved by at least one protease selected from the group consisting of a kallikrein, thrombin, chymase, carboxypeptidase A, cathepsin G, an elastase,

a FAP, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, PR-3, granzyme M, a calpain, a matrix metalloproteinase (MMP), a plasminogen activator, a cathepsin, a caspase, a tryptase, and a tumor cell surface protease. In another embodiment, the cytokine or fragment or mutein thereof is substantially dissociated from the cytokine blocking moiety after the protease-cleavable polypeptide linker is cleaved by a protease.

[17] Disclosed herein are fusion polypeptide comprising at least one of each of: a cytokine polypeptide or functional fragment or mutein thereof [A]; a cytokine blocking moiety [B]; and an optimized protease-cleavable polypeptide linker [L]; wherein the blocking moiety is selected from the group consisting of an antibody, an antibody fragment, and an albumin, and wherein the cytokine comprises a circularly permuted cytokine. In some embodiments, the fusion protein further comprises a tumor antigen binding component and/or a serum half-life extension domain. In some embodiment, the fusion polypeptide wherein the cytokine peptide or functional fragment or mutein thereof is selected from the group consisting of helix bundle proteins and cytokines (including, but not limited to, growth hormone, IL-2, IL-4, IL-5, IL-6, IL-10, IL-22, IL-23p19, IL-11, IL-13, IL-15, IL-12p35, IL-21, IL-30 (IL27p28), IL-34, IL-35, IL-35p35, IFN- β , IFN γ , LIF, CNTF, oncostatin M, CLCF-1, GCSF, GM-CSF, EPO, ferritin, leptin, placental lactogen, prolactin, apolipoprotein e), a FAP (e.g., Fap α), an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, b-trefoil proteins (including, but not limited to, IL-1 α , IL-1 β , IL-1Ra, IL18, IL-33, IL-36Ra, IL-36a, IL-36b, IL-36g, IL-37, IL-38, IL1Hy2, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8a, FGF-8b, FGF-8e, FGF-8f, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, FGF-16, FGF-17, FGF-18, FGF-19, FGF-20, FGF-21, FGF-22, FGF-23), α/β (TIM) barrel proteins (including, but not limited to, triosephosphate isomerase), beta sandwich proteins (including, but not limited to, galectin-1, galectin-3, TNF- β , seven β -propeller proteins, class 1 MHC $\alpha 1\alpha 2$ domain, integrin I domain, GYF domain, C1 domain, C2 domain (for example, from cPLA2, PKC, synaptotagmin), PDZ domains, C3d, C5a. In one embodiment, the cytokine peptide or functional fragment or mutein thereof comprises IL-2. In another embodiment, the cytokine blocking moiety comprises a ligand binding domain or fragment or mutein of a cognate receptor for the cytokine, a single domain antibody or scFv that binds the cytokine polypeptide or functional fragment or mutein thereof, or an antibody or antibody fragment that binds a receptor of the cytokine. In another embodiment, antibody is a single domain antibody or scFv. In another embodiment, the blocking moiety extends the serum half-life of the cytokine or fragment thereof.

[18] Disclosed herein are fusion polypeptides comprising a protease cleavable moiety, wherein the sequence is catalytically optimized for cleavage by certain proteases and wherein protease cleavage renders the composition inducible in a tumor microenvironment. In one embodiment, the fusion protein further comprises a biologically inactive polypeptide, wherein cleavage of the

cleavable moiety by the protease converts the biologically inactive polypeptide to a biologically active polypeptide. In one embodiment, the biologically inactive polypeptide comprises a cytokine, chemokine, growth factor, or soluble receptor. In another embodiment, the biologically inactive polypeptide comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain. In another embodiment, the biologically inactive polypeptide comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit. In another embodiment, the biologically inactive polypeptide comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.

[19] The disclosure further relates to a nucleic acid encoding any of the polypeptides disclosed herein, a vector comprising any of the nucleic acids encoding any of the polypeptides disclosed herein, and a host cell comprising said vector.

[20] Methods of making a pharmaceutical composition, comprising culturing the host cell comprising a vector comprising a nucleic acid encoding any of the polypeptides disclosed herein, under suitable conditions for expression and collection of desired polypeptides are provided herein. Method of using any of the polypeptides disclosed herein comprising administering an effective amount of a pharmaceutical composition comprising such polypeptides to a subject in need thereof are described. For example, use for treating a subject with a disease or disorder disclosed herein.

[21] Provided herein are pharmaceutical compositions comprising an effective amount of any of the recombinant polypeptides, any of the pro-proteins, any of the fusion proteins, any of the fusion polypeptides, any of the nucleic acids, any of the vectors, or any of the host cells comprising such vectors disclosed herein. For example, a pharmaceutical composition for treating a subject with a disease or disorder disclosed herein.

[22] Also disclosed is the use of the recombinant polypeptides, pro-proteins, fusion proteins, fusion polypeptides, nucleic acids, vectors, or host cells comprising such vectors disclosed herein for the manufacture of a medicament for treating a disease or disorder disclosed herein.

4. BRIEF DESCRIPTION OF THE DRAWINGS

[23] FIG. 1A is a schematic illustrating a protease-activated cytokine or chemokine that includes a blocking moiety. The blocking moiety may optionally function as a serum half-life extending domain. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety via a protease-cleavable linker, thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment a protease cleaves at a protease-cleavage site on the linker, releasing the blocking moiety and allowing the cytokine to bind to its receptor.

[24] FIG. 1B is a schematic illustrating a protease-activated cytokine or chemokine wherein

HSA (blocking moiety) is directly bound to the cytokine or chemokine of interest, with a protease cleavage site between the HSA and a cytokine or chemokine of interest. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety via a protease-cleavable linker, thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment, the protease cleaves at a protease-cleavage site on linker, releasing the blocking moiety and allowing the cytokine to bind to its receptor.

[25] FIG. 1C is a schematic illustrating a protease-activated cytokine or chemokine wherein more than one HSA (blocking moiety) is bound directly to the molecule of interest. If desired, one or more of the HSA can be bonded to the cytokine or chemokine through a linker, such as a linker that contains a protease cleavage site. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety via a protease-cleavable linker, thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment, protease cleaves at protease-cleavage site on linker, releasing the blocking moiety and allowing cytokine to bind receptor. The cytokine now has similar pK properties as compared to the native cytokine (e.g., has a short half-life).

[26] FIG. 1D is a schematic illustrating a protease-activated cytokine or chemokine comprising more than one cytokine, of the same type or different type, each of which is bonded to a binding domain through a protease-cleavable linker. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety via a protease-cleavable linker, thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment a protease cleaves at a protease cleavage site on linker, releasing the blocking moiety and allowing the cytokine to bind to its receptor.

[27] FIG. 2 is a schematic illustrating a protease-activated cytokine or chemokine comprising a cytokine or chemokine polypeptide, a blocking moiety, and a serum half-life extending domain connected by at least one protease-cleavable linker. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety via protease-cleavable linkers, thus blocking its ability to bind to its receptor. It is also bound to a separate half-life extension element, which extends half-life in serum. To the right of the arrow the drawing shows that in an inflammatory or tumor environment a protease cleaves at a protease-cleavage site on linker, thus releasing the serum half-life extension element and the blocking moiety and allowing the cytokine to bind to its receptor. The cytokine now has similar pK properties as compared to the native cytokine (e.g., a short half-life).

[28] FIG. 3 is a schematic illustrating a protease-activated cytokine or chemokine comprising a cytokine or chemokine polypeptide, a blocking moiety, and a targeting domain connected by at least one protease-cleavable linker. To the left of the arrow the drawing shows that a cytokine is connected to a blocking moiety and a targeting domain via a protease-cleavable linker, thus

blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor microenvironment a protease cleaves at the protease cleavage site in the linker, releasing the targeting domain and the blocking moiety and allowing the cytokine to bind to its receptor.

[29] FIG. 4A is a schematic illustrating a protease-activated cytokine or chemokine comprising a cytokine or chemokine polypeptide, a blocking moiety, a targeting domain, and a serum half-life extending domain connected by at least one protease-cleavable linker, wherein the cytokine polypeptide and the targeting domain are connected by a protease-cleavable linker. To the left of the arrow, the drawing shows that a cytokine or chemokine is connected to targeting domain, blocking moiety, and half-life extension element via protease-cleavable linker(s), thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment, the protease cleaves at a protease-cleavage site on linker(s), releasing the half-life extension element, the targeting domain, and the blocking moiety, and allowing the cytokine to bind to its receptor. The cytokine now has similar pK properties as compared to the native cytokine (e.g., short half-life).

[30] FIG. 4B is a schematic illustrating a protease-activated cytokine or chemokine comprising a cytokine or chemokine polypeptide, a blocking moiety, a targeting domain, and a serum half-life extending domain connected by at least one protease-cleavable linker. To the left of the arrow, the drawing shows that a cytokine is connected to targeting domain, a blocking moiety, and a half-life extension element via protease-cleavable linker(s), thus blocking its ability to bind to its receptor. To the right of the arrow the drawing shows that in an inflammatory or tumor environment, the protease cleaves at a protease-cleavage site on linker(s), releasing the half-life extension element and the blocking moiety and allowing the cytokine to bind to the receptor. The targeting moiety remains bound, keeping the cytokine in the tumor microenvironment. The cytokine now has similar pK properties as compared to the native cytokine (e.g., a short half-life).

[31] FIG. 5 depicts a graph showing that Linkers-2 (GPAGLYAQ, SEQ ID NO: 195) and Linkers-3 (ALFKSSFP, SEQ ID NO: 198) are minimally cleaved in lung, kidney, and liver cells.

[32] FIGs. 6A-6B show graphs that polypeptides containing recombinant human IL-2 and the sequence for Linker-1 (GPAGMKGL, SEQ ID NO: 196), Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) are not processed by healthy lung fibroblasts.

[33] FIGs. 7A-7H is a series of graphs showing activity of exemplary IL-2 fusion proteins in IL-2 dependent cytotoxic T lymphocyte cell line CTLL-2. Each graph shows results of the IL-2 proliferation assay as quantified by CellTiter-Glo® (Promega) luminescence-based cell viability assay. Each proliferation assay was performed with HSA (FIGs. 7B, 7D, 7F, 7H) or without (FIGs. 7A, 7C, 7E, 7G). Each fusion protein comprises an anti-HSA binder, and both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[34] FIGs. 8A-8F is a series of graphs showing activity of exemplary IL-2 fusion proteins in IL-2 dependent cytotoxic T lymphocyte cell line CTLL-2. Each graph shows results of the IL-2 proliferation assay as quantified by CellTiter-Glo (Promega) luminescence-based cell viability assay. Both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[35] FIGs. 9A-9Z is a series of graphs showing activity of exemplary IL-2 fusion proteins in IL-2 dependent cytotoxic T lymphocyte cell line CTLL-2. Each graph shows results of the IL-2 proliferation assay as quantified by CellTiter-Glo (Promega) luminescence-based cell viability assay. Both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[36] FIG. 10 shows results of protein cleavage assay. Fusion protein ACP16 was run on an SDS-PAGE gel in both cleaved and uncleaved form. As can be seen in the gel, cleavage was complete.

[37] FIGs. 11A-11B are a series of graphs depicting results from a HEK-Blue IL-12 reporter assay performed on human p40/murine p35 IL12 fusion proteins and recombinant human IL12 (Rec hIL-12). Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue® (InvivoGen). Results confirm that IL12 protein fusion proteins are active.

[38] FIGs. 12A-12F show a series of graphs depicting the results of HEK-blue assay of four IL-12 fusion proteins, before and after cleavage by MMP9. Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen). The data show greater activity in the cleaved IL12 than in the full fusion protein. Constructs tested were ACP06 (FIG. 12A), ACP07 (FIG. 12C), ACP08 (FIG. 12B), ACP09 (FIG. 12D), ACP10 (FIG. 12E), ACP11 (FIG. 12F)

[39] FIG. 13 shows results of protein cleavage assay. Fusion protein ACP11 was run on an SDS-PAGE gel in both cleaved and uncleaved form. As can be seen in the gel, cleavage was complete.

[40] FIG. 14 is a schematic which depicts a non-limiting example of an inducible cytokine protein, wherein the construct is activated upon protease cleavage of a linker attached between two subunits of the cytokine.

[41] FIGs. 15A-15D are graphs depicting results from a HEK-Blue assay performed on human p40/murine p35 IL12 fusion proteins and recombinant human IL12 (Rec hIL-12). Results confirm that IL12 protein fusion proteins are active. Each proliferation assay was performed with HSA or without HSA.

[42] FIGs. 16A-16F are a series of graphs showing activity of exemplary IFN γ fusion proteins compared to activity of mouse IFN γ control using WEHI 279 cell survival assay. Each assay was

performed with medium containing HSA (+HSA) or not containing HSA (-HSA). Each fusion protein comprises an anti-HSA binder, and both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[43] FIGs. 17A-17F are a series of graphs showing activity of exemplary IFN γ fusion proteins compared to activity of mouse IFN γ control using B16 reporter assay. Each assay was performed with medium containing HSA (+HSA) or not containing HSA (-HSA). Each fusion protein comprises an anti-HSA binder, and both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[44] FIGs. 18A-18B shows results of protein cleavage assay. Two constructs, ACP31 (IFN- α fusion protein; FIG. 18A) and ACP55 (IFN- γ fusion protein; 18B), were run on an SDS-PAGE gel in both cleaved and uncleaved form. As can be seen in the gel, cleavage was complete.

[45] FIGs. 19A-19B are a series of graphs showing activity of exemplary IFN γ fusion proteins compared to activity of mouse IFN γ control using B16 reporter assay. Each assay was performed with culture medium containing HSA, and each fusion protein comprises an anti-HSA binder. Both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[46] FIGs. 20A-20B are a series of graphs showing activity of exemplary IFN α fusion proteins compared to activity of mouse IFN α control using a B16 reporter assay. Each assay was performed with medium containing HSA, and each fusion protein comprises an anti-HSA binder. Both uncleaved and MMP9 protease cleaved versions of the fusion protein were used in each assay.

[47] FIGs. 21A-21D are a series of graphs depicting the results of tumor growth studies using the MC38 cell line. FIG. 21A-21C show the effect of IFN γ and IFN γ fusion proteins on tumor growth when injected intraperitoneally (IP) using different dosing levels and schedules (μ g=micrograms, BID=twice daily, BIW= twice weekly, QW= weekly). FIG. 21D shows the effect of intratumoral (IT) injection of IFN γ and IL-2 on tumor growth.

[48] FIGs. 22A-22B are a series of graphs showing activity of exemplary IFN γ fusion proteins (ACP51 and ACP52) cleaved by MMP9 protease compared to activity of uncleaved fusion proteins using B16 reporter assay. Each fusion protein comprises an anti-HSA binder and a tumor targeting domain.

[49] FIGs. 23A-23B are a series of graphs showing activity of exemplary IFN γ fusion proteins (ACP53 and ACP54) cleaved by MMP9 protease compared to activity of uncleaved fusion proteins using B16 reporter assay. Each fusion protein comprises IFN γ directly fused to albumin.

[50] FIGs. 24A-24B are two graphs showing the stability of IL-2 fusion proteins containing Linker-1 (GPAGMKGL, SEQ ID NO: 196), Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) in human serum in normal patient and a cancer patient.

FIG. 24A depicts the stability of the IL-2 fusion proteins containing Linker-1 (GPAGMKGL, SEQ ID NO: 196), Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) at 24 hours. FIG. 24B depicts the stability of the IL-2 fusion proteins containing Linker-1 (GPAGMKGL, SEQ ID NO: 196), Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) at 72 hours.

[51] FIG. 25A and 25B are two graphs showing analysis of ACP16 (FIG. 25A) and ACP124 (FIG. 25B) in a HEKBlue IL-2 reporter assay in the presence of HSA. Circles depict the activity of the uncut polypeptide, squares depict activity of the cut polypeptide, and triangles depict IL-2 alone as a control. FIG. 25C is a graph showing results of a CTLL-2 proliferation assay. CTLL2 cells (ATCC) were plated in suspension at a concentration of 500,000 cells/well in culture media with or without 40 mg/ml human serum albumin (HSA) and stimulated with a dilution series of recombinant hIL2 or activatable hIL2 for 72 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved activatable ACP16 was tested. Cleaved activatable hIL2 was generated by incubation with active MMP9. Cell activity was assessed using a CellTiter-Glo (Promega) luminescence-based cell viability assay. Triangles show wild-type cytokine, circles depict intact fusion protein, and squares depict protease-cleaved fusion protein.

[52] FIGs. 26A-26C are a series of graphs showing activity of fusion proteins in an HEKBlue IL-12 reporter assay. FIG. 26A is a graph showing activity of cut and uncut ACP11 (a human p40/murine p35 IL12 fusion protein). FIG. 26B is a graph showing analysis of ACP91 (a chimeric IL-12 fusion protein). Squares depict activity of the uncut ACP91 polypeptide, and triangles depict the activity of the cut polypeptide (ACP91 + MMP9). EC50 values for each are shown in the table. FIG. 26C is a graph showing analysis of ACP136 (a chimeric IL-12 fusion protein). Squares depict activity of the uncut ACP136 polypeptide, and triangles depict the activity of the cut polypeptide (ACP136 + MMP9). EC50 values for each are shown in the table insert.

[53] FIGs. 27A-27F are a series of graphs showing that cleaved IL-12 polypeptides are active in a HEKBlue IL2 reporter assay. Fusion proteins are evaluated both uncut (circles) and cut (squares) form, and wild type IL2 is used as a control +HSA for FIGs. 27A-C; ACP131 is used as a control (triangles) for FIGs. 27D-27F. Shown are data for APC31 + HSA (FIG. 27A), ACP125 + HSA (FIG. 27B), ACP126 + HSA (FIG. 27C), ACP127 (FIG. 27D), ACP128 (FIG. 27E), and ACP129 (FIG. 27F). The EC50 values for each are shown in the table below each graph.

[54] FIGs. 28A-28N are a series of graphs depicting the activity of APC56 (FIG. 28A), APC57 (FIG. 28B), APC58 (FIG. 28C), APC59 (FIG. 28D), APC60 (FIG. 28E), APC61 + HSA (FIG. 28F), ACP30 + HSA (FIG. 28G), ACP73 (FIG. 28H), ACP70 + HSA (FIG. 28I), ACP71 (FIG. 28J), ACP72 (FIG. 28K), ACP 73 (FIG. 28L), ACP74 (FIG. 28M), and ACP75 (FIG. 28N) in a HEKBlue IFN α reporter assay. Each fusion was tested for its activity when cut (squares) and uncut (circles). Analysis of murine IFN γ is included in each graph as a comparator.

[55] FIGs. 29A-29B is two graphs showing results of analyzing ACP31 (mouse IFN α 1 fusion protein) and ACP11 (a human p40/murine p35 IL12 fusion protein) in a tumor xenograft model. FIG. 29A shows tumor volume over time in mice treated with 33 μ g ACP31 (circles), 110 μ g ACP31 (triangles), 330 μ g ACP31 (diamonds), and as controls 1 μ g murine wild type IFN α 1 (dashed line, squares) and 10 μ g mIFN α 1 (dashed line, small circles). Vehicle alone is indicated by large open circles. The data show tumor volume decreasing over time in a dose-dependent manner in mice treated with ACP31. FIG. 29B shows tumor volume over time in mice treated with 17.5 μ g ACP11 (squares), 175 μ g ACP31 (triangles), 525 μ g ACP31 (circles), and as controls 2 μ g ACP04 (dashed line, triangles) and 10 μ g ACP04 (dashed line, diamonds). Vehicle alone is indicated by large open circles. The data show tumor volume decreasing over time in a dose-dependent manner in mice treated with both ACP11 and ACP04 (a human p40/murine p35 IL12 fusion protein).

[56] FIGs. 30A-30F are a series of spaghetti plots showing tumor volume over time in a mouse xenograft tumor model in mice each treated with vehicle alone (FIG. 30A), 2 μ g ACP04 (FIG. 30B), 10 μ g ACP04 (FIG. 30C), 17.5 μ g ACP11 (FIG. 30D), 175 μ g ACP11 (FIG. 30E), and 525 μ g ACP11 (FIG. 30F). Each line represents a single mouse.

[57] FIGs. 31A-C depicts three graphs showing results of analyzing ACP16 and ACP124 in a tumor xenograft model. FIG. 31A shows tumor volume over time in mice treated with 4.4 μ g ACP16 (squares), 17 μ g ACP16 (triangles), 70 μ g ACP16 (downward triangles), 232 μ g ACP16 (dark circles), and as a comparator 12 μ g wild type IL-2 (dashed line, triangles) and 36 μ g wild type IL-2 (dashed line, diamonds). Vehicle alone is indicated by large open circles. The data show tumor volume decreasing over time in a dose-dependent manner in mice treated with ACP16 at higher concentrations. FIG. 31B shows tumor volume over time in mice treated with 17 μ g ACP124 (squares), 70 μ g ACP124 (triangles), 230 μ g ACP124 (downward triangles), and 700 μ g ACP124. Vehicle alone is indicated by large open circles. FIG. 31C shows tumor volume over time in mice treated with 17 μ g ACP16 (triangles), 70 μ g ACP16 (circles), 232 μ g ACP16 (dark circles), and as a comparator 17 μ g ACP124 (dashed line, triangles) 70 μ g ACP124 (dashed line, diamonds), 230 μ g ACP124 (dashed line, diamonds). Vehicle alone is indicated by dark downward triangles. The data show tumor volume decreasing over time in a dose-dependent manner in mice treated with ACP16, but not ACP124.

[58] FIGs. 32A-32C are a series of spaghetti plots showing activity of fusion proteins in an MC38 mouse xenograft model. Each line in the plots is a single mouse.

[59] FIG. 33 is a graph showing tumor volume over time in a mouse xenograft model showing tumor growth in control mice (open circles) and AP16-treated mice (squares).

[60] FIGs. 34A-34D are a series of survival plots showing survival of mice over time after treatment with cleavable fusion proteins. FIG. 34A shows data for mice treated with vehicle alone (gray line), 17 μ g ACP16 (dark line), and 1 μ g ACP124 (dashed line). FIG. 34B shows data for

mice treated with vehicle alone (gray line), 70 μ g ACP16 (dark line), and 70 μ g ACP124 (dashed line). FIG. 34C shows data for mice treated with vehicle alone (gray line), 232 μ g ACP16 (dark line), and 230 μ g ACP124 (dashed line). FIG. 34D shows data for mice treated with vehicle alone (gray line), 232 μ g ACP16 (dark line), and 700 μ g ACP124 (dashed line).

[61] FIG. 35 a series of spaghetti plots showing activity of fusion proteins in an MC38 mouse xenograft model. All mouse groups were given four doses total except for the highest three doses of APC132, wherein fatal toxicity was detected after 1 week/2 doses. Shown are vehicle alone (top), 17, 55, 70, and 230 μ g ACP16 (top full row), 9, 28, 36, and 119 μ g ACP132 (middle full row), and 13, 42, 54, and 177 μ g ACP21 (bottom full row). Each line in the plots represents an individual animal.

[62] FIG. 36 is a schematic illustrating a substrate cleavage activity in conditioned complete (+FBS) media by FRET endpoint assay across four cell lines. The ratio of tumor vs control activity was approximated by averaging the three tumor cell lines and comparing to the control myofibroblast cell line where signal was detectable. FIG. 36 discloses SEQ ID NOs: 201, 198, 197, 196 and 195, respectively, in order of appearance.

[63] FIG. 37 is a schematic illustrating ADAM17_2 substrate kinetics in cell culture conditioned media. FIG. 37 discloses SEQ ID NO: 235.

[64] FIG. 38 is a schematic illustrating FAP α _1 substrate kinetics in conditioned media. FIG. 38 discloses SEQ ID NO: 197.

[65] FIG. 39 is a schematic illustrating FAP α _1 substrate kinetics in cell lysates. FIG. 39 discloses SEQ ID NO: 197.

[66] FIG. 40 is a schematic illustrating MMP9_1 substrate kinetics in cell lysates. FIG. 40 discloses SEQ ID NO: 196.

[67] FIG. 41 is a schematic illustrating a substrate cleavage activity in cell lysates by FRET endpoint assay. FIG. 41 discloses SEQ ID NOS 198 and 197, respectively, in order of appearance.

[68] FIG. 42 is a schematic illustrating CTSL1_1 substrate kinetics in cell lysates. FIG. 42 discloses SEQ ID NO: 198.

[69] FIG. 43 is a schematic illustrating MMP14_1 substrate kinetics in cell lysates. FIG. 43 discloses SEQ ID NO: 195.

[70] FIG. 44 is a schematic illustrating calculated concentration of enzyme equivalents per cell culture-derived sample. FIG. 44 discloses SEQ ID NOS 201, 198, 197, 197, 196 and 195, respectively, in order of appearance.

[71] FIG. 45 is a schematic illustrating an enzyme progress curve for CTSL1 cleavage of CSTL1_2 vs CTSL1_1. FIG. 45 discloses SEQ ID NOS 198, 199 and 236, respectively, in order of appearance.

[72] FIG. 46 is a schematic illustrating 30-mer cleavage of CTSL1_1 (ALFKSSFP, SEQ ID

NO: 198) vs CTSL1_2 (ALFFSSPP, SEQ ID NO: 199).

[73] FIG. 47 is a schematic illustrating susceptibility of the CTSL1 FRET substrates to CTSK cleavage. Rates of product formation were measured as a specific activity in units $\text{pmol min}^{-1} \mu\text{g}^{-1}$. The threshold value for the reference substrate, Z-LR-AMC is shown as a dashed line. FIG. 47 discloses SEQ ID NOS 198 and 199, respectively, in order of appearance.

[74] FIG. 48 is a schematic illustrating 30-mer Substrate Degradation by MMP9. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 48 discloses SEQ ID NOS 204, 205, 214, 216, 202, 217, 203, 211, 219, 207, 215, 212, 213, 206, 208, 209, 210, 218 and 220, respectively, in order of appearance.

[75] FIG. 49 is a schematic illustrating tandem MMP14_1 Motif degradation by MMP9. Top: substrate degradation traces, modeled with first-order kinetics. Bottom: product formation traces showing complex kinetics. FIG. 49 discloses SEQ ID NOS 202-205, respectively, in order of appearance.

[76] FIG. 50 is a schematic illustrating 30-mer Substrate Degradation by FAP α . Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 50 discloses SEQ ID NOS 205, 204, 206, 217, 203, 218, 219, 213, 216, 207, 214, 210, 202, 211, 208, 209, 212, 215 and 220, respectively, in order of appearance.

[77] FIG. 51 is a schematic illustrating 30-mer Substrate Degradation by CTSL1. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 51 discloses SEQ ID NOS 207, 208, 202, 218, 219, 212, 215, 217, 211, 209, 214, 206, 213, 210, 216, 203, 204, 205 and 220, respectively, in order of appearance.

[78] FIG. 52 is a schematic illustrating 30-mer Substrate Degradation by ADAM17. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 52 discloses SEQ ID NOS 208, 209, 211, 214, 217, 219, 213, 218, 215, 210, 212, 216, 207, 206, 202, 203, 204, 205 and 220, respectively, in order of appearance.

[79] FIG. 53 is a schematic illustrating 30-mer Substrate Degradation by Factor Xa. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 53 discloses SEQ ID NOS 220, 206, 202, 214, 208, 209, 215, 210, 218, 217, 207, 213, 216, 211, 212, 219, 203, 204 and 205, respectively, in order of appearance.

[80] FIG. 54 is a schematic illustrating 30-mer Substrate Degradation by Thrombin. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 54 discloses SEQ ID NOS 220, 204, 202, 207, 205, 211, 212, 215, 209, 218, 219, 217, 210, 213, 216, 214, 208, 206 and 203, respectively, in order of appearance.

[81] FIG. 55 is a schematic illustrating 30-mer Substrate Degradation by hepsin. Ranking of the substrates by relative rates of degradation are shown with “+”; uncleaved substrates are indicated as “-“. FIG. 55 discloses SEQ ID NOS 220, 209, 216, 215, 210, 213, 206, 214, 212, 207,

217, 208, 211, 218, 219, 202, 203, 204 and 205, respectively, in order of appearance.

[82] FIGs. 56A-56C show western blots probed with an IL-2 antibody demonstrating the stability of ACP16 (FIG. 56A), ACP153 (FIG. 56B), and ACP157 (FIG. 56C) in 90% serum. Serum was pooled from three human donors. Constructs of interest were incubated with PBS, Serum, or MMP9 protease and cleavage was assessed at T=0 hours and at T=24 hours.

[83] FIG. 57A-57B show western blots probed with an IL-2 antibody demonstrating the stability of ACP153, ACP155, ACP156, ACP16, and ACP372 in 90% serum. Serum was pooled from three human donors. Constructs of interest were incubated with PBS, Serum, or MMP9 protease and cleavage was assessed at T=24 hours and at T=72 hours. FIG. 57A shows the result using serum from a human donor and FIG. 57B shows the result using serum from a mouse donor.

[84] FIGs. 58A-58D show a series of spaghetti plots showing activity of fusion proteins in an MC38 mouse xenograft model. Shown are vehicle alone (FIG. 58A, top), 17, 55, and 230 μ g ACP16 (FIG. 58A), 55 and 230 μ g ACP153 (FIG. 58B), 55 and 230 μ g ACP155 (FIG. 58C), and 55 and 230 μ g ACP156 (FIG. 58D). Each line in the plots represents an individual animal.

[85] FIG. 59 shows a graph depicting results from a STAT activation reporter assay performed on IL-2 fusion proteins and recombinant human IL2 (Rec hIL-2). Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen).

[86] FIG. 60 shows a graph depicting results from a STAT activation reporter assay performed on IL-2 fusion proteins and recombinant human IL2 (Rec hIL-2). Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen).

[87] FIG. 61 shows a graph depicting results from a STAT activation reporter assay performed on IL-2 fusion proteins and recombinant human IL2 (Rec hIL-2). Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen).

[88] FIG. 62 shows a graph depicting results from a STAT activation reporter assay performed on IL-2 fusion proteins and recombinant human IL2 (Rec hIL-2). Analysis was performed based on quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen).

[89] FIG. 63 shows a table reporting engineered cleavage substrates described herein and the extent of cleavage observed using relevant proteases. Flanking sequences are lower case, the first cleavable sequence is underlined, the second is in bold font, and the third cleavable sequence is in italics. In some cases, there is overlap between cleavable sequences, which are indicated accordingly. FIG. 63 discloses SEQ ID NOS 202-220, respectively, in order of appearance.

[90] FIG. 64 is a schematic of an inducible tetravalent antibody format.

[91] FIG. 65A-65B show that multivalent 4-1BB antibodies are able to inducibly agonize 4-1BB.

5. DETAILED DESCRIPTION

[92] This disclosure relates to novel separation moieties or linkers and to polypeptides, such as fusion proteins, that contain the linkers. The linkers are preferably protease cleavable and link a first amino acid sequence of interest (e.g. a first domain of interest) to a second amino acid sequence of interest (e.g. a second domain of interest).

[93] The disclosed separation moieties confer site-selectivity with regard to the action of the attached payload or payloads. The payload can be a therapeutic agent, a half-life extender, a blocking agent and the like, or any combination thereof. The separation moieties may be used to attach any payload of interest, including e.g. cytokines, antibodies, cell-based therapies, etc. The separation moieties may be used individually or be used in tandem, triplicate, quadruplicate, and so forth, as long as the separation moiety is smaller than about 100 amino acids. Individual separation moieties may be directly joined to each other, or may be interspersed with non-cleavable linkers, whichever promotes high efficiency and site-specificity.

[94] The various embodiments of the present disclosure are further described in detail in the paragraphs below.

[95] Unless otherwise defined, all terms of art, notations and other scientific terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a difference over what is generally understood in the art. The techniques and procedures described or referenced herein are generally well understood and commonly employed using conventional methodologies by those skilled in the art, such as, for example, the widely utilized molecular cloning methodologies described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* 4th ed. (2012) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. As appropriate, procedures involving the use of commercially available kits and reagents are generally carried out in accordance with manufacturer-defined protocols and conditions unless otherwise noted.

[96] "Cytokine" is a well-known term of art that refers to any of a class of immunoregulatory proteins (such as interleukin or interferon) that are secreted by cells especially of the immune system and that are modulators of the immune system. Cytokine polypeptides that can be used in the fusion proteins disclosed herein include, but are not limited to transforming growth factors, such as TGF- α and TGF- β (e.g., TGF β 1, TGF β 2, TGF β 3); interferons, such as interferon- α , interferon- β , interferon- γ , interferon-kappa and interferon-omega; interleukins, such as IL-1, IL-

1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-21 and IL-25; tumor necrosis factors, such as tumor necrosis factor alpha and lymphotoxin; transforming growth factor beta (TGF β) family proteins, chemokines (e.g., C-X-C motif chemokine 10 (CXCL10), CCL19, CCL20, CCL21), and granulocyte macrophage-colony stimulating factor (GM-CS), as well as fragments of such polypeptides that active the cognate receptors for the cytokine (i.e., functional fragments of the foregoing). "Chemokine" is a term of art that refers to any of a family of small cytokines with the ability to induce directed chemotaxis in nearby responsive cells.

[97] Cytokines are well-known to have short serum half-lives that frequently are only a few minutes. Even forms of cytokines that have altered amino acid sequences intended to extend the serum half-life yet retain receptor agonist activity typically also have short serum half-lives. As used herein, a "short-half-life cytokine" refers to a cytokine that has a substantially brief half-life circulating in the serum of a subject, such as a serum half-life that is less than 10, less than 15, less than 30, less than 60, less than 90, less than 120, less than 240, or less than 480 minutes. As used herein, a short half-life cytokine includes cytokines which have not been modified in their sequence to achieve a longer than usual half-life in the body of a subject and polypeptides that have altered amino acid sequences intended to extend the serum half-life yet retain receptor agonist activity. This latter case is not meant to include the addition of heterologous protein domains, such as a bona fide half-life extension element, such as serum albumin.

[98] A "conservative" amino acid substitution, as used herein, generally refers to substitution of one amino acid residue with another amino acid residue from within a recognized group which can change the structure of the peptide but biological activity of the peptide is substantially retained. Conservative substitutions of amino acids are known to those skilled in the art. Conservative substitutions of amino acids can include, but not limited to, substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D. For instance, a person of ordinary skill in the art reasonably expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the biological activity of the resulting molecule.

[99] "Sortases" are transpeptidases that modify proteins by recognizing and cleaving a carboxyl-terminal sorting signal embedded in or terminally attached to a target protein or peptide. Sortase A catalyzes the cleavage of the LPXTG motif (where X is any standard amino acid) (SEQ ID NO: 237) between the Thr and Gly residue on the target protein, with transient attachment of the Thr residue to the active site Cys residue on the enzyme, forming an enzyme-thioacyl intermediate. To complete transpeptidation and create the peptide-monomer conjugate, a biomolecule with an N-terminal nucleophilic group, typically an oligoglycine motif, attacks the

intermediate, displacing Sortase A and joining the two molecules.

[100] As used herein, the term “steric blocker” refers to a polypeptide or polypeptide moiety that can be covalently bonded to a cytokine polypeptide directly or indirectly through other moieties such as linkers, for example in the form of a chimeric polypeptide (fusion protein), but otherwise does not covalently bond to the cytokine polypeptide. A steric blocker can non-covalently bond to the cytokine polypeptide, for example through electrostatic, hydrophobic, ionic or hydrogen bonding. A steric blocker typically inhibits or blocks the activity of the cytokine moiety due to its proximity to the cytokine moiety and comparative size.

[101] As used and described herein, a “half-life extension element” is a part of the chimeric polypeptide that increases the serum half-life and improve pK, for example, by altering its size (e.g., to be above the kidney filtration cutoff), shape, hydrodynamic radius, charge, or parameters of absorption, biodistribution, metabolism, and elimination.

[102] The term “separation moiety” or “linker” as used herein refers to an amino acid sequence typically less than about 100 amino acids that connects or links a first amino acid sequence of interest (e.g., an amino acid sequence that folds to form a first protein domain) to a second amino acid sequence of interest (e.g., an amino acid sequence that folds to form a second protein domain) in a contiguous polypeptide chain. The separation moiety or linker typically include one or more protease cleavage sites and thus is protease cleavable. A “tandem linker” refers to a linker that comprises two or more protease cleavages sites which can be cleaved by the same or different proteases, and which can be arranged in any desired orientation, such as one cleavage site adjacent to another cleavage site, one cleavage site overlapping another cleavage site, one cleavage site following by another cleavage site with intervening amino acids between the two cleavage sites.

[103] As used herein, the terms “activatable,” “activate,” “induce,” and “inducible” refer to the ability of a protein, i.e. a cytokine, that is part of a conjugate, to bind its receptor and effectuate activity upon cleavage of additional elements from the conjugate.

[104] As used herein, “plasmids” or “viral vectors” are agents that transport the disclosed nucleic acids into the cell without degradation and include a promoter yielding expression of the nucleic acid molecule and/or polypeptide in the cells into which it is delivered.

[105] As used herein, the terms “peptide”, “polypeptide”, or “protein” are used broadly to mean two or more amino acids linked by a peptide bond. Protein, peptide, and polypeptide are also used herein interchangeably to refer to amino acid sequences. It should be recognized that the term polypeptide is not used herein to suggest a particular size or number of amino acids comprising the molecule and that a peptide of the invention can contain up to several amino acid residues or more.

[106] As used throughout, “subject” can be a vertebrate, more specifically a mammal (e.g. a human, horse, cat, dog, cow, pig, sheep, goat, mouse, rabbit, rat, and guinea pig), birds, reptiles,

amphibians, fish, and any other animal. The term does not denote a particular age or sex. Thus, adult and newborn subjects, whether male or female, are intended to be covered.

[107] As used herein, “patient” or “subject” may be used interchangeably and can refer to a subject with a disease or disorder (e.g. cancer). The term patient or subject includes human and veterinary subjects.

[108] As used herein the terms “treatment”, “treat”, “treating,” or grammatically related terms refer to a method of reducing the effects of a disease or condition or symptom of the disease or condition. Thus, in the disclosed method, treatment can refer to at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or substantially complete reduction in the severity of an established disease or condition or symptom of the disease or condition. For example, a method for treating a disease is considered to be a treatment if there is a 10% reduction in one or more symptoms of the disease in a subject as compared to a control. Thus, the reduction can be a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or any percent reduction in between 10% and 100% as compared to native or control levels. It is well understood in the art that treatment does not necessarily refer to a cure or complete ablation of the disease, condition, or symptoms of the disease or condition. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis.

[109] As used herein, the terms “prevent”, “preventing”, and “prevention” of a disease or disorder refers to an action, for example, administration of the chimeric polypeptide or nucleic acid sequence encoding the chimeric polypeptide, that occurs before or at about the same time a subject begins to show one or more symptoms of the disease or disorder, which inhibits or delays onset or exacerbation of one or more symptoms of the disease or disorder.

[110] As used herein, references to “decreasing”, “reducing”, or “inhibiting” include a change of at least about 10%, of at least about 20%, of at least about 30%, of at least about 40%, of at least about 50%, of at least about 60%, of at least about 70%, of at least about 80%, of at least about 90% or greater as compared to a suitable control level. Such terms can include but do not necessarily include complete elimination of a function or property, such as agonist activity.

[111] An “attenuated cytokine receptor agonist” is a cytokine receptor agonist that has decreased receptor agonist activity as compared to the cytokine receptor’s naturally occurring agonist. An attenuated cytokine agonist may have at least about 10X, at least about 50X, at least about 100X, at least about 250X, at least about 500X, at least about 1000X or less agonist activity as compared to the receptor’s naturally occurring agonist. When a fusion protein that contains a cytokine

polypeptide as described herein is described as “attenuated” or having “attenuated activity”, it is meant that the fusion protein is an attenuated cytokine receptor agonist.

[112] An “intact fusion protein” is a fusion protein in which no domain has been removed from the fusion protein, for example by protease cleavage. A domain may be removable by protease cleavage or other enzymatic activity, but when the fusion protein is “intact”, this has not occurred.

[113] As used herein “moiety” refers to a portion of a molecule that has a distinct function within that molecule, and that function may be performed by that moiety in the context of another molecule. A moiety may be a chemical entity with a particular function, or a portion of a biological molecule with a particular function. For example, a “blocking moiety” within a fusion protein is a portion of the fusion protein which is capable of blocking the activity of some or all of the fusion polypeptide. This may be a protein domain, such as serum albumin.

A. Separation Moiety or Linker

[114] The disclosure relates to novel protease cleavable separation moieties. As described herein, the protease cleavable separation moieties were designed so that the separation moieties are cleaved with high efficiency by proteases at a desired location (e.g., proteases that are selectively expressed or expressed at high levels in the tumor microenvironment) but are stable and not cleaved or cleaved with low efficiency in other locations (e.g., in the periphery, for example healthy tissue or serum).

[115] The protease cleavable separation moieties were designed using a process that included prioritizing proteases that would be suitable for cleaving the separation moieties based on expression in target indications, such as, expression in particular types of tumors (e.g., colon, lung, breast, melanoma). Multiple data sources for increased expression or specific expression of proteases in the target indications were used, including mRNA, proteomics and tissue staining data. The proteases were also prioritized based on their specific activity as well as intrinsic specificity, with high specific activity and high intrinsic activity preferred. Stability in the serum is an important design consideration, and to avoid potential off-target cleavage of the separation moieties by serum proteases, proteases that are not dependent on arginine in their substrate were selected, since many off-target enzymes are active towards arginine residues.

[116] Starting sequences for the design process were selected using a diverse peptide library as substrates for proteases with mass spectrometric detection of protease cleaved products to identify preferred sequence motifs for each candidate protease. For selected initial motifs, a new peptide library that was tailored to the preferred sequence motif for the candidate protease was designed, created and analyzed. The peptide motifs were also counter-screened for cleavage by the serum proteases thrombin and Factor Xa as well as the liver/kidney protease hepsin. This process yielded peptides containing sequence motifs that are cleaved by certain tumor-associated proteases (e.g.,

Matrix Metalloprotease 9 (MMP9), MMP14 and/or Cathepsin L) with high efficiency, but are stable (not cleaved or cleaved with low efficiency) in the serum or normal healthy tissue (e.g., by thrombin, Factor Xa, hepsin and the like). The separation moieties disclosed herein are efficient cleavage by human tumors and minimal cleavage by normal tissues or serum.

[117] This disclosure relates to separation moieties or linkers that connect a first amino acid sequence of interest (e.g. a first domain of interest) to a second amino acid sequence of interest (e.g. a second domain of interest). Typically, the first amino acid sequence of interest and the second amino acid sequence of interest are not found together in a naturally occurring protein. For example, the separation moiety can connect or link a first domain of interest to a second domain of interest in a fusion protein. The separation moiety is an amino acid sequence that can be of any suitable length, and preferably can be cleaved by a protease.

[118] The separation moieties disclosed herein can confer functionality, including flexibility as well as the ability to be cleaved. Flexible linkers are usually applied when joined domains requires a certain degree of movement or interaction. Cleavable linkers are introduced to release free and functional domains *in vivo* at a target site. The separation moieties disclosed herein serve to connect at least two domains of interest. The separation moieties can maintain cooperative inter-domain interactions or preserving biological activity. The separation moieties can join functional domains (e.g., a payload and half-life extension element) that are released from the separation moiety at a target site (e.g. a tumor microenvironment).

[119] In a preferred embodiment, the separation moiety is cleavable by a cleaving agent, e.g., an enzyme. Preferably, the separation moiety comprises a protease cleavage site. In some cases, the separation moiety comprises one or more cleavage sites. The separation moiety can comprise a single protease cleavage site. The separation moiety can also comprise 2 or more protease cleavage sites. For example, 2 cleavage sites, 3 cleavage sites, 4, cleavage sites, 5 cleavage sites, or more. In cases the separation moiety comprises 2 or more protease cleavage sites, the cleavage sites can be cleaved by the same protease or different proteases. A separation moiety comprising two or more cleavage sites is referred to as a "tandem linker." The two or more cleavage sites can be arranged in any desired orientation, including, but not limited to one cleavage site adjacent to another cleavage site, one cleavage site overlapping another cleavage site, or one cleavage site following by another cleavage site with intervening amino acids between the two cleavage sites.

[120] Of particular interest in the present invention are disease specific protease-cleavable linkers. Also preferred are protease-cleavable linkers that are preferentially cleaved at a desired location in the body, such as the tumor microenvironment, relative to the peripheral circulation. For example, the rate at which the protease-cleavable linker is cleaved in the tumor microenvironment can be at least about 10 times, at least about 100 times, at least about 1000 times or at least about 10,000 times faster in the desired location in the body, e.g., the tumor

microenvironment, in comparison to in the peripheral circulation (e.g., in plasma).

[121] Proteases known to be associated with diseased cells or tissues include but are not limited to serine proteases, cysteine proteases, aspartate proteases, threonine proteases, glutamic acid proteases, metalloproteases, asparagine peptide lyases, serum proteases, cathepsins, Cathepsin B, Cathepsin C, Cathepsin D, Cathepsin E, Cathepsin G, Cathepsin K, Cathepsin L, kallikreins, hK1, hK10, hK15, plasmin, collagenase, Type IV collagenase, stromelysin, Factor Xa, chymotrypsin-like protease, trypsin-like protease, elastase-like protease, subtilisin-like protease, actinidain, bromelain, calpain, caspases, caspase-3, Mifl-CP, papain, HIV-1 protease, HSV protease, CMV protease, chymosin, renin, pepsin, matriptase, legumain, plasmepsin, nepenthesin, metalloexopeptidases, metalloendopeptidases, matrix metalloproteases (MMP), MMP1, MMP2, MMP3, MMP8, MMP9, MMP13, MMP11, MMP14, urokinase plasminogen activator (uPA), enterokinase, prostate-specific antigen (PSA, hK3), interleukin-1 β converting enzyme, thrombin, FAP (FAP α), dipeptidyl peptidase, meprins, granzymes and dipeptidyl peptidase IV (DPPIV/CD26). Proteases capable of cleaving linker amino acid sequences (which can be encoded by the chimeric nucleic acid sequences provided herein) can, for example, be selected from the group consisting of a prostate specific antigen (PSA), a matrix metalloproteinase (MMP), an A Disintegrin and a Metalloproteinase (ADAM), a plasminogen activator, a cathepsin, a caspase, a tumor cell surface protease, and an elastase. The MMP can, for example, be matrix metalloproteinase 2 (MMP2), matrix metalloproteinase 9 (MMP9), matrix metalloproteinase 14 (MMP14). In addition, or alternatively, the linker can be cleaved by a cathepsin, such as, Cathepsin B, Cathepsin C, Cathepsin D, Cathepsin E, Cathepsin G, Cathepsin K and/or Cathepsin L. Preferably, the linker can be cleaved by MMP14 or Cathepsin L.

[122] Proteases useful for cleavage of linkers and for use in the methods disclosed herein are presented in Table 1, and exemplary proteases and their cleavage site are presented in Table 1a:

Table 1. Proteases relevant to inflammation and cancer

Protease	Specificity	Other aspects
<i>Secreted by killer T cells:</i>		
Granzyme B (grB)	Cleaves after Asp residues (asp-ase)	Type of serine protease; strongly implicated in inducing perforin-dependent target cell apoptosis
Granzyme A (grA)	trypsin-like, cleaves after basic residues	Type of serine protease;
Granzyme H (grH)	Unknown substrate specificity	Type of serine protease; <i>Other granzymes are also secreted by killer T cells, but not all are present in humans</i>
Caspase-8	Cleaves after Asp residues	Type of cysteine protease; plays essential role in TCR-induced cellular expansion-exact molecular role unclear

Protease	Specificity	Other aspects
Mucosa-associated lymphoid tissue (MALT1)	Cleaves after arginine residues	Type of cysteine protease; likely acts both as a scaffold and proteolytically active enzyme in the CBM-dependent signaling pathway
Tryptase	Targets: angiotensin I, fibrinogen, prourokinase, TGF β ; preferentially cleaves proteins after lysine or arginine residues	Type of mast cell-specific serine protease; trypsin-like; resistant to inhibition by macromolecular protease inhibitors expressed in mammals due to their tetrameric structure, with all sites facing narrow central pore; <i>also associated with inflammation</i>
<i>Associated with inflammation:</i>		
Thrombin	Targets: FGF-2, HB-EGF, Osteo-pontin, PDGF, VEGF	Type of serine protease; modulates activity of vascular growth factors, chemokines and extracellular proteins; strengthens VEGF-induced proliferation; induces cell migration; angiogenic factor; regulates hemostasis
Chymase	Exhibit chymotrypsin-like specificity, cleaving proteins after aromatic amino acid residues	Type of mast cell-specific serine protease
Carboxypeptidase A (MC-CPA)	Cleaves amino acid residues from C-terminal end of peptides and proteins	Type of zinc-dependent metalloproteinase
Kallikreins	Targets: high molecular weight kininogen, pro-urokinase	Type of serine protease; modulate relaxation response; contribute to inflammatory response; fibrin degradation
Elastase	Targets: E-cadherin, GM-CSF, IL-1, IL-2, IL-6, IL8, p38 ^{MAPK} , TNF α , VE-cadherin	Type of neutrophil serine protease; degrades ECM components; regulates inflammatory response; activates pro-apoptotic signaling
Cathepsin G	Targets: EGF, ENA-78, IL-8, MCP-1, MMP-2, MT1-MMP, PAI-1, RANTES, TGF β , TNF α	Type of serine protease; degrades ECM components; chemo-attractant of leukocytes; regulates inflammatory response; promotes apoptosis
PR-3	Targets: ENA-78, IL-8, IL-18, JNK, p38 ^{MAPK} , TNF α	Type of serine protease; promotes inflammatory response; activates pro-apoptotic signaling
Granzyme M (grM)	Cleaves after Met and other long, unbranched hydrophobic residues	Type of serine protease; only expressed in NK cells
Calpains	Cleave between Arg and Gly	Family of cysteine proteases; calcium-dependent; activation is involved in the process of numerous inflammation-associated diseases

Table 1a: Exemplary Proteases and Protease Recognition Sequences

Protease	Cleavage Domain Sequence	SEQ ID NO:
MMP7	KRALGLPG	3
MMP7	(DE)₈RPLALWRS(DR)₈	4
MMP9	PR(S/T)(L/I)(S/T)	5
MMP9	LEATA	6
MMP11	GGAANLVRGG	7
MMP14	SGRIGFLRTA	8
MMP	PLGLAG	9
MMP	PLGLAX	10
MMP	PLGC(me)AG	11
MMP	ESPAYYTA	12
MMP	RLQLKL	13
MMP	RLQLKAC	14
MMP2, MMP9, MMP14	EP(Cit)G(Hof)YL	15
Urokinase plasminogen activator (uPA)	SGRSA	16
Urokinase plasminogen activator (uPA)	DAFK	17
Urokinase plasminogen activator (uPA)	GGRR	18
Lysosomal Enzyme	GFLG	19
Lysosomal Enzyme	ALAL	20
Lysosomal Enzyme	FK	21
Cathepsin B	NLL	22
Cathepsin D	PIC(Et)FF	23
Cathepsin K	GGPRGLPG	24
Prostate Specific Antigen	HSSKLQ	25
Prostate Specific Antigen	HSSKLQL	26
Prostate Specific Antigen	HSSKLQEDA	27
Herpes Simplex Virus Protease	LVLASSSFGY	28
HIV Protease	GVSQNYPIVG	29
CMV Protease	GVVQASCRLA	30
Thrombin	F(Pip)RS	31
Thrombin	DPRSFL	32
Thrombin	PPRSFL	33
Caspase-3	DEVD	34
Caspase-3	DEVDP	35

Protease	Cleavage Domain Sequence	SEQ ID NO:
Caspase-3	KGSGDVEG	36
Interleukin 1 β converting enzyme	GWEHDG	37
Enterokinase	EDDDDKA	38
FAP	KQEQNPGST	39
Kallikrein 2	GKAFFR	40
Plasmin	DAFK	41
Plasmin	DVLK	42
Plasmin	DAFK	43
TOP	ALLLALL	44
	GPLGVRG	221
	IPVSLRSG	222
	VPLSLYSG	223
	SGESPAYTA	224

[123] Exemplary protease linkers include, but are not limited to kallikrein cleavable linkers, thrombin cleavable linkers, chymase cleavable linkers, carboxypeptidase A cleavable linkers, cathepsin cleavable linkers, elastase cleavable linkers, FAP cleavable linkers, ADAM cleavable linkers, PR-3 cleavable linkers, granzyme M cleavable linkers, a calpain cleavable linkers, a matrix metalloproteinase (MMP) cleavable linkers, a plasminogen activator cleavable linkers, a caspase cleavable linkers, a tryptase cleavable linkers, or a tumor cell surface protease. Specifically, MMP9 cleavable linkers, ADAM cleavable linkers, CTSL1 cleavable linkers, FAP α cleavable linkers, and cathepsin cleavable linkers. Some preferred protease-cleavable linkers are cleaved by a MMP and/or a cathepsin.

[124] The separation moieties disclosed herein are typically less than 100 amino acids. Such separation moieties can be of different lengths, such as from 1 amino acid (e.g., Gly) to 30 amino acids, from 1 amino acid to 40 amino acids, from 1 amino acid to 50 amino acids, from 1 amino acid to 60 amino acids, from 1 to 70 amino acids, from 1 to 80 amino acids, from 1 to 90 amino acids, and from 1 to 100 amino acids. In some embodiments, the linker is at least about 1, about 2, about 3, about 4, about 5, about 10, about 15, about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 55, about 60, about 65, about 70, about 75, about 80, about 85, about 90, about 95, or about 100 amino acids in length. Preferred linkers are typically from about 5 amino acids to about 30 amino acids.

[125] Preferably the lengths of linkers vary from 2 to 30 amino acids, optimized for each condition so that the linker does not impose any constraints on the conformation or interactions of

the linked domains.

[126] In some embodiments, the separation moiety comprises the sequence GPAGLYAQ (SEQ ID NO: 195); GPAGMKGL (SEQ ID NO: 196); PGGPAGIG (SEQ ID NO: 197); ALFKSSFP (SEQ ID NO: 198); ALFFSSPP (SEQ ID NO: 199); LAQRLRSS (SEQ ID NO: 200); LAQKLKSS (SEQ ID NO: 201); GALFKSSFPSSGGGPAGLYAQGGSGKGGSGK (SEQ ID NO: 202); RSGGGGPAGLYAQGGSGGPAGLYAQGGSGK (SEQ ID NO: 203); KGGGPAGLYAQGPAGLYAQGPAGLYAQGSR (SEQ ID NO: 204); RGGPAGLYAQGGPAGLYAQGGGPAGLYAQK (SEQ ID NO: 205); KGGALFKSSFPGGPAGIGPLAQKLKSSGGG (SEQ ID NO: 206); SGGPGGPAGIGALFKSSFPPLAQKLKSSGGG (SEQ ID NO: 207); RGPLAQKLKSSALFKSSFPGGPAGIGGGGK (SEQ ID NO: 208); GGGALFKSSFPPLAQKLKSSPGGPAGIGGGR (SEQ ID NO: 209); RGPGGPAGIGPLAQKLKSSALFKSSFPGGG (SEQ ID NO: 210); RGGPLAQKLKSSPGGPAGIGALFKSSFP GK (SEQ ID NO: 211); RSGGPAGLYAQALFKSSFPPLAQKLKSSGGG (SEQ ID NO: 212); GGPLAQKLKSSALFKSSFPGPAGLYAQGGG (SEQ ID NO: 213); GGALFKSSFPGPAGLYAQPLAQKLKSSGGGK (SEQ ID NO: 214); RGGALFKSSFPPLAQKLKSSGPAGLYAQGGK (SEQ ID NO: 215); RGGGPAGLYAQPLAQKLKSSALFKSSFP GG (SEQ ID NO: 216); SGPLAQKLKSSGPAGLYAQALFKSSFP GSK (SEQ ID NO: 217); KGGPGGPAGIGPLAQRLRSSALFKSSFPGR (SEQ ID NO: 218); KSGPGGPAGIGALFFSSPPLAQKLKSSGGR (SEQ ID NO: 219); or SGGFPRSGGSFNPRTFGSKRKRRGSRGGGG (SEQ ID NO: 220)

[127] Certain preferred separation moieties comprises the sequence GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198). The separation moieties disclosed herein can comprise one or more cleavage motif or functional variants that are the same or different. The separation moieties can comprise 1, 2, 3, 4, 5, or more cleavage motifs or functional variants. Separation moieties comprising 30 amino acids can contain 2 cleavage motifs or functional variants, 3 cleavage motifs or functional variants or more. A “functional variant” of a separation moiety retains the ability to be cleaved with high efficiency at a target site (e.g., a tumor microenvironment that expresses high levels of the protease) and are not cleaved or cleaved with low efficiency in the periphery (e.g., serum). For example, the functional variants retain at least about 50%, about 55%, about 60%, about 70%, about 80%, about 85%, about 95% or more of the cleavage efficiency of a separation moiety comprising any one of SEQ ID NOs. 195-220.

[128] The separation moieties comprising more than one cleavage motif can be selected from SEQ ID NOs: 195-201 and combinations thereof. Preferred separation moieties comprising more

than one cleavage motif comprise the amino acids selected from SEQ ID NO: 202-220.

[129] The separation moiety can comprise both ALFKSSFP (SEQ ID NO: 198) and GPAGLYAQ (SEQ ID NO: 195). The separation moiety can comprise two cleavage motifs that each have the sequence GPAGLYAQ (SEQ ID NO: 195). Alternatively or additionally, the separation moiety can comprise two cleavage motifs that each have the sequence ALFKSSFP (SEQ ID NO: 198). The separation moiety can comprise a third cleavage motif that is the same or different.

[130] In some embodiments, the separation moiety comprises an amino acid sequence that is at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least 99% identical to SEQ ID NOs: 195 to SEQ ID NO: 220 over the full length of SEQ ID NO: 195-220.

[131] The disclosure also relates to functional variants of separation moieties comprising SEQ ID NOs. 195-220. The functional variants of separation moieties comprising SEQ ID NOs: 195-220 generally differ from SEQ ID NOs. 195-220 by one or a few amino acids (including substitutions, deletions, insertions, or any combination thereof), and substantially retain their ability to be cleaved by a protease.

[132] The functional variants can contain at least one or more amino acid substitutions, deletions, or insertions relative to the separation moieties comprising SEQ ID NOs. 195-220. The functional variant can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acid alterations compared to the separation moieties comprising SEQ ID NOs. 195-220. In some preferred embodiments, the functional variant differs from the separation moiety comprising SEQ ID NOs. 195-220 by less than 10, less than 8, less than 5, less than 4, less than 3, less than 2, or one amino acid alterations, e.g., amino acid substitutions or deletions. In other embodiments, the functional variant may comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acid substitutions compared to SEQ ID NOs. 195-220. The amino acid substitution can be a conservative substitution or a non-conservative substitution, but preferably is a conservative substitution.

[133] In other embodiments, the functional variants of the separation moieties may comprise 1, 2, 3, 4, or 5 or more non-conservative amino acid substitutions compared the separation moieties comprising SEQ ID NOs: 195-220. Non-conservative amino acid substitutions could be recognized by one of skill in the art. The functional variant of the separation moiety preferably contains no more than 1, 2, 3, 4, or 5 amino acid deletions.

[134] The amino acid sequences disclosed in the separation moieties can be described by the relative linear position in the separation moiety with respect to the scissile bond. As will be well-understood by persons skilled in the art, separation moieties comprising 8 amino acid protease substrates (e.g., SEQ ID Nos: 195-201) contain amino acid at positions P4, P3, P2, P1, P1', P2', P3', P4', wherein the scissile bond is between P1 and P1'. For example, amino acid positions for

the separation moiety comprising the sequence GPAGLYAQ (SEQ ID NO: 195) can be described as follows:

G	P	A	G	L	Y	A	Q
P4	P3	P2	P1	P1'	P2'	P3'	P4'

[135] Amino acids positions for the separation moiety comprising the sequence ALFKSSFP (SEQ ID NO: 198) can be described as follows:

A	L	F	K	S	S	F	P
P4	P3	P2	P1	P1'	P2'	P3'	P4'

[136] Preferably, the amino acids surrounding the cleavage site (e.g., positions P1 and P1' for SEQ ID NOs: 195-201) are not substituted.

[137] In embodiments, the separation moiety comprises the sequence GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198) or a functional variant of SEQ ID NO: 195 or a function variant of SEQ ID NO: 198. As described herein, a functional variant of PAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198) can comprise one or more amino acid substitutions, and substantially retain their ability to be cleaved by a protease. Specifically, the functional variants of GPAGLYAQ (SEQ ID NO: 195) is cleaved by MMP14, and the functional variant of ALFKSSFP (SEQ ID NO: 198) is cleaved by Cathepsin L (CTSL1). The functional variants also retain their ability to be cleaved with high efficiency at a target site (e.g., a tumor microenvironment that expresses high levels of the protease). For example, the functional variants of GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198) retain at least about 50%, about 55%, about 60%, about 70%, about 80%, about 85%, about 95% or more of the cleavage efficiency of a separation moiety comprising amino acid sequence GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198), respectively.

[138] Preferably, the functional variant of GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198) comprise no more than 1, 2, 3, 4, or 5 conservative amino acid substitutions compared to GPAGLYAQ (SEQ ID NO: 195) or ALFKSSFP (SEQ ID NO: 198). Preferably, the amino acids at position P1 and P1' are not substituted. The amino acids at positions P1 and P1' in SEQ ID NO: 195 are G and L, and the amino acids at positions P1 and P1' in SEQ ID NO: 198 are K and S.

[139] The functional variant of GPAGLYAQ (SEQ ID NO: 195) can preferably comprise one or more of the following: a) an arginine amino acid substitution at position P4, b) a leucine, valine, asparagine, or proline amino acid substitution at position P3, c) a asparagine amino acid substitution at position P2, d) a histidine, asparagine, or glycine amino acid substitution at position

P1, e) a asparagine, isoleucine, or leucine amino acid substitution at position P1', f) a tyrosine or arginine amino acid substitution at position P2', g) a glycine, arginine, or alanine amino acid substitution at position P3', h) or a serine, glutamine, or lysine amino acid substitution at position P4'. The following amino acid substitutions are disfavored in functional variants of GPAGLYAQ (SEQ ID NO: 195): a) arginine or isoleucine at position P3, b) alanine at position P2, c) valine at position P1, d) arginine, glycine, asparagine, or threonine at position P1', e) aspartic acid or glutamic acid at position P2', f) isoleucine at position P3', g) valine at position P4'. In some embodiments, the functional variant of GPAGLYAQ (SEQ ID NO: 195) does not comprise an amino acid substitution at position P1 and/or P1'.

[140] The amino acid substitution of the functional variant of GPAGLYAQ (SEQ ID NO: 195) preferably comprises an amino acid substitution at position P4 and/or P4'. For example, the functional variant of GPAGLYAQ (SEQ ID NO: 195) can comprises a leucine at position P4, or serine, glutamine, lysine, or phenylalanine at position P4. Alternatively or additionally, the functional variant of GPAGLYAQ (SEQ ID NO: 195) can comprises a glycine, phenylalanine, or a proline at position P4'.

[141] In some embodiments, the amino acid substitutions at position P2 or P2' of GPAGLYAQ (SEQ ID NO: 195) are not preferred.

[142] In some embodiments, the functional variant of GPAGLYAQ (SEQ ID NO: 195) comprises the amino acid sequence selected from SEQ ID NOs: 258-331. Specific functional variants of GPAGLYAQ (SEQ ID NO: 195) include GPLGLY AQ (SEQ ID NO: 295), and GPAGLKGA (SEQ ID NO: 285).

[143] The functional variants of LFKSSFP (SEQ ID NO: 198) preferably comprises hydrophobic amino acid substitutions. The functional variant of LFKSSFP (SEQ ID NO: 198) can preferably comprise one or more of the following: (a) lysine, histidine, serine, glutamine, leucine, proline, or phenylalanine at position P4; (b) lysine, histidine, glycine, proline, asparagine, phenylalanine at position P3; (c) arginine, leucine, alanine, glutamine, or histidine at position P2; (d) phenylalanine, histidine, threonine, alanine, or glutamine at position P1; (e) histidine, leucine, lysine, alanine, isoleucine, arginine, phenylalanine, asparagine, glutamic acid, or glycine at position P1', (f) phenylalanine, leucine, isoleucine, lysine, alanine, glutamine, or proline at position P2'; (g) phenylalanine, leucine, glycine, serine, valine, histidine, alanine, or asparagine at position P3'; and phenylalanine, histidine, glycine, alanine, serine, valine, glutamine, lysine, or leucine.

[144] The inclusion of aspartic acid and/or glutamic acid in functional variants of SEQ ID NO:198 are generally disfavored and avoided. The following amino acid substitutions are also disfavored in functional variants of LFKSSFP (SEQ ID NO: 198): (a) alanine, serine, or glutamic acid at position P3; (b) proline, threonine, glycine, or aspartic acid at position P2; (c) proline at

position P1; (d) proline at position P1'; (e) glycine at position P2'; (f) lysine or glutamic acid at position P3'; (g) aspartic acid at position P4'.

[145] The amino acid substitution of the functional variant of LFKSSFP (SEQ ID NO: 198) preferably comprises an amino acid substitution at position P4 and/or P1. In some embodiments, an amino acid substitution of the functional variant of LFKSSFP (SEQ ID NO: 198) at position P4' is not preferred.

[146] In some embodiments, the functional variant of LFKSSFP (SEQ ID NO: 198) comprises the amino acid sequence selected from SEQ ID NOs: 332-408. Specific functional variants of LFKSSFP (SEQ ID NO: 198) include ALFFSSPP (SEQ ID NO: 199), ALFKSFPP (SEQ ID NO: 381), ALFKSLPP (SEQ ID NO: 382); ALFKHSPP (SEQ ID NO: 370); ALFKSIPP (SEQ ID NO: 383); ALFKSSLP (SEQ ID NO: 390); or SPFRSSRQ (SEQ ID NO: 333).

[147] The separation moieties disclosed herein can form a stable complex under physiological conditions with the amino acid sequences (e.g. domains) that they link, while being capable of being cleaved by a protease. For example, the separation moiety is stable (e.g., not cleaved or cleaved with low efficiency) in the circulation and cleaved with higher efficiency at a target site (i.e. a tumor microenvironment). Accordingly, fusion polypeptides that include the linkers disclosed herein can, if desired, have a prolonged circulation half-life and/or lower biological activity in the circulation in comparison to the components of the fusion polypeptide as separate molecular entities. Yet, when in the desired location (e.g., tumor microenvironment) the linkers can be efficiently cleaved to release the components that are joined together by the linker and restoring or nearly restoring the half-life and biological activity of the components as separate molecular entities.

[148] The separation moiety desirably remains stable in the circulation for at least 2 hours, at least 5 hours, at least 10 hours, at least 15 hours, at least 20 hours, at least 24 hours, at least 30 hours, at least 35 hours, at least 40 hours, at least 45 hours, at least 50 hours, at least 60 hours, at least 65 hours, at least 70 hours, at least 80 hours, at least 90 hours, or longer.

[149] In some embodiments, the separation moiety is cleaved by less than 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 20%, 5%, or 1% in the circulation as compared to the target location. The separation moiety is also stable in the absence of an enzyme capable of cleaving the linker. However, upon exposure to a suitable enzyme (i.e., a protease), the separation moiety is cleaved resulting in separation of the linked domain.

B. Polypeptides and Compositions Comprising a Separation Moiety

[150] The separation moieties disclosed herein can be used in a wide range of applications. Without limitation, they are suitable for use in fusion proteins. As further described herein, the separation moieties are particularly useful for preparing therapeutic fusion proteins in which the

therapeutic of biological activity of the fusion protein is attenuated and the attenuation is removed upon cleavage of the separation moiety. The separation moieties can also be used to conjugate a variety of payloads such as therapeutic and/or diagnostic agents to carriers or targeting agents (e.g., antibodies and fragments of antibodies, nanoparticles). Suitable methods for prepare such conjugates are well-known in the art, see for example, *Bioconjugate Techniques*, Third Ed., G.T. Hermanson (Ed.) Academic Press 2013. Exemplary payloads include, but are not limited to cytokines, antibodies, cell based-therapies, antibiotics, cytotoxic drugs, or other recombinant polypeptide complexes. Of specific interest are separation moieties that are suitable for use in conjugation with payloads that target or are targeted to tumor microenvironments.

[151] This disclosure relates recombinant polypeptides in which a separation moiety as disclosed herein links a first amino acid sequence of interest (e.g. a first domain of interest) to a second amino acid sequence of interest (e.g. a second domain of interest). Typically, the first amino acid sequence of interest and the second amino acid sequence of interest are not found together in a naturally occurring protein. Preferred linkers are SEQ ID NOS:195 – 220. In embodiments, at least one of the first amino acid sequence of interest and the second amino acid sequence of interest is the amino acid sequence of a therapeutic polypeptide. In some embodiments in which at least one of the first amino acid sequence of interest and the second amino acid sequence of interest is the amino acid sequence of a therapeutic polypeptide, the other amino acid sequence of interest can be the amino acid sequence of a targeting polypeptide, a half-life extending polypeptide and/or a blocking polypeptide.

[152] The polypeptides that contain a separation moiety can be represented by Formula I: [D1]-[L1]-[D2], wherein D1 is a first amino acid sequence of interest (e.g., a domain of interest), L1 is a separation moiety that connects or links D1 to D2; and D2 is a second amino acid sequence of interest (e.g., a second domain of interest). Preferably, L1 is a protease-cleavable separation moiety, and more preferably L1 comprises or consists of any of SEQ ID NOS: 195-220.

[153] The polypeptides can also be represented by Formula II: [D1]-[L1]-[D2]-[L2]-[D3], wherein D1 is a first amino acid of interest (e.g., a domain of interest), L1 and L2 are each, independently, a linker; D2 is a second amino acid sequence of interest (e.g., a domain of interest), and D3 is a third amino acid of interest (e.g., a domain of interest), wherein at least one of L1 and L2 is a protease-cleavable separation moiety, and preferably at least one of L1 and L2 comprises or consists of any of SEQ ID NOS: 195-220.

[154] The polypeptide can also be represented by Formula III: [D1]-[L1]-[D2]-[L2]-[D3]-[L3]-[D4], wherein D1 is a first amino acid of interest (e.g., a domain of interest), L1, L2 and L3 are each, independently, a linker; D2 is a second amino acid sequence of interest (e.g., a domain of interest), D3 is a third amino acid of interest (e.g., a domain of interest); and wherein D4 is a fourth amino acid of interest (e.g., a domain of interest), wherein at least one of L1, L2 and L3 is a

protease-cleavable separation moiety, and preferably at least one of L1, L2 and L3 comprises or consists of any of SEQ ID NOS: 195-220.

[155] Additional specific applications of the separation moieties are described in more detail herein.

i. Delivery of payloads

[156] The separation moieties as described herein can be used to attach a therapeutic drug-moiety. In this approach, a therapeutic drug moiety is attached to the separation moiety to make a therapeutic drug moiety complex. The separate drug moiety complex can be a prodrug that is inactive until the target protease cleaves the prodrug, releasing the free drug.

ii. Antibody-drug conjugates

[157] Another example of use of the separation moiety is in the field of antibody-drug conjugates (ADCs) that are mainly directed toward the treatment of cancer. ADCs typically are an antibody that is linked to a cytotoxic moiety, such as a cytotoxic drug. The ADCs discriminate between the healthy and diseased cells and provide targeted delivery of drug (e.g. cytotoxic drug) to diseased cells. ADCs typically comprise an antibody that targets a tumor marker that is specific to tumor cells, whereupon the antibody attaches itself to the tumor cell, causing the ADC to be absorbed into the cell, which enables the cytotoxic component to be released to kill the tumor cell. A key aspect of ADCs is the provision of a stable linker between the antibody component and the cytotoxic agent. In such applications, linkers may be cleavable or non-cleavable. For non-cleavable linkers, the antibody, linker and cytotoxic unit is incorporated into the tumor cell. The nature of the linker typically determines the release profile of the cytotoxic agent. For example, cleavable linkers between the antibody and the cytotoxic agent are typically catalyzed by enzymes in the tumor cell or in the tumor microenvironment, wherein the antibody and the cytotoxic agents are cleaved to release the cytotoxic agent.

[158] In certain embodiments, the separation moiety disclosed herein links or connects a drug moiety to an antibody moiety.

iii. Peptide-drug conjugates

[159] The separation moieties disclosed herein are suitable for use in peptide-drug conjugates. These compounds typically comprise a cytotoxic payload and linker; however, instead of antibodies, peptide-drug conjugates are equipped with peptides that have the ability to penetrate tumors, thus allowing the cargo to be delivered inside the tumor. In some embodiments, the separation moiety links or connects a cytotoxic payload with a peptide. The peptide-drug conjugate remains stable and has no biological activity until the target protease cleaves the separation moiety.

iv. Inducible Adoptive Cell Therapy

[160] The separation moieties disclosed herein are suitable for use in constructs engineered for use in adoptive cell transfer (ACT) therapies. The field of adoptive cell transfer (ACT) is currently comprised of chimeric antigen receptor (CAR) engineered T cells (and next generation therapies) that target T cells to cell surface expressed targets (e.g., tumor cells expressing surface targets), and T cell receptor (TCR) engineered T cells that can target intracellular antigens.

[161] In one embodiment, the separation moiety is used to tether a targeting moiety to a CAR construct. A CAR replaces the endogenous TCR complex with a new receptor that uses a fragment of a human or mouse antibody to bind to targets outside of a cancer cell. The antibody fragment is linked to various signaling proteins inside T-cell which mediate receptor activation when the CAR binds to its target. Porter et al., (2011) NEJM, 365:725-733; Grupp et al., (2013) NEJM, 368:1509-1518; US10221245/ WO/2014/153270, Treatment of cancer using humanized anti-CD19 chimeric antigen receptor.

[162] In another embodiment, the separation moiety is used to tether a targeting moiety to a TCR construct. A TCR is based on the gene for the protein receptor that is already naturally present in T-cells. The gene for a desired TCR can be discovered in a single patient, e.g., a patient that is able to mount an effective immune response against a type of cancer. This gene can then be introduced into other patients by incorporating it into TCR T cell constructs or reengineered to improve the binding interaction with its MHC target. Guy et al., (2013) Nat Immunol., 14(3):262-70; Kuhns et al., (2012) Front Immunol., 25;3:159; Fesnak et al., (2016) Nat Rev Cancer, 16(9): 566-581.

[163] These types of engineered T cells, whether autologous or allogenic, comprise engineered T cell receptor components that comprise a targeting agent such as an isolated human or humanized antibody. In CAR-T and TCR-T cells, the binding affinity of the targeting moiety may be affected by the steric, chemical, or flexibility properties of the separation moiety tethering the targeting moiety to the rest of the construct and the T cell. The separation moieties disclosed herein are suitable for use with engineered constructs for making CAR T and TCR T cells.

v. Antigen-binding Proteins

[164] The separation moieties disclosed herein are suitable for use in antigen-binding proteins. An "antigen-binding protein" (ABP) is a protein comprising one or more antigen-binding domains that specifically bind to an antigen or epitope. In some embodiments, the antigen-binding domain binds the antigen or epitope with specificity and affinity similar to that of naturally occurring antibodies. Typically, the separation moieties link a polypeptide that blocks the antigen-binding site of the ABP from binding to its cognate antigen. But when the separation moiety is cleaved the blocking polypeptide can diffuse away from the ABP antigen binding site and the ABP can bind to

its cognate antigen. Exemplary binding polypeptides that can block the antigen binding site of an ABP include steric blocker such as human serum albumin, and peptides that interact with one or more of the complementarity determining regions (CDRs) in the antigen binding site of the ABP. Such blocking peptides can be obtained by screening libraries or by screening peptide fragments of the cognate antigen of an ABP of interest. Typically, when the ABP contains an antigen binding site of an antibody, the separation moiety and blocker will be bonded to the amino terminus of the antibody light chain, or the amino terminus of the antibody heavy chain, so that the blocker is tethered close to and will readily block the antigen binding site. Suitable variations of this approach are used when the ABP contains an alternative scaffold for the binding site. Similarly, when single chain antibody binding sites are used, such as scFV of dAb, the blocker-separation moiety is typically bonded to the amino terminus near the antigen binding site. In certain embodiments, the ABP comprises an antibody binding site that comprises a VH and a VL, and the blocker-separation moiety is bonded to the amino terminus of the VL.

[165] The ABP can be an antibody (e.g., the first and second antigen binding domains are in the form of an antibody). Preferably, at least one antigen binding domain of the ABP is in the form of an antibody. In another preferred embodiment, a first or second antigen binding domain is in the form of an antibody, and a first or second antigen binding domain is in the form of an antigen-binding fragment (e.g., the first antigen binding domain is an antibody and the second antigen binding domain is an antigen-binding fragment. Alternatively, the first antigen binding domain is an antigen binding fragment and the second antigen binding domain is an antibody).

[166] In some embodiments, the ABP consists of an antibody. In some embodiments, the ABP consists essentially of an antibody. In some embodiments, the ABP comprises an alternative scaffold. In some embodiments, the ABP consists of an alternative scaffold. In some embodiments, the ABP consists essentially of an alternative scaffold. In some embodiments, the ABP comprises an antibody fragment. In some embodiments, the ABP consists of an antibody fragment. In some embodiments, the ABP consists essentially of an antibody fragment.

[167] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with an antibody. The term “antibody” is used herein in its broadest sense and includes certain types of immunoglobulin molecules comprising one or more antigen-binding domains that specifically bind to an antigen or epitope. An antibody specifically includes intact antibodies (e.g., intact immunoglobulins), antibody fragments, and multi-specific antibodies. An antibody is one type of ABP.

[168] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with an antigen binding protein comprising an alternative scaffold. The term “alternative scaffold” refers to a molecule in which one or more regions may be diversified to produce one or more antigen-binding domains that specifically bind to an antigen or epitope.

[169] In some embodiments, the antigen-binding domain binds the antigen or epitope with specificity and affinity similar to that of an antibody. Exemplary alternative scaffolds include those derived from fibronectin (e.g., Adnectins™), the β -sandwich (e.g., iMab), lipocalin (e.g., Anticalins®), EETI-II/AGRP, BPTI/LACI-D1/TTI-D2 (e.g., Kunitz domains), thioredoxin peptide aptamers, protein A (e.g., Affibody®), ankyrin repeats (e.g., DARPins), gamma-B-crystallin/ubiquitin (e.g., Affilins), CTLD3 (e.g., Tetranectins), Fynomers, and (LDLR-A module) (e.g., Avimers). Additional information on alternative scaffolds is provided in Binz et al., *Nat. Biotechnol.*, 2005 23:1257-1268; Skerra, *Current Opin. in Biotech.*, 2007 18:295-304; and Silacci et al., *J. Biol. Chem.*, 2014, 289:14392-14398; each of which is incorporated by reference in its entirety. An alternative scaffold is one type of ABP.

[170] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with an antibody fragment. An “antibody fragment” comprises a portion of an intact antibody, such as the antigen-binding or variable region of an intact antibody. Antibody fragments include, for example, Fv fragments, Fab fragments, F(ab')₂ fragments, Fab' fragments, scFv (sFv) fragments, and scFv-Fc fragments.

[171] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with one or more Fv, Fab, or F(ab')₂ fragments. “Fv” fragments comprise a non-covalently-linked dimer of one heavy chain variable domain and one light chain variable domain. “Fab” fragments comprise, in addition to the heavy and light chain variable domains, the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments may be generated, for example, by recombinant methods or by papain digestion of a full-length antibody. “F(ab')₂” fragments contain two Fab' fragments joined, near the hinge region, by disulfide bonds. F(ab')₂ fragments may be generated, for example, by recombinant methods or by pepsin digestion of an intact antibody. The F(ab') fragments can be dissociated, for example, by treatment with 1-mercaptoethanol.

[172] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with an scFv or scFv-Fc. “Single-chain Fv” or “sFv” or “scFv” antibody fragments comprise a VH domain and a VL domain in a single polypeptide chain. The VH and VL are generally linked by a peptide linker. See Plückthun A. (1994). Any suitable linker may be used.

[173] In some embodiments, the linker is a (GGGGS)_n (SEQ ID NO: 231). In some embodiments, n=1, 2, 3, 4, 5, or 6. See *Antibodies from Escherichia coli*. In Rosenberg M. & Moore G. P. (Eds.), *The Pharmacology of Monoclonal Antibodies* vol. 113 (pp. 269-315). Springer-Verlag, New York, incorporated by reference in its entirety. “scFv-Fc” fragments comprise an scFv attached to an Fc domain. For example, an Fc domain may be attached to the C-terminal of the scFv. The Fc domain may follow the VH or VL, depending on the orientation of the variable domains in the scFv (i.e., VH-VL or VL-VH). Any suitable Fc domain known in the

art or described herein may be used. In some cases, the Fc domain comprises an IgG4 Fc domain.

[174] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a single domain antibody. The term “single domain antibody” refers to a molecule in which one variable domain of an antibody specifically binds to an antigen without the presence of another variable domain. Single domain antibodies, and fragments thereof, are described in Arabi Ghahroudi et al., *FEBS Letters*, 1998, 414:521-526 and Muyldermans et al., *Trends in Biochem. Sci.*, 2001, 26:230-245, each of which is incorporated by reference in its entirety. Single domain antibodies are also known as sdAbs or nanobodies.

[175] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a monospecific ABP. A “monospecific ABP” is an ABP that comprises one or more binding sites that specifically bind to the same epitope. An example of a monospecific ABP is a naturally occurring IgG molecule which, while divalent (i.e., having two antigen-binding domains), recognizes the same epitope at each of the two antigen-binding domains. The binding specificity may be present in any suitable valency.

[176] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a monoclonal antibody. The term “monoclonal antibody” refers to an antibody from a population of substantially homogeneous antibodies. A population of substantially homogeneous antibodies comprises antibodies that are substantially similar and that bind the same epitope(s), except for variants that may normally arise during production of the monoclonal antibody. Such variants are generally present in only minor amounts. A monoclonal antibody is typically obtained by a process that includes the selection of a single antibody from a plurality of antibodies. For example, the selection process can be the selection of a unique clone from a plurality of clones, such as a pool of hybridoma clones, phage clones, yeast clones, bacterial clones, or other recombinant DNA clones. The selected antibody can be further altered, for example, to improve affinity for the TNFR superfamily member proteins (“affinity maturation”), to humanize the antibody, to improve its production in cell culture, and/or to reduce its immunogenicity in a subject.

[177] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a chimeric antibody. The term “chimeric antibody” refers to an antibody in which a portion of the heavy and/or light chain is derived from a particular source or species, while the remainder of the heavy and/or light chain is derived from a different source or species.

[178] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a humanized antibody. “Humanized” forms of non-human antibodies are chimeric antibodies that contain minimal sequence derived from the non-human antibody. A humanized antibody is generally a human antibody (recipient antibody) in which residues from one or more CDRs are replaced by residues from one or more CDRs of a non-human antibody

(donor antibody). The donor antibody can be any suitable non-human antibody, such as a mouse, rat, rabbit, chicken, or non-human primate antibody having a desired specificity, affinity, or biological effect. In some instances, selected framework region residues of the recipient antibody are replaced by the corresponding framework region residues from the donor antibody. Humanized antibodies may also comprise residues that are not found in either the recipient antibody or the donor antibody. Such modifications may be made to further refine antibody function. For further details, see Jones et al., *Nature*, 1986, 321:522-525; Riechmann et al., *Nature*, 1988, 332:323-329; and Presta, *Curr. Op. Struct. Biol.*, 1992, 2:593-596, each of which is incorporated by reference in its entirety.

[179] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a human antibody. A “human antibody” is one which possesses an amino acid sequence corresponding to that of an antibody produced by a human or a human cell, or derived from a non-human source that utilizes a human antibody repertoire or human antibody-encoding sequences (e.g., obtained from human sources or designed de novo). Human antibodies specifically exclude humanized antibodies.

[180] In some embodiments, the ABPs provided herein specifically bind to the extracellular domain of a TNFR Superfamily Protein. In some embodiments, the TNFR superfamily protein is CD27, CD137, CD40, GITR, LT-betaR, CD30, HVEM, TNFR1, TNFR2, or OX-40. The TNFR superfamily protein may be expressed on the surface of any suitable target cell. In some embodiments, the target cell is a T cell. In some embodiments, the target cell is an effector T cell. In some embodiments, the target cell is a regulatory T cell. In some embodiments, the target cell is a natural killer (NK) cell. In some embodiments, the target cell is a natural killer T (NKT) cell. In some embodiments, the target cell is a B cell. In some embodiments, the target cell is a myeloid derived cell. In some embodiments, the target cell is a myeloid derived suppressor cell. In some embodiments, the target cell is a dendritic cell.

[181] In some embodiments, an ABP provided herein is an antibody. In some embodiments, an ABP provided herein is an antibody fragment. In some embodiments, an ABP provided herein is an alternative scaffold.

[182] In some embodiments, the ABPs provided herein comprise an immunoglobulin molecule. In some embodiments, the ABPs provided herein consist of an immunoglobulin molecule. In some embodiments, the ABPs provided herein consist essentially of an immunoglobulin molecule. In some aspects, the immunoglobulin molecule comprises an antibody. In some aspects, the immunoglobulin molecule consists of an antibody. In some aspects, the immunoglobulin molecule consists essentially of an antibody.

[183] In some embodiments, the ABPs provided herein comprise a light chain. In some aspects, the light chain is a kappa light chain. In some aspects, the light chain is a lambda light chain.

[184] In some embodiments, the ABPs provided herein comprise a heavy chain. In some aspects, the heavy chain is an IgA. In some aspects, the heavy chain is an IgD. In some aspects, the heavy chain is an IgE. In some aspects, the heavy chain is an IgG. In some aspects, the heavy chain is an IgM. In some aspects, the heavy chain is an IgG1. In some aspects, the heavy chain is an IgG2. In some aspects, the heavy chain is an IgG3. In some aspects, the heavy chain is an IgG4. In some aspects, the heavy chain is an IgA1. In some aspects, the heavy chain is an IgA2.

[185] In some embodiments, the separation moieties connect to a further blocking moiety.

[186] In some embodiments, the separation moieties are positioned such that movement of the antigen-binding protein subunits is restricted.

a. Multispecific and Monospecific Multivalent Antigen Binding Proteins

[187] In some embodiments, the separation moieties disclosed herein are suitable for use in conjunction with a multi-specific antigen binding protein (ABP). Multispecific ABP's provided herein bind more than one antigen. For instance, a multispecific antibody can bind 2 antigens, 3 antigens, binds 4 antigens, 5 antigens, or more. Alternatively, the multispecific ABP can bind 2 or more different epitopes. When the ABP binds two or more epitopes, the two or more different epitopes may be epitopes on the same antigen (e.g., a single TNFR superfamily protein molecule expressed by a cell) or on different antigens (e.g., different TNFR superfamily member protein molecules expressed by the same cell, or a TNFR superfamily member protein molecule and a non-TNFR Superfamily member protein molecule). In some aspects, a multi-specific ABP binds two different epitopes (i.e., a "bispecific ABP"). In some aspects, a multi-specific ABP binds three different epitopes (i.e., a "trispecific ABP"). In some aspects, a multi-specific ABP binds four different epitopes (i.e., a "quadspecific ABP"). In some aspects, a multi-specific ABP binds five different epitopes (i.e., a "quintspecific ABP"). In some aspects, a multi-specific ABP binds 6, 7, 8, or more different epitopes. Each binding specificity may be present in any suitable valency.

[188] In various embodiments, the antigen binding protein comprises a blocking domain. The separation moiety disclosed herein links the blocking domain to a first antigen binding domain and consequently the blocking domain inhibits (a) binding affinity or avidity of the antigen binding protein on the epitope and/or (b) agonist or antagonist activity of the antigen binding protein on the epitope. Preferably the separation moiety comprises a cleavage site. Upon cleavage of the separation moiety by a disease-specific enzyme (i.e., a protease) (a) the binding affinity or avidity of the antigen binding protein on the epitope and/or (b) agonist or antagonist activity of the antigen binding protein on the epitope is increased. In some embodiments, the antigen binding protein comprises an additional separation moiety that links or connects an additional blocking domain to an antigen binding domain. The additional separation moiety comprises a cleavage site.

[189] In various embodiments, the antigen binding protein comprises an additional blocking

domain (e.g., a second, third, or fourth blocking domain) that is/are operably linked to an antigen binding domain (e.g., the second, third, or fourth antigen binding domain) by a cleavable separation moiety as disclosed herein. The blocking domain can be a steric blocker or a specific blocker. In some embodiments, the steric blocker is independently selected from the group consisting of a fragment of the extracellular portion of the antibody binding protein, serum albumin, a fragment of serum albumin, and an antibody or antigen binding fragment thereof that binds serum albumin. In some embodiments, the specific blocker is independently selected from a fragment of an antibody or antigen-binding fragment thereof that binds to the first, second, third or fourth antigen binding domain of the antigen binding protein described herein. Preferably, the blocking domain is serum albumin or a fragment thereof, or an antibody or antigen binding fragment thereof that binds serum albumin.

[190] The antigen binding domain may further comprise a third antigen binding domain and a fourth antigen binding domain each with binding specificity for a target antigen. TNFR superfamily target antigens are well known in the art and are encompassed by this disclosure. Exemplary TNFR superfamily members include, but are not limited to CD27, CD30, CD137 (4-1BB), TNFR1 (CD120a), TNFR2 (CD120b), CD40, CD95 (Fas/Apo-1), HVEM, LT-betaR, GITR, nerve growth factor receptor or OX-40 (CD34). Preferred TNFR superfamily target antigens are CD27, CD137, and OX40.

[191] The antigen binding domains with specificity for an antigen can have binding specificity for the same epitope or for a different epitope on the same antigen. For example, the first, second and third antigen binding domains can have binding specificity for the same epitope, for a different epitope, two of the antigen binding domains may have binding specificity for the same epitope, or three of the antigen binding domains may have binding specificity for the same epitope. In some embodiments, the antigen binding protein may further comprise a fifth antigen binding domain with specificity for a tumor antigen.

[192] In some embodiments, the antigen binding polypeptide comprises an Fc region. In one format, two antigen binding domains can be located on each end of the Fc region. Another format comprises two antibody arms attached to the N terminus of the Fc region, each arm comprising 2 antigen binding domains. The antigen binding domain can also comprise a half-life extension domain. The half-life extension domain can be albumin, an antigen binding domain that recruits albumin, or an immunoglobulin Fc, or a fragment thereof. In some embodiments, the half-life extension domain is operably linked to the antigen binding polypeptide by a cleavable linker.

[193] The disclosure also relates to an antigen binding protein comprising at least a first polypeptide and a second polypeptide. The first polypeptide comprises at least a portion of an antibody heavy chain constant region and an antibody heavy chain variable region (VH). The second polypeptide comprises at least a portion of an antibody light chain constant region and an

antibody light chain variable region (VL). At least one of the first and second polypeptide further comprises a blocking domain that is operably linked to the VH or VL through a protease cleavable linker. The first polypeptide associates with the second polypeptide, and VH and VL form an antigen binding site with binding specificity for a target antigen (e.g., 4-1BB or CD27). The blocking domain inhibits binding of the antigen binding site to the target antigen (e.g., 4-1BB or CD27). In some embodiments, the first polypeptide further comprises a second VH, and the first polypeptide associates with two of said second polypeptides to form two VH/VL antigen binding sites that each have specificity for human 4-1BB. In some embodiments, the antigen binding protein is a dimer of the first polypeptide with the associated second polypeptides. The antigen binding protein can further comprise a third, fourth, fifth, or a sixth polypeptide.

[194] In some embodiments the ABP is monospecific multivalent ABP. Such formats can have a variety of structures and be prepared using suitable antibody engineering techniques. For example, a dual Fab antibody that contains a heavy chain with the structure VH-CH1-noncleavable linker-VH-CH1-CH2-CH3 can be prepared. Such a heavy chain can be expressed and can pair with two light chains. The heavy chain can dimerize through conventional interchain disulfide bonds to form an antibody format that contains 4 Fab antigen binding sites. In such formats, the separation moieties described herein can be bonded to the amino terminus of the light chain polypeptides to link a blocker to each of the Fab antigen binding sites. Other suitable monospecific, multivalent ABP formats can be readily envisioned by those of skill in the art. In one such example a heavy chain is prepared that has the structure VH-CH1-CH2-CH3-VH-CH1. Two such heavy chains and dimerize and associate with four light chains to form a monospecific tetravalent ABP format. In such formats, the separation moieties described herein can be bonded to the amino terminus of the light chain polypeptides to link a blocker to each of the Fab antigen binding sites. The binding activity of such monospecific tetravalent ABPs is masked by the blocking domain, and the mask is removed upon cleavage of the separation domain (e.g., in a tumor microenvironment) and the ABP is able to bind its cognate antigen.

[195] The monospecific multivalent ABP formats can have binding specificity for any desired antigen. In some embodiments, the monospecific multivalent ABP formats specifically bind to the extracellular domain of a TNFR Superfamily Protein, such as CD27, CD30, CD137 (4-1BB), TNFR1 (CD120a), TNFR2 (CD120b), CD40, CD95 (Fas/Apo-1), HVEM, LT-betaR, GITR, nerve growth factor receptor or OX-40 (CD34).

[196] In embodiments, the multivalent antigen binding protein can comprise a first antigen binding site with specificity for a target antigen (e.g., CD27 or TNFR1), second antigen binding site with specificity for the target antigen, a blocking polypeptide, at least one protease cleavable linker, and an optional half-life extension element. In such embodiments, a first blocking polypeptide is operably linked to the first antigen binding site by a protease cleavable linker and,

optionally, a second blocking polypeptide is operably linked to the second antigen binding site by a protease cleavable linker. Preferably, the blocking polypeptide is operably linked to the second antigen binding site by a protease cleavable linker. The blocking polypeptide inhibits (a) binding affinity or avidity of the antigen binding protein on the target antigen and/or (b) agonist or antagonist activity of the antigen binding protein on the target antigen and upon cleavage of the protease cleavable linker (a) the binding affinity or avidity of the antigen binding protein on the target antigen and/or (b) agonist activity of the antigen binding protein on the target antigen is increased. The optional half-life extension element can be operably linked to the first antigen binding site and/or second antigen binding site through a optionally protease cleavable linker. In such embodiments, the first and second binding sites can have specificity for the same epitope or different epitopes.

[197] In embodiments, the antigen binding protein further comprises a third antigen binding site with specificity for the same target antigen. The antigen binding protein can further comprise a fourth antigen binding site with specificity for the same target antigen as the first and second and third antigen binding sites. The third and fourth antigen binding sites can each further comprise a blocking polypeptide that is operable linked to the antigen-binding sight through a protease cleavable linker.

[198] This disclosure also relates to a tetravalent antigen binding protein that can comprise a first polypeptide that comprises at least a portion of an antibody heavy chain constant region and an antibody heavy chain variable region (VH) and a second polypeptide that comprises at least a portion of an antibody light chain constant region, an antibody light chain variable region (VL). The first polypeptide associates with the second polypeptide and the VH and VL form an antigen binding site with binding specificity for a target antigen. The tetravalent antigen binding protein further comprises a blocking polypeptide that is operably linked to the VH or VL through a protease cleavable linker. The blocking polypeptide inhibits binding of the antigen binding site to the target antigen. In some embodiments, the first polypeptide can further comprise a second VH. The first polypeptide associates with two of the second polypeptides to form two VH/VL antigen binding site that each have specificity for the target antigen.

[199] The tetravalent antigen binding protein can further comprise a third polypeptide, a fourth polypeptide, a fifth polypeptide, a sixth polypeptide, a seventh polypeptide, and an eight polypeptide. The third polypeptide can comprise at least a portion of an antibody heavy chain constant region and an antibody heavy chain variable region (VH). The fourth polypeptide can comprise at least a portion of an antibody light chain constant region, an antibody light chain variable region (VL). The fifth polypeptide can comprise at least portion of an antibody heavy chain constant region and an antibody heavy chain variable region (VH). The sixth polypeptide can comprise at least a portion of an antibody light chain constant region, an antibody light chain

variable region (VL). The seventh polypeptide can comprise at least portion of an antibody heavy chain constant region and an antibody heavy chain variable region (VH). The eighth polypeptide can comprise at least a portion of an antibody light chain constant region, an antibody light chain variable region (VL).

[200] In some embodiments at least one of the third and fourth polypeptide further comprises a blocking polypeptide that is operably linked to the VH or VL through a protease cleavable linker. The third polypeptide associates with the fourth polypeptide and VH and VL form an antigen binding site with binding specificity for a target antigen, and the blocking domain inhibits binding of the antigen binding site to the target antigen.

[201] In some embodiments, the fifth and sixth polypeptide further comprises a blocking polypeptide that is operably linked to the VH or VL through a protease cleavable linker. The fifth polypeptide associates with the sixth polypeptide and VH and VL form an antigen binding site with binding specificity for a target antigen, and the blocking domain inhibits binding of the antigen binding site to the target antigen.

[202] In some embodiments, the seventh and eighth polypeptide further comprises a blocking polypeptide that is operably linked to the VH or VL through a protease cleavable linker. The seventh polypeptide associates with the eighth polypeptide and VH and VL form an antigen binding site with binding specificity for a target antigen, and the blocking domain inhibits binding of the antigen binding site to the target antigen.

vi. Inducible Cytokines

[203] Disclosed herein are methods and compositions to engineer and use constructs comprising inducible cytokines. Cytokines are potent immune agonists, which lead to them being considered promising therapeutic agents for oncology. However, cytokines have a very narrow therapeutic window. Cytokines have short serum half-lives and are also considered to be highly potent.

Consequently, therapeutic administration of cytokines produces undesirable systemic effects and toxicities. These were exacerbated by the need to administer large quantities of cytokine in order to achieve the desired levels of cytokine at the intended site of cytokine action (e.g., a tumor).

Unfortunately, due to the biology of cytokines and inability to effectively target and control their activity, cytokines have not achieved the hoped-for clinical advantages in the treatment of tumors.

[204] Disclosed herein are fusion proteins that overcome the toxicity and short half-life problems that have severely limited the clinical use of cytokines in oncology. The fusion proteins contain cytokine polypeptides that have receptor agonist activity. But in the context of the fusion protein, the cytokine receptor agonist activity is attenuated, and the circulating half-life is extended. The fusion proteins include protease cleave sites, which are cleaved by proteases that are associated with a desired site of cytokine activity (e.g., a tumor), and are typically enriched or selectively present at the site of desired activity. Thus, the fusion proteins are preferentially (or

selectively) and efficiently cleaved at the desired site of activity to limit cytokine activity substantially to the desired site of activity, such as the tumor microenvironment. Protease cleavage at the desired site of activity, such as in a tumor microenvironment, releases a form of the cytokine from the fusion protein that is much more active as a cytokine receptor agonist than the fusion protein (typically at least about 100X more active than the fusion protein). The form of the cytokine that is released upon cleavage of the fusion protein typically has a short half-life, which is often substantially similar to the half-life of the naturally occurring cytokine, further restricting cytokine activity to the tumor microenvironment. Even though the half-life of the fusion protein is extended, toxicity is dramatically reduced or eliminated because the circulating fusion protein is attenuated, and active cytokine is targeted to the tumor microenvironment. The fusion proteins described herein, for the first time, enable the administration of an effective therapeutic dose of a cytokine to treat tumors with the activity of the cytokine substantially limited to the tumor microenvironment, and dramatically reduces or eliminates unwanted systemic effects and toxicity of the cytokine.

[205] In general, the therapeutic use of cytokines is strongly limited by their systemic toxicity. TNF, for example, was originally discovered for its capacity of inducing the hemorrhagic necrosis of some tumors, and for its *in vitro* cytotoxic effect on different tumoral lines, but it subsequently proved to have strong pro-inflammatory activity, which can, in case of overproduction conditions, dangerously affect the human body. As the systemic toxicity is a fundamental problem with the use of pharmacologically active amounts of cytokines in humans, novel derivatives and therapeutic strategies are now under evaluation, aimed at reducing the toxic effects of this class of biological effectors while keeping their therapeutic efficacy.

[206] IL-2 exerts both stimulatory and regulatory functions in the immune system and is, along with other members of the common γ chain (γc) cytokine family, central to immune homeostasis. IL-2 mediates its action by binding to IL-2 receptors (IL-2R), consisting of either trimeric receptors made of IL-2R α (CD25), IL-2R β (CD122), and IL-2R γ (γc , CD132) chains or dimeric $\beta\gamma$ IL-2Rs (1, 3). Both IL-2R variants are able to transmit signal upon IL-2 binding. However, trimeric $\alpha\beta\gamma$ IL-2Rs have a roughly 10–100 times higher affinity for IL-2 than dimeric $\beta\gamma$ IL-2Rs (3), implicating that CD25 confers high-affinity binding of IL-2 to its receptor but is not crucial for signal transduction. Trimeric IL-2Rs are found on activated T cells and CD4+ forkhead box P3 (FoxP3)+ T regulatory cells (Treg), which are sensitive to IL-2 *in vitro* and *in vivo*. Conversely, antigen-experienced (memory) CD8+, CD44 high memory-phenotype (MP) CD8+, and natural killer (NK) cells are endowed with high levels of dimeric $\beta\gamma$ IL-2Rs, and these cells also respond vigorously to IL-2 *in vitro* and *in vivo*.

[207] Expression of the high-affinity IL-2R is critical for endowing T cells to respond to low concentrations of IL-2 that is transiently available *in vivo*. IL-2R α expression is absent on naive

and memory T cells but is induced after antigen activation. IL-2R β is constitutively expressed by NK, NKT, and memory CD8⁺ T cells but is also induced on naive T cells after antigen activation. γ c is much less stringently regulated and is constitutively expressed by all lymphoid cells. Once the high-affinity IL-2R is induced by antigen, IL-2R signaling upregulates the expression of IL-2R α in part through Stat5-dependent regulation of *Il2ra* transcription (Kim et al., 2001). This process represents a mechanism to maintain expression of the high-affinity IL-2R and sustain IL-2 signaling while there remains a source of IL-2.

[208] IL-2 is captured by IL-2R α through a large hydrophobic binding surface surrounded by a polar periphery that results in a relatively weak interaction (K_d 10–8 M) with rapid on-off binding kinetics. However, the IL-2R α -IL-2 binary complex leads to a very small conformational change in IL-2 that promotes association with IL-2R β through a distinct polar interaction between IL-2 and IL-2R β . The pseudo-high affinity of the IL2/ α/β trimeric complex (i.e. K_d ~300 pM) clearly indicates that the trimeric complex is more stable than either IL2 bound to the α chain alone (K_d = 10 nM) or to the β chain alone (K_d = 450 nM). In any event, the IL2/ α/β trimer then recruits the γ chain into the quaternary complex capable of signaling, which is facilitated by the large composite binding site on the IL2-bound β chain for the γ chain.

[209] In other words, the ternary IL-2R α -IL-2R β -IL-2 complex then recruits γ c through a weak interaction with IL-2 and a stronger interaction with IL-2R β to produce a stable quaternary high-affinity IL-2R (K_d 10–11 M which is 10 pM). The formation of the high-affinity quaternary IL-2-IL-2R complex leads to signal transduction through the tyrosine kinases Jak1 and Jak3, which are associated with IL-2R β and γ c, respectively (Nelson and Willerford, 1998). The quaternary IL-2-IL-2R complex is rapidly internalized, where IL-2, IL-2R β , and γ c are rapidly degraded, but IL-2R α is recycled to the cell surface (Hémar et al., 1995; Yu and Malek, 2001). Thus, those functional activities that require sustained IL-2R signaling require a continued source of IL-2 to engage IL-2R α and form additional IL-2-IL-2R signaling complexes.

[210] Interleukin-15 (IL-15), another member of the 4-alpha-helix bundle family of cytokines, has also emerged as an immunomodulator for the treatment of cancer. IL-15 is initially captured via IL-15R α , which is expressed on antigen-presenting dendritic cells, monocytes and macrophages. IL-15 exhibits broad activity and induces the differentiation and proliferation of T, B and natural killer (NK) cells via signaling through the IL-15/IL-2-R- β (CD122) and the common γ chain (CD132). It also enhances cytolytic activity of CD8⁺ T cells and induces long-lasting antigen-experienced CD8⁺CD44 memory T cells. IL-15 stimulates differentiation and immunoglobulin synthesis by B cells and induces maturation of dendritic cells. It does not stimulate immunosuppressive T regulatory cells (Tregs). Thus, boosting IL-15 activity selectively in the tumor micro-environment could enhance innate and specific immunity and fight tumors (Waldmann et al., 2012). IL-15 was initially identified for its ability to stimulate T cell

proliferation in an IL-2-like manner through common receptor components (IL-2R/15R β - γ c) and signaling through JAK1/JAK3 and STAT3/STAT5. Like IL-2, IL-15 has been shown to stimulate proliferation of activated CD4-CD8-, CD4+CD8+, CD4+ and CD8+ T cells as well as facilitate the induction of cytotoxic T-lymphocytes, and the generation, proliferation and activation of NK cells (Waldmann et al., 1999). However, unlike IL-2 which is required to maintain forkhead box P3 (FOXP3)-expressing CD4+CD25+ Treg cells and for the retention of these cells in the periphery, IL-15 has little effect on Tregs (Berger et al., 2009). This is important as FOXP3-expressing CD4+CD25+ Tregs inhibit effector T cells, thereby inhibiting immune responses including those directed against the tumor. IL-2 also has a crucial role in initiating activation induced cell death (AICD), a process that leads to the elimination of self-reactive T cells, whereas IL-15 is an anti-apoptotic factor for T cells (Marks-Konczalik et al., 2000). IL-15 co-delivered with HIV peptide vaccines has been shown to overcome CD4+ T cell deficiency by promoting longevity of antigen-specific CD8+ T cells and blocking TRAIL-mediated apoptosis (Oh et al., 2008). Furthermore, IL-15 promotes the long-term maintenance of CD8+CD44^{hi} memory T cells (Kanegane et al., 1996).

[211] The importance of IL-15 and IL-15R α to T and NK cell development is further highlighted by the phenotype of IL-15R α ^{-/-} and IL-15^{-/-} mice. Knockout mice demonstrate decreased numbers of total CD8+ T cells, and are deficient in memory-phenotype CD8+ T cells, NK cells, NK/T cells and some subsets of intestinal intraepithelial lymphocytes, indicating that IL-15 provides essential positive homeostatic functions for these subsets of cells (Lodolce et al., 1996; Kennedy et al., 1998). The similarities in the phenotypes of these two strains of knockout mice suggest the importance of IL-15R α in maintaining physiologically relevant IL-15 signals.

[212] IL-15 is presented in trans by the IL-15 receptor alpha-chain to the IL-15R β γ c complex displayed on the surface of T cells and natural killer (NK) cells (Han et al., 2011). The IL-15R α -chain plays a role of chaperone protein, stabilizes, and increases IL-15 activity (Desbois et al., 2016). It has been shown that exogenous IL-15 may have a limited impact on patients with cancer due to its dependency on IL-15R α frequently downregulated in cancer patients. Therefore, the fusion protein RLI, composed of the sushi+ domain of IL15R α coupled via a linker to IL-15, has been suggested as an alternative approach to IL15 therapy (Bessard et al., 2009). It was found that administration of soluble IL-15/IL-15R α complexes greatly enhanced IL-15 serum half-life and bioavailability in vivo (Stoklasek et al., 2010).

[213] In addition to the effects on T and NK cells, IL-15 also has several effects on other components of the immune system. IL-15 protects neutrophils from apoptosis, modulates phagocytosis and stimulates the secretion of IL-8 and IL-1R antagonist. It functions through the activation of JAK2, p38 and ERK1/2 MAPK, Syk kinase and the NF- κ B transcriptional factor (Pelletier et al., 2002). In mast cells, IL-15 can act as a growth factor and an inhibitor of apoptosis.

In these cells IL-15 activates the JAK2/STAT5 pathway without the requirement of γ_c binding (Tagaya et al., 1996). IL-15 also induces B lymphocyte proliferation and differentiation, and increases immunoglobulin secretion (Armitage et al., 1995). It also prevents Fas-mediated apoptosis and allows induction of antibody responses partially independent of CD4-help (Demerci et al., 2004; Steel et al., 2010). Monocytes, macrophages and dendritic cells effectively transcribe and translate IL-15. They also respond to IL-15 stimulation. Macrophages respond by increasing phagocytosis, inducing IL-8, IL-12 and MCP-1 expression, and secreting IL-6, IL-8 and TNF α (Budagian et al., 2006). Dendritic cells incubated with IL-15 demonstrate maturation with increased CD83, CD86, CD40, and MHC class II expression, are also resistant to apoptosis, and show enhanced interferon- γ secretion (Anguille et al., 2009).

[214] IL-15 has also been shown to have effects on non-hematological cells including myocytes, adipocytes, endothelial and neural cells. IL-15 has an anabolic effect on muscle and may support muscle cell differentiation (Quinn et al., 1995). It stimulates myocytes and muscle fibers to accumulate contractile protein and is able to slow muscle wasting in rats with cancer-related cachexia (Figueras et al., 2004). IL-15 has also been shown to stimulate angiogenesis (Angiolillo et al., 1997) and induce microglial growth and survival (Hanisch et al., 1997).

[215] Interleukin-7 (IL-7), also of the IL-2/IL-15 family, is a well-characterized pleiotropic cytokine, and is expressed by stromal cells, epithelial cells, endothelial cells, fibroblasts, smooth muscle cells and keratinocytes, and following activation, by dendritic cells (Alpdogan et al., 2005). Although it was originally described as a growth and differentiation factor for precursor B lymphocytes, subsequent studies have shown that IL-7 is critically involved in T-lymphocyte development and differentiation. Interleukin-7 signaling is essential for optimal CD8 T-cell function, homeostasis and establishment of memory (Schluns et al., 2000); it is required for the survival of most T-cell subsets, and its expression has been proposed to be important for regulating T-cell numbers.

[216] IL-7 binds to a dimeric receptor, including IL-7R α and γ_c to form a ternary complex that plays fundamental roles in extracellular matrix remodeling, development, and homeostasis of T and B cells (Mazzucchelli and Durum, 2007). IL-7R α also cross-reacts to form a ternary complex with thymic stromal lymphopoietin (TSLP) and its receptor (TSLPR), and activates the TSLP pathway, resulting in T and dendritic cell proliferation in humans and further B cell development in mice (Leonard, 2002). Tight regulation of the signaling cascades activated by the complexes are therefore crucial to normal cellular function. Under-stimulation of the IL-7 pathway caused by mutations in the IL-7R α ectodomain inhibits T and B cell development, resulting in patients with a form of severe combined immunodeficiency (SCID) (Giliani et al., 2005; Puel et al., 1998).

[217] IL-7 has a potential role in enhancing immune reconstitution in cancer patients following cytotoxic chemotherapy. IL-7 therapy enhances immune reconstitution and can augment even

limited thymic function by facilitating peripheral expansion of even small numbers of recent thymic emigrants. Therefore, IL-7 therapy could potentially repair the immune system of patients who have been depleted by cytotoxic chemotherapy (Capitini et al., 2010).

[218] Interleukin-12 (IL-12) is a disulfide-linked heterodimer of two separately encoded subunits (p35 and p40), which are linked covalently to give rise to the so-called bioactive heterodimeric (p70) molecule (Lieschke et al., 1997; Jana et al., 2014). Apart from forming heterodimers (IL-12 and IL-23), the p40 subunit is also secreted as a monomer (p40) and a homodimer (p40₂). It is known in the art that synthesis of the heterodimer as a single chain with a linker connecting the p35 to the p40 subunit preserves the full biological activity of the heterodimer. IL-12 plays a critical role in the early inflammatory response to infection and in the generation of Th1 cells, which favor cell-mediated immunity. It has been found that overproduction of IL-12 can be dangerous to the host because it is involved in the pathogenesis of a number of autoimmune inflammatory diseases (e.g. MS, arthritis, type 1 diabetes).

[219] The IL-12 receptor (IL-12R) is a heterodimeric complex consisting of IL-12Rβ1 and IL-12Rβ2 chains expressed on the surface of activated T-cells and natural killer cells (Trinchieri et al., 2003). The IL-12Rβ1 chain binds to the IL-12p40 subunit, whereas IL-12p35 in association with IL-12Rβ2 confers an intracellular signaling ability (Benson et al., 2011). Signal transduction through IL-12R induces phosphorylation of Janus kinase (Jak2) and tyrosine kinase (Tyk2), that phosphorylate and activate signal transducer and activator of transcription (STAT)1, STAT3, STAT4, and STAT5. The specific cellular effects of IL-12 are due mainly to activation of STAT4. IL-12 induces natural killer and T-cells to produce cytokines, in particular interferon (IFN)γ, that mediate many of the proinflammatory activities of IL-12, including CD4+ T-cell differentiation toward the Th1 phenotype (Montepaone et al., 2014).

[220] Treg cells actively suppress activation of the immune system and prevent pathological self-reactivity and consequent autoimmune disease. Developing drugs and methods to selectively activate regulatory T cells for the treatment of autoimmune disease is the subject of intense research and, until the development of the present invention, which can selectively deliver active interleukins at the site of inflammation, has been largely unsuccessful. Treg are a class of CD4+CD25+ T cells that suppress the activity of other immune cells. Treg are central to immune system homeostasis and play a major role in maintaining tolerance to self-antigens and in modulating the immune response to foreign antigens. Multiple autoimmune and inflammatory diseases, including Type 1 Diabetes (T1D), Systemic Lupus Erythematosus (SLE), and Graft-versus-Host Disease (GVHD) have been shown to have a deficiency of Treg cell numbers or Treg function.

[221] Consequently, there is great interest in the development of therapies that boost the numbers and/or function of Treg cells. One treatment approach for autoimmune diseases being

investigated is the transplantation of autologous, ex vivo-expanded Treg cells (Tang, Q., et al, 2013, Cold Spring Harb. Perspect. Med., 3:1-15). While this approach has shown promise in treating animal models of disease and in several early stage human clinical trials, it requires personalized treatment with the patient's own T cells, is invasive, and is technically complex. Another approach is treatment with low dose Interleukin-2 (IL-2). Treg cells characteristically express high constitutive levels of the high affinity IL-2 receptor, IL2R $\alpha\beta\gamma$, which is composed of the subunits IL2R α (CD25), IL2R β (CD122), and IL2R γ (CD132), and Treg cell growth has been shown to be dependent on IL-2 (Malek, T. R., et al., 2010, Immunity, 33:153-65).

[222] Conversely, immune activation has also been achieved using IL-2, and recombinant IL-2 (Proleukin®) has been approved to treat certain cancers. High-dose IL-2 is used for the treatment of patients with metastatic melanoma and metastatic renal cell carcinoma with a long-term impact on overall survival.

[223] Clinical trials of low-dose IL-2 treatment of chronic GVHD (Koreth, J., et al., 2011, N Engl J Med., 365:2055-66) and HCV-associated autoimmune vasculitis patients (Saadoun, D., et al., 2011, N Engl J Med., 365:2067-77) have demonstrated increased Treg levels and signs of clinical efficacy. New clinical trials investigating the efficacy of IL-2 in multiple other autoimmune and inflammatory diseases have been initiated. The rationale for using so-called low dose IL-2 was to exploit the high IL-2 affinity of the trimeric IL-2 receptor which is constitutively expressed on Tregs while leaving other T cells which do not express the high affinity receptor in the inactivated state. Aldesleukin (marketed as Proleukin® by Prometheus Laboratories, San Diego, CA), the recombinant form of IL-2 used in these trials, is associated with high toxicity. Aldesleukin is approved for the treatment of metastatic melanoma and metastatic renal cancer, but its side effects are so severe that its use is only recommended in a hospital setting with access to intensive care (Web address: www.proleukin.com/assets/pdf/proleukin.pdf).

[224] The clinical trials of IL-2 in autoimmune diseases have employed lower doses of IL-2 in order to target Treg cells, because Treg cells respond to lower concentrations of IL-2 than many other immune cell types due to their expression of IL2R α (Klatzmann D, 2015 Nat Rev Immunol. 15:283-94). However, even these lower doses resulted in safety and tolerability issues, and the treatments used have employed daily subcutaneous injections, either chronically or in intermittent 5-day treatment courses. Therefore, there is a need for an autoimmune disease therapy that potentiates Treg cell numbers and function, that targets Treg cells more specifically than IL-2, that is safer and more tolerable, and that is administered less frequently.

[225] One approach that has been suggested for improving the therapeutic index of IL-2-based therapy is to use variants of IL-2 that are selective for Treg cells relative to other immune cells. IL-2 receptors are expressed on a variety of different immune cell types, including T cells, NK cells, eosinophils, and monocytes, and this broad expression pattern likely contributes to its

pleiotropic effect on the immune system and high systemic toxicity. In particular, activated T effector cells express IL2R $\alpha\beta\gamma$, as do pulmonary epithelial cells. But, activating T effector cells runs directly counter to the goal of down-modulating and controlling an immune response, and activating pulmonary epithelial cells leads to known dose-limiting side effects of IL-2 including pulmonary edema. In fact, the major side effect of high-dose IL-2 immunotherapy is vascular leak syndrome (VLS), which leads to accumulation of intravascular fluid in organs such as lungs and liver with subsequent pulmonary edema and liver cell damage. There is no treatment of VLS other than withdrawal of IL-2. Low-dose IL-2 regimens have been tested in patients to avoid VLS, however, at the expense of suboptimal therapeutic results.

[226] According to the literature, VLS is believed to be caused by the release of proinflammatory cytokines from IL-2-activated NK cells. However, there is strong evidence that pulmonary edema results from direct binding of IL-2 to lung endothelial cells, which expressed low to intermediate levels of functional $\alpha\beta\gamma$ IL-2Rs. And, the pulmonary edema associated with interaction of IL-2 with lung endothelial cells was abrogated by blocking binding to CD25 with an anti-CD25 monoclonal antibody (mAb), in CD25-deficient host mice, or by the use of CD122-specific IL-2/anti-IL-2 mAb (IL-2/mAb) complexes, thus preventing VLS.

[227] Treatment with interleukin cytokines other than IL-2 has been more limited. IL-15 displays immune cell stimulatory activity similar to that of IL-2 but without the same inhibitory effects, thus making it a promising immunotherapeutic candidate. Clinical trials of recombinant human IL-15 for the treatment of metastatic malignant melanoma or renal cell cancer demonstrated appreciable changes in immune cell distribution, proliferation, and activation and suggested potential antitumor activity (Conlon et. al., 2014). IL-15 is currently in clinical trials to treat various forms of cancer. However, IL-15 therapy is known to be associated with undesired and toxic effects, such as exacerbating certain leukemias, graft-versus-host disease, hypotension, thrombocytopenia, and liver injury. (Mishra A., et al., Cancer Cell, 2012, 22(5):645-55; Alpdogan O. et al., Blood, 2005, 105(2):866-73; Conlon KC et al., J Clin Oncol, 2015, 33(1):74-82.)

[228] IL-7 promotes lymphocyte development in the thymus and maintains survival of naive and memory T cell homeostasis in the periphery. Moreover, it is important for the organogenesis of lymph nodes (LN) and for the maintenance of activated T cells recruited into the secondary lymphoid organs (SLOs) (Gao et. al., 2015). In clinical trials of IL-7, patients receiving IL-7 showed increases in both CD4+ and CD8+ T cells, with no significant increase in regulatory T cell numbers as monitored by FoxP3 expression (Sportes et al., 2008). In clinical trials reported in 2006, 2008 and 2010, patients with different kinds of cancers such as metastatic melanoma or sarcoma were injected subcutaneously with different doses of IL-7. Little toxicity was seen except for transient fevers and mild erythema. Circulating levels of both CD4+ and CD8+ T cells increased significantly and the number of Treg reduced. TCR repertoire diversity increased after

IL-7 therapy. However, the anti-tumor activity of IL-7 was not well evaluated (Gao et. al., 2015). Results suggest that IL-7 therapy could enhance and broaden immune responses.

[229] IL-12 is a pleiotropic cytokine, the actions of which create an interconnection between the innate and adaptive immunity. IL-12 was first described as a factor secreted from PMA-induced EBV-transformed B-cell lines. Based on its actions, IL-12 has been designated as cytotoxic lymphocyte maturation factor and natural killer cell stimulatory factor. Due to bridging the innate and adaptive immunity and potently stimulating the production of IFN γ , a cytokine coordinating natural mechanisms of anticancer defense, IL-12 seemed ideal candidate for tumor immunotherapy in humans. However, severe side effects associated with systemic administration of IL-12 in clinical investigations and the very narrow therapeutic index of this cytokine markedly tempered enthusiasm for the use of this cytokine in cancer patients (Lasek et. al., 2014). Approaches to IL-12 therapy in which delivery of the cytokine is tumor-targeted, which may diminish some of the previous issues with IL-12 therapy, are currently in clinical trials for cancers.

[230] The direct use of IL-2 as an agonist to bind the IL-2R and modulate immune responses therapeutically has been problematic due its well-documented therapeutic risks, e.g., its short serum half-life and high toxicity. These risks have also limited the therapeutic development and use of other cytokines. New forms of cytokines that reduce these risks are needed. Disclosed herein are compositions and methods comprising IL-2 and IL-15 and other cytokines, functional fragments and muteins of cytokines as well as conditionally active cytokines designed to address these risks and provide needed immunomodulatory therapeutics.

[231] The present invention is designed to address the shortcomings of direct IL-2 therapy and therapy using other cytokines, for example using cytokine blocking moieties, e.g. steric blocking polypeptides, serum half-life extending polypeptides, targeting polypeptides, linking polypeptides, including protease cleavable linkers, and combinations thereof. Cytokines, including interleukins (e.g., IL-2, IL-7, IL-12, IL-15, IL-18, IL-21 IL-23), interferons (IFNs, including IFN α , IFN β and IFN γ), tumor necrosis factors (e.g., TNF α , lymphotoxin), transforming growth factors (e.g., TGF β 1, TGF β 2, TGF β 3), chemokines (C-X-C motif chemokine 10 (CXCL10), CCL19, CCL20, CCL21), and granulocyte macrophage-colony stimulating factor (GM-CS) are highly potent when administered to patients. As used herein, "chemokine" means a family of small cytokines with the ability to induce directed chemotaxis in nearby responsive cells. Cytokines can provide powerful therapy but are accompanied by undesired effects that are difficult to control clinically and which have limited the clinical use of cytokines. This disclosure relates to new forms of cytokines that can be used in patients with reduced or eliminated undesired effects. In particular, this disclosure relates to pharmaceutical compositions including chimeric polypeptides (fusion proteins), nucleic acids encoding fusion proteins and pharmaceutical formulations of the foregoing that contain cytokines or active fragments or muteins of cytokines that have decreased cytokine

receptor activating activity in comparison to the corresponding cytokine. However, under selected conditions or in a selected biological environment the chimeric polypeptides activate their cognate receptors, often with the same or higher potency as the corresponding naturally occurring cytokine. As described herein, this is typically achieved using a cytokine blocking moiety that blocks or inhibits the receptor activating function of the cytokine, active fragment or mutein thereof under general conditions but not under selected conditions, such as those present at the desired site of cytokine activity (e.g., an inflammatory site or a tumor).

[232] The chimeric polypeptides and nucleic acids encoding the chimeric polypeptides can be made using any suitable method. For example, nucleic acids encoding a chimeric polypeptide can be made using recombinant DNA techniques, synthetic chemistry or combinations of these techniques, and expressed in a suitable expression system, such as in CHO cells. Chimeric polypeptides can similarly be made, for example by expression of a suitable nucleic acid, using synthetic or semi-synthetic chemical techniques, and the like. In some embodiments, the blocking moiety can be attached to the cytokine polypeptide via sortase-mediated conjugation. "Sortases" are transpeptidases that modify proteins by recognizing and cleaving a carboxyl-terminal sorting signal embedded in or terminally attached to a target protein or peptide. Sortase A catalyzes the cleavage of the LPXTG motif (where X is any standard amino acid) (SEQ ID NO: 237) between the Thr and Gly residue on the target protein, with transient attachment of the Thr residue to the active site Cys residue on the enzyme, forming an enzyme-thioacyl intermediate. To complete transpeptidation and create the peptide-monomer conjugate, a biomolecule with an N-terminal nucleophilic group, typically an oligoglycine motif, attacks the intermediate, displacing Sortase A and joining the two molecules.

[233] To form the cytokine-blocking moiety conjugate, the cytokine polypeptide is first tagged at the N-terminus with a polyglycine sequence, or alternatively, with at the C-terminus with a LPXTG (SEQ ID NO: 237) motif. The blocking moiety or other element has respective peptides attached that serve as acceptor sites for the tagged polypeptides. For conjugation to domains carrying a LPXTG (SEQ ID NO: 237) acceptor peptide attached via its N-terminus, the polypeptide will be tagged with an N-terminal poly-glycine stretch. For conjugation to domain carrying a poly-glycine peptide attached via its C-terminus, the polypeptide will be tagged at its C-terminus with a LPXTG (SEQ ID NO: 237) sortase recognition sequence. Recognizing poly-glycine and LPXTG (SEQ ID NO: 237) sequences, sortase will form a peptide bond between polymer-peptide and tagged polypeptides. The sortase reaction cleaves off glycine residues as intermediates and occurs at room temperature.

[234] A variety of mechanisms can be exploited to remove or reduce the inhibition caused by the blocking moiety. For example, the pharmaceutical compositions can include a cytokine moiety and a blocking moiety, e.g. a steric blocking moiety, with a protease cleavable linker comprising a

protease cleavage site located between the cytokine and cytokine blocking moiety or within the cytokine blocking moiety. When the protease cleavage site is cleaved, the blocking moiety can dissociate from cytokine, and the cytokine can then activate cytokine receptor.

[235] Any suitable linker can be used. For example, the linker can comprise glycine-glycine, a sortase-recognition motif, or a sortase-recognition motif and a peptide sequence $(\text{Gly}_4\text{Ser})_n$ (SEQ ID NO: 238) or $(\text{Gly}_3\text{Ser})_n$ (SEQ ID NO: 239), wherein n is 1, 2, 3, 4 or 5. Typically, the sortase-recognition motif comprises a peptide sequence LPXTG (SEQ ID NO: 237), where X is any amino acid. In some embodiments, the covalent linkage is between a reactive lysine residue attached to the C-terminal of the cytokine polypeptide and a reactive aspartic acid attached to the N-terminal of the blocker or other domain. In other embodiments, the covalent linkage is between a reactive aspartic acid residue attached to the N-terminal of the cytokine polypeptide and a reactive lysine residue attached to the C-terminal of the blocker or another domain.

[236] Accordingly, as described in detail herein, the cytokine blocking moieties used can be steric blockers. As used herein, a “steric blocker” refers to a polypeptide or polypeptide moiety that can be covalently bonded to a cytokine polypeptide directly or indirectly through other moieties such as linkers, for example in the form of a chimeric polypeptide (fusion protein), but otherwise does not covalently bond to the cytokine polypeptide. A steric blocker can non-covalently bond to the cytokine polypeptide, for example through electrostatic, hydrophobic, ionic or hydrogen bonding. A steric blocker typically inhibits or blocks the activity of the cytokine moiety due to its proximity to the cytokine moiety and comparative size. The steric inhibition of the cytokine moiety can be removed by spatially separating the cytokine moiety from the steric blocker, such as by enzymatically cleaving a fusion protein that contains a steric blocker and a cytokine polypeptide at a site between the steric blocker and the cytokine polypeptide.

[237] As described in greater detail herein, the blocking function can be combined with or due to the presence of additional functional components in the pharmaceutical composition, such as a targeting domain, a serum half-life extension element, and protease-cleavable linking polypeptides. For example, a serum half-life extending polypeptide can also be a steric blocker.

[238] In the interest of presenting a concise disclosure of the full scope of the invention, aspects of the invention are described in detail using the cytokine IL-2 as an exemplary cytokine.

However, the invention and this disclosure are not limited to IL-2. It will be clear to a person of skill in the art that this disclosure, including the disclosed methods, polypeptides and nucleic acids, adequately describes and enables the use of other cytokines, fragments and muteins, such as IL-2, IL-7, IL-12, IL-15, IL-18, IL-21, IL-23, IFN α , IFN β , IFN γ , TNF α , lymphotoxin, TGF- β 1, TGF β 2, TGF β 3, GM-CSF, CXCL10, CCL19, CCL20, CCL21 and functional fragments or muteins of any of the foregoing.

[239] Various elements ensure the delivery and activity of IL-2 preferentially at the site of

desired IL-2 activity and to severely limit systemic exposure to the interleukin via a blocking and/or a targeting strategy preferentially linked to a serum half-life extension strategy. In this serum half-life extension strategy, the blocked version of interleukin circulates for extended times (preferentially 1-2 or more weeks) but the activated version has the typical serum half-life of the interleukin.

[240] By comparison to a serum half-life extended version, the serum half-life of IL-2 administered intravenously is only ~10 minutes due to distribution into the total body extracellular space, which is large, ~15 L in an average sized adult. Subsequently, IL-2 is metabolized by the kidneys with a half-life of ~2.5 hours. (Smith, K. "Interleukin 2 immunotherapy." *Therapeutic Immunology* 240 (2001)). By other measurements, IL-2 has a very short plasma half-life of 85 minutes for intravenous administration and 3.3 hours subcutaneous administration (Kirchner, G. I., et al., 1998, *Br J Clin Pharmacol.* 46:5-10). In some embodiments of this invention, the half-life extension element is linked to the interleukin via a linker which is cleaved at the site of action (e.g. by inflammation-specific proteases) releasing the interleukin's full activity at the desired site and also separating it from the half-life extension of the uncleaved version. In such embodiments, the fully active and free interleukin would have very different pharmacokinetic (pK) properties – a half-life of hours instead of weeks. In addition, exposure to active cytokine is limited to the site of desired cytokine activity (e.g., an inflammatory site or tumor) and systemic exposure to active cytokine, and associated toxicity and side effects, are reduced.

[241] Other cytokines envisioned in this invention have similar pharmacology (e.g. IL-15 as reported by Blood 2011 117:4787-4795; doi: doi.org/10.1182/blood-2010-10-311456) as IL-2 and accordingly, the designs of this invention address the shortcomings of using these agents directly, and provide chimeric polypeptides that can have extended half-life and/or be targeted to a site of desired activity (e.g., a site of inflammation or a tumor).

[242] If desired, IL-2 can be engineered to bind the IL-2R complex generally or one of the three IL-2R subunits specifically with an affinity that differs from that of the corresponding wild-type IL-2, for example to selectively activate Tregs or Teff (Effector T Cell). For example, IL-2 polypeptides that are said to have higher affinity for the trimeric form of the IL-2 receptor relative to the dimeric beta/gamma form of the IL-2 receptor in comparison to wild type IL-2 can have an amino acid sequence that includes one of the following sets of mutations with respect to SEQ ID NO: 1 (a mature IL-2 protein comprising amino acids 21–153 of human IL-2 having the UniProt Accession No. P60568-1): (a) K64R, V69A, and Q74P; (b) V69A, Q74P, and T101A; (c) V69A, Q74P, and I128T; (d) N30D, V69A, Q74P, and F103S; (e) K49E, V69A, A73V, and K76E; (f) V69A, Q74P, T101A, and T133N; (g) N30S, V69A, Q74P, and I128A; (h) V69A, Q74P, N88D, and S99P; (i) N30S, V69A, Q74P, and I128T; (j) K9T, Q11R, K35R, V69A, and Q74P; (k) A1T, M46L, K49R, E61D, V69A, and H79R; (l) K48E, E68D, N71T, N90H, F103S, and I114V; (m)

S4P, T10A, Q11R, V69A, Q74P, N88D, and T133A; (n) E15K, N30S Y31H, K35R, K48E, V69A, Q74P, and I92T; (o) N30S, E68D, V69A, N71A, Q74P, S75P, K76R, and N90H; (p) N30S, Y31C, T37A, V69A, A73V, Q74P, H79R, and I128T; (q) N26D, N29S, N30S, K54R, E67G, V69A, Q74P, and I92T; (r) K8R, Q13R, N26D, N30T, K35R, T37R, V69A, Q74P, and I92T; and (s) N29S, Y31H, K35R, T37A, K48E, V69A, N71R, Q74P, N88D, and I89V. This approach can also be applied to prepare muteins of other cytokines including interleukins (e.g., IL-2, IL-7, IL-12, IL-15, IL-18, IL-23), interferons (IFNs, including IFN α , IFN β and IFN γ), tumor necrosis factors (e.g., TNF α , lymphotoxin), transforming growth factors (e.g., TGF β 1, TGF β 2, TGF β 3) and granulocyte macrophage-colony stimulating factor (GM-CS). For example, muteins can be prepared that have desired binding affinity for a cognate receptor.

[243] As noted above, any of the mutant IL-2 polypeptides disclosed herein can include the sequences described; they can also be limited to the sequences described and otherwise identical to SEQ ID NO:1. Moreover, any of the mutant IL-2 polypeptides disclosed herein can optionally include a substitution of the cysteine residue at position 125 with another residue (e.g., serine) and/or can optionally include a deletion of the alanine residue at position 1 of SEQ ID NO:1.

[244] Another approach to improving the therapeutic index of an IL-2 based therapy is to optimize the pharmacokinetics of the molecule to maximally activate Treg cells. Early studies of IL-2 action demonstrated that IL-2 stimulation of human T cell proliferation *in vitro* required a minimum of 5-6 hours exposure to effective concentrations of IL-2 (Cantrell, D. A., et. al., 1984, Science, 224: 1312-1316). When administered to human patients, IL-2 has a very short plasma half-life of 85 minutes for intravenous administration and 3.3 hours subcutaneous administration (Kirchner, G. I., et al., 1998, Br J Clin Pharmacol. 46:5-10). Because of its short half-life, maintaining circulating IL-2 at or above the level necessary to stimulate T cell proliferation for the necessary duration necessitates high doses that result in peak IL-2 levels significantly above the EC50 for Treg cells or will require frequent administration. These high IL-2 peak levels can activate IL2R $\beta\gamma$ receptors and have other unintended or adverse effects, for example VLS as noted above. An IL-2 analog, or a multifunctional protein with IL-2 attached to a domain that enables binding to the FcRn receptor, with a longer circulating half-life than IL-2 can achieve a target drug concentration for a specified period of time at a lower dose than IL-2, and with lower peak levels. Such an IL-2 analog will therefore require either lower doses or less frequent administration than IL-2 to effectively stimulate Treg cells. Less frequent subcutaneous administration of an IL-2 drug will also be more tolerable for patients. A therapeutic with these characteristics will translate clinically into improved pharmacological efficacy, reduced toxicity, and improved patient compliance with therapy. Alternatively, IL-2 or muteins of IL-2 (herein, "IL-2*") can be selectively targeted to the intended site of action (e.g. sites of inflammation). This targeting can be achieved by one of several strategies, including the addition of domains to the administered agent

that comprise blockers of the IL-2 (or muteins) that are cleaved away or by targeting domains or a combination of the two.

[245] In some embodiments, IL-2* partial agonists can be tailored to bind with higher or lower affinity depending on the desired target; for example, an IL-2* can be engineered to bind with enhanced affinity to one of the receptor subunits and not the others. These types of partial agonists, unlike full agonists or complete antagonists, offer the ability to tune the signaling properties to an amplitude that elicits desired functional properties while not meeting thresholds for undesired properties. Given the differential activities of the partial agonists, a repertoire of IL-2 variants could be engineered to exhibit an even finer degree of distinctive signaling activities, ranging from almost full to partial agonism to complete antagonism.

[246] In some embodiments, the IL-2* has altered affinity for IL-2R α . In some embodiments, the IL-2* has a higher affinity for IL-2R α than wild-type IL-2. In other embodiments, the IL-2* has altered affinity for IL-2R β . In one embodiment, IL-2* has enhanced binding affinity for IL-2R β , e.g., the N-terminus of IL-2R β , that eliminates the functional requirement for IL-2R α . In another embodiment, an IL-2* is generated that has increased binding affinity for IL-2R β but that exhibited decreased binding to IL-2R γ , and thereby is defective IL-2R $\beta\gamma$ heterodimerization and signaling.

[247] Blocking moieties, described in further detail below, can also be used to favor binding to or activation of one or more receptors. In one embodiment, blocking moieties are added such that IL-2R $\beta\gamma$ binding or activation is blocked but IL-2R α binding or activation is not changed. In another embodiment, blocking moieties are added such that IL-2R α binding or activation is diminished. In another embodiment, blocking moieties are added such that binding to and or activation of all three receptors is inhibited. This blocking may be relievable by removal of the blocking moieties in a particular environment, for example by proteolytic cleavage of a linker linking one or more blocking moieties to the cytokine.

[248] A similar approach can be applied to improve other cytokines, particularly for use as immunostimulatory agents, for example for treating cancer. For example, in this aspect, the pharmacokinetics and/or pharmacodynamics of the cytokine (e.g., IL-2, IL-7, IL-12, IL-15, IL-18, IL-21, IL-23, IFN α , IFN β and IFN γ , TNF α , lymphotoxin, TGF- β 1, TGF β 2, TGF β 3, GM-CSF, CXCL10, CCL19, CCL20, and CCL21) can be tailored to maximally activate effector cells (e.g., effect T cells, NK cells) and/or cytotoxic immune response promoting cells (e.g., induce dendritic cell maturation) at a site of desired activity, such as in a tumor, but preferably not systemically.

[249] Thus, provided herein are pharmaceutical compositions comprising at least one cytokine polypeptide, such as interleukins (e.g., IL-2, IL-7, IL-12, IL-15, IL-18, IL-21, IL-23), interferons (IFNs, including IFN α , IFN β and IFN γ), tumor necrosis factors (e.g., TNF α , lymphotoxin), transforming growth factors (e.g., TGF- β 1, TGF β 2, TGF β 3), chemokines (e.g., CXCL10, CCL19,

CCL20, CCL21) and granulocyte macrophage-colony stimulating factor (GM-CS) or a functional fragment or mutein of any of the foregoing. The polypeptide typically also includes at least one linker amino acid sequence, wherein the amino acid sequence is in certain embodiments capable of being cleaved by an endogenous protease. In one embodiment, the linker comprises an amino acid sequence comprising HSKLQ (SEQ ID NO: 25), GPLGVRG (SEQ ID NO: 221), IPVSLRSG (SEQ ID NO: 222), VPLSLYSG (SEQ ID NO: 223), or SGESPAYYTA (SEQ ID NO: 224). In other embodiments, the chimeric polypeptide further contains a blocking moiety, e.g. a steric blocking polypeptide moiety, capable of blocking the activity of the interleukin polypeptide. The blocking moiety, for example, can comprise a human serum albumin (HSA) binding domain or an optionally branched or multi-armed polyethylene glycol (PEG). Alternatively, the pharmaceutical composition comprises a first cytokine polypeptide or a fragment thereof, and blocking moiety, e.g. a steric blocking polypeptide moiety, wherein the blocking moiety blocks the activity of the cytokine polypeptide on the cytokine receptor, and wherein the blocking moiety in certain embodiments comprises a protease cleavable domain. In some embodiments, blockade and reduction of cytokine activity is achieved simply by attaching additional domains with very short linkers to the N or C terminus of the interleukin domain. In such embodiments, it is anticipated the blockade is relieved by protease digestion of the blocking moiety or of the short linker that tethers the blocker to the interleukin. Once the domain is clipped or is released, it will no longer be able to achieve blockade of cytokine activity.

[250] The pharmaceutical composition e.g., chimeric polypeptide can comprise two or more cytokines, which can be the same cytokine polypeptide or different cytokine polypeptides. For example, the two or more different types of cytokines have complementary functions. In some examples, a first cytokine is IL-2 and a second cytokine is IL-12. In some embodiments, each of the two or more different types of cytokine polypeptides have activities that modulate the activity of the other cytokine polypeptides. In some examples of chimeric polypeptides that contain two cytokine polypeptides, a first cytokine polypeptide is T-cell activating, and a second cytokine polypeptide is non-T-cell-activating. In some examples of chimeric polypeptides that contain two cytokine polypeptides, a first cytokine is a chemoattractant, e.g., CXCL10, and a second cytokine is an immune cell activator.

[251] Preferably, the cytokine polypeptides (including functional fragments) that are included in the fusion proteins disclosed herein are not mutated or engineered to alter the properties of the naturally occurring cytokine, including receptor binding affinity and specificity or serum half-life. However, changes in amino acid sequence from naturally occurring (including wild type) cytokine are acceptable to facilitate cloning and to achieve desired expression levels, for example.

a. Blocking Moiety

[252] The blocking moiety can be any moiety that inhibits the ability of the cytokine to bind and/or activate its receptor. The blocking moiety can inhibit the ability of the cytokine to bind and/or activate its receptor sterically blocking and/or by noncovalently binding to the cytokine. Examples of suitable blocking moieties include the full length or a cytokine-binding fragment or mutein of the cognate receptor of the cytokine. Antibodies and fragments thereof including, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody a single chain variable fragment (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain of camelid-type nanobody (VHH), a sdAb and the like that bind the cytokine can also be used. Other suitable antigen-binding domain that bind the cytokine can also be used, include non-immunoglobulin proteins that mimic antibody binding and/or structure such as, anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPins, fynomers, kunitz domain peptides, monobodies, and binding domains based on other engineered scaffolds such as SpA, GroEL, fibronectin, lipocalin and CTLA4 scaffolds. Further examples of suitable blocking polypeptides include polypeptides that sterically inhibit or block binding of the cytokine to its cognate receptor. Advantageously, such moieties can also function as half-life extending elements. For example, a peptide that is modified by conjugation to a water-soluble polymer, such as PEG, can sterically inhibit or prevent binding of the cytokine to its receptor. Polypeptides, or fragments thereof, that have long serum half-lives can also be used, such as serum albumin (human serum albumin), immunoglobulin Fc, transferring and the like, as well as fragments and muteins of such polypeptides.

Antibodies and antigen-binding domains that bind to, for example, a protein with a long serum half-life such as HSA, immunoglobulin or transferrin, or to a receptor that is recycled to the plasma membrane, such as FcRn or transferrin receptor, can also inhibit the cytokine, particularly when bound to their antigen. Examples of such antigen-binding polypeptides include a single chain variable fragment (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain of camelid-type nanobody (VHH), a sdAb and the like. Other suitable antigen-binding domain that bind the cytokine can also be used, include non-immunoglobulin proteins that mimic antibody binding and/or structure such as, anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPins, fynomers, kunitz domain peptides, monobodies, and binding domains based on other engineered scaffolds such as SpA, GroEL, fibronectin, lipocalin and CTLA4 scaffolds.

[253] In illustrative examples, when IL-2 is the cytokine in the chimeric polypeptide, the blocking moiety can be the full length or fragment or mutein of the alpha chain of IL-2 receptor (IL-2R α) or beta (IL-2R β) or gamma chain of IL-2 receptor (IL-2R γ), an anti-IL-2 single-domain

antibody (sdAb) or scFv, an anti-CD25 antibody or fragment thereof, and anti-HSA sdAb or scFv, and the like.

b. In vivo Half-life Extension Elements

[254] Preferably, the chimeric polypeptides comprise an *in vivo* half-life extension element. Increasing the *in vivo* half-life of therapeutic molecules with naturally short half-lives allows for a more acceptable and manageable dosing regimen without sacrificing effectiveness. As used herein, a “half-life extension element” is a part of the chimeric polypeptide that increases the *in vivo* half-life and improve pK, for example, by altering its size (e.g., to be above the kidney filtration cutoff), shape, hydrodynamic radius, charge, or parameters of absorption, biodistribution, metabolism, and elimination. An exemplary way to improve the pK of a polypeptide is by expression of an element in the polypeptide chain that binds to receptors that are recycled to the plasma membrane of cells rather than degraded in the lysosomes, such as the FcRn receptor on endothelial cells and transferrin receptor. Three types of proteins, e.g., human IgGs, HSA (or fragments), and transferrin, persist for much longer in human serum than would be predicted just by their size, which is a function of their ability to bind to receptors that are recycled rather than degraded in the lysosome. These proteins, or fragments of them that retain the FcRn binding are routinely linked to other polypeptides to extend their serum half-life. In one embodiment, the half-life extension element is a human serum albumin (HSA) binding domain. HSA (SEQ ID NO:2) may also be directly bound to the pharmaceutical compositions or bound via a short linker. Fragments of HSA may also be used. HSA and fragments thereof can function as both a blocking moiety and a half-life extension element. Human IgGs can also carry out a similar function.

[255] The serum half-life extension element can also be antigen-binding polypeptide that binds to a protein with a long serum half-life such as serum albumin, transferrin and the like. Examples of such polypeptides include antibodies and fragments thereof including, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody a single chain variable fragment (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain of camelid-type nanobody (VHH), a sdAb and the like. Other suitable antigen-binding domain include non-immunoglobulin proteins that mimic antibody binding and/or structure such as, anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPin, fynomers, kunitz domain peptides, monobodies, and binding domains based on other engineered scaffolds such as SpA, GroEL, fibronectin, lipocalin and CTLA4 scaffolds. Further examples of antigen-binding polypeptides include a ligand for a desired receptor, a ligand-binding portion of a receptor, a lectin, and peptides that binds to or associates with one or more target antigens.

[256] Some preferred serum half-life extension elements are polypeptides that comprise complementarity determining regions (CDRs), and optionally non-CDR loops. Advantageously,

such serum half-life extension elements can extend the serum half-life of the cytokine, and also function as inhibitors of the cytokine (e.g., via steric blocking, non-covalent interaction or combination thereof) and/or as targeting domains. In some instances, the serum half-life extension elements are domains derived from an immunoglobulin molecule (Ig molecule) or engineered protein scaffolds that mimic antibody structure and/or binding activity. The Ig may be of any class or subclass (IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM etc.). A polypeptide chain of an Ig molecule folds into a series of parallel beta strands linked by loops. In the variable region, three of the loops constitute the "complementarity determining regions" (CDRs) which determine the antigen binding specificity of the molecule. An IgG molecule comprises at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding fragment thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs) which are hypervariable in sequence and/or involved in antigen recognition and/or usually form structurally defined loops, interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In some embodiments of this disclosure, at least some or all of the amino acid sequences of FR1, FR2, FR3, and FR4 are part of the "non-CDR loop" of the binding moieties described herein. A variable domain of an immunoglobulin molecule has several beta strands that are arranged in two sheets. The variable domains of both light and heavy immunoglobulin chains contain three hypervariable loops, or complementarity-determining regions (CDRs). The three CDRs of a V domain (CDR1, CDR2, CDR3) cluster at one end of the beta barrel. The CDRs are the loops that connect beta strands B-C, C'-C'', and F-G of the immunoglobulin fold, whereas the bottom loops that connect beta strands AB, CC', C''-D and E-F of the immunoglobulin fold, and the top loop that connects the D-E strands of the immunoglobulin fold are the non-CDR loops. In some embodiments of this disclosure, at least some amino acid residues of a constant domain, CH1, CH2, or CH3, are part of the "non-CDR loop" of the binding moieties described herein. Non-CDR loops comprise, in some embodiments, one or more of AB, CD, EF, and DE loops of a C1-set domain of an Ig or an Ig-like molecule; AB, CC', EF, FG, BC, and EC' loops of a C2-set domain of an Ig or an Ig-like molecule; DE, BD, GF, A(A1A2)B, and EF loops of I(Intermediate)-set domain of an Ig or Ig-like molecule.

[257] Within the variable domain, the CDRs are believed to be responsible for antigen recognition and binding, while the FR residues are considered a scaffold for the CDRs. However, in certain cases, some of the FR residues play an important role in antigen recognition and binding. Framework region residues that affect Ag binding are divided into two categories. The first are FR residues that contact the antigen, thus are part of the binding-site, and some of these residues are close in sequence to the CDRs. Other residues are those that are far from the CDRs in sequence but are in close proximity to it in the 3-D structure of the molecule, *e.g.*, a loop in heavy chain.

[258] The binding moieties are any kinds of polypeptides. For example, in certain instances the binding moieties are natural peptides, synthetic peptides, or fibronectin scaffolds, or engineered bulk serum proteins. The bulk serum protein comprises, for example, albumin, fibrinogen, or a globulin. In some embodiments, the binding moieties are engineered scaffolds. Engineered scaffolds comprise, for example, a sdAb, a scFv, a Fab, a VHH, a fibronectin type III domain, immunoglobulin-like scaffold (as suggested in Halaby et al., 1999. *Prot Eng* 12(7):563-571), DARPin, cystine knot peptide, lipocalin, three-helix bundle scaffold, protein G-related albumin-binding module, or a DNA or RNA aptamer scaffold.

[259] In some cases, the serum half-life extending element comprises a binding site for a bulk serum protein. In some embodiments, the CDRs provide the binding site for the bulk serum protein. The bulk serum protein is, in some examples, a globulin, albumin, transferrin, IgG1, IgG2, IgG4, IgG3, IgA monomer, Factor XIII, Fibrinogen, IgE, or pentameric IgM. In some embodiments, the CDR form a binding site for an immunoglobulin light chain, such as an Igκ free light chain or an Igλ free light chain.

[260] The serum half-life extension element can be any type of binding domain, including but not limited to, domains from a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody. In some embodiments, the binding moiety is a single chain variable fragment (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain (VHH) of camelid derived nanobody. In other embodiments, the binding moieties are non-Ig binding domains, *i.e.*, antibody mimetic, such as anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPins, fynomers, kunitz domain peptides, and monobodies.

[261] In other embodiments, the serum half-life extension element can be a water-soluble polymer or a peptide that is conjugated to a water-soluble polymer, such as PEG. "PEG," "polyethylene glycol" and "poly(ethylene glycol)" as used herein, are interchangeable and encompass any nonpeptidic water-soluble poly(ethylene oxide). The term "PEG" also means a polymer that contains a majority, that is to say, greater than 50%, of —OCH₂CH₂— repeating subunits. With respect to specific forms, the PEG can take any number of a variety of molecular weights, as well as structures or geometries such as "branched," "linear," "forked,"

“multifunctional,” and the like, to be described in greater detail below. The PEG is not limited to a particular structure and can be linear (e.g., an end capped, e.g., alkoxy PEG or a bifunctional PEG), branched or multi-armed (e.g., forked PEG or PEG attached to a polyol core), a dendritic (or star) architecture, each with or without one or more degradable linkages. Moreover, the internal structure of the PEG can be organized in any number of different repeat patterns and can be selected from the group consisting of homopolymer, alternating copolymer, random copolymer, block copolymer, alternating tripolymer, random tripolymer, and block tripolymer. PEGs can be conjugated to polypeptide and peptides through any suitable method. Typically a reactive PEG derivative, such as N-hydroxysuccinamidyl ester PEG, is reacted with a peptide or polypeptide that includes amino acids with a side chain that contains an amine, sulfhydryl, carboxylic acid or hydroxyl functional group, such as cysteine, lysine, asparagine, glutamine, threonine, tyrosine, serine, aspartic acid, and glutamic acid.

c. Targeting and Retention Domains

[262] For certain applications, it may be desirable to maximize the amount of time the construct is present in its desired location in the body. This can be achieved by including one further domain in the chimeric polypeptide (fusion protein) to influence its movements within the body. For example, the chimeric nucleic acids can encode a domain that directs the polypeptide to a location in the body, e.g., tumor cells or a site of inflammation; this domain is termed a “targeting domain” and/or encode a domain that retains the polypeptide in a location in the body, e.g., tumor cells or a site of inflammation; this domain is termed a “retention domain”. In some embodiments a domain can function as both a targeting and a retention domain. In some embodiments, the targeting domain and/or retention domain are specific to a protease-rich environment. In some embodiments, the encoded targeting domain and/or retention domain are specific for regulatory T cells (Tregs), for example targeting the CCR4 or CD39 receptors. Other suitable targeting and/or retention domains comprise those that have a cognate ligand that is overexpressed in inflamed tissues, e.g., the IL-1 receptor, or the IL-6 receptor. In other embodiments, the suitable targeting and/or retention domains comprise those who have a cognate ligand that is overexpressed in tumor tissue, e.g., Epcam, CEA or mesothelin. In some embodiments, the targeting domain is linked to the interleukin via a linker which is cleaved at the site of action (e.g. by inflammation or cancer specific proteases) releasing the interleukin full activity at the desired site. In some embodiments, the targeting and/or retention domain is linked to the interleukin via a linker which is not cleaved at the site of action (e.g. by inflammation or cancer specific proteases), causing the cytokine to remain at the desired site.

[263] Antigens of choice, in some cases, are expressed on the surface of a diseased cell or tissue, for example a tumor or a cancer cell. Antigens useful for tumor targeting and retention include but are not limited to EpCAM, EGFR, HER-2, HER-3, c-Met, FoIR, and CEA. Pharmaceutical

compositions disclosed herein, also include proteins comprising two targeting and/or retention domains that bind to two different target antigens known to be expressed on a diseased cell or tissue. Exemplary pairs of antigen binding domains include but are not limited to EGFR/CEA, EpCAM/CEA, and HER-2/HER-3.

[264] Suitable targeting and/or retention domains include antigen-binding domains, such as antibodies and fragments thereof including, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody a single chain variable fragment (scFv), single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain of camelid-type nanobody (VHH), a sdAb and the like. Other suitable antigen-binding domain include non-immunoglobulin proteins that mimic antibody binding and/or structure such as, anticalins, affilins, affibody molecules, affimers, affitins, alphabodies, avimers, DARPins, fynomers, kunitz domain peptides, monobodies, and binding domains based on other engineered scaffolds such as SpA, GroEL, fibronectin, lipocalin and CTLA4 scaffolds. Further examples of antigen-binding polypeptides include a ligand for a desired receptor, a ligand-binding portion of a receptor, a lectin, and peptides that binds to or associates with one or more target antigens.

[265] In some embodiments, the targeting and/or retention domains specifically bind to a cell surface molecule. In some embodiments, the targeting and/or retention domains specifically bind to a tumor antigen. In some embodiments, the targeting polypeptides specifically and independently bind to a tumor antigen selected from at least one of EpCAM, EGFR, HER-2, HER-3, cMet, CEA, and FoIR. In some embodiments, the targeting polypeptides specifically and independently bind to two different antigens, wherein at least one of the antigens is a tumor antigen selected from EpCAM, EGFR, HER-2, HER-3, cMet, CEA, and FoIR.

[266] The targeting and/or retention antigen can be a tumor antigen expressed on a tumor cell. Tumor antigens are well known in the art and include, for example, EpCAM, EGFR, HER-2, HER-3, c-Met, FoIR, PSMA, CD38, BCMA, and CEA. 5T4, AFP, B7-H3, Cadherin-6, CAIX, CD117, CD123, CD138, CD166, CD19, CD20, CD205, CD22, CD30, CD33, CD352, CD37, CD44, CD52, CD56, CD70, CD71, CD74, CD79b, DLL3, EphA2, FAP, FGFR2, FGFR3, GPC3, gpA33, FLT-3, gpNMB, HPV-16 E6, HPV-16 E7, ITGA2, ITGA3, SLC39A6, MAGE, mesothelin, Muc1, Muc16, NaPi2b, Nectin-4, P-cadherin, NY-ESO-1, PRLR, PSCA, PTK7, ROR1, SLC44A4, SLTRK5, SLTRK6, STEAP1, TIM1, Trop2, WT1.

[267] The targeting and/or retention antigen can be an immune checkpoint protein. Examples of immune checkpoint proteins include but are not limited to CD27, CD137, 2B4, TIGIT, CD155, ICOS, HVEM, CD40L, LIGHT, TIM-1, OX40, DNAM-1, PD-L1, PD1, PD-L2, CTLA-4, CD8, CD40, CEACAM1, CD48, CD70, A2AR, CD39, CD73, B7-H3, B7-H4, BTLA, IDO1, IDO2, TDO, KIR, LAG-3, TIM-3, or VISTA.

[268] The targeting and/or retention antigen can be a cell surface molecule such as a protein, lipid or polysaccharide. In some embodiments, a targeting and/or retention antigen is on a tumor cell, virally infected cell, bacterially infected cell, damaged red blood cell, arterial plaque cell, inflamed or fibrotic tissue cell. The targeting and/or retention antigen can comprise an immune response modulator. Examples of immune response modulator include but are not limited to granulocyte-macrophage colony stimulating factor (GM-CSF), macrophage colony stimulating factor (M-CSF), granulocyte colony stimulating factor (G-CSF), interleukin 2 (IL-2), interleukin 3 (IL-3), interleukin 12 (IL-12), interleukin 15 (IL-15), B7-1 (CD80), B7-2 (CD86), GITRL, CD3, or GITR.

[269] The targeting and/or retention antigen can be a cytokine receptor. Examples, of cytokine receptors include but are not limited to Type I cytokine receptors, such as GM-CSF receptor, G-CSF receptor, Type I IL receptors, Epo receptor, LIF receptor, CNTF receptor, TPO receptor; Type II Cytokine receptors, such as IFN-alpha receptor (IFNAR1, IFNAR2), IFN-beta receptor, IFN-gamma receptor (IFNGR1, IFNGR2), Type II IL receptors; chemokine receptors, such as CC chemokine receptors, CXC chemokine receptors, CX3C chemokine receptors, XC chemokine receptors; tumor necrosis receptor superfamily receptors, such as TNFRSF5/CD40, TNFRSF8/CD30, TNFRSF7/CD27, TNFRSF1A/TNFR1/CD120a, TNFRSF1B / TNFR2 / CD120b; TGF-beta receptors, such as TGF-beta receptor 1, TGF-beta receptor 2; Ig super family receptors, such as IL-1 receptors, CSF-1R, PDGFR (PDGFRA, PDGFRB), SCFR.

d. Linkers

[270] As stated above, the pharmaceutical compositions comprise one or more linker sequences. A linker sequence serves to provide flexibility between polypeptides, such that, for example, the blocking moiety is capable of inhibiting the activity of the cytokine polypeptide. The linker sequence can be located between any or all of the cytokine polypeptide, the serum half-life extension element, and/or the blocking moiety. As described herein, at least one of the linkers is protease cleavable, and contains a (one or more) cleavage site for a (one or more) desired protease. Preferably, the desired protease is enriched or selectively expressed at the desired site of cytokine activity (e.g., the tumor microenvironment). Thus, the fusion protein is preferentially or selectively cleaved at the site of desired cytokine activity.

[271] The orientation of the components of the pharmaceutical composition, are largely a matter of design choice and it is recognized that multiple orientations are possible, and all are intended to be encompassed by this disclosure. For example, a blocking moiety can be located C-terminally or N-terminally to a cytokine polypeptide.

[272] Provided herein are pharmaceutical compositions comprising polypeptide sequences. As with all peptides, polypeptides, and proteins, including fragments thereof, it is understood that additional modifications in the amino acid sequence of the chimeric polypeptides (amino acid sequence variants) can occur that do not alter the nature or function of the peptides, polypeptides,

or proteins. Such modifications include conservative amino acid substitutions and are discussed in greater detail below.

[273] The compositions provided herein have a desired function. The compositions are comprised of at least a cytokine polypeptide, such as IL-2, IL-7, IL-12, IL-15, IL-18, IL-21, or IFN γ , or a chemokine, such as CXCL10, CCL19, CCL20, CCL21, a blocking moiety, e.g. a steric blocking polypeptide, and an optional serum half-life extension element, and an optional targeting polypeptide, with one or more linkers connecting each polypeptide in the composition. The first polypeptide, e.g., an IL-2 mutein, is provided to be an active agent. The blocking moiety is provided to block the activity of the interleukin. The linker polypeptide, e.g., a protease cleavable polypeptide, is provided to be cleaved by a protease that is specifically expressed at the intended target of the active agent. Optionally, the blocking moiety blocks the activity of the first polypeptide by binding the interleukin polypeptide. In some embodiments, the blocking moiety, e.g. a steric blocking peptide, is linked to the interleukin via a protease-cleavable linker which is cleaved at the site of action (e.g. by inflammation specific proteases) releasing the cytokine full activity at the desired site.

[274] In some embodiments, the linker comprises glycine-glycine, a sortase-recognition motif, or a sortase-recognition motif and a peptide sequence (Gly₄Ser)_n (SEQ ID NO: 238) or (Gly₃Ser)_n (SEQ ID NO: 239), wherein n is 1, 2, 3, 4 or 5. In one embodiment, the sortase-recognition motif comprises a peptide sequence LPXTG, where X is any amino acid (SEQ ID NO: 237). In one embodiment, the covalent linkage is between a reactive lysine residue attached to the C-terminal of the cytokine polypeptide and a reactive aspartic acid attached to the N-terminal of the blocking or other moiety. In one embodiment, the covalent linkage is between a reactive aspartic acid residue attached to the N-terminal of the cytokine polypeptide and a reactive lysine residue attached to the C-terminal of the blocking or other moiety.

e. Cleavage and Inducibility

[275] As described herein, the activity of the cytokine polypeptide the context of the fusion protein is attenuated, and protease cleavage at the desired site of activity, such as in a tumor microenvironment, releases a form of the cytokine from the fusion protein that is much more active as a cytokine receptor agonist than the fusion protein. For example, the cytokine-receptor activating (agonist) activity of the fusion polypeptide can be at least about 10X, at least about 50X, at least about 100X, at least about 250X, at least about 500X, or at least about 1000x less than the cytokine receptor activating activity of the cytokine polypeptide as a separate molecular entity. The cytokine polypeptide that is part of the fusion protein exists as a separate molecular entity when it contains an amino acid that is substantially identical to the cytokine polypeptide and does not substantially include additional amino acids and is not associated (by covalent or non-covalent bonds) with other molecules. If necessary, a cytokine polypeptide as a separate molecular entity

may include some additional amino acid sequences, such as a tag or short sequence to aid in expression and/or purification.

[276] In other examples, the cytokine-receptor activating (agonist) activity of the fusion polypeptide is at least about 10X, at least about 50X, at least about 100X, at least about 250X, at least about 500X, or about 1000x less than the cytokine receptor activating activity of the polypeptide that contains the cytokine polypeptide that is produced by cleavage of the protease cleavable linker in the fusion protein. In other words, the cytokine receptor activating (agonist) activity of the polypeptide that contains the cytokine polypeptide that is produced by cleavage of the protease cleavable linker in the fusion protein is at least about 10X, at least about 50X, at least about 100X, at least about 250X, at least about 500X, or at least about 1000x greater than the cytokine receptor activating activity of the fusion protein. In other examples, a recombinant polypeptide is in conjugation with a cleavable moiety wherein the cleavable moiety is cleaved with reduced catalytic efficiency by one or more proteases than a reference polypeptide sequence.

[277] In some embodiments, the cleavable moiety is resistant to proteolytic cleavage by one or more proteases. A cleavable moiety is resistant to a protease if the sequence comprises a binding site that is altered from the canonical cleavable motif sequence for the specific protease. In some embodiments, a binding site is altered compared to a reference sequence by making one or more substitutions in the protease cleavage motif sequence. For example, the protease cathepsin S cleaves the sequence GAVVRGA (SEQ ID NO: 240); a sequence is made to substitution can be made to substitute the arginine (R) with a glutamine (Q), thus changing a charged residue to a shorter, polar residue, reducing the ability of cathepsin S to bind and cleave the sequence. Such semi-conservative amino acid substitutions in protease target sequence motifs can lead to reduced binding ability, and such altered sequence motifs are therefore protease resistant. An uncleavable moiety can be made by inserting a disruptive amino acid into the protease target sequence motif, such as a proline (causes curves in the secondary structure of the peptide) or histidine (causes steric interference with other amino acid side chains). As used herein, a "protease-resistant" peptide linker is one with reduced or undetectable cleavage by one or more specified proteases. Exemplary protease-resistant peptide linkers can be tested, e.g., in vitro by incubation with a specific protease followed by analysis of the digestion products by western blot.

f. Polypeptide substitutions

[278] The polypeptides described herein can include components (e.g., the cytokine, the blocking moiety) that have the same amino acid sequence of the corresponding naturally occurring protein (e.g., IL-2, IL-15, HSA) or can have an amino acid sequence that differs from the naturally occurring protein so long as the desired function is maintained. It is understood that one way to define any known modifications and derivatives or those that might arise, of the disclosed proteins and nucleic acids that encode them is through defining the sequence variants in terms of identity to

specific known reference sequences. Specifically disclosed are polypeptides and nucleic acids which have at least, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 percent identity to the chimeric polypeptides provided herein. For example, provided are polypeptides or nucleic acids that have at least, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 percent identity to the sequence of any of the nucleic acids or polypeptides described herein. This includes polypeptides or nucleic acids that have at least, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 percent identity to the sequence of the cleavable linkers provided herein. This also includes variants of the linker or inducible polypeptides that comprise 1, 2, 3, 4, 5, or 6 variants from the cleavage domain sequences. Those of skill in the art readily understand how to determine the identity of two polypeptides or two nucleic acids. For example, the identity can be calculated after aligning the two sequences so that the identity is at its highest level.

[279] Another way of calculating identity can be performed by published algorithms. Optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman Adv. Appl. Math. 2:482 (1981), by the identity alignment algorithm of Needleman and Wunsch, J. Mol. Biol. 48:443 (1970), by the search for similarity method of Pearson and Lipman, Proc. Natl. Acad. Sci. USA 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection.

[280] The same types of identity can be obtained for nucleic acids by, for example, the algorithms disclosed in Zuker, Science 244:48-52 (1989); Jaeger et al., Proc. Natl. Acad. Sci. USA 86:7706-7710 (1989); Jaeger et al., Methods Enzymol. 183:281-306 (1989), which are herein incorporated by reference for at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that in certain instances the results of these various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have the stated identity, and be disclosed herein.

[281] Protein modifications include amino acid sequence modifications. Modifications in amino acid sequence may arise naturally as allelic variations (e.g., due to genetic polymorphism), may arise due to environmental influence (e.g., by exposure to ultraviolet light), or may be produced by human intervention (e.g., by mutagenesis of cloned DNA sequences), such as induced point, deletion, insertion and substitution mutants. These modifications can result in changes in the amino acid sequence, provide silent mutations, modify a restriction site, or provide other specific mutations. Amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional or deletional modifications. Insertions include amino and/or carboxyl terminal fusions as well as intrasequence insertions of single or multiple amino acid residues.

Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site within the protein molecule. Amino acid substitutions are typically of single residues but can occur at a number of different locations at once; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs, i.e. a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. Substitutional modifications are those in which at least one residue has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Table 2 and are referred to as conservative substitutions.

Table 2. Exemplary amino acid substitutions

Amino Acid	Exemplary Substitutions
Ala	Ser, Gly, Cys
Arg	Lys, Gln, Met, Ile
Asn	Gln, His, Glu, Asp
Asp	Glu, Asn, Gln
Cys	Ser, Met, Thr
Gln	Asn, Lys, Glu, Asp
Glu	Asp, Asn, Gln
Gly	Pro, Ala
His	Asn, Gln
Ile	Leu, Val, Met
Leu	Ile, Val, Met
Lys	Arg, Gln, Met, Ile
Met	Leu, Ile, Val
Phe	Met, Leu, Tyr, Trp, His
Ser	Thr, Met, Cys
Thr	Ser, Met, Val
Trp	Tyr, Phe

Tyr	Trp, Phe, His
Val	Ile, Leu, Met

[282] Modifications, including the specific amino acid substitutions, are made by known methods. For example, modifications are made by site specific mutagenesis of nucleotides in the DNA encoding the polypeptide, thereby producing DNA encoding the modification, and thereafter expressing the DNA in recombinant cell culture. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example M13 primer mutagenesis and PCR mutagenesis.

[283] Modifications can be selected to optimize binding. For example, affinity maturation techniques can be used to alter binding of the scFv by introducing random mutations inside the complementarity determining regions (CDRs). Such random mutations can be introduced using a variety of techniques, including radiation, chemical mutagens or error-prone PCR. Multiple rounds of mutation and selection can be performed using, for example, phage display.

[284] The disclosure also relates to nucleic acids that encode the chimeric polypeptides described herein, and to the use of such nucleic acids to produce the chimeric polypeptides and for therapeutic purposes. For example, the invention includes DNA and RNA molecules (e.g., mRNA, self-replicating RNA) that encode a chimeric polypeptide and to the therapeutic use of such DNA and RNA molecules.

g. Exemplary Compositions

[285] Exemplary fusion proteins of the invention combine the above described elements in a variety of orientations. The orientations described in this section are meant as examples and are not to be considered limiting.

[286] In some embodiments, the fusion protein comprises a cytokine, a blocking moiety and a half-life extension element. In some embodiments, the cytokine is positioned between the half-life extension element and the blocking moiety. In some embodiments, the cytokine is N-terminal to the blocking moiety and the half-life extension element. In some such embodiments, the cytokine is proximal to the blocking moiety; in some such embodiments, the cytokine is proximal to the half-life extension element. At least one protease-cleavable linker must be included in all embodiments, such that the cytokine may be active upon cleavage. In some embodiments, the cytokine is C-terminal to the blocking moiety and the half-life extension element. Additional elements may be attached to one another by a cleavable linker, a non-cleavable linker, or by direct fusion.

[287] In some embodiments, the blocking domains used are capable of extending half-life, and the cytokine is positioned between two such blocking domains. In some embodiments, the cytokine is positioned between two blocking domains, one of which is capable of extending half-

life.

[288] In some embodiments, two cytokines are included in the same construct. In some embodiments, the cytokines are connected to two blocking domains each (three in total in one molecule), with a blocking domain between the two cytokine domains. In some embodiments, one or more additional half-life extension domains may be included to optimize pharmacokinetic properties.

[289] In some embodiments, three cytokines are included in the same construct. In some embodiments, the third cytokine may function to block the other two in place of a blocking domain between the two cytokines.

Preferred half-life extension elements for use in the fusion proteins are human serum albumin (HSA), an antibody or antibody fragment (e.g., scFV, dAb) which binds serum albumin, a human or humanized IgG, or a fragment of any of the foregoing. In some preferred embodiments, the blocking moiety is human serum albumin (HSA), or an antibody or antibody fragment which binds serum albumin, an antibody which binds the cytokine and prevents activation of binding or activation of the cytokine receptor, another cytokine, or a fragment of any of the foregoing. In preferred embodiments comprising an additional targeting domain, the targeting domain is an antibody which binds a cell surface protein which is enriched on the surface of cancer cells, such as EpCAM, FOLR1, and Fibronectin.

vii. Other uses

[290] The separation moieties disclosed herein can be used for antibody-antibiotic conjugates. The separation moiety disclosed herein connects or links an antimicrobial antibiotic to an antibody specific for a bacterial strain (e.g., *Staphylococcus aureus* Ab). The antibody-antibiotic conjugate does not display antibacterial activity when the antibody is linked to the antibiotic. However, upon internalization into host cells, the separation moiety is cleaved by proteases releasing the free antibiotic. The free antibiotic kills the intracellular bacteria.

[291] The separation moieties disclosed herein can be used for applications in chemical probes use for detection and isolation of proteins. Chemical probes are designed based on small molecule interaction with proteins. The probes typically comprise a covalent binding motif in order for the probe to interact with the target protein, a detection/purification tag for visualization/purification of the protein target and a linker group. The separation moieties described herein can be incorporated to enable the target protein to be detected and isolated.

C. Methods of Treatment and Pharmaceutical Compositions

[292] Further provided are methods of treating a subject with or at risk of developing an of a disease or disorder, such as proliferative disease, a tumorous disease, an inflammatory disease, an immunological disorder, an autoimmune disease, an infectious disease, a viral disease, an allergic

reaction, a parasitic reaction, or graft-versus-host disease. The methods administering to a subject in need thereof an effective amount of a fusion protein as disclosed herein that is typically administered as a pharmaceutical composition. In some embodiments, the method further comprises selecting a subject with or at risk of developing such a disease or disorder. The pharmaceutical composition preferably comprises a blocked cytokine, fragment or mutein thereof that is activated at a site of inflammation. In one embodiment, the chimeric polypeptide comprises a cytokine polypeptide, fragment or mutein thereof and a serum half-life extension element. In another embodiment, the chimeric polypeptide comprises a cytokine polypeptide, fragment or mutein thereof and a blocking moiety, e.g. a steric blocking polypeptide, wherein the steric blocking polypeptide is capable of sterically blocking the activity of the cytokine polypeptide, fragment or mutein thereof. In another embodiment, the chimeric polypeptide comprises a cytokine polypeptide, fragment or mutein thereof, a blocking moiety, and a serum half-life extension element.

[293] Inflammation is part of the complex biological response of body tissues to harmful stimuli, such as pathogens, damaged cells, or irritants, and is a protective response involving immune cells, blood vessels, and molecular mediators. The function of inflammation is to eliminate the initial cause of cell injury, clear out necrotic cells and tissues damaged from the original insult and the inflammatory process, and to initiate tissue repair. Inflammation can occur from infection, as a symptom or a disease, e.g., cancer, atherosclerosis, allergies, myopathies, HIV, obesity, or an autoimmune disease. An autoimmune disease is a chronic condition arising from an abnormal immune response to a self-antigen. Autoimmune diseases that may be treated with the polypeptides disclosed herein include but are not limited to lupus, celiac disease, diabetes mellitus type 1, Graves' disease, inflammatory bowel disease, multiple sclerosis, psoriasis, rheumatoid arthritis, and systemic lupus erythematosus.

[294] The pharmaceutical composition can comprise one or more protease-cleavable linker sequences. The linker sequence serves to provide flexibility between polypeptides, such that each polypeptide is capable of inhibiting the activity of the first polypeptide. The linker sequence can be located between any or all of the cytokine polypeptide, fragment or mutein thereof, the blocking moiety, and serum half-life extension element. Optionally, the composition comprises two, three, four, or five linker sequences. The linker sequence, two, three, or four linker sequences can be the same or different linker sequences. In one embodiment, the linker sequence comprises GGGGS (SEQ ID NO: 232), GSGSGS (SEQ ID NO: 233), or G(SGGG)₂SGGT (SEQ ID NO: 234). In another embodiment, the linker comprises a protease-cleavable sequence selected from group consisting of HSKLQ (SEQ ID NO: 25), GPLGVRG (SEQ ID NO: 221), IPVSLRSG (SEQ ID NO: 222), VPLSLYSG (SEQ ID NO: 223), or SGESPAYTA (SEQ ID NO: 224). In some embodiments, the linker is cleaved by a protease selected from the group consisting of a kallikrein,

thrombin, chymase, carboxypeptidase A, cathepsin G, an elastase, PR-3, granzyme M, a calpain, a matrix metalloproteinase (MMP), a plasminogen activator, a cathepsin, a caspase, a tryptase, or a tumor cell surface protease.

[295] Further provided are methods of treating a subject with or at risk of developing cancer. The methods comprise administering to the subject in need thereof an effective amount of a chimeric polypeptide (a fusion protein) as disclosed herein that is typically administered as a pharmaceutical composition. In some embodiments, the method further comprises selecting a subject with or at risk of developing cancer. The pharmaceutical composition preferably comprises a blocked cytokine, fragment or mutein thereof that is activated at a tumor site. Preferably, the tumor is a solid tumor. The cancer may be a colon cancer, a lung cancer, a melanoma, a sarcoma, a renal cell carcinoma, a breast cancer,

[296] The method can further involve the administration of one or more additional agents to treat cancer, such as chemotherapeutic agents (e.g., Adriamycin, Cerubidine, Bleomycin, Alkeran, Velban, Oncovin, Fluorouracil, Thiotepa, Methotrexate, Bisantrone, Noantrone, Thiguanine, Cytaribine, Procarabazine), immuno-oncology agents (e.g., anti-PD-L1, anti-CTLA4, anti-PD-1, anti-CD47, anti-GD2), cellular therapies (e.g., CAR-T, T-cell therapy), oncolytic viruses and the like.

[297] Provided herein are pharmaceutical formulations or compositions containing the chimeric polypeptides and a pharmaceutically acceptable carrier. The herein provided compositions are suitable for administration *in vitro* or *in vivo*. By pharmaceutically acceptable carrier is meant a material that is not biologically or otherwise undesirable, i.e., the material is administered to a subject without causing undesirable biological effects or interacting in a deleterious manner with the other components of the pharmaceutical formulation or composition in which it is contained. The carrier is selected to minimize degradation of the active ingredient and to minimize adverse side effects in the subject.

[298] Suitable carriers and their formulations are described in *Remington: The Science and Practice of Pharmacy*, 21st Edition, David B. Troy, ed., Lippicott Williams & Wilkins (2005). Typically, an appropriate amount of a pharmaceutically acceptable salt is used in the formulation to render the formulation isotonic, although the formulate can be hypertonic or hypotonic if desired. Examples of the pharmaceutically acceptable carriers include, but are not limited to, sterile water, saline, buffered solutions like Ringer's solution, and dextrose solution. The pH of the solution is generally about 5 to about 8 or from about 7 to 7.5. Other carriers include sustained release preparations such as semipermeable matrices of solid hydrophobic polymers containing the immunogenic polypeptides. Matrices are in the form of shaped articles, e.g., films, liposomes, or microparticles. Certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of composition being administered. Carriers are those suitable for

administration of the chimeric polypeptides or nucleic acid sequences encoding the chimeric polypeptides to humans or other subjects.

[299] The pharmaceutical formulations or compositions are administered in a number of ways depending on whether local or systemic treatment is desired and, on the area to be treated. The compositions are administered via any of several routes of administration, including topically, orally, parenterally, intravenously, intra-articularly, intraperitoneally, intramuscularly, subcutaneously, intracavity, transdermally, intrahepatically, intracranially, nebulization/inhalation, or by installation via bronchoscopy. In some embodiments, the compositions are administered locally (non-systemically), including intratumorally, intra-articularly, intrathecally, etc.

[300] Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives are optionally present such as, for example, antimicrobials, antioxidants, chelating agents, and inert gases and the like.

[301] Formulations for topical administration include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids, and powders. Conventional pharmaceutical carriers, aqueous, powder, or oily bases, thickeners and the like are optionally necessary or desirable.

[302] Compositions for oral administration include powders or granules, suspension or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders are optionally desirable.

[303] Optionally, the chimeric polypeptides or nucleic acid sequences encoding the chimeric polypeptides are administered by a vector. There are a number of compositions and methods which can be used to deliver the nucleic acid molecules and/or polypeptides to cells, either *in vitro* or *in vivo* via, for example, expression vectors. These methods and compositions can largely be broken down into two classes: viral based delivery systems and non-viral based delivery systems. Such methods are well known in the art and readily adaptable for use with the compositions and methods described herein. Such compositions and methods can be used to transfect or transduce cells *in vitro* or *in vivo*, for example, to produce cell lines that express and preferably secrete the encoded chimeric polypeptide or to therapeutically deliver nucleic acids to a subject. The components of the chimeric nucleic acids disclosed herein typically are operably linked in frame to encode a fusion protein.

[304] As used herein, plasmid or viral vectors are agents that transport the disclosed nucleic

acids into the cell without degradation and include a promoter yielding expression of the nucleic acid molecule and/or polypeptide in the cells into which it is delivered. Viral vectors are, for example, Adenovirus, Adeno-associated virus, herpes virus, Vaccinia virus, Polio virus, Sindbis, and other RNA viruses, including these viruses with the HIV backbone. Also preferred are any viral families which share the properties of these viruses which make them suitable for use as vectors. Retroviral vectors, in general are described by Coffin et al., *Retroviruses*, Cold Spring Harbor Laboratory Press (1997), which is incorporated by reference herein for the vectors and methods of making them. The construction of replication-defective adenoviruses has been described (Berkner et al., *J. Virol.* 61:1213-20 (1987); Massie et al., *Mol. Cell. Biol.* 6:2872-83 (1986); Haj-Ahmad et al., *J. Virol.* 57:267-74 (1986); Davidson et al., *J. Virol.* 61:1226-39 (1987); Zhang et al., *BioTechniques* 15:868-72 (1993)). The benefit and the use of these viruses as vectors is that they are limited in the extent to which they can spread to other cell types, since they can replicate within an initial infected cell, but are unable to form new infectious viral particles. Recombinant adenoviruses have been shown to achieve high efficiency after direct, in vivo delivery to airway epithelium, hepatocytes, vascular endothelium, CNS parenchyma, and a number of other tissue sites. Other useful systems include, for example, replicating and host-restricted non-replicating vaccinia virus vectors.

[305] The provided polypeptides and/or nucleic acid molecules can be delivered via virus like particles. Virus like particles (VLPs) consist of viral protein(s) derived from the structural proteins of a virus. Methods for making and using virus like particles are described in, for example, Garcea and Gissmann, *Current Opinion in Biotechnology* 15:513-7 (2004).

[306] The provided polypeptides can be delivered by subviral dense bodies (DBs). DBs transport proteins into target cells by membrane fusion. Methods for making and using DBs are described in, for example, Pepperl-Klindworth et al., *Gene Therapy* 10:278-84 (2003).

[307] The provided polypeptides can be delivered by tegument aggregates. Methods for making and using tegument aggregates are described in International Publication No. WO 2006/110728.

[308] Non-viral based delivery methods can include expression vectors comprising nucleic acid molecules and nucleic acid sequences encoding polypeptides, wherein the nucleic acids are operably linked to an expression control sequence. Suitable vector backbones include, for example, those routinely used in the art such as plasmids, artificial chromosomes, BACs, YACs, or PACs. Numerous vectors and expression systems are commercially available from such corporations as Novagen (Madison, Wis.), Clontech (Pal Alto, Calif.), Stratagene (La Jolla, Calif.), and Invitrogen/Life Technologies (Carlsbad, Calif.). Vectors typically contain one or more regulatory regions. Regulatory regions include, without limitation, promoter sequences, enhancer sequences, response elements, protein recognition sites, inducible elements, protein binding sequences, 5' and 3' untranslated regions (UTRs), transcriptional start sites, termination sequences,

polyadenylation sequences, and introns. Such vectors can also be used to make the chimeric polypeptides by expression in a suitable host cell, such as CHO cells.

[309] Preferred promoters controlling transcription from vectors in mammalian host cells may be obtained from various sources, for example, the genomes of viruses such as polyoma, Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis B virus, and most preferably cytomegalovirus (CMV), or from heterologous mammalian promoters, e.g. β -actin promoter or EF1 α promoter, or from hybrid or chimeric promoters (e.g., CMV promoter fused to the β -actin promoter). Of course, promoters from the host cell or related species are also useful herein.

[310] Enhancer generally refers to a sequence of DNA that functions at no fixed distance from the transcription start site and can be either 5' or 3' to the transcription unit. Furthermore, enhancers can be within an intron as well as within the coding sequence itself. They are usually between 10 and 300 base pairs (bp) in length, and they function in cis. Enhancers usually function to increase transcription from nearby promoters. Enhancers can also contain response elements that mediate the regulation of transcription. While many enhancer sequences are known from mammalian genes (globin, elastase, albumin, fetoprotein, and insulin), typically one will use an enhancer from a eukaryotic cell virus for general expression. Preferred examples are the SV40 enhancer on the late side of the replication origin, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

[311] The promoter and/or the enhancer can be inducible (e.g. chemically or physically regulated). A chemically regulated promoter and/or enhancer can, for example, be regulated by the presence of alcohol, tetracycline, a steroid, or a metal. A physically regulated promoter and/or enhancer can, for example, be regulated by environmental factors, such as temperature and light. Optionally, the promoter and/or enhancer region can act as a constitutive promoter and/or enhancer to maximize the expression of the region of the transcription unit to be transcribed. In certain vectors, the promoter and/or enhancer region can be active in a cell type specific manner.

Optionally, in certain vectors, the promoter and/or enhancer region can be active in all eukaryotic cells, independent of cell type. Preferred promoters of this type are the CMV promoter, the SV40 promoter, the β -actin promoter, the EF1 α promoter, and the retroviral long terminal repeat (LTR).

[312] The vectors also can include, for example, origins of replication and/or markers. A marker gene can confer a selectable phenotype, e.g., antibiotic resistance, on a cell. The marker product is used to determine if the vector has been delivered to the cell and once delivered is being expressed. Examples of selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase, neomycin, neomycin analog G418, hygromycin, puromycin, and blasticidin. When such selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell can survive if placed under selective pressure. Examples of other markers include, for example, the *E. coli* lacZ gene, green fluorescent protein (GFP), and

luciferase. In addition, an expression vector can include a tag sequence designed to facilitate manipulation or detection (e.g., purification or localization) of the expressed polypeptide. Tag sequences, such as GFP, glutathione S-transferase (GST), polyhistidine, c-myc, hemagglutinin, or FLAG™ tag (Kodak; New Haven, Conn.) sequences typically are expressed as a fusion with the encoded polypeptide. Such tags can be inserted anywhere within the polypeptide including at either the carboxyl or amino terminus.

[313] As used herein, the terms peptide, polypeptide, or protein are used broadly to mean two or more amino acids linked by a peptide bond. Protein, peptide, and polypeptide are also used herein interchangeably to refer to amino acid sequences. It should be recognized that the term polypeptide is not used herein to suggest a particular size or number of amino acids comprising the molecule and that a peptide of the invention can contain up to several amino acid residues or more. As used throughout, subject can be a vertebrate, more specifically a mammal (e.g. a human, horse, cat, dog, cow, pig, sheep, goat, mouse, rabbit, rat, and guinea pig), birds, reptiles, amphibians, fish, and any other animal. The term does not denote a particular age or sex. Thus, adult and newborn subjects, whether male or female, are intended to be covered. As used herein, patient or subject may be used interchangeably and can refer to a subject with a disease or disorder (e.g. cancer). The term patient or subject includes human and veterinary subjects.

[314] A subject at risk of developing a disease or disorder can be genetically predisposed to the disease or disorder, e.g., have a family history or have a mutation in a gene that causes the disease or disorder, or show early signs or symptoms of the disease or disorder. A subject currently with a disease or disorder has one or more than one symptom of the disease or disorder and may have been diagnosed with the disease or disorder.

[315] The methods and agents as described herein are useful for both prophylactic and therapeutic treatment. For prophylactic use, a therapeutically effective amount of the chimeric polypeptides or chimeric nucleic acid sequences encoding the chimeric polypeptides described herein are administered to a subject prior to onset (e.g., before obvious signs of cancer or inflammation) or during early onset (e.g., upon initial signs and symptoms of cancer or inflammation). Prophylactic administration can occur for several days to years prior to the manifestation of symptoms of cancer or inflammation. Prophylactic administration can be used, for example, in the preventative treatment of subjects diagnosed with a genetic predisposition to cancer. Therapeutic treatment involves administering to a subject a therapeutically effective amount of the chimeric polypeptides or nucleic acid sequences encoding the chimeric polypeptides described herein after diagnosis or development of cancer or inflammation (e.g., an autoimmune disease). Prophylactic use may also apply when a patient is undergoing a treatment, e.g., a chemotherapy, in which inflammation is expected.

[316] According to the methods taught herein, the subject is administered an effective amount of

the agent (e.g., a chimeric polypeptide). The terms effective amount and effective dosage are used interchangeably. The term effective amount is defined as any amount necessary to produce a desired physiologic response. Effective amounts and schedules for administering the agent may be determined empirically, and making such determinations is within the skill in the art. The dosage ranges for administration are those large enough to produce the desired effect in which one or more symptoms of the disease or disorder are affected (e.g., reduced or delayed). The dosage should not be so large as to cause substantial adverse side effects, such as unwanted cross-reactions, anaphylactic reactions, and the like. Generally, the dosage will vary with the age, condition, sex, type of disease, the extent of the disease or disorder, route of administration, or whether other drugs are included in the regimen, and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any contraindications. Dosages can vary and can be administered in one or more dose administrations daily, for one or several days. Guidance can be found in the literature for appropriate dosages for given classes of pharmaceutical products.

[317] As used herein the terms treatment, treat, or treating refers to a method of reducing the effects of a disease or condition or symptom of the disease or condition. Thus, in the disclosed method, treatment can refer to a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% reduction in the severity of an established disease or condition or symptom of the disease or condition. For example, a method for treating a disease is considered to be a treatment if there is a 10% reduction in one or more symptoms of the disease in a subject as compared to a control. Thus, the reduction can be a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, or any percent reduction in between 10% and 100% as compared to native or control levels. It is understood that treatment does not necessarily refer to a cure or complete ablation of the disease, condition, or symptoms of the disease or condition.

[318] As used herein, the terms prevent, preventing, and prevention of a disease or disorder refers to an action, for example, administration of the chimeric polypeptide or nucleic acid sequence encoding the chimeric polypeptide, that occurs before or at about the same time a subject begins to show one or more symptoms of the disease or disorder, which inhibits or delays onset or exacerbation of one or more symptoms of the disease or disorder. As used herein, references to decreasing, reducing, or inhibiting include a change of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater as compared to a control level. Such terms can include but do not necessarily include complete elimination.

[319] IL-2 variants have been developed that are selective for IL2R $\alpha\beta\gamma$ relative to IL2R $\beta\gamma$ (Shanafelt, A. B., et al., 2000, Nat Biotechnol.18:1197-202; Cassell, D. J., et. al., 2002, Curr Pharm Des., 8:2171-83). These variants have amino acid substitutions which reduce their affinity for IL2RB. Because IL-2 has undetectable affinity for IL2RG, these variants consequently have

reduced affinity for the IL2R $\beta\gamma$ receptor complex and reduced ability to activate IL2R $\beta\gamma$ -expressing cells but retain the ability to bind IL2RA and the ability to bind and activate the IL2R $\alpha\beta\gamma$ receptor complex.

[320] One of these variants, IL2/N88R (Bay 50-4798), was clinically tested as a low- toxicity version of IL-2 as an immune system stimulator, based on the hypothesis that IL2R $\beta\gamma$ -expressing NK cells are a major contributor to toxicity. Bay 50-4798 was shown to selectively stimulate the proliferation of activated T cells relative to NK cells, and was evaluated in phase I/II clinical trials in cancer patients (Margolin, K., et. al., 2007, Clin Cancer Res., 13:3312-9) and HIV patients (Davey, R. T., et. al., 2008, J Interferon Cytokine Res., 28:89-100). These clinical trials showed that Bay 50-4798 was considerably safer and more tolerable than aldesleukin, and also showed that it increased the levels of CD4+CD25+ T cells, a cell population enriched in Treg cells. Subsequent to these trials, research in the field more fully established the identity of Treg cells and demonstrated that Treg cells selectively express IL2R $\alpha\beta\gamma$ (reviewed in Malek, T. R., et al., 2010, Immunity, 33:153-65). Based on this new research, it can now be understood that IL2R $\alpha\beta\gamma$ selective agonists should be selective for Treg cells.

[321] In addition, mutants can be made that selectively alter the affinity for the CD25 chain relative to native IL-2.

[322] IL-2 can be engineered to produce mutants that bind the IL-2R complex generally or the IL-2R α subunit specifically with an affinity that differs from that of the corresponding wild-type IL-2 or of a presently available mutant (referred to as C125S, as the cysteine residue at position 125 is replaced with a serine residue).

[323] Accordingly, the present invention features mutant interleukin-2 (IL-2*) polypeptides that include an amino acid sequence that is at least 80% identical to wild-type IL-2 (e.g., 85, 87, 90, 95, 97, 98, or 99% identical) and that bind, as compared to WT IL-2, with higher to the IL-2 trimeric receptor relative to the dimeric IL-2 receptor. Typically, the muteins will also bind an IL-2 receptor α subunit (IL-2R α) with an affinity that is greater than the affinity with which wild type IL-2 binds the IL-2R α . The amino acid sequence within mutant IL-2 polypeptides can vary from SEQ ID NO:1 (UniProtKB accession number P60568) by virtue of containing (or only containing) one or more amino acid substitutions, which may be considered conservative or non-conservative substitutions. Non-naturally occurring amino acids can also be incorporated. Alternatively, or in addition, the amino acid sequence can vary from SEQ ID NO:1 (which may be considered the “reference” sequence) by virtue of containing and addition and/or deletion of one or more amino acid residues. More specifically, the amino acid sequence can differ from that of SEQ ID NO:1 by virtue of a mutation at least one of the following positions of SEQ ID NO:1: 1, 4, 8, 9, 10, 11, 13, 15, 26, 29, 30, 31, 35, 37, 46, 48, 49, 54, 61, 64, 67, 68, 69, 71, 73, 74, 75, 76, 79, 88, 89, 90, 92, 99, 101, 103, 114, 125, 128, or 133 (or combinations thereof). As noted, as few as one of these

positions may be altered, as may two, three, four, five, six, seven, eight, nine, ten, or 11 or more (including up to all) of the positions. For example, the amino acid sequence can differ from SEQ ID NO:1 at positions 69 and 74 and further at one or more of positions 30, 35, and 128. The amino acid sequence can also differ from SEQ ID NO:2 (as disclosed in US 7569215, incorporated herein by reference) at one of the following sets of positions: (a) positions 64, 69, and 74; (b) positions 69, 74, and 101; (c) positions 69, 74, and 128; (d) positions 30, 69, 74, and 103; (e) positions 49, 69, 73, and 76; (f) positions 69, 74, 101, and 133; (g) positions 30, 69, 74, and 128; (h) positions 69, 74, 88, and 99; (i) positions 30, 69, 74, and 128; (j) positions 9, 11, 35, 69, and 74; (k) positions 1, 46, 49, 61, 69, and 79; (l) positions 48, 68, 71, 90, 103, and 114; (m) positions 4, 10, 11, 69, 74, 88, and 133; (n) positions 15, 30, 31, 35, 48, 69, 74, and 92; (o) positions 30, 68, 69, 71, 74, 75, 76, and 90; (p) positions 30, 31, 37, 69, 73, 74, 79, and 128; (q) positions 26, 29, 30, 54, 67, 69, 74, and 92; (r) positions 8, 13, 26, 30, 35, 37, 69, 74, and 92; and (s) positions 29, 31, 35, 37, 48, 69, 71, 74, 88, and 89. Aside from mutations at these positions, the amino acid sequence of the mutant IL-2 polypeptide can otherwise be identical to SEQ ID NO:1. With respect to specific substitutions, the amino acid sequence can differ from SEQ ID NO:1 by virtue of having one or more of the following mutations: A1T, S4P, K8R, K9T, T10A, Q11R, Q13R, E15K, N26D, N29S, N30S, N30D, N30T, Y31H, Y31C, K35R, T37A, T37R, M46L, K48E, K49R, K49E, K54R, E61D, K64R, E67G, E68D, V69A, N71T, N71A, N71R, A73V, Q74P, S75P, K76E, K76R, H79R, N88D, I89V, N90H, I92T, S99P, T101A, F103S, I114V, I128T, I128A, T133A, or T133N. Our nomenclature is consistent with that of the scientific literature, where the single letter code of the amino acid in the wild-type or reference sequence is followed by its position within the sequence and then by the single letter code of the amino acid with which it is replaced. Thus, A1T designates a substitution of the alanine residue a position 1 with threonine. Other mutant polypeptides within the scope of the invention include those that include a mutant of SEQ ID NO:2 having substitutions at V69 (e.g. A) and Q74 (e.g., P). For example, the amino acid sequence can include one of the following sets of mutations with respect to SEQ ID NO:2: (a) K64R, V69A, and Q74P; (b) V69A, Q74P, and T101A; (c) V69A, Q74P, and I128T; (d) N30D, V69A, Q74P, and F103S; (e) K49E, V69A, A73V, and K76E; (f) V69A, Q74P, T101A, and T133N; (g) N30S, V69A, Q74P, and I128A; (h) V69A, Q74P, N88D, and S99P; (i) N30S, V69A, Q74P, and I128T; (j) K9T, Q11R, K35R, V69A, and Q74P; (k) A1T, M46L, K49R, E61D, V69A, and H79R; (l) K48E, E68D, N71T, N90H, F103S, and I114V; (m) S4P, T10A, Q11R, V69A, Q74P, N88D, and T133A; (n) E15K, N30S, Y31H, K35R, K48E, V69A, Q74P, and I92T; (o) N30S, E68D, V69A, N71A, Q74P, S75P, K76R, and N90H; (p) N30S, Y31C, T37A, V69A, A73V, Q74P, H79R, and I128T; (q) N26D, N29S, N30S, K54R, E67G, V69A, Q74P, and I92T; (r) K8R, Q13R, N26D, N30T, K35R, T37R, V69A, Q74P, and I92T; and (s) N29S, Y31H, K35R, T37A, K48E, V69A, N71R, Q74P, N88D, and I89V. SEQ ID NO:2 is disclosed in U.S. Patent No. 7,569,215, which is

incorporated herein by reference as an exemplary IL-2 polypeptide sequence that can be used in the invention.

[324] As noted above, any of the mutant IL-2 polypeptides disclosed herein can include the sequences described; they can also be limited to the sequences described and otherwise identical to SEQ ID NO:1. Moreover, any of the mutant IL-2 polypeptides described herein can optionally include a substitution of the cysteine residue at position 125 with another residue (e.g., serine) and/or can optionally include a deletion of the alanine residue at position 1 of SEQ ID NO:1.

[325] The mutant IL-2 polypeptides disclosed herein can bind to the IL-2R α subunit with a K_d of less than about 28 nM (e.g., less than about 25 nM; less than about 5 nM; about 1 nM; less than about 500 pM; or less than about 100 pM). More specifically, a mutant IL-2 polypeptide can have an affinity equilibrium constant less than 1.0 nM (e.g., about 0.8, 0.6, 0.4, or 0.2 nM). Affinity can also be expressed as a relative rate of dissociation from an IL-2R α subunit or from an IL-2 receptor complex (e.g., a complex expressed on the surface of a cell or otherwise membrane bound). For example, the mutant IL-2 polypeptides can dissociate from, e.g., IL-2R α , at a decreased rate relative to a wild-type polypeptide or to an IL-2 based therapeutic, e.g., IL-2*. Alternatively, affinity can be characterized as the time, or average time, an IL-2* polypeptide persists on, for example, the surface of a cell expressing an IL-2R. For example, an IL-2* polypeptide can persist on the receptor for at least about 2, 5, 10, 50, 100, or 250 times (or more).

[326] Disclosed are materials, compositions, and components that can be used for, can be used in conjunction with, can be used in preparation for, or are products of the disclosed methods and compositions. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutations of these compounds may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a method is disclosed and discussed and a number of modifications that can be made to a number of molecules including the method are discussed, each and every combination and permutation of the method, and the modifications that are possible are specifically contemplated unless specifically indicated to the contrary. Likewise, any subset or combination of these is also specifically contemplated and disclosed. This concept applies to all aspects of this disclosure including, but not limited to, steps in methods using the disclosed compositions. Thus, if there are a variety of additional steps that can be performed, it is understood that each of these additional steps can be performed with any specific method steps or combination of method steps of the disclosed methods, and that each such combination or subset of combinations is specifically contemplated and should be considered disclosed.

[327] Publications cited herein and the material for which they are cited are hereby specifically incorporated by reference in their entireties.

6. INCORPORATION BY REFERENCE

All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference. However, the citation of a reference herein should not be construed as an acknowledgement that such reference is prior art to the present invention. To the extent that any of the definitions or terms provided in the references incorporated by reference differ from the terms and discussion provided herein, the present terms and definitions control.

7. EXAMPLES

[328] The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided herein.

Example 1. Detection of IL-2, IL-2 mutein, IL-2R α and IL-2R γ in Fusion Proteins by ELISA

[329] IL-2 mutein is detected with a commercially available antibody, e.g., the anti-IL-2 monoclonal (JES6-1A12) (BD Pharmingen; San Jose, Calif.). A positive control is used to show whether the monoclonal antibody recognizes the cytokine or mutein. Antibodies against IL-2R α and IL-2R γ chain are also used. Wells of a 96-well plate are coated with an antibody (2.5 μ g/ml) in PBS. Wells are blocked with 5% non-fat milk in PBS with 0.2% Tween@20 (PBS-M-Tw) and fusion proteins are added for 1-2 hours at 37° C. After washing, an anti-IL-2 biotin-labeled antibody, e.g., JES5H4 (BD Pharmingen) is added and binding is detected using Streptavidin HRP (Southern Biotechnology Associates; Birmingham, Ala.). The ELISA plate is developed by adding 50 μ l O-phenylenediamine (OPD) (Sigma-Aldrich) in 0.1M Citrate pH 4.5 and 0.04% H₂O₂, stopped by adding 50 μ l/well 2N H₂SO₄ and the absorbance was read at 490 nm.

Example 2: Protease cleavage of fusion protein by MMP9 protease

[330] One of skill in the art would be familiar with methods of setting up protein cleavage assay. 100 μ g of protein in 1xPBS pH 7.4 were cleaved with 1 μ g active MMP9 (Sigma catalog # SAE0078-50 or Enzo catalog BML-SE360) and incubated at room temperature for up to 16 hours. Digested protein is subsequently used in functional assays or stored at -80°C prior to testing. Extent of cleavage was monitored by SDS PAGE using methods well known in the art. As shown in FIGs.10, 13, 18A, 18b, and 27A full cleavage of the fusion proteins by MMP9 protease is seen.

Example 3: CTLL-2 assay

[331] CTLL2 cells (ATCC) were plated in suspension at a concentration of 500,000 cells/well in culture media with or without 40mg/ml human serum albumin (HSA) and stimulated with a dilution series of recombinant hIL2 or activatable hIL2 for 72 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved activatable hIL2 was tested. Cleaved activatable hIL2 was generated by incubation with active MMP9. Cell activity was assessed using a CellTiter-Glo® (Promega) luminescence-based cell viability assay. Results are shown in FIGs. 8A-8F, FIGs. 9A-9Z, FIG. 25C.

Example 4: Protease Cleavage of the IL-2/IL-2R α /IL-2R γ Chimeric Polypeptide Results in Increased Accessibility to Antibodies and Biologically Active IL-2 mutein

[332] The IL-2 mutein fusion proteins are biochemically characterized before and after cleavage with a protease, e.g., PSA. Immunoblot analyses will show that the fusion proteins can be cleaved by PSA and that there is an increase in intensity of the predicted low molecular weight cleavage product of approximately 20 kDa reactive with an anti-IL-2 antibody after treatment of the samples with PSA. The degree of cleavage is dependent upon the amount of PSA as well as the time of incubation. Interestingly, when the fusion protein is analyzed before and after PSA treatment by ELISA, it was found that the apparent amount of IL-2 is increased after PSA cleavage. In this experiment, there is an approximately 2 or 4-fold increase in the apparent amount of IL-2 detected using this sandwich ELISA depending on the construct, suggesting that the antibody binding is partially hindered in the intact fusion protein. Aliquots of the same samples are also analyzed after PSA treatment using the CTLL-2 cell line that requires IL-2 for growth and survival and the viability of cells can be ascertained using the colorimetric MTT assay. In this assay, the more a supernatant can be diluted, the more biologically active IL-2 it contains, and there is an increase in the amount of biologically active IL-2 after PSA cleavage. The amount of IL-2 mutein increase will suggest that after PSA cleavage there is an increase in the predicted low molecular weight cleavage fragment of approximately 20 kDa reactive with an anti-IL-2 antibody, an increase in antibody accessibility, and most importantly, an increase in the amount of biologically active IL-2 mutein.

Example 5. *In Vivo* Delivery of a Protease Activated Fusion Protein Results in Decreased Tumor Growth

[333] The chimeric polypeptide is examined to determine if it could have biological effects *in vivo*. For these experiments a system is used in which tumor cells injected intraperitoneally rapidly and preferentially attach and grow initially on the milky spots, a series of organized immune aggregates found on the omentum (Gerber et al., Am. J. Pathol. 169:1739-52 (2006)). This system

offers a convenient way to examine the effects of fusion protein treatment on tumor growth since fusion proteins can be delivered intraperitoneally multiple times and tumor growth can be analyzed by examining the dissociated omental cells. For these experiments, the Colon 38 cell line, a rapidly growing tumor cell line that expresses both MMP2 and MMP9 *in vitro*, may be used. The omental tissue normally expresses a relatively small amount of MMP2 and MMP9, but, when Colon 38 tumor is present on the omentum, MMP levels increase. Using this tumor model, the ability of IL-2 mutein fusion proteins to affect tumor growth is examined. Colon 38 cells are injected intraperitoneally, allowed to attach and grow for 1 day, and then treated daily with fusion protein intraperitoneally. At day 7, the animals are sacrificed and the omenta examined for tumor growth using flow cytometry and by a colony-forming assay.

Example 6: Determination of antigen affinity by flow cytometry

[334] Activatable interleukin proteins are tested for their binding affinities to human CD20⁺ cells and cynomolgus CD20⁺ cells.

[335] CD20⁺ cells are incubated with 100 μ L of serial dilutions of the activatable interleukin proteins and at least one protease. After washing three times with FACS buffer the cells are incubated with 0.1 mL of 10 μ g/mL mouse monoclonal anti-idiotypic antibody in the same buffer for 45 min on ice. After a second washing cycle, the cells are incubated with 0.1 mL of 15 μ g/mL FITC-conjugated goat anti-mouse IgG antibodies under the same conditions as before. As a control, cells are incubated with the anti-His IgG followed by the FITC-conjugated goat anti-mouse IgG antibodies without the activatable IL2 proteins. The cells were then washed again and resuspended in 0.2 mL of FACS buffer containing 2 μ g/mL propidium iodide (PI) in order to exclude dead cells. The fluorescence of 1×10^4 living cells is measured using a Beckman-Coulter FC500 MPL flow cytometer using the MXP software (Beckman-Coulter, Krefeld, Germany) or a Millipore Guava EasyCyte flow cytometer using the Incyte software (Merck Millipore, Schwalbach, Germany). Mean fluorescence intensities of the cell samples are calculated using CXP software (Beckman-Coulter, Krefeld, Germany) or Incyte software (Merck Millipore, Schwalbach, Germany). After subtracting the fluorescence intensity values of the cells stained with the secondary and tertiary reagents alone the values are then used for calculation of the K_D values with the equation for one-site binding (hyperbola) of the GraphPad Prism (version 6.00 for Windows, GraphPad Software, La Jolla California USA).

[336] CD20 binding and crossreactivity are assessed on the human CD20⁺ tumor cell lines. The K_D ratio of crossreactivity is calculated using the K_D values determined on the CHO cell lines expressing either recombinant human or recombinant cynomolgus antigens.

Example 7: Cytotoxicity Assay

[337] Activatable interleukin protein is evaluated *in vitro* on its mediation of immune response to CD20⁺ target cells.

[338] Fluorescence labeled CD20⁺ REC-1 cells (a Mantle cell lymphoma cell line, ATCC CRL-3004) are incubated with isolated PBMC of random donors or CB15 T-cells (standardized T-cell line) as effector cells in the presence of the activatable IL2 protein and at least one protease. After incubation for 4 h at 37°C in a humidified incubator, the release of the fluorescent dye from the target cells into the supernatant is determined in a spectro-fluorimeter. Target cells incubated without the activatable IL2 protein and target cells totally lysed by the addition of saponin at the end of the incubation serve as negative and positive controls, respectively.

[339] Based on the measured remaining living target cells, the percentage of specific cell lysis is calculated according to the following formula: $[1 - (\text{number of living targets}_{\text{(sample)}} / \text{number of living targets}_{\text{(spontaneous)}})] \times 100\%$. Sigmoidal dose response curves and EC₅₀ values are calculated by non-linear regression/4-parameter logistic fit using the GraphPad Software. The lysis values obtained for a given antibody concentration are used to calculate sigmoidal dose-response curves by 4 parameter logistic fit analysis using the Prism® GraphPad® software.

Example 8: Pharmacokinetics of Activatable Interleukin Proteins

[340] Activatable interleukin protein is evaluated for half-time elimination in animal studies.

[341] The activatable IL2 protein is administered to cynomolgus monkeys as a 0.5 mg/kg bolus injection into the saphenous vein. Another cynomolgus monkey group receives a comparable IL2 construct in size but lacking a serum half-life extension element. A third and fourth group receive an IL2 construct with serum half-life extension element and a cytokine with CD20 and serum half-life extension elements respectively, and both comparable in size to the activatable interleukin protein. Each test group consists of 5 monkeys. Serum samples are taken at indicated time points, serially diluted, and the concentration of the proteins is determined using a binding ELISA to CD20.

[342] Pharmacokinetic analysis is performed using the test article plasma concentrations. Group mean plasma data for each test article conforms to a multi-exponential profile when plotted against the time post-dosing. The data are fit by a standard two-compartment model with bolus input and first-order rate constants for distribution and elimination phases. The general equation for the best fit of the data for i.v. administration is: $c(t) = Ae^{-\alpha t} + Be^{-\beta t}$, where $c(t)$ is the plasma concentration at time t , A and B are intercepts on the Y-axis, and α and β are the apparent first-order rate constants for the distribution and elimination phases, respectively. The α -phase is the initial phase of the clearance and reflects distribution of the protein into all extracellular fluid of the animal, whereas the second or β -phase portion of the decay curve represents true plasma clearance. Methods for fitting such equations are well known in the art. For example, $A = D/V(\alpha - k_{21})/(\alpha - \beta)$,

$B=D/V(\beta-k_{21})/(\alpha-\beta)$, and α and β (for $\alpha>\beta$) are roots of the quadratic equation: $r^2+(k_{12}+k_{21}+k_{10})r+k_{21}k_{10}=0$ using estimated parameters of V =volume of distribution, k_{10} =elimination rate, k_{12} =transfer rate from compartment 1 to compartment 2 and k_{21} =transfer rate from compartment 2 to compartment 1, and D =the administered dose.

[343] Data analysis: Graphs of concentration versus time profiles are made using KaleidaGraph (KaleidaGraph™ V. 3.09 Copyright 1986-1997. Synergy Software. Reading, Pa.). Values reported as less than reportable (LTR) are not included in the PK analysis and are not represented graphically. Pharmacokinetic parameters are determined by compartmental analysis using WinNonlin software (WinNonlin® Professional V. 3.1 WinNonlin™ Copyright 1998-1999. Pharsight Corporation. Mountain View, Calif.). Pharmacokinetic parameters are computed as described in Ritschel W A and Kearns G L, 1999, IN: *Handbook of Basic Pharmacokinetics Including Clinical Applications*, 5th edition, American Pharmaceutical Assoc., Washington, D.C.

[344] It is expected that the activatable interleukin protein has improved pharmacokinetic parameters such as an increase in elimination half-time as compared to proteins lacking a serum half-life extension element.

Example 9: Xenograft Tumor Model

[345] Activatable IL2 protein is evaluated in a xenograft model.

[346] Female immune-deficient NOD/scid mice are sub-lethally irradiated (2 Gy) and subcutaneously inoculated with 4×10^6 Ramos RA1 cells into the right dorsal flank. When tumors reach 100 to 200 mm³, animals are allocated into 3 treatment groups. Groups 2 and 3 (8 animals each) are intraperitoneally injected with 1.5×10^7 activated human T-cells. Three days later, animals from Group 3 are subsequently treated with a total of 9 intravenous doses of 50 µg activatable interleukin protein. Groups 1 and 2 are only treated with vehicle. Body weight and tumor volume are determined for 30 days.

[347] It is expected that animals treated with the activatable interleukin protein have a statistically significant delay in tumor growth in comparison to the respective vehicle-treated control group.

Example 10: Mouse IFNγ WEHI cell survival assay

[348] WEHI279 cells (ATCC) were plated in suspension at a concentration of 25,000 cells/well in culture media with or without 1.5% human serum albumin (HSA) and stimulated with a dilution series of recombinant mIFNγ or inducible mIFNγ for 72 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved inducible mIFNγ was tested. Cleaved inducible mIFNγ was generated by incubation with active MMP9. Cell survival was assessed using a CellTiter-Glo (Promega) luminescence-based cell viability assay. The EC50 values for cleaved inducible mIFNγ molecules

were at least 100x more potent than un-cleaved inducible mIFN γ molecules. As shown in FIGs. 16A-16, greater inducibility was seen in assays wherein the culture medium contained human serum albumin.

Example 11: *Reserved*

Example 12: Mouse IFN γ B16 reporter cell assay

[349] B16-Blue IFN γ cells (InvivoGen) were plated at a concentration of 75,000 cells/well in culture media with or without 1.5% human serum albumin (HSA) and stimulated with a dilution series of recombinant mIFN γ or inducible mIFN γ for 24 hours at 37°C and 5% CO $_2$. Activity of uncleaved and cleaved inducible mIFN γ was tested. Cleaved inducible mIFN γ was generated by incubation with active MMP9. Supernatants were harvested, and SEAP activation was assessed by adding QUANTI-Blue Reagent (InvivoGen), incubating at 37°C for 2 hours, and measuring absorbance at 620nm. The EC50 values for cleaved inducible mIFN γ molecules were at least 100x more potent than un-cleaved inducible mIFN γ molecules. Results are shown in e.g., FIG. 19A-19B, FIGs. 22A-22B, FIGs. 23A-23B. This experiment was repeated with for IFN α conjugates using B16-Blue IFN α/β cells. The EC50 values for cleaved inducible mIFN α molecules were at least 100x more potent than un-cleaved inducible mIFN α molecules. See FIGs. 20A-20B.

Example 13. *In Vivo* Delivery of a Protease Activated Fusion Protein Results in Decreased Tumor Growth

[350] The chimeric polypeptide is examined to determine if it could have biological effects *in vivo*. For these experiments a system is used in which tumor cells injected intraperitoneally rapidly and preferentially attach and grow initially on the milky spots, a series of organized immune aggregates found on the omentum (Gerber et al., Am. J. Pathol. 169:1739-52 (2006)). This system offers a convenient way to examine the effects of fusion protein treatment on tumor growth since fusion proteins can be delivered intraperitoneally multiple times and tumor growth can be analyzed by examining the dissociated omental cells. For these experiments, the Colon 38 cell line, a rapidly growing tumor cell line that expresses both MMP2 and MMP9 *in vitro*, may be used. The omental tissue normally expresses a relatively small amount of MMP2 and MMP9, but, when Colon 38 tumor is present on the omentum, MMP levels increase. Using this tumor model, the ability of IFN fusion proteins to affect tumor growth is examined. Colon 38 cells are injected intraperitoneally, allowed to attach and grow for 1 day, and then treated daily with fusion protein intraperitoneally. At day 7, the animals are sacrificed and the omenta examined for tumor growth using flow cytometry and by a colony-forming assay.

Example 14: The chimeric polypeptide was examined to determine its biological effects *in vivo*.

[351] The MC38 cell line, a rapidly growing colon adenocarcinoma cell line that expresses MMP9 *in vitro*, was used. Using this tumor model, the ability of IFN γ fusion proteins to affect tumor growth was examined. MC38 cells were injected subcutaneously, allowed to grow for 10-14 days, and then treated with fusion protein twice weekly intraperitoneally for a total of four doses. As a comparator, wild type mIFN γ was administered at the dose levels indicated, twice daily for 2 weeks on a 5 day on/2 day off schedule (10 total doses). Tumor growth and body weight were monitored approximately twice per week for two weeks.

Example 15: Construction of an Exemplary IFN γ Protein Targeting CD20

15.1 Generation of an activatable cytokine domain

[352] An IFN γ polypeptide capable of binding to CD20 polypeptide present in a tumor or on a tumor cell is produced as follows. A nucleic acid is produced that contains nucleic acid sequences: (1) encoding an IFN γ polypeptide sequence and (2) one or more polypeptide linkers. Activatable IFN γ plasmid constructs can have optional Flag, His or other affinity tags, and are electroporated into HEK293 or other suitable human or mammalian cell lines and purified. Validation assays include T cell activation assays using T cells responsive to IFN γ stimulation in the presence of a protease.

15.2 Generation of a scFv CD20 binding domain

[353] CD20 is one of the cell surface proteins present on B-lymphocytes. CD20 antigen is found in normal and malignant pre-B and mature B lymphocytes, including those in over 90% of B-cell non-Hodgkin's lymphomas (NHL). The antigen is absent in hematopoietic stem cells, activated B lymphocytes (plasma cells) and normal tissue. As such, several antibodies mostly of murine origin have been described: 1F5, 2B8/C2B8, 2H7, and 1H4.

[354] Human or humanized anti-CD20 antibodies are therefore used to generate scFv sequences for CD20 binding domains of an activatable IFN γ protein. DNA sequences coding for human or humanized VL and VH domains are obtained, and the codons for the constructs are, optionally, optimized for expression in cells from *Homo sapiens*. The order in which the VL and VH domains appear in the scFv is varied (i.e., VL-VH, or VH-VL orientation), and three copies of the "G4S" (SEQ ID NO: 241) or "G₄S" (SEQ ID NO: 241) subunit (G₄S)₃ (SEQ ID NO: 242) connect the variable domains to create the scFv domain. Anti-CD20 scFv plasmid constructs can have optional Flag, His or other affinity tags, and are electroporated into HEK293 or other suitable human or mammalian cell lines and purified. Validation assays include binding analysis by FACS, kinetic analysis using Proteon, and staining of CD20-expressing cells.

15.3 Cloning of DNA expression constructs encoding the activatable IFN γ protein

[355] The activatable IFN γ construct with protease cleavage site domains is used to construct an activatable IFN γ protein in combination with an anti-CD20 scFv domain and a serum half-life extension element (e.g., a HSA binding peptide or VH domain). For expression of an activatable IFN γ protein in CHO cells, coding sequences of all protein domains are cloned into a mammalian expression vector system. In brief, gene sequences encoding the activatable IFN γ domain, serum half-life extension element, and CD20 binding domain along with peptide linkers L1 and L2 are separately synthesized and subcloned. The resulting constructs are then ligated together in the order of CD20 binding domain – L1 – IFN γ subunit 1 – L2 – protease cleavage domain – L3 – IFN γ subunit 2 – L4 – anti-CD20 scFv – L5 – serum half-life extension element to yield a final construct. All expression constructs are designed to contain coding sequences for an N-terminal signal peptide and a C-terminal hexahistidine (6xHis)-tag (SEQ ID NO: 243) to facilitate protein secretion and purification, respectively.

15.4 Expression of activatable IFN γ proteins in stably transfected CHO cells

[356] A CHO cell expression system (Flp-In®, Life Technologies), a derivative of CHO-K1 Chinese Hamster ovary cells (ATCC, CCL-61) (Kao and Puck, Proc. Natl. Acad. Sci. USA 1968;60(4):1275-81), is used. Adherent cells are subcultured according to standard cell culture protocols provided by Life Technologies.

[357] For adaption to growth in suspension, cells are detached from tissue culture flasks and placed in serum-free medium. Suspension-adapted cells are cryopreserved in medium with 10% DMSO.

[358] Recombinant CHO cell lines stably expressing secreted activatable IFN γ proteins are generated by transfection of suspension-adapted cells. During selection with the antibiotic Hygromycin B viable cell densities are measured twice a week, and cells are centrifuged and resuspended in fresh selection medium at a maximal density of 0.1×10^6 viable cells/mL. Cell pools stably expressing activatable IFN γ proteins are recovered after 2-3 weeks of selection at which point cells are transferred to standard culture medium in shake flasks. Expression of recombinant secreted proteins is confirmed by performing protein gel electrophoresis or flow cytometry. Stable cell pools are cryopreserved in DMSO containing medium.

[359] Activatable IFN γ proteins are produced in 10-day fed-batch cultures of stably transfected CHO cell lines by secretion into the cell culture supernatant. Cell culture supernatants are harvested after 10 days at culture viabilities of typically >75%. Samples are collected from the production cultures every other day and cell density and viability are assessed. On day of harvest, cell culture supernatants are cleared by centrifugation and vacuum filtration before further use.

[360] Protein expression titers and product integrity in cell culture supernatants are analyzed by

SDS-PAGE.**15.5 Purification of activatable IFN γ proteins**

[361] Activatable IFN γ proteins are purified from CHO cell culture supernatants in a two-step procedure. The constructs are subjected to affinity chromatography in a first step followed by preparative size exclusion chromatography (SEC) on Superdex 200 in a second step. Samples are buffer-exchanged and concentrated by ultrafiltration to a typical concentration of >1 mg/mL. Purity and homogeneity (typically >90%) of final samples are assessed by SDS PAGE under reducing and non-reducing conditions, followed by immunoblotting using an anti-HSA or anti idiotypic antibody as well as by analytical SEC, respectively. Purified proteins are stored at aliquots at -80°C until use.

Example 16: Determination of antigen affinity by flow cytometry

[362] The activatable IFN γ proteins are tested for their binding affinities to human CD20⁺ cells and cynomolgus CD20⁺ cells.

[363] CD20⁺ cells are incubated with 100 μ L of serial dilutions of the activatable IFN γ proteins and at least one protease. After washing three times with FACS buffer the cells are incubated with 0.1 mL of 10 μ g/mL mouse monoclonal anti-idiotypic antibody in the same buffer for 45 min on ice. After a second washing cycle, the cells are incubated with 0.1 mL of 15 μ g/mL FITC-conjugated goat anti-mouse IgG antibodies under the same conditions as before. As a control, cells are incubated with the anti-His IgG followed by the FITC-conjugated goat anti-mouse IgG antibodies without the activatable IFN γ proteins. The cells were then washed again and resuspended in 0.2 mL of FACS buffer containing 2 μ g/mL propidium iodide (PI) in order to exclude dead cells. The fluorescence of 1×10^4 living cells is measured using a Beckman-Coulter FC500 MPL flow cytometer using the MXP software (Beckman-Coulter, Krefeld, Germany) or a Millipore Guava EasyCyte flow cytometer using the Incyte software (Merck Millipore, Schwalbach, Germany). Mean fluorescence intensities of the cell samples are calculated using CXP software (Beckman-Coulter, Krefeld, Germany) or Incyte software (Merck Millipore, Schwalbach, Germany). After subtracting the fluorescence intensity values of the cells stained with the secondary and tertiary reagents alone the values are then used for calculation of the K_D values with the equation for one-site binding (hyperbola) of the GraphPad Prism (version 6.00 for Windows, GraphPad Software, La Jolla California USA).

[364] CD20 binding and cross-reactivity are assessed on the human CD20⁺ tumor cell lines. The K_D ratio of cross-reactivity is calculated using the K_D values determined on the CHO cell lines expressing either recombinant human or recombinant cynomolgus antigens.

Example 17: Cytotoxicity Assay

[365] The activatable IFN γ protein is evaluated *in vitro* on its mediation of immune response to CD20⁺ target cells.

[366] Fluorescence labeled CD20⁺ REC-1 cells (a Mantle cell lymphoma cell line, ATCC CRL-3004) are incubated with isolated PBMC of random donors or CB15 T-cells (standardized T-cell line) as effector cells in the presence of the activatable IFN γ protein and at least one protease. After incubation for 4 h at 37°C in a humidified incubator, the release of the fluorescent dye from the target cells into the supernatant is determined in a spectrofluorometer. Target cells incubated without the activatable IFN γ protein and target cells totally lysed by the addition of saponin at the end of the incubation serve as negative and positive controls, respectively.

[367] Based on the measured remaining living target cells, the percentage of specific cell lysis is calculated according to the following formula: $[1 - (\text{number of living targets}_{\text{(sample)}} / \text{number of living targets}_{\text{(spontaneous)}})] \times 100\%$. Sigmoidal dose response curves and EC₅₀ values are calculated by non-linear regression/4-parameter logistic fit using the GraphPad Software. The lysis values obtained for a given antibody concentration are used to calculate sigmoidal dose-response curves by 4 parameter logistic fit analysis using the Prism software.

Example 18: Pharmacokinetics of activatable IFN γ Proteins

[368] The activatable IFN γ protein is evaluated for half-time elimination in animal studies.

[369] The activatable IFN γ protein is administered to cynomolgus monkeys as a 0.5 mg/kg bolus injection into the saphenous vein. Another cynomolgus monkey group receives a comparable cytokine in size but lacking a serum half-life extension element. A third and fourth group receive a cytokine with serum half-life extension elements and a cytokine with CD20 and serum half-life extension elements respectively, and both comparable in size to the activatable IFN γ protein. Each test group consists of 5 monkeys. Serum samples are taken at indicated time points, serially diluted, and the concentration of the proteins is determined using a binding ELISA to CD20.

[370] Pharmacokinetic analysis is performed using the test article plasma concentrations. Group mean plasma data for each test article conforms to a multi-exponential profile when plotted against the time post-dosing. The data are fit by a standard two-compartment model with bolus input and first-order rate constants for distribution and elimination phases. The general equation for the best fit of the data for i.v. administration is: $c(t) = Ae^{-\alpha t} + Be^{-\beta t}$, where $c(t)$ is the plasma concentration at time t , A and B are intercepts on the Y-axis, and α and β are the apparent first-order rate constants for the distribution and elimination phases, respectively. The α -phase is the initial phase of the clearance and reflects distribution of the protein into all extracellular fluid of the animal, whereas the second or β -phase portion of the decay curve represents true plasma clearance. Methods for fitting such equations are well known in the art. For example, $A = D/V(\alpha - k_{21})/(\alpha - \beta)$, $B = D/V(\beta - k_{21})/(\alpha - \beta)$, and α and β (for $\alpha > \beta$) are roots of the quadratic equation:

$r^2 + (k_{12} + k_{21} + k_{10})r + k_{21}k_{10} = 0$ using estimated parameters of V =volume of distribution, k_{10} =elimination rate, k_{12} =transfer rate from compartment 1 to compartment 2 and k_{21} =transfer rate from compartment 2 to compartment 1, and D =the administered dose.

[371] Data analysis: Graphs of concentration versus time profiles are made using KaleidaGraph (KaleidaGraph™ V. 3.09 Copyright 1986-1997. Synergy Software. Reading, Pa.). Values reported as less than reportable (LTR) are not included in the PK analysis and are not represented graphically. Pharmacokinetic parameters are determined by compartmental analysis using WinNonlin software (WinNonlin® Professional V. 3.1 WinNonlin™ Copyright 1998-1999. Pharsight Corporation. Mountain View, Calif.). Pharmacokinetic parameters are computed as described in Ritschel W A and Kearns G L, 1999, IN: *Handbook of Basic Pharmacokinetics Including Clinical Applications*, 5th edition, American Pharmaceutical Assoc., Washington, D.C.

[372] It is expected that the activatable IFN γ protein has improved pharmacokinetic parameters such as an increase in elimination half-time as compared to proteins lacking a serum half-life extension element.

Example 19: Xenograft Tumor Model

[373] The activatable IFN γ protein is evaluated in a xenograft model.

[374] Female immune-deficient NOD/scid mice are sub-lethally irradiated (2 Gy) and subcutaneously inoculated with 4×10^6 Ramos RA1 cells into the right dorsal flank. When tumors reach 100 to 200 mm³, animals are allocated into 3 treatment groups. Groups 2 and 3 (8 animals each) are intraperitoneally injected with 1.5×10^7 activated human T-cells. Three days later, animals from Group 3 are subsequently treated with a total of 9 intravenous doses of 50 μ g activatable IFN γ protein. Groups 1 and 2 are only treated with vehicle. Body weight and tumor volume are determined for 30 days.

[375] It is expected that animals treated with the activatable IFN γ protein have a statistically significant delay in tumor growth in comparison to the respective vehicle-treated control group.

[376] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

Example 20: HEK Blue assay

[377] HEK-Blue IL12 cells (InvivoGen) were plated in suspension at a concentration of 250,000

cells/well in culture media with or without 40mg/ml human serum albumin (HSA) and stimulated with a dilution series of recombinant hIL12, chimeric IL12 (mouse p35/human p40) or activatable hIL12 for 24 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved activatable hIL12 was tested. Cleaved inducible hIL12 was generated by incubation with active MMP9. IL12 activity was assessed by quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen), a colorimetric based assay.

[378] HEK-Blue IL2 cells (InvivoGen) were plated in suspension at a concentration of 50,000 cells/well in culture media with or without 15-40mg/ml human serum albumin (HSA) and stimulated with a dilution series of recombinant hIL2 or activatable hIL2 for 24 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved activatable hIL2 was tested. Cleaved inducible hIL2 was generated by incubation with active MMP9 or another protease. IL2 activity was assessed by quantification of Secreted Alkaline Phosphatase (SEAP) activity using the reagent QUANTI-Blue (InvivoGen), a colorimetric based assay. Results are shown in FIGs. 59-62.

Example 21: Splenocyte T-Blast assay

[379] T-Blasts were induced from murine splenocytes with a 6-day incubation with PHA and a 24hr incubation with recombinant hIL12. T-blasts were then plated in suspension at a concentration of 200,000 cells/well in culture media with or without 40mg/ml human serum albumin (HSA) and stimulated with a dilution series of recombinant hIL12 or chimeric IL12 (mouse p35/human p40) or mouse IL12 for 72 hours at 37°C and 5% CO₂. Activity of uncleaved and cleaved IL12 was tested. Cleaved inducible hIL12 was generated by incubation with active MMP9. IL12 activity was assessed by downstream quantification of IFN γ production using a mIFN γ alphaLISA.

Example 22: *In Vivo* Delivery of a Protease Activated Fusion Protein Results in Decreased Tumor Growth

[380] The chimeric polypeptide is examined to determine if it could have biological effects *in vivo*. For these experiments a system is used in which tumor cells injected intraperitoneally rapidly and preferentially attach and grow initially on the milky spots, a series of organized immune aggregates found on the omentum (Gerber et al., Am. J. Pathol. 169:1739-52 (2006)). This system offers a convenient way to examine the effects of fusion protein treatment on tumor growth since fusion proteins can be delivered intraperitoneally multiple times and tumor growth can be analyzed by examining the dissociated omental cells. For these experiments, the Colon 38 cell line, a rapidly growing tumor cell line that expresses both MMP2 and MMP9 *in vitro*, may be used. The omental tissue normally expresses a relatively small amount of MMP2 and MMP9, but, when Colon 38 tumor is present on the omentum, MMP levels increase. Using this tumor model, the ability of IL-

2 muetein fusion proteins to affect tumor growth is examined. Colon 38 cells are injected intraperitoneally, allowed to attach and grow for 1 day, and then treated daily with fusion protein intraperitoneally. At day 7, the animals are sacrificed and the omenta examined for tumor growth using flow cytometry and by a colony-forming assay.

Example 23A: Construction of an Exemplary Activatable Interleukin Protein Targeting CD20

23.1 Generation of an activatable interleukin domain

[381] The human IL-12p35 chain canonical sequence is UniProt Accession No. P29459. The human IL-12p40 chain canonical sequence is UniProt Accession No. P29460. IL-12p35 and IL-12p40 are cloned into an expression construct. A protease cleavage site is included between the IL-12p35 and IL-12p40 domains. An IL-12 polypeptide capable of binding to CD20 polypeptide present in a tumor or on a tumor cell is produced as follows. A nucleic acid is produced that contains nucleic acid sequences: (1) encoding an IFN γ polypeptide sequence and (2) one or more polypeptide linkers. Activatable interleukin plasmid constructs can have optional Flag, His or other affinity tags, and are electroporated into HEK293 or other suitable human or mammalian cell lines and purified. Validation assays include T cell activation assays using T cells responsive to IL-12 stimulation in the presence of a protease.

23.2 Generation of a scFv CD20 binding domain

[382] CD20 is one of the cell surface proteins present on B-lymphocytes. CD20 antigen is found in normal and malignant pre-B and mature B lymphocytes, including those in over 90% of B-cell non-Hodgkin's lymphomas (NHL). The antigen is absent in hematopoietic stem cells, activated B lymphocytes (plasma cells) and normal tissue. As such, several antibodies mostly of murine origin have been described: 1F5, 2B8/C2B8, 2H7, and 1H4.

[383] Human or humanized anti-CD20 antibodies are therefore used to generate scFv sequences for CD20 binding domains of an activatable interleukin protein. DNA sequences coding for human or humanized VL and VH domains are obtained, and the codons for the constructs are, optionally, optimized for expression in cells from Homo sapiens. The order in which the VL and VH domains appear in the scFv is varied (i.e., VL-VH, or VH-VL orientation), and three copies of the "G4S" (SEQ ID NO: 241) or "G₄S" (SEQ ID NO: 241) subunit (G₄S)₃ (SEQ ID NO: 242) connect the variable domains to create the scFv domain. Anti-CD20 scFv plasmid constructs can have optional Flag, His or other affinity tags, and are electroporated into HEK293 or other suitable human or mammalian cell lines and purified. Validation assays include binding analysis by FACS, kinetic analysis using Proteon, and staining of CD20-expressing cells.

23.3 Cloning of DNA expression constructs encoding the activatable interleukin protein

[384] The activatable interleukin construct with protease cleavage site domains are used to construct an activatable interleukin protein in combination with an anti-CD20 scFv domain and a serum half-life extension element (e.g., a HSA binding peptide or VH domain). For expression of an activatable interleukin protein in CHO cells, coding sequences of all protein domains are cloned into a mammalian expression vector system. In brief, gene sequences encoding the activatable interleukin domain, serum half-life extension element, and CD20 binding domain along with peptide linkers L1 and L2 are separately synthesized and subcloned. The resulting constructs are then ligated together in the order of CD20 binding domain – L1 – IL-12p35 – L2 – protease cleavage domain – L3 – IL-12p40 – L4 – anti-CD20 scFv – L5 – serum half-life extension element to yield a final construct. All expression constructs are designed to contain coding sequences for an N-terminal signal peptide and a C-terminal hexahistidine (6xHis)-tag (SEQ ID NO: 243) to facilitate protein secretion and purification, respectively.

23.4 Expression of activatable interleukin proteins in stably transfected CHO cells

[385] A CHO cell expression system (Flp-In®, Life Technologies), a derivative of CHO-K1 Chinese Hamster ovary cells (ATCC, CCL-61) (Kao and Puck, Proc. Natl. Acad. Sci. USA 1968;60(4):1275-81), is used. Adherent cells are subcultured according to standard cell culture protocols provided by Life Technologies.

[386] For adaption to growth in suspension, cells are detached from tissue culture flasks and placed in serum-free medium. Suspension-adapted cells are cryopreserved in medium with 10% DMSO.

[387] Recombinant CHO cell lines stably expressing secreted activatable interleukin proteins are generated by transfection of suspension-adapted cells. During selection with the antibiotic Hygromycin B viable cell densities are measured twice a week, and cells are centrifuged and resuspended in fresh selection medium at a maximal density of 0.1×10^6 viable cells/mL. Cell pools stably expressing activatable interleukin proteins are recovered after 2-3 weeks of selection at which point cells are transferred to standard culture medium in shake flasks. Expression of recombinant secreted proteins is confirmed by performing protein gel electrophoresis or flow cytometry. Stable cell pools are cryopreserved in DMSO containing medium.

[388] Activatable interleukin proteins are produced in 10-day fed-batch cultures of stably transfected CHO cell lines by secretion into the cell culture supernatant. Cell culture supernatants are harvested after 10 days at culture viabilities of typically >75%. Samples are collected from the production cultures every other day and cell density and viability are assessed. On day of harvest, cell culture supernatants are cleared by centrifugation and vacuum filtration before further use.

[389] Protein expression titers and product integrity in cell culture supernatants are analyzed by SDS-PAGE.

23.5 Purification of activatable interleukin proteins

[390] Activatable interleukin proteins are purified from CHO cell culture supernatants in a two-step procedure. The constructs are subjected to affinity chromatography in a first step followed by preparative size exclusion chromatography (SEC) on Superdex 200 in a second step. Samples are buffer-exchanged and concentrated by ultrafiltration to a typical concentration of >1 mg/mL. Purity and homogeneity (typically >90%) of final samples are assessed by SDS PAGE under reducing and non-reducing conditions, followed by immunoblotting using an anti-HSA or anti idotype antibody as well as by analytical SEC, respectively. Purified proteins are stored at aliquots at -80°C until use.

Example 23B: Determination of antigen affinity by flow cytometry

[391] Activatable interleukin proteins are tested for their binding affinities to human CD20⁺ cells and cynomolgus CD20⁺ cells.

[392] CD20⁺ cells are incubated with 100 µL of serial dilutions of the activatable interleukin proteins and at least one protease. After washing three times with FACS buffer the cells are incubated with 0.1 mL of 10 µg/mL mouse monoclonal anti-idiotypic antibody in the same buffer for 45 min on ice. After a second washing cycle, the cells are incubated with 0.1 mL of 15 µg/mL FITC-conjugated goat anti-mouse IgG antibodies under the same conditions as before. As a control, cells are incubated with the anti-His IgG followed by the FITC-conjugated goat anti-mouse IgG antibodies without the activatable interleukin proteins. The cells were then washed again and resuspended in 0.2 mL of FACS buffer containing 2 µg/mL propidium iodide (PI) in order to exclude dead cells. The fluorescence of 1×10^4 living cells is measured using a Beckman-Coulter FC500 MPL flow cytometer using the MXP software (Beckman-Coulter, Krefeld, Germany) or a Millipore Guava EasyCyte flow cytometer using the Incyte software (Merck Millipore, Schwalbach, Germany). Mean fluorescence intensities of the cell samples are calculated using CXP software (Beckman-Coulter, Krefeld, Germany) or Incyte software (Merck Millipore, Schwalbach, Germany). After subtracting the fluorescence intensity values of the cells stained with the secondary and tertiary reagents alone the values are then used for calculation of the K_D values with the equation for one-site binding (hyperbola) of the GraphPad Prism (version 6.00 for Windows, GraphPad Software, La Jolla California USA).

[393] CD20 binding and cross-reactivity are assessed on the human CD20⁺ tumor cell lines. The K_D ratio of cross-reactivity is calculated using the K_D values determined on the CHO cell lines expressing either recombinant human or recombinant cynomolgus antigens.

Example 24: Cytotoxicity Assay

[394] The activatable interleukin protein is evaluated *in vitro* on its mediation of immune response to CD20⁺ target cells.

[395] Fluorescence labeled CD20⁺ REC-1 cells (a Mantle cell lymphoma cell line, ATCC CRL-3004) are incubated with isolated PBMC of random donors or CB15 T-cells (standardized T-cell line) as effector cells in the presence of the activatable interleukin protein and at least one protease. After incubation for 4 h at 37°C in a humidified incubator, the release of the fluorescent dye from the target cells into the supernatant is determined in a spectrofluorometer. Target cells incubated without the activatable interleukin protein and target cells totally lysed by the addition of saponin at the end of the incubation serve as negative and positive controls, respectively.

[396] Based on the measured remaining living target cells, the percentage of specific cell lysis is calculated according to the following formula: $[1 - (\text{number of living targets}_{\text{(sample)}} / \text{number of living targets}_{\text{(spontaneous)}})] \times 100\%$. Sigmoidal dose response curves and EC₅₀ values are calculated by non-linear regression/4-parameter logistic fit using the GraphPad Software. The lysis values obtained for a given antibody concentration are used to calculate sigmoidal dose-response curves by 4 parameter logistic fit analysis using the Prism software.

Example 25: Pharmacokinetics of activatable interleukin Proteins

[397] The activatable interleukin protein is evaluated for half-time elimination in animal studies.

[398] The activatable interleukin protein is administered to cynomolgus monkeys as a 0.5 mg/kg bolus injection into the saphenous vein. Another cynomolgus monkey group receives a comparable cytokine in size but lacking a serum half-life extension element. A third and fourth group receive a cytokine with serum half-life extension elements and a cytokine with CD20 and serum half-life extension elements respectively, and both comparable in size to the activatable interleukin protein. Each test group consists of 5 monkeys. Serum samples are taken at indicated time points, serially diluted, and the concentration of the proteins is determined using a binding ELISA to CD20.

[399] Pharmacokinetic analysis is performed using the test article plasma concentrations. Group mean plasma data for each test article conforms to a multi-exponential profile when plotted against the time post-dosing. The data are fit by a standard two-compartment model with bolus input and first-order rate constants for distribution and elimination phases. The general equation for the best fit of the data for i.v. administration is: $c(t) = Ae^{-\alpha t} + Be^{-\beta t}$, where $c(t)$ is the plasma concentration at time t , A and B are intercepts on the Y-axis, and α and β are the apparent first-order rate constants for the distribution and elimination phases, respectively. The α -phase is the initial phase of the clearance and reflects distribution of the protein into all extracellular fluid of the animal, whereas the second or β -phase portion of the decay curve represents true plasma clearance. Methods for

fitting such equations are well known in the art. For example, $A=D/V(\alpha-k_{21})/(\alpha-\beta)$, $B=D/V(\beta-k_{21})/(\alpha-\beta)$, and α and β (for $\alpha>\beta$) are roots of the quadratic equation: $r^2+(k_{12}+k_{21}+k_{10})r+k_{21}k_{10}=0$ using estimated parameters of V =volume of distribution, k_{10} =elimination rate, k_{12} =transfer rate from compartment 1 to compartment 2 and k_{21} =transfer rate from compartment 2 to compartment 1, and D =the administered dose.

[400] Data analysis: Graphs of concentration versus time profiles are made using KaleidaGraph (KaleidaGraph™ V. 3.09 Copyright 1986-1997. Synergy Software. Reading, Pa.). Values reported as less than reportable (LTR) are not included in the PK analysis and are not represented graphically. Pharmacokinetic parameters are determined by compartmental analysis using WinNonlin software (WinNonlin® Professional V. 3.1 WinNonlin™ Copyright 1998-1999. Pharsight Corporation. Mountain View, Calif.). Pharmacokinetic parameters are computed as described in Ritschel W A and Kearns G L, 1999, IN: *Handbook of Basic Pharmacokinetics Including Clinical Applications*, 5th edition, American Pharmaceutical Assoc., Washington, D.C.

[401] It is expected that the activatable interleukin protein has improved pharmacokinetic parameters such as an increase in elimination half-time as compared to proteins lacking a serum half-life extension element.

Example 26: Xenograft Tumor Model

[402] Activatable interleukin protein is evaluated in a xenograft model.

[403] Female immune-deficient NOD/scid mice are sub-lethally irradiated (2 Gy) and subcutaneously inoculated with 4×10^6 Ramos RA1 cells into the right dorsal flank. When tumors reach 100 to 200 mm³, animals are allocated into 3 treatment groups. Groups 2 and 3 (8 animals each) are intraperitoneally injected with 1.5×10^7 activated human T-cells. Three days later, animals from Group 3 are subsequently treated with a total of 9 intravenous doses of 50 µg activatable interleukin protein. Groups 1 and 2 are only treated with vehicle. Body weight and tumor volume are determined for 30 days.

Table 3. Summary of the treatment modes

Gr.	N	Agent	Formulation dose	Route	Schedule
1 [#]	10	Vehicle	-	ip	biwk x 3
2	7	ACP16	700µg/animal	ip	biwk x 3
3	7	ACP16	230µg/animal	ip	biwk x 3
4	7	ACP16	70µg/animal	ip	biwk x 3
5	7	ACP16	55ug/animal	ip	biwk x 3

6	7	ACP16	17 μ g/animal	ip	biwk x 3
7	7	ACP132	361 μ g/animal	ip	biwk x 3
8	7	ACP132	119 μ g/animal	ip	biwk x 3
9	7	ACP132	36 μ g/animal	ip	biwk x 3
10	7	ACP132	28 μ g/animal	ip	biwk x 3
11	7	ACP132	9 μ g/animal	ip	biwk x 3
12	7	ACP21	540 μ g/animal	ip	biwk x 3
13	7	ACP21	177 μ g/animal	ip	biwk x 3
14	7	ACP21	54 μ g/animal	ip	biwk x 3
15	7	ACP21	42 μ g/animal	ip	biwk x 3
16	7	ACP21	13 μ g/animal	ip	biwk x 3
17	7	ACP133	210 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
18	7	ACP133	105 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
19	7	ACP133	40 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
20	7	ACP133	3 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
#	- Control Group				

[348] It is expected that animals treated with the activatable interleukin protein have a statistically significant delay in tumor growth in comparison to the respective vehicle-treated control group.

[404] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

[405] The MC38 cell line, a rapidly growing colon adenocarcinoma cell line that expresses MMP9 *in vitro*, was used. Using this tumor model, the ability of fusion proteins to affect tumor growth was examined.

Example 27A: MC38 IL-2POC

27A.1 Agents and Treatment

[406] Additional studies were carried out in non-tumor bearing animals as described below.

Table 4. Summarizes the treatment regime.

Gr.	N	Agent	Formulation dose	Route	Schedule
1	8	Vehicle	-	ip	biwk x 4
2	8	ACP16	700µg/animal	ip	biwk x 4
3	8	ACP16	230µg/animal	ip	biwk x 4
4	8	ACP16	70µg/animal	ip	biwk x 4
8	8	ACP153	700µg/animal	ip	biwk x 4
9	8	ACP153	230µg/animal	ip	biwk x 4
10	8	ACP153	70µg/animal	ip	biwk x 4
11	8	ACP154	700µg/animal	ip	biwk x 4
12	8	ACP154	230µg/animal	ip	biwk x 4
13	8	ACP154	70µg/animal	ip	biwk x 4
14	8	ACP155	700µg/animal	ip	biwk x 4
15	8	ACP155	230µg/animal	ip	biwk x 4
16	8	ACP155	70µg/animal	ip	biwk x 4
17	8	ACP156	700µg/animal	ip	biwk x 4
18	8	ACP156	230µg/animal	ip	biwk x 4
19	8	ACP156	70µg/animal	ip	biwk x 4
20	8	ACP157	700µg/animal	ip	biwk x 4
21	8	ACP157	230µg/animal	ip	biwk x 4
22	8	ACP157	70µg/animal	ip	biwk x 4

Table 5. Describes the constructs used in the MC38 IL-2POC animal study.

Construct Name	Description	MW
ACP16	IL2-X-HSA-LX-blocker Fusion protein-6xHis	58256
ACP133	IL-2 with C term 6x His	16462
ACP132	IL2-L-HSA	29996
ACP21	IL2-XL-blocker Fusion protein-6xHis	44843

Example 27B: MC38 IL-2

Table 6. Summarizes the treatment regime.

Gr.	N	Agent	Formulation dose	Route	Schedule
1 [#]	12	Vehicle	-	ip	biwk x 2
2	8	ACP16	4.4 µg/animal	ip	biwk x 2
3	8	ACP16	17 µg/animal	ip	biwk x 2
4	8	ACP16	70 µg/animal	ip	biwk x 2
5	8	ACP16	232 µg/animal	ip	biwk x 2
6	8	ACP130	19 µg/animal	ip	biwk x 2
7	8	ACP130	45 µg/animal	ip	biwk x 2
8	8	ACP130	180 µg/animal	ip	biwk x 2
9	8	ACP130	600 µg/animal	ip	biwk x 1
12	8	ACP124	17 µg/animal	ip	biwk x 2
13	8	ACP124	70 µg/animal	ip	biwk x 2
14	8	ACP124	230 µg/animal	ip	biwk x 2
15	8	ACP124	700 µg/animal	ip	biwk x 2
16	8	IL-2-WTI	12 µg/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
17	8	IL-2-	36 µg/animal	ip	bid x 5 then 2-day pause

		WTI			then bid x 5 then 2-day pause
#	- Control Group				

27B.1 Procedure

[407] Mice were anaesthetized with isoflurane for implant of cells to reduce the ulcerations. 308 CR female C57BL/6 mice were set up with 5×10^5 MC38 tumor cells in 0% Matrigel sc in flank. Cell Injection Volume was 0.1 mL/mouse. Mouse age at start date was 8 to 12 weeks. Pair matches were performed when tumors reach an average size of 100 - 150 mm³ and begin treatment. Body weights were taken at initiation and then biweekly to the end. Caliper measurements were taken biweekly to the end. Any adverse reactions were to be reported immediately. Any individual animal with a single observation of > than 30% body weight loss or three consecutive measurements of >25% body weight loss was euthanized. Any group with a mean body weight loss of >20 % or >10% mortality stopped dosing; the group was not euthanized and recovery is allowed. Within a group with >20% weight loss, individuals hitting the individual body weight loss endpoint were euthanized. If the group treatment related body weight loss is recovered to within 10% of the original weights, dosing resumed at a lower dose or less frequent dosing schedule. Exceptions to non-treatment body weight % recovery were allowed on a case-by-case basis. Endpoint was tumor growth delay (TGD). Animals were monitored individually. The endpoint of the experiment was a tumor volume of 1500 mm³ or 45 days, whichever comes first. Responders were followed longer. When the endpoint was reached, the animals are to be euthanized.

27B.2 Dosing Instructions

[408] No compounds in salt form were used. The amount needed per week was calculated, aliquoted accordingly, and stored at -20 C. For each week of dosing, one aliquot was thawed, stored at 4 C, and diluted in the required amount with PBS right before each injection.

IL-2-WTI required protection from light; pre-formulation stored at- 4° C, post-formulation – stored at-20° C. Lyophilized material was reconstituted as directed by instructions, similar to above. ACP16, ACP130, ACP124, and IL-2 WTI were prepared for dosing in PBS. IL-2-WTI indicates Proleukin (aldesleukin) in PBS; the vehicle was PBS.

[409] Dosing volume was 0.2 mL/mouse for IL-2-WTI; 0.3 mL for ACP16, ACP130; 0.5 mL for ACP124. Do not adjust for body weight.

27B.3 Special Instructions

[410] ACP16: current amount of required compound - 13.45 mg

[411] ACP130: current amount of required compound -25.83 mg

- [412] ACP124: current amount of required compound -42.31 mg
 [413] IL-2-WTI: current amount of required compound - 9.98 mg
 [414] Necropsy was to be performed in case of unexpected toxicity

Example 27C: MC38 IFN α and IL-12

27C.1 Agents and Treatment:

Table 7. Summarizes the treatment regime.

Gr.	N	Agent	Formulation dose	Route	Schedule
1 [#]	12	Vehicle	-	ip	biwk x 3
2	8	ACP11	17.5 μ g/animal	ip	biwk x 3
3	8	ACP11	175 μ g/animal	ip	biwk x 3
4	8	ACP11	525 μ g/animal	ip	biwk x 3
5	8	ACP31	33 μ g/animal	ip	biwk x 3
6	8	ACP31	110 μ g/animal	ip	biwk x 3
7	8	ACP31	330 μ g/animal	ip	biwk x 3
8	8	ACP131	1 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
9	8	ACP131	10 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
10	8	ACP131	30 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
11	8	mIFN α 1 -WTI	1 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
12	8	mIFN α 1 -WTI	10 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
13	8	IL-12- HM- WTI	2 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
14	8	IL-12- HM- WTI	10 μ g/animal	ip	bid x 5 then 2-day pause then bid x 5 then 2-day pause
15	8	ACP131	5 μ g/animal	itu	bid x 5 then 2-day pause then bid x 5 then 2-day pause
#	- Control Group				

27C.2 Procedures

[415] Mice were anaesthetized with isoflurane for implant of cells to reduce the ulcerations. 308 CR female C57BL/6 mice were set up with 5×10^5 MC38 tumor cells in 0% Matrigel sc in flank. Cell Injection Volume was 0.1 mL/mouse. Mouse age at start date was 8 to 12 weeks. Pair matches were performed when tumors reach an average size of 100 - 150 mm³ and begin treatment. Body weights were taken at initiation and then biweekly to the end. Caliper measurements were taken biweekly to the end. Any adverse reactions were to be reported immediately. Any individual animal with a single observation of > than 30% body weight loss or three consecutive measurements of >25% body weight loss was euthanized. Any group with a mean body weight loss of >20 % or >10% mortality stopped dosing; the group was not euthanized and recovery is allowed. Within a group with >20% weight loss, individuals hitting the individual body weight loss endpoint were euthanized. If the group treatment related body weight loss is recovered to within 10% of the original weights, dosing resumed at a lower dose or less frequent dosing schedule. Exceptions to non-treatment body weight % recovery were allowed on a case-by-case basis. Endpoint was tumor growth delay (TGD). Animals were monitored individually. The endpoint of the experiment was a tumor volume of 1500 mm³ or 45 days, whichever comes first. Responders were followed longer. When the endpoint was reached, the animals are to be euthanized.

27C.3 Dosing Instructions

[416] No compounds in salt form were used. Prepared dosing solutions were as follows: IL-12-HM-WTI was stored to provide protection from light; pre-formulation at -4° C, post-formulation at -20° C. mIFNa1-WTI was stored at -20° C, protected from light; pre-formulation stored at -20° C, post-formulation stored at -4° C. Lyophilized material was reconstituted as directed in instructions. The amount needed per week was calculated, aliquoted accordingly, and stored at -20 C. For each week of dosing, one aliquot was thawed and stored at 4 C, the required amount was diluted with PBS right before each injection.

[417] For ACP11, ACP31, ACP131, the amount needed per week was calculated, aliquoted accordingly, and stored at -20 C. For each week of dosing, one aliquot was thawed and stored at 4 C, the required amount was diluted with PBS right before each injection. PBS was used as the vehicle for all tests.

[418] Intraperitoneal (ip) dosing volume ACP131, mIFNa1-WTI, IL-12-HM-WTI = 0.2 mL/mouse. Dosing volume for ACP11 = 0.4 mL (before Day 18, 2/26/19) and 0.55 mL (starting on Day 18, 2/26/19).

[419] Dosing volume for ACP31 = 0.3 mL. Dosage was not adjusted for body weight. Intratumoral (itu) dosing volume for Gr.16 mIFNa1-WTI and Gr.17 ACP131 = 0.05 mL/mouse. Dosage was not adjusted for body weight.

Example 27D: MC38 rechallenge**27D.1 Agents and Treatment:**

Table 8. Summarizes the treatment regime.

Gr.	N	Agent	Formulation dose	Route	Schedule
1 [#]	33	No Treatment	-	-	-
2	7	ACP16	70 µg/animal	ip	MC38-e415 Gr 4 An 1-4, 6-8 (ACP16 biwx2)
3	8	ACP16	232 µg/animal	ip	MC38-e415 Gr 5 An 1-8 (ACP16 biwx2)
5	5	IL-2-WTI	12 µg/animal	ip	MC38-e415 Gr 16 An 1,3-6 (IL-2-WTI bid x 5 then 2-day pause then bid x 5 then 2-day pause)
6	7	IL-2-WTI	36 µg/animal	ip	MC38-e415 Gr 17 An 1-7 (IL-2-WTI bid x 5 then 2-day pause then bid x 5 then 2-day pause)
#	- Control Group				

27D.2 Procedures

[420] Mice were anaesthetized with isoflurane for implant of cells to reduce the ulcerations. This portion of the study began on the day of implant (Day 1). Group 1 consisted of 33 CR female C57BL/6 mice set up with 5x10⁵ MC38 tumor cells in 0% Matrigel subcutaneously in the flank. Groups 2-6 consisted of 33 CR female C57BL/6 mice set up with 5x10⁵ MC38 tumor cells in 0% Matrigel sc in the left flank. The tumors from the previous MC38 experiment (example 25x) were implanted in the right flank of each animal. Cell Injection Volume was 0.1 mL/mouse. Age of control mice at initiation was 14 to 17 weeks. These mice were age matched to mice from the previous MC38 experiment (example 25x). No dosing of active agent occurred during rechallenge. Body Weights were taken biweekly until end, as were caliper measurements. Any adverse reactions or death were reported immediately. Any individual animal with a single observation of > than 30% body weight loss or three consecutive measurements of >25% body weight loss was euthanized. Endpoint was tumor growth delay (TGD). Animals were monitored individually. The

endpoint of the experiment was a tumor volume of 1000 mm³ or 45 days, whichever comes first. Responders were followed longer when possible. When the endpoint is reached, the animals were euthanized.

Example 27E: Treatment with ACP16, ACP153, ACP155, and ACP156 (see FIG. 58)

27E.1 Agents and Treatment:

Table 9. Summarizes the treatment regime.

Gr.	N	Agent	Formulation dose	Route	Schedule
1 [#]	12	Vehicle	-	ip	biwk x 2
2	8	ACP16	17 µg/animal	ip	biwk x 2
3	8	ACP16	55 µg/animal	ip	biwk x 2
4	8	ACP16	230 µg/animal	ip	biwk x 2
5	8	ACP155	55 µg/animal	ip	biwk x 2
6	8	ACP155	230 µg/animal	ip	biwk x 2
7	8	ACP153	55 µg/animal	ip	biwk x 2
8	8	ACP153	230 µg/animal	ip	biwk x 2
9	8	ACP156	55 µg/animal	ip	biwk x 2
10	8	ACP156	230 µg/animal	ip	biwk x 2

27E.2 Procedures:

[421] Mice were anaesthetized with isoflurane for implant of cells to reduce the ulcerations. CR female C57BL/6 mice were set up with 5×10^5 MC38 tumor cells in 0% Matrigel sc in flank. Cell Injection Volume was 0.1 mL/mouse. Mouse age at start date was 8 to 12 weeks. Pair matches were performed when tumors reach an average size of 100 - 150 mm³ and begin treatment. ACP16 was dosed at 17, 55 or 230 µg/animal; ACP153, ACP155 and ACP156 were dosed at 55 or 230 µg/animal. Body weights were taken at initiation and then biweekly to the end. Caliper measurements were taken biweekly to the end. Any adverse reactions were to be reported immediately. Any individual animal with a single observation of > than 30% body weight loss or three consecutive measurements of >25% body weight loss was euthanized. Any group with a mean body weight loss of >20 % or >10% mortality stopped dosing; the group was not euthanized and recovery is allowed. Within a group with >20% weight loss, individuals hitting the individual body weight loss endpoint were euthanized. If the group treatment related body weight loss is recovered to within 10% of the original weights, dosing resumed at a lower dose or less frequent dosing schedule. Exceptions to non-treatment body weight % recovery were allowed on a case-by-

case basis. Endpoint was tumor growth delay (TGD). Animals were monitored individually. The endpoint of the experiment was a tumor volume of 1500 mm³ or 45 days, whichever comes first. Responders were followed longer. When the endpoint was reached, the animals are to be euthanized. Results are shown in Fig 58A-58D.

Example 28: FRET screens of conditioned media

[422] Proteolytic activity was screened in the conditioned media samples by fluorescent resonance energy transfer (FRET) assays with the substrates listed in Table 10.

Table 10. Substrate motif sequences.

Substrate Name	P4 – P4' Sequence	SEQ ID NO:	Experimentally Tested		
			in vitro kinetics	Conditioned media	30-mer MSP-MS
MMP14_1	GPAGLYAQ	195	✓	✓	✓
MMP9_1	GPAGMKGL	196	✓	✓	-
FAP α _1	PGGPAGIG	197	✓	✓	✓
CTSL1_1	ALFKSSFP	198	✓	✓	✓
CTSL1_2	ALFFSSPP	199	✓	✓	✓
ADAM17_1	LAQRLRSS	200	✓	✓	✓
ADAM17_2	LAQKLKSS	201	✓	✓	✓

28.1 Reaction conditions

[423] Protease specificity screening was performed using Multiplexed Substrate Profiling by Mass Spectrometry (MSP-MS), the method described in, e.g. O'Donoghue A. J. et al., *Nat Methods.*, 2012; 9(11): 1095–1100. This method employs a physico-chemically diverse peptide library as substrates for proteases, and reactions are monitored over time with mass spectrometric detection of cleaved products. The resulting cleavages are assessed for specific cleavage in the enzyme-treated sample by comparison with results from a no-enzyme control incubation. Sequence logos for motif analysis are generated with iceLogo software (v.1.2) (ionics.ugent.be/icelogoserver/).

[424] Recombinant human enzymes were sourced from R&D Systems (Bio-technie): CTSL1 (#952-CY), ADAM17 (a.k.a. TACE, TNF-alpha Converting Enzyme) (#930-ADB), FAP-alpha (#930-ADB-010), MMP-14 (#918-MP-010), furin (#1503-SE-010), MMP-9 (#911-MP-010), thrombin (#2196-SE-200), thermolysin (#3097-ZN), Factor Xa (#1063-SE-010), hepsin (#4776-

SE) and ADAM-TS1 (#2197-AD). Enzymes were activated and assayed following the manufacturer's recommendations.

[425] For CTSL1, enzyme was activated by pre-incubation in assay buffer (50mM MES, 5mM DTT, 1mM EDTA, pH = 6) for 15 min. MSP-MS reactions were then started with the mixing of enzyme and substrate, at final CTSL1 concentration of 0.04 ng/μl (0.77 nM), and peptide substrate concentration at 500 nM.

[426] ADAM17 activity was monitored following manufacturer's recommended conditions using the FRET substrate Mca-PLAQAV-Dpa-RSSSR-NH₂ (SEQ ID NO: 244), in the recommended assay buffer: 25mM TRIS, 2.5μM ZnCl₂ at the optimal pH 9.0, and also at physiological pH 7.4. ADAM17 enzyme specificity was profiled in the MSP-MS experiment at pH 9.0 using 10 nM enzyme, and at pH 7.4 using 50 nM enzyme.

[427] MMP-14 was activated following manufacturer's recommended conditions by pre-incubation with the enzyme furin, at a molar ratio of 1:100 furin to MMP-14, at 37° C and pH 9.0 for 1.5 hours. FRET assays were performed using 20 nM MMP-14 at 37° C in activation buffer: 50mM TRIS HCl, 3mM CaCl₂, 1μM ZnCl₂ at pH 8.5.

[428] MMP9 was activated by incubation with 1mM p-amino phenyl mercuric acetate (prepared from a stock at 100mM in DMSO) for 24h at 37° C in activation buffer: 50mM TRIS HCl (pH 7.5), 10mM CaCl₂, 150mM NaCl, 0.05%Brij35. The final enzyme concentration used for fluorescence assays was 2.5 nM.

[429] FAPα activity was tested with fluorescence detection using a generic Z-GP-AMC substrate. For FRET assays, the final enzyme concentration was 5 nM and the buffer was 50 mM TRIS HCl, 1 M NaCl, 1 mg/mL BSA, pH 7.5.

[430] Thrombin was activated following the manufacturer's recommended conditions by pre-incubation with the enzyme with thermolysin for 15 min, and then the thermolysin was quenched with 1,10 phenanthroline treatment. Activated thrombin was assayed at 1.2 nM in activation buffer: 50 mM TRIS HCl, 10 mM CaCl₂, 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5.

[431] Factor Xa was assayed at 4.7 nM enzyme, at 37° C in the manufacturer recommended buffer: 50 mM TRIS, 10 mM CaCl₂, 150 mM NaCl, pH 7.5.

[432] Hepsin was pre-activated overnight at 37° C in the manufacturer recommended buffer: 100 mM TRIS, 10 mM CaCl₂, 150 mM NaCl, 0.05% Brij-35, pH 8.0. Activity was measured with 2.4 or 24 nM enzyme in the buffer: 50mM TRIS HCl at pH 7.4.

[433] ADAM-TS1 was assayed at 5 μM enzyme concentration in the manufacturer recommended buffer: 50 mM TRIS, 10 mM CaCl₂, 150 mM NaCl at pH 7.5.

28.2 Samples

[434] Conditioned media and cell lysate samples were received from Charles River Laboratories for the murine colon cancer cell lines: MC38 (epithelial), CT26 (fibroblast), and CT26 transduced

with a lentivirus for doxycyclin-inducible expression of MMP9 (CT26 pLVX MMP9-5T4 +dox). An immortalized mouse intestinal myofibroblast cell line (ABM Good #T0565) was used as a control stromal cell line, grown under manufacturer recommended conditions. Conditioned media and cell lysate samples were also received for screening purposes from ABM Good for the immortalized mouse colonic epithelial cell line YAMC (ABM Good # T0567).

[435] Conditioned media samples were prepared according to the standard protocol using logarithmic phase cells. Briefly, cells were grown for 16h in either serum-free media or in complete media containing 10% fetal bovine serum (FBS). After conditioned media collection, the adherent cells were washed then lysed on plate with non-denaturing lysis buffer to produce a lysed “cell pellet” to capture any activity that remained associated with the cells. All cell culture-derived conditioned media samples, cell lysates from Charles River Laboratories, and cell lysates derived internally were processed with identical methods, to allow comparison between samples.

[436] The resulting conditioned media was buffer exchanged with PBS and concentrated to 10X the original titer as a stock solution.

28.3 Results

[437] Method development experiments with these samples showed that conditioned media containing fetal bovine serum (FBS) were amenable to FRET screening, although a small amount of background fluorescence for each substrate was obtained in the presence of FBS. FBS background controls were therefore used for baseline subtraction at each concentration of substrate throughout the FRET experiments.

[438] In the end-point screening assays, a fixed titer of 10x concentrated sample and a substrate concentration of 10 μ M was used. Values reported in the end-point experiments are initial velocity measurements, in relative fluorescence units (RFU) per second. End-point measurements are a starting point for a screen, but the rate of the reaction is non-linearly related to substrate concentration. Therefore, a more accurate representation of activity would be given by steady-state kinetic measurements covering a range of substrate concentrations. To compare activity across cell lines, the data were treated with a Michaelis-Menten model, and the concentration of equivalent enzyme units $[E_0]$ was solved.

[439] Accordingly, initial velocity was measured as a function of substrate concentration, using a 2-fold dilution series from 250 to 1 μ M. Data were fitted using non-linear least squares fitting with GraphPad Prism software v 8.0 to the Michaelis-Menten equation:

$$velocity = \frac{k_{cat} \cdot [E_0] \cdot [S]}{(K_M + [S])}$$

where k_{cat} is the rate of product formation under steady-state conditions in units (s^{-1}), and K_M is the Michaelis constant that gives the substrate concentration in molar units (M) at which half

maximum velocity of the reaction is produced for a given enzyme, $[E_0]$ is the enzyme concentration, and $[S]$ the substrate concentration in (M).

[440] In the assays where purified recombinant enzyme is used, the Michaelis-Menten parameters are as classically defined above. For conditioned media samples, the effective titer of conditioned media per assay was adjusted to approximately a 2x concentrate of the original unprocessed media samples to assure that the detection range matched the signal produced. The enzyme component in the calculation is redefined as a mixture of possible proteases that contribute to the cumulative observed activity, each with intrinsic reactivity toward a substrate. In this case, it is more appropriate to consider k_{cat} and K_M as macroscopic constants that represent the overall efficiency of catalysis, and E_0 is an enzyme equivalent, expressed in units of concentration. To solve the data fit, k_{cat}/K_M can be fixed at that of the recombinant enzyme and enzyme equivalents per cell culture volume can be output. The effective titer of enzyme equivalents in this case can be used to compare the activity produced by the different cell lines.

[441] The ratios of enzyme equivalents between tumor and control cell lines may be a contributing factor to therapeutic index in a pro-drug employing protease-cleavable linkers.

[442] For all six motif substrates, greater activity was detected in the conditioned media from the tumor cell lines than from the control myofibroblast cell line (FIG. 36).

[443] The ADAM17 substrate with the sequence LAQKLKSS (ADAM17_2) (SEQ ID NO: 201), based on the results in the end-point screen experiments, had approximately 3-fold greater activity in the conditioned media produced from the three tumor cell lines, compared to the myofibroblast cell line (FIG. 36, black bars). For reference, the activity of recombinant ADAM17 at 12 nM is shown in the same assay format. The steady state kinetic curves for ADAM17 substrate are shown in FIG. 37. There was essentially undetectable ADAM17_2 background activity in the myofibroblast sample.

[444] The CTSL1 substrate, ALFKSSFP (SEQ ID NO: 198) (CTSL1_1), based on the results in the end-point screen experiments, had undetectable activity in the control myofibroblast cell line, as well had low activity in the MC38 and CT26 parental cell lines, and had undetectable activity in the CT26-MMP9+ cell line (FIG. 36). This assay was benchmarked with CTSL1 at 1.5 nM, showing that the extracellular concentration of secreted CTSL1 activity was below this effective concentration across the four cell lines. Re-analysis of conditioned media with the CTSL1_1 substrate using the steady-state kinetics analysis yielded no measurable activity above background.

[445] The secreted activity for FAP α , based on the results in the end-point screen experiments, was very low, but was still detected in all three tumor cell lines (FIG. 36, blue bars). The FAP α _1 activity was benchmarked with 5 nM FAP α enzyme. This enzyme is also found in a plasma membrane-associated form, therefore FAP α activity was also tested in the cell pellet fraction that was collected after the media conditioning procedure. These results will be discussed below.

[446] As observed in the end-point screen, re-analysis of the conditioned media for activity toward the FAP α _1 substrate showed lower activity than that associated with the cell pellet, measured in cell lysates (FIGs. 38 and FIG. 39). Myofibroblasts had no measurable secreted activity but did have cell-associated activity (FIG. 38 and FIG. 39).

[447] The MMP substrates MMP9_1 and MMP14_1 all had high activity in the end point screen, and the activity was significantly higher in the tumor cell lines than in the myofibroblast control cells. MMP9_1 (second column from right, FIG. 38) had approximately 4-fold greater activity in the tumor lines than the myofibroblast control; and MMP14_1 had approximately 57-fold greater activity in the tumor lines. The MMP14_1 substrate had the lowest activity in the myofibroblast cells, thus contributing to the higher ratio of tumor vs control in this assay. Secreted activity was measured for the MMP9_1 substrate in the conditioned media for each of the tumor cell lines using the steady-state kinetics analysis (FIG. 36). The myofibroblast cell line had an artefact in analysis of MMP9_1 secreted activity (FIG. 36); this will be addressed with a repeat analysis at higher enzyme titer. As with the MMP9 substrates, the MMP14_1 substrate had significant measurable activity in the conditioned media from the tumor cells but not in the myofibroblast cell line.

[448] Taken together, the results of the FRET screens of conditioned media demonstrated tumor-specific activity for the enzymes ADAM17, CTSL1, MMP9 and MMP14. Soluble FAP-alpha activity was low.

Example 29: Tumor-specific activity toward the FAP α substrate in the cell lysates

[449] To test for cell pellet-associated FAP α activity, the cell lysates were clarified by ultracentrifugation and then assayed neat in the FRET assay. As shown in FIG. 41, FAP α _1 cleavage activity was detected in all four cell lines (blue bars, right hand bar of each pair). A modest 2- fold greater activity was detected in the three tumor cell lines over the myofibroblast cell line. For comparison, CTSL1 activity was also tested in the cell lysates. CTSL1 is a lysosomal enzyme; therefore, activity toward CTSL1_1 was expected to be similarly abundant in all four cell lines. The ratio of CTSL1_1 cleavage in tumor vs control cell lines in this screening format was ~1x (grey bars, left-hand bar of each pair, FIG. 41). FAP α (fibroblast activating protein-alpha) is a marker for fibroblast (CT26) and myofibroblast cell lines. Therefore, an additional non-fibroblast, epithelial cell line was obtained as an alternate negative control cell line, comparable to the MC38 epithelial cell line.

Example 30: Cell-associated activity toward CTSL1 substrate

[450] Cell-associated activity toward CTSL1 substrate was detected in all of the cell lysates, indicating cell-associated activity for this enzyme. The activity was significantly greater in MC38 cells over the myofibroblast, CT26 parental, and CT26 – MMP9+ cells (FIG. 42).

Example 31: Secreted activity in conditioned media from immortalized myofibroblast cell line

[451] Low background-corrected fluorescence was observed in all the myofibroblast reactions. Thus, it can be concluded that the immortalized myofibroblast cell line had low conditioned media activity.

Example 32: Screening of an additional control "normal" epithelial cell line

[452] An immortalized mouse colonic epithelial cell line (YAMC) was used to test activity in the presence of additional control "normal" epithelial cells. YAMC's are given the calculated enzyme equivalents (nM) resulting from all of these steady state experiments, to allow for comparison of the amount of equivalent activity per sample. For example, the YAMC control epithelial cell line produced activity with the ADAM17_2 substrate equivalent to that of 0.52 ± 0.03 nM of ADAM17 recombinant enzyme. This cell line compares directly to MC38, also of epithelial origin, which had activity equivalent to that of 0.32 ± 0.03 nM of ADAM17 recombinant enzyme. The results of the screening show that the YAMC cell line generally had more background activity than did the myofibroblast cell line with all of the substrates. This cell line required non-standard cell culture conditions, including growth with murine IFN γ , and at low temperature to maintain, and may not be fully representative of epithelial cell lines.

Example 33: Cleavage of CTSL1_2 motif in the context of an 8-mer FRET peptide and in the context of an extended tandem linker.

[453] Additional in vitro kinetic experiments were performed to characterize a new candidate motif for CTSL1 enzyme; the CTSL1_2 motif has the sequence ALFFSSPP (SEQ ID NO: 199). CTSL1 cleaves the FRET substrate bearing the CTSL1_1 motif highly efficiently (FIG. 46). Yet, initial experiments with the CTSL1_2 FRET substrate were inconclusive, possibly due to artifacts in the FRET assay; therefore, a new FRET peptide construct was designed with the 9-mer sequence ALFFSSPPS (SEQ ID NO: 236). This 9-mer FRET substrate bearing the CTSL1_2 motif produced comparable catalytic efficiency to the CTSL1_1 motif.

[454] All of the FRET substrates used in these studies have a methoxycoumarin (Mca) fluorophore and dinitrophenyl-lysyl (Dnp) quencher pair at the N- and C-termini of the sequences, respectively. The motif ALFFSSPP (SEQ ID NO: 199) was efficiently cleaved in previous

experiments at ALFFISSPP (SEQ ID NO: 199) (where | indicates the scissile bond) within 14-mer peptide constructs that were used in the MSP-MS tailored library assay. It was cleaved also at the ALFIFSSPP (SEQ ID NO: 199) and ALFFSISPP (SEQ ID NO: 199) sites within 14-mer peptides, indicating that the enzyme recognition in the CTSL1_2 motif may be shifted up- or down-sequence by +/-1 site in the 14-mer peptide experiments. This also suggests that the 8-mer design using Mca-ALFFSSPPK-Dnp (SEQ ID NO: 245) may have unfavorable interactions (steric or otherwise) with the fluor/quencher pair in the shifted P4 or P4' positions that were alleviated by the redesign as a 9-mer substrate using Mca-ALFFSSPPSK-Dnp (SEQ ID NO: 246). This FRET experiment indicated that cleavage of this sequence most likely requires binding recognition at the +/-1 positionally-shifted site and that the fluor or quencher may interfere with this binding.

[455] This hypothesis was tested in the context of the hybrid linker design experiment. Shown in FIG. 46 are the results of cleavage of two matched 30-mer peptides with the following sequences:

(CTSL1_1) SGGPGGPAGIGALFKSSFPLAQKLKSSGGG (SEQ ID NO:207)

(CTSL1_2) KSGPGGPAGIGALFFSSPPLAQKLKSSGGR (SEQ ID NO:219)

[456] Both peptides showed rapid disappearance of substrate (squares) and product formation from cleavage at the target scissile bond (site 0, blue triangles, FIG. 46). Where the peptides differed is in the permissive cleavage at the -1 and +1 sites. CTSL1_1 had a -1 site product at ALFKSSFP (SEQ ID NO: 198) (downward triangles) but CTSL1_2 did not, and the +1 site product was more rapidly formed from the CTSL1_2 substrate (ALFFSISPP (SEQ ID NO: 199)) than with the CTSL1_1 substrate (ALFKSISFP) (SEQ ID NO: 198).

[457] Counter-screening of CTSL1_1 and CTSL1_2 was performed with the enzyme Cathepsin K (CTSK). The substrate CTSL1_2, although lacking a basic P1 residue (ALFFSSPP (SEQ ID NO: 199)) was measurably cleaved by CTSK but at a 15x lower level than the CTSL1_1 substrate (ALFKSSFP (SEQ ID NO: 198)) FIG. 47. For comparison, the specific activity for reference substrates such as Z-LR-AMC and Z-FR-AMC is 15,000-25,000 pmol/min/ μ g, with the average value indicated by a dashed line in FIG. 47.

[458] Thus, these results show that the new CTSL1_2 motif was not cleaved in the context of an 8-mer FRET peptide, but it was cleaved as a 9-mer FRET peptide and it was rapidly cleaved in the context of an extended tandem linker.

Example 34: Tandem linker analysis with mass spectrometric detection

34.1 Substrate Profiling by Mass Spectrometry

[459] To analyze the catalytic efficiency of candidate tandem linker designs, a tailored library approach using MSP-MS was applied. For the tailored library, a set of nineteen synthetic peptides, 30 amino acids in length (30-mers), were designed to test whether sequence context affects the efficiency of cleavage. Peptides were assayed in a multiplex format together as a substrate library

with individual recombinant enzymes, using MSP-MS. For quantitative comparisons, kinetic analysis was performed over a time course, and results are reported either as a catalytic efficiency k_{cat}/K_M for well-behaved first-order kinetics, or as observed rates as k_{obs} for more complex kinetic behavior.

34.2 The library of tandem linker sequences

[460] The library of tandem linker sequences was designed to incorporate three individual protease motifs within the context of a longer, 30-mer peptide sequence. Table 11 lists the sequences and motif arrangement of the 30-mer peptides.

Table 11. 30-mer peptide sequences.

Substrate Name	Sequence	SEQ ID NO:
ALU30-1	GALFKSSFPSSGGGPAQLYAQGGSGKGGSGK	202
ALU30-2	RGSGGGPAQLYAQGGSGGGPAQLYAQGGSGK	203
ALU30-3	KGGGPAQLYAQGGPAQLYAQGGPAQLYAQGSR	204
ALU30-4	RGGPAQLYAQGGPAQLYAQGGGPAQLYAQK	205
ALU30-5	KGGALFKSSFPGGPAGIGPLAQKLKSSGGG	206
ALU30-6	SGGPGGPAGIGALFKSSFPPLAQKLKSSGGG	207
ALU30-7	RGPLAQKLKSSALFKSSFPGGPAGIGGGGK	208
ALU30-8	GGGALFKSSFPPLAQKLKSSPGGPAGIGGGR	209
ALU30-9	RGPGGPAGIGPLAQKLKSSALFKSSFPGGG	210
ALU30-10	RGGPLAQKLKSSPGGPAGIGALFKSSFP GK	211
ALU30-11	RSGGPAQLYAQALFKSSFPPLAQKLKSSGGG	212
ALU30-12	GGPLAQKLKSSALFKSSFPGPAGLYAQGGR	213
ALU30-13	GGALFKSSFPGPAGLYAQPLAQKLKSSGGK	214
ALU30-14	RGGALFKSSFPPLAQKLKSSGPAGLYAQGGK	215
ALU30-15	RGGGPAQLYAQPLAQKLKSSALFKSSFP GG	216
ALU30-16	SGPLAQKLKSSGPAGLYAQALFKSSFP GSK	217
ALU30-17	KGGPGGPAGIGPLAQRLRSSALFKSSFP GR	218
ALU30-18	KSGPGGPAGIGALFPSSPPLAQKLKSSGGR	219
ALU30-19	SGGFPRSGGSFNPRTEGSKKRRQSRGGGG	220

[461] To test for additive effects of having one, two, or three repeat MMP14_1 motifs with small differences in spacing between the motifs, a set of four tandem linkers were designed (ALU30-1 to -4).

[462] To test whether neighboring effects from adjacent sites could alter the catalytic efficiency of a given protease toward its target motif, the three motifs in a set were arranged in all permutations. For example, the combination of motifs A, B, and C could be arranged as: A–B–C, C–A–B, B–C–A, B–A–C, A–C–B, and C–B–A. These permutations were tested for the tandem combinations of MMP14_1, ADAM17_2, and CTSL1_1 as a broad set of motifs susceptible to matrix metalloprotease (MMP) activity (ALU30-11 to -16), and for FAPa_1 with ADAM17_2, and CTSL1_1 as a set with lower MMP sensitivity (ALU30-5 to -10).

[463] Also tested within the library were two alternate substrates for ADAM17 and CTSL1. Two additional peptides were designed, swapping the ADAM17_2 motif with ADAM17_1 (in ALU30-17), or CTSL1_1 with CTSL1_2 (in ALU30-18), demonstrating a strategy for testing further tandem linker designs with new individual protease motifs.

34.3 Counter-screening

[464] Counter-screening was performed against the enzymes Thrombin, Factor Xa and Hepsin using this library as well, and for calibration of activity, a positive-control peptide bearing favorable motifs for these enzymes was also designed (ALU30-19).

[465] Outside of these motifs, minor variations were made to the bracketing sequences at the N- or C-terminus of each tandem linker peptide to generate unique sequences that allow for differentiation of peptides using mass spectrometric detection.

[466] Catalytic efficiency (k_{cat}/K_M) estimations were used to rank the top cleaved substrates in the MSP-MS reaction. Peptides were quantified by mass spectrometric label-free quantitation from the MS1 precursor ion peak areas for each peptide. Enzyme progress curves were modeled from six-time point measurements in GraphPad Prism. Data were fitted using non-linear least squares fitting to the first order kinetics equation:

$$Y = e^{(-\frac{k_{cat}}{K_M}[E_0] \cdot t)}$$

where Y = percent product formation or substrate consumption, and t = time. The observed rate is a function of the enzyme concentration $[E_0]$, and an observed catalytic efficiency (k_{cat}/K_M) in units $M^{-1} s^{-1}$.

34.4 Results

[467] Analyses with the 30-mer tandem linker library showed specific cleavages with each of the recombinant enzymes. In this experiment, the rates of substrate degradation and of product formation were both useful comparisons for understanding the efficiency of linker cleavage.

[468] For example, the substrate degradation traces for the 30-mer library peptides are shown from MMP9-treatment in FIG. 55. The most efficient MMP9 substrates were those that bore three-repeats of the MMP14_1 motif. The next most efficient substrates contained the pair of motifs

where MMP14_1 was followed directly by ADAM17_2. The single or double MMP14_1 motif-bearing peptides were next in the order. An unanticipated result was that the FAPa_1 motif was also cleaved within the peptides bearing FAPa_1 directly followed by the CTSL1_1 motif at PGGPAGIIGALF (SEQ ID NO: 247); these were lower efficiency cleavages. The FAPa_1 motif was not cleaved by MMP9, however, when followed either by the ADAM17_2 motif or the spacing residues GG. The slowest peptides bear the MMP14_1 motif near the C-terminus of the 30-mer peptide. From this experiment, a trend emerged that MMP9 has additional sequence preferences in the downstream prime-side positions, and that the combination of MMP14_1 upstream of ADAM17_2 is most efficient.

[469] To better understand the intrinsic rates of product formation with MMP9 treatment, it is also possible to monitor the cleavage products at a specific peptide bond in these experiments. A comparison of peptides bearing one, two or three MMP14_1 tandem motifs is shown in FIG. 50. The most rapidly degraded peptides were those bearing three repeat motifs of MMP14_1 (top panel). Overall, the peptides bearing one or two MMP14_1 motifs were degraded at approximately the same rate (top panel). However, considering intrinsic rates of bond cleavage at individual sites, the cleavage product of Alu30-2 at bond 9 appeared more rapidly than the cleavage product of Alu30-1 at bond 16 or of Alu30-2 at bond 21 (bottom panel). Thus, the specific cleavage of the Alu30-2 peptide was more efficient than the Alu30-1 peptide. These individual bond cleavage events are monitored by tracking unique peptide fragments, and although too complex for data fitting, the ranking of products is possible.

[470] FAP α was able to cleave both FAP α _1 and MMP14_1 motifs, at PGGPIAGIG (SEQ ID NO: 197) and GPIAGLYAQ (SEQ ID NO: 195) respectively. Degradation of the 30-mer peptides by FAP α showed highest cleavage activity toward the peptides bearing tandem MMP14_1 motifs (FIG. 50, Alu30-4 or Alu30-3). The peptide bearing two MMP14_1 motifs (Alu30-2) was also efficiently cleaved. FAP α _1 motifs were cleaved more efficiently when followed by the ADAM17_1 or ADAM17_2 motifs than when followed by CTSL1_1 or CTSL1_2. Among these peptides, the Lys-bearing ADAM17_2 or CTSL1_1 motifs were lower efficiency than the new motifs tested in this experiment, ADAM17_1 and CTSL1_2.

[471] Another trend was a preference for cleavage of the FAP α _1 motif in the first or second position relative to the N-terminus of these peptides. When the FAP α _1 motif was in the third position closest to the C-terminus, these peptides were not cleaved (Alu30-7 or Alu30-8). The MMP14_1 motif was readily cleaved in the third position closest to the C-terminus, but this was most evident in the tandem MMP14_1 motif-bearing peptides, where N-terminal cleavages assist in shortening the 30-mer. Thus, it may be that FAP α could serve as a secondary-cleaving enzyme after activation with a first enzyme elsewhere in a hybrid linker design.

[472] CTSL1 treatment of the 30-mer peptide library produced cleavages in all peptides bearing a CTSL1_1 or CTSL1_2 motif (FIG. 51). The most efficiently cleaved peptides were those containing the FAPa_1, CTSL1_1 and ADAM17_2 motifs, in multiple orders. Peptides bearing the MMP14_1 motif with CTSL1_1 and ADAM17_2 were slightly lower efficiency. CTSL1 motifs in the middle of the 30-mer peptides were also favorable, thus CTSL1 may be able to serve as a first cleavage in the hybrid motifs.

[473] ADAM17 produced cleavages in all peptides bearing an ADAM17_1 or ADAM17_2 motif as well (FIG. 52). The most efficient peptide cleavages occurred in the peptides containing ADAM17_2, CTSL1_1 and FAPa_1 motifs. Slightly lower efficiency cleavages were obtained in peptides containing the MMP14_1 with ADAM17_2 and CTSL1_1. Peptides bearing only the MMP14_1 motif were not cleaved. Peptides Alu30-9 and Alu30-17, containing the ADAM17_2 and ADAM17_1 motifs respectively, had similar cleavage efficiencies.

[474] Counter-screening with Thrombin, Factor Xa and Hepsin was also performed following the same analysis. The library peptide Alu30-19 was designed to include authentic sites for each enzyme, for comparison of overall cleavage efficiency as well as cleavage site specificity within each motif. The substrate designed for Factor Xa has the motif FNPRITFGS (SEQ ID NO: 248), derived from the Factor Xa cleavage site in Thrombin. The substrate motif for Thrombin was FPR1, a common tool substrate motif. The substrate motif for hepsin was RKRRIGSRG (SEQ ID NO: 249) from filaggrin.

[475] Shown in FIG. 53 are the results of Factor Xa cleavage. The authentic cleavage site motif was Peptide Alu30-19 with this motif was completely degraded before the 5 min time point. Products of the Alu30-19 cleavage were formed from all three sites in this peptide. In comparison, the only library peptide with significant cleavage was Alu30-5, cleaved at the CTSL1_1 motif ALFKISFP (SEQ ID NO: 198), upstream of a FAPa_1 motif.

[476] Otherwise, all other peptides bearing a CTSL1_1 motif were cleaved by Factor Xa at either ALFKISSFP (SEQ ID NO: 198) or ALFKISFP (SEQ ID NO: 198). The ADAM17 motifs were also cleaved at KLKISSGP (SEQ ID NO: 250), KLKSSIALF (SEQ ID NO: 251), or RLRISALF SEQ ID NO: 252), but these cleavages were all lower efficiency. The peptide with the motif CTSL1_2 was cleaved at the lowest efficiency.

[477] Thrombin showed higher activity than Factor Xa, and it cleaved each of the 30-mer peptides. The MMP14_1 motifs were cleaved at GPAGILYAQ (SEQ ID NO: 195), the CTSL1_1 motifs at ALFKISSFP (SEQ ID NO: 198), and the ADAM17_2 motifs at LAQKILKSS (SEQ ID NO: 201) or LAQKLKISS (SEQ ID NO: 201) (or LAQRILRSS (SEQ ID NO: 200) and LAQLRISS (SEQ ID NO: 253) in ADAM17_1). The two alternate ADAM17 motifs were cleaved with similar efficiency (peptides Alu30-17 vs Alu30-9). However, the CTSL1_2 motif had lower cleavage activity than did the CTSL1_1 motif (peptides Alu30- 18 vs Alu30-6). The peptides

Alu30-1 and Alu30-3 were the most rapidly cleaved in the library, and Alu30-4 was also an efficient substrate, showing the susceptibility of MMP14_1 motifs in this experiment. The peptide Alu30-2, bearing two MMP14_1 motifs, was less susceptible to thrombin, however, potentially due to altered spacing with additional glycine residues between motifs. Also, the peptides Alu30-12, Alu30-13 and Alu30-15, with MMP14_1 motifs in combination with ADAM17_2 and CTSL1_1, were cleaved at lower efficiency by thrombin. Thus, increased spacing between the motifs or specific arrangements may rescue a thrombin-susceptible linker design (FIG. 54).

[478] The favored arrangements for reducing thrombin susceptibility include using ADAM17_2 upstream of CTSL1_1, and CTSL1_1 upstream of FAPa_1.

[479] Finally, hepsin treatment of the 30-mer library produced lower efficiency cleavages overall (FIG. 55). The kinetics of these peptide degradation reactions were complex due to the formation of multiple cleavage products; for example, Alu30-19 was cleaved at all three motifs within the first 15 min of the reaction at RKRR (SEQ ID NO: 254), FPR (SEQ ID NO: 255). The literature value for the catalytic efficiency of a FRET peptide bearing this motif was $3.5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$. The authentic peptide Alu30-19, as well as the top cleaved peptides Alu30-8, -9, -12, and -14 all had apparent catalytic efficiencies on the same order of $10^5 \text{ M}^{-1}\text{s}^{-1}$. The features that made peptides more susceptible to hepsin appear to be placement of an ADAM17_2 motif in the second motif position of the peptide, or in the combination of ADAM17_2 followed by CTSL1_1. In general, the library peptides were all cleaved at P1 Lys sites, QKLK (SEQ ID NO: 256) or ALFK (SEQ ID NO: 257). The ADAM17_1 motif was less efficiently cleaved than ADAM17_2, and the alternate CTSL1_2 motif was also cleaved at a much slower rate than CTSL1_1.

[480] To conclude, the tandem linker analysis with mass spectrometric detection revealed motif order preferences and unexpected side-reactions.

Example 35: 30-mer tandem linker design

[481] To generate the most efficient tandem linker, the cleavage efficiencies for each of the targeted enzymes toward the 30-mer library peptides can be compared (FIG. 64).

[482] The tandem linker with the arrangement of ADAM17_2 – MMP14_1 – CTSL1_1 in the peptide Alu30-16 had generally high activity toward the full set of five targeted enzymes, as well as the lowest susceptibility to thrombin, Factor Xa, and hepsin. The next best configuration was MMP14_1 – ADAM17_2 – CTSL1_1 in Alu30-15 which, although it had slightly higher hepsin susceptibility and lower CTSL1 activity may be rescued by the replacement with the CTSL1_2 motif. Replacement of the ADAM17_2 motif with ADAM17_1 may also enhance activity toward FAP-alpha and CTSL1, even if ADAM17 activity enhancement is minor, and hepsin susceptibility would be predicted to be reduced.

Example 36. Stability in Serum and Plasma

[483] The stability of constructs of interest was measured in human or mouse serum. Each construct (approximately 30mg/mL) was combined with serum (1:9 ratio of construct to serum), MMP9, or PBS. The mixture was incubated at 37°C for 24 or 72 hours. Samples were taken at T=0 hours for comparison. After incubation, samples were diluted 1:500 (in human serum, FIG 56) or 1:200 (in mouse serum, FIG 57) in PBS and run on an SDS PAGE gel for western blot analysis. A polyclonal anti-IL-2 antibody (R&D Systems) was used to probe the blots. Results are shown in FIG 56 and FIG 57.

Example 37. IL-2 Serum Stability

[484] The stability of IL2 fusion proteins in human serum (normal and cancer patient) was measured using capillary electrophoresis-based immunoassays (Jess instrument, Protein Simple). Fusion proteins were concentrated to 10mg/ml and incubated with serum (90%). A time zero sample was immediately stored on ice while 24 hour and 72 hour samples were placed at 37°C. Post incubation, samples were diluted 1:1000 with 0.1X sample buffer (Protein Simple) and loaded on the Jess cartridge per the manufacturer's protocol. The primary antibody was a monoclonal human IL2 antibody (R&D Systems, cat# AF-202-NA, stock concentration 0.2mg/ml, working concentration 1:100), and the secondary antibody was a peroxidase-conjugated AffiniPure Bovine Anti Goat IgG (H+L) (Jackson Immuno Research, cat# 805-035-18, reconstituted at 0.8mg/ml, working concentration 1:5000 dilution). All antibodies were diluted in milk free diluent (Protein Simple). Quantitation of IL2 containing species was quantitated using Compass software (Protein Simple) to determine the extent of cleavage (i.e. amount of IL2 containing species smaller than the input fusion protein). Results are shown in FIGs. 24A-24B.

Example 38: Tissue Stability

[485] Primary human lung epithelial cells and renal epithelial cells were obtained from ATCC. Primary hepatocytes were obtained from Lonza and Sigma. Cells were thawed, counted, and plated at 1e4 cells per well in a 96 well round bottom plate in their respective growth medias. Polypeptides containing recombinant human IL-2 and the sequence for Linker-2 and Linker-3 were incubated with cells or media alone for 24 and 72 hours at a concentration 5µg/mL. Cell culture supernatants were collected, and cells were discarded. Protein cleavage was measured by western blot for IL-2 using the protein simple JESS system and Compass software. Results are shown in FIG. 5.

[486] Primary human lung fibroblasts were obtained from ATCC. Prior to the processing assay, human PBMCs were stimulated with 5ug/mL PHA for 72 hours to generate T blasts, which were then frozen and used to measure polypeptide processing. Primary human fibroblasts were thawed, counted, and plated at 1e5 cells per well in a 96 well round bottom plate in X-Vivo 15 media. Polypeptides containing recombinant human IL-2 and the sequence for Linker-1 (GPAGMKGL, SEQ ID NO: 196), Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198), or a non-cleavable sequence were incubated with or without fibroblasts cells for 48 hours at a concentration of either A) 3.3nM or B) 0.33nM. As a positive control, some wells were also incubated with a polypeptide that was pre-cut in vitro overnight with 1µg pre-activated MMP9 enzyme at 37°C. After 48 hours, cell culture supernatants were collected and healthy fibroblasts were discarded. Polypeptide processing was measured either by A) western blot for IL-2 using the protein simple JESS system, or by B) measuring the capacity of the cell culture supernatants to stimulate T blast proliferation. Briefly, T blasts were incubated for 72 hours with cell culture supernatants containing polypeptides previously exposed to healthy human fibroblasts, before T blast proliferation was measured by Cell Titer Glow analysis. Results are shown in FIG. 6A-6B.

Example 39: Linker-2 is efficiently cleaved in human tumor cells

[487] Cleavage efficiency of Linker-2 (GPAGLYAQ, SEQ ID NO: 195) was evaluated in tumor tissue derived from human samples. A total of 66 samples were analyzed from seven solid tumor types. The tumor types are shown in Table 5 below.

[488] Briefly, frozen dissociated tumor cells (DTC) were thawed, counted and plated in X-Vivo media. Cells or “No Cell Control” wells were incubated with an inducible IL-2 cytokine containing Linker 1 (a cleavable linker that was not designed using the processes described herein) or Linker-2 (GPAGLYAQ, SEQ ID NO: 195). These inducible IL-2 proteins have no or minimal IL-2 biological activity when the linker is intact, and IL-2 biological activity is induced when the linker is cleaved. Cell culture supernatants were collected and frozen before analysis for IL-2 biological activity using T-Blast proliferation and Jess protein assays.

The minimum fold change in IL-2 activity is 1.00, and the maximum is 2.50. A 1.3 fold change is indicative of a significant increase in biological activity relative to the uncleaved control. Table 5 shows the average fold change for Linker-2 (GPAGLYAQ, SEQ ID NO: 195) in comparison to an uncleaved control, a cleaved control, and Linker-1. In Table 12 the symbol “-” indicate essentially no increase in activity; “+/-” indicates some increase in activity but not significant; “+,” “++,” and “+++” indicate the relative significant increase in activity compared to the uncleavable control. Linker-2 is sufficiently cleaved in human tumor types to induce IL-2 biological activity.

Table 12. Cleavage efficiency of Linker-2 in human tumor types

Diagnosis	Average Fold Change Over Non-Cleavable			
	Uncleavable	Linker 1	Linker 2	Pre-cut
Melanoma (n=8)	-	-	+/-	+++
Kidney Cancer, Renal Cell Carcinoma (n=11)	-	-	+	+++
Head and Neck Cancer, Squamous Cell (n=6)	-	+/-	+++	+++
Colorectal Cancer, Adenocarcinoma (n = 10)	-	+/-	+	+++
Lung Cancer, Squamous Cell Carcinoma (n=13)	-	-	++	+++
Lung Cancer, Adenocarcinoma (n=10)	-	-	+	+++
Breast Cancer (n = 5)	-	-	+/-	+++

Example 40: Linker-3 is efficiently cleaved in human tumor cells

[489] Cleavage efficiency of Linker-3 (ALFKSSFP, SEQ ID NO: 198) was evaluated in tumor tissue derived from human samples. A total of 66 samples were analyzed from seven solid tumor types. The tumor types are shown in Table 6 below.

[490] Briefly, frozen DTC cells were thawed, counted and plated in X-Vivo media. Cells or “No Cell Control” wells were incubated with an inducible IL-2 cytokine containing Linker 1 (a cleavable linker that was not designed using the processes described herein) or Linker-2 (GPAGLYAQ, SEQ ID NO: 195). These inducible IL-2 proteins have no or minimal IL-2 biological activity when the linker is intact, and IL-2 biological activity is induced when the linker is cleaved. Cell culture supernatants were collected and frozen before analysis for IL-2 biological activity using T-Blast proliferation and Jess protein assays.

The minimum fold change in IL-2 activity is 1.00, and the maximum is 2.50. A 1.3 fold change is indicative of a significant increase in biological activity relative to the uncleaved control. Table 6 shows the average fold change for Linker-3 (ALFKSSFP, SEQ ID NO: 198) in comparison to an uncleaved control, a cleaved control, and Linker 1. In Table 13 the symbol “-” indicate essentially no increase in activity; “+/-” indicates some increase in activity but not significant; “+,” “++,” and “+++” indicate the relative significant increase in activity compared to the uncleavable control. Linker-3 (ALFKSSFP, SEQ ID NO: 198) is sufficiently cleaved in human tumor types to induce IL-2 biological activity.

Table 13. Cleavage efficiency of Linker-3 in human tumor types

Diagnosis	Average Fold Change Over Non-Cleavable			
	Uncleavable	Linker 1	Linker 3	Pre-cut
Melanoma (n=8)	-	-	+	+++
Kidney Cancer, Renal Cell Carcinoma (n=11)	-	-	+	+++
Head and Neck Cancer, Squamous Cell (n=6)	-	+/-	+++	+++
Colorectal Cancer, Adenocarcinoma (n = 10)	-	+/-	+++	+++
Lung Cancer, Squamous Cell Carcinoma (n=13)	-	-	+++	+++
Lung Cancer, Adenocarcinoma (n=10)	-	-	+++	+++
Breast Cancer (n = 5)	-	-	+	+++

Example 41: Measuring Inducibility of Agonist Anti-4-1BB Antibodies

[491] A stable HT-1080 cell line expressing human 4-1BB was established. Agonism of human-4-1BB in these cells resulted in increased secretion of IL-8. Tetravalent monospecific antibodies, inducible format tetravalent monospecific antibodies (protease cleaved or uncleaved), or trimeric ligands were tested for ability to agonize 4-1BB.

[492] Before addition to the cultured cells, some samples of the antibody to be tested were incubated with an appropriate protease under suitable conditions for proteolysis. Some samples were maintained in an uncleaved state. The extent of inducibility was determined by comparison of uncleaved (uninduced) inducible format tetravalent monospecific antibody with the corresponding protease cleaved inducible format tetravalent monospecific antibody.

[493] After incubation at 37°C and 5% CO₂ for 6 hours, the agonistic activities of the cleaved and uncleaved antibodies were evaluated by the quantification of IL-8 production using an IL-8 AlphaLISA or ELISA. The EC₅₀s and maximum IL-8 levels were compared. Results are shown in FIG. 65A-65B.

Example 42: Protease Cleavage of Anti-4-1BB Antibodies by MMP9

[494] One of skill in the art would be familiar with methods of setting up protein cleavage assay. 100 µg of protein in 1xPBS pH 7.4 were cleaved with 1 µg active MMP9 (Sigma catalog # SAE0078-50 or Enzo catalog BML-SE360) and incubated at room temperature for up to 16 hours.

Digested protein is subsequently used in functional assays or stored at -80°C prior to testing. The extent of cleavage was monitored by SDS PAGE using methods well known in the art.

Example 43: Protease Cleavage of Anti-4-1BB Antibodies by MMP14 or CTSL1

[495] One of skill in the art can be replaced with Linker-3. The art would be familiar with methods of setting up protein cleavage assay. 100 µg of protein in 1xPBS pH 7.4 are cleaved with 1 µg active MMP14 (R&D Systems Catalog # 9518-MP-010) or CTSL1 (R&D Systems Catalog #952-CY) and incubated at room temperature for up to 16 hours. Digested protein are subsequently used in functional assays or stored at -80°C prior to testing. The extent of cleavage is monitored by SDS PAGE using methods well known in the art.

8. OTHER EMBODIMENTS

[496] The disclosure set forth above may encompass multiple distinct inventions with independent utility. Although each of these inventions has been disclosed in its preferred form(s), the specific embodiments thereof as disclosed and illustrated herein are not to be considered in a limiting sense, because numerous variations are possible. The subject matter of the inventions includes all novel and nonobvious combinations and subcombinations of the various elements, features, functions, and/or properties disclosed herein. The following claims particularly point out certain combinations and subcombinations regarded as novel and nonobvious. Inventions embodied in other combinations and subcombinations of features, functions, elements, and/or properties may be claimed in this application, in applications claiming priority from this application, or in related applications. Such claims, whether directed to a different invention or to the same invention, and whether broader, narrower, equal, or different in scope in comparison to the original claims, also are regarded as included within the subject matter of the inventions of the present disclosure.

[497] Exemplary polypeptide constructs are detailed herein in Appendix A. While the exemplary polypeptides that contain Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) or other cleavable linkers are disclosed in Appendix A, for each construct, the disclosed linker can be replaced with either Linker-2 (GPAGLYAQ, SEQ ID NO: 195), or Linker-3 (ALFKSSFP, SEQ ID NO: 198) or other cleavable linkers disclosed herein. For example construct

ACP355 (IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX

Blocker_ (Blocker=VHVL.F2.high.A02_VhVI_A46S;X=MMP14-1) can contain Linker-2,

however it can be replaced with Linker-3. Alternatively, construct ACPT464 (IgG4_Fc(S228P)-X-IL2-LX-blocker_ (Blocker=VHVL.F2.high.A02_VhVI_VH105-VL43_disulfidel;X=CTSL1-1) can contain Linker-3, however, it can be replaced with Linker-2.

[498] The elements of the polypeptide constructs provided in Appendix A contain the abbreviations as follows: “L,” “X,” “LX,” and “XL” each refer to a linker. “X” refers to a cleavable linker. “L” refers a linker that is optionally cleavable. When L is the only linker in a polypeptide, L is cleavable. “LX” or “XL” each refer to a cleavable linker with an extended non-cleavable sequence adjacent to it. Cleav. Lin. Also refers to a cleavable linker. Other abbreviations used include: “mIFN γ ” for mouse interferon gamma (IFN γ); (5) “hAlbumin” for human serum albumin (HSA); and (6) “mAlbumin” indicates mouse serum albumin.

APPENDIX A: CONSTRUCT PERMUTATION TABLE

Construct Name	Construct Description
ACP01	(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP02	(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP03	(anti-HSA)-(cleav. link.)-mouse IFN γ -mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP50	(anti-EpCAM)-(anti-HSA)-(cleav. link.)-mouse IFN γ -mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP51	(anti-EpCAM)-Linker-(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP52	(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-Linker-(anti-EpCAM)-6xHis
ACP53	mAlbumin-(cleav. link.)-mIFN γ -(cleav. link.)-mAlbumin-6xHis
ACP54	mAlbumin-(cleav. link.)-mIFN γ -Linker-mIFN γ -(cleav. link.)-mAlbumin-6xHis
ACP30	(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP55	(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-6xHis-C tag
ACP56	(anti-FOLR1)-Linker-(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP57	(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-Linker-(anti-FOLR1)-6xHis
ACP58	(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-Linker-(anti-EpCAM)-6xHis
ACP59	(anti-FOLR1)-Linker-(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-6xHis
ACP60	(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-Linker-(anti-FOLR1)-6xHis
ACP61	(anti-HSA)-(cleav. link.)-mIFN γ -(cleav. link.)-mIFN γ -(cleav. link.)-(anti-HSA)-Linker-FN(CGS-2)-6xHis
ACP63	anti-FN CGS-2 scFv (Vh/Vl)-6xHis
ACP69	(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)-(cleav. link.)-mouse IFN γ
ACP70	mouse IFN γ -(cleav. link.)-(anti-HSA)-(cleav. link.)-mouse IFN γ -(cleav. link.)-(anti-HSA)
ACP71	mouse IFN γ -(cleav. link.)-mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-

Construct Name	Construct Description
	mAlbumin
ACP72	mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-mAlbumin-(cleav. link.)-mouse IFN γ
ACP73	mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-mAlbumin
ACP74	mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-5mer linker-mAlbumin-5mer linker-(cleav. link.)-mouse IFN γ -(cleav. link.)-mAlbumin
ACP75	mAlbumin-(cleav. link.)-mouse IFN γ -(cleav. link.)-10mer linker-mAlbumin-10mer linker-(cleav. link.)-mouse IFN γ -(cleav. link.)-mAlbumin
ACP78	(anti-HSA)-Linker-mouse IFN γ -Linker-(anti-HSA)-Linker-mouse IFN γ -Linker-(anti-HSA)-(non-cleavable_control)
ACP134	Anti-HSA-X-mouse IFN γ -X-anti-HSA-X-mouse IFN γ -X-anti-HSA-L-anti-FOLR1
ACP135	Anti-FOLR1-L-HSA-X-mouse IFN γ -X-HSA-X-mouse IFN γ -X-HSA
ACP04	human p40-murine p35-6xHis
ACP05	human p40-human p35-6xHis
ACP34	mouse p35-(Cleavable Linker)-mouse p40-6xHis
ACP35	mouse p35-GS-(Cleavable Linker)-GS-mouse p40-6xHis
ACP36	(anti-HSA)-(Cleav. Linker)-mouse p40-mouse p35-(Cleav. Linker)-(anti-HSA)-6xHis
ACP37	(anti-EpCAM)-(anti-HSA)-(Cleav. Linker)-mouse p40-mouse p35-(Cleav. Linker)-(anti-HSA)-6xHis
ACP79	(anti-EpCAM)-Linker-(anti-HSA)-(Cleavable Linker)-mIL12-(Cleavable Linker)-(Anti-HSA)-6xHis
ACP80	(anti-HSA)-(Cleavable Linker)-mIL12-(Cleavable Linker)-(anti-HSA)-Linker-(anti-EpCAM)-6xHis
ACP06	Blocker12-Linker-(Cleavable Linker)-human p40-Linker-mouse p35-(Cleavable Linker)-(anti-HSA)-6xHis
ACP07	Blocker12-Linker-(Cleavable Linker)-human p40-Linker-mouse p35-(Cleavable Linker)-(anti-HSA)-Linker-(anti-FOLR1)-6xHis
ACP08	(anti-FOLR1)-Linker-Blocker12-Linker-(Cleavable Linker)-human p40-Linker-mouse p35-(Cleavable Linker)-(anti-HSA)-6xHis
ACP09	(anti-HSA)-Linker-Blocker12-Linker-(Cleavable Linker)-human p40-Linker-mouse p35-6xHis

Construct Name	Construct Description
ACP10	(anti-HSA)-(Cleavable Linker)-human p40-L-mouse p35-(Cleavable Linker)-Linker-Blocker12-6xHis
ACP11	hp40-Linker-mp35-(Cleavable Linker)-Linker-Blocker12-Linker-(anti-HSA)-6xHis
ACP91	human_p40-Linker-mouse_p35-Linker-Linker-Blocker-Linker-(anti-HSA)_(non-cleavable_control)
ACP136	human p40-L-mouse p35-XL-Blocker
ACP138	human_p40-L-mouse_p35-XL-Blocker-L-HSA-L-FOLR1
ACP139	FOLR1-L-human_p40-L-mouse_p35-XL-Blocker-L-HSA
ACP140	FOLR1-X-human_p40-L-mouse_p35-XL-Blocker-L-HSA
ACP12	(anti-EpCAM)-IL2-(cleav. link.)-(anti-HSA)-blocker-6xHis
ACP13	(anti-EpCAM)-Blocker2-(anti-HSA)-(cleav. link.)-IL2-6xHis
ACP14	Blocker2-Linker-(cleav. link.)-IL2-(cleav. link.)-(anti-HSA)-6xHis
ACP15	Blocker2-Linker-(anti-HSA)-Linker-(cleav. link.)-IL2-6xHis
ACP16	IL2-(cleav. link.)-(anti-HSA)-Linker-(cleav. link.)-Blocker2-6xHis
ACP17	(anti-EpCAM)-Linker-IL2-(cleav. link.)-(anti-HSA)-Linker-(cleav. link.)-Blocker2-6xHis
ACP18	(anti-EpCAM)-Linker-IL2-(cleav. link.)-(anti-HSA)-Linker-vh(cleav. link.)v1-6xHis
ACP19	IL2-(cleav. link.)-Linker-Blocker2-Linker-(anti-HSA)-Linker-(anti-EpCAM)-6xHis
ACP20	IL2-(cleav. link.)-Blocker2-6xHis
ACP21	IL2-(cleav. link.)-Linker-Blocker2-6xHis
ACP22	IL2-(cleav. link.)-Linker-blocker-(cleav. link.)-(anti-HSA)-Linker-(anti-EpCAM)-6xHis
ACP23	(anti-FOLR1)-(cleav. link.)-Blocker2-Linker-(cleav. link.)-(anti-HSA)-(cleav. link.)-IL2-6xHis
ACP24	(Blocker2)-(cleav. link.)-(IL2)-6xHis
ACP25	Blocker2-Linker-(cleav. link.)-IL2-6xHis
ACP26	(anti-EpCAM)-Linker-IL2-(cleav. link.)-(anti-HSA)-Linker-blocker(NARA1 Vh/Vl)
ACP27	(anti-EpCAM)-Linker-IL2-(cleav. link.)-(anti-HSA)-Linker-blocker(NARA1 Vl/Vh)
ACP28	IL2-(cleav. link.)-Linker-Blocker2-(NARA1 Vh/Vl)-Linker-(anti-HSA)-Linker-(anti-EpCAM)
ACP29	IL2-(cleav. link.)-Linker-Blocker2-(NARA1 Vl/Vh)-Linker-(anti-HSA)-Linker-(anti-EpCAM)
ACP38	IL2-(cleav. link.)-blocker-(anti-HSA)-(anti-EpCAM)-6xHis

Construct Name	Construct Description
ACP39	(anti-EpCAM)-(cleav. link.)-(anti-HSA)-(cleav. link.)-Blocker2-(cleav. link.)-IL2-6xHis
ACP40	CD25ecd-Linker-(cleav. link.)-IL2-6xHis
ACP41	IL2-(cleav. link.)-Linker-CD25ecd-6xHis
ACP42	(anti-HSA)-Linker-CD25ecd-Linker-(cleav. link.)-IL2-6xHis
ACP43	IL2-(cleav. link.)-Linker-CD25ecd-Linker-(anti-HSA)-6xHis
ACP44	IL2-(cleav. link.)-Linker-CD25ecd-(cleav. link.)-(anti-HSA)-6xHis
ACP45	(anti-HSA)-(cleav. link.)-Blocker2-Linker-(cleav. link.)-IL2-6xHis
ACP46	IL2-(cleav. link.)-linkerL-vh(cleav. link.)vI-Linker-(anti-HSA)-L-(anti-EpCAM)-6xHis
ACP47	(anti-EpCAM)-Linker-IL2-(Cleavable Linker)-(anti-HSA)-Linker-Blocker2-6xHis
ACP48	IL2-(cleav. link.)-Blocker2-Linker-(anti-HSA)-6xHis
ACP49	IL2-(cleav. link.)-Linker-Blocker2-Linker-(anti-HSA)-6xHis
ACP92	(anti-HSA)-(16mer Cleav. Link.)-IL2-(16mer Cleav. Link.)-(anti-HSA)-6xHis
ACP93	(anti-EpCAM)-(anti-HSA)-(anti-EpCAM)-Blocker2-(cleav. link.)-IL2-6xHis
ACP94	(anti-EpCAM)-(anti-HSA)-Blocker2-(cleav. link.)-IL2-6xHis
ACP95	(anti-EpCAM)-(anti-HSA)-(cleav. link.)-IL2-6xHis
ACP96	(anti-EpCAM)-(16mer cleav. link.)-IL2-(16mer cleav. link.)-(anti-HSA)
ACP38	IL2-(cleav. link.)-blocker-(anti-HSA)-(anti-EpCAM)-6xHis
ACP97	(anti-EpCAM)-(anti-HSA)-(cleav. link.)-IL2-(cleav. link.)-(anti-HSA)-6xHis
ACP99	(anti-EpCAM)-Linker-IL2-(cleav. link.)-(anti-HSA)-6xHis
ACP100	(anti-EpCAM)-Linker-IL2-6xHis
ACP101	IL2-(cleav. link.)-(anti-HSA)-6xHis
ACP102	(anti-EpCAM)-(cleav. link.)-IL2-(cleav. link.)-(anti-HSA)-Linker-blocker-6xHis
ACP103	IL2-(cleav. link.)-Linker-Blocker2-Linker-(anti-HSA)-Linker-(anti-FOLR1)-6xHis
ACP104	(anti-FOLR1)-IL2-(cleav. link.)-(anti-HSA)-Linker-Blocker2-6xHis
ACP105	Blocker2-Linker-(cleav. link.)-IL2-(cleav. link.)-(anti-HSA)-Linker-(anti-FOLR1)-6xHis
ACP106	(anti-FOLR1)-Linker-(anti-HSA)-(cleav. link.)-blocker-Linker-(cleav. link.)-IL2 -6xHis
ACP107	Blocker2-Linker-(anti-HSA)-(cleav. link.)-IL2-Linker-(anti-FOLR1)-6xHis
ACP108	(anti-EpCAM)-IL2-(Dually cleav. link.)-(anti-HSA)-Linker-blocker-6xHis
ACP117	anti-FN CGS-2 scFv (Vh/Vl)-6xHis

Construct Name	Construct Description
ACP118	NARA1 Vh/Vl non-cleavable
ACP119	NARA1 Vh/Vl cleavable
ACP120	NARA1 Vl/Vh non-cleavable
ACP121	NARA1 Vl/Vh cleavable
ACP124	IL2-Linker-(anti-HSA)-Linker-Linker-blocker_(non-cleavable_control)
ACP132	IL2-L-HSA
ACP141	IL2-L-hAlb
ACP142	IL2-X-hAlb
ACP144	IL2-X-HSA-LX-blocker-L-FOLR1
ACP145	FOLR1-L-IL2-X-HSA-LX-blocker
ACP146	FOLR1-X-IL2-X-HSA-LX-blocker
ACP133	IL-2-6x His
ACP147	IL2-X-HSA-LX-blocker-L-TAA
ACP148	TAA-L-IL2-X-HSA-LX-blocker
ACP149	TAA-X-IL2-X-HSA-LX-blocker
ACP31	(anti-HSA)-(cleav. link.)-mIFNa1-(cleav. link.)-(anti-HSA)
ACP32	(anti-HSA)-(cleav. link.)-mIFNa1(N+C trunc)-(cleav. link.)-(anti-HSA)
ACP33	(anti-HSA)-(cleav. link.)-mIFNa1(C trunc)-(cleav. link.)-(anti-HSA)
ACP131	mIFNa1
ACP125	HSA-X-mIFNa1
ACP126	mIFNa1-X-HSA
ACP127	mAlb-X-mIFNa1-X-mAlb
ACP128	mAlb-X-mIFNa1
ACP129	mIFNa1-X-mAlb
ACP150	FOLR1-L-HSA-X-mIFNa1-X-HSA
ACP151	FOLR1-L-HSA-X-mIFNa1-X-HSA-L-FOLR1
ACP152	HSA-L-mIFNa1-L-HSA_(non-cleavable_control)
ACP203	HSA-X-mIFNa1-X-HSA_(X=MMP14-1)
ACP204	HSA-X-mIFNa1-X-HSA_(X=CTSL1-1)
ACP205	HSA-X-mIFNa1-X-HSA_(X=ADAM17-2)
ACP206	HSA-X-Human_IFNA2b-X-HSA_(X=MMP14-1)
ACP207	HSA-X-Human_IFNA2b-X-HSA_(X=CTSL1-1)
ACP208	HSA-X-Human_IFNA2b-X-HSA_(X=ADAM17-2)

Construct Name	Construct Description
ACP336	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh-X-VL_A46S;X=MMP14-1)
ACP337	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VL_A46S;X=MMP14-1)
ACP338	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh-X-VL;X=MMP14-1)
ACP339	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VL;X=MMP14-1)
ACP340	IL2-X-anti-HSA-LX-blocker_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP341	IL2-X-anti-HSA-LX-blocker_(Blocker=Hu3TOW85_A;X=MMP14-1)
ACP342	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh-X-VL_A46S;X=MMP14-1)
ACP343	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VL_A46S;X=MMP14-1)
ACP344	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh-X-VL;X=MMP14-1)
ACP345	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VL;X=MMP14-1)
ACP346	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP347	CD25ecd_C213S-LX-IL2-X-anti-HSA-LX-blocker_(Blocker=Hu3TOW85_A;X=MMP14-1)
ACP348	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.A02_Vh-X-VL_A46S;X=MMP14-1)
ACP349	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.A02_Vh/VL_A46S;X=MMP14-1)
ACP350	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh-X-VL;X=MMP14-1)
ACP351	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh/VL;X=MMP14-1)
ACP352	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP353	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=Hu3TOW85_A;X=MMP14-1)
ACP354	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-Blocker_(Blocker=VHVL.F2.high.A02_Vh-X-VL_A46S;X=MMP14-1)
ACP355	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-

Construct Name	Construct Description
	Blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_A46S;X=MMP14-1)
ACP356	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh-X-VI;X=MMP14-1)
ACP357	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh\VI;X=MMP14-1)
ACP358	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-Blocker_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP359	IgG4_Fc(S228P)-X-CD25ecd_C213S-LX-IL2-LX-Blocker_(Blocker=Hu3TOW85_A;X=MMP14-1)
ACP360	MT204_Vh/VI_3xG4S_A46S
ACP361	MT204_Vh-X-VI_X=MMP14-1
ACP362	MT204_Vh-X-VI_X=MMP14-1, A46S
ACP365	VHVL.F2.high.A02_Vh-X-VI_X=MMP14-1
ACP366	VHVL.F2.high.A02_Vh-X-VI_X=MMP14-1, A46S
ACP368	VHVL.F2.high.F03_Vh-X-VI_X=MMP14-1
ACP371	IL2-X-anti-HSA-LX-blocker_(Blocker=MT204_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)
ACP372	IL2-X-anti-HSA-LX-blocker_(Blocker=MT204_Vh/VI_VH105-VL43_disulfide;X=MMP14-1)
ACP373	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)
ACP374	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VI_VH105-VL43_disulfide;X=MMP14-1)
ACP375	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)
ACP376	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VI_VH105-VL43_disulfideX=MMP14-1)
ACP377	IL2-X-anti-HSA-LX-blocker_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP378	IL2-X-anti-HSA-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=MMP14-1)
ACP379	IgG4_Fc(S228P)-X-IL2-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=MMP14-1)
ACP383	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=MT204_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)

Construct Name	Construct Description
ACP384	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=MT204_Vh/Vl_VH105-VL43_disulfide;X=MMP14-1)
ACP385	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_VH44-VL100_disulfide;X=MMP14-1)
ACP386	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_VH105-VL43_disulfide;X=MMP14-1)
ACP387	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/Vl_VH44-VL100_disulfide;X=MMP14-1)
ACP388	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/Vl_VH105-VL43_disulfide;X=MMP14-1)
ACP389	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP390	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46S_VH44-VL100_disulfide;X=MMP14-1)
ACP391	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46S_VH44-VL100_disulfide;X=MMP14-1)
ACP392	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh_G44C_Vl_A46S_G100C;X=MMP14-1)
ACP393	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh_Q105C_Vl_A43C;X=MMP14-1)
ACP394	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh_G44C_Vl_G100C;X=MMP14-1)
ACP395	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh_Q105C_Vl_A43C;X=MMP14-1)
ACP396	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP397	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-blocker_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP398	IL2-XL-CD25ecd_C2I3S-X-HSA-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=MMP14-1)
ACP399	Blocker-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=VHVL.F2.high.A02_Vh_G44C_Vl_A46S_G100C;X=MMP14-1)
ACP400	Blocker-XL-HSA-X-IL2(Nterm-41)-X-

Construct Name	Construct Description
	HSA_(Blocker=VHVL.F2.high.A02_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP401	Blocker-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=VHVL.F2.high.F03_Vh_G44C_VI_G100C;X=MMP14-1)
ACP402	Blocker-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=VHVL.F2.high.F03_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP403	Blocker-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP404	Blocker-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP405	Heavy_Blocker_Fab-XL-HSA-X-IL2(Nterm-41)-X-HSA_(Blocker=MT204_VH-CH1;X=MMP14-1)
ACP406	mIgG1_Fc(S228P)-X-IL2-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=MMP14-1)
ACP407	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)
ACP408	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VI_A46S_VH44-VL100_disulfide;X=MMP14-1)
ACP409	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/VI_VH105-VL43_disulfide;X=MMP14-1)
ACP410	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VI_VH44-VL100_disulfide;X=MMP14-1)
ACP411	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/VI_VH105-VL43_disulfide;X=MMP14-1)
ACP412	mIgG1_Fc(S228P)-X-IL2-LX-blocker_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP415	IL2-XL-blocker-L-CD25_213S-X-HSA_Blocker=VHVL.F2.high.A02_Vh_G44C_VI_A46S_G100C;X=MMP14-1)
ACP416	IL2-XL-blocker-L-CD25_213S-X-HSA_(Blocker=VHVL.F2.high.A02_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP417	IL2-XL-blocker-L-CD25_213S-X-HSA_(Blocker=VHVL.F2.high.F03_Vh_G44C_VI_G100C;X=MMP14-1)
ACP418	IL2-XL-blocker-L-CD25_213S-X-HSA_(Blocker=VHVL.F2.high.F03_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP419	IL2-XL-blocker-L-CD25_213S-X-HSA_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP420	IL2-XL-blocker-L-CD25_213S-X-HSA_(Blocker=Hu2TOW91_B;X=MMP14-1)

Construct Name	Construct Description
ACP421	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=VHVL.F2.high.A02_Vh_G44C_VI_A46S_G100C;X=MMP14-1)
ACP422	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=VHVL.F2.high.A02_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP423	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=VHVL.F2.high.F03_Vh_G44C_VI_G100C;X=MMP14-1)
ACP424	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=VHVL.F2.high.F03_Vh_Q105C_VI_A43C;X=MMP14-1)
ACP425	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=Hu2TOW91_A;X=MMP14-1)
ACP426	HSA-X-blocker-L-CD25_213S-LX-IL2_(Blocker=Hu2TOW91_B;X=MMP14-1)
ACP427	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.A02_Vh_G44C_VI_A46S_G100C, Blocker2=Hu2TOW91_A;X=MMP14-1)
ACP428	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.A02_Vh_Q105C_VI_A43C, Blocker2=Hu2TOW91_A;X=MMP14-1)
ACP429	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.F03_Vh_G44C_VI_G100C, Blocker2=Hu2TOW91_A;X=MMP14-1)
ACP430	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.F03_Vh_Q105C_VI_A43C, Blocker2=Hu2TOW91_A;X=MMP14-1)
ACP431	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.A02_Vh_G44C_VI_A46S_G100C, Blocker2=Hu2TOW91_B;X=MMP14-1)
ACP432	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.A02_Vh_Q105C_VI_A43C, Blocker2=Hu2TOW91_B;X=MMP14-1)
ACP433	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.F03_Vh_G44C_VI_G100C, Blocker2=Hu2TOW91_B;X=MMP14-1)
ACP434	IL2-X-anti-HSA-LX-Blocker1-L-Blocker2_(Blocker1=VHVL.F2.high.F03_Vh_Q105C_VI_A43C,

Construct Name	Construct Description
	Blocker2=Hu2TOW91_B;X=MMP14-1)
ACP439	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl;X=MMP14-1)
ACP440	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl_A46S;X=MMP14-1)
ACP441	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl_A46L;X=MMP14-1)
ACP442	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl_A46S_VH44-VL100_disulfide;X=MMP14-1)
ACP443	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl_A46L_VH44-VL100_disulfide;X=MMP14-1)
ACP444	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.C07_Vh/Vl_VH105-VL43_disulfide;X=MMP14-1)
ACP445	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh-X-Vl_A46L;X=MMP14-1)
ACP446	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46L;X=MMP14-1)
ACP447	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46L_VH44-VL100_disulfide;X=MMP14-1)
ACP451	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46S;X=CTSL1-1)
ACP452	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/Vl;X=CTSL1-1)
ACP453	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_A46S_VH44-VL100_disulfide;X=CTSL1-1)
ACP454	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh/Vl_VH105-VL43_disulfide;X=CTSL1-1)
ACP455	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/Vl_VH44-VL100_disulfide;X=CTSL1-1)
ACP456	IL2-X-anti-HSA-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh/Vl_VH105-VL43_disulfide;X=CTSL1-1)
ACP457	IL2-X-anti-HSA-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=CTSL1-1)
ACP458	IgG4_Fc(S228P)-X-IL2-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=CTSL1-1)
ACP459	IgG4_Fc(S228P)-X-IL2-LX-

Construct Name	Construct Description
	Blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_A46S;X=CTSL1-1)
ACP460	IgG4_Fc(S228P)-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh\VI;X=CTSL1-1)
ACP461	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_A46S_VH44-VL100_disulfide;X=CTSL1-1)
ACP462	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_VH105-VL43_disulfide;X=CTSL1-1)
ACP463	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh\VI_VH44-VL100_disulfide;X=CTSL1-1)
ACP464	IgG4_Fc(S228P)-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh\VI_VH105-VL43_disulfide;X=CTSL1-1)
ACP465	mIgG1_Fc-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_A46S;X=CTSL1-1)
ACP466	mIgG1_Fc-X-IL2-LX-Blocker_(Blocker=VHVL.F2.high.F03_Vh\VI;X=CTSL1-1)
ACP467	mIgG1_Fc-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_A46S_VH44-VL100_disulfide;X=CTSL1-1)
ACP468	mIgG1_Fc-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.A02_Vh\VI_VH105-VL43_disulfide;X=CTSL1-1)
ACP469	mIgG1_Fc-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh\VI_VH44-VL100_disulfide;X=CTSL1-1)
ACP470	mIgG1_Fc-X-IL2-LX-blocker_(Blocker=VHVL.F2.high.F03_Vh\VI_VH105-VL43_disulfide;X=CTSL1-1)
ACP471	mIgG1_Fc-X-IL2-LX-Heavy_blocker_Fab_(Blocker=MT204_VH-CH1;X=CTSL1-1)

APPENDIX B: Sequences

SEQ ID NO.	Name	Sequence
1	Human IL-2	MYRMQLLSCL ALSLALVTNS APTSSSTKKT QLQLEHLLLD LQMILNGINN YKNPKLTRML TFKFYMPKKA TELKHLQCLE EELKPLEEVL NLAQSKNFHL RPRDLISNIN VIVLELKGSE TTFMCEYADE TATIVEFLNR WITFCQSIISTLT
2	Human serum albumin	MKWVTFISLL FLSSAYSRG VFRRDAHKSE VAHRFKDLGE ENFKALVLIA FAQYLQQCPF EDHVKLNVNEV TEFAKTCVAD ESAENCCKSL HTLFGDKLCT VATLRETYGE MADCCAKQEP ERNECFLOHK DDNPNLRLV RPEVDVMCTA FHDNEETFLK KYLVEIARRH PYFYAPELLF FAKRYKAAFT ECCQAADKAA CLLPKLDELK DEGKASSAKQ GLKCSLQKF GERAFAWAV ARLSQRFPKA EFAEVSKLVT DLTQVHTECC HGDLLCADD RADLAKYICE NQDSISSKLK ECCEKPLEK SHCIAEVEND EMPADLPSLA ADFVGSKDVC KNYAEAKDVF LGMFLYEYAR RHPDYSVVLL LRLAKTYETT LEKCCAAADP HECYAKVFDE FKPLVEEPQN LIQNCSELF QLGEYKFQNA LLVRYTKKVP QVSTPTLVEV SRNLGKVGSK CCKHPEAKRM PCAEDCLSVF LNQLCVLHEK TPVSDRVTKC CTESLVNGRPCFSALEVDETYVPKEFNAETFTFHADICTLSEKERQIK KQTALV ELVKHK PKATKEQLKAVMDDFAAFVEKCKADDKET CFAEEGKKLVAASQAALGL
45	ACP12 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNKNTVYLQMNSLKPEDTGVYY CNALYGTIDYWGKGTQVTVSSggggsgggsgggsgggsaptssstktqlqlehlldlqmil nginnyknpkltrmltfkfymppkatelkhlqcleelkpleevlnlaqsknfhlrprdlisninvivlelkg settfmceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQLVESGGG LVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNKNTLYLQMNSLRPEDTAVYYCTIGGSLS VSSQGTLVTVSSggggsgggsgggsgggsEVQLVESGGGLVQPGGSLRLSCAA SGFTFSSYTLAWVRQAPGKGLEWVAIDSSSYTSPDVRGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCARDSDNWDALDYWGQGTITVTV SSGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTTTCASQNVG

SEQ ID NO.	Name	Sequence
		TNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPYTFGGGTKEIKHHHHH
46	ACPI3 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVYLQMNSLKPEDTGVYYCNALYGTDYWGKGTVTVSSgggsgggsggggsEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTTLAWVRQAPGKGLEWVAIDSSSYTYSPD TVRGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNALDYWGQGTTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPYTFGGGTKEIKgggsgggsggggsEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPE DTAVYYCTIGGSLSVSSQGTLVTVSSSGGPAGMKGLPGSaptssstktqlqlehlldlqmilnginnyknpkltmrmtfkfymppkatelkhlqcleelkpleevlnlaqsknfhlprdlisininvivlelkgettfmceyadetativeflnrwitfcqsisttHHHHH
47	ACPI4 (IL2 conjugate)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTTLAWVRQAPGKLEWVAIDSSSYTYSPD TVRGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNALDYWGQGTTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPYTFGGGTKEIKgggsgggsgggsgggsgggsgggsgggsgggsgggSGGPAGMKGLPGSa ptssstktqlqlehlldlqmilnginnyknpkltmrmtfkfymppkatelkhlqcleelkpleevlnlaq sknfhlprdlisininvivlelkgettfmceyadetativeflnrwitfcqsisttSGGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRP EDTAVYYCTIGGSLSVSSQGTLVTVSSH HHHHHH
48	ACPI5 (IL2 conjugate)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTTLAWVRQAPGKLEWVAIDSSSYTYSPD TVRGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNALDYWGQGTTVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPYTFGGGTKEIKgggsgggsgggsgggsgggsgggsgggsgggsgggEVQLVESGGGLVQPG NSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAE

SEQ ID NO.	Name	Sequence
		SVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGT LVTVSSggggsgggsgggsgggsgggSGGPGPAGMKGLPGSaptssstktqlqlehlldlqmil nginnyknpkltmrmltfkfympkkatelkhlqcleelkpleevlnlaqsknfhrlprdlisninvivlelkg settfmceyadetativeflnrwitfcqsiistltHHHHHH
49	ACP16 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltmrmltfkfympkkatelkhlqcleelkpleevlnla qsknfhrlprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMK GLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAP GKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRP EDTAVYYCTIGGSLSVSSQGT LVTVSSggggsgggsgggsgggsgggsgggsggg sgSGGPGPAGMKGLPGSEVQLVESGGGLVQPGSLRLSCAASGFTFS SYTLAWVRQAPGKGLEWVAIDSSSYTYSPD'TVRGRFTISRDNAAKN SLYLQMNSLRAEDTAVYYCARDSDALDYWGQGT'TVTVSSGGG GSGGGGSGGGGSDIQMTQSPSSLSASVGDRVITITCKASQNVGTNVG WYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISLQPED FATYYCQQYYTYTPYTFGGGTKVEIKHHHHHH
50	ACP17 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNAAKNTVYLQMNSLKPEDTGYYY CNALYGT'DYWGKGTQVTVSSggggsgggsgggsgggsgggsgggsgggsgggsggg nginnyknpkltmrmltfkfympkkatelkhlqcleelkpleevlnlaqsknfhrlprdlisninvivlelkg settfmceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQLVESGGG LVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSL VSSQGT LVTVSSggggsgggsgggsgggsgggsgggsgggsgggsgggSGGPGPAGMKGLP GSEVQLVESGGGLVQPGSLRLSCAASGFTFSYTLAWVRQAPGK LEWVAIDSSSYTYSPD'TVRGRFTISRDNAAKNSLYLQMNSLRAEDT VYYCARDSDALDYWGQGT'TVTVSSGGGGSGGGGSGGGGSDIQ MTQSPSSLSASVGDRVITITCKASQNVGTNVGWYQQKPGKAPKALIY SASFRYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYTPY FGGGTKVEIKHHHHHH
51	ACP18 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNAAKNTVYLQMNSLKPEDTGYYY CNALYGT'DYWGKGTQVTVSSggggsgggsgggsgggsgggsgggsgggsgggsggg nginnyknpkltmrmltfkfympkkatelkhlqcleelkpleevlnlaqsknfhrlprdlisninvivlelkg

SEQ ID NO.	Name	Sequence
		settfmceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQLVESGGG LVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLS VSSQGTLVTVSSggggsgggsgggsgggsgggsgggsgggsgggEVQLVESGGGLVQ PGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLEWVAADSSSYTYSP DTVRGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNWDALD YWGQGTTVTVSSsggpgpagmkglpgsDIQMTQSPSSLSASVGDRVTTTC ASQNVGTNVGWYQQKPGKAPKALIYSASFRTSGVPSRFSGSGSGTD FTLTISLQPEDFATYYCQQYYTYPTYTFGGGTKVEIKHHHHHHH
52	ACP19 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltrmltkfymppkatelkhlcleeekpleevlnla qsknfhlrprdlisninvivlelkgettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMK GLPGSgggsgggsgggsgggsgggsgggsgggsgggsgggEVQLVESGGGLVQPGGSLRL SCAASGFTFSSYTLAWVRQAPGKGLEWVAADSSSYTYSPDTVGRF TISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNWDALDYWGQGT TVTSSGGGGSGGGSGGGGSDIQMTQSPSSLSASVGDRVTTTCASQ NVGTNVGWYQQKPGKAPKALIYSASFRTSGVPSRFSGSGSGTDFTLT ISSLQPEDFATYYCQQYYTYPTYTFGGGTKVEIKgggsgggsgggsgggEVQ LVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTLVTVSSgggsgggsgggsgggQVQLQESGGGLVQAGGS LRLSCAASGRIFSIDMSWYRQAPGKQRELVARITRGGTISYDDSVKG RFTISRDNANKNTVYLQMNSLKPEDTGYYCNALYGTDYWGKGTV TVSSH HHHHHH**
53	ACP20 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltrmltkfymppkatelkhlcleeekpleevlnla qsknfhlrprdlisninvivlelkgettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMK GLPGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPG KGLEWVAADSSSYTYSPDTVGRFTISRDNANKNSLYLQMNSLRAED TAVYYCARDSDNWDALDYWGQGTTVTSSGGGGSGGGSGGGGSDI QMTQSPSSLSASVGDRVTTTCASQNVGTNVGWYQQKPGKAPKALI YSASFRTSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPY TFGGGTKVEIKHHHHHHH
54	ACP21 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltrmltkfymppkatelkhlcleeekpleevlnla qsknfhlrprdlisninvivlelkgettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMK

[illegible]

SEQ ID NO.	Name	Sequence
		yadetativeflnrwitfcqsiistltHHHHHHH
57	ACP24 (IL2 conjugate)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLE WVAADSSSYTYSPDTRGRFTISRDNAKNSLYLQMNSLRAEDTAV YYCARDSDNWALDYWGQGTTVTVS SGGGGSGGGGSGGGGSDIQM TQPSSLSASVGDRVITICKASQNVGTNVGWYQQKPGKAPKALIYS ASFRYSGVPSRFSGSGSGTDFTLTIS SLQPEDFATYYCQQYYTYPYTF GGGTKVEIKSGGPGPAGMKGLPGSa p t s s t k k t q l q l e h l l d l q m i n g i n n y k n p k l t r m l t f k y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p r d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e f l n r w i t f c q s i i s t l t H H H H H H
58	ACP25 (IL2 conjugate)	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLE WVAADSSSYTYSPDTRGRFTISRDNAKNSLYLQMNSLRAEDTAV YYCARDSDNWALDYWGQGTTVTVS SGGGGSGGGGSGGGGSDIQM TQPSSLSASVGDRVITICKASQNVGTNVGWYQQKPGKAPKALIYS ASFRYSGVPSRFSGSGSGTDFTLTIS SLQPEDFATYYCQQYYTYPYTF GGGTKVEIKg g g g s g g g s g g g s g g g s g g g s g g g s S G G P G P A G M K G L P G S a p t s s t k k t q l q l e h l l d l q m i n g i n n y k n p k l t r m l t f k y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p r d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e f l n r w i t f c q s i i s t l t H H H H H H
59	ACP26 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNAKNTVYLQMNSLKPEDTGVVY CNALYGTDYWKGKTQVTVS S g g g s g g g s g g g s a p t s s t k k t q l q l e h l l d l q m i l n g i n n y k n p k l t r m l t f k y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p r d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e f l n r w i t f c q s i i s t l t S G G P G P A G M K G L P G S E V Q L V E S G G G L V Q P G N S L R L S C A A S G F T F S K F G M S W V R Q A P G K G L E W V S S I S G S G R DTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLS VSSQGTLVTVSS g g g s g g g s g g g s g g g s Q V Q L Q Q S G A E L V R P G T S V K V SCKASGYAFTNYLIEWVKQRPGQGLEWIGVINPGSGGTNYNEKFEG KATLTADKSSSTAYMQLSSLTSDDS AVYFCARWRGDGYAYFDVW GAGTTVTVS S g g g s g g g s g g g s D I V L T Q S P A S L A V S L G Q R A T I S C K A S Q SVDYDGDSYMNWWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGT DFTLNHPVEEEDAATYYCQQSNEDPYTFGGGTKEIKHHHHHHHEPE A
60	ACP27 (IL2 conjugate)	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNAKNTVYLQMNSLKPEDTGVVY

SEQ ID NO.	Name	Sequence
		CNALYGT DYWGKGTQVT VSSggggsgggsgggsgaptssstktqlqlehlldlqmil nginnyknpkltrmltfkfympkkatelkhlqcleelkpleevlnlaqsknfhlrprdlisninvivlelkg settfmcceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQLVESGGG LVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLS VSSQGT LVT VSSggggsgggsgggsgggsggggDIVLTQSPASLAVSLGQRATIS CKASQSVDYDGDSYMNWYQQKPGQPPKLLIYAASNLESGIPARFSG SGSGTDFTLNIHPVEEEDAATYYCQQSNEDPYTFGGGT KLEIKggggsg gggsgggsgQVQLQQSGAELVRPGTSVKV SCKASGYAFTNYLIEWVKQ RPPGQGLEWIGVINPGSGGTNYNEKFKGKATLTADKSSSTAYMQLSSL TSDDSAVYFCARWRGDGYAYFDVWGAGTTVT VSSH HHHHHEPE A
61	ACP28 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltrmltfkfympkkatelkhlqcleelkpleevlnla qsknfhlrprdlisninvivlelkgsett fmcceyadetativeflnrwitfcqsiistltSGGPGPAGMK GLPGSggggsgggsgggsgggsgggsgggsgggsgQVQLQQSGAELVRPGTSVKV SCK ASGYAFTNYLIEWVKQRPGQGLEWIGVINPGSGGTNYNEKFKGKAT LTADKSSSTAYMQLSSLTSDDSAVYFCARWRGDGYAYFDVWGAG TTVTVSSggggsgggsgggsgggsgggsgggsgggsgDIVLTQSPASLAVSLGQRATISCKASQSD YDGDSYMNWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTL NIHPVEEEDAATYYCQQSNEDPYTFGGGT KLEIKggggsgggsgggsgggsgEV QLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGLEW VSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVY YCTIGGSLSVSSQGT LVT VSSggggsgggsgggsgggsgggsgggsgggsgQVQLQESGGGLVQAG GSLRLSCAASGRIFSIDIMSWYRQAPGKQRELVARITRGGTISYDDSV KGRFTISRDN AKNTVYLQMNSLKPEDTG VYYCNALYGT DYWGKGT QVT VSSH HHHHHEPEA
62	ACP29 (IL2 conjugate)	aptssstktqlqlehlldlqmilnginnyknpkltrmltfkfympkkatelkhlqcleelkpleevlnla qsknfhlrprdlisninvivlelkgsett fmcceyadetativeflnrwitfcqsiistltSGGPGPAGMK GLPGSggggsgggsgggsgggsgggsgggsgggsgDIVLTQSPASLAVSLGQRATISCKA SQSVDYDGDSYMNWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSG TDFTLNIHPVEEEDAATYYCQQSNEDPYTFGGGT KLEIKggggsgggsg gggsgggsgQVQLQQSGAELVRPGTSVKV SCKASGYAFTNYLIEWVKQRPGQ GLEWIGVINPGSGGTNYNEKFKGKATLTADKSSSTAYMQLSSLTSDD

SEQ ID NO.	Name	Sequence
		SAVYFCARWRGDGYAYFDVWGAGTTVTVSSggggsgggsgggsgEVQ LVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTLVTVSSggggsgggsgggsgQVQLQESGGGLVQAGGS LRLSCAASGRIFSIDIMSWYRQAPGKQRELVARITRGGTISYDDSVKG RFTISRDNAKNTVYLMNSLKPEDTGVYYCNALYGTDYWGKGTQV TVSSHHHHHHEPEA
63	IL2Ra	10 20 30 40 50 MDSYLLMWGL LTFIMVPGCQ AELCDDDPPE IPHATFKAMA YKEGTMLNCE 60 70 80 90 100 CKRGFRRIKS GSYMLCTGN SSHSSWDNQC QCTSSATRNT TKQVTPQPEE 110 120 130 140 150 QKERKTTEMQ SPMQPVDAQS LPGHCREPPP WENEATERIY HFVVGQMVYY 160 170 180 190 200 QCVQGYRALH RGPAESVCKM THGKTRWTQP QLICTGEMET SQFPGEKPKQ 210 220 230 240 250 ASPEGRPESE TSCLVTTTDF QIQTEMAATM ETSIFTTEYQ VAVAGCVFLL 260 270 ISVLLLSGLT WQRRQRKSRR TI
64	IL2Rb	10 20 30 40 50 MAAPALSWRL PLLILLPLA TSWASAAVNG TSQFTCFYNS RANISCVWSQ 60 70 80 90 100 DGALQDTSCQ VHAWPDRRRW NQTCELLPVS QASWACNLIL GAPDSQKLTT 110 120 130 140 150 VDIVTLRVLC REGVRWRVMA IQDFKPFENL RLMAPIQLQV VHVETHRCNI 160 170 180 190 200 SWEISQASHY FERHLEFEAR TLSPGHTWEE APLLTLKQKQ EWICLETLP 210 220 230 240 250 DTQYEFQVRV KPLQGEFTTW SPWSQPLAFR TKPAALGKDT

SEQ ID NO.	Name	Sequence
		IPWLGHLLVG 260 270 280 290 300 LSGAFGFIL VYLLINCRNT GPWLKKVLKC NTPDPSKFFS QLSSEHGGDV 310 320 330 340 350 QKWLSSPFPS SSFSPGGLAP EISPLEVLER DKVTQLLLQQ DKVPEPASLS 360 370 380 390 400 SNHSLTSCFT NQGYFFFHLP DALEIEACQV YFTYDPYSEE DPDEGVAGAP 410 420 430 440 450 TGSSPQPLQP LSGEDDAYCT FPSRDDLLF SPSLLGGPSP PSTAPGSGA 460 470 480 490 500 GEERMPPSLQ ERVPRDWDQP PLGPPTPGVP DLVDFQPPFE LVLREAGEEV 510 520 530 540 550 PDAGPREGVS FPWSRPPGQG EFRAFNARLP LNTDAYLSLQ ELQGQDPHTL V
65	IL2Rg	10 20 30 40 50 MLKPSLPFTS LLFLQLPLL GVLNTTILTP NGNEDTTADF FLTMTPTDSL 60 70 80 90 100 SVSTLPLPEV QCFVFNVEYM NCTWNSSESEP QPTNLTLYHW YKNSDNDKVQ 110 120 130 140 150 KCSHYLFSEE ITSGCQLQKK EIHLYQTFVV QLQDPREPRR QATQMLKLQN 160 170 180 190 200 LVIPWAPENL TLHKLSESQL ELNWNRRFLN HCLEHLVQYR TDWDHSWTEQ 210 220 230 240 250 SVDYRHKFSL PSVDGQKRYT FRVRSRFPNPL CGSAQHWSEW SHPIHWGSNT 260 270 280 290 300 SKENPFLFAL EAVVISVGSM GLIISLLCVY FWLERTMPRI PTLKNLEDLV 310 320 330 340 350 TEYHGNFSAW SGVSKGLAES LQPDYSERLC LVSEIPPKGG ALGEGPGASP 360 CNQHSPYWAP PCYTLKPET
66	ACP04 (human)	iwelkkdvyyvvelldwypdagemvvlctdtpedgitwldqssevlsgsklttiqvkefgdagqytch kggevlshslilhlhkkgedgiwstdilkdqkepkntflrceaknysgrftcwwlttistdlfsvkssrgssd

[illegible]

SEQ ID NO.	Name	Sequence
	p40/murine p35 IL12 conjugate)	THPALLFGTGTKVTVLgggsgggsgggsgggsgVQQLVESGGGVVQPGRSLR LSCAASGFTFSSYGMHWVRQAPGKGLEWVAFIRYDGSNKYYADSV KGRFTISRDN SKNTLYLQMNSLR AEDTAVYYCKTHGSHDNWGQGT MVT VSSggggsgggsgggsgggsgggsgggsgggsgSGGPGPAGMKGLPGSiwelk kdvyvveldwypdapgemvvlctdtpedgitwtldqssevlsgsklttiqvkefgdagqytchkgge vlshtlllhkkedgiwstdilkdkpknktflrceaknysgrftcwwlittistdltsvkssrgssdpqgv tgaatlsaervrgdnkeyeysvecqedsacpaeeslpievmdavhklkyenyttssffirdiikpdppk nlqlkplknsrqvevsweyptdwstphsysfsltfcvqqgkskrekkdrvf dtktsatvicrk nasisvra qdryyssswsewasvp csgggsgggsgggsgggsgggsvipvs gparclsqsrmlktddmvktareklkhy scta edidheditr dqtstlktclplelhknesclatre tsstrgsclppqktslmm tclgl siyedlkmyqte fqainaalqn hnhqqiildkg mlvaidelmqslnhngetlrqkp pvg eadpyrvkm klcillhafstrvv tinrmgy lssaSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLR LSCA ASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTI SRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLVS SSQGTLVT VSSggg sgggsgggsgggsgggsgggsgggsgggsgVQQLQESGGGLAQAGGSLSLSCAASGFTVS NSVMAWY RQT PGKQREFVA IINSVGSTNYADSVKGRFTISR DNAKNTVY LQMNN LKPEDTAVYYVCNRNFDRIYW GQGTQVT VSSH HHHHHEPEA
70	ACP08 (human p40/murine p35 IL12 conjugate)	QVQLQESGGGLAQAGGSLSLSCAASGFTVS NSVMAWY RQT PGKQR EFVA IINSVGSTNYADSVKGRFTISR DNAKNTVY LQMNNLKPEDTAV YVCNRNFDRIYW GQGTQVT VSSggggsgggsgggsgggsgQSVLTQPPS VSGAP GQRVTISC SGSR SNIGSN TVKWY QQLPGTAP KLLIY YNDQRPSGVPD RFSGSKSGTSASLAITGLQA EDEADYYCQSYDRY THPALLFGTGTKV TVLgggsgggsgggsgggsgVQQLVESGGGVVQPGRSLRLSCAASGFTFSSYG MHWRQAPGKGLEWVAFIRYDGSNKYYADSVKGRFTISR DN SKNT LYLQMNSLR AEDTAVYYCKTHGSHDNWGQGT MVT VSSggggsgggsg ggsgggsgggsgggsgggsgSGGPGPAGMKGLPGSiwelkdv yvveldwypdapgem vvlctdtpedgitwtldqssevlsgsklttiqvkefgdagqytchkgge vlshtlllhkkedgiwstdilk dkpknktflrceaknysgrftcwwlittistdltsvkssrgssdpqgv tgaatlsaervrgdnkeyeys vecqedsacpaeeslpievmdavhklkyenyttssffirdiikpdppknlqlkplknsrqvevswey ptdwstphsysfsltfcvqqgkskrekkdrvf dtktsatvicrk nasisvraqdryyssswsewasvp csgg ggsgggsgggsgggsgggsgggsgggsgggsgvipvs gparclsqsrmlktddmvktareklkhyscta edidheditr dqtstlktclplelhknesclatre tsstrgsclppqktslmm tclgl siyedlkmyqte fqainaalqn hnhqqiildkg mlvaidelmqslnhngetlrqkp pvg eadpyrvkm klcillhafstrvv tinrmgy lssaSGGPGP

SEQ ID NO.	Name	Sequence
		AGMKGLPGSEVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWV RQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMN SLRPEDTAVYYCTIGGSLSVSSQGTTLTVTSSHHHHHHEPEA
71	ACP09 (human p40/murine p35 IL12 conjugate)	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNLSLRPEDTAV YYCTIGGSLSVSSQGTTLTVTSSggggsgggsggggsQSVLTQPPSVSGAPG QRTVISCSSRSNIGSNTVKWYQQLPGTAPKLLIYYNDQRPSGVPDR FSGSKSGTSASLAITGLQAEDEADYYCQSYDRYTHPALLFGTGTKVT VLggggsgggsggggsQVQLVESGGGVVQPGSRSLRSLCAASGFTFSYG MHWVRQAPGKGLEWVAFIRYDGSNKYYADSVKGRFTISRDNASKNT LYLQMNLSRAEDTAVYYCKTHGSHDNWGQGTMTVTVSSggggsgggsgg ggsgggsgggsgggsgggsgSGGPGPAGMKGLPGSiwelkkdvyyvldwypdagm vvlctdtpedgitwtldqssevlsgsktltiqvkefgdagqytchkggevlshslllhk kedgiwstdilkdqkepnkflrceaknysgrftcwwltistdlfsvkssrgssdpqgvtcgaat saervrgdnkeyeysvecqedsacpaaeeslpievmdavhklkyenytsffirdiikpdp ppknqlkplknsrqvevsweypdtwstphsyfslfcvqvqgkskrekkdrvftdkt satvicrknaasisvraqdryssswsewasvpcsgggsgggsgggsgggsgsr vipvsgparclsqsmllktddmvktareklkhysctaedisiditrdqtsltklcl plelhknesclatretsstrgscldppqktslmmtlclgsiyedlkmyqtefqaina alqnhnhqqiildkgmlvaidelmqslnhngetlrqkppvgadpyrvkmkcllhafstr vvtinrvmgylssaHHHHHHEPEA
72	ACP10 (human p40/murine p35 IL12 conjugate)	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNLSLRPEDTAV YYCTIGGSLSVSSQGTTLTVTSSSGGPGPAGMKGLPGSiwelkkdvyyvld wypdagm vvlctdtpedgitwtldqssevlsgsktltiqvkefgdagqytchkgge vlshslllhk kedgiwstdilkdqkepnkflrceaknysgrftcwwltistdlfsvkss rgssdpqgvtcgaatlsaervrgdnkeyeysvecqedsacpaaeeslpievmdavhkl kyenytsffirdiikpdp ppknqlkplkn srqvevsweypdtwstphsyfslfcvqvq gkskrekkdrvftdkt satvicrknaasisvraqdryssswsewasvpcsgggsggg sgggsgggsgsr vipvsgparclsqsmllktddmvktareklkhysctaedisiditrd qtsltklclplelhknesclatretsstrgscldppqktslmmtlclgsiyedlkmyqtef qainaalqnhnhqqiildkgmlvaidelmqslnhngetlrqkppvgadpyrvkmkcllh afstrvvtinrvmgylssaSGGPGPAGMKGLPGSggggsgggsgggsgggsgggsgg sQSVLTQPPSVSGAPGQRTVISCSSRSNIGSNTVKWYQQLPGTAPKLLIYYNDQRP SGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDRYTHPALLFG

SEQ ID NO.	Name	Sequence
		TGTKVTVLggggsgggsgggsgggsgVQLVESGGGVVQPGRSLRLSCAASGF TFSSYGMHWVRQAPGKGLEWVAFIRYDGSNKYYADSVKGRFTISRDN NSKNTLYLQMNSLRAEDTAVYYCKTHGSHDNWGQGTMTVTVSSHH HHHHEPEA
73	ACPI1 (human p40/murine p35 IL12 conjugate)	iwelkkdvyyvvelwypdapgemvvlctdtpcedgitwtldqssevlsgskltliqvkefgdagqytch kggevlshslhllhkkedgiwstdilkdkqepknktflrceaknysgrftcwwlttistdltfsvkssrgssd pqgvtcgaatlsaervrgdnkeyesvecqedsacpaaeeslpievmvдавhklkyenytsffirdiik pdpknqlkplknsrqvevsweypdtwstphsyfsltfcvqvqgkskrekkdrvftdktsatvicrkna sisvraqdryyssswwsewasvpcsgggsgggsgggsgggsgvrvpsgparclsqsrnlkttddmvktar eklkhsctaedidheditrqtslktclplelhknesclatretsstrgsclppqktslmmtdclgsiyedlk myqtefqainaalqnhnhqqiilkkgmlvaideImqslnhngetlrqkppvgeadpyrvkmklcillha fstrvvtinrvmgylssaSGGPGPAGMKGLPGSggggsgggsgggsgggsgggsgggsgggsggg ggsQSVLTQPPSVSGAPGQRTISCSGSRNIGSNTVKWYQQLPGTAP KLLIYYNDQRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSY DRYTHPALLFGTGTKVTVLggggsgggsgggsgggsgVQLVESGGGVVQPGR SLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAFIRYDGSNKYYA DSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCKTHGSHDNWG QGTMTVTVSSggggsgggsgggsgggsgEVQLVESGGGLVQPGNSLRLSCAASG FTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN NAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVTVSSHHHH HHEPEA
74	IL12 p40 human (Uniprot Accession No. P29460)	1020304050 MCHQQLVISW FSLVFLASPL VAINELKKDV YVELDWYPD APGEMWVLT 68788090100 DTPEEDGITW TLDQSSEVLG SGKTLTIQVK EFGDAGQYTC HKGGEVLSHS 110120130140150 LLLLHKXEDG IKSTDIKDKQ KEPKNETFLR CEAKNYSGRF TCWWLTIST 160170180190200 DLTFSVKSSR GSSDPQGVTC GAATLSAERV RGDNKEYEYS VECQEDSACP 210220230240250 AAEESLPIEV MDAVHKLKY ENYTSSFFIR DIKPDPPKN LQLKPLKNSR 260270280290300 QVEVSWEYPD TWSTPHSYFS LTFCVQVQGK SKREKKDRVF TDKTSATVIC 310320 RKNASISVRA QDRYSSSSWS FWASVPCS

SEQ ID NO.	Name	Sequence
75	IL12 p35 mouse (Uniprot Accession No. P43431)	10 20 30 40 50 MCQSRYLFL ATLALLNHLS LARVIPVSGP ARCLSQSRNL LKTTODMVKT 60 70 80 90 100 AREKLKHYSC TAEDIDHEDI TRDQTSTLKT CLPLELHKNE SCLATRETSS 110 120 130 140 150 TTRGSCLPPO KTSLMMTLCL GSIYEDLKMY QTEFGAINAA LQNNHHQOII 160 170 180 190 200 LQKGMVAID ELMQSLNHNG ETLRQKPPVG EADPYRVKMK LCILLHAFST 210 RVVTINRVMG YLSSA
76	IL12Rb-2	10 20 30 40 50 MSHIERGGEL LNTALINLA LKAKLRACKS GRVYKRSKV ILLESTYKAI 60 70 80 90 100 GLKLRQCKE HYERENKLL YNEEDRINTH NCHSINSQNT SLRLGTLKLV 110 120 130 140 150 GLALINERK LQICGAKLEV GQAFGRNNL SCIGGKQQL VACAKRGGKH 160 170 180 190 200 THLYTETLD LSGFENKIDG KCKKQIVKRA LDGKMLLR RPSNTERAKV 210 220 230 240 250 KAKSLGSSS SLPSITFLD LURLEPEDI RIKTQKQVZ RCTLYKPDG 260 270 280 290 300 LVLRLRLRZ RSHRLNNMV NTKKAGGHR LLEKKEIKV RQIISKIHL 310 320 330 340 350 KKSSSSSSS SLRQCTPSS RICHLDKVM KHLIYSSQO LSLKKNLSV 360 370 380 390 400 SESRKILLY QTLQLRIGR KMTNHLIGR LSKILVLEKI GNVAVVYAA 410 420 430 440 450 SEKESLETR INDDLCSSG LARPCSSN SCGMNLIKZ NQRRKRRSA 460 470 480 490 500 VQENYVMDR LKGGGLQGR LKHLGRDYN VVALISENIK SXICKELIVY 510 520 530 540 550 ALSRGGSLA ALENSKHA FLGGHINAI IRRKSLIS KSHIVQSQM 560 570 580 590 600 QMLHYRIK KRRGSSSPQ LCLIEKRVG USHPINSLR EYIKLNLIA 610 620 630 640 650 LIRAGISGR NERFCLOCK LNMATVARS ICIALIMGI ESTEVQOKY 660 670 680 690 700 KVLLALRQ KSRRLKRA NSTCKKELI AEEKQLRLD RLIDKSTEL 710 720 730 740 750 REELVIAEV LKQIVFIS RCRNVSQRA KSLQGHQSE KSRHSAQSS 760 770 780 790 800 KRRALQSS SOLVELKYL EERGSSPER NRCCHVLR AGDRTKDEK 810 820 830 840 850 LRNIGLRS MEALRSLR ELRPHALS VERRASLMR IESCGKHAL 860 QQLNRCDEL ML

SEQ ID NO.	Name	Sequence
77	IL12Rb-1	<pre> 10 20 30 40 50 MEPLVETWVVD LILFLFLLSRQ GAACTSTSECC FQDPFVFDAS SSSRSGSRRL 60 70 80 90 100 RCYRISSERY EGSWQYLSPT AGVSHFLRCC LSSGRCCYFA AGSAILGFS 110 120 130 140 150 DQACUSVLYT VTLWVESWAR NQTEKSEVY LQLYNSUYE DFLGDIKVSX 160 170 180 190 200 LACQLRMWEV TDENQVGRVY QFRHRIPSSR WKLCDCGQDQ DYTESCLCPL 210 220 230 240 250 EMFVAGEFCL RRACLGSGGS SWSWSSPVC VPPENTPDPQ VRFVTECLGQ 260 270 280 290 300 SGARRLTKE QPTQLELSEG CQGLAPGTEV TVRLQLMLLS CPCKAKAIRI 310 320 330 340 350 LHLGKMDYLE GAAYNVAVIS SNQFGGCLNQ TWHIDADTHT EDVALNIEVG 360 370 380 390 400 TNGITMWFPA RAQSMYVCI E WQFVQDGGCL ATCSLTADQS PEPACMAIYS 410 420 430 440 450 WSRESGAMGQ EKCYYITIFA SAHPEKLTLE STVLSTYHFG GNASARGIPH 460 470 480 490 500 HVSUKWMSLD ESEVENADEL LETCPGVLRK YVVMCRDEDE KQVSEHFWDP 510 520 530 540 550 TETQVILSGL RAGVAYTVGV RADTRSLGV NSQDQRFQIR VVYSWNLIFE 560 570 580 590 600 ASLGSFLSIL LVCVLGVLGL NRAAPRLCPD LDTPCASSRI EFPGGMHMQ 610 620 630 640 650 WINFVDFQEE ASLQEALVVE MSWDKGERTE FLEKIELSEG APELALDTL 660 SLIDEDRCKA RM </pre>
79	IL-12 p35 human (Uniprot accession no. P29459)	<pre> 10 20 30 40 50 NCHQQLVISW FSLVFLASPL VAIWELKKDV YVVELDWYPD APGENWVLTC 60 70 80 90 100 DTPEEDGITW TLDQSSEVLG SGKTLTIQVK EFGDAGQYTC HKGGEVLSHS 110 120 130 140 150 ILLUNKKEGG IWSYDILKQK KEPKNKTFLE CEAKNYSGRF TCMWLTITST 160 170 180 190 200 DLTFSVKSSR GSSDPQGVTC GAATLSAERV RGNKKEYEYS VECQEDSACP 210 220 230 240 250 AAECESLPIEV MVDVHKLKY ENVTSSFFIR DIKPPPPKN LQLKPLKNSR 260 270 280 290 300 QVEVSWEVPD TWSTPHSYFS LIFCVQVQCK SKREKKDRVF YDKTSATVIC 310 320 RKNASISVRA QDRVYSSSW S EWSVPC5 </pre>

SEQ ID NO.	Name	Sequence
80	IL-12 p40 mouse (Uniprot accession no. P43432)	10 20 30 40 50 MCPQKLTISW FAIVLLVSPL NAWMELEKDV YVEVDWTPD APGETVNLTC 60 70 80 90 100 DTPEEDDITW TSDQRHGVIG SGKTLTITVK EFLDAGQYTC HKGGETLSHS 110 120 130 140 150 HLLLHKKENG IWSTEILKNF KSKTFLKCEA PHYSGRFTCS WLVRNMDLK 160 170 180 190 200 FNKSSSSSP DSRAVTCGMA SLSAEKVTLD QRDYEKYSVS CQEDVTCPTA 210 220 230 240 250 EETLPIELAL EARQQWNYEN YSTSFFIRDI IKPDPPKNLQ MKPLENSQVE 260 270 280 290 300 VSLWYPDWNS TPHSYFSLKF FVRIQRKKEK MKETEEGCNQ KGAFIVEKTS 310 320 330 TEVQCKGGNV CVQAQDRYYN SSSCKWACVP CRVRS
81	ACP01 (mouse IFN γ conjugate)	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTTLVTVSSSGGPGPAGMKGLPGShgtviesleslnnyfn ssgidveekslfldiwmwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffnskakkd afmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPAGMKGLPGSEVQ LVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTTLVTVSSHHHHHH
82	ACP02 (mouse IFN γ conjugate)	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTTLVTVSSSGGPGPAGMKGLPGShgtviesleslnnyfn ssgidveekslfldiwmwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffnskakkd afmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPAGMKGLPGShgtvi esleslnnyfnssgidveekslfldiwmwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlitt ffnskakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPAGMKG LPGSEVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPG KGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPE DTAVYYCTIGGSLSVSSQGTTLVTVSSHHHHHH
83	ACP03 (mouse IFN γ conjugate)	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTTLVTVSSSGGPGPAGMKGLPGShgtviesleslnnyfn

SEQ ID NO.	Name	Sequence
		ssgidveekslfldiwrnwqkdgdmkilqsqiisfylrlfevlkdnqaisnnisvieshlittffsnskakkd afmsiakfevnpqvqrqafnelirrvvhqllpesslrkrkrsrcggggsgggsgggshgtvieslesln nyfnssgidveekslfldiwrnwqkdgdmkilqsqiisfylrlfevlkdnqaisnnisvieshlittffsnsk akkdafmsiakfevnpqvqrqafnelirrvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGS EVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTLTVSSHHHHHH
84	Human IFN- g (Uniprot Accession No. P01579)	1020304050 MKYTSYILAF QLCIVLGSLG CYCQDPYVKE AENLKRYFNA GHSDVADNGT 60708090100 LFLGILKNWK EESDRKIMQS QIVSFYFKLF KNFKODQSIQ KSVETIKEDM 110120130140150 NVKFFNSNKK KRDDFEKLTH YSVTDLNVQR KAIHELIQVM AELSPAAKTG 160 KRKRSQMLFR GRRASQ
85	Mouse IFN-g (Uniprot Accession No. P01580)	1020304050 MNATHCILAL QLFMAVSGC YCHGTVIESL ESLNNYFNSS GIDVEEKSLF 60708090100 LDINRWQKD GDMKILQSQI ISFYLRLEFV LKDNQAISMN ISVIESHLIT 110120130140150 TFFSNSKAKK DAFMSIAKFE VNNPQVQRQA FNELIRRVHQLLPESSLRKR KRSRG
86	ACP30 (mouse IFN γ conjugate)	mdmrvpaqllgllllwlgarcEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfldiwrnwqkdgdmkilqsqiisfylrlfevlkdnqai snnisvieshlittffsnskakkdafmsiakfevnpqvqrqafnelirrvvhqllpesslrkrkrsrcSGGP GPAGMKGLPGSEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMS WVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYL QMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSSSGGPGPAGMKGLP GShgtviesleslnnyfnssgidveekslfldiwrnwqkdgdmkilqsqiisfylrlfevlkdnqaisnnis vieshlittffsnskakkdafmsiakfevnpqvqrqafnelirrvvhqllpesslrkrkrsrcSGGPGPA

SEQ ID NO.	Name	Sequence
		GMKGLPGSEVQLVESGGGLVQPGNSLRLSAASGFTFSKFGMSWVR QAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNS LRPEDTAVYYCTIGGSLSVSSQGTLVTVSSHHHHHH
87	ACP31 (mouse IFN α 1 conjugate)	EVQLVESGGGLVQPGNSLRLSAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTLVTVSSSGGPGPAGMKGLPGScdlpqthnlrnkraltl lvqmrrlspisclskdrkdfgfpqekvdaqqikkaqaipvlseItqqilniftskdssaawntllldsfendlh qqIndlqgclmqvgvqefplqtedallavrkyfhritylrekkhspcawevvraevwralsssanvlg rlreekSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSAASGFT FSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAA KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLVTVSSHHHHHHHE PEA
88	ACP32 (mouse IFN α 1 conjugate)	EVQLVESGGGLVQPGNSLRLSAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTLVTVSSSGGPGPAGMKGLPGScdlpqthnlrnkraltl lvqmrrlspisclskdrkdfgfpqekvdaqqikkaqaipvlseItqqilniftskdssaawntllldsfendlh qqIndlqgclmqvgvqefplqtedallavrkyfhritylrekkhspcawevvraevwralsssanvS GGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLVTVSSHHHHHHHEPEA
89	IFN γ R1	1520354050 MALLFLPLV MQGVSPAENG IADLGFSVF TPINVTIESY NNNPIVYWEY 60708090100 QIMFQVPVET VEVKNYGVEN SEWIDACINI SHMYCNISEH VEDDPNSLWV 110120130140150 KVKARVQKE SAYAKSEFA VCRDGRIGPF KLEIAKEEQ IMIDIFHSV 160170180190200 FVNGDEQEVY YDEETCYIR VYNVYVDSNG SEIQYKILTQ KEDDCDPIQC 210220230240250 QLAIFVSSLN SQYCVREGV LEVNGVYTER SEWVCITIFN SSINGSLWIE 260270280290300 VVRALLPLV LSLVFIQFYI KNINPLKENS IILPKSLISV VRSATLETWF 310320330340350 ESKYVELITS YQPPSLEKEV VCKEPLSDAT VPQMRTEDNP GKVENTEELS 360370380390400 SITEVYTER NIDDUVEGSE LTPDERSES SLSNQSEEG SIALNAYHSE 410420430440450 NCSESCHSPN GEDTSGCLE SHSLSGSEF PPNNGEINT EGQELITVIE 460470480490 ASTFEGYKQ NVLVELLVDG SKRESLISYR PTELSKEFE

SEQ ID NO.	Name	Sequence
90	IFN γ R2	<pre> 10 20 30 40 50 MRPTILNSLL LLIGVFRAAR AAPDFELSQL EAPQMPKIAL YNAEGWLSWE 60 70 80 90 100 SVALSNSIRP VVYQVQEKYT LSKNFTADIM SIGUNCTQIT ATECDFTAAS 110 120 130 140 150 PSAGFENDFN VTLRLAAILG ALNSAWVNF WFGWVRNIVY EFPENIEVTP 160 170 180 190 200 GEGSLIIRAF SDFDIADIST AFFCYVHYW ERGQIQVKS EYFNSISLB 210 220 230 240 250 NLKFSRVYCL QVQAQLWUK SWIKVGHLS NISCIETNAD ASTELQQVIL 260 270 280 290 300 ISVGTFSLLS VLASACFELV LKYSGLIKW FHIPPSSIPQ IEEYLKSTQ 310 320 330 EILEALDKDS SPEDDVWSSV SIISFPEZEQ EDVLQTL </pre>
91	ACP51 Mouse IFG conjugate	QVQLQESGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQREL VARITRGGTISYDDSVKGRFTISRDNNAKNTVYLQMNSLKPEDTGVYY CNALYGTDYWGKGTQVTVSSggggsgggsggggsEVQLVESGGGLVQP GNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLY AESVKGRFTISRDNNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQ GTLVTVSSSGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwq kdgdmkilqsqisfyllrfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvrqaf nelirvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGSEVQLVESGGGLVQPGN SLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAES VKGRFTISRDNNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTL VTVSSHHHHHH
92	ACP52 Mouse IFG conjugate	EVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLE WVSSISGSGRDTLYAESVKGRFTISRDNNAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTLVTVSSSGGPGPAGMKGLPGShgtviesleslnnyfn ssgidveekslfdiwrnwqkdgdmkilqsqisfyllrfevlkdnqaisnnisvieshlittffsnkakkd afmsiakfevnnpqvrqafnelirvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGSEVQ LVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDTLYAESVKGRFTISRDNNAKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTLVTVSSSGGPGPAGMKGLPGSEVQLVESGGGLVQ PGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLY AESVKGRFTISRDNNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQ GTLVTVSSggggsgggsggggsQVQLQESGGGLVQAGGSLRLSCAASGRI FSIDIMSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNNAK NTVYLQMNSLKPEDTGVYYCNALYGTDYWGKGTQVTVSSHHHHHH
93	ACP53 Mouse IFG	cahkseiahryndlgeqhfkgvliafsqylqkcsydehaklvqevtdfaktcvadesaancdkslhtlfg dklcaipnlrenygeladccctkqepemecflqhkdndpslppferpeaeamctsfkenpttfmgthylhe

SEQ ID NO.	Name	Sequence
	conjugate	varrhpyfyapellyyaeqyneiltqccaeadkescltpkldgvkekalvssvrqrmkcsmmqkfgeraf kawavarlsqtfpnadfaeitklatltdtkvnkecchgdlllecaddraclakymcenqatissklqtccdkpl lkkahclsevehdtmpadlpaiaadfvedqevcknyaekdvflgtflyeysrrhpdysvslllrlakkye atlekccaeapppacygtvlaefqplveepknlvktncdlyeklgeygfqnailvrytqkapqvstptlve aarnlgrvgtkcctlpedqrlpcvedylsailnrvcllhektpvsehvtkccsgslverrrpcfsaltvdetyvp kefkaetftfhsdictlpekekqikkqatalaelvkhkpkataeqlktvmddfaqlfdtcckaadkdtcfste gpnlvtrckdalaSGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfldiwrnw qkdgdmkilqsqiisfyrlfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrq afnelirvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGSeahkseiahryndlgeqhfkgvlv iafsqylqkcsydehaklvqevtdfaktcvadesaancdkslhtlfgdklcaipnlrenygeladccctkqep ernecflqhkddnpslppferpeaeamctsfkenpttfmghylhevarrhpyfyapellyyaeqyneilt qccaeadkescltpkldgvkekalvssvrqrmkcsmmqkfgerafkawavarlsqtfpnadfaeitklatl dtkvnkecchgdlllecaddraclakymcenqatissklqtccdkpllkkahclsevehdtmpadlpaia adfvedqevcknyaekdvflgtflyeysrrhpdysvslllrlakkyeatlekccaeapppacygtvlaefq plveepknlvktncdlyeklgeygfqnailvrytqkapqvstptlveaarnlgrvgtkcctlpedqrlpcve dylsailnrvcllhektpvsehvtkccsgslverrrpcfsaltvdetyvpkefkaetftfhsdictlpekekqik kqatalaelvkhkpkataeqlktvmddfaqlfdtcckaadkdtcfstegpnlvtrckdalaHHHHHH
94	ACP54 Mouse IFG conjugate	eahkseiahryndlgeqhfkgvlvlfiafsqylqkcsydehaklvqevtdfaktcvadesaancdkslhtlfg dklcaipnlrenygeladccctkqepernecflqhkddnpslppferpeaeamctsfkenpttfmghylhe varrhpyfyapellyyaeqyneiltqccaeadkescltpkldgvkekalvssvrqrmkcsmmqkfgeraf kawavarlsqtfpnadfaeitklatltdtkvnkecchgdlllecaddraclakymcenqatissklqtccdkpl lkkahclsevehdtmpadlpaiaadfvedqevcknyaekdvflgtflyeysrrhpdysvslllrlakkye atlekccaeapppacygtvlaefqplveepknlvktncdlyeklgeygfqnailvrytqkapqvstptlve aarnlgrvgtkcctlpedqrlpcvedylsailnrvcllhektpvsehvtkccsgslverrrpcfsaltvdetyvp kefkaetftfhsdictlpekekqikkqatalaelvkhkpkataeqlktvmddfaqlfdtcckaadkdtcfste gpnlvtrckdalaSGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfldiwrnw qkdgdmkilqsqiisfyrlfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrq afnelirvvhqllpesslrkrkrsrcggggsgggsgggshgtviesleslnnyfnssgidveekslfldiwrnw qkdgdmkilqsqiisfyrlfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqv qrqafnelirvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGSeahkseiahryndlgeqhfkgvlv iafsqylqkcsydehaklvqevtdfaktcvadesaancdkslhtlfgdklcaipnlrenygeladccct kqepernecflqhkddnpslppferpeaeamctsfkenpttfmghylhevarrhpyfyapellyyaeqy neiltqccaeadkescltpkldgvkekalvssvrqrmkcsmmqkfgerafkawavarlsqtfpnadfaeit

SEQ ID NO.	Name	Sequence
		klatdltkvnkeechgdllcaddraelakymcenqatissklqtccdkplllkkaclsevehdtmpadlp aiaadfvedqevcknyaeakdvflgtflyeysrrhpdysvslllrakkyeatleccaeanppacygtvla efqplveepknlvktncdlyeklgeygfqnailvrytqkapqvstptlveaamlgrvgtkcctlpedqrlp cvedylsailnrvcillhektpvsehvtkccsgslvenrpfcsaltvdetyvpkefkaetffthsdicltpekek qikkqtalaelvkhpkataeqlktvmddfaqlfdtcekaadkdtcfstegpnlvtreckdalaHHHHH H
95	ACP50 Mouse IFG conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANTVY LQMNSLKPEDTGYYCYCNALYGTDYWGKGTQVTVSSggggsgggsgggsggg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLPGShgtviesleslnnyf nssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffsnkakk dafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcgggsgggsgggsgggshgtvieslesl nnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffsnk kakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPAGMKGLPG SEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSHHHHHH
96	ACP55 Mouse IFG conjugate	mdmrvpaqllgllllwlrgrcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqai snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGG GPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMS WVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYL QMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLP GShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqaisnnis vieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPA GMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVR QAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNS LRPEDTAVYYCTIGGSLSVSSQGTLLTVSSHHHHHH
97	ACP56 Mouse IFG conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLAQAGGSLRLSCAASGFTVSNS VMAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDNANTV YLQMNNLKPEDTAVYYVCNRRNFDRIYWGQGTQVTVSSggggsgggsgggsggg gsEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLPGShgtviesleslnnyf nssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffsnkakk dafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPGPAGMKGLPGSEV QLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEW VSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVY YCTIGGSLSVSSQGTLLTVSSHHHHHHHEPEA
98	ACP57 Mouse IFG conjugate	mdmrvpaqllgllllwlrgrcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllrfevlkdnqai

SEQ ID NO.	Name	Sequence
		snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGP GPAGMKGLPGSEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMS WVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYL QMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsggggsQV QLQESGGGLAQAGGSLSLSCAASGFTVSNVMAWYRQTPGKQREFV AIINSVGSTNYADSVKGRFTISRDNAAKNTVYLQMNNLKPEDTAVYV CNRNFDRIYWGQGTQVTVSSHHHHHHEPEA
99	ACP58 Mouse IFG conjugate	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllrfevlkdnqai snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGP GPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllr fevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkr krsrcSGGPAGMKGLPGSEVQLVESGGGLVQPGNSLRRLSCAASGFT FSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAA KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsggggs ggggsQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDMSWYRQAPGK QRELVARITRGGTISYDDSVKGRFTISRDNAAKNTVYLQMNSLKPEDT GVYYCNALYGTDYWGKGTQVTVSSHHHHHHEPEA
100	ACP59 Mouse IFG conjugate	mdmrvpaqlglglllwlrgarcQVQLQESGGGLAQAGGSLSLSCAASGFTVSNV MAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDNAAKNTV YLQMNNLKPEDTAVYVCNRNFDRIYWGQGTQVTVSSggggsgggsgggsggg gsEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAV YYCTIGGSLSVSSQGTLLTVSSGGPGPAGMKGLPGShgtviesleslnnyf nssgidveekslfdiwrnwqkdgmkilqsqiisfyllrfevlkdnqaisnnisvieshlittffsnkakk dafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPAGMKGLPGShgt viesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllrfevlkdnqaisnnisvieshl ittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGPAGMK GLPGSEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKFGMSWVRQAP GKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRP EDTAVYYCTIGGSLSVSSQGTLLTVSSHHHHHHEPEA
101	ACP60 Mouse IFG conjugate	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllrfevlkdnqai snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGP GPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllr fevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkr krsrcSGGPAGMKGLPGSEVQLVESGGGLVQPGNSLRRLSCAASGFT FSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAA KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsggggs ggggsQVQLQESGGGLAQAGGSLSLSCAASGFTVSNVMAWYRQTPG KQREFVAIINSVGSTNYADSVKGRFTISRDNAAKNTVYLQMNNLKPED TAVYVCNRNFDRIYWGQGTQVTVSSHHHHHHEPEA
102	ACP61 Mouse IFG conjugate	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGNSLRRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgmkilqsqiisfyllrfevlkdnqai

SEQ ID NO.	Name	Sequence
		snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGP GPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyll fevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkr krsrcSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFT FSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsggggs ggggsEVQLVESGGGLVQPGSLRLSCAASGFTFSSYAMSWVRQAPG KGLEWVSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAE DTAVYYCARGVGAFRPYRKHEWGQGTLLTVSRggggsgggsggggsSSE LTQDPAVSVALGQTVRITCQGDSLRSYYASWYQQKPGQAPVLVIYG KNNRPSGIPDRFSGSSSGNTASLTTTGAQAEDADYYCNSSPFEHNL VVFGGGTKLTVLHHHHHHHEPEA
103	ACP63 Anti-FN CGS-2 scFv	mdmrvpaqllglllwlrgarcEVQLVESGGGLVQPGSLRLSCAASGFTFSSY AMSWVRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISRDN SKNT LYLQMNSLRAEDTAVYYCARGVGAFRPYRKHEWGQGTLLTVSRgg ggsgggsggggsSSELTQDPAVSVALGQTVRITCQGDSLRSYYASWYQQ KPGQAPVLVIYGKNNRPSGIPDRFSGSSSGNTASLTTTGAQAEDAD YYCNSSPFEHNLVVFGGGTKLTVLHHHHHHHEPEA
104	ACP69 Mouse IFG conjugate	mdmrvpaqllglllwlrgarcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllfevlkdnqai snnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcSGGP GPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMS WVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYL QMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLP GShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyllfevlkdnqaisnnis vieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkrkrsrcHHHHHH EPEA
105	ACP70 Mouse IFG conjugate	mdmrvpaqllglllwlrgarchgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqi sfyllfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpes slrkrkrsrcSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAAS GFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISR DN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGP GPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqiisfyll fevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpesslrkr krsrcSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFT FSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSHHHHHHHE PEA
106	ACP71 Mouse IFG conjugate	mdmrvpaqllglllwlrgarchgtviesleslnnyfnssgidveekslfdiwrnwqkdgdmkilqsqi sfyllfevlkdnqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirrvhqlpes slrkrkrsrcSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHFQKGLVIA FSQYLQKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGD KLCAIPNLRENYGELADCTKQEPERNECFQHKDDNPSLPPFERPE AEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEIL TQCCAEADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERA KAWAVARLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDR AELAKYMCENQATISSKLQTCCKPLKKAHCLSEVEHDTMPADLP AIAADFVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRL

SEQ ID NO.	Name	Sequence
		AKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLY EKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTKCCTLPE DQRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFS ALTVDETYVPKEFKAETFTFHSDICTLPEKEKQIKKQTALAELVKHKP KATAEQLKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDA LASGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfldiwmwqkdgmkil qsqiisfyrlfevlkdnqaisnnisvieshlittffnskakkdafmsiakfevnnpqvqrqafnelirvvhq llpesslrkrkrsrcSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHFHFKGL VLIAFSQYLQKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHT LFGDKLCAIPNLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPF ERPEAEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQY NEILTQCCAEADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFG ERAFKAWAVARLSQTFPNADFAEITKLATDLTKVNKECCHGDLLEC ADDRAELAKYMCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMP ADLPAIAADFVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSL LLRLAKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTN CDLYEKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTKC CTLPEDQRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVER RPCFSALTVDETYVPKEFKAETFTFHSDICTLPEKEKQIKKQTALAEL VKHKPKATAEQLKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTR CKDALAHHHHHHEPEA
107	ACP72 Mouse IFG conjugate	mdmrvpaqlglglllwlrgarcEAHKSEIAHRYNDLGEQHFHFKGLVLIAFSQYLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAEA DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAKAWAVA RLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAKY MCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEY GFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTKCCTLPEQRLPCV EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHSDICTLPEKEKQIKKQTALAELVKHKPKATAEQL KTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGShgtviesleslnnyfnssgidveekslfldiwmwqkdgmkilqsqiisfyrlfev lkdqaisnnisvieshlittffnskakkdafmsiakfevnnpqvqrqafnelirvvhqllpesslrkrkrsrc cSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHFHFKGLVLIAFSQYL QKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAI PNLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFERPEAEAMC TSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAE ADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAKAWAV ARLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAK YMCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADF VEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEY GFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTKCCTLPEQRLPC VEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDET YVPKEFKAETFTFHSDICTLPEKEKQIKKQTALAELVKHKPKATAEQ LKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPG PAGMKGLPGShgtviesleslnnyfnssgidveekslfldiwmwqkdgmkilqsqiisfyrlfe

SEQ ID NO.	Name	Sequence
		vlkdnqaisnnisvieshlittffsnkakddafmsiakfevnnpqvrqafnelirvvhqllpesslrkrkr srcHHHHHHEPEA
108	ACP73 Mouse IFG conjugate	mdmrvpaqllgllllwlgarcEAHKSEIAHRYNDLGEQHFHGLVLIASFQYQLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFHQHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAEA DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAKAWAVA RLSQTFPNADFAETTKLATDLTKVNKECCHGDLLECADDRAELAKY MCENQATISSKLQTCCKDLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEYG FQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCCCTLPEDQRLPCV EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHSDICTLPEKEKQIKKQTAALAEVVKHKPKATAEQ KTVMDDFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGShgtviesleslnnyfnssgidveekslfidiwmwqkdgmklqsqisfyrfev lkdnqaisnnisvieshlittffsnkakddafmsiakfevnnpqvrqafnelirvvhqllpesslrkrkr cSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHFHGLVLIASFQYQL QKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAI PNLRENYGELADCCTKQEPERNECFHQHKDDNPSLPPFERPEAEAMC TSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAE ADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAKAWAV ARLSQTFPNADFAETTKLATDLTKVNKECCHGDLLECADDRAELAK YMCENQATISSKLQTCCKDLLKKAHCLSEVEHDTMPADLPAIAADF VEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT TLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEY GFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCCCTLPEDQRLPC VEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDET YVPKEFKAETFTFHSDICTLPEKEKQIKKQTAALAEVVKHKPKATAEQ LKTVMDDFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPG PAGMKGLPGShgtviesleslnnyfnssgidveekslfidiwmwqkdgmklqsqisfyrfe vlkdnqaisnnisvieshlittffsnkakddafmsiakfevnnpqvrqafnelirvvhqllpesslrkrkr srcSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHFHGLVLIASFQY LQKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLC AIPNLRENYGELADCCTKQEPERNECFHQHKDDNPSLPPFERPEAEA MCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQC CAEADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAK WAVARLSQTFPNADFAETTKLATDLTKVNKECCHGDLLECADDRAE LAKYMCENQATISSKLQTCCKDLLKKAHCLSEVEHDTMPADLPAI AADFVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAK KYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEK LGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCCCTLPED QRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSAL TVDETYVPKEFKAETFTFHSDICTLPEKEKQIKKQTAALAEVVKHKPK ATAEQLKTVMDDFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDAL AHHHHHHEPEA
109	ACP74 Mouse IFG conjugate	mdmrvpaqllgllllwlgarcEAHKSEIAHRYNDLGEQHFHGLVLIASFQYQLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFHQHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAEA

SEQ ID NO.	Name	Sequence
		DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFKAWAVA RLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAKY MCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEYG FQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCTLPCV EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHSDICTLPEKEKQIKKQTALAELVKHKPKATAEQL KTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGShgtviesleslnnyfnssgidveekslfldiwnwqkdgmkilqsqiisfyllfev lkdqaisnnisvieshlittffsnkakkdafmsiakfevnpqvrqafnelirvqhllpesslrkrkrsrc SGGPGPAGMKGLPGSggggsEAHKSEIAHRYNDLGEQHFHFKGLVLI AF SQYLQKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGD KLCAIPNLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFERPE AEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEIL TQCCA EADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAF KAWAVARLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADD AELAKYMCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLP AIAADFVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRL AKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLY EKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCTLPE DQRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFS ALTVDETYVPKEFKAETFTFHSDICTLPEKEKQIKKQTALAELVKHKP KATAEQLKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDA LAggggsSGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfldiwnwqkdg dmkilqsqiisfyllfevlkdqaisnnisvieshlittffsnkakkdafmsiakfevnpqvrqafneli rvvqhllpesslrkrkrsrcSGGPGPAGMKGLPGSEAHKSEIAHRYNDLGEQHF KGLVLI AF SQYLQKCSYDEHAKLVQEVTDFAKTCVADESAANCDK SLHTLFGDKLCAIPNLRENYGELADCCTKQEPERNECFLOHKDDNPSL PPFERPEAEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAE QYNEILTQCCA EADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQK FGERAFKAWAVARLSQTFPNADFAEITKLATDLTKVNKECCHGDL LECADDRAELAKYMCENQATISSKLQTCCDKPLLKKAHCLSEVEHDT MPADLPAIAADFVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSV SLLRLAKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKT NCDLYEKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGT CKCTLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVE RRPCFSALTVDETYVPKEFKAETFTFHSDICTLPEKEKQIKKQTALAE LVKHKPKATAEQLKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLV TRCKDALAHHHHHHEPEA
110	ACP75 Mouse IFG conjugate	mdmrvpaqlglglwlgarcEAHKSEIAHRYNDLGEQHFHFKGLVLI AF SQYLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCA E DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFKAWAVA RLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAKY MCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEYG FQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCTLPCV

SEQ ID NO.	Name	Sequence
		EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHS DICTLPEKEKQIKKQ TALAE LVKHKPKATAEQ L KTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGShgtviesleslnnyfnssgidveekslfdiwrnwqkdgm kilqsqiisfylrlfev lkd nqaisnnisvieshlittffsnkakkdafmsiakfevnnpqvqrqafnelirvvhqllpesslrkrkrsr cSGGPGPAGMKGLPGSggggsggggsEAHKSEIAHRYNDLGEQHFKGLV LIAFSQYLQKCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTL FGDKLCAIPNLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFE RPEAEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYN EILTQCCAEADKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGE RAFKAWAVARLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECA DDRAELAKYMCENQATISSKLQTCCKPLLKKAHCLSEVEHDTMPA DLPAAAD FVEDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLL RLAKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCD LYEKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCT LPEDQRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPC FSALTVDETYVPKEFKAETFTFHS DICTLPEKEKQIKKQ TALAE LVKH KPKATAEQ LKTVMD DFAQFLDTCCKAADKDTCFSTEGPNLVTRCKD ALAgggsggggsSGGPGPAGMKGLPGShgtviesleslnnyfnssgidveekslfdiwr nwqkdgm kilqsqiisfylrlfev lkd nqaisnnisvieshlittffsnkakkdafmsiakfevnnpqv qrqafnelirvvhqllpesslrkrkrsrcSGGPGPAGMKGLPGSEAHKSEIAHRYNDL GEQHFKGLV LIAFSQYLQKCSYDEHAKLVQEVTDFAKTCVADESA A NCDKSLHTLFGDKLCAIPNLRENYGELADCCTKQEPERNECFLOHKD DNPSLPPFERPEAEAMCTSFKENPTTFMGHYLHEVARRHPYFYAPEL LYYAEQYNEILTQCCAEADKESCLTPKLDGVKEKALVSSVRQRMK SSMQKFGERAFKAWAVARLSQTFPNADFAEITKLATDLTKVNKEC HGDLECA DDRAELAKYMCENQATISSKLQTCCKPLLKKAHCLSE VEHDTMPADLPAAAD FVEDQEVCKNYAEAKDVFLGTFLYEYSRRH PDYSVSLLLRLAKKYEATLEKCCAEANPPACYGTVLAEFQPLVEEPK NLVKTNCDLYEKLGEYGFQNAILVRYTQKAPQVSTPTLVEAARNLG RVGTCKCTL PEDQRLPCVEDYLSAILNRVCLLHEKTPVSEHVTKCCS GSLVERRPCFSALTVDETYVPKEFKAETFTFHS DICTLPEKEKQIKKQ TALAE LVKHKPKATAEQ LKTVMD DFAQFLDTCCKAADKDTCFSTE GPNLVTRCKDALAHHHHHHEPEA
111	ACP78 Mouse IFG conjugate	mdmrvpaqlglglwlrgrcEVQLVESGGGLVQPGNSLR LSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGT LTVTVSSggggsgggsgggg shgtviesleslnnyfnssgidveekslfdiwrnwqkdgm kilqsqiisfylrlfev lkd nqaisnnisvi eshlittffsnkakkdafmsiakfevnnpqvqrqafnelirvvhqllpesslrkrkrsrcggggsgggsgg gggsEVQLVESGGGLVQPGNSLR LSCAASGFTFSKFGMSWVRQAPGK GLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMNSLRPED TAVYYCTIGGSLSVSSQGT LTVTVSSggggsgggsggggshgtviesleslnnyfnss gidveekslfdiwrnwqkdgm kilqsqiisfylrlfev lkd nqaisnnisvieshlittffsnkakkdaf msiakfevnnpqvqrqafnelirvvhqllpesslrkrkrsrcggggsgggsggggsEVQLVESG GGLVQPGNSLR LSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISG GRDTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGG LSVSSQGT LTVTVSSHHHHHHEPEA
112	ACP134 Mouse IFG	mdmrvpaqlglglwlrgrcEVQLVESGGGLVQPGNSLR LSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGT LTVTVSSGGPGPAGMK

[illegible]

SEQ ID NO.	Name	Sequence
		agqytchkggetlshshlllhkengiwsteilkknfkntflkceapnysgrftcswlvqrnmldkfnikss ssspdsravtcgmaslsaekvtldqrdyekysvscqedvtcptaeetlpielalearqqnkyenyystsffird iikpdpknqmkplknsqvevswepdswstphsyfslkffvriqrkkekmketeegcnqkgafive ktstevqckggncvcvqaqdrynsscskwacvpcrvrsgggsgggsgggsgggsvipvsgparclsqs mllktiddmvtareklkhysctaedidheditrdqstlktclplelhknesclatretssttrgscldppqktsl mmtlclgsiyedlkmyqtefqainaalqnhnhqqiildkgmlvaidelmqslnhngetlrqkppvgea dpyrvkmkclilhafstrvvtinrmgylssaSGGPGPAGMKGLPGSEVQLVESGGG LVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSLS VSSQGTTLTVSSHHHHHH
117	ACP37 Mouse IL-12 conjugate	mdmrvpaqllgllllwlgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNAAKTTLVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSggggsgggsgggg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTTLTVSSSGGPGPAGMKGLPGSmwelekdvyvve vdwtpdapgetvnlctdtpcedditwtsdqrhgvigsgkltitvkefldagqytchkggetlshshllhk kengiwsteilkknfkntflkceapnysgrftcswlvqrnmldkfnikssspdsravtcgmaslsaekv tldqrdyekysvscqedvtcptaeetlpielalearqqnkyenyystsffirdiikpdpknqmkplknsq evswepdswstphsyfslkffvriqrkkekmketeegcnqkgafivektstevqckggncvcvqaqdr ynsscskwacvpcrvrsgggsgggsgggsgggsvipvsgparclsqsmllktiddmvtareklkhy sctaedidheditrdqstlktclplelhknesclatretssttrgscldppqktslmmtlclgsiyedlkmyqte fqainaalqnhnhqqiildkgmlvaidelmqslnhngetlrqkppvgeadpyrvkmkclilhafstrv tinrmgylssaSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLS ASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTI SRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTTLTVSSHH HHHH
118	ACP79 Mouse IL-12 conjugate	mdmrvpaqllgllllwlgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNAAKTTLVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSggggsgggsgggg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTTLTVSSSGGPGPAGMKGLPGSmwelekdvyvve vdwtpdapgetvnlctdtpcedditwtsdqrhgvigsgkltitvkefldagqytchkggetlshshllhk kengiwsteilkknfkntflkceapnysgrftcswlvqrnmldkfnikssspdsravtcgmaslsaekv tldqrdyekysvscqedvtcptaeetlpielalearqqnkyenyystsffirdiikpdpknqmkplknsq evswepdswstphsyfslkffvriqrkkekmketeegcnqkgafivektstevqckggncvcvqaqdr ynsscskwacvpcrvrsgggsgggsgggsgggsvipvsgparclsqsmllktiddmvtareklkhy sctaedidheditrdqstlktclplelhknesclatretssttrgscldppqktslmmtlclgsiyedlkmyqte fqainaalqnhnhqqiildkgmlvaidelmqslnhngetlrqkppvgeadpyrvkmkclilhafstrv tinrmgylssaSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLS ASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTI SRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTTLTVSSHH HHHH
119	ACP80 Mouse IL-12 conjugate	mdmrvpaqllgllllwlgarcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTTLTVSSSGGPGPAGMK GLPGSmwelekdvyvvevdwtpdapgetvnlctdtpcedditwtsdqrhgvigsgkltitvkefld agqytchkggetlshshlllhkengiwsteilkknfkntflkceapnysgrftcswlvqrnmldkfnikss ssspdsravtcgmaslsaekvtldqrdyekysvscqedvtcptaeetlpielalearqqnkyenyystsffird

[illegible]

[illegible]

SEQ ID NO.	Name	Sequence
		SRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSH HHHHEPEA
125	ACP38 IL-2 conjugate	mnmrvpaqlglglllwlrgarcptssstsktkqlehlldlqmilnginnyknpkltmrltlfkfympkka telkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgssettfmceyadetativeflnrwifc qsiiistltSGGGPGPAGMKGLPGSEVQLVESGGGLVQPGGSLRLSCAASGF TFSSYTLAWVRQAPGKGLEWVAADIDSSSYTYSPDTRGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARSDNDALDYWGQTTVTVSSG GGGSGGGSGGGGSDIOMTQSPSSLSASVGDRVTITCKASQNVGTN VGWYQQKPKGKAPKALIYSASFYSGVPSRFSGSGSGTDFTLTISLQP EDFATYYCQYYITYPYTFGGGTKVEIKggggsgggsgggsgggsgggEVQLVESG GGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGS GRDTLYAESVKGRFTISRDNKTTLYLQMNSLRPEDTAVYYCTIGGS LSVSSQGTLLTVSSggggsgggsgggsgggsgggVVQLQESGGGLVQAGGSLRLSC AASGRIFSIDIMSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTIS RDNAKNTVYLQMNSLKPEDTGVIYCINALYGTDYWGKGTQVTVSS HHHHHH
126	ACP39 IL-2 conjugate	mnmrvpaqlglglllwlrgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSI DSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNKNTVY LQMNSLKPEDTGVIYCINALYGTDYWGKGTQVTVSSSGGGPGPAGM KGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQA PGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNKTTLYLQMNSLR PEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGGPGPAGMKGLPGSEVQL VESGGGLVQPGGSLRLSCAASGFTFSYTLAWVRQAPGKGLEWVAAD IDSSSYTYSPDTRGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAR DSNDALDYWGQTTVTVSSGGGGSGGGSGGGGSDIOMTQSPSS LSASVGDRVTITCKASQNVGTNVGWYQQKPKGKAPKALIYSASFYSG VPSRFSGSGSGTDFTLTISLQPEDFATYYCQYYITYPYTFGGGTKV EIKSGGGPGPAGMKGLPGSaptssstsktkqlehlldlqmilinginnyknpkltmrltlfky mpkkatelkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgssettfmceyadetativefln rwifcqsiiistltHHHHHH**
127	ACP40 IL-2 conjugate	mnmrvpaqlglglllwlrgarceldddppeiphatfkamaykegtmlnceckrgfrriksgslmictg nsshswdnqcctssatnttkqvtpqpeeekerkttmqspmqpvdaqslpgchrepppweneate riyhfvvgqmvyyqcvqgyrallhrgpaeavckmthgktrwtqpqlietgemetsqfpgeekpqaspe grpesetclvtutdfqiqtemaatmetstiftteyqggsgggsgggsgggsgggsgggsgggsgggSG GGPGPAGMKGLPGSaptssstsktkqlehlldlqmilinginnyknpkltmrltlfkfympkkate lkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgssettfmceyadetativeflnrwifcqs iistltHHHHHH
128	ACP41 IL-2 conjugate	mnmrvpaqlglglllwlrgarcptssstsktkqlehlldlqmilnginnyknpkltmrltlfkfympkka telkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgssettfmceyadetativeflnrwifc qsiiistltSGGGPGPAGMKGLPGSGggggsgggsgggsgggsgggsgggsgggsgggseledddp eiphatfkamaykegtmlnceckrgfrriksgslmictgnsshswdnqcctssatnttkqvtpqpe eqkerktmqspmqpvdaqslpgchrepppweneaterihfvgqmvyyqcvqgyrallhrgpae svckmthgktrwtqpqlietgemetsqfpgeekpqaspegresetclvtutdfqiqtemaatmetstift eyqHHHHHH
129	ACP42 IL-2 conjugate	mnmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgggsggg seledddppeiphatfkamaykegtmlnceckrgfrriksgslmictgnsshswdnqcctssatntt kqvtpqpeeekerkttmqspmqpvdaqslpgchrepppweneaterihfvgqmvyyqcvqgy rallhrgpaeavckmthgktrwtqpqlietgemetsqfpgeekpqaspegresetclvtutdfqiqtema

SEQ ID NO.	Name	Sequence
		atmetsiftteyqgggsgggsgggsgggsgggsgggsgggsgGGGPGPAGMKGLPGSapt ssstktkqlqlehlldlqmilnginnyknpkltmrmtfkfymppkkatelkhqlcleeelkpleevlnlaqsk nfhlrprdlisininvivlelkgssettfmceyadetativeflnrwitfcqsiistltHHHHHHH
130	ACP43 IL-2 conjugate	mmdmrvpaqlglglllwlgarcaptsstktkqlqlehlldlqmilnginnyknpkltmrmtfkfymppkka telkhqlcleeelkpleevlnlaqsknfhlrprdlisininvivlelkgssettfmceyadetativeflnrwitfc qsiistltSGGPGPAGMKGLPGSgggsgggsgggsgggsgggsgggsgggsgggseledddp eiphatfkamaykegtmlnceckrgfrriksgslymcltgnsshswdnqcqtssatmnttkqvtpqe eqkerkttemqspmqpvvdqaslpghcrepppweneateriyhfvgqmvyvycvgyralhrpgae svckmthgktrwtqpqlctgemetsqfpgeekpqaspegrpesetclvttdfqiqtemaatmetsifft eyqgggsgggsgggsgggsgEVQLVESGGGLVQPGNSLRSLSCAASGFTFSKFG MSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKTTTL YLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVTVSSHSHHHHHH
131	ACP44 IL-2 conjugate	mmdmrvpaqlglglllwlgarcaptsstktkqlqlehlldlqmilnginnyknpkltmrmtfkfymppkka telkhqlcleeelkpleevlnlaqsknfhlrprdlisininvivlelkgssettfmceyadetativeflnrwitfc qsiistltSGGPGPAGMKGLPGSgggsgggsgggsgggsgggsgggsgggsgggseledddp eiphatfkamaykegtmlnceckrgfrriksgslymcltgnsshswdnqcqtssatmnttkqvtpqe eqkerkttemqspmqpvvdqaslpghcrepppweneateriyhfvgqmvyvycvgyralhrpgae svckmthgktrwtqpqlctgemetsqfpgeekpqaspegrpesetclvttdfqiqtemaatmetsifft eyqSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRSLSCAASGFTF SKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVTVSSHSHHHHHH
132	ACP45 IL-2 conjugate	mmdmrvpaqlglglllwlgarcEVQLVESGGGLVQPGNSLRSLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVTVSSSGGPGPAGMK GLPGSEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPG KGLEWVAADSSSYTYSPDTPVRGRFTISRDNAKNSLYLQMNSLRAED TAVYYCARDNWDALDYWGQGTITVTVSSGGGGSGGGGSGGGGSDI QMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQQKPKGAPKALI YSASFRTYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPY TFGGGTKEIKgggsgggsgggsgggsgggsgggsgggsgggsgGGGPGPAGMKGLPG Saptssstktkqlqlehlldlqmilnginnyknpkltmrmtfkfymppkkatelkhqlcleeelkpleevlnl aqsknfhlrprdlisininvivlelkgssettfmceyadetativeflnrwitfcqsiistltHHHHHHH
133	ACP46 IL-2 conjugate	mmdmrvpaqlglglllwlgarcaptsstktkqlqlehlldlqmilnginnyknpkltmrmtfkfymppkka telkhqlcleeelkpleevlnlaqsknfhlrprdlisininvivlelkgssettfmceyadetativeflnrwitfc qsiistltSGGPGPAGMKGLPGSgggsgggsgggsgggsgggsgggsgggsgggsgEVQLV ESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLEWVAAI DSSSYTYSPDTPVRGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR DSNWDALDYWGQGTITVTVSSsggpgpagmkglpgsDIQMTQSPSSLSASV GDRVTITCKASQNVGTNVGWYQQKPKGAPKALITYSASFRTYSGVPSR FSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPTYTFGGGTKEIKggg sgggsgggsgggsgEVQLVESGGGLVQPGNSLRSLSCAASGFTFSKFGMSWVR QAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNS LRPEDTAVYYCTIGGSLSVSSQGTLTVTVSSgggsgggsgggsgggsgVQLQE SGGGLVQAGGSLRLSCAASGRIFSIDIMSWYRQAPGKQRELVARITR GGTISYDDSVKGRFTISRDNAKNTVYLQMNSLPEDTGVIYCNALEY GTDYWGKGQTQVTVSSHSHHHHHH
134	ACP47 IL-2 conjugate	mmdmrvpaqlglglllwlgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNAKNTVY LQMNSLPEDTGVIYCNALEYGTDYWGKGQTQVTVSSgggsgggsgggsgggsg saptssstktkqlqlehlldlqmilnginnynknokltmrmtfkfymppkkatelkhqlcleeelkpleevlnl

[illegible]

SEQ ID NO.	Name	Sequence
		NSLYLQMNSLRAEDTAVYYCARDSDNWDALDYWGQGTTVTVSSGG GGSGGGGGSGGGGSDIOMTQSPSSLSASVGDRTITCKASQNVGTNV GWYQQKPGKAPKALIYSASFYSGVPSRFSGSGSGTDFLTISLQPE DFATYYCQQYYTYPTYTFGGGTKEIKSGGPGPAGMKGLPGSaptssstk ktqlqlehlldlqmilnginnyknkplrmltfkfympkkatelkhlqleceelkpleevlnlaqsknfhlr prdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltHHHHHHH
139	ACP94 IL-2 conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSsgsgsgsgsgsgsg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSsgsgsgsgsgsgsgsgsEVQLVESGGGLVQ PGGSLRLSCAASGFTFSSTYTLAWVRQAPGKGLEWVAIDSSSYTYSP DTRGRFTISRDNANKNSLYLQMNSLRAEDTAVYYCARDSDNWDALD YWGQGTTVTVSSGGGGSGGGGSGGGGSDIOMTQSPSSLSASVGDRT ITCKASQNVGTNVGWYQQKPGKAPKALIYSASFYSGVPSRFSGSG SGTDFLTISLQPEDFATYYCQQYYTYPTYTFGGGTKEIKSGGPGPA GMKGLPGSaptssstkktqlqlehlldlqmilnginnyknkplrmltfkfympkkatelkhlqle eelkpleevlnlaqsknfhlrprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltHH HHHH
140	ACP95 IL-2 conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSsgsgsgsgsgsgsg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLPGSaptssstkktqlqle hlldlqmilnginnyknkplrmltfkfympkkatelkhlqleceelkpleevlnlaqsknfhlrprdlisn invivlelkgsettfmceyadetativeflnrwitfcqsiistltHHHHHHH
141	ACP96 IL-2 conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSSGGPGPAGM KGLPGSaptssstkktqlqlehlldlqmilnginnyknkplrmltfkfympkkatelkhlqleceelk pleevlnlaqsknfhlrprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPG PAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSW VRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQM NSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSHHHHHHH
142	ACP97 IL-2 conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSggggsgggsgggsg sEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLPGSaptssstkktqlqle hlldlqmilnginnyknkplrmltfkfympkkatelkhlqleceelkpleevlnlaqsknfhlrprdlisn invivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQL VESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSS ISGSGRDTLYAESVKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYCTI GGSLSVSSQGTLLTVSSHHHHHHH
143	ACP99 IL-2	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSggggsgggsgggsg

SEQ ID NO.	Name	Sequence
	conjugate	saptssstkkqlqlehlldlqmilnginnyknkplrmltkfymppkkelkhlcleeelkpleevnl aqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPGPAGM KGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQA PGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMNLSLR PEDTAVYYCTIGGSLSVSSQGTLLTVSSH HHHHHH
144	ACPI00 IL-2 conjugate	mdmrvpaqlglglllwlrgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDN AKNTVY LQMNLSLPEDTG VYYCNALYGT DYWGKGTVTVSSggggsgggsgggsggg saptssstkkqlqlehlldlqmilnginnyknkplrmltkfymppkkelkhlcleeelkpleevnl aqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltHH HHHHHH
145	ACPI01 IL-2 conjugate	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknkplrmltkfymppkka telkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfc qsiistltSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGF TFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMNLSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSH HHHHHH H
146	ACPI02 IL-2 conjugate	mdmrvpaqlglglllwlrgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDN AKNTVY LQMNLSLPEDTG VYYCNALYGT DYWGKGTVTVSSGGGPGPAGM KGLPGSaptssstkkqlqlehlldlqmilnginnyknkplrmltkfymppkkelkhlcleeelk pleevlnlaqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPG PAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSW VRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMN NSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgggsgggsgggsg ggggsgggsgEVQLVESGGGLVQPGGSLRLSCAASGFTFSYTLAWVRQA PGKGLEWVAADSSSYTYSPD TVRGRFTISRDN AKNSLYLQMNLSRA EDTAVYYCARDSDNWDALDYWGQGT TVTVSSGGGGSGGGSGGGSGGGG SDIQMTQSPSSLSASVGDRVTTITCKASQNVGTNVGWYQQKPGKAPK ALIYSASFRYSGVPSRFSGSGSGDFTLTISLQPEDFATYYCQYYTY YPYTFGGGTKEIKHH HHHHHH
147	ACPI03 IL-2 conjugate	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknkplrmltkfymppkka telkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfc qsiistltSGGPGPAGMKGLPGSggggsgggsgggsgggsgggsgggsgggsgggsgEVQLV ESGGGLVQPGGSLRLSCAASGFTFSYTLAWVRQAPGKGLEWVAAD DSSSYTYSPD TVRGRFTISRDN AKNSLYLQMNLSRAEDTAVYYCAR DSNWDALDYWGQGT TVTVSSGGGGSGGGSGGGSGGGSGDIQMTQSPSS LSASVGDRVTTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFRYS GVPSRFSGSGSGDFTLTISLQPEDFATYYCQYYTYYPYTFGGGTKEIK EIKggggsgggsgggsgggsgEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFG MSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTL YLQMNLSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgggsgggsg QVQLQESGGGLAQAGGSLSLSCAASGFTVSNVMAWYRQTPGKQR EFVAIINSVGSTNYADSVKGRFTISRDN AKNTVYLQMNLSLPEDTAV YVCNRFNFDRIYWGQGTQVTVSSH HHHHHH
148	ACPI04 IL-2 conjugate	mdmrvpaqlglglllwlrgarcQVQLQESGGGLAQAGGSLSLSCAASGFTVSNV VMAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDN AKNTV YLQMNLSLPEDTAVYYVCNRFNFDRIYWGQGTQVTVSSaptssstkkqlqle hlldlqmilnginnyknkplrmltkfymppkkelkhlcleeelkpleevlnlaqsknfhlprdlisn invivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPGPAGMKGLPGSEVQL VESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSS

SEQ ID NO.	Name	Sequence
		ISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTI GGSLSVSSQGTTLVTVSSggggsgggsgggsgggsgggsgggsgggsgggEVQLVESG GGLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLEWVAADSS SYTYSPTVVRGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDNSN WDALDYWGQGTTLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSA SVGDRVTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFYSGVPSR FSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPTYTFGGGTKVEIK HHHHHH
149	ACP105 IL-2 conjugate	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYT LAWVRQAPGKGLEWVAADSSSYTYSPTVVRGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCARDNSNWDALDYWGQGTTLVTVSSGGGGSG GGGSGGGGSDIQMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQ QKPGKAPKALIYSASFYSGVPSRFSGSGSGTDFTLTISLQPEDFATY YCQQYYTYPTYTFGGGTKVEIKggggsgggsgggsgggsgggsgggsgggsgggSGGP GPAGMKGLPGSaptssstktqlqlehlldlqmilnginnyknpltrmltkfymppkkelkh lqcleelkpleevlnlaqsknfhlprdlisninvilelkgsettfmceyadetativeflnrwitfcqsiistl tSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSK FGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAKT TLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTTLVTVSSggggsgggsggg gsQVQLQESGGGLAQAGGSLSLSCAASGFTVSNSVMAWYRQTPGKQ REFVAIINSVGSTNYADSVKGRFTISRDNAKNTVYLQMNNLKPEDTA VYVCNRRNFDRYWGQGTQVTVSSHHHHHH
150	ACP106 IL-2 conjugate	mdmrvpaqlglglllwlrgarcQVQLQESGGGLAQAGGSLSLSCAASGFTVSNS VMAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDNAKNTV YLQMNNLKPEDTAVYVCNRRNFDRYWGQGTQVTVSSggggsgggsgggsggg gsEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTTLVTVSSGGPGPAGMKGLPGSEVQLVESGG GLVQPGGSLRLSCAASGFTFSSYTLAWVRQAPGKGLEWVAADSSS YTYSPTVVRGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDNSN DALDYWGQGTTLVTVSSGGGGSGGGGSGGGGSDIQMTQSPSSLSASV GDRVTITCKASQNVGTNVGWYQQKPGKAPKALIYSASFYSGVPSR FSGSGSGTDFTLTISLQPEDFATYYCQQYYTYPTYTFGGGTKVEIKggg sgggsgggsgggsgggsgggsgggsgggsgggSGGPGPAGMKGLPGSaptssstktqlqlehlldl lqmilnginnyknpltrmltkfymppkkelkhkhqcleelkpleevlnlaqsknfhlprdlisninv vilelkgsettfmceyadetativeflnrwitfcqsiistlHHHHHH
151	ACP107 IL-2 conjugate	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYT LAWVRQAPGKGLEWVAADSSSYTYSPTVVRGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCARDNSNWDALDYWGQGTTLVTVSSGGGGSG GGGSGGGGSDIQMTQSPSSLSASVGDRVTITCKASQNVGTNVGWYQ QKPGKAPKALIYSASFYSGVPSRFSGSGSGTDFTLTISLQPEDFATY YCQQYYTYPTYTFGGGTKVEIKggggsgggsgggsgggsgggsgggsgggsgggEVQ LVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTTLVTVSSSGPGPAGMKGLPGSaptssstktqlqlehlldlq milnginnyknpltrmltkfymppkkelkhkhqcleelkpleevlnlaqsknfhlprdlisninvile lkgsettfmceyadetativeflnrwitfcqsiistlggggsgggsgggsgggsgggsgggsgggsgggQVQLQESGGGLA QAGGSLSLSCAASGFTVSNSVMAWYRQTPGKQREFVAIINSVGSTNY ADSVKGRFTISRDNAKNTVYLQMNNLKPEDTAVYVCNRRNFDRYWG GOGTOVTVSSHHHHHH

SEQ ID NO.	Name	Sequence
152	ACP108 IL-2 conjugate	mdmrvpaqllgllllwlgarcQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANTVY LQMNSLKPEDTGVIYCNALYGTDYWGKGTQVTVSSggggsgggsgggg saptssstktqlqlehlldlqmilnginnyknpltrmltkfymppkkelkhlcleeelkpleevlnl aqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfcqsiistltSGGPGPAGM KGLPGSRgetgpaaPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFG MSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNANTL YLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVTVSSggggsgggsgggsgggg ggggsgggsgggsggggEVQLVESGGGLVQPGGSLRLSCAASGFTFSYTLAW VRQAPGKGLEWVAADSSSYTYSPDTRGRFTISRDNANKSLYLQM NSLRAEDTAVYYCARDNWDALDYWGQGTITVTVSSGGGGSGGGG SGGGSDIQMTQSPSSLSASVGDRTTTCASQNVGTNVGWYQQKP GKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQ QYYTYPTYTFGGGTKEIKHHHHHH
153	ACP117 Anti-FN CGS-2 scFv	mdmrvpaqllgllllwlgarcEVQLVESGGGLVQPGGSLRLSCAASGFTFSY AMSWVRQAPGKGLEWVSAISGSGSTYYADSVKGRFTISRDNANKT LYLQMNSLRAEDTAVYYCARGVGAFRPYRKHEWGQGTLLTVTVSRgg ggsgggsgggsggggSSELTQDPAVSVALGQTVRITCQGDSLRSYYASWYQQ KPGQAPVLVIYGKNNRPSGIPDRFSGSSSGNTASLTITGAQAEDAD YYCNSSPFEHNLVVFGGGTKLTVLHHHHHHHEPEA
154	ACP118 NARA1 Vh/Vl non- cleavable	mdmrvpaqllgllllwlgarcQVQLQQSGAELVRPGTSVKVSKASGYAFTN YLIEWVKQRPGQGLEWIGVINPGSGGTNYNEKFKGKATLTADKSSST AYMQLSSLTSDDSAVYFCARWRGDGYAYFDVWGAGTTVTVSSgg ggsgggsgggsggggDIVLTQSPASLAVSLGQRATISCKASQSVDDYDGSYM NWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHPVEEE DAATYYCQQSNEDPYTFGGGTKEIKHHHHHHHEPEA
155	ACP119 NARA1 Vh/Vl cleavable	mdmrvpaqllgllllwlgarcQVQLQQSGAELVRPGTSVKVSKASGYAFTN YLIEWVKQRPGQGLEWIGVINPGSGGTNYNEKFKGKATLTADKSSST AYMQLSSLTSDDSAVYFCARWRGDGYAYFDVWGAGTTVTVSSSG GPGPAGMKGLPGSDIVLTQSPASLAVSLGQRATISCKASQSVDDYDGD SYMNWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHPV EEEDAATYYCQQSNEDPYTFGGGTKEIKHHHHHHHEPEA
156	ACP120 NARA1 VI/Vh non- cleavable	mdmrvpaqllgllllwlgarcDIVLTQSPASLAVSLGQRATISCKASQSVDDYD DSYMNWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHP VEEEDAATYYCQQSNEDPYTFGGGTKEIKggggsgggsgggsggggQVQLQ QSGAELVRPGTSVKVSKASGYAFTNYLIEWVKQRPGQGLEWIGVI NPGSGGTNYNEKFKGKATLTADKSSSTAYMQLSSLTSDDSAVYFCA RWRGDGYAYFDVWGAGTTVTVSSHHHHHHHEPEA
157	ACP121 NARA1 VI/Vh cleavable	mdmrvpaqllgllllwlgarcDIVLTQSPASLAVSLGQRATISCKASQSVDDYD DSYMNWYQQKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHP VEEEDAATYYCQQSNEDPYTFGGGTKEIKSGGPGPAGMKGLPGSQ VQLQQSGAELVRPGTSVKVSKASGYAFTNYLIEWVKQRPGQGLE WIGVINPGSGGTNYNEKFKGKATLTADKSSSTAYMQLSSLTSDDSAV YFCARWRGDGYAYFDVWGAGTTVTVSSHHHHHHHEPEA
158	ACP124 IL-2 conjugate	mdmrvpaqllgllllwlgarcaptssstktqlqlehlldlqmilnginnyknpltrmltkfymppkka telkhlcleeelkpleevlnlaqsknfhlprdlisninvivlelkgsettfmceyadetativeflnrwitfc qsiistltggggsgggsgggsggggEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNANTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVTVSSHHHHHHHEPEA

SEQ ID NO.	Name	Sequence
159	ACP132 IL-2 conjugate	m d m r v p a q l g l l l w l r g a r c a p t s s s t k t q l q l e h l l l d l q m i n g i n n y k n p k l r m l t f k f y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e l n r w i t f c q s i u s t l t g g g g s g g g g s g g g g s d a h k s e v a h r f k d l g e e n f k a l v l i a f a q y l q q c p f e d h v k l v n e v t e f a k t c v a d e s a e n c d k s l h t l f g d k l c t v a t l r e t y g e m a d c c a k q e p e r n e c f l q h k d d n p n l p r l v r p e v d v m c t a f h d n e e t f l k k y l y e i a r r h p y f y a p e l l f f a k r y k a a f t e c c q a a d k a a c l l p k d e l r d e g k a s s a k q r l k c a s l q k f g e r a f k a w a v a r l s q r f p k a e a v s k l v t d l t k v h t e c c h g d l l e c a d d r a d l a k y i c e n q d s i s s k l k e c c e k p l l e k s h c i a e v e n d e m p a d l p s l a a d f v e s k d v c k n y a e a k d v f l g m f l y e y a r r h p d y s v l l l r l a k t y e t t l e k c c a a a d p h e c y a k v f d e f k p l v e e p q n l i k q n c e l f e q l g e y k f q n a l l v r y t k k v p q v s t p t l v e v s r n l g k v g s k c c k h p e a k r m p c a e d y l s v v l n q l e v l h e k t p v s d r v t k c c t e s l v n r r p c f s a l e v d e t y v p k e f n a e t f t h a d i c t l s e k e r q i k k q t a l v e l v k h k p k a t k e q l k a v m d d f a a f v e k c c k a d d k e t c f a e e g k k l v a a s q a a l g l H H H H H H E P E A
160	ACP141 IL-2 conjugate	m d m r v p a q l g l l l w l r g a r c a p t s s s t k t q l q l e h l l l d l q m i n g i n n y k n p k l r m l t f k f y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e l n r w i t f c q s i u s t l t g g g g s g g g g s g g g g s d a h k s e v a h r f k d l g e e n f k a l v l i a f a q y l q q c p f e d h v k l v n e v t e f a k t c v a d e s a e n c d k s l h t l f g d k l c t v a t l r e t y g e m a d c c a k q e p e r n e c f l q h k d d n p n l p r l v r p e v d v m c t a f h d n e e t f l k k y l y e i a r r h p y f y a p e l l f f a k r y k a a f t e c c q a a d k a a c l l p k d e l r d e g k a s s a k q r l k c a s l q k f g e r a f k a w a v a r l s q r f p k a e a v s k l v t d l t k v h t e c c h g d l l e c a d d r a d l a k y i c e n q d s i s s k l k e c c e k p l l e k s h c i a e v e n d e m p a d l p s l a a d f v e s k d v c k n y a e a k d v f l g m f l y e y a r r h p d y s v l l l r l a k t y e t t l e k c c a a a d p h e c y a k v f d e f k p l v e e p q n l i k q n c e l f e q l g e y k f q n a l l v r y t k k v p q v s t p t l v e v s r n l g k v g s k c c k h p e a k r m p c a e d y l s v v l n q l e v l h e k t p v s d r v t k c c t e s l v n r r p c f s a l e v d e t y v p k e f n a e t f t h a d i c t l s e k e r q i k k q t a l v e l v k h k p k a t k e q l k a v m d d f a a f v e k c c k a d d k e t c f a e e g k k l v a a s q a a l g l H H H H H H E P E A
161	ACP142 IL-2 conjugate	m d m r v p a q l g l l l w l r g a r c a p t s s s t k t q l q l e h l l l d l q m i n g i n n y k n p k l r m l t f k f y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e l n r w i t f c q s i u s t l t S G G P G P A G M K G L P G S d a h k s e v a h r f k d l g e e n f k a l v l i a f a q y l q q c p f e d h v k l v n e v t e f a k t c v a d e s a e n c d k s l h t l f g d k l c t v a t l r e t y g e m a d c c a k q e p e r n e c f l q h k d d n p n l p r l v r p e v d v m c t a f h d n e e t f l k k y l y e i a r r h p y f y a p e l l f f a k r y k a a f t e c c q a a d k a a c l l p k d e l r d e g k a s s a k q r l k c a s l q k f g e r a f k a w a v a r l s q r f p k a e a v s k l v t d l t k v h t e c c h g d l l e c a d d r a d l a k y i c e n q d s i s s k l k e c c e k p l l e k s h c i a e v e n d e m p a d l p s l a a d f v e s k d v c k n y a e a k d v f l g m f l y e y a r r h p d y s v l l l r l a k t y e t t l e k c c a a a d p h e c y a k v f d e f k p l v e e p q n l i k q n c e l f e q l g e y k f q n a l l v r y t k k v p q v s t p t l v e v s r n l g k v g s k c c k h p e a k r m p c a e d y l s v v l n q l e v l h e k t p v s d r v t k c c t e s l v n r r p c f s a l e v d e t y v p k e f n a e t f t h a d i c t l s e k e r q i k k q t a l v e l v k h k p k a t k e q l k a v m d d f a a f v e k c c k a d d k e t c f a e e g k k l v a a s q a a l g l H H H H H H E P E A
162	ACP144 IL-2 conjugate	m d m r v p a q l g l l l w l r g a r c a p t s s s t k t q l q l e h l l l d l q m i n g i n n y k n p k l r m l t f k f y m p k k a t e l k h l q c l e e l k p l e e v l n l a q s k n f h l r p d l i s n i n v i v l e l k g s e t t f m c e y a d e t a t i v e l n r w i t f c q s i u s t l t S G G P G P A G M K G L P G S E V Q L V E S G G G L V Q P G N S L R L S C A A S G F T F S K F G M S W V R Q A P G K G L E W V S S I S G S G R D T L Y A E S V K G R F T I S R D N A K T T L Y L Q M N S L R P E D T A V Y Y C T I G G S L S V S S Q G T L V T V S S g g g g s g g g g s g g g s g g g g s g g g g s g g g s S G G P G P A G M K G L P G S E V Q L V E S G G G L V Q P G G S L R L S C A A S G F T F S S Y T L A W V R Q A P G K G L E W V A A I D S S S Y T Y S P D T V R G R F T I S R D N A K N S L Y L Q M N S L R A E D T A V Y Y C A R D S N W D A L D Y W G Q G T T V T V S S G G G S G G G G S G G G S D I Q M T Q S P S S L S A S V G D R V T T T C K A S Q N V G T N V G W Y Q Q K P G K A P K A L I Y S A S F R Y S G V P S R F S G S G S G T D F I L T I S S L Q P E D F A T Y Y C Q Q Y Y T Y P Y T F G G G T K V E I K g g g g s g g g s g g g g s Q V Q L Q E S G G G L A Q A G G S L S L S C A A S G F T V S N S V M A W Y R Q T P G K Q R E F V A I N S V G S T N Y A D S V K G R F T I S R D N A K N T V Y L Q M N N L K P E D T A V Y V C N R N F D R I Y W G O G T Q V T V S S H H H H H H E P E A
163	ACP145 IL-2	m d m r v p a q l g l l l w l r g a r c Q V Q L E S G G G L A Q A G G S L R L S C A A S G F T V S N S V M A W Y R Q T P G K Q R E F V A I N S V G S T N Y A D S V K G R F T I S R D N A K N T V Y L O M N N L K P E D T A V Y V C N R N F D R I Y W G O G T Q V T V S S g g g g s g g g s g g g

[illegible]

SEQ ID NO.	Name	Sequence
		ggsSGGPGPAGMKGLPGSEVQLVESGGGLVQPGGSLRLSCAASGFTFS SYTLAWVRQAPGKGLEWVAAIDSSSYTSPDTRGRFTISRDNANKN SLYLQMNSLRAEDTAVYYCARDSDNWDALDYWGQGTFTVTVSSGGG GSGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVGTNVG WYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISSLQPED FATYYCQQYYTYPTYTFGGGTKEIKHHHHHHEPEA
168	ACPI49 IL-2 conjugate	mdmrvpaqllgllllwlrgrcQVQLQESGGGLVQAGGSLRLSCAASGRIFSIDI MSWYRQAPGKQRELVARITRGGTISYDDSVKGRFTISRDNANKNTVY LQMNSLKPEDTGYYCNAALYGTQDYWGKGTQVTVSSSGGPGPAGM KGLPGSaptssstktqlqllehlldlqmilnginnynkpkltmrltkfymppkkatelkhlqeleelk pleevlnlaqsknfhrlprdlisninvivlelkgsettfmceyadetativeflorwitfcqsiistltSGGPG PAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSW VRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNANKTTLYLQM NSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgggsgggsg ggsgggsgsgSGGPGPAGMKGLPGSEVQLVESGGGLVQPGGSLRLSCAAS GFTFSYTLAWVRQAPGKGLEWVAAIDSSSYTSPDTRGRFTISRDN NAKNSLYLQMNSLRAEDTAVYYCARDSDNWDALDYWGQGTFTVTVS SGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRTITCKASQNVGT NVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFTLTISSLQ PEDFATYYCQQYYTYPTYTFGGGTKEIKHHHHHHEPEA
169	ACP33 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrgrcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNANKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGScdlpqthnlrnkraltllvqmrrlspisclskdrkdfgfpqekvdaqqikkaqipvlseftqqiln iftskdssaawnttllsfcndlhqqIndlqgclmqvgvgqefplqtedallavrkyfhritylrekkhsp cawevvraevwralsssanvSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNS LRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESV KGRFTISRDNANKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLL TVSSHHHHHHEPEA
170	ACP131 Mouse IFNa	mdmrvpaqllgllllwlrgrcddlpqthnlrnkraltllvqmrrlspisclskdrkdfgfpqekvdaqqikk aqaipvlseftqqilniftskdssaawnttllsfcndlhqqIndlqgclmqvgvgqefplqtedallavrky fhritylrekkhspcawevvraevwralsssanvlgrrlrekkHHHHHHEPEA
171	ACP125 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrgrcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNANKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGScdlpqthnlrnkraltllvqmrrlspisclskdrkdfgfpqekvdaqqikkaqipvlseftqqiln iftskdssaawnttllsfcndlhqqIndlqgclmqvgvgqefplqtedallavrkyfhritylrekkhsp cawevvraevwralsssanvlgrrlrekkHHHHHHEPEA
172	ACP126 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrgrcddlpqthnlrnkraltllvqmrrlspisclskdrkdfgfpqekvdaqqikk aqaipvlseftqqilniftskdssaawnttllsfcndlhqqIndlqgclmqvgvgqefplqtedallavrky fhritylrekkhspcawevvraevwralsssanvlgrrlrekkSGGPGPAGMKGLPGSEVQ LVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVS SISGSGRDTLYAESVKGRFTISRDNANKTTLYLQMNSLRPEDTAVYYC TIGGSLSVSSQGTLLTVSSHHHHHHEPEA
173	ACP127 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrgrcEAHKSEIAHRYNDLGEQHFGLVLIAFSQQYLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFLOHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRRHPYFYAPELLYYAEQYNEILTQCCAEA DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFAKAWAVA RLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAKY

SEQ ID NO.	Name	Sequence
		MCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEYG FQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCTLPEDQRLPCV EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHSDICTLPEKEKQIKKQATALAELVKHKPKATAEQL KTMDDFAQFLDTCCAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGScdlpqthnlrnlraltllvqmrrlspisclkrkdfgfpqekvdaqqikkaqipvls eltqqilniftskdssaawntllsdfcndlhqqIndlqgclmqvgvqefplqtedallavrkyfhrityl ekkhspcawevvraevwralsssanvlgrrlreekSGGPGPAGMKGLPGSEAHKSEIAH RYNDLGEQHFKGLVLIAPSYLQKCSYDEHAKLVQEVTDFAKTCVA DESAANCDKSLHTLFGDKLCAIPNLRENYGELADCCTKQEPERNECF LQHKDDNPSLPPFERPEAEAMCTSFKENPTTFMGHYLHEVARRHPYF YAPELLYYAEQYNEILTQCCAEADKESCLTPKLDGVKEKALVSSVR QRMKCSSMQKFGERAFKAWAVARLSQTFPNADFAEITKLATDLTKV NKECCHGDLLECADDRAELAKYMCENQATISSKLQTCCDKPLLKKA HCLSEVEHDTMPADLPAIAADFVEDQEVCKNYAEAKDVFLGTFLYE YSRRHPDYSVSLLLRLAKKYEATLEKCCAEANPPACYGTVLAEFQPL VEEPKNLVKTNCDLYEKLGEYGFQNAILVRYTQKAPQVSTPTLVEA ARNLGRVGTCKCTLPEDQRLPCVEDYLSAILNRVCLLHEKTPVSEHV TKCCSGSLVERRPCFSALTVDETYVPKEFKAETFTFHSDICTLPEKEK QIKKQATALAELVKHKPKATAEQLKTMDDFAQFLDTCCAADKDT CFSTEGPNLVTRCKDALAHHHHHEPEA
174	ACPI28 Mouse IFNa- conjugate	mdmrvpaqllglllwlrgarcEAAHKSEIAHRYNDLGEQHFKGLVLIAPSYLQ KCSYDEHAKLVQEVTDFAKTCVADESAANCDKSLHTLFGDKLCAIP NLRENYGELADCCTKQEPERNECFLQHKDDNPSLPPFERPEAEAMCT SFKENPTTFMGHYLHEVARRHPYFYAPELLYYAEQYNEILTQCCAEA DKESCLTPKLDGVKEKALVSSVRQRMKCSSMQKFGERAFKAWAVA RLSQTFPNADFAEITKLATDLTKVNKECCHGDLLECADDRAELAKY MCENQATISSKLQTCCDKPLLKKAHCLSEVEHDTMPADLPAIAADFV EDQEVCKNYAEAKDVFLGTFLYEYSRRHPDYSVSLLLRLAKKYEAT LEKCCAEANPPACYGTVLAEFQPLVEEPKNLVKTNCDLYEKLGEYG FQNAILVRYTQKAPQVSTPTLVEAARNLGRVGTCKCTLPEDQRLPCV EDYLSAILNRVCLLHEKTPVSEHVTKCCSGSLVERRPCFSALTVDETY VPKEFKAETFTFHSDICTLPEKEKQIKKQATALAELVKHKPKATAEQL KTMDDFAQFLDTCCAADKDTCFSTEGPNLVTRCKDALASGGPGP AGMKGLPGScdlpqthnlrnlraltllvqmrrlspisclkrkdfgfpqekvdaqqikkaqipvls eltqqilniftskdssaawntllsdfcndlhqqIndlqgclmqvgvqefplqtedallavrkyfhrityl ekkhspcawevvraevwralsssanvlgrrlreekHHHHHHEPEA
175	ACPI29 Mouse IFNa- conjugate	mdmrvpaqllglllwlrgarcddlpqthnlrnlraltllvqmrrlspisclkrkdfgfpqekvdaqqikk aqaipvlseltqqilniftskdssaawntllsdfcndlhqqIndlqgclmqvgvqefplqtedallavrky fhritylrekkhspcawevvraevwralsssanvlgrrlreekSGGPGPAGMKGLPGSEAH KSEIAHRYNDLGEQHFKGLVLIAPSYLQKCSYDEHAKLVQEVTDF AKTCVADESAANCDKSLHTLFGDKLCAIPNLRENYGELADCCTKQE PERNECFQHKDDNPSLPPFERPEAEAMCTSFKENPTTFMGHYLHEV ARRHPYFYAPELLYYAEQYNEILTQCCAEADKESCLTPKLDGVKEK ALVSSVRQRMKCSSMQKFGERAFKAWAVARLSQTFPNADFAEITKL ATDLTKVNKECCHGDLLECADDRAELAKYMCENQATISSKLQTCCD KPLLKKAHCLSEVEHDTMPADLPAIAADFVEDQEVCKNYAEAKDVFL GTFLYEYSRRHPDYSVSLLLRLAKKYEATLEKCCAEANPPACYGTV

SEQ ID NO.	Name	Sequence
		LAEFQPLVEEPKNLVKTNCDLYEKLGEYGFQNAILVRYTQKAPQVST PTLVEAARNLGRVGTKCCTLPEDQRLPCVEDYLSAILNRVCLLHEKT PVSEHVTKCCSGSLVERRPCFSALTVDETYVPKEFKAETFTFHSDICT LPEKEKQIKKQTALAEVLVHKPKATAEQLKTVMDDFQAQFLDTCCKA ADKDTCFSTEGPNLVTRCKDALAHHHHHHEPEA
176	ACPI50 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrarcQVQLQESGGGLAQAGGSLSLSCAASGFTVSNS VMAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDNKNTV YLQMNNLKPEDTAVVVCNRRNFDRIYWGQGTQVTVSSggggsgggsggg gsEVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGL EWVSSISGSGRDTLYAESVKGRFTISRDNKTTLYLQMNSLRPEDTA VYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMKGLPGScdlpqthnlrnkral dlvqmrrlspisclskdrkdfgfpqekvdaqqikkaqaipvlsetlqqilniftskdssawntllsdfcndl hqqIndlqgclmqvgvqefpltedallavrkyfhritylrekkhspcawevvraevwralsssanvl grlreekSGGPGPAGMKGLPGSEVQLVESGGGLVQPGNSLRLSCAASGF TFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSHHHHH HEPEA
177	ACPI51 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrarcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSSGGPGPAGMK GLPGScdlpqthnlrnkraldlvqmrrlspisclskdrkdfgfpqekvdaqqikkaqaipvlsetlqqiln iftskdssawntllsdfcndlhqqIndlqgclmqvgvqefpltedallavrkyfhritylrekkhsp cawevvraevwralsssanvlgrlreekSGGPGPAGMKGLPGSEVQLVESGGGLV QPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTL YAESVKGRFTISRDNKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSS QGTLLTVSSggggsgggsggggsQVQLQESGGGLAQAGGSLSLSCAASGF TVSNSVMAWYRQTPGKQREFVAIINSVGSTNYADSVKGRFTISRDN KNTVYLQMNNLKPEDTAVVVCNRRNFDRIYWGQGTQVTVSSHHHHH HEPEA
178	ACPI52 Mouse IFNa- conjugate	mdmrvpaqllgllllwlrarcEVQLVESGGGLVQPGNSLRLSCAASGFTFSKF GMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNKTT LYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgggg scdlpqthnlrnkraldlvqmrrlspisclskdrkdfgfpqekvdaqqikkaqaipvlsetlqqilniftskds saawntllsdfcndlhqqIndlqgclmqvgvqefpltedallavrkyfhritylrekkhspcawevv raevwralsssanvlgrlreekggggsgggsggggsEVQLVESGGGLVQPGNSLRLSC AASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRF TISRDNKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSS HHHHHHEPEA
179	ACPI53 (IL-2 Conju- gate)	mdmrvpaqllgllllwlrarcaptssstskktqlqlehlldlqmilnginnyknpkltmrmltkfympkka telkhqlcecelkpleevlnlaqsknfhlrprdlisninivlelkgsettfmceyadetativeflnrwitfc qsiiistltsggpGPAGLYAQpgsEVQLVESGGGLVQPGNSLRLSCAASGFTFS KFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNK TTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsgg ggsgggsgggsgggsgggsggpGPAGLYAQpgsEVQLVESGGGLVQPGSLR LSCAASGFTFSSYTLAWVRQAPGKGLEWVAIDSSSYTSPDTPVRGR FTISRDNKNSLYLQMNSLRAEDTAVYYCARDSDNDALDYWGQGT

SEQ ID NO.	Name	Sequence
		TVTVSSGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVITITCKAS QNVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFLT LTISSLQPEDFATYYCQQYYTYPYTFGGGGTKVEIKHHHHHHEPEA
180	ACPI54 (IL-2 Conju- gate)	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknpkltlrmlltkfymppkka telkhlqcleelkpleevlnlaqsknfhlprdlisininvivlelkgsettfmceyadetativeflnrwitfc qsüistltsggpPGGPAGIGpgsEVQLVESGGGLVQPGNSLRRLSCAASGFTFS KFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAK TTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsg ggsgggsgggsgggsgggsggpPGGPAGIGpgsEVQLVESGGGLVQPGGSLRL SCAASGFTFSSYTLAWVRQAPGKGLEWVAIDSSSYTYSPDTVRGRF TISRDNAAKNSLYLQMNSLRAEDTAVYYCARDNSWDALDYWGQGT TVTVSSGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVITITCKASQ NVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFLT ISSLQPEDFATYYCQQYYTYPYTFGGGGTKVEIKHHHHHHEPEA
181	ACPI55 (IL-2 Conju- gate)	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknpkltlrmlltkfymppkka telkhlqcleelkpleevlnlaqsknfhlprdlisininvivlelkgsettfmceyadetativeflnrwitfc qsüistltsggpALFKSSFPpgsEVQLVESGGGLVQPGNSLRRLSCAASGFTFS KFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAK TTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsg ggsgggsgggsgggsgggsggpALFKSSFPpgsEVQLVESGGGLVQPGGSLRL SCAASGFTFSSYTLAWVRQAPGKGLEWVAIDSSSYTYSPDTVRGRF TISRDNAAKNSLYLQMNSLRAEDTAVYYCARDNSWDALDYWGQGT TVTVSSGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVITITCKASQ NVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFLT ISSLQPEDFATYYCQQYYTYPYTFGGGGTKVEIKHHHHHHEPEA
182	ACPI56 (IL-2 Conju- gate)	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknpkltlrmlltkfymppkka telkhlqcleelkpleevlnlaqsknfhlprdlisininvivlelkgsettfmceyadetativeflnrwitfc qsüistltsggpPLAQKLKSSpgsEVQLVESGGGLVQPGNSLRRLSCAASGFTF SKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISRDNAAK KTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLLTVSSggggsgggsg ggsgggsgggsgggsgggsggpPLAQKLKSSpgsEVQLVESGGGLVQPGGSL RLSCAASGFTFSSYTLAWVRQAPGKGLEWVAIDSSSYTYSPDTVRGRF TISRDNAAKNSLYLQMNSLRAEDTAVYYCARDNSWDALDYWGQGT TTTVTVSSGGGSGGGGSGGGGSDIQMTQSPSSLSASVGDRVITITCKA SQNVGTNVGWYQQKPGKAPKALIYSASFRYSGVPSRFSGSGSGTDFLT LTISSLQPEDFATYYCQQYYTYPYTFGGGGTKVEIKHHHHHHEPEA
183	ACPI57 (IL-2 Conju- gate)	mdmrvpaqlglglllwlrgarcaptssstkkqlqlehlldlqmilnginnyknpkltlrmlltkfymppkka telkhlqcleelkpleevlnlaqsknfhlprdlisininvivlelkgsettfmceyadetativeflnrwitfc qsüistltsggpPGGPAGIGalfkssfpPLAQKLKSSpgsEVQLVESGGGLVQPGN SLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAES VKGRFTISRDNAAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLL TVTVSSggggsgggsgggsgggsgggsgggsgggsggpPGGPAGIGalfkssfpPLAQ KLKSSpgsEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTLAWVRQ APGKGLEWVAIDSSSYTYSPDTVRGRFTISRDNAAKNSLYLQMNSLR AEDTAVYYCARDNSWDALDYWGQGTTVTVSSGGGSGGGGSGGG GSDIQMTQSPSSLSASVGDRVITITCKASQNVGTNVGWYQQKPGKAP KALIYSASFRYSGVPSRFSGSGSGTDFLTISSLQPEDFATYYCQQYY TYPYTFGGGGTKVEIKHHHHHHEPEA

SEQ ID NO.	Name	Sequence
184		Place hold
185		Place hold
186		Place hold
187		Place hold
188		Place hold
189		Place hold
190		Place hold
191		Place hold
192	Blocker 2 (IL2 blocker)	mdmrvpaqlglglllwlrgarcEVQLVESGGGLVQPGGSLRLSCAASGFTFSSYT LAWVRQAPGKGLEWVA AIDSSSYTYSPDTRVGRFTISRDN AKNSLY LQMNSLRAEDTAVYYCARDSDNDALDYWGQGTITVTVSSgggsgggg sggggsDIQMTQSPSSLSASVGDRTITCKASQNVGTNVGWYQQKPGK APKALIYSASFRYSQVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQY YTPYTFGGGTKEIKHHHHHH
193	Blocker 12 (IL-12 blocker)	mdmrvpaqlglglllwlrgarcQSVLTQPPSVSGAPGQRVTISCSGSRSNIGSNTV KWYQQLPGTAPKLLIYYNDQRPSGVPDRFSGSKSGTSASLAITGLQA EDEADY YCQSYDRYTHPALLFGTGTKVTVLgggsgggsgggsggggQVQLV ESGGGVVQPGRLSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAFI RYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC KTHGSHDNWGQGTMTVTVSSHHHHHH
194		EVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQGGGGGLD GNEEPGGLEWVSSISGSGRDTLYADSVKGRFTISRDN AKTTLYLQMN SLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS
225	ACP203	EVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS sggpGPAGLYAQpgscdlpqthnlrnrkaltllvqmrrlspisclckdrkdfgfpqekvdaqqikkaqai pviseltqqilniftskdssaawnttldsfcdnlhqqIndlqgclmqqvgvqefplqtqedaallavrkfhrityl vylrekhhspcawevvraevwralssanvlgrlreeksggpGPAGLYAQpgsEVQLVESGGGLV QPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFT ISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS
226	ACP204	EVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS sggpALFKSSFPpgscdlpqthnlrnrkaltllvqmrrlspisclckdrkdfgfpqekvdaqqikkaqai pviseltqqilniftskdssaawnttldsfcdnlhqqIndlqgclmqqvgvqefplqtqedaallavrkfhrityl vylrekhhspcawevvraevwralssanvlgrlreeksggpALFKSSFPpgsEVQLVESGGGLVQ PGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTIS RDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS
227	ACP205	EVQLVESGGGLVQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS sggpPLAQKLKSSpgscdlpqthnlrnrkaltllvqmrrlspisclckdrkdfgfpqekvdaqqikkaqai pviseltqqilniftskdssaawnttldsfcdnlhqqIndlqgclmqqvgvqefplqtqedaallavrkfhrityl vylrekhhspcawevvraevwralssanvlgrlreeksggpPLAQKLKSSpgsEVQLVESGGGL VQPGNSLRLSCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGR FTISRDN AKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTITVTVSS

SEQ ID NO.	Name	Sequence
228	ACP206	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS sggpGPAGLYAQpgscdlpqthslgsrrtlllaqmrrislsclkdrrhdfgfpqeefgnqfqketipvl hemiiqqifnlfstkdsaaawdetlldkfyteyqqIndleacviqgvgtetplmkedsilavrkfgritlyl kekkyspcawewvraeimrfslstnlqeslrskesggpGPAGLYAQpgsEVQLVESGGGLVQPGN SLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISR DNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS
229	ACP207	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS sggpALFKSSFPpgscdlpqthslgsrrtlllaqmrrislsclkdrrhdfgfpqeefgnqfqketipvlh emiiqqifnlfstkdsaaawdetlldkfyteyqqIndleacviqgvgtetplmkedsilavrkfgritlylk ekkkyspcawewvraeimrfslstnlqeslrskesggpALFKSSFPpgsEVQLVESGGGLVQPGN SLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISR DNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS
230	ACP208	EVQLVESGGGLVQPGNSLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGR DTLYAESVKGRFTISRDNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS sggpPLAQKLKSSpgscdlpqthslgsrrtlllaqmrrislsclkdrrhdfgfpqeefgnqfqketipv lhemiiqqifnlfstkdsaaawdetlldkfyteyqqIndleacviqgvgtetplmkedsilavrkfgritly lkekkyspcawewvraeimrfslstnlqeslrskesggpPLAQKLKSSpgsEVQLVESGGGLVQPGN SLRSLCAASGFTFSKFGMSWVRQAPGKGLEWVSSISGSGRDTLYAESVKGRFTISR DNAKTTLYLQMNSLRPEDTAVYYCTIGGSLSVSSQGTLTVSS
258	MMP14 substrate motif sequence	GPLGLKAQ
259	MMP14 substrate motif sequence	LPLGLKAQ
260	MMP14 substrate motif sequence	SPLGLKAQ
261	MMP14 substrate motif sequence	QPLGLKAQ
262	MMP14 substrate	KPLGLKAQ

SEQ ID NO.	Name	Sequence
	motif sequence	
263	MMP14 substrate motif sequence	FPLGLKAQ
264	MMP14 substrate motif sequence	HPLGLKAQ
265	MMP14 substrate motif sequence	PPLGLKAQ
266	MMP14 substrate motif sequence	APLGLKAQ
267	MMP14 substrate motif sequence	DPLGLKAQ
268	MMP14 substrate motif sequence	GPHGLKAQ
269	MMP14 substrate motif sequence	GPSGLKAQ
270	MMP14 substrate motif	GPQGLKAQ

SEQ ID NO.	Name	Sequence
	sequence	
271	MMP14 substrate motif sequence	GPPGLKAQ
272	MMP14 substrate motif sequence	GPEGLKAQ
273	MMP14 substrate motif sequence	GPFGLKAQ
274	MMP14 substrate motif sequence	GPRGLKAQ
275	MMP14 substrate motif sequence	GPGGLKAQ
276	MMP14 substrate motif sequence	GPAGLKAQ
277	MMP14 substrate motif sequence	LPAGLKGA
195	MMP14 substrate motif sequence	GPAGLYAQ

SEQ ID NO.	Name	Sequence
278	MMP14 substrate motif sequence	GPANLVAQ
279	MMP14 substrate motif sequence	GPAALVGA
280	MMP14 substrate motif sequence	GPANLRAQ
281	MMP14 substrate motif sequence	GPAGLRAQ
282	MMP14 substrate motif sequence	GPAGLVAQ
283	MMP14 substrate motif sequence	GPAGLRGA
284	MMP14 substrate motif sequence	LPAGLVGA
285	MMP14 substrate motif sequence	GPAGLKGA
286	MMP14	GPLALKAQ

SEQ ID NO.	Name	Sequence
	substrate motif sequence	
287	MMP14 substrate motif sequence	GPLNLKAQ
288	MMP14 substrate motif sequence	GPLHLKAQ
289	MMP14 substrate motif sequence	GPLYLKAQ
290	MMP14 substrate motif sequence	GPLPLKAQ
291	MMP14 substrate motif sequence	GPLELKAQ
292	MMP14 substrate motif sequence	GPLRLKAQ
293	MMP14 substrate motif sequence	GPLLLKAQ
294	MMP14 substrate	GPLSLKAQ

SEQ ID NO.	Name	Sequence
	motif sequence	
295	MMP14 substrate motif sequence	GPLGLYAQ
296	MMP14 substrate motif sequence	GPLGLFAQ
297	MMP14 substrate motif sequence	GPLGLLAQ
298	MMP14 substrate motif sequence	GPLGLHAQ
299	MMP14 substrate motif sequence	GPLGLRAQ
300	MMP14 substrate motif sequence	GPLGLAAQ
301	MMP14 substrate motif sequence	GPLGLEAQ
302	MMP14 substrate motif	GPLGLGAQ

SEQ ID NO.	Name	Sequence
	sequence	
303	MMP14 substrate motif sequence	GPLGLPAQ
304	MMP14 substrate motif sequence	GPLGLQAQ
305	MMP14 substrate motif sequence	GPLGLSAQ
306	MMP14 substrate motif sequence	GPLGLVAQ
307	MMP14 substrate motif sequence	GPLGLKLQ
308	MMP14 substrate motif sequence	GPLGLKFQ
309	MMP14 substrate motif sequence	GPLGLKEQ
310	MMP14 substrate motif sequence	GPLGLKKQ

SEQ ID NO.	Name	Sequence
311	MMP14 substrate motif sequence	GPLGLKQQ
312	MMP14 substrate motif sequence	GPLGLKSQ
313	MMP14 substrate motif sequence	GPLGLKGQ
314	MMP14 substrate motif sequence	GPLGLKHQ
315	MMP14 substrate motif sequence	GPLGLKPQ
316	MMP14 substrate motif sequence	GPLGLKAG
317	MMP14 substrate motif sequence	GPLGLKAF
318	MMP14 substrate motif sequence	GPLGLKAP
319	MMP14	GPLGLKAL

SEQ ID NO.	Name	Sequence
	substrate motif sequence	
320	MMP14 substrate motif sequence	GPLGLKAE
321	MMP14 substrate motif sequence	GPLGLKAA
322	MMP14 substrate motif sequence	GPLGLKAH
323	MMP14 substrate motif sequence	GPLGLKAK
324	MMP14 substrate motif sequence	GPLGLKAS
325	MMP14 substrate motif sequence	GPLGLFGA
326	MMP14 substrate motif sequence	GPLGLQGA
327	MMP14 substrate	GPLGLVGA

SEQ ID NO.	Name	Sequence
	motif sequence	
328	MMP14 substrate motif sequence	GPLGLAGA
329	MMP14 substrate motif sequence	GPLGLLGA
330	MMP14 substrate motif sequence	GPLGLRGA
331	MMP14 substrate motif sequence	GPLGLYGA
332	CTSL1 substrate motif sequence	ALFKSSPP
333	CTSL1 substrate motif sequence	SPFRSSRQ
334	CTSL1 substrate motif sequence	KLFKSSPP
335	CTSL1 substrate motif	HLFKSSPP

SEQ ID NO.	Name	Sequence
	sequence	
336	CTSL1 substrate motif sequence	SLFKSSPP
337	CTSL1 substrate motif sequence	QLFKSSPP
338	CTSL1 substrate motif sequence	LLFKSSPP
339	CTSL1 substrate motif sequence	PLFKSSPP
340	CTSL1 substrate motif sequence	FLFKSSPP
341	CTSL1 substrate motif sequence	GLFKSSPP
342	CTSL1 substrate motif sequence	VLFKSSPP
343	CTSL1 substrate motif sequence	ELFKSSPP

SEQ ID NO.	Name	Sequence
344	CTSL1 substrate motif sequence	AKFKSSPP
345	CTSL1 substrate motif sequence	AHFKSSPP
346	CTSL1 substrate motif sequence	AGFKSSPP
347	CTSL1 substrate motif sequence	APFKSSPP
348	CTSL1 substrate motif sequence	ANFKSSPP
349	CTSL1 substrate motif sequence	AFFKSSPP
350	CTSL1 substrate motif sequence	AAFKSSPP
351	CTSL1 substrate motif sequence	ASFKSSPP
352	CTSL1	AEFKSSPP

SEQ ID NO.	Name	Sequence
	substrate motif sequence	
353	CTSL1 substrate motif sequence	ALRKSSPP
354	CTSL1 substrate motif sequence	ALLKSSPP
355	CTSL1 substrate motif sequence	ALAKSSPP
356	CTSL1 substrate motif sequence	ALQKSSPP
357	CTSL1 substrate motif sequence	ALHKSSPP
358	CTSL1 substrate motif sequence	ALPKSSPP
359	CTSL1 substrate motif sequence	ALTKSSPP
360	CTSL1 substrate	ALGKSSPP

SEQ ID NO.	Name	Sequence
	motif sequence	
361	CTSL1 substrate motif sequence	ALDKSSPP
199	CTSL1 substrate motif sequence	ALFFSSPP
362	CTSL1 substrate motif sequence	ALFHSSPP
363	CTSL1 substrate motif sequence	ALFTSSPP
364	CTSL1 substrate motif sequence	ALFASSPP
365	CTSL1 substrate motif sequence	ALFQSSPP
366	CTSL1 substrate motif sequence	ALFLSSPP
367	CTSL1 substrate motif	ALFGSSPP

SEQ ID NO.	Name	Sequence
	sequence	
368	CTSL1 substrate motif sequence	ALFESSPP
369	CTSL1 substrate motif sequence	ALFPSSPP
370	CTSL1 substrate motif sequence	ALFKHSPP
371	CTSL1 substrate motif sequence	ALFKLSPP
372	CTSL1 substrate motif sequence	ALFKKSPP
373	CTSL1 substrate motif sequence	ALFKASPP
374	CTSL1 substrate motif sequence	ALFKISPP
375	CTSL1 substrate motif sequence	ALFKGSPP

SEQ ID NO.	Name	Sequence
376	CTSL1 substrate motif sequence	ALFKNSPP
377	CTSL1 substrate motif sequence	ALFKRSPP
378	CTSL1 substrate motif sequence	ALFKESPP
379	CTSL1 substrate motif sequence	ALFKFSPP
380	CTSL1 substrate motif sequence	ALFKPSPP
381	CTSL1 substrate motif sequence	ALFKSFPP
382	CTSL1 substrate motif sequence	ALFKSLPP
383	CTSL1 substrate motif sequence	ALFKSIPP
384	CTSL1	ALFKSKPP

SEQ ID NO.	Name	Sequence
	substrate motif sequence	
385	CTSL1 substrate motif sequence	ALFKSAPP
386	CTSL1 substrate motif sequence	ALFKSQPP
387	CTSL1 substrate motif sequence	ALFKSPPP
388	CTSL1 substrate motif sequence	ALFKSEPP
389	CTSL1 substrate motif sequence	ALFKSGPP
198	CTSL1 substrate motif sequence	ALFKSSFP
390	CTSL1 substrate motif sequence	ALFKSSLP
391	CTSL1 substrate	ALFKSSGP

SEQ ID NO.	Name	Sequence
	motif sequence	
392	CTSL1 substrate motif sequence	ALFKSSSP
393	CTSL1 substrate motif sequence	ALFKSSVP
394	CTSL1 substrate motif sequence	ALFKSSH P
395	CTSL1 substrate motif sequence	ALFKSSAP
396	CTSL1 substrate motif sequence	ALFKSSNP
397	CTSL1 substrate motif sequence	ALFKSSKP
398	CTSL1 substrate motif sequence	ALFKSSEP
399	CTSL1 substrate motif	ALFKSSPF

SEQ ID NO.	Name	Sequence
	sequence	
400	CTSL1 substrate motif sequence	ALFKSSPH
401	CTSL1 substrate motif sequence	ALFKSSPG
402	CTSL1 substrate motif sequence	ALFKSSPA
403	CTSL1 substrate motif sequence	ALFKSSPS
404	CTSL1 substrate motif sequence	ALFKSSPV
405	CTSL1 substrate motif sequence	ALFKSSPQ
406	CTSL1 substrate motif sequence	ALFKSSPK
407	CTSL1 substrate motif sequence	ALFKSSPL

200

SEQ ID NO.	Name	Sequence
408	CTSL1 substrate motif sequence	ALFKSSPD

CLAIMS

1. A polypeptide comprising a protease cleavable amino acid sequence selected from the group consisting of SEQ ID NOS:195-220, or an amino acid sequence that has at least about 90% identity to SEQ ID NOS:195-220.

2. A polypeptide comprising SEQ ID NO:195 or a functional variant of SEQ ID NO:195 or SEQ ID 198 or a functional variant of SEQ ID NO:198.


3. The polypeptide of claim 2 wherein the functional variant of SEQ ID NO:195 comprises any of SEQ ID NOS: 258-331.

4. The polypeptide of claim 2 wherein the functional variant of SEQ ID NO:198 comprises SEQ ID NOS: 199 or any of SEQ ID NOS: 332-408.

5. The polypeptide of claim 2 wherein the polypeptide comprises SEQ ID NO:195.

6. The polypeptide of claim 2, wherein the polypeptide comprises SEQ ID NO:198.

7. A polypeptide comprising Formula I:



wherein D1 is a first domain of interest;

L1 is a separation moiety that connects or links D1 to D2, wherein the separation moiety comprises an amino acid sequence selected from SEQ ID NOS: 195-220 or an amino acid sequence that has at least about 90% identity to SEQ ID NOS:195-220;

D2 is a second domain of interest.

8. The polypeptide as in claim 1 or 7, wherein the separation moiety comprises at least one proline residue adjacent to the amino acid sequence selected from the group consisting of SEQ ID NOS: 195-220 or an amino acid sequence at least at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS: 195-220.

9. The polypeptide as in any one of claims 1-8, wherein the separation moiety comprises a cleavable moiety that is a substrate for two or more proteases.

10. The polypeptide as in any one of claims 1-9, wherein the protease is selected from the group consisting of FAPa, CTSL1, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 and MMP14.
11. The polypeptide as in any one of claims 1-9, wherein the proteases is selected from MMP1, MMP2, MMP9, MMP4, cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin K or cathepsin L.
12. The polypeptide of as in any one of claims 1-7, wherein the separation moiety comprises an amino acid sequence that is a substrate for at least one protease present in the tumor microenvironment of a human tumor.
13. The polypeptide as in any one of claims 1-8, wherein the separation moiety comprises two or more cleavable moieties, each of which is a substrate for a protease.
14. The polypeptide as in any one of claims 1-8, comprising a first cleavable moiety comprising a first amino acid sequence that is a substrate for a first protease and a second cleavable moiety comprising a second amino acid sequence that is a substrate for a second protease.
15. The polypeptide of any one of claims 1-8, further comprising a non-cleavable linker sequence.
16. The polypeptide of any of claims 1-8, wherein the polypeptide comprises a cytokine, chemokine, growth factor, a soluble receptor, or any combination thereof.
17. The polypeptide of any of claims 1-8, wherein the polypeptide comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain.
18. The polypeptide of any of claims 1-8, wherein the polypeptide comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit.

19. The polypeptide of any of claims claim 1-8, wherein the polypeptide comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.
20. The polypeptide of any one of claims 1-14, wherein the cleavable moiety is cleaved with either (a) greater catalytic efficiency or (b) greater specificity or (c) both (a) and (b), by one or more proteases than a reference polypeptide sequence.
21. The polypeptide of claim 20, wherein the one or more proteases are selected from the group consisting of FAPa, CTSL1, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 and MMP14.
22. The polypeptide of claim 21, wherein the proteases is selected from MMP1, MMP2, MMP9, MMP4, cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin K or cathepsin L.
23. The polypeptide of claim 21, wherein the reference polypeptide sequence is present in a naturally occurring polypeptide substrate for FAPa, CTSL1, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, and an MMP selected from MMP1, MMP2, MMP9 and MMP14, or a combination thereof.
24. The polypeptide of any one of claims 1-23, wherein the cleavable moiety is cleaved with reduced catalytic efficiency by one or more proteases than a reference polypeptide sequence.
25. The polypeptide of claim 24, wherein the cleavable moiety is cleaved with reduced catalytic efficiency by one or more serum proteases.
26. The polypeptide of claim 25, wherein the cleavable moiety is cleaved with reduced catalytic efficiency by one or more hepatic proteases.
27. The polypeptide of claim 25, wherein the cleavable moiety is cleaved with reduced catalytic efficiency by one or more Factor Xa, hepsin or thrombin.

28. The polypeptide of any one of claims 1-25, wherein the polypeptide is operably linked to a moiety selected from the group consisting of a polypeptide moiety, a lipid moiety, a nucleic acid moiety, a detectable moiety, and a small molecule.
29. A recombinant pro-protein comprising:
- a. a recombinant polypeptide comprising a cleavable moiety that is a substrate for a protease, wherein the cleavable moiety comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220 or an amino acid sequence at least at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220; and
 - b. a polypeptide with biological activity,
- wherein the pro-protein has attenuated biological activity and wherein cleavage of the cleavable moiety by the protease produces a polypeptide with biological activity that is not attenuated.
30. The recombinant pro-protein of claim 29, wherein the polypeptide with biological activity comprises a cytokine, chemokine, growth factor, a soluble receptor or a combination thereof.
31. The recombinant pro-protein of claim 29, wherein the polypeptide with biological activity comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain.
32. The recombinant pro-protein of claim 29, wherein the polypeptide with biological activity comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit.
33. The recombinant pro-protein of claim 29, wherein the polypeptide with biological activity comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.
34. A fusion protein comprising:
- a. a signaling protein or molecule; and
 - b. a blocking moiety selected from a steric blocking moiety, a specific blocking moiety, and the combination thereof; and
 - c. a peptide linker that comprises a cleavable moiety as defined in claim 29.

35. The fusion protein of claim 34, wherein (a) and (b) are operably linked by (c).
36. The fusion protein of claim 34, wherein the peptide linker comprises two or more copies of the same cleavable moiety.
37. The fusion protein of claim 34, wherein the steric blocking moiety comprises human serum albumin (HSA) or an anti-HSA antibody.
38. The fusion protein of claim 34, wherein the signaling protein is an interleukin-2 amino acid sequence comprising: (i) a non-native N terminus and/or (ii) a non-native C terminus.
39. The fusion protein of claim 34, wherein the signaling protein is an interleukin-2 amino acid sequence.
40. The fusion protein of claim 34, further comprising one or more half-life extension domains that are not also a specific blocker.
41. The fusion protein of claim 40, wherein the non-native N and/or C termini are generated by circular permutation.
42. A fusion polypeptide comprising a first polypeptide fusion partner linked to a ligand by a protease cleavable linker, wherein the cleavable linker has been optimized for catalytic efficiency, and wherein the ligand has been optionally modified, wherein the first polypeptide fusion partner is a blocking moiety which prevents binding of the modified ligand to a target receptor or a subunit of a target receptor until cleavage of the protease cleavable linker.
43. A fusion polypeptide comprising a first polypeptide fusion partner linked to a ligand by a protease cleavable linker, wherein the cleavable linker has been optimized for catalytic efficiency, and wherein the ligand has been optionally modified, including by circularly permutation to create a non-native N-terminus and a new C-terminus as compared to a native ligand, and wherein at least one of the new N-terminus or the new C-terminus of the modified ligand is operably linked to a first polypeptide fusion partner to form a fusion polypeptide wherein the first polypeptide fusion

partner is a blocking moiety which prevents binding of the modified ligand to a target receptor or a subunit of a target receptor until cleavage of the protease cleavable linker.

44. The fusion polypeptide as in claim 42 or 43, wherein the first polypeptide fusion partner is selected from the group consisting of an antibody, an antibody fragment, and an albumin molecule.

45. The fusion polypeptide as in claim 42 or 43, further comprising a second polypeptide fusion partner comprising a second blocking moiety.

46. The fusion polypeptide as in claim 42 or 43, wherein the second polypeptide fusion partner is a different kind of blocking moiety than the first polypeptide fusion partner.

47. The fusion polypeptide as in claim 42 or 43, wherein the first polypeptide fusion partner is albumin and the second polypeptide fusion partner is a domain comprising a complementary amino acid sequence that blocks activity of the cytokine.

48. The fusion polypeptide as in claim 42 or 43, wherein the first polypeptide fusion partner is a steric blocker, such as albumin, and the second polypeptide is a specific blocker, such as a cytokine receptor, portion of a cytokine receptor, a de novo affinity peptide specific for the cytokine, or an antibody or antibody fragment that specifically binds the cytokine of the fusion polypeptide.

49. The fusion polypeptide of claim 47, wherein the second polypeptide fusion partner is the same kind of blocking moiety as the first polypeptide fusion partner.

50. The fusion polypeptide of any one of claims 34, 42, or 43, further comprising a tumor antigen binding component.

51. The fusion polypeptide of any one of claims 34, 42, or 43, further comprising a serum half-life extension domain.

52. The fusion polypeptide as in claim 42 or 43, wherein the ligand is selected from the group consisting of helix bundle proteins and cytokines (including, but not limited to, growth hormone, IL-2, IL-4, IL-5, IL-6, IL-10, IL-22, IL-23p19, IL-11, IL-13, IL-15, IL-12p35, IL-12p40, IL-12p70, IL-21, IL-30 (IL27p28), IL-34, IL-35, IL-35p35, IFN- α , IFN- β , IFN γ , LIF, CNTF, oncostatin M, CLCF-1, GCSF, GM-CSF, EPO, ferritin, leptin, placental lactogen, prolactin, apolipoprotein e), b-trefoil proteins (including, but not limited to, IL-1 α , IL-1 β , IL-1Ra, IL18, IL-33, IL-36Ra, IL-36a, IL-36b, IL-36g, IL-37, IL-38, IL1Hy2, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8a, FGF-8b, FGF-8e, FGF-8f, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, FGF-16, FGF-17, FGF-18, FGF-19, FGF-20, FGF-21, FGF-22, FGF-23), α/β (TIM) barrel proteins (including, but not limited to, triosephosphate isomerase), beta sandwich proteins (including, but not limited to, galectin-1, galectin-3, TNF-beta, seven β -propeller proteins, class I MHC $\alpha 1\alpha 2$ domain, integrin I domain, GYF domain, C1 domain, C2 domain (for example, from cPLA2, PKC, synaptotagmin), PDZ domains, C3d, C5a.

53. The fusion polypeptide as in claim 42 or 43, wherein the ligand comprises IL-2 polypeptide or a fragment or fragments thereof.

54. The fusion polypeptide of any one of 34, 42, or 43, wherein the protease-cleavable linker polypeptide comprises a sequence that is capable of being cleaved by at least one protease selected from the group consisting of a kallikrein, thrombin, chymase, carboxypeptidase A, cathepsin G, cathepsin L, an elastase, a FAP, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, PR-3, granzyme M, a calpain, a matrix metalloproteinase (MMP), a plasminogen activator, a cathepsin, a caspase, a tryptase, and a tumor cell surface protease.

55. The fusion polypeptide of any one of 34, 42, or 43, wherein the cytokine or fragment or mutein thereof is substantially dissociated from the cytokine blocking moiety after the protease-cleavable polypeptide linker is cleaved by a protease.

56. A fusion polypeptide comprising at least one of each of:
- a cytokine polypeptide or functional fragment or mutein thereof [A];
 - a cytokine blocking moiety [B]; and
 - an optimized protease-cleavable polypeptide linker [L];

wherein the blocking moiety is selected from the group consisting of an antibody, an antibody fragment, and an albumin, and wherein the cytokine comprises a circularly permuted cytokine.

57. The fusion polypeptide of claim 56, further comprising a tumor antigen binding component.

58. The fusion polypeptide of claim 56, further comprising a serum half-life extension domain.

59. The fusion polypeptide of claim 56, wherein the cytokine peptide or functional fragment or mutein thereof is selected from the group consisting of helix bundle proteins and cytokines (including, but not limited to, growth hormone, IL-2, IL-4, IL-5, IL-6, IL-10, IL-22, IL-23p19, IL-11, IL-13, IL-15, IL-12p35, IL-12p40, IL-17p70, IL-21, IL-30 (IL27p28), IL-34, IL-35, IL-35p35, IFN- β , IFN γ , LIF, CNTF, Oncostatin M, CLCF-1, GCSF, GM-CSF, EPO, ferritin, leptin, placental lactogen, prolactin, apolipoprotein e), a FAP, an ADAM selected from ADAM 8, ADAM 9, ADAM 10, ADAM12 ADAM17, and ADAMTS1, b-trefoil proteins (including, but not limited to, IL-1 α , IL-1 β , IL-1Ra, IL18, IL-33, IL-36Ra, IL-36a, IL-36b, IL-36g, IL-37, IL-38, IL1Hy2, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8a, FGF-8b, FGF-8e, FGF-8f, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, FGF-16, FGF-17, FGF-18, FGF-19, FGF-20, FGF-21, FGF-22, FGF-23), α/β (TIM) barrel proteins (including, but not limited to, triosephosphate isomerase), beta sandwich proteins (including, but not limited to, galectin-1, galectin-3, TNF-beta, seven β -propeller proteins, class 1 MHC $\alpha 1\alpha 2$ domain, integrin I domain, GYF domain, C1 domain, C2 domain (for example, from cPLA2, PKC, synaptotagmin), PDZ domains, C3d, C5a.

60. The fusion polypeptide of claim 56, wherein the cytokine peptide or functional fragment or mutein thereof comprises IL-2.

61. The fusion polypeptide of claim 56, wherein the cytokine blocking moiety comprises a ligand binding domain or fragment or mutein of a cognate receptor for the cytokine, a single domain antibody or scFv that binds the cytokine polypeptide or functional fragment or mutein thereof, or an antibody or antibody fragment that binds a receptor of the cytokine.

62. The fusion polypeptide of claim 56, wherein the antibody is a single domain antibody or scFv.
63. The fusion polypeptide of claim 56, wherein the blocking moiety extends the serum half-life of the cytokine or fragment thereof.
64. A fusion polypeptide comprising a protease cleavable moiety, wherein the sequence is catalytically optimized for cleavage by certain proteases and wherein protease cleavage renders the composition inducible in a tumor microenvironment.
65. The fusion polypeptide of claim 64, further comprising a biologically inactive polypeptide, wherein cleavage of the cleavable moiety by the protease converts the biologically inactive polypeptide to a biologically active polypeptide
66. The fusion polypeptide of claim 65, wherein the biologically inactive polypeptide comprises a cytokine, chemokine, growth factor, or soluble receptor.
67. The fusion polypeptide of claim 65, wherein the biologically inactive polypeptide comprises at least one of an extracellular domain, a transmembrane domain, and an intracellular domain.
68. The fusion polypeptide of claim 65, wherein the biologically inactive polypeptide comprises a cell surface receptor, a chimeric antigen receptor (CAR), or a T Cell Receptor (TCR) subunit.
69. The fusion polypeptide of claim 65, wherein the biologically inactive polypeptide comprises an antigen-binding polypeptide, an antibody or an antigen-binding portion thereof.
70. A nucleic acid encoding the polypeptide of any one of claims 7-69.
71. A vector comprising the nucleic acid of claim 70.

72. A host cell comprising the vector of claim 71.
73. A method of making a pharmaceutical composition, comprising culturing the host cell of claim 72 under suitable conditions for expression and collection of desired polypeptides.
74. A method of using a polypeptide of any one of claims 1-69 comprising administering an effective amount of the pharmaceutical composition to a subject in need thereof.




FIG. 1A

2/94




FIG. 1B

3/94




FIG. 1C




FIG. 1D

5/94




FIG. 2

6/94




FIG. 3

7/94




FIG. 4A

8/94





FIG. 4B

9/94

**FIG. 5**

10/94




FIG. 6A




FIG. 6B

11/94




FIG. 7B




FIG. 7D

FIG. 7C

12/94




FIG. 7E




FIG. 7G




FIG. 7F




FIG. 7H

13/94




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 8E




FIG. 8F

14/94




FIG. 9A

FIG. 9B

FIG. 9C




FIG. 9D

FIG. 9E

FIG. 9F

15/94




FIG. 9H




FIG. 9J




FIG. 9G




FIG. 9I

16/94




FIG. 9M




FIG. 9P

FIG. 9L

FIG. 9O

FIG. 9K

FIG. 9N

17/94




FIG. 9Q




FIG. 9R




FIG. 9S

18/94




FIG. 9U




FIG. 9V

19/94




FIG. 9X




FIG. 9W

20/94




FIG. 9Z




FIG. 9Y

21/94




FIG. 10

22/94




FIG. 11B




FIG. 11A

23/94




FIG. 12C




FIG. 12D




FIG. 12A




FIG. 12B

24/94




FIG. 12E




FIG. 12F

25/94




FIG. 13

26/94




FIG. 14

27/94




FIG. 15A




FIG. 15B

28/94




FIG. 15C




FIG. 15D

29/94




FIG. 16E




FIG. 16C

FIG. 16A

FIG. 16F

FIG. 16D

FIG. 16B

30/94




FIG. 17A

FIG. 17C

FIG. 17E




FIG. 17B

FIG. 17D

FIG. 17F

31/94




FIG. 18A

FIG. 18B

32/94




FIG. 19B

FIG. 19A

33/94




FIG. 20B




FIG. 20A




FIG. 21A




FIG. 21C




FIG. 21B




FIG. 21D

35/94




FIG. 22B




FIG. 22A

36/94




FIG. 23B




FIG. 23A

37/94




FIG. 24A




FIG. 24B

38/94




FIG. 25B




FIG. 25A

39/94




FIG. 25C

40/94




FIG. 26A




FIG. 26B




FIG. 26C

41/94




FIG. 27A

42/94




FIG. 27C

FIG. 27B

43/94




FIG. 27D

FIG. 27E




FIG. 27F

44/94




FIG. 28B




FIG. 28D




FIG. 28A




FIG. 28C

45/94




FIG. 28F




FIG. 28H




FIG. 28E




FIG. 28G

46/94




FIG. 28I




FIG. 28J

FIG. 28K

FIG. 28L

FIG. 28M

FIG. 28N

47/94




FIG. 29B

FIG. 29A

48/94







FIG. 30

49/94



50/94




FIG. 32A

51/94




FIG. 32B

52/94




FIG. 32B
Continued

53/94




FIG. 32C

54/94





FIG. 32C
Continued

55/94

**FIG. 33**

56/94




FIG. 34A




FIG. 34B




FIG. 34C

57/94




FIG. 35

58/94




FIG. 35 Continued




FIG. 36

60/94




FIG. 37

61/94




FIG. 38

62/94




FIG. 39

63/94




FIG. 40




FIG. 41

65/94




FIG. 42

66/94





FIG. 43

67/94

Enzyme Equivalents by Substrate across Cell Lines



Substrate & Sample	ADAM17_2		CTSL1_1		FAPa_1		FAPa_1		MMP9_1		MMP14_1	
	cond media		cell lysate		cell lysate		cond media		cond media		cond media	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Myofibroblasts	0.00	0.00	13.65	0.61	0.58	0.05	0.00	0.00	0.00	0.00	0.21	0.11
CT26 Parental	0.26	0.02	9.63	0.64	1.22	0.11	1.42	0.24	0.21	0.01	1.10	0.05
CT26-MMP9+	0.30	0.02	8.55	0.65	1.23	0.09	1.71	0.30	0.28	0.01	1.34	0.05
YAMC	0.52	0.03	5.29	0.05	0.45	0.06	2.27	0.32	0.34	0.01	1.19	0.10
MC38	0.32	0.03	17.34	0.82	1.26	0.08	1.10	0.35	0.21	0.01	1.23	0.07

FIG. 44

68/94




FIG. 45

69/94




FIG. 46

70/94




FIG. 47

71/94




FIG. 48

72/94




FIG. 49

73/94




FIG. 50

74/94




FIG. 51

75/94




FIG. 52

76/94




FIG. 53

77/94





FIG. 54

78/94

**FIG. 55**

79/94

SERUM ONLY
ACP153 MMP9 T24
ACP153 SERUM T24
ACP153 PBS T24
ACP153 PBS T0
IL2




FIG. 56B

SERUM ONLY
ACP16 MMP9 T24
ACP16 SERUM T24
ACP16 PBS T24
ACP16 PBS T0
IL2




FIG. 56A

Intact fusion
Intermediate cleavage
IL2 Dimer
O-linked glycan IL2
IL2

80/94




FIG. 56C





FIG. 57A

82/94



Western Blot probed with IL2 antibody

FIG. 57B

83/94




FIG. 58A

84/94




FIG. 58B




FIG. 58C

85/94




FIG. 58D

86/94





FIG. 59



87/94




FIG. 60

88/94




FIG. 61

89/94




FIG. 62

90/94




FIG. 62
Continued

91/94

SUBSTRATE				
NAME	SEQUENCE	motif #1	motif #2	motif #3
ALU30-1	g <u>ALFKSSFP</u> sgg GPALYAQ ggsgkggsgk	CTSL1_1	MMP14_1	-
ALU30-2	rgsggGPAGLYAQgsgg GPAGLYAQ ggsgk	MMP14_1	MMP14_1	-
ALU30-3	kkgGPAGLYAQ GPAGLYAQ GPAGLYAQgsr	MMP14_1	MMP14_1	MMP14_1
ALU30-4	rgGPAGLYAQg GPAGLYAQ ggGPAGLYAQk	MMP14_1	MMP14_1	MMP14_1
ALU30-5	kkgALFKSSFP PGGPAGIG PLAQKLKSSggs	CTSL1_1	FAPa_1	ADAM17_2
ALU30-6	sggPGGPAGIG ALFKSSFP LAQKLKSSggg	FAPa_1	CTSL1_1	ADAM17_2
ALU30-7	rgPLAQKLKSS ALFKSSFP GGPAGIGgggk	ADAM17_2	CTSL1_1	FAPa_1
ALU30-8	gggALFKSSFP PLAQKLKSS PGGPAGIGggr	CTSL1_1	ADAM17_2	FAPa_1
ALU30-9	rgPGGPAGIG PLAQKLKSS ALFKSSFPggg	FAPa_1	ADAM17_2	CTSL1_1
ALU30-10	rggPLAQKLKSS PGGPAGIG ALFKSSFPgk	ADAM17_2	FAPa_1	CTSL1_1
ALU30-11	rsG GPAGLYAQ ALFKSSFP LAQKLKSSggg	MMP14_1	CTSL1_1	ADAM17_2
ALU30-12	ggPLAQKLKSS ALFKSSFP GPAGLYAQggr	ADAM17_2	CTSL1_1	MMP14_1
ALU30-13	ggALFKSSEP GPAGLYAQ PLAQKLKSSggk	CTSL1_1	MMP14_1	ADAM17_2
ALU30-14	rggALFKSSFP PLAQKLKSS GPAGLYAQgk	CTSL1_1	ADAM17_2	MMP14_1
ALU30-15	rggGPAGLYAQ PLAQKLKSS ALFKSSFPgg	MMP14_1	ADAM17_2	CTSL1_1
ALU30-16	sgPLAQKLKSS GPAGLYAQ ALFKSSFPgsk	ADAM17_2	MMP14_1	CTSL1_1
ALU30-17	kkgPGGPAGIG PLAQRLRSS ALFKSSFPgr	FAPa_1	ADAM17_1	CTSL1_1
ALU30-18	ksgPGGPAGIG ALFFSSPP LAQKLKSSggr	FAPa_1	CTSL1_2	ADAM17_2
ALU30-19	sggFPRSGGS FNPRTFGSK RKRGRGSRGggg	thrombin	factor Xa	hepsin

FIG. 63

92/94

CLEAVAGE ENZYME							
MMP9	MMP14	FAPa	CTSL1	ADAM17	Factor Xa	Thrombin	Hepsin
+++	++	+	++++	-	+	++++	-
+++	+	++++	-	-	-	+	-
++++	++	++++	-	-	-	++++	-
++++	++	++++	-	-	-	+++	-
-	-	++++	++	+	+++	+	+
++	-	++	++++	++	+	+++	+
+	-	-	++++	++++	+	+	+
-	-	-	++	++++	+	++	+++
-	-	++	++	+++	+	++	+++
+++	-	++	++	++++	+	++	+
+	++++	-	+++	+++	+	++	+
+	++	+++	++	+++	+	+	+++
++++	++	++	++	++++	+	+	+
+	++++	-	+++	+++	+	++	+++
++++	++++	+++	+	++	+	+	+++
+++	++++	++++	++	++++	+	++	+
-	-	++++	++++	+++	+	++	+
+++	-	++++	++++	+++	+	++	+
-	-	-	-	-	++++	++++	+++

FIG. 63
Continued

93/94





FIG. 64



94/94

Human 4-1BB HT-1080




FIG. 65A

Human 4-1BB HT-1080




FIG. 65B

WO 2020/232305
INTERNATIONAL SEARCH REPORT

PCT/US2020/032988

International application No.

PCT/US2020/032988

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item I.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

- a. ☒ forming part of the international application as filed:
 ☒ in the form of an Annex C/ST.25 text file.
 ☐ on paper or in the form of an image file.
- b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
- c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 ☐ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).
 ☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

SEQ ID NOs: 195 and 201 were searched.

WO 2020/232305

INTERNATIONAL SEARCH REPORT

PCT/US2020/032988

International application No.

PCT/US2020/032988

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 10, 11, 20-28, 70-74
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See extra sheet(s).

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
claims 1, 2, 5, 7, 8, 12, 29, 30, 34, 35, 37-39, 42, 52, 53, 56, 59, and 60 to the extent that they read on a protease cleavable amino acid sequence/separation moiety of SEQ ID NO:195; and claims 1, 7-9, 12-14, 29, 30, 34-36, 42, 43, 54, and 56 to the extent that they read on a protease cleavable amino acid sequence/separation moiety of SEQ ID NO:201.
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☒ No protest accompanied the payment of additional search fees.




FIG. 1A