

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2002242531 B2**

(54) Title
Antibodies for preventing and treating attaching and effacing escherichia coli (AEEC) associated diseases

(51) International Patent Classification(s)
C07K 16/12 (2006.01)

(21) Application No: **2002242531** (22) Date of Filing: **2002.03.14**

(87) WIPO No: **WO02/074812**

(30) Priority Data

(31) Number	(32) Date	(33) Country
2,339,436	2001.03.15	CA

(43) Publication Date: **2002.10.03**

(43) Publication Journal Date: **2003.03.27**

(44) Accepted Journal Date: **2007.09.20**

(71) Applicant(s)
Valorisation-Recherche, Societe En Commandite

(72) Inventor(s)
Guimond, Marie-Pierre; Fairbrother, John M.; Girard, Francis; Harel, Josee; Batisson, Isabelle

(74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(56) Related Art
Li et al, Infection and Immunity, vol.68 (2000), no.9, pages 5090-5095

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074812 A3

- (51) International Patent Classification⁷: C07K 16/12, 16/02
- (21) International Application Number: PCT/CA02/00353
- (22) International Filing Date: 14 March 2002 (14.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
2,339,436 15 March 2001 (15.03.2001) CA
- (71) Applicant (for all designated States except US): VALORISATION-RECHERCHE, SOCIÉTÉ EN COMMANDITE [CA/CA]; 6e Etage, 3744 Rue Jean-Brillant, Montréal, Québec H3T 1P1 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): FAIRBROTHER, John, M. [CA/CA]; 615 Duquesne, Saint-Hyacinthe, Québec J2S 7A6 (CA). HAREL, Josée [CA/CA]; 347 Jetté, Saint-Bruno, Québec J3V 5N5 (CA). BATAISSON, Isabelle [FR/FR]; 8, rue Dombasle, F-63000 Clermont-Ferrand (FR). GIRARD, Francis [CA/CA]; 2610 Nichols, App. 3, Saint-Hyacinthe, Québec J2S 2W8 (CA). GUIMOND, Marie-Pierre [CA/CA]; 2055 Bennett, App. 202, Montréal, Québec H1V 2T2 (CA).
- (74) Agent: ROBIC; 55, St-Jacques, Montréal, Quebec H2Y 3X2 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
21 August 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/074812 A3

(54) Title: ANTIBODIES FOR PREVENTING AND TREATING ATTACHING AND EFFACING *ESCHERICHIA COLI* (AEEC) ASSOCIATED DISEASES

(57) Abstract: The present invention relates to antibodies immunologically specific for an attaching and effacing *Escherichia coli* (AEEC) virulence-associated protein, products, compositions and methods and to their use thereof in the prevention of an AEEC infection in a mammal. The antibody of the invention is immunologically specific for an AEEC virulence-associated protein and is capable of preventing an *in vivo* AEEC intestinal infection when administered to a mammal. The antibody of the invention is preferably useful for preventing the development of A/E intestinal lesions associated with the AEEC. This is achieved by preferably using IgY antibodies immunologically specific for one or more AEEC virulence-associated proteins, such as Eae, Tir, EspA and Paa.

**ANTIBODIES FOR PREVENTING AND TREATING
ATTACHING AND EFFACING ESCHERICHIA COLI (AEEC) ASSOCIATED
DISEASES**

5

FIELD OF THE INVENTION

The present invention relates to antibodies immunologically specific for an attaching and effacing *Escherichia coli* (AEEC) virulence-associated protein and to the use thereof in the prevention of an AEEC infection in a mammal.

10

BACKGROUND OF THE INVENTION

Escherichia coli is associated with a wide variety of intestinal diseases in human and in animals. Some pathogenic *E. coli* produces attaching and effacing (A/E) lesions, characterized by intimate bacterial adherence to enterocytes and disruption of the underlying cytoskeleton. Such isolates have been termed A/E *E. coli* (AEEC).

15

Reports have shown that A/E lesions are characteristic of enteric pathogens of humans such as enteropathogenic *E. coli* (EPEC) responsible for severe childhood diarrhea in the developing countries, and enterohemorrhagic *E. coli* (EHEC) causing hemorrhagic colitis and hemolytic uremic syndrome (HUS). A/E lesions have also been associated with diarrhea in different animal species such as rabbits, calves, dogs, cats, lambs, and pigs.

20

A/E lesions result from the intimate bacterial adherence to the apical surface of the enterocytes and activation of several chromosomal gene products that interact with components of the host cell. Such gene products are commonly called AEEC virulence-associated proteins. Examples of these virulence-associated proteins are the intimin (Eae) and the secreted proteins Tir (translocated intimin receptor), EspA, EspD, and EspB.

25

30

2.

Currently, the only approach to the control and treatment of AEEC-associated diseases is the use of antibiotics, an approach which is becoming less and less desirable due to problems of bacterial resistance and antibiotic residues. Therefore, alternative approaches to control post-weaning diarrhea must be sought.

5 Alternatively, another approach known in the art suggests the use of AEEC virulence-associated proteins as antigens to immunize an animal in order to stimulate antibody production against the related pathogen. However, no indication of the efficacy of this approach has been shown to prevent AEEC infection in a mammal. Examples of such an alternative approach are shown in US patent 6,204,004 and in
10 international patent applications WO 99/41614; WO 97/40063 and WO 99/24576.

Also known in the art is the use of avian antibodies (IgY) in passive immunization. For instance, WO 00/52055 discloses the use of specific egg yolk IgY antibodies for immunotherapy in animal breeding and animal production. It also discloses the use of IgY antibodies in kits for diagnostics.

15 US patent 5,932,250 is directed to the treatment of vascular disorders particularly arteriosclerosis and atherosclerosis in warm-blooded animals. More specifically, this U.S. patent discloses methods of controlling cholesterol levels, lipid deposits, and the development of atheromatous lesions in warm-blooded animals by the ingestion of egg products containing IgY antibodies raised against *Escherichia coli*
20 proteins.

US patent 6,162,441 discloses a method for producing anti-*E. coli* O157 IgY antibodies in egg-laying hens.

25 The problem with WO 00/52055, 5,932,250 and 6,162,441 is that the *in vivo* efficacy of the IgY produced has not been demonstrated. Indeed, none of these contain evidence with respect to the feasibility of an approach consisting of administering to a mammal, an antibody immunologically specific for an AEEC virulence-associated protein for preventing an *in vivo* AEEC intestinal infection. Therefore, there is still a need for antibodies, egg products and methods for the prevention or treatment of AEEC-mediated diseases.

30

SUMMARY OF THE INVENTION

The present invention relates to antibodies immunologically specific for an attaching and effacing Escherichia coli (AEEC) virulence-associated protein and to the use thereof in the prevention of an AEEC infection in a mammal.

There is described herein an antibody immunologically specific for an AEEC virulence-associated protein, the antibody being capable of preventing an *in vivo* AEEC intestinal infection when administered to a mammal.

In a preferred embodiment, the antibody of the invention is resistant to gastrointestinal digestion. More preferably, the antibody of the invention is an IgY antibody.

The antibody of the invention is preferably capable of preventing the adhesion of the AEEC to the intestine of the mammal. Even more preferably, the antibody of the invention is capable of preventing the development of attaching and effacing (A/E) intestinal lesions associated with the AEEC. This aspect of the invention is achieved by particularly using antibodies immunologically specific for one or more AEEC virulence-associated proteins, such as Eae, Tir, EspA and Paa.

There is also described herein a fowl egg and an isolated yolk of such egg containing an antibody of the invention.

There is also described herein a composition which comprises a biologically acceptable vehicle or carrier and an antibody of the invention above, a fowl egg of the invention, and/or its isolated yolk of the invention.

There is also described herein a food additive which comprises an antibody of the invention, a fowl egg of the invention, a yolk fraction as defined above, and/or a composition of the invention.

There is also described herein a process for obtaining an antibody of the invention, the process comprising the steps of:

(a) actively immunizing a fowl hen for eliciting the production of antibodies in an egg of the hen; and

(b) recovering the antibodies from the egg.

There is also described herein a method for preventing an attaching and effacing *Escherichia coli* (AEEC) infection in a mammal. The method comprises the step of orally administering to the mammal an antibody of the invention, a fowl egg of the invention and/or an isolated yolk of the invention.

5 Accordingly, in a first embodiment of the invention there is provided an isolated IgY antibody immunologically specific for an attaching and effacing *Escherichia coli* (AEEC) virulence-associated protein selected from the group consisting of Eae, Tir, EspA and Paa, said antibody preventing an *in vivo* AEEC intestinal infection when orally administered to a mammal.

10 Accordingly, in a second embodiment of the invention there is provided a fowl egg containing an isolated IgY antibody as defined in the invention.

Accordingly, in a third embodiment of the invention there is provided an isolated yolk of a fowl egg according to the invention.

15 Accordingly, in a fourth embodiment of the invention there is provided a composition comprising at least one element selected from the group consisting of:

- (i) an isolated IgY antibody according to the invention;
 - (ii) a fowl egg according to the invention; and
 - (iii) an isolated yolk according to the invention;
- together with a biologically acceptable vehicle or carrier.

20 Accordingly, in a fifth embodiment of the invention there is provided a food additive comprising at least one element selected from the group consisting of:

- an isolated IgY antibody according to the invention;
- a fowl egg according to the invention;
- an isolated yolk according to the invention; and
- 25 - a composition according to the invention.

Accordingly, in a sixth embodiment of the invention there is provided a process for obtaining an IgY antibody according to the invention, the process comprising the steps of:

(a) actively immunizing a fowl hen for eliciting the production of
30 antibodies in an egg of said hen; and

(b) recovering said antibodies from said egg.

Accordingly, in a seventh embodiment of the invention there is provided use of at least one element selected from the group consisting of:

- an isolated IgY antibody according to the invention;
- a fowl egg according to the invention; and
- an isolated yolk according to the invention;

in the preparation of a composition for preventing a mammalian AEEC infection.

5 Accordingly, in a eighth embodiment of the invention there is provided a method for preventing an attaching and effacing *Escherichia coli* (AEEC) infection in a mammal, said method comprising the step of orally administering to said mammal at least one element selected from the group consisting of :

- an isolated IgY antibody according to the invention;
- 10 - a fowl egg according to the invention; and
- an isolated yolk according to the invention.

A major advantage of the invention is that it provides antibodies, products, compositions and methods useful and efficient for the prevention of an AEEC infection in a mammal.

15 Furthermore, it is particularly an advantage that the antibodies, products, compositions and methods of the invention prevent specifically the development of A/E intestinal lesions associated with the AEEC.

Other objects and advantages of the present invention will be apparent upon reading the following non-restrictive description of several preferred embodiments, made with reference to the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a bar graph showing that AEEC strains originating from the rabbit, pig, dog, bovine, and human (including non-0157 EHEC and EPEC), and producing Eae of the α , β , δ or ϵ subtypes, induced A/E lesions equally well on newborn pig ileal explants.

25 **Figure 2** is a bar graph showing that a replacement of the Eae of the γ subtype by the Eae of the α subtype in an 0157: H7 strain resulted in induction of A/E lesions to a similar extent as observed for the homologous porcine AEEC strain.

Figure 3 is a bar graph showing that antibodies according to a preferred embodiment of the invention are able to significantly block the development of A/E lesions in ileal 30 explants caused by the homologous strain 1390.

Figure 4 is a bar graph showing that antibodies according to a preferred embodiment of the invention are able to significantly block the development of A/E lesions in ileal explants caused by various calf AEEC strains.

5 **Figures 5 and 6** are bar graphs showing that antibodies according to a preferred embodiment of the invention are able to significantly block the development of A/E lesions in ileal explants caused by various human AEEC strains.

10 **Figure 7** is a bar graph showing that antibodies according to a preferred embodiment of the invention are able to significantly block the development of A/E lesions in ileal explants caused by various EPEC strains.

15 **Figures 8A and 8B** are bar graphs showing that oral administration of egg yolk antibodies according to a preferred embodiment of the invention significantly inhibit the development of A/E lesions in the cecum and colon of piglets challenged with an homologous porcine AEEC strain.

20 **Figures 9A and 9B** are bar graphs showing that oral administration of egg yolk antibodies according to a preferred embodiment of the invention inhibit the development of A/E lesions in the cecum and colon of piglets challenged with an O157:H7 AEEC strain.

DETAILED DESCRIPTION OF THE INVENTION

25 As mentioned previously, the present invention relates to a new alternative approach as opposed to antibiotics for the prevention of an AEEC infection in a mammal.

30 More precisely, the present invention relates to antibodies immunologically specific for an AEEC virulence-associated protein, these antibodies being capable of preventing an *in vivo* AEEC infection when administered in a mammal. More particularly, the antibodies of the invention are capable of preventing the adhesion

of the AEEC to the intestine of the mammal, and even more preferably, they prevent the development of A/E intestinal lesions associated with the AEEC.

As used herein, the term "**preventing**" refers to a process by which the AEEC infection is obstructed or delayed.

5 As used herein, the term "**mammal**" refers to any mammal that has the possibility of being infected by an AEEC. Among the mammals which are known to be potentially infected by an AEEC, there are humans, pigs, bovines, ovines, caprines, rabbits, dogs and cats. It will be apparent to one skilled in the art that the antibodies of the invention are intended to be immunologically specific to virulence
10 proteins isolated from a variety of AEEC strains that may cause intestinal lesions, such as those of enteropathogenic *E. coli* (EPEC) and enterohemorrhagic *E. coli* (EHEC). Some examples of EPEC and EHEC strains are shown in Table I.

With respect to antibodies of the invention, the term "**immunologically specific**" refers to antibodies that bind with a relatively high affinity to one or more
15 epitopes of a protein of interest, but which do not substantially recognize and bind molecules other than the one(s) of interest. As used herein, "**antibody**" and "**antibodies**" include all of the possibilities mentioned hereinafter: antibodies or fragments thereof obtained by purification, proteolytic treatment or by genetic engineering, artificial constructs comprising antibodies or fragments thereof and
20 artificial constructs designed to mimic the binding of antibodies or fragments thereof. Such antibodies are discussed in Colcher *et al.* (*Q J Nucl Med* 1998; 42: 225-241). They include complete antibodies, F(ab')₂ fragments, Fab fragments, Fv fragments, scFv fragments, other fragments, CDR peptides and mimetics. These can easily be obtained and prepared by those skilled in the art. For example, enzyme digestion can
25 be used to obtain F(ab')₂ and Fab fragments by subjecting an IgG molecule to pepsin or papain cleavage respectively. Recombinant antibodies are also covered by the present invention.

Alternatively, the antibody of the invention may be an antibody derivative. Such an antibody may comprise an antigen-binding region linked or not to a non-immunoglobulin region. The antigen binding region is an antibody light chain variable
30 domain or heavy chain variable domain. Typically, the antibody comprises both light

and heavy chain variable domains, that can be inserted in constructs such as single chain Fv (scFv) fragments, disulfide-stabilized Fv (dsFv) fragments, multimeric scFv fragments, diabodies, minibodies or other related forms (Colcher *et al. Q J Nucl Med* 1998; 42: 225-241). Such a derivatized antibody may sometimes be preferable since
5 it is devoid of the Fc portion of the natural antibody that can bind to several effectors of the immune system and elicit an immune response when administered to a human or an animal. Indeed, derivatized antibody normally do not lead to immuno-complex disease and complement activation (type III hypersensitivity reaction).

Alternatively, a non-immunoglobulin region is fused to the antigen-binding
10 region of the antibody of the invention. The non-immunoglobulin region is typically a non-immunoglobulin moiety and may be an enzyme, a region derived from a protein having known binding specificity, a region derived from a protein toxin or indeed from any protein expressed by a gene, or a chemical entity showing inhibitory or blocking activity(ies) against the AEEC virulence-associated proteins. The two regions of that
15 modified antibody may be connected via a cleavable or a permanent linker sequence.

Since the AEEC infection is generally localized in the intestinal tract, it is highly preferable that the antibody of the invention be resistant to gastrointestinal digestion. As used herein, the term "**resistant**" refers to an antibody that will substantially retain its immunological function even after being in contact with gastric acids for a period
20 of time necessary to prevent an *in vivo* AEEC infection. Preferably, the antibody of the invention is an avian immunoglobulin, such as IgY. Indeed, it is well known that IgY antibodies show great acid and heat resistance. The antibody of the invention may be also a human or animal immunoglobulin such as IgG1, IgG2, IgG3, IgG4, IgM, IgA, IgE or IgD carrying rat or mouse variable regions (chimeric) or CDRs
25 (humanized or "animalized"). However, the invention is not restricted to IgY antibodies since it is conceivable that the resistance capacity of the antibody, if lacking, can be provided or increased by genetic engineering or by any other way known to one skilled in the art. Furthermore, the antibody of the invention may also be conjugated to any suitable carrier known to one skilled in the art in order to increase its resistance
30 to gastric acids or to provide, for instance, a specific delivery and prolonged retention

of the antibody, either in a targeted local area, such as the intestine, or for a systemic application.

As mentioned above, the antibody of the invention is immunologically specific to one or more AEEC virulence-associated proteins. The preferred AEEC virulence-associated proteins contemplated by the invention are Eae, Tir, EspA, Paa and immunological derivatives thereof. As used herein, the term "**immunological derivative**" refers to a protein/peptide that possesses an immunological activity that is substantially similar to the immunological activity of the whole protein/peptide, and such immunological activity refers to the capacity of stimulating the production of antibodies immunologically specific to an AEEC virulence-associated protein or derivative thereof. The term "immunological derivative" therefore encompasses "fragments", "segments", "variants", or "analogs" of a protein/peptide.

In a highly preferred embodiment, the present invention uses IgY antibodies since and as mentioned above, they advantageously show gastric acid resistance. Furthermore, IgY antibodies have the advantage of not reacting with mammalian complement, Fc receptor, protein A or protein G. Also, IgY antibodies produced in eggs and their extraction from egg yolks can be performed on a large scale without costly investment.

In this connection and according to another aspect, the present invention is also directed to a process for obtaining the above mentioned antibody immunologically specific for an AEEC virulence-associated protein. Although, many processes known in the art may be suitable to obtain the antibodies contemplated by the inventors, it is preferable that the process of the present invention comprises the steps of : a) actively immunizing a fowl hen for eliciting the production of antibodies in an egg of the hen; and b) recovering the antibodies from the egg. A person skilled in the art will understand that the immunizing step is achieved by well known methods. For instance, the AEEC-virulence protein may be given parenterally, for example intravenously, intramuscularly, subcutaneously. As used herein, the term "**fowl**" refers to any birds capable of being immunized. Among those, the common domesticated chicken is preferred. The process of the invention may also comprises a step of administering at least one booster of the proteins to maintain a

hyperimmune state in the hen. Furthermore, the process of the invention preferably comprises a step of purifying the antibodies from a yolk fraction of the egg. Again, the purification step is achieved with methods well known to one skilled in the art. Such a person will also understand that the boosting step is not limited to being done
5 before the step of laying the egg. Indeed, a booster may also be given to the same hen even during or after the egg laying step.

Consequently, the present invention also relates to a fowl egg which contains an antibody of the invention. The present invention further relates to an isolated yolk of the egg.

10 The present invention also describes a method and compositions for the prevention of AEEC-associated diseases. Many methods could be used to reduce to practice the present invention. However, it is particularly believed that a method which comprises the step of orally administering to a mammal an antibody of the invention, a fowl egg and/or an isolated yolk as previously defined is particularly advantageous
15 from a commercial point of view. Nevertheless, methods that involve a parenteral administration of the antibody of the invention may be considered by one skilled in the art.

According to a preferred embodiment, the composition of the invention comprises at least one element immunologically active against AEEC and a
20 biologically acceptable vehicle or carrier. Such an element may either be an antibody, a fowl egg or an isolated yolk as defined above. For preparing the compositions of the present invention, methods well-known in the art may be used. As used herein, the term "**immunologically active**", or reference to the immunological activity of an element, such as an antibody of the invention, refers in that instance to the ability of
25 such antibody to prevent an AEEC infection in a mammal by binding to an AEEC virulence-associated protein. As used herein, the term "**biologically acceptable**" refers to a vehicle or a carrier that can be safely administered to a mammal, particularly humans and animals, without overly negative or toxic side effects. Such vehicle or carrier may be used for various purposes, such as preserving agents,
30 solubilizing agents, stabilizing agents, wetting agents, emulsifiers, sweeteners,

colorants, odorants, salts, buffers, coating agents or antioxidants and the like. They may be readily prepared by those of skill in this art using well known methods.

According to another embodiment, the composition of the invention is formulated under the form of a pharmaceutical or a nutraceutical composition. The present invention also provides a food additive comprising at least one of the above mentioned elements and further comprises a composition as defined above. It will be clear to one skilled in the art that although the compositions of the invention are preferably administered orally, they may be administered by any other suitable route. Indeed, it is conceivable that they could be given by other means. In the case that the compositions are given orally, they may be in the form of tablets, capsules, powder, syrups, etc.

The compositions of the invention may be used in conjunction with pharmaceutical compositions known in the art. For instance, one may find it advantageous to combine one of the elements of the composition of the invention with other active agents that may be used to treat or prevent diseases others than those induced by AEEC.

The amount of specific antibodies that is administered to a human or an animal or that is present in the composition of the invention is a therapeutically effective amount. A therapeutically effective amount of antibody is that amount necessary for obtaining beneficial results without causing overly negative secondary effects in the host to which the antibody or composition is administered. Moreover, an effective amount of an antibody for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such an amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease.

The exact amount of antibody or of each of the components in the composition and amount of the composition to be administered will vary according to factors such as the type of the condition to be treated, the other ingredients in the composition, the mode of administration, the age and weight of the mammal, etc. Without being bound

by any particular dosage, it is believed that for instance for oral administration, a daily dosage of **about 100 to about 600 mg/kg** of lyophilised egg yolk containing antibodies immunologically specific for an AEEC virulence-associated protein (usually present as part of a composition as indicated above) may be suitable for preventing a mammalian AEEC infection in a typical adult. This dosage may be repeated as often as appropriate. Typically, administration may be 1 to 21 times a week. If side effects develop, the amount and/or frequency of the dosage can be reduced.

EXAMPLE

The following example is illustrative of the wide range of applicability of the present invention and is not intended to limit its scope. Modifications and variations can be made therein without departing from the spirit and scope of the invention. Although any method and material similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred methods and materials are described.

Summary

The inventors have successfully produced egg yolk antibodies against purified fusion proteins of the known attaching and effacing virulence factors Eae, EspA, EspB, EspD, and Tir and of a new putative attaching and effacing virulence factor Paa. They have demonstrated that only the anti-Eae, anti-Tir, and anti-Paa antibodies were able to significantly block the development of attaching and effacing (A/E) lesions due to the homologous porcine *E. coli* strain *ex vivo* in the pig ileal organ culture model. These antibodies were also able to block the development of A/E lesions due to attaching and effacing *E. coli* originating from various other animal species such as the bovine and dog, and from humans, including both O157:H7 and non-O157:H7 *E. coli*. Finally, the inventors have demonstrated that the anti-Eae antibodies were able to significantly reduce the development of A/E lesions due to the homologous pig and O157:H7 human attaching and effacing *E. coli in vivo* in a

newborn pig infection model. These results clearly demonstrate that chicken egg yolk antibodies specific for certain of the virulence factors involved in attachment and effacement are able to block bacterial infection and development of intestinal lesions in the pig and thus are a potential candidate for use in the treatment of infections
5 caused by attaching and effacing *E. coli*. They are also a potential candidate as a feed additive for oral administration to cattle in order to eliminate O157:H7 and other attaching and effacing *E. coli* from the intestine and prevent contamination of meat and consequent infection of humans.

10 **Materials and Methods.**

Bacterial strains, plasmids and media. *E. coli* M155(pREP4TM) (Qiagen) was used as the host strain for recombinant proteins. The porcine attaching and effacing *E. coli* (AEEC) P86-1390 (serogroup O45, streptomycin resistant, Sm^R) was isolated at the
15 Faculté de Médecine Vétérinaire, Saint-Hyacinthe, Québec, Canada from a 4-week-old pig with postweaning diarrhea. O45 strain P86-1390 induces typical attaching and effacing (A/E) lesions (Zhu *et al.*, 1994 *Infect Immun* 62: 4153-4159; Zhu *et al.*, 1995 *Can J Vet Res* 59: 118-123) and its genomic DNA was used as template to amplify the virulence factor genes carried on the locus of enterocyte effacement (LEE).

20 For explant model challenge (see Table 2 for strains characteristics), bacteria were grown overnight in Trypticase Soy Broth (TSB, Difco) with agitation (150 rpm) at 37°C, then transferred in Dubelcco's Modified Eagle's Medium (DMEM, GibcoBRL) and grown to early exponential phase prior to use (OD₆₀₀ 0.7, corresponding to approximately 2.0 X 10⁸ CFU, determined by use of specific growth curves). 1% final
25 D-Mannose was added to each broth culture to minimize Type-1 fimbriae-mediated adherence prior to infection.

Construction of fusion genes. The 3' end (carboxy) or the entire (mature) *eae* gene, *espA*, *espB*, and *espD* genes, *tir* gene, and the 3' end (carboxy) or the entire
30 (mature) *paa* gene were amplified by PCR using primer pairs listed in Table 3. The amplicons were inserted into the pGEM-TTM vector (Promega) and then introduced

into the pQE-30TM expression vector (Qiagen) using the appropriate cloning site (between *Bam*HI and *Sall* sites for *eae carboxy*, *espA*, *espB*, *espD*, *paa carboxygenes*, between *Bam*HI and *Sph*I for *eae* mature gene, and between *Hind*III and *Sac*I for *tir* gene). Gene fusions were checked by sequencing.

5

Production and purification of fusion proteins. An overnight LB broth preculture was used to inoculate a 1 liter LB broth culture. Cells were grown at 37°C with shaking until the OD_{600nm} was 0.7-0.8. Isopropylthiogalactoside (IPTG) was then added to a final concentration of 1mM. After incubation for about 4 hours, cells were
10 harvested by centrifugation at 4000 x g for 10 minutes and resuspended in two volumes of buffer A (Qiagen). Samples can be centrifuged and the pellet stored frozen at -70°C until use. Samples were thawed at room temperature and resuspended into buffer A. One mg/ml of lysozyme and a final concentration of 100 µM of PMSF were added. The overall suspension was incubated 30 minutes on ice
15 and sonicated. For each ml of sample, 10 µg of Rnase A and 5 µg of Dnase 1 were added and after incubation for 15 minutes on ice, the sample was clarified by centrifugation at 15000 x g for 20 minutes. The supernatant was mixed with a 50% slurry of Ni-NTA resin (Qiagen) in buffer A (8 ml of slurry per liter of cells). After gentle mixing in a rotating tube for one night, the resin was added to a Qiagen column,
20 washed with buffer A until the OD_{280nm} was less than 0.01, washed with buffer B (Qiagen) until the OD_{280nm} was less than 0.02, washed with 0.1 M imidazole in buffer B (20 ml per liter of cells), and eluted with 0.25 M imidazole in buffer B (10 ml per liter of cells). Fractions (1-1.5 ml), collected upon strating the 0.25 M imidazole elution, were analyzed by SDS-PAGE through 15% acrylamide. Those containing proteins
25 of interest were pooled, quantified by the Lowry method, and stored at -70°C.

Western immunoblotting. His-tagged affinity column purified proteins were mixed with an equal volume of 2X Laemmli buffer, boiled for 5 minutes, applied to a 15% SDS-PAGE, and electrotransferred onto 0.2 µm nitrocellulose membrane (Bio Rad).
30 Blots were blocked and washed with 1% bovine serum albumin (BSA) -0.1% TweenTM 20 in Tris-buffered saline (TBS), and incubated overnight with primary

antibodies, including RGS-His antibody (Qiagen) and the anti-Eae, anti-EspA, anti-EspB, anti-EspD, and anti-Tir kindly provided by Dr. Gad Frankel and Dr. Frank Ebel. For chicken antibody (IgY) production analysis, purified IgY specifically directed against each of the LEE virulence factors were used as primary antibodies. Filters were then
5 developed with secondary goat anti-rabbit (1/1000), goat anti-mouse (1/1000) or rabbit anti-chicken (1/5000) HRP-conjugated IgG and with H₂O₂- α -chloronaphthol as the substrate.

Immunization of animals and antibodies purification. Two to eight laying hens
10 were immunized in their pectoral muscle at multiple sites with 50 μ g of purified proteins and emulsified in incomplete Freund's adjuvant on days 1, 14, 28, 42 and 56. M15 (pREP4TM) total proteins was used as negative control for anti-virulence factor IgY production. Eggs were collected from day 28 and kept at 4°C until purification of antibodies. For the purification of antibodies, the yolks were separated
15 from the yolk membrane and egg white, pooled and one volume of PBS 1X was added. The overall mixture was homogenized, mixed with one volume of chloroform, and centrifuged at 15000 x g for 10 minutes. An orange coloured solution containing the vitellus, a yellow semi-solid emulsion of the lipids in chloroform, and a watery phase containing chicken IgY were then observed from the bottom to the top of the
20 tube. Purified IgY were analyzed by SDS-PAGE stained with Coomassie blue and by Western immunoblotting as explained above.

ELISA. Anti-virulence factor IgY titers in yolk eggs were determined using microtiter plates (Immulon 2HB, Dynec) precoated with 100 ng of purified proteins in carbonate
25 buffer (pH 9.6). Purified IgY serially diluted in PBS pH 7.4 containing 1% BSA and 0.05% Tween 20 was added to the wells and incubated for 2 hours at 37°C. After washing thrice with PBS containing 0.05% TweenTM 20, bound antibodies were detected by adding rabbit anti-chicken IgG (1/25000) conjugated to peroxidase (Jackson Immuno Research Laboratories, Inc). After a 10 minutes enzyme-substrate
30 reaction, the absorbance at 405 nm was read and antibody titers were expressed as the log₁₀ of the reciprocal dilution. To monitor non-specific reactions, absorbances

measured with IgY from chickens immunized with the total protein extract from the M14 (pREP4TM) strain were subtracted from absorbances obtained with test samples.

Antibody extraction from egg yolks. Virulence factor-specific IgY was extracted from egg yolks by a method described by [Akita and Nakai, (1993) *Immunol. Methods* 18: 162(e): 155-164; 1993 *Immunol. Methods* 2: 160(2): 207-214], with some modifications. Briefly, egg yolks were separated from albumin then placed onto a towel. Egg yolks were gently rolled onto the towel to removed albumin residues, then punctured to aspirate the yolk without the vitellus. An equal volume of Phosphate Buffered Saline (PBS) was added to the yolks, then homogenized by Vortex agitation. An equal volume of chloroform was added to the solution, then mixed until there was a solid homogenate. The preparation was centrifuged for 5 minutes at 14 000 rpm, and the supernatant containing IgY was removed. The supernatant was lyophilised prior to use.

15

Collection and culture of explant tissues. The explant culture technique was derived from *Zhu et al.*, 1994 (*supra*). Briefly, mucosal tissues from ileum were obtained from newborn colostrum-deprived piglets from a conventional herd. Piglets were tranquilized with ketamin hydrochloride before being euthanized with a pentobarbital overdose. The time lapse between death and initiation of explant cultures was approximately 1 h and each strain was tested in at least 2 piglets. Upon collection, the serosa was carefully removed and mucus was gently discarded with steril swab. Tissues were immersed in sterile complete RPMI 1640TM media (GibcoBRL), transported to the laboratory on ice, and placed on a rocking platform for 30 minutes. Prior to culture, tissues were cut into 5 X 5 mm pieces and placed mucosal side up onto biopsy foam pads (Curtin Matheson Scientific, Inc.) in multidish four-well Nunclon DeltaTM Surface tissue culture plates (Nalge Nunc International). One tissue (now called an explant) was placed on each sponge with 1 sponge per well. Complete RPMI 1640 medium was added to wells without submerging the explants. Plates were incubated at 37°C on a rocker (position 2,5) in 95% O₂ and 5% CO₂ atmosphere.

30

Inoculation and examination of explants. The explants were infected three times at hourly intervals with 50 µl of broth culture (approximately 1×10^7 CFU) applied to the mucosal surface and incubated for 8 hours. To prevent bacterial overgrowth and acidic pH, hourly changes with sterile fresh complete RPMI 1640™ medium were carried out during culture starting 2 hours after initial explant infection.

Treatment with antibodies in the explant model. Broth cultures were incubated at 37°C with an equal volume of lyophilised antibody preparation, previously reconstituted with PBS, for 30 minutes prior to explant infection.

Light microscopy. After culture, explants were fixed in 10% buffered formalin for microscopic examination. The same protocol was used for sections from *in vivo* experimentally infected newborn piglets. Formalin-fixed sections were further placed into Nylon™ tissue biopsy bags (Shandon Inc.), processed, paraffin-embedded, sectioned at 5 µm, and stained with hematoxylin, phloxine, and safranin (HPS) according to standard techniques. Sections stained with HPS were examined by light microscopy for epithelial intimate-adherent bacteria. Each intact villus was examined for the presence of intimate-adherent bacteria, and evaluation of mean epithelial surface covered with intimate-adherent bacteria was performed, according to a scale from 0 to 100%.

Immunofluorescence. Sections on glass slides were deparaffined into xylene for 5 minutes before rehydration into progressive ethanol solutions. Sections were rinsed in PBS then antigenic binding sites were blocked in PBS-1% BSA-0.1% Tween20 solution at 37°C for 20 minutes. Thereafter, sections were rinsed in PSB and incubated at 37°C for 20 minutes with 1:50 dilution of primary antibodies (*Escherichia coli* Serotyping Laboratory, Canada) according to the specific strain serotype. After rinsing out into PSB, sections were incubated at 37°C with 1:200 dilution of goat anti-rabbit FITC-conjugated secondary antibody (Jackson ImmunoResearch Laboratories

inc., USA) for 20 minutes. Sections were finally counterstained with 0.2% Evan's BlueTM (Fisher Scientific Company, USA). Mounted sections were examined with a Leitz DiaplanTM microscope equipped with epifluorescence.

5 **Transmission electron microscopy (TEM).** Small ileum, cecum, or colon sections (3mm x 3mm) were fixed for 2 hours at room temperature in 2.5% (v/v) glutaraldehyde, then rinsed in cacodylate buffer (0.1M cacodylate, pH 7.3) for 1.5 hours with regular changes. Thereafter, tissues were post-fixed for 1 hour at room temperature into 2% osmium tetroxide (OsO₄), then rinsed into water for 1.5 hours
10 with regular changes, dehydrated in graded ethanol series, and finally embedded in Spurr resin (Marivac, Nova Scotia, Canada). Thin sections were mounted on copper grids, stained with uranyl acetate and lead citrate, and examined for AE lesions with a Philips 420TM transmission electron microscope at 80kV (Philips Electronics, The Netherlands).

15

Piglet infection and antibody challenges *in vivo*. 22 newborn piglets from a conventional herd were used to assay effects of anti-eaeM antibodies on bacterial adherence *in vivo*. 12 piglets were colostrum-deprived prior to infection with porcine strain 1390, while the other 10 piglets were colostrum-fed prior to infection with
20 human EHEC strain 85-170. All piglets were kept in cages, and fed with evaporated milk during the experiment. Group 1 (1390 infection) piglets were infected with 1×10^{10} CFU of 1390 EPEC strain in 2 ml of TSB broth for 2 days, and received twice a day egg yolks from hens immunized with sonicate of PREp15 strain as a adherence positive control (n=6), or egg yolks from hens immunized with EaeM fusion protein
25 (n=6). Group 2 (85-170 infection) piglets were infected with 1×10^{10} CFU of 85-170 EHEC strain in 2 ml of TSB broth for 2 days, and received three times a day, lyophilized egg yolks reconstituted in milk from hens immunized with a sonicate of PREp4 strain as an adherence positive control (n=5), or lyophilized egg yolks reconstituted in milk from hens immunized with EaeM fusion protein (n=5). Piglets
30 were monitored daily for any clinical signs of diarrhea, and were necropsied at 48

hours after the initial infection. Piglets were tranquilized with ketamine hydrochloride, then euthanized with an overdose of pentobarbital solution.

Results and Discussion

5

a) Production of fusion proteins

It has clearly been demonstrated in the literature that the intimin (Eae) and the secreted proteins Tir (Translocated intimin receptor), EspA, EspD and EspB are virulence factors playing an important role in the pathogenesis of various AEEC infections and could elicit an antibody response. Furthermore, the inventors have discovered a new protein called Paa (for Porcine attaching and effacing associated) that is also involved in AEEC adhesion to host cells (see section d). These different virulence factors were thus considered as good candidates for protective immunity and/or as markers in a diagnostic test. Also, it was considered that it would be important to produce a fusion protein by using only the C-terminal end of intimin (Eae carboxy) which is involved in receptor recognition. Thus, seven (7) fusion proteins were produced corresponding to the listed proteins in the pQE30 expression vector which links in frame a His6 tag at the N-terminal end of the proteins. Primer pairs specific for each of the virulence factors were chosen to amplify by PCR the entire (Eae, EspA, EspB, EspD, Paa, Tir) or a part (Eae carboxy) of the proteins from genomic DNA of enteropathogenic *E. coli* porcine strain 1390. The fusion proteins were detected by Western blot analysis revealed with anti-histidine antibodies and with antibodies specifically directed against each of the virulence factors. However, the His-Paa protein was only detected in low quantity. Since Paa is predicted to be unstable by the EXPASY program, a fusion with the stable C-terminal end of Paa (Paa carboxy) was carried out.

All fusion proteins were then produced on a large scale and purified on a nickel affinity column in sufficient quantities for immunization of chickens and for use as antigens in the ELISA. The purified His-Eae carboxy, His-Eae, His-EspA, His-EspB, His-EspD, His-Paa and His-Tir proteins were obtained.

30

b) Production and purification of the egg yolk antibodies

The immunization of chickens and the antibody purification technique were carried out from modified techniques as described in the Materials and Methods. SDS-PAGE analysis demonstrated production of IgY as early as 28 days after initial
5 immunization with proteins His-Eae carboxy, His-Eae, His-EspA, His-EspB, His-EspD, His-Paa and His-Tir.

c) Sensitivity and the specificity of the egg yolk antibodies against each of the proteins

10 Western blot analysis showed that purified IgY recognized homologous fusion proteins. The titer of the specific IgY for each protein was then measured by ELISA using homologous purified proteins as the antigen. The ELISA test conditions were determined for each antigen and results showed a high titer ($>1/25000$) for each specific IgY after 42 days. The capacity of each antibody to detect the corresponding
15 native antigen in the homologous strain 1390 and in AEEC strains from different animal species was also examined. However it appeared that the tested conditions and/or the antigen presentation did not favor the detection by ELISA of some virulence factors such as Eae or Paa in wild type bacteria.

20 d) Capacity of specific antibodies to prevent the development of A/E lesions *in vitro* in the pig ileal organ culture model

• *Elaboration of the model*

The first step of this work was to determine the explant culture conditions which
25 would allow AEEC adherence to ileal and cecal epithelial cells of weaned piglets. Different conditions were tested and a rapid technique for microscopic analysis of tissue sections by light microscopy and confirmation by electron microscopy, was set up (see Materials and Methods). A greater and more consistent adherence of the 1390 strain was observed on ileal explants from newborn piglets than on explants
30 from weaned pigs. Hence, the newborn pig explant model was used in subsequent experiments.

The inventors have demonstrated that AEEC strains originating from the rabbit, pig, dog, bovine, and human (including non-O157 EHEC and EPEC), and producing Eae of the α , β , δ , or ϵ subtypes, induced A/E lesions equally well on newborn pig ileal explants (Figure 1). More specifically, figure 1 shows the mean percentage of intact villi showing bacterial adherence on ileal explant sections. Strains from various animal species and from humans show a similar percentage adherence, when compared to the homologous porcine 1390 AEEC strain, except those from O157:H7 serotype (EC505, 43888, STJ348, 85-170, 43895, STJ854, and STJ919 on the graph). All except O157:H7 strains are significantly different from *eae*-mutant strain ICC-170 (negative control). These data validate our model for the study of the attaching and effacing phenotype expression for both homologous and heterologous strains, except for those belonging to O157:H7 serotype. Results are presented as the mean \pm the standard deviation of the mean.

On the other hand, O157:H7 AEEC strains producing Eae of the γ subtype adhered to a much lesser extent to the ileal epithelium. However, replacement of the Eae of the γ subtype by the Eae of the α subtype in an O157:H7 strain resulted in induction of A/E lesions to a similar extent as observed for the homologous porcine AEEC strain (Figure 2). More specifically, Figure 2 shows the effects of intimin subtype switch from gamma to alpha on the mean percentage of intact villi showing bacterial adherence of the human O157:H7 strain 85-170 on ileal explant sections. Complemented double-mutant strain PCVD-438 (intimin alpha from human E2348/69 EPEC strain) shows similar adherence as homologous porcine strain 1390. PICC-55 is a gamma-intimin subtype complemented mutant strain (similar to the wild strain 85-170). These data confirm the problem of gamma-intimin subtype in adhering to the ileal explant model. Results are presented as the mean \pm the standard deviation of the mean. These results also suggest that Eae of the γ subtype of O157:H7 strains recognizes receptors on porcine ileal epithelial cells less well than the Eae of the other known subtypes. Nevertheless, these results confirm that pig ileal explants are an appropriate model for the examination of the effect of specific antibodies on the formation of A/E lesions by AEEC of diverse origin.

- *Blocking of A/E lesions*

The capacity of the specific antibodies to prevent the development of A/E lesions *ex vivo* in the pig explant culture model and *in vivo* in the newborn pig infection model was tested with the homologous porcine strain 1390 and different heterologous AEEC strains. The results showed that the antibodies specific for the mature and carboxy terminal of Eae, Tir and Paa, in the case of the homologous porcine strain 1390, are able to significantly block the development of A/E lesions in ileal explants, for the homologous strain 1390 (Figure 3). More specifically, Figure 3 shows the effect of antibodies on mean percentage of intact villi showing adherence when infected by porcine strain 1390. 1390 anti-T(-) is a positive control for adherence. Results show a significant decrease in bacterial adherence (* on top of column) with anti-EaeC, anti-EaeM, anti-Tir, and anti-PaaM. Results are presented as the mean \pm the standard deviation of the mean. A Kruskal-Wallis test was performed with commercially available software, and post hoc 2-by-2 comparisons were done to assess differences between the groups; $P < 0.0001$ was taken to be significant.

The results show that the antibodies specific for the mature and carboxy terminal of Eae, Tir and Paa, in the case of the homologous porcine strain 1390, are able to significantly block (* on top of column) the development of A/E lesions in ileal explants for all of the tested AEEC strains from the calf (Figure 4), and human, the latter including both O157:H7 EHEC (Figures 5 and 6) and EPEC strains (Figure 7). The anti-EspA, anti-EspB, and anti-EspD antibodies did not affect the development of A/E lesions by any of the tested AEEC strains, with the exception of a human EPEC strain, for which the anti-EspA antibodies did significantly block the development of A/E lesions (Figure 7). Anti-Paa antibodies did not block the development of A/E lesions due to this human EPEC strain which produced Eae but not Paa. Hence, the anti-Eae mature, anti-Tir, and anti-Paa antibodies, and possibly the anti-EspA antibodies, were considered as potential candidates for the blocking of development of A/E lesions *in vivo*.

More specifically, Figure 4 shows the effect of antibodies on mean percentage of intact villi showing adherence when infected by bovine strain B00-H854. B00-H854 represents the strain alone, whereas B00-H854 anti-T(-) is a positive control for adherence. Results show a significant decrease in bacterial adherence (*on top of column) with anti-EaeC, anti-EaeM, and anti-Tir, and, to a lesser extent, with anti-PaaM. Results are presented as the mean \pm the standard deviation of the mean. A Kruskal-Wallis test was performed with commercially available software, and post hoc 2-by-2 comparisons were done to assess difference between the groups; $P < 0.0001$ was taken to be significant.

Figure 5 shows the effect of antibodies on mean percentage of intact villi showing adherence when infected by human O157:H7 EHEC strain STJ348. STJ348 represents the strain alone, whereas STJ348 anti-T(-) is a positive control for adherence, and ICC-170 is an *eae*- mutant strain used as a negative control. Results show a significant decrease in bacterial adherence (*on top of column) with anti-Tir, anti-EaeM, and, to a lesser extent, with anti-PaaC and anti-EaeC. Results are presented as the mean \pm the standard deviation of the mean. A Kruskal-Wallis test was performed with commercially available software, and post hoc 2-by-2 comparisons were done to assess difference between the groups; $P < 0.0001$ was taken to be significant.

Figure 6 shows the effect of antibodies on mean percentage of intact villi showing adherence when infected by human O157:H7 EHEC strain 85-170. 85-170 represents the strain alone, whereas 85-170 anti-T(-) is a positive control for adherence, and ICC-170 is an *eae*- mutant strain used as a negative control. Results show a significant decrease in bacterial adherence (*on top of column) with anti-EaeM, anti-Tir, and, to a lesser extent, with anti-EaeC and anti-PaaM. Results are presented as the mean \pm the standard deviation of the mean. A Kruskal-Wallis test was performed with commercially available software, and post hoc 2-by-2 comparisons were done to assess difference between the groups; $P < 0.0001$ was taken to be significant.

Moreover, Figure 7 shows the effect of antibodies on mean percentage of intact villi showing adherence when infected by human EPEC strain E2348/69. E2348/69 represents the strain alone, whereas E2348/69 anti-T(-) is a positive control for adherence. Results show a significant decrease in bacterial adherence (*on top of column) with anti-EspA, anti-EaeC, anti-Tir, and anti-EaeM. Results are presented as the mean \pm the standard deviation of the mean. A Kruskal-Wallis test was performed with commercially available software, and post hoc 2-by-2 comparisons were done to assess difference between the groups; $P < 0.0001$ was taken to be significant.

10

e) Ability of appropriate egg yolk antibodies to prevent the development of intestinal colonization and diarrhea *in vivo* in the newborn and weaned pig experimental infection models

Since *in vivo* challenge experiments in pigs are costly and time-consuming, the inventors have decided to focus on the evaluation of the effect of the anti-Eae antibodies on the development of A/E lesions *in vivo*. Using the colostrum-deprived newborn piglet model, it was demonstrated that oral administration of the purified anti-Eae egg yolk antibodies significantly inhibited the development of A/E lesions in the cecum and colon of piglets challenged with the homologous porcine AEEC strain (Figures 8A and 8B). More specifically, Figure 8A shows antibody effects in the cecum, whereas Figure 8B shows antibody effects in the colon. Anti-T(-) results are from piglets (n=6) that received egg yolks from non-immunized hens, and represent a positive control for adherence, whereas anti-EaeM represent piglets (n=6) that received egg yolks containing anti-EaeM antibodies directed against mature intimin. Results show a significant decrease in bacterial adherence with anti-EaeM, when compared with anti-T(-). Results are presented as the mean \pm the standard deviation of the mean. Statistical analysis is still in progress.

Since O157:H7 AEEC strains are an important cause of problems in humans, it was decided to also focus on the evaluation of the effect of the anti-Eae antibodies on the development of A/E lesions *in vivo* in an O157:H7 pig challenge model. It has been previously shown that O157:H7 strains induce A/E lesions to a greater extent

30

in colostrum-fed rather than in colostrum-deprived piglets. Using a colostrum-fed 3-day-old pig model, the inventors have demonstrated that oral administration of the purified anti-Eae egg yolk antibodies also inhibited the development of A/E lesions in the cecum and colon of piglets challenged with an O157:H7 AEEC strain (Figures 9a and 9B). More specifically, Figure 9A shows antibody effects in the cecum, whereas Figure 9B shows antibody effects in the colon. Anti-T(-) results are from piglets (n=5) that received lyophilised egg yolks from non-immunized hens, and represent a positive control for adherence, whereas anti-EaeM represents piglets (n=5) that received lyophilised egg yolks containing anti-EaeM antibodies directed against mature intimin. Results show a decrease in bacterial adherence with anti-EaeM, when compared with anti-T(-). Results are presented as the mean \pm the standard deviation of the mean. Statistical analysis is still in progress.

Conclusion

Hence, the inventors have shown, for the first time, that orally administered antibodies specific for AEEC virulence-associated protein, such as the adhesin Eae, are able to inhibit the development of A/E lesions in the live animal. In addition, they have showed that antibodies specific for the Eae of the β subtype and produced by an AEEC of pig origin are able to inhibit the development of A/E lesions due to an O157:H7 AEEC of human origin that produces Eae of the γ subtype. Altogether, these *ex vivo* and *in vivo* results show that antibodies specific for the Eae and Tir of the porcine AEEC are useful for the inhibition of development of A/E lesions due to AEEC infections in the various animal species and in humans. Antibodies specific for Paa are useful, at least, for the inhibition of development of A/E lesions due to the AEEC homologous to the strain used for preparation of the Paa antibodies.

Table 1. Serogroups and serotypes of AEEC

Human EHEC	
Serogroup	O157, O111, O26
Serotype	O4.H-; O5.H-; O16.H6; O26.H11; O26.H21; O26.H32; O46.H31; O48.H21; O55.H7; O91.H10; O91.H21; O98.H-; O111.H2; O111.H8; O111.H-; O113.H21; O117.H14; O118.H12; O119.H6; O125.H-; O126.H8; O128.H2; O145.H-; O157.H7; O157.H-; O172.H-
Human EPEC	
Serogroup	O26, O55, O111, O119, O125, O126, O127, and O128
Serotype	O18a,c.H7; O20.H26; O20.H34; O25.H1; O26.H11; O26.H-; O44.H34; O55.H6; O55.H7; O55.H-; O86.H27; O86.H34; O86.H-; O91.H7; O91.H-; O111.H2; O111.H12; O111.H-; O114.H10; O114.H32; O119.H6; O119.H-; O125.H21; O125.H-; O126.H2; O126.H-; O127.H9; O127.H21; O127.H40; O127.H-; O128.H2; O128.H7; O128.H8; O128.H12; O128.H-; O142.H6; O158.H23.
Rabbit EPEC	
Newborn rabbit serogroup	O2, O8, O15, O103, O109
Weaned rabbit serogroup	O15, O20, O25, O103, O109, O128, O132
Pig EPEC	
Serogroup	O45, O103, O108, NT
Cattle EPEC	
Serogroup	O5, O26, O11, O118

Table 2. Bacterial strains, and characteristics

5

Bacterial strains	Description and origin	Serotype	Genetic characteristics
P86-1390	Porcine EPEC	O45	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
C89-4221	Canine EPEC	O112ab	<i>eaεα+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
C86-4225	Canine EPEC	O49	<i>eaεδ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
RDEC-1	Rabbit EPEC	O15 :NM	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
E22	Rabbit EPEC	O103:H2	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
97-5899-175	Rabbit EPEC	Non-typable	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa-</i>
96-1744-174	Rabbit EPEC	Non-specific typing	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
97-1746-175	Rabbit EPEC	O55	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
E2348/69	Human EPEC	O127:H6	<i>eaεα+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa-</i>
STJ348	EHEC	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
EC505	EHEC	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
STJ919	EHEC	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
STJ854	EHEC	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
43888	EHEC, negative for vt genes	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
43895	EHEC	O157:H7	<i>eaεγ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
85-170	EHEC isolate from United States, cured of phage encoding verotoxin	O157:H7	<i>eaεγ+</i> , <i>Δvt-</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
ICC-170	EHEC	O157:H7	<i>Δeae-</i> , <i>Δvt-</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
PCVD-438	EHEC	O157:H7	<i>eaεα^c</i> , <i>Δvt-</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
PICC-55	EHEC	O157:H7	<i>eaεγ^c</i> , <i>Δvt-</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
B00-H854	Bovine EPEC	O45	<i>eaεε+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
B00-5999	Bovine EPEC	O26	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
FH1299	Non-O157 EHEC	O26	<i>eaefβ+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
FH894	Non-O157 EHEC	O45	<i>eaεε+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>
FH303	Non-O157 EHEC	O103	<i>eaεε+</i> , <i>espA+</i> , <i>tir+</i> , <i>paa+</i>

Table 3. Primers and PCR products

Genes		Sequence of primers 5'-3'	Amplicon size (bp)
<i>Eae</i> carboxy	F	<u>GGATCCGCAACAACCGATCAGAAT</u>	702
	R	<u>CTCGAGTTTTACACAAACAGGAAA</u>	
<i>Eae</i> mature	F	<u>GGATCCAATGGTGAAAAT</u>	2712
	R	<u>AAGCTTTTTTACACAAACAGG</u>	
<i>EspA</i>	F	<u>GGATCCATGGATACATCAACTGCA</u>	585
	R	<u>CTCGAGTTTACCAAGGGATA</u>	
<i>EspB</i>	F	<u>GGATCCATGAATACTATTGATTAT</u>	954
	R	<u>CTCGAGACCAGCTAAGCGAACCGA</u>	
<i>EspD</i>	F	<u>GGATCCATGCTTAATGTAAATAGC</u>	1152
	R	<u>CTCGAGAACTCGACCACTAACAAT</u>	
<i>Tir</i>	F	<u>GAGCTCATGCCTATTGGTAAT</u>	1626
	R	<u>AAGCTTAACGAAACGTGCGG</u>	
<i>Paa</i> carboxy	F	<u>GGATCCCTTTATCTGCGAAAAA</u>	488
	R	<u>CTCGAGAGTGCCTTTCCTGG</u>	
<i>Paa</i> mature	F	<u>GGATCCATGAGGAACATAA</u>	765
	R	<u>CTCGAGAGTGCCTTTCCTGG</u>	

- 5 Although preferred embodiments of the present invention have been described in detail herein and illustrated in the accompanying drawings, it is to be understood that the invention is not limited to these precise embodiments and that various changes and modifications may be effected therein without departing from the scope or spirit of the present invention as defined in the claims.

The claims defining the invention are as follows:

1. An isolated IgY antibody immunologically specific for an attaching and effacing *Escherichia coli* (AEEC) virulence-associated protein selected from the group consisting of Eae, Tir, EspA and Paa, said antibody preventing an *in vivo* AEEC intestinal infection when orally administered to a mammal.
2. The isolated antibody of claim 1, wherein said antibody prevents the adhesion of said AEEC to the intestine of said mammal.
3. The isolated antibody of claim 1 or 2, wherein said antibody prevents the development of attaching and effacing (A/E) intestinal lesions associated with said AEEC.
4. The isolated antibody of any one of claims 1 to 3, wherein said antibody is resistant to gastrointestinal digestion.
5. The isolated antibody of any one of claims 1 to 4, wherein the mammal is selected from the group consisting of humans, pigs, bovines, ovines, caprines, rabbits, dogs and cats.
6. The isolated antibody of any one of claims 1 to 5, wherein the mammal is a human.
7. The isolated antibody of any one of claims 1 to 6, wherein the AEEC is selected from the group consisting of enteropathogenic *E. coli* (EPEC) strains and enterohemorrhagic *E. coli* (EHEC) strains.
8. An isolated antibody according to claim 1, substantially as hereinbefore described with reference to any one of the examples.
9. A fowl egg containing an isolated IgY antibody as defined in any one of claims 1 to 8.
10. The fowl egg of claim 9, wherein said egg is obtained from a fowl immunized against at least one AEEC virulence-associated protein.
11. The fowl egg of claim 9 or 10, wherein said fowl is a chicken.
12. An isolated yolk of a fowl egg according to any one of claims 9 to 11.
13. A composition comprising at least one element selected from the group consisting of:
 - (i) an isolated IgY antibody according to any one of claims 1 to 8;
 - (ii) a fowl egg according to any one of claims 9 to 11; and
 - (iii) an isolated yolk according to claim 12;together with a biologically acceptable vehicle or carrier.

14. The composition of claim 13, wherein said composition is formulated to be administered orally to a mammal.

15. The composition of claim 13 or 14, wherein said composition is formulated under the form of a pharmaceutical composition.

5 16. The composition of claim 13 or 14, wherein said composition is formulated under the form of a nutraceutical composition.

17. A food additive comprising at least one element selected from the group consisting of:

- an isolated IgY antibody according to any one of claims 1 to 8;
- 10 - a fowl egg according to any one of claims 9 to 11;
- an isolated yolk according to claim 12; and
- a composition according to any one of claims 13 to 16.

18. A process for obtaining an IgY antibody according to any one of claims 1 to 8, the process comprising the steps of:

15 (a) actively immunizing a fowl hen for eliciting the production of antibodies in an egg of said hen; and

(b) recovering said antibodies from said egg.

19. The process according to claim 18, further comprising the step of administering at least one booster of at least one AEEC virulence-associated protein to maintain a hyperimmune state in said hen.

20 20. The process according to claim 18 or 19, further comprising the step of purifying said antibodies from a yolk fraction of said egg.

21. A process for obtaining an IgY antibody according to any one of claims 1 to 8, the process substantially as hereinbefore described with reference to any one of the examples.

22. Use of at least one element selected from the group consisting of:

- an isolated IgY antibody according to any one of claims 1 to 8;
- a fowl egg according to any one of claims 9 to 11; and
- an isolated yolk according to claim 12;

30 in the preparation of a composition for preventing a mammalian AEEC infection.

23. A method for preventing an attaching and effacing *Escherichia coli* (AEEC) infection in a mammal, said method comprising the step of orally administering to said mammal at least one element selected from the group consisting of:

- an isolated IgY antibody according to any one of claims 1 to 8;

- a fowl egg according to any one of claims 9 to 11; and
- an isolated yolk according to claim 12.

Dated 13 August, 2007

Valorisation-Recherche, Societe en Commandite

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

5




FIGURE 1




FIGURE 2





FIGURE 3

FIGURE 4






FIGURE 5

FIGURE 6






FIGURE 7



FIGURES 8A AND 8B



FIGURES 9A AND 9B

