



(86) Date de dépôt PCT/PCT Filing Date: 2012/09/18
(87) Date publication PCT/PCT Publication Date: 2013/03/28
(85) Entrée phase nationale/National Entry: 2014/02/24
(86) N° demande PCT/PCT Application No.: GB 2012/052296
(87) N° publication PCT/PCT Publication No.: 2013/041844
(30) Priorités/Priorities: 2011/09/19 (GB1116122.1);
2011/09/19 (GB1116120.5); 2012/02/24 (GB1203257.9);
2012/03/15 (GB1204592.8); 2012/03/29 (GB1205702.2);
2012/05/18 (GB1208749.0); 2012/07/02 (GB1211692.7)

(51) Cl.Int./Int.Cl. *C12N 15/85* (2006.01),
A01K 67/027 (2006.01), *C07K 16/46* (2006.01)

(71) Demandeur/Applicant:
KYMAB LIMITED, GB

(72) Inventeurs/Inventors:
BRADLEY, ALLAN, GB;
FRIEDRICH, GLENN, GB;
LEE, E-CHIANG, GB;
STRIVENS, MARK, GB;
ENGLAND, NICHOLAS, GB

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : ANTICORPS, DOMAINES VARIABLES & CHAINES ADAPTEES POUR UNE UTILISATION HUMAINE
(54) Title: ANTIBODIES, VARIABLE DOMAINS & CHAINS TAILORED FOR HUMAN USE

Recombineered BAC Vectors to add Polymorphic V-regions to the Mouse Genome

1. Cassette Exchange – Replace existing V-regions




Figure 1

(57) Abrégé/Abstract:

The invention relates to the provision of antibody therapeutics and prophylactics that are tailored specifically for human use. The present invention provides libraries, vertebrates and cells, such as transgenic mice or rats or transgenic mouse or rat cells. Furthermore, the invention relates to methods of using the vertebrates to isolate antibodies or nucleotide sequences encoding antibodies. Antibodies, heavy chains, polypeptides, nucleotide sequences, pharmaceutical compositions and uses are also provided by the invention.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number

WO 2013/041844 A3

(43) International Publication Date
28 March 2013 (28.03.2013)

(51) International Patent Classification:

C12N 15/85 (2006.01) C07K 16/46 (2006.01)
A01K 67/027 (2006.01)

(21) International Application Number:

PCT/GB2012/052296

(22) International Filing Date:

18 September 2012 (18.09.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1116122.1	19 September 2011 (19.09.2011)	GB
1116120.5	19 September 2011 (19.09.2011)	GB
1203257.9	24 February 2012 (24.02.2012)	GB
1204592.8	15 March 2012 (15.03.2012)	GB
1205702.2	29 March 2012 (29.03.2012)	GB
1208749.0	18 May 2012 (18.05.2012)	GB
1211692.7	2 July 2012 (02.07.2012)	GB

(71) Applicant (for all designated States except US): **KYMAB LIMITED** [GB/GB]; Meditrina (B260), Babraham Research Campus, Cambridge Cambridgeshire CB22 3AT (GB).

(72) Inventors; and

(71) Applicants (for US only): **BRADLEY, Allan** [GB/GB]; c/o Kymab Limited, Meditrina (B260), Babraham Research Campus, Cambridge Cambridgeshire CB22 3AT (GB). **FRIEDRICH, Glenn** [GB/GB]; c/o Kymab Limited, Meditrina (B260), Babraham Research Campus, Cambridge Cambridgeshire CB22 3AT (GB). **LEE, E-Chiang** [US/GB]; c/o Kymab Limited, Meditrina (B260), Babraham Research Campus, Cambridge Cambridgeshire CB22 3AT (GB). **STRIVENS, Mark** [—/GB]; c/o Kymab Limited, Meditrina (B260), Babraham Research Campus, Cambridge c CB22 3AT (GB). **ENGLAND, Nicholas** [GB/GB]; c/o Kymab Limited, Meditrina (B260), Babraham Research Campus, Cambridge Cambridgeshire CB22 3AT (GB).

(74) Agents: **STEPHEN, Robert** et al.; Olswang LLP, 90 High Holborn, London Greater London WC1V 6XX (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,

[Continued on next page]

(54) Title: ANTIBODIES, VARIABLE DOMAINS & CHAINS TAILORED FOR HUMAN USE

Recombineered BAC Vectors to add Polymorphic V-regions to the Mouse Genome

1. Cassette Exchange — Replace existing V-regions




Figure 1

(57) Abstract: The invention relates to the provision of antibody therapeutics and prophylactics that are tailored specifically for human use. The present invention provides libraries, vertebrates and cells, such as transgenic mice or rats or transgenic mouse or rat cells. Furthermore, the invention relates to methods of using the vertebrates to isolate antibodies or nucleotide sequences encoding antibodies. Antibodies, heavy chains, polypeptides, nucleotide sequences, pharmaceutical compositions and uses are also provided by the invention.

WO 2013/041844 A3

RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:

11 July 2013

ANTIBODIES, VARIABLE DOMAINS & CHAINS TAILORED FOR HUMAN USE**FIELD OF THE INVENTION**

The present invention relates to the provision of antibody therapeutics and prophylactics that are tailored specifically for human use.

The present invention provides libraries, vertebrates and cells, such as transgenic mice or rats or transgenic mouse or rat cells. Furthermore, the invention relates to methods of using the vertebrates to isolate antibodies or nucleotide sequences encoding antibodies. Antibodies, heavy chains, polypeptides, nucleotide sequences, pharmaceutical compositions and uses are also provided by the invention.

BACKGROUND

The state of the art provides non-human vertebrates (eg, mice and rats) and cells comprising transgenic immunoglobulin loci, such loci comprising human variable (V), diversity (D) and/or joining (J) segments, and optionally human constant regions. Alternatively, endogenous constant regions of the host vertebrate (eg, mouse or rat constant regions) are provided in the transgenic loci. Methods of constructing such transgenic vertebrates and use of these to generate antibodies and nucleic acids thereof following antigen immunisation are known in the art, eg, see US7501552 (Medarex), US5939598 (Abgenix), US6130364 (Abgenix), WO02/066630 (Regeneron), WO2011004192 (Genome Research Limited), WO2009076464, WO2009143472 and WO2010039900 (Ablexis), the disclosures of which are explicitly incorporated herein. Such transgenic loci in the art include varying amounts of the human V(D) J repertoire. Existing transgenic immunoglobulin loci are based on a single human DNA source. The potential diversity of human antibody variable regions in non-human vertebrates bearing such transgenic loci is thus confined.

The inventors considered that it would be desirable to tailor the genomes of these transgenic non-human vertebrates (and thus antibody and antibody chain products of these) to address the variability – and commonality – in the natural antibody gene usage of humans. The inventors wanted to do this in order to better address human use of antibody-based therapeutic and prophylactic drugs.

It would be desirable also to provide for novel and potentially expanded repertoire and diversity of human variable regions in transgenic immunoglobulin loci and non-human vertebrates harbouring these, as well as in antibodies produced following immunisation of such animals.

SUMMARY OF THE INVENTION

The present invention has been developed from extensive bioinformatics analysis of natural antibody gene segment distributions across a myriad of different human populations and across more than two thousand samples from human individuals. The inventors have undertaken this huge task to more thoroughly understand and design non-human vertebrate systems and resultant antibodies to better address human medical therapeutics as a whole, as well as to enable rational design to address specific ethnic populations of humans. Using such rational design, the inventors have constructed transgenic non-human vertebrates and isolated antibodies, antibody chains and cells expressing these in a way that yields products that utilise gene segments that have been purposely included on the basis of the human bioinformatics analysis. The examples illustrate worked experiments where the inventors isolated many cells and antibodies to this effect.

The invention also relates to synthetically-extended & ethnically-diverse superhuman immunoglobulin gene repertoires. The present invention thus provides for novel and potentially expanded synthetic immunoglobulin diversities, thus providing a pool of diversity from which human antibody therapeutic leads can be selected. This expanded pool is useful when seeking to find antibodies with desirable characteristics, such as relatively high affinity to target antigen without the need for further affinity maturation (eg, using laborious *in vitro* techniques such as phage or ribosome display), or improved biophysical characteristics, or to address targets and new epitopes that have previously been difficult to address with antibodies are not reached by prior antibody binding sites.

The invention also provides for diversity that is potentially biased towards variable gene usage common to members of a specific human population, which is useful for generating antibodies for treating and/or preventing diseases or conditions within such population. This ability to bias the

antibody repertoire allows one to tailor antibody therapeutics with the aim of more effectively treating and/or preventing disease or medical conditions in specific human populations.

The present inventors realised the possibility of providing immunoglobulin gene segments from disparate sources in transgenic loci, in order to provide for novel and potentially-expanded antibody diversities from which antibody therapeutics (and antibody tool reagents) could be generated. This opens up the potential of transgenic human-mouse/rat technologies to the possibility of interrogating different and possibly larger antibody sequence-spaces than has hitherto been possible.

In rationally designing transgenic antibody loci, as well as antibodies and antibody chains, the inventors also realised that a relatively long HCDR3 length (at least 20 amino acids) is often desirable to address epitopes. For example, naturally-occurring antibodies have been isolated from humans infected with infectious disease pathogens, such antibodies having a long HCDR3 length. Neutralising antibodies have been found in this respect. A long HCDR3 length would be desirable to address other antigens (eg, receptor clefts or enzyme active sites), not just limited to infectious disease pathogens, and thus the inventors realised the general desirability of the possibility of engineering transgenic loci to be able to produce long HCDR3 antibodies and heavy chains. The inventors, through laborious execution of bioinformatics on in excess of 2000 human DNA samples *via* the 1000 Genomes project together with rational sequence choices, identified that the inclusion of the specific human gene segment variant JH6*02 is desirable for producing long HCDR3 antibodies and chains.

Additional rational design and bioinformatics has led the inventors to realise that specific human constant region variants are conserved across many diverse human populations. The inventors realised that this opens up the possibility of making a choice to humanise antibodies, chains and variable domains by using such specific constant regions in products, rather than arbitrarily choosing the human constant region (or a synthetic version of a human constant region). This aspect of the invention also enables one to tailor antibody-based drugs to specific human ethnic populations, thereby more closely matching drug to patient (and thus disease setting) than has hitherto been performed. It can be a problem in the state of the art that antibodies are humanised with an arbitrary choice of human constant region (presumably derived from one (often unknown) ethnic

population or non-naturally occurring) that does not function as well in patients of a different human ethnic population. This is important, since the constant region has the major role in providing antibody effector functions, eg, for antibody recycling, cellular and complement recruitment and for cell killing.

To this end, in a first configuration of the invention, there is provided

First Configuration

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) comprising a genome having a superhuman immunoglobulin heavy chain human VH and/or D and/or J gene repertoire.

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) comprising a genome having a superhuman immunoglobulin light chain human VL gene repertoire; optionally wherein the vertebrate or cell is according to the first configuration.

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus (eg, a heavy chain locus or a light chain locus), said locus comprising immunoglobulin gene segments according to the first and second human immunoglobulin gene segments (optionally V segments) as mentioned below operably connected upstream of an immunoglobulin constant region; optionally wherein the genome is homozygous for said transgenic immunoglobulin locus;
optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or
optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or
optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region; wherein the transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, a first (optionally a V segment) of said gene segments and a second (optionally a V segment) of said gene segments being different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related; optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises first and second transgenic immunoglobulin loci, each locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region;

wherein (i) the first transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, (ii) the second transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments; and (iii) wherein a first (optionally a V) gene segment of said first locus and a second (optionally a V) gene segment of said second gene locus are different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

optionally wherein the first and second loci are on different chromosomes (optionally chromosomes with the same chromosome number) in said genome;

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

A method of constructing a cell (eg, an ES cell) according to the invention, the method comprising

- (a) identifying functional V and J (and optionally D) gene segments of the genome sequence of a (or said) first human individual;
- (b) identifying one or more functional V and/or D and/or J gene segments of the genome sequence of a (or said) second human individual, wherein these additional gene segments are not found in the genome sequence of the first individual;
- (c) and constructing a transgenic immunoglobulin locus in the cell, wherein the gene segments of (a) and (b) are provided in the locus operably connected upstream of a constant region.

In one embodiment, the gene segment(s) in step (b) are identified from an immunoglobulin gene database selected from the 1000 Genomes, Ensembl, Genbank and IMGT databases.

Throughout this text, Genbank is a reference to Genbank release number 185.0 or 191.0; the 1000 Genomes database is Phase 1, release v3, 16th March 2012; the Ensembl database is assembly GRCh37.p8 (10/04/2012); the IMGT database is available at www.imgt.org.

In one embodiment, the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different, optionally wherein the human

immunoglobulin gene segment derived from the genome sequence of the second individual is low-frequency (optionally rare) within the second ethnic population.

This configuration of the invention also provides a method of making a transgenic non-human vertebrate (eg, a mouse or rat), the method comprising

- (a) constructing an ES cell (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain ES cell) by carrying out the method above;
- (b) injecting the ES cell into a donor non-human vertebrate blastocyst (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain blastocyst);
- (c) implanting the blastocyst into a foster non-human vertebrate mother (eg, a C57BL/6N, C57BL/6J, 129S5 or 129Sv strain mouse); and
- (d) obtaining a child from said mother, wherein the child genome comprises a transgenic immunoglobulin locus.

In one embodiment, the invention provides a method of isolating an antibody that binds a predetermined antigen (eg, a bacterial or viral pathogen antigen), the method comprising immunising a non-human vertebrate according to the invention.

Second Configuration

A library of antibody-producing transgenic cells whose genomes collectively encode a repertoire of antibodies, wherein

- (a) a first transgenic cell expresses a first antibody having a chain encoded by a first immunoglobulin gene, the gene comprising a first variable domain nucleotide sequence produced following recombination of a first human unrearranged immunoglobulin gene segment;
- (b) a second transgenic cell expresses a second antibody having a chain encoded by a second immunoglobulin gene, the second gene comprising a second variable domain nucleotide sequence produced following recombination of a second human unrearranged immunoglobulin gene segment, the first and second antibodies being non-identical;
- (c) the first and second gene segments are different and derived from the genome sequences of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

(d) wherein the cells are non-human vertebrate (eg, mouse or rat) cells.

In one embodiment, the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different; optionally wherein the ethnic populations are selected from those identified in the 1000 Genomes database.

In another embodiment, the second human immunoglobulin gene segment is a polymorphic variant of the first human immunoglobulin gene segment; optionally wherein the second gene segment is selected from the group consisting of a gene segment in any of Tables 1 to 7 and 9 to 14 below (eg, selected from Table 13 or Table 14), eg, the second gene segment is a polymorphic variant of VH1-69.

Third Configuration

An isolated antibody having

(a) a heavy chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unrearranged human immunoglobulin V gene segment with a human D and human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments is derived from the genome of an individual from a first human ethnic population; and the other two gene segments are derived from the genome of an individual from a second, different, human ethnic population, and wherein the antibody comprises heavy chain constant regions of said non-human vertebrate (eg, rodent, mouse or rat heavy chain constant regions); and/or

(b) a light chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unrearranged human immunoglobulin V gene segment with a human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments is derived from the genome of an individual from a first human ethnic population (optionally the same as the first population in (a)); and the other gene segment is derived from the genome of an individual from a second, different, human ethnic population (optionally the same as the second population in (a)), and wherein the antibody comprises light chain constant regions of said non-human vertebrate (eg, rodent, mouse or rat heavy light constant regions);

- (c) Optionally wherein each variable domain of the antibody is a human variable domain.
- (d) Optionally wherein the heavy chain constant regions are gamma-type constant regions.

The invention also provides an isolated nucleotide sequence encoding the antibody, optionally wherein the sequence is provided in an antibody expression vector, optionally in a host cell.

The invention also provides a method of producing a human antibody, the method comprising replacing the non-human vertebrate constant regions of the antibody of the third configuration with human antibody constant regions.

The invention also provides a pharmaceutical composition comprising an antibody according to the third configuration, or an antibody produced according to the method above and a diluent, excipient or carrier; optionally wherein the composition is provided in a container connected to an IV needle or syringe or in an IV bag.

The invention also provides an antibody-producing cell that expresses the second antibody recited in any one of the configurations.

In an alternative configuration, the invention contemplates the combination of nucleotide sequences of first and second immunoglobulin gene segments (eg, two or more polymorphic variants of a particular human germline VH or VL gene segment) to provide a synthetic gene segment. Such synthetic gene segment is used, in one embodiment, to build a transgenic immunoglobulin locus, wherein the synthetic gene segment is provided in combination with one or more human variable and J regions (and optionally one or more human D regions) operably connected upstream of a constant region. When provided in the genome of a non-human vertebrate or cell (eg, mouse or rat cell, eg, ES cell), the invention provides for superhuman gene segment diversity. The sequences to be combined can be selected from gene segments that have been observed to be commonly used in human antibodies raised against a particular antigen (eg, a flu antigen, such as haemagglutinin). By combining the sequences, the synthetic gene segment may

recombine *in vivo* to produce an antibody that is well suited to the treatment and/or prevention of a disease or condition (eg, influenza) mediated by said antigen.

Fourth Configuration

A non-human vertebrate (optionally a mouse or a rat) or vertebrate cell whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof so that the mouse is capable of producing an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.

A non-human vertebrate cell (optionally a mouse cell or a rat cell) whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof for producing (eg, in a subsequent progeny cell) an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.

A heavy chain (eg, comprised by an antibody) isolated from a vertebrate of the invention wherein the heavy chain comprises a HCDR3 of at least 20 amino acids.

A method for producing a heavy chain, VH domain or an antibody specific to a target antigen, the method comprising immunizing a non-human vertebrate according to the invention with the antigen and isolating the heavy chain, VH domain or an antibody specific to a target antigen or a cell producing the heavy chain, VH domain or an antibody, wherein the heavy chain, VH domain or an antibody comprises a HCDR3 that is derived from the recombination of human JH6*02 with a VH gene segment and a D gene segment.

A heavy chain, VH domain or an antibody produced by the method.

A B-cell or hybridoma expressing a heavy chain VH domain that is identical to the VH domain of the heavy chain.

A nucleic acid encoding the VH domain of the heavy chain of claim 22, 23 or 28, or encoding the heavy chain.

A vector (eg, a CHO cell or HEK293 cell vector) comprising the nucleic acid; optionally wherein the vector is in a host cell (eg, a CHO cell or HEK293 cell).

A pharmaceutical composition comprising the antibody, heavy chain or VH domain (eg, comprised by an antibody), together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, heavy chain or antibody).

The antibody, heavy chain or VH domain (eg, comprised by an antibody) as above for use in medicine.

The use of an antibody, heavy chain or VH domain (eg, comprised by an antibody) as above in the manufacture of a medicament for treating and/or preventing a medical condition in a human.

Fifth Configuration

A method of producing an antibody heavy chain, the method comprising

- (a) providing an antigen-specific heavy chain variable domain; and
- (b) combining the variable domain with a human heavy chain constant region to produce an antibody heavy chain comprising (in N- to C-terminal direction) the variable domain and the constant region;

wherein

the human heavy chain constant region is an IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region.

An antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region . Optionally, the variable domain comprises mouse-pattern AID somatic mutations.

A polypeptide comprising (in N- to C- terminal direction) a leader sequence, a human variable domain that is specific for an antigen and a human constant region that is an IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region wherein (i) the leader sequence is not the native human variable domain leader sequence; and/or (ii) the variable domain comprises mouse AID-pattern somatic mutations and/or mouse Terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations.

A nucleotide sequence encoding (in 5' to 3' direction) a leader sequence and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region; and the leader sequence being operable for expression of the heavy chain and wherein the leader sequence is not the native human variable domain leader sequence.

A nucleotide sequence encoding (in 5' to 3' direction) a promoter and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region ; and the promoter being operable for expression of the heavy chain and wherein the promoter is not the native human promoter.

A vector (eg, a CHO cell or HEK293 cell vector) comprising a IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region nucleotide sequence that is 3' of a cloning site for the insertion of a human antibody heavy chain variable domain nucleotide sequence, such

that upon insertion of such a variable domain sequence the vector comprises (in 5' to 3' direction) a promoter, a leader sequence, the variable domain sequence and the constant region sequence so that the vector is capable of expressing a human antibody heavy chain when present in a host cell.

Sixth Configuration

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and optionally a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.

A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human variable region gene segments, wherein the plurality comprises at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), a first of said different gene segments is provided in the genome of a

first vertebrate of the population, and a second of said different gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second gene segment.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.

A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.

A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and

optionally a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.

A method of providing an enhanced human immunoglobulin variable region gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human variable region gene segments, wherein the method comprises providing at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein a first of said different gene segments is provided in the genome of a first vertebrate of the population, and a second of said different gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second gene segment.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same Ig locus in said genome.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 and 9 to 14 (eg, selected from Table 13 or Table 14) (eg,IGHJ6-a) and the second gene segment is the corresponding reference sequence.

A population of non-human vertebrates (eg, mice or rats) comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 and 9 to 14 (eg, selected from Table 13 or Table 14) (eg,IGHJ6-a) and the second gene segment is the corresponding reference sequence, wherein the first gene segment is provided in the genome of a first vertebrate of the population, and the second gene segment is provided in the genome of a second vertebrate of the population.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 and 9 to 14 (eg, selected from Table 13 or Table 14) (eg,IGHJ6-a) and the second gene segment is the corresponding reference sequence.

In one aspect of this configuration, the invention relates to human D gene segment variants as described further below.

In one aspect of this configuration, the invention relates to human V gene segment variants as described further below.

In one aspect of this configuration, the invention relates to human J gene segment variants as described further below.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1 to 3: Schematic illustrating a protocol for producing recombineered BAC vectors to add V gene segments into a mouse genome;

Figure 4: Schematic illustrating a protocol for adding V gene segments to a mouse genome using sequential recombinase mediated cassette exchange (sRMCE); and

Figure 5 (in 4 parts): Alignment of 13 IGHV1-69 variants showing the variable (V) coding region only. Nucleotides that differ from VH1-69 variant *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above the corresponding triplet. Boxed regions correspond to CDR1, CDR2 and CDR3 as indicated.

Figure 6 is a schematic illustrating gene segment diversity and the effect of including variant variants in *cis* according to the invention:-

- (a) Situation in a normal person: Recombination on the same chromosome limits combinations of variants, for instance the antibody gene V4-4 can only be recombined within variant 1 to form for instance for instance V4-4-D-J6 or V4-4-D-J2^A. Similarly the variant V4-4^A can't be recombined with either J6 or J2^A from variant 1 and can only be joined with J-genes from variant 2 to form V4-4^A-D-J6^A and V4-4^A-D-J2. V4-4-J2/J6 complexity = 4.
- (b) Situation in a transgenic mouse: Only one variant is provided so the genome is limited. V4-4-J6/J2 complexity = 2.
- (c) Supra mouse of the invention: The variants are added in *cis* and thus can be recombined in every combination, expanding the repertoire. For instance V4-4 can be combined with J6A, J6, J2A or J2 and similarly V4-4A can be recombined with these same J-genes. The V4-4-J6/J2 complexity = 8, which in this simple example is double that of a person and 4X that of a mouse with a single variant.

Figure 7: Alignment of human JH6*02 variants. Nucleotides that differ from JH6 *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above. Accession numbers (eg, J00256) are shown to the left of the IMGT variant name.

Figure 8: Alignment of JH sequences from various species.

Figure 9: Codon Table

Figure 10: BAC database extract

BRIEF DESCRIPTION OF THE TABLES

<u>Table 1:</u>	Human IgH V Polymorphic Variants
<u>Table 2:</u>	Human IgH D Polymorphic Variants
<u>Table 3:</u>	Human IgH J Polymorphic Variants
<u>Table 4:</u>	Human Ig Vk Polymorphic Variants
<u>Table 5:</u>	Human Ig Vλ Polymorphic Variants
<u>Table 6:</u>	Human IgH Jk Polymorphic Variants
<u>Table 7:</u>	Human IgH Jλ Polymorphic Variants
<u>Table 8:</u>	1000 Genomes Project Human Populations
<u>Table 9:</u>	Immunoglobulin Gene Usage in Human Antibody Responses to Infectious Disease Pathogens
<u>Table 10A:</u>	Human IgH JH5 Variant Occurrences
<u>Table 10B:</u>	Non-Synonymous Human IgH JH5 Variants
<u>Table 11A:</u>	Human IgH JH6 Variant Occurrences
<u>Table 11B:</u>	Non-Synonymous Human IgH JH6 Variants
<u>Table 12A:</u>	Human IgH JH2 Variant Occurrences
<u>Table 12B:</u>	Non-Synonymous Human IgH JH2 Variants
<u>Table 13:</u>	Variant Frequency Analyses & Human Population Distributions

<u>Table 14:</u>	Frequent Human Variant Distributions
<u>Table 15:</u>	Human Gene Segment Usage: Heavy Chain Repertoires From Naïve Non-Human Vertebrates
<u>Table 16:</u>	Human Gene Segment Usage: Heavy Chain Repertoires From Immunised Non-Human Vertebrates
<u>Table 17:</u>	Human Gene Segment Usage: Heavy Chain Repertoires From Antigen-Specific Hybridomas
<u>Table 18:</u>	Sequence Correlation Table
<u>Table 19:</u>	Summary Of Function Correlated With Human Gamma Constant Region Sub-Type
<u>Table 20:</u>	Gene Segments Prevalent In Few Human Populations
<u>Table 21:</u>	Genomic and sequence information

DETAILED DESCRIPTION OF THE INVENTION

A suitable source of JH6*02 and other human DNA sequences for use in the invention will be readily apparent to the skilled person. For example, it is possible to collect a DNA sample from a consenting human donor (eg, a cheek swab sample as per the Example herein) from which can be obtained suitable DNA sequences for use in constructing a locus of the invention. Other sources of human DNA are commercially available, as will be known to the skilled person. Alternatively, the skilled person is able to construct gene segment sequence by referring to one or more databases of human Ig gene segment sequences disclosed herein.

An example source for human V, D and J gene segments according to the invention are Bacterial Artificial Chromosomes (RPCI-11 BACs) obtained from Roswell Park Cancer Institute (RPCI)/Invitrogen. See <http://bacpac.chori.org/hmale11.htm>, which describes the BACs as follows:-

"RPCI - 11 Human Male BAC Library

The RPCI-11 Human Male BAC Library (Osoegawa et al., 2001) was constructed using improved cloning techniques (Osoegawa et al., 1998) developed by Kazutoyo Osoegawa. The library was generated by Kazutoyo Osoegawa. Construction was funded by a grant from the National Human Genome Research Institute (NHGRI, NIH) (#1R01RG01165-03). This library was generated according to the new NHGRI/DOE "Guidance on Human Subjects in Large-Scale DNA Sequencing...

"Male blood was obtained via a double-blind selection protocol. Male blood DNA was isolated from one randomly chosen donor (out of 10 male donors)".

- Osoegawa K, Mammoser AG, Wu C, Frengen E, Zeng C, Catanese JJ, de Jong PJ; Genome Res. 2001 Mar;11(3):483-96; "A bacterial artificial chromosome library for sequencing the complete human genome";
- Osoegawa, K., Woon, P.Y., Zhao, B., Frengen, E., Tateno, M., Catanese, J.J., and de Jong, P.J. (1998); "An Improved Approach for Construction of Bacterial Artificial Chromosome Libraries"; Genomics 52, 1-8.

Superhuman Immunoglobulin Gene Repertoires

The invention relates to synthetically-extended & ethnically-diverse superhuman immunoglobulin gene repertoires. The human immunoglobulin repertoires are beyond those found in nature (ie, "Superhuman"), for example, they are more diverse than a natural human repertoire or they comprise combinations of human immunoglobulin gene segments from disparate sources in a way that is non-natural. Thus, the repertoires of the invention are "superhuman" immunoglobulin repertoires, and the invention relates to the application of these in transgenic cells and non-human vertebrates for utility in producing chimaeric antibodies (with the possibility of converting these into fully-human, isolated antibodies using recombinant DNA technology). The present invention thus provides for novel and potentially expanded synthetic immunoglobulin diversities, which provides for a pool of diversity from which antibody therapeutic leads (antibody therapeutics and antibody tool reagents) can be selected. This opens up the potential of transgenic human-mouse/rat technologies to the possibility of interrogating different and possibly larger antibody sequence-spaces than has hitherto been possible. To this end, in one embodiment, the invention provides a SUPERHUMAN MOUSETM (aka SUPRA-MOUSETM) and a SUPERHUMAN RATTM (aka SUPRA-RATTM)

In developing this thinking, the present inventors have realised the possibility of mining the huge genetics resources now available to the skilled person thanks to efforts such as the HapMap Project, 1000 Genomes Project and sundry other immunoglobulin gene databases (see below for more details). Thus, in some embodiments, the inventors realised the application of these genome sequencing developments in the present invention to generate synthetically-produced and ethnically-diverse artificial immunoglobulin gene repertoires. In one aspect, the inventors realised that such repertoires are useful for the production of antibodies having improved affinity and/or biophysical characteristics, and/or wherein the range of epitope specificities produced by means of such repertoire is novel, provides for antibodies to epitopes that have hitherto been intractable by prior transgenic immunoglobulin loci or difficult to address.

The present invention provides libraries, vertebrates and cells, such as transgenic mice or rats or transgenic mouse or rat cells. Furthermore, the invention relates to methods of using the vertebrates to isolate antibodies or nucleotide sequences encoding antibodies. Antibodies, nucleotide sequences, pharmaceutical compositions and uses are also provided by the invention.

Variation Analysis

The present inventors have realized methods and antibody loci designs that harness the power of genetic variation analysis. The reference human genome provides a foundation for experimental work and genetic analysis of human samples. The reference human is a compilation of the genomes from a small number of individuals and for any one segment of the genome a high quality single reference genome for one of the two chromosomes is available. Because the reference genome was assembled from a series of very large insert clones, the identity of these clones is known. Accordingly, experimental work with human genomic DNA is usually conducted on the clones from which the reference sequence was derived.

Individual humans differ in their sequence and recently several individuals have had their genomes sequenced, for instance James Watson and Craig Venter. Comparison of the genome sequence of these individuals has revealed differences between their sequences and the reference genome in both coding and non-coding parts of the genome, approximately 1 in 1000 bases are different. Some variants will be significant and contribute to differences between individuals. In extreme cases

these will result in genetic disease. Variation can be implicated in differing responses to drugs administered to human patients, eg, yielding an undesirable lowering of patient response to treatment.

The 1000-Genomes Project has the objective of identifying the most frequent variations in the human genome. This public domain project involved sequencing the genomes of more than 1000 individuals from diverse ethnic groups, comparing these sequences to the reference and assembling a catalogue of variants. This has enabled the annotation of variants in coding regions, but because this sequence wasn't derived from large clones of DNA, the analysis of the sequence from diploid individuals can't discriminate the distribution of the variation between the maternal and paternally inherited chromosomes. Where more than one variant is identified in a protein coding gene, it is not possible to illuminate the distribution of the pattern of variants in each version of the protein. For example, if two variants are detected in different positions of the same protein in an individual, this could have resulted from one copy with two variants and none in the other or each copy could have just one variant. To illuminate the sequence of real proteins, the 1000-Genome Project has sequenced mother-father-child trios. This allows one to "phase" the sequence variants, in other words identify blocks of sequence that are inherited from one or other parent and deconvolute the variants.

To further understand the variation within the 1000-genome set a tool has been developed that can identify the significant variants (defined as non-synonymous amino acid changes) from a region of DNA from the phased data in the 1000-genome data set. This tool has been made available online <http://www.1000genomes.org/variation-pattern-finder>. This tool allows an investigator to download non-synonymous variation delimited between specific coordinates. The downloaded files are configured as individual genotypes, but the data is phased so the haplotype information and the frequencies of specific halotypes in different populations can be extracted.

The inventors' analysis of the 1000-genome data for the individual human coding segments of the C, V D and J genes from the heavy and light chains reveals that there is significant variation in these segments. Individuals will usually have two different heavy chain alleles and also different light chain alleles at both kappa and lambda loci. The repertoire of antibodies that can be generated

from each allele will be different. This variation will contribute to a better or differing immune response to certain antigens.

Humanized mice that have hitherto been generated with immunoglobulin heavy and light chain loci contain just one type of immunoglobulin locus. Even if these mice contain a full human heavy chain locus, the variation will be less than contained in a typical human because only one set of C, V, D and J genes are available, while a typical human would have two sets.

The inventors have devised ways to improve on this limitation when constructing transgenic non-human vertebrates and cells for human antibody and variable region production *in vivo*.

Mice can be generated with two different loci, each engineered to have a different repertoire of V, D and J segments. This could be in a single mouse or two or more separate mouse strains and would be analogous to or beyond the repertoire found in a normal human. The engineering of such a mouse would go beyond the repertoire described in humanized mice to date which only have one set of alleles.

However, the inventors also realized that this also has limitations, because the different loci would not normally interact to shuffle V, D and J variants between loci. This same limitation is also inherent in a human, thus this system does not utilize the advantage of recombining variants in all combinations.

To go beyond the normal repertoire in humans and take advantage of combinations of C, V, D and J variants the inventors decided, in one embodiment, to provide these on the same chromosome in *cis*. See figure 6. These loci would be characterized by having more than the normal number of J, D or V genes. For example n=6 for the J genes, but including one J6 variant and one J2 variant would increase this to n=8. This could be combined with additional variants for the D and V genes, for example. By detailed analysis of the 1000- Genomes database, the inventors have devised a collection of candidate polymorphic human variant gene segments, eg, JH gene segments (eg, see the examples), that can be built into the design of transgenic heavy and light chain loci in mice for

expressing increasingly diverse and new, synthetic repertoires of human variable regions. Moreover, by utilizing naturally-occurring human variant gene segments, as per embodiments of the invention, this addresses compatibility with human patients since the inventor's analysis has drawn out candidate variants that are naturally conserved and sometimes very prevalent amongst human ethnic populations. Additionally this enables one to tailor the configurations of the invention to provide for antibody-based drugs that better address specific human ethnic populations.

In an example according to any configuration of the invention, loci (and cells and vertebrates comprising these) are provided in which gene segments from different human populations are used. This is desirable to increase antibody gene diversity to better address more diverse human patients. In an example, the gene segments are from first and second different human populations respectively, and thus the second gene segment is found in the second human population, but not so (or rarely) in the first human population. Rarely means, for example, that the gene segment is found in 5, 4, 3, 2, or 1 or zero individuals in the first population in the 1000 Genomes database. For example, the first gene segment may be shown as present in a first population by reference to Table 13 or 14 herein, the second gene segment may be shown as present in the second population by reference to Table 13 and not in the first population. Optionally, the first gene segment may also be shown as being present in the second population by reference to Table 13 or 14.

In any configuration or aspect of the invention, where a V gene segment is used, this may be used optionally with the native leader sequence. For example, use of genomic DNA (eg, from BACs as in the examples) will mean that the native leader will be used for each V gene segment incorporated into the locus and genomes of the invention. In an alternative, the skilled person may wish to inert a non-native leader sequence together with one or more of the V gene segments. Similarly, in any configuration or aspect of the invention, where a V gene segment is used, this may be used optionally with the native 5' UTR sequence. For example, use of genomic DNA (eg, from BACs as in the examples) will mean that the native 5' UTR sequence will be used for each V gene segment incorporated into the locus and genomes of the invention. In an alternative, the skilled person may wish to exclude the native 5' UTR sequence.

The present invention provides, in a first configuration

(a) Superhuman heavy chain gene repertoires

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) comprising a genome having a superhuman immunoglobulin heavy chain human VH and/or D and/or J gene repertoire.

In one aspect the cell of the invention is an embryonic stem cell. For example, the ES cell is derived from the mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain. In one aspect the non-human vertebrate is a rodent, suitably a mouse, and cells of the invention, are rodent cells or ES cells, suitably mouse ES cells. The ES cells of the present invention can be used to generate animals using techniques well known in the art, which comprise injection of the ES cell into a blastocyst followed by implantation of chimaeric blastocysts into females to produce offspring which can be bred and selected for homozygous recombinants having the required insertion. In one aspect the invention relates to a transgenic animal comprised of ES cell-derived tissue and host embryo derived tissue. In one aspect the invention relates to genetically-altered subsequent generation animals, which include animals having a homozygous recombinants for the VDJ and/or VJ regions.

The natural human immunoglobulin gene segment repertoire consists of (see eg, www.imgt.org):-

VH: total-125 ; functional-41

DH: total-27; functional-23

JH: total-8; functional-6

Vk: total-77; functional-38

Jk: total-5; functional-5

V lambda: total-75; functional-31

J lambda: total-7; functional-5

In one embodiment, the vertebrate or cell genome comprises a transgenic immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene segments, one or more human D gene segments and one or more human J gene segments, wherein the plurality of VH gene segments consists of more than the natural human repertoire of functional VH gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

In one embodiment of the vertebrate or cell, the VH gene repertoire consists of a plurality of VH gene segments derived from the genome sequence of a first human individual, supplemented with one or more different VH gene segments derived from the genome sequence of a second, different human individual. Optionally the D and J segments are derived from the genome sequence of the first human individual. Optionally the VH gene segments from the genome sequence of the second individual are selected from the VH gene segments listed in Table 1, 13 or 14. In this way, the locus provides a superhuman repertoire of D gene segments.

Optionally the individuals are not related. Individuals are "not related" in the context of any configuration or aspect of the invention, for example, if one of the individuals does not appear in a family tree of the other individual in the same generation or going back one, two, three or four generations. Alternatively, are not related, for example, if they do not share a common ancestor in the present generation or going back one, two, three or four generations.

In one embodiment of the vertebrate or cell, the transgenic locus comprises more than 41 functional human VH gene segment species, and thus more than the natural human functional repertoire. Optionally the locus comprises at least 42, 43, 44, 45, 46, 47, 48, 49 or 50 functional human VH gene segment species (eg, wherein the locus comprises the full functional VH repertoire of said first individual supplemented with one or more VH gene segments derived from the genome sequence of the second human individual and optionally with one or more VH gene segments derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of VH gene segments that is useful for generating a novel gene and antibody diversity for

use in therapeutic and tool antibody selection.

In one embodiment of the vertebrate or cell, the transgenic locus comprises a first VH gene segment derived from the genome sequence of the first individual and a second VH gene segment derived from the genome sequence of the second individual, wherein the second VH gene segment is a polymorphic variant of the first VH gene segment. For example, the VH gene segments are polymorphic variants of VH1-69 as illustrated in the examples below. Optionally the locus comprises a further polymorphic variant of the first VH gene segment (eg, a variant derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of VH gene segments.

In one embodiment of the vertebrate or cell, the genome (alternatively or additionally to the superhuman VH diversity) comprises a transgenic immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene segments, a plurality of human D gene segments and one or more human J gene segments, wherein the plurality of D gene segments consists of more than the natural human repertoire of functional D gene segments. Optionally the genome is homozygous for said transgenic heavy chain locus.

In one embodiment of the vertebrate or cell, the D gene repertoire consists of a plurality of D gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more different D gene segments derived from the genome sequence of a (or said) second, different human individual. Optionally the individuals are not related. Optionally the J segments are derived from the genome sequence of the first human individual. Optionally the D gene segments from the genome sequence of the second individual are selected from the D gene segments listed in Table 2, 13 or 14. In this way, the locus provides a superhuman repertoire of D gene segments.

In one embodiment of the vertebrate or cell, the transgenic locus comprises more than 23 functional human D gene segment species; optionally wherein the locus comprises at least 24, 25, 26, 27, 28, 29, 30 or 31 functional human D gene segment species (eg, wherein the locus comprises the full functional D repertoire of said first individual supplemented with one or more D gene segments derived from the genome sequence of the second human individual and optionally with one or more

D gene segments derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of D gene segments.

In one embodiment of the vertebrate or cell, the transgenic locus comprises a first D gene segment derived from the genome sequence of the first individual and a second D gene segment derived from the genome sequence of the second individual, wherein the second D gene segment is a polymorphic variant of the first D gene segment. Optionally the locus comprises a further polymorphic variant of the first D gene segment (eg, a variant derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of D gene segments.

In one embodiment of the vertebrate or cell (alternatively or additionally to the superhuman VH and/or JH diversity), the genome comprises a (or said) transgenic immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene segments, one or more human D gene segments and a plurality of human JH gene segments, wherein the plurality of J gene segments consists of more than the natural human repertoire of functional J gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

In one embodiment of the vertebrate or cell, the JH gene repertoire consists of a plurality of J gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more different J gene segments derived from the genome sequence of a (or said) second, different human individual. Optionally the individuals are not related. Optionally D segments are derived from the genome sequence of the first human individual. Optionally the J gene segments from the genome sequence of the second individual are selected from the J gene segments listed in Table 3 13 or 14. In this way, the locus provides a superhuman repertoire of JH gene segments.

In one embodiment of the vertebrate or cell, the transgenic locus comprises more than 6 functional human JH gene segment segments. Optionally the locus comprises at least 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 functional human JH gene segments (eg, wherein the locus comprises the full functional JH repertoire of said first individual supplemented with one or more JH gene segments derived from

the genome sequence of the second human individual and optionally with one or more JH gene segments derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of JH gene segments.

In one embodiment of the vertebrate or cell, the transgenic locus comprises a first JH gene segment derived from the genome sequence of the first individual and a second JH gene segment derived from the genome sequence of the second individual, wherein the second JH gene segment is a polymorphic variant of the first JH gene segment. Optionally the locus comprises a further polymorphic variant of the first JH gene segment (eg, a variant derived from the genome sequence of a third human individual). In this way, the locus provides a superhuman repertoire of JH gene segments.

(b) Superhuman light chain gene repertoires

The first configuration of the invention also provides:-

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) comprising a genome having a superhuman immunoglobulin light chain human VL gene repertoire. Optionally the vertebrate or cell comprises a heavy chain transgene according to aspect (a) of the first configuration. Thus, superhuman diversity is provided in both the heavy and light chain immunoglobulin gene segments in the cell and vertebrate. For example, the genome of the cell or vertebrate is homozygous for the heavy and light chain transgenes and endogenous antibody expression is inactivated. Such a vertebrate is useful for immunisation with a predetermined antigen to produce one or more selected antibodies that bind the antigen and have human variable regions resulting from recombination within the superhuman gene segment repertoire. This provides potentially for a novel antibody and gene sequence space from which to select therapeutic, prophylactic and tool antibodies.

In one embodiment of aspect (b) of the first configuration, the vertebrate or cell genome comprises

- (i) a transgenic immunoglobulin kappa light chain locus comprising a plurality of human

immunoglobulin V κ gene segments and one or more human J gene segments, wherein the plurality of V κ gene segments consists of more than the natural human repertoire of functional V κ gene segments; optionally wherein the genome is homozygous for said transgenic kappa light chain locus; and/or

(ii) a transgenic immunoglobulin lambda light chain locus comprising a plurality of human immunoglobulin V λ gene segments and one or more human J gene segments, wherein the plurality of V λ gene segments consists of more than the natural human repertoire of functional V λ gene segments; optionally wherein the genome is homozygous for said transgenic lambda light chain locus.

In this way, the locus provides a superhuman repertoire of VL gene segments.

In one embodiment of the vertebrate or cell,

(i) the V κ gene repertoire consists of a plurality of V κ gene segments derived from the genome sequence of a first human individual, supplemented with one or more V κ gene segments derived from the genome sequence of a second, different human individual; optionally wherein the individuals are not related; optionally wherein the J segments are derived from the genome sequence of the first human individual; and optionally wherein the V κ gene segments from the genome sequence of the second individual are selected from the V κ gene segments listed in Table 4, 13 or 14; and

(i) the V λ gene repertoire consists of a plurality of V λ gene segments derived from the genome sequence of a first human individual, supplemented with one or more V λ gene segments derived from the genome sequence of a second, different human individual; optionally wherein the individuals are not related; optionally wherein the J segments are derived from the genome sequence of the first human individual; and optionally wherein the V λ gene segments from the genome sequence of the second individual are selected from the V λ gene segments listed in Table 5, 13 or 14.

In this way, the locus provides a superhuman repertoire of VL gene segments.

In one embodiment of the vertebrate or cell,

-the kappa light transgenic locus comprises more than 38 functional human V κ gene segment species; optionally wherein the locus comprises at least 39, 40, 41, 42, 43, 44, 45, 46, 47 or 48 functional human V κ gene segment species (eg, wherein the locus comprises the full functional V κ repertoire of said first individual supplemented with one or more V κ gene segments derived from the genome sequence of the second human individual and optionally with one or more V κ gene segments derived from the genome sequence of a third human individual); and

-the lambda light transgenic locus comprises more than 31 functional human V λ gene segment species; optionally wherein the locus comprises at least 32, 33, 34, 35, 36, 37, 38, 39, 40 or 41 functional human V λ gene segment species (eg, wherein the locus comprises the full functional V λ repertoire of said first individual supplemented with one or more V λ gene segments derived from the genome sequence of the second human individual and optionally with one or more V λ gene segments derived from the genome sequence of a third human individual).

In this way, the locus provides a superhuman repertoire of VL gene segments.

In one embodiment of the vertebrate or cell,

-the kappa light transgenic locus comprises a first V κ gene segment derived from the genome sequence of the first individual and a second V κ gene segment derived from the genome sequence of the second individual, wherein the second V κ gene segment is a polymorphic variant of the first V κ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first V κ gene segment (eg, a variant derived from the genome sequence of a third human individual); and

-the lambda light transgenic locus comprises a first V λ gene segment derived from the genome sequence of the first individual and a second V λ gene segment derived from the genome sequence of the second individual, wherein the second V λ gene segment is a polymorphic variant of the first V λ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first V λ gene segment (eg, a variant derived from the genome sequence of a third human individual).

In this way, the locus provides a superhuman repertoire of VL gene segments.

In one embodiment of the vertebrate or cell, the genome comprises a (or said) transgenic immunoglobulin light chain locus comprising a plurality of human immunoglobulin VL gene segments and a plurality of human JL gene segments, wherein the plurality of J gene segments consists of more than the natural human repertoire of functional J gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

In one embodiment of the vertebrate or cell,

(i) the Jk gene repertoire consists of a plurality of Jk gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more Jk gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein the Vk segments are derived from the genome sequence of the first human individual; optionally wherein the Jk gene segments from the genome sequence of the second individual are selected from the Jk gene segments listed in Table 6, 13 or 14; and

(ii) the Jk gene repertoire consists of a plurality of Jλ gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more Jλ gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein the Vλ segments are derived from the genome sequence of the first human individual; optionally wherein the Jλ gene segments from the genome sequence of the second individual are selected from the Jλ gene segments listed in Table 7, 13 or 14.

In this way, the locus provides a superhuman repertoire of JL gene segments.

In one embodiment of the vertebrate or cell,

(i) the transgenic light chain locus comprises more than 5 functional human Jk gene segment species; optionally wherein the locus comprises at least 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 functional human Jk gene segment species (eg, wherein the locus comprises the full functional Jk repertoire of said first individual supplemented with one or more Jk gene segments derived from the genome sequence of the second human individual and optionally with one or more Jk gene segments derived

from the genome sequence of a third human individual); and/or

(i) the transgenic light chain locus comprises more than 5 functional human J λ gene segment species; optionally wherein the locus comprises at least 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 functional human J λ gene segment species (eg, wherein the locus comprises the full functional J λ repertoire of said first individual supplemented with one or more J λ gene segments derived from the genome sequence of the second human individual and optionally with one or more J λ gene segments derived from the genome sequence of a third human individual).

In this way, the locus provides a superhuman repertoire of JL gene segments.

In one embodiment of the vertebrate or cell,

(i) the kappa light transgenic locus comprises a first J κ gene segment derived from the genome sequence of the first individual and a second J κ gene segment derived from the genome sequence of the second individual, wherein the second J κ gene segment is a polymorphic variant of the first J κ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first J κ gene segment (eg, a variant derived from the genome sequence of a third human individual); and

(ii) the lambda light transgenic locus comprises a first J λ gene segment derived from the genome sequence of the first individual and a second J λ gene segment derived from the genome sequence of the second individual, wherein the second J λ gene segment is a polymorphic variant of the first J λ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first J λ gene segment (eg, a variant derived from the genome sequence of a third human individual).

In this way, the locus provides a superhuman repertoire of JL gene segments.

Further aspects of the first configuration are described below.

The present invention provides, in a second configuration

A library of antibody-producing transgenic cells whose genomes collectively encode a repertoire of antibodies, wherein

- (a) a first transgenic cell expresses a first antibody having a chain (eg, heavy chain) encoded by a first immunoglobulin gene, the gene comprising a first variable domain nucleotide sequence produced following recombination of a first human unrearranged immunoglobulin gene segment (eg, a VH);
- (b) a second transgenic cell expresses a second antibody having a chain (eg, a heavy chain) encoded by a second immunoglobulin gene, the second gene comprising a second variable domain nucleotide sequence produced following recombination of a second human unrearranged immunoglobulin gene segment (eg, a VH), the first and second antibodies being non-identical;
- (c) the first and second gene segments are different and derived from the genome sequences of first and second human individuals respectively, wherein the individuals are different; and optionally not related;
- (d) wherein the cells are non-human vertebrate (eg, mouse or rat) cells (eg, B-cells or hybridomas).

In one embodiment, the library is provided *in vitro*. In another embodiment, the library is provided *in vivo* by one or a plurality of transgenic non-human vertebrates. For example, the or each vertebrate is according to any aspect of the first configuration of the invention.

In one embodiment, the library encodes an antibody repertoire of from 10 to 10^9 antibodies, for example, 10, 20, 30, 40, 50, 100 or 1000 to 10^8 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^7 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^6 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^5 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^4 antibodies. In an example, library encodes an antibody repertoire of at least 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 or 10^{10} antibodies.

The first variable domain nucleotide sequence is produced following recombination of the first human unrearranged immunoglobulin gene segment with one or more other immunoglobulin gene segments (for example, human immunoglobulin gene segments). For example, where the first gene segment is a VH, the first variable domain nucleotide sequence (a VH domain) is produced following recombination of the VH with a human D and JH segments *in vivo*, optionally with somatic hypermutation, in the first transgenic cell or an ancestor thereof. For example, where the first gene segment is a VL, the first variable domain nucleotide sequence (a VL domain) is produced following recombination of the VL with a human JL segment *in vivo*, optionally with somatic hypermutation, in

the first transgenic cell or an ancestor thereof.

The second variable domain nucleotide sequence is produced following recombination of the second human unrearranged immunoglobulin gene segment with one or more other immunoglobulin gene segments (for example, human immunoglobulin gene segments). For example, where the second gene segment is a VH, the second variable domain nucleotide sequence (a VH domain) is produced following recombination of the VH with a human D and JH segments *in vivo*, optionally with somatic hypermutation, in the second transgenic cell or an ancestor thereof. For example, where the second gene segment is a VL, the second variable domain nucleotide sequence (a VL domain) is produced following recombination of the VL with a human JL segment *in vivo*, optionally with somatic hypermutation, in the second transgenic cell or an ancestor thereof.

The first and second gene segments are respectively derived from genome sequences of first and second human individuals. In one example, such a gene segment is isolated or cloned from a sample cell taken from said individual using standard molecular biology techniques as known to the skilled person. The sequence of the gene segment may be mutated (eg, by the introduction of up to 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 nucleotide changes) prior to use in the present invention. In another example, a gene segment is derived by identifying a candidate human immunoglobulin gene segment in a database (see guidance below) and a nucleotide sequence encoding a gene segment for use in the present invention is made by reference (eg, to be identical or a mutant with up to 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 nucleotide changes to the reference sequence) to the database sequence. The skilled person will be aware of methods of obtaining nucleotide sequences by reference to databases or by obtaining from cellular samples.

In one embodiment of the vertebrate, cell or library of any configuration of the invention, the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different. This, therefore, provides for superhuman gene diversity in transgenic loci, cells and vertebrates as per the invention.

Human Populations

Optionally the ethnic populations are selected from those identified in the 1000 Genomes Project of database. In this respect, see Table 8 which provides details of the ethnic populations on which the 1000 Genomes database is based.

N A Rosenberg *et al* (Science 20 December 2002: vol. 298 no. 5602 2342-2343) studied the genetic structure of human populations of differing geographical ancestry. In total, 52 populations were sampled, these being populations with:

African ancestry

(Mbuti Pygmies, Biaka Pygmies, San peoples, and speakers of Niger-Kordofanian languages (Bantu, Yoruba or Mandenka populations),

Eurasian ancestry

(European ancestry (Orkadian, Adyghe, Basque, French, Russians, Italians, Sardinian, Tuscan),

Middle Eastern ancestry (Mozabite, Bedouin, Druze, Palestinians),

Central/South Asian ancestry (Balochi, Brahui, Makrani, Sindhi, Pathan, Burusho, Hazara, Uyghur, Kalash)),

East Asian ancestry

(Han, Dai, Daur, Hezhen, Lahu, Miao, Oroqen, She, Tujia, Tu, Xibo, Yi, Mongolian, Naxi, Cambodian, Japanese, Yakut), Oceanic ancestry (Melanesian, Papuan); or

Americas ancestry

(Karitiana, Surui, Colombian, Maya, Pima).

The International HapMap Project, Nature, 2003 Dec 18;426(6968):789-96, discloses that goal of the HapMap Project: to determine the common patterns of DNA sequence variation in the human genome by determining the genotypes of one million or more sequence variants, their frequencies and the degree of association between them in DNA samples from populations with ancestry from parts of Africa, Asia and Europe. The relevant human populations of differing geographical ancestry include Yoruba, Japanese, Chinese, Northern European and Western European populations. More specifically:-

Utah population with Northern or Western European ancestry (samples collected in 1980 by the Centre d'Etude du Polymorphisme Humain (CEPH));
population with ancestry of Yoruba people from Ibadan, Nigeria;
population with Japanese ancestry; and
population with ancestry of Han Chinese from China.

The authors, citing earlier publications, suggest that ancestral geography is a reasonable basis for sampling human populations.

A suitable sample of human populations from which the populations used in the present invention are selected is as follows:-

(a) European ancestry

- (b) Northern European ancestry; Western European ancestry; Toscani ancestry; British ancestry, Finnish ancestry or Iberian ancestry.
- (c) More specifically, population of Utah residents with Northern and/or Western European ancestry; Toscani population in Italia; British population in England and/or Scotland; Finnish population in Finland; or Iberian population in Spain.

(a) East Asian ancestry

- (b) Japanese ancestry; Chinese ancestry or Vietnamese ancestry.
- (c) More specifically, Japanese population in Toyko, Japan; Han Chinese population in Beijing, China; Chinese Dai population in Xishuangbanna; Kinh population in Ho Chi Minh City, Vietnam; or Chinese population in Denver, Colorado, USA.

(a) West African ancestry

- (b) Yoruba ancestry; Luhya ancestry; Gambian ancestry; or Malawian ancestry.

- (c) More specifically, Yoruba population in Ibadan, Nigeria; Luhya population in Webuye, Kenya; Gambian population in Western Division, The Gambia; or Malawian population in Blantyre, Malawi.

(a) Population of The Americas

- (b) Native American ancestry; Afro-Caribbean ancestry; Mexican ancestry; Puerto Rican ancestry; Columbian ancestry; or Peruvian ancestry.
- (c) More specifically, population of African Ancestry in Southwest US; population of African American in Jackson, MS; population of African Caribbean in Barbados; population of Mexican Ancestry in Los Angeles, CA; population of Puerto Rican in Puerto Rico; population of Colombian in Medellin, Colombia; or population of Peruvian in Lima, Peru.

(a) South Asian ancestry

- (b) Ahom ancestry; Kayadtha ancestry; Reddy ancestry; Maratha; or Punjabi ancestry.
- (c) More specifically, Ahom population in the State of Assam, India; Kayadtha population in Calcutta, India; Reddy population in Hyderabad, India; Maratha population in Bombay, India; or Punjabi population in Lahore, Pakistan.

In any configuration of the invention, in one embodiment, each human population is selected from a population marked “(a)” above.

In any configuration of the invention, in another embodiment, each human population is selected from a population marked “(b)” above.

In any configuration of the invention, in another embodiment, each human population is selected from a population marked “(c)” above.

In one embodiment of the library of the vertebrate, cell or library of the invention, the first and second ethnic populations are selected from the group consisting of an ethnic population with

European ancestry, an ethnic population with East Asian, an ethnic population with West African ancestry, an ethnic population with Americas ancestry and an ethnic population with South Asian ancestry.

In one embodiment of the library of the vertebrate, cell or library of the invention, the first and second ethnic populations are selected from the group consisting of an ethnic population with Northern European ancestry; or an ethnic population with Western European ancestry; or an ethnic population with Toscani ancestry; or an ethnic population with British ancestry; or an ethnic population with Icelandic ancestry; or an ethnic population with Finnish ancestry; or an ethnic population with Iberian ancestry; or an ethnic population with Japanese ancestry; or an ethnic population with Chinese ancestry; or an ethnic population Vietnamese ancestry; or an ethnic population with Yoruba ancestry; or an ethnic population with Luhya ancestry; or an ethnic population with Gambian ancestry; or an ethnic population with Malawian ancestry; or an ethnic population with Native American ancestry; or an ethnic population with Afro-Caribbean ancestry; or an ethnic population with Mexican ancestry; or an ethnic population with Puerto Rican ancestry; or an ethnic population with Columbian ancestry; or an ethnic population with Peruvian ancestry; or an ethnic population with Ahom ancestry; or an ethnic population with Kayadtha ancestry; or an ethnic population with Reddy ancestry; or an ethnic population with Maratha; or an ethnic population with Punjabi ancestry.

In one embodiment of any configuration of the vertebrate, cell or library of the invention, the human immunoglobulin gene segment derived from the genome sequence of the second individual is low-frequency (optionally rare) within the second ethnic population. Optionally human immunoglobulin gene segment has a Minor Allele Frequency (MAF) (cumulative frequency) of between 0.5% - 5%, optionally less than 0.5%, in the second human population, eg, as in the 1000 Genomes database.

In one embodiment of any configuration of the vertebrate, cell or library of the invention, the first variable region nucleotide sequence is produced by recombination of the first human immunoglobulin gene segment with a first J gene segment and optionally a first D gene segment, wherein the first human immunoglobulin gene segment is a V gene segment and the V, D and J segments are derived from the first human population, optionally from the genome of one individual

of the first human population.

In one embodiment of the library of the vertebrate, cell or library of the invention, the second variable region nucleotide sequence is produced by recombination of the second human immunoglobulin gene segment with a second J gene segment and optionally a second D gene segment, wherein the second human immunoglobulin gene segment is a V gene segment derived from the second population and the D and/or J segments are derived from the first human population, optionally the D and J gene segments being from the genome of one individual of the first human population.

In one embodiment of the library of the vertebrate, cell or library of the invention, all of the D and J segments that have been recombined with the first and second V gene segments are D and J segments derived from the first human population, optionally the D and J gene segments being from the genome of one individual of the first human population.

In one embodiment of the library, the second human immunoglobulin gene segment is a polymorphic variant of the first human immunoglobulin gene segment; optionally wherein the second gene segment is selected from the group consisting of a gene segment in any of Tables 1 to 7 and 9 to 14 (eg, selected from Table 13 or 14).

In one embodiment of the library, the first and second human immunoglobulin gene segments are both (i) V_H gene segments; (ii) D segments; (iii) J segments (optionally both J_H segments, both J_K segments or both J_λ segments); (iv) constant regions (optionally both a gamma constant region, optionally both a C gamma-1 constant region); (v) CH1 regions; (vi) CH2 regions; or (vii) CH3 regions.

The library is, for example, a naive and optionally has a library size of from 10 or 10^2 to 10^9 cells. For example, from 10, 20, 30, 40, 50, 100 or 1000 to 10^8 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^7 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^6 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^5 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^4 cells.

The library has, for example, been selected against a predetermined antigen and optionally has a library size of from 10 or 10^2 to 10^9 cells. For example, from 10, 20, 30, 40, 50, 100 or 1000 to 10^8 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^7 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^6 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^5 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^4 cells.

In one embodiment of the library of the invention, said first and second cells are progeny of first and second ancestor non-human vertebrate cells respectively, wherein the first ancestor cell comprises a genome comprising said first human immunoglobulin gene segment; and the second ancestor cell comprises a genome comprising said second human immunoglobulin gene segment.

The invention further provides a library of antibody-producing transgenic cells whose genomes collectively encode a repertoire of antibodies, wherein the library comprises the first and second ancestor cells described above.

The invention further provides a library of hybridoma cells produced by fusion of the library of the invention (eg, a B-cell library) with fusion partner cells and optionally has a library size of from 10 or 10^2 to 10^9 cells. For example, from 10, 20, 30, 40, 50, 100 or 1000 to 10^8 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^7 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^6 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^5 ; or 10, 20, 30, 40, 50, 100 or 1000 to 10^4 cells. Production of hybridomas is well known to the skilled person. Examples of fusion partners are SP2/0-g14 (obtainable from ECACC), P3X63-Ag8.653 (obtainable from LGC Standards; CRL-1580), NS1 and NS0 cells. PEG fusion or electrofusion can be carried out, as is conventional.

The invention provides, in a third configuration:-

An isolated antibody having

- (a) a heavy chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unrearranged human immunoglobulin V gene segment with a human D and human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments (eg, VH) is derived from the genome of an individual from a first human ethnic population; and the other two gene segments (eg, D and JH) are derived from the genome of an individual from a second (eg, a second and third respectively), different, human ethnic population,

and wherein the antibody comprises heavy chain constant regions (eg, C gamma) of said non-human vertebrate (eg, rodent, mouse or rat heavy chain constant regions); and/or

- (b) a light chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unrearranged human immunoglobulin V gene segment with a human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments (eg, VL) is derived from the genome of an individual from a first human ethnic population (optionally the same as the first population in (a)); and the other gene segment (eg, JL) is derived from the genome of an individual from a second, different, human ethnic population (optionally the same as the second population in (a)), and wherein the antibody comprises light chain constant regions of said non-human vertebrate (eg, rodent, mouse or rat heavy light constant regions);
- (c) Optionally wherein each variable domain of the antibody is a human variable domain.
- (d) Optionally wherein the heavy chain constant regions are mu- or gamma-type constant regions.

The invention also provides an isolated nucleotide sequence encoding the antibody of the third configuration, optionally wherein the sequence is provided in an antibody expression vector, optionally in a host cell. Suitable vectors are mammalian expression vectors (eg, CHO cell vectors or HEK293 cell vectors), yeast vectors (eg, a vector for expression in *Picchia pastoris*, or a bacterial expression vector, eg, a vector for *E. coli* expression).

The invention also provides a method of producing a human antibody, the method comprising replacing the non-human vertebrate constant regions of the antibody of the third configuration with human antibody constant regions (eg, a C variant disclosed in table 13 or 18). The skilled person will be aware of standard molecular biology techniques to do this. For example, see Harlow, E. & Lane, D. 1998, 5th edition, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Lab. Press, Plainview, NY; and Pasqualini and Arap, *Proceedings of the National Academy of Sciences* (2004) 101:257-259 for standard immunisation. Joining of the variable regions of an antibody to a human constant region can be effected by techniques readily available in the art, such as using conventional recombinant DNA and RNA technology as will be apparent to the skilled person. See e.g. Sambrook, J and Russell, D. (2001, 3^d edition) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY).

In one embodiment, the method comprises further making a mutant or derivative of the antibody.

The invention also provides a pharmaceutical composition comprising an antibody according to the third configuration, or a human antibody of the invention and a diluent, excipient or carrier; optionally wherein the composition is provided in a container connected to an IV needle or syringe or in an IV bag.

The invention also provides an antibody-producing cell (eg, a mammalian cell, eg, CHO or HEK293; a yeast cell, eg, *P pastoris*; a bacterial cell, eg, *E coli*; a B-cell; or a hybridoma) that expresses the second antibody of the third configuration or the isolated antibody of the invention.

The first configuration of the invention also provides:-

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus (eg, a heavy chain locus or a light chain locus), said locus comprising immunoglobulin gene segments according to the first and second human immunoglobulin gene segments (optionally V segments) described above in connection with the third configuration. The gene segments are operably connected upstream of an immunoglobulin constant region; optionally wherein the genome is homozygous for said transgenic immunoglobulin locus.

Optionally the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

Optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

Optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

In this way, a superhuman immunoglobulin gene repertoire is provided in a transgenic non-human vertebrate or vertebrate cell according to the invention.

The first configuration also provides:-

A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region; wherein the transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, a first (optionally a V segment) of said gene segments and a second (optionally a V segment) of said gene segments being different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

In this way, a superhuman immunoglobulin gene repertoire is provided in a transgenic non-human vertebrate or vertebrate cell according to the invention.

The first configuration also provides:-

A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises first and second transgenic immunoglobulin loci, each locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region;

wherein (i) the first transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, (ii) the second transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments; and (iii) wherein a first (optionally a V) gene segment of said first locus and a second (optionally a V) gene segment of said second gene locus are different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

optionally wherein the first and second loci are on different chromosomes (optionally chromosomes with the same chromosome number) in said genome;

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

In this way, a superhuman immunoglobulin gene repertoire is provided in a transgenic non-human vertebrate or vertebrate cell according to the invention.

In these embodiments of the first configuration, the immunoglobulin gene segments are optionally as described for the third configuration.

In these embodiments of the first configuration, the genome optionally comprises a third immunoglobulin gene segment (optionally a V segment), the third gene segment being derived from a human individual that is different from the individual from which the first (and optionally also the second) gene segment is derived; optionally wherein the first, second and third gene segments are polymorphic variants of a human immunoglobulin gene segment (eg, VH1-69 – see the examples for

further description).

In these embodiments of the first configuration, the genome of the vertebrate or cell is optionally homozygous for the first, second and optional third gene segment, wherein a copy of the first, second and optional third gene segments are provided together on the same chromosome operably connected upstream of a common non-human vertebrate constant region.

For example, each first, second and optional third gene segment is a V gene segment.

In one example, the library of the invention is provided by a collection of non-human vertebrates (optionally a collection of rodents, mice or rats); optionally, wherein a first member of said collection produces said first antibody but not said second antibody, and a second member of the collection produces said second antibody (but optionally not said first antibody). It is therefore contemplated to make non-human vertebrates where different human genomes have been used as a source for building the transgenic loci in the vertebrates. For example, a first vertebrate comprises a transgenic heavy chain locus having gene segments only from a first (and optionally a second) human population or individual; a second vertebrate comprises a transgenic heavy chain locus having gene segments only from a third (and optionally a fourth) human population or individual; and optionally third and more vertebrates can be built similarly based on unique or overlapping human population genomes. However, when provided as a mixed population of transgenic vertebrates, the mixed population provides a collective pool of human immunoglobulin genes that is greater than found in a natural human repertoire. This is useful to extend the antibody and gene sequence space beyond those possible with prior transgenic mice and rats bearing human immunoglobulin loci. As explained above, these have been based on a single human genome.

In one embodiment, the collection of non-human vertebrates bear human immunoglobulin genes confined to human populations that are together grouped under the same population genus "(a)" mentioned above. This provides for a gene repertoire that is biased to producing human antibody variable regions prevalent in the population genus (a) and thus useful for generating antibody therapeutics/prophylactics for members of said population. Alternatively, where gene segments from different human populations are provided in a single transgene according to the invention (not necessarily in a collection of vertebrates), the different human populations are for example together

grouped under the same population genus “(a)” mentioned above.

The invention also provides a repertoire of antibodies expressed from a library of cells according to the invention.

In the non-human vertebrate or cell of any configuration of the invention, the constant region of the transgenic locus is, in one example, an endogenous constant region of said vertebrate (eg, endogenous mouse or rat constant region, eg, from the same strain of mouse or rat as the non-human vertebrate itself).

The invention also provides a method of constructing a cell (eg, an ES cell) according to the invention, the method comprising

- (a) identifying functional V and J (and optionally D) gene segments of the genome sequence of a (or said) first human individual;
- (b) identifying one or more functional V and/or D and/or J gene segments of the genome sequence of a (or said) second human individual, wherein these additional gene segments are not found in the genome sequence of the first individual;
- (c) and constructing a transgenic immunoglobulin locus in the cell, wherein the gene segments of (a) and (b) are provided in the locus operably connected upstream of a constant region.

Optionally the cell comprises a heavy chain locus constructed according to steps (a) to (c) and/or a light chain locus (kappa and/or lambda loci) constructed according to steps (a) to (c).

Optionally the cell is homozygous for the or each transgenic locus; optionally wherein antibody expression from loci endogenous to said cell has been inactivated. This is useful for confining the functional antibody gene repertoire, and thus antibody production, to antibodies bearing human variable regions.

Optionally the gene segment(s) in step (b) are identified from an immunoglobulin gene database selected from the 1000 Genomes, Ensembl, Genbank and IMGT databases.

Optionally the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different, optionally wherein the human immunoglobulin gene segment derived from the genome sequence of the second individual is low-frequency (optionally rare) within the second ethnic population.

The invention also provides a method of making a transgenic non-human vertebrate (eg, a mouse or rat), the method comprising

- (a) constructing an ES cell (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain ES cell) by carrying out the method above;
- (b) injecting the ES cell into a donor non-human vertebrate blastocyst (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain blastocyst);
- (c) implanting the blastocyst into a foster non-human vertebrate mother (eg, a C57BL/6N, C57BL/6J, 129S5 or 129Sv strain mouse); and
- (d) obtaining a child from said mother, wherein the child genome comprises a transgenic immunoglobulin locus.

The invention provides a transgenic non-human vertebrate (eg, a mouse or rat) made by the method or a progeny thereof. The invention also provides a population of such non-human vertebrates.

Microinjection of ES cells into blastocysts and generation of transgenic mice thereafter are conventional practices in the state of the art, and the skilled person is aware of techniques useful to effect this. C57BL/6N, C57BL/6J, 129S5 or 129Sv mouse strains and ES cells are readily and publicly available.

The invention also provides a method of isolating an antibody that binds a predetermined antigen (eg, a bacterial or viral pathogen antigen), the method comprising

- (a) providing a vertebrate (optionally a mouse or rat) according to the invention;
- (b) immunising (eg, using a standard prime-boost method) said vertebrate with said antigen (optionally wherein the antigen is an antigen of an infectious disease pathogen);
- (c) removing B lymphocytes from the vertebrate and selecting one or more B lymphocytes expressing antibodies that bind to the antigen;
- (d) optionally immortalising said selected B lymphocytes or progeny thereof, optionally by

producing hybridomas therefrom; and

(e) isolating an antibody (eg, and IgG-type antibody) expressed by the B lymphocytes; and

(f) optionally producing a derivative or variant of the antibody.

This method optionally further comprises after step (e) the step of isolating from said B lymphocytes nucleic acid encoding said antibody that binds said antigen; optionally exchanging the heavy chain constant region nucleotide sequence of the antibody with a nucleotide sequence encoding a human or humanised heavy chain constant region and optionally affinity maturing the variable region of said antibody; and optionally inserting said nucleic acid into an expression vector and optionally a host.

Bioinformatics Analysis & Selection of Immunoglobulin Gene Segments

See also the discussion on variation analysis above.

The skilled person will know of sources of human antibody gene sequences, such as IMGT (www.imgt.org), GenBank (www.ncbi.nlm.nih.gov/genbank) Bioinformatics tools for database manipulation are also readily available and known to the skilled person, eg, as publicly available from the 1000 Genomes Project/EBI (www.1000genomes.org)

As a source of antibody gene segment sequences, the skilled person will also be aware of the following available databases and resources (including updates thereof):-

1.1. The Kabat Database (G. Johnson and T. T. Wu, 2002; <http://www.kabatdatabase.com>).

Created by E. A. Kabat and T. T. Wu in 1966, the Kabat database publishes aligned sequences of antibodies, T-cell receptors, major histocompatibility complex (MHC) class I and II molecules, and other proteins of immunological interest. A searchable interface is provided by the SeqhuntII tool, and a range of utilities is available for sequence alignment, sequence subgroup classification, and the generation of variability plots. See also Kabat, E. A., Wu, T. T., Perry, H., Gottesman, K., and Foeller, C. (1991) *Sequences of Proteins of Immunological Interest*, 5th ed., NIH Publication No. 91-

3242, Bethesda, MD, which is incorporated herein by reference, in particular with reference to human gene segments for use in the present invention.

1.2. KabatMan (A. C. R. Martin, 2002; <http://www.bioinf.org.uk/abs/simkab.html>). This is a web interface to make simple queries to the Kabat sequence database.

1.3. *IMGT, the International ImMunoGeneTics Information System*[®]; M.-P. Lefranc, 2002; <http://imgt.cines.fr>). IMGT is an integrated information system that specializes in antibodies, T cell receptors, and MHC molecules of all vertebrate species. It provides a common portal to standardized data that include nucleotide and protein sequences, oligonucleotide primers, gene maps, genetic polymorphisms, specificities, and two-dimensional (2D) and three-dimensional (3D) structures. IMGT includes three sequence databases (*IMGT/LIGM-DB*, *IMGT/MHC-DB*, *IMGT/PRIMERDB*), one genome database (*IMGT/GENE-DB*), one 3D structure database (*IMGT/3Dstructure-DB*), and a range of web resources (“*IMGT Marie-Paule page*”) and interactive tools.

1.4. *V-BASE* (I. M. Tomlinson, 2002; <http://www.mrc-cpe.cam.ac.uk/vbase>). V-BASE is a comprehensive directory of all human antibody germline variable region sequences compiled from more than one thousand published sequences. It includes a version of the alignment software DNAPLOT (developed by Hans-Helmar Althaus and Werner Müller) that allows the assignment of rearranged antibody V genes to their closest germline gene segments.

1.5. *Antibodies—Structure and Sequence* (A. C. R. Martin, 2002; <http://www.bioinf.org.uk/abs>). This page summarizes useful information on antibody structure and sequence. It provides a query interface to the Kabat antibody sequence data, general information on antibodies, crystal structures, and links to other antibody-related information. It also distributes an automated summary of all antibody structures deposited in the Protein Databank (PDB). Of particular interest is a thorough description and comparison of the various numbering schemes for antibody variable regions.

1.6. *AAAAA—AHO's Amazing Atlas of Antibody Anatomy* (A. Honegger, 2001; <http://www.unizh.ch/~antibody>). This resource includes tools for structural analysis, modeling, and engineering. It adopts a unifying scheme for comprehensive structural alignment of antibody and T-

cell-receptor sequences, and includes Excel macros for antibody analysis and graphical representation.

1.7. *WAM—Web Antibody Modeling* (N. Whitelegg and A. R. Rees, 2001; <http://antibody.bath.ac.uk>). Hosted by the Centre for Protein Analysis and Design at the University of Bath, United Kingdom. Based on the AbM package (formerly marketed by Oxford Molecular) to construct 3D models of antibody Fv sequences using a combination of established theoretical methods, this site also includes the latest antibody structural information.

1.8. *Mike's Immunoglobulin Structure/Function Page* (M. R. Clark, 2001; <http://www.path.cam.ac.uk/~mrc7/mikeimages.html>) These pages provide educational materials on immunoglobulin structure and function, and are illustrated by many colour images, models, and animations. Additional information is available on antibody humanization and Mike Clark's Therapeutic Antibody Human Homology Project, which aims to correlate clinical efficacy and anti-immunoglobulin responses with variable region sequences of therapeutic antibodies.

1.9. *The Antibody Resource Page* (*The Antibody Resource Page*, 2000; <http://www.antibodyresource.com>). This site describes itself as the “complete guide to antibody research and suppliers.” Links to amino acid sequencing tools, nucleotide antibody sequencing tools, and hybridoma/cell-culture databases are provided.

1.9. *Humanization bY Design* (J. Saldanha, 2000; <http://people.cryst.bbk.ac.uk/~ubcq07s>). This resource provides an overview on antibody humanization technology. The most useful feature is a searchable database (by sequence and text) of more than 40 published humanized antibodies including information on design issues, framework choice, framework back-mutations, and binding affinity of the humanized constructs.

See also Antibody Engineering Methods and Protocols, Ed. Benny K C Lo, Methods in Molecular BiologyTM, Human Press. Also at <http://www.blogsua.com/pdf/antibody-engineering-methods-and-protocolsantibody-engineering-methods-and-protocols.pdf>

As a source of genomic sequence variation data, the skilled person will also be aware of the following available databases and resources (including updates thereof):-

1. HapMap (The International HapMap Consortium. 2003; <http://hapmap.ncbi.nlm.nih.gov/index.html.en>). The HapMap Project is an international project that aims to compare the genetic sequences of different individuals to identify chromosomal regions containing shared genetic variants. The HapMap www site provides tools to identify chromosomal regions and the variant therein, with options to drill down to population level frequency data.
2. 1000 Genomes ([The 1000 Genomes Project Consortium](http://www.1000genomes.org/) 2010; <http://www.1000genomes.org/>). This resource provides complete genomic sequence for 2500 unidentified individuals from one of 25 distinct population groups, with the aim of identifying genomic variants of >1%. The site provides the ability to interrogate data utilizing online tools (e.g. 'Variation Pattern Finder') and to download variant data for individual population groups.
3. Japanese SNP Database (H.Haga et al. 2002; <http://snp.ims.u-tokyo.ac.jp/index.html>). Based on a study identifying 190,562 human genetic variants this site catalogues genomic variants with useful features for searching and summarizing data.

It is possible to identify variants in immunoglobulin genes classed as low-frequency or rare variants that segregate with specific human ethnic populations. For the purpose of this analysis, a low-frequency immunoglobulin gene segment is classed as one with 'Minor Allele Frequency' (MAF) (cumulative frequency) of between 0.5% - 5%, rare variants are those classed as having a MAF of less than 0.5% in a particular human population.

The following bioinformatics protocol is envisaged to identify human immunoglobulin gene segments for use in the present invention:

- (a) Identify one or more genomic regions containing gene segments of interest ('target genomic regions') and calculate the genomic coordinates, using coordinates that match the sequence assembly build used by either the 1000 Genomes project or International HapMap project (or another selected human gene database of choice).
- (b) Identify genomic variants mapped to the genomic regions previously identified in (a). Retrieve variant frequencies for variants for each super population and preferably sub-population where such data is available. Tools readily available on the HapMap WWW site and the VWC tools for the 1000Genomes Project are useful for this step.
- (c) Filter list of genomic variants from target genomic regions to contain only variants classed as either 'Non-synonymous' single nucleotide polymorphisms (SNPs) or genomic 'insertions or deletions' (indels). Filter further to include those that are present in exonic sequences only.
- (d) Correlate population frequency data for each of the identified variants for each of the super populations (for example 'European Ancestry', 'East Asian ancestry', 'West African ancestry', 'Americas', and 'South Asian ancestry') to identify those variants that segregate with less than two super-populations. Further correlate all identified variants with each of the sub-populations (for example, 'European ancestry' super-population might be subdivided into groups such as 'CEU – Utah residents with Northern or Western European ancestry', 'TSI Toscani in Italia' and 'British from England and Scotland') and produce a second score for rarity of variants in within a super-population.
- (e) Collect one or more gene segments that show segregation to specific sub-populations for construction of synthetic loci according to the invention.

In one embodiment throughout the present text, “germline” refers to the canonical germline gene segment sequence.

By detailed analysis of the 1000 Genomes database, the inventors have devised a collection of candidate polymorphic antibody gene segment variants, eg, human variant JH gene segments (eg, see Example 4), that can be built into the design of transgenic heavy chain loci in mice for expressing increasingly diverse and new, synthetic repertoires of human variable regions. To this end, the invention provides the following embodiments.

The present invention provides in a fourth configuration:-

Selection of Human JH6*02 Variant

Transgenic IgH Loci, Non-Human Vertebrates, Cells & Antibodies Based on Human JH6*02

As explained above, in designing transgenic Ig heavy chain loci the present inventors have considered the huge amount of data available from the 1000 Genomes project (see www.1000genomes.org) that analyses gene distributions amongst many human populations, and in particular data on Ig gene segments. The inventors were also aware of human gene segments disclosed in the IMGT database (see www.imgt.org) and in Ensembl (see www.ensembl.org). The inventors needed to make choices about which human gene segments to include amongst the large number of human gene segments presented in these databases and the other sources of human Ig gene segment information known in the art, including those other databases disclosed herein. When choosing human JH gene segments, the inventors were aware that human JH6 encodes a relatively long amino acid sequence, and thus the inventors thought it desirable to include this for increasing the chances of producing IgH chains with relatively long HCDR3 regions. Antibodies with long HCDR3 (at least 20 amino acids according to IMGT nomenclature) have been shown to neutralise a variety of pathogens effectively including HIV, Influenza virus, malaria and Africa trypanosomes. Reference is also made to naturally-occurring Camelid (eg, llama or camel) heavy chain-only antibodies which bear long HCDR3s for reaching relatively inaccessible epitopes (see, eg, EP0937140). Long HCDR3s can form unique stable subdomains with extended loop structure that towers above the antibody

surface to confer fine specificity. In some cases, the long HCDR3 itself is sufficient for epitope binding and neutralization (Liu, L *et al*; Journal of Virology. 2011. 85: 8467-8476, incorporated herein by reference). The unique structure of the long HCDR3 allows it to bind to cognate epitopes within inaccessible structure or extensive glycosylation on a pathogen surface. In human peripheral blood, there is around 3.5% of naïve B antibodies or 1.9% of memory B IgG antibodies containing the HCDR3s with lengths of more than 24 amino acids (PLoS One. 2012;7(5):e36750. Epub 2012 May 9; “Human peripheral blood antibodies with long HCDR3s are established primarily at original recombination using a limited subset of germline genes”; Briney BS *et al*, incorporated herein by reference) (Fig. 1). The usage analysis indicates that these antibodies have the preference to use human JH6 with human D2-2, D3-3 or D2-15 (Brinley, BS *et al*, Figs. 2-5). See also PLoS One. 2011 Mar 30;6(3):e16857; Comparison of antibody repertoires produced by HIV-1 infection, other chronic and acute infections, and systemic autoimmune disease”; Breden F *et al*, incorporated herein by reference. Around 20% of all HCDR3 of antibodies use JH6. However, in those antibodies with HCDR3 of more than 24 amino acids, 70% use JH6 (Brinley, BS *et al*, Fig.2).

There is a need in the art for genetically modified non-human vertebrates and cells that can make antibodies and heavy chains that have long human HCDR3s, as well as antibodies, chains and VH domains that can be selected from such vertebrates and cells wherein these can address target epitopes better accessed by long HCDR3s.

The inventors, therefore, chose in this configuration of the invention to include a human JH6 gene segment as a mandatory human gene segment in their IgH locus design. Several different naturally-occurring human JH6 variants are known (eg, JH6*01 to *04 as well as others; IMGT nomenclature). The inventors considered this when deciding upon which human JH6 variant should be included in the transgenic IgH locus design. An alignment of some human JH6 variants is shown in Figure 7 (from www.imgt.org; dashes indicate identical nucleotides; nucleotide changes versus the *01 variant are shown by underlined nucleotides and corresponding amino acid changes are shown by underlined amino acids; Genbank accession numbers (release 185.0) are shown prefixed by J, X, M or A). The inventors used sequencing of human genomic DNA samples, inspection of public IgH DNA databases as well as informed choices on the basis of variant sequences as means to arrive at a rational choice of which JH6 variant to use.

The 1000 Genomes database uses human JH6*03 as the reference sequence, which would be a possible choice for the skilled person wishing to construct a transgenic IgH locus. The inventors noticed (eg, Figure 7 herein) that position 6 in JH6*03 is a tyrosine (Y) encoded by a TAC codon, whereas some other naturally-occurring human variants have a glycine (G) encoded by a GGT codon (the glycine being present as a YYG motif, forming part of a larger YYGXDX motif). To understand the potential significance of this, the inventors carried out analysis of JH sequences from other vertebrate species. The inventors surprisingly noticed that YYG and YYGXDX motifs are conserved across many vertebrate species (see Figures 7 & 8). This suggested to the inventors, therefore, that preservation of this motif might be desirable, which could guide the choice of JH6 variant for use in the present invention.

Another pointer arose when the inventors considered the TAC codon versus the GGT codon encoding Y or G respectively. The inventors considered the impact of these nucleotide sequences on the action of activation-induced cytidine deaminase (AID). The inventors knew that activation-induced cytidine deaminase (AID) is believed to initiate Ig somatic hypermutation (SHM) in a multi-step mechanism and they addressed this activity when rationally designing the locus. AID catalyses the deamination of C to U in DNA, generating mutations at C bases. Cytidines located within hotspot motifs are preferentially deaminated. Certain motifs are hotspots for AID activity (DGYW, WRC, WRCY, WRCH, RGYW, AGY, TAC, WGCW, wherein W=A or T, Y=C or T, D=A, G or T, H=A or C or T, and R=A or G). The presence of a TAC codon encoding Y at position 6 in JH6*03 creates AID mutation hotspots (the cytidine being the substrate of AID), these hotspots being the underlined motifs in the previous sentence. The inventors considered the impact of this and in doing so they considered possible mutants created by AID activity at the cytidine. Reference is made to Figure 9. The inventors noticed that a mutation at the third base of the TAC codon would yield 3 possible outcomes: Y, stop or stop. Thus, out of the three stop codons possible in the genetic code (the other being encoded by TGA – see Figure 9), two of them would be provided by mutation of the cytidine in the TAC codon encoding position 6 in JH6*03. The inventors, therefore, considered that this might increase the chances of non-productive IgH variable region production in transgenic loci based on JH6*03. Moreover, the inventors noticed that provision of a GGT codon instead (as per the other human JH6 variants) seemed preferable since mutation of the third base would never yield a stop codon (see Figure 9), and furthermore would retain coding, and thus conservation, of glycine at position 6, which the inventors also noticed was in the YYG and YYGXDX motifs conserved across species.

Having decided against using JH6*03, the inventors needed to make a choice from other possible human variants. The MDV motif is at the C-terminus of HCDR3 based on human JH6, the adjacent framework 4 (FW4) starting with the WGQ motif (with reference to the sequence shown encoded by JH6*01; Figure 7). In making their choices for locus design, the inventors wished to maximise conservation of this HCDR3/FW4 junction in product IgH chains and antibodies including these. The inventors believed this to be desirable for heavy chain variable domain functionality and conformation. The inventors thought that this might in some cases be desirable to minimise immunogenicity (suitable for human pharmaceutical use). Consistent with these considerations, the inventors wanted to make a choice that would minimise mutation around the HCDR3/FW4 junction as a result of SHM *in vivo* to conserve junction configuration. See Rogozin & Diaz; "Cutting Edge: DGYW/WRCH Is a Better Predictor of Mutability at G:C Bases in Ig Hypermutation Than the Widely Accepted RGYW/WRCY Motif and Probably Reflects a Two-Step Activation-Induced Cytidine Deaminase-Triggered Process"; Journal of Immunology; March 15, 2004 vol. 172 no. 6 3382-3384. An example of a DGYW motif is GGCA. The inventors had this in mind when analysing the variant sequences.

With these considerations in mind, the inventors decided specifically to use human JH6*02 as the mandatory human JH6 for their IgH locus design. JH6*01 was rejected as the mandatory JH6 gene segment since the nucleotide sequence GGG CAA (encoding G and Q) contains a GGCA motif which is an AID recognition hotspot. The inventors realised that JH6*04 also contains such a motif due to the presence of the sequence GGC AAA encoding G and K (positions 11 and 12 respectively). The inventors also realised that the *02 variant has a C instead of a G that is in the *01 variant, the C desirably being a synonymous change (ie, not changing the encoded amino acid sequence around the CDR3/FW4 junction) and also this does not provide a GGCA AID hotspot motif. The inventors, therefore, decided that the mandatory JH6 should have this C base and this too pointed them to using the human JH6*02 variant.

In one example of any configuration of the invention herein, the only JH6 species included in the locus or genome is human JH6*02.

The inventors obtained 9 anonymised DNA samples from cheek swabs of 9 consenting human adults. Sequencing was performed on IgH locus DNA to confirm natural JH6 variant usage. It was found that the genome of all 9 humans contained a JH6*02 variant gene segment. In 7 out of the 9 humans, the genome was homozygous for JH6*02 (ie, each chromosome 14 had JH6*02 as its JH6 gene segment in the IgH locus). The inventors also inspected the publicly-available sequence information from the genomes of well-known scientists Craig Venter and Jim Watson. Both of these genomes contain JH6*02 too. This indicated to the inventors that this variant is common in humans.

So, the inventors made a choice of human JH6*02 on the basis of

- (i) Containing the YYG and YYGXDX motifs that is conserved across several vertebrate species;
- (ii) Provision of one less TAC codon (an AID hotspot that risks stop codons) and a choice instead of a codon that preserves the YYG and YYGXDX motifs;
- (iii) Avoidance of a GGCA AID hotspot in the region of the HCDR3/FW4 junction; and
- (iv) Common occurrence (and thus conservation and acceptability) in humans of the JH6*02 variant.

This rationale was tested by the inventors in laboratory examples, in order to see if human JH6*02 could desirably participate in antibody gene segment recombination and heavy chain production in a foreign (non-human vertebrate) setting, and moreover to assess if long HCDR3s based on human JH6*02 could be produced *in vivo* (in naïve and immunised settings) in such non-human systems. It was noted that in some non-human settings, such as a mouse, the YYG and YYGXDX motifs are not conserved, and thus the inventors decided that it was important to test whether or not JH6*02 (having the YYG and YYGXDX motifs) could function properly in such a foreign setting to participate in VDJ recombination and selection against antigen.

Thus, as explained further in the examples, the inventors constructed transgenic JH6*02-containing IgH loci in ES cells, generated transgenic non-human vertebrates from the ES cells (both naïve and immunised with a range of different target antigen types), isolated antibodies and heavy chain sequences based on JH6*02 as well as B-cells expressing these and made hybridomas expressing antigen-specific antibodies that are based on the chosen JH6*02 variant. The inventors found that the JH6*02 variant was extensively used and could contribute to the production of HCDR3 of at least

20 amino acids in many different heavy chains (including antigen-specific heavy chains). The chosen variant was preferably used over other JH gene segments in all settings (naïve, immunised and antigen-specific) for the production of HCDR3 of at least 20 amino acids.

Thus, the present invention provides an IgH locus including human JH6*02 (IMGT nomenclature) as a mandatory JH gene segment. In one embodiment, the locus comprises non-human vertebrate (eg, mouse or rat) constant region gene segments downstream (ie, 3' of) the human JH6*02; and one or more VH gene segments (eg, a plurality of human VH gene segments) and one or more D gene segments (eg, a plurality of human D gene segments) upstream of (ie, 5' of) the human JH6*02. For example, the locus is comprised by a vector (eg, a DNA vector, eg, a yeast artificial chromosome (YAC), BAC or PAC). Such a vector (eg, YAC) can be introduced into a non-human vertebrate (eg, mouse or rat) cell using standard techniques (eg, pronuclear injection) so that the locus is integrated into the cell genome for expression of IgH chains comprising at least one chain whose variable domain is a product of the recombination of human JH6*02 with a VH and a D gene segment.

In another example, the locus (eg, with a completely human, rat or mouse constant region, or a human/mouse chimaeric constant region) can be provided in the genome of a non-human vertebrate (eg, mouse or rat) cell. For example, the cell is an ES cell or an antibody-producing cell (eg, an isolated B-cell, an iPS cell or a hybridoma).

In another example, the invention provides a non-human vertebrate (eg, a mouse or a rat) comprising an IgH locus of the invention which comprises a human JH6*02 gene segment, wherein the locus can express an IgH chain whose variable domain is a product of the recombination of human JH6*02 with a VH and a D gene segment. As shown in the examples, the inventors have successfully produced such mice which produce such IgH chains with VH domains based on human JH6*02. The inventors isolated and sequenced IgH chains from the mice before (naïve) and after (immunised) exposure to a range of target antigens and confirmed by comparison to IMGT IgH gene segment sequences that the isolated chains (and antibodies containing these) were produced based on JH6*02. Such chains were found in naïve mice, as well as in antigen-specific antibodies from immunised mice. B-cells were isolated from immunised mice, wherein the B-cells express antibodies based on JH6*02 and hybridomas were generated from the B-cells, the hybridomas expressing antigen-specific antibodies based on JH6*02. The inventors, therefore, provided the locus,

vertebrate, cell and hybridoma of the invention based on the use of human JH6*02 and showed that antibodies based on JH6*02 and B-cells expressing these can be successfully produced and isolated following immunisation of the vertebrates, corresponding hybridomas being a good source of antibodies whose VH domains are based on JH6*02, eg for administration to a patient, eg, for human medicine. Furthermore, it was found possible to produce and isolated antigen-specific antibodies whose VH domains are based on JH6*02 and which had a relatively long HCDR3 (eg, 20 amino acids).

Thus, the present invention provides embodiments as in the following clauses:-

1. A non-human vertebrate (optionally a mouse or a rat) or vertebrate cell whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof so that the mouse is capable of producing an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.

In another example, the invention provides

A non-human vertebrate (optionally a mouse or a rat) or vertebrate cell whose genome comprises an immunoglobulin heavy chain locus comprising one, more or all of human IGHV gene segments selected from V3-21, V3-13, V3-7, V6-1, V1-8, V1-2, V7-4-1, V1-3, V1-18, V4-4, V3-9, V3-23, V3-11 and V3-20 (eg, one, more or all of V3-21*03, V3-13*01, V3-7*01, V6-1*01, V1-8*01, V1-2*02, V7-4-1*01, V1-3*01, V1-18*01, V4-4*01, V3-9*01 and V3-23*04). These segments were found in naive repertoires to be productive to produce HCDR3s of at least 20 amino acids in length. In an embodiment, the locus comprises a human JH6, eg, JH6*02.

The invention also provides a HCDR3, VH domain, antibody heavy chain or antibody having a HCDR3 size of at least 20 amino acids. Optionally, the HCDR3 or VH domain (or VH domain of the heavy chain or antibody) comprises mouse AID-pattern somatic hypermutations and/or mouse dTd-pattern mutations. This can be provided, for example, wherein VH domain is produced in a mouse comprising mouse AID and/or mouse TdT (eg, endogenous AID or TdT). See also Annu. Rev. Biochem. 2007. 76:1–22; Javier M. Di Noia and Michael S. Neuberger, “Molecular Mechanisms of Antibody Somatic Hypermutation” (in particular figure 1 and

associated discussion on AID hotspots in mouse); and Curr Opin Immunol. 1995 Apr;7(2):248-54, "Somatic hypermutation", Neuberger MS and Milstein C (in particular, discussion on hotspots in mouse), the disclosures of which are incorporated herein by reference.

These segments were found in naive repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the vertebrate is naïve. In another embodiment, the vertebrate instead is immunised with a target antigen.

In an example, the vertebrate or cell mentioned below is capable of so producing an antibody heavy chain upon immunisation with a target antigen. In an example, the vertebrate is an immunised vertebrate that produces antibody heavy chains specific for a target antigen and wherein the variable domains of the heavy chains are the product of recombination between a VH, D and JH6*02. For example, the D is selected from human D3-3, D2-15, D3-9; D4-17; D3-10; D2-2; D5-24; D6-19; D3-22; D6-13; D5-12; D1-26; D1-20; D5-18; D3-16; D2-21; D1-14; D7-27; D1-1; D6-25; D2-14 and D4-23 (eg, selected from D3-9*01; D4-17*01; D3-10*01; D2-2*02; D5-24*01; D6-19*01; D3-22*01; D6-13*01; D5-12*01; D1-26*01; D1-20*01; D5-18*01; D3-16*02; D2-21*02; D1-14*01; D7-27*02; D1-1*01; D6-25*01; D2-15*01; and D4-23*01). For example, the D is human D3-9 or D3-10. In an example, the HCDR3 length is at least 20 amino acids (eg, 20, 21, 23 or 24).

In an example of the vertebrate or cell, the genome comprises additional human JH gene segments (eg, JH2, 3, 4 and 5 gene segments).

In an example of the vertebrate or cell, the genome comprises an immunoglobulin light chain locus comprising one or more human V gene segments and one or more human J gene segments upstream of a constant region (eg, a human or a mouse lambda or kappa constant region).

For rearrangement and expression of heavy chains, the locus comprises control elements, such as an E μ and S μ between the J gene segment(s) and the constant region as is known by the skilled person. In one example, a mouse E μ and S μ is included in the heavy chain locus between the JH6*02 and the constant region (ie, in 5' to 3' order the locus comprises the JH6*02, E μ and S μ and constant region). In an example, the E μ and S μ are E μ and S μ of a mouse 129-derived

genome (eg, a 129Sv-derived genome, eg, 129Sv/EV (such as 129S7Sv/Ev (such as from AB2.1 or AB2.2 cells obtainable from Baylor College of Medicine, Texas, USA) or 129S6Sv/Ev)); in another example, the E μ and S μ are E μ and S μ of a mouse C57BL/6 -derived genome. In this respect, the locus can be constructed in the IgH locus of the genome of a cell selected from AB2.1, AB2.2, VGF1, CJ7 and FH14. VGF1 cells were established and described in Auerbach W, Dunmore JH, Fairchild-Huntress V, *et al*; Establishment and chimera analysis of 129/SvEv- and C57BL/6-derived mouse embryonic stem cell lines. Biotechniques 2000; 29:1024–8, 30, 32, incorporated herein by reference.

Additionally or alternatively, the constant region (or at least a C μ ; or C μ and gamma constant regions thereof) is a constant region (or C μ ; or C μ and gamma constant regions thereof) is of a genome described in the paragraph immediately above.

A suitable source of JH6*02 and other human DNA sequences will be readily apparent to the skilled person. For example, it is possible to collect a DNA sample from a consenting human donor (eg, a cheek swab sample as per the Example herein) from which can be obtained suitable DNA sequences for use in constructing a locus of the invention. Other sources of human DNA are commercially available, as will be known to the skilled person. Alternatively, the skilled person is able to construct gene segment sequence by referring to one or more databases of human Ig gene segment sequences disclosed herein.

2. The vertebrate of clause 1, wherein the vertebrate has been immunised with a target antigen and wherein the variable domain of the heavy chain is the product of recombination between a VH, D and JH6*02 and wherein the HCDR3 length is at least 20 amino acids (eg, 20, 21, 23 or 24).

Optionally, the immunised vertebrate produces an antibody heavy chain specific for a target antigen and wherein the variable domain of the heavy chain is the product of recombination between a VH, D and JH6*02 and wherein the HCDR3 length is at least 20 amino acids (eg, 20, 21, 23 or 24).

3. A non-human vertebrate cell (optionally a mouse cell or a rat cell) whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof for

producing (eg, in a subsequent progeny cell) an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.

4. The cell of clause 3, which is an ES cell capable of differentiation into a progeny antibody-producing cell that expresses said heavy chain.
5. The vertebrate or cell of any preceding clause, wherein the heavy chain locus comprises a human JH6*02 recombination signal sequence (RSS) operably connected 5' to the JH6*02 gene segment.

For example, the native RSS-JH6*02 sequence can be used to advantageously maintain the natural pairing between RSS and the JH gene segment. In this respect, the following sequence is used:-

ggttttgtggggtgaggatggacattctgccattgtgattactactactactacggtatggacgtctggggccaagggaccacgggtcaccg
tctcctcag (SEQ ID NO: 238)

RSSs have a common architecture: 9mer (eg, first underlined sequence above) followed by a 22bp spacer and then a 7mer (eg, second underlined sequence above). Spacers are 23bp +/- 1 normally, while the 9 and 7mer are more conserved.

6. The vertebrate or cell of clause 5, wherein the RSS is SEQ ID NO: 238 or a sequence having an identical 9mer and 7mer sequence flanking a sequence that is at least 70% identical to the 22mer sequence of SEQ ID NO: 238.
7. The vertebrate or cell of clause 6, wherein the RSS and JH6*02 are provided as SEQ ID NO: 237.
8. The vertebrate or cell of any preceding clause, wherein the JH6*02 is the only JH6-type gene segment in the genome.
9. The vertebrate or cell of any preceding clause, wherein the JH6*02 is the closest JH gene segment to the constant region in the locus.

10. The vertebrate or cell of any preceding clause, wherein the locus comprises one, more or all human D gene segments D3-9; D4-17; D3-10; D2-2; D5-24; D6-19; D3-22; D6-13; D5-12; D1-26; D1-20; D5-18; D3-16; D2-21; D1-14; D7-27; D1-1; D6-25; D2-14; and D4-23.

For example, the locus comprises one, more or all of human D gene segments D3-9*01; D4-17*01; D3-10*01; D2-2*02; D5-24*01; D6-19*01; D3-22*01; D6-13*01; D5-12*01; D1-26*01; D1-20*01; D5-18*01; D3-16*02; D2-21*02; D1-14*01; D7-27*02; D1-1*01; D6-25*01; D2-15*01; and D4-23*01.

11. The vertebrate or cell of clause 10, wherein the locus comprises one, more or all human D gene segments D3-9, D3-10, D6-19, D4-17, D6-13, D3-22, D2-2, D2-25 and D3-3.

These D segments were found to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the locus comprises one, more or all human D gene segments D3-9, D3-10, D6-19, D4-17, D6-13 and D3-22 (for example one, more or all of D3-9*01, D3-10*01, D6-19*01, D4-17*01, D6-13*01 and D3-22*01). These D segments were found in naïve repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the locus comprises one, more or all human D gene segments D3-10, D6-19 and D1-26 (for example, one, more or all of D3-10*01, D6-19*01 and D1-26*01). These D segments were found in immunised repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the locus comprises one, more or all human D gene segments D3-9 and D3-10 (for example, one, more or all of D3-9*01 and D3-10*01). These D segments were found in antigen-specific repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

12. The vertebrate or cell of any preceding clause, wherein the locus comprises a plurality of human D gene segments and the JH6*02 is in human germline configuration with respect to the 3'-most human D gene segment (or all of the human D segments comprised by the locus).

In an example, the 3'-most D gene segment is D7-27. In an example, the locus comprises all of human D gene segments from D1-1 to D7-27 as present in a germline human IgH locus (eg, as shown in the IMGT database).

Alternatively or additionally, the JH6*02 is in human germline configuration with respect to one, more or all of the E μ , S μ and constant region (eg, C μ).

13. The vertebrate or cell of any preceding clause, wherein the locus comprises one, more or all of IGHV gene segments selected from V3-21, V3-13, V3-7, V6-1, V1-8, V1-2, V7-4-1, V1-3, V1-18, V4-4, V3-9, V3-23, V3-11 and V3-20.

In an example, the locus comprises one, more or all human IGHV gene segments V3-21, V3-13, V3-7, V6-1, V1-8, V1-2, V7-4-1, V1-3, V1-18, V4-4, V3-9, V3-23 (for example, one, more or all of V3-21*03, V3-13*01, V3-7*01, V6-1*01, V1-8*01, V1-2*02, V7-4-1*01, V1-3*01, V1-18*01, V4-4*01, V3-9*01 and V3-23*04). These segments were found in naive repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the locus comprises one, more or all human IGHV gene segments V3-7, V3-11 and V4-4 (for example, one, more or all of V3-7*01, V3-11*01 and V4-4*02). These segments were found in immunised repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

In an example, the locus comprises one, more or all human IGHV gene segments V4-4, V1-8, V3-9, V3-11 and V3-20 (for example, one, more or all of V4-4*02, V1-8*01, V3-9*01, V3-11*01 and V3-20 (eg, *d01). These segments were found in antigen-specific repertoires to be productive in recombination with human JH6*02 to produce HCDR3s of at least 20 amino acids in length.

14. The vertebrate or cell of any preceding clause, wherein the locus comprises one, more or all of human D3-9*01, D3-10*01, D6-19*01, D6-13*01, D1-26*01, IGHV1-8*01, IGHV4-61*01, IGHV6-1*01, IGHV4-4*02, IGHV1-3*01, IGHV3-66*03, IGHV3-7*01 and IGHV3-9*01.

These are gene segments that very frequently combine with JH6*02 to produce productive heavy chains and antibodies.

For example, the locus comprises one, more or all of human IGHV1-8*01, D3-9*01 and D3-10*01. These gene segments were productive with JH6*02 to produce HCDR3s of at least 20 amino acids in more than 10 antibodies.

15. An antibody-producing cell (eg, a B-cell) that is a progeny of the cell of any one of clauses 3 to 14, wherein the antibody-producing cell comprises a heavy chain locus comprising a rearranged variable region produced by recombination of human JH6*02 with a D segment and a VH segment (eg, JH6*02 with human VH3-11 (eg, VH3-11*01) and D3-9; VH3-20 (eg, VH3-20*01) and D3-10; VH4-4 (eg, VH4-4*02) and D3-10; VH3-9 (eg, VH3-9*01) and D3-10; or VH1-8 (eg, VH1-8*01) and D310).

Such a variable region would be the product of *in vivo* somatic hypermutation in a non-human vertebrate or cell of the invention.

16. The cell of clause 15, which is a B-cell or hybridoma that expresses a target antigen-specific antibody comprising a heavy chain that comprises a rearranged variable region produced by recombination of human JH6*02 with a D segment and a VH segment (eg, JH6*02 with human VH3-11 (eg, VH3-11*01) and D3-9; VH3-20 (eg, VH3-20*01) and D3-10; VH4-4 (eg, VH4-4*02) and D3-10; VH3-9 (eg, VH3-9*01) and D3-10; or VH1-8 (eg, VH1-8*01) and D310).

Such a variable region would be the product of *in vivo* somatic hypermutation in a non-human vertebrate or cell of the invention

17. The vertebrate or cell of any preceding clause, wherein the antibody heavy chain specifically binds a target antigen.
18. The vertebrate or cell of any preceding clause, wherein the antibody heavy chain has a HCDR3 length of at least 20 amino acids.

Optionally, the HCDR3 length is at least 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 amino acids. Additionally, in one example the length is no more than 35, 34, 33, 32 or 31 amino acids. For example, the HCDR3 length is 20, 21, 22, 23 or 24 amino acids.

19. The vertebrate or cell of any preceding clause, wherein the antibody heavy chain is a product of the recombination of JH6*02 with a human VH gene segment recited in clause 13 or 14 and/or a D gene segment recited in clause 10, 11 or 14.
20. The vertebrate or cell of any preceding clause, wherein all endogenous non-human vertebrate heavy chain variable region gene segments have been inactivated in the genome (Eeg, by gene segment deletion or inversion).
21. The vertebrate or cell of any preceding clause, wherein the genome is homozygous for said heavy chain locus.
22. A heavy chain (eg, comprised by an antibody) isolated from a vertebrate of any one of clauses 1, 2, 5 to 14 and 17 to 21 wherein the heavy chain comprises a HCDR3 of at least 20 amino acids.
23. The heavy chain of clause 22, wherein the HCDR3 is the product of recombination of human JH6*02 with a human VH gene segment recited in clause 13 or 14 and/or a D gene segment recited in clause 10, 11 or 14.

In an example, the heavy chain is chimaeric where the C region is non-human. In an example, the heavy chain is human where the C region is human.

24. A heavy chain (eg, comprised by an antibody) whose VH variable domain is identical to the VH variable domain of the heavy chain of clause 22 or 23, and which comprises a human constant region or a human-mouse chimaeric constant region (eg, CH1 is human and the other constant domains are mouse).
25. The heavy chain of clause 22, 23 or 24, whose VH variable domain is specific for a target antigen.
26. A method for producing a heavy chain, VH domain or an antibody specific to a target antigen, the method comprising immunizing a non-human vertebrate according to any one of clauses 1, 2, 5 to 14 and 17 to 21 with the antigen and isolating the heavy chain, VH domain or an antibody specific to a target antigen or a cell producing the heavy chain, VH domain or an antibody, wherein the heavy chain, VH domain or an antibody comprises a HCDR3 that is derived from the recombination of human JH6*02 with a VH gene segment and a D gene segment.

27. A method for producing a human heavy chain or antibody comprising carrying out the method of clause 26, wherein the constant region of the locus is a non-human vertebrate (eg, mouse or rat) constant region, and then replacing the non-human constant region of the isolated heavy chain or antibody with a human constant region (eg, by engineering of the nucleic acid encoding the antibody).

28. A heavy chain, VH domain or an antibody produced by the method of clause 26 or 27.

Optionally the HCDR3 length is at least 20 amino acids as herein described.

29. A B-cell or hybridoma expressing a heavy chain VH domain that is identical to the VH domain of the heavy chain of clause 22, 23 or 28.

30. A nucleic acid encoding the VH domain of the heavy chain of clause 22, 23 or 28, or encoding the heavy chain of clause 22, 23, 24, 25 or 28.

31. A vector (eg, a CHO cell or HEK293 cell vector) comprising the nucleic acid of clause 30; optionally wherein the vector is in a host cell (eg, a CHO cell or HEK293 cell).

32. A pharmaceutical composition comprising the antibody, heavy chain or VH domain (eg, comprised by an antibody) of any one of clauses 22 to 25 and 28, together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, heavy chain or antibody).

33. The antibody, heavy chain or VH domain (eg, comprised by an antibody) of any one of clauses 22 to 25 and 28 for use in medicine (eg, human medicine).

For example, the locus comprises the following human VH gene segments

IGHV6-1
IGHV3-7
IGHV1-8
IGHV3-9
IGHV3-11
IGHV3-13
IGHV1-18
IGHV3-30
IGHV4-31

IGHV4-39
IGHV4-59

Optionally also (i) and/or (ii)

(i)
IGHV1-2
IGHV2-5 and
IGHV3-21

(ii)
IGHV1-2
IGHV2-5
IGHV3-21
IGHV1-24

For example, the locus comprises the following human VH gene segment variants

IGHV6-1*01
IGHV3-7*01
IGHV1-8*01
IGHV3-9*01
IGHV3-11*01
IGHV3-13*01
IGHV1-18*01
IGHV3-30*18
IGHV4-31*03
IGHV4-39*01 and
IGHV4-59*01;

Optionally also (iii) or (iv)

(ii)
IGHV1-2*04
IGHV2-5*10 and
IGHV3-21*03

(iv)

IGHV1-2*02
IGHV2-5*01
IGHV3-21*01 and
IGHV1-24*01

For example, the locus comprises the following human JH gene segment variants

IGHJ2*01
IGHJ3*02

IGHJ4*02
IGHJ5*02 and
IGHJ6*02

For example, the locus comprises the following human D gene segments

IGHD1-1
IGHD2-2
IGHD3-9
IGHD3-10
IGHD5-12
IGHD6-13
IGHD1-14
IGHD2-15
IGHD3-16
IGHD4-17
IGHD6-19
IGHD2-21
IGHD5-24
IGHD1-26 and
IGHD7-27

and optionally also (v) or (vi)

(v)
IGHD3-3

(vi)
IGHD3-3
IGHD4-4
IGHD5-5
IGHD6-6
IGHD1-7
IGHD2-8 and
IGHD2-8

The present invention provides in a fifth configuration:-

Constant Regions Tailored to Human Use & Antibody Humanisation

Additional rational design and bioinformatics has led the inventors to realise that specific human constant region variants are conserved across many diverse human populations. The inventors realised that this opens up the possibility of making a choice to humanise antibodies, chains and variable domains by using such specific constant regions in products, rather than arbitrarily choosing the human constant region (or a synthetic version of a human constant region). This aspect of the invention also enables one to tailor antibody-based drugs to specific human ethnic populations, thereby more closely matching drug to patient (and thus disease setting) than has hitherto been performed. It can be a problem in the state of the art that antibodies are humanised with an arbitrary choice of human constant region (presumably derived from one (often unknown) ethnic population or non-naturally occurring) that does not function as well in patients of a different human ethnic population. This is important, since the constant region has the major role in providing antibody effector functions, eg, for antibody recycling, cellular and complement recruitment and for cell killing.

As discussed further in WO2011066501, human IgG sub-types IgG1, IgG2, IgG3 and IgG4 exhibit differential capacity to recruit immune functions, such as antibody-dependent cellular cytotoxicity (ADCC, e.g., IgG1 and IgG3), antibody-dependent cellular phagocytosis (ADCP, e.g., IgG1, IgG2, IgG3 and IgG4), and complement dependent cytotoxicity (CDC, e.g., IgG1, IgG3). Sub-type-specific engagement of such immune functions is based on selectivity for Fc receptors on distinct immune cells and the ability to bind C1q and activate the assembly of a membrane attack complex (MAC). Among the various types, relative affinity for Fc γ receptors (e.g., Fc γ RI, Fc γ RIIa/b/c, Fc γ RIIIa/b) is high for IgG1 and IgG3, however, there is minimal affinity for IgG2 (restricted to the Fc γ RIIa 131H polymorphism), and IgG4 only has measurable affinity for Fc γ RI. Using comparative sequence analysis and co-crystal structures, the key contact residues for receptor binding have been mapped to the amino acid residues spanning the lower hinge and CH2 region. Using standard protein engineering techniques, some success in enhancing or reducing the affinity of an antibody preparation for Fc receptors and the C1q component of complement has been achieved.

Among the isotypes, IgG2 is least capable of binding the family of Fc receptors. Using IgG2 as the starting point, efforts have been made to find a mutant with diminished effector functions but which retains FcRn binding, prolonged stability, and low immunogenicity. Improved mutants of this nature may provide improved antibody therapeutics with retained safety. Human IgG1 therapeutic

antibodies that bind to cell surface targets are able to engage effector cells that may mediate cell lysis of the target cell by antibody-dependent cellular cytotoxicity (ADCC) or complement dependent cytotoxicity (CDC). These mechanisms occur through interaction of the CH2 region of the antibody Fc domain to FcγR receptors on immune effector cells or with C1q, the first component of the complement cascade. Table 19 shows the activities of different human gamma sub-types. The skilled person may choose accordingly to promote or dampen-down activity depending upon the disease setting in humans of interest. For example, use of a human gamma-1 constant region is desirable when one wishes to isolated totally human heavy chains and antibodies that have relatively high complement activation activity by the classical pathway and FcγR1 recognition in human patients. See also Mol Immunol. 2003 Dec;40(9):585-93; "Differential binding to human FcγRIIIa and FcγRIIIb receptors by human IgG wild type and mutant antibodies"; Armour KL *et al*, which is incorporated herein by reference.

IgG2 constant regions are well suited to producing antibodies and heavy chains according to the invention for binding to cytokines or soluble targets in humans, since IgG2 is essentially FcγRI,III-silent, FcγRIIIa-active and has little Complement activity.

IgG1 constant regions have wide utility for human therapeutics, since IgG1 antibodies and heavy chains are FcγRI,II,III- active and have complement activity. This can be enhanced by using a human gamma-1 constant region that has been activated by engineering as is known in the art.

The work of the inventors has therefore identified a collection of human constant region of different isotypes from which an informed choice can be made when humanising chimaeric antibody chains (or conjugating V domains, such as dAbs or *Camelid* VHH, to constant regions). The collection was identified on the basis of bioinformatics analysis of the 1000 Genomes database, the inventors selecting constant region variants that are frequently occurring across several human ethnic populations, as well as those that appear with relatively high frequency within individual populations (as assessed by the number of individuals whose genomes comprise the variant). By sorting through the myriad possible sequences on this basis, the inventors have provided a collection of human constant region variants that are naturally-occurring and which can be used when rationally designing

antibodies, heavy chains and other antibody-based formats that bear a human constant region. In particular, this is useful when humanising chimaeric heavy chains to produce totally human chains in which both the variable and constant regions are human. This is useful for compatibility with human patients receiving antibody-based drugs.

To this end, the invention provides the following aspects:-

1. A method of producing an antibody heavy chain, the method comprising
 - (a) providing an antigen-specific heavy chain variable domain (eg, VH (such as a human VH or dAb) or VHH or a humanised heavy chain variable domain); and
 - (b) combining the variable domain with a human heavy chain constant region to produce an antibody heavy chain comprising (in N- to C-terminal direction) the variable domain and the constant region;
 wherein
 the human heavy chain constant region is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region.

Step (b) can be carried out, eg, using recombinant DNA technology using the corresponding nucleotide sequences.

For the constant region according to any aspect of this configuration, either genomic DNA or equivalent (ie, having introns and exons and optionally also 5' UTR sequences, eg, with native or a non-native leader sequence) can be used for the constant region. For example, any of the "GENOMIC" sequences disclosed as SEQ ID NO: 365 onwards herein. Alternatively, an intronless sequence can be used, for example any of the "CDS" sequences disclosed as SEQ ID NO: 365 onwards herein (eg, with native or a non-native leader sequence).

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHAref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHA1a constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHA2a constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHA2b constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is IGHG1ref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG2ref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG2a constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG3ref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG3a constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG3b constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG4ref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHG4a constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHDref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHEref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHMref constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHMa constant region.

Optionally for any aspect of this configuration of the invention, the human heavy chain constant region is an IGHMb constant region.

Optionally, a derivative (eg, a mutant or conjugate) of the heavy chain or an antibody containing the heavy chain is produced. For example, a toxic payload can be conjugated (eg, for oncology applications). For example, one or more mutations can be introduced, as is known in the art, to inactivate or enhance Fc effector function.

2. The method of aspect 1, wherein the variable domain is a human variable domain.

A human variable domain is, for example, the product of recombination in a transgenic non-human vertebrate of human VH, D and JH gene segments. Alternatively, the variable domain is identified using *in vitro* display technology from a human VH library, eg, using phage display, ribosome display or yeast display, as is known in the art.

In another embodiment, the variable domain is a humanised variable domain, eg, comprising human frameworks with non-human (eg, mouse or rat) CDRs). Humanisation technology is conventional in the art, and will be readily known to the skilled person.

3. The method of any preceding aspect, wherein the variable domain has previously been selected from a non-human vertebrate that has been immunised with the antigen.

For example, the vertebrate (such as a mouse or rat) genome comprises a chimaeric heavy chain locus comprising a human variable region (human V, D and JH gene segments) operably connected upstream of a non-human vertebrate constant region so that the locus is able to rearrange for the expression of heavy chains comprising human variable domains and non-human vertebrate constant regions.

In alternative embodiments, the variable domain is selected using an *in vitro* technology such as phage display, ribosome display or yeast display. In this case the variable domain may be displayed with or without an constant region, provided that it is later combined with a human constant region as per the invention.

4. The method of any preceding aspect, comprising providing an expression vector (Eg, a mammalian expression vector, such as a CHO or HEK293 vector) comprising a nucleotide sequence encoding the constant region; inserting a nucleotide sequence encoding the variable domain into the vector 5' of the constant region sequence; inserting the vector into a host cell and expressing the heavy chain by the host cell; the method further comprising isolating a heavy chain (eg, as part of an antibody) comprising the variable domain and the human constant region.

The vector comprises regulatory elements sufficient to effect expression of the heavy chain when the vector is harboured by a host cell, eg, a CHO or HEK293 cell.

5. The method of any preceding aspect, further comprising obtaining a nucleotide sequence encoding the heavy chain.

6. An antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region.
7. A polypeptide comprising (in N- to C- terminal direction) a leader sequence, a human variable domain that is specific for an antigen and a human constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; wherein (i) the leader sequence is not the native human variable domain leader sequence (eg, the leader sequence is another human leader sequence or a non-human leader sequence); and/or (ii) the variable domain comprises mouse AID-pattern somatic mutations or mouse terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations.
8. A nucleotide sequence encoding (in 5' to 3' direction) a leader sequence and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; and the leader sequence being operable for expression (eg, in a mammalian CHO or HEK293 cell) of the heavy chain and wherein the leader sequence is not the native human variable domain leader sequence (eg, the leader sequence is another human leader sequence or a non-human leader sequence).

In an example, the leader sequence is

ATGGGCTGGTCCTGCATCATCCTGTTTCTGGTGGCCACCGCCACCGGCGTGACAGC

Which translates to

MGWSCILFLVATATGVHS

9. A nucleotide sequence encoding (in 5' to 3' direction) a promoter and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref,

IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; and the promoter being operable for expression (eg, in a mammalian CHO or HEK293 cell) of the heavy chain and wherein the promoter is not the native human promoter.

In one embodiment, the promoter sequence is a human IGK 3-15 promoter.

10. The antibody, polypeptide or nucleotide sequence of any one of aspects 6 to 9, wherein the variable domain comprises mouse AID-pattern somatic mutations and/or mouse terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations.

For example, one way, in any aspect of this configuration of the invention, to provide mouse AID-pattern somatic mutations and/or mouse terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations is to select a variable domain from a non-human vertebrate or cell. For example, a vertebrate or cell as disclosed herein.

11. A vector (eg, a CHO cell or HEK293 cell vector) comprising the nucleic acid of aspect 8, 9 or 10; optionally wherein the vector is in a host cell (eg, a CHO cell or HEK293 cell).
12. A pharmaceutical composition comprising the antibody or polypeptide of any one of aspects 6, 7 and 10, together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).
13. The antibody or polypeptide of any one of aspects 6, 7 and 10 for use in treating and/or preventing a medical condition in a human patient.
14. Use of the antibody or polypeptide of any one of aspects 6, 7 and 10 for the manufacture of a medicament for treating and/or preventing a medical condition in a human patient.
15. The antibody, polypeptide or use of aspect 13 or 14, wherein the human is a member of a human population selected from population numbers 1-14, wherein the populations are numbered as follows (population labels being according to 1000 Genomes Project nomenclature)

1= ASW;

2=CEU;
3=CHB;
4=CHS;
5=CLM;
6=FIN;
7=GBR;
8=IBS;
9=JPT;
10=LWK;
11=MXL;
12=PUR;
13=TSI;
14=YRI.

16. The antibody, polypeptide or use of aspect 15, wherein the constant region is a

- (i) IGHA1a constant region and the human population is selected from any population number 1-14;
- (ii) IGHA2a constant region and the human population is selected from any population number 1-14;
- (iii) IGHA2b constant region and the human population is selected from any population number 1-14;
- (iv) IGHG2a constant region and the human population is selected from any population number 1-9 and 11-13;
- (v) IGHG3a constant region and the human population is selected from any population number 1-14;
- (vi) IGHG3b constant region and the human population is selected from any population number 1-8 and 11-13;
- (vii) IGHG4a constant region and the human population is selected from any population number 1-9 and 11-13;
- (viii) IGHMa constant region and the human population is selected from any population number 1-14; or
- (ix) IGHMb constant region and the human population is selected from any population number 1-14;

Wherein the populations are numbered as follows (population labels being according to 1000 Genomes Project nomenclature)

1= ASW;

2= CEU;

3=CHB;

4=CHS;

5=CLM;

6=FIN;

7=GBR;

8=IBS;

9=JPT;

10=LWK;

11=MXL;

12=PUR;

13=TSI;

14=YRI.

17. A vector (eg, a CHO cell or HEK293 cell vector) comprising a IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region nucleotide sequence that is 3' of a cloning site for the insertion of a human antibody heavy chain variable domain nucleotide sequence, such that upon insertion of such a variable domain sequence the vector comprises (in 5' to 3' direction) a promoter, a leader sequence, the variable domain sequence and the constant region sequence so that the vector is capable of expressing a human antibody heavy chain when present in a host cell.

The present invention provides in a Sixth configuration:-

Multiple Variants in the Same Genome *Cis* or *Trans*

The inventors' analysis has revealed groupings of naturally-occurring human antibody gene segment variants as set out in Table 13 and Table 14. This revealed the possibility of producing transgenic genomes in non-human vertebrates and cells wherein the genomes contain more than the natural

human complement of specific human gene segments. In one example, this can be achieved by providing more than the natural human complement of a specific gene segment type on one or both of the respective Ig locus (eg, one or both chromosomes harbouring IgH in a mouse genome or mouse cell genome).

To this end, this configuration of the invention provides the following (as set out in numbered paragraphs):-

1. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.

For example, the genome comprises a variable region that comprises V, D and J gene segments (for the variable region of a heavy chain locus) or V and J gene segments (for the variable region of a light chain locus) upstream of a constant region for expression of heavy or light chains respectively.

In an alternative, the skilled person can choose to provide more than the wild type human complement of a specific gene segment type by providing several copies of one variant type of the human gene segment. Thus, there is provided

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein the human gene segments are identical variants.

For example, the genome comprises a variable region that comprises V, D and J gene segments (for the variable region of a heavy chain locus) or V and J gene segments (for the variable region of a light chain locus) upstream of a constant region for expression of heavy or light chains respectively.

2. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.

In an alternative, the skilled person can choose to provide more than the wild type human complement of a specific gene segment type by providing several copies of one variant type of the human gene segment. Thus, there is provided

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 non-endogenous variable region gene segments of the same variant type (eg, at least 2 human JH6*02 gene segments) *cis* at the same Ig locus.

3. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and optionally a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.

In an alternative, the skilled person can choose to provide more than the wild type human complement of a specific gene segment type by providing several copies of one variant type of the human gene segment. Thus, there is provided

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human variable region gene segments of the same variant type (eg, at least 2 human JH6*02 gene segments) *trans* at the same Ig locus; and optionally a third human gene segment of the same variant type, wherein the third gene segment is *cis* with one of said 2 different gene segments.

4. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human variable region gene segments, wherein the plurality comprises at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), a first of said different gene segments is provided in the genome of a first vertebrate of the population, and a second of said different gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second gene segment.
5. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
6. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.
7. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.

8. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and optionally a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.
9. A method of providing an enhanced human immunoglobulin variable region gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human variable region gene segments, wherein the method comprises providing at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein a first of said different gene segments is provided in the genome of a first vertebrate of the population, and a second of said different gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second gene segment.
10. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
11. The vertebrate, cell or method of any preceding paragraph, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
12. The vertebrate, cell or method of any preceding paragraph, wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically

related over at least 3 generations.

13. The vertebrate, cell or method of any preceding paragraph, wherein the gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.
14. The vertebrate, cell or method of paragraph 13, wherein the individuals are not genetically related.
15. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same Ig locus in said genome.
16. The method of paragraph 15, wherein the different human individuals are from different human populations.
17. The method of paragraph 15, wherein the individuals are not genetically related.
18. The vertebrate, cell or method of preceding paragraph, wherein at least one of the different segments is a synthetic mutant of a human germline gene segment.
19. The vertebrate, cell or method of any preceding paragraph, wherein each of said gene segments occurs in 10 or more different human populations.

20. The vertebrate, cell or method of preceding paragraph, wherein each of said gene segments has a human frequency of 5% or greater (eg, 10, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, or 95% or greater).

In this respect, the skilled person can be guided by the information provided in Table 14.

Frequency can, for example, be cumulative frequency in the 1000 Genomes database.

21. The vertebrate, cell or method of paragraph 20, wherein each of said gene segments occurs in 10 or more different human populations.
22. The vertebrate, cell or method of any preceding paragraph, wherein each of said gene segments occurs in the 1000 Genomes database in more than 50 individuals.
23. The vertebrate, cell or method of preceding paragraph, wherein each of said gene segments (i) has a human frequency of 5% or greater (eg, 10, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, or 95% or greater); and (ii) occurs in 10 or more different human populations.

In this respect, the skilled person can be guided by the information provided in Table 14.

Frequency can, for example, be cumulative frequency in the 1000 Genomes database.

24. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from Table 14 (eg, IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, IGHJ6 ref; SEQ ID NO: 244).

Table 14 lists commonly-occurring natural human variants. It can be seen that these occur across many human populations and thus usefully have wide applicability for human antibody-based drugs.

For example, the gene segments are provided as targeted insertions into an endogenous non-human vertebrate Ig locus. Alternatively, random integration (eg, using YACs) as is known in the art can be performed.

For example, the genome comprises a variable region that comprises V, D and J gene segments (for the variable region of a heavy chain locus) or V and J gene segments (for the variable region of a light chain locus) upstream of a constant region for expression of heavy or light chains respectively.

In another embodiment, the invention enables the skilled person to select two or more different naturally-occurring human gene segment variants for combination into the genome of a non-human vertebrate or cell. A reference sequence need not be included. It may be desirable to use one or more rare gene segments to increase diversity of the repertoire. Additionally or alternatively, it may be desirable to include a mixture of frequent and rare variants of the same type to provide repertoire diversity. The variants may be chosen additionally or alternatively to tailor the gene segment inclusion to one or more specific human populations as indicated by the information provided in Table 13 or Table 14.

Thus, the invention provides

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the gene segments are gene segments selected from Table 13 or Table 14; and optionally wherein one or more of the gene segments appears in Table 14 (eg,IGHJ6-a) or is a reference sequence (eg, IGHJ6 ref; SEQ ID NO: 244).

25. The vertebrate or cell of paragraph 24, wherein the genome comprises a third human gene segment of said type, the third gene segment being different from the first and second gene segments.

26. The vertebrate or cell of paragraph 24 or 25, wherein the first and second gene segments are *cis* on the same chromosome; and optionally the third gene segment is also *cis* on said chromosome.

27. The vertebrate or cell of paragraph 26, wherein the gene segments are targeted insertions into an endogenous non-human Ig locus.

For example, the gene segments are heavy chain gene segments and the non-human locus is an IgH locus. For example, the gene segments are light chain (kappa or lambda) gene segments and the non-human locus is an IgL locus.

28. The vertebrate or cell of paragraph 24 or 25, wherein the first and second gene segments are *trans* on different chromosomes.

Thus, the chromosomes are the same type (eg, both mouse chromosome 6 or rat chromosome 4).

29. The vertebrate or cell of any one of paragraphs 24 to 28, wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, selected from Table 13 or 14) and the second gene segment is the corresponding reference sequence.

30. A population of non-human vertebrates (eg, mice or rats) comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, Table 13 or 14) (eg,IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, SEQ ID NO: 7), wherein the first gene segment is provided in the genome of a first vertebrate of the population, and the second gene segment is provided in the genome of a second vertebrate of the population.

31. The population of paragraph 30, wherein the genome of the first vertebrate does not comprise the second gene segment.
32. The population of paragraph 30 or 31, wherein the population comprises a third human gene segment of said type, the third gene segment being different from the first and second gene segments and optionally wherein the first and third gene segments are present in the genome of the first vertebrate.
33. The population of paragraph 30, 31 or 32, wherein the gene segments are targeted insertions into an endogenous non-human Ig locus in the respective genome.

For example, the gene segments are heavy chain gene segments and the non-human locus is an IgH locus. For example, the gene segments are light chain (kappa or lambda) gene segments and the non-human locus is an IgL locus.

34. The population of of any one of paragraphs 30 to 33, wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, Table 13 or 14) and the second gene segment is the corresponding reference sequence.
35. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, Table 13 or 14) (eg,IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, SEQ ID NO: 7).
36. A method of providing an enhanced human immunoglobulin gene segment repertoire, the method comprising providing a population according to any one of paragraphs 30 to 33.

Variants Prevalent in Few Populations

In another aspect, it is of note that certain human gene segment variants may appear relatively frequently in one or a small number of populations, but is not found prevalently across many different human populations. There is thinking that specific germline gene segment repertoires have evolved in individual human ethnic populations due to iterative exposure to antigens (eg, disease pathogen antigens) to which the population is often exposed. Repeated exposure and mutation may have lead to the evolution of gene segment variants that can provide an effective response to the antigen (pathogen) in the population, and this may explain the conservation of the gene segments in those populations (as opposed to other human ethnic populations that may not have frequently encountered the antigen). With this in mind, the inventors identified gene segment variants from their analysis that are relatively prevalent in a small number of human populations, and not across many populations. The inventors realized that inclusion of one or more of such gene segments in the configurations of the invention (eg, in transgenic Ig loci, vertebrates and cells) would be useful for producing antibodies, Ig chains and variable domains that can address antigens (eg, disease-causing antigens or pathogens) to which the small number of human populations may become exposed. Such products would be useful for treating and/or preventing disease or medical conditions in members of such a population. This aspect could also be useful for addressing infectious disease pathogens that may have been common in the small number of populations, but which in the future or relatively recently in evolution has become a more prevalent disease-causing pathogen in other human populations (ie, those not listed in Table 13 against the gene segment variant(s) in question). To this end, from the 1000 Genomes database the inventors have identified the gene segment variants listed in Table 20.

Thus, according to any configuration or aspect described herein, one, more or all of the gene segments used in the present invention can be a gene segment listed in Table 20A, 20B, 20C or 20D.

Multiple JH Gene Segment Variants

A specific application of this configuration is the provision of multiple human JH gene segments as follows.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein at least two of the human JH gene segments are variants that are not

identical to each other.

In an example, any cell of the invention is an isolated cell. An “isolated” cell is one that has been identified, separated and/or recovered from a component of its production environment (eg, naturally or recombinantly). Preferably, the isolated cell is free of association with all other components from its production environment, eg, so that the cell can produce an antibody to an FDA-approvable or approved standard. Contaminant components of its production environment, such as that resulting from recombinant transfected cells, are materials that would typically interfere with research, diagnostic or therapeutic uses for the resultant antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified: (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments, to greater than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Ordinarily, however, an isolated cell will be prepared by at least one purification step.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous JH gene segments (eg, human gene segments) of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *cis* at the same Ig (eg, IgH, eg, endogenous IgH, eg, mouse or rat IgH) locus. In an example, the genome comprises a human VH, D and JH repertoire comprising said different JH gene segments. Optionally the non-endogenous JH gene segments are non-mouse or non-rat, eg, human JH gene segments. In an example one or more or all of the non-endogenous gene segments are synthetic.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *trans* at the same Ig (eg, IgH, eg, endogenous IgH, eg, mouse or rat IgH) locus; and optionally a third human JH gene segments of the same type, wherein the third JH is *cis* with one of said 2 different JH gene segments.

A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human JH gene segments, wherein the plurality comprises at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), a first of said different JH gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JH gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JH gene segment.

A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous (eg, human) JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations (eg, 3, 4, 5 or 6 generations). Optionally the non-endogenous JH gene segments are human JH gene segments. In an example one or more or all of the non-endogenous gene segments are synthetic.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein at least two of the human JH gene segments are variants that are not identical to each other.

A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous (eg, human) JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *cis* at the same Ig (eg, IgH, eg, endogenous IgH, eg, mouse or rat IgH) locus). Optionally the non-endogenous JH gene segments are non-mouse or non-rat, eg, human JH gene segments. In an example one or more or all of the non-endogenous gene segments are synthetic.

A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *trans* at the same Ig (eg, IgH, eg, endogenous IgH, eg, mouse or rat IgH) locus; and optionally a third human JH

gene segments of the same type, wherein the third JH is *cis* with one of said 2 different JH gene segments.

A method of providing an enhanced human immunoglobulin JH gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human JH gene segments, wherein the method comprises providing at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein a first of said different JH gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JH gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JH gene segment.

A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous (eg, human) JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations (eg, 3, 4, 5, or 6 generations). Optionally the non-endogenous JH gene segments are human JH gene segments. In an example one or more or all of the non-endogenous gene segments are synthetic.

In an example of the vertebrate or cell or the method of the invention at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.

In an example of the vertebrate or cell or the method of the invention the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations (eg, 3, 4, 5, or 6 generations).

In an example of the vertebrate or cell or the method of the invention the JH gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.

In an example of the vertebrate or cell or the method of the invention the individuals are not genetically related (eg, going back 3, 4, 5, or 6 generations).

In an example of the vertebrate or cell or the method of the invention at least one of the different JH segments is a synthetic mutant of a human germline JH gene segment.

The invention also provides a method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations (eg, 3, 4, 5, or 6 generations); optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgH locus in said genome.

In an example of the vertebrate or cell or the method of this embodiment of the invention the genome comprises a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

In an example of the population of the invention, the population comprises a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

In an example of the vertebrate or the population, at least one of said JH gene segments is SEQ ID NO: 1, 2, 3 or 4. For example, at least one of said JH gene segments is SEQ ID NO: 1 and at least one, two or more of said supplementary JH gene segments is a variant according to any example above. For example, at least one of said JH gene segments is SEQ ID NO: 2 and at least one, two or more of said supplementary JH gene segments is a variant according to any one of the examples above. For example, at least one of said JH gene segments is SEQ ID NO: 2 and at least one, two or more of said supplementary JH gene segments is a variant according to any one of the examples above.

In an embodiment, the non-human vertebrate or vertebrate cell of the invention comprises a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the JH gene repertoire (eg, a human JH gene segment repertoire) comprising

a plurality of JH1 gene segments provided by at least 2 different JH1 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH2 gene segments provided by at least 2 different JH2 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH3 gene segments provided by at least 2 different JH3 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH4 gene segments provided by at least 2 different JH4 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH5 gene segments provided by at least 2 different JH5 gene segments in *cis* at the same Ig locus in said genome; and/or

a plurality of JH6 gene segments provided by at least 2 different JH6 gene segments in *cis* at the same Ig locus in said genome;

optionally wherein the JH gene segments are derived from the genome sequence of two or more different human individuals.

Optionally said at least 2 different JH gene segments are human gene segments or synthetic gene segments derived from human gene segments.

Optionally, the Ig locus is a IgH locus, eg, an endogenous locus, eg, a mouse or rat IgH locus.

In an embodiment, the non-human vertebrate or vertebrate cell of the invention comprises a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the JH gene repertoire (eg, a human JH gene segment repertoire) comprising

- a plurality of JH1 gene segments provided by at least 3 different JH1 gene segments;
- a plurality of JH2 gene segments provided by at least 3 different JH2 gene segments;
- a plurality of JH3 gene segments provided by at least 3 different JH3 gene segments;
- a plurality of JH4 gene segments provided by at least 3 different JH4 gene segments;
- a plurality of JH5 gene segments provided by at least 3 different JH5 gene segments; and/or
- a plurality of JH6 gene segments provided by at least 3 different JH6 gene segments;

optionally wherein the JH gene segments are derived from the genome sequence of two or three different human individuals;

optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same Ig locus in said genome.

Optionally said at least 3 different JH gene segments are human gene segments or synthetic gene segments derived from human gene segments.

Optionally, the Ig locus is a IgH locus, eg, an endogenous locus, eg, a mouse or rat IgH locus.

Optionally in the vertebrate or cell the different human individuals are from different human populations.

Optionally in the vertebrate or cell the individuals are not genetically related (eg, Going back 3, 4, 5 or 6 generations).

Optionally in the vertebrate or cell at least one of the different JH segments is a synthetic mutant of a human germline JH gene segment.

In an embodiment of a non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to the invention, the vertebrate or cell genome comprises human VH, D and JH gene repertoires, the JH gene repertoire (eg, a human JH gene repertoire) comprising a plurality of JH1 gene segments provided by at least 2 different human JH1 gene segments, optionally in *cis* at the same Ig locus in said genome;
 a plurality of JH2 gene segments provided by at least 2 different human JH2 gene segments, optionally in *cis* at the same Ig locus in said genome;
 a plurality of JH3 gene segments provided by at least 2 different human JH3 gene segments, optionally in *cis* at the same Ig locus in said genome;
 a plurality of JH4 gene segments provided by at least 2 different human JH4 gene segments, optionally in *cis* at the same Ig locus in said genome;
 a plurality of JH5 gene segments provided by at least 2 different human JH5 gene segments, optionally in *cis* at the same Ig locus in said genome; and/or
 a plurality of JH6 gene segments provided by at least 2 different human JH6 gene segments, optionally in *cis* at the same Ig locus in said genome;
 wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations (eg, 3, 4, 5 or 6 generations).

Optionally said at least 2 different JH gene segments are human gene segments or synthetic gene segments derived from human gene segments.

Optionally, the Ig locus is a IgH locus, eg, an endogenous locus, eg, a mouse or rat IgH locus.

Optionally in the vertebrate or cell the human individuals are from different human populations.

JH5

An embodiment provides a vertebrate, cell or population of the invention whose genome comprises a plurality of JH5 gene segments, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a nucleotide mutation at one or more positions

corresponding to positions

106,330,024

106,330,027

106,330,032

106,330,041

106,330,044

106,330,045

106,330,062

106,330,063

106,330,065

106,330,066

106,330,067

106,330,068 and

106,330,071

on human chromosome 14.

In the vertebrate, cell or population optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,067 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,071 on human chromosome 14 (optionally the additional mutation being a guanine); (ii) position 106,330,066 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (iii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,071 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,063 on human chromosome 14 (optionally the additional mutation being an adenine); and/or (ii) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,045 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine at a position corresponding to position 106,330,044 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,066 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,066 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,068 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,066 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,027 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine at a position corresponding to position 106,330,024 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,032 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,041 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine or thymine at a position corresponding to position 106,330,063 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the variant comprises additionally a mutation at a position corresponding to position 106,330,071 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,062 on human chromosome

14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

Optionally the genome comprises SEQ ID NO:1; optionally in *cis* at the same Ig locus as one, two or more of the variants.

JH6

An embodiment provides a vertebrate, cell or population of the invention whose genome comprises a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a nucleotide mutation at one or more positions corresponding to positions

106,329,411

106,329,413

106,329,414

106,329,417

106,329,419

106,329,426

106,329,434

106,329,435, and

106,329,468

on human chromosome 14.

Optionally the genome of the vertebrate, cell or population comprises a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a guanine at a position corresponding to position 106,329,435 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,329,468 on human chromosome 14 (optionally the additional mutation being a guanine); (ii) position 106,329,419 on human chromosome 14 (optionally the additional mutation being an adenine); (iii) position 106,329,434 on human chromosome 14 (optionally the additional mutation

being a cytosine) and/or position 106,329,414 on human chromosome 14 (optionally the additional mutation being a guanine); (iv) position 106,329,426 on human chromosome 14 (optionally the additional mutation being an adenine); (v) position 106,329,413 on human chromosome 14 (optionally the additional mutation being an adenine); (vi) position 106,329,417 on human chromosome 14 (optionally the additional mutation being a thymine); (vii) position 106,329,411 on human chromosome 14 (optionally the additional mutation being a thymine); (viii) position 106,329,451 on human chromosome 14 (optionally the additional mutation being an adenine); (ix) position 106,329,452 on human chromosome 14 (optionally the additional mutation being a cytosine); and/or (x) position 106,329,453 on human chromosome 14 (optionally the additional mutation being a cytosine).

Optionally the variant comprises additionally mutations at positions corresponding to position 106,329,451 on human chromosome 14, the additional mutation being an adenine; position 106,329,452 on human chromosome 14, the additional mutation being a cytosine; and position 106,329,453 on human chromosome 14, the additional mutation being a cytosine.

The vertebrate, cell or population optionally comprises a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a guanine at a position corresponding to position 106,329,468 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

Optionally the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the vertebrate, cell or population comprises a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a thymine at a position corresponding to position 106,329,417 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

Optionally the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the vertebrate, cell or population comprises a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a

cytosine at a position corresponding to position 106,329,434 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,329,414 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the vertebrate, cell or population comprises a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a thymine at a position corresponding to position 106,329,411 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

Optionally the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

Optionally the vertebrate, cell or population comprises a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant that is an antisense sequence of a variant described above.

Optionally the genome comprises SEQ ID NO:2; optionally *cis* at the same Ig locus as one, two or more of the JH6 variants.

JH2

An embodiment provides a vertebrate, cell or population of the invention whose genome comprises a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises a nucleotide mutation at one or more positions corresponding to positions

106,331,455

106,331,453, and

106,331,409

on human chromosome 14.

Optionally the vertebrate, cell or population comprises said plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises a guanine at a position corresponding to position 106,331,455 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

Optionally the variant comprises additionally a mutation at a position corresponding to (i) position 106,331,453 on human chromosome 14 (optionally the additional mutation being an adenine); and/or (ii) position 106,331,409 on human chromosome 14 (optionally the additional mutation being an adenine); (iii) position 106,329,434 on human chromosome 14 (optionally the additional mutation being an adenine).

Optionally the vertebrate, cell or population comprises a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises an adenine at a position corresponding to position 106,331,453 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

Optionally the variant comprises additionally a mutation at a position corresponding to position 106,331,409 on human chromosome 14 (optionally the additional mutation being an adenine).

Optionally the vertebrate, cell or population comprises a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises an adenine at a position corresponding to position 106,331,409 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

Optionally the vertebrate, cell or population comprises a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant that is an antisense sequence of a variant described above.

Optionally the genome comprises SEQ ID NO:3; optionally *cis* at the same Ig locus as one, two or more of the JH2 variants.

Optionally the vertebrate, cell or population genome comprises two or more different JH gene segments selected from SEQ ID NOs: 1 to 3 and variants described above; optionally wherein said JH gene segments are *cis* at the same immunoglobulin Ig locus.

Multiple Human D Gene Segment Variants

A specific application of this configuration is the provision of multiple human D gene segments as follows (as set out in numbered clauses, starting at clause number 154).

154. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human D gene segments of the same type (eg, D2-2 gene segments), wherein at least two of the human D gene segments are variants that are not identical to each other (eg, D2-2ref and D2-2a).

In an example of any aspect of the sixth configuration of the invention (V, D, J or C), one or more or all of the variants are naturally-occurring human gene segments.

In an example of any aspect of the sixth configuration of the invention (V, D, J or C), one or more of the variants may be a synthetic variant of a human gene segment.

155. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous D gene segments of the same type type (eg, D2-2ref and D2-2a) *cis* at the same Ig locus.

156. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a) *trans* at the same Ig locus; and optionally a third human D gene segment (eg, (eg, D2-2ref, D2-2a or D2-2b) of the same type, wherein the third D is *cis* with one of said 2 different D gene segments.

157. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human D gene segments, wherein the plurality comprises at least 2 different human D gene segments of the same type (eg, D2-2 gene segments), a first of said different D gene segments (eg, D2-2ref) is provided in the genome of a first vertebrate of the population, and a second of said different D gene segment (eg, D2-2a) being provided in the genome of a second vertebrate of the

population, wherein the genome of the first vertebrate does not comprise the second D gene segment.

158. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, human D2-2 gene segments), wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
159. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human D gene segments of the same type (eg, D2-2 gene segments), wherein at least two of the human D gene segments are variants that are not identical to each other (eg, D2-2ref and D2-2a).
160. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, human D2-2 gene segments) *cis* at the same Ig locus.
161. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a) *trans* at the same Ig locus; and optionally a third human D gene segment (eg, D2-2ref, D2-2a or D2-2b) of the same type, wherein the third D is *cis* with one of said 2 different D gene segments.
162. A method of providing an enhanced human immunoglobulin D gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human D gene segments, wherein the method comprises providing at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a), wherein a first of said different D gene segments is provided in the genome of a first vertebrate of the population, and a second of said different D gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not

comprise the second D gene segment.

163. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, D2-2ref and D2-2a), wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
164. The vertebrate or cell of clause 154, 156 or 158, or the method of clause 159, 161 or 163, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
165. The vertebrate or cell of clause 154, 155 or 156, or the method of any one of clauses 159 to 162 and 164, wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
166. The vertebrate or cell of any one of clauses 154 to 157, or the method of any one of clauses 159 to 162 and 165, wherein the D gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.
167. The vertebrate, cell or method of clause 166, wherein the individuals are not genetically related.
168. The vertebrate, cell or method of any one of clauses 154 to 167, wherein at least one of the different D segments is a synthetic mutant of a human germline D gene segment.
169. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human D gene segments of the same type (eg, D2-2ref and D2-2a), wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgH locus in said genome.

170. The vertebrate or cell of any one of clauses 154 to 158 and 164 to 168, wherein the genome comprises a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.
171. The population of clause 157, wherein the population comprises a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.
172. A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.
173. A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.
174. The vertebrate or cell of clause 172 or the population of clause 173, comprising first and second D gene segments selected from
D2-2ref and D2-2a; or
D2-21ref and D2-21a; or
D3-10ref and D3-10a; or
D3-16ref and D3-16a; or
D2-8ref and D2-8a; or
D3-3ref and D3-3a; or
D4-23ref and D4-23a; or
D6-13ref and D6-13a; or

D3-9ref and D3-9a; or

D4-4ref and D4-4a; or

D7-27ref and D7-27a;

Optionally wherein the first and/or second D gene segment is present in two or more copies.

For example, there are provided two or three copies of the first gene segment, optionally with one, two or three copies of the second gene segment. Copies can be arranged in *cis* or *trans*.

175. The vertebrate, cell or population of clause 174, comprising human gene segments D2-2ref and D2-2a; and D3-3ref and D3-3a; and optionally also D2-15.

In an example, the vertebrate, cell or population comprises one or more D segments selected from human D3-3, D2-15, D3-9; D4-17; D3-10; D2-2; D5-24; D6-19; D3-22; D6-13; D5-12; D1-26; D1-20; D5-18; D3-16; D2-21; D1-14; D7-27; D1-1; D6-25; D2-14 and D4-23 (eg, selected from D3-9*01; D4-17*01; D3-10*01; D2-2*02; D5-24*01; D6-19*01; D3-22*01; D6-13*01; D5-12*01; D1-26*01; D1-20*01; D5-18*01; D3-16*02; D2-21*02; D1-14*01; D7-27*02; D1-1*01; D6-25*01; D2-15*01; and D4-23*01), together with the reference sequence(s) of said selected segment(s). These were found in variable domains having a HCDR3 length of at least 20 amino acids (see examples herein).

176. A non-human vertebrate or vertebrate cell according to clause 155, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene segments in *cis* at the same Ig locus in said genome; and

a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene segments in *cis* at the same Ig locus in said genome;

optionally wherein the D gene segments are derived from the genome sequence of two or more different human individuals.

177. A non-human vertebrate or vertebrate cell according to clause 155, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene segments in *trans* in said genome;

a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene segments in *trans* in said genome;

a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene segments in *trans* in said genome;

a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene segments in *trans* in said genome;

a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene segments in *trans* in said genome;

a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene segments in *trans* in said genome;

a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene segments in *trans* in said genome;

a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene segments in *trans* in said genome;

a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene segments in *trans* in said genome;

a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene segments in *trans* in said genome; and

a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene segments in *trans* in said genome;

optionally wherein the D gene segments are derived from the genome sequence of two or more different human individuals.

178. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell), according to clause 154, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

a plurality of D2-2 gene segments provided by at least 3 different D2-2 gene segments;

a plurality of D2-21 gene segments provided by at least 3 different D2-21 gene segments;

a plurality of D3-10 gene segments provided by at least 3 different D3-10 gene segments;

a plurality of D3-16 gene segments provided by at least 3 different D3-16 gene segments;

a plurality of D2-8 gene segments provided by at least 3 different D2-8 gene segments;

a plurality of D3-3 gene segments provided by at least 3 different D3-3 gene segments;

a plurality of D4-23 gene segments provided by at least 3 different D4-23 gene segments;

a plurality of D6-13 gene segments provided by at least 3 different D6-13 gene segments;

a plurality of D3-9 gene segments provided by at least 3 different D3-9 gene segments;

a plurality of D4-4 gene segments provided by at least 3 different D4-4 gene segments; and

a plurality of D7-27 gene segments provided by at least 3 different D7-27 gene segments;

optionally wherein the D gene segments are derived from the genome sequence of two or three different human individuals;

optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same

Ig locus in said genome.

179. The vertebrate or cell of clause 176, 177 or 178, wherein the different human individuals are from different human populations.
180. The vertebrate or cell of any one of clauses 176 to 179, wherein the individuals are not genetically related.
181. The vertebrate or cell of any one of clauses 176 to 180, wherein at least one of the different D segments is a synthetic mutant of a human germline D gene segment.
182. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to clause 158, comprising a genome comprising human VH, D and JH gene repertoires, the D gene repertoire comprising of one or more of
 - a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene ; optionally in *cis* in said genome;
 - a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene ; optionally in *cis* in said genome;
 - a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene ; optionally in *cis* in said genome;
 - a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene ; optionally in *cis* in said genome;
 - a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene ; optionally in *cis* in said genome;
 - a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene ; optionally in *cis* in said genome;
 - a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene ; optionally in *cis* in said genome;
 - a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene ; optionally in *cis* in said genome;
 - a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene ; optionally in *cis* in said genome;
 - a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene ; optionally in *cis* in

said genome; and

a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene ; optionally in *cis* in said genome;

wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

183. The vertebrate or cell of clause 182, wherein the human individuals are from different human populations.

184. The vertebrate, cell or population of any one of clauses 154 to 183, wherein one or more of the D gene segments is a variant of a human germline D gene segment, wherein the variant gene segment encodes an amino acid sequence that differs by 1, 2 or 3 amino acids from the corresponding amino acid sequence encoded by the human germline D gene segment, provided in that said amino acid sequence encoded by the variant does not include a stop codon when said corresponding amino acid sequence does not include a stop codon.

Optionally, the variant and germline D gene segments encode the respective amino acid sequences in reading frame 2 (IMGT numbering). See Briney *et al* 2012.

185. The vertebrate, cell or population of clause 184, wherein said corresponding amino acid sequence encoded by the human germline D gene segment is a hydrophilic or hydrophobic sequence (according to J Mol Biol. 1997 Jul 25;270(4):587-97; Corbett SJ *et al*; Table 2).

186. The vertebrate, cell or population of clause 184 or 185, comprising said variant and said germline human D gene segments; optionally wherein the variant and germline human D gene segments are *cis* on the same chromosome.

187. The vertebrate, cell or population of any one of clauses 184 to 186, wherein germline human D gene segment is a D2, D3, D5 or D6 family gene segment; optionally a D2-2, D2-15, D3-3, D3-9, D3-10, D3-22, D5-5, D5-18, D6-6, D6-13, D6-19 gene segment.

These D segments are usable in all three reading frames.

Optionally a variant of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or all of these human germline D gene segments is used.

188. The vertebrate, cell or population of any one of clauses 154 to 187, comprising a plurality of D2-2 gene segments, wherein the plurality comprises D2-2 gene segments that vary from each other at one or more nucleotide positions corresponding to positions

106,382,687 and

106,382,711

on human chromosome 14.

189. The vertebrate, cell or population of clause 188, wherein the plurality comprises a human D2-2 gene segment ((optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,382,687 on human chromosome 14; and optionally no further mutation from the sequence of D2-2ref.

190. The vertebrate, cell or population of clause 188 or 189, wherein the plurality comprises a human D2-2 gene segment comprising a cytosine at a position corresponding to position 106,382,687 on human chromosome 14; and optionally no further mutation from the sequence of D2-2a.

191. The vertebrate, cell or population of any one of clauses 188 to 190, wherein the plurality comprises a human D2-2 gene segment comprising an adenine at a position corresponding to position 106,382,711 on human chromosome 14; and optionally no further mutation from the sequence of D2-2b.

192. The vertebrate, cell or population of any one of clauses 188 to 191, wherein the plurality comprises a human D2-2 gene segment comprising an thymine at a position corresponding to position 106,382,711 on human chromosome 14; and optionally no further mutation from the sequence of D2-2ref.

193. The vertebrate, cell or population of any one of clauses 154 to 192, comprising a plurality of D7-27 gene segments, wherein the plurality comprises D7-27 gene segments that vary from each other at a nucleotide position corresponding to position 106,331,767 on human chromosome 14.

194. The vertebrate, cell or population of clause 193, wherein the plurality comprises a human D7-27 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,331,767 on human chromosome 14; and optionally no further mutation from the sequence of D7-27ref.
195. The vertebrate, cell or population of clause 193 or 194, wherein the plurality comprises a human D7-27 gene segment comprising a guanine at a position corresponding to position 106,331,767 on human chromosome 14; and optionally no further mutation from the sequence of D7-27a.
196. The vertebrate, cell or population of any one of clauses 154 to 195, comprising a plurality of D4-23 gene segments, wherein the plurality comprises D4-23 gene segments that vary from each other at a nucleotide position corresponding to position 106,350,740 on human chromosome 14.
197. The vertebrate, cell or population of clause 196, wherein the plurality comprises a human D4-23 gene segment (optionally two copies and/or in homozygous state) comprising an adenine at a position corresponding to position 106,350,740 on human chromosome 14; and optionally no further mutation from the sequence of D4-23ref.
198. The vertebrate, cell or population of clause 196 or 197, wherein the plurality comprises a human D4-23 gene segment (optionally two copies and/or in homozygous state) comprising an guanine at a position corresponding to position 106,350,740 on human chromosome 14; and optionally no further mutation from the sequence of D4-23a.
199. The vertebrate, cell or population of any one of clauses 154 to 197, comprising a plurality of D2-21 gene segments, wherein the plurality comprises D2-21 gene segments that vary from each other at a nucleotide position corresponding to position 106,354,418 on human chromosome 14.
200. The vertebrate, cell or population of clause 199, wherein the plurality comprises a human D2-21 gene segment (optionally two copies and/or in homozygous state) comprising an adenine

at a position corresponding to position 106,354,418 on human chromosome 14; and optionally no further mutation from the sequence of D2-21ref.

201. The vertebrate, cell or population of clause 199 or 200, wherein the plurality comprises a human D2-21 gene segment (optionally two copies and/or in homozygous state) comprising a guanine at a position corresponding to position 106,354,418 on human chromosome 14; and optionally no further mutation from the sequence of D2-21a.
202. The vertebrate, cell or population of any one of clauses 154 to 201, comprising a plurality of D3-16 gene segments, wherein the plurality comprises D3-16 gene segments that vary from each other at a nucleotide position corresponding to position 106,354,418 on human chromosome 14.
203. The vertebrate, cell or population of clause 202, wherein the plurality comprises a human D3-16 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,361,515 on human chromosome 14; and optionally no further mutation from the sequence of D3-16ref.
204. The vertebrate, cell or population of clause 202 or 203, wherein the plurality comprises a human D3-16 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,361,515 on human chromosome 14; and optionally no further mutation from the sequence of D3-16a.
205. The vertebrate, cell or population of any one of clauses 154 to 204, comprising a plurality of D6-13 gene segments, wherein the plurality comprises D6-13 gene segments that vary from each other at a nucleotide position corresponding to position 106,367,013 on human chromosome 14.
206. The vertebrate, cell or population of clause 205, wherein the plurality comprises a human D6-13 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,367,013 on human chromosome 14; and optionally no further mutation from the sequence of D6-13ref.

207. The vertebrate, cell or population of clause 205 or 206, wherein the plurality comprises a human D6-13 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,367,013 on human chromosome 14; and optionally no further mutation from the sequence of D6-13a.
208. The vertebrate, cell or population of any one of clauses 154 to 207, comprising a plurality of D3-10 gene segments, wherein the plurality comprises D3-10 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,370,370 and 106,370,371 on human chromosome 14.
209. The vertebrate, cell or population of clause 208, wherein the plurality comprises a human D3-10 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,370,370 on human chromosome 14; and optionally no further mutation from the sequence of D3-10ref.
210. The vertebrate, cell or population of clause 208 or 209, wherein the plurality comprises a human D3-10 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,370,370 on human chromosome 14; and optionally no further mutation from the sequence of D3-10a.
211. The vertebrate, cell or population of clause 208, 209 or 210 wherein the plurality comprises a human D3-10 gene segment (optionally two copies and/or in homozygous state) comprising an adenine at a position corresponding to position 106,370,371 on human chromosome 14; and optionally no further mutation from the sequence of D3-10ref.
212. The vertebrate, cell or population of any one of clauses 208 to 211, wherein the plurality comprises a human D3-10 gene segment (optionally two copies and/or in homozygous state) comprising a guanine at a position corresponding to position 106,370,371 on human chromosome 14; and optionally no further mutation from the sequence of D3-10b.
213. The vertebrate, cell or population of any one of clauses 154 to 212, comprising a plurality of D3-9 gene segments, wherein the plurality comprises D3-9 gene segments that vary from each

other at a nucleotide position corresponding to position 106,370,567 on human chromosome 14.

214. The vertebrate, cell or population of clause 213, wherein the plurality comprises a human D3-9 gene segment (optionally two copies and/or in homozygous state) comprising an adenine at a position corresponding to position 106,370,567 on human chromosome 14; and optionally no further mutation from the sequence of D3-9ref.
215. The vertebrate, cell or population of clause 213 or 214, wherein the plurality comprises a human D3-9 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,370,567 on human chromosome 14; and optionally no further mutation from the sequence of D3-9a.
216. The vertebrate, cell or population of any one of clauses 154 to 215, comprising a plurality of D2-8 gene segments, wherein the plurality comprises D2-8 gene segments that vary from each other at one or more nucleotide positions corresponding to positions
106,373,085;
106,373,086 and
106,373,089
on human chromosome 14.
217. The vertebrate, cell or population of clause 216, wherein the plurality comprises a human D2-8 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,373,085 on human chromosome 14.
218. The vertebrate, cell or population of clause 216 or 217, wherein the plurality comprises a human D2-8 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,373,085 on human chromosome 14; and optionally no further mutation from the sequence of D2-8b.
219. The vertebrate, cell or population of clause 216, 217 or 218 wherein the plurality comprises a human D2-8 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,373,086 on human chromosome 14; and

optionally no further mutation from the sequence of D2-8ref.

220. The vertebrate, cell or population of any one of clauses 216 to 219, wherein the plurality comprises a human D2-8 gene segment comprising a thymine at a position corresponding to position 106,373,086 on human chromosome 14; and optionally no further mutation from the sequence of D2-8ref.
221. The vertebrate, cell or population of any one of clauses 154 to 220, comprising a plurality of D4-4 gene segments, wherein the plurality comprises D4-4 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,379,086; and 106,379,089 on human chromosome 14.
222. The vertebrate, cell or population of clause 221, wherein the plurality comprises a D4-4 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,379,086 on human chromosome 14; and optionally no further mutation from the sequence of D4-4ref.
223. The vertebrate, cell or population of clause 221 or 222, wherein the plurality comprises a human D4-4 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,379,086 on human chromosome 14; and optionally no further mutation from the sequence of D4-4a.
224. The vertebrate, cell or population of clause 221, 222 or 223 wherein the plurality comprises a human D4-4 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,379,089 on human chromosome 14; and optionally no further mutation from the sequence of D4-4ref or a cytosine at a position corresponding to position 106,379,086 on human chromosome 14.
225. The vertebrate, cell or population of any one of clauses 221 to 224, wherein the plurality comprises a human D4-4 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,373,089 on human

chromosome 14; and optionally no further mutation from the sequence of D4-4a.

226. The vertebrate, cell or population of any one of clauses 154 to 225, comprising a plurality of D3-3 gene segments, wherein the plurality comprises D3-3 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,380,241; and 106,380,246 on human chromosome 14.

227. The vertebrate, cell or population of clause 226, wherein the plurality comprises a D3-3 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,380,241 on human chromosome 14; and optionally no further mutation from the sequence of D3-3ref.

228. The vertebrate, cell or population of clause 226 or 227, wherein the plurality comprises a human D3-3 gene segment (optionally two copies and/or in homozygous state) comprising a cytosine at a position corresponding to position 106,380,241 on human chromosome 14; and optionally no further mutation from the sequence of D3-3a.

229. The vertebrate, cell or population of clause 226, 227 or 228 wherein the plurality comprises a human D3-3 gene segment (optionally two copies and/or in homozygous state) comprising an adenine at a position corresponding to position 106,380,246 on human chromosome 14; and optionally no further mutation from the sequence of D3-3ref.

230. The vertebrate, cell or population of any one of clauses 226 to 229, wherein the plurality comprises a human D3-3 gene segment (optionally two copies and/or in homozygous state) comprising a thymine at a position corresponding to position 106,380,246 on human chromosome 14; and optionally no further mutation from the sequence of D3-3a.

Multiple Human JL Gene Segment Variants

A specific application of this configuration is the provision of multiple human JL gene segments (Jk and/or Jλ) as follows (as set out in numbered paragraphs, starting at paragraph number 80).

80. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human JL gene segments of the same type (eg, Jk1), wherein at least two of the human JL gene segments are variants that are not identical to each other.
81. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1) *cis* at the same Ig locus.
82. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human JL gene segments of the same type (eg, Jk1) *trans* at the same Ig locus; and optionally a third human JL gene segment of the same type, wherein the third JL is *cis* with one of said 2 different JL gene segments.
83. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human JL gene segments, wherein the plurality comprises at least 2 different human JL gene segments of the same type (eg, Jk1), a first of said different JL gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JL gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JL gene segment.
84. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
85. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human JL gene segments of the same type (eg, Jk1), wherein at least two of the human JL gene segments are variants that are not identical to each other.
86. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1) *cis* at the same Ig

locus.

87. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human JL gene segments of the same type(eg, Jk1) *trans* at the same Ig locus; and optionally a third human JL gene segment of the same type, wherein the third JL is *cis* with one of said 2 different JL gene segments.
88. A method of providing an enhanced human immunoglobulin JL gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human JL gene segments, wherein the method comprises providing at least 2 different human JL gene segments of the same type (eg, Jk1), wherein a first of said different JL gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JL gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JL gene segment.
89. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
90. The vertebrate or cell of paragraph 80, 82 or 84, or the method of paragraph 85, 82 or 89, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
91. The vertebrate or cell of paragraph 80, 81 or 82, or the method of paragraph 85, 86 or 87, wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
92. The vertebrate or cell of paragraph 80, 81 or 82, or the method of paragraph 85, 86 or 87, wherein the JL gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human

populations.

93. The vertebrate, cell or method of paragraph 92, wherein the individuals are not genetically related.
94. The vertebrate, cell or method of any one of paragraphs 80 to 93, wherein at least one of the different JL segments is a synthetic mutant of a human germline JL gene segment.
95. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgL locus in said genome.
96. The vertebrate or cell of any one of paragraphs paragraph 80 to 82 and 84, wherein the genome comprises a substantially complete functional repertoire of human Jk and/or Jλ gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.
97. The population of paragraph 83, wherein the population comprises a substantially complete functional repertoire of human JL gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.
98. A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human Jk and/or Jλ gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.

99. A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human J κ and/or J λ gene segment types supplemented with one, two or more human J κ and/or J λ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.
100. A non-human vertebrate or vertebrate cell according to paragraph 81, comprising a genome that comprises VL and JL gene repertoires comprising human gene segments, the JL gene repertoire comprising
- a plurality of human J κ 1 gene segments provided by at least 2 different human J κ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J κ 2 gene segments provided by at least 2 different human J κ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J κ 3 gene segments provided by at least 2 different human J κ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J κ 4 gene segments provided by at least 2 different human J κ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J κ 5 gene segments provided by at least 2 different human J κ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 1 gene segments provided by at least 2 different human J λ 1 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 2 gene segments provided by at least 2 different human J λ 2 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 3 gene segments provided by at least 2 different human J λ 3 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 4 gene segments provided by at least 2 different human J λ 4 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 5 gene segments provided by at least 2 different human J λ 5 gene segments in *cis* at the same Ig locus in said genome;
 - a plurality of human J λ 6 gene segments provided by at least 2 different human J λ 6 gene segments in *cis* at the same Ig locus in said genome; or
 - a plurality of human J λ 7 gene segments provided by at least 2 different human J λ 7 gene segments in *cis* at the same Ig locus in said genome;

optionally wherein the JL gene segments are derived from the genome sequence of two or more different human individuals.

101. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell), according to paragraph 80, comprising a genome that comprises VL and JL gene repertoires comprising human gene segments, the JL gene repertoire comprising

a plurality of human J κ 1 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J κ 1 gene segments;

a plurality of human J κ 2 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J κ 1 gene segments;

a plurality of human J κ 3 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J κ 1 gene segments;

a plurality of human J κ 4 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J κ 1 gene segments;

a plurality of human J κ 5 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J κ 1 gene segments;

a plurality of human J λ 1 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 1 gene segments;

a plurality of human J λ 2 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 2 gene segments;

a plurality of human J λ 3 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 3 gene segments;

a plurality of human J λ 4 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 4 gene segments;

a plurality of human J λ 5 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 5 gene segments;

a plurality of human J λ 6 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 6 gene segments; or

a plurality of human J λ 7 gene segments provided by at least 3 (eg, 3, 4, 5, 6, or 7) different human J λ 7 gene segments;

optionally wherein the JL gene segments are derived from the genome sequence of two or three

different human individuals;

optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same Ig locus in said genome.

102. The vertebrate or cell of paragraph 104 or 105, wherein the different human individuals are from different human populations.

103. The vertebrate or cell of any one of paragraphs 104 to 106, wherein the individuals are not genetically related.

104. The vertebrate or cell of any one of paragraphs 104 to 107, wherein at least one of the different JL segments is a synthetic mutant of a human germline JL gene segment.

105. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to paragraph 84, comprising a genome comprising human VL and JL gene repertoires, the JL gene repertoire comprising

a plurality of human J κ 1 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 2 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 3 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 4 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 5 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 1 gene segments provided by at least 2 different human J λ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 2 gene segments provided by at least 2 different human J λ 2 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 3 gene segments provided by at least 2 different human J λ 3 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 4 gene segments provided by at least 2 different human J λ 4 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 5 gene segments provided by at least 2 different human J λ 5 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 6 gene segments provided by at least 2 different human J λ 6 gene segments, optionally in *cis* at the same Ig locus in said genome; or

a plurality of human J λ 7 gene segments provided by at least 2 different human J λ 7 gene segments, optionally in *cis* at the same Ig locus in said genome;

wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

106. The vertebrate or cell of paragraph 109, wherein the human individuals are from different human populations.

The skilled person will realise that standard molecular biology techniques can be used to provide vectors comprising synthetic combinations of immunoglobulin gene segments (eg, V, D and/or J) for use in the invention, such that the vectors can be used to build a transgenic immunoglobulin locus (eg, using homologous recombination and/or recombinase mediated cassette exchange as known in the art, eg, see US7501552 (Medarex), US5939598 (Abgenix), US6130364 (Abgenix), WO02/066630 (Regeneron), WO2011004192 (Genome Research Limited), WO2009076464, WO2009143472 and WO2010039900 (Ablexis), the disclosures of which are explicitly incorporated herein. For example, such synthetic combinations of gene segments can be made using standard recombineering techniques in *E coli* to construct BAC vectors harbouring the synthetic combination prior to insertion in embryonic stem cells using homologous recombination or RMCE (eg, using cre/lox site-specific recombination). Details of recombineering can be found at www.genebridges.com and in EP1034260 and EP1204740 the disclosures of which are explicitly incorporated herein.

In one embodiment, it is useful to bias the immune response of the vertebrate (and thus resultant lead antibodies) to a predetermined gene segment, eg, one known to be commonly used in natural human immune responses to antigens, such as antigens of infectious disease pathogens. For example, VH1-69 is commonly used to produce antibodies in humans against Influenza virus; it is possible, therefore, to include two or more polymorphic DNA versions of the VH segment VH1-69 in

the locus of the invention. The examples below illustrate how such a transgenic locus can be constructed in which diversity is extended by extending the VH1-69 gene segment repertoire based on naturally-occurring VH1-69 polymorphic variants.

In one embodiment in any configuration of the invention, the genome has been modified to prevent or reduce the expression of fully-endogenous antibody. Examples of suitable techniques for doing this can be found in PCT/GB2010/051122, US7501552, US6673986, US6130364, WO2009/076464, EP1399559 and US6586251, the disclosures of which are incorporated herein by reference. In one embodiment, the non-human vertebrate VDJ region of the endogenous heavy chain immunoglobulin locus, and optionally VJ region of the endogenous light chain immunoglobulin loci (lambda and/or kappa loci), have been inactivated. For example, all or part of the non-human vertebrate VDJ region is inactivated by inversion in the endogenous heavy chain immunoglobulin locus of the mammal, optionally with the inverted region being moved upstream or downstream of the endogenous Ig locus (see, eg, WO2011004192, the disclosure of which is incorporated herein by reference). For example, all or part of the non-human vertebrate VJ region is inactivated by inversion in the endogenous kappa chain immunoglobulin locus of the mammal, optionally with the inverted region being moved upstream or downstream of the endogenous Ig locus. For example, all or part of the non-human vertebrate VJ region is inactivated by inversion in the endogenous lambda chain immunoglobulin locus of the mammal, optionally with the inverted region being moved upstream or downstream of the endogenous Ig locus. In one embodiment the endogenous heavy chain locus is inactivated in this way as is one or both of the endogenous kappa and lambda loci.

Additionally or alternatively, the vertebrate has been generated in a genetic background which prevents the production of mature host B and T lymphocytes, optionally a RAG-1-deficient and/or RAG-2 deficient background. See US5859301 for techniques of generating RAG-1 deficient animals.

Thus, in one embodiment of any configuration or aspect of the invention herein, endogenous heavy and light chain expression has been inactivated.

In one embodiment each said locus constant region is a heavy chain endogenous non-human vertebrate (optionally host mouse or rat) constant region.

In one embodiment each said locus constant region is a light chain endogenous non-human vertebrate (optionally host mouse or rat) constant region.

The invention provides a monoclonal or polyclonal antibody composition prepared by immunisation of at least one vertebrate (eg, mouse or rat) according to the invention, optionally wherein the antigen is an antigen of an infectious disease pathogen (eg, a bacterial or viral pathogen antigen), optionally wherein the same antigen is used to immunise all the vertebrates; optionally wherein the antibody or antibodies are IgG-type (eg, IgG1).

The invention also provides a monoclonal or polyclonal antibody mixture produced by the method of the invention or a derivative antibody or mixture thereof, eg, where one or more constant region has been changed (eg, replaced with a different constant region such as a human constant region; or mutated to enhance or ablate Fc effector function). In an aspect of the invention, the monoclonal or polyclonal antibody mixture is provided for therapy and/or prophylaxis of a disease or condition in a human, eg, for the treatment and/or prevention of an infectious disease, wherein optionally wherein each antibody binds an antigen of an infectious disease pathogen, preferably the same antigen.

In an aspect of the invention, there is provided the use of an isolated, monoclonal or polyclonal antibody according to the invention, or a mutant or derivative antibody thereof in the manufacture of a medicament for the treatment and/or prevention of a disease or condition in a human, eg, an infectious disease, optionally wherein the infectious disease is a disease caused by a bacterial or viral pathogen.

An example of a mutant antibody is one that bears up to 15 or 10 amino acid mutations in its variable regions relative to an isolated antibody (eg, IgG-type, such as IgG1-type, antibody) obtainable or obtained by the method of the invention. An example of a derivative is one that has been modified to replace a constant region with a different constant region such as a human constant region; or mutated to enhance or ablate Fc effector function.

Examples of infectious diseases are diseases caused or mediated by a bacterial or viral pathogen. For example, the infectious disease is selected from the group consisting of a disease caused by a pathogen selected from the group consisting of Haemophilus influenza, E coli, Neisseria meningitidis, a herpes family virus, cytomegalovirus (CMV), HIV and influenza virus.

Tailoring V(D)J Incorporation Into Immunoglobulin Loci For The Generation of Antibodies Against Infectious Disease

The inventors realised that it would be desirable to provide for vertebrates, cells, methods etc for the production of therapeutic and/or prophylactic antibodies based on natural human immune responses to antigens, such as antigens of infectious disease pathogens. In this respect, the literature observes frequently used immunoglobulin gene segments to raise anti-infective responses in humans (Table 9).

In the various configurations, aspects, embodiments and examples above, the invention provides the skilled addressee with the possibility of choosing immunoglobulin gene segments in a way that tailors or biases the repertoire for application to generating antibodies to treat and/or prevent infectious diseases. The inventors have categorised the following groups of gene segments for use in the invention according to the desired application of resultant antibodies.

List A:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by a Pathogen

- (a) a VL gene segment selected from the group consisting of a V_λII gene family member, V_λVII 4A, V_λII 2.1, V_λVII 4A, a V_λ1 gene family member, a V_λ3 gene family member, IGLV1S2, V_λ3-cML70, Iah2, Ialvl, Ia3h3, Kv325, a VκI gene family member, κI-15A (KL012), VκII family member, a VκIII family member, a VκI gene family member, κI-15A (KL012), VκII A2 (optionally the A2a variant), Vκ A27 (Humkv325) and a gene segment at least 80% identical thereto.
- (b) a V_λ gene segment selected from a V_λII gene family member, V_λVII 4A, V_λII 2.1, V_λVII 4A, a V_λ1 gene family member, a V_λ3 gene family member, IGLV1S2, V_λ3-cML70, Iah2, Ialvl, Ia3h3 and a gene segment at least 80% identical thereto.
- (c) a Vκ gene segment selected from Kv325, a VκI gene family member, κI-15A (KL012), VκII family member, a VκIII family member, a VκI gene family member, κI-15A (KL012), VκII A2 (optionally the A2a variant), Vκ A27 (Humkv325) and a gene segment at least 80% identical thereto.
- (d) a V_H gene segment a V_HIII gene family member (optionally, a VHIIIa or VHIIIb family member), a V_HIV gene family member, V_HIII 9.1 (VH3-15), V_HIII VH26 (VH3-23), V_H3-21, LSG6.1, LSG12.1, DP77 (V3-21), V_H H11, VH1GRR, ha3h2, V_HI-ha1c1, V_HIII-VH2-1, VH4.18, ha4h3, Hv1051, 71-2, Hv1f10, VH4.11, 71-4, VH251, VH1-69 and a gene segment at least 80% identical thereto.
- (e) a J_λ gene segment selected from J_λ2, J_λ3 and a gene segment at least 80% identical thereto.

- (f) a D gene segment selected from Dk1, Dxp'1, Dn4r, D2r and a gene segment at least 80% identical thereto.

List A1:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by a Bacterial Pathogen

- (a) a V_{λ} gene segment selected from a V_{λ} II gene family member, V_{λ} VII 4A, V_{λ} II 2.1, V_{λ} VII 4A and a gene segment at least 80% identical thereto.
- (b) a V_{κ} gene segment selected from a V_{κ} I gene family member, κ I-15A (KL012), V_{κ} II family member, a V_{κ} III family member, a V_{κ} I gene family member, κ I-15A (KL012), V_{κ} II A2 (optionally the A2a variant), V_{κ} A27 (Humkv325) and a gene segment at least 80% identical thereto.
- (c) a V_H gene segment a VH3 gene family member (optionally, a VHIIIa or VHIIIb family member), V_H III 9.1 (VH3-15), V_H III VH26 (VH3-23), V_H 3-21, LSG6.1, LSG12.1, DP77 (V3-21), V_H H11 and a gene segment at least 80% identical thereto.
- (d) a J_{λ} gene segment selected from J_{λ} 2, J_{λ} 3 and a gene segment at least 80% identical thereto.
- (e) a J_H gene segment selected from J_H 2, J_H 3, J_H 4 and a gene segment at least 80% identical thereto.

List A1.1:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by *H influenza*

- (a) a V_{λ} gene segment selected from a V_{λ} II gene family member, V_{λ} VII 4A, V_{λ} II 2.1, V_{λ} VII 4A and a gene segment at least 80% identical thereto.
- (b) a V_{κ} gene segment selected from a V_{κ} II family member, a V_{κ} III family member, a V_{κ} I gene family member, κ I-15A (KL012), V_{κ} II A2 (optionally the A2a variant), V_{κ} A27 (Humkv325) and a gene segment at least 80% identical thereto.
- (c) a V_H gene segment a VH3 gene family member (optionally, a VHIIIb family member), V_H III 9.1 (VH3-15), V_H III VH26 (VH3-23), V_H 3-21, LSG6.1, LSG12.1, DP77 (V3-21) and a gene segment at least 80% identical thereto.
- (d) a J_{λ} gene segment selected from J_{λ} 2, J_{λ} 3 and a gene segment at least 80% identical thereto.

List A1.2:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by E Coli or Neisseria meningitidis

- (a) a V_H gene segment a VH3 gene family member (optionally a VHIIIa or VHIIIb member), V_{HIII} 9.1 (VH3-15), V_H H11, V_{HIII} VH26 (VH3-23) a gene segment at least 80% identical thereto, eg, V_{HIII} 9.1 + J_H3 ; or V_H H11 + J_H4 ; or V_{HIII} VH26 + J_H2 .
- (b) a V_K gene segment selected from a V_{KI} gene family member, κ I-15A (KL012) and a gene segment at least 80% identical thereto.
- (c) a V_λ gene segment selected from a $V_{\lambda II}$ gene family member, $V_{\lambda II}$ 2.1 and a gene segment at least 80% identical thereto.
- (d) a J_H gene segment selected from J_H2 , J_H3 , J_H4 and a gene segment at least 80% identical thereto.

A2:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by a viral Pathogen

- (a) a V_H gene segment selected from a V_{HIII} gene family member, a V_{HIV} gene family member, V_{HIII} -VH26 (VH3-23), VH1GRR, ha3h2, V_{HI} -ha1c1, V_{HIII} -VH2-1, VH4.18, ha4h3, Hv1051, 71-2, Hv1f10, VH4.11, 71-4, VH251, VH1-69 and a gene segment at least 80% identical thereto.
- (b) a V_λ gene segment selected from a $V_{\lambda 1}$ gene family member, a $V_{\lambda 3}$ gene family member, IGLV1S2, $V_{\lambda 3}$ -cML70, la1h2, la1v1, la3h3 and a gene segment at least 80% identical thereto.
- (c) a V_K gene segment selected from Kv325 and a gene segment at least 80% identical thereto.
- (d) a J_H gene segment selected from J_H3 , J_H5 , J_H6 and a gene segment at least 80% identical thereto.
- (e) a D gene segment selected from Dk1, Dxp'1, Dn4r, D2r and a gene segment at least 80% identical thereto.
- (f) a J_λ gene segment selected from $J_{\lambda 2}$, $J_{\lambda 3}$ and a gene segment at least 80% identical thereto.

A2.1:

Immunoglobulin gene segments for antibodies that bind an antigen expressed by Herpes Virus Family (eg, VZV or HSV)

- (a) a V_H gene segment selected from a V_{HIII} gene family member, a V_{HIV} gene family member, V_{HIII} -VH26 (VH3-23), VH1GRR, ha3h2, V_{HI} -ha1c1, V_{HIII} -VH2-1, VH4.18, ha4h3, and a gene segment at least 80% identical thereto.

- (b) a V_{λ} gene segment selected from a $V_{\lambda}1$ gene family member, a $V_{\lambda}3$ gene family member, IGLV1S2, $V_{\lambda}3$ -cML70, la1h2, la1v1, la3h3 and a gene segment at least 80% identical thereto.
- (c) a J_H gene segment selected from J_H3 , J_H5 , J_H6 and a gene segment at least 80% identical thereto.
- (d) a D gene segment selected from Dk1, Dxp'1, Dn4r, D2r and a gene segment at least 80% identical thereto.
- (e) a J_{λ} gene segment selected from $J_{\lambda}2$, $J_{\lambda}3$ and a gene segment at least 80% identical thereto.

A2.2:Immunoglobulin gene segments for antibodies that bind an antigen expressed by CMV

- (a) a V_H gene segment selected from Hv1051 and a gene segment at least 80% identical thereto.
- (b) a V_k gene segment selected from Kv325 and a gene segment at least 80% identical thereto.

A2.3:Immunoglobulin gene segments for antibodies that bind an antigen expressed by HIV

- (a) a V_H gene segment selected from 71-2, Hv1f10, VH4.11, 71-4, VH251, VH1-69 and a gene segment at least 80% identical thereto.

A2.4:Immunoglobulin gene segments for antibodies that bind an antigen expressed by Influenza Virus

- (a) a V_H gene segment selected from VH1-69 and a gene segment at least 80% identical thereto.

Thus,

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease, one or more V, D and/or or all J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A1. Thus, for example in (a) of the first configuration of the invention, the recited heavy chain V gene segment is selected from the VH gene segments in List A, optionally with a D in that list.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by a bacterial pathogen, one or more or all V, D and/or J gene

segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A1.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by a viral pathogen, one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A2.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by *H influenza*, one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A1.1.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by *E Coli* or *Neisseria meningitidis*, one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A1.2.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by Herpes Virus Family (eg, VZV or HSV), one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A2.1.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by CMV, one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A2.2.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by HIV, one or more or all V, D and/or J gene segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A2.3.

Where one wishes to generate an antibody or antibody mixture to treat and/or prevent an infectious disease caused or mediated by Influenza Virus, one or more or all V, D and/or J gene

segments used in any configuration, aspect, method, example or embodiment of the invention can be selected from List A2.4.

Optionally each VH segment in the locus of the invention is selected from List A1, A2, A1.1, A1.2, A2.1, A2.2, A2.3 or A2.4.

Optionally each VL segment in the locus of the invention is selected from List A1, A2, A1.1, A1.2, A2.1, A2.2, A2.3 or A2.4

Optionally each D segment in the locus of the invention is selected from List A1, A2, A1.1, A1.2, A2.1, A2.2, A2.3 or A2.4.

Optionally each J_L segment in the locus of the invention is selected from List A1, A2, A1.1, A1.2, A2.1, A2.2, A2.3 or A2.4.

Antibodies For Therapy & Prophylaxis of Patients of Specific Ancestry

The inventors, having undertaken the extensive Bioinformatics analysis exercise described herein, realised that the output of that analysis has made it possible to identify specific gene segments that are useful to produce antibody- and VH domain-based drugs that are tailored specifically to a patient's ancestry (ie, genotype). That is, antibodies can be selected on the basis that they are made *in vivo* in a transgenic non-human vertebrate (eg, mouse or rat with transgenic IgH loci) and particularly derived from gene segments that are relatively prevalent in members of the patient's population, ie, from individuals of the same human ancestry. Since variant distributions differ across different populations (see Table 13), this presumably reflects the effects of evolution, adaptation and conservation of useful variant gene types in those populations. Thus, by tailoring the antibody-based drugs according to the invention, it is possible to match the drug to the population gene biases, thus with the aim of making better drugs for that specific population of humans. Better can, for example, mean more efficacious, better neutralising, higher target antigen affinity, less immunogenic, less patient reactions to the drug etc. This can be determined empirically, as is standard in drug research and development processes.

Thus, the invention provides the following embodiments (numbered from clause 345 onwards):-

345. An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen

and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) in Table 13 found in a Chinese population and with a cumulative frequency of at least 1 or 5%;, and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); and/or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In another embodiment, the invention provides

An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) present in a Chinese population with a cumulative frequency of at least 5%;, and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); and/or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In an example, the constant region is found in the 1000 Genomes database. In an example, the constant region is found in Table 13.

346. The antibody of clause 345 wherein the constant region is a IGHG1a, IGHG2a, IGHG3a, IGHG3b or IGHG4a constant region.

347. The antibody of clause 345 or 346, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

In another embodiment, the invention provides

The antibody of clause 345 or 346, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

348. The antibody of clause 345, 346 or 347, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

In another embodiment, the invention provides

The antibody of clause 345, 346 or 347, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

349. The antibody of clause 345, 346, 347 or 348 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

In another embodiment, the invention provides

The antibody of clause 345, 346, 347 or 348 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

350. An isolated VH domain identical to a variable domain as recited in any one of clauses 347 to 349, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

In an embodiment, there is provided an isolated VH domain identical to a variable domain as recited in any one of clauses 347 to 349 which is part of a conjugate, conjugated with a label (eg, for imaging in the patient) or a toxin (eg, a radioactive toxic payload, such as for cancer treatment in the patient) or a half-life-extending moiety (eg, PEG of human serum albumin).

351. A pharmaceutical composition comprising the antibody or variable domain of any one of clauses 345 to 350 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

352. An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen

and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%; and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); and/or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In another embodiment, the invention provides

An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in a Chinese population with a cumulative frequency of at least 5%; and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); and/or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

353. The antibody of clause 352, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

In another embodiment, the invention provides

The antibody of clause 352, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

354. The antibody of clause 352 or 353, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

In another embodiment, the invention provides

The antibody of clause 352 or 353, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

355. An isolated VH domain identical to a variable domain as recited in any one of clauses 352 to 354, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

In an embodiment, there is provided a VH domain identical to a variable domain as recited in any one of clauses 352 to 354 which is part of a conjugate, conjugated with a label (eg, for imaging in the patient) or a toxin (eg, a radioactive toxic payload, such as for cancer treatment in the patient) or a half-life-extending moiety (eg, PEG of human serum albumin).

356. A pharmaceutical composition comprising the antibody or variable domain of any one of clauses 352 to 355 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

357. An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a Chinese patient, wherein the heavy chain is a heavy chain produced by the following steps (or is a copy of such a heavy chain):-

(a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%;

(b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

In another embodiment, the invention provides

An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a Chinese patient, wherein the heavy chain is a heavy chain produced by the following steps (or is a copy of such a heavy chain):-

(a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived from the

recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in a Chinese population with a cumulative frequency of at least 5%;

(b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

In an example, the VH gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

358. The antibody heavy chain or VH domain of clause 357, wherein the human constant region is as recited in clause 345 or 346.

359. An antibody heavy chain or VH domain as recited in clause 357 or 358 for use in a medicament for therapy and/or prophylaxis of a disease or medical condition in a Chinese patient.

360. A method of treating and/or preventing a disease or medical condition in a Chinese patient, the method comprising administering to the patient a therapeutically or prophylactically-effective amount of the antibody heavy chain or VH domain as recited in clause 357 or 358.

361. An isolated antibody for administration to a patient of European, East Asian, West African, South Asian or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) in Table 13 found in a population of European, East Asian, West African, South Asian or Americas ancestry respectively and with a cumulative frequency of at least 1 or 5%, and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In another embodiment, the invention provides

An isolated antibody for administration to a patient of European, East Asian, West African, South Asian or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) present in a population of European, East Asian, West African, South Asian or Americas ancestry respectively with a cumulative frequency of at least 1 or 5%, and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a

non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In an example, the constant region is found in the 1000 Genomes database. In an example, the constant region is found in Table 13.

362. The antibody of clause 361 wherein the constant region is a IGHG1a, IGHG2a, IGHG3a, IGHG3b or IGHG4a constant region and the patient is of European ancestry.

363. The antibody of clause 361 or 362, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%.

In another embodiment, the invention provides

The antibody of clause 361 or 362, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

364. The antibody of clause 361, 362 or 363, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%.

In another embodiment, the invention provides

The antibody of clause 361, 362 or 363, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

365. The antibody of clause 361, 362, 363 or 364 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%.

In another embodiment, the invention provides

The antibody of clause 361, 362, 363 or 364 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH present in a Chinese population with a cumulative frequency of at least 5%.

In an example, the gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

366. An isolated VH domain identical to a variable domain as recited in any one of clauses 363 to 365, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).
367. A pharmaceutical composition comprising the antibody or variable domain of any one of clauses 361 to 366 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).
368. An isolated antibody for administration to a patient of European, East Asian, West African or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a population of European, East Asian, West African, South Asian or Americas ancestry respectively and with a cumulative frequency of at least 1 or 5%; and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In another embodiment the invention provides:-

An isolated antibody for administration to a patient of European, East Asian, West African or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in a population of European, East Asian, West African, South Asian or Americas ancestry respectively with a cumulative frequency of at least 1 or 5%; and wherein

- (i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg,

mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

In an example, the VH gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

369. The antibody of clause 368, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%.

In another example there is provided

The antibody of clause 368, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D present in said population with a cumulative frequency of at least 1 or 5%.

In an example, the D gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

370. The antibody of clause 368 or 369, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%.

In another example there is provided

The antibody of clause 368 or 369, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH present in said population and with a cumulative frequency of at least 1 or 5%.

In an example, the JH gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

371. An isolated VH domain identical to a variable domain as recited in any one of clauses 368 to 370, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

372. A pharmaceutical composition comprising the antibody or variable domain of any one of clauses 368 to 371 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

373. An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a patient of European, East Asian, West African, South Asian or Americas ancestry, wherein the heavy chain is a heavy chain produced by

the following steps (or is a copy of such a heavy chain):-

(a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in said population and with a cumulative frequency of at least 1 or 5%;

(b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

In another embodiment, there is provided:-

An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a patient of European, East Asian, West African, South Asian or Americas ancestry, wherein the heavy chain is a heavy chain produced by the following steps (or is a copy of such a heavy chain):-

(a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH present in said population with a cumulative frequency of at least 1 or 5%;

(b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

In an example, the VH gene segment is found in the 1000 Genomes database. In an example, the gene segment is found in Table 13.

374. The antibody heavy chain or VH domain of clause 373, wherein the human constant region is as recited in clause 361 or 362.

375. An antibody heavy chain or VH domain as recited in clause 373 or 374 for use in a medicament for therapy and/or prophylaxis of a disease or medical condition in a patient of said ancestry.

376. A method of treating and/or preventing a disease or medical condition in a patient of European, East Asian, West African, South Asian or Americas ancestry, the method comprising administering to the patient a therapeutically or prophylactically-effective amount of the antibody heavy chain or VH domain as recited in clause 373 or 374.

In embodiments herein, a Chinese patient can be a Han Chinese patient.

In embodiments herein, a patient of European ancestry can be a patient of Northern or Western European ancestry, Italian ancestry, British or Scottish ancestry, Finnish ancestry or Iberian ancestry.

In embodiments herein, a patient of East Asian ancestry can be a patient of Han Chinese ancestry, Japanese ancestry Chinese Dai ancestry, Vietnamese ancestry or Kinh ancestry.

In embodiments herein, a patient of West African ancestry can be a patient of Yoruba ancestry, Luhya ancestry, Gambian ancestry or Malawian ancestry.

In embodiments herein, a patient of Americas ancestry can be a patient of African American ancestry, African Caribbean ancestry, Mexican ancestry, Puerto Rican ancestry, Colombian ancestry or Peruvian ancestry.

In embodiments herein, a patient of South Asian ancestry can be a patient of Ahom ancestry, Kayadtha ancestry, Reddy ancestry, Maratha ancestry, or Punjabi ancestry.

In an example of any aspect, the cumulative frequency is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90 or 95%.

It will be understood that particular embodiments described herein are shown by way of illustration and not as limitations of the invention. The principal features of this invention can be employed in various embodiments without departing from the scope of the invention. Those skilled in the art will recognize, or be able to ascertain using no more than routine study, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the claims. All publications and patent applications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference. The use of the word "a" or "an" when used in conjunction with the term "comprising" in the claims and/or the specification may mean "one," but it is also consistent with the meaning of "one or more," "at least one," and "one or more than one." The use

of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or." Throughout this application, the term "about" is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

As used in this specification and claim(s), the words "comprising" (and any form of comprising, such as "comprise" and "comprises"), "having" (and any form of having, such as "have" and "has"), "including" (and any form of including, such as "includes" and "include") or "containing" (and any form of containing, such as "contains" and "contain") are inclusive or open-ended and do not exclude additional, unrecited elements or method steps

The term "or combinations thereof" as used herein refers to all permutations and combinations of the listed items preceding the term. For example, "A, B, C, or combinations thereof" is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, MB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

Any part of this disclosure may be read in combination with any other part of the disclosure, unless otherwise apparent from the context.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or

methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

The present invention is described in more detail in the following non limiting prophetic Examples.

EXAMPLES

EXAMPLE 1

“Recombineered BAC Vectors to add Polymorphic V-regions to the Mouse Genome”

Figure 1 through 3 depict recombineering methods (see references above) that can be used to introduce polymorphic V-gene regions into genomic DNA. In one embodiment, a genomic fragment from the human heavy chain region is inserted into a bacterial artificial chromosome (BAC) vector by standard techniques. Preferably, such a BAC, which can range in size from 20-kb to 200-kb or more, can be isolated from libraries of BACs by standard techniques including sequence searches of commercially available libraries or by hybridization to bacterial colonies containing BACs to identify those with a BAC of interest.

A BAC is chosen that has several VH gene segments; in Figure 1, these are generically identified as VH[a] through VH[z] for example. One skilled in the art will readily identify appropriate genomic fragments, for example, an approximately 120-kb fragment from human VH5-78 through VH1-68 which includes 5 endogenous active VH gene segments and 7 VH psuedogenes. Using recombineering techniques, the endogenous VH gene segments can be replaced by polymorphic VH or VL gene segments. In this example, two steps are required. The first step replaces the V-region coding exon of an endogenous VH gene segment with a positive-negative selection operon, in this example, an operon encoding an ampicillin resistance gene (*Amp*) and a streptomycin-sensitizing ribosomal protein (*rpsL*). Certain strains of bacteria can be selected for the absence of the *rpsL* gene by resistance to streptomycin. Short stretches of DNA homologous to sequences flanking the

endogenous VH gene exon are placed 5' and 3' of the *rpsL*-Amp operon. In the presence of appropriate recombination factors per standard recombineering techniques (see references above) recombination between the operon fragment and the BAC will result in replacement of the endogenous VH gene exon with the operon (Figure 1a) which are selected by resistance to ampicillin. The second step uses the same homologous sequences in order to replace the inserted operon with a desired polymorphic VH gene segment. In this example, a human VH1-69 gene is inserted (Figure 1b and 1c). In particular the *02 variant of VH1-69 is used [ref IMGT and Figure 5]. Successful integrations of the polymorphic VH gene segment are selected in bacteria that become resistant to streptomycin due to the loss of the operon, specifically the *rpsL* portion.

In this example, the two step process as described can be repeated for each of the endogenous VH gene segments or for as many endogenous gene segments that one wishes to replace with polymorphic V gene segments (Figure 1d).

As is apparent, any polymorphic V gene segment can be inserted in this manner and any endogenous V gene segment can act as a target, including pseudogenes. V gene segments in each of the heavy chain and two light chain loci can be replaced using this technique with appropriate genomic fragments available as BAC inserts.

Figure 2 depicts another method for creating a genomic fragment encoding polymorphic V gene segments. In this example, polymorphic V gene segments are inserted into a region of genomic DNA devoid of other genes, control elements or other functions. Such 'desert' regions can be selected based on sequence analysis and corresponding DNA fragments cloned into BACs or identified in existing BAC libraries. Starting with such a genomic fragment, recombineering techniques can be used to insert polymorphic V gene segments at intervals of, for example, 10-kb. In this example, a 150-kb genomic fragment might accommodate insertion of up to 15 polymorphic V gene segments. Insertion of the segments is a two-step process. The first recombineering step inserts the *rpsL*-Amp operon at a specific site. Sequences homologous to a specific site are used to flank the operon. These are used by the recombineering system to insert the element specifically into the BAC genomic fragment and positive events are selected by resistance to ampicillin (Figure 2a). The second step replaces the operon in the genomic fragment with a polymorphic V gene segment by a similar recombineering step using the same sequence homology (Figure 2b). In this example, both

exons and promoter element of a polymorphic VH gene segment are inserted, resulting in replacement of the *rpsL-Amp* operon and therefore resistance to streptomycin (Figure 2c).

The two step technique for inserting polymorphic V gene segments into a specific site on the genomic fragment can be repeated multiple times resulting in a BAC genomic fragment with several polymorphic gene segments, including their promoter elements. It is apparent that the examples shown in Figures 1 and 2 can be combined wherein the technique for insertion can be used to add *extra* polymorphic V gene segments to a BAC genomic fragment as depicted in Figure 1. One might choose to add these extra segments to an IG genomic fragment since such a fragment would be more amenable to proper IG gene expression once inserted into a non-human mammal's genome. It is known that a genomic fragment can have elements such as enhancers or elements that contribute to certain chromatin conformations, both important in wild-type gene expression.

Figure 3 depicts an additional method to create genomic fragments with polymorphic V gene segments. This method depends upon the efficiency with which short (around 50 to 150 bases, preferably 100 bases) single stranded DNA fragments recombine with a homologous sequence using recombineering (Nat Rev Genet. 2001 Oct;2(10):769-79; Recombineering: a powerful new tool for mouse functional genomics; Copeland NG, Jenkins NA, Court DL). The recombinases used in recombineering preferentially bind and use such short single-stranded fragments of DNA as a substrate for initiating homologous recombination. The efficiency can be as high as 10^{-2} , that is, a positive event can be found in approximately 100 randomly picked (not selected) clones resulting from recombineering. A positive event in this example occurring when one or more single nucleotide changes introduced into the single-stranded fragment get transferred to the BAC insert containing V gene segments and surrounding genomic DNA, said nucleotide change or changes occurring at a homologous sequence on the BAC.

Polymorphic V gene segments can differ from endogenous V gene segments by only 1 or 2, or up to 10 or 15 nucleotide changes, for example. An example of such nucleotide polymorphisms are depicted in Figure 5. Short single stranded regions that encompass the polymorphic nucleotide changes can be chemically synthesized using standard techniques. The resulting single stranded DNA fragments are introduced into bacteria and *via* recombineering techniques approximately 1 in 100 BAC fragments will have incorporated the polymorphic nucleotides *via* homologous incorporation of

the single stranded fragment (Figure 3a). BACs with the desired nucleotide change can be identified by screening for example several hundred individual clones by polymerase chain reaction (PCR) amplification and sequencing, both by standard techniques. In the example, two nucleotide changes will convert a VH1-69*01 gene segment into a VH1-69*02 gene segment (Figure 3b).

It is clear that this process can be repeated for multiple endogenous V gene segments contained on a single BAC genomic fragment. In addition, the techniques depicted in Figure 2 can be used to add additional polymorphic V gene segments by insertion into regions between existing V gene segments. As would be evident to one skilled in the art, a combination of these techniques can be used to create numerous variations of both polymorphic and endogenous human V gene segments. And it would be evident that several different genomic fragments with engineered polymorphic V gene segments and endogenous human V gene segments can be combined to create even more variations.

EXAMPLE 2

“Adding Polymorphic V-regions to the Genome using SRMCE of Modified BACs”

Modified BACs with polymorphic V gene segments created using the methods described in Example 1 can be used to alter the genome of non-human mammals. These alterations can result in an intact IG locus in which normal immunoglobulin region recombination results in VDJ or VJ combinations which includes the human V gene segments. An example of how such an animal can be created is by altering the genome of, for example, mouse embryonic stem (ES) cells using the strategy outlined in Figure 4.

One technique to integrate modified BACs with polymorphic V gene segments into a genome is sequential recombinase mediated cassette exchange (SRMCE). The technique is described in WO2011004192 (Genome Research Limited), which is incorporated here in its entirety by reference.

SRMCE provides for a locus modified with a 'landing pad' inserted at a specific location. This insertion can either be *de novo* via homologous recombination or as a consequence of a previous BAC insertion. In this example, the landing pad is inserted in the mouse IGH locus between the most 3' J gene segment and the C μ gene segment and a previous BAC insertion *via* SRMCE techniques have resulted in the addition of 5 human V gene segments and 2 V region pseudogenes. The landing pad has elements as shown in Figure 4 that will allow the selection of correct insertion of a second targeting BAC fragment. The specificity of this insertion is provided by cre recombinase-mediated exchange between permissive *lox* sites. A *lox* site is permissive for recombination only with a compatible *lox* site. In this example, the *loxP* site will only recombine with *loxP* and *lox2272* will only recombine with *lox2272*. This provides directionality to the insertion of the BAC fragment as depicted in Figure 4b and 4c.

ES cell clones with correct insertions are selected from a pool of clones without insertions or with non-productive insertions by resistance to puromycin. Resistance to puromycin results from the juxtaposition of an active promoter element, PGK, with the *puroTK* coding region. Correct insertions are verified by standard techniques including PCR of junctions, PCR of internal elements, Southern blotting, comparative genomic hybridization (CGH), sequencing and *etc.* In the example, correct *lox2272-lox2272* and *loxP-loxP* recombination also results in two intact sets of *piggyBac* elements that did not exist prior to insertion. An intact *piggyBac* element is comprised of a set of inverted repeats which are depicted in the figure by "PB5'" and "PB3'". An appropriately oriented set of *piggyBac* elements are the substrate of *piggyBac* transposase which can catalyse recombination between the elements, resulting in deletion of intervening sequences as well as both elements. The DNA remaining after a *piggyBac* transposition is left intact and is lacking any remnant of the *piggyBac* element. In the example, ES cell clones with successful *piggyBac* transposition are selected by loss of the active *puroTK* element which renders the cells resistant to the drug FIAU (Figure 4c and 4d).

The final product of the SRMCE method in this example is a IGH locus with several polymorphic V gene segments inserted along with a set of endogenous unmodified VH gene segments between sequences of the mouse genome on the 5' side and the mouse IGH constant region gene segments on the 3' side. The polymorphic V gene segments are positioned such that they can participate in the recombination events associated with B cell maturation yielding VDJ gene segments. These gene

segments can then be transcribed and spliced to the mouse constant region. Translation of these transcripts will result in the production of an antibody heavy chain encoded by the polymorphic V gene segment, a human DH gene segment, a human JH gene segment and a mouse constant heavy chain gene segment.

As is well known to those skilled in the art, an ES cell clone can be used to create a line of genetically modified mice *via* injection of said cells into a mouse blastocyst embryo, transferring the injected embryo to a suitable recipient and breeding the chimeric offspring that result. The modified gene locus can be propagated through breeding and made either heterozygous or homozygous depending on the genetic cross.

It is evident from the structure of the IGH locus provided in this example and by knowledge of the mechanisms involved in B cell receptor (BCR) and antibody gene rearrangements that a large set of different combinations of polymorphic V gene segments with various DH and JH gene segments will result and these can contribute to a large repertoire of functional antibody genes in a population of B cells in genetically modified animals. In this example, several different human VH1-69 polymorphs are incorporated to provide superhuman VH diversity. This particular VH gene segment is known to be prevalent in antibodies that bind infectious disease pathogens (such as influenza virus) and therefore the antibody repertoire of a mouse with the genetic modification of this example would be expected to produce antibodies with a bias in favour of those that bind infectious disease pathogens. The repertoire, in other words, would have a larger subset of antibodies with superior affinities for pathogen antigens. Examples of such pathogens include influenza virus, hepatitis C virus (HCV) and human immunodeficiency virus-1 (HIV-1) (see also table above).

EXAMPLE 3

“Alignment of 13 VH1-69 Alleles”

Building a more diverse antibody repertoire by incorporating additional V gene segment polymorphs requires availability of polymorphic variants of V gene segments. One source of such variants include sequence databases. In this example, 13 distinct variants of the VH1-69 gene segment are provided.

These variant sequences and comparisons are drawn from the “IMmunoGeneTics” IMGT Information System (www.imgt.com) database. Figure 5 is a diagram of the alignment of variants *02 through *13 with the *01 variant. The VH1-69*01 nucleotide and amino acid sequence is provided at the top of the figure. Where the remaining variants are identical to the *01 variant sequence a dash is inserted below the sequence. Nucleotide differences are noted alongside the appropriate variant and if the sequence change results in a protein coding change, the amino acid change is indicated above the triplet.

Figure 5 depicts between 1 and 4 amino acid changes for each variant in comparison to the *01 variant. All of the amino acid changes occur in the part of the heavy chain protein encoding the complementarity determining regions (CDRs). These regions are responsible for antigen specificity and the affinity of the antibody for the antigen. It is evident that providing additional polymorphic CDRs in a repertoire of antibodies will increase the likelihood of there being an antibody with superior binding characteristics for various antigens. In several reports, it has been observed that the VH1-69-encoded variable region of the heavy chain is often found in antibodies that bind influenza virus, HCV and HIV-1 antigens (see table above). Therefore incorporating the polymorphic V gene segments of this example into a transgenic animal model using the methods of Examples 1 and 2 would likely result in an antibody repertoire in said transgenic animal with more antibodies that bind to antigens associated with these and other pathogens. And as is known in the art, a larger repertoire increases the probability of finding monoclonal antibodies using, for example, hybridoma technology, that bind with high affinity and specificity to a desired antigen.

This disclosure therefore describes in these examples a transgenic mouse model which can be immunized with pathogen or other antigens. Plasma B cells from such an immunized mouse can be used to make a hybridoma library that can be screened for production of antibodies that bind the pathogen antigens. This library will be superior to libraries from traditional transgenic mice for finding such antibodies given the addition of polymorphic VH1-69 gene segments to the IGH locus in said transgenic mouse.

These examples are not limiting to the human polymorphic V gene segments that can be chosen or to the methods used to introduce them into an animal model. The method can be used to construct a transgenic locus with immunoglobulin D and/or J segments. The V, D, J segments can be from a plurality of human sources (optionally more than one human ethnic population).

EXAMPLE 4**Human IgH JH Gene Variants Selected from the 1000 Genomes Database**

Data is presented for human JH2, 5 and 6 variants. In Tables 10A, 11A and 12A samples from humans from various populations are listed where the sequence analysis of the inventors has revealed the presence of polymorphisms in one or both IgH JH alleles. The population codes are explained in Table 8 above. The polymorphisms are nucleotide variants from JH2, 5 and 6 reference sequences (SEQ ID NOs: 1, 2 and 3 respectively; see below). All references are sequences taken from the Ensembl database (www.ensembl.org). The JH5 reference is human IgH J5-001 disclosed in that database. The JH6 reference is human IgH J6-001 disclosed in that database. The JH2 reference is human IgH J2-001 disclosed in that database.

The reference nucleotide and encoded amino acid sequences are shown on the next page. Alignments with encoded amino acid sequences are also provided, including the corresponding position numbers on human chromosome 14.

Variant Frequencies are shown in Tables 10A, 11A and 12A and these relate to the frequency of the variants in the 1000 Genomes Database (release current at October 2011).

Tables 10B, 11B and 12B show the non-synonymous nucleotide polymorphisms in the human JH variants, as sorted by the present inventors from the 1000 Genomes database. Position numbers corresponding to nucleotide positions on human chromosome 14 are shown for variant positions (chromosome 14 being the chromosome bearing the IgH locus in humans). Thus, for example, the first entry in Table 11B is "14:106330027:A/C" which refers to a position in a variant JH5 sequence wherein the position corresponds to position 106,330,027 on human chromosome 14, such position being A (adenine) in the reference sequence. The "C" indicates that the present inventors observed a mutation to cytosine at this position in the variants found in the 1000 Genomes database. This change leads to a change at the amino acid level of the encoded sequence (ie, a "non-synonymous" change), in this case a change from a serine (found in the reference) to an alanine in the variant.

Example 5:

Human Antibody Gene Segment Variant Identification & Population Analysis

The genomic coding region coordinates for each target gene for variant analysis were identified from the Ensembl WWW site (www.ensembl.org) using coordinates from the GRCh.p8 Human Genome assembly (www.ncbi.nlm.nih.gov/projects/genome/assembly/grc). Using the collected gene location coordinates, variant data was extracted from the public ftp site of the 1000 Genomes Project using the Perl 'Variant Pattern Finder' (VPF - www.1000genomes.org/variation-pattern-finder-api-documentation).

Data extracted by VPF was post processed using software to extract all non-synonymous (NSS) variants with their associated genotype calls. Genotypes calls were assembled to form unique haplotypes, representing groups of NSS variants associated with 1000 Genome population groups and frequency of occurrence within those populations.

The output of the analysis results in tables such as in Table 13. The main body of the table describes each haplotype in turn giving a unique ID for that gene (in the range a-z,aa-zz), the population frequencies and occurrence in individuals and unique population groups; one or more subsequent columns describe the DNA base calls at each location that form the haplotype giving both the base from the reference sequence or the variant base call.

Table 13 was constructed in this manner. The table can be read as follows:

The first four columns (left to right) consist of (1) the haplotype ID letter ('ref' indicates reference - the DNA base call at each genomic location from the GRCh37 Human Reference Assembly) (2) the observed cumulative frequency of the haplotype among the different populations (3) the number of individuals in which a specific haplotype was observed (4) the number of unique population groups that the identified individuals belong to (the actual population group identifiers are displayed as a string of ID's in the most right hand column for each haplotype. For example haplotype 'a' has a population ID string of '3,4,9,13').

The populations are numbered as follows (population labels being according to 1000 Genomes Project nomenclature)

- 1= ASW;
- 2= CEU;
- 3=CHB;
- 4=CHS;
- 5=CLM;
- 6=FIN;
- 7=GBR;
- 8=IBS;

9=JPT;
 10=LWK;
 11=MXL;
 12=PUR;
 13=TSI;
 14=YRI.

Subsequent columns detail a single point variant and have the following format (top to bottom) (1) the human genomic location of the variant (format [chromosome number]: [location] e.g. '14:106204113'); (2) The identifier for the point variant as defined in DbSNP (www.ncbi.nlm.nih.gov/projects/SNP/); (3) One or additional rows show the amino acid change as result of the variant for a specific transcript (denoted by the Ensembl transcript ID in the most right-hand column for each row), the format is the amino acid in the reference sequence followed by '->' and the amino acid caused by the substitution of the variant in the reference sequence (e.g. 'Gly->Arg' means a that the translated reference sequence would result in a glycine at that location, whereas the substitution of the identified variant would result in translated protein containing arginine) using the IUPAC three letter amino acid codes (<http://pac.iupac.org/publications/pac/pdf/1972/pdf/3104x0639.pdf>). Subsequent rows (one per haplotype) show the DNA base at each location, bases matching the reference sequence are shown in black on white back ground, bases varying from the reference are shown as white text on a black background.

The most right-hand column contains the Ensembl transcript ID's (e.g. 'ENST00000390542') for each of the gene transcript and relates to the amino acid changes to the left of this column. Because the transcripts are differing lengths each variant position may or may not have an associated amino acid change at the that position.

Example 6:

Transgenic Mice, B-cells, Hybridomas, Antibodies & Heavy Chains Based on Human JH6*02

A functional human gene segment repertoire (from V_H2-26 to J_H6, see the IMGT database for the structure of the human IgH locus;

<http://www.imgt.org/IMGTrepertoire/index.php?section=LocusGenes&repertoire=locus&species=human&group=IGK>) was sectored by the inventors to produce two different transgenic heavy chain alleles (denoted S2F and S3F) and corresponding mice. The transgenic alleles were expressed in the mice and the heavy chain repertoires were assessed at the RNA transcript level. Deep sequence

analysis was carried out using Bioinformatics methods to assess V, D and JH gene usage, including in variable domain sequences having a HCDR3 length of at least 20 amino acids. Endogenous, mouse variable region gene segments were inactivated by inversion (as per the method described in WO2011004192, this disclosure being incorporated herein by reference).

Sequencing of Human Donor DNA Samples: Identification of Conserved JH6*02 Variant

DNA samples from 9 anonymised consenting human donors were obtained by taking cheek swabs.

The samples were processed and the DNA Samples were extracted follow the protocol of QIAamp DNA Mini Kit (Cat.No.51304, Qiagen).

PCR reactions were set up to amplify the JH6 region and PCR products were sequenced (PCR Oligos sequence: Fwd. 5'-AGGCCAGCAGAGGGTTCCATG-3' (SEQ ID NO: 444), Rev. 5'-GGCTCCCAGATCCTCAAGGCAC-3' (SEQ ID NO: 445)).

Sequence analysis was carried out by comparing to the JH6 reference sequence from IMGT annotated database (<http://www.imgt.org/>), and this identified that all 9 donor genomes contained the human JH6*02 variant, with this variant being in the homozygous state in 7 out of the 9 donors. The inventors also consulted the genomic sequences publicly available for Jim Watson and Craig Venter at Ensembl human genome database [<http://www.ensembl.org/>]. These too contained the human JH6*02 variant. This confirmed to the inventors that human JH6*02 is a common, conserved variant in humans, and thus a good candidate for construction of a transgenic IgH locus as per the invention

Identification of Suitable Human DNA Sequence BACs

A series of human bacterial artificial chromosome (BAC) clones were identified from Ensemble (<http://www.ensembl.org/index.html>) or UCSC (<http://genome.ucsc.edu/>) human database searches based on gene name (IGH) or location (chromosome 14: 106026574-107346185). Seven human RP11 BAC clones (see an extract of the UCSC databse in Figure 10, identified BACs being circled) were selected, RP11-1065N8 BAC carrying human JH6*02. In total, the following BACs were

identified as sources of human IgH locus DNA: RP11-1065N8, RP11-659B19, RP11-141I7, RP-112H5, RP11-101G24, RP11-12F16 and RP11-47P23.

With a similar approach, different BAC clones (eg, different RP11 clone IDs or different sources from RP11) or genetically engineered BACs can be selected for insertion into the mouse IGH locus to provide different sets of human repertoires in the transgenic mouse.

Construction of Transgenic IgH Loci

Insertion of human heavy gene segments from a 1st IGH BAC (RP11-1065N8) into the IGH locus of mouse AB2.1 ES cells (Baylor College of Medicine) was performed to create a heavy chain allele denoted the S1 allele. The inserted human sequence corresponds to the sequence of human chromosome 14 from position 106494908 to position 106328951 and comprises functional heavy gene segments V_H2-5, V_H7-4-1, V_H4-4, V_H1-3, V_H1-2, V_H6-1, D1-1, D2-2, D3-9, D3-10, D4-11, D5-12, D6-13, D1-14, D2-15, D3-16, D4-17, D5-18, D6-19, D1-20, D2-21, D3-22, D4-23, D5-24, D6-25, D1-26, D7-27, J_H1, J_H2, J_H3, J_H4, J_H5 and J_H6 (in 5' to 3' order), wherein the JH6 was chosen to be the human JH6*02 variant. The insertion was made between positions 114666435 and 114666436 on mouse chromosome 12, which is upstream of the mouse C_μ region. The mouse V_H, D and J_H gene segments were retained in the locus, immediately upstream of (5' of) the inserted human heavy chain DNA.

A second allele, S2 was constructed in which more human functional V_H gene segments were inserted upstream (5') of the 5'-most V_H inserted in the S1 allele by the sequential insertion of human DNA from a second BAC (BAC2). The inserted human sequence from BAC2 corresponds to the sequence of human chromosome 14 from position 106601551 to position 106494909 and comprises functional heavy chain gene segments V_H3-13, V_H3-11, V_H3-9, V_H1-8, V_H3-7. The mouse V_H, D and J_H gene segments were retained in the locus, immediately upstream of (5' of) the inserted human heavy chain DNA. In a subsequent step, these were inverted to inactivate them, thereby producing S2F mice in which only the human heavy chain variable region gene segments are active.

A third allele, S3 was constructed in which more human functional V_H gene segments were inserted upstream (5') of the 5'-most V_H inserted in the S2 allele by the sequential insertion of human DNA

from a third BAC (BAC3). The inserted sequence corresponds to the sequence of human chromosome 14 from position 106759988 to position 106609301, and comprises functional heavy chain gene segments, V_H2-26, V_H1-24, V_H3-23, V_H3-21, V_H3-20, V_H1-18, and V_H3-15. The mouse V_H, D and J_H gene segments were retained in the locus, immediately upstream of (5' of) the inserted human heavy chain DNA. In a subsequent step, these were inverted to inactivate them, thereby producing S3F mice in which only the human heavy chain variable region gene segments are active.

Mice bearing either the S2F or S3F insertion into an endogenous heavy chain locus were generated from the ES cells using standard procedures. The other endogenous heavy chain locus was inactivated in the mice by insertion of an inactivating sequence comprising neo^R into the mouse J_H-C_μ intron (to produce the "HA" allele).

Immunisation procedure

Transgenic mice of the S2F or S3F genotype were primed with 20-40ug recombinant proteins obtained commercially or produced in house with Antigen 1 (OVA (Sigma A7641); Antigen 2 (a human infectious disease pathogen antigen) and Antigen 3 (a human antigen) via the ip route in complete Freund's adjuvant (Sigma F 5881) and 10ug/animal CpG (CpG oligo; Invivogen, San Diego, California, USA) and then boosted twice in about two weekly intervals with about half the amount of antigen in incomplete Freund's adjuvant (Sigma F 5506) and 10ug/animal CpG. Final boosts were administered two weeks later iv without any adjuvant and contained 5-10 ug protein in PBS.

Hybridoma fusion procedure

Spleens were taken 3 days after the final boost and spleenocytes were treated with CpG (25 μm final concentration) for and left until the following day. Cells were then fused with SP0/2 Ag14 myeloma cells (HPA Cultures Cat No 85072401) using a BTX ECM2001 electrofusion instrument. Fused cells were left to recover for 20 minutes then seeded in a T75 flask until next morning. Then the cells were spun down and plated out by dilution series on 96-well culture plates and left for about 10 days before screening. Media was changed 1-3 times during this period.

Screening

Screening

Culture supernatants of the hybridoma wells above were screened using homogenous time resolved fluorescence assay (htfrf) using Europium cryptate labelled anti-mouse IgG (Cisbio anti-mouse Ig Europium Cryptate) and a biotin tagged target antigen with a commercially available streptavidin conjugated donor (Cisbio; streptavidin conjugated D2) or by IgG-specific 384 well ELISA. Positive wells identified by htfrf were scaled to 24-well plates or immediately counterscreened using an IgG-specific detection ELISA method. Positives identified by primary ELISA screen were immediately expanded to 24-well plates. Once cultures were expanded to 24-well stage and reached confluency, supernatants were re-tested using htfrf or IgG-specific ELISA to confirm binding to target antigen. Supernatant of such confirmed cultures were then also analysed by surface plasmon resonance using a BioRad ProteOn XPR36 instrument. For this, antibody expressed in the hybridoma cultures was captured on a biosensor GLM chip (BioRad 176-512) which had an anti-mouse IgG (GE Healthcare BR-1008-38) covalently coupled the biosensor chip surface. The antigen was then used as the analyte and passed over the captured hybridoma antibody surface. For Antigen 2 and Antigen 3, concentrations of 256nM, 64nM, 16nM, 4nM and 1nM were typically used, for Antigen 1, concentrations of 1028nM, 256nM, 64nM, 16nM and 4nM were typically used, binding curves were double referenced using a 0nM injection (i.e. buffer alone). Kinetics and overall affinities were determined using the 1:1 model inherent to the BioRad ProteOn XPR36 analysis software.

Any clones with confirmed binding activity were used for preparing total RNA and followed by PCR to recover the heavy chain variable region sequences. Standard 5'-RACE was carried out to analyse RNA transcripts from the transgenic heavy chain loci in the S2F and S3F mice. Additionally, deep sequence analysis of almost 2000 sequences produced by the mice was carried out.

Bionformatics Analysis

Sequences for analysis were obtained from two different methods:

- The first is from RNA extracted from the spleen: first cDNA strand was synthesized using an oligo based on the Cmu region of the mouse IGH locus as a PCR template. PCR was performed using this oligo with an oligo dT-anchor primer. Then PCR product was cloned into pDrive vector (Qiagen) and then sequenced.

- The second is from hybridomas generated through electro-fusion: total RNA was extracted from hybridoma lines of interest using standard Trizol methods and frozen at -80 °C for long term storage. cDNA was generated from 100ng total RNA using standard Superscript III reverse transcriptase and a gene-specific reverse primer binding to all mouse IgG isotypes for heavy chain and a mouse kappa constant region primer for the light chain amplification. 2-3 ul of cDNA were then used as template in a PCR reaction using Pfu DNA polymerase and a panel of degenerate forward primers annealing to the leader sequence of the human immunoglobulin variable domain as well as one mouse pan-IgG reverse primer. PCR products were run out of a 1% agarose gel and bands of approximately 350-450 basepairs extracted and purified. DNA was then sequenced.

The sequences from the first method can either be from IgM from Naïve mice or IgG from immunised mice. The samples from the second method are all from IgG from immunised mice, and specific to the immunising antigen. Almost 2000 sequences were analysed.

The sequences were obtained as a pair of forward and reverse reads. These were first trimmed to remove low-quality base calls from the ends of the reads (trimmed from both ends until a 19 nucleotide window had an average quality score of 25 or more). The reads were combined together by taking the reverse complement of the reverse read, and aligning it against the forward read. The alignment scoring was 5 for a match, -4 for a mismatch, a gap open penalty of 10 and a gap extension penalty of 1. A consensus sequence was then produced by stepping through the alignment and comparing bases. When there was a disagreement the base with the highest quality value from sequencing was used.

The BLAST+ (Basic Local Alignment Search Tool) (Camacho C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., & Madden T.L. (2008) "BLAST+: architecture and applications." BMC Bioinformatics 10:421 <http://www.ncbi.nlm.nih.gov/pubmed/20003500>) program 'blastn' was then used to find the germline J and V segments used in each sequence. A wordsize of 30 was used for V matching, and 15 for J matching. The database searched against was constructed from the NGS sequencing of the BACs which were used to generate the Kymouse.

If a sequence matched both a V and a J segment, the sequence between the two was then compared to a database of germline D segments in the mouse using 'blastn' with a wordsize of 4 and the options 'blastn-short' and 'ungapped'. This was used to assign a D segment, if possible. The CDR3 was identified by searching for the conserved "TATTACTGT" sequence in the V segment, and the "CTGGGG" in the J segment. If these motifs were not found, then up to 4 mismatches were allowed. The IMGT definition of CDR3 was used, so the CDR3 length is calculated from after the "TGT" in the V to before the "TGG" in the J. Sequences with an out of frame junction (those which do not have a CDR3 nucleotide length divisible by 3) or which contained a stop codon ("TAA", "TAG" or "TGA") were excluded.

The identity of the matching V, J and D segments as well as the CDR3 length from this assignment were then saved as a table for downstream analysis. The ratio of JH6*02 used increased from the naïve to immunised mice, as well as being enriched in the sub-population of sequences with a long HCDR3 (defined as consisting of 20 or more amino acids):

	All		HCDR3>20		% HCDR3>20
	JH6*02%	Total Count	JH6*02%	Total Count	
Naïve	22.31%	1340	91.11%	45	3.36%
Immunised	37.50%	256	66.67%	9	3.52%
Hybridoma	36.13%	119	63.64%	11	9.24%

This shows that the JH6*02 gene segment is selected for by immunisation, as the proportion of JH6*02 usage increases after immunisation. JH6*02 is also used in the majority of antibodies with a long HCDR3 length, which is desirable for targets which are specifically bound by long HCDR3 length antibodies.

WO 2013/041844

SEQ ID NO: 1 (JH5 Reference)

T	T	G	A	C	C	A	A	G	C	T	G	G	G	A	C	C	C
G	G	T	C	C	C	T	T	G	G	G	A	C	C	A	T	G	C
A	G	A	G	G	A	G	T	C									C

JH5 Alignment:

(top line=SEQ ID NO: 1, Middle line=-SEQ ID NO:5, Bottom line=SEQ ID NO:6)

[illegible]

164

[illegible]

TABLES

In the tables, the notation is illustrated by the following example

IGLV1-40	V1-40*02	X53936	g9>c c10>g,L4>V
----------	----------	--------	-----------------

5

Polymorphic variant IGV lambda V1-40*02 has Genbank Accession No. X53936 and when compared to the *01 variant, the V1-40*02 variant has mutations at positions 9, 10 and 4. For example, at position 9, a “C” appears instead of a “G” that is present in the *01 variant. The “|” is simply a notation separator, and does not indicate any mutation. For example the “g282 I” notation indicates no change (ie, position 282 is a g). “del#” means that the residue at that position is absent.

166

10

Table 1: Human IgH V Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

15

--	--	--	--	--

subgroup	IGHV	IGHV	Accession number	Description of mutations
1	IGHV1-2	IGHV1-2*01	X07448	SEQ ID NO: 11
		IGHV1-2*02	X62106	c163>t,R55>W g233>t,t234>g,S78>M t299>c,V100>A
		IGHV1-2*03	X92208	c44>t,P15>L c163>t,R55>W g233>t,t234>g,S78>M t299>c,V100>A
		IGHV1-2*04	Z12310	c163>t,R55>W a223>t,R75>W g233>t,t234>g,S78>M t299>c,V100>A
	IGHV1-3	IGHV1-3*01	X62109	SEQ ID NO: 12 c6 t12 t167 ,I56 a208 ,K70 a291 c296 ,T99
		IGHV1-3*02	X62107	c6>t t12>g t167>g,I56>S a208>g,K70>E a291>g c296>t,T99>M
	IGHV1-8	IGHV1-8*01	M99637	SEQ ID NO: 13
	IGHV1-18	IGHV1-18*01	M99641	g282
		IGHV1-18*02	X60503	g282>a

				SEQ ID NO: 14	
IGHV1-24	IGHV1-24*01	M99642		SEQ ID NO: 15	
				g139 ,G47 t237	
				t237>c	
				g139>a,G47>R t237>c	
IGHV1-45	IGHV1-45*01	X92209		SEQ ID NO: 16	
				c92 ,T31 g315	
				c92>a,T31>N	
				g315>t	
IGHV1-46	IGHV1-46*01	X92343		SEQ ID NO: 17	
				g115 ,V39	
				g115>a,V39>M	
IGHV1-58	IGHV1-58*01	M29809			
IGHV1-58	IGHV1-58*02	AB019438			

IGHV1-69	IGHV1-69*01	L22582	SEQ ID NO: 18
	IGHV1-69*02	Z27506	g6>c g18>a g100>a,A34>T g163>a,G55>R t178>c,F60>L c185>t,T62>I g244>a,E82>K
	IGHV1-69*03	X92340	g291>t,E97>D
	IGHV1-69*04	M83132	g6>c g163>a,G55>R t178>c,F60>L c185>t,T62>I g244>a,E82>K
	IGHV1-69*05	X67905	g6>c g238>a,A80>T
	IGHV1-69*06	L22583	g244>a,E82>K
	IGHV1-69*07	Z29978	g163>a,G55>R
	IGHV1-69*08	Z14309	g6>c g18>a g100>a,A34>T g163>a,G55>R t178>c,F60>L g244>a,E82>K
	IGHV1-69*09	Z14307	g163>a,G55>R t178>c,F60>L c185>t,T62>I g244>a,E82>K
	IGHV1-69*10	Z14300	g6>c g54>a t178>c,F60>L c185>t,T62>I g244>a,E82>K
	IGHV1-	Z14296	g6>c g163>a,G55>R t178>c,F60>L

		69*11			
		IGHV1-69*12	Z14301		g6>c
		IGHV1-69*13	Z14214(st)		g6>c g54>a
	IGHV1-c	IGHV1-c*01	Z18904		SEQ ID NO: 19
		IGHV1-f*01	Z12305		SEQ ID NO: 20 c201
	IGHV1-f	IGHV1-f*02	Z29977		c201>t
		IGHV2-5*01	X62111		SEQ ID NO: 21
	IGHV2-5	IGHV2-5*02	Z14072		a175>g,N59>D
		IGHV2-5*03	X93619		a175>g,N59>D c234>t
		IGHV2-5*04	L21963		c299>g,A100>G c314>t,A105>V a317>g,c318>g,H106>R
		IGHV2-	L21964		a175>g,N59>D a202>g,S68>G

		5*05			
		IGHV2-5*06	L21966	g39>a	a175>g,N59>D a202>g,S68>G
		IGHV2-5*07	L21968	c314>t,A105>V	c299>g,A100>G
		IGHV2-5*08	L21971	a4>g,l2>V a31>g,T11>A	g60>a g103>a,V35>M g106>c,G36>R g118>a,G40>S a175>g,N59>D
		IGHV2-5*09	L21972	a4>g,l2>V G	a175>g,N59>D a202>g,S68>G
		IGHV2-5*10	X69690	a317>g,c318>g,H106>R	a175>g,N59>D
		IGHV2-26*01	M99648	SEQ ID NO: 22	
		IGHV2-70*01	L21969	SEQ ID NO: 23	22 a2 ,Q1 g4 ,V2 g14 ,R5 g31 ,A11 a60 t67 ,C23 a70 ,T24 t10 6 ,C36 t116 ,V39 a138 c157 ,L53 t164 ,L55 a197 ,Y66 t210 g2 16 ,K72 a297 c299 ,A100 a301 ,c302 ,T101 g303 c309 c314 ,A1 05 g317 ,g318 ,R106 t320 ,I107
		IGHV2-70*02	X92241	a297>g	a301>g,c302>t,T101>V

IGHV2-70*03	X92238	g14>a,R5>K t164>g,L55>R a197>t,Y66>F 02>t,T101>V	t106>c,C36>R a297>g a301>g,c3
IGHV2-70*04	Z12330	g14>a,R5>K t164>g,L55>R a197>t,Y66>F	t106>c,C36>R
IGHV2-70*05	Z27502	t164>g,L55>R a197>t,Y66>F	t106>c,C36>R t116>c,V39>A
IGHV2-70*06	X92239	g14>a,R5>K t164>g,L55>R a197>t,Y66>F t210>c g,c302>t,T101>V	t106>c,C36>R a297>g a301>
IGHV2-70*07	X92243	a297>g a301>g,c302>t,T101>V	a138>g
IGHV2-70*08	X92245	a70>g,T24>A a297>g 164>g,L55>R	t a301>g,c302>t,T101>V
IGHV2-70*09	L21962	g4>a,V2>I g14>a,R5>K g31>a,A11>T a60>g t67>c,C23>R G g303>a c314>t,A105>V	g216>c,K72>N c299>g,A100>
IGHV2-70*10	L21965	g14>a,R5>K c157>a,L53>I t164>g,L55>R	t106>c,C36>R
IGHV2-70*11	L21967	a2>g,Q1>R 64>g,L55>R	t1

3			IGHV2-70*12	L21970	g4>a,V2>I g14>a,R5>K g31>a,A11>T a60>g g317a,g318>c,R106>H t320>g,I107>R g303>a
			IGHV2-70*13	AB019437	c309>t
	IGHV3-7		IGHV3-7*01	M99649	SEQ ID NO: 24
			IGHV3-7*02	X92288	g144>a
	IGHV3-9		IGHV3-9*01	M99651	SEQ ID NO: 25
	IGH3-11		IGHV3-11*01	M99652	SEQ ID NO: 26 g13 ,V5 g32 ,G11 g47 ,G16 g178 ,G60 a184 ,c185 ,T62 t188 ,I63 t196 ,Y66 a206 ,D69 g240 c243 ,D81 c296 ,g297 ,T99 g301 ,g303 ,V101 t304 ,a305 ,Y102 t307 ,a308 ,c309 ,Y103 t312 c314 ,A105
			IGHV3-11*02	M15496	9^10>ins^t g32>del#,G11>del# g47>del#,G16>del# a206>del#,D69>del# c243>g,D81>E c296>del#,g297>del#,T99>del# g301>t,g303>a,V101>L t304>c,a305>t,Y102>L t307>a,a308>c,c309>t,Y103>T t312>c c314>a,A105>E

IGHV3-13	IGHV3-11*03	X92287	Y t188>c,l63>T t196>a,Y66>N g13>t,V5>L g178>a,G60>S a184>t,c185>a,T62> g240>a
	IGHV3-13*01	X92217	SEQ ID NO: 27 g9 ,Q3 t52 ,S18 g77 ,S26 g95 ,S32 t165 t167 ,I56 c222 g224 , R75
	IGHV3-13*02	M99653	g9>t,Q3>H t52>g,S18>A g g95>a,S32>N t165>c t167>a,I56>N c222>
	IGHV3-13*03	U29582	c77>g,S26>C g224>a,R75>Q
IGHV3-15	IGHV3-15*01	X92216	SEQ ID NO: 28 g23 ,g24 ,G8 g32 ,G11 a40 ,K14 a81 t89 ,F30 g119 ,S40 t159 c163 ,R55 a169 ,K57 a178 ,T60 g181 ,D61 g196 ,D66 c210 a242 ,D81 a275 ,N92 c279 a297 a320
	IGHV3-15*02	M99654	g32>c,G11>A
	IGHV3-15*03	M99408	g23>c,g24>c,G8>A g32>c,G11>A a40>c,K14>Q a178>g,T60>A g181>a,D61>N >V t89>g,F30>C c210>t a242>t,D81
	IGHV3-15*04	M99402	a169>g,K57>E

	IGHV3-15*05	M99403	c279>t
	IGHV3-15*06	M99404	g196>a,D66>N t159>c
	IGHV3-15*07	M99406	a81>t g119>a,S40>N t159>c
	IGHV3-15*08	M99400	g23>c ,G8>A a40>c,K14>Q t89>g,F30>C c163>t, R55>C a178>g,T60>A g181>a,D61>N c210>t a275 >t,N92>I a297>g a320>g
IGHV3-16	IGHV3-16*01	M99655	SEQ ID NO: 29 a6 a9
	IGHV3-16*02	AB019440	a6>g a9>g
IGHV3-19	IGHV3-19*01	M99656	SEQ ID NO: 30
IGHV3-20	IGHV3-20*01	M99657	SEQ ID NO: 31
IGHV3-21	IGHV3-21*01	AB019439	SEQ ID NO: 32 g9
	IGHV3-	M99658	g9>a

	21*02			
IGHV3-23	IGHV3-23*01	M99660	SEQ ID NO: 33	c164 ,A55 a169 ,g170 ,S57 g172 ,t174 ,G58 a175 ,S59 g181 ,G61 c201 c203 ,A68 c237 c243
	IGHV3-23*02	M35415		c203>g,A68>G c237>a
	IGHV3-23*03	U29481	c164>t,A55>V a169>t,g170>a,S57>Y g172>a,t174>c,G58>S a175>g,S59>G g181>a,G61>S c201>t	c243>t
	IGHV3-23*04	AJ879486	t13>g	
	IGHV3-23*05	AY757302	a154>t, g155>a, S>Y g157>a, t159>a, G>S g166>a, G>S	
IGHV3-30	IGHV3-30*01	M83134	SEQ ID NO: 34	g18 ,E6 a27 a49 ,R17 c75 g80 ,G27 c101 ,t102 ,A34 t135 a138 a150 g163 ,V55 t169 ,c170 ,a171 ,S57 c201 g202 ,A68 a229 ,T77 c257 ,T86 g267 a275 ,N92 t288 a293 ,D98 g317 ,R106
	IGHV3-30*02	L26401		a49>g,R17>G c75>g c101>g,t102>c,A34>G a150 >g g163>t,V55>F t169>c,c170>g,a171>g,S57>R c201>t g317>a,R106>K
	IGHV3-30*03	M99663		c101>g,t102>c,A34>G a150>g c201>t

IGHV3-30*04	L06615			a150>g
IGHV3-30*05	M77323			c101>g,t102>c,A34>G a293>g,D98>G
IGHV3-30*06	L06617			c75>g c101>c,t102>c,A34>G
IGHV3-30*07	L06614			t288>c
IGHV3-30*08	M62737			g18>c,E6>D g80>c,G27>A
IGHV3-30*09	M77300			a229>g,T77>A a150>g
IGHV3-30*10	M77326			g202>a,A68>T
IGHV3-30*11	M77331			c75>g
IGHV3-30*12	M77338			a27>g c75>g c101>g,t102>c,A34>G t288>c
IGHV3-30*13	M77339			c101>g,t102>c,A34>G c257>g,T86>R

	IGHV3-30*14	M77324	g267>t	a150>g
		M77327	a275>g,N92>S	
		M77328	t135>c	
		M77329	a138>g	
		X92214	c101>g,t102>g,A34>G	a150>g
			c201>t	g317>a,R
	IGHV3-30*19	L06616	c75>g	a150>g
		X92283	SEQ ID NO: 35	
			c75 g317 ,R106	
IGHV3-30-3	IGHV3-30-3*01	M77302	c75>g g317>a,R106>K	
	IGHV3-33	AB019439	SEQ ID NO: 36	
			g6 g150 g170 ,g171 ,W57 t177 t212 ,V71 t246 a251 ,K84 a263 ,Y88 g317 ,R106	

	IGHV3-33*02	M99665	g6>a	t212>c,V71>A	a251>c,K84>T a263>t,Y88>F
	IGHV3-33*03	M77305		t246>c	g317>a,R106>K
	IGHV3-33*04	M77335	g150>a	t177>c	
	IGHV3-33*05	M77334		g170>c,g171>a,W57>S	
IGHV3-35	IGHV3-35*01	M99666	SEQ ID NO: 37		
IGHV3-38	IGHV3-38*01	M99669	SEQ ID NO: 38 c302 ,A101		
	IGHV3-38*02	AB019439	c302>t,A101>V		
IGHV3-43	IGHV3-43*01	M99672	SEQ ID NO: 39 t32 ,V11 a100 ,T34 g138 t172 ,W58		
	IGHV3-43*02	Z18901	t32>g,V11>G a100>g,T34>A g138>a t172>g,W58>G		
IGHV3-47	IGHV3-47*01	Z18900	SEQ ID NO: 40 c58 c101 ,A34 t149 ,L50 t267 ,L89 t270 ,H90 t310		

	IGHV3-47*02	AB019438	c58>a c101>t,A34>V t149>c,L50>P t270>a,H90>Q
	IGHV3-47*03	M99674	c58>a c101>t,A34>V t149>c,L50>P t267>del#,L89>del# t270>a,H90>Q t310>g
IGHV3-48	IGHV3-48*01	M99675	SEQ ID NO: 41 g48 c96 a100 ,g101 ,c102 ,S34 a178 ,S60 t246 c287 ,A96 g303
	IGHV3-48*02	AB019438	c287>a,A96>D
	IGHV3-48*03	U03893	g48>a c96>t a100>g,g101>a,c102>a,S34>E a178>g,S60>G t246>c g303>t
IGHV3-49	IGHV3-49*01	M99676	SEQ ID NO: 42 g50 ,R17 t93 g94 ,D32 g100 ,A34 t124 ,F42 a202 ,T68 g245 ,G82
	IGHV3-49*02	M99401	g50>c,R17>P t93>g g94>t,D32>Y g100>c,A34>P t124>g,F42>V a202>g,T68>A g245>a,G82>D
	IGHV3-49*03	AB019438	a202>g,T68>A g245>a,G82>D
	IGHV3-49*04	AM940220	t124>g ,F42>V a202>g,T68>A g245>a,G82>D

		IGHV3-49*05	AM940221	a202>g,T68>A g245>a,G82>D
IGHV3-53		IGHV3-53*01	M99679	t19 ,S7 g133 ,A45 c210 g315 a318
		IGHV3-53*02	Z12342	SEQ ID NO: 43 t19>a,S7>T
		IGHV3-53*03	J03617	g133>c,A45>P c210>t g315>t a318>g
		IGHV3-64*01	M99682	SEQ ID NO: 44 g1 ,E1 g26 ,G9 g70 ,A24 t198 t201 a205 ,N69 t210 c265 ,t267 ,L89 g274 ,g275 ,G92 c279 t296 ,M99 c314 ,A105 g317 ,R106
IGHV3-64		IGHV3-64*02	AB019437	g26>a,G9>E a205>g,N69>D
		IGHV3-64*03	M77298	g70>t,A24>S t198>c t201>c a205>g,N69>D t210>a c265>g,t267>c,L89>V g274>a ,G92>S c279>t t296>c,M99>T c314>t,A105>V g317>a,R106>K
		IGHV3-64*04	M77299	g1>c,E1>Q g70>t,A24>S t198>c t201>c a205>g,N69>D t210>a ,t267>g, g274>a,g275>a,G92>N t296>c,M99>T
		IGHV3-64*05	M77301	g70>t,A24>S t198>c t201>c a205>g,N69>D t210>a c265>g, ,L89>V g274>a ,G92>S c279>t t296>c,M99>T c314>t,A105>V g317>a,R106>K

IGHV3-66	IGHV3-66*01	X92218	SEQ ID NO: 45 g24 g37 ,V13 a81 g175 ,G59 a223 c288 g319
	IGHV3-66*02	Z27504	a223>c c288>t
	IGHV3-66*03	AB019437	g24>a g37>a,V13> a81>g g175>t,G59>C a223>c c288>t
	IGHV3-66*04	X70208	g319>c
IGHV3-72	IGHV3-72*01	X92206	SEQ ID NO: 46 t186
	IGHV3-72*02	Z29979	t186>c
IGHV3-73	IGHV3-73*01	X70197	SEQ ID NO: 47 t21
	IGHV3-73*02	AB019437	t21>c
IGHV3-74	IGHV3-74*01	L33851	SEQ ID NO: 48 c21 g197 ,c198 ,S66
	IGHV3-	Z17392	c21>t

4		74*02			
		IGHV3-74*03	J00239		g197>c,c198>g,S66>T
	IGHV3-d	IGHV3-d*01	Z18898		SEQ ID NO: 49
	IGHV3-h	IGHV3-h*01	AJ879484		SEQ ID NO: 50
		IGHV3-h*02	AJ879485		g303>t
	IGHV4-4	IGHV4-4*01	X05713		SEQ ID NO: 51
		IGHV4-4*02	X92232		c46>t,P16>S >Y g308>a,C103
		IGHV4-4*03	X92252		g308>a,C103>Y
		IGHV4-4*04	X92253		g73>a,V25>I C103>Y g308>a,

	IGHV4-4*05	X92254	c20>t,S7>L g308>a,C103>Y
	IGHV4-4*06	Z75355	a234>g,I78>M a245>c,K82>T g308>a,C103>Y
	IGHV4-4*07	X62112	c46>t,P16>S g50>a,G17>E g70>a,A24>T c93>t a97>t,g98>a, t99>c,S33>Y a100>t,N34>Y t120>c g124>a,V42>I c129>g c136>g,a138>c, P46>A g147>a g163>c,a164>g,a165>t,E55>R c172>a,a173>c,t174>c,H58> T g207>c a234>g,I78>M a245>c,K82>T g308>a,C103>Y a234>g,I78>M a245>c,K82>T g308>a,C103>Y
IGHV4-28	IGHV4-28*01	X05714	SEQ ID NO: 52 g48 g49 ,c51 ,D17 c185 ,T62 g297 c299 ,A100 a319
	IGHV4-28*02	M83133	g48>a g49>c,c51>g,D17>Q c185>t,T62>I
	IGHV4-28*03	X92233	a319>g
	IGHV4-28*04	X56358	g297>c c299>g,A100>G
	IGHV4-28*05	X92260	c185>t,T62>I

IGHV4-30-2	IGHV4-30-2*01	L10089	SEQ ID NO: 53 t163 ,a164 ,c165 ,Y55 c172 ,H58 a237 g245 ,R82 c288 g291 c300 c315 g319
	IGHV4-30-2*02	M95122	c288>t c315>g
	IGHV4-30-2*03	X92229	t163>a,a164>g,c165>t,Y55>S c172>t,H58>Y a237>c g245>c,R82>T c288>t g291>a c300>t c315>g g319>c
	IGHV4-30-2*04	Z75351	g245>c,R82>T g291>a c315>g
IGHV4-30-4	IGHV4-30-4*01	Z14238	SEQ ID NO: 54 g18 ,E6 a48 c49 ,g51 ,Q17 c134 ,P45 a166 ,I56 t228 c288 a291
	IGHV4-30-4*02	Z14239	a48>g c49>g,g51>c,Q17>D c288>a
	IGHV4-30-4*03	X92274	a291>g
	IGHV4-30-4*04	X92275	g18>c,E6>D a166>t,I56>F
	IGHV4-30-4*05	Z75353	t228>c

	IGHV4-30-4*06	Z75360	c134>a,P45>H
IGHV4-31	IGHV4-31*01	L10098	SEQ ID NO: 55 a8 ,Q3 g37 ,V13 c69 c84 g100 ,G34 a164 ,Y55 t224 ,L75 a237 a244 ,c245 ,T82 t249 t285 t285-c287 a295 ,T99 t304 ,Y102
	IGHV4-31*02	M99683	c69>t t224>g,L75>R
	IGHV4-31*03	Z14237	t224>g,L75>R
	IGHV4-31*04	M95120	a8>g,Q3>R t224>g,L75>R
	IGHV4-31*05	M95121	t224>g,L75>R t28 5-c287>del(3nt)# a295>g,T99>A
	IGHV4-31*06	X92270	g100>a,G34>S t224>g,L75>R
	IGHV4-31*07	X92271	c84>a t224>g,L75>R
	IGHV4-31*08	X92272	t224>g,L75>R a237>c t249>c
	IGHV4-	X92273	t224>g,L75>R ,c245>a,T82>K t24

IGHV4-34	31*09		9>c t285>c
	IGHV4-31*10	Z14235	g37>t,V13>L c, T82>P t249>c a164>g,Y55>C t224>g,L75>R a244> t304>g,Y102>D
	IGHV4-34*01	AB019439	SEQ ID NO: 56 a12 g15 c16 ,Q6 g20 ,W7 g25 ,A9 t37 ,L13 g48 g49 ,E17 t85 ,S29 t88 ,F30 a118 ,S40 c129 c137 ,P46 g147 g163 ,a165 ,E55 a169 ,a170 ,t171 ,N57 c172 ,H58 a180 t199 ,Y67 g207 g226 ,t2 27 ,c228 ,V76 a234 ,I78 c249 c263 ,S88 g270 ,K90 a274 ,S92 c2 85 t300 a308 ,Y103
	IGHV4-34*02	M99684	g15>a
	IGHV4-34*03	X92255	t300>c
	IGHV4-34*04	X92236	V76>A t199>a,Y67>N ,t227>c ,
	IGHV4-34*05	X92237	c137>t,P46>L a118>t,S40>C ,t227>c ,V76>A t199>a,Y67>N

188

IGHV4-39	IGHV4-34*12	X56591	a170>t ,N57>I	
	IGHV4-34*13	Z75356	g207>c	
	IGHV4-39*01	AB019439	SEQ ID NO: 57 a2 ,Q1 t56 ,L19 c237 g258 ,Q86 t262 ,S88 a291 t300 c319	
	IGHV4-39*02	X05715	g258>c,Q86>H c319>g	
	IGHV4-39*03	X92259	t300>c	
	IGHV4-39*04	X92297	a291>g	
	IGHV4-39*05	M95116	t56>c,L19>P	
	IGHV4-39*06	Z14236	a2>g,Q1>R c237>a t262>c,S88>P a291>g t300>c	
	IGHV4-39*07	AM940222	c237>a a291>g t300>c c319>g	

IGHV4-55	IGHV4-55*01	M99685	SEQ ID NO: 58 c44 ,P15 c237 a251 ,K84 c255 ,N85 a263 ,Y88 c288 c290 ,A97 g291 t319
	IGHV4-55*02	X92223	c237>a
	IGHV4-55*03	X92263	c237>a a263>c,Y88>S
	IGHV4-55*04	X92265	c44>t,P15>L c237>a
	IGHV4-55*05	X92266	c44>t,P15>L
	IGHV4-55*06	X92267	c255>g,N85>K c288>t
	IGHV4-55*07	X92268	a251>g,K84>R a263>c,Y88>S g291>a
	IGHV4-55*08	X92234	c237>a t319>g
	IGHV4-55*09	X92235	a263>c,Y88>S c290>t,A97>V t319>a

IGHV4-59	IGHV4-59*01	AB019438	SEQ ID NO: 59 g12 g16 ,E6 c20 ,S7 c25 ,P9 g37 g51 ,E17 a70 ,T24 t76 ,c77 , S26 a88 ,I30 c135 c136 ,a138 ,P46 a147 t163 ,a164 ,t165 ,Y55 t172 ,a173 ,c174 ,Y58 a196 ,N66 c207 c228 a234 ,I78 a237 c24 9 g258 c285 t288 g291 c300 g319 a320
	IGHV4-59*02	M29812	a88>g,I30>V
	IGHV4-59*03	M95114	g258>a
	IGHV4-59*04	M95117	,c174>t a196>t,N66>Y c207>g a234>g,I78>M t288>c g291>a c300>t
	IGHV4-59*05	M95118	c135>g ,a138>g t163>c,a164>g, ,Y55>R ,c174>t a196>t,N66>Y c207> g a237>c t288>c g291>a c300>t
	IGHV4-59*06	M95119	t76>a ,S26>T c136>g, a138>t,P46>A a147>c ,t165>c, a196>t,N66>Y c2 07>g c228>t c249>t c285>t t288>c
	IGHV4-59*07	X56360	g51>c,E17>D

	IGHV4-59*08	X87091	g291>a	t288>c
	IGHV4-59*09	Z75359	a320>g	
	IGHV4-59*10	Z14243	g12>a g16>c,E6>Q c20>g,S7>W c25>g,P9>A g37>t	a70>g,T24>A ,c77>a,S26>Y c136>g,a138>c,P46>A a147>g t163>c,a164>g, ,Y55>R t172>a,a173>c, Y58>T a234>g,I78>M t288>c g319>t
IGHV4-61	IGHV4-61*01	M29811	SEQ ID NO: 60 g4 ,V2 g48 g49 ,E17 g88 ,V30 g97 ,G33 a100 ,S34 a118 ,S40 c 136 ,a138 ,P46 g162 t163 ,a164 ,Y55 t172 ,a173 ,Y58 c245 ,T82 g258 ,Q86 t288 g289-g291 g291 g315 g319	
	IGHV4-61*02	L10097	g48>a g49>c,E17>Q g88>a,V30>I >c,P46>A t163>c,a164>g,Y55>R t172>a,a173>c,Y58>T t288>c g291>a	c136>g,a138
	IGHV4-61*03	X92230	g258>c,Q86>H	
	IGHV4-61*04	X92250	g289-g291>del(3nt)	g162>a

		IGHV4-61*05	X56356	g4>c,V2>L g88>a,V30>I g97>a,G33>S a118>g,S40>G c245>a,T82>K t288>c	
		IGHV4-61*06	Z75347	t288>c g315>c	
		IGHV4-61*07	Z75348	g319>c	
		IGHV4-61*08	AB019437	a100>g,S34>G	
	IGHV4-b	IGHV4-b*01	Z12367	SEQ ID NO: 61	
		IGHV4-b*02	X56365	g70>a,A24>T	
	5	IGHV5-51	IGHV5-51*01	M99686	SEQ ID NO: 62
			IGHV5-51*02	M18806	t116>c,l39>T c148>t
IGHV5-51*03			X56368	c45>g	
IGHV5-51*04			X56367	c45>g t247>c,S83>P	

		IGHV5-51*05	Z27449	g139>a,G47>R
	IGHV5-a	IGHV5-a*01	X92227	SEQ ID NO: 63 t21 c148 c225 ,H75 c288 ,A96
		IGHV5-a*02	X92279	c148>t c288>del#,A96>del#
		IGHV5-a*03	X56375	t21>c
		IGHV5-a*04	X56376	c225>g,H75>Q
6	IGHV6-1	IGHV6-1*01	X92224/J04097	SEQ ID NO: 64
		IGHV6-1*02	Z14223	a27>g
7	IGHV7-4-1	IGHV7-4-1*01	L10057	SEQ ID NO: 65
		IGHV7-4-1*02	X62110	t274>a,C92>S
		IGHV7-4-1*03	X92290	t274>a,C92>S g278>c,c279>g,S93>T

		IGHV7- 81	IGHV7- 81*01	AB019437	SEQ ID NO: 66
--	--	--------------	-----------------	----------	---------------

Table 2: Human IgH D Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGHD subgro up	IGHD name	Variant name	Genbank Accession number	Sequence
1	IGHD1-1	IGHD1-1*01	X97051	SEQ ID NO: 67
	IGHD1-7	IGHD1-7*01	X13972	SEQ ID NO: 68
	IGHD1-14	IGHD1-14*01	X13972	SEQ ID NO: 69
	IGHD1-20	IGHD1-20*01	X97051	SEQ ID NO: 70
	IGHD1-	IGHD1-	X97051	SEQ ID NO: 71

	26	26*01		
2	IGHD2- 2	IGHD2- 2*01	J00232	SEQ ID NO: 72 AGGATATTGTAGTAGTACCAGCTGCTATGCC
		IGHD2- 2*02	X97051	G>A
		IGHD2- 2*03	M35648	A>T
	IGHD2- 8	IGHD2- 8*01	X13972	SEQ ID NO: 73 AGGATATTGTACTAA
		IGHD2- 8*02	J00233	AA>GG
	IGHD2- 15	IGHD2- 15*01	J00234	SEQ ID NO: 74
	IGHD2- 21	IGHD2- 21*01	J00235	SEQ ID NO: 75 AGCATATTGGTGGTGA
		IGHD2-	X97051	T>C

	21*02			
3	IGHD3-3*01	X13972	SEQ ID NO: 76 GTATTACGATTTTGGAGTGGTTATTATACC	
			CG>GC	
	IGHD3-3*02	X93618		
	IGHD3-9*01	X13972	SEQ ID NO: 77	
	IGHD3-10*01	X13972	SEQ ID NO: 78 GTATTACTATGGTTCGGGGGAGTTATTATAAC	
	IGHD3-10*02	X93615	G>[OMITTED]	
	IGHD3-16*01	X93614	SEQ ID NO: 79 GTATTATGATTACGTTTGGGGGAGTTATGCTTATACC	
	IGHD3-16*02	X97051	GC>CG	
	IGHD3-	X93616	SEQ ID NO: 80	

	22	22*01		
4	IGHD4-4	IGHD4-4*01	X13972	SEQ ID NO: 81
	IGHD4-11	D4-11*01	X13972	SEQ ID NO: 82
	IGHD4-17	IGHD4-17*01	X97051	SEQ ID NO: 83
	IGHD4-23	IGHD4-23*01	X97051	SEQ ID NO: 84
5	IGHD5-5	IGHD5-5*01	X13972	SEQ ID NO: 85
	IGHD5-12	IGHD5-12*01	X13972	SEQ ID NO: 86
	IGHD5-18	IGHD5-18*01	X97051</A<	SEQ ID NO: 87
	IGHD5-24	IGHD5-24*01	X97051	SEQ ID NO: 88
6	IGHD6-6	IGHD6-6*01	X13972	SEQ ID NO: 89

	IGHD6-13	IGHD6-13*01	X13972	SEQ ID NO: 90
	IGHD6-19	IGHD6-19*01	X97051	SEQ ID NO: 91
	IGHD6-25	IGHD6-25*01	X97051	SEQ ID NO: 92
	IGHD7-27	IGHD7-27*01	J00256	SEQ ID NO: 93
7				

Underlined nucleotides in the *01 variant are positions that are different in other variants, eg, GC>CG for IGH3-16*02 indicates that GC in the *01 variant is instead CG in the *02 variant. All other positions are identical to between the *01 and other variant. Similar notation is used in other tables below to denote changes between variants of gene segments.

Table 3: Human IgH J Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGHV name	Variant name	Genbank Accession number	Sequence
IGHJ1	J1*01	J00256	SEQ ID NO: 94
IGHJ2	J2*01	J00256	SEQ ID NO: 95
IGHJ3	J3*01	J00256	SEQ ID NO: 96 T GAT GCT TTT GAT <u>G</u> TCT TGG GGC CAA GGG ACA ATG GTC ACC GTC TCT TCA G
	J3*02	X86355	<u>G</u> >A
IGHJ4	J4*01	J00256	SEQ ID NO: 97 <u>AC</u> TAC TTT GAC TAC TGG GGC CAA <u>GGA</u> ACC CTG GTC ACC GTC TCC TCA G

	J4*02	X86355	<u>AA</u> >AG	
	J4*03	M25625	<u>AC</u> >GC <u>GA</u> >GG	
IGHJ5	J5*01	J00256	SEQ ID NO: 98 AC AAC TGG TTC GAC <u>T</u> CC TGG GGC CAA <u>A</u> GGA ACC CTG GTC ACC GTC TCC TCA G	
	J5*02	X86355	<u>T</u> >C <u>A</u> >G	
IGHJ6	J6*01	J00256	SEQ ID NO: 99 AT TAC TAC TAC TAC <u>G</u> GT ATG GAC GTC TGG <u>G</u> GG <u>CAA</u> GGG ACC ACG GTC ACC GTC TCC TCA G	
	J6*02	X86355	<u>GG</u> >GC	
	J6*03	X86356	<u>GGT</u> >TAC <u>GG</u> >GC	

				CA> <u>AA</u>
	J6*04	AJ8794887		<u>GG>GC</u> CA> <u>AA</u>

Table 4: Human Ig Vk Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGKV subgroup	IGKV name	Variant name	Accession number	Sequence
1	IGKV1-5	IGKV1-5*01	Z00001	SEQ ID NO: 100 GAC ATC CAG ATG ACC CAG TCT CCT TCC ACC CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC <u>ACT</u> TGC CGG GCC AGT CAG AGT ATT AGT AGC TGG TTG GCC TGG TAT CAG CAG AAA CCA GGG AAA GCC CCT AAG

				CTC CTG ATC TAT <u>GAT</u> GCC TCC AGT TTG GAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAT GAT TTT GCA ACT TAT TAC TGC CAA CAG TAT AAT AGT TAT TCT CC
	IGKV1-5*02	M23851		<u>AC</u> >AT
	IGKV1-5*03	X72813		<u>GAT</u> >AAG <u>GCC</u> >GCG <u>TCC</u> >TCT <u>TTG</u> >TTA
IGKV1-6	IGKV1-6*01	M64858		SEQ ID NO: 101
IGKV1-8	IGKV1-8*01	Z00014		SEQ ID NO: 102
IGKV1D-8	IGKV1D-8*01	Z00008		SEQ ID NO: 103
IGKV1-9	IGKV1-9*01	Z00013		SEQ ID NO: 104
IGKV1-12	IGKV1-	V01577		SEQ ID NO: 105

		12*01		GAC ATC CAG ATG ACC CAG TCT CCA TCT <u>TCC</u> GTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGT CGG GCG AGT CAG GGT ATT AGC AGC TGG TTA GCC TGG TAT CAG CAG AAA CCA GGG AAA GCC CCT AAG CTC CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC <u>ACC</u> ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAC TAT TGT CAA CAG GCT AAC AGT TTC CCT <u>CC</u>
	IGKV1D-12	IGKV1D-12*01	X17263	<u>TCC</u> >TCT <u>ACC</u> >ACT
	IGKV1-12/IGKV1D-12 (1)	IGKV1-12*02/1D-12*02	V01576	<u>CC</u> >TC
	IGKV1-13	IGKV1-13*01	Z00010	SEQ ID NO: 106 GCC ATC CAG TTG ACC CAG TCT CCA TCC TCC CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGC CGG GCA AGT CAG GGC ATT AGC AGT GCT TTA GCC <u>IGA</u> TAT CAG CAG AAA CCA GGG AAA GCT CCT AAG CTC CTG ATC TAT GAT GCC TCC AGT TTG GAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGT CAA CAG TTT AAT <u>AAT</u> TAC CCT CA

	IGKV1-13*02	Z00006			TGA><u>TGG</u> AAT><u>AGT</u>
IGKV1D-13	IGKV1D-13*01	X17262			TGA><u>TGG</u>
	IGKV1-16*01	J00248			SEQ ID NO: 107 GAC ATC CAG ATG ACC CAG TCT CCA TCC TCA CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGT CGG GCG AGT CAG GGC ATT AGC AAT TAT TTA GCC TGG TTT CAG CAG AAA CCA GGG AAA GCC CCT AAG TCC CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA <u>AGG</u> TTC AGC GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGC CAA CAG TAT AAT AGT TAC CCT CC
IGKV1D-16	IGKV1-16*02	FM164406			<u>AGG</u>>AAG
	IGKV1D-16*01	K01323			SEQ ID NO: 108 GAC ATC CAG ATG ACC CAG TCT CCA TCC TCA CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGT CGG GCG <u>AGT</u> CAG GGT ATT AGC AGC TGG TTA GCC TGG TAT CAG CAG AAA CCA GAG AAA GCC CCT AAG TCC CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC

				TGC CAA CAG TAT AAT AGT TAC CCT CC
	IGKV1D-16*02	V00558		<u>AGT</u> >AGG
	IGKV1-17	IGKV1-17*01	X72808	SEQ ID NO: 109 GAC ATC CAG ATG ACC CAG TCT CCA TCC TCC CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGC CGG GCA AGT CAG GGC ATT AGA AAT GAT TTA GGC TGG TAT CAG CAG AAA CCA GGG AAA GCC CCT AAG CGC CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC ACA ATC AGC <u>AGC</u> CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGT CTA CAG CAT AAT AGT TAC CCT CC
		IGKV1-17*02	D88255	<u>AGC</u> > AAC
IGKV1D-17	IGKV1D-17*01	X63392		SEQ ID NO: 110 <u>AAC</u> ATC CAG ATG ACC CAG TCT CCA TCT GCC ATG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGT CGG GCG AGG CAG GGC ATT AGC AAT TAT TTA GCC TGG TTT CAG CAG AAA CCA GGG AAA GTC CCT AAG <u>CAC</u> CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA AGG TTC AGC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC ACA ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGT CTA CAG CAT AAT AGT TAC CCT CC
	IGKV1D-	FM164407		<u>AAC</u> >GAC

				17*02			<u>CAC</u> >CGC
IGKV1-27	IGKV1-27*01	X63398	SEQ ID NO: 111				
IGKV1-33	IGKV1-33*01	M64856	SEQ ID NO: 112				
IGKV1D-33	IGKV1D-33*01	M64855	SEQ ID NO: 113				
IGKV1-37	IGKV1-37*01	X59316	SEQ ID NO: 114				
IGKV1D-37	IGKV1D-37*01	X71893	SEQ ID NO: 115				
IGKV1-39	IGKV1-39*01	X59315	SEQ ID NO: 116			GAC ATC CAG ATG ACC CAG TCT CCA TCC <u>TCC</u> CTG TCT GCA TCT GTA GGA GAC AGA GTC ACC ATC ACT TGC CGG GCA AGT CAG AGC ATT AGC AGC TAT TTA AAT TGG TAT CAG CAG AAA CCA GGG AAA GCC CCT AAG CTC CTG ATC TAT GCT GCA TCC AGT TTG CAA AGT GGG GTC CCA ... TCA AGG TTC AGT GGC AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGT CTG CAA CCT GAA GAT TTT GCA ACT <u>TAC</u> TAC TGT <u>CAA CAG AGT TAC</u> AGT <u>ACC</u> CCT CC	
	IGKV1-39*02	X59318				<u>TCC</u> >TTC <u>TAC</u> >TAT	

2					<u>CAA CAG AGT</u> > CAG TGT GGT <u>ACC</u> >ACA
	IGKV1D-39	IGKV1D-39*01	X59312	SEQ ID NO: 117	
	IGKV1D-42	IGKV1D-42*01	X72816	SEQ ID NO: 118	
	IGKV1D-43	IGKV1D-43*01	X72817	SEQ ID NO: 119	
	IGKV2-24	IGKV2-24*01	X12684	SEQ ID NO: 120	
	IGKV2D-24	IGKV2D-24*01	X63401	SEQ ID NO: 121	
	IGKV2-28	IGKV2-28*01	X63397	SEQ ID NO: 122	
	IGKV2D-28	IGKV2D-28*01	X12691	SEQ ID NO: 123	
	IGKV2-29	IGKV2-29*01	X63396	SEQ ID NO: 124	GAT ATT GTG ATG ACC CAG ACT CCA CTC TCT CTG TCC GTC ACC CCT GGA CAG CCG GCC TCC ATC TCC TGC AAG TCT AGT CAG AGC CTC CTG CAT AGT GAT GGA AAG ACC TAT ... TTG TAT TGG TAC CTG CAG AAG CCA GGC CAG

				TCT CCA CAG CTC <u>CTG</u> ATC TAT GAA GTTTTCC AGC CGG TTC TCT GGA GTG CCA ... GAT AGG TTC AGT GGC AGC GGG TCA GGG ACA GAT TTC ACA CTG AAA ATC AGC CGG GTG GAG GCT GAG GAT GTT GGG GTT TAT TAC <u>TGA</u> ATG CAA GGT ATA CAC CTT CCT CC
IGKV2D-29	IGKV2-29*02	U41645	<u>CTG</u> > CTA <u>TGA</u> > TGC	
	IGKV2-29*03	AJ783437	<u>TGA</u> > TGC	
	IGKV2D-29*01	M31952	SEQ ID NO: 125 GAT ATT GTG ATG ACC CAG ACT CCA CTC TCT CTG TCC GTC ACC CCT GGA CAG CCG GCC TCC ATC TCC TGC AAG TCT AGT CAG AGC CTC CTG CAT AGT GAT GGA AAG ACC TAT ... TTG TAT TGG TAC CTG CAG AAG CCA GGC CAG <u>CCT</u> CCA CAG CTC CTG ATC TAT GAA GTT TCC AAC CGG TTC TCT GGA GTG CCA ... GAT AGG TTC AGT GGC AGC GGG TCA GGG ACA GAT TTC ACA CTG AAA ATC AGC CGG GTG GAG GCT GAG GAT GTT GGG GTT TAT TAC TGC ATG CAA AGT ATA CAG CTT CCT CC	
	IGKV2D-29*02	U41644	<u>CCT</u> >TCT	
IGKV2-30	IGKV2-30*01	X63403	SEQ ID NO: 126 GAT GTT GTG ATG ACT CAG TCT CCA CTC TCC CTG CCC GTC ACC CTT GGA CAG CCG GCC TCC ATC TCC TGC AGG TCT AGT CAA AGC CTC GTA <u>TAC</u> AGT GAT GGA AAC ACC TAC ... TTG AAT TGG TTT CAG CAG AGG CCA GGC CAA	

				TCT CCA AGG CGC CTA ATT TAT AAG GTT TCT AAC CGG GAC TCT GGG GTC CCA ... GAC AGA TTC AGC GGC AGT GGG TCA GGC ACT GAT TTC ACA CTG AAA ATC AGC AGG GTG GAG GCT GAG GAT GTT GGG GTT TAT TAC TGC ATG CAA GGT ACA CAC TGG CCT CC
	IGKV2- 30*02	FM164408		<u>TAC</u> >CAC
	IGKV2D- 30*01	X63402		SEQ ID NO: 127
	IGKV2- 40*01	X59314		SEQ ID NO: 128 GAT ATT GTG ATG ACC CAG ACT CCA CTC TCC CTG CCC GTC ACC CCT GGA GAG CCG GCC TCC ATC TCC TGC AGG TCT AGT CAG AGC CTC TTG GAT AGT GAT GAT GGA AAC ACC TAT TTG <u>GAC TGG</u> TAC CTG CAG AAG CCA GGG CAG TCT CCA CAG CTC CTG ATC TAT ACG CTT TCC TAT CGG GCC TCT GGA GTC CCA ... GAC AGG TTC AGT <u>GGC</u> AGT GGG TCA GGC ACT GAT TTC ACA CTG AAA ATC AGC AGG GTG GAG GCT GAG GAT GTT GGA GTT TAT TAC TGC ATG CAA CGT ATA GAG TTT CCT TC
	IGKV2- 40*02	X59317		<u>GAC TGG</u> >GAT TGT <u>GGC</u> >GAC
	IGKV2D-40			SEQ ID NO: 129
	IGKV2D- 40*01	X59311		

3	IGKV3-7	IGKV3-7*01	X02725	SEQ ID NO: 130 GAA ATT GTA ATG ACA CAG TCT CCA CCC ACC CTG TCT TTG TCT CCA GGG GAA AGA GTC ACC CTC TCC TGC AGG GCC AGT CAG AGT GTT AGC AGC AGC TAC TTA ACC TGG TAT CAG CAG AAA CCT GGC CAG GCG CCC AGG CTC CTC ATC TAT GGT GCA TCC ACC AGG GCC ACT AGC ATC CCA ... GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA GTT TAT TAC TGT CAG CAG GAT CAT AAC TTA CCT CC
		IGKV3-7*02	X72812	<u>ACC>TCC</u> <u>TAT>TAC</u> <u>GGC>GGG</u> <u>GCG>GCT</u> <u>AGC>GGC</u> <u>ACA>AGA</u> <u>CAT>TAT</u>
		IGKV3-7*03	K02769	SEQ ID NO: 131 GAA ATT GTA ATG ACA CAG TCT CCA CCC ACC CTG TCT TTG TCT CCA GGG GAA AGA GTC ACC CTC TCC TGC AGG GCC AGT CAG AGT GTT AGC AGC AGC TAC TTA ACC TGG TAT CAG CAG AAA CCT GGC CAG GCG CCC AGG CTC CTC ATC TAT GGT GCA TCC ACC AGG GCC ACT AGC ATC CCA ... GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACT CTC ACC ATC AGC AGC CTG CAG CCT GAA GAT TTT GCA GTT TAT

TAC TGT CAG CAG GAT CAT AAC TTA CCT CC							
IGKV3D-7	IGKV3D-7*01	X72820	SEQ ID NO: 132	FM164409	IGKV3-7*04		CAT>TAT
IGKV3D-11	IGKV3-11*01	X01668	SEQ ID NO: 133				GAA ATT GTG TTG ACA CAG TCT CCA GCC ACC CTG TCT TTG TCT CCA GGG GAA AGA GCC ACC CTC TCC TGC AGG GCC AGT CAG AGT GTT AGC AGC TAC TTA GCC TGG TAC CAA CAG AAA CCT GGC CAG GCT CCC AGG CTC CTC ATC TAT GAT GCA TCC AAC AGG GCC ACT GGC ATC CCA ... GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACT CTC ACC ATC AGC AGC CTA GAG CCT GAA GAT TTT GCA GTT TAT TAC TGT CAG CAG CGT AGC AAC TGG CCT CC
				K02768	IGKV3-11*02		ACA>AGA
IGKV3D-11	IGKV3D-11*01	X17264	SEQ ID NO: 134				
IGKV3-15	IGKV3-15*01	M23090	SEQ ID NO: 135				
IGKV3D-15	IGKV3D-	X72815	SEQ ID NO: 136				

	15*01		GAA ATA GTG ATG <u>ACG</u> CAG TCT CCA GCC ACC CTG TCT GTG TCT CCA GGG GAA AGA GCC ACC CTC TCC TGC AGG GCC AGT CAG AGT GTT AGC AGC AAC TTA GCC TGG TAC CAG CAG AAA CCT GGC CAG GCT CCC AGG CTC CTC ATC TAT GGT GCA TCC ACC AGG GCC ACT GGC ATC CCA ... GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAG TTC ACT CTC ACC ATC AGC AGC CTG CAG TCT GAA GAT TTT GCA GTT TAT TAC TGT CAG CAG TAT AAT AAC <u>TGG</u> CCT CC
		M23091	<u>ACG</u> >ATG <u>TGG</u> >TGA
	IGKV3D- 15*02		
	IGKV3- 20*01	X12686	SEQ ID NO: 137 GAA ATT GTG TTG <u>ACG</u> CAG TCT CCA GGC <u>ACC</u> CTG TCT TTG TCT CCA GGG GAA AGA GCC ACC CTC TCC TGC AGG GCC AGT CAG AGT GTT AGC AGC AGC TAC TTA GCC TGG TAC CAG CAG AAA CCT GGC CAG GCT CCC AGG CTC CTC ATC TAT GGT GCA TCC AGC AGG GCC ACT GGC ATC CCA ... <u>GAC</u> AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACT CTC ACC ATC AGC AGA CTG GAG CCT GAA GAT TTT GCA GTG TAT TAC TGT CAG CAG TAT GGT AGC TCA CCT CC
IGKV3-20	IGKV3- 20*01, Tou-kv325	X56593	<u>ACC</u> >CCC <u>CTG</u> >TGT <u>TCT</u> > CTT <u>TTG</u> > TGT <u>TCT</u> > CTC

					IGKV3-20*02	L37729	ACG>ACA GGC>GCC GAC>GCA
4		IGKV3D-20	IGKV3D-20*01		X12687	SEQ ID NO: 138	
					Z00023		
					X02485		
5		IGKV5-2	IGKV5-2*01		X63399	SEQ ID NO: 141	
6		IGKV6D-21	IGKV6D-21*01		X12683	SEQ ID NO: 142	
					M27751 (3)	SEQ ID NO: 143	

Table 5: Human Ig Vλ Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGLV subgroup	IGLV Gene name	IGLV variant name	Accession number	Sequences & Description of mutations
1	IGLV1-36	V1-36*01	Z73653	SEQ ID NO: 144
	IGLV1-40	V1-40*01	M94116	SEQ ID NO: 145
		V1-40*02	X53936	g9>c c10>g,L4>V
		V1-40*03	Z22192	g9>c c10>g,L4>V a253>g,T85>A
	IGLV1-41	V1-41*01	M94118	SEQ ID NO: 146
		V1-41*02	D87010	g295>t,E99>* c332>t,P111>L

	IGLV1-44	V1-44*01	Z73654	SEQ ID NO: 147		
	IGLV1-47	V1-47*01	Z73663	SEQ ID NO: 148		
		V1-47*02	D87016	g168>t,R56>S		
	IGLV1-50	V1-50*01	M94112	SEQ ID NO: 149		
	IGLV1-51	V1-51*01	Z73661	SEQ ID NO: 150		
		V1-51*02	M30446	t162>c c168>a,D56>E		
	IGLV2-8	V2-8*01	X97462	SEQ ID NO: 151		
		V2-8*02	L27695	g37>a,G13>R		
	2					

		V2-8*03	Y12418	c230>t,S77>F		
IGLV2-11		V2-11*01	Z73657	SEQ ID NO: 152		
		V2-11*02	Z22198	t96>g		
		V2-11*03	Y12415	g132>a		
IGLV2-14		V2-14*01	Z73664	SEQ ID NO: 153		
		V2-14*02	L27822	c87>t t93>g g94>a,G32>S t103>c,a104>t,Y35>L t170>g,V57>G t198>g,N66>K		
		V2-14*03	Y12412	g132>a g168>t,E56>D		
		V2-14*04	Y12413	g168>t,E56>D		
		V2-18*01	Z73642	SEQ ID NO: 154		

18				
	V2-18*02	L27697	t317>c,L106>S	
	V2-18*03	L27694	t272>c,I91>T t317>c,L106>S	
	V2-18*04	L27692	t227>c,F76>S t249>c t317>c,L106>S	
IGLV2- 23	V2-23*01	X14616	SEQ ID NO: 155	
	V2-23*02	Z73665	g170>t,G57>V a339>c,L113>F	
	V2-23*03	D86994	a339>c,L113>F	
IGLV2- 33	V2-33*01	Z73643	SEQ ID NO: 156	
	V2-33*02	L27823	a3>g	

		V2-33*03	L27691	t96>c a256>g,M86>V	
3	IGLV3-1	V3-1*01	X57826	SEQ ID NO: 157	
	IGLV3-9	V3-9*01	X97473	a52 ,T18 a82 ,l28 a88 ,g89 ,S30 a95 ,N32 g173 ,S58	
		V3-9*02	X74288	a52>g,T18>A a82>c,l28>L a88>t,g89>a,S30>Y a95>g ,N32>S g173>a,S58>N	
		V3-9*03	X51754	a52>g,T18>A a82>c,l28>L a88>t,g89>a,S30>Y a95>del#,N32>del# g173>a,S58>N	
	IGLV3-10	V3-10*01	X97464	SEQ ID NO: 158	
		V3-10*02	L29166	g166>a,E56>K c207>a t270>c c299>a,A100>D a319>g,T107>A a325> t,g326>a,S109>Y	
	IGLV3-12	V3-12*01	Z73658	SEQ ID NO: 159	
		V3-12*02	D86998	a259>g,T87>A	

IGLV3-16	V3-16*01	X97471	SEQ ID NO: 160		
IGLV3-19	V3-19*01	X56178	SEQ ID NO: 161		
IGLV3-21	V3-21*01	X71966	SEQ ID NO: 162		
	V3-21*02	D87007	a27>g a49>c,K17>Q a160>g,l54>V t166>g,Y56>D c324>t		
IGLV3-22	V3-21*03	M94115	a27>g a160>g,l54>V t166>g,Y56>D c324>t		
	V3-22*01	Z73666	SEQ ID NO: 163		
IGLV3-25	V3-22*02	X71967	c53>a,T18>K g88>a,E30>K g140>del#,G47>del# c148>t,c149>g,t150>a,P50>* 150^151>ins^tatac#,50^51>ins^Y# g322>a,D108>N c330>t		
	V3-25*01	X97474	SEQ ID NO: 164		

		V3-25*02	D86994	t14>c,g15>a,M5>T	
		V3-25*03	L29165	t14>c,g15>a,M5>T t294>c	
	IGLV3-27	V3-27*01	D86994	SEQ ID NO: 165	
	IGLV3-32	V3-32*01	Z73645	SEQ ID NO: 166	
	IGLV4-3	V4-3*01	X57828	SEQ ID NO: 167	
	IGLV4-60	V4-60*01	Z73667	SEQ ID NO: 168	
		V4-60*02	D87000	a288>t,L96>F	
	IGLV4-	V4-60*03	AF073885	t287>c,a288>t,L96>S	
		V4-69*01	Z73648	SEQ ID NO: 169	

	69		V4-69*02		U03868	c198>t	
5	IGLV5-37		V5-37*01		Z73672	SEQ ID NO: 170	
	IGLV5-39		V5-39*01		Z73668	t81 ,g135>t	
			V5-39*02		AF216776	t81>c,g135>t	
	IGLV5-45		V5-45*01		Z73670	SEQ ID NO: 171	
			V5-45*02		Z73671	g22>t,A8>S g75>a	
			V5-45*03		D86999	g22>t,A8>S	
	IGLV5-48		V5-48*01		Z73649	SEQ ID NO: 172	

	IGLV5-52	V5-52*01	Z73669	SEQ ID NO: 173	
6	IGLV6-57	V6-57*01	Z73673	SEQ ID NO: 174	
7	IGLV7-43	V7-43*01	X14614	SEQ ID NO: 175	
	IGLV7-46	V7-46*01	Z73674	SEQ ID NO: 176	
		V7-46*02	D86999	c275>t,\$92>L	
8	IGLV8-61	V7-46*03	Z22210	c271>del# L91>del#	
		V8-61*01	Z73650	SEQ ID NO: 177	
		V8-61*02	U03637	c223>t,R75>C	
		V8-61*03	AF266511	g210>c	

9	IGLV9-49	V9-49*01	Z73675	SEQ ID NO: 178	
		V9-49*02	Z73675	g291>a	
		V9-49*03	U03869	g153>a	
10	IGLV10-54	V10-54*01	Z73676	SEQ ID NO: 179	
		V10-54*02	D86996	a86>t,N29>I a228>c,L76>F g320>t,W107>L	
		V10-54*03	S70116	g33>c t125>c,g126>t,L42>P c127>g,Q43>E	
11	IGLV11-55	V11-55*01	D86996	SEQ ID NO: 180	

Table 6: Human IgH Jk Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGKJ gene name	Variant name	Accession number	Sequence
IGKJ1	IGKJ1*01	J00242	SEQ ID NO: 181
	IGKJ2*01	J00242	SEQ ID NO: 182
IGKJ2	IGKJ2*02	Z70260 #c	TG TAC <u>ACT</u> TTT GGC CAG GGG ACC AAG CTG GAG ATC AAA C
	IGKJ2*03	U95246 #c	<u>TAC</u> >TGC
			<u>ACT</u> >AGT

	IGKJ2*04		L40735 #c		TAC> <u>TGC</u> ACT> <u>AGT</u>	
IGKJ3	IGKJ3*01		J00242		SEQ ID NO: 183	
IGKJ4	IGKJ4*01		J00242		SEQ ID NO: 184 G CTC <u>ACT</u> TTC GGC GGA GGG ACC AAG GTG GAG ATC AAA C	
	IGKJ4*02		AF103571 #c		<u>ACT</u> >ACG	
IGKJ5	IGKJ5*01		J00242		SEQ ID NO: 185	

Table 7: Human IgH Jλ Polymorphic Variants

All variant sequences appearing in Genbank under the listed Accession Numbers are incorporated herein in their entirety by reference as though explicitly recited herein and for possible inclusion in any of the claims herein.

IGLJ gene name	Variant name	Accession number	Sequence
IGLJ1	J1*01	X04457	SEQ ID NO: 186
IGLJ2	J2*01	M15641	SEQ ID NO: 187
IGLJ3	J3*01	M15642	SEQ ID NO: 188 T <u>GTG</u> <u>GTA</u> TTC GGC GGA GGG ACC AAG CTG ACC GTC CTA G
	J3*02	D87023(1)	<u>GTG</u> > TGG <u>GTA</u> > GTG
IGLJ4	J4*01	X51755	SEQ ID NO: 189
IGLJ5	J5*01	X51755	SEQ ID NO: 190 C TGG GTG TTT GGT GAG GGG <u>ACC</u> GAG CTG ACC GTC CTA G
	J5*02	D87017	<u>ACC</u> >ACG

IGLJ6	J6*01	M18338	SEQ ID NO: 191
IGLJ7	J7*01	X51755	SEQ ID NO: 192 T GCT GTG TTC GGA GGA GGC ACC CAG CTG ACC <u>GTC</u> CTC G
	J7*02	D87017	<u>GTC</u> >GCC

TABLE 8: 1000 GENOMES PROJECT HUMAN POPULATIONS

Below is a summary of the ethnic population origin of samples that the 1000 Genomes Project sequences.

Population

European ancestry

Utah residents (CEPH) with Northern and Western European ancestry (CEU)

Toscani in Italia (TSI)

British from England and Scotland (GBR)

Finnish from Finland (FIN)

Iberian populations in Spain (IBS)

East Asian ancestry

Han Chinese in Beijing, China (CHB)

Japanese in Toyko, Japan (JPT)

Han Chinese South (CHS)

Chinese Dai in Xishuangbanna (CDX)

Kinh in Ho Chi Minh City, Vietnam (KHV)

Chinese in Denver, Colorado (CHD) (pilot 3 only)

West African ancestry

Yoruba in Ibadan, Nigeria (YRI)

Luhya in Webuye, Kenya (LWK)

Gambian in Western Division, The Gambia (GWD)

Malawian in Blantyre, Malawi (MAB)

K00004-1 WO

Population

West African Population (TBD)

Americas

African Ancestry in Southwest US (ASW)

African American in Jackson, MS (AJM)

African Caribbean in Barbados (ACB)

Mexican Ancestry in Los Angeles, CA (MXL)

Puerto Rican in Puerto Rico (PUR)

Colombian in Medellin, Colombia (CLM)

Peruvian in Lima, Peru (PEL)

South Asian ancestry

Ahom in the State of Assam, India

Kayadtha in Calcutta, India

Reddy in Hyderabad, India

Maratha in Bombay, India

Punjabi in Lahore, Pakistan

[illegible]

• LSG6.1, LSG12.1, V_HIII VH26, V_HIII 9.1 • <u>VH and VL COMBINATIONS</u> • V_HIII 9.1 + V_λVII 4A • V_HIII 9.1 + V_λII 2.1 • V_HIII 9.1 + V_κII A2 • V_HIII VH26 + V_λII 2.1				
• V_HIII 9.1; V_HIII H11; V_HIII VH26 • κI 15A • Vλ2.1	• Polysaccharide capsule of E coli K1 • Meningococcal B polysaccharide; Poly[α(2→8)-N-acetylneuramic acid	• E coli K1 • Neisseria meningitidis Group B	9. Azmi et al, Infect Immun 1994	
• VHIII or VHIV family member • VλI or Vλ3 member • VH26 + Dk1 + JH6 with IGLV1S2 + Jλ2 • VH4.18 • VH2-1 (VH3) + D region Dxp'1 + JH5 with Vλ3 cML70 + Jλ3 • VH1GRR+ JH3 + Dn4r or D2r with IGLV1S2 + Jλ2	• HSV 120-kD glycoprotein • 116-, 105-, 64-kD glycoproteins of VZV	• Herpes family virus • Herpes simplex virus (HSV); HSV-1; HSV-2 • Varicella zoster virus (VZV)	10. Huang et al, J Clin Invest 1992; [human tonsils]	

- For VZV Abs: ha3h2 (VH3) with la1h2 (Vλ); or ha1c1 (VH1) with la1v1 (Vλ1) - For VZV Abs: ha4h3 (VH4) with la3h3 (Vλ3)				
- Hv1051 (VH) - Kv325 (Vk)			Cytomegalovirus (CMV)	10. Huang et al, J Clin Invest 1992;
71-2 (VH) - Hv1f10 (VH) - VH4.11 71-4 (VH) - VH251 - VH1-69			HIV	10. Huang et al, J Clin Invest 1992; 11. Wang & Palese, Science 2011
VH1-69	Haemagglutinin (HA)		Influenza virus, eg, Group 1 and/or Group 2 Influenza A virus; eg, H1N1, H2N2, or H3N2 or H7N2 or H7N7 influenza virus	12. Ekiert et al, Science 2009 13. Throsby et al, PLoS One 2008 14. Sui et al, Nat Struct Mol Biol 2009 15. Ekiert et al, Science 2011

Table 10A:Human IgH JH5 Variant Occurrences

HUMAN POPULATION														
NA19835	NA12814; NA12763 and 1 other(s)	NA18617; NA18636 and 2 other(s)	HG00683; HG00590	HG01359	HG00189	HG00261; HG00136	NA18987	NA19448; NA19331			HG00740	NA20799	0.017	
	NA11994		HG00595; HG00584		HG00267; HG00276	HG00137		NA19456; NA19332				NA20542	NA18934; NA19147 and 1 other(s)	0.011
				HG01494	HG00351	HG00234	NA19005		NA19719				0.005	
NA19921	NA12815		HG00650		HG00177							NA18517	0.005	
	NA12829; NA12890 and 1 other(s)				HG00358							NA20503	NA19093	0.005

236

237

674884000000TS IE	A/S GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	D/V GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	Q/L GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	I/T GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	A/T GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	Q/D GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	H/D GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	C/F GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	L/F GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	C/W GN DDC S UO W NONS _NON	1	1	
674884000000TS IE	*C/W DE MAG POS	1	1	
674884000000TS IE	T/N GN DDC S UO W NONS _NON	1	1	
PYT T NA RAV				

239

	A A	A T	A A	G G	T T	T T	C C	A A	C C	C C	C C	T T
	A A	A A	T A	G G	T T	T T	C C	A A	C C	C C	C C	T T
	A A	A A	A A	G G	T T	T T	C C	A A	C G	C C	C C	T T
	A A	A A	A A	G G	T T	T T	C C	A A	G C	C C	C C	T T
	A A	A A	A A	G G	T T	T T	C C	A A	C C	C C	C C	T G
	A A	A A	A A	G G	T T	T T	C C	A A	C G	C C	C C	T T
	A A	A A	A A	G G	T T	T T	C C	A A	C G	C C	C C	T G
	A A	A A	A A	A A	T T	T T	C C	A A	C C	T C	T T	
	A A	A A	A A	G G	T T	T T	C C	A A	C C	C C	C C	T T

Table 11A:Human IgH JH6 Variant Occurrences

HUMAN POPULATION														H9	A1
NA20289; NA20294 and 19 other(s)	NA12546; NA12046 and 41 other(s)	NA18532; NA18562 and 58 other(s)	HG00628; HG00690 and 57 other(s)	HG01456; HG01275 and 33 other(s)	HG00278; HG00313 and 57 other(s)	HG00259; HG00129 and 53 other(s)	HG01619; HG01626 and 7 other(s)	NA19083; NA18986 and 48 other(s)	NA19395; NA19321 and 20 other(s)	NA19777; NA19795 and 48 other(s)	HG00640; HG01167 and 22 other(s)	NA20798; NA20513 and 58 other(s)	NA19172; NA18909 and 8 other(s)		0.509
NA20127; NA20414 and 11 other(s)	NA12348; NA10851 and 17 other(s)	NA18558; NA18579 and 9 other(s)	HG00464; HG00683 and 16 other(s)	HG01495; HG01550 and 7 other(s)	HG00358; HG00362 and 12 other(s)	HG00135; HG00245 and 6 other(s)	HG01620; HG01625 and 1 other(s)	NA18942; NA19004 and 11 other(s)	NA19038; NA19451 and 17 other(s)	NA19725; NA19773 and 6 other(s)	HG01108; HG01072 and 12 other(s)	NA20787; NA20754 and 14 other(s)	NA19116; NA18517 and 22 other(s)		0.173
NA20314; NA20317 and 11 other(s)	NA06989; NA12044 and 10 other(s)	NA18608; NA18624 and 16 other(s)	HG00436; HG00404 and 13 other(s)	HG01357; HG01461 and 10 other(s)	HG00188; HG00282 and 10 other(s)	HG00116; HG00102 and 16 other(s)		NA18974; NA18978 and 17 other(s)	NA19372; NA19474 and 16 other(s)	NA19758; NA19749 and 4 other(s)	HG01191; HG01048 and 7 other(s)	NA20522; NA20815 and 14 other(s)	NA18520; NA19209 and 16 other(s)		0.171
	NA12827		HG00584		HG00336										0.003
									NA19471; NA19401				NA18868		0.003
													NA19121; NA19175		0.002
	NA07346								NA19376						0.002
													NA18934; NA18856		0.002
															0.001

242

(i) Non-Synonymous Human IgH JH6 Variants

243

	A1A	G1B	A1A	C1C	G1B	TTT	TTT	TTT	A1B
	A1A	G1C	A1A	G1C	A1C	TTT	TTT	C1C	C1A
	A1A	A1B	A1A	G1C	A1B	TTT	TTT	G1B	A1A
	A1	A1B	A1A	C1C	A1B	TTT	TTT	G1A	A1A
	A1A	A1B	A1A	C1C	A1C	TTT	TTT	G1C	A1A
	A1A	A1B	A1B	C1C	A1B	TTT	TTT	G1B	A1A
	A1A	C1C	A1A	C1C	A1C	TTT	TTT	C1C	A1A
	A1A	G1A	A1A	C1C	A1B	TTT	TTT	A1B	A1A
	A1A	G1B	A1A	G1C	A1A	TTT	TTT	A1B	A1A
	A1A	G1B	A1A	C1C	A1A	TTT	TTT	TTT	C1A
	A1A	G1C	A1A	G1C	A1C	TTT	TTT	C1C	A1A
	A1A	G1C	A1A	TTT	A1A	TTT	TTT	TTT	A1A

[illegible]

[illegible]

SEQ ID NO: 4 (Nucleotide sequence of JH6 variant)

[illegible]

HUMAN POPULATION	

247

[illegible]




Table 13.1 - IGHA1

h.type	cum.freq	# indivs	pops	14:106173524	14:106173573	14:106173749	14:106174188	14:106174191	14:106174211	14:106174221	14:106174261
				rs1130877	rs11546792	rs182183329	rs150528881	rs1140166	rs185815870	rs117775520	rs1407
ref				Val->Ala	Leu->Met	Arg->His	Thr->Ala	Lys->Glu	Ala->Val	Pro->Ser	Glu->Asp
a	0.5884	-	-	A	A	C	T	T	G	G	C
b	0.31385	556	14	A	A	C	T	T	G	G	G
c	0.05755	115	8	A	A	T	T	T	G	G	C
e	0.01455	31	8	A	A	C	T	T	G	■	G
g	0.0072	16	4	A	A	C	C	T	G	G	C
d	0.00405	9	4	A	T	C	T	T	G	G	C
h	0.00315	7	3	A	A	C	T	T	G	G	C
j	0.0018	4	3	A	A	C	T	C	G	G	G
f	0.00135	3	2	G	T	C	T	T	G	G	C
i	0.00135	3	1	A	T	T	T	T	G	G	C
n	0.0009	2	2	A	A	C	T	T	G	G	C
m	0.0009	2	2	A	A	T	T	T	G	G	G
p	0.0009	2	1	A	T	C	C	T	G	G	C
q	0.0009	2	2	A	T	C	T	T	G	G	C
r	0.00045	1	1	A	A	T	T	C	G	G	C
k	0.00045	1	1	G	A	T	T	T	G	G	C
l	0.00045	1	1	G	A	T	T	T	G	G	C
o	0.00045	1	1	A	A	C	T	T	■	G	C

Table 13.1 - IGHA1

h.type	cum.fr	# indivs	pops	14:106174368 rs188231401 Ser->Thr	14:106174744 rs193173336 Val->Met	14:106174984 rs112706225 Lys->Glu	
ref	0.5884	-	-	A	C	T	ENST00000390547
a	0.31385	556	14	A	C	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.05755	115	8	A	C	T	1,5,7,10,11,12,13,14
c	0.01455	31	8	A	C	T	1,2,5,6,7,8,11,13
e	0.0072	16	4	A	C	T	1,10,12,14
g	0.00405	9	4	A	C	T	4,5,10,14
d	0.00315	7	3	A	C	C	1,10,14
h	0.0018	4	3	A	C	T	2,6,7
j	0.00135	3	2	A	C	C	10,14
f	0.00135	3	2	A	C	T	1,10
i	0.00135	3	1	A	C	C	14
n	0.0009	2	2	A	T	T	9,10
m	0.0009	2	2	A	C	T	6,9
p	0.0009	2	1	A	C	T	10
q	0.0009	2	2	A	C	C	10,12
r	0.00045	1	1	A	C	T	1
k	0.00045	1	1	A	C	T	10
l	0.00045	1	1	T	C	T	10
o	0.00045	1	1	A	C	T	4

Table 13.2 - IGHA2

h.type	cum.fr	# indivs	pops	14:106053293	14:106053409	14:106053441	14:106053461	14:106053491	14:106053930	14:106053975	14:106053978
				rs111788346	rs11628783	rs192043389	rs11970	rs138896525	rs139611644	rs72713340	rs142827225
ref	0.2023	-	-	G	C	G	T	C	C	G	G
a	0.35655	528	14		G	G		C	C	G	G
b	0.1647	331	14	G	C	G	T	C	C	G	G
c	0.07815	133	3	G	C	G	T	T	C		G
g	0.03195	65	7	G	G	G	T	C	C	G	G
f	0.02645	58	13	G	C	G		C	C	G	G
i	0.0228	45	7	G	C	G		C	C	G	G
d	0.0227	48	11	G	G	G		C	C	G	G
j	0.01865	41	14	G	G	G	T	C	C	G	G
e	0.01465	31	10		C	G		C	C	G	G
k	0.01045	23	11		C	G	T	C	C	G	G
h	0.00955	20	9		G	G	T	C	C	G	G
m	0.00955	21	11	G	C	G	T	C	C	G	G
l	0.0041	9	3	G	C	G	T	T	C	G	G
p	0.0036	8	5	G	G	G	T	C	C	G	G
o	0.00315	7	3		G	G		C	C		G
n	0.0027	6	4	G	C	G		C	C		G
q	0.0018	4	2	G	C	G	T	T	C	G	G
r	0.0018	4	4		C	G	T	C	C	G	G
ai	0.0018	4	1	G	G	G	T	T	T	G	G
ag	0.00135	3	3	G	C	G	T	C	C		G
s	0.00135	3	1	G	C	G		C	C	G	G
y	0.0009	2	2		G	G	T	C	C	G	G
v	0.0009	2	2		G	G		C	C	G	G
ab	0.0009	2	2	G	C	G	T	C	C	G	G
ak	0.0009	2	2	G	C	G	T	C	C	G	G
ad	0.0009	2	2	G	C	G	T	C	T	G	G
ac	0.0009	2	2	G	G	G		C	C	G	G
ah	0.00045	1	1	G	C	G	T	C	C		G
u	0.00045	1	1	G	C	G		C	C	G	G

Table 13.2 - IGHA2

aj	0.00045	1	G	G	G	T	T	T	C	■	G	G
aa	0.00045	1	G	C	G	■	C	C	T	G	G	G
t	0.00045	1	■	C	G	■	C	C	C	■	G	G
z	0.00045	1	■	C	G	T	T	C	C	■	G	G
w	0.00045	1	G	G	G	■	C	C	C	■	G	G
af	0.00045	1	G	C	■	T	C	C	C	G	G	G
x	0.00045	1	■	G	G	■	C	C	C	G	C	C
ae	0.00045	1	■	G	G	T	C	C	C	■	G	G

Table 13.2 - IGHA2

h.type	cum.fr	# indivs	pops	14:106054166	14:106054428	14:106054456
ref	0.2023	-	-	C	C	A
a	0.35655	528	14	C	G	G
b	0.1647	331	14	C	G	G
c	0.07815	133	3	C	C	A
g	0.03195	65	7	C	C	A
f	0.02645	58	13	C	G	G
i	0.0228	45	7	C	C	A
d	0.0227	48	11	C	G	G
j	0.01865	41	14	C	G	G
e	0.01465	31	10	C	G	G
k	0.01045	23	11	C	G	G
h	0.00955	20	9	C	G	G
m	0.00955	21	11	C	C	G
l	0.0041	9	3	C	C	A
p	0.0036	8	5	C	C	G
o	0.00315	7	3	C	G	G
n	0.0027	6	4	C	G	G
q	0.0018	4	2	C	G	G
r	0.0018	4	4	C	C	A
ai	0.0018	4	1	C	C	A
ag	0.00135	3	3	C	G	G
s	0.00135	3	1	C	C	G
y	0.0009	2	2	C	C	A
v	0.0009	2	2	C	C	G
ab	0.0009	2	2	C	G	A
ak	0.0009	2	2	T	C	A
ad	0.0009	2	2	C	C	A
ac	0.0009	2	2	C	C	A
ah	0.00045	1	1	C	C	A
u	0.00045	1	1	C	G	A

ENST000000390539

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

3,4,9

1,3,5,10,11,12,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,4,5,9,10,12,14

2,3,4,5,6,7,9,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,3,5,6,7,8,9,11,12,13

2,3,5,6,7,8,9,10,11,12,13

1,3,5,6,7,9,11,13,14

1,2,3,4,7,9,10,11,12,13,14

3,4,9

6,7,10,11,14

2,7,13

2,5,6,13

4,9

2,5,11,13

14

1,5,7

1

10,14

1,9

4,10

10,12

7,10

1,9

5

5

Table 13.2 - IGHA2

aj	0.00045	1	C		C		A	3
aa	0.00045	1	C		C		A	5
t	0.00045	1	C		G		G	11
z	0.00045	1	T		G		G	3
w	0.00045	1	C		G		G	13
af	0.00045	1	C		C		A	14
x	0.00045	1	C		G		G	13
ae	0.00045	1	C		G		G	2

Table 13.3 - IGHD2-2

h.type	cum.freq	# indvs	pops	14:106382687	14:106382711	
ref	0.6095	-	-	rs2857339	rs191409821	
a	0.3896	590	14	Cys->Tyr	Ile->Lys	ENST000000390591
b	0.0009	2	2	T	A	1,2,3,4,5,6,7,8,9,10,11,12,13,14
				C	T	5,13

Table 13.4 - IGH2-8

h.type	cum.freq	# indvs	pops	14:106373089		ENST00000390585
				rs181619638		
ref	0.99955	-	-	Tyr->Phe	T	
a	0.00045	1	1			2

Table 13.5 - IGHD3-10

h.type	cum.freq	# indvs	pops	14:106370370	ENST00000390583
				rs141575023	
				Gly->Arg	
ref	0.9011	-	-	C	
a	0.0989	173	14	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14

Table 13.6 - IGHD3-16

h.type	cum.freq	# indivs	pops	14:106361515	ENST00000390577
				rs143676405	
ref	0.9061	-	-	Arg->His	1,2,3,4,5,6,7,8,9,10,11,12,13,14
	0.0939	155	14	C	
				T	

Table 13.7 - IGHD3-3

h.type	cum.freq	# indvs	pops	14:106380241	ENST000000390590
				rs112802424	
ref	0.9766	-	-	C	
a	0.0234	40	6	T	
				1,7,10,12,13,14	

Table 13.8 - IGHD4-23

h.type	cum.freq	# indvs	pops	14:106350740	Arg->Trp	ENST00000437320
				rs145157618		
ref	0.9767	-	-	G		
a	0.0233	41	10			1,2,3,5,7,10,11,12,13,14

Table 13.9 - IGHD4-4

h.type	cum.freq	# indivs	pops	14:106379089		ENST000000414852
				rs190861327		
ref	0.9973	-	-	Gln->Arg	T	
a	0.0027	5	4	C		1,4,9,10

Table 13.10 - IGHD6-13

h.type	cum.freq	# indivs	pops	14:106367013 rs185385671	ENST000000390580
ref	0.9982	-	-	Ser->Asn	
a	0.0018	4	4	C	2,3,5,6
				T	

Table 13.11 - IGHD7-27

h.type	cum.freq	# indvs	pops	14:106331767	Thr->Ser	ENST000000439842
				rs187067877		
ref	0.99955	-	-		G	
a	0.00045	1	1		C	1

Table 13.12 - IGHD

h.type	cum.freq	# indivs	pops	14:106303836	14:106303846	14:106306841	14:106307012	14:106307300	14:106307432	14:106311758	14:106311961
ref	0.003	-	-	rs146595719	rs1005582	rs709589	rs184889907	rs139530086	rs111902251	rs188179725	rs149054178
a	0.735	951	14								
b	0.225	369	14								
e	0.017	32	5								
c	0.013	24	4								
d	0.005	11	3								
f	0.002	5	3								
g	5E-04	1	1								
h	5E-04	1	1								

Table 13.12 - IGHD

h.type	cum.freq	# indvs	pops	
ref	0.003	-	-	ENST000000390556
a	0.735	951	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.225	369	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
e	0.017	32	5	1,5,10,12,14
c	0.013	24	4	1,5,11,12
d	0.005	11	3	3,4,9
f	0.002	5	3	2,6,7
g	5E-04	1	1	13
h	5E-04	1	1	6

Table 13.13 - IGHEP1

h.type	cum.freq	# indvs	pops	14:106188166	
				rs142072061	
ref	0.99775	-	-	Val->Ile	ENST000000558023
a	0.00225	5	4	C	
				T	2,6,7,9

Table 13.14 - IGHE

h.type	cum.freq	# indivs	pops	14:106066524	14:106066668	14:106066690	14:106066701	14:106067096	14:106067118	14:106067815
ref	0.87655	-	-	rs189365488	rs148738013	rs61675149	rs57724657	rs182957708	rs150823665	rs144154566
a	0.08375	144	7	Arg->His	Arg->His	Trp->Gly	Ala->Val	Pro->Leu	Ala->Thr	Thr->Ile
b	0.03205	66	5	C	C	C	C	G	C	G
d	0.00225	5	2	C	C	C	C	G	C	G
c	0.00135	3	2	T	C	A	G	G	T	G
h	0.00135	3	1	C	C	A	G	G	T	G
g	0.0009	2	2	C	T	A	G	G	C	G
e	0.0009	2	2	C	C	C	G	■	C	G
f	0.0009	2	2	C	C	A	G	G	C	■

ENST000000390541

3,4,5,9,10,11,14

1,5,10,12,14

10,14

1,10

14

7,13

3,4

6,13

Table 13.15 - IGHG1

h.type	cum.freq	# indvs	pops	14:106204113	14:106204131	14:106208326	14:106208411	
				rs117518546	rs138262373	rs1043109	rs184594692	
							Asp->Glu	ENST000000390542
				Gly->Arg	Asp->Asn	Leu->Val	Asp->Glu	ENST000000390548
						Leu->Val	Asp->Glu	ENST000000390549
ref	0.89185	-	-	C	C	G	G	
a	0.09825	158	4	T	C	G	G	3,4,9,13
b	0.0063	14	6	C	T	G	G	2,5,6,7,11,13
d	0.0018	4	2	C	C	C	G	3,9
c	0.0018	4	1	C	C	G	T	14

Table 13.16 - IGHG2

h.type	cum.freq	# indivs	pops	14:106109830	14:106110017	14:106110056	14:106110137	14:106110209	14:106110914	14:106110926
ref	0.6966	-	-	rs185308159	rs182705016	rs113678609	rs8009156	rs190914111	rs11627594	rs139934867
a	0.2868	499	12	Pro->Thr	Lys->Glu	Val->Leu	Val->Met	Glu->Lys	Pro->Thr	Val->Met
b	0.0045	9	1	G	T	C	C	C	G	C
f	0.0036	8	3	G	T	C	C	C	T	T
h	0.0027	6	6	G	T	G	C	C	T	C
d	0.0023	5	3	G	T	C	C	T	G	C
e	0.0018	4	3	G	T	C	T	C	G	C
c	0.0009	2	2	T	T	C	C	C	G	C
g	0.0005	1	1	G	C	C	C	C	G	C
i	0.0005	1	1	G	T	C	C	T	T	C

ENST000000390545

1,2,3,4,5,6,7,8,9,11,12,13

10

7,11,13

1,6,7,9,10,13

3,4,9

2,4,5

1,10

10

4

Table 13.17 - IGHG3

h.type	cum.fre	# indivs	pops	14:106235611 rs1051112 Phe->Tyr	14:106235663 rs190174000 Gln->Glu	14:106235692 rs187517394 Lys->Arg	14:106235729 rs139413052 Met->Val	14:106235742 rs149653267 Asn->Lys	14:106235783 rs113169458 Val->Met	14:106236000 rs141959627 Thr->Ala	14:106236128 rs12890621 Tyr->Phe
ref	0.2495	-	-	A	G	T	T	G	C	T	T
a	0.2813	479	14	T	G	T	T	G	C	T	T
b	0.19955	345	11	A	G	T	T	G	C	T	
j	0.02345	50	8	A	G	T	T	C	C	T	T
d	0.02095	44	6	T	G	T	C	C	T	T	T
c	0.0209	46	10	T	G	T	T	G	C	T	
i	0.019	41	10	T	G	T	T	G	C	T	T
g	0.01735	38	10	T	G	T	C	C	C	T	T
f	0.01725	36	7	T	G	T	T	G	C	C	T
h	0.0164	36	11	T	G	T	T	C	C	T	T
e	0.015	33	9	A	G	T	T	G	C	T	T
k	0.01315	27	5	A	C	C	C	G	C	T	T
t	0.0126	28	6	T	G	T	T	C	C	T	T
q	0.0126	27	3	A	G	T	C	G	C	T	T
v	0.0072	16	8	T	G	T	C	G	C	T	T
o	0.00675	15	7	A	G	T	T	C	C	T	
r	0.0063	14	5	T	G	T	C	C	C	T	T
m	0.00585	13	5	A	C	T	T	C	C	T	T
n	0.00495	11	4	T	G	T	T	C	T	T	T
am	0.0045	10	5	T	G	T	C	C	T	T	T
l	0.0036	8	6	T	G	T	T	C	C	T	
p	0.0036	8	2	A	C	T	T	G	C	T	T
u	0.0036	8	4	T	G	T	C	C	C	T	
x	0.00315	7	3	T	C	T	C	C	C	T	
ae	0.00225	5	3	T	C	C	C	G	C	T	T
ai	0.00225	5	2	T	G	T	T	C	T	T	T
aj	0.0018	4	2	A	G	T	C	C	C	T	T
w	0.0018	4	1	T	G	T	T	G	C	T	T
bd	0.00135	3	1	T	G	T	C	C	C	T	T
al	0.00135	3	2	A	G	T	C	C	T	T	T

Table 13.17 - IGHG3

s	0.00135	3	T	C	T	T	T	G	C	T	T	■
ab	0.00135	3	A	G	T	T	G	G	C	T	T	T
ak	0.00135	3	A	G	T	T	C	C	T	T	T	T
ag	0.0009	2	T	G	T	C	G	G	C	T	T	T
y	0.0009	2	A	G	T	T	G	G	C	T	■	
az	0.0009	2	T	G	T	T	G	G	T	T	■	
aa	0.0009	2	A	G	T	C	C	C	C	T	■	
ap	0.0009	2	A	C	T	C	G	G	C	T	T	T
av	0.0009	2	T	G	T	T	C	C	C	T	T	T
af	0.0009	2	T	C	C	C	C	C	C	T	T	T
at	0.0009	2	T	G	C	C	G	G	C	T	T	T
aw	0.0009	2	T	G	C	T	C	C	C	C	T	T
ay	0.0009	2	A	G	T	C	G	G	C	T	■	
ar	0.0009	2	A	G	C	C	G	G	C	T	T	T
an	0.0009	2	T	G	T	T	C	C	C	T	T	T
ah	0.00045	1	A	G	T	T	G	G	C	T	T	T
ba	0.00045	1	A	G	T	T	G	G	C	C	■	
ax	0.00045	1	A	C	T	C	C	C	C	T	T	T
bb	0.00045	1	T	G	C	C	C	C	C	T	T	T
aq	0.00045	1	A	G	T	T	G	G	C	C	T	T
be	0.00045	1	T	G	C	T	G	G	C	T	T	T
z	0.00045	1	A	G	T	T	G	G	C	T	■	
ao	0.00045	1	T	C	C	T	C	C	C	T	T	T
au	0.00045	1	T	G	T	T	G	G	T	T	T	T
ad	0.00045	1	T	C	T	T	C	C	C	T	T	T
bc	0.00045	1	A	G	T	T	C	C	T	T	T	T
ac	0.00045	1	T	C	T	T	C	C	C	T	■	
as	0.00045	1	A	G	T	T	G	G	T	T	T	T

Table 13.17 - IGHG3

h.type	cum.fre	# indivs	pops	14:106236141 rs60746425 Arg->Trp	14:106236143 rs74093865 Pro->Leu	14:106236174 rs184265224 Gly->Ser	14:106236187 rs141238286 Lys->Asn	14:106236195 rs145035200 Gln->Lys	14:106236315 rs189074626 Leu->Phe	14:106237516 rs186135943 Leu->Phe
ref	0.2495	-	-	G	G	C	C	G	G	C
a	0.2813	479	14	G		C	C	G	G	C
b	0.19955	345	11	G	G	C	C	G	G	C
j	0.02345	50	8	G	G	C	C	G	G	C
d	0.02095	44	6		G	C	C	G	G	C
c	0.0209	46	10	G	G	C	C	G	G	C
i	0.019	41	10	G	G	C	C	G	G	C
g	0.01735	38	10	G		C	C	G	G	C
f	0.01725	36	7	G		C	C	G	G	C
h	0.0164	36	11	G		C	C	G	G	C
e	0.015	33	9	G		C	C	G	G	C
k	0.01315	27	5	G	G	C	C	G	G	C
t	0.0126	28	6	G	G	C	C	G	G	C
q	0.0126	27	3	G	G	C	C	G	G	C
v	0.0072	16	8	G		C	C	G	G	C
o	0.00675	15	7	G	G	C	C	G	G	C
r	0.0063	14	5	G	G	C	C	G	G	C
m	0.00585	13	5	G	G	C	C	G	G	C
n	0.00495	11	4		G	C	C	G	G	C
am	0.0045	10	5	G	G	C	C	G	G	G
l	0.0036	8	6	G	G	C	C	G	G	C
p	0.0036	8	2	G	G	C	C	G	G	C
u	0.0036	8	4	G	G	C	C	G	G	C
x	0.00315	7	3	G	G	C	C	G	G	C
ae	0.00225	5	3	G	G	C	C	G	G	C
ai	0.00225	5	2	G	G	C	C	G	G	G
aj	0.0018	4	2	G	G	C	C	G	G	C
w	0.0018	4	1	G		C	C	T	G	C
bd	0.00135	3	1	G		C	C	T	G	C
al	0.00135	3	2	G	G	C	C	G	G	G

Table 13.17 - IGHG3

s	0.00135	3	G	G	C	C	C	G	C	G	G	C
ab	0.00135	3	G	G	C	C	G	G	G	G	G	C
ak	0.00135	3	G	G	C	C	C	C	C	G	G	C
ag	0.0009	2	G	G	C	C	C	C	C	G	G	C
y	0.0009	2	G	G	T	T	C	C	C	G	G	C
az	0.0009	2	G	G	C	C	C	C	C	G	G	C
aa	0.0009	2	G	G	C	C	C	C	C	G	G	C
ap	0.0009	2	G	G	C	C	C	C	C	G	G	C
av	0.0009	2	G	G	C	C	C	C	C	T	G	C
af	0.0009	2	G	G	C	C	C	C	C	G	G	C
at	0.0009	2	G	G	C	C	C	C	C	G	G	C
aw	0.0009	2	G	G	C	C	C	C	C	G	G	C
ay	0.0009	2	G	G	C	C	C	C	C	G	G	C
ar	0.0009	2	G	G	C	C	C	C	C	G	G	C
an	0.0009	2	G	G	C	C	C	C	C	G	G	C
ah	0.00045	1	G	G	C	C	C	C	C	G	G	C
ba	0.00045	1	G	G	C	C	C	C	C	G	G	C
ax	0.00045	1	G	G	C	C	C	C	C	G	G	C
bb	0.00045	1	G	G	C	C	C	C	C	G	G	C
aq	0.00045	1	G	G	C	C	C	C	C	G	G	C
be	0.00045	1	G	G	C	C	C	C	C	G	G	C
z	0.00045	1	G	G	C	C	C	C	C	G	G	C
ao	0.00045	1	G	G	C	C	C	C	C	G	G	C
au	0.00045	1	G	G	C	C	C	C	C	G	G	C
ad	0.00045	1	G	G	C	C	C	C	C	G	G	C
bc	0.00045	1	G	G	C	C	C	C	C	G	G	C
ac	0.00045	1	G	G	C	C	C	C	C	G	G	C
as	0.00045	1	G	G	C	C	C	C	C	G	G	C

Table 13.17 - IGHG3

h.type	cum.fr	# indivs	pops	
ref	0.2495	-	-	ENST00000390551
a	0.2813	479	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.19955	345	11	1,2,3,4,5,6,7,8,11,12,13
j	0.02345	50	8	1,2,3,4,5,10,12,14
d	0.02095	44	6	1,3,4,8,9,11
c	0.0209	46	10	1,2,3,5,6,7,8,11,12,13
i	0.019	41	10	1,2,3,4,7,8,10,12,13,14
g	0.01735	38	10	2,3,4,5,6,7,9,11,12,13
f	0.01725	36	7	1,2,5,6,7,11,13
h	0.0164	36	11	2,3,4,5,6,7,8,9,11,12,13
e	0.015	33	9	2,4,5,6,7,9,11,12,13
k	0.01315	27	5	1,5,10,12,14
t	0.0126	28	6	1,2,5,10,13,14
q	0.0126	27	3	1,10,14
v	0.0072	16	8	2,3,5,6,7,9,11,12
o	0.00675	15	7	2,5,7,8,11,12,13
r	0.0063	14	5	1,4,10,12,14
m	0.00585	13	5	1,5,10,12,14
n	0.00495	11	4	3,4,9,11
am	0.0045	10	5	1,5,10,12,14
l	0.0036	8	6	2,6,7,8,12,13
p	0.0036	8	2	1,10
u	0.0036	8	4	1,2,6,11
x	0.00315	7	3	2,7,13
ae	0.00225	5	3	1,10,14
ai	0.00225	5	2	1,14
aj	0.0018	4	2	10,14
w	0.0018	4	1	4
bd	0.00135	3	1	9
al	0.00135	3	2	1,14

Table 13.17 - IGHG3			
s	0.00135	3	2 7,11
ab	0.00135	3	2 1,14
ak	0.00135	3	2 10,14
ag	0.0009	2	2 1,10
y	0.0009	2	2 2,7
az	0.0009	2	2 2,13
aa	0.0009	2	2 2,6
ap	0.0009	2	2 10,14
av	0.0009	2	1 9
af	0.0009	2	1 14
at	0.0009	2	2 10,14
aw	0.0009	2	2 5,9
ay	0.0009	2	1 11
ar	0.0009	2	2 5,14
an	0.0009	2	1 9
ah	0.00045	1	1 10
ba	0.00045	1	1 13
ax	0.00045	1	1 10
bb	0.00045	1	1 1
aq	0.00045	1	1 5
be	0.00045	1	1 7
z	0.00045	1	1 7
ao	0.00045	1	1 14
au	0.00045	1	1 5
ad	0.00045	1	1 1
bc	0.00045	1	1 1
ac	0.00045	1	1 13
as	0.00045	1	1 4

Table 13.18 - IGHG4

h.type	cum.freq	# indivs	pops	14:106090959	14:106091083	14:106091281	14:106091329	14:106091332	14:106091341	14:106091347	14:106091450
ref	0.77355	-	-	rs190355823	rs146589796	rs112773794	rs8015545	rs138186657	rs143666134	rs145943426	rs138777601
				Ser->Ala	Met->Ile	Asn->His	Leu->Val	Val->Ile	Val->Ile	Val->Ile	Gln->His
a	0.19825	389	12	A	C	T	G	C	C	C	C
b	0.01005	22	4	A	C	G	C	T	C	C	C
c	0.0055	12	4	A	T	T	G	C	C	C	C
d	0.0041	8	4	A	C	G	G	T	C	C	C
f	0.0027	6	3	A	C	T	G	C	T	C	C
e	0.00225	4	2	A	C	T	G	C	C	C	G
g	0.0018	4	2	A	C	T	G	C	C	C	C
j	0.00045	1	1	A	C	G	G	C	C	C	C
k	0.00045	1	1	C	C	T	G	C	C	C	C
h	0.00045	1	1	A	C	T	G	C	C	T	C
i	0.00045	1	1	A	C	T	C	C	C	C	G

Table 13.18 - IGHG4

h.type	cum.freq	# indvs	pops	14:106091714 rs139754373	14:106092363 rs183713766	ENST00000390543
ref	0.77355	-	-	A	C	
a	0.19825	389	12	A	C	1,2,3,4,5,6,7,8,9,11,12,13
b	0.01005	22	4	G	C	1,10,12,14
c	0.0055	12	4	A	C	1,5,10,14
d	0.0041	8	4	A	C	1,5,10,14
f	0.0027	6	3	A	C	1,10,14
e	0.00225	4	2	A	C	10,11
g	0.0018	4	2	A	T	1,14
j	0.00045	1	1	A	C	1
k	0.00045	1	1	A	C	1
h	0.00045	1	1	A	C	10
i	0.00045	1	1	A	C	4

Table 13.19 - IGHJ1

h.type	cum.fre	# indivs	pops	14:106331628	
				rs181379404	ENST000000390565
ref	0.9982	-	-	Thr->Ser	
a	0.0018	3	3	G	
				C	2,3,13

Table 13.20 - IGHJ2

h.type	cum.fre	# indivs	pops	14:106331453	14:106331455	ENST00000390564
				rs181475274	rs185806009	
				Tyr->Phe	Trp->Cys	
ref	0.9936	-	-	T	C	
a	0.00415	9	6	T	G	1,3,4,7,10,13
b	0.00225	4	3		C	1,10,14

Table 13.21 - IGHJ3

h.type	cum.freq	# indivs	pops	14:106330799	14:106330814	14:106330815	14:106330833	14:106330841
				rs190137342	rs181332275	rs185836312	rs190698203	rs183030857
				Ser->Leu	Met->Thr	Met->Val	Ile->Phe	Ala->Val
ref	0.95755	-	-	G	A	T	T	G
a	0.01645	34	12	G	A	T	T	■
b	0.0114	23	10	G	A	C	T	G
c	0.0064	14	11	G	A	T	■	G
d	0.0046	9	7	G	G	T	T	G
e	0.00315	6	5	■	A	T	T	G
f	0.00045	1	1	G	A	C	T	■
				ENST000000463911				
				1,2,3,4,5,6,7,9,10,12,13,14				
				2,3,4,6,7,9,10,11,13,14				
				1,2,5,6,7,8,9,10,11,13,14				
				5,7,9,10,11,13,14				
				1,2,3,5,10				
				5				

[illegible]

Table 13.23 - IGHJ5

h.type	cum.freq	# indivs	pops	14:106330027	14:106330035	14:106330067
				rs147551384	rs141950820	rs78872461
				Ser->Ala	Thr->Ile	Trp->Cys
ref	0.9497	-	-	A	G	C
a	0.0357	78	13	A	G	G
b	0.011	24	10	A		C
c	0.0027	5	4	C	G	C
d	0.0009	2	2	A		G
				ENST000000488476		
				1,2,3,4,5,6,7,9,10,11,12,13,14		
				1,2,3,5,6,7,8,9,10,11		
				2,3,7,13		
				4,14		

Table 13.24 - IGHJ6

h.type	cum.freq	# indivs	pops	14:106329414 rs111729691	14:106329417 rs142191790	14:106329434 rs140411866	14:106329435 rs1950943	
ref	0.3083	-	-	Ser->Pro	Val->Ile	Lys->Arg	Lys->Gln	ENST000000390560
a	0.6845	945	14	A	C	T	T	
b	0.00315	7	2	A	T	T	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14
e	0.0009	2	2	A	C	C	G	10,14
h	0.0009	2	2	G	C	T	G	4,13
f	0.0009	2	2	A	C	C	T	2,6
d	0.00045	1	1	A	T	T	G	3,10
g	0.00045	1	1	G	C	C	G	9
c	0.00045	1	1	G	C	T	T	4
								10

Table 13.25 -IGHM

h.type	cum.freq	# indvs	pops	14:106320732	14:106321313	14:106321330	14:106321590	14:106321663	14:106322093
ref	0.0249	-	-	rs146456238	rs186069870	rs45471499	rs12365	rs1059216	rs113762053
a	0.7527	974	14	Asp->Asn	Ala->Thr	Thr->Ile	Gly->Val	Ser->Gly	Asp->Glu
b	0.15245	287	14	C	C	G	C	T	G
d	0.03795	74	9	C	C	G	C	C	C
c	0.02705	54	3	C	C			C	G
e	0.0018	4	2	C	C	G	C	C	G
g	0.00135	3	3	C	T	G	C	T	G
f	0.0009	2	1	C	C	G	C	T	C
h	0.00045	1	1	C	T	G		T	G
i	0.00045	1	1	T	C	G	C	C	C

ENST000000390559

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,5,7,10,11,12,13,14

3,4,9

10,14

5,12,14

11

5

10

Table 13.26 - IGHV1-18

h.type	cum.freq	# indivs	pops	14:106641643	14:106641660	14:106641668	14:106641688	14:106641769	14:106641801	14:106641832	14:106641981
				rs181586608	rs146311654	rs184544516	rs114910155	rs139305030	rs188509150	rs181324183	rs184535212
				Thr->Ala	Gly->Asp	Lys->Asn	Asn->Asp	Thr->Pro	Lys->Arg	Ala->Pro	Ser->Asn
ref	0.97595	-	-	T	C	C	T	T	T	C	C
a	0.01185	23	6	T	C	C	C	T	T	C	C
c	0.0045	7	4	T	T	C	T	T	T	C	C
b	0.00275	6	5	T	C	G	T	T	T	C	C
d	0.00135	3	3	T	C	C	T	T	C	C	C
e	0.00135	3	3	T	C	C	T	G	T	C	C
f	0.0009	2	2	T	C	C	T	T	T	C	T
g	0.00045	1	1	C	T	C	T	T	T	C	C
h	0.00045	1	1	T	C	C	T	T	T	G	C
i	0.00045	1	1	C	C	C	T	T	T	C	C

Table 13.26 - IGHV1-18

h.type	cum.freq	# indvs	pops	
ref	0.97595	-	-	ENST00000390605
a	0.01185	23	6	1,6,9,10,11,14
c	0.0045	7	4	1,10,11,14
b	0.00275	6	5	6,10,11,13,14
d	0.00135	3	3	4,11,14
e	0.00135	3	3	2,6,11
f	0.0009	2	2	8,9
g	0.00045	1	1	13
h	0.00045	1	1	10
i	0.00045	1	1	10

Table 13.27 - IGHV1-24

h.type	cum.freq	# indivs	pops	14:106733154	14:106733158	14:106733166	14:106733192	14:106733263	14:106733269	14:106733274	14:106733336
				rs141629050	rs150510876	rs139845206	rs141655646	rs183704154	rs114334184	rs187836940	rs139283249
ref	0.97175	-	-	Tyr->Phe	Tyr->His	Thr->Met	Glu->Asp	Ile->Phe	Glu->Lys	Asp->Gly	Met->Ile
a	0.01185	24	4	T	A	G	C	T	C	T	C
b	0.0082	18	4	T	A	G	C	T	C	T	C
c	0.00415	9	3	T	A	G	G	T	C	T	C
g	0.0009	2	1	T	A	G	C	T	C	T	T
d	0.00045	1	1	T	G		C	T	C	T	C
j	0.00045	1	1	T	A	G	C	T	C	T	C
k	0.00045	1	1	T	A	G	C	T	C	C	C
e	0.00045	1	1	T	A	G	C		C	T	C
h	0.00045	1	1	T	A	G	C	T	T	T	C
f	0.00045	1	1	T	A	G	C	T	C	T	C
i	0.00045	1	1		A	G	C	T	C	T	C

Table 13.27 - IGHV1-24

h.type	cum.freq	# indivs	pops	14:106733337	14:106733341	14:106733359	14:106733446	14:106733570
				rs8012805	rs186604864	rs184694183	rs147700118	rs192824042
ref	0.97175	-	-	A	A	A	T	T
a	0.01185	24	4	C	A	A	T	T
b	0.0082	18	4	A	A	A	C	T
c	0.00415	9	3	A	A	A	T	T
g	0.0009	2	1	A	A	A	T	T
d	0.00045	1	1	A	A	A	T	T
j	0.00045	1	1	A	A	A	T	C
k	0.00045	1	1	A	G	T	C	T
e	0.00045	1	1	A	G	A	T	T
h	0.00045	1	1	A	A	A	T	T
f	0.00045	1	1	A	A	T	T	T
i	0.00045	1	1	A	A	A	T	T

ENST000000390610

1,10,11,14

1,10,12,14

1,10,14

2

14

5

5

5

14

13

3

Table 13.28 - IGHV1-2

h.type	cum.freq	# indivs	pops	14:106452690	14:106452735	14:106452766	14:106452784	14:106452793	14:106452799	14:106452817	14:106452868
				rs12588974	rs188093056	rs112806369	rs180944534	rs186314384	rs189402450	rs1065059	rs192043188
ref	0.43185 -	-	-	Ala->Val	Ser->Asn	Arg->Trp	Ala->Ser	Thr->Ser	Gly->Ser	Trp->Arg	Tyr->Asn
a	0.3268	511	14	G	C	T	C	T	C	A	A
b	0.1304	228	14	G	C	■	C	T	C	A	A
c	0.0797	158	14	■	C	T	C	T	C	G	A
d	0.01135	25	10	G	C	T	C	T	C	G	A
e	0.0073	16	6	■	C	T	C	T	C	A	A
g	0.0018	4	3	G	T	T	C	T	C	A	A
h	0.00135	3	2	■	C	■	C	T	C	A	A
f	0.00135	3	3	G	C	■	C	T	C	A	A
i	0.0009	2	2	■	C	T	C	T	C	G	A
m	0.0009	2	2	G	C	T	C	T	C	G	A
l	0.0009	2	2	G	C	T	C	T	C	A	A
j	0.00045	1	1	G	C	T	■	T	C	A	A
u	0.00045	1	1	G	T	T	C	T	C	A	A
k	0.00045	1	1	G	C	T	C	T	C	A	A
t	0.00045	1	1	G	C	T	C	T	C	G	T
v	0.00045	1	1	G	C	T	C	T	C	A	A
s	0.00045	1	1	G	T	T	C	■	C	A	A
q	0.00045	1	1	G	C	T	C	T	T	A	A
w	0.00045	1	1	G	C	■	C	T	C	A	A
r	0.00045	1	1	G	C	T	C	T	C	A	A
n	0.00045	1	1	G	C	T	C	T	C	A	A
p	0.00045	1	1	G	T	T	C	T	C	G	A
o	0.00045	1	1	G	C	T	C	T	C	A	A

Table 13.28 - IGHV1-2

h.type	cum.freq	# indvs	pops	14:106452873	14:106452882	14:106452897	14:106452903	14:106452956	14:106452972
				rs1065058	rs11553010	rs185487229	rs189604828	rs182256840	rs186783545
				Gly->Asp	Thr->Ser	Lys->Met	Ser->Cys	Gln->His	Ala->Gly
ref	0.43185	-	-	C	G	T	G	C	G
a	0.3268	511	14	C	G	T	G	C	G
b	0.1304	228	14	C	G	T	G	C	G
c	0.0797	158	14	C	G	T	G	C	G
d	0.01135	25	10	C	G	T	G	C	G
e	0.0073	16	6	T	G	T	G	C	G
g	0.0018	4	3	C	G	T	G	C	G
h	0.00135	3	2	C	G	T	G	C	G
f	0.00135	3	3	T	G	T	G	C	G
i	0.0009	2	2	T	G	T	G	C	G
m	0.0009	2	2	T	G	T	G	C	G
l	0.0009	2	2	C	C	T	G	C	G
j	0.00045	1	1	C	G		G	C	G
u	0.00045	1	1	T	G	T	G	C	G
k	0.00045	1	1	C	C	T	G	C	G
t	0.00045	1	1	C	G	T	G	C	G
v	0.00045	1	1	T	G		G	C	G
s	0.00045	1	1	C	G	T	G	C	G
q	0.00045	1	1	C	G	T	G	C	G
w	0.00045	1	1	C	G	T	C	C	G
r	0.00045	1	1	C	G	T	G	C	C
n	0.00045	1	1	T	G	T	G	G	G
p	0.00045	1	1	C	G	T	G	C	G
o	0.00045	1	1	C	G	T	G	G	G

ENST000000390594

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,9,10,11,13,14

1,2,5,10,12,14

6,12,14

3,4

2,6,7

11,13

3,6

9,10

2

2

2

2

10

2

10

11

2

1

1

10

Table 13.29 - IGHV1-3

h.type	cum.freq	# indvs	pops	14:106471253	14:106471268	14:106471280	14:106471296	14:106471310	14:106471353	14:106471361	14:106471365
				rs190814253	rs1143505	rs141136074	rs113948114	rs191691305	rs34069216	rs187612074	rs191777242
ref	0.34155	-	-	Cys->Tyr	Met->Thr	Arg->Thr	Glu->Gln	Ser->Asn	Glu->Lys	Tyr->Cys	Lys->Gln
a	0.3564	555	14	C	A	C	C	C	C	T	T
b	0.0724	155	13	C	G	C	C	C	T	T	T
d	0.0439	97	13	T	A	G	C	C	C	T	T
c	0.02865	63	12	T	A	C	C	C	T	T	T
h	0.019	42	12	C	G	C	C	C	T	T	T
i	0.0113	25	11	C	A	C	C	C	T	T	T
g	0.0105	23	8	C	A	C	G	C	T	T	T
f	0.00995	22	9	C	A	C	C	C	T	T	T
j	0.0077	16	8	C	A	C	C	C	C	T	T
e	0.0073	16	4	T	A	C	C	C	C	T	T
n	0.0072	16	10	C	G	C	C	C	C	T	T
t	0.00675	15	8	T	A	G	C	C	T	T	T
r	0.00585	13	9	C	G	C	C	C	T	T	T
m	0.00585	13	6	T	A	G	C	C	T	T	T
l	0.00585	13	8	C	A	C	C	C	C	T	T
o	0.00495	11	5	C	G	C	C	C	T	T	T
p	0.0045	10	4	T	A	G	C	C	C	T	T
ab	0.00405	9	6	C	A	C	C	C	T	T	T
k	0.0036	8	5	C	G	C	C	C	T	T	T
x	0.00315	7	6	C	A	C	G	C	C	T	T
ah	0.0027	6	5	C	A	C	G	C	T	T	T
u	0.00225	5	2	C	G	C	C	C	T	T	T
v	0.00225	5	2	T	A	G	C	C	T	T	T
ap	0.00225	4	3	C	G	C	C	C	T	T	T
ac	0.00225	5	4	T	A	G	C	C	C	T	T
y	0.0018	4	3	C	A	G	C	C	C	T	T
ar	0.0018	4	3	C	G	C	C	C	T	T	T
ai	0.0018	4	3	C	A	C	G	C	T	T	T

Table 13.29 - IGHV1-3

aa	0.00135	3	C	G	C	C	C	C	T	T	T	T
ak	0.00135	3	C	A	C	C	C	C	C	C	T	T
w	0.00135	3	C	G	C	C	C	C	C	C	T	T
at	0.00135	3	T	A	G	C	C	C	T	T	T	T
ag	0.0009	2	C	A	G	C	C	C	T	T	T	T
aj	0.0009	2	T	G	C	C	C	C	T	T	T	T
aq	0.0009	2	T	A	G	C	C	C	C	T	T	T
s	0.0009	2	T	A	G	C	G	C	C	T	T	T
q	0.0009	2	C	G	C	C	G	C	T	T	T	T
z	0.0009	2	C	G	C	C	C	C	C	T	T	T
ad	0.0009	2	C	G	C	C	C	C	C	T	T	T
aw	0.0009	2	C	A	C	C	G	C	T	T	T	T
ae	0.0009	2	C	G	C	C	G	C	C	T	T	T
an	0.0009	2	T	A	C	C	C	C	T	T	T	T
bd	0.00045	1	C	G	C	C	C	C	C	T	T	T
ba	0.00045	1	C	G	C	C	C	C	C	T	T	T
ax	0.00045	1	C	A	C	C	C	C	C	T	T	T
bb	0.00045	1	C	G	C	C	C	C	T	T	T	T
az	0.00045	1	C	G	C	C	G	C	T	T	T	T
al	0.00045	1	T	A	G	C	C	C	C	T	T	T
be	0.00045	1	C	G	C	C	C	C	T	T	T	T
bf	0.00045	1	C	A	C	C	G	C	T	T	T	T
av	0.00045	1	C	A	C	C	C	C	C	C	G	G
ao	0.00045	1	T	A	G	C	C	C	C	C	T	T
af	0.00045	1	C	A	C	C	C	C	C	T	T	T
au	0.00045	1	C	G	C	C	C	C	T	T	T	T
bh	0.00045	1	T	A	C	C	C	C	T	T	T	T
am	0.00045	1	C	A	C	C	C	T	C	T	T	T
bc	0.00045	1	C	G	C	C	C	C	T	T	T	T
bg	0.00045	1	T	A	G	C	C	C	C	T	T	G
ay	0.00045	1	C	G	C	C	C	T	T	T	T	T
as	0.00045	1	T	A	G	C	C	C	C	T	T	G

Table 13.29 - IGHV1-3

h.type	cum.freq	# indvs	pops	14:106471383	14:106471388	14:106471421	14:106471503	14:106471508	14:106471547
				rs184320740	rs34874585	rs74091631	rs186640498	rs192340916	rs1143496
ref	0.34155	-	-	Ala->Thr	Ser->Ile	Ala->Asp	Lys->Gln	Val->Glu	Val->Ala
a	0.3564	555	14	C	C	G	T	A	A
b	0.0724	155	13	C		G	T	A	G
d	0.0439	97	13	C	C	G	T	A	A
c	0.02865	63	12	C		G	T	A	G
h	0.019	42	12	C	C	G	T	A	A
i	0.0113	25	11	C		G	T	A	G
g	0.0105	23	8	C		G	T	A	G
f	0.00995	22	9	C		G	T	A	A
j	0.0077	16	8	C		G	T	A	A
e	0.0073	16	4	C	C	G	T	A	A
n	0.0072	16	10	C	C	G	T	A	A
t	0.00675	15	8	C	C	G	T	A	A
r	0.00585	13	9	C	C	G	T	A	A
m	0.00585	13	6	C		G	T	A	A
l	0.00585	13	8	C	C	G	T	A	G
o	0.00495	11	5	C	C	G	T	A	G
p	0.0045	10	4	C		G	T	A	A
ab	0.00405	9	6	C	C	G	T	A	G
k	0.0036	8	5	C		G	T	T	G
x	0.00315	7	6	C	C	G	T	A	A
ah	0.0027	6	5	C		G	T	A	G
u	0.00225	5	2	C		T	T	A	G
v	0.00225	5	2	C		T	T	A	G
ap	0.00225	4	3	C		G	G	A	G
ac	0.00225	5	4	C		G	T	A	G
y	0.0018	4	3	C	C	G	T	A	A
ar	0.0018	4	3	T		G	T	A	G
ai	0.0018	4	3	C	C	G	T	A	G

ENST000000390595

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,4,5,6,7,9,11,12,13,14

1,2,3,4,5,6,7,9,10,11,13,14

1,2,3,4,5,6,7,9,10,11,13,14

1,2,3,5,6,7,10,11,12,13,14

2,3,4,6,7,9,11,13

2,3,5,7,8,9,11,13,14

1,4,5,6,7,10,12,14

1,10,12,14

2,4,5,6,7,9,10,12,13,14

3,4,6,9,10,11,13,14

3,4,5,6,7,10,12,13,14

1,2,10,11,13,14

1,3,5,6,7,9,10,13

2,3,6,7,13

1,9,10,14

4,6,8,9,10,13

2,6,7,12,13

3,4,7,9,10,11

3,5,6,9,13

1,14

1,14

3,4,9

1,10,12,14

1,3,12

1,10,13

2,4,7

Table 13.29 - IGHV1-3

aa	0.00135	3	C		T	T	A	A	14
ak	0.00135	3	C		G	T	A	G	1,9,10
w	0.00135	3	C		G	T	A	A	9
at	0.00135	3	C	C	G	T	A	G	6,10
ag	0.0009	2	C		G	T	A	A	10,14
aj	0.0009	2	T	C	G	T	A	G	3,4
aq	0.0009	2	C	C	G	T	A	A	10
s	0.0009	2	C	C	G	T	A	A	7,10
q	0.0009	2	T	C	G	T	A	A	2
z	0.0009	2	C		G	T	A	G	5
ad	0.0009	2	C	C	G	T	A	A	5,9
aw	0.0009	2	C	C	G	T	A	A	7,13
ae	0.0009	2	C		G	T	A	G	5,10
an	0.0009	2	C	C	T	T	A	A	12,14
bd	0.00045	1	C	C	G	G	A	G	10
ba	0.00045	1	C		G	T	A	G	3
ax	0.00045	1	C		G	T	A	A	5
bb	0.00045	1	C	C	G	T	A	A	8
az	0.00045	1	C	C	G	T	A	G	4
al	0.00045	1	T	C	G	T	A	A	10
be	0.00045	1	C		G	T	A	A	14
bf	0.00045	1	C	C	G	T	A	A	2
av	0.00045	1	C		T	T	A	G	13
ao	0.00045	1	T	C	G	T	A	A	1
af	0.00045	1	C	C	T	T	A	A	13
au	0.00045	1	C		G	T	A	A	14
bh	0.00045	1	C	C	G	T	A	A	14
am	0.00045	1	T		G	T	A	A	10
bc	0.00045	1	C		G	T	A	A	10
bg	0.00045	1	C		G	T	A	A	10
ay	0.00045	1	C		G	T	A	G	4
as	0.00045	1	C		G	T	A	G	10

Table 13.30 - IGHV1-45

h.type	cum.freq	# indivs	pops	14:106962947 rs186738824	14:106962984 rs145326338	14:106963101 rs7141669	14:106963167 rs182373995	14:106963326 rs12884037	
				Met->Thr	Met->Leu	Gly->Arg	Val->Phe	Ala->Val	ENST00000390621
ref	0.75745	-	-	A	T	C	C	G	
a	0.1286	251	12	A	T	T	C	G	1,2,3,4,5,6,7,9,10,11,12,13
b	0.1126	220	12	A	T	C	C		1,2,3,5,6,7,8,10,11,12,13,14
d	0.00045	1	1	G	T	C	C	G	1
e	0.00045	1	1	A	G	C	C	G	9
c	0.00045	1	1	A	T	C		G	3

Table 13.31 - IGHV1-46

h.type	cum.freq	# indivs	pops	14:106967122	14:106967151	14:106967155	14:106967171	14:106967173	14:106967177	14:106967241	14:106967249
				rs147211698	rs55801711	rs149338091	rs181189514	rs185595166	rs190309173	rs144704015	rs182132309
ref				Thr->Met	Phe->Leu	Lys->Arg	Thr->Ala	Ser->Thr	Gly->Ser	Met->Ile	Tyr->Asn
a	0.9778	-	-	G	G	T	T	C	C	C	A
b	0.0082	16	6	G	C	T	T	C	C	C	A
c	0.0032	7	3		G	T	T	C	C	C	A
d	0.00225	5	3	G	G	T	T	C	C	C	A
e	0.0018	4	4	G	G	T	T	C	C	T	A
k	0.0009	2	2	G	G	C	T	C	C	C	A
n	0.0009	2	2	G	G	T	T	G	C	C	A
j	0.0009	2	2	G	G	T	T	C	C	C	A
g	0.00045	1	1	G	G	T	T	C	C	C	A
h	0.00045	1	1	G	G	T	T	G	C	C	T
f	0.00045	1	1	G	G	T	T	C	C	C	A
i	0.00045	1	1	G	G	T	T	C	C	C	T
m	0.00045	1	1	G	G	C	T	C	T	C	A
l	0.00045	1	1	G	G	T	C	C	C	C	A
p	0.00045	1	1	G	G	T	T	C	C	C	A
o	0.00045	1	1	G	G	T	T	G	C	T	A

Table 13.31 - IGHV1-46

h.type	cum.freq	# indivs	pops	14:106967264	14:106967273	14:106967275	14:106967285	14:106967306	14:106967451	14:106967453
ref			-	rs187613260	rs192285778	rs61995748	rs61747196	rs188566927	rs191694446	rs143810901
				Tyr->Asp	Ala->Ser	Lys->Met	Val->Ile	Lys->Gln	Ala->Thr	Leu->Pro
a	0.9778	16	-	A	C	T	C	T	C	A
b	0.0082	7	1,5,7,8,12,13	A	C	T	C	T	C	A
c	0.0032	5	3,10,14	A	C	T	C	T	C	A
d	0.00225	4	1,10,14	A	C	■	C	T	C	A
k	0.0018	2	2,10,11,14	A	C	T	C	T	C	A
e	0.0009	2	1,10	A	C	T	C	T	C	A
n	0.0009	2	6,10	A	C	T	C	T	C	A
j	0.0009	1	4,6	A	C	T	T	T	C	A
g	0.00045	1	1	A	■	T	C	T	C	A
h	0.00045	1	5	A	C	T	C	T	C	A
f	0.00045	1	5	A	C	T	T	G	C	A
i	0.00045	1	10	A	C	T	T	T	T	A
m	0.00045	1	9	A	C	T	C	T	C	A
l	0.00045	1	3	A	C	T	C	T	C	A
p	0.00045	1	3	A	C	T	C	T	C	A
o	0.00045	1	4	A	C	T	C	G	C	A
	0.00045	1	10	C	C	T	C	T	C	G

ENST000000390622

Table 13.32 - IGHV1-58

h.type	cum.freq	# indivs	pops	14:107078380	14:107078395	14:107078567	14:107078570	14:107078578	14:107078591	14:107078609	14:107078661
				rs189473179	rs141236708	rs2516904	rs145028255	rs189734868	rs146830036	rs113248643	rs180889109
ref				Cys->Tyr	Thr->Met	Met->Val	Ala->Pro	Thr->Ser	Gly->Arg	Val->Phe	Met->Ile
a	0.16105	-	-	C	G	T	C	G	C	C	C
b	0.4503	718	14	C	G	C	C	G	C	C	C
c	0.3144	568	14	C	G	T	C	G	C	C	C
d	0.05525	105	6	C	G	T	C	G	C		C
e	0.0068	15	6	C		T	C	G	C	C	C
f	0.0068	15	4	C	G	C	C	G	C	C	C
g	0.00225	5	3	T	G	T	C	G	C	C	C
h	0.0009	2	1	C	G	C	C	G	C		C
i	0.00045	1	1	C	G	C	C	C	C	C	C
j	0.00045	1	1	C	G	T	C	G	T	C	C
k	0.00045	1	1	C	G	T	C	G	C	C	
l	0.00045	1	1	C	G	C	C	G	C	C	C
m	0.00045	1	1	C	G	C	G	G	C	C	C
n	0.00045	1	1	C	G	T	C	G	C	C	C

Table 13.32 - IGHV1-58

h.type	cum.freq	# indvs	pops	14:107078674	14:107078775	14:107078790
				rs186300171	rs148981028	rs1858692
ref	0.16105	-	-	Ala->Val	Gly->Arg	Val->Ile
a	0.4503	718	14	G	C	C
b	0.3144	568	14	G	C	T
c	0.05525	105	6	G	C	T
d	0.0068	15	6	G	C	C
e	0.0068	15	4	G	T	T
f	0.00225	5	3	G	C	T
g	0.0009	2	1	G	C	T
j	0.00045	1	1	G	C	T
k	0.00045	1	1	G	C	C
h	0.00045	1	1	■	C	C
i	0.00045	1	1	■	C	T
l	0.00045	1	1	G	C	C

ENST000000390628

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,5,10,12,13,14

3,4,5,9,11,12

1,10,13,14

3,4,9

10

10

13

10

1

13

Table 13.33 - IGHV1-69

h.type	cum.freq	# indivs	pops	14:107169948	14:107169992	14:107169995	14:107170004	14:107170011	14:107170014	14:107170019	14:107170025
				rs190613018	rs183124973	rs1064565	rs191670994	rs74089691	rs11557979	rs192735419	rs185522094
ref	0.49795	-	-	Val->Leu	Thr->Arg	Ser->Thr	Lys->Arg	Ala->Thr	Thr->Ser	Thr->Lys	Arg->Thr
a	0.1689	323	14	C	G	C	T	C	T	G	C
b	0.1543	312	14	C	G	C	T	C	T	G	C
c	0.0561	118	13	C	G	C	T	C	T	G	C
g	0.0348	76	7	C	G	C	T	T	T	G	C
e	0.014	31	4	C	G	C	T	T	T	G	C
l	0.0108	24	6	C	G	C	T	C	T	G	C
d	0.00995	22	3	C	G	C	T	C	T	G	C
h	0.00765	17	10	C	G	C	T	C	T	G	C
f	0.00685	15	8	C	G	C	T	C	T	G	C
i	0.00675	15	6	C	G	C	T	C	T	G	C
m	0.0027	6	3	C	G	C	T	C	T	G	C
j	0.00225	5	3	C	G	C	T	C	T	G	C
u	0.00225	5	4	C	G	C	T	C	T	G	C
s	0.0018	4	3	C	G	C	T	C	T	G	C
k	0.00135	3	1	C	G	C	T	T	T	G	C
ac	0.00135	3	3	C	G	C	T	C	T	G	C
ah	0.0009	2	2	C	G	G	T	C	T	G	C
t	0.0009	2	2	C	G	C	T	C	T	G	C
ao	0.0009	2	1	C	G	C	T	T	T	G	C
af	0.0009	2	2	C	G	G	T	C	T	G	C
ad	0.0009	2	2	C	G	C	T	C	T	G	C
n	0.0009	2	2	C	G	C	T	C	T	G	C
an	0.0009	2	2	C	G	C	T	T	T	G	C
o	0.0009	2	2	C	G	C	T	C	T	G	C
ba	0.00045	1	1	C	G	C	T	T	T	G	C
ax	0.00045	1	1	C	G	C	T	T	T	G	C
ag	0.00045	1	1	C	G	C	T	C	T	G	C
y	0.00045	1	1	C	G	C	T	C	T	G	C

Table 13.33 - IGHV1-69

az	0.00045	1	C		G		C	T	T	T	G	C	C
aj	0.00045	1	C		G		C	T	T	G	G	G	G
aa	0.00045	1	C		G		C	T	T	C	G	C	G
aq	0.00045	1	C		G		C	T	T	C	G	C	C
v	0.00045	1	C		G		C	T	T	C	G	C	C
al	0.00045	1	C		G		C	T	T	C	G	C	C
ab	0.00045	1	C		G		C	T	T	C	G	C	C
ak	0.00045	1	G		G		C	T	T	C	G	C	C
q	0.00045	1	C		G		C	T	T	C	G	C	C
ap	0.00045	1	C		G		C	C	T	C	T	C	C
z	0.00045	1	C		G		C	T	T	C	G	C	C
w	0.00045	1	C		G		C	T	T	C	G	C	C
av	0.00045	1	C		C		C	T	T	C	G	C	C
r	0.00045	1	C		G		C	T	T	C	G	C	C
x	0.00045	1	C		G		C	T	T	C	G	C	C
au	0.00045	1	G		G		C	T	T	C	G	C	C
am	0.00045	1	C		G		C	T	T	T	G	C	C
at	0.00045	1	C		G		G	T	T	C	G	C	C
aw	0.00045	1	C		G		C	T		C	G	C	C
ay	0.00045	1	C		G		C	T	T	C	G	C	C
ar	0.00045	1	C		G		C	T	T	C	G	C	C
p	0.00045	1	C		G		C	T	T	C	G	C	C
as	0.00045	1	C		G		C	T	T	C	G	C	C
ae	0.00045	1	C		G		C	T	T	C	G	C	C
ai	0.00045	1	C		G		G	T	T	C	G	C	C

Table 13.33 - IGHV1-69

h.type	cum.freq	# indivs	pops	14:107170028	14:107170034	14:107170055	14:107170056	14:107170062	14:107170077	14:107170091	14:107170121
				rs11557995	rs181467093	rs8009570	rs184214708	rs55891010	rs11845244	rs189019973	rs1064564
ref	0.49795	-	-	Gly->Ala	Phe->Ser	Thr->Ile	Thr->Ala	Phe->Leu	Gly->Arg	Leu->Pro	Ser->Asn
a	0.1689	323	14	C	A	G	T	A	C	A	C
b	0.1543	312	14	C	A		T	G	T	A	C
c	0.0561	118	13	C	A		T	G	C	A	C
g	0.0348	76	7	C	A	G	T	A	C	A	C
e	0.014	31	4	C	A	G	T	A	T	A	C
l	0.0108	24	6	C	A	G	T	A	T	A	C
d	0.00995	22	3	C	A	G	T	A	T	A	C
h	0.00765	17	10	C	A		T	A	C	A	C
f	0.00685	15	8	C	A		T	A	T	A	C
i	0.00675	15	6	C	A		T	A	T	A	C
m	0.0027	6	3	C	A	G	T	A	C	A	C
j	0.00225	5	3	C	A	G	T	G	T	A	C
u	0.00225	5	4	C	A		T	G	C	A	C
s	0.0018	4	3	C	A	G	T	A	C	A	T
k	0.00135	3	1	C	A		T	G	T	A	C
ac	0.00135	3	3	C	A	G	T	A	C	A	C
ah	0.0009	2	2	C	A	G	T	A	C	A	C
t	0.0009	2	2	C	A	G	T	A	C	A	C
ao	0.0009	2	1	C	A		T	G	C	A	C
af	0.0009	2	2	C	A		T	G	C	A	C
ad	0.0009	2	2	C	A	G	T	A	C	A	C
n	0.0009	2	2	C	A		T	G	T	A	T
an	0.0009	2	2	C	A		T	G	T	A	C
o	0.0009	2	2	C	A		T	G	T	A	C
ba	0.00045	1	1	C	A	G	T	A	C	A	C
ax	0.00045	1	1	C	A	G	T	A	C	A	C
ag	0.00045	1	1	C	A		T	G	T	A	C
y	0.00045	1	1	C	A		T	G	T	G	C

Table 13.33 - IGHV1-69

az	0.00045	1	C	A	G	T	A	T	A	C	C
aj	0.00045	1	C	A	G	C	A	C	A	T	T
aa	0.00045	1	C	A		T	A	C	A	C	C
aq	0.00045	1	C	A	G	C	A	C	A	C	C
v	0.00045	1	C	A		T	G	T	A	C	C
al	0.00045	1	C	A		T	G	T	A	C	C
ab	0.00045	1	C	A		T	G	C	A	T	T
ak	0.00045	1	C	A	G	T	A	C	A	C	C
q	0.00045	1	C	A	G	T	A	C	A	C	C
ap	0.00045	1	C	A	G	T	A	C	A	C	C
z	0.00045	1	C	A	G	T	A	C	A	C	C
w	0.00045	1	C	A		T	G	T	A	T	T
av	0.00045	1	C	A		T	G	T	A	C	C
r	0.00045	1	C	A		T	G	T	A	C	C
x	0.00045	1	C	A		T	G	T	A	C	C
au	0.00045	1	C	A		T	G	T	A	C	C
am	0.00045	1	C	A	G	T	A	C	A	C	C
at	0.00045	1	C	A		T	G	T	A	C	C
aw	0.00045	1	C	G		T	G	T	A	T	T
ay	0.00045	1	C	A	G	C	A	C	A	T	T
ar	0.00045	1	C	A		T	G	T	A	C	C
p	0.00045	1	C	A		T	G	T	A	C	C
as	0.00045	1	G	A		T	G	C	A	C	C
ae	0.00045	1	C	A	G	T	A	C	A	C	C
ai	0.00045	1	C	A		T	G	T	A	C	C

Table 13.33 - IGHV1-69

h.type	cum.freq	# indivs	pops	14:107170127	14:107170128	14:107170130	14:107170133	14:107170139	14:107170143	14:107170145	14:107170173
				rs11557984	rs11558005	rs140392997	rs138194290	rs11558018	rs186595111	rs189122962	rs150196675
ref	0.49795	-	-	Ala->Val	Ala->Thr	Tyr->Phe	Ser->Asn	Phe->Ser	Thr->Pro	Gly->Asp	Val->Leu
a	0.1689	323	14	G	C	T	C	A	T	C	C
b	0.1543	312	14	G	T	T	C	A	T	C	C
c	0.0561	118	13	G	C	T	C	A	T	C	C
g	0.0348	76	7	G	C	T	C	A	T	C	C
e	0.014	31	4	G	C	T	C	A	T	C	C
l	0.0108	24	6	G	C	T	C	A	T	C	C
d	0.00995	22	3	G	T	T	C	A	T	C	C
h	0.00765	17	10	G	C	T	C	A	T	C	C
f	0.00685	15	8	G	T	T	C	A	T	C	C
i	0.00675	15	6	G	C	T	C	A	T	C	C
m	0.0027	6	3	G	C	T	C	A	T	C	C
j	0.00225	5	3	G	C	T	C	A	T	C	C
u	0.00225	5	4	G	T	T	C	A	T	C	C
s	0.0018	4	3	G	C	T	C	A	T	C	C
k	0.00135	3	1	G	T	T	C	A	T	C	C
ac	0.00135	3	3	G	T	T	C	A	T	C	C
ah	0.0009	2	2	G	C	T	C	A	T	C	C
t	0.0009	2	2	■	C	T	C	A	T	C	C
ao	0.0009	2	1	G	C	T	C	A	T	C	C
af	0.0009	2	2	G	C	T	C	A	T	C	C
ad	0.0009	2	2	G	C	T	T	A	T	C	C
n	0.0009	2	2	G	C	T	C	A	T	C	C
an	0.0009	2	2	G	C	T	C	A	T	C	C
o	0.0009	2	2	■	T	T	C	A	T	C	C
ba	0.00045	1	1	G	C	T	C	A	T	C	C
ax	0.00045	1	1	G	C	T	C	A	T	C	G
ag	0.00045	1	1	G	T	■	T	A	T	C	C
y	0.00045	1	1	G	C	T	C	A	T	C	C

Table 13.33 - IGHV1-69

az	0.00045	1	G		C	T	T	T	A	T	C	C	C
aj	0.00045	1	G		C	T	T	T	A	T	C	C	C
aa	0.00045	1	G		T		T	T	A	T	C	C	C
aq	0.00045	1	G		C		T	T	A	T	C	C	C
v	0.00045	1	G		C		T	T	A	G	C	C	C
al	0.00045	1	G		C		T	T	A	T	C	C	C
ab	0.00045	1	G		C		T	T	A	T	T	C	C
ak	0.00045	1	G		C		T	T	A	T	C	C	C
q	0.00045	1	G		C		T	T	G	T	C	C	C
ap	0.00045	1	G		C		T	T	A	T	C	C	C
z	0.00045	1	G		C		T	T	A	T	C	C	C
w	0.00045	1	G		T		T	T	A	T	C	C	C
av	0.00045	1	G		C		T	T	A	T	C	C	C
r	0.00045	1			C		T	T	A	T	C	C	C
x	0.00045	1	G		C		T	T	A	T	T	C	C
au	0.00045	1	G		C		T	T	A	T	C	C	C
am	0.00045	1	G		T		T	T	A	T	C	C	C
at	0.00045	1	G		C		T	T	A	T	C	C	C
aw	0.00045	1	G		C		T	T	A	T	C	C	C
ay	0.00045	1	G		C		T	T	A	T	C	C	C
ar	0.00045	1	G		C		T	T	A	T	C	C	C
p	0.00045	1	G		T		T	T	A	T	T	C	C
as	0.00045	1	G		C		T	T	A	T	C	C	C
ae	0.00045	1	G		C		T	T	A	T	T	T	C
ai	0.00045	1	G		T		T	T	A	T	C	T	C

Table 13.33 - IGHV1-69

h.type	cum.freq	# indvs	pops	14:107170185	14:107170326	14:107170343
				rs181067247	.	rs190658595
ref	0.49795	-	-	Pro->Thr	->Ala	Phe->Val
a	0.1689	323	14	G	AGCT	A
b	0.1543	312	14	G	AGCT	A
c	0.0561	118	13	G	AGCT	A
g	0.0348	76	7	G	AGCT	A
e	0.014	31	4	G	AGCT	A
l	0.0108	24	6	G	AGCT	A
d	0.00995	22	3	G	AGCT	A
h	0.00765	17	10	G	AGCT	A
f	0.00685	15	8	G	AGCT	A
i	0.00675	15	6	G	AGCT	A
m	0.0027	6	3	G		A
j	0.00225	5	3	G	AGCT	A
u	0.00225	5	4	G	AGCT	A
s	0.0018	4	3	G	AGCT	A
k	0.00135	3	1	G	AGCT	A
ac	0.00135	3	3	G	AGCT	A
ah	0.0009	2	2	G	AGCT	A
t	0.0009	2	2	G	AGCT	A
ao	0.0009	2	1	G	AGCT	A
af	0.0009	2	2	G	AGCT	A
ad	0.0009	2	2	G	AGCT	A
n	0.0009	2	2	G	AGCT	A
an	0.0009	2	2	G	AGCT	A
o	0.0009	2	2	G	AGCT	A
ba	0.00045	1	1	T	AGCT	A
ax	0.00045	1	1	G	AGCT	A
ag	0.00045	1	1	G	AGCT	A
y	0.00045	1	1	G	AGCT	A

ENST00000390633

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,5,10,11,12,13,14

1,10,12,14

1,3,8,10,12,14

1,10,14

1,2,3,4,6,7,10,11,12,13

1,3,5,6,9,10,11,14

3,4,5,9,10,12

10,12,14

5,10,14

1,2,8,14

6,8,10

14

2,10,14

11,13

2,9

10

10,14

2,7

2,3

10,14

3,10

13

10

13

10

Table 13.33 - IGHV1-69

az	0.00045	1	G	AGCT	A	10
aj	0.00045	1	G	AGCT	A	7
aa	0.00045	1	G	AGCT	A	2
aq	0.00045	1	G	AGCT	A	9
v	0.00045	1	G	AGCT	A	11
al	0.00045	1	G	■	A	10
ab	0.00045	1	G	AGCT	A	14
ak	0.00045	1	G	AGCT	A	4
q	0.00045	1	G	AGCT	A	4
ap	0.00045	1	G	AGCT	A	13
z	0.00045	1	G	AGCT	A	14
w	0.00045	1	G	AGCT	A	14
av	0.00045	1	G	AGCT	A	4
r	0.00045	1	G	AGCT	A	8
x	0.00045	1	G	AGCT	A	2
au	0.00045	1	G	AGCT	A	2
am	0.00045	1	G	AGCT	A	1
at	0.00045	1	G	AGCT	A	3
aw	0.00045	1	G	AGCT	A	2
ay	0.00045	1	G	AGCT	A	10
ar	0.00045	1	G	AGCT	C	3
p	0.00045	1	G	AGCT	A	13
as	0.00045	1	G	AGCT	A	12
ae	0.00045	1	G	AGCT	A	13
ai	0.00045	1	G	AGCT	A	11

Table 13.34 - IGHV1-8

h.type	cum.freq	# indivs	pops	14:106539147	14:106539163	14:106539171	14:106539278	14:106539284	14:106539291	14:106539476	14:106539497
				rs148477632	rs73371428	rs187649961	rs184294529	rs151035485	rs187522932	rs189071216	rs180817521
ref				Ile->Val	Met->Ile	Val->Ile	Tyr->Phe	Thr->Ile	Thr->Pro	Ala->Thr	Ile->Phe
a	0.8844	-	-	T	C	C	T	G	T	C	T
c	0.1016	160	9	T		C	T	G	T	C	T
b	0.00405	8	2	C	C	C	T	G	T	C	T
d	0.0032	6	6	T	C	C	T	G	T	C	
e	0.0018	4	2	T		C	T		T	C	T
f	0.00135	3	2	T		C	T	G	T	C	
h	0.00135	3	2	T	C	T	T	G	T	C	T
j	0.0009	2	2	T	C	C		G	T	C	T
g	0.00045	1	1	T	C	C	T	G	G	C	T
i	0.00045	1	1	T		C	T	G	T	T	T
	0.00045	1	1	T	C	C	T		T	C	T

Table 13.34 - IGHV1-8

h.type	cum.fr	# indivs	pops	
ref	0.8844	-	-	ENST000000390599
a	0.1016	160	9	1,5,7,8,10,11,12,13,14
c	0.00405	8	2	10,13
b	0.0032	6	6	2,5,6,11,12,13
d	0.0018	4	2	1,14
e	0.00135	3	2	2,11
f	0.00135	3	2	7,11
h	0.0009	2	2	3,5
j	0.00045	1	1	2
g	0.00045	1	1	14
i	0.00045	1	1	13

Table 13.35 - IGHV2-26

h.type	cum. freq	# indivs	pops	14:106757655 rs12586893	14:106757706 rs185534639	14:106757798 rs181476099	14:106757831 rs111777969	14:106757865 rs185965403	14:106757880 rs146488407	14:106758074 rs186573875
ref	0.9245	-	-	Arg->Trp	Leu->Phe	Ala->Glu	Arg->His	Leu->Val	Val->Ile	Thr->Ala
				G	G	G	C	G	C	T
a	0.06135	126	13		G	G	C	G	C	T
b	0.0087	15	5	G	G	G	T	G	C	T
c	0.0032	7	2	G	G	T	C	C	C	T
f	0.0009	2	2	G	G	G	C	G	T	T
d	0.00045	1	1	G		G	C	G	C	T
g	0.00045	1	1	G	G	G	C	C	C	T
e	0.00045	1	1	G	G	G	C	G	C	C

ENST000000390611

1,3,4,5,6,7,8,9,10,11,1
2,13,14
1,6,10,11,14
10,14
13,14
1
5
10

Table 13.36 - IGHV2-5

h.type	cum.fr	# indivs	pops	14:106494234	14:106494245	14:106494248	14:106494251	14:106494269	14:106494325	14:106494346	14:106494404
ref	0.3179	-	-	rs1065554	rs1065552	rs181774040	rs141264800	rs12895651	rs150364725	rs185589849	rs190103562
				Ser->Arg	Ser->Thr	Pro->Thr	Ser->Gly	Asn->Asp	Gly->Ala	Ser->Thr	Leu->Val
a	0.6657	872	14	G	A	G	T	T	C	C	G
b	0.01055	18	3	G	A	G	C	C	C	C	G
d	0.0018	4	2	G	A	T	T	C	C	C	G
c	0.0009	2	2	G	A	G	T	C	G	C	G
j	0.00045	1	1	G	T	T	T	C	C	C	G
k	0.00045	1	1	G	A	G	T	T	C	G	G
g	0.00045	1	1	C	A	G	T	C	G	C	G
e	0.00045	1	1	G	A	G	T	C	C	G	G
h	0.00045	1	1	C	A	T	T	C	C	C	G
f	0.00045	1	1	G	A	G	T	C	C	C	C
i	0.00045	1	1	C	A	G	T	C	C	C	G

ENST00000390597

1,2,3,4,5,6,7,8,9,10,11,1
2,13,14
1,10,14
10,14
4,13
6
1
9
3
5
10
12

Table 13.37 - IGHV2-70

h.type	cum.freq	# indvs	pops	14:107178820	14:107178837	14:107178849	14:107178875	14:107178878	14:107178890	14:107178899	14:107178911	14:107178938
				rs139383686	rs144976346	rs185317654	rs188731942	rs192620545	rs138836314	rs112655724	rs188243654	rs17113976
ref	0.36095	-	-	Ile->Met	Tyr->Asp	Asp->Tyr	Leu->Pro	Val->Ala	Lys->Arg	Asp->Ala	Thr->Ile	Tyr->Cys
a	0.31	584	14	T	A	C	A	A	T	T	G	T
b	0.16595	353	14	T	A	C	A	A	T	T	G	T
d	0.06035	132	13	T	A	C	A	A	T	G	G	C
c	0.04295	94	11	T	A	C	A	A	T	T	G	T
e	0.0319	70	6	T	A	C	A	A	T	T	G	C
f	0.00405	9	5	T	A	C	A	A	T	T	G	C
l	0.00225	5	4	T	A	C	G	A	T	T	G	T
y	0.00135	3	3	T	A	C	A	A	T	T	G	T
z	0.00135	3	3	C	A	C	A	A	T	T	G	T
af	0.00135	3	2	T	A	C	A	A	T	G	G	C
h	0.00135	3	3	T	A	C	A	A	T	G	G	C
g	0.0009	2	2	T	A	C	A	A	T	T	G	T
s	0.0009	2	2	T	A	C	A	A	T	T	G	T
ab	0.0009	2	2	T	A	C	A	A	C	T	G	T
ah	0.00045	1	1	C	C	C	A	A	T	T	G	C
ag	0.00045	1	1	T	A	C	A	A	T	G	G	T
j	0.00045	1	1	T	C	C	A	A	T	T	G	C
u	0.00045	1	1	T	A	C	A	A	T	T	G	C
k	0.00045	1	1	T	A	C	A	A	T	T	G	T
aj	0.00045	1	1	C	A	C	A	A	T	T	G	T
aa	0.00045	1	1	T	A	C	A	A	T	T	G	C
t	0.00045	1	1	T	A	C	A	A	T	T	G	C
aq	0.00045	1	1	T	A	C	A	A	T	G	G	C
v	0.00045	1	1	C	A	C	A	A	T	T	G	T
al	0.00045	1	1	T	A	C	A	A	C	T	G	T
ak	0.00045	1	1	T	A	C	A	A	T	T	G	C
q	0.00045	1	1	T	A	C	A	A	T	T	G	T
ap	0.00045	1	1	T	A	C	A	A	T	G	G	C

Table 13.37 - IGHV2-70

w	0.00045	1	T	C	C	A	A	T	T	G	T	T
ao	0.00045	1	T	A	C	A	A	T	T	G	G	C
r	0.00045	1	T	A	C	A	A	T	T	G	G	T
x	0.00045	1	C	A	C	A	A	T	T	T		T
ad	0.00045	1	T	A	C	A	G	T	T	G	G	C
am	0.00045	1	T	A	C	A	A	C	T	T	G	C
i	0.00045	1	T	A	C	A	A	T	T	T	G	T
n	0.00045	1	T	A		A	A	T	T	T	G	T
ac	0.00045	1	C	A	C	A	A	T	T	G	G	C
m	0.00045	1	T	A	C	A	A	T	T	T		T
ar	0.00045	1	T	A	C	A	A	T	T	T	G	T
p	0.00045	1	T	A	C	A	A	T	T	T	G	T
ae	0.00045	1	T	A	C	A	A	T	T	T		T
ai	0.00045	1	T	A	C	A	A	T	T	T	G	C
an	0.00045	1	T	A	C	A	A	T	T	T	G	T
o	0.00045	1	T	C	C	A	A	T	T	G	G	C

Table 13.37 - IGHV2-70

h.type	cum.freq	# indvs	pops	14:107178959	14:107178965	14:107178976	14:107179098	14:107179103	14:107179106	14:107179115	14:107179123	14:107179243
				rs184814603	rs2073669	rs148856487	rs191880827	rs185635448	rs10144703	rs182292946	rs187704106	rs61734101
ref	0.36095	-	-	Asp->Gly	Leu->Arg	Glu->Asp	Gly->Ser	Glu->Val	Arg->Lys	Val->Gly	Leu->Phe	Thr->Met
a	0.31	584	14	T	A	C	C	T	C	A	T	G
b	0.16595	353	14	T	C	C	C	T	C	A	T	G
d	0.06035	132	13	T	C	C	C	T	T	A	T	
c	0.04295	94	11	T	A	C	C	T	T	A	T	G
e	0.0319	70	6	T	C	C	C	T	T	A	T	
f	0.00405	9	5	T	C	C	C	T	T	A	T	G
l	0.00225	5	4	T	C	C	C	T	C	A	T	G
y	0.00135	3	3	C	C	C	C	T	C	A	T	G
z	0.00135	3	3	T	C	C	C	T	T	A	T	G
af	0.00135	3	2	T	C	C	C	T	T	A	T	G
h	0.00135	3	3	T	C	C	C	T	T	A	T	
g	0.0009	2	2	T	A	G	C	T	C	A	T	G
s	0.0009	2	2	T	C	C	T	T	C	A	T	G
ab	0.0009	2	2	T	A	C	C	T	C	A	T	G
ah	0.00045	1	1	T	C	C	C	T	C	A	T	G
ag	0.00045	1	1	C	C	C	C	T	T	A	T	
j	0.00045	1	1	T	C	C	C	T	C	A	T	G
u	0.00045	1	1	T	A	C	C	T	C	A	T	G
k	0.00045	1	1	T	A	G	C	T	T	A	T	G
aj	0.00045	1	1	T	C	C	C	T	C	A	T	G
aa	0.00045	1	1	T	C	C	C	T	T	A	T	
t	0.00045	1	1	T	A	C	C	T	T	A	T	G
aq	0.00045	1	1	T	C	C	C		T	A	T	
v	0.00045	1	1	T	A	C	C	T	C	A	T	G
al	0.00045	1	1	T	C	C	C	T	C	A	T	G
ak	0.00045	1	1	T	G	C	C	T	T	A		
q	0.00045	1	1	T	C	C	C	T	C	A		G
ap	0.00045	1	1	T	C	C	C	T	C	A	T	

Table 13.37 - IGHV2-70

w	0.00045	1	T	A	C	C	C	T	C	A	T	G
ao	0.00045	1	T	C	C	C	C	T	T	A	T	G
r	0.00045	1	T	A	C	C	C	T	T	A	T	G
x	0.00045	1	T	C	C	C	C	T	C	A	T	G
ad	0.00045	1	T	C	C	C	C	T	T	A	T	
am	0.00045	1	T	C	C	C	C	T	T	A	T	
i	0.00045	1	T	A	C	C	C	T	C	C	T	G
n	0.00045	1	T	A	C	C	C	T	C	A	T	G
ac	0.00045	1	T	C	C	C	C	T	T	A	T	
m	0.00045	1	T	C	C	C	C	T	T	A	T	G
ar	0.00045	1	C	C	G	C	C	T	C	A	T	G
p	0.00045	1	T	C	C	C	C		C	A	T	G
ae	0.00045	1	C	C	C	C	C	T	T	A	T	G
ai	0.00045	1	T	C	C	C	C	T	C	A	T	G
an	0.00045	1	C	A	C	C	C	T	C	A	T	G
o	0.00045	1	T	C	C	C	C	T	T	A	T	

Table 13.37 - IGHV2-70

h.type	cum.freq	# indivs	pops	14:107179254	ENST00000390634
				rs61734098	
				Ile->Met	
ref	0.36095	-	-	T	
a	0.31	584	14	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.16595	353	14	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14
d	0.06035	132	13	C	1,2,3,4,5,6,7,9,10,11,12,13,14
c	0.04295	94	11	T	1,2,5,6,7,8,10,11,12,13,14
e	0.0319	70	6	C	1,5,10,12,13,14
f	0.00405	9	5	T	1,5,9,10,14
l	0.00225	5	4	T	2,3,4,10
y	0.00135	3	3	T	7,9,12
z	0.00135	3	3	T	10,13,14
af	0.00135	3	2	T	6,10
h	0.00135	3	3	T	2,10,13
g	0.0009	2	2	T	4,6
s	0.0009	2	2	T	5,6
ab	0.0009	2	2	T	2,5
ah	0.00045	1	1	T	2
ag	0.00045	1	1	C	10
j	0.00045	1	1	T	10
u	0.00045	1	1	T	10
k	0.00045	1	1	T	13
aj	0.00045	1	1	T	10
aa	0.00045	1	1	T	14
t	0.00045	1	1	T	10
aq	0.00045	1	1	C	4
v	0.00045	1	1	T	13
al	0.00045	1	1	T	14
ak	0.00045	1	1	C	10
q	0.00045	1	1	T	1
ap	0.00045	1	1	C	10

Table 13.37 - IGHV2-70

w	0.00045	1	T	13
ao	0.00045	1	C	1
r	0.00045	1	T	2
x	0.00045	1	T	4
ad	0.00045	1	C	12
am	0.00045	1	C	1
i	0.00045	1	T	4
n	0.00045	1	T	5
ac	0.00045	1	T	13
m	0.00045	1	T	12
ar	0.00045	1	T	11
p	0.00045	1	T	4
ae	0.00045	1	T	13
ai	0.00045	1	T	9
an	0.00045	1	T	13
o	0.00045	1	C	14

[illegible]

Table 13.39 - IGHV3-13

h.type	cum.freq	# indvs	pops	14:106586159	14:106586165	14:106586168	14:106586231	14:106586259	14:106586267	14:106586354	14:106586389
ref	0.69725	-	-	rs192397984	rs140926945	rs184992069	rs10151633	rs11625174	rs189564282	rs10131161	rs181545751
a	0.15245	276	13	Thr->Met	Gly->Glu	Ala->Gly	Arg->Gln	Thr->Pro	Ala->Val	Ser->Cys	Gln->His
b	0.1011	167	9	G	C	G	C	G	G	G	C
c	0.02925	59	10	G	T	G	C	T	G	G	C
d	0.0105	19	4		C	G	T	T	G	C	C
e	0.00225	5	3	G	C	G	C	T	G	C	C
j	0.0018	4	2	G	C	G	T	T	G	G	C
g	0.00135	3	2	G	C	G	C	T	G	G	
f	0.0009	2	2	G	C	G	T	T		C	C
k	0.00045	1	1	G	T	G	C	G	G	G	C
h	0.00045	1	1		C	G	C	T	G	G	C
i	0.00045	1	1		C	G	T	T	G	C	
n	0.00045	1	1	G	C	G	C	G	G	G	C
m	0.00045	1	1	G	C	G	C	T	G	G	C
l	0.00045	1	1	G	C	C	C	T	G	G	C
o	0.00045	1	1	G	T	G	T	T	G	C	C

Table 13.39 - IGHV3-13

h.type	cum.freq	# indvs	pops	14:106586557	14:106586563
ref	0.69725	-	-	Val->Ile	Phe->Leu
a	0.15245	276	13	C	A
b	0.1011	167	9	C	A
c	0.02925	59	10	C	A
d	0.0105	19	4	C	A
e	0.00225	5	3	C	A
j	0.0018	4	2	C	A
g	0.00135	3	2	C	A
f	0.0009	2	2	C	A
k	0.00045	1	1	C	A
h	0.00045	1	1	C	A
i	0.00045	1	1	C	A
n	0.00045	1	1	C	G
m	0.00045	1	1	T	G
l	0.00045	1	1	C	A
o	0.00045	1	1	C	A

ENST000000390602

1,2,3,4,5,6,7,8,9,10,11,12,13

1,5,7,8,10,11,12,13,14

1,2,5,6,7,10,11,12,13,14

1,10,12,14

1,10,14

1,10

6,14

10,12

2

1

14

11

14

3

10

Table 13.40 - IGHV3-15

h.type	cum.freq	# indvs	pops	14:106610377	14:106610467	14:106610509	14:106610570	14:106610736
				rs182550955	rs186087290	rs147758697	rs184104714	rs183000727
				Asn->Thr	Gly->Ala	Ser->Asn	Gly->Arg	Ile->Phe
ref	0.91495	-	-	T	C	C	C	T
a	0.0572	104	11	T	C	T	C	T
b	0.0265	56	5	T	C	C	T	T
d	0.00045	1	1	T	C	C	C	■
e	0.00045	1	1	G	C	C	C	T
c	0.00045	1	1	T	G	C	C	T
				ENST000000390603				
				1,2,3,5,6,7,10,11,12,13,14				
				1,3,4,9,13				
				6				
				6				
				9				

Table 13.41 - IGHV3-16

h.type	cum.fre	# indivs	pops	14:106622052	14:106622073	14:106622076	14:106622132
				rs12323556	rs142354696	rs12433709	rs12323559
				Glu->Lys	Lys->Glu	Arg->Cys	Arg->Ile
ref	0.8039	-	-	C	T	G	C
a	0.11075	200	13	C	T		C
b	0.07085	122	8	T	T	G	
c	0.01	21	4	C	C	G	
d	0.00315	7	3	T	T	G	C
e	0.00135	3	2	C	C	G	C

ENST000000390604

1,2,3,4,5,6,7,8,9,11,12,13,14

1,5,7,10,11,12,13,14

1,10,11,14

10,11,14

10,12

Table 13.42 - IGHV3-20

h.type	cum.freq	# indivs	pops	14:106667592	14:106667756	14:106667785	14:106667786	14:106667810	14:106668011
				rs3751514	rs190163710	rs181860901	rs3751513	rs112170273	rs180685266
				His->Tyr	Ala->Val	Asp->Glu	Asp->Gly	Cys->Phe	Val->Ala
ref	0.5693	-	-	G	G	A	T	C	A
a	0.2829	470	14		G	A	T	C	A
b	0.1209	221	13	G	G	A	T		A
c	0.02285	50	6		G	A	C	C	A
f	0.00135	3	3	G	G	A	C	C	A
d	0.0009	2	2		G	A	T		A
g	0.00045	1	1		G	A	T	C	G
e	0.00045	1	1	G	G	A	T		G
h	0.00045	1	1	G		A	T	C	A
i	0.00045	1	1	G		T	T	C	A

ENST000000390606

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13

1,3,4,9,10,14

9,11,14

5,11

10

2

14

1

Table 13.43 - IGHV3-21

h.type	cum.freq	# indivs	pops	14:106691762	14:106691807	14:106691824	14:106691833	14:106691849	14:106691865	14:106691867	14:106691879
				rs183314182	rs188058888	rs143336904	rs192148312	rs184550642	rs192949902	rs184209619	rs180974163
				Thr->Ala	Ser->Gly	Val->Ala	Leu->Pro	Ala->Thr	Met->Ile	Met->Leu	Ser->Gly
ref	0.9892	-	-	T	T	A	A	C	C	T	T
a	0.00225	5		T	C	A	A	C	C	T	T
d	0.0018	4		T	T	A	A	C	C	T	T
c	0.0009	2		T	T	G	A	C	C	T	T
b	0.0009	1		T	T	A	A	C	C	T	T
h	0.0009	2		C	T	A	A	C	C	T	T
j	0.00045	1		T	T	A	G	C	C	T	T
k	0.00045	1		T	T	A	A	C	C	T	T
g	0.00045	1		T	C	A	A	C	C	T	T
e	0.00045	1		T	T	A	A	C	C	T	T
f	0.00045	1		T	T	A	A	C	C	T	T
i	0.00045	1		T	T	A	A	C	G	T	C
n	0.00045	1		T	C	A	A	T	C	T	T
m	0.00045	1		T	T	A	A	C	C	T	T
l	0.00045	1		T	T	A	A	C	C	T	T

Table 13.43 - IGHV3-21

h.type	cum.freq	# indivs	pops	14:106691884	14:106691896	14:106691899	14:106691909
				rs184687518	rs187020320	rs191067794	rs181216108
ref	0.9892	-	-	Thr->Ile	Ala->Gly	Ala->Val	Leu->Phe
a	0.00225	5	5	G	G	G	G
d	0.0018	4	3		G	G	G
c	0.0009	2	2	G	G	G	G
b	0.0009	1	1		C	G	G
h	0.0009	2	2	G	G	G	G
j	0.00045	1	1		G	G	G
k	0.00045	1	1	G	G		G
g	0.00045	1	1		G	G	G
e	0.00045	1	1	G	G	G	G
f	0.00045	1	1	G	C		
i	0.00045	1	1	G	G	G	G
n	0.00045	1	1	G	G	G	G
m	0.00045	1	1	G	C		G
l	0.00045	1	1	G	C	G	G

ENST000000390607

7,9,11,13,14

1,5,14

10,14

1

1,13

2

1

5

2

4

14

3

11

1

Table 13.44 - IGHV3-23

h.type	cum.freq	# indvs	pops	14:106725320	14:106725328	14:106725329	14:106725346	14:106725357	14:106725397	14:106725400	14:106725433
ref	0.83265	-	-	rs189958807	rs1064091	rs1064090	rs1055799	rs143624256	rs61752504	rs183319819	rs187384491
				Tyr->His	Gly->Asp	Gly->Ser	Ala->Val	Glu->Asp	Ala->Gly	Tyr->Ser	Ser->Phe
a	0.09825	190	12	A	C	C	G	C	G	T	G
b	0.02695	58	9	A	C	T	G	C	G	T	G
c	0.01365	29	9	A	C	T	G	C	G	T	G
d	0.00865	18	4	A	C	C	G	C	C	T	G
e	0.00455	10	4	A	C	T		C	G	T	G
f	0.00225	5	3	A	C	C	G	C	C	T	G
k	0.00225	5	5	A	T	C	G	C	G	T	G
g	0.00225	5	3	A	C	C		C	G	T	G
n	0.0018	4	3	A	C	C	G	C	C	T	G
j	0.0009	2	1	A	C	C		C	G	T	G
h	0.0009	2	2	A	C	C	G	C	G	G	G
u	0.00045	1	1	A	T	C	G	C	G	T	G
t	0.00045	1	1	A	T	C	G	G	G	T	G
v	0.00045	1	1	A	C	C	G	C	G	T	G
s	0.00045	1	1	A	T	C	G	C	C	T	G
q	0.00045	1	1	A	C	C	G	C	G	T	G
r	0.00045	1	1	A	C	T	G	C	G	T	G
i	0.00045	1	1	A	C	C	G	C	G	T	G
m	0.00045	1	1	A	C	C	G	C	G	T	G
l	0.00045	1	1	G	C	C	G	C	G	T	G
p	0.00045	1	1	A	T	C	G	C	G	G	G
o	0.00045	1	1	A	C	C	G	C	G	T	

Table 13.44 - IGHV3-23

h.type	cum.freq	# indvs	pops	14:106725446 rs61750837	14:106725482 rs56069819	14:106725490 rs191754191	14:106725493 rs182117479	14:106725621 rs188250527	14:106725630 rs145913230	14:106725634 rs181072648
ref	0.83265	-	-	Ser->Thr	Leu->Val	Val->Gly	Glu->Gly	Ala->Pro	Phe->Val	Trp->Cys
a	0.09825	190	12	A	A	A	T	C	A	C
b	0.02695	58	9	A	C	A	T	C	A	C
c	0.01365	29	9	A	A	A	T	C	A	C
d	0.00865	18	4	T	C	A	T	C	A	C
e	0.00455	10	4	A	C	A	T	C	A	C
f	0.00225	5	3	T	A	A	T	C	A	C
k	0.00225	5	5	A	A	A	T	C	A	C
g	0.00225	5	3	A	A	A	T	C	A	C
n	0.0018	4	3	A	A	A	T	C	A	C
j	0.0009	2	1	A	C	A	T	C	A	C
h	0.0009	2	2	A	A	A	T	C	A	C
u	0.00045	1	1	A	C	A	T	C	A	C
t	0.00045	1	1	A	A	A	T	C	A	C
v	0.00045	1	1	A	A	A	C	G	A	C
s	0.00045	1	1	T	C	A	T	C	A	C
q	0.00045	1	1	A	A	A	T	C	A	G
r	0.00045	1	1	A	C	A	T	C	A	C
i	0.00045	1	1	A	A	A	T	C	C	C
m	0.00045	1	1	A	A	C	T	C	A	C
l	0.00045	1	1	A	A	A	T	C	A	C
p	0.00045	1	1	A	A	A	T	C	A	C
o	0.00045	1	1	A	A	A	T	C	A	C

ENST000000390609

1,2,3,5,6,7,8,10,11,12,13,14

1,3,4,5,9,10,11,12,14

1,3,4,5,6,9,10,11,14

1,5,10,14

1,10,11,14

1,7,10

1,5,9,10,11

1,5,11

2,9,14

14

6,10

14

13

8

10

14

14

9

6

10

2

2

[illegible]

Table 13.46 - IGHV3-33

h.type		cum.fre	# indvs	pops	14:106815855	14:106815858	14:106815862	14:106815868	14:106815924	14:106815933	14:106815948	14:106815970
ref	0.82905 -	-			Asp->Gly	Tyr->Ser	Trp->Arg	Val->Phe	Ser->Asn	Thr->Ile	Ala->Val	Arg->Gly
					rs183940317	rs187379689	rs112373679	rs111734923	rs192341521	rs187304239	rs192234669	rs138488134
a	0.145	243	12		T	T	A	C	C	G	G	T
b	0.00965	21	10		T	T	G		C	G	G	T
c	0.0082	18	6		T	T	A	C	C	G	G	C
e	0.0027	6	3		C	T	A	C	C	G	G	T
d	0.0018	4	3		T	T	A	C	C		G	T
h	0.00135	3	3		T	T	A	C	T	G	G	T
f	0.0009	2	2		T	G	A	C	C	G	G	T
j	0.00045	1	1		C	T	A	C	T	G		T
g	0.00045	1	1		C	T	G		C	G	G	C
i	0.00045	1	1		T	T	G		T	G	G	C

ENST000000390
615

1,2,4,5,6,7,8,10,
11,12,13,14
1,2,5,6,7,10,11,
12,13,14
1,2,7,10,12,14
1,10,14
2,13,14
1,6,10
5,8
10
10
1

h.type	cum.frec	# indvs	pops	14:106845417	14:106845487	14:106845753
ref	0.78085	-	-	C	C	C
a	0.11765	227	14	T	C	C
b	0.09695	190	10	C	G	C
c	0.0041	9	5	C	C	G
d	0.00045	1	1	T	G	C
				Arg->Gln	Gly->Arg	Val->Leu
				ENST00000390617		
				1,2,3,4,5,6,7,8,9,10,11,12,13,14		
				1,3,4,5,6,7,9,10,12,14		
				1,2,5,7,8		
				14		

Table 13.48 - IGHV3-38

h.type	cum.freq	# indvs	pops	14:106866449				14:106866451		14:106866664		14:106866828	
				rs78850219	Asn->Ser	Asn->Lys	Leu->Phe	rs144366955	rs150283006	rs112981911	Leu->Pro	ENST000000390618	
ref	0.8889	-	-	T	T	G	C	G	C	A			
a	0.0723	158	14	C	C	G	C	G	C	A		1,2,3,4,5,6,7,8,9,10,11,12,13,14	
b	0.02515	52	6	C	C	C	C	C	C	A		1,5,10,11,12,14	
c	0.00725	16	5	T	T	G	G	G	G	A		1,10,11,12,14	
d	0.0064	13	3	T	T	G	C	G	C	G		2,7,13	

Table 13.49 - IGHV3-43

h.type	cum.freq	# indivs	pops	14:106926211	14:106926223	14:106926231	14:106926323	14:106926328	14:106926388	14:106926435	14:106926454
				rs116899367	rs2467912	rs182842897	rs186760979	rs61999676	rs2467910	rs182650845	rs188914311
ref	0.57745	-	-	Ala->Thr	Thr->Ala	Ser->Asn	Asp->Glu	Trp->Gly	Thr->Ala	Ser->Phe	Val->Met
a	0.2428	512	14	C	T	C	A	A	T	G	C
b	0.08365	164	13	C	C	C	A	A	C	G	C
c	0.0302	66	10	C	T	C	A	C	T	G	C
e	0.0182	40	11	C	T	C	A	A	C	G	C
d	0.01455	30	4	T	T	C	A	A	T	G	C
g	0.00995	22	7	C	T	C	A	C	T	G	C
h	0.0064	13	5	C	C	C	A	C	C	G	C
f	0.0055	12	5	C	T	C	A	A	T		C
i	0.00365	8	3	C	C	C	A	A	C		C
j	0.00225	5	4	C	T	C	A	C	C	G	C
k	0.00135	3	2	C	T	T	A	A	T	G	C
m	0.00135	3	3	C	C	C	A	A	T	G	C
p	0.0009	2	2	C	C	C	A	A	C	G	T
q	0.00045	1	1	C	T	C	T	A	T	G	C
n	0.00045	1	1	C	T	C	A	A	C		C
l	0.00045	1	1	C	T	C	A	A	C	G	C
o	0.00045	1	1	C	C	C	A	A	C	G	C

Table 13.49 - IGHV3-43

h.type	cum.freq	# indivs	pops	14:106926456	14:106926481
				rs79008247	rs111739001
ref	0.57745	-	-	Val->Gly	Val->Met
a	0.2428	512	14	A	C
b	0.08365	164	13	C	C
c	0.0302	66	10	C	C
e	0.0182	40	11	A	C
d	0.01455	30	4	A	C
g	0.00995	22	7	A	C
h	0.0064	13	5	A	T
f	0.0055	12	5	A	C
i	0.00365	8	3	A	C
j	0.00225	5	4	A	C
k	0.00135	3	2	A	C
m	0.00135	3	3	A	C
p	0.0009	2	2	A	C
q	0.00045	1	1	A	C
n	0.00045	1	1	A	C
l	0.00045	1	1	C	C
o	0.00045	1	1	C	C

ENST000000434710

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,3,4,5,6,7,9,11,12,13

1,3,4,5,6,8,10,11,12,13,14

2,5,6,7

2,3,4,6,9,10,11

1,2,6,7,13

1,7,10,12,14

1,10,14

3,4,9,10

10,14

2,7,13

1,10

13

10

10

9

Table 13.50 - IGHV3-48

h.type	cum.freq	# indvs	pops	14:106993845 rs7148408	14:106993938 .	14:106993945 rs7148607	14:106993996 rs183130922	14:106994231 rs192884486	14:106994237 rs184894692	14:106994262 rs188639638
ref	0.3919	-	-	Asp->Ala T	->Thr GTAC	Ser->Gly T	Arg->Cys G	Ile->Val T	Val->Leu C	Glu->Asp C
a	0.3355	589	14	G	GTAC	T	G	T	C	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.20455	404	14	G	GTAC	C	G	T	C	1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.027	59	11	G	G	T	G	T	C	1,2,4,5,6,7,10,11,12,13,14
d	0.01595	35	11	T	GTAC	C	G	T	C	1,2,3,4,5,6,7,8,9,10,13
f	0.0128	28	9	G	G	C	G	T	C	1,3,4,6,7,9,10,11,14
e	0.0105	23	10	T	G	T	G	T	C	2,3,4,5,6,7,10,11,13,14
j	0.00045	1	1	T	GTAC	T		T	C	4
g	0.00045	1	1	T	GTAC	T	G	T	G	11
h	0.00045	1	1	G	GTAC	T	G	C	C	5
i	0.00045	1	1	G	GTAC	T	G	T	C	2
										1

Table 13.51 - IGHV3-49

h.type	cum.frec	# indivs	pops	14:107012994	14:107013065	14:107013072	14:107013093	14:107013129	14:107013146	14:107013171	14:107013201
				rs140678750	rs140027079	rs191590980	rs187070171	rs2073674	rs184357967	rs193139111	rs2073673
ref			-	Tyr->His	Gly->Glu	Tyr->Asp	Gly->Ser	Phe->Val	Asp->Ala	Thr->Ala	Gln->Lys
a	0.322	663	14	A	C	A	C	A	T	T	G
b	0.38705	499	13	A	C	A	C	C	T	T	G
c	0.28555	6	3	A	C	A	C	A	T	T	T
d	0.0027	1	1	G	C	C	T	C	G	T	G
g	0.00045	1	1	A	C	A	C	A	T	C	G
e	0.00045	1	1	A	C	A	C	C	T	T	G
h	0.00045	1	1	G	C	A	C	C	T	C	G
f	0.00045	1	1	A	C	A	C	C	T	T	G
i	0.00045	1	1	A	T	A	C	A	T	T	G

Table 13.51 - IGHV3-49

h.type	cum.freq	# indivs	pops	14:107013209	14:107013210	
				rs184512071	rs189168917	
ref	0.322	-	-	Gly->Asp	Gly->Ser	ENST00000390625
a	0.38705	663	14	C	C	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.28555	499	13	C	C	1,2,3,4,5,6,7,8,9,10,11,12,13
c	0.0027	6	3	C	C	7,11,13
d	0.00045	1	1	C	C	2
g	0.00045	1	1	C	C	1
e	0.00045	1	1	C	C	4
h	0.00045	1	1	C	C	1
f	0.00045	1	1	T	T	2
i	0.00045	1	1	C	C	10

Table 13.52 - IGHV3-53

h.type	cum.freq	# indivs	pops	14:107048717	14:107048749	14:107048795	14:107048796	14:107048929	14:107048944	14:107049080
ref	0.4053	-	-	Met->Ile	Asp->His	Ser->	Ser->Asn	Ile->Val	Ser->Thr	Leu->Ser
a	0.2535	483	14	C	C	G	C	T	A	A
b	0.2264	399	13	C	G	G	C	C	A	A
c	0.0453	99	13	C	C	G	C	T	T	G
d	0.0247	54	6	T	C	G	C	C	A	A
e	0.015	33	9	C	C	G	C	T	T	G
f	0.0092	20	6	C	C		C	T	A	A
g	0.0046	10	5	C	C	G	C	C	A	G
k	0.0041	9	8	C	G	G	C	T	A	A
h	0.0036	8	2	T	C	G	C	C	A	A
j	0.0023	5	4	C	G		C	C	A	A
i	0.0018	4	4	C	C	G	C	T	A	G
n	0.0014	3	2	C	C		C	T	T	G
m	0.0014	3	3	C	C		C	C	A	A
q	0.0005	1	1	C	C	G	C	C	T	A
l	0.0005	1	1	C	G	G	T	C	A	A
p	0.0005	1	1	C	G		C	T	A	A
o	0.0005	1	1	T	C	G	C	C	T	G

ENST000000390627

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,5,10,12,13,14

1,2,5,7,8,10,11,12,14

3,4,9,10,11,12

7,10,11,13,14

3,6,7,9,11,12,13,14

10,14

1,3,4,9

8,11,13,14

2,10

5,9,10

14

3

4

14

Table 13.53 - IGHV3-64

h.type	cum.freq	# indivs	pops	14:107113770 rs187347757	14:107113851 rs2072045	14:107113877 rs190575084	14:107113902 rs138428007	14:107114009 rs11846079	14:107114176 rs182438044	14:107114189 rs139161493	14:107114193 rs2073670
ref	0.4221	-	-	Glu->Gln	Asp->Asn	Ser->Asn	Leu->Val	Glu->Gly	Trp->Gly	Glu->Asp	Met->Thr
a	0.47965	793	14	C	C	C	G	T	A	C	A
b	0.06225	131	13	C	T	C	G	C	A	C	G
c	0.02785	61	14	C	C	C	G	C	A	C	A
d	0.00275	6	4	C	T	C	G	T	A	C	G
e	0.0018	4	2	C	T	C	G	C	C	C	G
f	0.00135	3	1	C	T	C	C	C	C	C	G
h	0.0009	2	1	C	T	C	G	C	A	C	A
j	0.00045	1	1	C	C	C	G	T	A	G	A
g	0.00045	1	1	G	C	C	G	T	A	C	A
i	0.00045	1	1	C	T	T	G	T	A	C	A

Table 13.53 - IGHV3-64

h.type	cum.frec	# indvs	pops	
ref	0.4221	-	-	ENST00000454421
a	0.47965	793	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.06225	131	13	1,2,3,4,5,6,7,9,10,11,12,13,14
c	0.02785	61	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
d	0.00275	6	4	2,4,13,14
e	0.0018	4	2	1,10
f	0.00135	3	1	10
h	0.0009	2	1	12
j	0.00045	1	1	2
g	0.00045	1	1	3
i	0.00045	1	1	8

Table 13.54 - IGHV3-66

h.type	cum.freq	# indivs	pops	14:107131089	14:107131164	14:107131223	14:107131290
				rs77593173	rs6423677	rs183864069	rs149638514
ref	0.43755	-	-	Tyr->His	Cys->Gly	Met->Lys	Ile->Val
a	0.27865	502	14	A	A	A	T
b	0.12575	252	14	A	C	A	C
c	0.0649	140	13	A	A	A	C
d	0.03975	87	13	G	C	A	C
e	0.02745	60	14	G	A	A	T
f	0.021	46	14	A	C	A	T
g	0.0027	6	3	G	C	A	T
h	0.0009	2	2	G	C	T	C
j	0.00045	1	1	A	A	T	T
k	0.00045	1	1	A	C	T	C
i	0.00045	1	1	A	A	T	C

ENST000000390632

1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,9,10,11,12,13,14
1,2,3,4,5,6,7,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
10,13,14
3,4
5
4
3

Table 13.55 - IGHV3-72

h.type	cum.freq	# indivs	pops	14:107198966	14:107199062	14:107199136	14:107199164	14:107199222	14:107199355
				rs193196315	rs184752334	rs139291045	rs191373077	rs188380722	rs192371094
ref				Lys->Arg	Ser->Asn	His->Gln	Ala->Gly	Leu->Val	Ile->Leu
				T	C	G	G	G	T
a	0.9973	-	-	C	T	G	G	G	T
d	0.00045	1	1	T	C	C	C	G	T
e	0.00045	1	1	C	C	G	G	G	T
c	0.00045	1	1	T	C	G	G	C	T
b	0.00045	1	1	T	C	G	C	G	T
f	0.00045	1	1	T	C	G	G	G	G

ENST000000433072

5
2
11
2
1
10

Table 13.56 - IGHV3-73

h.type	cum.freq	# indivs	pops	14:107211000	14:107211135	14:107211176	14:107211199	14:107211371
				rs142927366	rs184224767	rs61750833	rs188470998	rs61752554
				Lys->Glu	Ala->Pro	Lys->Arg	Leu->Phe	Trp->Cys
ref	0.91625	-	-	T	C	T	C	C
a	0.05265	96	6	T	C	T	C	G
b	0.0293	60	5	T	C	C	C	C
c	0.00135	3	3	T	C	T		C
d	0.00045	1	1	C	G	T	C	C
				ENST000000390636				
				1,5,10,12,13,14				
				1,7,9,10,14				
				1,10,14				
				2				

[illegible]

Table 13.58 - IGHV3-7

h.type	cum.freq	# indivs	pops	14:106518545 rs185279906 Asn->Ser	14:106518602 rs185538237 Ser->Asn	14:106518686 rs180765266 Gln->Arg	14:106518840 rs190666315 Leu->Arg	
ref	0.9808	-	-	T	C	T	A	ENST000000390598
a	0.016	29	5	T	C	C	A	1,5,10,11,14
b	0.0023	5	4	T	T	T	A	5,9,13,14
d	0.00045	1	1	C	C	T	A	4
c	0.00045	1	1	T	C	T	C	13

Table 13.59 - IGHV3-9

h.type	cum.fr	# indivs	pops	14:106552299	14:106552310	14:106552409	14:106552412	14:106552415	14:106552436	14:106552463
				rs188663119	rs8020204	rs140101036	rs10141136	rs185242411	rs192281729	rs182143003
ref				Tyr->His	Thr->Met	Ile->Thr	Ser->Asn	Gly->Val	Ser->Leu	Ala->Val
a	0.84305	-	-	A	G	A	C	C	G	G
b	0.10975	167	9	A		A	C	C	G	G
c	0.0228	46	7	A	G	A	C	C	G	G
e	0.0064	12	5	A	G	G	C	C	G	G
d	0.00405	6	4	A	G	A	C	C	G	G
f	0.00315	7	7	A	G	A	T	C	G	G
g	0.00225	5	4	A	G	A	C	C	G	
h	0.0018	4	3	A	G	A	C	C	G	G
r	0.00135	2	1	A	G	A	C	C		G
o	0.0009	1	1	A	G	A	T	C	G	G
j	0.0009	2	1	G	G	A	C	C	G	G
k	0.00045	1	1	A		A	T	C	G	G
q	0.00045	1	1	A		A	C	C	G	G
i	0.00045	1	1	A		A	C	C	G	G
n	0.00045	1	1	A	G	A	C	C	G	G
m	0.00045	1	1	A	G	A	C	T	G	G
l	0.00045	1	1	A		A	C	T	G	G
p	0.00045	1	1	A	G	A	T	C	G	G

Table 13.59 - IGHV3-9

h.type	cum.fr	# indivs	pops	14:106552464	14:106552472	14:106552491	14:106552493	14:106552713	
				rs186818749	rs191640600	rs182569652	rs141844186	rs116484141	
ref				Ala->Pro	Val->Ala	Asp->Asn	Asp->Gly	Ser->Asn	ENST000000390600
a	0.84305	-	-	C	A	C	T	C	1,5,7,8,10,11,12,13,14
b	0.10975	167	9	C	A	C	T	C	3,4,6,7,9,11,13
c	0.0228	46	7	C	A	C	C	C	1,7,10,12,14
e	0.0064	12	5	C	A	C	T	C	1,3,12,14
d	0.00405	6	4	C	A	T	T	C	1,2,3,5,6,8,10
f	0.00315	7	7	C	A	C	T	C	6,7,9,13
g	0.00225	5	4	C	A	C	T	C	2,10,14
h	0.0018	4	3	G	A	C	T	C	1
r	0.00135	2	1	C	A	C	T	C	2
o	0.0009	1	1	C	A	C	T	C	13
j	0.0009	2	1	C	A	C	T	C	10
k	0.00045	1	1	C	A	C	T	C	10
q	0.00045	1	1	G	A	C	T	T	2
i	0.00045	1	1	C	A	C	T	C	14
n	0.00045	1	1	C	G	C	T	C	6
m	0.00045	1	1	C	A	C	T	C	11
l	0.00045	1	1	C	A	C	T	C	14
p	0.00045	1	1	C	A	C	C	C	3

Table 13.60 - IGHV4-28

h.type	cum.fr	# indivs	pops	14:106780542	14:106780632	14:106780634	14:106780667	14:106780760	14:106780797	14:106780817
ref	0.22145	-	-	rs113785338	rs112792995	rs34629512	rs8010702	rs181692209	rs186483759	rs8009554
a	0.41435	691	14	Val->Leu	Tyr->Asn	Thr->Ile	Glu->Ala	Asp->Gly	Leu->Val	Leu->TrpLeu->Trp
b	0.13715	250	14	C	A	G	T	T	G	A
c	0.07385	161	14	C	A		T	T	G	C
d	0.06485	136	13	C	A	G	T	T	G	C
f	0.0195	42	5		T	G	T	T	G	C
e	0.01685	37	10	C	A	G	T	T	G	A
g	0.01355	29	4		T	G	T	T	G	C
i	0.01185	22	6	C	A	G	G	T	G	C
h	0.01175	25	11	C	A		T	T	G	A
j	0.00405	9	5	C	A		T	T	G	A
k	0.00315	7	3		T	G	T	C	G	C
p	0.00225	5	3	C	A	G	G	T	G	C
s	0.0009	2	2	C	A	G	T	C	G	C
n	0.0009	2	2		A	G	T	T	G	C
u	0.00045	1	1	C	T	G	T	T	G	C
t	0.00045	1	1		A	G	T	T	G	A
v	0.00045	1	1	C	A	G	T	T	C	C
q	0.00045	1	1		T	G	T	C	G	A
r	0.00045	1	1		T	G	T	C	G	C
m	0.00045	1	1		T	G	G	T	G	C
l	0.00045	1	1		T	G	T	T	G	A
o	0.00045	1	1		T	G	T	T	G	A

Table 13.60 - IGHV4-28

h.type	cum. fre	# indivs	pops	
ref	0.22145	-	-	ENST00000390612
a	0.41435	691	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.13715	250	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.07385	161	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
d	0.06485	136	13	1,2,3,5,6,7,8,9,10,11,12,13,14
f	0.0195	42	5	1,10,11,12,14
e	0.01685	37	10	1,2,5,6,7,10,11,12,13,14
g	0.01355	29	4	1,5,10,14
i	0.01185	22	6	1,8,10,11,13,14
h	0.01175	25	11	1,2,3,5,7,8,9,10,11,13,14
j	0.00405	9	5	2,6,7,10,13
k	0.00315	7	3	1,10,14
p	0.00225	5	3	1,10,14
s	0.0009	2	2	1,11
n	0.0009	2	2	10,14
u	0.00045	1	1	10
t	0.00045	1	1	11
v	0.00045	1	1	13
q	0.00045	1	1	11
r	0.00045	1	1	1
m	0.00045	1	1	10
l	0.00045	1	1	10
o	0.00045	1	1	14

Table 13.61 - IGHV4-31

h.type	cum.fr	# indivs	pops	14:106805289	14:106805303	14:106805321	14:106805345	14:106805358	14:106805381	14:106805388	14:106805408
				rs188101179	rs145562667	rs192498675	rs185638910	rs190512954	rs77489245	rs149858616	rs61995642
ref				Val->Leu	Arg->Leu	Asn->Ser	Tyr->Ser	Ile->Leu	His->Pro	Arg->Cys	Gly->Asp
a	0.01175	-	-	C	C	T	T	T	T	G	C
b	0.5475	886	14	C	C	T	T	T	T	G	C
c	0.2035	420	14	C	C	T	T	T	G	G	T
d	0.1554	303	14	C	C	T	T	T	G	G	C
e	0.04565	83	11	C	■	T	T	T	T	G	C
f	0.00865	18	11	C	C	T	T	T	T	G	T
g	0.00865	19	7	C	C	T	T	T	G	G	T
h	0.0054	10	3	C	C	T	T	T	T	■	C
i	0.00315	7	4	C	■	T	T	T	T	G	C
j	0.00225	5	3	C	C	T	T	T	G	G	C
k	0.00135	3	2	C	C	T	T	T	T	■	C
l	0.0009	2	2	C	C	T	G	T	T	G	C
m	0.0009	2	2	C	■	T	T	T	G	G	C
n	0.0009	2	2	C	■	T	T	T	G	G	T
o	0.00045	1	1	C	C	T	T	G	T	G	C
p	0.00045	1	1	C	■	C	G	T	T	G	C
q	0.00045	1	1	C	C	T	T	T	G	G	T
r	0.00045	1	1	C	C	T	T	T	G	G	C
s	0.00045	1	1	C	C	T	T	T	G	G	C
t	0.00045	1	1	G	■	T	T	T	T	G	C
u	0.00045	1	1	C	C	T	T	T	G	G	C
v	0.00045	1	1	C	■	C	G	T	T	G	T
w	0.00045	1	1	C	C	T	T	T	G	G	T
x	0.00045	1	1	C	C	T	T	T	T	G	C
y	0.00045	1	1	C	C	T	T	T	G	G	C
z	0.00045	1	1	C	C	T	T	T	G	G	C
aa	0.00045	1	1	C	C	T	T	T	G	G	C
ab	0.00045	1	1	C	C	T	T	T	G	G	C
ac	0.00045	1	1	C	C	T	T	T	G	G	C
ad	0.00045	1	1	C	C	T	T	T	G	G	C
ae	0.00045	1	1	C	C	T	T	T	G	G	C
af	0.00045	1	1	C	C	T	T	T	G	G	C
ag	0.00045	1	1	C	C	T	T	T	G	G	C
ah	0.00045	1	1	C	C	T	T	T	G	G	C
ai	0.00045	1	1	C	C	T	T	T	G	G	C
aj	0.00045	1	1	C	C	T	T	T	G	G	C
ak	0.00045	1	1	C	C	T	T	T	G	G	C
al	0.00045	1	1	C	C	T	T	T	G	G	C
am	0.00045	1	1	C	C	T	T	T	G	G	C
an	0.00045	1	1	C	C	T	T	T	G	G	C
ao	0.00045	1	1	C	C	T	T	T	G	G	C
ap	0.00045	1	1	C	C	T	T	T	G	G	C
aq	0.00045	1	1	C	C	T	T	T	G	G	C
ar	0.00045	1	1	C	C	T	T	T	G	G	C
as	0.00045	1	1	C	C	T	T	T	G	G	C
at	0.00045	1	1	C	C	T	T	T	G	G	C
au	0.00045	1	1	C	C	T	T	T	G	G	C
av	0.00045	1	1	C	C	T	T	T	G	G	C
aw	0.00045	1	1	C	C	T	T	T	G	G	C
ax	0.00045	1	1	C	C	T	T	T	G	G	C
ay	0.00045	1	1	C	C	T	T	T	G	G	C
az	0.00045	1	1	C	C	T	T	T	G	G	C
ba	0.00045	1	1	C	C	T	T	T	G	G	C
bb	0.00045	1	1	C	C	T	T	T	G	G	C
bc	0.00045	1	1	C	C	T	T	T	G	G	C
bd	0.00045	1	1	C	C	T	T	T	G	G	C
be	0.00045	1	1	C	C	T	T	T	G	G	C
bf	0.00045	1	1	C	C	T	T	T	G	G	C
bg	0.00045	1	1	C	C	T	T	T	G	G	C
bh	0.00045	1	1	C	C	T	T	T	G	G	C
bi	0.00045	1	1	C	C	T	T	T	G	G	C
bj	0.00045	1	1	C	C	T	T	T	G	G	C
bk	0.00045	1	1	C	C	T	T	T	G	G	C
bl	0.00045	1	1	C	C	T	T	T	G	G	C
bm	0.00045	1	1	C	C	T	T	T	G	G	C
bn	0.00045	1	1	C	C	T	T	T	G	G	C
bo	0.00045	1	1	C	C	T	T	T	G	G	C

Table 13.61 - IGHV4-31

h.type	cum.fr	# indivs	pops	14:106805411 rs185736587 Gly->Asp	14:106805421 rs188982921 Ile->Phe	14:106805497 rs181045179 Gln->His	14:106805508 rs4462488 Pro->Ser	
ref	0.01175	-	-	C	T	C	G	ENST000000438142
a	0.5475	886	14	C	T	C		1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.2035	420	14	C	T	C		1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.1554	303	14	C	T	C		1,2,3,4,5,6,7,8,9,10,11,12,13,14
d	0.04565	83	11	C	T	C		1,2,5,6,7,8,10,11,12,13,14
e	0.00865	18	11	C	T	C		1,2,3,4,6,9,10,11,12,13,14
f	0.00865	19	7	C	T	C	G	1,3,6,10,11,12,14
i	0.0054	10	3	T		C		1,10,14
g	0.00315	7	4	C	T	C	G	1,2,5,7
j	0.00225	5	3	C	T	C	G	2,10,14
h	0.00135	3	2	C	T	C		1,10
t	0.0009	2	2	C	T	C		4,14
m	0.0009	2	2	C	T	C		1,10
l	0.0009	2	2	C	T	C		1,10
u	0.00045	1	1	C	T	C		4
k	0.00045	1	1	C	T	C		13
v	0.00045	1	1	C	T	C		1
s	0.00045	1	1	C	T	G		5
q	0.00045	1	1	C	T	C		5
r	0.00045	1	1	C	T	G		2
n	0.00045	1	1	C	T	C		7
p	0.00045	1	1	C	T	C		10
o	0.00045	1	1	C	T	C		5

Table 13.62 - IGHV4-34

h.type	cum.freq	# indivs	pops	14:106829649	14:106829717	14:106829767	14:106829775	14:106829793	14:106829817	14:106829874
				rs183223225	rs11546808	rs188280613	rs191592421	rs11546811	rs1064183	rs181039074
ref	0.9955	-	-	Ser->Phe	Ser->Arg	Pro->Ser	Ile->Ser	Gly->Ala	Ala->Asp	Leu->Pro
d	0.0009	2	2	G	G	G	A	C	G	A
g	0.0009	2	2	G	G	G	A	G	G	A
b	0.0009	2	2	G	T	G	A	C	G	A
a	0.00045	1	1	G	G	G	A	C	G	G
e	0.00045	1	1	G	G	G	C	C	G	A
c	0.00045	1	1	G	G	G	A	C	G	A
f	0.00045	1	1	G	G	G	A	C	T	A

ENST000000390616

2,14
11,14
3,10
6
5
7
2

Table 13.63 - IGHV4-39

h.type	cum.freq	# indivs	pops	14:106877626 rs188639181	14:106877648 rs191739990	14:106877705 rs187288552	14:106877730 rs191860623	14:106877753 rs35281264	14:106877755 rs186830319	14:106877761 rs115523671	14:106877831 rs190205925
ref	0.6445	-	-	Cys->Tyr	Ala->Thr	Ile->Leu	Asn->Lys	Tyr->His	Tyr->Ser	Ser->Asn	Ile->Phe
a	0.10295	220	14	C	C	T	G	A	T	C	T
b	0.0918	197	14	C	C	T	G	G	T	C	T
d	0.04425	93	13	C	C	T	G	A	T	C	T
c	0.02555	56	10	C	C	T	G	G	T	C	T
e	0.01595	35	12	C	C	T	G	A	T	T	T
f	0.0132	29	10	C	C	T	G	G	T	C	T
g	0.0105	23	9	C	C	T	G	G	T	C	T
h	0.00595	13	8	C	C	T	G	G	T	C	T
i	0.0059	13	7	C	C	T	G	A	T	C	T
j	0.0055	12	8	C	C	T	G	A	T	C	T
m	0.00415	9	5	C	C	T	G	G	T	C	T
k	0.00365	8	5	C	C	T	G	G	T	C	T
l	0.00275	6	6	C	C	T	G	G	T	T	T
y	0.0018	4	3	C	C	T	G	A	T	C	T
n	0.0018	4	3	C	C	T	G	G	T	C	T
o	0.0018	4	3	C	C	T	G	G	T	C	T
u	0.00135	3	3	C	C	T	G	A	T	C	T
t	0.00135	3	2	C	C	T	G	G	T	C	T
q	0.00135	3	3	C	C	T	G	A	T	C	T
aa	0.0009	2	2	C	C	T	G	G	T	C	T
s	0.0009	2	2	C	C	T	G	G	T	C	T
x	0.0009	2	2	C	C	T	G	G	T	C	T
p	0.0009	2	2	C	C	T	G	A	T	T	T
ah	0.00045	1	1	C	C	T	G	A	T	C	T
ag	0.00045	1	1	C	C	T	G	A	T	T	T
aj	0.00045	1	1	C	C	T	G	G	T	C	T
aq	0.00045	1	1	C	T	T	G	A	T	C	T
v	0.00045	1	1	C	C	T	G	G	T	C	T
al	0.00045	1	1	C	C	T	C	A	T	C	T

Table 13.63 - IGHV4-39

ab	0.00045	1	C		T	C		C		A	G	C		T
ak	0.00045	1	C		T	C		G		G	T	T		T
ap	0.00045	1	C		T	C		G		A	T	T		T
z	0.00045	1	C		T	C		G		A	T	T		T
w	0.00045	1	C		T	C		G		G	T	C		T
ao	0.00045	1	C		T	C		G		A	T	C		T
r	0.00045	1	C		T	C		G		A	T	C		T
af	0.00045	1	T		T	C		G		A	T	C		T
ad	0.00045	1	C		T	C		G		A	T	C		T
am	0.00045	1	C		T	C		G		A	T	C		T
at	0.00045	1	C		T	C		G		A	G	C		■
ac	0.00045	1	C		T	C		G		A	T	C		T
ar	0.00045	1	C	T	T			G		A	T	C		T
as	0.00045	1	C		G	C		G		A	T	T		T
ae	0.00045	1	C		T	C		G		A	T	C		T
ai	0.00045	1	C		T	C		G		A	T	C		T
an	0.00045	1	C		T	C		G		A	T	C		T

Table 13.63 - IGHV4-39

h.type	cum.freq	# indivs	pops	14:106877836	14:106877837	14:106877848	14:106877877	14:106877888	14:106877907	14:106877912	14:106878019
				rs143740707	rs145759866	rs181859654	rs181228803	rs190955722	rs144078801	rs74093494	rs140461651
ref				Gly->Asp	Gly->Cys	Thr->Ser	Lys->Asn	Gly->Arg	Gln->His	Leu->Val	Ala->Thr
	0.6445	-	-	C	C	G	C	C	C	G	C
a	0.10295	220	14	C	C	G	C	C	C	C	C
b	0.0918	197	14	T		G	C	C	C	C	C
d	0.04425	93	13	C	C	G	C	C	C	G	C
c	0.02555	56	10	C	C	G	C	C	C	C	C
e	0.01595	35	12	C	C	G	C	C	C	G	C
f	0.0132	29	10	C	C	G	C	C	C	G	C
g	0.0105	23	9	C	C	G	C	C	C	G	C
h	0.00595	13	8	C	C	G	C	C	C	C	C
i	0.0059	13	7	C	C	G	C	C	C	C	C
j	0.0055	12	8	C	C	G	C	C	C	C	C
m	0.00415	9	5	C	C	G	C	C	C	C	C
k	0.00365	8	5	T		G	C	C	C	G	C
l	0.00275	6	6	C	C	G	C	C	C	C	C
y	0.0018	4	3	T	C	G	C	C	C	G	C
n	0.0018	4	3	C	C	G	C	C	C	G	C
o	0.0018	4	3	T		G	C	C	C	C	C
u	0.00135	3	3	C	C	G	C	C	C	G	C
t	0.00135	3	2	T		G	C	C	C	G	C
q	0.00135	3	3	C	C	G	C	C	C	G	C
aa	0.0009	2	2	C	C	G	C	C	C	G	C
s	0.0009	2	2	C		G	C	C	C	C	C
x	0.0009	2	2	T	C	G	C	C	C	C	C
p	0.0009	2	2	C	C	G	C	C	C	C	C
ah	0.00045	1	1	C	C	G	C	C	C	G	C
ag	0.00045	1	1	C	C	G	C	C	C	C	C
aj	0.00045	1	1	T		G	C	C	C	C	C
aq	0.00045	1	1	C	C	G	C	C	C	G	C
v	0.00045	1	1	T	C	G	C	C	C	G	C
al	0.00045	1	1	C	C	G	C	T	C	G	C

Table 13.63 - IGHV4-39

ab	0.00045	1	C		C	G	C	C	C	C	G	C	C
ak	0.00045	1	C		C	G	C	C	C	C	G	C	C
ap	0.00045	1	C		C	G	G	C	C	C	G	C	C
z	0.00045	1	C		C	G	C	C	C	C	G	C	C
w	0.00045	1	T		C	G	C	C	C	C	C	C	C
ao	0.00045	1	T		C	G	C	C	C	C	C	C	C
r	0.00045	1	C		C	G	C	C	C	■	G	C	C
af	0.00045	1	C		C	G	C	C	C	C	G	C	C
ad	0.00045	1	T		■	G	C	C	C	C	C	C	C
am	0.00045	1	C		C	G	C	C	C	C	G	C	C
at	0.00045	1	C		C	C	C	C	C	C	G	C	C
ac	0.00045	1	C		C	C	C	C	C	C	G	C	C
ar	0.00045	1	C		C	G	C	C	C	C	G	C	C
as	0.00045	1	T		C	G	C	T	C	C	C	T	C
ae	0.00045	1	T		■	G	C	C	C	C	G	C	C
ai	0.00045	1	C		C	G	C	C	C	C	G	C	C
an	0.00045	1	T		■	G	C	C	C	C	C	C	C

Table 13.63 - IGHV4-39

h.type	cum.freq	# indivs	pops	14:106878048	14:106878052	14:106878056	14:106878057	14:106878067	
				rs187579450	rs138364008	rs4774113	rs190089302	rs4774114	
ref				His->Arg	Lys->Gln	Lys->Asn	Lys->Thr	Met->Leu	ENST00000390619
	0.6445	-	-	T	T	T	T	T	
a	0.10295	220	14	T	T	G	T	G	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.0918	197	14	T	T	G	T	G	1,2,3,4,5,6,7,8,9,10,11,12,13,14
d	0.04425	93	13	T	T	G	T	G	1,2,3,4,5,6,7,9,10,11,12,13,14
c	0.02555	56	10	T	T	T	T	T	1,2,3,5,6,7,10,11,13,14
e	0.01595	35	12	T	T	T	T	T	1,2,3,4,5,6,9,10,11,12,13,14
f	0.0132	29	10	T	T	T	T	T	1,2,3,4,5,6,7,10,11,14
g	0.0105	23	9	T	T	G	T	G	1,2,4,5,10,11,12,13,14
h	0.00595	13	8	T	T	T	T	G	1,2,6,7,8,10,13,14
i	0.0059	13	7	T	T	G	T	G	1,2,4,8,10,13,14
j	0.0055	12	8	T	T	T	T	T	1,2,3,4,5,9,11,12
m	0.00415	9	5	T	T	G	T	T	2,3,7,10,14
k	0.00365	8	5	T	T	G	T	G	2,3,4,9,13
l	0.00275	6	6	T	T	G	T	G	1,2,5,7,12,13
y	0.0018	4	3	T	T	T	T	T	2,5,13
n	0.0018	4	3	T	T	G	T	T	3,13,14
o	0.0018	4	3	T	T	T	T	T	4,6,13
u	0.00135	3	3	T	T	T	T	G	1,4,14
t	0.00135	3	2	T	T	T	T	T	3,4
q	0.00135	3	3	T	T	G	T	T	2,3,6
aa	0.0009	2	2	T	T	T	T	G	1,6
s	0.0009	2	2	T	T	G	T	G	7,8
x	0.0009	2	2	T	T	G	T	G	3,4
p	0.0009	2	2	T	T	G	T	G	5,14
ah	0.00045	1	1	C	T	T	T	T	13
ag	0.00045	1	1	T	T	T	T	T	11
aj	0.00045	1	1	T	T	T	T	G	3
aq	0.00045	1	1	T	T	G	T	G	2
v	0.00045	1	1	T	T	T	T	T	7
al	0.00045	1	1	T	T	G	T	G	14

Table 13.63 - IGHV4-39

ab	0.00045	1	T	T	T	T	T	T	4
ak	0.00045	1	T	T	T	T	T	T	13
ap	0.00045	1	T	T	T	T	T	T	13
z	0.00045	1	T	T	T	G	T	T	10
w	0.00045	1	T	T	T	G	T	T	10
ao	0.00045	1	T	T	T	T	T	T	12
r	0.00045	1	T	T	T	T	T	T	3
af	0.00045	1	T	T	T	T	T	T	12
ad	0.00045	1	T	T	T	T	T	T	5
am	0.00045	1	T	T	T	T	G	T	13
at	0.00045	1	T	T	T	T	T	T	1
ac	0.00045	1	T	T	T	T	T	T	13
ar	0.00045	1	T	T	T	T	T	T	6
as	0.00045	1	T	T	T	G	T	G	2
ae	0.00045	1	T	T	T	G	T	G	9
ai	0.00045	1	T	T	G	T	T	T	13
an	0.00045	1	T	T	T	G	T	G	2

Table 13.64 - IGHV4-4

h.type	cum.frec	# indvs	pops	14:106478120	14:106478183	14:106478194	14:106478252	14:106478311	14:106478354
				rs150123115	rs77628366	rs76899215	rs73365408	.	rs188216744
ref				Tyr->Cys	Thr->Lys	Met->Ile	Arg->His	Ser->	Glu->Gly
a	0.34 -	222	-	T	G	C	C	A	T
d	0.1136	166	14	T	G	C	C		C
f	0.0817	138	14	T	G	T	C		T
b	0.06675	140	13	T	G	T	C		C
c	0.06385	127	14	T	G	C	C	A	T
e	0.0599	110	12	T	T	T	C		T
g	0.05215	65	10	C	T	T	C		C
m	0.03255	58	12	T	T	T	T		C
i	0.0267	58	14	T	T	T	C		T
j	0.0262	38	11	T	T	T	C	A	T
h	0.01715	33	11	T	G	C	C	A	C
k	0.01535	25	10	C	G	C	C		C
o	0.0113	19	9	T	T	C	C		C
q	0.00855	17	7	C	G	C	C	A	T
n	0.00765	16	5	T	G	T	T	A	T
p	0.00765	16	5	T	G	T	T		T
l	0.00685	15	4	C	G	C	C	A	C
s	0.00585	13	4	T	G	C	T	A	T
aj	0.0054	12	8	T	G	T	C	A	C
ah	0.00495	11	7	T	T	T	T		T
t	0.0045	10	6	C	G	T	C		C
w	0.00405	9	7	C	T	T	C		T
r	0.0036	8	7	T	G	T	T		C
v	0.00315	7	5	C	G	C	C		T
ag	0.0027	6	5	T	T	T	C	A	C
y	0.0027	6	4	C	T	C	C		C
u	0.0027	6	3	T	T	T	T	A	T
z	0.0027	6	4	C	T	T	T		C

ENST00000390596

1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,9,11,12,13,14
1,2,3,4,5,9,10,11,12,14
1,2,3,4,5,6,7,9,10,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,4,5,6,7,9,10,11,12,14
2,3,4,5,6,7,9,10,11,12,13
2,3,4,5,6,8,9,11,12,13
1,2,3,4,5,6,7,8,13
1,3,9,10,11,12,14
1,5,10,13,14
1,5,9,10,14
3,4,9,10
1,10,12,14
2,3,4,5,9,12,13,14
1,2,3,9,10,13,14
3,4,6,8,9,11
3,4,7,9,11,13,14
2,4,9,11,12,13,14
7,9,11,13,14
2,5,6,10,14
3,4,5,9
1,10,14
2,4,9,10

Table 13.64 - IGHV4-4

ak	0.0018	4	T	G	C	T	T		T	3,4,5,10
ao	0.00135	3	C	G	T	C	C		T	3,9,11
ac	0.00135	3	T	G	C	T	T		C	10
ae	0.00135	3	T	T	C	C	C	A	C	2,4
aa	0.0009	2	C	T	T	C	C	A	T	9,14
af	0.0009	2	T	T	C	T	T		C	6,9
x	0.0009	2	C	G	T	C	C	A	T	9,12
ad	0.0009	2	T	T	C	C	C		T	3,7
ai	0.0009	2	T	T	C	C	C	A	T	7
al	0.00045	1	C	G	C	T	T		C	4
ab	0.00045	1	C	T	C	T	T		C	10
am	0.00045	1	C	G	T	T	T	A	C	9
an	0.00045	1	C	G	C	T	T	A	C	1

	14:107082722	14:107083453	14:107083457
rs181388142		rs112607697	rs190037133
		Tyr->His	Ser->Arg
Ser->Leu		Tyr->His	Ser->Arg
G	A	A	A
G	G	G	A
G	G	A	C
		A	A
		A	A

Table 13.66 - IGHV4-61

h.type	cum.freq	# indvs	pops	14:107095319	14:107095325	14:107095326	14:107095338
				rs111864170	rs186969565	rs1064309	rs2072046
ref				Tyr->Ser	Gly->Asp	Gly->Ser	Val->Ile
a	0.0109	-	-	T	C	C	C
b	0.5701	806	14	T	C	T	C
c	0.18115	339	14	T	C	T	T
d	0.13535	275	13	G	C	C	T
e	0.05855	120	14	T	C	C	T
f	0.0191	42	12	G	C	C	C
g	0.0109	24	10	G	C	T	C
h	0.00675	15	4	T	T	C	T
i	0.00585	13	7	G	C	T	T
j	0.00045	1	1	T	T	C	C
k	0.00045	1	1	T	T	T	T
l	0.00045	1	1	G	T	C	T

ENST000000390630

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,6,7,9,10,11,12,13

1,9,10,14

2,3,6,7,9,10,14

7

10

14

Table 13.67 - IGHV5-51

h.type	cum.freq	# indivs	pops	14:107034771	14:107034793	14:107034796	14:107034813	14:107034850	14:107034851	14:107034868	14:107034910
				rs180672762	rs181977500	rs186695491	rs191700780	rs142572296	rs186314656	rs139665108	rs117410356
ref	0.97045	-	-	Ser->Arg	Ser->Asn	Ile->Ser	Ile->Met	Thr->Ser	Thr->Ala	Tyr->Phe	Arg->His
a	0.0164	35	3	G	C	A	G	G	T	T	C
b	0.0055	11	3	G	C	A	G	G	T	T	T
g	0.00135	3	3	G	T	A	G	G	T	T	C
d	0.0009	2	2	G	C	A	G	C	T	T	C
k	0.0009	2	2	G	C	A	G	G	T	T	C
c	0.0009	2	2	G	T	A	G	G	T	T	C
j	0.00045	1	1	G	C	A	G	G	T	■	C
e	0.00045	1	1	G	C	A	C	G	T	T	C
h	0.00045	1	1	G	C	A	G	G	T	T	C
f	0.00045	1	1	G	C	A	G	G	T	T	C
i	0.00045	1	1	G	C	A	G	G	T	T	C
n	0.00045	1	1	G	C	C	G	G	T	T	C
m	0.00045	1	1	T	C	A	C	G	C	T	C
l	0.00045	1	1	G	C	A	G	G	T	T	C

Table 13.67 - IGHV5-51

h.type	cum.freq	# indivs	pops	14:107034931	14:107034940	14:107034998	14:107035123	14:107035141
				rs113154616	rs113988349	rs183518874	rs116300992	rs147695427
ref	0.97045	-	-	Ser->Thr	Ser->Asn	Ala->Pro	Leu->Ile	Ala->Thr
a	0.0164	35	3	C	C	C	G	C
b	0.0055	11	3	C	C	C	T	C
g	0.00135	3	3	C	C	C	G	C
d	0.0009	2	2	C	C	C	G	C
k	0.0009	2	2	G	C	C	G	C
c	0.0009	2	2	C	T	C	G	C
j	0.00045	1	1	C	C	C	G	C
e	0.00045	1	1	C	C	C	G	C
h	0.00045	1	1	C	C	C	G	T
f	0.00045	1	1	G	T	C	G	C
i	0.00045	1	1	C	T	C	G	C
n	0.00045	1	1	C	C	C	G	C
m	0.00045	1	1	G	C	C	G	C
l	0.00045	1	1	C	C	G	G	C

ENST000000390626

3,4,9

5,10,14

6,7,13

2,13

6,10

2,10

13

10

7

10

10

4

4

6

Table 13.68 - IGHV6-1

h.type	cum.freq	# indvs	pops	14:106405816 rs186349163	14:106405877 rs72715486	14:106405910 rs188294175	ENST000000390593
ref	0.99775	-	-	Ser->Asn	Lys->Glu	Val->Ile	
a	0.0009	2	2	C	T	C	
c	0.0009	2	1	C	C	C	
b	0.00045	1	1	T	T	T	
							3,8 4 8

h.type	cum.freq	# indivs	pops	14:107282845	14:107282853	14:107282887	14:107283009	14:107283018	14:107283073
ref	0.9127	-	-	G	G	G	C	T	C
a	0.0732	136	6	T		G	C	T	C
b	0.01005	21	3	G	G	G		T	C
c	0.0018	4	3	G	G		C	T	C
e	0.00135	3	1	G	G	G	C		C
d	0.0009	2	2	G	G	G	C	T	
				Leu->Met	Thr->Ile	Arg->Trp	Gly->Val	Lys->Met	Val->Leu
				rs61741324	rs61741323	rs61741308	rs11845591	rs144292230	rs148758470
				14:107282845	14:107282853	14:107282887	14:107283009	14:107283018	14:107283073
				ENST00000390639					

Table 13.70 - IGKJ2

h.type	cum.freq	# indivs	pops	2:89161040	2:89161044	2:89161045	2:89161050	2:89161055	2:89161067	2:89161069	2:89161072
ref	0.921	-	-	Lys->Glu	Glu->Asp	Glu->Gly	Lys->Asn	Thr->Pro	Phe->Val	Thr->Ser	Tyr->Cys
a	0.0449	86	13	T	C	T	C	T	A	G	T
b	0.0243	48	10	T	C	T	C	T	A	C	T
d	0.0032	6	3	T	C	T	C	T	A	G	C
e	0.0018	4	3	T	C	C	C	T	A	G	T
c	0.0018	4	4	T	C	T	G	T	A	G	T
f	0.0014	3	2	T	C	T	C	T	C	G	T
g	0.0009	2	2	T	■	T	C	T	A	G	T
h	0.0005	1	1	C	C	T	C	T	C	G	C
i	0.0005	1	1	T	C	T	C	G	A	G	T

ENST000000390241

1,2,3,4,5,6,7,9,10,11,12,13,14

2,4,5,6,7,8,10,11,12,13

2,6,7

2,10,13

1,2,7,9

2,10

11,14

9

10

Table 13.71 - IGKJ3

h.type	cum.frec	# indivs	pops	2:89160742	2:89160748	2:89160762	2:89160769
				rs188533155	rs181878952	rs190475170	rs139877244
ref				Asp->His	Lys->Glu	Phe->Cys	Phe->Leu
a	0.99045	-	-	C	T	A	A
d	0.00685	15	10	C	T	A	G
b	0.00135	3	3	C	C	A	A
c	0.00009	2	2	G	T	A	A
	0.00045	1	1	C	T	C	A
				ENST000000390240			
				2,4,5,6,9,10,11,12,13,14			
				5,11,12			
				2,3			
				5			

Table 13.72 - IGKJ4

h.type	cum.fre	# indivs	pops	2:89160399	2:89160404	2:89160405	2:89160406	2:89160407	2:89160409	2:89160412	2:89160415	2:89160427
				rs78964890	rs139461207	rs180910647	rs185131270	rs145002796	rs191518742	rs147540982	rs139402168	rs186462161
ref	0.9883	-	-	Lys->Asn	Ile->Leu	Glu->Asp	Glu->Gly	Glu->Gln	Val->Ala	Lys->Arg	Thr->Ile	Phe->Ser
b	0.00225	5	4	T	T	C	T	C	A	T	G	A
a	0.0018	4	4	T	T	C	T	C	A	C	G	A
g	0.00135	3	3	■	T	C	T	C	A	T	G	A
j	0.0009	2	1	T	T	C	C	C	A	T	G	A
k	0.0009	2	2	T	G	C	T	C	A	T	G	A
c	0.0009	2	2	T	T	C	T	C	A	T	G	A
h	0.0009	2	2	T	T	C	T	G	A	T	G	A
d	0.00045	1	1	T	T	C	T	C	A	T	■	A
e	0.00045	1	1	T	T	C	T	C	A	T	G	G
f	0.00045	1	1	T	T	C	T	C	G	T	G	A
i	0.00045	1	1	T	T	G	T	C	A	T	G	A
m	0.00045	1	1	T	T	C	T	C	A	C	G	A
l	0.00045	1	1	■	T	C	T	C	A	T	G	A

Table 13.72 - IGKJ4

h.type	cum.fre	# indivs	pops	2:89160428	2:89160430	ENST00000390239
				rs191712435	rs187905157	
ref	0.9883	-	-	Phe->Leu	Thr->Ile	
b	0.00225	5	4	A	G	2,5,13,14
a	0.0018	4	4	A		2,5,6,10
g	0.00135	3	3	A	G	3,4,10
j	0.0009	2	1	A	G	10
k	0.0009	2	2	A	G	9,10
c	0.0009	2	2	G	G	2,14
h	0.0009	2	2	A	G	2,10
d	0.00045	1	1	A	G	5
e	0.00045	1	1	A	G	6
f	0.00045	1	1	A	G	14
i	0.00045	1	1	A	G	9
m	0.00045	1	1	G	G	9
l	0.00045	1	1	G	G	13

Table 13.73 - IGKJ5

h.type	cum.fre	# indivs	pops	2:89160082	2:89160087	2:89160106	2:89160109	2:89160110	2:89160113
				rs185692630	rs142632901	rs190985433	rs140799895	rs114105515	rs188703994
ref				Lys->Arg	Glu->Asp	Gly->Ala	Phe->Cys	Phe->Leu	Thr->Ser
a	0.9928	-	-	T	C	C	A	A	T
c	0.00225	5	4	C	C	C	A	A	T
b	0.00135	3	3	T	C	C	A	G	T
d	0.00135	3	3	T	G	C	A	A	T
g	0.00045	1	1	T	G	C	A	G	T
e	0.00045	1	1	T	C	C	A	A	
h	0.00045	1	1	T	C	G	A	A	T
f	0.00045	1	1	C	G	C	A	A	T

ENST000000390238

4,5,6,14

9,10,11

3,5,9

10

10

10

14

3

Table 13.74 - IGKV1-16

h.type	cum.freq	# indivs	pops	2:89399455	2:89399488	2:89399509	2:89399515	2:89399809	
ref	0.97625	-	-	rs193208674	rs184390186	rs116145759	rs2848410	rs188494311	
a	0.01145	25	8	Lys->Arg	Ala->Val	Ala->Val	Gly->Glu	Ala->Thr	ENST000000479981
b	0.00595	13	10	T	G	G	C	C	1,4,6,7,8,9,11,13
c	0.00455	10	7	C	G	G	C	C	2,4,5,6,7,8,9,10,13,14
d	0.0009	2	2	T	G	G	T	C	2,5,6,9,11,13,14
e	0.00045	1	1	T	G	■	C	C	1,14
f	0.00045	1	1	T	G	G	C	T	9
					■	G	C	C	7

Table 13.75 - IGKV1-17

h.type	cum.freq	# indivs	pops	2:89416951	2:89417000	2:89417011	2:89417047	2:89417087
				rs185657549	rs188985676	rs191430092	rs188555641	rs181054320
				Ser->Thr	Pro->Ser	Tyr->Phe	Arg->Gln	Leu->Met
ref	0.83295	-	-	C	G	T	C	G
a	0.08655	170	13	C	G	T	C	T
b	0.0545	111	12	C	G	■	C	T
c	0.0224	47	11	C	G	■	C	G
d	0.00225	4	1	C	■	T	C	G
e	0.0009	2	2	G	G	T	C	G
f	0.00045	1	1	C	G	T	T	G
				ENST000000490686				
				1,2,3,4,5,6,7,9,10,11,12,13,14				
				1,2,3,4,5,7,9,10,11,12,13,14				
				1,2,4,5,6,7,9,10,12,13,14				
				10				
				2,10				
				7				

Table 13.76 - IGKV1-5

h.type	cum.freq	# indivs	pops	2:89246874	2:89246904	2:89246946	2:89247012	2:89247045	2:89247051	2:89247106
				rs11546105	rs182050897	rs11546098	rs146302149	rs191761898	rs182340840	rs189714283
ref	0.9823	-	-	Ser->Asn	Ser->Tyr	Ser->Asn	Ser->Thr	Thr->Ser	Arg->Thr	Cys->Gly
a	0.0073	16	8	C	G	C	C	G	C	A
b	0.005	11	8	C	G	C	G	G	C	A
f	0.0018	4	4	T	G	C	C	G	C	A
c	0.00135	3	3	C	G	T	G	G	C	A
d	0.0009	2	1	C	G	C	C	G	C	C
g	0.00045	1	1	C	G	C	C	G	G	A
e	0.00045	1	1	C	G	C	C	C	C	A
h	0.00045	1	1	C	T	C	C	G	C	A

ENST000000496168

1,6,8,9,11,12,13,14

1,2,3,6,7,11,12,14

7,10,13,14

10,12,13

10

11

11

14

Table 13.77 - IGKV1-6

h.type	cum.fre	# indvs	pops	2:89265959	
				rs188876010	
				Tyr->Cys	
ref	0.99955	-	-	T	
a	0.00045	1	1	C	
				10	
				ENST000000464162	

Table 13.78 - IGKV1-8

h.type	cum.frec	# indivs	pops	2:89291968 rs187771921	2:89292080 rs190853230	2:89292130 rs183046670	2:89292149 rs186879210	2:89292182 rs59175543	2:89292188 rs192180011	2:89292201 rs35262290
ref	0.60715	-	-	Asp->Ala	Lys->Glu	Gly->Asp	Thr->Ser	Phe->Leu	Ser->Pro	Met->Ile
a	0.20315	348	14	T	T	C	T	A	A	C
b	0.1334	235	14	T	T	C	T	G	A	T
c	0.03325	70	14	T	C	C	T	A	A	C
e	0.01035	23	8	T	C	C	T	G	A	C
d	0.00955	21	8	T	C	C	T	G	A	T
h	0.0009	2	2	T	T	T	T	G	A	T
j	0.00045	1	1	G	T	C	T	A	A	C
k	0.00045	1	1	T	C	T	T	G	A	C
g	0.00045	1	1	T	T	C		G	A	C
f	0.00045	1	1	T	T	C	T	G	G	T
i	0.00045	1	1	T	T	T	T	A	A	C

ENST000000495489

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,6,7,10,11,12,13,14

1,3,4,5,7,9,10,14

9,10

7

1

14

5

11

Table 13.79 - IGKV1-9

h.type	cum.freq	# indvs	pops	2:89309554	2:89309694	2:89309735	2:89309762
				rs147109427	rs185997823	rs80322626	rs77800356
ref				Glu->Asp	Arg->Trp	Phe->Ser	Asp->Ala
a	0.7266	-	-	T	G	A	T
b	0.1854	317	14		G	G	G
c	0.02735	57	10	T	G	A	G
d	0.0197	43	13		G	A	G
e	0.01825	40	13		G	A	T
f	0.01095	24	11	T		A	T
g	0.00545	12	7	T	G	G	G
h	0.0018	4	4	T	G	G	T
i	0.0018	4	4	T		A	G
j	0.00135	3	3		G	G	T
k	0.0009	2	2			G	G
	0.00045	1	1			A	G

ENST00000493819

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,3,4,5,7,9,10,11,13,14

1,2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,4,5,7,8,9,10,11,12,13,14

2,4,5,6,7,8,9,10,11,12,13

1,3,4,5,7,10,11

1,10,11,14

3,4,10,11

5,12,13

4,14

4

Table 13.80 - IGKV1D-12

h.type	cum.freq	# indvs	pops	2:90198951 rs189385596	2:90199107 rs180730188	
ref	0.9955	-	-	Gly->Arg	Gly->Arg	ENST00000390276
b	0.0027	6	4	G	G	1,7,10,12
a	0.0018	4	4	G	G	2,8,13,14

Table 13.81 - IGKV1D-16

h.type	cum.freq	# indvs	pops	2:90139373 rs6760987	2:90139377 rs184550323	2:90139422 rs189369337	
ref	0.96165	-	-	Ser->Arg T	Gly->Ser G	Ala->Thr G	ENST00000492446
a	0.03655	71	13	G	G	G	1,2,3,4,5,6,7,8,10,11,12,13,14
c	0.0009	2	1	T	G		1
b	0.0009	2	1	T		G	10

Table 13.82 - IGKV1D-17

h.type	cum.fre	# indivs	pops	2:90121919				2:90121961		2:90121999		2:90122063	
				Arg->Gln	Gln->Arg	Ala->Ser	Thr->Ile	rs182773099	rs191806633	rs182889841	rs193008187	ENST000000483379	
ref	0.99315	-	-	G	A	G	C						
a	0.0037	8	4	G	A	T	C					1,5,10,14	
c	0.00135	3	2	G	G	G	C					7,12	
b	0.00135	3	2		A	G	C					1,10	
d	0.00045	1	1	G	A	G	T					6	

Table 13.83 - IGKV1D-42

h.type	cum.freq	# indvs	pops	2:90229083	2:90229247	2:90229260	2:90229265	2:90229292	2:90229314	2:90229471	2:90229477	2:90229491
				rs193096486	rs139321345	rs842173	rs145538558	rs147724164	rs192596253	rs184070665	rs188365115	rs181576535
ref	0.7584	-	-	Ala->Thr	Asp->Tyr	Ile->Thr	Ser->Pro	Gly->Arg	Cys->Phe	Ile->Met	Ser->Arg	Asp->Val
a	0.22345	383	14	G	G	T	T	G	G	C	C	A
b	0.01005	21	3	G	G	C	T	G	G	C	C	A
c	0.0027	6	1	G	G	T	C	G	G	C	C	A
d	0.00135	3	2	■	G	T	T	■	G	C	C	A
e	0.00135	3	1	G	T	T	T	G	G	C	C	A
g	0.0009	2	2	G	T	T	T	G	G	G	C	A
f	0.0009	2	1	G	G	T	T	G	T	C	C	A
h	0.00045	1	1	G	G	C	T	G	G	C	■	A
i	0.00045	1	1	G	G	T	T	G	G	C	C	T

Table 13.83 - IGKV1D-42

h.type	cum.freq	# indvs	pops	
ref	0.7584	-	-	ENST00000390278
a	0.22345	383	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.01005	21	3	1,10,14
c	0.0027	6	1	9
d	0.00135	3	2	10,14
e	0.00135	3	1	13
g	0.0009	2	2	12,14
f	0.0009	2	1	14
h	0.00045	1	1	13
i	0.00045	1	1	3

	h.type	cum.freq	# indivs	pops	2:90248957	2:90249120	2:90249147	2:90249151	2:90249237	2:90249247	2:90249348	2:90249372
ref	0.98095	-	-	-	rs116586875	rs188075270	rs184607123	rs138753337	rs142796670	rs192607496	rs184789392	rs188397844
b	0.00685	14	3		Leu->Gln	Met->Val	Ala->Thr	Ser->Phe	Ala->Thr	Leu->Arg	Pro->Ser	Cys->Arg
a	0.00635	12	4									ENST000000468879
c	0.0027	5	3		T	A						
d	0.0009	2	2		T	A	Ala->Thr	Ser->Phe	Ala->Thr	Leu->Arg	Pro->Ser	Cys->Arg
e	0.0009	2	1		T	A	G	C	G	T	T	
g	0.00045	1	1		T	G	G	C	G	T	T	
h	0.00045	1	1		T	A	G	C	G	G	C	
f	0.00045	1	1		T	A		C	G	T	C	

Table 13.85 - IGKV1D-8

h.type	cum.frec	# indivs	pops	2:90259956	2:90259965	2:90260022	2:90260033	2:90260085	2:90260091	2:90260178
ref	0.7813	-	-	Ala->Val	Val->Ala	Thr->Ile	Arg->Trp	Gly->Glu	Ala->Val	Thr->Ile
a	0.1872	319	14	C	T	C	C	G	C	C
b	0.01785	36	3	C	C	C	C	G	C	T
c	0.0105	22	6	C	T	T	C	G	C	C
d	0.0009	2	1	C	C	C	C	G	T	C
f	0.0009	2	2	T	T	C	C	G	C	C
g	0.00045	1	1	C	T	C	C		C	C
e	0.00045	1	1	C	T	C	T	G	C	C
h	0.00045	1	1	C	T	C	C	G	T	T

ENST000000471857

1,2,3,4,5,6,7,8,9,10,11,12,13,14

3,4,9

2,5,6,7,12,13

1

2,12

1

1

9

Table 13.86 - IGKV2-24

h.type	cum.freq	# indivs	pops	2:89475831 rs149900655	2:89476044 rs183417517	
				Met->Thr	Cys->Phe	ENST000000484817
ref	0.8082	-	-	A	C	
a	0.1708	293	13	G	C	1,2,3,4,5,6,7,9,10,11,12,13,14
b	0.02055	44	13	G		1,2,3,4,5,6,7,8,9,10,12,13,14
c	0.00045	1	1	A		8

Table 13.87 - IGKV2-30

h.type	cum.fre	# indvs	pops	2:89544389					2:89544473		2:89545042	
				Arg->Trp	Asn->Asp	Tyr->His	Leu->Arg	rs141762272	rs182126464	rs147079411	rs186725977	
ref	0.75005	-	-	G	T	A	A					ENST00000468494
a	0.24085	437	13	G	T	G	A					1,2,3,4,5,6,7,9,10,11,12,13,14
b	0.0082	18	7		T	A	A					2,4,6,7,9,12,13
d	0.00045	1	1	G	T	A	C					11
c	0.00045	1	1	G	C	A	A					11

Table 13.88 - IGKV2D-24

h.type	cum.freq	# indivs	pops	2:90044207 rs150942262	2:90044207 rs190167237	2:90044207 rs190167237	ENST000000462693
ref	0.9863	-	-	Phe->Cys	Ala->Val	C	
a	0.01235	27	12	G	C	C	1,2,3,4,6,7,8,9,11,12,13,14
b	0.00135	3	2	T	T	T	5,10

Table 13.89 - IGKV2D-26

h.type	cum.freq	# indvs	pops	2:90025217	2:90025275	2:90025397	2:90025483
ref	0.6664	-	-	Ile->Asn	Met->Ile	Gly->Glu	Tyr->Asn
a	0.325	556	14	T	G	G	T
b	0.00455	9	3	T	G	G	
c	0.0018	4	1		G	G	T
d	0.0009	2	2	T	C	G	
f	0.0009	2	2	T	G		T
e	0.00045	1	1		C	G	T

ENST000000390268

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,11,12

10

3,4

1,10

10

Table 13.90 - IGKV2D-29

h.type	cum.frec	# indivs	pops	2:89986777 rs140693346	2:89986852 rs191653248	2:89986921 rs424406	
ref	0.6969	-	-	Ala->Thr	Ser->Pro	Pro->Ser	ENST00000491977
a	0.30175	481	14	G	T	C	
b	0.0009	2	1	G	T	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.00045	1	1			C	13
				G	C	T	12

Table 13.91 - IGKV2D-30

h.type	cum.freq	# indivs	pops	2:89975711	2:89976341	2:89976353	2:89976364
				rs185844838	rs148937425	rs184425514	rs147056147
				Leu->His	Arg->His	Lys->Thr	Trp->Arg
				->	Arg->His	Lys->Thr	Trp->Arg
ref	0.9689	-	-	T	G	A	T
a	0.02335	51	14	T	G	A	C
b	0.00595	13	4		G	A	T
c	0.00135	3	2	T		A	T
d	0.00045	1	1	T	G	C	T
							1,2,3,4,5,6,7,8,9,10,11,12,13,14
							1,5,10,14
							1,14
							14

Table 13.92 - IGKV3-11

h.type	cum.fr	# indivs	pops	2:89326669	2:89326681	2:89326754	2:89326858	2:89326861	2:89326865	2:89326872
ref	0.94435	-	-	Pro->His	Arg->His	Ser->Pro	Tyr->Ser	Ser->Thr	Ser->Gly	Gln->His
a	0.02835	60	14	T	C	A	T	C	T	C
b	0.0096	21	11	G	C	G	T	C	T	C
d	0.0073	14	3	G	T	A	T	C	T	C
c	0.00545	12	7	T	C	A	T	C	T	C
e	0.0018	4	4	G	C	A	T	G	T	C
g	0.0009	2	1	G	T	G	T	C	T	C
f	0.0009	2	2	G	C	A	T	C	T	G
j	0.00045	1	1	G	C	A	G	C	C	C
h	0.00045	1	1	G	C	A	T	C	C	C
i	0.00045	1	1	G	C	A	G	C	T	C

ENST000000483158

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,3,4,5,6,7,8,9,11,13,14

1,10,14

1,6,7,9,11,13,14

2,5,7,9

14

9,13

14

13

9

Table 13.93 - IGKV3-20

h.type	cum.fr	# indivs	pops	2:89442183	2:89442191	2:89442193	2:89442196	2:89442253	2:89442255	2:89442262	2:89442342
ref	0.9847	-	-	Ser->Arg	Ala->Thr	Gly->Ala	Tyr->Phe	Ser->Asn	Ser->Arg	Ser->Thr	Glu->Asp
a	0.00225	5	3	G	C	C	T	C	G	C	T
b	0.00225	5	4	G	C	C	T	C	G	G	T
d	0.00135	3	3	G	C	C	T	C	G	C	T
f	0.00135	3	2	G	C	C	T	C	G	C	T
j	0.0009	2	2	G	C	C	T	C	C	C	T
e	0.0009	2	2	G	C	C	T	C	C	C	T
c	0.0009	2	2	G	C	G	T	C	G	C	T
i	0.0009	2	2	G	C	C	T	C	G	C	T
k	0.00045	1	1	C	T	C	T	C	G	G	T
g	0.00045	1	1	G	C	C	■	C	G	C	T
q	0.00045	1	1	G	T	C	T	C	G	C	T
r	0.00045	1	1	G	C	C	T	C	G	C	T
h	0.00045	1	1	G	C	C	■	C	G	C	T
n	0.00045	1	1	G	C	C	T	C	G	C	T
m	0.00045	1	1	G	C	C	T	C	G	C	T
l	0.00045	1	1	G	C	G	T	C	G	G	T
p	0.00045	1	1	G	C	C	T	T	C	C	T
o	0.00045	1	1	G	C	C	T	C	G	C	■

Table 13.93 - IGKV3-20

h.type	cum.fr	# indivs	pops	2:89442344	2:89442350	2:89442352	2:89442354	2:89442355
ref	0.9847	-	-	rs183373822	Thr->Ser	Thr->Ile	Asp->GluAsp->Glu	rs192831943
a	0.00225	5	3	C	T	G	A	Asp->AlaAsp->Ala
b	0.00225	5	4	C	T	G	A	T
d	0.00135	3	3	C	T		A	T
f	0.00135	3	2	C		G	A	T
j	0.0009	2	2	C	T		A	T
e	0.0009	2	2	C	T	G	A	T
c	0.0009	2	2	C	T	G	A	T
i	0.0009	2	2	C	T	G	T	T
k	0.00045	1	1	C	T	G	A	T
g	0.00045	1	1	G	T	G	A	T
q	0.00045	1	1	C	T	G	A	T
r	0.00045	1	1	C	T	G	A	T
h	0.00045	1	1	C	T	G	A	T
n	0.00045	1	1	C	T	G	A	G
m	0.00045	1	1	G	T	G	A	T
l	0.00045	1	1	C	T	G	A	T
p	0.00045	1	1	C	T	G	A	T
o	0.00045	1	1	C	T	G	A	T

ENST000000492167

1,9,14
5,7,10,14
2,6,14
1,14
1,10
9,13
10,11
9,14
11
14
13
9
14
10
13
10
2
4

Table 13.94 - IGKV3-7

h.type	cum.freq	# indivs	pops	2:89278103	2:89278106	2:89278219	2:89278262
				rs144594870	rs189211039	rs2458828	rs181802934
ref				Ser->Gly	Thr->Ser	Val->Ala	Thr->Ala
a	0.78325	-	-	T	T	A	T
b	0.2018	349	14	C	T	A	T
c	0.01	19	3	T	T	G	T
d	0.00225	5	3	T		A	T
e	0.00225	5	3	T	T	G	C
	0.00045	1	1	C	T	G	T
				ENST000000390247			
				1,2,3,4,5,6,7,8,9,10,11,12,13,14			
				1,10,14			
				1,10,14			
				1,10,14			
				9			

Table 13.95 - IGKV3D-11

h.type	cum.fr	# indivs	pops	2:90212050	2:90212051	2:90212126	2:90212138	2:90212167	2:90212198	2:90212240	2:90212243
ref	0.635	-	-	Gly->Ser	Gly->Val	Asn->Thr	Gly->Val	Pro->Ser	Ser->Asn	Arg->His	Ser->Ile
a	0.3511	561	14		G	A	G	C	G	G	G
c	0.0036	8	5		G	A	G	C	G	G	G
b	0.0023	5	3	G	G	A	G	T	G	G	G
g	0.0018	3	2	G	G	A	G	C	G		G
f	0.0018	4	3		G	A	G	C	G	G	T
d	0.0014	3	2	G	G	A	G	C	G		G
h	0.0009	2	1	G	G	C	G	C	G	G	G
i	0.0009	2	1	G	G	A	T	C	G	G	G
j	0.0005	1	1		G	A	G	T	G		G
k	0.0005	1	1	G	T	A	G	C	G	G	G
e	0.0005	1	1		G	A	G	T	G	G	G

ENST000000390277

1,2,3,4,5,6,7,8,9,10,11,12,13,14
4,6,7,8,9
3,4,9
1,10
1,3,4
10,14
4
10
14
9
1

Table 13.96 - IGKV3D-15

h.type	cum.freq	# indivs	pops	2:90154131	ENST00000417279
				rs146414056	
				Thr->Ile	
ref	0.7939	-	-	C	
a	0.2061	368	14	T	1,2,3,4,5,6,7,8,9,10,11,12,13,14

Table 13.97 - IGKV3D-20

h.type	cum.freq	# indivs	pops	2:90077781	2:90078037	2:90078079	2:90078093
				rs183765055	rs189772344	rs182769673	rs2555982
				Glu->Gly	Thr->Met	Ala->Gly	Gly->Arg
ref	0.9845	-	-	A	C	C	G
a	0.0073	16	9	A	C	C	
b	0.0064	12	4	G	C	C	G
d	0.0009	2	2	A	T	C	
e	0.00045	1	1	A	T	C	G
c	0.00045	1	1	A	C	G	G
				ENST000000390270			
				1,2,4,6,7,8,9,10,11			
				1,10,11,14			
				7,12			
				11			
				13			

Table 13.98 - IGKV4-1

h.type	cum.freq	# indvs	pops	2:89185362 rs180851255	2:89185392 rs116328981	2:89185400 rs148071875	2:89185431 rs141793384	2:89185450 rs189675840	2:89185451 rs138603647	2:89185455 rs181816666	2:89185457 rs141709704	2:89185461 rs78280375
ref	0.9516	-	-	Tyr->Phe	Asp->Gly	Ala->Thr	Asn->Ser	Ser->Arg	Val->Ile	Leu->Ser	Tyr->His	Ser->Asn
a	0.01185	26	11	A	A	G	A	T	G	T	T	G
b	0.00545	11	9	A	A	G	A	T		T	T	G
c	0.00365	8	7	A	A	G	A	T	G	T	T	G
d	0.00315	7	6	A	A		A	T	G	T	T	G
k	0.0018	4	4	A	A	G	A	T	G	T	T	G
w	0.0018	3	3	A	A	G	A	T	G	T	T	
h	0.0018	4	4	T	A	G	A	T	G	T	T	G
f	0.0018	4	4	A	A	G	A	T	G	T	C	G
e	0.00135	3	3	A	A	G	A	T	G	T	T	G
r	0.00135	3	3	A	A	G	A	T	G	T	T	G
x	0.00135	3	2	A	A	G	A	T	G	T	T	G
o	0.00135	3	2	A	A	G	A	T	G	T	T	G
y	0.0009	2	2	A	A	G	A	T	G	T	T	G
g	0.0009	2	2	A	A	G	A	T	G	T	T	G
t	0.0009	2	2	A	A	G	A	T	G	T	T	G
i	0.0009	2	2	A	A	G	A	T	G	T	T	G
n	0.0009	2	2	A	A	G	A	T	G	T	T	G
l	0.0009	2	2	A	A	G	G	T	G	T	T	G
j	0.00045	1	1	A	A	G	A	T	G	T	T	G
u	0.00045	1	1	A	A	G	A	T	G	T	T	G
aa	0.00045	1	1	A	A	G	A	T	G	C	T	G
v	0.00045	1	1	A	A		A	T		T	T	G
s	0.00045	1	1	A	A	G	A		G	T	T	G
ab	0.00045	1	1	A	A	G	A	T	G	T	T	G
q	0.00045	1	1	T	A	G	A	T	G	T	T	G
z	0.00045	1	1	A	A	G	A	T	G	T	T	G
af	0.00045	1	1	A	A	G	A	T	G	T	T	G
ad	0.00045	1	1	A	A	G	A	T	G	T	T	G

Table 13.98 - IGKV4-1

ac	1	A	A	G	A	T	G	T	T	G	T	G
m	1	A	A	G	A	T	G	T	T		T	
p	1	A	G	G	A	T	G	T	T	G	T	G
ae	1	A	A	G	A	T	G	T	T	G	T	G

Table 13.98 - IGKV4-1

h.type	cum.freq	# indvs	pops	2:89185464	2:89185466	2:89185474	2:89185476	2:89185482	2:89185484	2:89185485	2:89185491	2:89185511
				rs139176691	rs186837913	rs142402063	rs191518466	rs143877691	rs148599313	rs142986022	rs140954172	rs186251122
ref	0.9516	-	-	Ser->Phe	Asn->His	Lys->Asn	Asn->Ser	Leu->Ser	Ala->Pro	Ala->Val	Tyr->Phe	Pro->Thr
a	0.01185	26	11	C	A	G	A	T	G	C	A	C
b	0.00545	11	9	C	A	G	A	T	G	C	A	C
c	0.00365	8	7	C	A	G	A	T	G	T	A	C
d	0.00315	7	6	C	A	G	A	T	G	C	A	C
k	0.0018	4	4	C	A	G	A	T	G	C	A	C
w	0.0018	3	3	C	A	G	A	T	C	C	A	C
h	0.0018	4	4	C	A	G	A	T	G	C	A	C
f	0.0018	4	4	C	A	G	A	T	G	C	A	C
e	0.00135	3	3	C	A	G	A	T	G	C	T	C
r	0.00135	3	3	C	A	G	A	T	G	C	A	C
x	0.00135	3	2	C	A	G	A	T	G	C	A	C
o	0.00135	3	2	C	A	G	G	T	G	C	A	C
y	0.0009	2	2	C	A	C	A	T	C	C	A	C
g	0.0009	2	2	C	A	G	A	T	G	C	A	C
t	0.0009	2	2	C	A	C	A	T	G	C	A	C
i	0.0009	2	2	C	A	G	A	T	G	C	A	C
n	0.0009	2	2	C	A	G	A	T	G	C	A	C
l	0.0009	2	2	C	A	G	A	T	G	C	A	C
j	0.00045	1	1	C	A	G	A	T	G	C	A	C
u	0.00045	1	1	C	A	C	A	T	C	C	A	C
aa	0.00045	1	1	C	A	G	A	T	G	C	A	C
v	0.00045	1	1	C	A	G	A	T	G	C	A	C
s	0.00045	1	1	C	A	G	A	T	G	C	A	C
ab	0.00045	1	1	C	A	G	A	T	G	T	A	C
q	0.00045	1	1	C	C	G	A	T	G	C	A	C
z	0.00045	1	1	C	A	G	A	T	G	C	A	C
af	0.00045	1	1	C	A	G	A	T	C	C	A	C
ad	0.00045	1	1	C	A	G	A	T	G	C	A	C

Table 13.98 - IGKV4-1

ac 1 0.00045
m 1 0.00045
p 1 0.00045
ae 1 0.00045

1	C	A	G	A	T	G	C	A	C
1	C	A	G	A	T	G	C	T	C
1	C	A	G	A	T	G	C	A	C
1	T	A	C	A	T	G	C	A	C

Table 13.98 - IGKV4-1

h.type	cum.freq	# indvs	pops	2:89185518	2:89185529	2:89185530	2:89185563	2:89185577	2:89185586	2:89185607	2:89185622	2:89185637
				rs141640067	rs145197543	rs188057788	rs192839829	rs138751137	rs141777505	rs182919936	rs188769218	rs145449257
ref	0.9516	-	-	Lys->Arg	Tyr->Asp	Tyr->Cys	Asp->Gly	Ser->Arg	Gly->Arg	Ile->Phe	Ala->Thr	Val->Leu
a	0.01185	26	11	A	T	A	A	A	G	A	G	G
b	0.00545	11	9	A	T	A	A	A	G	A		G
c	0.00365	8	7	A	T	A	A	A	G	A	G	G
d	0.00315	7	6	A	T	A	A	A	G	A	G	G
k	0.0018	4	4	A	T	A	A	A	G	A	G	G
w	0.0018	3	3	A	T	A	A	A	G	A	G	G
h	0.0018	4	4	A	T	A	A	A	G	A	G	G
f	0.0018	4	4	A	T	A	A	A	G	A	G	G
e	0.00135	3	3	A	T	A	A	A	G	A	G	G
r	0.00135	3	3	A	T	A	A	A	G	A	G	G
x	0.00135	3	2	A	T	A	A	A	G	A	G	C
o	0.00135	3	2	A	T	A	A	A	G	A	G	G
y	0.0009	2	2	A	T	A	A	A	G	A	G	G
g	0.0009	2	2	A	T	G	A	A	G	A	G	G
t	0.0009	2	2	A	T	A	A	A	G	A	G	G
i	0.0009	2	2	A	G	A	A	A	G	A	G	G
n	0.0009	2	2	G	T	A	A	A	G	A	G	G
l	0.0009	2	2	A	T	A	A	A	G	A	G	G
j	0.00045	1	1	A	T	A	A	A	G	A	G	G
u	0.00045	1	1	A	T	A	A	C	G	A	G	G
aa	0.00045	1	1	A	T	A	A	A	G	A	G	G
v	0.00045	1	1	A	T	A	A	A	G	A	G	G
s	0.00045	1	1	A	T	A	A	A	G	A	G	G
ab	0.00045	1	1	A	T	A	A	A	G	A		G
q	0.00045	1	1	A	T	A	A	A	G	A	G	G
z	0.00045	1	1	A	T	A	A	A	G	T	G	G
af	0.00045	1	1	A	T	A	A	A	G	A	G	G
ad	0.00045	1	1	A	T	A	G	A	G	A	G	G

Table 13.98 - IGKV4-1

ac 1 0.00045
m 1 0.00045
p 1 0.00045
ae 1 0.00045

1	A	T	A	A	A	A	A	A		A	G	G
1	A	T	A	A	A	A	A	A	G	A	G	G
1	A	T	A	A	A	A	A	A	G	A	G	G
1	A	T	A	A	A	A	A	A	G	A	G	G

Table 13.98 - IGKV4-1

h.type	cum.freq	# indvs	pops	2:89185664	
				rs192141151	
ref	0.9516	-	-	Thr->Ser	ENST000000390243
a	0.01185	26	11	A	1,2,3,4,5,7,9,10,11,13,14
b	0.00545	11	9	A	2,3,4,5,6,10,12,13,14
c	0.00365	8	7	A	2,3,5,6,7,11,13
d	0.00315	7	6	A	2,5,6,7,9,11
k	0.0018	4	4	A	2,5,7,11
w	0.0018	3	3	A	2,5,14
h	0.0018	4	4	A	6,7,11,14
f	0.0018	4	4	A	4,5,6,14
e	0.00135	3	3	A	5,9,13
r	0.00135	3	3	T	5,7,8
x	0.00135	3	2	A	1,4
o	0.00135	3	2	A	9,11
y	0.0009	2	2	A	11,13
g	0.0009	2	2	A	5,6
t	0.0009	2	2	A	13,14
i	0.0009	2	2	A	9,14
n	0.0009	2	2	A	1,2
l	0.0009	2	2	A	2,11
j	0.00045	1	1	A	13
u	0.00045	1	1	A	13
aa	0.00045	1	1	A	7
v	0.00045	1	1	A	10
s	0.00045	1	1	A	11
ab	0.00045	1	1	A	13
q	0.00045	1	1	A	5
z	0.00045	1	1	A	11
af	0.00045	1	1	A	3
ad	0.00045	1	1	A	2

Table 13.98 - IGKV4-1

ac	0.00045	1	A	1
m	0.00045	1	A	12
p	0.00045	1	A	14
ae	0.00045	1	A	9

Table 13.99 - IGKV5-2

h.type	cum.freq	# indvs	pops	2:89197020	2:89197105	2:89197190
ref				rs183891175	rs189117859	rs55661410
				Thr->Arg	Asp->Glu	Pro->Ser
	0.6714	-	-	C	T	C
a	0.32315	501	14	C	T	T
b	0.005	11	6	C	G	T
c	0.00045	1	1	G	T	C

ENST00000390244

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,5,6,7,11,13

11

Table 13.100 - IGKV6-21

h.type	cum.freq	# indvs	pops	2:89459266		2:89459267	ENST00000390256
				rs181879031	Thr->Met	rs395483	
ref	0.9638	-	-		G	T	
a	0.03575	78	14		G	C	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.00045	1	1			T	14

Table 13.101 - IGKV6D-21

h.type	cum.freq	# indvs	pops	2:90060955	2:90060957	2:90060963
				rs190780280	rs141599034	rs150509294
ref	0.9735	42	13	Asp->Gly	Ala->Thr	Ala->Thr
				Asp->Gly	Ala->Thr	Ala->Thr
				Asp->Gly	Ala->Thr	Ala->Thr
				Asp->Gly	Ala->Thr	Ala->Thr
a	0.0192	14	4	A	G	G
b	0.00685	1	1	A	G	G
c	0.00045			G		

ENST000000436451
ENST000000557853
ENST000000558079

1,2,3,4,5,6,7,8,9,10,11,12,14
1,5,10,14
3

Table 13.102 - IGKV6D-41

h.type	cum.freq	# indvs	pops	2:90108534 rs191316044	2:90108797 .	2:90109001 rs192662836	
				Val->Ala	->Asp	Thr->Ile	ENST000000390271
ref	0.99865	-	-	T	ATGT	C	
a	0.00045	1	1	C	ATGT	C	3
c	0.00045	1	1	T	ATGT	T	6
b	0.00045	1	1	T		C	9

Table 13.103 - IGLJ1

h.type	cum_freq	# indvs	pops	22:23235930	
				rs182236201	
ref	0.9982	4	-	Arg->Gln	ENST000000390320
				Arg->Gln	ENST000000526893
				Arg->Gln	ENST000000532223
a	0.0018	2	-	G	
					3,4

Table 13.104 - IGLJ2

h.type	cum.freq	# indvs	pops	22:23241713	22:23241772	22:23241773	22:23241777	22:23241802	22:23241802
				Ala->Gly	Phe->Leu	Phe->Tyr	Leu->Phe	->Val	Val->Ile
ref	0.90495	-	-	C	T	T	G	GTAT	G
a	0.0741	160	13	G	T	T	G	GTAT	G
b	0.0178	38	12	C	C	T	G	GTAT	G
e	0.0009	2	2	G	C	T	G	GTAT	G
c	0.0009	2	2	C	T	T	C	GTAT	G
d	0.00045	1	1	C	T	T	G	GTAT	■
g	0.00045	1	1	C	T	■	G	GTAT	G
f	0.00045	1	1	C	T	T	G	G	G

ENST000000390322

1,2,3,4,5,6,7,9,10,11,12,13,14

1,2,3,4,5,7,9,10,11,12,13,14

4,9

1,10

2

13

2

Table 13.105 - IGLJ3

h.type	cum.freq	# indvs	pops	22:23247139	22:23247164	
				rs1852223887	rs139014240	
				Gly-> Arg	Gln->Arg	ENST000000390324
					->	ENST000000448061
ref	0.9973	-	-	G	A	
a	0.00225	4	2	G	G	2,13
b	0.00045	1	1	■	A	2

Table 13.106 - IGLJ5

h.type	cum.freq	# indivs	pops	22:23256422 rs189343862	22:23256422 rs192396255	22:23256465 rs192396255	ENST000000390327
ref	0.99545	-	-	Leu->Phe G	Glu->Lys G		
a	0.00365	8	5	T	G		1,6,7,12,13
b	0.0009	2	1	G			14

Table 13.107 - IGLJ6

h.type	cum.freq	# indvs	pops	22:23260329 rs76121504	22:23260345 rs114500104	22:23260367 rs184683407	
ref	0.9895	-	-	Ser->Leu C	Phe->Leu C	Val->Ile G	ENST000000390328
a	0.00595	13	3	T	C	G	1,12,14
b	0.0032	7	3	C	G	G	1,10,14
c	0.00135	3	2	C	C		1,10

Table 13.108 - IGLJ7

h.type	cum.freq	# indvs	pops	22:23263601	22:23263602	
ref	0.0664	-	-	rs150162743	rs373405	ENST00000390330
a	0.93225	1084	14	Ala->Thr	Ala->Val	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.00135	3	3	G	C	1,10,14
				G	T	
					T	

Table 13.109 - IGLV10-54

h.type	cum.fr	# indivs	pops	22:22569242	22:22569357	22:22569442	22:22569448	22:22569451	22:22569468	22:22569554	22:22569559
ref			-								
a	0.3126	812	14	G	T	G	T	T	G	C	C
b	0.50725	196	14	G	T	G		T	G		C
c	0.09425	109	9	G	T	G		T	G	C	C
d	0.05525	27	5	G	T	G		C	G	C	C
e	0.01265	8	3	G	T	G		T	G		C
f	0.0036	7	2	G	T	G		T	G		C
h	0.00315	6	2	G	T	G		T	G		C
i	0.0027	6	2	G	T	G		T	G		C
k	0.0027	4	2	G	T	G		T	G		C
g	0.0018	3	1	G	T	G		T	G		T
j	0.00135	2	2	G		G		T	G	C	C
n	0.0009	2	2	G	T	G		T	G	C	C
m	0.0009	1	1	G	T	C		T	G	C	C
l	0.00045	1	1	C	T	G		T	G	C	C
	0.00045			G	T	G		T	G	C	C

Table 13.109 - IGLV10-54

h.type	cum.fr	# indivs	pops	22:22569562	22:22569609	22:22569618	22:22569640	22:22569649
ref	0.3126	-	-					
a	0.50725	812	14	C	G	G	T	G
b	0.09425	196	14	C	G	G	G	G
c	0.05525	109	9	C	G	G	G	G
d	0.01265	27	5	C	G	G	G	G
e	0.0036	8	3	C		G	G	G
f	0.00315	7	2	C	G	G	G	
h	0.0027	6	2	C	G	G	G	G
i	0.0027	6	2	C	G	C	G	G
k	0.0018	4	2	C	G	G	G	G
g	0.00135	3	1	C	G	G	G	G
j	0.0009	2	2	C	G	G	T	G
n	0.0009	2	2	C	G	G	G	G
m	0.00045	1	1	C	G	G	G	G
l	0.00045	1	1	G	G	G	T	G

ENST000000390287

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,3,5,6,9,10,11,12,14

1,5,10,12,14

10,11,14

10,14

10,14

1,14

10,14

10

11,13

2,14

13

11

Table 13.110 - IGLV11-55

h.type	cum.frec	# indivs	pops	22:22556076	22:22556114	22:22556368	22:22556443	22:22556479	22:22556517	22:22556577
ref	0.4994	-	-	Leu->Gln	Gly->ArgGly->Arg	Pro->Leu	Arg->Gln	Ala->Val	Ala->Pro	Val->Phe
a	0.4463	728	14	T	G	C	G	C	G	G
b	0.0325	71	12	T	C	T	G	C	G	G
c	0.0096	21	6	T	G	T	G	T	G	G
d	0.00545	12	6	T	C	C	G	C	G	G
g	0.00225	5	3	T	G	T	G	T	G	T
e	0.0018	4	3	■	G	T	G	C	G	G
h	0.00135	3	2	T	G	C	G	C	C	G
f	0.00135	3	2	T	G	T	■	C	G	G

ENST000000390286

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,5,6,7,8,9,10,11,13,14

3,4,6,9,12,13

2,4,9,10,13,14

1,10,14

2,5,13

1,10

4,9

Table 13.111 - IGLV1-36

h.type	cum.freq	# indvs	pops	22:22786604	22:22786673	22:22786674	22:22786692	22:22786709	22:22786750	22:22786792	22:22786797
ref	0.8795	-	-	rs61735471	rs190488947	rs118006119	rs182633696	rs139155569	rs59686180	rs57337788	rs61735468
a	0.0942	185	14	Ala->Thr	Pro->Thr	Pro->Leu	Arg->Gln	Ser->Pro		Ser->Arg	Asn->Ser
b	0.00685	15	5	G	C	C	G	T	TGAG	C	A
d	0.0068	15	4	G	C	T	G	T	TGAG	G	A
c	0.00455	10	2	G	C	C	G	T	TGAG	C	G
f	0.0036	8	2	G	■	C	G	C	TGAG	C	A
g	0.0027	6	3	G	C	C	G	T	T	C	A
e	0.0018	4	3	G	C	C	■	T	TGAG	C	A

ENST000000390301

1,2,3,4,5,6,7,8,9,10,11,12,13,14
2,6,7,12,13
1,10,12,14
10,14
10,14
1,4,10
2,6,7

Table 13.112 - IGLV1-40

h.type	cum.freq	# indivs	pops	22:22764185	22:22764350	22:22764375	22:22764392	22:22764400	22:22764410	22:22764421	22:22764436	22:22764455
				rs146878832	rs186944811	rs183685135	rs143677402	rs187953594	rs141905358	rs146585623	rs185509404	rs190358290
ref	0.94895	-	-	His->Tyr	Gly->Glu	Ile->Met	Ser->Thr	Ile->Phe	Gly->Asp	His->Asp	Leu->Phe	Lys->Arg
a	0.0302	66	10	C	G	C	G	A	G	C	C	A
b	0.0046	10	5	C	G	C	G	T	G	C	C	A
d	0.0036	7	7	C	G	C	G	A	G	C	C	A
c	0.00275	6	2	C	G	C	G	A	G	C	C	A
f	0.0018	4	3	T	G	C	G	A	G	C	C	A
g	0.00135	3	3	C	■	C	G	A	G	C	C	A
e	0.0009	2	2	C	G	C	G	A	G	C	T	A
h	0.0009	2	2	C	G	C	G	A	■	C	C	A
j	0.00045	1	1	C	G	C	C	A	G	C	C	G
k	0.00045	1	1	C	G	C	G	A	G	C	T	A
s	0.00045	1	1	C	G	C	G	A	G	C	C	A
q	0.00045	1	1	C	G	G	G	A	G	C	C	G
r	0.00045	1	1	C	G	C	G	A	G	G	C	A
i	0.00045	1	1	C	G	C	C	A	G	C	C	A
n	0.00045	1	1	C	G	C	G	A	G	C	T	A
m	0.00045	1	1	C	G	C	G	A	G	C	C	A
l	0.00045	1	1	C	G	C	C	A	G	C	C	A
p	0.00045	1	1	C	G	C	G	A	■	C	C	A
o	0.00045	1	1	C	G	C	G	A	G	C	C	G

Table 13.112 - IGLV1-40

h.type	cum.freq	# indvs	pops	22:22764472	22:22764473	22:22764476	22:22764487	22:22764512	22:22764541	22:22764548
				rs181685335	rs185969886	rs111666735	rs138088831	rs181089093	rs190749695	rs143634461
ref	0.94895	-	-	Asn->Tyr	Asn->Ser	Ser->Asn	Ser->Ala	Gly->Ala	Ala->Thr	Thr->Ser
a	0.0302	66	10	A	A	G	T	G	G	C
b	0.0046	10	5	A	A	G	T	G	G	C
d	0.0036	7	7	A	A	■	T	G	G	G
c	0.00275	6	2	A	A	G	G	G	G	C
f	0.0018	4	3	A	A	G	T	G	G	C
g	0.00135	3	3	A	A	G	T	G	G	C
e	0.0009	2	2	A	A	G	T	G	G	C
h	0.0009	2	2	A	A	G	T	G	G	C
j	0.00045	1	1	A	A	G	T	G	G	C
k	0.00045	1	1	A	G	G	T	G	G	C
s	0.00045	1	1	T	A	G	T	G	G	C
q	0.00045	1	1	A	A	G	T	G	G	C
r	0.00045	1	1	A	A	G	T	G	G	C
i	0.00045	1	1	A	A	■	T	G	G	C
n	0.00045	1	1	A	A	■	T	G	G	C
m	0.00045	1	1	A	A	G	T	C	G	C
l	0.00045	1	1	A	A	G	T	G	G	C
p	0.00045	1	1	A	A	G	T	G	■	C
o	0.00045	1	1	A	A	G	T	G	G	C

ENST000000390299

1,2,3,4,5,6,7,10,13,14

2,6,7,12,13

1,2,5,10,11,13,14

10,14

1,10,14

2,7,12

10,13

2,3

1

1

13

10

10

10

2

2

10

1

5

Table 13.113 - IGLV1-44

h.type	cum.fre	# indvs	pops	22:22735255	22:22735448	22:22735517	22:22735551	22:22735573	22:22735661	22:22735678	22:22735710
ref	0.9531	-	-	rs183817639	rs180819320	rs61731361	rs190708617	rs111733579	rs6002086	rs181994204	rs149125249
				Ser->Gly	Ala->Thr	Thr->Ala	Ala->Gly	Ser->Arg	Ser->Ala	Asp->Glu	Asn->Ser
a	0.0292	60	12	A	G	A	C	T	T	T	A
b	0.00775	16	2	A	G	A	C	G	T	T	A
c	0.00365	8	4	A	G	A	C	G	G	T	A
d	0.0018	4	3	G	G	A	C	T	T	T	A
e	0.0018	4	3	A	G	A	C	T	T	■	A
f	0.00135	3	2	A	G	A	C	T	T	T	G
g	0.00045	1	1	A	G	G	C	T	T	T	A
h	0.00045	1	1	A	■	A	C	T	T	T	A
i	0.00045	1	1	A	G	A	C	G	T	T	G
											ENST000000390297
											1,2,3,4,5,6,7,9,10,11,13,14
											10,14
											6,7,11,13
											4,11,13
											7,13,14
											1,14
											9
											9
											3

Table 13.114 - IGLV1-47

h.type	cum.fr	# indivs	pops	22:22712158	22:22712333	22:22712403	22:22712410	22:22712444	22:22712467	22:22712552	22:22712562
ref	0.11455	-	-	rs12160269 Leu->Ile	rs185378771 Pro->Ala	rs181574862 Gly->Glu	rs185842267 Asn->Lys	rs8142074 Ala->Thr	rs5757973 Ser->Arg	rs61231611 Arg->Trp	rs184225865 Asp->Ala
a	0.8413	1059	14	C	C	G	T	G	T	C	A
b	0.0182	40	4	C	C	G	T	G	G	C	A
c	0.01695	33	4	C	C	G	T	G	G	C	A
d	0.00405	9	3	C	C	G	T	G	G	T	A
g	0.0018	4	2	C	C	G	G	G	G	C	A
f	0.00135	3	2	C	C	G	T	G	G	C	A
e	0.0009	2	2	C	C	G	T	G	T	C	A
h	0.00045	1	1	C	G	G	T	G	G	C	A
i	0.00045	1	1	C	C	G	T	G	G	C	C

ENST000000390294

1,2,3,4,5,6,7,8,9,10,11,
12,13,14
1,10,12,14
1,5,10,14
1,10,14
10,14
8,14
1,14
13
10

h.type	cum.freq	# indivs	pops						
				22:22681721	22:22681967	22:22682039	22:22682063		
ref	0.7922	-	-	T	C	A	A	Gln->Arg	ENST00000390291
a	0.19325	373	14	C	C	A	A		1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.0105	23	5	T	C	A	G		1,10,11,12,14
c	0.00225	5	3	T	C	G	A		10,12,13
d	0.0018	4	2	T	T	A	A		3,9

Table 13.116 - IGLV1-51

h.type	cum.freq	# indvs	pops	22:22677182	22:22677235	22:22677285	22:22677298	22:22677307	22:22677309	22:22677318
ref	0.8737	-	-	Asp->Glu	Gly->Asp	Asp->Asn	Gly->Val	Asp->Gly	Ser->Arg	Ser->Arg
a	0.1177	221	11		G	G	G	A	A	A
b	0.00275	6	3	C	G		G	A	A	A
d	0.0018	4	2		G	G	G	A	A	C
c	0.0018	4	3	C	G	G	G	A	A	C
g	0.0009	2	2	C		G	G	A	A	A
e	0.00045	1	1	C	G	G	G	A	C	A
h	0.00045	1	1	C	G	G	T	A	A	A
f	0.00045	1	1	C	G	G	G	G	A	A

ENST000000390290

1,2,5,6,7,8,10,11,12,13,14

2,5,12

3,4

3,4,9

6,10

6

13

6

Table 13.117 - IGLV2-11

h.type	cum.freq	# indivs	pops	22:23135047 rs185062160	22:23135342 rs187815581	22:23135358 rs193000225	22:23135360 rs185237664	22:23135376 rs189822640	22:23135379 rs181371196	22:23135382 rs184627661	22:23135466 rs142176771
ref	0.9745	-	-	Ser->Asn	Met->Ile	Lys->Gln	Lys->Asn	Pro->Ser	Asp->Asn	Arg->Cys	Cys->Ser
a	0.00775	17	3	G	G	A	G	C	G	C	T
b	0.00455	10	7	G	G	A	T	C	G	C	
d	0.0037	8	5	G	G	A	G	T		C	T
c	0.0032	7	5	G	G	A	T	T		C	T
g	0.00225	5	3	G	G	A	G	C		C	T
j	0.0009	2	2	G	G	A	T	C	G	C	
e	0.0009	2	1		G	A	G	C	G	C	T
f	0.0009	2	2	G	G	C	G	C	G	C	T
i	0.0009	2	2	G	G	A	G	C	G	T	T
h	0.00045	1	1	G	C	A	G	C	G	C	T

ENST000000390314

3,4,9
2,6,7,10,11,12,13
1,5,10,11,14
2,3,9,12,14
2,10,14
3,9
6
6,13
2,11
14

Table 13.118 - IGLV2-14

h.type	cum.freq	# indivs	pops	22:23101248	22:23101255	22:23101480	22:23101494	22:23101497	22:23101559	22:23101568	22:23101600	22:23101615
				rs4822304	rs183579067	rs188164895	rs191388769	rs182533580	rs4134484	rs28561583	rs184972365	rs188991923
ref	0.26855	-	-	Leu->Phe	Thr->Asn	Ser->Thr	Gly->Ser	Tyr->Asn	Glu->Asp	Asn->Lys	Gly->Val	Asn->Ser
a	0.48365	750	14	T	C	G	G	T	G	T	G	A
c	0.08855	191	14	C	C	G	G	T	T	T	G	A
b	0.0804	159	10	C	C	G	G	T	T	G	G	A
d	0.02365	51	9	C	C	G	G	T	T	T	G	A
g	0.0104	22	3	C		G	G	T	G	T	G	A
e	0.00995	19	3	C	C	G	G	T	T	G	G	A
h	0.0068	15	4	C	C	G	G	T	G	G	G	A
f	0.0055	12	6	C	C	G	G	T	G	T	G	A
l	0.0036	8	4	T	C	G	G	T	T	G	G	A
j	0.0027	6	5	T	C	G	G	T	G	T	G	A
m	0.0027	6	5	T	C	G	G	T	T	T	G	A
n	0.00225	5	3	C	C	G	G	T	G	G	G	A
i	0.00185	4	2	T	C	G		T	T	T	G	A
k	0.0018	4	4	T	C	G	G	T	G	G	G	A
r	0.0018	4	1	T		G	G	T	T	T	G	A
q	0.0009	2	1	C	C	G	G	T	G	T	G	G
p	0.0009	2	2	C		G	G	T	T	T	G	A
o	0.0009	2	1	C		G		T	G	T	G	A
y	0.00045	1	1	T	C	G	G	T	T	G	T	A
u	0.00045	1	1	C	C	C	G	T	G	T	G	A
t	0.00045	1	1	C	C	G		T	G	T	G	A
v	0.00045	1	1	C	C	G	G	T	G	T	G	A
s	0.00045	1	1	C	C	G	G	T	G	T	G	A
w	0.00045	1	1	C	C	G		T	T	G	G	A
x	0.00045	1	1	T	C	G	G		T	T	G	A

Table 13.118 - IGLV2-14

h.type	cum.freq	# indivs	pops	22:23101680	22:23101690	22:23101698
				rs183884653	rs188224577	rs16989343
ref				Tyr->Asn	Ser->Thr	Leu->Phe
				T	G	C
a	0.26855	-	-	T	G	C
	0.48365	750	14	T	G	C
c	0.08855	191	14	T	G	C
b	0.0804	159	10	T	G	T
d	0.02365	51	9	T	G	T
g	0.0104	22	3	T	G	C
e	0.00995	19	3	T	G	C
h	0.0068	15	4	T	G	T
f	0.0055	12	6	T	G	T
l	0.0036	8	4	T	G	C
j	0.0027	6	5	T	G	C
m	0.0027	6	5	T	G	T
n	0.00225	5	3	T	G	C
i	0.00185	4	2	T	G	C
k	0.0018	4	4	T	G	C
r	0.0018	4	1	T	G	C
q	0.0009	2	1	T	G	C
p	0.0009	2	2	T	G	C
o	0.0009	2	1	T	G	C
y	0.00045	1	1	T	G	C
u	0.00045	1	1	T	G	C
t	0.00045	1	1	T	G	C
v	0.00045	1	1	T	C	C
s	0.00045	1	1	■	G	C
w	0.00045	1	1	T	G	T
x	0.00045	1	1	T	G	C

ENST00000390312

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,3,4,5,6,9,10,11,12,14

1,2,3,4,5,6,9,10,14

3,4,9

5,10,11

3,4,9,10

1,9,10,12,13,14

2,4,11,12

1,9,10,12,13

3,4,6,9,10

2,5,12

2,9

6,9,11,12

14

10

2,13

9

10

6

6

1

1

9

2

Table 13.119 - IGLV2-18

h.type	cum.freq	# indvs	pops	22:23077436	22:23077555	ENST00000390310
				rs149656926	rs4822296	
				Glu->Asp	Leu->Ser	
ref	0.2279	-	-	G	T	
a	0.763	998	14	G	C	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.0091	19	6	T	C	1,3,4,7,10,14

Table 13.120 - IGLV2-23

h.type	cum.freq	# indivs	pops	22:23040461	22:23040637	22:23040638	22:23040685	22:23040697	22:23040700	22:23040724	22:23040765	22:23040767
				rs12166262	rs192197296	rs185785000	rs190209511	rs193192358	rs61749282	rs188648111	rs61732325	rs5751516
ref	0.58265	-	-									
a	0.34655	593	14	C	G	T	A	G	A	C	G	G
b	0.0405	82	12		G	T	A	G	A	C	G	T
c	0.01545	31	5	C	G	T	A	G	A	C	T	T
d	0.00315	7	5	C	G		A	G	A	C	G	T
g	0.00315	7	5	C	G	T	A	G	A	C	G	G
f	0.00225	5	2		G	T	A	G	A	C	T	T
e	0.00135	3	3	C	G	T	A	G	A	C	G	G
k	0.0009	2	2		G	T	A	G	A	C	G	G
o	0.0009	2	2	C	G	T	A	G	A	C	T	G
j	0.00045	1	1	C	G	T	T	G	A	C	G	G
h	0.00045	1	1	C		T	A	G	A	C	G	G
i	0.00045	1	1	C	G	T	A		A	C	G	G
n	0.00045	1	1	C	G	T	A	G	A	G	G	G
m	0.00045	1	1	C		T	A	G	A	C	G	T
l	0.00045	1	1	C	G	T	A	G	C	C	G	G
p	0.00045	1	1	C	G	T	A	G	A	C	T	G

Table 13.120 - IGLV2-23

h.type	cum.freq	# indivs	pops	22:23040887	
				rs5759376	
ref	0.58265	-	-	Tyr->Cys	ENST00000390306
a	0.34655	593	14	A	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.0405	82	12	A	1,2,3,4,5,7,9,10,11,12,13,14
c	0.01545	31	5	G	3,4,5,9,11
d	0.00315	7	5	A	2,5,6,7,12
g	0.00315	7	5	A	1,2,9,10,14
f	0.00225	5	2	A	10,13
e	0.00135	3	3	G	3,9,11
k	0.0009	2	2	A	8,10
o	0.0009	2	2	A	3,10
j	0.00045	1	1	A	9
h	0.00045	1	1	A	3
i	0.00045	1	1	A	9
n	0.00045	1	1	A	14
m	0.00045	1	1	A	13
l	0.00045	1	1	A	5
p	0.00045	1	1	G	11

Table 13.121 - IGLV2-33

h.type	cum.freq	# indvs	pops					
				22:22930684	22:22930896	22:22931059		
				rs11914178	rs143638085	rs185353299		
				Met->Val			Gly->Val	ENST000000390302
ref	0.98955	-	-	A	G		G	
a	0.00775	16	4	G	G		G	10,11,12,14
b	0.0018	4	3	A	G		T	2,5,13
c	0.0009	2	1	A			G	14

Table 13.122 - IGLV2-8

h.type	cum.freq	# indvs	pops	22:23165340	22:23165526	22:23165610	22:23165680	22:23165701	22:23165757
ref	0.8422	-	-	rs5996397	rs191286893	rs185026152	rs149347098	rs144694239	rs11558656
				Thr->Ser	Pro->Ser	Pro->Ser	Ser->Phe	Thr->Met	Ser->Cys
a	0.1451	267	11	C	C	C	C	C	A
b	0.0046	10	7	G	C	C	C	C	A
c	0.00315	7	3	C	C	C	T	C	A
d	0.0027	6	3	G	C	C	C	C	T
e	0.0018	4	2	C	C	C	C	T	A
f	0.00045	1	1	C	C	T	C	C	A
									ENST000000390317
									1,2,5,6,7,8,10,11,12,13,14
									2,5,6,7,11,12,13
									3,4,9
									1,10,14
									4,7
									2

Table 13.123 - IGLV3-10

h.type	cum.freq	# indivs	pops	22:23154549	22:23154570	22:23154580	22:23154582	22:23154583	22:23154592	22:23154646	22:23154659	22:23154667
				rs182282578	rs192547759	rs183865794	rs1065464	rs188439978	rs184893196	rs141938356	rs188537134	rs191450268
ref	0.92545	-	-	Thr->Ala	Pro->Thr	Tyr->Cys	Ala->Pro	Ala->Val	Tyr->Cys	Arg->Gln	Ile->Met	Arg->Lys
a	0.049	97	10	A	C	A	G	C	A	G	C	G
b	0.0206	41	4	A	C	A	C	C	A	G	C	G
c	0.00315	6	2	A	C	A	G	C	A	■	C	G
d	0.00045	1	1	A	C	G	G	C	A	G	C	G
g	0.00045	1	1	A	■	A	G	C	A	G	G	G
e	0.00045	1	1	G	C	A	G	C	G	G	C	■
f	0.00045	1	1	A	C	A	G	T	A	G	C	G

Table 13.123 - IGLV3-10

h.type	cum.freq	# indivs	pops	22:23154736	rs802223369	ENST000000390315
ref	0.92545	-	-	Ala->Asp	C	
a	0.049	97	10	C		1,2,5,6,7,8,11,12,13,14
b	0.0206	41	4			1,10,12,14
c	0.00315	6	2	C		1,10
d	0.00045	1	1	C		2
g	0.00045	1	1	C		11
e	0.00045	1	1	C		2
f	0.00045	1	1	C		11

Table 13.124 - IGLV3-12

h.type	cum.freq	# indvs	pops	22:23114327	22:23114350	22:23114938	22:23114993	22:23115059
				rs6003299	rs5996382	rs2073450	rs2073451	rs190626294
ref				Met->Thr	Ser->Gly	Ser->Arg	Ala->Thr	Ser->Gly
a	0.42115	-	-	T	A	C	G	A
b	0.38715	653	14	C	G	C		A
c	0.1116	220	14	C	G	C	G	A
d	0.0774	158	14	C	G		G	A
e	0.00135	3	3	T	A	C		A
f	0.00045	1	1	C	G	C		G
g	0.00045	1	1	C	A	C	G	A
h	0.00045	1	1	C	A	C		A
i	0.00045	1	1	C	A	C		A

ENST000000390313

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

2,4,14

1

4

10

Table 13.125 - IGLV3-16

h.type	cum.fre	# indivs	pops	22:23090155	22:23090222	22:23090242	22:23090243	22:23090248	22:23090260	22:23090327	22:23090381
ref	0.9395	-	-	Ser->Thr	Tyr->Phe	Phe->Val	Phe->Ser	Val->Met	Tyr->Asn	Val->Asp	Leu->Gln
a	0.04295	82	6	T	A	T	T	G	T	T	T
b	0.0081	18	9	T	A	G	C	G	T	T	T
c	0.0045	10	5	T	A	T	T	G	T	T	
h	0.0018	4	3	T	A	G	C	G	T	T	T
d	0.00135	3	3	T	A	T	T	G	T		
g	0.00045	1	1	T	A	T	T		T	T	T
e	0.00045	1	1	T	T	T	T	G		T	T
f	0.00045	1	1		A	T	T	G	T	T	T
i	0.00045	1	1	T	T	G	C	G	T	T	T

ENST000000390311

1,5,10,11,12,14

1,2,5,7,9,10,12,13,14

1,9,10,11,14

1,10,14

2,5,9

5

10

13

10

Table 13.126 - IGLV3-19

h.type	cum.freq	# indvs	pops	22:23063178	22:23063348	22:23063367	22:23063406	22:23063411	22:23063421	22:23063436	22:23063448	22:23063480
ref	0.86895	-	-	Leu->Phe	Val->Ile	Gln->Pro	Arg->Thr	Thr->Ala	Gly->Glu	Ser->Asn	Ser->Asn	Val->Ile
a	0.10015	190	12	T	G	A	G	A	G	G	G	G
b	0.01785	38	7	C	G	A	G	A	G	G	G	G
c	0.00225	5	2	C	G	A	G	A	■	G	G	G
d	0.00135	3	1	C	G	A	G	A	G	G	G	G
l	0.00135	3	2	C	G	A	G	A	G	G	■	G
j	0.0009	2	2	C	■	A	G	A	G	G	G	G
k	0.0009	2	2	C	G	A	G	A	G	■	G	G
h	0.0009	2	1	C	G	A	G	A	G	G	G	■
i	0.0009	2	1	C	G	A	G	A	G	G	■	■
p	0.0009	2	2	C	G	A	G	A	G	■	G	■
g	0.00045	1	1	C	G	A	G	A	G	■	■	G
e	0.00045	1	1	C	G	A	G	A	G	G	G	G
q	0.00045	1	1	C	G	C	G	A	G	G	G	G
r	0.00045	1	1	C	G	A	G	A	G	G	G	G
f	0.00045	1	1	C	G	A	G	A	G	G	G	G
n	0.00045	1	1	C	G	A	G	A	G	G	G	G
m	0.00045	1	1	C	G	A	G	G	G	■	■	G
o	0.00045	1	1	C	G	A	C	A	G	G	G	G

Table 13.126 - IGLV3-19

h.type	cum.freq	# indvs	pops	22:23063484	22:23063531	22:23063577	22:23063612	22:23063618
ref	0.86895	-	-	rs79338593	rs181016062	rs184342792	rs182162941	rs77723964
a	0.10015	190	12	Leu->Arg	Phe->Ile	Gly->Glu	Asn->Asp	Arg->Trp
b	0.01785	38	7	G	T	G	A	T
c	0.00225	5	2	T	T	G	A	C
d	0.00135	3	1	T	T	G	A	T
l	0.00135	3	2	T	T	G	A	C
j	0.0009	2	2	T	T	G	A	C
k	0.0009	2	2	T	T	G	A	C
h	0.0009	2	1	T	T	G	A	C
i	0.0009	2	1	T	T	G	A	C
p	0.0009	2	2	T	T	G	A	C
g	0.00045	1	1	G	T	G	A	T
e	0.00045	1	1	T	T	■	A	C
q	0.00045	1	1	T	T	G	A	C
r	0.00045	1	1	G	T	G	A	C
f	0.00045	1	1	T	T	G	G	C
n	0.00045	1	1	T	■	G	A	C
m	0.00045	1	1	T	T	G	A	C
o	0.00045	1	1	T	T	G	A	C

ENST000000390309

1,2,3,4,5,6,9,10,11,12,13,14

1,3,4,9,10,12,14

1,14

10

10,14

2,14

4,13

9

10

2,6

9

4

14

4

4

13

13

10

Table 13.127 - IGLV3-1

h.type	cum.fre	# indivs	pops	22:23223277	22:23223280	22:23223283	22:23223292	22:23223297	22:23223301	22:23223306	22:23223330
				rs144850905	rs184648956	rs61736464	rs145382269	rs181792449	rs186574328	rs61736463	rs145228815
ref	0.9255	-	-	Gly->AlaGly->Ala	Ser->Cys	Val->Glu	Tyr->Ser	Leu->Val	Thr->Ile	Pro->Ala	Gly->Arg
a	0.01865	40	13	G	C	T	A	C	C	C	G
c	0.00585	13	8	G	C	T	A	C	C	C	G
b	0.005	11	8	G	C	T	A	C	C	C	G
h	0.00315	7	7	G	C	T	A	C	C	C	G
o	0.00315	7	4	G	C	T	A	C	C	C	G
e	0.0027	6	5	G	C	T	C	C	C	C	G
t	0.00225	5	5	G	C	T	A	C	C	C	G
v	0.00225	5	5	G	C	T	A	C	C	C	G
f	0.00225	5	4	G	C	T	A	G	C	C	G
i	0.0018	4	4	G	C	T	A	C	C	C	G
n	0.0018	4	4	G	C	T	A	C	C	C	G
ag	0.00135	3	3	G	C	T	A	C	C	C	G
y	0.00135	3	3	G	C	T	A	C	C	C	G
s	0.00135	3	3	G	C	T	A	C	C	G	G
x	0.00135	3	3	G	C	T	A	C	T	C	G
ad	0.00135	3	2	G	C	T	A	C	C	C	G
l	0.00135	3	3	G	C	T	A	C	C	C	G
p	0.00135	3	2	G	C	T	A	C	C	C	G
ah	0.0009	2	2	G	C	T	A	C	C	C	G
d	0.0009	2	2	G	C	T	A	C	C	C	G
j	0.0009	2	1	G	C	T	A	C	C	C	G
g	0.0009	2	2	G	C	T	A	C	C	C	G
z	0.0009	2	2	G	C	T	A	C	C	C	G
ac	0.0009	2	2	G	C	T	A	C	C	C	G
ae	0.0009	2	2	G	C	T	A	C	C	C	G
u	0.00045	1	1	G	C	T	A	G	C	C	G
k	0.00045	1	1	G	C	T	A	C	C	C	G
aj	0.00045	1	1	G	C	T	A	C	C	C	C

Table 13.127 - IGLV3-1

aa	0.00045	1	C	C	T	A	C	C	C	G
aq	0.00045	1	G	C	■	A	C	C	C	G
al	0.00045	1	G	C	T	A	C	C	C	G
ab	0.00045	1	C	C	T	A	C	C	C	G
ak	0.00045	1	G	C	T	A	C	C	C	G
q	0.00045	1	G	C	T	A	C	C	C	G
ap	0.00045	1	G	C	T	A	C	C	C	G
w	0.00045	1	G	C	T	A	C	C	C	G
ao	0.00045	1	G	C	T	A	C	C	C	G
r	0.00045	1	G	C	T	A	C	C	C	G
af	0.00045	1	G	C	T	A	C	C	C	G
au	0.00045	1	G	C	T	A	C	C	C	G
am	0.00045	1	G	C	T	A	C	C	C	G
at	0.00045	1	G	C	T	A	C	C	C	G
m	0.00045	1	G	C	T	A	C	C	C	G
ar	0.00045	1	G	C	T	A	C	C	C	G
as	0.00045	1	G	C	T	A	C	C	C	G
ai	0.00045	1	G	G	T	A	C	C	C	G
an	0.00045	1	G	C	T	A	C	C	C	G

Table 13.127 - IGLV3-1

h.type	cum.fre	# indivs	pops	22:23223348	22:23223355	22:23223363	22:23223364	22:23223369	22:23223375	22:23223376	22:23223382	22:23223385
				rs189772470	rs183495921	rs187892119	rs191755097	rs188197731	rs192957059	rs61731694	rs138666033	rs61731693
ref	0.9255	-	-	Thr->Pro	Ser->Cys	Lys->Glu	Lys->Ile	Gly->Arg	Lys->Glu	Lys->Thr	Ala->Val	Cys->Tyr
a	0.01865	40	13	A	C	A	A	G	A	A	C	G
c	0.00585	13	8	A	C	A	A	G	A	A	T	G
b	0.005	11	8	A	C	A	A	G	A	A	C	
h	0.00315	7	7	A	C	A	A	G	A	A	C	G
o	0.00315	7	4	A	C	A	A	G	A	A	C	G
e	0.0027	6	5	A	C	A	A	G	A	A	C	G
t	0.00225	5	5	A	C	A	A	G	A	A	C	G
v	0.00225	5	5	A	C	A	A	G	A	A	C	G
f	0.00225	5	4	A	C	A	A	G	A	A	C	G
i	0.0018	4	4	C	C	A	A	G	A	A	C	G
n	0.0018	4	4	A	C	A	A	G	A	A	C	G
ag	0.00135	3	3	A	C	A	A	G	A	A	C	G
y	0.00135	3	3	A	C	A	A	G	A	C	C	G
s	0.00135	3	3	A	C	A	A	G	A	A	C	G
x	0.00135	3	3	A	C	A	A	G	A	A	C	G
ad	0.00135	3	2	A	C	A	A	G	A	A	C	G
l	0.00135	3	3	A	C	A	A	G	A	A	C	
p	0.00135	3	2	A	C	A	A	G	A	A	C	G
ah	0.0009	2	2	A	C	G	A	G	A	A	C	G
d	0.0009	2	2	A	C	A	A	G	A	A	C	G
j	0.0009	2	1	A	C	A	A	G	A	A	C	G
g	0.0009	2	2	A	C	A	A	G	A	A	C	G
z	0.0009	2	2	A	C	A	A	G	A	A	C	G
ac	0.0009	2	2	A	C	A	A	G	A	A	C	G
ae	0.0009	2	2	A	C	A	A	G	A	A	C	G
u	0.00045	1	1	A	C	A	A	G	A	A	C	G
k	0.00045	1	1	A	C	A	A	G	G	A	C	G
aj	0.00045	1	1	A	C	A	A	G	A	A	C	G

Table 13.127 - IGLV3-1

aa	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
aq	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
al	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
ab	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
ak	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
q	0.00045	1	C	C	A	A	A	A	G	A	A	T	G	G
ap	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
w	0.00045	1	A	G	A	A	A	A	G	A	A	C	G	G
ao	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
r	0.00045	1	A	C	A	A	A	T	G	A	A	C	G	G
af	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
au	0.00045	1	A	C	A	A	A	A	G	A	A	T	G	G
am	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
at	0.00045	1	A	C	A	A	A	T	G	A	A	C	G	G
m	0.00045	1	A	C	A	A	A	A	G	A	A	T	G	G
ar	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
as	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
ai	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G
an	0.00045	1	A	C	A	A	A	A	G	A	A	C	G	G

Table 13.127 - IGLV3-1

h.type	cum.fre	# indivs	pops	22:23223386	22:23223387	22:23223400	22:23223402	22:23223417	22:23223432	22:23223438	22:23223439	22:23223525
				rs61735526	rs186777494	rs61731688	rs61731687	rs190667156	rs73164939	rs143830449	rs61731685	rs61731681
ref	0.9255	-	-	Cys->Trp	Trp->Arg	Lys->Arg	Pro->Ala	Val->Met	Gln->Glu	Ser->Gly	Ser->Asn	Met->Leu
a	0.01865	40	13	C	T	A	C	G	C	A	G	A
c	0.00585	13	8	C	T	A	C	G	C	A		A
b	0.005	11	8	C	T	A	C	G	C	A	G	A
h	0.00315	7	7	C	T	A	C		C	A	G	A
o	0.00315	7	4	C	T	G	C	G	C	A	G	A
e	0.0027	6	5	C	T	A	C	G	C	A	G	A
t	0.00225	5	5	C	T	A	G	G	C	A	G	A
v	0.00225	5	5	C	T	A	C	G	G	A	G	A
f	0.00225	5	4	C	T	A	C	G	C	A	G	A
i	0.0018	4	4	C	T	A	C	G	C	A	G	A
n	0.0018	4	4	C	T	A	C	G	C	G	G	A
ag	0.00135	3	3	C	T	A	C	G	C	G		A
y	0.00135	3	3	C	T	A	C	G	C	A	G	A
s	0.00135	3	3	C	T	A	C	G	C	A	G	A
x	0.00135	3	3	C	T	A	C	G	C	A	G	A
ad	0.00135	3	2	C	T	A	C	G	C	A	G	A
l	0.00135	3	3	C	T	A	C	G	C	A		A
p	0.00135	3	2	C	T	A	C	G	C	A	G	A
ah	0.0009	2	2	C	T	A	C	G	C	A	G	A
d	0.0009	2	2	C	T	A	C	G	C	A	G	C
j	0.0009	2	1	C	T	A	C	G	C	A	G	A
g	0.0009	2	2	G	T	A	C	G	C	A		A
z	0.0009	2	2	C	T	A	C	G	C	A	G	A
ac	0.0009	2	2	G	T	A	C	G	C	A	G	A
ae	0.0009	2	2	C	T	A	C	G	C	A	G	A
u	0.00045	1	1	C	T	A	C	G	C	G	G	A
k	0.00045	1	1	C	T	G	C	G	C	G	G	A
aj	0.00045	1	1	C	T	A	C	G	C	A	G	A

Table 13.127 - IGLV3-1

aa	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
aq	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
al	0.00045	1	C	T	A	C	G	C	A		A		A
ab	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
ak	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
q	0.00045	1	G	T	A	C	G	C	A	G	A	G	A
ap	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
w	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
ao	0.00045	1	C		A	C	G	C	A	G	A	G	A
r	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
af	0.00045	1	C	T	G	C	G	C	A		A		A
au	0.00045	1	C	T	A	C	G	C	A		A		A
am	0.00045	1	G	T	A	C	G	G	A	G	A	G	A
at	0.00045	1	C	T	A	G	G	C	A	G	A	G	A
m	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
ar	0.00045	1	C	T	A	C	G	C	G	G	A	G	A
as	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
ai	0.00045	1	C	T	A	C	G	C	A	G	A	G	A
an	0.00045	1	C	T	A	C	G	C	A	G	A	G	A

Table 13.127 - IGLV3-1

h.type	cum.fre	# indvs	pops	22:23223531	22:23223535	22:23223540	22:23223541	22:23223544	22:23223550	22:23223573
				rs190043877	rs145774047	rs180853064	rs142180628	rs185803370	rs192317432	rs145570740
ref	0.9255	-	-	Glu->Lys	Ala->Gly	Tyr->Asp	Tyr->Ser	Tyr->Ser	Gln->Pro	His->Tyr
a	0.01865	40	13	G	C	T	A	A	A	C
c	0.00585	13	8	G	C	T	A	A	A	C
b	0.005	11	8	G	C	T	A	A	A	C
h	0.00315	7	7	G	C	T	A	A	A	C
o	0.00315	7	4	G	C	T	A	A	A	C
e	0.0027	6	5	G	C	T	A	A	A	C
t	0.00225	5	5	G	C	T	A	A	A	C
v	0.00225	5	5	G	C	T	A	A	A	C
f	0.00225	5	4	G	C	T	A	A	A	C
i	0.0018	4	4	G	C	T	A	A	A	C
n	0.0018	4	4	G	C	T	A	A	A	C
ag	0.00135	3	3	G	C	T	A	A	A	C
y	0.00135	3	3	G	C	T	A	A	A	C
s	0.00135	3	3	G	C	T	A	A	A	C
x	0.00135	3	3	G	C	T	A	A	A	C
ad	0.00135	3	2	G	C	T	A	A	A	T
l	0.00135	3	3	G	C	T	A	A	A	C
p	0.00135	3	2	G	C	T	A	C	A	C
ah	0.0009	2	2	G	C	T	A	A	A	C
d	0.0009	2	2	G	C	T	A	A	A	C
j	0.0009	2	1	G	C	T	C	A	A	C
g	0.0009	2	2	G	C	T	A	A	A	C
z	0.0009	2	2	G	C	T	A	A	C	C
ac	0.0009	2	2	G	C	T	A	A	A	C
ae	0.0009	2	2	G	G	T	A	A	A	C
u	0.00045	1	1	G	C	T	A	A	A	C
k	0.00045	1	1	G	C	T	A	A	A	C
aj	0.00045	1	1	G	C	T	A	A	A	C

ENST000000390319

2,3,4,5,6,7,8,9,10,11,12,13,14

2,5,6,7,9,10,13,14

1,4,5,6,10,11,13,14

1,3,4,5,9,10,14

3,7,10,14

1,2,3,10,11

2,3,5,10,14

1,5,6,7,9

2,4,10,13

1,7,10,11

6,12,13,14

2,5,13

5,10,12

2,9,10

3,10,14

2,13

2,6,13

5,9

9,14

2,13

2

6,12

1,4

6,11

13,14

14

11

2

Table 13.127 - IGLV3-1

aa	0.00045	1	G	C	T	A	A	A	C	5
aq	0.00045	1	G	C	T	A	A	A	C	13
al	0.00045	1	G	C	T	C		A	C	10
ab	0.00045	1	G	C	T	A	A	A	C	5
ak	0.00045	1	G	C	G	A	A	A	C	13
q	0.00045	1	G	C	T	A	A	A	C	2
ap	0.00045	1	G	C	T	A	A	A	C	5
w	0.00045	1	G	C	T	A	A	A	C	14
ao	0.00045	1	G	C	T	A	A	A	C	3
r	0.00045	1	G	C	T	A	A	A	C	2
af	0.00045	1	G	C	T	A	A	A	C	3
au	0.00045	1	G	C	T	A	A	A	C	1
am	0.00045	1	G	C	T	A	A	A	C	10
at	0.00045	1	G	C	T	A	A	A	C	10
m	0.00045	1	G	C	T	A	A	A	C	4
ar	0.00045	1	G	C	T	C		A	C	13
as	0.00045	1	G	C	T	A	A	A	C	1
ai	0.00045	1	G	C	T	A	A	A	C	2
an	0.00045	1		C	T	A	A	A	C	1

Table 13.128 - IGLV3-21

h.type	cum.fre	# indivs	pops	22:23054916	22:23054919	22:23054937	22:23055389	22:23055407	22:23055440	22:23055497	22:23055522	22:23055527
ref	0.2714	-	-	Val->Phe	Leu->Ile	Ser->Ala	Val->Met	Thr->Ala	Gln->Lys	Tyr->Asp	Pro->Leu	Leu->Val
a	0.31195	556	14	G	C	T	G	A	C	T	C	C
b	0.178	338	14	G	C	T	G	A		T	C	C
c	0.1613	296	14	T	C	T	G	A		T	C	C
f	0.02165	45	9	G	C	T	G	A		T	C	C
d	0.0173	36	10	T	C	T	G	A		T	C	C
g	0.01045	21	5	G	C	G	G	A	C	T	C	C
e	0.00995	22	5	G	C	T	G	A		G	C	C
h	0.0045	10	7	G	C	T	G	A	C	T	C	C
i	0.0027	6	4	T	C	T	G	A	C	T	C	C
j	0.0018	4	3	G	C	T	G	A	C	T	C	C
s	0.0018	4	4	T	C	T	G	A	C	T	C	C
t	0.00135	3	2	G	C	G	G	A		T	C	C
k	0.0009	2	2	G	C	T	G	A	C	T	T	C
r	0.0009	2	1	G	C	G	G	A	C	T	C	C
i	0.0009	2	2	G	C	G	G	A		T	C	C
u	0.00045	1	1	G	C	T		A		T	C	C
v	0.00045	1	1	G		T	G	A		T	C	C
q	0.00045	1	1	G	C	T	G	A		T	C	C
n	0.00045	1	1	G	C	T	G	A		T	C	G
m	0.00045	1	1	G	C	T	G	A		T	C	G
p	0.00045	1	1	G	C	T	G	G		T	C	C
o	0.00045	1	1	G	C	T	G	A		T	C	C

Table 13.128 - IGLV3-21

h.type	cum.fre	# indvs	pops	22:23055533	22:23055539	22:23055543	22:23055611	22:23055674
				rs4446155	rs1985791	rs188081944	rs192901531	rs189479110
ref	0.2714	-	-	Val->Ile	Asp->Tyr	Asp->Gly	Thr->Ser	Ser->Cys
a	0.31195	556	14	G	G	A	A	A
b	0.178	338	14		T	A	A	A
c	0.1613	296	14	G	G	A	A	A
f	0.02165	45	9		T	A	A	A
d	0.0173	36	10		G	A	A	A
g	0.01045	21	5	G	G	A	A	A
e	0.00995	22	5		T	A	A	A
h	0.0045	10	7		T	A	A	A
i	0.0027	6	4	G	G	A	A	A
j	0.0018	4	3		G	A	A	A
s	0.0018	4	4		T	A	A	A
t	0.00135	3	2		G	A	A	A
k	0.0009	2	2	G	G	A	A	A
r	0.0009	2	1		G	A	A	A
i	0.0009	2	2		T	A	A	A
u	0.00045	1	1	G	G	A	A	A
v	0.00045	1	1		T	A	A	A
q	0.00045	1	1	G	G	A	T	A
n	0.00045	1	1	G	G	A	A	A
m	0.00045	1	1		T	G	A	A
p	0.00045	1	1		T	A	A	A
o	0.00045	1	1	G	G	A	A	T

ENST000000390308

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

1,2,5,7,8,10,12,13,14

1,3,4,6,7,9,10,12,13,14

1,10,11,12,14

2,4,7,11,13

2,4,6,9,10,13,14

2,5,11,14

1,2,10

2,6,9,13

1,14

7,12

12

1,10

2

13

4

6

6

13

13

Table 13.129 - IGLV3-22

		h.type		cum.freq		# indivs	pops										
ref		0.6639	-					22:23046829	22:23047053	22:23047069	22:23047094	22:23047104	22:23047177	22:23047198	22:23047244	22:23047290	
a	14	0.2962	529	C	G			C			T	G	G	G	C	G	
b	7	0.01545	33	C	T			C			T		G	G	C		
c	5	0.00825	18	C	G	C		C			T	G	G	G	C		
e	1	0.0027	6	C	T			C			T		G	G	C	G	
d	3	0.0027	6	C	G	C		C			T		G	G	C	G	
g	3	0.0018	4	C	G			C			T		G	G	C	G	
h	2	0.0018	4	T	G	C		C			T	G	G	G	C	G	
f	3	0.00135	3	C	G	C		C			T		G	G	C		
i	1	0.00135	3	C	T			C			T	G	G		C		
j	1	0.0009	2	C	G	C		C			G	G	G	G	C	G	
n	2	0.0009	2	C	G	C		C			T	G	G		C	G	
m	2	0.0009	2	C	T			C			T	G	G		C	G	
o	2	0.0009	2	C	T			C			T	G	G	G	C	G	
k	1	0.00045	1	C	G			C			T		G	G	G		
l	1	0.00045	1	C	G			C			T			G	C		

Table 13.129 - IGLV3-22

h.type	cum.freq	# indivs	pops	
ref	0.6639	-	-	ENST00000390307
a	0.2962	529	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.01545	33	7	1,3,4,9,10,12,14
c	0.00825	18	5	3,4,6,9,14
e	0.0027	6	1	10
d	0.0027	6	3	1,10,14
g	0.0018	4	3	3,4,10
h	0.0018	4	2	10,14
f	0.00135	3	3	2,10,12
i	0.00135	3	1	10
j	0.0009	2	1	4
n	0.0009	2	2	10,12
m	0.0009	2	2	1,14
o	0.0009	2	2	10,14
k	0.00045	1	1	11
l	0.00045	1	1	14

Table 13.130 - IGLV3-25

h.type	cum.freq	# indvs	pops	22:23029258	22:23029263	22:23029478	22:23029526	22:23029537	22:23029586	22:23029588	22:23029595	22:23029597
ref	0.1086	-	-	rs12628782	rs443102	rs190375718	rs3205084	rs182727015	rs188154997	rs5751507	rs182651541	rs188941516
a	0.6581	943	14	Pro->His	Phe->Leu	Pro->His	Gly->Ala	Pro->Ser	Val->Glu	Leu->Val	Ile->Lys	Tyr->His
b	0.19965	367	14	C	C	C	C	C	T	C	T	T
c	0.02105	41	6	C	C	C	C	C	T	C	T	T
d	0.00405	9	5	C	C	C	C	C	T	G	T	T
e	0.00135	3	3	C	C	C	C	C	T	C	T	T
g	0.0009	2	1	C	C	■	G	C	T	C	T	T
l	0.0009	2	2	C	C	C	G	C	T	C	T	C
j	0.00045	1	1	C	C	C	G	C	T	C	T	T
k	0.00045	1	1	C	C	C	C	C	T	C	T	C
s	0.00045	1	1	C	T	C	G	T	T	G	T	T
q	0.00045	1	1	C	C	C	G	C	T	G	T	C
r	0.00045	1	1	C	C	C	C	C	T	C	T	T
h	0.00045	1	1	C	C	C	G	C	T	C	T	T
f	0.00045	1	1	■	C	C	G	C	T	C	T	T
i	0.00045	1	1	C	C	C	G	C	T	C	T	T
n	0.00045	1	1	C	C	C	C	C	T	C	T	T
m	0.00045	1	1	C	C	C	C	T	T	C	T	T
p	0.00045	1	1	C	T	C	G	C	■	C	T	T
o	0.00045	1	1	C	C	C	G	C	T	C	■	T

Table 13.130 - IGLV3-25

h.type	cum.freq	# indvs	pops	22:23029598 rs192490624	22:23029607 rs145801212	22:23029658 rs184773494	22:23029682 rs189679814	22:23029690 rs181030260	22:23029719 rs189572281	22:23029723 rs181874336
ref	0.1086 -	-	-	Tyr->Ser	Ser->Ile	Thr->Arg	Gly->Ala	Ala->Ser	Gln->His	Ala->Thr
a	0.6581	943	14	A	G	C	G	G	A	G
b	0.19965	367	14	A	G	C	G	G	A	G
c	0.02105	41	6	A	G	C	G	G	A	G
d	0.00405	9	5	A	G	C	G	G	A	G
e	0.00135	3	3	A	T	C	G	G	A	G
g	0.0009	2	1	A	G	C	G	G	A	G
l	0.0009	2	2	A	G	C	G	G	A	G
j	0.00045	1	1	A	G	G	G	G	A	G
k	0.00045	1	1	A	G	C	G	G	A	G
s	0.00045	1	1	A	G	C	G	G	A	G
q	0.00045	1	1	A	G	C	G	G	C	G
r	0.00045	1	1	A	G	C	G	G	A	■
h	0.00045	1	1	A	G	C	G	T	A	G
f	0.00045	1	1	A	T	C	G	G	A	G
i	0.00045	1	1	A	G	C	C	G	A	G
n	0.00045	1	1	A	G	C	C	G	A	G
m	0.00045	1	1	A	G	C	G	G	A	G
p	0.00045	1	1	C	G	C	G	G	A	G
o	0.00045	1	1	A	G	C	G	G	A	G

Table 13.130 - IGLV3-25

h.type	cum.freq	# indivs	pops	
ref	0.1086	-	-	ENST00000390305
a	0.6581	943	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.19965	367	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.02105	41	6	2,3,4,5,9,11
d	0.00405	9	5	1,2,5,6,10
e	0.00135	3	3	2,5,14
g	0.0009	2	1	2
l	0.0009	2	2	3,5
j	0.00045	1	1	6
k	0.00045	1	1	5
s	0.00045	1	1	10
q	0.00045	1	1	2
r	0.00045	1	1	7
h	0.00045	1	1	10
f	0.00045	1	1	4
i	0.00045	1	1	6
n	0.00045	1	1	14
m	0.00045	1	1	8
p	0.00045	1	1	10
o	0.00045	1	1	10

Table 13.131 - IGLV3-27

h.type	cum.freq	# indivs	pops	22:23011202 rs185761809	22:23011253 rs147724316	
ref	0.99865	-	-	Val->Leu G	Cys->Arg T	ENST000000390304
a	0.0009	2	2	G	C	3,9
b	0.00045	1	1	C	T	8

[illegible]

Table 13.133 - IGLV3-9

h.type	cum.fre	# indivs	pops	22:23162047 rs187387271	22:23162116 rs111894401	ENST00000390316
				Gly->Glu	Ser->Asn	
ref	0.9982	-	-	G	G	
a	0.0009	2	2	G		1,13
b	0.0009	2	2		G	3,13

Table 13.134 - IGLV4-3

h.type	cum.freq	# indivs	pops	22:23213939 rs191969136	22:23213987 rs183930410	22:23214006 rs188425943	22:23214144 rs141025913	22:23214190 rs149834282	ENST000000390318
ref	0.9872	-	-	Leu->Phe	Ser->Arg	Tyr->His	Leu->Phe	Thr->Lys	
a	0.01055	22	4	G	C	T	C	C	
d	0.0009	2	2	G	C	T	C	■	
e	0.00045	1	1	G	C	C	C	C	
c	0.00045	1	1	C	C	T	T	C	
b	0.00045	1	1	G	■	T	C	C	
									1,10,12,14 7,14 2 14 3

Table 13.135 - IGLV4-60

h.type	cum.freq	# indivs	pops	22:22516624	22:22516785	22:22516825	22:22516832	22:22516864	22:22516869	22:22516922	22:22516929	22:22516933
				rs190843139	Leu->Val	Ser->Leu	Lys->Asn	Ser->Asn	Ile->Phe	Lys->Asn	Gly->Arg	Ser->Asn
ref	0.0851	-	-		C	C	G	G	A	G	G	G
a	0.4868	785	14	TCTC	C	C	G	G	A	G	G	G
b	0.3607	628	14	TCTC	C	C	G	G	A	G	G	G
c	0.03615	74	5	TCTC	C	C	G	G	A	G	G	G
d	0.015	32	4	TCTC	C	T	G	G	A	G	G	G
e	0.00635	14	4	TCTC	C	C	G	G	A	G	G	■
f	0.00225	5	4	T	C	C	G	G	A	G	G	G
g	0.0009	2	2	TCTC	C	C	G	G	A	G	G	G
n	0.0009	2	2	TCTC	C	C	G	G	A	G	G	G
o	0.0009	2	2	TCTC	C	C	G	G	A	G	C	G
j	0.00045	1	1	TCTC	C	C	G	G	A	G	G	G
k	0.00045	1	1	TCTC	G	C	G	G	A	G	G	G
t	0.00045	1	1	TCTC	C	C	G	G	A	C	G	G
s	0.00045	1	1	TCTC	C	C	G	G	A	G	G	G
q	0.00045	1	1	TCTC	C	C	G	G	T	G	G	G
r	0.00045	1	1	TCTC	C	C	G	G	A	G	C	G
h	0.00045	1	1	TCTC	C	C	G	■	A	G	G	G
i	0.00045	1	1	TCTC	C	C	G	■	A	G	G	G
m	0.00045	1	1	TCTC	C	T	G	■	A	G	G	G
l	0.00045	1	1	TCTC	C	C	G	G	A	G	G	G
p	0.00045	1	1	TCTC	C	C	C	G	A	G	G	G

Table 13.135 - IGLV4-60

h.type	cum.freq	# indivs	pops	22:22516948	22:22516984	22:22516990	22:22516998	22:22517026	22:22517065	22:22517071
				rs182062284	rs187948661	rs115153265	rs2073453	rs738885	rs192463945	rs183795322
ref	0.0851	-	-	Lys->Thr	Ser->Asn	Gly->Val	Arg->Cys	Phe->Ser	Ser->Asn	Thr->Ile
a	0.4868	785	14	A	G	G	C	T	G	C
b	0.3607	628	14	A	G	G	T	C	G	C
c	0.03615	74	5	A	G	T	C	C	G	C
d	0.015	32	4	A	G	G	C	C	G	C
e	0.00635	14	4	A	G	G	C	C	G	C
f	0.00225	5	4	A	G	G	C	C	G	C
g	0.0009	2	2	A	G	G	C	C	G	T
n	0.0009	2	2	A		G	C	C	G	C
o	0.0009	2	2	A	G	G	C	C	G	C
j	0.00045	1	1	A	G	G	C	C		C
k	0.00045	1	1	A	G	G	C	C	G	C
t	0.00045	1	1	A	G	G	T	C	G	C
s	0.00045	1	1	C	G	G	T	C	G	C
q	0.00045	1	1	A	G	G	T	C	G	C
r	0.00045	1	1	A	G	G	T	C	G	C
h	0.00045	1	1	A	G	G	C	C	G	C
i	0.00045	1	1	A	G	G	T	C	G	C
m	0.00045	1	1	A	G	G	C	C	G	C
l	0.00045	1	1	A	G	G	C	T	G	T
p	0.00045	1	1	A	G	G	C	C	G	C

Table 13.135 - IGLV4-60

h.type	cum.frec	# indivs	pops	
ref	0.0851	-	-	ENST000000390284
a	0.4868	785	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
b	0.3607	628	14	1,2,3,4,5,6,7,8,9,10,11,12,13,14
c	0.03615	74	5	1,5,10,12,14
d	0.015	32	4	1,5,10,14
e	0.00635	14	4	1,10,11,14
f	0.00225	5	4	1,3,10,14
g	0.0009	2	2	5,11
n	0.0009	2	2	2,9
o	0.0009	2	2	5,10
j	0.00045	1	1	10
k	0.00045	1	1	5
t	0.00045	1	1	6
s	0.00045	1	1	13
q	0.00045	1	1	13
r	0.00045	1	1	13
h	0.00045	1	1	14
i	0.00045	1	1	10
m	0.00045	1	1	10
l	0.00045	1	1	5
p	0.00045	1	1	10

Table 13.136 - IGLV4-69

h.type	cum.freq	# indvs	pops	22:22385618	22:22385671	22:22385747	22:22385851	
ref	0.98945	-	-	rs150132122	rs185429538	rs140354453	rs188851121	
a	0.0092	20	4	Ala->Asp	Ala->Pro	Gly->Glu	Thr->Pro	ENST00000390282
d	0.00045	1	1	C	G		A	
c	0.00045	1	1	C	C	G	A	1,8,10,14
b	0.00045	1	1		G	G	C	5
							A	10
							A	2

Table 13.137 - IGLV5-37

h.type	cum.frec	# indivs	pops	22:22782150	22:22782240	22:22782258	
				rs56060925	rs181085584	rs111480336	
				Gly->Ser	Gly->Cys	Arg->Cys	ENST00000390300
ref	0.86435	-	-	G	G	C	
a	0.1266	251	13		G	C	1,2,3,4,5,6,7,8,9,10,11,12,13
b	0.0068	13	7	G	G	T	2,6,10,11,12,13,14
c	0.00225	4	2	G	T	C	10,14

Table 13.138 - IGLV5-45

h.type	cum.freq	# indvs	pops	22:22730582	22:22730584	22:22730635	22:22730636	22:22730663	22:22730725	22:22730732	22:22730786
ref	0.8017	-	-	rs12157664	rs1985918	rs61731372	rs149965450	rs191752433	rs192713801	rs117029072	rs147644645
				Pro->Leu	Ser->Ala	Arg->Cys	Arg->His	Arg->Lys	Asp->His	Asp->Gly	Ser->Leu
a	0.1706	312	12	C	T	C	G	G	G	A	C
b	0.0164	36	14	C	T	T	G	G	G	A	C
c	0.00545	12	2	T	G	C	G	G	G	A	C
d	0.0036	8	3	C	T	C	G	G	G	A	T
g	0.0009	2	2	C	T	C		G	G	A	C
e	0.00045	1	1	C	T	C	G		G	G	C
h	0.00045	1	1	C	G	C	G	G	C	A	C
f	0.00045	1	1	C	T	T	G		G	A	C

ENST000000390296

1,2,3,5,6,7,8,10,11,12,13,14

1,2,3,4,5,6,7,8,9,10,11,12,13,14

10,14

3,4,9

2,13

2

10

9

Table 13.139 - IGLV5-48

h.type	cum.freq	# indivs	pops	22:22707473	22:22707555	22:22707576	22:22707621	22:22707698	22:22707713	22:22707773
				rs74549497	rs79612806	rs117042445	rs118041409	rs191708293	rs73880637	rs138488444
ref	0.79525	-	-	Pro->Ala	Asn->Ser	Phe->Tyr	Ser->Asn	Asn->Tyr	Val->Phe	Ser->Gly
a	0.0873	174	14	C	A	T	G	A	G	A
c	0.0301	66	13	G	G			A	T	A
b	0.02875	63	14	G	G	T	G	A	G	A
d	0.0123	27	10	G	G		G	A	T	A
e	0.0082	18	9	C	G		G	A	T	A
i	0.00595	13	7	C	G			A	T	A
f	0.0055	12	9	G	G	T		A	T	A
h	0.00545	12	8	C	A	T	G	A	T	A
g	0.0041	9	5	G	G			A	G	A
o	0.00315	7	6	C	A		G	A	T	A
k	0.00225	5	3	C	G	T	G	A	G	A
j	0.0018	4	3	C	A		G	A	G	A
t	0.00135	3	2	G	A	T	G	A	T	A
s	0.00135	3	3	G	A	T	G	A	G	A
u	0.0009	2	2	C	A			A	T	A
q	0.0009	2	2	C	A	T		A	G	A
r	0.0009	2	2	G	G		G	T	T	A
n	0.0009	2	2	C	G	T	G	A	T	A
m	0.0009	2	1	G	A	T		A	T	A
l	0.0009	2	2	C	G	T		A	T	A
v	0.00045	1	1	C	G		G	A	G	A
w	0.00045	1	1	G	G	T		A	G	A
x	0.00045	1	1	G	G		G	A	G	A
p	0.00045	1	1	C	A	T	G	A	G	G

ENST000000390293

1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,4,5,6,9,10,12,13,14
2,4,5,6,7,10,11,13,14
2,3,4,7,9,11,13
2,3,4,6,7,9,10,11,14
1,2,3,6,10,12,13,14
3,6,10,12,13
1,2,7,8,11,13
2,6,13
4,10,12
3,4
2,5,13
6,9
5,12
5,11
2,10
6
12,13
10
2
10
3

Table 13.140 - IGLV5-52

h.type	cum.freq	# indvs	pops	22:22673101	22:22673305	22:22673463	22:22673551
				rs142368630	rs150977964	rs187205544	
				Ser->Phe	Arg->Cys	Cys->Tyr	ENST00000390289
ref	0.9914	-	-	TCTC	C	C	G
a	0.005	11	5	T	C	C	G
d	0.00135	3	2	TCTC	C	G	5,7,8,10,13
c	0.00135	3	1	TCTC	T	G	10,13
b	0.0009	2	1	TCTC	C	G	9
							6

Table 13.141 - IGLV6-57

h.type	cum.freq	# indvs	pops	22:22550370	22:22550381	22:22550387	22:22550399	22:22550405	22:22550415	22:22550436	22:22550450
				rs181723895	rs186024609	rs191529840	rs183090103	rs186363981	rs190816426	rs149903410	rs2073447
ref	0.2701	-	-	Gly->AlaGly->Ala	Asn->Asp	Met->Leu	Pro->Ser	Ser->Ala	Glu->Ala	Thr->Ser	Arg->Gly
a	0.3734	650	14	G	A	A	C	T	A	C	C
b	0.2802	500	14	G	A	A	C	T	A	C	C
c	0.0325	71	12	G	A	A	T	T	A	C	G
d	0.02015	44	11	G	A	A	T	T	A	C	G
e	0.0187	41	12	G	A	A	T	T	A	C	C
j	0.00045	1	1	G	A	A	T	T	A	C	C
k	0.00045	1	1	G	G	A	T	T	A	C	C
g	0.00045	1	1	G	A	T	C	T	A	C	C
h	0.00045	1	1	C	A	A	C	T	A	C	C
f	0.00045	1	1	G	A	A	C	T	A	C	C
i	0.00045	1	1	G	A	A	C	T	A	C	C
n	0.00045	1	1	G	A	A	C	T	A	C	C
m	0.00045	1	1	G	A	A	C	T	A	C	C
l	0.00045	1	1	G	A	A	C	T	A	G	C
p	0.00045	1	1	G	A	A	C	G	C	C	C
o	0.00045	1	1	G	A	A	C	T	A	C	C

Table 13.141 - IGLV6-57

h.type	cum.freq	# indvs	pops	22:22550466						22:22550510						22:22550525						22:22550565					
				22:22550490						22:22550510						22:22550525						22:22550565					
ref				rs144967736	rs182196598	rs2073448	rs187919298	rs192435787	rs188470766																		
a	0.2701	650	-	Ile->Asn	Tyr->Phe	Ser->Ala	Ile->Leu	Arg->Gln	Ser->Asn																		
b	0.3734	500		T	A	T	A	G	G																		
c	0.2802	71		T	A	G	A	G	G																		
d	0.0325	44		T	A	G	A	G	G																		
e	0.02015	41		T	A	T	A	G	G																		
j	0.0187	1		T	A	T	A	G	G																		
k	0.00045	1		T	A	T	A	G	G																		
g	0.00045	1		T	A	G	A	G	G																		
h	0.00045	1		T	A	T	A	G	G																		
f	0.00045	1		T	T	G	C	G	G																		
i	0.00045	1		T	A	T	C	G	G																		
n	0.00045	1		T	A	T	A	G	G																		
m	0.00045	1		T	A	G	A	G	G																		
l	0.00045	1		T	A	T	A	G	G																		
p	0.00045	1		T	T	T	A	G	G																		
o	0.00045	1		T	A	T	A	G	G																		

ENST000000390285

1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,8,9,10,11,12,13,14
1,2,3,4,5,6,7,9,10,11,13,14
1,2,3,4,6,7,9,10,11,13,14
1,2,3,4,5,6,7,9,10,11,13,14

Table 13.142 - IGLV7-43

h.type	cum.freq	# indvs	pops	22:22749426	22:22749555	22:22749576	22:22749580	22:22749591	22:22749676	22:22749693	22:22749704
ref	0.97765	-	-								
a	0.01005	22		T	C	G	G	C	C	A	C
b	0.0087	19		T	C		G	C	C	A	C
c	0.00135	3		T	C	G	G	C	T	A	C
d	0.00045	1		T	C	G	G	C	C	A	G
g	0.00045	1		T		G	G	C	C	A	C
e	0.00045	1		T	C	G	G	G	C	G	C
h	0.00045	1		C	C	G	G	C	C	A	C
f	0.00045	1		T	C	G		C	C	A	C

ENST000000390298

2,3,4,6,7,9,10,11,14

1,5,10,14

4,13

1

6

1

1

1

Table 13.143 - IGLV7-46

h.type	cum.freq	# indvs	pops	22:22724244	22:22724306	22:22724313	22:22724332	22:22724394	22:22724424
ref	0.0807	-	-	Ala->Gly	Leu->Met	Tyr->Cys	His->Gln	Leu->Ser	Tyr->Phe
a	0.9166	1080	14	C	C	A	C	T	A
c	0.0009	2	2	G	C	A	C	C	A
d	0.00045	1	1	C	C	G	C	C	A
e	0.00045	1	1	C	C	A	C	C	T
b	0.00045	1	1	C	■	A	C	C	A
f	0.00045	1	1	C	C	A	G	T	A

ENST000000390295

1,2,3,4,5,6,7,8,9,10,11,12,13,14,5,10,10,9,13,2

[illegible]

Table 13.145 - IGLV9-49

h.type	cum.fre	# indivs	pops	22:22697585	22:22697782	22:22697788	22:22697825	22:22697923	22:22697939	22:22697983	22:22697998	22:22698031
ref	0.98455	-	-	Ala->Pro	Thr->Ala	Pro->Ser	Thr->Lys	Thr->Ala	Gly->Glu	Ser->Pro	Tyr->His	Glu->Lys
a	0.00865	18	3	G	A	C	C	A	G	T	T	G
b	0.0032	7	2	G	G	C	C	A	G	T	T	G
c	0.0009	2	2	G	A	C	C	A	G	T	T	■
d	0.00045	1	1	G	A	T	C	A	G	T	T	G
g	0.00045	1	1	C	A	C	■	A	G	T	T	G
e	0.00045	1	1	G	A	C	C	A	G	C	T	G
h	0.00045	1	1	G	A	C	C	G	G	T	T	G
f	0.00045	1	1	G	A	C	C	A	G	T	C	G
i	0.00045	1	1	G	A	C	C	A	■	T	T	G

Table 13.145 - IGLV9-49

h.type	cum.fre	# indivs	pops	22:22698059	rs187595925	ENST00000427632
ref	0.98455	-	-	Hls->Leu	A	
a	0.00865	18	3	A	A	3,4,9
b	0.0032	7	2	A	A	6,13
c	0.0009	2	2	A	A	7,9
d	0.00045	1	1	A	A	1
g	0.00045	1	1	A	A	3
e	0.00045	1	1	T		13
h	0.00045	1	1	A	A	2
f	0.00045	1	1	A	A	13
i	0.00045	1	1	A	A	2

Table 14A: Variants Having a Cumulative Frequency Greater than 5% or 10%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>CUMULATIVE FREQUENCY >10%</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>				
<u>IGHA</u>	<u>IGHA1-a</u>	<u>0.31385</u>	<u>Yes</u>	<u>14</u>
<u>IGHA</u>	<u>IGHA2-a</u>	<u>0.35655</u>	<u>Yes</u>	<u>14</u>
<u>IGHA</u>	<u>IGHA2-b</u>	<u>0.1647</u>	<u>Yes</u>	<u>14</u>
<u>IGHG</u>	<u>IGHG2-a</u>	<u>0.28675</u>	<u>Yes</u>	<u>12</u>
<u>IGHG</u>	<u>IGHG3-a</u>	<u>0.2813</u>	<u>Yes</u>	<u>14</u>
<u>IGHG</u>	<u>IGHG3-b</u>	<u>0.19955</u>	<u>Yes</u>	<u>11</u>
<u>IGHG</u>	<u>IGHG4-a</u>	<u>0.19825</u>	<u>Yes</u>	<u>12</u>
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>Yes</u>	<u>14</u>
<u>IGHM</u>	<u>IGHM-b</u>	<u>0.15245</u>	<u>Yes</u>	<u>14</u>
<u>Heavy J</u>				
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>Yes</u>	<u>14</u>
<u>Heavy V</u>				
<u>IGHV</u>	<u>IGHV1-2-a</u>	<u>0.3268</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-2-b</u>	<u>0.1304</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-2-c</u>	<u>0.0797</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-3-a</u>	<u>0.3564</u>	<u>Yes</u>	<u>14</u>

<u>IGHV</u>	<u>IGHV1-3-b</u>	<u>0.0724</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV1-45-a</u>	<u>0.1286</u>	<u>Yes</u>	<u>12</u>
<u>IGHV</u>	<u>IGHV1-45-b</u>	<u>0.1126</u>	<u>Yes</u>	<u>12</u>
<u>IGHV</u>	<u>IGHV1-58-a</u>	<u>0.4503</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-58-b</u>	<u>0.3144</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-69-a</u>	<u>0.1689</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-69-b</u>	<u>0.1543</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-69-c</u>	<u>0.0561</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV2-26-a</u>	<u>0.06135</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-70-a</u>	<u>0.31</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-70-b</u>	<u>0.16595</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-70-d</u>	<u>0.06035</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-13-a</u>	<u>0.15245</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-15-a</u>	<u>0.0572</u>	<u>No</u>	<u>11</u>
<u>IGHV</u>	<u>IGHV3-16-a</u>	<u>0.11075</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-20-a</u>	<u>0.2829</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-20-b</u>	<u>0.1209</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-23-a</u>	<u>0.09825</u>	<u>No</u>	<u>12</u>
<u>IGHV</u>	<u>IGHV3-33-a</u>	<u>0.145</u>	<u>Yes</u>	<u>12</u>
<u>IGHV</u>	<u>IGHV3-35-a</u>	<u>0.11765</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-35-b</u>	<u>0.09695</u>	<u>No</u>	<u>10</u>
<u>IGHV</u>	<u>IGHV3-38-a</u>	<u>0.0723</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-43-a</u>	<u>0.2428</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-43-b</u>	<u>0.08365</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-48-a</u>	<u>0.3355</u>	<u>Yes</u>	<u>14</u>

<u>IGHV</u>	<u>IGHV3-48-b</u>	<u>0.20455</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-49-a</u>	<u>0.38705</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-49-b</u>	<u>0.28555</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-53-a</u>	<u>0.2535</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-53-b</u>	<u>0.2264</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-64-a</u>	<u>0.47965</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-64-b</u>	<u>0.06225</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-66-a</u>	<u>0.27865</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-66-b</u>	<u>0.12575</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-66-c</u>	<u>0.0649</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV4-28-a</u>	<u>0.41435</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-b</u>	<u>0.13715</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-c</u>	<u>0.07385</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-d</u>	<u>0.06485</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV4-31-a</u>	<u>0.5475</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-b</u>	<u>0.2035</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-c</u>	<u>0.1554</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-39-a</u>	<u>0.10295</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-39-b</u>	<u>0.0918</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-4-a</u>	<u>0.1136</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-4-d</u>	<u>0.0817</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-4-f</u>	<u>0.06675</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-4-b</u>	<u>0.06385</u>	<u>No</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV4-4-c</u>	<u>0.0599</u>	<u>No</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-4-e</u>	<u>0.05215</u>	<u>No</u>	<u>12</u>
<u>IGHV</u>	<u>IGHV4-61-a</u>	<u>0.5701</u>	<u>Yes</u>	<u>14</u>

<u>IGHV</u>	<u>IGHV4-61-b</u>	<u>0.18115</u>	<u>Yes</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-61-c</u>	<u>0.13535</u>	<u>Yes</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV4-61-f</u>	<u>0.05855</u>	<u>No</u>	<u>14</u>
<u>Kappa C</u>				
<u>IGKC</u>	<u>IGKC-a</u>	<u>0.2427</u>	<u>Yes</u>	<u>14</u>
<u>Kappa V</u>				
<u>IGKV</u>	<u>IGKV1-17-a</u>	<u>0.08655</u>	<u>No</u>	<u>13</u>
<u>IGKV</u>	<u>IGKV1-17-b</u>	<u>0.0545</u>	<u>No</u>	<u>12</u>
<u>IGKV</u>	<u>IGKV1-8-a</u>	<u>0.20315</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV1-8-b</u>	<u>0.1334</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV1-9-a</u>	<u>0.1854</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV1D-42-a</u>	<u>0.22345</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV1D-8-a</u>	<u>0.1872</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2-24-a</u>	<u>0.1708</u>	<u>Yes</u>	<u>13</u>
<u>IGKV</u>	<u>IGKV2-30-a</u>	<u>0.24085</u>	<u>Yes</u>	<u>13</u>
<u>IGKV</u>	<u>IGKV2D-26-a</u>	<u>0.325</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2D-29-a</u>	<u>0.30175</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3-7-a</u>	<u>0.2018</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3D-11-a</u>	<u>0.35105</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3D-15-a</u>	<u>0.2061</u>	<u>Yes</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV5-2-a</u>	<u>0.32315</u>	<u>Yes</u>	<u>14</u>
<u>Lambda C</u>				
<u>IGLC</u>	<u>IGLC7-a</u>	<u>0.0751</u>	<u>No</u>	<u>14</u>

<u>Lambda J</u>				
<u>IGLJ</u>	<u>IGLJ2-a</u>	<u>0.0741</u>	<u>No</u>	<u>13</u>
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>Yes</u>	<u>14</u>
<u>Lambda V</u>				
<u>IGLV</u>	<u>IGLV10-54-a</u>	<u>0.50725</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV10-54-b</u>	<u>0.09425</u>	<u>No</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV11-55-a</u>	<u>0.4463</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-36-a</u>	<u>0.0942</u>	<u>No</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-50-a</u>	<u>0.19325</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-51-a</u>	<u>0.1177</u>	<u>Yes</u>	<u>11</u>
<u>IGLV</u>	<u>IGLV2-14-a</u>	<u>0.48365</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-14-c</u>	<u>0.08855</u>	<u>No</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-14-b</u>	<u>0.0804</u>	<u>No</u>	<u>10</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-23-a</u>	<u>0.34655</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-8-a</u>	<u>0.1451</u>	<u>Yes</u>	<u>11</u>
<u>IGLV</u>	<u>IGLV3-12-a</u>	<u>0.38715</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-12-b</u>	<u>0.1116</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-12-c</u>	<u>0.0774</u>	<u>No</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-19-a</u>	<u>0.10015</u>	<u>Yes</u>	<u>12</u>
<u>IGLV</u>	<u>IGLV3-21-a</u>	<u>0.31195</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-21-b</u>	<u>0.178</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-21-c</u>	<u>0.1613</u>	<u>Yes</u>	<u>14</u>

<u>IGLV</u>	<u>IGLV3-22-a</u>	<u>0.2962</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-b</u>	<u>0.19965</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-a</u>	<u>0.4868</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-b</u>	<u>0.3607</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV5-37-a</u>	<u>0.1266</u>	<u>Yes</u>	<u>13</u>
<u>IGLV</u>	<u>IGLV5-45-a</u>	<u>0.1706</u>	<u>Yes</u>	<u>12</u>
<u>IGLV</u>	<u>IGLV5-48-a</u>	<u>0.0873</u>	<u>No</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV6-57-a</u>	<u>0.3734</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV6-57-b</u>	<u>0.2802</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>Yes</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV8-61-a</u>	<u>0.2193</u>	<u>Yes</u>	<u>12</u>

Table 14B: Variants Having a Cumulative Frequency Greater than 20%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHA</u>	<u>IGHA1-a</u>	<u>0.31385</u>	<u>14</u>
<u>IGHA</u>	<u>IGHA2-a</u>	<u>0.35655</u>	<u>14</u>
<u>IGHG</u>	<u>IGHG2-a</u>	<u>0.28675</u>	<u>12</u>
<u>IGHG</u>	<u>IGHG3-a</u>	<u>0.2813</u>	<u>14</u>
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Heavy J</u>			
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>14</u>

<u>Heavy V</u>			
<u>IGHV</u>	<u>IGHV1-2-a</u>	<u>0.3268</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-3-a</u>	<u>0.3564</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-58-a</u>	<u>0.4503</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-58-b</u>	<u>0.3144</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-70-a</u>	<u>0.31</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-20-a</u>	<u>0.2829</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-43-a</u>	<u>0.2428</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-48-a</u>	<u>0.3355</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-48-b</u>	<u>0.20455</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-49-a</u>	<u>0.38705</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-49-b</u>	<u>0.28555</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-53-a</u>	<u>0.2535</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-53-b</u>	<u>0.2264</u>	<u>13</u>
<u>IGHV</u>	<u>IGHV3-64-a</u>	<u>0.47965</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-66-a</u>	<u>0.27865</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-a</u>	<u>0.41435</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-a</u>	<u>0.5475</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-b</u>	<u>0.2035</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-61-a</u>	<u>0.5701</u>	<u>14</u>
<u>Kappa C</u>			
<u>IGKC</u>	<u>IGKC-a</u>	<u>0.2427</u>	<u>14</u>
	<u>IGKV1-8-a</u>	<u>0.20315</u>	<u>14</u>

<u>Kappa V</u>			
<u>IGKV</u>			
<u>IGKV</u>	<u>IGKV1D-42-a</u>	<u>0.22345</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2-30-a</u>	<u>0.24085</u>	<u>13</u>
<u>IGKV</u>	<u>IGKV2D-26-a</u>	<u>0.325</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2D-29-a</u>	<u>0.30175</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3-7-a</u>	<u>0.2018</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3D-11-a</u>	<u>0.35105</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3D-15-a</u>	<u>0.2061</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV5-2-a</u>	<u>0.32315</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV10-54-a</u>	<u>0.50725</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV11-55-a</u>	<u>0.4463</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-14-a</u>	<u>0.48365</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-23-a</u>	<u>0.34655</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-12-a</u>	<u>0.38715</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-21-a</u>	<u>0.31195</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-22-a</u>	<u>0.2962</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-a</u>	<u>0.4868</u>	<u>14</u>

<u>IGLV</u>	<u>IGLV4-60-b</u>	<u>0.3607</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV6-57-a</u>	<u>0.3734</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV6-57-b</u>	<u>0.2802</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV8-61-a</u>	<u>0.2193</u>	<u>12</u>

Table 14C: Variants Having a Cumulative Frequency Greater than 30%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHA</u>	<u>IGHA1-a</u>	<u>0.31385</u>	<u>14</u>
<u>IGHA</u>	<u>IGHA2-a</u>	<u>0.35655</u>	<u>14</u>
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Heavy J</u>			
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>14</u>
<u>Heavy V</u>			
<u>IGHV</u>	<u>IGHV1-2-a</u>	<u>0.3268</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-3-a</u>	<u>0.3564</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-58-a</u>	<u>0.4503</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV1-58-b</u>	<u>0.3144</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-70-a</u>	<u>0.31</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-48-a</u>	<u>0.3355</u>	<u>14</u>

<u>IGHV</u>	<u>IGHV3-49-a</u>	<u>0.38705</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-64-a</u>	<u>0.47965</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-a</u>	<u>0.41435</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-a</u>	<u>0.5475</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-61-a</u>	<u>0.5701</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2D-26-a</u>	<u>0.325</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV2D-29-a</u>	<u>0.30175</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV3D-11-a</u>	<u>0.35105</u>	<u>14</u>
<u>IGKV</u>	<u>IGKV5-2-a</u>	<u>0.32315</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV10-54-a</u>	<u>0.50725</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV11-55-a</u>	<u>0.4463</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-14-a</u>	<u>0.48365</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-23-a</u>	<u>0.34655</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-12-a</u>	<u>0.38715</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-21-a</u>	<u>0.31195</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-a</u>	<u>0.4868</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-b</u>	<u>0.3607</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV6-57-a</u>	<u>0.3734</u>	<u>14</u>

<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>
-------------	-------------------	---------------	-----------

Table 14D: Variants Having a Cumulative Frequency Greater than 40%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Heavy J</u>			
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>14</u>
<u>Heavy V</u>			
<u>IGHV</u>	<u>IGHV1-58-a</u>	<u>0.4503</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV3-64-a</u>	<u>0.47965</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-28-a</u>	<u>0.41435</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-a</u>	<u>0.5475</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-61-a</u>	<u>0.5701</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV10-54-a</u>	<u>0.50725</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV11-55-a</u>	<u>0.4463</u>	<u>14</u>

<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-14-a</u>	<u>0.48365</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV4-60-a</u>	<u>0.4868</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>

Table 14E: Variants Having a Cumulative Frequency Greater than 50%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Heavy J</u>			
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>14</u>
<u>Heavy V</u>			
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-31-a</u>	<u>0.5475</u>	<u>14</u>
<u>IGHV</u>	<u>IGHV4-61-a</u>	<u>0.5701</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>	<u>IGLV10-54-a</u>	<u>0.50725</u>	<u>14</u>

<u>IGLV</u>			
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>

Table 14F: Variants Having a Cumulative Frequency Greater than 60%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Heavy J</u>			
<u>IGHJ</u>	<u>IGHJ6-a</u>	<u>0.6845</u>	<u>14</u>
<u>Heavy V</u>			
<u>IGHV</u>	<u>IGHV2-5-a</u>	<u>0.6657</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV3-25-a</u>	<u>0.6581</u>	<u>14</u>

<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>
-------------	-------------------	---------------	-----------

Table 14G: Variants Having a Cumulative Frequency Greater than 70%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Heavy C</u>			
<u>IGHM</u>	<u>IGHM-a</u>	<u>0.7527</u>	<u>14</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV2-18-a</u>	<u>0.763</u>	<u>14</u>
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>

Table 14H: Variants Having a Cumulative Frequency Greater than 80%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV1-47-a</u>	<u>0.8413</u>	<u>14</u>

<u>IGLV</u>			
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>

Table 14l: Variants Having a Cumulative Frequency Greater than 90%

<u>GENE SEGMENT</u>	<u>VARIANT NAME</u>	<u>CUMULATIVE FREQUENCY</u>	<u>NO UNIQUE POPULATIONS</u>
<u>Lambda J</u>			
<u>IGLJ</u>	<u>IGLJ7-a</u>	<u>0.93225</u>	<u>14</u>
<u>Lambda V</u>			
<u>IGLV</u>	<u>IGLV7-46-a</u>	<u>0.9166</u>	<u>14</u>

Table 15: NAÏVE HEAVY CHAIN REPERTOIRES

<u>TABLE 15A:</u> <u>All Naïve – J</u> <u>Usage</u>			
	<u>Average HCDR3 Length</u>	<u>Count</u>	<u>%</u>
<u>J</u>			
IGHJ1*01	11	12	0.90%
IGHJ2*01	12	47	3.51%
IGHJ3*02	12	205	15.30%
IGHJ4*02	11	689	51.42%
IGHJ5*02	13	88	6.57%
<u>IGHJ6*02</u>	<u>16</u>	<u>299</u>	<u>22.31%</u>

<u>Naïve HCDR3>=20</u>			
	<u>Average HCDR3 Length</u>	<u>Count</u>	<u>%</u>
<u>J</u>			

IGHJ3*02	20	1	2.22%
IGHJ4*02	20	1	2.22%
IGHJ5*02	20	2	4.44%
IGHJ6*02	21	41	91.11%

Naïve – Long HCDR3 using IGHJ6*02

<u>HCDR3Length</u>	<u>Count</u>
20	19
21	10
22	7
23	3
24	1
26	1

<u>TABLE 15B:</u> <u>All Naïve – J & D Usage</u>			
<u>J</u>	<u>D</u>	<u>Average HCDR3 Length</u>	<u>Count</u>
IGHJ6*02	IGHD3-9*01	19	21
IGHJ6*02	IGHD4-17*01	18	7
IGHJ6*02	IGHD3-10*01	17	98
IGHJ6*02	IGHD2-2*02	17	4
IGHJ6*02	IGHD5-24*01	17	1
IGHJ6*02	IGHD6-19*01	16	30
IGHJ6*02	IGHD3-22*01	16	5
IGHJ6*02	IGHD6-13*01	16	33
IGHJ6*02	IGHD5-12*01	15	7
IGHJ6*02	IGHD1-26*01	15	25
IGHJ6*02	IGHD1-20*01	14	6
IGHJ6*02	IGHD5-18*01	14	3
IGHJ6*02	IGHD3-16*02	14	6
IGHJ6*02	IGHD2-21*02	13	5
IGHJ6*02	IGHD1-14*01	13	4
IGHJ6*02	IGHD7-27*02	12	8
IGHJ6*02	IGHD1-1*01	12	8
IGHJ6*02	IGHD6-25*01	12	3
IGHJ6*02	IGHD4-23*01	11	2

Naive HCDR3>=20			
J	D	Average HCDR3 Length	Count
IGHJ6*02	IGHD6-19*01	23	1
IGHJ6*02	IGHD4-17*01	22	2
IGHJ6*02	IGHD3-9*01	21	11
IGHJ6*02	IGHD3-10*01	21	25
IGHJ6*02	IGHD6-13*01	20	1
IGHJ6*02	IGHD3-22*01	20	1

TABLE 15C: All Naïve – V & J Usage			
V	J	Average HCDR3 Length	Count
IGHV3-21*03	IGHJ6*02	20	5
IGHV1-18*01	IGHJ6*02	20	10
IGHV1-2*02	IGHJ6*02	19	5
IGHV3-13*01	IGHJ6*02	19	3
IGHV3-7*01	IGHJ6*02	19	6
IGHV1-8*01	IGHJ6*02	19	21
IGHV7-4-1*01	IGHJ6*02	19	9
IGHV3-9*01	IGHJ6*02	18	13
IGHV4-61*01	IGHJ6*02	17	20
IGHV3-23*04	IGHJ6*02	17	6
IGHV4-4*02	IGHJ6*02	16	34
IGHV1-3*01	IGHJ6*02	16	39
IGHV3-20*d01	IGHJ6*02	16	1
IGHV6-1*01	IGHJ6*02	14	69
IGHV2-5*10	IGHJ6*02	14	8
IGHV3-11*01	IGHJ6*02	14	3
IGHV3-66*03	IGHJ6*02	13	21
IGHV3-15*01	IGHJ6*02	12	3

Naive HCDR3>=20			
-----------------	--	--	--

<u>V</u>	<u>J</u>	<u>Average HCDR3 Length</u>	<u>Count</u>
IGHV3-21*03	IGHJ6*02	23	3
IGHV3-13*01	IGHJ6*02	22	1
IGHV3-7*01	IGHJ6*02	22	3
IGHV6-1*01	IGHJ6*02	21	3
IGHV1-8*01	IGHJ6*02	21	10
IGHV1-2*02	IGHJ6*02	21	3
IGHV7-4-1*01	IGHJ6*02	21	3
IGHV1-3*01	IGHJ6*02	21	4
IGHV1-18*01	IGHJ6*02	21	5
IGHV4-4*02	IGHJ6*02	20	3
IGHV3-9*01	IGHJ6*02	20	2
IGHV3-23*04	IGHJ6*02	20	1

TABLE 16: IMMUNISED HEAVY CHAIN REPERTOIRES

<u>TABLE 16A: All Immunised – J Usage</u>			
<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>	<u>%</u>
IGHJ1*01	11	2	0.78%
IGHJ2*01	14	7	2.73%
IGHJ3*02	15	12	4.69%
IGHJ4*02	15	120	46.88%
IGHJ5*02	15	19	7.42%
<u>IGHJ6*02</u>	<u>17</u>	<u>96</u>	<u>37.50%</u>

<u>HCDR3>20</u>			
<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>	<u>%</u>
IGHJ4*02	22	2	22.22%
IGHJ5*02	25	1	11.11%
<u>IGHJ6*02</u>	<u>21</u>	<u>6</u>	<u>66.67%</u>

Immunised – Long HCDR3 using IGHJ6*02

<u>HCDR3Length</u>	<u>Count</u>
20	4

WO 2013/041844

PCT/GB2012/052296

21	1
24	1

TABLE 16B:
All Immunised – J & D Usage

<u>J</u>	<u>D</u>	<u>HCDR3 Length</u>	<u>Count</u>
IGHJ6*02	IGHD4-17*01	18	1
IGHJ6*02	IGHD1-26*01	18	5
IGHJ6*02	IGHD6-19*01	17	27
IGHJ6*02	IGHD3-10*01	17	42
IGHJ6*02	IGHD6-13*01	16	2
IGHJ6*02	IGHD5-18*01	15	2
IGHJ6*02	IGHD4-23*01	15	5
IGHJ6*02	IGHD5-12*01	14	3
IGHJ6*02	IGHD3-16*02	14	3
IGHJ6*02	IGHD3-22*01	13	3
IGHJ6*02	IGHD3-9*01	12	1
IGHJ6*02	IGHD1-20*01	11	1

<u>HCDR3>20</u>			
<u>J</u>	<u>D</u>	<u>HCDR3 Length</u>	<u>Count</u>
IGHJ6*02	IGHD3-10*01	21	4
IGHJ6*02	IGHD6-19*01	20	1
IGHJ6*02	IGHD1-26*01	20	1

TABLE 16C:
All Immunised – V & J Usage

<u>V</u>	<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>
IGHV3-7*01	IGHJ6*02	18	33
IGHV1-8*01	IGHJ6*02	17	3
IGHV3-9*01	IGHJ6*02	17	22
IGHV3-13*01	IGHJ6*02	17	5
IGHV1-2*02	IGHJ6*02	16	8
IGHV3-11*01	IGHJ6*02	16	3
IGHV4-4*02	IGHJ6*02	15	9
IGHV6-1*01	IGHJ6*02	15	6
IGHV1-3*01	IGHJ6*02	15	6

<u>HCDR3>20</u>			
<u>V</u>	<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>
IGHV3-7*01	IGHJ6*02	21	4
IGHV3-11*01	IGHJ6*02	21	1
IGHV4-4*02	IGHJ6*02	20	1

TABLE 17: ANTIGEN-SPECIFIC HEAVY CHAIN REPERTOIRES

<u>TABLE 17A:</u> <u>All Antigen-Specific – J</u> <u>Usage</u>			
<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>	<u>%</u>
IGHJ1*01	12	2	1.68%
IGHJ3*02	17	4	3.36%
IGHJ4*02	13	64	53.78%
IGHJ5*02	19	6	5.04%
<u>IGHJ6*02</u>	<u>17</u>	<u>43</u>	<u>36.13%</u>
<u>HCDR3>20</u>			
<u>J</u>	<u>HCDR3 Length</u>	<u>Count</u>	<u>%</u>
IGHJ4*02	22	1	9.09%
IGHJ5*02	21	3	27.27%
<u>IGHJ6*02</u>	<u>21</u>	<u>7</u>	<u>63.64%</u>

Immunised – Long HCDR3 using IGHJ6*02

<u>HCDR3Length</u>	<u>Count</u>
20	4
21	1
22	1
24	1

TABLE 17B: All Antigen-Specific – J & D Usage			
<u>J</u>	<u>D</u>	<u>HCDR3</u> <u>Length</u>	<u>Count</u>
IGHJ6*02	IGHD3-9*01	19	6
IGHJ6*02	IGHD3-10*01	19	12
IGHJ6*02	IGHD6-19*01	17	13
IGHJ6*02	IGHD5-12*01	16	2
IGHJ6*02	IGHD2-15*01	16	1
IGHJ6*02	IGHD1-26*01	15	7
IGHJ6*02	IGHD3-16*02	12	1
IGHJ6*02	IGHD5-18*01	11	1
<u>HCDR3>20</u>			
<u>J</u>	<u>D</u>	<u>HCDR3</u> <u>Length</u>	<u>Count</u>
IGHJ6*02	IGHD3-9*01	21	1
IGHJ6*02	IGHD3-10*01	21	6

TABLE 17C: All Antigen-Specific – V & J Usage			
<u>V</u>	<u>J</u>	<u>HCDR3</u> <u>Length</u>	<u>Count</u>
IGHV1-3*01	IGHJ6*02	18	1
IGHV3-7*01	IGHJ6*02	18	6
IGHV3-9*01	IGHJ6*02	18	6
IGHV1-8*01	IGHJ6*02	17	6
IGHV3-20*d01	IGHJ6*02	17	3
IGHV3-11*01	IGHJ6*02	17	8
IGHV4-4*02	IGHJ6*02	17	5
IGHV3-15*01	IGHJ6*02	15	5
IGHV3-13*01	IGHJ6*02	15	3
<u>HCDR3>20</u>			
<u>V</u>	<u>J</u>	<u>HCDR3</u> <u>Length</u>	<u>Count</u>
IGHV4-4*02	IGHJ6*02	20	2

IGHV1-8*01	IGHJ6*02	22	2
IGHV3-9*01	IGHJ6*02	20	1
IGHV3-11*01	IGHJ6*02	21	1
IGHV3-20*d01	IGHJ6*02	22	1

Table 18: Sequence Correlation Table

SEQ ID NO:	DESCRIPTION:
1	JH5 reference (nucleotide sequence)
2	JH6*03=JH6 reference (nucleotide sequence)
3	JH2 reference (nucleotide sequence)
4	JH6*02 (nucleotide sequence)
5	JH5 reference (nucleotide sequence)
6	JH5 reference (amino acid sequence)
7	JH6*03=JH6 reference (nucleotide sequence)
8	JH6*03=JH6 reference (amino acid sequence)

9	JH2 reference (nucleotide sequence)
10	JH2 reference (amino acid sequence)
11	IGHV1-2*01
12	IGHV1-3*01
13	IGHV1-8*01
14	IGHV1-24*01
15	IGHV1-45*01
16	IGHV1-46*01
17	IGHV1-58*01
18	IGHV1-69*01
19	IGHV1-c*01

WO 2013/041844

PCT/GB2012/052296

20	IGHV1-f*01
21	IGHV2-5*01
22	IGHV2-26*01
23	IGHV2-70*01
24	IGHV3-7*01
25	IGHV3-9*01
26	IGHV3-11*01
27	IGHV3-13*01
28	IGHV3-15*01
29	IGHV3-16*01
30	IGHV3-19*01
31	IGHV3-20*01
32	IGHV3-21*01
33	IGHV3-23*01
34	IGHV3-30*01

35	IGHV3-30-3*01
36	IGHV3-33*01
37	IGHV3-35*01
38	IGHV3-38*01
39	IGHV3-43*01
40	IGHV3-47*01
41	IGHV3-48*01
42	IGHV3-49*01
43	IGHV3-53*01
44	IGHV3-64*01
45	IGHV3-66*01
46	IGHV3-72*01
47	IGHV3-73*01
48	IGHV3-74*01
49	IGHV3-d*01

WO 2013/041844

PCT/GB2012/052296

50	IGHV3-h*01
51	IGHV4-4*01
52	IGHV4-28*01
53	IGHV4-30-2*01
54	IGHV4-30-4*01
55	IGHV4-31*01
56	IGHV4-34*01
57	IGHV4-39*01
58	IGHV4-55*01
59	IGHV4-59*01
60	IGHV4-61*01
61	IGHV4-b*01
62	IGHV5-51*01
63	IGHV5-a*01

64	IGHV6-1*01
65	IGHV7-4-1*01
66	IGHV7-81*01
67	IGHD1-1*01
68	IGHD1-7*01
69	IGHD1-14*01
70	IGHD1-20*01
71	IGHD1-26*01
72	IGHD2-2*01
73	IGHD2-8*01
74	IGHD2-15*01
75	IGHD2-21*01
76	IGHD3-3*01
77	IGHD3-9*01
78	IGHD3-10*01
79	IGHD3-16*01
80	IGHD3-22*01
81	IGHD4-4*01
82	IGHD4-11*01
83	IGHD4-17*01
84	IGHD4-23*01
85	IGHD5-5*01
86	IGHD5-12*01

WO 2013/041844

PCT/GB2012/052296

87	IGHD5-18*01
88	IGHD5-24*01
89	IGHD6-6*01
90	IGHD6-13*01
91	IGHD6-19*01
92	IGHD6-25*01
93	IGHD6-25*01
94	JH1*01
95	JH2*01
96	JH3*01
97	JH4*01
98	JH5*01
99	JH6*01
100	IGKV1-5*01
101	IGKV1-6*01
102	IGKV1-8*01
103	IGKV1D-8*01
104	IGKV1-9*01
105	IGKV1-12*01
106	IGKV1-13*01

107	IGKV1-16*01
108	IGKV1D-16*01
109	IGKV1-17*01
110	IGKV1D-17*01
111	IGKV1-27*01
112	IGKV1-33*01
113	IGKV1D-33*01
114	IGKV1-37*01
115	IGKV1D-37*01
116	IGKV1-39*01
117	IGKV1D-39
118	IGKV1D-42*01
119	IGKV1D-43*01
120	IGKV2-24*01
121	IGKV2D-24*01

WO 2013/041844

PCT/GB2012/052296

122	IGKV2-28*01
123	IGKV2D-28*01
124	IGKV2-29*01
125	IGKV2D-29*01
126	IGKV2-30*01
127	IGKV2D-30*01
128	IGKV2-40*01
129	IGKV2D-40*01
130	IGKV3-7*01
131	IGKV3-7*03
132	IGKV3D-7*01
133	IGKV3-11*01
134	IGKV3D-11*01
135	IGKV3-15*01
136	IGKV3D-15*01

137	IGKV3-20*01
138	IGKV3D-20*01
139	IGKV4-1*01
140	IGKV5-2*01
141	IGKV6-21*01
142	IGKV6D-21*01
143	IGKV6D-41*01
144	IGLV1-36*01
145	IGLV1-40*01
146	IGLV1-41*01
147	IGLV1-44*01
148	IGLV1-47*01
149	IGLV1-50*01
150	IGLV1-51*01
151	IGLV2-8*01

WO 2013/041844

PCT/GB2012/052296

152	IGLV2-11*01
153	IGLV2-14*01
154	IGLV2-18*01
155	IGLV2-23*01
156	IGLV2-33*01
157	IGLV3-1*01
158	IGLV3-10*01
159	IGLV3-12*01
160	IGLV3-16*01
161	IGLV3-19*01
162	IGLV3-21*01
163	IGLV3-22*01
164	IGLV3-25*01
165	IGLV3-27*01
166	IGLV3-32*01

167	IGLV4-3*01
168	IGLV4-60*01
169	IGLV4-69*01
170	IGLV5-37*01
171	IGLV5-45*01
172	IGLV5-48*01
173	IGLV5-52*01
174	IGLV6-57*01
175	IGLV7-43*01
176	IGLV7-46*01
177	IGLV8-61*01
178	IGLV9-49*01
179	IGLV10-54*01

WO 2013/041844

PCT/GB2012/052296

180	IGLV11-55*01
181	IGKJ1*01
182	IGKJ2*01
183	IGKJ3*01
184	IGKJ4*01
185	IGKJ5*01
186	IGLJ1*01
187	IGLJ2*01
188	IGLJ3*01
189	IGLJ4*01
190	IGLJ5*01
191	IGLJ6*01
192	IGLJ7*01
193	IGKV2D-26*01
194	IGLV1-36*01
195	IGLV1-47*01
196	IGLV1-50*01
197	IGLV1-51*01
198	IGLV10-54*01

199	IGLV11-55*01
200	IGLV2-14*01
201	IGLV2-18*01
202	IGLV2-23*01
203	IGLV2-8*01
204	IGLV3-1*01
205	IGLV3-12*01
206	IGLV3-19*01
207	IGLV3-21*01
208	IGLV3-22*01
209	IGLV3-25*01
210	IGLV4-60*01
211	IGLV5-37*01
212	IGLV5-45*01

WO 2013/041844

PCT/GB2012/052296

213	IGLV5-48*01
214	IGLV6-57*01
215	IGLV7-46*01
216	IGLV8-61*01
217	IGHG3
218	IGHA2
219	IGHG2

220	IGHE
221	IGHE
222	IGHA1
223	IGHG1
224	IGHG1

225	IGHM
226	IGHG1
227	IGHD

228	IGHG4
229	IGHE
230	IGKC
231	IGLC2
232	IGLC1
233	IGLC3
234	IGLC7
235	JH6*02 (nucleotide sequence)
236	JH6*01 & JH6*02(amino acid sequence)
237	Human RSS-JH6*02
238	Human RSS
239	IGHJ1 ref

WO 2013/041844

PCT/GB2012/052296

240	IGHJ2 ref
241	IGHJ3 ref
242	IGHJ4 ref
243	IGHJ5 ref
244	IGHJ6 ref
245	IGHV1-18 ref
246	IGHV1-2 ref
247	IGHV1-24 ref
248	IGHV1-3 ref
249	IGHV1-45 ref
250	IGHV1-46 ref
251	IGHV1-58 ref
252	IGHV1-69 ref
253	IGHV1-8 ref
254	IGHV2-26 ref
255	IGHV2-5 ref
256	IGHV2-70 ref
257	IGHV3-11 ref
258	IGHV3-13 ref
259	IGHV3-15 ref

260	IGHV3-16 ref
261	IGHV3-20 ref
262	IGHV3-21 ref
263	IGHV3-23 ref
264	IGHV3-30 ref
265	IGHV3-33 ref
266	IGHV3-35 ref
267	IGHV3-38 ref
268	IGHV3-43 ref
269	IGHV3-48 ref
270	IGHV3-49 ref
271	IGHV3-53 ref
272	IGHV3-64 ref
273	IGHV3-66 ref
274	IGHV3-7 ref

WO 2013/041844

PCT/GB2012/052296

275	IGHV3-72 ref
276	IGHV3-73 ref
277	IGHV3-74 ref
278	IGHV3-9 ref
279	IGHV4-28 ref
280	IGHV4-31 ref
281	IGHV4-34 ref
282	IGHV4-39 ref
283	IGHV4-4 ref
284	IGHV4-59 ref
285	IGHV4-61 ref
286	IGHV5-51 ref
287	IGHV6-1 ref
288	IGHV7-81 ref
289	IGKJ2 ref
290	IGKJ3 ref
291	IGKJ4 ref

292	IGKJ5 ref
293	IGKV1-16 ref
294	IGKV1-17 ref
295	IGKV1-5 ref
296	IGKV1-6 ref
297	IGKV1-8 ref
298	IGKV1-9 ref
299	IGKV1D-12 ref
300	IGKV1D-16 ref
301	IGKV1D-17 ref
302	IGKV1D-42 ref
303	IGKV1D-43 ref
304	IGKV1D-8 ref
305	IGKV2-24 ref
306	IGKV2-30 ref
307	IGKV2D-24 ref

WO 2013/041844

PCT/GB2012/052296

308	IGKV2D-26 ref
309	IGKV2D-29 ref
310	IGKV2D-30 ref
311	IGKV3-11 ref
312	IGKV3-20 ref
313	IGKV3-7 ref
314	IGKV3D-11 ref
315	IGKV3D-15 ref
316	IGKV3D-20 ref
317	IGKV4-1 ref
318	IGKV5-2 ref
319	IGKV6-21 ref
320	IGKV6D-21 ref
321	IGKV6D-41 ref
322	IGLJ1 ref

323	IGLJ2 ref
324	IGLJ3 ref
325	IGLJ5 ref
326	IGLJ6 ref
327	IGLJ7 ref
328	IGLV10-54 ref
329	IGLV11-55 ref
330	IGLV1-36 ref
331	IGLV1-40 ref
332	IGLV1-44 ref
333	IGLV1-47 ref
334	IGLV1-50 ref
335	IGLV1-51 ref
336	IGLV2-11 ref
337	IGLV2-14 ref
338	IGLV2-18 ref
339	IGLV2-23 ref
340	IGLV2-33 ref

WO 2013/041844

PCT/GB2012/052296

341	IGLV2-8 ref
342	IGLV3-1 ref
343	IGLV3-10 ref
344	IGLV3-12 ref
345	IGLV3-16 ref
346	IGLV3-19 ref
347	IGLV3-21 ref
348	IGLV3-22 ref
349	IGLV3-25 ref
350	IGLV3-27 ref
351	IGLV3-32 ref
352	IGLV3-9 ref
353	IGLV4-3 ref
354	IGLV4-60 ref

355	IGLV4-69 ref
356	IGLV5-37 ref
357	IGLV5-45 ref
358	IGLV5-48 ref
359	IGLV5-52 ref
360	IGLV6-57 ref
361	IGLV7-43 ref
362	IGLV7-46 ref
363	IGLV8-61 ref
364	IGLV9-49 ref

365	IGHG1 ref GENOMIC
366	IGHG1 ref - CDS (ensembl transcript ID ENST00000390542)
367	IGHG1 ref - CDS (Ensembl transcript ID ENST00000390549)
368	IGHG1 ref - CDS (ensembl transcript ID ENST00000390548)

369	IGHG2 ref - GENOMIC
370	IGHG2 ref - CDS (ensembl transcript ID ENST00000390545)
371	IGHG2-a CDS
372	IGHG2-a GENOMIC
373	IGHG3 ref - GENOMIC
374	IGHG3 ref - CDS (ensembl transcript ID ENST00000390551)
375	IGHG3-a CDS

376	IGHG3-a GENOMIC
377	IGHG3-b CDS
378	IGHG3-b GENOMIC
379	IGHG4 ref - GENOMIC
380	IGHG4 ref - CDS (ensembl transcript ID ENST00000390543)
381	IGHG4-a CDS

382	IGHG4-a GENOMIC
383	IGHA1 ref GENOMIC
384	IGHA1 ref- CDS (ensembl transcript ID ENST00000390547)
385	IGHA1-a GENOMIC
386	IGHA1-a CDS
387	IGHA2 ref GENOMIC
388	IGHA2 ref- CDS (ensembl transcript ID ENST00000390539)

389	IGHA2-a GENOMIC
390	IGHA2-a CDS
391	IGHA2-b GENOMIC
392	IGHA2-b CDS

393	IGHD ref GENOMIC
394	IGHD ref- CDS (ensembl transcript ID ENST00000390556)

395	IGHE ref GENOMIC
396	IGHE ref- CDS (ensembl transcript ID ENST00000390541)
397	IGHE ref- CDS (ensembl transcript ID ENST00000576077)
398	IGHE ref- CDS (ensembl transcript ID ENST00000577108)
399	IGHM ref GENOMIC
400	IGHM ref- CDS (ensembl transcript ID ENST00000390559)

401	IGHM-a GENOMIC
402	IGHM-a CDS
403	IGHM-b GENOMIC
404	IGHM-b CDS
405	IGHD7-27 GENOMIC
406	IGHD2-15 GENOMIC
407	IGHD3-16 GENOMIC
408	IGHD6-6 ref GENOMIC
409	IGHD5-18 ref GENOMIC
410	IGHD2-2 ref GENOMIC
411	IGHD4-11 ref GENOMIC
412	IGHD5-12 ref

	GENOMIC
413	IGHD3-3 ref GENOMIC
414	IGHD2-8 ref GENOMIC
415	IGHD4-4 ref GENOMIC
416	IGHD4-23 ref GENOMIC
417	IGHD1-14 ref GENOMIC
418	IGHD3-10 ref GENOMIC
419	IGHD1-26 ref GENOMIC
420	IGHD3-9 ref GENOMIC
421	IGHD1-1 ref GENOMIC
422	IGHD6-25 ref GENOMIC
423	IGHD5-24 ref GENOMIC
424	IGHD2-21 ref GENOMIC
425	IGHD1-20 ref GENOMIC
426	IGHD6-13 ref GENOMIC
427	IGHD4-17 ref GENOMIC

428	IGHD3-22 ref GENOMIC
429	IGHD5-5 ref GENOMIC
430	IGHD6-19 ref GENOMIC
431	IGHD1-7 ref GENOMIC
432	Rabbit JH6 (amino acid)
433	Rabbit JH6 (nucleotide)
434	Sheep JH6 (amino acid)
435	Sheep JH6 (nucleotide)
436	Bovine JH6 (amino acid)
437	Bovine JH6 (nucleotide)
438	Dog JH3 (amino acid)
439	Dog JH3 (nucleotide)
440	Human JH6*02 (amino acid)
441	Human JH6*02 (nucleotide)
442	Leader Sequence (nucleotide)
443	Leader Sequence (amino acid)
444	PRIMER
445	PRIMER

Table 19

	IgG1	IgG2	IgG3	IgG4
Complement activation				
Classical pathway	+++	+	+++	—
Alternative pathway	—	+	—	—
Fc receptor recognition				
FcγRI	+++	—	+++	++
FcγRIIa, 131R/R	++	—	++	—
FcγRIIa, 131H/H	+	+	++	—
FcγRIIb	++	—	++	+
FcγRIII	+	+/-	+	+/-

Table 20**Table 20A: Variants Found In No More Than 3 Human Populations**

TYPE	VARIANT	C.FREQ	C.FREQ (%)	#INDIVIDUALS	#POPULATIONS
IGKV	IGKV1D-8-b	0.01785	1.79%	36	3
IGHV	IGHV3-11-a	0.01695	1.70%	36	3
IGHV	IGHV5-51-a	0.0164	1.64%	35	3
IGHV	IGHV2-5-b	0.01055	1.06%	18	3
IGLV	IGLV2-14-g	0.0104	1.04%	22	3
IGHV	IGHV7-81-b	0.01005	1.01%	21	3
IGKV	IGKV1D-42-b	0.01005	1.01%	21	3
IGKV	IGKV3-7-b	0.01	1.00%	19	3
IGHV	IGHV1-69-d	0.00995	1.00%	22	3
IGLV	IGLV2-14-e	0.00995	1.00%	19	3
IGLV	IGLV9-49-a	0.00865	0.87%	18	3
IGLV	IGLV1-44-b	0.00775	0.78%	16	2
IGLV	IGLV2-11-a	0.00775	0.78%	17	3
IGKV	IGKV3-11-d	0.0073	0.73%	14	3
IGKV	IGKV1D-43-b	0.00685	0.69%	14	3
IGHV	IGHV3-38-d	0.0064	0.64%	13	3

IGLJ	IGLJ6-a	0.00595	0.60%	13	3
IGHV	IGHV5-51-b	0.0055	0.55%	11	3
IGLV	IGLV5-45-c	0.00545	0.55%	12	2
IGHV	IGHV4-31-i	0.0054	0.54%	10	3
IGKV	IGKV2D-26-b	0.00455	0.46%	9	3
IGLV	IGLV1-36-c	0.00455	0.46%	10	2
IGHV	IGHV1-24-c	0.00415	0.42%	9	3
IGHV	IGHV1-8-c	0.00405	0.41%	8	2
IGLV	IGLV1-47-d	0.00405	0.41%	9	3
IGHV	IGHV3-43-i	0.00365	0.37%	8	3
IGHV	IGHV3-53-h	0.0036	0.36%	8	2
IGLV	IGLV10-54-e	0.0036	0.36%	8	3
IGLV	IGLV1-36-f	0.0036	0.36%	8	2
IGLV	IGLV5-45-d	0.0036	0.36%	8	3
IGHV	IGHV1-46-b	0.0032	0.32%	7	3
IGHV	IGHV2-26-c	0.0032	0.32%	7	2
IGLJ	IGLJ6-b	0.0032	0.32%	7	3
IGLV	IGLV3-32-a	0.0032	0.32%	7	3
IGLV	IGLV9-49-b	0.0032	0.32%	7	2
IGHJ	IGHJ6-b	0.00315	0.32%	7	2
IGHV	IGHV3-16-d	0.00315	0.32%	7	3
IGHV	IGHV4-28-k	0.00315	0.32%	7	3
IGKJ	IGKJ2-d	0.00315	0.32%	6	3
IGLV	IGLV10-54-f	0.00315	0.32%	7	2
IGLV	IGLV2-8-c	0.00315	0.32%	7	3
IGLV	IGLV3-10-c	0.00315	0.32%	6	2
IGLV	IGLV1-40-c	0.00275	0.28%	6	2
IGLV	IGLV1-51-b	0.00275	0.28%	6	3
IGHV	IGHV1-69-m	0.0027	0.27%	6	3
IGHV	IGHV3-33-e	0.0027	0.27%	6	3
IGHV	IGHV3-49-c	0.0027	0.27%	6	3
IGHV	IGHV3-66-g	0.0027	0.27%	6	3
IGHV	IGHV4-4-u	0.0027	0.27%	6	3
IGKV	IGKV1D-42-c	0.0027	0.27%	6	1
IGKV	IGKV1D-43-c	0.0027	0.27%	5	3
IGLV	IGLV10-54-h	0.0027	0.27%	6	2
IGLV	IGLV10-54-i	0.0027	0.27%	6	2
IGLV	IGLV1-36-g	0.0027	0.27%	6	3
IGLV	IGLV2-8-d	0.0027	0.27%	6	3
IGLV	IGLV3-22-e	0.0027	0.27%	6	1
IGLV	IGLV3-22-d	0.0027	0.27%	6	3
IGHJ	IGHJ2-b	0.00225	0.23%	4	3
IGHV	IGHV1-3-u	0.00225	0.23%	5	2
IGHV	IGHV1-3-v	0.00225	0.23%	5	2
IGHV	IGHV1-3-ap	0.00225	0.23%	4	3
IGHV	IGHV1-46-c	0.00225	0.23%	5	3

IGHV	IGHV1-58-f	0.00225	0.23%	5	3
IGHV	IGHV1-69-j	0.00225	0.23%	5	3
IGHV	IGHV3-13-e	0.00225	0.23%	5	3
IGHV	IGHV3-23-f	0.00225	0.23%	5	3
IGHV	IGHV3-23-g	0.00225	0.23%	5	3
IGHV	IGHV4-28-p	0.00225	0.23%	5	3
IGHV	IGHV4-31-j	0.00225	0.23%	5	3
IGKV	IGKV1-17-d	0.00225	0.23%	4	1
IGKV	IGKV3-20-a	0.00225	0.23%	5	3
IGKV	IGKV3-7-c	0.00225	0.23%	5	3
IGKV	IGKV3-7-d	0.00225	0.23%	5	3
IGKV	IGKV3D-11-b	0.00225	0.23%	5	3
IGLJ	IGLJ3-a	0.00225	0.23%	4	2
IGLV	IGLV11-55-g	0.00225	0.23%	5	3
IGLV	IGLV1-50-c	0.00225	0.23%	5	3
IGLV	IGLV2-11-g	0.00225	0.23%	5	3
IGLV	IGLV2-14-n	0.00225	0.23%	5	3
IGLV	IGLV2-23-f	0.00225	0.23%	5	2
IGLV	IGLV3-19-c	0.00225	0.23%	5	2
IGLV	IGLV5-37-c	0.00225	0.23%	4	2
IGLV	IGLV5-48-k	0.00225	0.23%	5	3
IGLV	IGLV2-14-i	0.00185	0.19%	4	2
IGHJ	IGHJ1-a	0.0018	0.18%	3	3
IGHV	IGHV1-2-g	0.0018	0.18%	4	3
IGHV	IGHV1-3-y	0.0018	0.18%	4	3
IGHV	IGHV1-3-ar	0.0018	0.18%	4	3
IGHV	IGHV1-3-ai	0.0018	0.18%	4	3
IGHV	IGHV1-69-s	0.0018	0.18%	4	3
IGHV	IGHV1-8-d	0.0018	0.18%	4	2
IGHV	IGHV2-5-d	0.0018	0.18%	4	2
IGHV	IGHV3-13-j	0.0018	0.18%	4	2
IGHV	IGHV3-21-d	0.0018	0.18%	4	3
IGHV	IGHV3-23-n	0.0018	0.18%	4	3
IGHV	IGHV3-33-d	0.0018	0.18%	4	3
IGHV	IGHV3-64-e	0.0018	0.18%	4	2
IGHV	IGHV3-9-g	0.0018	0.18%	4	3
IGHV	IGHV4-39-y	0.0018	0.18%	4	3
IGHV	IGHV4-39-n	0.0018	0.18%	4	3
IGHV	IGHV4-39-o	0.0018	0.18%	4	3
IGHV	IGHV7-81-c	0.0018	0.18%	4	3
IGKJ	IGKJ2-e	0.0018	0.18%	4	3
IGKV	IGKV2D-26-c	0.0018	0.18%	4	1
IGKV	IGKV3D-11-g	0.0018	0.18%	3	2
IGKV	IGKV3D-11-f	0.0018	0.18%	4	3
IGKV	IGKV4-1-w	0.0018	0.18%	3	3
IGLJ	IGLJ1-a	0.0018	0.18%	4	2

IGLV	IGLV10-54-k	0.0018	0.18%	4	2
IGLV	IGLV11-55-e	0.0018	0.18%	4	3
IGLV	IGLV1-36-e	0.0018	0.18%	4	3
IGLV	IGLV1-40-f	0.0018	0.18%	4	3
IGLV	IGLV1-44-d	0.0018	0.18%	4	3
IGLV	IGLV1-44-e	0.0018	0.18%	4	3
IGLV	IGLV1-47-g	0.0018	0.18%	4	2
IGLV	IGLV1-50-d	0.0018	0.18%	4	2
IGLV	IGLV1-51-d	0.0018	0.18%	4	2
IGLV	IGLV1-51-c	0.0018	0.18%	4	3
IGLV	IGLV2-14-r	0.0018	0.18%	4	1
IGLV	IGLV2-33-b	0.0018	0.18%	4	3
IGLV	IGLV2-8-e	0.0018	0.18%	4	2
IGLV	IGLV3-16-h	0.0018	0.18%	4	3
IGLV	IGLV3-21-j	0.0018	0.18%	4	3
IGLV	IGLV3-22-g	0.0018	0.18%	4	3
IGLV	IGLV3-22-h	0.0018	0.18%	4	2
IGLV	IGLV5-48-j	0.0018	0.18%	4	3
IGHJ	IGHJ4-d	0.00135	0.14%	3	3
IGHV	IGHV1-18-d	0.00135	0.14%	3	3
IGHV	IGHV1-18-e	0.00135	0.14%	3	3
IGHV	IGHV1-2-h	0.00135	0.14%	3	2
IGHV	IGHV1-2-f	0.00135	0.14%	3	3
IGHV	IGHV1-3-aa	0.00135	0.14%	3	1
IGHV	IGHV1-3-ak	0.00135	0.14%	3	3
IGHV	IGHV1-3-w	0.00135	0.14%	3	1
IGHV	IGHV1-3-at	0.00135	0.14%	3	2
IGHV	IGHV1-69-k	0.00135	0.14%	3	1
IGHV	IGHV1-69-ac	0.00135	0.14%	3	3
IGHV	IGHV1-8-e	0.00135	0.14%	3	2
IGHV	IGHV1-8-f	0.00135	0.14%	3	2
IGHV	IGHV2-70-y	0.00135	0.14%	3	3
IGHV	IGHV2-70-z	0.00135	0.14%	3	3
IGHV	IGHV2-70-af	0.00135	0.14%	3	2
IGHV	IGHV2-70-h	0.00135	0.14%	3	3
IGHV	IGHV3-13-g	0.00135	0.14%	3	2
IGHV	IGHV3-16-e	0.00135	0.14%	3	2
IGHV	IGHV3-20-f	0.00135	0.14%	3	3
IGHV	IGHV3-30-d	0.00135	0.14%	3	2
IGHV	IGHV3-30-c	0.00135	0.14%	3	3
IGHV	IGHV3-33-h	0.00135	0.14%	3	3
IGHV	IGHV3-43-k	0.00135	0.14%	3	2
IGHV	IGHV3-43-m	0.00135	0.14%	3	3
IGHV	IGHV3-53-n	0.00135	0.14%	3	2
IGHV	IGHV3-53-m	0.00135	0.14%	3	3
IGHV	IGHV3-64-f	0.00135	0.14%	3	1

IGHV	IGHV3-73-c	0.00135	0.14%	3	3
IGHV	IGHV3-9-h	0.00135	0.14%	2	1
IGHV	IGHV4-31-h	0.00135	0.14%	3	2
IGHV	IGHV4-39-u	0.00135	0.14%	3	3
IGHV	IGHV4-39-t	0.00135	0.14%	3	2
IGHV	IGHV4-39-q	0.00135	0.14%	3	3
IGHV	IGHV4-4-ao	0.00135	0.14%	3	3
IGHV	IGHV4-4-ac	0.00135	0.14%	3	1
IGHV	IGHV4-4-ae	0.00135	0.14%	3	2
IGHV	IGHV5-51-g	0.00135	0.14%	3	3
IGHV	IGHV7-81-e	0.00135	0.14%	3	1
IGKJ	IGKJ2-f	0.00135	0.14%	3	2
IGKJ	IGKJ3-d	0.00135	0.14%	3	3
IGKJ	IGKJ4-g	0.00135	0.14%	3	3
IGKJ	IGKJ5-c	0.00135	0.14%	3	3
IGKJ	IGKJ5-b	0.00135	0.14%	3	3
IGKV	IGKV1-5-c	0.00135	0.14%	3	3
IGKV	IGKV1-9-j	0.00135	0.14%	3	3
IGKV	IGKV1D-17-c	0.00135	0.14%	3	2
IGKV	IGKV1D-17-b	0.00135	0.14%	3	2
IGKV	IGKV1D-42-e	0.00135	0.14%	3	1
IGKV	IGKV1D-42-d	0.00135	0.14%	3	2
IGKV	IGKV2D-24-b	0.00135	0.14%	3	2
IGKV	IGKV2D-30-c	0.00135	0.14%	3	2
IGKV	IGKV3-20-d	0.00135	0.14%	3	3
IGKV	IGKV3-20-f	0.00135	0.14%	3	2
IGKV	IGKV3D-11-d	0.00135	0.14%	3	2
IGKV	IGKV4-1-r	0.00135	0.14%	3	3
IGKV	IGKV4-1-x	0.00135	0.14%	3	2
IGKV	IGKV4-1-e	0.00135	0.14%	3	3
IGKV	IGKV4-1-o	0.00135	0.14%	3	2
IGLJ	IGLJ6-c	0.00135	0.14%	3	2
IGLJ	IGLJ7-b	0.00135	0.14%	3	3
IGLV	IGLV10-54-g	0.00135	0.14%	3	1
IGLV	IGLV11-55-h	0.00135	0.14%	3	2
IGLV	IGLV11-55-f	0.00135	0.14%	3	2
IGLV	IGLV1-40-g	0.00135	0.14%	3	3
IGLV	IGLV1-44-f	0.00135	0.14%	3	2
IGLV	IGLV1-47-f	0.00135	0.14%	3	2
IGLV	IGLV2-23-e	0.00135	0.14%	3	3
IGLV	IGLV3-12-d	0.00135	0.14%	3	3
IGLV	IGLV3-16-d	0.00135	0.14%	3	3
IGLV	IGLV3-19-d	0.00135	0.14%	3	1
IGLV	IGLV3-19-l	0.00135	0.14%	3	2
IGLV	IGLV3-1-ag	0.00135	0.14%	3	3
IGLV	IGLV3-1-y	0.00135	0.14%	3	3

IGLV	IGLV3-1-s	0.00135	0.14%	3	3
IGLV	IGLV3-1-x	0.00135	0.14%	3	3
IGLV	IGLV3-1-ad	0.00135	0.14%	3	2
IGLV	IGLV3-1-l	0.00135	0.14%	3	3
IGLV	IGLV3-1-p	0.00135	0.14%	3	2
IGLV	IGLV3-21-t	0.00135	0.14%	3	2
IGLV	IGLV3-22-f	0.00135	0.14%	3	3
IGLV	IGLV3-22-i	0.00135	0.14%	3	1
IGLV	IGLV3-25-e	0.00135	0.14%	3	3
IGLV	IGLV5-48-t	0.00135	0.14%	3	2
IGLV	IGLV5-48-s	0.00135	0.14%	3	3
IGLV	IGLV5-52-d	0.00135	0.14%	3	2
IGLV	IGLV5-52-c	0.00135	0.14%	3	1
IGLV	IGLV7-43-c	0.00135	0.14%	3	2
IGHJ	IGHJ5-d	0.0009	0.09%	2	2
IGHJ	IGHJ6-e	0.0009	0.09%	2	2
IGHJ	IGHJ6-h	0.0009	0.09%	2	2
IGHJ	IGHJ6-f	0.0009	0.09%	2	2
IGHV	IGHV1-18-f	0.0009	0.09%	2	2
IGHV	IGHV1-24-g	0.0009	0.09%	2	1
IGHV	IGHV1-2-i	0.0009	0.09%	2	2
IGHV	IGHV1-2-m	0.0009	0.09%	2	2
IGHV	IGHV1-2-l	0.0009	0.09%	2	2
IGHV	IGHV1-3-ag	0.0009	0.09%	2	2
IGHV	IGHV1-3-aj	0.0009	0.09%	2	2
IGHV	IGHV1-3-aq	0.0009	0.09%	2	1
IGHV	IGHV1-3-s	0.0009	0.09%	2	2
IGHV	IGHV1-3-q	0.0009	0.09%	2	1
IGHV	IGHV1-3-z	0.0009	0.09%	2	1
IGHV	IGHV1-3-ad	0.0009	0.09%	2	2
IGHV	IGHV1-3-aw	0.0009	0.09%	2	2
IGHV	IGHV1-3-ae	0.0009	0.09%	2	2
IGHV	IGHV1-3-an	0.0009	0.09%	2	2
IGHV	IGHV1-46-k	0.0009	0.09%	2	2
IGHV	IGHV1-46-e	0.0009	0.09%	2	2
IGHV	IGHV1-46-n	0.0009	0.09%	2	2
IGHV	IGHV1-58-g	0.0009	0.09%	2	1
IGHV	IGHV1-69-ah	0.0009	0.09%	2	2
IGHV	IGHV1-69-t	0.0009	0.09%	2	2
IGHV	IGHV1-69-ao	0.0009	0.09%	2	1
IGHV	IGHV1-69-af	0.0009	0.09%	2	2
IGHV	IGHV1-69-ad	0.0009	0.09%	2	2
IGHV	IGHV1-69-n	0.0009	0.09%	2	2
IGHV	IGHV1-69-an	0.0009	0.09%	2	2
IGHV	IGHV1-69-o	0.0009	0.09%	2	2
IGHV	IGHV1-8-h	0.0009	0.09%	2	2

IGHV	IGHV2-26-f	0.0009	0.09%	2	2
IGHV	IGHV2-5-c	0.0009	0.09%	2	2
IGHV	IGHV2-70-g	0.0009	0.09%	2	2
IGHV	IGHV2-70-s	0.0009	0.09%	2	2
IGHV	IGHV2-70-ab	0.0009	0.09%	2	2
IGHV	IGHV3-11-b	0.0009	0.09%	2	2
IGHV	IGHV3-13-f	0.0009	0.09%	2	2
IGHV	IGHV3-20-d	0.0009	0.09%	2	2
IGHV	IGHV3-21-c	0.0009	0.09%	2	2
IGHV	IGHV3-21-h	0.0009	0.09%	2	2
IGHV	IGHV3-23-j	0.0009	0.09%	2	1
IGHV	IGHV3-23-h	0.0009	0.09%	2	2
IGHV	IGHV3-33-f	0.0009	0.09%	2	2
IGHV	IGHV3-43-p	0.0009	0.09%	2	2
IGHV	IGHV3-64-h	0.0009	0.09%	2	1
IGHV	IGHV3-66-h	0.0009	0.09%	2	2
IGHV	IGHV3-74-c	0.0009	0.09%	2	2
IGHV	IGHV3-9-o	0.0009	0.09%	2	1
IGHV	IGHV4-28-s	0.0009	0.09%	2	2
IGHV	IGHV4-28-n	0.0009	0.09%	2	2
IGHV	IGHV4-31-t	0.0009	0.09%	2	2
IGHV	IGHV4-31-m	0.0009	0.09%	2	2
IGHV	IGHV4-31-l	0.0009	0.09%	2	2
IGHV	IGHV4-34-d	0.0009	0.09%	2	2
IGHV	IGHV4-34-g	0.0009	0.09%	2	2
IGHV	IGHV4-34-b	0.0009	0.09%	2	2
IGHV	IGHV4-39-aa	0.0009	0.09%	2	2
IGHV	IGHV4-39-s	0.0009	0.09%	2	2
IGHV	IGHV4-39-x	0.0009	0.09%	2	2
IGHV	IGHV4-39-p	0.0009	0.09%	2	2
IGHV	IGHV4-4-aa	0.0009	0.09%	2	2
IGHV	IGHV4-4-af	0.0009	0.09%	2	2
IGHV	IGHV4-4-x	0.0009	0.09%	2	2
IGHV	IGHV4-4-ad	0.0009	0.09%	2	2
IGHV	IGHV4-4-ai	0.0009	0.09%	2	1
IGHV	IGHV5-51-d	0.0009	0.09%	2	2
IGHV	IGHV5-51-k	0.0009	0.09%	2	2
IGHV	IGHV5-51-c	0.0009	0.09%	2	2
IGHV	IGHV6-1-a	0.0009	0.09%	2	2
IGHV	IGHV6-1-c	0.0009	0.09%	2	1
IGHV	IGHV7-81-d	0.0009	0.09%	2	2
IGKJ	IGKJ2-g	0.0009	0.09%	2	2
IGKJ	IGKJ3-b	0.0009	0.09%	2	2
IGKJ	IGKJ4-j	0.0009	0.09%	2	1
IGKJ	IGKJ4-c	0.0009	0.09%	2	2
IGKJ	IGKJ4-k	0.0009	0.09%	2	2

IGKJ	IGKJ4-h	0.0009	0.09%	2	2
IGKV	IGKV1-16-d	0.0009	0.09%	2	2
IGKV	IGKV1-17-e	0.0009	0.09%	2	2
IGKV	IGKV1-5-d	0.0009	0.09%	2	1
IGKV	IGKV1-8-h	0.0009	0.09%	2	2
IGKV	IGKV1-9-i	0.0009	0.09%	2	2
IGKV	IGKV1D-16-c	0.0009	0.09%	2	1
IGKV	IGKV1D-16-b	0.0009	0.09%	2	1
IGKV	IGKV1D-42-g	0.0009	0.09%	2	2
IGKV	IGKV1D-42-f	0.0009	0.09%	2	1
IGKV	IGKV1D-43-e	0.0009	0.09%	2	1
IGKV	IGKV1D-43-d	0.0009	0.09%	2	2
IGKV	IGKV1D-8-d	0.0009	0.09%	2	1
IGKV	IGKV1D-8-f	0.0009	0.09%	2	2
IGKV	IGKV2D-26-d	0.0009	0.09%	2	2
IGKV	IGKV2D-26-f	0.0009	0.09%	2	2
IGKV	IGKV2D-29-b	0.0009	0.09%	2	1
IGKV	IGKV3-11-g	0.0009	0.09%	2	1
IGKV	IGKV3-11-f	0.0009	0.09%	2	2
IGKV	IGKV3-20-j	0.0009	0.09%	2	2
IGKV	IGKV3-20-i	0.0009	0.09%	2	2
IGKV	IGKV3-20-e	0.0009	0.09%	2	2
IGKV	IGKV3-20-c	0.0009	0.09%	2	2
IGKV	IGKV3D-11-h	0.0009	0.09%	2	1
IGKV	IGKV3D-11-i	0.0009	0.09%	2	1
IGKV	IGKV3D-20-d	0.0009	0.09%	2	2
IGKV	IGKV4-1-y	0.0009	0.09%	2	2
IGKV	IGKV4-1-g	0.0009	0.09%	2	2
IGKV	IGKV4-1-t	0.0009	0.09%	2	2
IGKV	IGKV4-1-i	0.0009	0.09%	2	2
IGKV	IGKV4-1-n	0.0009	0.09%	2	2
IGKV	IGKV4-1-l	0.0009	0.09%	2	2
IGLJ	IGLJ2-e	0.0009	0.09%	2	2
IGLJ	IGLJ2-c	0.0009	0.09%	2	2
IGLJ	IGLJ5-b	0.0009	0.09%	2	1
IGLV	IGLV10-54-n	0.0009	0.09%	2	2
IGLV	IGLV10-54-j	0.0009	0.09%	2	2
IGLV	IGLV1-40-h	0.0009	0.09%	2	2
IGLV	IGLV1-40-e	0.0009	0.09%	2	2
IGLV	IGLV1-47-e	0.0009	0.09%	2	2
IGLV	IGLV1-51-g	0.0009	0.09%	2	2
IGLV	IGLV2-11-j	0.0009	0.09%	2	2
IGLV	IGLV2-11-f	0.0009	0.09%	2	2
IGLV	IGLV2-11-i	0.0009	0.09%	2	2
IGLV	IGLV2-11-e	0.0009	0.09%	2	1
IGLV	IGLV2-14-p	0.0009	0.09%	2	2

IGLV	IGLV2-14-q	0.0009	0.09%	2	1
IGLV	IGLV2-14-o	0.0009	0.09%	2	1
IGLV	IGLV2-23-k	0.0009	0.09%	2	2
IGLV	IGLV2-23-o	0.0009	0.09%	2	2
IGLV	IGLV2-33-c	0.0009	0.09%	2	1
IGLV	IGLV3-19-j	0.0009	0.09%	2	2
IGLV	IGLV3-19-k	0.0009	0.09%	2	2
IGLV	IGLV3-19-h	0.0009	0.09%	2	1
IGLV	IGLV3-19-i	0.0009	0.09%	2	1
IGLV	IGLV3-19-p	0.0009	0.09%	2	2
IGLV	IGLV3-1-ah	0.0009	0.09%	2	2
IGLV	IGLV3-1-d	0.0009	0.09%	2	2
IGLV	IGLV3-1-j	0.0009	0.09%	2	1
IGLV	IGLV3-1-g	0.0009	0.09%	2	2
IGLV	IGLV3-1-z	0.0009	0.09%	2	2
IGLV	IGLV3-1-ac	0.0009	0.09%	2	2
IGLV	IGLV3-1-ae	0.0009	0.09%	2	2
IGLV	IGLV3-21-k	0.0009	0.09%	2	2
IGLV	IGLV3-21-r	0.0009	0.09%	2	1
IGLV	IGLV3-21-i	0.0009	0.09%	2	2
IGLV	IGLV3-22-j	0.0009	0.09%	2	1
IGLV	IGLV3-22-n	0.0009	0.09%	2	2
IGLV	IGLV3-22-m	0.0009	0.09%	2	2
IGLV	IGLV3-22-o	0.0009	0.09%	2	2
IGLV	IGLV3-25-g	0.0009	0.09%	2	1
IGLV	IGLV3-25-l	0.0009	0.09%	2	2
IGLV	IGLV3-27-a	0.0009	0.09%	2	2
IGLV	IGLV3-32-c	0.0009	0.09%	2	2
IGLV	IGLV3-9-a	0.0009	0.09%	2	2
IGLV	IGLV3-9-b	0.0009	0.09%	2	2
IGLV	IGLV4-3-d	0.0009	0.09%	2	2
IGLV	IGLV4-60-g	0.0009	0.09%	2	2
IGLV	IGLV4-60-n	0.0009	0.09%	2	2
IGLV	IGLV4-60-o	0.0009	0.09%	2	2
IGLV	IGLV5-45-g	0.0009	0.09%	2	2
IGLV	IGLV5-48-u	0.0009	0.09%	2	2
IGLV	IGLV5-48-q	0.0009	0.09%	2	2
IGLV	IGLV5-48-r	0.0009	0.09%	2	2
IGLV	IGLV5-48-n	0.0009	0.09%	2	2
IGLV	IGLV5-48-m	0.0009	0.09%	2	1
IGLV	IGLV5-48-l	0.0009	0.09%	2	2
IGLV	IGLV5-52-b	0.0009	0.09%	2	1
IGLV	IGLV7-46-c	0.0009	0.09%	2	2
IGLV	IGLV9-49-c	0.0009	0.09%	2	2

Table 20B: Variants Found In No More Than 3 Human Populations & In At Least 10 Individuals

TYPE	VARIANT	C.FREQ	C.FREQ (%)	#INDIVIDUALS	#POPULATIONS
IGKV	IGKV1D-8-b	0.01785	1.79%	36	3
IGHV	IGHV3-11-a	0.01695	1.70%	36	3
IGHV	IGHV5-51-a	0.0164	1.64%	35	3
IGHV	IGHV2-5-b	0.01055	1.06%	18	3
IGLV	IGLV2-14-g	0.0104	1.04%	22	3
IGHV	IGHV7-81-b	0.01005	1.01%	21	3
IGKV	IGKV1D-42-b	0.01005	1.01%	21	3
IGKV	IGKV3-7-b	0.01	1.00%	19	3
IGHV	IGHV1-69-d	0.00995	1.00%	22	3
IGLV	IGLV2-14-e	0.00995	1.00%	19	3
IGLV	IGLV9-49-a	0.00865	0.87%	18	3
IGLV	IGLV1-44-b	0.00775	0.78%	16	2
IGLV	IGLV2-11-a	0.00775	0.78%	17	3
IGKV	IGKV3-11-d	0.0073	0.73%	14	3
IGKV	IGKV1D-43-b	0.00685	0.69%	14	3
IGHV	IGHV3-38-d	0.0064	0.64%	13	3
IGLJ	IGLJ6-a	0.00595	0.60%	13	3
IGHV	IGHV5-51-b	0.0055	0.55%	11	3
IGLV	IGLV5-45-c	0.00545	0.55%	12	2
IGHV	IGHV4-31-i	0.0054	0.54%	10	3
IGLV	IGLV1-36-c	0.00455	0.46%	10	2

Table 20C: Variants Found In No More Than 3 Human Populations & In At Least 20 Individuals

TYPE	VARIANT	C.FREQ	C.FREQ (%)	#INDIVIDUALS	#POPULATIONS
IGKV	IGKV1D-8-b	0.01785	1.79%	36	3
IGHV	IGHV3-11-a	0.01695	1.70%	36	3
IGHV	IGHV5-51-a	0.0164	1.64%	35	3
IGLV	IGLV2-14-g	0.0104	1.04%	22	3
IGHV	IGHV7-81-b	0.01005	1.01%	21	3
IGKV	IGKV1D-42-b	0.01005	1.01%	21	3
IGHV	IGHV1-69-d	0.00995	1.00%	22	3

Table 20D: Variants Found In No More Than 3 Human Populations & In At Least 30 Individuals

TYPE	VARIANT	C.FREQ	C.FREQ (%)	#INDIVIDUALS	#POPULATIONS
IGKV	IGKV1D-8-b	0.01785	1.79%	36	3
IGHV	IGHV3-11-a	0.01695	1.70%	36	3
IGHV	IGHV5-51-a	0.0164	1.64%	35	3

Table 21

<u>Seq ID #</u>	<u>Associated Gene Name</u>	<u>Ensembl Gene ID</u>	<u>Gene Location</u> [Chromosome Name:Gene Chr Start (bp)-GeneChr End (bp)]	<u>Ensembl Transcript ID</u>	<u>Exon Positions</u> [Chromosome Name:Exon Chr Start (bp)-Exon Chr End (bp)]
<u>239</u>	<u>IGHJ1</u>	<u>ENSG000000211905</u>	<u>14:1063331617-1063331668</u>	<u>ENST000000390565</u>	<u>14:1063331617-1063331668</u>
<u>240</u>	<u>IGHJ2</u>	<u>ENSG000000211904</u>	<u>14:1063331409-1063331460</u>	<u>ENST000000390564</u>	<u>14:1063331409-1063331460</u>
<u>241</u>	<u>IGHJ3</u>	<u>ENSG000000242887</u>	<u>14:1063330797-1063330845</u>	<u>ENST000000463911</u>	<u>14:1063330797-1063330845</u>
<u>242</u>	<u>IGHJ4</u>	<u>ENSG000000240041</u>	<u>14:1063330425-1063330470</u>	<u>ENST000000461719</u>	<u>14:1063330425-1063330470</u>
<u>243</u>	<u>IGHJ5</u>	<u>ENSG000000242472</u>	<u>14:1063330024-1063330072</u>	<u>ENST000000488476</u>	<u>14:1063330024-1063330072</u>
<u>244</u>	<u>IGHJ6</u>	<u>ENSG000000211900</u>	<u>14:106329408-106329468</u>	<u>ENST000000390560</u>	<u>14:106329408-106329468</u>
<u>245</u>	<u>IGHV1-18</u>	<u>ENSG000000211945</u>	<u>14:106641563-106642056</u>	<u>ENST000000390605</u>	<u>14:106641952-106642056 14:106641563-106641867</u>
<u>246</u>	<u>IGHV1-2</u>	<u>ENSG000000211934</u>	<u>14:106452671-106453170</u>	<u>ENST000000390594</u>	<u>14:106453061-106453170 14:106452671-106452975</u>
<u>247</u>	<u>IGHV1-24</u>	<u>ENSG000000211950</u>	<u>14:106733144-106733639</u>	<u>ENST000000390610</u>	<u>14:106733534-106733639 14:106733144-106733448</u>
<u>248</u>	<u>IGHV1-3</u>	<u>ENSG000000211935</u>	<u>14:106471246-106471723</u>	<u>ENST000000390595</u>	<u>14:106471636-106471723 14:106471246-106471550</u>
<u>249</u>	<u>IGHV1-45</u>	<u>ENSG000000211961</u>	<u>14:106962931-106963424</u>	<u>ENST000000390621</u>	<u>14:106963321-106963424 14:106962931-106963235</u>
<u>250</u>	<u>IGHV1-46</u>	<u>ENSG000000211962</u>	<u>14:106967049-106967788</u>	<u>ENST000000390622</u>	<u>14:106967439-106967788 14:106967049-106967353</u>
<u>251</u>	<u>IGHV1-58</u>	<u>ENSG000000211968</u>	<u>14:107078373-107078869</u>	<u>ENST000000390628</u>	<u>14:107078763-107078869 14:107078373-107078677</u>

<u>252</u>	<u>IGHV1-69</u>	<u>ENSG000000211973</u>	<u>14:107169931-107170428</u>	<u>ENST000000390633</u>	<u>14:107170322-107170428 14:107169931-107170235</u>
<u>253</u>	<u>IGHV1-8</u>	<u>ENSG000000211939</u>	<u>14:106539079-106539577</u>	<u>ENST000000390599</u>	<u>14:106539470-106539577 14:106539079-106539383</u>
<u>254</u>	<u>IGHV2-26</u>	<u>ENSG000000211951</u>	<u>14:106757650-106758116</u>	<u>ENST000000390611</u>	<u>14:106758047-106758116 14:106757650-106757960</u>
<u>255</u>	<u>IGHV2-5</u>	<u>ENSG000000211937</u>	<u>14:106494135-106494597</u>	<u>ENST000000390597</u>	<u>14:106494532-106494597 14:106494135-106494445</u>
<u>256</u>	<u>IGHV2-70</u>	<u>ENSG000000211974</u>	<u>14:107178820-107179338</u>	<u>ENST000000390634</u>	<u>14:107179217-107179338 14:107178820-107179130</u>
<u>257</u>	<u>IGHV3-11</u>	<u>ENSG000000211941</u>	<u>14:106573233-106573800</u>	<u>ENST000000390601</u>	<u>14:106573635-106573800 14:106573233-106573537</u>
<u>258</u>	<u>IGHV3-13</u>	<u>ENSG000000211942</u>	<u>14:106586137-106586667</u>	<u>ENST000000390602</u>	<u>14:106586542-106586667 14:106586137-106586438</u>
<u>259</u>	<u>IGHV3-15</u>	<u>ENSG000000211943</u>	<u>14:106610313-106610852</u>	<u>ENST000000390603</u>	<u>14:106610727-106610852 14:106610313-106610623</u>
<u>260</u>	<u>IGHV3-16</u>	<u>ENSG000000211944</u>	<u>14:106621894-106622419</u>	<u>ENST000000390604</u>	<u>14:106622300-106622419 14:106621894-106622198</u>
<u>261</u>	<u>IGHV3-20</u>	<u>ENSG000000211946</u>	<u>14:106667581-106668095</u>	<u>ENST000000390606</u>	<u>14:106667988-106668095 14:106667581-106667885</u>
<u>262</u>	<u>IGHV3-21</u>	<u>ENSG000000211947</u>	<u>14:106691673-106692203</u>	<u>ENST000000390607</u>	<u>14:106692079-106692203 14:106691673-106691977</u>
<u>263</u>	<u>IGHV3-23</u>	<u>ENSG000000211949</u>	<u>14:106725201-106725733</u>	<u>ENST000000390609</u>	<u>14:106725609-106725733 14:106725201-106725505</u>
<u>264</u>	<u>IGHV3-30</u>	<u>ENSG000000211953</u>	<u>14:106791005-106791536</u>	<u>ENST000000390613</u>	<u>14:106791411-106791536 14:106791005-106791309</u>
<u>265</u>	<u>IGHV3-33</u>	<u>ENSG000000211955</u>	<u>14:106815722-106816253</u>	<u>ENST000000390615</u>	<u>14:106816128-106816253 14:106815722-106816026</u>
<u>266</u>	<u>IGHV3-35</u>	<u>ENSG000000211957</u>	<u>14:106845323-106845789</u>	<u>ENST000000390617</u>	<u>14:106845729-106845789 14:106845323-106845627</u>
<u>267</u>	<u>IGHV3-38</u>	<u>ENSG000000211958</u>	<u>14:106866406-106866934</u>	<u>ENST000000390618</u>	<u>14:106866811-106866934 14:106866406-106866707</u>
<u>268</u>	<u>IGHV3-43</u>	<u>ENSG000000232216</u>	<u>14:106926188-106926724</u>	<u>ENST000000434710</u>	<u>14:106926599-106926724 14:106926188-106926495</u>

<u>269</u>	<u>IGHV3-48</u>	<u>ENSG000000211964</u>	<u>14:106993814-106994346</u>	<u>ENST000000390624</u>	<u>14:106994222-106994346 14:106993814-106994118</u>
<u>270</u>	<u>IGHV3-49</u>	<u>ENSG000000211965</u>	<u>14:107012938-107013477</u>	<u>ENST000000390625</u>	<u>14:107013352-107013477 14:107012938-107013248</u>
<u>271</u>	<u>IGHV3-53</u>	<u>ENSG000000211967</u>	<u>14:107048672-107049341</u>	<u>ENST000000390627</u>	<u>14:107049075-107049341 14:107048672-107048973</u>
<u>272</u>	<u>IGHV3-64</u>	<u>ENSG000000223648</u>	<u>14:107113741-107114274</u>	<u>ENST000000454421</u>	<u>14:107114149-107114274 14:107113741-107114045</u>
<u>273</u>	<u>IGHV3-66</u>	<u>ENSG000000211972</u>	<u>14:107131033-107131560</u>	<u>ENST000000390632</u>	<u>14:107131436-107131560 14:107131033-107131334</u>
<u>274</u>	<u>IGHV3-7</u>	<u>ENSG000000211938</u>	<u>14:106518400-106518932</u>	<u>ENST000000390598</u>	<u>14:106518808-106518932 14:106518400-106518704</u>
<u>275</u>	<u>IGHV3-72</u>	<u>ENSG000000225698</u>	<u>14:107198932-107199471</u>	<u>ENST000000433072</u>	<u>14:107199346-107199471 14:107198932-107199242</u>
<u>276</u>	<u>IGHV3-73</u>	<u>ENSG000000211976</u>	<u>14:107210932-107211471</u>	<u>ENST000000390636</u>	<u>14:107211346-107211471 14:107210932-107211242</u>
<u>277</u>	<u>IGHV3-74</u>	<u>ENSG000000224650</u>	<u>14:107218676-107219365</u>	<u>ENST000000424969</u>	<u>14:107219084-107219365 14:107218676-107218980</u>
<u>278</u>	<u>IGHV3-9</u>	<u>ENSG000000211940</u>	<u>14:106552285-106552809</u>	<u>ENST000000390600</u>	<u>14:106552684-106552809 14:106552285-106552592</u>
<u>279</u>	<u>IGHV4-28</u>	<u>ENSG000000211952</u>	<u>14:106780513-106781017</u>	<u>ENST000000390612</u>	<u>14:106780900-106781017 14:106780513-106780817</u>
<u>280</u>	<u>IGHV4-31</u>	<u>ENSG000000231475</u>	<u>14:106805209-106805716</u>	<u>ENST000000438142</u>	<u>14:106805599-106805716 14:106805209-106805516</u>
<u>281</u>	<u>IGHV4-34</u>	<u>ENSG000000211956</u>	<u>14:106829594-106830076</u>	<u>ENST000000390616</u>	<u>14:106829979-106830076 14:106829594-106829895</u>
<u>282</u>	<u>IGHV4-39</u>	<u>ENSG000000211959</u>	<u>14:106877619-106878126</u>	<u>ENST000000390619</u>	<u>14:106878010-106878126 14:106877619-106877926</u>
<u>283</u>	<u>IGHV4-4</u>	<u>ENSG000000211936</u>	<u>14:106478110-106478603</u>	<u>ENST000000390596</u>	<u>14:106478494-106478603 14:106478110-106478411</u>
<u>284</u>	<u>IGHV4-59</u>	<u>ENSG000000224373</u>	<u>14:107081806-107083830</u>	<u>ENST000000390629</u>	<u>14:107083640-107083830 14:107083256-107083557</u>
<u>284</u>	<u>IGHV4-59</u>	<u>ENSG000000224373</u>	<u>14:107081806-107083830</u>	<u>ENST000000455737</u>	<u>14:107083640-107083725 14:107083240-107083557</u>

					<u>14:107081806-107082728</u>	
<u>285</u>	<u>IGHV4-61</u>	<u>ENSG000000211970</u>		<u>14:107095126-107095662</u>	<u>ENST000000390630</u>	<u>14:107095516-107095662 14:107095126-107095433</u>
<u>286</u>	<u>IGHV5-51</u>	<u>ENSG000000211966</u>		<u>14:107034729-107035221</u>	<u>ENST000000390626</u>	<u>14:107035117-107035221 14:107034729-107035033</u>
<u>287</u>	<u>IGHV6-1</u>	<u>ENSG000000211933</u>		<u>14:106405611-106406108</u>	<u>ENST000000390593</u>	<u>14:106406008-106406108 14:106405611-106405924</u>
<u>288</u>	<u>IGHV7-81</u>	<u>ENSG000000211979</u>		<u>14:107282792-107283280</u>	<u>ENST000000390639</u>	<u>14:107283181-107283280 14:107282792-107283096</u>
<u>289</u>	<u>IGKJ2</u>	<u>ENSG000000211596</u>		<u>2:89161037-89161074</u>	<u>ENST000000390241</u>	<u>2:89161037-89161074</u>
<u>290</u>	<u>IGKJ3</u>	<u>ENSG000000211595</u>		<u>2:89160733-89160770</u>	<u>ENST000000390240</u>	<u>2:89160733-89160770</u>
<u>291</u>	<u>IGKJ4</u>	<u>ENSG000000211594</u>		<u>2:89160398-89160434</u>	<u>ENST000000390239</u>	<u>2:89160398-89160434</u>
<u>292</u>	<u>IGKJ5</u>	<u>ENSG000000211593</u>		<u>2:89160080-89160117</u>	<u>ENST000000390238</u>	<u>2:89160080-89160117</u>
<u>293</u>	<u>IGKV1-16</u>	<u>ENSG000000240864</u>		<u>2:89399352-89399854</u>	<u>ENST000000479981</u>	<u>2:89399773-89399854 2:89399352-89399647</u>
<u>294</u>	<u>IGKV1-17</u>	<u>ENSG000000240382</u>		<u>2:89416833-89417335</u>	<u>ENST000000490686</u>	<u>2:89417254-89417335 2:89416833-89417128</u>
<u>295</u>	<u>IGKV1-5</u>	<u>ENSG000000243466</u>		<u>2:89246819-89247475</u>	<u>ENST000000496168</u>	<u>2:89247240-89247475 2:89246819-89247114</u>
<u>296</u>	<u>IGKV1-6</u>	<u>ENSG000000239855</u>		<u>2:89265781-89266286</u>	<u>ENST000000464162</u>	<u>2:89266203-89266286 2:89265781-89266076</u>
<u>297</u>	<u>IGKV1-8</u>	<u>ENSG000000240671</u>		<u>2:89291928-89292450</u>	<u>ENST000000495489</u>	<u>2:89292349-89292450 2:89291928-89292223</u>
<u>298</u>	<u>IGKV1-9</u>	<u>ENSG000000241755</u>		<u>2:89309479-89310012</u>	<u>ENST000000493819</u>	<u>2:89309900-89310012 2:89309479-89309774</u>
<u>299</u>	<u>IGKV1D-12</u>	<u>ENSG000000240834</u>		<u>2:90198535-90199190</u>	<u>ENST000000390276</u>	<u>2:90198535-90198770 2:90198895-90199190</u>
<u>300</u>	<u>IGKV1D-16</u>	<u>ENSG000000241244</u>		<u>2:90139078-90139580</u>	<u>ENST000000492446</u>	<u>2:90139078-90139159 2:90139285-90139580</u>

<u>301</u>	<u>IGKV1D-17</u>	<u>ENSG000000242766</u>	<u>2:90121477-90122133</u>	<u>ENST000000483379</u>	<u>2:90121477-90121712 2:90121838-90122133</u>
<u>302</u>	<u>IGKV1D-42</u>	<u>ENSG000000211633</u>	<u>2:90229045-90229531</u>	<u>ENST000000390278</u>	<u>2:90229045-90229119 2:90229236-90229531</u>
<u>303</u>	<u>IGKV1D-43</u>	<u>ENSG000000242580</u>	<u>2:90248739-90249395</u>	<u>ENST000000468879</u>	<u>2:90248739-90248974 2:90249100-90249395</u>
<u>304</u>	<u>IGKV1D-8</u>	<u>ENSG000000239819</u>	<u>2:90259593-90260248</u>	<u>ENST000000471857</u>	<u>2:90259593-90259828 2:90259953-90260248</u>
<u>305</u>	<u>IGKV2-24</u>	<u>ENSG000000241294</u>	<u>2:89475812-89476644</u>	<u>ENST000000484817</u>	<u>2:89476566-89476644 2:89475812-89476122</u>
<u>306</u>	<u>IGKV2-30</u>	<u>ENSG000000243238</u>	<u>2:89544264-89545079</u>	<u>ENST000000468494</u>	<u>2:89545001-89545079 2:89544264-89544574</u>
<u>307</u>	<u>IGKV2D-24</u>	<u>ENSG000000241566</u>	<u>2:90043607-90044439</u>	<u>ENST000000462693</u>	<u>2:90043607-90043685 2:90044129-90044439</u>
<u>308</u>	<u>IGKV2D-26</u>	<u>ENSG000000211623</u>	<u>2:90024732-90025512</u>	<u>ENST000000390268</u>	<u>2:90024732-90024810 2:90025202-90025512</u>
<u>309</u>	<u>IGKV2D-29</u>	<u>ENSG000000243264</u>	<u>2:89986322-89987079</u>	<u>ENST000000491977</u>	<u>2:89986322-89986400 2:89986769-89987079</u>
<u>310</u>	<u>IGKV2D-30</u>	<u>ENSG000000239571</u>	<u>2:89975669-89976489</u>	<u>ENST000000474213</u>	<u>2:89975669-89975752 2:89976179-89976489</u>
<u>311</u>	<u>IGKV3-11</u>	<u>ENSG000000241351</u>	<u>2:89326668-89327228</u>	<u>ENST000000483158</u>	<u>2:89327133-89327228 2:89326668-89326963</u>
<u>312</u>	<u>IGKV3-20</u>	<u>ENSG000000239951</u>	<u>2:89442057-89442643</u>	<u>ENST000000492167</u>	<u>2:89442543-89442643 2:89442057-89442355</u>
<u>313</u>	<u>IGKV3-7</u>	<u>ENSG000000243063</u>	<u>2:89277987-89278600</u>	<u>ENST000000390247</u>	<u>2:89278455-89278600 2:89277987-89278285</u>
<u>314</u>	<u>IGKV3D-11</u>	<u>ENSG000000211632</u>	<u>2:90211643-90212253</u>	<u>ENST000000390277</u>	<u>2:90211643-90211788 2:90211958-90212253</u>
<u>315</u>	<u>IGKV3D-15</u>	<u>ENSG000000224041</u>	<u>2:90153696-90154258</u>	<u>ENST000000417279</u>	<u>2:90153696-90153793 2:90153963-90154258</u>
<u>316</u>	<u>IGKV3D-20</u>	<u>ENSG000000211625</u>	<u>2:90077680-90078311</u>	<u>ENST000000390270</u>	<u>2:90077680-90077825 2:90078013-90078311</u>
<u>317</u>	<u>IGKV4-1</u>	<u>ENSG000000211598</u>	<u>2:89184913-89185669</u>	<u>ENST000000390243</u>	<u>2:89184913-89185136 2:89185356-89185669</u>

<u>318</u>	<u>IGKV5-2</u>	<u>ENSG0000002111599</u>	<u>2:89196748-89197300</u>	<u>ENST000000390244</u>	<u>2:89196748-89196859 2:89197005-89197300</u>
<u>319</u>	<u>IGKV6-21</u>	<u>ENSG0000002111611</u>	<u>2:89459235-89459850</u>	<u>ENST000000390256</u>	<u>2:89459741-89459850 2:89459235-89459530</u>
<u>320</u>	<u>IGKV6D-21</u>	<u>ENSG000000225523</u>	<u>2:90060377-90060995</u>	<u>ENST000000436451</u>	<u>2:90060377-90060489 2:90060700-90060995</u>
<u>321</u>	<u>IGKV6D-41</u>	<u>ENSG0000002111626</u>	<u>2:90108504-90109080</u>	<u>ENST000000390271</u>	<u>2:90108504-90108578 2:90108785-90109080</u>
<u>322</u>	<u>IGLJ1</u>	<u>ENSG0000002111674</u>	<u>22:23235872-23235998</u>	<u>ENST000000390320</u>	<u>22:23235872-23235998</u>
<u>323</u>	<u>IGLJ2</u>	<u>ENSG0000002111676</u>	<u>22:23241661-23241835</u>	<u>ENST000000390322</u>	<u>22:23241661-23241835</u>
<u>324</u>	<u>IGLJ3</u>	<u>ENSG0000002111678</u>	<u>22:23247030-23247205</u>	<u>ENST000000390324</u>	<u>22:23247030-23247205</u>
<u>325</u>	<u>IGLJ5</u>	<u>ENSG0000002111681</u>	<u>22:23256408-23256479</u>	<u>ENST000000390327</u>	<u>22:23256408-23256479</u>
<u>326</u>	<u>IGLJ6</u>	<u>ENSG0000002111682</u>	<u>22:23260304-23260373</u>	<u>ENST000000390328</u>	<u>22:23260304-23260373</u>
<u>327</u>	<u>IGLJ7</u>	<u>ENSG0000002111684</u>	<u>22:23263562-23263607</u>	<u>ENST000000390330</u>	<u>22:23263562-23263607</u>
<u>328</u>	<u>IGLV10-54</u>	<u>ENSG0000002111642</u>	<u>22:22569184-22569660</u>	<u>ENST000000390287</u>	<u>22:22569184-22569242 22:22569355-22569660</u>
<u>329</u>	<u>IGLV11-55</u>	<u>ENSG0000002111641</u>	<u>22:22556057-22556600</u>	<u>ENST000000390286</u>	<u>22:22556057-22556114 22:22556233-22556600</u>
<u>330</u>	<u>IGLV1-36</u>	<u>ENSG0000002111655</u>	<u>22:22786296-22786802</u>	<u>ENST000000390301</u>	<u>22:22786296-22786381 22:22786497-22786802</u>
<u>331</u>	<u>IGLV1-40</u>	<u>ENSG0000002111653</u>	<u>22:22764098-22764614</u>	<u>ENST000000390299</u>	<u>22:22764098-22764194 22:22764305-22764614</u>
<u>332</u>	<u>IGLV1-44</u>	<u>ENSG0000002111651</u>	<u>22:22735135-22735715</u>	<u>ENST000000390297</u>	<u>22:22735135-22735294 22:22735410-22735715</u>
<u>333</u>	<u>IGLV1-47</u>	<u>ENSG0000002111648</u>	<u>22:22712087-22712608</u>	<u>ENST000000390294</u>	<u>22:22712087-22712188 22:22712304-22712608</u>
<u>334</u>	<u>IGLV1-50</u>	<u>ENSG0000002111645</u>	<u>22:22681658-22682172</u>	<u>ENST000000390291</u>	<u>22:22681658-22681754 22:22681865-22682172</u>

<u>335</u>	<u>IGLV1-51</u>	<u>ENSG000000211644</u>	<u>22:22676828-22677336</u>	<u>ENST000000390290</u>	<u>22:22676828-22676909 22:22677019-22677336</u>
<u>336</u>	<u>IGLV2-11</u>	<u>ENSG000000211668</u>	<u>22:23134980-23135496</u>	<u>ENST000000390314</u>	<u>22:23134980-23135067 22:23135185-23135496</u>
<u>337</u>	<u>IGLV2-14</u>	<u>ENSG000000211666</u>	<u>22:23101189-23101707</u>	<u>ENST000000390312</u>	<u>22:23101189-23101275 22:23101393-23101707</u>
<u>338</u>	<u>IGLV2-18</u>	<u>ENSG000000211664</u>	<u>22:23077095-23077584</u>	<u>ENST000000390310</u>	<u>22:23077095-23077153 22:23077270-23077584</u>
<u>339</u>	<u>IGLV2-23</u>	<u>ENSG000000211660</u>	<u>22:23040274-23040892</u>	<u>ENST000000390306</u>	<u>22:23040274-23040481 22:23040599-23040892</u>
<u>340</u>	<u>IGLV2-33</u>	<u>ENSG000000211656</u>	<u>22:22930626-22931145</u>	<u>ENST000000390302</u>	<u>22:22930626-22930729 22:22930852-22931145</u>
<u>341</u>	<u>IGLV2-8</u>	<u>ENSG000000211671</u>	<u>22:23165153-23165787</u>	<u>ENST000000390317</u>	<u>22:23165153-23165360 22:23165476-23165787</u>
<u>342</u>	<u>IGLV3-1</u>	<u>ENSG000000211673</u>	<u>22:23222886-23223576</u>	<u>ENST000000390319</u>	<u>22:23222886-23222983 22:23223277-23223576</u>
<u>343</u>	<u>IGLV3-10</u>	<u>ENSG000000211669</u>	<u>22:23154244-23154782</u>	<u>ENST000000390315</u>	<u>22:23154244-23154324 22:23154478-23154782</u>
<u>344</u>	<u>IGLV3-12</u>	<u>ENSG000000211667</u>	<u>22:23114317-23115079</u>	<u>ENST000000390313</u>	<u>22:23114317-23114371 22:23114775-23115079</u>
<u>345</u>	<u>IGLV3-16</u>	<u>ENSG000000211665</u>	<u>22:23089870-23090398</u>	<u>ENST000000390311</u>	<u>22:23089870-23089953 22:23090108-23090398</u>
<u>346</u>	<u>IGLV3-19</u>	<u>ENSG000000211663</u>	<u>22:23063108-23063630</u>	<u>ENST000000390309</u>	<u>22:23063108-23063193 22:23063340-23063630</u>
<u>347</u>	<u>IGLV3-21</u>	<u>ENSG000000211662</u>	<u>22:23054174-23055688</u>	<u>ENST000000390308</u>	<u>22:23054174-23054290 22:23054770-23054949</u> <u>22:23055384-23055688</u>
<u>348</u>	<u>IGLV3-22</u>	<u>ENSG000000211661</u>	<u>22:23046750-23047307</u>	<u>ENST000000390307</u>	<u>22:23046750-23046859 22:23047009-23047307</u>
<u>349</u>	<u>IGLV3-25</u>	<u>ENSG000000211659</u>	<u>22:23029190-23029735</u>	<u>ENST000000390305</u>	<u>22:23029190-23029278 22:23029445-23029735</u>
<u>350</u>	<u>IGLV3-27</u>	<u>ENSG000000211658</u>	<u>22:23010758-23011276</u>	<u>ENST000000390304</u>	<u>22:23010758-23010841 22:23010984-23011276</u>

<u>351</u>	<u>IGLV3-32</u>	<u>ENSG000000211657</u>	<u>22:22936998-22937501</u>	<u>ENST000000390303</u>	<u>22:22936998-22937079 22:22937226-22937501</u>
<u>352</u>	<u>IGLV3-9</u>	<u>ENSG000000211670</u>	<u>22:23161507-23162253</u>	<u>ENST000000390316</u>	<u>22:23161507-23161667 22:23161954-23162253</u>
<u>353</u>	<u>IGLV4-3</u>	<u>ENSG000000211672</u>	<u>22:23213686-23214214</u>	<u>ENST000000390318</u>	<u>22:23213686-23213763 22:23213890-23214214</u>
<u>354</u>	<u>IGLV4-60</u>	<u>ENSG000000211639</u>	<u>22:22516592-22517074</u>	<u>ENST000000390284</u>	<u>22:22516592-22516643 22:22516765-22517074</u>
<u>355</u>	<u>IGLV4-69</u>	<u>ENSG000000211637</u>	<u>22:22385332-22385870</u>	<u>ENST000000390282</u>	<u>22:22385332-22385440 22:22385561-22385870</u>
<u>356</u>	<u>IGLV5-37</u>	<u>ENSG000000211654</u>	<u>22:22781876-22782371</u>	<u>ENST000000390300</u>	<u>22:22781876-22781926 22:22782049-22782371</u>
<u>357</u>	<u>IGLV5-45</u>	<u>ENSG000000211650</u>	<u>22:22730355-22730874</u>	<u>ENST000000390296</u>	<u>22:22730355-22730428 22:22730552-22730874</u>
<u>358</u>	<u>IGLV5-48</u>	<u>ENSG000000211647</u>	<u>22:22707289-22707781</u>	<u>ENST000000390293</u>	<u>22:22707289-22707334 22:22707459-22707781</u>
<u>359</u>	<u>IGLV5-52</u>	<u>ENSG000000211643</u>	<u>22:22673082-22673581</u>	<u>ENST000000390289</u>	<u>22:22673082-22673132 22:22673254-22673581</u>
<u>360</u>	<u>IGLV6-57</u>	<u>ENSG000000211640</u>	<u>22:22550113-22550686</u>	<u>ENST000000390285</u>	<u>22:22550113-22550244 22:22550370-22550686</u>
<u>361</u>	<u>IGLV7-43</u>	<u>ENSG000000211652</u>	<u>22:22749356-22749827</u>	<u>ENST000000390298</u>	<u>22:22749356-22749435 22:22749523-22749827</u>
<u>362</u>	<u>IGLV7-46</u>	<u>ENSG000000211649</u>	<u>22:22723982-22724454</u>	<u>ENST000000390295</u>	<u>22:22723982-22724061 22:22724151-22724454</u>
<u>363</u>	<u>IGLV8-61</u>	<u>ENSG000000211638</u>	<u>22:22453110-22453622</u>	<u>ENST000000390283</u>	<u>22:22453110-22453216 22:22453316-22453622</u>
<u>364</u>	<u>IGLV9-49</u>	<u>ENSG000000223350</u>	<u>22:22697539-22698084</u>	<u>ENST000000427632</u>	<u>22:22697539-22697621 22:22697759-22698084</u>

References:

1. Nat Biotechnol. 2005 Sep;23(9):1117-25; Human antibodies from transgenic animals; Lonberg N.
2. J Clin Invest. 1992 Mar;89(3):729-38; Immunoglobulin light chain variable region gene sequences for human antibodies to Haemophilus influenzae type b capsular polysaccharide are dominated by a limited number of V kappa and V lambda segments and VJ combinations; Adderson EE, Shackelford PG, Insel RA, Quinn A, Wilson PM, Carroll WL.
3. J Immunol. 1993 Oct 15;151(8):4352-61; Clonal characterization of the human IgG antibody repertoire to Haemophilus influenzae type b polysaccharide. V. In vivo expression of individual antibody clones is dependent on Ig CH haplotypes and the categories of antigen; Chung GH, Scott MG, Kim KH, Kearney J, Siber GR, Ambrosino DM, Nahm MH.
4. J Immunol. 1998 Dec 1;161(11):6068-73; Decreased frequency of rearrangement due to the synergistic effect of nucleotide changes in the heptamer and nonamer of the recombination signal sequence of the V kappa gene A2b, which is associated with increased susceptibility of Navajos to Haemophilus influenzae type b disease; Nadel B, Tang A, Lugo G, Love V, Escuro G, Feeney AJ.
5. J Clin Invest. 1996 May 15;97(10):2277-82; A defective Vkappa A2 allele in Navajos which may play a role in increased susceptibility to haemophilus influenzae type b disease; Feeney AJ, Atkinson MJ, Cowan MJ, Escuro G, Lugo G.
6. Infect Immun. 1994 Sep;62(9):3873-80; Variable region sequences of a protective human monoclonal antibody specific for the Haemophilus influenzae type b capsular polysaccharide; Lucas AH, Larrick JW, Reason DC.
7. J Clin Invest. 1993 Jun;91(6):2734-43; Restricted immunoglobulin VH usage and VDJ combinations in the human response to Haemophilus influenzae type b capsular polysaccharide. Nucleotide sequences of monospecific anti-Haemophilus antibodies and polyspecific antibodies cross-reacting with self antigens; Adderson EE, Shackelford PG, Quinn A, Wilson PM, Cunningham MW, Insel RA, Carroll WL.
8. J Clin Invest. 1993 Mar;91(3):788-96; Variable region expression in the antibody responses of infants vaccinated with Haemophilus influenzae type b polysaccharide-protein conjugates. Description of a new lambda light chain-associated idiotypic and the relation between idiotypic expression, avidity, and vaccine formulation. The Collaborative Vaccine Study Group; Granoff DM, Shackelford PG, Holmes SJ, Lucas AH.
9. Infect Immun. 1994 May;62(5):1776-86; Variable region sequences and idiotypic expression of a protective human immunoglobulin M antibody to capsular polysaccharides of Neisseria meningitidis group B and Escherichia coli K1; Azmi FH, Lucas AH, Raff HV, Granoff DM.

10. J Clin Invest. 1992 Dec;90(6):2197-208; Sequence analyses of three immunoglobulin G anti-virus antibodies reveal their utilization of autoantibody-related immunoglobulin Vh genes, but not V lambda genes; Huang DF, Olee T, Masuho Y, Matsumoto Y, Carson DA, Chen PP.
11. Science. 2011 Aug 12;333(6044):834-5, Biochemistry. Catching a moving target, Wang TT, Palese P
12. Science. 2009 Apr 10;324(5924):246-51. Epub 2009 Feb 26; Antibody recognition of a highly conserved influenza virus epitope; Ekiert DC, Bhabha G, Elsliger MA, Friesen RH, Jongeneelen M, Throsby M, Goudsmit J, Wilson IA.
13. PLoS One. 2008;3(12):e3942. Epub 2008 Dec 16; Heterosubtypic neutralizing monoclonal antibodies cross-protective against H5N1 and H1N1 recovered from human IgM+ memory B cells; Throsby M, van den Brink E, Jongeneelen M, Poon LL, Alard P, Cornelissen L, Bakker A, Cox F, van Deventer E, Guan Y, Cinatl J, ter Meulen J, Lasters I, Carsetti R, Peiris M, de Kruif J, Goudsmit J.
14. Nat Struct Mol Biol. 2009 Mar;16(3):265-73. Epub 2009 Feb 22, Structural and functional bases for broad-spectrum neutralization of avian and human influenza A viruses, Sui J, Hwang WC, Perez S, Wei G, Aird D, Chen LM, Santelli E, Stec B, Cadwell G, Ali M, Wan H, Murakami A, Yammanuru A, Han T, Cox NJ, Bankston LA, Donis RO, Liddington RC, Marasco WA.
15. Science. 2011 Aug 12;333(6044):843-50. Epub 2011 Jul 7, A highly conserved neutralizing epitope on group 2 influenza A viruses, Ekiert DC, Friesen RH, Bhabha G, Kwaks T, Jongeneelen M, Yu W, Ophorst C, Cox F, Korse HJ, Brandenburg B, Vogels R, Brakenhoff JP, Kompier R, Koldijk MH, Cornelissen LA, Poon LL, Peiris M, Koudstaal W, Wilson IA, Goudsmit J.

CLAIMS:

1. A non-human vertebrate (optionally a mouse or a rat) or vertebrate cell whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof so that the mouse is capable of producing an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.
2. The vertebrate of claim 1, wherein the vertebrate has been immunised with a target antigen and wherein the variable domain of the heavy chain is the product of recombination between a VH, D and JH6*02 and wherein the HCDR3 length is at least 20 amino acids.
3. A non-human vertebrate cell (optionally a mouse cell or a rat cell) whose genome comprises an immunoglobulin heavy chain locus comprising human gene segment JH6*02, one or more VH gene segments and one or more D gene segments upstream of a constant region; wherein the gene segments in the heavy chain locus are operably linked to the constant region thereof for producing (eg, in a subsequent progeny cell) an antibody heavy chain produced by recombination of the human JH6*02 with a D segment and a VH segment.
4. The cell of claim 3, which is an ES cell capable of differentiation into a progeny antibody-producing cell that expresses said heavy chain.
5. The vertebrate or cell of any preceding claim, wherein the heavy chain locus comprises a human JH6*02 recombination signal sequence (RSS) operably connected 5' to the JH6*02 gene segment.
6. The vertebrate or cell of claim 5, wherein the RSS is SEQ ID NO: 238 or a sequence having an identical 9mer and 7mer sequence flanking a sequence that is at least 70% identical to the 22mer sequence of SEQ ID NO: 238.
7. The vertebrate or cell of claim 6, wherein the RSS and JH6*02 are provided as SEQ ID NO: 237.

8. The vertebrate or cell of any preceding claim, wherein the JH6*02 is the only JH6-type gene segment in the genome.
- 5 9. The vertebrate or cell of any preceding claim, wherein the JH6*02 is the closest JH gene segment to the constant region in the locus.
- 10 10. The vertebrate or cell of any preceding claim, wherein the locus comprises one, more or all human D gene segments D3-9; D4-17; D3-10; D2-2; D5-24; D6-19; D3-22; D6-13; D5-12; D1-26; D1-20; D5-18; D3-16; D2-21; D1-14; D7-27; D1-1; D6-25; D2-14; and D4-23.
11. The vertebrate or cell of claim 10, wherein the locus comprises one, more or all human D gene segments D3-9, D3-10, D6-19, D4-17, D6-13, D3-22, D2-2, D2-25 and D3-3.
- 15 12. The vertebrate or cell of any preceding claim, wherein the locus comprises a plurality of human D gene segments and the JH6*02 is in human germline configuration with respect to the 3'-most human D gene segment.
- 20 13. The vertebrate or cell of any preceding claim, wherein the locus comprises one, more or all of IGHV gene segments selected from V3-21, V3-13, V3-7, V6-1, V1-8, V1-2, V7-4-1, V1-3, V1-18, V4-4, V3-9, V3-23, V3-11 and V3-20.
- 25 14. The vertebrate or cell of any preceding claim, wherein the locus comprises one, more or all of human D3-9*01, D3-10*01, D6-19*01, D6-13*01, D1-26*01, IGHV1-8*01, IGHV4-61*01, IGHV6-1*01, IGHV4-4*02, IGHV1-3*01, IGHV3-66*03, IGHV3-7*01 and IGHV3-9*01.
- 30 15. An antibody-producing cell (eg, a B-cell) that is a progeny of the cell of any one of claims 3 to 14, wherein the antibody-producing cell comprises a heavy chain locus comprising a rearranged variable region produced by recombination of human JH6*02 with a D segment and a VH segment.
16. The cell of claim 15, which is a B-cell or hybridoma that expresses a target antigen-specific antibody comprising a heavy chain that comprises a rearranged variable region produced by recombination of human JH6*02 with a D segment and a VH segment.

17. The vertebrate or cell of any preceding claim, wherein the antibody heavy chain specifically binds a target antigen.
- 5 18. The vertebrate or cell of any preceding claim, wherein the antibody heavy chain has a HCDR3 length of at least 20 amino acids.
19. The vertebrate or cell of any preceding claim, wherein the antibody heavy chain is a product of the recombination of JH6*02 with a human VH gene segment recited in claim 13 or 14 and/or a D gene segment recited in claim 10, 11 or 14.
- 10 20. The vertebrate or cell of any preceding claim, wherein all endogenous non-human vertebrate heavy chain variable region gene segments have been inactivated in the genome.
- 15 21. The vertebrate or cell of any preceding claim, wherein the genome is homozygous for said heavy chain locus.
22. A heavy chain (eg, comprised by an antibody) isolated from a vertebrate of any one of claims 1, 2, 5 to 14 and 17 to 21 wherein the heavy chain comprises a HCDR3 of at least 20 amino acids.
- 20 23. The heavy chain of claim 22, wherein the HCDR3 is the product of recombination of human JH6*02 with a human VH gene segment recited in claim 13 or 14 and/or a D gene segment recited in claim 10, 11 or 14.
- 25 24. A heavy chain (eg, comprised by an antibody) whose VH variable domain is identical to the VH variable domain of the heavy chain of claim 22 or 23, and which comprises a human constant region or a human-mouse chimaeric constant region.
- 30 25. The heavy chain of claim 22, 23 or 24, whose VH variable domain is specific for a target antigen.
26. A method for producing a heavy chain, VH domain or an antibody specific to a target antigen, the method comprising immunizing a non-human vertebrate according to any one

- of claims 1, 2, 5 to 14 and 17 to 21 with the antigen and isolating the heavy chain, VH domain or an antibody specific to a target antigen or a cell producing the heavy chain, VH domain or an antibody, wherein the heavy chain, VH domain or an antibody comprises a HCDR3 that is derived from the recombination of human JH6*02 with a VH gene segment and a D gene segment.
- 5
27. A method for producing a human heavy chain or antibody comprising carrying out the method of claim 26, wherein the constant region of the locus is a non-human vertebrate (eg, mouse or rat) constant region, and then replacing the non-human constant region of the isolated heavy chain or antibody with a human constant region.
- 10
28. A heavy chain, VH domain or an antibody produced by the method of claim 26 or 27.
29. A B-cell or hybridoma expressing a heavy chain VH domain that is identical to the VH domain of the heavy chain of claim 22, 23 or 28.
- 15
30. A nucleic acid encoding the VH domain of the heavy chain of claim 22, 23 or 28, or encoding the heavy chain of claim 22, 23, 24, 25 or 28.
31. A vector (eg, a CHO cell or HEK293 cell vector) comprising the nucleic acid of claim 30; optionally wherein the vector is in a host cell (eg, a CHO cell or HEK293 cell).
- 20
32. A pharmaceutical composition comprising the antibody, heavy chain or VH domain (eg, comprised by an antibody) of any one of claims 22 to 25 and 28, together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, heavy chain or antibody).
- 25
33. The antibody, heavy chain or VH domain (eg, comprised by an antibody) of any one of claims 22 to 25 and 28 for use in medicine.
- 30
34. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 3 human variable region gene segments of the same type (eg, at least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or

at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.

- 5 35. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.
- 10 36. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and optionally
15 a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.
- 20 37. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human variable region gene segments, wherein the plurality comprises at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), a first of said different gene segments is provided in the genome of a first vertebrate of the population, and a second of said different gene segments being provided in the genome of a second vertebrate of the
25 population, wherein the genome of the first vertebrate does not comprise the second gene segment.
- 30 38. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
35
39. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 3 human variable region gene segments of the same type (eg, at

least 3 human VH6-1 gene segments, at least 3 human JH6 gene segments, at least 3 human Vk1-39 gene segments, at least 3 human D2-2 gene segments or at least 3 human Jk1 gene segments), wherein at least two of the human gene segments are variants that are not identical to each other.

5

40. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human
10 Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *cis* at the same Ig locus.

41. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different human variable region gene segments of the same type (eg, at least 2
15 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments) *trans* at the same Ig locus; and optionally a third human gene segment of the same type, wherein the third gene segment is *cis* with one of said 2 different gene segments.

20

42. A method of providing an enhanced human immunoglobulin variable region gene segment repertoire, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human variable region gene segments, wherein the method comprises providing at least 2 different human variable region gene segments of the
25 same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein a first of said different gene segments is provided in the genome of a first vertebrate of the population, and a second of said different gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of
30 the first vertebrate does not comprise the second gene segment.

43. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene
35 segments, at least 2 human Vk1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jk1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

44. The vertebrate, cell or method of any one of claims 34 to 43, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
- 5 45. The vertebrate, cell or method of any one of claims 34 to 44, wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
- 10 46. The vertebrate, cell or method of any one of claims 34 to 45, wherein the gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.
47. The vertebrate, cell or method of claim 46, wherein the individuals are not genetically related.
- 15 48. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human variable region gene segments of the same type (eg, at least 2 human VH6-1 gene segments, at least 2 human JH6 gene segments, at least 2 human Vκ1-39 gene segments, at least 2 human D2-2 gene segments or at least 2 human Jκ1 gene segments), wherein the gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same Ig locus in said genome.
- 20 49. The method of claim 48, wherein the different human individuals are from different human populations.
- 25 50. The method of claim 48, wherein the individuals are not genetically related.
- 30 51. The vertebrate, cell or method of preceding claim, wherein at least one of the different segments is a synthetic mutant of a human germline gene segment.

52. The vertebrate, cell or method of any one of claims 34 to 51, wherein each of said gene segments occurs in 10 or more different human populations.
53. The vertebrate, cell or method of preceding claim, wherein each of said gene segments has a human frequency of 5% or greater.
54. The vertebrate, cell or method of claim 53, wherein each of said gene segments occurs in 10 or more different human populations.
55. The vertebrate, cell or method of any one of claims 34 to 54, wherein each of said gene segments occurs in the 1000 Genomes database in more than 50 individuals.
56. The vertebrate, cell or method of preceding claim, wherein each of said gene segments (i) has a human frequency of 5% or greater; and (ii) occurs in 10 or more different human populations.
57. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) having a genome comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, IGHJ6 ref; SEQ ID NO: 244).
58. The vertebrate or cell of claim 57, wherein the genome comprises a third human gene segment of said type, the third gene segment being different from the first and second gene segments.
59. The vertebrate or cell of claim 57 or 58, wherein the first and second gene segments are *cis* on the same chromosome; and optionally the third gene segment is also *cis* on said chromosome.
60. The vertebrate or cell of claim 59, wherein the gene segments are targeted insertions into an endogenous non-human Ig locus.

61. The vertebrate or cell of claim 57 or 58, wherein the first and second gene segments are *trans* on different chromosomes.
- 5 62. The vertebrate or cell of any one of claims 57 to 61, wherein the first gene segment is a gene segment selected any one of Tables 1 to 7 and 9 to 14 and the second gene segment is the corresponding reference sequence.
- 10 63. A population of non-human vertebrates (eg, mice or rats) comprising first and second human Ig locus gene segments of the same type (eg, first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, SEQ ID NO: 244), wherein the first gene segment is provided in the genome of a first vertebrate of the population, and the second gene segment is provided in the genome of a second vertebrate of the population.
- 15
64. The population of claim 63, wherein the genome of the first vertebrate does not comprise the second gene segment.
- 20
65. The population of claim 63 or 64, wherein the population comprises a third human gene segment of said type, the third gene segment being different from the first and second gene segments and optionally wherein the first and third gene segments are present in the genome of the first vertebrate.
- 25
66. The population of claim 63, 64 or 65, wherein the gene segments are targeted insertions into an endogenous non-human Ig locus in the respective genome.
- 30 67. The population of any one of claims 63 to 66, wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 and the second gene segment is the corresponding reference sequence.
- 35 68. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising first and second human Ig locus gene segments of the same type (eg,

first and second human JH6 gene segments; or first and second IgG2 gene segments; or first and second human Jλ7 gene segments), wherein the first gene segment is a gene segment selected from any one of Tables 1 to 7 and 9 to 14 (eg, IGHJ6-a) and the second gene segment is the corresponding reference sequence (eg, SEQ ID NO: 244).

5

69. A method of providing an enhanced human immunoglobulin gene segment repertoire, the method comprising providing a population according to any one of claims 63 to 66.

10

70. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 34, having a genome comprising at least 3 human D gene segments of the same type (eg, D2-2 gene segments), wherein at least two of the human D gene segments are variants that are not identical to each other (eg, D2-2ref and D2-2a).

15

71. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 35, having a genome comprising at least 2 different non-endogenous D gene segments of the same type type (eg, D2-2ref and D2-2a) *cis* at the same Ig locus.

20

72. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 36, having a genome comprising at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a) *trans* at the same Ig locus; and optionally a third human D gene segment (eg, (eg, D2-2ref, D2-2a or D2-2b) of the same type, wherein the third D is *cis* with one of said 2 different D gene segments.

25

73. A population of non-human vertebrates (eg, mice or rats) according to claim 37, comprising a repertoire of human D gene segments, wherein the plurality comprises at least 2 different human D gene segments of the same type (eg, D2-2 gene segments), a first of said different D gene segments (eg, D2-2ref) is provided in the genome of a first vertebrate of the population, and a second of said different D gene segment (eg, D2-2a) being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not

30

comprise the second D gene segment.

35

74. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 38, having a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, human D2-2 gene segments), wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

75. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 39, the method comprising providing the vertebrate with

a genome comprising at least 3 human D gene segments of the same type (eg, D2-2 gene segments), wherein at least two of the human D gene segments are variants that are not identical to each other (eg, D2-2ref and D2-2a).

- 5 76. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 40, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, human D2-2 gene segments) *cis* at the same Ig locus.
- 10 77. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 41, the method comprising providing the vertebrate with a genome comprising at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a) *trans* at the same Ig locus; and optionally a third human D gene segment (eg, D2-2ref, D2-2a or D2-2b) of the same type, wherein the third D is *cis* with one of said 2 different D
15 gene segments.
78. A method of providing an enhanced human immunoglobulin D gene segment repertoire according to claim 42, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human D gene segments, wherein the method
20 comprises providing at least 2 different human D gene segments of the same type (eg, D2-2ref and D2-2a), wherein a first of said different D gene segments is provided in the genome of a first vertebrate of the population, and a second of said different D gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second D gene segment.
25
79. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 43, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous D gene segments of the same type (eg, D2-2ref and D2-2a), wherein the D gene segments are derived from the genome sequence
30 of different human individuals that are not genetically related over at least 3 generations.
80. The vertebrate or cell of claim 70, 71 or 74, or the method of claim 75, 77 or 79, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
- 35 81. The vertebrate or cell of claim 70, 71 or 71, or the method of any one of claims 75 to 78 and 80, wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
82. The vertebrate or cell of any one of claims 70 to 73, or the method of any one of claims 75 to 78
40 and 81, wherein the D gene segments are derived from the genome sequence of two or more

different human individuals; optionally wherein the different human individuals are from different human populations.

83. The vertebrate, cell or method of claim 82, wherein the individuals are not genetically related.

5

84. The vertebrate, cell or method of any one of claims 70 to 83, wherein at least one of the different D segments is a synthetic mutant of a human germline D gene segment.

85. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 48, the method comprising providing the vertebrate with a genome comprising at least 2 human D gene segments (eg, D2-2ref and D2-2a), wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgH locus in said genome.

10

15

86. The vertebrate or cell of any one of claims 70 to 74 and 80 to 84, wherein the genome comprises a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.

20

87. The population of claim 73, wherein the population comprises a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.

25

88. A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human D gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.

30

89. A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more variant human D gene segments, wherein said substantially complete functional repertoire and the supplementary D gene segments are not found together in the germline genome of a human individual.

35

90. The vertebrate or cell of claim 88 or the population of claim 89, comprising first and second D gene segments selected from

D2-2ref and D2-2a; or

D2-21ref and D2-21a; or

D3-10ref and D3-10a; or

D3-16ref and D3-16a; or

D2-8ref and D2-8a; or

D3-3ref and D3-3a; or

D4-23ref and D4-23a; or

D6-13ref and D6-13a; or

D3-9ref and D3-9a; or

D4-4ref and D4-4a; or

D7-27ref and D7-27a;

optionally wherein the first and/or second D gene segment is present in two or more copies.

91. The vertebrate, cell or population of claim 90, comprising human gene segments D2-2ref and D2-2a; and D3-3ref and D3-3a; and optionally also D2-15.

92. A non-human vertebrate or vertebrate cell according to claim 71, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene segments in *cis* at the same Ig locus in said genome;

a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene segments in *cis* at

the same Ig locus in said genome; and

a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene segments in *cis* at the same Ig locus in said genome;

5 optionally wherein the D gene segments are derived from the genome sequence of two or more different human individuals.

93. A non-human vertebrate or vertebrate cell according to claim 71, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

10 a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene segments in *trans* in said genome;

15 a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene segments in *trans* in said genome;

a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene segments in *trans* in said genome;

a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene segments in *trans* in said genome;

20 a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene segments in *trans* in said genome;

a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene segments in *trans* in said genome;

25 a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene segments in *trans* in said genome;

a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene segments in *trans* in said genome;

a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene segments in *trans* in said genome;

30 a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene segments in *trans* in said genome; and

a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene segments in *trans* in said genome;

35 optionally wherein the D gene segments are derived from the genome sequence of two or more different human individuals.

94. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell), according to claim 70, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the D gene repertoire comprising one or more of

a plurality of D2-2 gene segments provided by at least 3 different D2-2 gene segments;

- a plurality of D2-21 gene segments provided by at least 3 different D2-21 gene segments;
 a plurality of D3-10 gene segments provided by at least 3 different D3-10 gene segments;
 a plurality of D3-16 gene segments provided by at least 3 different D3-16 gene segments;
 a plurality of D2-8 gene segments provided by at least 3 different D2-8 gene segments;
 5 a plurality of D3-3 gene segments provided by at least 3 different D3-3 gene segments;
 a plurality of D4-23 gene segments provided by at least 3 different D4-23 gene segments;
 a plurality of D6-13 gene segments provided by at least 3 different D6-13 gene segments;
 a plurality of D3-9 gene segments provided by at least 3 different D3-9 gene segments;
 a plurality of D4-4 gene segments provided by at least 3 different D4-4 gene segments; and
 10 a plurality of D7-27 gene segments provided by at least 3 different D7-27 gene segments;
- optionally wherein the D gene segments are derived from the genome sequence of two or three different human individuals;
- 15 optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same Ig locus in said genome.
95. The vertebrate or cell of claim 92, 93 or 94, wherein the different human individuals are from different human populations.
 20
96. The vertebrate or cell of any one of claims 92 to 95, wherein the individuals are not genetically related.
97. The vertebrate or cell of any one of claims 92 to 96, wherein at least one of the different D
 25 segments is a synthetic mutant of a human germline D gene segment.
98. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to claim 74, comprising a genome comprising human VH, D and JH gene repertoires, the D gene repertoire comprising of one or more of
 30
- a plurality of D2-2 gene segments provided by at least 2 different D2-2 gene ; optionally in *cis* in said genome;
 a plurality of D2-21 gene segments provided by at least 2 different D2-21 gene ; optionally in *cis* in said genome;
 35 a plurality of D3-10 gene segments provided by at least 2 different D3-10 gene ; optionally in *cis* in said genome;
 a plurality of D3-16 gene segments provided by at least 2 different D3-16 gene ; optionally in *cis* in said genome;
 a plurality of D2-8 gene segments provided by at least 2 different D2-8 gene ; optionally in *cis* in
 40 said genome;
 a plurality of D3-3 gene segments provided by at least 2 different D3-3 gene ; optionally in *cis* in said genome;

a plurality of D4-23 gene segments provided by at least 2 different D4-23 gene ; optionally in *cis* in said genome;

a plurality of D6-13 gene segments provided by at least 2 different D6-13 gene ; optionally in *cis* in said genome;

5 a plurality of D3-9 gene segments provided by at least 2 different D3-9 gene ; optionally in *cis* in said genome;

a plurality of D4-4 gene segments provided by at least 2 different D4-4 gene ; optionally in *cis* in said genome; and

10 a plurality of D7-27 gene segments provided by at least 2 different D7-27 gene ; optionally in *cis* in said genome;

wherein the D gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

15 99. The vertebrate or cell of claim 98, wherein the human individuals are from different human populations.

20 100. The vertebrate, cell or population of any one of claims 70 to 99, wherein one or more of the D gene segments is a variant of a human germline D gene segment, wherein the variant gene segment encodes an amino acid sequence that differs by 1, 2 or 3 amino acids from the corresponding amino acid sequence encoded by the human germline D gene segment, provided in that said amino acid sequence encoded by the variant does not include a stop codon when said corresponding amino acid sequence does not include a stop codon.

25 101. The vertebrate, cell or population of claim 100, wherein said corresponding amino acid sequence encoded by the human germline D gene segment is a hydrophilic or hydrophobic sequence (according to J Mol Biol. 1997 Jul 25;270(4):587-97; Corbett SJ *et al*; Table 2).

30 102. The vertebrate, cell or population of claim 100 or 101, comprising said variant and said germline human D gene segments; optionally wherein the variant and germline human D gene segments are *cis* on the same chromosome.

35 103. The vertebrate, cell or population of any one of claims 100 to 102, wherein germline human D gene segment is a D2, D3, D5 or D6 family gene segment; optionally a D2-2, D2-15, D3-3, D3-9, D3-10, D3-22, D5-5, D5-18, D6-6, D6-13, D6-19 gene segment.

40 104. The vertebrate, cell or population of any one of claims 70 to 103, comprising a plurality of D2-2 gene segments, wherein the plurality comprises D2-2 gene segments that vary from each other at one or more nucleotide positions corresponding to positions

106,382,687 and

106,382,711

on human chromosome 14.

105. The vertebrate, cell or population of claim 104, wherein the plurality comprises a human D2-2 gene segment comprising a thymine at a position corresponding to position 106,382,687 on human chromosome 14; and optionally no further mutation from the sequence of D2-2ref.

106. The vertebrate, cell or population of claim 104 or 105, wherein the plurality comprises a human D2-2 gene segment comprising a cytosine at a position corresponding to position 106,382,687 on human chromosome 14; and optionally no further mutation from the sequence of D2-2a.

107. The vertebrate, cell or population of any one of claims 104 to 106, wherein the plurality comprises a human D2-2 gene segment comprising an adenine at a position corresponding to position 106,382,711 on human chromosome 14; and optionally no further mutation from the sequence of D2-2b.

108. The vertebrate, cell or population of any one of claims 104 to 107, wherein the plurality comprises a human D2-2 gene segment comprising an thymine at a position corresponding to position 106,382,711 on human chromosome 14; and optionally no further mutation from the sequence of D2-2ref.

109. The vertebrate, cell or population of any one of claims 70 to 108, comprising a plurality of D7-27 gene segments, wherein the plurality comprises D7-27 gene segments that vary from each other at a nucleotide position corresponding to position 106,331,767 on human chromosome 14.

110. The vertebrate, cell or population of claim 109, wherein the plurality comprises a human D7-27 gene segment comprising a cytosine at a position corresponding to position 106,331,767 on human chromosome 14; and optionally no further mutation from the sequence of D7-27ref.

111. The vertebrate, cell or population of claim 109 or 110, wherein the plurality comprises a human D7-27 gene segment comprising a guanine at a position corresponding to position 106,331,767 on human chromosome 14; and optionally no further mutation from the sequence of D7-27a.

112. The vertebrate, cell or population of any one of claims 70 to 111, comprising a plurality of D4-23 gene segments, wherein the plurality comprises D4-23 gene segments that vary from each other at a nucleotide position corresponding to position 106,350,740 on human chromosome 14.

113. The vertebrate, cell or population of claim 112, wherein the plurality comprises a human D4-23 gene segment comprising an adenine at a position corresponding to position 106,350,740 on human chromosome 14; and optionally no further mutation from the sequence of D4-23ref.

5

114. The vertebrate, cell or population of claim 112 or 113, wherein the plurality comprises a human D4-23 gene segment comprising an guanine at a position corresponding to position 106,350,740 on human chromosome 14; and optionally no further mutation from the sequence of D4-23a.

10

115. The vertebrate, cell or population of any one of claims 70 to 113, comprising a plurality of D2-21 gene segments, wherein the plurality comprises D2-21 gene segments that vary from each other at a nucleotide position corresponding to position 106,354,418 on human chromosome 14.

15

116. The vertebrate, cell or population of claim 115, wherein the plurality comprises a human D2-21 gene segment comprising an adenine at a position corresponding to position 106,354,418 on human chromosome 14; and optionally no further mutation from the sequence of D2-21ref.

20

117. The vertebrate, cell or population of claim 115 or 116, wherein the plurality comprises a human D2-21 gene segment comprising a guanine at a position corresponding to position 106,354,418 on human chromosome 14; and optionally no further mutation from the sequence of D2-21a.

25

118. The vertebrate, cell or population of any one of claims 70 to 117, comprising a plurality of D3-16 gene segments, wherein the plurality comprises D3-16 gene segments that vary from each other at a nucleotide position corresponding to position 106,354,418 on human chromosome 14.

30

119. The vertebrate, cell or population of claim 118, wherein the plurality comprises a human D3-16 gene segment comprising a thymine at a position corresponding to position 106,361,515 on human chromosome 14; and optionally no further mutation from the sequence of D3-16ref.

35

120. The vertebrate, cell or population of claim 118 or 119, wherein the plurality comprises a human D3-16 gene segment comprising a cytosine at a position corresponding to position 106,361,515 on human chromosome 14; and optionally no further mutation from the sequence of D3-16a.

121. The vertebrate, cell or population of any one of claims 70 to 120, comprising a plurality of D6-13 gene segments, wherein the plurality comprises D6-13 gene segments that vary from each other at a nucleotide position corresponding to position 106,367,013 on human chromosome 14.

5

122. The vertebrate, cell or population of claim 121, wherein the plurality comprises a human D6-13 gene segment comprising a thymine at a position corresponding to position 106,367,013 on human chromosome 14; and optionally no further mutation from the sequence of D6-13ref.

10 123. The vertebrate, cell or population of claim 121 or 122, wherein the plurality comprises a human D6-13 gene segment comprising a cytosine at a position corresponding to position 106,367,013 on human chromosome 14; and optionally no further mutation from the sequence of D6-13a.

15 124. The vertebrate, cell or population of any one of claims 70 to 123, comprising a plurality of D3-10 gene segments, wherein the plurality comprises D3-10 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,370,370 and 106,370,371
20 on human chromosome 14.

125. The vertebrate, cell or population of claim 124, wherein the plurality comprises a human D3-10 gene segment comprising a thymine at a position corresponding to position 106,370,370 on human chromosome 14; and optionally no further mutation from the sequence of D3-10ref.

25

126. The vertebrate, cell or population of claim 124 or 125, wherein the plurality comprises a human D3-10 gene segment comprising a cytosine at a position corresponding to position 106,370,370 on human chromosome 14; and optionally no further mutation from the sequence of D3-10a.

30

127. The vertebrate, cell or population of claim 124, 125 or 126 wherein the plurality comprises a human D3-10 gene segment comprising an adenine at a position corresponding to position 106,370,371 on human chromosome 14; and optionally no further mutation from the sequence of D3-10ref.

35

128. The vertebrate, cell or population of any one of claims 124 to 127, wherein the plurality comprises a human D3-10 gene segment comprising a guanine at a position corresponding to position 106,370,371 on human chromosome 14; and optionally no further mutation from the

sequence of D3-10b.

129. The vertebrate, cell or population of any one of claims 70 to 128, comprising a plurality of D3-9 gene segments, wherein the plurality comprises D3-9 gene segments that vary from each other at a nucleotide position corresponding to position 106,370,567 on human chromosome 14.

130. The vertebrate, cell or population of claim 129, wherein the plurality comprises a human D3-9 gene segment comprising an adenine at a position corresponding to position 106,370,567 on human chromosome 14; and optionally no further mutation from the sequence of D3-9ref.

131. The vertebrate, cell or population of claim 129 or 130, wherein the plurality comprises a human D3-9 gene segment comprising a thymine at a position corresponding to position 106,370,567 on human chromosome 14; and optionally no further mutation from the sequence of D3-9a.

132. The vertebrate, cell or population of any one of claims 70 to 131, comprising a plurality of D2-8 gene segments, wherein the plurality comprises D2-8 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,373,085; 106,373,086 and 106,373,089 on human chromosome 14.

133. The vertebrate, cell or population of claim 132, wherein the plurality comprises a human D2-8 gene segment comprising a cytosine at a position corresponding to position 106,373,085 on human chromosome 14.

134. The vertebrate, cell or population of claim 132 or 133, wherein the plurality comprises a human D2-8 gene segment comprising a thymine at a position corresponding to position 106,373,085 on human chromosome 14; and optionally no further mutation from the sequence of D2-8b.

135. The vertebrate, cell or population of claim 132, 133 or 134 wherein the plurality comprises a human D2-8 gene segment comprising a cytosine at a position corresponding to position 106,373,086 on human chromosome 14; and optionally no further mutation from the sequence of D2-8ref.

136. The vertebrate, cell or population of any one of claims 132 to 135, wherein the plurality comprises a human D2-8 gene segment comprising a thymine at a position corresponding to position 106,373,086 on human chromosome 14; and optionally no further mutation from the sequence of D2-8ref.

5

137. The vertebrate, cell or population of any one of claims 70 to 136, comprising a plurality of D4-4 gene segments, wherein the plurality comprises D4-4 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,379,086; and 106,379,089 on human chromosome 14.

10

138. The vertebrate, cell or population of claim 137, wherein the plurality comprises a D4-4 gene segment comprising a cytosine at a position corresponding to position 106,379,086 on human chromosome 14; and optionally no further mutation from the sequence of D4-4ref.

15

139. The vertebrate, cell or population of claim 137 or 138, wherein the plurality comprises a human D4-4 gene segment comprising a thymine at a position corresponding to position 106,379,086 on human chromosome 14; and optionally no further mutation from the sequence of D4-4a.

20

140. The vertebrate, cell or population of claim 137, 138 or 139 wherein the plurality comprises a human D4-4 gene segment comprising a cytosine at a position corresponding to position 106,379,089 on human chromosome 14; and optionally no further mutation from the sequence of D4-4ref or a cytosine at a position corresponding to position 106,379,086 on human chromosome 14.

25

141. The vertebrate, cell or population of any one of claims 137 to 140, wherein the plurality comprises a human D4-4 gene segment comprising a thymine at a position corresponding to position 106,373,089 on human chromosome 14; and optionally no further mutation from the sequence of D4-4a.

30

142. The vertebrate, cell or population of any one of claims 70 to 141, comprising a plurality of D3-3 gene segments, wherein the plurality comprises D3-3 gene segments that vary from each other at one or more nucleotide positions corresponding to positions 106,380,241; and 106,380,246 on human chromosome 14.

35

143. The vertebrate, cell or population of claim 142, wherein the plurality comprises a D3-3 gene segment comprising a thymine at a position corresponding to position 106,380,241 on human chromosome 14; and optionally no further mutation from the sequence of D3-3ref.
- 5 144. The vertebrate, cell or population of claim 142 or 143, wherein the plurality comprises a human D3-3 gene segment comprising a cytosine at a position corresponding to position 106,380,241 on human chromosome 14; and optionally no further mutation from the sequence of D3-3a.
- 10 145. The vertebrate, cell or population of claim 142, 143 or 144 wherein the plurality comprises a human D3-3 gene segment comprising an adenine at a position corresponding to position 106,380,246 on human chromosome 14; and optionally no further mutation from the sequence of D3-3ref.
- 15 146. The vertebrate, cell or population of any one of claims 142 to 145, wherein the plurality comprises a human D3-3 gene segment comprising a thymine at a position corresponding to position 106,380,246 on human chromosome 14; and optionally no further mutation from the sequence of D3-3a.
- 20 147. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 34, having a genome comprising at least 3 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein at least two of the human JH gene segments are variants that are not identical to each other.
- 25 148. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 35, having a genome comprising at least 2 different non-endogenous JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *cis* at the same Ig locus.
- 30 149. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 36, having a genome comprising at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *trans* at the same Ig locus; and optionally a third human JH gene segments of the same type, wherein the third JH is *cis* with one of said 2 different JH gene segments.
- 35 150. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human JH gene segments according to claim 37, wherein the plurality comprises at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), a first of said different JH gene segments is provided in the genome of a first vertebrate of the population, and
- 40 a second of said different JH gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JH gene segment.
- 45 151. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 38, having a genome comprising at least 2 different non-

endogenous JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

5 152. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 39, the method comprising providing the vertebrate with a genome comprising at least 3 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein at least two of the human JH gene segments are variants that are not identical to each other.

10 153. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 40, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *cis* at the same Ig locus.

15 154. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 41, the method comprising providing the vertebrate with a genome comprising at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6) *trans* at the same Ig locus; and optionally a third human JH gene segments of the same type, wherein the third JH is *cis* with one of said 2 different JH gene segments.

20 155. A method of providing an enhanced human immunoglobulin JH gene segment repertoire according to claim 42, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human JH gene segments, wherein the method comprises providing at least 2 different human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein a first of said different JH gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JH gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JH gene segment.

25 30 156. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 43, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

35 40 157. The vertebrate or cell of claim 147, 149 or 151, or the method of claim 152, 154, 155 or 156, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.

45 158. The vertebrate or cell of claim 147, 148 or 149, or the method of claim 152, 153, 154 or 155, wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

50 159. The vertebrate or cell of claim 147, 148 or 149, or the method of claim 152, 153, 154 or 155, wherein the JH gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.

160. The vertebrate, cell or method of claim 159, wherein the individuals are not genetically related.

161. The vertebrate, cell or method of any one of claims 147 to 160, wherein at least one of the different JH segments is a synthetic mutant of a human germline JH gene segment.

162. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human JH gene segments of the same type (JH1, JH2, JH3, JH4, JH5 or JH6), wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgH locus in said genome.

163. The vertebrate or cell of any one of claims 147 to 149 and 151, wherein the genome comprises a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

164. The population of claim 150, wherein the population comprises a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

165. A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

166. A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human JH gene segment types supplemented with one, two or more human JH gene segments, wherein said substantially complete functional repertoire and the supplementary JH gene segments are not found together in the germline genome of a human individual.

167. The vertebrate of claim 165 or the population of claim 166, wherein at least one of said JH gene segments is SEQ ID NO: 1, 2, 3 or 4.

168. The vertebrate or population of claim 167, wherein at least one of said JH gene segments is SEQ ID NO: 1 and at least one, two or more of said supplementary JH gene segments is a variant according to any one of claims 178 to 196.

169. The vertebrate or population of claim 167 or 168, wherein at least one of said JH gene segments is SEQ ID NO: 2 and at least one, two or more of said supplementary JH gene segments is a variant according to any one of claims 198 to 209.

170. The vertebrate or population of claim 167, 168 or 169, wherein at least one of said JH gene segments is SEQ ID NO: 2 and at least one, two or more of said supplementary JH gene segments

is a variant according to any one of claims 212 to 217.

171. A non-human vertebrate or vertebrate cell according to claim 148, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the JH gene repertoire comprising

a plurality of JH1 gene segments provided by at least 2 different JH1 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH2 gene segments provided by at least 2 different JH2 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH3 gene segments provided by at least 2 different JH3 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH4 gene segments provided by at least 2 different JH4 gene segments in *cis* at the same Ig locus in said genome;

a plurality of JH5 gene segments provided by at least 2 different JH5 gene segments in *cis* at the same Ig locus in said genome; or

a plurality of JH6 gene segments provided by at least 2 different JH6 gene segments in *cis* at the same Ig locus in said genome;

optionally wherein the JH gene segments are derived from the genome sequence of two or more different human individuals.

172. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell), according to claim 147, comprising a genome that comprises VH, D and JH gene repertoires comprising human gene segments, the JH gene repertoire comprising

a plurality of JH1 gene segments provided by at least 3 different JH1 gene segments;

a plurality of JH2 gene segments provided by at least 3 different JH2 gene segments;

a plurality of JH3 gene segments provided by at least 3 different JH3 gene segments;

a plurality of JH4 gene segments provided by at least 3 different JH4 gene segments;

a plurality of JH5 gene segments provided by at least 3 different JH5 gene segments; or

a plurality of JH6 gene segments provided by at least 3 different JH6 gene segments;

optionally wherein the JH gene segments are derived from the genome sequence of two or three different human individuals;

optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same Ig locus in said genome.

173. The vertebrate or cell of claim 171 or 172, wherein the different human individuals are from different human populations.

174. The vertebrate or cell of any one of claims 171 to 173, wherein the individuals are not genetically related.

175. The vertebrate or cell of any one of claims 171 to 174, wherein at least one of the different JH segments is a synthetic mutant of a human germline JH gene segment.

176. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to claim 151, comprising a genome comprising human VH, D and JH gene repertoires, the JH gene repertoire comprising

a plurality of JH1 gene segments provided by at least 2 different human JH1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of JH2 gene segments provided by at least 2 different human JH2 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of JH3 gene segments provided by at least 2 different human JH3 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of JH4 gene segments provided by at least 2 different human JH4 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of JH5 gene segments provided by at least 2 different human JH5 gene segments, optionally in *cis* at the same Ig locus in said genome; or

a plurality of JH6 gene segments provided by at least 2 different human JH6 gene segments, optionally in *cis* at the same Ig locus in said genome;

wherein the JH gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

177. The vertebrate or cell of claim 176, wherein the human individuals are from different human populations.

178. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 177, comprising a plurality of JH5 gene segments, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a nucleotide mutation at one or more positions corresponding to positions

106,330,024

106,330,027

106,330,032

106,330,041

106,330,044

106,330,045

106,330,062

106,330,063

106,330,065

106,330,066

106,330,067

106,330,068 and

106,330,071

on human chromosome 14.

179. The vertebrate, cell or population of claim 178, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,067 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

180. The vertebrate, cell or population of claim 179, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,071 on human chromosome 14 (optionally the additional mutation being a guanine); (ii) position 106,330,066 on human

chromosome 14 (optionally the additional mutation being a guanine); and/or (iii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

181. The vertebrate, cell or population of claim 178, 179 or 180, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,071 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

182. The vertebrate, cell or population of claim 181, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,063 on human chromosome 14 (optionally the additional mutation being an adenine); and/or (ii) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine).

183. The vertebrate, cell or population of claim 178 to 182, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,045 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

184. The vertebrate, cell or population of claim 178 to 183, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine at a position corresponding to position 106,330,044 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

185. The vertebrate, cell or population of claim 184, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,066 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

186. The vertebrate, cell or population of claim 178 to 185, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a guanine at a position corresponding to position 106,330,066 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

187. The vertebrate, cell or population of claim 186, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,068 on human chromosome 14 (optionally the additional mutation being a thymine).

188. The vertebrate, cell or population of claim 178 to 185, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,068 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

189. The vertebrate, cell or population of claim 188, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,330,067 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,330,066 on human chromosome 14 (optionally the additional mutation being a guanine).

190. The vertebrate, cell or population of claim 178 to 189, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,027 on human chromosome 14; and optionally no further

mutation from the sequence of SEQ ID NO: 1.

191. The vertebrate, cell or population of claim 178 to 190, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine at a position corresponding to position 106,330,024 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

192. The vertebrate, cell or population of claim 178 to 191, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,032 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

193. The vertebrate, cell or population of claim 178 to 192, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a thymine at a position corresponding to position 106,330,041 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

194. The vertebrate, cell or population of claim 178 to 193, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises an adenine or thymine at a position corresponding to position 106,330,063 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

195. The vertebrate, cell or population of claim 194, wherein the variant comprises additionally a mutation at a position corresponding to position 106,330,071 on human chromosome 14 (optionally the additional mutation being a guanine).

196. The vertebrate, cell or population of claim 178 to 195, wherein the plurality comprises a human JH5 gene variant of SEQ ID NO: 1, wherein the variant comprises a cytosine at a position corresponding to position 106,330,062 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 1.

197. The vertebrate, cell or population of claim 178 to 196, wherein the genome comprises SEQ ID NO:1; optionally in *cis* at the same Ig locus as one, two or more of the variants.

198. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 96 to 130, comprising a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a nucleotide mutation at one or more positions corresponding to positions

106,329,411
106,329,413
106,329,414
106,329,417
106,329,419
106,329,426
106,329,434
106,329,435, and
106,329,468

on human chromosome 14.

199. The vertebrate, cell or population of claim 198, comprising a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a guanine at a position corresponding to position 106,329,435 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

200. The vertebrate, cell or population of claim 199, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,329,468 on human chromosome 14 (optionally the additional mutation being a guanine); (ii) position 106,329,419 on human chromosome 14 (optionally the additional mutation being an adenine); (iii) position 106,329,434 on human chromosome 14 (optionally the additional mutation being a cytosine) and/or position 106,329,414 on human chromosome 14 (optionally the additional mutation being a guanine); (iv) position 106,329,426 on human chromosome 14 (optionally the additional mutation being an adenine); (v) position 106,329,413 on human chromosome 14 (optionally the additional mutation being an adenine); (vi) position 106,329,417 on human chromosome 14 (optionally the additional mutation being a thymine); (vii) position 106,329,411 on human chromosome 14 (optionally the additional mutation being a thymine); (viii) position 106,329,451 on human chromosome 14 (optionally the additional mutation being an adenine); (ix) position 106,329,452 on human chromosome 14 (optionally the additional mutation being a cytosine); and/or (x) position 106,329,453 on human chromosome 14 (optionally the additional mutation being a cytosine).

201. The vertebrate, cell or population of claim 200, wherein the variant comprises additionally mutations at positions corresponding to position 106,329,451 on human chromosome 14, the additional mutation being an adenine; position 106,329,452 on human chromosome 14, the additional mutation being a cytosine; and position 106,329,453 on human chromosome 14, the additional mutation being a cytosine.

202. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 201, comprising a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a guanine at a position corresponding to position 106,329,468 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

203. The vertebrate, cell or population of claim 202, wherein the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

204. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 203, comprising a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a thymine at a position corresponding to position 106,329,417 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

205. The vertebrate, cell or population of claim 204, wherein the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

206. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 205, comprising a plurality of JH6 gene segments,, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a cytosine at a position corresponding to position 106,329,434 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

207. The vertebrate, cell or population of claim 206, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,329,414 on human chromosome 14 (optionally the additional mutation being a guanine); and/or (ii) position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

208. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 207, comprising a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant of SEQ ID NO: 2, wherein the variant comprises a thymine at a position corresponding to position 106,329,411 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 2.

209. The vertebrate, cell or population of claim 208, wherein the variant comprises additionally a mutation at a position corresponding to position 106,329,435 on human chromosome 14 (optionally the additional mutation being a guanine).

210. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 209, comprising a plurality of JH6 gene segments, wherein the plurality comprises a human JH6 gene variant that is an antisense sequence of the variant of any one of claims 107 to 118.

211. The vertebrate, cell or population of any one of claims 198 to 210, wherein the genome comprises SEQ ID NO:2; optionally *cis* at the same Ig locus as one, two or more of the JH6 variants.

212. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 211, comprising a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises a nucleotide mutation at one or more positions corresponding to positions

106,331,455
106,331,453, and
106,331,409

on human chromosome 14.

213. The vertebrate, cell or population of claim 212, comprising said plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises a guanine at a position corresponding to position 106,331,455 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

214. The vertebrate, cell or population of claim 213, wherein the variant comprises additionally a mutation at a position corresponding to (i) position 106,331,453 on human chromosome 14 (optionally the additional mutation being an adenine); and/or (ii) position 106,331,409 on human chromosome 14 (optionally the additional mutation being an adenine); (iii) position 106,329,434 on human chromosome 14 (optionally the additional mutation being an adenine).

215. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 214, comprising a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises an adenine at a position corresponding to position 106,331,453 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

216. The vertebrate, cell or population of claim 215, wherein the variant comprises additionally a mutation at a position corresponding to position 106,331,409 on human chromosome 14 (optionally the additional mutation being an adenine).

217. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 216, comprising a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant of SEQ ID NO: 3, wherein the variant comprises an adenine at a position corresponding to position 106,331,409 on human chromosome 14; and optionally no further mutation from the sequence of SEQ ID NO: 3.

218. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 217, comprising a plurality of JH2 gene segments, wherein the plurality comprises a human JH2 gene variant that is an antisense sequence of the variant of any one of claims 188 to 193.

219. The vertebrate, cell or population of any one of claims 212 to 218, wherein the genome comprises SEQ ID NO:3; optionally *cis* at the same Ig locus as one, two or more of the JH2 variants.

220. The vertebrate, cell or population of any one of claims 147 to 151, 157 to 161 and 163 to 219, wherein the genome comprises two or more different JH gene segments selected from SEQ ID NOs: 1 to 3 and variants according to any one of claims 178 to 219; optionally wherein said JH gene segments are *cis* at the same immunoglobulin Ig locus.

221. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 34, having a genome comprising at least 3 human JL gene segments of the same type (eg, Jk1), wherein at least two of the human JL gene segments are variants that are not identical to each other.

222. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 35, having a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1) *cis* at the same Ig locus.

223. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 36, having a genome comprising at least 2 different human JL gene segments of the same type (eg, Jk1) *trans* at the same Ig locus; and optionally a third human JL gene segment of the same type, wherein the third JL is *cis* with one of said 2 different JL gene segments.

224. A population of non-human vertebrates (eg, mice or rats) comprising a repertoire of human JL gene segments according to claim 37, wherein the plurality comprises at least 2 different human JL gene segments of the same type (eg, Jk1), a first of said different JL gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JL gene segments being provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JL gene segment.
225. A non-human vertebrate (eg, a mouse or rat) or a non-human vertebrate cell (eg, an ES cell or a B-cell) according to claim 38, having a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
226. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 39, the method comprising providing the vertebrate with a genome comprising at least 3 human JL gene segments of the same type (eg, Jk1), wherein at least two of the human JL gene segments are variants that are not identical to each other.
227. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 40, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1) *cis* at the same Ig locus.
228. A method of enhancing the immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 41, the method comprising providing the vertebrate with a genome comprising at least 2 different human JL gene segments of the same type (eg, Jk1) *trans* at the same Ig locus; and optionally a third human JL gene segment of the same type, wherein the third JL is *cis* with one of said 2 different JL gene segments.
229. A method of providing an enhanced human immunoglobulin JL gene segment repertoire according to claim 42, the method comprising providing a population of non-human vertebrates (eg, a mouse or rat) comprising a repertoire of human JL gene segments, wherein the method comprises providing at least 2 different human JL gene segments of the same type (eg, Jk1), wherein a first of said different JL gene segments is provided in the genome of a first vertebrate of the population, and a second of said different JL gene segments is provided in the genome of a second vertebrate of the population, wherein the genome of the first vertebrate does not comprise the second JL gene segment.
230. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat) according to claim 43, the method comprising providing the vertebrate with a genome comprising at least 2 different non-endogenous JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.
231. The vertebrate or cell of claim 221, 223 or 225, or the method of claim 226, 228, 229 or 230, wherein at least 2 or 3 of said different gene segments are provided *cis* at the same Ig locus in said genome.
232. The vertebrate or cell of claim 221, 222 or 223, or the method of claim 226, 227, 228 or 229, wherein the JL gene segments are derived from the genome sequence of different human

individuals that are not genetically related over at least 3 generations.

233. The vertebrate or cell of claim 221, 222 or 223, or the method of claim 226, 227, 228 or 229, wherein the JL gene segments are derived from the genome sequence of two or more different human individuals; optionally wherein the different human individuals are from different human populations.

234. The vertebrate, cell or method of claim 233, wherein the individuals are not genetically related.

235. The vertebrate, cell or method of any one of claims 221 to 234, wherein at least one of the different JL segments is a synthetic mutant of a human germline JL gene segment.

236. A method of enhancing the human immunoglobulin gene diversity of a non-human vertebrate (eg, a mouse or rat), the method comprising providing the vertebrate with a genome comprising at least 2 human JL gene segments of the same type (eg, Jk1), wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations; optionally wherein at least 2 or 3 of said different gene segments are provided at the same IgL locus in said genome.

237. The vertebrate or cell of any one of claims claim 221 to 223 and 225, wherein the genome comprises a substantially complete functional repertoire of human Jk and/or Jλ gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.

238. The population of claim 224, wherein the population comprises a substantially complete functional repertoire of human JL gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.

239. A non-human vertebrate (eg, a mouse or rat) or a non-human cell (eg, an ES cell or a B-cell) having a genome comprising a substantially complete functional repertoire of human Jk and/or Jλ gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.

240. A population of non-human vertebrates (eg, mice or rats) comprising a substantially complete functional repertoire of human Jk and/or Jλ gene segment types supplemented with one, two or more human Jk and/or Jλ gene segments respectively, wherein said substantially complete functional repertoire and the supplementary gene segments are not found together in the germline genome of a human individual.

241. A non-human vertebrate or vertebrate cell according to claim 222, comprising a genome that comprises VL and JL gene repertoires comprising human gene segments, the JL gene repertoire comprising

a plurality of human Jk1 gene segments provided by at least 2 different human Jk1 gene segments in *cis* at the same Ig locus in said genome;
a plurality of human Jk2 gene segments provided by at least 2 different human Jk1 gene

segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jk3 gene segments provided by at least 2 different human Jk1 gene
 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jk4 gene segments provided by at least 2 different human Jk1 gene
 5 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jk5 gene segments provided by at least 2 different human Jk1 gene
 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jλ1 gene segments provided by at least 2 different human Jλ1 gene
 segments in *cis* at the same Ig locus in said genome;
 10 a plurality of human Jλ2 gene segments provided by at least 2 different human Jλ2 gene
 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jλ3 gene segments provided by at least 2 different human Jλ3 gene
 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jλ4 gene segments provided by at least 2 different human Jλ4 gene
 15 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jλ5 gene segments provided by at least 2 different human Jλ5 gene
 segments in *cis* at the same Ig locus in said genome;
 a plurality of human Jλ6 gene segments provided by at least 2 different human Jλ6 gene
 segments in *cis* at the same Ig locus in said genome; or
 20 a plurality of human Jλ7 gene segments provided by at least 2 different human Jλ7 gene
 segments in *cis* at the same Ig locus in said genome;

optionally wherein the JL gene segments are derived from the genome sequence of two or more
 different human individuals.

242. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell), according to claim
 221, comprising a genome that comprises VL and JL gene repertoires comprising human gene
 segments, the JL gene repertoire comprising

a plurality of human Jk1 gene segments provided by at least 3 different human Jk1 gene
 segments;
 a plurality of human Jk2 gene segments provided by at least 3 different human Jk1 gene
 segments;
 a plurality of human Jk3 gene segments provided by at least 3 different human Jk1 gene
 35 segments;
 a plurality of human Jk4 gene segments provided by at least 3 different human Jk1 gene
 segments;
 a plurality of human Jk5 gene segments provided by at least 3 different human Jk1 gene
 segments;
 40 a plurality of human Jλ1 gene segments provided by at least 3 different human Jλ1 gene
 segments;
 a plurality of human Jλ2 gene segments provided by at least 3 different human Jλ2 gene
 segments;
 a plurality of human Jλ3 gene segments provided by at least 3 different human Jλ3 gene
 45 segments;
 a plurality of human Jλ4 gene segments provided by at least 3 different human Jλ4 gene
 segments;
 a plurality of human Jλ5 gene segments provided by at least 3 different human Jλ5 gene
 segments;
 50 a plurality of human Jλ6 gene segments provided by at least 3 different human Jλ6 gene
 segments; or

a plurality of human J λ 7 gene segments provided by at least 3 different human J λ 7 gene segments;

optionally wherein the JL gene segments are derived from the genome sequence of two or three different human individuals;

optionally wherein at least 2 or 3 of said different gene segments are provided in *cis* at the same Ig locus in said genome.

243. The vertebrate or cell of claim 241 or 242, wherein the different human individuals are from different human populations.

244. The vertebrate or cell of any one of claims 241 to 243, wherein the individuals are not genetically related.

245. The vertebrate or cell of any one of claims 241 to 244, wherein at least one of the different JL segments is a synthetic mutant of a human germline JL gene segment.

246. A non-human vertebrate or vertebrate cell (optionally an ES cell or B-cell) according to claim 225, comprising a genome comprising human VL and JL gene repertoires, the JL gene repertoire comprising

a plurality of human J κ 1 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 2 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 3 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 4 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J κ 5 gene segments provided by at least 2 different human J κ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 1 gene segments provided by at least 2 different human J λ 1 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 2 gene segments provided by at least 2 different human J λ 2 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 3 gene segments provided by at least 2 different human J λ 3 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 4 gene segments provided by at least 2 different human J λ 4 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 5 gene segments provided by at least 2 different human J λ 5 gene segments, optionally in *cis* at the same Ig locus in said genome;

a plurality of human J λ 6 gene segments provided by at least 2 different human J λ 6 gene segments, optionally in *cis* at the same Ig locus in said genome; or

a plurality of human J λ 7 gene segments provided by at least 2 different human J λ 7 gene segments, optionally in *cis* at the same Ig locus in said genome;

wherein the JL gene segments are derived from the genome sequence of different human individuals that are not genetically related over at least 3 generations.

247. The vertebrate or cell of claim 246, wherein the human individuals are from different human populations.

248. A method of producing an antibody heavy chain, the method comprising

- (a) providing an antigen-specific heavy chain variable domain; and
- (b) combining the variable domain with a human heavy chain constant region to produce an antibody heavy chain comprising (in N- to C-terminal direction) the variable domain and the constant region;

wherein

the human heavy chain constant region is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region.

249. The method of claim 248, wherein the variable domain is a human variable domain.

250. The method of any claim 248 or 249, wherein the variable domain has previously been selected from a non-human vertebrate that has been immunised with the antigen.

251. The method of any one of claims 248 to 250, comprising providing an expression vector comprising a nucleotide sequence encoding the constant region; inserting a nucleotide sequence encoding the variable domain into the vector 5' of the constant region sequence; inserting the vector into a host cell and expressing the heavy chain by the host cell; the method further comprising isolating a heavy chain (eg, as part of an antibody) comprising the variable domain and the human constant region.

252. The method of any one of claims 248 to 251, further comprising obtaining a nucleotide sequence encoding the heavy chain.

253. An antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region.

254. A polypeptide comprising (in N- to C- terminal direction) a leader sequence, a human variable domain that is specific for an antigen and a human constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; wherein (i) the leader sequence is not the native human variable domain leader sequence; and/or (ii) the variable domain comprises mouse AID-pattern somatic mutations or mouse terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations.

255. A nucleotide sequence encoding (in 5' to 3' direction) a leader sequence and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; and the leader sequence being operable for expression of the heavy chain and wherein the leader sequence is not the native human variable domain leader sequence.
256. A nucleotide sequence encoding (in 5' to 3' direction) a promoter and a human antibody heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region that is an IGHAref, IGHA1a, IGHA2a, IGHA2b, IGHG1ref, IGHG2ref, IGHG2a, IGHG3ref, IGHG3a, IGHG3b, IGHG4ref, IGHG4a, IGHDref, IGHEref, IGHMref, IGHMa or IGHMb constant region; and the promoter being operable for expression of the heavy chain and wherein the promoter is not the native human promoter.
257. The antibody, polypeptide or nucleotide sequence of any one of claims 253 to 256, wherein the variable domain comprises mouse AID-pattern somatic mutations or mouse terminal deoxynucleotidyl transferase (TdT)- pattern junctional mutations.
258. A vector (eg, a CHO cell or HEK293 cell vector) comprising the nucleic acid of claim 255, 256 or 257; optionally wherein the vector is in a host cell (eg, a CHO cell or HEK293 cell).
259. A pharmaceutical composition comprising the antibody or polypeptide of any one of claims 253, 254 and 257, together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).
260. The antibody or polypeptide of any one of claims 253, 254 and 257 for use in treating and/or preventing a medical condition in a human patient.
261. Use of the antibody or polypeptide of any one of claims 253, 254 and 257 for the manufacture of a medicament for treating and/or preventing a medical condition in a human patient.
262. The antibody, polypeptide or use of claim 260 or 261, wherein the human is a member of a human population selected from population numbers 1-14, wherein the populations are numbered as follows (population labels being according to 1000 Genomes Project nomenclature)
- 1= ASW;
 - 2= CEU;
 - 3=CHB;
 - 4=CHS;
 - 5=CLM;

6=FIN;
 7=GBR;
 8=IBS;
 9=JPT;
 10=LWK;
 11=MXL;
 12=PUR;
 13=TSI;
 14=YRI.

263. The antibody, polypeptide or use of claim 262, wherein the constant region is a

- (i) IGHA1a constant region and the human population is selected from any population number 1-14;
- (ii) IGHA2a constant region and the human population is selected from any population number 1-14;
- (iii) IGHA2b constant region and the human population is selected from any population number 1-14;
- (iv) IGHG2a constant region and the human population is selected from any population number 1-9 and 11-13;
- (v) IGHG3a constant region and the human population is selected from any population number 1-14;
- (vi) IGHG3b constant region and the human population is selected from any population number 1-8 and 11-13;
- (vii) IGHG4a constant region and the human population is selected from any population number 1-9 and 11-13;
- (viii) IGHMa constant region and the human population is selected from any population number 1-14; or
- (ix) IGHMb constant region and the human population is selected from any population number 1-14;

Wherein the populations are numbered as follows (population labels being according to 1000 Genomes Project nomenclature)

1= ASW;
 2= CEU;
 3=CHB;
 4=CHS;
 5=CLM;
 6=FIN;
 7=GBR;
 8=IBS;
 9=JPT;
 10=LWK;
 11=MXL;
 12=PUR;

13=TSI;

14=YRI.

265. A vector (eg, a CHO cell or HEK293 cell vector) comprising a IGHG1ref, IGHG2ref, IGHG2a,
 5 IGHG3ref, IGHG3a, IGHG3b, IGHG4ref or IGHG4a constant region nucleotide sequence that is 3'
 of a cloning site for the insertion of a human antibody heavy chain variable domain nucleotide
 sequence, such that upon insertion of such a variable domain sequence the vector comprises (in
 5' to 3' direction) a promoter, a leader sequence, the variable domain sequence and the
 constant region sequence so that the vector is capable of expressing a human antibody heavy
 10 chain when present in a host cell.

A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell)
 comprising a genome having a superhuman immunoglobulin heavy chain human VH and/or D
 and/or J gene repertoire.

266. The vertebrate or cell of claim 265, wherein the genome comprises a transgenic
 immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene
 segments, one or more human D gene segments and one or more human J gene segments,
 wherein the plurality of VH gene segments consists of more than the natural human repertoire
 20 of functional VH gene segments; optionally wherein the genome is homozygous for said
 transgenic heavy chain locus.

267. The vertebrate or cell of claim 266, wherein VH gene repertoire consists of a plurality of VH
 gene segments derived from the genome sequence of a first human individual, supplemented
 25 with one or more different VH gene segments derived from the genome sequence of a second,
 different human individual; optionally wherein the individuals are not related; optionally
 wherein the D and J segments are derived from the genome sequence of the first human
 individual; and optionally wherein the VH gene segments from the genome sequence of the
 second individual are selected from the VH gene segments listed in Table 1.

268. The vertebrate or cell of claim 267, wherein the transgenic locus comprises more than 26
 functional human VH gene segments; optionally wherein the locus comprises at least
 27 or 28 functional human VH gene segments (eg, wherein the locus comprises the full functional
 VH repertoire of said first individual supplemented with one or more VH gene segments derived
 35 from the genome sequence of the second human individual and optionally with one or more VH
 gene segments derived from the genome sequence of a third human individual).

269. The vertebrate or cell of claim 267 or 268, wherein the transgenic locus comprises a first VH
 gene segment derived from the genome sequence of the first individual and a second VH gene
 40 segment derived from the genome sequence of the second individual, wherein the second VH
 gene segment is a polymorphic variant of the first VH gene segment; optionally wherein the
 locus comprises a further polymorphic variant of the first VH gene segment (eg, a variant derived
 from the genome sequence of a third human individual).

270. The vertebrate or cell of any one of claims 265 to 269, wherein the genome comprises a (or said) transgenic immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene segments, a plurality of human D gene segments and one or more human J gene segments, wherein the plurality of D gene segments consists of more than the natural human repertoire of functional D gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

271. The vertebrate or cell of claim 270, wherein D gene repertoire consists of a plurality of D gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more different D gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein J segments are derived from the genome sequence of the first human individual; and optionally wherein the D gene segments from the genome sequence of the second individual are selected from the D gene segments listed in Table 2.

272. The vertebrate or cell of claim 271, wherein the transgenic locus comprises more than 26 functional human D gene segments; optionally wherein the locus comprises at least 27 or 28 functional human D gene segments (eg, wherein the locus comprises the full functional D repertoire of said first individual supplemented with one or more D gene segments derived from the genome sequence of the second human individual and optionally with one or more D gene segments derived from the genome sequence of a third human individual).

273. The vertebrate or cell of claim 271 or 272, wherein the transgenic locus comprises a first D gene segment derived from the genome sequence of the first individual and a second D gene segment derived from the genome sequence of the second individual, wherein the second D gene segment is a polymorphic variant of the first D gene segment; optionally wherein the locus comprises a further polymorphic variant of the first D gene segment (eg, a variant derived from the genome sequence of a third human individual).

274. The vertebrate or cell of any one of claims 265 to 273, wherein the genome comprises a (or said) transgenic immunoglobulin heavy chain locus comprising a plurality of human immunoglobulin VH gene segments, one or more human D gene segments and a plurality of human JH gene segments, wherein the plurality of J gene segments consists of more than the natural human repertoire of functional J gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

275. The vertebrate or cell of claim 274, wherein the J gene repertoire consists of a plurality of J gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more different J gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein D segments are derived from the genome sequence of the first human individual; and optionally wherein the J gene segments from the genome sequence of the second individual are selected from the J gene segments listed in Table 3.

276. The vertebrate or cell of claim 275, wherein the transgenic locus comprises more than 26 functional human J gene segments; optionally wherein the locus comprises at least 27 or 28 functional human J gene segments (eg, wherein the locus comprises the full functional J repertoire of said first individual supplemented with one or more J gene segments derived from the genome sequence of the second human individual and optionally with one or more J gene segments derived from the genome sequence of a third human individual).

277. The vertebrate or cell of claim 275 or 276, wherein the transgenic locus comprises a first J gene segment derived from the genome sequence of the first individual and a second J gene segment derived from the genome sequence of the second individual, wherein the second J gene segment is a polymorphic variant of the first J gene segment; optionally wherein the locus comprises a further polymorphic variant of the first J gene segment (eg, a variant derived from the genome sequence of a third human individual).

278. A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) comprising a genome having a superhuman immunoglobulin light chain human VL gene repertoire; optionally wherein the vertebrate or cell is according to any one of claims 265 to 277.

279. The vertebrate or cell of claim 278, wherein the genome comprises

(i) a transgenic immunoglobulin kappa light chain locus comprising a plurality of human immunoglobulin Vk gene segments and one or more human J gene segments, wherein the plurality of Vk gene segments consists of more than the natural human repertoire of functional Vk gene segments; optionally wherein the genome is homozygous for said transgenic kappa light chain locus; and/or

(ii) a transgenic immunoglobulin lambda light chain locus comprising a plurality of human immunoglobulin Vλ gene segments and one or more human J gene segments, wherein the plurality of Vλ gene segments consists of more than the natural human repertoire of functional Vλ gene segments; optionally wherein the genome is homozygous for said transgenic lambda light chain locus.

280. The vertebrate or cell of claim 279, wherein

(i) the Vk gene repertoire consists of a plurality of Vk gene segments derived from the genome sequence of a first human individual, supplemented with one or more Vk gene segments derived from the genome sequence of a second, different human individual; optionally wherein the individuals are not related; optionally wherein the J segments are derived from the genome sequence of the first human individual; and optionally wherein the Vk gene segments from the genome sequence of the second individual are selected from the Vk gene segments listed in Table 4; and

(i) the Vλ gene repertoire consists of a plurality of Vλ gene segments derived from the genome sequence of a first human individual, supplemented with one or more Vλ gene segments derived from the genome sequence of a second, different human individual; optionally wherein the

individuals are not related; optionally wherein the J segments are derived from the genome sequence of the first human individual; and optionally wherein the V λ gene segments from the genome sequence of the second individual are selected from the V λ gene segments listed in Table 5.

5

281. The vertebrate or cell of claim 280, wherein

the kappa light transgenic locus comprises more than 26 functional human V κ gene segments; optionally wherein the locus comprises at least 27 or 28 functional human V κ gene segments (eg, wherein the locus comprises the full functional V κ repertoire of said first individual supplemented with one or more V κ gene segments derived from the genome sequence of the second human individual and optionally with one or more V κ gene segments derived from the genome sequence of a third human individual); and

10

the lambda light transgenic locus comprises more than 26 functional human V λ gene segments; optionally wherein the locus comprises at least 27 or 28 functional human V λ gene segments (eg, wherein the locus comprises the full functional V λ repertoire of said first individual supplemented with one or more V λ gene segments derived from the genome sequence of the second human individual and optionally with one or more V λ gene segments derived from the genome sequence of a third human individual);

15

20

282. The vertebrate or cell of claim 280 or 281, wherein

the kappa light transgenic locus comprises a first V κ gene segment derived from the genome sequence of the first individual and a second V κ gene segment derived from the genome sequence of the second individual, wherein the second V κ gene segment is a polymorphic variant of the first V κ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first V κ gene segment (eg, a variant derived from the genome sequence of a third human individual); and

25

30

the lambda light transgenic locus comprises a first V λ gene segment derived from the genome sequence of the first individual and a second V λ gene segment derived from the genome sequence of the second individual, wherein the second V λ gene segment is a polymorphic variant of the first V λ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first V λ gene segment (eg, a variant derived from the genome sequence of a third human individual).

35

283. The vertebrate or cell of any one of claims 278 to 282, wherein the genome comprises a (or said) transgenic immunoglobulin light chain locus comprising a plurality of human immunoglobulin VL gene segments and a plurality of human JL gene segments, wherein the plurality of J gene segments consists of more than the natural human repertoire of functional J gene segments; optionally wherein the genome is homozygous for said transgenic heavy chain locus.

40

284. The vertebrate or cell of claim 283, wherein

(i) the Jk gene repertoire consists of a plurality of Jk gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more Jk gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein the Vk segments are derived from the genome sequence of the first human individual; optionally wherein the Jk gene segments from the genome sequence of the second individual are selected from the Jk gene segments listed in Table 6; and

(ii) the Jk gene repertoire consists of a plurality of Jλ gene segments derived from the genome sequence of a (or said) first human individual, supplemented with one or more Jλ gene segments derived from the genome sequence of a (or said) second, different human individual; optionally wherein the individuals are not related; optionally wherein the Vλ segments are derived from the genome sequence of the first human individual; optionally wherein the Jλ gene segments from the genome sequence of the second individual are selected from the Jλ gene segments listed in Table 7.

285. The vertebrate or cell of claim 284, wherein

(i) the transgenic light chain locus comprises more than 26 functional human Jk gene segments; optionally wherein the locus comprises at least 27 or 28 functional human Jk gene segments (eg, wherein the locus comprises the full functional Jk repertoire of said first individual supplemented with one or more Jk gene segments derived from the genome sequence of the second human individual and optionally with one or more Jk gene segments derived from the genome sequence of a third human individual); and/or

(i) the transgenic light chain locus comprises more than 26 functional human Jλ gene segments; optionally wherein the locus comprises at least 27 or 28 functional human Jλ gene segments (eg, wherein the locus comprises the full functional Jλ repertoire of said first individual supplemented with one or more Jλ gene segments derived from the genome sequence of the second human individual and optionally with one or more Jλ gene segments derived from the genome sequence of a third human individual).

286. The vertebrate or cell of claim 284 or 285, wherein

(i) the kappa light transgenic locus comprises a first Jk gene segment derived from the genome sequence of the first individual and a second Jk gene segment derived from the genome sequence of the second individual, wherein the second Jk gene segment is a polymorphic variant of the first Jk gene segment; optionally wherein the locus comprises a further polymorphic variant of the first Jk gene segment (eg, a variant derived from the genome sequence of a third human individual); and

(ii) the lambda light transgenic locus comprises a first Jλ gene segment derived from the genome sequence of the first individual and a second Jλ gene segment derived from the genome

sequence of the second individual, wherein the second Jk gene segment is a polymorphic variant of the first Jλ gene segment; optionally wherein the locus comprises a further polymorphic variant of the first Jλ gene segment (eg, a variant derived from the genome sequence of a third human individual).

5

287. A library of antibody-producing transgenic cells whose genomes collectively encode a repertoire of antibodies, wherein

10

(a) a first transgenic cell expresses a first antibody having a chain encoded by a first immunoglobulin gene, the gene comprising a first variable domain nucleotide sequence produced following recombination of a first human unrearranged immunoglobulin gene segment;

15

(b) a second transgenic cell expresses a second antibody having a chain encoded by a second immunoglobulin gene, the second gene comprising a second variable domain nucleotide sequence produced following recombination of a second human unrearranged immunoglobulin gene segment, the first and second antibodies being non-identical;

20

(c) the first and second gene segments are different and derived from the genome sequences of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

(d) wherein the cells are non-human vertebrate (eg, mouse or rat) cells.

25

288. The library of claim 23, wherein in step (c) the individuals are not related.

30

289. The vertebrate, cell or library of any one of claims 265 to 288, wherein the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different; optionally wherein the ethnic populations are selected from those identified in the 1000 genomes database.

35

290. The vertebrate, cell or library of claim 289, wherein the human immunoglobulin gene segment derived from the genome sequence of the second individual is low-frequency (optionally rare) within the second ethnic population.

40

291. The library of claim 289 or 290, wherein the first variable region nucleotide sequence is produced by recombination of the first human immunoglobulin gene segment with a first J gene segment and optionally a first D gene segment, wherein the first human immunoglobulin gene segment is a V gene segment and the V, D and J segments are derived from the first human population, optionally from the genome of one individual of the first human population.

45

292. The library of claim 291, wherein the second variable region nucleotide sequence is produced by recombination of the second human immunoglobulin gene segment with a second J gene segment and optionally a second D gene segment, wherein the second human immunoglobulin gene segment is a V gene segment derived from the second population and the

D and/or J segments are derived from the first human population, optionally the D and J gene segments being from the genome of one individual of the first human population.

293. The library of claim 292, wherein all of the D and J segments that have been recombined with the first and second V gene segments are D and J segments derived from the first human population, optionally the D and J gene segments being from the genome of one individual of the first human population.

294. The vertebrate, cell or library of any one of claims 289 to 293, wherein the first and second ethnic populations are selected from the group consisting of an ethnic population with European ancestry, an ethnic population with East Asian, an ethnic population with West African ancestry, an ethnic population with Americas ancestry and an ethnic population with South Asian ancestry.

295. The vertebrate, cell or library of claim 294, wherein the first and second ethnic populations are selected from the group consisting of an ethnic population with Northern European ancestry; or an ethnic population with Western European ancestry; or an ethnic population with Toscani ancestry; or an ethnic population with British ancestry; or an ethnic population with Icelandic ancestry; or an ethnic population with Finnish ancestry; or an ethnic population with Iberian ancestry; or an ethnic population with Japanese ancestry; or an ethnic population with Chinese ancestry; or an ethnic population Vietnamese ancestry; or an ethnic population with Yoruba ancestry; or an ethnic population with Luhya ancestry; or an ethnic population with Gambian ancestry; or an ethnic population with Malawian ancestry; or an ethnic population with Native American ancestry; or an ethnic population with Afro-Caribbean ancestry; or an ethnic population with Mexican ancestry; or an ethnic population with Puerto Rican ancestry; or an ethnic population with Columbian ancestry; or an ethnic population with Peruvian ancestry; or an ethnic population with Ahom ancestry; or an ethnic population with Kayadtha ancestry; or an ethnic population with Reddy ancestry; or an ethnic population with Maratha; or an ethnic population with Punjabi ancestry.

296. The library of claim 287, wherein the second human immunoglobulin gene segment is a polymorphic variant of the first human immunoglobulin gene segment; optionally wherein the second gene segment is selected from the group consisting of a gene segment in any of Tables 1 to 7 and 9 to 14, eg, the second gene segment is a polymorphic variant of VH1-69.

297. The library of any one of claims 287 to 296, wherein the first and second human immunoglobulin gene segments are both (i) V_H gene segments; (ii) D segments; (iii) J segments (optionally both J_H segments, both J_K segments or both J_λ segments); (iv) constant regions (optionally both a gamma constant region, optionally both a C gamma-1 constant region); (v) CH1 regions; (vi) CH2 regions; or (vii) CH3 regions.

298. The library of any one of claims 287 to 297, wherein the library is naive and optionally has a library size of from 10^2 to 10^9 cells.

299. The library of any one of claims 287 to 298, wherein the library has been selected against a predetermined antigen optionally has a library size of from 10^2 to 10^9 cells.

300. The library of any one of claims 287 to 299, wherein said first and second cells are progeny of first and second ancestor non-human vertebrate cells respectively, wherein the first ancestor cell comprises a genome comprising said first human immunoglobulin gene segment; and the second ancestor cell comprises a genome comprising said second human immunoglobulin gene segment.

301. A library of antibody-producing transgenic cells whose genomes collectively encode a repertoire of antibodies, wherein the library comprises the first and second ancestor cells recited in claim 300.

302. A library of hybridoma cells produced by fusion of the library of any one of claims 265 to 301 with fusion partner cells; and optionally the hybridoma library has a size of from 10^2 to 10^9 cells

303. An isolated antibody having

(a) a heavy chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unarranged human immunoglobulin V gene segment with a human D and human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments is derived from the genome of an individual from a first human ethnic population; and the other two gene segments are derived from the genome of an individual from a second, different, human ethnic population, and wherein the antibody comprises heavy chain constant regions of said non-human vertebrate (eg, rodent, mouse or rat heavy chain constant regions); and/or

(b) a light chain encoded by a nucleotide sequence produced following recombination in a transgenic non-human vertebrate cell of an unarranged human immunoglobulin V gene segment with a human J segment, optionally with affinity maturation in said cell, wherein one of the gene segments is derived from the genome of an individual from a first human ethnic population (optionally the same as the first population in (a)); and the other gene segment is derived from the genome of an individual from a second, different, human ethnic population (optionally the same as the second population in (a)), and wherein the antibody comprises light chain constant regions of said non-human vertebrate (eg, rodent, mouse or rat heavy light constant regions);

(c) Optionally wherein each variable domain of the antibody is a human variable domain.

(d) Optionally wherein the heavy chain constant regions are gamma-type constant regions.

304. An isolated nucleotide sequence encoding the antibody of claim 303, optionally wherein the sequence is provided in an antibody expression vector, optionally in a host cell.

305. A method of producing a human antibody, the method comprising replacing the non-human vertebrate constant regions of the antibody of claim 303 with human antibody constant regions.

306. A pharmaceutical composition comprising an antibody according to claim 303, or an antibody produced according to the method of claim 305 and a diluent, excipient or carrier; optionally wherein the composition is provided in a container connected to an IV needle or syringe or in an IV bag.

307. An antibody-producing cell that expresses the second antibody recited in any one of claims 287 to 302 or the isolated antibody of claim 303.

308. A non-human vertebrate or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus (eg, a heavy chain locus or a light chain locus), said locus comprising immunoglobulin gene segments according to the first and second human immunoglobulin gene segments (optionally V segments) recited in any one of claims 287 to 302 operably connected upstream of an immunoglobulin constant region; optionally wherein the genome is homozygous for said transgenic immunoglobulin locus;

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

309. A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises a transgenic immunoglobulin locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region;

wherein the transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, a first (optionally a V segment) of said gene segments and a second (optionally a V segment) of said gene segments being different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related;

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

optionally wherein the immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

5

310. A transgenic non-human vertebrate (eg, a mouse or rat) or vertebrate cell (optionally an ES cell or antibody-producing cell) whose genome comprises first and second transgenic immunoglobulin loci, each locus comprising a plurality of human immunoglobulin gene segments operably connected upstream of a non-human vertebrate constant region for the production of a repertoire of chimaeric antibodies, or chimaeric light or heavy chains, having a non-human vertebrate constant region and a human variable region;

10

wherein (i) the first transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments, (ii) the second transgenic locus comprises one or more human immunoglobulin V gene segments, one or more human J gene segments and optionally one or more human D gene segments; and (iii) wherein a first (optionally a V) gene segment of said first locus and a second (optionally a V) gene segment of said second gene locus are different and derived from the genomes of first and second human individuals respectively, wherein the individuals are different; and optionally not related

15

20

optionally wherein the first and second loci are on different chromosomes (optionally chromosomes with the same chromosome number) in said genome;

25

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional V gene segments; and/or

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional D gene segments; and/or

30

optionally wherein each immunoglobulin locus comprises more than the natural human complement of functional J gene segments.

35

311. The non-human vertebrate or cell of claim 309 or 310, wherein the immunoglobulin gene segments are as recited in any one of claims 23 to 37.

312. The non-human vertebrate or cell of claim 308, 309, 310 or 311, wherein the genome comprises a third immunoglobulin gene segment (optionally a V segment), the third gene segment being derived from a human individual that is different from the individual from which the first (and optionally also the second) gene segment is derived; optionally wherein the first, second and third gene segments are polymorphic variants of a human immunoglobulin gene segment.

40

313. The non-human vertebrate or cell of any one of claims 308 to 312, wherein the genome of the vertebrate or cell is homozygous for the first, second and optional third gene segment,

45

wherein a copy of the first, second and optional third gene segments are provided together on the same chromosome operably connected upstream of a common non-human vertebrate constant region.

- 5 314. The non-human vertebrate or cell of any one of claims 308 to 313, wherein each first, second and optional third gene segment is a V gene segment.
- 10 315. The library of any one of claims 287 to 302, wherein the library is provided by a collection of non-human vertebrates (optionally a collection of rodents, mice or rats); optionally, wherein a first member of said collection produces said first antibody but not said second antibody, and a second member of the collection produces said second antibody (but optionally not said first antibody).
- 15 316. A repertoire of antibodies expressed from a library of cells according to any one of claims 287 to 302 and 315.
- 20 317. A method of constructing a cell (eg, an ES cell) according to any one of claims 265 to 286, the method comprising
 (a) identifying functional V and J (and optionally D) gene segments of the genome sequence of a (or said) first human individual;
 (b) identifying one or more functional V and/or D and/or J gene segments of the genome sequence of a (or said) second human individual, wherein these additional gene segments are not found in the genome sequence of the first individual;
 (c) and constructing a transgenic immunoglobulin locus in the cell, wherein the gene segments
 25 of (a) and (b) are provided in the locus operably connected upstream of a constant region.
- 30 318. The method of claim 317, wherein the cell comprises a heavy chain locus constructed according to steps (a) to (c) and/or a light chain locus (kappa and/or lambda loci) constructed according to steps (a) to (c).
- 35 319. The method of claim 317 or 318, wherein the cell is homozygous for the or each transgenic locus; optionally wherein antibody expression from loci endogenous to said cell has been inactivated.
- 40 320. The method of any one of claims 317 to 319, wherein the gene segment(s) in step (b) are identified from an immunoglobulin gene database selected from the 1000 genomes, Ensembl, Genbank and IMGT databases.
321. The method of any one of claims 317 to 320, wherein the first and second human individuals are members of first and second ethnic populations respectively, wherein the populations are different, optionally wherein the human immunoglobulin gene segment derived from the genome sequence of the second individual is low-frequency (optionally rare) within the second ethnic population.

322. A method of making a transgenic non-human vertebrate (eg, a mouse or rat), the method comprising
- (a) constructing an ES cell (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain ES cell) by carrying out the method of any one of claims 317 to 321;
 - 5 (b) injecting the ES cell into a donor non-human vertebrate blastocyst (eg, a mouse C57BL/6N, C57BL/6J, 129S5 or 129Sv strain blastocyst);
 - (c) implanting the blastocyst into a foster non-human vertebrate mother (eg, a C57BL/6N, C57BL/6J, 129S5 or 129Sv strain mouse); and
 - 10 (d) obtaining a child from said mother, wherein the child genome comprises a transgenic immunoglobulin locus.
323. A transgenic non-human vertebrate (eg, a mouse or rat) made by the method of claim 322 or a progeny thereof.
- 15 324. A population of non-human vertebrates according to claim 323.
325. A method of isolating an antibody that binds a predetermined antigen (eg, a bacterial or viral pathogen antigen), the method comprising
- (a) providing a vertebrate (optionally a mouse or rat) according to any one of claims 265 to 286, 289, 290, 294, 295, 308 to 314 and 323;
 - 20 (b) immunising (eg, using a standard prime-boost method) said vertebrate with said antigen (optionally wherein the antigen is an antigen of an infectious disease pathogen);
 - (c) removing B lymphocytes from the vertebrate and selecting one or more B lymphocytes expressing antibodies that bind to the antigen;
 - 25 (d) optionally immortalising said selected B lymphocytes or progeny thereof, optionally by producing hybridomas therefrom; and
 - (e) isolating an antibody (eg, and IgG-type antibody) expressed by the B lymphocytes; and
 - (f) optionally producing a derivative or variant of the antibody.
- 30 326. The method of claim 325, further comprising after step (e) the step of isolating from said B lymphocytes nucleic acid encoding said antibody that binds said antigen; optionally exchanging the heavy chain constant region nucleotide sequence of the antibody with a nucleotide sequence encoding a human or humanised heavy chain constant region and optionally affinity maturing the variable region of said antibody; and optionally inserting said nucleic acid into an expression vector and optionally a host.
- 35 327. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic heavy chain immunoglobulin locus, wherein the locus comprises at least 42 (optionally at least 43, 44, 45, 46, 47, 48, 49, 50) functional human VH gene segments, one or more functional human D gene segments, one or more functional human JH gene segments operably connected upstream of a non-human vertebrate constant region (eg, a mouse constant region, eg, a Cmu and/or a C gamma).
- 40 328. The vertebrate or cell of claim 327, wherein the locus comprises at least 23 functional human D gene segments and/or at least 6 functional human JH gene segments, optionally at
- 45

least a full human repertoire of functional VH, D and JH segments (optionally derived from the genome sequence of the same human individual).

329. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic heavy chain immunoglobulin locus, wherein the locus comprises a full human repertoire of functional VH, D and JH segments derived from the genome sequence of the same human individual, supplemented with one or more additional functional human VH, D and/or JH gene segment that is not found in the genome sequence of said human individual.

330. The vertebrate or cell of claim 329, wherein each additional VH, D and JH segment is selected from the group consisting of (i) a gene segment derived from a second, different individual, optionally wherein the first and second individuals are members of different ethnic populations; (ii) a mutant of a human gene segment (optionally a human germline gene segment having up to 5, 10 or 15 mutations); and (iii) a hybrid human gene segment comprising a first and second nucleotide sequences derived from genome sequences of different human individuals or different polymorphic variants of a human immunoglobulin gene segment (eg, variants of a germline human VH, D or JH gene segment).

331. The vertebrate or cell of any one of claims 328 to 330, wherein the genome is homozygous for said heavy chain transgene and optionally endogenous immunoglobulin heavy chain expression has been inactivated.

332. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic kappa light chain immunoglobulin locus, wherein the locus comprises at least 39 functional human Vk gene segments and one or more functional human Jk gene segments operably connected upstream of a non-human vertebrate constant region (eg, a mouse constant region).

333. The vertebrate or cell of claim 332, wherein the locus comprises at least a full human repertoire of functional Vk and Jk segments (optionally derived from the genome sequence of the same human individual).

334. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic kappa light chain immunoglobulin locus, wherein the locus comprises a full human repertoire of functional Vk and Jk segments derived from the genome sequence of the same human individual, supplemented with one or more additional functional human Vk and/or Jk gene segment that is not found in the genome sequence of said human individual.

335. The vertebrate or cell of claim 334, wherein each additional Vk and Jk segment is selected from the group consisting of (i) a gene segment derived from a second, different individual, optionally wherein the first and second individuals are members of different ethnic populations; (ii) a mutant of a human gene segment (optionally a human germline gene segment having up to 5, 10 or 15 mutations); and (iii) a hybrid human gene segment comprising a first and second nucleotide sequences derived from: genome sequences of different human individuals or

different polymorphic variants of a human immunoglobulin gene segment (eg, variants of a germline human Vk or Jk gene segment).

336. The vertebrate or cell of any one of claims 332 to 335, wherein the genome is homozygous for said kappa light chain transgene and optionally endogenous immunoglobulin kappa light chain expression has been inactivated.

337. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic lambda light chain immunoglobulin locus, wherein the locus comprises at least 32 functional human V lambda gene segments and one or more functional human J lambda gene segments operably connected upstream of a non-human vertebrate constant region (eg, a mouse constant region).

338. The vertebrate or cell of claim 337, wherein the locus comprises at least a full human repertoire of functional V lambda and J lambda segments (optionally derived from the genome sequence of the same human individual).

339. A non-human vertebrate (eg, a mouse or rat) or cell (eg, a mouse cell or rat cell) comprising a genome that comprises a transgenic lambda light chain immunoglobulin locus, wherein the locus comprises a full human repertoire of functional V lambda and J lambda segments derived from the genome sequence of the same human individual, supplemented with one or more additional functional human V lambda and/or J lambda gene segment that is not found in the genome sequence of said human individual.

340. The vertebrate or cell of claim 339, wherein each additional V lambda and J lambda segment is selected from the group consisting of (i) a gene segment derived from a second, different individual, optionally wherein the first and second individuals are members of different ethnic populations; (ii) a mutant of a human gene segment (optionally a human germline gene segment having up to 5, 10 or 15 mutations); and (iii) a hybrid human gene segment comprising a first and second nucleotide sequences derived from genome sequences of different human individuals or different polymorphic variants of a human immunoglobulin gene segment (eg, variants of a germline human V lambda or J lambda gene segment).

341. The vertebrate or cell of any one of claims 337 to 340, wherein the genome is homozygous for said lambda light chain transgene and optionally endogenous immunoglobulin lambda light chain expression has been inactivated.

342. A transgenic immunoglobulin locus comprising a synthetic immunoglobulin gene haplotype, the haplotype comprising first and second human gene segments (each being a V, D or J), a switch region and a constant region, wherein

- a) the second gene segment is a polymorphic variant of the first gene segment; or
- b) the first and second gene segments are derived respectively from genome sequence of individuals from different, first and second, ethnic populations (eg, according to the 1000 Genomes database) and the second gene segment is not found in the first population (eg,

according to the 1000 Genomes database), optionally wherein the second gene segment occurs at low-frequency (or is rare) in the second population; and

Wherein the constant region, and optionally the switch region, are non-human vertebrate (eg, mouse or rat) constant and switch regions (eg, mouse or rat Cmu; mouse or rat Smu; mouse or rat C gamma; or mouse or rat S gamma).

343. The transgenic immunoglobulin locus of claim 342 wherein the transgene comprises a third human immunoglobulin gene segment (a V, D or J) which is (a) a polymorphic variant of the first and second gene segments; or (b) derived from a genome sequence of the second population (and optionally is a low-frequency or rare gene in that population) or is from a third human ethnic population, wherein the third gene segment is not found in the first population (eg, according to the 1000 Genomes database).

344. A non-human vertebrate or cell comprising the transgene of claim 342 or 343; optionally wherein the vertebrate or cell is homozygous for said transgene.

345. An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%;, and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

346. The antibody of claim 345 wherein the constant region is a IGHG1a, IGHG2a, IGHG3a, IGHG3b or IGHG4a constant region.

347. The antibody of claim 345 or 346, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

348. The antibody of claim 345, 346 or 347, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%.

349. The antibody of claim 345, 346, 347 or 348 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in

a Chinese population and with a cumulative frequency of at least 5%.

350. An isolated VH domain identical to a variable domain as recited in any one of claims 347 to 349, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

5

351. A pharmaceutical composition comprising the antibody or variable domain of any one of claims 345 to 350 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

10 352. An isolated antibody for administration to a Chinese patient, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found
15 in a Chinese population and with a cumulative frequency of at least 5%; and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg,
20 mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

353. The antibody of claim 352, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in a Chinese
25 population and with a cumulative frequency of at least 5%.

354. The antibody of claim 352 or 353, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in a Chinese
30 population and with a cumulative frequency of at least 5%.

355. An isolated VH domain identical to a variable domain as recited in any one of claims 352 to 354, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

35 356. A pharmaceutical composition comprising the antibody or variable domain of any one of claims 352 to 355 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

40 357. An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a Chinese patient, wherein the heavy chain is a heavy chain produced by the following steps (or is a copy of such a heavy chain):-

45 (a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived

from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment , the VH gene segment being selected from a VH in Table 13 found in a Chinese population and with a cumulative frequency of at least 5%;

5 (b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

10 358. The antibody heavy chain or VH domain of claim 357, wherein the human constant region is as recited in claim 345 or 346.

15 359. An antibody heavy chain or VH domain as recited in claim 357 or 358 for use in a medicament for therapy and/or prophylaxis of a disease or medical condition in a Chinese patient.

20 360. A method of treating and/or preventing a disease or medical condition in a Chinese patient, the method comprising administering to the patient a therapeutically or prophylactically-effective amount of the antibody heavy chain or VH domain as recited in claim 357 or 358.

25 361. An isolated antibody for administration to a patient of European, East Asian, West African, South Asian or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the constant region is a human constant region selected from a constant region (eg, an IGHG constant region) in Table 13 found in a population of European, East Asian, West African, South Asian or Americas ancestry respectively and with a cumulative frequency of at least 5%;, and wherein

30 (i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

35 362. The antibody of claim 361 wherein the constant region is a IGHG1a, IGHG2a, IGHG3a, IGHG3b or IGHG4a constant region and the patient is of European ancestry.

40 363. The antibody of claim 361 or 362, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in said population and with a cumulative frequency of at least 5%.

364. The antibody of claim 361, 362 or 363, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment , the D gene segment being selected from a D in Table 13 found in

said population and with a cumulative frequency of at least 5%.

365. The antibody of claim 361, 362, 363 or 364 wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in said population and with a cumulative frequency of at least 5%.

366. An isolated VH domain identical to a variable domain as recited in any one of claims 363 to 365, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

367. A pharmaceutical composition comprising the antibody or variable domain of any one of claims 361 to 366 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

368. An isolated antibody for administration to a patient of European, East Asian, West African or Americas ancestry, the antibody comprising a human heavy chain, the heavy chain comprising a variable domain that is specific for an antigen and a constant region, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in a population of European, East Asian, West African, South Asian or Americas ancestry respectively and with a cumulative frequency of at least 5%; and wherein

(i) the variable domain is derived from the recombination of said human gene segments in a non-human vertebrate (eg, in a mouse or a rat); or (ii) the variable domain comprises non-human vertebrate (eg, mouse or rat) AID-pattern mutations and non-human vertebrate (eg, mouse or rat) terminal deoxynucleotidyl transferase (TdT)-pattern mutations.

369. The antibody of claim 368, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the D gene segment being selected from a D in Table 13 found in said population and with a cumulative frequency of at least 5%.

370. The antibody of claim 368 or 369, wherein the variable domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the JH gene segment being selected from a JH in Table 13 found in said population and with a cumulative frequency of at least 5%.

371. An isolated VH domain identical to a variable domain as recited in any one of claims 368 to 370, optionally fused at its C-terminus to a polypeptide (eg, an antibody Fc).

372. A pharmaceutical composition comprising the antibody or variable domain of any one of claims 368 to 371 together with a pharmaceutically-acceptable excipient, diluent or a medicament (eg, a further antigen-specific variable domain, antibody chain or antibody).

373. An antibody heavy chain or VH domain (eg, provided as part of an antibody) for therapy and/or prophylaxis of a disease or medical condition in a patient of European, East Asian, West African, South Asian or Americas ancestry, wherein the heavy chain is a heavy chain produced by the following steps (or is a copy of such a heavy chain):-

5

(a) Selection of an antigen-specific antibody heavy chain or VH domain from a non-human vertebrate (eg, a mouse or a rat), wherein the heavy chain or VH domain is derived from the recombination of a human VH gene segment with a human D gene segment and a human JH gene segment, the VH gene segment being selected from a VH in Table 13 found in said population and with a cumulative frequency of at least 5%;

10

(b) Optional humanisation of the heavy chain by combining the variable domain of the heavy chain with a human constant region; or optional humanisation of the selected VH domain by combining with a human constant region.

15

374. The antibody heavy chain or VH domain of claim 373, wherein the human constant region is as recited in claim 361 or 362.

375. An antibody heavy chain or VH domain as recited in claim 373 or 374 for use in a medicament for therapy and/or prophylaxis of a disease or medical condition in a patient of said ancestry.

20

376. A method of treating and/or preventing a disease or medical condition in a patient of European, East Asian, West African, South Asian or Americas ancestry, the method comprising administering to the patient a therapeutically or prophylactically-effective amount of the antibody heavy chain or VH domain as recited in claim 373 or 374.

25

Recombineered BAC Vectors to add Polymorphic V-regions to the Mouse Genome

1. Cassette Exchange – Replace existing V-regions




Figure 1

Recombineered BAC Vectors to add
Heterogeneous V-regions to the Mouse Genome

2. Insertion of V-regions into Heterologous Genomic DNA




Figure 2

Recombineered BAC Vectors to add
Heterogeneous V-regions to the Mouse Genome

3. Single Strand Nucleotide Changes





Figure 3

Adding Polymorphic V-regions to the Genome using SRMCE of Modified BACs

1. Insertion at a 'Landing Pad' site during SRMCE



00000

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
	Q	V	Q	L	V	Q	S	G	A	E	V	K	K	P	G	S	S	S	V	K	V	S	C	K	A	S
102582 ,IGHV1-69*01, hv1051	CAG	GTC	CAG	CTG	GTC	CAG	TCT	GCG	GCT	...	GAG	GTG	AAG	AAG	CCT	GCG	TCC	TCC	GTG	AAG	GTC	TCC	TGC	AAG	GCT	TCT
227505 ,IGHV1-69*02, yIGH6(YAC7)	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
N92340 ,IGHV1-69*02, 57GTA8	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
M83132 ,IGHV1-69*04, hv1263	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
X67805 ,IGHV1-69*05, RR.VR1.2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
522553 ,IGHV1-69*06, hv1051K	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
229978 ,IGHV1-69*07, DA-2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214309 ,IGHV1-69*08	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214307 ,IGHV1-69*09	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214300 ,IGHV1-69*10	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214296 ,IGHV1-69*11	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214301 ,IGHV1-69*12	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
214214 ,IGHV1-69*13	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...

Alignment of 13 IGHV1-69 alleles showing the variable (V) coding region only. Nucleotides that differ from VH1-69 allele *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above the corresponding triplet. Boxed regions correspond to CDR1, CDR2 and CDR3 as indicated.

Figure 5 (part 1 of 4)

	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53
	G	G	T	F	S	S	Y	A					I	S	W	Y	R	Q	A	P	G	Q	G	L	E	W	M
	GGA	GCG	ACC	TTC	AGC	AGC	TAT	GCT	...	...	...	...	ATC	AGC	TGG	GTC	CGA	CAG	GCC	CGT	GGA	CAA	GCG	CGT	GAG	TGG	ATG
L22582 ,IGHV1-69*01, hv1051	...	...	...	...	...	...	...	A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z27506 ,IGHV1-69*02, y13H6(YAC7)	...	...	...	...	...	...	...	A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
X92340 ,IGHV1-69*03, 5YCHAB	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
M83131 ,IGHV1-69*04, hv1263	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
X67905 ,IGHV1-69*05, BR.VH1.2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
L22563 ,IGHV1-69*06, hv1051N	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z29978 ,IGHV1-69*07, BA-2	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14309 ,IGHV1-69*08	...	...	...	...	...	...	...	A	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14307 ,IGHV1-69*09	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14300 ,IGHV1-69*10	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14296 ,IGHV1-69*11	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14301 ,IGHV1-69*12	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...
Z14214 ,IGHV1-69*13	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...

Alignment of 13 IGHV1-69 alleles showing the variable (V) coding region only. Nucleotides that differ from VH1-69 allele *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above the corresponding triplet. Boxed regions correspond to CDR1, CDR2 and CDR3 as indicated.

Figure 5 (part 2 of 4)

	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
	G	G	I	I	P	I	E	G	T	A			N	Y	A	Q	R	F	Q		G	R	V	F	I	I	A
L22582 ,IGHV1-69*01, hv1051	CGA	GCG	ATC	ACC	CCT	ATC	TTT	GCT	ACA	GCA	...	...	AAC	TAC	GCA	CAG	AAG	TTC	CAG	...	GCC	AGA	GTC	ACG	ATT	ACC	GCG
Z07506 ,IGHV1-69*02, yIGH6(YAC7	R					L	I																				
X92340 ,IGHV1-69*03, 57GTAS	R					L	I																				
M83132 ,IGHV1-69*04, hv1263	R					C																					
X67905 ,IGHV1-69*05, RR.VH1.2																											T
L22563 ,IGHV1-69*06, hv1051K	R																										R
Z29978 ,IGHV1-69*07, DA-2	R					L																					
Z14309 ,IGHV1-69*08	R					C																					
Z14307 ,IGHV1-69*09	R					L	I																				
Z14300 ,IGHV1-69*10						C																					
Z14296 ,IGHV1-69*11	R					L																					
Z14301 ,IGHV1-69*12						C																					
Z14014 ,IGHV1-69*13																											


Alignment of 13 IGHV1-69 alleles showing the variable (V) coding region only. Nucleotides that differ from VH1-69 allele *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above the corresponding triplet. Boxed regions correspond to CDR1, CDR2 and CDR3 as indicated.

Figure 5 (part 3 of 4)

Alignment of 13 IGHV1-69 alleles showing the variable (V) coding region only. Nucleotides that differ from VH1-69 allele *01 are indicated at the appropriate position whereas identical nucleotides are marked with a dash. Where nucleotide changes result in amino acid differences, the encoded amino acid is shown above the corresponding triplet. Boxed regions correspond to CDR1, CDR2 and CDR3 as indicated.

41016150

Figure 6



[illegible]

FIGURE 8


Rabbit JH6	Y	tac	Y	G	M	D	L
		at	tac	ggc	atg	gac	ctc
Sheep JH6	Y	tac	Y	G	V	D	V
		at	tac	ggt	gta	gat	gtc
Bovine JH6	Y	tac	Y	G	V	D	V
		at	tac	ggt	gta	gat	gtc
Dog JH3	Y	tac	Y	G	M	D	Y
		at	tac	ggt	atg	gac	tac
Human JH6*02	Y	tac	Y	G	M	D	V
		at	tac	ggt	atg	gac	gtc

FIGURE 9

	T			C			A			G		
T	TTT	Phe	F	TCT	Ser	S	TAT	Tyr	Y	TGT	Cys	C
	TTC	Phe	F	TCC	Ser	S	TAC	Tyr	Y	TGC	Cys	C
	TTA	Leu	L	TCA	Ser	S	TAA	stop	*	TGA	stop	*
	TTG	Leu	L	TCG	Ser	S	TAG	stop	*	TGG	Trp	W
C	CTT	Leu	L	CCT	Pro	P	CAT	His	H	CGT	Arg	R
	CTC	Leu	L	CCC	Pro	P	CAC	His	H	CGC	Arg	R
	CTA	Leu	L	CCA	Pro	P	CAA	Gln	Q	CGA	Arg	R
	CTG	Leu	L	CCG	Pro	P	CAG	Gln	Q	CGG	Arg	R
A	ATT	Ile	I	ACT	Thr	T	AAT	Asn	N	AGT	Ser	S
	ATC	Ile	I	ACC	Thr	T	AAC	Asn	N	AGC	Ser	S
	ATA	Ile	I	ACA	Thr	T	AAA	Lys	K	AGA	Arg	R
	ATG	Met	M	ACG	Thr	T	AAG	Lys	K	AGG	Arg	R
G	GTT	Val	V	GCT	Ala	A	GAT	Asp	D	GGT	Gly	G
	GTC	Val	V	GCC	Ala	A	GAC	Asp	D	GGC	Gly	G
	GTA	Val	V	GCA	Ala	A	GAA	Glu	E	GGA	Gly	G
	GTG	Val	V	GCG	Ala	A	GAG	Glu	E	GGG	Gly	G

Figure 10

chr14:106,026,574-107,346,185 1,319,612 bp. enter position, gene symbol or search term



Recombineered BAC Vectors to add Polymorphic V-regions to the Mouse Genome

1. Cassette Exchange – Replace existing V-regions




Figure 1