



US 20070148147A1

(19) **United States**(12) **Patent Application Publication**
Dennin et al.(10) **Pub. No.: US 2007/0148147 A1**(43) **Pub. Date: Jun. 28, 2007**(54) **BACTERIAL COMPOSITION AND ITS USE****Publication Classification**(76) Inventors: **Veronique Dennin**, Villeneuve D'ascq
(FR); **Gregory John Leyer**, Madison,
WI (US); **Annick Mercenier**, Epalinges
(CH); **Sophie Nutton**, Steenvoorde
(FR); **Bruno Pot**, St-Michiels Brugge
(BE)(51) **Int. Cl.****A61K 35/74** (2006.01)**C12N 1/20** (2006.01)(52) **U.S. Cl.** **424/93.45; 435/252.9**

Correspondence Address:

FAY SHARPE LLP**1100 SUPERIOR AVENUE, SEVENTH FLOOR**
CLEVELAND, OH 44114 (US)

(57)

ABSTRACT(21) Appl. No.: **11/594,542**(22) Filed: **Nov. 8, 2006****Related U.S. Application Data**(62) Division of application No. 10/310,549, filed on Dec.
5, 2002, now Pat. No. 7,179,460.

The subject of the present invention is a bacterial composition having immunomodulation properties comprising at least one strain selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802. An other subject of the invention is an immunomodulation method comprising the step of using the at least one strain selected from the preceding group.




Figure 1: schedule of probiotic administration




Figure 3: schedule of probiotic administration




Figure 5: schedule of probiotic administration




Figure 2: Wallace and Ameho scores, and weight evolution





Figure 4: Wallace and Ameho scores




Figure 6: Wallace scores

BACTERIAL COMPOSITION AND ITS USE

[0001] The subject of the present invention is a bacterial composition, an immunomodulation method and the use of this composition. Immunomodulation is the capacity to improve the global immune functions either in healthy or in pathologic situations.

[0002] Among bacteria, some have a positive influence on the immune system of the intestinal medium, in particular lactic acid bacteria and bifidobacteria, and are termed "probiotic" bacteria or strains.

[0003] The expression probiotic bacteria or strains is understood to mean a strain which, when ingested live, exerts a beneficial effect on the host by having an action on the balance of the intestinal flora. These probiotic strains have the capacity to survive following passage across the upper part of the digestive tube. They are non-pathogenic, non-toxic and exert a beneficial action on health through in one hand their ecological interactions with the resident flora of the digestive tract and in the other hand their ability to positively influence immune system, through their effects onto the GALT (gut associated immune tissue). According to the probiotic definition, those bacteria when given in sufficient numbers have the capacity to transit alive all along the gut. There they become part of the resident flora for the period of administration. The so called colonisation (or transient colonisation) allow the probiotic bacteria to exert beneficial effect, such as repression of potentially pathogen micro-organisms on the flora and interactions with the gut immune system.

[0004] The probiotic strains most widely used, in particular in dairy products, are mainly bacteria and yeasts, of the following genera *Lactobacillus* spp, *Streptococcus* spp, *Enterococcus* spp and *Bifidobacteriu* spp and *Sacharomyces* spp. Among the probiotic effects reported for those bacteria, one can cite for example, improvement of lactose tolerance, prevention and/or treatment of gastrointestinal and urogenital infections, reduction of some cancers, decrease of blood cholesterol. However, it should be highlighted that not all the individual strains from the genera described above have those effects, but only some carefully selected strains do.

[0005] Thus in order to satisfy the requirements for performant probiotic strains, it has become necessary to select strains or a mixture thereof which is efficient and which allow stimulation of the immune system (immunomodulation).

[0006] Accordingly, the problem which the present invention proposes to solve is to provide a bacterial composition having probiotic properties.

[0007] The aim of the present invention is to satisfy these requirements.

[0008] For this purpose, the present invention provides a bacterial composition having immunomodulation properties comprising at least one strain selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802.

[0009] Another subject of the invention is an, immunomodulation method.

[0010] The subject of the invention is also a food or pharmaceutical composition comprising the bacterial compositions described above.

[0011] Another subject of the invention is also the use of this composition in the preparation of a carrier administered to humans or to animals for a therapeutic or prophylactic purpose in the gastrointestinal system.

[0012] The composition of the invention has the advantage of providing unquestionable virtues which enrich the range of available strains.

[0013] Such a composition is particularly advantageous when it is administered to humans or to animals for a therapeutic or prophylactic purpose in the gastrointestinal system, in particular in the reduction of inflammatory and/or allergic reactions.

[0014] The advantage of the present invention is also to preserve all its properties when it is incorporated into a pharmaceutically acceptable carrier or into a food product.

[0015] Other advantages and characteristics of the present invention will emerge more clearly on reading the description and the examples given purely by way of illustration and without limitation, which follow.

[0016] First of all, the subject of the invention is a bacterial composition having immunomodulation properties comprising at least one strain selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802.

[0017] The *Lactobacillus acidophilus* used according to the invention is a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a homofermentative metabolism giving rise to the production of lactic acid.

[0018] The *Lactobacillus acidophilus* used according to the invention may also produce a bacteriocin, lactacin, which is active against other microorganisms.

[0019] Preferably, it is a *Lactobacillus acidophilus* exhibiting good resistance to pepsin, under acidic pH conditions (about at least 73% of the bacteria are still alive after 40 minutes of treatment).

[0020] More particularly, the *Lactobacillus acidophilus* used according to the invention exhibits a very good resistance to pancreatin (at least 100% of the bacteria are still alive after 40 minutes of treatment).

[0021] Advantageously, the *Lactobacillus acidophilus* used according to the invention exhibits good tolerance to bile salts.

[0022] Preferably, a *Lactobacillus acidophilus* described as being "hydrophobic", that is to say which exhibits a high affinity for hydrophobic organic solvents, polar or not polar, such as for example n-decane, chloroform, hexadecane or xylene, will be used.

[0023] The *Lactobacillus acidophilus* used according to the invention can induce the production of cytokines. This detection of the induction-of cytokines was made by means of a test for in vitro stimulation of isolated peripheral blood mononuclear cells (PBMC). Among the cytokines induced during this test, there may be mentioned interleukins 10

(IL10), γ -interferon (γ -IFN) and tumour necrosis factor α (TNF α). On the other hand, the *Lactobacillus acidophilus* used according to the invention induces little or no secretion of interleukins 12 (IL12) using this same test.

[0024] The *Lactobacillus acidophilus* used according to the invention is *Lactobacillus acidophilus* PTA-4797. This *Lactobacillus acidophilus* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number PTA-4797.

[0025] One embodiment of the composition according to the invention is a bacterial composition comprising *Lactobacillus acidophilus* PTA-4797.

[0026] The composition made of probiotic bacteria according to the invention may also comprise at least one *Lactobacillus plantarum* strain.

[0027] The *Lactobacillus plantarum* strain used according to the invention is preferably a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a homofermentative metabolism giving rise to the production of lactic acid.

[0028] Preferably, it is a *Lactobacillus plantarum* which is resistant to pepsin under acidic pH conditions (about at least 95% of the bacteria are still alive after 40 minutes of treatment). More particularly, the *Lactobacillus plantarum* used according to the invention exhibits good resistance to pancreatin (about at least 79% of the bacteria are still alive after 40 minutes of treatment).

[0029] Advantageously, the *Lactobacillus plantarum* used according to the invention exhibits good resistance to bile salts.

[0030] The *Lactobacillus plantarum* used according to the invention can induce the production of cytokines. This detection of the induction of cytokines was made by means of a test for in vitro stimulation of isolated peripheral blood mononuclear cells (PBMC). Among the cytokines induced during this test, there may be mentioned interleukins 10 (IL10), γ -interferon (γ -IFN) and tumour necrosis factor α (TNF α). On the other hand, the *Lactobacillus plantarum* used according to the invention induces little or no secretion of interleukins 12 (IL12) using this same test.

[0031] The *Lactobacillus plantarum* used according to the invention is *Lactobacillus plantarum* PTA-4799. This *Lactobacillus plantarum* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number PTA-4799.

[0032] One embodiment of the composition according to the invention is a bacterial composition comprising *Lactobacillus plantarum* PTA-4799.

[0033] The composition made of probiotic bacteria according to the invention may also comprise at least one *Lactobacillus salivarius* strain.

[0034] The *Lactobacillus salivarius* strain used according to the invention is a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a homofermentative metabolism giving rise to the production of lactic acid.

[0035] More particularly, the *Lactobacillus salivarius* used according to the invention exhibits good resistance to pancreatin (at least 100% of the bacteria are still alive after 40 minutes of treatment).

[0036] The *Lactobacillus salivarius* used according to the invention can induce the secretion of cytokines. This detection of the induction of cytokines was made by means of a test for in vitro stimulation of isolated peripheral blood mononuclear cells (PBMC). Among the cytokines induced during this test, there may be mentioned interleukins 10 (IL10) and tumour necrosis factor α (TNF α). On the other hand, the *Lactobacillus salivarius* used according to the invention induces little or no secretion of interleukin 12 (IL12) and γ -interferon (γ -IFN) using this same test.

[0037] The *Lactobacillus salivarius* used according to the invention is *Lactobacillus salivarius* PTA-4800. This *Lactobacillus salivarius* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number PTA-4800.

[0038] One embodiment of the composition according to the invention is a bacterial composition comprising *Lactobacillus salivarius* PTA-4800.

[0039] The composition made of probiotic bacteria according to the invention may also comprise at least one *Lactobacillus paracasei* strain.

[0040] The *Lactobacillus paracasei* used according to the invention is a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a homofermentative metabolism giving rise to the production of lactic acid.

[0041] Preferably, it is a *Lactobacillus paracasei* exhibiting poor resistance to pepsin, under acidic pH conditions (about at least 17.5% of the bacteria are still alive after 40 minutes of treatment).

[0042] More particularly, the *Lactobacillus paracasei* used according to the invention exhibits a very good resistance to pancreatin (about at least 100% of the bacteria are still alive after 40 minutes of treatment).

[0043] Advantageously, the *Lactobacillus paracasei* used according to the invention exhibits good tolerance to bile salts.

[0044] The *Lactobacillus paracasei* used according to the invention can induce the production of cytokines. This detection of the induction of cytokines was made by means of a test for in vitro stimulation of isolated peripheral blood mononuclear cells (PBMC). Among the cytokines induced during this test, there may be mentioned interleukins 10 (IL10), γ -interferon (γ -IFN) and tumour necrosis factor α (TNF α). On the other hand, the *Lactobacillus paracasei* used according to the invention induces little or no secretion of interleukins 12 (IL12) using this same test.

[0045] The *Lactobacillus paracasei* used according to the invention is *Lactobacillus paracasei* PTA-4798. This *Lactobacillus paracasei* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number PTA-4798.

[0046] One embodiment of the composition according to the invention is a bacterial composition comprising *Lactobacillus paracasei* PTA-4798.

[0047] The composition made of probiotic bacteria according to the invention may also comprise at least one *Bifidobacterium bifidum* strain.

[0048] The *Bifidobacterium bifidum* strain used according to the invention is a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a heterofermentative metabolism giving rise to the production of lactic acid and acetic acid.

[0049] The *Bifidobacterium bifidum* used according to the invention is *Bifidobacterium bifidum* PTA-4801. This *Bifidobacterium bifidum* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number ATCC PTA-4801.

[0050] One embodiment of the composition according to the invention is a bacterial composition comprising *Bifidobacterium bifidum* PTA-4801.

[0051] The composition made of probiotic bacteria according to the invention may also comprises at least one *Bifidobacterium lactis* strain.

[0052] The *Bifidobacterium lactis* strain used according to the invention is a Gram-positive strain. Advantageously, it is a catalase-negative strain, with a heterofermentative metabolism giving rise to the production of lactic acid and acetic acid.

[0053] The *Bifidobacterium lactis* used according to the invention is *Bifidobacterium lactis* PTA-4802. This *Bifidobacterium lactis* strain was deposited according to the Treaty of Budapest at the American Type Culture Collection where it is recorded under the deposit number ATCC PTA-4801.

[0054] One embodiment of the composition according to the invention is a bacterial composition comprising *Bifidobacterium lactis* PTA-4801.

[0055] Another embodiment of the composition according to the invention is a bacterial composition comprising at least one strain selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, and *Bifidobacterium lactis* PTA-4802.

[0056] More particularly, the composition according to the invention may comprise at least 2 strains selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802.

[0057] More particularly, the composition according to the invention may comprise 3 strains selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802.

[0058] Preferably, the composition according to the invention may comprise a blend of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium lactis* PTA-4802 and *Bifidobacterium bifidum* PTA-4801.

[0059] The composition according to the invention may be used in the form of a bacterial suspension, before or after freezing, or of a freeze-dried powder. Indeed, regardless of the form used, the composition may be frozen.

[0060] The relative proportion of each bacterium in the composition can vary in large limit for example from 1/99 to 99/1 in the case there is at least 2 strains.

[0061] The composition according to the invention may comprise from 10^6 to 10^{11} CFU of bacteria/g of composition, and more particularly from 10^8 to 10^{11} CFU of bacteria/g of composition. The term CFU means "colony forming units". The expression gram of composition is understood to mean the food product or pharmaceutical preparation, and preferably 10^9 to 10^{11} CFU/g if in a freeze-dried form.

[0062] The composition according to the invention is useful for the treatment, the primary prevention or the recurrence, and also the prevention and/or the reduction of inflammatory bowel disease.

[0063] The bacterial composition according to the invention is also useful for maintaining the homeostasis of the immune system.

[0064] The bacterial composition according to the invention is also useful for the prevention and/or the reduction of allergenicity.

[0065] The bacterial composition according to the invention is also useful as immunoadjuvant.

[0066] An other subject of the invention is also an immunomodulation method comprising the step of using at least one strain selected from the group consisting of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium bifidum* PTA-4801 and *Bifidobacterium lactis* PTA-4802.

[0067] The composition according to the invention may be used in the form of a food product or pharmaceutical preparation.

[0068] The expression pharmaceutical preparation is understood to mean for example a preparation in the form of capsule or tablet.

[0069] The subject of the invention is also a food, dietary supplement or pharmaceutical composition comprising the bacterial composition described above.

[0070] Preferably, the food composition comprising the composition according to the invention is a food supplements, a beverage product or a milk base powder.

[0071] More particularly, the food composition comprising the composition according to the invention is a dairy product.

[0072] More preferably the food composition comprising the composition according to the invention is an infant formula.

[0073] The food or pharmaceutical composition according to the invention may have immunomodulation properties.

[0074] Another subject- of the invention- is also the use of this composition in the preparation of a carrier administered to humans or to animals for a therapeutic or prophylactic purpose in the gastrointestinal system.

[0075] The invention provides also a pharmaceutical composition comprising the bacterial composition described above useful for the prevention of inflammatory bowel disease.

[0076] The invention provides also a pharmaceutical composition comprising the bacterial composition described above useful for immunomodulation.

[0077] The FIGS. 1, 3 and 5 are schedules of injection.

[0078] The FIGS. 2, 4 and 6 are Wallace and Ameho scores, and weight of loss.

[0079] Concrete but nonlimiting examples of the invention will now be described.

EXAMPLES

1/PBMC Test

[0080] PBMC (Peripheral Blood Mononuclear Cells) are prepared by centrifugation from human blood, derived from known donors and is further purified on a Ficoll gradient. Cells are harvested, washed, (red blood cells removed) and counted.

[0081] Bacteria are prepared according to standard conditions and counted-by plating on agarose medium according to proper dilutions (10^{-7} , 10^{-8} , 10^{-9} in Ringer solution).

[0082] Cells are washed 3 times and suspended (et non dissolved) in PBS buffer. A verifier par Dominique

[0083] PBMC cells are stimulated for 48 hours with the bacteria (allow negative control with Phosphate Buffer Saline solution (PBS)) under appropriate conditions of CO_2 . Then the supernatant containing the cytokines is frozen at -20°C .

[0084] Cytokines expression levels are determined by ELISA tests (<<Enzyme linked immuno sorbent assay>>). ELISA plates are coated with anti-cytokine antibody (overnight procedure) and the antibody is blocked with a surfactant, the tween 80.

[0085] A proper standard is prepared for known concentrations of cytokines which will cover the detection range of 15.62 till 2000 pg/ml (incubate overnight). Perform the anti-cytokine detection and quantify with streptavidine reaction on substrate (10 mg dABTS/10 ml of citric acid buffer 0.1 M, pH 4.35/20 μl H_2O_2).

[0086] Cytokines are either pro-inflammatory/Th1 ($\text{TNF}\alpha$, $\text{IFN}\gamma$, IL12) or anti-inflammatory (IL10).

Strains	IL-10	IL-12	$\text{IFN-}\gamma$	$\text{TNF-}\alpha$
<i>L. acidophilus</i> PTA-4797	309		1970	27591
<i>L. salivarius</i> PTA-4800	6881	31	1148	23509
<i>L. paracasei</i> PTA-4798	300	31	921	14046

The values are pg/ml.

2/Preparation of a Composition Made of *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799 and *Bifidobacterium lactis* PTA-4802.

[0087] A blend of 3 strains is prepared containing *Lactobacillus acidophilus* PTA-4797, *Lactobacillus plantarum* PTA-4799, et *Bifidobacterium lactis* PTA-4802. This blend is named LAB.

[0088] To show in vivo, the immuno modulation capacity of this blend according to the invention, animal test were performed aimed at decreasing the chemically induced gut inflammation of mice. Several models were set-up to mimic human inflammatory bowel- diseases.

[0089] In the first model, injections of 1 mg/mice of TNBS (trinitrobenzene sulfonic acid, which is a colitis inducing agent) on days 0, 7 and 14, and 2 mg/mice of TNBS on day 20 (double dose) will evoke a chronic colitis. The schedule of injection is drawn on FIG. 1. The intake of bacteria (LAB) on days -4 to -1, 6, 9, 13 and 19 will show macroscopic and histological. effects, which are scored on day 22, according to standard scores (Wallace and Ameho respectively).

[0090] The results obtained (see FIG. 2) clearly showed an improvement of the colitis symptoms evidenced by

[0091] reduction of the increase of the submucosa

[0092] reduction of the inflammation

[0093] less weight loss

[0094] A second model of chronic colitis was also used (based on Camoglio et al., *Eur J Immunol* 2000), 2 mg/mice of TNBS injected by intra-rectal route on days 0, 7, and 14 induce a serious chronic colitis. The schedule of injection is drawn on figure. 3. The intake of bacteria (LAB) on days -5 to -1, on days 5 and 12 (FIG. 3) will show macroscopic and histological effects, which are scored on day 16, according to standard scores (Wallace and Ameho respectively).

[0095] The results obtained clearly showed an improvement of the colitis symptoms evidenced (see FIG. 4) by

[0096] absence of necrotic lesions

[0097] reduction of the increase of the submucosa

[0098] reduction of the inflammation

[0099] A third acute model of mice colitis was also set up. 100 mg/kg mice of TNBS is injected on day 0. This will evoke an acute colitis. The schedule of injection is drawn on FIG. 5. The intake of *Bifidobacterium lactis* PTA-4802 or *Lactobacillus plantarum* PTA-4799 (10^7 bacteria/mouse) on days -5 to 0 show macroscopic and histological effects which are scored on day 2.

[0100] Clear macroscopic improvement of the colitis symptoms were evidenced for strains *Bifidobacterium lactis* PTA-4802 and *Lactobacillus plantarum* PTA-4799 (FIG. 6).

[0101] In general all the TNBS-induced colitis models in mice show that feeding LAB to these mice significantly reduced the level of intestinal translocation of indigenous bacteria from the intestinal flora into mesenteric lymph nodes and spleen after the induction of colitis. In addition, no translocation of probiotic bacterial strains according to the invention was observed.

3/Resistance to Pepsin

[0102] The aim of this test is to evaluate the resistance of bacteria to the passage of the gastric barrier by determining the number of bacteria that survive after cultivation in the presence of a pepsin. solution in acid environment. In short, 100 μl of a stock culture frozen at -80°C . is inoculated in 10 ml of MRS (Man Rogosa Sharp) culture medium and incubated overnight at 30°C . or 37°C . Then cells are

washed 3 times in PBS at pH7 and the pellet is suspended in 1 ml PBS buffered and inoculated in 200 μ l aliquots in 4 tubes with 1 ml of filtered pepsin solution at pH 2 supplemented with 300 μ l of NaCl (0.5%). Cell count is performed on 100 μ l aliquots taken from the tubes incubated at T0 (time of inoculation), T0+5min; T0+20min; T0+40min; T0+60min and dilutions are plated on MRS (100 μ l) of 10^5 , 10^6 et 10^7 (consecutive 10 fold dilutions made in 1 ml Ringer solution). The percentage surviving bacteria is. calculated for each incubation time from the reading of the dilution series.

[0103] All results displayed in the tables are the result of triplicate experiments

Strains	5 min	20 min	40 min	60 min
<i>L. acidophilus</i> PTA-4797	91	74	73	55
<i>L. plantarum</i> PTA-4799	100	100	95	42
<i>L. salivarius</i> PTA-4800	100	66	23	7
<i>L. paracasei</i> PTA-4798	100	17.5	17.5	1.5

Results as % of survival compared to T=0.

4/Resistance to Pancreatin

[0104] The aim of this test is to evaluate the resistance of bacteria to the passage of the intestinal transit by determining the number of bacteria that survive after cultivation in the presence of a pancreatin solution in basic environment. In short, 100 μ l of a stock culture frozen at -80° C. is inoculated in 10 ml of MRS (Man Rogosa Sharp) culture medium and incubated overnight at 30° C. or 37° C. Cells are washed 3 times in PBS at pH7 and the pellet is suspended in 1 ml PBS buffered and inoculated in 200 μ l aliquots in 4 tubes with 1 ml of filtered pancreatin. solution at pH 8 supplemented with 300 μ l of NaCl (0.5%). Cell count is performed on 100 μ l aliquots taken from the tubes incubated at T0(time of inoculation), T0+5min; T0+20min; T0+40min; T0+60min;

[0105] T0+120min and dilutions are plated on MRS (100 μ l) of 10^5 , 10^6 , 10^7 (consecutive 10 fold dilutions made in 1 ml Ringer solution). The percentage surviving bacteria is calculated for each incubation. time from the reading of the dilution series. All results displayed in the tables are the result of triplicate experiments.

Strains	5 min	20 min	40 min	60 min	120 min
<i>L. acidophilus</i> PTA-4797	83	100	100	170	170
<i>L. salivarius</i> PTA-4800	100	115	152	158	155
<i>L. plantarum</i> PTA-4799	86	72	79	84	87
<i>L. paracasei</i> PTA-4798	65	100	100	100	100

Results as % of survival compared to T=0.

5/Resistance to Bile Salts

[0106] The experimental is procedure is as follows. 100 μ l culture of a stock frozen at -80° C. is inoculated in 10 ml de

milieu de culture MRS (Man Rogosa Sharp)and incubated overnight at 30° C. or 37° C. The optical density(OD) is measured and a dilution is prepared (OD=0.05 to 0.1) in two bottles. One of the culture bottles is supplemented with a 0.3% solution of bile salts (750 μ l of a filtered 12% solution) and incubated at the correct temperature. The OD is regularly measured until a OD of 0.3 is reached in the control bottle without bile salts. At that point the OD of the supplemented bottle is measured and one starts to measure the time needed to obtain the same OD of 0.3. The resistance of-the strain is therefore evaluated as the delay in growth time.

[0107] Results are coded as sensitive (0 ; more than 60 minutes), tolerant (1 ; between 15 and 60 minutes) or resistant (2 ; less than 15 minutes).

[0108] All results displayed in the tables are the results of triplicate experiments

Strains	Resistance
<i>L. acidophilus</i> PTA-4797	1
<i>L. plantarum</i> PTA-4799	2
<i>L. paracasei</i> PTA-4798	2

6/Hydrophobicity and Affinity for Organic Solvents

[0109] The MATS test as described by Rosdenberg et al., 1980 (FEMS Microbiol. Letters 9 ;29-33) has been used. In short the test measures (grown overnight under normal in vitro growth conditions and washed twice in PBS) the hydrophobicity and polarity of the bacterial cell wall, based on the repartition (measured by optical density) of the bacteria into two non mixable liquid aqueous phases and five respective solvents. Growth medium is MRS (Man Rogosa Sharp) and the aqueous phase is a buffer as. PBS. The solvents were decane, hexadecane, ethyl acetate, chloroform and xylene.

[0110] The affinity for chloroform, acid solvent, reflects the electron donor nature of the bacterium

(Basic Reaction)

[0111] The affinity for Ethyl Acetate, basic solvent, reflects the electron acceptor nature of the bacterium

(Acid Reaction)

[0112] The affinity for apolar solvents (decane, hexadecane, xylene) reflects the hydrophobic character of the bacterium.

[0113] Possibly a high hydrophobicity is related to the presence of glycoproteins on the cell- wall surface, while a low hydrophobicity could be linked to the presence of polysaccharides on the cell wall surface.

[0114] % D'HYDROPHOBICITY=(1-A/AO) $\times$ 100 with

[0115] AO: Original OD at 540 nm (set more or less at OD=0.6 in PBS)

[0116] A: Optical density of aqueous phase (for 1 ml) after vortexing and stabilisation period

Strains	Decane	Hexa-decane	Ethyl acetate	Chloro-form	Xylene
<i>L. acidophilus</i> PTA-4797	81.9	60.4	44.2	90.3	90.9
<i>L. salivarius</i> PTA-4800	23.02	38.17	2.25	79.28	51.18
<i>L. paracasei</i> PTA-4798	36.5	37.4	28.8	45.2	30.6
<i>L. plantarum</i> PTA-4799	25.3	25.6	28.7	28.8	21.9

1-21. (canceled)

22. A bacterial composition having immunomodulation properties comprising *Lactobacillus acidophilus* PTA4797.

23. The bacterial composition according to claim 22 comprising at least one additional strain selected from the group consisting of *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798 and *Bifidobacterium bifidum* PTA-4801.

24. The bacterial composition according to claim 22 comprising at least two additional strains selected from the group consisting of *Lactobacillus acidophilus*, *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA4800, *Lactobacillus paracasei* PTA-4798 and *Bifidobacterium bifidum* PTA-4801.

25. The bacterial composition according to claim 22 comprising a blend of *Lactobacillus plantarum* PTA-4799, *Lactobacillus salivarius* PTA4800, *Lactobacillus paracasei* PTA-4798, *Bifidobacterium lactis* PTA-4802 and *Bifidobacterium bifidum* PTA-4801.

26. The bacterial composition according to claim 22, comprising from 10^6 to 10^{11} CFU of bacteria/g of composition.

27. The bacterial composition according to claim 26, comprising from 10^8 to 10^{11} CFU of bacteria/g of composition.

28. A process for the prevention or the reduction of inflammatory bowel disease, comprising the step of admin-

istering to a human or an animal an efficient preventing or reducing amount of inflammatory bowel disease, of a bacterial composition as defined in claim 22.

29. A process for maintaining the homeostatis of the immune system comprising the step of administering to a human or an animal an efficient maintaining amount of the homeostasis of the immune system, of a bacterial composition as defined in claim 22.

30. A process for the prevention or the reduction of allergenicity comprising the step of administering to a human or an animal an efficient preventing or reducing amount of allergenicity, of a bacterial composition as defined in claim 22.

31. An immunoadjuvant composition, comprising a bacterial composition as defined in claim 22.

32. An immunomodulation method comprising the step of using *Lactobacillus acidophilus* PTA-4797.

33. A food, dietary supplement or pharmaceutical composition comprising a composition as defined in claim 22.

34. The composition according to claim 33, being in the form of a dairy product.

35. The composition according to claim 34, being in the form of an infant formula.

36. The composition of claim 34, having immunomodulation properties.

37. A pharmaceutical composition comprising the composition as defined in claim 22 for the prevention of inflammatory bowel disease.

38. The immunomodulation method according to claim 32, further comprising at least one additional strain selected from the group consisting of *Lactobacillus plantarum* PTA4799, *Lactobacillus salivarius* PTA-4800, *Lactobacillus paracasei* PTA-4798, and *Bifidobacterium bifidum* PTA4801.

39. A pharmaceutical composition comprising the composition as defined in claim 22 for immunomodulation.

* * * * *