



- (51) **International Patent Classification:**
A61K 39/00 (2006.01) *A61K 39/395* (2006.01)
C12Q 1/00 (2006.01)
- (21) **International Application Number:**
PCT/US2013/021306
- (22) **International Filing Date:**
11 January 2013 (11.01.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/586,660 13 January 2012 (13.01.2012) US
- (71) **Applicant** (for all designated States except AL, AT, BE, BG, CH, CN, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IN, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR): **GENENTECH, INC.** [US/US]; 1 Dna Way, South San Francisco, CA 94080 (US).
- (71) **Applicant** (for AL, AT, BE, BG, CH, CN, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IN, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR only): **F. HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (72) **Inventors:** **BAIS, Carlos**; 1 DNA Way, South San Francisco, CA 94080 (US). **BRAUER, Matthew**; 1 DNA Way, South San Francisco, CA 94080 (US). **SCHMIDT, Maik**; 1 DNA Way, South San Francisco, CA 94080 (US). **SINGH, Mallika**; 1 DNA Way, South San Francisco, CA 94080 (US).
- (74) **Agent:** **MICHAUD, Susan M.**; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))



(54) **Title:** BIOLOGICAL MARKERS FOR IDENTIFYING PATIENTS FOR TREATMENT WITH VEGF ANTAGONISTS

(57) **Abstract:** Methods and compositions to detect expression of one or more biomarkers for identifying and treating patients who are likely to be responsive to VEGF antagonist therapy are disclosed. Also disclosed are kits and articles of manufacture for use in the methods.

5 **BIOLOGICAL MARKERS FOR IDENTIFYING PATIENTS FOR TREATMENT
WITH VEGF ANTAGONISTS**

Field of the Invention

The present invention is directed to methods for identifying patients that will benefit from treatment with a VEGF antagonist, *e.g.*, an anti-VEGF antibody.

10 **Background of the Invention**

Measuring expression levels of biomarkers (*e.g.*, secreted proteins in plasma) can be an effective means to identify patients and patient populations that will respond to specific therapies including, *e.g.*, treatment with VEGF antagonists, such as anti-VEGF antibodies.

15 There is a need for effective means for determining which patients will respond to which treatment and for incorporating such determinations into effective treatment regimens for patients with VEGF antagonist therapies, whether used as single agents or combined with other agents.

20 **Summary of the Invention**

The present invention provides methods for identifying patients who will benefit from treatment with a VEGF antagonist, such as an anti-VEGF antibody. These patients are identified based on expression levels of the genes set forth in Table 1 or 2.

25 Accordingly, one embodiment of the invention provides methods of determining whether a patient is likely to respond to treatment with the VEGF antagonist, the methods comprising (a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient prior to any administration of a VEGF antagonist to the patient; and (b) comparing the expression level of the at least one gene to a reference expression level of the at least one gene, wherein a change in the level of
30 expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist.

A further embodiment of the invention provides methods of optimizing therapeutic efficacy of a VEGF antagonist for a patient, the methods comprising (a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient
35 prior to any administration of a VEGF antagonist to the patient; and (b) comparing the

expression level of the at least one gene to a reference expression level of the at least one gene, wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist.

5 Referring to the embodiments set forth above, in some further embodiments, the patient is in a population of patients being tested for responsiveness to a VEGF antagonist and the reference level is the median level of expression of the at least one gene in the population of patients. In some embodiments, the change in level of expression of the at least one gene in the patient sample is an increase relative to the reference level. In some
10 embodiments, the change in level of expression of the at least one gene in the patient sample is a decrease relative to the reference level. In some embodiments, the at least one gene in the biological sample obtained from the patient is detected by measuring mRNA. In some embodiments, expression of the at least one gene in the biological sample obtained from the patient is detected by measuring plasma protein levels. In some embodiments, the biological
15 sample is tumor tissue. In some embodiments, the methods further comprise detecting expression of at least a second, third, fourth, or more gene set forth in Table 1 or 2 in the biological sample from the patient. In some embodiments, the at least one gene is selected from the group consisting of: Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1,
20 VEGFR1, VEGFR2, VEGFR3, and VIM. In some embodiments, the VEGF antagonist is an anti-VEGF antibody, such as bevacizumab. In some embodiments, the patient has an angiogenic disorder. In some embodiments, the patient has cancer selected from the group consisting of: colorectal cancer, breast cancer, lung cancer, glioblastoma, and combinations thereof.

25 Also referring to the embodiments set forth above, the methods can further include (c) selecting a VEGF antagonist for treatment of the patient when a change in the level of expression of the at least one gene in the patient sample relative to the reference level is detected. Further, the methods can include (d) administering a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab) to the patient.

30 Another embodiment of the invention provides methods for selecting a therapy for a particular patient in a population of patients being considered for therapy, the methods comprising: (a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient prior to any administration of a VEGF antagonist

to the patient; (b) comparing the expression level of the at least one gene to a reference expression level of the at least one gene, wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist, and (c) selecting a VEGF antagonist
5 as the therapy if the patient is identified as likely to respond to treatment with a VEGF antagonist; or (d) selecting a therapy that is not a VEGF antagonist if the patient is not identified as likely to respond to treatment with a VEGF antagonist.

In some embodiments, the reference level is the median level of expression of the at least one gene in the population of patients. In some embodiments, the change in level of
10 expression of the at least one gene in the patient sample is an increase relative to the reference level. In some embodiments, the change in level of expression of the at least one gene in the patient sample is a decrease relative to the reference level. In some embodiments, the methods further comprise detecting expression of at least a second, third, fourth, or more gene set forth in Table 1 or 2 in the biological sample from the patient. In some
15 embodiments, the therapy of (d) is an agent selected from the group consisting of: an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, and combinations thereof. In some embodiments, the methods further comprise (e) administering an effective amount of a VEGF antagonist to the patient if the patient is identified as likely to respond to treatment with a VEGF antagonist. In some embodiments, the VEGF antagonist is
20 an anti-VEGF antibody, such as bevacizumab. In some embodiments, the methods further comprising administering an effective amount of at least a second agent. In some embodiments, the second agent is selected from the group consisting of: an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, and combinations thereof.

25 A further embodiment of the invention provides methods of identifying a biomarker for determining responsiveness to a VEGF antagonist, the methods comprising: (a) detecting the expression of a candidate biomarker in a biological sample obtained from a patient prior to administration of a VEGF antagonist to the patient; and (b) comparing the expression of the candidate biomarker to a reference expression level of the candidate biomarker, wherein a
30 change in the level of expression of the candidate biomarker in the patient sample relative to the reference level identifies the candidate biomarker as a biomarker of a patient who will likely respond to treatment with a VEGF antagonist. In some embodiments, the reference level is the median level of expression of the at least one gene in a population of patients

being tested for the likelihood they will respond to a VEGF antagonist. In some embodiments, the reference level is the expression level of the at least one gene in a sample previously obtained from the patient. In some embodiments, the patient was previously treated with a VEGF antagonist and is currently experiencing metastasis. In some
 5 embodiments, the VEGF antagonist is an anti-VEGF antibody, such as bevacizumab. Further, the methods can further include (c) selecting a candidate biomarker having a change in level of expression relative to the reference for use as a biomarker for determining responsiveness to VEGF antagonist treatment.

In another embodiment, the invention provides methods for diagnosing an angiogenic
 10 disorder in a patient, the methods comprising the steps of: (a) detecting the expression level of at least one gene set forth in Table 1 or 2 or a biomarker identified according to methods such as those described above in a sample obtained from the patient prior to any administration of a VEGF antagonist to the patient; and (b) comparing the expression level of the at least one gene or biomarker to a reference level of the at least one gene; wherein a
 15 change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient having an angiogenic disorder. These methods can further include (c) selecting a VEGF antagonist for treatment of the patient when a change in the level of expression of the at least one gene in the patient sample relative to the reference level is detected. In addition, the methods can further include (d) administering a VEGF antagonist
 20 to the patient. In addition, the methods described herein can include a step of obtaining a sample from a patient, as described herein. Further, the methods described herein can be carried out on patients diagnosed with cancer, as described herein, to determine an optimal treatment regimen.

In any of the embodiments set forth above, the change in the level of expression of the
 25 at least one gene or biomarker in the patient sample relative to the reference level can be determined by calculating a VDV signature score (VDV_i) for the patient sample according to the algorithm:

$$VDV_i = \frac{1}{\sqrt{n}} \sum_{g=1}^n Z_{g,i}$$

in which $Z_{g=1,i}, Z_{g=2,i}, \dots, Z_{g=n,i}$ are standardized z-scores of expression values for each gene or biomarker g , from $g=1$ to $g=n$, of the sample i , and in which a VDV_i below a first defined
 30 threshold indicates a decrease relative to the reference level, and a VDV_i above a second

defined threshold indicates an increase relative to the reference level. In some embodiments, the expression values for each gene or biomarker g , from $g=1$ to $g=n$, are qRT-PCR values for each gene g , from $g=1$ to $g=n$. In some embodiments, the first defined threshold is from -4 to -0.5 (e.g., -4, -3.5, -3, -2.5, -2, -1.5, -1, or -0.5) and the second defined threshold is from 0.5 to 4 (e.g., 0.5, 1, 1.5, 2, 2.5, 3, 3.5, or 4). In some embodiments, the first defined threshold is from -4 to -1 (e.g., -4, -3.5, -3, -2.5, -2, -1.5, or -1) and the second defined threshold is from 1 to 4 (e.g., 1, 1.5, 2, 2.5, 3, 3.5, or 4). In some embodiments, the first defined threshold is from -4 to -1.5 (e.g., -4, -3.5, -3, -2.5, -2, or -1.5) and the second defined threshold is from 1.5 to 4 (e.g., 1.5, 2, 2.5, 3, 3.5, or 4). In other embodiments, the first defined threshold is from -4 to -2 (e.g., -4, -3.5, -3, -2.5, or -2) and the second defined threshold is from 2 to 4 (e.g., 2, 2.5, 3, 3.5, or 4).

These and other embodiments are further described by the detailed description that follows.

Brief Description of the Drawings

Fig. 1A shows histological and graphical analysis of micro-vascular density (MVD) and proliferation index determination of murine pancreatic neuro-endocrine tumor (PNET) cells at 72 hours, 7 days, and 14 days post-treatment with anti-VEGF mAb. Representative images from histological analysis of tumor vessel density via MECA-32 staining (left) and proliferative index via Ki67 (right) at various times following anti-VEGF treatment (20X magnification). Quantitation from 4-6 tumors in each case is shown in the bar graphs below as mean \pm SEM. * $P < 0.05$, NS = not significant.

Fig. 1B is a graph showing the kinetics of anti-VEGF effects on tumor burden in the *RIP-T β Ag* model. Tumor burden in anti-VEGF-treated mice (red bars) is significantly lower than control-treated mice (black bars) at 21 days, but not at 14 days on the study. * = $p < 0.05$ (t-test), $n = 5-8$ mice/group/time point.

Fig. 1C is a graph showing changes in VDV gene expression levels in response to treatment with a VEGF antagonist. Expression levels of genes (shown as red lines) decrease significantly relative to all genes (grey histogram). The dashed red line indicates the mean change for these selected genes. The black dashed line indicates the mean fold change for the rest of the genes.

Fig. 1D is a graph showing changes in gene expression levels in a subgroup of genes in response to treatment with a VEGF antagonist as assessed by qPCR. Bars represent mean expression from three independent biological replicates. Error bars = \log_2 standard deviation.

Fig. 2A is a set of graphs showing changes in VDV gene response to VEGF blockade in an established subcutaneous breast carcinoma tumor model (MDA-MB-231). Tumor samples were collected 24 hours after anti-VEGF or control treatment. Genes in the VDV signature (red lines) decrease significantly relative to all genes (shown as a grey histogram) in the stroma (upper graph, mouse chip, $p < 0.0001$), but not in the tumor cells (lower graph, human chip, no significant differences). Individual proxVDV transcript fold-changes are annotated in black letters in the microarray density plot. $n = 5-10$ instances for each treatment cohort.

Fig. 2B is a set of graphs showing changes in VDV gene response to VEGF blockade in an orthotopic (intracranial) U87 glioblastoma model. Tumor samples were collected after 13-42 days of anti-VEGF or control treatment. Genes in the VDV signature (red lines) decrease significantly relative to all genes (shown as a grey histogram) in the stroma (upper graph, mouse chip, $p < 0.0105$), but not in the tumor cells (lower graph, human chip, no significant differences). Individual proxVDV transcript fold-changes are annotated in black letters in the microarray density plot. $n = 5-10$ instances for each treatment cohort.

Fig. 2C is a set of graphs showing a down-modulation is observed upon topical anti-VEGF application to a skin wound (upper graph, $p = 0.0125$) and the converse up-regulation, when recombinant VEGF was applied for 12 hours ($p < 0.0001$). Individual proxVDV transcript fold-changes are annotated in black letters.

Fig. 2D presents histological data and graphs showing VEGF signaling induces VDV gene expression. In contrast to anti-VEGF downregulation of the VDV gene signature (right, upper graph, $p < 0.0001$), anti-Dll4 treatment causes a upregulation of a majority of VDV genes (right, lower graph, $p < 0.0001$) after 48 hours in an MDA-MB-231 model consistent with hypervascularization evident by immunofluorescent staining for CD31/PECAM as compared to control treatment (left). Individual proxVDV transcript fold-changes are annotated in black letters. $n = 5-10$ instances for each treatment cohort.

Fig. 3 is a heat map showing that most proxVDV genes are not obviously up-regulated by rVEGF *in vitro*. Gene expression analyses of rVEGF stimulation of HUVECs. H (hours) indicates length of rVEGF stimulation. The heat map shown here highlights the results of microarray expression analyses for selected VDV probes. Dark blue represents

maximal relative down-regulation and dark red the maximal degree of transcript up-regulation. ProxVDV genes are not markedly regulated by rVEGF *in vitro*. However, a yet uncharacterized small group of VDV genes (EHD3, PCHD17, and THBD) seem to be strongly up-regulated upon rVEGF stimulation of HUVECs. Expression data for each time-point is from three independent replicates.

Fig. 4A is a *in situ* hybridization (ISH) images showing that the proxVDV gene ESM1 is an *in vivo* VEGF target specifically expressed in tumor-associated vasculature. Top pictures (left and right): ISH negative controls with sense oligos show no significant background (non-specific staining). Bottom pictures show ESM1 mRNA expression (by ISH with anti-sense oligos) in HM7 tumor sections from anti-VEGF or control treated animals. Black arrows indicate several areas of strong ESM1 mRNA vascular expression (brown staining) in control treated tumor slides (bottom picture slide left). In contrast, ESM1 was almost undetectable in tumor slides from anti-VEGF treated animals (bottom picture, right). All slides were also counter stained with Haematoxylin Eosin (H&E).

Fig. 4B is a graph showing quantification of ESM1 (ISH) staining in tumor slides from control and anti-VEGF-treated animals. n=10.

Fig. 4C is a graph showing quantification of MECA32 (PLVAP) staining in tumor slides from control and anti-VEGF-treated animals. n=10.

Fig. 4D is a set of histological images and corresponding quantitative graph showing evidence for the *in vivo* activity of VEGF pathway inhibitors in MDA-MB-231 tumors. Anti-VEGF mAb, sunitinib, or axitinib *in vivo* treatment efficiently reduces MVD in tumors 72 hour post-treatment. Animals bearing MDA-MB-231 tumors were treated as indicated in materials and methods for 72 hours and then tumors were collected for histological and gene expression analyses. Top images: tumor vessel density via MECA-32 (PLVAP) and CD31 staining (red). Nuclei were counterstained with DAPI (blue). Images were taken at 20X magnification. Bottom graph shows quantification (as mean +/- SEM) from 8 tumors in each treatment group. *P<0.05.

Fig. 4E is a set of graphs showing that *in vivo* VEGF blockade or VEGFR-2 downstream signaling inhibition induces consistent downregulation of proxVDV genes. qRT-PCR analysis of gene expression in 400 mm³ MDA-MB-231 xenograft tumors collected 8 (bottom panel), 16 (middle panel), or 72 hours (top panel) after treatment with VEGF and VEGFR-2 inhibitors (sunitinib and axitinib). Values represent the mean log₂ fold change in relative gene expression induced by VEGF/VEGFR-2 inhibitor when compared to control

treatment mean gene expression. Non-vascular markers such as E-cadh and CD45 do not change significantly in response to these inhibitors. Bottom panel (8 hours post-treatment) shows only anti-VEGF treatment because Axitinib and sunitinib did not have evident activity at this particular time point. Gene expression data represent mean of 8 biological replicates
5 for each treatment. Error bars represent standard deviation.

Fig. 5A are graphs showing the consistent proxVDV downregulation by multiple VEGF pathway inhibitors. Analysis of gene expression in MDA-MB-231 xenograft tumors collected 8, 16, or 72 hours after treatment with VEGF and VEGFR-2 inhibitors (sunitinib and axitinib). Values represent the mean of the \log_2 fold change in relative gene expression
10 induced by VEGF/VEGFR-2 inhibitor compared to control treatment. Gene expression data represent the \log_2 mean of 8 biological replicates for each treatment. Error bars represent standard deviation.

Fig. 5B is a graph showing the quantification of proxVDV gene expression by qRT-PCR in endothelial cells sorted from MDA-MB-231 xenograft tumors treated with ragweed
15 or anti-VEGF mAb. Values represent the mean of the \log_2 fold change of 3 replicates. Error bars represent standard deviation.

Fig. 6 is a graph showing enrichment of VDV markers in Tumor-Associated Endothelial Cells (TAECs) by *ex vivo* enrichment expression analyses of VDV transcripts in freshly FACS isolated TAEC versus GFP-MDA-MB-231 tumor cells. Relative gene
20 expression in TAEC versus tumor cells was measured by qRT-PCR of selected genes. All VDV genes tested were highly enriched (25-200 fold) in TAECs. By contrast Zeb1 mRNA expression (an epithelial marker) was decreased in TAECs and enriched in tumor cells. Values represent the mean of the relative \log_2 fold gene enrichment when comparing TAECs to tumor cells. TAEC cells were sorted as CD31 positive, CD45 negative, and GFP negative
25 cells. Tumor cells were sorted by GFP positivity. Gene expression data represent the mean of six tumors that were pooled for each FACS sorting experiment. qRT-PCR was run in triplicates. Error bars represent standard deviation.

Fig. 7 is a graph showing the change in gene expression (post- vs pre-treatment) in biopsy samples from 19 inflammatory breast cancer patients. Genes in the VDV signature
30 (red lines) decrease significantly relative to all genes (grey histogram), $p=0.0275$.

Fig. 8A is a set of graphs showing progression-free (top) and overall survival (bottom) of 103 colorectal cancer patients with available pre-treatment mRNA in the NO16966 trial.

Fig. 8B is a schematic of the experiment overview for quantifying gene expression of 22 VDV genes using the angiogenesis Fluidigm qRT-PCR chip.

Fig. 8C is a graph showing the correlation of expression levels of the 22 VDV genes in colon cancer samples.

5 Fig. 8D is a set of graphs showing the 22-gene VDV signature stratifies the effect of bevacizumab (bev) treatment for patients with advanced colorectal cancer. Shown are progression-free (top) or overall survival (bottom) of “VDV-high” (solid lines) versus “VDV-low” patients (dashed lines), which were treated with XELOX (black) or XELOX+bevacizumab (Mullen et al. *Cell*. 147(3): 565-576, 2011). PFS (top) VDV gene set
10 effect, stratified by expression levels (interaction $p=0.036$), and OS (bottom) VDV gene set effect stratified by expression levels (interaction $p=0.37$).

Fig. 9A is a set of graphs verifying concordance between the full VDV signature “VDV” (x-axis) and the 22-gene representative subset “VDV-22” (y-axis) used to interrogate archival clinical material. A prevalence sample set is shown with archival samples from
15 patients that would qualify for metastatic first line trial setting (comparable to NO16966). Archival samples were assessed for whole genome RNA expression on Illumina DASL bead arrays; NSCLC = Non-small cell lung carcinoma, BR = Breast, CRC = Colorectal carcinoma.

Fig. 9B is a set of graphs showing the stratification of NO16966 patients by VEGF-A expression levels, PFS (upper), interaction $p=0.76$ and OS (below), interaction $p=0.33$.

20 Fig. 9C is a set of graphs showing the stratification of NO16966 patients by CD31 expression levels, PFS (upper) interaction $p=0.15$ and OS (lower), interaction $p=0.99$.

Detailed Description of the Preferred Embodiments

I. Introduction

The present invention provides methods and compositions for monitoring and/or identifying patients sensitive or responsive to treatment with VEGF antagonists, *e.g.*, an anti-VEGF antibody. The invention is based on the discovery that determination of expression levels of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 before treatment with a VEGF antagonist (such as an anti-VEGF antibody) is useful for identifying patients sensitive to or responsive to treatment with a VEGF antagonist, *e.g.*, an anti-VEGF antibody. Optionally, VEGF antagonist therapy can then be selected for the patients and, further, VEGF antagonist therapy can optionally be administered to the patients.

II. Definitions

The terms “biomarker” and “marker” are used interchangeably herein to refer to a DNA, RNA, protein, carbohydrate, or glycolipid-based molecular marker, the expression or presence of which in a subject’s or patient’s sample can be detected by standard methods (or methods disclosed herein) and is useful for monitoring the responsiveness or sensitivity of a mammalian subject to a VEGF antagonist. Such biomarkers include, but are not limited to, the genes set forth in Tables 1 and 2. Expression of such a biomarker may be determined to be higher or lower in a sample obtained from a patient sensitive or responsive to a VEGF antagonist than a reference level (including, *e.g.*, the median expression level of the biomarker in a samples from a group/population of patients being tested for responsiveness to a VEGF antagonist; the level in a sample previously obtained from the individual at a prior time; or the level in a sample from a patient who received prior treatment with a VEGF antagonist (such as an anti-VEGF antibody) in a primary tumor setting, and who now may be experiencing metastasis). Individuals having an expression level that is greater than or less than the reference expression level of at least one gene, such as those set forth in Tables 1 and 2, can be identified as subjects/patients likely to respond to treatment with a VEGF antagonist. For example, such subjects/patients who exhibit gene expression levels at the most extreme 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% relative to (*i.e.*, higher or lower than) the reference level (such as the median level, noted above), can be identified as subjects/patients likely to respond to treatment with a VEGF antagonist, such as an anti-VEGF antibody.

The terms “sample” and “biological sample” are used interchangeably to refer to any biological sample obtained from an individual including body fluids, body tissue (*e.g.*, tumor tissue), cells, or other sources. Body fluids are, *e.g.*, lymph, sera, whole fresh blood, peripheral blood mononuclear cells, frozen whole blood, plasma (including fresh or frozen),
5 urine, saliva, semen, synovial fluid and spinal fluid. Samples also include breast tissue, renal tissue, colonic tissue, brain tissue, muscle tissue, synovial tissue, skin, hair follicle, bone marrow, and tumor tissue. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art.

An “effective response” of a patient or a patient’s “responsiveness” or “sensitivity” to
10 treatment with a VEGF antagonist refers to the clinical or therapeutic benefit imparted to a patient at risk for or suffering from an angiogenic disorder from or as a result of the treatment with the VEGF antagonist, such as an anti-VEGF antibody. Such benefit includes cellular or biological responses, a complete response, a partial response, a stable disease (without progression or relapse), or a response with a later relapse of the patient from or as a result of
15 the treatment with the antagonist. For example, an effective response can be reduced tumor size or progression-free survival in a patient diagnosed as expressing one or more of the biomarkers set forth in Table 1 or 2 versus a patient not expressing one or more of the biomarkers. The expression of genetic biomarker(s) effectively predicts, or predicts with high sensitivity, such effective response.

20 “Antagonists” as used herein refer to compounds or agents which inhibit or reduce the biological activity of the molecule to which they bind. Antagonists include antibodies, synthetic or native-sequence peptides, immunoadhesins, and small-molecule antagonists that bind to VEGF, optionally conjugated with or fused to another molecule. A “blocking” antibody or an “antagonist” antibody is one which inhibits or reduces biological activity of
25 the antigen it binds.

An “agonist antibody,” as used herein, is an antibody which partially or fully mimics at least one of the functional activities of a polypeptide of interest.

The term “antibody” herein is used in the broadest sense and specifically covers monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.*, bispecific
30 antibodies) formed from at least two intact antibodies, and antibody fragments so long as they exhibit the desired biological activity.

An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its

natural environment are materials which would interfere with research, diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In some embodiments, an antibody is purified (1) to greater than 95% by weight of antibody as determined by, for example, the Lowry method, and in some embodiments, to greater than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of, for example, a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using, for example, Coomassie blue or silver stain. Isolated antibody includes the antibody *in situ* within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

"Native antibodies" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light-chain and heavy chain variable domains.

The "variable region" or "variable domain" of an antibody refers to the amino-terminal domains of the heavy or light chain of the antibody. The variable domain of the heavy chain may be referred to as "VH." The variable domain of the light chain may be referred to as "VL." These domains are generally the most variable parts of an antibody and contain the antigen-binding sites.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions (HVRs) both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the

framework regions (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a beta-sheet configuration, connected by three HVRs, which form loops connecting, and in some cases forming part of, the beta-sheet structure. The HVRs in each chain are held together in close proximity by the FR regions and, with the HVRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, Fifth Edition, National Institute of Health, Bethesda, MD (1991)). The constant domains are not involved directly in the binding of an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

The “light chains” of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

Depending on the amino acid sequences of the constant domains of their heavy chains, antibodies (immunoglobulins) can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), *e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgA₁, and IgA₂. The heavy chain constant domains that correspond to the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known and described generally in, for example, Abbas *et al.* *Cellular and Mol. Immunology*, 4th ed. (W. B. Saunders, Co., 2000). An antibody may be part of a larger fusion molecule, formed by covalent or non-covalent association of the antibody with one or more other proteins or peptides.

The terms “full-length antibody,” “intact antibody,” and “whole antibody” are used herein interchangeably to refer to an antibody in its substantially intact form, not antibody fragments as defined below. The terms particularly refer to an antibody with heavy chains that contain an Fc region.

A “naked antibody” for the purposes herein is an antibody that is not conjugated to a cytotoxic moiety or radiolabel.

“Antibody fragments” comprise a portion of an intact antibody, preferably comprising the antigen-binding region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields a $F(ab')_2$ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

5 "Fv" is the minimum antibody fragment which contains a complete antigen-binding site. In one embodiment, a two-chain Fv species consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv (scFv) species, one heavy- and one light-chain variable domain can be covalently linked by a flexible peptide linker such that the light and heavy chains can associate in a "dimeric"
10 structure analogous to that in a two-chain Fv species. It is in this configuration that the three HVRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six HVRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three HVRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower
15 affinity than the entire binding site.

The Fab fragment contains the heavy- and light-chain variable domains and also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from
20 the antibody-hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. $F(ab')_2$ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

"Single-chain Fv" or "scFv" antibody fragments comprise the VH and VL domains
25 of an antibody, wherein these domains are present in a single polypeptide chain. Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domains that enables the scFv to form the desired structure for antigen binding. For a review of scFv, see, e.g., Pluckthün, in *The Pharmacology of Mono-clonal Antibodies*, vol. 113, Rosenberg and Moore eds. (Springer-Verlag, New York: 1994), pp 269-315.

30 The term "diabodies" refers to antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) in the same polypeptide chain (VH-VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to

pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies may be bivalent or bispecific. Diabodies are described more fully in, for example, EP 404,097; WO 1993/01161; Hudson *et al.*, *Nat. Med.* 9:129-134 (2003); and Hollinger *et al.*, *PNAS USA* 90: 6444-6448 (1993). Triabodies and tetrabodies are also described in
5 Hudson *et al.*, *Nat. Med.* 9:129-134 (2003).

The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible mutations, *e.g.*, naturally occurring mutations, that may be present in minor amounts. Thus, the modifier
10 “monoclonal” indicates the character of the antibody as not being a mixture of discrete antibodies. In certain embodiments, such a monoclonal antibody typically includes an antibody comprising a polypeptide sequence that binds a target, wherein the target-binding polypeptide sequence was obtained by a process that includes the selection of a single target binding polypeptide sequence from a plurality of polypeptide sequences. For example, the
15 selection process can be the selection of a unique clone from a plurality of clones, such as a pool of hybridoma clones, phage clones, or recombinant DNA clones. It should be understood that a selected target binding sequence can be further altered, for example, to improve affinity for the target, to humanize the target-binding sequence, to improve its production in cell culture, to reduce its immunogenicity *in vivo*, to create a multispecific
20 antibody, *etc.*, and that an antibody comprising the altered target binding sequence is also a monoclonal antibody of this invention. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal-antibody preparation is directed against a single determinant on an antigen. In addition to their specificity, monoclonal-antibody preparations
25 are advantageous in that they are typically uncontaminated by other immunoglobulins.

The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of
30 techniques, including, for example, the hybridoma method (*e.g.*, Kohler and Milstein., *Nature*, 256:495-97 (1975); Hongo *et al.*, *Hybridoma*, 14 (3): 253-260 (1995), Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier,

N.Y., 1981)), recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567), phage-
display technologies (see, e.g., Clackson *et al.*, *Nature*, 352: 624-628 (1991); Marks *et al.*, *J.*
Mol. Biol. 222: 581-597 (1992); Sidhu *et al.*, *J. Mol. Biol.* 338(2): 299-310 (2004); Lee *et al.*,
J. Mol. Biol. 340(5): 1073-1093 (2004); Fellouse, *PNAS USA* 101(34): 12467-12472 (2004);
5 and Lee *et al.*, *J. Immunol. Methods* 284(1-2): 119-132(2004), and technologies for
producing human or human-like antibodies in animals that have parts or all of the human
immunoglobulin loci or genes encoding human immunoglobulin sequences (see, e.g., WO
1998/24893; WO 1996/34096; WO 1996/33735; WO 1991/10741; Jakobovits *et al.*, *PNAS*
USA 90: 2551 (1993); Jakobovits *et al.*, *Nature* 362: 255-258 (1993); Bruggemann *et al.*,
10 *Year in Immunol.* 7:33 (1993); U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126;
5,633,425; and 5,661,016; Marks *et al.*, *Bio/Technology* 10: 779-783 (1992); Lonberg *et al.*,
Nature 368: 856-859 (1994); Morrison, *Nature* 368: 812-813 (1994); Fishwild *et al.*, *Nature*
Biotechnol. 14: 845-851 (1996); Neuberger, *Nature Biotechnol.* 14: 826 (1996); and Lonberg
and Huszar, *Intern. Rev. Immunol.* 13: 65-93 (1995).

15 The monoclonal antibodies herein specifically include “chimeric” antibodies in which
a portion of the heavy and/or light chain is identical with or homologous to corresponding
sequences in antibodies derived from a particular species or belonging to a particular
antibody class or subclass, while the remainder of the chain(s) is identical with or
homologous to corresponding sequences in antibodies derived from another species or
20 belonging to another antibody class or subclass, as well as fragments of such antibodies, so
long as they exhibit the desired biological activity (e.g., U.S. Pat. No. 4,816,567 and
Morrison *et al.*, *PNAS USA* 81:6851-6855 (1984)). Chimeric antibodies include
PRIMATIZED® antibodies wherein the antigen-binding region of the antibody is derived
from an antibody produced by, e.g., immunizing macaque monkeys with the antigen of
25 interest.

“Humanized” forms of non-human (e.g., murine) antibodies are chimeric antibodies
that contain minimal sequence derived from non-human immunoglobulin. In one
embodiment, a humanized antibody is a human immunoglobulin (recipient antibody) in
which residues from a HVR of the recipient are replaced by residues from a HVR of a non-
30 human species (donor antibody) such as mouse, rat, rabbit, or nonhuman primate having the
desired specificity, affinity, and/or capacity. In some instances, FR residues of the human
immunoglobulin are replaced by corresponding non-human residues. Furthermore,
humanized antibodies may comprise residues that are not found in the recipient antibody or in

the donor antibody. These modifications may be made to further refine antibody performance. In general, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin, and all, or substantially all, of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally will also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see, *e.g.*, Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992). See also, for example, Vaswani and Hamilton, *Ann. Allergy, Asthma & Immunol.* 1:105-115 (1998); Harris, *Biochem. Soc. Transactions* 23:1035-1038 (1995); Hurle and Gross, *Curr. Op. Biotech.* 5:428-433 (1994); and U.S. Pat. Nos. 6,982,321 and 7,087,409.

A “human antibody” is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks *et al.*, *J. Mol. Biol.*, 222:581 (1991). Also available for the preparation of human monoclonal antibodies are methods described in Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boerner *et al.*, *J. Immunol.*, 147(1):86-95 (1991). See also van Dijk and van de Winkel, *Curr. Opin. Pharmacol.*, 5: 368-74 (2001). Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, *e.g.*, immunized xenomice (see, *e.g.*, U.S. Pat. Nos. 6,075,181 and 6,150,584 regarding XENOMOUSE™ technology). See also, for example, Li *et al.*, *PNAS USA*, 103:3557-3562 (2006) regarding human antibodies generated via a human B-cell hybridoma technology.

The term “hypervariable region,” “HVR,” or “HV,” when used herein refers to the regions of an antibody-variable domain which are hypervariable in sequence and/or form structurally defined loops. Generally, antibodies comprise six HVRs; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). In native antibodies, H3 and L3 display the most diversity of the six HVRs, and H3 in particular is believed to play a unique role in conferring fine specificity to antibodies. See, *e.g.*, Xu *et al. Immunity* 13:37-45 (2000); Johnson and Wu

in *Methods in Molecular Biology* 248:1-25 (Lo, ed., Human Press, Totowa, NJ, 2003).

Indeed, naturally occurring camelid antibodies consisting of a heavy chain only are functional and stable in the absence of light chain. See, e.g., Hamers-Casterman *et al.*, *Nature* 363:446-448 (1993) and Sheriff *et al.*, *Nature Struct. Biol.* 3:733-736 (1996).

5 A number of HVR delineations are in use and are encompassed herein. The HVRs that are Kabat complementarity-determining regions (CDRs) are based on sequence variability and are the most commonly used (Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991)). Chothia refers instead to the location of the structural loops (Chothia
10 and Lesk *J. Mol. Biol.* 196:901-917 (1987)). The AbM HVRs represent a compromise between the Kabat CDRs and Chothia structural loops, and are used by Oxford Molecular's AbM antibody-modeling software. The "contact" HVRs are based on an analysis of the available complex crystal structures. The residues from each of these HVRs are noted below.

	<u>Loop</u>	<u>Kabat</u>	<u>AbM</u>	<u>Chothia</u>	<u>Contact</u>
15	L1	L24-L34	L24-L34	L26-L32	L30-L36
	L2	L50-L56	L50-L56	L50-L52	L46-L55
	L3	L89-L97	L89-L97	L91-L96	L89-L96
	H1	H31-H35B	H26-H35B	H26-H32	H30-H35B (Kabat Numbering)
	H1	H31-H35	H26-H35	H26-H32	H30-H35 (Chothia Numbering)
20	H2	H50-H65	H50-H58	H53-H55	H47-H58
	H3	H95-H102	H95-H102	H96-H101	H93-H101

HVRs may comprise "extended HVRs" as follows: 24-36 or 24-34 (L1), 46-56 or 50-56 (L2), and 89-97 or 89-96 (L3) in the VL, and 26-35 (H1), 50-65 or 49-65 (H2), and 93-102, 94-102, or 95-102 (H3) in the VH. The variable-domain residues are numbered
25 according to Kabat *et al.*, *supra*, for each of these extended-HVR definitions.

"Framework" or "FR" residues are those variable-domain residues other than the HVR residues as herein defined.

The expression "variable-domain residue-numbering as in Kabat" or "amino acid-position numbering as in Kabat," and variations thereof, refers to the numbering system used
30 for heavy-chain variable domains or light-chain variable domains of the compilation of antibodies in Kabat *et al.*, *supra*. Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or HVR of the variable domain. For example, a heavy-chain variable

domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (*e.g.*, residues 82a, 82b, and 82c, *etc.* according to Kabat) after heavy-chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a

5 “standard” Kabat numbered sequence.

An “affinity-matured” antibody is one with one or more alterations in one or more HVRs thereof which result in an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In one embodiment, an affinity-matured antibody has nanomolar or even picomolar affinities for the target antigen. Affinity-matured antibodies are produced by procedures known in the art. 10 For example, Marks *et al.*, *Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH- and VL-domain shuffling. Random mutagenesis of HVR and/or framework residues is described by, for example: Barbas *et al. Proc Nat. Acad. Sci. USA* 91:3809-3813 (1994); Schier *et al. Gene* 169:147-155 (1995); Yelton *et al. J. Immunol.* 155:1994-2004 (1995); 15 Jackson *et al., J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al., J. Mol. Biol.* 226:889-896 (1992).

“Growth-inhibitory” antibodies are those that prevent or reduce proliferation of a cell expressing an antigen to which the antibody binds.

Antibodies that “induce apoptosis” are those that induce programmed cell death, as 20 determined by standard apoptosis assays, such as binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies).

Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native-sequence Fc region or amino acid-sequence-variant Fc region) of an 25 antibody, and vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement- dependent cytotoxicity (CDC); Fc-receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down-regulation of cell-surface receptors (*e.g.*, B-cell receptor); and B-cell activation.

The term “Fc region” herein is used to define a C-terminal region of an 30 immunoglobulin heavy chain, including native-sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine

(residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue.

Unless indicated otherwise herein, the numbering of the residues in an immunoglobulin heavy chain is that of the EU index as in Kabat *et al.*, *supra*. The “EU index as in Kabat” refers to the residue numbering of the human IgG1 EU antibody.

A “functional Fc region” possesses an “effector function” of a native-sequence Fc region. Exemplary “effector functions” include C1q binding; CDC; Fc-receptor binding; ADCC; phagocytosis; down-regulation of cell-surface receptors (*e.g.*, B-cell receptor; BCR), *etc.* Such effector functions generally require the Fc region to be combined with a binding domain (*e.g.*, an antibody-variable domain) and can be assessed using various assays as disclosed, for example, in definitions herein.

A “native-sequence Fc region” comprises an amino acid sequence identical to the amino acid sequence of an Fc region found in nature. Native-sequence human Fc regions include a native-sequence human IgG1 Fc region (non-A and A allotypes); native-sequence human IgG2 Fc region; native-sequence human IgG3 Fc region; and native-sequence human IgG4 Fc region, as well as naturally occurring variants thereof.

A “variant Fc region” comprises an amino acid sequence which differs from that of a native-sequence Fc region by virtue of at least one amino acid modification, preferably one or more amino acid substitution(s). Preferably, the variant Fc region has at least one amino acid substitution compared to a native-sequence Fc region or to the Fc region of a parent polypeptide, *e.g.*, from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native-sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% homology with a native-sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% homology therewith, more preferably at least about 95% homology therewith.

The term “Fc-region-comprising antibody” refers to an antibody that comprises an Fc region. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during purification of the antibody or by recombinant

engineering the nucleic acid encoding the antibody. Accordingly, a composition comprising an antibody having an Fc region according to this invention can comprise an antibody with K447, with all K447 removed, or a mixture of antibodies with and without the K447 residue.

“Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. In some embodiments, an FcR is a native-human FcR. In some embodiments, an FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of those receptors. FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see, e.g., Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed, for example, in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein.

The term “Fc receptor” or “FcR” also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.* 24:249 (1994)) and regulation of homeostasis of immunoglobulins. Methods of measuring binding to FcRn are known (see, e.g., Ghetie and Ward, *Immunology Today*, 18 (12):592-8 (1997); Ghetie *et al.*, *Nature Biotechnology*, 15 (7):637-40 (1997); Hinton *et al.*, *J. Biol. Chem.*, 279(8):6213-6 (2004); WO 2004/92219 (Hinton *et al.*).

Binding to human FcRn *in vivo* and serum half-life of human FcRn high-affinity binding polypeptides can be assayed, e.g., in transgenic mice or transfected human cell lines expressing human FcRn, or in primates to which the polypeptides with a variant Fc region are administered. WO 2000/42072 (Presta) describes antibody variants with improved or diminished binding to FcRs. See, also, for example, Shields *et al.* *J. Biol. Chem.* 9(2): 6591-6604 (2001).

“Human effector cells” are leukocytes which express one or more FcRs and perform effector functions. In certain embodiments, the cells express at least FcγRIII and perform ADCC effector function(s). Examples of human leukocytes which mediate ADCC include

peripheral blood mononuclear cells (PBMC), natural-killer (NK) cells, monocytes, cytotoxic T cells, and neutrophils. The effector cells may be isolated from a native source, *e.g.*, from blood.

“Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which secreted Ig bound onto Fc receptors (FcRs) present on certain cytotoxic cells (*e.g.*, NK cells, neutrophils, and macrophages) enables these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII, and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 or U.S. Patent No. 6,737,056 (Presta), may be performed. Useful effector cells for such assays include PBMC and NK cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, *e.g.*, in an animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998).

“Complement-dependent cytotoxicity” or “CDC” refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system (C1q) to antibodies (of the appropriate subclass), which are bound to their cognate antigen. To assess complement activation, a CDC assay, *e.g.*, as described in Gazzano-Santoro *et al., J. Immunol. Methods* 202:163 (1996), may be performed. Polypeptide variants with altered Fc region amino acid sequences (polypeptides with a variant Fc region) and increased or decreased C1q binding capability are described, *e.g.*, in US Patent No. 6,194,551B1 and WO 1999/51642. See, also, *e.g.*, Idusogie *et al. J. Immunol.* 164: 4178-4184 (2000).

“Binding affinity” generally refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (*e.g.*, an antibody) and its binding partner (*e.g.*, an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (*e.g.*, antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d). Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigen slowly and tend to dissociate readily, whereas high-affinity antibodies

generally bind antigen faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present invention. Specific illustrative and exemplary embodiments for measuring binding affinity are described in the following.

5 In one embodiment, the “Kd” or “Kd value” according to this invention is measured by a radiolabeled antigen-binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen as described by the following assay. Solution-binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (¹²⁵I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound
10 antigen with an anti-Fab antibody-coated plate (see, *e.g.*, Chen *et al.*, *J. Mol. Biol.* 293:865-881 (1999)). To establish conditions for the assay, microtiter plates (DYNEX Technologies, Inc.) are coated overnight with 5 µg/ml of a capturing anti-Fab antibody (Cappel Labs) in 50 mM sodium carbonate (pH 9.6), and subsequently blocked with 2% (w/v) bovine serum albumin in PBS for two to five hours at room temperature (approximately 23°C). In a non-adsorbent plate (Nunc #269620), 100 pM or 26 pM [¹²⁵I]-antigen are mixed with serial
15 dilutions of a Fab of interest (*e.g.*, consistent with assessment of the anti-VEGF antibody, Fab-12, in Presta *et al.*, *Cancer Res.* 57:4593-4599 (1997)). The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (*e.g.*, about 65 hours) to ensure that equilibrium is reached. Thereafter, the mixtures are transferred to the
20 capture plate for incubation at room temperature (*e.g.*, for one hour). The solution is then removed and the plate washed eight times with 0.1% TWEEN-20TM surfactant in PBS. When the plates have dried, 150 µl/well of scintillant (MICROSCINT-20TM; Packard) is added, and the plates are counted on a TOPCOUNTTM gamma counter (Packard) for ten minutes. Concentrations of each Fab that give less than or equal to 20% of maximal binding are
25 chosen for use in competitive binding assays.

According to another embodiment, the Kd or Kd value is measured by using surface-plasmon resonance assays using a BIACORE[®]-2000 or a BIACORE[®]-3000 instrument (BIAcore, Inc., Piscataway, NJ) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIAcore
30 Inc.) are activated with N-ethyl-N'- (3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately ten response units (RU) of coupled protein.

Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% TWEEN 20™ surfactant (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIAcore® Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (Kd) is calculated as the ratio k_{off}/k_{on} . See, *e.g.*, Chen *et al.*, *J. Mol. Biol.* 293:865-881 (1999). If the on-rate exceeds $10^6 \text{ M}^{-1}\text{s}^{-1}$ by the surface-plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence-emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow-equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCO™ spectrophotometer (ThermoSpectronic) with a stirred cuvette.

An “on-rate,” “rate of association,” “association rate,” or “ k_{on} ” according to this invention can also be determined as described above using a BIACORE®-2000 or a BIACORE®-3000 system (BIAcore, Inc., Piscataway, NJ).

The term “substantially similar” or “substantially the same,” as used herein, denotes a sufficiently high degree of similarity between two numeric values (for example, one associated with an antibody of the invention and the other associated with a reference/comparator antibody), such that one of skill in the art would consider the difference between the two values to be of little or no biological and/or statistical significance within the context of the biological characteristic measured by said values (*e.g.*, Kd values). The difference between said two values is, for example, less than about 50%, less than about 40%, less than about 30%, less than about 20%, and/or less than about 10% as a function of the reference/comparator value.

The phrase “substantially reduced,” or “substantially different,” as used herein, denotes a sufficiently high degree of difference between two numeric values (generally one associated with a molecule and the other associated with a reference/comparator molecule) such that one of skill in the art would consider the difference between the two values to be of statistical significance within the context of the biological characteristic measured by said values (*e.g.*, Kd values). The difference between said two values is, for example, greater than

about 10%, greater than about 20%, greater than about 30%, greater than about 40%, and/or greater than about 50% as a function of the value for the reference/comparator molecule.

In certain embodiments, the humanized antibody useful herein further comprises amino acid alterations in the IgG Fc and exhibits increased binding affinity for human FcRn over an antibody having wild-type IgG Fc, by at least 60 fold, at least 70 fold, at least 80 fold, 5 more preferably at least 100 fold, preferably at least 125 fold, even more preferably at least 150 fold to about 170 fold.

A “disorder” or “disease” is any condition that would benefit from treatment with a substance/molecule or method of the invention. This includes chronic and acute disorders or 10 diseases including those pathological conditions which predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include malignant and benign tumors; non-leukemias and lymphoid malignancies; neuronal, glial, astrocytal, hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, immunologic and other angiogenic disorders.

15 The terms “cell proliferative disorder” and “proliferative disorder” refer to disorders that are associated with some degree of abnormal cell proliferation. In one embodiment, the cell proliferative disorder is cancer. In one embodiment, the cell proliferative disorder is angiogenesis.

“Tumor”, as used herein, refers to all neoplastic cell growth and proliferation, 20 whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. The terms “cancer”, “cancerous”, “cell proliferative disorder”, “proliferative disorder” and “tumor” are not mutually exclusive as referred to herein.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell proliferation. Examples of 25 cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, lung cancer (including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung), cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer (including gastrointestinal cancer), pancreatic cancer, glioblastoma, 30 cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer, as well as B-cell lymphoma (including low

grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's Macroglobulinemia); chronic lymphocytic leukemia (CLL); acute lymphoblastic leukemia (ALL); Hairy cell leukemia; chronic myeloblastic leukemia; and post-transplant lymphoproliferative disorder (PTLD), as well as abnormal vascular proliferation associated with phakomatoses, edema (such as that associated with brain tumors), and Meigs' syndrome.

The term "anti-neoplastic composition" or "anti-cancer composition" or "anti-cancer agent" refers to a composition useful in treating cancer comprising at least one active therapeutic agent, *e.g.*, "anti-cancer agent." Examples of therapeutic agents (anti-cancer agents) include, but are limited to, *e.g.*, chemotherapeutic agents, growth inhibitory agents, cytotoxic agents, agents used in radiation therapy, anti-angiogenesis agents, apoptotic agents, anti-tubulin agents, and other-agents to treat cancer, such as anti-HER-2 antibodies, anti-CD20 antibodies, an epidermal growth factor receptor (EGFR) antagonist (*e.g.*, a tyrosine kinase inhibitor), HER1/EGFR inhibitor (*e.g.*, erlotinib (TarcevaTM), platelet derived growth factor inhibitors (*e.g.*, GleevecTM (Imatinib Mesylate)), a COX-2 inhibitor (*e.g.*, celecoxib), interferons, cytokines, antagonists (*e.g.*, neutralizing antibodies) that bind to one or more of the following targets ErbB2, ErbB3, ErbB4, PDGFR-beta, BlyS, APRIL, BCMA VEGF, or VEGF receptor(s), TRAIL/Apo2, and other bioactive and organic chemical agents, etc. Combinations thereof are also included in the invention.

An "angiogenic factor or agent" is a growth factor which stimulates the development of blood vessels, *e.g.*, promote angiogenesis, endothelial cell growth, stability of blood vessels, and/or vasculogenesis, etc. For example, angiogenic factors, include, but are not limited to, *e.g.*, VEGF and members of the VEGF family, PIGF, PDGF family, fibroblast growth factor family (FGFs), TIE ligands (Angiopoietins), ephrins, Del-1, fibroblast growth factors: acidic (aFGF) and basic (bFGF), Follistatin, Granulocyte colony-stimulating factor (G-CSF), Hepatocyte growth factor (HGF) /scatter factor (SF), Interleukin-8 (IL-8), Leptin, Midkine, Placental growth factor, Platelet-derived endothelial cell growth factor (PD-ECGF), Platelet-derived growth factor, especially PDGF-BB or PDGFR-beta, Pleiotrophin (PTN), Progranulin, Proliferin, Transforming growth factor-alpha (TGF-alpha), Transforming growth factor-beta (TGF-beta), Tumor necrosis factor-alpha (TNF-alpha), Vascular endothelial

growth factor (VEGF)/vascular permeability factor (VPF), etc. It would also include factors that accelerate wound healing, such as growth hormone, insulin-like growth factor-I (IGF-I), VIGF, epidermal growth factor (EGF), CTGF and members of its family, and TGF-alpha and TGF-beta. See, e.g., Klagsbrun and D'Amore, *Annu. Rev. Physiol.*, 53:217-39 (1991); Streit and Detmar, *Oncogene*, 22:3172-3179 (2003); Ferrara & Alitalo, *Nature Medicine* 5(12):1359-1364 (1999); Tonini et al., *Oncogene*, 22:6549-6556 (2003) (e.g., Table 1 listing known angiogenic factors); and, Sato *Int. J. Clin. Oncol.*, 8:200-206 (2003).

The term "VEGF" as used herein refers to the 165-amino acid human vascular endothelial cell growth factor and related 121-, 189-, and 206- amino acid human vascular endothelial cell growth factors, as described by Leung *et al. Science*, 246:1306 (1989), and Houck *et al. Mol. Endocrin.*, 5:1806 (1991), together with the naturally occurring allelic and processed forms thereof. The term "VEGF" also refers to VEGFs from non-human species such as mouse, rat or primate. Sometimes the VEGF from a specific species are indicated by terms such as hVEGF for human VEGF, mVEGF for murine VEGF, and etc. The term "VEGF" is also used to refer to truncated forms of the polypeptide comprising amino acids 8 to 109 or 1 to 109 of the 165-amino acid human vascular endothelial cell growth factor. Reference to any such forms of VEGF may be identified in the present application, e.g., by "VEGF (8-109)," "VEGF (1-109)" or "VEGF₁₆₅." The amino acid positions for a "truncated" native VEGF are numbered as indicated in the native VEGF sequence. For example, amino acid position 17 (methionine) in truncated native VEGF is also position 17 (methionine) in native VEGF. The truncated native VEGF has binding affinity for the KDR and Flt-1 receptors comparable to native VEGF. According to a preferred embodiment, the VEGF is a human VEGF.

A "VEGF antagonist" refers to a molecule capable of neutralizing, blocking, inhibiting, abrogating, reducing or interfering with VEGF activities including its binding to VEGF or one or more VEGF receptors or the nucleic acid encoding them. Preferably, the VEGF antagonist binds VEGF or a VEGF receptor. VEGF antagonists include anti-VEGF antibodies and antigen-binding fragments thereof, polypeptides that bind VEGF and VEGF receptors and block ligand-receptor interaction (e.g., immunoadhesins, peptibodies), anti-VEGF receptor antibodies and VEGF receptor antagonists such as small molecule inhibitors of the VEGFR tyrosine kinases, aptamers that bind VEGF and nucleic acids that hybridize under stringent conditions to nucleic acid sequences that encode VEGF or VEGF receptor (e.g., RNAi). According to one preferred embodiment, the VEGF antagonist binds to VEGF

and inhibits VEGF-induced endothelial cell proliferation *in vitro*. According to one preferred embodiment, the VEGF antagonist binds to VEGF or a VEGF receptor with greater affinity than a non-VEGF or non-VEGF receptor. According to one preferred embodiment, the VEGF antagonist binds to VEGF or a VEGF receptor with a K_d of between 1 μ M and 1pM.

5 According to another preferred embodiment, the VEGF antagonist binds to VEGF or a VEGF receptor between 500nM and 1pM.

According to a preferred embodiment, the VEGF antagonist is selected from a polypeptide such as an antibody, a peptibody, an immunoadhesin, a small molecule or an aptamer. In a preferred embodiment, the antibody is an anti-VEGF antibody such as the
10 AVASTIN[®] antibody or an anti-VEGF receptor antibody such as an anti-VEGFR2 or an anti-VEGFR3 antibody. Other examples of VEGF antagonists include: VEGF-Trap, Mucagen, PTK787, SU11248, AG-013736, Bay 439006 (sorafenib), ZD-6474, CP632, CP-547632, AZD-2171, CDP-171, SU-14813, CHIR-258, AEE-788, SB786034, BAY579352, CDP-791, EG-3306, GW-786034, RWJ-417975/CT6758 and KRN-633.

15 An “anti-VEGF antibody” is an antibody that binds to VEGF with sufficient affinity and specificity. Preferably, the anti-VEGF antibody of the invention can be used as a therapeutic agent in targeting and interfering with diseases or conditions wherein the VEGF activity is involved. An anti-VEGF antibody will usually not bind to other VEGF homologues such as VEGF-B or VEGF-C, nor other growth factors such as PlGF, PDGF or
20 bFGF. A preferred anti-VEGF antibody is a monoclonal antibody that binds to the same epitope as the monoclonal anti-VEGF antibody A4.6.1 produced by hybridoma ATCC HB 10709. More preferably the anti-VEGF antibody is a recombinant humanized anti-VEGF monoclonal antibody generated according to Presta et al. (1997) Cancer Res. 57:4593-4599, including but not limited to the antibody known as bevacizumab (BV; Avastin[®]). According
25 to another embodiment, anti-VEGF antibodies that can be used include, but are not limited to the antibodies disclosed in WO 2005/012359. According to one embodiment, the anti-VEGF antibody comprises the variable heavy and variable light region of any one of the antibodies disclosed in Figures 24, 25, 26, 27 and 29 of WO 2005/012359 (*e.g.*, G6, G6-23, G6-31, G6-23.1, G6-23.2, B20, B20-4 and B20.4.1). In another preferred embodiment, the anti-VEGF
30 antibody known as ranibizumab is the VEGF antagonist administered for ocular disease such as diabetic neuropathy and AMD.

The anti-VEGF antibody “Bevacizumab (BV)”, also known as “rhuMAb VEGF” or “Avastin[®]”, is a recombinant humanized anti-VEGF monoclonal antibody generated

according to Presta et al. (1997) *Cancer Res.* 57:4593-4599. It comprises mutated human IgG1 framework regions and antigen-binding complementarity-determining regions from the murine anti-hVEGF monoclonal antibody A4.6.1 that blocks binding of human VEGF to its receptors. Approximately 93% of the amino acid sequence of Bevacizumab, including most
5 of the framework regions, is derived from human IgG1, and about 7% of the sequence is derived from the murine antibody A4.6.1. Bevacizumab has a molecular mass of about 149,000 daltons and is glycosylated. Other anti-VEGF antibodies include the antibodies described in United States Patent No. 6884879 and WO 2005/044853.

The anti-VEGF antibody Ranibizumab or the LUCENTIS® antibody or rhuFab V2 is
10 a humanized, affinity-matured anti-human VEGF Fab fragment. Ranibizumab is produced by standard recombinant technology methods in *Escherichia coli* expression vector and bacterial fermentation. Ranibizumab is not glycosylated and has a molecular mass of ~48,000 daltons. See WO98/45331 and US20030190317.

Dysregulation of angiogenesis can lead to abnormal angiogenesis, *i.e.*, when
15 excessive, insufficient, or otherwise inappropriate growth of new blood vessels (*e.g.*, the location, timing or onset of the angiogenesis being undesired from a medical standpoint) in a diseased state or such that it causes a diseased state, *i.e.*, an angiogenic disorder. Excessive, inappropriate or uncontrolled angiogenesis occurs when there is new blood vessel growth that contributes to the worsening of the diseased state or causes a diseased state. The new blood
20 vessels can feed the diseased tissues, destroy normal tissues, and in the case of cancer, the new vessels can allow tumor cells to escape into the circulation and lodge in other organs (tumor metastases). Disease states involving abnormal angiogenesis (*i.e.*, angiogenic disorders) include both non-neoplastic and neoplastic conditions including, *e.g.*, cancer, especially vascularized solid tumors and metastatic tumors (including colon cancer, breast
25 cancer, lung cancer (especially small-cell lung cancer), brain cancer (especially glioblastoma) or prostate cancer), undesired or aberrant hypertrophy, arthritis, rheumatoid arthritis (RA), inflammatory bowel disease or IBD (Crohn's disease and ulcerative colitis), psoriasis, psoriatic plaques, sarcoidosis, atherosclerosis, atherosclerotic plaques, diabetic and other proliferative retinopathies including retinopathy of prematurity, retrolental fibroplasia,
30 neovascular glaucoma, age-related macular degeneration, diabetic macular edema, corneal neovascularization, corneal graft neovascularization, corneal graft rejection, retinal/choroidal neovascularization, neovascularization of the anterior surface of the iris (rubeosis), ocular neovascular disease, vascular restenosis, arteriovenous malformations (AVM), meningioma,

hemangioma, angiofibroma, thyroid hyperplasias (including Grave's disease), chronic inflammation, lung inflammation, acute lung injury/ARDS, sepsis, primary pulmonary hypertension, malignant pulmonary effusions, cerebral edema (*e.g.*, associated with acute stroke/ closed head injury/ trauma), synovial inflammation, myositis ossificans, hypertrophic bone formation, osteoarthritis (OA), refractory ascites, polycystic ovarian disease, 5 endometriosis, 3rd spacing of fluid diseases (pancreatitis, compartment syndrome, burns, bowel disease), uterine fibroids, premature labor, chronic inflammation such as IBD, renal allograft rejection, inflammatory bowel disease, nephrotic syndrome, undesired or aberrant tissue mass growth (non-cancer), hemophilic joints, hypertrophic scars, inhibition of hair 10 growth, Osler-Weber syndrome, pyogenic granuloma retroental fibroplasias, scleroderma, trachoma, vascular adhesions, synovitis, dermatitis, preeclampsia, ascites, pericardial effusion (such as that associated with pericarditis), and pleural effusion.

As used herein, "treatment" refers to clinical intervention in an attempt to alter the natural course of the individual or cell being treated, and can be performed either for 15 prophylaxis or during the course of clinical pathology. Desirable effects of treatment include preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. In some embodiments, antibodies of the invention are used 20 to delay development of a disease or disorder.

An "effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

A "therapeutically effective amount" of a substance/molecule of the invention, agonist or antagonist may vary according to factors such as the disease state, age, sex, and 25 weight of the individual, and the ability of the substance/molecule, agonist or antagonist to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the substance/molecule, agonist or antagonist are outweighed by the therapeutically beneficial effects. The term "therapeutically effective amount" refers to an amount of an antibody, polypeptide or antagonist of this invention 30 effective to "treat" a disease or disorder in a mammal (aka patient). In the case of cancer, the therapeutically effective amount of the drug can reduce the number of cancer cells; reduce the tumor size or weight; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop)

tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. To the extent the drug can prevent growth and/or kill existing cancer cells, it can be cytostatic and/or cytotoxic. In one embodiment, the therapeutically effective amount is a growth inhibitory amount. In another embodiment, the therapeutically effective amount is an amount that extends the survival of a patient. In another embodiment, the therapeutically effective amount is an amount that improves progression free survival of a patient.

A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically but not necessarily, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount is less than the therapeutically effective amount.

The term “cytotoxic agent” as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (*e.g.*, At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³² and radioactive isotopes of Lu), chemotherapeutic agents, *e.g.*, methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents, enzymes and fragments thereof such as nucleolytic enzymes, antibiotics, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof, and the various antitumor or anticancer agents disclosed below. Other cytotoxic agents are described below. A tumoricidal agent causes destruction of tumor cells.

A “chemotherapeutic agent” is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin);

bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin;

5 spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (*e.g.*, calicheamicin, especially

10 calicheamicin gamma1I and calicheamicin omega1I (see, *e.g.*, Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin,

15 daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex,

20 zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, encitabine, floxuridine; androgens such as calusterone, dromostanolone propionate,

25 epitiostanol, mepitiothane, testolactone; anti- adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as

30 maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes

(especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside (“Ara-C”); thiotepa; taxoids, *e.g.*, TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE™ Cremophor-free, 5 albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Illinois), and TAXOTERE® doxorubicin (Rhône-Poulenc Rorer, Antony, France); chloranbucil; gemcitabine (GEMZAR®); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); 10 oxaliplatin; leucovorin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine (XELODA®); pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of 15 cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovorin. Additional chemotherapeutic agents include the cytotoxic agents useful as antibody drug conjugates, such as maytansinoids (DM1, for example) and the auristatins MMAE and MMAF, for example.

20 “Chemotherapeutic agents” also include “anti-hormonal agents” that act to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer, and are often in the form of systemic, or whole-body treatment. They may be hormones themselves. Examples include anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), 25 EVISTA® raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® toremifene; anti-progesterones; estrogen receptor down-regulators (ERDs); agents that function to suppress or shut down the ovaries, for example, leutinizing hormone-releasing hormone (LHRH) agonists such as LUPRON® and ELIGARD® leuprolide acetate, goserelin acetate, busarelin acetate and triptorelin; other anti- 30 androgens such as flutamide, nilutamide and bicalutamide; and aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA®

letrozole, and ARIMIDEX® anastrozole. In addition, such definition of chemotherapeutic agents includes bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), DIDROCAL® etidronate, NE-58095, ZOMETA® zoledronic acid/zoledronate, FOSAMAX® alendronate, AREDIA® pamidronate, SKELID® tiludronate, or ACTONEL®
5 risedronate; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and
10 VAXID® vaccine; LURTOTECAN® topoisomerase I inhibitor; ABARELIX® rmRH; lapatinib ditosylate (an ErbB-2 and EGFR dual tyrosine kinase small-molecule inhibitor also known as GW572016); and pharmaceutically acceptable salts, acids or derivatives of any of the above.

A “growth inhibitory agent” when used herein refers to a compound or composition
15 which inhibits growth and/or proliferation of a cell (*e.g.*, a cell expressing Robo4) either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one which significantly reduces the percentage of Robo4-expressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas
20 (vincristine and vinblastine), taxanes, and topoisomerase II inhibitors such as the anthracycline antibiotic doxorubicin ((8S-cis)-10-[(3-amino-2,3,6-trideoxy- α -L-lyxo-hexapyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-5,12-naphthacenedione), epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as
25 tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in The Molecular Basis of Cancer, Mendelsohn and Israel, eds., Chapter 1, entitled “Cell cycle regulation, oncogenes, and antineoplastic drugs” by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13. The taxanes (paclitaxel and docetaxel) are anticancer drugs both derived from the yew tree. Docetaxel
30 (TAXOTERE®, Rhone-Poulenc Rorer), derived from the European yew, is a semisynthetic analogue of paclitaxel (TAXOL®, Bristol-Myers Squibb). Paclitaxel and docetaxel promote the assembly of microtubules from tubulin dimers and stabilize microtubules by preventing depolymerization, which results in the inhibition of mitosis in cells.

As used herein, the term “patient” refers to any single animal, more preferably a mammal (including such non-human animals as, for example, dogs, cats, horses, rabbits, zoo animals, cows, pigs, sheep, and non-human primates) for which treatment is desired. Most preferably, the patient herein is a human.

5 A “subject” herein is any single human subject, including a patient, eligible for treatment who is experiencing or has experienced one or more signs, symptoms, or other indicators of an angiogenic disorder. Intended to be included as a subject are any subjects involved in clinical research trials not showing any clinical sign of disease, or subjects involved in epidemiological studies, or subjects once used as controls. The subject may have
10 been previously treated with a VEGF antagonist, or not so treated. The subject may be naïve to a second medicament being used when the treatment herein is started, *i.e.*, the subject may not have been previously treated with, for example, an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent at “baseline” (*i.e.*, at a set point in time before the administration of a first dose of antagonist in the treatment
15 method herein, such as the day of screening the subject before treatment is commenced). Such “naïve” subjects are generally considered to be candidates for treatment with such second medicament.

The expression “effective amount” refers to an amount of a medicament that is effective for treating angiogenesis disorders.

20 The term “pharmaceutical formulation” refers to a sterile preparation that is in such form as to permit the biological activity of the medicament to be effective, and which contains no additional components that are unacceptably toxic to a subject to which the formulation would be administered.

A “sterile” formulation is aseptic or free from all living microorganisms and their
25 spores.

A “package insert” is used to refer to instructions customarily included in commercial packages of therapeutic products or medicaments, that contain information about the indications, usage, dosage, administration, contraindications, other therapeutic products to be combined with the packaged product, and/or warnings concerning the use of such therapeutic
30 products or medicaments, *etc.*

A “kit” is any manufacture (*e.g.*, a package or container) comprising at least one reagent, *e.g.*, a medicament for treatment of an angiogenic disorder, or a probe for specifically detecting a biomarker gene or protein of the invention. The manufacture is

preferably promoted, distributed, or sold as a unit for performing the methods of the present invention.

For purposes of non-response to medicament(s), a subject who experiences “a clinically unacceptably high level of toxicity” from previous or current treatment with one or more medicaments experiences one or more negative side-effects or adverse events associated therewith that are considered by an experienced clinician to be significant, such as, for example, serious infections, congestive heart failure, demyelination (leading to multiple sclerosis), significant hypersensitivity, neuropathological events, high degrees of autoimmunity, a cancer such as endometrial cancer, non-Hodgkin’s lymphoma, breast cancer, prostate cancer, lung cancer, ovarian cancer, or melanoma, tuberculosis (TB), *etc.*

By “reducing the risk of a negative side effect” is meant reducing the risk of a side effect resulting from treatment with the antagonist herein to a lower extent than the risk observed resulting from treatment of the same patient or another patient with a previously administered medicament. Such side effects include those set forth above regarding toxicity, and are preferably infection, cancer, heart failure, or demyelination.

By “correlate” or “correlating” is meant comparing, in any way, the performance and/or results of a first analysis or protocol with the performance and/or results of a second analysis or protocol. For example, one may use the results of a first analysis or protocol in carrying out a second protocols and/or one may use the results of a first analysis or protocol to determine whether a second analysis or protocol should be performed. With respect to various embodiments herein, one may use the results of an analytical assay to determine whether a specific therapeutic regimen using a VEGF antagonist, such as anti-VEGF antibody, should be performed.

The word “label” when used herein refers to a compound or composition that is conjugated or fused directly or indirectly to a reagent such as a nucleic acid probe or an antibody and facilitates detection of the reagent to which it is conjugated or fused. The label may itself be detectable (*e.g.*, radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable. The term is intended to encompass direct labeling of a probe or antibody by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a

fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin.

The terms “level of expression” or “expression level” are used interchangeably and generally refer to the amount of a polynucleotide or an amino acid product or protein in a biological sample. “Expression” generally refers to the process by which gene-encoded information is converted into the structures present and operating in the cell. Therefore, according to the invention “expression” of a gene may refer to transcription into a polynucleotide, translation into a protein, or even posttranslational modification of the protein. Fragments of the transcribed polynucleotide, the translated protein, or the post-translationally modified protein shall also be regarded as expressed whether they originate from a transcript generated by alternative splicing or a degraded transcript, or from a post-translational processing of the protein, *e.g.*, by proteolysis. “Expressed genes” include those that are transcribed into a polynucleotide as mRNA and then translated into a protein, and also those that are transcribed into RNA but not translated into a protein (for example, transfer and ribosomal RNAs).

As used herein, the term “covariate” refers to certain variables or information relating to a patient. The clinical endpoints are frequently considered in regression models, where the endpoints represent the dependent variable and the biomarkers represent the main or target independent variables (regressors). If additional variables from the clinical data pool are considered, they are denoted as (clinical) covariates.

The term “clinical covariate” is used herein to describe all clinical information about the patient, which is in general available at baseline. These clinical covariates comprise demographic information like sex, age, *etc.*, other anamnestic information, concomitant diseases, concomitant therapies, results of physical examinations, common laboratory parameters obtained, known properties of the angiogenic disorders, clinical disease staging, timing and result of pretreatments, disease history, as well as all similar information that may be associated with the clinical response to treatment.

As used herein, the term “raw analysis” or “unadjusted analysis” refers to regression analyses, wherein besides the considered biomarkers, no additional clinical covariates are used in the regression model, neither as independent factors nor as stratifying covariate.

As used herein, the term “adjusted by covariates” refers to regression analyses, wherein besides the considered biomarkers, additional clinical covariates are used in the regression model, either as independent factors or as stratifying covariate.

As used herein, the term “univariate” refers to regression models or graphical approaches wherein, as an independent variable, only one of the target biomarkers is part of the model. These univariate models can be considered with and without additional clinical covariates.

5 As used herein, the term “multivariate” refers to regression models or graphical approaches wherein, as independent variables, more than one of the target biomarkers is part of the model. These multivariate models can be considered with and without additional clinical covariates.

10 III. Methods to Identify Patients Responsive to VEGF Antagonists

The present invention provides methods for identifying and/or monitoring patients likely to be responsive to VEGF antagonist (*e.g.*, anti-VEGF antibody) therapy. The methods are useful, *inter alia*, for increasing the likelihood that administration of a VEGF antagonist (*e.g.*, an anti-VEGF antibody) to a patient will be efficacious. The methods comprise
15 detecting expression of one or more genetic biomarkers in a biological sample from a patient, wherein the expression of one or more such biomarkers is indicative of whether the patient will be sensitive or responsive to VEGF antagonists, such as anti-VEGF antibodies. More particularly, the expression of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23 genes set forth in Table 1 or 2 in a sample from a patient is useful for
20 monitoring whether the patient will be responsive or sensitive to a VEGF antagonist, such as an anti-VEGF antibody. In some embodiments, expression of at least one gene selected from the following group: Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM, is useful for monitoring whether the patient will be responsive
25 or sensitive to a VEGF antagonist, such as an anti-VEGF antibody. The methods further can, optionally, include selection of a VEGF antagonist (*e.g.*, an anti-VEGF antibody, such as bevacizumab) for administration to the patient and further include, optionally, administration of a VEGF antagonist (*e.g.*, an anti-VEGF antibody, such as bevacizumab) to the patient.

The disclosed methods and assays provide for convenient, efficient, and potentially
30 cost-effective means to obtain data and information useful in assessing appropriate or effective therapies for treating patients. For example, a patient could provide a tissue sample (*e.g.*, a tumor biopsy or a blood sample) before treatment with a VEGF antagonist and the

sample could be examined by way of various *in vitro* assays to determine whether the patient's cells would be sensitive to a VEGF antagonist, such as an anti-VEGF antibody.

The invention provides methods for monitoring the sensitivity or responsiveness of a patient to a VEGF antagonist, such as an anti-VEGF antibody. The methods may be
5 conducted in a variety of assay formats, including assays detecting genetic or protein expression (such as PCR and enzyme immunoassays) and biochemical assays detecting appropriate activity. Determination of expression or the presence of such biomarkers in the samples is predictive that the patient providing the sample will be sensitive to the biological effects of a VEGF antagonist, such as an anti-VEGF antibody. Applicants' invention herein is
10 that a change (*i.e.*, an increase or decrease) in the expression at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 in a sample from a patient correlates with the observed treatment efficacy of such a patient to a VEGF antagonist, such as an anti-VEGF antibody. Example 1 shows that increased levels of the genes in Table 2 correlate with such treatment efficacy, and thus in various embodiments
15 detection of such levels in the methods described herein are included in the invention. In other embodiments, the invention includes a test panel for analysis of expression of the genes of Table 1 or Table 2, e.g., a test panel including probes specific for these genes or a subset thereof (e.g., at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2). Such a test panel may comprise, for example,
20 probes on a microchip array for use in this analysis.

According to the methods of the invention, the likelihood that a particular individual (*e.g.*, a patient) is likely to respond to treatment with a VEGF antagonist can be determined by detecting the expression level of at least one gene set forth in Table 1 or 2 and comparing the expression level of the gene to a reference expression level. For example, the reference
25 expression level may be the median expression level of the at least one gene in a group/population of patients being tested for responsiveness to a VEGF antagonist. In some embodiments, the reference expression level is the expression level of the at least one gene in a sample previously obtained from the individual at a prior time. In other embodiments, the individuals are patients who received prior treatment with a VEGF antagonist in a primary
30 tumor setting. In some embodiments, the individuals are patients who are experiencing metastasis. Individuals who have an expression level that is greater than or less than the reference expression level of at least one gene set forth in Table 1 or 2 are identified as subjects/patients likely to respond to treatment with a VEGF antagonist. Subjects/patients

who exhibit gene expression levels at the most extreme 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, or 5% relative to (i.e., higher or lower than) the median are identified as patients likely to respond to treatment with a VEGF antagonist. The gene expression level can be determined using at least one of the genes set forth in Table 1 or 2 or any linear combination of genes set forth in Table 1 or 2 (e.g., mean, weighted mean, or median) using methods known in the art and described in, e.g., Sokal R.R. and Rohlf, F.J. (1995) "Biometry: the principles and practice of statistics in biological research," W.H. Freeman and Co. New York, NY. As noted above, the methods further can, optionally, include selection of a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab) for administration to the patient and further include, optionally, administration of a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab) to the patient.

In any of the methods described above, a VDV signature score (VDV_i) can be calculated that provides quantitative information regarding the extent to which the expression of a particular set of genes is collectively overexpressed or underexpressed relative to a centered mean. For example, a VDV_i can be calculated for each sample i for which all of the VDV genes (see, e.g., Table 1 or 2) were analyzed, which represents a weighted average of z-scores across the analyzed VDV genes and is given by the algorithm:

$$VDV_i = \frac{1}{\sqrt{n}} \sum_{g=1}^n Z_{g,i}$$

in which $Z_{g=1,i}, Z_{g=2,i}, \dots, Z_{g=n,i}$ are standardized z-scores of expression values for each gene or biomarker g , from $g=1$ to $g=n$, of the sample i , and for which a VDV_i below a first defined threshold indicates a decrease relative to the reference level (e.g., collective underexpression), and a VDV_i above a second defined threshold indicates an increase relative to the reference level (e.g., collective overexpression). The expression values for each gene or biomarker g , from $g=1$ to $g=n$, may be, for example, qRT-PCR values for each gene g or biomarker, from $g=1$ to $g=n$. The first defined threshold may be from -4 to -0.5 (e.g., -4, -3.5, -3, -2.5, -2, -1.5, -1, or -0.5) and the second defined threshold may be from 0.5 to 4 (e.g., 0.5, 1, 1.5, 2, 2.5, 3, 3.5, or 4). In some instances, the first defined threshold may be from -4 to -1 (e.g., -4, -3.5, -3, -2.5, -2, -1.5, or -1) and the second defined threshold may be from 1 to 4 (e.g., 1, 1.5, 2, 2.5, 3, 3.5, or 4). In other instances, the first defined threshold may be from -4 to -1.5 (e.g., -4, -3.5, -3, -2.5, -2, or -1.5) and the second defined threshold may be from 1.5 to 4 (e.g., 1.5, 2, 2.5, 3, 3.5, or 4). Alternatively, the first defined threshold may be from -4 to -2 (e.g., -4, -

3.5, -3, -2.5, or -2) and the second defined threshold may be from 2 to 4 (e.g., 2, 2.5, 3, 3.5, or 4).

In one aspect, this invention provides a method of monitoring whether a patient with an angiogenic disorder will respond to treatment with a VEGF antagonist, such as an anti-VEGF antibody, comprising assessing, as a biomarker, expression of at least one gene set forth in Table 1 or 2 (e.g., at least one of Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM) in a sample from the patient; obtained before any VEGF antagonist has been administered to the patient. A change (*i.e.*, increase or decrease) in the expression of the at least one gene set forth in Table 1 or 2 relative to a reference level (see above) indicates that the patient will respond to treatment with a VEGF antagonist, such as an anti-VEGF antibody.

In another embodiment, the present invention provides a method of monitoring the sensitivity or responsiveness of a patient to a VEGF antagonist, such as an anti-VEGF antibody. This method comprises assessing gene expression of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 from a patient sample and predicting the sensitivity or responsiveness of the patient to the VEGF antagonist, such as an anti-VEGF antibody, wherein a change (*i.e.*, increase or decrease) in the expression of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 correlates with sensitivity or responsiveness of the patient to effective treatment with the VEGF antagonist. According to this method, a biological sample is obtained from the patient before administration of any VEGF antagonist and subjected to an assay to evaluate whether the expression products of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 are present in the sample. If expression of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or more gene(s) set forth in Table 1 or 2 is changed (*i.e.*, increased or decreased) relative to a reference level (e.g., see above), the patient is determined to be sensitive or responsive to treatment with a VEGF antagonist, such as an anti-VEGF antibody. As noted above, the methods further can, optionally, include selection of a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab) for administration to the patient and further include, optionally, administration of a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab) to the patient.

One of skill in the medical arts, particularly pertaining to the application of diagnostic tests and treatment with therapeutics, will recognize that biological systems are somewhat variable and not always entirely predictable, and thus many good diagnostic tests or therapeutics are occasionally ineffective. Thus, it is ultimately up to the judgment of the attending physician to determine the most appropriate course of treatment for an individual patient, based upon test results, patient condition and history, and his or her own experience. There may even be occasions, for example, when a physician will choose to treat a patient with a VEGF antagonist, such as an anti-VEGF antibody, even when a patient is not predicted to be particularly sensitive to VEGF antagonists, based on data from diagnostic tests or from other criteria, particularly if all or most of the other obvious treatment options have failed, or if some synergy is anticipated when given with another treatment.

In further expressed embodiments, the present invention provides a method of predicting the sensitivity of a patient to treatment with a VEGF antagonist, such as an anti-VEGF antibody, or predicting whether a patient will respond effectively to treatment with a VEGF antagonist, comprising assessing the level of one or more of the genetic biomarkers identified herein expressed in the sample; and predicting the sensitivity of the patient to inhibition by a VEGF antagonist, wherein expression levels of one or more of these genetic biomarkers correlates with high sensitivity of the patient to effective response to treatment with a VEGF antagonist.

The present invention further provides a method of identifying a biomarker whose expression level is predictive of the sensitivity or responsiveness of a particular patient to a VEGF antagonist, such as an anti-VEGF antibody, comprising: (a) measuring the expression level of a candidate biomarker in a panel of cells that displays a range of sensitivities to a VEGF antagonist, and (b) identifying a correlation between the expression level of, seropositivity for, or presence of said candidate biomarker in the cells and the sensitivity or responsiveness of the patient to the VEGF antagonist, wherein the correlation indicates that the expression level, seropositivity, or presence of said biomarker is predictive of the responsiveness of the patient to treatment by a VEGF antagonist. In one embodiment of this method the panel of cells is a panel of samples prepared from samples derived from patients or experimental animal models. In an additional embodiment the panel of cells is a panel of cell lines in mouse xenografts, wherein responsiveness can, for example, be determined by monitoring a molecular marker of responsiveness, *e.g.*, at least one of Alk1, CD34, CD105,

CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM.

The present invention also provides a method of identifying a biomarker that is useful for monitoring sensitivity or responsiveness to a VEGF antagonist, such as an anti-VEGF
5 antibody, the method comprising: (a) measuring the level of a candidate biomarker in samples from patients with angiogenic disorders obtained before any dose of a VEGF antagonist is administered to the patients, wherein an change (*i.e.*, an increase or decrease) in the expression of the candidate biomarker relative to a control indicates that the biomarker is diagnostic for more effective treatment of the angiogenic disorder with a VEGF antagonist.

10 In some embodiments, the biomarker is genetic and its expression is analyzed.

The sample may be taken from a patient who is suspected of having, or is diagnosed as having an angiogenic disorder, and hence is likely in need of treatment, or from a normal individual who is not suspected of having any disorder. For assessment of marker expression, patient samples, such as those containing cells, or proteins or nucleic acids
15 produced by these cells, may be used in the methods of the present invention. In the methods of this invention, the level of a biomarker can be determined by assessing the amount (*e.g.*, the absolute amount or concentration) of the markers in a sample, preferably a tissue sample (*e.g.*, a tumor tissue sample, such as a biopsy). In addition, the level of a biomarker can be assessed in bodily fluids or excretions containing detectable levels of biomarkers. Bodily
20 fluids or secretions useful as samples in the present invention include, *e.g.*, blood, urine, saliva, stool, pleural fluid, lymphatic fluid, sputum, ascites, prostatic fluid, cerebrospinal fluid (CSF), or any other bodily secretion or derivative thereof. The word blood is meant to include whole blood, plasma, serum, or any derivative of blood. Assessment of a biomarker in such bodily fluids or excretions can sometimes be preferred in circumstances where an
25 invasive sampling method is inappropriate or inconvenient. However, in the case of samples that are bodily fluids, the sample to be tested herein is preferably blood, synovial tissue, or synovial fluid, most preferably blood.

The sample may be frozen, fresh, fixed (*e.g.*, formalin fixed), centrifuged, and/or embedded (*e.g.*, paraffin embedded), *etc.* The cell sample can, of course, be subjected to a
30 variety of well-known post-collection preparative and storage techniques (*e.g.*, nucleic acid and/or protein extraction, fixation, storage, freezing, ultrafiltration, concentration, evaporation, centrifugation, *etc.*) prior to assessing the amount of the marker in the sample.

Likewise, biopsies may also be subjected to post-collection preparative and storage techniques, *e.g.*, fixation.

As noted above, all of the methods further can, optionally, include selection of a VEGF antagonist (*e.g.*, an anti-VEGF antibody, such as bevacizumab) for administration to the patient and further include, optionally, administration of a VEGF antagonist (*e.g.*, an anti-VEGF antibody, such as bevacizumab) to the patient.

A. Detection of Gene Expression

The genetic biomarkers described herein can be detected using any method known in the art. For example, tissue or cell samples from mammals can be conveniently assayed for, *e.g.*, mRNAs or DNAs from a genetic biomarker of interest using Northern, dot-blot, or polymerase chain reaction (PCR) analysis, array hybridization, RNase protection assay, or using DNA SNP chip microarrays, which are commercially available, including DNA microarray snapshots. For example, real-time PCR (RT-PCR) assays such as quantitative PCR assays are well known in the art. In an illustrative embodiment of the invention, a method for detecting mRNA from a genetic biomarker of interest in a biological sample comprises producing cDNA from the sample by reverse transcription using at least one primer; amplifying the cDNA so produced; and detecting the presence of the amplified cDNA. In addition, such methods can include one or more steps that allow one to determine the levels of mRNA in a biological sample (*e.g.*, by simultaneously examining the levels a comparative control mRNA sequence of a “housekeeping” gene such as an actin family member). Optionally, the sequence of the amplified cDNA can be determined.

1. Detection of Nucleic Acids

In one specific embodiment, expression of the genes set forth in Table 1 or 2 can be performed by RT-PCR technology. Probes used for PCR may be labeled with a detectable marker, such as, for example, a radioisotope, fluorescent compound, bioluminescent compound, a chemiluminescent compound, metal chelator, or enzyme. Such probes and primers can be used to detect the presence of expressed genes set forth in Table 1 or 2 in a sample. As will be understood by the skilled artisan, a great many different primers and probes may be prepared based on the sequences provided in herein and used effectively to amplify, clone and/or determine the presence and/or levels of expressed genes set forth in Table 1 or 2.

Other methods include protocols that examine or detect mRNAs from at least one of

the genes set forth in Table 1 or 2 (e.g., Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM mRNAs), in a tissue or cell sample by microarray technologies. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. For example, a selection of genes that have potential to be expressed in certain disease states may be arrayed on a solid support. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. Differential gene expression analysis of disease tissue can provide valuable information. Microarray technology utilizes nucleic acid hybridization techniques and computing technology to evaluate the mRNA expression profile of thousands of genes within a single experiment (see, e.g., WO 2001/75166). See, for example, U.S. Patent 5,700,637, U.S. Patent 5,445,934, and U.S. Patent 5,807,522, Lockart, *Nature Biotechnology*, 14:1675-1680 (1996); and Cheung *et al.*, *Nature Genetics* 21(Suppl):15-19 (1999) for a discussion of array fabrication.

In addition, the DNA profiling and detection method utilizing microarrays described in EP 1753878 may be employed. This method rapidly identifies and distinguishes between different DNA sequences utilizing short tandem repeat (STR) analysis and DNA microarrays. In an embodiment, a labeled STR target sequence is hybridized to a DNA microarray carrying complementary probes. These probes vary in length to cover the range of possible STRs. The labeled single-stranded regions of the DNA hybrids are selectively removed from the microarray surface utilizing a post-hybridization enzymatic digestion. The number of repeats in the unknown target is deduced based on the pattern of target DNA that remains hybridized to the microarray.

One example of a microarray processor is the Affymetrix GENECHIP® system, which is commercially available and comprises arrays fabricated by direct synthesis of oligonucleotides on a glass surface. Other systems may be used as known to one skilled in the art.

Other methods for determining the level of the biomarker besides RT-PCR or another PCR-based method include proteomics techniques, as well as individualized genetic profiles that are necessary to treat angiogenic disorders based on patient response at a molecular level.

The specialized microarrays herein, *e.g.*, oligonucleotide microarrays or cDNA microarrays, may comprise one or more biomarkers having expression profiles that correlate with either sensitivity or resistance to one or more anti-VEGF antibodies. Other methods that can be used to detect nucleic acids, for use in the invention, involve high throughput RNA sequence expression analysis, including RNA-based genomic analysis, such as, for example, RNASeq.

Many references are available to provide guidance in applying the above techniques (Kohler *et al.*, *Hybridoma Techniques* (Cold Spring Harbor Laboratory, New York, 1980); Tijssen, *Practice and Theory of Enzyme Inimunoassays* (Elsevier, Amsterdam, 1985); Campbell, *Monoclonal Antibody Technology* (Elsevier, Amsterdam, 1984); Hurrell, *Monoclonal Hybridoma Antibodies: Techniques and Applications* (CRC Press, Boca Raton, FL, 1982); and Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-1 58 (CRC Press, Inc., 1987)). Northern blot analysis is a conventional technique well known in the art and is described, for example, in *Molecular Cloning, a Laboratory Manual*, second edition, 1989, Sambrook, Fritch, Maniatis, Cold Spring Harbor Press, 10 Skyline Drive, Plainview, NY 11803-2500. Typical protocols for evaluating the status of genes and gene products are found, for example in Ausubel *et al.* eds., 1995, *Current Protocols In Molecular Biology*, Units 2 (Northern Blotting), 4 (Southern Blotting), 15 (Immunoblotting) and 18 (PCR Analysis).

2. Detection of Proteins

As to detection of protein biomarkers such as at least one of Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM, for example, various protein assays are available including, for example, antibody-based methods as well as mass spectroscopy and other similar means known in the art. In the case of antibody-based methods, for example, the sample may be contacted with an antibody specific for said biomarker under conditions sufficient for an antibody-biomarker complex to form, and then detecting said complex. Detection of the presence of the protein biomarker may be accomplished in a number of ways, such as by Western blotting (with or without immunoprecipitation), 2-dimensional SDS-PAGE, immunoprecipitation, fluorescence activated cell sorting (FACS), flow cytometry, and ELISA procedures for assaying a wide variety of tissues and samples, including plasma or serum. A wide range of immunoassay techniques using such an assay format are available, see, *e.g.*, U.S. Patent Nos. 4,016,043,

4,424,279, and 4,018,653. These include both single-site and two-site or “sandwich” assays of the non-competitive types, as well as in the traditional competitive binding assays. These assays also include direct binding of a labeled antibody to a target biomarker.

Sandwich assays are among the most useful and commonly used assays. A number of
5 variations of the sandwich assay technique exist, and all are intended to be encompassed by the present invention. Briefly, in a typical forward assay, an unlabelled antibody is immobilized on a solid substrate, and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen complex, a second antibody specific to the antigen, labeled
10 with a reporter molecule capable of producing a detectable signal is then added and incubated, allowing time sufficient for the formation of another complex of antibody-antigen-labeled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, by simple observation of the visible signal, or may be quantitated by
15 comparing with a control sample containing known amounts of biomarker.

Variations on the forward assay include a simultaneous assay, in which both sample and labeled antibody are added simultaneously to the bound antibody. These techniques are well known to those skilled in the art, including any minor variations as will be readily apparent. In a typical forward sandwich assay, a first antibody having specificity for the
20 biomarker is either covalently or passively bound to a solid surface. The solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride, or polypropylene. The solid supports may be in the form of tubes, beads, discs of microplates, or any other surface suitable for conducting an immunoassay. The binding processes are well-known in the art and generally
25 consist of cross-linking covalently binding or physically adsorbing, the polymer-antibody complex is washed in preparation for the test sample. An aliquot of the sample to be tested is then added to the solid phase complex and incubated for a period of time sufficient (*e.g.*, 2-40 minutes or overnight if more convenient) and under suitable conditions (*e.g.*, from room temperature to 40°C such as between 25°C and 32°C inclusive) to allow binding of any
30 subunit present in the antibody. Following the incubation period, the antibody subunit solid phase is washed and dried and incubated with a second antibody specific for a portion of the biomarker. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody to the molecular marker.

An alternative method involves immobilizing the target biomarkers in the sample and then exposing the immobilized target to specific antibody which may or may not be labeled with a reporter molecule. Depending on the amount of target and the strength of the reporter molecule signal, a bound target may be detectable by direct labeling with the antibody.

5 Alternatively, a second labeled antibody, specific to the first antibody is exposed to the target-first antibody complex to form a target-first antibody-second antibody tertiary complex. The complex is detected by the signal emitted by the reporter molecule. By “reporter molecule”, as used in the present specification, is meant a molecule which, by its chemical nature, provides an analytically identifiable signal which allows the detection of
10 antigen-bound antibody. The most commonly used reporter molecules in this type of assay are either enzymes, fluorophores or radionuclide containing molecules (*i.e.*, radioisotopes) and chemiluminescent molecules.

In the case of an enzyme immunoassay, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. As will be readily recognized,
15 however, a wide variety of different conjugation techniques exist, which are readily available to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase, and alkaline phosphatase, amongst others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable color change. Examples of suitable enzymes
20 include alkaline phosphatase and peroxidase. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. In all cases, the enzyme-labeled antibody is added to the first antibody-molecular marker complex, allowed to bind, and then the excess reagent is washed away. A solution containing the appropriate substrate is then added to the complex of antibody-antigen-
25 antibody. The substrate will react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of biomarker which was present in the sample. Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light
30 of a particular wavelength, the fluorochrome-labeled antibody adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic color visually detectable with a light microscope. As in the EIA, the fluorescent labeled antibody is allowed to bind to the first antibody-molecular marker

complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to the light of the appropriate wavelength, the fluorescence observed indicates the presence of the molecular marker of interest. Immunofluorescence and EIA techniques are both very well established in the art. However, other reporter molecules, such as
5 radioisotope, chemiluminescent or bioluminescent molecules, may also be employed.

B. Kits

For use in detection of the biomarkers, kits or articles of manufacture are also provided by the invention. Such kits can be used to determine if a subject with an angiogenic
10 disorder will be effectively responsive to a VEGF antagonist. These kits may comprise a carrier means being compartmentalized to receive in close confinement one or more container means such as vials, tubes, and the like, each of the container means comprising one of the separate elements to be used in the method. For example, one of the container means may comprise a probe that is or can be detectably labeled. Such probe may be an antibody or
15 polynucleotide specific for a protein or message, respectively. Where the kit utilizes nucleic acid hybridization to detect the target nucleic acid, the kit may also have containers containing nucleotide(s) for amplification of the target nucleic acid sequence and/or a container comprising a reporter-means, such as a biotin-binding protein, *e.g.*, avidin or streptavidin, bound to a reporter molecule, such as an enzymatic, florescent, or radioisotope
20 label.

Such kit will typically comprise the container described above and one or more other containers comprising materials desirable from a commercial and user standpoint, including buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. A label may be present on the container to indicate that the composition is used for a specific
25 application, and may also indicate directions for either *in vivo* or *in vitro* use, such as those described above.

The kits of the invention have a number of embodiments. A typical embodiment is a kit comprising a container, a label on said container, and a composition contained within said container, wherein the composition includes a primary antibody that binds to a protein or
30 autoantibody biomarker, and the label on said container indicates that the composition can be used to evaluate the presence of such proteins or antibodies in a sample, and wherein the kit includes instructions for using the antibody for evaluating the presence of biomarker proteins in a particular sample type. The kit can further comprise a set of instructions and materials

for preparing a sample and applying antibody to the sample. The kit may include both a primary and secondary antibody, wherein the secondary antibody is conjugated to a label, *e.g.*, an enzymatic label.

Another embodiment is a kit comprising a container, a label on said container, and a composition contained within said container, wherein the composition includes one or more polynucleotides that hybridize to a complement of a biomarker set forth in Table 1 or 2 under stringent conditions, and the label on said container indicates that the composition can be used to evaluate the presence of a biomarker set forth in Table 1 or 2 in a sample, and wherein the kit includes instructions for using the polynucleotide(s) for evaluating the presence of the biomarker RNA or DNA in a particular sample type.

Other optional components of the kit include one or more buffers (*e.g.*, block buffer, wash buffer, substrate buffer, *etc.*), other reagents such as substrate (*e.g.*, chromogen) that is chemically altered by an enzymatic label, epitope retrieval solution, control samples (positive and/or negative controls), control slide(s), *etc.* Kits can also include instructions for interpreting the results obtained using the kit.

In further specific embodiments, for antibody-based kits, the kit can comprise, for example: (1) a first antibody (*e.g.*, attached to a solid support) that binds to a biomarker protein; and, optionally, (2) a second, different antibody that binds to either the protein or the first antibody and is conjugated to a detectable label.

For oligonucleotide-based kits, the kit can comprise, for example: (1) an oligonucleotide, *e.g.*, a detectably labeled oligonucleotide, which hybridizes to a nucleic acid sequence encoding a biomarker protein or (2) a pair of primers useful for amplifying a biomarker nucleic acid molecule. The kit can also comprise, *e.g.*, a buffering agent, a preservative, or a protein stabilizing agent. The kit can further comprise components necessary for detecting the detectable label (*e.g.*, an enzyme or a substrate). The kit can also contain a control sample or a series of control samples that can be assayed and compared to the test sample. Each component of the kit can be enclosed within an individual container and all of the various containers can be within a single package, along with instructions for interpreting the results of the assays performed using the kit.

C. Statistics

As used herein, the general form of a prediction rule consists in the specification of a function of one or multiple biomarkers potentially including clinical covariates to predict

response or non-response, or more generally, predict benefit or lack of benefit in terms of suitably defined clinical endpoints.

The simplest form of a prediction rule consists of a univariate model without covariates, wherein the prediction is determined by means of a cutoff or threshold. This can be phrased in terms of the Heaviside function for a specific cutoff c and a biomarker measurement x , where the binary prediction A or B is to be made, then if $H(x-c)=0$, then predict A. If $H(x-c)=1$, then predict B.

This is the simplest way of using univariate biomarker measurements in prediction rules. If such a simple rule is sufficient, it allows for a simple identification of the direction of the effect, *i.e.*, whether high or low expression levels are beneficial for the patient.

The situation can be more complicated if clinical covariates need to be considered and/or if multiple biomarkers are used in multivariate prediction rules. The two hypothetical examples below illustrate the issues involved:

Covariate Adjustment (Hypothetical Example):

For a biomarker X it is found in a clinical trial population that high expression levels are associated with a worse clinical response (univariate analysis). A closer analysis shows that there are two types of clinical response in the population, a first group which possesses a worse response than the second group and at the same time the biomarker expression for the first group is generally higher following administration of at least one dose of a VEGF antagonist. An adjusted covariate analysis reveals that for each of the groups the relation of clinical benefit and clinical response is reversed, *i.e.*, within the groups, lower expression levels are associated with better clinical response. The overall opposite effect was masked by the covariate type--and the covariate adjusted analysis as part of the prediction rule reversed the direction.

Multivariate Prediction (Hypothetical Example):

For a biomarker X it is found in a clinical trial population that high expression levels are slightly associated with a worse clinical response (univariate analysis). For a second biomarker Y a similar observation was made by univariate analysis. The combination of X and Y revealed that a good clinical response is seen if both biomarkers are low. This makes the rule to predict benefit if both biomarkers are below some cutoffs (AND--connection of a Heaviside prediction function). For the combination rule, a simple rule no longer applies in a univariate sense; for example, having low expression levels in X will not automatically predict a better clinical response.

These simple examples show that prediction rules with and without covariates cannot be judged on the univariate level of each biomarker. The combination of multiple biomarkers plus a potential adjustment by covariates does not allow assigning simple relationships to single biomarkers. Since the marker genes, in particular in serum, may be used in multiple-
5 marker prediction models potentially including other clinical covariates, the direction of a beneficial effect of a single marker gene within such models cannot be determined in a simple way, and may contradict the direction found in univariate analyses, *i.e.*, the situation as described for the single marker gene.

A clinician may use any of several methods known in the art to measure the
10 effectiveness of a particular dosage scheme of a VEGF antagonist. For example, *in vivo* imaging (*e.g.*, MRI) can be used to determine the tumor size and to identify any metastases to determine relative effective responsiveness to the therapy. Dosage regimens may be adjusted to provide the optimum desired response (*e.g.*, a therapeutic response). For example, a dose may be administered, several divided doses may be administered over time or the dose may
15 be proportionally reduced or increased as indicated by exigencies of the therapeutic situation.

A physician having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required, depending on such factors as the particular antagonist type. For example, the physician could start with doses of such antagonist, such as an anti-VEGF antibody, employed in the pharmaceutical composition at
20 levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. The effectiveness of a given dose or treatment regimen of the antagonist can be determined, for example, by assessing signs and symptoms in the patient using standard measures of efficacy.

In yet another embodiment, the subject is treated with the same antagonist, such as
25 anti-VEGF antibody at least twice. Thus, the initial and second antagonist exposures are preferably with the same antagonist, and more preferably all antagonist exposures are with the same antagonist, *i.e.*, treatment for the first two exposures, and preferably all exposures, is with one type of VEGF antagonist, for example, an antagonist that binds to VEGF, such as an anti-VEGF antibody, *e.g.*, all with bevacizumab.

30 In all the inventive methods set forth herein, the antagonist (such as an antibody that binds to VEGF) may be unconjugated, such as a naked antibody, or may be conjugated with another molecule for further effectiveness, such as, for example, to improve half-life.

The preferred antagonist antibody herein is a chimeric, humanized, or human antibody, more preferably, an anti-VEGF antibody, and most preferably bevacizumab.

In another embodiment, the VEGF antagonist (*e.g.*, an anti-VEGF antibody) is the only medicament administered to the subject.

5 In one embodiment, the antagonist is an anti-VEGF antibody that is administered at a dose of about 100 or 400 mg every 1, 2, 3, or 4 weeks or is administered a dose of about 1, 3, 5, 10, 15, or 20 mg/kg every 1, 2, 3, or 4 weeks. The dose may be administered as a single dose or as multiple doses (*e.g.*, 2 or 3 doses), such as infusions.

10 In yet another aspect, the invention provides, after the diagnosis step, a method of determining whether to continue administering a VEGF antagonist (*e.g.*, an anti-VEGF antibody) to a subject with an angiogenic disorder comprising measuring reduction in tumor size, using imaging techniques, such as radiography and/or MRI, after administration of the antagonist a first time, measuring reduction in tumor size in the subject, using imaging techniques such as radiography and/or MRI after administration of the antagonist a second
15 time, comparing imaging findings in the subject at the first time and at the second time, and if the score is less at the second time than at the first time, continuing administration of the antagonist.

In a still further embodiment, a step is included in the treatment method to test the subject's response to treatment after the administration step to determine that the level of
20 response is effective to treat the angiogenic disorder. For example, a step is included to test the imaging (radiographic and/or MRI) score after administration and compare it to baseline imaging results obtained before administration to determine if treatment is effective by measuring if, and by how much, it has been changed. This test may be repeated at various scheduled or unscheduled time intervals after the administration to determine maintenance of
25 any partial or complete remission. Alternatively, the methods herein comprise a step of testing the subject, before administration, to see if one or more biomarkers or symptoms are present for angiogenic disorders, as set forth above.

In one embodiment of the invention, no other medicament than VEGF antagonist such as anti-VEGF antibody is administered to the subject to treat an angiogenic disorder.

30 In any of the methods herein, the VEGF antagonist may be administered in combination with an effective amount of a second medicament (where the VEGF antagonist (*e.g.*, an anti-VEGF antibody) is a first medicament). Suitable second medicaments include,

for example, an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, or combinations thereof.

All these second medicaments may be used in combination with each other or by themselves with the first medicament, so that the expression "second medicament" as used
5 herein does not mean it is the only medicament in addition to the first medicament. Thus, the second medicament need not be a single medicament, but may constitute or comprise more than one such drug.

These second medicaments as set forth herein are generally used in the same dosages and with administration routes as used hereinbefore or about from 1 to 99% of the heretofore-
10 employed dosages. If such second medicaments are used at all, preferably, they are used in lower amounts than if the first medicament were not present, especially in subsequent dosings beyond the initial dosing with the first medicament, so as to eliminate or reduce side effects caused thereby.

For the re-treatment methods described herein, where a second medicament is
15 administered in an effective amount with an antagonist exposure, it may be administered with any exposure, for example, only with one exposure, or with more than one exposure. In one embodiment, the second medicament is administered with the initial exposure. In another embodiment, the second medicament is administered with the initial and second exposures. In a still further embodiment, the second medicament is administered with all exposures. It is
20 preferred that after the initial exposure, such as of steroid, the amount of such second medicament is reduced or eliminated so as to reduce the exposure of the subject to an agent with side effects such as prednisone, prednisolone, methylprednisolone, and cyclophosphamide.

The combined administration of a second medicament includes co-administration
25 (concurrent administration), using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents (medicaments) simultaneously exert their biological activities.

The antagonist herein is administered by any suitable means, including parenteral,
30 topical, subcutaneous, intraperitoneal, intrapulmonary, intranasal, and/or intralesional administration. Parenteral infusions include intramuscular, intravenous (i.v.), intraarterial, intraperitoneal, or subcutaneous administration. Intrathecal administration is also contemplated. In addition, the antagonist may suitably be administered by pulse infusion,

e.g., with declining doses of the antagonist. Preferably, the dosing is given intravenously or subcutaneously, and more preferably by intravenous infusion(s).

If multiple exposures of antagonist are provided, each exposure may be provided using the same or a different administration means. In one embodiment, each exposure is by intravenous administration. In another embodiment, each exposure is given by subcutaneous administration. In yet another embodiment, the exposures are given by both intravenous and subcutaneous administration.

In one embodiment, the antagonist such as an anti-VEGF antibody is administered as a slow intravenous infusion rather than an intravenous push or bolus. For example, a steroid such as prednisolone or methylprednisolone (*e.g.*, about 80-120 mg i.v., more specifically about 100 mg i.v.) is administered about 30 minutes prior to any infusion of the anti-VEGF antibody. The anti-VEGF antibody is, for example, infused through a dedicated line.

For the initial dose of a multi-dose exposure to anti-VEGF antibody, or for the single dose if the exposure involves only one dose, such infusion is preferably commenced at a rate of about 50 mg/hour. This may be escalated, *e.g.*, at a rate of about 50 mg/hour increments every about 30 minutes to a maximum of about 400 mg/hour. However, if the subject is experiencing an infusion-related reaction, the infusion rate is preferably reduced, *e.g.*, to half the current rate, *e.g.*, from 100 mg/hour to 50 mg/hour. Preferably, the infusion of such dose of anti-VEGF antibody (*e.g.*, an about 1000-mg total dose) is completed at about 255 minutes (4 hours 15 min.). Optionally, the subjects receive a prophylactic treatment of acetaminophen/paracetamol (*e.g.*, about 1 g) and diphenhydramine HCl (*e.g.*, about 50 mg or equivalent dose of similar agent) by mouth about 30 to 60 minutes prior to the start of an infusion.

If more than one infusion (dose) of anti-VEGF antibody is given to achieve the total exposure, the second or subsequent anti-VEGF antibody infusions in this infusion embodiment are preferably commenced at a higher rate than the initial infusion, *e.g.*, at about 100 mg/hour. This rate may be escalated, *e.g.*, at a rate of about 100 mg/hour increments every about 30 minutes to a maximum of about 400 mg/hour. Subjects who experience an infusion-related reaction preferably have the infusion rate reduced to half that rate, *e.g.*, from 100 mg/hour to 50 mg/hour. Preferably, the infusion of such second or subsequent dose of anti-VEGF antibody (*e.g.*, an about 1000-mg total dose) is completed by about 195 minutes (3 hours 15 minutes).

In a preferred embodiment, the antagonist is an anti-VEGF antibody and is administered in a dose of about 0.4 to 4 grams, and more preferably the antibody is administered in a dose of about 0.4 to 1.3 grams at a frequency of one to four doses within a period of about one month. Still more preferably, the dose is about 500 mg to 1.2 grams, and in other embodiments is about 750 mg to 1.1 grams. In such aspects, the antagonist is preferably administered in two to three doses, and/or is administered within a period of about 2 to 3 weeks.

In one embodiment, the subject has never been previously administered any drug(s) to treat the angiogenic disorder. In another embodiment, the subject or patient has been previously administered one or more medicaments(s) to treat the angiogenic disorder. In a further embodiment, the subject or patient was not responsive to one or more of the medicaments that had been previously administered. Such drugs to which the subject may be non-responsive include, for example, anti-neoplastic agents, chemotherapeutic agents, cytotoxic agents, and/or growth inhibitory agents. More particularly, the drugs to which the subject may be non-responsive include VEGF antagonists such as anti-VEGF antibodies. In a further aspect, such antagonists include an antibody or immunoadhesin, such that re-treatment is contemplated with one or more antibodies or immunoadhesins of this invention to which the subject was formerly non-responsive.

IV. Treatment with the Antagonist

Once the patient population most responsive or sensitive to treatment with the antagonist has been identified, treatment with the antagonist herein, alone or in combination with other medicaments, results in an improvement in the angiogenic disorder. For instance, such treatment may result in a reduction in tumor size or progression free survival.

Moreover, treatment with the combination of an antagonist herein and at least one second medicament(s) preferably results in an additive, more preferably synergistic (or greater than additive) therapeutic benefit to the patient. Preferably, in this combination method the timing between at least one administration of the second medicament and at least one administration of the antagonist herein is about one month or less, more preferably, about two weeks or less.

Administration of VEGF antagonists, as described herein, is optionally included in the invention. Thus, in a further embodiment, the invention provides a method of treating cancer (e.g., colorectal cancer, breast cancer, lung cancer, or glioblastoma) in a patient by administration of a VEGF antagonist (e.g., an anti-VEGF antibody, such as bevacizumab),

wherein the patient is or has been identified as being one that will benefit from such treatment, according to the methods described herein.

It will be appreciated by one of skill in the medical arts that the exact manner of administering to said patient a therapeutically effective amount of a VEGF antagonist following a diagnosis of a patient's likely responsiveness to the antagonist will be at the discretion of the attending physician. The mode of administration, including dosage, combination with other agents, timing and frequency of administration, and the like, may be affected by the diagnosis of a patient's likely responsiveness to such antagonist, as well as the patient's condition and history. Thus, even patients diagnosed with an angiogenic disorder who are predicted to be relatively insensitive to the antagonist may still benefit from treatment therewith, particularly in combination with other agents, including agents that may alter a patient's responsiveness to the antagonist.

The composition comprising an antagonist will be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular type of angiogenic disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the angiogenic disorder, the site of delivery of the agent, possible side-effects, the type of antagonist, the method of administration, the scheduling of administration, and other factors known to medical practitioners. The effective amount of the antagonist to be administered will be governed by such considerations.

As a general proposition, the effective amount of the antagonist administered parenterally per dose will be in the range of about 20 mg to about 5000 mg, by one or more dosages. Exemplary dosage regimens for antibodies such as anti-VEGF antibodies include 100 or 400 mg every 1, 2, 3, or 4 weeks or is administered a dose of about 1, 3, 5, 10, 15, or 20 mg/kg every 1, 2, 3, or 4 weeks. The dose may be administered as a single dose or as multiple doses (*e.g.*, 2 or 3 doses), such as infusions.

As noted above, however, these suggested amounts of antagonist are subject to a great deal of therapeutic discretion. The key factor in selecting an appropriate dose and scheduling is the result obtained, as indicated above. In some embodiments, the antagonist is administered as close to the first sign, diagnosis, appearance, or occurrence of the angiogenic disorder as possible.

The antagonist is administered by any suitable means, including parenteral, topical, subcutaneous, intraperitoneal, intrapulmonary, intranasal, and/or intralesional administration.

Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Intrathecal administration is also contemplated. In addition, the antagonist may suitably be administered by pulse infusion, *e.g.*, with declining doses of the antagonist. Most preferably, the dosing is given by intravenous injections.

5 One may administer a second medicament, as noted above, with the antagonists herein. The combined administration includes co-administration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

10 Aside from administration of antagonists to the patient by traditional routes as noted above, the present invention includes administration by gene therapy. Such administration of nucleic acids encoding the antagonist is encompassed by the expression "administering an effective amount of an antagonist". See, for example, WO 1996/07321 concerning the use of gene therapy to generate intracellular antibodies.

15 There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells; *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the site where the antagonist is required. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells and the modified cells are administered to the patient either directly or, for
20 *e.g.*, encapsulated within porous membranes which are implanted into the patient (see, *e.g.*, U.S. Patent Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro* or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro*
25 include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, *etc.* A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and
30 lipid-based systems (useful lipids for lipid-mediated transfer of the gene are DOTMA, DOPE and DC-Chol, for example). In some situations it is desirable to provide the nucleic acid source with an agent specific for the target cells, such as an antibody specific for a cell-surface membrane protein on the target cell, a ligand for a receptor on the target cell, *etc.*

Where liposomes are employed, proteins that bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, *e.g.*, capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins that undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu *et al.*, *J. Biol. Chem.* 262:4429-4432 (1987); and Wagner *et al.*, *PNAS USA* 87:3410-3414 (1990). Gene-marking and gene-therapy protocols are described, for example, in Anderson *et al.*, *Science* 256:808-813 (1992) and WO 1993/25673.

A VEGF antagonist may be combined in a pharmaceutical combination formulation, or dosing regimen as combination therapy, with at least one additional compound having anti-cancer properties. The at least one additional compound of the pharmaceutical combination formulation or dosing regimen preferably has complementary activities to the VEGF antagonist composition such that they do not adversely affect each other.

The at least one additional compound may be a chemotherapeutic agent, a cytotoxic agent, a cytokine, a growth inhibitory agent, an anti-hormonal agent, and combinations thereof. Such molecules are suitably present in combination in amounts that are effective for the purpose intended. A pharmaceutical composition containing an VEGF antagonist (*e.g.*, an anti-VEGF antibody) may also comprise a therapeutically effective amount of an anti-neoplastic agent, a chemotherapeutic agent a growth inhibitory agent, a cytotoxic agent, or combinations thereof.

In one aspect, the first compound is an anti-VEGF antibody and the at least one additional compound is a therapeutic antibody other than an anti-VEGF antibody. In one embodiment, the at least one additional compound is an antibody that binds a cancer cell surface marker. In one embodiment the at least one additional compound is an anti-HER2 antibody, trastuzumab (*e.g.*, Herceptin®, Genentech, Inc., South San Francisco, CA). In one embodiment the at least one additional compound is an anti-HER2 antibody, pertuzumab (Omnitarg™, Genentech, Inc., South San Francisco, CA, see US6949245). In an embodiment, the at least one additional compound is an antibody (either a naked antibody or an ADC), and the additional antibody is a second, third, fourth, fifth, sixth antibody or more, such that a combination of such second, third, fourth, fifth, sixth, or more antibodies (either naked or as an ADC) is efficacious in treating an angiogenic disorder.

Other therapeutic regimens in accordance with this invention may include administration of a VEGF-antagonist anticancer agent and, including without limitation

radiation therapy and/or bone marrow and peripheral blood transplants, and/or a cytotoxic agent, a chemotherapeutic agent, or a growth inhibitory agent. In one of such embodiments, a chemotherapeutic agent is an agent or a combination of agents such as, for example, cyclophosphamide, hydroxydaunorubicin, adriamycin, doxorubicin, vincristine
5 (ONCOVIN™), prednisolone, CHOP, CVP, or COP, or immunotherapeutics such as anti-PSCA, anti-HER2 (*e.g.*, HERCEPTIN®, OMNITARG™). The combination therapy may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in two or more administrations. The combined
10 administration includes coadministration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

In one embodiment, treatment with an anti-VEGF antibody involves the combined administration of an anticancer agent identified herein, and one or more chemotherapeutic
15 agents or growth inhibitory agents, including coadministration of cocktails of different chemotherapeutic agents. Chemotherapeutic agents include taxanes (such as paclitaxel and docetaxel) and/or anthracycline antibiotics. Preparation and dosing schedules for such chemotherapeutic agents may be used according to manufacturer's instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such
20 chemotherapy are also described in "Chemotherapy Service", (1992) Ed., M.C. Perry, Williams & Wilkins, Baltimore, Md.

Suitable dosages for any of the above coadministered agents are those presently used and may be lowered due to the combined action (synergy) of the newly identified agent and other chemotherapeutic agents or treatments.

25 The combination therapy may provide "synergy" and prove "synergistic", *i.e.* the effect achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined, unit dosage formulation; (2) delivered by alternation or in
30 parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are administered or delivered sequentially, *e.g.* by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is

administered sequentially, *i.e.* serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together.

For the prevention or treatment of disease, the appropriate dosage of the additional therapeutic agent will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the VEGF antagonist and additional agent are administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the VEGF antagonist and additional agent, and the discretion of the attending physician. The VEGF antagonist and additional agent are suitably administered to the patient at one time or over a series of treatments. The VEGF antagonist is typically administered as set forth above. Depending on the type and severity of the disease, about 20 mg/m² to 600 mg/m² of the additional agent is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about or about 20 mg/m², 85 mg/m², 90 mg/m², 125 mg/m², 200 mg/m², 400 mg/m², 500 mg/m² or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. Thus, one or more doses of about 20 mg/m², 85 mg/m², 90 mg/m², 125 mg/m², 200 mg/m², 400 mg/m², 500 mg/m², 600 mg/m² (or any combination thereof) may be administered to the patient. Such doses may be administered intermittently, *e.g.*, every week or every two, three weeks, four, five, or six (*e.g.*, such that the patient receives from about two to about twenty, *e.g.* about six doses of the additional agent). An initial higher loading dose, followed by one or more lower doses may be administered. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

V. Pharmaceutical Formulations

Therapeutic formulations of the antagonists used in accordance with the present invention are prepared for storage by mixing the antagonist having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients, or stabilizers in the form of lyophilized formulations or aqueous solutions. For general information concerning formulations, see, *e.g.*, Gilman *et al.*, (eds.) (1990), *The Pharmacological Bases of Therapeutics*, 8th Ed., Pergamon Press; A. Gennaro (ed.), Remington's Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.; Avis *et al.*,

(eds.) (1993) *Pharmaceutical Dosage Forms: Parenteral Medications* Dekker, New York; Lieberman *et al.*, (eds.) (1990) *Pharmaceutical Dosage Forms: Tablets* Dekker, New York; and Lieberman *et al.*, (eds.) (1990), *Pharmaceutical Dosage Forms: Disperse Systems* Dekker, New York, Kenneth A. Walters (ed.) (2002) *Dermatological and Transdermal Formulations* (Drugs and the Pharmaceutical Sciences), Vol 119, Marcel Dekker.

Acceptable carriers, excipients, or stabilizers are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™, or polyethylene glycol (PEG).

Exemplary anti-VEGF antibody formulations are described in U.S. Patent Nos. 6,884,879. In certain embodiments anti-VEGF antibodies are formulated at 25 mg/mL in single use vials. In certain embodiments, 100 mg of the anti-VEGF antibodies are formulated in 240 mg α,α -trehalose dihydrate, 23.2 mg sodium phosphate (monobasic, monohydrate), 4.8 mg sodium phosphate (dibasic anhydrous), 1.6 mg polysorbate 20, and water for injection, USP. In certain embodiments, 400 mg of the anti-VEGF antibodies are formulated in 960 mg α,α -trehalose dihydrate, 92.8 mg sodium phosphate (monobasic, monohydrate), 19.2 mg sodium phosphate (dibasic anhydrous), 6.4 mg polysorbate 20, and water for injection, USP.

Lyophilized formulations adapted for subcutaneous administration are described, for example, in U.S. Patent No. 6,267,958 (Andya *et al.*). Such lyophilized formulations may be reconstituted with a suitable diluent to a high protein concentration and the reconstituted formulation may be administered subcutaneously to the mammal to be treated herein.

Crystallized forms of the antagonist are also contemplated. See, for example, US 2002/0136719A1.

The formulation herein may also contain more than one active compound (a second medicament as noted above), preferably those with complementary activities that do not adversely affect each other. The type and effective amounts of such medicaments depend, for example, on the amount and type of VEGF antagonist present in the formulation, and clinical parameters of the subjects. The preferred such second medicaments are noted above.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the antagonist, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The formulations to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Table 1.

ABCC9	CD97	DLC1	GIMAP5	LEPREL2	PCDH18	RRAS	THSD1
ACVRL1	CD105	DLL4	GIMAP6	LHFP	PDGFB	S1PR1	THY1
ADAM12	CD144	DPYS	GJA4	LIX1L	PDGFD	S1PR3	TIMP3
ADAMTS1	CD276	DUSP6	GJC1	LIMS2	PDGFRB	SCARF1	TMEM88
ADAMTS2	CARD11	EBF2	GPR4	LMO2	PHACTR2	SEMA3F	TMEM204
ADAMTS12	CDC42EPI	EFNB2	GPR124	LPAR6	PLK2	SEPT4	TM4SF1
AFAPIL2	CDH11	EGFL7	HAPLN1	LRP4	PLVAP	SERPINE1	TNFAIP2
AHR	CDH5	EHD4	HEY2	LRRC33	PLXDC2	SERPINH1	TNNT2
Alk1	CDRT4	ELTD1	HEYL	LY6E	PPAP2A	SLC9A3R2	TP53
AMOTL1	CES2	EMCN	HIGD1B	MAPK12	PPAP2B	SLC11A1	TREML4
ANGPT2	CHST15	ENG	HLX	MCAM	PPFIA4	SLC5A2	TRDN
ANXA1	CLEC4D	ERG	HOXA3	MECOM	PPM1F	SLC22A9	TRIB2
APLNR	CLEC6A	ESAM	HOXD8	MED24	PTAFR	SLC22A25	TRIM5
ARAP3	CLEC14A	ESM1	ICAM2	MDFIC	PRICKLE2	SLC43A3	TSLP
ARHGAP29	CLEC1A	ETS1	ID1	MEF2C	PROCR	SLFN5	TSP1
ARHGAP31	CNN2	F2R	IFI44	MFGE8	PRKCDBP	SNCG	TUSC5
ART3	COL15A1	F2RL3	IGFBP3	MMRN2	PRKCH	SOD3	UBXN10
ASB2	COL18A1	FAM198B	IL2RG	MYCT1	PRND	SOX7	USHBP1
ATP1B4	COL1A2	FANCI	INHBB	NAALAD2	PRR5L	SOX18	VAMP5
BGN	COL4A1	FBLN5	ITPRIPL2	NFIB	PTPRB	SPARC	VEGFR1
BHMT2	COL4A2	FBN1	JUB	NHSL2	PTPRG	SPIC	VEGFR2
BTNL9	COL4A3	FKBP10	KANK3	NID1	PTPRM	SRGN	VEGFR3
BMPR2	COL8A1	FLI1	KCNE3	NID2	QKI	SPRY4	VIM
BPIFB1	COMMD7	FLT1	KDR	NNMT	RAPGEF3	STAT1	WISP1
BST2	CRTAM	FMOD	KIAA1274	NOS3	RAPGEF5	STEAP4	WWTR1
C3orf64	CSPG4	FREM1	KIAA1462	NOTCH1	RASIP1	STON1	ZFP36L1
C8orf4	CTGF	FSTL1	KLHL4	NRARP	RASGRP3	STON1- GTF2A1L	ZNF521
C13orf15	CTTNBP2NL	FXVD5	KLHL5	NRP1	RBP7	ST8SIA4	
C1S	CYYR1	GAS6	KLHL6	NRP2	RBMS1	ST8SIA6	
CAV1	DAPK2	GAS7	KLRB1	NRXN3	RGS5	SWAP70	
CCND1	DCHS1	GIMAP1	KPNA7	P2RX7	REST	TEK	
CD34	DDAH1	GIMAP2	LAMA4	PBX2	RHOJ	TFAP2C	
CD40	DEF6	GIMAP3	LAMB1	PCDH12	RIN3	THBS1	
CD93	DKK2	GIMAP4	LATS2	PCDH17	ROBO4	THBS2	

Table 2.

Alk1	EGFL7	RGS5
CD34	ESM1	Sema3f
CD105	LAMA4	TSP1
CD144	NG2	VEGFR1
Col4a1	Nid2	VEGFR2

Col4a2	Notch1	VEGFR3
Dll4	NRP1	VIM
EFNB2	NRP2	

EXAMPLES

The following examples are provided to illustrate, but not to limit the presently claimed invention.

5 **Example 1. Materials and Methods**

Mouse strains & models

We obtained *RIP-T β Ag* mice from Exelixis, Inc. and Beige Nude XID mice from Harlan. Animals were housed and cared for according to guidelines from the Institutional Animal Care and Use Committee (IACUC) at Genentech, Inc.

10 *Treatment regimens and dosing*

All dosing regimens were carried out according to IACUC guidelines. Study animals were monitored daily and body weights were measured at least twice weekly. Procedures used for intracranial tumor establishment, including monitoring of tumor growth and response to therapy by bioluminescence imaging, have previously been described (Ozawa and James. *J Vis Exp.* 41: 1-5, 2010). The anti-vasculature endothelial growth factor (VEGF) monoclonal antibody B20-4.1.1, anti-Ragweed (control) and anti-Dll4 were prepared and purified as previously described (Fuh et al. *J Biol Chem.* 281: 6625-6631, 2006), and dosed at 15 5 or 10mg/kg twice weekly by intraperitoneal (i.p.) injection for all experiments in immunocompromised mice and once weekly (at 5mg/kg) in *RIP-T β Ag* mice. Sunitinib was dosed at 20 60mg/kg daily by oral gavage. Wound healing assays were carried out as previously described in Bais et al. (*Cell.* 141: 166-177, 2010).

Immunofluorescent staining & histological quantitation

Whole pancreata dissected from tumor-bearing *RIP-T β Ag* mice were incubated in sucrose (30%) for 5-10 min at 4°C, followed by washing with PBS (twice 15 minutes each). 25 Pancreata were then placed in cryomolds containing Optimum Cutting Temperature (OCT, Sakura Finetek) medium and were maintained at -70°C. Sections (6 μ m) of pancreata were cut from each OCT block using a cryostat instrument (Leica Microsystems) and were maintained at -70°C until used for staining. For immunofluorescent staining, frozen sections were air-dried at room temperature and fixed with cold acetone for 5-10 minutes. Then 30 sections were dried again and blocked for 30-60 minutes with a buffer containing 2.5% BSA and 5% donkey serum in PBS. Staining with primary antibodies diluted in blocking buffer

(dilutions as per manufacturer's guidelines) was carried out overnight at 4°C. Sections were then washed with PBS and exposed to secondary antibodies diluted 1:300 in blocking buffer for 30-60 minutes, and then washed again in PBS. Finally sections were mounted in DAKO (DAKO) containing 4',6-diamidino-2-phenylindole (DAPI, Molecular Probes) for nuclear visualization.

The following antibodies were used for immunofluorescent staining: primary rat anti-mouse MECA-32 antibody (Invitrogen, Inc.), rabbit anti-Ki67, and clone SP6 (Research Diagnostics, Inc.). Secondary antibodies included Alexa-594 conjugated donkey anti-rat and anti-rabbit and Alexa-488 conjugated donkey anti-rabbit and goat anti-chicken (Invitrogen, Inc.). Tumors on stained slides were identified microscopically and photographed using an Axioskop and Axiovision Software (Zeiss, Inc.), and histological quantification on images was performed using customized algorithms in Metamorph software (Molecular Dynamics). At each time point, the mean measurement was calculated from at least 3-5 images per islet/tumor x 5 tumors from each mouse x 3 mice = 45-75 images/time point/treatment from 15 independent lesions. For micro-vascular density (MVD) analyses in implanted tumor models, tumors from 6 mice per treatment group were collected and embedded in O.C.T. blocks. Tissues were cryo-sectioned to 16- μ m thickness on Leica CM3050S, and stained with CD31 antibody (BD Biosciences). Images were acquired with Zeiss AxioImager Z1 fluorescence microscope controlled by TissueFAXS software. Image files were loaded into the TissueStudio analysis package (v1.5, Definiens). Necrotic tissues and staining artifacts such as skin tissues and folds were automatically identified and excluded based on nuclei staining. Vessel density was calculated as the ratio of CD31-positive pixels to the total viable tumor area.

Microarray experiments

Total RNA was extracted from control and anti-VEGF-treated *RIP-T β Ag* tumors after 7 days of treatment as follows: mice were anesthetized using 0.25% Avertin, injected intraperitoneally according to weight. The abdominal cavity was opened to access the pancreatic duct and perfuse the pancreas via the common bile duct. The pancreas was perfused with approximately 2.5 ml of Liberase TL (Roche) diluted as per the manufacturer's instructions. The pancreas was then dissected out of the abdominal cavity and tumor further macro-dissected away from the exocrine pancreas. Tumors were suspended in fresh perfusion solution and agitated at 37°C for 5-6 minutes and then re-examined under a dissection microscope and any remaining fragments of exocrine pancreas removed. Clean

tumors were then flash-frozen in RNAlater solution (Qiagen, Inc.). For all implant experiments in immune-deficient mice, animals were euthanized at the end of the study interval and tumors dissected out and snap frozen. Total RNA was extracted and microarrays run using Agilent Whole Mouse Genome 44K arrays, Affymetrix HGU133-plus2, or Agilent
5 Whole Human Genome arrays according to manufacturer's instructions.

Cell culture

D551 (ATCC) skin fibroblast cells were cultured in M199 medium (Invitrogen), supplemented with fetal bovine serum (Sigma-Aldrich), penicillin (100 units/mL), streptomycin (100 µg/mL). Primary human umbilical vascular endothelial cells (HUVEC)
10 were purchased from (Lonza Walkersville), and maintained in EGM-2 medium (Lonza Walkersville). Condition medium: D551 cells were grown to 90% confluence, changed medium to EGM-2, after 7 days of incubation, the supernatant was collected and stored at 4°C.

HUVEC transfection and sprouting assay

15 COL4A2, NID2, and MEST gene expression silencing: cells were grown to 70% confluence, and siRNA were transfected using the DharmaFECT1 according to the manufacturer's instructions (Thermo Scientific). The final concentration of all siRNA for transfection was 12.5 nM, and mRNA downregulation for each gene was confirmed by qRT-PCR. 24 hours after transfection, cells were trypsinized and mixed with Cytodex
20 microcarrier beads (Sigma-Aldrich) in a ratio of 1×10^6 cells per 2,500 beads. Coating was performed for four hours at 37°C and mixtures shaken by hand every 20 minutes. Coated beads were then transferred to a 6-well dish, and left for 18-20 hours in EGM-2 at 37°C and 5% CO₂. The following day, coated beads were washed with EGM-2, and dissolved in a solution of fibrinogen (2 mg ml⁻¹; Sigma-Aldrich) in EGM-2. The solution with about 200
25 HUVEC coated beads was added to 0.625 U ml⁻¹ of thrombin (Sigma) in one well of a 24-well tissue culture plate. 8×10^4 skin fibroblast cells (D551) were plated on top of the clot and incubated with 2 ml D551 condition medium/EGM-2 (1: 3) containing 20ug/ml of antibodies. The medium was replaced every 2 days and assays were terminated at day 4. HUVEC sprouts were visualized by immunostaining in fibrin gels fixed in 4%
30 paraformaldehyde (PFA) for 2 hrs at RT, then blocked with block buffer (DAKO) for 4 hrs at RT, incubated with Alexa Fluor 488 phalloidin (1:100) and Hoecsht 33258 (1:1000) (Invitrogen) overnight at 4 °C followed by imaging. Image Xpress Micro was used for capturing images and HUVEC sprouting was analyzed in MetaXpress software. Three

components of vessel growth were measured. The total outgrowth represents the cumulative length of all sprouts per bead, the mean outgrowth represents the number of sprout bifurcations per bead, and the total processes per beads were determined by counting the number of sprouts originating directly from the cells lining the surface of the bead. For 5 statistical analysis, 4 wells were assessed for each condition and each experiment was repeated 3 times.

Quantitative gene expression analyses

RNA was prepared with RNeasy Mini kit (Qiagen) according to the manufacturer's protocol. 500 ng total RNA was subjected to reverse transcription using the High Capacity 10 cDNA reverse transcription kit (Applied Biosystems). The real-time PCR was run on the Applied Biosystems 7500 machine. Taqman probes for all gene tested were acquired from Applied Biosystems. Relative expression levels of each gene were normalized to actin.

Determination of signature gene expression in clinical samples

Sections from archival tumor samples from patients enrolled in the XELOX treatment 15 arms of NO16966 were evaluated by a pathologist and high tumor content area macro-dissected for subsequent RNA isolation using the FFPE RNA isolation kit (Roche; 7-10 sections per patient). RNA levels were assessed after cDNA synthesis following standard techniques, with the qPCR protocol on the Fluidigm Biomark platform according to the manufacturer's protocol.

Gene signature derivation and application

The \log_2 ratio intensity values from Agilent WMG microarrays and the Robust Multi-array Average (RMA) normalized intensities (also on the logarithmic scale) from Affymetrix Mouse430.2 microarrays were imported into R as expression sets using the package "Biobase," and a linear model was fitted to each feature using the functions "lmFit" and 25 "eBayes" in the "limma" package. Features that were significantly ($p < 0.05$) down-regulated in anti-VEGF treated samples (as compared to anti-ragweed control treated samples) were retained and translated to EntrezGene identifiers. The union of these genes was taken as representative of the VEGF-responsive vasculature. The extent to which this VDV gene expression signature varied in other experiments was determined by fitting a linear model to 30 the microarray data using the "limma" package (as above) and calculating the mean of the t-statistics from gene in the signature according to the method of Falcon and Gentleman (*Bioinformatics*. 23: 257-258, 2007).

Statistical methods of biomarker selection

The statistical tasks can comprise the following steps:

1. Pre-selection of candidate biomarkers
2. Pre-selection of relevant clinical efficacy response predictive covariates
3. Selection of biomarker prediction functions at a univariate level
- 5 4. Selection of biomarker prediction functions including clinical covariates at a univariate level
5. Selection of biomarker prediction functions at a multivariate level
6. Selection of biomarker prediction functions including clinical covariates at a multivariate level

10 The following text details the different steps:

1: Pre-selection of candidate biomarkers

The statistical pre-selection of candidate biomarkers is oriented towards the strength of association with measures of clinical benefit. For this purpose the different clinical endpoints may be transformed in derived surrogate scores, as, *e.g.*, an ordinal assignment of the degree of clinical benefit scores regarding TTP that avoid censored observations. These surrogate transformed measures can be easily used for simple correlation analysis, *e.g.* by the non-parametric Spearman rank correlation approach. An alternative is to use the biomarker measurements as metric covariates in time-to-event regression models, as, *e.g.*, Cox proportional hazard regression. Depending on the statistical distribution of the biomarker values, this step may require some pre-processing, as, for example, variance-stabilizing transformations and the use of suitable scales or, alternatively, a standardization step such as using percentiles instead of raw measurements. A further approach is inspection of bivariate scatter plots, for example, by displaying the scatter of (x-axis=biomarker value, y-axis=measure of clinical benefit) on a single-patient basis. Some non-parametric regression line as achieved, for example, by smoothing splines can be useful to visualize the association of biomarker and clinical benefit.

The goal of these different approaches is the pre-selection of biomarker candidates that show some association with clinical benefit in at least one of the benefit measures employed, while results for other measures are not contradictory. When there are available control groups, then differences in association of biomarkers with clinical benefit in the different arms could be a sign of differential prediction that makes the biomarker(s) eligible for further consideration.

2: Pre-selection of relevant clinical efficacy response predictive covariates

The statistical pre-selection of clinical covariates as defined herein parallels the approaches for pre-selecting biomarkers and is also oriented towards the strength of association with measures of clinical benefit. So in principle the same methods apply as considered under 1 above. In addition to statistical criteria, criteria from clinical experience and theoretical knowledge may apply to pre-select relevant clinical covariates.

The predictive value of clinical covariates could interact with the predictive value of the biomarkers. They will be considered for refined prediction rules, if necessary.

3: Selection of biomarker prediction functions at a univariate level

The term “prediction function” will be used in a general sense to mean a numerical function of a biomarker measurement that results in a number scaled to imply the target prediction.

A simple example is the choice of the Heaviside function for a specific cutoff c and a biomarker measurement x , where the binary prediction A or B is to be made, then if $H(x-c)=0$, then predict A. If $H(x-c)=1$, then predict B.

This is probably the most common way of using univariate biomarker measurements in prediction rules. The definition of “prediction function” as noted above includes referral to an existing training data set that can be used to explore the prediction possibilities. Different routes can be taken to achieve a suitable cutoff c from the training set. First, the scatterplot with smoothing spline mentioned under 1 can be used to define the cutoff. Alternatively, some percentile of the distribution could be chosen, e.g., the median or a quartile. Cutoffs can also be systematically extracted by investigating all possible cutoffs according to their prediction potential with regard to the measures of clinical benefit. Then, these results can be plotted to allow for an either manual selection or to employ some search algorithm for optimality. This can be realized based on certain clinical endpoints using a Cox model, wherein at each test cutoff the biomarker is used as a binary covariate. Then the results for the clinical endpoints can be considered together to chose a cutoff that shows prediction in line with both endpoints.

Another uncommon approach for choosing a prediction function can be based on a fixed-parameter Cox regression model obtained from the training set with biomarker values (possibly transformed) as covariate. A further possibility is to base the decision on some likelihood ratio (or monotonic transform of it), where the target probability densities are pre-

determined in the training set for separation of the prediction states. Then the biomarker would be plugged into some function of predictive criteria.

4: Selection of biomarker prediction functions including clinical covariates at a univariate level

5 Univariate refers to using only one biomarker--with regard to clinical covariates, this can be a multivariate model. This approach parallels the search without clinical covariates, except that the methods should allow for incorporating the relevant covariate information. The scatterplot method of choosing a cutoff allows only a limited use of covariates, *e.g.*, a binary covariate could be color coded within the plot. If the analysis relies on some
10 regression approach, then the use of covariates (also many of them at a time) is usually facilitated. The cutoff search based on the Cox model described under 3 above allows for an easy incorporation of covariates and thereby leads to a covariate adjusted univariate cutoff search. The adjustment by covariates may be done as covariates in the model or via the inclusion in a stratified analysis.

15 Also the other choices of prediction functions allow for the incorporation of covariates.

This is straightforward for the Cox model choice as prediction function. This includes the option to estimate the influence of covariates on an interaction level, which means that, *e.g.*, for different age groups different predictive criteria apply.

20 For the likelihood ratio type of prediction functions, the prediction densities must be estimated including covariates. For this purpose, the methodology of multivariate pattern recognition can be used or the biomarker values can be adjusted by multiple regression on the covariates (prior to density estimation).

The CART technology (*Classification and Regression Trees*; Breiman *et al.*
25 (Wadsworth, Inc.: New York, 1984) can be used for this purpose, employing a biomarker (raw measurement level) plus clinical covariates and utilizing a clinical benefit measure as response. Cutoffs are searched and a decision-tree type of function will be found involving the covariates for prediction. The cutoffs and algorithms chosen by CART are frequently close to optimal and may be combined and unified by considering different clinical benefit
30 measures.

5: Selection of biomarker prediction functions at a multivariate level

When there are several biomarker candidates that maintain their prediction potential within the different univariate prediction function choices, then a further improvement may

be achieved by combinations of biomarkers, *i.e.*, considering multivariate prediction functions.

Based on the simple Heaviside function model, combinations of biomarkers may be evaluated, *e.g.*, by considering bivariate scatterplots of biomarker values where optimal cutoffs are indicated. Then a combination of biomarkers can be achieved by combining different Heaviside function by the logical “AND” and “OR” operators to achieve an improved prediction.

The CART technology can be used for this purpose, employing multiple biomarkers (raw measurement level) and a clinical benefit measure as response, to achieve cutoffs for biomarkers and decision-tree type of functions for prediction. The cutoffs and algorithms chosen by CART are frequently close to optimal and may be combined and unified by considering different clinical benefit measures.

The Cox-regression can be employed on different levels. A first way is to incorporate the multiple biomarkers in a binary way (*i.e.*, based on Heaviside functions with some cutoffs). The other option is to employ biomarkers in a metric way (after suitable transformations), or a mixture of the binary and metric approach. The evolving multivariate prediction function is of the Cox type as described under 3 above.

The multivariate likelihood ratio approach is difficult to implement, but presents another option for multivariate prediction functions.

6: Selection of biomarker prediction functions including clinical covariates at a multivariate level

When there are relevant clinical covariates, then a further improvement may be achieved by combining multiple biomarkers with multiple clinical covariates. The different prediction function choices will be evaluated with respect to the possibilities to include clinical covariates.

Based on the simple logical combinations of Heaviside functions for the biomarkers, further covariates may be included to the prediction function based on the logistic regression model obtained in the training set.

The CART technology and the evolving decision trees can be easily used with additional covariates, which would include these in the prediction algorithm.

All prediction functions based on the Cox-regression can use further clinical covariates. The option exists to estimate the influence of covariates on an interaction level, which means that, *e.g.*, for different age groups different predictive criteria apply.

The multivariate likelihood ratio approach is not directly extendible to the use of additional covariates.

Statistical methods of histological and clinical data analysis

Quantitative histological data from various animal experiments were plotted using Microsoft Excel software. Student's t test was applied to compare data sets of interest and differences with p values < 0.05 were considered significant. 103 biopsies out of 1017 patients from the XELOX (capecitabine and oxaliplatin)-containing arm in the NO16966 trial were analyzed for the gene expression of VDV genes. qRT-PCR values were normalized by the housekeeping genes and relative to a universal reference sample to derive delta-delta Ct values. Subsequently, the delta-delta Ct value of each of the 22 VDV genes was mean centered and variance scaled to a Z-score.

For each sample i for which all of the VDV genes were analyzed, a VDV signature score (VDV_i) was calculated. The VDV_i represents a weighted average of z-scores across the analyzed VDV genes and is given by the algorithm:

$$VDV_i = \frac{1}{\sqrt{n}} \sum_{g=1}^n Z_{g,i}$$

wherein $Z_{g,i}$ is the standardized z-score of the qRT-PCR value for gene g of sample i and n , in this instance, is 22. The VDV_i value provides quantitative information regarding the extent to which the expression of a particular set of genes is collectively overexpressed or underexpressed relative to a centered mean.

To compare the clinical outcomes (PFS and OS) between marker and/or treatment subgroups, log-rank tests and Cox regression were used, with median time calculated by the Kaplan-Meier analysis. All statistical tests were two-sided.

Example 2. Identification of genes expressed in the tumor vascular compartment targeted by anti-VEGF

As a first step towards identification of direct *in vivo* biomarkers of VEGF pathway inhibition activity, we characterized the biological consequences of VEGF neutralization in an established transgenic murine model of pancreatic neuroendocrine tumors (PNETs). In the highly vascularized *RIP-TβAg* genetically engineered tumor mouse model (GEMM), anti-VEGF monoclonal antibody (mAb) treatment has previously been shown to have anti-tumor efficacy and to increase overall survival (Singh et al. *J. Pathology* 227(4): 417-430, 2012).

Histological analyses of *RIP-TβAg* late-stage tumors showed that anti-VEGF treatment caused a rapid reduction in micro-vascular density (MVD), detectable at 72 hours post-treatment, and reaching a plateau of approximately 50% at day 7 (Fig. 1A, left). The anti-VEGF induced pruning of this fraction of the tumor vasculature (hereinafter referred to as “VEGF-dependent vasculature” or VDV) is not significantly reversed nor increased at later treatment time points (Fig. 1A, left). In these experiments, tumor vessel density and proliferative index was assessed histologically via MECA-32 staining (red, left) and Ki67 staining (red, right), respectively. Nuclei were counterstained with DAPI (blue). In contrast to the observed rapid anti-vascular effects, the indirect anti-tumor effects of VEGF neutralization progressed more slowly: a reduction in the tumor proliferative index in the anti-VEGF treated group compared to the control (anti-ragweed) treated group was not observed at day 7 but obvious at day 14 (Fig. 1A, right), and a consequent reduction in tumor burden was only evident at day 21 (Fig. 1B). This suggests that at early time-points, the biological consequences of VEGF blockade in this model are primarily vascular specific.

Expression microarray analysis of whole tumors from animals treated for seven days with anti-VEGF showed that the vast majority of genes were unchanged in expression, as compared to tumors from animals treated with a control antibody. However, a small population of genes responded to anti-VEGF treatment with a significant (adjusted $p < 0.01$) decrease in transcript abundance (Table 3). Interestingly, we observed no corresponding up-regulation of gene expression, suggesting that gene expression changes were primarily driven by physical elimination of VEGF-dependent tumor-associated endothelial cells (Fig. 1C and Table 3). As depicted in Table 3 below, the genes in the VDV signature were ordered by the extent of anti-VEGF response in the human IBC trial. Orthologs between human and mouse were mapped using the Ensembl Biomart, where “Log2FC” represents the \log_2 fold-change of the gene’s expression post-Avastin treatment as compared to pre-treatment. The 19 paired patient samples and the murine PNETs were analyzed on Agilent microarrays. A small number of the VDV genes did not have human probes on the Agilent microarrays and are denoted by NA at the end of the table. In addition, probes for some murine VDV genes were not present on the Agilent platform; for these genes, Log2Fc is shown from the repeat experiment on the murine Affymetrix array platform and marked with an asterisk*.

Characterization of the genes within this set having differential (decreased) expression revealed enrichment for known endothelial specific genes (Table 3) implicated in blood vessel development (Table 4). In addition, the median fold-change in expression for this

gene set was similar to that seen in the pan-vascular markers CD31 and PLVAP, and was also consistent with the extent of MVD decrease as measured by immunohistochemistry of the pan-vascular marker PLVAP (Fig. 1A and 1C).

Quantitative PCR (qRT-PCR) of whole tumors confirmed the microarray results, verifying that markers specific to: tumor (insulin); epithelial (E-Cad and Epcam); pan-hematopoietic (CD45); or macrophage (CD68) cells are not significantly changed by anti-VEGF treatment. Also confirming microarray data, qRT-PCR showed that while VEGF transcript levels are not significantly changed, multiple known endothelial markers are downregulated by this treatment (Fig. 1D).

Both microarray and qRT-PCR data identified a subset of genes whose downregulation in response to anti-VEGF was more pronounced than other genes in the signature (Fig. 1D), suggesting that some of the signature genes might be more selectively expressed in the vessels that are sensitive to anti-VEGF treatment. In contrast to low- and intermediate-responding genes (Fig. 1D, yellow and red bars, respectively), this set of extra-responsive genes (Fig. 1D, purple bars) include tip-cell markers and, in the context of developmental retinal angiogenesis, known VEGF targets (Toro et al. *Blood*. 116: 4025-4033, 2010; Roberts et al. *Mol Cell Biol*. 24: 10515-10528, 2004; Testori et al. *Blood*. 117: 2735-2744, 2011; Lobov et al. *Blood*. 117: 6728-6737, 2011). Thus, we postulated that these later genes are candidate proximal biomarkers of VEGF pathway inhibitor activity (proxVDV genes), and are likely to be VEGF targets that are more selectively expressed in the VEGF-dependent tumor vasculature. Taken together, these results suggest that VDV gene expression signature reflects at least two related biological processes: (i) direct VEGF downstream signaling inhibition, and (ii) the subsequent loss of vessels that are dependent on VEGF signaling for survival. According to this working hypothesis, VDV endothelial genes are likely to include proximal (proxVDV) as well as more distal (distVDV) downstream surrogate markers of VEGF signaling inhibition in tumor-associated endothelial cells.

Table 3. Genes in the VDV signature, ordered by extent of anti-VEGF response in human IBC trial.

Human Inflammatory Breast Cancer			Murine Pancreatic Neuroendocrine Tumors		
Entrezgene	Symbol	Log2FC	Entrezgene	Symbol	Log2FC
11082	ESM1	-1.928	71690	Esm1	-3.625
25780	RASGRP3	-1.278	240168	Rasgrp3	-1.777
10060	ABCC9	-1.229	20928	Abcc9	-1.232
3910	LAMA4	-1.219	16775	Lama4	-2.712

22795	NID2	-1.094	18074	Nid2	-3.818
4162	MCAM	-1.005	84004	Mcam	-1.731
8828	NRP2	-0.897	18187	Nrp2	-1.842
7127	TNFAIP2	-0.854	21928	Tnfaip2	-0.367
6678	SPARC	-0.843	20692	Sparc	-1.746
27253	PCDH17	-0.829	219228	Pcdh17	-1.909
23743	BHMT2	-0.808	64918	Bhmt2	-0.163
51313	FAM198B	-0.794	68659	Fam198b	-0.26
9444	QKI	-0.763	19317	Qk	-0.652
1306	COL15A1	-0.736	12819	Col15a1	-2.274
23627	PRND	-0.732	26434	Prnd	-2.48
6772	STAT1	-0.709	20846	Stat1	-0.227
8829	NRP1	-0.707	18186	Nrp1	-1.434
2200	FBN1	-0.701	14118	Fbn1	-1.139
28984	C13orf15	-0.698	66214	1190002H23Rik	-2.238
285	ANGPT2	-0.685	11601	Angpt2	-0.85
419	ART3	-0.675	109979	Art3	-0.643
85480	TSLP	-0.646	53603	Tslp	0.108
4208	MEF2C	-0.644	17260	Mef2c	-1.316
10008	KCNE3	-0.643	57442	Kcne3	-1.938
10395	DLC1	-0.64	50768	Dlc1	-0.745*
22918	CD93	-0.636	17064	Cd93	-2.406
25937	WWTR1	-0.631	97064	Wwtr1	-1.261
1848	DUSP6	-0.614	67603	Dusp6	-1.427
6649	SOD3	-0.578	20657	Sod3	0.144
4005	LMO2	-0.572	16909	Lmo2	-0.057
29969	MDFIC	-0.569	16543	Mdfic	-0.183
51088	KLHL5	-0.557	71778	Klhl5	-0.489*
83595	SOX7	-0.553	20680	Sox7	-0.765*
5414	SEPT4	-0.531	18952	Sept4	-0.814
55917	CTTNBP2NL	-0.531	80281	Ctnbp2nl	-0.501
1282	COL4A1	-0.501	12826	Col4a1	-1.998
301	ANXA1	-0.5	16952	Anxa1	-1.31
1284	COL4A2	-0.499	12827	Col4a2	-1.742
7431	VIM	-0.493	22352	Vim	-1.321
8613	PPAP2B	-0.488	67916	Ppap2b	-2.638
112464	PRKCDBP	-0.487	109042	Prkcdbp	-1.395
1009	CDH11	-0.487	12552	Cdh11	-1.326
1295	COL8A1	-0.483	12837	Col8a1	-1.433
23493	HEY2	-0.481	15214	Hey2	-0.327
10186	LHFP	-0.467	108927	Lhfp	-0.481
2331	FMOD	-0.462	14264	Fmod	0.007
7057	THBS1	-0.46	21825	Thbs1	-0.743
659	BMPR2	-0.442	12168	Bmpr2	-0.346*

10544	PROCR	-0.441	19124	Procr	-0.743
7070	THY1	-0.435	21838	Thy1	-0.757
5027	P2RX7	-0.431	18439	P2rx7	-0.227
5159	PDGFRB	-0.43	18596	Pdgfrb	-1.35
871	SERPINH1	-0.419	12406	Serpinh1	-2.022
8490	RGS5	-0.419	19737	Rgs5	-1.002
54510	PCDH18	-0.418	73173	Pcdh18	-0.074
5787	PTPRB	-0.409	19263	Ptprb	-2.028
857	CAV1	-0.404	12389	Cav1	-0.826
54538	ROBO4	-0.386	74144	Robo4	-1.605
8611	PPAP2A	-0.385	19012	Ppap2a	-0.933
10791	VAMP5	-0.378	53620	Vamp5	-0.815
5552	SRGN	-0.378	19073	Srgn	-0.902
162073	ITPRIPL2	-0.378	319622	Itpripl2	-0.707*
51294	PCDH12	-0.362	53601	Pcdh12	-1.515
4781	NFIB	-0.362	18028	Nfib	-0.945
716	C1S	-0.357	50908	C1s	-0.122
595	CCND1	-0.355	12443	Ccnd1	-1.116
1464	CSPG4	-0.355	121021	Cspg4	-1.713
54922	RASIP1	-0.348	69903	Rasip1	-1.423
3384	ICAM2	-0.347	15896	Icam2	-2.346
80781	COL18A1	-0.342	12822	Col18a1	-1.659
2828	GPR4	-0.341	319197	Gpr4	-0.14
2313	FLI1	-0.334	14247	Fli1	-1.218
57381	RHOJ	-0.33	80837	Rhoj	-1.463
79890	RIN3	-0.329	217835	Rin3	-0.137
51705	EMCN	-0.328	59308	Emcn	-2.347
11167	FSTL1	-0.324	14314	Fstl1	-1.146
79689	STEAP4	-0.321	117167	Steap4	0.26
3200	HOXA3	-0.319	15400	Hoxa3	0.271
25925	ZNF521	-0.319	225207	Zfp521	-0.482
5797	PTPRM	-0.308	19274	Ptprm	-0.479
8642	DCHS1	-0.301	233651	Dchs1	-0.367*
29015	SLC43A3	-0.301	58207	Slc43a3	-0.647
116362	RBP7	-0.299	63954	Rbp7	-2.883
1404	HAPLN1	-0.294	12950	Hapln1	0.2
1490	CTGF	-0.286	14219	Ctgf	-2.002
5937	RBMS1	-0.282	56878	Rbms1	-0.663
158326	FREM1	-0.281	329872	Frem1	-0.307*
56892	C8orf4	-0.279	69068	1810011O10Rik	-2.026
161198	CLEC14A	-0.273	66864	Clec14a	-0.563
10769	PLK2	-0.27	20620	Plk2	-1.322
51751	HIGD1B	-0.264	75689	Higd1b	-0.913
8038	ADAM12	-0.264	11489	Adam12	0.019

55901	THSD1	-0.261	56229	Thsd1	-1.379
51162	EGFL7	-0.259	353156	Egfl7	-0.934
25960	GPR124	-0.248	78560	Gpr124	-0.78
26524	LATS2	-0.247	50523	Lats2	-0.33
90952	ESAM	-0.244	69524	Esam	-1.027
1265	CNN2	-0.243	12798	Cnn2	-0.726
51676	ASB2	-0.243	65256	Asb2	-0.012
10536	LEPREL2	-0.242	14789	Leprel2	-0.925
2022	ENG	-0.235	13805	Eng	-2.167
28951	TRIB2	-0.235	217410	Trib2	-0.782*
64411	ARAP3	-0.232	106952	Arap3	-2.098
196	AHR	-0.228	11622	Ahr	-0.693
10003	NAALAD2	-0.227	72560	Naalad2	-0.356
256949	KANK3	-0.226	80880	Kank3	-0.888
81848	SPRY4	-0.225	24066	Spry4	-1.589
2122	MECOM	-0.223	14013	Mecom	-0.175
51363	CHST15	-0.221	77590	Chst15	-0.101
6623	SNCG	-0.215	20618	Sncg	-0.7
4071	TM4SF1	-0.215	17112	Tm4sf1	-0.752
116159	CYYR1	-0.211	224405	Cyyr1	0.114
2149	F2R	-0.21	14062	F2r	-0.815
79899	PRR5L	-0.209	72446	Prr5l	-0.617
84898	PLXDC2	-0.209	67448	Plxdc2	-1.16
53827	FXD5	-0.206	18301	Fxd5	-0.728
10561	IFI44	-0.203	99899	Ifi44	-1.481
166336	PRICKLE2	-0.203	243548	Prickle2	-0.003
80310	PDGFD	-0.2	71785	Pdgfd	-0.386
2321	FLT1	-0.2	14254	Flt1	-2.762
2621	GAS6	-0.198	14456	Gas6	-0.895
83878	USHBP1	-0.191	234395	Ushbp1	-2.261
3625	INHBB	-0.19	16324	Inhbb	-1.283
677	ZFP36L1	-0.187	12192	Zfp36l1	-1.117
84632	AFAP1L2	-0.186	226250	Afap1l2	-1.297
10516	FBLN5	-0.186	23876	Fbln5	-0.457*
23075	SWAP70	-0.184	20947	Swap70	-0.19
5724	PTAFR	-0.18	19204	Ptafr	-0.639*
84962	JUB	-0.178	16475	Jub	-0.318
92747	BPIFB1	-0.175	228801	Bpifb1	0.146
2078	ERG	-0.163	13876	Erg	-1.816
27123	DKK2	-0.159	56811	Dkk2	-2.214
26508	HEYL	-0.153	56198	Heyl	-0.699
51267	CLEC1A	-0.151	243653	Clec1a	-0.636
5793	PTPRG	-0.151	19270	Ptprg	-1.015
10161	LPAR6	-0.148	67168	Lpar6	-0.587

154810	AMOTL1	-0.147	75723	Amotl1	-0.114
286753	TUSC5	-0.146	237858	Tusc5	-0.358
9411	ARHGAP29	-0.145	214137	Arhgap29	-0.42
6237	RRAS	-0.143	20130	Rras	-0.164
57514	ARHGAP31	-0.141	12549	Arhgap31	-0.98
81792	ADAMTS12	-0.139	239337	Adamts12	-0.352*
947	CD34	-0.136	12490	Cd34	-2.707
149951	COMMD7	-0.135	99311	Commd7	0.073
9509	ADAMTS2	-0.13	216725	Adamts2	-1.263
7010	TEK	-0.13	21687	Tek	-0.57
9369	NRXN3	-0.128	18191	Nrxn3	0.422
94	ACVRL1	-0.123	11482	Acvrl1	-0.364
5978	REST	-0.123	19712	Rest	-0.851
64123	ELTD1	-0.122	170757	Eltld1	-1.681
128077	LIX1L	-0.12	280411	Lix1l	-0.488*
684	BST2	-0.119	69550	Bst2	-0.637
3397	ID1	-0.113	15901	Id1	-2.744
958	CD40	-0.109	21939	Cd40	-0.623*
4811	NID1	-0.103	18073	Nid1	-2.257
3486	IGFBP3	-0.101	16009	Igfbp3	-1.989
10345	TRDN	-0.1	76757	Trdn	-0.338*
114571	SLC22A9	-0.099	319800	Slc22a30	0.088
4837	NNMT	-0.097	18113	Nnmt	-0.29*
57608	KIAA1462	-0.091	240185	9430020K01Rik	-1.869
9749	PHACTR2	-0.09	215789	Phactr2	-0.597*
7078	TIMP3	-0.084	21859	Timp3	-1.573
2113	ETS1	-0.079	23871	Ets1	-1.224
80381	CD276	-0.078	102657	Cd276	-0.282*
30844	EHD4	-0.07	98878	Ehd4	-0.483
2701	GJA4	-0.068	14612	Gja4	-0.431
170575	GIMAP1	-0.066	16205	Gimap1	-1.332
285203	C3orf64	-0.06	101351	A130022J15Rik	-0.485
4240	MFGE8	-0.06	17304	Mfge8	-1.199
5089	PBX2	-0.057	18515	Pbx2	-0.127
3234	HOXD8	-0.052	15437	Hoxd8	0.089
9647	PPM1F	-0.052	68606	Ppm1f	-0.385
474344	GIMAP6	-0.052	231931	Gimap6	-1.191
83483	PLVAP	-0.051	84094	Plvap	-0.983
5155	PDGFB	-0.05	18591	Pdgfb	-0.884*
55340	GIMAP5	-0.049	317757	Gimap5	-1.194
7157	TP53	-0.045	22059	Trp53	-0.207*
4061	LY6E	-0.042	17069	Ly6e	-0.375
79812	MMRN2	-0.04	105450	Mmrn2	-2.415
23576	DDAH1	-0.031	69219	Ddah1	-0.14

153579	BTNL9	-0.031	237754	Btnl9	-0.913
633	BGN	-0.029	12111	Bgn	-0.797*
27143	KIAA1274	-0.027	27355	X99384	-0.52
375387	LRRC33	-0.025	224109	Lrrc33	-0.333
127733	UBXN10	-0.023	212190	Ubxn10	-0.293*
3820	KLRB1	-0.017	80782	Klr1b	-0.301*
80177	MYCT1	-0.015	68632	Myct1	-1.574
3791	KDR	-0.014	16542	Kdr	-1.778
9862	MED24	-0.012	23989	Med24	-0.242*
8522	GAS7	-0.011	14457	Gas7	-0.122
56062	KLHL4	-0.011	237010	Klhl4	-0.462
54345	SOX18	-0.008	20672	Sox18	-1.532
85363	TRIM5	-0.008	20128	Trim30a	-0.837
285852	TREML4	-0.006	224840	Trem14	-0.134
9510	ADAMTS1	-0.005	11504	Adamts1	-0.491
8840	WISP1	-0.005	22402	Wisp1	-0.866
10411	RAPGEF3	-0.004	223864	Rapgef3	-1.67
7903	ST8SIA4	-0.003	20452	St8sia4	-0.286
55303	GIMAP4	0.002	107526	Gimap4	-1.127
8824	CES2	0.003	234673	Ces2e	-2.514
89857	KLHL6	0.009	239743	Klhl6	-0.442
10052	GJC1	0.009	14615	Gjc1	-0.894
3561	IL2RG	0.01	16186	Il2rg	-0.028
56253	CRTAM	0.02	54698	Crtam	-0.022
5583	PRKCH	0.026	18755	Prkch	-0.104
6300	MAPK12	0.026	29857	Mapk12	-0.182
162394	SLFN5	0.029	327978	Slfn5	-0.515*
1807	DPYS	0.029	64705	Dpys	-0.085
4038	LRP4	0.029	228357	Lrp4	-1.971
387601	SLC22A25	0.038	319800	Slc22a30	0.088
3142	HLX	0.041	15284	Hlx	-0.189
6556	SLC11A1	0.044	18173	Slc11a1	-0.133
6524	SLC5A2	0.046	246787	Slc5a2	-0.306*
92162	TMEM88	0.048	67020	Tmem88	-1.771
64641	EBF2	0.048	13592	Ebf2	-0.12
6405	SEMA3F	0.048	20350	Sema3f	-0.581
9351	SLC9A3R2	0.054	65962	Slc9a3r2	-0.611
7022	TFAP2C	0.054	21420	Tcfap2c	-0.159
187	APLNR	0.055	23796	Aplnr	-0.024
7139	TNNT2	0.058	21956	Tnnt2	-1.175
55679	LIMS2	0.059	225341	Lims2	-0.69
11135	CDC42EP1	0.062	104445	Cdc42ep1	-1.568
976	CD97	0.067	26364	Cd97	-1.036
121599	SPIC	0.07	20728	Spic	-0.668

23604	DAPK2	0.072	13143	Dapk2	-1.32
1903	S1PR3	0.073	13610	S1pr3	0.028
23439	ATP1B4	0.083	67821	Atp1b4	0.141
338339	CLEC4D	0.091	17474	Clec4d	-0.149
9002	F2RL3	0.093	14065	F2rl3	-0.122
4846	NOS3	0.112	18127	Nos3	-0.819
1901	S1PR1	0.127	13609	S1pr1	-0.315
11037	STON1	0.144	77057	Ston1	-0.387*
50619	DEF6	0.146	23853	Def6	-0.068
1003	CDH5	0.149	12562	Cdh5	-2.001
55215	FANCI	0.15	208836	Fanci	0.342
84433	CARD11	0.15	108723	Card11	-0.205*
8497	PPFLA4	0.155	68507	Ppfia4	-0.241*
3912	LAMB1	0.181	16777	Lamb1	-2.313
5054	SERPINE1	0.194	18787	Serpine1	-1.055
286749	STON1- GTF2A1L	0.217	77057	Ston1	-0.387*
1285	COL4A3	0.223	12828	Col4a3	0.038
9771	RAPGEF5	0.269	217944	Rapgef5	-0.991
93978	CLEC6A	NA	56620	Clec4n	-0.421
284040	CDRT4	NA	66338	Cdrt4	-0.306
79652	TMEM204	NA	407831	Tmem204	-0.708
100527949	GIMAP1- GIMAP5	NA	317757	Gimap5	-1.194
7058	THBS2	NA	21826	Thbs2	-0.18
338596	ST8SIA6	NA	241230	St8sia6	-0.569
340527	NHSL2	NA	100042480	Nhsl2	0.383
8578	SCARF1	NA	380713	Scarf1	-1.018
402569	KPNA7	NA	381686	Kpna7	-0.362*
441478	NRARP	NA	67122	Nrarp	-0.218
60681	FKBP10	NA	14230	Fkbp10	-0.893

Table 4. Gene ontology terms over-represented in VDV signature.

5 a. Biological Process Ontology

GOBPID	P value	Odds Ratio	Exp Count	Count	Size	Term
GO:0001568	1.659e-30	10.57	6.785	51	385	blood vessel development
GO:0072358	9.829e-27	11.35	5.111	42	307	cardiovascular system development
GO:0032501	8.002e-20	3.465	78.43	145	4610	multicellular organismal process
GO:0001525	6.98e-17	9.984	3.435	27	205	angiogenesis
GO:0009653	7.513e-14	4.884	10.34	40	698	anatomical structure

						morphogenesis
GO:2000145	1.642e-12	6.281	5.146	27	292	regulation of cell motility
GO:2000026	3.997e-12	3.818	14.26	45	809	regulation of multicellular organismal development
GO:0009888	5.801e-12	3.667	15.58	47	901	tissue development
GO:0023052	7.066e-12	2.478	70.26	120	3987	signaling
GO:0007166	1.25e-10	2.676	33.91	72	1924	cell surface receptor linked signaling pathway
GO:0032879	1.428e-10	3.256	17.69	48	1004	regulation of localization
GO:0007155	4.022e-10	4.398	8.01	30	486	cell adhesion
GO:0000902	6.839e-10	3.51	13.06	39	741	cell morphogenesis
GO:0051716	1.061e-09	2.433	52.42	92	3224	cellular response to stimulus
GO:0022603	1.467e-09	4.37	7.419	28	421	regulation of anatomical structure morphogenesis
GO:0001570	8.696e-09	15.27	0.8459	10	48	vasculogenesis
GO:0048869	1.529e-08	3.118	15.44	40	1031	cellular developmental process
GO:0040013	1.842e-08	8.857	1.745	13	99	negative regulation of locomotion
GO:0048468	1.947e-08	2.812	19.23	46	1115	cell development
GO:0048731	2.693e-08	2.53	31.52	62	2288	system development
GO:0030336	3.752e-08	9.341	1.533	12	87	negative regulation of cell migration
GO:0042330	6.307e-08	3.522	9.376	29	532	taxis
GO:0009893	1.02e-07	2.496	24.48	52	1389	positive regulation of metabolic process
GO:0051271	1.021e-07	8.436	1.674	12	95	negative regulation of cellular component movement
GO:0044093	1.213e-07	2.808	16.41	40	931	positive regulation of molecular function
GO:0007411	3.51e-07	4.087	5.78	21	328	axon guidance
GO:0048812	3.608e-07	3.374	9.023	27	512	neuron projection morphogenesis
GO:0030182	3.819e-07	2.824	14.52	36	824	neuron differentiation
GO:0007154	5.952e-07	2.121	47.13	78	2982	cell communication
GO:0040017	6.315e-07	5.375	3.155	15	179	positive regulation of locomotion
GO:0016310	6.357e-07	2.528	19.53	43	1108	phosphorylation
GO:0051272	6.783e-07	5.342	3.172	15	180	positive regulation of cellular component movement
GO:0042221	7.279e-07	2.602	18.43	41	1141	response to chemical stimulus
GO:0048667	9.499e-07	3.275	8.9	26	505	cell morphogenesis involved in neuron differentiation

GO:0050789	1.172e-06	2.17	74.81	105	5270	regulation of biological process
GO:0032990	1.185e-06	3.084	10.19	28	578	cell part morphogenesis
GO:0006793	1.34e-06	2.38	22.21	46	1260	phosphorus metabolic process
GO:0030030	1.641e-06	2.758	13.48	33	765	cell projection organization
GO:0010646	1.784e-06	2.444	19.6	42	1112	regulation of cell communication
GO:0009966	2.756e-06	2.621	15.19	35	923	regulation of signal transduction
GO:0022008	3.002e-06	2.528	16.53	37	938	neurogenesis
GO:0048870	3.186e-06	3.659	6.098	20	382	cell motility
GO:0009887	4.196e-06	2.871	10.87	28	617	organ morphogenesis
GO:0007507	4.451e-06	3.866	5.164	18	293	heart development
GO:0045446	4.495e-06	17.35	0.4527	6	26	endothelial cell differentiation
GO:0001569	4.818e-06	17.13	0.4582	6	26	patterning of blood vessels
GO:0048584	4.947e-06	2.568	14.86	34	843	positive regulation of response to stimulus
GO:0043549	7.131e-06	3.048	8.724	24	495	regulation of kinase activity
GO:0045937	7.801e-06	4.308	3.86	15	219	positive regulation of phosphate metabolic process
GO:0001936	1.231e-05	8.512	1.093	8	62	regulation of endothelial cell proliferation
GO:0008284	1.364e-05	3.179	7.289	21	425	positive regulation of cell proliferation
GO:0030198	1.997e-05	5.91	1.898	10	109	extracellular matrix organization
GO:0008015	2.281e-05	3.708	4.741	16	269	blood circulation
GO:0009725	2.406e-05	2.638	11.3	27	641	response to hormone stimulus
GO:0048534	2.486e-05	3.138	6.997	20	397	hemopoietic or lymphoid organ development
GO:0019220	2.551e-05	2.954	8.206	22	489	regulation of phosphate metabolic process
GO:0010595	2.975e-05	11.81	0.6168	6	35	positive regulation of endothelial cell migration
GO:0010810	3.38e-05	7.291	1.251	8	71	regulation of cell-substrate adhesion
GO:0042698	3.434e-05	6.244	1.621	9	92	ovulation cycle
GO:0001934	3.78e-05	4.201	3.401	13	193	positive regulation of protein phosphorylation
GO:0048010	4.137e-05	11.04	0.6521	6	37	vascular endothelial growth factor receptor signaling pathway
GO:0006950	5.219e-05	1.815	44.18	69	2507	response to stress

GO:0032101	5.353e-05	3.606	4.547	15	258	regulation of response to external stimulus
GO:0061041	5.373e-05	8.175	0.9869	7	56	regulation of wound healing
GO:0032268	5.635e-05	2.661	9.924	24	594	regulation of cellular protein metabolic process
GO:0001945	5.934e-05	25.18	0.2291	4	13	lymph vessel development
GO:0007162	6.034e-05	8.011	1.005	7	57	negative regulation of cell adhesion
GO:0045766	6.761e-05	7.853	1.022	7	58	positive regulation of angiogenesis
GO:0008285	6.909e-05	2.899	7.525	20	427	negative regulation of cell proliferation
GO:0043627	6.995e-05	4.577	2.644	11	150	response to estrogen stimulus
GO:0050900	7.404e-05	3.915	3.63	13	206	leukocyte migration
GO:0021700	7.715e-05	4.973	2.221	10	126	developmental maturation
GO:0001974	7.921e-05	13.54	0.4582	5	26	blood vessel remodeling
GO:0032835	7.921e-05	13.54	0.4582	5	26	glomerulus development
GO:0032270	8.472e-05	2.943	7.032	19	399	positive regulation of cellular protein metabolic process
GO:0035556	9.397e-05	1.921	29.33	50	1664	intracellular signal transduction
GO:0001932	9.729e-05	3.013	6.515	18	391	regulation of protein phosphorylation
GO:0042060	9.976e-05	2.741	8.357	21	492	wound healing
GO:0016337	0.0001106	3.214	5.41	16	307	cell-cell adhesion
GO:0071363	0.0001128	5.282	1.886	9	107	cellular response to growth factor stimulus
GO:0010628	0.0001161	2.276	14.47	30	821	positive regulation of gene expression
GO:0050790	0.0001167	2.351	13.11	28	783	regulation of catalytic activity
GO:0051093	0.0001208	7.078	1.117	7	67	negative regulation of developmental process
GO:0045165	0.0001249	4.267	2.82	11	160	cell fate commitment
GO:0007220	0.0001449	18.88	0.282	4	16	Notch receptor processing
GO:0051345	0.000164	2.876	6.785	18	385	positive regulation of hydrolase activity
GO:0018212	0.0001647	4.503	2.432	10	138	peptidyl-tyrosine modification
GO:0070887	0.0001692	2.072	19.16	36	1087	cellular response to chemical stimulus
GO:0001886	0.0001795	42.33	0.1234	3	7	endothelial cell morphogenesis
GO:0001541	0.0001858	8.144	0.8459	6	48	ovarian follicle development

GO:0010557	0.0002096	2.164	15.68	31	890	positive regulation of macromolecule biosynthetic process
GO:0043086	0.0002104	2.728	7.543	19	428	negative regulation of catalytic activity
GO:0030195	0.0002225	10.52	0.564	5	32	negative regulation of blood coagulation
GO:0034446	0.0002369	16.18	0.3172	4	18	substrate adhesion-dependent cell spreading
GO:0006897	0.0002547	3.098	5.234	15	297	endocytosis
GO:0050679	0.0002688	5.27	1.674	8	95	positive regulation of epithelial cell proliferation
GO:0030168	0.0002748	3.396	4.142	13	235	platelet activation
GO:0065008	0.0002799	1.783	34.8	55	2016	regulation of biological quality
GO:0070168	0.0002835	33.86	0.141	3	8	negative regulation of biomineral tissue development
GO:0051128	0.0002838	2.469	9.684	22	574	regulation of cellular component organization
GO:0007178	0.0002866	6.079	1.281	7	75	transmembrane receptor protein serine/threonine kinase signaling pathway
GO:0090101	0.0002906	7.434	0.9164	6	52	negative regulation of transmembrane receptor protein serine/threonine kinase signaling pathway
GO:0045596	0.0002939	3.054	5.305	15	301	negative regulation of cell differentiation
GO:0042246	0.0002987	9.796	0.5992	5	34	tissue regeneration
GO:0002682	0.0003018	2.271	12.44	26	706	regulation of immune system process
GO:0070482	0.0003078	3.56	3.648	12	207	response to oxygen levels
GO:0001300	0.0003093	Inf	0.03525	2	2	chronological cell aging
GO:0007521	0.0003093	Inf	0.03525	2	2	muscle cell fate determination
GO:0048522	0.0003187	1.841	30.24	49	1922	positive regulation of cellular process
GO:0050818	0.000323	7.275	0.934	6	53	regulation of coagulation
GO:0045860	0.0003382	3.01	5.375	15	305	positive regulation of protein kinase activity
GO:0007599	0.0003383	2.549	8.477	20	481	hemostasis
GO:0001933	0.0003581	7.123	0.9517	6	54	negative regulation of protein phosphorylation
GO:0034329	0.0003606	4.054	2.679	10	152	cell junction assembly
GO:0014706	0.0003664	3.487	3.719	12	211	striated muscle tissue

						development
GO:0035239	0.0003664	3.487	3.719	12	211	tube morphogenesis
GO:0071841	0.0003674	1.663	49.66	72	2818	cellular component organization or biogenesis at cellular level
GO:0030509	0.0003759	5.797	1.339	7	76	BMP signaling pathway
GO:0051384	0.000388	4.418	2.221	9	126	response to glucocorticoid stimulus
GO:0008354	0.0004197	28.22	0.1586	3	9	germ cell migration
GO:0022617	0.0004197	28.22	0.1586	3	9	extracellular matrix disassembly
GO:0045601	0.0004197	28.22	0.1586	3	9	regulation of endothelial cell differentiation
GO:0051918	0.0004197	28.22	0.1586	3	9	negative regulation of fibrinolysis
GO:0060627	0.0004433	3.942	2.749	10	156	regulation of vesicle-mediated transport
GO:0009968	0.0004471	2.717	6.742	17	391	negative regulation of signal transduction
GO:0048732	0.0004527	3.401	3.807	12	216	gland development
GO:0002009	0.0004607	3.205	4.371	13	248	morphogenesis of an epithelium
GO:0051130	0.0004695	2.8	6.151	16	349	positive regulation of cellular component organization
GO:0002576	0.0004767	5.554	1.392	7	79	platelet degranulation
GO:0002064	0.0004815	6.702	1.005	6	57	epithelial cell development
GO:0017015	0.0004815	6.702	1.005	6	57	regulation of transforming growth factor beta receptor signaling pathway
GO:0030855	0.0005169	4.232	2.308	9	136	epithelial cell differentiation
GO:0009612	0.0005305	4.723	1.85	8	105	response to mechanical stimulus
GO:0048762	0.0005305	4.723	1.85	8	105	mesenchymal cell differentiation
GO:0050730	0.0005305	4.723	1.85	8	105	regulation of peptidyl-tyrosine phosphorylation
GO:0061061	0.0005315	2.766	6.221	16	353	muscle structure development
GO:0030335	0.0005443	4.201	2.325	9	135	positive regulation of cell migration
GO:0002274	0.000555	5.403	1.427	7	81	myeloid leukocyte activation
GO:0001503	0.0005782	3.302	3.912	12	222	ossification
GO:0006936	0.0005782	3.302	3.912	12	222	muscle contraction
GO:0048660	0.0005806	6.448	1.04	6	59	regulation of smooth muscle cell proliferation
GO:0008585	0.0005979	5.331	1.445	7	82	female gonad

						development
GO:0014910	0.0006398	11.92	0.4053	4	23	regulation of smooth muscle cell migration
GO:2000379	0.0006398	11.92	0.4053	4	23	positive regulation of reactive oxygen species metabolic process
GO:0019538	0.0006644	1.598	56.89	79	3228	protein metabolic process
GO:0071900	0.0006835	2.921	5.146	14	292	regulation of protein serine/ threonine kinase activity
GO:0051347	0.0006995	2.793	5.763	15	327	positive regulation of transferase activity
GO:0016044	0.0007038	2.34	9.658	21	548	cellular membrane organization
GO:0051704	0.0007045	2.023	16.09	30	913	multi-organism process
GO:0009790	0.0007198	2.202	11.75	24	667	embryo development
GO:0033002	0.0007424	5.124	1.498	7	85	muscle cell proliferation
GO:0045944	0.0007803	2.432	8.389	19	476	positive regulation of transcription from RNA polymerase II promoter
GO:0032496	0.0007913	3.639	2.961	10	168	response to lipopolysaccharide
GO:0001775	0.0007992	2.571	7.103	17	419	cell activation
GO:0001953	0.0008032	21.16	0.1939	3	11	negative regulation of cell-matrix adhesion
GO:0017187	0.0008032	21.16	0.1939	3	11	peptidyl-glutamic acid carboxylation
GO:0080090	0.0008098	1.558	67.27	90	3817	regulation of primary metabolic process
GO:0050673	0.0008897	4.954	1.543	7	90	epithelial cell proliferation
GO:0010757	0.0009171	112.4	0.05287	2	3	negative regulation of plasminogen activation
GO:0001819	0.0009233	3.881	2.503	9	142	positive regulation of cytokine production
GO:0031328	0.0009347	1.962	17.13	31	972	positive regulation of cellular biosynthetic process
GO:0008360	0.000975	5.79	1.146	6	65	regulation of cell shape
GO:0030099	0.0009911	3.526	3.049	10	173	myeloid cell differentiation

b. Cellular Component Ontology

GOCCID	Pvalue	OddsRatio	ExpCount	Count	Size	Term
GO:0044421	6.134e-11	6.593	3.923	22	252	extracellular region part
GO:0005581	1.2e-08	10.43	1.384	12	81	collagen
GO:0005615	6.927e-08	3.082	12.87	35	753	extracellular space
GO:0005886	2.337e-07	2.486	24.72	51	1707	plasma membrane
GO:0044459	3.383e-07	2.294	28.98	57	1748	plasma membrane

						part
GO:0005604	3.837e-07	11.44	0.9494	9	57	basement membrane
GO:0030141	7.723e-06	4.297	3.846	15	225	stored secretory granule
GO:0005887	1.712e-05	2.235	19.92	40	1174	integral to plasma membrane
GO:0005796	5.637e-05	14.64	0.4273	5	25	Golgi lumen
GO:0009897	6.485e-05	5.071	2.171	10	127	external side of plasma membrane
GO:0005587	9.504e-05	58.16	0.1026	3	6	collagen type IV
GO:0031093	0.0001243	8.805	0.7862	6	46	platelet alpha granule lumen
GO:0005605	0.0001291	19.45	0.2735	4	16	basal lamina
GO:0005606	0.0001642	43.62	0.1196	3	7	laminin-1 complex
GO:0060205	0.0001778	8.189	0.8375	6	49	cytoplasmic membrane-bounded vesicle lumen
GO:0045121	0.0004315	3.947	2.735	10	160	membrane raft
GO:0005788	0.0005969	4.621	1.88	8	110	endoplasmic reticulum lumen
GO:0005912	0.0006938	3.697	2.906	10	170	adherens junction
GO:0043259	0.0008633	115.9	0.05128	2	3	laminin-10 complex
GO:0070022	0.0008633	115.9	0.05128	2	3	transforming growth factor beta receptor complex
GO:0005578	0.0009301	3.548	3.018	10	191	proteinaceous extracellular matrix

c. Molecular Function Ontology

GOMFID	Pvalue	OddsRatio	ExpCount	Count	Size	Term
GO:0005201	4.706e-11	13.3	1.329	14	80	extracellular matrix structural constituent
GO:0005021	5.369e-10	182.3	0.1329	6	8	vascular endothelial growth factor-activated receptor activity
GO:0005509	1.059e-08	3.583	10.28	32	619	calcium ion binding
GO:0005178	3.02e-08	10.98	1.213	11	73	integrin binding
GO:0019199	1.52e-07	9.187	1.412	11	85	transmembrane

						receptor protein kinase activity
GO:0050431	5.162e-07	50.39	0.1827	5	11	transforming growth factor beta binding
GO:0060089	2.742e-06	2.354	21.36	44	1286	molecular transducer activity
GO:0005518	8.458e-06	11.21	0.7476	7	45	collagen binding
GO:0038023	6.846e-05	2.287	15.45	32	930	signaling receptor activity
GO:0017154	8.724e-05	59.96	0.09968	3	6	semaphorin receptor activity
GO:0005024	0.000149	18.52	0.2824	4	17	transforming growth factor beta-activated receptor activity
GO:0008289	0.0001585	2.793	7.376	19	444	lipid binding
GO:0030246	0.0001911	2.937	6.263	17	377	carbohydrate binding
GO:0005539	0.0002787	4.192	2.592	10	156	glycosaminoglycan binding
GO:0019899	0.0003593	2.176	14	28	843	enzyme binding
GO:0001871	0.0005776	3.797	2.841	10	171	pattern binding
GO:0005520	0.0006073	12.03	0.3987	4	24	insulin-like growth factor binding
GO:0048407	0.0006767	22.48	0.1827	3	11	platelet-derived growth factor binding
GO:0019834	0.0008154	119.4	0.04984	2	3	phospholipase A2 inhibitor activity

Example 3. Vascular responses to VEGF signaling blockade are conserved across multiple tumor models

VDV Responses to VEGF signaling blockade are stromal specific and conserved across multiple tumor models

We sought to determine whether the VDV transcriptional signature identified in the murine late-stage PNET model is also detectable in other tumor models.

First, we analyzed whole-tumor responses to anti-VEGF treatment in samples from an established (400 mm³) subcutaneous human breast carcinoma tumor model (MDA-MB-231). While unsupervised expression analysis failed to distinguish between anti-VEGF and control treatment samples, short-term (i.e., 24 hours) anti-VEGF treatment was sufficient to induce a significant downward shift in expression of the vast majority of the VDV genes relative to all other genes ($p < 0.0001$; Fig. 2A, upper panel). Consistent with the hypothesis that VDV genes are endothelial specific, expression changes in these genes were detected only by probes on the murine, but not the human microarrays, corresponding to the distinction between stromal and tumor cells (Fig. 2A). Also, in agreement with the data from the *RIP-TβAg* GEMM, treatment of the MDA-MB231 subcutaneous xenograft breast cancer tumors

with anti-VEGF induced a steeper downregulation of candidate proxVDV gene candidates, relative to pan-vascular markers and other VDV genes.

We found that the effects of long-term anti-VEGF treatment in a third model, an orthotopic, intracranial U87 glioblastoma, resulted also in a pattern of decreased VDV gene expression, with exclusive detection by the mouse-specific probes; and with more marked downregulation of candidate proxVDV genes (Fig. 2B). Comparable VDV patterns of response to anti-VEGF antibody were also consistently observed across multiple anti-VEGF treated human xenograft and murine tumor models tested.

Thus, regardless of the tumor model and implantation site, and independent of the length of antibody treatment, the VDV signature (comprised by proxVDV and distVDV genes) enables consistent detection of gene expression changes that reflect vascular downstream biological consequences of VEGF pathway inhibition in whole tumor mRNA samples.

VEGF signaling induces VDV gene expression

Next we assessed the extent to which VEGF stimulation could, by reciprocally increasing endothelial VDV gene expression, mark the formation of immature neo-vasculature in two different pathological contexts: wound healing and increased tumor angiogenesis in response to blockade of the Dll4/Notch1 signaling pathway.

In an *in vivo* mouse skin-wounding assay, topical addition of recombinant VEGF (rVEGF) for 12 hours increased the expression of a majority of VDV genes at the skin wound site, while anti-VEGF treatment had the expected opposite effect (Fig. 2C). Consistent with the response seen in tumors, the effects of VEGF blockade and topical VEGF were most marked in the proxVDV gene candidates.

The consequences of inducing neo-vascularization through inhibition of the Dll4/Notch1 signaling pathway were assessed by treating MDA-MB-231 tumor bearing mice with anti-Dll4, anti-VEGF, or control anti-ragweed antibody for 48 hours. The Dll4/Notch1 pathway increases nonproductive angiogenesis partly via enhancement of VEGF signaling (Jakobsson et al. *Biochem Soc Trans.* 37: 1233-1236, 2009; Ridgway et al. *Nature.* 444: 1083-1087, 2006; Jakobsson et al. *Nat. Cell Biol.* 12(10): 943-953, 2010). Histological analysis showed that, as expected, anti-VEGF treatment induced vascular pruning of tumor vasculature, while tumors treated with anti-Dll4 showed increased MVD when compared to controls (Fig. 2D, left). The expression of VDV genes changed concordantly, decreasing

upon anti-VEGF treatment and increasing in response to blockade of Dll4 (Fig. 2D, right). Again, as seen in other models, the change in expression of the proxVDV gene candidates is more pronounced than of the other VDV genes.

Overall, these data suggest that most VDV genes are expressed in VEGF-driven neo-vasculature, and that their collective expression likely reflects VEGF biological activity as well as the relative abundance of VEGF-dependent vasculature.

Of note, the transcriptional effects of VEGF pathway activation have been previously investigated, in both *in vitro* and *in vivo* systems. However, the transcriptional consequences of rVEGF stimulation in endothelial cells *in vitro* does not provide sufficient information to determine if a gene is indeed a bona-fide VEGF *in vivo* target. Indeed, we have found under many conditions that rVEGF stimulation of HUVECs *in vitro* does not induce transcription of most of the proxVDV genes we have identified (Fig. 3). This discrepancy should not be surprising, considering the systemic role of the vasculature as an integrator of many stimuli, including blood flow and mechanical forces, and that VEGF responses are dose as well as context dependent.

On the other hand, some of the genes identified here (including ESM1) as well as genes that we did not identify as likely targets (including Pcdh17, EHD3, PRDM1, and THBD among others; see Fig. 3) are reliably upregulated by VEGF stimulation *in vitro*. Thus, the list of proxVDV genes presented here is unlikely to be comprehensive, and we are currently integrating *in vivo* and *in vitro* data-sets in an effort to identify additional *in vivo* VEGF targets.

Example 4. proxVDV genes are in vivo proximal markers of VEGF/VEGFR-2 downstream bioactivity

Data from *in situ* hybridization (ISH) indicated that Esm1 is highly expressed in a significant fraction of vessels from HM7 colon xenograft tumors, while being nearly undetectable in vessels from anti-VEGF treated tumors (Fig. 4A-4C). This is consistent with Esm1 being a *bona fide* proxVDV gene. However, because ESM1 is regulated by other stimuli besides VEGF (Scherpereel et al. *Crit. Care Med.* 34(2): 532-537, 2006) and since ESM1 may also be occasionally expressed in tumor cells, we wanted to validate additional proxVDV gene candidates as a collective and more specific mean to gauge, *in vivo*, VEGF downstream signaling bioactivity in response to FDA approved VEGF/VEGFR-2 inhibitors. Thus, in these experiments mice bearing established MDA-MB-231 tumors with treated with a control mAb, anti-VEGF mAb (as a surrogate for bevacizumab), sunitinib (Sutent®;

Escudier. *Expert Rev. Anticancer Ther.* 10(3): 305-317, 2010), a small-molecule TKI that targets VEGFR-2 among other RTKs, or with axitinib (Kindler et al. *Lancet Oncol.* 12(3): 256-262, 2011; Grunwald et al. *Onco. Targets Ther.* 5:111-117, 2012), a more specific VEGFR-2 inhibitor. Subsequently, tumors were collected at 8, 16, or 72 hours after treatment for analyses (Fig. 5A, upper panel). Consistent with prior observations, all three inhibitors induced a significant reduction of MVD in tumors collected 72 hours after treatment (Fig. 4D). At the gene expression level, we found that in all cases, the proxVDV genes showed a greater downregulation than that seen for the pan-vascular markers Cd31 and Plvap (Fig. 5A). Importantly, the expression of Vegfa and of the non-vascular markers, E-cad (epithelial) and Cd45 (hematopoietic), was not markedly affected by any of the VEGF pathway inhibitors tested (Fig. 4E), suggesting that the changes in proxVDV gene expression are likely “on-target” and endothelial specific. At 72 hours, sunitinib seems to be the strongest VEGF pathway inhibitor (Fig. 5A).

The dynamics of the transcriptional response differed between small molecule inhibitor and anti-VEGF antibody treatment. No significant changes in gene expression were induced by sunitinib or axitinib after 8 hours of treatment. Only at 16 hours post-treatment were the effects of the two SMIs on downregulating proxVDV gene expression apparent, and this effect increased at 72 hours after treatment (Fig. 5A). In contrast, downregulation of proxVDV (but not distVDV) genes by anti-VEGF mAb (intraperitoneal administration) is obvious at 8 hours after dosing, reaching its peak by 16 hours post treatment.

To confirm whether this proxVDV gene set was indeed regulated by VEGF signaling in tumor-associated endothelial cells (TAECs), we treated MDA-MB-231-GFP tumor-bearing animals with anti-VEGF or control antibody, and then isolated TAECs (Cd31+, Cd45-, GFP-) by fluorescence-activated cell sorting (FACS), and compared gene expression *ex vivo* among the two different treatment groups. We found that all pan-vascular markers and proxVDV genes tested were highly enriched on endothelial cells when compared to other sorted cell populations (Fig. 6). The proxVDV genes Prnd, Esm1, Nid2, Kcne3, Apj, Apln, and Mest were consistently downregulated in TAEC cells isolated from anti-VEGF treated animals relative to controls, while other VDV genes such as Cd31, Plvap, Ets-1 and Hlx were not (Fig. 5B). This data confirms that Prnd, Esm1, Nid2, Kcne3, Apln, Apj, and Mest are proximal and sensitive biomarkers of VEGF bioactivity and are candidate reporters for the direct inhibition of the pathway in tumor-associated endothelial cells *in vivo*.

Example 5. Vascular transcriptional responses to VEGF signaling blockade are conserved between mouse and human tumors

To test whether the responses to VEGF neutralization we identified in murine models were conserved in human tumor vasculature, we investigated the effect of VEGF blockade on VDV signature expression in tumor biopsies from bevacizumab (Avastin®) treated patients. Here, we took advantage of published micro-array data from matched pre- and 21-days-post-treatment paired biopsies from 19 inflammatory breast cancer patients treated with one dose of bevacizumab as a single agent (Wedam et al. *Journal of Clin. Oncology*. 24: 769-777, 2006). Although conventional bioinformatics analyses had failed to distinguish most specific vascular gene expression changes in response to bevacizumab treatment in these biopsies *a priori* (Yang et al. *Clin Cancer Res*. 14: 5893-5899, 2008), focused expression analyses of the human orthologues of the VDV gene set showed a clear down regulation of VDV transcripts in post-bevacizumab treatment clinical samples as compared with pre-treatment biopsies (Fig. 7). As in the murine tumor models, we observed that the expression of several proxVDV genes such as ESM1, NID2, PRND, KCNE3, and MEST decreases more markedly upon VEGF treatment. The VDV gene set we identified in preclinical models thus enables the detection of an evolutionary conserved vascular response to VEGF signaling inhibition in clinical tumor samples.

Example 6. Pre-treatment expression levels of VDV genes in human colorectal cancers correlate with clinical responses to bevacizumab

Our data indicate that human tumor vessels enriched in VDV genes is uniquely responsive to VEGF signaling inhibition. We next tested the hypothesis that the relative enrichment of these markers in pre-treatment tumor samples could actually predict responsiveness to anti-VEGF therapy.

Bevacizumab in combination with chemotherapy has previously been shown to increase progression free survival (PFS) and overall survival (OS) in patients with metastatic colorectal carcinoma (CRC) (Hurwitz et al. *N Engl J Med*. 350: 2335-2342, 2004). NO16966 was a first-line metastatic CRC trial where patients received oxaliplatin-based chemotherapy (XELOX or FOLFOX-4) in combination with either placebo or bevacizumab; the addition of bevacizumab to chemotherapy in this particular study significantly improved the primary endpoint of PFS, albeit the OS differences (a secondary endpoint) did not reach statistical significance (Saltz et al. *Journal of Clin. Oncology*. 26: 2013-2019, 2008). To test if VDV gene expression could correlate with clinical outcomes, we analyzed VDV gene expression

levels in available pre-treatment archival tumor tissue from 103 patients (biomarker evaluable subpopulation) that were enrolled in the XELOX-containing arms. Due to the limited quality and amount of RNA from these clinical samples, gene expression was analyzed in a previously designed and validated “angiogenesis” Fluidigm qRT-PCR chip (Fig. 8B) that included 4 different housekeeping genes (for gene expression normalization), VEGF (as a control), and 22 representative proximal and distal VDV genes comprising: (i) pan-vascular markers such as CD31, CD34, and VE-CADH as indicators of MVD; (ii) key VEGF pathway components and proxVDV genes, including VEGFR-1, VEGFR-2, VEGFR-3, EPHRINB2, NRPs, ESM1, NID2, COL4a2, and LAMA4a; and (iii) additional VDV genes that are components of endothelial signaling pathways and potential modulators of VDV biology, including DLL-4, NOTCH1, ALK1, and EGFL7 (see Fig. 9A for validation of this compacted 22 VDV gene signature).

Figure 8A shows the Kaplan-Meier analysis of OS and PFS in the 103 patients from the NOI6966 trial examined in this study. In these patients, the addition of bevacizumab to chemotherapy provided a statistically significant PFS (HR, 0.59; 95% CI, 0.37 to 0.93; $p=0.024$) and OS benefit (HR, 0.45; 95% CI, 0.23 to 0.85; $p=0.015$). Since preliminary experiments in colon cancer prevalence samples indicated that the expression of these 22 VDV genes is highly correlated (Fig. 8C), we decided to use the median expression level of either an individual gene or a median expression score for the 22 VDV gene subset as a basis to classify the CRC samples as either “high” or “low” in each case (Fig. 8B). We then tested the correlation between clinical outcomes and “high” or “low” pre-treatment gene expression. As previously reported, the stratification of the treatment cohorts by pre-treatment VEGF mRNA levels alone did not show differential effects on clinical outcomes in the bevacizumab treated patients (Fig. 9B). Also, bevacizumab-treated patients classified into “high” and “low” subsets by the expression levels of a single distVDV gene (CD31) did not show differences in PFS; although there was a trend in OS benefit (Fig. 9C). In either case, the interaction between VEGF or CD31 and treatment did not show any predictive effect. Notably, when the same patient population was stratified into “VDV high” vs. “VDV low” populations (Fig. 8D), the effect size and significance altered substantially as compared to the baseline analysis: in “VDV low” patients (dotted lines in Fig. 8D), the combination of bevacizumab and chemotherapy conferred modest gains as compared to chemotherapy alone in both PFS (HR, 0.88; 95% CI, 0.47 to 1.62; $p=0.67$) and OS (HR, 0.58; 95% CI, 0.25 to 1.33; $p=0.2$). In contrast, in “VDV high” patients (Fig. 8D, solid lines), the addition of

bevacizumab to chemotherapy vs chemotherapy alone provided a marked and significant PFS (HR, 0.36; 95% CI, 0.17 to 0.77; $p=0.0079$) and OS (HR, 0.31; 95% CI, 0.11 to 0.93; $p=0.036$) benefit. The interaction between treatment and marker status shows a significant predictive effect for PFS ($p=0.036$), but despite the observed improvement in relative risk in the “VDV high” patients, did not reach statistical significance for OS ($p=0.37$). Thus, in this relatively small 103-patient sample set, the higher expression of a collective VDV gene subset correlates specifically with the improved clinical outcome provided by bevacizumab addition to chemotherapy, and has a predictive effect with respect to PFS.

Example 7. Monitoring patient responsiveness or sensitivity to a VEGF antagonist

This example describes an assay to monitor whether a patient will be responsive or sensitive to a VEGF antagonist. A sample (*e.g.*, blood or tissue biopsy) is obtained, with informed consent, from one or more patients before treatment with a VEGF antagonist (*e.g.*, an anti-VEGF antibody). DNA and serum/plasma are isolated, according to well known procedures. The samples may be pooled or maintained as individual samples.

The expression of at least one gene set forth in Table 1 or 2 is assessed by measuring mRNA for the at least one gene or by detecting protein encoded by the at least one gene using an ELISA as described above, with the following substitutions: (1) human gene (*e.g.*, Nid2) standards for murine gene (*e.g.*, Nid2) standards; (2) biotinylated goat anti-human gene (*e.g.*, Nid2) polyclonal antibodies for biotinylated goat anti-mouse gene (*e.g.*, Nid2) polyclonal Ab; and (3) 10% FBS for 0.5% BSA. Patients whose samples exhibit at least a two-fold increase in expression of the at least one gene relative to a control as described herein are identified as patients responsive or sensitive to treatment with VEGF antagonists.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patents, patent applications, scientific references, and Genbank Accession Nos. cited herein are expressly incorporated by reference in their entirety for all purposes as if each patent, patent application, scientific reference, and Genbank Accession No. were specifically and individually incorporated by reference. Such patent applications specifically include United States Provisional Patent Application No. 61/586,660, filed on January 13, 2012, from which this application claims benefit.

WHAT IS CLAIMED IS:

1. A method of determining whether a patient is likely to respond to treatment with the VEGF antagonist, the method comprising:

5 (a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient prior to any administration of a VEGF antagonist to the patient; and

(b) comparing the expression level of the at least one gene to a reference expression level of the at least one gene,

10 wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist.

2. A method of optimizing therapeutic efficacy of a VEGF antagonist for a patient, the method comprising:

15 (a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient prior to any administration of a VEGF antagonist to the patient; and

(b) comparing the expression level of the at least one gene to a reference expression level of the at least one gene,

20 wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist.

25 3. The method of claim 1 or 2, wherein the patient is in a population of patients being tested for responsiveness to a VEGF antagonist and the reference level is the median level of expression of the at least one gene in the population of patients.

4. The method of claim 1 or 2, wherein the change in level of expression of the at least one gene in the patient sample is an increase relative to the reference level.

30 5. The method of claim 1 or 2, wherein the change in level of expression of the at least one gene in the patient sample is a decrease relative to the reference level.

6. The method of claim 1 or 2, wherein expression of the at least one gene in the biological sample obtained from the patient is detected by measuring mRNA.

5 7. The method of claim 1 or 2, wherein expression of the at least one gene in the biological sample obtained from the patient is detected by measuring plasma protein levels.

8. The method of claim 1 or 2, wherein the biological sample is tumor tissue.

10 9. The method of claim 1 or 2, further comprising detecting expression of at least a second gene set forth in Table 1 or 2 in the biological sample from the patient.

10. The method of claim 9, further comprising detecting expression of at least a third gene set forth in Table 1 or 2 in the biological sample from the patient.

15

11. The method of claim 10, further comprising detecting expression of at least a fourth gene set forth in Table 1 or 2 in the biological sample from the patient.

12. The method of claim 1 or 2, wherein the at least one gene is selected from the
20 group consisting of: Alk1, CD34, CD105, CD144, Col4a1, Col4a2, Dll4, EFNB2, EGFL7, ESM1, LAMA4, NG2, Nid2, Notch1, NRP1, NRP2, RGS5, Sema3f, TSP1, VEGFR1, VEGFR2, VEGFR3, and VIM.

13. The method of claim 1 or 2, wherein the VEGF antagonist is an anti-VEGF
25 antibody.

14. The method of claim 13, wherein the anti-VEGF antibody is bevacizumab.

15. The method of claim 1 or 2, wherein the patient has an angiogenic disorder.

30

16. The method of claim 15, wherein the patient has cancer selected from the group consisting of: colorectal cancer, breast cancer, lung cancer, glioblastoma, and combinations thereof.

17. The method of claim 1 or 2, further comprising (c) selecting a VEGF antagonist for treatment of said patient when a change in the level of expression of the at least one gene in the patient sample relative to the reference level is detected.

5

18. The method of claim 17, further comprising (d) administering a VEGF antagonist to the patient.

19. The method of claim 17 or 18, wherein said VEGF antagonist is an anti-VEGF antibody.

10

20. The method of claim 19, wherein the anti-VEGF antibody is bevacizumab.

21. A method for selecting a therapy for a particular patient in a population of patients being considered for therapy, the method comprising:

15

(a) detecting expression of at least one gene set forth in Table 1 or 2 in a biological sample obtained from the patient prior to any administration of a VEGF antagonist to the patient;

20

(b) comparing the expression level of the at least one gene to a reference expression level of the at least one gene,

wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient who is likely to respond to treatment with a VEGF antagonist, and

25

(c) selecting a VEGF antagonist as the therapy if the patient is identified as likely to respond to treatment with a VEGF antagonist; or

(d) selecting a therapy that is not a VEGF antagonist if the patient is not identified as likely to respond to treatment with a VEGF antagonist.

30

22. The method of claim 21, wherein the reference level is the median level of expression of the at least one gene in the population of patients.

23. The method of claim 21, wherein the change in level of expression of the at least one gene in the patient sample is an increase relative to the reference level.

24. The method of claim 21, wherein the change in level of expression of the at least one gene in the patient sample is a decrease relative to the reference level.

5 25. The method of claim 21, further comprising detecting expression of at least a second gene set forth in Table 1 or 2 in the biological sample from the patient.

26. The method of claim 25, further comprising detecting expression of at least a third gene set forth in Table 1 or 2 in the biological sample from the patient.

10 27. The method of claim 26, further comprising detecting expression of at least a fourth gene set forth in Table 1 or 2 in the biological sample from the patient.

28. The method of claim 21, wherein the therapy of (c) is an agent selected from
15 the group consisting of: an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, and combinations thereof.

29. The method of claim 21, further comprising:
(e) administering an effective amount of a VEGF antagonist to the patient if the
20 patient is identified as likely to respond to treatment with a VEGF antagonist.

30. The method of claim 29, wherein the VEGF antagonist is an anti-VEGF antibody.

25 31. The method of claim 30, wherein the anti-VEGF antibody is bevacizumab.

32. The method of claim 31, further comprising administering an effective amount of at least a second agent.

30 33. The method of claim 32, wherein the second agent is selected from the group consisting of: an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, and combinations thereof.

34. The method of claim 21, wherein the therapy of (d) is an agent selected from the group consisting of: an anti-neoplastic agent, a chemotherapeutic agent, a growth inhibitory agent, a cytotoxic agent, and combinations thereof.

5 35. A method of identifying a biomarker for determining responsiveness to a VEGF antagonist, the method comprising:

(a) detecting the expression of a candidate biomarker in a biological sample obtained from a patient prior to administration of a VEGF antagonist to the patient; and

10 (b) comparing the expression of the candidate biomarker to a reference expression level of the candidate biomarker, wherein a change in the level of expression of the candidate biomarker in the patient sample relative to the reference level identifies the candidate biomarker as a biomarker of a patient who will likely respond to treatment with a VEGF antagonist.

15 36. The method of claim 35, wherein the reference level is the median level of expression of the at least one gene in a population of patients being tested for the likelihood they will respond to a VEGF antagonist.

20 37. The method of claim 35, wherein the reference level is the expression level of the at least one gene in a sample previously obtained from the patient.

38. The method of claim 37, wherein the patient was previously treated with a VEGF antagonist and is currently experiencing metastasis.

25 39. The method of claim 35, wherein the VEGF antagonist is an anti-VEGF antibody.

40. The method of claim 39, wherein the anti-VEGF antibody is bevacizumab.

30 41. The method of claim 35, further comprising (c) selecting a candidate biomarker having a change in level of expression relative to the reference for use as a biomarker for determining responsiveness to VEGF antagonist treatment.

42. A method for diagnosing an angiogenic disorder in a patient, the method comprising the steps of:

(a) detecting the expression level of at least one gene set forth in Table 1 or 2 or a biomarker identified according to the method of claim 31 in a sample obtained from the patient prior to any administration of a VEGF antagonist to the patient; and

(b) comparing the expression level of the at least one gene or biomarker to a reference level of the at least one gene;

wherein a change in the level of expression of the at least one gene in the patient sample relative to the reference level identifies a patient having an angiogenic disorder.

43. The method of claim 42, further comprising (c) selecting a VEGF antagonist for treatment of said patient when a change in the level of expression of the at least one gene in the patient sample relative to the reference level is detected.

44. The method of claim 43, further comprising (d) administering a VEGF antagonist to the patient.

45. The method of any one of claims 1, 2, 21, 35, or 42, wherein the change in the level of expression of the at least one gene or biomarker in the patient sample relative to the reference level is determined by calculating a VDV signature score (VDV_i) for the patient sample according to the algorithm:

$$VDV_i = \frac{1}{\sqrt{n}} \sum_{g=1}^n Z_{g,i}$$

wherein $Z_{g=1,i}, Z_{g=2,i}, \dots, Z_{g=n,i}$ are standardized z-scores of expression values for each gene or biomarker g , from $g=1$ to $g=n$, of the sample i , and

wherein a VDV_i below a first defined threshold indicates a decrease relative to the reference level, and a VDV_i above a second defined threshold indicates an increase relative to the reference level.

46. The method of claim 45, wherein the expression values for each gene or biomarker g , from $g=1$ to $g=n$, are qRT-PCR values for each gene g , from $g=1$ to $g=n$.

47. The method of claim 45, wherein the first defined threshold is from -4 to -0.5 and the second defined threshold is from 0.5 to 4.

48. The method of claim 47, wherein the first defined threshold is from -4 to -1
5 and the second defined threshold is from 1 to 4.

49. The method of claim 48, wherein the first defined threshold is from -4 to -1.5 and the second defined threshold is from 1.5 to 4.

10 50. The method of claim 49, wherein the first defined threshold is from -4 to -2 and the second defined threshold is from 2 to 4.

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 1D




FIGURE 2A
MDA-MB-231 Human breast cancer model
(Subcutaneous, 24 Hours of Treatment)




FIGURE 2B
U87 Human glioblastoma model
(Intracranial, Long term Treatment)





FIGURE 2C

Wound Healing
(12 Hours of Treatment)



8/24

FIGURE 2D




FIGURE 3





FIGURE 4A



FIGURES 4B-4C




Figure 4B

Figure 4C

FIGURE 4D

MDA-MB-231 tumor sections (72 hours post-treatment)





FIGURE 4E




FIGURE 5A




FIGURE 5B




FIGURE 6

Specific Enrichment of VDV Markers in Tumor-Associated Endothelial Cells





FIGURE 7



Effect of bevacizumab on VDV expression in human tumor biopsies






FIGURE 8A

Effect of bev on PFS



	HR	95% CI	p-value
— XELOX			
- - - XELOX + bevacizumab	0.59	0.37 to 0.93	0.024

Effect of bev on OS



	HR	95% CI	p-value
— XELOX			
- - - XELOX + bev	0.45	0.23 to 0.85	0.015

FIGURE 8B





FIGURE 8C


VDV gene expression is highly correlated



Spearman's correlation coefficient


FIGURE 8D

Effect of bev & VDV gene expression on PFS



		HR	95% CI	p-value	
—	XELOX				
—	XELOX + bev	VDV High	0.36	0.17 to 0.77	0.0079
---	XELOX				
---	XELOX + bev	VDV Low	0.88	0.47 to 1.62	0.67

Effect of bev & VDV gene expression on OS



		HR	95% CI	p-value	
—	XELOX				
—	XELOX + bev	VDV High	0.31	0.11 to 0.93	0.036
---	XELOX				
---	XELOX + bev	VDV Low	0.58	0.25 to 1.33	0.2

FIGURE 9A




FIGURE 9B







FIGURE 9C



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/21306

A. CLASSIFICATION OF SUBJECT MATTER IPC(8) - A61K 39/00; C12Q 1/00; A61K 39/395 (2013.01) USPC - 424/133.1 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC(8) - A61K 39/00; C12Q 1/00; A61K 39/395 (2013.01) USPC - 424/133.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched IPC(8) - A61K 39/00; C12Q 1/00; A61K 39/395 (2013.01), USPC - 424/133.1, 435/4, 424/158.1, 435/7.96: keyword search, as below Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) PatBase, USPTO PubWest, Google Scholar -- Search Terms: Bais, Brauer, Schmidt, Singh, z-score, anti-VEGF, bevacizumab, VEGF, VEGFR, inhibitor, antagonist, antibody, gene, expression, threshold, diagnose, diagnosis, angiogenesis, angiogenic, disorder, disease, condition, marker, biomarker, abcc9, adam12, adamts2, adamts12, adamts1, serpine1, serpi		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 2011/0117083 A1 (BAIS et al.) 19 May 2011 (19.05.2011) entire document, especially abstract; Table 1; claim 2; para [0005]; [0006]; [0008]; [0009]; [0018]; [0016]; [0078]; [0079]; [0245]; [0149]; [0003]; [0007]; [0173]; [0260];	1-18, 21-41 ----- 42-44, (45-50)/(1,2,21,35,42)
Y	SPEERS et al., Novel Strategies for the Treatment of Estrogen Receptor-Negative Breast Cancer. Annual Summary; U.S. Army Medical Research and Materiel Command, April 2009, pg. 1-300 [online]. [Retrieved on 1 March 2013] Retrieved from the internet: <URL: http://www.dtic.mil/dtic/tr/fulltext/u2/a512945.pdf > on 1 March 2013. Entire document, especially pg. 278, para 2; pg 279, para 1; pg. 16, para 2; pg. 77, para 1 - pg. 78, para 2.	(45-50)/(1,2,21,35,42)
Y	US 2006/0204951 A1 (FOLKMAN et al.) 14 September 2006 (14.09.2006) para [0065]; [0068].	42-44,, (45-50)/42
A	CHEADLE et al., Analysis of Microarray Data Using Z Score Transformation. J Mol Diagnost, May 2003, Vol. 5, No. 2, pg. 73-81. Entire document.	1-18, 21-50
A	US 2011/0151468 A1 (WU et al.) 23 June 2011 (23.06.2011) entire document	1-18, 21-50
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/>		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 1 March 2013 (01.03.2013)		Date of mailing of the international search report 27 MAR 2013
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201		Authorized officer: Lee W. Young PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 13/21306

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 19, 20
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.