



(86) **Date de dépôt PCT/PCT Filing Date:** 2004/12/21
(87) **Date publication PCT/PCT Publication Date:** 2005/07/14
(45) **Date de délivrance/Issue Date:** 2015/06/23
(85) **Entrée phase nationale/National Entry:** 2006/06/28
(86) **N° demande PCT/PCT Application No.:** EP 2004/014555
(87) **N° publication PCT/PCT Publication No.:** 2005/063820
(30) **Priorité/Priority:** 2003/12/30 (US60/533,406)

(51) **Cl.Int./Int.Cl.** **C07K 14/54** (2006.01),
A61K 38/04 (2006.01), **C12N 15/62** (2006.01)
(72) **Inventeurs/Inventors:**
LAUDER, SCOTT, US;
GILLIES, STEPHEN D., US
(73) **Propriétaire/Owner:**
MERCK PATENT GESELLSCHAFT MIT
BESCHRAENKTER HAFTUNG, DE
(74) **Agent:** FETHERSTONHAUGH & CO.

(54) **Titre : PROTEINES DE FUSION DE IL-7**
(54) **Title: IL-7 FUSION PROTEINS**

(57) **Abrégé/Abstract:**

The invention relates to interleukin-7 (IL-7) fusion proteins, methods of their production and uses thereof. The fusion proteins comprise an immunoglobulin portion fused directly or indirectly to IL-7, which was modified at specific positions as compared to the wild-type IL-7 in order to improve biological and pharmaceutical properties. The proteins of the invention are particularly useful in treating disorders accompanied by immune deficiencies and particularly diseases which involve T-cell deficiencies.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 July 2005 (14.07.2005)

PCT

(10) International Publication Number
WO 2005/063820 A2

(51) International Patent Classification⁷: **C07K 19/00**,
C12N 15/12, 5/10, A61P 35/00, A61K 38/17 // C07K
14/54, 16/00

(21) International Application Number:
PCT/EP2004/014555

(22) International Filing Date:
21 December 2004 (21.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/533,406 30 December 2003 (30.12.2003) US

(71) Applicant (for all designated States except US): **MERCK
PATENT GMBH** [DE/DE]; Frankfurter Strasse 250,
64293 Darmstadt (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LAUDER, Scott**
[CA/US]; 44 Coolidge Farm Road, Boxborough, MA
01719 (US). **GILLIES, Stephen, D.** [US/US]; 47 Swan-
son Lane, Carlisle, MA 01741 (US).

(74) Common Representative: **MERCK PATENT GMBH**;
Frankfurter Strasse 250, 64293 Darmstadt (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: IL-7 FUSION PROTEINS

(57) Abstract: The invention relates to interleukin-7 (IL-7) fusion proteins, methods of their production and uses thereof. The fu-
sion proteins comprise an immunoglobulin portion fused directly or indirectly to IL-7, which was modified at specific positions as
compared to the wildtype IL-7 in order to improve biological and pharmaceutical properties. The proteins of the invention are partic-
ularly useful in treating disorders accompanied by immune deficiencies and particularly diseases which involve T-cell deficiencies.



WO 2005/063820 A2

IL-7 FUSION PROTEINS

FIELD OF THE INVENTION

[0001] The invention relates to interleukin-7 (IL-7) fusion proteins, methods of their
5 production and uses thereof. The fusion proteins comprise an immunoglobulin portion
fused directly or indirectly to IL-7, which was modified at specific positions as compared
to the wild-type IL-7 in order to improve biological and pharmaceutical properties. The
proteins of the invention are particularly useful in treating disorders accompanied by
immune deficiencies and particularly diseases which involve T-cell deficiencies.

10

BACKGROUND OF THE INVENTION

[0002] A variety of disorders and therapies involve a deficiency of immune cells. For
example, HIV infection results in a loss of CD4+ T-cells, while therapies such as
chemotherapy and radiation therapy generally result in a loss of a wide variety of blood
15 cells. Attempts have been made to provide specific protein drugs that can replenish
specific types of immune cells that are lost as a result of a disease or therapy. For
example, in cancer chemotherapy, erythropoietin is used to replenish red blood cells,
granulocyte colony-stimulating factor (G-CSF) is used to replenish neutrophils, and
granulocyte macrophage colony stimulating factor (GM-CSF) is used to replenish
20 granulocytes and macrophages. These protein drugs, although beneficial, have relatively
short serum half-lives such that immune cell replenishment is often insufficient.
Moreover, no specific treatment is currently in clinical use to specifically stimulate T or
B-cell development, even though loss of these cells as a result of disease or after certain
myeloablative treatments is known to be particularly deleterious to a patient's health.
25 Thus, there exists a need in the art to develop immune system stimulators and
restoratives, particularly of lymphocytes, that have extended serum half-lives.

SUMMARY OF THE INVENTION

[0003] The present invention is directed to interleukin-7 (IL-7) fusion proteins which
30 have improved biological properties compared to corresponding wild-type IL-7 proteins.
Moreover, the present invention is based, in part, on the finding that IL-7 fusion proteins
having particular structural features have improved biological properties compared to
wild-type recombinant IL-7.

[0004] Accordingly, in one aspect, the invention features a fusion protein including a first portion comprising an immunoglobulin (Ig) chain and a second portion comprising interleukin-7 (IL-7), wherein the IL-7 fusion protein has an increased biological activity, such as an extended-serum half-life or in promoting the survival or expansion of immune cells, as compared to wild-type IL-7.

[0005] In one embodiment, the invention features a fusion protein including a first portion including an Ig chain and a second portion including IL-7, wherein the amino acid residues at positions 70 and 91 of IL-7 are glycosylated and the amino acid residue at position 116 of IL-7 is non-glycosylated. Throughout this document, amino acid positions of IL-7 refer to the corresponding positions in the mature, human IL-7 sequence. In one embodiment, the amino acid residue at position 116 of IL-7 is asparagine. In another embodiment, the amino acid residue at position 116 of IL-7 is altered such that it does not serve as a glycosylation site. In one embodiment, the IL-7 moiety comprises disulfide bonds between Cys2 and Cys92, Cys34 and Cys129, and Cys47 and Cys141 of IL-7.

[0006] In another embodiment, the invention includes a fusion protein including a first portion including an Ig chain and a second portion including IL-7, wherein the IL-7 comprises disulfide bonding between Cys2 and Cys92, Cys34 and Cys129, and Cys47 and Cys141 of IL-7. In one embodiment, the amino acid residue at position 116 of IL-7 is non-glycosylated. In another embodiment, the amino acid residue at position 116 of IL-7 is asparagine or is altered such that it does not serve as a glycosylation site. In another embodiment, the amino acid residues at positions 70 and 91 of IL-7 are glycosylated.

[0007] The Ig chain is generally an intact antibody or portion thereof, such as an Fc region. The Ig chain of the IL-7 fusion protein can be derived from any known Ig isotype and can include at least a portion of one or more constant domains. For example, the constant domain can be selected from the group consisting of a CH1 region, a hinge region, a CH2 region, and a CH3 region. In one embodiment, the Ig moiety includes a hinge region, a CH2 region and a CH3 region. The Ig chain is optionally connected to the IL-7 portion by a linker.

[0008] Ig moieties of a single antibody isotype, such as IgG1 or IgG2, and hybrid Ig moieties are permitted in the present invention. For example, in one embodiment, the Ig moiety includes a hinge region derived from one isotype (*i.e.* IgG2) and a CH region from another isotype (*i.e.* IgG1). An Ig chain including an Fc portion of IgG1 can advantageously be modified to include the mutations Asn297Gln and Tyr296Ala.

26474-1011

3

Furthermore, an Ig chain including an Fc portion of IgG2 can be advantageously modified to include the mutations Asn297Gln and Phe296Ala.

[0009] The IL-7 portion of the IL-7 fusion protein described above may comprise the mature portion of the IL-7 portion. In one embodiment, the IL-7 portion
5 can further include a deletion, such as an internal deletion. In one example, IL-7 can include an eighteen amino acid deletion of amino acids 96 to 114 of SEQ ID NO:1.

[0010] In other embodiments, the invention includes purified nucleic acids encoding the IL-7 fusion proteins described above and cultured host cells including these nucleic acids.

10 **[0011]** In another aspect, the invention includes a method of preparing an IL-7 fusion protein including expressing in a host cell the nucleic acid described above and harvesting the fusion protein.

[0012] In another aspect, the invention includes a composition such as a pharmaceutical composition which includes the fusion protein described above.

15 **[0013]** In another aspect, the invention includes a method of treating a patient by administering Fc-IL-7 fusion proteins.

According to one aspect of the present invention, there is provided an IL-7 fusion protein having improved biological properties compared to wild-type mature human IL-7, comprising: a first portion comprising an Fc region from IgG1 or
20 IgG2 wherein the Fc region comprises a hinge region, a CH2 domain, and a CH3 domain; and a second portion comprising a polypeptide having an amino acid sequence comprising at least 98% sequence identity with wild-type mature human interleukin-7 (IL-7) wherein wild-type mature human IL-7 consists of amino acid residues 26-176 of SEQ ID NO:1, and wherein amino acid residues corresponding to
25 positions 70 and 91 of wild-type mature human IL-7 are glycosylated and the amino acid residue corresponding to position 116 of wild-type mature human IL-7 is non-glycosylated.

26474-1011

3a

According to another aspect of the present invention, there is provided a fusion protein comprising: a first portion comprising an Fc region from IgG1 or IgG2 wherein the Fc region comprises a hinge region, a CH2 domain, and a CH3 domain; and a second portion comprising wild-type mature human IL-7 consisting of amino acid residues 26-176 of SEQ ID NO:1 except that residues 121 to 138 of SEQ ID NO:1 are deleted in said second portion, and wherein the amino acid residues corresponding to positions 70 and 91 of wild-type mature human IL-7 are glycosylated and the amino acid residue corresponding to position 116 of wild-type mature human IL-7 is non-glycosylated.

10 BRIEF DESCRIPTION OF THE DRAWINGS

[0014] Figure 1 depicts the amino acid sequence of human IL-7 (SEQ ID NO:1). The signal sequence is shown in bold. Also depicted in bold and italics is a stretch of eighteen amino acids which can be deleted from the IL-7 sequence.

[0015] Figure 2 depicts the amino acid sequence of cow IL-7 (SEQ ID NO:2). The signal sequence is shown in bold.

[0016] Figure 3 depicts the amino acid sequence of sheep IL-7 (SEQ ID NO:3). The signal sequence is shown in bold.

[0017] Figure 4 depicts the amino acid sequence of mature human Fc γ 1-IL-7 (SEQ ID NO:4).

[0018] Figure 5 depicts the amino acid sequence of mature human Fc γ 2(h)(FN>AQ)-IL-7 (SEQ ID NO:5).

[0019] Figure 6 depicts the amino acid sequence of mature human Fc γ 1(linker1)-IL-7 (SEQ ID NO:6).

[0020] Figure 7 depicts the amino acid sequence of mature human Fc γ 1(YN>AQ)(linker2)-IL-7 (SEQ ID NO:7).

[0021] Figure 8 depicts the amino acid sequence of mature human Fcγ1(YN>AQ,d)(linker2)-IL-7 (SEQ ID NO:8).

[0022] Figure 9 depicts the nucleic acid sequence for the Fc region of human Fcγ1-IL-7 (SEQ ID NO:22).

5 [0023] Figure 10 depicts the nucleic acid sequence for the Fc region of human Fcγ1(YN>AQ)-IL-7 (SEQ ID NO:21).

[0024] Figure 11 is the nucleic acid sequence for the Fc region of human Fcγ2(h)-IL-7 (SEQ ID NO:20).

10 [0025] Figure 12 is the nucleic acid sequence for the Fc region of human Fcγ2(h)(FN>AQ)-IL-7 (SEQ ID NO:19)

[0026] Figure 13 is a graphical representation of the pharmacokinetic profile of recombinant human IL-7 (open squares) and the fusion protein Fcγ2(h)(FN>AQ)-IL-7 (open diamonds) of Example 7. The serum concentration of the administered IL-7 fusion proteins (in ng/ml) was measured over time (in hours).

15 [0027] Figure 14 is a graphical representation of B-cell reconstitution in irradiated, bone marrow transplanted mice treated with recombinant human IL-7 (open symbols), human Fc-IL-7 (filled symbols) or PBS (X). Proteins were administered every other day (squares) or once a week (triangles). The stippled line represents B-cell concentration in donor mice.

20 [0028] Figure 15 is a graphical representation of T-cell reconstitution in irradiated, bone marrow transplanted mice treated with recombinant human IL-7 (open symbols), human Fc-IL-7 (filled symbols) or PBS (X). Proteins were administered every other day (squares) or once a week (triangles). The stippled line represents T-cell concentration in donor mice.

25 [0029] Figure 16 is a dot plot representing lymphocyte populations of samples from the blood (top row) and spleen (bottom row) of irradiated, bone marrow transplanted mice treated with huFcγ2(h)(FN>AQ)-IL-7 (first two columns), and untreated controls (last column). The first column represents reconstituted endogenous lymphocytes (CD45.2+), and the second column represents reconstituted donor lymphocytes (CD45.1+). T-
30 Lymphocytes were detected as CD3 positive cells shown in the lower right quadrant. B-lymphocytes were detected as B220 positive cells, shown in the upper left quadrant.

DETAILED DESCRIPTION OF THE INVENTION

[0030] The invention provides IL-7 fusion proteins that have enhanced biological activity compared to wild-type IL-7 proteins. In particular, the invention provides IL-7 fusion

proteins that include an immunoglobulin (Ig) portion. These Ig-IL-7 fusion proteins have enhanced biological activity, such as extended serum half-life as compared to the wild type IL-7 proteins, which makes them suitable for use in the treatment of conditions accompanied by immune cell deficiencies such as lymphocyte deficiencies.

5 [0031] The invention is further based in part on the finding that IL-7 fusion proteins that have particular structural characteristics also have enhanced biological properties. While the amino acid sequence of mammalian IL-7 is well known, information about the structure of eukaryotically derived IL-7 proteins, including, for example, how the protein folds and the effects of its predicted N-linked glycosylation sites on its biological activity, 10 remain ill-defined. For example, human IL-7 protein has a cysteine at positions 2, 34, 47, 92, 129, and 141 of the mature protein and three potential N-linked glycosylation sites at positions asparagine (Asn)70, Asn91, and Asn116. However, the precise structure of IL-7 synthesized under eukaryotic conditions is unknown.

[0032] The present invention includes IL-7 fusion proteins having particular structural 15 forms and enhanced biological activity. For example, IL-7 fusion proteins having the disulfide bonding pattern of Cys2-Cys92, Cys34-Cys129 and Cys47-141 are more active *in vivo* than a wild-type recombinant IL-7 protein.

[0033] Moreover, the invention provides a form of an IL-7 fusion protein in which only two of the three potential N-linked glycosylation sites of IL-7 are glycosylated. 20 Specifically, Asn70 and Asn91 of the mature protein are glycosylated, while the predicted N-linked glycosylation site at IL-7 Asn116 is not glycosylated. Such an IL-7 fusion protein is more active *in vivo* than a wild-type recombinant IL-7.

[0034] The invention also includes IL-7 fusion proteins wherein the IL-7 moiety contains a deletion and which retain comparable activity compared to the corresponding 25 unmodified IL-7 fusion proteins. For example, the invention provides a form of Ig-IL-7 in which the IL-7 moiety contains an eighteen amino acid internal deletion corresponding to the sequence VKGRKPAALGGEAQPTKSL (SEQ ID NO:9).

Interleukin-7 Fusion Proteins

[0035] Typically, the IL-7 protein portion is fused to a carrier protein. In one 30 embodiment, the carrier protein is disposed towards the N-terminus of the fusion protein and the IL-7 protein is disposed towards the C-terminus. In another embodiment, the IL-7 fusion protein is disposed towards the N-terminus of the fusion protein and the carrier protein is disposed towards the C-terminus.

[0036] As used herein, the term “interleukin-7” or “IL-7” mean IL-7 polypeptides and derivatives and analogs thereof having substantial amino acid sequence identity to wild-type mature mammalian IL-7s and substantially equivalent biological activity, *e.g.*, in standard bioassays or assays of IL-7 receptor binding affinity. For example, IL-7 refers to an amino acid sequence of a recombinant or non-recombinant polypeptide having an amino acid sequence of: i) a native or naturally-occurring allelic variant of an IL-7 polypeptide, ii) a biologically active fragment of an IL-7 polypeptide, iii) a biologically active polypeptide analog of an IL-7 polypeptide, or iv) a biologically active variant of an IL-7 polypeptide. IL-7 polypeptides of the invention can be obtained from any species, *e.g.*, human, cow or sheep. IL-7 nucleic acid and amino acid sequences are well known in the art. For example, the human IL-7 amino acid sequence has a Genbank accession number of NM 000880 (SEQ ID NO:1) and is shown in Figure 1; the mouse IL-7 amino acid sequence has a Genbank accession number of NM 008371; the rat IL-7 amino acid sequence has a Genbank accession number of AF 367210; the cow IL-7 amino acid sequence has a Genbank accession number of NM 173924 (SEQ ID NO:2) and is shown in Figure 2; and the sheep IL-7 amino acid sequence has a Genbank accession number of U10089 (SEQ ID NO:3) and is shown in Figure 3. The signal sequence for each of the polypeptide species is shown in bold in each of the figures and is typically not included where the IL-7 portion is fused C-terminal to the carrier protein.

[0037] A “variant” of an IL-7 protein is defined as an amino acid sequence that is altered by one or more amino acids. The variant can have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, *e.g.*, replacement of leucine with isoleucine. More rarely, a variant can have “nonconservative” changes, *e.g.*, replacement of a glycine with a tryptophan. Similar minor variations can also include amino acid deletions or insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological activity can be found using computer programs well known in the art, for example software for molecular modeling or for producing alignments. The variant IL-7 proteins included within the invention include IL-7 proteins that retain IL-7 activity. IL-7 polypeptides which also include additions, substitutions or deletions are also included within the invention as long as the proteins retain substantially equivalent biological IL-7 activity. For example, truncations of IL-7 which retain comparable biological activity as the full length form of the IL-7 protein are included within the invention. The activity of the IL-7 protein can be measured using *in vitro* cellular proliferation assays such as

described in Example 6 below. The activity of IL-7 variants of the invention maintain biological activity of at least 10 %, 20%, 40%, 60%, but more preferably 80%, 90%, 95% and even more preferably 99% as compared to wild type IL-7.

[0038] Variant IL-7 proteins also include polypeptides that have at least about 70%, 75%, 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98%, 99%, or more sequence identity with wild-type IL-7. To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino acid or nucleic acid sequence). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions.times.100). The determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, *et al.* (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used.

[0039] Potential T-cell or B-cell epitopes in the IL-7 moiety can be removed or modified in the Fc-IL-7 fusion proteins of the invention. Exemplary deimmunized IL-7 moieties are disclosed in the U.S. Provisional Patent Application entitled "IL-7 Variants with Reduced Immunogenicity" (Attorney Docket No. LEX-035PR), which was filed in the United States Patent and Trademark Office on December 9, 2004.

30 Carrier Protein

[0040] The carrier protein can be any moiety covalently fused to the IL-7 protein. In one embodiment, the carrier protein is albumin, for example, human serum albumin. In another embodiment, the carrier protein is an immunoglobulin (Ig) moiety, such as an Ig heavy chain. The Ig chain may be derived from IgA, IgD, IgE, IgG, or IgM. According

to the invention, the Ig moiety may form an intact antibody and may direct the IL-7 fusion protein to specific target sites in the body. Fusion proteins making use of antibody targeting are known to those in the art. In another embodiment, the carrier Ig moiety further comprises an Ig light chain.

5 [0041] In one embodiment, the Ig moiety comprises an Fc region. As used herein, "Fc portion" encompasses domains derived from the constant region of an immunoglobulin, preferably a human immunoglobulin, including a fragment, analog, variant, mutant or derivative of the constant region. Suitable immunoglobulins include IgG1, IgG2, IgG3, IgG4, and other classes. The constant region of an immunoglobulin is defined as a
10 naturally-occurring or synthetically-produced polypeptide homologous to the immunoglobulin C-terminal region, and can include a CH1 domain, a hinge, a CH2 domain, a CH3 domain, or a CH4 domain, separately or in any combination. In the present invention, the Fc portion typically includes at least a CH2 domain. For example, the Fc portion can include hinge-CH2-CH3. Alternatively, the Fc portion can include all
15 or a portion of the hinge region, the CH2 domain and/or the CH3 domain and/or the CH4 domain.

[0042] The constant region of an immunoglobulin is responsible for many important antibody functions including Fc receptor (FcR) binding and complement fixation. There are five major classes of heavy chain constant region, classified as IgA, IgG, IgD, IgE,
20 and IgM. For example, IgG is separated into four subclasses: 1, 2, 3, and 4, also known as IgG1, IgG2, IgG3, and IgG4, respectively.

[0043] IgG molecules interact with multiple classes of cellular receptors including three classes of Fcγ receptors (FcγR) specific for the IgG class of antibody, namely FcγRI, FcγRII, and FcγRIII. The important sequences for the binding of IgG to the FcγR
25 receptors have been reported to be located in the CH2 and CH3 domains. The serum half-life of an antibody is influenced by the ability of that antibody to bind to an Fc receptor (FcR). Similarly, the serum half-life of immunoglobulin fusion proteins is also influenced by the ability to bind to such receptors (Gillies *et al.*, (1999) Cancer Res. 59:2159-66). Compared to those of IgG1, CH2 and CH3 domains of IgG2 and IgG4 have
30 biochemically undetectable or reduced binding affinity to Fc receptors. It has been reported that immunoglobulin fusion proteins containing CH2 and CH3 domains of IgG2 or IgG4 had longer serum half-lives compared to the corresponding fusion proteins containing CH2 and CH3 domains of IgG1 (U.S. Patent No. 5,541,087; Lo *et al.*, (1998) Protein Engineering, 11:495-500). Accordingly, in certain embodiments of the invention,

preferred CH2 and CH3 domains are derived from an antibody isotype with reduced receptor binding affinity and effector functions, such as, for example, IgG2 or IgG4. More preferred CH2 and CH3 domains are derived from IgG2.

5 [0044] The hinge region is normally located C-terminal to the CH1 domain of the heavy chain constant region. In the IgG isotypes, disulfide bonds typically occur within this hinge region, permitting the final tetrameric antibody molecule to form. This region is dominated by prolines, serines and threonines. When included in the present invention, the hinge region is typically at least homologous to the naturally-occurring immunoglobulin region that includes the cysteine residues to form disulfide bonds linking
10 the two Fc moieties. Representative sequences of hinge regions for human and mouse immunoglobulins are known in the art and can be found in Borrebaeck, ed., (1992) Antibody Engineering, A Practical Guide, W. H. Freeman and Co. Suitable hinge regions for the present invention can be derived from IgG1, IgG2, IgG3, IgG4, and other immunoglobulin classes.

15 [0045] The IgG1 hinge region has three cysteines, the second and third of which are involved in disulfide bonds between the two heavy chains of the immunoglobulin. These same two cysteines permit efficient and consistent disulfide bonding of an Fc portion. Therefore, a preferred hinge region of the present invention is derived from IgG1, more preferably from human IgG1, wherein the first cysteine is preferably mutated to another
20 amino acid, preferably serine.

[0046] The IgG2 isotype hinge region has four disulfide bonds that tend to promote oligomerization and possibly incorrect disulfide bonding during secretion in recombinant systems. A suitable hinge region can be derived from an IgG2 hinge; the first two cysteines are each preferably mutated to another amino acid.

25 [0047] The hinge region of IgG4 is known to form interchain disulfide bonds inefficiently. However, a suitable hinge region for the present invention can be derived from the IgG4 hinge region, preferably containing a mutation that enhances correct formation of disulfide bonds between heavy chain-derived moieties (Angal *et al.* (1993) Mol. Immunol., 30:105-8).

30 [0048] In accordance with the present invention, the Fc portion can contain CH2 and/or CH3 and/or CH4 domains and a hinge region that are derived from different antibody isotypes, *i.e.*, a hybrid Fc portion. For example, in one embodiment, the Fc portion contains CH2 and/or CH3 domains derived from IgG2 or IgG4 and a mutant hinge region derived from IgG1. As used in this application, Fcγ2(h) refers to an embodiment wherein

the hinge is derived from IgG1 and the remaining constant domains are from IgG2. Alternatively, a mutant hinge region from another IgG subclass is used in a hybrid Fc portion. For example, a mutant form of the IgG4 hinge that allows efficient disulfide bonding between the two heavy chains can be used. A mutant hinge can also be derived from an IgG2 hinge in which the first two cysteines are each mutated to another amino acid. Such hybrid Fc portions facilitate high-level expression and improve the correct assembly of the Fc-IL-7 fusion proteins. Assembly of such hybrid Fc portions is known in the art and has been described in U.S. Published Patent Application No. 2003-0044423. [0049] In some embodiments, the Fc portion contains amino acid modifications that generally extend the serum half-life of an Fc fusion protein. Such amino acid modifications include mutations substantially decreasing or eliminating Fc receptor binding or complement fixing activity. For example, the glycosylation site within the Fc portion of an immunoglobulin heavy chain can be removed. In IgG1, the glycosylation site is Asn297 within the amino acid sequence Gln-Tyr-Asn-Ser (SEQ ID NO:30). In other immunoglobulin isotypes, the glycosylation site corresponds to Asn297 of IgG1. For example, in IgG2 and IgG4, the glycosylation site is the asparagine within the amino acid sequence Gln-Phe-Asn-Ser (SEQ ID NO:29). Accordingly, a mutation of Asn297 of IgG1 removes the glycosylation site in an Fc portion derived from IgG1. In one embodiment, Asn297 is replaced with Gln. In other embodiments, the tyrosine within the amino acid sequence Gln-Tyr-Asn-Ser (SEQ ID NO:30) is further mutated to eliminate a potential non-self T-cell epitope resulting from asparagine mutation. As used herein, a T-cell epitope is a polypeptide sequence in a protein that interacts with or binds an MHC class II molecule. For example, the amino acid sequence Gln-Tyr-Asn-Ser (SEQ ID NO:30) within an IgG1 heavy chain can be replaced with a Gln-Ala-Gln-Ser (SEQ ID NO:28) amino acid sequence. Similarly, in IgG2 or IgG4, a mutation of asparagine within the amino acid sequence Gln-Phe-Asn-Ser (SEQ ID NO:29) removes the glycosylation site in an Fc portion derived from IgG2 or IgG4 heavy chain. In one embodiment, the asparagine is replaced with a glutamine. In other embodiments, the phenylalanine within the amino acid sequence Gln-Phe-Asn-Ser (SEQ ID NO:29) is further mutated to eliminate a potential non-self T-cell epitope resulting from asparagine mutation. For example, the amino acid sequence Gln-Phe-Asn-Ser (SEQ ID NO:29) within an IgG2 or IgG4 heavy chain can be replaced with a Gln-Ala-Gln-Ser (SEQ ID NO:28) amino acid sequence.

[0050] It has also been observed that alteration of amino acids near the junction of the Fc portion and the non-Fc portion can dramatically increase the serum half-life of the Fc fusion protein. (U.S. Published Patent Application No. 2002-0147311). Accordingly, the junction region of an Fc-IL-7 or IL-7-Fc fusion protein of the present invention can
5 contain alterations that, relative to the naturally-occurring sequences of an immunoglobulin heavy chain and IL-7, preferably lie within about 10 amino acids of the junction point. These amino acid changes can cause an increase in hydrophobicity by, for example, changing the C-terminal lysine of the Fc portion to a hydrophobic amino acid such as alanine or leucine. In yet another embodiment of the invention, the C-terminal
10 lysine and preceding glycine of the Fc portion is deleted.

[0051] In other embodiments, the Fc portion contains amino acid alterations of the Leu-Ser-Leu-Ser segment near the C-terminus of the Fc portion of an immunoglobulin heavy chain. The amino acid substitutions of the Leu-Ser-Leu-Ser (SEQ ID NO:27) segment eliminate potential junctional T-cell epitopes. In one embodiment, the Leu-Ser-Leu-Ser
15 (SEQ ID NO:27) amino acid sequence near the C-terminus of the Fc portion is replaced with an Ala-Thr-Ala-Thr (SEQ ID NO:26) amino acid sequence. In other embodiments, the amino acids within the Leu-Ser-Leu-Ser (SEQ ID NO:27) segment are replaced with other amino acids such as glycine or proline. Detailed methods of generating amino acid substitutions of the Leu-Ser-Leu-Ser (SEQ ID NO:27) segment near the C-terminus of an
20 IgG1, IgG2, IgG3, IgG4, or other immunoglobulin class molecules, as well as other exemplary modifications for altering junctional T-cell epitopes, have been described in U.S. Published Patent Application No. 2003-0166877.

Spacer

[0052] In one embodiment, a spacer or linker peptide is inserted between the carrier
25 protein and the IL-7 fusion protein. For example, the spacer is placed immediately C-terminal to the last amino acid of an Ig constant region. The spacer or linker peptide is preferably non-charged and more preferably non-polar or hydrophobic. The length of a spacer or linker peptide is preferably between 1 and about 100 amino acids, more preferably between 1 and about 50 amino acids, or between 1 and about 25 amino acids,
30 and even more preferably between 1 and about 15 amino acids, and even more preferably less than 10 amino acids. In one embodiment, the spacer contains a sequence $(G_4S)_n$, where n is less than 5. In a preferred embodiment, the spacer contains the sequence G_4SG_4 (SEQ ID NO:25). In yet another embodiment, the spacer contains a motif that is recognized as an N-linked glycosylation site. In yet another embodiment, the spacer

contains a motif that is recognized by a site specific cleavage agent. In an alternative embodiment of the invention, the carrier protein and IL-7 fusion protein are separated by a synthetic spacer, for example a PNA spacer, that is preferably non-charged, and more preferably non-polar or hydrophobic.

5 Production of IL-7 Fusion Proteins

[0053] Non-limiting methods for synthesizing useful embodiments of the invention are described in the Examples herein, as well as assays useful for testing the *in vitro* properties, and pharmacokinetic and *in vivo* activities in animal models.

[0054] The IL-7 fusion proteins of the invention can be produced using recombinant
10 expression vectors known in the art. The term "expression vector" refers to a replicable DNA construct used to express DNA which encodes the desired IL-7 fusion protein and which includes a transcriptional unit comprising an assembly of (1) genetic element(s) having a regulatory role in gene expression, for example, promoters, operators, or enhancers, operatively linked to (2) a DNA sequence encoding the desired IL-7 fusion
15 protein which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. The choice of promoter and other regulatory elements generally varies according to the intended host cell. A preferred expression vector of the invention is an Fc expression vector derived from the PdCs-huFc expression vector described in Lo *et al.*, Protein Engineering (1998) 11:495.

20 [0055] In a preferred example, the nucleic acid encoding the IL-7 fusion protein is transfected into a host cell using recombinant DNA techniques. In the context of the present invention, the foreign DNA includes a sequence encoding the inventive proteins. Suitable host cells include prokaryotic, yeast or higher eukaryotic cells. Preferred host cells are eukaryotic cells.

25 [0056] The recombinant IL-7 fusion proteins can be expressed in yeast hosts, preferably from *Saccharomyces* species, such as *S. cerevisiae*. Yeast of other genera such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of replication from a yeast plasmid or an autonomously replicating sequence (ARS), a promoter, DNA encoding the IL-7 fusion protein, sequences for polyadenylation and
30 transcription termination and a selection gene. Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-4-phosphate

isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase and glucokinase.

[0057] Various mammalian or insect cell culture systems can be employed to express recombinant protein. Baculovirus systems for production of proteins in insect cells are well known in the art. Examples of suitable mammalian host cell lines include NS/O cells, L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa, and BHK cell lines. Additional suitable mammalian host cells include CV-1 cells (ATCC CCL70) and COS-7 cells both derived from monkey kidney. Another suitable monkey kidney cell line, CV-1/EBNA, was derived by transfection of the CV-1 cell line with a gene encoding Epstein-Barr virus nuclear antigen-1 (EBNA-1) and with a vector containing CMV regulatory sequences (McMahan *et al.*, (1991) EMBO J. 10:2821). The EBNA-1 gene allows for episomal replication of expression vectors, such as HAV-EO or pDC406, that contain the EBV origin of replication.

[0058] Mammalian expression vectors may comprise non-transcribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a poly-adenylation site, splice donor and acceptor sites, and transcriptional termination sequences. Commonly used promoters and enhancers are derived from Polyoma, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence.

[0059] For secretion of the IL-7 fusion protein from the host cell, the expression vector comprises DNA encoding a signal or leader peptide. In the present invention, DNA encoding the native signal sequence of IL-7 can be used, or alternatively, a DNA encoding a heterologous signal sequence may be used, such as the signal sequence from another interleukin or from a secreted Ig molecule.

[0060] The present invention also provides a process for preparing the recombinant proteins of the present invention including culturing a host cell transformed with an expression vector comprising a DNA sequence that encodes the IL-7 fusion protein under conditions that promote expression. The desired protein is then purified from culture media or cell extracts. For example, supernatants from expression systems that secrete recombinant protein into the culture medium can be first concentrated using a

commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix, as known in the art. For example, Fc-IL-7 fusion proteins are conveniently captured using a matrix coupled to Protein A.

5 [0061] An "isolated" or "purified" IL-7 fusion protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the IL-7 fusion protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of IL-7 fusion protein in
10 which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations of IL-7 fusion protein having less than about 30% (by dry weight) of non-IL-7 fusion protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-IL-7 fusion
15 protein, still more preferably less than about 10% of non-IL-7 fusion protein, and most preferably less than about 5% non-IL-7 fusion protein. When the IL-7 fusion protein or biologically active portion thereof is purified from a recombinant source, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5%
20 of the volume of the protein preparation.

[0062] The term "substantially pure Ig-IL-7 fusion protein" refers to a preparation in which the Ig-IL-7 fusion protein constitutes at least 60%, 70%, 80%, 90%, 95% or 99% of the proteins in the preparation. In one embodiment, the invention includes substantially pure preparations of Ig-IL-7 fusion proteins having a disulfide bonding
25 pattern between Cys2 and Cys92, Cys34 and Cys129, and Cys47 and Cys141. In another embodiment, the invention features substantially pure preparations of Ig-IL-7 fusion proteins where Asn116 is non-glycosylated but Asn70 and Asn91 are glycosylated.

Methods of Treatment Using Fc-IL-7 Proteins

[0063] The IL-7 fusion proteins of the invention are useful in treating immune
30 deficiencies and in accelerating the natural reconstitution of the immune system that occurs, for example, after diseases or treatments that are immunosuppressive in nature. For example, IL-7 fusion proteins can be used to treat infectious pathogens, immune disorders, and to enhance the growth (including proliferation) of specific immune cell types. Moreover, the IL-7 fusion proteins can be used in the treatment of cancers such as

bladder cancer, lung cancer, brain cancer, breast cancer, skin cancer, and prostate cancer. In one example, it is useful to treat patients who have undergone one or more cycles of chemotherapy with IL-7 fusion proteins as described above to help their immune cells replenish. Alternatively, IL-7 fusion proteins are useful in adoptive T-cell
5 transplantations. For example, IL-7 fusion proteins may be administered to facilitate the expansion and survival of transplanted T-cells, or to expand isolated T-cell populations *ex vivo*. Alternatively, it is also useful to administer the IL-7 fusion proteins described above to patients with HIV, the elderly, patients receiving a transplant or other patients with suppressed immune system function.

10 Administration

[0064] The IL-7 fusion proteins of the invention can be incorporated into a pharmaceutical composition suitable for administration. Such compositions typically comprise the IL-7 fusion protein and a pharmaceutically-acceptable carrier. As used herein the language "pharmaceutically-acceptable carrier" is intended to include any and
15 all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art.

[0065] A pharmaceutical composition of the invention is formulated to be compatible
20 with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils,
25 polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or
30 sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0066] Medicaments that contain the IL-7 fusion proteins of the invention can have a concentration of 0.01 to 100% (w/w), though the amount varies according to the dosage form of the medicaments.

[0067] Administration dose depends on the body weight of the patients, the seriousness of the disease, and the doctor's opinion. However, it is generally advisable to administer about 0.01 to about 10 mg/kg body weight a day, preferably about 0.02 to about 2 mg/kg, and more preferably about 0.5mg/kg in case of injection. The dose can be administered
5 once or several times daily according to the seriousness of the disease and the doctor's opinion.

[0068] Compositions of the invention are useful when co-administered with one or more other therapeutic agents, for example, a molecule also known to be useful to replenish blood cells. For example, the molecule may be erythropoietin which is known to be used
10 to replenish red blood cells, G-CSF which is used to replenish neutrophils or GM-CSF which is used to replenish granulocytes and macrophages.

EXAMPLE 1.

Cloning of human (hu) Fc-IL-7 and huFc-IL-7 variants.

15 [0069] The nucleic acid encoding the mature form of human IL-7 (*i.e.* lacking its N-terminal signal sequence) is amplified by Polymerase Chain Reaction (PCR), using forward and reverse primers that incorporated the restriction sites for Sma I and Xho I, respectively. The amplified PCR product is cloned into a pCRII vector (Invitrogen, Carlsbad, CA), and its sequence verified. The amino acid sequence of mature IL-7 is
20 shown as SEQ ID NO:1. The Sma I/Xho I digested IL-7 fragment is transferred into a likewise treated pdCs-huFc derived expression vector, resulting in a chimeric sequence between huFc and IL-7, with IL-7 placed in-frame, directly downstream of the sequence encoding a CH3 moiety of Fc (see Lo *et al.*, Protein Engineering (1998) 11:495).

[0070] A series of expression vectors are derived from the pdCs-huFc vector which
25 encodes an Fc fragment that generally includes a hinge, a CH2 domain, and a CH3 domain of Ig, and which have been engineered to incorporate specific alterations in the Fc region. Thus, by shuttling the IL-7 fragment between these vectors, a series of huFc-IL-7 fusion proteins were generated which differ in their Fc backbone. In order to create the various backbones, the appropriate mutations can first be introduced into the Fc sequence
30 by methods known in the art. Since the Fc region of the pdCs-huFc derived vector is flanked by an AflII restriction site and a SmaI restriction site, by subjecting the nucleic acid of the appropriately modified backbone to PCR using primers that incorporate the restriction sites for AflII and SmaI respectively, the resultant Fc encoding nucleic acid fragment can then be substituted into the pdCs-huFc derived vector as an AflII to SmaI

fragment. The AflIII sequence CTTAAGC (SEQ ID NO:24) is upstream of the Fc sequence beginning GAGCCCAA (SEQ ID NO:23), which represents the beginning of the hinge region as shown in Figure 20. The SmaI site CCCGGGT (SEQ ID NO:17) is towards the end of the CH3 region as shown by the underlined nucleic acids in Figure 12 and encodes the Pro-Gly amino acids proceeding the alanine residue from the lysine to alanine mutation at the end of the CH3 region.

[0071] For example, huFcγ1-IL-7 is constructed having the hinge region, the CH2 and the CH3 domains derived from the IgG 1 subclass. In the context of an Fc fusion protein, the IgGγ1 hinge region in addition contains a mutation substituting the first cysteine for serine. The sequence of the encoded fusion protein is depicted in Figure 4 (SEQ ID NO:4), while the sequence in SEQ ID NO:22 encodes the mature huFcγ1 backbone of the vector.

[0072] In addition, Fcγ1-IL-7 fusion proteins were generated that included the dipeptide mutation YN to AQ to eliminate the glycosylation site on Fc (corresponding to N297 in IgGγ1) as well as a potential immunogenic T-cell epitope, according to the methods described above. The mature Fc backbone sequence for huFcγ1(YN>AQ) is disclosed in SEQ ID NO:21. The substitution of alanine and glycine in place of tyrosine and asparagine was accomplished by first introducing mutations into the Fc backbone by an overlap PCR approach. Two overlapping complementary mutagenic primers were used to generate two PCR fragments, which were used as the template in a second round of amplification to produce a single fragment containing the appropriate codon substitutions. The mutagenic primer in the sense direction was 5'-AGCAGGCCCAGAGCACGTACCGTGTGGT-3' (mutation underlined) (SEQ ID NO:36). The complementary strand was 5'-GTACGTGCTCTGGGCCTGCTCCTCCCGC-3' (SEQ ID NO:37). The flanking forward primer was 5'-CTCTCTGCAGAGCCCAA TCT-3' (SEQ ID NO:38), which also contains a PstI site. In the antisense direction, the flanking reverse primer was 5'CAGGGTGTACACCTGTGGTTC-3' (SEQ ID NO:33), which also contains a BsrGI site. After amplification, the sequence was verified through standard methods and was subject to restriction by BsrGI and PstI. The resultant fragment was then substituted for the non-mutant fragment of the Fc region.

[0073] The huFcγ2(h)(FN>AQ)-IL-7 was also constructed using the techniques previously described. This fusion protein includes an altered hinge region which was derived from IgGγ1 subclass, while the CH2 and CH3 domains were derived from IgGγ2

subclass. Additionally, the dipeptide mutation FN to AQ was included to eliminate the glycosylation site on Fc (corresponding to N297 in IgG_{Y1}) as well as a potential immunogenic T-cell epitope. The sequence of the encoded fusion protein is depicted in Figure 5 (SEQ ID NO:5). The sequence of the mature Fc backbone huFc_{Y2}(h)(FN>AQ) is shown in SEQ ID NO:19).

[0074] In addition, Fc-IL-7 fusion proteins were generated which included a flexible linker sequence between the Fc moiety and the IL-7 moiety. For example, a linker polypeptide with the sequence GGGGSGGGGSGGGGS (linker1, SEQ ID NO:34) was inserted. To generate huFc_{Y1}(linker1)-IL-7, a synthetic oligonucleotide duplex of the sequence 5' - G GGT GCA GGG GGC GGG GGC AGC GGG GGC GGA GGA TCC GGC GGG GGC TC- 3' (SEQ ID NO:18) was inserted by blunt-end ligation at the unique SmaI site of the expression vector pdCs-huFc-IL-7 and the orientation of the duplex was verified. The forward primer was designed such that the amino acid residues Pro -Gly encoded by the codons spanning the SmaI site (C CCG GGT) (SEQ ID NO:17), and the ensuing Ala residue (resulting from the encoded lysine to alanine substitution) of the CH3 region, were maintained. The amino acid sequence of the encoded fusion protein is shown in Figure 6 (SEQ ID NO:6).

[0075] Additional Fc-IL-7 fusion proteins were constructed that included a shorter linker polypeptide with the sequence GGGGSGGGG (linker2, SEQ ID NO:25). To generate huFc_{Y1}(YN>AQ)(linker2)-IL-7, an amplified PCR product, obtained from the primer pair 5'- CCCGGGCGCCGGCGGTGGAGGATCAGGTGGTGGCGGTGAT TGTGA TATTGAAGGTAAAGATG-3' (containing the encoded linker sequence, SEQ ID NO:15) and 5'- ATCATGTCTGGATCCCTCGA-3' (SEQ ID NO:14) on an appropriate pdCs-Fc-IL-7 template plasmid, was cloned into a pCRII vector (Invitrogen, Carlsbad, CA), and its sequence verified. A Xma I/Xho I digested fragment encoding linker2/IL-7 was then transferred into a likewise treated pdCs-huFc derived expression vector. The vector was modified to contain the mature Fc backbone huFc_{Y1}(YN>AQ) of SEQ ID NO:21. The amino acid sequence of the encoded fusion protein is shown in Figure 7 (SEQ ID NO:7).

[0076] Similarly, huFc_{Y1}(YN>AQ,d)(linker2)-IL-7 was generated, using the primer pair 5'-CCCGGGCGGTGGAGGATCAGGTGGTGGCGGTGATTGTGATAT TGAAGGTAAAGATG-3' (SEQ ID NO:16) and 5'-ATCATGTCTGGATCCCTCGA-3' (SEQ ID NO:12). huFc_{Y1}(YN>AQ,d)(linker2)-IL-7 differs from the preceding fusion

protein huFcγ1(YN>AQ)(linker2)-IL-7 in that it lacks the terminal two amino acid residues of the Fc portion of the fusion protein. Specifically, rather than terminating in with the sequence ...ATATPGA (SEQ ID NO:11), the Fc portion ends with the sequence ATATP (SEQ ID NO:10). The amino acid sequence of the encoded fusion protein is
5 shown in Figure 8 (SEQ ID NO:8).

EXAMPLE 2.

Transfection and Expression of Fc-IL-7 Fusion Proteins.

[0077] Electroporation was used to introduce the DNA encoding the IL-7 fusion proteins
10 described above into a mouse myeloma NS/0 cell line. To perform electroporation NS/0 cells were grown in Dulbecco's modified Eagle's medium supplemented with 10% heat-inactivated fetal bovine serum, 2 mM glutamine and penicillin/streptomycin. About 5×10^6 cells were washed once with PBS and resuspended in 0.5 ml PBS. 10 μg of linearized plasmid DNA for huFcγ1-IL-7 was then incubated with the cells in a Gene
15 Pulser Cuvette (0.4 cm electrode gap, BioRad) on ice for 10 min. Electroporation was performed using a Gene Pulser (BioRad, Hercules, CA) with settings at 0.25 V and 500 μF. Cells were allowed to recover for 10 min on ice, after which they were resuspended in growth medium and plated onto two 96 well plates.

[0078] Stably transfected clones were selected by their growth in the presence of 100 nM
20 methotrexate (MTX), which was added to the growth medium two days post-transfection. The cells were fed every 3 days for two to three more times, and MTX-resistant clones appeared in 2 to 3 weeks. Supernatants from clones were assayed by anti-Fc ELISA to identify clones that produced high amounts of the IL-7 fusion proteins. High producing clones were isolated and propagated in growth medium containing 100 nM MTX.
25 Typically, a serum-free growth medium, such as H-SFM or CD medium (Life Technologies), was used.

EXAMPLE 3.

Biochemical analysis of huFc-IL-7 fusion proteins

30 [0079] Routine SDS-PAGE characterization was used to assess the integrity of the fusion proteins. Differences between the huFc-IL-7 variants huFcγ1-IL-7, huFcγ2(h)(FN>AQ)-IL-7, huFcγ1(linker1)-IL-7, huFcγ1(YN>AQ)(linker2)-IL-7 and huFcγ1(YN>AQ,d)-IL-7 were investigated. The huFc-IL-7 fusion proteins, expressed from NS/0 cells, were

captured on Protein A Sepharose beads (Repligen, Needham, MA) from the tissue culture medium into which they were secreted, and were eluted by boiling in protein sample buffer, with or without a reducing agent such as β -mercaptoethanol. The samples were separated by SDS-PAGE and the protein bands were visualized by Coomassie staining. By SDS-PAGE the tested huFc-IL-7 fusion proteins were generally well expressed, as they were present substantially as a single band on the gel; it was found that in samples of huFc-IL-7 variants that included a linker, secondary bands, which may represent clipped material, were noticeably reduced.

[0080] Purified huFc-IL-7 fusion proteins were also analyzed by size exclusion chromatography (SEC) to assess the extent to which the huFc-IL-7 variants were aggregated. Briefly, the cell culture supernatant was loaded onto a pre-equilibrated Fast-Flow Protein A Sepharose column, the column was washed extensively in a physiological buffer (such as 100 mM Sodium Phosphate, 150 mM NaCl at neutral pH), and the bound protein was eluted at about pH 2.5 to 3 in same salt buffer as above. Fractions were immediately neutralized.

[0081] It was found that for each of the fusion proteins tested at least 50% of the product was monomeric, and generally more than 65%. "Monomeric," as used herein, refers to non-aggregated proteins. It is understood that proteins with an Fc portion normally form a disulfide-bonded complex which normally include two polypeptide chains (unless the two Fc portions are present within the same polypeptide) and may be thought of as a "unit-dimer". "Monomeric" is not intended to exclude such disulfide-bonded species, but only to connote that the proteins are non-aggregated. To obtain a virtually monomeric huFc-IL-7 fusion protein preparation (around 98%), the eluate from a Sepharose-ProteinA purification was loaded onto a preparative SEC column (Superdex) and the monomeric peak fraction was collected. Typically, the concentration of the recovered protein was around 1mg/ml. If required, the sample was concentrated, for example by spin dialysis (*e.g.* VivaSpin) with a molecular weight cut-off of 10–30 kDa.

Disulfide Bonding

[0082] IL-7 contains six Cys residues which could make disulfide bonds, at positions Cys2, Cys34, Cys47, Cys92, Cys129 and Cys141 of the mature IL-7 protein sequence. The folding of huFcyl-IL-7 was assessed by determining the pattern of disulfide bonds present in the IL-7 moiety of the fusion protein. Briefly, peptide maps of huFcyl-IL-7 were generated from trypsinized material and analyzed for the presence of signature peptide fragments. huFcyl-IL-7 protein was trypsinized either in a native form or after

reduction and alkylation. To account for peptide fragments that may be glycosylated, samples of the native and denatured proteins were additionally treated with PNGaseF to remove glycosyl chains prior to tryptic digestion. Peptide fragments were fractionated by HPLC, and their mass was determined by mass spectroscopy.

5 [0083] In the context of Fcγ1-IL-7, a peptide fragment containing the disulfide bond Cys47-Cys141 ("3-6") would be predicted to have a mass of 1447.6, whereas a peptide fragment containing the disulfide bond Cys2-Cys141 ("1-6") would be predicted to have a mass of 1426.6. Similarly, a peptide fragment containing the disulfide bond Cys34-Cys129 ("2-5") would be predicted to have a mass of 2738.3. Indeed, peptide fragments
10 of a mass of 1447.6 ("3-6") and of 2738.3 ("2-5") were identified in samples derived from the native Fc-IL-7 protein regardless of whether the samples were treated with PNGaseF or not, but not in samples from reduced Fc-IL-7. Conversely, the peptide fragment of a mass of 1426.6 ("1-6") was not found in any sample. Thus, Fcγ1-IL-7 contained the disulfide bonds Cys47-Cys141 and Cys34-Cys129, but not Cys2-Cys141. It was noted
15 that a peptide fragment of the predicted mass of 2439.2, corresponding to a fragment containing Cys2-Cys92 ("1-4") was identified only in the sample from the native fusion protein treated with PNGaseF. In fact, Cys92 lies within the tripeptide motif Asn91Cys92Thr93, indicating that Asn91 was glycosylated in huFcγ1-IL-7. Thus, in huFcγ1-IL-7 the disulfide bonding pattern was consistent with Cys2-Cys92, Cys34-Cys129, and Cys47-Cys141. This experimentally determined configuration of disulfide
20 bonds of Fc-IL-7 stands in contrast to the experimentally determined configuration reported for bacterially produced and re-folded IL-7 (Cosenza *et al.* (1997) JBC 272:32995).

N-linked Glycosylation Sites

25 [0084] Human IL-7 contains three potential glycosylation sites, at positions Asn70, Asn91 and Asn116 of the mature IL-7 protein sequence. Peptide maps of huFcγ1-IL-7 (reduced/alkylated) were analyzed for the presence of signature peptide fragments. If glycosylated, these signature fragments would only be revealed in samples treated with PNGaseF. Masses of 1489.7, 1719.9 and 718.3 would be predicted for tryptic peptide
30 fragments containing the unmodified residues for Asn70, Asn91, and Asn116, respectively.

[0085] Indeed, peptide fragments of a mass of 1489.7 and of 1719.9 were identified in samples that had been treated with PNGaseF, but were absent in the untreated sample,

indicating that Asn70 (contained in the sequence ...MNSTG...) (SEQ ID NO:31), and Asn91 (contained in the sequence ...LNCTG...) (SEQ ID NO:32), were indeed glycosylated. Surprisingly, a tryptic fragment of a mass of 718.3, corresponding to SLEENK (SEQ ID NO:35), was identified in both the PNGaseF treated sample and the untreated sample, indicating that Asn116 was not glycosylated. This was not expected because Asn116 in the human IL-7 sequence ...PTKSLEENKSLKE ... (SEQ ID NO:13) (see SEQ ID NO:1) is predicted to be an N-linked glycosylation site. The NKS putative glycosylation site is conserved in sheep and cows as well as humans.

[0086] The analysis of disulfide bonding patterns and N-linked glycosylation sites, repeated with samples of Fcγ1-(linker1)-IL-7 and of Fcγ2h(FN>AQ)-IL-7, gives similar results.

EXAMPLE 4.

ELISA Procedures.

[0087] The concentrations of protein products in the supernatants of MTX-resistant clones and other test samples was determined by anti-huFc ELISA, as described in detail below. ELISA plates were coated with AffiniPure Goat anti-Human IgG (H+L) (Jackson Immuno Research Laboratories, West Grove, PA) at 5 µg/mL in PBS, and 100 µL/well in 96-well plates. Coated plates were covered and incubated at 4°C overnight. Plates were washed 4 times with 0.05% Tween (Tween 20) in PBS, and blocked with 1% BSA/1% goat serum in PBS, 200 µL/well. After incubation with the blocking buffer at 37°C for 2 hrs, the plates were washed 4 times with 0.05% Tween in PBS and tapped dry. Test samples were diluted as appropriate in sample buffer (1% BSA/1% goat serum/0.05% Tween in PBS). A standard curve was prepared using a chimeric antibody (with a human Fc) of known concentration. To prepare a standard curve, serial dilutions were made in the sample buffer to give a standard curve ranging from 125 ng/mL to 3.9 ng/mL. The diluted samples and standards were added to the plate, 100 µL/well and the plate incubated at 37°C for 2 hr. After incubation, the plate was washed 8 times with 0.05% Tween in PBS. To each well 100 µL of the secondary antibody horseradish peroxidase-conjugated anti-human IgG was added, diluted to around 1:120,000 in sample buffer. The exact dilution of the secondary antibody was determined for each lot of the HRP-conjugated anti-human IgG. After incubation at 37°C for 2 hr, the plate was washed 8 times with 0.05% Tween in PBS.

[0088] The substrate solution was added to the plate at 100 μ L /well. This solution was prepared by dissolving 30 mg of OPD (o-phenylenediamine dihydrochloride (OPD), (1 tablet) into 15 mL of 0.025 M citric acid/0.05 M Na_2HPO_4 buffer, pH 5, which contained 0.03% of freshly added hydrogen peroxide. The color was allowed to develop for about 30 minutes at room temperature in the dark. The reaction was stopped by adding 4N sulfuric acid, 100 μ L/well. The plate was read by a plate reader, which was set at both 490 and 650 nm and programmed to subtract the background OD at 650 nm from the OD at 490 nm.

[0089] The concentration of human IL-7 in serum samples of animals treated with huFc-IL-7 fusion proteins or recombinant human IL-7 was determined by ELISA, essentially as described above. Human IL-7 was captured via a mouse anti-human IL-7 antibody (R&D Systems, Minneapolis, MN) and detected with a goat anti-human IL-7 biotin antibody (R&D Systems, Minneapolis, MN).

EXAMPLE 5.

Purification of huFc-IL-7 proteins.

[0090] A standard purification of Fc-containing fusion proteins was performed based on the affinity of the Fc protein moiety for Protein A. Briefly, NS/0 cells expressing the appropriate fusion protein, were grown in tissue culture medium and the supernatant containing the expressed protein was collected and loaded onto a pre-equilibrated Past Flow Protein A Sepharose column. The column was then washed extensively with buffer (such as 100 mM Sodium Phosphate, 150 mM NaCl at neutral pH). Bound protein was eluted at a low pH (pH 2.5 - 3) in same buffer as above and fractions were immediately neutralized.

[0091] To obtain a non-aggregated huFc-IL-7 fusion protein preparation (around 98% monomer), the eluate was loaded onto a preparative SEC column (Superdex) and the monomeric peak fraction was collected. Typically, the concentration of the recovered protein was around 0.5 mg/ml to 2 mg/ml, and where appropriate the sample was concentrated by spin dialysis (e.g. Viva Spin with a molecular weight cut-off of 30 kDa).

EXAMPLE 6.

In vitro activity of huFc-IL-7 proteins.

[0092] The cytokine activity of the purified huFc-IL-7 fusion proteins was determined *in vitro* in a cellular proliferation bioassay. Human PBMC (Peripheral Blood Mononuclear

Cells) were activated by PHA-P to produce cells which were responsive to IL-7. Proliferation was measured in a standard thymidine incorporation assay. Briefly, PBMC's were first incubated for five days with 10 microgram/ml PHA-P, cells were washed and then incubated in a medium supplemented with the huFc-IL-7 fusion proteins, in a dilution series, for a total of 48 hours. During the final 12 hours, the samples were pulsed with 0.3 μ Ci of [methyl-3H]thymidine (Dupont-NEN-027). Cells were then washed extensively, harvested and lysed onto glass filters. 3H-thymidine incorporated into DNA was measured in a scintillation counter. As a standard, wild type huIL-7 protein, obtained from R&D Systems (Minneapolis, MN), or obtained from the National Institute for Biological Standards and Control (NIBSC), was assayed.

[0093] An ED50 value of cell proliferation for huFc-IL-7 fusion proteins was obtained from plotting a dose response curve according to standard techniques, and determining the protein concentration that resulted in half-maximal response. The fusion proteins huFc γ 1-IL-7, huFc γ 2(h)(FN>AQ)-IL-7, and huFc γ 1(linker1)-IL-7 were evaluated. The ED50 values of the fusion proteins were fairly similar to one another, falling within a 3-fold range from one another. Therefore, it was found that these alterations in the Fc moiety have little influence on IL-7 activity of the fusion protein.

[0094] In addition, it was found that the ED50 values of these fusion proteins were about 3- to 10-fold higher than the ED50 value obtained for huIL-7 commercially available from R&D Systems. Since this commercial preparation is produced in bacteria and is not glycosylated, enzymatically deglycosylated huFc γ 1-IL-7 protein, by treatment with PNGaseF, was evaluated. It was found to have similar activity to the untreated form. Without wishing to be bound by theory, the somewhat decreased activity of the fusion proteins may have been due not to glycosylation of the IL-7 moiety but instead to a steric effect resulting from a constrained N-terminus of the IL-7 moiety.

EXAMPLE 7.

Pharmacokinetics of huFc-IL-7 proteins.

[0095] The pharmacokinetic (PK) profiles of an huFc-IL-7 fusion protein and of recombinant human IL-7 (Peprotech, Rocky Hill, NJ) were evaluated, and the results are depicted in Figure 13. A single subcutaneous injection of equimolar amounts of huFc γ 2(h)(FN>AQ)-IL-7 or of recombinant human IL-7 (50 micrograms) was administered to groups of C57BL6/J mice. Blood samples were obtained by retro-orbital

bleeding at injection (*i.e.*, at $t = 0$ min), and at 30 min, 1 hr, 2 hrs, 4 hrs, 8 hrs, 24, 48, 72, 96, 120 and 144 hrs post-injection. Samples were collected in heparin-tubes to prevent clotting, and cells were removed by centrifugation in a high-speed Eppendorf microcentrifuge for 4 min at 12,500 g. PK values were calculated with the PK solutions 2.0TM software package (Summit Research Services, Montrose, CO).

[0096] The concentration of administered IL-7 was determined in quadruplicate plasma samples at each time point by an ELISA specific for human IL-7. It was found that the pharmacokinetic behavior of the huFc-IL-7 and recombinant IL-7 differed dramatically. For recombinant human IL-7, the maximum concentration (C_{max}) was 23.5 ng/ml at 2.0 hours post injection (T_{max}), whereas for huFc-IL-7 C_{max} was 1588.7 ng/ml 24 hours post injection. In addition, while recombinant human IL-7 was absorbed more rapidly than huFc-IL-7 (β -phase half-life 0.9 hours vs. 12.4 hours), huFc-IL-7 was eliminated approximately 9-fold more slowly from circulation during the β -phase. Thus, in terms of AUC (area under the curve) as a measure of total drug exposure, mice receiving huFc-IL-7 had a 572-fold higher exposure to the administered protein than mice receiving recombinant human IL-7. These data demonstrate a significant improvement of huFc-IL-7 fusion proteins relative to free recombinant human IL-7 with regards to their PK. It was further found that the PK profiles of huFc-IL-7 fusion proteins, such as huFc γ 1-IL-7 and huFc γ 2(h)(FN>AQ)-IL-7, huFc γ 1(YN>AQ)(linker2)-IL-7, and huFc γ 1(YN>AQ,d)(linker2)-IL-7, which were administered to the mice by intravenous injection, were similar to one another.

EXAMPLE 8.

Efficacy of huFc-IL-7 in lymphopenic mice after bone marrow (BM) transplantation.

[0097] The efficacy of huFc-IL-7 fusion proteins compared to recombinant human IL-7 was evaluated *in vivo*. For example, huFc γ 2(h)(FN>AQ)-IL-7 or recombinant human IL-7 (Peprotech, Rocky Hill, New Jersey) was administered to lymphopenic mice after transplantation of T-cell-depleted bone marrow (BM), and the recovery of immune cell populations was assessed.

[0098] Essentially, recipient mice were lethally irradiated prior to BM transplantation with two doses of 600 cGy total body irradiation at a 4 hr interval, and BM cells re-suspended in PBS were infused into tail veins of recipient mice. At regular intervals from Day 5 onwards, an equimolar amount of huFc-IL-7 (7 μ g) or recombinant human IL-7

(2.5 µg) (Peprotech, Rocky Hill, NJ) was administered subcutaneously to the recipient mice. Over the course of the experiment, recipient mouse blood samples were taken, and lymphocyte cell concentrations in the samples were measured.

[0099] For BM cell transplantations, BM cells were obtained aseptically from femurs and tibias of BL/6.SJL (H2^b, CD45.1) mice (Jackson Labs, Bar Harbor, ME) and depleted of T-cells by removing magnetically labeled T-cells over MACS[®] columns (Miltenyi Biotec, Auburn, CA). The degree of T-cell depletion was monitored by FACS analysis with fluorescently labeled antibodies against CD45, αβ-TCR (T-cells) and 7-Amino Actinomycin D (7-AAD, apoptotic cells) (Calbiochem, X). 10 x 10⁶ live (7-AAD-negative) BM cells (containing less than 1 % T-cells) were used per recipient mouse. In congenic BM transplantations, B6 (H2^b, CD45.2) mice were used as the recipient mouse strain, and in allogeneic BM transplantations B6C3F1 (H2^{b/k}, CD45.2) mice were chosen.

[00100] Lymphocyte cell concentrations (as presented in Table 1) were measured essentially as described by Brocklebank and Sparrow (Brocklebank and Sparrow (2001) Cytometry 46:254). Briefly, fluorescent beads (TruCOUNT[™] Tubes, BD Biosciences, San Jose, CA) were dissolved in 40 µl of PBS containing a mixture of lymphocyte-specific antibodies. Subsequently, 10 µl of anti-coagulated blood was added, mixed and incubated for 30 minutes in the dark at room temperature. Red blood cells were lysed in 450 µl of Red Blood Cell lysis solution (BD Biosciences, San Jose, CA) and samples were analyzed by flow cytometry (BD FACSCalibur[™], BD Biosciences, San Jose, CA). The concentration of a particular lymphocyte population (*e.g.* B-cells, T-cells or total leukocytes) was determined by creating separate gates around lymphocytes and fluorescent beads and reading the number of events within each gate. The number of gated lymphocytes per microliter was calculated by dividing the number of events in a gated lymphocyte region by the number of events in the gated bead region. This number was multiplied by the fraction of the number of beads per TruCOUNT[™] tube (provided by the supplier) over the sample volume and finally multiplied by the sample dilution factor.

[00101] In one experiment, lymphocyte reconstitution was assessed in a congenic BM transplantation setting, using materials and methods specified above. Recipient mice were injected with huFcγ2(h)(FN>AQ)-IL-7 at a dose of 7 µg (125 µg IL-7/kg body weight), and lymphocyte cells were measured as described. Donor lymphocytes were detected as CD45.1 positive cells, whereas endogenous lymphocytes of the recipient mice

were detected as CD45.2 positive cells. Lymphocyte B-cells and T-cells were identified using B220 and CD3 lymphocyte markers respectively. It was found that by Day 49, donor lymphocytes (CD45.1 positive cells) had repopulated the recipient mice to levels comparable to non-irradiated control mice, while endogenous lymphocytes (CD45.2 positive cells) had not significantly expanded. In addition, the treatment with the huFc-IL-7 fusion protein did not cause significant toxicity. These results demonstrated the efficacy of the fusion protein in expanding adoptively transferred lymphocyte populations. These results are depicted in Figure 16.

[00102] In another experiment, an allogeneic BM transplantation model, which may better simulate a clinical transplantation setting, was used to compare a huFc-IL-7 fusion protein to recombinant human IL-7, and the results are shown in Table 1. Again, methods and materials described above were used. huFc γ 2(h)(FN>AQ)-IL-7 and human IL-7 (equivalent of 125 μ g IL-7/kg body weight) were administered either every other day (q2d) or once a week (q7d) from Day 5 to Day 56 after transplantation. PBS-treated donor mice and irradiated, bone marrow recipient mice treated with PBS served as controls.

[00103] In recipient mice treated with the fusion protein, donor-derived B-cells (CD45.1⁺, B220⁺, CD19⁺) reached baseline levels (as defined by the blood concentration of B-cells in donor control mice) 14 or 16 days after transplantation, when given q2d or q7d, respectively. In contrast, in recipient mice treated with recombinant human IL-7, neither dosing regimen had an effect; B-cell numbers in PBS-treated and human IL-7-treated recipient mice required about the same time to reach baseline levels, about 28 days. In addition to accelerated B-cell reconstitution, huFc-IL-7 treatment promoted the continuous expansion of B-cells until Day 33: huFc-IL-7 administered q2d resulted in a 7-fold increase, whereas administered q7d resulted in a 2.5-fold increase in B-cell numbers compared to control mice. After Day 33, B-cell numbers declined, but were still approximately 2-fold higher than in control mice. Also, after Day 33, levels of administered IL-7 proteins declined in the blood, which partially may be due to the formation of neutralizing antibodies to the human fusion protein. Figure 14 represents these results of B-cell reconstitution in irradiated, bone marrow transplanted mice treated with recombinant human IL-7 and huFC-IL-7.

[00104] A similar result was observed regarding donor-derived T-cells (CD45.1⁺, CD3⁺, TCR α β ⁺). Treatment with the huFc-IL-7 fusion protein resulted in accelerated T-cell reconstitution, whereas treatment with recombinant human IL-7 did not. Maximum

T-cell levels were reached around Day 49. However, T-cell numbers above baseline (*i.e.*, blood concentration of T-cells in donor mice) were only achieved with a q2d dosing schedule of the huFc-IL-7 fusion protein, reaching about 1.5 fold the number of T-cells in the control mice. Figure 15 represents these results of T-cell reconstitution in irradiated bone marrow transplanted mice treated with recombinant human IL-7 and huFc-IL-7.

[00105] Despite the transiently high numbers of donor B-cells and T-cells in the recipient mice under certain conditions, none of the experimental mice showed any signs of morbidity during the course of the experiment. Analysis of internal organs at day 55 did not reveal pathological abnormalities in liver, kidney, lung, spleen, thymus, lymph nodes, stomach, small intestine and colon. Thus, this allogeneic transplantation experiment demonstrated that the huFc-IL-7 fusion protein was significantly superior *in vivo* over recombinant human IL-7 in reconstituting lymphocytes after myeloablative conditioning.

Table 1: Effect of huFc-IL-7 fusion protein on immune cell reconstitution.

			Number of cells per microliter of blood								
			Day								
			7	13	19	27	33	41	47	55	
I. Donor Derived Leukocytes (CD45.1 ⁺) a.) Allogeneic BMT into B6C3 mice	PBS	AVG	580.8	3576.8	4384.0	16805.6	20732.8	16395.2	21462.4	20329.6	
		SEM	324.5	1717.0	1474.2	8367.6	10065.3	7997.3	11230.2	9693.2	
	huFc-IL-7, q2d	AVG	248.8	26597.0	130253.6	148048.0	194266.4	168794.0	127848.0	79175.2	
		SEM	89.3	4786.2	25106.8	18786.3	17131.1	25949.9	981.0	10740.2	
	huFc-IL-7, q7d	AVG	278.4	5848.8	32308.0	64256.0	74103.2	69042.4	55989.6	58519.2	
		SEM	106.3	762.9	2786.3	6081.4	8345.9	14497.4	5989.8	6583.8	
	human IL-7, q7d	AVG	270.7	6020.0	6922.7	24278.7	37353.3	42166.7	33570.7	38138.7	
		SEM	84.9	1229.2	1089.7	5235.3	2697.0	4691.4	764.5	3273.6	
	human IL-7, q7d	AVG	229.0	4603.0	4963.0	27443.0	27027.0	27797.0	32675.0	33205.0	
		SEM	31.9	435.6	654.1	2513.8	1813.4	4084.6	6307.8	4894.7	
	b.) B6.SJL Control mice	PBS	AVG	30744.5							
			SEM	3018.6							
			Number of cells per microliter of blood								
			Day								
			7	13	19	27	33	41	47	55	
II. Donor Derived B-cells (CD45.1 ⁺ CD19 ⁺ B220 ⁺) a.) Allogeneic BMT into B6C3 mice	PBS	AVG	32.0	1013.3	4257.3	19294.7	22549.3	17842.7	23841.3	22296.0	
		SEM	5.7	25.7	661.1	2858.5	2625.4	1691.6	3783.7	1474.5	
	huFc-IL-7, q2d	AVG	16.0	17830.0	122122.4	131862.4	169745.6	142972.0	98571.0	58661.6	
		SEM	5.7	3755.4	24028.4	16974.5	15413.6	22834.6	2599.1	9352.9	

	huFc-IL-7, q7d	AVG	14.4	2559.0	28124.0	53573.6	59999.2	52932.8	41489.6	42899.2		
		SEM	6.5	408.9	2464.1	4911.5	6337.2	10998.2	4821.1	5013.0		
		human IL-7, q7d	AVG	5.3	1550.7	4316.0	17802.7	28061.3	30177.3	24720.0	26104.0	
			SEM	3.5	403.5	542.7	3689.5	1484.5	2861.6	633.8	2085.3	
		human IL-7, q7d	AVG	4.0	748.0	3179.0	19303.0	19203.0	17930.0	22460.0	21473.0	
			SEM	1.6	56.1	443.3	1666.1	1429.5	2511.6	4458.6	3199.5	
	b.) B6.SJL Control mice	PBS	AVG	15572.6								
			SEM	1631.4								
		Treatment	Number of cells per microliter of blood									
			Day									
	III. Donor Derived T-cells (CD45.1*CD3*abTCR*) a.) Allogeneic BMT into B6C3 mice	PBS	AVG	7	13	19	27	33	41	47	55	
0.0				206.7	192.0	1808.0	3928.0	4453.3	6178.7	5622.7		
huFc-IL-7, q2d		SEM	0.0	121.6	46.2	245.0	546.7	533.0	538.1	210.0		
		AVG	0.0	124.0	839.0	5260.8	12252.8	15204.8	16628.0	13365.6		
huFc-IL-7, q7d		SEM	0.0	24.9	155.5	1183.1	1734.8	1641.2	1524.5	1423.9		
		AVG	0.0	38.4	228.0	2428.0	5492.0	9988.0	9252.8	8598.4		
human IL-7, q7d		SEM	0.0	6.1	38.5	326.2	918.1	2475.8	2748.6	1185.9		
		AVG	0.0	56.0	144.0	1057.3	3068.0	5368.0	5081.3	7112.0		
human IL-7, q7d		SEM	0.0	3.3	42.5	160.1	154.0	645.5	113.7	409.9		
		AVG	0.0	41.0	88.0	1493.0	2579.0	3918.0	5021.0	6043.0		
		SEM	0.0	2.5	20.2	187.5	238.0	694.6	724.2	1013.1		
		b.) B6.SJL Control mice	PBS	AVG	10832.0							
SEM				1500.4								

EXAMPLE 9.Efficacy of huFc-IL-7 on T-cell transplantations in lymphopenic mice.

- 5 [00106] The efficacy of huFc-IL-7 fusion proteins was also evaluated in a T-cell transplantation model. In essence, a homogeneous (clonal) population of T-cells was transferred into immunodeficient, irradiated mice, the recipient mice were administered the huFc-IL-7 fusion protein, and the degree of T-cell reconstitution and, eventually, T-cell function, was assessed.
- 10 [00107] To obtain a homogeneous population of T-cells, splenocytes were taken from P14 TCR-tg/RAG mice (Charles River Laboratories, Wilmington, MA), which are devoid of B-cells. In addition, all T-cells of these mice express the transgenic T-cell receptor (TCR), P14, which is specific for a viral epitope (gp33 of LCMV).

[00108] Single cell suspensions of splenocytes were injected intravenously into the tails of RAG Cy+ immunodeficient mice (Charles River Laboratories) that had been irradiated once with 650 Rads (sub-lethal dose) 4 hrs prior to transplantation. On alternate days starting at Day 2, recipient mice were administered 7 µg of the fusion protein huFcγ2(h)(FN>AQ)-IL-7. A control group of recipient mice were administered PBS. The degree of T-cell reconstitution in response to the huFc-IL-7 fusion protein or PBS was determined by measuring the presence of P14 T-cells (CD8⁺Vβ8.1⁺Vα2⁺ cells) in the blood by flow cytometry.

[00109] It was found that at Day 35, mice that were administered the huFc-IL-7 fusion protein had a 17-fold increase in T-cell numbers (35,000 cells/µl) compared to control mice (2,000 cells/µl). Indeed, the levels of T-cell reconstitution exceeded those seen in untreated P14 TCR mice (23,000 cells/µl). In addition, in these huFc-IL-7-treated mice a significant fraction of reconstituted T-cells had up-regulated the IL-2Rα receptor subunit, CD25, on the cell surface. Thus, not only was the huFc-IL-7 fusion protein useful in expanding transplanted T-cells, but in addition may have preconditioned the transferred T-cells to be responsive to cytokines, such as IL-2.

EXAMPLE 10.

huFc-IL-7 adjuvant therapy for immunocompromised patients.

[00110] Numerous clinical settings are envisaged in which patients may benefit from huFc-IL-7 adjuvant therapy. For example, new treatment modalities are being developed for pediatric patients with malignant disease such as lymphoblastic or myeloid leukemias, who, following a myeloablative therapy, are treated by allogeneic hematopoietic stem cell transplantation to reconstitute the immune system.

[00111] To increase markedly the potential donor pool for these patients, it has been found that G-CSF mobilized peripheral blood stem cells (PBSCs) from matched unrelated donors or haplo-identical donors with 1 – 3 HLA loci mismatches may be a source of cells, provided that the transplant is depleted of T-cells (see Handgretinger *et al.* (2001) Annals NY Acad. Sciences 938:340-357). This depletion drastically reduced the occurrence of acute graft-to-host transplant rejection (GvHD); however, it is believed that because of the low concentration of T-cells, there was a significant delay in immunoreconstitution. Patients were at high risk of viral infections for at least the first 6 month post transplant, and T-cells did not return to normal levels for a year

(Handgretinger *et al.*, (2001) Annals NY Acad. Sciences 938:340-357; Lang *et al.*, (2003) Blood 101:1630-6). Therefore, it would be advantageous to increase the rate of repopulation of T-cells and of other immune cells in these patients.

[00112] Patients that will benefit huFc-IL-7 therapy include patients with a
5 childhood leukemia, such as a lymphoblastic leukemia or a myeloid leukemia. Children having this disorder will first undergo a myeloablative conditioning therapy which may be based either on the chemotherapeutic agent busulfan or total body irradiation combined with chemotherapy. For instance, according to the patient's diagnosis and age, the patient is treated with total body irradiation (typically 6 treatments of 2 Gy each),
10 rabbit anti-thymocyte globulin (10 mg/kg daily for 3 days), etoposide (40 mg/kg) and cyclophosphamide (120 mg/kg).

[00113] To obtain, CD34 positive (pos) stem cells for transplant, peripheral blood stem cells (PBSCs) of a histocompatible (allogeneic) donor are mobilized with a daily dose of 10 micrograms/kg of G-CSF for 6 days, and are harvested by leukapheresis on
15 days 5 and 6. Generally, about 20×10^6 /kg CD34 pos stem cells are obtained and transplanted. CD34 pos stem cells are purified from the PBSCs by positive selection with an antiCD34 antibody in a SuperMACS system (Magnetic activated cell sorting, Miltenyi Biotec) and eluted. T-cell depletion is typically around 5 logs, to about 10×10^3 cells/kg. Aggregates and other debris are excluded from the graft by FACS sorting. The cell
20 suspension is infused into the patient via a central venous catheter. Optionally, the graft may include purified populations of other immune cells, such as haploidentical NK cells, DCs, monocytes, as well as CD34neg stem cells.

[00114] To assess engraftment, an absolute neutrophil count is performed. Engraftment is considered successful once neutrophil levels remain above 50
25 cells/microliter. Reconstitution of immune cells is monitored by FACS analysis, weekly at first and, once T-cell recovery begins, every 3 months.

[00115] To augment immunoreconstitution, the patient is treated with a huFc-IL-7 fusion protein such as huFc γ 2h(FN>AQ)-IL-7 or huFc γ 1(YN>AQ)(linker2)-IL-7. Approximately 3 weeks after transplant (or after engraftment is established), the patient
30 receives a subcutaneous administration of huFc γ 2h(FN>AQ)-IL-7 or huFc γ 1(YN>AQ)(linker2)-IL-7 at a dose of about 1.5 mg/m^2 (or a dose in the range of 0.15 mg/m^2 to 15 mg/m^2), about 2 times a week for 6 months – 12 months, until T-cell counts reach 50% of normal levels. It is found that the prognosis of the patient is improved due to lowered risk of viral infection, one of the main post-transplant

complications. It is further found that this treatment does not significantly increase the risk of acute GvHD.

[00116] In addition to administration of huFc-IL-7 protein, other medications are optimally given prophylactically. These include, for example, acyclovir, metronidazole, 5 flucanazole and co-trimoxazole. For the first three months, the patient may receive weekly administration of immunoglobulins, as well as of G-CSF.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 26474-1011 Seq 25-Jul-14 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110>	Lauder, Scott Gillies, Stephen
<120>	IL-7 Fusion Proteins
<130>	26474-1011
<140>	CA 2551915
<141>	2006-06-28
<150>	US 60/533,406
<151>	2003-12-30
<160>	38
<170>	PatentIn version 3.3
<210>	1
<211>	177
<212>	PRT
<213>	Homo sapiens

complications. It is further found that this treatment does not significantly increase the risk of acute GvHD.

[00116] In addition to administration of huFc-IL-7 protein, other medications are optimally given prophylactically. These include, for example, acyclovir, metronidazole, 5 flucanazole and co-trimoxazole. For the first three months, the patient may receive weekly administration of immunoglobulins, as well as of G-CSF.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 26474-1011 Seq 25-Jul-14 v1.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

The sequences in the sequence listing in electronic form are reproduced in the following table.

SEQUENCE TABLE

<110>	Lauder, Scott Gillies, Stephen
<120>	IL-7 Fusion Proteins
<130>	26474-1011
<140>	CA 2551915
<141>	2006-06-28
<150>	US 60/533,406
<151>	2003-12-30
<160>	38
<170>	PatentIn version 3.3
<210>	1
<211>	177
<212>	PRT
<213>	Homo sapiens

32a

<400> 1

Met	Phe	His	Val	Ser	Phe	Arg	Tyr	Ile	Phe	Gly	Leu	Pro	Pro	Leu	Ile
1				5					10					15	
Leu	Val	Leu	Leu	Pro	Val	Ala	Ser	Ser	Asp	Cys	Asp	Ile	Glu	Gly	Lys
			20					25					30		
Asp	Gly	Lys	Gln	Tyr	Glu	Ser	Val	Leu	Met	Val	Ser	Ile	Asp	Gln	Leu
		35					40					45			
Leu	Asp	Ser	Met	Lys	Glu	Ile	Gly	Ser	Asn	Cys	Leu	Asn	Asn	Glu	Phe
	50					55					60				
Asn	Phe	Phe	Lys	Arg	His	Ile	Cys	Asp	Ala	Asn	Lys	Glu	Gly	Met	Phe
65					70					75					80
Leu	Phe	Arg	Ala	Ala	Arg	Lys	Leu	Arg	Gln	Phe	Leu	Lys	Met	Asn	Ser
				85					90					95	
Thr	Gly	Asp	Phe	Asp	Leu	His	Leu	Leu	Lys	Val	Ser	Glu	Gly	Thr	Thr
			100					105					110		
Ile	Leu	Leu	Asn	Cys	Thr	Gly	Gln	Val	Lys	Gly	Arg	Lys	Pro	Ala	Ala
		115					120					125			
Leu	Gly	Glu	Ala	Gln	Pro	Thr	Lys	Ser	Leu	Glu	Glu	Asn	Lys	Ser	Leu
	130					135					140				
Lys	Glu	Gln	Lys	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Leu	Lys	Arg	Leu	Leu
145					150					155					160
Gln	Glu	Ile	Lys	Thr	Cys	Trp	Asn	Lys	Ile	Leu	Met	Gly	Thr	Lys	Glu
				165					170					175	

His

<210> 2

<211> 176

<212> PRT

<213> Bos taurus

<400> 2

Met	Phe	His	Val	Ser	Phe	Arg	Tyr	Ile	Phe	Gly	Ile	Pro	Pro	Leu	Ile
1				5					10					15	
Leu	Val	Leu	Leu	Pro	Val	Ala	Ser	Ser	Asp	Cys	Asp	Ile	Ser	Gly	Arg
			20					25					30		
Asp	Gly	Gly	Ala	Tyr	Gln	Asn	Val	Leu	Met	Val	Asn	Ile	Asp	Asp	Leu
		35					40					45			
Asp	Asn	Met	Ile	Asn	Phe	Asp	Ser	Asn	Cys	Leu	Asn	Asn	Glu	Pro	Asn
	50					55					60				
Phe	Phe	Lys	Lys	His	Ser	Cys	Asp	Asp	Asn	Lys	Glu	Ala	Ser	Phe	Leu
65					70					75					80
Asn	Arg	Ala	Ser	Arg	Lys	Leu	Arg	Gln	Phe	Leu	Lys	Met	Asn	Ile	Ser
				85					90					95	
Asp	Asp	Phe	Lys	Leu	His	Leu	Ser	Thr	Val	Ser	Gln	Gly	Thr	Leu	Thr
			100					105					110		
Leu	Leu	Asn	Cys	Thr	Ser	Lys	Gly	Lys	Gly	Arg	Lys	Pro	Pro	Ser	Leu
		115					120					125			
Ser	Glu	Ala	Gln	Pro	Thr	Lys	Asn	Leu	Glu	Glu	Asn	Lys	Ser	Ser	Arg
	130					135					140				
Glu	Gln	Lys	Lys	Gln	Asn	Asp	Leu	Cys	Phe	Leu	Lys	Ile	Leu	Leu	Gln
145					150					155					160
Lys	Ile	Lys	Thr	Cys	Trp	Asn	Lys	Ile	Leu	Arg	Gly	Ile	Lys	Glu	His
				165					170					175	

32b

<210> 3
 <211> 176
 <212> PRT
 <213> Ovis aries

<400> 3
 Met Phe His Val Ser Phe Arg Tyr Ile Phe Gly Ile Pro Pro Leu Ile
 1 5 10 15
 Leu Val Leu Leu Pro Val Ala Ser Ser Asp Cys Asp Phe Ser Gly Lys
 20 25 30
 Asp Gly Gly Ala Tyr Gln Asn Val Leu Met Val Ser Ile Asp Asp Leu
 35 40 45
 Asp Asn Met Ile Asn Phe Asp Ser Asn Cys Leu Asn Asn Glu Pro Asn
 50 55 60
 Phe Phe Lys Lys His Ser Cys Asp Asp Asn Lys Glu Ala Ser Phe Leu
 65 70 75 80
 Asn Arg Ala Ala Arg Lys Leu Lys Gln Phe Leu Lys Met Asn Ile Ser
 85 90 95
 Asp Asp Phe Lys Leu His Leu Ser Thr Val Ser Gln Gly Thr Leu Thr
 100 105 110
 Leu Leu Asn Cys Thr Ser Lys Gly Lys Gly Arg Lys Pro Pro Ser Leu
 115 120 125
 Gly Glu Ala Gln Pro Thr Lys Asn Leu Glu Glu Asn Lys Ser Leu Lys
 130 135 140
 Glu Gln Arg Lys Gln Asn Asp Leu Cys Phe Leu Lys Ile Leu Leu Gln
 145 150 155 160
 Lys Ile Lys Thr Cys Trp Asn Lys Ile Leu Arg Gly Ile Thr Glu His
 165 170 175

<210> 4
 <211> 384
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Human Fc-gamma1-IL-7

<400> 4
 Glu Pro Lys Ser Ser Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
 1 5 10 15
 Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
 20 25 30
 Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
 35 40 45
 Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
 50 55 60
 Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
 65 70 75 80
 Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
 85 90 95
 Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
 100 105 110
 Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
 115 120 125

<400>	5															
Glu	Pro	Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	
1				5					10					15		
Pro	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	
			20					25					30			
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	
		35					40					45				
Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	
	50					55					60					
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Ala	
65					70					75					80	
Gln	Ser	Thr	Phe	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Val	His	Gln	Asp	
				85					90					95		

32d

Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu
			100					105					110		
Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg
		115					120					125			
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys
	130					135					140				
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp
145					150					155					160
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys
			165					170						175	
Thr	Thr	Pro	Pro	Met	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser
		180					185						190		
Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser
	195					200						205			
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
	210					215					220				
Ala	Thr	Ala	Thr	Pro	Gly	Ala	Asp	Cys	Asp	Ile	Glu	Gly	Lys	Asp	Gly
225					230					235					240
Lys	Gln	Tyr	Glu	Ser	Val	Leu	Met	Val	Ser	Ile	Asp	Gln	Leu	Leu	Asp
			245					250					255		
Ser	Met	Lys	Glu	Ile	Gly	Ser	Asn	Cys	Leu	Asn	Asn	Glu	Phe	Asn	Phe
		260					265					270			
Phe	Lys	Arg	His	Ile	Cys	Asp	Ala	Asn	Lys	Glu	Gly	Met	Phe	Leu	Phe
	275					280						285			
Arg	Ala	Ala	Arg	Lys	Leu	Arg	Gln	Phe	Leu	Lys	Met	Asn	Ser	Thr	Gly
	290					295					300				
Asp	Phe	Asp	Leu	His	Leu	Leu	Lys	Val	Ser	Glu	Gly	Thr	Thr	Ile	Leu
305					310					315					320
Leu	Asn	Cys	Thr	Gly	Gln	Val	Lys	Gly	Arg	Lys	Pro	Ala	Ala	Leu	Gly
			325					330						335	
Glu	Ala	Gln	Pro	Thr	Lys	Ser	Leu	Glu	Glu	Asn	Lys	Ser	Leu	Lys	Glu
			340					345					350		
Gln	Lys	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Leu	Lys	Arg	Leu	Leu	Gln	Glu
		355				360						365			
Ile	Lys	Thr	Cys	Trp	Asn	Lys	Ile	Leu	Met	Gly	Thr	Lys	Glu	His	
	370					375					380				

<210> 6
 <211> 398
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Human FC-gamma1(linker1)-IL-7

<400> 6

Glu	Pro	Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala
1				5				10					15		
Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro
		20					25					30			
Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val
	35					40					45				
Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val
	50					55					60				

32e

```

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65          70          75          80
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
          85          90          95
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
          100         105         110
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
          115         120         125
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr
          130         135         140
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
145          150         155         160
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
          165         170         175
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          180         185         190
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
          195         200         205
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
210          215         220
Ser Ala Thr Ala Thr Pro Gly Ala Gly Gly Gly Ser Gly Gly Gly
225          230         235         240
Gly Ser Gly Gly Gly Ser Asp Cys Asp Ile Glu Gly Lys Asp Gly Lys
          245         250         255
Gln Tyr Glu Ser Val Leu Met Val Ser Ile Asp Gln Leu Leu Asp Ser
          260         265         270
Met Lys Glu Ile Gly Ser Asn Cys Leu Asn Asn Glu Phe Asn Phe Phe
          275         280         285
Lys Arg His Ile Cys Asp Ala Asn Lys Glu Gly Met Phe Leu Phe Arg
          290         295         300
Ala Ala Arg Lys Leu Arg Gln Phe Leu Lys Met Asn Ser Thr Gly Asp
305          310         315         320
Phe Asp Leu His Leu Leu Lys Val Ser Glu Gly Thr Thr Ile Leu Leu
          325         330         335
Asn Cys Thr Gly Gln Val Lys Gly Arg Lys Pro Ala Ala Leu Gly Glu
          340         345         350
Ala Gln Pro Thr Lys Ser Leu Glu Glu Asn Lys Ser Leu Lys Glu Gln
          355         360         365
Lys Lys Leu Asn Asp Leu Cys Phe Leu Lys Arg Leu Leu Gln Glu Ile
          370         375         380
Lys Thr Cys Trp Asn Lys Ile Leu Met Gly Thr Lys Glu His
385          390         395

```

```

<210> 7
<211> 393
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Human Fc-gamma1(YN>AQ)(linker2)-IL-7

```

```

<400> 7
Glu Pro Lys Ser Ser Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1          5          10          15

```

32f

Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro
			20					25					30		
Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val
		35					40					45			
Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val
	50					55					60				
Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
65					70					75					80
Ala	Gln	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln
				85					90					95	
Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala
			100					105					110		
Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro
		115					120						125		
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr
	130					135						140			
Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser
145					150					155					160
Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr
				165					170					175	
Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr
			180				185						190		
Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe
	195					200						205			
Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys
	210					215					220				
Ser	Ala	Thr	Ala	Thr	Pro	Gly	Ala	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly
225					230					235					240
Gly	Asp	Cys	Asp	Ile	Glu	Gly	Lys	Asp	Gly	Lys	Gln	Tyr	Glu	Ser	Val
				245					250					255	
Leu	Met	Val	Ser	Ile	Asp	Gln	Leu	Leu	Asp	Ser	Met	Lys	Glu	Ile	Gly
			260				265						270		
Ser	Asn	Cys	Leu	Asn	Asn	Glu	Phe	Asn	Phe	Phe	Lys	Arg	His	Ile	Cys
	275						280					285			
Asp	Ala	Asn	Lys	Glu	Gly	Met	Phe	Leu	Phe	Arg	Ala	Ala	Arg	Lys	Leu
	290					295					300				
Arg	Gln	Phe	Leu	Lys	Met	Asn	Ser	Thr	Gly	Asp	Phe	Asp	Leu	His	Leu
305					310					315					320
Leu	Lys	Val	Ser	Glu	Gly	Thr	Thr	Ile	Leu	Leu	Asn	Cys	Thr	Gly	Gln
				325					330					335	
Val	Lys	Gly	Arg	Lys	Pro	Ala	Ala	Leu	Gly	Glu	Ala	Gln	Pro	Thr	Lys
			340					345					350		
Ser	Leu	Glu	Glu	Asn	Lys	Ser	Leu	Lys	Glu	Gln	Lys	Lys	Leu	Asn	Asp
		355					360					365			
Leu	Cys	Phe	Leu	Lys	Arg	Leu	Leu	Gln	Glu	Ile	Lys	Thr	Cys	Trp	Asn
	370					375					380				
Lys	Ile	Leu	Met	Gly	Thr	Lys	Glu	His							
385					390										

<210> 8
 <211> 391
 <212> PRT
 <213> Artificial Sequence

<400> 8

Glu 1	Pro	Lys	Ser	Ser 5	Asp	Lys	Thr	His	Thr 10	Cys	Pro	Pro	Cys	Pro	Ala
Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro
			20					25					30		
Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val
		35					40					45			
Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val
	50					55					60				
Asp 65	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln
					70					75					80
Ala	Gln	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln
				85					90					95	
Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala
			100					105					110		
Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro
		115					120					125			
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr
	130					135					140				
Lys 145	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser
					150					155					160
Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr
				165					170					175	
Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr
			180					185					190		
Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe
		195				200						205			
Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys
	210					215					220				
Ser 225	Ala	Thr	Ala	Thr	Pro	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Asp
					230					235					240
Cys	Asp	Ile	Glu	Gly	Lys	Asp	Gly	Lys	Gln	Tyr	Glu	Ser	Val	Leu	Met
				245					250					255	
Val	Ser	Ile	Asp	Gln	Leu	Leu	Asp	Ser	Met	Lys	Glu	Ile	Gly	Ser	Asn
			260					265					270		
Cys	Leu	Asn	Asn	Glu	Phe	Asn	Phe	Phe	Lys	Arg	His	Ile	Cys	Asp	Ala
		275					280					285			
Asn	Lys	Glu	Gly	Met	Phe	Leu	Phe	Arg	Ala	Ala	Arg	Lys	Leu	Arg	Gln
	290					295					300				
Phe 305	Leu	Lys	Met	Asn	Ser	Thr	Gly	Asp	Phe	Asp	Leu	His	Leu	Leu	Lys
					310					315					320
Val	Ser	Glu	Gly	Thr	Thr	Ile	Leu	Leu	Asn	Cys	Thr	Gly	Gln	Val	Lys
				325					330					335	
Gly	Arg	Lys	Pro	Ala	Ala	Leu	Gly	Glu	Ala	Gln	Pro	Thr	Lys	Ser	Leu
			340					345					350		
Glu	Glu	Asn	Lys	Ser	Leu	Lys	Glu	Gln	Lys	Lys	Leu	Asn	Asp	Leu	Cys
		355													

32h

<210> 9
 <211> 18
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Eighteen amino acid internal deletion in the IL-7 moiety

<400> 9
 Val Lys Gly Arg Lys Pro Ala Ala Leu Gly Glu Ala Gln Pro Thr Lys
 1 5 10 15
 Ser Leu

<210> 10
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> End of Fc portion of huFc-gamma1 (YN>AQ,d)(linker2)-IL-7

<400> 10
 Ala Thr Ala Thr Pro
 1 5

<210> 11
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> End of Fc portion of huFc-gamma1 (YN>AQ)(linker2)-IL-7

<400> 11
 Ala Thr Ala Thr Pro Gly Ala
 1 5

<210> 12
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer 2 for huFc-gamma1 (YN>AQ,d)(linker2)-IL-7

<400> 12
 atcatgtctg gatccctcga

20

<210> 13
 <211> 13
 <212> PRT
 <213> Homo sapiens

<220>
<223> Primer 2 for huFc-gamma1(YN>AQ) (linker2)-IL-7

```
<400> 14
atcatgtctg gatccctcga 20
```

<210>	15
<211>	62
<212>	DNA
<213>	Artificial Sequence

<220>
<223> Primer 1 for huFc-gamma1(YN>AQ)(linker2)-IL-7

```
<400> 15
cccgggcgcc ggcggtggag gatcaggtgg tggcggatgat tgtgatattg aaggtaaaga 60
tg 62
```

<210>	16
<211>	56
<212>	DNA
<213>	Artificial Sequence

<220>
<223> Primer 1 for huFc-gamma1(YN>AQ,d)(linker2)-IL-7

```
<400> 16
cccggggcggg ggaggatcag gtggtggcgg tgattgtgat attgaaggta aagatg          56
```

```
<210> 17
<211> 7
<212> DNA
<213> Artificial Sequence
```

<220>
<223> SmaI site on a primer

```
<400> 17
ccccgggt
```

$\langle 210 \rangle$	18
$\langle 211 \rangle$	48

32j

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic oligonucleotide duplex for generation of
huFc-gamma1(linker1)-IL-7

<400> 18

gggtgcaggg ggcggggggca gcggggggcgg aggatccggc gggggctc

48

<210> 19

<211> 908

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic acid sequence for the Fc region of human Fc-gamma2(h)
(FN>AQ)

<400> 19

gagcccaa	aat	cttctg	acaa	aactca	caca	tgcccac	ccgt	gcccagg	taa	gccagccc	cag	60	
gcctcgcc	ct	ccagct	caag	gcgggac	agg	tgccctag	ag	tagcctg	cat	ccagggac	ag	120	
gccccagc	tg	ggtgct	gaca	cgtccac	cctc	catctctt	cc	tcagcacc	ac	ctgtggc	cagg	180	
accgtcag	tc	ttcctctt	cc	ccccaaa	aacc	caaggac	acc	ctcatgat	tct	cccggacc	ccc	240	
tgaggtc	acg	tgcg	tgg	tgg	acgtg	ag	ccacga	aagac	ccc	gaggtcc	agttca	actg	300
gtacgtg	gac	ggcgtg	gagg	tgcata	aatgc	caagaca	aaag	ccacggg	agg	agcagg	ccca	360	
gagcacgt	tc	cgtgtg	gtca	gcgtcct	cac	cgttgtg	cac	caggact	ggc	tgaacgg	caa	420	
ggagtaca	ag	tgcaagg	tct	ccaaca	aaagg	cctccc	agcc	ccc	atcgaga	aaaccat	ctc	480	
caaaacca	aaa	ggtggg	accc	gcgggg	tatg	agggcc	acat	ggacag	aggc	cggctc	ggcc	540	
caccctct	gc	cctgag	agtg	accgct	gtac	caacct	ctgt	ccctac	aggg	cagcccc	gag	600	
aaccacag	gt	gtacac	cctg	ccccat	cac	gggagg	agat	gaccaaga	aac	caggtc	agcc	660	
tgacctgc	ct	ggtcaa	aggc	ttctac	ccca	gcgacat	cgc	cgtggag	tg	gagagca	atg	720	
ggcagcc	gga	gaaca	actac	aagacc	acac	ctcccat	gct	ggactcc	gac	ggctcct	tct	780	
tcctctac	ag	caagct	cacc	gtggaca	aga	gcaggtg	gca	gcagggg	aac	gtcttct	cat	840	
gctccgtg	at	gcatgag	gct	ctgcaca	aacc	actacac	aca	gaagagc	gcc	accgcg	accc	900	
cgggtg	ca											908	

<210> 20

<211> 908

<212> DNA

<213> Artificial Sequence

<220>

<223> Nucleic acid sequence for the Fc region of human Fc-gamma2(h)

<400> 20

gagcccaa	aat	cttctg	acaa	aactca	caca	tgcccac	ccgt	gcccagg	taa	gccagccc	cag	60	
gcctcgcc	ct	ccagct	caag	gcgggac	agg	tgccctag	ag	tagcctg	cat	ccagggac	ag	120	
gccccagc	tg	ggtgct	gaca	cgtccac	cctc	catctctt	cc	tcagcacc	ac	ctgtggc	cagg	180	
accgtcag	tc	ttcctctt	cc	ccccaaa	aacc	caaggac	acc	ctcatgat	tct	cccggacc	ccc	240	
tgaggtc	acg	tgcg	tgg	tgg	acgtg	ag	ccacga	aagac	ccc	gaggtcc	agttca	actg	300
gtacgtg	gac	ggcgtg	gagg	tgcata	aatgc	caagaca	aaag	ccacggg	agg	agcagtt	caa	360	
cagcacgt	tc	cgtgtg	gtca	gcgtcct	cac	cgttgtg	cac	caggact	ggc	tgaacgg	caa	420	
ggagtaca	ag	tgcaagg	tct	ccaaca	aaagg	cctccc	agcc	ccc	atcgaga	aaaccat	ctc	480	

32k

```

caaaaccaa ggtgggaccc gcggggtatg agggccacat ggacagaggc cggctcggcc 540
caccctctgc cctgagagt accgctgtac caacctctgt ccctacagg cagccccgag 600
aaccacaggt gtacaccctg ccccatcac gggaggagat gaccaagaac caggtcagcc 660
tgacctgect ggtcaaaggc ttctaccca gcgacatcgc cgtggagtgg gagagcaatg 720
ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac ggctccttct 780
tctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac gtcttctcat 840
gctccgtgat gcatgaggct ctgcacaacc actacacaca gaagagcgcc accgcgaccc 900
cgggcgcc 908

```

<210> 21
 <211> 911
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Nucleic acid sequence for the Fc region of human Fc-gamma1
 (YN>AQ)

```

<400> 21
gagcccaaat cttctgacaa aactcacaca tgcccaccgt gcccaggtaa gccagcccag 60
gcctcgccct ccagctcaag gcgggacagg tgccctagag tagcctgcat ccagggacag 120
gccccagccg ggtgctgaca cgtccacctc catctcttcc tcagcacctg aactcctggg 180
gggaccgtca gtcttctct tcccccaaa acccaaggac accctcatga tctcccggac 240
ccctgaggtc acatgcgtgg tggaggacgt gagccacgaa gacctgagg tcaagttcaa 300
ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcaggc 360
ccagagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg 420
caaggagtac aagtgcaagg tctccaacaa agccctccca gccccatcg agaaaaccat 480
ctccaaagcc aaaggtggga cccgtggggg gcgagggcca catggacaga ggccggctcg 540
gcccaccctc tgccctgaga gtgaccgtg taccaacctc tgctcctaca gggcagcccc 600
gagaaccaca ggtgtacacc ctgcccccat cacgggagga gatgaccaag aaccagggtca 660
gcctgacctg cctggtcaaa ggcttctatc ccagcgacat cgccgtggag tgggagagca 720
atgggcagcc ggagaacaac tacaagacca cgctcccggt gctggactcc gacggctcct 780
tcttctctta tagcaagctc accgtggaca agagcagggt gcagcagggg aacgtcttct 840
catgctccgt gatgcatgag gctctgcaca accactacac gcagaagagc gccaccgcga 900
ccccgggcgc c 911

```

<210> 22
 <211> 911
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Nucleic acid for the Fc region of human Fc-gamma1

```

<400> 22
gagcccaaat cttctgacaa aactcacaca tgcccaccgt gcccaggtaa gccagcccag 60
gcctcgccct ccagctcaag gcgggacagg tgccctagag tagcctgcat ccagggacag 120
gccccagccg ggtgctgaca cgtccacctc catctcttcc tcagcacctg aactcctggg 180
gggaccgtca gtcttctct tcccccaaa acccaaggac accctcatga tctcccggac 240
ccctgaggtc acatgcgtgg tggaggacgt gagccacgaa gacctgagg tcaagttcaa 300
ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta 360
caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg 420
caaggagtac aagtgcaagg tctccaacaa agccctccca gccccatcg agaaaaccat 480
ctccaaagcc aaaggtggga cccgtggggg gcgagggcca catggacaga ggccggctcg 540

```

321

```

gccaccctc tgcctgaga gtgaccgtg taccaacctc tgtccctaca gggcagcccc 600
gagaaccaca ggtgtacacc ctgcccccat cacgggagga gatgaccaag aaccaggtca 660
gcctgacctg cctgggtcaaa ggcttctatc ccagcgacat cgccgtggag tgggagagca 720
atgggcagcc ggagaacaac tacaagacca cgcctcccgt gctggactcc gacggctcct 780
tcttcctcta tagcaagctc accgtggaca agagcaggtg gcagcagggg aacgtcttct 840
catgctccgt gatgcatgag gctctgcaca accactacac gcagaagagc gccaccgoga 900
ccccgggcgc c 911

```

<210> 23
 <211> 9
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Beginning of the hinge region as shown in Figure 20

<400> 23
 gagcccaaa 9

<210> 24
 <211> 7
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> AflIII sequence

<400> 24
 cttaagc 7

<210> 25
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Linker 2 for Fc-IL-7 fusion proteins

<400> 25
 Gly Gly Gly Gly Ser Gly Gly Gly Gly
 1 5

<210> 26
 <211> 4
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Replacement sequence near the C-terminus of the Fc portion of an immunoglobulin heavy chain

32m

<400> 26
Ala Thr Ala Thr
1

<210> 27
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Sequence near the C-terminus of the Fc portion of an
immunoglobulin heavy chain

<400> 27
Leu Ser Leu Ser
1

<210> 28
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Replacement sequence within an IgG1 heavy chain

<400> 28
Gln Ala Gln Ser
1

<210> 29
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Sequence within an IgG1 heavy chain

<400> 29
Gln Phe Asn Ser
1

<210> 30
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Sequence from IgG1

<400> 30
Gln Tyr Asn Ser
1

32n

<210> 31
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Sequence from IL-7

<400> 31
 Met Asn Ser Thr Gly
 1 5

<210> 32
 <211> 5
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Sequence from IL-7

<400> 32
 Leu Asn Cys Thr Gly
 1 5

<210> 33
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Flanking reverse primer for generation of Fc-gammal-IL-7 fusion proteins

<400> 33
 cagggtgtac acctgtggtt c

21

<210> 34
 <211> 15
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Linker 1 for generation of Fc-IL-7 fusion proteins

<400> 34
 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15

<210> 35
 <211> 6
 <212> PRT
 <213> Artificial Sequence

32o

<220>

<223> Sequence from IL-7

<400> 35

Ser Leu Glu Glu Asn Lys

1

5

<210> 36

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Mutagenic primer in the sense direction

<400> 36

agcaggccca gagcacgtac cgtgtggt

28

<210> 37

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Complementary strand of the mutagenic primer in the sense
direction

<400> 37

gtacgtgctc tgggcctgct cctcccgc

28

<210> 38

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Flanking forward primer

<400> 38

ctctctgcag agcccaaata t

21

26474-1011

33

CLAIMS:

1. An IL-7 fusion protein having improved biological properties compared to wild-type mature human IL-7, comprising:
 - a first portion comprising an Fc region from IgG1 or IgG2 wherein the Fc region comprises a hinge region, a CH2 domain, and a CH3 domain; and
 - a second portion comprising a polypeptide having an amino acid sequence comprising at least 98% sequence identity with wild-type mature human interleukin-7 (IL-7) wherein wild-type mature human IL-7 consists of amino acid residues 26-176 of SEQ ID NO:1, and wherein amino acid residues corresponding to positions 70 and 91 of wild-type mature human IL-7 are glycosylated and the amino acid residue corresponding to position 116 of wild-type mature human IL-7 is non-glycosylated.
2. The fusion protein of claim 1, wherein the amino acid residue corresponding to position 116 of wild-type mature human IL-7 is asparagine.
3. The fusion protein of claim 1, wherein the polypeptide having 98% sequence identity with wild-type mature human IL-7 comprises disulfide bonding between residues corresponding to positions Cys2 and Cys92, Cys34 and Cys129, and Cys47 and Cys141 of wild-type mature human IL-7.
4. The fusion protein of claim 1, further comprising a linker between the first portion and the second portion.
5. The fusion protein of claim 1, wherein the Fc region is from IgG1.
6. The fusion protein of claim 5, wherein the Fc region further comprises an amino acid substituted in place of Asn297 at the glycosylation site of the Fc region.
7. The fusion protein of claim 6, wherein the Fc region further comprises an amino acid substituted in place of Tyr296 at the glycosylation site of the Fc region.

26474-1011

34

8. The fusion protein of claim 7, wherein Ala is substituted in place of Tyr296 and Gln is substituted in place of Asn297 at the glycosylation site of the Fc region.
9. The fusion protein of claim 1, wherein the Fc region is from IgG2.
- 5 10. The fusion protein of claim 9, wherein the Fc region further comprises an IgG1 hinge.
11. The fusion protein of claim 9, wherein the Fc region further comprises an amino acid substituted in place of Asn297 at the glycosylation site of the Fc region.
- 10 12. The fusion protein of claim 11, wherein the Fc region further comprises an amino acid substituted in place of the Phe296 residue adjacent the Asn297 residue at the glycosylation site of the Fc region.
13. The fusion protein of claim 12, wherein Ala is substituted in place of Phe296 and Gln is substituted in place of Asn297 at the glycosylation site of the Fc
15 region.
14. The fusion protein of claim 1, wherein said second portion comprises wild-type mature human IL-7 consisting of amino acid residues 26-176 of SEQ ID NO:1, and wherein the amino acid residues corresponding to positions 70 and 91 of said wild-type mature human IL-7 are glycosylated and the amino acid residue
20 corresponding to position 116 of said wild-type mature human IL-7 is non-glycosylated.
15. A purified nucleic acid encoding the fusion protein of claim 1.
16. A cultured host cell comprising the nucleic acid of claim 15.
17. A fusion protein comprising:

26474-1011

35

a first portion comprising an Fc region from IgG1 or IgG2 wherein the Fc region comprises a hinge region, a CH2 domain, and a CH3 domain; and

a second portion comprising wild-type mature human IL-7 consisting of amino acid residues 26-176 of SEQ ID NO:1 except that residues 121 to 138 of SEQ ID NO:1 are deleted in said second portion, and wherein the amino acid residues corresponding to positions 70 and 91 of wild-type mature human IL-7 are glycosylated and the amino acid residue corresponding to position 116 of wild-type mature human IL-7 is non-glycosylated.

Figure 1Human IL-7

MFHVSFRYIFGLPPLILVLLPVASSDCDIEGKDGKQYESVLMVSIQQLDSMKEIGSNC
LNNEFNFFKRHICDANKEGMFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTTILLNCT
GQVKGRKPAALGEAQPTKSLEENKSLKEQKKLNDLCFLKRLQLQKTCWNKILMGTKEH

(SEQ ID NO:1)

Figure 2*Cow IL-7*

MFHVSFRYIFGIPPLILVLLPVASSDCDISGRDGGAYQNVLMVNIDDLDNMINFDSNCL
NNEPNFFKKHSCDDNKEASFLNRASRKLQFLKMNISDDFKLHLSTVSQGTLLNCTS
KGKGRKPPSLSEAQPTKNLEENKSSREQKKQNDLCFLKILLQKIK TCWNKILRGIKEH

(SEQ ID NO:2)

Figure 3*Sheep IL-7*

MFHVSFRYIFGIPPLILVLLPVASSDCDFSGKDGGAYQNVLMVSIDDLDNMINFDSNCL
NNEPNFFKKHSCDDNKEASFLNRAARKLKQFLKMNISDDFKLHLSTVSQGTLTLLNCTS
KGKGRKPPSLGEAQPTKNLEENKSLKEQRKQNDLCFLKILLQKIKTCWUNKILRGITEH

(SEQ ID NO:3)

Figure 4*Human Fcγ1-IL-7*

EPKSSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSATATPGADCD
IEGKDGKQYESVLMVSIQQLDSMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAARK
LRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVKGRKPAALGEAQPTKSLEENKS LKE
QKKLNDLCFLKRLLQEIKTCWNKILMGTKEH

(SEQ ID NO:4)

Figure 5

Human Fcy2(h)(FN>AO)-IL-7

EPKSSDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFN
WYVDGVEVHNAKTKPREEQAQSTFRVVS VLT VVHQDWLNGKEYKCKVSNKGLPAPIE
KTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSATATPGADCD
IEGKDGKQYESVLMV SIDQLLD SMKEIGSNCLNNEFNFFKRHICDANKEGMFLFRAARK
LRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVKGRKPAALGEAQPTKSLEENKSLKE
QKKLNDLCFLKRLLQEIKTCWNKILMGTKEH

(SEQ ID NO:5)

Figure 6Human FCγ1(linker1)-IL-7

EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSATATPGAGGG
GSGGGGSGGGGSDCDIEGKD GKQYESVLMVSDQLLD SMKEIGSNCLNNEFNFFKRHICD
ANKEGMFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVKGRKPAALGE
AQPTKSLEENKSLKEQKKLNDLCFLKRLLQEIKTCWNKILMGTKEH

(SEQ ID NO:6)

Figure 7

Human Fcγ1 (YN>AQ)(linker2)-IL-7

EPKSSDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQAQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSA TATPGAGGG
GSGGGGDCDIEGKD GKQYESVLMV SIDQLLD SMKEIGSNCLNNEFNFFKR HICDANKEG
MFLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVKGRKPAALGEAQPTK
SLEENKSLKEQKKLNDLCFLKRLLEIKTCWNKILMGTKEH

(SEQ ID NO:7)

Figure 8

Human Fcγ1 (YN>AQ,d)(linker2)-IL-7

EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVTVVDVSHEDPEVKFN
WYVDGVEVHNAKTKPREEQAQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSATATPGGGGS
GGGGDCDIEGKDQYESVLMVSIQQLDSMKEIGSNCLNNEFNFFKRHICDANKEGM
FLFRAARKLRQFLKMNSTGDFDLHLLKVSEGTTILLNCTGQVKGRKPAALGEAQPTKSL
EENKSLKEQKKLNDLCFLKRLLEIKTCWNKILMGTKEH

(SEQ ID NO:8)

Figure 9*Human Fcγ1*

GAGCCCAAATCTTCTGACAAAACCTCACACATGCCACCGTGCCCAGGTAAGCCAGCCCAGGCCTCGCC
CTCCAGCTCAAGGCGGGACAGGTGCCCTAGAGTAGCCTGCATCCAGG GACAGGCCCCAGCCGGGTGC
TGACACGTCCACCTCCATCTCTTCCTCAGCACCTGAACTCCTGGGGGG ACCGTCAGTCTTCCTCTTCCC
CCCAAACCCAAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCA CATGCGTGGTGGTGGACGTGA
GCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTG GAGGTGCATAATGCCAAGAC
AAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAG
GACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGC CCTCCCAGCCCCCATCGAGA
AAACCATCTCCAAAGCCAAAGGTGGGACCCGTGGGGTGCGAGGGCCA CATGGACAGAGGCCGGCTCG
GCCACCCCTCTGCCCTGAGAGTGACCGCTGTACCAACCTCTGTCCCTA CAGGGCAGCCCCGAGAACCA
CAGGTGTACACCCTGCCCCCATCACGGGAGGAGATGACCAAGAACCA GGTCAGCCTGACCTGCCTGGT
CAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATG GGCAGCCGGAGAACAACACTAC
AAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTAT AGCAAGCTCACCGTGGACAA
GAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATG AGGCTCTGCACAACCACTACA
CGCAGAAGAGCGCCACCGCGACCCCGGGCGCC

(SEQ ID NO:22)

Figure 10

Human Fcγ1 (YN>AO)

GAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCCCAGGTAAGCCAGCCCAGG
CCTCGCCCTCCAGCTCAAGGCGGGACAGGTGCCCTAGAGTAGCCTGCATCCAGGGACAGGC
CCCAGCCGGGTGCTGACACGTCCACCTCCATCTCTTCCTCAGCACCTGAACTCCTGGGGGGA
CCGTCAGTCTTCCTCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGA
GGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC
GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGGCCCAGAGC
ACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAGCC
AAAGGTGGGACCCGTGGGGTGCGAGGGGCCACATGGACAGAGGCCGGCTCGGCCCACCCTCT
GCCCTGAGAGTGACCGCTGTACCAACCTCTGTCCCTACAGGGCAGCCCCGAGAACCACAGGT
GTACACCCTGCCCCCATCACGGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTG
GTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGA GTGGGAGAGCAATGGGCAGCCGGAGA
ACA ACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTATAGCAAG
CTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATG
AGGCTCTGCACAACCACTACACGCAGAAGAGCGCCACCGCGACCCCGGGCGCC

(SEQ ID NO:21)

Figure 11*Human Fcγ2(h)*

GAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCCCAGGTAAGCCAGCCCAGG
CCTCGCCCTCCAGCTCAAGGCGGGACAGGTGCCCTAGAGTAGCCTGCATCCAGGGACAGGC
CCCAGCTGGGTGCTGACACGTCCACCTCCATCTCTTCCTCAGCACCACTGTGGCAGGACCG
TCAGTCTTCCTCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTC
ACGTGCGTGGTGGTGGACGTGAGCCACGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGG
ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGGAGGAGCAGTTCAACAGCACGTT
CCGTGTGGTCAGCGTCCTCACCGTTGTGCACCAGGACTGGCTGAACGGCAAGGAGTACAAGT
GCAAGGTCTCCAACAAAGGCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAACCAAAGG
TGGGACCCGCGGGGTATGAGGGGCCACATGGACAGAGGCCGGCTCGGCCCACCCTCTGCCCT
GAGAGTGACCGCTGTACCAACCTCTGTCCCTACAGGGCAGCCCCGAGAACCACAGGTGTAC
ACCCTGCCCCCATCACGGGAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCA
AAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA
CTACAAGACCACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCA
CCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGC
TCTGCACAACCACTACACACAGAAGAGCGCCACCGCGACCCCGGGCGCC


(SEQ ID NO:20)

Figure 12

Human Fcγ2(h)(FN>AO)

GAGCCCAAATCTTCTGACAAAACCTCACACATGCCCCACCGTGCCCAGGTAAGCCAGCCCAGG
CCTCGCCCTCCAGCTCAAGGCGGGACAGGTGCCCTAGAGTAGCCTGCATCCAGGGACAGGC
CCCAGCTGGGTGCTGACACGTCCACCTCCATCTCTTCCTCAGCACCACTGTGGCAGGACCG
TCAGTCTTCCTCTTCCCCCCTCAAGGACACCTCATGATCTCCCGGACCCCTGAGGTC
ACGTGCGTGGTGGTGGACGTGAGCCACGAAAGACCCCGAGGTCCAGTTCAACTGGTACGTGG
ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGGAGGAGCAGGCCAGAGCACGT
TCCGTGTGGTCAGCGTCCTCACCGTTGTGCAACAGGACTGGCTGAACGGCAAGGAGTACAAG
TGCAAGGTCTCCAACAAAGGCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAACCAAAG
GTGGGACCCGCGGGGTATGAGGGCCACATGGACAGAGGCCGGCTCGGCCCACCCTCTGCCC
TGAGAGTGACCGCTGTACCAACCTCTGTCCCTACAGGGCAGCCCCGAGAACCACAGGTGTAC
ACCCTGCCCCCATCACGGGAGGAGATGACCAAGAACCAGGTTCAGCCTGACCTGCCTGGTCA
AAGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA
CTACAAGACCACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCA
CCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGC
TCTGCACAACCACTACACACAGAAGAGCGCCACCGCGACCCCGGGTGCA

(SEQ ID NO:19)

Figure 13

14 / 16


Figure 14


Figure 15

Figure 16