

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199868068 B2**
(10) Patent No. **748461**

(54) Title
Purification of high order transcription complexes from transgenic non-human animals

(51)⁶ International Patent Classification(s)
C12N 015/11 C12N 015/12

(21) Application No: 199868068 (22) Application Date: 1998 .05 .25

(30) Priority Data

(31) Number (32) Date (33) Country
97108433 1997 .05 .26 EP

(43) Publication Date : 1998 .11 .26
(43) Publication Journal Date : 1998 .11 .26
(44) Accepted Journal Date : 2002 .06 .06

(71) Applicant(s)
Hoechst Aktiengesellschaft

(72) Inventor(s)
Bernd Kirschbaum; Erick Berglund; Michael Meisterernst; Greg Polites

(74) Agent/Attorney
WATERMARK PATENT and TRADEMARK ATTORNEYS,Locked Bag 5,HAWTHORN VIC 3122

(56) Related Art
BIOTECHNIQUES 21, 718-25
GENES DEV 6, 304-15, 10 AND 711-24

Abstract

Purification of higher order transcription complexes from transgenic non-human animals

5

The invention relates to transgenes comprising DNA encoding for epitope-tagged TATA-box binding protein (TBP), the production of transgenic animals which express epitope-tagged TBP and use thereof for affinity purification of (novel) transcription factors and transcription complexes from a variety of eukaryotic tissues and cell-types.

10



AUSTRALIA

Patents Act 1990

**ORIGINAL
COMPLETE SPECIFICATION
STANDARD PATENT**

Application Number:

Lodged:

Invention Title:

PURIFICATION OF HIGHER ORDER TRANSCRIPTION COMPLEXES
FROM TRANSGENIC NON-HUMAN ANIMALS

The following statement is a full description of this invention, including the
best method of performing it known to us :-

Purification of higher order transcription complexes from transgenic non-human
5 animals.

The invention relates to a transgene that comprises DNA encoding for epitope-
tagged TATA-box binding protein (TBP), the production of transgenic animals which
express epitope-tagged TBP and the use thereof for affinity purification of (novel)
10 transcription factors and transcription complexes from a variety of eukaryotic tissues
and cell-types.

Central to regulation of the eukaryotic transcription event is the stepwise formation
and activity of the pre-initiation complex. This is a large, multi-subunit complex
15 which is necessary for the correct positioning and initiation of the RNA polymerase II
enzyme at the transcription start site. Many of the general transcription factors
(GTFs) of this complex have been characterized from eukaryotic nuclei in recent
years including TFIIA, TFIIB, TFIID, TFIIE, TFIIF and TFIIH (Zawel and Reinberg
(1995) Ann. Rev. Biochem. 64, 533-561; Serizawa et al. (1994) In: Transcription:
20 Mechanisms and regulation, 45-66, Raven press). In TATA-containing promoters,
TFIID has a specific affinity for the TATA-box sequence. It is through this sequence
recognition that TFIID is the first element to bind the promoter in basal RNA
polymerase II transcription, thus nucleating complex formation (Lewin, B. (1990) Cell
61: 1161-1164). The other GTFs bind in a defined, stepwise manner, resulting in a
25 completed pre-initiation complex. Subsequently, this large complex recruits and
correctly positions the RNA Polymerase II (Pol II) at the transcription start site to
initiate basal transcription. It has become clear that TFIID, itself a multi-subunit
complex, plays a key role in the regulation of "activated transcription", loosely
defined as elevated levels of mRNA production in the presence of transcriptional
30 activators. Such activators can be naturally occurring enhancer-binding
determinants, such as the E-box binding USF (Sawadogo and Roeder (1985) Cell

43: 165-175; Kirschbaum et al. (1992) *Mol. Cell. Biol.* 12: 5094-5100), or viral factors such as VP16 (Stringer et al. (1990) *Nature* 345: 783-786).

TFIID is composed of the TATA-binding protein (TBP) and several TBP-associated factors (TAF_{II}s) (In the application "TAF_{II}s" include any kind of transcription factor, transcription activator, transcription inhibitor). As many as 20 different TAF_{II}s have been characterized to date, and TFIID complexes containing different combinations of TAF_{II}s have been observed (Zawel and Reinberg (1995) *Ann. Rev. Biochem.* 64: 533-561; Hori, R. and Carey, M. (1994) *Curr. Opin. Gen. Dev.* 4: 236-244).

Moreover, different combinations of TAF_{II}s lend distinct properties to the TFIID complex. In *Drosophila*, for example, the pattern formation proteins Hunchback (HB) and Bicoid (BCD) are absolutely reliant on the presence of TAF_{II}60, TAF_{II}110, and TAF_{II}250 in the TFIID complex (Sauer, F. et al. (1995) *Science* 270, 1783-1788). These TFIID components act as co-activators to the upstream enhancer-bound HB and BCD proteins. The neurogenic factor NTF-1 has been shown to require a minimum complex of TBP, TAF_{II}150 (to which it binds) and TAF_{II}250 for activated transcription, whereas SP1 requires the additional factor TAF_{II}110 for its activation (Chen, J.-L. et al. (1994) *Cell* 79: 93-105). Another study has identified TAF_{II}28, whose presence is necessary for transcriptional activation by the estrogen and vitamin D3 nuclear receptors (May, M. et al. (1996) *EMBO* 15: 3093-3104). It is likely that the TAF_{II}s act as transcriptional adapters, transmitting regulatory information from activator/repressor factors to the core initiation complex by way of protein-protein interactions.

As regulated transcription is currently viewed, the expression of individual genes and/or small groups of closely related loci are controlled by definable sets of transcription complex subunits. Though some of the factors are ubiquitous and present in most transcription events, for example GTFs, increasing numbers of gene- and cell-specific elements of regulated transcription are now being described.

There are several proven methods which have been used to identify transcription factors. The early strategies, which uncovered RNA Pol II and the seven GTFs (TFIIA, TFIIB, TFIID, TFIIIE, TFIIF, TFIIH, and TFIIJ), mainly involved column fractionation of nuclear preparations from cell lines (Zawel and Reinberg (1995); Serizawa et al. (1994); Roeder, R.G. (1996) TIBS 21: 327-335). These fractions yielded semi-purified proteins with various amounts of transcriptional ability. Most of these fractions appeared to be absolutely necessary for basal transcription. At this point, it was known that the TFIID fraction was responsible for TATA-box recognition. However, attempts to isolate a single protein with TATA-binding ability were not successful. A breakthrough occurred when a single component of yeast was shown to be able to replace TFIID in reconstituted basal transcription assays, which led to the isolation and cloning of a 27kD TATA-binding protein (TBP) (Buratowski, S. et al. (1988) Nature 334: 37-42; Cavallini, B. et al. (1988) Proc. Nat. Acad. Sci. 86: 9803-9809). Use of degenerated primers led to the further identification of genes for the TBP subunit of human (Kao, C.C. et al. (1990) Science 248: 1646-1649; Hoffmann, A. et al. (1990) Nature 346: 387-390; Peterson, M.G. et al. (1990) Science 248: 1625-1630), Drosophila (Hoey, T. et al. (1990) Cell 61: 1179-1186; Muhich, M.L. et al. (1990) Proc. Nat. Acad. Sci. 87, 9148-9152), and mouse TFIID (Tamura, T. et al. (1991) Nuc. Acids Res. 19: 3861-3865).

The availability of the cDNA for TBP from different species made possible a wide range of investigations including the over-expression of these proteins in cell culture. HeLa cell lines were produced which constitutively expressed a TBP protein with an FLAG-tag or the influenza virus hemagglutinin (HA) epitope-tag added to its amino terminus (Zhou, Q. et al. (1993) Genes & Development 7, 180-187; Chiang et al. (1993) EMBO 12, 2749-2762). The FLAG-tag is an epitope consisting of a synthetic sequence of eight amino acids. The HA-tag is a natural epitope with the amino acid sequence SEQ ID NO. 1 "MGYPYDVPDYAV" (one letter code).

Also a shorter peptide from the natural HA tag (10 amino acids from the influenza virus hemagglutinin) has been used for the expression of a fusion-protein containing TBP in a *Drosophila* cell line (Colgan and Manley (1992), *Genes Dev.* 6, 304-331; Trivrdi et al., (1996) *Mol. Cel. Biol.* 16, 6909-6916).

5

TBP proteins with two epitopes tagged to its amino terminus, the FLAG- and the HA-epitope have been expressed in bacteria (Chiang et al. (1993) *EMBO* 12: 2749-2762). FLAG-tagged TBP proteins have also been expressed under the control of an inducible promoter (Wu et al., (1996) *BioTechniques* 21: 718-725). However, monoclonal antibodies against the epitope/epitopes were used to purify TBP-associated complexes from nuclear extracts, thus co-purifying TBP-associated factors of the TFIID complex (TAF_{II}s) (Zhou et al, (1993) *Genes Dev.* 7: 180-187).

15

Research continues along these lines in many laboratories, with a recent wave of new TAFs and TAF-interacting factors being identified and characterized from HeLa nuclear extracts and yeast (Hori and Carey (1994) *Curr. Op. Gen. Dev.* 4: 236-244; Zawel and Reinberg (1995) *Ann. Rev. Biochem.* 64: 533-561; Roeder, R.G. (1996) *Trends Biochem. Sci.* 21, 327-335).

20

From human the TAFs TAF_{II}68, TAF_{II}55, TAF_{II}30, TAF_{II}28, TAF_{II}20 and TAF_{II}18 are known (Mengus et al. (1995) *EMBO* 14: 1520-1531; Bertolotti et al. (1996) *EMBO* 15: 5022-5031; Wu and Chiang (1996) *Biotechniques* 21: 718-725).

25

It cannot be overemphasized that although a wealth of factors have been found with these methods, the research is confined to transcription complexes, TAFs and TAF-interacting factors that are particular to the cell-line type or yeast strain that is being used.

30

The invention relates to a more "universal system", wherein epitope-tagged TBP is applied for the affinity purification of novel transcription complexes and transcription

factors, TAFs and TAF-interacting factors in the context of a whole animal, and therefore, from a variety of different eukaryotic tissues and cell types.

The invention relates to a transgenic non-human animal having the ability to express
5 epitope-tagged TATA-box binding protein (TBP). In a further aspect, the invention relates to the use of the transgenic non-human animal, preferably for the identification and isolation of higher order transcription complexes and for identification and isolation of proteins associated in the higher order transcription complex (TAFs and TAF-interacting factors, e.g. transcription factors). In another
10 aspect the invention relates to the preparation of the non-human transgenic animal by introducing a transgene into the germline and/or into somatic cells of the non-human transgenic animal, preferably at a particular stage of development. The invention further relates to a transgene that can be used for making the transgenic non-human animals.

15

Thus an embodiment of the invention provides a transgene that encodes an epitope-tagged TBP. Preferably the transgene comprises a first DNA sequence that encodes for one or more epitope-tags and that comprises a second DNA sequence that encodes a TBP. The transgene may comprise further DNA-sequence(s) which
20 encode epitope-tag(s). Preferably, the DNA that encodes TBP is a cDNA. The DNA that encodes TBP could be any naturally occurring DNA, a derivative or a part thereof. The DNA, preferably cDNA can e.g. be from eukaryotes, including birds, amphibians, reptiles, yeast, *C. elegans*, mammals, etc. For example the TBP-encoding DNA from rodents, sheep, dog, cow, pig and primates, human or a part thereof can be used. Preferred is the use of human TBP (hTBP) -cDNA. The
25 invention also comprises the use of any non-naturally occurring DNA, e.g. a TBP-cDNA derivative. A derivative of the DNA might for example have a altered sequence, e.g. a mutated or modified sequence and/or might comprise modified nucleotides. A derivative of the DNA can also be a salt, preferably a physiological
30 tolerable salt.

The transgene comprises one or more DNA sequences that encode for one, two, three, four, five or more epitope-tags. Preferably, the transgene comprise the DNA that encodes for two epitope-tags. The DNA encoding the individual epitope-tags can be located at the 5'- and/or the 3'-end of the DNA that encodes TBP and/or at
 5 any suitable position in between the sequence of the DNA that encodes TBP. The DNA encoding individual epitope-tags can be separated and/or arranged in tandem or can be directly adjacent respectively.

As epitope-tag, any natural or synthetic peptide can be used. Each epitope-tag is
 10 expressed as fusion protein with TBP, the epitope tag may e.g. be connected directly to the TBP or by a spacer peptide. Preferably, an epitope-tag should offer the opportunity to affinity purify TBP or the fusion protein and to affinity purify proteins which are associated with the TBP or the fusion protein, like TAFs and TAF-interacting factors. Furthermore, an epitope-tag should not destroy the
 15 functional activity of the TBP when expressed as fusion protein with TBP. For this purpose preferably short peptides are employed as epitope-tags. They can comprise about 1 to 50 or more amino acids, in particular peptides are employed that comprise 5 to 15 amino acids. Non-limiting examples of peptides that can be used as epitope-tags are the FLAG-epitope, the HA-epitope, multiple Histidine residues
 20 (6 to 10 histidine residues or more, preferably 6 histidine residues) (His tag), the Myc tag (Stone et al. (1996) Nature 384: 129-134), streptavidin tags and others. Also shorter peptides of natural epitopes can be used for this purpose. For example the use of the HA-epitope include the use of the epitopes "MGYPYDVPDYA" (SEQ ID NO. 2), "GYPDVPDYA" (SEQ ID NO. 3), "YPYDVPDYA" (SEQ ID NO. 4) or
 25 other peptides derived from the HA-epitope.

In one embodiment of the invention the transgene contains the cDNA of human TBP and two DNA-sequences which encode an epitope-tag. Preferably the first DNA sequence encodes the HA epitope, which may serve as an epitope for
 30 immunoreaction e.g. with a commercially available monoclonal antibody (Kolodziej and Young (1991) Meth. Enzym. 194: 508-519). Just 3' to the sequence that

- encodes the HA tag a DNA sequence is located, which encodes for a stretch of 6 histidine residues (His tag). The His tag can form a non-covalent, reversible complex with Ni^{2+} ions. For example, commercially available Ni^{2+} agarose affinity column material is routinely used to purify His-tagged proteins (Hochuli, E. et al. (1987) J. Chromatography 411: 177-184; Janknecht, R. et al. (1991) Proc. Nat. Acad. Sci. 74: 4835). In a special embodiment of the invention the transgene comprises the DNA sequence SEQ ID NO. 13. The SEQ ID NO. 13 provides a transgene which encodes for a fusion protein consisting of double tagged hTBP.
- 10 In a preferred embodiment the invention provides a transgene that encodes for epitope-tagged TBP and that comprises a promoter for the expression of the fusion protein. The transgene can comprise one or more gene regulatory sequences in addition to a DNA which encodes TBP (e.g. cDNA of TBP or a derivative thereof) and DNA-sequence(s) encoding epitope-tags. Such gene regulatory sequences are,
- 15 for example, natural or synthetic promoters or parts thereof and/or cis-acting elements (e.g. enhancer, silencer). For this purpose mammalian promoters, for example the promoter of the mouse transferrin gene, the promoter of the neuron-specific enolase (NSE) gene (Forss-Petter, S. et al. (1990) Neuron 5: 187-197) or the promoter of the thymidine kinase gene can be used. Also viral promoters, like
- 20 for example the promoters of the cytomegalovirus genes or the SV 40 early gene can be used. Preferably inducible or constitutive promoters or derivatives thereof are used. As constitutive promoter for example the promoter of the human elongation factor-1 alpha gene (EF) (Uetsuki, T. et al. (1989) J. Biol. Chem. 264: 5791-5798) can be used. As inducible promoter for example the metallothionein
- 25 promoter (MT), which has several cis-elements that are responsive to heavy metals (Palmiter, R.D. (1987) Experimentia Supplementum 52: 63-80, Birkhäuser Verlag) can be used. In addition, a promoter of a gene which is expressed e.g. cell-cycle specific, cell-type specific or developmental specific can be used for this purpose.
- 30 In a special embodiment of the invention the transgene comprises the cDNA of hTBP and DNA-sequences that encode for HA epitope (e.g. 9 amino acids of the

natural HA epitope) and His epitope (e.g. 6xhis tag) and a constitutive promoter. For example, TBP expression is controlled by the promoter for human elongation factor-1 alpha (EF) (Uetsuki et al. (1989) J. Biol. Chem. 264: 5791-5798). The EF-promoter is a TATA-less promoter that has been used to express transgenes in mice at

5 moderate but constant levels (Hanaoka, K. et al. (1991) Differentiation, 183-189).

In a special embodiment of the invention the transgene comprises the DNA sequence which encodes for double-tagged (HA and His epitope) hTBP and the sequence of the EF-promoter, in particular the transgene has the DNA sequence SEQ ID NO. 14.

10

In another special embodiment of the invention the transgene comprises a DNA that encodes for TBP, preferably the cDNA of hTBP and DNA-sequences that encode for HA (e.g. 9 amino acids of the natural HA epitope) and His epitope (e.g. 6xhis) and an inducible promoter. Preferably, the inducible promoter is the metallothionine

15 promoter (MT). For example, in the case that the introduction of an additional TBP encoding sequence into an animals genome and the subsequent expression of TBP

or a TBP fusion protein respectively is toxic, this embodiment of the invention will allow the animal to come to term with the transgene encoding for the TBP fusion protein lying silent until the promoter is introduced. The promoter can for example be induced when the animal is full grown or at any developmental stage of interest, e.g. in the case of the MT promoter with an interperitoneal injection containing divalent cations like Zn^{2+} , Mg^{2+} , Mn^{2+} or Cd^{2+} . As has been shown in other models, the MT directed gene will then express at elevated levels (Palmiter, R.D. et al. (1982) Cell 29: 701-710). In a particular embodiment of the invention the transgene

25 comprises a DNA sequence which encodes for double-tagged hTBP (HA and HIS epitope) and the MT-promoter, in particular the transgene has the DNA sequence SEQ ID NO. 15.

30 The invention further relates to a method of making a transgene by connecting the DNA sequence(s) that encode for one or more epitope-tags to the DNA sequence that encodes for the TBP protein.

The invention further relates to the use of the transgene. For example the transgene can be used for the preparation of a recombinant vector.

The invention also relates to a method of preparing a recombinant vector. This
5 method comprises the integration of the transgene into an appropriate vector, e.g. a vector that contains regulatory sequences. Examples for vectors are expression vectors and retroviruses or derivatives thereof.

The invention further relates to the use of recombinant vectors which comprise the
10 transgene. In particular the invention relates to the use of a recombinant vector that comprises a transgene (that e.g. contains a cDNA of TBP or a derivative thereof, in particular hTBP and DNA-sequences encoding epitope-tags, for example HA and His epitope encoding DNA-sequences). The invention relates to the use of the vector for introducing the transgene into an eukaryotic cell, in particular for the
15 introduction into a mammalian cell. The invention further relates to the use of a vector comprising such transgene for the amplification of the transgenic DNA in bacteria or in eukaryotic cells. An eukaryotic cell into which a vector that contains the transgene has been introduced can also be used for the heterologous/
transgenic expression of the transgene. The eukaryotic cell (host cell) might be part
20 of a transgenic animal.

In a major embodiment of the invention the transgene is used for the preparation of a transgenic non-human animal. Therefore, a transgene or a recombinant vector comprising the transgene is introduced into a host cell and/or an animal. In another
25 aspect the invention relates to a transgenic non-human animal that has been produced by introducing the transgene into the animal or a cell thereof. A transgenic animal according to the invention has the ability to express or to overexpress TPB or a fusion protein comprising or consisting of epitope-tagged TBP.

30 For the preparation of a transgenic animal a non-human animal is used as host animal. Such non-human animal include vertebrates such as rodents, non-human

primates, sheep, goat, dog, cow, pig, birds, amphibians, reptiles, etc. Preferred animals are selected from non-human mammalian species of animals, preferably, animals from the rodent family including rats and mice, most preferably mice.

- 5 A transgenic non-human animal according to the invention comprises any animal into the genome of which one or more copies of a transgene(s) that directs the expression of or which encodes for TBP or derivatives thereof, like a fusion protein consisting of or comprising TBP and epitope-tags. The transgenic animal should have the ability to express the epitope-tagged TBP protein. In a particular
10 embodiment of the invention the transgenic animal can have a transgenic interruption or alteration of the endogenous TBP gene(s) (knock-out animal).

- The transgenic animal according to the invention is an animal into which by nonnatural means (i.e. by human manipulation), one or more TBP genes
15 (transgenes according to the invention) that do not occur naturally in the animal e.g., foreign TBP gene or a derivative thereof, like genetically engineered endogenous or foreign TBP gene have been introduced. The non-naturally occurring TBP gene is called transgene. The transgene may be from the same or a different species as the animal but in any case the transgene is not naturally found in the animal in the
20 configuration and/or at the chromosomal locus conferred by the transgene.

- The transgene (transgenetic DNA) may comprise a foreign gene encoding for TBP, i.e. sequences not normally found in the genome of the host animal, like a TBP gene or a cDNA obtained from a different animal species. Alternatively or additionally, a
25 transgene may comprise an endogenous gene encoding for TBP, e.g. DNA sequences that are abnormal in that they have been rearranged or mutated in vitro in order to alter the normal in vivo pattern of expression of the TBP gene, or to alter or eliminate the biological activity of the endogenous TBP. The invention also relates to expression vectors that comprise the transgene and that can be used to
30 prepare the transgenic animal.

The invention further relates to a method of preparing a transgenic animal according to the invention. A transgenic animal according to the invention can be produced by introducing a transgene and/or a vector, e.g. an expression vector which comprises the transgene into the germline or a germline cell respectively and/or into a somatic cell of the non-human animal. For example embryonic target cells at various developmental stages can be used to introduce the transgene of the invention. Different methods can be applied depending on the stage of development of the embryonic target cell(s). Some examples are:

1. Microinjection of zygotes is a preferred method for incorporating a transgene into an animal's genome in the course of practicing the invention. Microinjection involves the isolation of embryos at the single cell stage. Therefore a zygote, a fertilized ovum that has not undergone pronuclei fusion or subsequent cell division, is the preferred target cell for microinjection of transgenic DNA (DNA of the transgene). The murine male pronucleus reaches a size of approximately 20 micrometers in diameter, a feature which allows for the reproducible injection of 1-2 picoliters of a solution containing transgenic DNA. The use of a zygote for introduction of a transgene has the advantage that, in most cases, the injected transgenic DNA will be incorporated into the host animal's genome before the first cell division (Brinster, et al. (1985) Proc. Natl. Acad. Sci. 82: 4438-4442). As a consequence, all cells of the resultant transgenic animals (founder animals) stably carry an incorporated transgene at a particular genetic locus, referred to as a transgenic allele. The transgenic allele demonstrates Mendelian inheritance: half of the offspring resulting from the cross of a transgenic animal with a non-transgenic animal will inherit the transgenic allele, in accordance with Mendel's rules of random assortment.

Commonly used procedures for manipulation of embryo and for microinjection of transgenic DNA are described in detail in „Transgenic Animal Technology - A Laboratory Handbook“ edited by Carl A. Pinkert, Academic Press, Inc. (1994).

2. Viral integration can also be used to introduce a transgene according to the invention into an animal. The developing embryos are cultured in vitro to the developmental stage known as a blastocyst. At this time, the blastomeres may be
- 5 infected with vectors containing the transgene (transgenic DNA/DNA-constructs), for example an appropriate viral or retroviral vector can be used for this purpose (Jaenich, R. (1976) *Proc. Natl. Sci. (USA)* 73: 1260-1264). Transformation or infection of the blastomeres can be enhanced by enzymatic removal of the zona pellucida (Hogan, et al. (1986) *Manipulating the Mouse Embryo*, Cold Spring Harbor
- 10 Press, Cold Spring Harbor, N.Y.). If the transgene is introduced into blastomeres via viral vectors, such vectors are typically replication-defective but they remain competent for the integration of transgenic DNA sequences which are linked to vector sequences, into the host animal's genome (Jahner et al. (1985) *Proc. Natl. Acad. Sci. (USA)* 82: 6927-6931; Van der Putten et al. (1985) *Proc. Natl. Acad. Sci. (USA)* 82: 6148-6152). Transfection is easily and efficiently obtained by culture of
- 15 blastomeres on a mono-layer of cells producing the transgene-containing vector (Van der Putten et al. (1985) *Proc. Natl. Acad. Sci. (USA)* 82: 6148-6152; Stewart et al. (1987) *EMBO* 6: 383-388).
- 20 Alternatively, infection may be performed at a later stage, such as a blastocoele (Jahner, D. et al. (1982) *Nature* 298: 623-628). In any event, most transgenic founder animals produced by retroviral or viral integration into only a subset of all the cells that form the transgenic founder animal. Moreover, multiple (retro)viral integration events may occur in a single founder animal, generating multiple
- 25 transgenic alleles which will segregate in future generations of offspring. Introduction of a transgene into germline cells by this method is possible but probably occurs at a low frequency (Jahner, D. et al. (1982) *Nature* 298:623-628). However, once a transgene has been introduced into germline cells by this method, offspring may be produced in which the transgenic allele is present in all of the
- 30 animal's cells, i.e., in both somatic and germline cells.

3. Embryonic stem (ES) cells can also serve as target cells for introduction of a transgene according to the invention into animals. ES cells are obtained from pre-implantation embryos that can be cultured in vitro (Evens, M.J. et al. (1981) *Nature* 292: 154-156; Bradley, M.O. et al. (1984) *Nature* 309: 255-258; Gossler, et al. (1986) *Proc. Natl. Acad. Sci. (USA)* 83: 9065-9069; Robertson et al. (1986) *Nature* 322: 455-448; Robertson, E.J., in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, E.J., ed., IRL Press, Oxford (1987), pages 71-112). ES cells, which are commercially available (from, e.g., Genome Systems, Inc., St. Louis, MO), can be transformed with one or more transgenes by established methods (Lovell-Badge, R.H., in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, E.J., ed., IRL Press, Oxford (1987), pages 153-182). Transformed ES cells can be combined with an animal blastocyst, whereafter the ES cells colonize the embryo and contribute to the germline of the resulting animal, which is a chimera (composed of cells derived from two or more animals) (Jaenisch, R. (1988) *Science* 240: 1468-1474; Bradley, A., in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, E.J., ed., IRL Press, Oxford (1987), pages 113-151). Again, once a transgene has been introduced into germline cells by this method, offspring may be produced in which the transgenic allele is present in all of the animal's cells, i.e., in both somatic and germline cells.
- 20 However it occurs, the initial introduction of a transgene is a non-Mendelian event. However, a transgene of the invention may be stably integrated into germ line cells and transmitted to offspring of the transgenic animal as Mendelian loci. Germline integration is essential for the production of transgenic animals that can transmit the genetic information to their progeny in a Mendelian fashion and in order to utilize these transgenic animals as perpetual animal models.
- 25 Other transgenic techniques result in mosaic transgenic animals, in which some cells carry the transgene and other cells do not. In mosaic transgenic animals in which germ line cells do not carry the transgene, transmission of the transgene to offspring does not occur. Nevertheless, mosaic transgenic animals are capable of

demonstrating phenotypes associated with the transgene and may be used for affinity purification of particular transcription complexes, TAFs and TAF interacting factors. The invention relates to mosaic transgenic animals that contain the described transgene of the invention.

5

The invention further relates to transgenically introduced mutations, this comprises null („knock-out“) alleles in which the DNA sequence encoding for the species-specific TBP is deleted and/or substituted by a genetically altered TBP sequence (e.g. a transgene according to the invention) of the same or a different species under the control of the promoter(s)/enhancer(s) of choice.

10

The invention relates to a transgenic animal into which a transgene comprising the DNA encoding epitope-tagged TBP, if necessary, in the context of a constitutive or an inducible promoter has been introduced. In particular, the invention relates to a transgenic animal into which a transgene that comprises the DNA encoding HA and His epitope-tagged hTBP and the DNA sequence of the EF promoter has been introduced. In another special embodiment the invention relates to a transgenic animal into which a transgene that comprises the DNA encoding HA and His epitope-tagged hTBP and the DNA sequence of the MT promoter has been introduced. In a special embodiment of the invention the transgenic animal contains the transgene with the sequence SEQ ID NO. 13. Another transgenic animal of the invention contains the transgene with the sequence SEQ ID NO. 14. Another transgenic animal of the invention contains the transgene with the sequence SEQ ID NO. 15. In particular the invention relates to a transgenic animal which has stably integrated into its genome the transgene, for example the DNA sequence SEQ ID NO. 13, SEQ ID No. 14 and/or SEQ ID NO. 15.

20

25

30

Offspring that have inherited the transgene, can be distinguished from littermates that have not inherited the transgene by analysis of genetic material from the offspring for the presence of biomolecules that comprise unique sequences corresponding to sequences of, or encoded by, the transgene. Therefore for

example, biological fluids that contain the polypeptides (e.g. the epitope-tagged TBP) uniquely encoded by a transgene according to the invention may be immunoassayed for the presence of the polypeptide encoded by the transgene, e.g. the epitope-tagged TBP. A more simple and reliable means of identifying transgenic offspring comprises obtaining a tissue sample from an extremity of an animal, e.g., a tail and analyzing the sample for the presence of nucleic acid sequences corresponding to the DNA sequence of a unique portion or portions of the transgene of the invention. The presence of such nucleic acid sequence may be determined by, e.g., hybridization („Southern", „Northern") analysis with DNA sequences corresponding to unique portions of the transgene, analysis of the products of PCR reactions using DNA sequences in a sample as substrates and oligonucleotides derived from the transgene's DNA sequence, etc.

Therefore, the invention also relates to tests wherein possible first generation transgenic animals (G_0) as well as all further generation transgenic animals (G_1 , G_2 , G_3 , G_4 , ...) or animals of transgenic animal lines can be tested for presence of the transgene, e.g. with standard PCR reactions. For this purpose genomic DNA can be extracted from animal tissue, for example from tail tissue after Proteinase K and RNase treatment. For such PCR reactions the sequence of the primers should correspond to parts of the sequence of the transgene, e.g. within the promoter regions, the DNA region(s) encoding epitope-tag(s) and/or DNA regions unique for the particular TBP construct. With such PCR reactions in mice for example in about 25% of the injected eggs the corresponding PCR product can be detected - i.e. about 25 % of the mice produce positive offspring.

Another method to verify whether or not the animals carry the transgene relates to test for the presence of transgenic mRNA in possible transgenic animals/transgenic animal lines. Initial testing can for example be carried out with S1 analysis, which is very sensitive to small levels of mRNA (Berk, A.J. and Sharp, P.A. (1977) Cell 12, 721). For this purpose, labeled antisense oligonucleotides are hybridized with total RNA that has been isolated from tissue of the animal. Subsequent treatment with S1

nuclease digests all single-stranded nucleic acids, DNA and RNA. Double-stranded DNA or DNA-RNA hybrids are left intact. If any transgene mRNA is present, it hybridizes with the antisense oligonucleotide and thus "protects" it from S1 nuclease digestion.

5

The invention comprises that the presence of transgenic mRNA is detected in tissue preparations, e.g. in preparations of total liver mRNA by S1 protection assays.

Antisense oligonucleotides can be synthesized which are complementary to a part of the transgenic mRNA (e.g. the mRNA corresponding to the DNA sequence of epitope-tagged TBP). Oligonucleotides are labeled, e.g. 5'-end-labeled, for example radioactively with ^{32}S , ^{33}P , ^{35}P , ^3H or ^{14}C or with fluorescence markers or other types of markers, like biotin or digoxigenin. Labeled oligonucleotides are mixed with mRNA from the transgenic mouse under selective hybridizing conditions according to standard protocols (Sambrook et al., 1989). The mixture is then treated with S1

15 nuclease. A short region of non-matching sequence, e.g. at the 3'-end of the oligonucleotide should always be digested, and provides an internal control to show that the S1 nuclease indeed digests all available single-stranded nucleic acid. In the presence of transgenic mRNA, the labeled oligonucleotide should be protected from digestion and its presence and size could be easily determined e.g. by sequencing-style denaturing gel (e.g. by 8M urea PAGE) electrophoresis. The presence of

20 undigested, labeled oligonucleotide can then be detected e.g. by exposing the gel to a film. Absence of transgenic mRNA would give no band, as the unprotected oligonucleotide would be digested by the S1 nuclease. The invention includes that different tissues and cells relating to different cell types were prepared from the

25 animals and tested for the presence of transgenic mRNA.

Another embodiment of the invention relates to test transgenic animals with Northern analysis. Most preferably transgenic animals that were found to be positive for transgenic mRNA by PCR or S1 nuclease mapping were further tested with

30 Northern analysis (McMaster, G.K. and Carmichael (1977) Proc. Nat. Acad. Sci. 74: 4835). For this purpose total RNA from different tissues and/or cell types, can be

- isolated and size separated e.g. under denaturing conditions on an agarose gel. The RNA can then be bound for example onto a membrane or filter using e.g. capillary transfer and UV cross-linked. The bound RNA can be probed using conventional Northern blot conditions with a labeled probe corresponding to the coding region of the transgene or parts thereof, e.g. with a DNA-probe corresponding to the 5'end of the transgene according to SEQ ID NO. 13. Such DNA-probes have preferably a length of about 20 to 1000 basepairs (bp). Most preferably they have a length of about 100, 200, 300, 400 and 500 bp. If necessary, excess of the labeled probe is rinsed away and the membrane is exposed to film.
- The probes can be labeled as described above for oligonucleotides. Oligonucleotides may sometimes also be used for those Northern blot experiments.

- A transgenic animal, preferably such transgenic animal which has been positive in any other molecular biological test, are being tested for presence of the transgenic TBP protein with specific immunoreaction e.g. with Western blotting or ELISA. Therefore from tissue or particular cells of the animals, preferably from liver tissue or any other soft tissue nuclei can be isolated by standard procedures, e.g. on an ultracentrifuged sucrose gradient. From such nuclei, total nuclear protein can be collected, e.g. by treating the nuclei with high salt conditions (e.g. 400mM KCl) and non-ionic surfactant (e.g. NP-40). Afterwards nuclear protein can be size separated e.g. with appropriate denaturing gel electrophoresis, preferably by SDS-PAGE. Such gels can be transferred, e.g. electro-transferred on a solid surface, e.g. on membranes or filters, preferably onto nitrocellulose membrane. Membranes or filters can then be preblocked and probed with suitable antibodies according to standard protocols (Sambrook et al. "Molecular Cloning" Second Edition (1989), Cold Spring Harbor Laboratory Press).

- For such purpose polyclonal and/or monoclonal antibodies can be applied, e.g. generated in mouse, rabbit, rat, sheep, goat, horse, birds etc. The detection could be performed directly with antibodies that recognize the TBP fusion protein and which are coupled to an enzyme or a marker, e.g. alkaline phosphatase or

fluorescence marker or biotin or digoxigenin or radiolabel. The detection can also be performed indirectly by using a second antibody which recognizes a conserved region of the primary antibody, for example when the first antibody is generated in mouse the second antibody has to be an anti-mouse antibody generated for
 5 example in sheep. Such second antibody can also be coupled to an enzyme or other markers for detection.

The invention further relates to the use of the transgene or a part thereof or the encoded fusion protein or a part thereof for the production of antibodies which bind
 10 the epitope-tagged TBP encoded by the transgene, e.g. to the preparation of monoclonal or polyclonal antibodies.

Antibodies with respect to the invention are antibodies that recognize one or more epitope(s) of the TBP fusion protein. Such antibodies could be directed against
 15 individual epitope(s) belonging to the TBP and/or could be directed against the epitope-tag(s). One embodiment of the invention relates to an antibody which recognizes the amino-terminal region of the TBP fusion protein. Another embodiment of the invention relates to an antibody which recognize the carboxy-terminal region of the TBP fusion protein. Another embodiment of the invention
 20 relates to an antibody which recognize the epitope neighbouring the part of the amino acid sequence where the TBP and the epitope-tag(s) and/or where the two epitope-tags are connected together.

One embodiment of the invention relates to an antibody that recognizes an epitope
 25 of the fusion protein that consists of hTBP and two epitope-tags. In particular, the antibody recognizes an epitope of the fusion protein consisting of His- and HA- tag and hTBP. Most preferably the antibody recognizes an epitope(s) at the amino-terminal end of HA- and His-tagged TBP. A special embodiment of the invention relates to an antibody that recognizes the amino acid sequence SEQ ID
 30 NO. 16 or an epitope of the correctly folded fusion protein having the amino acid sequence SEQ ID NO. 16 or a part thereof. The invention relates to an antibody

(anti-TBP antibodies) that recognizes the amino acid sequence SEQ ID NO. 17 or the correctly folded epitope thereof, preferably in the context of a TBP fusion protein.

- 5 Antibodies of the invention can be polyclonal or monoclonal. An antibody can be generated in all species of non-human animals, preferably from mouse, rat, rabbit, sheep, goat, horse, birds (e.g. from their eggs). Such antibody can be generated according to standard protocols (Hurlow and Lane "Antibodies: A Laboratory Manual" (1988) Cold Spring Harbor Press). Antibodies can, if necessary be affinity
10 purified using the original immunization peptide (epitope). A special embodiment of the invention relates to polyclonal antibodies produced in rabbit which are generated by immunization of the rabbit with the peptide having sequence SEQ ID NO. 17 (e.g. coupled to a suitable carrier-protein). The polyclonal antibody (anti-TBP antibody) recognizes the hTBP fusion protein (HA- and His-tag).

15

An antibody relating to the invention can be applied for Western blot analysis as well as for the affinity purification of TBP fusion protein and of higher order transcription complexes which are associated to the TBP fusion protein. Higher order complexes comprise TAFs which are associated with the TBP fusion protein and TAF-interacting factors.

20

The invention relates to the use of the transgenic animal. A transgenic animal according to the invention can be used for affinity co-purification of higher order transcription complexes, TAFs and TAF-interacting factors from transgenic animal
25 tissue and/or cultured cells of the transgenic animal. Such higher order transcription complexes can be isolated and purified from a variety of different tissues and/or cell types. Preferably, nuclear preparations from such tissue/cell types are performed, most preferably from homogenized cells according to standard protocols (Dignam et al. (1983) Nuc. Acids Res. 11: 1575; Lichtenstein et al. (1987) Cell 51: 963-973; Gorsky et al. (1986) Cell 47: 767-776). If necessary, nuclear proteins can then be
30 accumulated by standard methods. Therefore, the invention relates also to methods

of affinity co-purify higher or to transcription complexes, TAFs and TAF-interacting factors from transgenic animals.

For example, the invention relates to the affinity-purification of higher-order
 5 transcription complexes, TAFs and TAF-interacting factors by using epitope-specific
 antibodies or charged (positive or negative charged) materials. For example the
 antibodies already described in detail can be used for this purpose. One of the most
 preferable co-purification methods for higher order transcription complexes which
 comprise HA and His epitope-tagged TBPs, preferably hTBPs, include affinity
 10 purification using Ni^{2+} and/or anti-HA-antibodies (e.g. commercially available
 antibodies) and/or antibodies recognizing the epitope according to sequence SEQ
 ID NO. 17. Such antibodies or Ni^{2+} might be coupled to a suitable column material
 so that the affinity purification can be performed by using e.g. Ni^{2+} -columns or
 columns with specific antibodies, e.g. with anti-HA antibodies or anti-TBP
 15 antibodies.

In another embodiment the invention relates to a TBP fusion protein, the epitope-
 tagged TBP. Preferably the TBP fusion protein is expressed in a transgenic animal.
 In particular the invention relates to a hTBP fusion protein. A TBP fusion protein can
 20 be isolated and purified from a transgenic animal. One embodiment of the invention
 is a HA and His tagged TBP protein, in particular the HA and His tagged hTBP
 protein. An other embodiment of the invention is the protein that has the amino acid
 sequence SEQ ID NO. 16. In another embodiment of the invention the TBP fusion
 protein comprises one or more cleavage side(s) for a proteinase/peptidase, e.g. a
 25 thrombin cleavage site. Preferably the cleavage sites within the amino acid
 sequence of the fusion protein are located between the epitope-tag(s) and the TBP-
 protein.

The invention relates to a method of preparing a TBP fusion protein, wherein a
 30 transgene according to the invention is expressed in a suitable host cell, who is
 preferably part of a transgenic animal.

Transgenic mRNA and/or a TBP fusion protein encoded by the transgene and/or higher order transcription complexes associated with the TBP fusion protein can be isolated from different types of tissue and/or cell types that are found in the
 5 transgenic animal e.g. brain, heart, kidney, liver, lung, nervous system, muscle, glands, bone marrow, cells belonging to the immunosystem, skin etc.

The invention relates to the use of the TBP fusion protein, in particular for the isolation of higher order transcription complexes from the transgenic animal and to
 10 the characterisation of isolated higher order transcription complexes obtained from different species, from different tissues and/or different cell-types. Therefore, the TBP fusion protein and a transgenic animal according to the invention can be used for the isolation and characterization of individual proteins such as TAFs and TAF-interacting factors which are associated in the different higher order complexes. For
 15 example, the proteins associated in a particular higher order transcription complex can be dissociated and separated so that individual TAFs and TAF-associated factors can be identified. The composition of TAFs and TAF-interacting factors are different to at least some extent in the different higher order complexes depending on the tissue type and/or the cell type and/or the developmental stage and/or the
 20 transgene which is expressed.



TAFs and TAF-interacting factors, in particular tissue-specific factors which have already been characterized and for which antibodies already exist can be quickly identified and assessed for degree of association with the transcription complex. An



25 example of this would be the Bob-1/OCA-B factor, which is thought to be a tissue-specific co-activator responsible for B-cell restricted activation (Gstaiger, M. et al. (1996) EMBO 15, 2781-2790). Although without intrinsic DNA-binding capacity and a requirement for the nearly ubiquitous Oct factors, the tissue-restricted appearance of this factor confers B-cell specific transcriptional activation through a protein-
 30 protein mechanism.




Further, novel TAFs and TAF-interacting factors, in particular tissue- and/or cell type-specific and/or developmental (stage) specific and/or cell cycle specific TAFs and TAF-interacting factors, can be identified in a higher order transcription complex. As mentioned above, there is a wealth of data which strongly suggests that

5 many enhancer-binding tissue-specific factors are able to exert influence and, thus, tissue specificity on a gene's transcriptional activity. Therefore "unique", tissue-specific, cell-type specific, cell-cycle specific, developmental stage specific factors might be identified, which regulate the specific expression of genes.

10 This "universal" transgenic system offers furthermore a powerful tool to investigate the degree to which such TAFs and TAF-interacting factors (e.g. coactivators), associate with TBP and/or TAFs and the transcription complex in a range of tissues and cell types.

15 The invention also relates to a method for the identification and characterization of different higher order transcription complexes, wherein epitope-tagged TBP to which higher order transcription complexes are associated is isolated from a transgenic animal. A method of characterizing the composition of different higher order transcription complexes can for example comprise a) the introduction of a transgene

20 according to the invention into a non-human animal, b) the isolation of the epitope-tagged TBP from different animal tissue and/or different cell types of the animal, optionally at different developmental stages of the animal and c) the determination of the composition of the higher order transcription complexes.



25 One method of isolating a higher order transcription complex from a transgenic non human animal comprises the affinity purification by using at least one of the epitopes tagged to TBP. Preferably the higher order transcription complex and the TAFs and TAF-interacting factors associated in this complex are co-purified, when the epitope-tagged TBP is isolated. For example, a higher order transcription

30 complex can be isolated, when epitope-tagged TBP is purified by binding of one of

ist epitopes, preferably an epitope-tag to a material to which the epitope specifically binds. This material can for example be a Ni^{2+} -column (to which a His-epitope binds) or antibodies, e.g. anti-HA antibodies (bind to a HA epitope) or antibodies which bind to an epitope of epitope-tagged TBP that has the sequence SEQ ID NO. 17 or
 5 a part thereof (e.g. epitope of sequence SEQ ID NO. 17).

The invention also relates to a method of identifying a new and/or a specific TAF and/or TAF-interacting factor. Preferably, a higher order transcription complex is isolated from a transgenic animal according to the invention. Such a
 10 method may for example comprise a) the introduction of a transgene according to the invention into a non-human animal, b) the isolation of epitope-tagged TBP from a particular animal tissue and/or a particular cell type of the animal, optionally at a particular developmental stage of the animal and c) dissociation and separation of a TAF and/or a TAF-interacting factor associated with the epitope-tagged TBP in the
 15 higher order transcription complex and d) if necessary determination of the aminoacid sequence of the TAF and/or the TAF-interacting factor.

In addition, to corroborating already known mechanisms, the transgenic model has an advantage over current cell culture systems in finding and characterizing new
 20 TAFs or TAF-associated factors. As mentioned above, the TAF proteins that have been affinity co-purified to date have come from studies in HeLa and yeast. TAF cDNAs for other organisms have also been found, but with time consuming interaction library screening methods. The "whole organism" aspect of the transgenic model makes this universal transgenic system especially responsive to
 25 the recognition of novel tissue-specific activation elements by remaining very "close" to the true in vivo process of transcription.

It is clear that such a tagged-TBP-expressing transgenic animal, e.g. mouse would be able to contribute to the area of drug development. Considerable pharmaceutical
 30 interest can be attributed to any 'unique', tissue-specific, cell-type specific, cell-cycle specific, developmental stage specific factors (TAFs, TAF-interacting factors) of a

given gene's transcription complex, especially if it is a disease-related gene. The identification and characterization of "key" TAFs and TAF-interacting factors which are involved in transcriptional control of one or a few disease-related genes is a sound strategy to develop therapeutic compounds which alleviate such genetic diseases. Once such a TAF or TAF-interacting factor has been characterized and cloned, any number of screening procedures can be undertaken to identify molecular species/substances which specifically interact with the TAF or TAF-associated factor. A pharmaceutically useful species/substance would be one which enhances or represses the TAF's or TAF-interacting factor's natural activity in vitro and/or in vivo, thus allowing therapeutic manipulation of a related gene.

Identified TAFs and TAF-interacting factors might then also be applied as tools in biochemistry and in molecular biology, for example such proteins from transgenic non-human animals might then be used as probes for the isolation of the corresponding human TAFs and TAF-interacting factors or their cDNAs. Human TAFs and TAF-interacting factors might then also be applied to screen for highly specific new drugs.

The present invention is described in further detail in the following non-limiting examples.

Example 1: Construction of transgenic animals.

Potential animal sources:

Animals suitable for transgenic experiments were obtained from standard commercial sources, Charles River (Wilmington, MA), Taconic (Germantown, NY), and The Jackson Laboratory (Bar Harbor, Maine). B6SJL/F1 mice were used for embryo retrieval and transfer. B6SJL/F1 males can be used for mating and vasectomized Swiss Webster studs can be used to stimulate pseudopregnancy.

Transgenic mice:

Female mice six weeks of age are induced to superovulate with a 5 IU injection (0.1 cc, intraperitoneal) of pregnant mare serum gonadotropin (PMSG; e.g. Sigma, Saint Louis, Missouri, USA) followed 48 hours later by a 5 IU injection (0.1 cc,

- 5 intraperitoneal) of human chorionic gonadotropin (hCG; e.g. Sigma). Females are placed with males immediately after hCG injection. Twenty-one hours after hCG, the mated females are sacrificed by CO₂ asphyxiation or cervical dislocation and embryos are recovered from excised oviducts and placed in M2 media (e.g. Sigma). Surrounding cumulus cells are removed with hyaluronidase (1 mg/ml). Pronuclear
- 10 embryos are then washed and placed in M16 media (e.g. Sigma) and then put in a 37° C incubator with a humidified atmosphere at 5% CO₂, O₂, and 90% N₂ until the time of injection.

15 Example 2: Preparation of constructs for transfections and microinjections

DNA clones for microinjection were cleaved with appropriate enzymes, DNA clones comprising the MT-hTBP transgene (double-tagged, according to the sequence in table 3) with Cla I and BamHI or EF-hTBP transgene (double-tagged, according to the sequence in table 2) with Eco RI/Eco RI and the appropriate size DNA fragments

- 20 electrophoresed on 1% agarose gels in TBE buffer (Sambrook et al. (1989)). The DNA bands are visualized by staining with ethidium bromide, excised, and placed in dialysis bags containing 0.3 M sodium acetate, pH 7.0. DNA is electroeluted into the dialysis bags, extracted with phenol-chloroform (1:1), and precipitated by two
- 25 volumes of ethanol. The DNA is redissolved in TE buffer (10 mM Tris, pH 7.4 and 1 mM EDTA) and purified on a DEAE sephacel (e.g. Pharmacia, Uppsala, Sweden) column. The column is first primed with 0.5 ml of high salt buffer (1.5 M NaCl, 10 mM Tris, pH 7.4, and 1 mM EDTA) followed by washing with 3 ml of low salt buffer (0.15M NaCl, 10mM Tris pH 8.0, and 1mM EDTA). The DNA solutions are adjusted
- 30 in salt to 0.15M NaCl then passed through the column to bind DNA to the column matrix. After three washes with 3 ml of low salt buffer, the DNA is eluted in aliquots

of 4 x 0.3 ml of high salt buffer and precipitated by two volumes of ethanol. The fractions were pooled by dissolving in 200µl of TE, phenol:chloroform extracted once, chloroform extracted twice, then the DNA was precipitated with ethanol overnight. The DNA was resuspended in microinjection buffer TE (10mM Tris pH 7.4, 0.1mM EDTA) and the DNA concentration was adjusted to 2ng/µl and visualized against known DNA standards by electrophoresing on an agarose gel.

Microinjection:

DEAE purified transgene DNA (MT-hTBP or EF-hTBP) was dissolved in microinjection buffer at 2ng/µl. Microneedles and holding pipettes were pulled on a Flaming Brown micropipette puller e.g. Model P87 (Sutter Inst. Co.). Holding pipettes were then broken and fire polished to on a deFonbrune-type microforge (e.g. Technical Product Inst. Inc.). Pipettes were mounted micromanipulators (e.g. on Leitz) which were attached to a Zeiss Axiovert® 135 microscope. The air-filled injection pipette (e.g. Medical System Corp.) was filled with DNA solution through the tip. Embryos in groups of 40-50 were placed in 200µl of M2 media under silicone oil for micromanipulation. The embryo was oriented and held with the holding pipette and then the injection pipette was inserted into the pronucleus closest to the injection pipette. The injection was monitored by the swelling of the pronucleus. Following injection, the group of embryos was placed in M16 media until transfer to recipient females.

Example 3: Embryo transfer by microinjection.

Randomly cycling adult female mice are paired with vasectomized males. Swiss Webster or other comparable strains can be used for this purpose. Recipient females are mated at the same time as donor females. At the time of embryo transfer, the recipient females are anesthetized with an intraperitoneal injection of 0.015 ml of 2.5% avertin per gram of body weight. The oviducts are exposed by a single midline dorsal incision. An incision is then made through the body wall

directly over the oviduct. The ovarian bursa is then torn with watchmakers forceps. Embryos to be transferred are placed in M16 and then in the tip of a transfer pipette (about 10-12 embryos). The pipette tip is inserted into the infundibulum and the embryos transferred. After the transfer, the incision is closed by two sutures and the skin stapled. The recipient recovered for three hours on a warming tray then was placed in the colony for delivery.

A total of 469 pronuclear embryos were microinjected for MT-hTBP and 170 pups were born and for EF-hTBP a total 407 pronuclear embryos yield 76 pups.

Example 4: Detection of transgenic DNA in founder mice.

4a) DNA Extraction:

- 15 Genomic DNA of possible founder mice was extracted from a small piece of tail tissue (ca. 1 cm) cut from 2-4 week old offspring. Tail sections were incubated overnight on a shaker in 500-750 μ l Tail Buffer at 54°C (Tail Buffer: 10mM Tris pH 7.5, 100mM NaCl, 10mM EDTA, 0.5% SDS, 30 μ g/ml Proteinase K). To each tail sample was subsequently added an equivolume of phenol/chloroform/isoamyl alcohol (25:24:1). Samples were shaken gently by hand or automatically on a Vortex
- 20 micer with low setting. Samples were all centrifuged in tabletop centrifuge for 10 minutes. Aqueous phase was transferred to 5ml polypropylene tube (e.g. falcon tube®). An equivolume of ethanol was slowly added (dropwise) to the tube to allow the DNA to gradually precipitate. A second volume of ethanol was added forcefully
- 25 to mix the contents of the tube. Tubes were then inverted several times to ensure mixture. Genomic DNA was then spooled around a flat pipette-tip (sequencing-gel tip) and transferred to an individual well of a microtiter dish. DNA was then air dried in dish for 4 hours. 200 μ l of 1x TE was then added to each well (1x TE: 10mM Tris pH 7.4, 1mM EDTA). Genomic DNA was then allowed to dissolve for 15-30 minutes
- 30 at room temperature, then mixed gently by carefully pipetting up and down. Microtiter plates were often stored at -20°C before PCR testing. Prior to testing, 10 μ l

were removed and digested with EcoRI (e.g. 10µl genomic DNA, 2µl 10x EcoRI buffer, 7µl H₂O, 1µl EcoRI; enzyme and buffer from Boehringer-Mannheim, Mannheim, Germany).

5 4b) PCR Reactions:

Digested genomic DNA was diluted 1:3 with water. Samples were then heated to 100°C on a heating block for 10 minutes. 2µl was then used for PCR reactions. Reactions were carried out in a 50µl volume consisting of 2µl genomic DNA, 1x reaction buffer, 1.5mM MgCl, 0.8µM forward oligonucleotide primer, 0.8µM reverse oligonucleotide primer, 8µl of nucleotide mixture containing 0.2mM of each nucleotide (dATP, dCTP, dTTP, and dGTP), and 2.5 units of Taq polymerase. PCR was performed using e.g. AmpliTaq® enzyme and buffer from Perkin Elmer (Perkin Elmer Norwalk, Connecticut, USA).

- 15 Detection of the MT-hTBP transgene was accomplished using the forward primer oligonucleotide (sense primer) 5' GGAGCA ACC GCC TGC TGG GTG C 3' (SEQ ID NO. 5) and the reverse primer oligonucleotide (antisense primer) 5' CCT GTG TTG CCT GCT GGG ACG 3' (SEQ ID NO. 6).

- 20 Detection of the EF-hTBP transgene was accomplished with the forward oligonucleotide primer 5' GGA GAC TGA AGT TAG GCC AGC 3' (SEQ ID NO. 7). The same reverse primer as in the MT-hTBP detection was used (5' CCT GTG TTG CCT GCT GGG ACG 3').

- 25 Temperature cycling was carried forth using a robotic temperature cycler (e.g. from Stratagene, La Jolla, CA, USA).

The cycling temperatures were:

cycle 1:	94°C 5min	60°C 3min	72°C 2min;
cycle 2-25:	94°C 1min	60°C 2min	72°C 3min;
cycle 26:	94°C 1min	60°C 2min	72°C 5min.

or

cycle 1-40: 95°C 2 min 55°C 1 min 72°C 1 min.

5

The presence of amplified product was detected by running standard agarose gel electrophoresis (e.g. 1.2-1,5% agarose) and staining the gel with ethidium bromide for visualization of DNA with UV light.

- 10 Positive MT-hTBP samples were distinguished by the presence of a about 580 bp amplified DNA product. Positive EF-hTBP samples produced a about 500bp amplified DNA fragment.

- 15 The DNA-analysis from 170 pups for MT-hTBP by PCR-reaction indicated that 35 genomic DNA-samples were positive for the transgene. The analysis of the genomic DNA of the 76 EF-hTBP pups by PCR analysis indicated that the EF-hTBP transgene was contained in 9 mice.

4d) Transgenic model expansion:

- 20 DNA positive G₀ founders were bred to non-transgenic B6SJL mates and the resulting litters were PCR genotyped. G₁ offspring were used for continued breeding and maintenance of the individual lines and for further mRNA and protein expression analysis.

25

Example 5: Detection of transgenic mRNA by S1 nuclease protection assay:

5a) Specifics:

Oligonucleotides (oligo) complementary to the 5' end and the 3' end of the transgenic transcript were produced:

30

Sequence of 5'oligo (sense primer) (SEQ ID NO. 8):

5'GCGGCACCAGGCCGCTGCTGTGATGATGATGATGATGGCTGCTGCCCATGA
CTGCGTAATGCGGTCATGACGCTTT 3'

5 Sequence of 3'oligo (antisense primer) (SEQ ID NO. 9):

5'GAAGGGGGTGGGGGAGGCAAGGGTACATGAGAGCCATTACGTCGTCTTCCT
GAATCCCTTTAGCCGCTTTGCTCG 3'

Underlined regions are non-hybridizing sequence. 40ng of oligo were 5'-labeled with
10 (³²P gamma)ATP (5000cpm/mM) using T4 polynucleotide kinase and buffer e.g.
from Boehringer-Mannheim (Mannheim). The reaction was performed in 50µl
reaction volume at 30°C for 1 hour. Labeled oligo was isolated from unincorporated
(³²P gamma)ATP using size exclusion chromatography (e.g. Push-Columns®,
Stratagene, La Jolla, CA, USA).

15

5b) Promoter induction:

Prior to RNA analysis, it was necessary to induce the MT-hTBP mice, since the
promoter is activated in the presence of Zn²⁺. On each MT-hTBP mouse, two
interperitoneal injections of ZnSO₄ in H₂O were made at 18 hours and again at 4
20 hours prior to liver removal (dose was 0.1mg ZnSO₄/10g mouse weight).

5c) RNA Extraction:

Total RNA was extracted from 1-5g of tissue using commercially available Trizol
reagent (e.g. Gibco-BRL, Paisly, UK). Tissue was homogenized with an Ultra-Turrax
25 in 5ml of Trizol solution in a 12ml polypropylene tube (e.g. falcon tube®). 1ml
chloroform was added, and tubes were capped and shaken vigorously by hand for
15 seconds. Solutions were incubated at RT for 3 min, then centrifuged at 12,000xg
for 15 mins at 4°C in Sorvall SS-34 rotor. Supernatant was transferred to a new
tube; 2.5ml isopropanol was added and mixed. Incubation at RT for 10 min followed.
30 The samples were centrifuged with 12,000xg for 10 min at 4°C. Supernatant was
removed and RNA pellet were washed with 5ml 75% ethanol. Samples were mixed

and centrifuged at 7,500xg for 10 min. RNA pellets were resuspended in RNase-free water and subsequently quantitated by measuring the absorbance at 260nm.

5d) S1 Nuclease protection:

- 5 Uniform amounts of RNA (10-20µg) were brought up to 100µl with RNase-free water. 50,000-150,000 dpm of labeled oligo was added (0.1-1ng, depending on labeling efficiency). RNA/oligo mixture was precipitated with 0.3M sodium acetate and ethanol. The RNA/oligo pellet was washed and air dried. 23µl hybridization solution was added (80% formamide, 10mM PIPES pH 6.4 (Sambrook et al. (1989)),
- 10 1mM EDTA, 0.05% SDS). Reaction was mixed and denatured at 65°C for 20 min. and then 2µl of 5M NaCl were added to each sample at 65°C. Samples were incubated 1 additional hour at 65°C in a H₂O bath, then the temperature of the bath was reset to 37°C. Gradual temperature decrease from 65°C to 37°C (overnight) facilitated the specific oligo-mRNA hybridization. The next day, 300µl S1 buffer were
- 15 added to each sample (S1 buffer: 167U/ml S1 nuclease (e.g. Gibco BRL, Paisly, UK), 0.3M NaCl, 30mM NaOAc (pH 4.5), 3mM ZnSO₄). Samples were incubated at room temperature for 1 hour, then 1 ml ethanol was added (cold) to precipitate all nucleic acid. The pellet was centrifuged and washed and dried briefly in a Speed-Vac. Pellet was then resuspended in 12µl S1 loading buffer (85% formamide, 0.01%
- 20 bromphenol blue, 0.01% xylene cyanol, 1x TBE). Samples were heated to 70°C for 5 min before being loaded and size separated using denaturing (6M urea) thin polyacrylamide gel electrophoresis. Autoradiography of the dried gel facilitated detection of protected, undigested bands.
- 25 From 35 founders for MT-hTBP, 7 founders showed detectable mRNA levels as analyzed by S1 protection assay. For EF-hTBP, 3 founders showed detectable mRNA from S1 analysis.
- 30 Example 6: Northern blot detection of transgenic mRNA.

6a) Synthesis of the hybridization probe by PCR amplification of TBP and labeling of the TBP probe:

Oligonucleotides/primers bracketing a 498bp fragment of the double-tagged hTBP construct were designed and synthesized.

5

The forward oligo began 10 bases downstream from "AUG" start site (ATG site in Table 1). The sequence of the sense primer was:

SEQ ID NO. 10: 5' CCCTATGACGTCCCGGATTACG 3'.

10 The reverse primer ended at 507 bp downstream of the "AUG" start site (ATG site in Table 1). The sequence of the antisense primer was:

SEQ ID NO. 11: 5' GTGGAGTGGTGCCCGCAAGGG 3'.

15 PCR reactions were carried out with 0.2µg pAG-17, 0.5mM MgCl₂, 0.8µM of each primer, 1x PCR buffer (Perkin-Elmer), 0.2mM dATP, dTTP, dCTP, dGTP, and 2.5U of AmpliTaq® enzyme (Perkin Elmer Norwalk, Connecticut, USA).

Thermocycler program:

cycles 1-35: 94°C 1 min 55°C 1min 72°C 1min;

20 then 10 min at 72°C, then 4°C storage. The amplified bands were purified from a 0.7% agarose gel.

The TBP-DNA- probe was labeled with random primers and Klenow fragment enzyme from Megaprime labeling kit® (Amersham, UK). 25-50ng probe DNA was
 25 combined with 5µl random hexamer primer (e.g. Amersham) and 20 µl H₂O. DNA was denatured at 100°C for 5 minutes. Labeling mix was added to a final volume of 50µl with 1x reaction buffer, 1x dATP, 1x dTTP, 1x dGTP, 4µl 3000Ci/mmol alpha-³²P dCTP (e.g. Amersham) and 2U Klenow enzyme. Reaction was allowed to proceed at RT for 1 hour. Labeled DNA was isolated from unincorporated
 30 nucleotides using size exclusion chromatography (e.g. Push Columns®).

6b) Gel electrophoresis and transfer:

10-20µg of total RNA was ethanol precipitated, the pellet was denatured 10 min at 65°C in RNA loading buffer (65% formamide, 20% formaldehyde (37% solution), 1x MOPS buffer (Sambrook et al. (1989)), 5% glycerol, 0.01% bromphenol blue, 0.01% xylene cyanol, 0.1mg/ml ethidium bromide). RNA was size fractionated on 1% agarose gel containing 18% formaldehyde (37% solution) and 1x MOPS running buffer (40mM MOPS pH 7.0, 10mM sodium acetate, 1mM EDTA). Gels were run slowly overnight at 1V/cm in 1xMOPS running buffer containing 18% formaldehyde (37% solution). Gel was then treated in 0.05M NaOH/1.5M NaCl for 30 minutes, followed by 20 minutes in 0.5M Tris (pH 7.4)/1.5M NaCl. RNA was transferred onto nylon membrane (e.g. Hybond-N+, Amersham, UK) using capillary techniques. Membrane-bound RNA was crosslinked, e.g. UV-crosslinked using a Stratalinker® (Stratagene, La Jolla, CA, USA).

6c) Hybridization and washing:

Pre-hybridization was performed for example in "Rapid-Hyb" buffer (Amersham, UK) in rotating hybridization oven (e.g. Hybaid) at 65°C for 1-2 hours. Hybridization was performed for example in 6-10 ml "Rapid-Hyb" buffer containing 10^6 - 10×10^6 cpm of denatured probe at 65°C for 2-3 hours. Membrane was washed twice at RT (10 min/wash) in 2x SSC/0.1%SDS (1xSSC: 150mM NaCl, 15mM Na₃citrate, pH 7.0). Membrane was further washed 2 times at 65°C (15min/wash) in 1xSSC/0.1% SDS. The next 2 washes were performed at 65°C for 15 min in 0.5xSSC/0.1%SDS, followed by 1 or 2 final washes (as necessary) for 15 min/wash at 65°C in 0.2xSSC/0.1% SDS. Autoradiography of washed membrane.

Example 7: Generation of Polyclonal Antibody Specific for tagged-hTBP

General Strategy:

A significant amount of sequence homology exists between the mouse and human TBP. Therefore, most commercially available antibodies will cross react, detecting

both endogenous and transgenic TBP. To differentiate between the two, a polyclonal antibody was generated against the tagged region of the transgenic TBP (corresponding to the thrombin cleavage site and the His tag of the tagged hTBP):
SEQ ID NO. 12: H₂N-- MGSSHHHHHHSSGLVPRGC--COOH

5

This peptide was coupled to carrier protein and injected into rabbits using standard protocols. Serum was collected at regular intervals and tested for anti-hTBP titer using ELISA assays.

10

Example 8: Western-Blot Detection of Transgenic Protein:

8a) Preparation of liver nuclear extracts:

MT-hTBP mice were subjected to 2 intraperitoneal injections of ZnSO₄ (see 5b)).

15

Mice were killed with cervical dislocation and livers removed. Livers were homogenized immediately in 10ml homogenization buffer (1.8M sucrose, 10mM HEPES pH 7.4 (Sambrook et al. (1989), 25mM KCL, 1mM EDTA, 5% glycerol, 0.15 mM spermine, 0.5mM spermidine, 0.4mM PMSF). After homogenization, volume was increased to 25 ml with the same buffer. The homogenate was carefully layered onto 7 ml homogenization buffer in centrifuge tubes, e.g. in Ultra-Clear SW-28 tubes® (Beckman, Palo Alto, CA, USA). Samples were centrifuged in SW-28 rotor for 1 hour at 25,000 rpm (4°C). Supernatant was carefully removed, and pelleted nuclei were resuspended in 200µl NEXB buffer (20mM HEPES pH 7.9, 400mM NaCl, 1mM EDTA, 1mM EGTA, 1mM DTT, 1mM PMSF, 10% glycerol, 2µg/ml aprotinin, 2µg/ml leupeptin, 2µg/ml pepstatin-A). Resuspended nuclei were incubated on ice for 15-30 min with frequent mixing and pipetting up and down. Samples were submitted to 5 freeze/thaw cycles with dry ice/ethanol bath and 37°C H₂O bath. Samples were centrifuged for 30 seconds at high speed and supernatant was measured for protein content, e.g. with protein assay reagent (e.g. Bio-Rad, Hercules, CA, USA).⁵

30

8b) Electrophoresis and transfer:

20-75µg of extract was size separated by denaturing gel electrophoresis. Resolving gel: 10% acrylamide (1:37 ratio acrylamide:bis-acrylamide), 0.1% SDS, 375mM Tris pH 8.8. Stacking gel: 5% acrylamide, 0.1% SDS, 125mM Tris pH 8.3. Running
 5 buffer: 25mM Tris pH 8.3, 192mM glycine, 0.1% SDS. Protein was transferred onto pure nitrocellulose membrane (e.g. Bio-Rad) with semi-dry blotter (e.g. Hoefer, San Francisco, CA, USA) using modified Bjerrum transfer buffer (48mM Tris, 39mM glycine, 10% methanol, 0.0375% SDS, pH 9.2). Transfer was allowed to run at 0.8mA/cm for 2-4 hours.

10

8c) Immunodetection:

Membranes were blocked 1-2 hours at room temperature on rocking surface in 1x TBS (20mM Tris pH 7.5, 500mM NaCl) with 3% gelatin (e.g. Bio-Rad). Membranes were washed 10 min in 1xTTBS (1xTBS with 0.05% Tween-20) at RT. Hybridization
 15 with primary antibody in 1% gelatin/1xTTBS was performed for 4 hours to overnight on rocking surface at RT. Membranes were washed 2-3 times (5 min/wash) with 1x TTBS. Hybridization with secondary antibody coupled with alkaline phosphatase (AP) in 1% gelatin/1xTTBS was done for 2-3 hours. Membranes were washed again 2-3 times (5 min/wash) in 1xTTBS at RT followed by 5 min wash in 1x TBS buffer.

20

Membranes were then incubated in 1x development buffer (e.g. BioRad) containing NBT/BCIP reagents at RT until sufficient appearance of bands occurred. AP reaction was stopped by H₂O wash.

25

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Hoechst Aktiengesellschaft
- (B) STREET: -
- (C) CITY: Frankfurt
- (D) STATE: -
- (E) COUNTRY: Germany
- (F) POSTAL CODE (ZIP): 65926
- (G) TELEPHONE: 069-305-7072
- (H) TELEFAX: 069-35-7175
- (I) TELEX: -

(ii) TITLE OF INVENTION: Purification of higher order transcription complexes from transgenic non-human animals

(iii) NUMBER OF SEQUENCES: 17

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..12

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Val
1 5 10

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

```
(ix) FEATURE:
      (A) NAME/KEY: Peptide
      (B) LOCATION: 1..11
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..9

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
1 5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon

(B) LOCATION: 1..22



(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GGAGCAACCG CCTGCTGGGT GC

22

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon
 (B) LOCATION: 1..21

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CCTGTGTGC CTGCTGGGAC G

21

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon
 (B) LOCATION: 1..21

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GGAGACTGAA GTTAGGCCAG C

21

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 76 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

The image shows four 5x5 dot patterns arranged vertically. The first pattern represents the digit '8', the second '2', the third '9', and the fourth '5'. Each pattern is composed of black dots on a white background.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCGGCACCAG GCCGCTGCTG TGATGATGAT GATGATGGCT GCTGCCCATG ACTGCGTAAT 60
 GCGGTCATGA CGCTTT 76

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 75 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 1..75

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

GAAGGGGGTG GGGGAGGCAA GGGTACATGA GAGCCATTAC GTCGTCTTCC TGAATCCCTT 60
 TAGCCGCTTT GCTCG 75

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 1..22

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

CCCTATGACG TCCCGGATTA CG 22

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon
(B) LOCATION: 1..22

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GTGGAGTGGT GCCCGGCAAG GG

22

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

(A) NAME/KEY: Peptide
(B) LOCATION: 1..19

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
1 5 10 15
Arg Gly Cys

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1310 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon
(B) LOCATION: 1..1310

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CCATGGGCTA TCCCTATGAC GTCCCGGATT ACGCAGTCAT GGGCAGCAGC CATCATCATC 60
ATCATCACAG CAGCGGCCTG GTGCCGCGCG GCAGCCATAT GGATCAGAAC AACAGCCTGC 120
CACCTTACGC TCAGGGCTTG GCCTCCCCCTC AGGGTGCCAT GACTCCCGGA ATCCCTATCT 180
TTAGTCCAAT GATGCCTTAT GGCCTGGAC TGACCCACACA GCCTATTTCAG AACACCAATA 240
GTCTGTCTAT TTTGGAAGAG CAACAAAGGC AGCAGCAGCA ACAACAACAG CAGCAGCAGC 300
AGCAGCAGCA GCAGCAACAG CAACAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC 360
AGCAGCAGCA GCAGCAGCAA CAGGCAGTGG CAGCTGCAGC CGTTCAGCAG TCAACGTCCC 420
AGCAGGCAAC ACAGGGAACC TCAGGCCAGG CACCACAGCT CTTCCTCA CAGACTCTCA 480
CAACTGCACC CTTGCCGGGC ACCACTCCAC TGTATCCCTC CCCCATGACT CCCATGACCC 540
CCATCACTCC TGCCACGCCA GCTTCGGAGA GTTCTGGGAT TGTACCGCAG CTGCAAAATA 600
TTGTATCCAC AGTGAATCTT GGTGTAAAC TTGACCTAAA GACCATTGCA CTTCTGCCCC 660
GAAACGCCGA ATATAATCCC AAGCGSTTTG CTGCGGTAAT CATGAGGATA AGAGAGCCAC 720
GAACCACGGC ACTGATTTTC AGTTCCTGGA AAATGGTGTG CACAGGAGCC AAGAGTGAAG 780
AACAGTCCAG ACTGGCAGCA AGAAAATATG CTAGAGTTGT ACAGAAGTTG GGTTTTCCAG 840
CTAAGTTCTT GGAATTCAAG ATTCAGAACA TGGTGGGAG CTGTGATGTG AAGTTTCTTA 900
TAAGGTTAGA AGGCCTTGTG CTCACCCACC AACAATTTAG TAGTTATGAG CCAGAGTTAT 960
TTCTGGGTTT AATCTACAGA ATGATCAAAC CCAGAATTGT TCTCCTTATT TTTGTTTCTG 1020
GAAAAGTTGT ATTAACAGGT GCTAAAGTCA GAGCAGAAAT TTATGAAGCA TTTGAAAACA 1080
TCTACCTAT TCTAAGGGA TTCAGGAAGA CGACGTAATG GCTCTCATGT ACCCTTGCCT 1140
CCCCGACCCC CTTCTTTTTT TTTTTTAAA CAAATCAGTT TGTTTTGGTA CCTTTAAATG 1200
GTGGTGTGTG GAGAAGATGG ATGTTGAGTT GCAGGGTGTG GCACCAGGTG ATGCCCTTCT 1260
GTAAGTGCCC CTCCGGCAT CCCGGAATTC CTGCAGCCCA ACGCGGCCGC 1310

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4286 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: exon

(B) LOCATION: 1..4286

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GAATTCCTCCCT GCAGGTCCT TAGCGTTGGC CACATAGTAG GTTCTCAAAT ACTTGTTAAT	60
AAATAAGTTT GTTCGAGAAG CTGGGCAATG ATATTCTACA GCTGGAAGAA GAAACATAAT	120
GATCTAGTAA TTAGCTCAAT TAAAAATAAA CGTTCTTCTT TCCTCAGAGG AGCATTTCCT	180
AAGGCCTGCC TTGATAGCCA TCCAAAAGG CCAAGCTCAT CCAATCTTGC CCTAGATTTA	240
TGCTAAAATG CAGTTACAAT CGATAGGATG ACAGAAAACG ACAGCACTTA TTTAAATATA	300
ATAGGCACCTT ATTTAAATAG GAGAAGCTGT GACTTCATAG CAAGTGTGG GGTAGGAAA	360
CTGGGTGGAT AAAGTGTGCT ATGCTGTAGA TCTTAGCCTC TACATGAGAT CATGTGGAAA	420
ATCTGAAAGC ATTTTAGGTT CCTTATGTTT GCAATCAAAT AACTGTACAC CTTTAAATTT	480
AAAAAGTACC ATGAGGCACA CACACACACT CGCAGGAAGT TTTGGCGTA ACAAAGTAG	540
AATTAGATCT AAAAGCTAAC TGTAGGACTG AGTCTATTCT AAAGTGAAG CCTGGACATC	600
TGGAGTACCA GGGGGAGATG ACGTGTACG GGCTTCCATA AAAGCAGCTG GCTTTGAATG	660
GAAGGAGCCA AGAGGCCAGC ACAGGAGCGG ATTCGTCGCT TTCACGGCCA TCGAGCCGAA	720
CCTCTCGCAA GTCCGTGAGC CGTTAAGGAG GCCCCAGTC CCGACCTTC GCCCAAGCC	780
CCTCGGGGTC CCCGGGCTG GTACTCCTTG CCACACGGGA GGGGCGCGGA AGCCGGGGCG	840
GAGGAGGAGC CAACCCCGGG CTGGGCTGAG ACCCGCAGAG GAAGACGCTC TAGGGATTG	900
TCCCGGACTA GCGAGATGGC AAGGCTGAGG ACGGGAGGCT GATTGAGAGG CGAAGGTACA	960
CCCTAATCTC AATACAACCT TTGGAGCTAA GCCAGCAATG GTAGAGGGAA GATTCTGCAC	1020
GTCCCTTCCA GCGCGCTCC CCGTCACCAC CCCCCCAAC CCGCCCCGAC CGGAGCTGAG	1080
AGTAATTCAT ACAAAGGAC TCGCCCTGC CTTGGGGAAT CCCAGGACC GTCGTTAAAC	1140
TCCCACTAAC GTAGAACCA GAGATCGCTG CGTTCCCGCC CCCTCACCAG CCCGCTCTCG	1200
TCATCACTGA GGTGGAGAAG AGCATGCGTG AGGCTCCGGT GCCCGTCAGT GGGCAGAGCG	1260
CACATCGCCC ACAGTCCCCG AGAAGTTGGG GGGAGGGGTC GGCAATTGAA CCGGTGCCTA	1320
GAGAAGGTGG CGCGGGGTAA ACTGGGAAAG TGATGTCGTG TACTGGCTCC GCCTTTTTCC	1380
CGAGGGTGGG GGAGAACCGT ATATAAGTGC AGTAGTCGCC GTGAACGTTC TTTTCGCAA	1440
CGGGTTTGCC GCCAGAACAC AGGTAAGTGC CGTGTGTGGT TCCCGCGGGC CTGGCCTCTT	1500

TACGGGTAT	GGCCCTTGCG	TGCCTTGAAT	TACTTCCACG	CCCCTGGCTG	CAGTACGTGA	1560
TTCTTGATCC	CGAGCTTCGG	GTTGGAAGTG	GGTGGGAGAG	TTCGAGGCCT	TGCGCTTAAG	1620
GAGCCCTTC	GCCTCGTGCT	TGAGTTGAGG	CCTGGCCTGG	GCGCTGGGGC	CGCCGCGTGC	1680
GAATCTGGTG	GCACCTTCGC	GCCTGTCTCG	CTGCTTTCGA	TAAGTCTCTA	GCCATTTAAA	1740
ATTTTGTATG	ACCTGCTGCG	ACGCTTTTTT	TCTGGCAAGA	TAGTCTTGTA	AATGCGGGCC	1800
AAGATCTGCA	CACTGGTATT	TCGGTTTTTG	GGGCCGCGGG	CGGCGACGGG	GCCCGTGCGT	1860
CCCAGCGCAC	ATGTTGCGCG	AGGCGGGGCC	TGCGAGCGCG	GCCACCGAGA	ATCGGACGGG	1920
GGTAGTCTCA	AGCTGGCCGG	CCTGCTCTGG	TGCCTGGCCT	GCGCCGCGCG	TGTATCGCCC	1980
CGCCCTGGGC	GGCAAGGCTG	GCCCGGTGCG	CACCAGTTGC	GTGAGCGGAA	AGATGGCCGC	2040
TTCCCGGCCC	TGCTGCAGGG	AGCTCAAAAT	GGAGGACGCG	GCGCTCGGGA	GAGCGGGCGG	2100
GTGAGTCACC	CACACAAAGG	AAAAGGGCCT	TTCCGTCCTC	AGCCGTCGCT	TCATGTGACT	2160
CCACGGAGTA	CCGGGCGCCG	TCCAGGCACC	TCGATTAGTT	CTCGAGCTTT	TGGAGTACGT	2220
CGTCTTTAGG	TTGGGGGGAG	GGGTTTTATG	CGATGGAGTT	TCCCCACACT	GAGTGGGTGG	2280
AGACTGAAGT	TAGGCCAGCT	TGGCACTTGA	TGTAATTCTC	CTTGGAATTT	GCCCTTTTTG	2340
AGTTTGGATC	TTGGTTCATT	CTCAAGCCTC	AGACAGTGGT	TCAAAGTTTT	TTTCTTCCAT	2400
TTCAGGTGTC	GTGAGGAATT	GCCCCGGGGA	TCCATGGGCT	ATCCCTATGA	CGTCCCGGAT	2460
TACGCAGTCA	TGGGCAGCAG	CCATCATCAT	CATCATCACA	GCAGCGGCCT	GGTGCCGCGC	2520
GGCAGCCATA	TGGATCAGAA	CAACAGCCTG	CCACCTTACG	CTCAGGGCTT	GGCCTCCCCT	2580
CAGGGTGCCA	TGACTCCCGG	AATCCCTATC	TTTAGTCCAA	TGATGCCTTA	TGGCACTGGA	2640
CTGACCCAC	AGCCTATTCA	GAACACCAAT	AGTCTGTCTA	TTTTGGAAGA	GCAACAAAGG	2700
CAGCAGCAGC	AACAACAACA	GCAGCAGCAG	CAGCAGCAGC	AGCAGCAACA	GCAACAGCAG	2760
CAGCAGCAGC	AGCAGCAGCA	GCAGCAGCAG	CAGCAGCAGC	AGCAGCAGCA	ACAGGCAGTG	2820
GCAGCTGCAG	CCGTTTCAGCA	GTCACCGTCC	CAGCAGGCAA	CACAGGGAAC	CTCAGGCCAG	2880
GCACCACAGC	TCTTCCACTC	ACAGACTCTC	ACAACTGCAC	CCTTGCCGGG	CACCACTCCA	2940
CTGTATCCCT	CCCCCATGAC	TCCCATGACC	CCCATCACTC	CTGCCACGCC	AGCTTCGGAG	3000
AGTTCTGGGA	TTGTACCGCA	GCTGCAAAAT	ATTGTATCCA	CAGTGAATCT	TGTTGTAAA	3060
CTTGACCTAA	AGACCATTGC	ACTTCGTGCC	CGAAACGCCG	AATATAATCC	CAAGCGGTTT	3120
GCTGCGGTAA	TCATGAGGAT	AAGAGAGCCA	CGAACCACGG	CACTGATTTT	CAGTCTGGG	3180
AAAAATGGTGT	GCACAGGAGC	CAAGAGTGAA	GAACAGTCCA	GACTGGCAGC	AAGAAAATAT	3240

GCTAGAGTTG TACAGAAGTT GGGTTTTCCA GCTAAGTTCT TGGACTTCAA GATTCAGAAC 3300
 ATGGTGGGGA GCTGTGATGT GAAGTTTCCT ATAAGGTTAG AAGGCCTTGT GCTCACCAC 3360
 CAACAATTTA GTAGTTATGA GCCAGAGTTA TTTCCTGGTT TAATCTACAG AATGATCAAA 3420
 CCCAGAATTG TTCTCCTTAT TTTGTTTCT GGAAGTTG TATTAACAGG TGCTAAAGTC 3480
 AGAGCAGAAA TTTATGAAGC ATTTGAAAC ATCTACCCTA TTCTAAAGG ATTGAGGAAG 3540
 ACGACGTAAT GGCTCTCATG TACCCTTGCC TCCCCACCC CCTTCTTTT TTTTTTTAA 3600
 ACAAATCAGT TTGTTTGGT ACCTTTAAAT GGTGGTGTG TGAGAAGATG GATGTTGAGT 3660
 TGCAGGGTGT GGCACCAGGT GATGCCCTC TGTAAGTGCC CCTTCCGGCA TCCCGGATAT 3720
 CCTGCAGCCC AACACGGCCG CTCGAGCATG CATCTAGAGA ACGTCACGGC CGCGATCCCC 3780
 CTGTGCCTTC TAGTTGCCAG CCATCTGGTT GTTTGCCCT CCCCCTGCC TTCCTTGACC 3840
 CTGGAAGGTG CCACTCCAC TGTCCTTCC TAATAAAATG AGCAAATTGC ATCGCATTGT 3900
 CTGAGTAGGT GTCATTCTAT TCTGGGGGT GGGGTGGGC AGGACAGCAA GGGGGAGGAT 3960
 TGGGAAGACA ATAGCAGGCA TGCTGGGGAT GCGGTGGGCT CTATGGGTAC CCAGGTGCTG 4020
 AAGAATTGAC CCGGTTCTC CTGGGCCAGA AAGAAGCAGG CACATCCCT TCTCTGTGAC 4080
 ACACCCTGTC CACGCCCTG GTTCTTAGT CCAGCCCCAC TCATAGGACA CTCAACTTGG 4140
 AGCGGTCTCT CCCTCCCTCA TCAGCCCACC AAACCAAACC TAGCCTCAA GAGTGGGAAG 4200
 AAATTAAAGC AAGAAGGCTA TTAAGTGAG AGGGAGAGAA AATGCCTCCA ACATGTGAGG 4260
 AAGTAATGAT AGAAATCATA GAATTC 4286

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 3263 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 1..3263

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

ATCGATAAGC TGAGATCCCG CTAGAACTG CTGAGGGCTG GACCGCATCT GGGGACCATC 60

TGTTCTTGGC	CCTGAGCGGG	GCAGGAAGTG	CTTACCGCAG	ATATCCTGTT	TGCCCCAATT	120
CAGCTGTTC	ATCTGTTCTT	GGCCCTGAGC	GGGGCAGGAA	CTGCTTACCA	CAGATATCCT	180
GTTTGCCCA	TATTCAGCTG	TCTCTCTGTT	CCTGACCTTG	ATCTGAACCT	CTCTATTCTC	240
AGTTATGTAT	TTTTCCCATG	CCTTGCAAAA	TGGCGTTACT	TAAGCTAGCT	TGCCAAACCT	300
ACGGCTGGGG	TCTTTCACGT	TTATATCTAT	GAGGGGAAGG	ACCCAGAGTG	GGGAAGCTGG	360
GATCTTGGGA	ACACGCTTCT	CTACATGGCA	TTGTCTGCAC	GGTGGAGTCC	GGATCTGAGC	420
TTGGCTTGGT	TTTTAAAACC	AGCCTGGAGT	AGAGCAGATG	GGTTAAGGTG	AGTGACCCCT	480
CAGCCCTGGA	CATTCTTAGA	TGAGCCCCCT	CAGGAGTAGA	GAATAATGTT	GAGATGAGTT	540
CTGTTGGCTA	AAATAATCAA	GGCTAGTCTT	TATAAACTG	TCTCCTCTTC	TCCTAGCTTC	600
GATCCAGAGA	GAGACCTGGG	CGGAGCTGGT	CGCTGCTCAG	GAAGTCCAGG	AAAGGAGAAG	660
CTGAGGTTAC	CACGCTGCGA	ATGGGTTTAC	GGAGATAGCT	GGCTTTCCGG	GGTGAGTTCT	720
CGTAAACTCC	AGAGCAGCGA	TAGGCCGTAA	TATCGGGGAA	AGCACTATAG	GGACATGATG	780
TTCCACACGT	CACATGGGTC	GTCCTATCCG	AGCCAGTCGT	GCCAAAGGGG	CGGTCCCGCT	840
GTGCACACTG	GCGCTCCAGG	GAGCTCTGCA	CTCCGCCCGA	AAAGTGGGCT	CGGCTCTGCC	900
AGGACGCGGG	GCGCGTGACT	ATGCGTGGGC	TGGAGCAACC	GCCTGCTGGG	TGCAAAACCT	960
TTGCGCCCGG	ACTCGTCCAA	CGACTATAAA	GAGGGCAGGC	TGCTCTCTAA	GCGTCACCAC	1020
GACTTCAACG	TCCTGAGTAC	CTTCTCCTCA	CTTACTCCGT	AGCTCCAGCT	TCACCAGATC	1080
CTCGAGAACG	TCTCCCATGG	GCTATCCCTA	TGACGTCCCG	GATTACGCAG	TCATGGGCAG	1140
CAGCCATCAT	CATCATCATC	ACAGCAGCGG	CCTGGTGCCG	CGCGGCAGCC	ATATGGATCA	1200
GAACAACAGC	CTGCCACCTT	ACGCTCAGGG	CTTGGCCTCC	CCTCAGGGTG	CCATGACTCC	1260
CGGAATCCCT	ATCTTTAGTC	CAATGATGCC	TTATGGCACT	GGACTGACCC	CACAGCCTAT	1320
TCAGAACACC	AATAGTCTGT	CTATTTTGGA	AGAGCAACAA	AGGCAGCAGC	AGCAACAACA	1380
ACAGCAGCAG	CAGCAGCAGC	AGCAGCAGCA	ACAGCAACAG	CAGCAGCAGC	AGCAGCAGCA	1440
GCAGCAGCAG	CAGCAGCAGC	AGCAGCAGCA	GCAACAGGCA	GTGGCAGCTG	CAGCCGTTCA	1500
GCAGTCAACG	TCCCAGCAGG	CAACACAGGG	AACCTCAGGC	CAGGCACCAC	AGCTCTTCCA	1560
CTCACAGACT	CTCACAACTG	CACCCTTGCC	GGGCACCACT	CCACTGTATC	CCTCCCCCAT	1620
GACTCCCATG	ACCCCATCA	CTCCTGCCAC	GCCAGCTTCG	GAGAGTTCTG	GGATTGTACC	1680
GCAGCTGCAA	AATATTGTAT	CCACAGTGAA	TCTTGGTTGT	AAACTTGACC	TAAAGACCAT	1740

TGCACTTCGT GCCCGAAACG CCGAATATAA TCCCAAGCGG TTTGCTGCGG TAATCATGAG	1800
GATAAGAGAG CCACGAACCA CGGCACTGAT TTTCAGTTCT GGGAAAATGG TGTGCACAGG	1860
AGCCAAGAGT GAAGAACAGT CCAGACTGGC AGCAAGAAAA TATGCTAGAG TTGTACAGAA	1920
GTTGGGTTTT CCAGCTAAGT TCTTGGACTT CAAGATTCAG AACATGGTGG GGAGCTGTGA	1980
TGTGAAGTTT CCTATAAGGT TAGAAGGCCT TGTGCTCACC CACCAACAAT TTAGTAGTTA	2040
TGAGCCAGAG TTATTTCCCTG GTTTAATCTA CAGAATGATC AAACCCAGAA TTGTTCTCCT	2100
TATTTTTGTT TCTGGAAGG TTGTATTAAC AGGTGCTAAA GTCAGAGCAG AAATTATGA	2160
AGCATTTGAA AACATCTACC CTATTCTAAA GGGATTCAGG AAGACGACGT AATGGCTCTC	2220
ATGTACCCTT GCCTCCCCCA CCCCCTTCTT TTTTTTTTTT TAAACAAATC AGTTTGTTTT	2280
GGTACCTTTA AATGGTGGTG TTGTGAGAAG ATGGATGTTG AGTTGCAGGG TGTGGCACCA	2340
GGTGATGCCC TTCTGTAAGT GCCCCTTCCG GCATCCCGGA ATTCCTGCAG CCCAACGCGG	2400
CCGCTTCGAG GGATCTTTGT GAAGGAACCT TACTTCTGTG GTGTGACATA ATTGGACAAA	2460
CTACCTACAG AGATTTAAAG CTCTAAGGTA AATATAAAAT TTTTAAGTGT ATAATGTGTT	2520
AAACTACTGA TTCTAATTGT TTGTGTATTT TAGATTCCAA CCTATGGAAC TGATGAATGG	2580
GAGCAGTGGT GGAATGCCTT TAATGAGGAA AACCTGTTTT GCTCAGAAGA AATGCCATCT	2640
AGTGATGATG AGGCTACTGC TGA CTCTCAA CATTCTACTC CTCCAAAAA GAAGAGAAAG	2700
GTAGAAGACC CCAAGGACTT TCCTTCAGAA TTGCTAAGTT TTTTGAGTCA TGCTGTGTTT	2760
AGTAATAGAA CTCTTGCTTG CTTTGCTATT TACACCACAA AGGAAAAAGC TGA CTGCTA	2820
TACAAGAAAA TTATGGAAAA ATATTCTGTA ACCTTTATAA GTAGGCATAA CAGTTATAAT	2880
CATAACATAC TGTTTTTTCT TACTCCACAC AGGCATAGAG TGTCTGCTAT TAATAACTAT	2940
GCTCAAAAAT TGTGTACCTT TAGCTTTTTA ATTTGTAAAG GGGTTAATAA GGAATATTTG	3000
ATGTATAGTG CCTTGACTAG AGATCATAAT CAGCCATACC ACATTGTAG AGGTTTTACT	3060
TGCTTTAAAA AACCTCCCAC ACCTCCCCCT GAACCTGAAA CATAAAATGA ATGCAATTGT	3120
TGTTGTTAAC TTGTTTATTG CAGCTTATAA TGTTTACAAA TAAAGCAATA GCATCACAAA	3180
TTTCACAAAT AAAGCATTTT TTCACTGCA TTCTAGTTGT GGTGTGTTCA AACTCATCAA	3240
TGTATCTTAT CATGTCTGGA TCC	3263

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 371 amino acids

(B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

(A) NAME/KEY: Protein
 (B) LOCATION: 1..371

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

```

Met Gly Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Val Met Gly Ser Ser
1           5           10           15
His His His His His His Ser Ser Gly Leu Val Pro Arg Gly Ser His
20           25           30
Met Asp Gln Asn Asn Ser Leu Pro Pro Tyr Ala Gln Gly Leu Ala Ser
35           40           45
Pro Gln Gly Ala Met Thr Pro Gly Ile Pro Ile Phe Ser Pro Met Met
50           55           60
Pro Tyr Gly Thr Gly Leu Thr Pro Gln Pro Ile Gln Asn Thr Asn Ser
65           70           75           80
Leu Ser Ile Leu Glu Glu Gln Gln Arg Gln Gln Gln Gln Gln Gln
85           90           95
Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln
100          105          110
Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Ala
115          120          125
Val Ala Ala Ala Ala Val Gln Gln Ser Thr Ser Gln Gln Ala Thr Gln
130          135          140
Gly Thr Ser Gly Gln Ala Pro Gln Leu Phe His Ser Gln Thr Leu Thr
145          150          155          160
Thr Ala Pro Leu Pro Gly Thr Thr Pro Leu Tyr Pro Ser Pro Met Thr
165          170          175
Pro Met Thr Pro Ile Thr Pro Ala Thr Pro Ala Ser Glu Ser Ser Gly
180          185          190
Ile Val Pro Gln Leu Gln Asn Ile Val Ser Thr Val Asn Leu Gly Cys
195          200          205
Lys Leu Asp Leu Lys Thr Ile Ala Leu Arg Ala Arg Asn Ala Glu Tyr
210          215          220
Asn Pro Lys Arg Phe Ala Ala Val Ile Met Arg Ile Arg Glu Pro Arg

```

225					230					235				240	
Thr	Thr	Ala	Leu	Ile	Phe	Ser	Ser	Gly	Lys	Met	Val	Cys	Thr	Gly	Ala
				245					250					255	
Lys	Ser	Glu	Glu	Gln	Ser	Arg	Leu	Ala	Ala	Arg	Lys	Tyr	Ala	Arg	Val
			260					265					270		
Val	Gln	Lys	Leu	Gly	Phe	Pro	Ala	Lys	Phe	Leu	Asp	Phe	Lys	Ile	Gln
			275				280					285			
Asn	Met	Val	Gly	Ser	Cys	Asp	Val	Lys	Phe	Pro	Ile	Arg	Leu	Glu	Gly
290						295					300				

Leu Val Leu Thr His Gln Gln Phe Ser Ser Tyr Glu Pro Glu Leu Phe
 305 310 315 320
 Pro Gly Leu Ile Tyr Arg Met Ile Lys Pro Arg Ile Val Leu Leu Ile
 325 330 335
 Phe Val Ser Gly Lys Val Val Leu Thr Gly Ala Lys Val Arg Ala Glu
 340 345 350
 Ile Tyr Glu Ala Phe Glu Asn Ile Tyr Pro Ile Leu Lys Gly Phe Arg
 355 360 365
 Lys Thr Thr
 370

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 18 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (ix) FEATURE:
- (A) NAME/KEY: Protein
 - (B) LOCATION: 1..18

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Met Gly Ser Ser His His His His His Ser Ser Gly Leu Val Pro
 1 5 10 15
 Arg Gly

"Comprises/comprising" when used in this specification is taken to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Transgenic non-human vertebrate having the ability to express epitope-tagged TATA-box binding protein (TBP).
2. Transgenic non-human vertebrate according to claim 1, wherein TBP is expressed as fusion protein with two epitope-tags.
3. Transgenic non-human vertebrate according to any one of claims 1 and 2, wherein the fusion protein comprises human TBP (hTBP).
4. Transgenic non-human vertebrate according to any one of claims 1 to 3, wherein the fusion protein comprises a HA- and a His-epitope.
5. Transgenic non-human vertebrate according to claim 4, wherein the fusion protein has the sequence SEQ ID NO. 16.
6. Transgenic non-human vertebrate according to any one of claims 1 to 5, wherein the transgenic vertebrate is a mice.
7. A method of making a transgenic non-human vertebrate as claimed in any one of claims 1 to 6 by introducing a transgene into the germline and/or into somatic cells of a non-human vertebrate.
8. A method of making a non-human transgenic vertebrate as claimed in claim 7 by microinjecting transgenic DNA into a non-human vertebrate zygote.
9. A method of making a non-human transgenic vertebrate as claimed in claim 7 by transfecting a blastocyst with a vector that contains a transgene.
10. A method of making a non-human transgenic vertebrate as claimed in claim 7 by introducing a transgene into an embryonic stem cell.



11. A transgene that encodes for epitope-tagged TBP when used to make a transgenic non-human vertebrate as claimed in any one of claims 1 to 6.

12. A transgene as claimed in claim 11, which comprises a first DNA sequence that encodes for TBP and a second DNA sequence that encodes for one or more epitope-tags.

13. A transgene as claimed in any one of claims 11 and 12, wherein the DNA sequence that encodes for TBP is a cDNA.

14. A transgene as claimed in claim 13, wherein the DNA sequence is the cDNA of human TBP (hTBP).

15. A transgene as claimed in any one of claims 11 to 14, wherein the second DNA sequence encodes for two epitope-tags.

16. A transgene as claimed in claim 15, wherein the epitope-tags are a HA-epitope and a His-epitope.

17. A transgene as claimed in any one of claims 15 and 16, wherein the HA-epitope has one of the sequences SEQ ID NO. 1, SEQ ID NO. 2, SEQ ID NO. 3 or SEQ ID NO. 4.

18. A transgene as claimed in any one of claims 11 to 17, which comprises the sequence SEQ ID NO. 13.

19. A transgene as claimed in any one of claims 11 to 18, which comprises the DNA of an inducible or a constitutive promoter.

20. A transgene as claimed in claim 19, wherein the promoter is the promoter of the EF gene.

21. A transgene as claimed in claim 19, wherein the promoter is the promoter of the MT gene.



22. A transgene as claimed in any one of claims 11 to 22, wherein the transgene has the sequence SEQ ID NO. 14 or SEQ ID NO. 15.

23. The use of a transgene as claimed in any one of claims 11 to 22 for making a non-human transgenic vertebrate.

24. A method of making a transgene as claimed in any one of claims 11 to 22, by connecting the DNA sequence(s) that encode for one or more epitope-tags to the DNA sequence that encodes for the TBP protein.

25. The use of a transgene as claimed in any one of claims 11 to 22 for making epitope-tagged TBP.

26. The use of a transgenic vertebrate as claimed in any one of claims 1 to 6 for expressing epitope-tagged TBP.

27. An epitope-tagged TBP expressed in a transgenic vertebrate as claimed in any one of claims 1 to 6.

28. An epitope-tagged TBP as claimed in claim 27 that comprises two epitopes.

29. An epitope-tagged TBP as claimed in any one of claims 27 and 28, wherein TBP is hTBP.

30. An epitope-tagged TBP as claimed in any one of claims 27 to 29, wherein TBP is linked to a HA-epitope and a His-epitope.

31. An epitope-tagged TBP as claimed in any one of claims 27 to 30, wherein the HA epitope has one of the sequences SEQ ID NO. 1, SEQ ID NO. 2, SEQ ID NO. 3 or SEQ ID NO. 4.

32. An epitope-tagged TBP as claimed in any one of claims 27 to 31 which has the sequence SEQ ID NO. 16.



33. A method of making an epitope-tagged TBP as claimed in any one of claims 27 to 32 by introducing a transgene as claimed in any one of claims 11 to 22 into the germline and/or somatic cells of a non-human vertebrate.

34. A method of making an epitope-tagged TBP as claimed in claim 33, by introducing a transgene that comprises a constitutive promoter into a non-human vertebrate, whereupon epitope-tagged TBP is expressed in particular cell types and/or tissue of the animal, optionally at a particular developmental stage of the animal.

35. A method of making an epitope-tagged TBP as claimed in claim 33, by introducing a transgene that comprises an inducible promoter into a non-human vertebrate and expression of the epitope-tagged TBP upon induction of the promoter.

36. The use of epitope-tagged TBP for the isolation of higher order transcription complexes from a non-human transgenic vertebrate as claimed in any one of claims 1 to 6 and for the identification of TAFs and TAF-interacting factors from such non-human transgenic vertebrate.

37. Use of a transgenic non-human vertebrate as claimed in any one of claims 1 to 6, for expressing epitope-tagged TBP.

38. The use of a transgenic non-human vertebrate as claimed in any one of claims 1 to 6 for the isolation of higher order transcription complexes from different tissues and/or cell types, optionally at different developmental stages of the animal.

39. The use of a transgenic non-human vertebrate as claimed in any one of claims 1 to 6 for identifying new and/or specific TAFs and/or TAF-interacting factors.

40. A method for the identification and characterization of different higher order transcription complexes, wherein epitope-tagged TBP to which higher order



transcription complexes are associated is isolated from a transgenic non-human vertebrate.

41. A method of characterizing the composition of different higher order transcription complexes, wherein a) a transgene as claimed in any one of claims 11 to 22 is introduced into a non-human vertebrate, b) epitope-tagged TBP is isolated from different animal tissue and/or different cell types of the animal, optionally at different developmental stages of the animal and c) the composition of the higher order transcription complexes is determined.

42. A method of identifying a new and/or a specific TAF and/or TAF-interacting factor, wherein a) a transgene as claimed in any one of claims 11 to 22 is introduced into a non-human vertebrate, b) epitope-tagged TBP is isolated from a particular animal tissue and/or a particular cell type of the animal, optionally at a particular developmental stage and c) the TAF and/or TAF-interacting factors associated with the epitope-tagged TBP in the higher order transcription complex are dissociated and separated and d) optionally the amino acid sequence of a TAF and/or a TAF-interacting factor is determined.

43. A method of isolating different higher order transcription complexes from a transgenic non-human vertebrate as claimed in any one of claims 1 to 6, wherein a higher order transcription complex associated with epitope-tagged TBP is affinity co-purified, when the epitope-tagged TBP is affinity purified by using at least one of the epitopes tagged to TBP.

44. A method of isolating a higher order transcription complex as claimed in claim 43, wherein epitope-tagged TBP is purified by binding of its His epitope to a Ni^{2+} -column.

45. A method of isolating a higher order transcription complex as claimed in any one of claims 43 and 44, wherein epitope-tagged TBP is purified by binding of its HA epitope to anti-HA antibodies.



46. A method of isolating a higher order transcription complex as claimed in any one of claims 43 to 45, wherein the higher order transcription complex is isolated by using an affinity column with antibodies that recognize an epitope of an epitope-tagged TBP that has the sequence SEQ ID NO. 17.

47. An antibody that recognizes an epitope-tagged TBP that has the sequence SEQ ID NO. 17 when used to study transgenic non-human vertebrates as claimed in any one of claims 1 to 6.

48. Transgenic non-human vertebrates whose genome comprises a nucleic acid sequence encoding an epitope-tagged TATA-box binding protein (TBP), wherein the expressed epitope-tagged TBP is used for the isolation of developmental- or tissue-specific higher order transcription complexes from said transgenic animal and for the identification of developmental- or tissue-specific TAFs and TAF-interacting factors.

DATED this 10th day of April 2002

HOECHST AKTIENGESELLSCHAFT

WATERMARK PATENT & TRADE MARK ATTORNEYS
290 BURWOOD ROAD
HAWTHORN VICTORIA 3122
AUSTRALIA

P12452AU00 KJS:BJD:SLB

