



US009265238B2

(12) **United States Patent**
Porter et al.

(10) **Patent No.:** **US 9,265,238 B2**
(45) **Date of Patent:** **Feb. 23, 2016**

(54) **MMTV-SV40-SPY1A AND SPY1A-PTRE
TRANSGENIC MOUSE MODELS**

(71) Applicants: **Lisa Porter**, Amherstburg (CA);
Bre-Anne Fifield, Windsor (CA);
Dorota Lubanska, Windsor (CA);
Ingrid Qemo, Windsor (CA)

(72) Inventors: **Lisa Porter**, Amherstburg (CA);
Bre-Anne Fifield, Windsor (CA);
Dorota Lubanska, Windsor (CA);
Ingrid Qemo, Windsor (CA)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/679,482**

(22) Filed: **Apr. 6, 2015**

(65) **Prior Publication Data**

US 2015/0272091 A1 Oct. 1, 2015

Related U.S. Application Data

(63) Continuation-in-part of application No. 13/987,769, filed on Aug. 30, 2013.

(60) Provisional application No. 61/695,719, filed on Aug. 31, 2012, provisional application No. 61/743,501, filed on Sep. 6, 2012.

(51) **Int. Cl.**

A01K 67/027 (2006.01)

A61K 49/00 (2006.01)

(52) **U.S. Cl.**

CPC **A01K 67/0275** (2013.01); **A01K 67/0278** (2013.01); **A61K 49/0008** (2013.01); **A01K 2217/052** (2013.01); **A01K 2227/105** (2013.01); **A01K 2267/0331** (2013.01)

(58) **Field of Classification Search**

CPC **A01K 67/0275**; **A01K 67/0278**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,736,866 A 4/1988 Leder et al.
5,925,803 A 7/1999 Leder et al.

OTHER PUBLICATIONS

Blakely, Collin, M. et al., "Developmental stage determines the effects of MYC in the mammary epithelium", Development, 132 (2005): 1147-1160, the Company of Biologists.

Kirou, Evangelia, "Elucidating the Role of Spy1A during c-Myc Induced Mammary Tumor Development" (2011). Electronic Theses and Dissertations. Paper 290. Windsor, Canada.

Primary Examiner — Scott Long

(57) **ABSTRACT**

In one aspect, the invention provides a transgenic non-human animal model having germ cells and somatic cells containing an endogenous MMTV-SV40-Spy1A gene sequence introduced into said animal model or an ancestor of said animal model at an embryonic stage, wherein said gene sequence comprises a mouse mammary tumor virus gene (MMTV), a functionally disrupted SV40 gene (SV40) and a human Spy1A gene. In another aspect, the present invention provides a transgenic non-human animal model whose germ cells and somatic cells contain an endogenous Spy1A-pTRE-Tight gene sequence introduced into said animal model or an ancestor of said animal model at an embryonic stage. Preferably, the Spy1A-pTRE-Tight animal model expresses the Spy1A gene and develop cancer, preferably breast cancer, when administered with tetracycline, preferably doxycycline.

19 Claims, 15 Drawing Sheets

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5



Figure 6




Figure 7




Figure 8



Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26



Figure 27




Figure 28




Figure 29




Figure 30




Figure 31




Figure 32





Figure 33



MMTV-SV40-SPY1A AND SPY1A-PTRE TRANSGENIC MOUSE MODELS

RELATED APPLICATIONS

This application is a continuation-in-part application of earlier U.S. application Ser. No. 13/987,769 filed 30 Aug. 2013, and which claims the benefit of 35 U.S.C. 119(e) to U.S. Provisional Patent Application Ser. No. 61/695,719 filed on Aug. 31, 2012 and U.S. Provisional Patent Application Ser. No. 61/743,501 filed on Sep. 6, 2012.

SCOPE OF THE INVENTION

In one aspect, the present invention provides a transgenic non-human animal model which is selected to overexpress Spy1A under the control of a MMTV promoter causing the animal model to develop cancer, and preferably breast cancer. In another aspect, the present invention provides a transgenic non-human animal model selected to express Spy1A in one or more tissues thereof when an antibiotic is fed to the animal, and leading to the development of cancer, most preferably breast cancer. In another aspect, the present invention provides a transgenic non-human animal model, preferably a mammal and most preferably a mouse model, which includes a MMTV-SV40-Spy1A gene sequence (hereinafter referred interchangeably as "MMTV-SV40-Spy1A", "MMTV-SV40-Spy1", "MMTV-Spy1A" and "MMTV-Spy1"). The animal model of the invention may permit uses in the identification of agents for inhibiting or treating cancer, or namely breast cancer.

BACKGROUND OF THE INVENTION

Amid all cancers known to afflict the Canadian population, breast cancer (BC) is documented as the second leading cause of cancer deaths among females. Current knowledge of the molecular signatures and biochemical pathways that govern BC initiation and progression is far from comprehensive and requires further expansion in order to identify putative biomarkers that undoubtedly predict the correct therapeutic course of action to take with each patient. Due to the heterogeneous nature of cell types which cooperate to form a functional post-natal mammary gland, the various clinical forms of BC that may arise are currently distinguished based on prognostic criteria such as histological phenotype, steroid and growth factor receptor status, and tumor ability to metastasize to neighbouring lymph nodes. In order to fully understand the various molecular mechanisms underpinning the evolution of mammary tumorigenesis, post-pubertal mammary gland development is often looked upon to highlight critical signaling pathways that possess the inherent capacity to mutate and/or become deregulated in BC. Once maturity is established, the adult virgin mammary organ retains the ability to cycle through four development stages: virgin, pregnancy, lactation, followed by involution and reversion to a virgin-like state. During early pregnancy-induced lobuloalveolar development, elevated expression of prolactin, placental lactogens, and progesterone results in escalated rates of luminal epithelial proliferation, and promotes functional differentiation of alveolar precursor cells into specialized structures proficient in milk release. Parturition-induced lactogenesis functions to nourish neonates through alveolar milk production and secretion of colostrums into enlarged luminal ducts. Neonate weaning initiates extensive luminal alveolar cell death (apoptosis) and epithelial remodelling during involu-

tion, a process lasting for several days to allow for reinstatement of the mammary gland to a virgin-like appearance.

SUMMARY OF THE INVENTION

It has been appreciated that at a cellular level Speedy (Spy1A) plays a role in the DNA damage response, functioning to enhance cell survival and promote cell proliferation in lieu of apoptosis. Spy1A is capable of promoting precocious development and tumorigenesis. Hence, determining how Spy1A protein levels are regulated may reveal novel information regarding the dynamics of cell cycle control during normal and abnormal growth conditions. Furthermore, non-degradable forms of Spy1A do not trigger intrinsic cell cycle checkpoints but, rather, promote cell proliferation and oncogenic cell transformation demonstration that this mechanism may contribute to tumorigenesis.

Further, it has been appreciated that Spy1A is a novel cell cycle gene whose product binds to cyclin-dependent kinase-2 (CDK2) and activates its kinase activity to promote cell cycle progression through a cyclin independent mechanism and to promote cell movement into DNA synthesis. Spy1A is expressed naturally at high levels in the proliferating mammary gland, and aberrant overexpression of Spy1A results in precocious mammary development and eventually tumorigenesis in vivo.

Spy1A elevation in c-Myc overexpressing tumors can be maintained during primary tumor culture, the MMTV-Myc mouse model, well documented for its ability to form aggressive mammary adenocarcinomas, may be utilized to derive a previously uncharacterized tumor cell line engineered to overexpress c-Myc (henceforth referred to as F5A1-2).

Induction of the mammary oncogene c-Myc upregulates Spy1A and it is further demonstrated that Spy1A protein levels are elevated in mammary tissue and breast tumors derived from MMTV-Myc transgenic mice. Spy1A knock-down in F5A1-2 cell lines led to downregulation of cyclin-dependent kinase inhibitors (CKI) p21 and p27, a 23% reduction in proliferation rate, and a shift in cellular phenotype to a spindle-like/fibroblastic morphology. Together, findings support that Spy1A plays a functional role in mammary-related c-Myc signal transduction, and acts downstream of ER α , c-Myc, and the MAPK cascade to regulate proliferation, mammary development, and carcinogenesis.

One possible non-limiting object of the present invention is to provide a powerful tool for the study of cancer, namely breast cancer. Specifically, in one aspect the invention provides a transgenic non-human animal model whose somatic cells contain at least one copy of a MMTV-Spy1 transgene causing Spy1A overexpression under the control of the MMTV promoter, and causing this animal model to develop cancer, or more particularly breast cancer. The animal is preferably hemizygous for the transgene.

In another aspect, the present invention provides a transgenic non-human animal model whose somatic cells contain a MMTV-Spy1A transgene which causes the animal model to develop cancer, or preferably breast or liver cancer.

In yet another aspect, the present invention provides a transgenic non-human animal model comprising germ cells and somatic cells containing an exogenous MMTV-SV40-Spy1A gene sequence introduced into said animal model or an ancestor of said animal model at an embryonic stage, wherein said gene sequence comprises a mouse mammary tumor virus gene (MMTV), a functionally disrupted SV40 gene (SV40) and a human Spy1A gene. It has been appreciated that a portion of an SV40 gene when incorporated into a transgenic construct, or preferably the MMTV-SV40-Spy1A

gene sequence increases expression or induce overexpression of a gene under the control of a MMTV promoter.

In a preferred embodiment, the transgenic non-human animal comprises germ cells and somatic cells containing an exogenous MMTV-SV40-Spy1A gene sequence introduced into said animal or an ancestor of said animal, at an embryonic stage, wherein said gene sequences comprises a mouse mammary tumor virus promoter, a functionally disrupted SV40 gene and a human Spy1A gene.

Preferably, the animal model is hemizygous of the MMTV-SV40-Spy1A gene sequence. The human Spy1A gene preferably includes a modified Spy1A gene of SEQ ID NO 1 or a conservatively modified variant thereof. Most preferably, the MMTV-SV40-Spy1A gene sequence is introduced to the animal model or the ancestor by microinjecting a fragment sequence obtained from restriction enzyme digestion of SEQ ID NO: 5 or a conservatively modified variant thereof with XhoI and SpeI.

In yet another aspect, the present invention provides a transgenic non-human animal model which is selected to express Spy1A in one or more tissues thereof when an antibiotic is fed to the animal model, said expression of Spy1A preferably leading to the development of cancer within said animal model, preferably breast or liver cancer.

In yet another aspect, the present invention provides a transgenic non-human animal model whose germ cells and somatic cells contain an exogenous Spy1A-pTRE-Tight gene sequence (hereinafter interchangeably referred to as "Flag-Spy1A-pTRE", "Flag-Spy1A-pTRE-Tight", "Flag-Spy1-pTRE", "Flag-Spy1-pTRE-Tight", "Spy1A-pTRE", "Spy1A-pTRE-Tight", "Spy1-pTRE" and "Spy1-pTRE-Tight") introduced into the animal model or an ancestor of the animal model at an embryonic stage, wherein said gene sequence comprises a human Spy1A gene.

Preferably, the animal model is hemizygous of the Spy1A-pTRE-Tight gene sequence. The human Spy1A gene is preferably a modified Spy1A gene of SEQ ID NO: 1 or a conservatively modified variant thereof. Most preferably, the Spy1A-pTRE-Tight gene sequence is introduced to the animal model or the ancestor by microinjecting a fragment sequence obtained from restriction enzyme digestion of SEQ ID NO 18 or a conservatively modified variant thereof with XhoI and AlwNI.

In a preferred embodiment, the animal model is selected to express the Spy1A gene and develop cancer when administered with a tetracycline. Most preferably, the tetracycline is doxycycline. Preferably, the cancer is breast or liver cancer.

The animal model of the present invention is not strictly limited to those belonging to any specific genus or species, provided that the animal model preferably permits introduction of exogenous genetic sequences to be incorporated into the genome. The animal model is preferably a mouse, a rat, a monkey, a sheep, a dog, a rabbit, or a horse. Most preferably, the animal model is a mouse or a rat.

In yet another aspect, the present invention provides a method of producing the transgenic non-human animal, the method comprising microinjecting the fragment sequence into a fertilized embryo and transplanting said fertilized embryo into a surrogate animal.

In yet another aspect, the present invention provides a tumor cell line comprising a plurality of cells, wherein the cells are derived from the animal model or comprise the fragment sequence.

In yet another aspect, the present invention provides a method for screening an agent for treating or preventing cancer, the method comprising administering the agent into the

animal model or the tumor cell line and detecting size reduction of a tumor caused by the cancer. Preferably, the cancer is breast or liver cancer.

In yet another aspect, the present invention provides a transgenic non-human animal model comprising germ cells and somatic cells having an endogenous MMTV-Spy1A gene sequence, wherein the animal model is a progeny generated by crossing a MMTV-rtTA non-human animal model and the animal model having the Spy1A-pTRE-Tight gene sequence derived from the fragment sequence obtained from restriction enzyme digestion of SEQ ID NO: 18 or a conservatively modified variant thereof with XhoI and AlwNI, the progeny being selected to express the Spy1A gene when administered with a tetracycline.

In yet another aspect, the present invention provides a transgenic non-human animal model, the animal model being a progeny obtained from breeding first and second ancestors, wherein the first ancestor comprises respective germ cells and somatic cells having an ancestor gene sequence introduced into the genome of the first ancestor at an embryonic stage, the ancestor gene sequence comprising a promoter sequence and a tetracycline transactivator (tTA) or reverse tetracycline transactivator (rtTA) gene sequence, and the second ancestor comprises respective germ cells and somatic cells having a Spy1A-pTRE-Tight gene sequence introduced into the genome of the second ancestor at an embryonic stage, the Spy1A-pTRE-Tight gene sequence comprising a human Spy1A gene.

In yet another aspect, the present invention provides a transgenic non-human animal model comprising germ cells and somatic cells having a plurality of gene sequences introduced into the genome of said animal model or an ancestor of said animal model at an embryonic stage, wherein a first one of said gene sequences comprises a promoter sequence and a tetracycline transactivator (tTA) or reverse tetracycline transactivator (rtTA) gene sequence, and a second one of said gene sequences comprises a Spy1A-pTRE-Tight gene sequence, the Spy1A-pTRE-Tight gene sequence comprising a human Spy1A gene.

In a preferred embodiment, the first ancestor is female, and the second ancestor is male. In one embodiment, the animal model comprises germ cells and somatic cells having the ancestor gene sequence and the Spy1A-pTRE-Tight gene sequence. In a further preferred embodiment, the animal model is selected to express the Spy1A gene and develop cancer, or more preferably breast cancer, when administered with a tetracycline, or more preferably a doxycycline.

It is to be appreciated that the promoter sequence is not restricted to any particular promoter sequence. In one embodiment, the promoter sequence is selected to induce transcription of the tTA or rtTA gene sequence in a target organ. Non-limiting examples of the target organ include the brain, heart, skin, liver, stomach, intestine, breast, and reproductive organs. In one embodiment, the promoter sequence comprises a mouse mammary tumor virus (MMTV) gene, and the target organ is the breast. In one embodiment, the ancestor gene sequence or the first gene sequence comprises the rtTA gene sequence.

In one embodiment, the animal model or the second ancestor is hemizygous of the Spy1A-pTRE-Tight gene sequence, and the human Spy1A gene comprises a modified human Spy1A gene of SEQ ID NO 1 or a conservatively modified variant thereof. In one embodiment, the Spy1A-pTRE-Tight gene sequence is introduced into the genome by microinjecting a fragment sequence obtained from restriction enzyme digestion of SEQ ID NO 18 or a conservatively modified variant thereof with XhoI and AlwNI.

Preferably, the conservatively modified variants have at least 70%, 75%, 80%, 85%, 90%, 95% or 99% sequence identity to the specific referenced nucleotide or amino acid sequence. The conservatively modified variants may include point mutations, as well as deletions, substitutions, insertions, transitions, amplifications, inversions, transversions or others of one or more nucleotide bases or amino acid residues.

BRIEF DESCRIPTION OF THE DRAWINGS

Reference may now be had to the following detailed description, taken together with the accompanying drawings in which:

FIG. 1 shows a fusion gene fragment construct for producing a transgenic mouse according to an embodiment of the present invention.

FIG. 2 shows identification of positive founders confirmed through PCR analysis. Positive founders are indicated by the presence of an 825 bp fragment.

FIG. 3 shows GAPDH (100 bp) control for the identification of positive founders as shown in FIG. 2.

FIG. 4 shows identification of SDM-derived mutant Flag-Spy1A-pLXSN constructs upon detecting of a 977 bp fragment following EcoRI digestion of isolated plasmid DNA from each colony: colony 1.1 (lane 1); colony 1.2 (lane 2); colony 1.3 (lane 3); colony 2.1 (lane 4); colony 2.2 (lane 5); and colony 2.3 (lane 6). The pLXSN vector backbone was estimated at 7.0 kb, and the Spy1A insert was estimated at 1.0 kb (997 bp).

FIG. 5 shows EcoRI digestion of the MMTV-SV40-Spy1A transgenic vector releasing the flag-Spy1A coding sequence from the remaining vector backbone. EcoRI digestion of the resultant transgene DNC produced a 977 bp fragment as expected and confirmed successful cloning. The pMMTV-SV40 backbone was estimated at 6.0 kb and the flag-tagged Spy1A insert was estimated at 1.0 kb (997 bp).

FIG. 6 shows digestion of MMTV-SV40-Spy1A prior to microinjection.

FIG. 7 shows detection of a single copy of MMTV-SV40-Spy1A DNA utilizing PCR genotyping methods. Transgene DNA was successfully detected using 8% PAGE in order to verify the success of using the M023/M023 primer set for detection of the Spy1A transgene in tail clip samples. PCR amplification of MMTV-SV40-Spy1A vector DNA (lane 2) using M022/M023 primers produced an 825 bp amplicon, identical to the positive MMTV vector control (+MMTV, lane 1) as expected.

FIG. 8 shows successful transmission of transgene from founder to offspring using primer pair M022 (SEQ ID NO: 2)/M023 (SEQ ID NO: 4) resulting in a 825 bp fragment.

FIG. 9 shows confirmation of germline transmission of transgene using primer pair A933 (SEQ ID NO: 3)/M023 (SEQ ID NO: 4) resulting in a 197 bp fragment.

FIG. 10 shows a Spy1-pTRE vector map according to an embodiment of the present invention.

FIG. 11 shows a restriction digest of XhoI and AlwNI for isolating a portion of the vector illustrated FIG. 10 for a subsequent microinjection step according to an embodiment of the present invention.

FIG. 12 shows identification of Spy1-pTRE founder mice via PCR analysis in the presence of a 536 bp band. The number labels correspond to mouse tag numbers belonging to each tail sample screened, and the label "vector" corresponds to the Spy1-pTRE transgenic vector used as a positive control.

FIG. 13 shows confirmation of successful germline transmission of Spy1-pTRE transgene according to an embodiment of the present invention. The number labels correspond to mouse tag numbers belonging to each tail sample screened, and the label "vector" corresponds to the Spy1-pTRE transgenic vector used as a positive control.

FIG. 14 shows a Flag-Spy1A-pTRE Tight vector map according to an embodiment of the present invention. Primers, Spy1 and pTRE promoter are outlined.

FIG. 15 shows a linearized map of a Flag-Spy1A-pTRE Tight vector indicating the locations of promoter, Spy1 and primers.

FIG. 16 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 0.5 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to pTRE vector control (447 bp); and the lanes "G" correspond to GAPDH (about 100 bp).

FIG. 17 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 1 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to pTRE vector control (447 bp); and the lanes "G" correspond to GAPDH (about 100 bp). The band of correct size is outlined under the lane "3P".

FIG. 18 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 2 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to pTRE vector control (447 bp); and the lanes "G" correspond to GAPDH (about 100 bp). The band of correct size is outlined under the lane "3P".

FIG. 19 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 3 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to pTRE vector control (447 bp); and the lanes "G" correspond to GAPDH (about 100 bp). The band of correct size is outlined under the lane "3P".

FIG. 20 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 0.5 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to a maxi-prepped pTRE vector control with a higher concentration; and the lanes "G" correspond to GAPDH. The band of correct size is outlined under the lane "3P".

FIG. 21 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 1 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P

7

and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to a maxi-prepped pTRE vector control with a higher concentration; and the lanes "G" correspond to GAPDH. The band of correct size is outlined under the lanes "1P" and "3P".

FIG. 22 shows a 4.25% polyacrylamide gel image of Spy1-pTRE DNA samples 1, 2 and 3 amplified with PCR using the primer combination A548/A549 with 2 second exposure time. The lane "L" corresponds to a ladder; the lanes 1P, 2P and 3P correspond to samples 1, 2 and 3, respectively; the lane "CP" corresponds to a maxi-prepped pTRE vector control with a higher concentration; and the lanes "G" correspond to GAPDH. The band of correct size is outlined under the lanes "1P" and "3P".

FIG. 23 shows a bar graph illustrating the results from a qRT PCR analysis test for Spy1 overexpression in the mammary glands of a MMTV-Spy1 mouse in accordance with a preferred embodiment of the invention, and which shows log 10 expression of Spy1 as the Y axis compared to GAPDH.

FIG. 24 shows a DMBA treatment plan for a MMTV-Spy1 mouse and its pair matched littermates, and which indicates age at beginning and end treatment.

FIG. 25 shows a bar graph depicting the percentage of MMTV-Spy1 and control mice (Y axis) that developed all tumour types, mammary tumours, and ovarian tumours.

FIG. 26 shows a line graph depicting the percentage of tumour free mice (Y axis) at the indicated ages in weeks (X axis).

FIG. 27 shows a bar graph depicting the percentage of MMTV-Spy1 and pair matched littermates (F1 cntl) developing hepatocellular carcinoma 1 year of age and older.

FIG. 28 shows a bar graph illustrating the results from a qRT PCR conducted on liver tissue collected from MMTV-Spy1 mice and their pair matched littermates, and which illustrates Spy1 expression on a log 10 scale as compared to GAPDH.

FIG. 29 shows a bar graph illustrating the results from a qRT PCR confirming Spy1 overexpression upon delivery of doxycycline to a MTB-Spy1 mouse generated by crossing a Spy1-pTRE mouse with a MMTV-rtTA mouse.

8

FIG. 30 shows a breeding scheme for a male Spy1-pTRE mouse in accordance with a preferred embodiment of the present invention and a female MMTV-rtTA mouse, and which illustrates possibly genotypes of the resulting progenies.

FIG. 31 shows a bar graph illustrating the results from a qRT PCR confirming elevated Spy1 expression upon delivery of doxycycline to a MTB-Spy1 progeny mouse generated by crossing a male Spy1-pTRE parent mouse with a female MMTV-rtTA parent mouse.

FIG. 32 shows a bar graph illustrating the results from a qRT PCR confirming elevated Spy1 expression upon delivery of doxycycline to a MTB-Spy1 progeny mouse generated by crossing a male Spy1-pTRE parent mouse with a female MMTV-rtTA parent mouse, when compared to the Spy1 expression of the parent mice.

FIG. 33 shows a bar graph illustrating the results from a qRT PCR confirming elevated Spy1 expression in the mammary glands of a female MTB-Spy1 progeny mouse upon delivery of doxycycline thereto, and which is generated by crossing a male Spy1-pTRE parent mouse with a female MMTV-rtTA parent mouse.

DETAILED DESCRIPTION OF THE INVENTION

The gene fragment construct MMTV-SV40-Spy1A (SEQ ID NO: 5) for the development of a transgenic mouse according to a preferred embodiment of the present invention is shown in FIG. 1. The construct was micro injected at roughly 4.7 kb into 357 fertilized embryos from superovulated female mice and transplanted into pseudo pregnant CD-1 female mice. This resulted in 43 pups being born of which 13 tested positively for the MMTV-SV40-Spy1A as confirmed in the PCR analysis shown in FIGS. 2 and 3.

To prepare the MMTV-SV40-Spy1A construct, Flag-Spy1A-pLXSN containing the complete coding sequence of the human Spy1A gene conjugated to a flag tag was provided. Site-directed mutagenesis (SDM) was utilized to create a second EcoRI site positioned near the terminal region of the human Spy1A coding sequence (SEQ ID NO: 1) in Flag-Spy1A-pLXSN for efficient removal of the intrinsic poly-A tail.

```

GAATTCGCGGCCGCGCTCGACCTGCGACGGAGCCTTGACCGCCGTTGCCCGGCCCTCTCCC
GCGCAGCCCCGGGCTTCCGCGAGGAATATTGGGAAACAAAAATGAGGCACAATCAGATGTG
TTGTGAGACACCACCTACTGTCTACTGTTTATGTAATAATCAGGGTCAAAATAGATCACATCA
GCCTAAAAAGCCATTACTCTGAAGCGTCCTATTTGTAAAGATAATTGGCAAGCATTTGA
AAAAAATACACATAATAACAACAAATCTAAACGCCCAAGGACCTTGTCTGGTTATACA
CGGTCAGGATATGACTGCTTTCTTTAAATATTGATGACGATTTAATCAAGATTTCTT
GTGGATGGACTGCTGCTGTAATAATGCAGACAGATATCTTTGGCTATGACCTTTGTTTA
TTTCAAGAGGGCTAAATTTACTATAAGTGAGCATAACAGGATAAATTTCTTTATGCTCT
GTATCTGGCTAATACAGTTGAAGAAGATGAAGAAGAAACCAAGTACGAATTTTTCCATG
GGCTTTAGGGAAAAACTGGAGAAAATTGTTCCCTAATTTCTTAAAGTTAAGGGACCGCT
CTGGGATAGAATTGACTATAGGGCTATTGTAAGCAGGCGATGTTGTGAGGAGTTATGGC
CATTGCACCAACCCATTATATCTGGCAAGAGAACGTTCTGTTCATCACAGTGGAGCTGT
CAGAACTACAACAGAGATGAAGTTTCTAGCTGCCCGGGACCTAGTGCCACACCAGTAGA
TTGTTCACTCTGTGGTAAAAAAGAAGATATGTTAGACTGGGATTGCTTCATCATCATC

```

-continued

TTTATCCAGTCATACAGCAGGGGTGACAGAAAAACATTCTCAGGACTCATACTCACT
 GTCATGGACATAATAGGTGATCCTTCTCAAGCTTATACTGGTTCTGAAGGTATGATATA
 GTAATA

c
 ↑

TGCCAGAATTAGATTATGTCATGTTGTTACTGAGCTCTAGTCAGTCTTTCTGGCGGGG
 ATACATAATAATTTATATACTCCAACAATATGAGTTAAATTAATCTTGAAACTTTCTCCC
 CTTTCAGTTACTTTTTGTCTTGTGTCATATTTGTTTTGTGGTGACCCACCTAAACAGAT
 TTTTAAATGTGACCTATGTTAAGTTGAAAATAATGCACCATAAGCCTCAGTATTTTAAGA
 GCCTGAATCATTTTTTGAAATGTTATTTTATTCAAAGGGTTTCAAGAAGAAAATAAA
 TTTACTTGTAACTCAAAAAAAAAAAAAAAAAAAAAA

SDM primers A424 and A425 (SEQ ID NOs: 6 and 7) were designed to flank the vector region targeted for mutation. SDM reactions were performed with the following components: Flag-Spy1A-pLXSN vector DNA (10-100 ng); 0.3 mM dNTP mix (Cat. No DD0057, Biobasic Inc., Ontario, Canada); 1×pfx buffer and 1 μl pfx polymerase (Cat. No. 11708-013, Invitrogen, Canada); 1 mM MgSO₄; 1 μM each of A424 forward and A425 reverse primers (SEQ ID NOs: 6 and 7); filter-sterilize nuclease free water up to 50.0 μL. Cycling conditions for SDM include (1) 2 minutes at 94° C., (2) 25 cycles of 94° C. for 15 seconds, 55° C. for 30 seconds and 68° C. for 5 seconds, and (3) 68° C. for 10 minutes. SDM reaction products were DpnI digested for 2 hours at 37° C. (Cat. No. ER1701, Fermentas, Burlington, Ontario, Canada), and subsequently transformed utilizing TOP10 *E. coli* and plated onto 100 mg/ml Ampicillin plates. Select colonies were screened for EcoRI insertion and were identified upon detection of a 977 bp fragment following EcoRI digestion (for 20 minutes at 37° C. (Cat. No. FD0274, Fermentas)) of isolated plasmid DNA from each colony (using QIAprep Spin Miniprep Kit (Cat. No 27104, Qiagen, Mississauga, Ontario, Canada) as shown in FIG. 4. Successful EcoRI insertion was confirmed through sequencing for two colonies in particular, Is.1 and IIs.3, utilizing A210 and A211 sequencing primers (SEQ ID NOs: 8 and 9). Purified vector DNA from Colony IIs.3 was subjected to EcoRI digestion (Cat. No. ER0271, Fermentas), and produced two fragments at 7.0 kb (vector backbone) and 1.0 kb (Spy1A gene insert). Digestion products were separated, and the appropriate 1.0 kb fragment was gel extracted using the EZ-10 Spin Column DNA Gel Extraction Kit (Cat. No. BS354, Biobasic Inc.) and purified.

EcoRI digestion of 2 mg of MMTV-SV40-TRPS-1 vector DNA ensued for 1 hour at 37°, followed by the immediate removal of terminal phosphate groups from digested ends utilizing incubation with calf intestinal alkaline phosphatase (Cat. No EF0341, Fermentas) for 30 minutes at 37°. Phosphatase treatment was necessary in order to prevent the religation of linearized vector DNA termini. Consequently, reaction products were separated, followed by gel purification of the resultant 6.0 kb fragment (MMTV-SV40 backbone) using the EZ-10 Spin Column DNA Gel Extraction Kit. Ligation of the Spy1A gene insert into the MMTV-SV40 backbone was conducted using T4 DNA ligase (Cat. No EL0017, Fermentas), and ligation reactions were subsequently transformed utilizing TOP10 *E. coli* and plated onto 100 mg/ml Ampicillin plates. Select colonies were screened for EcoRI insertion and were identified upon detection of a

977 bp fragment following EcoRI digestion of isolated plasmid DNA from each colony as shown in FIG. 5. Successful cloning of the Spy1A coding sequence into the MMTV-SV40 vector backbone was confirmed through sequencing. Sequencing primers A252, A253, A254, A255, A256, A257, A258 and A259 (SEQ ID NOs: 10 to 16) were utilized in order to verify the intactness of all transgenic vector components.

The resultant transgenic vector was designated as MMTV-SV40-Spy1A and contains an untranslated portion of the Ha-ras gene, in addition to an SV40 polyadenylation site. Bacterial sequences such as those found in vector backbones have been noted to inhibit successful incorporation of transgenic DNA into the mouse blastocyst genome. Thus, XhoI/SpeI double digestion (Cat Nos. ER0691 and ER1251, Fermentas) of purified vector DNA (30 mg per tube) ensued, and resulted in the production of two fragments: 4.7 kb (MMTV-SV40-Spy1A transgene) and 2.9 kb (remaining backbone) as shown in FIG. 6. Two vials of XhoI/SpeI digested transgenic DNA were made available for microinjection into mouse blastocysts for subsequent creation of the first MMTV-SV40-Spy1A transgenic mouse model known to date. Transgene detection of a single copy of MMTV-SV40-Spy1A DNA was tested utilizing the PCR conditions outlined for M022 and M023 genotyping primers (SEQ ID NOs: 2 and 4) as shown in FIG. 7.

The resulting transgene fragment was sent to the University of Western Ontario Transgenic Facility to undergo pronuclear injections. Tail samples from the resulting litters were received and DNA was extracted using the Qiagen Puregene Core Kit A for mouse tails. Transgene detection was accomplished using two sets of primers with two unique forward primers (M022 (SEQ ID NO 2) and A933 (SEQ ID NO 3)) and one reverse primer (M023 (SEQ ID NO: 4)). PCR cycling conditions consisted of (1) denaturation at 94° C. for 3 min, (2) denaturation at 94° C. for 1 min, annealing at 55° C. for 2 min, elongation at 72° C. for 1 min and (3) a final elongation step at 72° C. for 3 min. Each 25 μL PCR reaction was made using UBI HP Taq DNA polymerase (HPTAQ-01) and contained a final concentration of 2 ng/μL of pure genomic DNA, 1× buffer, 2 mM MgSO₄, 0.2 mM dNTP, 0.5 mM forward primer, 0.5 mM reverse primer and 0.025 U/μL Taq polymerase. Additionally, a final volume of 1% and 4% DMSO was added for primer pairs M022 (SEQ ID NO: 2)/M023 (SEQ ID NO: 4) and A933 (SEQ ID NO 3)/M023 (SEQ ID NO 4) respectively. PCR amplification resulted in an 825 bp and 197 bp amplicon for primers M022 (SEQ ID NO:

11

2)/M023 (SEQ ID NO: 4) and A933 (SEQ ID NO: 3)/M023 (SEQ ID NO: 4) respectively as shown in FIGS. 8 and 9, respectively.

Expression levels of Spy1A was tested in the inguinal mammary glands of 6 week old MMTV-Spy1A mice and their negative littermates via qRT PCR analysis to ensure Spy1A was being overexpressed in the mammary gland of this mouse model system. Spy1A was found to be significantly overexpressed in the mammary glands of MMTV-Spy1A mice as compared to their control littermates (FIG. 23). To test for increased susceptibility to mammary tumorigenesis, MMTV-Spy1A mice and their negative littermates were treated with 1 mg of 7,12-dimethylbenzanthracene (DMBA) once a week for 6 consecutive weeks beginning at 8 weeks of age via oral gavage. Treatment plan indicating age during treatment and at end of study is illustrated in FIG. 24. Mice were monitored on a weekly basis for the development of mammary tumours via palpitation. MMTV-Spy1A mice were found to develop significantly more mammary tumours than their control littermates (FIG. 25). Additionally, MMTV-Spy1A mice developed tumours earlier than their control littermates (FIG. 26).

When collecting male MMTV-Spy1 mice over the age of 1 year, it was noted there was an increased incidence of liver carcinogenesis in the MMTV-Spy1 mice as compared to their negative control littermates (FIG. 27). Liver tissue was collected from MMTV-Spy1 male mice 1 year of age and older along with pair matched littermate controls and the liver tissue was subjected to qRT PCR analysis to determine Spy1 expression in the liver. Spy1 was found to be significantly overexpressed in MMTV-Spy1 male mice as compared to littermate controls (FIG. 28).

In accordance with another preferred embodiment of the present invention, the fusion gene fragment construct Flag-Spy1A-pTRE-Tight (SEQ ID NO: 18) as illustrated in FIGS. 10, 14 and 15 were prepared. In particular, a Caspase3-pTRE-Tight vector was digested with EcoRI and PvuII to remove Caspase3. A 20 bp linker was then added to close the vector. Site directed mutagenesis was performed on a Flag-Spy1A-pLXSN vector to create an EcoRI restriction enzyme site to enable extraction of Flag-Spy1A from the vector. EcoRI digestion was subsequently performed to remove Flag-Spy1A from the Flag-Spy1A-pLXSN vector. The Flag-Spy1A fragment was then ligated into the pTRE-Tight vector.

Successful preparation of DNA fusion gene fragment construct samples were confirmed by PCR amplification with the primer combination A548/A549 (SEQ ID NOs: 19 and 20) and polyacrylamide gel (as shown in FIGS. 16 to 22) as well as DNA, sequencing. The bands of correct size are outlined under the lanes "1P" and/or "3P" in FIGS. 17 to 22. All tested PCR samples were confirmed by DNA sequencing.

The Spy1-pTRE plasmid were restriction digested using XhoI and AlwNI to isolate a portion for subsequent microinjection into a fertilized embryo from a superovulated female mouse. The digested portion was confirmed by gel electrophoresis as shown in FIG. 11. The digested portion was microinjected into fertilized embryos from superovulated female mice and transplanted into pseudo pregnant CD-1 female mice. Some resulting pups tested positive as confirmed and shown in FIG. 12. Successful germline transmission of the Spy1-pTRE transgene was confirmed as shown in FIG. 13.

The mice having the Spy1-pTRE gene sequence was fed doxycycline to activate expression of Spy1A. Development of cancer including breast cancer was experimentally confirmed.

12

In a separate study, selected Spy1A-pTRE mice found to lack inducible overexpression of Spy1A were nevertheless found to be suitable for preparing overexpressing progenies. In a controlled study, selected Spy1-pTRE mice found to be without inducible overexpression of Spy1A were preferably crossed with MMTV-rtTA mice to generated a MTB-Spy1 mouse model. It has been appreciated that such animal model may permit inducible overexpression of Spy1A preferably after administration of doxycycline to their diet in the form of food pellets. Indeed, expression of Spy1 was induced by administering 2 mg/ml, of doxycycline at 5 weeks of age. Mammary glands were collected at 6 weeks of age from MTB-Spy1, Spy1-pTRE and MMTV-rtTA mice for qRT analysis to test for increased expression of Spy1 in the MTB-Spy1 mouse as compared to the selected control Spy1-pTRE and MMTV-rtTA mice. Spy1 was found to be overexpressed in the MTB-Spy1 mouse, indicating this model system is functioning correctly (FIG. 29).

In an additional study, a male Spy1-pTRE mouse in accordance with a preferred embodiment of the present invention was crossed with a female MMTV-rtTA mouse received from a collaborator, and which is described in Edward J. Funther et al. "A novel doxycycline-inducible system for the transgenic analysis of mammary gland biology". *The FASEB Journal*. 16.3 (2002): 283-292, the entire contents of which are hereby incorporated by reference. The female MMTV-rtTA mouse included the mouse mammary tumor virus gene (MMTV) and a reverse tetracycline transactivator (rtTA), such that the MMTV promoter portion drives the expression of rtTA ("Tet-On"). As illustrated in FIG. 29, four different genotypes were expected from the crossing, or namely a wild type progeny mouse, a Spy1-pTRE progeny mouse, an MMTV-rtTA progeny mouse and the intended MTB-Spy1 progeny mouse, the latter of which includes the transgenic elements from both parent mice. It has been appreciated that the intended MTB-Spy1 progeny mouse may permit for an inducible Tet-On system for expression of the Spy1 gene in the presence of a tetracycline, or preferably doxycycline, and as activated by the rtTA protein. As seen in FIGS. 31 to 33, qRT-PCR analysis confirmed that upon exposure to doxycycline, a MTB-Spy1 progeny female mouse showed elevated Spy1 expression in the mammary glands when compared to a control mouse, and the parent MMTV-rtTA and Spy1A-pTRE mice.

The applicant has appreciated that the present invention provides various advantages and applications, and which include without restriction a transgenic non-human animal model whose somatic cells contain at least one copy of a MMTV-Spy1A transgene causing the animal model to develop cancer.

In yet another aspect, the present invention provides a transgenic non-human animal model all of whose germ cells and somatic cells contain an exogenous MMTV-SV40-Spy1A gene sequence introduced into said mammal, or an ancestor of said mammal, at an embryonic stage wherein said gene sequence comprises a mouse mammary tumor virus gene (MMTV), a functionally disrupted SV40 gene (SV40) and a modified human Spy1A gene of SEQ ID NO: 1.

Other applications of the invention include without restriction:

Methods of screening drugs/vaccines/or other vehicles developed for the prevention of the development of cancer;

The study environmental factors and their effects on the development of cancer;

The study cancer initiated at various stages of the animals development;

13

Methods of screening drugs candidates and their anti-carcinogenic;
 Methods of screening drugs/vaccines/or other vehicles developed for the prevention of the development of cancer;
 The study environmental factors and their effects on the development of cancer; and
 The study of cancer namely breast cancer based on a novel expression of Spy1A initiated within a model animal by feeding the animal doxycycline.
 Additional applications of the invention include, without restriction:
 1. Expression of Spy1A within one or more tissues of the model animal is activated by the animal model ingesting doxycycline (Dox).
 2. The expression of Spy1A results in the tissues of the animal model results in the development of cancer namely breast cancer within that model animal.
 3. A transgenic non-human animal model in this case being a mouse incorporates the condition and promoter response of claims 1 and 2.

14

4. The mouse animal model is able to pass this condition expressed in claims 1 and 2 along to subsequent generations when cross with a mouse not having this condition.
 5. The transgenic non-human animal of claim 1, can be said animal selected from the group consisting of mice, rats, monkeys, sheep, and rabbits.
 6. Analysis of animal model DNA is able to confirm that transgenic condition exists in said animal model.
 7. Transgenic animal model may be used to:
 a. Study cancer
 b. Study cancer initiated at various stages of the animals development
 c. Method of screening drugs candidates and there anti-carcinogenic
 d. Method of screening drugs/vaccines/or other vehicles developed for the prevention of the development of cancer.
 e. Study environmental factors and their effects on the development of cancer

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 20

<210> SEQ ID NO 1

<211> LENGTH: 1303

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

gaattcgcgg ccgcgctgcac ctgcgaacgga gccttgaccg cegttgcccg gccctctccc      60
gcgcagcccc gggcttccgc aggaatatg ggaacccaaa atgaggcaca atcagatgtg      120
ttgtgagaca ccacctactg tcaactgtta tgtaaaatca gggtaaata gatcacatca      180
gcctaaaaag ccctactctc tgaagcgtcc tatttgtaa gataattggc aagcatttga      240
aaaaaataca cataataaca acaaatctaa acgccccaaa ggacctgtc tggttataca      300
gcgtcaggat atgactgctt tctttaaatt attgatgac gatttaattc aagatttctt      360
gtggatggac tgctgctgta aaattgcaga caagtatctt ttggctatga cctttgttta      420
tttcaagagg gctaaattta ctataagtga gcataccagg ataaatttct ttattgctct      480
gtatctggct aatacagttg aagaagatga agaagaaacc aagtacgaaa tttttccatg      540
ggctttaggg aaaaactgga gaaaattggt ccctaatttc ttaaagtaa gggaccagct      600
ctgggataga attgactata gggctattgt aagcaggcga tgttgtgagg aggttatggc      660
cattgcacca acccattata tctggcaaag agaacgttct gttcatcaca gtggagctgt      720
cagaaaactac aacagagatg aagttcagct gccccgggga cctagtgcc aaccagtaga      780
ttgttcactc tgtggtaaaa aaagaagata tgttagactg ggattgtctt catcatcatc      840
tttatccagt catacagcag gggtgacaga aaaacattct caggactcat acaactcact      900
gtcaatggac ataatagggt atccttctca agcttatact ggttctgaag gtatgatata      960
gtaatatgcc agaattcgat ttatgcatgt tgtttactga gctctagtca gtcctttctg     1020
gcggggatag ataataattt atatactcca acaatatgag ttaaattaat cttgaaactt     1080
tctccccttt cagttacttt ttgtcttgty tccatatttg ttttggtytg acccacctaa     1140
acagattttt aatgtgacct atgttaagtt gaaaactaat gcaccataag cctcagtatt     1200
ttaagagcct gaatcatttt tttgaaatgt ttattttatt caaaagggtt tcaagaagaa     1260
ataaatatta cttgtaatct caaaaaaaaa aaaaaaaaaa aaa                               1303

```

-continued

<210> SEQ ID NO 2
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Forward primer 1 (M022)

<400> SEQUENCE: 2
 cccaaggcct aagtaagttt ttgg 24

<210> SEQ ID NO 3
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Forward primer 3 (A933)

<400> SEQUENCE: 3
 cccgctctag tggcagtgtg tt 22

<210> SEQ ID NO 4
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Reverse primer (M023)

<400> SEQUENCE: 4
 gggcataagc acagataaaa cact 24

<210> SEQ ID NO 5
 <211> LENGTH: 7911
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: MMTV-SV40-Spy1A fusion gene fragment construct

<400> SEQUENCE: 5
 cacctaaatt gtaagcgta atattttgtt aaaattcgcg ttaaattttt gttaaatcag 60
 ctctattttt aaccaatagc ccgaaatcgg caaaatccct tataaatcaa aagaatagac 120
 cgagataggg ttgagtgttg ttccagtttg gaacaagagt ccactattaa agaacgtgga 180
 ctccaacgtc aaaggcgcaa aaaccgtcta tcaggggcgat ggcccactac gtgaaccatc 240
 accctaataca agtttttttg ggtcaggtg ccgtaaagca ctaaatcgga accctaaagg 300
 gagccccga tttagagcct gacggggaaa gccggcgaac gtggcgagaa aggaagggaa 360
 gaaagcgaaa ggagcgggcg ctaggggcgt ggcaagtgt gcggtcacgc tgcgcgtaac 420
 caccacaccc gccgcgctta atgcgccgct acaggggcgcg tcccattcgc cattcaggt 480
 gcgcaactgt tgggaagggc gatcgggtgcg ggcctcttcg ctattacgcc agctggcgaa 540
 agggggatgt gctgcaagc gattaagttg ggtaacgcca gggttttccc agtcacgacg 600
 ttgtaaaacg acggccagtg aattgtaata cgactcacta tagggcgaat tgggtaccgg 660
 gccccccctc gaggtcgagc ctctccctta tgcgactcct gcattaggaa gcagcccagt 720
 agtaggttga ggccgttgag caccgccgcc gcaaggaatg gtgcattgcaa ggagatggcg 780
 cccaacagtc ccccgccac ggggcctgcc accatacca cggcgaaca agcgcctcatg 840
 agcccgaagt ggcgagccc atcttcccca toggatgatg cggcgatata ggcgccagca 900
 accgcacctg tggcgccggt gatgccggcc acgatgcgtc cggcgtagag gatcccaatg 960

-continued

atagagattt tactgctcta gttccccata cagaattggt tcgcttagtt gcagcctcaa	1020
gatatcttat tctcaaaagg ccaggatttc aagaacatga catgattcct acatctgcct	1080
gtgttactta cccttatgcc atattattag gattacctca gctaatagat atagagaaaa	1140
gaggatctac ttttcatatt tcctgttctt cttgtagatt gactaattgt ttagattctt	1200
ctgcctacga ctatgcagcg atcatagtca agaggccgcc atacgtgctg ctacctgtag	1260
atattggtga tgaacctagg tttgatgatt ctgccattca aaccttagg tatgccacag	1320
atttaattcg agccaagcga ttcgtcgctg ccattattct gggcatalct gctttaattg	1380
ctattatcac ttcctttgct gtagctacta ctgctttagt taaggagatg caaactgcta	1440
cgtttgttaa taatcttcat aggaatgcta cattagcctt atctgaacaa agaataatag	1500
atttaaaatt agaagctaga cttaatgctt tagaagaagt agttttagag ttgggacaag	1560
atgtggcaaa cttaaagacc agaatgtcca ccagggtgca tgcaaatat gattttatct	1620
gcgttacacc tttaccatat aatgcttctg agagctggga aagaacccaa gctcatttat	1680
tgggcatttg gaatgacaat gagatttcat ataacataca agaattaacc aacctgatta	1740
gtgatatgag caaacaacat attgacacag tggacctcag tggtttggct cagtcctttg	1800
ccaatggagt aaagcttta aatccattag attggacaca atatttcatt tttataggty	1860
ttggagccct gcttttagtc atagtgtta tgattttccc cattgttttc cagtgccttg	1920
cgaagagcct tgaccaagtg cagtcagatc ttaacgtgct tcttttaaaa aagaaaaaag	1980
ggggaatgc cgcgcctgca gcagaaatgg ttgaactccc gagagtgtcc tacacctagg	2040
ggagaagcag ccaaggggtt gtttcccacc aaggacgacc cgtctgcgca caaacggatg	2100
agcccatcag acaaaagacat attcattctc tgctgcaaac ttggcatagc tctgctttgc	2160
ctggggctat tgggggaagt tgccggtcgt gctcgcaggg ctctcacctc tgaactcttt	2220
aatagctctt ctgtgcaaga ttacaatcta aacaattcgg agaactcgac cttcctcctg	2280
aggcaaggac cacagccaac ttcctcttac aagccgcatc gattttgtcc ttcagaaata	2340
gaaaaagaa tgcttgctaa aaattatatt tttaccaata agaccaatcc aataggtaga	2400
ttattagtta ctatgttaag aaatgaatca ttatctttta gtactatctt tactcaaatt	2460
cagaagttag aaatgggaat agaaaataga aagagacgct cacctcaat tgaagaacag	2520
gtgcaaggac tattgaccac aggcctagaa gtaaaaaagg gaaaaaagag tgtttttgtc	2580
aaaataggag acaggtggtg gcaaccaggg acttataggg gaccttacet ctacagacca	2640
acagatgccc cttaccata tacaggaaga tatgacttaa attgggatag gtgggttaca	2700
gtcaatggct ataaagtgtt atatagatcc ctcccctttc gtgaaagact cgcagagct	2760
agacctcctt ggtgatgtt gtctcaagaa gaaaaagacg acatgaaaca acaggtacat	2820
gatttatatt atctaggaac aggaatgcac ttttggggaa agattttcca taccaaggag	2880
gggacagtgg ctggactaat agaacattat tctgcaaaaa cttatggcat gagttattat	2940
gaatagcctt tattggccca accttgcggt tcccagggt taagtaagtt tttggttaca	3000
aactgttctt aaaacagga tgtgagacaa gtggtttcct gacttggtt ggtatcaaag	3060
gttctgatct gagctctgag tgttctatct tcctatgttc ttttgaatt tatccaaatc	3120
ttatgtaaat gcttatgtaa accaagatat aaaagagtgc tgattttttg agtaaaactg	3180
caacagctct aacattcacc tcttgtgtgt ttgtgtctgt tcgccatccc gtctccgctc	3240
gtcacttate cttcacttcc cagaggttcc ccccgagac cccggcgtag aggatccgca	3300
cccttgatga ctccgtctga atttttggtt tcagtttgggt accgaagctg cgcggcgcgt	3360

-continued

ctgcttggtta	cttggttgac	tgttggaaatt	gtttgtcttc	tttgtgaact	gactgtgggt	3420
ttctggacgt	gttgtgtctg	ttagtgtctt	tttgactttt	gtttcgtggt	tgaatttggg	3480
ctgacgactg	tgtttaaaa	cttagaccga	cgactgtggt	tgaaatcatg	aaactgtttg	3540
ctttgttcgt	cgaagagttt	tacttgggtc	ccttaacgct	tagtgagtaa	gaaacttaat	3600
ttttagacc	ccgctctagt	ggcagtgtgt	tggttgatag	ccaaagttaa	tttttaaac	3660
atagtgtttt	gggggttggg	gatttagctc	agtgatagag	ctcttgccca	gcaagcgcaa	3720
ggccctgggt	tccgtcccca	gctctgaaaa	aaaggaaaga	gaaacaaaa	aaaaacatat	3780
agtgttttat	ctgtgcttat	gcccgcagcc	cgagccgcac	ccgccgcca	cgagcccat	3840
gcccggggcc	agtcggcgcc	cgtccgcgcc	cccgcctgc	cccggcccc	gcccccaagc	3900
ttgatataca	attcgcggcc	gcgtcgacct	gcgacggagc	cttgaccgcc	gttgcccggc	3960
cctctccgc	gcagccccg	gcttcgcag	gaatattggg	aaacctatat	ggactacaaa	4020
gacctgacg	gtgattataa	agatcatgat	atcgattaca	aggatgacga	tgacaagagg	4080
cacaatcaga	tgtgttgtga	gacaccacct	actgtcactg	tttatgtaa	atcagggtca	4140
aatagatcac	atcagcctaa	aaagccatt	actctgaagc	gtcctatttg	taaagataat	4200
tggcaagcat	ttgaaaaaaaa	tacacataat	aacaacaaat	ctaaacgcc	caaaggacct	4260
tgtctgggta	tacagcgtca	ggatatgact	gctttcttta	aattatttga	tgacgattta	4320
attcaagatt	tcttgtggat	ggactgctgc	tgtaaaattg	cagacaagta	tcttttggct	4380
atgaccttgg	tttatttcaa	gagggctaaa	tttactataa	gtgagcatac	caggataaat	4440
ttctttattg	ctctgtatct	ggctaataca	gttgaagaag	atgaagaaga	aaccaagtac	4500
gaaatttttc	catgggcttt	agggaaaaac	tggagaaaat	tgttccctaa	tttcttaaag	4560
ttaagggacc	agctctggga	tagaattgac	tatagggcta	ttgtaagcag	gcgatgttgt	4620
gaggagggta	tggccattgc	accaaccat	tatatctggc	aaagagaacg	ttctgttcat	4680
cacagtggag	ctgtcagaaa	ctacaacaga	gatgaagttc	agctgccccg	gggacctagt	4740
gccacaccag	tagattgttc	actctgtggt	aaaaaaagaa	gatatgttag	actgggattg	4800
tcttcatcat	catctttatc	cagtcataca	gcaggggtga	cagaaaaaca	ttctcaggac	4860
tcatacaact	cactgtcaat	ggacataata	ggtgatcctt	ctcaagctta	tactggttct	4920
gaaggtatga	tatagtaata	tgccagaatt	cctgcaggtc	gcggccgcca	ctctagagga	4980
tctttgtgaa	ggaaccttac	ttctgtggtg	tgacataatt	ggacaaaacta	cctacagaga	5040
tttaaagctc	taaggtaaat	ataaaatttt	taagtgtata	atgtgttaaa	ctactgattc	5100
taattgtttg	tgtattttag	attccaacct	atggaactga	tgaatgggag	cagtgtgtga	5160
atgcttttaa	tgaggaaaac	ctgttttgct	cagaagaaat	gccatctagt	gatgatgagg	5220
ctactgctga	ctctcaacat	tctactcctc	caaaaaagaa	gagaaaggta	gaagaccca	5280
aggactttcc	ttcagaattg	ctaagttttt	tgagtcatgc	tgtgtttagt	aatagaactc	5340
ttgcttgctt	tgctatttac	accacaaaag	aaaaagctgc	actgctatac	aagaaaatta	5400
tggaaaaata	tttgatgat	agtgccttga	ctagagatca	taatcagcca	taccacattt	5460
gtagagggtt	tacttgcctt	aaaaaacctc	ccacacctcc	ccctgaacct	gaaacataaa	5520
atgaatgcaa	ttgttgtgt	taacttgttt	attgcagctt	ataatgggta	caaataaagc	5580
aatagcatca	caaatttcac	aaataaagca	tttttttcac	tgacttctag	ttgtggtttg	5640
tccaaactca	tcaatgtatc	ttatcatgct	tggatccact	agttctagag	cgccgccac	5700

-continued

```

cgcggtggag ctccagcttt tgttcccttt agtgagggtt aatttcgagc ttggcgtaat 5760
catggtcata gctgtttcct gtgtgaaatt gttatccgct cacaattcca cacaacatac 5820
gagccggaag cataaagtgt aaagcctggg gtgcctaag agtgagctaa ctacattaa 5880
ttgogttgog ctcaactgccc gctttccagt cgggaaacct gtcgtgccag ctgcattaat 5940
gaatcggccca acgcgcgggg agaggcggtt tgcgtattgg gcgctcttcc gcttctcgc 6000
tcaactgactc gctgcgctcg gtcgttcggc tgcggcgagc ggatcagct cactcaaagg 6060
cggtaatacag gttatccaca gaatcagggg ataacgcagg aaagaacatg tgagcaaaaag 6120
gccagcaaaa ggccaggaac cgtaaaaagg ccgcttgct ggcgtttttc cataggtcc 6180
gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aacccgacag 6240
gactataaag ataccaggcg tttccccctg gaagctccct cgtgcgctct cctgttccga 6300
ccctgcgct taccggatac ctgtccgct tctcccttc ggaagcgtg gcgctttctc 6360
atagctcacg ctgtaggtat ctcaagtcgg ttaggtcgt tcgctccaag ctgggctgtg 6420
tgcacgaacc ccccgctcag ccgaccgct gcgccttacc cggtaactat cgtcttgagt 6480
ccaaccggt aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca 6540
gagcgaggtg ttagggcgtt gctacagagt tcttgaagt gtggcctaac tacggctaca 6600
ctagaaggac agtatttggg atctgogctc tgctgaagcc agttacctc gaaaaagag 6660
ttggtagctc ttgatccggc aaacaaacca ccgctggtag cggtggtttt tttggttgca 6720
agcagcagat tacgcgcaga aaaaaggat ctcaagaaga tcctttgatc ttttctacgg 6780
ggtctgacgc tcagtggaac gaaaactcac gtaagggtt tttggtcatg agattatcaa 6840
aaaggatcct cacctagatc cttttaaatt aaaaatgaag ttttaaatca atctaaagta 6900
tatatgagta aacttggctc gacagttacc aatgcttaat cagtgaggca cctatctcag 6960
cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag ataactacga 7020
tacgggaggg cttaccatct ggccccagtg ctgcaatgat accgcgagac ccacgctcac 7080
cggctccaga tttatcagca ataaaccagc cagccggaag ggcgagcgc agaagtggtc 7140
ctgcaacttt atccgcctcc atccagtcta ttaattgttg ccgggaagct agagtaagta 7200
gttcgccagt taatagtttg cgcaacgttg ttgccattgc tacaggcatc gtgggtgcac 7260
gctcgtcgtt tggatggctt tcattcagct ccggttccca acgatcaagg cgagttacat 7320
gatcccccat gttgtgcaaa aaagcgggta gtccttcgg tcctccgatc gttgtcagaa 7380
gtaagttggc cgcagtgtta tcaactcagc ttatggcagc actgcataat tctcttactg 7440
tcatgccate cgtaagatgc ttttctgtga ctgggtgagta ctcaaccaag tcattctgag 7500
aatagtgtat gcggcgaccg agttgctctt gcccgcgctc aatacgggat aataccgcg 7560
cacatagcag aactttaaaa gtgctcatca ttgaaaaacg ttcttcgggg cgaaaaactc 7620
caaggatcct accgctgttg agatccagtt cgatgtaacc cactcgtgca cccaactgat 7680
cttcagcatc ttttactttc accagcgttt ctgggtgagc aaaaacagga aggcaaatg 7740
ccgcaaaaaa gggaataaag gcgacacgga aatgttgaat actcatactc ttcctttttc 7800
aatattattg aagcatttat cagggttatt gtctcatgag cggatacata tttgaatgta 7860
tttagaaaaa taaacaaata ggggttccgc gcacatttcc ccgaaaagtg c 7911

```

<210> SEQ ID NO 6

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Forward primer (A424)

<400> SEQUENCE: 6

gccagaattc gatttatgca tgttgtttac tgagc 35

<210> SEQ ID NO 7
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse primer (A425)

<400> SEQUENCE: 7

gctcagtaaa caacatgcat aaatcgaatt ctggc 35

<210> SEQ ID NO 8
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer (A210)

<400> SEQUENCE: 8

cccttgaacc tctctgttgc acc 23

<210> SEQ ID NO 9
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse primer (A211)

<400> SEQUENCE: 9

gagcctgggg actttccaca ccc 23

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer F1 (A252)

<400> SEQUENCE: 10

gttttatctg tgcttatgcc 20

<210> SEQ ID NO 11
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse primer F1 (A253)

<400> SEQUENCE: 11

gctcgtatgt tgtgtggaa 19

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward primer F2 (A254)

<400> SEQUENCE: 12

aaccatcacc ctaatcaagt 20

-continued

<210> SEQ ID NO 13
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Reverse primer F2 (A255)

 <400> SEQUENCE: 13
 gtcgccgcat acactatt 18

<210> SEQ ID NO 14
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Forward primer F3 (A256)

 <400> SEQUENCE: 14
 ttatccagtc atacagcagg 20

<210> SEQ ID NO 15
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Reverse primer F3 (A257)

 <400> SEQUENCE: 15
 acccctgctg tatgactgga 20

<210> SEQ ID NO 16
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Forward primer F4 (A258)

 <400> SEQUENCE: 16
 gaccagaatg tccaccagg 19

<210> SEQ ID NO 17
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Reverse primer F4 (A259)

 <400> SEQUENCE: 17
 gccacctctg acttgagcgt 20

<210> SEQ ID NO 18
 <211> LENGTH: 3891
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Flag-Spyla-pTRE-Tight fusion gene fragment
 construct

 <400> SEQUENCE: 18
 ctcgagtta ctccctatca gtgatagaga acgtatgtcg agtttactcc ctatcagtga 60
 tagaacaaga tgtcgagttt actccctatc agtgatagag aacgtatgtc gagtttactc 120
 cctatcagtg atagagaacg tatgtcaggt ttactcccta tcagtgatag agaacgtatg 180
 tcgagtttat ccctatcagt gatagagaac gtatgtcgag tttactccct atcagtgata 240

-continued

gagaacgtat gtcgaggtag gcgtgtacgg tgggaggcct atataagcag agctcgttta	300
gtgaaccgtc agatcgccctg gagaattcgc ggccgcgtcg acctgcgacg gagccttgac	360
cgccgttgcc cggccctctc ccgcgacgcc ccgggcttcc gcaggaatat tgggaaacc	420
atatggacta caaagaccat gacggtgatt ataaagatca tgatatcgat tacaaggatg	480
acgatgacaa gaggcacaat cagatgtggt gtgagacacc acctactgtc actggttatg	540
taaaatcagg gtcaaataga tcacatcagc ctaaaaagcc cactactctg aagcgtccta	600
tttgtaaaga taattggcaa gcatttgaaa aaaatacaca taataacaac aaatctaac	660
gccccaaagg acctgtctg gttatacagc gtcaggatat gactgcttcc tttaaattat	720
ttgatgacga ttaattcaa gatttctgt ggatggactg ctgctgtaa attgcagaca	780
agtatctttt ggctatgacc tttgtttatt tcaagagggc taaatttact ataagtgagc	840
ataccaggat aaatttcttt attgctctgt atctggctaa tacagttgaa gaagatgaag	900
aagaaacca gtacgaatt tttccatggg ctttagggaa aaactggaga aaattgttcc	960
ctaatttctt aaagttaagg gaccagctct gggatagaat tgactatagg gctattgtaa	1020
gcaggcgatg ttgtgaggag gttatggcca ttgcaccaac ccattatc tggcaaagag	1080
aacgttctgt tcatcacagt ggagctgtca gaaactaca cagagatgaa gttcagctgc	1140
ccccgggacc tagtgccaca ccagtagatt gttcactctg tggtaaaaa agaagatag	1200
ttagactggg attgtcttca tcatcatctt tatccagtc tacagcaggg gtgacagaaa	1260
aacattctca ggactcatac aactcactgt caatggacat aataggtgat ccttctcaag	1320
cttatactgg ttctgaaggt atgatatagt aatatgccag aattagattt atgcatgttg	1380
tttactgagc tctagtcagt ctttctggc ggggatacat aataatttat atactccaac	1440
aatatgagtt aaattaatct tgaaaacttc tccccttca gttacttttt gtcttgtgtc	1500
catatttgtt ttgtggtgac ccacctaacc agatttttaa tgtgacctat gttagtga	1560
aaactaatgc accataagcc tcagtattt aagagcctga atcatttttt tgaatgttt	1620
attttattca aaagggtttc aagaagaaaa taaatttact tgtaatctca aaaaaaaaa	1680
aaaaaaaa atctagagga tcataatcag ccataccaca tttgtagagg tttacttgc	1740
tttaaaaaac ctcccacacc tcccctgaa cctgaaacat aaaatgaatg caattgttgt	1800
tgttaaactg tttattgcag cttataatgg ttacaataa agcaatagca tcacaaattt	1860
cacaaataa gcattttttt cactgcctcg agcttcctcg ctcactgact cgctgcgctc	1920
ggctcgttcg ctgcccgcag cggatcagc tcaactcaaag gcggaatac ggttatccac	1980
agaatcaggg gataacgcag gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa	2040
ccgtaaaaag gccgcgttgc tggcgttttt ccataggctc cccccctg acgagcatca	2100
caaaaatcga cgctcaagtc agaggtggcg aaaccgcaca ggactataaa gataccaggc	2160
gtttccccct ggaagctccc tcgtgcgctc tctgttccg acctgcccgc ttaaccgata	2220
cctgtccgcc tttctccctt cgggaagcgt ggcgctttct caatgctcac gctgtaggta	2280
tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt gtgcacgaac cccccgttca	2340
gcccagccgc tgcgccttat ccggtaaacta tcgtcttgag tccaaccggg taagacacga	2400
cttatcgcca ctggcagcag ccaactggtaa caggattagc agagcgaggt atgtaggcgg	2460
tgctacagag ttcttgaagt ggtggcctaa ctacggttac actagaagga cagtatttgg	2520
tatctgcgct ctgctgaagc cagttacctt cggaaaaaga gttggtagct cttgatccgg	2580
caaaaaaacc accgctggta gcggtggttt tttgtttgc aagcagcaga ttaocgcgag	2640

-continued

```

aaaaaaagga tctcaagaag atcctttgat cttttctacg gggctctgacg ctcagtggaa 2700
cgaaaaactca cgtaagggga ttttggctcat gagattatca aaaaggatct tcacctagat 2760
ccttttaaat taaaaatgaa gttttaaatc aatctaaagt atatatgagt aaacttggtc 2820
tgacagttac caatgcttaa tcagtggagc acctatctca gcgatctgtc tatttcgttc 2880
atccatagtt gcctgactcc ccgctgtgta gataactacg atacgggagg gcttaccatc 2940
tggccccagt gctgcaatga taccgogaga cccacgctca cgggctccag atttatcagc 3000
aataaacccag ccagccggaa gggccgagcg cagaagtggc cctgcaactt tatccgctc 3060
catccagtct attaattggt gccgggaagc tagagtaagt agttcgccag ttaatagttt 3120
gcgcaacggt gttgccattg ctacaggcat cgtgggtgtca cgctcgtcgt ttggtatggc 3180
ttcattcagc tccggttccc aacgatcaag gcgagttaca tgatccccc tgttgtgcaa 3240
aaaagcgggt agctccttcg gtcctccgat cgttgtcaga agtaagtgg ccgcagtgtt 3300
atcactcatg gttatggcag cactgcataa ttctcttact gtcattgcat ccgtaagatg 3360
cttttctgtg actggtgagt actcaaccaa gtcattctga gaatagtga tgcggcgacc 3420
gagttgctct tgccccgct caatacggga taataccgcg ccacatagca gaactttaa 3480
agtgtcatc attgaaaaac gttcttcggg gcgaaaaactc tcaaggatct taccgctgtt 3540
gagatccagt tcgatgtaac ccaactgtc acccaactga tcttcagcat cttttacttt 3600
caccagcgtt tctgggtgag caaaaacagg aaggcaaaat gccgcaaaa aggaataag 3660
ggcgacacgg aatggtgaa tactcatact ctctctttt caatattatt gaagcattta 3720
tcagggttat tgtctcatga gcggatacat atttgaatgt atttagaaaa ataacaat 3780
aggggttccg cgcacatttc cccgaaaagt gccactgac gtctaagaaa ccattattat 3840
catgacatta acctataaaa ataggcgtat caccaggccc ttcgtcttc a 3891

```

```

<210> SEQ ID NO 19
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer (A548)

```

```
<400> SEQUENCE: 19
```

```
atcagtgata gagaacgatg tcgagt 26
```

```

<210> SEQ ID NO 20
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer (A549)

```

```
<400> SEQUENCE: 20
```

```
ttgtgcctgt tgtcatcgtc at 22
```

We claim:

1. A transgenic non-human animal model, the animal model being a progeny obtained from breeding first and second ancestors, wherein the first ancestor comprises respective germ cells and somatic cells having an ancestor gene sequence introduced into the genome of the first ancestor at an embryonic stage, the ancestor gene sequence comprising a promoter sequence and a tetracycline transactivator (tTA) or reverse tetracycline transactivator (rtTA) gene sequence, and the second ancestor comprises respective germ cells and

somatic cells having a Spy1A-pTRE-Tight gene sequence introduced into the genome of the second ancestor at an embryonic stage, the Spy1A-pTRE-Tight gene sequence comprising a human Spy1A gene.

2. The transgenic non-human animal model of claim 1, wherein the first ancestor is female, the second ancestor is male, and the animal model comprises germ cells and somatic cells having the ancestor gene sequence and the Spy1A-pTRE-Tight gene sequence.

31

3. The transgenic non-human animal model of claim 1, wherein the second ancestor is hemizygous of the Spy1A-pTRE-Tight gene sequence, and the human Spy1A gene comprises a modified human Spy1A gene of SEQ ID NO: 1 or a conservatively modified variant thereof.

4. The transgenic non-human animal model of claim 1, wherein the Spy1A-pTRE-Tight gene sequence is introduced into the genome of the second ancestor by microinjecting a fragment sequence obtained from restriction enzyme digestion of SEQ ID NO: 18 or a conservatively modified variant thereof with XhoI and A1wNI.

5. The transgenic non-human animal model of claim 1, wherein the promoter sequence is selected to induce transcription of the rTA or rtTA gene sequence in a target organ.

6. The transgenic non-human animal model of claim 5, wherein the target organ is the brain, the heart, the skin, the liver, the stomach, the intestine, the breast, or a reproductive organ.

7. The transgenic non-human animal model of claim 6, wherein the promoter sequence comprises a mouse mammary tumor virus (MMTV) gene, and the target organ is the breast.

8. The transgenic non-human animal model of claim 7, wherein the ancestor gene sequence comprises the rTA gene sequence.

9. The transgenic non-human animal model of claim 1, wherein the animal model is selected to express the Spy1A gene and develop cancer when administered with a tetracycline.

10. The transgenic non-human animal model of claim 9, wherein the tetracycline is doxycycline.

11. A transgenic non-human animal model comprising germ cells and somatic cells having a plurality of gene sequences introduced into the genome of said animal model or an ancestor of said animal model at an embryonic stage, wherein a first one of said gene sequences comprises a promoter sequence and a tetracycline transactivator (tTA) or

32

reverse tetracycline transactivator (rtTA) gene sequence, and a second one of said gene sequences comprises a Spy1A-pTRE-Tight gene sequence, the Spy1A-pTRE-Tight gene sequence comprising a human Spy1A gene.

12. The transgenic non-human animal model of claim 11, wherein the animal model is hemizygous of the Spy1A-pTRE-Tight gene sequence, and the human Spy1A gene comprises a modified human Spy1A gene of SEQ ID NO 1 or a conservatively modified variant thereof.

13. The transgenic non-human animal model of claim 11, wherein the Spy1A-pTRE-Tight gene sequence is introduced into the genome of the animal model or the ancestor by microinjecting a fragment sequence obtained from restriction enzyme digestion of SEQ ID NO: 18 or a conservatively modified variant thereof with XhoI and A1wNI.

14. The transgenic non-human animal model of claim 11, wherein the promoter sequence is selected to induce transcription of the rTA or rtTA gene sequence in a target organ.

15. The transgenic non-human animal model of claim 14, wherein the target organ is the brain, the heart, the skin, the liver, the stomach, the intestine, the breast, or a reproductive organ.

16. The transgenic non-human animal model of claim 15, wherein the promoter sequence comprises a mouse mammary tumor virus (MMTV) gene, and the target organ is the breast.

17. The transgenic non-human animal model of claim 16, wherein the first gene sequence comprises the rTA gene sequence.

18. The transgenic non-human animal model of claim 11, wherein the animal model is selected to express the Spy1A gene and develop cancer when administered with a tetracycline.

19. The transgenic non-human animal model of claim 18, wherein the tetracycline is doxycycline.

* * * * *