

(19) **DANMARK**

(10) **DK/EP 2361012 T3**



(12)

## Oversættelse af europæisk patentskrift

Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **A 01 N 25/02 (2006.01)** **A 01 N 43/40 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-08-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-05-11**
- (86) Europæisk ansøgning nr.: **09744567.0**
- (86) Europæisk indleveringsdag: **2009-10-28**
- (87) Den europæiske ansøgnings publiceringsdag: **2011-08-31**
- (86) International ansøgning nr.: **US2009062360**
- (87) Internationalt publikationsnr.: **WO2010053784**
- (30) Prioritet: **2008-10-29 US 109196 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, IN 46268-1054, USA**
- (72) Opfinder: **LI, Mei, 13679 Cunningham Drive, Westfield, IN 46074, USA**  
**TANK, Holger, 1172 Huntington Woods Point, Zionsville, IN 46077, USA**  
**OLDS, Melissa, 2132 Williams Glen Boulevard, Zionsville, IN 46077, USA**  
**HOPKINS, Derek, 503 Mangorei Road, New Plymouth 4312, New Zealand**  
**BUTTIMOR, Robert, 65 Marama Cres., Spotswood, New Plymouth, 4310, New Zealand**
- (74) Fuldmægtig i Danmark: **Chas. Hude A/S, H.C. Andersens Boulevard 33, 1780 København V, Danmark**
- (54) Benævnelse: **STABILE EMULGERBARE KONCENTRATER INDEHOLDENDE ET FØRSTE HERBICIDT CARBOXYLSYRESALT OG EN ANDEN HERBICID CARBOXYLSYREESTER**
- (56) Fremdragne publikationer:  
**WO-A1-2007/094836**  
**US-A1- 2007 117 721**  
**MANN R K ET AL: "Biology and Control of Tall Ironweed (Vernonia altissima)", WEED SCIENCE, WEED SCIENCE SOCIETY OF AMERICA, CHAMPAIGN, IL, US, vol. 31, no. 3, 1 May 1983 (1983-05-01), pages 324-328, XP009144997, ISSN: 0043-1745**  
**ANDERSON T M D ET AL: "Fireweed response to boomspray applications of different herbicides and adjuvants", PLANT PROTECTION QUARTERLY, INKATA PRESS, NORTH CLAYTON, AU, vol. 10, no. 4, 1 January 1995 (1995-01-01), pages 152-153, XP009144994, ISSN: 0815-2195**



## DESCRIPTION


**[0001]** This application claims the benefit of United States Provisional Application Serial Number 61/109,196 filed on October 29, 2008. This invention concerns a novel stable emulsifiable concentrate containing a first herbicidal carboxylic acid in the salt form and a second herbicidal carboxylic acid in the ester form.

**[0002]** To design an agricultural formulation product, the most important question to be answered is its stability. Failure to meet a set of stability requirements which usually depend on the specific market, application and regulations will certainly lead to failure of its commercialization. There are many causes of formulation instabilities, such as a) chemical instabilities due to reactions between ingredients (actives and/or inerts, etc.), photo-degradations, and oxidations, etc., b) physical instabilities due to phase separations (Oswald ripening, crystallization, sedimentations, creamings, etc.) and c) environmental factors (temperature, humidity/moisture, etc.). In today's agrochemical market, it becomes increasingly common to design formulations to contain multiple active ingredients and their required solvents, safeners, and/or adjuvants, etc., in order to achieve the optimal spectrum, efficacy, and delivery efficiency, which consequently makes formulation stability more and more challenging. Therefore, technologies that can effectively isolate, hinder, or eliminate, adverse reactions or interactions between incompatible ingredients are often critical for a successful product.

**[0003]** The emulsifiable concentrate, a liquid homogeneous formulation to be applied as an emulsion after dilution in water, is one of the most common formulation types for many agricultural products. Emulsifiable concentrates are mixtures of an oil-soluble active ingredient and emulsifying agents dissolved in an organic solvent. The emulsifying agent enables the emulsifiable concentrate to disperse easily in water, thereby forming a "milky" and homogenous emulsion. Emulsifiable concentrates require tank agitation to form the emulsion and maintain it during spraying. However, many challenges may exist, when, for instance, active ingredients may react with one another or with other ingredients in the formulation. For example, a composition containing triclopyr butoxyethyl ester and aminopyralid acid or its potassium salt has been found to be extremely useful for the control of brush and woody plants in range and pasture and industrial vegetation management applications. In some typical emulsifiable concentrate formulations with the hydrophobic ester herbicide and the hydrophilic acid or salt herbicide, only small amounts of the hydrophilic herbicide may dissolve in the organic solvent. In typical emulsifiable concentrate formulations with the hydrophobic ester herbicide and the hydrophilic acid herbicide dissolved in the oil phase, the acid herbicide may decarboxylate upon storage at elevated temperatures. In typical emulsifiable concentrate formulations with the hydrophobic ester herbicide and the hydrophilic salt herbicide dissolved in the oil phase, the hydrophilic salt herbicide may react with the oil-soluble ester herbicide, leading to hydrolysis or transesterification. It would be desirable to have a more stable emulsifiable concentrate containing a first herbicidal carboxylic acid in the salt form and a second herbicidal carboxylic acid in the ester form.

**[0004]** The present invention concerns a stable emulsifiable concentrate which comprises:

1. a) a first carboxylic acid herbicide in the form of a tertiary amine salt in which the tertiary amine has the formula




wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently represent a C<sub>1</sub>-C<sub>18</sub> alkyl, optionally substituted with hydroxyl, amino or alkoxy groups;

2. b) a second carboxylic acid herbicide in the form of a C<sub>4</sub>-C<sub>8</sub> alkyl or alkoxy-substituted alkyl ester;
3. c) an alcohol solvent or co-solvent which is the same as the ester portion of the second carboxylic acid herbicide ester; and
4. d) optionally emulsifiers.

**[0005]** Another aspect of the present invention concerns a method of preparing the emulsifiable concentrate which comprises contacting the first carboxylic acid herbicide in the acid form with the tertiary amine in the alcohol solvent and with the second carboxylic acid herbicide ester and optionally any emulsifiers with stirring until a homogeneous solution is obtained.

**[0006]** This invention provides a stable emulsifiable concentrate containing a first herbicidal carboxylic acid in the salt form and a second herbicidal carboxylic acid in the ester form. The first carboxylic acid herbicide is in the form of a tertiary amine salt in which the tertiary amine has the formula



wherein R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently represent a C<sub>1</sub>-C<sub>18</sub> alkyl, optionally substituted with hydroxyl, amino or alkoxy groups.

Preferably, at least one of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> represents a C<sub>12</sub>-C<sub>18</sub> alkyl group. The tertiary amines include, but are not necessarily limited to, a tertiary alkylamine, a tertiary alkanolamine, a tertiary ethoxylated alkylamine, a tertiary propoxylated alkylamine, a tertiary alkyldiamine, a tertiary ethoxylated alkyldiamine, a tertiary propoxylated alkyldiamine and mixtures thereof. The solvent or co-solvent is an alcohol which is the same as the ester portion of the second carboxylic acid herbicide ester.

**[0007]** The term "alkyl", as used herein, includes within its scope straight chain, branched chain and cyclic hydrocarbon groups, which may be saturated or unsaturated. Unsaturated hydrocarbon groups may contain more than one unsaturated bond.

**[0008]** The herbicidal carboxylic acids useful in the emulsifiable concentrates of the present invention are well known in the art and are described, for example, in The Pesticide Manual, Fourteenth Edition, 2006. Preferred examples of herbicidal carboxylic acids include phosphorus-containing carboxylic acids such as glufosinate and glyphosate; benzoic acid herbicides such as dicamba; phenoxyalkanoic acid herbicides such as 2,4-D, MCPA or 2,4-DB; aryloxyphenoxypropionic acid herbicides such as clodinafop, cyhalofop, fenoxaprop, fluazifop, haloxyfop and quizalofop; and pyridinecarboxylic acid and picolinic acid herbicides such as aminopyralid, clopyralid, fluroxypyr, picloram and triclopyr.

**[0009]** Preferred examples of the first carboxylic acid herbicide in the form of an amine salt include the amine salts of aminopyralid, picloram, clopyralid, and triclopyr.

**[0010]** Preferred examples of the second carboxylic acid herbicide in the form of a C<sub>4</sub>-C<sub>8</sub> alkyl or alkoxy-substituted alkyl ester include triclopyr butotyl (butoxyethyl ester), fluroxypyr meptyl (1-methylheptyl ester), and picloram isooctyl (2-ethylhexyl ester).

**[0011]** The amine salts of a tertiary amine include but are not limited to a tertiary alkylamine, a tertiary alkanolamine, a tertiary ethoxylated alkylamine, a tertiary propoxylated alkylamine, a tertiary alkyldiamine, a tertiary ethoxylated alkyldiamine, a tertiary propoxylated alkyldiamine and mixtures thereof. The tertiary alkylamines as used herein refer to trialkyl amines in which at least one of the alkyl groups is a C<sub>12</sub>-C<sub>18</sub> alkyl group. The Armeen<sup>™</sup> products of Akzo Nobel, such as Armeen DMTD, Armeen DMCD, Armeen DMOD, Armeen DMSD, Armeen M2C, Armeen M2HT, Armeen 312, Armeen 316 and Armeen 380 are typical tertiary alkylamines. The tertiary alkanolamines as used herein refer to trialkyl alkanolamines, such as dimethylethanolamine, diethylethanolamine, triethanolamine, methyldiethanolamine, triisopropanolamine. The tertiary ethoxylated alkylamines as used herein refer to trialkyl amines in which at least one of the alkyl groups is a C<sub>12</sub>-C<sub>18</sub> alkyl group and the remaining alkyl groups are hydroxyethyl groups. The Ethomeen<sup>™</sup> products of Akzo Nobel, such as Ethomeen C/12, Ethomeen C/15, Ethomeen C/25, Ethomeen S/12, Ethomeen S/15, Ethomeen S/25, Ethomeen T/12, Ethomeen T/15, Ethomeen T/20, Ethomeen T/25 and Ethomeen T/30 are typical tertiary ethoxylated alkylamines. The tertiary propoxylated alkylamines as used herein refer to trialkyl amines in which at least one of the alkyl groups is a C<sub>12</sub>-C<sub>18</sub> alkyl group and the remaining alkyl groups are hydroxypropyl groups. The Propomeen<sup>™</sup> products of Akzo Nobel, such as Propomeen O/12 and Propomeen T/12, are typical tertiary propoxylated alkylamines. The tertiary alkyldiamines as used herein refer to tetraalkyl diamines in which at least one of the alkyl groups is a C<sub>12</sub>-C<sub>18</sub> alkyl group. The Duomeen<sup>™</sup> products of Akzo Nobel, such as Duomeen TTM are typical tertiary alkyldiamines. The tertiary ethoxylated alkyldiamines as used herein refer to tetraalkyl diamines in which at least one of the alkyl groups is a C<sub>12</sub>-C<sub>18</sub> alkyl group and the remaining alkyl groups are hydroxyethyl groups. The Ethoduomeen<sup>™</sup> products of Akzo Nobel, such as Ethoduomeen T/13, Ethoduomeen T/20 and Etoduomeen T/25 are typical tertiary ethoxylated alkyldiamines.

**[0012]** The solvent or co-solvent is an alcohol which is the same as the ester portion of the second carboxylic acid herbicide ester. Suitable alcohols may be a C<sub>4</sub>-C<sub>8</sub> alkyl or alkoxy-substituted alkyl alcohols, such as butanol, 2-methylpropanol, 1-methyl heptanol, 2-ethylhexanol, 2-butoxyethanol, 2-butoxy-1-methylethanol, 2-methoxy-1-methylethanol or cyclohexanol.

**[0013]** The emulsifiers used in the present invention are optional and can be anionic, cationic or nonionic in character. Typical emulsifiers include salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C<sub>18</sub> ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C<sub>16</sub> ethoxylate; soaps, such as sodium stearate; alkylnaphthalene-sulfonate salts, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride;

polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono and dialkyl phosphate esters.

**[0014]** The emulsifiable concentrates of the present invention are generally comprised of from 0.01 to 40 weight percent of the first carboxylic acid herbicide in the form of the amine salt, from 0.01 to 70 weight percent of the second carboxylic acid herbicide in the form of the ester, from 1 to 60 weight percent of the alcoholic solvent and from 0 to 20 weight percent of the emulsifiers. More preferably, these emulsifiable concentrates are generally comprised of from 1 to 20 weight percent of the first carboxylic acid herbicide in the form of the amine salt, from 10 to 60 weight percent of the second carboxylic acid herbicide in the form of the ester, from 5 to 50 weight percent of the alcoholic solvent and from 2 to 10 weight percent of the emulsifiers.

**[0015]** In addition to the compositions and uses set forth above, the present invention also embraces the composition and use of these emulsifiable concentrates in combination with one or more additional compatible ingredients. Other additional ingredients may include, for example, one or more other pesticides, dyes, and any other additional ingredients providing functional utility, such as, for example, stabilizers, fragrances, viscosity-modifying additives, suspension aids, dispersants, and freeze-point depressants.

**[0016]** The emulsifiable concentrates are generally prepared by contacting the first carboxylic acid herbicide in the acid form with the tertiary amine in the alcohol solvent and with the second carboxylic acid herbicide ester and optionally any emulsifiers with stirring until a homogeneous solution is obtained. More particularly, while the order of addition is not critical, the tertiary amine can be conveniently added with stirring to a suspension of the first carboxylic acid herbicide in the acid form in the alcohol solvent. Once a clear solution is obtained, the second carboxylic acid herbicide ester and emulsifiers can be added and the resulting mixture stirred until a single phase is obtained.

**[0017]** Examples 1-18 illustrate the present invention.

**Example 1:**

**[0018]** Ethoxylated tallowalkylamine (7.4 g, Ethomeen T/12, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (4.0 g, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (62.3 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (16.5 g, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.3 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.6 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 2:**

**[0019]** Ethoxylated soyaalkylamine (610.8 g, Ethomeen S/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (344.1, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (4698 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (1848.7 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (266.7 g, Ninate 411, from Stepan), EO-PO block copolymer (293.3 g, Toximul 8320, from Stepan), and ethoxylated castor oil (240 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 3:**

**[0020]** Ethoxylated cocoalkylamine (12.58 g, Ethomeen C/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (8.39, purity 95.3%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (49.0 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (34.66 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan) and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 4:**

**[0021]** Ethoxylated cocoalkylamine (12.58 g, Ethomeen C/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (8.39 g, purity 95.3%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (39.1 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (46.22 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan) and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 5:**

**[0022]** Ethoxylated soyaalkylamine (3.34 g, Ethomeen S/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (1.88 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (31.99 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 6:**

**[0023]** Ethoxylated soyaalkylamine (6.68 g, Ethomeen S/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.76 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (27.84 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 7:**

**[0024]** Ethoxylated cocoalkylamine (5.93 g, Ethomeen C/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical clopyralid acid (3.64 g, purity 92.2%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (27.95 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 8:**

**[0025]** Ethoxylated cocoalkylamine (4.72 g, Ethomeen C/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical picloram acid (4.49 g, purity 77.9%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (28.20 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 9:**

**[0026]** Ethoxylated oleylamine (6.70 g, Ethomeen O/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.67 g, purity 95.3%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (27.05 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 10:**

**[0027]** Ethoxylated soyaalkylamine (9.20 g, Ethomeen S/15, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.67 g, purity 95.3%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (27.84 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 11:**

**[0028]** Propoxylated oleylamine (7.23 g, Propomeen O/12, purity 98%, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.67 g, purity 95.3%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (26.51 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.33 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.67 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 12:**

**[0029]** Ethoxylated tallowdiamine (4.37 g, Ethoduomeen T/13, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.76 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (20.0 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences), Aromatic 150 (12.2 g, from ExxonMobil), and emulsifiers including amine salt of dodecylbenzene sulfonate (3.3 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.60 g, Toximul 8320, from Stepan) and ethoxylated castor oil (3.0 g, Agnique CSO 40, from Cognis) were added with stirring until a single phase was obtained.

**Example 13:**

**[0030]** Trimethyl tallowdiamine (3.7 g, Duomeen TTM, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.76 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (17.0 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclopyr butoxyethyl ester (69.32 g, purity 96.3%, from Dow AgroSciences), Aromatic 150 (16.9 g, from ExxonMobil), and emulsifiers including EO-PO block copolymer (3.60 g, Toximul 8320, from Stepan) and ethoxylated castor oil (3.0 g, Agnique CSO 40, from Cognis) were added with stirring until a single phase was obtained.

**Example 14:**

**[0031]** Tallowalkyl-dimethylamine (5.4 g, Armeen DMTD, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.76 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (17.0 g,

Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To above mixture, technical triclop pyr butoxyethyl ester (69.3 g, purity 96.3%, from Dow AgroSciences), Aromatic 150 (15.2 g, from ExxonMobil), and emulsifiers including EO-PO block copolymer (3.60 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Agnique CSO 40, from Cognis) were added with stirring until a single phase was obtained.

**Example 15:**

[0032] Ethoxylated tallowalkylamine (5.47 g, Ethomeen T/12, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.23 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (25 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclop pyr butoxyethyl ester (26 g, purity 96.3%, from Dow AgroSciences), picloram isooctyl (9.25 g, purity 95%, from Dow AgroSciences), methylated vegetable oil (27.7 g, Edenor ME C12-18, from Cognis), and emulsifiers including amine salt of dodecylbenzene sulfonate (3.0 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (4.0 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 16:**

[0033] Ethoxylated tallowalkylamine (5.47 g, Ethomeen T/12, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (3.23 g, purity 93%, from Dow AgroSciences) and ethylene glycol monobutyl ether solvent (25 g, Butyl Cellosolve, from The Dow Chemical Company). Stirring was continued until a clear solution was obtained. To the above mixture, technical triclop pyr butoxyethyl ester (26 g, purity 96.3%, from Dow AgroSciences), picloram isooctyl (9.25 g, purity 95%, from Dow AgroSciences), Aromatic solvent (28.1 g, Aromatic 150ND, from Exxon), and emulsifiers including amine salt of dodecylbenzene sulfonate (3.0 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (4.0 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 17:**

[0034] Ethoxylated soyaalkylamine (8.68 g, Ethomeen S/12, from Akzo Nobel) was added with stirring at  $45 \pm 5$  °C to a suspension of technical aminopyralid acid (5.25 g, purity 95.3%, from Dow AgroSciences), 2-ethylhexanol (20.53 g, from Aldrich), and dimethylalkyl amide (30 g, Agnique KE 3658, from Cognis). Stirring was continued until a clear solution was obtained. To the above mixture, pre-melted technical fluroxypyr meptyl technical (22.1 g, purity 97.7%, from Dow AgroSciences), picloram isooctyl (15.4 g, purity 95%, from Dow AgroSciences) and emulsifiers including amine salt of dodecylbenzene sulfonate (3.3 g, Bio-Soft N-411, from Stepan), EO-PO block copolymer (3.7 g, Toximul 8320, from Stepan), and ethoxylated castor oil (3.0 g, Toximul 8242, from Stepan) were added with stirring until a single phase was obtained.

**Example 18:**

[0035] The stability of the emulsifiable concentrates prepared in Examples 1-17 were assessed in accelerated storage tests after 2 weeks at 54 °C and after 8 weeks at 40 °C. The results are summarized in Table 1. All examples demonstrate active ingredient stability of at least 95.0% after 2 weeks accelerated storage at 54 °C or 8 weeks accelerated storage at 40 °C.

Table 1: Chemical Stability of Examples 1-17 During Storage at Accelerated Storage Temperature

| Example        | % Retention of Active 1 |               | % Retention of Active 2 |               | % Retention of Active 3 |               |
|----------------|-------------------------|---------------|-------------------------|---------------|-------------------------|---------------|
|                | 2-wk at 54 °C           | 8-wk at 40 °C | 2-wk at 54 °C           | 8-wk at 40 °C | 2-wk at 54 °C           | 8-wk at 40 °C |
| 1 <sup>a</sup> | 96.7                    | 96.2          | 99.9                    | 99.9          |                         |               |
| 2 <sup>a</sup> | 99.9                    | 99.5          | 99.5                    | 98.1          |                         |               |
| 3 <sup>a</sup> | 97.9                    | 96.6          | 98.0                    | 96.7          |                         |               |
| 4 <sup>a</sup> | 99.7                    | 95.1          | 96.4                    | 95.2          |                         |               |
| 5 <sup>a</sup> | 96.7                    | 98.0          | 98.6                    | 97.6          |                         |               |

| Example                                                                             | % Retention of Active 1 |               | % Retention of Active 2 |               | % Retention of Active 3 |               |
|-------------------------------------------------------------------------------------|-------------------------|---------------|-------------------------|---------------|-------------------------|---------------|
|                                                                                     | 2-wk at 54 °C           | 8-wk at 40 °C | 2-wk at 54 °C           | 8-wk at 40 °C | 2-wk at 54 °C           | 8-wk at 40 °C |
| 6 <sup>a</sup>                                                                      | 99.9                    | 99.5          | 99.5                    | 98.1          |                         |               |
| 7 <sup>a</sup>                                                                      | 99.3                    | 97.7          | 99.1                    | 95.7          |                         |               |
| 8 <sup>a</sup>                                                                      | 98.7                    | n/a           | 98.8                    | n/a           |                         |               |
| 9 <sup>a</sup>                                                                      | 96.0                    | 95.3          | 95.9                    | 95.1          |                         |               |
| 10 <sup>a</sup>                                                                     | 96.2                    | 95.8          | 95.7                    | 95.1          |                         |               |
| 11 <sup>a</sup>                                                                     | 96.7                    | 96.7          | 95.9                    | 95.1          |                         |               |
| 15 <sup>b</sup>                                                                     | 97.9                    | n/a           | 95.0                    | n/a           | 97.0                    | n/a           |
| 16 <sup>b</sup>                                                                     | 99.9                    | n/a           | 95.5                    | n/a           | 97.8                    | n/a           |
| 17 <sup>c</sup>                                                                     | 99.9                    | n/a           | 99.9                    | n/a           | 99.9                    | n/a           |
| a Active 1: aminopyralid; Active 2: triclopyr butotyl;                              |                         |               |                         |               |                         |               |
| b Active 1: aminopyralid; Active 2: picloram isooctyl; Active 3: triclopyr butotyl; |                         |               |                         |               |                         |               |
| c Active 1: aminopyralid; Active 2: picloram isooctyl; Active 3: fluroxypyr meptyl. |                         |               |                         |               |                         |               |

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- US61109196B [0001]


### Non-patent literature cited in the description

- The Pesticide Manual20060000 [0008]

**Patentkrav**

1. Stabilt emulgerbart koncentrat, der indeholder:

- 5 a) et første carboxylsyreherbicid i form af et tertiært aminsalt, hvori den tertiære amin har formlen



i hvilken  $R^1$ ,  $R^2$  og  $R^3$  uafhængigt betyder en  $C_1$ - $C_{18}$ -alkylgruppe, eventuelt substitueret med hydroxyl-, amino- eller alkoxygrupper;


10

- b) et andet carboxylsyreherbicid i form af en  $C_4$ - $C_8$ -alkyl- eller alkoxysubstitueret alkylester;

- c) et alkoholopløsningsmiddel eller co-opløsningsmiddel, der er det samme  
15 som esterdelen af den anden carboxylsyreherbicidester; og

- d) eventuelt emulgatorer.

2. Stabilt emulgerbart koncentrat ifølge krav 1, i hvilket den tertiære amin  
20 har formlen



i hvilken mindst én af  $R^1$ ,  $R^2$  og  $R^3$  betyder en  $C_{12}$ - $C_{18}$ -alkylgruppe.

3. Stabilt emulgerbart koncentrat ifølge krav 1, i hvilket det første  
25 carboxylsyreherbicid i form af et tertiært aminsalt er et tertiært aminsalt af aminopyralid, picloram, clopyralid eller triclopyr.

4. Stabilt emulgerbart koncentrat ifølge krav 1, i hvilket det tertiære aminsalt  
30 er en tertiær alkylamin, en tertiær alkanolamin, en tertiær ethoxileret alkylamin, en tertiær propoxyleret alkylamin, en tertiær alkylldiamin, en tertiær ethoxileret

alkyldiamin, en tertiær propoxyleret alkyldiamin og blandinger deraf.

5.      Stabilt emulgerbart koncentrat ifølge krav 1, i hvilket det andet  
carboxylsyreherbicide i form af en C<sub>4</sub>-C<sub>8</sub> alkyl- eller alkoxy-substitueret alkylester  
5 er en ester af triclopyr, fluroxypyr eller picloram.

6.      Stabilt emulgerbart koncentrat ifølge krav 1, i hvilket det andet  
carboxylsyreherbicide i form af en C<sub>4</sub>-C<sub>8</sub>-alkyl- eller alkoxy-substitueret alkylester  
er en butotyl-, meptyl- eller isooctylester.  
10

7.      Stabilt emulgerbart koncentrat ifølge krav 1, der indeholder fra 0,01 til 40  
vægt-% af det første carboxylsyreherbicide i form af aminsaltet, fra 0,01 til 70  
vægt-% af det andet carboxylsyreherbicide i form af esteren, fra 1 til 60 vægt-%  
af det alkoholiske opløsningsmiddel og fra 0 til 20 vægt-% af emulgatorerne.  
15

8.      Stabilt emulgerbart koncentrat ifølge krav 1, der indeholder fra 1 til 20  
vægt-% af det første carboxylsyreherbicide i form af aminsaltet, fra 10 til 60  
vægt-% af det andet carboxylsyreherbicide i form af esteren, fra 5 til 50 vægt-%  
af det alkoholiske opløsningsmiddel og fra 2 til 10 vægt-% af emulgatorerne.  
20

9.      Fremgangsmåde til fremstilling af det emulgerbare koncentrat ifølge krav  
1, hvilken fremgangsmåde omfatter kontakt mellem det første  
carboxylsyreherbicide på syreformen og den tertiære amin i  
alkoholopløsningsmidlet og den anden carboxylsyreherbicidester og eventuelt  
25 tilstedeværende emulgatorer under omrøring, indtil der opnås en homogen  
opløsning.