



HU000028230T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 028 230**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 12 716818**
(22) A bejelentés napja: **2012. 03. 28.**
(96) Az európai bejelentés bejelentési száma:
EP 20120716818
(97) Az európai bejelentés közzétételi adatai:
EP 2691416 A1 **2012. 10. 04.**
(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2691416 B1 **2016. 05. 18.**
(51) Int. Cl.: **C07K 16/00** (2006.01)
C07K 16/24 (2006.01)
C07K 16/28 (2006.01)
C07K 16/32 (2006.01)
C07K 16/46 (2006.01)
(86) A nemzetközi (PCT) bejelentési szám:
PCT/US 12/030948
(87) A nemzetközi közzétételi szám:
WO 12135345

(30) Elsőbbségi adatok: 201161468276 P 2011. 03. 28. US 1160311 2011. 11. 14. FR	(73) Jogosult(ak): SANOFI, 75008 Paris (FR)
(72) Feltaláló(k): BAURIN, Nicolas, 75008 Paris (FR) BEIL, Christian, 65926 Frankfurt am Main (DE) CORVEY, Carsten, 65926 Frankfurt am Main (DE) LANGE, Christian, 65926 Frankfurt am Main (DE) LI, Danxi, Bridgewater, NJ 08807 (US) MIKOL, Vincent, 75008 Paris (FR) STEINMETZ, Anke, 75008 Paris (FR) RAO, Ercole, 65926 Frankfurt am Main (DE)	(74) Képviselő: Danubia Szabadalmi és Jogi Iroda Kft., Budapest

- (54) **Kettős variábilis régióval rendelkező ellenanyagszerű fehérjék keresztezett kötőrégiós orientációval**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 691 416 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
18.05.2016 Bulletin 2016/20

(51) Int Cl.:
C07K 16/00 ^(2006.01) **C07K 16/24** ^(2006.01)
C07K 16/28 ^(2006.01) **C07K 16/32** ^(2006.01)
C07K 16/46 ^(2006.01)

(21) Application number: **12716818.5**

(86) International application number:
PCT/US2012/030948

(22) Date of filing: **28.03.2012**

(87) International publication number:
WO 2012/135345 (04.10.2012 Gazette 2012/40)

(54) **DUAL VARIABLE REGION ANTIBODY-LIKE BINDING PROTEINS HAVING CROSS-OVER
BINDING REGION ORIENTATION**

ANTIKÖRPERÄHNLICHE BINDUNGSPROTEINE MIT ZWEIFACH VARIABLEN REGIONEN UND
ÜBERKREUZTER AUSRICHTUNG DER BINDUNGSREGION

PROTÉINES DE LIAISON À DES ANTICORPS-LIKE À DOUBLE RÉGION VARIABLE AYANT UNE
ORIENTATION CROISÉE DE RÉGION DE LIAISON

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME

- **LI, Danxi**
Bridgewater, NJ 08807 (US)
- **MIKOL, Vincent**
75008 Paris (FR)
- **STEINMETZ, Anke**
75008 Paris (FR)
- **RAO, Ercole**
65926 Frankfurt am Main (DE)

(30) Priority: **28.03.2011 US 201161468276 P**
14.11.2011 FR 1160311

(43) Date of publication of application:
05.02.2014 Bulletin 2014/06

(74) Representative: **Zwicker, Jörk**
ZSP Patentanwälte PartG mbB
Radtkoferstrasse 2
81373 München (DE)

(73) Proprietor: **SANOFI**
75008 Paris (FR)

(56) References cited:
EP-A1- 2 050 764 WO-A1-97/14719
WO-A2-2011/014659

(72) Inventors:

- **BAURIN, Nicolas**
75008 Paris (FR)
- **BEIL, Christian**
65926 Frankfurt am Main (DE)
- **CORVEY, Carsten**
65926 Frankfurt am Main (DE)
- **LANGE, Christian**
65926 Frankfurt am Main (DE)

- **CUESTA A M ET AL: "Multivalent antibodies:
when design surpasses evolution", TRENDS IN
BIOTECHNOLOGY, ELSEVIER PUBLICATIONS,
CAMBRIDGE, GB, vol. 28, no. 7, 1 July 2010
(2010-07-01), pages 355-362, XP027102411, ISSN:
0167-7799 [retrieved on 2010-05-04]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 691 416 B1

Description**FIELD OF THE INVENTION**

5 **[0001]** The invention relates to antibody-like binding proteins comprising four polypeptide chains that form four antigen binding sites, wherein each pair of polypeptides forming the antibody-like binding protein possesses dual variable domains having a cross-over orientation. Also disclosed are methods for making such antigen-like binding proteins.

BACKGROUND OF THE INVENTION

10 **[0002]** Naturally occurring IgG antibodies are bivalent and monospecific. Bispecific antibodies having binding specificities for two different antigens can be produced using recombinant technologies and are projected to have broad clinical applications. It is well known that complete IgG antibody molecules are Y-shaped molecules comprising four polypeptide chains: two heavy chains and two light chains. Each light chain consists of two domains, the N-terminal domain being known as the variable or V_L domain (or region) and the C-terminal domain being known as the constant (or C_L) domain (constant kappa (C_K) or constant lambda (C_λ) domain). Each heavy chain consists of four or five domains, depending on the class of the antibody. The N-terminal domain is known as the variable (or V_H) domain (or region), which is followed by the first constant (or C_{H1}) domain, the hinge region, and then the second and third constant (or C_{H2} and C_{H3}) domains. In an assembled antibody, the V_L and V_H domains associate together to form an antigen binding site. Also, the C_L and C_{H1} domains associate together to keep one heavy chain associated with one light chain. The two heavy-light chain heterodimers associate together by interaction of the C_{H2} and C_{H3} domains and interaction between the hinge regions on the two heavy chains.

20 **[0003]** It is known that proteolytic digestion of an antibody can lead to the production of antibody fragments (Fab and Fab2). Such fragments of the whole antibody can exhibit antigen binding activity. Antibody fragments can also be produced recombinantly. Fv fragments, consisting only of the variable domains of the heavy and light chains associated with each other may be obtained. These Fv fragments are monovalent for antigen binding. Smaller fragments such as individual variable domains (domain antibodies or dABs; Ward et al., 1989, Nature 341(6242): 544-46), and individual complementarity determining regions or CDRs (Williams et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86(14): 5537-41) have also been shown to retain the binding characteristics of the parent antibody, although most naturally occurring antibodies generally need both a V_H and V_L to retain full binding potency.

30 **[0004]** Single chain variable fragment (scFv) constructs comprise a V_H and a V_L domain of an antibody contained in a single polypeptide chain wherein the domains are separated by a flexible linker of sufficient length (more than 12 amino acid residues), that forces intramolecular interaction, allowing self-assembly of the two domains into a functional epitope binding site (Bird et al., 1988, Science 242(4877): 423-26). These small proteins (MW ~25,000 Da) generally retain specificity and affinity for their antigen in a single polypeptide and can provide a convenient building block for larger, antigen-specific molecules.

35 **[0005]** An advantage of using antibody fragments rather than whole antibodies in diagnosis and therapy lies in their smaller size. They are likely to be less immunogenic than whole antibodies and more able to penetrate tissues. A disadvantage associated with the use of such fragments is that they have only one antigen binding site, leading to reduced avidity. In addition, due to their small size, they are cleared very fast from the serum, and hence display a short half-life.

40 **[0006]** It has been of interest to produce bispecific antibodies (BsAbs) that combine the antigen binding sites of two antibodies within a single molecule, and therefore, would be able to bind two different antigens simultaneously. Besides applications for diagnostic purposes, such molecules pave the way for new therapeutic applications, e.g., by redirecting potent effector systems to diseased areas (where cancerous cells often develop mechanisms to suppress normal immune responses triggered by monoclonal antibodies, like antibody-dependent cellular cytotoxicity (ADCC) or complement-dependent cytotoxicity (CDC)), or by increasing neutralizing or stimulating activities of antibodies. This potential was recognized early on, leading to a number of approaches for obtaining such bispecific antibodies. Initial attempts to couple the binding specificities of two whole antibodies against different target antigens for therapeutic purposes utilized chemically fused heteroconjugate molecules (Staerz et al., 1985, Nature 314(6012): 628-31).

45 **[0007]** Bispecific antibodies were originally made by fusing two hybridomas, each capable of producing a different immunoglobulin (Milstein et al., 1983, Nature 305(5934): 537-40), but the complexity of species (up to ten different species) produced in cell culture made purification difficult and expensive (George et al., 1997, THE ANTIBODIES 4: 99-141 (Capra et al., ed., Harwood Academic Publishers)). Using this format, a mouse IgG2a and a rat IgG2b antibody were produced together in the same cell (e.g., either as a quadroma fusion of two hybridomas, or in engineered CHO cells). Because the light chains of each antibody associate preferentially with the heavy chains of their cognate species, three major species of antibody are assembled: the two parental antibodies, and a heterodimer of the two antibodies comprising one heavy/light chain pair of each, associating via their Fc portions. The desired heterodimer can be purified

from this mixture because its binding properties to Protein A are different from those of the parental antibodies: rat IgG2b does not bind to Protein A, whereas the mouse IgG2a does. Consequently, the mouse-rat heterodimer binds to Protein A but elutes at a higher pH than the mouse IgG2a homodimer, and this makes selective purification of the bispecific heterodimer possible (Lindhofer et al., 1995, J. Immunol. 155(1): 219-25). The resulting bispecific heterodimer is fully non-human, hence highly immunogenic, which could have deleterious side effects (e.g., "HAMA" or "HARA" reactions), and/or neutralize the therapeutic. There remained a need for engineered bispecifics with superior properties that can be readily produced in high yield from mammalian cell culture.

[0008] Despite the promising results obtained using heteroconjugates or bispecific antibodies produced from cell fusions as cited above, several factors made them impractical for large scale therapeutic applications. Such factors include: rapid clearance of heteroconjugates *in vivo*, the laboratory intensive techniques required for generating either type of molecule, the need for extensive purification of heteroconjugates away from homoconjugates or mono-specific antibodies, and the generally low yields obtained.

[0009] Genetic engineering has been used with increasing frequency to design, modify, and produce antibodies or antibody derivatives with a desired set of binding properties and effector functions. A variety of recombinant methods have been developed for efficient production of BsAbs, both as antibody fragments (Carter et al., 1995, J. Hematother. 4(5): 463-70; Pluckthun et al., 1997, Immunotechnology 3(2): 83-105; Todorovska et al., 2001, J. Immunol. Methods 248(1-2): 47-66) and full length IgG formats (Carter, 2001, J. Immunol. Methods 248(1-2): 7-15).

[0010] Combining two different scFvs results in BsAb formats with minimal molecular mass, termed sc-BsAbs or Ta-scFvs (Mack et al., 1995, Proc. Natl. Acad. Sci. U.S.A. 92(15): 7021-25; Mallender et al., 1994, J. Biol. Chem. 269(1): 199-206). BsAbs have been constructed by genetically fusing two scFvs to a dimerization functionality such as a leucine zipper (Kostelny et al., 1992, J. Immunol. 148(5): 1547-53; de Kruif et al., 1996, J. Biol. Chem. 271(13): 7630-34).

[0011] Diabodies are small bivalent and bispecific antibody fragments. The fragments comprise a V_H connected to a V_L on the same polypeptide chain, by using a linker that is too short (less than 12 amino acid residues) to allow pairing between the two domains on the same chain. The domains are forced to pair intermolecularly with the complementary domains of another chain and create two antigen-binding sites. These dimeric antibody fragments, or "diabodies," are bivalent and bispecific (Holliger et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90(14): 6444-48). Diabodies are similar in size to a Fab fragment. Polypeptide chains of V_H and V_L domains joined with a linker of between 3 and 12 amino acid residues form predominantly dimers (diabodies), whereas with a linker of between 0 and 2 amino acid residues, trimers (triabodies) and tetramers (tetrabodies) predominate. In addition to the linker length, the exact pattern of oligomerization seems to depend on the composition as well as the orientation of the variable domains (Hudson et al., 1999, J. Immunol. Methods 231(1-2): 177-89). The predictability of the final structure of diabody molecules is very poor.

[0012] Although sc-BsAb and diabody-based constructs display interesting clinical potential, it was shown that such non-covalently associated molecules are not sufficiently stable under physiological conditions. The overall stability of a scFv fragment depends on the intrinsic stability of the V_L and V_H domains as well as on the stability of the domain interface. Insufficient stability of the V_H - V_L interface of scFv fragments has often been suggested as a main cause of irreversible scFv inactivation, since transient opening of the interface, which would be allowed by the peptide linker, exposes hydrophobic patches that favor aggregation and therefore instability and poor production yield (Wörn et al., 2001, J. Mol. Biol. 305(5): 989-1010).

[0013] An alternative method of manufacturing bispecific bivalent antigen-binding proteins from V_H and V_L domains is described in U.S. Patent No. 5,989,830. Such double head and dual Fv configurations are obtained by expressing a bicistronic vector, which encodes two polypeptide chains. In the Dual-Fv configuration, the variable domains of two different antibodies are expressed in a tandem orientation on two separate chains (one heavy chain and one light chain), wherein one polypeptide chain has two times a V_H in series separated by a peptide linker (V_{H1} -linker- V_{H2}) and the other polypeptide chain consists of complementary V_L domains connected in series by a peptide linker (V_{L1} -linker- V_{L2}). In the cross-over double head configuration, the variable domains of two different antibodies are expressed in a tandem orientation on two separate polypeptide chains (one heavy chain and one light chain), wherein one polypeptide chain has two times a V_H in series separated by a peptide linker (V_{H1} -linker- V_{H2}) and the other polypeptide chain consists of complementary V_L domains connected in series by a peptide linker in the opposite orientation (V_{L2} -linker- V_{L1}). Molecular modeling of the constructs suggested the linker size to be long enough to span 30-40 Å (15-20 amino acid residues).

[0014] Increasing the valency of an antibody is of interest as it enhances the functional affinity of that antibody due to the avidity effect. Polyvalent protein complexes (PPC) with an increased valency are described in U.S. Patent Application Publication No. US 2005/0003403 A1. PPCs comprise two polypeptide chains generally arranged laterally to one another. Each polypeptide chain typically comprises three or four "v-regions," which comprise amino acid sequences capable of forming an antigen binding site when matched with a corresponding v-region on the opposite polypeptide chain. Up to about six "v-regions" can be used on each polypeptide chain. The v-regions of each polypeptide chain are connected linearly to one another and may be connected by interspersed linking regions. When arranged in the form of the PPC, the v-regions on each polypeptide chain form individual antigen binding sites. The complex may contain one or several binding specificities.

[0015] A strategy was proposed by Carter *et al.* (Ridgway *et al.*, 1996, Protein Eng. 9(7): 617-21; Carter, 2011, J. Immunol. Methods 248(1-2): 7-15) to produce a Fc heterodimer using a set of "knob-into-hole" mutations in the C_{H3} domain of Fc. These mutations lead to the alteration of residue packing complementarity between the C_{H3} domain interface within the structurally conserved hydrophobic core so that formation of the heterodimer is favored as compared with homodimers, which achieves good heterodimer expression from mammalian cell culture. Although the strategy led to higher heterodimer yield, the homodimers were not completely suppressed (Merchant *et al.*, 1998, Nat. Biotechnol. 16(7): 677-81).

[0016] Gunasekaran *et al.* explored the feasibility of retaining the hydrophobic core integrity while driving the formation of Fc heterodimer by changing the charge complementarity at the C_{H3} domain interface (Gunasekaran *et al.*, 2010, J. Biol. Chem. 285(25): 19637-46). Taking advantage of the electrostatic steering mechanism, these constructs showed efficient promotion of Fc heterodimer formation with minimum contamination of homodimers through mutation of two pairs of peripherally located charged residues. In contrast to the knob-into-hole design, the homodimers were evenly suppressed due to the nature of the electrostatic repulsive mechanism, but not totally avoided.

[0017] Davis *et al.* describe an antibody engineering approach to convert Fc homodimers into heterodimers by interdigitating β -stand segments of human IgG and IgA C_{H3} domains, without the introduction of extra interchain disulfide bonds (Davis *et al.*, 2010, Protein Eng. Des. Sel. 23(4): 195-202). Expression of SEEDbody (Sb) fusion proteins by mammalian cells yields Sb heterodimers in high yield that are readily purified to eliminate minor by-products.

[0018] U.S. Patent Application Publication No. US 2010/331527 A1 describes a bispecific antibody based on heterodimerization of the C_{H3} domain, introducing in one heavy chain the mutations H95R and Y96F within the C_{H3} domain. These amino acid substitutions originate from the C_{H3} domain of the IgG3 subtype and will heterodimerize with an IgG1 backbone. A common light chain prone to pair with every heavy chain is a prerequisite for all formats based on heterodimerization through the C_{H3} domain. A total of three types of antibodies are therefore produced: 50% having a pure IgG1 backbone, one-third having a pure H95R and Y96F mutated backbone, and one-third having two different heavy chains (bispecific). The desired heterodimer can be purified from this mixture because its binding properties to Protein A are different from those of the parental antibodies: IgG3-derived C_{H3} domains do not bind to Protein A, whereas the IgG1 does. Consequently, the heterodimer binds to Protein A, but elutes at a higher pH than the pure IgG1 homodimer, and this makes selective purification of the bispecific heterodimer possible.

[0019] U.S. Patent No. 7,612,181 describes a Dual-Variable-Domain IgG (DVD-IgG) bispecific antibody that is based on the Dual-Fv format described in U.S. Patent No. 5,989,830. A similar bispecific format was also described in U.S. Patent Application Publication No. US 2010/0226923 A1. The addition of constant domains to respective chains of the Dual-Fv (C_{H1}-Fc to the heavy chain and kappa or lambda constant domain to the light chain) led to functional bispecific antibodies without any need for additional modifications (*i.e.*, obvious addition of constant domains to enhance stability). Some of the antibodies expressed in the DVD-Ig/TBTI format show a position effect on the second (or innermost) antigen binding position (Fv2). Depending on the sequence and the nature of the antigen recognized by the Fv2 position, this antibody domain displays a reduced affinity to its antigen (*i.e.*, loss of on-rate in comparison to the parental antibody). One possible explanation for this observation is that the linker between V_{L1} and V_{L2} protrudes into the CDR region of Fv2, making the Fv2 somewhat inaccessible for larger antigens.

[0020] The second configuration of a bispecific antibody fragment described in U.S. Patent No. 5,989,830 is the cross-over double head (CODH), having the following orientation of variable domains expressed on two chains:


V_{L1}-linker-V_{L2}, for the light chain, and
V_{H2}-linker-V_{H1}, for the heavy chain

The '830 patent discloses that a bispecific cross-over double-head antibody fragment (construct GOSA.E) retains higher binding activity than a Dual-Fv (see page 20, lines 20-50 of the '830 patent), and further discloses that this format is less impacted by the linkers that are used between the variable domains (see page 20-21 of the '830 patent). EP 2 050 764 and WO 2010/014659 disclose bispecific antibody formats.


SUMMARY OF THE INVENTION

[0021] The scope of the present invention is defined in the appended claims.

[0022] The invention provides an antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites, wherein two polypeptide chains have a structure represented by the formula:




and two polypeptide chains have a structure represented by the formula:




wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains;
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers as defined in the appended claims;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.
[0023] The invention also provides an antibody-like binding protein comprising two polypeptide chains that form two antigen binding sites, wherein a first polypeptide chain has a structure represented by the formula:



and a second polypeptide chain has a structure represented by the formula:





wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers as defined in the appended claims;

and wherein the first and second polypeptides form a cross-over light chain-heavy chain pair.

[0024] Also disclosed is a method of making an antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites, comprising identifying a first antibody variable domain that binds a first target antigen and a second antibody variable domain that binds a second target antigen, each containing a V_L , and a V_H ; assigning either the light chain or the heavy chain as template chain; assigning the V_L of the first antibody variable domain or the second antibody variable domain as V_{L1} ; assigning a V_{L2} , a V_{H1} , and a V_{H2} according to formulas [I] and [II] below:




determining maximum and minimum lengths for L_1 , L_2 , L_3 , and L_4 ; generating the polypeptide structures of formulas I and II; selecting polypeptide structures of formulas I and II that bind the first target antigen and the second target antigen when combined to form the antibody-like binding protein;
 wherein:

V_{L1} is a first immunoglobulin light chain variable domain;

V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0025] Also disclosed is a method of making an antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites, comprising identifying a first antibody variable domain that binds a first target antigen and a second antibody variable domain that binds a second target antigen, each containing a V_L , and a V_H ; assigning either the light chain or the heavy chain as template chain; assigning the V_L of the first antibody variable domain or the second antibody variable domain as V_{L1} ; assigning a V_{L2} , a V_{H1} , and a V_{H2} according to formulas [I] and [II] below:



determining maximum and minimum lengths for L_1 , L_2 , L_3 , and L_4 ; generating polypeptide structures of formulas I and II; selecting polypeptide structures of formulas I and II that bind the first target antigen and the second target antigen when combined to form the antibody-like binding protein; wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0026] Specific embodiments of the invention will become evident from the following more detailed description of certain embodiments and the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0027]

Figure 1. Schematic representation of the antigen binding domains Fv1 and Fv2 within the dual V region configuration and arrangement of their respective peptide linkers L_L and L_H in the TBTI format.

Figure 2. Schematic diagram (2D) of the antigen binding domains Fv1 (anti-IL4) and Fv2 (anti-IL13) within the cross-over dual variable (CODV) configuration and the arrangement of their respective peptide linkers.

Figure 3. Schematic representation of the Fv anti-IL4 and Fab anti-IL13 showing one possible spatial arrangement obtained by protein-protein docking of Fv of anti-IL4 and the Fv of anti-IL13.

Figure 4. Assessment of tetravalent and bispecific binding ability of the CODV protein in a BIACORE assay by injecting the two antigens sequentially or simultaneously over a DVD-Ig protein-coated chip. The maximal signal observed by sequential injection can be obtained by co-injection of both antigens, demonstrating saturation of all binding sites.

Figure 5. Schematic diagram (2D) of the antigen binding domains within the CODV configuration and arrangement

of their respective peptide linker L_L (L_1 and L_2) and L_H (L_3 and L_4). In panel A, the light chain is kept in a "linear or template" alignment, whereas the heavy chain is in the "cross-over" configuration. In panel B, the heavy chain is kept in a "linear or template" alignment and the light chain is in the "cross-over" configuration.

Figure 6. Schematic representation of CODV-Ig design based on whether the light chain or heavy chain is used as "template."

Figure 7. Comparison of TBTI/DVD-Ig or CODV-Ig molecules incorporating anti-IL4 and anti-IL13 sequences.

Figure 8. Comparison of CODV-Fab and B-Fab formats in a cytotoxic assay using NALM-6 cells.

DETAILED DESCRIPTION OF THE INVENTION

[0028] The invention provides antibody-like binding proteins as defined in the appended claims, comprising four polypeptide chains that form four antigen binding sites, wherein each pair of polypeptides forming an antibody-like binding protein possesses dual variable domains having a cross-over orientation. Also disclosed are methods for making such antigen-like binding proteins.

[0029] Computer modeling predicted that the cross-over double-head (CODH) design of U.S. Patent No. 5,989,830 would yield a complex in which both binding sites face in the opposite direction, without the restraints suggested for the Dual-Fv configuration of U.S. Patent No. 7,612,181. In particular, computer modeling indicated that the length of the amino acid linkers between the variable domains was not critical for the CODH design, but was important for permitting full access to both antigen binding sites in the Dual-Fv design. As with the DVD-Ig/TBTI format, antibody-like binding protein constructs were prepared in which constant domains were attached to the CODH configuration to form antibody-like binding proteins comprising four polypeptide chains that form four antigen binding sites, wherein each pair of polypeptides forming an antibody-like binding protein possesses dual variable domains having a cross-over orientation (*i.e.*, CODH-Ig). CODH-Ig molecules are expected to possess significantly improved stability as compared with CODH molecules (as DVD-Ig/TBTI possessed improved stability over Dual-Fv molecules).

[0030] In order to test the above hypothesis, a CODH-Ig molecule was prepared using the anti-IL4 and anti-IL13 antibody sequences described in U.S. Patent Application Publication No. US 2010/0226923 A1. The CODH-Ig molecule differed from the CODH molecule of US 2010/0226923 with respect to the lengths of amino acid linkers separating the variable domains on the respective polypeptide chains. The CODH-Ig molecules were expressed in cells following transient transfection and were then purified by Protein A chromatography. Although their size-exclusion chromatography (SEC) profiles showed aggregation levels of 5-10%, none of the CODH-Ig molecules were functional, and thus none of the CODH-Ig molecules was able to bind all of its target antigens. The lack of antigen binding activity may have been due to a perturbed dimerization of the Fv-regions of the heavy and light chains due to unsuitable linker lengths compromising correct paratope formation. As a result, a protocol was developed to identify suitable amino acid linkers for insertion between the two variable domains and the second variable domain and constant domain on both the heavy and light polypeptide chains of an antibody-like binding protein. This protocol was based on protein-protein docking of homology and experimental models of the FvIL4 and FvIL13 regions, respectively, inclusion of the Fc1 domain the model, and construction of appropriate linkers between the FvIL4 and FvIL13 regions and between the Fv and constant Fc1 regions.

[0031] Standard recombinant DNA methodologies are used to construct the polynucleotides that encode the polypeptides which form the antibody-like binding proteins of the invention, incorporate these polynucleotides into recombinant expression vectors, and introduce such vectors into host cells. See *e.g.*, Sambrook et al., 2001, MOLECULAR CLONING: A LABORATORY MANUAL (Cold Spring Harbor Laboratory Press, 3rd ed.). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications, as commonly accomplished in the art, or as described herein. Unless specific definitions are provided, the nomenclature utilized in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Similarly, conventional techniques may be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, delivery, and treatment of patients.

1. General Definitions

[0032] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings. Unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular.

[0033] The term "polynucleotide" as used herein refers to single-stranded or double-stranded nucleic acid polymers

of at least 10 nucleotides in length. In certain embodiments, the nucleotides comprising the polynucleotide can be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. Such modifications include base modifications such as bromuridine, ribose modifications such as arabinoside and 2',3'-dideoxyribose, and internucleotide linkage modifications such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate and phosphoroamidate. The term "polynucleotide" specifically includes single-stranded and double-stranded forms of DNA.

[0034] An "isolated polynucleotide" is a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, which by virtue of its origin the isolated polynucleotide: (1) is not associated with all or a portion of a polynucleotide in which the isolated polynucleotide is found in nature, (2) is linked to a polynucleotide to which it is not linked in nature, or (3) does not occur in nature as part of a larger sequence.

[0035] An "isolated polypeptide" is one that: (1) is free of at least some other polypeptides with which it would normally be found, (2) is essentially free of other polypeptides from the same source, *e.g.*, from the same species, (3) is expressed by a cell from a different species, (4) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is associated in nature, (5) is not associated (by covalent or noncovalent interaction) with portions of a polypeptide with which the "isolated polypeptide" is associated in nature, (6) is operably associated (by covalent or noncovalent interaction) with a polypeptide with which it is not associated in nature, or (7) does not occur in nature. Such an isolated polypeptide can be encoded by genomic DNA, cDNA, mRNA or other RNA, of synthetic origin, or any combination thereof. Preferably, the isolated polypeptide is substantially free from polypeptides or other contaminants that are found in its natural environment that would interfere with its use (therapeutic, diagnostic, prophylactic, research or otherwise).

[0036] The term "human antibody" as used herein includes antibodies having variable and constant regions substantially corresponding to human germline immunoglobulin sequences. In some embodiments, human antibodies are produced in non-human mammals, including, but not limited to, rodents, such as mice and rats, and lagomorphs, such as rabbits. In other embodiments, human antibodies are produced in hybridoma cells. In still other embodiments, human antibodies are produced recombinantly.

[0037] Naturally occurring antibodies typically comprise a tetramer. Each such tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one full-length "light" chain (typically having a molecular weight of about 25 kDa) and one full-length "heavy" chain (typically having a molecular weight of about 50-70 kDa). The terms "heavy chain" and "light chain" as used herein refer to any immunoglobulin polypeptide having sufficient variable domain sequence to confer specificity for a target antigen. The amino-terminal portion of each light and heavy chain typically includes a variable domain of about 100 to 110 or more amino acids that typically is responsible for antigen recognition. The carboxy-terminal portion of each chain typically defines a constant domain responsible for effector function. Thus, in a naturally occurring antibody, a full-length heavy chain immunoglobulin polypeptide includes a variable domain (V_H) and three constant domains (C_{H1} , C_{H2} , and C_{H3}), wherein the V_H domain is at the amino-terminus of the polypeptide and the C_{H3} domain is at the carboxyl-terminus, and a full-length light chain immunoglobulin polypeptide includes a variable domain (V_L) and a constant domain (C_L), wherein the V_L domain is at the amino-terminus of the polypeptide and the C_L domain is at the carboxyl-terminus.

[0038] Human light chains are typically classified as kappa and lambda light chains, and human heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to, IgG1, IgG2, IgG3, and IgG4. IgM has subclasses including, but not limited to, IgM1 and IgM2. IgA is similarly subdivided into subclasses including, but not limited to, IgA1 and IgA2. Within full-length light and heavy chains, the variable and constant domains typically are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See, *e.g.*, FUNDAMENTAL IMMUNOLOGY (Paul, W., ed., Raven Press, 2nd ed., 1989). The variable regions of each light/heavy chain pair typically form an antigen binding site. The variable domains of naturally occurring antibodies typically exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair typically are aligned by the framework regions, which may enable binding to a specific epitope. From the amino-terminus to the carboxyl-terminus, both light and heavy chain variable domains typically comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4.


[0039] The term "native Fc" as used herein refers to a molecule comprising the sequence of a non-antigen-binding fragment resulting from digestion of an antibody or produced by other means, whether in monomeric or multimeric form, and can contain the hinge region. The original immunoglobulin source of the native Fc is preferably of human origin and can be any of the immunoglobulins, although IgG 1 and IgG2 are preferred. Native Fc molecules are made up of monomeric polypeptides that can be linked into dimeric or multimeric forms by covalent (*i.e.*, disulfide bonds) and non-covalent association. The number of intermolecular disulfide bonds between monomeric subunits of native Fc molecules ranges from 1 to 4 depending on class (*e.g.*, IgG, IgA, and IgE) or subclass (*e.g.*, IgG1, IgG2, IgG3, IgA1, and IgA2). One example of a native Fc is a disulfide-bonded dimer resulting from papain digestion of an IgG. The term "native Fc"

as used herein is generic to the monomeric, dimeric, and multimeric forms.


[0040] The term "Fc variant" as used herein refers to a molecule or sequence that is modified from a native Fc but still comprises a binding site for the salvage receptor, FcRn (neonatal Fc receptor). Exemplary Fc variants, and their interaction with the salvage receptor, are known in the art. Thus, the term "Fc variant" can comprise a molecule or sequence that is humanized from a non-human native Fc. Furthermore, a native Fc comprises regions that can be removed because they provide structural features or biological activity that are not required for the antibody-like binding proteins of the invention. Thus, the term "Fc variant" comprises a molecule or sequence that lacks one or more native Fc sites or residues, or in which one or more Fc sites or residues has been modified, that affect or are involved in: (1) disulfide bond formation, (2) incompatibility with a selected host cell, (3) N-terminal heterogeneity upon expression in a selected host cell, (4) glycosylation, (5) interaction with complement, (6) binding to an Fc receptor other than a salvage receptor, or (7) antibody-dependent cellular cytotoxicity (ADCC).

[0041] The term "Fc domain" as used herein encompasses native Fc and Fc variants and sequences as defined above. As with Fc variants and native Fc molecules, the term "Fc domain" includes molecules in monomeric or multimeric form, whether digested from whole antibody or produced by other means.

[0042] The term "antibody-like binding protein" as used herein refers to a non-naturally occurring (or recombinant) molecule that specifically binds to at least one target antigen, and which comprises four polypeptide chains that form four antigen binding sites, wherein two polypeptide chains have a structure represented by the formula:



and two polypeptide chains have a structure represented by the formula:



wherein:

V_{L1} is a first immunoglobulin light chain variable domain;

V_{L2} is a second immunoglobulin light chain variable domain;

V_{H1} is a first immunoglobulin heavy chain variable domain;

V_{H2} is a second immunoglobulin heavy chain variable domain;


C_L is an immunoglobulin light chain constant domain;

C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;


Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains;

L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair. The term "antibody-like binding protein" as used herein also refers to a non-naturally occurring (or recombinant) molecule that specifically binds to at least one target antigen, and which comprises two polypeptide chains that form two antigen binding sites, wherein a first polypeptide chain has a structure represented by the formula:



and a second polypeptide chain has a structure represented by the formula:



wherein:

V_{L1} is a first immunoglobulin light chain variable domain;

V_{L2} is a second immunoglobulin light chain variable domain;

V_{H1} is a first immunoglobulin heavy chain variable domain;

V_{H2} is a second immunoglobulin heavy chain variable domain;

C_L is an immunoglobulin light chain constant domain;

C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
L₁, L₂, L₃, and L₄ are amino acid linkers;

and wherein the first and second polypeptides form a cross-over light chain-heavy chain pair. A "recombinant" molecule is one that has been prepared, expressed, created, or isolated by recombinant means.

[0043] One embodiment of the invention provides antibody-like binding proteins as defined in the appended claims having biological and immunological specificity to between one and four target antigens. Another embodiment of the invention provides nucleic acid molecules comprising nucleotide sequences encoding polypeptide chains that form such antibody-like binding proteins. Another embodiment of the invention provides expression vectors comprising nucleic acid molecules comprising nucleotide sequences encoding polypeptide chains that form such antibody-like binding proteins. Yet another embodiment of the invention provides host cells that express such antibody-like binding proteins (*i.e.*, comprising nucleic acid molecules or vectors encoding polypeptide chains that form such antibody-like binding proteins).

[0044] The term "swapability" as used herein refers to the interchangeability of variable domains within the CODV format and with retention of folding and ultimate binding affinity. "Full swapability" refers to the ability to swap the order of both V_{H1} and V_{H2} domains, and therefore the order of V_{L1} and V_{L2} domains, in a CODV-Ig (*i.e.*, to reverse the order) or CODV-Fab while maintaining full functionality of the antibody-like binding protein as evidenced by the retention of binding affinity. Furthermore, it should be noted that the designations V_H and V_L within a particular CODV-Ig or CODV-Fab refer only to the domain's location on a particular protein chain in the final format. For example, V_{H1} and V_{H2} could be derived from V_{L1} and V_{L2} domains in parent antibodies and placed into the V_{H1} and V_{H2} positions in the antibody-like binding protein. Likewise, V_{L1} and V_{L2} could be derived from V_{H1} and V_{H2} domains in parent antibodies and placed in the V_{H1} and V_{H2} positions in the antibody-like binding protein. Thus, the V_H and V_L designations refer to the present location and not the original location in a parent antibody. V_H and V_L domains are therefore "swappable."

[0045] An "isolated" antibody-like binding protein is one that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the antibody-like binding protein, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the antibody-like binding protein will be purified: (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody-like binding proteins include the antibody-like binding protein *in situ* within recombinant cells since at least one component of the antibody-like binding protein's natural environment will not be present.

[0046] The terms "substantially pure" or "substantially purified" as used herein refer to a compound or species that is the predominant species present (*i.e.*, on a molar basis it is more abundant than any other individual species in the composition). In some embodiments, a substantially purified fraction is a composition wherein the species comprises at least about 50% (on a molar basis) of all macromolecular species present. In other embodiments, a substantially pure composition will comprise more than about 80%, 85%, 90%, 95%, or 99% of all macromolecular species present in the composition. In still other embodiments, the species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

[0047] The term "antigen" or "target antigen" as used herein refers to a molecule or a portion of a molecule that is capable of being bound by an antibody-like binding protein, and additionally is capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. A target antigen may have one or more epitopes. With respect to each target antigen recognized by an antibody-like binding protein, the antibody-like binding protein is capable of competing with an intact antibody that recognizes the target antigen. A "bivalent" antibody-like binding protein, other than a "multispecific" or "multifunctional" antibody-like binding protein, is understood to comprise antigen binding sites having identical antigenic specificity.

[0048] A bispecific or bifunctional antibody typically is an artificial hybrid antibody having two different heavy chain/light chain pairs and two different binding sites or epitopes. Bispecific antibodies may be produced by a variety of methods including, but not limited to, fusion of hybridomas or linking of F(ab') fragments.

[0049] A F(ab) fragment typically includes one light chain and the V_H and C_{H1} domains of one heavy chain, wherein the V_H-C_{H1} heavy chain portion of the F(ab) fragment cannot form a disulfide bond with another heavy chain polypeptide. As used herein, a F(ab) fragment can also include one light chain containing two variable domains separated by an amino acid linker and one heavy chain containing two variable domains separated by an amino acid linker and a C_{H1} domain.

[0050] A F(ab') fragment typically includes one light chain and a portion of one heavy chain that contains more of the constant region (between the C_{H1} and C_{H2} domains), such that an interchain disulfide bond can be formed between two

heavy chains to form a $F(ab')_2$ molecule.

[0051] The phrases "biological property," "biological characteristic," and the term "activity" in reference to an antibody-like binding protein of the invention are used interchangeably herein and include, but are not limited to, epitope affinity and specificity, ability to antagonize the activity of the antigen target (or targeted polypeptide), the *in vivo* stability of the antibody-like binding protein, and the immunogenic properties of the antibody-like binding protein. Other identifiable biological properties or characteristics of an antibody-like binding protein include, for example, cross-reactivity, (*i.e.*, with non-human homologs of the antigen target, or with other antigen targets or tissues, generally), and ability to preserve high expression levels of protein in mammalian cells. The aforementioned properties or characteristics can be observed or measured using art-recognized techniques including, but not limited to ELISA, competitive ELISA, surface plasmon resonance analysis, *in vitro* and *in vivo* neutralization assays, and immunohistochemistry with tissue sections from different sources including human, primate, or any other source as the need may be.

[0052] The term "immunologically functional immunoglobulin fragment" as used herein refers to a polypeptide fragment that contains at least the CDRs of the immunoglobulin heavy or light chains from which the polypeptide fragment was derived. An immunologically functional immunoglobulin fragment is capable of binding to a target antigen.

[0053] A "neutralizing" antibody-like binding protein as used herein refers to a molecule that is able to block or substantially reduce an effector function of a target antigen to which it binds. As used herein, "substantially reduce" means at least about 60%, preferably at least about 70%, more preferably at least about 75%, even more preferably at least about 80%, still more preferably at least about 85%, most preferably at least about 90% reduction of an effector function of the target antigen.

[0054] The term "epitope" includes any determinant, preferably a polypeptide determinant, capable of specifically binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl groups, or sulfonyl groups, and, in certain embodiments, may have specific three-dimensional structural characteristics and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody or antibody-like binding protein. In certain embodiments, an antibody-like binding protein is said to specifically bind an antigen when it preferentially recognizes its target antigen in a complex mixture of proteins and/or macromolecules. In preferred embodiments, an antibody-like binding protein is said to specifically bind an antigen when the equilibrium dissociation constant is $\leq 10^{-8}$ M, more preferably when the equilibrium dissociation constant is $\leq 10^{-9}$ M, and most preferably when the dissociation constant is $\leq 10^{-10}$ M.

[0055] The dissociation constant (K_D) of an antibody-like binding protein can be determined, for example, by surface plasmon resonance. Generally, surface plasmon resonance analysis measures real-time binding interactions between ligand (a target antigen on a biosensor matrix) and analyte (an antibody-like binding protein in solution) by surface plasmon resonance (SPR) using the BIAcore system (Pharmacia Biosensor; Piscataway, NJ). Surface plasmon analysis can also be performed by immobilizing the analyte (antibody-like binding protein on a biosensor matrix) and presenting the ligand (target antigen). The term " K_D ," as used herein refers to the dissociation constant of the interaction between a particular antibody-like binding protein and a target antigen.

[0056] The term "specifically binds" as used herein refers to the ability of an antibody-like protein or an antigen-binding fragment thereof to bind to an antigen containing an epitope with a K_D of at least about 1×10^{-6} M, 1×10^{-7} M, 1×10^{-8} M, 1×10^{-9} M, 1×10^{-10} M, 1×10^{-11} M, 1×10^{-12} M, or more, and/or to bind to an epitope with an affinity that is at least two-fold greater than its affinity for a nonspecific antigen.

[0057] The term "linker" as used herein refers to one or more amino acid residues inserted between immunoglobulin domains to provide sufficient mobility for the domains of the light and heavy chains to fold into cross over dual variable region immunoglobulins. A linker is inserted at the transition between variable domains or between variable and constant domains, respectively, at the sequence level. The transition between domains can be identified because the approximate size of the immunoglobulin domains are well understood. The precise location of a domain transition can be determined by locating peptide stretches that do not form secondary structural elements such as beta-sheets or alpha-helices as demonstrated by experimental data or as can be assumed by techniques of modeling or secondary structure prediction. The linkers described herein are referred to as L_1 , which is located on the light chain between the N-terminal V_{L1} and V_{L2} domains; L_2 , which is also on the light chain is located between the V_{L2} and C-terminal C_L domains. The heavy chain linkers are known as L_3 , which is located between the N-terminal V_{H2} and V_{H1} domains; and L_4 , which is located between the V_{H1} and C_{H1} -Fc domains. The linkers L_1 , L_2 , L_3 , and L_4 are independent, but they may in some cases have the same sequence and/or length.

[0058] The term "vector" as used herein refers to any molecule (*e.g.*, nucleic acid, plasmid, or virus) that is used to transfer coding information to a host cell. The term "vector" includes a nucleic acid molecule that is capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double-stranded DNA molecule into which additional DNA segments may be inserted. Another type of vector is a viral vector, wherein additional DNA segments may be inserted into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and

episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell and thereby are replicated along with the host genome. In addition, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. The terms "plasmid" and "vector" may be used interchangeably herein, as a plasmid is the most commonly used form of vector. However, the invention is intended to include other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses, and adeno-associated viruses), which serve equivalent functions.

[0059] The term "operably linked" is used herein to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably linked to a coding sequence may be capable of effecting the replication, transcription, and/or translation of the coding sequence. For example, a coding sequence is operably linked to a promoter when the promoter is capable of directing transcription of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

[0060] The phrase "recombinant host cell" (or "host cell") as used herein refers to a cell into which a recombinant expression vector has been introduced. A recombinant host cell or host cell is intended to refer not only to the particular subject cell, but also to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but such cells are still included within the scope of the term "host cell" as used herein. A wide variety of host cell expression systems can be used to express the antibody-like binding proteins of the invention, including bacterial, yeast, baculoviral, and mammalian expression systems (as well as phage display expression systems). An example of a suitable bacterial expression vector is pUC19. To express an antibody-like binding protein recombinantly, a host cell is transformed or transfected with one or more recombinant expression vectors carrying DNA fragments encoding the polypeptide chains of the antibody-like binding protein such that the polypeptide chains are expressed in the host cell and, preferably, secreted into the medium in which the host cells are cultured, from which medium the antibody-like binding protein can be recovered.

[0061] The term "transformation" as used herein refers to a change in a cell's genetic characteristics, and a cell has been transformed when it has been modified to contain a new DNA. For example, a cell is transformed where it is genetically modified from its native state. Following transformation, the transforming DNA may recombine with that of the cell by physically integrating into a chromosome of the cell, or may be maintained transiently as an episomal element without being replicated, or may replicate independently as a plasmid. A cell is considered to have been stably transformed when the DNA is replicated with the division of the cell. The term "transfection" as used herein refers to the uptake of foreign or exogenous DNA by a cell, and a cell has been "transfected" when the exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are well known in the art. Such techniques can be used to introduce one or more exogenous DNA molecules into suitable host cells.

[0062] The term "naturally occurring" as used herein and applied to an object refers to the fact that the object can be found in nature and has not been manipulated by man. For example, a polynucleotide or polypeptide that is present in an organism (including viruses) that can be isolated from a source in nature and that has not been intentionally modified by man is naturally-occurring. Similarly, "non-naturally occurring" as used herein refers to an object that is not found in nature or that has been structurally modified or synthesized by man.

[0063] As used herein, the twenty conventional amino acids and their abbreviations follow conventional usage. Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids; unnatural amino acids such as α -, α -disubstituted amino acids, N-alkyl amino acids, lactic acid, and other unconventional amino acids may also be suitable components for the polypeptide chains of the antibody-like binding proteins of the invention. Examples of unconventional amino acids include: 4-hydroxyproline, γ -carboxyglutamate, ϵ -N,N,N-trimethyllysine, ϵ -N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, σ -N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxyl-terminal direction, in accordance with standard usage and convention.

[0064] Naturally occurring residues may be divided into classes based on common side chain properties:

- (1) hydrophobic: Met, Ala, Val, Leu, Ile, Phe, Trp, Tyr, Pro;
- (2) polar hydrophilic: Arg, Asn, Asp, Gln, Glu, His, Lys, Ser, Thr ;
- (3) aliphatic: Ala, Gly, Ile, Leu, Val, Pro;
- (4) aliphatic hydrophobic: Ala, Ile, Leu, Val, Pro;
- (5) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

- (6) acidic: Asp, Glu;
 (7) basic: His, Lys, Arg;
 (8) residues that influence chain orientation: Gly, Pro;
 (9) aromatic: His, Trp, Tyr, Phe; and
 (10) aromatic hydrophobic: Phe, Trp, Tyr.

[0065] Conservative amino acid substitutions may involve exchange of a member of one of these classes with another member of the same class. Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and other reversed or inverted forms of amino acid residues. Non-conservative substitutions may involve the exchange of a member of one of these classes for a member from another class.

[0066] A skilled artisan will be able to determine suitable variants of the polypeptide chains of the antibody-like binding proteins of the invention using well-known techniques. For example, one skilled in the art may identify suitable areas of a polypeptide chain that may be changed without destroying activity by targeting regions not believed to be important for activity. Alternatively, one skilled in the art can identify residues and portions of the molecules that are conserved among similar polypeptides. In addition, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

[0067] The term "patient" as used herein includes human and animal subjects.

[0068] A "disorder" is any condition that would benefit from treatment using the antibody-like binding proteins of the invention. "Disorder" and "condition" are used interchangeably herein and include chronic and acute disorders or diseases, including those pathological conditions that predispose a patient to the disorder in question.

[0069] The terms "treatment" or "treat" as used herein refer to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those having the disorder as well as those prone to have the disorder or those in which the disorder is to be prevented.

[0070] The terms "pharmaceutical composition" or "therapeutic composition" as used herein refer to a compound or composition capable of inducing a desired therapeutic effect when properly administered to a patient.


[0071] The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to one or more formulation materials suitable for accomplishing or enhancing the delivery of an antibody-like binding protein.

[0072] The terms "effective amount" and "therapeutically effective amount" when used in reference to a pharmaceutical composition comprising one or more antibody-like binding proteins refer to an amount or dosage sufficient to produce a desired therapeutic result. More specifically, a therapeutically effective amount is an amount of an antibody-like binding protein sufficient to inhibit, for some period of time, one or more of the clinically defined pathological processes associated with the condition being treated. The effective amount may vary depending on the specific antibody-like binding protein that is being used, and also depends on a variety of factors and conditions related to the patient being treated and the severity of the disorder. For example, if the antibody-like binding protein is to be administered *in vivo*, factors such as the age, weight, and health of the patient as well as dose response curves and toxicity data obtained in preclinical animal work would be among those factors considered. The determination of an effective amount or therapeutically effective amount of a given pharmaceutical composition is well within the ability of those skilled in the art.


[0073] One embodiment of the invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of an antibody-like binding protein as defined in the appended claims.

2. Antibody-like Binding Proteins

[0074] In one embodiment of the invention, the antibody-like binding proteins comprise four polypeptide chains that form four antigen binding sites, wherein two polypeptide chains have a structure represented by the formula:



and two polypeptide chains have a structure represented by the formula:




wherein:


V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains;
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers as defined in the appended claims;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0075] In another embodiment of the invention, the antibody-like binding proteins comprise two polypeptide chains that form two antigen binding sites, wherein a first polypeptide chain has a structure represented by the formula:



and a second polypeptide chain has a structure represented by the formula:



wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers as defined in the appended claims;

and wherein the first and second polypeptides form a cross-over light chain-heavy chain pair.

[0076] The antibody-like binding proteins of the invention may be prepared using domains or sequences obtained or derived from any human or non-human antibody, including, for example, human, murine, or humanized antibodies.

[0077] In some antibody-like binding proteins of the invention, the length of L_3 is at least twice the length of L_1 . In other antibody-like binding proteins of the invention, the length of L_4 is at least twice the length of L_2 . In some antibody-like binding proteins of the invention, the length of L_1 is at least twice the length of L_3 . In other antibody-like binding proteins of the invention, the length of L_2 is at least twice the length of L_4 .

[0078] In some antibody-like binding proteins of the invention, L_1 is 3 to 12 amino acid residues in length, L_2 is 3 to 14 amino acid residues in length, L_3 is 1 to 8 amino acid residues in length, and L_4 is 1 to 3 amino acid residues in length. In other antibody-like binding proteins, L_1 is 5 to 10 amino acid residues in length, L_2 is 5 to 8 amino acid residues in length, L_3 is 1 to 5 amino acid residues in length, and L_4 is 1 to 2 amino acid residues in length. In a preferred antibody-like binding protein, L_1 is 7 amino acid residues in length, L_2 is 5 amino acid residues in length, L_3 is 1 amino acid residues in length, and L_4 is 2 amino acid residues in length.

[0079] In some antibody-like binding proteins of the invention, L_1 is 1 to 3 amino acid residues in length, L_2 is 1 to 4 amino acid residues in length, L_3 is 2 to 15 amino acid residues in length, and L_4 is 2 to 15 amino acid residues in length. In other antibody-like binding proteins, L_1 is 1 to 2 amino acid residues in length, L_2 is 1 to 2 amino acid residues in length, L_3 is 4 to 12 amino acid residues in length, and L_4 is 2 to 12 amino acid residues in length. In a preferred antibody-like binding protein, L_1 is 1 amino acid residue in length, L_2 is 2 amino acid residues in length, L_3 is 7 amino acid residues in length, and L_4 is 5 amino acid residues in length.

[0080] Also disclosed herein are antibody-like binding proteins, in which L_1 , L_3 , or L_4 may be equal to zero. However, in antibody-like binding proteins wherein L_1 , L_3 , or L_4 is equal to zero, the corresponding transition linker between the variable region and constant region or between the dual variable domains on the other chain cannot be zero. In some embodiments, L_1 is equal to zero and L_3 is 2 or more amino acid residues, L_3 is equal to zero and L_1 is equal to 1 or more amino acid residues, or L_4 is equal to 0 and L_2 is 3 or more amino acid residues.

[0081] In some antibody-like binding proteins of the invention, at least one of the linkers selected from the group consisting of L_1 , L_2 , L_3 , and L_4 contains at least one cysteine residue.

[0082] Examples of suitable linkers include a single glycine (Gly) residue; a diglycine peptide (Gly-Gly); a tripeptide (Gly-Gly-Gly); a peptide with four glycine residues (Gly-Gly-Gly-Gly; SEQ ID NO: 25); a peptide with five glycine residues (Gly-Gly-Gly-Gly-Gly; SEQ ID NO: 26); a peptide with six glycine residues (Gly-Gly-Gly-Gly-Gly-Gly; SEQ ID NO: 27); a peptide with seven glycine residues (Gly-Gly-Gly-Gly-Gly-Gly-Gly; SEQ ID NO: 28); a peptide with eight glycine residues (Gly-Gly-Gly-Gly-Gly-Gly-Gly-Gly; SEQ ID NO: 29). Other combinations of amino acid residues may be used such as the peptide Gly-Gly-Gly-Gly-Ser (SEQ ID NO: 30) and the peptide Gly-Gly-Gly-Gly-Ser-Gly-Gly-Gly-Gly-Ser (SEQ ID NO: 31). Other suitable linkers include a single Ser, and Val residue; the dipeptide Arg-Thr, Gln-Pro, Ser-Ser, Thr-Lys, and Ser-Leu; Thr-Lys-Gly-Pro-Ser (SEQ ID NO: 52), Thr-Val-Ala-Ala-Pro (SEQ ID NO: 53), Gln-Pro-Lys-Ala-Ala (SEQ ID NO: 54), Gln-Arg-Ile-Glu-Gly (SEQ ID NO: 55); Ala-Ser-Thr-Lys-Gly-Pro-Ser (SEQ ID NO: 48), Arg-Thr-Val-Ala-Ala-Pro-Ser (SEQ ID NO: 49), Gly-Gln-Pro-Lys-Ala-Ala-Pro (SEQ ID NO: 50), and His-Ile-Asp-Ser-Pro-Asn-Lys (SEQ ID NO: 51). The examples listed above are not intended to limit the scope of the invention in any way, and linkers comprising randomly selected amino acids selected from the group consisting of valine, leucine, isoleucine, serine, threonine, lysine, arginine, histidine, aspartate, glutamate, asparagine, glutamine, glycine, and proline have been shown to be suitable in the antibody-like binding proteins of the invention (see Example 12).

[0083] The identity and sequence of amino acid residues in the linker may vary depending on the type of secondary structural element necessary to achieve in the linker. For example, glycine, serine, and alanine are best for linkers having maximum flexibility. Some combination of glycine, proline, threonine, and serine are useful if a more rigid and extended linker is necessary. Any amino acid residue may be considered as a linker in combination with other amino acid residues to construct larger peptide linkers as necessary depending on the desired properties.


[0084] In some antibody-like binding proteins of the invention, V_{L1} comprises the amino acid sequence of SEQ ID NO: 1; V_{L2} comprises the amino acid sequence of SEQ ID NO: 3; V_{H1} comprises the amino acid sequence of SEQ ID NO: 2; and V_{H2} comprises the amino acid sequence of SEQ ID NO: 4.

[0085] In some embodiments of the invention, the antibody-like binding protein is capable of specifically binding one or more antigen targets. In preferred embodiments of the invention, the antibody-like binding protein is capable of specifically binding at least one antigen target selected from the group consisting of B7.1, B7.2, BAFF, BlyS, C3, C5, CCL11 (eotaxin), CCL15 (MIP-1d), CCL17 (TARC), CCL19 (MIP-3b), CCL2 (MCP-1), CCL20 (MIP-3a), CCL21 (MIP-2), SLC, CCL24 (MIP-2/eotaxin-2), CCL25 (TECK), CCL26 (eotaxin-3), CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (MCP-2), CD3, CD19, CD20, CD24, CD40, CD40L, CD80, CD86, CDH1 (E-cadherin), Chitinase, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CX3CL1 (SCYD1), CXCL12 (SDF1), CXCL13, EGFR, FCER1A, FCER2, HER2, IGF1R, IL-1, IL-12, IL13, IL4, IL15, IL17, IL18, IL1A, IL1B, IL1F10, IL1 β , IL2, IL4, IL6, IL7, IL8, IL9, IL12/23, IL22, IL23, IL25, IL27, IL35, ITGB4 (b 4 integrin), LEP (leptin), MHC class II, TLR2, TLR4, TLR5, TNF, TNF- α , TNFSF4 (OX40 ligand), TNFSF5 (CD40 ligand), Toll-like receptors, TREM1, TSLP, TWEAK, XCR1 (GPR5/CCXCR1), DNNGR-1(CLEC91), and HMGB1. In other embodiments of the invention, the antibody-like binding protein is capable of inhibiting the function of one or more of the antigen targets.

[0086] In some embodiments of the invention, the antibody-like binding protein is bispecific and capable of binding two different antigen targets or epitopes. In a preferred embodiment of the invention, the antibody-like binding protein is bispecific and each light chain-heavy chain pair is capable of binding two different antigen targets or epitopes. In a more preferred embodiment, the antibody-like binding protein is capable of binding two different antigen targets that are selected from the group consisting of IL4 and IL13, IGF1R and HER2, IGF1R and EGFR, EGFR and HER2, BK and IL13, PDL-1 and CTLA-4, CTLA4 and MHC class II, IL-12 and IL-18, IL-1 α and IL-1 β , TNF α and IL12/23, TNF α and IL-12p40, TNF α and IL-1 β , TNF α and IL-23, and IL17 and IL23. In an even more preferred embodiment, the antibody-like binding protein is capable of binding the antigen targets IL4 and IL13.

[0087] In some embodiments of the invention, the antibody-like binding protein specifically binds IL4 with an on-rate of 2.97×10^7 and an off-rate of 3.30×10^{-4} and specifically binds IL13 with an on-rate of 1.39×10^6 and an off-rate of 1.63×10^{-4} . In other embodiments of the invention, the antibody-like binding protein specifically binds IL4 with an on-rate of 3.16×10^7 and an off-rate of 2.89×10^{-4} and specifically binds IL13 with an on-rate of 1.20×10^6 and an off-rate of 1.12×10^{-4} .

[0088] Also disclosed is a method, in which an antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites is prepared by identifying a first antibody variable domain that binds a first target antigen and a second antibody variable domain that binds a second target antigen, each containing a V_L , and a V_H ; assigning either the light chain or the heavy chain as template chain; assigning the V_L of the first antibody variable domain or the second antibody variable domain as V_{L1} ; assigning a V_{L2} , a V_{H1} , and a V_{H2} according to formulas [I] and [II] below:





determining maximum and minimum lengths for L_1 , L_2 , L_3 , and L_4 ; generating polypeptide structures of formulas I and II; selecting polypeptide structures of formulas I and II that bind the first target antigen and the second target antigen when combined to form the antibody-like binding protein;
wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0089] Also disclosed is a method, in which an antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites is prepared by identifying a first antibody variable domain that binds a first target antigen and a second antibody variable domain that binds a second target antigen, each containing a V_L and a V_H ; assigning either the light chain or the heavy chain as template chain; assigning the V_L of the first antibody variable domain or the second antibody variable domain as V_{L1} ; assigning a V_{L2} , a V_{H1} , and a V_{H2} according to formulas [I] and [II] below:





determining maximum and minimum lengths for L_1 , L_2 , L_3 , and L_4 ; generating polypeptide structures of formulas I and II; selecting polypeptide structures of formulas I and II that bind the first target antigen and the second target antigen when combined to form the antibody-like binding protein;
wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain; and
 L_1 , L_2 , L_3 , and L_4 are an amino acid linkers;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0090] In other embodiments, an antibody-like binding protein in which the first antibody variable domain and the second antibody variable domain are the same is prepared.

[0091] Also disclosed is a method for making an antibody-like binding protein, comprising expressing in a cell one or more nucleic acid molecules encoding polypeptides having structures represented by the formulas [I] and [II] below:




wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;

V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;
 and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

[0092] Also disclosed is a method for making an antibody-like binding protein, comprising expressing in a cell one or more nucleic acid molecules encoding polypeptides having structures represented by the formulas [I] and [II] below:



wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

and wherein the polypeptide of formula I and the polypeptide of formula II form a cross-over light chain-heavy chain pair.

3. Uses for Antibody-like Binding Proteins

[0093] The antibody-like binding proteins of the invention can be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays for the detection and quantitation of one or more target antigens. The antibody-like binding proteins will bind the one or more target antigens with an affinity that is appropriate for the assay method being employed.

[0094] For diagnostic applications, in certain embodiments, antibody-like binding proteins can be labeled with a detectable moiety. The detectable moiety can be any one that is capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety can be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , ^{125}I , ^{99}Tc , ^{111}In , or ^{67}Ga ; a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase, β -galactosidase, or horseradish peroxidase.

[0095] The antibody-like binding proteins of the invention are also useful for *in vivo* imaging. An antibody-like binding protein labeled with a detectable moiety can be administered to an animal, preferably into the bloodstream, and the presence and location of the labeled antibody in the host assayed. The antibody-like binding protein can be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

[0096] Also disclosed is a kit comprising an antibody-like binding protein and other reagents useful for detecting target antigen levels in biological samples. Such reagents can include a detectable label, blocking serum, positive and negative control samples, and detection reagents.

4. Antibody-Like Binding Protein Therapeutic Compositions and Administration Thereof

[0097] Therapeutic or pharmaceutical compositions comprising antibody-like binding proteins as defined in the appended claims are within the scope of the invention. Such therapeutic or pharmaceutical compositions can comprise a therapeutically effective amount of an antibody-like binding protein, or antibody-like binding protein-drug conjugate, in admixture with a pharmaceutically or physiologically acceptable formulation agent selected for suitability with the mode of administration.

[0098] Acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations em-

ployed.

[0099] The pharmaceutical composition can contain formulation materials for modifying, maintaining, or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption, or penetration of the composition. Suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine, or lysine), antimicrobials, antioxidants (such as ascorbic acid, sodium sulfite, or sodium hydrogen-sulfite), buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates, or other organic acids), bulking agents (such as mannitol or glycine), chelating agents (such as ethylenediamine tetraacetic acid (EDTA)), complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin, or hydroxypropyl-beta-cyclodextrin), fillers, monosaccharides, disaccharides, and other carbohydrates (such as glucose, mannose, or dextrans), proteins (such as serum albumin, gelatin, or immunoglobulins), coloring, flavoring and diluting agents, emulsifying agents, hydrophilic polymers (such as polyvinylpyrrolidone), low molecular weight polypeptides, salt-forming counterions (such as sodium), preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid, or hydrogen peroxide), solvents (such as glycerin, propylene glycol, or polyethylene glycol), sugar alcohols (such as mannitol or sorbitol), suspending agents, surfactants or wetting agents (such as pluronics; PEG; sorbitan esters; polysorbates such as polysorbate 20 or polysorbate 80; triton; tromethamine; lecithin; cholesterol or tyloxapal), stability enhancing agents (such as sucrose or sorbitol), tonicity enhancing agents (such as alkali metal halides - preferably sodium or potassium chloride - or mannitol sorbitol), delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants (*see, e.g.*, REMINGTON'S PHARMACEUTICAL SCIENCES (18th Ed., A.R. Gennaro, ed., Mack Publishing Company 1990), and subsequent editions of the same).

[0100] The optimal pharmaceutical composition will be determined by a skilled artisan depending upon, for example, the intended route of administration, delivery format, and desired dosage. Such compositions can influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the antibody-like binding protein.

[0101] The primary vehicle or carrier in a pharmaceutical composition can be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier for injection can be water, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. Other exemplary pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which can further include sorbitol or a suitable substitute. Antibody-like binding protein compositions can be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents in the form of a lyophilized cake or an aqueous solution. Further, the antibody-like binding protein can be formulated as a lyophilizate using appropriate excipients such as sucrose.

[0102] The pharmaceutical compositions of the invention can be selected for parenteral delivery. Alternatively, the compositions can be selected for inhalation or for delivery through the digestive tract, such as orally. The preparation of such pharmaceutically acceptable compositions is within the skill of the art.

[0103] The formulation components are present in concentrations that are acceptable to the site of administration. For example, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

[0104] When parenteral administration is contemplated, the therapeutic compositions can be in the form of a pyrogen-free, parenterally acceptable, aqueous solution comprising the desired antibody-like binding protein in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which an antibody-like binding protein is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads, or liposomes, that provides for the controlled or sustained release of the product which can then be delivered via a depot injection. Hyaluronic acid can also be used, and this can have the effect of promoting sustained duration in the circulation. Other suitable means for the introduction of the desired molecule include implantable drug delivery devices.

[0105] In one embodiment, a pharmaceutical composition can be formulated for inhalation. For example, an antibody-like binding protein can be formulated as a dry powder for inhalation. Antibody-like binding protein inhalation solutions can also be formulated with a propellant for aerosol delivery. In yet another embodiment, solutions can be nebulized.

[0106] It is also contemplated that certain formulations can be administered orally. Antibody-like binding proteins that are administered in this fashion can be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule can be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be included to facilitate absorption of the antibody-like binding protein. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders can also be employed.

[0107] Another pharmaceutical composition can involve an effective quantity of antibody-like binding proteins in a mixture with non-toxic excipients that are suitable for the manufacture of tablets. By dissolving the tablets in sterile water,

or another appropriate vehicle, solutions can be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

[0108] Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving antibody-like binding proteins in sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. Additional examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices can include polyesters, hydrogels, polylactides, copolymers of L-glutamic acid and gamma ethyl-L-glutamate, poly(2-hydroxyethylmethacrylate), ethylene vinyl acetate, or poly-D(-)-3-hydroxybutyric acid. Sustained-release compositions can also include liposomes, which can be prepared by any of several methods known in the art.

[0109] Pharmaceutical compositions of the invention to be used for *in vivo* administration typically must be sterile. This can be accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using this method can be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration can be stored in lyophilized form or in a solution. In addition, parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0110] Once the pharmaceutical composition has been formulated, it can be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or as a dehydrated or lyophilized powder. Such formulations can be stored either in a ready-to-use form or in a form (e.g., lyophilized) requiring reconstitution prior to administration.

[0111] Also disclosed are kits for producing a single-dose administration unit. The kits can each contain both a first container having a dried protein and a second container having an aqueous formulation. Also disclosed are kits containing single and multi-chambered pre-filled syringes (e.g., liquid syringes and lysyringes).

[0112] The effective amount of an antibody-like binding protein pharmaceutical composition to be employed therapeutically will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will thus vary depending, in part, upon the molecule delivered, the indication for which the antibody-like binding protein is being used, the route of administration, and the size (body weight, body surface, or organ size) and condition (the age and general health) of the patient. Accordingly, the clinician can titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage can range from about 0.1 $\mu\text{g/kg}$ to up to about 100 mg/kg or more, depending on the factors mentioned above. In other embodiments, the dosage can range from 0.1 $\mu\text{g/kg}$ up to about 100 mg/kg ; or 1 $\mu\text{g/kg}$ up to about 100 mg/kg ; or 5 $\mu\text{g/kg}$, 10 $\mu\text{g/kg}$, 15 $\mu\text{g/kg}$, 20 $\mu\text{g/kg}$, 25 $\mu\text{g/kg}$, 30 $\mu\text{g/kg}$, 35 $\mu\text{g/kg}$, 40 $\mu\text{g/kg}$, 45 $\mu\text{g/kg}$, 50 $\mu\text{g/kg}$, 55 $\mu\text{g/kg}$, 60 $\mu\text{g/kg}$, 65 $\mu\text{g/kg}$, 70 $\mu\text{g/kg}$, 75 $\mu\text{g/kg}$, up to about 100 mg/kg .

[0113] Dosing frequency will depend upon the pharmacokinetic parameters of the antibody-like binding protein in the formulation being used. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition can therefore be administered as a single dose, as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages can be ascertained through use of appropriate dose-response data.

[0114] The route of administration of the pharmaceutical composition is in accord with known methods, e.g., orally; through injection by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, intraportal, or intralesional routes; by sustained release systems; or by implantation devices. Where desired, the compositions can be administered by bolus injection or continuously by infusion, or by implantation device.

[0115] The composition can also be administered locally via implantation of a membrane, sponge, or other appropriate material onto which the desired molecule has been absorbed or encapsulated. Where an implantation device is used, the device can be implanted into any suitable tissue or organ, and delivery of the desired molecule can be via diffusion, timed-release bolus, or continuous administration.

5. Examples

[0116] The Examples that follow are illustrative of specific embodiments of the invention, and various uses thereof. They are set forth for explanatory purposes only, and should not be construed as limiting the scope of the invention in any way.

Example 1. Design and Engineering of Bispecific Cross-over Dual Variable Region Antibody-like Binding Proteins

[0117] The cross-over dual variable region in an Fv format was described in U.S. Patent No. 5,989,830 and was referred to as a cross-over double head (CODH) configuration. Molecular modeling predicted that the cross-over double-head (CODH) design results in a complex with both binding sites facing in opposite directions, without the restraints suggested for the Dual-Fv configuration. The CODH Fv format was examined to determine whether it could be converted into complete antibody-like molecules by adding a C_L domain to the light chain and an Fc region to the heavy chain. A similar conversion was successful for the corresponding dual variable domains (DVD-Ig) and TBTI as described in U.S. Patent No. 7,612,181 and International Publication No. WO 2009/052081. The arrangement of the variable regions in the CODH format is shown in the structures below, which indicate the amino to carboxyl orientation of the peptide chains:

- (a) light chain: NH₂-V_{L1}-Linker-V_{L2}-COOH
 (b) heavy chain: NH₂-V_{H2}-Linker-V_{H1}-COOH

[0118] The amino to carboxyl terminal arrangement of the variable regions in (a) and (b) above can be distinguished from the arrangement in the Dual-Fv configuration shown in (c) and (d) below:

- (c) light chain: NH₂-V_{L1}-Linker-V_{L2}-COOH
 (d) heavy chain: NH₂-V_{H1}-Linker-V_{H2}-COOH

[0119] The main difference to note is the distinct placement of the corresponding light chain and heavy chain variable regions (V_{H1}/V_{L1} and V_{H2}/V_{L2}) with respect to each other in the two dual variable region configurations. The corresponding V_{L1} and V_{H1} domains were both at the N-terminus of the light and heavy chains in dual variable region configuration. In contrast, in the cross-over configuration, one half of one pair of an antibody variable region was separated spatially within the protein chain in the cross-over configuration. In the cross-over configuration, the V_{L1} domain would be at the N-terminus of the protein light chain but the pairing V_{H1} domain is at the C-terminus of the cross-over configuration heavy chain. The spatial relationship between V_{L1} and V_{H1} found in the dual variable region configuration is the arrangement found in natural antibodies.

[0120] One potential disadvantage of the dual Fv configuration is that the linker L_L separating the two variable regions protrudes into the antigen binding site of the Fv2 domain (see Figure 1). This protrusion may interfere with antigen binding and result in a perturbed accessibility of Antigen 2 to Fv2. This perturbed accessibility or interference may prevent antigen binding. In addition, this interference might be more pronounced where the size of Antigen 2 is larger. Indeed, it has been documented in U.S. Patent No. 7,612,181 that the binding affinity and neutralization ability of a DVD-Ig molecule depends on which antigen specificity is presented at the N-terminus or C-terminus. See U.S. Patent No. 7,612,181, Example 2.

[0121] Therefore, to create more stable antibody-like binding proteins that are not subject to loss of antigen affinity as compared to the parental antibody, cross-over dual variable region molecules having a C_L domain on the light chain and an Fc region on the heavy chain were designed and constructed. The polypeptides that form these antibody-like proteins have the structures shown below, in which the amino to carboxyl terminal orientation of the polypeptide chains is indicated:

- (e) light chain: NH₂-V_{L1}-Linker-V_{L2}-C_L-COOH
 (f) heavy chain: NH₂-V_{H2}-Linker-V_{H1}-C_{H1}-Fc-COOH

[0122] To evaluate whether this bispecific antibody-like protein design would bind to two different antigens, two previously generated and humanized variable regions from antibodies specific for IL4 (parental humanized anti-IL4) and IL13 (parental humanized anti-IL13) were used to construct the bispecific antibody-like molecules shown in Table 1. Sequencing of the mouse antibodies and the humanization process were described in International Publication No. WO 2009/052081 (TBTI). Briefly, amino acid sequences of the variable heavy and light chains of the murine anti-IL13 clone B-B13 and the murine anti-IL4 clone 8D4-8 were determined by amino acid sequencing. The murine sequences were humanized and then back-translated into nucleotide sequences as described in Example 5 of International Publication No. WO 2009/052081. The parental humanized anti-IL4 V_H and V_L, and parental humanized anti-IL13 V_H and V_L sequences were combined and arranged as shown in Table 1. The shorthand codes in column one of Table 1 were created to simplify discussion of these antibody-like binding proteins. The antibody-like binding proteins differ in the size of the linker inserted between the two variable regions as shown in Table 1. DNA molecules encoding the polypeptides shown in Table 1 were generated from the back-translated parental anti-IL4 and anti-IL13 antibodies. C_{H1}, C_L, and Fc domains were obtained from IGHG1 (GenBank Accession No. 569F4) and IGKC (GenBank Accession No. Q502W4).

Table 1. Cross-over Double Head Immunoglobulins

Shorthand Code for Protein	Protein Description	SEQ ID NO:
Parental anti-IL4 Light Chain	anti-IL4 V _L	1
Parental anti-IL4 Heavy Chain	anti-IL4 V _H	2
Parental anti-IL13 Light Chain	anti-IL13 V _L	3
Parental anti-IL13 Heavy Chain	anti-IL13 V _H	4
Heavy Chain Codes		
IL13(G4S)IL4CH1-Fc	anti-IL13 V _H -(G4S)-anti-IL4 V _H -C _{H1} -Fc	5
IL13(G4S2)IL4CH1-Fc	anti-IL13 V _H -(G4S) ₂ -anti-IL4 V _H -C _{H1} -Fc	6
IL4(G4S)IL13CH1-Fc	anti-IL4 V _H -(G4S)-anti-IL13 V _H -C _{H1} -Fc	7
IL4(G4S2)IL13CH1-Fc	anti-IL4 V _H -(G4S) ₂ -anti-IL13 V _H -C _{H1} -Fc	8
Light Chain Codes		
IL13(G4S)IL4CL	anti-IL13 V _L -(G4S)-anti-IL4 V _L -CL	9
IL13 (G4S2)IL4CL	anti-IL13 V _L -(G4S) ₂ -anti-IL4 V _L -CL	10
IL4(G4S)IL13CL	anti-IL4 V _L -(G4S)-anti-IL13 V _L -CL	11
IL4(G4S2)IL13CL	anti-IL4 V _L -(G4S) ₂ -anti-IL13 V _L -CL	12

[0123] The protein combinations shown in Table 2 were expressed by transient transfection and purified by Protein A chromatography. In each case, size exclusion chromatography revealed less than 12% aggregation, with most having less than 7% aggregation; but none of the cross-over double head immunoglobulins were found to display any ability to bind either IL4 or IL13. However, no functional antibody-like binding could be detected and the reasons for this lack of activity could not be ascertained. It was previously predicted that this arrangement would show superior stability over the dual variable region domain antibodies described in U.S. Patent No. 7,612,181 and International Publication No. WO 2009/052081.

Table 2. Binding of CODH-Ig to IL4 and IL13

Protein Combination	Aggregation	IL4 binding	IL13 binding
anti-IL13 V _H -(G4S)-anti-IL4 V _H -C _{H1} -Fc anti-IL4 V _L -(G4S)-anti-IL13 V _L -CL	5.4%	ND*	ND
anti-IL13 V _H -(G4S)-anti-IL4 V _H -C _{H1} -Fc anti-IL4 V _L -(G4S) ₂ -anti-IL13 V _L -CL	6.3%	ND	ND
anti-IL13 V _H -(G4S) ₂ -anti-IL4 V _H -C _{H1} -Fc anti-IL4 V _L -(G4S)-anti-IL13 V _L -CL	11.5%	ND	ND
anti-IL13 V _H -(G4S) ₂ -anti-IL4 V _H -C _{H1} -Fc anti-IL4 V _L -(G4S) ₂ -anti-IL13 V _L -CL	10.1%	ND	ND
anti-IL4 V _H -(G4S)-anti-IL13 V _H -C _{H1} -Fc anti-IL13 V _L -(G4S)-anti-IL4 V _L -CL	2.7%	ND	ND
anti-IL4 V _H -(G4S)-anti-IL13 V _H -C _{H1} -Fc anti-IL13 V _L -(G4S) ₂ -anti-IL4 V _L -CL	3.6%	ND	ND
anti-IL4 V _H -(G4S) ₂ -anti-IL13 V _H -C _{H1} -Fc anti-IL13 V _L -(G4S)-anti-IL4 V _L -CL	2.9%	ND	ND
anti-IL4 V _H -(G4S) ₂ -anti-IL13 V _H -C _{H1} -Fc anti-IL13 V _L -(G4S) ₂ -anti-IL4 V _L -CL	10.8%	ND	ND
* ND means none detected			

Example 2. Design of CODV-Ig Proteins by Molecular Modeling

[0124] To obtain fully functional antibody-like proteins utilizing the cross-over double head configuration that are amendable to incorporation of the Fc and C_{L1} domains, a molecular modeling protocol was developed for the inclusion and evaluation of different linkers between the constant and variable domains and between the dual variable domains on both the heavy and light chains. The question was whether the addition of unique linkers between each constant/variable domain interface and between the two variable/variable domain interfaces on both the heavy and light chains would allow proper protein folding to occur and produce functional antibody-like molecules in the cross-over dual variable region configuration (see Figure 2). In other words, a total of four independent and unique linkers were evaluated (see Figure 2). This molecular modeling protocol was based on protein-protein docking of homology models and experimental models of the Fv_{IL4} and Fv_{IL13} regions, respectively, in combination with appropriate linkers between the Fv_{IL4} and Fv_{IL13} regions and between the Fv and constant or Fc regions.

[0125] The independent linkers were assigned unique names as follows: L₁ refers to the linker between N-terminal V_L and the C-terminal V_L on the light chain; L₂ refers to the linker between the C-terminal V_L and C_L on the light chain; L₃ refers to the linker between N-terminal V_H and the C-terminal V_H on the heavy chain; L₄ refers to the linker between the C-terminal V_H and C_{H1} (and Fc) on the heavy chain. It should be noted that the designations V_H and V_L refer only to the domain's location on a particular protein chain in the final format. For example, V_{H1} and V_{H2} could be derived from V_{L1} and V_{L2} domains in parent antibodies and placed into the V_{H1} and V_{H2} positions in a CODV-Ig. Likewise, V_{L1} and V_{L2} could be derived from V_{H1} and V_{H2} domains in parent antibodies and placed into the V_{H1} and V_{H2} positions in a CODV-Ig. Thus, V_H and V_L designations refer to present location and not the original location in a parent antibody.

[0126] In more detail, a homology model of Fv_{IL4} was constructed on PDB entries 1YLD (light chain) and 1IQW (heavy chain). The Fv_{IL4} dimer was recomposed on an in-house crystal structure of the IL13/anti-IL13 Fab_{IL13} complex and optimized. In order to obtain an estimate of the volume required by IL4 when bound to Fv_{IL4}, the crystal structure of IL4 (1RCB.pdb) was docked to the homology model of Fv_{IL4}. Next, twenty-two putative models of the complex were generated that merited further consideration.

[0127] In parallel, the homology model of Fv_{IL4} was docked to Fv_{IL13} extracted from an in-house crystal structure of the IL13/Fab_{IL13} complex. One superior solution was found that permitted construction of relatively short linkers while showing no steric interference for antigen binding and placement of the constant domains as was the case for dual variable region immunoglobulins (see Figure 3). In this arrangement Fv_{IL4} (V_{L1}) was placed at the N-terminus of the light chain, followed by Fv_{IL13} (V_{L2}) and Fc (C_{L1}) on the light chain C-terminus. On the heavy chain, Fv_{IL13} (V_{H2}) was placed N-terminally, followed by Fv_{IL4} (V_{H1}) and the constant regions (C_{H1} - C_{H2} - C_{H3}).

[0128] As shown in Table 3, the models of the light chain suggested that the linker L₁ between the V_{L1} and V_{L2} domains and the linker L₂ between the V_{L2} and C_{L1} domains should be between one to three and zero to two glycine residues long, respectively. Models of the heavy chain suggested that the linker L₃ between the V_{H2} and V_{H1} domains and the linker L₄ between the V_{H1} and C_{H1} domains should be between two to six and four to seven glycine residues long, respectively (see Table 3 and Figure 2). In this example, glycine was used as a prototypical amino acid for the linkers but other amino acid residues may also serve as linkers. The structural stability of the proposed models was verified by optimization of the linker conformations, minimization, and molecular dynamics calculations. Systematic combination between four light chain and six heavy chain constructs resulted in 24 possible cross-over dual variable region bispecific anti-IL4 and anti-IL13 antibody-like binding proteins (see Table 4).

Table 3. Proposed Linker Lengths

Linker Between	Maximal Linker Insertion	Minimal Linker Insertion	Linker Name
V _{L1} -V _{L2}	Gly ₃	Gly	L ₁
V _{L2} -C _L	Gly ₂	None	L ₂
V _{H2} -V _{H1}	Gly ₆	Gly ₂	L ₃
V _{H1} -C _{H1}	Gly ₇	Gly ₄	L ₄

Table 4. CODV-Ig for Expression

Code*	Heavy Chains (N to C terminal)	SEQ ID NO:
HC1	IL13 V _H -(Gly6)-IL4 V _H -(Gly7)-C _{H1} -Fc	13
HC2	IL13 V _H -(Gly6)-IL4 V _H -(Gly4)-C _{H1} -Fc	14

(continued)

Code*	Heavy Chains (N to C terminal)	SEQ ID NO:
HC3	IL13 V _H -(Gly2)-IL4 V _H -(Gly7)- C _{H1} -Fc	15
HC4	IL13 V _H -(Gly2)-IL4 V _H -(Gly4)- C _{H1} -Fc	16
HC5	IL13 V _H -(Gly4)-IL4 V _H -(Gly7)- C _{H1} -Fc	17
HC6	IL13 V _H -(Gly4)-IL4 V _H -(Gly4)- C _{H1} -Fc	18
Code*	Light Chains (N to C terminal)	
LC1	IL4 V _L -(Gly3)-IL13 V _L -C _{L1}	19
LC2	IL4 V _L -(Gly)-IL13 V _L - C _{L1}	20
LC3	IL4 V _L -(Gly3)-IL13 V _L -(Glv2)- C _{L1}	21
LC4	IL4 V _L -(Gly)-IL13 V _L -(Gly2)- C _{L1}	22
C _{L1}	C _{L1} human light chain constant domain	23
C _{H1} -Fc	C _{H1} human heavy chain constant domain and Fc region	24
Gly4	peptide linker with 4 glycines (GGGG)	25
Gly5	peptide linker with 5 glycines (GGGGG)	26
Glv6	peptide linker with 6 glycines (GGGGGG)	27
Gly7	peptide linker with 7 glycines (GGGGGGG)	28
Gly8	peptide linker with 8 glycines (GGGG GGGG)	29
*A short hand code was devised to represent the associated structures. Codes beginning with HC represent the adjacent heavy chain and codes beginning with LC represent the adjacent light chains.		

[0129] In Table 4, the prefix "anti" is not included but it is intended to mean that IL13 refers to anti-IL13 and IL4 refers to anti-IL4.

Example 3. Generation of CODV-Ig Expression Plasmids

[0130] Nucleic acid molecules encoding the variable heavy and light chains of the six heavy chains and four light chains described in Table 4 were generated by gene synthesis at Geneart (Regensburg, Germany). The variable light chain domains were fused to the constant light chain (IGKC, GenBank Accession No. Q502W4) by digestion with the restriction endonucleases ApaI and BsiWI and subsequently ligated into the ApaI/BsiWI sites of the episomal expression vector pFF, an analogon of the pTT vector described by Durocher et al., (2002, Nucl. Acids Res. 30(2): E9), creating the mammalian expression plasmid for expression of the light chains.

[0131] The variable heavy chain domains were fused to the "Ted" variant of the human constant heavy chain (IGHG1, GenBank Accession No. 569F4), or alternatively, to a 6x His tagged C_{H1} domain from the human constant IGHG1 in order to create a bispecific Fab. Next, the V_H domain was digested with the restriction endonucleases ApaI and Apal and then fused to the IGHG1 or His tagged C_{H1} domain respectively, by ligation into the ApaI/Apal sites of the episomal expression vector pFF, creating the mammalian expression plasmids for expression of the heavy chains (IgG1 or Fab respectively).

Example 4. Expression of CODV-Ig

[0132] The expression plasmids encoding the heavy and light chains of the corresponding constructs were propagated in *E. coli* DH5a cells. Plasmids used for transfection were prepared from *E. coli* using the Qiagen EndoFree Plasmid Mega Kit.

[0133] HEK 293-FS cells growing in Freestyle Medium (Invitrogen) were transfected with indicated LC and HC plasmids encoding the heavy chains and light chains shown in Table 4 using 293fectin (Invitrogen) transfection reagent as described

by the manufacturer. After 7 days, cells were removed by centrifugation and the supernatant was passed over a 0.22 μm filter to remove particles.

[0134] CODV-IgG1 constructs were purified by affinity chromatography on Protein A columns (HiTrap Protein A HP Columns, GE Life Sciences). After elution from the column with 100 mM acetate buffer and 100 mM NaCl, pH 3.5, the CODV-IgG1 constructs were desalted using HiPrep 26/10 Desalting Columns, formulated in PBS at a concentration of 1 mg/mL and filtered using a 0.22 μm membrane.

[0135] Bispecific CODV Fab constructs were purified by IMAC on HiTrap IMAC HP Columns (GE Life Sciences). After elution from the column with a linear gradient (Elution buffer: 20 mM sodium phosphate, 0.5 M NaCl, 50 - 500 mM imidazole, pH 7.4), the protein containing fractions were pooled and desalted using HiPrep 26/10 Desalting Columns, formulated in PBS at a concentration of 1 mg/mL and filtered using a 0.22 μm membrane.

[0136] Protein concentration was determined by measurement of absorbance at 280 nm. Each batch was analyzed by SDS-PAGE under reducing and non-reducing conditions to determine the purity and molecular weight of each subunit and of the monomer.

[0137] A Nunc F96-MaxiSorp-Immuno plate was coated with goat anti-Human IgG (Fc specific) [NatuTec A80-104A]. The antibody was diluted to 10 $\mu\text{g}/\text{ml}$ in carbonate coating buffer (50 mM sodium carbonate, pH 9.6) and dispensed at 50 μL per well. The plate was sealed with adhesive tape, and stored overnight at 4° C. The plate was washed three times with Wash buffer (PBS, pH 7.4 and 0.1% Tween20). 150 μL of blocking solution (1% BSA / PBS) was dispensed into each well to cover the plate. After 1 hour at room temperature, the plate was washed three times with Wash buffer. 100 μL of sample or standards (in a range from 1500 ng/ml to 120 ng/ml) were added and allowed to sit for 1 hour at room temperature. The plate was washed three times with Wash buffer. 100 μL of goat anti-Human IgG-FC - HRP conjugate [NatuTec A80-104P-60] diluted 1:10.000 were added using incubation solution (0.1% BSA, PBS, pH 7.4, and 0.05% Tween20). After 1 hour incubation at room temperature, the plate was washed three times with Wash buffer. 100 μL of ABTS substrate (10 mg ABTS tablet (Pierce 34026) in 0.1 M Na_2HPO_4 0.05 M citric acid solution, pH 5.0). Addition of 10 μL of 30% H_2O_2 / 10 ml Substrate buffer prior to use) were dispensed to each well, and the color was allowed to develop. After the color developed (approximately 10 to 15 minutes), 50 μL of 1% SDS solution were added to stop the reaction. The plate was read at A_{405} .

Example 5. Characterization of CODV-Ig Variants

[0138] To determine whether the CODV-Ig antibody-like protein heavy and light chains were pairing and folding properly, the aggregation level was measured by analytical size-exclusion chromatography (SEC). Analytical SEC was performed on assembled pairs using an AKTA explorer 10 (GE Healthcare) equipped with a TSKgel G3000SWXL column (7.8 mm x 30 cm) and TSKgel SWXL guard column (Tosoh Bioscience). The analysis was run at 1 ml/min using 250 mM NaCl, 100 mM Na-phosphate, pH 6.7, with detection at 280 nm. 30 μL of protein sample (at 0.5-1 mg/ml) were applied onto the column. For estimation of the molecular size, the column was calibrated using a gel filtration standard mixture (MWGF-1000, SIGMA Aldrich). Data evaluation was performed using UNICORN software v5.11.

[0139] Table 5 shows the results of the first set of 24 different CODV-Ig molecules made using the anti-IL4 and anti-IL13 variable region combinations described in Table 4. The codes assigned in Table 4 represent the adjacent structures shown in Table 4. For the pairs of light chain and heavy chains where protein was produced, aggregation levels were measured using SEC. The results are shown in Table 5 where LC4 ($L_1 = 1$; $L_2 = 2$) was most successful in pairing with all six heavy chains. LC4 corresponds to the structure IL4 V_L -(Gly)-IL13 V_L -(Gly2)- C_{L1} having linker L_1 equal to 1, where a single amino acid residue separated the two V_L domains of the dual variable region light chain. In addition, LC4 had L_2 equal to 2, which contained a Gly-Gly dipeptide linker between the central V_L and the C-terminal C_{H1} .

Table 5. Levels of Aggregation Among Pairs of Heavy and Light Chains

	HC1	HC2	HC3	HC4	HC5	HC6
LC1	>50%	>50%	ND*	ND	ND	ND
LC2	ND	ND	ND	>50%	ND	ND
LC3	ND	ND	ND	ND	ND	ND
LC4	7.2%	6.8%	6.8%	7.1%	6.3%	5.9%
*ND indicates that no protein was produced						

[0140] Where CODV-Ig molecules were produced, a single-concentration BIACORE experiment at intermediate IL13 and IL4 concentrations was performed to verify binding to target antigens. CODV-Ig antibody-like molecules correspond-

ing to the LC4:HC4 and LC4:HC6 combinations described in Table 4 were chosen for assessment of a full kinetic analysis using surface plasmon resonance.

[0141] As depicted in Table 5, most of the CODV-Ig molecules could not be produced at all or only as aggregates (up to 90%). The heavy chain/light chain combinations giving rise to acceptable aggregation levels (5-10%) after one chromatography step were the ones combined with the light chain IL4 V_L-(Gly)-IL13 V_L-(Gly2)-C_L1. The light chain was the most finicky chain within these CODV-Ig variants and served as platform to accept different heavy chains with different linker compositions.

1. Kinetic Analysis

[0142] Two pairs of heavy and light chains were selected for full kinetic analysis. Recombinant human IL13 and IL4 was purchased from Chemicon (USA). Kinetic characterization of purified antibodies was performed using surface plasmon resonance technology on a BIACORE 3000 (GE Healthcare). A capture assay using a species specific antibody (e.g., human-Fc specific MAB 1302, Chemicon) for capture and orientation of the investigated antibodies was used. The capture antibody was immobilized via primary amine groups (11000 RU) on a research grade CM5 chip (GE Life Sciences) using standard procedures. The analyzed antibody was captured at a flow rate of 10 µL/min with an adjusted RU value that would result in maximal analyte binding of 30 RU. Binding kinetics were measured against recombinant human IL4 and IL13 over a concentration range between 0 to 25 nM in HBS EP (10 mM HEPES, pH 7.4, 150 mM NaCl, 3 mM EDTA, and 0.005 % Surfactant P20) at a flow rate of 30 µL/min. Chip surfaces were regenerated with 10 mM glycine, pH 2.5. Kinetic parameters were analyzed and calculated in the BIAevaluation program package v4.1 using a flow cell without captured antibody as reference.

[0143] Table 6 below shows the comparison of kinetics of the parental BB13 (anti-IL13) and 8D4 (anti-IL4) antibodies (expressed as IgGs) with the respective domains within the CODV-Ig format (Table 4, Codes LC4:HC4 and LC4:HC6). As shown in Table 6, the CODV-Ig constructs did not exhibit decreased binding properties against the corresponding antigens, when compared to the parental anti-IL13 and anti-IL4 antibodies. The loss in the on-rate observed in the DVD-Ig/TBTI format using the same Fv sequences did not occur with the CODV-Ig configuration. The opposite facing binding sites should allow for binding large antigens or bridging different cells with a bispecific antibody-like configuration, and would also be suitable for a wider selection of parental antibodies. A further advantage of the CODV-Ig was that no linker residues protrude into the antigen binding site and reduce accessibility of the antigen.

Table 6. Kinetic Analysis of LC4:HC4 and LC4:HC6

Code		On-rate [1/Ms]	Off-rate [1/s]	KD [M]
	parental anti-IL4 mAb	2.49E+07	1.95E-04	7.83E-12
	parental anti-IL13 mAb	1.59E+06	1.30E-04	8.18E-11
LC4:HC4	CODV-Ig with IL4	3.16E+07	2.89E-04	9.14E-12
LC4:HC4	CODV-Ig with IL13	1.20E+06	1.12E-04	9.34E-11
LC4:HC6	CODV-Ig with IL4	2.97E+07	3.30E-04	1.11E-11
LC4:HC6	CODV-Ig with IL13	1.39E+06	1.63E-04	1.18E-10

2. Co-injection of IL4 and IL13 for Demonstration of Additive Antigen Binding by CODV-Ig

[0144] To investigate additive binding of both antigens, a wizard-driven co-inject method was applied in which one antigen was injected immediately followed by the other antigen after a lag time (IL4 then IL13 and vice versa). The resulting binding level could be compared to that achieved with a 1:1 mixture of both antigens in the same concentration. In order to show additive binding of both the IL4 and IL13 antigens by the CODV-Ig molecules, a BIACORE experiment was performed with a CODV-Ig combination [HC4:LC4] by co-injection of both antigens in three separate analysis cycles (see Figure 4). The co-injection was done with 3.125 nM IL4/25 nM IL13 (and vice versa) and with a 1:1 mixture of 3.125 nM IL4 and 25 nM IL13. A co-injection of HBS-EP buffer was done as a reference. At a time point of 800 seconds, an identical binding level of 63 RU was achieved after injection of the antigen mixture or co-injection of the antigens, regardless of the co-injection sequence. When CODV-Ig protein had been saturated by the first antigen (IL4), the second antigen (IL13) was injected and a second binding signal was observed. This observation was reproduced when the antigen injection sequence was reversed. This demonstrates the additive and non-inhibition of binding of both antigens by the CODV-Ig. Therefore, the CODV-Ig construct was able to bind both antigens simultaneously (*i.e.*, exhibit bispecificity) saturating all binding sites (*i.e.*, exhibited tetravalency).

Example 6. Tolerance of Linker Lengths for CODV-Ig

[0145] The tolerance for linkers of various lengths was evaluated by constructing CODV-Ig molecules having different combinations of linker lengths for L_1 , L_2 on the light chain and L_3 and L_4 on the heavy chain. CODV-Ig constructs were generated with heavy chain linkers L_3 and L_4 varying between 1 through 8 residues for L_3 and either 0 or 1 residues for L_4 . The heavy chain contained anti-IL4 as the N-terminal binding domain and anti-IL13 as the C-terminal binding domain followed by C_{H1} -Fc. The light chain linkers L_1 and L_2 were varied from 3 to 12 residues for L_1 and from 3 to 14 residues for L_2 . The light chain contained anti-IL13 as the N-terminal binding domain and anti-IL4 as the C-terminal binding domain followed by C_{L1} .

1. Characterization of CODV-Ig Variants

[0146] Determination of aggregation level was by analytical size-exclusion chromatography (SEC). Analytical SEC was performed using an ÄKTA explorer 10 (GE Healthcare) equipped with a TSKgel G3000SWXL column (7.8 mm x 30 cm) and TSKgel SWXL guard column (Tosoh Bioscience). The analysis was run at 1 ml/min using 250 mM NaCl, 100 mM Na-phosphate pH 6.7, with detection at 280 nm. 30 μ L of protein sample (at 0.5-1 mg/ml) was applied onto the column. For estimation of the molecular size, the column was calibrated using a gel filtration standard mixture (MWGF-1000, SIGMA Aldrich). Data evaluation was performed using UNICORN software v5.11.

[0147] Recombinant human IL13 and IL4 were purchased from Chemicon (USA). Recombinant human TNF- α was purchased from Sigma Aldrich (H8916-10 μ g), recombinant human IL-1 β (201-LB/CF), recombinant human IL-23 (1290-IL/CF), recombinant human EGFR (344 ER), and recombinant human HER2 (1129-ER-50) were purchased from R&D Systems.

[0148] Kinetic binding analysis by Biacore was performed as follows. Surface plasmon resonance technology on a Biacore 3000 (GE Healthcare) was used for detailed kinetic characterization of purified antibodies. A capture assay using a species-specific antibody (e.g., human-Fc specific MAB 1302, Chemicon) for capture and orientation of the investigated antibodies was used. For determination of IL4 and IL13 binding kinetics, the corresponding CODV Fabs as in Example 10, Table 12 were captured using the anti-human Fab capture Kit (GE Healthcare). The capture antibody was immobilized via primary amine groups (11000 RU) on a research grade CM5 chip (GE Life Sciences) using standard procedures. The analyzed antibody was captured at a flow rate of 10 μ L/min with an adjusted RU value that would result in maximal analyte binding of 30 RU. Binding kinetics were measured against recombinant human IL4 and IL13 over a concentration range between 0 to 25 nM in HBS EP (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA, 0.005% Surfactant P20) at a flow rate of 30 μ L/min. Chip surfaces were regenerated with 10 mM glycine pH 2.5. Kinetic parameters were analyzed and calculated in the BIAevaluation program package v4.1 using a flow cell without captured antibody as a reference.

[0149] Binding affinities of CODV-Ig, CODV-Fab, and TBTI against EGFR and HER2 were measured using a Proteon XPR36 protein interaction array system (Biorad). The antigens were immobilized by amine reactive coupling on GLC sensor chips (Biorad). Dilution series of the bispecific antibody variants in PBSET buffer (Biorad) were analyzed in parallel in one-shot kinetics mode with double referencing. Data were analyzed using Proteon Manager Software v3.0 (Biorad) with either Langmuir 1:1 model with mass transfer or bivalent analyte model.

[0150] Table 7 summarizes the results for yield, aggregation (as measured by size exclusion chromatography), and binding affinity for CODV-Ig having different size combinations of linkers. The results revealed that CODV-Ig molecules in which L_2 was zero generally could not be produced, or where protein was produced, there was a high level of aggregation (see Batch ID Nos. 101, 102, 106-111, and 132-137 in Table 7). Therefore, in contrast to the molecular modeling prediction from Example 2, where L_2 equal to zero was within the acceptable range, these results indicate that the V_{L2} - C_L transition (or L_2) requires a linker of at least one residue (see Table 7).

Table 7. Optimization of Linker Sizes for CODV-Ig

Batch ID	Alignment on LC1*	L_1	L_2	L_3	L_4	Yield ¹ [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
101	IL4 x IL13	0	0	0	0	n.p.	-	-	-
102	IL4 x IL13	0	0	2	4	1.0	66.4	35	125
103	IL4 x IL13	0	1	2	4	2.0	10.8	4	124
104	IL4 x IL13	0	2	2	4	1.0	10.0	3	137
105	IL4 x IL13	0	2	4	4	2.0	8.5	6	94

EP 2 691 416 B1

(continued)

	Batch ID	Alignment on LC1*	L ₁	L ₂	L ₃	L ₄	Yield ¹ [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
5	106	IL4 x IL13	1	0	2	4	4.2	61.0	-	-
	107	IL4 x IL13	1	0	2	7	n.p.	-	-	-
	108	IL4 x IL13	1	0	4	4	n.p.	-	-	-
10	109	IL4 x IL13	1	0	4	7	n.p.	-	-	-
	110	IL4 x IL13	1	0	6	4	3.0	75.0	-	-
	111	IL4 x IL13	1	0	6	7	5.6	91.0	-	-
15	112	IL4 x IL13	1	1	2	4	5.3	8.8	5	55
	113	IL4 x IL13	1	2	0	0	2.0	1.9	6	59
	114	IL4 x IL13	1	2	0	2	1.0	7.2	4	75
	115	IL4 x IL13	1	2	0	4	2.0	8.0	11	62
20	116	IL4 x IL13	1	2	1	4	3.0	5.7	5	74
	117	IL4 x IL13	1	2	12	12	4.0	9.2	7	75
	118	IL4 x IL13	1	2	12	9	7.0	11.8	7	74
25	119	IL4 x IL13	1	2	15	15	10.0	7.9	6	64
	120	IL4 x IL13	1	2	2	2	2.0	5.9	6	70
	121	IL4 x IL13	1	2	2	4	7.0	7.1	11	118
	122	IL4 x IL13	1	2	2	7	6.6	6.8	1	50
30	123	IL4 x IL13	1	2	4	4	5.8	5.9	9	93
	124	IL4 x IL13	1	2	4	7	5.6	6.3	6	51
	125	IL4 x IL13	1	2	6	4	7.6	6.8	8	64
35	126	IL4 x IL13	1	2	6	7	8.2	7.2	9	56
	127	IL4 x IL13	1	2	9	12	4.0	8.7	2	83
	128	IL4 x IL13	1	2	9	9	10.2	14.0	12	94
	129	IL4 x IL13	1	3	2	4	4.0	4.7	5	70
40	130	IL4 x IL13	1	4	2	4	7.1	6.0	4	57
	131	IL4 x IL13	2	2	2	4	1.0	7.3	5	50
	132	IL4 x IL13	3	0	2	4	n.p.	-	-	-
45	133	IL4 x IL13	3	0	2	7	n.p.	-	-	-
	134	IL4 x IL13	3	0	4	4	n.p.	-	-	-
	135	IL4 x IL13	3	0	4	7	n.p.	-	-	-
	136	IL4 x IL13	3	0	6	4	9.8	93.0	-	-
50	137	IL4 x IL13	3	0	6	7	5.8	90.0	-	-
	138	IL4 x IL13	3	2	2	4	n.p.	-	-	-
	139	IL4 x IL13	3	2	2	7	n.p.	-	-	-
55	140	IL4 x IL13	3	2	4	4	n.p.	-	-	-
	141	IL4 x IL13	3	2	4	7	n.p.	-	-	-
	142	IL4 x IL13	3	2	6	4	n.p.	-	-	-

(continued)

Batch ID	Alignment on LC1*	L ₁	L ₂	L ₃	L ₄	Yield ¹ [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
143	IL4 x IL13	3	2	6	7	n.p.	-	-	-
* Alignment on the heavy chain must be IL13V _H -L ₃ -IL4V _H -L ₄ -C _{H1} -Fc									
¹ n.p. means the construct was not producible.									

[0151] In addition, the CODV-Ig linker lengths described above were found to be more sensitive to increases in 1 amino acid residue than increases in 2 amino acid residues. For example, while Batch ID Nos. 103 and 104 differ by 1 amino acid residue in L₂, Batch ID No. 103 shows 6 fold more aggregation and Batch ID No. 104 displays less aggregation and twice the yield. In contrast, Batch ID Nos. 104 and 105, which differ by two residues in L₂, displayed similar profiles with respect to yield, aggregation, and binding.

Example 7. Heavy Chain as Template Chain for CODV-Ig

[0152] In examples 1 through 5, the optimal short linker sizes on the light chain suggested that the light chain was serving as a template by remaining in a linear arrangement and that larger linkers were required on the heavy chain in order for the heavy chain to fold properly into the cross-over configuration to conform to the template light chain (see Figure 5, Panel A). Whether the short linkers specifically placed on the heavy chain to maintain a linear arrangement on the heavy chain rendered the heavy chain the "template" chain, and whether the pattern would repeat itself and larger linkers would be required to allow the non-template chain to fold properly and accommodate the now template heavy chain was evaluated next (see Figure 5, Panel B).

[0153] Figure 6 illustrates these principles of CODV-Ig design based on having either the light chain or the heavy chain as the "template." To evaluate the generic nature of this concept, CODV-Ig constructs were generated with heavy chain linkers L₃ and L₄ varying between 1 through 8 residues for L₃ and either 0 or 1 residues for L₄. The heavy chain contained anti-IL4 as the N-terminal binding domain and anti-IL13 as the C-terminal binding domain followed by C_{H1}-Fc. The light chain linkers L₁ and L₂ were varied from 3 to 12 residues for L₁ and from 3 to 14 residues for L₂. The light chain contained anti-IL13 as the N-terminal binding domain and anti-IL4 as the C-terminal binding domain followed by C_{L1}.

[0154] Table 8 summarizes the results for yield, aggregation (as measured by size exclusion chromatography), and binding affinity for CODV-Ig having different size combinations of linkers and where the heavy chain is maintained in a linear arrangement as the template chain and the light chain is allowed to fold in a cross-over configuration. The results revealed that CODV-Ig molecules in which L₄ was zero generally could not be produced, or where protein was produced, there was a high level of aggregation (similar to molecules in which L₂ was equal to zero) (see Batch ID Nos. 207-209, 211-212, 219-224, 231-236, 243-252, and 263-266 in Table 8). One exception was Batch ID No. 210, in which L₁ was 7, L₂ was 5, L₃ was 2, and L₄ was zero. This arrangement produced a sufficient amount of protein and had an acceptable level of aggregation and binding, which suggested that some combination of linker sizes could be found to compensate for a zero length linker at L₄ in some circumstances.

Table 8. Optimization of Linker Sizes with Heavy Chain as Template

Batch ID	Alignment on HC*	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
201	IL4 x IL13	5	3	1	2	6.3	12	4	72
202	IL4 x IL13	5	5	1	2	10.5	7.0	9	54
203	IL4 x IL13	7	3	1	2	19.3	9.4	80	46
204	IL4 x IL13	7	5	1	2	15.3	5.2	3	25
205	IL4 x IL13	10	3	1	2	4.7	4.0	8	58
206	IL4 x IL13	10	5	1	2	9.1	3.9	4	58
207	IL4 x IL13	5	3	2	0	6.7	25.3	3	33
208	IL4 x IL13	5	5	2	0	10.2	18.4	10	77
209	IL4 x IL13	7	3	2	0	16.2	22.2	5	47

EP 2 691 416 B1

(continued)

	Batch ID	Alignment on HC*	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
5	210	IL4 x IL13	7	5	2	0	14.7	9.7	4	47
	211	IL4 x IL13	10	3	2	0	2.1	12.8	7	53
	212	IL4 x IL13	10	5	2	0	7.0	36.3	10	29
10	213	IL4 x IL13	5	3	2	2	4.0	13.2	5	27
	214	IL4 x IL13	5	5	2	2	8.0	7.9	10	53
	215	IL4 x IL13	7	3	2	2	14.9	11.5	4	50
	216	IL4 x IL13	7	5	2	2	7.5	3.6	11	40
15	217	IL4 x IL13	10	3	2	2	2.4	4.4	8	79
	218	IL4 x IL13	10	5	2	2	4.6	6.6	4	36
	219	IL4 x IL13	3	6	3	0	2.1	51.8	8	71
20	220	IL4 x IL13	3	10	3	0	3.9	59.4	1	42
	221	IL4 x IL13	3	14	3	0	1.9	57.6	35	81
	222	IL4 x IL13	6	6	3	0	4.0	11.8	7	53
	223	IL4 x IL13	6	10	3	0	10.3	16.6	6	23
25	224	IL4 x IL13	6	14	3	0	5.1	13.5	9	52
	225	IL4 x IL13	3	6	3	2	2.8	71.6	6	68
	226	IL4 x IL13	3	10	3	2	7.3	65.8	6	64
30	227	IL4 x IL13	3	14	3	2	1.6	53.6	7	39
	228	IL4 x IL13	6	6	3	2	4.0	19.1	7	44
	229	IL4 x IL13	6	10	3	2	2.2	15.4	3	14
	230	IL4 x IL13	6	14	3	2	4.0	16.2	6	76
35	231	IL4 x IL13	5	3	5	0	n.p.	-	-	-
	232	IL4 x IL13	5	5	5	0	0.6	24.9	8	70
	233	IL4 x IL13	7	3	5	0	0.4	15.1	3	113
40	234	IL4 x IL13	7	5	5	0	1.3	30.7	3	122
	235	IL4 x IL13	10	3	5	0	0.1	11.3	2	82
	236	IL4 x IL13	10	5	5	0	0.4	18.6	11	112
	237	IL4 x IL13	5	3	5	2	2.1	45.3	8.1	101.0
45	238	IL4 x IL13	5	5	5	2	0.6	45.4	9.3	67.2
	239	IL4 x IL13	7	3	5	2	n.p.	-	-	-
	240	IL4 x IL13	7	5	5	2	1.6	31.7	4	65
50	241	IL4 x IL13	10	3	5	2	0.2	14.7	7	119
	242	IL4 x IL13	10	5	5	2	1.1	17.6	10	37
	243	IL4 x IL13	3	6	6	0	1.6	54.3	5	50
	244	IL4 x IL13	3	10	6	0	1.5	63.9	10	10
55	245	IL4 x IL13	3	14	6	0	1.0	61.5	10	69
	246	IL4 x IL13	6	6	6	0	1.1	16.2	6	57

EP 2 691 416 B1

(continued)

Batch ID	Alignment on HC*	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen1) IL4 [pM]	KD (Antigen2) IL13 [pM]
247	IL4 x IL13	6	10	6	0	4.7	27.9	2	41
248	IL4 x IL13	6	14	6	0	0.9	18.1	10	79
249	IL4 x IL13	10	6	6	0	0.3	8.7	3	87
250	IL4 x IL13	10	8	6	0	0.7	21.3	8	53
251	IL4 x IL13	12	6	6	0	1.3	9.7	8	70
252	IL4 x IL13	12	8	6	0	1.3	11.7	7	85
253	IL4 x IL13	3	6	6	2	5.1	66.8	6	66
254	IL4 x IL13	3	10	6	2	2.4	62.4	6	80
255	IL4 x IL13	3	14	6	2	2.0	72.1	2	60
256	IL4 x IL13	6	6	6	2	2.0	32.4	4	81
257	IL4 x IL13	6	10	6	2	1.9	29.8	7	30
258	IL4 x IL13	6	14	6	2	2.5	24.6	5	70
259	IL4 x IL13	10	6	6	2	1.4	16.4	8	71
260	IL4 x IL13	10	8	6	2	0.8	16.6	10	71
261	IL4 x IL13	12	6	6	2	1.2	12.3	5	265
262	IL4 x IL13	12	8	6	2	1.1	13.2	4	111
263	IL4 x IL13	10	6	8	0	2.4	10.8	2	74
264	IL4 x IL13	10	8	8	0	0.8	8.0	7	22
265	IL4 x IL13	12	6	8	0	1.0	9.5	8	66
266	IL4 x IL13	12	8	8	0	2.0	9.3	3	69
267	IL4 x IL13	10	6	8	2	1.4	15.0	9	170
268	IL4 x IL13	10	8	8	2	1.0	12.9	4	52
269	IL4 x IL13	12	6	8	2	1.2	8.8	5	66
270	IL4 x IL13	12	8	8	2	2.4	11.7	3	72
* Alignment on the light chain must be IL13V _L -L ₁ -IL4V _L -L ₂ -C _{L1}									

[0155] The results from Tables 7 and 8 clearly show that linkers are required between the variable and constant domains to allow optimal folding. Only in rare arrangements was a linker equal to zero tolerated (see Batch ID Nos. 103-105, in which L₁ (LC) was zero, and Batch No. 210, in which L₄ was zero). However, in each case, the corresponding transition linker between the variable region and constant region on the other chain could not be zero.

[0156] The results above indicated that the combinations L₁=7, L₂=5, L₃=1, and L₄=2 were a good starting point for optimizing a new CODV-Ig in which the heavy chain is the template. The ranges in Table 9 were shown to be reasonable ranges for successfully engineering a new CODV-Ig from two parent antibodies.

Table 9. Ranges of Linker Sizes for Either LC or HC as Template

Linker	LC as Template		HC as Template	
	Max Range	Min Range	Max Range	Min Range
L ₁	1-3	1-2	3-12	5-10
L ₂	1-4	1-2	3-14	5-8
L ₃	2-15	4-12	1-8	1-5

(continued)

Linker	LC as Template		HC as Template	
	Max Range	Min Range	Max Range	Min Range
L ₄	2-15	2-12	1-3	1-2

Example 8. Universal Applicability of CODV-Ig Format

[0157] To evaluate the suitability of the CODV-Ig format for engineering new antibody-like binding proteins the variable regions from numerous existing human and humanized antibodies having specificity for insulin-like growth factor 1 receptor (IGF1R(1)), a second antibody to insulin-like growth factor 1 receptor (IGF1R(2)), human epidermal growth factor receptor 2 (HER2), epidermal growth factor receptor (EGFR), tumor necrosis factor - alpha (TNF α), Interleukin 12 and 23 (IL-12/23) and interleukin 1beta (IL-1 β) were incorporated into the CODV-Ig format (see Table 10).

Table 10. Descriptive Codes for Heavy and Light Chains Used in Bispecific CODV-Ig

Code*	Heavy Chains (N to C terminal)	SEQ ID NO:
HC10	IGF1R(1) V _H -(Gly)-HER2 V _H -(Gly2)-C _{H1} -Fc	32
HC11	HER2 V _H -(Gly)- IGF1R(1) V _H -(Gly2)-C _{H1} -Fc	33
HC12	IGF1R(2) V _H -(Gly)-EGFR V _H -(Gly2)-C _{H1} -Fc	34
HC13	EGFR V _H -(Gly)- IGF1R(2) V _H -(Gly2)- C _{H1} -Fc	35
HC14	TNF α V _H -(Gly)-IL12/23 V _H -(Gly2)- C _{H1} -Fc	36
HC15	IL12/23 V _H -(Gly)-TNF α V _H -(Gly2)- C _{H1} -Fc	37
HC16	TNF α V _H -(Gly)-IL1 β V _H -(Gly2)-C _{H1} -Fc	38
HC17	IL1 β V _H -(Gly)-TNF α V _H -(Gly2)- C _{H1} -Fc	39
Code*	Light Chains (N to C terminal)	
LC10	HER2 V _L -(Gly7)-IGF1R(1) V _L -(Gly5)-C _{L1}	40
LC11	IGF1R(1) V _L -(Gly7)- HER2 V _L -(Gly5)- C _{L1}	41
LC12	EGFR V _L -(Gly7)-IGF1R(2) V _L -(Gly5)-C _{L1}	42
LC13	IGF1R(2) V _L -(Gly7)- EGFR V _L -(Gly5)- C _{L1}	43
LC14	IL12/23 V _L -(Gly7)- TNF α V _L -(Gly5)-C _{L1}	44
LC15	TNF α V _L -(Gly7)- IL12/23 V _L -(Gly5)- C _{L1}	45
LC16	IL1 β V _L -(Gly7)- TNF α V _L -(Gly5)-C _{L1}	46
LC17	TNF α V _L -(Gly7)- IL1 β V _L -(Gly5)-C _{L1}	47
C _{L1}	C _{L1} human light chain constant domain	23
C _{H1} -Fc	C _{H1} human heavy chain constant domain and Fc region	24
Gly4	peptide linker with 4 glycines (GGGG)	25
Gly5	peptide linker with 5 glycines (GGGGG)	26
Gly6	peptide linker with 6 glycines (GGGGGG)	27
Gly7	peptide linker with 7 glycines (GGGGGGG)	28
Gly8	peptide linker with 8 glycines (GGGG GGGG)	29

*A short hand code was devised to represent the associated structures. Codes beginning with HC represent the adjacent heavy chain and codes beginning with LC represent the adjacent light chains.

[0158] The antibody variable regions from known human and humanized antibodies were used to test the universal applicability of the CODV-Ig format in designing bispecific antibody-like binding proteins. In addition, the possibility of positional effects with regard to the placement of certain antibody variable regions either N-terminal or C-terminal on either the heavy chain or the light chain was examined. Based on the design of CODV-Ig molecules having a linker composition of $L_1 = 7$, $L_2 = 5$, $L_3 = 1$, and $L_4 = 2$, different sequences of antibodies were introduced into the CODV-Ig format.

[0159] Activities of bispecific antibodies or derivatives against IL1 β and TNF α were determined by using commercially available HEK-Blue TNF α /IL1 β reporter cells (InvivoGen). To determine antibody activities against TNF α and IL1 β , the cytokines were pre-incubated for 1 hour with different concentrations of the antibodies and added to 50,000 HEK Blue TNF α /IL1 β cells. Cytokine mediated induction of SEAP was measured after 24 hours in the culture supernatant with the QUANTI-Blue assay (InvivoGen).

[0160] As shown in Table 11, all constructs showed good to excellent protein yield and acceptable levels of aggregation (see, in particular, Batch ID Nos. 301 and 302 in Table 11). The measured affinity for each antibody variable domain was within the published or expected affinity. In cases where affinity was assessed, no positional effects were detected. In summary, as shown in the following tables, no positional effects were seen with any of the antibody variable domains used or with the use of these domains on either antibody chain.

Table 11. Universal Use of CODV-Ig Format for Bispecific Antibody-like Binding Proteins

Batch ID DC/LC Codes ¹	Alignment on HC	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen 1) [pM]	KD (Antigen 2) [pM]	IC50 [pM] TNF α Cellular Assay
301 HC10/LC10	IGF1R(1) x HER2	7	5	1	2	70	5.9	n.m.	153 (HER2)	-
302 HC11/LC11	HER2 x IGF1R(1)	7	5	1	2	60	1.8	163 (HER2)	n.m.	-
303 HC12/LC12	IGF1R(2) x EGFR	7	5	1	2	17	2.7	n.m.	-	-
304 HC13/LC13	EGFR x IGF1R(2)	7	5	1	2	9.5	4.3	-	n.m.	-
305 HC14/LC14	TNF α x IL12/23	7	5	1	2	7.1	7.5	321 (TNF α)	65 (IL23)	95
306 HC15/LC15	IL12/23 x TNF α	7	5	1	2	11.9	7.1	118 (IL23)	543 (TNF α)	138
307 HC16/LC16	TNF α x IL1 β	7	5	1	2	6.6	13.6	340 (TNF α)	155 (IL1 β)	136
308 HC17/LC17	IL1 β x TNF α	7	5	1	2	2.4	5.7	97.5 (IL1 β)	358 (TNF α)	138

*n.m.= not measurable by Biacore.
1 - Heavy chain and light chains corresponding to codes can be found in Table 10.

Example 9. Retention of Parental Antibody Affinity in CODV-Ig Format

[0161] The identical antibody sequences for anti-IL4 and anti-IL13 were incorporated into either the TBTI/DVD-Ig or CODV-Ig formats for a direct comparison of these configurations, the positioning of the linkers, and affinities of the resulting molecules. As shown in Figure 7, the parental affinity of each of the antibodies was maintained in the CODV format. As shown in the upper panel of Figure 7, when the variable regions were placed in the TBTI/DVD-Ig format, a drop in affinity of the IL4 antibody positioned at the inner Fv2 position was manifested as a reduction of the on-rate of antibody binding to the antigen. In contrast, there was no loss in affinity for the CODV-Ig format as compared to the parental antibodies (see Figure 7, lower panel).

Example 10. Adaptability of CODV-Ig to Fab Format

[0162] The ability of the CODV-Ig format to provide fragments such as Fab fragments was evaluated next. Two different variable heavy chains were fused to each other through linker L_3 and elongated C-terminally by linker L_4 . This V_H complex was then fused to the C_{H1} domain of IGHG1 (GenBank Accession No. Q569F4) harboring C-terminally the five amino acid sequence DKTHT (SEQ ID NO: 60) from the hinge region followed by six histidine residues. Two different variable light chains were fused to each other in a cross-over configuration to the corresponding heavy chain through linker L_1 and extended C-terminally by linker L_2 and subsequently fused to the constant kappa chain (IGKC, GenBank Accession No. Q502W4).

[0163] Fab fragments were expressed by transient transfection as described previously. Seven days post-transfection, cells were removed by centrifugation, 10 % vol/vol 1M Tris-HCl, pH 8.0, was added and the supernatant was passed over a 0.22 μ m filter to remove particles. The Fab proteins were captured using HisTrap High Performance columns (GE Healthcare) and eluted via imidazole gradient. The protein containing fractions were pooled and desalted using PD-10 or Sephadex columns. Concentrated and sterile filtered (0.22 μ m) protein solutions were adjusted to 1 mg/ml and kept at 4°C until use.

[0164] Immediate advantages were observed in that the Fab-like molecules in a CODV orientation showed no tendency to aggregate and retained the affinities of the parental antibodies (see Table 12). Binding protein constructs from Batch ID Nos. 401-421 directly compared antibody-like proteins in which antibody variable regions were arranged as in CODV-Ig molecules with the heavy chain as the template (401, 402, 406, and 407), CODV Fab-like fragments (402, 408, 413, 418, and 421), four domain antibody-like molecules in TBTI/DVD-Ig format (404, 409, 414, and 419), and CODV-Ig with no linkers (405, 410, 415, and 420). As shown in Table 12, the results of this comparison indicated that there is more likely to be a loss in affinity as compared to the parent antibodies when the variable region is incorporated into a TBTI or DVD-Ig format. In contrast, both the CODV-Ig and CODV-Ig Fab-like formats were better able to maintain parental affinities. The results further confirmed that CODV-Ig molecules require linkers between the variable regions and between the variable regions and the constant domains (see Table 12).

Table 12. Use of CODV-Ig Format for Fab-like Fragments

Batch ID	Sample-ID	Format	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen 1) [pM]	KD (Antigen 2) [pM]	IC50 [pM] TNF α Cellular Assay
401	TNF α x IL12/23	CODV-Ig	7	5	1	2	7.1	7.5	321 (TNF α)	90 (IL23)	95
402	IL12/23 x TNF α	CODV-Ig	7	5	1	2	11.9	7.1	118 (IL23)	543 (TNF α)	138
403	TNF α x IL12/23	CODV-Fab	7	5	1	2	18.7	1.7	232 (TNF α)	41 (IL23)	785
404	TNF α x IL12/23	TBTI	(G4S)2	0	(G4S)2	0	26.0	8.7	219 (TNF α)	399 (IL23)	-
405	TNF α x IL12/23	CODV-Ig	0	0	0	0	3.5	71	-	-	-
406	IL1 β x TNF α	CODV-Ig	7	5	1	2	2.4	5.7	98 (IL1 β)	358 (TNF α)	139
407	TNF α x IL1 β	CODV-Ig	7	5	1	2	6.6	13.6	340 (TNF α)	155 (IL1 β)	122
408	IL1 β x TNF α	CODV-Fab	7	5	1	2	8.6	0	179 (IL1 β)	-	-
409	IL1 β x TNF α	TBTI	(G4S)2	0	(G4S)2	0	1.3	40.5	133 (IL1 β)	456 (TNF α)	-
410	TNF α x IL1 β	CODV-Ig	0	0	0	0	n.p.	-	-	-	-
411	EGFR x IGF1R(2)	CODV-Ig	7	5	1	2	9.5	4.3	124nM (EGFR)	n.m.	-
412	IGF1R(2) x EGFR	CODV-Ig	7	5	1	2	17	2.7	n.m.	-	-
413	EGFR x IGF1R(2)	CODV-Fab	7	5	1	2	13.3	0	42nM (EGFR)	n.m.	-
414	EGFR x IGF1R(2)	TBTI	(G4S)2	0	(G4S)2	0	2.1	2.9	7nM (EGFR)	n.m.	-

(continued)

Batch ID	Sample-ID	Format	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen 1) [pM]	KD (Antigen 2) [pM]	IC50 [pM] TNF α Cellular Assay
415	EGFR x IGF1R(2)	CODV-Ig	0	0	0	0	4.4	100	-	-	-
416	HER2 x IGF1R(1)	CODV-Ig	7	5	1	2	60.0	1.8	163 (HER2)	n.m.	-
417	IGF1R(1) x HER2	CODV-Ig	7	5	1	2	70	5.9	n.m.	41 (HER2)	-
418	HER2 x IGF1R(1)	CODV-Fab	7	5	1	2	34.4	0	190 (HER2)	n.m.	-
419	HER2 x IGF1R(1)	TBTI	(G4S)2	0	(G4S)2	0	6.3	6.8	56 (HER2)	n.m.	-
420	IGF1R(1) x HER2	CODV-Ig	0	0	0	0	0.35	54.5	-	-	-
421	IL4 x IL13	CODV-Fab	7	5	1	2	8.7	14	12 (IL4)	48 (IL13)	-

Example 11. Substitution of Variable Domains within CODV-Ig and CODV-Fab

[0165] To characterize the CODV format in a T-cell engaging approach, bispecific CODV Fab-like binding proteins (CODV-Fab) having a TCR binding site (CD3epsilon) and a CD19 binding site were generated and compared to a bispecific Fab derived from the TBTI/DVD-Ig format (B-Fab). To investigate the importance of the orientation of the binding sites (TCR x CD19 vs. CD19 x TCR), both orientations were evaluated for each of the binding proteins.

[0166] The binding proteins were characterized in a cytotoxic assay using NALM-6 (CD 19 expressing) cells as target cells and primary human T-cells as effector cells. CD3 positive cells were isolated from freshly prepared human PBMC's. Effector and target cells were mixed at a ratio of 10:1 and incubated for 20 hours with the indicated concentrations of bispecific binding proteins (see Figure 8). Apoptotic target cells were determined in a FACS-based assay using 7-Aminoactinomycin staining.

[0167] The B-Fab format in the configuration CD3-CD19 (1060) was shown to be active in inducing T-cell mediated cytotoxicity towards NALM-6 cells with an EC50 of 3.7 ng/ml. A similarly high activity was observed for the CD19-CD3 CODV-Fab (1109) with an EC50 of 3.2 ng/ml (see Figure 8).

[0168] A swap of the configuration of the B-Fab molecule (Fab of the TBTI / DVD-Ig format) to a CD19-CD3 orientation resulted in a significant loss of activity (see Figure 8). The swapped B-Fab molecule showed no activity at concentrations that were maximal for both orientations of CODV-Ig Fab and the other orientation of B-Fab. For the CODV-Ig Fabs and one orientation of B-Fab, a maximum response was observed (ranging between 1 and 100 ng/ml). For the CD19-CD3 orientation of B-Fab, even at the maximal concentration (30 µg/ml), the optimal cytotoxic response was not reached. In sharp contrast, a change in the orientation of the domains in the CODV-Fab towards CD3-CD19 (1108), resulted in a molecule with significant activity in this assay (see Figure 8). Although domain swapping in the CODV-Fab also reduced the induction of T-cell mediated cytotoxicity (increase of EC50 by factor ~100), this effect was far less pronounced than was seen in the B-Fab format and the molecule was able to induce cytotoxicity up to the maximal level. The data were representative and obtained from three independent experiments.

Example 12. Influence of Amino Acid Sequence Identity on CODV-Ig Linkers

[0169] The optimized construct corresponding to Batch ID 204 (see Example 7 and Table 8) was chosen to investigate the influence of linker composition on the linkers L₁ to L₄. Linker lengths were set at 7, 5, 1, and 2 residues in length for L₁, L₂, L₃, and L₄, respectively (see Table 13). Test sequences were derived from naturally occurring linkers at the transitions between natural antibody V_H and C_{H1} domains or between antibody Fv and C_L domains of kappa or lambda light chains. The candidate sequences were ASTKGPS (SEQ ID NO: 48), which is derived from the V_H and C_{H1} domain transition, RTVAAPS (SEQ ID NO: 49) and GQPKAAP (SEQ ID NO: 50), which were derived from the Fv and C_L domain transitions of kappa and lambda light chains, respectively. Furthermore, one construct was generated with an arbitrary linker composition to show that any sequence can be potentially used in linkers L₁ to L₄. This linker composition was obtained by randomly distributing the amino acids valine, leucine, isoleucine, serine, threonine, lysine, arginine, histidine, aspartate, glutamate, asparagine, glutamine, glycine, and proline at the 15 positions of the four linkers. The aromatic amino acids phenylalanine, tyrosine, and tryptophan, as well as the amino acids methionine and cysteine were deliberately excluded to avoid potential increases in aggregation.

[0170] A three-dimensional model of the construct for Batch ID No. 204 was generated to assure suitability or refine the choices of linker composition. Thus, serine was chosen for linker L₃ as positively and negatively charged residues are observed nearby in the three-dimensional model. The residues in linker L₄ were selected to be compatible with solvent exposure of these positions as suggested by the model. Similarly, no problems were anticipated or predicted for the linker compositions of L₁ and L₂. Three-dimensional models of selected proposals for linker composition were constructed.

[0171] As shown in Table 12, linker composition may have a dramatic influence on yield. Sequences that were derived from lambda chain on L₁ (comparing Batch ID Nos. 505-507 with Batch ID Nos. 501-503) were more productive protein generators (up to 8 fold increase). Indeed, the linkers based on random generation also produced good yields, as shown in Table 13, Batch ID No. 508. Therefore, linker composition should be one parameter considered during CODV-Ig optimization.

Table 13. Effect of Linker Composition on CODV-Ig

Batch ID	Alignment on HC	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (IL4) [pM]	KD (IL13) [pM]	IC50 Cellular Assay IL4 [nM]	IC50 Cellular Assay IL13 [nM]
501	IL4 x IL13	ASTKGPS (SEQ ID NO: 48)	TKGPS (SEQ ID NO: 52)	S	RT	10.8	1.2	1	55	0.034	4.5
502	IL4 x IL13	ASTKGPS (SEQ ID NO: 48)	TVAAP (SEQ ID NO: 53)	S	QP	15.8	2.9	3	61	0.049	2.4
503	IL4 x IL13	ASTKGPS (SEQ ID NO: 48)	TVAAP (SEQ ID NO: 53)	S	SS	11.6	3.5	3	52	0.047	2.1
504	IL4 x IL13	RTVAAPS (SEQ ID NO: 49)	QPKAA (SEQ ID NO: 54)	S	TK	15	1.9	8	71	0.042	1.4
505	IL4 x IL13	GQPKAAP (SEQ ID NO: 50)	TKGPS (SEQ ID NO: 52)	S	RT	71.7	1.9	6	68	0.033	0.9
506	IL4 x IL13	GQPKAAP (SEQ ID NO: 50)	TVAAP (SEQ ID NO: 53)	S	TK	49.3	1.7	7	55	0.045	1.8
507	IL4 x IL13	GQPKAAP (SEQ ID NO: 50)	QPKAA (SEQ ID NO: 54)	S	RT	62.4	2	1	69	0.040	2.0
508	IL4 x IL13	HIDSPNK (SEQ ID NO: 51)	QRIEG (SEQ ID NO: 55)	V	SL	37.7	2.1	1	44	0.054	1.3

[0172] Activities of bispecific antibodies or derivatives against cytokines IL4 and IL13 were determined in commercially available HEK-Blue IL-4/IL-13 reporter cells (InvivoGen). HEK-Blue IL-4/IL-13 cells are designed to monitor the activation of the STAT6 pathway by IL-4 or IL13. Stimulation of the cells with either cytokine results in production of the reporter gene secreted embryonic alkaline phosphatase (SEAP), which can be measured in the culture supernatant with the QUANTI-Blue assay. To test antibody activities against IL4 or IL13, the cytokines were pre-incubated for 1 hour with different concentrations of the antibodies and added to 50,000 HEK-Blue IL-4/IL-13 cells. Cytokine-mediated induction of SEAP was measured after 24 hours incubation in the cell culture supernatant with the QUANTI-Blue assay (InvivoGen).

Example 13. Introduction of Cysteines into CODV-Ig Linkers

[0173] Published data suggest that the stability of antibodies and antibody-derived proteins can be increased by the introduction of non-natural disulfide bridges (see Wozniak-Knopp et al., 2012, "Stabilisation of the Fc Fragment of Human IgG1 by Engineered Intradomain Disulfide Bonds," PLoS ONE 7(1): e30083). To examine whether the equivalent Fc fragment derived from a human IgG1 antibody and engineered into a CODV-Ig molecule can be stabilized by the introduction of inter- and intra-chain disulfide bridges, the equivalent Fc positions of the CODV-Ig construct Batch ID No. 204 (from Example 7) were mutated to cysteine residues, and the mutant proteins were overproduced, purified, and characterized (see Table 14).

[0174] As shown in Table 14, each of the mutated CODV-Ig molecules containing additional cysteine residues had melting temperatures that were the same as the melting temperature for the CODV-Ig construct Batch ID No. 204.

[0175] In addition, two simultaneous cysteines were introduced at Kabat positions 100 for the light chain and 44 for the heavy chain on each of the variable domains as described in Brinkmann et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90: 7538-42. These positions have been shown to be structurally conserved within antibody folds, and therefore tolerable of cysteine substitution without interfering with the integrity of the individual domains.

[0176] As shown in Table 15, CODV and CODV-Ig constructs in which cysteine residues were introduced at Kabat positions 100 for the light chain and 44 for the heavy chain on each of the variable domains had higher melting temperatures than the CODV and CODV-Ig constructs in which cysteine residues were not introduced at these positions (see, e.g., Batch ID Nos. 704 and 706 and Batch ID Nos. 713 and 714).

1. Thermostability Measurements of CODV and TBTI Variants

[0177] Melting points (T_m) of CODV and TBTI variants were determined using differential scanning fluorimetry (DSF). Samples were diluted in D-PBS buffer (Invitrogen) to a final concentration of 0.2 $\mu\text{g}/\mu\text{l}$ and added to 2 μl of a 40x concentrated solution of SYPRO-Orange dye (Invitrogen) in D-PBS in white semi-skirt 96-well plates. All measurements were done in duplicate using a MyiQ2 real time PCR instrument (Biorad). T_m values were extracted from the negative first derivative of the melting curves using iQ5 Software v2.1.

[0178] Next the effect of introducing cysteine residues directly into the linkers or within the variable region was examined. In this example, Batch ID No. 204 (from Example 7 and Table 8) was used as the model CODV-Ig binding protein, and cysteine was substituted for glycine in L_1 , L_3 , or the variable region on the basis of the three-dimensional model. As shown in Table 16 below, the results indicate how the introduction of cysteine pairs would affect yield and aggregation. The envisioned mutations were all modeled to ascertain that disulfide bonds were properly formed and correct geometry was maintained for the linkers and their environment in the models. Nevertheless, Batch ID No. 808 showed good yield and little aggregation, thus suggesting that a proper cysteine bridge could be formed.

[0179] Each embodiment herein described may be combined with any other embodiment or embodiments unless clearly indicated to the contrary. In particular, any feature or embodiment indicated as being preferred or advantageous may be combined with any other feature or features or embodiment or embodiments indicated as being preferred or advantageous, unless clearly indicated to the contrary.

Table 14. Effect of Disulfide Bridge Stabilization on CODV-Ig

Batch ID	Alignment on HC	Introduced Mutations IGHG1	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (IL4) [pM]	KD (IL13) [pM]	IC50 Cellular Assay IL4 [nM]	IC50 Cellular Assay IL13 [nM]	T _m [°C]
601	IL4 x IL13	P344C & A432C	7	5	1	2	21.6	0.9	n.d.	n.d.	0.032	1.7	64
602	IL4 x IL13	S376C & P397C	7	5	1	2	13.9	0.8	10	100	0.034	1.5	64
603	IL4 x IL13	P446G447 to G446E447C448	7	5	1	2	15.7	1.3	4	64	0.036	1.4	64

Table 15. Effect of Disulfide Bridge Stabilization on CODV-Ig

Batch ID	Alignment on HC	Format	Introduced Mutations LC	Introduced Mutations HC	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (Antigen 1) [pM]	KD (Antigen 2) [pM]	T _m [°C]
704	IL13 x IL4	CODV	-	-	7	5	1	2	15.3	5.2	3	25	64
706	IL4 x IL13	CODV	G100C & G100C	G44C & G44C	7	5	1	2	3.2	17.7	8	61	68
713	IGF1R(1) x HER2	CODV-Ig	-	-	7	5	1	2	60	1.8	163 (HER2)	-	64
714	IGF1R(1) x HER2	CODV-Ig	G100C & Q100C	G44C & G44C	7	5	1	2	3.4	7.9	362 (HER22)	-	68

Table 16A. Effect of Disulfide Bridge Stabilization on CODV-Ig

Batch ID	Alignment on HC	Introduced Mutations LC	Introduced Mutations HC	L ₁	L ₂	L ₃	L ₄	Yield [mg/L]	Aggregation [%]	KD (IL4) [pM]	KD (IL13) [pM]	T _m
801	IL4 x IL13			7	5 GGGGG (SEQ ID NO: 56)	1	2	15.3	5.2	3	25	64
802	IL4 x IL13	G 114C	Q43C	GGCGGG (SEQ ID NO: 56)	GGGGG (SEQ ID NO: 26)	G	GG	3.0	11.3	6	34	64
803	IL4 x IL13	G115C	Q43C	GGCGGG (SEQ ID NO: 57)	GGGGG (SEQ ID NO: 26)	G	GG	4.0	9.4	6	20	63
804	IL4 x IL13	G116C	Q43C	GGGGCGG (SEQ ID NO: 58)	GGGGG (SEQ NO: 26)	G	GG	11.1	9.6	2	33	64
805	IL4 x IL13	G117C	Q43C	GGGGCGG (SEQ ID NO: 59)	GGGGG (SEQ ID NO: 26)	G	GG	5.8	10.6	7	45	64
806	IL4 x IL13	S60C	S122C	GGGGGGG (SEQ ID NO: 28)	GGGGG (SEQ ID NO: 26)	G	GG	n.p.	-	-	-	-
807	IL4 x IL13	S60C	S123C	GGGGGGG (SEQ ID NO: 28)	GGGGG (SEQ NO: 26)	G	GG	7.5	27.9	2	202	66
808	IL4 x IL13	S60C	G124C	GGGGGGG (SEQ ID NO: 28)	GGGGG (SEQ NO: 26)	C	GG	1.1	5.8	5	70	64
809	IL4 x IL13	G61C	S122C	GGGGGGG (SEQ ID NO: 28)	GGGGG (SEQ ID NO: 26)	G	GG	8.7	6.9	6	81	65

Table 16B. Effect of Disulfide Bridge Stabilization on CODV-Ig

Batch ID	Alignment on HC	Introduced Mutations LC	Introduced Mutations HC	L ₁	L ₂	L ₃	L ₄	IC50 Cellular Assay IL4 [nM]	IC50 Cellular Assay IL13 [nM]
801	IL4 x IL13			7	5	1	2	0.045	1.8
802	IL4 x IL13	G114C	Q43C	GGCGGG (SEQ ID NO: 56)	GGGG (SEQ ID NO: 26)	G	GG	0.027	1.2
803	IL4 x IL13	G115C	Q43C	GGCGGG (SEQ ID NO: 57)	GGGG (SEQ ID NO: 26)	G	GG	0.036	1.5
804	IL4 x IL13	G116C	Q43C	GGGCGG (SEQ ID NO: 58)	GGGG (SEQ ID NO: 26)	G	GG	0.040	1.8
805	IL4 x IL13	G117C	Q43C	GGGCGG (SEQ ID NO: 59)	GGGG (SEQ ID NO: 26)	G	GG	0.047	1.9
806	IL4 x IL13	S60C	S122C	GGGGGG (SEQ ID NO: 28)	GGGG (SEQ ID NO: 26)	G	GG	-	-
807	IL4 x IL13	S60C	S123C	GGGGGG (SEQ ID NO: 28)	GGGG (SEQ ID NO: 26)	G	GG	0.048	10.1
808	IL4 x IL13	S60C	G124C	GGGGGG (SEQ ID NO: 28)	GGGG (SEQ ID NO: 26)	C	GG	0.023	3.8
809	IL4 x IL13	G61C	S122C	GGGGGG (SEQ ID NO: 28)	GGGG (SEQ ID NO: 26)	G	GG	0.035	2.8

SEQUENCE LISTING

[0180]

5 <110> Rao, Ercole
 Mikol, Vincent
 Corvey, Carsten
 Beil, Christian
 Lange, Christian
 10 Steinmetz, Anke
 Baurin, Nicolas
 Li, Danxi

 <120> Dual Variable Region Antibody-Like Binding Proteins Having Cross-over Binding Region Orientation

15 <130> US2011/026

 <140> TBD
 <141> 2011-03-11

20 <160> 59

 <170> PatentIn version 3.5

25 <210> 1
 <211> 108
 <212> PRT
 <213> Artificial Sequence

30 <220>
 <223> Humanized antibody light chain (VL) from anti-IL4

 <400> 1

35

40

45

50

55

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly
	1				5					10				15		
5	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp
				20					25					30		
10	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile
			35					40					45			
15	Tyr	Lys	Ala	Ser	Asn	Leu	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
		50					55					60				
20	Ser	Gly	Ser	Gly	Thr	Gly	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
	65					70					75				80	
25	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ala	His	Ser	Tyr	Pro	Phe
					85					90					95	
30	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg				
				100					105							

<210> 2

<211> 124

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized heavy chain V region (VH) from anti-IL4 antibody

<400> 2

EP 2 691 416 B1

1 Gln Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 5 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Ser Tyr
 10 Trp Ile His Trp Ile Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 15 Gly Met Ile Asp Pro Ser Asp Gly Glu Thr Arg Leu Asn Gln Arg Phe
 20 Gln Gly Arg Ala Thr Leu Thr Val Asp Glu Ser Thr Ser Thr Ala Tyr
 25 Met Gln Leu Arg Ser Pro Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 30 Thr Arg Leu Lys Glu Tyr Gly Asn Tyr Asp Ser Phe Tyr Phe Asp Val
 35 Trp Gly Ala Gly Thr Leu Val Thr Val Ser Ser Ala
 40 <210> 3
 <211> 111
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Humanized light chain V region (VL) from an anti-IL13 antibody
 50 <400> 3
 55 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 60 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr
 65 Gly Gln Ser Tyr Met His Trp Tyr Gln Gln Lys Ala Gly Gln Pro Pro
 70 Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala

EP 2 691 416 B1

	50	55	60
5	Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp 65 70 75 80		
10	Pro Val Gln Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asn Ala 85 90 95		
15	Glu Asp Ser Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys 100 105 110		
20	<210> 4 <211> 118 <212> PRT <213> Artificial Sequence		
25	<220> <223> Humanized heavy chain V region (VH) from an anti-IL-13 antibody. <400> 4		
30	Glu Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Gly Gly 1 5 10 15		
35	Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Asp Ser 20 25 30		
40	Ser Ile Asn Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu 35 40 45		
45	Gly Met Ile Trp Gly Asp Gly Arg Ile Asp Tyr Ala Asp Ala Leu Lys 50 55 60		
50	Ser Arg Leu Ser Ile Ser Lys Asp Ser Ser Lys Ser Gln Val Phe Leu 65 70 75 80		
55	Glu Met Thr Ser Leu Arg Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala 85 90 95		
	Arg Asp Gly Tyr Phe Pro Tyr Ala Met Asp Phe Trp Gly Gln Gly Thr 100 105 110		
	Ser Val Thr Val Ser Ser 115		
	<210> 5 <211> 575 <212> PRT <213> Artificial Sequence		

EP 2 691 416 B1

<220>

<223> Heavy chain of cross over double head construct IL13(G4S)IL4CHFc

<400> 5

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser	
				20					25					30			
10	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
			35					40					45				
15	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
		50					55					60					
20	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
	65					70					75					80	
25	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
					85					90					95		
30	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
35	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	Gln	Val	Gln	Leu	Gln	
				115				120					125				
40	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	
		130					135					140					
45	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	Ile	
	145					150					155					160	
50	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	Pro	
				165						170					175		
55	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	Thr	
				180					185					190			
60	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	Ser	
			195					200					205				
65	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	Glu	
		210					215					220					
70	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	Thr	
	225					230					235					240	

EP 2 691 416 B1

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
 245 250 255

5 Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
 260 265 270

10 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 275 280 285

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 290 295 300

15 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 305 310 315 320

20 Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
 325 330 335

25 Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr
 340 345 350

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
 355 360 365

30 Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 370 375 380

35 Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
 385 390 395 400

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 405 410 415

40 Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val
 420 425 430

45 Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 435 440 445

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
 450 455 460

50 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 465 470 475 480

55 Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
 485 490 495

EP 2 691 416 B1

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
500 505 510

5 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
515 520 525

10 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
530 535 540

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
545 550 555 560

15 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
565 570 575

<210> 6
20 <211> 580
<212> PRT
<213> Artificial Sequence

<220>
25 <223> heavy chain of cross over double head construct IL13(G4S2)IL4CHFc

<400> 6

30 Glu Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Gly Gly
1 5 10 15

Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Asp Ser
35 20 25 30

Ser Ile Asn Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
35 40 45

40 Gly Met Ile Trp Gly Asp Gly Arg Ile Asp Tyr Ala Asp Ala Leu Lys
50 55 60

45 Ser Arg Leu Ser Ile Ser Lys Asp Ser Ser Lys Ser Gln Val Phe Leu
65 70 75 80

Glu Met Thr Ser Leu Arg Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala
50 85 90 95

Arg Asp Gly Tyr Phe Pro Tyr Ala Met Asp Phe Trp Gly Gln Gly Thr
100 105 110

55 Ser Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
115 120 125

EP 2 691 416 B1

	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	
	130						135					140					
5	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	
	145					150					155					160	
	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	
10					165					170					175		
	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	
				180					185					190			
15	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	
			195					200					205				
	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	
20		210					215					220					
	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	
	225					230					235					240	
25	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	
					245					250					255		
30	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	
				260					265					270			
	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	
35			275					280					285				
	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	
		290					295					300					
40	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	
	305					310					315					320	
	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	
45					325					330					335		
	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	
50				340					345					350			
	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	
			355					360					365				
55	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	
	370						375					380					

EP 2 691 416 B1

	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	
	385					390					395					400	
5	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	
					405					410					415		
10	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	
				420					425					430			
15	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	
			435					440					445				
20	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	
	450						455					460					
25	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	
	465					470					475					480	
30	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	
					485					490					495		
35	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	
				500					505					510			
40	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	
			515					520					525				
45	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	
	530						535					540					
50	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	
	545					550					555					560	
55	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	
					565					570					575		
60	Leu	Ser	Pro	Gly													
				580													
65	<210>	7															
70	<211>	698															
75	<212>	PRT															
80	<213>	Artificial Sequence															
85	<220>																
90	<223>	Heavy chain cross over double head construct IL4(G4S)IL13CHFc															
95	<400>	7															

EP 2 691 416 B1

	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	
	1				5					10					15		
5	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	
				20					25					30			
	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	
10			35					40					45				
	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	
		50					55					60					
15	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	
	65					70					75					80	
	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	
20					85					90					95		
	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	
25				100					105					110			
	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	
			115					120					125				
30	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly	
		130					135					140					
	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser	
35	145					150					155					160	
	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
				165						170					175		
40	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
				180					185					190			
	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
45			195					200					205				
	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
50		210					215					220					
	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
	225					230					235					240	
55	Ser	Val	Thr	Val	Ser	Ser	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	
					245					250					255		

EP 2 691 416 B1

	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	
				260					265					270			
5	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	Pro	Gly	
			275					280					285				
10	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly	Glu	Thr	
		290					295					300					
15	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val	Asp	Glu	
	305					310					315					320	
20	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser	Glu	Asp	
					325					330					335		
25	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn	Tyr	Asp	
				340					345					350			
30	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr	Val	Ser	
			355					360					365				
35	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	
		370					375					380					
40	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	
	385					390					395					400	
45	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	
					405					410						415	
50	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	
				420					425					430			
55	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	
			435					440					445				
60	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	
	450						455					460					
65	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	
	465					470					475					480	
70	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	
					485					490					495		
75	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	

EP 2 691 416 B1

	500	505	510
5	Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn 515 520 525		
10	Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg 530 535 540		
15	Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val 545 550 555 560		
20	Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser 565 570 575		
25	Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys 580 585 590		
30	Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp 595 600 605		
35	Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe 610 615 620		
40	Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu 625 630 635 640		
45	Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe 645 650 655		
50	Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly 660 665 670		
55	Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr 675 680 685		
	Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly 690 695		

<210> 8

<211> 703

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain cross over double head construct IL4(G4S2)IL13CHFc

<400> 8

EP 2 691 416 B1

Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala
1				5					10					15	

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	
				20					25					30			
5	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	
			35					40					45				
10	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	
		50					55					60					
15	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	
	65					70					75					80	
20	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
25	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	
				100					105					110			
30	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ser	
			115					120					125				
35	Gly	Gly	Gly	Gly	Ser	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	
		130					135					140					
40	Val	Ala	Pro	Gly	Gly	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	
	145					150					155					160	
45	Ser	Leu	Thr	Asp	Ser	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	
					165					170					175		
50	Gly	Leu	Glu	Trp	Leu	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	
				180					185					190			
55	Ala	Asp	Ala	Leu	Lys	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	
			195					200					205				
60	Ser	Gln	Val	Phe	Leu	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	
		210					215					220					
65	Thr	Tyr	Tyr	Cys	Ala	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	
	225					230					235					240	
70	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Gln	Val	Gln	Leu	Gln	
					245					250					255		
75	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	

EP 2 691 416 B1

	260	265	270
5	Cys Lys Ala Ser Gly Tyr Ser Phe Thr Ser Tyr Trp Ile His Trp Ile 275 280 285		
10	Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Met Ile Asp Pro 290 295 300		
15	Ser Asp Gly Glu Thr Arg Leu Asn Gln Arg Phe Gln Gly Arg Ala Thr 305 310 315 320		
20	Leu Thr Val Asp Glu Ser Thr Ser Thr Ala Tyr Met Gln Leu Arg Ser 325 330 335		
25	Pro Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Thr Arg Leu Lys Glu 340 345 350		
30	Tyr Gly Asn Tyr Asp Ser Phe Tyr Phe Asp Val Trp Gly Ala Gly Thr 355 360 365		
35	Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro 370 375 380		
40	Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly 385 390 395 400		
45	Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn 405 410 415		
50	Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln 420 425 430		
55	Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser 435 440 445		
	Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser 450 455 460		
	Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr 465 470 475 480		
	His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser 485 490 495		
	Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg 500 505 510		

EP 2 691 416 B1

	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	
			515					520					525				
5	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	
		530					535					540					
10	Lys	Thr	Lys	Pro	Arg	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val		
	545					550				555					560		
	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	
					565					570					575		
15	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	
				580					585					590			
20	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	
			595					600					605				
25	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	
		610					615					620					
	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	
	625					630				635						640	
30	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	
					645					650					655		
35	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	
				660					665					670			
40	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	
			675					680					685				
	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
		690					695					700					
45	<210> 9																
	<211> 330																
	<212> PRT																
	<213> Artificial Sequence																
50	<220>																
	<223> Light chain cross over double head construct IL13(G4S)IL4CL																
	<400> 9																
55																	

EP 2 691 416 B1

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1 5 10 15

5 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	20	25	30
5	Gly Gln Ser Tyr Met His Trp Tyr Gln Gln Lys Ala Gly Gln Pro Pro 35 40 45		
10	Lys Leu Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala 50 55 60		
15	Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp 65 70 75 80		
20	Pro Val Gln Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asn Ala 85 90 95		
25	Glu Asp Ser Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly 100 105 110		
30	Gly Gly Gly Ser Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser 115 120 125		
35	Val Ser Val Gly Asp Thr Ile Thr Leu Thr Cys His Ala Ser Gln Asn 130 135 140		
40	Ile Asp Val Trp Leu Ser Trp Phe Gln Gln Lys Pro Gly Asn Ile Pro 145 150 155 160		
45	Lys Leu Leu Ile Tyr Lys Ala Ser Asn Leu His Thr Gly Val Pro Ser 165 170 175		
50	Arg Phe Ser Gly Ser Gly Ser Gly Thr Gly Phe Thr Leu Thr Ile Ser 180 185 190		
55	Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Ala His 195 200 205		
	Ser Tyr Pro Phe Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg 210 215 220		
	Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln 225 230 235 240		
	Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr 245 250 255		
	Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser 260 265 270		

EP 2 691 416 B1

	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr
			275					280					285			
5	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys
		290					295					300				
10	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro
	305					310					315					320
	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys						
					325					330						
15																
	<210> 10															
	<211> 335															
	<212> PRT															
	<213> Artificial Sequence															
20																
	<220>															
	<223> Light chain cross over double head construct IL13(G4S2)IL4CL															
	<400> 10															
25																
30																
35																
40																
45																
50																
55																

EP 2 691 416 B1

	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ala	Val	Ser	Leu	Gly
	1				5					10					15	
5	Gln	Arg	Ala	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Glu	Ser	Val	Asp	Ser	Tyr
				20					25					30		
	Gly	Gln	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Ala	Gly	Gln	Pro	Pro
10			35					40					45			
	Lys	Leu	Leu	Ile	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Ala
		50					55					60				
15	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Arg	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Asp
	65					70					75					80
	Pro	Val	Gln	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Asn	Ala
20					85					90					95	
	Glu	Asp	Ser	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly
25				100					105					110		
	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile	Gln	Met	Thr	Gln	Ser
			115					120					125			
30	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly	Asp	Thr	Ile	Thr	Leu	Thr	Cys
		130					135						140			
35	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp	Leu	Ser	Trp	Phe	Gln	Gln	Lys
40																
45																
50																
55																

EP 2 691 416 B1

	145		150		155		160
5	Pro Gly Asn Ile	Pro Lys Leu Leu Ile	Tyr Lys Ala Ser Asn Leu His				
		165	170	175			
10	Thr Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Gly Phe						
		180	185	190			
15	Thr Leu Thr Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr						
		195	200	205			
20	Cys Gln Gln Ala His Ser Tyr Pro Phe Thr Phe Gly Gly Gly Thr Lys						
		210	215	220			
25	Leu Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro						
		225	230	235			240
30	Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu						
		245	250	255			
35	Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp						
		260	265	270			
40	Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp						
		275	280	285			
45	Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys						
		290	295	300			
50	Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln						
		305	310	315			320
55	Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys						
		325	330	335			
60	<210> 11						
	<211> 330						
	<212> PRT						
	<213> Artificial Sequence						
65	<220>						
	<223> Light chain cross over double head construct IL4(G4S)IL13CL						
70	<400> 11						

EP 2 691 416 B1

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly
1				5					10					15	

5	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp
			20						25					30		

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile	
			35					40					45				
5	Tyr	Lys	Ala	Ser	Asn	Leu	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
		50					55					60					
	Ser	Gly	Ser	Gly	Thr	Gly	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
10	65					70					75					80	
	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ala	His	Ser	Tyr	Pro	Phe	
					85					90					95		
15	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser	
				100					105					110			
	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ala	Val	Ser	Leu	Gly	
20			115					120					125				
	Gln	Arg	Ala	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Glu	Ser	Val	Asp	Ser	Tyr	
25		130					135					140					
	Gly	Gln	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Ala	Gly	Gln	Pro	Pro	
	145					150					155					160	
30	Lys	Leu	Leu	Ile	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Ala	
				165						170					175		
	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Arg	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Asp	
35				180					185					190			
	Pro	Val	Gln	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Asn	Ala	
			195					200					205				
40	Glu	Asp	Ser	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	
	210						215					220					
45	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	
	225					230					235					240	
	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	
50					245					250					255		
	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	
				260					265					270			
55	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	

EP 2 691 416 B1

	275	280	285
5	Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys 290 295 300		
10	His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro 305 310 315 320		
	Val Thr Lys Ser Phe Asn Arg Gly Glu Cys 325 330		
15	<210> 12 <211> 335 <212> PRT <213> Artificial Sequence		
20	<220> <223> Light chain cross over double head construct IL4(G4S2)IL13CL		
	<400> 12		
25			
30			
35			
40			
45			
50			
55			

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly
	1				5					10					15	
5	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp
				20					25					30		
10	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile
			35					40					45			
15	Tyr	Lys	Ala	Ser	Asn	Leu	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50						55					60				
20	Ser	Gly	Ser	Gly	Thr	Gly	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
	65					70					75				80	
25	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ala	His	Ser	Tyr	Pro	Phe
					85					90					95	
30	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Ser
				100					105					110		
35	Gly	Gly	Gly	Gly	Ser	Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ser	Leu
			115					120					125			
40	Ala	Val	Ser	Leu	Gly	Gln	Arg	Ala	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Glu
	130						135					140				
45	Ser	Val	Asp	Ser	Tyr	Gly	Gln	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys
	145					150					155					160

EP 2 691 416 B1

	Ala	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	
					165					170					175		
5	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Arg	Thr	Asp	Phe	
				180					185					190			
10	Thr	Leu	Thr	Ile	Asp	Pro	Val	Gln	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	
			195					200					205				
15	Cys	Gln	Gln	Asn	Ala	Glu	Asp	Ser	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	
		210					215					220					
20	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	
	225					230					235					240	
25	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	
					245					250					255		
30	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	
				260					265					270			
35	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	
			275				280						285				
40	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	
		290					295					300					
45	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	
	305					310					315					320	
50	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys		
					325					330					335		
	<210>	13															
	<211>	583															
	<212>	PRT															
	<213>	Artificial Sequence															
	<220>																
	<223>	Heavy chain sequence for cross over dual V-Ig having code HC1															
55	<400>	13															
	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly	
	1				5				10					15			
60	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser	
				20					25					30			

EP 2 691 416 B1

	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
			35					40					45				
5	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
		50					55					60					
	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
10		65				70					75					80	
	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
					85					90					95		
15	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gln	Val	Gln	Leu	
20				115				120					125				
	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	
25			130				135					140					
	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	
		145				150					155					160	
30	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	
					165					170					175		
	Pro	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	
35				180					185					190			
	Thr	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	
				195				200					205				
40	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	
			210				215					220					
	Glu	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	
45			225			230					235					240	
	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ala	Ser	
50					245					250					255		
	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	
				260					265					270			
55	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	
			275					280					285				

EP 2 691 416 B1

	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	
	290						295					300					
5	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	
	305					310					315					320	
	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	
10					325					330					335		
	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	
				340					345					350			
15	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	
			355					360					365				
	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	
20		370					375					380					
	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	
25		385				390					395					400	
	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	
					405					410					415		
30	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	
				420					425					430			
	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	
35			435					440					445				
	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	
40		450					455					460					
	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	
	465					470					475					480	
45	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	
					485					490					495		
	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	
50				500					505					510			
	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	
		515						520					525				
55	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	

EP 2 691 416 B1

530

535

540

5

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
545 550 555 560

10

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
565 570 575

Ser Leu Ser Leu Ser Pro Gly
580

15

<210> 14
<211> 580
<212> PRT
<213> Artificial Sequence

20

<220>
<223> Heavy chain sequence for cross over dual V-Ig having code HC2

<400> 14

25

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser	
				20					25					30			
10	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
			35					40					45				
15	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
		50					55					60					
20	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
	65					70					75					80	
25	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
					85					90					95		
30	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
			100						105					110			
35	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gln	Val	Gln	Leu	
			115					120					125				
40	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	
		130					135					140					
45	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	
	145					150					155					160	

EP 2 691 416 B1

	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	
					165					170					175		
5	Pro	Ser	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	
				180					185					190			
	Thr	Leu	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	
10				195				200					205				
	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	
		210					215					220					
15	Glu	Tyr	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	
	225					230					235					240	
	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ala	Ser	Thr	Lys	Gly	
20					245					250					255		
	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	
25				260					265					270			
	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	
			275					280					285				
30	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	
		290					295					300					
	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	
35	305					310					315					320	
	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	
					325					330					335		
40	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	
				340					345					350			
45	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	
			355					360					365				
	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	
50		370					375					380					
	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	
	385					390					395					400	
55	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	

EP 2 691 416 B1

	405	410	415
5	Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 420 425 430		
10	Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 435 440 445		
15	Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 450 455 460		
20	Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 465 470 475 480		
25	Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 485 490 495		
30	Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 500 505 510		
35	Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 515 520 525		
40	Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 530 535 540		
45	Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 545 550 555 560		
50	Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 565 570 575		
55	Leu Ser Pro Gly 580		

<210> 15

<211> 579

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain sequence for cross over dual V-Ig having code HC3

<400> 15

EP 2 691 416 B1

Glu Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Gly Gly
 1 5 10 15

5 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Asp Ser
 20 25 30

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
			35					40					45				
5	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
		50					55					60					
10	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
	65					70					75					80	
15	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
					85					90					95		
20	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
25	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	
			115					120					125				
30	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	
		130					135					140					
35	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	
	145					150					155					160	
40	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly	
					165					170					175		
45	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val	
				180					185					190			
50	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser	
			195					200					205				
55	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn	
		210					215					220					
60	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr	
	225					230					235					240	
65	Val	Ser	Ser	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	
				245					250						255		
70	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	
				260					265					270			
75	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	

EP 2 691 416 B1

	275	280	285
5	Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro 290 295 300		
10	Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr 305 310 315 320		
15	Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn 325 330 335		
20	His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser 340 345 350		
25	Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu 355 360 365		
30	Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu 370 375 380		
35	Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser 385 390 395 400		
40	His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu 405 410 415		
45	Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr 420 425 430		
50	Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn 435 440 445		
55	Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro 450 455 460		
	Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln 465 470 475 480		
	Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val 485 490 495		
	Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val 500 505 510		
	Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro 515 520 525		

EP 2 691 416 B1

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
530 535 540

5 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
545 550 555 560

10 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
565 570 575

Ser Pro Gly

15 <210> 16
<211> 576
<212> PRT
<213> Artificial Sequence

20 <220>
<223> Heavy chain sequence for cross over dual V-Ig having code HC4

<400> 16

25

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser	
				20					25					30			
	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu	
10			35					40					45				
	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys	
		50					55					60					
15	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu	
	65					70					75					80	
	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala	
20					85					90					95		
	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr	
25				100					105					110			
	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	
			115					120					125				
30	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	
		130					135					140					
35	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	Ile	Lys	Gln	Arg	
40																	
45																	
50																	
55																	

EP 2 691 416 B1

	145		150		155		160									
5	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	Pro	Ser	Asp	Gly
					165					170					175	
10	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	Thr	Leu	Thr	Val
				180					185					190		
15	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	Ser	Pro	Thr	Ser
			195					200					205			
20	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	Glu	Tyr	Gly	Asn
		210					215					220				
25	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	Thr	Leu	Val	Thr
	225					230					235					240
30	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
				245						250					255	
35	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
				260					265					270		
40	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
			275					280					285			
45	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
		290					295					300				
50	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
	305					310					315				320	
55	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
				325						330					335	
60	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
				340					345					350		
65	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro
			355					360					365			
70	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
		370					375					380				
75	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
	385					390					395					400

EP 2 691 416 B1

	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	
					405					410					415		
5	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	
				420					425					430			
	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	
10				435				440					445				
	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	
		450					455					460					
15																	
	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	
	465					470					475					480	
20	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	
					485					490					495		
25	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	
				500					505					510			
	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	
			515					520					525				
30																	
	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	
		530					535					540					
35	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	
	545					550					555					560	
	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	
40					565					570					575		

<210> 17

<211> 581

<212> PRT

45 <213> Artificial Sequence

<220>

<223> Heavy chain sequence for cross over dual V-Ig having code HC5

50 <400> 17

55

EP 2 691 416 B1

Glu Val Gln Leu Lys Glu Ser Gly Pro Gly Leu Val Ala Pro Gly Gly
 1 5 10 15

5 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Asp Ser
 20 25 30

10 Ser Ile Asn Trp Val Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	35	40	45
5	Gly Met Ile Trp Gly Asp Gly Arg Ile Asp Tyr Ala Asp Ala Leu Lys 50 55 60		
10	Ser Arg Leu Ser Ile Ser Lys Asp Ser Ser Lys Ser Gln Val Phe Leu 65 70 75 80		
15	Glu Met Thr Ser Leu Arg Thr Asp Asp Thr Ala Thr Tyr Tyr Cys Ala 85 90 95		
20	Arg Asp Gly Tyr Phe Pro Tyr Ala Met Asp Phe Trp Gly Gln Gly Thr 100 105 110		
25	Ser Val Thr Val Ser Ser Gly Gly Gly Gly Gln Val Gln Leu Gln Gln 115 120 125		
30	Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser Val Lys Ile Ser Cys 130 135 140		
35	Lys Ala Ser Gly Tyr Ser Phe Thr Ser Tyr Trp Ile His Trp Ile Lys 145 150 155 160		
40	Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Met Ile Asp Pro Ser 165 170 175		
45	Asp Gly Glu Thr Arg Leu Asn Gln Arg Phe Gln Gly Arg Ala Thr Leu 180 185 190		
50	Thr Val Asp Glu Ser Thr Ser Thr Ala Tyr Met Gln Leu Arg Ser Pro 195 200 205		
55	Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Thr Arg Leu Lys Glu Tyr 210 215 220		
	Gly Asn Tyr Asp Ser Phe Tyr Phe Asp Val Trp Gly Ala Gly Thr Leu 225 230 235 240		
	Val Thr Val Ser Ser Gly Gly Gly Gly Gly Gly Gly Ala Ser Thr Lys 245 250 255		
	Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly 260 265 270		
	Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro 275 280 285		

EP 2 691 416 B1

	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	
	290						295					300					
5	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	
	305					310					315					320	
	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	
10					325					330					335		
	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	
				340					345					350			
15	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	
			355					360					365				
	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	
20		370					375					380					
	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	
25	385					390					395					400	
	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	
					405					410					415		
30	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	
				420					425					430			
	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	
35			435					440					445				
	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	
		450					455					460					
40	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	
	465					470					475					480	
	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	
45					485					490					495		
	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	
50				500					505					510			
	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	
			515					520					525				
55	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	
	530						535					540					

EP 2 691 416 B1

	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
	545					550					555					560
5	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu
					565					570					575	
10	Ser	Leu	Ser	Pro	Gly											
					580											
	<210> 18															
	<211> 578															
	<212> PRT															
15	<213> Artificial Sequence															
	<220>															
	<223> Heavy chain for cross over dual V-Ig having code HC6															
20	<400> 18															
	Glu	Val	Gln	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Leu	Val	Ala	Pro	Gly	Gly
	1				5					10					15	
25	Ser	Leu	Ser	Ile	Thr	Cys	Thr	Val	Ser	Gly	Phe	Ser	Leu	Thr	Asp	Ser
				20					25					30		
30	Ser	Ile	Asn	Trp	Val	Arg	Gln	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Leu
			35					40					45			
35	Gly	Met	Ile	Trp	Gly	Asp	Gly	Arg	Ile	Asp	Tyr	Ala	Asp	Ala	Leu	Lys
	50						55					60				
40	Ser	Arg	Leu	Ser	Ile	Ser	Lys	Asp	Ser	Ser	Lys	Ser	Gln	Val	Phe	Leu
	65					70					75					80
45	Glu	Met	Thr	Ser	Leu	Arg	Thr	Asp	Asp	Thr	Ala	Thr	Tyr	Tyr	Cys	Ala
					85					90					95	
50	Arg	Asp	Gly	Tyr	Phe	Pro	Tyr	Ala	Met	Asp	Phe	Trp	Gly	Gln	Gly	Thr
				100					105					110		
55	Ser	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Gln	Val	Gln	Leu	Gln	Gln
			115					120					125			
60	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser	Val	Lys	Ile	Ser	Cys
	130						135					140				
65	Lys	Ala	Ser	Gly	Tyr	Ser	Phe	Thr	Ser	Tyr	Trp	Ile	His	Trp	Ile	Lys
	145					150					155					160

EP 2 691 416 B1

	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Met	Ile	Asp	Pro	Ser	
					165					170					175		
5	Asp	Gly	Glu	Thr	Arg	Leu	Asn	Gln	Arg	Phe	Gln	Gly	Arg	Ala	Thr	Leu	
				180					185					190			
10	Thr	Val	Asp	Glu	Ser	Thr	Ser	Thr	Ala	Tyr	Met	Gln	Leu	Arg	Ser	Pro	
			195					200					205				
	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Thr	Arg	Leu	Lys	Glu	Tyr	
		210					215					220					
15	Gly	Asn	Tyr	Asp	Ser	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Ala	Gly	Thr	Leu	
	225					230					235					240	
20	Val	Thr	Val	Ser	Ser	Gly	Gly	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	
					245					250					255		
25	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	
				260					265					270			
30	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	
			275					280					285				
35	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	
		290					295					300					
40	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	
	305					310					315					320	
45	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	
					325					330					335		
50	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	
				340					345					350			
55	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	
			355					360					365				
	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	
		370					375					380					
	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	
	385					390					395					400	
	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	
					405					410					415		

EP 2 691 416 B1

	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	
				420					425					430			
5	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	
			435					440					445				
10	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	
		450					455					460					
	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	
	465					470					475					480	
15	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	
					485					490					495		
20	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	
				500					505					510			
25	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	
			515					520					525				
	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	
	530						535					540					
30	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	
	545					550					555					560	
35	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	
					565					570					575		
40	Pro Gly																
	<210> 19																
	<211> 328																
	<212> PRT																
	<213> Artificial Sequence																
45	<220>																
	<223> Light chain sequence for cross over dual V-Ig having code LC1																
50	<400> 19																
	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly	
	1				5					10					15		
55	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp	
				20					25					30			

EP 2 691 416 B1

	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile	
			35					40					45				
5	Tyr	Lys	Ala	Ser	Asn	Leu	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
		50					55					60					
10	Ser	Gly	Ser	Gly	Thr	Gly	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
		65				70					75					80	
15	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ala	His	Ser	Tyr	Pro	Phe	
					85					90					95		
20	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Asp	Ile	
				100					105						110		
25	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ala	Val	Ser	Leu	Gly	Gln	Arg	
			115					120					125				
30	Ala	Thr	Ile	Ser	Cys	Arg	Ala	Ser	Glu	Ser	Val	Asp	Ser	Tyr	Gly	Gln	
		130					135					140					
35	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Ala	Gly	Gln	Pro	Pro	Lys	Leu	
		145				150					155					160	
40	Leu	Ile	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	
					165					170					175		
45	Ser	Gly	Ser	Gly	Ser	Arg	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Asp	Pro	Val	
				180					185					190			
50	Gln	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Asn	Ala	Glu	Asp	
			195					200					205				
55	Ser	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	
		210					215					220					
60	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	
		225				230					235					240	
65	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	
					245					250					255		
70	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	
				260					265					270			
75	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	
			275					280					285				

EP 2 691 416 B1

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
 290 295 300

5 Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
 305 310 315 320

10 Lys Ser Phe Asn Arg Gly Glu Cys
 325

<210> 20
 <211> 326
 <212> PRT
 15 <213> Artificial Sequence

<220>
 <223> Light chain sequence for cross over dual V-Ig having code LC2

20 <400> 20

Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Val Ser Val Gly
 1 5 10 15

25 Asp Thr Ile Thr Leu Thr Cys His Ala Ser Gln Asn Ile Asp Val Trp
 20 25 30

30 Leu Ser Trp Phe Gln Gln Lys Pro Gly Asn Ile Pro Lys Leu Leu Ile
 35 40 45

35 Tyr Lys Ala Ser Asn Leu His Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Gly Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

40 Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Ala His Ser Tyr Pro Phe
 85 90 95

45 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly Asp Ile Val Leu
 100 105 110

Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg Ala Thr
 115 120 125

50 Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr Gly Gln Ser Tyr
 130 135 140

55 Met His Trp Tyr Gln Gln Lys Ala Gly Gln Pro Pro Lys Leu Leu Ile
 145 150 155 160

EP 2 691 416 B1

Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala Arg Phe Ser Gly
 165 170 175
 5 Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp Pro Val Gln Ala
 180 185 190
 10 Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asn Ala Glu Asp Ser Arg
 195 200 205
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
 210 215 220
 15 Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 225 230 235 240
 20 Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 245 250 255
 25 Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 260 265 270
 Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 275 280 285
 30 Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 290 295 300
 35 Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 305 310 315 320
 40 Phe Asn Arg Gly Glu Cys
 325

<210> 21

<211> 330

<212> PRT

<213> Artificial Sequence

<220>

<223> Light chain sequence for cross over dual V-Ig having code LC3

<400> 21

EP 2 691 416 B1

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly
1				5					10					15	

5	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp
			20						25					30		

10	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile
----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	35	40	45
5	Tyr Lys Ala Ser Asn Leu His Thr Gly Val Pro Ser Arg Phe Ser Gly	50	55 60
10	Ser Gly Ser Gly Thr Gly Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro	65	70 75 80
15	Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Ala His Ser Tyr Pro Phe	85	90 95
20	Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Asp Ile	100	105 110
25	Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg	115	120 125
30	Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr Gly Gln	130	135 140
35	Ser Tyr Met His Trp Tyr Gln Gln Lys Ala Gly Gln Pro Pro Lys Leu	145	150 155 160
40	Leu Ile Tyr Leu Ala Ser Asn Leu Glu Ser Gly Val Pro Ala Arg Phe	165	170 175
45	Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp Pro Val	180	185 190
50	Gln Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asn Ala Glu Asp	195	200 205
55	Ser Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly Gly Arg	210	215 220
	Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln	225	230 235 240
	Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr	245	250 255
	Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser	260	265 270
	Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr	275	280 285

EP 2 691 416 B1

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
290 295 300

5 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
305 310 315 320

10 Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
325 330

<210> 22

<211> 328

<212> PRT

15 <213> Artificial Sequence

<220>

<223> Light chain sequence for cross over dual V-Ig having code LC4

20 <400> 22

25

30

35

40

45

50

55

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ser	Val	Ser	Val	Gly
	1				5					10				15		
5	Asp	Thr	Ile	Thr	Leu	Thr	Cys	His	Ala	Ser	Gln	Asn	Ile	Asp	Val	Trp
				20					25					30		
10	Leu	Ser	Trp	Phe	Gln	Gln	Lys	Pro	Gly	Asn	Ile	Pro	Lys	Leu	Leu	Ile
			35					40					45			
15	Tyr	Lys	Ala	Ser	Asn	Leu	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
		50					55					60				
20	Ser	Gly	Ser	Gly	Thr	Gly	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
	65					70					75				80	
25	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Ala	His	Ser	Tyr	Pro	Phe
					85					90					95	
30	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Asp	Ile	Val	Leu
				100					105					110		
35	Thr	Gln	Ser	Pro	Ala	Ser	Leu	Ala	Val	Ser	Leu	Gly	Gln	Arg	Ala	Thr
			115					120					125			
40	Ile	Ser	Cys	Arg	Ala	Ser	Glu	Ser	Val	Asp	Ser	Tyr	Gly	Gln	Ser	Tyr
		130					135					140				
45	Met	His	Trp	Tyr	Gln	Gln	Lys	Ala	Gly	Gln	Pro	Pro	Lys	Leu	Leu	Ile
	145					150					155				160	
50	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly
55																

EP 2 691 416 B1

	165	170	175
5	Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asp Pro Val Gln Ala 180 185 190		
10	Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Asn Ala Glu Asp Ser Arg 195 200 205		
15	Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly Gly Arg Thr Val 210 215 220		
20	Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys 225 230 235 240		
25	Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg 245 250 255		
30	Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn 260 265 270		
35	Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser 275 280 285		
40	Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys 290 295 300		
45	Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr 305 310 315 320		
50	Lys Ser Phe Asn Arg Gly Glu Cys 325		

<210> 23
 <211> 107
 <212> PRT
 <213> Homo sapiens

<400> 23

EP 2 691 416 B1

1	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu
				5					10					15		
5	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe
			20						25					30		
10	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln
			35					40					45			
	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser
15																
			50					55					60			
20	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu
	65					70					75				80	
	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser
				85						90					95	
25																
	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys					
				100					105							

<210> 24
 <211> 329
 <212> PRT
 <213> Homo sapiens
 <400> 24

EP 2 691 416 B1

	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys
	1				5					10					15	
5	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
				20					25					30		
10	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
			35					40					45			
15	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
	50						55					60				
20	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr
	65					70				75					80	
25	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
				85					90						95	
30	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys
				100					105					110		
35	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro
				115				120					125			
40	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys
	130						135					140				
45	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp
	145					150				155					160	
50	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
55																

EP 2 691 416 B1

	165	170	175
5	Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu 180 185 190		
10	His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn 195 200 205		
15	Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly 210 215 220		
20	Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu 225 230 235 240		
25	Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr 245 250 255		
30	Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn 260 265 270		
35	Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe 275 280 285		
40	Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn 290 295 300		
45	Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr 305 310 315 320		
50	Gln Lys Ser Leu Ser Leu Ser Pro Gly 325		
55	<210> 25 <211> 4 <212> PRT <213> Artificial Sequence		
	<220> <223> Peptide linker with four glycine residues		
	<400> 25		
	Gly Gly Gly Gly 1		
	<210> 26 <211> 5 <212> PRT <213> Artificial Sequence		

EP 2 691 416 B1

<220>

<223> Peptide linker with 5 glycine amino acid residues

<400> 26

5

Gly Gly Gly Gly Gly
1 5

<210> 27

10

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

15

<223> Peptide linker with 6 glycine amino acid residues.

<400> 27

20

Gly Gly Gly Gly Gly Gly
1 5

<210> 28

<211> 7

<212> PRT

25

<213> Artificial Sequence

<220>

<223> Peptide linker with seven glycine amino acid residues.

30

<400> 28

Gly Gly Gly Gly Gly Gly Gly
1 5

35

<210> 29

<211> 8

<212> PRT

<213> Artificial Sequence

40

<220>

<223> Peptide linker with eight glycine amino acid residues.

<400> 29

45

Gly Gly Gly Gly Gly Gly Gly Gly
1 5

<210> 30

<211> 5

50

<212> PRT

<213> Artificial Sequence

<220>

<223> artificial peptide linker

55

<400> 30

EP 2 691 416 B1

Gly Gly Gly Gly Ser
1 5

5 <210> 31
<211> 10
<212> PRT
<213> Artificial Sequence

10 <220>
<223> artificial peptide linker

<400> 31

15 Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10

<210> 32
<211> 582
<212> PRT
20 <213> Artificial Sequence

<220>
<223> Heavy chain sequence for cross over dual V-Ig having code HC10

25 <400> 32

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser	
	1				5					10					15		
5	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Gly	Thr	Phe	Ser	Ser	Tyr	
				20					25					30			
10	Ala	Ile	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35					40					45				
15	Gly	Gly	Ile	Ile	Pro	Ile	Phe	Gly	Thr	Ala	Asn	Tyr	Ala	Gln	Lys	Phe	
		50					55					60					
20	Gln	Gly	Arg	Val	Thr	Ile	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr	
	65					70					75					80	
25	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		
30	Ala	Arg	Ala	Pro	Leu	Arg	Phe	Leu	Glu	Trp	Ser	Thr	Gln	Asp	His	Tyr	
				100					105					110			
35	Tyr	Tyr	Tyr	Tyr	Met	Asp	Val	Trp	Gly	Lys	Gly	Thr	Thr	Val	Thr	Val	
			115					120					125				
40	Ser	Ser	Gly	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	
		130					135					140					
45	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Asn	Ile	
	145					150					155					160	

EP 2 691 416 B1

	Lys	Asp	Thr	Tyr	Ile	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
					165					170					175		
5	Glu	Trp	Val	Ala	Arg	Ile	Tyr	Pro	Thr	Asn	Gly	Tyr	Thr	Arg	Tyr	Ala	
				180					185					190			
10	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Ala	Asp	Thr	Ser	Lys	Asn	
			195					200					205				
	Thr	Ala	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	
		210					215					220					
15	Tyr	Tyr	Cys	Ser	Arg	Trp	Gly	Gly	Asp	Gly	Phe	Tyr	Ala	Met	Asp	Tyr	
	225					230				235						240	
20	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Gly	Ala	Ser	Thr	
					245					250					255		
25	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	
				260					265					270			
	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	
			275					280					285				
30	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	
		290					295					300					
35	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	
	305					310					315					320	
40	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	
					325					330					335		
45	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	
				340					345					350			
50	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	
			355					360					365				
	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	
		370					375					380					
55	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	
	385					390				395						400	
	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	

EP 2 691 416 B1

	405	410	415
5	Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr 420 425 430		
10	Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp 435 440 445		
15	Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu 450 455 460		
20	Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg 465 470 475 480		
25	Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys 485 490 495		
30	Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp 500 505 510		
35	Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys 515 520 525		
40	Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser 530 535 540		
45	Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser 545 550 555 560		
50	Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser 565 570 575		
55	Leu Ser Leu Ser Pro Gly 580		

<210> 33

<211> 582

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain sequence for cross over dual V-Ig having code HC11

<400> 33

EP 2 691 416 B1

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

5

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Asn	Ile	Lys	Asp	Thr
			20					25					30		

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

Tyr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 5 Ala Arg Ile Tyr Pro Thr Asn Gly Tyr Thr Arg Tyr Ala Asp Ser Val
 50 55 60
 10 Lys Gly Arg Phe Thr Ile Ser Ala Asp Thr Ser Lys Asn Thr Ala Tyr
 65 70 75 80
 15 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 20 Ser Arg Trp Gly Gly Asp Gly Phe Tyr Ala Met Asp Tyr Trp Gly Gln
 100 105 110
 25 Gly Thr Leu Val Thr Val Ser Ser Gly Glu Val Gln Leu Val Gln Ser
 115 120 125
 30 Gly Ala Glu Val Lys Lys Pro Gly Ser Ser Val Lys Val Ser Cys Lys
 130 135 140
 35 Ala Ser Gly Gly Thr Phe Ser Ser Tyr Ala Ile Ser Trp Val Arg Gln
 145 150 155 160
 40 Ala Pro Gly Gln Gly Leu Glu Trp Met Gly Gly Ile Ile Pro Ile Phe
 165 170 175
 45 Gly Thr Ala Asn Tyr Ala Gln Lys Phe Gln Gly Arg Val Thr Ile Thr
 180 185 190
 50 Ala Asp Lys Ser Thr Ser Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg
 195 200 205
 55 Ser Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Ala Pro Leu Arg Phe
 210 215 220
 60 Leu Glu Trp Ser Thr Gln Asp His Tyr Tyr Tyr Tyr Tyr Met Asp Val
 225 230 235 240
 65 Trp Gly Lys Gly Thr Thr Val Thr Val Ser Ser Gly Gly Ala Ser Thr
 245 250 255
 70 Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser
 260 265 270
 75 Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu

EP 2 691 416 B1

	275	280	285
5	Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His 290 295 300		
	Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser 305 310 315 320		
10	Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys 325 330 335		
15	Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu 340 345 350		
20	Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro 355 360 365		
25	Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys 370 375 380		
30	Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val 385 390 395 400		
35	Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp 405 410 415		
	Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr 420 425 430		
40	Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp 435 440 445		
45	Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu 450 455 460		
	Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg 465 470 475 480		
50	Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys 485 490 495		
55	Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp 500 505 510		
	Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys 515 520 525		

EP 2 691 416 B1

	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser
	530						535					540				
5	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser
	545					550					555					560
10	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
					565					570					575	
	Leu	Ser	Leu	Ser	Pro	Gly										
					580											
15	<210> 34															
	<211> 571															
	<212> PRT															
	<213> Artificial Sequence															
20	<220>															
	<223> Heavy chain sequence for cross over dual V-Ig having code HC12															
25	<400> 34															
30																
35																
40																
45																
50																
55																

EP 2 691 416 B1

	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	
	1				5					10					15		
5	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Phe	
				20					25					30			
10	Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile	
			35					40					45				
	Ser	Val	Ile	Asp	Thr	Arg	Gly	Ala	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	Lys	
		50					55					60					
15	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	
	65					70					75					80	
20	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	
					85					90					95		
25	Arg	Leu	Gly	Asn	Phe	Tyr	Tyr	Gly	Met	Asp	Val	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
	Thr	Val	Thr	Val	Ser	Ser	Gly	Gln	Val	Gln	Leu	Gln	Glu	Ser	Gly	Pro	
				115				120					125				
30	Gly	Leu	Val	Lys	Pro	Ser	Gln	Thr	Leu	Ser	Leu	Thr	Cys	Thr	Val	Ser	
		130					135					140					
35	Gly	Gly	Ser	Ile	Ser	Ser	Gly	Asp	Tyr	Tyr	Trp	Ser	Trp	Ile	Arg	Gln	
40																	
45																	
50																	
55																	

EP 2 691 416 B1

	145		150		155		160									
5	Pro	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile	Gly	Tyr	Ile	Tyr	Tyr	Ser	Gly
					165					170					175	
10	Ser	Thr	Asp	Tyr	Asn	Pro	Ser	Leu	Lys	Ser	Arg	Val	Thr	Met	Ser	Val
				180					185					190		
15	Asp	Thr	Ser	Lys	Asn	Gln	Phe	Ser	Leu	Lys	Val	Asn	Ser	Val	Thr	Ala
			195					200					205			
20	Ala	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Val	Ser	Ile	Phe	Gly	Val
		210					215					220				
25	Gly	Thr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser
	225					230					235					240
30	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser
					245					250					255	
35	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys
				260					265					270		
40	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu
			275					280					285			
45	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu
	290						295					300				
50	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr
	305					310				315					320	
55	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val
					325					330					335	
60	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro
				340					345					350		
65	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe
		355						360					365			
70	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
		370				375						380				
75	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
	385					390					395					400

EP 2 691 416 B1

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 405 410 415
 5 Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
 420 425 430
 10 Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 435 440 445
 Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala
 450 455 460
 15 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 465 470 475 480
 20 Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 485 490 495
 25 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 500 505 510
 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
 515 520 525
 30 Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
 530 535 540
 35 Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 545 550 555 560
 40 Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 565 570

<210> 35

<211> 571

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain sequence for cross over dual V-Ig having code HC13

<400> 35

EP 2 691 416 B1

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

5 Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
 20 25 30

10 Asp Tyr Tyr Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	35	40	45
5	Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asp Tyr Asn Pro Ser 50 55 60		
	Leu Lys Ser Arg Val Thr Met Ser Val Asp Thr Ser Lys Asn Gln Phe 65 70 75 80		
10	Ser Leu Lys Val Asn Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95		
15	Cys Ala Arg Val Ser Ile Phe Gly Val Gly Thr Phe Asp Tyr Trp Gly 100 105 110		
20	Gln Gly Thr Leu Val Thr Val Ser Ser Gly Glu Val Gln Leu Val Gln 115 120 125		
25	Ser Gly Gly Gly Leu Val Lys Pro Gly Gly Ser Leu Arg Leu Ser Cys 130 135 140		
30	Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe Ala Met His Trp Val Arg 145 150 155 160		
35	Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser Val Ile Asp Thr Arg 165 170 175		
40	Gly Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser 180 185 190		
45	Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln Met Asn Ser Leu Arg 195 200 205		
50	Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Leu Gly Asn Phe Tyr 210 215 220		
55	Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser 225 230 235 240		
	Gly Gly Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser 245 250 255		
	Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys 260 265 270		
	Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu 275 280 285		

EP 2 691 416 B1

	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	
	290						295					300					
5	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	
	305					310					315					320	
	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	
10					325					330					335		
	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	
				340					345					350			
15	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	
			355					360					365				
	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	
20			370				375					380					
	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	
25						390					395					400	
	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	
					405					410					415		
30	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	
				420					425					430			
	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	
35			435					440					445				
	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	
		450					455					460					
40	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	
	465					470					475					480	
	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	
45					485					490					495		
	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	
50				500					505					510			
	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	
			515					520					525				
55	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	
	530						535					540					

EP 2 691 416 B1

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
545 550 555 560

5 Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
565 570

<210> 36

<211> 572

10 <212> PRT

<213> Artificial Sequence

<220>

15 <223> Heavy chain sequence for cross over dual V-Ig having code HC14

<400> 36

20 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
20 25 30

25 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

30 Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
50 55 60

35 Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

40 Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
100 105 110

45 Gln Gly Thr Leu Val Thr Val Ser Ser Gly Glu Val Gln Leu Val Gln
115 120 125

50 Ser Gly Ala Glu Val Lys Lys Pro Gly Glu Ser Leu Lys Ile Ser Cys
130 135 140

Lys Gly Ser Gly Tyr Ser Phe Thr Thr Tyr Trp Leu Gly Trp Val Arg
145 150 155 160

55 Gln Met Pro Gly Lys Gly Leu Asp Trp Ile Gly Ile Met Ser Pro Val
165 170 175

EP 2 691 416 B1

	Asp	Ser	Asp	Ile	Arg	Tyr	Ser	Pro	Ser	Phe	Gln	Gly	Gln	Val	Thr	Met	
				180					185					190			
5	Ser	Val	Asp	Lys	Ser	Ile	Thr	Thr	Ala	Tyr	Leu	Gln	Trp	Asn	Ser	Leu	
			195					200					205				
	Lys	Ala	Ser	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	Ala	Arg	Arg	Arg	Pro	Gly	
10		210					215					220					
	Gln	Gly	Tyr	Phe	Asp	Phe	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	
	225					230					235					240	
15	Ser	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	
					245					250					255		
	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	
20				260					265					270			
	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	
25			275					280					285				
	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	
	290						295					300					
30	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	
	305					310					315					320	
	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	
35					325					330					335		
	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	
				340					345					350			
40	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	
			355					360					365				
45	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	
	370						375					380					
	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	
50	385					390					395					400	
	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	
					405				410						415		
55	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	
				420					425					430			

EP 2 691 416 B1

Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
435 440 445

5 Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys
450 455 460

10 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
465 470 475 480

Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
485 490 495

15 Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
500 505 510

20 Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
515 520 525

25 Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln
530 535 540

Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
545 550 555 560

30 His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
565 570

<210> 37
35 <211> 572
<212> PRT
<213> Artificial Sequence

<220>
40 <223> Heavy chain sequence for cross over dual V-Ig having code HC15

<400> 37

45 Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Thr Tyr
20 25 30

50 Trp Leu Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu Asp Trp Ile
35 40 45

55 Gly Ile Met Ser Pro Val Asp Ser Asp Ile Arg Tyr Ser Pro Ser Phe
50 55 60

EP 2 691 416 B1

	Gln	Gly	Gln	Val	Thr	Met	Ser	Val	Asp	Lys	Ser	Ile	Thr	Thr	Ala	Tyr	65	70	75	80
5	Leu	Gln	Trp	Asn	Ser	Leu	Lys	Ala	Ser	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	85	90	95	
10	Ala	Arg	Arg	Arg	Pro	Gly	Gln	Gly	Tyr	Phe	Asp	Phe	Trp	Gly	Gln	Gly	100	105	110	
	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	115	120	125	
15	Gly	Gly	Leu	Val	Gln	Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	130	135	140	
20	Ser	Gly	Phe	Thr	Phe	Asp	Asp	Tyr	Ala	Met	His	Trp	Val	Arg	Gln	Ala	145	150	155	160
25	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Ala	Ile	Thr	Trp	Asn	Ser	Gly	165	170	175	
	His	Ile	Asp	Tyr	Ala	Asp	Ser	Val	Glu	Gly	Arg	Phe	Thr	Ile	Ser	Arg	180	185	190	
30	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	195	200	205	
35	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Lys	Val	Ser	Tyr	Leu	Ser	Thr	210	215	220	
40	Ala	Ser	Ser	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	225	230	235	240
	Ser	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	245	250	255	
45	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	260	265	270	
50	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	275	280	285	
	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	290	295	300	
55	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	305	310	315	320

EP 2 691 416 B1

	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	
					325					330					335		
5	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	
				340					345					350			
10	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	
			355					360					365				
	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	
		370					375					380					
15	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	
	385					390					395					400	
20	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	
					405					410					415		
25	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	
				420					425					430			
	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	
			435					440					445				
30	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	
		450					455					460					
35	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	
	465					470					475					480	
	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	
					485					490					495		
40	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	
				500					505					510			
45	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	
			515					520					525				
50	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	
		530					535					540					
	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	
	545					550					555					560	
55	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly					
					565					570							

EP 2 691 416 B1

<210> 38
 <211> 571
 <212> PRT
 <213> Artificial Sequence

5

<220>
 <223> Heavy chain sequence for cross over dual V-Ig having code HC16

<400> 38

10

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Arg
 1 5 10 15

15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asp Asp Tyr
 20 25 30

20

Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

25

Ser Ala Ile Thr Trp Asn Ser Gly His Ile Asp Tyr Ala Asp Ser Val
 50 55 60

Glu Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

30

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

35

Ala Lys Val Ser Tyr Leu Ser Thr Ala Ser Ser Leu Asp Tyr Trp Gly
 100 105 110

Gln Gly Thr Leu Val Thr Val Ser Ser Gly Gln Val Gln Leu Val Glu
 115 120 125

40

Ser Gly Gly Gly Val Val Gln Pro Gly Arg Ser Leu Arg Leu Ser Cys
 130 135 140

45

Ala Ala Ser Gly Phe Thr Phe Ser Val Tyr Gly Met Asn Trp Val Arg
 145 150 155 160

50

Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Ile Ile Trp Tyr Asp
 165 170 175

Gly Asp Asn Gln Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile
 180 185 190

55

Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Gly Leu
 195 200 205

EP 2 691 416 B1

	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	Asp	Leu	Arg	Thr	
	210						215					220					
5	Gly	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	
	225					230					235					240	
	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	
10					245					250					255		
	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	
				260					265					270			
15	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	
			275					280					285				
	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	
20							295					300					
	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	
25						310					315					320	
	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	
					325					330					335		
30	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	
				340					345					350			
	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	
35				355				360					365				
	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	
						375						380					
40	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	
	385					390					395					400	
	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	
45					405					410					415		
	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	
50				420					425					430			
	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	
			435					440					445				
55	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	
	450						455					460					

EP 2 691 416 B1

Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
 465 470 475 480
 5 Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 485 490 495
 10 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 500 505 510
 15 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
 515 520 525
 Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
 530 535 540
 20 Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 545 550 555 560
 25 Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 565 570
 <210> 39
 <211> 571
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> Heavy chain sequence for cross over dual V-Ig having code HC17
 35 <400> 39
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Val Tyr
 20 25 30
 45 Gly Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 50 Ala Ile Ile Trp Tyr Asp Gly Asp Asn Gln Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 55 Leu Gln Met Asn Gly Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

EP 2 691 416 B1

	Ala	Arg	Asp	Leu	Arg	Thr	Gly	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
5	Leu	Val	Thr	Val	Ser	Ser	Gly	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	
			115					120					125				
	Gly	Leu	Val	Gln	Pro	Gly	Arg	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	
10		130					135					140					
	Gly	Phe	Thr	Phe	Asp	Asp	Tyr	Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	
	145					150					155					160	
15	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	Ala	Ile	Thr	Trp	Asn	Ser	Gly	His	
				165						170					175		
	Ile	Asp	Tyr	Ala	Asp	Ser	Val	Glu	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	
20				180					185					190			
	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	
25			195					200					205				
	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Lys	Val	Ser	Tyr	Leu	Ser	Thr	Ala	
	210						215					220					
30	Ser	Ser	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	
	225					230					235					240	
	Gly	Gly	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	
35					245					250					255		
	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	
				260					265					270			
40	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	
			275					280					285				
45	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	
	290						295					300					
	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	
50	305					310					315					320	
	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	
				325						330					335		
55	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	
				340					345					350			

EP 2 691 416 B1

	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe
			355					360					365			
5	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
		370					375					380				
10	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
	385					390					395					400
15	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro
				405						410					415	
20	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr
				420					425					430		
25	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val
		435						440					445			
30	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala
	450						455					460				
35	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg
	465					470					475					480
40	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly
					485					490					495	
45	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro
				500					505					510		
50	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser
			515					520					525			
55	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln
	530						535					540				
60	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His
	545					550					555					560
65	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly					
					565					570						

<210> 40

<211> 334

<212> PRT

<213> Artificial Sequence

<220>

EP 2 691 416 B1

<223> Light chain sequence for cross over dual V-Ig having code L10

<400> 40

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
	1				5					10					15		
5	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Val	Asn	Thr	Ala	
				20					25					30			
10	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	
			35					40					45				
15	Tyr	Ser	Ala	Ser	Phe	Leu	Tyr	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
		50					55					60					
20	Ser	Arg	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
	65					70					75					80	
25	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	His	Tyr	Thr	Thr	Pro	Pro	
					85					90					95		
30	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly	
				100					105					110			
35	Gly	Gly	Ser	Ser	Glu	Leu	Thr	Gln	Asp	Pro	Ala	Val	Ser	Val	Ala	Leu	
			115					120					125				
40	Gly	Gln	Thr	Val	Arg	Ile	Thr	Cys	Gln	Gly	Asp	Ser	Leu	Arg	Ser	Tyr	
		130					135					140					
45	Tyr	Ala	Thr	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Ile	Leu	Val	
	145					150					155					160	
50	Ile	Tyr	Gly	Glu	Asn	Lys	Arg	Pro	Ser	Gly	Ile	Pro	Asp	Arg	Phe	Ser	
					165					170					175		
55	Gly	Ser	Ser	Ser	Gly	Asn	Thr	Ala	Ser	Leu	Thr	Ile	Thr	Gly	Ala	Gln	
				180					185					190			
60	Ala	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Lys	Ser	Arg	Asp	Gly	Ser	Gly	
			195					200					205				
65	Gln	His	Leu	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gly	
		210					215					220					
70	Gly	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	
	225					230					235					240	

EP 2 691 416 B1

	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu
					245					250					255	
5	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn
				260					265					270		
10	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser
			275					280					285			
15	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala
		290					295					300				
20	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly
	305					310					315					320
25	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys		
					325					330						
30	<210> 41															
	<211> 334															
	<212> PRT															
	<213> Artificial Sequence															
	<220>															
	<223> Light chain sequence for cross over dual V-Ig having code LC11															
	<400> 41															
35	Ser	Ser	Glu	Leu	Thr	Gln	Asp	Pro	Ala	Val	Ser	Val	Ala	Leu	Gly	Gln
	1				5					10					15	
40	Thr	Val	Arg	Ile	Thr	Cys	Gln	Gly	Asp	Ser	Leu	Arg	Ser	Tyr	Tyr	Ala
				20					25					30		
45	Thr	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Ile	Leu	Val	Ile	Tyr
			35					40					45			
50	Gly	Glu	Asn	Lys	Arg	Pro	Ser	Gly	Ile	Pro	Asp	Arg	Phe	Ser	Gly	Ser
		50					55					60				
55	Ser	Ser	Gly	Asn	Thr	Ala	Ser	Leu	Thr	Ile	Thr	Gly	Ala	Gln	Ala	Glu
	65				70					75					80	
	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Lys	Ser	Arg	Asp	Gly	Ser	Gly	Gln	His
				85						90					95	
	Leu	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gly	Gly	Gly
				100					105					110		

EP 2 691 416 B1

	Gly	Gly	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	
			115					120					125				
5	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Val	
		130					135					140					
10	Asn	Thr	Ala	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	
	145					150					155					160	
	Leu	Leu	Ile	Tyr	Ser	Ala	Ser	Phe	Leu	Tyr	Ser	Gly	Val	Pro	Ser	Arg	
					165					170					175		
15	Phe	Ser	Gly	Ser	Arg	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	
				180					185					190			
20	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	His	Tyr	Thr	
			195					200					205				
25	Thr	Pro	Pro	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	
	210					215						220					
30	Gly	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	
	225				230					235						240	
	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	
				245					250					255			
35	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	
				260				265					270				
40	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	
			275					280					285				
	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	
	290				295					300							
45	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	
	305				310					315						320	
50	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys			
				325					330								

<210> 42

<211> 333

55 <212> PRT

<213> Artificial Sequence

<220>

EP 2 691 416 B1

<223> Light chain sequence for cross over dual V-Ig having code LC12

<400> 42

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Leu	Ser	Pro	Gly	
	1				5					10					15		
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Tyr	
				20					25					30			
10	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	
			35					40					45				
15	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala	Arg	Phe	Ser	Gly	
	50						55					60					
20	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Glu	Pro	
	65					70					75					80	
25	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	His	Gln	Tyr	Gly	Ser	Thr	Pro	Leu	
					85					90					95		
30	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Ala	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly	
				100					105					110			
35	Gly	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Gly	Thr	Leu	Ser	Val	Ser	
			115					120					125				
40	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Gly	
		130					135					140					
45	Ser	Ser	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	
	145					150					155					160	
50	Leu	Ile	Lys	Tyr	Ala	Ser	Gln	Ser	Leu	Ser	Gly	Ile	Pro	Asp	Arg	Phe	
					165					170					175		
55	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Arg	Leu	
				180					185					190			
60	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	His	Gln	Ser	Ser	Arg	Leu	
			195					200					205				
65	Pro	His	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	
		210					215					220					
70	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	
	225					230					235					240	

EP 2 691 416 B1

	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn
					245					250					255	
5	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala
				260					265					270		
10	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys
			275					280					285			
15	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp
	290						295					300				
20	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu
	305					310					315					320
25	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys			
					325					330						
	<210> 43															
	<211> 333															
	<212> PRT															
	<213> Artificial Sequence															
	<220>															
	<223> Light chain sequence for cross over dual V-Ig having code LC13															
30	<400> 43															
35	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Gly	Thr	Leu	Ser	Val	Ser	Pro	Gly
	1				5					10					15	
40	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Gly	Ser	Ser
				20					25					30		
45	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile
			35				40						45			
50	Lys	Tyr	Ala	Ser	Gln	Ser	Leu	Ser	Gly	Ile	Pro	Asp	Arg	Phe	Ser	Gly
	50						55					60				
55	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Arg	Leu	Glu	Pro
	65				70						75				80	
60	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	His	Gln	Ser	Ser	Arg	Leu	Pro	His
					85					90					95	
65	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly
				100					105					110		

EP 2 691 416 B1

	Gly	Gly	Glu	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Leu	Ser	
			115					120					125				
5	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Ser	
		130					135					140					
10	Ser	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	
	145					150					155					160	
	Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Thr	Gly	Ile	Pro	Ala	Arg	Phe	
				165						170					175		
15	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	
				180					185					190			
20	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	His	Gln	Tyr	Gly	Ser	Thr	
			195					200					205				
25	Pro	Leu	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Ala	Glu	Ile	Lys	Gly	Gly	Gly	
	210						215					220					
	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	
	225					230					235					240	
30	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	
				245						250					255		
35	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	
				260					265					270			
40	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	
			275					280					285				
	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	
	290						295					300					
45	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	
	305					310					315					320	
50	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys				
					325					330							

<210> 44

<211> 333

55 <212> PRT

<213> Artificial Sequence

<220>

EP 2 691 416 B1

<223> Light chain sequence for cross over dual V-Ig having code LC14

<400> 44

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
	1				5					10					15		
5	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Ser	Ser	Trp	
				20					25					30			
10	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Glu	Lys	Ala	Pro	Lys	Ser	Leu	Ile	
			35					40					45				
15	Tyr	Ala	Ala	Ser	Ser	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
		50					55					60					
20	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
	65					70					75					80	
25	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Tyr	Asn	Ile	Tyr	Pro	Tyr	
					85					90					95		
30	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly	
				100					105					110			
35	Gly	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	
			115					120					125				
40	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Arg	
		130					135					140					
45	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	
	145					150					155					160	
50	Leu	Ile	Tyr	Ala	Ala	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	
				165						170					175		
55	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	
			180					185						190			
60	Gln	Pro	Glu	Asp	Val	Ala	Thr	Tyr	Tyr	Cys	Gln	Arg	Tyr	Asn	Arg	Ala	
			195					200					205				
65	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	
		210				215						220					
70	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	
	225					230					235					240	

EP 2 691 416 B1

	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn
					245					250					255	
5	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala
				260					265					270		
10	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys
			275					280					285			
15	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp
	290						295					300				
20	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu
	305					310					315					320
25	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys			
					325					330						
30	<210> 45															
	<211> 333															
35	<212> PRT															
	<213> Artificial Sequence															
40	<220>															
	<223> Light chain sequence for cross over dual V-Ig having code LC15															
45	<400> 45															

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
	1				5					10					15	
5	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Arg	Asn	Tyr
				20					25					30		
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
10				35				40					45			
	Tyr	Ala	Ala	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
		50					55					60				
15	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
	65					70					75					80
20	Glu	Asp	Val	Ala	Thr	Tyr	Tyr	Cys	Gln	Arg	Tyr	Asn	Arg	Ala	Pro	Tyr
					85					90					95	
	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly
25				100					105					110		
	Gly	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser

EP 2 691 416 B1

	115	120	125
5	Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser 130	135	140
10	Ser Trp Leu Ala Trp Tyr Gln Gln Lys Pro Glu Lys Ala Pro Lys Ser 145	150	155
15	Leu Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe 165	170	175
20	Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu 180	185	190
25	Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ile Tyr 195	200	205
30	Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly 210	215	220
35	Gly Gly Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser 225	230	235
40	Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn 245	250	255
45	Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala 260	265	270
50	Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys 275	280	285
55	Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp 290	295	300
60	Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu 305	310	315
65	Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys 325	330	

<210> 46

<211> 333

<212> PRT

<213> Artificial Sequence

<220>

<223> Light chain sequence for cross over dual V-Ig having code LC16

<400> 46

5

10

15

20

25

30

35

40

45

50

55

EP 2 691 416 B1

	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Asp	Phe	Gln	Ser	Val	Thr	Pro	Lys	
	1				5					10					15		
5	Glu	Lys	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Gly	Ser	Ser	
				20					25					30			
	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gln	Ser	Pro	Lys	Leu	Leu	Ile	
10			35					40					45				
	Lys	Tyr	Ala	Ser	Gln	Ser	Phe	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
		50					55					60					
15	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Asn	Ser	Leu	Glu	Ala	
	65					70					75					80	
	Glu	Asp	Ala	Ala	Ala	Tyr	Tyr	Cys	His	Gln	Ser	Ser	Ser	Leu	Pro	Phe	
20					85					90					95		
	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys	Gly	Gly	Gly	Gly	Gly	
25				100					105					110			
	Gly	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	
			115					120					125				
30	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Arg	
		130					135					140					
	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	
35		145				150					155					160	
	Leu	Ile	Tyr	Ala	Ala	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	
				165						170					175		
40	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	
				180					185					190			
	Gln	Pro	Glu	Asp	Val	Ala	Thr	Tyr	Tyr	Cys	Gln	Arg	Tyr	Asn	Arg	Ala	
45			195					200					205				
	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	
50		210					215					220					
	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	
	225					230					235					240	
55	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	

EP 2 691 416 B1

		245		250		255
5	Asn Phe Tyr	Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala				
		260		265		270
10	Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys					
		275		280		285
15	Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp					
		290		295		300
20	Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu					
		305		310		315
25	Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys					
		325		330		

<210> 47

<211> 333

<212> PRT

<213> Artificial Sequence

<220>

<223> Light chain sequence for cross over dual V-Ig having code LC17

<400> 47

EP 2 691 416 B1

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
	1				5					10					15	
5	Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Gly	Ile	Arg	Asn	Tyr
				20					25					30		
	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
10			35					40					45			
	Tyr	Ala	Ala	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
		50					55					60				
15	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
	65					70					75					80
20	Glu	Asp	Val	Ala	Thr	Tyr	Tyr	Cys	Gln	Arg	Tyr	Asn	Arg	Ala	Pro	Tyr
					85					90					95	
	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Gly	Gly	Gly	Gly	Gly
25				100					105					110		
	Gly	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Asp	Phe	Gln	Ser	Val	Thr
			115					120					125			
30																
35																
40																
45																
50																
55																

EP 2 691 416 B1

	Pro	Lys	Glu	Lys	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Ser	Ile	Gly	
	130						135					140					
5	Ser	Ser	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gln	Ser	Pro	Lys	Leu	
	145					150					155					160	
10	Leu	Ile	Lys	Tyr	Ala	Ser	Gln	Ser	Phe	Ser	Gly	Val	Pro	Ser	Arg	Phe	
					165					170					175		
15	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Asn	Ser	Leu	
				180					185					190			
20	Glu	Ala	Glu	Asp	Ala	Ala	Ala	Tyr	Tyr	Cys	His	Gln	Ser	Ser	Ser	Leu	
			195					200					205				
25	Pro	Phe	Thr	Phe	Gly	Pro	Gly	Thr	Lys	Val	Asp	Ile	Lys	Gly	Gly	Gly	
	210						215					220					
30	Gly	Gly	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	
	225					230					235					240	
35	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	
					245					250					255		
40	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	
				260					265					270			
45	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	
			275					280					285				
50	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	
	290						295					300					
55	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	
	305					310					315					320	
60	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys				
					325					330							

<210> 48

50 <211> 7

<212> PRT

<213> Artificial Sequence

<220>

55 <223> Peptide linker

<400> 48

EP 2 691 416 B1

Ala Ser Thr Lys Gly Pro Ser
1 5

5 <210> 49
 <211> 7
 <212> PRT
 <213> Artificial Sequence

10 <220>
 <223> Peptide linker

<400> 49

15 **Arg Thr Val Ala Ala Pro Ser**
1 5

<210> 50
 <211> 7
 <212> PRT
 20 <213> Artificial Sequence

<220>
 <223> Peptide linker

25 <400> 50

Gly Gln Pro Lys Ala Ala Pro
1 5

30 <210> 51
 <211> 7
 <212> PRT
 <213> Artificial Sequence

35 <220>
 <223> Peptide linker

<400> 51

40 **His Ile Asp Ser Pro Asn Lys**
1 5

<210> 52
 <211> 5
 45 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Peptide linker

50 <400> 52

Thr Lys Gly Pro Ser
1 5

55 <210> 53
 <211> 5
 <212> PRT

EP 2 691 416 B1

<213> Artificial Sequence

<220>

<223> Peptide linker

5

<400> 53

Thr Val Ala Ala Pro
1 5

10

<210> 54

<211> 5

<212> PRT

15

<213> Artificial Sequence

<220>

<223> Peptide linker

20

<400> 54

Gln Pro Lys Ala Ala
1 5

25

<210> 55

<211> 5

<212> PRT

<213> Artificial Sequence

30

<220>

<223> Peptide linker

<400> 55

35

Gln Arg Ile Glu Gly
1 5

<210> 56

<211> 7

40

<212> PRT

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Artificial peptide linker

45

<400> 56

Gly Gly Cys Gly Gly Gly Gly
1 5

50

<210> 57

<211> 7

<212> PRT

55

<213> Artificial Sequence

<220>

<223> Artificial peptide linker

<400> 57

5 Gly Gly Gly Cys Gly Gly Gly
1 5

<210> 58

<211> 7

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Artificial peptide linker

15 <400> 58

Gly Gly Gly Gly Cys Gly Gly
1 5

20 <210> 59

<211> 7

<212> PRT

<213> Artificial Sequence

25 <220>

<223> Artificial peptide linker

<400> 59

30 Gly Gly Gly Gly Gly Cys Gly
1 5

Claims

35

1. An antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites, wherein two polypeptide chains have a structure represented by the formula:

40
$$V_{L1}-L_1-V_{L2}-L_2-C_L \quad [I]$$

and two polypeptide chains have a structure represented by the formula:

45
$$V_{H2}-L_3-V_{H1}-L_4-C_{H1}-Fc \quad [II]$$

wherein:

50 V_{L1} is a first immunoglobulin light chain variable domain;

V_{L2} is a second immunoglobulin light chain variable domain;

V_{H1} is a first immunoglobulin heavy chain variable domain;

V_{H2} is a second immunoglobulin heavy chain variable domain;

C_L is an immunoglobulin light chain constant domain;

C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;

55 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains; and


L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

wherein:

- L₁ is 3 to 12 amino acid residues in length;
- L₂ is 3 to 14 amino acid residues in length;
- L₃ is 1 to 8 amino acid residues in length; and
- L₄ is 1 to 3 amino acid residues in length;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

2. An antibody-like binding protein comprising two polypeptide chains that form two antigen binding sites, wherein a first polypeptide chain has a structure represented by the formula:



and a second polypeptide chain has a structure represented by the formula:



wherein:

- V_{L1} is a first immunoglobulin light chain variable domain;
- V_{L2} is a second immunoglobulin light chain variable domain;
- V_{H1} is a first immunoglobulin heavy chain variable domain;
- V_{H2} is a second immunoglobulin heavy chain variable domain;
- C_L is an immunoglobulin light chain constant domain;
- C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain; and
- L₁, L₂, L₃, and L₄ are amino acid linkers;

wherein:

- L₁ is 3 to 12 amino acid residues in length;
- L₂ is 3 to 14 amino acid residues in length;
- L₃ is 1 to 8 amino acid residues in length; and
- L₄ is 1 to 3 amino acid residues in length;

and wherein the first and second polypeptides form a cross-over light chain-heavy chain pair.


3. The antibody-like binding protein of either claim 1 or 2, wherein:

- L₁ is 5 to 10 amino acid residues in length;
- L₂ is 5 to 8 amino acid residues in length;
- L₃ is 1 to 5 amino acid residues in length; and
- L₄ is 1 to 2 amino acid residues in length.


4. The antibody-like binding protein of either claim 1 or 2, wherein:

- L₁ is 7 amino acid residues in length;
- L₂ is 5 amino acid residues in length;
- L₃ is 1 amino acid residues in length; and
- L₄ is 2 amino acid residues in length.

5. An antibody-like binding protein comprising four polypeptide chains that form four antigen binding sites, wherein two polypeptide chains have a structure represented by the formula:



and two polypeptide chains have a structure represented by the formula:



wherein:


V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain;
 Fc is the immunoglobulin hinge region and C_{H2} , C_{H3} immunoglobulin heavy chain constant domains; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

wherein:


L_1 is 1 to 3 amino acid residues in length;
 L_2 is 1 to 4 amino acid residues in length;
 L_3 is 2 to 15 amino acid residues in length; and
 L_4 is 2 to 15 amino acid residues in length;

and wherein the polypeptides of formula I and the polypeptides of formula II form a cross-over light chain-heavy chain pair.

6. An antibody-like binding protein comprising two polypeptide chains that form two antigen binding sites, wherein a first polypeptide chain has a structure represented by the formula:



and a second polypeptide chain has a structure represented by the formula:



wherein:

V_{L1} is a first immunoglobulin light chain variable domain;
 V_{L2} is a second immunoglobulin light chain variable domain;
 V_{H1} is a first immunoglobulin heavy chain variable domain;
 V_{H2} is a second immunoglobulin heavy chain variable domain;
 C_L is an immunoglobulin light chain constant domain;
 C_{H1} is the immunoglobulin C_{H1} heavy chain constant domain; and
 L_1 , L_2 , L_3 , and L_4 are amino acid linkers;

wherein:

L_1 is 1 to 3 amino acid residues in length;
 L_2 is 1 to 4 amino acid residues in length;
 L_3 is 2 to 15 amino acid residues in length; and
 L_4 is 2 to 15 amino acid residues in length;

and wherein the first and second polypeptides form a cross-over light chain-heavy chain pair.

7. The antibody-like binding protein of either claim 5 or 6, wherein:

- 5 L₁ is 1 to 2 amino acid residues in length;
 L₂ is 1 to 2 amino acid residues in length;
 L₃ is 4 to 12 amino acid residues in length; and
 L₄ is 2 to 12 amino acid residues in length.

10 8. The antibody-like binding protein of either claim 5 or 6, wherein:

- L₁ is 1 amino acid residue in length;
 L₂ is 2 amino acid residues in length;
 L₃ is 7 amino acid residues in length; and
 15 L₄ is 5 amino acid residues in length.

9. The antibody-like binding protein of any one of claims 1, 2, 5 or 6, wherein the binding protein is capable of specifically binding one or more antigen targets, wherein, preferably, the one or more antigen targets is selected from the group consisting of B7.1, B7.2, BAFF, BlyS, C3, C5, CCL11 (eotaxin), CCL15 (MIP-1d), CCL17 (TARC), CCL19 (MIP-3b),
 20 CCL2 (MCP-1), CCL20 (MIP-3a), CCL21 (MIP-2), SLC, CCL24 (MIP-2/eotaxin-2), CCL25 (TECK), CCL26 (eotaxin-3), CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (mcp-2), CD3, CD19, CD20, CD24, CD40, CD40L, CD80, CD86, CDH1 (E-cadherin), Chitinase, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CX3CL1 (SCYD1), CXCL12 (SDF1), CXCL13, EGFR, FCER1A, FCER2, HER2, IGF1R, IL-1, IL-12, IL13, IL15, IL17, IL18, IL1A, IL1B, IL1F10, IL1β, IL2, IL4, IL6, IL7, IL8, IL9, IL12/23, IL22, IL23, IL25, IL27, IL35, ITGB4
 25 (b 4 integrin), LEP (leptin), MHC class II, TLR2, TLR4, TLR5, TNF, TNFα, TNFSF4 (OX40 ligand), TNFSF5 (CD40 ligand), Toll-like receptors, TREM1, TSLP, TWEAK, XCR1 (GPR5/CCXCR1), DNGR-1(CLEC91), and HMGB1.

10. The antibody-like binding protein of any one of claims 1, 2, 5 or 6, wherein the binding protein is bispecific and capable of binding two different antigen targets, wherein, preferably, the two different antigen targets are selected from the group consisting of IL4 and IL13, IGF1R and HER2, IGF1R and EGFR, EGFR and HER2, BK and IL13, PDL-1 and CTLA-4, CTLA4 and MHC class II, IL-12 and IL-18, IL-1α and IL-1β, TNFα and IL12/23, TNFα and IL-12p40, TNFα and IL1β, TNFα and IL-23, and IL17 and IL23.

11. The antibody-like binding protein of any one of claims 1, 2, 5 or 6, wherein the binding protein is capable of inhibiting the function of one or more of the antigen targets.

12. The antibody-like binding protein of any one of claims 1, 2, 5 or 6, wherein at least one of the linkers selected from the group consisting of L₁, L₂, L₃, and L₄ contains at least one cysteine residue.

40 13. An isolated nucleic acid molecule comprising a nucleotide sequence encoding the antibody-like binding protein of any one of claims 1, 2, 5 or 6.


14. An expression vector comprising the nucleic acid molecule of claim 13.

45 15. An isolated host cell comprising the nucleic acid molecule of claim 13 or the expression vector of claim 14, wherein, preferably, the host cell is a mammalian cell or an insect cell.


16. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the antibody-like binding protein of any one of claims 1, 2, 5 or 6.

Patentansprüche

55 1. Antikörperähnliches Bindeprotein, das vier Polypeptidketten umfasst, die vier Antigenbindungsstellen bilden, wobei zwei Polypeptidketten eine Struktur aufweisen, die dargestellt wird durch die Formel:



und zwei Polypeptidketten eine Struktur aufweisen, die dargestellt wird durch die Formel:



wobei:


V_{L1} eine erste variable Domäne einer leichten Immunglobulinkette ist;
 V_{L2} eine zweite variable Domäne einer leichten Immunglobulinkette ist;
 V_{H1} eine erste variable Domäne einer schweren Immunglobulinkette ist;
 V_{H2} eine zweite variable Domäne einer schweren Immunglobulinkette ist;
 C_L eine konstante Domäne einer leichten Immunglobulinkette ist;
 C_{H1} die konstante Domäne der schweren Immunglobulin- C_{H1} - Kette ist;
 Fc die Immunglobulin-Gelenkregion ist und C_{H2} , C_{H3} konstante Domänen einer schweren Immunglobulinkette sind; und
 L_1 , L_2 , L_3 und L_4 Aminosäurelinker sind;

wobei:


L_1 3 bis 12 Aminosäurereste lang ist;
 L_2 3 bis 14 Aminosäurereste lang ist;
 L_3 1 bis 8 Aminosäurereste lang ist; und
 L_4 1 bis 3 Aminosäurereste lang ist;

und wobei die Polypeptide der Formel I und die Polypeptide der Formel II ein überkreuztes Paar einer leichten Kette-schweren Kette bilden.

2. Antikörperähnliches Bindeprotein, das zwei Polypeptidketten umfasst, die zwei Antigenbindungsstellen bilden, wobei eine erste Polypeptidkette eine Struktur aufweist, die dargestellt wird durch die Formel:



und eine zweite Polypeptidkette eine Struktur aufweist, die dargestellt wird durch die Formel:



wobei:

V_{L1} eine erste variable Domäne einer leichten Immunglobulinkette ist;
 V_{L2} eine zweite variable Domäne einer leichten Immunglobulinkette ist;
 V_{H1} eine erste variable Domäne einer schweren Immunglobulinkette ist;
 V_{H2} eine zweite variable Domäne einer schweren Immunglobulinkette ist;
 C_L eine konstante Domäne einer leichten Immunglobulinkette ist;
 C_{H1} die konstante Domäne der schweren Immunglobulin- C_{H1} - Kette ist; und
 L_1 , L_2 , L_3 und L_4 Aminosäurelinker sind;

wobei:

L_1 3 bis 12 Aminosäurereste lang ist;
 L_2 3 bis 14 Aminosäurereste lang ist;
 L_3 1 bis 8 Aminosäurereste lang ist; und
 L_4 1 bis 3 Aminosäurereste lang ist;

und wobei die ersten und zweiten Polypeptide ein überkreuztes Paar einer leichten Kette-schweren Kette bildet.

3. Antikörperähnliches Bindeprotein nach entweder Anspruch 1 oder 2, wobei:

5 L₁ 5 bis 10 Aminosäurereste lang ist;
L₂ 5 bis 8 Aminosäurereste lang ist;
L₃ 1 bis 5 Aminosäurereste lang ist; und
L₄ 1 bis 2 Aminosäurereste lang ist.

- 10 4. Antikörperähnliches Bindeprotein nach entweder Anspruch 1 oder 2, wobei:

L₁ 7 Aminosäurereste lang ist
L₂ 5 Aminosäurereste lang ist;
L₃ 1 Aminosäurerest lang ist; und
15 L₄ 2 Aminosäurereste lang ist.

5. Antikörperähnliches Bindeprotein, das vier Polypeptidketten umfasst, die vier Antigenbindungsstellen bilden, wobei zwei Polypeptidketten eine Struktur aufweisen, die dargestellt wird durch die Formel:

20
$$V_{L1}-L_1-V_{L2}-L_2-C_L \quad [I]$$

und zwei Polypeptidketten eine Struktur aufweisen, die dargestellt wird durch die Formel:

25
$$V_{H2}-L_3-V_{H1}-L_4-C_{H1}-Fc \quad [II]$$

wobei:

30 V_{L1} eine erste variable Domäne einer leichten Immunglobulinkette ist;
V_{L2} eine zweite variable Domäne einer leichten Immunglobulinkette ist;
V_{H1} eine erste variable Domäne einer schweren Immunglobulinkette ist;
V_{H2} eine zweite variable Domäne einer schweren Immunglobulinkette ist;
C_L eine konstante Domäne einer leichten Immunglobulinkette ist;
35 C_{H1} die konstante Domäne der schweren Immunglobulin- C_{H1} - Kette ist;
Fc die Immunglobulin-Gelenkregion ist und C_{H2}, C_{H3} konstante Domänen einer schweren Immunglobulinkette sind; und
L₁, L₂, L₃ und L₄ Aminosäurelinker sind;

40 wobei:


L₁ 1 bis 3 Aminosäurereste lang ist;
L₂ 1 bis 4 Aminosäurereste lang ist;
L₃ 2 bis 15 Aminosäurereste lang ist; und
45 L₄ 2 bis 15 Aminosäurereste lang ist;

und wobei die Polypeptide der Formel I und die Polypeptide der Formel II ein überkreuztes Paar einer leichten Kette-schweren Kette bilden.

- 50 6. Antikörperähnliches Bindeprotein, das zwei Polypeptidketten umfasst, die zwei Antigenbindungsstellen bilden, wobei eine erste Polypeptidkette eine Struktur aufweist, die dargestellt wird durch die Formel:

55
$$V_{L1}-L_1-V_{L2}-L_2-C_L \quad [I]$$

und eine zweite Polypeptidkette eine Struktur aufweist, die dargestellt wird durch die Formel:



wobei:

V_{L1} eine erste variable Domäne einer leichten Immunglobulinkette ist;
 V_{L2} eine zweite variable Domäne einer leichten Immunglobulinkette ist;
 V_{H1} eine erste variable Domäne einer schweren Immunglobulinkette ist;
 V_{H2} eine zweite variable Domäne einer schweren Immunglobulinkette ist;
 C_L eine konstante Domäne einer leichten Immunglobulinkette ist;
 C_{H1} die konstante Domäne der schweren Immunglobulin- C_{H1} - Kette ist; und
 L_1 , L_2 , L_3 und L_4 Aminosäurelinker sind;

wobei:

L_1 1 bis 3 Aminosäurereste lang ist;
 L_2 1 bis 4 Aminosäurereste lang ist;
 L_3 2 bis 15 Aminosäurereste lang ist; und
 L_4 2 bis 15 Aminosäurereste lang ist;

und wobei die ersten und zweiten Polypeptide ein überkreuztes Paar einer leichten Kette-schweren Kette bildet.

7. Antikörperähnliches Bindeprotein nach entweder Anspruch 5 oder 6, wobei:

L_1 1 bis 2 Aminosäurereste lang ist;
 L_2 1 bis 2 Aminosäurereste lang ist;
 L_3 4 bis 12 Aminosäurereste lang ist; und
 L_4 2 bis 12 Aminosäurereste lang ist.

8. Antikörperähnliches Bindeprotein nach entweder Anspruch 5 oder 6, wobei:

L_1 1 Aminosäurerest lang ist;
 L_2 2 Aminosäurereste lang ist;
 L_3 7 Aminosäurereste lang ist; und
 L_4 5 Aminosäurereste lang ist.

9. Antikörperähnliches Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6, wobei das Bindeprotein fähig ist, speziell ein oder mehrere Antigenziele zu binden, wobei, vorzugsweise das eine oder die mehreren Antigenziele ausgewählt sind aus der Gruppe bestehend aus B7.1, B7.2, BAFF, BlyS, C3, C5, CCL11 (Eotaxin), CCL15 (MIP-Id), CCL17 (TARC), CCL19 (MIP-3b), CCL2 (MCP-1), CCL20 (MIP-3a), CCL21 (MIP-2), SLC, CCL24 (MPIF-2/Eotaxin-2), CCL25 (TECK), CCL26 (Eotaxin-3), CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (mcp-2), CD3, CD 19, CD20, CD24, CD40, CD40L, CD80, CD86, CDH1 (E-Cadherin), Chitinase, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CX3CL1 (SCYD1), CXCL12 (SDF1), CXCL13, EGFR, FCER1A, FCER2, HER2, IGF1R, IL-1, IL-12, IL13, IL15, IL17, IL18, IL1A, IL1B, IL1F10, IL1 β , IL2, IL4, IL6, IL7, IL8, IL9, IL12/23, IL22, IL23, IL25, IL27, IL35, ITGB4 (b 4 Integrin), LEP (Leptin), MHC Klasse II, TLR2, TLR4, TLR5, TNF, TNF α , TNFSF4 (OX40 Ligand), TNFSF5 (CD40 Ligand), Toll-ähnlichen Rezeptoren, TREM1, TSLP, TWEAK, XCR1 (GPR5/CCXCR1), DNGR-1(CLEC91) und HMGB1.


10. Antikörperähnliches Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6, wobei das Bindeprotein bispezifisch und fähig ist, zwei unterschiedliche Antigenziele zu binden, wobei vorzugsweise die zwei unterschiedlichen Antigenziele ausgewählt sind aus der Gruppe bestehend aus IL4 und IL13, IGF1R und HER2, IGF1R und EGFR, EGFR und HER2, BK und IL13, PDL-1 und CTLA-4, CTLA4 und MHC Klasse II, IL-12 und IL-18, IL-1 α und IL-1 β , TNF α und IL12/23, TNF α und IL-12p40, TNF α und IL1 β , TNF α und IL-23 und IL17 und IL23.

11. Antikörperähnliches Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6, wobei das Bindeprotein fähig ist, die Funktion von einem oder mehreren von den Antigenzielen zu hemmen.


12. Antikörperähnliches Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6, wobei mindestens einer von den Linkern, der ausgewählt ist aus der Gruppe bestehend aus L₁, L₂, L₃, und L₄, mindestens einen Cysteinrest enthält.
13. Isoliertes Nukleinsäuremolekül, das eine Nukleotidsequenz umfasst, die das antikörperähnliche Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6 codiert.
14. Expressionsvektor, der das Nukleinsäuremolekül nach Anspruch 13 umfasst.
15. Isolierte Wirtszelle, die das Nukleinsäuremolekül nach Anspruch 13 oder den Expressionsvektor nach Anspruch 14 umfasst, wobei vorzugsweise die Wirtszelle eine Säugetierzelle oder eine Insektenzelle ist.
16. Pharmazeutische Zusammensetzung, die einen pharmazeutisch verträglichen Träger und einen therapeutisch wirksamen Betrag an antikörperähnlichem Bindeprotein nach irgendeinem der Ansprüche 1, 2, 5 oder 6 umfasst.

Revendications

1. Protéine de liaison de type anticorps comprenant quatre chaînes polypeptide qui forment quatre sites de liaison antigènes, dans laquelle deux chaînes polypeptide ont une structure représentée par la formule :



et deux chaînes polypeptidiques ont une structure représentée par la formule :



dans laquelle:


V_{L1} est un premier domaine variable d'une chaîne légère d'immunoglobine ;
V_{L2} est un second domaine variable d'une chaîne légère d'immunoglobine ;
V_{H1} est un premier domaine variable d'une chaîne lourde d'immunoglobine ;
V_{H2} est un second domaine variable d'une chaîne lourde d'immunoglobine ;
C_L est un domaine constant d'une chaîne légère d'immunoglobine ;
C_{H1} est le domaine constant d'une chaîne lourde CH1 d'immunoglobine ;
Fc est the zone charnière d'immunoglobine et des domaines constants d'une chaîne lourde d'immunoglobine CH2, CH3 ; et
L₁, L₂, L₃, et L₄ sont des agents de liaison d'acides aminés ;

dans laquelle :


L₁ est de 3 à 12 résidus d'acides aminés en longueur ;
L₂ est de 3 à 14 résidus d'acides aminés en longueur ;
L₃ est de 1 à 8 résidus d'acides aminés en longueur ; et
L₄ est de 1 à 3 résidus d'acides aminés en longueur ;

et dans laquelle les polypeptides de la formule I et les polypeptides de la formule II forment une paire de chaînes lourde et chaîne légère croisée.

2. Protéine de liaison de type anticorps comprenant deux chaînes polypeptidiques qui forment deux sites de liaison antigènes, dans laquelle une première chaîne polypeptidiques a une structure représentée par la formule :



et une seconde chaîne polypeptidique a une structure représentée par la formule :



dans laquelle :

V_{L1} est un premier domaine variable d'une chaîne légère d'immunoglobine ;
 V_{L2} est un second domaine variable d'une chaîne légère d'immunoglobine ;
 V_{H1} est un premier domaine variable d'une chaîne lourde d'immunoglobine ;
 V_{H2} est un second domaine variable d'une chaîne lourde d'immunoglobine ;
 C_L est un domaine constant d'une chaîne légère d'immunoglobine ;
 C_{H1} est le domaine constant d'une chaîne lourde d'immunoglobuline CH1 ; et
 L_1 , L_2 , L_3 , et L_4 sont des agents de liaison d'acides aminés ;

dans laquelle :

L_1 est de 3 à 12 résidus d'acides aminés en longueur ;
 L_2 est de 3 à 14 résidus d'acides aminés en longueur ;
 L_3 est de 1 à 8 résidus d'acides aminés en longueur ; et
 L_4 est de 1 à 3 résidus d'acides aminés en longueur ;

et dans laquelle le premier et le second polypeptides forment une paire de chaînes lourde et chaîne légère croisée.


3. Protéine de liaison de type anticorps selon la revendication 1 ou 2, dans laquelle :

L_1 est de 5 à 10 résidus d'acides aminés en longueur ;
 L_2 est de 5 à 8 résidus d'acides aminés en longueur ;
 L_3 est de 1 à 5 résidus d'acides aminés en longueur ; et
 L_4 est de 1 à 2 résidus d'acides aminés en longueur.


4. Protéine de liaison de type anticorps selon la revendication 1 ou 2, dans laquelle :

L_1 est de 7 résidus d'acides aminés en longueur ;
 L_2 est de 5 résidus d'acides aminés en longueur ;
 L_3 est de 1 résidus d'acides aminés en longueur ; et
 L_4 est de 2 résidus d'acides aminés en longueur.

5. Protéine de liaison de type anticorps comprenant quatre chaînes polypeptidiques qui forment quatre sites de liaison antigènes, dans laquelle deux chaînes polypeptidiques ont une structure représentée par la formule :



et deux chaînes polypeptidiques ont une structure représentée par la formule :



dans laquelle :

V_{L1} est un premier domaine variable d'une chaîne légère d'immunoglobine ;
 V_{L2} est un second domaine variable d'une chaîne légère d'immunoglobine ;
 V_{H1} est un premier domaine variable d'une chaîne lourde d'immunoglobine ;
 V_{H2} est un second domaine variable d'une chaîne lourde d'immunoglobine ;
 C_L est un domaine constant d'une chaîne légère d'immunoglobine ;
 C_{H1} est le domaine constant d'une chaîne lourde d'immunoglobine CH1 ;
 Fc est la zone charnière d'immunoglobine et des domaines constants d'une chaîne lourde d'immunoglobine CH2, CH3 ; et


L_1 , L_2 , L_3 et L_4 sont des agents de liaison d'acides aminés ;

dans laquelle :


- 5 L_1 est de 1 à 3 résidus d'acides aminés en longueur ;
 L_2 est de 1 à 4 résidus d'acides aminés en longueur ;
 L_3 est de 2 à 15 résidus d'acides aminés en longueur ; et
 L_4 est de 2 à 15 résidus d'acides aminés en longueur ;

10 et dans laquelle les polypeptides de la formule I et les polypeptides de la formule II forment une paire de chaînes lourde et chaîne légère croisée.

6. Protéine de liaison de type anticorps comprenant deux chaînes polypeptidiques qui forment deux sites de liaison antigènes, dans lesquels une première chaîne polypeptidique a une structure représentée par la formule :



20 et une seconde chaîne polypeptidique a une structure représentée par la formule :



25 dans laquelle :

- V_{L1} est un premier domaine variable d'une chaîne légère d'immunoglobine ;
 V_{L2} est un second domaine variable d'une chaîne légère d'immunoglobine ;
 V_{H1} est un premier domaine variable d'une chaîne lourde d'immunoglobine ;
 V_{H2} est un second domaine variable d'une chaîne lourde d'immunoglobine ;
 C_L est un domaine constant d'une chaîne légère d'immunoglobine ;
 C_{H1} est le domaine constant d'une chaîne lourde d'immunoglobuline CH1 ; et
 L_1 , L_2 , L_3 , et L_4 sont des agents de liaison d'acides aminés ;

35 dans laquelle :

- L_1 est de 1 à 3 résidus d'acides aminés en longueur ;
 L_2 est de 1 à 4 résidus d'acides aminés en longueur ;
 L_3 est de 2 à 15 résidus d'acides aminés en longueur ; et L_4 est de 2 à 15 résidus d'acides aminés en longueur ;

et dans laquelle le premier et le second polypeptides forment une paire de chaînes lourde et chaîne légère croisée.

7. Protéine de liaison de type anticorps selon la revendication 5 ou 6, dans laquelle :

- 45 L_1 est de 1 à 2 résidus d'acides aminés en longueur ;
 L_2 est de 1 à 2 résidus d'acides aminés en longueur ;
 L_3 est de 4 à 12 résidus d'acides aminés en longueur ; et
 L_4 est de 2 à 12 résidus d'acides aminés en longueur.

- 50 8. Protéine de liaison de type anticorps selon la revendication 5 ou 6, dans laquelle :

- L_1 est de 1 résidu d'acide aminé en longueur ;
 L_2 est de 2 résidus d'acides aminés en longueur ;
 L_3 est de 7 résidus d'acides aminés en longueur ; et
 L_4 est de 5 résidus d'acides aminés en longueur.

9. Protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6, dans laquelle la protéine

de liaison est capable de se lier spécifiquement à une ou plusieurs cibles antigènes, dans laquelle, de préférence, la ou plusieurs cibles antigènes est choisie parmi le groupe constitué de B7.1, B7.2, BAFF, BlyS, C3, C5, CCL11 (eotaxine), CCL15 (MIP-1d), CCL17 (TARC), CCL19 (MIP-3b), CCL2 (MCP-1), CCL20 (MIP-3a), CCL21 (MIP-2), SLC, CCL24 (MPIF-2/eotaxine-2), CCL25 (TECK), CCL26 (eotaxine-3), CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (mcp-2), CD3, CD 19, CD20, CD24, CD40, CD40L, CD80, CD86, CDH1 (E-cadherine), Chitinase, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CX3CL1 (SCYD1), CXCL12 (SDF1), CXCL13, EGFR, FCER1A, FCER2, HER2, IGF1R, IL-1, IL-12, IL13, IL15, IL17, IL18, IL1A, IL1B, IL1F10, IL1 β , IL2, IL4, IL6, IL7, IL8, IL9, IL12/23, IL22, IL23, IL25, IL27, IL35, ITGB4 (b 4 intégrine), LEP (leptine), MHC classe II, TLR2, TLR4, TLR5, TNF, TNF α , TNFSF4 (OX40 ligand), TNFSF5 (CD40 ligand), récepteurs Toll-like, TREM1, TSLP, TWEAK, XCR1 (GPR5/CCXCR1), DNGR-1(CLEC91), et HMGB1.

10. Protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6, dans laquelle la protéine de liaison est bispécifique et est capable de se lier à deux cibles antigènes différentes, dans laquelle, de préférence, les deux cibles antigènes différentes sont choisies parmi le groupe constitué de IL4 et IL13, IGF1R et HER2, IGF1R et EGFR, EGFR et HER2, BK et IL13, PDL-1 et CTLA-4, CTLA4 et MHC classe II, IL-12 et IL-18, IL-1 α et IL-1 β , TNF α et IL12/23, TNF α et IL-12p40, TNF α et IL1 β , TNF α et IL-23, et IL17 et IL23.

11. Protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6, dans laquelle la protéine de liaison est capable d'inhiber la fonction d'une ou plusieurs cibles antigènes.

12. Protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6, dans laquelle au moins un des agents de liaison choisi parmi le groupe constitué de L₁, L₂, L₃, et L₄ contient au moins un résidu cystéine.

13. Molécule d'acide nucléique isolée, comprenant une séquence de nucléotides codant la protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6.

14. Vecteur d'expression comprenant la molécule d'acide nucléique selon la revendication 13.

15. Cellule hôte isolée, comprenant la molécule d'acide nucléique selon la revendication 13 ou le vecteur d'expression selon la revendication 14, dans lesquels, de préférence, la cellule hôte est une cellule de mammifère ou une cellule d'insecte.

16. Composition pharmaceutique comprenant un support pharmaceutiquement acceptable et une quantité efficace thérapeutiquement de protéine de liaison de type anticorps selon l'une quelconque des revendications 1, 2, 5 ou 6.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6






FIG. 7




	On-rate [1/Ms]	Off-rate [1/s]	KD [M]
Parental anti-IL4 mAb	2.49E+07	1.96E-04	7.83E-12
Parental anti-IL13 mAb	1.58E+06	1.30E-04	8.18E-11
TB11 (IL4)	6.70E+06	1.50E-04	2.63E-11
TB11 (IL13)	3.28E+06	1.70E-04	5.18E-11



	On-rate [1/Ms]	Off-rate [1/s]	KD [M]
Parental anti-IL4 mAb	2.49E+07	1.96E-04	7.83E-12
Parental anti-IL13 mAb	1.58E+06	1.30E-04	8.18E-11
CODH (IL4)	3.16E+07	2.63E-04	8.14E-12
CODH (IL13)	1.20E+06	1.12E-04	9.34E-11

FIG. 8



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5989830 A [0013] [0019] [0020] [0029] [0117]
- US 20050003403 A1 [0014]
- US 2010331527 A1 [0018]
- US 7612181 B [0019] [0029] [0117] [0120] [0123]
- US 20100226923 A1 [0019] [0030]
- EP 2050764 A [0020]
- WO 2010014659 A [0020]
- US 20100226923 A [0030]
- WO 2009052081 A [0117] [0122] [0123]
- US 2011026 A [0180]

Non-patent literature cited in the description

- **WARD et al.** *Nature*, 1989, vol. 341 (6242), 544-46 [0003]
- **WILLIAMS et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1989, vol. 86 (14), 5537-41 [0003]
- **BIRD et al.** *Science*, 1988, vol. 242 (4877), 423-26 [0004]
- **STAERZ et al.** *Nature*, 1985, vol. 314 (6012), 628-31 [0006]
- **MILSTEIN et al.** *Nature*, 1983, vol. 305 (5934), 537-40 [0007]
- **GEORGE et al.** *THE ANTIBODIES*. Harwood Academic Publishers, 1997, vol. 4, 99-141 [0007]
- **LINDHOFER et al.** *J. Immunol.*, 1995, vol. 155 (1), 219-25 [0007]
- **CARTER et al.** *J. Hematother.*, 1995, vol. 4 (5), 463-70 [0009]
- **PLUCKTHUN et al.** *Immunotechnology*, 1997, vol. 3 (2), 83-105 [0009]
- **TODOROVSKA et al.** *J. Immunol. Methods*, 2001, vol. 248 (1-2), 47-66 [0009]
- **CARTER.** *J. Immunol. Methods*, 2001, vol. 248 (1-2), 7-15 [0009]
- **MACK et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1995, vol. 92 (15), 7021-25 [0010]
- **MALLENDER et al.** *J. Biol. Chem.*, 1994, vol. 269 (1), 199-206 [0010]
- **KOSTELNY et al.** *J. Immunol.*, 1992, vol. 148 (5), 1547-53 [0010]
- **DE KRUIF et al.** *J. Biol. Chem.*, 1996, vol. 271 (13), 7630-34 [0010]
- **HOLLIGER et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1993, vol. 90 (14), 6444-48 [0011]
- **HUDSON et al.** *J. Immunol. Methods*, 1999, vol. 231 (1-2), 177-89 [0011]
- **WÖRN et al.** *J. Mol. Biol.*, 2001, vol. 305 (5), 989-1010 [0012]
- **RIDGWAY et al.** *Protein Eng.*, 1996, vol. 9 (7), 617-21 [0015]
- **CARTER.** *J. Immunol. Methods*, 2011, vol. 248 (1-2), 7-15 [0015]
- **MERCHANT et al.** *Nat. Biotechnol.*, 1998, vol. 16 (7), 677-81 [0015]
- **GUNASEKARAN et al.** *J. Biol. Chem.*, 2010, vol. 285 (25), 19637-46 [0016]
- **DAVIS et al.** *Protein Eng. Des. Sel.*, 2010, vol. 23 (4), 195-202 [0017]
- **SAMBROOK et al.** *MOLECULAR CLONING: A LABORATORY MANUAL*. Cold Spring Harbor Laboratory Press, 2001 [0031]
- *FUNDAMENTAL IMMUNOLOGY*. Raven Press, 1989 [0038]
- *REMINGTON'S PHARMACEUTICAL SCIENCES*. Mack Publishing Company, 1990 [0099]
- **DUROCHER et al.** *Nucl. Acids Res.*, 2002, vol. 30 (2), E9 [0130]
- **WOZNIAK-KNOPP et al.** Stabilisation of the Fc Fragment of Human IgG1 by Engineered Intradomain Disulfide Bonds. *PLoS ONE*, 2012, vol. 7 (1), e30083 [0173]
- **BRINKMANN et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1993, vol. 90, 7538-42 [0175]

KETTÓS VARIÁBILIS RÉGIÓVAL RENDELKEZŐ ELLENANYAG-SZERŰ FEHÉRJÉK KERESZTEZETT KÖTŐRÉGIÓS ORIENTÁCIÓVAL

Szabadalmi igénypontok

1. Ellenanyag-szerű kötőfehérje, amely négy polipeptid-láncot tartalmaz, amelyek négy antigénkötő helyet alkotnak, ahol két polipeptid-lánc szerkezetét a következő képlet szemlélteti:



és két polipeptid-lánc szerkezetét a következő képlet szemlélteti:



ahol:

V_{L1} jelentése egy első immunglobulin könnyűlánc variábilis domén;

V_{L2} jelentése egy második immunglobulin könnyűlánc variábilis domén;

V_{H1} jelentése egy első immunglobulin nehézlánc variábilis domén;

V_{H2} jelentése egy második immunglobulin nehézlánc variábilis domén;

C_L jelentése immunglobulin könnyűlánc konstans domén;

C_{H1} jelentése az immunglobulin C_{H1} nehézlánc konstans domén;

Fc jelentése az immunglobulin csuklórégió és C_{H2} , C_{H3} immunglobulin nehézlánc konstans domének; és

L_1 , L_2 , L_3 és L_4 jelentése aminosav kapcsolómolekulák;

ahol:

L_1 3-12 aminosav-oldallánc hosszúságú;


L_2 3-14 aminosav-oldallánc hosszúságú;

L_3 1-8 aminosav-oldallánc hosszúságú; és

L_4 1-3 aminosav-oldallánc hosszúságú;

és ahol az I képletű polipeptid és a II képletű polipeptidek keresztezett („cross-over”) könnyűlánc-nehézlánc párt alkotnak.

2. Ellenanyag-szerű kötőfehérje, amely két polipeptid-láncot tartalmaz, amelyek két antigénkötő helyet alkotnak, ahol az első polipeptid-lánc szerkezetét a következő képlet szemlélteti:



és a második polipeptid-lánc szerkezetét a következő képlet szemlélteti:



ahol:

V_{L1} jelentése egy első immunglobulin könnyűlánc variábilis domén;

V_{L2} jelentése egy második immunglobulin könnyűlánc variábilis domén;

V_{H1} jelentése egy első immunglobulin nehézlánc variábilis domén;

V_{H2} jelentése egy második immunglobulin nehézlánc variábilis domén;

C_L jelentése immunglobulin könnyűlánc konstans domén;

C_{H1} jelentése az immunglobulin C_{H1} nehézlánc konstans domén; és

L_1 , L_2 , L_3 és L_4 jelentése aminosav kapcsolómolekulák;

ahol:

- L_1 3-12 aminosav-oldallánc hosszúságú;
- L_2 3-14 aminosav-oldallánc hosszúságú;
- L_3 1-8 aminosav-oldallánc hosszúságú; és
- L_4 1-3 aminosav-oldallánc hosszúságú;

és ahol az első és második polipeptid keresztezett könnyűlánc-nehézlánc párt alkotnak.


3. Az 1. vagy 2. igénypont szerinti ellenanyag-szerű kötőfehérje, ahol:

- L_1 5-10 aminosav-oldallánc hosszúságú;
- L_2 5-8 aminosav-oldallánc hosszúságú;
- L_3 1-5 aminosav-oldallánc hosszúságú; és
- L_4 1-2 aminosav-oldallánc hosszúságú.

4. Az 1. vagy 2. igénypont szerinti ellenanyag-szerű kötőfehérje, ahol:

- L_1 7 aminosav-oldallánc hosszúságú;
- L_2 5 aminosav-oldallánc hosszúságú;
- L_3 1 aminosav-oldallánc hosszúságú; és
- L_4 2 aminosav-oldallánc hosszúságú.

5. Ellenanyag-szerű kötőfehérje, amely négy polipeptid-láncot tartalmaz, amelyek négy antigénkötő helyet alkotnak, ahol két polipeptid-lánc szerkezetét a következő képlet szemlélteti:



és két polipeptid-lánc szerkezetét a következő képlet szemlélteti:



ahol:

- V_{L1} jelentése egy első immunglobulin könnyűlánc variábilis domén;
- V_{L2} jelentése egy második immunglobulin könnyűlánc variábilis domén;
- V_{H1} jelentése egy első immunglobulin nehézlánc variábilis domén;
- V_{H2} jelentése egy második immunglobulin nehézlánc variábilis domén;
- C_L jelentése immunglobulin könnyűlánc konstans domén;
- C_{H1} jelentése az immunglobulin C_{H1} nehézlánc konstans domén;
- Fc jelentése az immunglobulin csuklórégió és C_{H2} , C_{H3} immunglobulin nehézlánc konstans domének; és

L_1 , L_2 , L_3 és L_4 jelentése aminosav kapcsolómolekulák;

ahol:

- L_1 1-3 aminosav-oldallánc hosszúságú;
- L_2 1-4 aminosav-oldallánc hosszúságú;
- L_3 2-15 aminosav-oldallánc hosszúságú; és
- L_4 2-15 aminosav-oldallánc hosszúságú;

és ahol az I képletű polipeptid és a II képletű polipeptidek keresztezett könnyűlánc-nehézlánc párt alkotnak.

6. Ellenanyag-szerű kötőfehérje, amely két polipeptid-láncot tartalmaz, amelyek két antigénkötő helyet alkotnak, ahol az első polipeptid-lánc szerkezetét a következő képlet szemlélteti:



és a második polipeptid-lánc szerkezetét a következő képlet szemlélteti:



ahol:

- V_{L1} jelentése egy első immunglobulin könnyűlánc variábilis domén;
- V_{L2} jelentése egy második immunglobulin könnyűlánc variábilis domén;
- V_{H1} jelentése egy első immunglobulin nehézlánc variábilis domén;
- V_{H2} jelentése egy második immunglobulin nehézlánc variábilis domén;
- C_L jelentése immunglobulin könnyűlánc konstans domén;
- C_{H1} jelentése az immunglobulin C_{H1} nehézlánc konstans domén; és
- L_1, L_2, L_3 és L_4 jelentése aminosav kapcsolómolekulák;

ahol:

- L_1 1-3 aminosav-oldallánc hosszúságú;
- L_2 1-4 aminosav-oldallánc hosszúságú;
- L_3 2-15 aminosav-oldallánc hosszúságú; és
- L_4 2-15 aminosav-oldallánc hosszúságú;

és ahol az első és második polipeptid keresztezett könnyűlánc-nehézlánc párt alkotnak.

7. Az 5. vagy 6. igénypont szerinti ellenanyag-szerű kötőfehérje, ahol:

- L_1 1-2 aminosav-oldallánc hosszúságú;
- L_2 1-2 aminosav-oldallánc hosszúságú;
- L_3 4-12 aminosav-oldallánc hosszúságú; és
- L_4 2-12 aminosav-oldallánc hosszúságú.

8. Az 5. vagy 6. igénypont szerinti ellenanyag-szerű kötőfehérje, ahol:

- L_1 1 aminosav-oldallánc hosszúságú;
- L_2 2 aminosav-oldallánc hosszúságú;
- L_3 7 aminosav-oldallánc hosszúságú; és
- L_4 5 aminosav-oldallánc hosszúságú.

9. Az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérje, ahol a kötőfehérje képes specifikusan kötődni egy vagy több antigén-célponthoz, ahol előnyösen az egy vagy több antigén-célpont a következők által alkotott csoportból van kiválasztva: B7.1, B7.2, BAFF, BlyS, C3, C5, CCL11 (eotaxin), CCL15 (MIP-1d), CCL17 (TARC), CCL19 (MIP-3b), CCL2 (MCP-1), CCL20 (MIP-3a), CCL21 (MIP-2), SLC, CCL24 (MIP-2/eotaxin-2), CCL25 (TECK), CCL26 (eotaxin-3), CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (mcp-2), CD3, CD19, CD20, CD24, CD40, CD40L, CD80, CD86, CDH1 (E-kadherin), kitináz, CSF1 (M-CSF), CSF2 (GM-CSF), CSF3 (G-CSF), CTLA4, CX3CL1 (SCYD1), CXCL12 (SDF1), CXCL13, EGFR, FCER1A, FCER2, HER2, IGF1R, IL-1, IL-12, IL13, IL15, IL17, IL18, IL1A, IL1B, IL1F10, IL1β, IL2, IL4, IL6, IL7, IL8, IL9, IL12/23, IL22, IL23, IL25, IL27, IL35, ITGB4 (b 4 integrin), LEP (leptin), MHC II. osztály, TLR2, TLR4, TLR5, TNF, TNFα, TNFSF4 (OX40 ligandum), TNFSF5 (CD40 ligandum), Toll-szerű receptorok, TREM1, TSLP, TWEAK, XCR1 (GPR5/CCXCR1), DNCR-1 (CLEC91) és HMGB1.

10. Az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérje, ahol a kötőfehérje bispecifikus és képes kötődni két különböző antigén-célponthoz, ahol előnyösen a két különböző antigén-célpont a következők által alkotott csoportból van kiválasztva: IL4 és IL13, IGF1R és HER2, IGF1R és EGFR, EGFR és HER2, BK és IL13, PDL-1 és CTLA-4, CTLA4 és MHC II. osztály, IL-12 és IL-18, IL-1 α és IL-1 β , TNF α és IL12/23, TNF α és IL-12p40, TNF α és IL1 β , TNF α és IL-23 és IL17 és IL23.

11. Az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérje, ahol a kötőfehérje képes gátolni az egy vagy több antigén-célpont működését.

12. Az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérje, ahol a L₁, L₂, L₃ és L₄ által alkotott csoportból kiválasztott kapcsolómolekulák legalább egyike legalább egy cisztein-oldalláncot tartalmaz.

13. Izolált nukleinsav-molekula, amely az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérjét kódoló nukleotid-szekvenciát tartalmaz.

14. Expressziós vektor, amely 13. igénypont szerinti nukleinsav-molekulát tartalmaz.

15. Izolált gazdasejt, amely 13. igénypont szerinti nukleinsav-molekulát vagy 14. igénypont szerinti expressziós vektort tartalmaz, ahol előnyösen a gazdasejt emlőssejt vagy rovarsejt.

16. Gyógyászati készítmény, amely gyógyászatilag elfogadható hordozót és az 1., 2., 5. és 6. igénypontok bármelyike szerinti ellenanyag-szerű kötőfehérje terápiásan hatásos mennyiségét tartalmazza.