



(51) International Patent Classification:

C07D 471/04 (2006.01) *A61P 31/04* (2006.01)
A61K 31/437 (2006.01) *C07D 487/04* (2006.01)

(21) International Application Number:

PCT/IB2018/052649

(22) International Filing Date:

17 April 2018 (17.04.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

201721013819 18 April 2017 (18.04.2017) IN

(71) Applicant: **WOCKHARDT LIMITED** [IN/IN]; D-4, MIDC Area, Chikalthana, Aurangabad 431006 (IN).

(72) Inventors: **BHAGWAT, Sachin Subhash**; Row House No.C-2, "Nilgiris", Khivansara Park(E), Behind Ramayana Cultural Hall, Ulkanagari, Aurangabad-431005, Maharashtra, India Aurangabad 431005 (IN). **JOSHI, Prashant Ratanakar**; C/o R.S. Joshi, Audit Nagar, Old Pedgaon Road, Parbhani-431401, Maharashtra, India, Parbhani 431401 (IN). **UMARKAR, Kushal**; C/o V.R.Umarkar, Ramgitai, Behind Waghmadi, Malipura, Akola-444001, Maharashtra, India, Akola 444001 (IN). **SATAV, Jaykumar Satwaji**; C/o S.N.Satav, Mali Galli, Kalamnuri, Tal – Kalamnuri, Hingoli-431702, Maharashtra, India, Hingoli 431702 (IN). **TAKALKAR, Swapna Shripad**; "Purvesh Apartment", Flat No 3, Plot No 45, Khivansara Park, Opposite Ganesh Temple, Ulka Nagari, Garkheda Parisar, Aurangabad -431005 Maharashtra, India Aurangabad 431005 (IN). **PATIL, Vijaykumar Jagdishwar**; Prem Nagar-2, Near Bharati Vidhyapeeth, Solapur-413004, Maharashtra, India, Solapur 413004 (IN). **TADIPARTHI, Ravikumar**; Door no: 26-3-95, NRP Road, Gandhinagar, Vijayawada-520003, Andhra Pradesh, India, Vijaywada 520003 (IN). **PATEL, Mahesh Vithalbhaji**; Plot No.157, Opp. Saint Meera School, N-3, CIDCO, Aurangabad-431003, Maharashtra, India, Aurangabad 431003 (IN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):

AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: ANTIBACTERIAL COMPOSITIONS



(57) Abstract: Pharmaceutical compositions comprising meropenem or a pharmaceutically acceptable derivative, and a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof are disclosed. They are useful for treating or preventing bacterial infection.



ANTIBACTERIAL COMPOSITIONS

RELATED PATENT APPLICATIONS

This application claims the priority to and benefit of Indian Provisional Patent Application No. 201721013819 filed on April 18, 2017; the disclosures of which are incorporated herein by reference in its entirety as if fully rewritten herein.

FIELD OF THE INVENTION

The invention relates to antibacterial compositions and methods for treating or preventing bacterial infections.

BACKGROUND OF THE INVENTION

Bacterial infections continue to remain one of the major causes contributing towards human diseases. One of the key challenges in treatment of bacterial infections is the ability of bacteria to develop resistance to one or more antibacterial agents over time. Examples of such bacteria that have developed resistance to typical antibacterial agents include: Penicillin-resistant *Streptococcus pneumoniae*, Vancomycin-resistant *Enterococci*, and Methicillin-resistant *Staphylococcus aureus*. The problem of emerging drug-resistance in bacteria is often tackled by switching to newer antibacterial agents, which can be more expensive and sometimes more toxic. Additionally, this may not be a permanent solution as the bacteria often develop resistance to the newer antibacterial agents as well in due course. In general, bacteria are particularly efficient in developing resistance, because of their ability to multiply very rapidly and pass on the resistance genes as they replicate.


Treatment of infections caused by resistant bacteria remains a key challenge for the clinician community. One example of such challenging pathogen is *Acinetobacter baumannii* (*A. baumannii*), which continues to be an increasingly important and demanding species in healthcare settings. The multidrug resistant nature of this pathogen and its unpredictable susceptibility patterns make empirical and therapeutic decisions more difficult. *A. baumannii* is associated with infections such as pneumonia, bacteremia, wound infections, urinary tract infections and meningitis.

Therefore, there is a need for development of newer ways to treat infections that are becoming resistant to known therapies and methods. Surprisingly, it has been found that a

compositions comprising meropenem and certain nitrogen containing bicyclic compounds (disclosed in PCT/IB2012/054706) exhibit unexpectedly synergistic antibacterial activity, even against highly resistant bacterial strains.

SUMMARY OF THE INVENTION

Accordingly, there are provided pharmaceutical compositions comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



In one general aspect, there are provided pharmaceutical compositions comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof; wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

In yet another general aspect, there are provided methods for treating or preventing a bacterial infection in a subject, said methods comprising administering to said subject an effective amount of a pharmaceutical composition comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof; and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof.

In another general aspect, there are provided methods for treating or preventing a bacterial infection in a subject, said methods comprising administering to said subject an effective amount of a pharmaceutical composition comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof; wherein a compound of

Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

In yet another general aspect, there are provided methods for treating or preventing a bacterial infection in a subject, said methods comprising administering to said subject an effective amount of: (a) meropenem or a pharmaceutically acceptable derivative thereof; and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof.

In another general aspect, there are provided methods for treating or preventing a bacterial infection in a subject, said methods comprising administering to said subject an effective amount of: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof; wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is administered in an amount from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

The details of one or more embodiments of the invention are set forth in the description below. Other features, objects and advantages of the invention will be apparent from the following description including claims.

DETAILED DESCRIPTION OF THE INVENTION

Reference will now be made to the exemplary embodiments, and specific language will be used herein to describe the same. It should nevertheless be understood that no limitation of the scope of the invention is thereby intended. Alterations and further modifications of the inventive features illustrated herein, which would occur to one skilled in the relevant art and having possession of this disclosure, are to be considered within the scope of the invention. It must be noted that, as used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the content clearly dictates otherwise. All references including patents, patent applications, and literature cited in the specification are expressly incorporated herein by reference in their entirety as if fully rewritten herein.

The inventors have surprisingly discovered that a pharmaceutical composition comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof exhibits unexpectedly improved antibacterial efficacy, even against highly resistant bacteria, including those producing extended spectrum beta-lactamase enzymes (ESBLs).

The term “infection” or “bacterial infection” as used herein includes presence of bacteria, in or on a subject, which, if its growth were inhibited, would result in a benefit to the subject. As such, the term “infection” in addition to referring to the presence of bacteria also refers to presence of other floras, which are not desirable. The term “infection” includes infection caused by bacteria.

The term “treat”, “treating” or “treatment” as used herein refers to administration of a medicament, including a pharmaceutical composition, or one or more pharmaceutically active ingredients, for prophylactic and/or therapeutic purposes. The term “prophylactic treatment” refers to treating a subject who is not yet infected, but who is susceptible to, or otherwise at a risk of infection (preventing the bacterial infection). The term “therapeutic treatment” refers to administering treatment to a subject already suffering from infection. The terms “treat”, “treating” or “treatment” as used herein also refer to administering compositions, or one or more of pharmaceutically active ingredients discussed herein, with or without additional pharmaceutically active or inert ingredients, in order to: (i) reduce or eliminate either a bacterial infection, or one or more symptoms of a bacterial infection, or (ii) retard progression of a bacterial infection, or one or more symptoms of a bacterial infection, or (iii) reduce severity of a bacterial infection, or one or more symptoms of a bacterial infection, or (iv) suppress clinical manifestation of a bacterial infection, or (v) suppress manifestation of adverse symptoms of a bacterial infection.

The terms “pharmaceutically effective amount” or “therapeutically effective amount” or “effective amount” as used herein refer to an amount, which has a therapeutic effect or is the amount required to produce a therapeutic effect in a subject. For example, a “therapeutically effective amount” or “pharmaceutically effective amount” or “effective amount” of an antibacterial agent or a pharmaceutical composition is the amount of the antibacterial agent or the pharmaceutical composition required to produce a desired therapeutic effect as may be judged by clinical trial results, model animal infection studies, and/or in vitro studies (e.g. in agar or broth media). Such effective amount depends on several factors, including but not limited to,

the microorganism (e.g. bacteria) involved, characteristics of the subject (for example height, weight, sex, age and medical history), severity of infection and particular type of the antibacterial agent used. For prophylactic treatments, a prophylactically effective amount is that amount which would be effective in preventing the bacterial infection.

The term “administration” or “administering” refers to and includes delivery of a composition, or one or more pharmaceutically active ingredients to a subject, including for example, by any appropriate method, which serves to deliver the composition or its active ingredients or other pharmaceutically active ingredients to the site of infection. The method of administration may vary depending on various factors, such as for example, the components of the pharmaceutical composition or type/nature of the pharmaceutically active or inert ingredients, site of the potential or actual infection, the microorganism involved, severity of the infection, age and physical condition of the subject and a like. Some non-limiting examples of ways to administer a composition or a pharmaceutically active ingredient to a subject according to this invention include oral, intravenous, topical, intraspiratory, intraperitoneal, intramuscular, parenteral, sublingual, transdermal, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, gene gun, dermal patch, eye drop and mouthwash. In case of a pharmaceutical composition comprising more than one ingredients (active or inert), one of the ways of administering such composition is by admixing the ingredients (e.g. in the form of a suitable unit dosage form such as tablet, capsule, solution, powder or a like) and then administering the dosage form. Alternatively, the ingredients may also be administered separately (simultaneously or one after the other) as long as these ingredients reach beneficial therapeutic levels such that the composition as a whole provides a synergistic and/or desired effect.

The term “growth” as used herein refers to a growth of one or more microorganisms and includes reproduction or population expansion of the microorganism (e.g. bacteria). The term “growth” also includes maintenance of on-going metabolic processes of the microorganism, including the processes that keep the microorganism alive.

The term, “effectiveness” as used herein refers to ability of a treatment, or a composition, or one or more pharmaceutically active ingredients to produce a desired biological effect in a subject. For example, the term “antibacterial effectiveness” of a composition or of an antibacterial agent refers to the ability of the composition or the antibacterial agent to prevent or treat bacterial infection in a subject.

The term “synergistic” or “synergy” as used herein refers to the interaction of two or more agents so that their combined effect is greater than their individual effects.

The term “antibacterial agent” as used herein refers to any substance, compound, a combination of substances, or a combination of compounds capable of: (i) inhibiting, reducing or preventing growth of bacteria; (ii) inhibiting or reducing ability of a bacteria to produce infection in a subject; or (iii) inhibiting or reducing ability of bacteria to multiply or remain infective in the environment. The term “antibacterial agent” also refers to compounds capable of decreasing infectivity or virulence of bacteria.

The term “beta-lactam antibacterial agent” as used herein refers to compounds with antibacterial properties and containing a beta-lactam nucleus in their molecular structure.

The term “beta-lactamase” or “beta-lactamase enzyme” as used herein refers to any enzyme or protein or any other substance that breaks down a beta-lactam ring. The term “beta-lactamase” includes enzymes that are produced by bacteria and have the ability to hydrolyse the beta-lactam ring in a beta-lactam compound, either partially or completely.

The term “extended spectrum beta-lactamase” (ESBL) as used herein includes those beta-lactamase enzymes, which are capable of conferring bacterial resistance to various beta-lactam antibacterial agents such as penicillins, cephalosporins, aztreonam and the like.

The term “beta-lactamase inhibitor” as used herein refers to a compound capable of inhibiting activity of one or more beta-lactamase enzymes, either partially or completely.

The term “colony forming units” or “CFU” as used herein refers to an estimate of number of viable bacterial cells per ml of the sample. Typically, a “colony of bacteria” refers to a mass of individual bacteria growing together.

The term “pharmaceutically inert ingredient” or “carrier” or “excipient” refers to and includes compounds or materials used to facilitate administration of a compound, for example, to increase the solubility of the compound. Typical, non-limiting examples of solid carriers include starch, lactose, dicalcium phosphate, sucrose, and kaolin. Typical, non-limiting examples of liquid carriers include sterile water, saline, buffers, non-ionic surfactants, and edible oils. In

addition, various adjuvants commonly used in the art may also be included. These and other such compounds are described in literature, e.g., in the Merck Index (Merck & Company, Rahway, N.J.). Considerations for inclusion of various components in pharmaceutical compositions are described, e.g., in Gilman et al. (Goodman and Gilman's: The Pharmacological Basis of Therapeutics, 8th Ed., Pergamon Press., 1990), which is incorporated herein by reference in its entirety.

The term "subject" as used herein refers to vertebrate or invertebrate, including a mammal. The term "subject" includes human, animal, a bird, a fish, or an amphibian. Typical, non-limiting examples of a "subject" include humans, cats, dogs, horses, sheep, bovine cows, pigs, lambs, rats, mice and guinea pigs.

The term "pharmaceutically acceptable derivative" as used herein refers to and includes any pharmaceutically acceptable salt, pro-drug, metabolite, ester, ether, hydrate, polymorph, solvate, complex, and adduct of a compound described herein which, upon administration to a subject, is capable of providing (directly or indirectly) the parent compound. For example, the term "antibacterial agent or a pharmaceutically acceptable derivative thereof" includes all derivatives of the antibacterial agent (such as salts, pro-drugs, metabolites, esters, ethers, hydrates, polymorphs, solvates, complexes, and adducts) which, upon administration to a subject, are capable of providing (directly or indirectly) the antibacterial agent.


The term "pharmaceutically acceptable salt" as used herein refers to one or more salts of a given compound which possesses desired pharmacological activity of the free compound and which is neither biologically nor otherwise undesirable. In general, the term "pharmaceutically acceptable salts" refer to salts that are suitable for use in contact with the tissues of human and animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, et al. (*J. Pharmaceutical Sciences*, 66; 1-19, 1977), incorporated herein by reference in its entirety, describes various pharmaceutically acceptable salts in details.

The term "stereoisomer" as used herein refers to and includes isomeric molecules that have the same molecular formula but differ in positioning of atoms and/or functional groups in the space. Stereoisomers may further be classified as enantiomers (where different isomers are mirror-images of each other) and diastereomers (where different isomers are not mirror-images

of each other). Diastereomers include isomers such as conformers, meso compounds, cis-trans (E-Z) isomers, and non-enantiomeric optical isomers.

A person of skills in the art would appreciate that various compounds described herein (including, for example a compound of Formula (I) and meropenem) can exist and are often used as their pharmaceutically acceptable derivatives (such as salts, pro-drugs, metabolites, esters, ethers, hydrates, polymorphs, solvates, complexes, and adducts).

In one general aspect, there are provided pharmaceutical compositions comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



Compound of Formula (I), according to the invention can be used in various forms including as such, a stereoisomer or a pharmaceutically acceptable derivative thereof. A compound of Formula (I) (CAS Registry Number: 1427462-70-1) may also be known chemically by different names including the following: (a) “trans-7-oxo-6-(sulphooxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”; (b) “(2S, 5R)-7-oxo-6-(sulphooxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”; or (c) “Sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”. A reference to “a compound of Formula (I)” is intended to include compounds chemically known as: (a) “trans-7-oxo-6-(sulphooxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”; (b) “(2S, 5R)-7-oxo-6-(sulphooxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”; or (c) “Sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”.

Compound of Formula (I) may also be used in the form of its stereoisomer or a pharmaceutically acceptable derivative thereof. Typical, non-limiting examples of suitable pharmaceutically acceptable derivatives of a compound of Formula (I) include its sodium salt (also known as “sodium salt of sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-

diazabicyclo[3.2.1]oct-6-yl] ester” or “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester, sodium salt (1:1); CAS Registry Number: 1427462-59-6”); potassium salt (also known as “potassium salt of sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester” or “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester, potassium salt (1:1); CAS Registry Number: 1427462-60-9”); and other salts such as “1-Butanaminium, N,N,N-tributyl-, (1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl sulphate (1:1); CAS Registry Number: 1427462-72-3”.

In another general aspect, there are provided pharmaceutical compositions comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof; wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

Both, meropenem and a compound of Formula (I) may be present in the composition in their free forms or in the form of their pharmaceutically acceptable derivatives (such as salts, pro-drugs, metabolites, esters, ethers, hydrates, polymorphs, solvates, complexes, or adducts).

Individual amounts of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or pharmaceutically acceptable derivative thereof in the composition may vary depending on clinical requirements. In some embodiments, a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof in the composition is present in an amount from about 0.01 gram to about 10 gram. In some other embodiments, meropenem or a pharmaceutically acceptable derivative thereof in the composition is present in an amount from about 0.01 gram to about 10 gram.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.25 gram of a compound of Formula (I) or a stereoisomer or a

pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some other embodiments, the pharmaceutical composition according to the invention comprises about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some other embodiments, the pharmaceutical composition according to the invention comprises about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some other embodiments, the pharmaceutical composition according to the invention comprises about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically

acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.250 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically

acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, the pharmaceutical composition according to the invention comprises about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

The pharmaceutical compositions according to the invention may include one or more pharmaceutically acceptable carriers or excipients or the like. Typical, non-limiting examples of such carriers or excipients include mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, sodium crosscarmellose, glucose, gelatine, sucrose, magnesium carbonate, wetting agents, emulsifying agents, solubilizing agents, buffering agents, lubricants, preservatives, stabilizing agents, binding agents and the like.

The pharmaceutical compositions or the active ingredients according to the present invention may be formulated into a variety of dosage forms, such as solid, semi-solid, liquid and aerosol dosage forms. Typical, non-limiting examples of some dosage forms include tablets, capsules, powders, solutions, suspensions, suppositories, aerosols, granules, emulsions, syrups, elixirs and the like.

In some embodiments, pharmaceutical compositions according to the invention are in the form of a powder or a solution. In some other embodiments, pharmaceutical compositions according to the invention are present in the form of a powder or a solution that can be reconstituted by addition of a compatible reconstitution diluent prior to administration. In some other embodiments, pharmaceutical compositions according to the invention are in the form of a frozen composition that can be diluted with a compatible reconstitution diluent prior to administration. Typical, non-limiting example of suitable compatible reconstitution diluent includes water.

In some other embodiments, pharmaceutical compositions according to the invention are present in the form ready to use for parenteral administration.

The compositions according to the invention can be formulated into various dosage forms wherein the active ingredients and/or excipients may be present either together (e.g. as an

admixture) or as separate components. When the various ingredients in the composition are formulated as a mixture, such compositions can be delivered by administering such a mixture to a subject using any suitable route of administration. Alternatively, pharmaceutical compositions according to the invention may also be formulated into a dosage form wherein one or more ingredients (such as active or inactive ingredients) are present as separate components. The composition or dosage form wherein the ingredients do not come as a mixture, but come as separate components, such composition/dosage form may be administered in several ways. In one possible way, the ingredients may be mixed in the desired proportions and the mixture is reconstituted in suitable reconstitution diluent and is then administered as required. Alternatively, the components or the ingredients (active or inert) may be separately administered (simultaneously or one after the other) in appropriate proportion so as to achieve the same or equivalent therapeutic level or effect as would have been achieved by administration of the equivalent mixture.


In some embodiments, pharmaceutical compositions according to the invention are formulated into a dosage form such that a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or a pharmaceutically acceptable derivative thereof, are present in the composition as admixture or as a separate components. In some other embodiments, pharmaceutical compositions according to the invention are formulated into a dosage form such that a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or a pharmaceutically acceptable derivative thereof, are present in the composition as separate components.

In one general aspect, pharmaceutical compositions according to the invention are used in treatment or prevention of a bacterial infection.

In another general aspect, there are provided methods for treating or preventing a bacterial infection in a subject, said method comprising administering to said subject effective amount of a pharmaceutical composition according to the invention. In case of dosage forms wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or a pharmaceutically acceptable derivative thereof, are present in the composition as separate components; a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof may be administered before, after or

simultaneously with the administration of meropenem or a pharmaceutically acceptable derivative thereof.

In yet another general aspect, there are provided methods for treating or preventing bacterial infections in a subject, said methods comprising administering to said subject an effective amount of: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



In another general aspect, there are provided methods for treating or preventing bacterial infections in a subject, said methods comprising administering to said subject an effective amount of: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof; wherein amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof administered is from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, there is provided a method for treating or preventing a bacterial infection in a subject, said method comprising administering to said subject: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, in any of the following amounts:

(i) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(ii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(iii) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(iv) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(v) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(vi) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(vii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(viii) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(ix) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(x) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xi) about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xii) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xiii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xiv) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xv) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xvi) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof; or

(xvii) about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

In some embodiments, in the methods according to the invention, a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is administered in an amount from about 0.01 gram to about 10 gram. In some other embodiments, in the methods according to the invention, meropenem or a pharmaceutically acceptable derivative thereof is administered in an amount from about 0.01 gram to about 10 gram.

In some embodiments, in the methods according to the invention, a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is administered before, after or simultaneously with the administration of meropenem or a pharmaceutically acceptable derivative thereof.

In the methods according to the invention, the pharmaceutical composition and/or other pharmaceutically active ingredients disclosed herein may be administered by any appropriate method, which serves to deliver the composition, or its constituents, or the active ingredients to the desired site. The method of administration can vary depending on various factors, such as for example, the components of the pharmaceutical composition and the nature of the active ingredients, the site of the potential or actual infection, the microorganism (e.g. bacteria) involved, severity of infection, age and physical condition of the subject. Some non-limiting examples of administering the composition to a subject according to this invention include oral, intravenous, topical, intraspiratory, intraperitoneal, intramuscular, parenteral, sublingual, transdermal, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, gene gun, dermal patch, eye drop, ear drop or mouthwash. In some embodiments, the compositions or one or more active ingredients according to the invention are administered parenterally.

In some embodiments, in the compositions and methods according to the invention, a compound of Formula (I) is “(2S, 5R)-7-oxo-6-(sulphooxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”. In some other embodiments, in the compositions and methods according to the invention, a compound of Formula (I) is: “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”. In some embodiments, in compositions and methods according to the invention, a compound of Formula (I) is present as a sodium or potassium salt of “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”.

In some embodiments, there is provided a method for increasing antibacterial effectiveness of meropenem or a pharmaceutically acceptable derivative thereof in a subject, said method comprising co-administering meropenem or a pharmaceutically acceptable derivative thereof, with a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof. In some other embodiments, there is provided a method for increasing antibacterial effectiveness of meropenem or a pharmaceutically acceptable derivative thereof in a subject, said method comprising co-administering meropenem or a pharmaceutically acceptable derivative thereof, with a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, wherein the amount of a compound of Formula (I) or a

stereoisomer or a pharmaceutically acceptable derivative thereof is from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

A wide variety of bacterial infections can be treated or prevented using compositions and methods according to the invention. Typical, non-limiting examples of bacterial infections that can be treated or prevented using methods and/or pharmaceutical compositions according to the invention include *E. coli* infections, *Yersinia pestis* (pneumonic plague), staphylococcal infection, mycobacteria infection, bacterial pneumonia, *Shigella* dysentery, *Serratia* infections, *Candida* infections, *Cryptococcal* infection, anthrax, tuberculosis or infections caused by *Pseudomonas aeruginosa*, *Acinetobacter baumannii* or methicillin resistant *Staphylococcus aureus* (MRSA) etc.

The pharmaceutical compositions and methods according to the invention are useful in treatment or prevention of several infections, including for example, skin and soft tissue infections, febrile neutropenia, urinary tract infection, intraabdominal infections, respiratory tract infections, pneumonia (nosocomial), bacteremia meningitis, surgical infections and the like.

In some embodiments, pharmaceutical compositions and methods according to the invention are used in treatment or prevention of infections caused by resistant bacteria. In some other embodiments, the compositions and methods according to the invention are used in treatment or prevention of infections caused by bacteria producing one or more beta-lactamase enzymes.

In general, the pharmaceutical compositions and methods disclosed herein are also effective in preventing or treating infections caused by bacteria that are considered to be less or not susceptible to one or more of known antibacterial agents or their known compositions. Some non-limiting examples of such bacteria known to have developed resistance to various antibacterial agents include *Acinetobacter*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Enterobacter*, *Klebsiella*, *Citrobacter* and a like.

EXAMPLES

The following examples illustrate embodiments of the invention that are presently best known. However, it is to be understood that the following are only exemplary or illustrative of

the application of the principles of the present invention. Numerous modifications and alternative compositions, methods, and systems may be devised by those skilled in the art without departing from the spirit and scope of the present invention. The appended claims are intended to cover such modifications and arrangements. Thus, while the present invention has been described above with particularity, the following examples provide further detail in connection with what are presently deemed to be the most practical embodiments of the invention.

The antibacterial activity of combinations according to the invention was investigated against various bacterial strains. In a typical study, minimum inhibitory concentrations (MIC) were determined using Muller Hinton Agar (MHA) (BD, USA) according to Clinical and Laboratory Standards Institute (CLSI) recommendations, (Clinical and Laboratory Standards Institute (CLSI), Performance Standards for Antimicrobial Susceptibility Testing, 20th Informational Supplement, M 100-S20, Volume 30, No. 1, 2010). In short, the inocula were adjusted to deliver about 10^4 colony forming units (CFU) per spot with a multipoint inoculator (Applied Quality Services, UK). The plates were pored with doubling concentration range of the test combinations according to invention containing MHA. The plates were inoculated and were incubated at 35°C for 18 hours. MICs were read as the lowest concentration of drug that completely inhibited bacterial growth.

Example 1

The results of antibacterial activity of combination of meropenem and the compound of Formula (I) against highly resistant strains like *Acinetobacter baumannii* harbouring carbapenem hydrolyzing (CHDL) OXA beta-lactamases, *K. pneumonia* harbouring Klebsiella pneumonia carbapenemases (KPC) and *Enterobacter cloacae* harbouring Klebsiella pneumonia carbapenemases are given in Table 1. As may be seen, the data in Table 1 reveals that both meropenem and compound of Formula (I) when used alone exhibited higher MIC values. However, the MIC values of meropenem lowered significantly in presence of the compound of Formula (I) (at 4 and 8 mcg/ml). The MIC values of meropenem reduced from 8 - 256 mcg/ml to about 0.12 - 2 mcg/ml in presence of the compound of Formula (I) (at 4 and 8 mcg/ml).

The results given in the Table 1, clearly and surprisingly demonstrate the potent antibacterial activity of combinations comprising meropenem and a compound of Formula (I) against highly resistant bacterial strains. Thus, combination of meropenem and a compound of Formula (I) has tremendous beneficial effect in inhibiting highly resistant bacterial strains

demonstrating the noteworthy therapeutic advance in the treatment of infections caused by such pathogens.

Table 1: Antimicrobial activity of the composition according to invention against various resistant bacterial strains

Organisms	Resistance mechanisms	MIC ($\mu\text{g/ml}$)			
		Meropenem	Compound of Formula (I)	MIC of Meropenem in presence of compound of Formula (I) (4 $\mu\text{g/ml}$)	MIC of Meropenem in presence of compound of Formula (I) (8 $\mu\text{g/ml}$)
NCTC Acinetobacter baumannii 13301	CHDL OXA-23	16	> 64	2	2
NCTC Acinetobacter baumannii 13304	CHDL OXA-26	16	> 64	1	1
NCTC Acinetobacter baumannii 13305	CHDL OXA-27	8	> 64	1	1
Acinetobacter baumannii SL98	CHDL OXA-23	16	> 64	1	1
Acinetobacter baumannii 1466116	CHDL OXA-23	32	> 64	2	2
<i>K. pneumoniae</i> J234	KPC	256	> 64	0.5	0.25
<i>K. pneumoniae</i> J235	KPC	128	> 64	0.25	0.25
<i>K. pneumoniae</i> J236	KPC	256	> 64	0.5	0.5
<i>K. pneumoniae</i> J237	KPC	32	> 64	0.12	0.12
<i>K. pneumoniae</i> J238	KPC	128	> 64	0.25	0.25
<i>Enterobacter cloacae</i> J222	KPC	64	> 64	0.25	0.25

Example 2


Sterile Meropenem was aseptically blended or mixed with sterile compound of Formula (I) for about 30 minutes at 8 rpm to obtain sterile dry powder for injection (compositions shown

in Table 1). Each single vial of this formulation may be reconstituted with suitable reconstitution diluent prior to administration.

Sr.	Ingredient	Quantity per vial (1.5 g/vial)	Quantity per vial (2 g/vial)	Quantity per vial (3 g/vial)	Quantity per vial (4 g/vial)
1.	Sterile Meropenem for Injection USP	1 g	1 g	2 g	2 g
2.	Sterile compound of Formula (I)	0.5 g	1 g	1 g	2 g
3.	Buffering agent	q.s	q.s	q.s.	q.s.

CLAIMS

1. A pharmaceutical composition comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



2. A pharmaceutical composition according to Claim 1, wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.125 gram to about 4 gram per gram of meropenem or a pharmaceutically acceptable derivative thereof.

3. A pharmaceutical composition according to any of the Claims 1 or 2, wherein a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.01 gram to about 10 gram.

4. A pharmaceutical composition according to any of the Claims 1 or 2, wherein meropenem or a pharmaceutically acceptable derivative thereof is present in the composition in an amount from about 0.01 gram to about 10 gram.

5. A pharmaceutical composition according to any of the Claims 1 to 4, comprising: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, in any of the following amounts:

(i) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(ii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(iii) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(iv) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(v) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(vi) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(vii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(viii) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(ix) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(x) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xi) about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xii) about 0.125 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xiii) about 0.25 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xiv) about 0.5 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xv) about 1 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof;

(xvi) about 2 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof; or

(xvii) about 4 gram of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or a pharmaceutically acceptable derivative thereof.

6. A pharmaceutical composition according to any of the Claims 1 to 5, wherein a compound of Formula (I) is: “(2S, 5R)-7-oxo-6-(sulphoxy)-1,6-diazabicyclo[3.2.1]octane-2-carbonitrile”.

7. A pharmaceutical composition according to any of the Claims 1 to 5, wherein a compound of Formula (I) is: “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”.

8. A pharmaceutical composition according to any of the Claims 1 to 5, wherein a compound of Formula (I) is present as a sodium or potassium salt of “sulphuric acid, mono[(1R,2S,5R)-2-cyano-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl] ester”.

9. A pharmaceutical composition according to any of the Claims 1 to 8, wherein the composition is formulated into a dosage form such that a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or a pharmaceutically acceptable derivative thereof, are present in the composition as admixture or as separate components.

10. A pharmaceutical composition according to Claim 9, wherein the composition is formulated into a dosage form such that a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof, and meropenem or a pharmaceutically acceptable derivative thereof, are present in the composition as separate components.


11. A pharmaceutical composition according to any of the Claims 1 to 10, wherein the composition is in form of a powder or a solution.

12. A pharmaceutical composition according to Claim 11, wherein the composition is in the form of a powder or a solution that can be reconstituted by addition of a compatible reconstitution diluent for use in parenteral administration.

13. A pharmaceutical composition according to any of the Claims 1 to 12 for use in treatment or prevention of a bacterial infection.

14. A method for preventing or treating a bacterial infection in a subject, said method comprising administering to said subject an effective amount of a pharmaceutical composition according to any of the Claims 1 to 12.

15. A method of treating or preventing a bacterial infection in a subject, said method comprising administering to said subject an effective amount of: (a) meropenem or a pharmaceutically acceptable derivative thereof, and (b) a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



16. A method according to Claim 15, wherein the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof is administered in an amount from about 0.125 gram to about 4 gram per gram of meropenem or the pharmaceutically acceptable derivative thereof.

17. A method according to any one of the Claims 15 or 16, wherein meropenem or the pharmaceutically acceptable derivative thereof and the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, are administered in any one of the following amounts:

(i) about 0.125 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(ii) about 0.25 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(iii) about 0.5 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(iv) about 1 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(v) about 2 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 0.5 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(vi) about 0.125 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(vii) about 0.25 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(viii) about 0.5 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(ix) about 1 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(x) about 2 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(xi) about 4 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 1 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(xii) about 0.125 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(xiii) about 0.25 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(xiv) about 0.5 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof;


(xv) about 1 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof;

(xvi) about 2 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof; or

(xvii) about 4 gram of the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof, and about 2 gram of meropenem or the pharmaceutically acceptable derivative thereof.

18. The method according to any one of the Claims 15 to 17, wherein the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof is administered before, after or simultaneously with the administration of meropenem or the pharmaceutically acceptable derivative thereof.

19. A method of increasing antibacterial effectiveness of meropenem or the pharmaceutically acceptable derivative thereof in a subject, said method comprising co-administering meropenem or a pharmaceutically acceptable derivative thereof, with a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable derivative thereof:



Formula (I)

20. A method according to Claim 19, wherein the compound of Formula (I) or the stereoisomer or the pharmaceutically acceptable derivative thereof is administered in an amount from about 0.125 gram to about 4 gram per gram of meropenem or the pharmaceutically acceptable derivative thereof.

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2018/052649

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D471/04 A61K31/437 A61P31/04 C07D487/04
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2015/059642 A1 (WOCKHARDT LTD [IN]) 30 April 2015 (2015-04-30) entries 8 and 9; table 1	1-20
X	WO 2013/038330 A1 (WOCKHARDT LTD [IN]; PATIL VIJAYKUMAR JAGDISHWAR [IN]; TADIPARTHI RAVIK) 21 March 2013 (2013-03-21) pages 12, 14	1,8-15, 18,19
	----- -/--	

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 27 June 2018	Date of mailing of the international search report 06/07/2018
--	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Grégoire, Ariane
--	---

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2018/052649

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CASTANHEIRA M ET AL: "Antimicrobial activities of doripenem and other carbapenems against Pseudomonas aeruginosa, other nonfermentative bacilli, and Aeromonas spp", DIAGNOSTIC MICROBIOLOGY AND INFECTIOUS DISEASE, ELSEVIER, AMSTERDAM, NL, vol. 63, no. 4, 1 April 2009 (2009-04-01), pages 426-433, XP026007294, ISSN: 0732-8893, DOI: 10.1016/J.DIAGMICROBIO.2009.01.026 [retrieved on 2009-02-26] abstract</p> <p style="text-align: center;">-----</p>	1-20
X,P	<p>KRISZTINA M. PAPP-WALLACE ET AL: "Strategic Approaches to Overcome Resistance against Gram-Negative Pathogens Using [beta]-Lactamase Inhibitors and [beta]-Lactam Enhancers: Activity of Three Novel Diazabicyclooctanes WCK 5153, Zidebactam (WCK 5107), and WCK 4234", JOURNAL OF MEDICINAL CHEMISTRY, vol. 61, no. 9, 7 April 2018 (2018-04-07), pages 4067-4086, XP055487958, ISSN: 0022-2623, DOI: 10.1021/acs.jmedchem.8b00091 abstract; table 3</p> <p style="text-align: center;">-----</p>	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2018/052649

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2015059642	A1	30-04-2015	AU 2014338611 A1	12-05-2016
			CA 2928358 A1	30-04-2015
			CN 106132411 A	16-11-2016
			EP 3060208 A1	31-08-2016
			JP 2016538262 A	08-12-2016
			RU 2016115374 A	28-11-2017
			US 2016303094 A1	20-10-2016
			WO 2015059642 A1	30-04-2015

WO 2013038330	A1	21-03-2013	AU 2012310136 A1	19-12-2013
			BR 112014003476 A2	01-03-2017
			CA 2846107 A1	21-03-2013
			CN 103781787 A	07-05-2014
			EP 3052497 A1	10-08-2016
			ES 2640520 T3	03-11-2017
			JP 6114275 B2	12-04-2017
			JP 2014521739 A	28-08-2014
			JP 2016047853 A	07-04-2016
			KR 20140023401 A	26-02-2014
			MX 349827 B	14-08-2017
			NZ 618286 A	31-07-2015
			RU 2014109447 A	10-11-2015
			US 2014088070 A1	27-03-2014
			US 2015018312 A1	15-01-2015
			US 2015141394 A1	21-05-2015
			WO 2013038330 A1	21-03-2013
			ZA 201308930 B	25-02-2015
