

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
29 January 2009 (29.01.2009)

PCT

(10) International Publication Number
WO 2009/014910 A2

(51) International Patent Classification:

C07D 401/14 (2006.01) **C07D 413/14** (2006.01)
C07D 401/04 (2006.01) **C07D 417/14** (2006.01)

(21) International Application Number:

PCT/US2008/069714

(22) International Filing Date:

10 July 2008 (10.07.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/950,859 19 July 2007 (19.07.2007) US

(71) Applicant (for all designated States except US):

METABOLEX, INC. [US/US]; 3876 Bay Center Place, Hayward, California 94545 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MA, Jingyuan** [CN/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **RABBAT, Christopher J.** [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **SONG, Jiangao** [CN/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **CHEN, Xin** [CN/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **NASHASHIBI, Imad** [JO/US];

c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **ZHAO, Zuchun** [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **NOVACK, Aaron** [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **SHI, Dong Fang** [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **CHENG, Peng** [CN/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **ZHU, Yan** [CN/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US). **MURPHY, Alison** [GB/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, California 94545 (US).

(74) Agents: **TANNER, Lorna L.** et al.; **FOLEY & LARDNER LLP**, 975 Page Mill Road, Palo Alto, California 94304-1013 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ,

[Continued on next page]

(54) Title: N-LINKED HETEROCYCLIC RECEPTOR AGONISTS FOR THE TREATMENT OF DIABETES AND METABOLIC DISORDERS

Figure 1

Tissue map of rat IC-GPCR2



WO 2009/014910 A2

(57) Abstract: Compounds and methods are provided for the treatment of, inter alia, Type II diabetes and other diseases associated with poor glycemic control.



TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FT, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL,

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

N-LINKED HETEROCYCLIC RECEPTOR AGONISTS FOR THE TREATMENT OF DIABETES AND METABOLIC DISORDERS

CROSS-REFERENCES TO RELATED APPLICATIONS

5

This application claims the benefit under 35 U.S.C. 119(e) of U.S. Provisional Application No. 60/950,859, filed 19 July 2007, which is incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

10 Diabetes mellitus can be divided into two clinical syndromes, Type I and Type II diabetes mellitus. Type I diabetes, or insulin-dependent diabetes mellitus, is a chronic autoimmune disease characterized by the extensive loss of beta cells in the pancreatic islets of Langerhans (hereinafter referred to as "pancreatic islet cells" or "islet cells"), which produce insulin. As these cells are progressively destroyed, the amount of secreted insulin
15 decreases, eventually leading to hyperglycemia (abnormally high level of glucose in the blood) when the amount secreted drops below the level required for euglycemia (normal blood glucose level). Although the exact trigger for this immune response is not known, patients with Type I diabetes have high levels of antibodies against pancreatic beta cells (hereinafter "beta cells"). However, not all patients with high levels of these antibodies
20 develop Type I diabetes.

Type II diabetes, or non-insulin-dependent diabetes mellitus, develops when muscle, fat and liver cells fail to respond normally to insulin. This failure to respond (called insulin resistance) may be due to reduced numbers of insulin receptors on these cells, or a dysfunction of signaling pathways within the cells, or both. The beta cells initially
25 compensate for this insulin resistance by increasing their insulin output. Over time, these cells become unable to produce enough insulin to maintain normal glucose levels, indicating progression to Type II diabetes (Kahn SE, *Am J Med* (2000) 108 Suppl 6a, 2S-8S).

The fasting hyperglycemia that characterizes Type II diabetes occurs as a consequence of the combined lesions of insulin resistance and beta cell dysfunction. The

beta cell defect has two components: the first component, an elevation of basal insulin release (occurring in the presence of low, non-stimulatory glucose concentrations), is observed in obese, insulin-resistant pre-diabetic stages as well as in Type II diabetes. The second component is a failure to increase insulin release above the already elevated basal output in response to a hyperglycemic challenge. This lesion is absent in prediabetes and appears to define the transition from normo-glycemic insulin-resistant states to frank diabetes. There is currently no cure for diabetes. Conventional treatments for diabetes are very limited, and focus on attempting to control blood glucose levels in order to minimize or delay complications. Current treatments target either insulin resistance (metformin, thiazolidinediones ("TZDs")), or insulin release from the beta cell (sulphonylureas, exanatide). Sulphonylureas, and other compounds that act by depolarizing the beta cell, have the side effect of hypoglycemia since they cause insulin secretion independent of circulating glucose levels. One approved drug, Byetta (exanatide) stimulates insulin secretion only in the presence of high glucose, but is not orally available and must be injected. Januvia (sitagliptin) is another recently approved drug that increases blood levels of incretin hormones, which can increase insulin secretion, reduce glucagon secretion and have other less well characterized effects. However, Januvia and other dipeptidyl peptidases IV inhibitors may also influence the tissue levels of other hormones and peptides, and the long-term consequences of this broader effect have not been fully investigated. There is an unmet need for oral drugs that stimulate insulin secretion in a glucose dependent manner.

Progressive insulin resistance and loss of insulin secreting pancreatic beta cells are primary characteristics of Type II diabetes. Normally, a decline in the insulin sensitivity of muscle and fat is compensated for by increases in insulin secretion from the beta cell. However, loss of beta cell function and mass results in insulin insufficiency and diabetes (Kahn BB, *Cell* 92:593-596, 1998; Cavaghan MK, et al, *J Clin Invest* 106:329-333, 2000; Saltiel AR, *Cell* 104:517-529, 2001; Prentki M and Nolan CJ, *J Clin Invest* 116:1802-1812 (2006); and Kahn SE, *J Clin Endocrinol Metab* 86:4047-4058, 2001). Hyperglycemia further accelerates the decline in beta cell function (UKPDS Group, *JAMA* 281 :2005-2012, 1999; Levy J, et al., *Diabetes Med* 15:290-296, 1998; and Zhou YP, et al., *J Biol Chem* 278:13 16-23, 2003). Several of the genes in which allelic variation is associated with an

increased risk of Type II diabetes are expressed selectively in the beta cell (Bell GI and Polonsky KS, *Nature* 414:788-791 (2001); Saxena R, et al, *Science* (2007) Apr 26; and Valgerdur Steinhorsdottir, et al., *Nature Genetics* (2007) Apr 26).

Insulin secretion from the beta cells of pancreatic islets is elicited by increased
5 levels of blood glucose. Glucose is taken up into the beta cell primarily by the beta cell and liver selective transporter GLUT2 (Thorens B, *Mol Membr Biol* 2001 Oct-Dec;18(4):265-73). Once inside the cell, glucose is phosphorylated by glucokinase, which is the primary glucose sensor in the beta cell since it catalyzes the irreversible rate limiting step for glucose metabolism (Matschinsky FM, *Curr Diab Rep* 2005 Jun;5(3): 171-6). The rate of glucose-6-
10 phosphate production by glucokinase is dependent on the concentration of glucose around the beta cell, and therefore this enzyme allows for a direct relationship between level of glucose in the blood and the overall rate of glucose oxidation by the cell. Mutations in glucokinase produce abnormalities in glucose dependent insulin secretion in humans giving further evidence that this hexokinase family member plays a key role in the islet response to
15 glucose (Gloyn AL, et al., *J Biol Chem* 2005 Apr 8;280(14): 14105-13, Epub 2005 Jan 25). Small molecule activators of glucokinase enhance insulin secretion and may provide a route for therapeutic exploitation of the role of this enzyme (Guertin KR and Grimsby J, *Curr Med Chem* 2006;13(15): 1839-43; and Matschinsky FM, et al., *Diabetes* 2006 Jan;55(1):1-
12) in diabetes. Glucose metabolism via glycolysis and mitochondrial oxidative
20 phosphorylation ultimately results in ATP production, and the amount of ATP produced in a beta cell is directly related to the concentration of glucose to which the beta cell is exposed.

Elevated ratios of ATP to ADP that occur in the presence of higher glucose result in the closure of the Kir6.2 channel via interaction with the SUR1 subunit of the channel complex. Closure of these channels on the plasma membrane of the beta cell results in depolarization of the membrane and subsequent activation of voltage dependent calcium channels (VDCCs) (Ashcroft FM, and Gribble FM, *Diabetologia* 42:903-919, 1999; and Seino S, *Annu Rev Physiol* 61:337-362, 1999). Calcium ion entry as well as release of calcium from intracellular stores triggers exocytosis of insulin granules, resulting in secretion of insulin into the blood stream. Agents which close the Kir6.2 channel such as
25 sulphonylureas and metaglitinides (Rendell M, *Drugs* 2004; 64(12): 1339-58; and Bickle JF, *Diabetes Metab* 2006 Apr;32(2):1 13-20) also cause membrane depolarization, and

therefore these agents stimulate insulin secretion in a glucose independent fashion. Potassium channel openers, such as diazoxide, inhibit insulin secretion by preventing elevated ATP/ADP ratios from closing the Kir6.2 channel (Hansen JB, *CurrMed Chem* 2006;13(4):361-76). Calcium channel blockers, such as verapamil and nifedipine, can also 5 inhibit insulin secretion (Henquin J C, (2004) *Diabetes* 53, S48-S58). Although sulfonylureas and metaglitinides are effective glucose lowering agents in the clinic, they act independently of blood glucose levels. Because they act independently of glucose levels, these drugs may result in hypoglycemia.

Glucose dependent insulin secretion from the beta cell is dependent on numerous 10 neurotransmitters and blood-borne hormones, as well as local, intra-islet factors. CNS activation of the vagal innervation of the islet can lead to the release of small molecules such as acetylcholine and peptides such as vasoactive intestinal polypeptide (VIP), gastrin releasing peptide (GRP) and Pituitary Adenylate Cyclase Activating Peptide (PACAP). Acetylcholine activation of phospholipase C through the G_{αq}-coupled GPCR M3 muscarinic 15 receptor leads to release of Ca++ from intracellular stores (Gilon P and Henquin JC, *Endocr Rev* 2001 Oct;22(5):565-604). Cholinergic agonists also lead to a subtle Na+ -dependent plasma membrane depolarization that can work in concert with glucose-initiated depolarization to enhance insulin release (Gilon P and Henquin JC, *Endocr Rev* 2001 Oct;22(5):565-604). VIP and PACAP each bind to an overlapping set of G_α-coupled 20 GPCRs (PAC1, VIPR1 , and VIPR2) on the beta cell that lead to stimulation of adenylate cyclase and an increase in intracellular cAMP (Filipsson K, et al., *Diabetes* 2001 Sep;50(9): 1959-69; Yamada H, et al., *Regul Pept* 2004 Dec 15;123(1-3):147-53; and Qader SS, et al., *Am J Physiol Endocrinol Metab* 2007 May;292(5):E1447-55).

Elevation of beta cell cAMP has a substantial potentiating effect on insulin 25 secretion in the presence of stimulatory levels of glucose (see below). Unfortunately, many potentiators of glucose-stimulated insulin secretion also have effects outside of the islet which limit their ability to be used as diabetes therapeutics. For example, the best available selective muscarinic agonists which stimulate insulin secretion also stimulate multiple undesirable responses in multiple tissues (Rhoades RA and Tanner GA, eds. (2003) *Medical Physiology*, 2nd ed. Lippincott, Williams and Wilkins, ISBN 0-7817-1936-4). Likewise, 30 VIP and PACAP receptors are present in multiple organ systems and mediate effects on the

reproductive, immune and other diverse systems that make them less attractive as specific enhancers of glucose dependent insulin secretion.

Incretin hormones such as Glucagon-Like Peptide 1 (GLP-I) and Glucose-dependent Insulinotropic Polypeptide (GIP, also known as Gastric Inhibitory Polypeptide) also bind to specific $G\alpha_{i/\gamma}\alpha_s$ -coupled GPCRs receptors on the surface of islet cells, including beta cells, and raise intracellular cAMP (Drucker DJ, *J Clin Invest* 2007 Jan;117(1):24-32). Although the receptors for these hormones are present in other cells and tissues, the overall sum of effects of these peptides appear to be beneficial to control of glucose metabolism in the organism (Hansotia T, et al, *J Clin Invest* 2007 Jan; 117(1): 143-52, Epub 2006 Dec 21). GIP and GLP-I are produced and secreted from intestinal K and L cells, respectively, and these peptide hormones are released in response to meals by both direct action of nutrients in the gut lumen and neural stimulation resulting from food ingestion. GIP and GLP-I have short half-lives in human circulation due to the action of the protease dipeptidyl-peptidase IV (DPPIV), and inhibitors of this protease can lower blood glucose due to their ability to raise the levels of active forms of the incretin peptides. The glucose lowering that can be obtained with DPPIV inhibitors, however, is somewhat limited since these drugs are dependent on the endogenous release of the incretin hormones. Peptides (e.g., exanatide (Byetta)) and peptide-conjugates that bind to the GIP or GLP-I receptors but are resistant to serum protease cleavage can also lower blood glucose substantially (Gonzalez C, et al., *Expert Opin Investig Drugs* 2006 Aug;15(8):887-95), but these incretin mimetics must be injected and tend to induce a high rate of nausea and therefore are not ideal therapies for general use in the Type II diabetic population. The clinical success of DPPIV inhibitors and incretin mimetics, though far from ideal, do point to the potential utility of compounds that increase incretin activity in the blood or directly stimulate cAMP in the beta cell. Some studies have indicated that beta cell responsiveness to GIP is diminished in Type II diabetes (Nauck MA, et al., *J Clin Invest* 91:301-307 (1993); and Elahi D, et al., *Regul Pept* 51:63-74 (1994)). Restoration of this responsiveness (Meneilly GS, et al., *Diabetes Care* 1993 Jan;16(1):1 10-4) may be a promising way to improve beta cell function *in vivo*.

Since increased incretin activity has a positive effect on glucose dependent insulin secretion and perhaps other mechanisms that lead to lower blood glucose, it is also of

interest to explore therapeutic approaches to increasing incretin release from intestinal K and L cells. GLP-I secretion appears to be attenuated in Type II diabetes (Vilsboll T, et al, *Diabetes* 50:609-613), so improving incretin release may ameliorate this component of metabolic dysregulation. Nutrients such as glucose and fat in the gut lumen prompt incretin secretion by interaction with apical receptors (Vilsboll T, et al., *Diabetes* 50:609-613).

5 GLP-I and GIP release can also result from neural stimulation; acetylcholine and GRP can enhance incretin release in a manner perhaps analogous to the effects of these neurotransmitters on the beta cell in regard to insulin secretion (Brubaker P, *Ann N Y Acad Sci* 2006 Jul; 1070: 10-26; and Reimann F, et al., *Diabetes* 2006 Dec;55(Suppl 2):S78-S85).

10 Somatostatin, leptin and free fatty acids also appear to modulate incretin secretion (Brubaker P, *Ann N Y Acad Sci* 2006 Jul; 1070: 10-26; and Reimann F, et al., *Diabetes* 2006 Dec;55(Suppl 2):S78-S85). To date, however, there does not appear to be a way to selectively impact these pathways to promote incretin secretion for therapeutic benefit. There is a need for oral drugs that stimulate incretin secretion in the treatment of diabetes.

15 Incretins can also increase the rate of beta cell proliferation and decrease the apoptotic rates of beta cells in animal models (Farilla L, et al., *Endocrinology* 2002 Nov; 143(11):4397-408) and human islets *in vitro* (Farilla L, et al., *Endocrinology* 2003 Dec; 144(12):5149-58). The net result of these changes is an increase in beta cell number and islet mass, and this should provide for increased insulin secretory capacity, which is another desired aim of anti-diabetic therapies. GLP-I has also been shown to protect islets from the destructive effects of agents such as streptozotocin by blocking apoptosis (Li Y, et al., *J Biol Chem* 2003 Jan 3;278(1):471-8). Cyclin D1, a key regulator of progression through the cell cycle, is up-regulated by GLP-I, and other agents that increase cAMP and PKA activity also have a similar effect (Friedrichsen BN, et al., *J Endocrinol* 2006 Mar;188(3):481-92; and Kim MJ, et al., *J Endocrinol* 2006 Mar;188(3):623-33). Increased transcription of the cyclin D1 gene occurs in response to PKA phosphorylation of CREB (cAMP-response element binding) transcription factors (Hussain MA, et al., *Mol Cell Biol* 2006 Oct;26(20):7747-59). There is a need for oral drugs that increase beta cell number and islet mass in the treatment of diabetes.

25 Beta cell cAMP levels may also be raised by inhibiting the degradation of this second messenger by phosphodiesterases to AMP (Furman B and Pyne N, *Curr Opin*

Investig Drugs 2006 Oct;7(10):898-905). There are several different cAMP phosphodiesterases in the beta cell, and many of these have been shown to serve as a brake on glucose-dependent insulin secretion. Inhibitors of cAMP phosphodiesterases have been shown to increase insulin secretion *in vitro* and *in vivo*, including PDE1C, PDE3B, PDE1O
5 (Han P, et al, *J Biol Chem* 1999 Aug 6;274(32):22337-44; Harndahl L, et al, *J Biol Chem* 2002 Oct 4;277
(40):37446-55; Walz HA, et al., *J Endocrinol* 2006 Jun;189(3):629-41; Choi YH, et al., *J Clin Invest* 2006 Dec;1 16(12):3240-51; and Cantin LD, et al., *BioorgMed Chem Lett* 2007 May 15;17(10):2869-73), but so far, no PDEs have been found to have the cell type
10 selectivity necessary to avoid undesirable effects. However, this remains an area of active investigation due to the potential for amplification of the effects of incretins and other agents that stimulate adenylate cyclase.

There appear to be multiple mechanisms by which cAMP elevation in the beta cell can enhance glucose dependent insulin secretion. Classically, many of the intracellular effects of cAMP are mediated by the cAMP-dependent protein kinase (protein kinase A, PKA) (Hatakeyama H, et al., *J Physiol* 2006 Jan 15;570(Pt 2):271-82). PKA consists of a complex of two regulatory and two catalytic domains; binding of cAMP to the catalytic domains releases the catalytic domains and results in increased protein phosphorylation activity. One of the downstream effects of this kinase activity is enhanced efficiency of
20 insulin exocytosis (Gromada J, et al., *Diabetes* 1998 Jan;47(1):57-65). Another cAMP binding protein is Epac, a guanine nucleotide exchange factor (GEF) (Kashima Y, et al., *J Biol Chem* 2001 Dec 7;276(49):46046-53, Epub 2001 Oct 11; and Shibasaki T, et al., *J Biol Chem* 2004 Feb 27;279(9):7956-61), which mediates a cAMP-dependent, but PKA-independent, increase in insulin exocytosis. Epac activated by cAMP may also enhance release of intracellular Ca++ (Holz GG, *Diabetes* 2004 Jan;53(1):5-13). The effects of
25 cAMP on insulin secretion are dependent on elevated glucose levels, so raising cAMP in the pancreatic beta cell is an important goal for therapeutics of Type II diabetes.

Agents that raise intracellular cAMP levels in the beta cell increase insulin secretion in a glucose dependent manner (MiuraY and Matsui H, *Am J Physiol Endocrinol Metab* (2003) 285, E1001-E1009). One mechanism for raising cAMP is by the action of G-protein coupled cell surface receptors, which stimulate the enzyme adenylate cyclase to

produce more cAMP. The GLP-I receptor, which is the target of exanatide, is an example of such a receptor (Thorens B, et al., *Diabetes* (1993) 42, 1678-1682). There is a need for oral drugs that increase intracellular levels of cAMP in the treatment of diabetes.


5

BRIEF SUMMARY OF THE INVENTION

Quite surprisingly, we now find that novel agonists of another G-protein coupled receptor ("GPCR"), IC-GPCR2 can raise intracellular cAMP levels (*see In Vitro* Activity Table 1 in Biological Example 1). IC-GPCR2 is also referred to as RUP3 and GPRI 19. Such raised cAMP levels increase insulin secretion in a glucose dependent manner (*see* 10 Biological Examples 2 and 3) and thus provide a useful treatment for, *inter alia*, Type II diabetes. The novel agonists described in this invention are orally active (*see* Biological Example 3), providing a significant differentiating feature to exanatide. Biological Example 4 shows the tissue specific expression of GPRI 19. Additionally, Biological Example 5 describes methods to determine the effect of GPRI 19 agonists on the secretion of incretins. 15 Biological Example 6 shows methods of determining improvements in diabetes parameters widely accepted by skilled artisans in an animal diabetes model using ZDF rats. We have also found that nucleic acid probes corresponding to IC-GPCR2 are highly enriched in pancreatic islets (the majority of which are beta cells), and are not detected in any other tissue examined (*see* Figures 1 and 2). This surprising occurrence means that the novel 20 agonists described in the current invention will be useful in diagnosing diseases effecting pancreatic islets (including beta cells) such as diabetes. Agonists of IC-GPCR2 capable of raising intracellular cAMP levels have now been identified using a cell-based screen (*see* Biological Example 1).


The present invention provides compounds represented by Formula I:

25



as well as pharmaceutical compositions containing those compounds.

The present invention further provides compounds represented by Formula II:



as well as pharmaceutical compositions containing compounds of Formula II.

Also provided are methods of treating diseases such as Type II diabetes and other diseases and conditions using one or more of these compounds or compositions, as described in further detail below. The invention also provides methods of raising intracellular levels of cyclic AMP (cAMP) by using one or more of the compounds described herein. Further, the compounds may be used to stimulate insulin production and stimulate secretion of insulin, glucagon-like peptide 1 (GLP1), and glucose dependent insulinotropic polypeptide (GIP) in a mammal, in particular a human. Additionally, the compounds described herein are useful in lowering blood glucose when administered to a subject in need of treatment to lower blood glucose.

In a related aspect, the present invention provides methods of diagnosing a number of diseases and conditions using labeled compounds of Formula I.

15

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates rat islet chip hybridization results demonstrating the islet enrichment of IC-GPCR2 mRNA relative to other tissues. Chips were hybridized with equivalent amounts of cRNA from five sets of isolated rat islets, as well as the rat tissues: brain, duodenum, adipose (fat), kidney, liver, skeletal muscle, pancreas and spleen. The "Average Difference" score reflects the relative abundance of the IC-GPCR2 mRNA in each of the tissues. The Affymetrix GeneChip analysis package called the IC-GPCR2 mRNA are "Present" in four of five islet samples and "Absent" in each of the other tissues.

Figure 2 illustrates mouse islet chip hybridization results demonstrating the islet enrichment of IC-GPCR2 mRNA relative to other tissues. Chips were hybridized with equivalent amounts of cRNA from a pancreatic beta cell line (betaHC9), four sets of isolated mouse islets, as well as the mouse tissues: adipose (fat), brain, heart, kidney, liver, lung, pituitary, skeletal muscle, small intestine, thymus, hypothalamus, adrenal, thyroid and

pancreas. The "Signal" score reflects the relative abundance of the IC-GPCR2 mRNA in each of the tissues. The Affymetrix GeneChip analysis package called the IC-GPCR2 mRNA are "Present" in betaHC9, and three of four islet samples. The Affymetrix GeneChip analysis package called the IC-GPCR2 mRNA "Absent" in each of the other 5 tissues.

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations and Definitions

The abbreviations used herein are conventional, unless otherwise defined: AcOH: acetic acid; nBuLi: n-butyllithium; Cs₂CO₃: cesium carbonate; CH₂Cl₂ or DCM: dichloromethane; CH₂MgI: methyl magnesium iodide; CuCl₂: copper chloride; DAST: (diethylamino)sulfur trifluoride; DEAD: diethyl azodicarboxylate; DIBAL: diisobutylaluminum hydride; DIPEA: diisopropylethylamine; DMSO: dimethyl sulfoxide; Et₃N: triethylamine; EtOAc: ethyl acetate; H₂: hydrogen; HBr: hydrogen bromide; HCl: hydrogen chloride; H₂O: water; H₂O₂: hydrogen peroxide; HPLC: high performance 15 liquid chromatography; KCN: potassium cyanide; LHMDS: lithium hexamethyldisilazide; LiAlH₄: lithium aluminum hydride; LiOH: lithium hydroxide; MeCN: acetonitrile; MeI: methyl iodide; MeOH: methanol; MgSO₄: magnesium sulfate; MgCO₃: magnesium carbonate; MsCl: mesyl chloride; NaHSO₃: sodium hydrogen sulfite; mCPBA: meta-chloroperoxybenzoic acid; N₂: nitrogen; Na₂CO₃: sodium carbonate; NaHCO₃: sodium 20 bicarbonate; NaNO₂: sodium nitrite; NaOH: sodium hydroxide; Na₂S₂O₃: sodium bisulfate; Na₂SO₄: sodium sulfate; NBS: N-bromosuccinimide; NH₄Cl: ammonium chloride; NH₄OAc: ammonium acetate; NMR: nuclear magnetic resonance; Pd/C: palladium on carbon; PPh₃: triphenyl phosphine; iPrOH: isopropyl alcohol; SOCl₂: thionyl chloride; THF: tetrahydrofuran; TLC: thin layer chromatography.

25 Unless otherwise stated, the following terms used in the specification and claims have the meanings given below.

"Alkyl" refers to monovalent saturated aliphatic hydrocarbyl groups having from 1 to 10 carbon atoms and, in some embodiments, from 1 to 6 carbon atoms. "C_{u-v}alkyl" refers to alkyl groups having from u to v carbon atoms. This term includes, by way of example, 30 linear and branched hydrocarbyl groups such as methyl (CH₃-), ethyl (CH₃CH₂-), n-propyl

(CH₃CH₂CH₂-), isopropyl ((CH₃)₂CH-), *i*-butyl (CH₃CH₂CH₂CH₂-), isobutyl ((CH₃)₂CHCH₂-), sec-butyl ((CH₃)(CH₃CH₂)CH-), *t*-butyl ((CH₃)₃C-), *n*-pentyl (CH₃CH₂CH₂CH₂CH₂-), and neopentyl ((CH₃)₃CCH₂-).

"Substituted alkyl" refers to an alkyl group having from 1 to 5 and, in some 5 embodiments, 1 to 3 or 1 to 2 substituents selected from the group consisting of alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted 10 aryloxy, arylthio, substituted arylthio, azido, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, guanidino, substituted guanidino, halo, hydroxy, hydroxyamino, alkoxyamino, hydrazino, substituted hydrazino, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, 15 substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclithio, substituted heterocyclithio, nitro, spirocycloalkyl, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiocyanate, thiol, alkylthio, and substituted alkylthio, wherein said substituents are as defined herein.

"Alkylidene" or "alkylene" refers to divalent saturated aliphatic hydrocarbyl 20 groups having from 1 to 10 carbon atoms and, in some embodiments, from 1 to 6 carbon atoms. "(Cu-v)alkylene" refers to alkylene groups having from u to v carbon atoms. The alkylidene and alkylene groups include branched and straight chain hydrocarbyl groups. For example "(C^ualkylene)" is meant to include methylene, ethylene, propylene, 2-methypropylene, pentylene, and the like.

"Substituted alkylidene" or "substituted alkylene" refers to an alkylidene group 25 having from 1 to 5 and, in some embodiments, 1 to 3 or 1 to 2 substituents selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, azido, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted 30

cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, guanidino, substituted guanidino, halo, hydroxy, hydroxyamino, alkoxyamino, hydrazino, substituted hydrazino, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, 5 heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclthio, substituted heterocyclthio, nitro, oxo, thione, spirocycloalkyl, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiocyanate, thiol, alkylthio, and substituted alkylthio, wherein said substituents are as defined herein.

"Alkenyl" refers to a linear or branched hydrocarbyl group having from 2 to 10 carbon atoms and in some embodiments from 2 to 6 carbon atoms or 2 to 4 carbon atoms and having at least one site of vinyl unsaturation (>C=C<). For example, (C_u-v)alkenyl refers to alkenyl groups having from u to v carbon atoms and is meant to include for example, ethenyl, propenyl, 1,3-butadienyl, and the like.

"Substituted alkenyl" refers to alkenyl groups having from 1 to 3 substituents and, 15 in some embodiments, 1 to 2 substituents, selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, alkyl, substituted alkyl, alkynyl, substituted alkynyl, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted 20 aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, guanidino, substituted guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, 25 heterocyclyloxy, substituted heterocyclyloxy, heterocyclthio, substituted heterocyclthio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined as herein and with the proviso that any hydroxy or thiol substitution is not attached to an acetylenic carbon atom.

"Alkenylene" refers to divalent alkenyl groups having from 2 to 10 carbon atoms 30 and, in some embodiments, from 2 to 6 carbon atoms. "(C_u-v)alkenylene" refers to alkenylene groups having from u to v carbon atoms.

"Alkynyl" refers to a linear monovalent hydrocarbon radical or a branched monovalent hydrocarbon radical containing at least one triple bond. The term "alkynyl" is also meant to include those hydrocarbyl groups having one triple bond and one double bond. For example, (C₂-Ce)alkynyl is meant to include ethynyl, propynyl, and the like.

5 "Substituted alkynyl" refers to alkynyl groups having from 1 to 3 substituents and, in some embodiments, from 1 to 2 substituents, selected from the group consisting of alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, 10 substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, cycloalkenyl, substituted cycloalkenyl, cycloalkenyloxy, substituted cycloalkenyloxy, cycloalkenylthio, substituted cycloalkenylthio, guanidino, substituted 15 guanidino, halo, hydroxy, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclithio, substituted heterocyclithio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined herein and with the proviso 20 that any hydroxy or thiol substitution is not attached to an acetylenic carbon atom.

"Alkynylene" refers to divalent alkynyl groups having from 2 to 10 carbon atoms and, in some embodiments, from 2 to 6 carbon atoms. "(C_u-v)alkynylene" refers to alkynylene groups having from u to v carbon atoms.

25 "Alkoxy" refers to the group -O-alkyl wherein alkyl is defined herein. Alkoxy includes, by way of example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, t-butoxy, sec-butoxy, and n-pentoxy.

"Substituted alkoxy" refers to the group -O-(substituted alkyl) wherein substituted alkyl is as defined herein.

30 "Acyl" refers to the groups H-C(O)-, alkyl-C(O)-, substituted alkyl-C(O)-, alkenyl-C(O)-, substituted alkenyl-C(O)-, alkynyl-C(O)-, substituted alkynyl-C(O)-,

cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, substituted aryl-C(O)-, substituted hydrazino-C(O)-, heteroaryl-C(O)-, substituted heteroaryl-C(O)-, heterocyclic-C(O)-, and substituted heterocyclic-C(O)-, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, 5 aryl, substituted aryl, substituted hydrazino, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein. Acyl includes the "acetyl" group $\text{CH}_3\text{C}(\text{O})-$.

"Acylamino" refers to the groups -NR²⁰C(O)H, -NR²⁰C(O)alkyl, -NR²⁰C(O)substituted alkyl, -NR²⁰C(O)cycloalkyl, -NR²⁰C(O)substituted cycloalkyl, 10 -NR²⁰C(O)alkenyl, -NR²⁰C(O)substituted alkenyl, -NR²⁰C(O)alkynyl, -NR²⁰C(O)substituted alkynyl, -NR²⁰C(O)aryl, -NR²⁰C(O)substituted aryl, -NR²⁰C(O)heteroaryl, -NR²⁰C(O)substituted heteroaryl, -NR²⁰C(O)heterocyclic, and -NR²⁰C(O)substituted heterocyclic wherein R²⁰ is hydrogen or alkyl and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, 15 substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Acyloxy" refers to the groups H-C(O)O-, alkyl-C(O)O-, substituted alkyl-C(O)O-, alkenyl-C(O)O-, substituted alkenyl-C(O)O-, alkynyl-C(O)O-, substituted alkynyl-C(O)O-, aryl-C(O)O-, substituted aryl-C(O)O-, cycloalkyl-C(O)O-, substituted cycloalkyl-C(O)O-, heteroaryl-C(O)O-, substituted heteroaryl-C(O)O-, 20 heterocyclic-C(O)O-, and substituted heterocyclic-C(O)O- wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

25 "Amino" refers to the group -NH₂.

"Substituted amino" refers to the group -NR²¹R²² where R²¹ and R²² are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted 30 heterocyclic, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-alkenyl, -S(O)₂-substituted

alkenyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocycl, and -S(O)₂-substituted heterocycl and wherein R²¹ and R²² are optionally joined together with the nitrogen bound thereto to form a heterocycl or substituted heterocycl group,

5 provided that R²¹ and R²² are both not hydrogen, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein. When R⁰¹ is hydrogen and R⁰² is alkyl, the substituted amino group is sometimes referred to herein as alkylamino. When R⁰¹ and R⁰² are alkyl, the 10 substituted amino group is sometimes referred to herein as dialkylamino. When referring to a monosubstituted amino, it is meant that either R²¹ or R²² is hydrogen but not both. When referring to a disubstituted amino, it is meant that neither R²¹ nor R²² are hydrogen.

"Hydroxyamino" refers to the group -NHOH.

"Alkoxyamino" refers to the group -NHO-alkyl wherein alkyl is defined herein.

15 "Aminocarbonyl" refers to the group -C(O)NR²³R²⁴ where R²³ and R²⁴ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, hydroxy, alkoxy, substituted alkoxy, amino, substituted amino, and acylamino, and where R²³ and R²⁴ are optionally joined together with the nitrogen bound thereto to 20 form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

25 "Aminothiocarbonyl" refers to the group -C(S)NR²³R²⁴ where R²³ and R²⁴ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R²³ and R²⁴ are optionally joined together with the nitrogen bound 30 thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl,

substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminocarbonylamino" refers to the group -NR²⁰C(O)NR²³R²⁴ where R²⁰ is
5 hydrogen or alkyl and R²³ and R²⁴ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R²³ and R²⁴ are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.
10

"Aminothiocarbonylamino" refers to the group -NR²⁰C(S)NR²³R²⁴ where R²⁰ is hydrogen or alkyl and R²³ and R²⁴ are independently selected from the group consisting of
15 hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R²³ and R²⁴ are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.
20

"Aminocarbonyloxy" refers to the group -O-C(O)NR²³R²⁴ where R²³ and R²⁴ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R²³ and R²⁴ are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.
25
30

"Aminosulfonyl" refers to the group $-S(O)_2NR^{23}R^{24}$ where R^{23} and R^{24} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R^{23} and R^{24} are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminosulfonyloxy" refers to the group $-O-S(O)_2NR^{23}R^{24}$ where R^{23} and R^{24} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R^{23} and R^{24} are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aminosulfonylamino" refers to the group $-NR^{20}-S(O)_2NR^{23}R^{24}$ where R^{20} is hydrogen or alkyl and R^{23} and R^{24} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic and where R^{23} and R^{24} are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Amidino" refers to the group $-C(=NR^{25})NR^{23}R^{24}$ where R^{25} , R^{23} , and R^{24} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic

heterocyclic and where R²³ and R²⁴ are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Aryl" refers to an aromatic group of from 6 to 14 carbon atoms and no ring heteroatoms and having a single ring (*e.g.*, phenyl) or multiple condensed (fused) rings (*e.g.*, naphthyl or anthryl). For multiple ring systems, including fused, bridged, and spiro ring systems having aromatic and non-aromatic rings that have no ring heteroatoms, the term "Aryl" or "Ar" applies when the point of attachment is at an aromatic carbon atom (*e.g.*, 5,6,7,8 tetrahydronaphthalene-2-yl is an aryl group as its point of attachment is at the 2-position of the aromatic phenyl ring).

"Substituted aryl" refers to aryl groups which are substituted with 1 to 8 and, in some embodiments, 1 to 5, 1 to 3 or 1 to 2 substituents selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, azido, carboxyl, carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, guanidino, substituted guanidino, halo, hydroxy, hydroxyamino, alkoxyamino, hydrazino, substituted hydrazino, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclthio, substituted heterocyclthio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiocyanate, thiol, alkylthio, and substituted alkylthio, wherein said substituents are defined herein.

"Arylalkyl" or "Aryl(C₁-C_z)alkyl" refers to the radical -R^uR^v where R^u is an alkylene group (having eight or fewer main chain carbon atoms) and R^v is an aryl group as defined herein. Thus, "arylalkyl" refers to groups such as, for example, benzyl, and phenylethyl, and the like. Similarly, "Arylalkenyl" means a radical -R^uR^v where R^u is an

alkenylene group and R^v is an aryl group as defined herein, *e.g.*, styrenyl, 3-phenyl-2-propenyl, and the like.

"Aryloxy" refers to the group -O-aryl, where aryl is as defined herein, that includes, by way of example, phenoxy and naphthoxy.

5 "Substituted aryloxy" refers to the group -O-(substituted aryl) where substituted aryl is as defined herein.

"Arylthio" refers to the group -S-aryl, where aryl is as defined herein.

"Substituted arylthio" refers to the group -S-(substituted aryl), where substituted aryl is as defined herein.

10 "Azido" refers to the group -N₃.

"Hydrazino" refers to the group -NHNH₂.

"Substituted hydrazino" refers to the group -NR²⁶NR²⁷R²⁸ where R²⁶, R²⁷, and R²⁸ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl, carboxyl ester, cycloalkyl, substituted cycloalkyl, heteroaryl, substituted heteroaryl, heterocyclic, substituted heterocyclic, -S(O)₂-alkyl, -S(O)₂-substituted alkyl, -S(O)₂-alkenyl, -S(O)₂-substituted alkenyl, -S(O)₂-cycloalkyl, -S(O)₂-substituted cycloalkyl, -S(O)₂-aryl, -S(O)₂-substituted aryl, -S(O)₂-heteroaryl, -S(O)₂-substituted heteroaryl, -S(O)₂-heterocyclic, and -S(O)₂-substituted heterocyclic and wherein R²⁷ and R²⁸ are 15 optionally joined, together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, provided that R²⁷ and R²⁸ are both not hydrogen, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as 20 defined herein.

25 "Cyano" or "carbonitrile" refers to the group -CN.

"Carbonyl" refers to the divalent group -C(O)- which is equivalent to -C(=O)-.

"Carboxyl" or "carboxy" refers to -C(O)OH or salts thereof.

"Carboxyl ester" or "carboxy ester" refers to the groups -C(O)O-alkyl, -C(O)O-substituted alkyl, -C(O)O-alkenyl, -C(O)O-substituted alkenyl, -C(O)O-alkynyl, -C(O)O-substituted alkynyl, -C(O)O-aryl, -C(O)O-substituted aryl, -C(O)O-cycloalkyl, -C(O)O-substituted cycloalkyl, -C(O)O-heteroaryl, -C(O)O-substituted heteroaryl, 5 -C(O)O-heterocyclic, and -C(O)O-substituted heterocyclic wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

"(Carboxyl ester)amino" refers to the group -NR²⁰-C(O)O-alkyl, 10 -NR²⁰-C(O)O-substituted alkyl, -NR²⁰-C(O)O-alkenyl, -NR²⁰-C(O)O-substituted alkenyl, -NR²⁰-C(O)O-alkynyl, -NR²⁰-C(O)O-substituted alkynyl, -NR²⁰-C(O)O-aryl, -NR²⁰-C(O)O-substituted aryl, -NR²⁰-C(O)O-cycloalkyl, -NR²⁰-C(O)O-substituted cycloalkyl, -NR²⁰-C(O)O-heteroaryl, -NR²⁰-C(O)O-substituted heteroaryl, 15 -NR²⁰-C(O)O-heterocyclic, and -NR²⁰-C(O)O-substituted heterocyclic wherein R²⁰ is alkyl or hydrogen, and wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

"(Carboxyl ester)oxy" refers to the group -O-C(O)O-alkyl, -O-C(O)O-substituted alkyl, -O-C(O)O-alkenyl, -O-C(O)O-substituted alkenyl, -O-C(O)O-alkynyl, 20 -O-C(O)O-substituted alkynyl, -O-C(O)O-aryl, -O-C(O)O-substituted aryl, -O-C(O)O-cycloalkyl, -O-C(O)O-substituted cycloalkyl, -O-C(O)O-heteroaryl, -O-C(O)O-substituted heteroaryl, -O-C(O)O-heterocyclic, and -O-C(O)O-substituted heterocyclic wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, substituted cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic, and substituted heterocyclic are as defined herein.

"Cycloalkyl" refers to a saturated or partially saturated cyclic group of from 3 to 14 carbon atoms and no ring heteroatoms and having a single ring or multiple rings including fused, bridged, and spiro ring systems. For multiple ring systems having aromatic 30 and non-aromatic rings that have no ring heteroatoms, the term "cycloalkyl" applies when the point of attachment is at a non-aromatic carbon atom (*e.g.*, 5,6,7,8,-

tetrahydronaphthalene-5-yl). The term "cycloalkyl" includes cycloalkenyl groups. Examples of cycloalkyl groups include, for instance, adamantyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and cyclohexenyl. " $C_u\sqrt{v}$ cycloalkyl" refers to cycloalkyl groups having u to v carbon atoms as ring members. " $C_u\sqrt{v}$ Cycloalkenyl" refers to cycloalkenyl groups having u to v carbon atoms as ring members.

5 "Cycloalkenyl" refers to a partially saturated cycloalkyl ring having at least one site of $>C=C<$ ring unsaturation.

"Substituted cycloalkyl" refers to a cycloalkyl group, as defined herein, having from 1 to 8, or 1 to 5, or in some embodiments 1 to 3 substituents selected from the group 10 consisting of oxo, thione, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, alkoxy, substituted alkoxy, acyl, acylamino, acyloxy, amino, substituted amino, aminocarbonyl, aminothiocarbonyl, aminocarbonylamino, aminothiocarbonylamino, aminocarbonyloxy, aminosulfonyl, aminosulfonyloxy, aminosulfonylamino, amidino, aryl, substituted aryl, aryloxy, substituted aryloxy, arylthio, substituted arylthio, azido, carboxyl, 15 carboxyl ester, (carboxyl ester)amino, (carboxyl ester)oxy, cyano, cycloalkyl, substituted cycloalkyl, cycloalkyloxy, substituted cycloalkyloxy, cycloalkylthio, substituted cycloalkylthio, guanidino, substituted guanidino, halo, hydroxy, hydroxyamino, alkoxyamino, hydrazino, substituted hydrazino, heteroaryl, substituted heteroaryl, heteroaryloxy, substituted heteroaryloxy, heteroarylthio, substituted heteroarylthio, 20 heterocyclic, substituted heterocyclic, heterocyclyloxy, substituted heterocyclyloxy, heterocyclthio, substituted heterocyclthio, nitro, SO₃H, substituted sulfonyl, sulfonyloxy, thioacyl, thiocyanate, thiol, alkylthio, and substituted alkylthio, wherein said substituents are as defined herein. The term "substituted cycloalkyl" includes substitutes cycloalkenyl groups.

25 "Cycloalkyloxy" refers to -O-cycloalkyl wherein cycloalkyl is as defined herein.

"Substituted cycloalkyloxy" refers to -O-(substituted cycloalkyl) wherein substituted cycloalkyl is as defined herein.

"Cycloalkylthio" refers to -S-cycloalkyl wherein substituted cycloalkyl is as defined herein.

"Substituted cycloalkylthio" refers to -S-(substituted cycloalkyl) wherein substituted cycloalkyl is as defined herein.

"Guanidino" refers to the group -NHC(=NH)NH₂.

"Substituted guanidino" refers to -NR²⁹C(=NR²⁹)N(R²⁹)₂ where each R²⁹ is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclyl, and substituted heterocyclyl and two R²⁹ groups attached to a common guanidino nitrogen atom are optionally joined together with the nitrogen bound thereto to form a heterocyclic or substituted heterocyclic group, provided that at least one R²⁹ is not hydrogen, and wherein said substituents are as defined herein.

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo.

"Haloalkyl" refers to substitution of alkyl groups with 1 to 5 or in some embodiments 1 to 3 halo groups.

"Haloalkoxy" refers to substitution of alkoxy groups with 1 to 5 or in some embodiments 1 to 3 halo groups.

"Hydroxy" or "hydroxyl" refers to the group -OH.

"Heteroaryl" refers to an aromatic group of from 1 to 14 carbon atoms and 1 to 6 heteroatoms selected from the group consisting of oxygen, nitrogen, and sulfur and includes a 5 to 18 member ring or ring system that includes a single ring (*e.g.*, imidazolyl) or multiple rings (*e.g.*, benzimidazol-2-yl and benzimidazol-6-yl). For multiple ring systems, including fused, bridged, and spiro ring systems having aromatic and non-aromatic rings, the term "heteroaryl" applies if there is at least one ring heteroatom and the point of attachment is at an atom of an aromatic ring (*e.g.*, 1,2,3,4-tetrahydroquinolin-6-yl and 5,6,7,8-tetrahydroquinolin-3-yl). In one embodiment, the nitrogen and/or the sulfur ring atom(s) of the heteroaryl group are optionally oxidized to provide for the N-oxide (N→O), sulfmyl, or sulfonyl moieties. More specifically the term heteroaryl includes, but is not limited to, pyridyl, furanyl, thienyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyridazinyl, pyrimidinyl, benzofuranyl, tetrahydrobenzofuranyl, isobenzofuranyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl,

benzoxazolyl, quinolyl, tetrahydroquinolinyl, isoquinolyl, quinazolinonyl, benzimidazolyl, benzisoxazolyl, or benzothienyl.

"Substituted heteroaryl" refers to heteroaryl groups that are substituted with from 1 to 8, or in some embodiments 1 to 5, or 1 to 3, or 1 to 2 substituents selected from the group consisting of the substituents defined for substituted aryl.

"Heteroaryloxy" refers to -O-heteroaryl wherein heteroaryl is as defined herein.

"Substituted heteroaryloxy" refers to the group -O-(substituted heteroaryl) wherein heteroaryl is as defined herein.

"Heteroarylthio" refers to the group -S-heteroaryl wherein heteroaryl is as defined herein.

"Substituted heteroarylthio" refers to the group -S-(substituted heteroaryl) wherein heteroaryl is as defined herein.

"Heterocycle" or "heterocyclic" or "heterocyclo" or "heterocycloalkyl" or "heterocyclyl" refers to a saturated or partially saturated cyclic group having from 1 to 14 carbon atoms and from 1 to 6 heteroatoms selected from the group consisting of nitrogen, sulfur, or oxygen and includes single ring and multiple ring systems including fused, bridged, and spiro ring systems. For multiple ring systems having aromatic and/or non-aromatic rings, the term "heterocyclic", "heterocycle", "heterocyclo", "heterocycloalkyl" or "heterocyclyl" applies when there is at least one ring heteroatom and the point of attachment is at an atom of a non-aromatic ring (*e.g.*, 1,2,3,4-tetrahydroquinoline-3-yl, 5,6,7,8-tetrahydroquinoline-6-yl, and decahydroquinolin-6-yl). In one embodiment, the nitrogen and/or sulfur atom(s) of the heterocyclic group are optionally oxidized to provide for the N-oxide, sulfmyl, sulfonyl moieties. More specifically the heterocyclyl includes, but is not limited to, tetrahydropyranyl, piperidinyl, N-methylpiperidin-3-yl, piperazinyl, N-methylpyrrolidin-3-yl, 3-pyrrolidinyl, 2-pyrrolidone-1-yl, morpholinyl, and pyrrolidinyl. A prefix indicating the number of carbon atoms (*e.g.*, C₃-C₁₀) refers to the total number of carbon atoms in the portion of the heterocyclyl group exclusive of the number of heteroatoms.

"Substituted heterocycle" or "substituted heterocyclic" or "substituted heterocyclo'Or "substituted heterocycloalkyl" or "substituted heterocyclyl" refers to heterocyclic groups, as defined herein, that are substituted with from 1 to 5 or in some embodiments 1 to 3 of the substituents as defined for substituted cycloalkyl.

5 "Heterocyclyloxy" refers to the group -O-heterocyclyl wherein heterocyclyl is as defined herein.

"Substituted heterocyclyloxy" refers to the group -O-(substituted heterocyclyl) wherein heterocyclyl is as defined herein.

10 "Heterocyclylthio" refers to the group -S-heterocyclyl wherein heterocyclyl is as defined herein.

"Substituted heterocyclylthio" refers to the group -S-(substituted heterocyclyl) wherein heterocyclyl is as defined herein.

15 Examples of heterocycle and heteroaryl groups include, but are not limited to, azetidine, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, dihydroindole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, phthalimide, 1,2,3,4-tetrahydroisoquinoline, 4,5,6,7-tetrahydrobenzo[b]thiophene, 20 thiazole, thiazolidine, thiophene, benzo[b]thiophene, morpholinyl, thiomorpholinyl (also referred to as thiamorpholinyl), 1,1-dioxothiomorpholinyl, piperidinyl, pyrrolidine, and tetrahydrofuranyl.


"Nitro" refers to the group -NO₂.

"Oxo" refers to the atom (=O).

25 "Oxide" refers to products resulting from the oxidation of one or more heteroatoms. Examples include N-oxides, sulfoxides, and sulfones.

"Spirocycloalkyl" refers to a 3 to 10 member cyclic substituent formed by replacement of two hydrogen atoms at a common carbon atom with an alkylene group

having 2 to 9 carbon atoms, as exemplified by the following structure wherein the methylene group shown here attached to bonds marked with wavy lines is substituted with a spirocycloalkyl group:



5 "Sulfonyl" refers to the divalent group $-S(O)_2^-$.

"Substituted sulfonyl" refers to the group $-S(O)_2$ -alkyl, $-S(O)_2$ -substituted alkyl, $-S(O)_2$ -alkenyl, $-S(O)_2$ -substituted alkenyl, $-S(O)_2$ -alkynyl, $-S(O)_2$ -substituted alkynyl, $-S(O)_2$ -cycloalkyl, $-S(O)_2$ -substituted cycloalkyl, $-S(O)_2$ -aryl, $-S(O)_2$ -substituted aryl, $-S(O)_2$ -heteroaryl, $-S(O)_2$ -substituted heteroaryl, $-S(O)_2$ -heterocyclic, $-S(O)_2$ -substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

10 Substituted sulfonyl includes groups such as methyl- $S(O)_2^-$, phenyl- $S(O)_2^-$, and 4-methylphenyl- $S(O)_2^-$.

15 "Sulfonyloxy" refers to the group $-OS(O)_2$ -alkyl, $-OS(O)_2$ -substituted alkyl, $-OS(O)_2$ -alkenyl, $-OS(O)_2$ -substituted alkenyl, $-OS(O)_2$ -cycloalkyl, $-OS(O)_2$ -substituted cycloalkyl, $-OS(O)_2$ -aryl, $-OS(O)_2$ -substituted aryl, $-OS(O)_2$ -heteroaryl, $-OS(O)_2$ -substituted heteroaryl, $-OS(O)_2$ -heterocyclic, $-OS(O)_2$ -substituted heterocyclic, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

20 "Thioacyl" refers to the groups H-C(S)-, alkyl-C(S)-, substituted alkyl-C(S)-, alkenyl-C(S)-, substituted alkenyl-C(S)-, alkynyl-C(S)-, substituted alkynyl-C(S)-, cycloalkyl-C(S)-, substituted cycloalkyl-C(S)-, aryl-C(S)-, substituted aryl-C(S)-, heteroaryl-C(S)-, substituted heteroaryl-C(S)-, heterocyclic-C(S)-, and substituted heterocyclic-C(S)-, wherein alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, heterocyclic and substituted heterocyclic are as defined herein.

"Thiol" refers to the group -SH.

"Alkylthio" refers to the group -S-alkyl wherein alkyl is as defined herein.

"Substituted alkylthio" refers to the group -S-(substituted alkyl) wherein substituted alkyl is as defined herein.

5 "Thiocarbonyl" refers to the divalent group -C(S)- which is equivalent to -C(=S)-.

"Thione" refers to the atom (=S).

"Thiocyanate" refers to the group -SCN.

"Compound" and "compounds" as used herein refers to a compound encompassed by the generic formulae disclosed herein, any subgenus of those generic formulae, and any forms of the compounds within the generic and subgeneric formulae, such as an oxide, ester, prodrug, pharmaceutically acceptable salt, or solvate. Unless specified otherwise, the term further includes the racemates, stereoisomers, and tautomers of the compound or compounds.

"Racemates" refers to a mixture of enantiomers.

15 "Solvate" or "solvates" of a compound refer to those compounds, where compounds are as defined above, that are bound to a stoichiometric or non-stoichiometric amount of a solvent. Solvates of a compound includes solvates of all forms of the compound such as the oxide, ester, prodrug, or pharmaceutically acceptable salt of the disclosed generic and subgeneric formulae. Preferred solvents are volatile, non-toxic, 20 and/or acceptable for administration to humans.

"Stereoisomer" or "stereoisomers" refer to compounds that differ in the chirality of one or more stereocenters. Stereoisomers include enantiomers and diastereomers. The compounds of this invention may exist in stereoisomeric form if they possess one or more asymmetric centers or a double bond with asymmetric substitution and, therefore, can be 25 produced as individual stereoisomers or as mixtures. Unless otherwise indicated, the description is intended to include individual stereoisomers as well as mixtures. The methods for the determination of stereochemistry and the separation of stereoisomers are

well-known in the art (*see* discussion in Chapter 4 of ADVANCED ORGANIC CHEMISTRY, 4th ed., J March, John Wiley and Sons, New York, 1992).

"Tautomer" refers to alternate forms of a compound that differ in the position of a proton, such as enol-keto and imine-enamine tautomers, or the tautomeric forms of 5 heteroaryl groups containing a ring atom attached to both a ring -NH- moiety and a ring =N-moiety such as pyrazoles, imidazoles, benzimidazoles, triazoles, and tetrazoles.

"Prodrug" refers to any derivative of a compound of the embodiments that is capable of directly or indirectly providing a compound of the embodiments or an active metabolite or residue thereof when administered to a patient. Prodrugs of a compound of 10 the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications may be cleaved *in vivo* to release the parent compound, or an active metabolite. For example, prodrugs include compounds wherein a hydroxy, amino, or sulfhydryl group in a compound I is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Particularly 15 favored derivatives and prodrugs are those that increase the bioavailability of the compounds of the embodiments when such compounds are administered to a patient (*e.g.*, by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (*e.g.*, the brain or lymphatic system) relative to the parent species. Prodrugs include ester, amide, 20 carbamate (*e.g.*, N,N-dimethylaminocarbonyl) forms of hydroxy functional groups of compounds of the invention. Examples of ester prodrugs include formate, acetate, propionate, butyrate, acrylate, and ethylsuccinate derivatives. An general overview of prodrugs is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the ACS Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers 25 in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

"Pharmaceutically acceptable salt" refers to pharmaceutically acceptable salts derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, and 30 tetraalkylammonium. When the molecule contains a basic functionality, pharmaceutically acceptable salts include acid addition salts of organic or inorganic acids, such as

hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, oxalic acid, 4-toluenesulfonic acid, camphorsulfonic acid, methanesulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like. Salts can also be formed when an acidic proton present in the parent compound is either replaced by a metal ion, *e.g.*, an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, trimethylamine, N-methylglucamine, and the like. Pharmaceutically acceptable salts are suitable for administration in a patient and possess desirable pharmacological properties. Suitable salts further include those described in P. Heinrich Stahl, Camille G. Wermuth (Eds.), *Handbook of Pharmaceutical Salts Properties, Selection, and Use*; 2002.

Unless indicated otherwise, the nomenclature of substituents that are not explicitly defined herein are arrived at by naming the terminal portion of the functionality followed by the adjacent functionality toward the point of attachment. For example, the substituent "arylalkyloxycarbonyl" refers to the group (aryl)-(alkyl)-O-C(O)-.

It is understood that in all substituted groups defined above, polymers arrived at by defining substituents with further substituents to themselves (*e.g.*, substituted aryl having a substituted aryl group as a substituent which is itself substituted with a substituted aryl group, which is further substituted by a substituted aryl group, etc.) are not intended for inclusion herein. In such cases, the maximum number of such substitutions is three. For example, serial substitutions of substituted aryl groups with two other substituted aryl groups are limited to -substituted aryl-(substituted aryl)-substituted aryl.

Similarly, it is understood that the above definitions are not intended to include impermissible substitution patterns (*e.g.*, methyl substituted with 5 fluoro groups). Such impermissible substitution patterns are well known to the skilled artisan.

- "Patient" refers to mammals and includes humans and non-human mammals.
- 5 Examples of patients include, but are not limited to mice, rats, hamsters, guinea pigs, pigs, rabbits, cats, dogs, goats, sheep, cows, and humans.

The term "mammal" includes, without limitation, humans, domestic animals (*e.g.*, dogs or cats), farm animals (cows, horses, or pigs), and laboratory animals (mice, rats, hamsters, guinea pigs, pigs, rabbits, dogs, monkeys).

- 10 The terms "optional" or "optionally" as used throughout the specification means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "heterocyclo group optionally mono- or di- substituted with an alkyl group" means that the alkyl may but need not be present, and the description
- 15 includes situations where the heterocyclo group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

- "Protecting group" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in T.W. Greene and P.G. Wuts, PROTECTIVE GROUPS IN ORGANIC
- 20 CHEMISTRY, (Wiley, 2nd ed. 1991) and Harrison and Harrison, et al, COMPENDIUM OF SYNTHETIC ORGANIC METHODS, VOIS. 1-8 (John Wiley and Sons 1971-1996).

- Representative amino protecting groups include formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethyl silyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (FMOC), nitro-veratryloxycarbonyl (NVOC) and the like.
- 25 Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as benzyl and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

- Turning next to the compositions of the invention, the term "pharmaceutically acceptable carrier or excipient" means a carrier or excipient that is useful in preparing a

pharmaceutical composition that is generally safe, and possesses acceptable toxicities. Acceptable carriers or excipients include those that are acceptable for veterinary use as well as human pharmaceutical use. A "pharmaceutically acceptable carrier or excipient" as used in the specification and claims includes both one and more than one such carrier or
5 excipient.

With reference to the methods of the present invention, the following terms are used with the noted meanings:

The terms "treating" or "treatment" of a disease includes:

- (1) preventing or reducing the risk of developing the disease, *i.e.*, causing the
10 clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease,
- (2) inhibiting the disease, *i.e.*, arresting or reducing the development of the disease or its clinical symptoms, or
- (3) relieving the disease, *i.e.*, causing regression of the disease or its clinical
15 symptoms.

A preferred embodiment of the invention is treatment of a disease that consists of relieving the disease.

The term "diagnosing" refers to determining the presence or absence of a particular disease or condition. Additionally, the term refers to determining the level or
20 severity of a particular disease or condition, as well as monitoring of the disease or condition to determine its response to a particular therapeutic regimen.

The term "therapeutically effective amount" means the amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.
25 "A therapeutically effective amount" includes the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

The term "insulin resistance" can be defined generally as a disorder of glucose metabolism. More specifically, insulin resistance can be defined as the diminished ability of insulin to exert its biological action across a broad range of concentrations producing less than the expected biologic effect (*see, e.g.*, Reaven GM, *J Basic & Clin Phys & Pharm* 5 (1998) 9:387-406 and Flier J, *Ann Rev Med* (1983) 34:145-60). Insulin resistant persons have a diminished ability to properly metabolize glucose and respond poorly, if at all, to insulin therapy. Manifestations of insulin resistance include insufficient insulin activation of glucose uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue and of glucose production and secretion in liver. Insulin 10 resistance can cause or contribute to polycystic ovarian syndrome, impaired glucose tolerance, gestational diabetes, metabolic syndrome, hypertension, obesity, atherosclerosis and a variety of other disorders. Eventually, the insulin resistant individuals can progress to a point where a diabetic state is reached.

The term "diabetes mellitus" or "diabetes" means a disease or condition that is 15 generally characterized by metabolic defects in production and utilization of glucose that result in the failure to maintain appropriate blood sugar levels in the body. The result of these defects is elevated blood glucose, referred to as "hyperglycemia." Two major forms of diabetes are Type I diabetes and Type II diabetes. As described above, Type I diabetes is generally the result of an absolute deficiency of insulin, the hormone that regulates glucose 20 utilization. Type II diabetes often occurs in the presence of normal, or even elevated levels of insulin and can result from the inability of tissues to respond appropriately to insulin. Most Type II diabetic patients are insulin resistant and have a relative deficiency of insulin, in that insulin secretion can not compensate for the resistance of peripheral tissues to 25 respond to insulin. In addition, many Type II diabetics are obese. Other types of disorders of glucose homeostasis include impaired glucose tolerance, which is a metabolic stage intermediate between normal glucose homeostasis and diabetes, and gestational diabetes mellitus, which is glucose intolerance in pregnancy in women with no previous history of Type I or Type II diabetes.

The term "metabolic syndrome" refers to a cluster of metabolic abnormalities 30 including abdominal obesity, insulin resistance, glucose intolerance, diabetes, hypertension

and dyslipidemia. These abnormalities are known to be associated with an increased risk of vascular events.

The term "abdominal obesity" is defined by a cutoff point of waist circumference ≥ 102 cm in men and ≥ 80 cm in women, as recommended by the third report of the national cholesterol education program expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (NCEP/ATP Panel III).

The guidelines for diagnosis of Type II diabetes, impaired glucose tolerance, and gestational diabetes have been outlined by the American Diabetes Association (*see, e.g.,* The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus, *Diabetes Care*, (1999) VoI 2 (Suppl 1):S5-19).

The term "secretagogue" means a substance or compound that stimulates secretion. For example, an insulin secretagogue is a substance or compound that stimulates secretion of insulin.

The term "symptom" of diabetes, includes, but is not limited to, polyuria, polydipsia, and polyphagia, as used herein, incorporating their common usage. For example, "polyuria" means the passage of a large volume of urine during a given period; "polydipsia" means chronic, excessive thirst; and "polyphagia" means excessive eating. Other symptoms of diabetes include, *e.g.*, increased susceptibility to certain infections (especially fungal and staphylococcal infections), nausea, and ketoacidosis (enhanced production of ketone bodies in the blood).

The term "complication" of diabetes includes, but is not limited to, microvascular complications and macrovascular complications. Microvascular complications are those complications that generally result in small blood vessel damage. These complications include, *e.g.*, retinopathy (the impairment or loss of vision due to blood vessel damage in the eyes); neuropathy (nerve damage and foot problems due to blood vessel damage to the nervous system); and nephropathy (kidney disease due to blood vessel damage in the kidneys). Macrovascular complications are those complications that generally result from large blood vessel damage. These complications include, *e.g.*, cardiovascular disease and peripheral vascular disease. Cardiovascular disease refers to diseases of blood vessels of the heart. *See, e.g.*, Kaplan RM, et al, "Cardiovascular diseases" in *HEALTH AND HUMAN*

BEHAVIOR, pp. 206-242 (McGraw-Hill, New York 1993). Cardiovascular disease is generally one of several forms, including, *e.g.*, hypertension (also referred to as high blood pressure), coronary heart disease, stroke, and rheumatic heart disease. Peripheral vascular disease refers to diseases of any of the blood vessels outside of the heart. It is often a
5 narrowing of the blood vessels that carry blood to leg and arm muscles.

The term "atherosclerosis" encompasses vascular diseases and conditions that are recognized and understood by physicians practicing in the relevant fields of medicine. Atherosclerotic cardiovascular disease, coronary heart disease (also known as coronary artery disease or ischemic heart disease), cerebrovascular disease and peripheral vessel
10 disease are all clinical manifestations of atherosclerosis and are therefore encompassed by the terms "atherosclerosis" and "atherosclerotic disease".

The term "antihyperlipidemic" refers to the lowering of excessive lipid concentrations in blood to desired levels.

The term "modulate" refers to the treating, prevention, suppression, enhancement
15 or induction of a function or condition. For example, compounds can modulate Type II diabetes by increasing insulin in a human, thereby suppressing hyperglycemia.

The term "triglyceride(s)" ("TGs"), as used herein, incorporates its common usage. TGs consist of three fatty acid molecules esterified to a glycerol molecule. TGs serve to store fatty acids that are used by muscle cells for energy production or are taken up
20 and stored in adipose tissue.

Because cholesterol and TGs are water insoluble, they must be packaged in special molecular complexes known as "lipoproteins" in order to be transported in the plasma. Lipoproteins can accumulate in the plasma due to overproduction and/or deficient removal. There are at least five distinct lipoproteins differing in size, composition, density, and
25 function. In the cells of the small intestine, dietary lipids are packaged into large lipoprotein complexes called "chylomicrons", which have a high TG and low-cholesterol content. In the liver, TG and cholesterol esters are packaged and released into plasma as TG-rich lipoprotein called very low density lipoprotein ("VLDL"), whose primary function is the endogenous transport of TGs made in the liver or released by adipose tissue. Through
30 enzymatic action, VLDL can be either reduced and taken up by the liver, or transformed

into intermediate density lipoprotein ("IDL"). IDL, is in turn, either taken up by the liver, or is further modified to form low density lipoprotein ("LDL"). LDL is either taken up and broken down by the liver, or is taken up by extrahepatic tissue. High density lipoprotein ("HDL") helps remove cholesterol from peripheral tissues in a process called reverse
5 cholesterol transport.

The term "dyslipidemia" refers to abnormal levels of lipoproteins in blood plasma including both depressed and/or elevated levels of lipoproteins (*e.g.*, elevated levels of LDL and/or VLDL, and depressed levels of HDL).

The term "hyperlipidemia" includes, but is not limited to, the following:

10 (1) *Familial Hyperchylomicronemia*, a rare genetic disorder that causes a deficiency in an enzyme, LP lipase, that breaks down fat molecules. The LP lipase deficiency can cause the accumulation of large quantities of fat or lipoproteins in the blood;

15 (2) *Familial Hypercholesterolemia*, a relatively common genetic disorder caused where the underlying defect is a series of mutations in the LDL receptor gene that result in malfunctioning LDL receptors and/or absence of the LDL receptors. This brings about ineffective clearance of LDL by the LDL receptors resulting in elevated LDL and total cholesterol levels in the plasma;

20 (3) *Familial Combined Hyperlipidemia*, also known as multiple lipoprotein-type hyperlipidemia; an inherited disorder where patients and their affected first-degree relatives can at various times manifest high cholesterol and high triglycerides. Levels of HDL cholesterol are often moderately decreased;

25 (4) *Familial Defective Apolipoprotein B-100* is a relatively common autosomal dominant genetic abnormality. The defect is caused by a single nucleotide mutation that produces a substitution of glutamine for arginine, which can cause reduced affinity of LDL particles for the LDL receptor. Consequently, this can cause high plasma LDL and total cholesterol levels;

(5) *Familial Dysbetalipoproteinemia*, also referred to as Type III Hyperlipoproteinemia, is an uncommon inherited disorder resulting in moderate to severe

elevations of serum TG and cholesterol levels with abnormal apolipoprotein E function. HDL levels are usually normal; and

- 5 (6) *Familial Hypertriglyceridemia*, is a common inherited disorder in which the concentration of plasma VLDL is elevated. This can cause mild to moderately elevated TG levels (and usually not cholesterol levels) and can often be associated with low plasma HDL levels.

10 Risk factors for hyperlipidemia include, but are not limited to, the following: (1) disease risk factors, such as a history of Type I diabetes, Type II diabetes, Cushing's syndrome, hypothyroidism and certain types of renal failure; (2) drug risk factors, which include, birth control pills; hormones, such as estrogen, and corticosteroids; certain diuretics; and various β blockers; (3) dietary risk factors include dietary fat intake per total calories greater than 40%; saturated fat intake per total calories greater than 10%; cholesterol intake greater than 300 mg per day; habitual and excessive alcohol use; and obesity.

15 The terms "obese" and "obesity" refers to, according to the World Health Organization, a Body Mass Index ("BMI") greater than 27.8 kg/m^2 for men and 27.3 kg/m^2 for women (BMI equals weight (kg)/height (m^2)). Obesity is linked to a variety of medical conditions including diabetes and hyperlipidemia. Obesity is also a known risk factor for the development of Type II diabetes (see, e.g., Barrett-Conner E, *Epidemol Rev* (1989) 20 11:172-181; and Knowler, et al, *Am J Clin Nutr* (1991) 53:1543-1551).

The term "pancreas" refers to a gland organ in the digestive and endocrine system of vertebrates, including mammals. The pancreas secretes both digestive enzymes and hormones such as insulin, GLP-I and GIP as well as other hormones.

25 The term "islet" or "islet of Langerhans" refers to endocrine cells of the pancreas that are grouped together in islets and secrete insulin and other hormones.

The term "beta cell" refers to cells found in the islet of Langerhans that secrete insulin, amylin, and other hormones.

The term "endocrine cell" refers to cells that secrete hormones into the blood stream. Endocrine cells are found in various glands and organ systems of the body including the pancreas, intestines, and other organs.

The term "L cell" refers to gut endocrine cells that produce GLP-I.

5 The term "K cell" refers to gut endocrine cells that produce GIP.

The term "incretin" refers to a group of hormones that increases insulin secretion in response to food intake. Incretins include GLP-I and GIP.

10 The term "insulin" refers to a polypeptide hormone that regulates glucose metabolism. Insulin binds to insulin receptors in insulin sensitive cells and mediates glucose uptake. Insulin is used to treat Type I diabetes and may be used to treat Type II diabetes.

15 The term "GLP-I" or "glucagon-like peptide" is a peptide hormone primarily produced by L cells. GLP-I increases insulin secretion, decreases glucagon secretion, increases beta cell mass and insulin gene expression, inhibits acid secretion and gastric emptying in the stomach, and decreases food intake by increasing satiety.

The term "GIP" or "gastric inhibitory peptide" or "glucose dependent insulinotropic polypeptide" refers to a peptide hormone produced primarily by K cells. GIP stimulates insulin secretion. GIP also has significant effects on lipid metabolism.

20 The term "cAMP" or "cyclic AMP" or "cyclic adenosine monophosphate" refers to an intracellular signaling molecule involved in many biological processes, including glucose and lipid metabolism.

25 The term "agonist" refers to a compound that binds to a receptor and triggers a response in a cell. An agonist mimics the effect of an endogenous ligand, a hormone for example, and produces a physiological response similar to that produced by the endogenous ligand.

The term "partial agonist" refers to a compound that binds to a receptor and triggers a partial response in a cell. A partial agonist produces only a partial physiological response of the endogenous ligand.


General

The present invention derives from the discovery of compounds that act as agonists of IC-GPCR2 (Seq. ID 1) using a cell-based screen. A stable CHO cell line expressing IC-GPCR2 under the control of the CMV promoter was used and cAMP levels were measured in the cells using a homogeneous time resolved fluorescence assay. With a parental CHO cell line as a control, increased cAMP levels could be measured and compounds identified that, like exanatide, raise cAMP in cells (*see In Vitro* Activity Table in Biological Example 1). Since elevated intracellular cAMP levels in the beta cell increase insulin secretion in a glucose dependant manner (*see* Biological Examples 2 and 3), the present invention is useful for the treatment of, *inter alia*, Type II diabetes and other diseases associated with poor glycemic control. The novel agonists described in this invention are orally active (*see* Biological Example 3), providing a significant differentiating feature to exanatide. Additionally, the islet specific expression of the receptor for the novel agonists of the present invention (*see* Biological Example 4) also make the present invention useful for the diagnosis of, *inter alia*, diabetes and other diseases associated with beta cell health.

Embodiments of the Invention

Compounds

The compounds of the present invention are represented by Formula I:



20

wherein, the letters X, Y and Z are each independently selected from N and C(R³). The subscript q is 0, 1, 2, 3 or 4; the subscript r is 0, 1, 2 or 3; the subscript s is 0, 1, 2, or 3, and the sum of r + s is ≤ 4. The letter L is -(CH₂)_n- wherein n is 2, 3 or 4 and optionally at least one CH₂ is replaced by O, N(R⁵), S, S(O) or S(O)₂, and any remaining CH₂ is optionally substituted with one or two substituents selected from the group consisting of halo, Ci-₄alkyl, C₃₋₆cycloalkyl and C^haloalkyl. Ar is an aryl or heteroaryl group, wherein the aryl or heteroaryl group is optionally substituted with from one, two, three, four or five R⁶ substituents.

In one embodiment, Ar is a six-membered aryl optionally substituted with from 1 to 3 R⁶ substituents or a heteroaryl, optionally substituted with from 1 to 3 R⁶ substituents. In another embodiment, Ar is selected from the group consisting of optionally substituted phenyl, optionally substituted pyridyl, optionally substituted pyrimidinyl, optionally substituted pyrazinyl, and optionally substituted pyridazinyl.

Turning next to the R groups, R¹ is selected from the group consisting of H, C₁₋₁₀alkyl, C[^]substituted alkyl, C₃₋₇cycloalkyl, C₃₋₇substituted cycloalkyl, C₂₋₁₀alkenyl, C₂ioalkynyl, -X[^]C(O)R", -X[^]C(O)OR", -X^{1-C(0)NR^aR^b}, -S(O)₂R^a, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said 10 cycloalkyl group, heterocyclyl group, aryl group and heteroaryl group is optionally substituted with from one, two, three or four substituents independently selected from the group consisting of halo, C₁₋₁₀alkyl, C₁₋₁₀osubstituted alkyl, C₃₋₇cycloalkyl, C₂ioalkenyl, C₂₋₁₀alkynyl, aryl, heteroaryl, -CN, -NR^aC(0)R^b, -NR^aC(0)NR^aR^b, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)₂R^b, and - 15 S(O)₂NR^aR^b, or optionally R^a and R^b are combined to form a A-, 5- or 6-membered ring, and X¹ is selected from the group consisting of a bond, C[^]alkylene, C₂₋₆alkenylene, C₂₋₆alkynylene, -C(O)-, and -C(O)(CH₂)₁₋₄- , wherein the aliphatic portions of X¹ are 20 optionally substituted with one, two or three groups selected from the group consisting of halo, Ci₁₋₄alkyl, and Ci₁₋₄substituted alkyl. In one embodiment, R¹ is selected from the group consisting of -X^{1-C(0)R^a}, -X^{1-C(0)OR^a}, optionally substituted aryl group and optionally substituted heteroaryl group. In another embodiment, R¹ is -C(O)OR^a, pyrimidine 25 optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁₀alkyl, and Ci-iosubstituted alkyl, or oxadiazole optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁₀alkyl, and C₁₋₁₀substituted alkyl.

Each R² is independently selected from the group consisting of halo, C₁salkyl, C[^]substituted alkyl, C₃₋₇cycloalkyl, C₃₋₇substituted cycloalkyl, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -OR^a, -NR^aR^b, -NR^aC(0)R^b, -S(O)R^a, -S(O)₂R^a and -S(O)₂NR^aR^b, and wherein when the subscript q is 2, 3, or 4, two R² groups can optionally cyclize to form a 30 ring.

Each R³ is independently selected from the group consisting of H, halo, C₁₋₄alkyl, C^haloalkyl, C₃₋₇cycloalkyl, aryl and -OR^a.

R⁴ is selected from the group consisting of H, halo, C₁₋₆alky!, and Ci₆-substituted alkyl.

5 R⁵ is selected from the group consisting of H, Ci₄alkyl, C^haloalkyl, C₃₋₆cycloalkyl, -C(O)R^a and -S(O)₂R^a.


Each R⁶ is independently selected from the group consisting of H, halo, Ci-ioalkyl, Ci-iosubstituted alkyl, C₃₋₇cycloalkyl, C₃₋₇substituted cycloalkyl, C₂_ioalkenyl, C₂_ioalkynyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, S(O)₂R^a, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said heterocyclyl groups, said aryl and heteroaryl groups are optionally substituted with from one to four substituents independently selected from halo, oxo, C^halkyl, Ci₄haloalkyl, C₃₋₇cycloalkyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, 10 -NR^aC(O)R^b, -NR^aC(O)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)R^b, -NR^aS(O)₂R^b and -S(O)₂NR^aR^b or optionally R^a and R^b are combined to form a A-, 5- or 6-membered ring. In one embodiment, each R⁶ is independently selected from the group consisting of H, halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, C^halkyl, C^haloalkyl, substituted heteroaryl and unsubstituted heteroaryl. In one preferred embodiment, each R⁶ independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a. 15 20

For each of the above groups, each R^a and R^b is independently selected from the group consisting of H, Ci-ioalkyl, Ci-ihaloalkyl, C₃-iocycloalkyl, heterocyclyl, C₂_ioalkenyl, C₂_ioalkynyl, aryl, 5- to 6-membered heteroaryl and arylC^halkyl; and wherein the aliphatic portions of each of said R^a and R^b is optionally substituted with from one to three groups selected from the group consisting of halo, -ORⁿ, -OC(O)Rⁿ, -OC(0)N(Rⁿ)₂, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -S(O)₂N(Rⁿ)₂, -NRⁿS(O)₂Rⁿ, -C(0)N(Rⁿ)₂, -C(O)Rⁿ, -NRⁿC(O)Rⁿ, -NRⁿC(O)N(Rⁿ)₂, -C(O)ORⁿ, -NRⁿC(O)ORⁿ, -CN, -NO₂, -N(Rⁿ)₂ and -NRⁿS(O)₂N(Rⁿ)₂, 25 wherein each Rⁿ is independently H, C^haloalkyl or an unsubstituted C^halkyl; and wherein the aryl and heteroaryl portions are optionally substituted with from one to three 30

groups selected from halo, -OR^m, -OC(O)N(R^m)₂, -SR^m, -S(O)R^m, -S(O)₂R^m, -S(O)₂N(R^m)₂, -NR^mS(O)₂R^m, -C(O)N(R^m)₂, -C(O)R^m, -NR^mC(O)R^m, -NR^mC(O)N(R^m)₂, -C(O)OR^m, -NR^mC(O)OR^m, -CN, -NO₂, -N(R^m)₂ and -NR^mS(O)₂N(R^m)₂, wherein each R^m is independently H, C_{i-3}haloalkyl or an unsubstituted C^aHcyl.


5 The compounds provided herein also include any pharmaceutically acceptable salts, solvates, stereoisomers, tautomers, and esters of the compounds as well as any isotopically labeled isomers thereof. In general, the compounds useful in the methods described herein are those compounds of the formula above, wherein the molecular weight of the compound is less than 1200, more preferably less than about 1000, still more 10 preferably less than about 800 and still more preferably from about 200 to about 600.

The ring having X, Y and Z as ring substituents will, in one group of embodiments, be a ring in which one of X, Y or Z is N. In another group of embodiments, two of X, Y and Z are N. In still another group of embodiments, the ring is one in which all three of X, Y and Z are N. A further group of embodiments comprises compounds in which 15 X, Y, and Z are all C(R³). Representations of these embodiments are provided as:



wherein the wavy lines indicate the positions of attachment to either L or to the carbon atom bearing R⁴.

20 In one group of preferred embodiments, the ring having X, Y and Z is selected from:




For each of the above groups of embodiments, an additional set of embodiments are those in which subscript r is 1, subscript s is 0 or 1, subscript q is 0, 1 or 2 and Ar is phenyl, optionally substituted with from 1 to 3 R⁶ substituents. Still another set of embodiments are those in which subscript r is 1, subscript s is 0 or 1, subscript q is 0 and Ar is selected from the group consisting of pyridyl, pyrimidinyl and pyrazinyl, each of which is optionally substituted with from 1, 2 or 3 R⁶ substituents. In one embodiment, the compounds of Formula I are those compounds in which the subscript n is 2. Yet another set of embodiments are those in which subscript r is 1, subscript s is 0 or 1, subscript q is 0, 1 or 2, and subscript n is 2. In still another set of embodiments, subscript r is 1, subscript s is 0 or 1, subscript q is 0, 1 or 2, subscript n is 2 and one CH₂ is replaced by O.

In another group of embodiments of Formula I, subscript r is 1; subscript s is 0 or 1; subscript q is 0, 1 or, 2; subscript n is 2 and one of CH₂ is replaced by O, S or N(R⁵); R⁴ is selected from H, F, CH₃ and OH; and the ring having X, Y and Z is selected from pyrazole, imidazole, 1,2,3-triazole and 1,2,4-triazole. Preferably, Ar is phenyl, optionally substituted with from 1 to 3 R⁶ substituents. More preferably, Ar is substituted with from 1 to 2 R⁶ substituents independently selected from the group consisting of halo, C₁₋₁₀alkyl, Ci_idhaloalkyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aSR^b, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 5-membered heterocyclo group, aryl, and a 5- to 6-membered heteroaryl group. In some embodiments, each R⁶ is independently selected from the group consisting of halo, -OR^a, -NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)OR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aSR^b, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 5-membered heterocyclo group, aryl, and a 5- to 6-membered heteroaryl group. Within each of the groups of embodiments and preferred embodiments, one group of further preferred embodiments are those in which R¹ is a 5- to 10-membered heteroaryl group, and is optionally substituted with from one to two substituents independently selected from halo, Ci_ioalkyl, Ci_iohaloalkyl, C₃₋₇cycloalkyl, C₂_ioalkenyl, C₂_ioalkynyl, aryl, heteroaryl, CN, NO₂, -OR^a, -NR^aR^b, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)OR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aSR^b, -NR^aS(O)R^b, -NR^aS(O)₂R^b, and -S(O)₂NR^aR^b. Still further preferred are those embodiments in which R¹ is a pyridine or pyrimidine, and is optionally substituted with from one to two substituents independently selected from halo, C₁₋₁₀alkyl, Ci_idhaloalkyl, C₃₋₇cycloalkyl, C₂₋₁₀alkenyl,

C_2 -ioalkynyl, aryl, heteroaryl, CN, NO_2 , -OR^a, -NR^aR^b, -C(0)0R^a, -C(O)NR^aR^b, -NR^aC(0)R^b, -NR^aC(0)0R^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aSR^b, -NR^aS(O)R^b, -NR^aS(O)₂R^b, and -S(O)₂NR^aR^b. In still another group of embodiments, R¹ is selected from the group consisting of -X¹C(O)R^a, -X¹C(O)OR^a, -X¹-C(0)NR^aR^b and -S(O)₂R^a.

5 In another aspect, this invention provides a compound represented by Formula II:



(H)

wherein, the letters X, Y, Z, D and G are each independently selected from the group consisting of N and C(R³). The subscript q is 0, 1, 2, 3 or 4. The subscript k is 1, 2 or 3.

10 R¹ is selected from the group consisting of H, C¹⁻¹⁰substituted alkyl, C_{3-7} cycloalkyl, C_{3-7} substituted cycloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, -X¹C(O)R^a, -X¹C(O)OR^a, -X¹-C(0)NR^aR^b, -S(O)₂R^a, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said cycloalkyl group, heterocyclyl group, aryl group and heteroaryl group is optionally substituted with from 1 to 4
15 substituents independently selected from the group consisting of halo, C_{1-10} alkyl, Ci- i osubstituted alkyl, C_{3-7} cycloalkyl, C_2 -ioalkenyl, C_2 -ioalkynyl, aryl, heteroaryl, -CN, -NR^aC(0)R^b, -NR^aC(0)NR^aR^b, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)₂R^b, and -S(O)₂NR^aR^b, or optionally R^a and R^b are combined to form a 4-, 5- or 6-membered ring, and X¹ is selected from the group consisting
20 of a bond, C¹alkylene, C_{2-6} alkenylene, C_{2-6} alkynylene, -C(O)-, and -C(O)(CH₂)₁₋₄-⁻, wherein the aliphatic portions of X¹ are optionally substituted with one to three groups selected from the group consisting of halo, Ci-₄alkyl, and C¹substituted alkyl.

Each R² is independently selected from the group consisting of halo, C¹alkyl, C¹substituted alkyl, C_{3-7} cycloalkyl, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -OR^a, -NR^aR^b, -NR^aC(0)R^b, -S(O)R^aR^b, -S(O)₂R^a and -S(O)₂NR^aR^b, and wherein when the subscript q is 2, 3 or 4, two R² groups can optionally cyclize to form a ring.

Each R³ is independently selected from the group consisting of H, halo, C^{alkyl}, C₁₋₄haloalkyl, C₃₋₇cycloalkyl, aryl and OR^a.


Turning now to R⁶, each R⁶ is independently selected from the group consisting of H, halo, Ci-alkyl, C^{alkyl}, C₁₋₄haloalkyl, C₃₋₇cycloalkyl, C₂-ioalkenyl, C₂-ioalkynyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, S(O)₂R^a, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said heterocyclyl groups, said aryl and heteroaryl groups are optionally substituted with from one to four substituents independently selected from halo, oxo, Ci₁₋₄alkyl, C^{alkyl}, C₃₋₇cycloalkyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)₂R^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)₂R^b, and -S(O)₂NR^aR^b or optionally R^a and R^b are combined to form a 4-, 5- or 6-membered ring.

Each R^a and R^b is independently selected from the group consisting of H, Ci-alkyl, C₁₋₄haloalkyl, C₃₋₇cycloalkyl, heterocyclyl, C₂-ioalkenyl, C₂-ioalkynyl, aryl, 5- to 6-membered heteroaryl and arylCi₁₋₄alkyl; and wherein the aliphatic portions of each of said R^a and R^b is optionally substituted with from one to three groups selected from the group consisting of halo, -ORⁿ, -OC(O)Rⁿ, -OC(O)N(Rⁿ)₂, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -S(O)₂N(Rⁿ)₂, -NRⁿS(O)₂Rⁿ, -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NRⁿC(O)Rⁿ, -NRⁿC(O)N(Rⁿ)₂, -C(O)ORⁿ, -NRⁿC(O)ORⁿ, -CN, -NO₂, -N(Rⁿ)₂ and -NRⁿS(O)₂N(Rⁿ)₂, wherein each Rⁿ is independently H, C^{alkyl} or an unsubstituted Ci₁₋₆alkyl.

Each R⁷ and R⁸ is independently selected from the group consisting of hydrogen, Ci₁₋₆alkyl, C^{alkyl}, C₃₋₇cycloalkyl.

One embodiment of the present invention are compounds in which q is 0 or 1, and/or one or both R⁷ and R⁸ are hydrogen.

In one embodiment, the ring having X, Y, and Z in Formula II is a ring in which X, Y, and Z are all CR³. Alternatively, one of X, Y, and Z, is N or two of X, Y, and Z are N, or three of X, Y, and Z are N. Representations of some of these embodiments are provided below:



wherein, the wavy lines indicate the position of attachment to either the piperidinyl group or the aryloxymethyl group.

A further embodiment provides compounds in which R³ is hydrogen and/or one or 5 both R⁷ and R⁸ of Formula II are hydrogen.

Another embodiment provides compounds of Formula II wherein R¹ is selected from the group consisting of -X¹-C(O)R^a, -X¹-C(O)OR^a, optionally substituted aryl and optionally substituted heteroaryl group. Alternatively, R¹ is -C(O)OR^a, pyrimidine 10 optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₃₋₆substituted cycloalkyl, Ci₁₋₆alkyl, and C¹⁻⁶alkyl, or oxadiazole optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁alkyl, and Ci₁₋₆alkyl, or oxadiazole optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁alkyl, and Ci₁₋₆alkyl.

Yet another group of embodiments provides compounds in which subscript k is 1 15 or 2. Additionally, in an embodiment of Formula II wherein subscript k is 1 or 2, each R⁶ is independently selected from the group consisting of halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, Ci₃₋₆alkyl, C¹⁻⁶haloalkyl, substituted heteroaryl and unsubstituted heteroaryl. Further, in an embodiment of Formula II wherein subscript k is 1 or 2, each R⁶ is 20 independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

One embodiment of the invention provides compounds of Formula II wherein R¹ is a heteroaryl group optionally substituted with from one to two substituents selected from the group consisting of halo, Ci₁₋₆alkyl, and Ci₁₋₆haloalkyl, and each R⁶ is independently selected from the group consisting of halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, C₁₋₆alkyl, C¹⁻⁶haloalkyl substituted heteroaryl and unsubstituted heteroaryl.

Compounds of Formula II are also provided wherein R¹ is a substituted 25 pyrimidinyl or substituted oxadiazolyl group, and each R⁶ is independently selected from

the group consisting of fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

Yet another embodiment provides compounds of Formula II wherein k is 2, R¹ is a pyrimidinyl group optionally substituted with from one to two substituents selected from the group consisting of halo, C^aHcyl, and Ci₆haloalkyl, and each R⁶ is independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

In one embodiment of the present invention are compounds of Formula II wherein q is 0, k is 2, both R⁷ and R⁸ are hydrogen, R¹ is a heteroaryl group optionally substituted with from one to two substituents selected from the group consisting of halo, C₁₋₆alkyl, and Ci₆haloalkyl, and each R⁶ is independently selected from the group consisting of hydrogen, halo, -SR^a, -S(O)R^a, -S(O)₂R^a, -S(O)₂NR^aR^b, substituted heteroaryl and unsubstituted heteroaryl.

Yet another embodiment of the present invention are compounds of Formula II wherein q is 0, k is 2, both R⁷ and R⁸ are hydrogen, R¹ is an unsubstituted pyrimidinyl or a substituted pyrimidinyl substituted with from one to two substituents selected from the group consisting of halo, C^aalkyl, and C^ahaloalkyl, and each R⁶ is independently selected from the group consisting of hydrogen, halo, -SR^a, -S(O)R^a, -S(O)₂R^a, -S(O)₂NR^aR^b, substituted heteroaryl and unsubstituted heteroaryl.

Still another embodiment of the present invention are compounds of Formula II wherein q is 0, k is 2, both R⁷ and R⁸ are hydrogen, R¹ is an unsubstituted pyrimidinyl or a substituted pyrimidinyl substituted with from one to two substituents selected from the group consisting of halo, Ci₆alkyl, and Ci₆haloalkyl, and each R⁶ is independently selected from the group consisting of hydrogen, halo, -SR^a, -S(O)R^a, -S(O)₂R^a, -S(O)₂NR^aR^b, and tetrazolyl.

Preparation of Compounds of the Invention

The compounds of the present invention can be prepared in a number of ways familiar to one skilled in the art of organic synthesis. The synthetic route of compounds in the present invention is not limited to the methods outlined below or as provided in the Examples. Individual compounds may require manipulation of the conditions in order to

accommodate various functional groups and may require appropriate use of protecting groups. Purification, if necessary, can be accomplished on a silica gel column eluted with the appropriate organic solvent system. Also, reverse phase HPLC or recrystallization may be employed.

5 *Compositions and Methods of Treatment*

In accordance with the present invention methods of treating a disease or condition selected from the group consisting of Type I diabetes, Type II diabetes and metabolic syndrome are provided. The method comprises administering to a subject in need of such treatment an effective amount of a compound of the present invention.

10 In another aspect, methods of raising intracellular levels of cyclic AMP in a cell expressing GPR 119 are provided. The method comprises exposing a cell that expresses GPRI 19 to a compound of the invention. Cyclic AMP levels are determined by the methods disclosed in the Example sections herein.

15 In one embodiment, the cell that expresses GPRI 19 is a pancreatic cell, an islet cell, or a beta cell, an intestinal endocrine cell, an L cell or a K cell.

Another aspect of the invention provides a method of stimulating insulin production in a mammal, in particular a human. The method comprises administering an effective amount of a compound of the invention to the mammal. In response to administration of a compound to the subject, insulin is produced by the beta cells.

20 Biological Example 2 provides detailed methods by which a skilled artisan can measure insulin secretion in laboratory animals in response to administration of a compound of the invention.

In another aspect, the invention provides a method of stimulating insulin secretion in a mammal, in particular a human. The method comprises administering an effective amount of a compound of the invention to the mammal. In response to administration of a compound to the subject, insulin is secreted into the blood stream by the beta cells.

Biological Example 2 provides methods of determining insulin secretion in rats.

A further aspect of the invention provides a method of stimulating glucose-dependent insulin secretion in a mammal, in particular a human. The method comprises

administering an effective amount of a compound of the invention to the mammal. After administration to the subject, insulin is secreted into the blood stream by the beta cells in a glucose-dependent manner. Biological Example 3 provides methods and data that show the blood glucose lowering effects of the compounds of the invention.

5 In another embodiment, the invention provides methods of lowering blood glucose in a mammal, preferably a human. The method comprises administering an effective amount of a compound of the invention to the mammal. In response to administration of a compound to the subject, blood glucose levels are lowered. The method further comprises steps to measure blood glucose levels before and after administration of a compound of the
10 invention. Blood glucose levels are easily measured by numerous commercially available glucose monitoring devices that measure blood glucose from samples of blood or urine. Blood glucose can also be measured by commercially available glucometers that do not require blood or urine samples. Biological Example 5 provides methods that teach how to measure improvements in diabetes paramaters, including blood glucose monitoring.

15 Another aspect of the invention provides a method of stimulating incretin production in a mammal, in particular a human. The method comprises administering an effective amount of a compound of the invention to the mammal. In response to administration of a compound to the subject, glucagon-like peptide 1 and glucose-dependent insulinotropic polypeptide is produced by the intestinal endocrine cells. Biological
20 Example 4 provides detailed methods by which a skilled artisan can measure incretin production in laboratory animals in response to administration of a compound of the invention.

25 In accordance with the present invention, a therapeutically effective amount of a compound of Formula I can be used for the preparation of a pharmaceutical composition useful for treating Type II diabetes and/or lowering the plasma level of glucose. In addition, a therapeutically effective amount of a compound of Formula I can be used for the preparation of a pharmaceutical composition useful for treating other indications that include diabetes as a component, such as metabolic syndrome, as well as indications that can be improved as a result of increased insulin production (such as the early stages of Type
30 I diabetes).

The compositions of the invention can include compounds of Formula I, pharmaceutically acceptable salts thereof, or a hydrolysable precursor thereof. In general, the compound is mixed with suitable carriers or excipient(s) in a therapeutically effective amount. By a "therapeutically effective dose", "therapeutically effective amount", or, 5 interchangeably, "pharmacologically acceptable dose" or "pharmacologically acceptable amount", it is meant that a sufficient amount of the compound of the present invention and a pharmaceutically acceptable carrier, will be present in order to achieve a desired result, *e.g.*, alleviating a symptom or complication of Type II diabetes.

The compounds of Formula I that are used in the methods of the present invention 10 can be incorporated into a variety of formulations for therapeutic administration. More particularly, the compounds of Formula I can be formulated into pharmaceutical compositions by combination with appropriate, pharmaceutically acceptable carriers or diluents, and can be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, pills, powders, granules, dragees, gels, slurries, ointments, 15 solutions, suppositories, injections, inhalants and aerosols. As such, administration of the compounds can be achieved in various ways, including oral, buccal, rectal, parenteral, intraperitoneal, intradermal, transdermal, and/or intratracheal administration. Moreover, the compound can be administered in a local rather than systemic manner, in a depot or sustained release formulation. In addition, the compounds can be administered in a 20 liposome.

The compounds of Formula I can be formulated with common excipients, diluents or carriers, and compressed into tablets, or formulated as elixirs or solutions for convenient oral administration, or administered by the intramuscular or intravenous routes. The compounds can be administered transdermally, and can be formulated as sustained release 25 dosage forms and the like. Compounds of Formula I can be administered alone, in combination with each other, or they can be used in combination with other known compounds (see *Combination Therapy* below).

Suitable formulations for use in the present invention are found in *Remington's Pharmaceutical Sciences* (Mack Publishing Company (1985) Philadelphia, PA, 17th ed.), 30 which is incorporated herein by reference. Moreover, for a brief review of methods for drug delivery, see, Langer, *Science* (1990) 249:1527-1533, which is incorporated herein by

reference. The pharmaceutical compositions described herein can be manufactured in a manner that is known to those of skill in the art, *i.e.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. The following methods and excipients are merely exemplary and
5 are in no way limiting.

For injection, the compounds can be formulated into preparations by dissolving, suspending or emulsifying them in an aqueous or nonaqueous solvent, such as vegetable or other similar oils, synthetic aliphatic acid glycerides, esters of higher aliphatic acids or propylene glycol; and if desired, with conventional additives such as solubilizers, isotonic
10 agents, suspending agents, emulsifying agents, stabilizers and preservatives. Preferably, the compounds of the present invention can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in
15 the art.

For oral administration, the compounds of Formula I can be formulated readily by combining with pharmaceutically acceptable carriers that are well known in the art. Such carriers enable the compounds to be formulated as tablets, pills, dragees, capsules, emulsions, lipophilic and hydrophilic suspensions, liquids, gels, syrups, slurries,
20 suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by mixing the compounds with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose
25 preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone. If desired, disintegrating agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

30 Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions can be used, which can optionally contain gum arabic, talc, polyvinyl

pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

5 Pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved 10 or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers can be added. All formulations for oral administration should be in dosages suitable for such administration.

For buccal administration, the compositions can take the form of tablets or lozenges formulated in conventional manner.

15 For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas, or from propellant-free, dry-powder inhalers. In the case of a 20 pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, *e.g.*, 25 by bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, *e.g.*, in ampules or in multidose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulator agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension can also contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds can also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter, carbowaxes, polyethylene glycols or other glycerides, all of which melt at body temperature, yet are solidified at room temperature.

In addition to the formulations described previously, the compounds can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

Alternatively, other delivery systems for hydrophobic pharmaceutical compounds can be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. In a presently preferred embodiment, long-circulating, *i.e.*, stealth liposomes can be employed. Such liposomes are generally described in Woodle, et al., U.S. Patent No. 5,013,556. The compounds of the present invention can also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719.

Certain organic solvents such as dimethyl sulfoxide ("DMSO") also can be employed, although usually at the cost of greater toxicity. Additionally, the compounds can

be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules can, depending on their chemical nature, release the compounds for a few
5 hours up to over 100 days.

The pharmaceutical compositions also can comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

10 Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in a therapeutically effective amount. The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician. Determination of an effective
15 amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any compound used in the method of the present invention, a therapeutically effective dose can be estimated initially from cell culture assays, animal models, or microdosing of human subjects.

20 Moreover, toxicity and therapeutic efficacy of the compounds described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, by determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effect is the therapeutic index and can be expressed as the ratio
25 between LD₅₀ and ED₅₀. Compounds that exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a dosage range that is not toxic for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage can vary within this range depending upon the dosage
30 form employed and the route of administration utilized. The exact formulation, route of

administration and dosage can be chosen by the individual physician in view of the patient's condition (see, e.g., Fingl, et al., 1975 In: *The Pharmacological Basis of Therapeutics*, Ch. 1).

The amount of active compound that can be combined with a carrier material to produce a single dosage form will vary depending upon the disease treated, the mammalian species, and the particular mode of administration. However, as a general guide, suitable unit doses for the compounds of the present invention can, for example, preferably contain between 0.1 mg to about 1000 mg of the active compound. A preferred unit dose is between 1 mg to about 100 mg. A more preferred unit dose is between 1 mg to about 20 mg. Such unit doses can be administered more than once a day, for example 2, 3, 4, 5 or 6 times a day, but preferably 1 or 2 times per day, so that the total dosage for a 70 kg adult is in the range of 0.001 to about 15 mg per kg weight of subject per administration. A preferred dosage is 0.01 to about 1.5 mg per kg weight of subject per administration, and such therapy can extend for a number of weeks or months, and in some cases, years. It will be understood, however, that the specific dose level for any particular patient will depend on a variety of factors including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the individual being treated; the time and route of administration; the rate of excretion; other drugs that have previously been administered; and the severity of the particular disease undergoing therapy, as is well understood by those of skill in the area.

A typical dosage can be one 1 mg to about 20 mg tablet taken once a day, or, multiple times per day, or one time-release capsule or tablet taken once a day and containing a proportionally higher content of active ingredient. The time-release effect can be obtained by capsule materials that dissolve at different pH values, by capsules that release slowly by osmotic pressure, or by any other known means of controlled release.

It can be necessary to use dosages outside these ranges in some cases as will be apparent to those skilled in the art. Further, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

Combination Therapy

As noted above, the compounds of the present invention will, in some instances, be used in combination with other therapeutic agents to bring about a desired effect.

Selection of additional agents will, in large part, depend on the desired target therapy (*see, e.g.*,

5 Turner N, et al, *Prog Drug Res* (1998) 51:33-94; Haffner S, *Diabetes Care* (1998) 21:160-178; and DeFronzo R, et al. (eds.), *Diabetes Reviews* (1997) Vol. 5 No. 4). A number of studies have investigated the benefits of combination therapies with oral agents (*see, e.g.*, Mahler R, *J Clin Endocrinol Metab* (1999) 84:1 165-71; United Kingdom Prospective Diabetes Study Group: UKPDS 28, *Diabetes Care* (1998) 21:87-92; Bardin 10 CW,(ed.), CURRENT THERAPY IN ENDOCRINOLOGY AND METABOLISM, 6th Edition (Mosby - Year Book, Inc., St. Louis, MO 1997); Chiasson J, et al., *Ann Intern Med* (1994) 121:928-935; Coniff R, et al., *Clin Ther* (1997) 19:16-26; Coniff R, et al., *Am J Med* (1995) 98:443-451; and Iwamoto Y, et al., *Diabet Med* (1996) 13:365-370; Kwiterovich P, *Am J Cardiol* (1998) 82(12A):U-17U). These studies indicate that diabetes modulation can be further 15 improved by the addition of a second agent to the therapeutic regimen. Combination therapy includes administration of a single pharmaceutical dosage formulation that contains a compound having the general structure of Formula I and one or more additional active agents, as well as administration of a compound of Formula I and each active agent in its own separate pharmaceutical dosage formulation. For example, a compound of Formula I 20 and a DPPIV inhibitor can be administered to the human subject together in a single oral dosage composition, such as a tablet or capsule, or each agent can be administered in separate oral dosage formulations. Where separate dosage formulations are used, a compound of Formula I and one or more additional active agents can be administered at essentially the same time (*i.e.*, concurrently), or at separately staggered times (*i.e.*, sequentially). Combination therapy is understood to include all these regimens.

An example of combination therapy can be seen in modulating (preventing the onset of the symptoms or complications associated with) diabetes (or treating, preventing or reducing the risk of developing, diabetes and its related symptoms, complications, and disorders), wherein the compounds of Formula I can be effectively used in combination 30 with, for example, biguanides (such as metformin); thiazolidinediones (such as ciglitazone, pioglitazone, troglitazone, and rosiglitazone); dipeptidyl-peptidase-4 ("DPPIV") inhibitors

(such as vildagliptin and sitagliptin); glucagonlike peptide- 1 ("GLP-I") receptor agonists (such as exanatide) (or GLP-I mimetics); PPAR gamma agonists or partial agonists; dual PPAR alpha, PPAR gamma agonists or partial agonists; dual PPAR delta, PPAR gamma agonists or partial agonists; pan PPAR agonists or partial agonists; dehydroepiandrosterone
5 (also referred to as DHEA or its conjugated sulphate ester, DHEA-SO₄); antiglucocorticoids; TNF α inhibitors; α -glucosidase inhibitors (such as acarbose, miglitol, and voglibose); sulfonylureas (such as chlorpropamide, tolbutamide, acetohexamide, tolazamide, glyburide, gliclazide, glynase, glimepiride, and glipizide); pramlintide (a synthetic analog of the human hormone amylin); other insulin secretagogues (such as
10 repaglinide, gliquidone, and nateglinide); insulin (or insulin mimetics); glucagon receptor antagonists; gastric inhibitory peptide ("GIP"); or GIP mimetics; as well as the active agents discussed below for treating obesity, hyperlipidemia, atherosclerosis and/or metabolic syndrome.

Another example of combination therapy can be seen in treating obesity or
15 obesity-related disorders, wherein the compounds of Formula I can be effectively used in combination with, for example, phenylpropanolamine, phenteramine; diethylpropion; mazindol; fenfluramine; dexfenfluramine; phentiramine, β -3 adrenoceptor agonist agents; sibutramine; gastrointestinal lipase inhibitors (such as orlistat); and leptins. Other agents used in treating obesity or obesity-related disorders wherein the compounds of Formula I
20 can be effectively used in combination with, for example, cannabinoid-1 ("CB-I") receptor antagonists (such as rimonabant); PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR delta agonists or partial agonists; dual PPAR delta, PPAR gamma agonists or partial agonists; pan PPAR agonists or partial agonists; neuropeptide Y; enterostatin; cholecytokinin; bombesin; amylin; histamine H₃ receptors; dopamine D₂ receptors;
25 melanocyte stimulating hormone; corticotrophin releasing factor; galanin; and gamma amino butyric acid (GABA).

Still another example of combination therapy can be seen in modulating hyperlipidemia (treating hyperlipidemia and its related complications), wherein the compounds of Formula I can be effectively used in combination with, for example, statins
30 (such as atorvastatin, fluvastatin, lovastatin, pravastatin, and simvastatin), CETP inhibitors (such as torcetrapib); a cholesterol absorption inhibitor (such as ezetimibe); PPAR alpha

agonists or partial agonists; PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR gamma agonists or partial agonists; dual PPAR delta, PPAR gamma agonists or partial agonists; pan PPAR agonists or partial agonists; fenofibric acid derivatives (such as gemfibrozil, clofibrate, fenofibrate, and bezafibrate); bile acid-binding resins (such as colestipol or cholestyramine); nicotinic acid; probucol; betacarotene; vitamin E; or vitamin C.

A further example of combination therapy can be seen in modulating atherosclerosis, wherein a compound of Formula I is administered in combination with one or more of the following active agents: an antihyperlipidemic agent; a plasma HDL-raising agent; an antihypercholesterolemic agent, such as a cholesterol biosynthesis inhibitor, *e.g.*, an hydroxymethylglutaryl (HMG) CoA reductase inhibitor (also referred to as statins, such as lovastatin, simvastatin, pravastatin, fluvastatin, and atorvastatin); an HMG-CoA synthase inhibitor; a squalene epoxidase inhibitor; or a squalene synthetase inhibitor (also known as squalene synthase inhibitor); an acyl-coenzyme A cholesterol acyltransferase (ACAT) inhibitor, such as melinamide; probucol; nicotinic acid and the salts thereof and niacinamide; a cholesterol absorption inhibitor, such as β -sitosterol; a bile acid sequestrant anion exchange resin, such as cholestyramine, colestipol or dialkylaminoalkyl derivatives of a cross-linked dextran; an LDL receptor inducer; fibrates, such as clofibrate, bezafibrate, fenofibrate, and gemfibrozil; vitamin B₆ (also known as pyridoxine) and the pharmaceutically acceptable salts thereof, such as the HCl salt; vitamin B₁₂ (also known as cyanocobalamin); vitamin B₃ (also known as nicotinic acid and niacinamide); anti-oxidant vitamins, such as vitamin C and E and beta carotene; a β -blocker; an angiotensin II antagonist; an angiotensin converting enzyme inhibitor; PPAR alpha agonists or partial agonists; PPAR delta agonists or partial agonists; PPAR gamma agonists or partial agonists; dual PPAR alpha, PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR gamma agonists or partial agonists; dual PPAR delta, PPAR gamma agonists or partial agonists; pan PPAR agonists or partial agonists; and a platelet aggregation inhibitor, such as fibrinogen receptor antagonists (*i.e.*, glycoprotein IIb/IIIa fibrinogen receptor antagonists) and aspirin. As noted above, the compounds of Formula I can be administered in combination with more than one additional active agent, for example, a combination of a compound of Formula I with an HMG-CoA reductase inhibitor (*e.g.*, atorvastatin, fluvastatin, lovastatin, pravastatin,

and simvastatin) and aspirin, or a compound of Formula I with an HMG-CoA reductase inhibitor and a β -blocker.

Additionally, an effective amount of a compound of Formula I and a therapeutically effective amount of one or more active agents selected from the group consisting of: an antihyperlipidemic agent; a plasma HDL-raising agent; an antihypercholesterolemic agent, such as a cholesterol biosynthesis inhibitor, for example, an HMG-CoA reductase inhibitor; an HMG-CoA synthase inhibitor; a squalene epoxidase inhibitor, or a squalene synthetase inhibitor (also known as squalene synthase inhibitor); an acyl-coenzyme A cholesterol acyltransferase inhibitor; probucol; nicotinic acid and the salts thereof; CETP inhibitors such as torcetrapib; a cholesterol absorption inhibitor such as ezetimibe; PPAR alpha agonists or partial agonists; PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR delta agonists or partial agonists; dual PPAR alpha, PPAR gamma agonists or partial agonists; dual PPAR delta, PPAR gamma agonists or partial agonists; pan PPAR agonists or partial agonists; niacinamide; a cholesterol absorption inhibitor; a bile acid sequestrant anion exchange resin; a LDL receptor inducer; clofibrate, fenofibrate, and gemfibrozil; vitamin *B*_e and the pharmaceutically acceptable salts thereof; vitamin B₁₂; an anti-oxidant vitamin; a β -blocker; an angiotensin II antagonist; an angiotensin converting enzyme inhibitor; a platelet aggregation inhibitor; a fibrinogen receptor antagonist; aspirin; phentiramines, β -3 adrenergic receptor agonists; sulfonylureas, biguanides, α -glucosidase 10
inhibitors, other insulin secretagogues, and insulin can be used together for the preparation 15
of a pharmaceutical composition useful for the above-described treatments.
20

An additional example of combination therapy can be seen in modulating metabolic syndrome (or treating metabolic syndrome and its related symptoms, complications and disorders), wherein the compounds of Formula I can be effectively used 25
in combination with, for example, the active agents discussed above for modulating or treating diabetes, obesity, hyperlipidemia, atherosclerosis, and/or their respective related symptoms, complications and disorders.

In a further embodiment, a compound of the present invention can be administered in combination with halofenic acid, an ester of halofenic acid, or another prodrug of

halofenic acid, preferably with (-)-(4-chlorophenyl)-(3-trifluoromethylphenoxy)-acetic acid 2-acetylaminooethyl ester.

Methods of Diagnosis and/or Imaging

Compounds of the present invention are also useful in methods of diagnosis and/or imaging. Many direct methods are available to evaluate an agent's biodistribution in the body such as magnetic resonance imaging ("MRI"), positron emission tomography ("PET"), and single photon emission computed tomography ("SPECT"). Each of these methods can detect the distribution of a compound within the body if that compound contains an atom with the appropriate nuclear properties. MRI detects paramagnetic nuclei; PET and SPECT detect the emission of particles from the decay of radionuclei.

Most therapeutic agents are not able to be detected by these techniques without modification. Thus, for PET it is necessary to incorporate an appropriate positron-emitting radionuclide. There are relatively few positron-emitting isotopes that are suitable for labeling a therapeutic agent. The carbon isotope, ¹¹C, has been used for PET, but has a short half-life of 20.5 minutes. Accordingly, the facilities for synthesis and use are typically near to a cyclotron where the precursor ¹¹C starting material is generated. Other isotopes have even shorter half-lives. ¹³N has a half-life of 10 minutes and ¹⁵O has an even shorter half-life of 2 minutes. The emissions of both are more energetic, however, than those of ¹¹C and PET studies have been carried out with these isotopes (*see*, Clinical Positron Emission Tomography, Mosby Year Book, 1992, KF Hubner, et al, Chapter 2). Another useful isotope, ¹⁸F, has a half-life of 110 minutes. This allows sufficient time for incorporation into a radiolabeled tracer, for purification and for administration into a human or animal subject. ¹⁸F labeled compounds have been used in studies of glucose metabolism and localization of glucose uptake associated with brain activity. For example, ¹⁸F-L-fluorodopa and other dopamine receptor analogs have also been used in mapping dopamine receptor distribution.

SPECT imaging employs isotope tracers that emit high energy photons (γ -emitters). The range of useful isotopes is greater than for PET, but SPECT provides lower three-dimensional resolution. Nevertheless, SPECT is widely used to obtain clinically significant information about analog binding, localization and clearance rates. A

useful isotope for SPECT imaging is ^{123}I , a γ -emitter with a 13.3 hour half life. Compounds labeled with ^{123}I can be shipped up to about 1000 miles from the manufacturing site, or the isotope itself can be transported for on-site synthesis. Eighty-five percent of the isotope's emissions are 159 KeV photons, which are readily measured by SPECT instrumentation currently in use. Other halogen isotopes can serve for PET or SPECT imaging, or for conventional tracer labeling. These include ^{75}Br , ^{76}Br , ^{77}Br and ^{82}Br as having usable half-lives and emission characteristics. In general, the chemical means exist to substitute any halogen moiety for the described isotopes. Therefore, the biochemical or physiological activities of any halogenated homolog of the described compounds are now available for use by those skilled in the art, including stable isotope halogen homologs.

In the context of the present invention, methods are provided for diagnosing a disease or condition selected from Type I diabetes and Type II diabetes, the method comprising

- (a) administering to a subject having, or at risk for, such a disease or condition an imaging amount of a compound of the invention, wherein the compound is isotopically labeled; and
- (b) imaging the subject to determine the number, mass or volume of pancreatic beta cells or islet endocrine cells; or to assess the function of pancreatic beta cells or islet endocrine cells.

Preferably, the compound is labeled with ^{11}C or ^{14}C . In other preferred embodiments, the imaging is conducted via PET or SPECT.

Kits

In addition, the present invention provides for kits with unit doses of the compounds of Formula I, either in oral or injectable doses. In addition to the containers containing the unit doses will be an informational package insert describing the use and attendant benefits of the drugs in treating Type II diabetes, obesity, hyperlipidemia, atherosclerosis and metabolic syndrome, and/or their respective related symptoms, complications and disorders. Preferred compounds and unit doses are those described herein above.

For the compositions, methods and kits provided above, one of skill in the art will understand that preferred compounds for use in each are those compounds that are noted as preferred above. Still further preferred compounds for the compositions, methods and kits are those compounds provided in the non-limiting Examples below.

5

EXAMPLES

General Methods. All operations involving moisture and/or oxygen sensitive materials were conducted under an atmosphere of dry nitrogen in pre-dried glassware. Unless noted otherwise, materials were obtained from commercially available sources and used without further purification.

10 Flash chromatography was performed on E. Merck silica gel 60 (240-400 mesh) according to the protocol of Still, Kahn, and Mitra (*J. Org. Chem.* **1978**, *43*, 2923). Thin layer chromatography was performed using precoated plates purchased from E. Merck (silica gel 60 PF₂₅₄, 0.25 mm) and spots were visualized with long-wave ultraviolet light followed by an appropriate staining reagent.

15 Nuclear magnetic resonance ("NMR") spectra were recorded on a Varian Inova-400 resonance spectrometer. ¹H NMR chemical shifts are given in parts per million (δ) downfield from tetramethylsilane ("TMS") using TMS or the residual solvent signal (CHCl₃ = δ 7.24, DMSO = δ 2.50) as internal standard. ¹H NMR information is tabulated in the following format: number of protons, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet), coupling constant(s) (J) in Hertz, and, in selected cases, position assignment. 20 The prefix app is occasionally applied in cases where the true signal multiplicity was unresolved and br indicates the signal in question was broadened.


The compounds were named using ChemBioDraw Ultra Version 11.0.

LCMS analysis was performed using a PE SCIEX API 2000 spectrometer with a 25 Phenomenex Luna 5 micron C₁₈ column.


The compounds of the present invention can be prepared by methodology in the Reaction Scheme below, and with specific reagents and conditions provided in each of the examples below.

Reaction Schemes

Scheme 1




Reagents and conditions: a. DMF, 90 °C, 72 h; b. LiAlH₄, THF; c. MsCl, NEt₃, CH₂Cl₂; d. CsCO₃, HA-Ar, CH₃CN, 82 °C, 4 h.


Scheme 2

Reagents and conditions: a. sodium ascorbate, cat. CuSO₄, ^tBuOH/H₂O (1:1 v/v), 72 h, rt; b. NaOH, 10:1 MeOH/THF, 1-4 h; c. PPh₃, DIAD, THF, 0 °C-rt, 4h.

5 **Preparation of Intermediate 1: 4-(3-Methanesulfonyloxymethyl-[1,2,4]triazol-1-yl)-piperidine-1-carboxylic acid *tert*-butyl ester**




10 **Step 1: 4-(3-Methoxycarbonyl-[1,2,4]triazol-1-yl)-piperidine-1-carboxylic acid *tert*-butyl ester**



To a solution of *IH*-[1,2,4]triazole-3-carboxylic acid ethyl ester (1.05 g, 8.23 mmol) in dimethylformamide (50 mL) was added sodium hydride (60%, 0.395 g, 9.88 mmol). The solution was stirred at room temperature for 20 minutes followed by 1 hour at 70 °C. 4-methanesulfonyloxy-piperidine-1-carboxylic acid tert-butyl ester (2.3 g, 8.23 mmol) was then added in a single portion and heated at 70 °C for 40 hours. The solution was cooled to 0 °C and the salt precipitate was removed by filtration. The filtrate was diluted with dichloromethane and washed with water, brine, dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was purified by flash chromatograph on silica gel with Hexanes and EtOAc to afford the desired product. ¹H NMR (CDCl₃): δ 8.18 (1H, s), 4.40 (1H, m), 4.22 (2H, m), 3.96 (3H, s), 2.82 (2H, m), 2.12 (2H, m), 1.94 (2H, m), 1.41 (9H, s).


Step 2: 4-(3-Hydroxymethyl-[1,2,4]triazol-1-yl)-piperidine-1-carboxylic acid tert-butyl ester



To a solution of 4-(3-ethoxycarbonyl-[1,2,4]triazol-1-yl)-piperidine-1-carboxylic acid tert-butyl ester (from **Step 1**, 0.64 g, 2.06 mmol) in THF (10 mL) at 0 °C was added


lithium aluminum hydride (0.078 g, 2.06 mmol). The solution was allowed to warm to room temperature and stirred for 4 hours. The solution was diluted with THF, cooled to 0 °C, and 1 mL of water was added followed by 1 mL of 15% NaOH and 3 mL of water. The solution was allowed to warm to room temperature and stirred for 15 minutes. MgSO₄ was 5 then added and the solution was stirred an additional 15 minutes, filtered and concentrated *in vacuo*. The residue was purified by flash chromatograph on silica gel with MeOH and CH₂Cl₂ to afford the desired product. ¹H NMR (CDCl₃): δ 8.03 (1H, s), 4.79 (2H, d), 4.25 (3H, m), 2.95 (2H, m), 2.58 (1H, t), 2.19 (2H, m), 1.98 (2H, m), 1.43 (9H, s).

Step 3: 4-(3-Methanesulfonyloxyethyl-[1,2,4]triazol-1-yl)-piperidine-1-carboxylic acid *tert*-hvXy\ ester
10




To a solution of 4-(3-hydroxymethyl-[1,2,4]triazol-1-yl)-piperidine-1 -carboxylic acid *tert*-hvXy\ ester (from **Step 2** above, 0.181 g, 0.641 mmol) and triethylamine (0.134 mL, 0.961 mmol) in dichloromethane (5 mL) at 0 °C was added methanesulfonylchloride (0.055 mL, 0.705 mmol). The resulting solution was stirred at room temperature for 1 hour. The solvent was removed *in vacuo* to yield the desired product as a light yellow oil that was used without further purification.
15

Preparation of Intermediate 2: 4-(4-Methanesulfonyloxyethyl-[1,2,3]triazol-2-yl)-piperidine-1 -carboxylic acid *tert*-butyl ester
20



Intermediate 2 was prepared manner similar to that described in **Intermediate 1** above from 1,2,3-triazole-4-carboxylic acid methyl ester which can be prepared from azidotrimethylsilane and methyl propiolate as described in patent EP1422228 A1.

Preparation of Intermediate 3: 4-(4-Methanesulfonyloxymethyl-pyrazol-1-yl)-piperidine-1-carboxylic acid *tert-butyl* ester




10

Intermediate 3 was prepared manner similar to that described in **Intermediate 1** above from commercial available ethyl 4-pyrazolecarboxylate.


Preparation of Intermediate 4: 4-(3-Methanesulfonyloxymethyl-pyrazol-1-yl)-piperidine-1-carboxylic acid *tert-butyl* ester

15



Intermediate 4 was prepared manner similar to that described in **Intermediate 1** above from 1*H*-pyrazole-3-carboxylic acid methyl ester which was prepared from
5 commercial available 1*H*-pyrazole-3-carboxylic acid.

Preparation of Intermediate 5: 4-(5-Methanesulfonyloxymethyl-tetrazol-2-yl)-piperidine-1-carboxylic acid *tert*-butyl ester




10

Intermediate 5 was prepared manner similar to that described in **Intermediate 1** above from commercially available 1*H*-tetrazole-5-carboxylic acid ethyl ester sodium salt.


Preparation of Intermediate 6: 4-[3-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,4]triazol-1-yl]-piperidine hydrochloride

15



A solution of 4-[3-(2-fluoro-4-methanesulfonyl-phenoxyethyl)-[1,2,4]triazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester (see **Example 1**, 0.249 g, 2.74 mmol) in methanol (10 mL) was treated with 10 mL of 4N HCl in dioxane. The resulting solution was stirred at room temperature for 30 minutes and the solvent was removed *in vacuo* to afford the desired product as an HCl salt which was used without further purification.


Preparation of Intermediate 7: 4-[4-(2-Fluoro-4-methanesulfonyl-phenoxyethyl)-[1,2,3]triazol-2-yl]-piperidine hydrochloride



10

Intermediate 7 was prepared manner similar to that described in **Intermediate 6**.

Preparation of Intermediate 8: 4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxyethyl)-[1,2,3]triazol-2-yl]-piperidine hydrochloride




15

Intermediate 8 was prepared manner similar to that described in **Intermediate 6**.

Preparation of Intermediate 9: 4-[4-(4-Tetrazol-1-yl-phenoxyethyl)-[1,2,3]triazol-2-yl]-piperidine hydrochloride


20



Intermediate 9 was prepared manner similar to that described in **Intermediate 6**.

Preparation of Intermediate 10: 4-[5-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-tetrazol-2-yl]-piperidine hydrochloride


5



Intermediate 10 was prepared in a manner similar to that described in


Intermediate 6 using.

10 **Preparation of Intermediate 11:** 4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxy)methyl]-pyrazol-1-yl]-piperidine hydrochloride



15 **Intermediate 11** was prepared in a manner similar to that described in
Intermediate 6.

Preparation of Intermediate 12: 4-[4-(4-Tetrazol-1-yl-phenoxy)methyl]-pyrazol-1-yl]-piperidine hydrochloride




20

Intermediate 12 was prepared in a manner similar to that described in **Intermediate 6**.


Preparation of Intermediate 13: 5-((2*H*-1,2,3-Triazol-4-yl)methoxy)-2-(1 *H*-tetrazol-1-yl)pyridine

5




Step 1: 6-(1 *H*-Tetrazol-1-yl)pyridin-3-ol

10



NaN₃ (2.89g, 44 mmol) and triethylorthoformate (7.4 ml, 44 mmol) were added to commercially available 6-amino-pyridin-3-ol (3.5g, 32 mmol) in AcOH (50mL). The reaction mixture was heated to 100 °C for 1-3 hours. The reaction was cooled to room temperature and left to stand overnight. The precipitate was filtered, washed with EtOAc (150 mL) and dried *in vacuo*. The crude product was obtained as a beige solid and used in the next step without further purification.


Step 2 : 5-(Prop-2-ynyoxy)-2-(1 *H*-tetrazol-1-yl)pyridine



To a flask under N₂ fitted with a reflux condenser was added 6-tetrazol-1-yl-pyridin-3-ol (2.25 g, 13.7 mmol) and *N,N*-dimethylformamide (8.2 mL). Potassium carbonate (3.79 g, 27.4 mmol, 2 eq.), potassium iodide (0.143 g, 0.86 mmol, 0.06 eq.) and propargyl chloride (1.98 mL, 17.9 mmol, 1.3 eq.) were added and heated at 80 °C for three hours. The reaction was cooled to room temperature and 10 mL of water was added to affect precipitation. The solid was filtered off, washed with water and dried overnight. The expected product was used in the next step without further purification. ¹H NMR (DMSO-*d*₆): δ 10.08 (1H, s), 8.36 (1H, d, *J* = 3.0 Hz), 8.01 (1H, d, *J* = 9 Hz), 7.79 (1H, dd, *J* = 3.0, 9.0 Hz), 5.01 (2H, s), 3.71 (1H, s)

Step 3: (4-((1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-1*H*-1,2,3-triazol-1-yl)methyl pivalate

15




To a solution of azidomethyl pivalate (4.26 g, 27.1 mmol) in 1:1 *tert*-butanol water (90 mL total) was added 5-(prop-2-ynyl)oxy-2-(1*H*-tetrazol-1-yl)pyridine (5.45 g, 27.1 mmol), sodium ascorbate (1.4 mL of a 1M solution in water) and copper sulfate (1.4 mL of a 1M solution in water). The solution was stirred at room temperature for 72 hours. Water was added and the suspension was extracted with ethyl acetate. The organic layer was separated, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford the expected product which was used in the next step without further purification. ¹H NMR

(DMSCM λ): δ 10.07 (IH, s), 8.43-8.41 (2H, m), 8.00 (IH, d, $J = 8.8$ Hz), 7.88 (IH, dd, $J = 3.2, 8.8$ Hz), 6.32 (2H, s), 5.38 (2H, s), 1.08 (9H, s).

Step 4: 5-((2*H*-1,2,3-Triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine


5



To a solution of (4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-1*H*-1,2,3-triazol-1-yl)methyl pivalate (10 g, 27.9 mmol) in MeOH (60 mL) and THF (6 mL) was added sodium hydroxide (61.5 mL of a 1M solution in water). The mixture was stirred for 30 minutes and hydrochloric acid was added (61.5 mL of a 1M solution in water). The resulting solution was extracted with ethyl acetate, separated, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford the expected product which was used in subsequent steps without further purification. ^1H NMR (DMSO- \wedge): δ 10.07 (IH, s), 8.43-8.41 (2H, m), 8.00 (IH, d, $J = 8.8$ Hz), 7.88 (IH, dd, $J = 3.2, 8.8$ Hz), 6.32 (2H, s), 5.38 (2H, s), 1.08 (9H, s).


Preparation of Intermediate 14: 1-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)piperidin-4-ol

20




The title compound was synthesized in a manner similar to that described in WO2008/008887 A2, example 16, page 84.

5 **Preparation of Intermediate 15:** 5-((2-(Piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine hydrochloride



10 **Intermediate 15** was prepared in a manner similar to that described in
Intermediate 6.



15 **Preparation of Intermediate 16:** 1-(4-((2*H*-1,2,3-Triazol-4-yl)methoxy)-3-fluorophenyl)-1*H*-tetrazole



15

This intermediate was synthesized from commercially available 4-amino-2-fluorophenol in a manner similar to that described for **Intermediate 13** above.


20 **Preparation of Intermediate 17:** *cis*-1-(5-Ethylpyrimidin-2-yl)-3-fluoropiperidin-4-ol

**Step 1:** 1-(5-Ethylpyrimidin-2-yl)piperidin-4-one

5

The title compound was synthesized from piperidin-4-one and 2-chloro-5-ethylpyrimidine manner similar to that described in **Example 9**.

10

Step 2: *cis*-1-(5-Ethylpyrimidin-2-yl)-3-fluoropiperidin-4-ol


15

The title compound was synthesized from piperidin-4-one and 2-chloro-5-ethylpyrimidine in a manner similar to that described in *J. Med. Chem.* **1998**, *41*, 2667-2670.

20

Preparation of Intermediate 18: *cis*-tert-Butyl 3-fluoro-4-hydroxypiperidine-1-


carboxylate



The title compound was synthesized as described in *J. Med. Chem.* **1998**, *41*, 2667-2670.

5

Preparation of Intermediate 19: *trans*-3-Fluoro-4-((2-fluoro-4-(1*f*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidine hydrochloride




10

Intermediate 19 was prepared in a manner similar to that described in **Intermediate 6**.


Preparation of Intermediate 20: 1-(5-(Trifluoromethyl)pyrimidin-2-yl)piperidin-4-ol

15



To a solution of 4-hydroxypiperidine (6.18 g, 61 mmol), and potassium carbonate (11.0 g, 80 mmol) in *N,N* dimethylformamide (150 mL) was added 2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidine (12.3 g, 54 mmol) and stirred at room temperature for 18 h. To 20 the reaction mixture was added 300 mL of water and the heterogeneous mixture was stirred for an additional 10 minutes. The white solid was filtered and washed with water to afford the desired product.


Preparation of Intermediate 21: 5-((2-(3-Methylpiperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine



5

Intermediate 21 was prepared in a manner similar to that described in **Intermediate 6** above in a mixture of four stereoisomers.

Preparation of Intermediate 22: 1-(5-Chloropyrimidin-2-yl)piperidin-4-ol




10

Intermediate 22 was prepared from 2, 5-dichloropyrimidine and 4-hydroxypiperidine in a manner similar to that described in **Intermediate 20**.

Preparation of Intermediate 23: 1-(5-Chloropyrimidin-2-yl)piperidin-4-yl methanesulfonate


15



Intermediate 23 was prepared from **Intermediate 22** in a manner similar to that described in **Intermediate 1, Step 3.**


Preparation of Intermediate 24: (2-(1-(5-Chloropyrimidin-2-yl)piperidin-4-yl)-
2*H*-1,2,3-triazol-4-yl)methyl methanesulfonate

5



Intermediate 24 was prepared from 1-(5-chloropyrimidin-2-yl)piperidin-4-yl
methanesulfonate (**Intermediate 23**) and 1,2,3-triazole-4-carboxylic acid methyl ester
10 (prepared from azidotrimethylsilane and methyl propiolate as described in patent
EP1422228 A1) in a manner similar to that described in **Intermediate 1.**

Preparation of Intermediate 25: 1-(5-(Trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl methanesulfonate




15

Intermediate 25 was prepared from **Intermediate 20** in a manner similar to that described in **Intermediate 1, Step 3.**


Preparation of Intermediate 26: (2-(1-(5-(Trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methyl methanesulfonate

20



Intermediate 26 was prepared from 1-(5-(trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl methanesulfonate (**Intermediate 25**) and 1,2,3-triazole-4-carboxylic acid 5 methyl ester (prepared from azidotrimethylsilane and methyl propiolate as described in patent EP1422228 A1) in a manner similar to that described in **Intermediate 1**.


Preparation of Intermediate 27: (2-(1-(5-(Trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methanol



10


Intermediate 27 was prepared from 1-(5-(trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl methanesulfonate (**Intermediate 25**) and 1,2,3-triazole-4-carboxylic acid methyl ester (prepared from azidotrimethylsilane and methyl propiolate as described in patent EP1422228 A1) in a manner similar to that described in **Intermediate 1** through Step 2.

Preparation of Intermediate 28: 1-(5-Ethylpyrimidin-2-yl)piperidin-4-ol




Intermediate 28 was prepared from 2-chloro-5-ethylpyrimidine and 4-hydroxypiperidine in a manner similar to that described in **Intermediate 20**.

Preparation of **Intermediate 29**: 1-(5-Ethylpyrimidin-2-yl)piperidin-4-yl
5 methanesulfonate



Intermediate 29 was prepared from **Intermediate 28** in a manner similar to that
10 described in **Intermediate 1, Step 3**.

Preparation of **Intermediate 30**: (2-(1-(5-Ethylpyrimidin-2-yl)piperidin-4-yl)-
2H-1,2,3-triazol-4-yl)methyl methanesulfonate




15

Intermediate 30 was prepared from 1-(5-ethylpyrimidin-2-yl)piperidin-4-yl methanesulfonate (**Intermediate 29**) and 1,2,3-triazole-4-carboxylic acid methyl ester (prepared from azidotrimethylsilane and methyl propiolate as described in patent EP1422228 A1) in a manner similar to that described in **Intermediate 1**.

20

Preparation of Intermediate 31: 4-(3-((2-fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)- 1H-pyrazol- 1-yl)piperidine hydrochloride




5

Intermediate 31 was prepared in a manner similar to that described in
Intermediate 6.

Example 1

4-[3-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,4]triazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester




A mixture of 4-(3-methanesulfonyloxy)methyl-[1,2,4]triazol- 1-yl)-piperidine-1 -
15 carboxylic acid *tert*-butyl ester (**Intermediate 1**, 0.222 g, 0.641 mmol), 2-Fluoro-4-
methanesulfonyl-phenol (0.122 g, 0.641 mmol) and CsCO₃ (0.250 g, 0.769 mmol) in
acetonitrile (10 mL) was heated under reflux for 4 hours. After cooling, the solution was
filtered through a pad of celite and the solvent was removed *in vacuo*. The residue was
purified by flash chromatography on silica gel with Hexanes and EtOAc to afford the
20 desired product. ¹H NMR (CDCl₃): δ 8.15 (1H, s), 7.62 (2H, m), 7.37 (1H, m), 5.25 (2H,
s), 4.25 (3H, m), 3.05 (3H, s), 2.85 (2H, m), 2.18 (2H, m), 1.97 (2H, m), 1.43 (9H, s).

The compounds in **Examples 2-8** were synthesized from one of **Intermediates 2-5** with the corresponding phenol in a manner similar to that described in **Example 1**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ⁱPr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields.

5 Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 2

4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxymethyl)-[1,2,3]triazol-2-yl]-piperidine-1-carboxylic acid *tert-butyl*| ester




¹H NMR (CDCl₃): δ 8.92 (1H, s), 7.72 (1H, s), 7.52 (1H, d), 7.42 (1H, d), 7.31

15 (IH, m), 5.30 (2H, s), 4.61 (IH, m), 4.20 (2H, m), 2.99 (2H, m), 2.16-2.08 (4H, m), 1.48 (9H, s).

Example 3

4-[4-(4-Tetrazol-1-yl-phenoxymethyl)-[1,2,3]triazol-2-yl]-piperidine-1-carboxylic acid *tert-butyl*| ester




¹H NMR (CDCl₃): δ 8.98 (1H, s), 7.7 (1H, s), 7.64 (2H, m), 7.18 (2H, m), 5.23 (2H, s), 4.63 (1H, m), 4.2 (2H, m), 3 (2H, m), 2.22-2.08 (4H, m), 1.48 (9H, s).

5

Example 4

4-[4-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,3]triazol-2-yl]-piperidine-1-carboxylic acid *tert-butyl* ester




10

¹H NMR (CDCl₃): δ 7.73-7.65 (3H, m), 7.27 (1H, m), 5.3 (2H, s), 4.61 (1H, m), 4.2 (2H, m), 3.05 (3H, s), 2.95 (2H, m), 2.2-2.1 (4H, m), 1.48 (9H, s).

Example 5

15


4-[5-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-tetrazol-2-yl]-piperidine-1-carboxylic acid *tert-butyl* ester



¹H NMR (CDCl₃): δ 7.72-7.62 (2H, m), 7.33 (1H, t), 5.46 (2H, s), 4.87 (1H, m), 4.2 (2H, m), 3.03 (3H, s), 2.95 (2H, m), 2.35-2.1 (4H, m), 1.46 (9H, s).

Example 6

5 4-[3-(2-Fluoro-4-tetrazol-1-yl-phenoxyethyl)-pyrazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester




10 ¹H NMR (CDCl₃): δ 8.79 (1H, s), 7.28-7.36 (4H, m), 6.24 (1H, s), 5.08 (2H, s), 4.11 (3H, m), 2.74 (2H, m), 1.97 (2H, m), 1.77 (2H, m), 1.36 (9H, s).

Example 7

4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxyethyl)-pyrazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester

15




18 ¹H NMR (CDCl₃): δ 8.92 (1H, s), 7.59 (1H, s), 7.55 (1H, s), 7.50 (1H, m), 7.42 (1H, m), 7.20 (1H, m), 5.10 (2H, s), 4.25 (3H, m), 2.87 (2H, m), 2.12 (2H, m), 1.89 (2H, m), 1.46 (9H, s).

Example 8

4-[4-(4-Tetrazol-1-yl-phenoxyethyl)-pyrazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester

5




¹H NMR (CDCl₃): δ 8.92 (1H, s), 7.60 (3H, m), 7.54 (1H, s), 7.11 (1H, m), 5.02 (2H, s), 4.25 (3H, m), 2.87 (2H, m), 2.12 (2H, m), 1.88 (2H, m), 1.46 (9H, s).

10

Example 9

5-Ethyl-2-{4-[3-(2-fluoro-4-methanesulfonyl-phenoxyethyl)-[1,2,4]triazol-1-yl]-piperidin-1-yl}-pyrimidine



15

A mixture of 4-[3-(2-fluoro-4-methanesulfonyl-phenoxyethyl)-[1,2,4]triazol-1-yl]-piperidine hydrochloride salt (**Intermediate 6**, 0.100g, 0.256 mmol), potassium carbonate (0.106g, 0.768 mmol) and 2-chloro-5-ethyl-pyrimidine (0.062 mL, 0.512 mmol) in dimethylformamide (2 mL) was heated at 90 °C for 4 hours or until the starting material was consumed. The reaction was cooled to room temperature, the solution was diluted with water and extracted with ethyl acetate. The organic layer was separated, washed with water and brine, dried over sodium sulfate, filtered and concentration *in vacuo*. The residue was chromatographed on silica gel (1:1 Hex/EtOAc) to afford the desired product. ¹H NMR (CDCl₃): δ 8.22 (2H, s), 8.17 (1H, s), 7.68 (2H, m), 7.38 (1H, m), 5.28 (2H, s), 4.92 (2H, d), 4.47 (1H, m), 3.08 (3H, s), 2.94 (2H, m), 2.48 (2H, q), 2.27 (1H, m), 2.18 (1H, m), 1.98 (2H, m), 1.20 (3H, t).


The compounds in **Examples 10-20** were synthesized from one of **Intermediates 7-12** with the corresponding substituted 2-fluoropyrimidine, 2-chloropyrimidine, 2-bromopyrimidine, 2-iodopyrimidine, 2-(methylsulfinyl)pyrimidine or 2-(methylsulfonyl)pyrimidine in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields.

5 Additionally, alternative coupling methods can be used that are well known in the art of
10 organic synthesis.

Example 10

5-Ethyl-2-{4-[4-(2-fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine


15



¹H NMR (CDCl₃): δ 8.21 (2H, s), 7.68 (3H, m), 7.28 (1H, m), 5.28 (2H, s), 4.82 (3H, m), 3.19 (2H, t), 3.02 (3H, s), 2.45 (2H, q), 2.23 (2H, m), 2.18 (2H, m), 1.20 (3H, t).

Example 11

20 5-Butyl-2-{4-[4-(2-fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine




¹H NMR (CDCl₃): δ 8.17 (2H, s), 7.68 (3H, m), 7.27 (1H, m), 5.30 (2H, s), 4.81-4.70 (3H, m), 3.16 (2H, t), 3.05 (3H, s), 2.44 (2H, t), 2.26 (2H, m), 2.16 (2H, m), 1.53 (2H, m), 1.32 (2H, m), 0.95 (3H, q).

Example 12

5-Ethyl-2-{4-[4-(2-fluoro-4-tetrazol-1-yl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine

10




¹H NMR (CDCl₃): δ 8.92 (1H, s), 8.20 (2H, s), 7.71 (1H, s), 7.52 (1H, dd), 7.42 (1H, m), 7.30 (1H, d), 5.30 (2H, s), 4.80-4.70 (3H, m), 3.17 (2H, m), 2.48 (2H, q), 2.28-2.11 (4H, m), 1.21 (3H, t).

15

Example 13


5-Chloro-2-{4-[4-(2-fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine



¹H NMR (CDCl₃): δ 8.25 (2H, s), 7.69 (3H, m), 7.28 (1H, m), 5.30 (2H, s), 4.76 (3H, m), 3.22 (2H, m), 3.04 (3H, s), 2.28-2.13 (4H, m).

Example 14

5 2-{4-[4-(2-Fluoro-4-methanesulfonyl-phenoxyethyl)-[1,2,3]triazol-2-yl]-piperidin-1-yl}-5-trifluoromethyl-pyrimidine




10 ¹H NMR (CDCl₃): δ 8.50 (2H, s), 7.68 (3H, m), 7.27 (1H, m), 5.30 (2H, s), 4.78 (3H, m), 3.29 (2H, m), 3.04 (3H, s), 2.31-2.20 (4H, m).

Example 15

15 2-{4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxyethyl)-[1,2,3]triazol-2-yl]-piperidin-1-yl}-5-trifluoromethyl-pyrimidine

15




¹H NMR (CDCl₃): δ 8.92 (1H, s), 8.49 (2H, s), 7.71 (1H, s), 7.52 (1H, dd), 7.42 (1H, m), 7.29 (1H, d), 5.29 (2H, s), 4.9-4.70 (3H, m), 3.3 (2H, m), 2.35-2.1 (4H, m).

20

Example 16

5-Chloro-2-{4-[4-(2-fluoro-4-tetrazol-1-yl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine




5

¹H NMR (CDCl₃): δ 8.93 (1H, s), 8.22 (2H, s), 7.7 (1H, s), 7.5 (1H, dd), 7.4 (1H, m), 7.29 (1H, d), 5.29 (2H, s), 4.8-4.65 (3H, m), 3.19 (2H, m), 2.3-2.1 (4H, m).

Example 17

10 5-Chloro-2-{4-[5-(2-fluoro-4-methanesulfonyl-phenoxy)methyl]-tetrazol-2-yl}-piperidin-1-yl}-pyrimidine




15

¹H NMR (CDCl₃): δ 8.24 (2H, s), 7.72-7.6 (2H, m), 7.33 (1H, t), 5.47 (2H, s), 5 (1H, m), 4.8-4.7 (2H, m), 3.24 (2H, m), 3.03 (3H, s), 2.36-2.28 (2H, m), 2.25-2.14 (2H, m).

Example 18

2-(4-((4-(1H-tetrazol-1-yl)phenoxy)methyl)-1 H-pyrazol-1-yl)piperidin-1-yl)-5-
ethylpyrimidine

20




¹H NMR (CDCl₃): δ 8.90 (1H, s), 8.19 (2H, s), 7.60-7.55 (3H, m), 7.54 (1H, s), 7.10 (2H, *J* = 9.2 Hz, d), 5.02 (2H, s), 4.87 (2H, m), 4.40 (1H, m), 3.04 (2H, m), 2.47 (2H, *J* = 7.2 Hz, q), 2.23 (2H, m), 1.96 (2H, m), 1.20 (3H, *J* = 7.2 Hz, t).

Example 19

5-Ethyl-2-(4-((2-fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-1H-pyrazol-1-yl)piperidin-1-ylpyrimidine

10




¹H NMR (CDCl₃): δ 8.92 (1H, s), 8.18 (2H, s), 7.58 (1H, s), 7.55 (1H, s), 7.49 (1H, m), 7.40 (1H, m), 7.19 (1H, m), 5.10 (2H, s), 4.86 (2H, m), 4.39 (1H, m), 3.03 (2H, m), 2.47 (2H, *J* = 7.6 Hz, q), 2.21 (2H, m), 1.95 (2H, m), 1.19 (3H, *J* = 7.6 Hz, t).

Example 20

2-{4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-5-methyl-pyrimidine

20




¹H NMR (CDCl₃): δ 8.86 (IH, s), 8.14 (2H, s), 7.64 (IH, s), 7.44 (IH, d), 7.35 (IH, d), 7.23 (IH, d), 5.23 (2H, s), 4.67 (2H, m), 4.15 (IH, m), 3.14 (2H, m), 2.21-2.12 (4H, m), 2.09 (3H, s).

5


Example 21

4-[4-(4-Methanesulfonyl-phenoxy-methyl)-[1,2,3]triazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester



10

Step 1: Preparation of 1-methanesulfonyl-4-prop-2-ynyoxy-benzene




15

A mixture of 4-methylsulfonyl-phenol, propargyl bromide and K₂CO₃ was refluxed overnight. After cooling, the mixture was filtered through a pad of celite. The filtrate was concentrated *in vacuo*. The crude solid product was purified on silica gel to give the desired product as a pale yellow solid. ¹H NMR (CDCl₃): δ 7.9 (2H, d), 7.12 (2H, d), 4.79 (2H, d), 3.05 (3H, s), 2.58 (IH, t).


20

Step 2: Preparation of 4-azido-piperidine-1-carboxylic acid *tert*-butyl ester



A mixture of 4-methanesulfonyloxy-piperidine-1-carboxylic acid *tert*-butyl ester and sodium azide in DMF was stirred at 60°C overnight. The mixture was poured into water and the resulting mixture was extracted with EtOAc. The organic phase was separated and washed with water and brine, dried over Na_2SO_4 , filtered and concentrated *in vacuo* to give the desired product as a pale yellow oil, which was used without further purification. ^1H NMR (CDCl_3): δ 7.9 (2H, d), 7.12 (2H, d), 4.79 (2H, d), 3.05 (3H, s), 2.58 (1H, t).

Step 3: Preparation of 4-[4-(4-Methanesulfonyl-phenoxymethyl)-[1,2,3]triazol-1-yl]-piperidine-1-carboxylic acid *tert*-butyl ester




4-Azido-piperidine-1-carboxylic acid *tert*-butyl ester (1 eq) and 1-methanesulfonyl-4-prop-2-ynylbenzene (1 eq) were suspended in $\text{H}_2\text{O}/\text{t-BuOH}$ (1:1). Sodium ascorbate (0.1 eq) was added followed by $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (0.01 eq). The mixture was stirred vigorously overnight. The mixture was diluted with water and a precipitate formed. The precipitate was collected by filtration and purified on silica gel to give the desired product. ^1H NMR (CDCl_3): δ 7.88 (2H, d), 7.67 (1H, s), 7.14 (2H, d), 5.29 (2H, s), 4.64 (1H, m), 4.28 (2H, m), 3.05 (3H, s), 2.94 (2H, m), 2.2 (2H, m), 1.96 (2H, m), 1.48 (9H, s).

The compounds in **Examples 22-23** were synthesized from **Intermediate 15** with the corresponding substituted 2-methanesulfonyl-pyrimidine or 2-iodopyrimidine in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, 'Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 22


10 2-{4-[4-(6-Tetrazol-1-yl-pyridin-3-yloxymethyl)-[1,2,3]triazol-2-yl]-piperidin-1-yl}-5-trifluoromethyl-pyrimidine



15 ¹H NMR (DMSO-^δ): δ 10.06 (1H, s), 8.7 (2H, s), 8.4 (1H, d), 7.99 (1H, d), 7.94 (1H, s), 7.86 (1H, dd), 5.34 (2H, s), 4.92 (1H, m), 4.68 (2H, m), 3.33 (2H, m), 2.22 (2H, m), 1.94 (2H, m).

Example 23

20 5-Chloro-2- {4-[4-(6-tetrazol-1-yl-pyridin-3-yloxyethyl)-[1,2,3]triazol-2-yl] -piperidin-1-yl}-pyrimidine




¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.27 (1H, s), 8.26 (2H, s), 8.02 (1H, d), 7.71 (1H, s), 7.56 (1H, dd), 5.28 (2H, s), 4.75 (3H, m), 3.23 (2H, m), 2.29 (2H, m), 2.15 (2H, m).

The compounds listed in **Examples 24-28** are synthesized in a manner similar to
5 that described in **Example 9** by using substituted 5-chloro-oxadiazoles in place of
substituted 2-chloropyrimidines.

Example 24

1-[3-(1-Fluoro-1-methyl-ethyl)-[1,2,4]oxadiazol-5-yl]-4-[4-(2-fluoro-4-tetrazol-1-yl-phenoxy)methyl]- [1,2,3]triazol-2-yl] -piperidine


10



Example 25

1-(3-Cyclopropyl-[1,2,4]oxadiazol-5-yl)-4-[4-(2-fluoro-4-tetrazol-1-yl-phenoxy)methyl]- [1,2,3]triazol-2-yl] -piperidine


15



Example 26

4-[4-(2-Fluoro-4-tetrazol-1-yl-phenoxy)methyl]- [1,2,3]triazol-2-yl] -1-(3-isopropyl-[1,2,4]oxadiazol-5-yl)-piperidine


20



Example 27

5-{2-[1-(3-Isopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yl]-2H-[1,2,3]triazol-4-ylmethoxy}-2-tetrazol-1-yl-pyridine


5



Example 28

5-(2-{1-[3-(1-Fluoro-1-methyl-ethyl)-[1,2,4]oxadiazol-5-yl]-piperidin-4-yl}-2H-[1,2,3]triazol-4-ylmethoxy)-2-tetrazol-1-yl-pyridine


10



Example 29

5-{2-[1-(3-Cyclopropyl-[1,2,4]oxadiazol-5-yl)-piperidin-4-yl]-2H-[1,2,3]triazol-4-ylmethoxy}-2-tetrazol-1-yl-pyridine

15



To a solution of 5-((2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (**0.500 g, 2.05 mmol**) (**Intermediate 13**), in THF (4 mL) was added 4-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)cyclohexanol (**Intermediate 14**), polymer supported triphenylphosphine (1.05 g, 3.07 mmol) and diisopropylazodicarboxylate (0.603 mL, 3.07 mmol). The suspension was stirred for 16 hours and filtered through a pad of celite. The filtrate was concentrated *in vacuo* and chromatographed on silica gel (1:1 hexanes/ethyl acetate) to obtain the expected product. ¹H NMR (CDCl₃): δ 9.45 (1H, s), 8.28 (1H, s), 7.03 (1H, d), 7.72 (1H, s), 7.58 (1H, dd), 5.28 (2H, s), 4.72 (1H, m), 4.17 (2H, m), 3.34 (2H, m), 2.27 (4H, m), 1.89 (1H, m).


10

The compounds in **Examples 30-34** were synthesized from one of **Intermediates 6-12, 15 or 19** with the corresponding substituted 2-fluoropyrimidine, 2-chloropyrimidine, 2-bromopyrimidine, 2-iodopyrimidine, 2-(methylsulfinyl)pyrimidine or 2-(methylsulfonyl)pyrimidine in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

20

Example 30

5-Fluoro-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine




25

¹H NMR (CDCl₃): δ 8.91 (1H, s), 8.21 (2H, s), 7.72 (1H, s), 7.46 (1H, d), 7.41 (1H, d), 7.23 (1H, m), 5.23 (2H, s), 4.72 (3H, m), 3.19 (2H, m), 2.22 (2H, m), 2.19 (2H, m).

Example 31

2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-fluoropyrimidine




5

¹H NMR (CDCl₃): δ 9.43 (1H, s), 8.23 (1H, d), 8.21 (2H, s), 8.01 (1H, d), 7.71 (1H, s), 7.59 (1H, dd), 5.23 (2H, s), 4.77 (3H, m), 3.20 (2H, m), 2.29 (2H, m), 2.18 (2H, m).

Example 32

10 2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-ethylpyrimidine




15

¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.27 (1H, d), 8.20 (2H, s), 8.02 (1H, d), 7.70 (1H, s), 7.59 (1H, dd), 5.28 (2H, s), 4.77 (3H, m), 3.20 (2H, m), 2.50 (2H, q), 2.22 (2H, m), 2.14 (2H, m), 1.20 (3H, t).

Example 33


20 5-Chloro-2-((^trans)-3-fluoro-4-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine



¹H NMR (CDCl₃): δ 8.93 (1H, s), 8.28 (2H, s), 7.77 (1H, s), 7.52 (1H, dd), 7.41 (1H, dd), 7.27 (1H, m), 5.31 (2H, s), 5.08 (1H, m), 4.91 (1H, m), 4.75 (2H, m), 3.23 (2H, m), 2.26 (2H, m).

Example 34

2-((^{trans}-3-Fluoro-4-(4-((2-fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)-5-(trifluoromethyl)pyrimidine




¹H NMR (CDCl₃): δ 8.91 (1H, s), 8.54 (2H, s), 7.77 (1H, s), 7.52 (1H, dd), 7.41 (1H, d), 7.27 (1H, m), 5.31 (2H, s), 5.10 (1H, m), 4.93 (1H, m), 4.82 (2H, m), 3.34 (2H, m), 2.26 (2H, m).

15

Example 35

Isopropyl 4-((6-(1H-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2H-1,2,3-triazol-2-yl)piperidine-1-carboxylate




To a suspension of 5-((2-(piperidin-4-yl)-2H-1,2,3-triazol-4-yl)methoxy)-2-(1H-tetrazol-1-yl)pyridine (**Intermediate 15**) (0.300 g, 0.917 mmol) in ethyl acetate (3 mL) was added triethylamine (0.169 mL, 1.22 mmol) and isopropyl chloroformate (1.10 mL, 1M solution in toluene, 1.10 mmol). The resulting solution was stirred for 1 hour. Water was added and the solution was extracted with ethyl acetate. The organic layer was separated, dried over sodium sulfate, filtered and concentrated *in vacuo*. The residue was chromatographed on silica gel (1:1 hexanes/ethyl acetate) to obtain the expected product.

5 ¹H NMR (CDCl_3): δ 9.45 (1H, s), 8.28 (1H, d), 8.03 (1H, d), 7.71 (1H, s), 7.58 (1H, dd), 5.28 (2H, s), 4.95 (1H, m), 4.64 (1H, m), 4.22 (1H, m), 2.21 (2H, m), 2.14 (2H, m), 1.26 (6H, d).

Example 36

15 Isopropyl 4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidine-1-carboxylate




20 The title compound was synthesized using 4-[4-(2-Fluoro-4-methanesulfonyl-phenoxy)methyl]-[1,2,3]triazol-2-yl]-piperidine (**Intermediate 7**) and isopropyl chloroformate in a manner similar to that described in **Example 35**. ¹H NMR (CDCl_3): δ

7.66 (3H, m), **7.27** (IH, m), **5.30** (2H, s), **4.94** (IH, m), **4.63** (IH, m), **4.20** (2H, m), **3.04** (3H, s), 2.19 (2H, m), 2.17 (2H, m), 1.25 (6H, d).

Example 37


Isopropyl 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-
5 yl)piperidine-1-carboxylate



The title compound was synthesized using 1-((2*H*-1,2,3-triazol-4-yl)methoxy)-
10 3-fluorophenyl-1*H*-tetrazole (**Intermediate 8**) and isopropyl chloroformate in a manner
similar to that described in **Example 35**. **¹H NMR** (CDCl₃): δ **8.93** (IH, s), 7.71 (IH, s),
7.52 (IH, dd), **7.43** (IH, m), 7.31 (IH, t), **5.3** (2H, s), **4.94** (IH, m), **4.63** (IH, m), **4.2** (2H,
br), **3.03** (2H, m), 2.15 (4H, m), 1.26 (6H, d).

Example 38

15 5-Ethyl-2-((^rans)-3-fluoro-4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-
1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine




20 The title compound was synthesized using 1-((2*H*-1,2,3-triazol-4-yl)methoxy)-
3-fluorophenyl-1*H*-tetrazole (**Intermediate 16**) and *cis*-1-(5-ethylpyrimidin-2-yl)-3-
fluoropiperidin-4-ol (**Intermediate 17**) in a manner similar to that described in **Example**

29. ^1H NMR (CDCl_3): δ 8.92 (IH, s), 8.21 (2H, s), 7.76 (IH, s), 7.51 (IH, dd), 7.41 (IH, d), 7.26 (IH, m), 5.31 (2H, s), 5.17 (IH, m), **5.03** (IH, m), 4.83 (2H, m), 3.11 (2H, m), **2.50** (2H, q), 2.22 (2H, m), 1.22 (3H, t).

Example 39

5 5-(4-((4-((1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-ylsulfonyl)-2,4-dimethylthiazole




10 To a solution of 5-((2-(piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine hydrochloride (**Intermediate 15**, 0.050 g, 0.153 mmol), dichloromethane (1.5 mL), and triethyl amine (0.043 mL, 0.306 mmol) was added 2,4-dimethyl-thiazole-5-sulfonyl chloride (0.049 g, 0.229 mmol) and the reaction stirred at room temperature for 2 hours. The solution was concentrated and purified by flash column chromatography on silica gel eluting with 0-100% ethyl acetate in hexanes to afford the desired product.

15 ^1H NMR ($\text{DMSO}-\text{C}_6\text{F}_5$): δ 10.01 (IH, s), 8.41 (IH, d), 8.00 (IH, d), 7.94 (IH, s), 7.85 (IH, dd), 5.34 (2H, s), 4.66 (IH, m), 3.69 (2H, m), 2.81 (2H, m), 2.64 (3H, s), 2.51 (3H, s), 2.24 (2H, m), 2.02 (2H, m).

20

Example 40


5-((2-(1*H*-Imidazol-4-ylsulfonyl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine



The title compound was synthesized using **Intermediate 15** and 1*H*-imidazole-4-sulfonyl chloride in a similar manner as described in **Example 39**. **¹H NMR** (DMSO-*d*₆): δ 12.89 (1H, **bs**), 10.07 (1H, **s**), 8.41 (1H, **d**), **7.99** (1H, **d**), **7.92** (1H, **s**), **7.86** (2H, **m**), **7.79**, (1H, **dd**), **5.32** (2H, **s**), **4.59** (1H, **m**), **3.62** (2H, **m**), **2.77** (2H, **m**), 2.17 (2H, **m**), 1.95 (2H, **M**).

Example 41


10 5-((2-(1-(Cyclopropylsulfonyl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine



15 The title compound was synthesized using **Intermediate 15** and cyclopropanesulfonyl chloride in a similar manner as described in **Example 39**. **¹H NMR** (DMSO-*d*₆): δ **9.45** (1H, **s**), **8.28** (1H, **d**), **8.03** (1H, **d**), **7.73** (1H, **s**), **7.58** (1H, **dd**), **5.28** (2H, **s**), **4.62** (1H, **m**), **3.87** (2H, **m**), 3.12 (2H, **m**), 2.31 (**5H, m**), 1.86 (2H, **m**), 1.02 (2H, **m**).

Example 42


20 5-((2-(1-Methyl-1*H*-imidazol-4-ylsulfonyl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine



The title compound was synthesized using **Intermediate 15** and 1-methyl- 1-imidazole-4-sulfonyl chloride in a similar manner as described in **Example 39**. ¹H NMR (*DMSO-Cl₆*) : δ 10.07 (IH, s), 8.41 (IH, d), 8.00 (IH, d), 7.93 (IH, s), 7.87 (IH, dd), 7.82 (2H, s), 5.34 (2H, s), 4.59 (IH, m), 3.69 (3H, s), 3.36 (2H, m), 2.74 (2H, m), 2.19 (2H, m), 2.00 (2H, m).

Example 43

10 4-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-ylsulfonyl)-3,5-dimethylisoxazole




15 The title compound was synthesized using **Intermediate 15** and 3,5-dimethylisoxazole -4-sulfonyl chloride in a similar manner as described in **Example 39**. ¹H NMR (*DMSO-Cl₆*) : δ 10.07 (IH, s), 8.41 (IH, d), 8.00 (IH, d), 7.94 (IH, s), 7.86 (IH, dd), 5.34 (2H, s), 4.65 (IH, m), 3.68 (2H, m), 2.86 (2H, m), 2.61 (3H, s), 2.33 (3H, s), 2.24 (2H, m), 2.00 (2H, m).

20

Example 44


5-((2-(1-(Cyclohexylsulfonyl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine



5 The title compound was synthesized using **Intermediate 15** and cyclohexanesulfonyl chloride in a similar manner as described in **Example 39**. ¹H NMR (DMSO-d₆): δ 10.08 (IH, s), 8.42 (IH, d), 8.01 (IH, d), 7.95 (IH, s), 7.86 (IH, dd), 3.52 (2H, s), 4.74 (IH, m), 3.68 (2H, m), 3.11 (3H, m) 2.51 (IH, m), 2.17 (2H, m) 1.92 (3H, m), 1.73 (IH, m), 1.23 (7H, m).

Example 45


10 5-((2-(1-(Isopropylsulfonyl)piperidin-4-yl)-2-(1H-1,2,3-triazol-4-yl)methoxy)-2-(1H-tetrazol-1-yl)pyridine



15 The title compound was synthesized using **Intermediate 15** and isopropylsulfonyl chloride in a similar manner as described in **Example 39**. ¹H NMR (DMSO-d₆): δ 10.08 (IH, s), 8.41 (IH, d), 8.01 (IH, d), 7.95 (IH, s), 7.86 (IH, dd), 5.35 (2H, s), 4.75 (IH, m), 3.69 (2H, m), 3.31 (IH, m), 3.13 (2H, m), 2.17 (2H, m), 1.97 (2H, m), 1.20 (6H, d).

Example 46


20 5-((2-(1-(Butylsulfonyl)piperidin-4-yl)-2-(1H-1,2,3-triazol-4-yl)methoxy)-2-(1H-tetrazol-1-yl)pyridine



The title compound was synthesized using **Intermediate 15** and 1-butanesulfonyl chloride in a similar manner as described in **Example 39**. **¹H NMR** (DMSO-*d*₆): δ 10.08 (1H, s), 8.41 (1H, d), **8.00** (1H, d), 7.96 (1H, s), 7.86 (1H, dd), 5.36 (2H, s), 4.71 (1H, m), 3.63 (2H, m), 3.01 (4H, m), **2.2** (2H, m), 1.99 (2H, m), 1.64 (2H, m), 1.39 (2H, m), **0.89** (3H, t).

Example 47

10 5-((2-(1-(Benzylsulfonyl)piperidin-4-yl)-2 *H*-1,2,3-triazol-4-yl)methoxy)-2-(1 *H*-tetrazol-1-yl)pyridine




15 The title compound was synthesized using **Intermediate 15** and phenylmethanesulfonyl chloride in a similar manner as described in **Example 39**. **¹H NMR** (DMSO-*d*₆): δ 10.07 (1H, s), 8.41 (1H, d), 8.00 (1H, d), 7.96 (1H, s), 7.86 (1H, dd), 7.36 (5H, m), 5.35 (2H, s), 4.68 (1H, m), 4.43 (2H, s), 3.54 (2H, m), 2.99 (2H, m), 2.12 (2H, m), 1.91 (2H, m).

20


Example 48

2-(4-((2-Methylpyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-ylsulfonyl)-5-(trifluoromethyl)pyrimidine



5

Step 1: 2-Methyl-3-(prop-2-ynyloxy)pyridine




10

The title compound was synthesized using 3-hydroxy-2-methylpyridine and propargyl chloride in a manner similar to that described in **Intermediate 13, Step 2.**


Step 2: (5-((2-Methylpyridin-3-yloxy)methyl)-1*H*-1,2,3-triazol-1-yl)methyl

15 pivalate




The title compound was synthesized using 2-methyl-3-(prop-2-ynyloxy)pyridine and 20 azidomethyl pivalate in a manner similar as that described in **Intermediate 13, Step 3.**

Step 3: 3-((1*H*-1,2,3-Triazol-5-yl)methoxy)-2-methylpyridine



The title compound was synthesized using (5-((2-methylpyridin-3-yloxy)methyl)-1H-1,2,3-triazol-1-yl)methyl pivalate and sodium hydroxide in a manner similar as that described in **Intermediate 13, Step 4.**

Step 4: 2-(4-((2-Methylpyridin-3-yloxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)-5-(trifluoromethyl)pyrimidine




10

To a solution of 3-((1H-1,2,3-triazol-5-yl)methoxy)-2-methylpyridine (0.20 g, 1.05 mmol), in THF (2 mL) at 0 °C, was added 1-(5-(trifluoromethyl)pyrimidin-2-yl)piperidin-4-ol (0.391 g, 1.58 mmol) (**Intermediate 20**), polymer bound triphenylphosphine (0.527 g, 1.58 mmol), and di-*tert*-butylazodicarboxylate (0.363, 1.58 mmol). The reaction was allowed to warm to room temperature overnight and was then filtered through a pad of celite washing with THF. The solution was concentrated *in vacuo* and purified by flash column chromatography on silica gel eluting with 0-60% ethyl acetate in hexane to afford the desired product. ¹H NMR (DMSO-d₆): δ 8.71 (2H, s), 8.00 (1H, dd), 7.89 (1H, s), 7.47 (1H, dd), 7.19 (1H, dd), 5.18 (2H, s), 4.90 (1H, m), 4.68 (2H, s), 3.31 (2H, m), 2.31 (3H, s), 2.19 (2H, m), 1.92 (2H, m).

Example 49

(*ta ns*)-*tert*-Butyl 3-fluoro-4-(4-((2-fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidine-1-carboxylate

25




The title compound was synthesized using **Intermediate 18** and **Intermediate 16** in a similar manner to that described in Example 29. ^1H NMR (CDCl_3): δ 8.98 (1H, s), 7.74 (1H, s), 7.51 (1H, dd), 7.43 (1H, m), 7.26 (1H, t), 5.29(2H, s), 4.94 (1H, m), 4.7 (1H, m), 4.43 (1H, m), 4.1 (1H, m), 2.96 (2H, m), 2.16 (2H, m), 1.45 (9H, s).

Example 50

tert-Butyl 4-(2-((6-(1 *H*-tetrazol-1-yl)pyridine-3-yloxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)-2-oxoethyl)piperidine-1-carboxylate

10




To a stirred solution of 5-((2-(Piperidin-4-yl)-2 *H*-1,2,3-triazol-4-yl)methoxy)-2-(1 *H*-tetrazol-1-yl)pyridine (225 mg, 0.688 mmol) (**Intermediate 15**) dissolved in DMF (10mL) was added triethylamine (4 eq.) and o-benzotriazol-1-yl- N,N,N',N' -tetramethyluronium tetrafluoroborate (TBTU) (2 eq.). The solution was stirred for 5 minutes and 2- $\{\text{O}\text{-}\text{tert}$ -butoxycarbonyl)piperidin-4-yl)acetic acid (1.5 eq.) was added and was stirred at room temperature for 3 h. The reaction was quenched with water and extracted with ethyl acetate. The organic layer was washed with brine, and dried over sodium sulfate. The solution was filtered and concentrated *in vacuo*. The crude product was purified by flash column chromatography on silica gel eluting with ethyl acetate and hexanes to afford the desired product. ^1H NMR (DMSO- \wedge): δ 10.07 (1H, s), 8.40 (1H, d), 8.00 (1H, d), 7.93 (1H, s), 7.86 (1H, dd), 5.34 (2H, s), 4.78 (1H, m), 4.35 (1H, m), 3.85-3.94 (3H, m), 3.21 (1H, m),

2.64-2.82 (3H, m), 2.27 (2H, d), 2.10 (2H, m), 1.86 (3H, m), 1.60 (2H, d), 1.35 (9H, s), 1.99 (2H, m).

Example 51


5 1-(4-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-2-(piperidin-4-yl)ethanone hydrochloride



10 To a solution of *tert*-butyl 4-(2-(4-((6-(1*H*-tetrazol-1-yl)pyridine-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-2-oxoethyl)piperidine-1-carboxylate (110 mg, 0.20 mmol) in dichloromethane (5 mL) cooled to 0 °C was added hydrochloric acid (4M in 1,4-dioxane, 5 eq.) dropwise. After the reaction was stirred for 3 h the solvents were removed *in vacuo* to afford the desired HCl salt. ¹H NMR (DMSO- δ): δ 10.06 (1H, s), 8.78 (1H, br), 8.40 (1H, d), 7.98 (1H, d), 7.94 (1H, s), 7.87 (1H, dd), 5.35 (2H, s), 4.79 (1H, m), 4.34 (1H, m), 3.92 (1H, m), 3.20 (3H, m), 2.83 (2H, m), 2.31 (2H, d), 2.10 (2H, m), 1.93 (2H, m), 1.77 (3H, m), 1.32 (2H, m).

Example 52

20 Methyl 2-(2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidin-5-yl)acetate




To a solution of 5-((2-(Piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (85 mg, 0.26 mmol) in DMF (5 mL) was added triethylamine (3 eq.)

and methyl 2-(2-(methylsulfonyl)pyrimidin-5-yl)acetate (2 eq.). The reaction was stirred overnight at room temperature, then was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated *in vacuo*. The crude material was purified by flash chromatography on silica gel eluting with ethyl acetate and hexanes to afford the title compound. ^1H NMR (CDCl_3): δ 9.44 (1H, s), 8.27 (1H, d), 8.25 (2H, s), 8.02 (1H, d), 7.70 (1H, s), 7.59 (1H, dd), 5.27 (2H, s), 4.78 (3H, m), 3.45 (2H, s), 3.36 (3H, s), 3.20 (2H, m), 2.27 (2H, m), 2.17 (2H, m).

Example 53

(S)-*tert*-Butyl 1-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-1-oxopropan-2-ylcarbamate




The title compound was prepared in a manner analogous to **Example 50** above. ^1H

15 NMR (CDCl_3): δ 9.45 (1H, s), 8.28 (1H, d), 8.03 (1H, d), 7.72 (1H, s), 7.58 (1H, dd), 5.52 (1H, m), 5.28 (2H, s), 4.71 (2H, m), 4.41-4.67 (1H, m), 4.04 (1H, m), 2.97-3.40 (2H, m), 2.25 (4H, m), 1.44 (9H, s), 1.32 (3H, m).

Example 54

(S)-*tert*-Butyl 1-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-1-oxopropan-2-ylcarbamate




The title compound was prepared in a manner similar to that described in

Example 50. $^1\text{H NMR}$ (CDCl_3): δ 9.44 (1H, s), 8.27 (1H, d), 8.03 (1H, d), 7.72 (1H, s), 7.58 (1H, dd), 5.52 (1H, m), 5.29 (2H, s), 4.71 (2H, m), 4.45 (1H, m), 3.99 (1H, m), 5 2.96-3.1 1 (2H, m), 2.19 (4H, m), 1.44 (9H, s), 1.32 (3H, m).

Example 55

(*S*)-1-(4-((6-(1 *H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)-2-aminopropan-1-one hydrochloride


10



The title compound was prepared from (*S*)-*tert*-butyl 1-(4-((6-(1 *H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)-1-oxopropan-2-ylcarbamate (**Example 54**) in a manner similar to that described in Example 51. LCMS: 15 (ES⁺-MS found: 398.6).

Example 56

(*S*)-*tert*-Butyl 4-(4-((6-(1 *H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)-4-oxo-1-(2,4,5-trifluorophenyl)butan-2-ylcarbamate




The title compound was prepared in a manner similar to that described in

Example 50. ^1H NMR (CDCl_3): δ 9.44 (1H, s), 8.27 (1H, s), 8.03 (1H, d), 7.72 (1H, d), 7.58 (1H, dd), 7.08 (1H, m), 6.90 (1H, m), 5.58 (1H, m), 5.28 (2H, s), 4.71 (1H, m), 4.54 (1H, m), 4.13 (1H, m), 3.87 (1H, m), 3.25 (1H, m), 2.96 (3H, m), 2.60 (2H, m), 2.1 1-2.27 (4H, m), 1.37 (9H, s).

Example 57


(*S*)-1-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-3-amino-4-(2,4,5-trifluorophenyl)butan-1-one hydrochloride



The title compound was prepared from (*S*-*tert*-Butyl 4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-4-oxo-1-(2,4,5-trifluorophenyl)butan-2-ylcarbamate (**Example 56**) in a manner similar to that described in Example 51. LCMS: (ES⁺-MS found: 542.6).

Example 58

Ethyl 4-(2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidin-5-yl)butanoate




The title compound was prepared from **Intermediate 15** in a manner similar to
5 that described in **Example 9**. LCMS (ES⁺-MS found: 520.0).

Example 59

(ta ns)-1-*tert*-Butyl 2-methyl 4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidine-1,2-dicarboxylate


10



To a solution of 5-((2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (600 mg, 2.45 mmol), triphenylphosphine (1.5 eq.), and (2*R*,4*S*)-*N*-Boc-4-hydroxypiperidine-2-carboxylic acid methyl ester (1.5 eq.) dissolved in THF (20mL) at 0 °C
15 was added a solution of di-*tert*butylazodicarboxylate (1.5eq.) in THF (5 mL) drop wise. The reaction was allowed to warm to room temperature overnight and the solvents removed *in vacuo*. The crude material was purified by flash column chromatography on silica gel to afford the desired product. ¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.27 (1H, d), 8.03 (1H, d), 7.71 (1H, s), 7.58 (1H, dd), 5.27 (2H, s), 4.99-5.19 (1H, m), 4.57 (1H, m), 4.11-4.31 (1H, m), 3.79 (3H, s), 3.04-3.24 (1H, m), 2.75 (1H, m), 2.18-2.32 (2H, m), 2.05 (1H, m), 1.50 (9H, s). ES⁺-MS found: 486.1.

Example 60

Ethyl 4-(2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylate




5

The title compound was prepared in a manner similar to that described in

Example 52 above. ¹H NMR (CDCl₃): δ 9.45 (1H, s), 8.86 (2H, s), 8.27 (1H, d), 8.03 (1H, d), 7.72 (1H, s), 7.58 (1H, dd), 5.27 (2H, s), 4.92 (2H, m), 4.80 (1H, m), 4.35 (2H, q), 3.31 (2H, m), 2.32 (2H, m), 2.18 (2H, m), 1.37 (3H, t). LCMS: (ES⁺MS found: 478.2).


Example 61

2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidin-5-yl)(morpholino)methanone




15

Step 1: Preparation of 2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylic acid



Ethyl 4-(2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylate (0.8g, 0.00167mol) (**Example 60**) was dissolved in MeOH (20mL). An aqueous solution of NaOH (10eq., 40 nL) was added and the reaction mixture was heated at 60 °C for three hours. The methanol was removed *in vacuo* and the remaining reaction mixture was cooled to 0 °C and concentrated HCl was added drop wise until the pH reached 2-3. The aqueous solution was extracted with DCM, dried over sodium sulfate, filtered, and concentrated *in vacuo* to afford the title compound.

Step 2 : Preparation of 2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidin-5-yl)(morpholino)methanone




The title compound was prepared from 2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylic acid (**Step 1**) in a manner similar to that described in **Example 50** above. ¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.46 (2H, s), 8.27 (1H, d), 8.02 (1H, d), 7.71 (1H, s), 7.58 (1H, dd), 5.27 (2H, s), 4.82 (3H, m), 3.72 (8H, m), 3.28 (2H, m), 2.30 (2H, m), 2.23 (2H, m). LCMS: (ES⁺-MS found: 519.0).

20

Example 62


(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)(5-bromopyrimidin-2-yl)methanone



The title compound was prepared from 2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylic acid (**Example 61, Step 1**) in a manner similar to that described in **Example 50** above. LCMS: (ES⁺-MS found: 512.0 & 514.0).

Example 63

tert-Butyl 4-((6(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)octahydroquinoline-1(2*H*)-carboxylate




10

The title compound was prepared in a manner similar to that described in **Example 59**. ¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.25 (1H, s), 8.03 (1H, d), 7.65 (1H, s), 7.61 (1H, dd), 5.27 (2H, s), 4.93 (1H, m), 4.11-4.13 (1H, m), 3.40 (1H, m), 3.12 (1H, m), 2.49 (1H, m), 2.41 (1H, m), 2.15 (1H, m), 1.96 (1H, m), 1.50-1.74 (3H, m), 1.50 (9H, s), 1.08-1.34 (3H, m), 0.49 (1H, m).

Example 64

2-(4-((2-Methyl-6-(1*H*-1,2,4-triazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-(trifluoromethyl)pyrimidine

20




The title compound was prepared from **Intermediate 26** and 2-methyl-6-(1*H*-1,2,4-triazol-1-yl)pyridin-3-ol in a manner similar to that described in **Example 1**.

¹H NMR (CDCl₃): δ 9.08 (1H, s), 8.50 (2H, s), 8.06 (1H, s), 7.69 (1H, s), 7.67 (1H, m), 7.40 (1H, d), 5.21 (2H, s), 4.87-4.79 (3H, m), 3.31 (2H, m), 2.49 (3H, s), 2.29 (2H, m), 2.17 (2H, m). LCMS: (ES⁺-MS found: 487.1).

Example 65

5-Ethyl-2-(4-((2-methyl-6-(1*H*-1,2,4-triazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine




The title compound was prepared from **Intermediate 30** and 2-methyl-6-(1*H*-1,2,4-triazol-1-yl)pyridin-3-ol in a manner similar to that described in **Example 1**.

¹H NMR (CDCl₃): δ 9.07 (1H, s), 8.20 (2H, s), 8.05 (1H, s), 7.67 (1H, s), 7.65 (1H, d), 7.40 (1H, d), 5.20 (2H, s), 4.79-4.71 (3H, m), 3.18 (2H, m), 2.49 (2H, m), 2.48 (3H, s), 2.28 (2H, m), 2.21 (2H, m), 1.20 (3H, m). LCMS: (ES⁺-MS found: 446.8).


Example 66

5-Ethyl-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)pyrimidine




Step 1: Ethyl 1-(l-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxylate

5




The compound was synthesized using ethyl 3-(trifluoromethyl)-1*H*-pyrazole-4-carboxylate and 1-(5-ethylpyrimidin-2-yl)piperidin-4-yl methanesulfonate in a manner similar to that described in **Intermediate 1, Step 1**. The reaction afforded two regioisomers, ethyl 1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-3 -(trifluoromethyl)- 1*H*-pyrazole-4-carboxylate and ethyl 1-(I -(5-ethylpyrimidin-2-yl)piperidin-4-yl)-5 -(trifluoromethyl)- 1*H*-pyrazole-4-carboxylate which were separated by flash column chromatography on silica gel.

15 **Step 2:** (l-(l-(5-Ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazol-4-yl)methanol




The compound was synthesized using ethyl 1-(1-(5-Ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxylate and sodium borohydride in a manner similar to that described in *Journal of Medicinal Chemistry*, **1996**, **29**, 341.

5 **Step 3:** (1-(1-(5-Ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazol-4-yl)methyl methanesulfonate



10 The title compound was synthesized using (1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazol-4-yl)methanol in a manner similar to that described in **Intermediate 1, Step 3**.

15 **Step 4:** 5-Ethyl-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)pyrimidine




20 The compound was synthesized using (1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-3-(trifluoromethyl)-1*H*-pyrazol-4-yl)methyl methanesulfonate and 2-fluoro-4-(1*H*-tetrazol-1-yl)phenol in a manner similar to that described in **Example 1**. ¹H NMR (CDCl₃): δ 8.99 (1H, s), 8.15 (2H, s), 7.64 (1H, s), 7.50 (1H, m), 7.41 (1H, m), 7.17 (1H, m), 5.13

(2H, s), **4.86** (2H, m), **4.42** (1H, m), **2.99** (2H, m), **2.44** (2H, q), **2.20** (2H, m), 1.93 (2H, m), 1.16 (3H, t).

Example 67

5-Ethyl-2-(4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-3-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)pyrimidine




The compound was synthesized using (1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-10 3-(trifluoromethyl)-1*H*-pyrazol-4-yl)methyl methanesulfonate and 2-fluoro-4-(methylsulfonyl)phenol in a manner similar to that described in **Example 1**. **¹H NMR** (CDCl_3): δ **8.25** (2H, s), 8.17 (1H, s), 7.71 (2H, m), **7.52** (1H, m), **5.30** (2H, s), **4.90** (2H, m), **4.65** (1H, m), 3.13 (3H, s), **3.05** (2H, m), **2.47** (2H, q), **2.20** (2H, m), 1.98 (2H, m), 1.18 (3H, t).

15


Example 68

5-Ethyl-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-5-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)pyrimidine




20

Step 1: ethyl 1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-5-(trifluoromethyl)-1*H*-pyrazole-4-carboxylate




The title compound was obtained in **Example 66, Step 1.**

5 **Step 2:** (1-(1-(5-Ethylpyrimidin-2-yl)piperidin-4-yl)-5-(trifluoromethyl)-1 *H*-pyrazol-4-yl)methanol



10 The title compound was synthesized using ethyl 1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-5-(trifluoromethyl)-1 *H*-pyrazole-4-carboxylate and sodium borohydride in a manner similar to that described in **Example 66, Step 2.**


15 **Step 3:** 2-(4-(4-(chloromethyl)-5 -(trifluoromethyl)-1*H*-pyrazol- 1-yl)piperidin- 1-yl)-5-ethylpyrimidine



20 To a solution (1-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-5 -(trifluoromethyl)-1*H*-pyrazol-4-yl)methanol (0.13 g, 0.366 mmol) in dichloromethane (3 mL) was added thionyl chloride (1.2 mL) and the solution was stirred at room temperature for 2 h. The reaction was concentrated *in vacuo* and the crude product was used in the next step without further purification.

Step 4: 5-Ethyl-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-5-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)pyrimidine


5




The title compound was synthesized using 2-(4-(4-chloromethyl)-5-(trifluoromethyl)-1*H*-pyrazol-1-yl)piperidin-1-yl)-5-ethylpyrimidine and 2-fluoro-4-(1*H*-tetrazol-1-yl)phenol in manner similar to that described in **Example 1**. **¹H NMR** (CDCl₃): δ 10 8.95 (1H, s), 8.18 (2H, s), 7.64 (1H, s), 7.52 (1H, m), 7.43 (1H, m), 7.19 (1H, m), 5.16 (2H, s), 4.92 (2H, m), 4.50 (1H, m), 3.02 (2H, m), 2.47 (2H, q), 2.22 (2H, m), 2.01 (2H, m), 1.19 (3H, t).

Example 69

15 5-Chloro-2-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidin-1-yl)pyrimidine




20 **Step 1:** *tert-EvXy* 4-hydroxy-2-methylpiperidine-1-carboxylate



To a solution of *tert*-butyl 2-methyl-4-oxopiperidine-1-carboxylate (0.94 g, 4.41 mmol) in methanol (5 mL) was added sodium borohydride (0.22 g, 5.73 mmol) and the reaction was stirred at room temperature for 1h. Water was added slowly, followed by ethyl acetate. The layers were separated and the aqueous phase was extracted with ethyl acetate. The combined organic phases were dried over sodium sulfate, filtered and concentrated *in vacuo* to afford the desired product as a mixture of four stereoisomers, which was used in the next step without further purification.

10

Step 2: *tert*-Butyl 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2-*f*-1,2,3-triazol-2-yl)-2-methylpiperidine-1-carboxylate




15

The title compound was synthesized from *tert*-butyl 4-hydroxy-2-methylpiperidine-1-carboxylate and 1-(4-((2*H*-1,2,3-triazol-4-yl)methoxy)-3-fluorophenyl)-1*H*-tetrazole (**Intermediate 16**) in a manner similar to that described in **Example 48, Step 4**. The title compound was isolated as a mixture of four stereoisomers.

20


Step 3: 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2-*f*-1,2,3-triazol-2-yl)-2-methylpiperidine hydrochloride



The title compound was synthesized from *tert*-butyi 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidine-1-carboxylate in a manner similar to that described in **Intermediate 6**. The title compound was isolated as a mixture of four stereoisomers.

Step 4 : 5-Chloro-2-(4-((2-fiuoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidin- 1-yl)pyrimidine


10



The title compound was synthesized from 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidine hydrochloride and 6-chloro-2-iodopyrimidine in a manner similar to that described in **Example 9**. The title compound was isolated as a mixture of four stereoisomers.

Example 70


2-(4-((2-Fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidin-1-yl)-5-(trifluoromethyl)pyrimidine



The title compound was synthesized from 4-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)-2-methylpiperidine hydrochloride and 2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidine in a manner similar to that described in Example 9. The title compound was isolated as a mixture of four stereoisomers.


Example 71

exo-3-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo [3.2.1]octane




Step 1: 8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octan-3-one

15



The title compound was synthesized from 2-(methylsulfonyl)-5-(trifluoromethyl)pyrimidine and 8-azabicyclo[3.2.1]octan-3-one in a manner similar to that described in Example 9.


Step 2: en</o-8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octan-3-ol



The title compound was synthesized from 8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octan-3-one in a manner similar to that described in **Example 69, Step 1** to afford *endo* and *exo* 8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octan-3-ol. The two compounds were separated by flash column chromatography on silica gel eluting with 0-100% ethyl acetate in hexanes.

Step 3: *exo*-3-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octane


10




The title compound was synthesized from 8-(5-(trifluoromethyl)pyrimidin-2-yl)-8-azabicyclo[3.2.1]octan-3-ol and 5-((2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (**Intermediate 13**) in a manner similar to that described in **Example 48, Step 4**. ¹H NMR (CDCl₃): δ 9.42 (1H, s), 8.51 (2H, s), 8.23 (1H, d), 8.00 (1H, d), 7.65 (1H, s), 7.54 (1H, m), 5.22 (2H, s), 5.15 (1H, m), 5.00 (2H, s), 2.40-1.90 (8H, m).

Example 72

en/o-tert-Butyl 3-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo [3.2.1]octane-8-carboxylate



Step 1: *tert-Butyl 3-oxo-8-azabicyclo[3.2.1]octane-8-carboxylate*




5

The title compound was synthesized from 8-azabicyclo[3.2.1]octan-3-one hydrochloride and *di-tert-butyl* dicarbonate in a manner similar to that described in *Journal of Organic Chemistry* **2005**, *70*, 10872-10874

Step 2: *exo-tert-Butyl 3-hydroxy-8-azabicyclo[3.2.1]octane-8-carboxylate*


10



The title compound was synthesized from tert-butyl 3-oxo-8-azabicyclo[3.2.1]octane-8-carboxylate in a manner similar to that described in **Example 69**,

15 **Step 2.**

Step 3: *endo-tert-Butyl 3-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo[3.2.1]octane-8-carboxylate*




20

The title compound was synthesized from *exo-tert-butyl* 3-hydroxy-8-azabicyclo[3.2.1]octane-8-carboxylate and 5-((2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (**Intermediate 13**) in a manner similar to that described in **Example 48, Step 4.** ^1H NMR (CDCl_3): δ 9.46 (1*H*, s), 8.29 (1*H*, d), 8.02 (1*H*, d), 7.75 (1*H*, s), 7.62 (1*H*, dd), 5.33 (2*H*, s), 4.64 (1*H*, m), 4.26 (2*H*, m), 2.97 (2*H*, m), 2.43 (2*H*, m), 1.77 (2*H*, m), 1.50 (9*H*, s), 1.38 (2*H*, m).

Example 73


exo-tert-Butyl 3-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo [3.2.1]octane-8-carboxylate




The title compound was synthesized from *endo-tert-butyl* 3-hydroxy-8-azabicyclo[3.2.1]octane-8-carboxylate and 5-((2*H*-1,2,3-triazol-4-yl)methoxy)-2-(1*H*-tetrazol-1-yl)pyridine (**Intermediate 13**) in a manner similar to that described in **Example 48, Step 4.** ^1H NMR (CDCl_3): δ 9.48 (1*H*, s), 8.28 (1*H*, d), 8.02 (1*H*, d), 7.71 (1*H*, s), 7.63 (1*H*, m), 5.30 (2*H*, s), 5.02 (1*H*, m), 4.39 (2*H*, m), 2.40-1.85 (8*H*, m), 1.51 (9*H*, s).

Example 74

en</o-3-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-(5-fluoropyrimidin-2-yl)-8-azabicyclo[3.2.1] octane



Step 1: *endo*-3-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo[3.2.1]octane hydrochloride




5

The title compound was synthesized from *endo-tert-butyl* 3-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo[3.2.1]octane-8-carboxylate and 4N hydrochloric acid in dioxane in a manner similar to that described in **Intermediate 6**.

10

Step 2 : *e/?<io*-3-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-(5-fluoropyrimidin-2-yl)-8-azabicyclo[3.2.1]octane

15




The compound was synthesized from *e/?<io*-3-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-azabicyclo[3.2.1]octane hydrochloride and 2-chloro-5-fluoro-pyrimidine in a manner similar to that described in **Example 9**. **1H NMR** (CDCl₃): δ **9.45** (1*H*, s), **8.29** (1*H*, d), **8.25** (2*H*, s), **8.03** (1*H*, d), **7.75** (1*H*, s), 7.61 (1*H*, dd), **5.33** (2*H*, s), **4.59** (2*H*, s), **4.57** (1*H*, m), 3.01 (2*H*, m), **2.49** (2*H*, m), 1.88 (2*H*, m), 1.55 (2*H*, m).

Example 75

exo-3-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-8-(5-fluoropyrimidin-2-yl)-8-azabicyclo[3.2.1]octane

25




The title compound was synthesized in a manner similar to that described in

Example 74. $^1\text{H NMR}$ (CDCl_3): δ **9.45** (1H, s), **8.24** (3H, m), 8.01 (1H, m), **7.65** (1H, s), **7.56** (1H, m), **5.23** (2H, s), 5.14 (1H, m), **4.85** (2H, m), 2.40-1.95 (8H, m).

Example 76

Allyl 4-((4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-
y1)piperidine-1-carboxylate




10

The title compound was synthesized from **Intermediate 8** and allyl chloroformate in a manner similar to that described in *Journal of Medicinal Chemistry* **1998**, *41*, 4983-4994.

$^1\text{H NMR}$ (CDCl_3): δ **8.95** (1H, s), **7.72** (1H, s), **7.53** (1H, m), **7.45** (1H, m), **7.32** (1H, m), **5.95** (1H, m), 5.40-5.20 (4H, m), **4.64** (3H, m), **4.23** (2H, m), **3.09** (2H, m), 2.14 (4H, m).

Example 77

Allyl 4-((4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-
y1)piperidine-1-carboxylate




The title compound was synthesized from **Intermediate 15** and allyl chloroformate in a manner similar to that described in *Journal of Medicinal Chemistry* **1998**, *41*, 4983-4994.

⁵ ^1H NMR (CDCl_3): δ 9.46 (1H, s), 8.28 (1H, m), 8.04 (1H, m), 7.73 (1H, s), 7.61 (1H, m), 5.95 (1H, m), 5.40-5.20 (4H, m), 4.64 (3H, m), 4.24 (2H, m), 3.10 (2H, m), 2.16 (4H, m).

The compounds in **Examples 78-82** were synthesized from **Intermediates 15** with the corresponding 2-fluoropyrimidine, 2-chloropyrimidine, 2-bromopyrimidine, 2-iodopyrimidine, 2-(methylsulfinyl)pyrimidine, 2-(methylsulfonyl)pyrimidine or 2-chlorothiazole in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH_3CN), temperature, base (NEt_3 , $^1\text{Pr}_2\text{NEt}$, K_2CO_3 , NaHCO_3 , Na_2CO_3 , Cs_2CO_3) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 78

2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-pentylpyrimidine




²⁰ ^1H NMR (CDCl_3): δ 8.17 (2H, s), 7.76 (1H, s), 7.66 (1H, s), 7.42 (1H, m), 7.27 (1H, s), 7.07 (1H, m), 5.12 (2H, s), 4.78 (3H, m), 3.16 (2H, m), 2.42 (2H, m), 2.25 (2H, m), 2.16 (2H, m), 1.28 (6H, m), 0.88 (3H, m).

²⁵

Example 79

Methyl 2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-6-methylpyrimidine-4-carboxylate




5

¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.27 (1H, s), 8.02 (1H, m), 7.70 (1H, s), 7.59 (1H, m), 7.05 (1H, s), 5.28 (2H, s), 4.92 (2H, m), 4.75 (1H, m), 3.95 (3H, s), 3.21 (2H, m), 2.43 (3H, s), 2.28 (2H, m), 2.17 (2H, m).

10

Example 80

2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-(4-methoxyphenyl)pyrimidine




15

¹H NMR (CDCl₃): δ 9.44 (1H, s), 8.53 (2H, s), 8.27 (1H, d), 8.03 (1H, d), 7.71 (1H, s), 7.59 (1H, dd), 7.41 (2H, d), 6.98 (2H, d), 5.28 (2H, s), 4.86 (2H, m), 4.76 (1H, m), 3.85 (3H, s), 3.25 (2H, m), 2.29 (2H, m), 2.21 (2H, m).

20

Example 81

Ethyl 2-(4-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)thiazole-5-carboxylate




¹H NMR (CDCl₃): δ 9.44 (IH, s), 8.27 (IH, d), 8.03 (IH, d), 7.87 (IH, s), 7.73 (IH, s), 7.58 (IH, dd), 5.28 (2H, s), 4.76 (IH, m), 4.30 (2H, q), 4.13 (2H, m), 3.39 (2H, m), 3.33 (4H, m), 1.34 (3H, t).

Example 82

2-(4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-(trifluoromethyl)pyridine

10




¹H NMR (CDCl₃): δ 9.44 (IH, s), 8.41 (IH, s), 8.27 (IH, d), 8.03 (IH, d), 7.71 (IH, s), 7.65 (IH, d), 7.58 (IH, dd), 6.72 (IH, d), 5.28 (2H, s), 4.77 (IH, m), 4.47 (2H, m), 3.23 (2H, m), 2.30 (2H, m), 2.20 (2H, m).

Example 83

tert-Butyl 4-((6-(1*H*-tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-3-methylpiperidine-1-carboxylate

20




The title compound was synthesized from **Intermediate 13** and *tert*-butyl 4-hydroxy-3-methylpiperidine-1-carboxylate in a manner similar to that described in **Example 69**. The compounds were isolated as a mixture of four stereoisomers. Characterized by LCMS (ES⁺-MS found: 464 (M+Na), 442 (M+H)).

5 The compounds in **Examples 84-87** were synthesized from **Intermediate 21** with the corresponding substituted 2-fluoropyrimidine, 2-chloropyrimidine, 2-bromopyrimidine, 2-iodopyrimidine, 2-(methylsulfinyl)pyrimidine, 2-(methylsulfonyl)pyrimidine or 2-chlorothiazole in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN),
10 temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 84


15 2-(4-((4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-3-methylpiperidin-1-yl)-5-(trifluoromethyl)pyrimidine



The compounds were isolated as a mixture of four stereoisomers. LCMS:
20 (ES⁺-MS found: 488).

Example 85


2-(4-((4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-3-methylpiperidin-1-yl)-5-ethylpyrimidine



The compounds were isolated as a mixture of four stereoisomers. LCMS:
(ES⁺-MS found: 448).

Example 86

5 2-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-3-methylpiperidin-1-yl)-5-methylpyrimidine




10 The compounds were isolated as a mixture of four stereoisomers. LCMS:
(ES⁺-MS found: 434).

Example 87

2-(4-((6-(1*H*-Tetrazol-1-yl)pyridin-3-yloxy)methyl)-2*H*-1,2,3-triazol-2-yl)-3-methylpiperidin-1-yl)-5-chloropyrimidine

15




The compounds were isolated as a mixture of four stereoisomers. LCMS:
(ES⁺-MS found: 454).

20

The compounds in **Examples 88-98** were synthesized from (2-(1-(5-chloropyrimidin-2-yl)piperidin-4-yl)-2 *H*-1,2,3-triazol-4-yl)methyl methanesulfonate (**Intermediate 24**) and the corresponding phenol, pyrazol-3-ol or thiazol-2-ol in a similar manner to that described in **Example 1**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN); temperature, base (NEt₃, 'Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 88


10 2-(4-(4-((4-(1 *H*-Tetrazol-1-yl)phenoxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-chloropyrimidine



15 ¹H NMR (CDCl₃): δ 8.90 (1H, s), 8.24 (2H, s), 7.68 (1H, s), 7.61 (2H, d), 7.16 (2H, d), 5.21 (2H, s), 4.75 (3H, m), 3.21 (2H, m), 2.26 (2H, m), 2.17 (2H, m).

Example 89

20 5-chloro-2-(4-(4-((2-Methyl-4-(1 *H*-tetrazol-1-yl)phenoxy)methyl)-2 *H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine




25 ¹H NMR (CDCl₃): δ 8.91 (1H, s), 8.21 (2H, s), 7.66 (1H, s), 7.45 (2H, m), 7.07 (1H, m), 5.21 (2H, s), 4.72 (3H, m), 3.19 (2H, m), 2.26 (5H, m), 2.15 (2H, m).

Example 90

2-((2-(1-(5-Chloropyrimidin-2-yl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-5-methyl-4-(pyridin-3-yl)thiazole

5




¹H NMR (CDCl₃): δ 8.58 (2H, m), 8.17 (2H, s), 7.63 (2H, m), 7.60 (1H, s), 5.42 (2H, s), 4.66 (3H, m), 3.14 (2H, m), 2.26 (3H, s), 2.18 (2H, m), 2.11 (2H, m).

10

Example 91

2-((2-(1-(5-Chloropyrimidin-2-yl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methoxy)-5-methyl-4-(pyridin-4-yl)thiazole

15




The title compound was isolated as a TFA salt. ¹H NMR (CDCl₃): δ 8.78 (2H, m), 8.24 (2H, s), 8.09 (2H, m), 7.66 (1H, s), 5.52 (2H, s), 4.75 (1H, br.s), 4.71 (3H, m), 3.21 (2H, m), 2.38 (3H, s), 2.24 (2H, m), 2.15 (2H, m).

20

Example 92

2-(4-((4-(1*H*-1,2,3-Triazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-chloropyrimidine




¹H NMR (CDCl₃): δ 8.23 (2H, s), 7.91 (1H, s), 7.82 (1H, s), 7.68 (1H, s), 7.64 (2H, d), 7.11 (2H, d), 5.19 (2H, s), 4.72 (3H, m), 3.20 (2H, m), 2.25 (2H, m), 2.14 (2H, m).

5

Example 93

2-(4-((4-(1*H*-1,2,4-Triazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-chloropyrimidine




¹H NMR (CDCl₃): δ 8.45 (1H, s), 8.22 (2H, s), 8.07 (1H, s), 7.67 (1H, s), 7.56 (2H, d), 7.08 (2H, d), 5.17 (2H, s), 4.72 (3H, m), 3.19 (2H, m), 2.24 (2H, m), 2.13 (2H, m).

15

Example 94

2-(4-((4-(1*H*-Imidazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)-5-chloropyrimidine




20

¹H NMR (CDCl₃): δ 8.73 (1H, s), 8.24 (2H, s), 7.68 (1H, s), 7.48 (1H, s), 7.40 (1H, s), 7.36 (2H, d), 7.15 (2H, d), 5.20 (2H, s), 4.74 (3H, m), 3.21 (2H, m), 2.25 (2H, m), 2.17 (2H, m).

Example 95

5-Chloro-2-(4-(4-((Methylsulfonyl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine

5




¹H NMR (CDCl₃): δ 8.24 (2H, s), 7.87 (2H, d), 7.67 (1H, s), 7.11 (2H, d), 5.21 (2H, s), 4.71 (3H, m), 3.20 (2H, m), 3.03 (3H, s), 2.25 (2H, m), 2.14 (2H, m).

10

Example 96

5-Chloro-2-{4-[4-(2-[1,3,4]oxadiazol-2-yl-phenoxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine

15




¹H NMR (DMSO-J₆): δ 9.31 (1H, s), 8.43 (2H, s), 7.87-7.85 (2H), 7.60 (1H, t, J=7.4 Hz), 7.42 (1H, d, J=8.4 Hz), 7.15 (1H, t, J=7.6 Hz), 5.32 (2H, s), 4.84-4.82 (1H, m), 4.57-4.53 (2H, m), 3.26-3.21 (2H, m), 2.18-2.14 (2H, m), 1.93-1.85 (2H, m).

20

Example 97

5-Chloro-2-{4-[4-(1-methyl-5-trifluoromethyl-1H-pyrazol-3-yloxy)methyl]-[1,2,3]triazol-2-yl}-piperidin-1-yl}-pyrimidine

25




¹H NMR (DMSO- J_6): δ 8.43 (2H, s), 7.85 (1H, s), 6.41 (1H, s), 5.19 (2H, s), 4.88-4.83 (1H, m), 4.57-4.54 (2H, m), 3.26-3.20 (2H, m), 2.17-2.15 (2H, m), 1.93-1.85 (2H, m).

5

Example 98

5-Chloro-2-{4-[4-(6-chloro-pyridin-3-yloxymethyl)-[1,2,3]triazol-2-yl]-piperidin-1-yl}-pyrimidine




¹H NMR (DMSO- J_6): δ 8.42 (2H, s), 8.19 (1H, d, $J=2.8$ Hz), 7.89 (1H, s), 7.57 (1H, dd, $J=2.8, 8.8$ Hz), 7.43 (1H, d, $J=8.8$ Hz), 5.23 (2H, s), 4.89-4.84 (1H, m), 4.58-4.54 (2H, m), 3.26-3.20 (2H, m), 2.17-2.15 (2H, m), 1.94-1.85 (2H, m).

15

Example 99

1-(4-{2-[1-(5-Trifluoromethyl-pyrimidin-2-yl)-piperidin-4-yl]-2-H-[1,2,3]triazol-4-ylmethoxy}-phenyl)-pyrrolidin-2-one




20

The title compound was synthesized from 1-(4-hydroxyphenyl)pyrrolidin-2-one and **Intermediate 26** in a manner similar to that described in **Example 1**. ¹H NMR (DMSO- J_6): δ 8.71 (2H, s), 7.86 (1H, s), 7.53 (2H, d, $J=8.8$ Hz), 7.01 (2H, d, $J=8.8$ Hz), 5.11 (2H, s),

4.93-4.86 (1H, **m**), 4.70-4.67 (2H, **m**), **3.76** (2H, t, *J*=6.8 Hz), 3.36-3.31(2H, **m**), **2.43** (2H, t, *J*=8.0 Hz), 2.23-2.20 (2H, **m**), 2.06-1.87 (4H, **m**).

Example 100

5 Synthesis **of** 2-[4-(4-benzyloxymethyl-[1,2,3]triazol-2-yl)-piperidin-1-yl]-5-trifluoromethyl-pyrimidine




10 The title compound was synthesized from (2-(1-(5-(trifluoromethyl)pyrimidin-2-yl)piperidin-4-yl)-2*H*-1,2,3-triazol-4-yl)methanol (**Intermediate 27**) and benzyl alcohol in a manner similar to that described in **Example 29**. ¹H NMR (CDCl₃): δ 8.50 (2H, s), 7.60 (1H, s), 7.36-7.30(5H, m), 4.86-4.82 (2H, m), 4.79-4.72 (1H, m), 4.62 (2H, s), 4.59 (2H, s), 3.33-3.26(2H, m), 2.30-2.16 (4H, m).

15 The compounds in **Examples 101-104** were synthesized from one of **Intermediates 7** or **8** with the corresponding chloroformate in a manner similar to that described in **Example 35**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 101

Isobutyl 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidine-1-carboxylate




¹H NMR (CDCl₃): δ 8.92 (1H, s), 7.72 (1H, s), 7.51 (1H, m), 7.42 (1H, m), 7.30 (1H, m), 5.30 (2H, s), 4.64 (1H, m), 4.21 (2H, m), 3.89 (2H, d, *J* = 6.80 Hz), 3.07 (2H, m), 5 2.10-2.18 (4H, m), 1.95 (1H, m), 0.95 (6H, t, d, *J* = 6.80 Hz). LCMS: (ES⁺-MS found: 445.3).

Example 102

Benzyl 4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidine-1-carboxylate

10




¹H NMR (CDCl₃): δ 8.92 (1H, s), 7.72 (1H, s), 7.51 (1H, m), 7.42 (1H, m), 7.29-7.38 (6H, m), 5.30 (2H, s), 5.15 (2H, s), 4.64 (1H, m), 4.23 (2H, m), 3.10 (2H, m), 15 2.10-2.18 (4H, m). LCMS: (ES⁺-MS found: 479.1).

Example 103

Isobutyl 4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidine-1-carboxylate


20



¹H NMR (CDCl₃): δ 7.65-7.70 (3H, m), 7.24-7.28 (1H, m), 5.30 (2H, s), 4.63 (1H, m), 4.22 (2H, m), 3.89 (2H, d, *J* = 6.80 Hz), 3.07 (2H, m), 3.05 (3H, s), 2.10-2.21 (4H, m), 1.95 (1H, m), 0.95 (6H, t, d, *J* = 6.80 Hz). LCMS: (ES⁺-MS found: 455.3).

Example 104

5 Benzyl 4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-2 *H*-1,2,3-triazol-2-
y1)piperidine-1-carboxylate




10 ¹H NMR (CDCl₃): δ 7.65-7.70 (3H, m), 7.33-7.37 (4H, m), 7.24-7.28 (2H, m),
5.30 (2H, s), 5.15 (2H, s), 4.63 (1H, m), 4.23 (2H, m), 3.09 (2H, m), 3.04 (3H, s), 2.10-2.18
(4H, m). LCMS: (ES⁺-MS found: 489.0).

The compounds in **Examples 105-107** were synthesized from one of **Intermediates 7 or 8** with the corresponding substituted 2-fluoropyrimidine, 2-chloropyrimidine, 2-bromopyrimidine, 2-iodopyrimidine, 2-(methylsulfinyl)pyrimidine or 15 2-(methylsulfonyl)pyrimidine in a manner similar to that described in **Example 9**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of 20 organic synthesis.

Example 105

Ethyl 2-(4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-2 *H*-1,2,3-triazol-2-
y1)piperidin-1-yl)pyrimidine-5-carboxylate




¹H NMR (CDCl₃): δ 8.86 (1H, s), 7.65-7.70 (3H, m), 7.24-7.28 (2H, m), 5.30 (2H, s), 4.90 (2H, m), 4.78 (1H, m), 4.35 (2H, q), 3.31 (2H, m), 2.32 (2H, m), 2.19 (2H, m), 1.38 (3H, t). LCMS: (ES⁺-MS found: 504.9).

Example 106

Ethyl 2-(4-(4-((2-fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine-5-carboxylate

10




¹H NMR (CDCl₃): δ 8.91 (1H, s), 8.86 (2H, s), 7.72 (1H, s), 7.51 (1H, m), 7.41 (1H, m), 7.29 (1H, m), 5.30 (2H, s), 4.90 (2H, m), 4.79 (1H, m), 4.34 (2H, q), 3.32 (2H, m), 2.32 (2H, m), 2.19 (2H, m), 1.38 (3H, t). LCMS: (ES⁺-MS found: 495.0).

Example 107

5-Fluoro-2-(4-(4-((2-fluoro-4-(methylsulfonyl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)pyrimidine

20




¹H NMR (CDCl₃): δ 8.22 (2H, s), 7.65-7.70 (3H, m), 7.24-7.28 (1H, m), 5.30 (2H, s), 4.72 (3H, m), 3.19 (2H, m), 3.04 (3H, s), 2.25 (2H, m), 2.16 (2H, m). LCMS: (ES⁺-MS found: 451.2).

The compounds in **Examples 108-112** were synthesized from **Intermediates 8** and the corresponding carboxylic acid in a manner similar to that described in **Example 50**. One skilled in the art of organic synthesis will appreciate that conditions such as solvent (DMF, CH₃CN), temperature, base (NEt₃, ¹Pr₂NEt, K₂CO₃, NaHCO₃, Na₂CO₃, Cs₂CO₃) and concentration can be selected through routine experimentation to optimize yields. Additionally, alternative coupling methods can be used that are well known in the art of organic synthesis.

Example 108

(4-(4-((2-Fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)(pyridin-2-yl)methanone




15

¹H NMR (DMSO-d₆): δ 10.00 (1H, s), 8.58 (1H, d), 7.94 (1H, s), 7.92 (1H, d), 7.89 (1H, m), 7.72 (1H, d), 7.61 (1H, m), 7.58 (1H, d), 7.46 (1H, m), 5.34 (2H, s), 4.87 (1H, m), 4.48 (1H, m), 3.73 (1H, m), 3.29 (1H, m), 3.12 (1H, m), 2.22 (1H, m), 2.06 (1H, m), 1.97 (2H, m). LCMS: (ES⁺-MS found: 450.2).

Example 109

(4-(4-((2-Fluoro-4-(1*H*-tetrazol-1-yl)phenoxy)methyl)-2*H*-1,2,3-triazol-2-yl)piperidin-1-yl)(6-hydroxypyridin-2-yl)methanone

25




¹H NMR (*DMSO-Cd*): δ 10.00 (IH, s), 7.94 (IH, s), 7.90 (IH, m), 7.73 (IH, d), 7.59 (IH, m), 7.52 (IH, d), 6.45 (2H, m), 5.34 (2H, s), 4.86 (2H, m), 4.37 (1H, m), 3.62 (IH, m), 5 3.06 (IH, m), 1.99-2.19 (4H, m). LCMS: (ES⁺-MS found: 466.1).

Example 110

(4-((2-Fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)(3-hydroxypyridin-2-yl)methanone

10




¹H NMR (*OMSO-d* β): δ 10.39 (IH, br), 9.99 (IH, s), 8.03 (IH, d), 7.93 (IH, s), 7.91 (IH, m), 7.71 (IH, m), 7.59 (IH, d), 7.29 (2H, m), 5.34 (2H, s), 4.86 (IH, m), 4.49 (IH, m), 15 3.22 (IH, m), 3.07 (IH, m), 2.19 (IH, m), 1.93-2.04 (4H, m). LCMS: (ES⁺-MS found: 466.2).

Example 111

(4-((2-Fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)(3-methylpyridin-2-yl)methanone

20




¹H NMR (*DMSO-d₆*): δ 10.00 (IH, s), 8.38 (IH, d), 7.94 (IH, s), 7.90 (IH, m), 7.72 (2H, m), 7.59 (IH, d), 7.34 (IH, m), 5.34 (2H, s), 4.87 (IH, m), 4.49 (IH, m), 3.22 (IH, m), 5 3.07 (IH, m), 2.23 (3H, s), 2.04 (IH, m), 1.93 (4H, m). LCMS: (ES⁺-MS found: 464.4).

Example 112

(4-(4-((2-Fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-2H-1,2,3-triazol-2-yl)piperidin-1-yl)(pyridin-3-yl)methanone

10




¹H NMR (*OMSO-d₃*): δ 10.00 (IH, s), 8.66 (2H, d), 7.94 (IH, s), 7.90 (2H, m), 7.72 (IH, m), 7.59 (IH, d), 7.34 (IH, m), 5.34 (2H, s), 4.86 (IH, m), 4.45 (IH, m), 3.59 (IH, m), 15 3.07 (IH, m), 1.99-2.35 (5H, m). LCMS: (ES⁺-MS found: 450.0).

Example 113

5-Ethyl-2-(4-((2-fluoro-4-(1H-tetrazol-1-yl)phenoxy)methyl)-1H-pyrazol-1-yl)piperidin-1-yl)pyrimidine

20



The title compound was synthesized using **Intermediate 31** and 2-chloro-5-ethylpyrimidine in a similar manner as that described in **Example 9**.

¹H NMR (CDCl₃): δ 8.19 (2H, s), 7.41 (1H, m), 7.05 (1H, m), 6.77 (1H, m), 6.67 (1H, m), 6.35 (1H, s), 6.26 (1H, m), 5.09 (2H, m), 4.86 (2H, m), 4.37 (1H, m), 3.02 (2H, m), 2.47 (2H, m), 2.20 (2H, m), 1.95 (2H, m), 1.20 (3H, t).

Biological Example 1

Stimulation of cAMP

10 Generation of IC-GPCR2 stable cell line

The compounds of the present invention were evaluated in an assay demonstrating agonism of IC-GPCR2. This assay was developed using a stable cell line expressing IC-GPCR-2, generated as follows. IC-GPCR2 (Seq. ID No. 1) was cloned into Gateway pDEST 40vector (Invitrogen), using the Gateway cloning system (invitrogen) according to the manufacturers instructions. A stable cell line was generated by transfecting a 10cm plate of CHO cells (source) with 8ug of this construct using Transit-CHO transfection kit (Minis). CHO cells were plated the day prior to transfection at a density of 3,000,000 cells/plate. Clones were selected using the antibiotic G418 at 500ug/ml. 23 clones were picked and assayed for the expression of the receptor by measuring changes in intracellular cAMP levels in response to an IC-GPCR2 agonist (Arena 51).

To measure cAMP activity in response to IC-GPCR2 agonist, the clones were plated in 96 well plates at 17500 cells per well. On the day after plating, cells were incubated with the IC-CPCR2 agonist at 10uM for 30 minutes in Ham's F12 Media (Gibco) with 0.04% DMSO. cAMP was measured using the cAMP dynamic kit from Cis Bio (Bedford, MA) according to the manufacturers instructions. Briefly, cells were lysed, and cAMP levels determined by competitive immunoassay using D2 labeled cAMP, and europium cryptate tagged anti cAMP antibody. When in close proximity, the D2 and europium cryptate undergo fluorescence resonance energy transfer (FRET), which is measured as a fluorescence ratio (665 nm/620 nm). Unlabelled cAMP in the cell lysate

competed with the D2 labeled cAMP for the europium cryptate labeled antibody. The resulting decrease in FRET signal corresponded to intracellular cAMP levels. Fluorescence was read on a BMG Labtech PHERAstar, software version 1.50.

5 The clone with the greatest response to IC-GPCR2 agonist was selected for the screening assay.

Determination of activity of compounds

Compounds were dissolved in 100% DMSO to a concentration of 10⁻⁶M to provide stock solutions. To determine activity against IC-GPCR2, compounds were 10 incubated with IC-GPCR2 stably expressing cells (described above), at 6-8 concentrations ranging from 0.00003 to 10 micromolar, in 96 well plates, in 50^l of Hams F12 media for 30 minutes. Cells were plated at 17500 cells per well 1 day before running the assay. All compounds were also screened against the parental CHO cells. cAMP was measured using the cAMP dynamic kit from Cis Bio (Bedford, MA), according to the manufacturer's 15 instructions. Briefly, cells were lysed and cAMP levels determined by competitive immunoassay using D2 labeled cAMP, and europium cryptate tagged anti cAMP antibody. When in close proximity, the D2 and europium cryptate undergo fluorescence resonance energy transfer (FRET), which is measured as a fluorescence ratio (665 nm/620 nm). Unlabelled cAMP in the cell lysate competed with the D2 labeled cAMP for the europium 20 cryptate labeled antibody. The resulting decrease in FRET signal corresponded to intracellular cAMP levels

To determine percent activity for a tested compound, the FRET signal value obtained at a particular concentration are compared to the Maximal FRET signal value obtained for 5-Ethyl-2- {4-[4-(4-tetrazol- 1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1 -yl} -pyrimidine. The maximal activity of 5-Ethyl-2-{4-[4-(4-tetrazol- 1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine is designated as 100% activity. Typically, the concentration of 5-Ethyl-2- {4-[4-(4-tetrazol- 1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin- 1-yl} -pyrimidine in the assay was approximately 0.1 μ M. The synthesis of 5-Ethyl-2-{4-[4- 25 (4-tetrazol-1-yl-phenoxyethyl)-thiazol-2-yl]-piperidin-1-yl}-pyrimidine is disclosed in co- 30 owned pending U.S. patent application serial no. 11/964,461, herein incorporated by

reference. Activities of compounds that were tested are shown in Table 1 below and are expressed as % activity at 3 μ M compound compared to the maximal activity of 5-Ethyl-2-{4-[4-(4-tetrazol-1-yl-phenoxy)methyl]-thiazol-2-yl}-piperidin-1-yl}-pyrimidine at 3 μ M. Examples 39, 42, 43, 53, 79, and 97 were tested did not exhibit activity when tested at 3 μ M
5 but are contemplated to have activity at higher concentrations and therefore are not included in Table 1.

Table 1

Example No.	% activity at 3 μ M
1	64.46
3	79.02
4	79.18
5	75.78
6	79.51
7	78.16
8	77.16
9	65.75
10	71.04
13	61.30
14	74.80
15	75.63
16	70.31
17	68.23
18	77.67
19	80.10
21	63.66
22	78.44
29	48.76
30	53.41
31	59.95
32	75.55
33	88.32
34	87.61
35	72.14
36	70.06
37	62.88
38	85.32
40	7.10
41	42.63
44	70.92
45	67.58

Example No.	% activity at 3 µM
46	79.78
47	74.90
48	83.24
49	78.48
50	49.51
51	57.93
52	52.86
54	36.82
55	31.89
56	7.31
57	27.80
58	79.55
59	33.25
60	27.86
61	43.59
62	32.06
64	66.18
65	70.03
66	80.00
67	57.87
68	74.11
69	55.22
70	66.77
71	70.78
72	60.55
73	84.07
74	61.88
75	47.94
76	56.39
77	69.86
78	57.13
80	77.53
81	32.98
82	64.90
83	63.14
84	78.05
85	76.12
86	71.39
87	75.63
88	60.96
89	48.60
90	13.60
91	5.82
92	58.28
93	50.84
94	49.73
95	51.03

Example No.	% activity at 3 µM
96	28.90
98	17.16
99	56.81
100	10.69
101	50.67
102	38.98
103	51.30
104	28.40
105	10.65
106	11.46
107	36.83
108	41.78
109	7.45
110	1.30
111	24.12
112	42.30
113	57.71

Biological Example 2

Insulin Secretion (Islet Perfusion)

To determine the effect of IC-GPCR2 agonists on insulin secretion from islets, 5 islets from Sprague Dawley rats are isolated. 200-250g Sprague Dawley rats (Charles River laboratories) are maintained on regular chow (Purina 5001). Before the procedure rats are anesthetized with intra peritoneal injection of pentobarbital at 200mg/kg . The bile duct is clamped where it enters the duodenum, then a catheter is placed in the bile duct between the liver and the pancreas. The pancreas is infused through the catheter with a solution of 10 0.75mg/ml collagenase P (Roche) in HBSS buffer (Biowhitaker) supplemented with 0.1% glucose and 0.02% BSA. The pancreas is then excised from the rat and placed in 5ml of the collagenase P solution in a 37°C waterbath for 8 minutes. After 8 minutes the digested pancreas is shaken vigorously by hand for 30 seconds. The resulting digest is washed four times in the HBSS buffer, then applied to a discontinuous ficoll gradient. To make the 15 gradient, the digest is resuspended in 7.5ml of ficoll DL400 solution (Sigma) density 1.108, in a 15ml tube. Three 2ml layers of ficoll solution of decreasing density (1.096, 1.069, 1.037) are then added to the tube to create a density gradient. The gradient is centrifuged at 1500rpm for 15 minutes after which islets are picked from the top two layers. Islets are

washed four times in HBSS buffer, then cultured in RPMI 1640 media (Gibco) supplemented with 1% fetal bovine serum. The following day, 25 size-matched islets are placed in a perfusion chamber and exposed to Krebs Ringer Buffer (KRB; 119mM NaCl, 4.7mM KCl, 25mM NaHCO₃, 2.5mM CaCl₂, 1.2 mM MgSO₄, 1.2mM KH₂PO₄) at a rate of 5 1ml/minute, using a Cellex Acu-sys S perfusion culture system. The islets are exposed to KRB containing glucose at 2mM for 30 minutes, followed with buffer containing 16mM glucose for 30 minutes, then returned to 2mM glucose for a further 30 minutes, in the presence of 0.1-100μM of the IC-GPCR2 agonist or vehicle (DMSO). Perifusate is collected at 1 minute intervals using a fraction collector, and assayed for insulin using an 10 ELISA kit (Mercodia Ultrasensitive Rat Insulin ELISA Kit, ALPCO). Insulin secretion rate in response to glucose is plotted against time, and the AUC of the curve determined in order to quantify the insulin secretory response to 16mM glucose during the 30 minute perfusion. Statistical significance of differences in AUC between treated and untreated islets are determined by paired Students t test.

15

Biological Example 3

Oral Glucose Tolerance

8-10 week old male C57/6J mice (Harlan) were maintained on regular chow diet (Purina 5001). The day of the experiment mice were fasted for 6 hours, then randomized 20 into groups (n=8) to receive the tested IC-GPCR2 agonist at doses ranging from 0.3-30mg/kg or the vehicle (1% CMC, 2% TWEEN 80). Compounds were delivered orally via gavage at 10ml/kg. Blood glucose levels were measured by glucometer (Ascensia Elite XL, Bayer) at time 0, before administration of compound. Blood glucose was measured again after 30 minutes, and then the mice were dosed orally with 2g/kg glucose at 10ml/kg. 25 Blood glucose measurements were taken 15, 30, 60, 90 and 120 minutes after glucose administration, by glucometer (Ascensia Elite XL, Bayer).

Glucose levels were plotted against time, and the incremental area under the curve (AUC) of the glucose excursion was determined from time 0 using Graphpad Prism 5.0. Outliers were excluded using Tukey's box plot outlier test, and statistical significance of

differences in AUC of compound treatment compared to vehicle was determined by non-parametric Kruskal-Wallis test with Dunn's post test.

Table 2 below shows the mean percentage inhibition of the glucose excursion for the eight animals tested in each group. The compounds were tested at 30 mg/kg and the levels of blood glucose were determined in the presence and absence of the tested compounds. The percent of glucose reduction is reported. The tested compounds were selected as examples from the exemplified compounds. These results demonstrate that the IC-GPCR2 agonists can lower blood glucose in response to an oral glucose challenge.

Table 2

Compound	% reduction AUC glucose excursion at 30mg/kg
Compound 1	40.8
Compound 1	58.8

10

Biological Example 4

Tissue Specific Expression

RNA was extracted from isolated rat and mouse islets, and used to prepare double stranded cDNA using standard techniques (*see* Sambrook et al., Molecular Cloning, A Laboratory Manual (3rd ed. 2001); Current Protocols in Molecular Biology (Ausubel et al., eds., 1994)). The cDNA was cloned into the pZL1 vector (Invitrogen) and the 3' ends of individual clones were sequenced in multiple rounds of sequencing reactions. Sequence data representing approximately 12,000 independent clones were used to construct oligonucleotide probes synthesized on a GENECHIP® (Affymetrix Inc., Santa Clara, CA), producing mouse and rat islet chips. RNA from five rat islet preparations (each preparation from a different mouse), and preparations from a panel of rat tissues, were hybridized to the rat chips. Expression data was analyzed with the Affymetrix MAS 4.0 algorithm to give relative gene expression values as average difference scores, and presence/absence calls. RNA from two preparations from a mouse beta cell line BHC-9, four mouse islet

preparations (each preparation from a different mouse), and preparations from a panel of mouse tissues, were hybridized to the mouse chips. Expression data was analyzed with the Affymetrix MAS 5.0 algorithm to give relative gene expression values as signal, and presence/absence calls.

5 Figures 1 (rat) and 2 (mouse) show the tissue specific expression of the receptor for the novel agonists of the present invention, showing tissue specificity to pancreatic islet cells (including the beta cells therein).

Sequence ID No. 1

atggaatcatttctcattggagtgatcctgcgtcctggcccccattgtc
actaacacactagtggtggctgtgtgttgcataacaagaatgtggcgtc
5 ctctgcttcacccatgttgcacccatgtggctgacacccattgttgc
ctactcacagaccaggctccagcccttcggccacacagaagaccctgtgc
cgatggcattgtcacttcctccgcagctgcctgtcctcacggcatgtgatc
tttgacaggtacccatcaagcagccctccgcacttgcataaggatcatgatc
gtggccggggcctgcattgccggctgtggtagtgtcttacccattggcttc
10 ctgcgaatccccatgtccagcagactgcctacaaagggcagtgcagcttgc
ttcacccctacttcgtgtgcaccctcctgcgttgcattccagccatgc
tttgtcttctactgcgacatgctcaagattgcctccatgcacagccagattc
aagatgaaacatgcaggagccatggctggaggattcgtatccccacggactcc
ttcaaagctctccgtactgtgtcttcattggagctttgcattctgc
15 ttccattatcactggcattgtgcagggtggccaggagtgtcaccttacccatgt
gaacggtagctgtggctgcggcggtggcaactccctgcataccactcatatgc
tattggcagaaggagggtgcgactgcagctaccatggccctaggagtgaagaagg
ctcacccattccctttcttgcaggattgtggccagagaggcccaggaa
agttccctgtcacatcgtaactatccagctcagattgtatggctaa

20

Sequence ID No. 2

MESSFSFGVILAVLASLIIATNTLVAVAVLLLIHKNDGVSLCFTLNLAADTLIGVAISG
LLTDQLSSPSRPTQKTLCSLRMAFVTSSAAASVLTVMLITFDRYLAIKQPFRYLKIMSGF
VAGACIAGLWLVSYLIQLPLGIPMFQQTAYKGQCSFFAVFHPHFVLTSCVGFFPAMLL
25 FVFFYCDMLKIASMHSQQIRKMEHAGAMAGGYRSPRTPSDFKALRTVSVLIGSFALSWTP
FLITGIVQVACQECHLYVLERYLWLLGVGNSSLNPLIYAYWQKEVRLQYHMALGVKKV
LTSFLLFLLARNCGPERPRESSCHIVTISSSEFDG

Biological Example 5**Incretin measurement**

5 The effect of IC-GPCR2 agonists on the secretion of insulin, Glucagon-like peptide-1 (GLP-I) and GIP in C57/6J mice are determined as follows.

8-10 week old male C57/6J mice (Harlan) are maintained on a regular chow diet (Purina 5001). On the day of the experiment mice are fasted for 6 hours then randomized into groups (n=8). All groups are treated with the DPPIV inhibitor sitagliptin at 100mg/kg 10 to prevent degradation of active GLP-I . IC-GPCR-2 agonist compounds are dosed at concentrations ranging from 0.3-300mg/kg in 1% CMC, 2% TWEEN 80 at -30 minutes. Sitagliptin is administered in the same dosing solution. Oral glucose at 2g/kg is administered at 0 minutes. At 10 minutes after glucose administration, animals are 15 anesthetized with pentobarbital (40mg/ml in 10% ethanol) and blood collected by heart puncture in microtainer tubes (BD) with potassium EDTA. For GLP-I assay, the collection tubes also contain a DPP-IV inhibitor provided in the GLP-I assay kit.

Insulin is measured using the Mercodia mouse Insulin ELISA Kit (ALPCO) according to the manufacturer's instructions. Bioactive GLP-I is measured using Glucagon-like peptide- 1 (active) ELISA assay kit (Linco) according to the manufacturers instructions. 20 GIP is measured using rat/mouse GIP total ELISA assay kit (Linco), according to the manufacturer's instructions.

Biological Example 6**Improvement of diabetes parameters in female ZDF rat**

25 Female ZDF rats (Charles River laboratories) are obtained at 6 weeks of age and acclimatized for 1 week before being placed on a high fat diet (RD 13004, Research Diets). Compounds are administered to the rats by daily gavage at concentrations ranging from 0.3-300 mg/kg in 1% CMC, 2% TWEEN 80. Body weight and food intake is monitored daily. After 14 days of dosing, blood samples are taken from overnight fasted animals to measure 30 glucose and insulin. Glucose is measured using a glucometer (Ascensis Elite XL, Bayer),


insulin is measured using rat insulin ELISA kit (ALPCO). Insulin and glucose levels are compared to those of vehicle treated animals to determine efficacy.

All patents, patent applications, publications and presentations referred to herein are incorporated by reference in their entirety. Any conflict between any reference cited
5 herein and the teaching of this specification is to be resolved in favor of the latter.

Similarly, any conflict between an art-recognized definition of a word or phrase and a definition of the word or phrase as provided in this specification is to be resolved in favor of the latter.

WHAT IS CLAIMED IS:

- ## 1. A compound of Formula (I):



(I)

wherein,

X, Y and Z are each independently selected from the group consisting of N and C(R^3):

the subscript r is 0, 1, 2 or 3;

the subscript s is 0, 1, 2 or 3, and the sum of r + s is ≤ 4 ;

the subscript q is 0, 1, 2, 3 or 4;

L is $-(\text{CH}_2)_n-$ wherein the subscript n is 2, 3 or 4, and optionally at least one CH_2 is replaced by O, C(O), NR⁵, S, S(O) or S(O)₂, and any remaining CH_2 is optionally substituted with one or two substituents selected from the group consisting of halo, C₁₋₄alkyl, C₃₋₆cycloalkyl and C¹haloalkyl;

Ar is an aryl or heteroaryl group, optionally substituted with from one, two, three, four or five R⁶ substituents;

R^1 is selected from the group consisting of H, C_{1-10} alkyl, C^{11} substituted alkyl, C_{3-7} cycloalkyl, C_{3-7} substituted cycloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, $-X^1-C(O)R^a$, $-X^1-C(O)OR^a$, $-X^1-C(O)NR^aR^b$, $-S(O)_2R^a$, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said cycloalkyl group, heterocyclyl group, aryl group and heteroaryl group is optionally substituted with from one, two, three or four substituents independently selected from the group consisting of halo, C_{1-10} alkyl, C_{1-10} iosubstituted alkyl, C_{3-7} cycloalkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, aryl, heteroaryl, $-CN$, $-NR^aC(O)R^b$, $-NR^aC(O)NR^aR^b$, $-NO_2$, $-OR^a$, $-NR^aR^b$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)NR^aR^b$, $-SR^a$, $-S(O)R^a$, $-S(O)_2R^a$, $-NR^aS(O)_2R^b$, and $-S(O)_2NR^aR^b$, or optionally R^a and R^b are combined to form a 4-, 5- or 6-membered ring, and X^1 is selected from the group consisting of a bond, C_{1-4} alkylene, C_{2-6} alkenylene, C_{2-6}

ϵ alkynylene, -C(O)-, and -C(O)(CH₂)₁₋₄-, wherein the aliphatic portions of X¹ are optionally substituted with one, two or three groups selected from the group consisting of halo, C₁₋₄alkyl, and C[^]substituted alkyl;

each R² is independently selected from the group consisting of halo, C₁₋₅alkyl, C[^]substituted alkyl, C₃₋₇cycloalkyl, C₃₋₇substituted cycloalkyl, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -OR^a, -NR^aR^b, -NR^aC(0)R^b, -S(O)R^a, -S(O)₂R^a and -S(O)₂NR^aR^b, and wherein when the subscript q is 2, 3, or 4, two R² groups can optionally cyclize to form a ring;

each R³ is independently selected from the group consisting of H, halo, C₁₋₄alkyl, C₁₋₄haloalkyl, C₃₋₇cycloalkyl, aryl and -OR^a;

R⁴ is selected from the group consisting of H, halo, C₁₋₆alkyl, and Ci₆substituted alkyl;


R⁵ is selected from the group consisting of H, C₁₋₄alkyl, C[^]haloalkyl, C₃₋₆cycloalkyl, -C(O)R^a and -S(O)₂R^a;

each R⁶ is independently selected from the group consisting of H, halo, Ci-alkyl, C₁₋₁₀substituted alkyl, C₃₋₇cycloalkyl, C₃₋₇substituted cycloalkyl, C₂-ioalkenyl, C₂-ioalkynyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -NR^aC(0)R^b, -NR^aC(0)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, S(O)₂R^a, -NR^aSR^b, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said heterocyclyl groups, said aryl and heteroaryl groups are optionally substituted with from one to four substituents independently selected from halo, oxo, Ci₄alkyl, C₁₋₄haloalkyl, C₃₋₇cycloalkyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(0)R^a, -C(0)OR^a, -C(O)NR^aR^b, -NR^aC(0)R^b, -NR^aC(0)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)R^b and -S(O)₂NR^aR^b or optionally R^a and R^b are combined to form a A-, 5- or 6-membered ring;


and each R^a and R^b is independently selected from the group consisting of H, C₁₋₁₀alkyl, C₁₋₁haloalkyl, C₃₋₁iocycloalkyl, heterocyclyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, aryl, 5- to 6-membered heteroaryl and arylC₁₋₄alkyl; and wherein the aliphatic portions of each of said R^a and R^b is optionally substituted with from one to three groups selected from the group consisting of halo, -ORⁿ,

-OC(O)Rⁿ, -OC(O)N(Rⁿ)₂, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -S(O)₂N(Rⁿ)₂, -NRⁿS(O)₂Rⁿ, -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NRⁿC(O)Rⁿ, -NRⁿC(O)N(Rⁿ)₂, -C(O)ORⁿ, -NRⁿC(O)ORⁿ, -CN, -NO₂, -N(Rⁿ)₂ and -NRⁿS(O)₂N(Rⁿ)₂, wherein each Rⁿ is independently H, C₁₋₃haloalkyl or an unsubstituted C₁₋₆ alkyl; and wherein the aryl and heteroaryl portions are optionally substituted with from one to three groups selected from halo, -OR^m, -OC(O)N(R^m)₂, -SR^m, -S(O)R^m, -S(O)₂R^m, -S(O)₂N(R^m)₂, -NR^mS(O)₂R^m, -C(O)N(R^m)₂, -C(O)R^m, -NR^mC(O)R^m, -NR^mC(O)N(R^m)₂, -C(O)OR^m, -NR^mC(O)OR^m, -CN, -NO₂, -N(R^m)₂ and -NR^mS(O)₂N(R^m)₂, wherein each R^m is independently H, C¹⁻³haloalkyl or an unsubstituted C₁₋₆ alkyl; and pharmaceutically acceptable salts, solvates, stereoisomers, and esters thereof.

2. A compound of claim 1, wherein the ring having X, Y and Z is selected from the group consisting of



3. A compound of claim 2, wherein the ring having X, Y and Z is selected from the group consisting of



4. A compound of claim 1, wherein each R⁶ is independently selected from the group consisting of H, halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, C¹⁻³alkyl, C₁₋₃haloalkyl, substituted heteroaryl and unsubstituted heteroaryl.

5. A compound of claim 4 wherein each R⁶ is independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

6. A compound of claim 1, wherein Ar is a six-membered aryl optionally substituted with from 1 to 3 R⁶ substituents or a heteroaryl, optionally substituted with from 1 to 3 R⁶ substituents.


7. A compound of claim 6, wherein Ar is selected from the group consisting of optionally substituted six-membered aryl, optionally substituted pyridyl, optionally substituted pyrimidinyl, optionally substituted pyrazinyl, and optionally substituted pyridazinyl.

8. A compound of claim 1, wherein the subscript n is 2.

9. A compound of claim 1, wherein R¹ is selected from the group consisting of -X¹-C(O)R^a, -X¹-C(O)OR^a, optionally substituted aryl group and optionally substituted heteroaryl group.

10. A compound of claim 9, wherein R¹ is -C(O)OR^a, pyrimidine optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁₀alkyl, and C₁₋₁₀osubstituted alkyl, or oxadiazole optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, Ci-alkyl, and Ci-iosubstituted alkyl.

11. A compound of Formula II:



wherein,

X, Y, Z, D, and G are each independently selected from the group consisting of N and C(R³);

the subscript q is 0, 1, 2, 3 or 4;

the subscript k is 0, 1, 2 or 3;

R¹ is selected from the group consisting of H, C₁₋₁₀alkyl, Ci-iosubstituted alkyl, C₃₋₇cycloalkyl, C₃₋₇Subsited cycloalkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, -X^aC(O)R^a, -X^aC(O)OR^b, -X^{1-C(O)NR^aR^b, -S(O)₂R^a, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said cycloalkyl group, heterocyclyl group, aryl group and heteroaryl group is optionally substituted with from 1 to 4 substituents independently selected from the group consisting of halo, C₁₋₁₀alkyl, C₁₋₁₀iosubstituted alkyl, C₃₋₇cycloalkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, aryl, heteroaryl, -CN, -NR^aC(O)R^b, -NR^aC(0)NR^aR^b, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)R^b, and -S(O)₂NR^aR^b, or optionally R^a and R^b are combined to form a A-, 5- or 6-membered ring, and X¹ is selected from the group consisting of a bond, Ci-4alkylene, C₂₋₆alkenylene, C₂₋₆alkynylene, -C(O)-, and -C(O)(CH₂)₁₋₄-; wherein the aliphatic portions of X¹ are optionally substituted with one to three groups selected from the group consisting of halo, C₁₋₄alkyl, and C^asubstituted alkyl;}

each R² is independently selected from the group consisting of halo, C₁₋₅alkyl, C^asubstituted alkyl, C₃₋₇cycloalkyl, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -OR^a, -NR^aR^b, -NR^aC(0)R^b, -S(O)R^aR^b, -S(O)₂R^a and -S(O)₂NR^aR^b, and wherein when the subscript q is 2, 3, or 4, two R² groups can optionally cyclize to form a ring;

each R³ is independently selected from the group consisting of H, halo,

Ci-4 alkyl, C₁₋₄ haloalkyl, C₃₋₇ cycloalkyl, aryl and OR^a;

each R⁶ is independently selected from the group consisting of H, halo, C₁₋₁₀alkyl, C₁₋₁₀iosubstituted alkyl, C₃₋₇cycloalkyl, C₂₋₁₀ioalkenyl, C₂₋₁₀ioalkynyl, CN, NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(0)NR^aR^b, -NR^aC(0)R^b, -NR^aC(0)OR^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, S(O)₂R^a, -NR^aS(O)R^b, -NR^aS(O)₂R^b, -S(O)₂NR^aR^b, a 4- to 7-membered heterocyclyl group, aryl and a 5- to 10-membered heteroaryl group, wherein each of said heterocyclyl groups, said aryl and heteroaryl groups are optionally substituted with from one to four


substituents independently selected from halo, oxo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₃₋₇ cycloalkyl, -CN, -NO₂, -OR^a, -NR^aR^b, -C(O)R^a, -C(O)OR^a, -C(O)NR^aR^b, -NR^aC(O)R^b, -NR^aC(O)₂R^b, -NR^aC(O)NR^aR^b, -SR^a, -S(O)R^a, -S(O)₂R^a, -NR^aS(O)₂R^b, and -S(O)₂NR^aR^b or optionally R^a and R^b are combined to form a 4-, 5- or 6-membered ring;

and each R^a and R^b is independently selected from the group consisting of H, C₁₋₁₀ alkyl, Ci-iohaloalkyl, C₃₋₁₀iocycloalkyl, heterocyclyl, C₂₋₁₀ioalkenyl, C₂₋₁₀ioalkynyl, aryl, 5- to 6-membered heteroaryl and arylCi₁₋₄alkyl; and wherein the aliphatic portions of each of said R^a and R^b is optionally substituted with from one to three groups selected from the group consisting of halo, -ORⁿ, -OC(O)Rⁿ, -OC(O)N(Rⁿ)₂, -SRⁿ, -S(O)Rⁿ, -S(O)₂Rⁿ, -S(O)₂N(Rⁿ)₂, -NRⁿS(O)₂Rⁿ, -C(O)N(Rⁿ)₂, -C(O)Rⁿ, -NRⁿC(O)Rⁿ, -NRⁿC(O)N(Rⁿ)₂, -C(O)ORⁿ, -NRⁿC(O)ORⁿ, -CN, -NO₂, -N(Rⁿ)₂ and -NRⁿS(O)₂N(Rⁿ)₂, wherein each Rⁿ is independently H, C¹⁻⁶haloalkyl or an unsubstituted C₁₋₆ alkyl; each R⁷ and R⁸ is independently selected from the group consisting of H, Ci₁₋₆ alkyl, Ci₁₋₆ haloalkyl, C₃₋₇ cycloalkyl;

and pharmaceutically acceptable salts, solvates, stereoisomers, and esters thereof;

12. A compound of claim 11, wherein q is 0 or 1 and, both R⁷ and R⁸ are H.

13. A compound of claim 11, wherein the ring having X, Y, and Z is selected from the group consisting of



14. A compound of claim 13, wherein R³ is H.

15. A compound of claim 11, wherein R¹ is selected from the group consisting of -X¹-C(O)R^a, -X¹-C(O)OR^a, optionally substituted aryl and optionally substituted heteroaryl group.

16. A compound of claim 15, wherein R¹ is -C(O)OR^a, pyrimidine optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₃₋₆Substituted cycloalkyl, C₁₋₁₀alkyl, and C₁iosubstituted alkyl, or oxadiazole optionally substituted with from one to two substituents independently selected from the group consisting of halo, C₃₋₆cycloalkyl, C₁₋₁₀alkyl, and C₁₋₁₀osubstituted alkyl.

17. A compound of claim 11, wherein the subscript k is 1 or 2;

18. A compound of claim 17, wherein each R⁶ is independently selected from the group consisting of halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, Ci₁₋₃alkyl, C^haloalkyl substituted heteroaryl and unsubstituted heteroaryl.

19. A compound of claim 18 wherein k is 2, and each R⁶ is independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

20. A compound of claim 17 wherein R¹ is a heteroaryl group optionally substituted with from one to two substituents selected from the group consisting of halo, Ci₁₋₆alkyl, and Ci₁₋₆haloalkyl, and each R⁶ is independently selected from the group consisting of halo, -NR^aR^b, -NR^aC(O)R^b, -S(O)R^a, -S(O)₂R^a, C^halkyl, C^haloalkyl substituted heteroaryl and unsubstituted heteroaryl.

21. A compound of claim 20, wherein R¹ is a substituted pyrimidinyl or substituted oxadiazolyl group, and each R⁶ is independently selected from the group consisting of fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

22. A compound of claim 17, wherein k is 2, R¹ is a pyrimidinyl group optionally substituted with from one to two substituents selected from the group consisting of halo, C₁₋₆alkyl, and C^haloalkyl, and each R⁶ is independently selected from the group consisting of H, fluoro, chloro, methyl, ethyl, -CF₃, imidazolyl, triazolyl, tetrazolyl, -S(O)R^a, and -S(O)₂R^a.

23. A compound or a pharmaceutically acceptable salt, solvate, tautomer, stereoisomer, or ester thereof as disclosed in examples 1-1 13.

24. A method of raising intracellular levels of cyclic AMP (cAMP), said method comprising exposing a cell that expresses GPR1 19 to a compound of claim 11.

25. The method of claim 24, wherein said cell is a cell selected from the group consisting of a pancreatic cell, an islet cell, a beta cell, an intestinal endocrine cell, an L cell and a K cell.

26. A method of stimulating insulin production in a mammal, said method comprising administering an effective amount of a compound of claim 11 to said mammal.

27. A method of lowering blood glucose in a mammal, said method comprising administering an effective amount of a compound of claim 11 to said mammal.

28. A method of stimulating production of glucagon-like peptide 1 (GLP- 1) or glucose dependent insulinotropic polypeptide (GIP) in a mammal, said method comprising administering an effective amount of a compound of claim 11 to said mammal.

29. A method of treating diabetes in a mammal, said method comprising administering an effective amount of a compound of claim 11 to said mammal.

30. A method of treating type 2 diabetes in a mammal, said method comprising administering an effective amount of a compound of claim 11 to said mammal.


Figure 1**Tissue map of rat IC-GPCR2**

Figure 2**Tissue Map of mouse IC-GPCR2**