



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2563261 C 2013/04/09

(11)(21) **2 563 261**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2005/04/13
(87) Date publication PCT/PCT Publication Date: 2005/11/10
(45) Date de délivrance/Issue Date: 2013/04/09
(85) Entrée phase nationale/National Entry: 2006/10/06
(86) N° demande PCT/PCT Application No.: US 2005/012398
(87) N° publication PCT/PCT Publication No.: 2005/104846
(30) Priorité/Priority: 2004/04/14 (US60/562,071)

(51) Cl.Int./Int.Cl. *A01N 43/40* (2006.01),
A01N 25/14 (2006.01), *A01N 25/30* (2006.01),
A01P 7/04 (2006.01)

(72) Inventeurs/Inventors:
LIU, HONG, US;
DEXTER, ROBIN, US;
MARTIN, TIMOTHY, US;
MARTIN, CRAIG, US

(73) Propriétaire/Owner:
FMC CORPORATION, US

(74) Agent: SMART & BIGGAR

(54) Titre : COMPOSITIONS PESTICIDES DISPERSIBLES
(54) Title: DISPERSIBLE PESTICIDAL COMPOSITIONS

(57) Abrégé/Abstract:

Soluble granule pesticidal compositions comprise Flonicamid, a dispersant, and a wetting agent are disclosed.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 November 2005 (10.11.2005)

PCT

(10) International Publication Number
WO 2005/104846 A1

(51) International Patent Classification⁷: **A01N 43/40**

(21) International Application Number:
PCT/US2005/012398

(22) International Filing Date: 13 April 2005 (13.04.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/562,071 14 April 2004 (14.04.2004) US

(71) Applicant (for all designated States except US): **FMC CORPORATION** [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LIU, Hong** [US/US]; 19 Chicory Lane, Pennington, NJ 08534 (US). **DEXTER, Robin** [US/US]; 541 Freedom Drive, Yardley, PA 19067 (US). **MARTIN, Timothy** [US/US]; 1 High Mowing Road, Ringoes, NJ 08543 (US). **MARTIN, Craig** [US/US]; 109 Rice Drive, Morrisville, PA 19067 (US).

(74) Agents: **SHEEHAN, John, M.** et al.; FMC Corporation, 1735 Market Street, Philadelphia, PA 19103 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DISPERSIBLE PESTICIDAL COMPOSITIONS

(57) Abstract: Soluble granule pesticidal compositions comprise Flonicamid, a dispersant, and a wetting agent are disclosed.

WO 2005/104846 A1

76785-34

DISPERSIBLE PESTICIDAL COMPOSITIONS

5

Field of the Invention

The present invention relates to pesticidal compositions. In particular, the invention provides a soluble granule formulation of Flonicamid, a water-soluble pesticide, which is easily dispersed in an aqueous medium.

10

Background of the Invention

To enable the efficient elimination or controlling of unwanted pests, it is desirable to use effective chemical pesticides on these unwanted pests. Soluble granule formulations containing water-soluble pesticides are desirable in agricultural and related endeavors due to ease of handling and the elimination of organic solvents, which can be costly and dangerous to the environment. Safety, specifically eye irritation, caused by solvents, dispersants, clays and wetting agents in traditional formulations are also of concern. In addition, the process used to make the soluble granule formulation can be conducted in relatively simple equipment using relatively simple process steps.

20

The formulation of soluble granule pesticides has been disclosed in the prior art, for example in U.S. Patent 6,387,388 (Misselbrook et al.). Challenges in formulating a soluble granule of a water-soluble pesticide include the requirements that the final product successfully achieves (a) good dispersibility/solubility in an aqueous medium and (b) chemical stability once dispersed.

25

Good dispersibility/solubility in an aqueous medium is important in a soluble granule formulation since ineffective mixing of the pesticide into solution can result in reduced or inconsistent efficacy and, ultimately, costly returns of product and loss of sales. Chemical degradation is also important because there may be no physical sign of the degradation and resultant loss of efficacy until used on a pest. Examples of problematic chemical degradation include hydrolysis, oxidation, dehalogenation and bond cleavage.

30

76785-34

Summary of the Invention

The present invention is directed to a soluble granule composition comprising Flonicamid, a dispersant, and a wetting agent. This soluble granule composition has improved dispersibility/solubility as compared to other formulations.

5 According to one aspect of the present invention, there is provided a soluble granule composition, comprising: (a) Flonicamid; (b) a dispersant selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer sodium salt, alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate sodium salt; and (c) a wetting agent selected
10 from the group consisting of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent, polyalkylencoxide modified heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3-20 moles of ethylene oxide.

15 According to another aspect of the present invention, there is provided a soluble granule composition, comprising: (a) Flonicamid; (b) a dispersant selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer sodium salt, alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate sodium salt; (c) a wetting agent selected from
20 the group consisting of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent, polyalkylencoxide modified heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3-20 moles of ethylene oxide; and (d) a water-soluble carrier selected from the
25 group consisting of lactose, sucrose, glucose, starch, sodium tripolyphosphate and water-soluble salts.

 According to still another aspect of the present invention, there is provided a soluble granule composition described herein, comprising: (i) from 45% to 55% of Flonicamid; (j) from 2% to 6% of sodium alkyl naphthalene sulfonate

76785-34

formaldehyde condensate; (k) from 0.5% to 3% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85%; and (l) from 35% to 45% of lactose; wherein all % are % by weight based upon the total weight of all components in the

5 composition.

According to yet another aspect of the present invention, there is provided a process for preparing a soluble granule composition, said process comprising: (a) forming a mixture of Flonicamid, a dispersant, a wetting agent and, optionally, a water-soluble carrier; wherein the dispersant and the wetting agent are

10 as described herein; (b) kneading the mixture; (c) extruding the kneaded mixture from step (b); and (d) drying the extruded mixture from step (c).

76785-34

Detailed Description of the Invention

As used in this specification and unless otherwise indicated the term "Flonicamid" refers to N-cyanomethyl-4-trifluoromethyl nicotinamide, or salts thereof. The term "pesticide" refers to a molecule or combination of molecules that repels, retards, or kills pests, such as, but not limited to, deleterious or annoying insects, weeds, worms, fungi, bacteria, and the like, and can be used for crop protection, edifice protection, turf protection, or protection of a person; pesticide as used herein includes, but is not limited to, herbicides, insecticides, acaricides, fungicides, nematocides, ectoparasiticides, and growth regulators, either used to encourage growth of a desired plant species or retard growth of an undesired pest.

The modifier "about" is used herein to indicate that certain preferred operating ranges, such as ranges for molar ratios for reactants, material amounts, and temperature, are not fixedly determined. The meaning will often be apparent to one of ordinary skill. For example, a recitation of a temperature range of about 120° C to about 135° C in reference to, for example, an organic chemical reaction would be interpreted to include other like temperatures that can be expected to favor a useful reaction rate for the reaction, such as 105° C or 150° C. Where guidance from the experience of those of ordinary skill is lacking, guidance from the context is lacking, and where a more specific rule is not recited below, the "about" range shall be not more than 10% of the absolute value of an end point or 10% of the range recited, whichever is less.

As used in this specification and unless otherwise indicated the substituent terms "alkyl", "cycloalkyl", "alkoxy", "aryloxy", and "alkoxyarylamino", used alone or as part of a larger moiety, includes straight or branched chains of at least one or two carbon atoms, as appropriate to the substituent, and preferably up to 20 carbon atoms, more preferably up to ten carbon atoms, most preferably up to seven carbon atoms. "Halogen" or "halo" refers to fluorine, bromine, iodine, or chlorine. "Aryl" refers to an aromatic ring structure having 5 to 10 carbon atoms.

The term "ambient temperature" as utilized herein shall mean any suitable temperature found in a laboratory or other working environment, and is generally not below about 15° C nor above about 30° C.

The present invention is directed to a soluble granule composition comprising
5 Flonicamid, a dispersant, and a wetting agent. The invention can further comprise a water-soluble carrier.

Flonicamid can be present in a concentration of from 1% by weight to 94% by weight, more particularly 45% by weight to 55% by weight, based upon the total weight of all components in the composition.

10 The dispersant can be selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer (sodium salt), alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate (Na) salt. Preferably, the dispersant is sodium alkyl naphthalene sulfonate formaldehyde condensate. The dispersant can be present in a concentration of from 0.5% by
15 weight to 20% by weight, more particularly 2% by weight to 6% by weight, based upon the total weight of all components in the composition.

The wetting agent can be selected from the group consisting of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is up to 85 weight percent, polyalkylencoxide modified
20 heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3 – 20 moles of ethylene oxide. Preferably, the wetting agent is a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent. The wetting agent can be present in a concentration of from 0.1% by weight to 20%
25 by weight, more particularly 0.5% by weight to 3% by weight, based upon the total weight of all components in the composition.

The water-soluble carrier can be selected from the group consisting of lactose, sucrose, glucose, starch, sodium tripolyphosphate and water-soluble salts. Preferably, the water-soluble carrier is lactose. The water-soluble carrier can be
30 present in a concentration of from 0.001% by weight to 97% by weight, more particularly 35% by weight to 45% by weight, based upon the total weight of all components in the composition.

Another embodiment of the present invention is a soluble granule composition

comprising Flonicamid; a dispersant selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer (sodium salt), alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate (Na) salt; a wetting agent selected from the group consisting of a mixture of sodium
5 dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent, polyalkylencoxide modified heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3 – 20 moles of ethylene oxide; and a water-soluble carrier selected from the group consisting of lactose, sucrose, glucose, starch, sodium
10 tripolyphosphate and water-soluble salts.

Another embodiment of the present invention is a soluble granule composition comprising from 1% to 94% of Flonicamid, from 0.5% to 20% of sodium alkyl naphthalene sulfonate formaldehyde condensate, from 0.1% to 20% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium
15 dioctylsufosuccinate is present in a concentration up to 85%, and from 0.001% to 97% of lactose, wherein all % are % by weight based upon the total weight of all components in the composition.

Yet another embodiment of the present invention is a soluble granule composition comprising from 25% to 88% of Flonicamid, from 0.5% to 20% of
20 sodium alkyl naphthalene sulfonate formaldehyde condensate, from 0.4% to 5.0% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85%, and from 6% to 70% of lactose, wherein all % are % by weight based upon the total weight of all components in the composition.

25 Yet another embodiment of the present invention is a soluble granule composition comprising from 45% to 55% of Flonicamid, from 2% to 6% of sodium alkyl naphthalene sulfonate formaldehyde condensate, from 0.5% to 3% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium
30 dioctylsufosuccinate is present in a concentration up to 85%, and from 35% to 45% of lactose, wherein all % are % by weight based upon the total weight of all components in the composition.

The composition may further comprise an anti-foam agent and/or an anti-bacteria agent.

The present invention also encompasses a method of controlling unwanted pests, comprising applying to an area infested with such pests an effective amount of the soluble granule composition, discussed above, in an aqueous solvent.

5 The present invention also encompasses a process for preparing the soluble granule composition comprising a) forming a mixture of Flonicamid, a dispersant, a wetting agent and, optionally, a water-soluble carrier; b) kneading the mixture; c) extruding the kneaded mixture from step b); and d) drying the extruded mixture from step c). Preferably, the process further comprises milling the mixture prior to kneading.

10

The pesticidal compositions of the present invention are further illustrated by the examples below. Flonicamid used in all examples contains 97% active ingredient. The examples serve only to illustrate the invention and should not be interpreted as limiting since further modifications of the disclosed invention will be
15 apparent to those skilled in the art. All such modifications are deemed to be within the scope of the invention as defined in the claims.

In the examples below, testing to demonstrate the dispersibility/solubility of the composition was carried out in the following manner: 1) At ambient temperature, 1.0 gram of the composition prepared in the example was added to 100 ml of water
20 in a cylinder of appropriate size; 2) the cylinder was repeatedly inverted 180 degrees to effect dispersion/solution of the composition into the water; 3) the number of inversions required for full dispersion/solution to a clear state was determined visually and recorded. The results of these tests are presented in each example.

25

Example 1

This example illustrates the preparation of a 50SG composition (Composition A) of the present invention.

30 A mixture of 52.0 grams of Flonicamid, 5.0 grams of sodium alkyl naphthalene sulfonate formaldehyde condensate, 1.0 gram of a combination of sodium dioctylsulfosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) and 42.0 grams of lactose were combined and agitated under high shear. To this mixture was added 14.0 grams of water and then

the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

5 Composition A required 3 inversions for full dispersion into a clear solution.

Example 2

This example illustrates the preparation of a 50SG composition (Composition B) of the present invention.

10 A mixture of 52.0 grams of Flonicamid, 5.0 grams of sodium alkyl naphthalene sulfonate formaldehyde condensate, 1.0 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) and 42.0 grams of ammonium sulfate were combined and agitated under high shear. To this mixture was added 14.0 grams of
15 water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

20 Composition B required 8 inversions for full dispersion into a clear solution.

Example 3

This example illustrates the preparation of a 50SG composition (Composition C) of the present invention.

25 A mixture of 52.0 grams of Flonicamid, 4.0 grams of modified acrylate copolymer (sodium salt), 1.0 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) and 43.0 grams of sodium tripolyphosphate were combined and agitated under high shear. To this mixture was added 14.0 grams of water and then
30 the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

Composition C required 7 inversions for full dispersion into a clear solution.

Example 4

5 This example illustrates the preparation of a 24SG composition (Composition D) of the present invention.

A mixture of 25.0 grams of Flonicamid, 5.0 grams of sodium alkyl naphthalene sulfonate formaldehyde condensate, 0.5 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ), 68.45 grams of lactose, 1.0 gram of an anti-
10 bacteria agent (Dowcide A available from Dow Chemical Company in Midland Michigan) and 0.05 gram of an anti-foam agent were combined and agitated under high shear. To this mixture was added 13.0 grams of water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The
15 resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

Composition D required 5 inversions for full dispersion into a clear solution.

Example 5

20 This example illustrates the preparation of a 24SG composition (Composition E) of the present invention.

A mixture of 25.0 grams of Flonicamid, 5.0 grams of sodium alkyl naphthalene sulfonate formaldehyde condensate, 0.5 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec
25 Industries, Inc. in West Paterson, NJ) and 69.5 grams of lactose were combined and agitated under high shear. To this mixture was added 13.0 grams of water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to
30 15 minutes until the moisture content of the product was less than 1%.

Composition E required 3 inversions for full dispersion into a clear solution.

Example 6

This example illustrates the preparation of an 85SG composition (Composition F) of the present invention.

5 A mixture of 87.6 grams of Flonicamid, 5.0 grams of sodium alkylnaphthalene sulfonate formaldehyde condensate, 1.0 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) and 6.4 grams of lactose were combined and agitated under high shear. To this mixture was added 15.0 grams of water and then
10 the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

Composition F required 3 inversions for full dispersion into a clear solution.

15

Example 7

This example illustrates the preparation of a 50SG composition (Composition G) of the present invention.

A mixture of 52.0 grams of Flonicamid, 3.0 grams of sodium alkylnaphthalene
20 sulfonate formaldehyde condensate, 1.0 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ), 41.5 grams of lactose and 2.5 grams of a mixture of polyalkyleneoxide modified heptamethyl trisiloxane and alkyloxypolyethylene glycol methyl ether (Silwet L-77 available from OSi
25 Specialties, a Crompton Business in Greenwich, CT) were combined and agitated under high shear. To this mixture was added 14.0 grams of water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to
30 15 minutes until the moisture content of the product was less than 1%.

Composition G (after being stored at -10 C) required 2 inversions for full dispersion into a clear solution.

76785-34

Example 8

This example illustrates the preparation of a 70SG composition (Composition H) of the present invention.

5 A mixture of 72.0 grams of Flonicamid, 5.0 grams of sodium alkylnaphthalene sulfonate formaldehyde condensate, 1.0 gram of a combination of sodium dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) and 22.0 grams of lactose were combined and agitated under high shear. To this mixture was added 15.0 grams of water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded
10 mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant product was then dried at 60 °C with 80% airflow for 10 to 15 minutes until the moisture content of the product was less than 1%.

Composition H required 3 inversions for full dispersion into a clear solution.

15

Example 9

This example illustrates the preparation of a 91SG composition (Composition I) of the present invention.

A mixture of 94.0 grams of Flonicamid, 5.0 grams of sodium alkylnaphthalene sulfonate formaldehyde condensate and 1.0 gram of a combination of sodium
20 dioctylsufosuccinate and sodium benzoate (Aerosol OTB available from Cytec Industries, Inc. in West Paterson, NJ) were combined and agitated under high shear. To this mixture was added 30.0 grams of water and then the mixture was kneaded in a Hobart mixer for about 2 minutes. The kneaded mixture was then extruded in a Fuji Paudal DGL-1 using a dome with 0.8-millimeter apertures. The resultant
25 product was then dried at 60 °C with 80% airflow for 5 to 10 minutes until the moisture content of the product was less than 1%.

Composition I required 3 inversions for full dispersion into a clear solution.

76785-34

CLAIMS:

1. A soluble granule composition, comprising:
 - (a) Flonicamid;
 - (b) a dispersant selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer sodium salt, alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate sodium salt; and
 - (c) a wetting agent selected from the group consisting of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent, polyalkylencoxide modified heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3-20 moles of ethylene oxide.
2. The composition of claim 1, further comprising a water-soluble carrier.
3. The composition of claim 1 or 2, wherein the dispersant is sodium alkyl naphthalene sulfonate formaldehyde condensate.
4. The composition of any one of claims 1 to 3, wherein the dispersant is present in a concentration of from 0.5% by weight to 20% by weight based upon the total weight of all components in the composition.
5. The composition of claim 4, wherein the dispersant is present in a concentration of from 2% by weight to 6% by weight based upon the total weight of all components in the composition.
6. The composition of any one of claims 1 to 5, wherein the wetting agent is a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent.

76785-34

7. The composition of any one of claims 1 to 6, wherein the wetting agent is present in a concentration of from 0.1% by weight to 20% by weight based upon the total weight of all components in the composition.

8. The composition of claim 7, wherein the wetting agent is present in a
5 concentration of from 0.5% by weight to 3% by weight based upon the total weight of all components in the composition.

9. The composition of any one of claims 2 to 8, wherein the water-soluble carrier is selected from the group consisting of lactose, sucrose, glucose, starch, sodium tripolyphosphate and water-soluble salts.

10 10. The composition of claim 9, wherein the water-soluble carrier is lactose.

11. The composition of any one of claims 2 to 10, wherein the water-soluble carrier is present in a concentration of from 0.001% by weight to 97% by weight based upon the total weight of all components in the composition.

12. The composition of claim 11, wherein the water-soluble carrier is
15 present in a concentration of from 35% by weight to 45% by weight based upon the total weight of all components in the composition.

13. The composition of any one of claims 1 to 12, wherein Flonicamid is present in a concentration of from 1% by weight to 94% by weight based upon the total weight of all components in the composition.

20 14. The composition of claim 13, wherein Flonicamid is present in a concentration of from 45% by weight to 55% by weight based upon the total weight of all components in the composition.

15. The composition of any one of claims 1 to 14, further comprising an anti-foam agent.

76785-34

16. The composition of any one of claims 1 to 15, further comprising an anti-bacteria agent.

17. A soluble granule composition, comprising:

(a) Flonicamid;

5 (b) a dispersant selected from the group consisting of sodium alkyl naphthalene sulfonate formaldehyde condensate, modified acrylate copolymer sodium salt, alkyl naphthalene sulfonate salts and dodecyl benzene sulfonate sodium salt;

10 (c) a wetting agent selected from the group consisting of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85 weight percent, polyalkylencoxide modified heptamethyl trisiloxane, alkyloxypolyethylene glycol methyl ether and alcohol (C8-C16) ethoxylates with 3-20 moles of ethylene oxide; and

15 (d) a water-soluble carrier selected from the group consisting of lactose, sucrose, glucose, starch, sodium tripolyphosphate and water-soluble salts.

18. A soluble granule composition of claim 17, comprising:

(a) from 1% to 94% of Flonicamid;

20 (b) from 0.5% to 20% of sodium alkyl naphthalene sulfonate formaldehyde condensate;

(c) from 0.1% to 20% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85%; and

(d) from 0.001% to 97% of lactose;

76785-34

wherein all % are % by weight based upon the total weight of all components in the composition.

19. A soluble granule composition of claim 18, comprising:

(e) from 25% to 88% of Flonicamid;

5 (f) from 0.5% to 20% of sodium alkyl naphthalene sulfonate formaldehyde condensate;

(g) from 0.4% to 5.0% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85%; and

10 (h) from 6% to 70% of lactose;

wherein all % are % by weight based upon the total weight of all components in the composition.

20. A soluble granule composition of claim 19, comprising:

(i) from 45% to 55% of Flonicamid;

15 (j) from 2% to 6% of sodium alkyl naphthalene sulfonate formaldehyde condensate;

(k) from 0.5% to 3% of a mixture of sodium dioctylsufosuccinate and sodium benzoate wherein the sodium dioctylsufosuccinate is present in a concentration up to 85%; and

20 (l) from 35% to 45% of lactose;

wherein all % are % by weight based upon the total weight of all components in the composition.

76785-34

21. A method for controlling unwanted pests, said method comprising applying to an area infested with such pests an effective amount of the composition of any one of claims 1 to 20, in an aqueous solvent.

22. A process for preparing a soluble granule composition, said process
5 comprising:

(a) forming a mixture of Flonicamid, a dispersant, a wetting agent and, optionally, a water-soluble carrier; wherein the dispersant and the wetting agent are as defined in claim 1;

(b) kneading the mixture;

10 (c) extruding the kneaded mixture from step (b); and

(d) drying the extruded mixture from step (c).

23. The process of claim 22, further comprising milling the mixture prior to kneading.