



(51) International Patent Classification:

C07K 14/475 (2006.01) C07K 16/28 (2006.01)  
A61K 38/00 (2006.01) C07K 14/765 (2006.01)  
C07K 16/22 (2006.01) C07K 14/47 (2006.01)

(21) International Application Number:

PCT/EP2016/057272

(22) International Filing Date:

1 April 2016 (01.04.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

15162502.7 2 April 2015 (02.04.2015) EP  
15162511.8 2 April 2015 (02.04.2015) EP

(71) Applicant: MOLECULAR PARTNERS AG [CH/CH];  
Wagistrasse 14, 8952 Zürich-Schlieren (CH).

(72) Inventors: BAKKER, Talitha; Schürenstrasse 1b, 8903  
Birmensdorf (CH). STUMPP, Michael T.; Chratzstrasse  
6, 8954 Geroldswil (CH). BINZ, Hans Kaspar; Bach-  
strasse 27, 8903 Birmensdorf (CH). PHILLIPS, Douglas;  
Eulenweg 37, 5400 Baden (CH). DOLADO, Ignacio;  
Quellenstrasse 39A, 4310 Rheinfelden (CH). FORRER,  
Patrik; Rainackerstrasse 17, 8953 Dietikon (CH). MERZ,  
Frieder W.; Im Geerig 79, 5507 Mellingen (CH). SON-  
DEREGGER, Ivo; Im Baurenacker 9, 8902 Urdorf (CH).  
STEINER, Daniel; Langmauerstrasse 28, 8006 Zürich  
(CH). GULOTTI-GEORGIEVA, Maya; Niederweg 36,  
8907 Wettswil a.A. (CH). ABRAM SALIBA, Johan;  
Chemin du Calvaire 19, 1005 Lausanne (CH).

(74) Agents: MICHALSKI HÜTTERMANN & PARTNER  
PATENTANWÄLTE MBB et al.; Speditionstraße 21,  
40221 Düsseldorf (DE).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,  
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,  
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,  
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,  
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,  
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,  
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,  
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: DESIGNED ANKYRIN REPEAT DOMAINS WITH BINDING SPECIFICITY FOR SERUM ALBUMIN

(57) Abstract: New designed ankyrin repeat domains with binding specificity for serum albumin, recombinant binding proteins comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, as well as recombinant binding proteins comprising at least one designed ankyrin repeat domain with binding specificity for hepatocyte growth factor (HGF), at least one designed ankyrin repeat domain with binding specificity for vascular endothelial growth factor (VEGF-A), and at least two designed ankyrin repeat domain with binding specificity for serum albumin are described, as well as nucleic acids encoding such designed ankyrin repeat domains and recombinant binding proteins, pharmaceutical compositions comprising such designed ankyrin repeat domains, recombinant binding proteins or nucleic acids and the use of such designed ankyrin repeat domains, recombinant binding proteins, nucleic acids or pharmaceutical compositions in the treatment of diseases.



## DESIGNED ANKYRIN REPEAT DOMAINS WITH BINDING SPECIFICITY FOR SERUM ALBUMIN

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application claims the benefit of and the priority to European patent application EP 15162502 filed on 2 April 2015 with the European Patent Office. The present application also claims the benefit of and the priority to European patent application EP 15162511 filed on 2 April 2015 with the European Patent Office. The contents of European patent applications EP 15162502 and EP 15162511 is incorporated herein by reference for all purposes in its entirety including all tables, figures, and claims - as well as including an incorporation of any element or part of the description, claims or drawings not contained herein and referred to in Rule 20.5(a) of the PCT, pursuant to Rule 4.18 of the PCT.

FIELD OF THE DISCLOSURE

Provided is a new designed ankyrin repeat domain with binding specificity for serum albumin exhibiting improved storage stability properties. Provided are also recombinant binding proteins comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, which exhibit improved pharmacokinetic properties compared to the recombinant binding proteins comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. Particularly provided are recombinant binding proteins comprising at least one designed ankyrin repeat domain with binding specificity for hepatocyte growth factor (HGF), comprising at least one designed ankyrin repeat domain with binding specificity for vascular endothelial growth factor A (VEGF-A), and comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin. Furthermore provided are nucleic acids encoding such designed ankyrin repeat domains and/or recombinant binding proteins, pharmaceutical compositions comprising such designed ankyrin repeat domains, recombinant binding proteins or nucleic acids, and the use of such designed ankyrin repeat domains, recombinant binding proteins, nucleic acids, or pharmaceutical compositions in the treatment of a disease.

## BACKGROUND

The following discussion of the background is merely provided to aid the reader in understanding the invention and is not admitted to describe or constitute prior art to the present invention.

5 There are, beside antibodies, novel binding proteins or binding domains that can be used to specifically bind a target molecule (e.g. Binz, H.K., Amstutz, P., Plückthun, A., Nat. Biotechnol. 23, 1257-1268, 2005). One such novel class of binding proteins or binding domains not possessing an Fc are based on designed repeat proteins or designed repeat domains, such as designed ankyrin repeat proteins or designed ankyrin repeat domains  
10 (WO 2002/020565; Binz, H.K., Amstutz, P., Kohl, A., Stumpp, M.T., Briand, C., Forrer, P., Grütter, M.G., Plückthun, A., Nat. Biotechnol. 22, 575-582, 2004). WO 2002/020565 describes how large libraries of repeat proteins, such as ankyrin repeat proteins, can be constructed, and their general application. WO 2012/069654 describes recombinant binding proteins comprising a designed ankyrin repeat domain with binding specificity for  
15 serum albumin. WO 2010/060748 describes recombinant binding proteins comprising designed ankyrin repeat domains with binding specificity for VEGF-A, and WO 2011/135067 describes modified versions of such recombinant binding proteins specific for binding to VEGF-A. WO 2014/191574 describes recombinant binding proteins comprising designed ankyrin repeat domains with binding specificity for HGF. None of  
20 these patent applications discloses a recombinant binding protein comprising a designed ankyrin repeat domain with binding specificity for VEGF-A and a designed ankyrin repeat domain with binding specificity for HGF.

Unlike e.g. IgG antibodies, which exhibit long systemic half-lives mediated by FcRn recycling, proteins comprising designed ankyrin repeat domains typically exhibit a fast  
25 pharmacokinetic clearance and short terminal half-lives, unless the protein comprises elements that improve the pharmacokinetic properties, such as e.g. a designed ankyrin repeat domain with binding specificity to serum albumin described in WO 2012/069654. Using serum albumin binding for improving pharmacokinetic properties of proteins is a process well-known in the art (see e.g. WO 9101743; Frejd F.Y., 2012 (in Kontermann, R  
30 (Ed.) "Therapeutic proteins: strategies to modulate their plasma half-lives", Wiley-VCH Verlag GmbH, 2012, ISBN 978-3-527-32849-9); Nguyen, A., Reyes, A.E.II., Zhang, M., McDonald, P., Wong, W.L., Damico, L.A., Dennis, M.S. Protein Eng. Des. Sel. 19, 291-297, 2006; WO 2008/096158; WO 2006/122787; WO 2011/095545; and  
35 WO 2012/069654). In order to be able to use designed ankyrin repeat domains with binding specificity for serum albumin in clinical drug candidates, the storage stability of

known designed ankyrin repeat domains with binding specificity for serum albumin has to be improved. Disclosed herein are designed ankyrin repeat domains with binding specificity for serum albumin with improved properties.

The effect of valency of designed ankyrin repeat domains with binding specificity for serum albumin on the pharmacokinetic properties of recombinant binding proteins has not  
5 been investigated. Based on findings of the albumin binding domain (Hopp, J., Horning, N., Zettlitz, K.A., Schwarz, A., Fuss, N., Müller, D., Kontermann, R.E. *Protein Eng. Des. Sel.* 23, 827-834, 2010), one skilled in the art would expect that a recombinant binding protein comprising two albumin binding protein domains such as designed ankyrin repeat  
10 domains with binding specificity for serum albumin would not have improved pharmacokinetic properties compared to a recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. Surprisingly, we found that this is not the case. Disclosed are thus recombinant binding proteins comprising at least two designed ankyrin repeat domains with binding specificity  
15 for serum albumin that exhibit improved pharmacokinetic properties (i.e. prolonged terminal half-lives, increased exposures, reduced clearance, and/or increased percentages of injected dose) compared to recombinant binding proteins comprising only one designed ankyrin repeat domain with binding specificity for serum albumin.

Neovascularisation (new blood vessel formation) is widely known to play an important role  
20 in the development and maintenance of tumors (Ferrera, N., and Kerbel, R.S., *Nature* 438, 967-974, 2005). Accordingly, the inhibition of angiogenesis has become a main cornerstone in modern clinical oncology; especially the targeting of vascular endothelial growth factor (VEGF) and its receptors (Hurwitz, H., *Clin. Colorectal Cancer, Suppl.* 2, 62-68, 2004; Escudier, B., *Clin. Adv. Hematol. Oncol.* 5, 530-531, 2007). The mammalian  
25 VEGF family consists of five glycoproteins referred to as VEGF-A, VEGF-B, VEGF-C, VEGF-D (also known as FIGF) and placenta growth factor (PlGF, also known as PGF). VEGF-A has been shown to be an effective target for anti-angiogenic therapy (Weis, S.M., and Cheresh, D.A., *Nat. Med.* 17, 1359-1370, 2011). The VEGF-A ligands bind to and activate three structurally similar type III receptor tyrosine kinases, designated VEGFR-1  
30 (also known as FLT1), VEGFR-2 (also known as KDR) and VEGFR-3 (also known as FLT4). Several angiogenesis inhibitors have received regulatory approval to date showing a prolonged progression-free survival (PFS) and/or overall survival in various cancer types in combination with chemotherapy. Unfortunately, resistance inevitably occurs during the course of treatment with VEGF/VEGFR inhibitors, such as the VEGF-A inhibitor  
35 bevacizumab (Avastin®), suggesting that concomitant inhibition of additional targets and

resistance pathways may be necessary to achieve superior clinical results (Kerbel, R.S., *N. Engl. J. Med.* 358, 2039-2049, 2008; Hurwitz, 2004, loc. cit.; Escudier, 2007, loc. cit.).

cMet tyrosine kinase is a cell surface receptor for hepatocyte growth factor (HGF, also known as scatter factor, SF) primarily expressed on epithelial cells (Comoglio, P.M., Giordano, S., and Trusolino, L., *Nat. Rev. Drug Discov.* 7, 504-516, 2008). While cMet and HGF are expressed at low levels in normal adult tissues, their expression is frequently up regulated in a broad range of human tumors, which has been correlated in preclinical models with tumor cell survival, growth, angiogenesis, invasion and metastasis (Rong, S., Segal, S., Anver, M., Resau, J.H., Vande Woude, G.F., *Proc. Natl. Acad. Sci. USA* 91, 4731-4735, 1994; Michieli, P., Mazzone, M., Basilico, C., Cavassa, S., Sottile, A., Naldini, L., Comoglio, P.M., *Cancer Cell* 6, 61-73, 2004). Up-regulation of HGF and/or cMet expression and signaling has been found to be associated with poor prognosis and drug resistance in many tumor types in the clinic (Fasolo, A., Sessa, C., Gianni, L., Broggin, M., *Ann. Oncol.* 24,14-20, 2013). Altogether this indicates that the HGF-cMet axis is an important target for therapeutic intervention (Comoglio, 2008 loc. cit.; Fasolo et al., 2013, loc. cit.). Through binding to its receptor, HGF mediates a number of cellular responses, including scattering of various cell types, the formation of tubules and lumens, epithelial-mesenchymal transition, angiogenesis, liver regeneration, wound healing and embryological development. The HGF/c-Met signaling pathway has also been shown to play a role in various diseases, including many human solid tumors, in which it participates in tumor development, invasion and metastasis. Current HGF/cMet pathway inhibitors in phase II or III clinical development comprise monoclonal antibodies (mAbs) targeting the extracellular domain of cMet (i.e. MetMab from Genentech-Roche) or small molecule inhibitors of its intracellular kinase domain. Small molecule inhibitors such as tivantinib (ArQule®) and cabozantinib (Cometriq®) are very potent but less specific than mAbs and bear the potential for higher toxicity. Biological agents against HGF/SF include rilotumumab (AMG102), a humanised mAb against HGF, and ficlatuzumab (AV-299), a humanised anti-HGF IgG1. The use of HGF/cMet inhibitors in combination with other targeted agents is an active field of investigation which aims to simultaneously inhibit various signaling pathways that have redundant or synergistic tumor functions. HGF/cMet triggers potent angiogenic signals that act synergistically with VEGF in inducing new tumor blood vessels and can induce resistance to anti-angiogenic therapy such as Avastin® and Sutent® (sunitinib) in glioblastoma (Jahangiri, A., De Lay, M., Miller, L.M., Carbonell, W.S., Hu, Y.L., Lu, K., Tom, M.W., Paquette, J., Tokuyasu, T.A., Tsao, S., Marshall, R., Perry, A., Bjorgan, K.M., Chaumeil, M.M., Ronen, S.M., Bergers, G., Aghi, M.K., *Clin. Cancer Res.* 19, 1773-1783, 2013) and renal cell cancer, respectively.

There are currently a number of anti-HGF/cMet compounds under investigation in combination with other targeted agents such as anti-VEGF receptor inhibitors, which have demonstrated a favorable safety profile in a variety of tumor types (Sharma, P.S., Sharma, R., Tyagi, T. *Curr. Cancer Drug Targets.* 11, 624-653, 2011). However, such combination therapy approaches imply that the patient must receive two separate treatments, each with a different safety profile, which may lead to increased undesirable toxicities, which in turn may limit the medical treatment options. Furthermore, different treatments may be subjected to different administration schemes, which could make the dosing more burdensome for the patient. Last but not least, the dosing of various agents simultaneously may significantly increase the costs associated to treatment and patient care.

One commercially available drug with dual cMet and VEGF inhibitory activity is cabozantinib (Cometriq®; a small molecule drug), an oral, multi-specific tyrosine kinase inhibitor targeting cMet and VEGFR 1-3 (in addition to RET, KIT, AXL and FLT3). Cabozantinib has validated the clinical approach of simultaneously inhibiting HGF and VEGF in tumors with a single agent (Yakes, F.M., Chen, J., Tan, J., Yamaguchi, K., Shi, Y., Yu, P., Qian, F., Chu, F., Bentzien, F., Cancilla, B., Orf, J., You, A., Laird, A.D., Engst, S., Lee, L., Lesch, J., Chou, Y.C., Joly, A.H., *Mol. Cancer Ther.* 10, 2298-2308, 2011; Castellone, M.D., Carlomagno, F., Salvatore, G., Santoro, M., *Best Pract. Res. Clin. Endocrinol. Metab.* 22, 1023-1038, 2008). For instance, in castration resistant prostate cancer, an indication where the anti-HGF mAb rilotumumab failed to demonstrate efficacy as single agent in phase II studies, cabozantinib showed anti-tumor activity in a high percentage of patients in phase II (Smith, D.C., Smith, M.R., Sweeney, C., Elfiky, A.A., Logothetis, C., Corn, P.G., Vogelzang, N.J., Small, E.J., Harzstark, A.L., Gordon, M.S., Vaishampayan, U.N., Haas, N.B., Spira, A.I., Lara, P.N. Jr., Lin, C.C., Srinivas, S., Sella, A., Schöffski, P., Scheffold, C., Weitzman, A.L., Hussain, M., *J. Clin. Oncol.* 31, 412-419, 2013). However, activity was paralleled with a high incidence of adverse events that led to dose reductions in 62% of patients, raising doubts on the safety and tolerability of such pleiotropic modes of action.

Simultaneous targeting of VEGF-A and HGF/cMet may beneficially disrupt angiogenesis and tumor progression. As described hereinbefore, current therapies acting simultaneously on the VEGF-A/VEGFR-2 and the HGF/cMet-pathways either are based on single therapeutics that are unspecific and lead to safety findings, or involve several specific therapeutics that have to be combined, resulting in a need of co-administration or multiple administrations. Furthermore, some of the current drugs exhibit short systemic

half-lives. Thus, there is a need to provide improved drugs blocking the VEGF-A/VEGFR-2 and the HGF/cMet pathways. This is technically difficult to achieve with antibody drugs, which further suffer from the need of laborious production in mammalian cells. Provided herein are recombinant binding proteins that address these issues. In some embodiments  
5 a recombinant binding protein provided herein comprises at least one designed ankyrin repeat domain with binding specificity for VEGF-A, at least one designed ankyrin repeat domain with binding specificity for HGF, and, for pharmacokinetic property improvement, at least two designed ankyrin repeat domains with binding specificity for serum albumin.

## 10 SUMMARY

The present invention relates to a new designed ankyrin repeat domain with binding specificity for serum albumin comprising the amino acid sequence of SEQ ID NO: 50, which exhibits improved storage stability over known designed ankyrin repeat domains with binding specificity for serum albumin. In one embodiment, the invention relates to a  
15 recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said designed ankyrin repeat domains with binding specificity for serum albumin each comprise SEQ ID NO: 50. In one embodiment, the invention relates to a recombinant binding protein comprising a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat  
20 domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin and comprise the amino acid sequence of SEQ ID NO: 50. In one embodiment, said first, second, third and fourth designed ankyrin repeat domains of said recombinant binding protein are in the  
25 order third-second-first-fourth from N terminus to C terminus. In one embodiment, said first designed ankyrin repeat domain of said recombinant binding protein comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21 and amino acid sequences in which up to 10 amino acids of SEQ ID NOs: 12 to 21 are exchanged by any amino acid, and said second designed ankyrin  
30 repeat domain of said recombinant binding protein comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37 and amino acid sequences in which up to 10 amino acids of SEQ ID NOs: 23 to 37 are exchanged by any amino acid, and said third and fourth designed ankyrin repeat domains of said recombinant binding protein each comprise amino acid sequence SEQ ID NO: 50,  
35 and said designed ankyrin repeat domains are linked by polypeptide linkers comprising

amino acid sequences selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9 and amino acid sequences in which up to 4 amino acids of SEQ ID NOs: 2 to 9 are exchanged by any amino acid. In one embodiment, said first designed ankyrin repeat domain of said recombinant binding protein comprises amino acid sequences SEQ ID NO: 18, and said second designed ankyrin repeat domain of said recombinant binding protein comprises amino acid sequence SEQ ID NO: 26, and said third and fourth designed ankyrin repeat domains of said recombinant binding protein each comprise amino acid sequence SEQ ID NO: 50, and said designed ankyrin repeat domains are linked by polypeptide linkers consisting of amino acid sequence SEQ ID NO: 9. In one embodiment, the invention relates to a recombinant binding protein comprising an amino acid sequence that has at least 90% amino acid sequence identity with the amino acid sequence of SEQ ID NO: 134. In one embodiment, the invention relates to a recombinant binding protein comprising the amino acid sequence consisting of the amino acid sequence of SEQ ID NO: 134. In a preferred embodiment, the invention relates to a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134.

The invention further relates to a nucleic acid encoding the amino acid sequence of a designed ankyrin repeat domain or a recombinant binding protein of the invention.

The invention also relates to a pharmaceutical composition comprising a recombinant binding protein and/or a designed ankyrin repeat domain or a nucleic acid of the present invention, and optionally a pharmaceutical acceptable carrier and/or diluent.

The invention also relates to the use of the pharmaceutical composition of the invention for the treatment of a disease. In one embodiment, it relates to the use of the pharmaceutical composition of the invention for the treatment of cancer, gastric cancer, or renal cancer.

25

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Illustration of recombinant binding proteins comprising designed ankyrin repeat domains with binding specificity for serum albumin.

(a) Illustration of a designed ankyrin repeat domain with binding specificity for serum albumin. Examples of such ankyrin repeat domains are designed ankyrin repeat domains with an amino acid sequence selected from the group consisting of SEQ ID NOs: 40 to 56, in particular the designed ankyrin repeat domain with amino acid sequence of SEQ ID NO: 50. (b) Illustration of a designed ankyrin repeat domain with binding specificity for another target than serum albumin. Examples of such ankyrin repeat domains are



designed ankyrin repeat domains with an amino acid sequence selected from the group consisting of SEQ ID NOs: 12 to 39. (c) Illustration of a polypeptide linker (for example a polypeptide with an amino acid sequence corresponding to any of SEQ ID NOs: 2 to 9). (d) Illustration of an N-terminal amino acid sequence. Examples for such N-terminal amino acid sequences are for example the sequences MGS or GS, or polypeptide tags, as exemplified by the amino acid sequence corresponding to SEQ ID NO: 1. (e) Illustration of a bioactive compound. Such a moiety can for example be a protein or protein domain with e.g. agonistic (e.g. hormone, or enzyme), antagonistic (e.g. receptor domain or antibody fragment), or toxic (e.g. toxin) activity. Such a moiety can for example also be a small molecule compound exhibiting e.g. agonistic, antagonistic, or toxic activity. (f) Illustration of a recombinant binding protein provided herein comprising two designed ankyrin repeat domains with binding specificity for serum albumin, and one designed ankyrin repeat domain with binding specificity for another target than serum albumin, linked by polypeptide linkers and having an N-terminal amino acid sequence. For example a recombinant binding protein with an amino acid sequence corresponding to any of SEQ ID NOs: 73 to 81 consist of such three designed ankyrin repeat domains, wherein SEQ ID NOs: 73, 75, 78, and 80 have the two designed ankyrin repeat domains with binding specificity for serum albumin flanking the respective third designed ankyrin repeat domain as shown in the illustration. (g) Illustration of a recombinant binding protein provided herein comprising two designed ankyrin repeat domains with binding specificity for serum albumin, and two designed ankyrin repeat domains with binding specificities for other targets than serum albumin, linked by polypeptide linkers and having an N-terminal amino acid sequence. The two designed ankyrin repeat domains with binding specificity for serum albumin are flanking the two other designed ankyrin repeat domains. For example a recombinant binding protein with an amino acid sequence corresponding to any of SEQ ID NOs: 95 to 107, 110, 116, 122, 129 to 131, 134 to 144, 149 to 172, and 175 to 179, in particular SEQ ID NO: 134 corresponds to this illustration. (h) Illustration of a recombinant binding protein provided herein comprising two designed ankyrin repeat domains with binding specificity for serum albumin, and two designed ankyrin repeat domains with binding specificities for other targets than serum albumin, linked by polypeptide linkers and having an N-terminal amino acid sequence. The two designed ankyrin repeat domains with binding specificity for serum albumin are N-terminal to the two other designed ankyrin repeat domains. For example a recombinant binding protein with an amino acid sequence corresponding to any of SEQ ID NOs: 112, 119, 124, 128, 132, and 133 corresponds to this illustration. (i) Illustration of a pharmaceutical compound comprising two designed ankyrin repeat domains with binding specificity for serum

albumin and a bioactive compound. The bioactive compound can be covalently linked to the two designed ankyrin repeat domains with binding specificity for serum albumin by means of chemical coupling or, in the case of polypeptides, protein fusion.

5 Figure 2. Improved storage stability of recombinant binding proteins comprising SEQ ID NO: 50. SDS 15% PAGE analysis of Proteins #49 and #50 (corresponding to SEQ ID NOs: 49 and 50, respectively, additionally having SEQ ID NO: 1 at the N terminus; prepared as described in Example 4) stored at 10 mg/ml in PBS for 1 week at 4°C (1), 25°C (2), 40°C (3), and 60°C (4), respectively. M: Marker (lower band: 6.5 kDa; band at  
10 Protein #50 level: 14.4 kDa; upper band in case of Protein #50 PAGE: 21.5 kDa).

Figure 3. Mouse pharmacokinetic studies illustrating the benefit of having two designed ankyrin repeat domains with binding specificity for serum albumin in a recombinant binding protein. Mouse pharmacokinetic studies were performed using  $m^{99}Tc$  labeled proteins as  
15 described in Example 5. The percentage injected dose (% ID), referenced to an early measurement time point (a: 4h; b-d:1h) is shown over time (t; hours). Proteins used comprised an N-terminal His-tag (SEQ ID NO: 1) in addition to the sequence indicated unless stated otherwise. (a) Pharmacokinetic profile comparison of Protein #57 (single  
20 designed ankyrin repeat domain with binding specificity for serum albumin; SEQ ID NO: 57, which comprises SEQ ID NO: 51; filled circles) with Proteins #62 and #63 (proteins comprising two designed ankyrin repeat domains with binding specificity for serum albumin (twice SEQ ID NO: 51), linked by GS- (SEQ ID NO: 63; filled diamonds) or PT-rich (SEQ ID NO: 62; filled squares) polypeptide linkers). Having two designed ankyrin  
25 repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+57% GS; +59% PT), 48 h (+76% GS; +82% PT) or 72 h (+79% GS; +94% PT) post-injection, and leads to an improved terminal half-life (+38% GS; +48% PT) compared to the protein comprising only a single designed ankyrin repeat domain with binding  
30 specificity for serum albumin. (b) Pharmacokinetic profile comparison of Protein #64 (filled circles), comprising SEQ ID NOs: 22 (designed ankyrin repeat domain with binding specificity for another target than serum albumin) and 51 (designed ankyrin repeat domain with binding specificity for serum albumin), with Proteins #73 (filled squares) and #74 (filled diamonds), comprising each SEQ ID NOs: 22 and two times 51. Protein #73 has  
35 SEQ ID NOs: 51 flanking SEQ ID NO: 22, and Protein #74 has twice SEQ ID NOs: 51 N-terminal of SEQ ID NO: 22. Having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+62% N-terminal; +89%

flanking), or 48 h (+136% N-terminal; +175% flanking) post-injection, and leads to an improved terminal half-life (>63% for both N-terminal or flanking) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin. (c) Pharmacokinetic profile comparison of Protein #82 (filled circles), comprising  
5 SEQ ID NOs: 11 (twice; designed ankyrin repeat domain with no known binding specificity) and 51 (designed ankyrin repeat domain with binding specificity for serum albumin), with Protein #109 (filled squares) comprising SEQ ID NOs: 11 (twice) and 51 (twice; N-terminal). Having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+12%), or 48 h (+35%) post-injection,  
10 and leads to an improved terminal half-life (+71%) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin. (d) Pharmacokinetic profile comparison of Protein #83 (filled circles), comprising SEQ ID NOs: 38 and 39 (designed ankyrin repeat domains each with binding specificity for another target than serum albumin) and 50 (designed ankyrin repeat domain with binding  
15 specificity for serum albumin), with Proteins #110 (filled squares), comprising each SEQ ID NOs: 38, 39 and 50 (twice; flanking SEQ ID NOs: 38 and 39). Having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+198%), 48 h (+198%), or 72 h (+228%) post-injection, and leads to an improved terminal half-life (+19%) compared to the protein comprising only a single  
20 designed ankyrin repeat domain with binding specificity for serum albumin. Note that the measurement of Protein #83 was close to the lower limit of quantification.

Figure 4. Cynomolgus monkey pharmacokinetic studies illustrating the benefit of having two designed ankyrin repeat domains with binding specificity for serum albumin in a recombinant binding protein. Cynomolgus monkey pharmacokinetic studies were  
25 performed as described in Example 6. The concentration of the respective protein is shown in nM (a), or as relative value referenced to the measurement point at 10 minutes post-injection (b), over time indicated in days (a) or hours (b). Proteins used comprised an N-terminal His-tag (SEQ ID NO: 1) in addition to the sequence indicated unless stated otherwise. (a) Pharmacokinetic profile comparison of Protein #57 (0.5 mg/Kg; 27.7 nmol/kg; single designed ankyrin repeat domain with binding specificity for serum  
30 albumin; SEQ ID NO: 57, which comprises SEQ ID NO: 51; filled circles) with Protein #62 (1.04 mg/Kg; 34.5 nmol/kg; protein comprising two designed ankyrin repeat domains with binding specificity for serum albumin (twice SEQ ID NO: 51), linked by a PT-rich  
35 polypeptide linker). Having two designed ankyrin repeat domains with binding specificity

for serum albumin leads to higher exposure (2138 d\*nmol/L vs. 4676 d\*nmol/L, i.e. +119% calculated up to day 7), leads to a reduced clearance (0.0108 L/(d\*kg) vs. 0.0031 L/(d\*kg); i.e. -71%), and leads to an improved terminal half-life (4.57 d vs. 9.00 d, i.e. +97% calculated from day 1 to day 7) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin. (b) Pharmacokinetic profile of Protein #97 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 97; filled squares) and Protein #134 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134, with no additional sequence tag; filled circles) administered at 1 mg/kg i.v. to cynomolgus monkeys are shown. Protein #134 has an improved pharmacokinetic profile compared to Protein #97.

Figure 5. Size exclusion chromatography coupled to static light scattering of a recombinant binding protein (Protein #134) comprising two designed ankyrin repeat domains with binding specificity for serum albumin. The experiment was performed as described in Example 7 using Protein #134 (a recombinant binding protein consisting of SEQ ID NO: 134; solid line), human serum albumin (dotted line), and a mixture of the two (dashed line). The experiment indicates that Protein #134 comprising two designed ankyrin repeat domains (twice SEQ ID NO: 50) with binding specificity for serum albumin is able to bind two human serum albumin molecules simultaneously.

Figure 6. ELISA analysis of recombinant binding proteins. Analysis of Protein #134 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134) in various binding ELISA, as outlined in Example 10. Protein #134 comprises one designed ankyrin repeat domain with binding specificity for VEGF-A, one designed ankyrin repeat domain with binding specificity for HGF, and two designed ankyrin repeat domains with binding specificity for serum albumin, and, correspondingly, interaction of Protein #134 with these target proteins is expected. (a) VEGF-A ELISA. The binding signal of various concentrations of Protein #134 to immobilized VEGF-A of human (filled circles), rat (filled squares), and mouse (filled rhombus), as well as human VEGF-C (open inverse triangles), and human PDGF-AB (open circles), and the corresponding fitting inhibition curves are shown. Protein #134 binds VEGF-A of these species with high affinity and is not binding VEGF-C and PDGF-AB. (b) HGF ELISA. The binding signal of various concentrations of Protein #134 to immobilized HGF of human (filled circles), cynomolgus monkey (filled triangles), and mouse (filled rhombus), and the corresponding fitting inhibition curves are shown. Protein #134 binds HGF of these species with high affinity. (c) Serum albumin

ELISA. The binding signal of various concentrations of Protein #134 to immobilized serum albumin of human (filled circles), cynomolgus monkey (open inverse triangle), rat (open triangle), mouse (open squares), and dog (open rhombus) and the corresponding fitting inhibition curves are shown. Protein #134 binds serum albumin of these species with high affinity. OD, optical density at 450 nm minus OD at 620 nm; c [pM], concentration of recombinant binding protein in pM in logarithmic scale.

Figure 7. VEGF-A/VEGF-R2 and HGF/cMet receptor competition assays Analysis of Protein #134 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134) in various competition assays as described in Example 11. (a) VEGF-A/VEGFR-2 HTRF binding competition assay. Protein #134 inhibits the VEGF-A/VEGFR-2 interaction. Baseline is indicated by the dashed line, no competition signal is indicated by the circular symbol. R: ratio 665 nm signal to 620 nm signal, c: concentration of Protein #134 in nM. (b) HGF/cMet competition binding assay. Protein #134 inhibits the HGF/cMet interaction. OD: OD at 450 nm minus OD at 620 nm, c: concentration of Protein #134 in nM. (c) VEGF-A competition binding ELISA. Protein #134 binds VEGF-A with an  $IC_{50}$  of better than 10 pM. OD: OD at 450 nm minus OD at 620 nm, c: concentration of Protein #134 in pM.

Figure 8. SPR analysis of recombinant binding proteins. Analysis of the binding of VEGF-A, HGF, and HSA by Protein #134 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134) using a ProteOn instrument as described in Example 12. Human HGF is immobilized on the biosensor chip, and Protein #134, human VEGF-A, or human serum albumin were injected according to the following injection schemes: (1) Protein #134 – hVEGF-A – HSA, (2) Protein #134 – hVEGF-A – PBST, (3; dotted line) Protein #134 – PBST – HSA, (4) PBST – PBST – PBST, (5) PBST – hVEGF-A – PBST, (6) PBST – PBST – HSA. Curves 1 and 2 indicate that Protein #134 can bind human HGF and human VEGF-A simultaneously. Furthermore, since VEGF-A binding reaches saturation in curve 1, curve 1 indicates that Protein #134 can bind human HGF, human VEGF-A and human serum albumin simultaneously. The control injections indicate that no unspecific interaction occurs. RU: resonance units; t: time in seconds.

Figure 9. Effect of recombinant binding proteins cell proliferation and cell migration. The effect of Protein #134 (a recombinant binding protein consisting of the amino acid

sequence of SEQ ID NO: 134) in different cellular assays was assessed as described in Example 13. (a) Inhibition of proliferation of HUVECs by Protein #134. HUVECs ( $3 \times 10^3$  cells/well) were stimulated by 8 ng/mL human VEGF-A. Proliferative status of HUVECs was analyzed in the absence (open circle) or presence of increasing concentrations of Protein #134 (filled circles). After 3 days of cultivation at 37°C and 5% CO<sub>2</sub>, inhibition was quantified by addition of BrdU for the last 24h of incubation. Protein #134 exhibits an IC<sub>50</sub> in the range of 100 - 150 pM. Error bars reflect standard deviation of independent duplicates. OD: OD at 450 nm minus OD at 620 nm, c: concentration of Protein #134 in ng/ml. The X axis is shown in logarithmic scale. (b) Effect of Protein #134 in an Oris cell migration assay with A549 cells. Protein #134 significantly inhibits the HGF-induced cell migration. Cells were seeded 24 h prior to stimulation with HGF (500 pM; H and D) or PBS (N), in the presence (D) and absence (H, N) of Protein #134 (5 μM). The stoppers were removed and migration was detected and quantified 48 h later after staining of cells with Calcein. Images were taken and the uncovered area in the cell culture plate well quantified. U: Uncovered area in 8 independent wells in μm<sup>2</sup>, H: HGF, no Protein #134, N: No HGF, no Protein #134, D: HGF & Protein #134. (c) Inhibition of cMet phosphorylation in A549 cells by Protein #134. A549 cells were starved overnight and stimulated with 1 nM human HGF (no HGF for negative control) in presence of PBS or increasing concentrations of Protein #134 for 10 minutes. P-cMet was detected in cell lysates by ELISA measuring OD450-620. Relative signals (% phosphorylation) were calculated using maximal signal (HGF, no Protein #134) and minimal signal (no HGF, no Protein #134). Protein #134 inhibits cMet phosphorylation with an IC<sub>50</sub> of better than 1 nM. %P: % phosphorylation, c: concentration of Protein #134 in nM.

Figure 10. Effect of recombinant binding proteins on tumor growth in vivo. The efficacy of Protein #134 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134; comprises (i) one designed ankyrin repeat domain with binding specificity for VEGF-A, (ii) one designed ankyrin repeat domain with binding specificity for HGF, and (iii) two designed ankyrin repeat domains with binding specificity for serum albumin; see Figure 1) and of other recombinant binding proteins was assessed in tumor xenograft mouse models as described in Example 14. (a) Quantification of proliferative cells and mean vascular area in tumor tissue of a U87M mouse model treated with Protein #134, Protein #60 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 60 and additionally having SEQ ID NO: 1 at the N-terminus; comprises one designed ankyrin repeat domain with binding specificity for HGF (identical to the one

in Protein #134), and one designed ankyrin repeat domain with binding specificity for serum albumin (identical to the one in Protein #134)), or Protein #61 (a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 61 and additionally having SEQ ID NO: 1 at the N-terminus; comprises one designed ankyrin repeat domain with binding specificity for VEGF-A (identical to the one in Protein #134), and one designed ankyrin repeat domain with binding specificity for serum albumin (identical to the one in Protein #134)), as described in Example 14. Regarding inhibition of the proliferation of U87M tumor xenograft cells (P; measured as percent proliferative cells, %pc), Protein #60 exhibits a slight inhibition, similar as Protein #61, whereas Protein #134 has a significantly stronger effect. Likewise, regarding inhibition of vascular growth (A; measured as mean vascular area percentage; %mva), Protein #60 exhibits a slight inhibition, Protein #61 exhibits an intermediate inhibition, and Protein #134 exhibits the strongest effect. PBS (white bars), Protein #60 (horizontally striped bars), Protein #61 (vertically striped bars), Protein #134 (black bars). (b) Effect of Protein #134 on tumor growth in a patient derived renal tumor xenograft mouse model in comparison to sorafenib and PBS. Details of the model are described in Example 14. Sorafenib suppresses tumor growth as expected. Interestingly, Protein #134 suppresses tumor growth beyond the levels of sorafenib, controlling the tumor volume at its initial levels. V: tumor volume in  $\text{mm}^3$ , d: days of treatment, open circles: vehicle, closed circles: Protein #134 (4 mg/kg), open square: sorafenib (200 mg/kg). (c) Effect of Protein #134 on tumor growth in a patient derived gastric tumor xenograft mouse model in comparison to paclitaxel and a combination of Protein #134 and paclitaxel. Details of the model are described in Example 14. Paclitaxel and Protein #134 suppress tumor growth to about the same level. The combination of Paclitaxel and Protein #134 suppresses tumor growth even more, controlling the tumor volume at its initial levels. V: tumor volume in  $\text{mm}^3$ , d: days post therapy, open circles: vehicle, closed circles: Protein #134 (4 mg/kg i.v.), open square: paclitaxel (15 mg/kg i.v.), open triangles: Protein #134 (4 mg/kg) and paclitaxel (15 mg/kg).

#### DETAILED DESCRIPTION

In the context of the present invention the term "protein" refers to a polypeptide, wherein at least part of the polypeptide has, or is able to acquire a defined three-dimensional arrangement by forming secondary, tertiary, or quaternary structures within a single polypeptide chain and/or between multiple polypeptide chains. If a protein comprises two or more polypeptide chains, the individual polypeptide chains may be linked non-covalently or covalently, e.g. by a disulfide bond between two polypeptides. A part of a

protein, which individually has, or is able to acquire, a defined three-dimensional arrangement by forming secondary or tertiary structure, is termed "protein domain". Such protein domains are well known to the practitioner skilled in the art.

The term "recombinant" as used in recombinant protein, recombinant protein domain,  
5 recombinant binding protein and the like, means that said polypeptides are produced by the use of recombinant DNA technologies well known by the practitioner skilled in the relevant art. For example, a recombinant DNA molecule (e.g. produced by gene synthesis) encoding a polypeptide can be cloned into a bacterial expression plasmid (e.g. pQE30, QIAGEN), yeast expression plasmid, mammalian expression plasmid, or plant  
10 expression plasmid, or a DNA enabling in vitro expression. If, for example, such a recombinant bacterial expression plasmid is inserted into an appropriate bacteria (e.g. *Escherichia coli*), this bacteria can produce the polypeptide encoded by this recombinant DNA. The correspondingly produced polypeptide is called a recombinant polypeptide or recombinant protein.

15 In the context of the present invention, the term "binding protein" refers to a protein comprising two or more, preferably three or more, more preferably four or more binding domains. Preferably, said binding protein is a recombinant binding protein. Preferably, said binding protein comprises two or more repeat domains. More preferably, said binding protein comprises three repeat domains. More preferably, said binding protein comprises  
20 four repeat domains. Also preferably, said binding protein comprises three or more designed ankyrin repeat domains. Further preferably, said binding protein comprises four or more designed ankyrin repeat domains. More preferably, said binding protein comprises four designed ankyrin repeat domains. Optionally, said binding protein comprises one or more bioactive compound. Said binding domains of said binding protein  
25 each have a target specificity. Preferably, two or more of said binding domains of said binding protein each have a target specificity for serum albumin. Preferably, said binding protein comprises at least three binding domains binding to at least two different targets. More preferably, said binding protein comprises at least four binding domains binding to at least three different targets.

30 Furthermore, any such binding protein may comprise additional polypeptides (such as e.g. polypeptide tags, or polypeptide linkers, well known to the person skilled in the art.

The term "bioactive compound" refers to a compound that is disease modifying when applied to a mammal having said disease. A bioactive compound may have antagonistic or agonistic properties and can be a proteinaceous bioactive compound or a non-  
35 proteinaceous bioactive compound. Such proteinaceous bioactive compounds can be



covalently attached to, for example, a binding domain of the invention by the generation of genetic fusion polypeptides using standard DNA cloning technologies, followed by their standard expression and purification. Non-proteinaceous bioactive compounds can be covalently attached to, for example, a binding domain of the invention by chemical means, 5 e.g., by coupling to a cysteine thiol via a maleimide linker with a cysteine being coupled via a polypeptide linker to the N or C terminus of a binding domain as described hereinbefore. Examples of proteinaceous bioactive compounds are binding domains having a distinct target specificity (e.g. neutralizing a growth factor by binding to it), cytokines (e.g. interleukins), growth factors (e.g. human growth hormone), antibodies and 10 fragments thereof, hormones (e.g. GLP-1), or a proteinaceous drug. Examples of non-proteinaceous bioactive compounds are toxins (e.g. DM1 from ImmunoGen), small molecules targeting GPCRs, antibiotics or a non-proteinaceous drug.

The term "binding domain" means a protein domain exhibiting the same "fold" (i.e. secondary, tertiary, and/or quaternary structure) as a protein scaffold and having a 15 predetermined property, as defined below. Such a binding domain may be obtained by rational, or most commonly, combinatorial protein engineering techniques, skills which are known in the art (Binz et al., 2005, loc. cit.). For example, a binding domain having a predetermined property can be obtained by a method comprising the steps of (a) providing a diverse collection of protein domains exhibiting the same fold as a protein 20 scaffold as defined further below; and (b) screening said diverse collection and/or selecting from said diverse collection to obtain at least one protein domain having said predetermined property. The diverse collection of protein domains may be provided by several methods in accordance with the screening and/or selection system being used, and may comprise the use of methods well known to the person skilled in the art, such as 25 phage display or ribosome display. Preferably, said binding domain is a recombinant binding domain.

The term "protein scaffold" means a protein with exposed surface areas in which amino acid insertions, substitutions or deletions are highly tolerable. Examples of protein scaffolds that can be used to generate binding domains of the present invention are 30 antibodies or fragments thereof such as single-chain Fv or Fab fragments, protein A from *Staphylococcus aureus*, the bilin binding protein from *Pieris brassicae* or other lipocalins, ankyrin repeat proteins or other repeat proteins, and human fibronectin. Protein scaffolds are known to the person skilled in the art (Binz et al., 2005, loc. cit.; Binz et al., 2004, loc. cit.).

The term "target" refers to an individual molecule such as a nucleic acid molecule, a polypeptide or protein, a carbohydrate, or any other naturally occurring molecule, including any part of such individual molecule, or complexes of two or more of such molecules. A target may be a whole cell or a tissue sample, or it may be any non-natural  
5 compound. Preferably, a target is a naturally occurring or non-natural polypeptide or a polypeptide containing chemical modifications, for example modified by natural or non-natural phosphorylation, acetylation, or methylation. In the particular application of the present invention, the targets are serum albumin, HGF and VEGF-A.

The term "predetermined property" refers to a property such as binding to a target,  
10 blocking of a target, activation of a target-mediated reaction, enzymatic activity, and related further properties. Depending on the type of desired property, one of ordinary skill will be able to identify format and necessary steps for performing screening and/or selection of a binding domain with the desired property. Preferably, said predetermined property is specifically binding to a target.

15 In the context of the present invention, the term "polypeptide" relates to a molecule consisting of a chain of multiple, i.e. two or more, amino acids linked via peptide bonds. Preferably, a polypeptide consists of more than eight amino acids linked via peptide bonds. The term "polypeptide" also includes multiple chains of amino acids, linked together by S-S bridges of cysteines. Polypeptides are well-known to the person skilled in  
20 the art.

The term "polypeptide tag" refers to an amino acid sequence attached to a polypeptide/protein, wherein said amino acid sequence is useful for the purification, detection, or "targeting" (i.e. localization to the site of a target) of said polypeptide/protein, or wherein said amino acid sequence improves the physicochemical behavior of the  
25 polypeptide/protein, or wherein said amino acid sequence possesses an effector function. The individual polypeptide tags of a binding protein may be connected to other parts of the binding protein directly or via polypeptide linkers. These polypeptide tags are all well known in the art and are fully available to the person skilled in the art. Examples of polypeptide tags are small polypeptide sequences, for example, His (e.g. the His-tag of  
30 SEQ ID NO: 1), myc, FLAG, or Strep-tags, or polypeptides such as enzymes (for example alkaline phosphatase), which allow the detection of said polypeptide/protein, or polypeptides which can be used for targeting (such as immunoglobulins or fragments thereof) and/or as effector molecules.

The term "polypeptide linker" refers to an amino acid sequence, which is able to link, for  
35 example, two protein domains, a polypeptide tag and a protein domain, a protein domain

and a non-proteinaceous compound or polymer such as polyethylene glycol, or two sequence tags. Such additional domains, tags, non-proteinaceous compounds or polymers and linkers are known to the person skilled in the relevant art. A list of examples is provided in the description of patent application WO 2002/020565. Particular examples  
5 of such linkers are glycine-serine-linkers and proline-threonine-linkers of variable lengths; preferably, said linkers have a length between 2 and 30 amino acids; more preferably, said linkers have a length between 2 and 24 amino acids. Examples of glycine-serine-linkers are GS and amino acid sequences provided in SEQ ID NOs: 2 to 6, and examples of proline-threonine-linkers are provided in amino acid sequences SEQ ID NOs: 7 to 9.

10 Patent application WO 2002/020565 and Forrer et al., 2003 (loc. cit.), contain a general description of repeat protein features and repeat domain features, techniques and applications. The term "repeat protein" refers to a protein comprising one or more repeat domains. Preferably, a repeat protein comprises up to six repeat domains. More preferably, a repeat protein comprises up to five repeat domains. More preferably, a  
15 repeat protein comprises up to four repeat domains. Furthermore, said repeat protein may comprise additional non-repeat protein domains, polypeptide tags and/or polypeptide linkers. The repeat domains can be binding domains as described hereinbefore.

The term "repeat domain" refers to a protein domain comprising two or more consecutive repeat modules as structural units, wherein said structural units have the same fold, and  
20 stack tightly to create a superhelical structure having a joint hydrophobic core. Next to a structural homology, such repeat modules further have a sequence homology. Preferably, a repeat domain further comprises an N-terminal and/or a C-terminal capping repeat. For clarity, a capping repeat can be a repeat module. Such repeat domains, repeat modules, and capping repeats, sequence motives, as well as structural homology and sequence  
25 homology are well known to the practitioner in the art from examples of designed ankyrin repeat domains (WO 2002/020565), leucine-rich repeat domains (WO 2002/020565), tetratricopeptide repeat domains (Main, E.R., Xiong, Y., Cocco, M.J., D'Andrea, L., Regan, L., Structure 11(5), 497-508, 2003), and armadillo repeat domains (WO 2009/040338). It is further well known to the practitioner in the art, that such repeat domains are different  
30 from proteins comprising repeated amino acid sequences, where every repeated amino acid sequence is able to form an individual domain (for example FN3 domains of Fibronectin), or where the repeated amino acid sequences are no structural units, i.e. said repeated amino acid sequences do not stack tightly to create a superhelical structure having a joint hydrophobic core. Methods for identifying and determining repeat modules  
35 or repeat sequence motifs or for identifying families of related proteins comprising such

repeat units or motifs, such as homology searches (BLAST etc.), are well established in the field of bioinformatics, and are well known to the practitioner in the art.

The term "designed repeat protein" and "designed repeat domain" refer to a repeat protein or repeat domain, respectively, obtained as the result of an inventive procedure, e.g. as explained in patent application WO 2002/020565. The term "designed" refers to the property that such repeat proteins and repeat domains, respectively, are man-made, synthetic and not from nature. The designed repeat proteins or designed repeat domains of WO 2002/020565 include designed ankyrin repeat proteins or designed ankyrin repeat domains, respectively. Accordingly, a designed ankyrin repeat protein herein corresponds to protein of the invention comprising at least one designed ankyrin repeat domain. Further, the term "not from nature" means that the sequence of said binding protein or said binding domain is not present as a non-artificial sequence entry in a sequence database, for example in GenBank, EMBL-Bank or Swiss-Prot. These databases and other similar sequence databases are well known to the person skilled in the art. The recombinant binding proteins or designed ankyrin repeat domains of the invention are non-naturally occurring.

The terms "repeat module", "repeat unit", "capping repeat", "capping module", and further terms relating to repeat proteins and repeat domains, are defined in WO 2002/020565, and the definitions are incorporated by reference.

The term "has binding specificity for a target", "specifically binding to a target", "binding to a target with high specificity", "specific for a target" or "target specificity" and the like means that a binding protein or binding domain binds in PBS to a target with a lower dissociation constant (i.e. it binds with higher affinity) than it binds to an unrelated protein such as the *E. coli* maltose binding protein (MBP). Preferably, the dissociation constant ("Kd") in PBS for the target is at least  $10^2$ ; more preferably, at least  $10^3$ ; more preferably, at least  $10^4$ ; or more preferably, at least  $10^5$  times lower than the corresponding dissociation constant for MBP.

Methods to determine dissociation constants of protein-protein interactions, such as surface plasmon resonance (SPR) based technologies (e.g. SPR equilibrium analysis) or isothermal titration calorimetry (ITC) are well known to the person skilled in the art. The measured Kd values of a particular protein-protein interaction can vary if measured under different conditions (e.g., salt concentration, pH). Thus, measurements of Kd values are preferably made with standardized solutions of protein and a standardized buffer, such as PBS.

The term "PBS" means a phosphate buffered water solution containing 137 mM NaCl, 10 mM phosphate and 2.7 mM KCl and having a pH of 7.4.

The term "inhibits the binding" in the context of the binding domains of the present invention refers to the ability of said binding domains to prevent the binding of its target to another protein, typically a natural ligand of the target or another antagonist. The strength of inhibition is typically measured by assessing the concentration of half-maximal inhibition (IC<sub>50</sub>). The term inhibition and the assessment of IC<sub>50</sub> values are well established in the field. For example, the designed ankyrin repeat domain of SEQ ID NO: 18 inhibits the binding of VEGF-A to its natural ligand VEGFR-2.

5 The invention relates to designed ankyrin repeat domains with binding specificity for serum albumin, and to recombinant binding proteins comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, and to recombinant binding proteins comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding  
15 specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin.

In one embodiment, the invention relates to designed ankyrin repeat domains with binding specificity for serum albumin. Examples of designed ankyrin repeat domains with binding  
20 specificity for serum albumin are given in SEQ ID NOs: 40 to 56 (see also Examples) and further examples are described in WO 2012/069654. In particular, the invention relates to designed ankyrin repeat domains with binding specificity for serum albumin selected from the group of SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50, in which up to 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0 amino acids are  
25 exchanged by any amino acid. In one embodiment, the invention relates to designed ankyrin repeat domains with binding specificity for serum albumin that have 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with a designed ankyrin repeat domain selected from the group of SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50. In one embodiment, the  
30 invention relates to designed ankyrin repeat domains with binding specificity for serum comprising an amino acid sequence selected from the group of SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50. In one embodiment, the invention relates to designed ankyrin repeat domains with binding specificity for serum selected that consist of an amino acid sequence selected from the  
35 group of SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably

SEQ ID NO: 50. In one embodiment, the invention relates to a designed ankyrin repeat domain with binding specificity for serum albumin comprising the amino acid sequence of SEQ ID NO: 50. The preferred designed ankyrin repeat domain with binding specificity for serum albumin of the invention is SEQ ID NO: 50. Preferably, said designed ankyrin repeat domain with binding specificity for serum albumin binds serum albumin of mouse, rat, dog, cynomolgus monkey, or human origin, more preferably serum albumin of mouse, cynomolgus monkey or human origin, more preferably serum albumin of cynomolgus monkey or human origin, more preferably serum albumin of human origin, in PBS with a dissociation constant ( $K_d$ ) below  $10^{-5}M$ ; preferably below  $10^{-6}M$ ; or more preferably below  $10^{-7}M$ . The term "mouse serum albumin" refers to UniProt accession number P07724, the term "cynomolgus monkey serum albumin" (i.e. *macaca fascicularis*) refers to UniProt accession number A2V9Z4, and the term "human serum albumin" refers to UniProt accession number P02768. In one embodiment, the invention relates to a designed ankyrin repeat domain with binding specificity for serum albumin comprising, more preferably consisting of, an amino acid sequence selected from the group of SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50, which exhibit improved storage stabilities compared to SEQ ID NO: 51. "Improved storage stability" in the context of the present invention means an improved midpoint of denaturation temperature (i.e. midpoint of the cooperative unfolding upon temperature increase) by  $0.5^{\circ}C$ ,  $1^{\circ}C$ ,  $1.5^{\circ}C$ ,  $2^{\circ}C$ ,  $2.5^{\circ}C$ ,  $3^{\circ}C$ ,  $3.5^{\circ}C$ , or  $4^{\circ}C$ , and/or the reduction of the amounts of a degradation band, preferably the reduction of the amount of degradation products, as detected by a Coomassie-stained SDS-PAGE occurring after storage at  $40^{\circ}C$  for 1 month at 10 mg/ml in PBS, by at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%. Methods to assess storage stability by SDS-PAGE and methods to determine the midpoint of denaturation by using fluorimetric methods or circular dichroism are well known to the person skilled in the art. In one embodiment, the invention relates to a designed ankyrin repeat domain with binding specificity for serum albumin comprising, more preferably consisting of amino acid sequence SEQ ID NO: 50, which exhibits improved storage stability compared to SEQ ID NO: 49, preferably which exhibits reduced amounts of degradation products, as detected by SDS-PAGE, occurring after storage at  $40^{\circ}C$  for 1 month at 10 mg/ml in PBS, by at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%, compared to SEQ ID NO: 49. Examples of designed ankyrin repeat domains and recombinant binding proteins with improved storage stability properties are given in Example 9.

In one embodiment, the invention relates to a designed ankyrin repeat domain with binding specificity for serum albumin selected from the group consisting of amino acid

sequences SEQ ID NOs: 44 to 49, 51 and 52, more preferably 48, 49, 51, and 52, more preferably 48 and 49, more preferably 49, comprising Glutamate at position 78. In one embodiment, the invention relates to SEQ ID NO: 49 wherein Aspartate at position 78 has been exchanged by Glutamate, corresponding to SEQ ID NO: 50. SEQ ID NO: 49  
5 comprises a high number of potential degradation sites. Degradation may for example occur in the vicinity of any one of the 5 asparagines (including asparagine-glycine dipeptides), 13 aspartates, or 10 glycines of SEQ ID NO: 49, amongst additional potential degradation sites. SEQ ID NO: 49 further comprises a number of potential oxidation sites. Surprisingly, a major effect on storage stability can be achieved by mutating only position  
10 78 of SEQ ID NO: 49. Furthermore, the functionality of the designed ankyrin repeat with binding specificity for serum albumin can be preserved by mutating position 78 of SEQ ID NO: 49 from aspartate to glutamine. The designed ankyrin repeat domain consisting of SEQ ID NO: 49 comprising Glutamate in position 78 exhibits higher storage stability compared to the designed ankyrin repeat domain comprising Aspartate in that position.

15 In one embodiment, the present invention relates to a recombinant binding protein comprising at least two, preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin. The preferred recombinant binding protein of the invention comprises two designed ankyrin repeat domains with binding specificity for serum albumin. Examples of such recombinant binding proteins are given in the amino  
20 acid sequences SEQ ID NOs: 62, 63, 73 to 81, and 95 to 179.

In one embodiment, the present invention relates to a recombinant binding protein comprising at least two, more preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin, wherein said recombinant binding protein exhibits improved pharmacokinetic properties compared to the recombinant  
25 binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. The examples of the present invention disclose such recombinant binding proteins.

The expression "the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin", means a recombinant binding  
30 that has the composition of a recombinant binding protein of the present invention in which the number of designed ankyrin repeat domains with binding specificity for serum albumin is reduced to one, by removing all designed ankyrin repeat domains with binding specificity for serum albumin but one, and the corresponding polypeptide linkers. Preferably, said remaining one designed ankyrin repeat domain with binding specificity for  
35 serum albumin is located at a position in the recombinant binding protein corresponding to

a position that was comprising a designed ankyrin repeat domain with binding specificity for serum albumin in the recombinant binding protein of the present invention, and the remaining one designed ankyrin repeat domain with binding specificity for serum albumin is identical to the designed ankyrin repeat domain with binding specificity for serum albumin that was at the corresponding position in the recombinant binding protein of the present invention. For example, the recombinant binding protein consisting of SEQ ID NO: 85 is the recombinant binding protein consisting of SEQ ID NO: 95, in which the C-terminal designed ankyrin repeat domain with binding specificity for serum albumin (in this case SEQ ID NO: 50) as well as the adjacent polypeptide linker (in this case SEQ ID NO: 9) have been removed. Importantly, the remaining designed ankyrin repeat domain with binding specificity for serum albumin (SEQ ID NO: 50) of SEQ ID NO: 85 is N-terminal, and SEQ ID NO: 95 comprises the same SEQ ID NO: 50 at the same position. Likewise, the recombinant binding protein consisting of SEQ ID NO: 83 is the recombinant binding protein consisting of SEQ ID NO: 110, in which the C-terminal designed ankyrin repeat domain with binding specificity for serum albumin (in this case SEQ ID NO: 50) as well as the adjacent polypeptide linker (in this case SEQ ID NO: 9) have been removed.

The expression “exhibits improved pharmacokinetic properties”, “improved pharmacokinetic properties”, or “pharmacokinetic property improvement” in this invention has the meaning that a pharmacokinetic parameter of a recombinant binding protein is improved compared to the corresponding pharmacokinetic parameter of a protein it is compared with. Corresponding examples are shown in Examples 5 and 6 and Figures 3 and 4. For example, when comparing Protein #110 (a protein consisting of SEQ ID NO: 110 and additionally SEQ ID NO: 1 at the N terminus) with Protein #83 (a protein consisting of SEQ ID NO: 83 and additionally SEQ ID NO: 1 at the N terminus) in cynomolgus monkey pharmacokinetic studies Protein #110 has a higher exposure (+32%), a reduced clearance (-47%) as well as a higher terminal half-life (+168%, calculated from day 1 to day 6) as Protein #83. As another example, when comparing Protein #62 (a protein consisting of SEQ ID NO: 62 and additionally SEQ ID NO: 1 at the N terminus) with Protein #57 (a protein consisting of SEQ ID NO: 57 and additionally SEQ ID NO: 1 at the N terminus) in cynomolgus monkey pharmacokinetic studies Protein #62 has a higher exposure (+119%), a reduced clearance (-71%) as well as a higher terminal half-life (+97%, calculated from day 1 to day 7) as Protein #57. Or when comparing Protein #109 (a protein consisting of SEQ ID NO: 109 and additionally SEQ ID NO: 1 at the N terminus) with Protein #82 (a protein consisting of SEQ ID NO: 82 and additionally SEQ ID NO: 1 at the N terminus) in cynomolgus monkey pharmacokinetic studies Protein #109 has a higher exposure (+19%), a reduced clearance (-37%) as well as a higher



terminal half-life (+55%, calculated from day 1 to day 7) as Protein #82. As yet another example, when comparing Protein #97 (a protein consisting of SEQ ID NO: 97 and additionally SEQ ID NO: 1 at the N terminus) with Protein #68 (a protein consisting of SEQ ID NO: 68 and additionally SEQ ID NO: 1 at the N terminus) in cynomolgus monkey  
5 pharmacokinetic studies Protein #97 has a higher terminal half-life (+264%, calculated from day 1 to day 7) as Protein #68. Further examples are given in Examples 5 and 6 as well as Figures 3 and 4. Preferably, an improved pharmacokinetic property is a reduced clearance, and/or an increased exposure, and/or an increased terminal half-life. More preferably, an improved pharmacokinetic property is an increased terminal half-life. In one  
10 embodiment, a recombinant binding protein of the present invention, comprising at least two, more preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin exhibits an increased terminal half-life, and/or a reduced clearance, and/or an increased exposure of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, or 250% compared to  
15 the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. In one embodiment, a recombinant binding protein of the present invention, comprising at least two, more preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin exhibits an increased terminal half-life, preferably an increased terminal half-life of at least 5%, 10%,  
20 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, or 250% compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin.

Preferably, clearance, and/or exposure, and/or terminal half-life are assessed in a mammal, more preferably mouse and/or cynomolgus monkey, more preferably  
25 cynomolgus monkey. Preferably, when measuring the clearance, and/or exposure, and/or terminal half-life in mouse, the evaluation is done considering the data up to 48 h post-injection. More preferably, the evaluation of terminal half-life in mouse is calculated from 24 h to 48 h. Preferably, when measuring the clearance, and/or exposure, and/or terminal  
30 half-life in cynomolgus monkey, the evaluation is done considering the data up to day 7 post-injection. More preferably, the evaluation of terminal half-life in cynomolgus monkey is calculated from day 1 to day 7. The term "terminal half-life" of a drug such as a recombinant binding protein of the invention refers to the time required to reach half the plasma concentration of the drug applied to a mammal after reaching pseudo-equilibrium (for example calculated from 24h to 48h in mouse or calculated from day 1 to day 7 in  
35 cynomolgus monkey). Terminal half-life is not defined as the time required to eliminate half the dose of the drug administered to the mammal. The term terminal half-life is well

known to the person skilled in the art. Preferably, pharmacokinetic comparison is done at any dose, more preferably at equivalent dose (i.e. same mg/kg dose) or equimolar dose (i.e. same mol/kg dose), more preferably at equimolar dose (i.e. same mol/kg dose). It is understood by the person skilled in the art that equivalent and/or equimolar dosing in  
5 animals is subject to experimental dose variations of at least 20%, more preferably 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. Preferably, a dose used for pharmacokinetic measurement is selected from 0.001 to 1000 mg/kg, more preferably 0.01 to 100 mg/kg, more preferably 0.1 to 50 mg/kg, more preferably 0.5 to 10 mg/kg.

In one embodiment, a recombinant binding protein of the present invention comprising at  
10 least two, more preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin, exhibits a higher percentage of injected dose in mouse 24 h and/or 48 h and/or 72 h post injection, preferably 24 h post-injection, preferably 48 h post-injection, more preferably 72 h post-injection, more preferably 72 h and 48 h post-injection, more preferably 24 h, 48 h and 72 h post-injection, compared to the recombinant  
15 binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. Preferably, the percentage of injected dose in mouse is calculated by comparison to the concentration measurement 1 h or 4 h, preferably 1 post-injection. In one embodiment, the recombinant binding protein of the present invention comprising at least two, more preferably comprising two, designed ankyrin repeat  
20 domains with binding specificity for serum albumin, exhibits a higher percentage of injected dose in cynomolgus monkey 4 days and/or 5 days and/or 6 days post-injection, preferably 4 days, preferably 5 days, more preferably 6 days, more preferably 5 and 6 days post-injection, more preferably 4, 5, and 6 days post-injection, compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with  
25 binding specificity for serum albumin. Preferably, the percentage of injected dose in cynomolgus monkey is calculated by comparison to the concentration measurement 10 min or 1 h, preferably 10 min post-injection. A higher percentage of injected dose refers to an increased percentage of dose of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, or 250%.

30 In one embodiment, the recombinant binding protein of the present invention comprises at least three designed ankyrin repeat domains, wherein at least two designed ankyrin repeat domains are designed ankyrin repeat domains with binding specificity for serum albumin. Examples of such recombinant binding proteins are given in the amino acid sequences SEQ ID NOs: 73 to 81 and 95 to 179.

In one embodiment, the recombinant binding protein comprises at least four designed ankyrin repeat domains, wherein at least two designed ankyrin repeat domains are designed ankyrin repeat domains with binding specificity for serum albumin. Examples of such recombinant binding proteins are given in the amino acid sequences SEQ ID NOs:  
5 95 to 179.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein each of said designed ankyrin repeat domains with binding specificity for serum albumin in PBS has binding specificity for serum albumin of mammalian origin, more preferably mouse, rat, dog, cynomolgus monkey, or human origin, more preferably serum albumin of mouse, cynomolgus monkey or human origin, more preferably serum albumin of cynomolgus monkey or human origin, more preferably serum albumin of human origin.  
10

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein each of said designed ankyrin repeat domains with binding specificity for serum albumin binds serum albumin, more preferably serum albumin of mammalian origin, more preferably serum albumin of mouse, rat, dog, cynomolgus monkey, or human origin, more preferably serum albumin of mouse, cynomolgus monkey or human origin, more preferably serum albumin of cynomolgus monkey or human origin, preferably serum albumin of human origin, in PBS with a dissociation constant ( $K_d$ ) below  $10^{-5}M$ , preferably below  $10^{-6}M$ , more preferably below  $10^{-7}M$ . Examples of such designed ankyrin repeat domains with binding specificity for serum albumin are given in Example 2 and in SEQ ID NOs: 40 to 56.  
15  
20

In one embodiment, the present invention relates to a recombinant binding protein comprising two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said two designed ankyrin repeat domains with binding specificity for serum albumin are at any position compared to any other protein domain, preferably any other designed ankyrin repeat domain, comprised in said recombinant binding protein, preferably wherein said two designed ankyrin repeat domains with binding specificity for serum albumin are both N-terminal of any other protein domain, preferably any other designed ankyrin repeat domain, comprised in said recombinant binding protein, or wherein said two designed ankyrin repeat domains with binding specificity for serum albumin are one N-terminal and one C-terminal of any other protein domain, preferably any other designed ankyrin repeat domain, comprised in said recombinant binding protein, or, more preferably, wherein said two designed ankyrin repeat domains with binding  
25  
30  
35

specificity for serum albumin are one N-terminal and one C-terminal of any other protein domain, preferably any other designed ankyrin repeat domain, comprised in said recombinant binding protein. Preferably, said two designed ankyrin repeat domains with binding specificity for serum albumin are not both C-terminal of any other protein domain,  
5 preferably any other designed ankyrin repeat domain, comprised in said recombinant binding protein. Examples of different arrangements of designed ankyrin repeat domains within a recombinant binding protein are given in SEQ ID NOs: 95 to 179, and are described in the Examples. SEQ ID NOs: 134 illustrates the preferred arrangement of the two designed ankyrin repeat domains with binding specificity for serum albumin in a  
10 recombinant binding protein of the present invention.

In one embodiment, the invention relates to a recombinant binding protein comprising at least three, preferably comprising at least four, more preferably comprising four designed ankyrin repeat domains, wherein two of said at least three, preferably at least four, more preferably four designed ankyrin repeat domains each have binding specificity for serum  
15 albumin, and/or, preferably and wherein said at least three, preferably at least four, more preferably four designed ankyrin repeat domains are linked by polypeptide linkers.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said designed ankyrin repeat domains with binding specificity for serum albumin  
20 each have at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identity with any amino acid sequence selected from the group of SEQ ID NOs: 44 to 52, preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin,  
25 wherein said designed ankyrin repeat domains with binding specificity for serum albumin are selected from any amino acid sequence selected from the group of SEQ ID NOs: 44 to 52, preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50, and wherein in each of said two designed ankyrin repeat  
30 domains with binding specificity for serum albumin up to 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 0 amino acids are exchanged by any amino acid.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin,  
35 wherein said designed ankyrin repeat domains with binding specificity for serum albumin are selected from any amino acid sequence selected from the group of SEQ ID NOs: 44

to 52, preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50. In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said designed ankyrin repeat domains with binding specificity for serum albumin are at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical in amino acid sequence. In one embodiment, said designed ankyrin repeat domains with binding specificity for serum albumin of said recombinant binding protein are identical in amino acid sequence. For example, the two designed ankyrin repeat domains with binding specificity for serum albumin comprised in SEQ ID NO: 130 are at least 95% identical (6 residues difference on 124 amino acids). In another example, the two designed ankyrin repeat domains with binding specificity for serum albumin comprised in SEQ ID NO: 129 are at least 80% identical (24 residues difference on 124 amino acids).

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein the designed ankyrin repeat domains with binding specificity for serum albumin are able to simultaneously bind one serum albumin molecule each. Preferably, said serum albumin is of human origin. Examples for simultaneous binding of two human serum albumin molecules by recombinant binding proteins of the present invention, comprising two designed ankyrin repeat domains with binding specificity for serum albumin, are shown in Example 7.

In one embodiment, the invention relates to a recombinant binding protein comprising at least 3, 4, 5, 6, 7, 8, 9, 10 designed ankyrin repeat domains with binding specificity for serum albumin.

In one embodiment, the invention relates to a recombinant binding protein comprising at least three, preferably at least four, more preferably comprising four, designed ankyrin repeat domains, wherein at least two, more preferably two, of said designed ankyrin repeat domains are designed ankyrin repeat domains with binding specificity for serum albumin, more preferably human serum albumin, and wherein said at least two, more preferably two, designed ankyrin repeat domains with binding specificity for serum

albumin are at least, preferably are, one N-terminal and one C-terminal of any other designed ankyrin repeat domain, preferably the other two designed ankyrin repeat domains, and wherein said at least two, more preferably two, designed ankyrin repeat domains with binding specificity for serum albumin are each binding serum albumin, preferably serum albumin of human origin, in PBS with a dissociation constant ( $K_d$ ) of at least  $10^{-5}M$ , preferably below  $10^{-6}M$ , or more preferably below  $10^{-7}M$ , and wherein said recombinant binding protein exhibits an increased terminal half-life, preferably an increased terminal half-life of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50% compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin, and wherein said at least three, preferably at least four, more preferably four, designed ankyrin repeat domains are linked by polypeptide linkers. In one embodiment, said at least two, more preferably two, designed ankyrin repeat domains with binding specificity for serum albumin of said recombinant binding protein are at least 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical, more preferably identical.

In one embodiment, protein domains or the designed ankyrin repeat domains present in a recombinant binding protein of the present invention are linked by polypeptide linkers of any amino acid sequence composition. In one embodiment, the polypeptide linkers linking protein domains or the designed ankyrin repeat domains present in a recombinant binding protein of the present invention comprise amino acid sequences selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9, in which up to 4, 3, 2, 1, 0 amino acids are exchanged by any amino acid. In one embodiment, said polypeptide linkers comprise an amino acid sequence chosen from any of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9. In one embodiment, the flanking N-terminal Gly Ser of SEQ ID NOs: 7 to 9 and/or the flanking C-terminal Gly Ser of SEQ ID NOs: 2 to 9 are optionally missing. In one embodiment, SEQ ID NOs: 7 to 9 additionally comprises Arg Ser C-terminally (as e.g. present in SEQ ID NOs: 68 and 109). In one embodiment, the second-to-C-terminal amino acid glycine of said polypeptide linkers of SEQ ID NOs: 2 to 6 may be exchanged by arginine (as e.g. present in SEQ ID NOs: 70 and 88). In one embodiment, the polypeptide linkers linking the designed ankyrin repeat domains present in a recombinant binding protein of the present invention consist of an amino acid sequence selected from of any of amino acid sequences SEQ ID NOs: 2 to 9, more

preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9. In one embodiment, said polypeptide linkers present in a recombinant binding protein of the present invention are 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical, preferably identical.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said polypeptide linkers comprise amino acid sequences selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9, in which up to 4, 3, 2, 1, 0 amino acids are exchanged by any amino acid, and wherein said designed ankyrin repeat domains with binding specificity for serum albumin each have at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% identity with any amino acid sequence selected from the group of SEQ ID NOs: 44 to 52, preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NOs: 49 and 50, more preferably SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said polypeptide linkers consist of an amino acid sequences selected from the amino acid sequences SEQ ID NOs: 6 or 9, in which up to 4, 3, 2, 1, 0 amino acids are exchanged by any amino acid, and wherein said designed ankyrin repeat domains with binding specificity for serum albumin each consist of an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% identity with any amino acid sequence selected from the group of SEQ ID NOs: 48 to 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 50, and wherein said recombinant binding protein exhibits improved storage stability, preferably reduced amounts of degradation products after storage at 40°C for 1 month at 10 mg/ml in PBS,

compared to the recombinant binding protein, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 49, and/or compared to the recombinant binding protein, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 51, preferably compared to the recombinant binding protein, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 49.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence of SEQ ID NO: 9.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each consist of the amino acid sequence of SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least two designed ankyrin repeat domains with binding specificity for serum albumin, wherein said at least two designed ankyrin repeat domains with binding specificity for serum albumin each consist of the amino acid sequence of SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers each consisting of the amino acid sequence of SEQ ID NO: 9.

In one embodiment the invention relates to a recombinant binding protein comprising four designed ankyrin repeat domains, wherein two of said designed ankyrin repeat domains are designed ankyrin repeat domains with binding specificity for serum albumin, wherein said two designed ankyrin repeat domains with binding specificity for serum albumin each comprise the amino acid sequence of SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence of SEQ ID NO: 9, and wherein said designed ankyrin repeat domains are arranged (from N-terminal side to C-terminal side): SEQ ID NO: 50 – SEQ ID NO: 9 – XXX – SEQ ID NO: 9 – YYY – SEQ ID NO: 9 – SEQ ID NO: 50, wherein XXX and YYY each represent a designed ankyrin repeat domain with binding specificity for another target than serum albumin.



In one embodiment, the present invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and  
5 wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin. Preferably, said recombinant binding protein consists of a single polypeptide chain. More preferably, said first, second, third and fourth designed ankyrin repeat domain are linked by polypeptide linkers. In one embodiment, the present  
10 invention relates to a recombinant binding protein comprising a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin. Examples of  
15 such recombinant binding proteins are given in amino acid sequences SEQ ID NOs: 95 to 108 and 116 to 179.

Preferably, the designed ankyrin repeat domain with binding specificity for VEGF-A binds VEGF-A of mouse, rat, dog, rabbit, cynomolgus monkey, or human origin, more preferably VEGF-A of mouse, cynomolgus monkey or human origin, more preferably VEGF-A of  
20 cynomolgus monkey or human origin, more preferably VEGF-A of human origin. Preferably, VEGF-A is human VEGF-A165. Examples of designed ankyrin repeat domains with binding specificity to VEGF-A are given herein (SEQ ID NOs: 12 to 21; see examples) and further examples are described in WO 2010/060748 and WO 2011/135067.

25 Preferably, the designed ankyrin repeat domain with binding specificity for HGF binds HGF of mouse, rat, dog, rabbit, cynomolgus monkey, or human origin, more preferably HGF of mouse, cynomolgus monkey or human origin, more preferably HGF of cynomolgus monkey or human origin, more preferably HGF of human origin. Examples of designed ankyrin repeat domains with binding specificity to HGF are given herein (SEQ ID  
30 NOs: 23 to 37; see examples) and further examples are described in WO 2014/191574.

In one embodiment the recombinant binding protein or designed ankyrin repeat domain is devoid of a free Cys residue. A "free Cys residue" is not involved in the formation of a disulfide bond. In one embodiment, the invention relates to a binding protein or binding domain free of any Cys residue. In one embodiment, the designed ankyrin repeat domain  
35 and/or recombinant binding protein devoid of any disulfide bond. The disulfide bonds of

antibody fragments for example are known to the person skilled in the art to hamper the simple production of the antibody fragments in bacteria.

The techniques to modify a recombinant binding protein of the present invention are well known to the person skilled in the art.

5 In particular, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain binds VEGF-A in PBS with a dissociation constant (Kd) below  $10^{-7}$ M; preferably below  $10^{-8}$ M; more preferably below  $10^{-9}$ M; or more preferably below  $10^{-10}$ M; and wherein said second designed ankyrin repeat domain binds HGF in  
10 PBS with a Kd below  $10^{-7}$ M; preferably below  $10^{-8}$ M; more preferably below  $10^{-9}$ M; or more preferably below  $10^{-10}$ M; and wherein said third and fourth designed ankyrin repeat domains each bind serum albumin in PBS with a Kd below  $10^{-5}$ M; preferably below  $10^{-6}$ M; or more preferably below  $10^{-7}$ M. Examples of designed ankyrin repeat domains with binding specificity to VEGF-A, designed ankyrin repeat domains with binding specificity to  
15 HGF, and designed ankyrin repeat domains with binding specificity to serum albumin are given herein (SEQ ID NOs: 12 to 56; see examples).

Furthermore, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain inhibits the binding of human VEGF-A to human VEGFR-  
20 2 in PBS with an  $IC_{50}$  value below  $10^{-7}$ M, preferably  $10^{-8}$ M, more preferably  $10^{-9}$ M, and wherein said second designed ankyrin repeat domain inhibits the binding of human HGF to human cMet in PBS with an  $IC_{50}$  value below  $10^{-7}$ M, preferably  $10^{-8}$ M, more preferably  $10^{-9}$ M. Different examples of designed ankyrin repeat domains selected from SEQ ID NOs: 12 to 37 are given in the examples.

25 In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain comprises an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% amino acid sequence identity with one designed ankyrin repeat domain selected from the group consisting of  
30 amino acid sequences SEQ ID NOs: 12 to 21, more preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more preferably SEQ ID NO: 18. In one embodiment, said first designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21, more preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more  
35 preferably SEQ ID NO: 18, and amino acid sequences in which up to 10, 9, 8, 7, 6, 5, 4, 3,

2, 1, 0 amino acids of SEQ ID NOs: 12 to 21, more preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more preferably SEQ ID NO: 18, are exchanged by any amino acid. In one embodiment, said first designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 14 to 21, in which individual amino acids are replaced by any amino acid occurring at the same position of an alignment of the amino acid sequences of SEQ ID NOs: 14 to 21. In one embodiment, said first designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21, more preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more preferably SEQ ID NO: 18. In one embodiment, said first designed ankyrin repeat domain consists of an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21, more preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more preferably SEQ ID NO: 18. Furthermore, said second designed ankyrin repeat domain of said recombinant binding protein preferably comprises an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% amino acid sequence identity with one designed ankyrin repeat domain selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26. In one embodiment, said second designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26, and amino acid sequences in which up to 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0 amino acids of SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26, are exchanged by any amino acid. In one embodiment, said second designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 27, in which individual amino acids are replaced by any amino acid occurring at the same position of an alignment of the amino acid sequences of SEQ ID NOs: 23 to 27. In one embodiment, said second designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26. In one embodiment, said second designed ankyrin repeat domain consists of an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26. In one embodiment, said second designed ankyrin repeat domain consists of an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26.

SEQ ID NO: 26. In one embodiment, said third and fourth designed ankyrin repeat domains of said recombinant binding protein each comprise an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 100% amino acid sequence identity with one ankyrin repeat domain selected from the group consisting of amino acid sequences SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50. In one embodiment, said third and fourth designed ankyrin repeat domains each comprise an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50, and amino acid sequences in which up to 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0 amino acids of SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50, are exchanged by any other amino acid. In one embodiment, said third and fourth designed ankyrin repeat domains each comprise an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 42 to 51, in which individual amino acids are replaced by any amino acid occurring at the same position of an alignment of the amino acid sequences of SEQ ID NOs: 42 to 51. In one embodiment, said third and fourth designed ankyrin repeat domains each comprise an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50. In one embodiment, said third and fourth designed ankyrin repeat domains each consist of an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50. In one embodiment, said third and fourth designed ankyrin repeat domains are identical. Further in this embodiment, said designed ankyrin repeat domains are linked by polypeptide linkers selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9, and amino acid sequences in which up to 4, 3, 2, 1, 0 amino acids of SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9, are exchanged by any amino acid. In one embodiment, said polypeptide linkers comprise an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9. In one embodiment, the flanking N-terminal Gly Ser of SEQ ID NOs: 7 to 9 and/or the flanking C-terminal Gly

Ser of SEQ ID NOs: 2 to 9 are optionally missing. In one embodiment, SEQ ID NOs: 7 to 9 additionally comprises Arg Ser C-terminally (as e.g. present in SEQ ID NOs: 97 and 98). In one embodiment, the second-to-C-terminal amino acid glycine of said polypeptide linkers of SEQ ID NOs: 2 to 6 may be exchanged by arginine (as e.g. present in SEQ ID NOs: 99 and 100). In one embodiment, the polypeptide linkers linking the designed ankyrin repeat domains present in a recombinant binding protein of the present invention consist of an amino acid sequence selected from any of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9. In one embodiment, said polypeptide linkers present in a recombinant binding protein of the present invention are identical. Examples of such polypeptide linkers, variations thereof, and the use of such polypeptide linkers in recombinant binding proteins are given in the examples.

In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21, preferably SEQ ID NOs: 17 to 21, more preferably SEQ ID NOs: 18 to 20, more preferably SEQ ID NO: 18, and wherein said second designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37, more preferably SEQ ID NOs: 23 to 27, more preferably SEQ ID NOs: 25 to 27, more preferably SEQ ID NO: 26, and wherein said third and fourth designed ankyrin repeat domains each comprise an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 40 to 56, preferably SEQ ID NOs: 48 to 52, more preferably SEQ ID NOs: 48 to 50, more preferably SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9, more preferably SEQ ID NOs: 3 to 9, more preferably SEQ ID NOs: 4 to 9, more preferably SEQ ID NOs: 6 or 9, more preferably SEQ ID NO: 9.

In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers, and wherein said recombinant binding protein can bind VEGF-A and HGF, more

preferably VEGF-A, HGF, and serum albumin, more preferably human VEGF-A, human HGF and human serum albumin, simultaneously. In one embodiment, said recombinant binding protein can bind two serum albumin molecules, more preferably two human serum albumin molecules, simultaneously.

- 5 The terms “first”, “second”, “third”, and optionally “fourth”, used in “first designed ankyrin repeat domain”, “second designed ankyrin repeat domain”, “third designed ankyrin repeat domain”, and “fourth designed ankyrin repeat domain”, do not indicate or imply any positional arrangement of said designed ankyrin repeat domains within the recombinant binding protein.
- 10 In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain which are linked by polypeptide linkers. In one embodiment, said first designed ankyrin repeat domain is N-terminal of the C-terminal designed ankyrin repeat domain and C-terminal of the other two designed ankyrin repeat domains. In one embodiment, said second
- 15 designed ankyrin repeat domain, having a binding specificity for HGF, is C-terminal of the N-terminal designed ankyrin repeat domain and N-terminal of the other two designed ankyrin repeat domains. In one embodiment, said third and fourth designed ankyrin repeat domains, each having a binding specificity for serum albumin, are one N-terminal and one C-terminal of the other two designed ankyrin repeat domains, or they are N-terminal of the
- 20 other two designed ankyrin repeat domains, more preferably said third and fourth designed ankyrin repeat domains are one N-terminal and one C-terminal of the other two designed ankyrin repeat domains. In one embodiment, said third designed ankyrin repeat domain is N-terminal of the other three designed ankyrin repeat domains, said fourth designed ankyrin repeat domain is C-terminal of the other three designed ankyrin repeat
- 25 domains, said second designed ankyrin repeat domain is C-terminal of said third designed ankyrin repeat domain and N-terminal of said first designed ankyrin repeat domain, and said first designed ankyrin repeat domain is C-terminal of said second designed ankyrin repeat domain and N-terminal of said fourth designed ankyrin repeat domain.
- In one embodiment, the invention relates to a recombinant binding protein comprising at
- 30 least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin, and wherein said designed ankyrin repeat domains are linked by polypeptide
- 35 linkers. In one embodiment, said first, second, third and fourth designed ankyrin repeat

domains are in the order (from N terminus to C terminus) third-second-first-fourth, third-fourth-second-first, fourth-second-first-third, or fourth-third-second-first, even more preferably third-second-first-fourth, or fourth-second-first-third, more preferably third-second-first-fourth.

5 In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum  
10 albumin, wherein said recombinant binding protein binds VEGF-A, preferably human VEGF-A, with an  $EC_{50}$  of less than  $10^{-7}M$ , preferably less than  $10^{-8} M$ , more preferably less than  $10^{-9}M$ , more preferably less than  $10^{-10}M$ .

In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said  
15 third and fourth designed ankyrin repeat domains each comprise the amino acid sequence of SEQ ID NO: 50. In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain comprises the amino acid sequence of SEQ ID NO: 18, and wherein said second designed ankyrin repeat domain  
20 comprises the amino acid sequence of SEQ ID NO: 26, and wherein said third and fourth designed ankyrin repeat domains each comprise the amino acid sequence of SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence of SEQ ID NO: 9.

In one embodiment the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said  
25 first designed ankyrin repeat domain comprises the amino acid sequence of SEQ ID NO: 18, and wherein said second designed ankyrin repeat domain comprises the amino acid sequence of SEQ ID NO: 26, and wherein said third and fourth designed ankyrin repeat domains each comprise the amino acid sequence of SEQ ID NO: 50, and wherein said  
30 designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence of SEQ ID NO: 9, and wherein said designed ankyrin repeat domains are arranged (from N-terminal side to C-terminal side): SEQ ID NO: 50 – SEQ ID NO: 9 – SEQ ID NO: 26 – SEQ ID NO: 9 – SEQ ID NO: 18 – SEQ ID NO: 9 – SEQ ID NO: 50.

In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin, and wherein said designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence of SEQ ID NO: 9, and wherein said recombinant binding protein binds VEGF-A, and/or HGF, preferably VEGF-A with a lower, i.e. better,  $EC_{50}$  compared to the recombinant binding protein wherein said designed ankyrin repeat domains are linked by polypeptide linkers each comprising the amino acid sequence SEQ ID NO: 6. Examples of the influence of the linker on  $EC_{50}$  are given in Example 8 and the term "lower  $EC_{50}$ " is well known to the person skilled in the art. Preferably, the term "lower  $EC_{50}$ " means an  $EC_{50}$  value which is improved by a factor of 1.1, more preferably 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0.

In one embodiment the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin, wherein said recombinant binding protein binds VEGF-A, and/or HGF, preferably VEGF-A with a lower, i.e. better,  $EC_{50}$  compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. Examples are given in Example 8.

In one embodiment, the invention relates to a recombinant binding protein comprising an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% amino acid sequence identity with an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 134 to 179, preferably SEQ ID NOs: 134 to 158, more preferably SEQ ID NOs: 134 to 149, more preferably SEQ ID NOs: 134 to 140, more preferably SEQ ID NO: 134.

In one embodiment, the invention relates to a recombinant binding protein comprising an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 134 to 179, preferably SEQ ID NOs: 134 to 158, more preferably SEQ ID NOs: 134 to 149, more preferably SEQ ID NOs: 134 to 140, more preferably SEQ ID NO: 134, in which up to 50, 49, 48, 47, 46, 45, 44, 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31,



30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0 amino acids are exchanged by any amino acid.

In any embodiment of the present invention relating to a designed ankyrin repeat domain or a recombinant binding protein comprising an amino acid sequence that has at least  
5 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identity to a given amino acid sequence, the non-identical amino acids may be located at any position of the designed ankyrin repeat domain or the recombinant binding protein.

Likewise, in any embodiment of the present invention relating to a designed ankyrin repeat domain or a recombinant binding protein in which up to 50, 49, 48, 47, 46, 45, 44,  
10 43, 42, 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0 amino acids are exchanged by any amino acid, the exchanged amino acids amino acids may be located at any position of the designed ankyrin repeat domain.

In one embodiment, the invention relates to a recombinant binding protein comprising an  
15 amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 134 to 179, preferably SEQ ID NOs: 134 to 158, more preferably SEQ ID NOs: 134 to 149, more preferably SEQ ID NOs: 134 to 140, more preferably SEQ ID NO: 134.

The invention particularly relates to a recombinant binding protein comprising an amino acid sequence that has at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%,  
20 or 100% amino acid sequence identity with the amino acid sequence of SEQ ID NO: 134.

In one embodiment, the invention relates to a recombinant binding protein comprising the amino acid sequence of SEQ ID NO: 134, in which up to 50, 49, 48, 47, 46, 45, 44, 43, 42,  
25 41, 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, or 0 amino acids are exchanged by any amino acid.

The invention further particularly relates to a recombinant binding protein comprising the amino acid sequence consisting of the amino acid sequence of SEQ ID NO: 134. In one embodiment, the invention relates to a recombinant binding protein comprising the amino acid sequence of SEQ ID NO: 134.

30 In one embodiment, the invention relates to a recombinant binding protein consisting of an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 134 to 179, preferably SEQ ID NOs: 134 to 158, more preferably SEQ ID NOs: 134 to 149, more preferably SEQ ID NOs: 134 to 140, more preferably SEQ ID NO: 134.

Preferred is SEQ ID NO: 134. Preferred is a recombinant binding protein, wherein the amino acid sequence is SEQ ID NO: 134. Preferred is a protein, wherein the amino acid sequence is SEQ ID NO: 134. Preferred is a recombinant binding protein consisting of the amino acid sequence of SEQ ID NO: 134.

5 Multiple features make SEQ ID NO: 134 the preferred recombinant binding protein of the invention. It comprises two designed ankyrin repeat domains with binding specificity for serum albumin each consisting of SEQ ID NO: 50, which shows improved storage stability properties (see Example 9; Figure 2) compared to known designed ankyrin repeat domain with binding specificity for serum albumin. It comprises two designed ankyrin repeat  
10 domains with binding specificity for serum albumin, which surprisingly leads to improved pharmacokinetic properties (Examples 5 and 6, Figures 3 and 4). The two designed ankyrin repeat domains with binding specificity for serum albumin are flanking the other designed ankyrin repeat domains leading to the best pharmacokinetic properties observed (Example 6). The designed ankyrin repeat domains and with binding specificity for VEGF-  
15 A and HGF as well as their structural arrangement were chosen to maximize activity of the compound (Example 8). The designed ankyrin repeat domains are connected using a PT-rich linker, surprisingly leading to improved activity of the individual designed ankyrin repeat domains (Example 8) and surprisingly leading to improved pharmacokinetic properties (Example 5).

20 In one embodiment, the invention relates to a recombinant binding protein comprising at least a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum  
25 albumin, and wherein said first, second, third, and fourth designed ankyrin repeat domain are linked by polypeptide linkers, and wherein said recombinant binding protein exhibits an increase in terminal half-life, preferably an increase in terminal half-life of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, or 45%, compared to the recombinant binding protein lacking said fourth designed ankyrin repeat domain with binding specificity for  
30 serum albumin. Examples of such an increase in terminal half-life are given in Examples 5 and 6 and Figures 3 and 4.

In one embodiment, the invention relates to a nucleic acid encoding the amino acid sequence of a designed ankyrin repeat domain or a recombinant binding protein of the present invention, more preferably a recombinant binding protein of the present invention.

35 In one embodiment, the invention relates to a nucleic acid encoding the amino acid

sequence of any recombinant binding protein of the present invention comprising at least two, more preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin. In one embodiment, the invention relates to a nucleic acid encoding the amino acid sequence of a recombinant binding protein of the present invention. Furthermore, the invention relates to vectors comprising any nucleic acid of the invention. Nucleic acids are well known to the skilled person. In the examples, nucleic acids were used to produce designed ankyrin repeat domains or recombinant binding proteins of the invention in *E. coli*.

In one embodiment, the invention relates to a pharmaceutical composition comprising a recombinant binding protein and/or a designed ankyrin repeat domain of the present invention, or a nucleic acid encoding a recombinant binding protein and/or a designed ankyrin repeat domain of the present invention, and optionally a pharmaceutically acceptable carrier and/or diluent.

In one embodiment, the invention relates to a pharmaceutical composition comprising a recombinant binding protein or a nucleic acid encoding a recombinant binding protein, and optionally a pharmaceutically acceptable carrier and/or diluent.

Pharmaceutical acceptable carriers and/or diluents are known to the person skilled in the art and are explained in more detail below. Even further, a diagnostic composition comprising one or more of the above mentioned recombinant binding proteins and/or designed ankyrin repeat domains, and/or nucleic acids, in particular recombinant binding proteins, is considered.

A pharmaceutical composition comprises a recombinant binding protein, and/or a designed ankyrin repeat domain, and/or a nucleic acid as described herein and a pharmaceutically acceptable carrier, excipient or stabilizer, for example as described in Remington's Pharmaceutical Sciences 16<sup>th</sup> edition, Osol, A. Ed., 1980. Suitable carriers, excipients or stabilizers known to the skilled man are saline, Ringer's solution, dextrose solution, Hank's solution, fixed oils, ethyl oleate, 5% dextrose in saline, substances that enhance isotonicity and chemical stability, buffers and preservatives. Other suitable carriers include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids and amino acid copolymers. A pharmaceutical composition may also be a combination formulation, comprising an additional active agent, such as an anti-cancer agent or an anti-angiogenic agent, or an additional bioactive compound.

One embodiment of the present invention relates to the use of a recombinant binding protein of the present invention comprising at least two, preferably comprising two, designed ankyrin repeat domains with binding specificity for serum albumin for manufacturing a pharmaceutical composition, wherein said recombinant binding protein  
5 exhibits an increased terminal half-life, preferably an increased terminal half-life of at least 5%, preferably 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, or 250%, compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin.

In one embodiment, a pharmaceutical composition comprises at least one recombinant  
10 binding protein as described herein and a detergent such as nonionic detergent, a buffer such as phosphate, and a sugar such as sucrose. In one embodiment, such a composition comprises recombinant binding proteins as described above and PBS.

In one embodiment, the invention relates to the use of a pharmaceutical composition, or a recombinant binding protein according to the present invention for the treatment of a  
15 disease. For that purpose, the pharmaceutical composition, or the recombinant binding protein according to the present invention is administered, to a patient in need thereof, in a therapeutically effective amount. Administration may include topical administration, oral administration, and parenteral administration. The typical route of administration is parenteral administration. In parental administration, the medicament of this invention will  
20 be formulated in a unit dosage injectable form such as a solution, suspension or emulsion, in association with the pharmaceutically acceptable excipients as defined above. The dosage and mode of administration will depend on the individual to be treated and the particular disease.

Further, any of the above mentioned pharmaceutical composition or recombinant binding  
25 protein is considered for the treatment of a disorder.

A pharmaceutical composition of the present invention may be administered by e.g. parenteral administration. In parental administration, the medicament of this invention will be formulated in a unit dosage injectable form such as a solution, suspension or emulsion, in association with the pharmaceutically acceptable excipients as defined above. The  
30 dosage and mode of administration will depend on the individual to be treated and the particular disease. In one embodiment, said recombinant binding protein or such other pharmaceutical composition described herein is applied intravenously. For parenteral application, the recombinant binding protein or said pharmaceutical composition can be injected as bolus injection or by slow infusion at a therapeutically effective amount.

In one embodiment, the invention relates to a method of treatment of a medical condition, the method comprising the step of administering, to a patient in need of such a treatment, a therapeutically effective amount of a recombinant binding protein of the invention. In one embodiment, the invention relates to a method of treatment of a medical condition, the method comprising the step of administering, to a patient in need of such a treatment, a therapeutically effective amount of a pharmaceutical composition of the invention. Example 14 (Figure 10) illustrates the utility of the use of a recombinant binding protein consisting of SEQ ID NO: 134 for the treatment of cancer. In one embodiment, the invention relates to the use of a pharmaceutical composition of the present invention for the treatment of a disease. In one embodiment, the invention relates to a pharmaceutical composition for use in the treatment of a disease.

A "medical condition" (or disorder) may be one that is characterized by inappropriate angiogenesis. A medical condition may be a hyperproliferative condition. Examples of medical conditions suitable for treatment include autoimmune disorders, inflammatory disorders, retinopathies (particularly proliferative retinopathies), neurodegenerative disorders, infections, and neoplastic diseases. Any of the recombinant binding proteins described herein may be used for the preparation of a medicament for the treatment of such a disorder, particularly a disorder selected from the group consisting of: an autoimmune disorder, an inflammatory disorder, a retinopathy, and a neoplastic disease. The invention particularly relates to a method of treating a medical condition, the method comprising the step of administering, to a patient in need of such treatment, a therapeutically effective amount of a recombinant binding protein or said pharmaceutical composition of the invention. In some embodiments said medical condition is a neoplastic disease. The term "neoplastic disease", as used herein, refers to an abnormal state or condition of cells or tissue characterized by rapidly proliferating cell growth or neoplasm. In a more specific meaning, the term relates to cancer. In a more specific meaning, the term may relate to renal cancer and/or gastric cancer and/or multiple myeloma. The term "therapeutically effective amount" means an amount that is sufficient to produce a desired effect on a patient.

In particular, the invention relates to the treatment of a medical condition using a pharmaceutical composition of the present invention, wherein said medical condition is cancer.

The use of a recombinant binding protein of the present invention or said pharmaceutical compositions for the treatment of cancer diseases can also be in combination with any other therapy known in the art. The term "use in combination with", as used herein, shall

refer to a coadministration, which is carried out under a given regimen. This includes synchronous administration of the different compounds as well as time-shifted administration of the different compounds (e.g. compound A is given once and compound B is given several times thereafter, or vice versa, or both compounds are given  
5 synchronously and one of the two is also given at later stages).

The use of a recombinant protein for the treatment of a disease including pathological angiogenesis is further considered. The term "pathological angiogenesis" refers to the formation and growth of blood vessels during the maintenance and the progression of several disease states.

10 In a further embodiment, the invention relates to the use of a recombinant binding protein of the invention for the manufacture of a medicament that is used for the treatment of a medical condition, preferably a neoplastic disease, more preferably cancer.

In one embodiment, the invention relates to the use of a pharmaceutical composition of the invention for the manufacture of a medicament that is used for the treatment of a  
15 medical condition, which may be a neoplastic disease, in particular cancer.

The formulations to be used for *in vivo* administration must be aseptic or sterile. This is readily accomplished by filtration through sterile filtration membranes.

The term "selected from the group consisting of" in connection with a single choice in this invention has the meaning of that particular choice. For example, in one embodiment, "the  
20 invention relates to a recombinant binding protein comprising an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NO: 134", which has the meaning "the invention relates to a recombinant binding protein comprising the amino acid sequence of SEQ ID NO: 134".

In one embodiment the invention relates to a recombinant binding protein comprising any  
25 of the above mentioned repeat domains. In one embodiment, the invention relates to a recombinant binding protein comprising any of the above mentioned SEQ ID NO: 134 to 179.

The invention is not restricted to the particular embodiments described in the Examples. Other sources may be used and processed following the general outline described below.

30 A number of documents are cited throughout this specification. The disclosure content of these documents is herewith incorporated by reference.

This specification refers to a number of amino acid sequences of the amino acid sequence listing of this specification named "MD41211\_Sequence\_Listing.txt" and the amino acid sequences of the sequence listing are herewith incorporated by reference.

## EXAMPLES

All of the standard materials and reagents disclosed here are known to those skilled in the art, and are available commercially or can be prepared using well-known techniques.

### 5 Materials

Chemicals were purchased from Sigma-Aldrich (Switzerland). Oligonucleotides were from Microsynth (Switzerland). Unless stated otherwise, DNA polymerases, restriction enzymes and buffers were from New England Biolabs (USA) or Thermo Fisher Scientific Fermentas (Lithuania). The cloning and protein production strain was *E. coli* XL1-blue (Stratagene, USA) or BL21 (Novagen, USA). Recombinant VEGF-A (human, mouse, rat), VEGF-C, PDGF-AB, and HGF (human, cynomolgus monkey, mouse) were from R&D Systems (Biotechne; Minneapolis, USA), Peprotech (Rocky Hill, USA), Sino Biological (Beijing, China), ReliaTech (Wolfenbüttel, Germany) or produced in Chinese Hamster Ovary Cells or in *Pichia pastoris* and purified according to standard protocols. Serum albumin of different species were from Sigma-Aldrich, Innovative Research (Novi, USA), CSL Behring (Switzerland), or collected from animals directly using standard methods. Biotinylated VEGF-A or HGF were obtained chemically via coupling of the biotin moiety to primary amines of the protein using standard biotinylation reagents and methods (Thermo Fisher Scientific Inc., USA). Antibodies were from Thermo Fisher Scientific or QIAGEN (Germany), or were generated using standard immunization and hybridoma procedures in mice or rabbits, procedures well known to the person skilled in the relevant art. Cell culture reagents were from Lonza (Switzerland), Roche (Switzerland), Thermo Fisher Scientific, and Promocell (Germany).

### Molecular Biology

25 Unless stated otherwise, methods are performed according to described protocols (Sambrook J., Fritsch E.F. and Maniatis T., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory 1989, New York).

### Designed ankyrin repeat domains, libraries and selections

30 Methods to generate designed ankyrin repeat protein libraries, examples of designed ankyrin repeat protein libraries, and methods to select designed ankyrin repeat proteins from libraries of designed ankyrin repeat proteins are described (WO 2002/020565; WO 2010/060748; WO 2012/069654; WO 2012/069655; WO 2014/001442; Binz et al. 2004, loc. cit.).

Example 1: Selection, expression, purification, and analysis of designed ankyrin repeat domains with binding specificity for VEGF-A, HGF, or serum albumin

Using ribosome display (Binz et al., 2004, loc. cit.) designed ankyrin repeat domains with binding specificity for VEGF-A, HGF, or serum albumin were selected from combinatorial  
5 libraries by methods described in WO 2010/060748 for the generation of designed ankyrin repeat domains with binding specificity for VEGF-A, and by methods described in WO 2014/191574 for the generation of designed ankyrin repeat domains with binding specificity for HGF, and by methods described in WO 2012/069654 for the generation of designed ankyrin repeat domains with binding specificity to serum albumin. The binding of  
10 the selected clones toward specific (VEGF-A, HGF, or serum albumin, respectively) and unspecific (e.g. MBP, *E. coli* maltose binding protein) targets was assessed by crude extract ELISA, indicating that hundreds of designed ankyrin repeat domains with binding specificity to VEGF-A, HGF, or serum albumin, respectively, were successfully selected in each selection for the respective target. For example, the designed ankyrin repeat domains of SEQ ID NO: 12 to 22 constitute amino acid sequences of ankyrin repeat domains with binding specificity for VEGF-A, the designed ankyrin repeat domains of SEQ ID NO: 23 to 37 constitute amino acid sequences of designed ankyrin repeat domains with binding specificity for HGF, and the designed ankyrin repeat domains of SEQ ID NO: 40 to 56 constitute amino acid sequences of designed ankyrin repeat domains with binding  
20 specificity for serum albumin.

These designed ankyrin repeat domains with binding specificity for VEGF-A, HGF, or serum albumin, and negative control designed ankyrin repeat domains with no known binding specificity (i.e. Proteins #10 and #11) were cloned into a pQE (QIAGEN, Germany) based expression vector providing an N-terminal His-tag to facilitate simple protein  
25 purification. The proteins were produced and purified with methods known to the person skilled in the art such as described for example in WO 2010/060748.

Example 2: Characterization of designed ankyrin repeat domains using surface plasmon resonance

SPR was measured using a ProteOn instrument (BioRad) and measurement was  
30 performed according standard procedures known to the person skilled in the art. Kd values that were measured for selected proteins are listed in Tables 1 to 3.



Table 1: Examples of dissociation constants of designed ankyrin repeat domains binding to human VEGF-A

Protein #*	Kd [pM]
12	94
13	96
16	141

\* Protein #12, #13, and #16 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NO: 12, 13 and 16, and additionally an N-terminal His-tag (SEQ ID NO: 1).

Similar VEGF-A dissociation constant values are obtained for Proteins #14, #15, and #17 to #22.

Table 2: Examples of dissociation constants of designed ankyrin repeat domains binding to human HGF

Protein #*	Kd [pM]
23	16
24	163
25	66
26	51
27	129
28	26
29	25

\* Protein #23 to #29 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NO: 23 to 29, and additionally an N-terminal His-tag (SEQ ID NO: 1).

Similar HGF dissociation constant values are obtained for Proteins #30 to #37.

Table 3: Examples of dissociation constants of designed ankyrin repeat domains binding to human HSA

Protein #*	Kd [nM]
44	26
45	13
46	22
47	27
48	20
49	15
50	27
51	14
52	6
54	11
55	15

\* Protein #44, to #55 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NO: 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, and 55, and additionally an N-terminal His-tag (SEQ ID NO: 1).

Similar human serum albumin dissociation constant values are obtained for Proteins #40 to #43, #53, #56, and #57.

Example 3: Competition binding assays and receptor competition binding assays

5 Characterization of designed ankyrin repeat domains with binding specificity for VEGF-A, HGF, or serum albumin, respectively, by competition assays. Such assays are well known to the person skilled in the art. For designed ankyrin repeat domains with binding specificity to VEGF-A, a quantitative sandwich enzyme immunoassay technique was used according to the manufacturer (VEGF-A Quantikine kit DVE00, R&D Systems). A monoclonal antibody specific for VEGF-A was pre-coated onto a microplate. VEGF-A

standards and mixtures of VEGF-A (20 pM) and Protein #18, #19, or #20 at varying concentrations were applied to the wells and any free VEGF-A present (i.e. not bound to the designed ankyrin repeat domain) is bound by the immobilized antibody. After washing away any unbound substances, an enzyme-linked polyclonal antibody specific for VEGF-A is added to the wells. Following a wash to remove any unbound antibody-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of VEGF-A bound in the initial step. The color development is stopped and the intensity of the color is measured. In this assay, the tested designed ankyrin repeat proteins showed high VEGF-A inhibition potency. IC<sub>50</sub> values were calculated from such titration curves obtained as described above using Graph Pad Prism software and standard procedures known to the person skilled in the art. For designed ankyrin repeat domains with binding specificity to HGF, a cMet receptor competition assay was performed. For that purpose, 5 nM of human cMet receptor in PBS were immobilized on Maxisorp plates overnight at 4°C. After washing with PBS 0.05% TWEEN 20, the plate was blocked for 2 h with shaking at 300 rpm using PBS containing 0.05% TWEEN 20 and 0.25% Casein at RT. A constant concentration of 5 nM human HGF was pre-incubated for 30min at RT with a 1000 nM-1 pM of Proteins #23, #26, #28, and #29 (1:4 dilution series each) on a dilution plate in PBS 0.05% TWEEN 20. After washing of the ELISA plate with PBS 0.05% TWEEN, pre-incubated samples were transferred to the ELISA plate and plate was incubated for 2 h at RT with shaking at 300 rpm. After washing with PBS 0.25% TWEEN, 200 ng/ml anti human HGF antibody was added for 1h at RT with shaking at 300 rpm. After washing with PBS 0.05% TWEEN, 100 ng/ml HRP conjugated polyclonal anti HGF species antibody was added for 30min at RT with shaking at 300 rpm. Detection was using 1:3 diluted BM blue POD (Roche). Color reaction was stopped after 15 min by addition of 1 M H<sub>2</sub>SO<sub>4</sub>. Readout was done at A450 using A620 as reference wavelength.

Example IC<sub>50</sub> values obtained by these assays are given in Tables 4 and 5. Similar VEGF-A IC<sub>50</sub> values are obtained with Proteins #12 to #17, #21, and similar HGF IC<sub>50</sub> values are obtained with Proteins #24, #25, #27, and #30 to #37.

Table 4: Inhibition of the VEGF-A-binding of an antibody by designed ankyrin repeat domains (mean IC<sub>50</sub> values)

Protein #*	IC <sub>50</sub> [pM]
#18	13
#19	13
#20	4

\* Protein #18 to #20 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NO: 18 to 20, and additionally an N-terminal His-tag (SEQ ID NO: 1).

Table 5: Inhibition of HGF binding to cMET by designed ankyrin repeat domains (mean IC<sub>50</sub> values)

Protein #*	IC <sub>50</sub> [pM]
23	915
26	623
28	955
29	1357

\* Protein #23, #26, #28, and #29 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NO: 23, 26, 28, and 29, and additionally an N-terminal His-tag (SEQ ID NO: 1).

Example 4: Generation of recombinant binding proteins, in particular recombinant proteins comprising two, three or four designed ankyrin repeat domains, and other repeat proteins

- DNA encoding designed ankyrin repeat domains or recombinant binding proteins was generated by genetic means well known to the person skilled in the art. Recombinant binding proteins selected from the group of amino acid sequences SEQ ID NOs: 58 to 133, additionally having SEQ ID NO: 1 or the amino acids GS at the N terminus or recombinant binding proteins selected from the group of amino acid sequences SEQ ID

NOs: 134 to 179, or designed ankyrin repeat domains selected from the group of amino acid sequences SEQ ID NOs: 10 to 57, additionally having SEQ ID NO: 1 or the amino acids GS at the N terminus, were expressed in the cytoplasm of *Escherichia coli* using standard techniques using the pQE expression system from Qiagen (Germany). In case  
5 the amino acids GS were at the N terminus, the Met residue additionally encoded by the expression vector was efficiently cleaved off in the cytoplasm of *E. coli* from the expressed polypeptide since the start Met is followed by a small Gly residue (i.e. the amino acid at position 1 of SEQ ID NOs: 134 to 179). The cells were lysed by using a French press, and the proteins were purified to near homogeneity from the crude cell extract by using  
10 standard chromatographic techniques well known to the person in the art.

Example 5. Improving pharmacokinetic properties with increasing numbers of designed ankyrin repeat domains with binding specificity for serum albumin comprised in a recombinant binding protein – mouse pharmacokinetic studies.

15 For mouse pharmacokinetic studies Proteins #57, #62, #63, #64, #68, #73, #74, #82, #83, #97, #109, and #110 (proteins corresponding to SEQ ID NOs: 57, 62, 63, 64, 68, 73, 74, 82, 83, 97, 109, and 110, additionally having SEQ ID NO: 1 at the N terminus) prepared as described in Example 4, were labeled radioactively as described (Zahnd, C., Kawe, M., Stumpp, M.T., de Pasquale, C., Tamaskovic, R., Nagy-Davidescu, G., Dreier, B., Schibli,  
20 R., Binz, H.K., Waibel, R., Plückthun, A., Cancer Res.70, 1595-1605, 2010) and administered at 10 µg in 100 µl as a single intravenous bolus injection into the tail vein of female BALB/c mice, respectively. Serum samples from each mouse were collected at various time points and the accumulated radioactivity was determined using a gamma-scintillation counter.

25 In these experiments, proteins comprising two designed ankyrin repeat domains with binding specificity for serum albumin consistently exhibited improved pharmacokinetic properties compared to comparable constructs comprising only one designed ankyrin repeat domain with binding specificity for serum albumin (Figure 3). For example, the comparison of Protein #57, which comprises a single designed ankyrin repeat domain with  
30 binding specificity for serum albumin (SEQ ID NO: 57, comprising SEQ ID NO: 51 plus a C-terminal polypeptide) with Proteins #62 and #63, which comprise two designed ankyrin repeat domains with binding specificity for serum albumin (two times SEQ ID NO: 51, linked by GS- (SEQ ID NO: 63) or PT-rich (SEQ ID NO: 62) polypeptide linkers) shows that having two designed ankyrin repeat domains with binding specificity for serum  
35 albumin leads to higher % ID at e.g. 24 h (+57% GS; +59% PT), 48 h (+76% GS; +82%

PT) or 72 h (+79% GS; +94% PT) post-injection, and leads to an improved terminal half-life (+38% GS; +48% PT) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin (Figure 3a). In particular, using a PT-rich linker, in particular SEQ ID NO: 9, leads to improved pharmacokinetic properties (Figure 3a). The following three examples show that the method of improving pharmacokinetic properties by having two (instead of one) designed ankyrin repeat domains with binding specificity for serum albumin present in a protein, is transferable to different proteins comprising different designed ankyrin repeat domains. For example, the comparison of the pharmacokinetic profile of Protein #64, comprising SEQ ID NOs: 22 (designed ankyrin repeat domain with binding specificity for another target than serum albumin) and 51 (designed ankyrin repeat domain with binding specificity for serum albumin), with Proteins #73 and #74, comprising each SEQ ID NOs: 22 and two times 51 (Protein #73 has SEQ ID NOs: 51 flanking SEQ ID NO: 22, and Protein #74 has SEQ ID NOs: 51 N-terminal of SEQ ID NO: 22) shows that having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+62% N-terminal; +89% flanking), or 48 h (+136% N-terminal; +175% flanking) post-injection, and leads to an improved terminal half-life (>63% for both N-terminal or flanking) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin (Figure 3b). Likewise, a pharmacokinetic profile comparison of Protein #82, comprising twice SEQ ID NO: 11 (designed ankyrin repeat domain with no known binding specificity) and once SEQ ID NO: 51 (designed ankyrin repeat domain with binding specificity for serum albumin), with Protein #109, comprising twice SEQ ID NO: 11 and twice SEQ ID NO: 51 (N-terminal), indicates that having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+12%), or 48 h (+35%) post-injection, and leads to an improved terminal half-life (+71%) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin (Figure 3c). Furthermore, the pharmacokinetic profile comparison of Protein #83, comprising SEQ ID NOs: 38 and 39 (designed ankyrin repeat domains each with binding specificity for another target than serum albumin) and 50 (designed ankyrin repeat domain with binding specificity for serum albumin), with Proteins #110, comprising each SEQ ID NOs: 38 and 39 and twice SEQ ID NO: 50 (flanking SEQ ID NOs: 38 and 39), shows that having two designed ankyrin repeat domains with binding specificity for serum albumin leads to higher % ID at e.g. 24 h (+198%), 48 h (+198%), or 72 h (+228%) post-injection, and leads to an improved terminal half-life (+19%) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin

(Figure 3d). Furthermore, Protein #97 exhibited significantly improved pharmacokinetic properties over Protein #68, e.g. the terminal half-lives were 21 and 16 hours, respectively, indicating that having two designed ankyrin repeat domains with binding specificity for serum albumin is beneficial over having only one such domain .

- 5 These results indicate, surprisingly, that using two designed ankyrin repeat domains with binding specificity for serum albumin instead of one in a recombinant binding protein leads to improved pharmacokinetic properties, as discussed further in Example 6.

10 Example 6. Improving pharmacokinetic properties with increasing numbers of designed ankyrin repeat domains with binding specificity for serum albumin comprised in a recombinant binding protein – cynomolgus monkey pharmacokinetic studies.

For cynomolgus monkey pharmacokinetic studies Proteins #57, #62, and #97 (proteins corresponding to SEQ ID NOs: 57, 62, and 97, additionally having SEQ ID NO: 1 at the N terminus) and Protein #134 (a protein corresponding to SEQ ID NOs: 134), prepared as described in Example 4, were administered via intravenous infusion for 30 min at a target dose level of between 0.5-100 mg/kg to cynomolgous monkeys. Blood samples were collected pre-dose and again at selected time points, for example 5 min, 10 min, 0.5h, 1h, 2h, 4h, 8h, 12h, 24h, 48h, 72h, 96h, 120h and 168h post-end of infusion (i.e. post-injection). The blood samples were allowed to stand at room temperature and were centrifuged to generate serum, followed by storage at -80°C pending analyses. Pharmacokinetic parameters were determined using procedures well known to the person skilled in the art. Serum concentrations of Proteins #57, #62, #97 and #134 were determined by sandwich ELISA using a rabbit monoclonal anti-designed ankyrin repeat domain antibody as capture reagent and murine monoclonal anti-designed ankyrin repeat domain antibody as detection reagent, and using a standard curve. Pharmacokinetic parameters were determined using standard software such as Phoenix WinNonLin (Certara, Princeton, USA) or GraphPadPrism (GraphPad Software, La Jolla, USA) and standard analyses such as non-compartmental analyses. The resulting pharmacokinetic profiles are shown in Figure 4. Proteins comprising two designed ankyrin repeat domains with binding specificity for serum albumin consistently exhibited improved pharmacokinetic properties compared to comparable constructs comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. For example by comparing Protein #57 (0.5 mg/Kg; 27.7 nmol/kg), comprising a single designed ankyrin repeat domain with binding specificity for serum albumin (SEQ ID NO: 51), with Protein #62 (1.04 mg/Kg; 34.5 nmol/kg), a protein comprising two designed ankyrin repeat domains

with binding specificity for serum albumin (twice SEQ ID NO: 51, linked by a PT-rich polypeptide linker), the results show that having two designed ankyrin repeat domains with binding specificity for serum albumin in a protein leads to higher exposure (2138 d\*nmol/L vs. 4676 d\*nmol/L, i.e. +119% calculated up to day 7), leads to a reduced clearance (0.0108 L/(d\*kg) vs. 0.0031 L/(d\*kg); i.e. -71%), and leads to an improved terminal half-life (4.57 d vs. 9.00 d, i.e. +97% calculated from day 1 to day 7) compared to the protein comprising only a single designed ankyrin repeat domain with binding specificity for serum albumin (Figure 4a). Also, the percentage injected dose, normalized to the concentration measured 10 min post-injection, is increased comparing Protein #57 with Protein #62 at day 4 (23.39% vs. 57.72%; +148%), day 5 (19.00% vs. 48.41%; +155%), and day 6 (18.5% vs. 51.94%; +175%). As a further example in cynomolgus monkey, Protein #134 (a protein corresponding to SEQ ID NOs: 134, produced as described in Example 4, was tested at different doses in 10 animals each (5 male, 5 female, each dose) and the terminal half-life was evaluated using WinNonLin considering concentration values up to day 7. Protein #134 exhibited an average terminal half-life of 4.0 days (95 h) when given at 1 mg/kg (compared to Protein #97 exhibiting 2.7 days (65 h) terminal half-life at 1 mg/kg (+46%)), 5.3 days (127 h) when given at 10 mg/kg, and 5.8 days (139 h) when given at 100 mg/kg to cynomolgus monkey. The pharmacokinetic profile of Protein #134 in comparison to Protein #97 in cynomolgus monkey is shown in Figure 4b. Like in Example 5, these results indicate, surprisingly, that using two designed ankyrin repeat domains with binding specificity for serum albumin instead of one in a recombinant binding protein leads to improved pharmacokinetic properties. These results are discussed in the following.

In the absence of any albumin binding activity, a recombinant binding protein has a terminal half-life in the range of minutes both in mouse and cynomolgus monkey (See WO 2012/069654). Proteins comprising at least one designed ankyrin repeat domain with binding specificity for serum albumin show terminal half-lives which are far greater than if no designed ankyrin repeat domain with binding specificity for serum albumin is present. A pharmacokinetic profile of a protein comprising one designed ankyrin repeat domain with binding specificity for serum albumin are shown in Figures 3a and 4a.

The art contains a study, in which the effect of valency of another serum albumin binding protein domain, the albumin binding domain (ABD) derived from streptococcal protein G, was investigated (Hopp et al., 2010; loc. cit.). Another study uses C-terminally fused peptides (WO 2011/095545), which are no protein domains. ABD is a helical protein domain with binding specificity for serum albumin. Importantly, Hopp et al. 2010 (loc. cit.)



show that having two such ABDs (one N-terminal and one C-terminal) in a recombinant binding protein does not lead to a significantly improved terminal half-life in mouse compared to a recombinant binding protein comprising only one ABD (C-terminal;  $37.9 \pm 1.1$  h vs.  $36.4 \pm 4.8$  h). In particular, at 24 h and 72 h post injection, the recombinant binding protein comprising one ABD showed identical percentage injected doses as the recombinant binding protein comprising two ABD, indicating equivalent pharmacokinetic properties of the two recombinant binding proteins. Based on the findings with ABD, one skilled in the art would expect that a recombinant binding protein comprising two albumin binding protein domains such as designed ankyrin repeat domains with binding specificity for serum albumin would not have improved pharmacokinetic properties compared to a recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. Surprisingly, we found that this is not the case. In contrast to Hopp et al. (loc. cit.), recombinant binding proteins comprising two designed ankyrin repeat domains with binding specificity for serum albumin surprisingly exhibited clearly prolonged terminal half-lives compared to recombinant binding proteins comprising only one designed ankyrin repeat domain with binding specificity for serum albumin.

These examples illustrate a number of additional findings. For example, the pharmacokinetic properties of Protein #134 are superior to the ones of Protein #97, illustrating the importance of the choice of the individual designed ankyrin repeat domains. SEQ ID NO: 134 was chosen to be composed of components that lead to maximal activity and optimal pharmacokinetic properties. Also, the arrangement of the designed ankyrin repeat domains within Protein #134 was chosen to lead to optimal pharmacokinetic properties. When analyzing recombinant binding proteins comprising four designed ankyrin repeat domains including two designed ankyrin repeat domains with binding specificity for serum albumin regarding mouse and cynomolgus monkey pharmacokinetics, the most favorable pharmacokinetic properties were observed for recombinant binding proteins having the two designed ankyrin repeat domains with binding specificity for serum albumin flanking the other two designed ankyrin repeat domains.

As the examples of this example comprise different combinations of designed ankyrin repeat domains with binding specificity for another target than serum albumin, the approach of using at least two designed ankyrin repeat domains with binding specificity for serum albumin for improving the pharmacokinetic properties appears to be generally applicable to proteins comprising several designed ankyrin repeat domains.

Example 7. Simultaneous binding of two human serum albumin molecules by Protein #134.

Protein #134 (A recombinant binding protein consisting of SEQ ID NO: 134, additionally having GS at the N terminus) was prepared as described in Example 4. Protein #134, purified human serum albumin (HSA), as well as Protein #134/HSA mixture (1:2 stoichiometry) were analyzed by size exclusion chromatography coupled to multi-angle static light scattering (SEC-MALS). SEC-MALS was performed using the proteins of Table 6 at 30  $\mu$ M (Protein #134) or 60  $\mu$ M (HSA) concentration on a Agilent 1200 system (Life Technologies, USA) connected to a Wyatt (USA) MALS and refractive index detector (flow rate: 0.6 ml/min; injection volume: 100  $\mu$ l; column: GE Healthcare (USA) Superdex200 10/300GL). The Protein #134/HSA mixture was pre-incubated for 3 hours at 20°C prior injection. The chromatograms are shown in Figure 5 and the molecular masses of the eluates were determined and compared to the theoretical molecular masses, as shown in Table 6. For this experiment, 100 mg HSA (CSL Behring 20% solution) were purified using a Superdex 200\_26.60 column on a AEKTA prime system (GE Healthcare; 2.0ml/min, PBS, isocratic flow, injection volume 10ml of 1:20 in PBS diluted HSA, collecting 4ml fractions). The peak fraction of the main peak was used for performance of the SEC-MALS experiment.

Protein #134 at 30  $\mu$ M was monodisperse with the elution fractions containing protein of the expected molecular weight (Figure 5). Likewise, purified HSA at 60  $\mu$ M was monodisperse with the elution fractions containing protein of the expected molecular weight (Table 6). The mixture of 30  $\mu$ M Protein #134 and 60  $\mu$ M HSA resulted in two peaks in SEC. One peak contained protein complexes of a molecular weight corresponding to a 1:2 (Protein #134/HSA) complex, indicating that the two designed ankyrin repeat domains with binding specificity for serum albumin are functional simultaneously. Additionally, in the tail of this peak, protein complexes of a molecular weight corresponding to a 1:1 (Protein #134/HSA) complex could be detected. Additionally, free HSA could be detected. As this amount is minor, one can rule out that the main peak is a 2:1 (Protein #134/HSA) complex, which would theoretically be consistent with the observed weight, yet a large fraction corresponding to 75% of free HSA would be expected. No free Protein #134/HSA could be detected. No peaks with a molecular weight corresponding larger than the one of the 1:2 (Protein #134/HSA) complex were detected. SEC-MALS measurements and the variations observed in SEC-MALS measurements are well known to the person skilled in the art.

Table 6. Size-exclusion chromatography coupled to static light scattering of Protein #134 and HSA, as well as the complex Protein #134/HSA.

Peaks	Theoretical stoichiometry & MW	MW measured
HSA	69366.6 Da	63350 Da
Protein #134	62397.0 Da	58700 Da
Protein #134/HSA tail	1:1, 131763.6 Da	132500 Da
Protein #134/HSA middle	mixture of 1:1 & 1:2, MW depends on ratio	173700 Da
Protein #134/HSA front	1:2, 201130.2 Da	197500 Da

The simultaneous binding of two human serum albumin molecules by one recombinant binding protein is similarly observed when analyzing Protein #97, and Protein #102, Protein #109, Protein #110 in size exclusion chromatography coupled to static light scattering.

Example 8: Maximizing target binding activity by choosing linker composition and by choosing the number of designed ankyrin repeat domains with binding specificity for serum albumin

10 Polypeptide linkers that link protein domains are well-known to the person skilled in the art. Gly-Ser-rich linkers are well-known from single-chain Fv antibody fragments, where they are used to link the two Fv polypeptide chains. Various other polypeptide linkers exist, including e.g. antibody hinge regions, or unstructured polypeptides such as sequences comprising mostly the amino acids Ala, Glu, Lys, Pro, Ser, Thr  
 15 (WO 2007/103515) or Ala, Pro, and Ser (WO 2008/155134). Furthermore, Pro-Thr-rich linkers have been disclosed (WO 2014/191574). The effect of such a linker on the properties of protein domains linked by such linker needs to be assessed for every linker/domain combination. Next to the nature of a polypeptide linker, we surprisingly found that the number of serum albumin-binding domains can influence the functionality of  
 20 a protein. To maximize target binding activity of the recombinant binding proteins of the present invention, recombinant binding proteins comprising Gly-Ser-rich and Pro-Thr-rich polypeptide linkers were compared as well as recombinant binding proteins comprising one or two designed ankyrin repeat domains with binding specificity for serum albumin. For that purpose, Proteins #69, #71, and, #107, each additionally having SEQ ID NO: 1 at  
 25 the N terminus and prepared as described in Example 4, were analyzed for binding to

VEGF-A and HGF, respectively, by ELISA (for methods see Example 4). The results are shown in Table 7. The comparison of the EC<sub>50</sub> values of Protein #69 with Protein #71 indicates that the recombinant binding protein with Pro-Thr-rich linkers is more potent with respect to binding of VEGF-A (factor 2) and HGF (factor 1.3), respectively, compared to the recombinant binding proteins with Gly-Ser-rich linkers. The comparison of the EC<sub>50</sub> values of Protein #69 with Protein #107 indicates that the recombinant binding protein comprising two designed ankyrin repeat domains with binding specificity for serum albumin is more potent with respect to binding of VEGF-A (factor 1.4) and HGF (factor 1.1), respectively, compared to the recombinant binding protein comprising only one designed ankyrin repeat domain with binding specificity for serum albumin. This is surprising in view of previous results (Hopp et al., 2010), where the presence of two albumin binding domains in a construct has a negative impact on the functionality of the molecule. This result indicates that recombinant binding proteins preferentially comprises Pro-Thr-rich linkers and two designed ankyrin repeat domains with binding specificity for serum albumin, rather than Gly-Ser-rich linkers and one designed ankyrin repeat domains with binding specificity for serum albumin.

Table 7: ELISA analysis of recombinant binding proteins with different linkers and different numbers of designed ankyrin repeat domains with binding specificity for serum albumin

Protein #*	Linker	Number of SABD†	EC <sub>50</sub> [nM]	
			VEGF-A	HGF
69	PT	1	0.102	0.117
71	GS	1	0.205	0.179
107	PT	2	0.073	0.107

\* Protein #69, #71, and #107 in this table represent designed ankyrin repeat domains consisting of the corresponding amino acid sequence of SEQ ID NOs: 69, 71, and 107, and additionally an N-terminal His-tag (SEQ ID NO: 1).

†Number of designed ankyrin repeat domains with binding specificity for serum albumin

Example 9. Improvement of protein stability when using SEQ ID NO: 50.

Proteins #48, #49, and #51 were further characterized for their midpoint of denaturation temperature (i.e. midpoint of the cooperative unfolding upon temperature increase) by

mixing the Proteins (25  $\mu$ l; 100  $\mu$ M in PBS) with a fluorescent dye (25  $\mu$ l Sypro orange (Life Technologies) diluted 1/2500 in PBS) and measuring a melting curve with a thermal cycler comprising a fluorescence reader (CFX96 Real-Time PCR Detection System; Biorad; 25 seconds holding time every 0.5°C followed by fluorescence read), essentially  
5 as described by Niesen et al. 2007 (Niesen, F.H., Berglund, H., Vedadi, M., Nature Protocols 2, 2212-2221, 2007). In PBS, Protein #48 exhibited a midpoint of denaturation of 83.5°C, and Protein #49 exhibited a midpoint of denaturation of 84.5°C, while Protein #51 exhibited a midpoint of denaturation of 79.5°C.

In order to identify the designed ankyrin repeat domain with binding specificity for serum  
10 albumin having the best storage stability properties, Proteins #49, #50 and #51 (corresponding to SEQ ID NOs: 49, 50 and 51, respectively, additionally having SEQ ID NO: 1 at the N terminus) were prepared as described in Example 4, and samples were concentrated to 10 mg/ml in PBS. Proteins #50 and #51 were then stored for 1 month at -80°C or at 40°C in glass vials, followed by analysis on SDS 15% PAGE. While Proteins  
15 #50 and #51 showed equivalent stability upon storage at -80°C, Proteins #50 showed significantly reduced amounts of degradation products by >50% reduction compared to Protein #51 on SDS 15% PAGE after 1 month storage at 40°C. Similarly, when stored at 4°C, 25°C, 40°C and 60°C for one week at 10mg/ml in PBS, Protein #50 showed significantly reduced amounts of degradation products compared to Protein #49. In  
20 particular, Protein #50 showed >50% reduction of degradation products compared to Protein #49 on SDS 15% PAGE both when stored at 40°C or 60°C, respectively (Figure 2). These findings illustrate that Protein #50 has an improved storage stability compared to Proteins #49 and #51. Similarly, when comparing the storage stability of Proteins #48 to #51 (corresponding to SEQ ID NOs: 48 to 51, additionally having SEQ ID NO: 1 at the N  
25 terminus; produced as described in Example 4), by incubating the proteins at 10 mg/ml in PBS in glass vials for 1 month at 40°C, Proteins #48 to #50 exhibit >30% reduction of degradation products compared to Protein #51.

These findings are corroborated by testing the storage stability of Protein #102 and Protein #103 (recombinant binding proteins consisting of the amino acid sequences  
30 corresponding to SEQ ID NOs: 102 and 103, both additionally having SEQ ID NO: 1 at the N terminus). Protein #102 and Protein #103 were prepared as described in Example 4, samples were concentrated to 10mg/ml in PBS and stored for 1 month at -80°C in or at 40°C in glass vials, followed by analysis on standard size-exclusion chromatography. While Protein #102 and #103 showed equivalent elution profiles upon storage at -80°C,  
35 Protein #102 showed 98.72% monomeric species and Protein #103 showed 100%

monomeric species upon storage at 40°C. This indicates that having SEQ ID NO: 50 present in the recombinant binding protein is more favorable regarding storage stability than having SEQ ID NO: 49 present. Similarly, Protein #103 exhibits lower amounts of degradation products than Protein #102 when analyzed by SDS-PAGE after 1 month storage at 40°C in glass vials in PBS at 10 mg/ml, confirming the higher storage stability of a recombinant binding protein comprising SEQ ID NO: 50 in comparison to the recombinant binding protein comprising SEQ ID NO: 49.

Similar results are obtained when comparing Protein#134 with Protein #143 or Protein #150 (recombinant binding proteins consisting of the amino acid sequences corresponding to SEQ ID NOs: 134, 143 and 150, respectively), prepared as described in Example 4. When stored at 40°C for one month at 10 mg/ml in PBS in glass vials, Protein #134 shows >50% reduction of degradation products compared to Proteins #143 and #150 when analyzed on SDS 15% PAGE. This indicates that having SEQ ID NO: 50 present in the recombinant binding protein is more favorable regarding storage stability than having either SEQ ID NO: 49 or SEQ ID NO: 51 present.

Example 10: Characterization of recombinant binding proteins using ELISA

Purified recombinant binding protein consisting of the amino acid sequence SEQ ID NO: 134, prepared as described in Example 4, was subjected to ELISA analyses. 100 µl or 50 µl of 20 nM target (VEGF-A, HGF, or serum albumin) in PBS per well were immobilized in a Maxisorp plate (Nunc, Denmark) overnight at 4°C. After washing 5 times with 300 µl PBST (PBS supplemented with 0.1% Tween 20), the wells were blocked with 300 µl PBST-C (PBST supplemented with 0.25% casein) for 2 h at room temperature with shaking at 450 rpm on a Titramax 1000 shaker (Heidolph, Germany). After washing 5 times as described above, 100 µl/well or 50 µl/well Protein #134 (concentrations ranging from 100 nM to 0.01 pM) in PBST-C was applied and incubated for 1 h to 2 h at room temperature with shaking at 450 rpm. After washing 5 times as described above, binding of Protein #134 was detected using 100 µl or 50 µl/well rabbit anti-designed ankyrin repeat domain monoclonal antibody in PBST-C for 1 h at room temperature with shaking at 450 rpm. After washing 5 times as described above, bound anti-designed ankyrin repeat domain antibody was detected using 100 µl or 50 µl/well goat anti-rabbit IgG-HRP conjugate in PBST-C for 1 h at room temperature with shaking at 450 rpm. After washing 5 times as described above, the ELISA was then developed using 100 µl BM soluble blue

POD substrate (Roche, Switzerland), diluted 1:4 in water. The reaction was stopped after 5 min using 100  $\mu$ l 1 M H<sub>2</sub>SO<sub>4</sub>. The OD (OD 450 nm - OD 620 nm) was then recorded.

The ELISA results indicate that Protein #134 binds human, cynomolgus monkey, rat and mouse VEGF-A with equivalent potency (Table 8 and Figure 6a). Cynomolgus monkey VEGF-A is identical to human VEGF-A and was thus not tested separately. No binding of Protein #134 to VEGF-C and PDGF-AB was detected (Table 9 and Figure 6a). Human, cynomolgus and mouse HGF is bound by Protein #134 with equivalent potency (EC<sub>50</sub> values in the 20 – 50 pM range; Table 8 and Figure 6b). Furthermore, Protein #134 binds serum albumin of human, cynomolgus monkey, rat, dog and mouse with equivalent potency (EC<sub>50</sub> values in the 10 – 20 pM range; Table 8 and Figure 6c). A comparison of Protein #134 (i.e. protein consisting of SEQ ID NO: 134) with Protein #60 or Protein #61 (i.e. proteins consisting of SEQ ID NOs: 60 or 61, additionally having SEQ ID NO: 1 at the N terminus, produced as described in Example 4), revealed that the EC<sub>50</sub> of Protein #134 observed for the binding of human serum albumin is significantly better than the ones observed for Proteins #60 or #61 (225 pM or 322 pM, respectively).

Table 8: Apparent EC<sub>50</sub> values of Protein #134 for binding VEGF-A, HGF and serum albumin of different species

Species	EC <sub>50</sub> [pM] VEGF-A (95% C.I.)*	EC <sub>50</sub> [pM] HGF (95% C.I.)*	EC <sub>50</sub> [pM] SA (95% C.I.)*
Human	24 (20-27)	24 (20-29)	13 (10-16)
Mouse	21 (19-23)	45 (38-52)	15 (13-17)
Rat	22 (18-26)	n.a.	17 (13-21)
Dog	n.a.	n.a.	23 (20-25)
Cynomolgus monkey	24 (20-27) <sup>†</sup>	40 (36-44)	17 (13-21)

\* C.I. confidence interval, <sup>†</sup> 100% sequence identity to human VEGF-A, thus the value for human VEGF-A is listed, n.a. not analyzed

Table 9. Apparent  $EC_{50}$  values of Protein #134 for binding different human VEGFs and PDGFs

Target	$EC_{50}$ [pM] (95% C.I.)*
Human VEGF-A	24 (20-27)
Human VEGF-C	No binding detected
Human PDGF-AB	No binding detected

\* C.I. confidence interval

Example 11: Characterization of recombinant binding proteins using competition assays

Purified recombinant binding protein consisting of the amino acid sequence SEQ ID NO: 134, prepared as described in Example 4, was subject to competition ELISA and FRET analyses. Such competition FRET and ELISA assays are well known to the person skilled in the art. Protein #134 was measured in a VEGF-A/VEGFR-2 competition FRET assay. For that purpose, Protein #134 and biotinylated VEGF-A165 (Reliatech, #300-076Bi-L) were prepared as eight-fold concentration stocks in PBS containing 0.2% BSA and 0.01% Tween (PBST-BSA). A competition mixture of 5  $\mu$ l eight-fold concentration Protein #134 and 5  $\mu$ l of eight-fold concentration biotinylated VEGF-A165 were pre-incubated for 1 hour at room temperature (competition mixture). In parallel, 5  $\mu$ l of Streptavidin-Tb (streptavidin-Lumi4-terbium cryptate donor, Cisbio #610SATLB) and 5  $\mu$ l of PAb anti-hIgG-de (D2-conjugated goat anti-Human IgG, Cisbio #61HFCDA) were added to 500  $\mu$ l of PBST-BSA buffer and incubated for 20 minutes (2x reagent). Ten  $\mu$ l/well of 2x reagent were dispensed in a 384-well HTRF white-plate (Thermo Fisher Scientific Inc.) and 5  $\mu$ l/well of four-time concentration hVEGF-R2-Fc fusion (Reliatech #SFC-008) were added. Five  $\mu$ l of the preincubated competition mixture were then added to wells. The complete reaction mix was incubated in the dark for 1 hour at room temperature before the fluorescence read out using a fluorescence reader. The final mixture contained 10 nM soluble VEGF-R2-Fc fusion, 10 nM biotinylated VEGF-A, and varying concentrations of Protein #134. The read-out was done for A665 nm and A595 nm wavelength (Excitation 340nm). The results of the assay are shown in Figure 7a. In this assay, Protein #134 inhibits the VEGF-A/VEGFR-2 interaction with an  $IC_{50}$  value of 0.6 nM. Protein #134 was further measured in a HGF/cMet competition ELISA experiment as described in Example 3. The results of



the assay are shown in Figure 7b. In this assay, Protein #134 inhibits the HGF/cMet interaction with an  $IC_{50}$  value of 0.92 nM. Protein #134 was also measured in a VEGF-A competition ELISA experiment as described in Example 3. The results are shown in Figure 7c. In this assay, Protein #134 inhibits VEGF-binding with an  $IC_{50}$  value in the  
5 single-digit pM range ( $IC_{50}$  4.5 pM).

Example 12: Characterization of simultaneous target binding of recombinant binding proteins using surface plasmon resonance

SPR was measured in a similar way as described in Example 2, with the following setup.  
10 2700 RU human HGF were immobilized on a sensor chip. Then 100 nM Protein #134 or PBST were injected for 180 seconds followed by a PBST wash of 360 seconds. Following this, 100 nM human VEGF-A or PBST were injected for 180 seconds (leading to saturation) followed by a PBST wash of 360 seconds. Finally, 100 nM human serum albumin or PBST were injected for 180 seconds followed by a PBST wash of 600  
15 seconds. The resulting signals are shown in Figure 8. The results indicate that Protein #134 can bind HGF, VEGF-A, and serum albumin. Furthermore, the results indicate that Protein #134 can bind HGF and VEGF-A, as well as HGF, VEGF-A, and serum albumin at the same time.

20 Example 13: Characterization of recombinant binding proteins in cell culture

Purified recombinant binding protein consisting of the amino acid sequence SEQ ID NO: 134, prepared as described in Example 4, was further subjected to cellular assays including a HUVEC proliferation assay to assess VEGF-A inhibition, and an A549 cell migration assay as well as a cMet phosphorylation assay, both to assess HGF inhibition,  
25 assays well known to the person in the art.

Inhibition of VEGF-A-induced HUVEC proliferation was determined by titrating increasing Protein #134 concentrations in the HUVEC proliferation assay. Human VEGF-A was used at a concentration of 8 ng/ml (corresponding to EC80 as determined in a proliferation assay). Protein #134 was titrated from 200 ng/ml to 0.195 ng/ml. Cells were seeded in  
30 50  $\mu$ l assay medium. Protein dilutions (in assay medium) were made by serial dilution 1:2 fold in a dilution plate; the concentration was four times the final concentration. Protein #134 dilutions were mixed with four-fold VEGF-A concentrations (32 ng/ml; final 8 ng/ml) in a ratio 1:1. 50  $\mu$ l of the mixtures were added to the cells for 72 h. Cell proliferation was determined either by BrdU incorporation in the replicating DNA or by monitoring metabolic

activity using WST-1. The results are shown in Figure 9a, indicating that Protein #134 exhibits an  $IC_{50}$  of 5.7 ng/ml (91.35 pM).

Inhibition of the HGF/cMet interaction was determined using Protein #134 in an Oris cell migration assay (Platypus Technologies, USA). The assay was performed according to the manufacturers' protocol. Briefly, cells were seeded with 100,000 A549 cells in serum-free DMEM. Cells adhered after 24 hours and medium was exchanged to assay medium; DMEM with and without 0.5 nM HGF with and without 5  $\mu$ M Protein. HGF and the neutralizing Protein were preincubated for 1h at RT before addition to cells. The Oris™ stoppers were removed. The assay was then incubated for 48 hours to permit cell migration. Cells were stained with Calcein (2.5 ng/ml) for 40 minutes and images were taken. The migration zone was measured using an inverse microscope Olympus and its software CellSens Dimension. The migration area was calculated as the covered area by subtracting the uncovered area of the pre-migration well with the uncovered area of respective samples wells. The uncovered area was the cell-free area and was measured using the diameter function in the processing folder of the software. The results are shown in Figure 9b and indicate that Protein #134 can suppress the HGF-induced cell migration of A549 cells.

Inhibition of cMet phosphorylation by Protein #134 was measured using A549 cells and a DuoSet P-cMet-ELISA (RnD Systems). Cells were seeded in complete medium in 96 well plates at 200,000 cells per well in complete medium. 24h later medium was replaced by serum-free medium. Cells were incubated for another 24h and stimulated by 1 nM human HGF (or PBS for negative control) in the presence and absence of Protein #134. HGF and Protein #134 were preincubated for at least 30 min at room temperature prior to addition to cells. Cells were stimulated for 10 minutes at room temperature. Stimulation was terminated by removing the cell supernatant (by flicking) and addition of cell lysis buffer according to protocol. Cell lysates were kept at -20°C until the ELISA experiment. The results are shown in Figure 9c and indicate that Protein #134 can suppress HGF-mediated cMet phosphorylation with an  $IC_{50}$  of 184 pM.

30 Example 14: Effect of recombinant binding proteins on tumor growth in vivo

A U87MG xenograft mouse model was used to assess the benefit of having a designed ankyrin repeat domain with binding specificity for VEGF-A combined with a designed ankyrin repeat domain with binding specificity for HGF compared to having them separate. Protein #134 consisting of SEQ ID NO: 134 (comprising two designed ankyrin repeat

domains with binding specificity for serum albumin each consisting of amino acids of SEQ ID NO: 50, comprising one designed ankyrin repeat domain with binding specificity for VEGF-A consisting of SEQ ID NO: 18, and comprising one designed ankyrin repeat domain with binding specificity for HGF consisting of amino acids of SEQ ID NO: 26),  
5 Protein #61 consisting of SEQ ID NO: 61 (comprising a designed ankyrin repeat domain with binding specificity for serum albumin, consisting of amino acids of SEQ ID NO: 50, and comprising a designed ankyrin repeat domain with binding specificity for VEGF-A, consisting of amino acids of SEQ ID NO: 18) and additionally having SEQ ID NO: 1 at the N-terminus, or Protein #60 consisting of SEQ ID NO: 60 (comprising a designed ankyrin  
10 repeat domain with binding specificity for serum albumin, consisting of amino acids of SEQ ID NO: 50, and comprising a designed ankyrin repeat domain with binding specificity for HGF, consisting of amino acids of SEQ ID NO: 26) and additionally having SEQ ID NO: 1 at the N-terminus, were prepared as described in Example 4. For the in vivo analysis,  $2 \times 10^6$  U87MG cells per mouse were implanted subcutaneously into the right  
15 flank of female NMRI nu/nu mice (Harlan) and the mice were grouped in groups with equivalent tumor volumes each. On day 29 and day 32, mice were treated with PBS or 4 mg/kg protein i.v. On day 35, tumors were harvested and cryo-frozen. On every day of treatment the tumor volume of each tumor was measured using the formula: Volume = (width)<sup>2</sup> x length/2. Body weight measurements indicated no significant difference  
20 between the four treatment groups. Tumor cross-sections were then stained using an antibody for Ki67 (ab66155; Abcam, U.K.) for the quantification of proliferation, or using an antibody for CD-31 (BD550274; BD Biosciences, USA) for the quantification of angiogenesis using standard IHC methods. The percentage of proliferative cells and the percentage of mean vascular areas were measured using the software Image J. The  
25 results are shown in Figure 10a. Compared to the PBS, Protein #60 as well as Protein #61 inhibit proliferation, and Protein #60 (slightly) as well as Protein #61 inhibit angiogenesis, as expected. The combination of the two, resulting in Protein #134, however, leads to improved inhibition of both proliferation and angiogenesis. This indicates that the combination of anti-VEGF-A and anti-HGF activity is key for good  
30 efficacy.

Protein #134 was further characterized in two patient-derived xenograft mouse models, a gastric cancer model and a renal cancer model. Patient-derived tumor xenograft mouse models are well known to the person skilled in the art. Protein #134 was prepared as described in Example 4.

For the renal cancer patient-derived xenograft mouse model, renal cell cancer specimens from surgical specimens were implanted s.c. in NMRI nu/nu mice and passaged three to five times until establishment of stable growth patterns. After removal from donor mice, tumors were cut into fragments of 4-5 mm diameter, which were implanted s.c. in NMRI  
5 nu/nu mice. Upon obvious onset of solid tumor growth, mice were randomized to groups of three animals each, and test articles were administered as follows to one animal group each: PBS was given i.v. at 10 ml/kg three times weekly for three times; Protein #134 was given i.v. at 4 mg/kg three times weekly for three times; sorafenib was given p.o. at 200 mg/kg daily for 21 days. Tumor volumes were assessed as described above at the  
10 day of treatment start as well as on days 3, 7, 10, 14, 18, and 21. The results are shown in Figure 10b. In this model, Protein #134 is more efficacious than sorafenib, today's standard of care for the treatment of renal cell carcinoma.

For the gastric cancer patient-derived xenograft mouse model, gastric cancer specimens from surgical specimens were implanted to in NMRI nu/nu mice and passaged three to  
15 five times until establishment of stable growth patterns. After removal from donor mice, tumors were cut into fragments of 4-5 mm diameter, which were implanted s.c. in NMRI nu/nu mice. Upon obvious onset of solid tumor growth, mice were randomized to groups of eight animals each, and test articles were administered as follows to one animal group each: PBS was given i.v. at 10 ml/kg on days 0, 3, 6, 9, 12, 15, and 18; Protein #134 was  
20 given i.v. at 4 ml/kg on days 0, 3, 6, 9, 12, 15, and 18; Paclitaxel was given i.v. at 15 mg/kg on days 0, 7, and 14; Protein #134 plus paclitaxel were given i.v. at 4 mg/kg on days 0, 3, 6, 9, 12, 15, and 18, and i.v. at 15 mg/kg on days 0, 7, and 14. Tumor volumes were assessed as described above at the day of treatment start as well as on days 2, 6, 13, 16, and 20. The results are shown in Figure 10c. In this model, Protein #134 was at  
25 least as efficacious as paclitaxel, and a combination of the two was significantly more efficacious than the individual components.

CLAIMS

1. A designed ankyrin repeat domain with binding specificity for serum albumin comprising the amino acid sequence of SEQ ID NO: 50.
- 5
2. A recombinant binding protein comprising at least two designed ankyrin repeat domains of claim 1.
3. The recombinant binding protein of claim 2, wherein said recombinant binding protein comprises a first, a second, a third, and a fourth designed ankyrin repeat domain, wherein said first designed ankyrin repeat domain has binding specificity for VEGF-A, and wherein said second designed ankyrin repeat domain has binding specificity for HGF, and wherein said third and fourth designed ankyrin repeat domains each have binding specificity for serum albumin.
- 10
4. The recombinant binding protein of claim 3, wherein said recombinant binding protein comprises an amino acid sequence that has at least 90% amino acid sequence identity with the amino acid sequence of SEQ ID NO: 134.
- 15
5. The recombinant binding protein of claim 3, wherein said recombinant binding protein comprises the amino acid sequence consisting of the amino acid sequence of SEQ ID NO: 134.
- 20
6. The recombinant binding protein of claim 3, wherein said recombinant binding protein consists of the amino acid sequence of SEQ ID NO: 134.
- 25
7. The recombinant binding protein of claim 3, wherein said first, second, third and fourth designed ankyrin repeat domains are in the order third-second-first-fourth from N terminus to C terminus.
- 30

8. The recombinant binding protein of claim 3, wherein said first designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 12 to 21 and amino acid sequences in which up to 10 amino acids of SEQ ID NOs: 12 to 21 are exchanged by any amino acid, and wherein said  
5 second designed ankyrin repeat domain comprises an amino acid sequence selected from the group consisting of amino acid sequences SEQ ID NOs: 23 to 37 and amino acid sequences in which up to 10 amino acids of SEQ ID NOs: 23 to 37 are exchanged by any amino acid, and wherein said third and fourth designed ankyrin repeat domains each comprise amino acid sequence SEQ ID NO: 50, and wherein said designed ankyrin  
10 repeat domains are linked by polypeptide linkers comprising amino acid sequences selected from the group consisting of amino acid sequences SEQ ID NOs: 2 to 9 and amino acid sequences in which up to 4 amino acids of SEQ ID NOs: 2 to 9 are exchanged by any amino acid.
- 15 9. The recombinant binding protein of claim 3, wherein said first designed ankyrin repeat domain comprises amino acid sequences SEQ ID NO: 18, and wherein said second designed ankyrin repeat domain comprises amino acid sequence SEQ ID NO: 26, and wherein said third and fourth designed ankyrin repeat domains each comprise amino acid sequence SEQ ID NO: 50, and wherein said designed ankyrin repeat domains are linked  
20 by polypeptide linkers consisting of amino acid sequence SEQ ID NO: 9.
10. A recombinant binding protein comprising an amino acid sequence that has at least 90% amino acid sequence identity with the amino acid sequence of SEQ ID NO: 134.
- 25 11. The recombinant binding protein of claim 10, wherein said recombinant binding protein comprises the amino acid sequence of SEQ ID NO: 134.
12. The recombinant binding protein of any one of claims 10 to 11, consisting of the amino acid sequence of SEQ ID NO: 134.
- 30 13. A nucleic acid encoding the amino acid sequence of a designed ankyrin repeat domain or a recombinant binding protein of any one of claims 1 to 12.

14. A pharmaceutical composition comprising a recombinant binding protein and/or a designed ankyrin repeat domain of any one of claims 1 to 12 or the nucleic acid of claim 13, and optionally a pharmaceutical acceptable carrier and/or diluent.
- 5 15. The pharmaceutical composition of claim 14 for use in the treatment of a disease.

Figure 1






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6a




Figure 6b




Figure 6c




Figure 7a




Figure 7b




Figure 7c




Figure 8




Figure 9a




Figure 9b




Figure 9c




Figure 10a






Figure 10b





Figure 10c



INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2016/057272

A. CLASSIFICATION OF SUBJECT MATTER  
 INV. C07K14/475 A61K38/00 C07K16/22 C07K16/28 C07K14/765  
 C07K14/47  
 ADD.  
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
 Minimum documentation searched (classification system followed by classification symbols)  
 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
 EPO-Internal, BIOSIS, Sequence Search, EMBASE, FSTA, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2014/191574 A1 (MOLECULAR PARTNERS AG [CH]) 4 December 2014 (2014-12-04) cited in the application	1-15
Y	the whole document pages 33-35; examples; sequences 51, 48 -& DATABASE Geneseq [Online]  15 January 2015 (2015-01-15), "DARPin preparation related ankyrin repeat domain, SEQ 51.", XP002744973, retrieved from EBI accession no. GSP:BBQ86961 Database accession no. BBQ86961 sequence  -/--	1-15

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  25 May 2016	Date of mailing of the international search report  09/06/2016
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Madruga, Jaime
--	--

## INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2016/057272

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	-& DATABASE Geneseq [Online]  15 January 2015 (2015-01-15), "DARPin preparation related ankyrin repeat domain, SEQ 48.", XP002744974, retrieved from EBI accession no. GSP:BBQ86958 Database accession no. BBQ86958 sequence  -----	
Y	WO 2010/060748 A1 (MOLECULAR PARTNERS AG [CH]; BINZ HANS KASPAR [CH]; FORRER PATRIK [CH];) 3 June 2010 (2010-06-03) cited in the application the whole document page 26; claims; sequence 18 -& DATABASE Geneseq [Online]  5 August 2010 (2010-08-05), "Ankyrin repeat domain protein, SEQ ID 18.", XP002744975, retrieved from EBI accession no. GSP:AYD11297 Database accession no. AYD11297 sequence  -----	1-15
X	WO 2012/069654 A1 (MOLECULAR PARTNERS AG [CH]; STEINER DANIEL [CH]; BINZ HANS KASPAR [CH]) 31 May 2012 (2012-05-31) cited in the application the whole document page 32; claims; examples; sequences 46, 24 -& DATABASE Geneseq [Online]	1,2,10, 13-15
Y	19 July 2012 (2012-07-19), "Designed ankyrin repeat protein (DARPin) protein DARPin #46, SEQ:46.", XP002744976, retrieved from EBI accession no. GSP:AZW76566 Database accession no. AZW76566 sequence -& DATABASE Geneseq [Online]  19 July 2012 (2012-07-19), "Designed ankyrin repeat protein (DARPin) protein DARPin #24, SEQ:24.", XP002744977, retrieved from EBI accession no. GSP:AZW76544 Database accession no. AZW76544 sequence  -----	1-15
		-/--

## INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2016/057272

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>Ulrike Fiedler, Michael Tobias Stumpp, Andreas Cornelius, Savira Ekawardhani, Keith Martin Dawson, Pat Gilboy, Ignacio Dolado: "Antitumor activity of MP0250, a bispecific VEGF- and HGF-targeting darpin, in patient-derived xenograft models.", J Clin Oncol 32:5s, 2014 (suppl; abstr 11039) -2014 ASCO Annual Meeting J CLIN ONCOL, 11039, 30 May 2014 (2014-05-30), XP002744986, Retrieved from the Internet: URL:http://meetinglibrary.asco.org/print/1739339 [retrieved on 2015-09-24] the whole document</p>	1-15
Y	<p>-----            WO 2012/069557 A1 (GLAXO GROUP LTD [GB]; GRIGGS JEREMY [GB]; PARMAR SHAH RADHA [GB]; STEW) 31 May 2012 (2012-05-31) the whole document pages 30-31; claims; examples</p>	1-15
A	<p>-----            J. HOPP ET AL: "The effects of affinity and valency of an albumin-binding domain (ABD) on the half-life of a single-chain diabody-ABD fusion protein", PROTEIN ENGINEERING DESIGN AND SELECTION, vol. 23, no. 11, 3 September 2010 (2010-09-03), pages 827-834, XP055054496, ISSN: 1741-0126, DOI: 10.1093/protein/gzq058 cited in the application the whole document</p>	1-15
A	<p>-----            PETER-CHRISTIAN KLÖHN ET AL: "IBC's 23rd Annual Antibody Engineering, 10th Annual Antibody Therapeutics International Conferences and the 2012 Annual Meeting of The Antibody Society: December 3-6, 2012, San Diego, CA", MABS, vol. 5, no. 2, 1 March 2013 (2013-03-01), pages 178-201, XP055137110, ISSN: 1942-0862, DOI: 10.4161/mabs.23655 page 195</p> <p>-----            -/--</p>	1-15

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2016/057272

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	SKRLEC KATJA ET AL: "Non-immunoglobulin scaffolds: a focus on their targets", TRENDS IN BIOTECHNOLOGY, vol. 33, no. 7, 27 April 2015 (2015-04-27) , pages 408-418, XP029175810, ISSN: 0167-7799, DOI: 10.1016/J.TIBTECH.2015.03.012 the whole document page 412 - page 413; table 2 -----	1-15

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2016/057272

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2014191574	A1	04-12-2014	CN 105209483 A	30-12-2015
			EP 3004152 A1	13-04-2016
			WO 2014191574 A1	04-12-2014
-----				
WO 2010060748	A1	03-06-2010	AU 2009319204 A1	03-06-2010
			BR PI0921469 A2	12-01-2016
			CA 2742241 A1	03-06-2010
			CN 102272148 A	07-12-2011
			EP 2358746 A1	24-08-2011
			JP 2012507271 A	29-03-2012
			JP 2015133978 A	27-07-2015
			KR 20110082528 A	19-07-2011
			NZ 592591 A	27-04-2012
			RU 2011122201 A	10-12-2012
			US 2011207668 A1	25-08-2011
			US 2015344539 A1	03-12-2015
			WO 2010060748 A1	03-06-2010
-----				
WO 2012069654	A1	31-05-2012	AU 2011333665 A1	06-06-2013
			AU 2011333666 A1	06-06-2013
			CA 2818969 A1	31-05-2012
			CA 2818990 A1	31-05-2012
			CN 103403024 A	20-11-2013
			CN 103459415 A	18-12-2013
			EP 2643348 A2	02-10-2013
			EP 2643349 A1	02-10-2013
			JP 2014501509 A	23-01-2014
			JP 2014501510 A	23-01-2014
			KR 20140002657 A	08-01-2014
			KR 20140032356 A	14-03-2014
			RU 2013128895 A	10-01-2015
			RU 2013128896 A	10-01-2015
			US 2013244940 A1	19-09-2013
			US 2013296221 A1	07-11-2013
			US 2016075767 A1	17-03-2016
			WO 2012069654 A1	31-05-2012
			WO 2012069655 A2	31-05-2012
-----				
WO 2012069557	A1	31-05-2012	AU 2011333738 A1	11-07-2013
			CA 2818621 A1	31-05-2012
			CN 103459422 A	18-12-2013
			EA 201390611 A1	30-01-2014
			EP 2643351 A1	02-10-2013
			EP 2853542 A1	01-04-2015
			JP 2014501725 A	23-01-2014
			KR 20130121886 A	06-11-2013
			SG 190362 A1	28-06-2013
			US 2013236467 A1	12-09-2013
			WO 2012069557 A1	31-05-2012
-----				