



(51) International Patent Classification:

A61K 33/32 (2006.01) A61K 39/395 (2006.01)
A61K 39/00 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2018/027588

(22) International Filing Date:

13 April 2018 (13.04.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/485,061 13 April 2017 (13.04.2017) US
62/572,377 13 October 2017 (13.10.2017) US

(71) Applicant: GALERA LABS, LLC [US/US]; 1100 Corporate Square Drive, Suite 223, Creve Coeur, Missouri 63132 (US).


(72) Inventors: BEARDSLEY, Robert A.; 6935 Waterman Ave, University City, Missouri 63130 (US). KEENE, Jeffery L.; 8100 Leafland Court, St. Louis, Missouri 63123 (US). RILEY, Dennia P.; 1722 Claymont Estates Drive, Chesterfield, Missouri 63107 (US).

(74) Agent: COTTON, Abigail; c/o Bryan Cave LLP, 211 North Broadway, Suite 3600, St. Louis, Missouri 63102 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: COMBINATION CANCER IMMUNOTHERAPY WITH PENTAAZA MACROCYCLIC RING COMPLEX




Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

COMBINATION CANCER IMMUNOTHERAPY WITH PENTAАЗA MACROCYCLIC RING COMPLEX


5 [0001] The present disclosure generally relates to combination therapies for cancer treatment, including administration of a pentaaza macrocyclic ring complex in combination with an immunotherapy treatment.

[0002] Transition metal-containing pentaaza macrocyclic ring complexes having the macrocyclic ring system corresponding to Formula A have been shown to be effective in a number of animal and cell models of human disease, as well as in
10 treatment of conditions afflicting human patients.





FORMULA A

For example, in a rodent model of colitis, one such compound, GC4403, has been reported to very significantly reduce the injury to the colon of rats subjected to an experimental model of colitis (see Cuzzocrea et al., *Europ. J. Pharmacol.*, 432, 79-89
15 (2001)).



GC4403 has also been reported to attenuate the radiation damage arising both in a clinically relevant hamster model of acute, radiation-induced oral mucositis (Murphy et al., *Clin. Can. Res.*, 14(13), 4292 (2008)), and lethal total body irradiation of adult mice
20 (Thompson et al., *Free Radical Res.*, 44(5), 529-40 (2010)). Similarly, another such compound, GC4419, has been shown to attenuate VEGFr inhibitor-induced pulmonary disease in a rat model (Tuder, et al., , *Am. J. Respir. Cell Mol. Biol.*, 29, 88-97 (2003)).

Additionally, another such compound, GC4401 has been shown to provide protective effects in animal models of septic shock (S. Cuzzocrea, et.al., *Crit. Care Med.*, 32(1), 157 (2004) and pancreatitis (S. Cuzzocrea, et.al., *Shock*, 22(3), 254-61 (2004)).



- 5 [0003] Certain of these compounds have also been shown to possess potent anti-inflammatory activity and prevent oxidative damage *in vivo*. For example, GC4403 has been reported to inhibit inflammation in a rat model of inflammation (Salvemini, et.al., *Science*, 286, 304 (1999)), and prevent joint disease in a rat model of collagen-induced arthritis (Salvemini et al., *Arthritis & Rheumatism*, 44(12), 2009-2021 (2001)).
- 10 Yet others of these compounds, MdPAM and MnBAM, have shown *in vivo* activity in the inhibition of colonic tissue injury and neutrophil accumulation into colonic tissue (Weiss et al., *The Journal of Biological Chemistry*, 271(42), 26149-26156 (1996)). In addition, these compounds have been reported to possess analgesic activity and to reduce inflammation and edema in the rat-paw carrageenan hyperalgesia model, see, e.g.,
- 15 U.S. Pat. No. 6,180,620.

- [0004] Compounds of this class have also been shown to be safe and effective in the prevention and treatment of disease in human subjects. For example, GC4419 has been shown to reduce oral mucositis in head-and-neck cancer patients undergoing chemoradiation therapy (Anderson, C., *Phase 1 Trial of Superoxide*
- 20 *Dismutase (SOD) Mimetic GC4419 to Reduce Chemoradiotherapy (CRT)-Induced Mucositis (OM) in Patients (pts) with Mouth or Oropharyngeal Carcinoma (OCC)*, Oral Mucositis Research Workshop, MASCC/ISOO Annual Meeting on Supportive Care in Cancer, Copenhagen, Denmark (June 25, 2015)).

- [0005] In addition, transition metal-containing pentaaza macrocyclic ring
- 25 complexes corresponding to this class have shown efficacy in the treatment of various cancers. For example, certain compounds corresponding to this class have been provided in combination with agents such as paclitaxel and gemcitabine to enhance

cancer therapies, such as in the treatment of colorectal cancer and lung cancer (non-small cell lung cancer) (see, e.g., U.S. Patent No. 9,998,893). The 4403 compound above has also been used for treatment in *in vivo* models of Meth A spindle cell squamous carcinoma and RENCA renal carcinoma (Samlowski et al., *Nature Medicine*, 9(6), 750-755 (2003), and has also been used for treatment in *in vivo* models of spindle-cell squamous carcinoma metastasis (Samlowski et al., *Madame Curie Bioscience Database (Internet)*, 230-249 (2006)). The 4419 compound above has also been used in combination with cancer therapies such as cisplatin and radiation therapy to enhance treatment in *in vivo* models (Sishc et al., poster for Radiation Research Society (2015)).

[0006] Various cancer immunotherapies have also been developed that recruit the immune system to attack cancer cells to provide treatment. For example, recent immunotherapies have included the administration of immune checkpoint inhibitors, which help the immune system bypass the “checks” that may otherwise inhibit full activation and/or attack of the immune system against cancer cells. The drug ipilimumab is an example of such an immune checkpoint inhibitor, and has been approved for treatment of melanoma (Cameron et al., *Ipilimumab; First Global Approval*, *Drugs* (2011) 71(8), 1093-1094).

[0007] However, a need remains for enhanced methods for cancer treatment that provide improved efficacy in the killing of cancer cells.

[0008] Briefly, therefore, aspects of the present disclosure are directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient prior to, concomitantly with, or after an inhibitor of immune response checkpoint inhibitor therapy for cancer, increasing the response of the tumors to the checkpoint inhibitor dose.

[0009] Another aspect of the present disclosure is directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient prior to, concomitantly with or after an adoptive T-cell transfer therapy for cancer, increasing the response of the tumors to the adoptive T-cell transfer treatment.


[0010] Another aspect of the present disclosure is directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient prior to, concomitantly with or after a therapeutic vaccine, increasing the response of the tumors to the therapeutic vaccine.

[0011] Another aspect of the present disclosure is directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient prior to, concomitantly with or after a immunologic treatment for cancer, including those comprised of a compound, a composition, a device, or a procedure, increasing the response of the tumors to the immunologic treatment.

[0012] Another aspect of the present disclosure is directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient suffering from a viral infection or other infectious disease, alone or in combination with one or more of an immune response checkpoint inhibitor, a T-cell transfer therapy, a therapeutic vaccine.

[0013] Another aspect of the present disclosure is directed to a method wherein a transition metal pentaaza-macrocyclic ring complex is administered to a patient for the purpose of increasing numbers of CD4+ or CD8+ T-cells, producing or increasing an immune response to a tumor or a viral infection.

[0014] Among the various aspects of the present disclosure, therefore, is method of treating a cancer in a mammalian subject afflicted with the cancer, the method including administering to the subject an immune checkpoint inhibitor, and administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after administration of the immune checkpoint inhibitor, to increase the response of the cancer to the immune checkpoint inhibitor:



[0015] wherein

[0016] M is Mn^{2+} or Mn^{3+} ;

[0017] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

5 of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

[0018] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or
10 heterocycle having 3 to 20 ring carbon atoms;

[0019] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[0020] W, together with the nitrogen of the macrocycle and the carbon atoms
15 of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of
20 the heterocycle and the macrocycle are absent;

[0021] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;


[0022] Z is a counterion;

25 [0023] n is an integer from 0 to 3; and

[0024] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[0025] According to yet another aspect of the present disclosure, a method of treating a cancer in a mammalian subject afflicted with the cancer includes
30 administering to the subject an adoptive T-cell transfer therapy, and administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below,

prior to, concomitantly with, or after the adoptive T-cell transfer therapy, to increase the response of the cancer to the adoptive T-cell transfer therapy,



[0026]

[0027] wherein

5 [0028] M is Mn^{2+} or Mn^{3+} ;

[0029] $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

10

[0030] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

15 [0031] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[0032] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle

20

and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;


[0033] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding
5 anion thereof;

[0034] Z is a counterion;

[0035] n is an integer from 0 to 3; and

[0036] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

10 [0037] According to yet another aspect of the disclosure, a method of treating a cancer in a mammalian subject afflicted with the cancer includes administering to the subject a cancer vaccine, and administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after
15 administration of the cancer vaccine, to increase the response of the cancer to the cancer vaccine,



[0038] wherein

20 [0039] M is Mn^{2+} or Mn^{3+} ;

[0040] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino

acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

5 [0041] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

 [0042] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or
10 heterocycle having 3 to 20 ring carbon atoms;

 [0043] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic
15 heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

 [0044] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding
20 anion thereof;


 [0045] Z is a counterion;

 [0046] n is an integer from 0 to 3; and

 [0047] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

25 [0048] According to yet another aspect of the disclosure, a method of treating a viral infection in a mammalian subject in need thereof includes administering to the subject at least one of an immune checkpoint inhibitor, an adoptive T-cell transfer therapy, and a cancer vaccine, and administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after
30 administration of the at least one immune checkpoint inhibitor, adoptive T-cell transfer therapy, and cancer vaccine, to increase the effectiveness of the at least one immune

checkpoint, adoptive T-cell transfer therapy, and cancer vaccine in treating the viral infection,



[0049] (I)

5 [0050] wherein

[0051] M is Mn²⁺ or Mn³⁺;

[0052] R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting
 10 of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or alkyl;

[0053] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or
 15 heterocycle having 3 to 20 ring carbon atoms;

[0054] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[0055] W, together with the nitrogen of the macrocycle and the carbon atoms
 20 of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle

and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;


[0056] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[0057] Z is a counterion;

[0058] n is an integer from 0 to 3; and

[0059] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[0060] According to yet another aspect of the present disclosure, a kit for treating cancer includes at least one of an immune checkpoint inhibitor, T-cells for an adoptive T-cell transfer therapy, and a cancer vaccine, and a pentaaza macrocyclic ring complex according to formula (I),



[0061] (I)

[0062] wherein

[0063] M is Mn^{2+} or Mn^{3+} ;

[0064] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R$

₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or alkyl;

5 [0065] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[0066] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [0067] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of
15 the heterocycle and the macrocycle are absent;

[0068] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[0069] Z is a counterion;

20 [0070] n is an integer from 0 to 3; and

[0071] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[0072] Other objects and features will be in part apparent and in part pointed out hereinafter.

25 Brief Description of the Drawings

[0073] Figure 1 shows median tumor volumes over a duration of treatment in a colon 26 cancer model using GC4419 and anti-PD1.

[0074] Figure 2 shows mean tumor volumes over a duration of treatment in a colon 26 cancer model using GC4419 and anti-PD1.

[0075] Figure 3A shows median tumor volumes during treatment in a CT26 cancer model using GC4419 and anti-PDL1 through day 16 post-implantation.

[0076] Figure 3B depicts intratumoral leukocytes assessed by flow cytometry for treatment in a CT26 cancer model using GC4419 and anti-PDL1.

5 [0077] Figure 4A shows average tumor volumes over a duration of treatment in a 4T1 breast metastatic cancer model with GC4419 in radiation therapy.

[0078] Figure 4B shows average tumor volumes over a duration of treatment in a 4T1 metastatic breast cancer model with radiation therapy, GC4419 and anti-CTLA4.

10 [0079] Figure 4C shows a number of surface lung metastases for treatment in a 4T1 metastatic breast cancer model with radiation therapy, GC4419 and anti-CTLA4.

[0080] Figure 5A shows mean tumor volumes over a duration of treatment in a 4T1 metastatic breast cancer model with GC4419 and anti-CTLA4.

15 [0081] Figure 5B shows normalized mean tumor volumes for treatment in a 4T1 metastatic breast cancer model with GC4419 and anti-CTLA4, where the GC4419 start date is day 13 after the initial anti-CTLA4 treatment.

[0082] Figures 5C-5D show mean tumor volumes over a duration of treatment in a 4T1 metastatic breast cancer model with GC4419 and anti-CTLA4.

20 [0083] Figure 6A shows the sensitizing effect of GC4419 on Lewis Lung Carcinoma tumors to ionizing radiation

[0084] Figure 6B shows changes in tumor infiltrating lymphocyte populations in Lewis Lung Carcinoma tumors post ionizing radiation and GC4419.

[0085] Figure 7 shows mean tumor volumes over a duration of treatment in a 4T1 metastatic breast cancer model with GC4419 and anti-PD-1.

25 [0086] Figures 8A-8E show tumor volumes for an abscopal study.

[0087] Figure 9 shows average tumor volumes over a duration of treatment with GC4419 and anti-PDL-1.

[0088] Figures 10A-10E show individual tumor volumes over a duration of treatment with GC4419 and anti-PDL-1

30 Abbreviations and Definitions

[0089] The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

5 [0090] "Acyl" means a -COR moiety where R is alkyl, haloalkyl, optionally substituted aryl, or optionally substituted heteroaryl as defined herein, e.g., acetyl, trifluoroacetyl, benzoyl, and the like.

[0091] "Acyloxy" means a -OCOR moiety where R is alkyl, haloalkyl, optionally substituted aryl, or optionally substituted heteroaryl as defined herein, e.g.,
10 acetyl, trifluoroacetyl, benzoyl, and the like.

[0092] "Alkoxy" means a -OR moiety where R is alkyl as defined above, e.g., methoxy, ethoxy, propoxy, or 2-propoxy, n-, iso-, or tert-butoxy, and the like.

[0093] "Alkyl" means a linear saturated monovalent hydrocarbon moiety such as of one to six carbon atoms, or a branched saturated monovalent hydrocarbon
15 moiety, such as of three to six carbon atoms, e.g., C₁-C₆ alkyl groups such as methyl, ethyl, propyl, 2-propyl, butyl (including all isomeric forms), pentyl (including all isomeric forms), and the like.

[0094] Moreover, unless otherwise indicated, the term "alkyl" as used herein is intended to include both "unsubstituted alkyls" and "substituted alkyls," the latter of
20 which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Indeed, unless otherwise indicated, all groups recited herein are intended to include both substituted and unsubstituted options.

[0095] The term "C_{x-y}" when used in conjunction with a chemical moiety, such as alkyl and aralkyl, is meant to include groups that contain from x to y carbons in the
25 chain. For example, the term C_{x-y} alkyl refers to substituted or unsubstituted saturated hydrocarbon groups, including straight chain alkyl and branched chain alkyl groups that contain from x to y carbon atoms in the chain.

[0096] "Alkylene" means a linear saturated divalent hydrocarbon moiety, such as of one to six carbon atoms, or a branched saturated divalent hydrocarbon moiety,
30 such as of three to six carbon atoms, unless otherwise stated, e.g., methylene, ethylene, propylene, 1-methylpropylene, 2-methylpropylene, butylene, pentylene, and the like.

[0097] "Alkenyl" a linear unsaturated monovalent hydrocarbon moiety, such as of two to six carbon atoms, or a branched saturated monovalent hydrocarbon moiety, such as of three to six carbon atoms, e.g., ethenyl (vinyl), propenyl, 2-propenyl, butenyl (including all isomeric forms), pentenyl (including all isomeric forms), and the like.

5 [0098] "Alkaryl" means a monovalent moiety derived from an aryl moiety by replacing one or more hydrogen atoms with an alkyl group.

[0099] "Alkenylcycloalkenyl" means a monovalent moiety derived from an alkenyl moiety by replacing one or more hydrogen atoms with a cycloalkenyl group.

10 [00100] "Alkenylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with an alkenyl group.

[00101] "Alkylcycloalkenyl" means a monovalent moiety derived from a cycloalkenyl moiety by replacing one or more hydrogen atoms with an alkyl group.

[00102] "Alkylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with an alkyl group.

15 [00103] "Alkynyl" means a linear unsaturated monovalent hydrocarbon moiety, such of two to six carbon atoms, or a branched saturated monovalent hydrocarbon moiety, such as of three to six carbon atoms, e.g., ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like.

20 [00104] "Alkoxy" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a hydroxy group.

[00105] "Amino" means a $-NR^aR^b$ group where R^a and R^b are independently hydrogen, alkyl or aryl.

[00106] "Aralkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an aryl group.

25 [00107] "Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon moiety of 6 to 10 ring atoms e.g., phenyl or naphthyl.

[00108] "Cycle" means a carbocyclic saturated monovalent hydrocarbon moiety of three to ten carbon atoms.

[00109] "Cycloalkyl" means a cyclic saturated monovalent hydrocarbon moiety of three to ten carbon atoms, e.g., cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, and the like.

[00110] "Cycloalkylalkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a cycloalkyl group, e.g., cyclopropylmethyl, cyclobutylmethyl, cyclopentylethyl, or cyclohexylethyl, and the like.

[00111] "Cycloalkylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with a cycloalkyl group.

[00112] "Cycloalkenyl" means a cyclic monounsaturated monovalent hydrocarbon moiety of three to ten carbon atoms, e.g., cyclopropenyl, cyclobutenyl, cyclopentenyl, or cyclohexenyl, and the like.

[00113] "Cycloalkenylalkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a cycloalkenyl group, e.g., cyclopropenylmethyl, cyclobutenylmethyl, cyclopentenylethyl, or cyclohexenylethyl, and the like.

[00114] "Ether" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an alkoxy group.

[00115] "Halo" means fluoro, chloro, bromo, or iodo, preferably fluoro or chloro.

[00116] "Heterocycle" or "heterocyclyl" means a saturated or unsaturated monovalent monocyclic group of 4 to 8 ring atoms in which one or two ring atoms are heteroatom selected from N, O, or S(O)_n, where n is an integer from 0 to 2, the remaining ring atoms being C. The heterocyclyl ring is optionally fused to a (one) aryl or heteroaryl ring as defined herein provided the aryl and heteroaryl rings are monocyclic. The heterocyclyl ring fused to monocyclic aryl or heteroaryl ring is also referred to in this Application as "bicyclic heterocyclyl" ring. Additionally, one or two ring carbon atoms in the heterocyclyl ring can optionally be replaced by a -CO- group. More specifically the term heterocyclyl includes, but is not limited to, pyrrolidino, piperidino, homopiperidino, 2-oxopyrrolidinyl, 2-oxopiperidinyl, morpholino, piperazino, tetrahydropyranyl, thiomorpholino, and the like. When the heterocyclyl ring is unsaturated it can contain one or two ring double bonds provided that the ring is not aromatic. When the heterocyclyl group is a saturated ring and is not fused to aryl or heteroaryl ring as stated above, it is also referred to herein as saturated monocyclic heterocyclyl.

[00117] "Heteroaryl" means a monovalent monocyclic or bicyclic aromatic moiety of 5 to 10 ring atoms where one or more, preferably one, two, or three, ring atoms are heteroatom selected from N, O, or S, the remaining ring atoms being carbon. Representative examples include, but are not limited to, pyrrolyl, pyrazolyl, thienyl, thiazolyl, imidazolyl, furanyl, indolyl, isoindolyl, oxazolyl, isoxazolyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, quinolinyl, isoquinolinyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl, tetrazolyl, and the like.

[00118] "Nitro" means --NO_2 .

[00119] "Organosulfur" means a monovalent moiety a --SR group where R is hydrogen, alkyl or aryl.

[00120] "Substituted alkyl," "substituted cycle," "substituted phenyl," "substituted aryl," "substituted heterocycle," and "substituted nitrogen heterocycles" means an alkyl, cycle, aryl, phenyl, heterocycle or nitrogen-containing heterocycle, respectively, optionally substituted with one, two, or three substituents, such as those independently selected from alkyl, alkoxy, alkoxyalkyl, halo, hydroxy, hydroxyalkyl, or organosulfur.

[00121] "Thioether" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an --SR group wherein R is alkyl.

[00122] As used herein, (i) the compound referred to herein and in the Figures as compound 401, 4401 or GC4401 is a reference to the same compound, (ii) the compound referred to herein and in the Figures as compound 403, 4403 or GC4403 is a reference to the same compound, (iii) the compound referred to herein and in the Figures as compound 419, 4419 or GC4419 is a reference to the same compound, and (iv) the compound referred to herein and in the Figures as compound 444, 4444 or GC4444 is a reference to the same compound.

Detailed Description


[00123] Aspects of the present disclosure are directed to the treatment of cancer by administration of a pentaaza macrocyclic ring complex according to Formula (I), described below, in combination with an immunotherapeutic agent, to a subject suffering from cancer, to enhance response of the cancer to the immunotherapeutic agent.

[00124] In general, the immunotherapeutic agent may be an agent that is capable of stimulating or otherwise facilitating attack of the immune system on cancer cells or other cells. Examples of suitable immunotherapeutic agents can include, for example, immune checkpoint inhibitors, adoptive T-cell transfer therapy materials, and cancer vaccines. By providing the pentaaza macrocyclic ring complex in combination with the immunotherapeutic agent, it has been discovered that immune system activity can be enhanced to impart improved treatment of cancer in a subject suffering therefrom.

[00125] Accordingly, in one embodiment, aspects of the present disclosure comprise a method of treating a cancer in a mammalian subject by administering an immune checkpoint inhibitor, and a pentaaza macrocyclic ring complex corresponding to the formula (I) below. In yet another embodiment, aspects of the present disclosure comprise a method of treating a cancer in a mammalian subject by administering an adoptive T-cell transfer therapy, and a pentaaza macrocyclic ring complex corresponding to formula (I) below. In yet another embodiment, aspects of the present disclosure comprise a method of treating cancer in a mammalian subject by administering a cancer vaccine, and a pentaaza macrocyclic ring complex corresponding to formula (i) below. In yet another embodiment, a method of treatment of a viral infection by any of a checkpoint inhibitor, adoptive T-cell transfer, and cancer vaccine, can be enhanced by providing the pentaaza macrocyclic ring complex in combination with the treatment. Accordingly, the combination therapy can impart benefits in the treatment of cancer and viral infections, such as by facilitating the immunotherapeutic effects of the immunotherapeutic agent being provided as a part of the combination.

Transition Metal Pentaaza Macrocyclic Ring Complex

[00126] In one embodiment, the pentaaza macrocyclic ring complex corresponds to the complex of Formula (I):



wherein

[00127] M is Mn^{2+} or Mn^{3+} ;

[00128] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

[00129] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00130] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00131] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

[00132] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00133] Z is a counterion;

5 [00134] n is an integer from 0 to 3; and

[00135] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00136] As noted above in connection with the pentaaza macrocyclic ring complex of Formula (I), M is Mn^{2+} or Mn^{3+} . In one particular embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I), M is Mn^{2+} . In another particular embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I), M is Mn^{3+} .

[00137] In the embodiments in which one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are hydrocarbyl, for example, suitable hydrocarbyl moieties include, but are not limited to alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and aralkyl. In one embodiment, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heterocyclyl. More preferably in this embodiment, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen or lower alkyl (e.g., C_1 - C_6 alkyl, more typically C_1 - C_4 alkyl). Thus, for example, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} may be independently hydrogen, methyl, ethyl, propyl, or butyl (straight, branched, or cyclic). In one preferred embodiment, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen or methyl.

[00138] In one preferred embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I), R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen and one of R_6 and R'_6 is hydrogen and the other of R_6 and R'_6 is methyl. In this embodiment, for example, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} may each be hydrogen while R'_6 is methyl. Alternatively, for example, R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} may each be hydrogen while R_6 is methyl. In another preferred embodiment in which the pentaaza macrocyclic ring

complex corresponds to Formula (I), R₁, R₃, R₄, R₅, R'₅, R'₆, R₇, R₈, and R₁₀ are each hydrogen, one of R₂ and R'₂ is hydrogen and the other of R₂ and R'₂ is methyl, and one of R₉ and R'₉ is hydrogen and the other of R₉ and R'₉ is methyl. In this embodiment, for example, R₁, R'₂, R₃, R₄, R₅, R'₅, R₇, R₈, R₉, and R₁₀ may each be hydrogen while R₂ and R'₉ are methyl. Alternatively, for example, R₁, R₂, R₃, R₄, R₅, R'₅, R₇, R₈, R'₉, and R₁₀ may each be hydrogen while R'₂ and R₉ are methyl. In another embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I), R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each hydrogen.

[00139] In certain embodiments the U and V moieties are independently substituted or unsubstituted fused cycloalkyl moieties having 3 to 20 ring carbon atoms, more preferably 4 to 10 ring carbon atoms. In a particular embodiment, the U and V moieties are each trans-cyclohexanyl fused rings.

[00140] In certain embodiments the W moiety is a substituted or unsubstituted fused heteroaromatic moiety. In a particular embodiment, the W moiety is a substituted or unsubstituted fused pyridino moiety. Where W is a substituted fused pyridino moiety, for example, the W moiety is typically substituted with a hydrocarbyl or substituted hydrocarbyl moiety (e.g., alkyl, substituted alkyl) at the ring carbon atom positioned para to the nitrogen atom of the heterocycle. In a one preferred embodiment, the W moiety is an unsubstituted fused pyridino moiety.

[00141] As noted above, X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof (for example benzoic acid or benzoate anion, phenol or phenoxide anion, alcohol or alkoxide anion). For example, X and Y may be selected from the group consisting of halo, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite,

bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof, among other possibilities.

In one embodiment, X and Y if present, are independently selected from the group consisting of halo, nitrate, and bicarbonate ligands. For example, in this embodiment, X and Y, if present, are halo ligands, such as chloro ligands.

[00142] Furthermore, in one embodiment X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

[00143] each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

[00144] each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

[00145] each X_4 is independently hydrogen or together with X_3 is $(=O)$.

[00146] In yet another embodiment, X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or X and Y are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} .

[00147] In the pentaaza macrocyclic ring complex corresponding to Formula (I), Z is a counterion (e.g., a charge-neutralizing anion), wherein n is an integer from 0 to 3. In general, Z may correspond to counterions of the moieties recited above in connection for X and Y.

5 [00148] In combination, among certain preferred embodiments are pentaaza macrocyclic ring complexes corresponding to Formula (I) wherein

[00149] M is Mn^{2+} or Mn^{3+} ;

[00150] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen or lower alkyl;

10 [00151] U and V are each trans-cyclohexanyl fused rings;


[00152] W is a substituted or unsubstituted fused pyridino moiety;

[00153] X and Y are ligands; and

[00154] Z, if present, is a charge-neutralizing anion.

[00155] More preferably in these embodiments, M is Mn^{2+} ; R_1 , R_2 , R'_2 , R_3 , R_4 ,
 15 R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen or methyl; U and V are each trans-cyclohexanyl fused rings; W is an unsubstituted fused pyridino moiety; and X and Y are independently halo ligands (e.g., fluoro, chloro, bromo, iodo). Z, if present, may be a halide anion (e.g., fluoride, chloride, bromide, iodide).

[00156] In yet another embodiment, the pentaaza macrocyclic ring complex is
 20 represented by formula (II) below:




[00157]

[00158] wherein

[00159] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

5 [00160] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein
10 R_{11} and R_{12} are independently hydrogen or alkyl.

[00161] Furthermore, in one embodiment, the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):









15 [00162] wherein


[00163] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

[00164] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$
20


, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or alkyl.

[00165] In yet another embodiment, the pentaaza macrocyclic ring complex is a compound represented by a formula selected from the group consisting of formulae (V)-5 (XVI):









(XI)



(XII)






(XVI)

5 [00166] In one embodiment, X and Y in any of the formulas herein are independently selected from the group consisting of fluoro, chloro, bromo and iodo anions. In yet another embodiment, X and Y in any of the formulas herein are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates. In yet another embodiment, X and Y in any of
10 the formulas herein are independently amino acids.

[00167] In one embodiment, the pentaaza macrocyclic ring complex has the following Formula (IA):



(IA)

wherein

15 [00168] M is Mn^{2+} or Mn^{3+} ;

[00169] R_{1A} , R_{1B} , R_2 , R_3 , R_{4A} , R_{4B} , R_5 , R_6 , R_{7A} , R_{7B} , R_8 , R_9 , R_{10A} , and R_{10B} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino

acid side chain moiety, or a moiety independently

selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-C(=O)NR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(=O)(OR_{11})(OR_{12})$, $-P(=O)(OR_{11})(R_{12})$, and $-OP(=O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

[00170] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00171] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00172] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_5 and R_6 attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; wherein

[00173] each X_1 is independently substituted or unsubstituted phenyl or $-C(X_2)(-X_3)(-X_4)$;

[00174] each X_2 is independently substituted or unsubstituted phenyl or alkyl;

[00175] each X_3 is independently hydrogen, hydroxyl, alkyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C_1 - C_{18} alkyl, substituted or unsubstituted aryl or C_1 - C_{18} aralkyl, or $-OR_{14}$, where R_{14} is C_1 - C_{18} alkyl, substituted or unsubstituted aryl or C_1 - C_{18} aralkyl, or together with X_4 is $(=O)$;

[00176] each X_4 is independently hydrogen or together with X_3 is $(=O)$; and

[00177] the bonds between the transition metal M and the macrocyclic nitrogen atoms and the bonds between the transition metal M and the oxygen atoms of the axial ligands $-OC(=O)X_1$ are coordinate covalent bonds.

[00178] In one embodiment, within Formula (IA), and groups contained therein, in one group of compounds X_1 is $-C(-X_2)(-X_3)(-X_4)$ and each X_2 , X_3 , and X_4 , in combination, corresponds to any of the combinations identified in the following table:

Combination	X_2	X_3	X_4
1	Ph	H	H
2	Ph	OH	H
3	Ph	NH ₂	H
4	Ph	=O (X_3 and X_4 in combination)	
5	Ph	CH ₃	H
6	CH ₃	H	H
7	CH ₃	OH	H
8	CH ₃	NH ₂	H
9	CH ₃	=O (X_3 and X_4 in combination)	

5







[00179] Furthermore, within embodiment (IA), and groups contained therein, in one group of compounds X_1 is $C(-X_2)(-X_3)(-X_4)$, and X_3 is $-X_5C(=O)R_{13}$, such that the combinations of X_2 , X_3 and X_4 include any of the combinations identified in the following table:

10

Combination	X_2	X_3	X_4
1	Ph	NHC(=O) R_{13}	H
2	Ph	OC(=O) R_{13}	H
3	CH ₃	NHC(=O) R_{13}	H
4	CH ₃	OC(=O) R_{13}	H

[00180] where R_{13} is C_1 - C_{18} alkyl, substituted or unsubstituted aryl or C_1 - C_{18} aralkyl, or $-OR_{14}$, where R_{14} is C_1 - C_{18} alkyl, substituted or unsubstituted aryl or C_1 - C_{18} aralkyl.

[00181] In one embodiment, the pentaaza macrocyclic ring complex corresponding to Formula (IA) is one of the complexes Formula (IE), such as (IE_{R1}), (IE_{S1}), (IE_{R2}), (IE_{S2}), (IE_{R3}), or (IE_{S3}):



wherein

M is Mn^{+2} or Mn^{+3} ;

each X_1 is independently substituted or unsubstituted phenyl or $-C(X_2)(X_3)(X_4)$;

5 each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl, or propyl;

each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, or together with X_4 is $=O$;

each X_4 is independently hydrogen or together with X_3 is $=O$; and


10 the bonds between the manganese and the macrocyclic nitrogen atoms and the bonds between the manganese and the oxygen atoms of the axial ligands $-OC(O)X_1$ are coordinate covalent bonds.

[00182] In one embodiment, each X_1 is $-C(X_2)(X_3)(X_4)$ and each $-C(X_2)(X_3)(X_4)$ corresponds to any of combinations 1 to 9 appearing in the table for Formula (IA)




15 above.

[00183] In yet another embodiment, the X and Y in pentaaza macrocyclic ring complex of Formula (I) correspond to the ligands in Formulas (IA) or (IE). For example, X and Y in the complex of Formula (I) may correspond to $-O-C(O)-X_1$, where X_1 is as defined for the complex of Formula (IA) and (IE) above.


20 [00184] In one embodiment, pentaaza macrocyclic ring complexes corresponding to Formula (I) (e.g., of Formula (I) or any of the subsets of Formula (I) corresponding to Formula (II)-(XIV), (IA) and (IE)), can comprise any of the following structures:




25




5






GC4711


[00185] In one embodiment, pentaaza macrocyclic ring complexes for use in
5 the methods and compositions described herein include those corresponding to
Formulae (2), (3), (4), (5), (6), and (7):




2




3




4



5




6



7


wherein X and Y in each of Formulae (2), (3), (4), (5), (6), and (7) are independently ligands. For example, according to one embodiment, the pentaaza macrocyclic ring complex for use in the methods and compositions described herein include those corresponding to Formulae (2), (3), (4), (5), (6), and (7) with X and Y in each of these formulae being halo, such as chloro. Alternatively, X and Y may be ligands other than chloro, such as any of the ligands described above.

[00186] In another embodiment, the pentaaza macrocyclic ring complex corresponds to Formula (6) or Formula (7):





[00187] The chemical structures of 6 (such as the dichloro complex form described, for example, in Riley, D.P., Schall, O.F., 2007, *Advances in Inorganic Chemistry*, 59: 233-263) and of 7 herein (such as the dichloro complex form of 7), are identical except that they possess mirror image chirality; that is, the enantiomeric structures are non-superimposable.

[00188] For example, the pentaaza macrocyclic ring complex may correspond to at least one of the complexes below:



[00189] In yet another embodiment, the pentaaza macrocyclic ring complex may correspond to at least one of the complexes below, and/or an enantiomer thereof:



20 [00190] In one embodiment, the enantiomeric purity of the pentaaza macrocyclic ring complex is greater than 95%, more preferably greater than 98%, more preferably greater than 99%, and most preferably greater than 99.5%. As used herein, the term “enantiomeric purity” refers to the amount of a compound having the depicted absolute stereochemistry, expressed as a percentage of the total amount of the depicted
25 compound and its enantiomer. In one embodiment, the diastereomeric purity of the pentaaza macrocyclic ring complex is greater than 98%, more preferably greater than 99%, and most preferably greater than 99.5%. As used herein, the term “diastereomeric purity” refers to the amount of a compound having the depicted absolute stereochemistry,

expressed as a percentage of the total amount of the depicted compound and its diastereomers. Methods for determining diastereomeric and enantiomeric purity are well-known in the art. Diastereomeric purity can be determined by any analytical method capable of quantitatively distinguishing between a compound and its diastereomers, such as high performance liquid chromatography (HPLC). Similarly, enantiomeric purity can be determined by any analytical method capable of quantitatively distinguishing between a compound and its enantiomer. Examples of suitable analytical methods for determining enantiomeric purity include, without limitation, optical rotation of plane-polarized light using a polarimeter, and HPLC using a chiral column packing material.

[00191] In one embodiment, a therapeutically effective amount of the pentaaza macrocyclic ring complex may be an amount sufficient to provide a peak plasma concentration of at least 0.1 μM when administered to a patient. For example, in one embodiment, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration of at least 1 μM when administered to a patient. In yet another embodiment, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration of at least 10 μM when administered to a patient. Generally, the pentaaza macrocyclic ring complex will not be administered in an amount that would provide a peak plasma concentration greater than 40 μM when administered to a patient. For example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 0.1 μM to 40 μM in a patient. As another example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 0.5 μM to 20 μM in a patient. As another example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 1 μM to 10 μM in a patient.

[00192] In yet another embodiment, a dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be at least 0.1 mg/kg, such as at least 0.2 mg/kg. For example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be at least 0.5 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be at least 1 mg/kg. In another example, the pentaaza macrocyclic compound that is administered per kg body weight

may be at least 2 mg/kg, such as at least 3 mg/kg, and even at least about 15 mg/kg, such as at least 24 mg/kg and even at least 40 mg/kg. Generally, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient will not exceed 1000 mg/kg. For example the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 0.1 to 1000 mg/kg, such as from 0.2 mg/kg to 40 mg/kg, such as 0.2 mg/kg to 24 mg/kg, and even 0.2 mg/kg to 10 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight may be in a range of from 1 mg/kg to 1000 mg/kg, such as from 3 mg/kg to 1000 mg/kg, and even from 5 mg/kg to 1000 mg/kg, such as 10 mg/kg to 1000 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight may be in a range of from 2 mg/kg to 15 mg/kg. As yet another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight may be in a range of from 3 mg/kg to 10 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 0.5 to 5 mg/kg. As yet a further example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 1 to 5 mg/kg.

[00193] In one embodiment, the dosages and/or plasma concentrations discussed above may be particularly suitable for the pentaaza macrocyclic ring complex corresponding to GC4419, although they may also be suitable for other pentaaza macrocyclic ring complexes. In addition, one of ordinary skill in the art would recognize how to adjust the dosages and/or plasma concentrations based on factors such as the molecular weight and/or activity of the particular compound being used. For example, for a pentaaza macrocyclic ring complex having an activity twice that of GC4419, the dosage and/or plasma concentration may be halved, or for a pentaaza macrocyclic ring complex having a higher molecular weight than GC4419, a correspondingly higher dosage may be used.

[00194] The dosing schedule of the pentaaza macrocyclic ring complex can similarly be selected according to the intended treatment. For example, in one embodiment, a suitable dosing schedule can comprise dosing a patient at least once per week, such as at least 2, 3, 4, 5, 6 or 7 days per week (e.g., daily), during a course of treatment. As another example, in one embodiment, the dosing may be at least once a

day (qd), or even at least twice a day (bid). In one embodiment, the course of treatment with the pentaaza macrocyclic ring complex may last at least as long as a course of treatment with an immunotherapeutic agent, such as an immune checkpoint inhibitor, and may even exceed the duration during which the immunotherapeutic agent is provided.

- 5 The course of therapy with the pentaaza macrocyclic ring complex may also start on the same date as treatment with the immunotherapeutic agent, or may start sometime after initial dosing with the immunotherapeutic agent, as is discussed in more detail below. For example, in one embodiment, for a checkpoint inhibitor that is administered for a course of therapy lasting 9 weeks, the pentaaza macrocyclic ring complex may be administered
10 for a course of therapy lasting at least 3 weeks, and even at least 4 weeks, such as at least 6 weeks and even up to at least 9 weeks.

Immune Checkpoint Inhibitors

[00195] According to one embodiment, an immune checkpoint inhibitor is provided as a part of a method of treatment herein, in combination with the pentaaza
15 macrocyclic compound. Immune checkpoints are inhibitory pathways in the immune system that maintain self-tolerance and modulate the duration and amplitude of immune response to minimize damage that could otherwise be inflicted by an excessive immune response. Without being limited by any specific theory, it is believed that cancer cells can co-opt the immune checkpoints to provide immune resistance, such as against T cells
20 that are specific for tumor antigens. That is, cancer cells may be capable of activating an immune system checkpoint to inhibit immune response to the cancer cells. Accordingly, by providing an immune checkpoint inhibitor that is capable of inhibiting the immune checkpoint, the immune response against the cancer cells can be facilitated.

[00196] Accordingly, in one embodiment, an immune checkpoint inhibitor can
25 comprise any agent that blocks or inhibits a checkpoint on the immune system or immune response. For example, many immune checkpoints are regulated by interactions between a specific receptor and a ligand, such as the interaction between the PD-1 receptor expressed on the surface of activated T cells, and its ligands PDL-1 and PDL-2 that are expressed on the surface of antigen-presenting cells. Cancer cells can co-opt
30 this interaction by presenting high levels of PDL-1 on their surface to interact with the PD-1 receptor of T-cells, thus activating this "checkpoint" of the immune system and suppressing the immune response. Accordingly, in one embodiment, an immune checkpoint inhibitor can be any one or more of a small molecule inhibitor (generally, an

inhibitor having a molecular weight of <900 daltons), an antibody, an antigen binding fragment of an antibody, and an Ig fusion protein that is capable of blocking or inhibiting an immune checkpoint, such as by blocking or inhibiting immune checkpoint receptors or blocking or inhibiting immune checkpoint receptor ligands.

5 [00197] In one embodiment, the immune checkpoint inhibitor interacts with (e.g., by inhibiting) one or more of cytotoxic T-lymphocyte antigen 4 (CTLA4), programmed death 1 (PD-1), programmed death ligand 1 (PDL-1), PDL-2, lymphocyte activation genes-3 (LAG3), B7 homolog 3 (B7-H3), B7 homolog 4 (B7-H4), indoleamine (2,3)-dioxygenase (IDO), adenosine A2a receptor (A2AR), neuritin, B- and T-lymphocyte
10 attenuator (BTLA), killer immunoglobulin-like receptors (KIR), T cell immunoglobulin and mucin domain-containing protein 3 (TIME-3), inducible T cell costimulator (ICOS), CD27, CD28, CD40, CD137, CD160, CD244, HVEM, GAL9, VISTA, 2B4, CGEN-15049, CHK 1, CHK 2, GATR, CD47 and combinations thereof. In one embodiment, the immune checkpoint inhibitor is a T-cell checkpoint inhibitor. For example, in one embodiment, the
15 checkpoint inhibitor may interact with one or more of CTLA4, PD-1 and PDL-1 or PDL-2.

 [00198] For example, the checkpoint inhibitor may be at least one of an anti-CTLA4 antibody, an anti-PD-1 antibody, and anti-PDL-1 antibody, and an anti-PDL-2 antibody. As used herein “antibody” and “antigen-binding fragments” include naturally occurring immunoglobulins (e.g., IgM, IgG, IgD, IgA, IgE) as well as non-naturally
20 occurring immunoglobulins, such as single chain antibodies, chimeric antibodies (e.g., humanized antibodies), heteroconjugate antibodies, Fab’, F(ab’)2, Fab, Fv and rIgG. An “antigen-binding fragment” is a portion of an antibody that is capable of recognizing an antigen. Furthermore, antibodies or antigen-binding fragments can include but are not limited to polyclonal, monoclonal, multispecific, human, humanized, primatized and/or
25 chimeric antibodies.

 [00199] In one embodiment, the immune checkpoint inhibitor is selected from the group consisting of ipilimumab (YERVOY (Bristol-Myers Squibb)), nivolumab (Bristol-Meyers Squibb), pembrolizumab (Merck), pidilizumab (Curetech), arelumab (Merck Serono), tremelimumab (Pfizer), atezolizumab, AMP-224
30 (GlaxoSmithKline/Amplimmune), MPDL3280A (Roche), MDX-1105 (Medarex, Inc/Bristol-Meyer Squibb), MDX-1106, MEDI-4736, IMP321, INCB024360, NLG-919, indoximod, AUNP 12, galiximab (Biogen Idec), avelumab (EMD Serono), varlilumab (CellDex Therapeutics), mogamulizumab (Kyowa Hakko Kirin), CP-870,893, MEDI-6469

(MedImmune), IPH2101 (Innate Pharma/Bristol-Meyers Squibb), urelumab (Bristol-Meyers Squibb), lirilumab (Bristol-Meyers Squibb), BMS-986016 (Bristol-Meyers Squibb), MGA271, IMP321, BMS-936559, MSB0010718C, anti-OX40, MK-3475, CT-011, BY55, AMP224, and BGB-A317.

5 [00200] The dose of the immune checkpoint inhibitor can be selected according to the treatment to be provided and the particular immune checkpoint inhibitor being used. For example, a suitable dose of an immune checkpoint inhibitor may be at least at least 0.1 mg/kg. For example, the dose of the immune checkpoint inhibitor that is administered per kg body weight of the patient may be at least 0.5 mg/kg. As another
10 example, the dose of the immune checkpoint inhibitor that is administered per kg body weight of the patient may be at least 1 mg/kg. In another example, the immune checkpoint inhibitor that is administered per kg body weight may be at least 2 mg/kg, such as at least 3 mg/kg, and even at least 10 mg/kg, such as at least 15 mg/kg. Generally, the dose of the immune checkpoint inhibitor that is administered per kg body
15 weight of the patient will not exceed 20 mg/kg, such as a dose that does not exceed 15 mg/kg, and even that does not exceed 10 mg/kg. For example the dose of the immune checkpoint inhibitor that is administered per kg body weight of the patient may be in the range of from 0.1 to 15 mg/kg. As another example, the dose of the immune checkpoint inhibitor that is administered per kg body weight may be in a range of from 2 mg/kg to 15
20 mg/kg. As yet another example, the dose of the immune checkpoint inhibitor that is administered per kg body weight may be in a range of from 3 mg/kg to 10 mg/kg.

 [00201] The dosing schedule of the immune checkpoint inhibitor can similarly be selected according to the intended treatment and the particular immune checkpoint inhibitor being provided. For example, in one embodiment, a suitable dosing schedule in
25 one embodiment can comprise dosing a patient once every 2 or 3 weeks, for a total of 4 doses (9 weeks of treatment total). That is, in some embodiments treatment may involve a course of therapy that lasts at least 9 weeks and even 10 weeks, but in some embodiments may not extend past 16 weeks. In particular, the package insert for Yervoy (ipilimumab) indicates that a dose of 3 mg/kg should be given every 3 weeks for 4 doses,
30 as given by IV over the course of 90 minutes. Dosage regimens for Opdivo (nivolumab) and Keytruda (pembrolizumab) similarly indicate dosing once every 2 or 3 weeks.

Adoptive T-Cell Transfer Therapies

[00202] According to one embodiment, an adoptive T-cell transfer therapy is provided as a part of a method of treatment herein, in combination with the pentaaza macrocyclic compound. In adoptive cell therapy, cells are removed from a donor and
5 cultured and/or manipulated in vivo, after which they are administered to the patient for treatment. For example, cancer-specific cytotoxic T-cells can be cultured and/or modified to provide for the targeting and destroying of cancer cells in a patient. In one embodiment, an adoptive T-cell transfer therapy comprises administering to the subject cancer-specific autologous T-cells (i.e., cells originally obtained from the same patient).
10 In another embodiment, the adoptive T-cell transfer therapy comprises administering to the subject cancer-specific allogeneic T-cells (i.e., cells originally obtained from a donor). The cancer specific T-cells can facilitate immune system attack of the cancer cells to provide for treatment of the cancer in the subject.

[00203] In one embodiment, the adoptive T-cell transfer therapy comprises
15 providing autologous tumor infiltrating lymphocytes. For example, in one embodiment, tumor infiltrating lymphocytes (TILs) can be expanded ex vivo from tumor fragments from a patient, and transplanted back into the subject. In one embodiment, the TILs are expanded by placing in a growth medium and exposing to a high dose of IL-2. Once the TILs have been sufficiently expanded, a patient may receive the cells via infusions, such
20 as via 1 to 2 infusions separated by 1-2 weeks.

[00204] In yet another embodiment, the adoptive T-cell transfer therapy comprises providing antigen-expanded CD8+ and/or CD4+ T cells. For example, in one embodiment, peripheral blood lymphocytes can be harvested and expanded in vitro through antigen-specific expansion to produce tumor-specific T cells. In yet another
25 embodiment, the adoptive T-cell transfer therapy comprises providing genetically modified T cells that express T-cell receptors (TCR) that recognize tumor antigens. For example, in one embodiment, peripheral blood lymphocytes can be harvested and genetically engineered to produce tumor-specific T cells with TCRs that specifically recognize cancer antigens, such as by transducing lymphocytes with a retrovirus that
30 contains genes encoding the tumor-antigen-specific TCR. The tumor-specific T-cells can be provided to the patient by one or more infusions of the cells, as discussed above.

[00205] According to one aspect, the dosing regimen and schedule for the adoptive T-cell transfer process can be selected according to the treatment to be provided and the type of cells to be transferred, and the dosing regimen and schedule may further be coordinated with dosing with the pentaaza macrocyclic ring complex, as is discussed in more detail below.

Cancer Vaccines

[00206] According to one embodiment, a cancer vaccine is provided as a part of a method of treatment herein, in combination with the pentaaza macrocyclic compound. Cancer vaccines may help prime and mobilize the immune system to attack cancer cells in the body, and may use, for example, cancer cells or parts of cancer cells, or antigens, to invoke or increase the immune response to cancer cells in the patient.

[00207] In one embodiment, the cancer vaccine is selected from the group consisting of tumor cell vaccines, antigen vaccines, dendritic cell vaccines, DNA vaccines and vector based vaccines. For example, a tumor cell vaccine can comprise cancer cells that have been removed from a subject and then modified so they cannot reproduce, such as by exposing to radiation, as well as optionally by modifying to make the cells more visible to the immune system. The modified tumor cells can then be provided to a subject to train the subject's immune system to recognize the cancer cells and go after other such cancer cells in the subject's body. The tumor cell vaccines can be either autologous (from the subject themselves) or allogeneic (from a donor). Antigen vaccines provide one or more antigens, typically specific for a certain type of cancer, to train the immune system to recognize the cancer-specific antigens. Dendritic cell vaccines involve exposing immune cells in vitro to antigens and other chemicals that convert them into dendritic cells, after which the dendritic cells are injected back into a subject to provoke an immune response. DNA vaccines and vector vaccines can be used to program cells to express specific antigens to provoke an immune response.

[00208] In one embodiment, a cancer vaccine for use in treatment can be selected from the group consisting of M-Vax (Avax Technologies), Provenge (Dendreon), GRNVAC1 (Geron), Bexidex (IDM Pharma), Uvidex (IDM Pharma), Collidex (IDM Pharma), INGN 225 (Introgen Therapeutics), M3Tk (MolMed), DC-Vax (Northwest Biotherapeutics), CVac (Prima Biomed), GVAX (Cell Genesys), Lucanix (NovaRx), Onyvax-P (Onyvax), HSPP-96 Oncophage (Antigenics), BiovaxID (Biovest International),

NeuVax (Apthera), CDX-110 (CeppDex), GV1001 (Pharmexa), CYT004-MelQbG10 (Cytos Biotechnology), li-Key/HER2/neu (Generiex Biotechnology), MAGE-A3 (Glaxo-SmithKline Biologicals), IDM-2101 (IDM Pharma), IMA901/IMA910 (Immatics Biotechnologies), melanoma cancer vaccine (Norwood Immunology), inCVAX (Immunophotonics)) and Stimuvax (Oncothyreon).

Timing of Administration

[00209] In one embodiment, a treatment regimen can comprise administering an initial dose of the pentaaza macrocyclic complex after a predetermined period of time has elapsed since administration of an initial dose of an immunotherapeutic agent. That is, the treatment regimen can comprise administering an initial dose and optionally one or more subsequent doses of the immunotherapeutic agent, with the onset of dosing with the pentaaza macrocyclic ring complex being delayed for a predetermined period of time after the initial immunotherapeutic agent dose. Unexpectedly, it has been discovered that delaying the initial administration of the pentaaza macrocyclic ring complex until a predetermined time after treatment with the immunotherapeutic agent has begun, can provide significantly improved results over treatment where dosing with the immunotherapeutic agent and pentaaza macrocyclic ring complex is started closer to the same time.

[00210] For example, in an embodiment where an immune checkpoint inhibitor is being administered, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be performed after a predetermined period of time has elapsed since an initial administration of the immune checkpoint inhibitor to start a course of therapy. In one embodiment, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be no less than 3 days after the initial administration of the immune checkpoint inhibitor (for example, if the immune checkpoint inhibitor is administered on day 1 of treatment, the pentaaza macrocyclic ring complex is administered no sooner than on day 4 of treatment). In another embodiment, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be no less than 6 days after the initial administration of the immune checkpoint inhibitor (for example, if the immune checkpoint inhibitor is administered on day 1 of treatment, the pentaaza macrocyclic ring complex is administered no sooner than on day 7 of treatment). In yet another embodiment, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be no less than 2 weeks after the

initial administration of the immune checkpoint inhibitor. In yet another embodiment, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be no less than 3 weeks after the initial administration of the immune checkpoint inhibitor. In yet another embodiment, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy may be no less than 6 weeks after the initial administration of the immune checkpoint inhibitor. Generally, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy will be within 9 weeks of the initial administration of the immune checkpoint inhibitor. For example, the initial administration of the pentaaza macrocyclic ring complex in a course of therapy can be in the range of from 3 days to 9 weeks after the initial administration of the immune checkpoint inhibitor. In one embodiment, an initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows at least two doses of the immune checkpoint inhibitor. In another embodiment, an initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows at least three doses of the immune checkpoint inhibitor. In another embodiment, an initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows at least four doses of the immune checkpoint inhibitor. In another embodiment, an initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows at least five doses of the immune checkpoint inhibitor. As an example, in one embodiment where a course of treatment with a checkpoint inhibitor involves dosing once every 3 weeks for 4 total doses, the initial administration of the pentaaza macrocyclic ring complex can be provided not less than 3 days after the initial immune checkpoint inhibitor dose, but no more than 9 weeks after the initial immune checkpoint inhibitor dose, meaning that administration of the pentaaza macrocyclic ring complex may be delayed until before the final dose of the immune checkpoint inhibitor given during a course of therapy. Furthermore, in one embodiment, the initial administration of the pentaaza macrocyclic ring complex may be delayed with respect to an initial administration of the immune checkpoint inhibitor until after at least a second dose of the immune checkpoint inhibitor has been administered, such as after a third dose of the immune checkpoint inhibitor has been administered, and even after a fourth dose of the immune checkpoint inhibitor has been administered. Furthermore, other dosing schemes other than those specifically mentioned herein may also be provided.

[00211] It yet another embodiment, it has been unexpectedly found that improved results in terms of treatment can be provided by dosing with the pentaaza macrocyclic ring complex on a day that is other than a day on which dosing with the immunotherapeutic agent is provided. For example, dosing with a pentaaza macrocyclic ring complex on separate days from the days on which immune checkpoint inhibitors are dosed, that is, skipping administration of the pentaaza macrocyclic ring complex on days when the immune checkpoint inhibitor is being administered, provides improved benefits in terms of the immune response. Accordingly, in one embodiment, doses of the pentaaza macrocyclic ring complex that are provided in a course of cancer treatment are provided on separate days from any dose of an immune checkpoint inhibitor that is provide in the course of cancer therapy.

[00212] Similarly, in an embodiment where an adoptive T-cell transfer therapy is being administered, the initial administration of the pentaaza macrocyclic ring complex may be performed after a predetermined period of time has elapsed since an initial administration of the T-cells being provided as a part of the start of the adoptive T-cell transfer therapy. The predetermined period of time may be, for example, the same time period described for delay between the pentaaza macrocyclic ring complex and immune checkpoint inhibitor above, or different delay in the administration of the pentaaza macrocyclic complex may also be provided. Furthermore, administration of the pentaaza macrocyclic ring complex may “skip” days on which an infusion of cells as a part of the adoptive T-cell transfer therapy is being provided, similarly to the combination with the immune checkpoint inhibitor therapy, as discussed above. Also, in an embodiment where a cancer vaccine is being administered, the initial administration of the pentaaza macrocyclic ring complex may be performed after a predetermined period of time has elapsed since an initial administration of the cancer vaccine. The predetermined period of time may be, for example, the same time period described for delay between the pentaaza macrocyclic ring complex and immune checkpoint inhibitor above, or different delay in the administration of the pentaaza macrocyclic complex may also be provided. Furthermore, administration of the pentaaza macrocyclic ring complex may “skip” days on which a cancer vaccine is being administered to the patient, similarly to the combination with the immune checkpoint inhibitor therapy, as discussed above.

[00213] Furthermore, in one embodiment, a treatment regimen may involve administration of multiple immunotherapeutic agents. For example, in one embodiment,

the administration of a checkpoint inhibitor and pentaaza macrocyclic ring complex may be further supplemented with the administration of one or more of adoptive T-cell transfer and cancer vaccine, either prior to, concomitantly with, or after administration of one or more of the immune checkpoint inhibitor and pentaaza macrocyclic ring complex. In yet another embodiment, the administration of an adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex may be further supplemented with the administration of one or more of an immune checkpoint inhibitor and cancer vaccine, either prior to, concomitantly with, or after administration of one or more of the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex. In yet another embodiment, the administration of a cancer vaccine and pentaaza macrocyclic ring complex may be further supplemented with the administration of one or more of adoptive T-cell transfer and immune checkpoint inhibitor, either prior to, concomitantly with, or after administration of one or more of the cancer vaccine and pentaaza macrocyclic ring complex. Furthermore, other dosing schemes other than those specifically mentioned herein may also be provided.

Other Cancer Therapies

[00214] In one embodiment, the treatment provided herein can comprise further comprise treatment with another therapy other than those specifically described above, such as for example a radiation therapy, a chemotherapy, or other immunotherapeutic treatment. For example, in one embodiment, one or more of radiation therapy and chemotherapy is administered to the subject prior to, concomitantly with, or after administration of one or more of the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine) and the pentaaza macrocyclic ring complex. Further detailed description of radiation therapies and chemotherapies suitable for the treatment of cancer are provided below.

[00215] In one embodiment, one or more of radiation therapy and chemotherapy can be administered concomitantly with administration of one or more of the immunotherapeutic agent and pentaaza macrocyclic ring complex. For example, one or more of the immunotherapeutic agent and pentaaza macrocyclic ring complexes may be administered during a course of radiation therapy and/or chemotherapy, such as in between, before or after, or on the same day as dosing with radiation and/or chemotherapy. In one embodiment, as is further demonstrated in the Examples below, it has been found that administering a pentaaza macrocyclic ring complex such as GC4419

can improve a subject's response to radiation therapy, including when such radiation therapy is combined with administration of an immunotherapy agent, such as the checkpoint inhibitor anti-CTLA4. Without being limited by any theory, it is believed that pentaaza macrocyclic ring complexes such as GC4419 can sensitize cancer cells to
5 radiation to improve treatment therewith.

[00216] In yet another embodiment, the combination therapy of the pentaaza macrocyclic ring complex and immunotherapeutic agent (e.g. immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine), can be administered in the absence of any other cancer treatment. As demonstrated further in the examples below, it has been
10 unexpectedly discovered that the pentaaza macrocyclic ring complexes are capable of enhancing the response to and/or efficacy of immunotherapeutic agents such as immune checkpoint inhibitors, even when administered without radiation therapy or chemotherapy. Accordingly, in one embodiment, the cancer treatment provided to the subject may consist essentially of the pentaaza macrocyclic ring complex and immunotherapeutic
15 agent, without the administration of a chemotherapeutic agent or radiation exposure (i.e. without administering a radiation dose or dose fraction). For example, the combination of the pentaaza macrocyclic ring complex and immunotherapeutic agent may be administered to a subject that is not receiving radiation therapy, and/or that is not receiving chemotherapy. That is, in one embodiment, the treatment comprises
20 administering the pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy. In yet another embodiment, the treatment comprises administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy. In yet another embodiment, where a course of therapy comprises administration of the pentaaza macrocyclic ring complex and the immune
25 checkpoint inhibitor, they are administered to a subject that does not receive radiation therapy during the course of therapy.

[00217] In one embodiment, the subject receiving the combination of pentaaza macrocyclic ring complex and immunotherapeutic agent (e.g. immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine), may be one that has not been exposed
30 to radiation (i.e., received a dose or dose fraction of radiation) and/or has not received a dose of chemotherapeutic agent for at least one day, such as at least one week, and even at least one month, and even at least 6 months, and/or that has not ever received such treatment at all before initial treatment with one or more of the pentaaza macrocyclic ring

complex and immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine). In yet another embodiment, any radiation therapy and/or chemotherapy that is administered to the subject after the combination treatment with the pentaaza macrocyclic ring complex and immunotherapeutic agent (e.g., immune

5 checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine) is delayed by at least one day, such as at least one week, and even at least one month, such as at least 6 months, after a final dose of one or more of the pentaaza macrocyclic ring complex and immunotherapeutic agent provided during the course of the combination therapy treatment. That is, the combination therapy of the pentaaza macrocyclic ring complex

10 and immunotherapeutic agent (e.g. immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine) can be administered to a subject that has never before received radiation therapy and/or chemotherapy, or that has received such therapy only in the distant past. Furthermore, the combination therapy of the pentaaza macrocyclic ring complex and immunotherapeutic agent (e.g. immune checkpoint inhibitor, adoptive T-cell transfer,

15 cancer vaccine) can be administered to provide a course of treatment that does not include any exposure to radiation or doses of chemotherapeutic agent. As yet a further embodiment, the combination therapy of the pentaaza macrocyclic ring complex and immunotherapeutic agent (e.g. immune checkpoint inhibitor, adoptive T-cell transfer, cancer vaccine) can be provided to form a course of treatment substantially without

20 performing any radiation therapy or chemotherapy after the course of treatment, or with such radiation or chemotherapeutic treatment being performed only after a significant period of time has elapsed after the course of combination treatment has ended. In one embodiment, the treatment comprises administering one or more of the pentaaza macrocyclic ring complex and immune checkpoint inhibitor to the subject on a day other

25 than a day that the subject is receiving radiation therapy.

METHODS OF ADMINISTRATION

[00218] According to one embodiment, the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), is

30 administered as a co-therapy or combination therapy with the pentaaza macrocyclic ring complex. Co-therapy or combination therapy according to the methods described herein is intended to embrace administration of each compound in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as

well to embrace co-administration of these agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent, or single or multiple parenteral administrations, or other routes of administration and dosage forms. When administered in combination, therefore, the therapeutic agents (i.e., the pentaaza macrocyclic ring complex and/or the immunotherapeutic agent) can be formulated as separate compositions that are administered at the same time or sequentially at different times, or the therapeutic agents can be given as a single composition. Pharmaceutical compositions and formulations are discussed elsewhere herein. Furthermore, while the immunotherapeutic agent is referred to as including one or more of an immune checkpoint inhibitor, adoptive T-cell transfer therapy, and cancer vaccine, it is noted that all combinations of these are also explicitly included herein. Furthermore, other immunotherapeutic agents such as anti-cancer antibodies, cytokines such as IL-2, and other cancer treating agents, can also be administered as a co-therapy or combination therapy with the pentaaza macrocyclic ring complex and the specific immunotherapeutic agents described herein.

[00219] It is not necessary that the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) be administered simultaneously or essentially simultaneously; the agents and compounds may be administered in sequence. The advantage of a simultaneous or essentially simultaneous administration, or sequential administration, is well within the determination of the skilled clinician. For instance, while a pharmaceutical composition or formulation comprising an immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be advantageous for administering first in the combination for one particular treatment, prior to administration of the pentaaza macrocyclic ring complex, prior administration of the pentaaza macrocyclic ring complex may be advantageous in another treatment. It is also understood that the instant combination of pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be used in conjunction with other methods of treating cancer (typically cancerous tumors) including, but not limited to, radiation therapy and surgery, or other chemotherapy. It is further understood that another active agent, such

as a cytostatic or quiescent agent, or antiemetic agent, if any, may be administered sequentially or simultaneously with any or all of the other synergistic therapies.

[00220] Thus, embodiments of the therapeutic method include wherein a pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune
5 checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), and combinations thereof, are administered simultaneously or sequentially. For instance, the present disclosure encompasses a method for the treatment of cancer wherein a pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered
10 simultaneously or sequentially. Other active agents can also be administered simultaneously or sequentially with the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine).

[00221] As noted above, if the pentaaza macrocyclic ring complex and the
15 immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are not administered simultaneously or essentially simultaneously, then the initial order of administration of the components may be varied.

[00222] Thus, for example, the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be
20 administered first, followed by the administration of the pentaaza macrocyclic ring complex; or the pentaaza macrocyclic ring complex may be administered first, followed by the administration of the immunotherapeutic agent. This alternate administration may be repeated during a single treatment protocol. Other sequences of administration to exploit the effects described herein are contemplated, and other sequences of administration of
25 other active agents can also be provided.

[00223] In one embodiment, the subject is pre-treated with the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), followed by administration of the pentaaza macrocyclic ring complex, or vice versa. In accordance with such embodiments, the pentaaza macrocyclic
30 ring complex may be administered at least 1 hour, and even at least 3 days, after administration of the immunotherapeutic agent, or vice versa. For example, in one embodiment, the pentaaza macrocyclic ring complex is administered between 1 hour

and 3 days after administration of the immunotherapeutic agent, or vice versa. In another embodiment, for example, the pentaaza macrocyclic ring complex is administered between 1 hour and 1 day after administration of the immunotherapeutic agent, or vice versa. For example, the pentaaza macrocyclic ring complex may be administered within
5 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 18 hours, 24 hours, 36 hours, 48 hours, one week, 2 weeks, 3 weeks, 4 weeks, 6 weeks, 8 weeks, 9 weeks, 10 weeks or 12 weeks after administration of the immunotherapeutic agent, or vice versa. In these and other embodiments, the immunotherapeutic agent may be administered in multiple doses leading up to administration of the pentaaza macrocyclic ring complex, or
10 vice versa.

[00224] Alternatively, the subject may be pre-treated with the pentaaza macrocyclic ring complex, followed by administration of the immunotherapeutic agent, or vice versa. In accordance with such embodiments, the pentaaza macrocyclic ring complex may be administered within at least 1 plasma half-life of the immunotherapeutic
15 agent, such as within 4 plasma half-lives of the immunotherapeutic agents, or vice versa. For example, the pentaaza macrocyclic ring complex may be administered within 1, 2, or 3 plasma half-lives of the other immunotherapeutic agents, or vice versa.

[00225] In other alternative embodiments, the subject may be pre-treated with the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer
20 therapy, cancer vaccine), followed by administration of the pentaaza macrocyclic ring complex, which is further followed by one or more additional administrations of the immunotherapeutic agent, or vice versa. For example, the subject could be pre-treated with a dose of immunotherapeutic agent, followed by administration of a dose of pentaaza macrocyclic ring complex, which is then followed by the administration of additional (or
25 partial) dose of the same or different immunotherapeutic agent, which may be further followed by another dose of pentaaza macrocyclic ring complex. Further, the subject could be pre-treated with a partial or full dose of pentaaza macrocyclic ring complex, followed by administration of an immunotherapeutic agent, which is then followed by administration of an additional (or partial) dose of pentaaza macrocyclic complex.

30 [00226] As described in further detail below, the combinations of the disclosure may also be co-administered with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. Combinations

may alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a multiple combination formulation is inappropriate.

[00227] In one embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer
5 therapy, cancer vaccine) can generally be administered according to therapeutic protocols that may be known for these agents in the art. For example, the administration of the various components can be varied depending on the disease being treated and the effects of pentaaza macrocyclic ring complex and immunotherapeutic agent on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic
10 protocols (e.g., dosage amounts and times of administration) can be varied in view of the observed effects of the administered therapeutic agents (i.e., pentaaza macrocyclic ring complex, immunotherapeutic agent) on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

[00228] Also, in general, the pentaaza macrocyclic ring complex and the
15 immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) do not have to be administered in the same pharmaceutical composition, and may, because of different physical and chemical characteristics, have to be administered by different routes. For example, the pentaaza macrocyclic ring complex may be administered orally to generate and maintain good blood levels thereof, while the
20 immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered intravenously or via transfusion, or vice versa. The mode of administration may include, where possible, in the same pharmaceutical composition, or in separate pharmaceutical compositions (e.g., two or three separate compositions). Furthermore, once the initial administration has been
25 made, then based upon the observed effects, the dosage, modes of administration and times of administration can be modified.

[00229] The particular choice of pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), and other related therapies (such as chemotherapy or
30 radiation), will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

[00230] Thus, in accordance with experience and knowledge, the practicing physician can modify each protocol for the administration of a component (pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) of the treatment according to
5 the individual patient's needs, as the treatment proceeds.

[00231] The attending clinician, in judging whether treatment is effective at the dosage administered, will consider the general well-being of the patient as well as more definite signs such as relief of disease-related symptoms, inhibition of tumor growth, actual shrinkage of the tumor, or inhibition of metastasis. Size of the tumor can be
10 measured by standard methods such as radiological studies, e.g., CAT or MRI scan, and successive measurements can be used to judge whether or not growth of the tumor has been retarded or even reversed. Relief of disease-related symptoms such as pain, and improvement in overall condition can also be used to help judge effectiveness of treatment.

[00232] The products of which the combination are composed may be administered simultaneously, separately or spaced out over a period of time so as to obtain the maximum efficacy of the combination; it being possible for each administration to vary in its duration from a rapid administration to a relatively continuous perfusion of either component (in separate formulations or in a single formulation). As a result, for the
15 purposes of the present disclosure, the combinations are not exclusively limited to those which are obtained by physical association of the constituents, but also to those which permit a separate administration, which can be simultaneous or spaced out over a period of time.

[00233] Accordingly, administration of the components described herein can
25 occur as a single event or over a time course of treatment. For example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can be administered (simultaneously or in sequence) hourly (e.g., every hour, every two hours, every three hours, every four hours, every five hours, every six hours, and so on), daily, weekly, bi-
30 weekly, or monthly. For treatment of acute conditions, the time course of treatment may be at least several hours or days. Certain conditions could extend treatment from several days to several weeks. For example, treatment could extend over one week, two weeks, or three weeks. For more chronic conditions, treatment could extend from several weeks

to several months, a year or more, or the lifetime of the patient in need of such treatment. Alternatively, the compounds and agents can be administered hourly, daily, weekly, bi-weekly, or monthly, for a period of several weeks, months, years, or over the lifetime of the patient as a prophylactic measure.

5 [00234] The dose or amount of pharmaceutical compositions including the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) administered to the patient should be an effective amount for the intended purpose, i.e., treatment or prophylaxis of one or more of the diseases, pathological disorders, and medical
10 conditions discussed herein, particularly cancer. Generally speaking, the effective amount of the composition administered can vary according to a variety of factors such as, for example, the age, weight, sex, diet, route of administration, and the medical condition of the patient in need of the treatment. Specifically preferred doses are discussed more fully herein. It will be understood, however, that the total daily usage of
15 the compositions described herein will be decided by the attending physician or veterinarian within the scope of sound medical judgment.

 [00235] As noted above, the combinations can be co-administered (via a co-formulated dosage form or in separate dosage forms administered at about the same time). The combinations can also be administered separately, at different times, with each
20 agent in a separate unit dosage form. Numerous approaches for administering the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) and pentaaza macrocyclic ring complex can be readily adapted for use in the present disclosure. The pharmaceutical compositions may be delivered orally, e.g., in a tablet or capsule unit dosage form, or parenterally, e.g., in an injectable
25 unit dosage form, or by some other route. For systemic administration, for example, the drugs can be administered by, for example, intravenous infusion (continuous or bolus). The compositions can be used for any therapeutic or prophylactic treatment where the patient benefits from treatment with the combination.

 [00236] The specific therapeutically effective dose level for any particular
30 patient will depend upon a variety of factors including the disorder being treated and the

severity of the disorder; activity of the specific compound(s) employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the route of administration; the rate of excretion of the specific compound(s) employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound(s) employed and like factors well known in the medical and/or veterinary arts. For example, it is well within the skill of the art to start doses of the compound(s) at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. If desired, the effective daily doses may be divided into multiple doses for purposes of administration. Consequently, single dose compositions may contain such amounts or submultiples to make up the daily dose.

[00237] In one embodiment, suitable or preferred doses for each of the components are employed in the methods or included in the compositions described herein. Preferred dosages for the pentaaza macrocyclic ring complex, for instance, may be within the range of 10 to 500 mg per patient per day. However, the dosage may vary depending on the dosing schedule, which can be adjusted as necessary to achieve the desired therapeutic effect. It should be noted that the ranges of effective doses provided herein are not intended to limit the disclosure and represent exemplary dose ranges. The most preferred dosage will be tailored to the individual subject, taking into account, among other things, the particular combinations employed, and the patient's age, sex, weight, physical condition, diet, etc., as is understood and determinable by one of ordinary skill in the art without undue experimentation.

[00238] Treatment of cancer, or cancer therapies, described herein includes achieving a therapeutic benefit, however the therapy may also be administered to achieve a prophylactic benefit. Therapeutic benefits generally refer to at least a partial eradication or amelioration of the underlying disorder being treated. For example, in a cancer patient, therapeutic benefit includes (partial or complete) eradication or amelioration of the underlying cancer. Also, a therapeutic benefit is achieved with at least partial, or complete, eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the patient, notwithstanding the fact that the patient may still be afflicted with the underlying disorder. For prophylactic benefit, a method of the disclosure may be performed on, or a composition of the invention administered to, a patient at risk of

developing cancer, or to a patient reporting one or more of the physiological symptoms of such conditions, even though a diagnosis of the condition may not have been made.

Cancer Treatment Methods

[00239] In general, any subject having, or suspected of having, a cancer or
5 other proliferative disorder may be treated using the compositions and methods of the present disclosure. Subjects receiving treatment according to the methods described herein are mammalian subjects, and typically human patients. Other mammals that may be treated according to the present disclosure include companion animals such as dogs and cats, farm animals such as cows, horses, and swine, as well as birds and
10 more exotic animals (e.g., those found in zoos or nature preserves). In one embodiment of the disclosure, a method is provided for the treatment of cancerous tumors, particularly solid tumors. Advantageously, the methods described herein may reduce the development of tumors, reduce tumor burden, or produce tumor regression in a mammalian host. Cancer patients and individuals desiring cancer prophylaxis can be
15 treated with the combinations described herein.

[00240] Cancer and tumors generally refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. By means of the pharmaceutical combinations, co-formulations, and combination therapies of the present disclosure, various tumors can be treated such as tumors of the breast, heart,
20 lung, small intestine, colon, spleen, kidney, bladder, head and neck, ovary, prostate, brain, pancreas, skin, bone, bone marrow, blood, thymus, uterus, testicles, cervix, and liver.

[00241] In one embodiment, the tumor or cancer is chosen from adenoma, angio-sarcoma, astrocytoma, epithelial carcinoma, germinoma, glioblastoma, glioma,
25 hamartoma, hemangioendothelioma, hemangiosarcoma, hematoma, hepato-blastoma, leukemia, lymphoma, medulloblastoma, melanoma, neuroblastoma, osteosarcoma, retinoblastoma, rhabdomyosarcoma, sarcoma, and teratoma. The tumor can be chosen from acral lentiginous melanoma, actinic keratoses, adenocarcinoma, adenoid cystic carcinoma, adenomas, adenosarcoma, adenosquamous carcinoma, astrocytic tumors,
30 bartholin gland carcinoma, basal cell carcinoma, bronchial gland carcinomas, capillary, carcinoids, carcinoma, carcinosarcoma, cavernous, cholangio-carcinoma, chondrosarcoma, choroid plexus papilloma/carcinoma, clear cell carcinoma,

cystadenoma, endodermal sinus tumor, endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, ependymal, epitheloid, Ewing's sarcoma, fibrolamellar, focal nodular hyperplasia, gastrinoma, germ cell tumors, glioblastoma, glucagonoma, hemangioblastomas, hemangioendothelioma, hemangiomas, hepatic
 5 adenoma, hepatic adenomatosis, hepatocellular carcinoma, insulinoma, intraepithelial neoplasia, interepithelial squamous cell neoplasia, invasive squamous cell carcinoma, large cell carcinoma, leiomyosarcoma, lentigo maligna melanomas, malignant melanoma, malignant mesothelial tumors, medulloblastoma, medulloepithelioma, melanoma, meningeal, mesothelial, metastatic carcinoma, mucoepidermoid carcinoma,
 10 neuroblastoma, neuroepithelial adenocarcinoma nodular melanoma, oat cell carcinoma, oligodendroglial, osteosarcoma, pancreatic, papillary serous adeno-carcinoma, pineal cell, pituitary tumors, plasmacytoma, pseudo-sarcoma, pulmonary blastoma, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, small cell carcinoma, soft tissue carcinomas, somatostatin-secreting tumor, squamous carcinoma,
 15 squamous cell carcinoma, submesothelial, superficial spreading melanoma, undifferentiated carcinoma, uveal melanoma, verrucous carcinoma, vipoma, well differentiated carcinoma, and Wilm's tumor.

[00242] Thus, for example, the present disclosure provides methods for the treatment of a variety of cancers, including, but not limited to, the following: carcinoma
 20 including that of the bladder (including accelerated and metastatic bladder cancer), breast, colon (including colorectal cancer), kidney, liver, lung (including small and non-small cell lung cancer and lung adenocarcinoma), ovary, prostate, testes, genitourinary tract, lymphatic system, rectum, larynx, pancreas (including exocrine pancreatic carcinoma), esophagus, stomach, gall bladder, cervix, thyroid, and skin (including squamous cell
 25 carcinoma); hematopoietic tumors of lymphoid lineage including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma, histiocytic lymphoma, and Burketts lymphoma; hematopoietic tumors of myeloid lineage including acute and chronic myelogenous leukemias, myelodysplastic syndrome, myeloid leukemia,
 30 and promyelocytic leukemia; tumors of the central and peripheral nervous system including astrocytoma, neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and

other tumors including melanoma, xenoderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer, and teratocarcinoma.

[00243] For example, particular leukemias that can be treated with the combinations and methods described herein include, but are not limited to, acute
5 nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemic leukemia, basophylic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia,
10 hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia,
15 myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, and undifferentiated cell leukemia.

[00244] Lymphomas can also be treated with the combinations and methods described herein. Lymphomas are generally neoplastic transformations of cells that
20 reside primarily in lymphoid tissue. Lymphomas are tumors of the immune system and generally are present as both T cell- and as B cell-associated disease. Among lymphomas, there are two major distinct groups: non-Hodgkin's lymphoma (NHL) and Hodgkin's disease. Bone marrow, lymph nodes, spleen and circulating cells, among others, may be involved. Treatment protocols include removal of bone marrow from the
25 patient and purging it of tumor cells, often using antibodies directed against antigens present on the tumor cell type, followed by storage. The patient is then given a toxic dose of radiation or chemotherapy and the purged bone marrow is then re-infused in order to repopulate the patient's hematopoietic system.

[00245] Other hematological malignancies that can be treated with the
30 combinations and methods described herein include myelodysplastic syndromes (MDS), myeloproliferative syndromes (MPS) and myelomas, such as solitary myeloma and multiple myeloma. Multiple myeloma (also called plasma cell myeloma) involves the skeletal system and is characterized by multiple tumorous masses of neoplastic plasma

cells scattered throughout that system. It may also spread to lymph nodes and other sites such as the skin. Solitary myeloma involves solitary lesions that tend to occur in the same locations as multiple myeloma.

[00246] In one embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of breast cancer, melanoma, oral squamous cell carcinoma, lung cancer including non-small cell lung cancer, renal cell carcinoma, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, urothelial cancer, bladder cancer, colorectal cancer, head and neck cancers such as squamous cell carcinoma, and pancreatic cancer. In yet another embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of head and neck cancer and lung cancer.

Pharmaceutical Formulations

[00247] Another aspect of the present disclosure relates to the pharmaceutical compositions comprising the combinations described herein, together with a pharmaceutically acceptable excipient. The pharmaceutical compositions include the pentaaza macrocyclic ring complex (e.g., those corresponding to Formula (I)), and at least one immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), and combinations thereof, as discussed above, typically formulated as a pharmaceutical dosage form, optionally in combination with a pharmaceutically acceptable carrier, additive or excipient. In one embodiment, for example, the pharmaceutical composition comprises a pentaaza macrocyclic ring complex, the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) and a pharmaceutically acceptable excipient. Pharmaceutical compositions according to the present disclosure may be used in the treatment of cancer.

[00248] The pharmaceutical compositions described herein are products that result from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. Fixed combinations are those in which the active ingredients, e.g., a pentaaza macrocyclic ring complex and an immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), are administered to a patient simultaneously in the form of a single entity or dosage. Other active agents may also be administered as a part of the

single entity or dosage, or may be separately administered. Non-fixed combinations are those in which the active ingredients, e.g., a pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), are administered to a patient as separate entities either
5 simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the compounds in the body of the patient. The latter also applies to cocktail therapy, e.g., the administration of three or more active ingredients.

[00249] The above-described pentaaza macrocyclic ring complex and the
10 immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be dispersed in a pharmaceutically acceptable carrier prior to administration to the mammal; i.e., the components described herein are preferably co-formulated. The carrier, also known in the art as an excipient, vehicle, auxiliary, adjuvant, or diluent, is typically a substance which is pharmaceutically inert, confers a suitable
15 consistency or form to the composition, and does not diminish the efficacy of the compound. The carrier is generally considered to be "pharmaceutically or pharmacologically acceptable" if it does not produce an unacceptably adverse, allergic or other untoward reaction when administered to a mammal, especially a human.

[00250] The selection of a pharmaceutically acceptable carrier will also, in part,
20 be a function of the route of administration. In general, the compositions of the described herein can be formulated for any route of administration so long as the blood circulation system is available via that route, and in accordance with the conventional route of administration. For example, suitable routes of administration include, but are not limited to, oral, parenteral (e.g., intravenous, intraarterial, subcutaneous, rectal, subcutaneous,
25 intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, or intrasternal), topical (nasal, transdermal, intraocular), intravesical, intrathecal, enteral, pulmonary, intralymphatic, intracavitary, vaginal, transurethral, intradermal, aural, intramammary, buccal, orthotopic, intratracheal, intralesional, percutaneous, endoscopic, transmucosal, sublingual and intestinal administration.

[00251] Pharmaceutically acceptable carriers for use in combination with the
30 compositions of the present disclosure are well known to those of ordinary skill in the art and are selected based upon a number of factors: the particular compound(s) and agent(s) used, and its/their concentration, stability and intended bioavailability; the

subject, its age, size and general condition; and the route of administration. Suitable nonaqueous, pharmaceutically-acceptable polar solvents include, but are not limited to, alcohols (e.g., α -glycerol formal, 6-glycerol formal, 1,3-butyleneglycol, aliphatic or aromatic alcohols having 2 to 30 carbon atoms such as methanol, ethanol, propanol, isopropanol, butanol, t-butanol, hexanol, octanol, amylene hydrate, benzyl alcohol, glycerin (glycerol), glycol, hexylene glycol, tetrahydrofurfuryl alcohol, lauryl alcohol, cetyl alcohol, or stearyl alcohol, fatty acid esters of fatty alcohols such as polyalkylene glycols (e.g., polypropylene glycol, polyethylene glycol), sorbitan, sucrose and cholesterol); amides (e.g., dimethylacetamide (DMA), benzyl benzoate DMA, dimethylformamide, N-(6-hydroxyethyl)-lactamide, N,N-dimethylacetamide amides, 2-pyrrolidinone, 1-methyl-2-pyrrolidinone, or polyvinylpyrrolidone); esters (e.g., 1-methyl-2-pyrrolidinone, 2-pyrrolidinone, acetate esters such as monoacetin, diacetin, and triacetin, aliphatic or aromatic esters such as ethyl caprylate or octanoate, alkyl oleate, benzyl benzoate, benzyl acetate, dimethylsulfoxide (DMSO), esters of glycerin such as mono, di-, or tri-glyceryl citrates or tartrates, ethyl benzoate, ethyl acetate, ethyl carbonate, ethyl lactate, ethyl oleate, fatty acid esters of sorbitan, fatty acid derived PEG esters, glyceryl monostearate, glyceride esters such as mono, di-, or tri-glycerides, fatty acid esters such as isopropyl myristate, fatty acid derived PEG esters such as PEG-hydroxyoleate and PEG-hydroxystearate, N-methyl pyrrolidinone, pluronic 60, polyoxyethylene sorbitol oleic polyester, polyoxyethylene sorbitan esters such as polyoxyethylene-sorbitan monooleate, polyoxyethylene-sorbitan monopalmitate, polyoxyethylene-sorbitan monolaurate, polyoxyethylene-sorbitan monostearate, and Polysorbate® 20, 40, 60 or 80 from ICI Americas, Wilmington, DE, polyvinylpyrrolidone, alkyleneoxy modified fatty acid esters such as polyoxyl 40 hydrogenated castor oil and polyoxyethylated castor oils (e.g., Cremophor® EL solution or Cremophor® RH 40 solution), saccharide fatty acid esters (i.e., the condensation product of a monosaccharide (e.g., pentoses such as ribose, ribulose, arabinose, xylose, lyxose and xylulose, hexoses such as glucose, fructose, galactose, mannose and sorbose, trioses, tetroses, heptoses, and octoses), disaccharide (e.g., sucrose, maltose, lactose and trehalose) or oligosaccharide or mixture thereof with a C_{4} to C_{22} fatty acid(s) (e.g., saturated fatty acids such as caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid and stearic acid, and unsaturated fatty acids such as palmitoleic acid, oleic acid, elaidic acid, erucic acid and linoleic acid)), or steroidal esters); alkyl, aryl, or cyclic ethers having 2 to 30 carbon atoms (e.g., diethyl ether, tetrahydrofuran, dimethyl isosorbide, diethylene glycol monoethyl ether); glycofuro

(tetrahydrofurfuryl alcohol polyethylene glycol ether); ketones having 3 to 30 carbon atoms (e.g., acetone, methyl ethyl ketone, methyl isobutyl ketone); aliphatic, cycloaliphatic or aromatic hydrocarbons having 4 to 30 carbon atoms (e.g., benzene, cyclohexane, dichloromethane, dioxolanes, hexane, n-decane, n-dodecane, n-hexane, sulfolane, tetramethylenesulfon, tetramethylenesulfoxide, toluene, di methylsulfoxide (DMSO), or tetramethylenesulfoxide); oils of mineral, vegetable, animal, essential or synthetic origin (e.g., mineral oils such as aliphatic or wax-based hydrocarbons, aromatic hydrocarbons, mixed aliphatic and aromatic based hydrocarbons, and refined paraffin oil, vegetable oils such as linseed, tung, safflower, soybean, castor, cottonseed, groundnut, rapeseed, coconut, palm, olive, corn, corn germ, sesame, persic and peanut oil and glycerides such as mono-, di- or triglycerides, animal oils such as fish, marine, sperm, cod-liver, haliver, squalene, squalane, and shark liver oil, oleic oils, and polyoxyethylated castor oil); alkyl or aryl halides having 1 to 30 carbon atoms and optionally more than one halogen substituent; methylene chloride; monoethanolamine; petroleum benzin; trolamine; omega-3 polyunsaturated fatty acids (e.g., alpha-linolenic acid, eicosapentaenoic acid, docosapentaenoic acid, or docosahexaenoic acid); polyglycol ester of 12-hydroxystearic acid and polyethylene glycol (Solutol® HS-15, from BASF, Ludwigshafen, Germany); polyoxyethylene glycerol; sodium laurate; sodium oleate; or sorbitan monooleate.

[00252] In some embodiments, oils or non-aqueous solvents may be employed in the formulations, e.g., to bring one or more of the compounds into solution, due to, for example, the presence of large lipophilic moieties. Alternatively, emulsions, suspensions, or other preparations, for example, liposomal preparations, may be used. With respect to liposomal preparations, for example, any known methods for preparing liposomes may be used. See, for example, Bangham et al., J. Mol. Biol, 23: 238-252 (1965) and Szoka et al., Proc. Natl Acad. Sci 75: 4194-4198 (1978), incorporated herein by reference. Thus, in one embodiment, one or more of the compounds are administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. Ligands may also be attached to the liposomes, for instance, to direct these compositions to particular sites of action.

[00253] Other pharmaceutically acceptable solvents for use in the pharmaceutical compositions described herein are well known to those of ordinary skill in

the art, and are identified in The Chemotherapy Source Book (Williams & Wilkins Publishing), The Handbook of Pharmaceutical Excipients, (American Pharmaceutical Association, Washington, D.C., and The Pharmaceutical Society of Great Britain, London, England, 1968), Modern Pharmaceutics, (G. Banker et al., eds., 3d ed.) (Marcel Dekker, Inc., New York, New York, 1995), The Pharmacological Basis of Therapeutics, (Goodman & Gilman, McGraw Hill Publishing), Pharmaceutical Dosage Forms, (H. Lieberman et al., eds.) (Marcel Dekker, Inc., New York, New York, 1980), Remington's Pharmaceutical Sciences (A. Gennaro, ed., 19th ed.) (Mack Publishing, Easton, PA, 1995), The United States Pharmacopeia 24, The National Formulary 19, (National Publishing, Philadelphia, PA, 2000), and A.J. Spiegel et al., Use of Nonaqueous Solvents in Parenteral Products, Journal of Pharmaceutical Sciences, Vol. 52, No. 10, pp. 917-927 (1963).

[00254] Formulations containing the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may take the form of solid, semi-solid, lyophilized powder, or liquid dosage forms such as, for instance, aerosols, capsules, creams, emulsions, foams, gels/jellies, lotions, ointments, pastes, powders, soaps, solutions, sprays, suppositories, suspensions, sustained-release formulations, tablets, tinctures, transdermal patches, and the like, preferably in unit dosage forms suitable for simple administration of precise dosages. If formulated as a fixed dose, such pharmaceutical compositions or formulation products employ the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) within accepted dosage ranges.

[00255] In one embodiment, a formulation is provided that contains the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) as a part of liquid dosage form, such as a sterile liquid dosage form suitable for injection. For example, the liquid form containing the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) in combination with one or more further ingredients, such as edetate disodium (EDTA). In one embodiment, the liquid form can comprise EDTA in an amount suitable to act as a preservative and/or metal-chelating agent, such as an amount of about 0.025%. The liquid form can further comprise water, and may also comprise a pH adjuster, such as sodium bicarbonate, for pH adjustment in the range of pH 5.5 to 7.0. In one embodiment, the pentaaza macrocyclic ring complex can also be provided as a part of a sterile liquid

dosage form suitable for injection, either in the same liquid dosage form with the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) or as a separate dosage form.

[00256] Formulations for certain pentaaza macrocyclic ring complexes are also described in, for example, in U.S. Patent Nos. 5,610,293, 5,637,578, 5,874,421, 5,976,498, 6,084,093, 6,180,620, 6,204,259, 6,214,817, 6,245,758, 6,395,725, and 6,525,041 (each of which is hereby incorporated herein by reference in its entirety).

[00257] It is contemplated that co-formulations of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may employ conventional formulation techniques for these components individually, or alternative formulation routes, subject to compatibility and efficacy of the various components, in combination.

[00258] The above-described pharmaceutical compositions including the pentaaza macrocyclic compound and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may additionally include one or more additional pharmaceutically active components. Suitable pharmaceutically active agents that may be included in the compositions of the present invention include, for instance, antiemetics, anesthetics, antihypertensives, antianxiety agents, ant clotting agents, anticonvulsants, blood glucose-lowering agents, decongestants, antihistamines, antitussives, antineoplastics, beta blockers, anti-inflammatory agents, antipsychotic agents, cognitive enhancers, cholesterol-reducing agents, antiobesity agents, autoimmune disorder agents, anti-impotence agents, antibacterial and antifungal agents, hypnotic agents, anti-Parkinsonism agents, anti-Alzheimer's Disease agents, antibiotics, anti-depressants, and antiviral agents. The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.

[00259] In yet another embodiment, a kit may be provided that includes both the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine), for treatment of a condition such as cancer or a viral infection. For example, the kit may comprise a first vessel or container having therein a formulation comprising the pentaaza macrocyclic ring complex, such as an oral or injectable formulation of the pentaaza macrocyclic ring

complex, and a second vessel or container having therein a formulation comprising the immunotherapeutic agent, such as an injectable formulation of an immune checkpoint inhibitor or other immunotherapeutic agent. The kit may further comprise a label or other instructions for administration of the active agents, recommended dosage amounts, durations and administration regimens, warnings, listing of possible drug-drug interactions, and other relevant instructions.

Combination Treatment with Cancer Therapy

[00260] In one embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can be administered in combination with another cancer therapy, to provide therapeutic treatment. For example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered as a part of at least one of a chemotherapy treatment and radiation therapy.

[00261] In general, the temporal aspects of the administration of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may depend for example, on the particular compound, radiation therapy, or chemotherapy that is selected, or the type, nature, and/or duration of the radiation exposure. Other considerations may include the disease or disorder being treated and the severity of the disease or disorder; activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors. For example, the compounds may be administered in various embodiments before, during, and/or after the administration of the cancer therapy (e.g., before, during or after exposure to and/or before, during or after a dose of chemotherapy, or before, during or after a course of radiation therapy or chemotherapy comprising multiple exposures and/or doses). By way of another example, the compounds may be administered in various embodiments before, during, and/or after an exposure to radiation.

[00262] If desired, the effective dose can be divided into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the dose.

[00263] In one embodiment, for example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient prior to or simultaneous with the cancer therapy corresponding to at least one of radiation therapy and chemotherapy, such as prior to or simultaneous with a dose or dose fraction of such treatment, or prior to or simultaneous with a course of such treatment comprising multiple doses. In another embodiment, for example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient prior to, but not after, the cancer therapy, such as before but not after a cancer therapy dose or dose fraction or prior to but not after a course of cancer therapy comprising multiple doses or dose fractions. In yet another embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient at least 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, prior to an initial dose or dose fraction of cancer therapy corresponding to at least one of radiation therapy and chemotherapy. In still other embodiments, for example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient after a dose or dose fraction of cancer therapy; thus, for example, the compound may be administered up to 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, or 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, after a single dose or dose fraction and/or final dose or dose fraction in a course of cancer treatment corresponding to one or more of radiation therapy and chemotherapy.

[00264] In another embodiment, for example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient prior to or

simultaneous with the radiation exposure. In another embodiment, for example, the components are administered to the patient prior to, but not after, the radiation exposure. In yet another embodiment, one or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient at least 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, prior to the radiation exposure, such as an initial radiation exposure in a course of radiation treatment, or prior to another dose or dose fraction of radiation that is one of the doses or dose fractions of radiation in the course of treatment. In still other embodiments, for example, pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered to the patient after the radiation exposure; thus, for example, the compound may be administered up to 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, or 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, after the radiation exposure, which may be a dose or dose fraction of radiation in a multi-dose course of radiation therapy, or may be the single or final dose or dose fraction of radiation in the radiation therapy.

[00265] In one embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered as a part of a course of therapy that includes the radiation therapy. In radiation therapy, a patient receives a dose or dose fraction of ionizing radiation to kill or control the growth of cancerous cells. The dose or dose fraction of radiation may be directed at a specific part of the body, and the beam of radiation may also be shaped according to a predetermined treatment regimen, to reduce deleterious effects on parts of the body not afflicted with cancer. A typical course of radiation therapy may include one or a plurality of doses or dose fractions of radiation, which can be administered over the course of days, weeks and even months. A total “dose” of radiation given during a course of radiation therapy typically refers to the amount of radiation a patient receives during the entire course of radiation therapy, which doses may be administered as dose “fractions” corresponding to multiple radiation

exposures in the case where the total dose is administered over several sessions, with the sum of the fractions administered corresponding to the overall dose. As is discussed in more detail in the Examples section below, the administration of pentaaza macrocyclic ring complex with the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can provide benefits treatment of cancer, thereby improving the efficacy of radiation treatment provided in combination with the immunotherapeutic agent.

[00266] In one embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined time period before or after a radiation exposure, such as a before or after a radiation dose or dose fraction. For example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered within 1 week, 48 hours, 24 hours, 12 hours, 6, hours, 2 hours, 1 hour or even within 30 minutes of the patient receiving the radiation exposure, such as the dose or dose fraction (either before or after the radiation exposure corresponding to the radiation dose or dose fraction). Other durations between the radiation exposure and administration of the compound that result in the enhanced the killing of cancer cells may also be suitable. In one embodiment, one or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered before the radiation exposure, and the remaining one or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can be administered after the radiation exposure. One or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may also be administered both before and after administration of a radiation exposure.

[00267] In one embodiment, a course of radiation therapy includes a plurality of radiation doses or dose fractions given over a predetermined period of time, such as over the course of hours, weeks, days and even months, with the plural doses or dose fractions being either of the same magnitude or varying. That is, a course of radiation therapy can comprise the administration of a series of multiple doses or dose fractions of

radiation. In one embodiment, pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can be administered before one or more radiation doses or dose fractions in the series, such as before each radiation dose or dose fraction, or before
5 some number of the radiation doses or dose fractions. Furthermore, the administration of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) during the course of radiation therapy can be selected to enhance the cancer treating effects of the radiation therapy, such as by sensitizing cancer cells to the radiation therapy. In one embodiment,
10 the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined duration before or after of each dose or dose fraction, such as the predetermined duration discussed above. In another embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint
15 inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined duration of time before or after only select doses or dose fractions. In yet another embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered within a predetermined duration of time before
20 the doses, while another of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered within a predetermined duration of time after the doses or dose fractions. In a further embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint
25 inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered only within the predetermined duration before or after select doses or dose fractions, while another of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered only within the predetermined duration before or after doses or dose fractions other than the
30 select doses or dose fractions.

[00268] A suitable overall dose to provide during a course of therapy can be determined according to the type of treatment to be provided, the physical characteristics of the patient and other factors, and the dose fractions that are to be provided can be

similarly determined. In one embodiment, a dose fraction of radiation that is administered to a patient may be at least 1.8 Gy, such as at least 2 Gy, and even at least 3 Gy, such as at least 5 Gy, and even at least 6 Gy. In yet another embodiment, a dose fraction of radiation that is administered to a patient may be at least 10 Gy, such as at least 12 Gy, and even at least 15 Gy, such as at least 18 Gy, and even at least 20 Gy, such as at least 24 Gy. In general, a dose fraction of radiation administered to a patient will not exceed 54 Gy. In some embodiments, the dose fraction of radiation administered to a patient may even be less than 10 Gy, and even less than 8 Gy, such as less than 5 Gy, or less than 3 Gy, including less than 2.5 Gy, less than 2 Gy, or about 1.8 Gy.

Furthermore, it should be noted that, in one embodiment, a dose fraction delivered to a subject may refer to an amount delivered to a specific target region of a subject, such as a target region of a tumor, whereas other regions of the tumor or surrounding tissue may be exposed to more or less radiation than that specified by the nominal dose fraction amount.

[00269] For example, in one embodiment, the overall dose of radiation provided during the course of therapy may be provided via a hypofractionation radiotherapy scheme, which typically involves providing relatively high dose fractions administered over relatively fewer sessions, as compared to lower dose fraction schemes. Examples of such hypofractionation radiotherapy methods can include, but are not limited to, stereotactic radiosurgery (SRS), which typically refers to a single-fraction treatment directed to targets such as intracranial and spinal targets, as well as stereotactic body radiation therapy (SBRT), which typically refers to multifractional treatment of targets such as intracranial and spinal targets, and also extracranial targets such as lung, liver, head and neck, pancreas and prostate. As an example, in one embodiment of a hypofractionation radiotherapy scheme, the overall dose of radiation provided during the course of therapy may be divided into less than 10 fractions, such as less than 8 fractions, less than 6 fractions, less than 5 fractions, less than 4 fractions, less than 3 fractions, less than 2 fractions and may even be provided in just one administration (single fraction). For example, in one embodiment, the overall dose of radiation provided during the course of therapy may be divided into from 1 to 10 fractions, such as from 1 to 6 fractions, and even from 1 to 5 fractions, such as from 2 to 5 fractions or even 2 to 4 fractions. As yet another example, the hypofractionation radiotherapy scheme can comprise dividing the overall dose of radiation provided during the course of therapy into

dose fractions that are at least 10% (1/10) of the overall dose provided during therapy, such as at least 12.5% (1/8) of the overall dose, at least 16% (~1/6) of the overall dose, at least 20% (1/5) of the overall dose, at least 25% (1/4) of the overall dose, at least 30% (1/3) of the overall dose, at least 50% of the overall dose, and/or at least 100% of the overall dose may be provided in a single administration (single fraction). For example, in one embodiment, the overall dose of radiation provided during the course of therapy may be divided into fractions that provide from 10% to 100% of the overall dose in each fraction, such as from 16% to 100% of the overall dose, and even from 20% to 100% of the overall dose, such as from 20% to 50% of the overall dose or even from 25% to 50% of the overall dose. For example a dose fraction size may be at least 5 Gy, such as at least 6 Gy, at least 8 Gy, at least 10 Gy, at least 12 Gy, and even at least 15 Gy, such as at least 18 Gy, and even at least 20 Gy, such as at least 24 Gy, and typically do not exceed 54 Gy, such as less than 40 Gy and even less than 30 Gy. In one embodiment, dose fraction sizes may be in the range of from 5 Gy to 30 Gy, such as from 6 Gy to 28 Gy, and even from 8 Gy to 25 Gy. Furthermore, in one embodiment, the dose fractions may be administered no more than three times per day, and even no more than twice per day, such as no more than once per day, on consecutive or non-consecutive days and/or some combination thereof, and may be administered over a period of a few days and up to a few weeks, such as over a period of 1 to 15 days, 1 to 12 days, 1 to 10 days, 1 to 5 days, and even 1 to 3 days. Typically, the dose fractions making up the overall course of therapy will be administered in no more than 20 days, no more than 15 days, no more than 10 days, no more than 5 days, and even no more than 3 days.

[00270] As yet another example, in one embodiment, the overall dose of radiation provided during the course of therapy may be provided via a radiotherapy scheme that provides relatively lower dose fractions administered over relatively more sessions, as compared to, e.g., hypofractionation schemes. Examples of such lower dose fraction radiotherapy methods can include, but are not limited to, intensity-modulated radiation therapy (IMRT) and image guided radiation therapy (IGRT), which typically involve three-dimensional conformal therapy (3D-CRT) to match the administered radiation to a target volume. As an example, in one embodiment of such a radiotherapy scheme, the overall dose of radiation provided during the course of therapy may be divided into at least 15 fractions, such as at least 18 fractions, at least 20 fractions, at least 22 fractions, at least 25 fractions, at least 28 fractions, at least 30

fractions, at least 32 fractions, at least 35 fractions, and even at least 38 fractions, although the total number of fractions will typically be less than 50, such as less than 45, and even less than 42. For example, in one embodiment, the overall dose of radiation provided during the course of therapy may be divided into from 15 to 38 fractions, such as from 20 to 38 fractions, and even from 20 to 35 fractions, such as from 25 to 35 fractions. As yet another example, the radiotherapy scheme can comprise dividing the overall dose of radiation provided during the course of therapy into dose fractions that are no more than 7% (1/15) of the overall dose provided during therapy, such as no more than 6% (1/18) of the overall dose, no more than 5% (1/20) of the overall dose, no more than 4.5% (1/22) of the overall dose, no more than 4% (1/25) of the overall dose, no more than 3.6% (1/28) of the overall dose, no more than 3.3% (1/30) of the overall dose, no more than 3.1% (1/32) of the overall dose, no more than 2.8% of the overall dose (1/35), and even no more than 2.6% (1/38) of the overall dose. For example, in one embodiment, the overall dose of radiation provided during the course of therapy may be divided into fractions that provide from 2.5% to 8% of the overall dose in each fraction, such as from 2.8% to 5% of the overall dose, and even from 2.8% to 4% of the overall dose. For example a dose fraction size may be less than 5 Gy, such as less than 4 Gy, less than 3.5 Gy, less than 3 Gy, less than 2.8 Gy, and even less than 2.5 Gy, such as less than 2.3 Gy, and even less than 2 Gy, such as less than 1.8 Gy, and typically is at least 0.5 Gy, such as at least 1 Gy and even at least 1.5 Gy. In one embodiment, dose fraction sizes may be in the range of from 1.5 Gy to 4.5 Gy, such as from 1.8 Gy to 3 Gy, and even from 2 Gy to 2.5 Gy. Furthermore, in one embodiment, the dose fractions may be administered no more than three times per day, and even no more than twice per day, such as no more than once per day, on consecutive or non-consecutive days, and/or a combination thereof (e.g., on consecutive weekdays), and in some embodiments may be administered over a period of a few days to a few weeks and even a few months, such as over a period of up to 3 weeks, up to 5 weeks, up to 6 weeks, up to 8 weeks and even up to 10 weeks, such as in a range of from 3 weeks to 10 weeks, or even in a range of from 5 weeks to 8 weeks. For example, the dose fractions making up the overall course of therapy can be administered in no more than 12 weeks, such as no more than 10 weeks and even no more than 8 weeks.

[00271] In yet another embodiment, the overall dose of radiation provided by the radiation scheme, whether in a relatively high dose fraction scheme or relatively low

dose fraction scheme such as those described above, or other scheme, is selected to provide suitable treatment of the cancer. The overall dose may also be provided according to the specific dose fractionation scheme being administered, along with other factors. For example, in certain embodiments, a relatively larger overall dose may be administered as relatively smaller individual dose fractions. In one embodiment, the overall dose provided over the course of the therapy (i.e., the sum of the administered dose fractions), is at least 50 Gy, such as at least 55 Gy, at least 58 Gy, at least 60 Gy, at least 65 Gy, at least 68 Gy, at least 70 Gy, at least 72 Gy, and even at least 75 Gy. In certain embodiments, the overall dose does not exceed 80 Gy, such as not exceeding 78 Gy and even not exceeding 75 Gy. For example, the overall dose may be in a range of from 50 Gy to 75 Gy, such as from 55 Gy to 75 Gy, and even from 60 Gy to 70 Gy.

[00272] In yet another embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered as a part of a course of therapy that includes chemotherapy. In chemotherapy, chemotherapeutic agents are administered to a patient to kill or control the growth of cancerous cells. A typical course of chemotherapy may include one or a plurality of doses of one or more chemotherapeutic agents, which can be administered over the course of days, weeks and even months.

Chemotherapeutic agents can include at least one of: alkylating antineoplastic agents such as nitrogen mustards (e.g. cyclophosphamide, chlorambucil), nitrosoureas (e.g. n-nitroso-n-methylurea, carmustine, semustine), tetrazines (e.g. dacarbazine, mitozolimide), aziridines (e.g. thiotepa, mytomicin), platinum-based antineoplastic agents (platinates) (e.g. cisplatin, carboplatin, oxaliplatin, neoplatin, platamin); anti-metabolites such as anti-folates (e.g. methotrexate and pemetrexed), fluoropyrimidines (e.g., fluorouracil, capecitabine), anthracyclines (e.g. doxorubicin, daunorubicin, epirubicin), deoxynucleoside analogs (e.g. cytarabine, gemcitabine, decitabine) and thiopurines (e.g., thioguanine, mercaptopurine); anti microtubule agents such as taxanes (e.g. paclitaxel, docetaxel); topoisomerase inhibitors (e.g. etoposide, doxorubicin, mitoxantrone, teniposide); and antitumor antibiotics (e.g. bleomycin, mitomycin). For example, the chemotherapeutic agent may be selected from the group consisting of all-trans retinoic acid, arsenic trioxide, azacitidine, azathioprine, bleomycin, carboplatin, capecitabine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, gemcitabine,

hydroxyurea, idarubicin, imatinib, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, teniposide, tiguane, valrubicin, vinblastine, vincristine, vindesine, and vinorelbine. The administration of many of the chemotherapeutic agents is described in the "Physicians' Desk Reference" (PDR), e.g.,
5 1996 edition (Medical Economics Company, Montvale, N.J. 07645-1742, USA).

[00273] In one embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered as a part of a course of therapy that includes a chemotherapeutic agent selected from the group consisting of cisplatin, doxorubicin,
10 bleomycin, and paclitaxel. Without being limited to any particular theory, it is believed that cisplatin, doxorubicin, bleomycin, and paclitaxel may contribute to the generation of superoxide radicals in cells, thereby leading when combined with a manganese pentaaza macrocyclic ring complex to increased oxidative stress and cytotoxicity of the cancer cells. Furthermore, in one embodiment, the chemotherapeutic agent may be selected
15 from the group consisting of a platinum-based antineoplastic agents, a taxane, an anticancer antibiotic, and an anthracycline, which categories of chemotherapeutic agents, without being limited to any particular theory or mechanism, may also be effective in providing chemotherapeutic activity at least in part due to generation of superoxide radicals in cells. Other chemotherapeutic agents that may increase superoxide levels can
20 include arsenic trioxide and 5-FU, which agents can also be used in the methods and compositions described herein. (Alexandre et al., Cancer Res. 67: (8), 3512-3517 (2007); Yen et al., J. Clin. Invest. 98 (5), 1253-1260 (1996); Masuda et al., Cancer Chemother. Pharmacol. 47(2), 155-160 (2001)).

[00274] According to yet another embodiment, a chemotherapeutic agent can
25 include at least one of an antimetabolite anti-cancer agents and antimitotic anti-cancer agents, and combinations thereof, which may include some of the agents described above and well as other agents described further herein. Various antimetabolite and antimitotic agents may be employed in the methods and compositions described herein.

[00275] Antimetabolic agents typically structurally resemble natural
30 metabolites, which are involved in normal metabolic processes of cancer cells such as the synthesis of nucleic acids and proteins. The antimetabolites, however, differ enough from the natural metabolites such that they interfere with the metabolic processes of cancer cells. In the cell, antimetabolites are mistaken for the metabolites they resemble,

and are processed by the cell in a manner analogous to the normal compounds. The presence of the “decoy” metabolites prevents the cells from carrying out vital functions and the cells are unable to grow and survive. For example, antimetabolites may exert cytotoxic activity by substituting these fraudulent nucleotides into cellular DNA, thereby
5 disrupting cellular division, or by inhibition of critical cellular enzymes, which prevents replication of DNA.

[00276] In one embodiment, therefore, the antimetabolite agent is a nucleotide or a nucleotide analog. In certain embodiments, for example, the antimetabolite agent may comprise purine (e.g., guanine or adenosine) or analogs thereof, or pyrimidine
10 (cytidine or thymidine) or analogs thereof, with or without an attached sugar moiety.

[00277] Suitable antimetabolite agents for use in the present disclosure may be generally classified according to the metabolic process they affect, and can include, but are not limited to, analogues and derivatives of folic acid, pyrimidines, purines, and cytidine. Thus, in one embodiment, the antimetabolite agent(s) is selected from the group
15 consisting of cytidine analogs, folic acid analogs, purine analogs, pyrimidine analogs, and combinations thereof.

[00278] In one particular embodiment, for example, the antimetabolite agent is a cytidine analog. According to this embodiment, for example, the cytidine analog may be selected from the group consisting of cytarabine (cytosine arabinoside), azacitidine
20 (5-azacytidine), and salts, analogs, and derivatives thereof.

[00279] In another particular embodiment, for example, the antimetabolite agent is a folic acid analog. Folic acid analogs or antifolates generally function by inhibiting dihydrofolate reductase (DHFR), an enzyme involved in the formation of nucleotides; when this enzyme is blocked, nucleotides are not formed, disrupting DNA
25 replication and cell division. According to certain embodiments, for example, the folic acid analog may be selected from the group consisting of denopterin, methotrexate (amethopterin), pemetrexed, pteropterin, raltitrexed, trimetrexate, and salts, analogs, and derivatives thereof.

[00280] In another particular embodiment, for example, the antimetabolite
30 agent is a purine analog. Purine-based antimetabolite agents function by inhibiting DNA synthesis, for example, by interfering with the production of purine containing nucleotides, adenine and guanine which halts DNA synthesis and thereby cell division. Purine

analogues can also be incorporated into the DNA molecule itself during DNA synthesis, which can interfere with cell division. According to certain embodiments, for example, the purine analogue may be selected from the group consisting of acyclovir, allopurinol, 2-aminoadenosine, arabinosyl adenine (ara-A), azacitidine, azathioprine, 8-aza-adenosine, 5 8-fluoro-adenosine, 8-methoxy-adenosine, 8-oxo-adenosine, cladribine, deoxycoformycin, fludarabine, gancyclovir, 8-aza-guanosine, 8-fluoro-guanosine, 8-methoxy-guanosine, 8-oxo-guanosine, guanosine diphosphate, guanosine diphosphate-beta-L-2-aminofucose, guanosine diphosphate-D-arabinose, guanosine diphosphate-2-fluorofucose, guanosine diphosphate fucose, mercaptopurine (6-MP), pentostatin, thiamiprine, thioguanine (6-TG), 10 and salts, analogues, and derivatives thereof.

[00281] In yet another particular embodiment, for example, the antimetabolite agent is a pyrimidine analogue. Similar to the purine analogues discussed above, pyrimidine-based antimetabolite agents block the synthesis of pyrimidine-containing nucleotides (cytosine and thymine in DNA; cytosine and uracil in RNA). By acting as "decoys," the 15 pyrimidine-based compounds can prevent the production of nucleotides, and/or can be incorporated into a growing DNA chain and lead to its termination. According to certain embodiments, for example, the pyrimidine analogue may be selected from the group consisting of ancitabine, azacitidine, 6-azauridine, bromouracil (e.g., 5-bromouracil), capecitabine, carmofur, chlorouracil (e.g. 5-chlorouracil), cytarabine (cytosine 20 arabinoside), cytosine, dideoxyuridine, 3'-azido-3'-deoxythymidine, 3'-dideoxycytidin-2'-ene, 3'-deoxy-3'-deoxythymidin-2'-ene, dihydrouracil, doxifluridine, enocitabine, floxuridine, 5-fluorocytosine, 2-fluorodeoxycytidine, 3-fluoro-3'-deoxythymidine, fluorouracil (e.g., 5-fluorouracil (also known as 5-FU), gemcitabine, 5-methylcytosine, 5-propynylcytosine, 5-propynylthymine, 5-propynyluracil, thymine, uracil, uridine, and salts, 25 analogues, and derivatives thereof. In one embodiment, the pyrimidine analogue is other than 5-fluorouracil. In another embodiment, the pyrimidine analogue is gemcitabine or a salt thereof.

[00282] In certain embodiments, the antimetabolite agent is selected from the group consisting of 5-fluorouracil, capecitabine, 6-mercaptopurine, methotrexate, 30 gemcitabine, cytarabine, fludarabine, pemetrexed, and salts, analogues, derivatives, and combinations thereof. In other embodiments, the antimetabolite agent is selected from the group consisting of capecitabine, 6-mercaptopurine, methotrexate, gemcitabine, cytarabine, fludarabine, pemetrexed, and salts, analogues, derivatives, and combinations

thereof. In one particular embodiment, the antimetabolite agent is other than 5-fluorouracil. In a particularly preferred embodiment, the antimetabolite agent is gemcitabine or a salt or thereof (e.g., gemcitabine HCl (Gemzar®)).

[00283] Other antimetabolite agents may be selected from, but are not limited to, the group consisting of acanthifolic acid, aminothiadiazole, brequinar sodium, Ciba-Geigy CGP-30694, cyclopentyl cytosine, cytarabine phosphate stearate, cytarabine conjugates, Lilly DATHF, Merrel Dow DDFC, dezaguanine, dideoxycytidine, dideoxyguanosine, didox, Yoshitomi DMDC, Wellcome EHNA, Merck & Co. EX-015, fazarabine, fludarabine phosphate, N-(2'-furanidyl)-5-fluorouracil, Daiichi Seiyaku FO-152, 5-FU-fibrinogen, isopropyl pyrrolizine, Lilly LY-188011; Lilly LY-264618, methobenzaprim, Wellcome MZPES, norspermidine, NCI NSC-127716, NCI NSC-264880, NCI NSC-39661, NCI NSC-612567, Warner-Lambert PALA, pentostatin, piritrexim, plicamycin, Asahi Chemical PL-AC, Takeda TAC-788, tiazofurin, Erbamont TIF, tyrosine kinase inhibitors, Taiho UFT and uricytin, among others.

[00284] In one embodiment, the chemotherapeutic agent comprises an antimitotic agent that is a microtubule inhibitor or a microtubule stabilizer. In general, microtubule stabilizers, such as taxanes (some of which are also described above) and epothilones, bind to the interior surface of the beta-microtubule chain and enhance microtubule assembly by promoting the nucleation and elongation phases of the polymerization reaction and by reducing the critical tubulin subunit concentration required for microtubules to assemble. Unlike microtubule inhibitors, such as the vinca alkaloids, which prevent microtubule assembly, the microtubule stabilizers, such as taxanes, decrease the lag time and dramatically shift the dynamic equilibrium between tubulin dimers and microtubule polymers towards polymerization. In one embodiment, therefore, the microtubule stabilizer is a taxane or an epothilone. In another embodiment, the microtubule inhibitor is a vinca alkaloid.

[00285] One element of the therapy described herein may include the use of a taxane or derivative or analog thereof, some of which have also been discussed above. In one embodiment, the taxane may be a naturally derived compound or a related form, or may be a chemically synthesized compound or a derivative thereof, with antineoplastic properties. The taxanes are a family of terpenes, including, but not limited to paclitaxel (Taxol®) and docetaxel (Taxotere®), which are derived primarily from the Pacific yew tree, *Taxus brevifolia*, and which have activity against certain tumors, particularly breast

and ovarian tumors. In one embodiment, the taxane is docetaxel or paclitaxel. Paclitaxel is a preferred taxane and is considered an antimitotic agent that promotes the assembly of microtubules from tubulin dimers and stabilizes microtubules by preventing depolymerization. This stability results in the inhibition of the normal dynamic
5 reorganization of the microtubule network that is essential for vital interphase and mitotic cellular functions.

[00286] Also included are a variety of known taxane derivatives, including both hydrophilic derivatives, and hydrophobic derivatives. Taxane derivatives include, but are not limited to, galactose and mannose derivatives described in International Patent
10 Application No. WO 99/18113; piperazino and other derivatives described in WO 99/14209; taxane derivatives described in WO 99/09021, WO 98/22451, and U.S. Patent No. 5,869,680; 6-thio derivatives described in WO 98/28288; sulfenamide derivatives described in U.S. Patent No. 5,821,263; deoxygenated paclitaxel compounds such as those described in U.S. Patent No. 5,440,056; and taxol derivatives described in U.S.
15 Patent No. 5,415,869. As noted above, it further includes prodrugs of paclitaxel including, but not limited to, those described in WO 98/58927; WO 98/13059; and U.S. Patent No. 5,824,701. The taxane may also be a taxane conjugate such as, for example, paclitaxel-PEG, paclitaxel-dextran, paclitaxel-xylose, docetaxel-PEG, docetaxel-dextran, docetaxel-xylose, and the like. Other derivatives are mentioned in "Synthesis and Anticancer
20 Activity of Taxol Derivatives," D. G. I. Kingston et al., Studies in Organic Chemistry, vol. 26, entitled "New Trends in Natural Products Chemistry" (1986), Atta-ur-Rabman, P. W. le Quesne, Eds. (Elsevier, Amsterdam 1986), among other references. Each of these references is hereby incorporated by reference herein in its entirety.

[00287] Various taxanes may be readily prepared utilizing techniques known to
25 those skilled in the art (see also WO 94/07882, WO 94/07881, WO 94/07880, WO 94/07876, WO 93/23555, WO 93/10076; U.S. Pat. Nos. 5,294,637; 5,283,253; 5,279,949; 5,274,137; 5,202,448; 5,200,534; 5,229,529; and EP 590,267) (each of which is hereby incorporated by reference herein in its entirety), or obtained from a variety of commercial sources, including for example, Sigma-Aldrich Co., St. Louis, MO.

30 [00288] Alternatively, the antimitotic agent can be a microtubule inhibitor; in one preferred embodiment, the microtubule inhibitor is a vinca alkaloid. In general, the vinca alkaloids are mitotic spindle poisons. The vinca alkaloid agents act during mitosis when chromosomes are split and begin to migrate along the tubules of the mitosis spindle

towards one of its poles, prior to cell separation. Under the action of these spindle poisons, the spindle becomes disorganized by the dispersion of chromosomes during mitosis, affecting cellular reproduction. According to certain embodiments, for example, the vinca alkaloid is selected from the group consisting of vinblastine, vincristine, vindesine, vinorelbine, and salts, analogs, and derivatives thereof.

[00289] The antimitotic agent can also be an epothilone. In general, members of the epothilone class of compounds stabilize microtubule function according to mechanisms similar to those of the taxanes. Epothilones can also cause cell cycle arrest at the G2-M transition phase, leading to cytotoxicity and eventually apoptosis. Suitable epothilones include epothilone A, epothilone B, epothilone C, epothilone D, epothilone E, and epothilone F, and salts, analogs, and derivatives thereof. One particular epothilone analog is an epothilone B analog, ixabepilone (Ixempra™).

[00290] In certain embodiments, the antimitotic anti-cancer agent is selected from the group consisting of taxanes, epothilones, vinca alkaloids, and salts and combinations thereof. Thus, for example, in one embodiment the antimitotic agent is a taxane. More preferably in this embodiment the antimitotic agent is paclitaxel or docetaxel, still more preferably paclitaxel. In another embodiment, the antimitotic agent is an epothilone (e.g., an epothilone B analog). In another embodiment, the antimitotic agent is a vinca alkaloid.

[00291] In one embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined time period before or after a dose of a chemotherapeutic agent is administered. For example, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered within 1 week, 48 hours, 24 hours, 12 hours, 6 hours, 2 hours, 1 hour or even within 30 minutes of the patient receiving the dose of chemotherapeutic agent (either before or after the dose of chemotherapeutic agent). Other durations between the chemotherapeutic agent dose and administration of the components that result in the enhanced the killing of cancer cells may also be suitable. In one embodiment, one or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may be administered before the dose of the chemotherapeutic agent, and the remaining one or

more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) can be administered after the dose of the chemotherapeutic agent. One or more of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint
5 inhibitor, adoptive T-cell transfer therapy, cancer vaccine) may also be administered both before and after administration of the dose of chemotherapeutic agent.

[00292] In one embodiment, a course of chemotherapy includes a singular dose of a chemotherapeutic agent. In another embodiment, a course of chemotherapy includes a plurality of doses of a chemotherapeutic agent given over a predetermined
10 period of time, such as over the course of hours, weeks, days and even months. The plural doses may be either of the same magnitude or varying, and can include doses of the same or different chemotherapeutic agents and/or a combination of chemotherapeutic agents. The administration of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer
15 therapy, cancer vaccine) during the course of chemotherapy can be selected to enhance the cancer treating effects of the chemotherapy, such as by increasing intracellular levels of hydrogen peroxide to promote oxidative stress in the cancer cells. In one embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered
20 within a predetermined duration before or after each dose, such as the predetermined duration discussed above. In another embodiment, the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined duration of time before or after only select doses. In yet another embodiment, at least one of the
25 pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a predetermined duration of time before the doses, while another of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) are administered within a
30 predetermined duration of time after the doses. In a further embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered only within the predetermined duration before or after select doses, while another of the

pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered only within the predetermined duration before or after doses other than the select doses.


[00293] In yet another embodiment, at least one of the pentaaza macrocyclic ring complex and the immunotherapeutic agent (e.g., immune checkpoint inhibitor, adoptive T-cell transfer therapy, cancer vaccine) is administered in combination with both a radiation therapy and chemotherapy.

[00294] Embodiments according to aspects of the disclosure are provided below, although the disclosure is not limited thereto.

[00295] Embodiment 1. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

[00296] administering to the subject an immune checkpoint inhibitor;

[00297] administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after administration of the immune checkpoint inhibitor, to increase the response of the cancer to the immune checkpoint inhibitor:



[00298]

[00299] wherein

[00300] M is Mn^{2+} or Mn^{3+} ;

[00301]

 $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9,$ and R_{10}

are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$

5 , $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein

R_{11} and R_{12} are independently hydrogen or alkyl;

[00302]

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [00303]

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00304]

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-
15 containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

20 [00305]

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00306]

Z is a counterion;

[00307]

n is an integer from 0 to 3; and

25 [00308]

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00309]


Embodiment 2. The method according to Embodiment 1, wherein $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9,$ and R_{10} are each hydrogen.

[00310]

Embodiment 3. The method according to Embodiment 1 or 2, wherein
30 W is an unsubstituted pyridine moiety.

[00311] Embodiment 4. The method according to any preceding Embodiment, wherein U and V are transcyclohexanyl fused rings.

[00312] Embodiment 5. The method according to any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is represented by formula (II):



5 [00313]



[00314] wherein

[00315] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

10 [00316] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein

15 R_{11} and R_{12} are independently hydrogen or alkyl.

[00317] Embodiment 6. The method according to any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):




[00318] wherein


[00319] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding
5 anion thereof; and

[00320] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting
of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$
10 , $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl.


[00321] Embodiment 7. The method according to any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is a compound represented by a formula selected from the group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)




(VIII)



(IX)




(X)




[00322]

(XI)




(XII)




[00323]

(XIII)




(XIV)

[00324]



(XV)

[00331]



[00332]

(XVI)

[00333] Embodiment 8. The method according to any preceding Embodiment,

5 wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl

10 nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl

15 thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate,

20 thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate,

25 bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino

acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

[00334] or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

5 [00335] each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

[00336] each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

[00337] each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl,
10 amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

[00338] each X_4 is independently hydrogen or together with X_3 is $(=O)$;

15 [00339] or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

[00340] or X and Y are independently attached to one or more of $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} .


20 [00341] Embodiment 9. The method according to any preceding Embodiment, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

[00342] Embodiment 10. The method according to any one of Embodiments 1-8, wherein X and Y are independently selected from the group consisting of alkyl
25 carboxylates, aryl carboxylates and arylalkyl carboxylates.

[00343] Embodiment 11. The method according to any one of Embodiments 1-8, wherein X and Y are independently amino acids.

[00344] Embodiment 12. The method according to any one of Embodiments 1-8 Embodiment, wherein the pentaaza macrocyclic ring complex is a compound
30 represented by the formula:


[00345]



[00346]


[00347] Embodiment 13. The method according to any one of Embodiments 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the 5 formula:

[00348]




[00349]

[00350] Embodiment 14. The method according to any one of Embodiment s 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the 10 formula:




[00351]

[00352] Embodiment 15. The method according to any one of Embodiments 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



[00353]

[00354] Embodiment 16. The method according to any one of Embodiments 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:


**GC4702**

5

[00355]

[00356] Embodiment 17. The method according to any one of Embodiments 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:

[00357]

**GC4711**

[00358]

[00359] Embodiment 18. The method according to any preceding Embodiment, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is administered a predetermined period of time after initial administration of the immune checkpoint inhibitor.

5 [00360] Embodiment 19. The method according to Embodiment 18, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is no less than 3 days after initial administration of the immune checkpoint inhibitor.

[00361] Embodiment 20. The method according to Embodiment 19, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is
10 no less than 6 days after initial administration of the immune checkpoint inhibitor.

[00362] Embodiment 21. The method according to Embodiment 19, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is in a range of from 3 days to 9 weeks after initial administration of the immune checkpoint inhibitor.

15 [00363] Embodiment 22. The method according to any preceding Embodiment, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows two doses of the immune checkpoint inhibitor.

[00364] Embodiment 23. The method according to Embodiment 22, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy
20 follows three doses of the immune checkpoint inhibitor.

[00365] Embodiment 24. The method according to Embodiment 23, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows four doses of the immune checkpoint inhibitor.

[00366] Embodiment 25. The method according to Embodiment 24, wherein
25 initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows five doses of the immune checkpoint inhibitor.

[00367] Embodiment 26. The method according to any preceding Embodiment, wherein doses of the pentaaza macrocyclic ring complex provided in a course of cancer therapy are provided on separate days from any dose of the immune
30 checkpoint inhibitor.

[00368] Embodiment 27. The method according to any preceding Embodiment, further comprising administering one or more of radiation therapy and chemotherapy to the subject, prior to, concomitantly with, or after administration of one or more of the immune checkpoint inhibitor and pentaaza macrocyclic ring complex.

5 [00369] Embodiment 28. The method according to Embodiment 27, wherein radiation therapy is administered concomitantly with administration of one or more of the immune checkpoint inhibitor and pentaaza macrocyclic ring complex.

[00370] Embodiment 29. The method according to any preceding Embodiment, comprising administering the pentaaza macrocyclic ring complex to a
10 subject that is not receiving radiation therapy.

[00371] Embodiment 30. The method according to any preceding Embodiment, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

[00372] Embodiment 31. The method according to any preceding
15 Embodiment, wherein a course of therapy comprising administration of the pentaaza macrocyclic ring complex and the immune checkpoint inhibitor, is administered to a subject that does not receive radiation therapy during the course of therapy.

[00373] Embodiment 32. The method according to any of Embodiments 1-28, comprising administering one or more of the pentaaza macrocyclic ring complex and
20 immune checkpoint inhibitor to the subject on a day other than a day that the subject is receiving radiation therapy.

[00374] Embodiment 33. The method according to any preceding Embodiment, comprising administering a course of therapy comprising administration of the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that
25 has not received radiation therapy for at least a day.

[00375] Embodiment 34. The method according to any preceding Embodiment, comprising administering a course of therapy comprising administration of the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that has not received radiation therapy for at least a week.

30 [00376] Embodiment 35. The method according to any preceding Embodiment, comprising administering a course of therapy comprising administration of

the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that has not received radiation therapy for at least a month.

[00377] Embodiment 36. The method according to any preceding Embodiment, comprising administering a course of therapy comprising administration of
5 the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that has not received radiation therapy for at least six months.

[00378] Embodiment 37. The method according to any preceding Embodiment, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject, and delaying any radiation therapy optionally
10 administered to the subject thereafter by at least one day after a final administration of the pentaaza macrocyclic ring complex.

[00379] Embodiment 38. The method according to any preceding Embodiment, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject, and delaying any radiation therapy optionally
15 administered to the subject thereafter by at least one week after a final administration of the pentaaza macrocyclic ring complex.

[00380] Embodiment 39. The method according to any preceding Embodiment, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject, and delaying any radiation therapy optionally
20 administered to the subject thereafter by at least one month after a final administration of the pentaaza macrocyclic ring complex.

[00381] Embodiment 40. The method according to any preceding Embodiment, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject, and delaying any radiation therapy optionally
25 administered to the subject thereafter by at least six months after a final administration of the pentaaza macrocyclic ring complex.

[00382] Embodiment 41. The method according to any preceding Embodiment, wherein the checkpoint inhibitor interacts with one or more of cytotoxic T-lymphocyte antigen 4 (CTLA4), programmed death 1 (PD-1), programmed death ligand 1
30 (PDL-1), PDL-2, lymphocyte activation genes-3 (LAG3), B7 homolog 3 (B7-H3), B7 homolog 4 (B7-H4), indoleamine (2,3)-dioxygenase (IDO), adenosine A2a receptor (A2AR), neuritin, B- and T-lymphocyte attenuator (BTLA), killer immunoglobulin-like

receptors (KIR), T cell immunoglobulin and mucin domain-containing protein 3 (TIME-3), inducible T cell costimulator (ICOS), CD27, CD28, CD40, CD137, CD160, CD244, HVEM, GAL9, VISTA, 2B4, CGEN-15049, CHK 1, CHK 2, GITR, CD47 and combinations thereof.

5 [00383] Embodiment 42. The method according to any preceding Embodiment, wherein the checkpoint inhibitor comprises one or more of small molecule inhibitor, an antibody, an antigen binding fragment, and an Ig fusion protein.

 [00384] Embodiment 43. The method according to any preceding Embodiment, wherein the checkpoint inhibitor is selected from the group consisting of
10 ipilimumab, nivolumab, pembrolizumab, pidilizumab, areluman, tremelimumab, atezolizumab, AMP-224, MPDL3280A, MDX-1105, MDX-1106, MEDI-4736, IMP321, INCB024360, NLG-919, indoximod, AUNP 12, galiximab, avelumab, varlilumab, mogamulizumab, CP-870,893, MEDI-6469, IPH2101, urelumab, lirilumab, BMS-986016, MGA271, IMP321, BMS-936559, MSB0010718C, anti-OX40, MK-3475, CT-011, BY55,
15 AMP224, and BGB-A317.

 [00385] Embodiment 44. The method according to any preceding Embodiment, wherein the checkpoint inhibitor is at least one of an anti-CTLA4 antibody, an anti-PD-1 antibody and an anti-PDL-1 antibody.

 [00386] Embodiment 45. The method according to any preceding
20 Embodiment, further comprising administering one or more of adoptive T-cell transfer therapy and a cancer vaccine to the subject, either prior to, concomitantly with, or after administration of one or more of the checkpoint inhibitor and pentaaza macrocyclic ring complex.

 [00387] Embodiment 46. The method according to any preceding
25 Embodiment, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.

 [00388] Embodiment 47. The method according to any preceding
30 Embodiment, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 40 mg/kg.

[00389] Embodiment 48. The method according to Embodiment 47, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 24 mg/kg.

[00390] Embodiment 49. The method according to Embodiment 48, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 10 mg/kg.

[00391] Embodiment 50. The method according to any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.


[00392] Embodiment 51. The method according to Embodiment 40, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.

[00393] Embodiment 52. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

[00394] administering to the subject an adoptive T-cell transfer therapy;

[00395] administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after the adoptive T-cell transfer therapy, to increase the response of the cancer to the adoptive T-cell transfer therapy,

[00396]



[00397]

[00398] wherein

[00399] M is Mn^{2+} or Mn^{3+} ;

[00400]

R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀

are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂

5 , -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein

R₁₁ and R₁₂ are independently hydrogen or alkyl;

[00401]

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [00402]

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00403]

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-
15 containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

20 [00404]

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00405]

Z is a counterion;

[00406]

n is an integer from 0 to 3; and

25 [00407]

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00408]


Embodiment 53. The method according to Embodiment 52, wherein R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each hydrogen.

[00409]

Embodiment 54. The method according to Embodiment 52 or 53,
30 wherein W is an unsubstituted pyridine moiety.

[00410] Embodiment 55. The method according to any of Embodiments 52-54, wherein U and V are transcyclohexanyl fused rings.

[00411] Embodiment 56. The method according to any of Embodiments 52-55, wherein the pentaaza macrocyclic ring complex is represented by formula (II):



5 [00412]



[00413] wherein

[00414] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

10 [00415] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein

15 R_{11} and R_{12} are independently hydrogen or alkyl.

[00416] Embodiment 57. The method according to any of Embodiments 52-56, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):




[00417] wherein


[00418] X and Y represent suitable ligands which are derived from any
5 monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

[00419] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting
10 of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl.


[00420] Embodiment 58. The method according to any of Embodiments 52-57, wherein the pentaaza macrocyclic ring complex is a compound represented by a formula
15 selected from the group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)




(VIII)



(IX)




(X)




[00421]

(XI)




(XII)

[00422]




[00423]

(XIII)




(XIV)

[00424]



[00431]

(XV)



[00432]

[00433]

(XVI)

[00434] Embodiment 59. The method according to any of Embodiments 52-58, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino

acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

[00435] or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

5 [00436] each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

[00437] each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

[00438] each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl,
10 amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

[00439] each X_4 is independently hydrogen or together with X_3 is $(=O)$;

15 [00440] or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

[00441] or X and Y are independently attached to one or more of $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} .


20 [00442] Embodiment 60. The method according to any of Embodiments 52-59, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

[00443] Embodiment 61. The method according to any one of Embodiments 52-59, wherein X and Y are independently selected from the group consisting of alkyl
25 carboxylates, aryl carboxylates and arylalkyl carboxylates.

[00444] Embodiment 62. The method according to any one of Embodiments 52-59, wherein X and Y are independently amino acids.

[00445] Embodiment 63. The method according to any one of Embodiments 52-59, wherein the pentaaza macrocyclic ring complex is a compound represented by the
30 formula:

[00446]




[00447]

[00448] Embodiment 64. The method according to any one of Embodiments 52-62, wherein the pentaaza macrocyclic ring complex is a compound represented by the

5 formula:


[00449]



[00450]


[00451] Embodiment 65. The method according to any one of Embodiments 52-62, wherein the pentaaza macrocyclic ring complex is a compound represented by the

10 formula:



[00452]

[00453] Embodiment 66. The method according to any one of Embodiments 52-62, wherein the pentaaza macrocyclic ring complex is represented by the formula:




[00454]

[00455] Embodiment 67. The method according to any one of Embodiments 52-62, wherein the pentaaza macrocyclic ring complex is represented by the formula:

5


[00456]



[00457]

[00458] Embodiment 68. The method according to any one of Embodiments 52-62, wherein the pentaaza macrocyclic ring complex is represented by the formula:

[00459]



[00460]

GC4711

[00461] Embodiment 69. The method according to any of Embodiments 52-68, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is a predetermined period of time after initial administration of the adoptive T-cell transfer therapy.

[00462] Embodiment 70. The method according to any of Embodiments 52-68, further comprising administering one or more of radiation therapy and chemotherapy to the subject, prior to, concomitantly with, or after administration of one or more of the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex.

10 [00463] Embodiment 71. The method according to any of Embodiments 52-68, comprising administering the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

[00464] Embodiment 72. The method according to any of Embodiments 52-71, wherein the adoptive T-cell transfer therapy comprises administering to the subject
15 cancer-specific autologous or allogeneic T-cells.

[00465] Embodiment 73. The method according to any of Embodiments 52-72, wherein the adoptive T-cell transfer therapy comprises providing autologous tumor infiltrating lymphocytes, antigen-expanded CD8⁺ and/or CD4⁺ T cells, and genetically modified T cells that express T-cell receptors (TCR) that recognize tumor antigens.

20 [00466] Embodiment 74. The method according to any of Embodiment s 52-73, further comprising administering one or more of an immune checkpoint inhibitor and a cancer vaccine to the subject, either prior to, concomitantly with, or after administration of

one or more of the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex.

[00467] Embodiment 75. The method according to any of Embodiments 52-74, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.

[00468] Embodiment 76. The method according to any of Embodiments 52-75, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.


[00469] Embodiment 77. The method according to Embodiment 76, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.

[00470] Embodiment 78. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

[00471] administering to the subject a cancer vaccine;

[00472] administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after administration of the cancer vaccine, to increase the response of the cancer to the cancer vaccine,

[00473]



[00474]

[00475] wherein

[00476] M is Mn^{2+} or Mn^{3+} ;

[00477]

R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀

are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂

5 , -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein

R₁₁ and R₁₂ are independently hydrogen or alkyl;

[00478]

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [00479]

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00480]

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-
15 containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

20 [00481]

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00482]

Z is a counterion;

[00483]

n is an integer from 0 to 3; and

25 [00484]

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00485] Embodiment 79. The method according to Embodiment 78, wherein


R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each hydrogen.

[00486] Embodiment 80. The method according to Embodiment 78 or 79,

30 wherein W is an unsubstituted pyridine moiety.

[00487] Embodiment 81. The method according to any of Embodiments 78-80, wherein U and V are transcyclohexanyl fused rings.

[00488] Embodiment 82. The method according to any of Embodiments 78-69, wherein the pentaaza macrocyclic ring complex is represented by formula (II):



5


[00489] wherein

[00490] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

10 [00491] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein

15 R_{11} and R_{12} are independently hydrogen or alkyl.

[00492] Embodiment 83. The method according to any of Embodiments 78-82, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):




[00493] wherein


[00494] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding
5 anion thereof; and

[00495] R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting
of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$
10 , $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl.


[00496] Embodiment 84. The method according to any of Embodiments 78-82, wherein the pentaaza macrocyclic ring complex is a compound represented by a formula selected from the group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)



(VIII)




(IX)




(X)

[00497]




[00498]

(XI)




(XII)




[00499]

(XIII)



(XIV)


[00500]



[00501]

(XV)

[00502]



[00503]

[00504]

(XVI)

5 [00505] Embodiment 85. The method according to any of Embodiments 78-84, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, 10 nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl 15 carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl 20 phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, 25 bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate,

tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

[00506] or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

[00507] each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

[00508] each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

10 [00509] each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

[00510] each X_4 is independently hydrogen or together with X_3 is $(=O)$;

[00511] or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

[00512] or X and Y are independently attached to one or more of
20 $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} .


[00513] Embodiment 86. The method according to any of Embodiments 78-85, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

[00514] Embodiment 87. The method according to any one of Embodiments
25 78-85, wherein X and Y are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates.

[00515] Embodiment 88. The method according to any one of Embodiments 78-85, wherein X and Y are independently amino acids.

[00516] Embodiment 89. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:


[00517]



5 [00518]


[00519] Embodiment 90. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:

[00520]




10 [00521]

[00522] Embodiment 91. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




[00523]

[00524] Embodiment 92. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is represented by the formula:



5 [00525] Embodiment 93. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is represented by the formula:


[00526]



[00527]

[00528] Embodiment 94. The method according to any one of Embodiments 78-85, wherein the pentaaza macrocyclic ring complex is represented by the formula:

[00529]



[00530]

GC4711

[00531] Embodiment 95. The method according to any of Embodiments 78-94, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is a predetermined period of time after initial administration of the cancer vaccine.

5 [00532] Embodiment 96. The method according to any of Embodiments 78-95, further comprising administering one or more of radiation therapy and chemotherapy to the subject, prior to, concomitantly with, or after administration of one or more of the cancer vaccine and pentaaza macrocyclic ring complex.

[00533] Embodiment 97. The method according to any of Embodiment s 78-10 96, comprising administering the cancer vaccine and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

[00534] Embodiment 98. The method according to any of Embodiments 78-97, wherein the cancer vaccine is selected from the group consisting of tumor cell vaccines, antigen vaccines, dendritic cell vaccines, DNA vaccines and vector based vaccines.

15 [00535] Embodiment 99. The method according to any of Embodiments 78-98, wherein the cancer vaccine is selected from the group consisting of M-Vax (Avax Technologies) , Provenge (Dendreon), GRNVAC1 (Geron), Bexidex (IDM Pharma), Uvidex (IDM Pharma), Collidex (IDM Pharma), INGN 225 (Introgen Therapeutics), M3Tk (MolMed), DC-Vax (Northwest Biotherapeutics), CVax (Prima Biomed), GVAX (Cell
20 Genesys), Lucanix (NovaRx), Onyvax-P (Onyvax), HSPP-96 Oncophage (Antigenics), BiovaxID (Biovest International), NeuVax (Apthera), CDX-110 (CeppDex), GV1001 (Pharmexa), CYT004-MelQbG10 (Cytos Biotechnology), li-Key/HER2/neu (Generex Biotechnology), MAGE-A3 (Glaxo-SmithKline Biologicals), IDM-2101 (IDM Pharma),

IMA901/IMA910 (Immatics Biotechnologies), melanoma cancer vaccine (Norwood Immunology), inCVAX (Immunophotonics) and Stimuvax (Oncothyreon).

[00536] Embodiment 100. The method according to any of Embodiments 78-99, further comprising administering one or more of an immune checkpoint inhibitor and an adoptive T-cell transfer therapy to the subject, either prior to, concomitantly with, or after administration of one or more of the cancer vaccine and pentaaza macrocyclic ring complex.

[00537] Embodiment 101. The method according to any of Embodiments 78-100, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.

[00538] Embodiment 102. The method according to any of Embodiments 78-101, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.


[00539] Embodiment 103. The method according to Embodiment 102, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.

[00540] Embodiment 104. A method of treating a viral infection in a mammalian subject in need thereof, comprising.

[00541] administering to the subject at least one of an immune checkpoint inhibitor, an adoptive T-cell transfer therapy, and a vaccine; and

[00542] administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after the at least one immune checkpoint inhibitor, adoptive T-cell transfer therapy, and vaccine, to increase the effectiveness of the at least one immune checkpoint, adoptive T-cell transfer therapy, and vaccine in treating the viral infection,

[00543]



[00544] W (I)

[00545] wherein

[00546] M is Mn^{2+} or Mn^{3+} ;

[00547] $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10}

5 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

10 [00548] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00549] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or
15 unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00550] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a
20 fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

[00551] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00552] Z is a counterion;

5 [00553] n is an integer from 0 to 3; and


[00554] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00555] Embodiment 105. A kit comprising:

[00556] at least one of an immune checkpoint inhibitor, T-cells for
10 an adoptive T-cell transfer therapy, and a cancer vaccine; and

[00557] a pentaaza macrocyclic ring complex according to formula (I),

[00558]



[00559]

15 [00560] wherein

[00561] M is Mn^{2+} or Mn^{3+} ;

[00562] R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting
20 of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

[00563]

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00564]

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00565]

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

[00566]

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00567]

Z is a counterion;

[00568]

n is an integer from 0 to 3; and

[00569]

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

Examples

[00570]

The following non-limiting examples are provided to further illustrate aspects of the present invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

[00571]

Pentaaza-macrocyclic ring complexes may protect cells including T-cells and other immunologically active cells, including CD8+, CD4+, natural killer (NK),

lymphokine-activated killer (LAK) and other cytotoxic or helper T lymphocytes from oxidative stressors including those within the tumor or tumor microenvironment. Here, we report evidence supporting that GC4419 (Galera Therapeutics, St. Louis, MO), a Mn(II) pentaaza-macrocyclic ring complex both alone and in combination with immune response
5 checkpoint inhibitors can increase the numbers of CD8+, and CD4+ (but not CD4+/CD25+/FoxP3+), T-cells. Such increases are believed to be beneficial in treating cancer, and in fact we also report here that GC4419 in combination with an immune response checkpoint inhibitor increases anti-tumor response versus treatment with the immune response checkpoint inhibitor as a single agent.

10 [00572] These results have significant implications with respect to combinations with immunotherapies other than immune checkpoint inhibitor treatments as well. This is because adoptive T-cell transfer therapies exogenously add T(effector)-cells, which are, or are similar to, CD8+ T-cells, and since, in addition, certain subsets of CD4+ (specifically excluding CD4+/CD25+/FoxP3) T-cells are believed to be important in
15 achieving good response with adoptive T-cell transfer therapies. Accordingly, as GC4419 increases CD4+ and/or CD8+ T-cell numbers, it is believed that GC4419 and other pentaaza macrocyclic ring complexes may also be beneficial in increasing the anti-tumor response to an adoptive T-cell transfer therapy.

[00573] Further, the results described herein are relevant to immunotherapies
20 such as treatments with cancer vaccines because the administration of a vaccine for treatment of cancer results in the generation of CD8+ and/or CD4+ T-cells. Accordingly, as GC4419 increases CD8+ and/or CD4+ T-cell numbers, it is believed that GC4419 and other macrocyclic ring complexes may also be beneficial in increasing the anti-tumor response to a therapeutic cancer vaccine.

25 [00574] Further, since therapeutic vaccines, T-cell transfer therapies and immune response checkpoint inhibitors may also be used to treat viral infections, both acute and chronic, by increasing CD8+ and/or CD4+ and/or similar T-cell numbers, and since GC4419 also increases CD8+ and/or CD4+ and/or similar T-cell numbers, it is believed that GC4419 and other pentaaza macrocyclic ring complexes may also be
30 beneficial in increasing the anti-viral response to therapeutic vaccines, T cell transfer therapies and immune response checkpoint inhibitors and be useful for the treatment viral disease in which increasing the immune system response is effective for treatment.

Example 1

[00575] GC4419 was administered in combination with the T-cell checkpoint inhibitor anti-PD-1 (RMP1-14) to female Balb/C mice implanted with the mouse colon cancer cell line, Colon 26 beginning on day 3 post-implantation. Tumors were allowed to grow for up to 52 days or until they exceeded 1000 mm³.

[00576] Treatments with the antibody and GC4419 are described in Table 1.

[00577] Table 1. Dosing Regimen for Colon 26 Syngeneic Tumor Model

		Treatment Regimen 1				Treatment Regimen 2			
Group	n	Agent	mg/kg	Route	Schedule	Agent	mg/kg	Route	Schedule
1	10	vehicle	-	ip	bid x 21 (start on day 3)	-	-	-	-
2	10	GC4419	10	ip	bid x 21 (start on day 3)	-	-	-	-
3	10	anti-PD1 RMP1-14	5	ip	biwk x 2 (start on day 3)	-	-	-	-
4	10	GC4419	1	ip	bid x 21 (start on day 3)	anti-PD1 RMP1-14	5	ip	biwk x 2 (start on day 3)
5	10	GC4419	3	ip	bid x 21 (start on day 3)	anti-PD1 RMP1-14	5	ip	biwk x 2 (start on day 3)
6	10	GC4419	10	ip	bid x 21 (start on day 3)	anti-PD1 RMP1-14	5	ip	biwk x 2 (start on day 3)

[00578] Tumor volumes were assessed and median and mean values are shown in Figures 1 (Median Tumor Volumes in Colon 26 Model) and 2 (Mean Tumor Volumes in Colon 26 Model).

[00579] Anti-PD1 monoclonal antibody treatment caused a modest decrease in tumor growth, and the addition of 1 and 3 mg/kg bid GC4419 caused a further decrease.

Example 2

[00580] GC4419 was administered in combination with the T-cell checkpoint inhibitor anti-PDL-1 (10F.9G2) to female Balb/C mice implanted with the mouse colon cancer cell line CT26 beginning on day 3 post-implantation. Tumors were allowed to grow for 17 days post-implantation and then collected.

[00581] Treatments with the antibody and GC4419 are described in Table 2.

[00582] Table 2. Dosing Regimen for CT26 Syngeneic Model

<u>Group</u>	<u>N =</u>	<u>Treatment</u>	<u>Dose</u> <u>(mg/kg/inj)</u>	<u>Schedule</u> <u>(start Day 3)</u>
<u>1</u>	<u>5</u>	<u>Control</u>	<u>-</u>	<u>Days 3,6,10,13</u>
<u>2</u>	<u>5</u>	<u>Anti-PD-L1</u> <u>(10F.9G2)</u>	<u>10</u>	<u>Days 3,6,10,13</u>
<u>3</u>	<u>5</u>	<u>GC4419</u>	<u>3</u>	<u>qd x 14</u>
<u>4</u>	<u>5</u>	<u>Anti-PD-L1</u> <u>GC4419</u>	<u>10</u> <u>10</u>	<u>Days 3,6,10,13</u> <u>qd x 14</u>
<u>5</u>	<u>5</u>	<u>Anti-PD-L1</u> <u>GC4419</u>	<u>10</u> <u>3</u>	<u>Days 3,6,10,13</u> <u>qd x 14</u>
<u>6</u>	<u>5</u>	<u>Anti-PD-L1</u> <u>GC4419</u>	<u>10</u> <u>1</u>	<u>Days 3,6,10,13</u> <u>qd x 14</u>

[00583] The mice were sacrificed on day 17 for analysis of the tumor for tumor infiltrating leukocytes and other immunologic cells by flow cytometry. The median tumor volumes are shown in Figure 3A (Median Tumor Volume Through Day 16 Post-Implantation).

[00584] To assess whether GC4419 was amplifying immune mediated tumor reduction, dissociated tumor cells were stained for markers of tumor infiltrating leukocytes and other immunologic cells, such as CD4+ (T helper class) and CD8+ (cytotoxic) T-cells,

myeloid derived suppressor Cells (MDSC) and Treg cells, with the results shown in Figure 3B (Intratumoral Leukocytes Assessed by Flow Cytometry).

[00585] When administered together, GC4419 and anti-PDL-1 antibody significantly increased CD4+ and CD8+ T-cells (but not CD4+/CD25+/FoxP3+ T(regulatory) cells) as compared to either GC4419 or anti-PDL-1 antibody alone, consistent with the hypothesis that GC4419, either increased the recruitment, survival or proliferation of T-cells produced as a result of checkpoint inhibition and involved in mounting an effective immune response to tumors.

Example 3

[00586] GC4419 enhances anti-tumor response in animals treated with ionizing radiation (IR). It is also shown herein that in immune competent animal models, GC4419 enhances the anti-tumor immune response to IR. The findings also show that the radiation therapy is enhanced by providing GC4419, even when radiation therapy is being used in combination with the immune checkpoint inhibitor anti-CTLA4. Other findings have indicated that GC4419 is suitable as a normal tissue radiation protector, and the above findings thus indicate the added advantage of enhancing radiation therapy.

[00587] Specifically, as shown in Figures 4A-4C, GC4419 decreases metastasis and enhances the efficacy of the combination of radiation and T-cell checkpoint inhibitor anti-CTLA4 (9D9) in the 4T1 metastatic breast cancer model.

Syngeneic animals were subcutaneously implanted with 4T1 cells to form a xenograft. On day 12, when no lung metastasis is present, animals were treated with GC4419 (24 mg/kg), 15 Gy of 250 kVp X-rays, and/or anti CTLA-4 antibodies (10 mg/kg) or IgG control antibodies, as described in Table 3 below. Tumor growth was tracked as a function of time and at day 35 post implantation (day 24 post start of treatment), animals were euthanized and lungs collected to count metastases.

[00588] Table 3. Dosing Regimen

Group	n	Treatment (s)	Dose	Schedule
1	10	Control	0	D12,13,14,15,16
2	10	GC4419	24 mg/kg	D12,13,14,15,16
3	10	RT	15 Gy	D12 x 1

4	10	GC4419	24 mg/kg	D12,13,14,15,16
		RT	15 Gy	D12 x 1
5	15	Anti-CTLA4 (9D9)	10 mg/kg	D10,13,16
		RT	15 Gy	D12 x 1
6	5	GC4419	24 mg/kg	D12,13,14,15,16
		Anti-CTLA4 (9D9)	10 mg/kg	D10,13,16
		RT	15 Gy	D12 x 1

[00589] GC4419 sensitized 4T1 tumors to radiation regardless of GC4419 dose (Figure 4A), enhanced the efficacy of combination anti CTLA-4 therapy and radiation therapy (Figure 4B), and significantly decreased the number of metastases 5 (Figure 4C).

[00590] Triple combination therapy of radiation, GC4419 and anti-CTLA-4, decreased the number of metastases per animal so much that it produced a significant number of animals without any lung metastases, as shown in Table 4 below.

[00591] Table 4. % Mice without lung metastases

10

Treatment	% Mice without Lung Metastases
Control	0
GC4419	0
RT	11
GC4419 + RT	22
aCTLA4 + RT	25
GC4419 + aCTLA4 + RT	80

[00592] Accordingly, the results shown in Figures 4A-4C and Table 4 demonstrate that GC4419 can be used favorably in triple combination therapies with

radiation treatment and cancer immune therapies such as immune checkpoint inhibitor treatment with anti-CTLA-4.

[00593] In separate studies, either syngeneic Balb/c mice or immunodeficient nu/nu mice were subcutaneously implanted with 4T1 cells to form a xenograft. On day 11 after transplant, animals were treated with GC4419 (24 mg/kg) and/or 15 Gy of 250 kVp X-rays, as described in Table 5 below.

[00594] Table 5. Dosing Regimen in nu/nu and Balb/c Mice

	Group	n	Treatment (s)	Dose	Schedule
Balb/c Mice	1	9	Control	0	D11,12,13,14,15
	2	6	GC4419	24 mg/kg	D11,12,13,14,15
	3	7	RT	15 Gy	D11 x 1
	4	6	GC4419	24 mg/kg	D11,12,13,14,15
			RT	15 Gy	D11 x 1
nu/nu Mice	1	6	Control	0	D11,12,13,14,15
	2	5	GC4419	24 mg/kg	D11,12,13,14,15
	3	5	RT	15 Gy	D11 x 1
	4	5	GC4419	24 mg/kg	D11,12,13,14,15
			RT	15 Gy	D11 x 1

10 [00595] Tumor volume was tracked as a function of time, and Day 17 (the last common day of measurement between the two studies given the rapid growth of tumors in the nu/nu mice) mean tumor volumes are described in Table 6 below. Additional measures such as Tumor Growth Delay and Tumor Doubling Time also showed similar trends.

15

[00596] Table 6. Mean Tumor Volume on Day 17

	Group	Treatment (s)	Day 17 Tumor Volume, V	V(x) / V(Group 3)
Balb/c Mice	1	Control	2263 mm ³	1.47
	2	GC4419	1923 mm ³	1.25
	3	RT	1535 mm ³	1.00
	4	GC4419 + RT	673 mm ³	0.47
nu/nu Mice	1	Control	All sac'd at Day 10	
	2	GC4419	All sac'd at Day 13	
	3	RT	1424 mm ³	1.00
	4	GC4419 + RT	902 mm ³	0.73

[00597] Both immunocompetent Balb/c mice and immunodeficient nu/nu mice
5 treated with radiation alone showed reduced tumor growth after treatment and further
tumor growth reduction was seen in both strains of mice treated with both radiation and
GC4419. Further, this increase in response to combined radiation and GC4419 treatment
was greater in immunocompetent mice than in immunodeficient mice, consistent with an
immunologic role for GC4419 being at least part of this increased response.

10 **Example 4**

[00598] The effects of GC4419 plus IR on intratumoral levels of key immune
cell populations were tested in Lewis lung carcinoma (LLC) tumors. In the study, 4 week
old C57.CL/6 mice were injected with LLC cells to form tumors. At 13 days post-injection,
animals were treated with GC4419 at doses of 3, 10 and 24 mg/kg and 15 Gy of 250 kVp
15 X-rays. Tumor growth was tracked until tumor size exceeded 2 cm in any one direction.
GC4419 sensitized tumors to ionizing radiation regardless of dose, as shown in Figure
6A. In addition, tumor infiltrating lymphocyte populations were assessed in separate
animals via flow cytometry at various time points post IR (15 Gy) in combination with 10
mg/kg GC4419 treatment. Intratumoral populations of neutrophils, macrophages and
20 activated cytotoxic T-cells were altered due to the presence of GC4419 either with or
without ionizing radiation, as shown in Figure 6B. However, it is unclear from this study
alone whether these transient alterations in the immune cell populations caused by

GC4419 contribute to improved results for the combination of GC4419 with checkpoint inhibitor therapy combined or in the absence of radiation therapy.

Example 5

[00599] GC4419 was administered in combination with the T-cell checkpoint inhibitor anti-CTLA-4 (9D9) to female Balb/C mice implanted subcutaneously with the mouse breast cancer cell line, 4T1. Tumors were allowed to grow for up to 45 days or until they exceeded 3000 mm³ (or Group mean of 2000 mm³).

[00600] Treatments with the antibody and GC4419 are described in Table 7.

[00601] Table 7. Dosing Regimen for 4T1 Syngeneic Tumor Model

10

Group	n	Treatment (s)	Dose (mg/kg)	Schedule
1	10	Vehicle (10 mM NaHCO ₃)	0	From D0;QDx21
2	10	GC4419	3	From D0;QDx21
3	10	Anti-CTLA4 (9D9)	10	BiW x 3 wk
4	10	Anti-CTLA4 (9D9)	10	BiW x 3 wk
		GC4419	3	From D0;QDx21
5	10	Anti-CTLA4 (9D9)	10	BiW x 3 wk
		GC4419	3	From D3;QDx21
6	10	Anti-CTLA4 (9D9)	10	BiW x 3 wk
		GC4419	3	From D6;QDx21

[00602] Tumor volumes were assessed and mean values are shown in Figure 5A (Mean Tumor Volumes in 4T1 Model).

[00603] Anti-CTLA4 monoclonal antibody treatment caused a significant decrease in tumor growth, and the addition of 3 mg/kg GC4419 started at least 3 days after antibody treatment caused a further decrease.

[00604] Notably, the results depicted for treatment with anti-CTLA4 alone are somewhat inconsistent with prior experience with this therapy, as the tumor growth decrease with anti-CTLA4 alone was somewhat higher than prior experience, and was

also higher than in other arms with anti-CTLA4 up until the point where GC4419 was added. In addition, the tumor growth decrease as compared to control when treating with anti-CTLA4 alone appeared to be improved even over a combination therapy with GC4419 where started on the same day as anti-CTLA4 treatment onset. This is despite the fact that combination of GC4419 with other checkpoint inhibitors, such as the anti-PD1 and anti-PDL1 therapies described above, demonstrate improved results when combined with GC4419 even for a same day start (or even the day prior). Accordingly, it is believed that the magnitude of this particular result for anti-CTLA4 treatment alone may be somewhat anomalous, and while not wishing to be limited by any theory, it is believed that treatment in combination with GC4419 provides good results over anti-CTLA4 alone, even for a same day start. Nonetheless, the results clearly demonstrate that the combination of GC4419 with anti-CTLA4 provided improved results over anti-CTLA4 alone, when a start of GC4419 treatment is delayed after the anti-CTLA4 treatment onset to a start on day 3 or day 6 after the first day of anti-CTLA4 treatment. That is, delaying the start of administration of GC4419, such as until day 3 or day 6, or even day 10 or day 13 after a start of anti-CTLA4 administration (such as, in this example until a day following the second, third, fourth or even fifth anti-CTLA4 dose), significantly improves treatment over anti-CTLA4 alone, as well as over a same-day start combination of anti-CTLA4 with GC4419.

[00605] Figure 5B depicts such an improvement occurring for dosing of GC4419 that is delayed until day 13 after a start of anti-CTLA4 administration (i.e., after the 5th dose of anti-CTLA4). In comparison of arms treated with (a) combination of GC4419 with anti-CTLA4 treatment and (b) anti-CTLA4 treatment alone, no difference should be apparent between the two arms before addition of GC4419 treatment. As a result tumor control was assessed by normalizing the tumor growth curves of each arm with respect to the day on which treatment with GC4419 was started in the combination arm (i.e., day 13 in Figure 5B). Such analysis shows the reduced tumor growth occurring after addition of GC4419 treatment as compared to anti-CTLA4 alone.

[00606] Accordingly, Figures 5A and 5B demonstrate the improved results in terms of tumor growth decrease that can be achieved with combinations of anti-CTLA4 and GC4419, including in dosing regimens where dosing with GC4419 is delayed for a period of time after dosing with anti-CTLA4 has begun, such as 3 to 6 days, and even

10 to 13 days after an anti-CTLA4 treatment onset (such as after the second, third or even fourth anti-CTLA4 dose).

Example 6

[00607] GC4419 was administered in combination with the T-cell checkpoint inhibitor anti-CTLA-4 (9D9) to female Balb/C mice implanted subcutaneously with the mouse breast cancer cell line, 4T1. Tumors were allowed to grow for up to 35 days or until they exceeded 3000 mm³ (or Group mean of 2000 mm³).

[00608] Treatments with the antibody and GC4419 are described in Table 8.

[00609] Table 8. Dosing Regimen for 4T1 Syngeneic Model

Group	N	Treatment	Dose (mg/kg)	Dosing days	Schedule
1	10	Vehicle(10mM NaHCO ₃)	0	day 4~ 21	QD
2	10	GC4419	10	day 4~ 21	QD
3	10	Anti-CTLA4(9D9)	10	day 1,4,8,11,15,18	BIW x 3 weeks
4	10	Anti-CTLA4(9D9)	10	day 1,4,8,11,15,18	BIW x 3 weeks
		GC4419	10	day 4~ 21	QD
5	10	Anti-CTLA4(9D9)	10	day 1,4,8,11,15,18	BIW x 3 weeks
		GC4419 or	3	day 5,6,7,9,10,12,13,14,16,17,19,20,21	QD
		Vehicle(10mM NaHCO ₃)	0	day 4,8,11,15,18	QD
6	10	Anti-CTLA4(9D9)	10	day 1,4,8,11,15,18	BIW x 3 weeks
		GC4419 or	10	day 5,6,7,9,10,12,13,14,16,17,19,20,21	QD
		Vehicle(10mM NaHCO ₃)	0	day 4,8,11,15,18	QD

10

[00610] Tumor volumes were assessed and mean values are shown in Figures 5C and 5D (Mean Tumor Volumes in 4T1 Model).

[00611] Figure 5C shows that the combination of anti-CTLA4 with GC4419 provided improved results in terms of decreased tumor volume both when administration of GC4419 started 3 days (day 9-26) and 4 days (day 10) following anti-

15

CTLA administration onset, over anti-CTLA4 treatment alone. Figure 5C further shows that skipping administration of GC4419 on those days when anti-CTLA4 was administered (day 10, skip) further improved the results, albeit slightly. Figure 5D further demonstrates that delaying administration of GC4419 until 4 days (day 10) after the first anti-CTLA4 administration provides improved results over anti-CTLA4 administration alone, including when administration of GC4419 is skipped on those days when anti-CTLA4 is administered. Also, the increased dose of 10 mg/kg of GC4419 provides improved results over a dose of 3 mg/kg, although significant improvements in treatment as compared to anti-CTLA4 alone are seen with both dose levels.

[00612] Accordingly, while anti-CTLA4 monoclonal antibody treatment caused a significant decrease in tumor growth, the addition of 3 mg/kg or 10 mg/kg GC4419, particularly when started at least 3 days after antibody treatment, caused a significant further decrease in tumor volumes, and skipping GC4419 administration on those days when anti-CTLA4 was administered further improved the results.

Example 7

[00613] In this example, the effects of treatment with GC4419 in combination with the immune checkpoint inhibitor anti-PD-1 (RMP1-14) were tested. 4T1 mouse breast cancer tumors were implanted subcutaneously in female mice. Dosing with controls, GC4419 and the anti-PD-1 antibodies was started on day 7, except where dosing with GC4419 was begun on day 6 or day 10, and continued until the time point as indicated in Table 9 below. Tumor volumes were measured approximately every 3 days through day 16.

[00614] Table 9.

Group	N	Treatment	Dose (mg/kg)	Planned Dosing Schedule/Days
Group-1	10	Vehicle(10mM NaHCO ₃)	10	QD x 3 weeks
Group-2	10	GC4419	10	QD x 3 weeks (start 1 day before first anti-PD-1; Day 6)
Group-3	10	Anti-PD-1 (RMP1-14)	10	BIW x 3 weeks (start day 7)
Group-4	10	GC4419	10	QD x 3 weeks (start 1 day before first anti-PD-1)
		Anti-PD-1 (RMP1-14)	10	BIW x 3 weeks
Group-5	10	GC4419	10	QD x 3 weeks (start 3 days post first anti-PD-1)
		Anti-PD-1 (RMP1-14)	10	BIW x 3 weeks

[00615] Figure 7 shows the result on mean tumor volume for the treatment regimens in Table 9 above. The Group 5 combination of anti-PD-1 antibody and GC4419 (started 3 days after first antibody injection) was the most effective in slowing 4T1 growth. However, the Group 4 combination of anti-PD-1 antibody and GC4419 started 1 day before the first anti-PD-1 treatment also provided good results. Accordingly, while not wishing to be limited by any theory, it is believed that treatment with the combination of GC4419 and anti-PD-1 provides good results over anti-PD-1 alone, even for a same day start, as good results are shown for both delaying administration of GC4419 after the anti-PD-1 start (such as 3 days as in Group 5), and for administration closer to the start of anti-PD-1 (e.g., on the day before administration as in Group 4), although the results further appear to show that delaying administration of GC4419 after the start of anti-PD-1 administration can provide improvements over administration closer to the start of anti-PD-1 administration (e.g., compare Group 5 and Group 4).

Example 8

[00616] Similar to the findings in Example 3, another experiment also showed that the radiation therapy is enhanced by providing GC4419, even when radiation therapy is being used in combination with the immune checkpoint inhibitor anti-CTLA4.

[00617] Specifically, as shown in Figures 8A-8E, GC4419 enhanced the efficacy of the combination of radiation and T-cell checkpoint inhibitor anti-CTLA4 (9D9) in the LLC squamous cell carcinoma breast cancer model. In addition to enhancing

efficacy against the irradiated primary tumor implanted in one flank of the animal, GC4419 also enhanced efficacy against an unirradiated second tumor implanted in the opposite flank.

[00618] Syngeneic animals (C57Bl/6 mice) were subcutaneously implanted with LLC cells in the right flank to form a xenograft (primary tumor). On day 2, the same animals were subcutaneously implanted with LLC cells in the left flank to form another xenograft (secondary tumor). On day 8, some animals began treatment with anti CTLA-4 antibodies (200 µg) as indicated below. On day 11, when both primary and secondary tumors were palpable, all animals were treated with GC4419 (24 mg/kg), 15 Gy of 250 kVp X-rays, and/or anti CTLA-4 antibodies (10 mg/kg), as described in Table 10 below. Tumor growth was tracked as a function of time for both primary and secondary tumors.

[00619] Table 10. Dosing Regimen

Group	n	Treatment (s)	Dose	Schedule
1	10	Control	0	D11,12,13,14,15
2	10	RT	15 Gy	D11 x 1
3	10	GC4419	10 mg/kg	D11,12,13,14,15
		RT	15 Gy	D11 x 1
4	10	Anti-CTLA4 (9D9)	200 µg	D8,11,13
		RT	15 Gy	D11 x 1
5	9	GC4419	10 mg/kg	D11,12,13,14,15
		Anti-CTLA4 (9D9)	200 µg	D8,11,13
		RT	15 Gy	D11 x 1

[00620] GC4419 sensitized primary and secondary tumors to radiation and primary tumors to the combination of anti CTLA-4 therapy and radiation therapy, delaying their growth (Figures 8A-8E). It also appeared to enhance the efficacy of the combination of anti CTLA-4 therapy and radiation therapy against secondary tumors, increasing the number of animals whose secondary tumors either stopped growing or disappeared at Day 73 post-implantation as described in Table 11 below.

[00621] Table 11. Secondary Tumor Responses at Day 73

Group	n	Treatment (s)	Complete Response + Stable Disease, n
1	10	Control	0
2	10	RT	0
3	10	GC4419 + RT	0
4	10	Anti-CTLA4 + RT	3
5	9	GC4419 + RT + Anti-CTLA4	6

5 [00622] The occasional ability of radiation treatment of one or more targeted tumors to produce an anti-tumor response in a distant unirradiated tumors is commonly known as the “abscopal” effect. This abscopal effect is widely believed due to a radiation-induced immune response against the unirradiated tumors. The combination of immunotherapy, such as anti CTLA4 therapy, and radiation therapy has been shown to increase the number of such abscopal effects. In the study described here the combination of anti CTLA-4 therapy and radiation therapy both slowed the growth of the unirradiated secondary tumors (Figure 8D) and generated apparent long-term responses (stable disease or better) in secondary tumors in some animals. Addition of GC4419 to the combination of anti CTLA-4 therapy and radiation therapy also slowed the growth of secondary tumors (Figure 8E) and resulted in a greater number of long-term responses (Table 11).

[00623] Accordingly, the results shown in Figures 8A-8E and Table 11 demonstrate that GC4419 can be used favorably in triple combination therapies with radiation treatment and cancer immune therapies such as immune checkpoint inhibitor treatment with anti-CTLA-4, to both increase efficacy in irradiated tumors and potentially generate efficacy in unirradiated tumors.

Example 9

[00624] Similar to the findings in Examples 3 and 8, another experiment showed that GC4419 enhances the combination of the immune checkpoint inhibitor anti-PD-L1 with radiation therapy.

[00625] Specifically, as shown in Figures 9 and 10A-10E and Table 12 below, GC4419 enhanced the efficacy of the combination of radiation and T-cell checkpoint inhibitor anti-PD-L1 (10F.9G2) in the LLC squamous cell carcinoma breast cancer model.

[00626] Syngeneic animals (C57Bl/6 mice) were subcutaneously implanted with LLC cells in the left flank to form a xenograft. On day 8, some animals began treatment with anti PD-L1 antibodies (200 µg) as indicated below. On day 11, all animals were treated with GC4419 (24 mg/kg), 15 Gy of 250 kVp X-rays, and/or anti PD-L1 antibodies (200 µg), as described in Table 12 below. Tumor growth was tracked as a function of time.

10 [00627] Table 12. Dosing Regimen

Group	n	Treatment (s)	Dose	Schedule
1	5	Control	0	D11,12,13,14,15
2	5	RT	15 Gy	D11 x 1
3	5	GC4419	10 mg/kg	D11,12,13,14,15
		RT	15 Gy	D11 x 1
4	5	GC4419	10 mg/kg	D11,12,13,14,15
		Anti-PD-L1 (10F.9G2)	200 µg	D8,11,14
5	9	Anti-PD-L1 (10F.9G2)	200 µg	D8,11,14
		RT	15 Gy	D11 x 1
6	9	GC4419	10 mg/kg	D11,12,13,14,15
		Anti-PD-L1 (10F.9G2)	200 µg	D8,11,14
		RT	15 Gy	D11 x 1

[00628] Through Day 31 post-implantation, the triple combination of GC4419 with anti PD-L1 therapy and radiation therapy, delayed tumor growth at least as well as the dual combination of anti PD-L1 therapy and radiation therapy (Figures 9 and 10A-10E). It also appeared to increase the number of animals whose tumors were “cured” (disappeared) as described in Figures 10A-10E and Table 13 below.

[00629] Table 13. % Animals Cured at Day 51


Group	n	Treatment (s)	Cures, n	% Cures
1	5	Control	0	0
2	5	RT	0	0
3	5	GC4419 + RT	0	0
4	5	GC4419 + Anti-PD-L1	0	0
5	9	Anti-PD-L1 + RT	5	56%
6	9	GC4419 +RT+ Anti-PD-L1	7	78%

WHAT IS CLAIMED IS:

1. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

- 5 administering to the subject an immune checkpoint inhibitor;
 administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after administration of the immune checkpoint inhibitor, to increase the response of the cancer to the immune checkpoint inhibitor:

10



wherein

M is Mn^{2+} or Mn^{3+} ;

- 15 R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

- 20 U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

5 W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of
10 the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and

15 the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.


2. The method according to claim 1, wherein R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each hydrogen.

20

3. The method according to claim 1 or 2, wherein W is an unsubstituted pyridine moiety.

4. The method according to any preceding claim, wherein U and V are
25 transcyclohexanyl fused rings.

5. The method according to any preceding claim, wherein the pentaaza macrocyclic ring complex is represented by formula (II):





wherein

5 X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

R_A, R_B, R_C, and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected

10 from the group consisting of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or alkyl.

15 6. The method according to any preceding claim, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):



wherein


X and Y represent suitable ligands which are derived from any monodentate or
 5 polydentate coordinating ligand or ligand system or the corresponding anion
 thereof; and

R_A, R_B, R_C, and R_D are independently hydrogen, hydrocarbyl, substituted
 hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected
 from the group consisting


10 of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO
 2NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂),

and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or
 alkyl.


15 7. The method according to any preceding claim, wherein the pentaaza
 macrocyclic ring complex is a compound represented by a formula selected from the
 group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)




(VIII)




(IX)




(X)




(XI)




(XII)




(XIII)



(XIV)



(XV)



(XVI)

5

8. The method according to any preceding claim, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate,

25

tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

5 each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

10 each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

each X_4 is independently hydrogen or together with X_3 is $(=O)$;

15 or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

or X and Y are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} .


20

9. The method according to any preceding claim, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

25 10. The method according to any one of claims 1-8, wherein X and Y are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates.


30 11. The method according to any one of claims 1-8, wherein X and Y are independently amino acids.

12. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




5

13. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




10

14. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




15


15. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



16. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



17. The method according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:

**GC4711**

18. The method according to any one of claims 1-8, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is administered a predetermined period of time after initial administration of the immune checkpoint inhibitor.

19. The method according to claim 18, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is no less than 3 days after initial administration of the immune checkpoint inhibitor.

20. The method according to claim 19, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is no less than 6 days after initial administration of the immune checkpoint inhibitor.

21. The method according to claim 19, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy is in a range of from 3 days to 9 weeks after initial administration of the immune checkpoint inhibitor.

22. The method according to any preceding claim, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows two doses of the immune checkpoint inhibitor.

23. The method according to claim 22, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows three doses of the immune checkpoint inhibitor.

24. The method according to claim 23, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows four doses of the immune checkpoint inhibitor.

25. The method according to claim 24, wherein initial administration of the pentaaza macrocyclic ring complex in the course of therapy follows five doses of the immune checkpoint inhibitor.

26. The method according to any preceding claim, wherein doses of the pentaaza macrocyclic ring complex provided in a course of cancer therapy are provided on separate days from any dose of the immune checkpoint inhibitor.

27. The method according to any preceding claim, comprising administering the immune checkpoint inhibitor and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

28. The method according to any preceding claim, wherein the checkpoint inhibitor interacts with one or more of cytotoxic T-lymphocyte antigen 4 (CTLA4), programmed death 1 (PD-1), programmed death ligand 1 (PDL-1), PDL-2, lymphocyte activation genes-3 (LAG3), B7 homolog 3 (B7-H3), B7 homolog 4 (B7-H4), indoleamine (2,3)-dioxygenase (IDO), adenosine A2a receptor (A2AR), neuritin, B- and T-lymphocyte attenuator (BTLA), killer immunoglobulin-like receptors (KIR), T cell immunoglobulin and mucin domain-containing protein 3 (TIME-3), inducible T cell costimulator (ICOS), CD27, CD28, CD40, CD137, CD160, CD244, HVEM, GAL9, VISTA, 2B4, CGEN-15049, CHK 1, CHK 2, GITR, CD47 and combinations thereof.

29. The method according to any preceding claim, wherein the checkpoint inhibitor comprises one or more of a small molecular inhibitor, an antibody, an antigen binding fragment, and an Ig fusion protein.

30. The method according to any preceding claim, wherein the checkpoint inhibitor is selected from the group consisting of ipilimumab, nivolumab, pembrolizumab, pidilizumab, areluman, tremelimumab, atezolizumab, AMP-224, MPDL3280A, MDX-1105, MDX-1106, MEDI-4736, IMP321, INCB024360, NLG-919, indoximod, AUNP 12, galiximab, avelumab, varlilumab, mogamulizumab, CP-870,893, MEDI-6469, IPH2101, urelumab, lirilumab, BMS-986016, MGA271, IMP321, BMS-936559, MSB0010718C, anti-OX40, MK-3475, CT-011, BY55, AMP224, and BGB-A317.

31. The method according to any preceding claim, wherein the checkpoint inhibitor is at least one of an anti-CTLA4 antibody, an anti-PD-1 antibody and an anti-PDL-1 antibody.

32. The method according to any preceding claim, further comprising administering one or more of adoptive T-cell transfer therapy and a cancer vaccine to the subject, either prior to, concomitantly with, or after administration of one or more of the checkpoint inhibitor and pentaaza macrocyclic ring complex.

33. The method according to any preceding claim, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.

34. The method according to any preceding claim, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 40 mg/kg.

35. The method according to claim 34, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 24 mg/kg.

36. The method according to claim 35, wherein the pentaaza macrocyclic ring complex is administered to the subject in a dose in a range of from 0.2 mg/kg to 10 mg/kg.

37. The method according to any preceding claim, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.

5

38. The method according to claim 37, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.


39. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

10

administering to the subject an adoptive T-cell transfer therapy;

administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after the adoptive T-cell transfer therapy, to increase the response of the cancer to the adoptive T-cell transfer therapy,

15



wherein

M is Mn^{2+} or Mn^{3+} ;

20

R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R$

12, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$,
wherein R_{11} and R_{12} are independently hydrogen or alkyl;

U, together with the adjacent carbon atoms of the macrocycle, forms a fused
substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or
5 heterocycle having 3 to 20 ring carbon atoms;

V, together with the adjacent carbon atoms of the macrocycle, forms a fused
substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or
heterocycle having 3 to 20 ring carbon atoms;

W, together with the nitrogen of the macrocycle and the carbon atoms of the
10 macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or
unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused
heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic
heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle
and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of
15 the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or
polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and


20 the dashed lines represent coordinating bonds between the nitrogen atoms of the
macrocycle and the transition metal, manganese.

40. The method according to claim 39, wherein R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 ,
 R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} are each hydrogen.
25

41. The method according to claim 39 or 40, wherein W is an unsubstituted
pyridine moiety.

42. The method according to any of claims 39-41, wherein U and V are
30 transcyclohexanyl fused rings.

43. The method according to any of claims 39-42, wherein the pentaaza macrocyclic ring complex is represented by formula (II):





wherein

X and Y represent suitable ligands which are derived from any monodentate or
 5 polydentate coordinating ligand or ligand system or the corresponding anion
 thereof; and

R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted
 hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected
 from the group consisting

10 of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$,
 and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or
 alkyl.

15 44. The method according to any of claims 39-43, wherein the pentaaza
 macrocyclic ring complex is represented by formula (III) or formula (IV):



wherein


X and Y represent suitable ligands which are derived from any monodentate or
 5 polydentate coordinating ligand or ligand system or the corresponding anion
 thereof; and

R_A, R_B, R_C, and R_D are independently hydrogen, hydrocarbyl, substituted
 hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected
 from the group consisting


10 of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO
 2NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂),

and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or
 alkyl.


15 45. The method according to any of claims 39-44, wherein the pentaaza
 macrocyclic ring complex is a compound represented by a formula selected from the
 group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)




(VIII)




(IX)




(X)




(XI)




(XII)




(XIII)



(XIV)



(XV)



(XVI)

5 46. The method according to any of claims 39-45, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, 10 nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, 15 aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, 20 thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate,

saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

each X_4 is independently hydrogen or together with X_3 is $(=O)$;

or X and Y are independently selected from the group consisting of

charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;


or X and Y are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} .

47. The method according to any of claims 39-46, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

48. The method according to any one of claims 39-46, wherein X and Y are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates.


49. The method according to any one of claims 39-46, wherein X and Y are independently amino acids.

50. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




5

51. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




10

52. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




15


53. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is represented by the formula:



54. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is represented by the formula:



55. The method according to any one of claims 39-46, wherein the pentaaza macrocyclic ring complex is represented by the formula:

**GC4711**

56. The method according to any of claims 39-55, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is a predetermined period of time after initial administration of the adoptive T-cell transfer therapy.

57. The method according to any of claims 39-56, comprising administering the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

58. The method according to any of claims 39-57, wherein the adoptive T-cell transfer therapy comprises administering to the subject cancer-specific autologous or allogeneic T-cells.

59. The method according to any of claims 39-58, wherein the adoptive T-cell transfer therapy comprises providing autologous tumor infiltrating lymphocytes, antigen-expanded CD8+ and/or CD4+ T cells, and genetically modified T cells that express T-cell receptors (TCR) that recognize tumor antigens.


60. The method according to any of claims 39-59, further comprising administering one or more of an immune checkpoint inhibitor and a cancer vaccine to the subject, either prior to, concomitantly with, or after administration of one or more of the adoptive T-cell transfer therapy and pentaaza macrocyclic ring complex.

61. The method according to any of claims 39-60, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.

62. The method according to any of claims 39-61, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.

63. The method according to claim 62, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.

64. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:
 administering to the subject a cancer vaccine;
 administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after administration of the cancer vaccine, to increase the response of the cancer to the cancer vaccine,



wherein

M is Mn^{2+} or Mn^{3+} ;

R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO₂NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂), and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or alkyl;

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and


the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

65. The method according to claim 64, wherein R₁, R₂, R'₂, R₃, R₄, R₅, R'₅, R₆, R'₆, R₇, R₈, R₉, R'₉, and R₁₀ are each hydrogen.

66. The method according to claim 64 or 65, wherein W is an unsubstituted pyridine moiety.

67. The method according to any of claims 64-66, wherein U and V are transcyclohexanyl fused rings.

- 5 68. The method according to any of claims 64-67, wherein the pentaaza macrocyclic ring complex is represented by formula (II):





wherein

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

R_A , R_B , R_C , and R_D are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl.

69. The method according to any of claims 64-68, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):



wherein


X and Y represent suitable ligands which are derived from any monodentate or
 5 polydentate coordinating ligand or ligand system or the corresponding anion
 thereof; and

R_A, R_B, R_C, and R_D are independently hydrogen, hydrocarbyl, substituted
 hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected
 from the group consisting


10 of -OR₁₁, -NR₁₁R₁₂, -COR₁₁, -CO₂R₁₁, -CONR₁₁R₁₂, -SR₁₁, -SOR₁₁, -SO₂R₁₁, -SO
 2NR₁₁R₁₂, -N(OR₁₁)(R₁₂), -P(O)(OR₁₁)(OR₁₂), -P(O)(OR₁₁)(R₁₂),

and -OP(O)(OR₁₁)(OR₁₂), wherein R₁₁ and R₁₂ are independently hydrogen or
 alkyl.


15 70. The method according to any of claims 64-69, wherein the pentaaza
 macrocyclic ring complex is a compound represented by a formula selected from the
 group consisting of formulae (V)-(XVI):




(V)




(VI)




(VII)




(VIII)




(IX)




(X)




(XI)




(XII)




(XIII)



(XIV)



(XV)



(XVI)

5

71. The method according to any of claims 64-70, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate,

tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

or X and Y correspond to $-O-C(O)-X_1$, where each X_1 is $-C(X_2)(X_3)(X_4)$, and

5 each X_1 is independently substituted or unsubstituted phenyl or $-C(-X_2)(-X_3)(-X_4)$;

each X_2 is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

10 each X_3 is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, $-X_5C(=O)R_{13}$ where X_5 is NH or O, and R_{13} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or $-OR_{14}$, where R_{14} is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with X_4 is $(=O)$; and

each X_4 is independently hydrogen or together with X_3 is $(=O)$;

15 or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

or X and Y are independently attached to one or more of R_1 , R_2 , R'_2 , R_3 , R_4 , R_5 , R'_5 , R_6 , R'_6 , R_7 , R_8 , R_9 , R'_9 , and R_{10} .


20

72. The method according to any of claims 64-71, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.

25 73. The method according to any one of claims 64-71, wherein X and Y are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates.


30 74. The method according to any one of claims 64-71, wherein X and Y are independently amino acids.

75. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




5

76. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




10

77. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




15


78. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is represented by the formula:



5 79. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is represented by the formula:



10 80. The method according to any one of claims 64-71, wherein the pentaaza macrocyclic ring complex is represented by the formula:

**GC4711**

81. The method according to any of claims 64-80, wherein initial administration of the pentaaza macrocyclic ring complex in a course of therapy is a predetermined period of time after initial administration of the cancer vaccine.

82. The method according to any of claims 64-81, comprising administering the cancer vaccine and pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy.

83. The method according to any of claims 64-82, wherein the cancer vaccine is selected from the group consisting of tumor cell vaccines, antigen vaccines, dendritic cell vaccines, DNA vaccines and vector based vaccines.

84. The method according to any of claims 64-83, wherein the cancer vaccine is selected from the group consisting of M-Vax (Avax Technologies), Provenge (Dendreon), GRNVAC1 (Geron), Bexidex (IDM Pharma), Uvidex (IDM Pharma), Collidex (IDM Pharma), INGN 225 (Introgen Therapeutics), M3Tk (MolMed), DC-Vax (Northwest Biotherapeutics), CVax (Prima Biomed), GVAX (Cell Genesys), Lucanix (NovaRx), Onyvax-P (Onyvax), HSPP-96 Oncophage (Antigenics), BiovaxID (Biovest International), NeuVax (Apthera), CDX-110 (CeppDex), GV1001 (Pharmexa), CYT004-MelQbG10 (Cytos Biotechnology), li-Key/HER2/neu (Generex Biotechnology), MAGE-A3 (Glaxo-SmithKline Biologicals), IDM-2101 (IDM Pharma), IMA901/IMA910 (Immatics Biotechnologies), melanoma cancer vaccine (Norwood Immunology), inCVAX (Immunophotonics) and Stimuvax (Oncothyreon).

85. The method according to any of claims 64-84, further comprising administering one or more of an immune checkpoint inhibitor and an adoptive T-cell transfer therapy to the subject, either prior to, concomitantly with, or after administration of one or more of the cancer vaccine and pentaaza macrocyclic ring complex.

86. The method according to any of claims 64-85, wherein the cancer is selected from the group consisting of breast cancer, non-small-cell lung cancer, melanoma, renal cell carcinoma, urothelial carcinoma, bladder cancer, pancreatic cancer, head and neck cancers, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, and oral squamous cell carcinoma.


87. The method according to any of claims 64-86, wherein the pentaaza macrocyclic ring complex is administered via at least one of parenteral route and oral route.

88. The method according to claim 87, wherein the pentaaza macrocyclic ring complex is administered intraperitoneally or intravenously.

89. A method of treating a viral infection in a mammalian subject in need thereof, comprising.

administering to the subject at least one of an immune checkpoint inhibitor, an adoptive T-cell transfer therapy, and a vaccine; and

administering to the subject a pentaaza macrocyclic ring complex corresponding to the formula (I) below, prior to, concomitantly with, or after the at least one immune checkpoint inhibitor, adoptive T-cell transfer therapy, and vaccine, to increase the effectiveness of the at least one immune checkpoint, adoptive T-cell transfer therapy, and vaccine in treating the viral infection,



wherein

M is Mn^{2+} or Mn^{3+} ;

$R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$, and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-OR_{11}$, $-NR_{11}R_{12}$, $-COR_{11}$, $-CO_2R_{11}$, $-CONR_{11}R_{12}$, $-SR_{11}$, $-SOR_{11}$, $-SO_2R_{11}$, $-SO_2NR_{11}R_{12}$, $-N(OR_{11})(R_{12})$, $-P(O)(OR_{11})(OR_{12})$, $-P(O)(OR_{11})(R_{12})$, and $-OP(O)(OR_{11})(OR_{12})$, wherein R_{11} and R_{12} are independently hydrogen or alkyl;

U , together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

V , together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

W , together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R_1 and R_{10} attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.


5

90. A kit comprising:

at least one of an immune checkpoint inhibitor, T-cells for an adoptive T-cell transfer therapy, and a cancer vaccine; and

a pentaaza macrocyclic ring complex according to formula (I),

10



wherein

M is Mn^{2+} or Mn^{3+} ;

15 $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9,$ and R_{10} are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of $-\text{OR}_{11}, -\text{NR}_{11}\text{R}_{12}, -\text{COR}_{11}, -\text{CO}_2\text{R}_{11}, -\text{CONR}_{11}\text{R}_{12}, -\text{SR}_{11}, -\text{SOR}_{11}, -\text{SO}_2\text{R}_{11}, -\text{SO}_2\text{NR}_{11}\text{R}_{12}, -\text{N}(\text{OR}_{11})(\text{R}_{12}), -\text{P}(\text{O})(\text{OR}_{11})(\text{OR}_{12}), -\text{P}(\text{O})(\text{OR}_{11})(\text{R}_{12}),$ and $-\text{OP}(\text{O})(\text{OR}_{11})(\text{OR}_{12}),$

20 wherein R_{11} and R_{12} are independently hydrogen or alkyl;

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

5 W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R₁ and R₁₀ attached to the carbon atoms which are both part of
10 the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and

15 the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.




FIG. 1




FIG. 2

2/16




FIG. 3A




FIG. 3B

3/16




FIG. 4A

4/16




FIG. 4B

5/16




FIG. 4C

6/16




FIG. 5A




FIG. 5B

7/16




FIG. 5C




FIG. 5D

8/16




FIG. 6A




FIG. 6B

10/16




FIG. 7

11/16




FIG. 8A




FIG. 8B

12/16




FIG. 8C




FIG. 8D

13/16




FIG. 8E




FIG. 9




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D

16/16




FIG. 10E

A. CLASSIFICATION OF SUBJECT MATTER**A61K 33/32(2006.01)i, A61K 39/00(2006.01)i, A61K 39/395(2006.01)i, A61P 35/00(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K 33/32; A61K 47/16; A61K 31/555; A61K 9/00; A61K 31/7068; A61K 31/675; C07D 259/00; C07D 257/00; C07F 13/00; A61K 39/00; A61K 39/395; A61P 35/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: cancer, immune checkpoint inhibitor, adoptive T cell transfer therapy, cancer vaccine, pentaaza macrocyclic ring complex

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2011-0136756 A1 (KEENE, JEFFERY L. et al.) 09 June 2011 See paragraphs [0143]-[0145]; claims 1-23.	90
X	US 2017-0044193 A1 (GALERA LABS, LLC) 16 February 2017 See paragraphs [0202], [0205] and [0224].	90
A	PARK, WONCHOUL et al., `Synthesis and SOD activity of manganese complexes of pentaaza macrocycles containing amino-and guanidino-auxiliary` Bulletin of the Korean Chemical Society, 2011, Vol.32, No.10, pages 3787-3789 See the whole document.	90
A	US 6214817 B1 (RILEY, DENNIS P. et al.) 10 April 2001 See the whole document.	90
A	US 5610293 A (RILEY, DENNIS P. et al.) 11 March 1997 See the whole document.	90



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

27 August 2018 (27.08.2018)

Date of mailing of the international search report

27 August 2018 (27.08.2018)

Name and mailing address of the ISA/KR

International Application Division

Korean Intellectual Property Office


189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

HEO, Joo Hyung

Telephone No. +82-42-481-8150



Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-89
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 1-89 pertain to methods for treatment of the human body by therapy, and thus relate to a subject matter which this International Searching Authority is not required, under PCT Article 17(2)(a)(i) and PCT Rule 39.1(iv), to search.
2. ☒ Claims Nos.: 19-21,23-25,35,36,38,63,88
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 19-21, 23-25, 35, 36, 38, 63 and 88 each refer to one of unsearchable claims which do not comply with PCT Rule 6.4(a).
3. ☒ Claims Nos.: 4-18,22,26-34,37,42-62,67-87
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2018/027588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2011-0136756 A1	09/06/2011	CA 2724550 A1	26/11/2009
		CA 2724550 C	03/01/2017
		EP 2296645 A2	23/03/2011
		EP 2296645 B1	19/11/2014
		EP 2859916 A2	15/04/2015
		EP 2859916 A3	29/04/2015
		EP 2859916 B1	29/11/2017
		US 9198893 B2	01/12/2015
		WO 2009-143454 A2	26/11/2009
		WO 2009-143454 A3	04/03/2010
US 2017-0044193 A1	16/02/2017	AU 2016-306568 A1	22/03/2018
		CA 2994845 A1	16/02/2017
		US 2017-0042907 A1	16/02/2017
		US 9738669 B2	22/08/2017
		US 9738670 B2	22/08/2017
		WO 2017-027728 A1	16/02/2017
US 6214817 B1	10/04/2001	AU 1998-80685 B2	17/05/2001
		AU 2000-77024 A1	17/04/2001
		AU 2000-77024 B2	02/02/2006
		AU 733415 B2	17/05/2001
		AU 7702400 A	17/04/2001
		AU 784078 B2	02/02/2006
		AU 8068598 A	04/01/1999
		CA 2294155 A1	30/12/1998
		CA 2382105 A1	22/03/2001
		CA 2382105 C	10/01/2012
		EP 1001752 A1	24/05/2000
		EP 1001752 B1	17/12/2003
		EP 1212323 A2	12/06/2002
		EP 1212323 B1	09/06/2004
		EP 1420019 A1	19/05/2004
		EP 1420019 B1	25/07/2007
		EP 1420022 A1	19/05/2004
		EP 1420022 B1	04/07/2007
		JP 2002-506445 A	26/02/2002
		JP 2003-509423 A	11/03/2003
		JP 2012-097097 A	24/05/2012
		US 2002-0128248 A1	12/09/2002
		US 2004-0147498 A1	29/07/2004
		US 2004-0219138 A1	04/11/2004
		US 2005-0171198 A1	04/08/2005
		US 2006-0270639 A1	30/11/2006
		US 2009-0131377 A1	21/05/2009
		US 6180620 B1	30/01/2001
		US 6395725 B1	28/05/2002
		WO 01-19823 A2	22/03/2001
		WO 01-19823 A3	07/09/2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2018/027588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5610293 A	11/03/1997	WO 2006-083508 A2	10/08/2006
		WO 2006-083508 A3	28/12/2006
		WO 98-58636 A1	30/12/1998
		CA 2072934 A1	20/01/1993
		CA 2072934 C	28/08/2007
		EP 0524161 A1	20/01/1993
		EP 0598753 A1	01/06/1994
		EP 0598753 B1	18/03/1998
		JP 06-509566 A	27/10/1994
		JP 3155552 B2	09/04/2001
		KR 10-0145953 B1	17/08/1998
		KR 10-0154346 B1	01/12/1998
		US 5637578 A	10/06/1997
		US 5874421 A	23/02/1999
		US 6084093 A	04/07/2000
		WO 93-02090 A1	04/02/1993