

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3514173 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 K 16/18 (2006.01)** **A 61 K 39/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2021-08-02**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2021-06-23**
- (86) Europæisk ansøgning nr.: **19156509.2**
- (86) Europæisk indleveringsdag: **2010-04-06**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-07-24**
- (30) Prioritet: **2009-04-10 US 16841109 P**
- (62) Stamansøgningsnr: **15159780.4**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **ELI LILLY AND COMPANY, Lilly Corporate Center, Indianapolis, IN 46285, USA**
- (72) Opfinder: **Chedid, Marcio, c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288, USA**
DARLING, Ryan James, c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288, USA
Galvin, Rachelle Jeanette, c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288, USA
Swanson, Barbara Anne, c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, IN 46206-6288, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **DKK-1-antistoffer**
- (56) Fremdragne publikationer:
WO-A1-2008/097510
WO-A1-2010/129752
WO-A2-2006/015373
WO-A2-2007/084344
FULCINITI MARIATERESA ET AL: "Anti-DKK1 mAb (BHQ880) as a potential therapeutic agent for multiple myeloma", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, UNITED STATES LNKD- DOI:10.1182/BLOOD-2008-11-191577, vol. 114, no. 2, 9 July 2009 (2009-07-09), pages 371-379, XP009134332, ISSN: 1528-0020 [retrieved on 2009-05-05]
GAVRIATOPOULOU MARIA ET AL: "Dickkopf-1: a suitable target for the management of myeloma bone disease.", EXPERT OPINION ON THERAPEUTIC TARGETS JUL 2009 LNKD- PUBMED:19530987, vol. 13, no. 7, July 2009 (2009-07), pages 839-848, XP009134320, ISSN: 1744-7631
YACCOBY SHMUEL ET AL: "Antibody-based inhibition of DKK1 suppresses tumor-induced bone resorption and multiple myeloma growth in vivo", BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US LNKD- DOI:10.1182/BLOOD-2006-09-047712, vol. 109, no. 5, 1 March 2007 (2007-03-01), pages 2106-2111, XP002486200, ISSN: 0006-4971 [retrieved on 2006-10-26]
ETTENBERG SETH A ET AL: "BHQ880, a novel anti-DKK1 neutralizing antibody, inhibits tumor-induced osteolytic bone disease", PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, NEW YORK, NY, US, vol. 49, 1 April 2008 (2008-04-01), page 947, XP009134334, ISSN:

Fortsættes ...

0197-016X

HEATH DEBORAH J ET AL: "Inhibiting Dickkopf-1 (Dkk1) removes suppression of bone formation and prevents the development of osteolytic bone disease in multiple myeloma", JOURNAL OF BONE AND MINERAL RESEARCH, AMERICAN SOCIETY FOR BONE AND MINERAL RESEARCH, NEW YORK, NY, US LNKD- DOI:10.1359/JBMR.081104, vol. 24, no. 3, 1 March 2009 (2009-03-01), pages 425-436, XP009134333, ISSN: 0884-0431 [retrieved on 2009-12-04]

FILIPPO CARACI ET AL: "The Wnt Antagonist, Dickkopf-1, as a Target for the Treatment of Neurodegenerative Disorders", NEUROCHEMICAL RESEARCH, KLUWER ACADEMIC PUBLISHERS-PLENUM PUBLISHERS, NE LNKD- DOI:10.1007/S11064-008-9710-0, vol. 33, no. 12, 22 April 2008 (2008-04-22), pages 2401-2406, XP019647587, ISSN: 1573-6903

HALL CHRISTOPHER L ET AL: "Dickkopf-1 expression increases early in prostate cancer development and decreases during progression from primary tumor to metastasis", PROSTATE, WILEY-LISS, NEW YORK, NY, US LNKD- DOI:10.1002/PROS.20805, vol. 68, no. 13, 15 September 2008 (2008-09-15), pages 1396-1404, XP009134318, ISSN: 0270-4137 [retrieved on 2008-06-16]

TERPOS E: "Antibodies to dickkopf-1 protein", EXPERT OPINION ON THERAPEUTIC PATENTS, INFORMA HEALTHCARE, GB LNKD- DOI:10.1517/13543776.16.10.1453, vol. 16, no. 10, 1 October 2006 (2006-10-01), pages 1453-1458, XP002486202, ISSN: 1354-3776

ZHOU XIAO-LEI ET AL: "Downregulation of Dickkopf-1 is responsible for high proliferation of breast cancer cells via losing control of Wnt/beta-catenin signaling", ACTA PHARMACOLOGICA SINICA, vol. 31, no. 2, February 2010 (2010-02), pages 202-210, XP009134321,

KRAJ MARIA: "Multiple myeloma bone disease and novel anti-myeloma agents", ACTA HAEMATOLOGICA POLONICA, vol. 39, no. 2, 2008, pages 163-178, XP009134337, ISSN: 0001-5814

DIARRA DANIELLE ET AL: "Dickkopf-1 is a master regulator of joint remodeling", NATURE MEDICINE, NATURE PUBLISHING GROUP, NEW YORK, NY, US LNKD- DOI:10.1038/NM1538, vol. 13, no. 2, 1 February 2007 (2007-02-01), pages 156-163, XP002486201, ISSN: 1078-8956 [retrieved on 2007-01-21]

MELODY MAZON ET AL: "Modulating Dickkopf-1: A Strategy to Monitor or Treat Cancer?", CANCERS, vol. 8, no. 7, 28 June 2016 (2016-06-28), page 62, XP55329263, DOI: 10.3390/cancers8070062

DESCRIPTION

[0001] The present invention relates to human engineered antibodies against DKK-1 and their use in treating diseases in which pathogenesis is mediated by DKK-1.

[0002] Dickkopf-1 (Dkk-1) is a member of the dickkopf family of proteins that have been shown to be negative regulators of the canonical Wnt-signaling pathway. The pathway plays a central role in bone development and formation. Dkk-1 inhibits Wnt signaling through its interaction with the Wnt co-receptor LRP5 or LRP6 and the kremen proteins. Dkk-1 prevents members of the Wnt pathway from interacting with either LRP5 or LRP6, thus preventing Wnt-mediated signal transduction. DKK1 has also been shown to be involved in bone metastasizing cancers, including multiple myeloma, breast carcinoma, renal carcinoma and non-small cell lung cancer.

[0003] WO2007/084344 describes antibodies that interfere with or neutralise the protein Dickkopf (Dkk-1) and/or Dickkopf (Dkk-4) mediated antagonism of Wnt signalling which are used to treat Dkk-related abnormalities of bone density, metabolism, diabetes and cancers.

[0004] The antibody BHQ880 (MOR-4910) described in Fulcintini Mariateresa et al: "Anti-DKK1 mAb (BHQ880) as a potential therapeutic agent for multiple myeloma" Blood, vol. 114, no. 2, 9 July 2009, pages 371-379, Gaviatopoulou Maria et al, "Dickkopf-1: a suitable target for the management of myeloma bone disease", Expert Opinion on Therapeutic Targets, vol. 13, no. 7, July 2009, pages 839-848 Yaccoby Shmeul et al: "Antibody-based inhibition of DKK1 suppresses tumor-induced bone resorption and multiple myeloma growth in vivo", Blood, American Society of Hematology, vol. 109, no. 5, 1 March 2007, pages 2106-211 and Ettenberg Seth et al: "BHQ880, a novel anti-DKK1 neutralizing antibody, inhibits tumor-induced osteolytic bone disease", Proceedings of the Annual Meeting of the American Association for Cancer Research, vol. 49, 1 April 2008, page 947 is reported to be an anti-DKK1/DKK4 antibody that neutralizes both DKK1 and DKK4.

[0005] Antibodies that bind Dkk-1 have been described (for example, see WO2006015373), however, there is still a need for therapeutic human engineered DKK-1 antibodies that will inhibit the interaction of Dkk-1 with LRP5 and LRP6. In addition, in view of the involvement of Dkk-1 in the regulation of bone formation, there is a need for therapeutic human engineered anti-Dkk-1 antibodies for use in bone healing. In addition, given the involvement of Dkk-1 in cancers, there is a need for human engineered anti-Dkk-1 antibodies for treating cancers, including multiple myeloma, breast and non-small cell lung cancers.

[0006] The antibodies of the present invention are therapeutically useful DKK-1 antagonists possessing a number of desirable properties. The present human engineered antibodies exhibit high affinity (Kd) to human DKK-1, cynomolgus DKK-1, rat DKK-1, mouse DKK-1, and rabbit DKK-1. Antibodies of the present invention block DKK-1-mediated inhibition of alkaline phosphatase, a marker of osteoblast activity. Furthermore, antibodies of the present invention exhibit an increase in bone mass density at both anterior and posterior cortices in a cortical

defect in vivo model and demonstrate significant growth inhibition of non-small cell lung xenografts in vivo.

[0007] According to a first aspect of the present invention, there is provided a DKK-1 antibody or antigen-binding fragment thereof, comprising a light chain variable region (LCVR) and a heavy chain variable region (HCVR), wherein the LCVR comprises the amino acid sequence of SEQ ID NO: 36 and the HCVR comprises the amino acid sequence of SEQ ID NO: 37.

[0008] Preferably, the DKK-1 antibody according to the first aspect of the present invention comprises a light chain comprising the amino acid sequence of SEQ ID NO: 34 and a heavy chain comprising the amino acid sequence of SEQ ID NO: 35.

[0009] More preferably, the DKK-1 antibody according to the first aspect of the present invention comprises two light chains wherein each light chain comprises the amino acid sequence of SEQ ID NO: 34, and two heavy chains wherein each heavy chain comprises the amino acid sequence of SEQ ID NO: 35.

[0010] According to a second aspect of the present invention, there is provided a pharmaceutical composition comprising the DKK-1 antibody or antigen-binding fragment thereof according to the present invention and a pharmaceutically acceptable carrier, diluent, or excipient.

[0011] The present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof that has a K_d at 37°C of less than 5.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32).

[0012] The present invention provides a human engineered DKK-1 antibody or a binding fragment thereof wherein LCDR1 has the amino sequence of SEQ ID NO:5, LCDR2 has the amino sequence of SEQ ID NO:47, LCDR3 has the amino sequence of SEQ ID NO:49, HCDR1 has the amino sequence of SEQ ID NO: 1, HCDR2 has the amino sequence of SEQ ID NO:2, and HCDR3 has the amino sequence of SEQ ID NO:44.

[0013] The present invention also provides a pharmaceutical composition comprising the human engineered DKK-1 antibody or antigen-binding fragment thereof of the present invention and a pharmaceutically acceptable carrier, diluent, or excipient.

[0014] The present invention provides a method of healing bone comprising administering a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention. The present invention also provides a method of treating cancer comprising administering a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention, wherein the cancer is selected from the group consisting of multiple myeloma, breast cancer and non-small cell lung cancer.

[0015] The present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof as described herein, wherein the antibody has a K_d of less than 1.5×10^{-11} M for human DKK-1 (SEQ ID NO: 29). More preferably, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d of less than 1.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29). Further preferred, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d of less than 5.0×10^{-12} M for human DKK-1 (SEQ ID NO: 29). More preferably, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d between 0.5×10^{-12} M and 1.5×10^{-11} M for human DKK-1 (SEQ ID NO: 29). Further preferred, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d between 1.0×10^{-12} M and 1.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29). The K_d values are established by a binding equilibrium at 37°C as described in Example 2.

[0016] The present invention also provides a human engineered DKK-1 antibody or antigen-binding fragment thereof as described herein, wherein the antibody has a K_d of less than 5.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). More preferably, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d of less than 3.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). Further preferred, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d of less than 2.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), rat Dkk-1 (SEQ ID NO: 31), and mouse DKK-1 (SEQ ID NO: 33). More preferably, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d between 1.0×10^{-11} M and 5.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). Further preferred, the present invention provides human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the antibody has a K_d between 1.5×10^{-11} M and 3.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), rat Dkk-1 (SEQ ID NO: 31), and mouse DKK-1 (SEQ ID NO: 33). The K_d values are established by a binding equilibrium at 37°C as described in Example 2.

[0017] More preferably, the present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12 and wherein the human engineered DKK-1 antibody or antigen-binding fragment thereof has a K_d of less than 3.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID

NO: 32). Further preferred, the present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12 and wherein the human engineered DKK-1 antibody or antigen-binding fragment thereof has a K_d of less than 2.5×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). More preferably, the present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12 and wherein the human engineered DKK-1 antibody or antigen-binding fragment thereof has a K_d of less than 2.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). Further preferred, the present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12 and wherein the human engineered DKK-1 antibody or antigen-binding fragment thereof has a K_d between 0.5×10^{-12} M and 3.0×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). More preferably, the present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12 and wherein the human engineered DKK-1 antibody or antigen-binding fragment thereof has a K_d between 1.0×10^{-12} M and 2.5×10^{-11} M for human DKK-1 (SEQ ID NO: 29), cynomolgus DKK-1 (SEQ ID NO: 30), rat Dkk-1 (SEQ ID NO: 31), mouse DKK-1 (SEQ ID NO: 33), and rabbit DKK-1 (SEQ ID NO: 32). The K_d values are established by a binding equilibrium at 37°C as described in Example 2.

[0018] The present invention provides a human engineered DKK-1 antibody or a binding fragment thereof, comprising a light chain variable region (LCVR) and a heavy chain variable region (HCVR), wherein the LCVR comprises complementarity determining regions (CDRs) LCDR1, LCDR2, and LCDR3 and the HCVR comprises CDRs HCDR1, HCDR2 and HCDR3, wherein LCDR1 has the amino sequence of SEQ ID NO:5, HCDR1 has the amino sequence of SEQ ID NO: 1, and HCDR2 has the amino sequence of SEQ ID NO:2.

[0019] The present invention provides a human engineered DKK-1 antibody or a binding fragment thereof wherein LCDR1 has the amino sequence of SEQ ID NO:46, LCDR2 has the amino sequence of SEQ ID NO:48, LCDR3 has the amino sequence of SEQ ID NO:50, HCDR1 has the amino sequence of SEQ ID NO: 1, HCDR2 has the amino sequence of SEQ ID NO:43, and HCDR3 has the amino sequence of SEQ ID NO:45. The present invention preferably provides a human engineered DKK-1 antibody or a binding fragment thereof wherein LCDR1 has the amino sequence of SEQ ID NO:5, LCDR2 has the amino sequence of SEQ ID NO:47, LCDR3 has the amino sequence of SEQ ID NO:49, HCDR1 has the amino sequence of SEQ ID NO: 1, HCDR2 has the amino sequence of SEQ ID NO:2, and HCDR3 has the amino sequence

of SEQ ID NO:44.

[0020] The present invention provides a human engineered DKK-1 antibody or a binding fragment thereof wherein LCDR1 has the amino sequence of SEQ ID NO:5, LCDR2 has an amino sequence selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, and SEQ ID NO:10, LCDR3 has an amino sequence selected from the group consisting of SEQ ID NO:7 and SEQ ID NO:9, HCDR1 has the amino sequence of SEQ ID NO:1, HCDR2 has the amino sequence of SEQ ID NO:2, and HCDR3 has an amino sequence selected from the group consisting of SEQ ID NO:3 and SEQ ID NO:4.

[0021] The present invention preferably provides a human engineered DKK-1 antibody or antigen-binding fragment thereof wherein LCDR1, LCDR2, LCDR3, HCDR1, HCDR2 and HCDR3 have amino acid sequences selected from the group consisting of:

1. (i) LCDR1 is SEQ ID NO: 5, LCDR2 is SEQ ID NO: 6, LCDR3 is SEQ ID NO: 7, HCDR1 is SEQ ID NO: 1, HCDR2 is SEQ ID NO: 2, and HCDR3 is SEQ ID NO: 3,
2. (ii) LCDR1 is SEQ ID NO: 5, LCDR2 is SEQ ID NO: 8, LCDR3 is SEQ ID NO: 7, HCDR1 is SEQ ID NO: 1, HCDR2 is SEQ ID NO: 2, and HCDR3 is SEQ ID NO: 4,
3. (iii) LCDR1 is SEQ ID NO: 5, LCDR2 is SEQ ID NO: 6, LCDR3 is SEQ ID NO: 9, HCDR1 is SEQ ID NO: 1, HCDR2 is SEQ ID NO: 2, and HCDR3 is SEQ ID NO: 3,
4. (iv) LCDR1 is SEQ ID NO: 5, LCDR2 is SEQ ID NO: 10, LCDR3 is SEQ ID NO: 9, HCDR1 is SEQ ID NO: 1, HCDR2 is SEQ ID NO: 2, and HCDR3 is SEQ ID NO: 3

[0022] The present invention provides a human engineered DKK-1 antibody or antigen-binding fragment thereof wherein the LCVR comprises the amino acid sequence of SEQ ID NO: 52 and the HCVR comprises the amino acid sequence of SEQ ID NO: 51.

[0023] The present invention preferably provides a human engineered DKK-1 antibody or antigen-binding fragment thereof wherein the LCVR comprises an amino acid sequence selected from the group consisting of SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16.

[0024] The present invention preferably provides a human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the HCVR comprises an amino acid sequence selected from the group consisting of SEQ ID NO:11 and SEQ ID NO:12.

[0025] The present invention preferably provides a human engineered DKK-1 antibody or antigen-binding fragment thereof, wherein the LCVR and HCVR comprise amino acid sequences selected from the group consisting of:

1. (i) a LCVR comprising the amino acid sequence of SEQ ID NO:13 and a HCVR comprising the amino acid sequence of SEQ ID NO:11;
2. (ii) a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR

- comprising the amino acid sequence of SEQ ID NO:12;
3. (iii) a LCVR comprising the amino acid sequence of SEQ ID NO:15 and a HCVR comprising the amino acid sequence of SEQ ID NO:11;
 4. (iv) a LCVR comprising the amino acid sequence of SEQ ID NO:16 and a HCVR comprising the amino acid sequence of SEQ ID NO:11.

[0026] The present invention preferably provides a human engineered DKK-1 antibody or antigen-binding fragment thereof comprising a LCVR comprising the amino acid sequence of SEQ ID NO:14 and a HCVR comprising the amino acid sequence of SEQ ID NO:12.

[0027] The present invention preferably provides a human engineered DKK-1 antibody wherein the human engineered DKK-1 antibody comprises a light chain wherein the light chain comprises an amino acid sequence selected from the group consisting of SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22.

[0028] The present invention preferably provides a human engineered DKK-1 antibody, wherein the human engineered DKK-1 antibody comprises a heavy chain wherein the heavy chain comprises an amino acid sequence selected from the group consisting of SEQ ID NO:17 and SEQ ID NO:18.

[0029] More preferably, the present invention provides a human engineered DKK-1 antibody, wherein the human engineered DKK-1 antibody comprises a heavy chain and a light chain amino acid sequence selected from the group consisting of

1. (i) a heavy chain comprising the amino acid sequence of SEQ ID NO:17 and light chain comprising the amino acid sequence of SEQ ID NO:19,
2. (ii) a heavy chain comprising the amino acid sequence of SEQ ID NO:18 and a light chain comprising the amino acid sequence of SEQ ID NO:20,
3. (iii) a heavy chain comprising the amino acid sequence of SEQ ID NO:17 and a light chain comprising the amino acid sequence of SEQ ID NO:21, and
4. (iv) a heavy chain comprising the amino acid sequence of SEQ ID NO:17 and a light chain comprising the amino acid sequence of SEQ ID NO: 22.

[0030] The present invention preferably provides a human engineered DKK-1 antibody comprising two light chains wherein each light chain comprises the amino acid sequence of SEQ ID NO: 19, and two heavy chains wherein each heavy chain comprises the amino acid sequence of SEQ ID NO:17.

[0031] The present invention preferably provides a human engineered DKK-1 antibody comprising two light chains wherein each light chain comprises the amino acid sequence of SEQ ID NO: 21, and two heavy chains wherein each heavy chain comprises the amino acid

sequence of SEQ ID NO:17.

[0032] The present invention preferably provides a human engineered DKK-1 antibody comprising two light chains wherein each light chain comprises the amino acid sequence of SEQ ID NO:22, and two heavy chains wherein each heavy chain comprises the amino acid sequence of SEQ ID NO:17.

[0033] The present invention preferably provides a human engineered DKK-1 antibody comprising two light chains wherein each light chain comprises the amino acid sequence of SEQ ID NO: 20, and two heavy chains wherein each heavy chain comprises the amino acid sequence of SEQ ID NO:18.

[0034] The present invention also provides an antibody or antigen-binding fragment thereof, which competes with the human engineered DKK-1 antibody or antigen-binding fragment thereof of the present invention for binding to human DKK-1 (SEQ ID NO: 29) as determined via an antibody competition assay.

[0035] The present invention also provides a pharmaceutical composition comprising the human engineered DKK-1 antibody or antigen-binding fragment thereof of the present invention and a pharmaceutically acceptable carrier, diluent, or excipient.

[0036] Furthermore, the present invention provides a pharmaceutical composition comprising the human engineered DKK-1 antibody or antigen-binding fragment thereof of the present invention together with a pharmaceutically acceptable carrier, diluent, or excipient and optionally other therapeutic ingredients.

[0037] In a further aspect, the present invention provides a method of healing bone comprising administering a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention.

[0038] In a further aspect, the present invention provides a method of treating cancer comprising administering a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention, wherein the cancer is preferably selected from the group consisting of multiple myeloma, breast cancer and non-small cell lung cancer.

[0039] Furthermore, the present invention provides a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention for use in therapy. Preferably, the present invention provides a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention for use in the treatment for healing bone or in the treatment of cancer, wherein the cancer is preferably selected from the group consisting of multiple myeloma, breast cancer and non-small cell lung cancer.

[0040] Furthermore, the present invention also provides for the use of a human engineered DKK-1 antibody or antigen binding fragment thereof of the present invention in the

manufacture of a medicament for therapy, healing bone or to treat cancer, wherein the cancer is preferably selected from the group consisting of multiple myeloma, breast cancer and non-small cell lung cancer.

Definitions

[0041] A full-length antibody as it exists naturally is an immunoglobulin molecule comprising 2 heavy (H) chains and 2 light (L) chains interconnected by disulfide bonds. The amino terminal portion of each chain includes a variable region of about 100-110 amino acids primarily responsible for antigen recognition via the complementarity determining regions (CDRs) contained therein. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function.

[0042] The CDRs are interspersed with regions that are more conserved, termed framework regions ("FR"). Each light chain variable region (LCVR) and heavy chain variable region (HCVR) is composed of 3 CDRs and 4 FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The 3 CDRs of the light chain are referred to as "LCDR1, LCDR2, and LCDR3" and the 3 CDRs of the heavy chain are referred to as "HCDR1, HCDR2, and HCDR3." The CDRs contain most of the residues which form specific interactions with the antigen. The numbering and positioning of CDR amino acid residues within the LCVR and HCVR regions is in accordance with the well-known Kabat numbering convention.

[0043] Light chains are classified as kappa or lambda, and are characterized by a particular constant region as known in the art. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the isotype of an antibody as IgG, IgM, IgA, IgD, or IgE, respectively. IgG antibodies can be further divided into subclasses, e.g., IgG1, IgG2, IgG3, IgG4. Each heavy chain type is characterized by a particular constant region with a sequence well known in the art.

[0044] As used herein, the term "monoclonal antibody" (Mab) refers to an antibody that is derived from a single copy or clone including, for example, any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Mabs of the present invention preferably exist in a homogeneous or substantially homogeneous population. Complete Mabs contain 2 heavy chains and 2 light chains. "Antigen-binding fragments" of such monoclonal antibodies include, for example, Fab fragments, Fab' fragments, F(ab')₂ fragments, and single chain Fv fragments. Monoclonal antibodies and antigen-binding fragments thereof of the present invention can be produced, for example, by recombinant technologies, phage display technologies, synthetic technologies, e.g., CDR-grafting, or combinations of such technologies, or other technologies known in the art. For example, mice can be immunized with human DKK-1 or fragments thereof, the resulting antibodies can be recovered and purified, and determination of whether they possess binding and functional properties similar to or the same as the antibody compounds disclosed herein can be assessed by the methods disclosed in

Examples below. Antigen-binding fragments can also be prepared by conventional methods. Methods for producing and purifying antibodies and antigen-binding fragments are well known in the art and can be found, for example, in Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, chapters 5-8 and 15, ISBN 0-87969-314-2.

[0045] The phrase "chimeric antibody" refers to an antibody containing domains from different species (generally 2 species). Antibody V is a chimeric antibody wherein the light chain and heavy chain variable domains contain residues from a murine antibody, whereas the constant region light chain domain contains residues comprising a rat kappa light chain and the constant region heavy chain domains contains residues comprising a rat IgG1 antibody. Antibody V is a murine/rat chimera and is used in studies to reduce the likelihood of immune response in long-term pre-clinical models.

[0046] The phrase "human engineered antibodies" refers to monoclonal antibodies that have binding and functional properties according to the invention, and that have framework regions that are substantially human or fully human surrounding CDRs derived from a non-human antibody. "Antigen-binding fragments" of such human engineered antibodies include, for example, Fab fragments, Fab' fragments, F(ab')₂ fragments, and single chain Fv fragments. "Framework region" or "framework sequence" refers to any one of framework regions 1 to 4. Human engineered antibodies and antigen-binding fragments thereof encompassed by the present invention include molecules wherein any one or more of framework regions 1 to 4 is substantially or fully human, i.e., wherein any of the possible combinations of individual substantially or fully human framework regions 1 to 4, is present. For example, this includes molecules in which framework region 1 and framework region 2, framework region 1 and framework region 3, framework region 1, 2, and 3, etc., are substantially or fully human. Substantially human frameworks are those that have at least about 80% sequence identity to a known human germline framework sequence. Preferably, the substantially human frameworks have at least about 85%, about 90%, about 95%, or about 99% sequence identity to a known human germline framework sequence.

[0047] Fully human frameworks are those that are identical to a known human germline framework sequence. Human framework germline sequences can be obtained from ImMunoGeneTics (IMGT) via their website <http://imgt.cines.fr>, or from *The Immunoglobulin FactsBook* by Marie-Paule Lefranc and Gerard Lefranc, Academic Press, 2001, ISBN 012441351. For example, germline light chain frameworks can be selected from the group consisting of: AI1, A17, A18, A19, A20, A27, A30, LI, L1I, L12, L2, L5, L15, L6, L8, O12, O2, and O8, and germline heavy chain framework regions can be selected from the group consisting of: VH2-5, VH2-26, VH2-70, VH3-20, VH3-72, VHI-46, VH3-9, VH3-66, VH3-74, VH4-31, VHI-18, VHI-69, VI-13-7, VH3-11, VH3-15, VH3-21, VH3-23, VH3-30, VH3-48, VH4-39, VH4-59, and VH5-51.

[0048] Human engineered antibodies in addition to those disclosed herein exhibiting similar functional properties according to the present invention can be generated using several

different methods. The specific antibody compounds disclosed herein can be used as templates or parent antibody compounds to prepare additional antibody compounds. In one approach, the parent antibody compound CDRs are grafted into a human framework that has a high sequence identity with the parent antibody compound framework. The sequence identity of the new framework will generally be at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% identical to the sequence of the corresponding framework in the parent antibody compound. This grafting may result in a reduction in binding affinity compared to that of the parent antibody. If this is the case, the framework can be back-mutated to the parent framework at certain positions based on specific criteria disclosed by Queen et al. (1991) Proc. Natl. Acad. Sci. USA 88:2869. Additional references describing methods useful in humanizing mouse antibodies include U.S. Patent Nos. 4,816,397; 5,225,539, and 5,693,761; computer programs ABMOD and ENCAD as described in Levitt (1983) J. Mol. Biol. 168:595-620; and the method of Winter and co-workers (Jones et al. (1986) Nature 321:522-525; Riechmann et al. (1988) Nature 332:323-327; and Verhoeyen et al. (1988) Science 239:1534-1536.

[0049] The identification of residues to consider for back-mutation can be carried out as follows:

When an amino acid falls under the following category, the framework amino acid of the human germ-line sequence that is being used (the "acceptor framework") is replaced by a framework amino acid from a framework of the parent antibody compound (the "donor framework"):

1. (a) the amino acid in the human framework region of the acceptor framework is unusual for human frameworks at that position, whereas the corresponding amino acid in the donor immunoglobulin is typical for human frameworks at that position;
2. (b) the position of the amino acid is immediately adjacent to one of the CDRs; or
3. (c) any side chain atom of a framework amino acid is within about 5-6 angstroms (center-to-center) of any atom of a CDR amino acid in a three dimensional immunoglobulin model.

[0050] When each of the amino acids in the human framework region of the acceptor framework and a corresponding amino acid in the donor framework is generally unusual for human frameworks at that position, such amino acid can be replaced by an amino acid typical for human frameworks at that position. This back-mutation criterion enables one to recover the activity of the parent antibody compound.

[0051] Another approach to generating human engineered antibodies exhibiting similar functional properties to the antibody compounds disclosed herein involves randomly mutating amino acids within the grafted CDRs without changing the framework, and screening the resultant molecules for binding affinity and other functional properties that are as good as or better than those of the parent antibody compounds. Single mutations can also be introduced at each amino acid position within each CDR, followed by assessing the effects of such mutations on binding affinity and other functional properties. Single mutations producing

improved properties can be combined to assess their effects in combination with one another.

[0052] Further, a combination of both of the foregoing approaches is possible. After CDR grafting, one can back-mutate specific framework regions in addition to introducing amino acid changes in the CDRs. This methodology is described in Wu et al. (1999) J. Mol. Biol. 294:151-162.

[0053] Applying the teachings of the present invention, a person skilled in the art can use common techniques, e.g., site-directed mutagenesis, to substitute amino acids within the presently disclosed CDR and framework sequences and thereby generate further variable region amino acid sequences derived from the present sequences. All alternative naturally occurring amino acids can be introduced at a specific substitution site. The methods disclosed herein can then be used to screen these additional variable region amino acid sequences to identify sequences having the indicated *in vivo* functions. In this way, further sequences suitable for preparing human engineered antibodies and antigen-binding portions thereof in accordance with the present invention can be identified. Preferably, amino acid substitution within the frameworks is restricted to one, two, or three positions within any one or more of the 4 light chain and/or heavy chain framework regions disclosed herein. Preferably, amino acid substitution within the CDRs is restricted to one, two, or three positions within any one or more of the 3 light chain and/or heavy chain CDRs. Combinations of the various changes within these framework regions and CDRs described above are also possible. Most preferably, these techniques are used to generate further variable region amino acid sequences using the variable heavy and light chain amino acid sequences describes in SEQ ID NO:12 and SEQ ID NO:14 respectively.

[0054] Human engineered antibodies or antigen-binding fragments thereof that "compete" with the molecules disclosed herein are those that bind human DKK-1 (SEQ ID NO:29) at site(s) that are identical to, or overlapping with, the site(s) at which the present molecules bind. Competing human engineered antibodies or antigen-binding fragments thereof can be identified, for example, via an antibody competition assay. For example, a sample of purified or partially purified human DKK-1 (SEQ ID NO:29) is bound to a solid support. An antibody disclosed herein and a test monoclonal antibody or antigen-binding fragment, with either the test or antibody of the present invention labeled, are then added. If the labeled antibody and the unlabeled antibody bind to separate and discrete sites on DKK-1, the labeled antibody will bind to the same level whether or not the suspected competing antibody is present. However, if the sites of interaction are identical or overlapping, the unlabeled antibody will compete, and the amount of labeled antibody bound to the antigen will be lowered. If the unlabeled antibody is present in excess, no labeled antibody will bind. For purposes of the present invention, competing human engineered antibodies or antigen-binding fragments thereof are those that decrease the binding of the present antibodies to DKK-1 by about 50%, about 60%, about 70%, about 80%, about 85%, about 90%, about 95%, or about 99%. Details of procedures for carrying out such competition assays are well known in the art and can be found, for example, in Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, pages 567-569, ISBN 0-87969-314-2. Such assays can

be made quantitative by using purified antibodies. A standard curve is established by titrating one antibody against a labeled version of itself. The capacity of an unlabeled competing monoclonal antibody or antigen-binding fragment thereof to inhibit the binding of the labeled molecule to the plate is titrated. The results are plotted, and the concentrations necessary to achieve the desired degree of binding inhibition are compared. Whether monoclonal antibodies or antigen-binding fragments thereof that compete with human engineered antibodies or antigen-binding fragments thereof of the present invention in such competition assays possess the same or similar functional properties of the present human engineered antibodies can be determined via the methods disclosed in Examples herein.

[0055] The term "inhibit" means the ability to substantially antagonize, prohibit, prevent, restrain, slow, disrupt, eliminate, stop, reduce, or reverse the biological effects of DKK-1.

[0056] The term "treating" (or "treat" or "treatment") refers to processes involving a slowing, interrupting, arresting, controlling, stopping, reducing, or reversing the progression or severity of a symptom, disorder, condition, or disease, but does not necessarily involve a total elimination of all disease-related symptoms, conditions, or disorders associated with DKK-1 activity.

[0057] The term "preventing" (or "prevent" or "prevention") means prohibiting, restraining, or inhibiting the incidence or occurrence of a symptom, disorder, condition, or disease. Acute events and chronic conditions may be treated and prevented. In an acute event, an antibody or antigen-binding fragment thereof is administered at the onset of a symptom, disorder, condition, or disease, and is discontinued when the acute event ends. In contrast, a chronic symptom, disorder, condition, or disease is treated over a more protracted time frame.

[0058] The term "effective amount" refers to the amount or dose of an antibody compound of the present invention which, upon single or multiple dose administration to a patient, provides the desired treatment or prevention. Therapeutically effective amounts of the present antibody compounds can comprise an amount in the range of from about 0.1 mg/kg to about 20 mg/kg per single dose. A therapeutically effective amount for any individual patient can be determined by the health care provider by monitoring the effect of the antibody compounds on a biomarker.

[0059] The term "bone healing" refers to the stimulation of bone formation at sites of injury by blocking DKK-1. Examples of bone healing indications include, but are not limited to, fracture healing, implant fixation/retention, and dental implant fixation/retention.

[0060] The human engineered antibodies of the present invention can be used as medicaments in human medicine, administered by a variety of routes. Most preferably, such compositions are for parenteral administration. Such pharmaceutical compositions can be prepared by methods well known in the art (See, e.g., Remington: The Science and Practice of Pharmacy, 19th ed. (1995), A. Gennaro et al., Mack Publishing Co.) and comprise a human engineered antibody as disclosed herein, and a pharmaceutically acceptable carrier, diluent, or

excipient.

[0061] The results of the following assays demonstrate that the monoclonal antibodies and antigen-binding fragments thereof, of the present invention are useful as a DKK-1 inhibitor. The antibodies of the present invention possess a number of desirable properties. For example, Antibody II of the present invention has increased chemical and physical stability, and solubility. Accelerated studies are performed to assess the chemical stability of antibodies by incubating the antibodies of the present invention under different buffer conditions (pH and NaCl variation) and incubating at 4°C, 25°C, and 40°C for 4 weeks. Chemical modifications of the antibodies of the present invention are detected by cation-exchange ("CEX") chromatography to separate charge variants (eg., deamidation of asparagine to aspartic acid) and LC-MS analysis to identify specific sites of degradation. Antibody II has the least amount of degradation detected by CEX and this result is further confirmed by LC-MS data showing that all three asparagine residues in the CDRs had the least amount of deamidation compared to the other antibodies described herein after 4-week incubation at 40°C in pH 8 buffer. Solubility for Antibody II is more favorable compared to Antibody I when formulated at pH 6 + 150 mM NaCl. Furthermore, Antibody II maintained solubility of >105 mg/mL when stored in 4°C, whereas solubility for Antibody I is only 48 mg/mL and precipitated under these same conditions. As used herein, "EC50" refers to the concentration of an agent which produces 50% of the maximal response possible for that agent. "Kd" refers to the equilibrium dissociation constant which can be calculated by the formula: $k_{\text{off}}/k_{\text{on}} = K_d$.

Example 1-Production of Antibodies

[0062] Antibodies I, II, III, and IV can be made and purified as follows. An appropriate host cell, such as HEK 293 EBNA or CHO, is either transiently or stably transfected with an expression system for secreting antibodies using an optimal predetermined HC:LC vector ratio or a single vector system encoding both HC, such as SEQ ID NO: 23, or SEQ ID NO: 24, and LC, such as SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, or SEQ ID NO: 28. Clarified media, into which the antibody has been secreted, is purified using any of many commonly-used techniques. For example, the medium may be conveniently applied to a Protein A or G Sepharose FF column that has been equilibrated with a compatible buffer, such as phosphate buffered saline (pH 7.4). The column is washed to remove nonspecific binding components. The bound antibody is eluted, for example, by pH gradient (such as 0.1 M sodium phosphate buffer pH 6.8 to 0.1 M sodium citrate buffer pH 3.0). Antibody fractions are detected, such as by SDS-PAGE, and then are pooled. Further purification is optional, depending on the intended use. The antibody may be concentrated and/or sterile filtered using common techniques. Soluble aggregate and multimers may be effectively removed by common techniques, including size exclusion, hydrophobic interaction, ion exchange, or hydroxyapatite chromatography. The purity of the antibody after these chromatography steps is greater than 99%. The product may be immediately frozen at -70°C or may be lyophilized. The amino acid sequences for these antibodies are provided below.

SEQ ID NOs

Antibody	Heavy Chain	Light Chain	HCVR	LCVR
I	17	19	11	13
II	18	20	12	14
III	17	21	11	15
IV	17	22	11	16
V	35	34	37	36

Antibody	HCDR1	HCDR2	HCDR3	LCDR1	LCDR2	LCDR3
I	1	2	3	5	6	7
II	1	2	4	5	8	7
III	1	2	3	5	6	9
IV	1	2	3	5	10	9
V	1	38	39	40	41	42

Example 2: Affinity (Kd) measurements for DKK-1 antibodies

[0063] To establish a binding equilibrium, a constant concentration of antibody is mixed with varying concentrations of His-tagged DKK-1 (SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, or SEQ ID NO:33) (1 nM - 1 pM range) in PBS (pH 7.4) + 1 mg/mL bovine serum albumin ("BSA") or buffer alone and incubated for several days at 37°C. Two sets of binding reactions are set up, one set at low concentration of antibody (3 pM) and one set at high concentration of antibody (30 or 50 pM).

[0064] After establishing equilibrium, a KinExA 3000 instrument (Sapidyne Inst. Inc.) is used to probe for the fraction of 'free' (unbound) antibody. Briefly, His-tagged DKK-1 is covalently coupled to NHS-activated Sepharose 4 Fast Flow beads (GE Healthcare) which are packed by the instrument to create a small column. The pre-equilibrated mixtures of antibody + His-tagged DKK-1 are passed over the beads to capture only free antibody. The amount of free antibody captured is proportional to the free concentration in the equilibrated samples, and is detected by injection of a fluorescently labeled secondary antibody. Data sets from low and high concentrations of antibody are fit globally using n-curve analysis in the KinExA software. This fitting returns a Kd value as well as the 95% confidence interval.

Table 1: Antibody II affinity to DKK-1 from various species

DKK-1 Species	Kd (pM)	95% Confidence Interval for Kd (pM)
Human	3.3	1.4 - 7.5
Cynomolgus	14	8.4 - 26
Rat	8.4	3.9 - 23
Murine	7.0	4.7 - 11
Rabbit	17	11 - 27

Example 3: Effect of Dkk-1 antibodies in C2C12 alkaline phosphatase assay

[0065] Canonical Wnt signaling is important for osteoblast differentiation and activity. Wnt-3a CM (Conditioned Media) combined with BMP-4 induces pluripotent mouse C2C12 cells to differentiate into osteoblasts with a measurable endpoint of alkaline phosphatase ("AP"), a marker of osteoblast activity. DKK-1, an inhibitor of canonical Wnt signaling, inhibits the differentiation and production of AP. Neutralizing DKK-1 antibodies prevent DKK-1-mediated inhibition of AP. Antibodies which block DKK-1 inhibitory activity prevent the loss of AP activity.

[0066] C2C12 cells are grown to 60% - 80% confluence in tissue culture flasks in growth medium (Dulbecco's Modified Eagle's Medium ("DMEM") containing L-glutamine, 10% heat-inactivated fetal bovine serum ("FBS"), 1x antibiotic/antimycotic, 1x sodium pyruvate). C2C12 cells are resuspended to a concentration of 30,000 cells/mL in growth medium and 100 μ L/well are added to a 96-well tissue culture plate and incubated overnight at 37°C, 95% humidity, 5% CO₂. Growth medium is replaced with 50 μ L assay medium (DMEM containing L-glutamine, 5% heat-inactivated FBS, 1x antibiotic/antimycotic, 1x sodium pyruvate). To stimulate differentiation (and thus induce AP production), 100 μ L assay medium plus 1.5x Wnt-3a CM + BMP-4 (R & D Systems catalogue #314-BP) are added. This yields a final concentration of 1x Wnt-3a CM and 25 ng/mL BMP-4. Negative controls contain only L-cell CM (no Wnt-3a or BMP-4). Cells are incubated at 37°C, 95% humidity, 5% CO₂ for 72 hours. Medium is removed, cells are washed with 200 μ L phosphate-buffered saline ("PBS"), and the PBS is removed. Cells are freeze-thawed 3 times. 100 μ L One-step pNPP substrate (Thermo Scientific catalogue #37621) is added per well, and the plate is incubated at room temperature. Absorbance is read at 405 nm. To determine the concentration of DKK-1 required to inhibit differentiation, Dkk-1 molecules from various species are titrated to identify the minimal concentration required to fully inhibit AP induction.

[0067] The minimal concentration of DKK-1 from each species which fully inhibits AP induction is as follows: human DKK-1=38 nM, cynomolgus DKK-1=11.4 nM, rat DKK-1=5.8 nM, and rabbit DKK-1 = 25.0 nM.

[0068] Having determined the concentration of DKK-1 which fully inhibits AP induction, increasing concentrations of an anti-DKK-1 antibody are pre-incubated in assay medium with

an inhibitory concentration of Dkk-1 for 30 minutes at room temperature. AP induction is determined as described above. Results are reported as an EC50 (nM $\pm$ standard error).

[0069] Antibody II blocks Dkk-1-mediated inhibition of C2C12 AP. For the DKK-1 of various species, the EC50's (given as nM $\pm$ standard error) are as follows: human = 9.8 ± 0.41 , cynomolgus = 6.4 ± 0.25 , rat = 2.9 ± 0.25 , and rabbit = 6.0 ± 0.32 .

Example 4: Cortical defect (CD) in vivo assay

[0070] Six month old female Sprague-Dawley rats are ovariectomized and allowed to lose bone for two months. 2 mm diameter defects are introduced into the left and right femurs with an electric drill with a 2 mm dental bit. This hole extends through both the anterior and posterior cortices. Healing of bone is monitored longitudinally by assessing bone mass density ("BMD") through the use of quantitative computed tomography ("qCT") for 35 days after surgery. At the end of the experiment, animals are sacrificed and whole femora are subjected to loaded-to-failure determinations to ascertain biomechanical strength of the whole diaphysis. Antibodies are administered subcutaneously at doses and intervals as indicated.

[0071] Antibody II is dosed as follows: 5 mg/kg once every two weeks starting one day after surgery (Group 1), 1 mg/kg (Group 2), 5 mg/kg (Group 3), or 15 mg/kg (Group 4) is administered once every two weeks starting nine days after surgery. BMD is assessed by qCT at day 35. Group 1 showed a statistically significant increase in BMD at both the anterior and posterior cortices. Group 4 showed a statistically significant increase in BMD at both cortices, although Groups 2 and 3 showed a non-significant increase in BMD.

Example 5: Cancer in vivo efficacy assay

[0072] Mice (female C.B-17 mice, Fox Chase severe combined immunodeficient model # CB 17SC-M) are acclimated for one week in an animal facility prior to experiment initiation. After acclimation, mice are randomized into groups of 10 per treatment. Cultured human A549 non-small cell lung carcinoma cells are implanted subcutaneously in the rear flank of the mice and tumor are allowed to reach a mean tumor volume ~ 100 mm³. Antibody II (1 mg/kg and 5 mg/kg), control IgG antibody (1 mg/kg and 5 mg/kg), or vehicle (citrate buffered saline supplemented with 0.02% Tween 80) is administered via subcutaneous injection. Animals receive 2 treatments separated by 7 days.

[0073] The tumors are measured 2 times per week by electronic calipers to plot growth curves. Animals are also monitored twice a week for fluctuations in body weight and signs of toxicity. Tumor volume measurements shown in table 3 are taken on day 29.

[0074] As shown in Table 2, treatment groups receiving Antibody II, demonstrated significant

growth inhibition of human A549 non-small cell lung xenografts in vivo.

Table 2: Efficacy of Antibody II in A549 human non-small cell lung carcinoma xenograft model

Treatment Group	Tumor Volume $\pm$ Standard Error (mm ³)	p-Value (relative to vehicle control group)
Citrate vehicle	402 $\pm$ 36	-
IgG control (1 mg/kg)	372 $\pm$ 45	-
IgG control (5 mg/kg)	492 $\pm$ 72	-
Antibody II (1 mg/kg)	279 $\pm$ 32	0.01 - 0.05
Antibody II (5 mg/kg)	202 $\pm$ 21	< 0.001

Example 6: Angiogenic re-normalization in vivo xenograft assay

[0075] To understand the mechanism of the anti-tumor efficacy, high content imaging analysis is carried out on A549 tumors treated with Antibody II or IgG control. Several phenotypic-based markers are analyzed both quantitatively and qualitatively to assess the cancer-relevant biological processes of angiogenesis (CD31 and smooth muscle actin, SMA), hypoxia induction (Glucose transporter 1, GLUT1), cell proliferation (Ki67), and apoptosis (Terminal UDP Nick-End Labeling, TUNEL).

[0076] Female C.B-17 (Fox Chase SCID) model # CB17SC-M mice age 7 to 8 weeks old are acclimated for one week and allowed to feed *ad libitum* on a normal low fat (4.5%) diet, which is continued for the duration of the study.

[0077] A549 cells from ATCC origin are grown and divided in F-12 Kaighn's media (Invitrogen #21127) supplemented with 10% FBS (Invitrogen #0, and 100x dilutions of sodium pyruvate, non-essential amino acids and pen-strep (Invitrogen #11360, #11140 and #15140 respectively). They are detached and prepared at a final concentration of 50×10^6 cells/ml in PBS at passage 19 with 95% viability.

[0078] 5×10^6 A549 human lung carcinoma cells are injected subcutaneously in the flank of subject mice in a 1:1 mixture of PBS and Matrigel (Becton Dickinson, Bedford, MA). Tumor and body weight measurements are performed twice weekly. Prior to treatment, mice are randomized based on tumor size using a randomization algorithm.

[0079] Starting when tumors reached 200 mm³, the randomized mice are separated into 2 groups of 10 animals and dosed subcutaneously on day 1 and again on day 8 with 5 mg/kg of either Antibody II or the IgG4 control. The study is terminated 10 days after administration of

the first dose of antibody.

[0080] Antibody II is prepared in Citrate Buffered Saline (CBS) (10 mM citrate pH6, 150 mM NaCl and 0.2% polysorbate). IgG4 control is provided at 6.0 mg/ml in Phosphate Buffered Saline (PBS).

[0081] Xenograft tumors are excised from mice after 17 days of dosing and placed into Zinc-Tris fixative (BD Pharmingen). Fixed tumors are processed, blocked in paraffin, and sectioned as 3 µm slices onto standard microscope slides. Slides are baked at 60°F for 1 hour and then deparaffinized in xylene (4 treatments, 10 minutes each). Slides are rehydrated through a series of ethanol/water immersions with final washes in Tris-buffered saline/Tween (TBST). Slides are then blocked with Protein Block (DAKO) for 30 minutes. For the Tumor Health Panel, slides are stained with a combination of Hoechst 33324 (Invitrogen), rat anti-human CD31 (Pharmingen)/anti-rat Alexa-488 (Invitrogen), rabbit anti-Ki67 (NeoMarkers)/anti-rabbit Alexa 647 (Invitrogen), and TUNEL-TMR red (diluted 1:5 in TUNEL dilution buffer; Roche). For the Angiogenesis Panel, slides are stained with a combination of Hoechst 33324 (Invitrogen), rat anti-human CD31 (Pharmingen)/anti-rat Alexa-488 (Invitrogen), rabbit anti-GLUT1 (Chemicon)/anti-rabbit Alexa 647 (Invitrogen), and mouse anti-Smooth Muscle Actin/Cy3 (Sigma). Slides are imaged using an iCys Laser Scanning Cytometer (CompuCyte) and a Marianas Digital Imaging Workstation configured with a Zeiss Axiovert 200M inverted fluorescence microscope (Intelligent Imaging Innovations). Quantitative data comparisons of treatment groups are performed using the Dunnet's analysis in JMP statistics software (SAS).

[0082] As shown in Table 3, A549 xenografts from control IgG-treated animals are modestly vascularized, have moderate levels of myofibroblasts and many focalized areas of hypoxia. Control IgG-treated tumors have an extended network of vessels consisting of both neoangiogenic vascular sprouts and mature vessels that are poorly covered by pericytes. Control IgG-treated tumors have hypoxic areas (marked by GLUT1) that are clearly demarcated zones some distance from perfused vessels and necrotic areas that are even further from vessels. Treatment with Antibody II results in decreased vessel area, decreased pericyte area, and decreased pericyte coverage of vessels. However, this treatment did not result in a significant difference in tumor hypoxia. Qualitatively, the Antibody II-treated tumor vasculature appears to consist of smaller, less-networked vessels and with less pericyte coverage compared with the control IgG-treated group.

[0083] As shown in Table 3, IgG-treated tumors have proliferating tumor cells evenly distributed throughout, but also frequently display one or more large areas of necrosis that are avascular. Treatment with Antibody II results in no change in apoptosis and only a slight, non-significant decrease in total proliferation area. However, there is a profound decrease in the percentage of area of high-intensity Ki67 cells in the Antibody II-treated tumors, which could indicate a decreased amount of cells in the G2 cell cycle phase.

Table 3: Antibody II in angiogenic re-normalization in vivo xenograft assay

Phenotypic Panel	Biological Parameter	Control IgG Value : Antibody II Treatment Value	Statistical Significance (Dunnet's p-value)
Tumor Angiogenesis Panel	% Total Vessel Area	10.26 : 7.88	0.001
Tumor Angiogenesis Panel	% Functional Vessel Area	9.84 : 7.59	0.0009
Tumor Angiogenesis Panel	% Non-Functional Vessel Area	0.42 : 0.30	0.1341
Tumor Angiogenesis Panel	% Pericyte Covered Vessels	42.44 : 30.88	0.0234
Tumor Angiogenesis Panel	% Tumor Hypoxic Area	14.90 : 11.77	0.3615
Tumor Proliferation Panel	% Tumor Proliferation Area	12.16 : 9.68	0.1739
Tumor Proliferation Panel	% High Ki67 Area	0.66 : 0.19	0.0012
Tumor Proliferation Panel	% Cycling Tumor Cells with High Ki67	5.24 : 1.68	<0.0001
Tumor Apoptosis Panel	% Tumor Apoptosis Area	8.63 : 8.24	0.7466
% Total Vessel Area - Percentage of tumor tissue area covered by all vessels regardless of their functional status.			
% Functional Vessel Area - Percentage of tumor tissue area covered by vessels in non-hypoxic regions.			
% Non-Functional Vessel Area - Percentage of tumor tissue area covered by vessels in hypoxic regions.			
% Pericyte Covered Vessels - Percentage of all tumor vessels that are associated with smooth muscle actin (SMA)-expressing cells.			
% Tumor Hypoxic Area - Percentage of tumor tissue area covered by cells expressing GLUT1, a surrogate marker for hypoxia.			
% Tumor Proliferation Area - Percentage of tumor tissue area that is positive for Ki67, a nuclear protein routinely used as a marker of actively proliferating cells.			
% High Ki67 Area - Percentage of tumor tissue area that stains intensely for Ki67. Cells expressing high levels of Ki67 are in later phases of cell cycle than those that express lower levels of Ki67.			
% Cycling Tumor cells with High Ki67 - Percentage of all Ki67 positive cells that are designated as having a high level of Ki67 expression. Cells expressing high levels of Ki67 are in later phases of cell cycle than those that express lower levels of Ki67.			
% Tumor Apoptosis Area - Percentage of tumor tissue area with high TUNEL			

Phenotypic Panel	Biological Parameter	Control IgG Value : Antibody II Treatment Value	Statistical Significance (Dunnett's p-value)
staining. TUNEL is a routinely used method to detect double-stranded DNA breakages which is indicative of terminal apoptosis.			

Example 7: Bone in vivo efficacy assay

[0084] To evaluate the ability of Antibody V (Light chain SEQ ID NO:34 and Heavy chain SEQ ID NO:35) accelerating the peri-implant bone formation and regenerating bone tissue, four months old male Sprague-Dawley rats (Harlan Sprague Dawley Inc) weighting around 425gram are utilized. Rats are surgically implanted with titanium screw (2 mm x 4mm) into the both legs of the media lateral side right above the tibial - fibular junction. Rats then are randomly divided into 2 groups according the body weight and receive either Antibody V 10mg/kg or IgG control 10mg/kg subcutaneously injection once weekly starting the same day of the surgery for 21 days. Quantitative computed tomography (Aloka LaTheta LTC-100 model CT scanner) is used to *ex vivo* scan and quantify the newly formed bone with a 60- μ m voxel size. Volume of interest (VOI) is defined as 28 slices at 0.1mm interval over the previous implant site. Group differences are assessed with JMP version 5.1. software (Cary, North Carolina), comparison against IgG-control using Dunnett's Method with a significance level of $P < 0.05$. Qualitatively evaluation by *ex vivo* faxitron x-ray images indicates that Antibody V treated rats have more new bone or callus around the implant. Quantitative CT analyses reveal a statistically significant increase cortical peri-implant bone mineral content (14%), new bone area (16%), cortical thickness (7%) and bone marrow cancellous peri-implant new bone mineral content (18%), and bone area (16%) in Antibody V treated rats when compared to IgG control. The data demonstrate that Antibody V stimulates new bone formation and accelerate regenerating bone tissue in titanium implanted rats.

Table 4: Micro-CT analyses of Peri-implant New Bone Formation

Group	n	Bone Marrow Peri-implant Analyses		Cortical Peri-implant Analyses		Cortical thickness (cm)
		Bone mineral Content (mg)	Bone Area (cm ²)	Bone mineral Content (mg)	Bone Area (cm ²)	
IgG control	10	0.84 $\pm$ 0.04	0.034 $\pm$ 0.001	6.70 $\pm$ 0.07	0.064 $\pm$ 0.001	0.054 $\pm$ 0.001
Antibody V	10	1.00 $\pm$ 0.04*	0.039 $\pm$ 0.002*	7.61 $\pm$ 0.06*	0.074 $\pm$ 0.001*	0.057 $\pm$ 0.001*

Data are presented as Mean $\pm$ SE. *, $p < 0.05$, vs IgG control, JMP Dunnett's test.

SEQ ID Listing**Heavy Chain CDRs****[0085]**

SEQ ID NO: 1	GFTFSSYTMS
SEQ ID NO:2	TISGGGFGTYYPDSVKG
SEQ ID NO:3	PGYHNYFFDI
SEQ ID NO:4	PGYNNYFFDI

Light Chain CDRs**[0086]**

SEQ ID NO:5	HASDSISNSLH
SEQ ID NO:6	YGRQSIQ
SEQ ID NO:7	QQSESWPLH
SEQ ID NO: 8	YARQSIQ
SEQ ID NO:9	QQSASWPLH
SEQ ID NO:10	YARQSEQ

Heavy Chain Variable Regions**[0087]**

SEQ ID NO:11

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTMSWVRQAPGKGLEWVATISGG
 GFGTYYPDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARPGYHNYFFDI
 WGQGTTTVTVSS

SEQ ID NO:12

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTMSWVRQAPGKGLEWVATISGG
 GFGTYYPDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARPGYNNYFFDI
 WGQGTTTVTVSS

Light Chain Variable Regions**[0088]**

SEQ ID NO:13

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYGRQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSESWPLHFGGGTKVEIK

SEQ ID NO:14

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYARQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSESWPLHFGGGTKVEIK

SEQ ID NO:15

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYGRQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSASWPLHFGGGTKVEIK

SEQ ID NO:16

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYARQSEQ
GIPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSASWPLHFGGGTKVEIK

Complete Heavy Chains**[0089]**

SEQ ID NO:17

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTMSWVRQAPGKGLEWVATISGG
GFGTYYPDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARPGYHNYFYDI
WGQGTITVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPTVSWNSG
ALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTKYTCNVDHKPSNTKVDKRVE
SKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQF
NWFYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGL
LPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYT
QKSLSLSLG

SEQ ID NO:18

EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTMSWVRQAPGKGLEWVATISGG
GFGTYYPDSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARPGYNYYFYDI
WGQGTITVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPTVSWNSG
ALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTKYTCNVDHKPSNTKVDKRVE
SKYGPPCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQF
NWFYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGL
LPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYT
QKSLSLSLG

Complete Light Chains

[0090]

SEQ ID NO:19

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYGRQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSESWPLHFGGGTKVEIKRTVAAPS
VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS
KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO:20

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYARQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSESWPLHFGGGTKVEIKRTVAAPS
VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS
KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO:21

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYGRQSIQG
IPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSASWPLHFGGGTKVEIKRTVAAP

SVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS
KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

SEQ ID NO:22

EIVLTQSPATLSLSPGERATLSCHASDSISNSLHWYQQKPGQAPRLLIYYARQSEQ
GIPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSASWPLHFGGGTKVEIKRTVAA
PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD
SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

Nucleotide Sequences - Heavy Chain Variable Regions**[0091]**

SEQ ID NO:23

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCC
TGAGACTCTCCTGTGCAGCCTCTGGATTCACTTTTCAGTAGCTATACCATGTCTT
GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCACCATTTCGGG
TGGTGGTTTCGGCACATACTATCCCGACAGTGTGAAGGGTCGATTACCATCT
CCAGAGACAACGCCAAGAACTCACTGTATCTGCAAATGAACAGCCTGAGAGC
CGAGGACACGGCTGTGTATTACTGTGCGAGACCTGGATATCACAATACTACT
TTGACATCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGCCTCCACCAA
GGGCCCATCGGTCTTCCCGCTAGCGCCCTGCTCCAGGAGCACCTCCGAGAGC
ACAGCCGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGG
TGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGCTGTC
CTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAG
CAGCTTGGGCACGAAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAAC
ACCAAGGTGGACAAGAGAGTTGAGTCCAAATATGGTCCCCCATGCCACCCT
GCCAGCACCTGAGGCCGCCGGGGGACCATCAGTCTTCCTGTTCCCCCAAA
ACCCAAGGACACTCTCATGATCTCCCGGACCCCTGAGGTACGTGCGTGGTG
GTGGACGTGAGCCAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGATG
GCGTGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTTCAACA
GCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAAC
GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCGTCTCCATCG
AGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAGCCACAGGTGTACA
CCCTGCCCCCATCCAGGAGGAGATGACCAAGAACCAGGTACGCTGACCTG
CCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAAAGCAAT

GGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACG
GCTCCTTCTTCTCTACAGCAGGCTAACCGTGGACAAGAGCAGGTGGCAGGA
GGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA
CACAGAAGAGCCTCTCCCTGTCTCTGGGT

SEQ ID NO:24

GAGGTGCAGCTGGTGGAGTCTGGGGGAGGCTTGGTCCAGCCTGGGGGGTCCC
TGAGACTCTCCTGTGCAGCCTCTGGATTCACTTTTCAGTAGCTATACCATGTCTT
GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGGCCACCATTTCGGG
TGGTGGTTTTCGGCACATACTATCCCGACAGTGTGAAGGGTCGATTACCATCT
CCAGAGACAACGCCAAGAACTCACTGTATCTGCAAATGAACAGCCTGAGAGC
CGAGGACACGGCTGTGTATTACTGTGCGAGACCTGGATATAATAACTACTACT

TTGACATCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGCCTCCACCAA
GGGCCCATCGGTCTTCCCGCTAGCGCCCTGCTCCAGGAGCACCTCCGAGAGC
ACAGCCGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGG
TGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGTGTC
CTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAG
CAGCTTGGGCACGAAGACCTACACCTGCAACGTAGATCACAAGCCCAGCAAC
ACCAAGGTGGACAAGAGAGTTGAGTCCAAATATGGTCCCCCATGCCACCCT
GCCAGCACCTGAGGCGCGGGGGGACCATCAGTCTTCTGTTCCCCCAAA
ACCCAAGGACACTCTCATGATCTCCCGGACCCCTGAGGTACGTGCGTGGTG
GTGGACGTGAGCCAGGAAGACCCCGAGGTCCAGTTCAACTGGTACGTGGATG
GCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTTCAACA
GCACGTACCGTGTGGTCAGCGTCCTACCGTCCTGCACCAGGACTGGCTGAAC
GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCGTCTTCCATCG
AGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAGCCACAGGTGTACA
CCCTGCCCCCATCCCAGGAGGAGATGACCAAGAACCAGGTACGCTGACCTG
CCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAAAGCAAT
GGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACG
GCTCCTTCTTCTCTACAGCAGGCTAACCGTGGACAAGAGCAGGTGGCAGGA
GGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA
CACAGAAGAGCCTCTCCCTGTCTCTGGGT

Nucleotide Sequences - Light Chain Variable Regions

[0092]

SEQ ID NO:25

GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAG
AGCCACCCTCTCCTGCCACGCCAGCGACAGTATTAGCAACAGCCTACACTGGT
ACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATTATGGCAGACA
GTCCATCCAGGGCATCCCAGCCAGGTTCAGTGGCAGTGGGTCTGGGACAGAC
TTCCTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTG
TCAACAGAGTGAGAGCTGGCCGCTCCACTTCGGCGGAGGGACCAAGGTGGAG
ATCAAACGAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGA
GCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATC
CCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAA
CTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
AGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTAC
GCCTGCGAAGTCAACCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCA
ACAGGGGAGAGTGC

SEQ ID NO:26

GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAG

GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAG
 AGCCACCCTCTCCTGCCACGCCAGCGACAGTATTAGCAACAGCCTACACTGGT
 ACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATTATGCTAGACA
 GTCCATCCAGGGCATCCCAGCCAGGTTCAAGTGGCAGTGGGTCTGGGACAGAC
 TTCCTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTG
 TCAACAGAGTGAGAGCTGGCCGCTCCACTTCGGCGGAGGGACCAAGGTGGAG
 ATCAAACGAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGA

GCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATC
 CCAGAGAGGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAA
 CTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
 AGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTAC
 GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCA
 ACAGGGGAGAGTGC

SEQ ID NO:27

GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAG
 AGCCACCCTCTCCTGCCACGCCAGCGACAGTATTAGCAACAGCCTACACTGGT
 ACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATTATGGCAGACA
 GTCCATCCAGGGCATCCCAGCCAGGTTCAAGTGGCAGTGGGTCTGGGACAGAC
 TTCCTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTG
 TCAACAGAGTGCCAGCTGGCCGCTCCACTTCGGCGGAGGGACCAAGGTGGAG
 ATCAAACGAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGA
 GCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATC
 CCAGAGAGGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAA
 CTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
 AGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTAC
 GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCA
 ACAGGGGAGAGTGC

SEQ ID NO:28

GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAG
 AGCCACCCTCTCCTGCCACGCCAGCGACAGTATTAGCAACAGCCTACACTGGT
 ACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATTATGCTAGACA
 GTCCGAGCAGGGCATCCCAGCCAGGTTCAAGTGGCAGTGGGTCTGGGACAGAC
 TTCCTCTCACCATCAGCAGCCTAGAGCCTGAAGATTTTGCAGTTTATTACTG
 TCAACAGAGTGCCAGCTGGCCGCTCCACTTCGGCGGAGGGACCAAGGTGGAG
 ATCAAACGAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGA
 GCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATC
 CCAGAGAGGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAA
 CTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTC
 AGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTAC
 GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCA
 ACAGGGGAGAGTGC

Human DKK-1

[0093]

SEQ ID NO:29

TLNSVLNSNAIKNLPPPLGGAAGHPGSAVSAAPGILYPGGNKEYQTIDNYQPYP
 CAEDECCTDEYCASPTRGGDAGVQICLACRKRRCMRHAMCCPGNYCKNGICV
 SSDQNHFRGEIETITESFGNDHSTLDGYSRRITLSSKMYHTKGQEGSVCLRSSD
 C ASGLCCARHFWSKICKPVLKEGQVCTKHRRKGSHGLEIFQRCYCGEGLS
 CRIQKDHHQASNSSRLHTCQRH

Cynomolgus DKK-1**[0094]**

SEQ ID NO:30

TLNSVLNSNAIKNLPPPLGGAAGHPGS AVSAAPGILYPGGNKYQTIDNYQPYP
CAEDEECGTDEY CASPTRGGDAGVQICLACRKRRCMRHAMCCPGNYCKNGICV
SSDQNNFRGEIEETITESFGNDHSTLDGYSRRTTLSSKMYHSGQEGSVCLRSSDC
ATGLCCARHFWSKICKPVLKEGQVCTKHRRKGSHGLEIFQRCYCGEGLSRIQKD
HHQASNSSRLHTCQR

Rat DKK-1**[0095]**

SEQ ID NO:31

TLNSVLINSNAIKNLPPPLGGAGGQPGSAVSVAPGVLYEGGNKYQTLDNYPYP
CAEDEECGTDEYCSSPSRGAAGVGGVQICLACRKRRCMRHAMCCPGNYCKNG
ICMPSDHSHLPRGEIEEGIIENLGNDHGAGDGYPRRTTLTSKIYHTKGQEGSVCLR
SSDCATGLCCARHFWSKICKPVLKEGQVCTKHRRKGSHGLEIFQRCYCGEGLAC
RIQKDHHQTSNSSRLHTCQRHAFIDYKDDDDKHV

Rabbit DKK-1**[0096]**

SEQ ID NO:32

TLNSVLVNSNAIKNLPPPLGGANGHPGS AVSATPGILYEGGNKYLPDNYQPYP
CATEDEECGTDEY CASPARGGGAGVQICLACRKRRCMRHAMCCPGNYCKNGIC
MPSDHNHFHRGEIEETIVESFGNDHSTSDGYSRRTTLSSKMYHAKGQEGSVCLRS
SDCATGLCCARHFWSKICKPVLKEGQVCTKHRRKGSHGLEIFQRCYCGDGLSCR
LQNDQHEASNSSRLHTCQR

Mouse DKK-1**[0097]**

SEQ ID NO:33

TLNSVLINSNAIKNLPPPLGGAGGQPGSAVSVAPGVLYEGGNKYQTLDNYPYP
CAEDEECGSDEYCSSPSRGAAGVGGVQICLACRKRRCMRHAMCCPGNYCKNG
ICMPSDHSHFPRGEIEESIIENLGNDHNAAAGDGYPRRTTLTSKIYHTKGQEGSV
CLRSSDCAAGLCCARHFWSKICKPVLKEGQVCTKHKRKGSHGLEIFQRCYCGEGL
ACRIQKDHHQASNSSRLHTCQRH

Antibody V - Complete Light Chain

[0098]

SEQ ID NO:34

DILLTQSPATLSVTPGDSVSLSCRASDSISGSLHWYQQKSHESPRLLIKYASQSISGI
PSRFSGSGSGTDFTLINSVETEDFGMYFCQQSNSWPLNFGAGTKLELKRADAAP

TVSIFPPSTEQLATGGASVVCLMNNFYPRDISVKWKIDGTERRDGVLDSVTDQDS
KDSTYSMSSTLSLSKADYESHNLYTCEVVHK TSSSPVVKSFNRNEC

Antibody V - Complete Heavy Chain

[0099]

SEQ ID NO:35

EVQLVESGGGLVKPGGSLKLSCAASGFTFSSYTMSWVRQTPEKRLEWVATISGG
GGNTYYPDSVKGRFTISRDNKNTLYLQLSSLRSEDALYYCARPGYNNYYFDY
WGQGTTLTVSSAKTTPPSVYPLAPGTALKSNSMVTLGCLVKGYFPEPVTVTWNS
GALSSGVHTFPAVLQSGLYTLTSSVTVPSSWPSQTVTCNV AHPASSTKVDKKIV
PRNCGGDCKPCICTGSEVSSVFIFPPKPKDVLITITLTPKVT CVVVDISQDDPEVHFS
WVVDVVEVHTAQTRPPEEQFNSTFRSVSELPILHQDWLNGRTFRCKVTSAAFPSPI
EKTISKPEGRTQVPHVYTMSPTKEEMTQNEVSITCMVKGFYPPDIYVEWQMNGQ
PQENYKNTPTMDTDGSYFLYSKLNVKKEKWQQGNTFTCSVLHEGLHNHHTK
SLSHSPGK

Antibody V - Light Chain Variable Region

[0100]

SEQ ID NO:36

DILLTQSPATLSVTPGDSVSLSCRASDSISGSLHWYQQKSHESPRLLIKYASQSISGI
PSRFSGSGSGTDFTLINSVETEDFGMYFCQQSNSWPLNFGAGTKLELK

Antibody V - Heavy Chain Variable Region

[0101]

SEQ ID NO:37

EVQLVESGGGLVKPGGSLKLSCAASGFTFSSYTMSWVRQTPEKRLEWVATISGG
GGNTYYPDSVKGRFTISRDNKNTLYLQLSSLRSEDALYYCARPGYNNYYFDY
WGQGTTLTVSS

Antibody V - Heavy Chain CDRs

[0102]

SEQ ID NO:38	TISGGGGNTYYPDSVKG
SEQ ID NO:39	PGYNNYYFDY

Antibody V - Light Chain CDRs

[0103]

SEQ ID NO:40	RASDSISGSLH
SEQ ID NO:41	YASQSIG
SEQ ID NO:42	QQSNSWPLN

Consensus Heavy Chain CDRs

[0104]

SEQ ID NO:43	TISGGGX ₁ X ₂ TYYPDSVKG
X ₁ is F, G	
X ₂ is G, N	
SEQ ID NO:44	PGYX ₃ NYFFDI
X ₃ is H, N	
SEQ ID NO:45	PGYX ₄ NYFFDX ₅
X ₄ is H, N	
X ₅ is I, Y	

Consensus Light Chain CDRs

[0105]

--	--

SEQ ID NO:46	X_6 ASDSIS X_7 SLH
X_6 is H, R	
X_7 is N, G	
SEQ ID NO:47	Y X_8 RQS X_9 Q
X_8 is G, A	
X_9 is I, E	
SEQ ID NO:48	Y X_{10} X_{11} QS X_{12} X_{13}
X_{10} is G, A	
X_{11} is R, S	
X_{12} is I, E	
X_{13} is Q, S	
SEQ ID NO:49	QQS X_{14} SWPLH
X_{14} is E, A	
SEQ ID NO:50	QQS X_{15} SWPL X_{16}
X_{15} is E, A, N	
X_{16} is H, N	

Consensus Heavy Chain Variable Region

[0106]

SEQ ID NO:51
 EVQLVESGGGLVQPGGSLRLSCAASGFTFSSYTMSWVRQAPGKGLEWVATISGG
 GFGTYYPDSVKGRFTISRDNKNSLYLQMNSLRAEDTAVYYCARPGY X_{17} NYFF
 DIWGQGTTVTVSS
 X_{17} is H, N

Consensus Light Chain Variable Region

[0107]

SEQ ID NO:52
 DIVLTQSRATLSLSPGERATLSCHASDSISNSLHWYQOKDGOADPLLVVY X_{18} POS

EIVLTQSFATLSLSFQERATLSCHASDSISNSLHWIQQKFGQAFKLLIITAI8KQ3
 X₁₉QGIPARFSGSGSGTDFTLTISSLEPEDFAVYYCQQSX₂₀SWPLHFGGGTKVEIK
 X₁₈ is G, A

X₁₉ is I, E

X₂₀ is E, A

[0108] The present invention is defined by the claims.

SEQUENCE LISTING

[0109]

<110> Eli Lilly and Company

<120> DKK-1 Antibodies

<130> X18097B

<150> 61/168411

<151> 2009-04-10

<150> PCT/US2010/030039

<151> 2010-04-06

<150> 10712681.5

<151> 2010-04-06

<160> 52

<170> PatentIn version 3.5

<210> 1

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 1

Gly	Phe	Thr	Phe	Ser	Ser	Tyr	Thr	Met	Ser
1				5					10

<210> 2

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 2

Thr	Ile	Ser	Gly	Gly	Gly	Phe	Gly	Thr	Tyr	Tyr	Pro	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 3

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 3

Pro Gly Tyr His Asn Tyr Tyr Phe Asp Ile

1				5					10
---	--	--	--	---	--	--	--	--	----

<210> 4

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 4

Pro	Gly	Tyr	Asn	Asn	Tyr	Tyr	Phe	Asp	Ile
1				5					10

<210> 5

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 5

His	Ala	Ser	Asp	Ser	Ile	Ser	Asn	Ser	Leu	His
1				5					10	

<210> 6

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 6

Tyr	Gly	Arg	Gln	Ser	Ile	Gln
1				5		

<210> 7

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 7

Gln	Gln	Ser	Glu	Ser	Trp	Pro	Leu	His
1				5				

<210> 8

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 8

Tyr	Ala	Arg	Gln	Ser	Ile	Gln
1				5		

<210> 9

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 9

Gln	Gln	Ser	Ala	Ser	Trp	Pro	Leu	His
1				5				

<210> 10

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 10

Tyr Ala Arg Gln Ser Glu Gln
1 5

<210> 11

<211> 119

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 11

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Thr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Thr Ile Ser Gly Gly Gly Phe Gly Thr Tyr Tyr Pro Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Pro Gly Tyr His Asn Tyr Tyr Phe Asp Ile Trp Gly Gln Gly
100 105 110

Thr Thr Val Thr Val Ser Ser
115

<210> 12

<211> 119

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 12

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Thr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Thr Ile Ser Gly Gly Gly Phe Gly Thr Tyr Tyr Pro Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Pro Gly Tyr Asn Asn Tyr Tyr Phe Asp Ile Trp Gly Gln Gly
100 105 110

Thr Thr Val Thr Val Ser Ser
115

<210> 13

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 13

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Tyr Gly Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Glu Ser Trp Pro Leu
85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 14

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 14

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
 20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Tyr Ala Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Glu Ser Trp Pro Leu
 85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 15

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 15

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
 20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Tyr Gly Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ala Ser Trp Pro Leu
 85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 16

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 16

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
 20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Tyr Ala Arg Gln Ser Glu Gln Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ala Ser Trp Pro Leu
 85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 17

<211> 445

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 17

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Thr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Thr Ile Ser Gly Gly Gly Phe Gly Thr Tyr Tyr Pro Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Pro Gly Tyr His Asn Tyr Tyr Phe Asp Ile Trp Gly Gln Gly
 100 105 110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125

Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
 130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190

Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro
 195 200 205

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro
 210 215 220

Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe
 225 230 235 240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
 245 250 255

Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val
 260 265 270

Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 275 280 285

Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val
 290 295 300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 305 310 315 320

Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser
 325 330 335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
 340 345 350

Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
 355 360 365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 370 375 380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
 385 390 395 400

Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp
 405 410 415

Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
 420 425 430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly
 435 440 445

<210> 18

<211> 445

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 18

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Thr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Thr Ile Ser Gly Gly Gly Phe Gly Thr Tyr Tyr Pro Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Pro Gly Tyr Asn Asn Tyr Tyr Phe Asp Ile Trp Gly Gln Gly
 100 105 110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125

Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu
 130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190

Ser Ser Leu Glv Thr Lvs Thr Tvr Thr Cvs Asn Val Asp His Lvs Pro

```

-----
      195              200              205
-----
Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro
 210              215              220

Cys Pro Pro Cys Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe
 225              230              235              240

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
              245              250              255

Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val
              260              265              270

Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 275              280              285

Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val
 290              295              300

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 305              310              315              320

Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser
              325              330              335

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
              340              345              350

Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
 355              360              365

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 370              375              380

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
 385              390              395              400

Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp
              405              410              415

Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
 420              425              430

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly
 435              440              445

```

<210> 19

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 19

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
 20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
 35 40 45

Tyr Tyr Gly Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Glu Ser Trp Pro Leu
 85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> 20

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 20

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Tyr Ala Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Glu Ser Trp Pro Leu
85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195 200 205

Phe Asn Arg Gly Glu Cys
210

<210> 21

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 21

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile

```

-----
      35              40              45
-----
Tyr Tyr Gly Arg Gln Ser Ile Gln Gly Ile Pro Ala Arg Phe Ser Gly
 50              55              60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65              70              75              80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ala Ser Trp Pro Leu
      85              90              95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
      100              105              110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
      115              120              125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
      130              135              140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145              150              155              160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
      165              170              175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
      180              185              190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
      195              200              205

Phe Asn Arg Gly Glu Cys
      210

```

<210> 22

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 22

```

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1              5              10              15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
      20              25              30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
      35              40              45

```

Tyr Tyr Ala Arg Gln Ser Glu Gln Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ala Ser Trp Pro Leu
85 90 95

His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195 200 205

Phe Asn Arg Gly Glu Cys
210

<210> 23

<211> 1335

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 23

gaggtgcagc tgggtggagtc tgggggaggc ttggtccagc ctgggggggtc cctgagactc	60
tctctgtgcag cctctggatt cactttcagt agctatacca tgtcttgggt ccgccaggct	120
ccagggaagg ggctggagtg ggtggccacc atttccggtg gtggtttcgg cacatactat	180
cccgacagtg tgaagggtcg attcaccatc tccagagaca acgccaagaa ctcaactgtat	240
ctgcaaatga acagcctgag agccgaggac acggctgtgt attactgtgc gagacctgga	300
tatcacaact actactttga catctggggc caagggaacca cggtcaccgt ctctcagcc	360
tccaccaagg gcccatcggt cttcccgcta gcgccctgct ccaggagcac ctccgagagc	420
acagccgccc tgggctgcct ggtcaaggac tacttccccg aaccggtgac ggtgtcgtgg	480
aactcaggcg cctgaccag cggcgtgcac accttccccg ctgtctaca gtctcagga	540

ctctactccc tcagcagcgt ggtgaccgtg ccctccagca gcttgggcac gaagacctac	600
acctgcaacg tagatcacaa gccagcaac accaagggtg acaagagagt tgagtccaaa	660
tatgggtcccc catgcccacc ctgccagca cctgaggccg ccggggggacc atcagtcttc	720
ctgttccccc caaaacccaa ggacactctc atgatctccc ggaccctga ggtcacgtgc	780
gtggtggtgg acgtgagcca ggaagacccc gaggtccagt tcaactggta cgtggatggc	840
gtggaggtgc ataatgccaa gacaaagccg cgggaggagc agttcaacag cacgtaccgt	900
gtggtcagcg tcctcaccgt cctgcaccag gactggctga acggcaagga gtacaagtgc	960
aaggctctcca acaaaagcct cccgtcctcc atcgagaaaa ccatctccaa agccaaaagg	1020
cagccccgag agccacaggt gtacaccctg ccccatccc aggaggagat gaccaagaac	1080
caggtcagcc tgacctgcct ggtcaaaggc ttctacccca gcgacatcgc cgtggagtgg	1140
gaaagcaatg ggagccgga gaacaactac aagaccacgc ctcccgtgct ggactccgac	1200
ggctccttct tcctctacag caggctaacc gtggacaaga gcaggtggca ggaggggaat	1260
gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacaca gaagagcctc	1320
tccctgtctc tgggt	1335

<210> 24

<211> 1335

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 24

gaggtgcagc tgggtggagtc tgggggaggc ttggtccagc ctggggggtc cctgagactc	60
tcctgtgcag cctctggatt cactttcagt agctatacca tgtcttgggt ccgccaggct	120
ccagggaagg ggctggagtg ggtggccacc atttcgggtg gtggtttcgg cacatactat	180
cccgacagtg tgaagggtcg attcaccatc tccagagaca acgccaagaa ctcaactgtat	240
ctgcaaatga acagcctgag agccgaggac acggctgtgt attactgtgc gagacctgga	300
tataataact actactttga catctggggc caagggacca cggtcaccgt ctctcagcc	360
tcaccaagg gcccatcggc ctcccgtcta gcgccctgct ccaggagcac ctccgagagc	420
acagccgccc tgggctgcct ggtcaaggac tacttccccg aaccggtgac ggtgtcgtgg	480
aactcaggcg ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtctcagga	540
ctctactccc tcagcagcgt ggtgaccgtg ccctccagca gcttgggcac gaagacctac	600
acctgcaacg tagatcacaa gccagcaac accaagggtg acaagagagt tgagtccaaa	660
tatgggtcccc catgcccacc ctgccagca cctgaggccg ccggggggacc atcagtcttc	720
ctgttccccc caaaacccaa ggacactctc atgatctccc ggaccctga ggtcacgtgc	780
gtggtggtgg acgtgagcca ggaagacccc gaggtccagt tcaactggta cgtggatggc	840
gtggaggtgc ataatgccaa gacaaagccg cgggaggagc agttcaacag cacgtaccgt	900

gtggtcagcg tcctcacgt cctgcaccag gactggctga acggcaagga gtacaagtgc	960
aaggtctcca acaaaggcct cccgtcctcc atcgagaaaa ccatctccaa agccaaagg	1020
cagccccgag agccacaggt gtacaccctg ccccatccc aggaggagat gaccaagaac	1080
caggtcagcc tgacctgcct ggtcaaaggc ttctacccca gcgacatcgc cgtggagtgg	1140
gaaagcaatg ggcagccgga gaacaactac aagaccacgc ctcccgctgt ggactccgac	1200
ggctccttct tcctctacag caggctaacc gtggacaaga gcaggtggca ggaggggaat	1260
gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacaca gaagagcctc	1320
tcctgtctc tgggt	1335

<210> 25

<211> 642

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 25

gaaatttgtg tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc	60
ctctcctgcc acgccagcga cagtattagc aacagcctac actggtacca acagaaacct	120
ggccaggctc ccaggctcct catctattat ggcagacagt ccatccaggg catcccagcc	180
aggttcagtg gcagtgggtc tgggacagac ttactctca ccatcagcag cctagagcct	240
gaagattttg cagttatta ctgtcaacag agtgagagct ggccgctcca cttcggcgga	300
gggaccaagg tggagatcaa acgaactgtg gctgcaccat ctgtcttcat cttcccgcca	360
tctgatgagc agttgaaatc tggaaactgcc tctgttgtgt gcctgctgaa taacttctat	420
cccagagagg ccaaagtaca gtggaagggtg gataacgcc tccaatcggg taactcccag	480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg	540
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc	600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gc	642

<210> 26

<211> 642

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 26

gaaatttgtg tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc	60
ctctcctgcc acgccagcga cagtattagc aacagcctac actggtacca acagaaacct	120
ggccaggctc ccaggctcct catctattat gctagacagt ccatccaggg catcccagcc	180
aggttcagtg gcagtgggtc tgggacagac ttactctca ccatcagcag cctagagcct	240

```

--  - - - - -
gaagattttg cagtttatta ctgtcaacag agtgagagct ggccgctcca cttcggcgga      300
gggaccaagg tggagatcaa acgaactgtg gctgcaccat ctgtcttcat cttcccgcca      360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taactttctat      420
cccagagagg ccaaagtaca gtggaagggtg gataacgccc tccaatcggg taactcccag      480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg      540
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc      600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gc                          642

```

<210> 27

<211> 642

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 27

```

gaaatttgtg tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc      60
ctctcctgcc acgccagcga cagtattagc aacagcctac actggtacca acagaaacct      120
ggccaggctc ccaggctcct catctattat ggcagacagt ccatccaggg catcccagcc      180
aggttcagtg gcagtgggtc tgggacagac ttactctca ccatcagcag cctagagcct      240
gaagattttg cagtttatta ctgtcaacag agtgccagct ggccgctcca cttcggcgga      300
gggaccaagg tggagatcaa acgaactgtg gctgcaccat ctgtcttcat cttcccgcca      360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taactttctat      420
cccagagagg ccaaagtaca gtggaagggtg gataacgccc tccaatcggg taactcccag      480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg      540
ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc      600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gc                          642

```

<210> 28

<211> 642

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 28

```

gaaatttgtg tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc      60
ctctcctgcc acgccagcga cagtattagc aacagcctac actggtacca acagaaacct      120
ggccaggctc ccaggctcct catctattat gctagacagt ccgagcaggg catcccagcc      180
aggttcagtg gcagtgggtc tgggacagac ttactctca ccatcagcag cctagagcct      240

```

```

gaagattttg cagtttatta ctgtcaacag agtgccagct ggccgctcca cttcggcgga      300
gggaccaagg tggagatcaa acgaactgtg gctgcaccat ctgtcttcat cttcccgccca      360
tctgatgagc agttgaaatc tggaactgcc tctgttgtgt gcctgctgaa taactttcat      420
cccagagagg ccaaagtaca gtggaaggtg gataacgcc tccaatcggg taactcccag      480
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg      540
ctgagcaaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc      600
ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gc                          642

```

<210> 29

<211> 235

<212> PRT

<213> Homo sapiens

<400> 29

```

Thr Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro
1           5           10           15

```

```

Pro Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala
20           25           30

```

```

Pro Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn
35           40           45

```

```

Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu
50           55           60

```

```

Tyr Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys
65           70           75           80

```

```

Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys
85           90           95

```

```

Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln
100          105          110

```

```

Asn His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly
115          120          125

```

```

Asn Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser
130          135          140

```

```

Ser Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg
145          150          155          160

```

```

Ser Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser
165          170          175

```

```

Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His
180          185          190

```

```

Arg Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys
195          200          205

```

195 200 205
 Gly Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser
 210 215 220

 Asn Ser Ser Arg Leu His Thr Cys Gln Arg His
 225 230 235

 <210> 30
 <211> 234
 <212> PRT
 <213> Cebus albifrons

 <400> 30
 Thr Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro
 1 5 10 15

 Pro Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala
 20 25 30

 Pro Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn
 35 40 45

 Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu
 50 55 60

 Tyr Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys
 65 70 75 80

 Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys
 85 90 95

 Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln
 100 105 110

 Asn Asn Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly
 115 120 125

 Asn Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser
 130 135 140

 Ser Lys Met Tyr His Ser Lys Gly Gln Glu Gly Ser Val Cys Leu Arg
 145 150 155 160

 Ser Ser Asp Cys Ala Thr Gly Leu Cys Cys Ala Arg His Phe Trp Ser
 165 170 175

 Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His
 180 185 190

 Arg Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys
 195 200 205

 Gly Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser
 210 215 220

Asn Ser Ser Arg Leu His Thr Cys Gln Arg
225 230

<210> 31

<211> 252

<212> PRT

<213> Rattus rattus

<400> 31

Thr Leu Asn Ser Val Leu Ile Asn Ser Asn Ala Ile Lys Asn Leu Pro
1 5 10 15

Pro Pro Leu Gly Gly Ala Gly Gly Gln Pro Gly Ser Ala Val Ser Val
20 25 30

Ala Pro Gly Val Leu Tyr Glu Gly Gly Asn Lys Tyr Gln Thr Leu Asp
35 40 45

Asn Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp
50 55 60

Glu Tyr Cys Ser Ser Pro Ser Arg Gly Ala Ala Gly Val Gly Gly Val
65 70 75 80

Gln Ile Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His
85 90 95

Ala Met Cys Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Met Pro
100 105 110

Ser Asp His Ser His Leu Pro Arg Gly Glu Ile Glu Glu Gly Ile Ile
115 120 125

Glu Asn Leu Gly Asn Asp His Gly Ala Gly Asp Gly Tyr Pro Arg Arg
130 135 140

Thr Thr Leu Thr Ser Lys Ile Tyr His Thr Lys Gly Gln Glu Gly Ser
145 150 155 160

Val Cys Leu Arg Ser Ser Asp Cys Ala Thr Gly Leu Cys Cys Ala Arg
165 170 175

His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val
180 185 190

Cys Thr Lys His Arg Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln
195 200 205

Arg Cys Tyr Cys Gly Glu Gly Leu Ala Cys Arg Ile Gln Lys Asp His
210 215 220

His Gln Thr Ser Asn Ser Ser Arg Leu His Thr Cys Gln Arg His Ala
225 230 235 240

Phe Ile Asp Tyr Lys Asp Asp Asp Asp Lys His Val
245 250

<210> 32

<211> 236

<212> PRT

<213> *Lepus americanus*

<400> 32

Thr Leu Asn Ser Val Leu Val Asn Ser Asn Ala Ile Lys Asn Leu Pro
1 5 10 15

Pro Pro Leu Gly Gly Ala Asn Gly His Pro Gly Ser Ala Val Ser Ala
20 25 30

Thr Pro Gly Ile Leu Tyr Glu Gly Gly Asn Lys Tyr Leu Pro Leu Asp
35 40 45

Asn Tyr Gln Pro Tyr Pro Cys Thr Glu Asp Glu Glu Cys Gly Thr Asp
50 55 60

Glu Tyr Cys Ala Ser Pro Ala Arg Gly Gly Gly Ala Gly Val Gln Ile
65 70 75 80

Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met
85 90 95

Cys Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Met Pro Ser Asp
100 105 110

His Asn His Phe His Arg Gly Glu Ile Glu Glu Thr Ile Val Glu Ser
115 120 125

Phe Gly Asn Asp His Ser Thr Ser Asp Gly Tyr Ser Arg Arg Thr Thr
130 135 140

Leu Ser Ser Lys Met Tyr His Ala Lys Gly Gln Glu Gly Ser Val Cys
145 150 155 160

Leu Arg Ser Ser Asp Cys Ala Thr Gly Leu Cys Cys Ala Arg His Phe
165 170 175

Trp Ser Lys Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr
180 185 190

Lys His Arg Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys
195 200 205

Tyr Cys Gly Asp Gly Leu Ser Cys Arg Leu Gln Asn Asp Gln His Glu
210 215 220

Ala Ser Asn Ser Ser Arg Leu His Thr Cys Gln Arg
225 230 235

<210> 33

<211> 241

<212> PRT

<213> Mus musculus

<400> 33

Thr Leu Asn Ser Val Leu Ile Asn Ser Asn Ala Ile Lys Asn Leu Pro
1 5 10 15

Pro Pro Leu Gly Gly Ala Gly Gly Gln Pro Gly Ser Ala Val Ser Val
20 25 30

Ala Pro Gly Val Leu Tyr Glu Gly Gly Asn Lys Tyr Gln Thr Leu Asp
35 40 45

Asn Tyr Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Ser Asp
50 55 60

Glu Tyr Cys Ser Ser Pro Ser Arg Gly Ala Ala Gly Val Gly Gly Val
65 70 75 80

Gln Ile Cys Leu Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Thr His
85 90 95

Ala Met Cys Cys Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Met Pro
100 105 110

Ser Asp His Ser His Phe Pro Arg Gly Glu Ile Glu Glu Ser Ile Ile
115 120 125

Glu Asn Leu Gly Asn Asp His Asn Ala Ala Ala Gly Asp Gly Tyr Pro
130 135 140

Arg Arg Thr Thr Leu Thr Ser Lys Ile Tyr His Thr Lys Gly Gln Glu
145 150 155 160

Gly Ser Val Cys Leu Arg Ser Ser Asp Cys Ala Ala Gly Leu Cys Cys
165 170 175

Ala Arg His Phe Trp Ser Lys Ile Cys Lys Pro Val Leu Lys Glu Gly
180 185 190

Gln Val Cys Thr Lys His Lys Arg Lys Gly Ser His Gly Leu Glu Ile
195 200 205

Phe Gln Arg Cys Tyr Cys Gly Glu Gly Leu Ala Cys Arg Ile Gln Lys
210 215 220

Asp His His Gln Ala Ser Asn Ser Ser Arg Leu His Thr Cys Gln Arg
225 230 235 240

His

<210> 34

<211> 214

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 34

Asp Ile Leu Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15

Asp Ser Val Ser Leu Ser Cys Arg Ala Ser Asp Ser Ile Ser Gly Ser
 20 25 30

Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45

Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Thr
 65 70 75 80

Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asn Ser Trp Pro Leu
 85 90 95

Asn Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg Ala Asp Ala Ala
 100 105 110

Pro Thr Val Ser Ile Phe Pro Pro Ser Thr Glu Gln Leu Ala Thr Gly
 115 120 125

Gly Ala Ser Val Val Cys Leu Met Asn Asn Phe Tyr Pro Arg Asp Ile
 130 135 140

Ser Val Lys Trp Lys Ile Asp Gly Thr Glu Arg Arg Asp Gly Val Leu
 145 150 155 160

Asp Ser Val Thr Asp Gln Asp Ser Lys Asp Ser Thr Tyr Ser Met Ser
 165 170 175

Ser Thr Leu Ser Leu Ser Lys Ala Asp Tyr Glu Ser His Asn Leu Tyr
 180 185 190

Thr Cys Glu Val Val His Lys Thr Ser Ser Ser Pro Val Val Lys Ser
 195 200 205

Phe Asn Arg Asn Glu Cys
 210

<210> 35

<211> 445

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 35

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	1	5	10	15
Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr	20	25	30	
Thr	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Glu	Lys	Arg	Leu	Glu	Trp	Val	35	40	45	
Ala	Thr	Ile	Ser	Gly	Gly	Gly	Gly	Asn	Thr	Tyr	Tyr	Pro	Asp	Ser	Val	50	55	60	
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	Tyr	65	70	75	80
Leu	Gln	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Leu	Tyr	Tyr	Cys	85	90	95	
Ala	Arg	Pro	Gly	Tyr	Asn	Asn	Tyr	Tyr	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	100	105	110	
Thr	Thr	Leu	Thr	Val	Ser	Ser	Ala	Lys	Thr	Thr	Pro	Pro	Ser	Val	Tyr	115	120	125	
Pro	Leu	Ala	Pro	Gly	Thr	Ala	Leu	Lys	Ser	Asn	Ser	Met	Val	Thr	Leu	130	135	140	
Gly	Cys	Leu	Val	Lys	Gly	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Thr	Trp	145	150	155	160
Asn	Ser	Gly	Ala	Leu	Ser	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	165	170	175	
Gln	Ser	Gly	Leu	Tyr	Thr	Leu	Thr	Ser	Ser	Val	Thr	Val	Pro	Ser	Ser	180	185	190	
Thr	Trp	Pro	Ser	Gln	Thr	Val	Thr	Cys	Asn	Val	Ala	His	Pro	Ala	Ser	195	200	205	
Ser	Thr	Lys	Val	Asp	Lys	Lys	Ile	Val	Pro	Arg	Asn	Cys	Gly	Gly	Asp	210	215	220	
Cys	Lys	Pro	Cys	Ile	Cys	Thr	Gly	Ser	Glu	Val	Ser	Ser	Val	Phe	Ile	225	230	235	240
Phe	Pro	Pro	Lys	Pro	Lys	Asp	Val	Leu	Thr	Ile	Thr	Leu	Thr	Pro	Lys	245	250	255	
Val	Thr	Cys	Val	Val	Val	Asp	Ile	Ser	Gln	Asp	Asp	Pro	Glu	Val	His	260	265	270	
Phe	Ser	Trp	Phe	Val	Asp	Asp	Val	Glu	Val	His	Thr	Ala	Gln	Thr	Arg	275	280	285	

Pro Pro Glu Glu Gln Phe Asn Ser Thr Phe Arg Ser Val Ser Glu Leu
290 295 300

Pro Ile Leu His Gln Asp Trp Leu Asn Gly Arg Thr Phe Arg Cys Lys
305 310 315 320

Val Thr Ser Ala Ala Phe Pro Ser Pro Ile Glu Lys Thr Ile Ser Lys
325 330 335

Pro Glu Gly Arg Thr Gln Val Pro His Val Tyr Thr Met Ser Pro Thr
340 345 350

Lys Glu Glu Met Thr Gln Asn Glu Val Ser Ile Thr Cys Met Val Lys
355 360 365

Gly Phe Tyr Pro Pro Asp Ile Tyr Val Glu Trp Gln Met Asn Gly Gln
370 375 380

Pro Gln Glu Asn Tyr Lys Asn Thr Pro Pro Thr Met Asp Thr Asp Gly
385 390 395 400

Ser Tyr Phe Leu Tyr Ser Lys Leu Asn Val Lys Lys Glu Lys Trp Gln
405 410 415

Gln Gly Asn Thr Phe Thr Cys Ser Val Leu His Glu Gly Leu His Asn
420 425 430

His His Thr Glu Lys Ser Leu Ser His Ser Pro Gly Lys
435 440 445

<210> 36

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 36

Asp Ile Leu Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
1 5 10 15

Asp Ser Val Ser Leu Ser Cys Arg Ala Ser Asp Ser Ile Ser Gly Ser
20 25 30

Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
35 40 45

Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Thr
65 70 75 80

Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asn Ser Trp Pro Leu
 85 90 95

Asn Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

<210> 37

<211> 119

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 37

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Thr Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45

Ala Thr Ile Ser Gly Gly Gly Gly Asn Thr Tyr Tyr Pro Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95

Ala Arg Pro Gly Tyr Asn Asn Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser
 115

<210> 38

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 38

Thr Ile Ser Gly Gly Gly Gly Asn Thr Tyr Tyr Pro Asp Ser Val Lys
 1 5 10 15

Gly

<210> 39

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 39

Pro	Gly	Tyr	Asn	Asn	Tyr	Tyr	Phe	Asp	Tyr
1				5					10

<210> 40

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 40

Arg	Ala	Ser	Asp	Ser	Ile	Ser	Gly	Ser	Leu	His
1				5						10

<210> 41

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 41

Tyr	Ala	Ser	Gln	Ser	Ile	Ser
1				5		

<210> 42

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<400> 42

Gln	Gln	Ser	Asn	Ser	Trp	Pro	Leu	Asn
1				5				

<210> 43

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (7)..(7)

<223> Xaa at position 7 = Phe or Gly

<220>

<221> MISC_FEATURE

<222> (8)..(8)

<223> Xaa at position 8 = Gly or Asn

<400> 43

Thr	Ile	Ser	Gly	Gly	Gly	Xaa	Xaa	Thr	Tyr	Tyr	Pro	Asp	Ser	Val	Lys
1				5					10					15	

Gly

<210> 44

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (4)..(4)

<223> Xaa at position 4 = His or Asn

<400> 44

Pro	Gly	Tyr	Xaa	Asn	Tyr	Tyr	Phe	Asp	Ile
1				5				10	

<210> 45

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (4)..(4)

<223> Xaa at position 4 = His or Asn

<220>

<221> MISC_FEATURE

<222> (10)..(10)

<223> Xaa at position 10 = Ile or Tyr

<400> 45

Pro	Gly	Tyr	Xaa	Asn	Tyr	Tyr	Phe	Asp	Xaa
1				5					10

<210> 46

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (1)..(1)

<223> Xaa at position 1 = His or Arg

<220>

<221> MISC_FEATURE

<222> (8)..(8)

<223> Xaa at position 8 = Asn or Gly

<400> 46

Xaa	Ala	Ser	Asp	Ser	Ile	Ser	Xaa	Ser	Leu	His
1				5					10	

<210> 47

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (2)..(2)

<223> Xaa at position 2 = Gly or Ala

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa at position 6 = Ile or Glu

<400> 47

Tyr	Xaa	Arg	Gln	Ser	Xaa	Gln
1				5		

<210> 48

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (2)..(2)

<223> Xaa at position 2 = Gly or Ala

<220>

<221> MISC_FEATURE

<222> (3)..(3)

<223> Xaa at position 3 = Arg or Ser

<220>

<221> MISC_FEATURE

<222> (6)..(6)

<223> Xaa at position 6 = Ile or Glu

<220>

<221> MISC_FEATURE

<222> (7)..(7)

<223> Xaa at position 7 = Gln or Ser

<400> 48

Tyr	Xaa	Xaa	Gln	Ser	Xaa	Xaa
1				5		

<210> 49

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (4)..(4)

<223> Xaa at position 4 = Glu or Ala

<400> 49

Gln Gln Ser Xaa Ser Trp Pro Leu His
1 5

<210> 50

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (4)..(4)

<223> Xaa at position 4 = Glu, Ala or Asn

<220>

<221> MISC_FEATURE

<222> (9)..(9)

<223> Xaa at position 9 = His or Asn

<400> 50

Gln Gln Ser Xaa Ser Trp Pro Leu Xaa
1 5

<210> 51

<211> 119

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (102)..(102)

<223> Xaa at position 102 = His or Asn

<400> 51

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Thr Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Ala Thr Ile Ser Gly Gly Gly Phe Gly Thr Tyr Tyr Pro Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Pro Gly Tyr Xaa Asn Tyr Tyr Phe Asp Ile Trp Gly Gln Gly
100 105 110

Thr Thr Val Thr Val Ser Ser 115

<210> 52

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic Construct

<220>

<221> MISC_FEATURE

<222> (51)..(51)

<223> Xaa at position 51 = Gly or Ala

<220>

<221> MISC_FEATURE

<222> (55)..(55)

<223> Xaa at position 55 = Ile or Glu

<220>

<221> MISC_FEATURE

<222> (92)..(92)

<223> Xaa at position 92 = Glu or Ala

<400> 52

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys His Ala Ser Asp Ser Ile Ser Asn Ser
20 25 30

Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile
35 40 45

Tyr Tyr Xaa Arg Gln Ser Xaa Gln Gly Ile Pro Ala Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
65 70 75 80

Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Xaa Ser Trp Pro Leu
 85 90 95
 His Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2007084344A \[0003\]](#)
- [WO2006015373A \[0005\]](#)
- [US4816397A \[0048\]](#)
- [US5225539A \[0048\]](#)
- [US5693761A \[0048\]](#)
- [US6116841B \[0109\]](#)
- [US2010030039W \[0109\]](#)
- [US10712681 \[0109\]](#)

Non-patent literature cited in the description

- **FULCINTINI MARIATERESA et al.** Anti-DKK1 mAb (BHQ880) as a potential therapeutic agent for multiple myeloma *Blood*, 2009, vol. 114, 2371-379 [\[0004\]](#)
- **GAVIATOPOULOU MARIA et al.** Dickkopf-1: a suitable target for the management of myeloma bone disease *Expert Opinion on Therapeutic Targets*, 2009, vol. 13, 7839-848 [\[0004\]](#)
- Antibody-based inhibition of DKK1 suppresses tumor-induced bone resorption and multiple myeloma growth in vivo **YACCOBY SHMEUL et al.** *Blood American Society of Hematology* 20070301 vol. 109, 2106-211 [\[0004\]](#)
- **ETTENBERG SETH et al.** BHQ880, a novel anti-DKK1 neutralizing antibody, inhibits tumor-induced osteolytic bone disease *Proceedings of the Annual Meeting of the*

- American Association for Cancer Research, 2008, vol. 49, 947- [0004]
- **HARLOWLANE**Antibodies, A Laboratory ManualCold Spring Harbor Laboratory Press19880000 [0044]
 - **MARIE-PAULE LEFRANCGERARD LEFRANC**The Immunoglobulin FactsBookAcademic Press20010000 [0047]
 - **QUEEN et al.**Proc. Natl. Acad. Sci. USA, 1991, vol. 88, 2869- [0048]
 - **LEVITTJ.** Mol. Biol., 1983, vol. 168, 595-620 [0048]
 - **JONES et al.**Nature, 1986, vol. 321, 522-525 [0048]
 - **RIECHMANN et al.**Nature, 1988, vol. 332, 323-327 [0048]
 - **VERHOEYEN et al.**Science, 1988, vol. 239, 1534-1536 [0048]
 - **WU et al.**J. Mol. Biol., 1999, vol. 294, 151-162 [0052]
 - **HARLOWLANE**Antibodies, A Laboratory ManualCold Spring Harbor Laboratory Press19880000567-569 [0054]
 - **A. GENNARO et al.**Remington: The Science and Practice of PharmacyMack Publishing Co.19950000 [0060]

P a t e n t k r a v

5 **1.** DKK-1-antistof eller antigen-bindende fragment deraf, omfattende en variabel region med let kæde (LCVR) og en variabel region med tung kæde (HCVR), hvor LCVR'en omfatter aminosyresekvensen med SEQ ID NO: 36, og HCVR'en omfatter aminosyresekvensen med SEQ ID NO: 37.

10 **2.** DKK-1-antistof ifølge krav 1, hvor DKK-1-antistoffet omfatter en let kæde omfattende aminosyresekvensen med SEQ ID NO: 34 og en tung kæde omfattende aminosyresekvensen med SEQ ID NO: 35.

15 **3.** A DKK-1-antistof ifølge krav 1 eller krav 2, omfattende to lette kæder, hvor hver let kæde omfatter aminosyresekvensen med SEQ ID NO: 34, og to tunge kæder, hvor hver tung kæde omfatter aminosyresekvensen med SEQ ID NO: 35.

4. Farmaceutisk sammensætning omfattende DKK-1-antistoffet eller det antigen-bindende fragment deraf ifølge et af kravene 1 til 3, og en farmaceutisk acceptabel bærer, fortyndingsmiddel eller hjælpestof.