



(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/US2017/034193

(22) International Filing Date:
24 May 2017 (24.05.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/341,140 25 May 2016 (25.05.2016) US

(71) Applicant: **ALEXION PHARMACEUTICALS, INC.**
[US/US]; 100 College Street, New Haven, CT 06510 (US).

(72) Inventors; and

(71) Applicants: **DAVIS, Jared** [US/US]; 310 Thornton Street,
Hamden, CT 06517 (US). **DEBEAR, Joanna** [US/US]; 11
Shire Court, Cheshire, CT 06410 (US). **CHENG, Christo-
pher** [US/US]; 114 Bedford Avenue, Hamden, CT 06517
(US).

(74) Agent: **YAMAUCHI, Michael**; Alexion Pharmaceuticals,
Inc., 100 College St., New Haven, CT 06510 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))

(54) Title: POLYNUCLEOTIDE PURIFICATION WITH MONOLITH COLUMNS

(57) Abstract: Described herein are methods of purifying polynucleotides,
e.g., imRNA and oligonucleotides, *e.g.*, probes, primers and siRNA, using
monolithic columns with immobilized ligands coupled to the monolithic col-
umn. Also described are monolithic columns for purifying polynucleotides
from a sample; and methods of preparing such columns.




FIG. 1

Published:

— *with international search report (Art. 21(3))*

Polynucleotide Purification with Monolith Columns

BACKGROUND

[0001] Messenger RNA, or mRNA, is a key intermediary in the conversion of genetic information into biologically active proteins. Many aspects of biomedical research and drug development depend on the ability to obtain high-quality, purified mRNA. Several properties of mRNA, however, make its purification challenging. Relative to total RNA, mRNA exists in very low copy numbers in cells. Furthermore, mRNA is highly sensitive to degradation by RNase enzymes, further compounding the difficulties of purification.

[0002] Current methods of isolating mRNA take advantage of the poly-adenine (poly-A) tail present on mRNA molecules. Oligonucleotides consisting of stretches of the nucleic acid base, deoxythymine (oligo-dT), are used to bind to the complementary poly-A tails of mRNA molecules. Oligo-dT-cellulose affinity chromatography has been used to purify mRNA from total RNA fractions.

[0003] Purification of mRNA using traditional chromatographic methods, however, is inefficient. Cellulose and other particle-based chromatography columns contain small pore sizes causing slow diffusion of large biomolecules such as mRNA and large contaminants or other particles. Consequently, mRNA purification over traditional particle-based columns is characterized by low flow rates, poor yields and extensive processing time. This problem is further exacerbated when mRNA, formulated for therapeutic delivery, are to be purified, as such formulated mRNA are even larger than naked mRNA.

[0004] There remains a need for improved methods of purifying mRNA molecules and other polynucleotides, both naked and formulated, to high purity with a high yield and lower processing time.

SUMMARY

[0005] Described herein are compositions and methods for purifying polynucleotides, both naked and formulated, *e.g.*, mRNA formulated in lipid nanoparticles (LNPs). Polynucleotides are purified from contaminants, such as, for example, other biomolecules, such as DNA, ribosomal and transfer RNA, and proteins, using monolithic column

chromatography. Where formulated polynucleotides are purified, contaminants also include unformulated polynucleotide ("free" or "naked" polynucleotides).

[0006] In one embodiment, the disclosure is directed to a method of separating a formulated polynucleotide from free polynucleotide, the method comprising: a) loading a sample onto a monolith matrix comprising a ligand comprising: i) a reactive moiety coupled to the monolith matrix, and ii) an affinity moiety that binds to the free polynucleotide but not the formulated polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety; and b) collecting the formulated polynucleotide from the column while the free polynucleotide remains immobilized on the monolith matrix. In a particular embodiment, the monolith matrix is contained in a column. In a particular embodiment, the formulated polynucleotide is a formulated mRNA. In a particular embodiment, the mRNA is formulated in a lipid nanoparticle. In a particular embodiment, the ligand is an oligo-dT probe. In a particular embodiment, the ligand is $\text{NH}_2\text{-C}_X\text{-dT}_Y$, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30. In a particular embodiment, the ligand is $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$. In a particular embodiment, the ligand further comprises a carbon linker positioned between the reactive moiety and the ligand. In a particular embodiment, the carbon linker is C_X , where X is a whole number between about 1 and about 50. In a particular embodiment, the methods described herein further comprise eluting the free polynucleotide from the monolith matrix by reducing the ionic strength of the liquid phase.

[0007] In one embodiment, the disclosure is directed to a method of purifying a polynucleotide from a sample, the method comprising: a) loading the sample onto a monolithic matrix comprising a ligand comprising: i) a reactive moiety coupled to the monolithic matrix, and ii) a ligand that binds to the polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety; b) allowing for the polynucleotide to bind to the ligand; and c) eluting the polynucleotide from the monolith matrix after one or more contaminants have been substantially separated from the bound polynucleotide. In a particular embodiment, the reactive moiety is a primary amine. In a particular embodiment, the monolithic matrix is activated with an activating agent selected from carbonyldiimidazole, epoxy, ethylenediamine (EDA), carbodiimide, aldehyde, anhydride, imidoester and NHS ester. In a particular embodiment, the ligand further

comprises a carbon linker positioned between the reactive moiety and the ligand. In a particular embodiment, the carbon linker is C_X , where X is a whole number between about 1 and about 50. In a particular embodiment, the polynucleotide is mRNA. In a particular embodiment, the ligand is an oligo-dT probe. In a particular embodiment, the ligand is $NH_2-C_X-dT_Y$, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30. In a particular embodiment, the ligand is $NH_2-C_{12}-dT_{18}$. In a particular embodiment, the methods described herein further comprise washing the column prior to eluting the polynucleotide, *e.g.*, wherein the wash buffer contains a salt concentration of at least 200 mM, and the elution buffer contains a salt concentration of 100 mM or less, wherein the wash buffer comprises one or more salts selected from sodium sulfate, sodium chloride and sodium phosphate. In a particular embodiment, the elution buffer is selected from water and Tris. In a particular embodiment, the flow rate of the column is at least 0.5 mL/min or 0.5 CV/min (*e.g.*, in a 1 mL column). Column volume is abbreviated "CV."

[0008] In one embodiment, the disclosure is directed to a column for purifying a polynucleotide from a sample, said column comprising: a) a monolithic matrix; and b) a ligand comprising a reactive moiety coupled to the monolithic matrix, and a ligand that binds to the polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety. In a particular embodiment, the reactive moiety is a primary amine. In a particular embodiment, the monolithic matrix is activated with an activating agent selected from carbonyldiimidazole, epoxy, ethylenediamine (EDA), carbodiimide, aldehyde, anhydride, imidoester and NHS ester. In a particular embodiment, the ligand is an oligo-dT probe. In a particular embodiment, the ligand further comprises a carbon linker positioned between the reactive moiety and the oligo-dT probe. In a particular embodiment, the carbon linker is C_X , where X is a whole number between 1 and 50. In a particular embodiment, the ligand is $NH_2-C_X-dT_Y$, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30. In a particular embodiment, the ligand is $NH_2-C_{12}-dT_{18}$.

[0009] In one embodiment, the disclosure is directed to a method of preparing a column described herein by a method comprising: a) treating the monolithic matrix with an activating agent to produce an activated monolithic matrix; and b) incubating the activated monolithic matrix in the presence of a ligand comprising a reactive moiety. In a particular

embodiment, the activating agent is selected from carbonyldiimidazole, epoxy, ethylenediamine (EDA), carbodiimide, aldehyde, anhydride, imidoester and NHS ester.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] FIG. 1 shows chromatogram traces from an RNA001-10-9 chromatography run at 200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7 over the $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$ immobilized monolith disk column, as described in Example 2.

[0011] FIG. 2 shows a chromatogram (top) and glyoxyl gel image (bottom) from the RNA001 chromatography run described in Example 2. M = markers, S = starting material, D = linearized DNA template, C6-C9 = main peak fractions.

[0012] FIG. 3 shows a chromatogram of RNA021 transcription (without poly-A tail) over the $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$ -immobilized monolith disk, as described in Example 2.

[0013] FIG 4 shows an overlay of chromatogram traces from RNA001 (with poly-A tail) and RNA021 (without poly-A tail) run under the same conditions on the $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$ -immobilized monolithic column, as described in Example 2.

[0014] FIG. 5 shows a chromatogram of the transcription reaction of RNA001 tested on the $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$ monolith disk, as described in Example 3.

[0015] FIG. 6 shows RNA transcription reactions purified on an $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$ immobilized monolithic column and then run on a 1% E-Gel[®], as described in Example 3. The transcription reactions were loaded onto the monolithic column without further purification following transcription.

[0016] FIG. 7. shows an overlay of RNA001 chromatograms tested at 2, 3 and 4 mL/minute flow rate, as described in Example 4.

[0017] FIG. 8 is a map of DNA001 with two additional linearization sites to remove the poly-A tail, as described in Example 5.

[0018] FIG. 9 shows a chromatogram of RNA001 samples with and without the poly-A tail, as described in Example 5.

[0019] FIG. 10 shows a 1% E-Gel[®] of fractions eluted from an oligo-dT ligand immobilized monolithic column. The sample applied to the column contained RNA transcripts with and without poly-A tails, as described in Example 5.

[0020] FIG. 11 shows an overlay of chromatogram traces from RNA001 purified over two monolithic columns with oligo-dT₁₈ ligands containing either a 6-carbon linker (C₆) or a 12-carbon linker (C₁₂), as described in Example 6. The trace labeled * is the A₂₆₀ of RNA001 purified over the C₁₂ oligo-dT monolith, and the trace labeled ** is the A₂₆₀ of RNA001 purified over the C₆ oligo-dT monolith.

[0021] FIG. 12 is an overlay of chromatograms of RNA001 purified using sodium sulfate- vs. sodium chloride-based buffers (overlapping traces are labeled with *).

[0022] FIG. 13 shows a chromatogram of RNA001 binding to the 1 mL NH₂-C₁₂-dT₁₈ monolith and elution using 10 mM Tris pH 7.5.

[0023] FIG. 14 is a set of gels showing separation of free mRNA, which is eluted (lane 10, top panel) and LNP-formulated mRNA (which comes off in the flow-through, lanes 1-6, bottom panel). The top panel shows intact, LNP-formulated mRNAs loaded onto a gel; the bottom panel shows the same LNP-formulated mRNAs after the LNP has been treated with detergent to lyse the LNP (note the “smiling” of the gel was due to high salt concentrations of the samples during lysis). For both panels, Lane M = Sample Load, Lanes 1-6 = flow-through, Lanes 7-8 = wash and Lane 10 = elution. Note mRNA in wells on the top gel, mRNA within LNPs have hindered electrophoretic mobility. Lysing of the LNPs shows that the load and flow-through fractions contained mRNA that was sequestered in LNPs.

[0024] FIG. 15 is a bar graph showing the improvement of encapsulation efficiency (EE) that results from LNP purification from free mRNA. LNP load materials with EE = 84-87% were purified to EE = 94-96%. Spiking LNPs with free mRNA (1:1) was purified from EE = 53% to EE = 79%. The yield for the purification procedure was ~70%.

[0025] FIGS. 16A-C are chromatograms (A₂₆₀) showing elution profiles for various buffer gradients that were tested for purifying mRNA (RNA025). The top line represents the gradient (% buffer B), and the bottom line is the chromatogram. Oligonucleotides complementary to the 5' end of the mRNA were designed following the T7 polymerase start site (18mer and 24mer oligos were tested; 18mer results are shown). For each oligonucleotide, a 6-carbon modifier was attached to the 3' end and then conjugated to the monolith. FIG. 16A shows the results of a gradient wash. FIG. 16B shows results using a step wash at a conductivity level just before material starts to elute in the gradient wash

(FIG. 16A). This step wash resulted in a significantly sharper elution peak in 10 mM Tris. A significant amount of material was still bound to the column, however, and only removed by a NaOH cleaning step for both of the chromatography runs. Therefore, a step elution at 2M, 4M, 6M and 8M urea was tested (FIG. 16C). As the A260 trace of the chromatogram shows, the RNA bound to the column was completely removed during the urea step elution before reaching the 10 mM NaOH cleaning step. Going forward, 4M urea was selected as the elution condition for all subsequent chromatography purification testing as the majority of the RNA was eluted from the monolith under these conditions. ALK2 = 5' oligo(18)C6dT 3'. Buffer B: 50 mM sodium phosphate, 10 mM EDTA, pH = 7.0.

DETAILED DESCRIPTION

[0026] Described herein are compositions and methods of purifying polynucleotides and formulated polynucleotides, *e.g.*, DNA, or RNA, *e.g.*, mRNA, oligonucleotides, *e.g.*, probes, primers and siRNA, or artificial or synthetic polynucleotides, from contaminants.

Contaminants include, for example, other biomolecules, such as DNA, ribosomal and transfer RNA and proteins. In the case of formulated nucleotides, *e.g.*, polynucleotides enveloped within a lipid nanoparticle (LNP), contaminants also included unformulated nucleotides ("free" polynucleotides). The materials and methods described herein comprise using monolithic column chromatography. The materials and methods described herein relate to unexpected findings that immobilization of polynucleotide ligands, *e.g.*, oligo-deoxythymine (oligo-dT) and sequence-specific or non-specific oligonucleotides or affinity moieties, on monolithic chromatography columns allows for improved purification of polynucleotides, *e.g.*, polynucleotides comprising poly-A. As described herein, any affinity moiety, *e.g.*, a sequence-specific polynucleotide, can be used in conjunction with monolith columns to achieve polynucleotide purification, *e.g.*, separation of formulated polynucleotides from free polynucleotides. The methods described herein are applicable to immobilizing a ligand via an active moiety to an activated monolith matrix, wherein the ligand specifically binds to the polynucleotide to be purified, *e.g.*, through sequence-specific binding, through hybridization or other base-pairing interactions, or through chemical and non-chemical interactions.

[0027] The present disclosure is not limited to the particular embodiments of the disclosure described below, as variations of the particular embodiments can be made that still fall within the scope of the appended Claims. The terminology employed is for the purpose of describing particular embodiments, and is not intended to be limiting. The singular forms “a,” “an” and “the” include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art.

[0028] Described herein is a solid support medium comprising attached polynucleotides or affinity ligands for the purification of biomolecules that specifically bind to the attached polynucleotides or affinity ligands. The solid support medium, for example, can be a column used to purify mRNA from a sample, said column comprising a monolithic matrix coupled to, for example, a ligand comprising an oligo-dT probe. The material of interest to be purified, for example, can be the material that binds to the ligand. Alternatively, the material of interest to be purified can be material that does not bind to the ligand, with a primary contaminant being bound to the ligands instead.

[0029] The terms “monolith,” “monolithic matrix” and “monolithic column” are used interchangeably herein to refer to a chromatography column composed of a continuous stationary phase made of a polymer matrix. In contrast to particle-based chromatography columns, monolithic columns are made of a porous polymer material with highly interconnected channels and large pore size. While particle-based columns rely on diffusion through pores, separation by monolithic columns occurs primarily by convective flow through relatively large channels (about 1 micron or more). Monolithic columns are commercially available and have been used to purify large biomolecules such as viruses, plasmid DNA, and proteins (Rajamanickam, V. *et al.*, *Chromatography*, 2:195-212, 2015).

[0030] The monolithic matrix may be derived from a variety of materials, such as but not limited to, polymethacrylate, polyacrylamide, polystyrene, silica and cryogels.

[0031] The monolithic matrix may be activated to promote coupling to a reactive moiety. Coupling to the activated monolithic matrix may occur, for example, through the formation of a covalent bond between the activated monolithic matrix and the reactive moiety. In some embodiments, the monolithic matrix is activated to couple to a primary amine group. Activation of the monolithic matrix can be accomplished through any appropriate methods

known in the art (see, *e.g.*, Pfaunmiller, E. *et al.*, *Anal. Bioanal. Chem.*, 405:2133-45, 2013; Hermanson, G., *Bioconjugate Techniques*, 3rd Ed., 2013). Non-limiting examples of activation agents include carbonyldiimidazole (CDI), epoxy ethylenediamine (EDA), carbodiimide, aldehyde, anhydride, imidoester and NHS ester.

[0032] As used herein, the term “ligand” refers to a molecule that preferentially binds, covalently or non-covalently, to a molecule or material of interest. The ligands described herein can further comprise a reactive moiety capable of coupling to a monolithic matrix. An “oligo-dT ligand” is an oligo-dT probe. A “probe” refers to a ligand that selectively interacts, *e.g.*, binds to or hybridizes with, a desired interaction partner, *e.g.*, a specific polynucleotide sequence. A ligand can itself be a polynucleotide, *e.g.*, an oligo-dT probe or an oligonucleotide, that, for example, specifically hybridizes to a sequence of interest, *e.g.*, a poly-A tail or a sequence specific to the polynucleotide to be purified.

[0033] An oligo-dT probe consists of a chain of thymine bases or uracil bases or chemically modified bases of any length appropriate to specifically bind to the poly-A tail of mRNA. Non-limiting examples of oligo-dT probes include oligomers of the formula dT_Y, wherein Y is a whole number between 5 and 30. In specific embodiments, the oligo-dT probe is dT₁₅, dT₁₈, dT₂₀, dT₂₅ or dT₃₀.

[0034] The ligands described herein are coupled or attached to the solid support monolith matrix via a reactive moiety. The monolith can be activated, thereby allowing for coupling to the ligand via the active moiety of the ligand. In a particular embodiment, the monolithic matrix is activated with an activation agent to allow coupling to amine groups, and the reactive moiety of the ligand is a primary amine. In one embodiment the activation agent is carbonyldiimidazole.

[0035] In some embodiments, the ligand further comprises a carbon linker positioned between the reactive moiety and the oligo-dT probe. Selection of the length of the carbon linker is within capabilities of the skilled person. Non-limiting examples of carbon linkers include linkers of the formula C_X, wherein X is any whole number between 5 and 50. In specific embodiments, the carbon linker is C₆, C₇, C₈, C₉, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄ or C₁₅.

[0036] In some embodiments, the ligand is NH₂-C_X-dT_Y, wherein C_X is a carbon chain of length X, and X is a whole number between 5 and 50; and dT_Y is an oligo-dT probe of length Y, and Y is a whole number between 1 and about 100, about 5 and about 50, about

10 and about 30, about 7 and about 26, about 18 and about 24, or between about 5 and about 25. In a particular embodiment the ligand is $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$.

[0037] Also described herein is a method of preparing a column for purifying mRNA from a sample, the method comprising treating a monolithic matrix with an activating agent to produce an activated monolithic matrix; and incubating the activated monolithic matrix in the presence of a ligand comprising a reactive moiety and a polynucleotide, *e.g.*, an oligonucleotide, *e.g.*, an oligo-dT probe. In some embodiments the ligand further comprises a carbon linker positioned between the reactive moiety and the polynucleotide probe. In some embodiments, the reactive moiety is a primary amine. In some embodiments, the activating agent is carbonyldiimidazole.

[0038] Also described herein are methods for purifying polynucleotides, *e.g.*, oligonucleotides, *e.g.*, mRNA or siRNA from a sample. Such methods include, for example, a) loading a sample onto a column comprising: i) a monolithic matrix with an attached ligand comprising: A) a reactive moiety coupled to the monolithic matrix, and B) a polynucleotide, *e.g.*, oligo-dT, probe; b) washing the column; c) eluting the polynucleotide from the column; and d) collecting at least one elution fraction from the column. In one embodiment, step b) comprises washing the column with at least one wash buffer. In another embodiment, step c) comprises eluting the polynucleotides from the column with at least one elution buffer. In another embodiment, the elution fractions of step d) contain mRNA. In some embodiments, the wash buffer contains a salt concentration between about 150 mM to about 1 M. In particular embodiments, the wash buffer contains a salt concentration of at least about 200 mM, at least 400 mM or at least about 750 mM. In some embodiments, the elution buffer contains a salt concentration between 0 and about 100 mM. As used herein, the term "about" means plus or minus 10% of the numerical value of the number with which it is being used. In particular embodiments the elution buffer has a salt concentration of 100 mM or less. In particular embodiments the wash buffer comprises one or more salts selected from sodium sulfate, sodium chloride and sodium phosphate.

[0039] In one embodiment, the elution buffer is water. In another embodiment, the elution buffer comprises Tris. Tris buffer may be used at a concentration from about 1 mM to about 20 mM. In a particular embodiment, the elution buffer comprises 10 mM Tris.

[0040] Selection of the flow rate of the column is within capabilities of the skilled person. In some embodiments, the flow rate of the column is from about 1 mL/min to about 5 mL/min. In particular embodiments the flow rate is at least 2 mL/min, at least 3 mL/min or at least 4 mL/min.

[0041] The molecule or material of interest is separated from contaminants, and can come off the column in any of the flow-through, wash or elution fraction, depending on the nature of the molecule or material of interest and the major contaminant(s).

[0042] The following examples are included for illustrative purposes only and are not intended to limit the scope of the claims.

EXAMPLES

EXAMPLE 1.

[0043] Polynucleotides can be applied to monolith matrices as described herein for mRNA. The mRNA transcripts used in this example are described in Table 1; additional mRNA transcripts purified by the methods described herein are described in Table 2. Transcripts used were either LiCl precipitated or used straight after the transcription reaction following EDTA treatment.

Table 1. Characteristics of mRNA used in purification studies.

RNA ID	Transcript length	Poly-A tail
RNA001	1837	Yes
RNA021	452	No
RNA023	971	Yes

[0044] CDI (carbonyldiimidazole or carboxydiimidazole)-activated monolith disk columns (0.34 mL) were purchased from BIA Separations through High Purity New England (Smithfield, RI). Ligands for immobilization on the CDI-monolithic columns were designed and purchased from Integrated DNA Technologies (Coralville, IA). Two ligands were used in these studies: an NH₂-C₁₂-dT₁₈ ligand, containing a primary amine followed by a 12-carbon linker chain and 18 deoxythymine bases; and an NH₂-C₆-dT₁₈ ligand, containing a primary amine followed by a 6-carbon linker chain and with 18 deoxythymine bases.

Experiments were run using a GE AKTA Avant 25 preparative chromatography system (GE Healthcare Life Sciences).

Table 2. Poly-A containing mRNA of various lengths purified to >90% purity by oligo-dT

ID	Transcript bases
RNA023	971
RNA025	1909
RNA027	1924
RNA034	832
RNA037	1432
RNA181	1467
RNA385	1642

Oligo-dT Immobilization to a monolithic matrix

[0045] A syringe was used to load the oligo-dT ligand onto the monolithic column. All steps were performed at room temperature. The CDI disk column was assembled in the housing according to the manufacturer's instructions. The assembled column was flushed with at least 10 column volumes (CV) of Milli-Q water. The column was then equilibrated with at least 10 CV of suitable buffer (0.5 M Na Phosphate pH 8.0).

[0046] The oligo-dT ligand was dissolved in 0.5 M sodium phosphate (pH 8.0) to a final stock concentration of about 100 mg/mL. Then, 1.5-2.0 mL of ligand was diluted to 3 mg/mL with equilibration buffer and was pushed through the column using a syringe to completely fill the monolith channels. The column was then disconnected from the syringe and sealed with blind fittings. The column was stored at room temperature for 20-24 hours.

[0047] Following incubation with the oligo-dT ligand, the column was rinsed with at least 10 CV of suitable buffer (0.5 M Na Phosphate pH 8.0), and the column was then flushed with at least 10 CV Milli-Q water. The column was equilibrated with loading buffer (50 mM

sodium phosphate, 750 mM sodium sulfate, 10 mM EDTA pH 7.0) for testing with samples of RNA.

Purification Testing

[0048] Initial testing of mRNA binding to the monolithic column with immobilized oligo-dT ligand was done as described in Table 3, in the order stated.

Table 3. Initial purification process with oligo dT-immobilized monolithic column

Step	Buffer	Flow rate, ml/min	CV
Clean	10 mM sodium hydroxide	2	10
Equilibrate	750 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	2	10
Load	RNA	1	
Wash	750 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	6
Wash 2	50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	25
Elution	Ultra-pure water	1	15
Clean	10 mM sodium hydroxide	1	15

[0049] The starting buffers used for purification testing were: Buffer A: 50 mM sodium phosphate, 1.0 M Na₂SO₄, 10 mM EDTA pH 7.0; and Buffer B: 50 mM sodium phosphate, 10 mM EDTA pH 7.0. Fractions from the flow-through were desalted as needed and analyzed appropriately.

EXAMPLE 2: Purification of mRNA using amino-linked oligo-dT probe immobilized on an activated monolithic column

Initial Binding experiments

[0050] Initial conditions for testing purification of mRNA on the NH₂-C₁₂-dT₁₈ immobilized monolithic column were designed using a high salt binding buffer. The RNA bound to the monolith in 750 mM sodium sulfate, 50 mM phosphate buffer, 10 mM EDTA at pH 7.0 and

was eluted with water. Various salt conditions were tested and are listed in Table 4. These experiments were completed using LiCl purified material.

Table 4. Initial binding results of RNA to C12-oligo d(T)18 immobilized monolith

Sample	Load and Wash 1 Buffer Components	Binding Result
RNA001	750 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA, pH 7.0	Bound and eluted with water
RNA001	400 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA, pH 7.0	Bound and eluted with water
RNA001	200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA, pH 7.0	Bound and eluted with water

[0051] A chromatogram of LiCl precipitated RNA001 bound at 200 mM sodium sulfate buffer is shown in FIG. 1. A glyoxyl gel was used to visualize fractions from the RNA001 chromatography, and can be seen in FIG. 2. The flow through fractions (1A1-1A4) were combined and concentrated ten-fold for this gel. The majority of the flow through material was not full-length RNA. The gel indicates that 50 mM sodium phosphate, 10 mM EDTA pH 7.0 (no sodium sulfate) eluted off a small amount of product. Once the flow was shifted to ultra-pure water, the majority of RNA was eluted in a single peak of RNA, and was seen in fractions 1C6 to 1C9 (labeled C6, C7, C8 and C9 in FIG. 2). Gel analysis of the chromatography fractions showed removal of impurities, specifically DNA template and abortives of the RNA that are missing the poly-A tail.

[0052] Following these initial experiments, RNA021 (which had no poly-A tail to interact with the oligo-dT ligand) was assessed using the immobilized monolith disk column. RNA021 was tested using the conditions described in Table 5, in the order stated, and compared to the RNA001 that contains a poly-A tail.

Table 5. Process conditions for testing of RNA021 on the C12 oligo dT(18) monolith.

Step	Buffer	Flow rate, ml/min	CV
Clean	10 mM sodium hydroxide	2	10
Equilibrate	200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	2	10
Load	RNA	1	

Wash	200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	6
Elute	50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	25
Water flush	Ultra-pure water	1	15
Clean	10 mM sodium hydroxide	1	15

[0053] The resulting chromatogram is shown in FIG. 3. RNA elution would be expected to start at approximately 17 mL. FIG. 4 shows an overlay of the RNA021 (without poly-A tail) and RNA001 (with poly-A tail) chromatograms run using the same conditions. RNA elution would be expected to start at approximately 17 mL.

EXAMPLE 3: Purification of transcription reactions.

[0054] The NH₂-C₁₂-dT₁₈ immobilized monolithic column was evaluated using transcription reactions that had not been purified further following *in vitro* transcription. Chromatography conditions for these samples are described in Table 6. The RNA loads in Table 6 were transcription reactions treated with EDTA only. There were slight adjustments made to the CV amount for the wash (increased from 6 to 10 CV) and elution (decreased from 25 to 15 CV).

Table 6. Chromatography process for RNA transcription reactions.

Step	Buffer	Flow rate, ml/min	CV
Clean	10 mM sodium hydroxide	2	10
Equilibrate	200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	2	10
Load	RNA	1	8ml
Wash	200 mM sodium sulfate, 50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	10
Elute	50 mM sodium phosphate, 10 mM EDTA pH 7.0	1	15
Water flush	Ultra-pure water	1	15
Clean	10 mM sodium hydroxide	1	10

[0055] The resulting chromatogram (FIG. 5) shows a large flow-through (FT) peak from the remaining reaction components and products such as excess NTPs that do not bind the

column. FT fractions were desalted and run over a 1% E-Gel[®], along with the peak fractions. FIG. 6 shows the results analyzed by agarose gel. The FT fractions (labeled 1A1 through 1B12 in FIG. 6) contain linearized DNA and what appear to be abortive RNA sequences. The peak fractions (labeled 1C1 through 1C4 in FIG. 6) showed a lower-running diffuse band, which disappears following treatment with RNase A, indicating that it is RNA.

[0056] The results indicate that a transcription reaction can be applied directly to an immobilized monolithic column and purified to the same degree as applying RNA that has been initially purified by LiCl precipitation and buffer exchange.

EXAMPLE 4: Increased flow rates

[0057] To test the influence of flow rate on mRNA purification over the ligand-immobilized monolithic column, flow rates up to 4 mL/minute were tested using the same samples and process conditions. Pressures were below acceptable levels for all flow rates. Overlays of chromatograms at 2, 3 and 4 mL/minute (FIG. 7) showed only minor differences between runs, indicating that increases in flow rate did not change the elution profile of the run. Exemplary column scales and operating parameters utilizing oligo-dT immobilization are described in Table 7.

Table 7. Monolith column sizes and operating parameters

Column volume (mL)	Recommended flow rates	Max mL/min	Max CV/min
0.34*	2-4 mL/min	6mL/min	18
1*	1-5 mL/min	16 mL/min	16
8*	8-60 mL/min	100 mL/min	12.5
80*	80-240 mL/min	400 mL min	5
800	200-1300 mL/min	2000 mL/min	2.5
8000	2000-10000 mL/min	10000mL/min	1.25

*denotes columns have been tested.

EXAMPLE 5: Testing RNA with and without a poly-A tail

[0058] Binding of RNA transcripts where the poly-A tail was absent from the RNA001 transcript was accomplished by digesting the DNA template (DNA001; FIG. 8) for RNA001 using restriction enzymes that cut upstream of the sequence coding for the poly-A tail.

Table 8 describes the resulting RNA transcripts following digestion and transcription of DNA001.

Table 8. Resulting transcripts from digested DNA001

Restriction Enzyme	RNA product length	Poly-A tail
EcoR1	1837	Yes
EcoN1	1662	No
BstB1	1150	No

[0059] Following transcription of these templates, the single RNA was loaded on the ligand-immobilized monolithic column using the conditions listed in Table 5. RNA transcripts without poly-A tails flowed through when applied to the column. An equal parts mixture of the three RNA transcripts listed in Table 8 were applied to the monolithic column. The resulting chromatogram is shown in FIG. 9. Peak fractions collected were applied to a 1% E-Gel[®] to visualize the FT components and elution peak (FIG. 10).

[0060] The shorter transcripts lacking a poly-A tail do not bind to the column and are found in FT fractions (FIG. 10, lanes 3-7). DNA is also observed in the FT (FIG. 10, lanes 3-7). Only the full-length RNA that has the poly-A tail is observed in the elution peak (FIG. 10, lanes 8 and 9). The results confirm that the poly-A tail is required for the bind/elute purification observed in EXAMPLE 2.

EXAMPLE 6: Alternate Linkers.

[0061] To evaluate the effect of the ligand linker on purification efficiency of RNA containing a poly-A tail, a shorter linker (C₆ vs C₁₂) between the amino group and oligo-dT probe was tested. The ligand was attached to a new monolithic column using the same method as described in EXAMPLE 1. Once the newly immobilized ligand was attached, the column was washed and tested with RNA001 to compare binding to the C₁₂ linker version of the ligand.

[0062] FIG. 11 shows an overlay of chromatograms of the same RNA run with the two different linkers. For the C₆-linker column, there was higher A₂₆₀ absorbance observed in the FT and the low salt wash as compared to the material run over the C₁₂-linker column. In addition, the C₁₂-linker purification resulted in about 100% yield (peak labeled * in

FIG. 11) vs. the C6 linker, which was about 65% yield (peak labeled ** in FIG. 11). The difference may be due to the shorter linker arm and proximity of the dT being closer to the monolith hindering full complementary binding of the poly-A tail to the dT stretch of nucleotides.

EXAMPLE 7: Salt Comparison.

[0063] To evaluate the effect of different salts on the purification efficiency, sodium sulfate was replaced with sodium chloride in the equilibration/loading and wash buffers, keeping the phosphate buffer, EDTA and pH the same. FIG. 12 shows an overlay of traces from each of the two salts (* indicates the overlapping traces in FIG. 12). With the same process conditions, the chromatograms rendered from the two salts tested showed no difference in purification of the RNA.

EXAMPLE 8: Elution conditions.

[0064] Experiments for the binding and elution conditions for the RNA from oligo-dT monolithic columns looked at loading and washing in a high salt (at least 200 mM) followed by removal of the salt component of the buffer system. The remaining phosphate and EDTA did not elute the RNA; however the subsequent ultra-pure water flush eluted the RNA in a single tight peak. The absence of conductivity proved to be a potent elution condition.

[0065] Analysis of the chromatograms indicated that the pH of the elution drifted upwards from pH 7 to as high as pH 9. To control the pH during the elution step, a low conductivity buffer (10 mM Tris pH 7.5) was tested on the 1 mL NH₂-C₁₂-oligo-dT₁₈ monolith and implemented for elution of the RNA. FIG. 13 shows a chromatogram trace of the elution step with Tris buffer. The results indicated that mRNA can be eluted from the ligand-immobilized monolithic column using either water or Tris buffer.

[0066] EXAMPLE 9:

[0067] Although mRNA has a short half-life *in vivo*, high doses of free mRNA can transfect cells and tissues. Additionally, unwanted systemic introduction of mRNA can trigger an immune response before degradation and clearance.

[0068] Lipid Nanoparticles (LNPs) can be used to encapsulate and deliver, for example, mRNA. LNPs typically have at least 80% encapsulation of mRNA, *i.e.*, mRNA that is located inside of an LNP as opposed to outside (“free” mRNA). This EXAMPLE evaluates the ability of monolith, oligo-dT purification to remove unencapsulated mRNA to produce a purified LNP.

[0069] The results (FIGS. 14 and 15) demonstrate the general ability of oligo-dT based chromatography to separate free mRNA (with and without chemical modifications) from mRNA formulated in LNPs. This work could extend to purification of LNPs from other impurities besides mRNA if different affinity or immobile phase conditions are used. These data also indicate the strategy for purifying formulated polynucleotides extends to any polynucleotide, *e.g.*, mRNA delivery system, not just LNPs.

[0070] LNPs were formulated with mRNA with encapsulation efficiency greater than 80%. Chromatograms and gels demonstrated that LNPs eluted in the flow-through fractions (FIG. 14). Free mRNA bound to the oligo dT column and eluted with salt adjustment to the mobile phase. Assessment of LNPs before and after purification showed no impact on size and polydispersity (Table 9). LNP yield was ~70% after purification (FIG. 15)

Table 9. LNP characterization before and after purification.

Sample	LNP Load		LNP purified	
	Diameter (nm)	PDI*	Diameter (nm)	PDI
N1-methylpseudouridine	92	0.04	96	0.05
Uridine	95	0.08	89	0.03
Pseudouridine	93	0.05	95	0.07

OTHER EMBODIMENTS

[0071] It is to be understood that the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims. References cited herein are herein incorporated by reference in their entireties.

CLAIMS

What is claimed is:

1. A method of purifying a polynucleotide from a sample, the method comprising:
 - a) loading the sample onto a monolithic matrix comprising a ligand comprising:
 - i) a reactive moiety coupled to the monolithic matrix, and
 - ii) a ligand that binds to the polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety;
 - b) allowing for the polynucleotide to bind to the ligand; and
 - c) eluting the polynucleotide from the monolith matrix after one or more contaminants have been substantially separated from the bound polynucleotide.
2. The method of Claim 1, wherein the reactive moiety is a primary amine.
3. The method of Claim 1, wherein the monolithic matrix is activated with an activating agent selected from carbonyldiimidazole, epoxy, ethylenediamine, carbodiimide, aldehyde, anhydride, imidoester and NHS ester.
4. The method of Claim 1, wherein the ligand further comprises a carbon-containing linker positioned between the reactive moiety and the ligand.
5. The method of Claim 4, wherein the carbon linker is C_X, where X is a whole number between about 1 and about 50.
6. The method of Claim 1, wherein the polynucleotide is mRNA.
7. The method of Claim 6, wherein the ligand is a poly-A binding probe.
8. The method of Claim 7, wherein the ligand is NH₂-C_X-dT_Y, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30.
9. The method of Claim 8, wherein the ligand is NH₂-C₁₂-dT₁₈.
10. The method of Claim 1, further comprising washing the column prior to eluting the polynucleotide.
11. The method of Claim 10, wherein the wash buffer contains a salt concentration of at least 200 mM, and the elution buffer contains a salt concentration of 100 mM or less.
12. The method of Claim 10, wherein the wash buffer comprises one or more salts

- selected from sodium sulfate, sodium chloride and sodium phosphate.
13. The method of Claim 11, wherein the elution buffer is selected from water and Tris.
 14. The method of Claim 11, wherein the flow rate of the column is at least about 0.5 CV/min.
 15. The method of Claim 15, further comprising eluting the free polynucleotide from the monolith matrix by reducing the ionic strength of the liquid phase.
 16. A method of separating a formulated polynucleotide from free polynucleotide, the method comprising:
 - a) loading a sample onto a monolith matrix comprising a ligand comprising:
 - i) a reactive moiety coupled to the monolithic matrix, and
 - ii) an affinity moiety that binds to the free polynucleotide but not the formulated polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety; and
 - b) collecting the formulated polynucleotide from the column while the free polynucleotide remains immobilized on the monolith matrix.
 17. The method of Claim 16, wherein the monolith matrix is contained in a column.
 18. The method of Claim 16, wherein the formulated polynucleotide is a formulated mRNA.
 19. The method of Claim 18, wherein the mRNA is formulated in a lipid nanoparticle.
 20. The method of Claim 18, wherein the ligand is a poly-A binding probe.
 21. The method of Claim 20, wherein the ligand is $\text{NH}_2\text{-C}_X\text{-dT}_Y$, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30.
 22. The method of Claim 21, wherein the ligand is $\text{NH}_2\text{-C}_{12}\text{-dT}_{18}$.
 23. The method of Claim 16, wherein the ligand further comprises a carbon-containing linker positioned between the reactive moiety and the ligand.
 24. The method of Claim 23, wherein the carbon linker is C_X , where X is a whole number between about 1 and about 50.
 25. The method of Claim 16, further comprising eluting the free polynucleotide from the monolith matrix by reducing the ionic strength of the liquid phase.

26. A column for purifying a polynucleotide from a sample, said column comprising:
 - a) a monolithic matrix; and
 - b) a ligand comprising a reactive moiety coupled to the monolithic matrix, and a ligand that binds to the polynucleotide, wherein the ligand is immobilized to the monolithic matrix via the reactive moiety.
27. The column of Claim 26, wherein the reactive moiety is a primary amine.
28. The column of Claim 26, wherein the monolithic matrix is activated with an activating agent selected from carbonyldiimidazole, epoxy, ethylenediamine, carbodiimide, aldehyde, anhydride, and imidoester and NHS ester.
29. The column of Claim 26, wherein the ligand is a poly-A binding probe.
30. The column of Claim 29, wherein the ligand further comprises a carbon linker positioned between the reactive moiety and the oligo-dT probe.
31. The column of Claim 30, wherein the carbon linker is C_X, where X is a whole number between 1 and 50.
32. The column of Claim 29, wherein the ligand is NH₂-C_X-dT_Y, where X is a whole number between 1 and 50 and Y is a whole number between 5 and 30.
33. The column of Claim 32, wherein the ligand is NH₂-C₁₂-dT₁₈.
34. A method of preparing the column of Claim 26, comprising:
 - a) treating the monolithic matrix with an activating agent to produce an activated monolithic matrix; and
 - b) incubating the activated monolithic matrix in the presence of a ligand comprising a reactive moiety.
35. The method of Claim 34, wherein the activating agent is selected from carbonyldiimidazole, epoxy, ethylenediamine, carbodiimide, aldehyde, anhydride, imidoester and NHS ester.

1/16




FIG. 1

2/16




FIG. 2




FIG. 3

4/16




FIG. 4




FIG. 5




FIG. 6

7/16




FIG. 7




FIG. 8

9/16




FIG. 9

10/16




FIG. 10

11/16




FIG. 11

12/16




FIG. 12

13/16




FIG. 13

14/16





FIG. 14

15/16




FIG. 15




FIG. 16

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/034193

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12Q1/68
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HANORA A ET AL: "Direct capture of plasmid DNA from non-clarified bacterial lysate using polycation-grafted monoliths", JOURNAL OF BIOTECHNOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 123, no. 3, 29 May 2006 (2006-05-29), pages 343-355, XP024956794, ISSN: 0168-1656, DOI: 10.1016/J.JBIOTEC.2005.11.017 [retrieved on 2006-05-29]	1,10-15, 26,34
A	abstract; figures 2-4; table 2 page 349, column 2, paragraph 2 - page 353 ----- -/--	2-9, 16-25, 27-33,35



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 July 2017

Date of mailing of the international search report

24/07/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Tilkorn, A

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/034193

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TETALA KISHORE K R ET AL: "A review on recent developments for biomolecule separation at analytical scale using microfluidic devices", ANALYTICA CHIMICA ACTA, vol. 906, 17 December 2015 (2015-12-17), pages 7-21, XP029380855, ISSN: 0003-2670, DOI: 10.1016/J.ACA.2015.11.037	1-3, 10-15, 26-28, 34,35
A	abstract; figure 4 page 11 - page 12, column 1, paragraph 1 page 15, column 1 page 17, column 1, paragraph 5 - page 18 -----	4-9, 16-25, 29-33
X	BRIAN E. ROOT ET AL: "Purification of HIV RNA from Serum Using a Polymer Capture Matrix in a Microfluidic Device", ANALYTICAL CHEMISTRY, vol. 83, no. 3, 1 February 2011 (2011-02-01), pages 982-988, XP055390428, US ISSN: 0003-2700, DOI: 10.1021/ac102736g abstract; figures 4,5 page 983, column 2 - page 984, column 2, paragraph 2 -----	1-15, 26-35
X	HAN Y ET AL: "Single step purification of plasmid DNA using peptide ligand affinity chromatography", JOURNAL OF CHROMATOGRAPHY B: BIOMEDICAL SCIENCES & APPLICATIONS, ELSEVIER, AMSTERDAM, NL, vol. 874, no. 1-2, 15 October 2008 (2008-10-15), pages 21-26, XP025506829, ISSN: 1570-0232, DOI: 10.1016/J.JCHROMB.2008.08.025 [retrieved on 2008-09-07]	1-5, 10-15, 26-28, 34,35
A	abstract; figures 2-7 page 23, column 1, paragraph 3 - column 2, paragraph 1 page 24 - page 25 -----	6-9, 16-25, 29-33
A	WO 2016/004318 A1 (SHIRE HUMAN GENETIC THERAPIES [US]) 7 January 2016 (2016-01-07) abstract paragraph [0100] - paragraph [0101] -----	1-35
A	WO 2010/068856 A1 (UNIV BOSTON [US]; FRAUNHOFER USA INC [US]; SHARON ANDRE [US]; CHATTERJ) 17 June 2010 (2010-06-17) ----- -/--	1-35

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/034193

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PODGORNIK ALES ET AL: "Fast separation of large biomolecules using short monolithic columns", JOURNAL OF CHROMATOGRAPHY B: BIOMEDICAL SCIENCES & APPLICATIONS, ELSEVIER, AMSTERDAM, NL, vol. 927, 11 February 2013 (2013-02-11), pages 80-89, XP028589564, ISSN: 1570-0232, DOI: 10.1016/J.JCHROMB.2013.02.004 page 87, column 2 - page 88; figure 17 -----	1-35
A	WO 2013/180654 A1 (AGENCY SCIENCE TECH & RES [SG]) 5 December 2013 (2013-12-05) the whole document -----	1-35

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/034193

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2016004318 A1	07-01-2016	AU 2015283954 A1 CN 106456547 A EP 3164112 A1 US 2016038432 A1 WO 2016004318 A1	22-12-2016 22-02-2017 10-05-2017 11-02-2016 07-01-2016
WO 2010068856 A1	17-06-2010	US 2011313145 A1 WO 2010068856 A1	22-12-2011 17-06-2010
WO 2013180654 A1	05-12-2013	CN 104470606 A EP 2854982 A1 JP 2015527052 A KR 20150023331 A SG 11201407808X A US 2015148528 A1 WO 2013180654 A1	25-03-2015 08-04-2015 17-09-2015 05-03-2015 30-12-2014 28-05-2015 05-12-2013