



HU000028581T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 028 581**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 11 715928**(22) A bejelentés napja: **2011. 04. 21.**

(96) Az európai bejelentés bejelentési száma:

EP 20110715928

(97) Az európai bejelentés közzétételi adatai:

EP 2560691 A2 **2011. 10. 27.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2560691 B1 **2015. 11. 04.**(51) Int. Cl.: **A61L 2/10**

(2006.01)

A61K 9/08

(2006.01)

A61P 31/00

(2006.01)

C07K 16/06

(2006.01)

A61P 37/00

(2006.01)

A61K 393/95

(2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/EP 11/056486

(87) A nemzetközi közzétételi szám:

WO 11131786

(30) Elsőbbségi adatok:

201006753**2010. 04. 22.****GB**

(73) Jogosult(ak):

Biotest AG, 63303 Dreieich (DE)

(72) Feltaláló(k):

MÖLLER, Wolfgang, 61440 Oberursel (DE)**RUDNICK, Dieter, 63303 Dreieich (DE)****MANEG, Oliver, 61352 Bad Homburg (DE)****RODEMÉR, Michael, 63110 Rodgau (DE)****DICHTELMUELLER, Herbert, 65843 Sulzbach (DE)****FLECHSIG, Eckhard, 63128 Dietzenbach (DE)**

(74) Képviselő:

Danubia Szabadalmi és Jogi Iroda Kft.,**Budapest**

(54)

Eljárás immunglobulin-készítmény előállítására

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 560 691 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
04.11.2015 Bulletin 2015/45

(51) Int Cl.:
A61L 2/10 ^(2006.01) **C07K 16/06** ^(2006.01)
A61K 9/08 ^(2006.01) **A61K 39/395** ^(2006.01)
A61P 31/00 ^(2006.01) **A61P 37/00** ^(2006.01)

(21) Application number: **11715928.5**

(86) International application number:
PCT/EP2011/056486

(22) Date of filing: **21.04.2011**

(87) International publication number:
WO 2011/131786 (27.10.2011 Gazette 2011/43)

(54) **PROCESS FOR PREPARING AN IMMUNOGLOBULIN COMPOSITION**

VERFAHREN ZUR HERSTELLUNG EINES IMMUNOGLOBULIN-PRÄPARATES

PROCÉDÉ POUR PRÉPARER UNE COMPOSITION D'IMMUNOGLOBULINE

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

- **DICHTELMUELLER, Herbert**
65843 Sulzbach (DE)
- **FLECHSIG, Eckhard**
63128 Dietzenbach (DE)

(30) Priority: **22.04.2010 GB 201006753**

(74) Representative: **Daniels, Jeffrey Nicholas**
Page White & Farrer
Bedford House
John Street
London WC1N 2BF (GB)

(43) Date of publication of application:
27.02.2013 Bulletin 2013/09

(73) Proprietor: **Biotest AG**
63303 Dreieich (DE)

(56) References cited:
EP-A1- 0 013 901 EP-A1- 0 413 187
EP-A1- 0 450 412 EP-A1- 0 835 880
EP-A2- 0 352 500 EP-A2- 0 413 188
US-A- 4 318 902 US-A1- 2003 152 966

(72) Inventors:

- **MÖLLER, Wolfgang**
61440 Oberursel (DE)
- **RUDNICK, Dieter**
63303 Dreieich (DE)
- **MANEG, Oliver**
61352 Bad Homburg (DE)
- **RODEMER, Michael**
63110 Rodgau (DE)

- **BURNOUF ET AL: "Modern Plasma
Fractionation", TRANSFUSION MEDICINE
REVIEWS, GRUNE AND STRATTON, ORLANDO,
FL, US, vol. 21, no. 2, 28 March 2007 (2007-03-28) ,
pages 101-117, XP022020853, ISSN: 0887-7963,
DOI: 10.1016/J.TMRV.2006.11.001**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 560 691 B1

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to a process for the preparation of a virus safe immunoglobulin composition, and antibody preparations and pharmaceutical compositions which can be prepared using the process.

BACKGROUND OF THE INVENTION

10 **[0002]** Immunoglobulin compositions prepared from human plasma and suitable for intravenous administration are known in the art and for several decades have played an important role in the treatment of a wide range of diseases. Immunoglobulins are used, for example, for the treatment of infections in humans and can be assigned to various classes with various biochemical and physiological properties. Immunoglobulin G participates in defending against viral antigens, whereas IgM is predominantly active in antibacterial and antitoxin immune responses.

15 **[0003]** The immunoglobulin solutions comprise IgG, IgA and IgM in various percentages, with different preparations having different treatment applications, e.g. preparations with a higher percentage of IgM are used in the prophylaxis or treatment of bacterial infections. In contrast to IgG preparations IgM antibodies easily aggregate in solution. IgM preparations are difficult to stabilize especially if they are enriched compared to plasma concentrations and stored in liquid solution.

20 **[0004]** The immunoglobulin solutions are usually prepared from fractions of blood plasma or serum, e.g. Cohn fractions. These fractions are then subjected to a number of purification steps to remove contaminants including viruses, denatured proteins, proteases and lipids. Human plasma for fractionation is collected from thousands of donors and may contain pathogen viruses despite thorough testing of the source plasma. Therefore process steps to inactivate or remove viruses are essential in order to achieve safe products for use in medicine. Several techniques for virus inactivation/removal are
25 known in the art, e.g. chemical treatments, irradiation with UVC light or nanometer filtration, which are performed in order to ensure overall virus safety. However, such steps can have a negative impact on the activity of the immunoglobulins; for example extended irradiation times with UVC can reduce the yield of native and active IgM obtained in the final immunoglobulin solution.

30 **[0005]** The virus removal or inactivation capacity of the process steps is validated using laboratory scale models of the production process and for each step a removal or inactivation factor is determined. An increase of the inactivation / removal factor adds additional viral safety to the pharmaceutical product. Today guidelines from regulatory authorities require at least two effective steps for enveloped and non-enveloped viruses in the manufacture of plasma-derived pharmaceuticals.

35 **[0006]** In addition to viruses which are potentially present it is also necessary to remove other contaminants like proteases, protein aggregates, and denatured immunoglobulins, to achieve a well tolerated product. Denatured immunoglobulins especially are a potential risk for the patients because they have a high capacity to activate complement unspecifically, leading to severe side effects in patients receiving these denatured immunoglobulins. This anticomplementary activity (ACA) is measured by a standardized test described in the European Pharmacopoeia.

40 **[0007]** The removal of all these contaminants is essential (1) in order for the product to be tolerated by the patient after intravenous administration, (2) to ensure the product complies with bio-safety guidelines regarding viral contamination, (3) to allow the product to be stable during long-term storage, and (4) to generate the desired compound mixture / pharmaceutical composition.

45 **[0008]** The initial purification of human IgM solutions has been carried out by classical Cohn plasma fractionation methods or its well known modifications (e.g. Cohn/Oncley, Kistler/Nitschmann). Using cold ethanol precipitation processes the IgM fraction is recovered in fraction III or fraction I/III (also called B or B+I). Starting from fraction III or I/III methods have been described for purification of protein solutions enriched in IgM. EP0013901 describes a purification method starting from fraction III including steps using octanoic acid, β -Propiolactone and an adsorption step using an anionic exchange resin. This method is used to produce Pentaglobin® - to date the only commercially available intravenous IgM product. EP0352500 describes the preparation of an IgM concentrate for intravenous application with a reduced
50 anti-complementary activity by using anionic exchange chromatography, β -Propiolactone, UVC light irradiation and an incubation step at increased temperature (40 °C to 60 °C). β -propiolactone is a well known chemical used in sterilization steps in order to inactivate viruses which are potentially present. As β -propiolactone is a very reactive substance which causes the chemical modification of proteins there is also substantial loss of the anti-viral and anti-bacterial activities of the immunoglobulins. The preparation produced by this method was stable in liquid solution for a limited time due to the
55 chemical modification. The IgM concentration was above 50 % from the total immunoglobulin content.

[0009] The preparation of protein solutions enriched in IgM without chemical modification by β -propiolactone has been described in EP0413187 (Biotest) and EP0413188 (Biotest). These methods involve subjecting a suitable protein solution to octanoic acid treatment and anionic exchange chromatography, starting from Cohn fraction III or I/III. In patent

EP0413187 (Biotest) the octanoic acid treatment is carried out by stirring for 15 min, in order to remove lipids being present in Cohn fraction III.

[0010] As large amounts of immunoglobulins are administered intravenously to patients a tolerable pharmaceutical preparation must be achieved. IgM preparations have been described as being difficult to prepare for intravenous application. IgM by nature is a vigorous activator of complement after the binding of antigens. Therefore unspecific anticomplementary activity of denatured IgM molecules is far more dangerous to patients than denatured IgG molecules. The preparation according to EP0413187 had a low anticomplementary activity, between 0,6 and 0,8 CH50/mg protein, but had to be stabilized and virus inactivated by β -propiolactone. Low anticomplementary activity is considered to be ≤ 1 CH50/mg protein according to EP monograph for immunoglobulins.

[0011] EP0413188B1 (Biotest) describes the preparation of an IgM-enriched preparation for intravenous administration by using an anion exchange chromatography in order to reduce the anti-complementary activity. Additionally a heat treatment at pH 4 - 4.5 at 40 to 60°C, preferably between 50 and 54°C, was described to reduce the anticomplementary activity. This preparation had to be lyophilized to ensure stability of the preparation for several months. Long term stability as a liquid solution could not be shown.

[0012] Another method describes the use of mild-heat treatment of IgM preparations at 40 to 62°C, preferably 45 to 55°C, at pH 4.0 to 5.0 (EP 0450412, Miles) to reduce the non-specific complement activation. In this patent application octanoic acid is added to a Cohn fraction III suspension in order to remove prekallikrein activator and lipoproteins by centrifugation. Nevertheless this treatment led to partial loss of antigenic determinants of IgM. This may increase the risk of generating neo-antigens leading to a increased immunogenicity in humans or the loss of activity.

[0013] The preparation of an IgM containing protein solution for intravenous application by using a protease treatment (e.g. with pepsin) after an octanoic acid precipitation step has been described in EP0835880 (US 6136312, ZLB). Protease treatment leads to partial fragmentation of the immunoglobulin molecule impairing the full functional activity of the Fab and Fc parts. Therefore protease-treated immunoglobulins cannot be regarded as unmodified. Also this preparation method leads to about 5 % fragments with a molecular weight of <100kD.

[0014] The described methods to carry out the octanoic acid treatment (EP0413187 and EP0835880) have the drawback that the octanoic acid treatment is not effective with respect to removal and inactivation of non-enveloped viruses, and does not remove substantially all proteolytic activity.

[0015] In EP 0345543 (Bayer, Miles) a highly concentrated IgM preparation with at least 33% IgM for therapeutic use is disclosed, the preparation being substantially free of isoagglutinin titres. In this patent application an octanoic acid precipitation is carried out by adding the octanoic acid and the isoagglutinins are removed by Synsorb affinity chromatography. The final preparation had to be freeze dried.

[0016] Altogether the preparation of a IgM containing preparation with low anticomplementary activity is possible if the immunoglobulins are chemically or enzymatically modified and/or further purified by chromatography and/or subjected to a mild heat treatment.

[0017] Nevertheless the methods of the prior art leading to an unmodified immunoglobulin preparation are not able to achieve the virus inactivation capacity for all viruses which are potentially present. Although several methods, such as solvent/detergent treatment, octanoic acid treatment, nanometer filtration and heat treatment, are effective to inactivate or remove enveloped viruses there are only a few methods known to inactivate or remove non-enveloped viruses, for example Parvo viruses. These non-enveloped viruses are mostly very small, usually passing through nanometer filters with pore sizes above 20 nm. This pore size is too small for IgM molecules having a diameter up to 30 nm. Non enveloped viruses are effectively inactivated by chemicals like β -propiolactone which, however, also leads to a modified immunoglobulin with impaired functions. Another effective treatment is UVC-irradiation (EP1842561, CAF-DCF). However, known solvent/detergent treatments, octanoic acid treatment and mild heat treatment have no substantial effect on non-enveloped viruses.

[0018] Therefore all the chemically unmodified IgM containing preparations, which are prepared by the methods of the prior art, and which have low anticomplementary activity, are not safe for human use with respect to non enveloped viruses e.g. Parvoviruses.

[0019] In summary the methods of the prior art which isolate an intravenous-tolerable IgM-containing preparations have certain drawbacks, such as an inability to effectively inactivate or remove non-enveloped viruses and the limited ability to remove proteolytic activity while keeping IgM with high yield in solution. (Proteolytic activity refers to the sum of proteases being present in the preparation). As a liquid protein preparation must be storable for long periods (e.g. 2 years), residual protease activities must be omitted, as these activities might lead to degradation of the pharmaceutical preparation.

[0020] Thus, it is the aim of the present invention to address these drawbacks.

SUMMARY OF THE INVENTION

[0021] In a first aspect the present invention provides a process for the preparation of an IgM immunoglobulin com-

position from a plasma fraction comprising immunoglobulins, the process comprising:

- (a) providing a plasma fraction as a solution containing the immunoglobulins;
- (b) mixing a C₇ to C₉ carboxylic acid with the solution and treating the mixed solution with a vibrating agitator to precipitate contaminating proteins ; and
- (c) separating the precipitated proteins from the solution to yield the IgM containing immunoglobulin composition.

[0022] The present applicants have surprisingly found that the use of a vibrating agitator during the step where the immunoglobulin solution is mixed with the carboxylic acid is extremely advantageous. This method step provides a more efficient removal of unwanted proteins (including proteases) and produces an intermediate product which is better suited to downstream processing steps utilised to produce an immunoglobulin medicament; the intermediate product allows these downstream processing steps to be more efficient. In particular, the IgM immunoglobulin containing composition obtained from step (c) can be combined with further treatment steps, such as treatment with mild acid conditions and treatment with UVC irradiation, to produce an IgM containing immunoglobulin product or antibody preparation which is suitable for intravenous administration and which has the following advantageous properties: (i) being chemically unmodified; (ii) being virus safe (iii) having low proteolytic activity (and therefore being stable during long term storage); (iv) having low anti-complementary activity; and (v) retaining a high level of native and active IgM. The level of virus safety achieved with the methods described herein has not previously been obtainable. Further, the use of the method of the present invention achieves an IgM containing immunoglobulin product or antibody preparation with combinations of these features which have not previously been obtainable.

[0023] In particular, the method steps of the present invention lead to a higher inactivation and removal of virus particles, especially very resistant, non-enveloped viruses such as Parvo viruses, which are usually not very susceptible to octanoic acid treatment. Furthermore, an improved removal of proteolytic activity is achieved in comparison to conventional stirring. These features are achieved while keeping a high yield of IgM that is chemically unmodified. This finding contrasts with the conventional view that the treatment with octanoic acid is not an effective step against non-enveloped viruses and improved viral safety must be achieved through inactivation of virus through harsher methods such as β -Propiolactone treatment. Also it was well known that increasing e.g. octanoic acid concentration to completely remove proteolytic activity results in a massive loss of IgM.

[0024] The results of the present invention are achieved through the use of mixing devices using a vibrating mode in combination with the octanoic acid treatment. This is particularly surprising since it is known that IgM is very susceptible to shear stress, which may lead to an undesired high anticomplementary activity. Accordingly, one would not consider using a vibrating mixer to prepare an IgM composition and would not expect such a favorable impact when using a vibrating mixing during processing of an IgM containing solution.

[0025] Furthermore with the method of the present invention the separation achieved by step (c), such as clarification by filtration of the octanoic acid treated solution resulting from step (b), is enhanced when a vibrating mixing device is used. Separation is more easily achieved, reducing processing time and manufacturing costs, and step (c) leads to a limpid solution which creates advantages for downstream processing. Conventional solutions, achieved by filtering the results of octanoic acid treated IgM containing solutions which have been stirred, are opalescent or opaque.

[0026] The resulting IgM containing composition obtained from step (c) is preferably subjected to treatment with mild acid conditions (e.g. pH 4) and an UVC-irradiation step to further improve virus safety and stabilize the final product. Due to the enhanced clarification of the IgM containing immunoglobulin composition obtained from step (c) it is possible to lower the necessary irradiation time with UVC to achieve a virus inactivation of non enveloped viruses of more than 3 or 4 log₁₀. This results in a higher yield of native and active IgM during UVC treatment.

[0027] Surprisingly these steps lead to a chemically and enzymatically unmodified IgM containing solution which has higher yields of native and active IgM, having low anticomplementary activity and low proteolytic activity and having high antibacterial and antiviral activity, with an outstanding virus safety concerning enveloped and non enveloped viruses; a key feature for pharmaceuticals which are for intravenous administration. Moreover, a treated IgM containing solution has improved long term stability being very stable in liquid solution for more than 12 months at 2 - 8°C.

[0028] Accordingly, in a further aspect the present invention provide an antibody preparation obtainable using the process of the present invention set out above. The antibody preparation is suitable for intravenous administration in humans and comprises IgG, IgA and IgM, wherein at least 5% of the total immunoglobulins are IgM. In an aspect the present invention provides an antibody preparation comprising immunoglobulins IgG, IgA and IgM, wherein at least 15% of the total immunoglobulins are IgM, wherein the antibody preparation is virus safe with respect to enveloped and non-enveloped virus, has an anticomplementary activity of ≤ 1 CH 50/mg protein, and is stable in liquid form for at least 6 months when stored at 2 to 8°C. The preparation preferably has a proteolytic activity of less than 8U/l.

[0029] Still further, the present invention provides an antibody preparation of the present invention for use in medicine. In one embodiment the antibody preparation is for use in the treatment of immunological disorders and bacterial infections.

[0030] A further aspect of the present invention is the use of an antibody preparation of the present invention for the

manufacture of a medicament for treatment of immunological disorders and bacterial infections.

[0031] The antibody preparation of the present invention is suitable for use with a method of treatment comprising administering the antibody preparation of the present invention to a patient.

[0032] The present invention will now be described in further detail by way of example only, with reference to the accompanying figures.

[0033] FIGURE 1 provides an overview of an embodiment of the invention in which the steps that can be utilised to form an antibody preparation suitable for intravenous administration are shown. The octanoic acid treatment step employing a vibromixer device, the pH4 treatment and the UVC treatment are highlighted. The starting material is generated from a standard cold ethanol precipitation process of human plasma.

DETAILED DESCRIPTION OF THE INVENTION

[0034] As described above, the present invention provides a process for the preparation of an IgM containing immunoglobulin composition from a plasma fraction comprising immunoglobulins. Plasma fractions suitable for the preparation of pharmaceutical immunoglobulin compositions, and methods for their production are well known in the art. In particular, where the pharmaceutical immunoglobulin compositions are for administration to humans, the plasma fraction is obtained from human plasma. The plasma fraction is preferably a precipitated plasma fraction and most preferably a precipitated plasma fraction obtained by the process of Cohn fractionation or its well known modifications (e.g. Kistler-Nitschmann). Most preferably the fraction is fraction I/III or fraction III (also known as fraction B+I or fraction B) out of cold ethanol fractionation. It is preferred that the immunoglobulins of the plasma fraction comprise at least 5% IgM.

[0035] Step (a) comprises providing a plasma fraction as a solution containing the immunoglobulins. In many cases the plasma fraction containing the immunoglobulins will be in solid or semi-solid form. Thus the aim of this step is to ensure or to bring the protein of the plasma fraction into solution such that it is in a suitable state for mixing with the carboxylic acid in step (b). This step may comprise mixing the plasma fraction with a suitable buffer. Preferably the buffer is of low molarity (i.e. less than 1M) and has a pH between 4.5 and 5.5 e.g. 0.1 M sodium acetate buffer pH 5.05 ± 0.1 . Mixing can be completed using a blade mixer or a vibrating agitator.

[0036] In step (b) the solution formed in step (a) is mixed using a vibrating agitator with a C₇ to C₉ carboxylic acid to precipitate contaminating proteins (e.g. proteases, viruses etc). The carboxylic acid may be branched and/or may include substituents which do not substantially alter the effect of step (b). The carboxylic acid is preferably octanoic acid. The acid is preferably added at a concentration of at least 0.075 kg/kg of plasma fraction, up to a concentration of 0.2 kg/kg. More preferably the acid is added at 0.8 to 0.15 kg/kg of plasma fraction, and most preferably between 0.09 kg/kg and 0.13 kg/kg. Acid of any convenient molarity may be used to provide the correct concentration.

[0037] Any type of commercially available vibrating agitator, suitable for use in the chemical/pharmaceutical industry, may be used. Examples of suitable vibrating agitators are available from Graber + Pfenninger GmbH. In particular, the "Labormodell Typ 1" vibromixer can be used for lab scale experiments, and the "Industriemixer Typ 4" can be used for production scale preparations. The vibrating mixers can be used according to manufacturer's instructions, and in particular at settings which are described by the manufacturers as suitable for mixing solutions containing proteins. For example the vibrating mixers can usually be operated at less than 100 Hz with an amplitude less than 10 mm, e.g. the vibration mixing using the "Labormodell Typ 1" at lab scale was carried out by the present inventors at 50 Hz, when 230 V power supply is used. The vibration amplitude of the mixing process was varied between 0 and 3 mm, and for the IgM preparation preferably 3 mm was used. Stirrer plates with a diameter between 23 mm and 65 mm were used for lab scale experiments. For production scale a stirrer plate diameter of 395 mm was used (hole diameters of 13.5 mm and 16 mm).

[0038] In step (b) the pH of the mixed solution is preferably between 4.5 to 5.5, and more preferably between pH 4.8 and pH 5.3. The step can be carried out in sodium acetate buffer, and, for example, with approximately 0.1 M sodium acetate buffer. The temperature at which step (b) is conducted is preferably between 10°C and 35°C, and more preferably 14 to 30 °C.

[0039] The mixing time using the vibrating agitator is not particularly limited but is preferably at least 30 minutes and not more than 3 hours, and more preferably from 40 - 110 minutes. Incubation times of less than 30 minutes can reduce the level of virus inactivation.

[0040] In one embodiment of step (b) tri-calcium phosphate is mixed with the solution in step (b). Preferably this is added at 0.01 to 0.02 kg/kg plasma fraction (as it is in solid or semi-solid form). The tri-calcium phosphate can be added simultaneously, separately or sequentially to the carboxylic acid. In a preferred embodiment the tri-calcium phosphate is added at least 20 minutes after the carboxylic acid.

[0041] In step (c) the contaminating proteins precipitated in step (b) are separated off from the solution to yield the IgM containing immunoglobulin composition (i.e. an immunoglobulin containing solution). This step of separation is not particularly limited but can be performed by any suitable method known in the art. However, the separating step is preferably performed using filtration, and more preferably ultrafiltration, and the result of step (c) is therefore a filtered solution.

[0042] As described above, the method of the present invention is advantageous in manufacturing terms since it appears to cause a more efficient precipitation of contaminating proteins, and, as a result, step (c) is easier to perform. When the mixture resulting from step (b) is separated, a transparently clear solution, i.e. the IgM containing immunoglobulin composition, is achieved. Filtration is therefore quicker and easier.

[0043] Further process steps are required to convert the IgM containing immunoglobulin composition obtained from step (c) into an immunoglobulin preparation suitable for intravenous administration. Accordingly, in a preferred embodiment the process of the present invention comprises the additional steps of treating the IgM containing immunoglobulin composition obtained from step (c) with mild acid conditions and, further, subjecting the acid treated composition to UVC irradiation.

[0044] For the treatment with mild acid conditions the IgM containing immunoglobulin composition obtained from step (c) is incubated at between pH 3.5 to pH 4.5, and preferably between pH3.8 and pH4.2, to form an incubated solution. The mild acid conditions can be created by adding a suitable acid to the IgM containing immunoglobulin composition, for example the pH can be adjusted by adding 0.2M HCl.

[0045] This incubation step is preferably carried out at between 32 and 42°C, and more preferably at between 35 and 39°C. The incubation time is preferably at least 2 hours and not greater than 24 hours, and more preferably at least 9 hours but not greater than 16 hours.

[0046] In the irradiation step the incubated solution obtained from the mild acid treatment described above is treated with UVC light to form a UVC irradiated solution. This step can be performed using devices which are commercially available, such as the UVivitec® device (Bayer Technology Services). It is preferred that the incubated solution is treated at 254 ± 10 nm between 200 and 500 J/m², more particularly between 200 and 300 J/m², in order to further inactivate viruses and proteases which are potentially present. It is noted that UVC treatment under gentle conditions than would normally be required is only possible with the water-clear filtrate which is obtained by the present invention after the octanoic acid treatment with vibromixing. More opalescent or opaque solutions normally received by standard stirring techniques would necessitate longer irradiation times which would lead to more denaturation of the IgM activity and lower virus inactivation rates.

[0047] In addition to the mild acid treatment and the UVC irradiation the additional steps to achieve an immunoglobulin preparation for intravenous administration can optionally also comprise one or more filtration steps. In one embodiment the protein solution being processed can be adsorbed onto DEAE-Sephadex and then separated from the Sephadex by depth filtration. For example, it may further be subjected to a batch adsorption with 75 mg per kg protein DEAE Sephadex at pH 5.8, in order to remove the unwanted accompanying protein Ceruloplasmin.

[0048] In a particularly preferred embodiment the incubated solution obtained from the mild acid treatment is subjected to adsorption onto DEAE-Sephadex and then separated from the Sephadex by depth filtration, before being treated to UVC irradiation.

[0049] In another embodiment the immunoglobulin solution being processed may be filtered through a nanometer filter. Filters of 75 ± 5 nm to 35 ± 5 nm pore size, or filters having a nominal pore size of 75 to 35 nm (for example Pall Ultipor DV50), can be used at various stages during the process. (A nominal pore size of e.g. 50 nm means retention rate of $\geq 4 \log_{10}$ for virus with size of 50 nm or larger). In a preferred embodiment the solution obtained from the DEAE-Sephadex step described in the above paragraph is filtered through a 0.2µm filter prior to UVC irradiation. In a further preferred embodiment the immunoglobulin solution obtained after UVC irradiation is subjected to nanofiltration, preferably through a filter having a 40 to 50 nm pore size. It is preferred that this step should be carried out under sterile conditions.

[0050] The final antibody preparation (i.e. the processed IgM containing immunoglobulin solution) obtained from the process defined above may be directly filled into a container under sterile conditions. Alternatively, the antibody preparation may be formulated with a stabilizing agent, such as glycine. In particular, the preparation may be formulated in a glycine-containing buffer at a pH between 4 and 5.5, and preferably between 4.1 to 4.5. The antibody preparation may also be diluted to a protein concentration between 40 and 80 g/L and preferably between 55 and 70 g/L.

[0051] Preferably the process of the present invention does not comprise a step involving one or more of chemical modification of the antibodies in the preparation, enzymatic modification of the antibodies, or heat treatment of the antibodies (e.g. treatment of the antibodies at a temperature of 40°C or more for 10 minutes or more). More particularly the process of the present invention does not include a step of contacting the antibodies with β-propiolactone and/or pepsin.

[0052] The present invention further provides an immunoglobulin preparation or antibody preparation obtainable by the process described above. The immunoglobulin preparation is polyclonal and can comprise at least 5% IgM, preferably at least 15% IgM and most preferably at least 20% IgM. The other immunoglobulins are IgG and IgA. Preferably the immunoglobulin preparation comprises between 5 and 30% IgM (most preferably between about 15 and 30%), between 5 and 30% IgA (most preferably between 15 and 30%) and between 40 and 70% IgG (most preferably between 45 and 70%). In the most preferred product the IgG content is about 50 % and the IgM and IgA content each are about 25 %. The values mean the percentage of e.g. IgM from the sum of IgG+IgA+IgM. Nevertheless it is also possible to enrich the IgM further by well known methods like e.g. anion exchange chromatography. The values can be determined by

nephelometry or by immunoprecipitation according Ph. Eur. (current edition (2010) 2.9.17).

[0053] In particular, the immunoglobulins in the immunoglobulin preparation are not chemically modified. The immunoglobulin preparation has not been treated during its process of production with a chemical such as β -propiolactone to sterilize the product. Similarly the preparation has not been treated with an added protease, such as pepsin, to sterilize the product. As such, the immunoglobulins are substantially in native form.

[0054] The antibody preparation has a lower proteolytic activity than the antibody preparations described in the prior art. In particular, no proteolytic activity is detectable in the preparation when it is stored at between 2 to 8°C. The proteolytic activity can be measured by standardized test methods known in the art, such as that using the chromogenic substrate which is described below in example 6. In such methods proteolytic activity is assessed by mixing a chromogenic substrate (in particular those sensitive to at least one serine protease) and a sample of the antibody preparation (usually diluted in buffer to meet the linear range of the assay) at 37°C and monitoring the absorption kinetics using a spectrophotometer. The proteolytic activity of the sample is calculated from the initial absorption difference ($\Delta\text{Abs}/\text{min}$) by using the equation $C \text{ (U/L)} = S \times \Delta\text{Abs}/\text{min} \times F$ (C = proteolytic activity; S = conversion factor relating to specific adsorption change of the chromogenic substrate; and F = dilution factor). Use of the substrate is according to manufacturer's instructions.

[0055] The proteolytic activity can in particular be assessed via the following steps:

- (a) 25 mg of the substrate S-2288 (Chromogenix) is dissolved in 7.2 ml of water-for-injection;
- (b) a sample of the antibody preparation is diluted into buffer (100mM Tris.HCl pH 8.4, 106 mM NaCl) to meet the linear range of the assay and temperature is adjusted to 37°C;
- (c) equal amounts (e.g. 200 μ l) of the diluted antibody preparation and the dissolved substrate are mixed;
- (d) the absorption kinetics are measured at 405 nm for 1 to 3 minutes at 37°C using a spectrophotometer;
- (e) the proteolytic activity of the sample is calculated from the initial absorption difference ($\Delta\text{Abs}/\text{min}$) by using the equation $C \text{ (U/L)} = 313 \times \Delta\text{Abs}/\text{min} \times F$ (C = proteolytic activity, F = dilution factor)

[0056] The limit of quantitation of this method is 8 U/l, and using a sample of the antibody preparation of the present invention proteolytic activity is undetectable. As such the level of the proteolytic activity in the final product of the present invention is below 8 U/l.

[0057] As with preparations known in the art the antibody preparation of the present invention can be stored at $5 \pm 3^\circ\text{C}$. However, due to the efficient purification with the method of the present invention the stability of the antibody preparation is extremely good. The final product is stable in liquid form for at least 3 months, preferably at least 6 months and most preferably at least two years at 2 to 8°C, which means that there is no fragmentation or polymerization of IgM above 5% measured in HPSEC, no increase of proteolytic activity, no decrease of IgM antibody activity against Escherichia coli and IgM antibody activity against Pneumococcus saccharide of more than 25 % and no increase in anticomplementary activity of more than 25 %, staying below 1 CH50/mg protein. Still further, the final product produced by the method of the present invention is stable in liquid form for at least 3 months, preferably at least 6 months, and most preferably at least one year at room temperature (between 23 and 27°C) as assessed by the same criteria.

[0058] The process of the present invention further provides a higher level of virus removal than the processes of the prior art resulting in an antibody preparation which is safer than the antibody preparations of the prior art, particularly with respect to active non enveloped viruses like, for example, parvoviruses. The method of the present invention is capable of removing/inactivating viruses, and in particular non-enveloped virus, by more than $3 \log_{10}$, preferably by more than $4 \log_{10}$, and most preferably by more than $5 \log_{10}$. This results in an antibody preparation that is substantially free of virus, and in particular substantially free of non-enveloped virus (i.e. is virus-safe). Still further, the method of the present invention is able to achieve this level of viral particle removal/inactivation without a significant impact on the amount of active IgM or on the anticomplementary activity of the antibody preparation. Accordingly, antibody preparations which are virus safe with respect to enveloped and non-enveloped virus, which contain at least 15%, more preferably at least 20% level of IgM and which have anticomplementary activity of ≤ 1 CH 50/mg protein are obtainable.

[0059] The antibody preparation of the present invention is suitable for use in medicine and can be used in the treatment of immunological disorders and infections, particularly IgM deficiency disorder and bacterial infections. Human IgM-enriched polyvalent immunoglobulin preparation for intravenous administration, which can be prepared by the process of the present invention, contain higher antibody titres against clinically relevant Gram-negative and Gram-positive bacteria as well as higher antibody titres against endotoxins of Gram-negative bacteria and exotoxins of Gram-negative and Gram-positive bacteria compared with polyvalent immunoglobulin G preparations.

[0060] In particular, the antibody preparations of the present invention are suitable for intravenous administration to patients, and in particular are suitable for intravenous injection in humans.

[0061] Thus the antibody preparation of the present invention is suitable for use in a method of treatment of a patient comprising a step of administering the antibody preparation of the present invention to the patient. In particular, the patient can be suffering from an immunological disorder or a bacterial infection.

[0062] The present invention will now be described further by way of example only.

Description of the test method for determination of the anti-complementary activity

[0063] The assay to determine the anti-complementary activity was carried out according to the method described in the European Pharmacopoeia (method 2.6.17, Ph. Eur. 6. Edition, 2008).

[0064] Hemolysin pre-treated sheep erythrocytes are hemolysed by complement. By complement-binding antibodies in the sample the hemolysis is suppressed. The amount of complement is determined, which is bound (inactivated) by 1 mg immunoglobulin.

[0065] A certain amount of immunoglobulin (10 mg) is mixed with complement of guinea pig and the free complement is titrated. The anti-complementary activity is expressed as used complement in respect to the used complement of a reference solution. The hemolytic unit of complement activity (CH_{50}) is the amount of complement leading to hemolysis of 2.5×10^8 optimally prepared erythrocytes of a total amount of 5×10^8 erythrocytes in optimal buffer conditions.

[0066] Optimally prepared erythrocytes (8ml stabilized erythrocytes from sheep, washed three times with gelatine-barbital-buffer, finally 1 ml erythrocyte sediment are suspended in 24 ml gelatine-barbital-buffer) are prepared by mixing 20 ml erythrocytes suspension with 20 ml hemolysine (adjusted to 2 MHE/ml - minimal hemolytic unit) and incubation for 15 min at 37°C.

[0067] An equivalent of 10 mg immunoglobulin is diluted in gelatine-barbital-buffer (1 g gelatine in 1 L barbital buffer pH 7.3, 5-fold barbital-buffer solution: 83 g sodium chloride, 10.192 g sodium barbital in 2 liters water, pH 7.3). To a final volume of 1 ml 200 µl complement 100 CH_{50} / ml are added. The test tubes are incubated under shaking for 1 h at 37 °C. The samples are diluted and titrated against optimally prepared erythrocytes. After an incubation for 1 h at 37°C the samples are centrifuged and the optical density is determined by using a spectrophotometer at a wavelength of 541 nm.

Example 1 -Preparation of an IgM enriched preparation from fraction I/III

[0068] 180 kg Cohn Fraction I/III, originating from cold ethanol fractionation of human plasma are suspended in 720 L 0.1 M sodium acetate buffer pH 5.05 and mixed for 15 - 30 minutes after the suspension temperature is reached (22 ± 4 °C).

[0069] The solution is treated by addition of a 19.8 kg octanoic acid (0.110 kg per kg paste I/III used) at room temperature and the protein solution is further mixed for 80 minutes, using a vibrating mixer (Vibromixer®, Size 4, Graber+Pfenniger GmbH, Vibromixer adjusted to level 2 - 3). The octanoic acid is added slowly over 30 min.

[0070] Approx. 3 kg tri-calcium phosphate ($Ca_3PO_4)_2$) are added and the protein solution is further mixed for at least 15 min. The precipitate is removed by clarifying filtration using a filter press. An additional 0.2 µm filtration is carried out and the protein solution is subjected to ultrafiltration with 10 kD membranes. The protein solution is diafiltered against 0.04 M NaCl solution and afterwards adjusted to a protein concentration of 40 g/L.

[0071] The protein solution is treated at $pH 4.0 \pm 0.1$ after dilution 1+1 with water for injection. pH adjustment is carried out by using 1 M HCl and the protein solution is incubated for 9 h at $37 \text{ °C} \pm 2 \text{ °C}$. After the pH 4 incubation the protein solution is adjusted to pH 5.8, using 1 M NaOH. The resulting protein solution is further purified by adding DEAE Sephadex in a batch mode (75 g DEAE Sephadex per kg protein). The protein solution is incubated under stirring for ≥ 60 min at room temperature. The DEAE Sephadex is removed by clarifying filtration. The protein solution is subjected to a 0.2 µm filtration.

[0072] The protein solution is filtered through a 0.1 µm filter and a Pall, Ultipor VF DV50, 20" filter. The filtrate is further processed by UVC light treatment at 254 nm, using a flow-through UVivatec® process device (Bayer Technology Services / Sartorius Stedim) at a UVC dose of 240 J/m². The flow velocity through the UVC reactor is calculated using the manufacture's instructions. The irradiated protein solution is concentrated to a protein concentration of 50 - 70 g/l by ultrafiltration and is subjected to diafiltration (10 kD membrane, using 0.32 M glycine buffer pH 4.3). The final product is filtered through a 0.2 µm filter and is stored at 2 to 8°C.

Example 2 - Investigation of Conditions in Octanoic Acid Treatment Step

[0073] For the octanoic acid treatment the following experimental ranges were tested, also in combination with each other using the method described in Example 1 (results not shown).

- Octanoic acid amount: 0.09 kg/kg to 0.13 kg/kg (Amount octanoic acid per kg used fraction I/III) (120 to 180 mM octanoic acid)
- pH of the octanoic acid treatment between pH 4.8 and 5.3
- Temperature range of the reaction: 14°C to 30°C
- Incubation time: 40 to 110 min

[0074] All conditions tested lead to intermediates being easy to clarify for further processing and with a extensive reduction of proteolytic activity from several thousand U/L in suspended Cohn fraction I/III). These intermediates result in a final product with a proteolytic activity below 8 U/l (calculated as described below in Example 6) which is the limit of quantitation.

Example 3 - Virus Reduction Through Use of Vibromixer - Determination of virus removal factors for the octanoic acid treatment with and without use of a vibromixer.

[0075] 250 ml of suspended faction I/III were homogenised for 30 min at pH 5.05 and 22°C. The suspension was spiked with 2.6 ml of the virus stock solution. Octanoic acid was added (110 g/kg) and homogenised for 60 min using a vibromixer. In a parallel experiment the same mixture was homogenised with standard stirring. After 60 min tri-calcium phosphate (0.15 g/kg octanoic acid) was added and the suspension stirred for 15 min. The suspension was cleared by depth filtration using a filter disc. The filter disc was pre-rinsed with 70 - 80 ml of buffer. After filtration, the filter was rinsed with 80 ml of buffer. Filtrate and wash were pooled and a sample was drawn for virus titration.

[0076] Virus titres from samples taken prior to addition of octanoic acid and after filtration were determined on appropriate indicator cells for SV40, Reo and PPV (CV-1, CCL7.1 and PK13). Finally, the removal factor was calculated in compliance with the current guidelines for virus validation studies.

[0077] In virus validations studies, non-enveloped viruses such as SV40 and Reo were effectively removed in the order of more than 4 log₁₀ and more than 5 log₁₀, respectively. Moreover, PPV was removed by more than 3 log₁₀. These values are more than 10 times and up to 1000 times higher than with the same octanoic acid treatment under standard stirring conditions without vibromixing.

Table 1: Comparison of the virus reduction factors (log₁₀) for the octanoic acid treatment with and without the use of a vibromixer.

	Octanoic acid reaction standard stirring [log ₁₀ reduction]	Octanoic acid reaction with vibromixing [log ₁₀ reduction]
PPV	2.15 ± 0.32	3.39 ± 0.36
REO	2.34 ± 0.38	5.46 ± 0.28
SV40	2.05 ± 0.4	4.71 ± 0.34

Example 4 - Evaluation of UVC treatment

[0078] The optimal range for the dosage of UVC radiation has been evaluated. There is a balance between the minimal necessary dosage to achieve at least 4 log₁₀ inactivation for non enveloped viruses and the maximum tolerable dosage to avoid denaturation of the IgM molecules leading to an impaired Fab function to bind antigens and impaired Fc function influencing complement activation. In the range of 200 to 400 J/m² one could observe only a slight increase of immunoglobulin aggregates and no significant impact on fragment content.

[0079] For the experiments the optical density (OD) of the original protein solution is used to calculate the flow rate in the UVivatec lab system with the vendor provided Excel-Sheet from BTS (customer Master Calculation Sheet UVivatec Lab II Version 3.0). The flow rate is calculated by taking into account the lamp performance, the set point of the UV signal lamp sensor and the desired UVC irradiation dose.

[0080] IgM containing solution with a protein content of about 55 g/l (Batch 86GB005BE07) was pumped at a flow rate of 5.8 l/h through the UVivatec system in order to achieve a dose of 200 J/m² for a single flow-through. A dose of 300 J/m² was achieved by pumping the protein solution at a flow rate of 3.9 L/m² through the system. 400 J/m² were achieved by pumping the protein solution at a flow rate of 2.9 L/m² through the system.

Table 2: Analytical results for the activity and titre determinations before and after UVC treatment in the concentrated final product

		IgM product no UVC-irradiation	IgM product after UVC 200 J/m ²	IgM product after UVC 300 J/m ²	IgM product after UVC 400 J/m ²
Protein content	g/l	56.3	56.2	57.6	54.4
IgG content (nephelometry)	%	56.1	55.5	55.7	54.9

EP 2 560 691 B1

(continued)

		IgM product no UVC-irradiation	IgM product after UVC 200 J/m ²	IgM product after UVC 300 J/m ²	IgM product after UVC 400 J/m ²
IgA content (nephelometry)	%	20.1	20.6	20.5	20.7
IgM content (nephelometry)	%	23.7	23.9	23.7	24.4
HPSEC aggregates > 1200 kD	area %	1.9	2.6	3.3	4.0
fragments < 100 kD	area %	0.66	0.73	0.76	0.79
ACA	CH50/mg protein	0.68	0.48	0.46	0.46
PA	U/l	<8	<8	<8	<8

[0081] No significant difference could be observed for immunoglobulin content, proteolytic activity or ACA in the range of 200 to 400 J/m². The preferred range for the dosage was set between 200 and 300 J/m² because 200 J/m² are well enough to inactivate non enveloped viruses and at 300 J/m² no significant impact could be seen on aggregate formation and antibody titres. The preferred dosage is 225 J/m²

[0082] Diluted IgM containing solution with a protein content of 8 to 12 g/l (Batch 86BB059BE07) was pumped at a flow rate of 5.8 l/h through the UVivatech system in order to achieve a doses between 200 and 300 J/m² for a single flow-through.

Table 3: Analytical results for IgM solutions before and after UVC irradiation at different UVC doses

Batch fraction I/III 86BB059BE07		before UVC	UVC: 200 J/m ²	UVC: 225 J/m ²	UVC: 250 J/m ²	UVC: 300 J/m ²
Protein	g/l	11.34	10.56	10.65	10.69	10.56
IgG content	%	59.2	59.1	58.5	58.6	57.1
IgA content	%	19.6	19.6	20.2	20.1	20.3
IgM content	%	21.1	21.3	21.2	21.4	22.6
HSEC						
aggregates > 1200 kD	%	0.20	0.39	0.54	0.3	0.47
fragments <100 kD	%	0.47	0.46	0.25	0.26	0.47
PA	U/l	<8	<8	n.t.	n.t.	n.t.
PKA	U/ml	3	3	3	3	3
ACA	CH50/ mg protein	0.1	0.08	0.1	0.1	0.18
Anti-E.coli O1:K1-IgG	U/mg	24.7	20.5	18.9	19.5	20.2
Anti-E.coli O1:K1-IgA	U/mg	9.4	9.5	9.5	9.1	8.9
Anti-E.coli O1:K1-IgM	U/mg	14.1	13.0	15.1	13.9	13.4
Anti-Candida albicans - IgG	U/mg	15.6	16.8	17.9	17.3	17.0
Anti-Candida albicans - IgA	U/mg	11.3	11.6	10.5	10.3	10.4
Anti-Candida albicans - IgM	U/mg	13.8	13.3	13.7	13.9	13.1
Anti-Enterococcusfaecalis - IgG	U/mg	13.0	15.5	13.5	14.8	15.0

(continued)

Batch fraction I/III 86BB059BE07		before UVC	UVC: 200 J/m ²	UVC: 225 J/m ²	UVC: 250 J/m ²	UVC: 300 J/m ²
Anti-Enterococcusfaecalis - IgA	U/mg	11.3	10.5	10.1	9.7	9.6
Anti-Enterococcusfaecalis - IgM	U/mg	17.2	14.1	16.7	14.0	13.9
Anti-Pneumococcus Saccharid -IgG	U/mg	23.2	24.1	24.7	24.0	25.7
Anti-Pneumococcus Saccharid -IgA	U/mg	13.3	12.1	18.0	16.5	14.8
Anti-Pneumococcus Saccharid -IgM	U/mg	17.5	15.1	18.0	16.4	16.6

[0083] The distribution between the immunoglobulin classes remains unaffected by the UV irradiation within this dosage range procedure. The molecular weight distribution pattern analyzed by HPSEC is also not changed. The level of purity analyzed by CZE remains unchanged. Proteolytic activity (PA), prekallikrein activator (PKA) and anti-complementary activity (ACA) are unchanged. Also the anti bacterial activity measured by an Elisa method are not significantly altered for all immunoglobulin classes.

[0084] The aliquots - irradiated with increasing UV intensities - were further processed until final product and subjected to the same panel of analytical tests. There was also no significant difference observable in the final products. All tested antibody titres are always in the range of 100 ± 10 % of the control preparation not UVC treated.

Example 5 - Overall Virus Reduction Through Use of Vibromixer/pH4 treatment and UVC treatment - Determination of virus removal factors

[0085] The validation of virus removal/inactivation of the three steps octanoic acid treatment with vibromixing, pH4 treatment and UVC treatment (215 J/m²) was performed using the following model viruses : Bovine Viral Diarrhea Virus (BVDV) as model virus for Hepatitis C Virus, Pseudorabies Virus (PRV) as model virus for Human Herpes Viruses, Human Immunodeficiency virus (HIV-1), Equine Arteritis Virus (EAV) as model virus for corona viruses, Sindbis Virus (SinV) as model virus for Flavi viruses, Murine Encephalomyelitis Virus (MEV) as model virus for Hepatitis A Virus, Reovirus (Reo) as model virus for other non enveloped viruses, Porcine Parvovirus (PPV) as model virus for human Parvovirus B 19.

[0086] The results of these studies with the three steps octanoic acid treatment, pH4 treatment and UVC treatment are listed in the following Table 2.

Table 4: Total virus reduction by the IgM production process

Model virus	BVDV	PRV	HIV-1	EAV	SinV	MEV	Reo	PPV
Total reduction (log ₁₀)	>12.5	>10.1	>12.7	>8.4 ^a	>13.7 ^a	9.2	>11.0	>8.4
^a Reduction factor without data for validation of the UVC irradiation step								

[0087] The optional nano filtration with filters with a nominal pore size of about 50 nm adds additional safety by increasing the total reduction up to more than 17 log₁₀ depending on the size of the virus. E.g >17.5 log₁₀ are then reached for HIV-1 whereas PPV was not further removed by nano filtration.

[0088] Therefore the purification procedure according to the invention leads to an outstanding virus safe IgM preparation with up to now for such an IgM containing preparation unreached virus inactivation / reduction rates of more than 8 log₁₀. This is especially important for the non enveloped viruses like MEV, Reo and PPV which are generally more resistant against virus inactivation and removal procedures due to their small size and the lack of a lipid envelope.

Example 6: Determination of residual proteolytic activity for the octanoic acid treatment with and without use of a vibromixer.

[0089] The octanoic acid treatment was performed like in example 1 and in a parallel experiment without a vibromixer

but with vigorous standard stirring with a blade stirrer. The proteolytic activity in samples after octanoic acid / tricalcium phosphate treatment and ultra-/diafiltration were determined using the chromogenic substrate S-2288 (Chromogenix), following the manufacturers instructions. 25 mg of the substrate S-2288 (Chromogenix) are dissolved in 7,2 ml water-for-injection. Samples are diluted into buffer (100 mM Tris/HCl pH 8.4, 106 mM NaCl) to meet the linear range of the assay, e.g. 200 µl buffer are mixed with 200 µl sample (mixing and temperature adjustment to 37 °C) and 200 µl chromogenic substrate solution. The absorption kinetics are measured at 405 nm (1-3 min) at 37 °C, using a spectrophotometer. The proteolytic activity of the sample is calculated from the initial absorption difference ($\Delta\text{Abs}/\text{min}$) by using the equation $C \text{ (U/L)} = 313 * \Delta\text{Abs}/\text{min} * F$ (C = proteolytic activity, F = dilution factor)

Table 5: Reduction of proteolytic activity by octanoic acid treatment

	Octanoic acid treatment without vibromixing	Octanoic acid treatment with vibromixing
Starting material (U/l)	5630	5630
Mean residual proteolytic activity after octanoic acid treatment (U/L)	42	< 8 (LOD)

[0090] The filtrate after octanoic acid treatment was limpid when vibromixing was employed. In the comparison experiment the filtrate after octanoic acid treatment with blade stirrer was very opaque and difficult to filtrate.

Example 7: Anti-bacterial titres in an IgM preparation according to the invention

[0091] For comparison with the only commercially available intravenous tolerable IgM containing preparation, Pentaglobin, the anti-bacterial activities were analyzed in three batches of this well established drug and compared to a preparation according to the invention. The determination of antibodies of the IgA or IgM class in the IgM preparation versus antibacterial or antifungal antigens was carried out by ELISA. Microtitre plates were coated with a corresponding antigen and incubated with a standard or the IgM preparation. Antibodies bound to the antigen were detected with an anti-human-IgA or anti-human-IgM conjugate. The detection was carried out by using an enzyme substrate. The resulting colour change is corresponding to the amount of antibodies present in IgM preparation.

Table 6 Comparison of anti bacterial binding activity of IgM in an preparation according to the invention and commercially available Pentaglobin

parameter	unit	IgM preparation invention mean	Pentaglobin commercial product mean
IgM antibodies against <i>Pneumococcus saccharide</i>	U/mg IgM	72	21
IgM antibodies against <i>Escherichia coli</i>	U/mg IgM	62	39
IgM antibodies against <i>Enterococcus faecalis</i>	U/mg IgM	69	27
IgM antibodies against <i>Candida albicans</i>	U/mg IgM	61	41
IgM antibodies against <i>Chlamydia</i>	U/mg IgM	71	6

Table 7 Comparison of anti bacterial binding activity of IgA in an preparation according to the invention and commercially available Pentaglobin

parameter	unit	IgM preparation invention mean	Pentaglobin commercial product mean
IgA antibodies against <i>Pneumococcus saccharide</i>	U/mg IgA	86	25

(continued)

parameter	unit	IgM preparation invention mean	Pentaglobin commercial product mean
IgA antibodies against <i>Escherichia coli</i>	U/mg IgA	83	26
IgA antibodies against <i>Enterococcus faecalis</i>	U/mg IgA	93	21
IgA antibodies against <i>Chlamydia</i>	U/mg IgA	65	38
IgA antibodies against <i>Helicobacter pylori</i>	U/mg IgA	59	24

[0092] The IgM and IgA mediated activities in the new preparation were typically at least 1.5 times as high as in Pentaglobin which can be explained by the fact that IgM and IgA in Pentaglobin is chemically modified with β -Propiolactone. This step is replaced by the more gentle procedures according to this invention.

[0093] Overall these data demonstrate that the binding region of the IgM molecules in the final preparation is functionally full active.

Example 8 Storage stability studies with liquid IgM product

[0094] Product according example 1 without UVC treatment was stored in 10 or 100 ml glass vials (filling volume 5 ml or 50 ml) at 2 - 8°C and analyzed for all parameter according to specification. The results are shown in Table 8. Parameter which are relevant to show stability are the aggregate and fragment content measured with high performance size exclusion chromatography (HPSEC), proteolytic activity (PA) and anticomplementary activity (ACA). These parameter are critical for intravenous tolerability and likely to change during long term storage. At 2 - 8 °C there was no significant change of these parameters. Even at storage at room temperature (23 - 27°C) these values remained within specification, although there is a slight increase of fragments after 24 months at room temperature. Other parameter like coloration, opalescence, pH value were also determined and stayed unchanged over the whole study period. IgM and IgA titres against various bacteria remain stable over 2 years at 2 - 8°C.

[0095] Product according example 1 with UVC treatment was also stored in 10 or 100 ml glass vials (filling volume 5 ml or 50 ml) at 2 - 8°C and room temperature and analyzed for all parameter according to specification. The results are shown in Table 9. In this ongoing stability study the currently available 12 month date show the same stability profile of the product as without UVC treatment which allows the extrapolation to a 24 month stability.

Table 8 Stability of the batch A586067 tested at 2-8 °C lying position Filling size: 5 ml

Parameters tested	Requirement (Tolerance)	Storage in months 2-8 °C							23 - 27°C
		0	3	6	9	12	18	24	24
protein (g/l)	45-55	50.3	51.4	50.3	50.4	50.5	49.6	50.8	49.8
HPSEC									
% aggregates > 1200 kD	≤5	0.9	0.6	0.5	0.8	0.6	1.0	1.3	1.7
% fragments < 100 kD	≤5	0.2	0.6	1.1	0.7	1.6	0.9	1.2	4.1
proteolytic activity (U/l)	<8	<8	<8	<8	n.t.	< 8	n.t.	< 8	<8
immunoglobulin content (%)	>95%	96.7	99.0	100	n.t.	99.5	n.t.	98.4	97.5
IgM content	≥ 20 %	21.6	22.1	22.1	n.t.	22.3	n.t.	20.9	20.5
anticomplementary activity (CH50/mg protein)	≤1.0	0.48	0.56	0.48	0.66	0.70	0.64	0.54	0.38
n.t. = not tested									

Table 9 Stability of the batch A586057 tested at 2-8 °C lying position Filling size: 50 ml

Parameters tested	Requirement (Tolerance)	Storage in months 2 - 8 °C							23 - 27°C
		0	3	6	9	12	18	24	
protein (g/l)	45-55	50.2	50.8	49.7	50.4	50.3	49.4	50.3	49.7
HPSEC									
% aggregates > 1200 kD		0.9	0.5	0.4	0.8	0.6	1.0	1.3	1.5
% fragments < 100 kD	≤5	0.3	0.6	1.0	0.9	1.4	1.2	1.2	4.2
proteolytic activity (U/l)	<8	<8	<8	<8	n.t.	<8	n.t.	<8	<8
immunoglobulin content (%)	>95%	98.6	98.9	100	n.t.	99.5	n.t.	98.5	98.0
IgM content	≥ 20 %	21.3	22.3	24.5	n.t.	22.0	n.t.	20.9	20.1
anticomplementary activity (CH50/mg protein)	≤ 1.0	0.48	0.82	0.52	0.64	0.68	0.48	0.60	0.40

Example 9 In vivo experiments with IgM product

[0096] To confirm safety and tolerability the effects of the IgM preparation on arterial blood pressure following repeated intravenous infusions over 5 days were studied in 8 conscious cynomolgus monkeys. A dose of 190 mg/IgM/kg/day of the IgM preparation prepared according to the methods described herein was administered. Pentaglobin, the commercially available intravenous tolerable IgM containing preparation was administered to some monkeys as a comparison substance. Pentaglobin was administered in such a way that the same IgM dose was administered. A control dose of 0.9% NaCl was administered to the animals sometime prior to the administration of the immunoglobulin preparations. Blood pressure was determined following injection to determine whether administration was associated with an intolerable level of non-specific complement activation.

[0097] The administration of the IgM preparation (15 ml/kg/day) had only minor effects on arterial blood pressure (mean, systolic and diastolic). The differences up to 4 hours after every infusion compared to pretest values did not exceed 4 mmHg. These differences can be considered not biologically relevant.

[0098] C3a levels were determined in plasma samples taken after injection as a marker for unspecific activation of the complement pathway. C3a levels [ng/ml] were only slightly increased by the administration of the IgM preparation (15 mL/kgBW) and were even lower than with the commercially available reference preparation Pentaglobin at equal amounts of IgM. Blood samplings were performed approximately 6 hours after treatment.

Table 10a C3a levels [ng/ml] after the administration of the IgM preparation

	Control (0.9% NaCl, pH 4.5) C3a [ng/ml]	Administration of IgM preparation C3a [ng/ml]
Mean	229	240
SD	83	37
N	8	8

[0099] No substantial toxicological findings could be attributed to the IgM preparation, and there were no relevant alterations that have not been observed with Pentaglobin. As the safety of Pentaglobin is well established in the clinical practice of many years it is reasonable to conclude that these alterations do not have any clinical relevance.

Table 10b C3a levels [ng/ml] after the administration of the reference preparation Pentaglobin

	Control (0.9% NaCl, pH 6.8) C3a [ng/ml]	Administration of Pentaglobin C3a [ng/ml]
Mean	204	263

(continued)

	Control (0.9% NaCl, pH 6.8) C3a [ng/ml]	Administration of Pentaglobin C3a [ng/ml]
SD	20	61
N	4	4

[0100] The good tolerability and safety of the IgM preparation was also verified in a human Phase I study in 24 healthy male and female volunteers. Systolic blood pressure in the first 4 hours after administration in the mean decreased only about 9 % (11.9 mmHg) after infusion of 91 to 274 mg of the IgM preparation per kg BW/d at 0.5 ml/min.

[0101] This was in the same range like the placebo 0.9 % NaCl-solution (9.4%, 11.7 mmHg). No serious adverse events were recorded and all non-serious adverse events were self limiting. Further, there was no evidence for the transmission of an infectious agent, as shown by PCT determinations.

[0102] It is noted that the usefulness of efficacy studies in animal models of relevant diseases is limited due to the immunogenicity and preformed Gal-antibodies in IgM preparations obtained from human plasma. However, given the prior art knowledge regarding the use of Pentaglobin in the treatment of disease and the anti-bacterial antibody titres of the IgM preparation prepared by the method of the present invention (as demonstrated in Example 7) it can be concluded that the IgM preparation have clinical efficacy.

Claims

1. A process for the preparation of an IgM containing immunoglobulin composition from a plasma fraction comprising immunoglobulins, the process comprising:

- (a) providing a plasma fraction as a solution containing the immunoglobulins;
- (b) mixing a C₇ to C₉ carboxylic acid with the solution and treating the mixed solution with a vibrating agitator to precipitate contaminating proteins; and
- (c) separating the precipitated proteins from the solution to yield the IgM containing immunoglobulin composition.

2. A process according to claim 1 wherein in step (b):

- (i) the concentration of the C₇ to C₉ carboxylic acid is at least 0.075 kg/kg of plasma fraction; and/or
- (ii) the pH of the mixed solution is between 4.5 to 5.5; and/or
- (iii) the temperature of the mixed solution is 10°C to 35°C; and/or
- (iv) the C₇ to C₉ carboxylic acid is incubated with the solution containing immunoglobulins for at least 30 minutes.

3. A process according to claim 1 or claim 2 wherein:

- (i) the C₇ to C₉ carboxylic acid is octanoic acid; and/or
- (ii) the immunoglobulins of the plasma fraction comprise at least 5% IgM; and/or
- (iii) the plasma fraction is a precipitation of Cohn fraction I/III or Kistler/Nitschmann fraction B or B+I; and/or
- (iv) step (c) comprises ultrafiltration and the immunoglobulin composition comprises a filtered solution.

4. A process according to any preceding claim further comprising a step of incubating the IgM containing immunoglobulin composition from step (c) at between pH 3.5 and pH 4.5 to form an incubated solution, and preferably wherein the step of incubating the IgM containing immunoglobulin composition from step (c) is conducted at between 32 and 42°C.

5. A process according to claim 4 further comprising the steps of subjecting the incubated solution to adsorption on DEAE-Sephadex and separating the DEAE Sephadex from the solution by depth filtration.

6. A process according to claim 5 further comprising the step of subjecting the filtrate from the depth filtration to nanofiltration, and preferably wherein the nanofiltration is conducted with a filter with a 35 to 75 nm nominal pore size and more preferably a 40 to 50 nm nominal pore size.

7. A process according to any one of claims 4 to 6 further comprising a step of treating the incubated solution of claim

4 or the filtrate of claim 5 or claim 6 with UVC irradiation to form a UVC irradiated solution, and preferably wherein the incubated solution or the filtrate is treated with UVC irradiation at 200 to 500 J/m², and more preferably at 200 to 300 J/m².

- 5 8. A process according to claim 7 further comprising the step of filtering the UVC irradiated solution under sterile conditions to produce an antibody preparation suitable for intravenous administration, and preferably further comprising formulating the antibody preparation in a glycine-containing buffer at a pH between pH 4 and 5.5.
9. A process according to claim 7 or claim 8, wherein:
 - 10 (i) the process further comprises a step of filling a container with the UVC irradiated solution of claim 7 or the antibody preparation of claim 8 under sterile conditions; and/or
 - (ii) the UVC irradiated solution of claim 7 or the antibody preparation of claim 8 has a proteolytic activity of less than 8U/l; and/or
 - 15 (iii) the process provides a more than 3 log₁₀ removal of non-enveloped viruses.
10. An antibody preparation comprising immunoglobulins obtainable by the process of claim 8 or claim 9, preferably wherein at least 15% of the total immunoglobulins are IgM.
- 20 11. An antibody preparation comprising immunoglobulins IgG, IgA and IgM, wherein at least 15% of the total immunoglobulins are IgM, wherein the antibody preparation is virus safe with respect to enveloped and non-enveloped virus, has an anticomplementary activity of ≤ 1 CH 50/mg protein, and is stable in liquid form for at least 6 months when stored at 2 to 8°C.
- 25 12. An antibody preparation according to claim 11, which has a proteolytic activity of less than 8U/l.
13. An antibody preparation according to any one of claims 10 to 12 wherein:
 - 30 (i) the preparation comprises at least 20% IgM and preferably 20% to 30% IgM; and/or
 - (ii) the preparation further comprising a stabilizing agent, preferably wherein the stabilizing agent is glycine; and/or
 - (iii) wherein the preparation is stable for at least 2 years when stored at 2 to 8°C.; and/or
 - (iv) wherein the immunoglobulins are not chemically modified, or wherein the preparation has not been treated during its process of production with β-propiolactone or with an added protease such as pepsin.
- 35 14. An IgM containing immunoglobulin composition obtainable by the method of any one of claims 1 to 3.
15. An antibody preparation according to any one of claims 10 to 13 for use in medicine.
- 40 16. An antibody preparation according to claim 15 for use in the treatment of immunological disorders or bacterial infections, preferably wherein the immunological disorder is an IgM deficiency disorder.
17. Use of an antibody preparation according to any one of claims 10 to 13 for the manufacture of a medicament for treatment of immunological disorders and bacterial infections.

Patentansprüche

- 50 1. Verfahren zur Herstellung einer IgM enthaltenden Immunglobulinzusammensetzung aus einer Immunglobuline umfassenden Plasmafraktion, wobei das Verfahren umfasst:
 - (a) Bereitstellen einer Plasmafraktion als eine Lösung, welche die Immunglobuline enthält;
 - (b) Mischen einer C₇- bis C₉-Carbonsäure mit der Lösung und Behandeln der gemischten Lösung mit einem Vibrationsmischer, um kontaminierende Proteine auszufällen; und
 - 55 (c) Abscheiden der ausgefällten Proteine aus der Lösung, um die IgM enthaltende Immunglobulinzusammensetzung zu ergeben.
2. Verfahren nach Anspruch 1, wobei in Schritt (b):

- (i) die Konzentration der C₇- bis C₉-Carbonsäure mindestens 0,075 kg/kg Plasmafraktion ist; und/oder
- (ii) der pH-Wert der gemischten Lösung zwischen 4,5 und 5,5 liegt; und/oder
- (iii) die Temperatur der gemischten Lösung 10°C bis 35°C beträgt; und/oder
- (iv) die C₇- bis C₉-Carbonsäure mit der Immunglobuline enthaltenden Lösung mindestens 30 Minuten lang inkubiert wird.

3. Verfahren nach Anspruch 1 oder Anspruch 2, wobei:

- (i) die C₇- bis C₉-Carbonsäure Octansäure ist; und/oder
- (ii) die Immunglobuline der Plasmafraktion mindestens 5% IgM umfassen; und/oder
- (iii) die Plasmafraktion eine Ausfällung der Cohn-Fraktion I/III oder der Kistler/Nitschmann-Fraktion B oder B+I ist; und/oder
- (iv) Schritt (c) eine Ultrafiltration umfasst und die Immunglobulinzusammensetzung eine filtrierte Lösung umfasst.

4. Verfahren nach einem vorhergehenden Anspruch, ferner umfassend einen Schritt des Inkubierens der IgM enthaltenden Immunglobulinzusammensetzung aus Schritt (c) bei einem pH zwischen 3,5 und 4,5, um eine inkubierte Lösung zu bilden, und wobei vorzugsweise der Schritt des Inkubierens der IgM enthaltenden Immunglobulinzusammensetzung aus Schritt (c) zwischen 32 und 42°C durchgeführt wird.

5. Verfahren nach Anspruch 4, ferner umfassend die Schritte des Unterziehens der inkubierten Lösung einer Adsorption an DEAE-Sephadex und des Abtrennens des DEAE Sephadex von der Lösung mittels Tiefenfiltration.

6. Verfahren nach Anspruch 5, ferner umfassend den Schritt des Unterziehens des Filtrats aus der Tiefenfiltration einer Nanofiltration und wobei vorzugsweise die Nanofiltration mit einem Filter mit einer nominalen Porengröße von 35 bis 75 nm und stärker bevorzugt mit einer nominalen Porengröße von 40 bis 50 nm durchgeführt wird.

7. Verfahren nach einem der Ansprüche 4 bis 6, ferner umfassend einen Schritt des Behandeln der inkubierten Lösung des Anspruchs 4 oder des Filtrats des Anspruchs 5 oder des Anspruchs 6 mit UVC-Bestrahlung, um eine UVC bestrahlte Lösung zu bilden, und wobei vorzugsweise die inkubierte Lösung oder das Filtrat mit UVC-Bestrahlung bei 200 bis 500 J/m² und stärker bevorzugt bei 200 bis 300 J/m² behandelt werden.

8. Verfahren nach Anspruch 7, ferner umfassend den Schritt des Filtrierens der UVC bestrahlten Lösung unter sterilen Bedingungen, um ein Antikörperpräparat zu erzeugen, das zur intravenösen Verabreichung geeignet ist, und vorzugsweise ferner umfassend das Formulieren des Antikörperpräparats in einem Glycin enthaltenden Puffer bei einem pH zwischen 4 und 5,5.

9. Verfahren nach Anspruch 7 oder Anspruch 8, wobei:

- (i) das Verfahren ferner einen Schritt des Befüllens eines Behälters mit der UVC bestrahlten Lösung des Anspruchs 7 oder des Antikörperpräparats des Anspruchs 8 unter sterilen Bedingungen umfasst; und/oder
- (ii) die UVC bestrahlte Lösung des Anspruchs 7 oder das Antikörperpräparat des Anspruchs 8 eine proteolytische Aktivität von weniger als 8 U/I aufweist; und/oder
- (iii) das Verfahren ein Entfernen von mehr als 3 log₁₀ der unbehüllten Viren bereitstellt.

10. Antikörperpräparat, das Immunglobuline umfasst, erhältlich durch das Verfahren des Anspruchs 8 oder Anspruchs 9, wobei vorzugsweise mindestens 15% der gesamten Immunglobuline IgM sind.

11. Antikörperpräparat, das die Immunglobuline IgG, IgA und IgM umfasst, wobei mindestens 15% der gesamten Immunglobuline IgM sind, wobei das Antikörperpräparat in Bezug auf die behüllten und unbehüllten Viren virussicher ist, eine antikomplementäre Aktivität von ≤ 1 CH 50/mg Protein aufweist und in flüssiger Form mindestens 6 Monaten lang stabil ist, wenn es bei 2 bis 8°C gelagert wird.

12. Antikörperpräparat nach Anspruch 11, das eine proteolytische Aktivität von weniger als 8 U/I aufweist.

13. Antikörperpräparat nach einem der Ansprüche 10 bis 12, wobei:

- (i) das Präparat mindestens 20% IgM und vorzugsweise 20% bis 30% IgM umfasst; und/oder
- (ii) das Präparat ferner einen Stabilisator umfasst, wobei der Stabilisator vorzugsweise Glycin ist; und/oder

- (iii) wobei das Präparat mindestens 2 Jahre lang stabil ist, wenn es bei 2 bis 8°C gelagert wird; und/oder
- (iv) wobei die Immunglobuline nicht chemisch modifiziert sind, oder wobei das Präparat während seines Produktionsverfahrens nicht mit β -Propiolacton oder mit einer zugesetzten Protease wie Pepsin behandelt wurde.

14. IgM enthaltende Immunglobulinzusammensetzung, erhältlich durch das Verfahren nach einem der Ansprüche 1 bis 3.
15. Antikörperpräparat nach einem der Ansprüche 10 bis 13 zur Verwendung in der Medizin.
16. Antikörperpräparat nach Anspruch 15 zur Verwendung bei der Behandlung von immunologischen Störungen oder bakteriellen Infektionen, wobei die immunologische Störung vorzugsweise eine IgM-Mangelkrankung ist.
17. Verwendung eines Antikörperpräparats nach einem der Ansprüche 10 bis 13 zur Herstellung eines Medikaments zur Behandlung von immunologischen Störungen und bakteriellen Infektionen.

Revendications

1. Procédé pour la préparation d'une composition d'immunoglobulines contenant des IgM à partir d'une fraction de plasma comprenant des immunoglobulines, le procédé comprenant :
 - (a) la fourniture d'une fraction de plasma sous forme d'une solution contenant les immunoglobulines ;
 - (b) le mélange d'un acide carboxylique en C₇ à C₉ avec la solution et le traitement de la solution mixte avec un agitateur vibratoire pour provoquer la précipitation des protéines contaminantes ; et
 - (c) la séparation des protéines précipitées de la solution pour obtenir la composition d'immunoglobulines contenant des IgM.
2. Procédé suivant la revendication 1, dans lequel, dans l'étape (b) :
 - (i) la concentration de l'acide carboxylique en C₇ à C₉ est d'au moins 0,075 kg/kg de fraction de plasma ; et/ou
 - (ii) le pH de la solution mixte est compris entre 4,5 et 5,5 ; et/ou
 - (iii) la température de la solution mixte va de 10°C à 35°C ; et/ou
 - (iv) l'acide carboxylique en C₇ à C₉ est mis en incubation avec la solution contenant les immunoglobulines pendant au moins 30 minutes.
3. Procédé suivant la revendication 1 ou la revendication 2, dans lequel :
 - (i) l'acide carboxylique en C₇ à C₉ est l'acide octanoïque ; et/ou
 - (ii) les immunoglobulines de la fraction de plasma comprennent au moins 5 % de IgM ; et/ou
 - (iii) la fraction de plasma est un précipité de fraction I/III de Cohn ou de fraction de Kistler/Nitschmann B ou B+I ; et/ou
 - (iv) l'étape (c) comprend une ultrafiltration et la composition d'immunoglobulines comprend une solution filtrée.
4. Procédé suivant l'une quelconque des revendications précédentes, comprenant en outre une étape d'incubation de la composition d'immunoglobulines contenant des IgM de l'étape (c) à un pH entre 3,5 et 4, pour former une solution incubée, et de préférence dans lequel l'étape d'incubation de la composition d'immunoglobulines contenant des IgM de l'étape (c) est mise en oeuvre à une température entre 32 et 42°C.
5. Procédé suivant la revendication 4, comprenant en outre les étapes consistant à soumettre la solution incubée à une adsorption sur DEAE-Sephadex et à séparer le DEAE-Sephadex de la solution par filtration profonde.
6. Procédé suivant la revendication 5, comprenant en outre l'étape consistant à soumettre le filtrat de la filtration profonde à une nanofiltration, et de préférence dans lequel la nanofiltration est effectuée avec un filtre ayant un diamètre nominal des pores de 35 à 75 nm et plus avantageusement un diamètre nominal des pores de 40 à 50 nm.
7. Procédé suivant l'une quelconque des revendications 4 à 6, comprend en outre une étape de traitement de la solution incubée de la revendication 4 ou du filtrat de la revendication 5 ou de la revendication 6 à une irradiation UVC pour former une solution ayant subi une irradiation UVC, et de préférence dans lequel la solution incubée ou

le filtrat est traité par une irradiation UVC à 200 à 500 J/m², et plus avantageusement à 200 à 300 J/m².

8. Procédé suivant la revendication 7, comprenant en outre l'étape de filtration de la solution ayant subi une irradiation UVC dans des conditions stériles pour produire une préparation d'anticorps apte à l'administration intraveineuse, et comprenant en outre de préférence la formulation de la préparation d'anticorps dans un tampon contenant de la glycine à un pH entre 4 et 5,5.

9. Procédé suivant la revendication 7 ou la revendication 8 :

(i) le procédé comprenant en outre une étape de remplissage d'un récipient avec la solution ayant subi une irradiation UVC de la revendication 7 ou la préparation d'anticorps de la revendication 8 dans des conditions stériles ; et/ou

(ii) dans lequel la solution ayant subi une irradiation UVC de la revendication 7 ou la préparation d'anticorps de la revendication 8 a une activité protéolytique inférieure à 8 U/l ; et/ou

(iii) le procédé provoquant une élimination supérieure à 3 log₁₀ des virus sans enveloppe.

10. Préparation d'anticorps comprenant des immunoglobulines pouvant être obtenues par le procédé de la revendication 8 ou de la revendication 9, de préférence dans laquelle une proportion d'au moins 15 % des immunoglobulines totale consiste en IgM.

11. Préparation d'anticorps comprenant des immunoglobulines IgG, IgA et IgM, dans laquelle une proportion d'au moins 15 % des immunoglobulines totale consiste en IgM, ladite préparation d'anticorps étant sans risque viral en rapport avec les virus à enveloppe et les virus sans enveloppe, possédant une activité anti-complémentaire inférieure ou égale à 1 CH 50/mg de protéine, et étant stable sous forme liquide pendant au moins 6 mois lors de son stockage à une température de 2 à 8°C.

12. Préparation d'anticorps suivant la revendication 11, qui a une activité protéolytique inférieure à 8 U/l.

13. Préparation d'anticorps suivant l'une quelconque des revendications 10 à 12 :

(i) la préparation comprenant au moins 20 % d'IgM et de préférence 20 % à 30 % d'IgM ; et/ou

(ii) la préparation comprenant en outre un agent stabilisant, l'agent stabilisant étant de préférence la glycine ; et/ou

(iii) la préparation étant stable pendant au moins 2 ans lors de son stockage à une température de 2 à 8°C ; et/ou

(iv) dans laquelle les immunoglobulines ne sont pas modifiées chimiquement, ou bien la préparation n'a pas été traitée au cours de son procédé de production avec de la β-propiolactone ou avec une protéase ajoutée telle que la pepsine.

14. Composition d'immunoglobulines contenant des IgM, pouvant être obtenue par le procédé de l'une quelconque des revendications 1 à 3.

15. Préparation d'anticorps suivant l'une quelconque des revendications 10 à 13 pour une utilisation en médecine.

16. Préparation d'anticorps suivant la revendication 15, pour une utilisation dans le traitement de troubles immunologiques ou d'affections bactériennes, le trouble immunologique étant de préférence un trouble de déficience en IgM.

17. Utilisation d'une préparation d'anticorps suivant l'une quelconque des revendications 10 à 13 pour la production d'un médicament pour le traitement de troubles immunologiques et d'affections bactériennes.




FIGURE 1

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- EP 0013901 A [0008]
- EP 0352500 A [0008]
- EP 0413187 A [0009] [0010] [0014]
- EP 0413188 A [0009]
- EP 0413188 B1 [0011]
- EP 0450412 A [0012]
- EP 0835880 A [0013] [0014]
- US 6136312 A [0013]
- EP 0345543 A [0015]
- EP 1842561 A [0017]

ELJÁRÁS IMMUNGLOBULIN-KÉSZÍTMÉNY ELŐÁLLÍTÁSÁRA

Szabadalmi igénypontok

1. Eljárás IgM-et tartalmazó immunglobulin-készítmény előállítására immunglobulinokat tartalmazó plazmafrakcióból, amely eljárás magában foglalja a következőket:

- (a) plazmafrakció biztosítása immunglobulinokat tartalmazó oldatként,
- (b) C₇-C₉ karbonsav összekeverése az oldattal és az összekevert oldat kezelése vibrációs keverővel, ezáltal a szennyező fehérjék kicsapása, és
- (c) a kicsapott fehérjék és az oldat szétválasztása, ezáltal az IgM-et tartalmazó immunglobulin-készítmény előállítása.

2. Az 1. igénypont szerinti eljárás, ahol a (b) lépés során:

- (i) a C₇-C₉ karbonsav koncentrációja legalább 0,075 kg a plazmafrakció kg-jára számítva, és/vagy
- (ii) az összekevert oldat pH-ja 4,5-5,5, és/vagy
- (iii) az összekevert oldat hőmérséklete 10-35°C, és/vagy
- (iv) a C₇-C₉ karbonsav legalább 30 percig van inkubálva az immunglobulint tartalmazó oldattal.

3. Az 1. vagy 2. igénypont szerinti eljárás, ahol:

- (i) a C₇-C₉ karbonsav oktánsav, és/vagy
- (ii) a plazmafrakcióban lévő immunglobulin legalább 5% IgM-et tartalmaz, és/vagy
- (iii) a plazmafrakció Cohn-frakció I/III vagy Kistler/Nitschmann-frakció B vagy B+I csapadéka, és/vagy
- (iv) a (c) lépés tartalmaz ultraszűrést és az immunglobulin-készítmény tartalmaz szűrt oldatot.

4. Az előző igénypontok bármelyike szerinti eljárás, amely tartalmazza a következő lépést is: a (c) lépésből származó, IgM-et tartalmazó immunglobulin-készítmény inkubálása 3,5-4,5-ös pH-n, ezáltal inkubált oldat előállítása, és előnyösen ahol a (c) lépésből származó, IgM-et tartalmazó immunglobulin-készítmény inkubálása 32-42°C-on történik.

5. A 4. igénypont szerinti eljárás, amely tartalmazza a következő lépéseket is: az inkubált oldat DEAE-Sephadexen történő adszorpciónak történő alávetése és a DEAE Sephadex elválasztása az oldattól mélységi szűréssel.

6. Az 5. igénypont szerinti eljárás, amely tartalmazza a következő lépést is: a mélységi szűrésből származó szűrlet alávetése nanoszűrésnek, és előnyösen ahol a nanoszűrés 35-75 nm nominális pórusméretű filterrel és előnyösebben 40-50 nm nominális pórusméretű filterrel történik.

7. A 4-6. igénypontok bármelyike szerinti eljárás, amely tartalmazza a következő lépést is: a 4. igénypont szerinti inkubált oldat vagy az 5. vagy 6. igénypont szerinti szűrlet kezelése UVC-sugárzással, ezáltal UVC-vel besugárzott oldat előállítása, ahol az inkubált oldat vagy a szűrlet 200-500 J/m², előnyösebben 200-300 J/m² UVC-sugárzással kezelt.

8. A 7. igénypont szerinti eljárás, amely tartalmazza a következő lépést is: az UVC-vel besugárzott oldat szűrése steril körülmények között, ezáltal intravénás beadásra alkalmas ellenanyag-készítmény előállítása, és előnyösen tartalmazza a következőt is: az ellenanyag-készítmény kiszűrése glicintartalmú pufferben, 4-5,5 pH-értéken.

9. A 7. vagy 8. igénypont szerinti eljárás, ahol:

- (i) az eljárás tartalmazza a következő lépést is: tároló megtöltése a 7. igénypont szerinti, UVC-vel besugárzott oldattal vagy a 8. igénypont szerinti ellenanyag-készítménnyel steril körülmények között, és/vagy
- (ii) a 7. igénypont szerinti, UVC-vel besugárzott oldat vagy a 8. igénypont szerinti ellenanyag-készítmény proteolitikus aktivitása kisebb, mint 8E/l, és/vagy
- (iii) az eljárás a burok nélküli vírusok nagyobb, mint $3 \log_{10}$ mértékű eltávolítását biztosítja.

10. Ellenanyag-készítmény, amely a 8. vagy 9. igénypont szerinti eljárással előállítható immunglobulint tartalmaz, előnyösen ahol az összes immunglobulin legalább 15%-a IgM.

11. Ellenanyag-készítmény, amely IgG, IgA és IgM immunglobulint tartalmaz, ahol az összes immunglobulin legalább 15%-a IgM, ahol az ellenanyag-készítmény burokkal rendelkező és burok nélküli vírusokat tekintve biztonságos, antikomplementer aktivitása ≤ 1 CH 50/mg fehérje, és folyadék formájában, 2-8°C-on tárolva legalább 6 hónapig stabil.

12. A 11. igénypont szerinti ellenanyag-készítmény, melynek proteolitikus aktivitása kisebb, mint 8E/l.

13. A 10-12. igénypontok bármelyike szerinti ellenanyag-készítmény, ahol

- (i) a készítmény legalább 20% IgM-et tartalmaz, és előnyösen 20-30% IgM-et tartalmaz, és/vagy
- (ii) a készítmény tartalmaz továbbá stabilizálószer, előnyösen ahol a stabilizálószer glicin, és/vagy
- (iii) ahol a készítmény 2-8°C-on tárolva legalább 2 évig stabil, és/vagy
- (iv) ahol az immunglobulinok nincsenek kémiai módon módosítva, vagy ahol a készítmény előállítási eljárása során nem lett kezelve β -propiolaktonnal vagy hozzáadott proteázzal, például pepszinnel.

14. IgM-et tartalmazó immunglobulin-készítmény, amely az 1-3. igénypontok bármelyike szerinti eljárással előállítható.

15. A 10-13. igénypontok bármelyike szerinti ellenanyag-készítmény gyógyászatban történő alkalmazására.

16. A 15. igénypont szerinti ellenanyag-készítmény immunológiai rendellenesség vagy baktériumfertőzés kezelésében történő alkalmazásra, előnyösen ahol az immunológiai rendellenesség IgM-hiányos rendellenesség.

17. A 10-13. igénypontok bármelyike szerinti ellenanyag-készítmény alkalmazása immunológiai rendellenesség és baktériumfertőzés kezelésére szolgáló gyógyszer előállítására.