

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199738856 B2**
(10) Patent No. **708572**

(54) Title
Preparation of (pichia methanolica) auxotrophic mutants

(51)⁶ International Patent Classification(s)
C12N 015/00

(21) Application No: 199738856 (22) Application Date: 1997 .07 .14

(87) WIPO No: W098/02536

(30) Priority Data

(31) Number	(32) Date	(33) Country
08/683500	1996 .07 .17	US
08/703808	1996 .08 .26	US

(43) Publication Date : 1998 .02 .09

(43) Publication Journal Date : 1998 .04 .02

(44) Accepted Journal Date : 1999 .08 .05

(71) Applicant(s)
ZymoGenetics, Inc.

(72) Inventor(s)
Christopher K. Raymond

(74) Agent/Attorney
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

OPI DATE 09/02/98 APPLN. ID 38856/97
AOJP DATE 02/04/98 PCT NUMBER PCT/US97/12582



AU9738856

CT)

(51) International Patent Classification 6 :
C12N 15/00

A2

(11) International Publication Number: WO 98/02536

(43) International Publication Date: 22 January 1998 (22.01.98)

(21) International Application Number: PCT/US97/12582

(22) International Filing Date: 14 July 1997 (14.07.97)

(30) Priority Data:

08/683,500 17 July 1996 (17.07.96) US
08/703,808 26 August 1996 (26.08.96) US

(71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US).

(72) Inventor: RAYMOND, Christopher, K.; 3103 N.E. 85th Street, Seattle, WA 98103 (US).


(74) Agent: PARKER, Gary, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: PREPARATION OF *PICHA METHANOLICA* AUXOTROPHIC MUTANTS



(57) Abstract

Methods for preparing *Pichia methanolica* cells having auxotrophic mutations are disclosed. The methods comprise the steps of: (a) exposing *P. methanolica* cells to mutagenizing conditions; (b) culturing the cells from step (a) in a rich medium to allow mutations to become established and replicated in at least a portion of the cells; (c) culturing the cells from step (b) in a culture medium deficient in assimilable nitrogen to deplete cellular nitrogen stores; (d) culturing the cells from step (c) in a defined culture medium comprising an inorganic nitrogen source and an amount of nystatin sufficient to kill growing *P. methanolica* cells to select for cells having a deficiency in a nutritional gene; and (e) culturing the selected cells from step (d) in a rich culture medium.

Description5 PREPARATION OF *PICHIA METHANOLICA* AUXOTROPHIC MUTANTSBackground of the Invention

10 Methylophilic yeasts are those yeasts that are able to utilize methanol as a sole source of carbon and energy. Species of yeasts that have the biochemical pathways necessary for methanol utilization are classified in four genera, *Hansenula*, *Pichia*, *Candida*, and *Torulopsis*. These genera are somewhat artificial, having been based on cell morphology and growth characteristics, and do not reflect close genetic relationships (Billon-Grand, *Mycotaxon* 35:201-204, 1989; Kurtzman, *Mycologia* 84:72-76, 1992). Furthermore, not all species within these genera are capable of utilizing methanol as a source of carbon and energy. As a consequence of this classification, there are great differences in physiology and metabolism between individual species of a genus.

15 Methylophilic yeasts are attractive candidates for use in recombinant protein production systems. Some methylophilic yeasts have been shown to grow rapidly to high biomass on minimal defined media. Certain genes of methylophilic yeasts are tightly regulated and highly expressed under induced or de-repressed conditions, suggesting that promoters of these genes might be useful for producing polypeptides of commercial value. See, for example, Faber et al., *Yeast* 11:1331, 1995; Romanos et al., *Yeast* 8:423, 1992; and Cregg et al., *Bio/Technology* 11:905, 1993.

20 Development of methylophilic yeasts as hosts for use in recombinant production systems has been slow, due in part to a lack of suitable materials (e.g., promoters, selectable markers, and mutant host cells) and methods (e.g., transformation techniques). The most highly developed methylophilic host systems utilize *Pichia pastoris* and *Hansenula polymorpha* (Faber et al., *Curr. Genet.* 25:305-310, 1994; Cregg et al., *ibid.*; Romanos et al., *ibid.*; U.S. Patent No. 4,855,242; U.S. Patent No. 4,857,467; U.S. Patent No. 4,879,231; and U.S. Patent No. 4,929,555).

25 There remains a need in the art for methods of transforming additional species of methylophilic yeasts and for using the transformed cells to produce polypeptides of economic importance, including industrial enzymes and pharmaceutical proteins. The present invention provides compositions and methods useful in these processes as well as other, related advantages.

Summary of the Invention

35 The present invention provides methods for preparing *Pichia methanolica* cells having an auxotrophic mutation. The methods comprise the steps of (a) exposing *P. methanolica* cells to mutagenizing conditions; (b) culturing the cells from step (a) in a rich medium to allow mutations to become established and replicated in at least a portion of the cells; (c) culturing the cells from step (b) in a culture medium deficient in assimilable nitrogen to deplete cellular nitrogen stores; (d) culturing the cells from step (c) in a defined culture medium comprising an inorganic nitrogen source and an amount of nystatin sufficient to kill growing *P. methanolica* cells to select

40

for cells having a deficiency in a nutritional gene; and (e) culturing the selected cells from step (d) in a rich culture medium. Within one embodiment of the invention, the selected cells from step (e) are replica plated to a defined medium and cultured to confirm the presence of an auxotrophic mutation. Within another embodiment, the selected cells are auxotrophic for adenine. Within a related embodiment, the selected cells are deficient in phosphoribosyl-5-aminoimidazole carboxylase. Within additional embodiments the mutagenizing conditions comprise exposure to ultraviolet light or exposure to a chemical mutagen. Within a further embodiment, the inorganic nitrogen source comprises ammonium ions.

These and other aspects of the invention will become evident upon reference to the following detailed description and the attached drawings.

Brief Description of the Drawings

Fig. 1 illustrates the effects of field strength and pulse duration on electroporation efficiency of *P. methanolicus*.

Fig. 2 is a schematic diagram of a recombination event between plasmid pCZR140 and *P. methanolicus* genomic DNA.

Fig. 3 is a schematic diagram of a recombination event between plasmid pCZR137 and *P. methanolicus* genomic DNA.

Detailed Description of the Invention

Prior to setting forth the invention in more detail, it will be useful to define certain terms used herein:

A "DNA construct" is a DNA molecule, either single- or double-stranded, that has been modified through human intervention to contain segments of DNA combined and juxtaposed in an arrangement not existing in nature.

"Early log phase growth" is that phase of cellular growth in culture when the cell concentration is from 2×10^6 cells/ml to 8×10^6 cells/ml.

"Heterologous DNA" refers to a DNA molecule, or a population of DNA molecules, that does not exist naturally within a given host cell. DNA molecules heterologous to a particular host cell may contain DNA derived from the host cell species so long as that host DNA is combined with non-host DNA. For example, a DNA molecule containing a non-host DNA segment encoding a polypeptide operably linked to a host DNA segment comprising a transcription promoter is considered to be a heterologous DNA molecule.

A "higher eukaryotic" organism is a multicellular eukaryotic organism. The term encompasses both plants and animals.

"Integrative transformants" are cells into which has been introduced heterologous DNA, wherein the heterologous DNA has become integrated into the genomic DNA of the cells.

"Linear DNA" denotes DNA molecules having free 5' and 3' ends, that is non-circular DNA molecules. Linear DNA can be prepared from closed circular DNA molecules, such as plasmids, by enzymatic digestion or physical disruption.

The term "operably linked" indicates that DNA segments are arranged so that they function in concert for their intended purposes. e.g., transcription initiates in the promoter and proceeds through the coding segment to the terminator.

5 The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes. Sequence elements within promoters that function in the initiation of transcription are often characterized by consensus nucleotide sequences. These promoter elements include RNA polymerase binding sites; TATA sequences; CAAT sequences; differentiation-specific elements (DSEs; McGehee et al., Mol. Endocrinol. 7:551-560, 1993); cyclic AMP response elements (CREs); 10 serum response elements (SREs; Treisman, Seminars in Cancer Biol. 1:47-58, 1990); glucocorticoid response elements (GREs); and binding sites for other transcription factors, such as CRE/ATF (O'Reilly et al., J. Biol. Chem. 267:19938-19943, 1992), AP2 (Ye et al., J. Biol. Chem. 269:25728-25734, 1994), SP1, cAMP response element binding protein (CREB; Loeken, Gene Expr. 3:253-264, 1993) and octamer factors. See, in general, Watson et al., eds., Molecular Biology of the Gene, 4th ed., The Benjamin/Cummings Publishing Company, Inc., Menlo Park, CA, 1987; and Lemaigre and Rousseau, Biochem. J. 303:1-14, 1994.

A "repressing carbon source" is a metabolizable, carbon-containing compound that, when not limited, suppresses the expression in an organism of genes required for the catabolism of 20 other carbon sources. By "limited" is meant that the carbon source is unavailable or becomes available at such a rate that it is immediately consumed and therefore the prevailing concentration of that carbon source in an organism's environment is effectively zero. Repressing carbon sources that can be used within the present invention include hexoses and ethanol. Glucose is particularly preferred.

25 "Rich" culture media are those culture media that are based on complex sources of nutrients, typically cell or tissue extracts or protein hydrolysates. Rich media will vary in composition from batch to batch due to variations in the composition of the nutrient sources.

As noted above, the present invention provides methods for preparing *Pichia methanolica* cells having an auxotrophic mutation. Auxotrophic mutants of *P. methanolica* can be 30 transformed with both homologous DNA (DNA from the host species) and heterologous DNA, and the resulting transformants can be used within a large number of diverse biological applications. The mutant cells of the present invention are particularly well suited for transformation with heterologous DNA, which transformed cells can be used for the production of polypeptides and proteins, including polypeptides and proteins of higher organisms, including humans. Auxotrophic 35 *P. methanolica* cells can be transformed with other DNA molecules, including DNA libraries and synthetic DNA molecules. The invention thus provides host cells that can be used to express genetically diverse libraries to produce products that are screened for novel biological activities, can be engineered for use as targets for the screening of compound libraries, and can be genetically modified to enhance their utility within other processes.

40 Cells to be transformed with heterologous DNA will commonly have a mutation that can be complemented by a gene (a "selectable marker") on the heterologous DNA molecule. This

selectable marker allows the transformed cells to grow under conditions in which untransformed cells cannot multiply ("selective conditions"). The general principles of selection are well known in the art. Commonly used selectable markers are genes that encode enzymes required for the synthesis of amino acids or nucleotides. Cells having mutations in these genes (auxotrophic mutants) cannot grow in media lacking the specific amino acid or nucleotide unless the mutation is complemented by the selectable marker. Use of such "selective" culture media ensures the stable maintenance of the heterologous DNA within the host cell. A preferred selectable marker of this type for use in *Pichia methanolica* is a *P. methanolica* ADE2 gene, which encodes phosphoribosyl-5-aminoimidazole carboxylase (AIRC; EC 4.1.1.21). The ADE2 gene, when transformed into an ade2 host cell, allows the cell to grow in the absence of adenine. The coding strand of a representative *P. methanolica* ADE2 gene sequence is shown in SEQ ID NO:1. The sequence illustrated includes 1006 nucleotides of 5' non-coding sequence and 442 nucleotides of 3' non-coding sequence, with the initiation ATG codon at nucleotides 1007-1009. Within a preferred embodiment of the invention, a DNA segment comprising nucleotides 407-2851 is used as a selectable marker, although longer or shorter segments could be used as long as the coding portion is operably linked to promoter and terminator sequences. Those skilled in the art will recognize that this and other sequences provided herein represent single alleles of the respective genes, and that allelic variation is expected to exist. Any functional ADE2 allele can be used within the present invention. Other nutritional markers that can be used within the present invention include the *P. methanolica* ADE1, HIS3, and LEU2 genes, which allow for selection in the absence of adenine, histidine, and leucine, respectively. Heterologous genes, such as genes from other fungi, can also be used as selectable markers. For large-scale, industrial processes where it is desirable to minimize the use of methanol, it is preferred to use host cells in which both methanol utilization genes (*AUG1* and *AUG2*) are deleted. For production of secreted proteins, host cells deficient in vacuolar protease genes (*PEP4* and *PRB1*) are preferred. Gene-deficient mutants can be prepared by known methods, such as site-directed mutagenesis. *P. methanolica* genes can be cloned on the basis of homology with their counterpart *Saccharomyces cerevisiae* genes. The ADE2 gene disclosed herein was given its designation on the basis of such homology.

Strains of *Pichia methanolica* are available from the American Type Culture Collection (Rockville, MD) and other repositories, and can be used as starting materials within the present invention. To prepare auxotrophic mutants of *P. methanolica*, cells are first exposed to mutagenizing conditions, i.e. environmental conditions that cause genetic mutations in the cells. Methods for mutagenizing cells are well known in the art and include chemical treatment, exposure to ultraviolet light, exposure to x-rays, and retroviral insertional mutagenesis. Chemical mutagens include ethylmethane sulfonate (EMS), *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine, 2-methoxy-6-chloro-9-[3-(ethyl-2-chloroethyl)aminopropylamino]acridine 2HCl, 5-bromouracil, acridine, and aflatoxin. See Lawrence, Methods Enzymol. 194:273-281, 1991. The proportion of mutagenized cells obtained is a function of the strength or amount of mutagenizing agent to which the cells are exposed. A low level of mutagen produces a small proportion of mutant cells. Higher levels of mutagen produce a higher proportion of mutant cells, but also kill more cells. It is therefore necessary to balance mutagenesis with killing so that a reasonable number of mutant cells is

obtained. Balancing is generally done empirically by exposing cells to different conditions to establish a killing curve. In general, the cells are exposed to mutagenizing conditions and cultured for one day, after which they are tested for viability according to standard assay methods. Within the present invention, it is preferred to use a level of mutagenesis that results in 20-50% mortality, although one skilled in the art will recognize that this value can be adjusted as necessary, for example if working with a very large number of cells.

Mutagenized cells are then cultured in a rich medium to allow mutations to become established and replicated in at least a portion of the cell population. This step allows cells in which the genome has been altered to replicate the mutation and pass it on to their progeny, thereby establishing the mutation within the population.

The cells are then transferred to a culture medium deficient in assimilable nitrogen so that cellular nitrogen stores are depleted. By "deficient in assimilable nitrogen" it is meant that the medium lacks an amount of nitrogen sufficient to support growth of the cells. Depletion of cellular nitrogen stores will generally require about 12 to 24 hours of incubation, with 16 hours being sufficient under common conditions. Following depletion of nitrogen stores, the cells are cultured in a defined culture medium comprising an inorganic nitrogen source and an amount of an antifungal antibiotic sufficient to kill growing *P. methanolica* cells. A preferred antibiotic is nystatin (mycostatin). Preferred inorganic nitrogen sources are those comprising ammonium ions, such as ammonium sulfate. In general, the medium will contain 10-200 mM ammonium, preferably about 60 mM ammonium. Nystatin is included at a concentration of 0.1 to 100 mg/l, preferably 0.5 to 20 mg/L, more preferably about 2 mg/L (10 units/L). Treatment with nystatin is carried out for ten minutes to six hours, preferably about 1 hour. Those skilled in the art will recognize that the actual antibiotic concentration and exposure time required to kill prototrophic cells can be readily determined empirically, and certain adjustments may be necessary to compensate for variations in specific activity between individual batches of antibiotic. By depleting cellular nitrogen stores and then culturing the cells in a defined medium containing an inorganic nitrogen source and antibiotic, cells that are auxotrophic for amino acid or nucleotide biosynthesis remain alive because they cannot grow in the defined medium. Growing cells are killed by the antibiotic. Following the antibiotic treatment, the cells are transferred to a rich culture medium.

Auxotrophic mutations are confirmed and characterized by determining the nutrient requirements of the treated cells. Replica plating is commonly used for this determination. Cells are plated on both rich medium and media lacking specific nutrients. Cells that do not grow on particularly plates are auxotrophic for the missing nutrient. Complementation analysis can be used for further characterization.

Heterologous DNA can be introduced into *P. methanolica* cells by any of several known methods, including lithium transformation (Hiep et al., Yeast 9:1189-1197, 1993; Tarutina and Tolstorukov, Abst. of the 15th International Specialized Symposium on Yeasts, Riga (USSR), 1991, 137; Ito et al., J. Bacteriol. 153:163, 1983; Bogdanova et al., Yeast 11:343, 1995), spheroplast transformation (Beggs, Nature 275:104, 1978; Hinnen et al., Proc. Natl. Acad. Sci. USA 75:1929, 1978; Cregg et al., Mol. Cell. Biol. 5:3376, 1985), freeze-thaw polyethylene glycol transformation (Pichia Expression Kit Instruction Manual, Invitrogen Corp., San Diego, CA. Cat. No. K1710-01).

or electroporation, the latter method being preferred. Electroporation is the process of using a pulsed electric field to transiently permeabilize cell membranes, allowing macromolecules, such as DNA, to pass into cells. Electroporation has been described for use with mammalian (e.g., Neumann et al., EMBO J. 1:841-845, 1982) and fungal (e.g., Meilhoc et al., Bio/Technology 8:223-227, 1990) host cells. However, the actual mechanism by which DNA is transferred into the cells is not well understood. For transformation of *P. methanolica*, it has been found that electroporation is surprisingly efficient when the cells are exposed to an exponentially decaying, pulsed electric field having a field strength of from 2.5 to 4.5 kV/cm and a time constant (τ) of from 1 to 40 milliseconds. The time constant τ is defined as the time required for the initial peak voltage V_0 to drop to a value of V_0/e . The time constant can be calculated as the product of the total resistance and capacitance of the pulse circuit, i.e., $\tau = R \times C$. Typically, resistance and capacitance are either preset or may be selected by the user, depending on the electroporation equipment selected. In any event, the equipment is configured in accordance with the manufacturer's instructions to provide field strength and decay parameters as disclosed above. Electroporation equipment is available from commercial suppliers (e.g., BioRad Laboratories, Hercules, CA).

DNA molecules for use in transforming *P. methanolica* will commonly be prepared as double-stranded, circular plasmids, which are preferably linearized prior to transformation. For polypeptide or protein production, the DNA molecules will include, in addition to the selectable marker disclosed above, an expression cassette comprising a transcription promoter, a DNA segment (e.g., a cDNA) encoding the polypeptide or protein of interest, and a transcription terminator. These elements are operably linked to provide for transcription of the DNA segment of interest. It is preferred that the promoter and terminator be that of a *P. methanolica* gene. Useful promoters include those from constitutive and methanol-inducible promoters. Promoter sequences are generally contained within 1.5 kb upstream of the coding sequence of a gene, often within 1 kb or less. In general, regulated promoters are larger than constitutive promoters due the presence of regulatory elements. Methanol-inducible promoters, which include both positive and negative regulatory elements, may extend more than 1 kb upstream from the initiation ATG. Promoters are identified by function and can be cloned according to known methods.

A particularly preferred methanol-inducible promoter is that of a *P. methanolica* alcohol utilization gene. A representative coding strand sequence of one such gene, *AUG1*, is shown in SEQ ID NO:2. Within SEQ ID NO:2, the initiation ATG codon is at nucleotides 1355-1357. Nucleotides 1-23 of SEQ ID NO:2 are non-*AUG1* polylinker sequence. It is particularly preferred to utilize as a promoter a segment comprising nucleotides 24-1354 of SEQ ID NO:2, although additional upstream sequence can be included. *P. methanolica* contains a second alcohol utilization gene, *AUG2*, the promoter of which can be used within the present invention. A partial DNA sequence of one *AUG2* clone is shown in SEQ ID NO:9. *AUG2* promoter segments used within the present invention will generally comprise nucleotides 91-169 of SEQ ID NO:9, although small truncations at the 3' end would not be expected to negate promoter function. Other useful promoters include those of the dihydroxyacetone synthase (DHAS), formate dehydrogenase (FMD), and catalase (CAT) genes. Genes encoding these enzymes from other species have been described, and their sequences are available (e.g., Janowicz et al., Nuc. Acids Res. 13:2043, 1985; Hollenberg

and Janowicz. EPO publication 0 299 108; Didion and Roggenkamp. FEBS Lett. 303:113, 1992). Genes encoding these proteins can be cloned by using the known sequences as probes, or by aligning known sequences, designing primers based on the alignment, and amplifying *P. methanolica* DNA by the polymerase chain reaction (PCR).

5 Constitutive promoters are those that are not activated or inactivated by environmental conditions; they are always transcriptionally active. Preferred constitutive promoters for use within the present invention include those from glyceraldehyde-3-phosphate dehydrogenase, triose phosphate isomerase, and phosphoglycerate kinase genes of *P. methanolica*. These genes can be cloned by complementation in a host cell, such as a *Saccharomyces cerevisiae* cell, having a
10 mutation in the counterpart gene. Mutants of this type are well known in the art. See, for example, Kawasaki and Fraenkel, Biochem. Biophys. Res. Comm. 108:1107-1112, 1982; McKnight et al., Cell 46:143-147, 1986; Aguilera and Zimmermann, Mol. Gen. Genet. 202:83-89, 1986.

The DNA molecules will further include a selectable marker to allow for identification, selection, and maintenance of transformants. The DNA molecules may further
15 contain additional elements, such as an origin of replication and a selectable marker that allow amplification and maintenance of the DNA in an alternate host (e.g., *E. coli*). To facilitate integration of the DNA into the host chromosome, it is preferred to have the entire expression segment, comprising the promoter--gene of interest--terminator plus selectable marker, flanked at both ends by host DNA sequences. This is conveniently accomplished by including 3' untranslated
20 DNA sequence at the downstream end of the expression segment and relying on the promoter sequence at the 5' end. When using linear DNA, the expression segment will be flanked by cleavage sites to allow for linearization of the molecule and separation of the expression segment from other sequences (e.g., a bacterial origin of replication and selectable marker). Preferred such cleavage sites are those that are recognized by restriction endonucleases that cut infrequently within
25 a DNA sequence, such as those that recognize 8-base target sequences (e.g., Not I).

Proteins that can be produced in *P. methanolica* include proteins of industrial and pharmaceutical interest. Such proteins include higher eukaryotic proteins from plants and animals, particularly vertebrate animals such as mammals, although certain proteins from microorganisms are also of great value. Proteins that can be prepared using methods of the present invention
30 include enzymes such as lipases, cellulases, and proteases; enzyme inhibitors, including protease inhibitors; growth factors such as platelet derived growth factor, fibroblast growth factors, and epidermal growth factor; cytokines such as erythropoietin and thrombopoietin; and hormones such as insulin, leptin, and glucagon.

For production of polypeptides, *P. methanolica* cells are cultured in a medium
35 comprising adequate sources of carbon, nitrogen and trace nutrients at a temperature of about 25°C to 35°C. Liquid cultures are provided with sufficient aeration by conventional means, such as shaking of small flasks or sparging of fermentors. A preferred culture medium is YEPD (Table 1). The cells may be passaged by dilution into fresh culture medium or stored for short periods on plates under refrigeration. For long-term storage, the cells are preferably kept in a 50% glycerol
40 solution at -70°C.

Table 1

YEPD

2% D-glucose

2% Bacto™ Peptone (Difco Laboratories, Detroit, MI)

1% Bacto™ yeast extract (Difco Laboratories)

5 0.004% adenine

0.006% L-leucine

Table 1. continuedADE D

- 5 0.056% -Ade -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution

ADE DS

- 10 0.056% -Ade -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution
18.22% D-sorbitol

LEU D

- 15 0.052% -Leu -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution

HIS D

- 20 0.052% -His -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution

URA D

- 25 0.056% -Ura -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution

URA DS

- 30 0.056% -Ura -Trp -Thr powder
0.67% yeast nitrogen base without amino acids
2% D-glucose
0.5% 200X tryptophan, threonine solution
18.22% D-sorbitol

Table 1, continued-Leu -Trp -Thr powder

powder made by combining 4.0 g adenine, 3.0 g arginine, 5.0 g aspartic acid, 2.0 g histidine, 6.0 g isoleucine, 4.0 g lysine, 2.0 g methionine, 6.0 g phenylalanine, 5.0 g serine, 5.0 g tyrosine, 4.0 g uracil, and 6.0 g valine (all L- amino acids)

-His -Trp -Thr powder

powder made by combining 4.0 g adenine, 3.0 g arginine, 5.0 g aspartic acid, 6.0 g isoleucine, 8.0 g leucine, 4.0 g lysine, 2.0 g methionine, 6.0 g phenylalanine, 5.0 g serine, 5.0 g tyrosine, 4.0 g uracil, and 6.0 g valine (all L- amino acids)

-Ura -Trp -Thr powder

powder made by combining 4.0 g adenine, 3.0 g arginine, 5.0 g aspartic acid, 2.0 g histidine, 6.0 g isoleucine, 8.0 g leucine, 4.0 g lysine, 2.0 g methionine, 6.0 g phenylalanine, 5.0 g serine, 5.0 g tyrosine, and 6.0 g valine (all L- amino acids)

-Ade -Trp -Thr powder

powder made by combining 3.0 g arginine, 5.0 g aspartic acid, 2.0 g histidine, 6.0 g isoleucine, 8.0 g leucine, 4.0 g lysine, 2.0 g methionine, 6.0 g phenylalanine, 5.0 g serine, 5.0 g tyrosine, 4.0 g uracil, and 6.0 g valine (all L- amino acids)

200X tryptophan, threonine solution

3.0% L-threonine, 0.8% L-tryptophan in H₂O

For plates, add 1.8% Bacto™ agar (Difco Laboratories)

Electroporation of *P. methanolicus* is preferably carried out on cells in early log phase growth. Cells are streaked to single colonies on solid media, preferably solid YEPD. After about 2 days of growth at 30°C, single colonies from a fresh plate are used to inoculate the desired volume of rich culture media (e.g., YEPD) to a cell density of about 5 - 10 x 10⁵ cells/ml. Cells are incubated at about 25 - 35°C, preferably 30°C, with vigorous shaking, until they are in early log phase. The cells are then harvested, such as by centrifugation at 3000 x g for 2-3 minutes, and resuspended. Cells are made electrocompetent by reducing disulfide bonds in the cell walls, equilibrating them in an ionic solution that is compatible with the electroporation conditions, and chilling them. Cells are typically made electrocompetent by incubating them in a buffered solution at pH 6-8 containing a reducing agent, such as dithiothreitol (DTT) or β-mercaptoethanol (BME), to reduce cell wall proteins to facilitate subsequent uptake of DNA. A preferred incubation buffer in this regard is a fresh solution of 50 mM potassium phosphate buffer, pH 7.5, containing 25 mM DTT. The cells are incubated in this buffer (typically using one-fifth the original culture volume) at about 30°C for about 5 to 30 minutes, preferably about 15 minutes. The cells are then harvested and washed in a suitable electroporation buffer, which is used ice-cold. Suitable buffers in this regard include pH 6-8 solutions containing a weak buffer, divalent cations (e.g., Mg⁺⁺, Ca⁺⁺) and an osmotic stabilizer (e.g., a sugar). After washing, the cells are resuspended in a small volume of the buffer, at which time they are electrocompetent and can be used directly or aliquotted and stored frozen (preferably at -70°C). A preferred electroporation buffer is STM (270 mM sucrose, 10 mM Tris, pH 7.5, 1 mM MgCl₂). Within a preferred protocol, the cells are subjected to two washes, first

in the original culture volume of ice-cold buffer, then in one-half the original volume. Following the second wash, the cells are harvested and resuspended, typically using about 3-5 ml of buffer for an original culture volume of 200 ml.

Electroporation is carried out using a small volume of electrocompetent cells (typically about 100 μ l) and up to one-tenth volume of linear DNA molecules. For example, 0.1 ml of cell suspension in a buffer not exceeding 50 mM in ionic strength is combined with 0.1-10 μ g of DNA (vol. $\leq$ 10 μ l). This mixture is placed in an ice-cold electroporation cuvette and subjected to a pulsed electric field of from 2.5 to 4.5 kV/cm, preferably about 3.75 kV/cm, and a time constant of from 1 to 40 milliseconds, preferably 10-30 milliseconds, more preferably 15-25 milliseconds, most preferably about 20 milliseconds, with exponential decay. The actual equipment settings used to achieve the desired pulse parameters will be determined by the equipment used. When using a BioRad (Hercules, CA) Gene Pulser™ electroporator with a 2 mm electroporation cuvette, resistance is set at 600 ohms or greater, preferably "infinite" resistance, and capacitance is set at 25 μ F to obtain the desired field characteristics. After being pulsed, the cells are diluted approximately 10X into 1 ml of YEPD broth and incubated at 30°C for one hour.

The cells are then harvested and plated on selective media. Within a preferred embodiment, the cells are washed once with a small volume (equal to the diluted volume of the electroporated cells) of 1X yeast nitrogen base (6.7 g/L yeast nitrogen base without amino acids; Difco Laboratories, Detroit, MI), and plated on minimal selective media. Cells having an *ade2* mutation that have been transformed with an *ADE2* selectable marker can be plated on a minimal medium that lacks adenine, such as ADE D (Table 1) or ADE DS (Table 1). In a typical procedure, 250 μ l aliquots of cells are plated on 4 separate ADE D or ADE DS plates to select for Ade⁺ cells.

P. methanolicus recognizes certain infrequently occurring sequences, termed autonomously replicating sequences (ARS), as origins of DNA replication, and these sequences may fortuitously occur within a DNA molecule used for transformation, allowing the transforming DNA to be maintained extrachromosomally. However, integrative transformants are generally preferred for use in protein production systems. Integrative transformants have a profound growth advantage over ARS transformants on selective media containing sorbitol as a carbon source, thereby providing a method for selecting integrative transformants from among a population of transformed cells. ARS sequences have been found to exist in the *ADE2* gene and, possibly, the *AUG1* gene of *P. methanolicus*. *ade2* host cells of *Pichia methanolicus* transformed with an *ADE2* gene can thus become Ade⁺ by at least two different modes. The ARS within the *ADE2* gene allows unstable extrachromosomal maintenance of the transforming DNA (Hiep et al., *Yeast* 9:1189-1197, 1993). Colonies of such transformants are characterized by slower growth rates and pink color due to prolific generation of progeny that are Ade⁻. Transforming DNA can also integrate into the host genome, giving rise to stable transformants that grow rapidly, are white, and that fail to give rise to detectable numbers of Ade⁻ progeny. ADE D plates allow the most rapid growth of transformed cells, and unstable and stable transformants grow at roughly the same rates. After 3-5 days of incubation on ADE D plates at 30°C stable transformant colonies are white and roughly twice the size of unstable, pink transformants. ADE DS plates are more selective for stable transformants, which form large ($\approx$ 5 mm) colonies in 5-7 days, while unstable (ARS-maintained)

colonies are much smaller (≈ 1 mm). The more selective ADE DS media is therefore preferred for the identification and selection of stable transformants. For some applications, such as the screening of genetically diverse libraries for rare combinations of genetic elements, it is sometimes desirable to screen large numbers of unstable transformants, which have been observed to outnumber stable transformants by a factor of roughly 100. In such cases, those skilled in the art will recognize the utility of plating transformant cells on less selective media, such as ADE D.

Integrative transformants are preferred for use in protein production processes. Such cells can be propagated without continuous selective pressure because DNA is rarely lost from the genome. Integration of DNA into the host chromosome can be confirmed by Southern blot analysis. Briefly, transformed and untransformed host DNA is digested with restriction endonucleases, separated by electrophoresis, blotted to a support membrane, and probed with appropriate host DNA segments. Differences in the patterns of fragments seen in untransformed and transformed cells are indicative of integrative transformation. Restriction enzymes and probes can be selected to identify transforming DNA segments (e.g., promoter, terminator, heterologous DNA, and selectable marker sequences) from among the genomic fragments.

Differences in expression levels of heterologous proteins can result from such factors as the site of integration and copy number of the expression cassette and differences in promoter activity among individual isolates. It is therefore advantageous to screen a number of isolates for expression level prior to selecting a production strain. A variety of suitable screening methods are available. For example, transformant colonies are grown on plates that are overlaid with membranes (e.g., nitrocellulose) that bind protein. Proteins are released from the cells by secretion or following lysis, and bind to the membrane. Bound protein can then be assayed using known methods, including immunoassays. More accurate analysis of expression levels can be obtained by culturing cells in liquid media and analyzing conditioned media or cell lysates, as appropriate. Methods for concentrating and purifying proteins from media and lysates will be determined in part by the protein of interest. Such methods are readily selected and practiced by the skilled practitioner.

For small-scale protein production (e.g., plate or shake flask production), *P. methanolicus* transformants that carry an expression cassette comprising a methanol-regulated promoter (such as the *AUG1* promoter) are grown in the presence of methanol and the absence of interfering amounts of other carbon sources (e.g., glucose). For small-scale experiments, including preliminary screening of expression levels, transformants may be grown at 30°C on solid media containing, for example, 20 g/L Bacto-agar (Difco), 6.7 g/L yeast nitrogen base without amino acids (Difco), 10 g/L methanol, 0.4 μ g/L biotin, and 0.56 g/L of -Ade -Thr -Trp powder. Because methanol is a volatile carbon source it is readily lost on prolonged incubation. A continuous supply of methanol can be provided by placing a solution of 50% methanol in water in the lids of inverted plates, whereby the methanol is transferred to the growing cells by evaporative transfer. In general, not more than 1 mL of methanol is used per 100-mm plate. Slightly larger scale experiments can be carried out using cultures grown in shake flasks. In a typical procedure, cells are cultivated for two days on minimal methanol plates as disclosed above at 30°C, then colonies are used to inoculate a small volume of minimal methanol media (6.7 g/L yeast nitrogen base without amino acids, 10 g/L

methanol. 0.4 µg/L biotin) at a cell density of about 1×10^6 cells/ml. Cells are grown at 30°C. Cells growing on methanol have a high oxygen requirement, necessitating vigorous shaking during cultivation. Methanol is replenished daily (typically 1/100 volume of 50% methanol per day).

For production scale culturing, fresh cultures of high producer clones are prepared in shake flasks. The resulting cultures are then used to inoculate culture medium in a fermenter. Typically, a 500 ml culture in YEPD grown at 30°C for 1-2 days with vigorous agitation is used to inoculate a 5-liter fermenter. The cells are grown in a suitable medium containing salts, glucose, biotin, and trace elements at 28°C, pH 5.0, and >30% dissolved O₂. After the initial charge of glucose is consumed (as indicated by a decrease in oxygen consumption), a glucose/methanol feed is delivered into the vessel to induce production of the protein of interest. Because large-scale fermentation is carried out under conditions of limiting carbon, the presence of glucose in the feed does not repress the methanol-inducible promoter. The use of glucose in combination with methanol under glucose-limited conditions produces rapid growth, efficient conversion of carbon to biomass and rapid changes in physiological growth states, while still providing full induction of methanol-inducible gene promoters. In a typical fermentation run, a cell density of from about 80 to about 400 grams of wet cell paste per liter is obtained. "Wet cell paste" refers to the mass of cells obtained by harvesting the cells from the fermentor, typically by centrifugation of the culture.

The invention is further illustrated by the following non-limiting examples.

Examples

Example 1

P. methanolica cells (strain CBS6515 from American Type Culture Collection, Rockville, MD) were mutagenized by UV exposure. A killing curve was first generated by plating cells onto several plates at approximately 200-250 cells/plate. The plates were then exposed to UV radiation using a G8T5 germicidal lamp (Sylvania) suspended 25 cm from the surfaces of the plates for periods of time as shown in Table 2. The plates were then protected from visible light sources and incubated at 30°C for two days.

Table 2

Viable Cells

Time	Plate 1	Plate 2	Average
0 sec.	225	229	227
1 sec.	200	247	223
2 sec.	176	185	181
4 sec.	149	86	118
8 sec.	20	7	14
16 sec.	0	2	1

10

Large-scale mutagenesis was then carried out using a 2-second UV exposure to provide about 20% killing. Cells were plated at approximately 10^4 cells/plate onto eight YEPD plates that were supplemented with 100 mg/L each of uracil, adenine, and leucine, which were added to supplement the growth of potential auxotrophs having the cognate deficiencies. Following UV exposure the plates were wrapped in foil and incubated overnight at 30°C. The following day the colonies on the plates ($\sim 10^5$ total) were resuspended in water and washed once with water. An amount of cell suspension sufficient to give an OD₆₀₀ of 0.1 - 0.2 was used to inoculate 500 ml of minimal broth made with yeast nitrogen base without amino acids or ammonia, supplemented with 1% glucose and 400 µg/L biotin. The culture was placed in a 2.8 L baffled Bell flask and shaken vigorously overnight at 30°C. The following day the cells had reached an OD₆₀₀ of ~ 1.0 - 2.0. The cells were pelleted and resuspended in 500 ml of minimal broth supplemented with 5 g/L ammonium sulfate. The cell suspension was placed in a 2.8 L baffled Bell flask and shaken vigorously at 30°C for 6 hours. 50 ml of the culture was set aside in a 250-ml flask as a control, and to the remainder of the culture was added 1 mg nystatin (Sigma Chemical Co., St. Louis, MO) to select for auxotrophic mutants (Snow, *Nature* 211:206-207, 1966). The cultures were incubated with shaking for an additional hour. The control and nystatin-treated cells were then harvested by centrifugation and washed with water three times. The washed cells were resuspended to an OD₆₀₀ of 1.0 in 50% glycerol and frozen. Titering of nystatin-treated cells versus the control cells for colony forming units revealed that nystatin enrichment had decreased the number of viable cells by a factor of 10^4 .

30

10^{-2} dilutions of nystatin-treated cells were plated on 15 YEPD plates. Colonies were replica-plated onto minimal plates (2% agar, 1 x YNB, 2% glucose, 400 µg/L biotin). The frequency of auxotrophs was about 2 - 4%. Approximately 180 auxotrophic colonies were picked to YEPD + Ade, Leu, Ura plates and replica-plated to various dropout plates. All of the auxotrophs were Ade⁻. Of these, 30 were noticeably pink on dropout plates (LEU D, HIS D, etc.; see Table 1). Of the 30 pink mutants, 21 were chosen for further study; the remainder were either leaky for growth on ADE D plates or contaminated with wild-type cells.

35

The Ade⁻ mutants were then subjected to complementation analysis and phenotypic testing. To determine the number of loci defined by the mutants, all 21 mutants were mated to a single pink, Ade⁻ tester strain (strain #2). Mating was carried out by mixing cell suspensions (OD₆₀₀ = 1) and plating the mixtures in 10 µl aliquots on YEPD plates. The cells were then

40

replicated to SPOR media (0.5% Na acetate, 1% KCl, 1% glucose, 1% agar) and incubated overnight at 30°C. The cells were then replica-plated to ADE D plates for scoring of phenotype. As shown in Table 3, some combinations of mutants failed to give Ade⁺ colonies (possibly defining the same genetic locus as in strain #2), while others gave rise to numerous Ade⁺ colonies (possibly defining a separate genetic locus). Because mutant #3 gave Ade⁺ colonies when mated to #2, complementation testing was repeated with mutant #3. If the group of mutants defined two genetic loci, then all mutants that failed to give Ade⁺ colonies when mated to strain #2 should give Ade⁺ colonies when mated to #3. Results of the crosses are shown in Table 3.

10

Table 3

Mutant	x Mutant #2	x Mutant #3
#1	+	-
#3	+	-
#10	+	-
#15	+	-
#18	+	-
#24	+	-
#28	+	-
#30	+	-
#2	-	+
#6	-	+
#8	-	+
#9	-	+
#11	-	+
#17	-	+
#19	-	+
#20	-	+

Table 3. continued

#22	-	+
#27	-	+
#4	+	+
#12	+	+
#16	+	+

15

As shown in Table 3, most mutants fell into one of two groups, consistent with the idea that there are two adenine biosynthetic genes that, when missing, result in pink colonies on limiting adenine media. Three colonies (#4, #12, and #16) may either define a third locus or exhibit intragenic complementation. Two intensely pigmented mutants from each of the two complementation groups (#3 and #10; #6 and #11) were selected for further characterization. Additional analysis indicated that Ade⁻ was the only auxotrophy present in these strains.

A *P. methanolicus* clone bank was constructed in the vector pRS426, a shuttle vector comprising 2 μ and *S. cerevisiae* *URA3* sequences, allowing it to be propagated in *S. cerevisiae*. Genomic DNA was prepared from strain CBS6515 according to standard procedures. Briefly, cells were cultured overnight in rich media, spheroplasted with zymolyase, and lysed with SDS. DNA was precipitated from the lysate with ethanol and extracted with a phenol/chloroform mixture, then precipitated with ammonium acetate and ethanol. Gel electrophoresis of the DNA preparation showed the presence of intact, high molecular weight DNA and appreciable quantities of RNA. The DNA was partially digested with Sau 3A by incubating the DNA in the presence of a dilution series of the enzyme. Samples of the digests were analyzed by electrophoresis to determine the size distribution of fragments. DNA migrating between 4 and 12 kb was cut from the gel and extracted from the gel slice. The size-fractionated DNA was then ligated to pRS426 that had been digested with Bam HI and treated with alkaline phosphatase. Aliquots of the reaction mixture were electroporated in *E. coli* MC1061 cells using a BioRad Gene Pulser™ device as recommended by the manufacturer.

The genomic library was used to transform *S. cerevisiae* strain HBY21A (*ade2 ura3*) by electroporation (Becker and Guarente, Methods Enzymol. 194:182-187, 1991). The cells were resuspended in 1.2 M sorbitol, and six 300- μ l aliquots were plated onto ADE D, ADE DS, URA D and URA DS plates (Table 1). Plates were incubated at 30°C for 4-5 days. No Ade⁺ colonies were recovered on the ADE D or ADE DS plates. Colonies from the URA D and URA DS plates were replica-plated to ADE D plates, and two closely spaced, white colonies were obtained. These colonies were restreaked and confirmed to be Ura⁺ and Ade⁺. These two strains, designated Ade1 and Ade6, were streaked onto media containing 5 FOA (5 fluoro orotic acid; Sikorski and Boeke, Methods Enzymol. 194:302-318). Ura⁺ colonies were obtained, which were found to be Ade⁻ upon replica plating. These results indicate that the Ade⁺ complementing activity is genetically linked to the plasmid-borne URA3 marker. Plasmids obtained from yeast strains Ade1 and Ade6 appeared to be identical by restriction mapping as described below. These genomic clones were designated pADE1-1 and pADE1-6, respectively.

Total DNA was isolated from the HBY21A transformants Ade1 and Ade6 and used to transform *E. coli* strain MC1061 to Amp^R. DNA was prepared from 2 Amp^R colonies of Ade1 and 3 Amp^R colonies of Ade6. The DNA was digested with Pst I, Sca I, and Pst I + Sca I and analyzed by gel electrophoresis. All five isolates produced the same restriction pattern.

PCR primers were designed from the published sequence of the *P. methanolicus* *ADE2* gene (also known as *ADE1*; Hiep et al., Yeast 9:1251-1258, 1993). Primer 9080 (SEQ ID NO:3) was designed to prime at bases 406-429 of the *ADE2* DNA (SEQ ID NO:1), and primer 9079 (SEQ ID NO:4) was designed to prime at bases 2852-2829. Both primers included tails to introduce Avr II and Spe I sites at each end of the amplified sequence. The predicted size of the resulting PCR fragment was 2450 bp.

PCR was carried out using plasmid DNA from the five putative *ADE2* clones as template DNA. The 100 μ l reaction mixtures contained 1x Taq PCR buffer (Boehringer Mannheim, Indianapolis, IN), 10-100 ng of plasmid DNA, 0.25 mM dNTPs, 100 pmol of each primer, and 1 μ l Taq polymerase (Boehringer Mannheim). PCR was run for 30 cycles of 30 seconds at 94°C, 60

seconds at 50°C, and 120 seconds at 72°C. Each of the five putative ADE2 genomic clones yielded a PCR product of the expected size (2.4 kb). Restriction mapping of the DNA fragment from one reaction gave the expected size fragments when digested with Bgl II or Sal I.

5 The positive PCR reactions were pooled and digested with Spe I. Vector pRS426 was digested with Spe I and treated with calf intestinal phosphatase. Four µl of PCR fragment and 1 µl of vector DNA were combined in a 10 µl reaction mix using conventional ligation conditions. The ligated DNA was analyzed by gel electrophoresis. Spe I digests were analyzed to identify plasmids carrying a subclone of the ADE2 gene within pRS426. The correct plasmid was designated pCZR118.

10 Because the ADE2 gene in pCZR118 had been amplified by PCR, it was possible that mutations that disabled the functional character of the gene could have been generated. To test for such mutations, subclones with the desired insert were transformed singly into *Saccharomyces cerevisiae* strain HBY21A. Cells were made electrocompetent and transformed according to standard procedures. Transformants were plated on URA D and ADE D plates. Three phenotypic
15 groups were identified. Clones 1, 2, 11, and 12 gave robust growth of many transformants on ADE D. The transformation frequency was comparable to the frequency of Ura⁺ transformants. Clones 6, 8, 10, and 14 also gave a high efficiency of transformation to both Ura⁺ and Ade⁺, but the Ade⁺ colonies were somewhat smaller than those in the first group. Clone 3 gave many Ura⁺ colonies, but no Ade⁺ colonies, suggesting it carried a non-functional *ade2* mutation. Clones 1, 2, 11, and 12
20 were pooled.

To identify the *P. methanolicus ade2* complementation group, two representative mutants from each complementation group (#3 and #10; #6 and #11), which were selected on the basis of deep red pigmentation when grown on limiting adenine, were transformed with the cloned ADE gene. Two hundred ml cultures of early log phase cells were harvested by centrifugation at
25 3000 x g for 3 minutes and resuspended in 20 ml of fresh KD buffer (50 mM potassium phosphate buffer, pH 7.5, containing 25 mM DTT). The cells were incubated in this buffer at 30°C for 15 minutes. The cells were then harvested and resuspended in 200 ml of ice-cold STM (270 mM sucrose, 10 mM Tris, pH 7.5, 1 mM MgCl₂). The cells were harvested and resuspended in 100 ml of ice-cold STM. The cells were again harvested and resuspended in 3-5 ml of ice-cold STM. 100-
30 µl aliquots of electrocompetent cells from each culture were then mixed with Not I-digested pADE1-1 DNA. The cell/DNA mixture was placed in a 2 mm electroporation cuvette and subjected to a pulsed electric field of 5 kV/cm using a BioRad Gene Pulser™ set to 1000Ω resistance and capacitance of 25 µF. After being pulsed, the cells were diluted by addition of 1 ml YEPD and incubated at 30°C for one hour. The cells were then harvested by gentle centrifugation and
35 resuspended in 400 µl minimal selective media lacking adenine (ADE D). The resuspended samples were split into 200-µl aliquots and plated onto ADE D and ADE DS plates. Plates were incubated at 30°C for 4-5 days. Mutants #6 and #11 gave Ade⁺ transformants. No Ade⁺ transformants were observed when DNA was omitted, hence the two isolates appeared to define the *ade2* complementation group. The ADE2 sequence is shown in SEQ ID NO:1.

40

Example 2

The *P. methanolica* clone bank disclosed in Example 1 was used as a source for cloning the Alcohol Utilization Gene (*AUG1*). The clone bank was stored as independent pools, each representing about 200-250 individual genomic clones. 0.1 µl of "miniprep" DNA from each pool was used as a template in a polymerase chain reaction with PCR primers (8784, SEQ ID NO:5; 8787, SEQ ID NO:6) that were designed from an alignment of conserved sequences in alcohol oxidase genes from *Hansenula polymorpha*, *Candida boidini*, and *Pichia pastoris*. The amplification reaction was run for 30 cycles of 94°C, 30 seconds; 50°C, 30 seconds; 72°C, 60 seconds; followed by a 7 minute incubation at 72°C. One pool (#5) gave a ~600 bp band. DNA sequencing of this PCR product revealed that it encoded an amino acid sequence with ~70% sequence identity with the *Pichia pastoris* alcohol oxidase encoded by the *AOX1* gene and about 85% sequence identity with the *Hansenula polymorpha* alcohol oxidase encoded by the *MOX1* gene. The sequence of the cloned *AUG1* gene is shown in SEQ ID NO:2.

Sub-pools of pool #5 were analyzed by PCR using the same primers used in the initial amplification. One positive sub-pool was further broken down to identify a positive colony. This positive colony was streaked on plates, and DNA was prepared from individual colonies. Three colonies gave identical patterns after digestion with *Cla* I.

Restriction mapping of the genomic clone and PCR product revealed that the *AUG1* gene lay on a 7.5 kb genomic insert and that sites within the PCR fragment could be uniquely identified within the genomic insert. Because the orientation of the gene within the PCR fragment was known, the latter information provided the approximate location and direction of transcription of the *AUG1* gene within the genomic insert. DNA sequencing within this region revealed a gene with very high sequence similarity at the amino acid level to other known alcohol oxidase genes.

Example 3

ade2 mutant *P. methanolica* cells are transformed by electroporation essentially as disclosed above with an expression vector comprising the *AUG1* promoter and terminator, human GAD65 DNA (Karlsen et al., *Proc. Natl. Acad. Sci. USA* 88:8337-8341, 1991), and *ADE2* selectable marker. Colonies are patched to agar minimal methanol plates (10 to 100 colonies per 100-mm plate) containing 20 g/L Bacto™-agar (Difco), 6.7 g/L yeast nitrogen base without amino acids (Difco), 10 g/L methanol, and 0.4 µg/L biotin. The agar is overlaid with nitrocellulose, and the plates are inverted over lids containing 1 ml of 50% methanol in water and incubated for 3 to 5 days at 30°C. The membrane is then transferred to a filter soaked in 0.2 M NaOH, 0.1% SDS, 35 mM dithiothreitol to lyse the adhered cells. After 30 minutes, cell debris is rinsed from the filter with distilled water, and the filter is neutralized by rinsing it for 30 minutes in 0.1 M acetic acid.

The filters are then assayed for adhered protein. Unoccupied binding sites are blocked by rinsing in TTBS-NFM (20 mM Tris pH 7.4, 0.1% Tween 20, 160 mM NaCl, 5% powdered nonfat milk) for 30 minutes at room temperature. The filters are then transferred to a solution containing GAD6 monoclonal antibody (Chang and Gottlieb, *J. Neurosci.* 8:2123-2130, 1988), diluted 1:1000 in TTBS-NFM. The filters are incubated in the antibody solution with gentle agitation for at least one hour, then washed with TTBS (20 mM Tris pH 7.4, 0.1% Tween 20, 160 mM NaCl) two times for five minutes each. The filters are then incubated in goat anti-mouse

antibody conjugated to horseradish peroxidase (1 µg/ml in TTBS-NFM) for at least one hour, then washed three times, 5 minutes per wash with TTBS. The filters are then exposed to commercially available chemiluminescence reagents (ECL™; Amersham Inc., Arlington Heights, IL). Light generated from positive patches is detected on X-ray film.

- 5 To more accurately detect the level of GAD₆₅ expression, candidate clones are cultured in shake flask cultures. Colonies are grown for two days on minimal methanol plates at 30 °C as disclosed above. The colonies are used to inoculate 20 ml of minimal methanol media (6.7 g/L yeast nitrogen base without amino acids, 10 g/L methanol, 0.4 µg/L biotin) at a cell density of 1 x 10⁶ cells/ml. The cultures are grown for 1-2 days at 30°C with vigorous shaking. 0.2 ml of 50%
 10 methanol is added to each culture daily. Cells are harvested by centrifugation and suspended in ice-cold lysis buffer (20 mM Tris pH 8.0, 40 mM NaCl, 2 mM PMSF, 1 mM EDTA, 1 µg/ml leupeptin, 1 µg/ml pepstatin, 1 µg/ml aprotinin) at 10 ml final volume per 1 g cell paste. 2.5 ml of the resulting suspension is added to 2.5 ml of 400-600 micron, ice-cold, acid-washed glass beads in a 15-ml vessel, and the mixture is vigorously agitated for one minute, then incubated on ice for 1
 15 minute. The procedure is repeated until the cells have been agitated for a total of five minutes. Large debris and unbroken cells are removed by centrifugation at 1000 x g for 5 minutes. The clarified lysate is then decanted to a clean container. The cleared lysate is diluted in sample buffer (5% SDS, 8 M urea, 100 mM Tris pH 6.8, 10% glycerol, 2 mM EDTA, 0.01% bromphenol glue) and electrophoresed on a 4-20% acrylamide gradient gel (Novex, San Diego, CA). Proteins are
 20 blotted to nitrocellulose and detected with GAD6 antibody as disclosed above.

- Clones exhibiting the highest levels of methanol-induced expression of foreign protein in shake flask culture are more extensively analyzed under high cell density fermentation conditions. Cells are first cultivated in 0.5 liter of YEPD broth at 30°C for 1 - 2 days with vigorous agitation, then used to inoculate a 5-liter fermentation apparatus (e.g., BioFlow III; New Brunswick
 25 Scientific Co., Inc., Edison, NJ). The fermentation vessel is first charged with mineral salts by the addition of 57.8 g (NH₄)₂SO₄, 68 g KH₂PO₄, 30.8 g MgSO₄·7H₂O, 8.6 g CaSO₄·2H₂O, 2.0 g NaCl, and 10 ml antifoam (PPG). H₂O is added to bring the volume to 2.5 L, and the solution is autoclaved 40 minutes. After cooling, 350 ml of 50% glucose, 250 ml 10 X trace elements (Table 4), 25 ml of 200 µg/ml biotin, and 250 ml cell inoculum are added.

Table 4

10 X trace elements:

	FeSO ₄ ·7H ₂ O	100mM	27.8 g/L
	CuSO ₄ ·5H ₂ O	2mM	0.5 g/L
35	ZnCl ₂	8mM	1.09 g/L
	MnSO ₄ ·H ₂ O	8mM	1.35 g/L
	CoCl ₂ ·6H ₂ O	2mM	0.48 g/L
	Na ₂ MoO ₄ ·2H ₂ O	1mM	0.24 g/L
	H ₃ BO ₃	8mM	0.5 g/L
40	KI	0.5mM	0.08 g/L
	biotin		5mg/L

thiamine

0.5 g/L

Add 1-2 mls H₂SO₄ per liter to bring compounds into solution.

5 The fermentation vessel is set to run at 28°C, pH 5.0, and >30% dissolved O₂. The cells will consume the initial charge of glucose, as indicated by a sharp demand for oxygen during glucose consumption followed by a decrease in oxygen consumption after glucose is exhausted. After exhaustion of the initial glucose charge, a glucose-methanol feed supplemented with NH₄⁺ and trace elements is delivered into the vessel at 0.2% (w/v) glucose, 0.2% (w/v) methanol for 5 hours followed by 0.1% (w/v) glucose, 0.4% (w/v) methanol for 25 hours. A total of 550 grams of
10 methanol is supplied through one port of the vessel as pure methanol using an initial delivery rate of 12.5 ml/hr and a final rate of 25 ml/hr. Glucose is supplied through a second port using a 700 ml solution containing 175 grams glucose, 250 ml 10X trace elements, and 99 g (NH₄)₂SO₄. Under these conditions the glucose and methanol are simultaneously utilized, with the induction of GAD₆₅ expression upon commencement of the glucose-methanol feed. Cells from the
15 fermentation vessel are analyzed for GAD₆₅ expression as described above for shake flask cultures.

Cells are removed from the fermentation vessel at certain time intervals and subsequently analyzed. Little GAD₆₅ expression is observed during growth on glucose. Exhaustion of glucose leads to low level expression of the GAD₆₅ protein; expression is enhanced by the addition of MeOH during feeding of the fermentation culture. The addition of methanol has
20 a clear stimulatory effect of the expresion of human GAD₆₅ driven by the methanol-responsive *AUG1* promoter.

Example 4

Transformation conditions were investigated to determine the electric field
25 conditions. DNA topology, and DNA concentration that were optimal for efficient transformation of *P. methanolic*. All experiments used *P. methanolic* *ade2* strain #11. Competent cells were prepared as previously described. Electroporation was carried out using a BioRad Gene Pulser™.

Three field parameters influence transformation efficiency by electroporation: capacitance, field strength, and pulse duration. Field strength is determined by the voltage of the
30 electric pulse, while the pulse duration is determined by the resistance setting of the instrument. Within this set of experiments, a matrix of field strength settings at various resistances was examined. In all experiments, the highest capacitance setting (25 µF) of the instrument was used. 100 µl aliquots of electrocompetent cells were mixed on ice with 10 µl of DNA that contained approximately 1 µg of the *ADE2* plasmid pCZR133 that had been linearized with the restriction
35 enzyme Not I. Cells and DNA were transferred to 2 mm electroporation cuvettes (BTX Corp., San Diego, CA) and electropulsed at field strengths of 0.5 kV (2.5 kV/cm), 0.75 kV (3.75 kV/cm), 1.0 kV (5.0 kV/cm), 1.25 kV (6.25 kV/cm), and 1.5 kV (7.5 kV/cm). These field strength conditions were examined at various pulse durations. Pulse duration was manipulated by varying the instrument setting resistances to 200 ohms, 600 ohms, or "infinite" ohms. Pulsed cells were
40 suspended in YEPD and incubated at 30°C for one hour, harvested, resuspended, and plated. Three separate sets of experiments were conducted. In each set, electroporation conditions of 0.75 kV

(3.75 kV/cm) at a resistance of "infinite" ohms was found to give a dramatically higher transformation efficiency than other conditions tested (see Fig. 1).

After the optimal pulse conditions were established, the influence of DNA topology on transformation efficiency was investigated. Electrocompetent cells were mixed with 1 µg of uncut, circular pCZR133 or with 1 µg of Not I-digested pCZR133. In three separate experiments, an average of roughly 25 transformants were recovered with circular DNA while linear DNA yielded an average of nearly 1×10^4 transformants. These data indicate that linear DNA transforms *P. methanolica* with much greater efficiency than circular DNA.

Finally, the relationship between DNA concentration and transformation efficiency was investigated. Aliquots of linear pCZR133 DNA (1 ng, 10 ng, 100 ng and 1 µg in 10 µl H₂O) were mixed with 100 µl electrocompetent cells, and electroporation was carried out at 3.75 kV/cm and "infinite" ohms. The number of transformants varied from about 10 (1 ng DNA) to 10^4 (1 µg DNA) and was found to be proportional to the DNA concentration.

Example 5

Integration of transforming DNA into the genome of *P. methanolica* was detected by comparison of DNA from wild-type cells and stable, white transformant colonies. Two classes of integrative transformants were identified. In the first, transforming DNA was found to have integrated into a homologous site. In the second class, transforming DNA was found to have replaced the endogenous *AUG1* open reading frame. While not wishing to be bound by theory, this second transformant is believed to have arisen by a "transplacement recombination event" (Rothstein, *Methods Enzymol.* 194:281-301, 1991) whereby the transforming DNA replaces the endogenous DNA via a double recombination event.

P. methanolica ade2 strain #11 was transformed to Ade⁺ with Asp I-digested pCZR140, a Bluescript® (Stratagene Cloning Systems, La Jolla, CA)-based vector containing the *P. methanolica ADE2* gene and a mutant of *AUG1* in which the entire open reading frame between the promoter and terminator regions has been deleted (Fig. 2). Genomic DNA was prepared from wild-type and transformant cells grown for two days on YEPD plates at 30°C. About 100-200 µl of cells was suspended in 1 ml H₂O, then centrifuged in a microcentrifuge for 30 seconds. The cell pellet was recovered and resuspended in 400 µl of SCE + DTT + zymolyase (1.2 M sorbitol, 10 mM Na citrate, 10 mM EDTA, 10 mM DTT, 1-2 mg/ml zymolyase 100T) and incubated at 37°C for 10-15 minutes. 400 µl of 1% SDS was added, and the solution was mixed until clear. 300 µl of 5 M potassium acetate, pH 8.9 was added, and the solution was mixed and centrifuged at top speed in a microcentrifuge for five minutes. 750 µl of the supernatant was transferred to a new tube and extracted with an equal volume of phenol/chloroform. 600 µl of the resulting supernatant was recovered, and DNA was precipitated by the addition of 2 volumes of ethanol and centrifugation for 15 minutes in the cold. The DNA pellet was resuspended in 50 µl TE (10 mM Tris pH 8, 1 mM EDTA) + 100 µg/ml RNAase for about 1 hour at 65°C. 10-µl DNA samples were digested with Eco RI (5 µl) in a 100 µl reaction volume at 37°C overnight. DNA was precipitated with ethanol, recovered by centrifugation, and resuspended in 7.5 µl TE + 2.5 µl 5X loading dye. The entire 10 µl volume was applied to one lane of a 0.7% agarose in 0.5 X TBE (10 X TBE is 108 g/L Tris base

7-9, 55 g/L boric acid, 8.3 g/L disodium EDTA) gel. The gel was run at 100 V in 0.5 X TBE containing ethidium bromide. The gel was photographed, and DNA was electrophoretically transferred to a positively derivatized nylon membrane (Nytran® N+, Schleicher & Schuell, Keene, NH) at 400 mA, 20 mV for 30 minutes. The membrane was then rinsed in 2 X SSC, blotted onto

5 denaturation solution for five minutes, neutralized in 2 X SSC, then cross-linked damp in a UV crosslinker (Stratalinker®, Stratagene Cloning Systems) on automatic setting. The blot was hybridized to a PCR-generated *AUG1* promoter probe using a commercially available kit (ECL™ kit, Amersham Corp., Arlington Heights, IL). Results indicated that the transforming DNA altered the structure of the *AUG1* promoter DNA, consistent with a homologous integration event (Fig. 2).

10 In a second experiment, *P. methanolicus* *ade 2* strain #11 was transformed to Ade⁺ with Not I-digested pCZR137, a vector containing a human GAD65 cDNA between the *AUG1* promoter and terminator (Fig. 3). Genomic DNA was prepared as described above from wild-type cells and a stable, white, Ade⁺ transformant and digested with Eco RI. The digested DNA was separated by electrophoresis and blotted to a membrane. The blot was probed with a PCR-

15 generated probe corresponding to either the *AUG1* open reading frame or the *AUG1* promoter. Results demonstrated that the *AUG1* open reading frame DNA was absent from the transformant strain, and that the *AUG1* promoter region had undergone a significant rearrangement. These results are consistent with a double recombination event (transplacement) between the transforming DNA and the host genome (Fig. 3).

20

Example 6

An *AUG1* strain of *P. methanolicus* is grown in high-density fermentation conditions. The fermentation vessel is charged with mineral salts by the addition of 57.8 g (NH₄)₂SO₄, 46.6 g KCl, 30.8 g MgSO₄·7H₂O, 8.6 g CaSO₄·2H₂O, 2.0 g NaCl, and 10 ml antifoam (PPG). H₂O is

25 added to bring the volume to 2.5 L, and the solution is autoclaved 40 minutes. After cooling, 350 ml of 50% glucose, 250 ml 10 X trace elements (Table 4), 210 ml of 30% NaPhosphate, 25 ml 200 µg/ml biotin, and 250 ml cell inoculum are added. Cells are batch-fed glucose or glucose/methanol in three phases. In phase 1, the cells receive 0.4%/L/hour glucose (w/v final fermentation volume) for 25 hours using 750 g glucose, 110 g (NH₄)₂SO₄, and 278 ml 10 X trace elements per 1.5 liter.

30 The cells are then given a transition feed of 0.2% glucose, 0.2% methanol/L/hour for 5 hours. The final glucose-supplemented methanol feed contains 0.1% glucose, 0.4% methanol/L/hr for 25 hours. Final biomass is about 300 g/L cell paste.

Example 7

For fermentation of a *P. methanolica aug1Δ* strain, the fermentation vessel is initially charged with mineral salts, glucose, phosphate, trace elements and biotin as disclosed in Example 6. 250 ml of cell inoculum is added. A glucose feed is prepared using 600 g glucose, 108 g (NH₄)₂SO₄, and 273 ml 10 X trace elements per 1.2 liter. The cells are batch-fed in three phases. In the first phase, the cells receive glucose for 12 to 25 hours at 0.4%/L/hour. The cells are then induced with a bolus addition of 1% methanol by weight and transitioned to methanol utilization with a mixed 0.2% glucose/0.1% methanol feed for 10 hours. In the third phase, a mixed feed of 0.2% glucose, 0.2% methanol is delivered for 15 hours.

Example 8

P. methanolica cells in which the *AUG1* gene had been disrupted by insertion of a GAD65 expression construct retained the ability to grow on methanol, indicating that a second alcohol oxidase gene was present. The second gene, designated *AUG2*, was identified by PCR. Sequence analysis of the 5' coding region of the gene showed that the N-terminus of the encoded protein was similar to those of known alcohol oxidase genes.

Strain MC GAD8, a transformant that grew very poorly on minimal methanol broth, was used as a source for cloning the *AUG2* gene. Genomic DNA was prepared from MC GAD8 and amplified with sense and antisense PCR primers specific for the *AUG1* open reading frame (8784, SEQ ID NO:5; 8787, SEQ ID NO:6). A product identical in size to the *AUG1* product but showing very low intensity on an analytical gel was obtained.

The putative *AUG2* PCR product was digested with a battery of restriction enzymes. Partial digestion by Eco RI and Pvu I, and the presence of several Bgl II sites suggested that the DNA was contaminated with small amounts of *AUG1*. To remove the contaminating *AUG1* DNA, the PCR mixture was cut with Eco RI and gel purified. Since the MC GAD 8 product did not appear to have an Eco RI site, it was unaffected. The resulting gel-purified DNA was reamplified and again analyzed by restriction digestion. The DNA gave a different restriction map from that of the *AUG1* PCR product.

Southern blot analysis was performed on genomic DNA from MC GAD8 and wild-type cells using either *AUG1* or *AUG2* open reading frame PCR fragments as probes. The *AUG2* probe hybridized at low stringency to the *AUG1* locus and at both low and high stringency to a second locus. The *AUG1* probe bound to both loci at low stringency, but bound predominantly to the *AUG1* locus at high stringency. These data indicated that the new PCR product from MC GAD8 was similar to but distinct from *AUG1*. Sequence analysis showed an 83% identity between *AUG1* and *AUG2* gene products.

To clone the *AUG2* genomic locus, PCR primers were designed from the original *AUG2* PCR fragment. Primers 9885 (SEQ ID NO:7) and 9883 (SEQ ID NO:8) were used to screen a *P. methanolica* genomic library. A positive clone bank pool was then probed with the original MC GAD8 PCR product. Cells were plated on 10 plates at about 5000 colonies/plate and grown overnight, then the plates were overlaid with filter discs (Hybond-N, Amersham Corp., Arlington Heights, IL). Colonies were denatured, neutralized, and UV cross-linked. Bacterial debris was

washed from the filters with 5X SSC, and the filters were again cross-linked. Blots were pre-hybridized in pairs at 42°C for 1 hour in 25 ml hybridization buffer. Approximately 250 ng of probe was then added to each pair of filters. Hybridization was conducted at 42°C for four hours. The blots were then washed in 500 ml of 0.1 X SSC, 6M urea, 0.4% SDS at 42°C for 10 minutes, four times. The blots were then neutralized with 500 ml of 2 X SSC at room temperature for 5 minutes, two rinses. The blots were then immersed in 100 ml development reagent (ECL, Amersham Corp.).

Positive colonies were picked and amplified using PCR primers 9885 (SEQ ID NO:7) and 9883 (SEQ ID NO:8) to confirm their identity. Positive pools were streaked on plates, and single colonies were rescreened by PCR. One colony was selected for further analysis (restriction mapping and sequencing). A partial sequence of the *AUG2* gene is shown in SEQ ID NO:9. As shown in SEQ ID NO:9, the *AUG2* sequence begins at the HindIII site a nucleotide 91. Nucleotides upstream from this position are vector sequence. The coding sequence begins at nucleotide 170.

Disruption of the *AUG2* gene had little effect on cell growth on methanol. Cells lacking both functional *AUG1* and *AUG2* gene products did not grow on methanol. Subsequent analysis showed that the *AUG1* gene product is the only detectable alcohol oxidase in cells grown in a fermentor.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: ZymoGenetics, Inc.
1201 Eastlake Avenue East
Seattle, Washington 98102
United States of America

(ii) TITLE OF INVENTION: PREPARATION OF PICHIA METHANOLICA AUXOTROPHIC
MUTANTS

(iii) NUMBER OF SEQUENCES: 9

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: ZymoGenetics, Inc.
(B) STREET: 1201 Eastlake Avenue East
(C) CITY: Seattle
(D) STATE: WA
(E) COUNTRY: USA
(F) ZIP: 98102

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parker, Gary E
(B) REGISTRATION NUMBER: 31-648
(C) REFERENCE/DOCKET NUMBER: 96-17

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206-442-6673
(B) TELEFAX: 206-442-6678

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3077 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CAGCTGCTCT GCTCCTTGAT TCGTAATTAA TGTTATCCTT TTA CTTTGAA CTCTTGTCGG	60
TCCCCAACAG GGATTCCAAT CGGTGCTCAG CGGGATTTC CATGAGGTTT TTGACAACCT	120
TATTGATGCT GCAAAAACCT TTTTAGCCGG GTTTAAGTAA CTGGGCAATA TTTCCAAAGG	180
CTGTGGGCGT TCCACACTCC TTGCTTTTCA TAATCTCTGT GTATTGTTTT ATTCGCATTT	240
TGATTCTCTT ATTACCAGTT ATGTAGAAAG ATCGGCAAAC AAAATATCAA CTTTTATCTT	300
GAACGCTGAC CCACGGTTTC AAATACTAT CAGAACTCTA TAGCTATAGG GGAAGTTTAC	360
TGCTTGCTTA AAGCGGCTAA AAAGTGTTC GCAAATTA AAAGCTGTGA CAAGTAGGAA	420
CTCCTGTAAA GGGCCGATTC GACTTCGAAA GAGCCTAAAA ACAGTGACTA TTGGTGACGG	480
AAAATTGCTA AAGGAGTACT AGGGCTGTAG TAATAAATAA TGGAACAGTG GTACAACAAT	540
AAAAGAATGA CGCTGTATGT CGTAGCCTGC ACGAGTAGCT CAGTGGTAGA GCAGCAGATT	600
GCAAATCTGT TGGTCACCGG TTCGATCCGG TCTCGGGCTT CCTTTTTTGC TTTTTCGATA	660
TTTGCCGGTA GGAAGCAAGG TCTAGTTTTC GTCGTTTCGG ATGGTTTACG AAAGTATCAG	720
CCATGAGTGT TTCCCTCTGG CTACCTAATA TATTATTGA TCGGTCTCTC ATGTGAATGT	780
TTCTTTCCAA GTTCGGCTTT CAGCTCGTAA ATGTGCAAGA AATATTTGAC TCCAGCGACC	840
TTTCAGAGTC AAATTAATTT TCGCTAACAA TTTGTGTTTT TCTGGAGAAA CCTAAAGATT	900

TAAGTGATAA GTCGAATCAA CATCTTTAAA TCCTTTAGTT AAGATCTCTG CAGCGGCCAG 960
TATTAACCAA TAGCATATTC ACAGGCATCA CATCGGAACA TTCAGAATGG ACTCGCAAAC 1020
TGTCGGGATT TTAGGTGGTG GCCAACTTGG TCGTATGATC GTTGAAGCTG CACACAGATT 1080
GAATATCAAA ACTGTGATTC TCGAAAATGG AGACCAGGCT CCAGCAAAGC AAATCAACGC 1140
TTTAGATGAC CATATTGACG GCTCATTCAA TGATCCAAAA GCAATTGCCG AATTGGCTGC 1200
CAAGTGTGAT GTTTTAACCG TTGAGATTGA ACATGTTGAC ACTGATGCGT TGGTTGAAGT 1260
TCAAAAGGCA ACTGGCATCA AAATCTTCCC ATCACCAGAA ACTATTTTCAT TGATCAAAGA 1320
TAAATACTTG CAAAAGAGC ATTTGATTAA GAATGGCATT GCTGTTGCCG AATCTTGTAG 1380
TGTTGAAAGT AGCGCAGCAT CTTTAGAAGA AGTTGGTGCC AAATACGGCT TCCATACAT 1440
GCTAAAATCT AGAACAATGG CCTATGACGG AAGAGGTAAT TTTGTTGTCA AAGACAAGTC 1500
ATATATACCT GAAGCTTTGA AAGTTTGA TGACAGGCCG TTATACGCCG AGAAATGGGC 1560
TCCATTTTCA AAGGAGTTAG CTGTTATGGT TGTGAGATCA ATCGATGGCC AAGTTTATTC 1620
CTACCCAAT GTTGAAACCA TCCACCAAAA CAACATCTGT CACACTGTCT TTGCTCCAGC 1680
TAGAGTTAAC GATACTGTCC AAAAGAAGGC CCAAATTTTG GCTGACAACG CTGTCAAATC 1740
TTTCCAGGT GCTGGTATCT TTGGTGTGA AATGTTTTTA TTACAAAATG GTGACTTATT 1800
AGTCAACGAA ATTGCCCCAA GACCTCACA TTCTGGTCAC TATACCATCG ACGCTTGTGT 1860
CACCTCGCAA TTTGAAGCTC ATGTTAGGGC CATTACTGGT CTACCCATGC CGAAGAACTT 1920
CACTTGTTTG TCGACTCCAT CTACCCAAGC TATTATGTTG AACGTTTTAG GTGGCGATGA 1980
GCAAAACGGT GAGTTCAAGA TGTGTAAG AGCACTAGAA ACTCCTCATG CTTCTGTTTA 2040
CTTATACGGT AAGACTACAA GACCAGGCAG AAAAATGGGT CACATTAATA TAGTTTCTCA 2100
ATCAATGACT GACTGTGAGC GTAGATTACA TTACATAGAA GGTACGACTA ACAGCATCCC 2160
TCTCGAAGAA CAGTACACTA CAGATTCCAT TCCGGGCACT TCAAGCAAGC CATTAGTCGG 2220

TGTCATCATG GGTTCGATT CGGACCTACC AGTCATGTCT CTAGGTTGTA ATATATTGAA 2280
GCAATTTAAC GTTCCATTG AAGTCACTAT CGTTTCCGCT CATAGAACCC CACAAAGAAT 2340
GGCCAAGTAT GCCATTGATG CTCCAAGAG AGGGTTGAAG TGCATCATTG CTGGTGCTGG 2400
TGGTGCCGCT CATTTACCGG GAATGGTTGC GGCATGACG CCGCTGCCTG TTATTGGTGT 2460
CCCTGTAAA GGCTCTACTT TGGATGGTGT TGATTTACTA CACTCCATCG TTCAAATGCC 2520
AAGAGGTATT CCTGTTGCTA CTGTGGCTAT TAACAATGCT ACTAACGCTG CCTTGCTAGC 2580
TATCACAATC TTAGGTGCCG GCGATCCAAA TACTTGTCTG CAATGGAAGT TTATATGAAC 2640
AATATGGAAT ATGAAGTTT GGGCAAGGCT GAAAAATTGG AAAATGGTGG ATATGAAGAA 2700
TACTTGAGTA CATACAAGAA GTAGAACCTT TTATAATTGA TATAGTACTT ACTCAAAGTC 2760
TTAATTGTTT TAACTGTAA TTTCTGCTT GCATTTCTGA AAAGTTTAA ACAAGAAATC 2820
TTGAAATTC TAGTTGCTCG TAAGAGGAAA CTTGCATTCA AATAACATTA ACAATAAATG 2880
ACAATAATAT ATTATTTCAA CACTGCTATA TGGTAGTTTT ATAGGTTTGG TTAGGATTTG 2940
AGATATTGCT AGCGCTTATC ATTATCCTTA ATTGTTTCATC GACGCAAATC GACGCATTTT 3000
CACAAAAATT TTCCGAACCT GTTTTTCATC TCTCCAGATC TTGGTTTAGT ATAGCTTTTG 3060
ACACCTAATA CCTGCAG 3077

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3386 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GAATTCCTGC AGCCCGGGGG ATCGGGTAGT GGAATGCACG GTTATACCCA CTCCAAATAA	60
AAGTGTAGTA GCCGGACTGA AAGGTTTTAG GAGTCTGTTT GTTTGTTTAT GTGCATCATT	120
CCCTAATCTG TTAACAGTCT CGGAGTATAC AAAAAAGTAA GTCAAATATC AAGGTGGCCG	180
GGGGCAGCAT CGAGACTCGA GATGGTACAT ACTTAAAAGC TGCCATATTG AGGAACTTCA	240
AAGTTTATC TGTTTTTAGA ATTAAAAGAC GATTGTTGTA ACAAACGTT GTGCCTACAT	300
AAACTCAAAT TAATGGAAT AGCCTGTTTT GAAAAATACA CCTTCTTAAG TACTGACAAA	360
GTTTTGTAA ATGACTATCG AACAAGCCAT GAAATAGCAC ATTTCTGCCA GTCACTTTTA	420
ACACTTTCCT GCTTGCTGGT TGAATCTCCT CATACAAACA CCCAAAAGGG AAACCTTCAG	480
TGTGGGACA CTTGACATCT CACATGCACC CCAGATTAAT TTCCCAGAC GATGCGGAGA	540
CAAGACAAA CAACCTTTG TCCTGCTCTT TTCTTTCTCA CACCGCGTGG GTGTGTGCGC	600
AGGCAGGCAG GCAGGCAGCG GGCTGCCTGC CATCTCTAAT CGCTGCTCCT CCCCCCTGGC	660
TTCAAATAAC AGCCTGCTGC TATCTGTGAC CAGATTGGGA CACCCCCCTC CCCTCCGAAT	720
GATCCATCAC CTTTGTGCT ACTCCGACAA TGATCCTTCC CTGTCATCTT CTGGCAATCA	780
GCTCCTTCAA TAATTAAATC AAATAAGCAT AAATAGTAAA ATCGCATACA AACGTCATGA	840
AAAGTTTAT CTCTATGGCC AACGGATAGT CTATCTGCTT AATCCATCC ACTTTGGGAA	900
CCGCTCTCTC TTTACCCAG ATTCTCAAAG CTAATATCTG CCCCTTGCTT ATTGTCCTTT	960
CTCCGTGTAC AAGCGGAGCT TTGCTCTCC ATCCTCTGC TTTGTTTCGG TTATTTTTTT	1020
TTCTTTTGAA ACTCTTGGTC AAATCAAATC AAACAAAACC AAACCTTCTA TTCCATCAGA	1080
TCAACCTTGT TCAACATTCT ATAAATCGAT ATAAATATAA CCTATCCCT CCCTGTTTTT	1140
TTACCAATTA ATCAATCTTC AAATTTCAA TATTTTCTAC TTGCTTTATT ACTCAGTATT	1200
AACATTTGTT TAAACCAACT ATAACTTTTA ACTGGCTTTA GAAGTTTAT TTAACATCAG	1260

TTTCAATTTA CATCTTTATT TATTAACGAA ATCTTTACGA ATTAAC TCAA AACTTT 1320
TACGAAAAAA AAATCTTACT ATTAATTTCT CAAAATGGCT ATTCCAGATG AATTGATAT 1380
TATTGTTGTC GGTGGTGGTT CCACCGGTTG TGCTCTTGCT GGTAGATTAG GTAAC TTGGA 1440
CGAAAACGTC ACAGTTGCTT TAATCGAAGG TGGTGAAAAC AACATCAACA ACCCATGGGT 1500
TTACTTACCA GGTGTTTATC CAAGAAACAT GAGATTAGAC TCAAAGACTG CTACTTTTIA 1560
CTCTTCAAGA CCATCACCAC ACTTGAACGG TAGAAGAGCT ATTGTTCCAT GTGCTAACAT 1620
CTTGGTGGT GGTTCCTCCA TCAACTTCTT GATGTACACC AGAGCCTCTG CCTCCGATTA 1680
CGATGATTGG GAATCTGAAG GTTGGACTAC CGATGAATTA TTACCACTAA TGAAGAAGAT 1740
TGAAACTTAT CAAAGACCAT GTAACAACAG AGAATTGCAC GGTTTCGATG GTCCAATTAA 1800
GGTTTCATTT GGTAACATA CTTATCCAAA CGGTCAAGAT TTCATTAGAG CTGCCGAATC 1860
TCAAGGTATT CCATTTGTTG ATGATGCTGA AGATTGAAA TGTTCCACG GTGCTGAGCA 1920
CTGGTTGAAG TGGATCAACA GAGACTTAGG TAGAAGATCC GATTCTGCTC ATGCTTACAT 1980
TCACCAACC ATGAGAAAA AGCAAACTT GTTCTTGATT ACTCCACCA AGTGTGAAAA 2040
GATTATCATT GAAAACGGTG TTGCTACTGG TGTTAAGACT GTTCCAATGA AGCCAAC TGG 2100
TTCTCAAAG ACCCAAGTTG CTAGAACTTT CAAGGCTAGA AAGCAAATTA TTGTTTCTTG 2160
TGGTACTATC TCATCACCAT TAGTTTTGCA AAGATCTGGT ATCGGTTCCG CTCACAAGTT 2220
GAGACAAGTT GGTATTAAAC CAATTGTTGA CTTACCAGGT GTTGGTATGA ACTTCCAAGA 2280
TCACTACTGT TTCTTCACTC CATACCATGT CAAGCCAGAT ACTCCATCAT TCGATGACTT 2340
TGTTAGAGGT GATAAAGCTG TTCAAAAATC TGCTTTCGAC CAATGGTATG CTAACAAGGA 2400
TGGTCCATTA ACCACTAATG GTATTGAGGC AGGTGTTAAG ATTAGACCAA CTGAAGAAGA 2460
ATTAGCCACT GCTGATGACG AATTCAGAGC TGCTTATGAT GACTACTTTG GTAACAAGCC 2520
AGATAAGCCA TTAATGCACT ACTCTCTAAT TTCTGGTTTC TTTGGTGACC ACACCAAGAT 2580

TCCAAACGGT AAGTACATGT GCATGTTCCA CTTCTTGGAA TATCCATTCT CCAGAGGTTT 2640
CGTTCACGTT GTTTCCTCAA ACCCATACGA TGCTCCTGAC TTTGATCCAG GTTTCATGAA 2700
CGATCCAAGA GATATGTGGC CAATGGTTTG GTCTTACAAG AAGTCCAGAG AAATGCCAG 2760
AAGAATGGAC TGTTTTGCCG GTGAAGTTAC TTCTCACCAC CCACACTACC CATACGACTC 2820
ACCAGCCAGA GCTGCTGACA TGGACTTGA AACTACTAAA GCTTATGCTG GTCCAGACCA 2880
CTTTACTGCT AACTTGTAAC ACGGTTTCATG GACTGTTCCA ATTGAAAAGC CAACTCCAAA 2940
GAACGCTGCT CACGTTACTT CTAACCAAGT TGAAAAACAT CGTGACATCG AATACACCAA 3000
GGAGGATGAT GCTGCTATCG AAGATTACAT CAGAGAACAC ACTGAAACCA CATGGCATTG 3060
TCTTGGTACT TGTTCATGG CTCCAAGAGA AGGTTCTAAG GTTGTCCCAA CTGGTGGTGT 3120
TGTGACTCC AGATTAAACG TTTACGGTGT TGAAAAGTTG AAGGTTGCTG ATTTATCAAT 3180
TTGCCCAGAT AATGTTGGTT GTAACACTTA CTCTACTGCT TTGTTAATCG GTGAAAAGGC 3240
TTCTACCTTA GTTGCTGAAG ACTTGGGCTA CTCTGGTGAT GCTTTGAAGA TGA CTGTTCC 3300
AAACTCAAA TTGGGTACTT ATGAAGAAGC TGGTCTAGCT AGATTCTAGG GCTGCCTGTT 3360
TGGATATTTT TATAATTTT GAGAGT 3386

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGATCACCTA GGACTAGTGA CAAGTAGGAA CTCCTGTA

38

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CAGCTGCCTA GGACTAGTTT CCTCTTACGA GCAACTAGA

39

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TGGTTGAAGT GGATCAA

17

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GTGTGGTCAC CGAAGAA

17

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC9885

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GTTGTTTCCTT CCAAACCATT GAAC

24

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:
(B) CLONE: ZC9883

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AAAGTAAGAA GCGTAGCCTA GTTG

24

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 329 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GACCATGATT ACGCCAAGCG CGCAATTAAC CCTCACTAAA GGGAACAAAA GCTGGGTACC 60

GGGCCCCCCC TCGAGGTCGA CGGTATCGAT AAGCTTTATT ATAACATTAA TATACTATTT 120

TATAACAGGA TTGAAAATTA TATTTATCTA TCTAAACTA AAATTCAAAA TGGCTATTCC 180

TGAAGAATTC GATATCATTG TTGTCGGTGG TGGTTCTGCC GGCTGTCCTA CTGCTGGTAG	240
ATTGGCTAAC TTAGACCCAA ATTTAACTGT TGCTTTAATC GAAGCTGGTG AAAACAACAT	300
TAACAACCCA TGGGTCTACT TACCAGGCG	329

Claims

What is claimed is:

1. A method for preparing *Pichia methanolica* cells having an auxotrophic mutation comprising:

- (a) exposing *P. methanolica* cells to mutagenizing conditions;
- (b) culturing the cells from step (a) in a rich medium to allow mutations to become established and replicated in at least a portion of said cells;
- (c) culturing the cells from step (b) in a culture medium deficient in assimilable nitrogen to deplete cellular nitrogen stores;
- (d) culturing the cells from step (c) in a defined culture medium comprising an inorganic nitrogen source and an amount of nystatin sufficient to kill growing *P. methanolica* cells to select for cells having a deficiency in a nutritional gene; and
- (e) culturing the selected cells from step (d) in a rich culture medium.

2. A method according to claim 1 wherein the selected cells from step (e) are replica plated to a defined medium and cultured to confirm the presence of an auxotrophic mutation.

3. A method according to claim 1 wherein the selected cells are auxotrophic for adenine.

4. A method according to claim 3 wherein the selected cells are deficient in phosphoribosyl-5-aminoimidazole carboxylase.

5. A method according to claim 1 wherein the defined culture medium contains 2 mg/L nystatin.

6. A method according to claim 1 wherein the mutagenizing conditions comprise exposure to ultraviolet light.

7. A method according to claim 1 wherein the mutagenizing conditions comprise exposure to a chemical mutagen.

8. A method according to claim 1 wherein the inorganic nitrogen source comprises ammonium ions.

9. A method according to claim 8 wherein the inorganic nitrogen source is ammonium sulfate.

10. A method for preparing *Pichia methanolica* cells having an auxotrophic mutation substantially as hereinbefore described with reference to any one of the
5 Examples.

11. *Pichia methanolica* cells having an auxotrophic mutation prepared by the method of any one of claims 1 to 10.

Dated 4 February, 1999

ZymoGenetics, Inc.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
219

1/3




Fig. 2




Fig. 3

